

# INTERIOR CONTROLLABILITY OF SURFACE GRAVITY WAVES AND THE SLOSHING PROBLEM

M. A. FONTELOS AND J. LÓPEZ-RÍOS

ABSTRACT. We study the problem of controlling the free surface, by fluid jets on the boundary, for a two dimensional solid container in the context of the gravity waves and the sloshing problem. By using conformal maps and the Dirichlet–Neumann operator, the problem is formulated as a second order evolutionary equation on the free surface involving a self-adjoint operator. We present then the appropriate Sobolev spaces where having solutions for the system and study the exact controllability through an observability inequality for the adjoint problem.

## 1. INTRODUCTION

Finding and controlling the frequencies for the oscillations of the free boundary in the context of gravity waves and studying related spectral problems is a classical, well known problem in the literature [5, 8, 9, 18]. Roughly speaking, controlling a system consist not only in testing that its behavior is satisfactory, but also in putting things in order to guarantee that it behaves as desired. In mathematical terms, controlling the state  $y$ , ruled by the state equation

$$A(y) = f(v),$$

where  $v$  is the control, consists in finding  $v \in U_{ad}$ , the set of admissible controls, such that the solution to the equation gets close to a desired prescribed state,  $\bar{y}$ .

In this paper, we study the interior exact controllability of a Partial Differential Equation (PDE), in the context of controlling the oscillations of a liquid free surface in a two-dimensional bounded container and the so-called sloshing problem. We formulate the geometrical problem in terms of an integrodifferential equation by using the Hilbert transform, then we establish the appropriate Sobolev spaces to study existence of solutions for the eigenvalue problem and, finally, we set up an observability inequality for the homogeneous adjoint problem. The sloshing problem adduces to an important difference with respect to the classical water-waves formulation: the presence of vertical walls and the contact with the free surface. Inspired in the developments for the classical wave equation, we introduce analytical tools to prove that it is possible to control the oscillations of the free surface, by injecting fluid on the rigid side walls.

The main strengths of our method lie in the use of the Hilbert transform to formulate the problem as an evolutionary equation involving a self-adjoint operator. This is known as the boundary integral method and has proved to be very fruitful in the study of water waves problems (see [7] and references therein, for instance). Moreover, the use of Tchebyshev polynomials provides an explicit orthogonal basis which allows to study, analytically, the associated eigenvalue problem. Then, an observability inequality arises naturally.

Generally speaking, the water-waves problem for an ideal liquid consists of describing the motion of a layer of incompressible, inviscid fluid, delimited below by a solid bottom, and above by a free surface under the influence of gravity. In mathematical terms, if  $\mathbf{u}(x, y)$  is the fluid velocity and  $\varphi$  is the velocity potential such that  $\mathbf{u} = \nabla\varphi$ , by the conservation laws [13]

$$\begin{cases} \Delta\varphi = 0, & \Omega_t, \\ \eta_t + \eta_x\varphi_x = \varphi_y, & y = \eta, \\ \varphi_t + \frac{1}{2}(\varphi_x^2 + \varphi_y^2) + g\eta = 0, & y = \eta, \\ \partial_n\varphi = 0, & y = b, \end{cases} \quad (1) \quad \boxed{ww}$$

where  $\eta$  and  $b$  are the free boundary and bottom parametrization, respectively,  $g$  is the acceleration due to gravity and  $\Omega_t = \{(x, y) \in \mathbb{R}^2 : b(x) < y < \eta(t, x)\}$ . We put ourselves in the general situation described in

Date: November 3, 2021.

Key words and phrases. Controllability; gravity waves; the sloshing problem; Hilbert transform.

M. A. Fontelos was partially supported by Ministerio de Economía, Industria y Competitividad, Gobierno de España (Grant No. MTM2017–89423–P).

(1). However, we consider the case of a bounded domain and the sloshing problem of describing the contact line between the free surface and the solid walls [3, 10]. Two conditions are customary, the pinned-end and the free-end boundary conditions. We will consider both and will impose conditions on the Cauchy problem to deal with it, accordingly.

When we talk about the controllability by fluid injection, we mean the condition

$$\frac{\partial \varphi}{\partial n} = j,$$

for a given function  $j = j(t, x)$ , with  $n$  being the outward normal vector. This boundary condition lead to think of a boundary control of the gravity waves problem; nevertheless, we will restate the problem as an integrodifferential equation on the free surface and the boundary condition becomes a source term. Then, we may use the classical approach of interior controllability by means of the adjoint problem and the observability inequality [15, 17].

Following the methods introduced in [8], after linearizing (1), the problem is restated through a conformal map into the lower half-plane. We link the normal derivative to the specific conformal map and rewrite the problem as a second order evolution equation on the interval  $[-1, 1]$ , where the Hilbert transform is involved. In [8], we used this approach to propose an efficient computational method to find the sloshing frequencies on general 2d domains. We mention [12] where the capillary version of the problem is developed, under the context of the oscillations in a nozzle of an inkjet printer. In these works, two possibilities for the contact line were considered: the ‘pinned-end edge condition’, where the contact line is always pinned to the solid surface, and the ‘free-end condition’ where the contact angle between the fluid-air interface and the side walls is fixed and the contact line is allowed to move, with contact angle  $\pi/2$ .

By addressing this problem, we give an answer to a question raised in [9]; namely, the problem of controlling undesirable splashing appearing in a cooper converter when air is injected into the molten matte. There, the authors studied the problem numerically by using triangular finite elements to mesh the bounded domain, on a damped linear gravity waves model. Controlling the surface by different methods is of practical interest in oceanography, controllability and inverse problems theory. Among others, we mention [16], where the authors designed the ‘best’ moving solid bottom generating a prescribed wave under the context of a BBM-type equation and [1] where the local exact controllability of the full water-waves system is obtained by controlling a localized portion of the free surface. We mention also the possibility of studying the inverse problem of detecting the source  $j$  by measuring the free surface as in [14].

With the present work, we have completed the numerical analysis started in [8]. Namely, we establish a Sobolev frame where the Cauchy problem is well-posed and study there the eigenvalue associated problem. We also prove an observability inequality for the adjoint problem and explore the possibility of finding explicit controls taking the oscillations at the free surface to zero.

The rest of the paper is organized as follows: In 2 we formulate the general equations to be considered, the linearization approach and corresponding formulation on the half-plane by the conformal mapping. In 3 we use the Hilbert transform to state a second order evolutionary PDE modeling the dynamics of the fluid interface on the bounded domain  $[-1, 1]$ . In 4 we make use of the Tchebyshev polynomials to study the stationary adjoint problem in suitable Sobolev spaces. In 5 we prove an observability inequality for the adjoint problem. Finally, in 6 we establish the controllability of the problem and explore the possibility of determining possible control functions, explicitly.

## 2. FORMULATION

S2

Let us consider a two dimensional container, filled with water, bounded from above by a free surface. In this context, the motion is governed by the incompressible Euler equations with zero surface tension:

$$\begin{aligned} \nabla \cdot \mathbf{u} &= 0, \\ \rho \left[ \frac{\partial \mathbf{u}}{\partial t} + (\mathbf{u} \cdot \nabla) \mathbf{u} \right] &= -\nabla p - g e_2, \end{aligned}$$

with  $-g e_2$  being the constant acceleration of gravity,  $g > 0$ , and  $e_2$  the unit upward vector in the vertical direction,  $\rho$  is the (constant) density of the fluid and  $p$  is the pressure inside. We use the classical notation  $(x, y) \in \mathbb{R}^2$  and  $z = x + iy$  for complex numbers.




FIGURE 1. Geometry of the problem.

F1

By considering the potential function  $\varphi$  of  $\mathbf{u}$ , so that  $\mathbf{u} = \nabla\varphi$ :

$$\Delta\varphi = 0, \quad (2) \quad \text{pf2}$$

$$\frac{\partial\varphi}{\partial t} + \frac{1}{2}|\nabla\varphi|^2 + \frac{1}{\rho}p + gy = \text{const.} \quad (3)$$

We complement system above with the nonzero Neumann condition at the solid walls,  $\mathbf{u} \cdot \mathbf{n} = j(t, x)$ , and a kinematic condition on the free boundary. In terms of  $\varphi$ :

$$\frac{\partial\varphi}{\partial n} = j(t, x), \quad \text{at the solid boundary} \quad (4) \quad \text{pf3}$$

$$\eta_t = \frac{\partial\varphi}{\partial n}, \quad |x| < 1. \quad (5) \quad \text{pf4}$$

ss1

**2.1. Linearized equations.** Next, we are going to linearize Euler's system around the zero state. This will allow us to obtain a single, explicit, evolutionary conservation law on the free surface modeling the fluid dynamics of the free surface side bounded by vertical solid walls. This equation must be complemented by boundary conditions as explained in 3.

Even if we consider the reference domain  $\Omega = \{(x, y) \in \mathbb{D} : y' < 0\}$ , other domains can be considered though. Let

$$\eta(t, x) = \epsilon\zeta(t, x).$$

Then if

$$\varphi = \text{const.} + \epsilon\phi,$$

conditions (2)-(4) in terms of  $\phi$ , at the first order for  $\epsilon \ll 1$ , become (after re-scaling to make  $g = 1$ )

$$\Delta\phi = 0, \quad (6) \quad \text{le1}$$

$$\phi_t + \zeta = 0, \quad \text{at } y = 0, \quad |x| \leq 1, \quad (7) \quad \text{le2}$$

$$\zeta_t = \frac{\partial\phi}{\partial n}, \quad \text{at } y = 0, \quad |x| \leq 1, \quad (8) \quad \text{le3}$$

$$\frac{\partial\phi}{\partial n} = \frac{1}{\epsilon}j(t, x), \quad \text{on the solid walls.} \quad (9) \quad \text{le4}$$

Therefore, from (7)-(8), on the free boundary and for  $|x| \leq 1$ ,

$$\phi_{tt} + \frac{\partial\phi}{\partial n} = 0. \quad (10) \quad \text{le5}$$

**2.2. Conformal transplants.** As is explained in [\[2, 8\]](#), let  $\psi$  be a real-valued function written as [\[asmarr2002applied, fontelos2020gravity\]](#)

$$\psi : (x, y) \rightarrow \psi(x, y) = \psi(z)$$

be defined in a domain  $D$ . Also, let  $\tilde{\psi}$  be defined in  $\tilde{D}$  as follows: for any  $w \in \tilde{D}$  we have

$$\tilde{\psi}(w) := \psi(f^{[-1]}(w)) = \psi(x(x', y'), y(x', y')).$$

Then the following relation between the normal derivatives holds:

$$\frac{\partial\tilde{\psi}}{\partial\tilde{n}} = \left| \frac{dz}{dw} \right| \frac{\partial\psi}{\partial n} = \frac{1}{|f'(z)|} \frac{\partial\psi}{\partial n}. \quad (11) \quad \text{ct8}$$

**2.3. Reformulation as an integrodifferential PDE.** In the particular case of the half-cylinder geometry,  $\Omega$ , we use the conformal map

$$w = f(z) = \frac{1}{\frac{1}{2} \left( z + \frac{1}{z} \right)},$$

into the half plane  $\Omega' := \{(x', y') \in \mathbb{R}^2 : y' < 0\}$ .

From (11), equations (6), (9), and (10), in variables  $w = x' + iy'$  become (see (1))

$$\Delta \tilde{\phi} = 0, \quad \text{for } y' < 0, \quad (12) \quad \text{ie5}$$

$$\tilde{\phi}_{tt} + |f'(x')| \frac{\partial \tilde{\phi}}{\partial \tilde{n}} = 0, \quad \text{at } y' = 0, |x'| \leq 1, \quad (13) \quad \text{ie52}$$

$$\frac{\partial \tilde{\phi}}{\partial \tilde{n}} = \frac{\tilde{j}(t, x')}{\epsilon |f'(x')|}, \quad \text{at } y' = 0, |x'| > 1, \quad (14) \quad \text{ie53}$$

where  $j = \tilde{j} \circ f$ . That is, for  $|x'| > 1$ ,  $x' = f(x)$ .

We complement system above with the following boundary conditions at infinity:

$$\partial_{x'} \tilde{\phi}, \partial_{y'} \tilde{\phi} \rightarrow 0, \quad \text{as } y' \rightarrow -\infty \text{ or } |x'| \rightarrow \infty.$$

By taking Fourier transform in  $x'$  of (12), denoted by  $\tilde{\Phi}$ , we have

$$\tilde{\Phi}_{y'y'} - k^2 \tilde{\Phi} = 0,$$

which implies

$$\begin{aligned} \tilde{\Phi}(t, k, y') &= \tilde{\Phi}(t, k, 0) e^{|k|y'} \\ &= \tilde{\Phi}(t, k, 0) \frac{\widehat{-\sqrt{2}y'}}{\sqrt{\pi}(x'^2 + y'^2)}. \end{aligned} \quad (15) \quad \text{ie6}$$

By taking inverse Fourier transform

$$\tilde{\phi}(t, x', y') = -\frac{y'}{\pi} \int_{-\infty}^{+\infty} \frac{\tilde{\phi}(t, \xi, 0)}{(x' - \xi)^2 + y'^2} d\xi.$$

Since we want to establish properties on the normal derivative,  $\tilde{\phi}_{y'}$ , taking the  $y'$  derivative in (15) instead, and evaluating at  $y' = 0$  we find

$$\begin{aligned} \tilde{\Phi}_{y'}(t, k, 0) &= \frac{1}{i} \operatorname{sgn}(k) (ik) \tilde{\Phi}(t, k, 0), \\ &= \frac{\widehat{\sqrt{2}}}{\sqrt{\pi} x'} \tilde{\Phi}_{x'}(t, k, 0). \end{aligned} \quad (16) \quad \text{ie7}$$

Then, taking inverse Fourier transform

$$\tilde{\phi}_{y'} \Big|_{y'=0} = \frac{1}{\pi} P.V. \int_{-\infty}^{+\infty} \frac{\tilde{\phi}_{x'}(t, \xi, 0)}{x' - \xi} d\xi = H \left( \tilde{\phi}_{x'} \Big|_{y'=0} \right). \quad (17) \quad \text{ie8}$$

Since  $HH = -I$  (see [\[11\]](#)):

$$\tilde{\phi}_{x'} \Big|_{y'=0} = -\frac{1}{\pi} P.V. \int_{-\infty}^{\infty} \frac{\tilde{\phi}_{y'}(t, \xi, 0)}{x' - \xi} d\xi = -\frac{1}{\pi} P.V. \int_{-\infty}^{\infty} \frac{\frac{\partial \tilde{\phi}}{\partial \tilde{n}}(t, \xi, 0)}{x' - \xi} d\xi. \quad (18) \quad \text{ie9}$$

**2.4. The Dirichlet-Neumann operator.** Let us consider in this section the basic situation when  $\frac{\partial \phi}{\partial n} = 0$ , on the solid walls of the domain  $\Omega$ . The value of  $\frac{\partial \phi}{\partial n}$  at the fluid interface may be viewed as the result of an operator (the so-called Dirichlet-Neumann operator) acting on the function  $\phi$  restricted to the interface. We are going to deduce a few simple consequences obtained from the properties of this operator. Keeping in mind the equation (8), we can easily deduce the following mass conservation relation:

$$\frac{d}{dt} \int_{-1}^1 \zeta dx = \int_{-1}^1 \frac{\partial \phi}{\partial n} dx = \int_{\partial \Omega} \frac{\partial \phi}{\partial n} = \int_{\Omega} \nabla \cdot (\nabla \phi) = 0. \quad (19) \quad \text{dn1}$$

Notice then that, in the mapped coordinates, by (11)

$$0 = \int_{-1}^1 \frac{\partial \phi}{\partial n} dx = \int_{-1}^1 |f'(x')| \frac{\partial \tilde{\phi}}{\partial \tilde{n}} \frac{dx'}{|f'(x')|} = \int_{-1}^1 \frac{\partial \tilde{\phi}}{\partial \tilde{n}} dx', \quad (20) \quad \boxed{\text{dn2}}$$

and hence, from (19) and (11)

$$\frac{d}{dt} \int_{-1}^1 \tilde{\zeta} \frac{dx'}{|f'(x')|} = 0.$$

Next, since  $\frac{\partial \phi}{\partial n} = 0$  at the solid boundaries of  $\Omega$ , we have

$$\int_{-1}^1 \psi \frac{\partial \phi}{\partial n} dx = \int_{\partial \Omega} \psi \frac{\partial \phi}{\partial n} dx$$

for any harmonic function  $\psi$  also vanishing at the solid boundaries of  $\Omega$ , and therefore by Green's identity we deduce

$$\int_{\partial \Omega} \psi \frac{\partial \phi}{\partial n} dx - \int_{\partial \Omega} \phi \frac{\partial \psi}{\partial n} dx = \int_{\Omega} (\psi \Delta \phi - \phi \Delta \psi) = 0,$$

which implies

$$\int_{\partial \Omega} \psi \frac{\partial \phi}{\partial n} dx = \int_{\partial \Omega} \phi \frac{\partial \psi}{\partial n} dx,$$

or, equivalently,

$$\int_{-1}^1 \tilde{\psi} \frac{\partial \tilde{\phi}}{\partial \tilde{n}} dx' = \int_{-1}^1 \tilde{\phi} \frac{\partial \tilde{\psi}}{\partial \tilde{n}} dx',$$

showing that the Dirichlet-Neumann operator, mapping  $\tilde{\phi}$  at the free surface into  $\frac{\partial \tilde{\phi}}{\partial \tilde{n}}$ , is selfadjoint.

Finally, from (7) and (19),

$$\int_{-1}^1 \phi_t dx = - \int_{-1}^1 \zeta dx = 0, \quad (21) \quad \boxed{\text{dn4}}$$

which implies

$$\frac{d}{dt} \int_{-1}^1 \phi(t, x) dx = 0.$$

Therefore, by choosing  $\phi(0, x)$  such that  $\int_{-1}^1 \phi(0, x) dx = 0$  (this can always be achieved by adding a suitable constant to a given  $\phi$ ), we have

$$\int_{-1}^1 \phi(t, x) dx = 0,$$

implying

$$\int_{-1}^1 \tilde{\phi}(t, x') \frac{dx'}{|f'(x')|} = 0. \quad (22) \quad \boxed{\text{dn5}}$$

### 3. AN ASSOCIATED CAUCHY PROBLEM

S3

As in the last section, let  $\frac{\partial \phi}{\partial n} = 0$  on the solid walls of the domain  $\Omega$ . Then, from the integrodifferential formulation (18), we obtain

$$\tilde{\phi}_{x'} \Big|_{y'=0} = -\frac{1}{\pi} P.V. \int_{-1}^1 \frac{\tilde{\phi}_{y'}(t, \xi, 0)}{x' - \xi} d\xi. \quad (23) \quad \boxed{\text{cp1}}$$

We observe two properties of  $\tilde{\phi}_x$ . First, making use of the identity

$$\frac{1}{\pi} P.V. \int_{-1}^1 \frac{1}{\sqrt{1-x'^2}} \frac{dx'}{x' - \xi} = 0, \text{ for } x' \in [-1, 1], \quad (24)$$

we prove

$$\begin{aligned} \int_{-1}^1 \frac{\tilde{\phi}_{x'}(x')}{\sqrt{1-x'^2}} dx' &= -\frac{1}{\pi} \int_{-1}^1 \frac{1}{\sqrt{1-x'^2}} \left( P.V. \int_{-1}^1 \frac{\tilde{\phi}_{y'}(\xi)}{x' - \xi} d\xi \right) dx' \\ &= -\frac{1}{\pi} \int_{-1}^1 \tilde{\phi}_{y'}(\xi) \left( P.V. \int_{-1}^1 \frac{1}{\sqrt{1-x'^2}(x' - \xi)} dx' \right) d\xi \\ &= 0 \end{aligned} \quad (25) \quad \boxed{\text{n1}}$$

and secondly, when  $\int_{-1}^1 \tilde{\phi}_{y'}(\xi) d\xi = 0$ , and using

$$\frac{1}{\pi} P.V. \int_{-1}^1 \frac{x'}{\sqrt{1-x'^2}} \frac{dx'}{x'-\xi} = 1, \text{ for } x' \in [-1, 1],$$

we also have

$$\begin{aligned} \int_{-1}^1 \frac{x' \tilde{\phi}_{x'}(x')}{\sqrt{1-x'^2}} dx' &= -\frac{1}{\pi} \int_{-1}^1 \frac{x'}{\sqrt{1-x'^2}} \left( P.V. \int_{-1}^1 \frac{\tilde{\phi}_{y'}(\xi)}{x'-\xi} d\xi \right) dx' \\ &= -\frac{1}{\pi} \int_{-1}^1 \tilde{\phi}_{y'}(\xi) \left( P.V. \int_{-1}^1 \frac{x'}{\sqrt{1-x'^2}(x'-\xi)} dx' \right) d\xi \\ &= -\int_{-1}^1 \tilde{\phi}_{y'}(\xi) d\xi = 0. \end{aligned} \quad (26) \quad \boxed{\text{n2}}$$

By using the inverse transform instead, from (23) we have, for  $|x'| < 1$ ,

$$\tilde{\phi}_{y'} \Big|_{y'=0} = \sqrt{1-x'^2} \frac{1}{\pi} P.V. \int_{-1}^1 \frac{1}{\sqrt{1-\xi^2}} \frac{\tilde{\phi}_{x'}(t, \xi, 0)}{x'-\xi} d\xi + \frac{C}{\sqrt{1-x'^2}}, \quad (27) \quad \boxed{\text{cp2}}$$

where  $C$  is an arbitrary constant that, due to mass conservation and using (26), must be chosen to be zero:

$$\begin{aligned} 0 &= \int_{-1}^1 \tilde{\phi}_{y'}(x') dx' = \int_{-1}^1 \frac{\tilde{\phi}_{x'}(\xi)}{\sqrt{1-\xi^2}} \left( \frac{1}{\pi} P.V. \int_{-1}^1 \frac{\sqrt{1-x'^2}}{x'-\xi} dx' \right) d\xi + \pi C \\ &= -\int_{-1}^1 \frac{\xi \tilde{\phi}_{x'}(\xi)}{\sqrt{1-\xi^2}} d\xi + \pi C = \pi C. \end{aligned}$$

By formulas (78) and (79) in the appendix and using (25), (26) we conclude that the following three expressions are equivalent for our problem provided  $\tilde{\phi}$  is sufficiently regular:

$$\tilde{\phi}_{y'} \Big|_{y'=0} = \sqrt{1-x'^2} \frac{1}{\pi} P.V. \int_{-1}^1 \frac{1}{\sqrt{1-\xi^2}} \frac{\tilde{\phi}_{x'}(t, \xi, 0)}{x'-\xi} d\xi \quad (28) \quad \boxed{\text{ex1}}$$

$$= \frac{1}{\sqrt{1-x'^2}} \frac{1}{\pi} P.V. \int_{-1}^1 \sqrt{1-\xi^2} \frac{\tilde{\phi}_{x'}(t, \xi, 0)}{x'-\xi} d\xi \quad (29) \quad \boxed{\text{ex2}}$$

$$= \partial_{x'} \left( \sqrt{1-x'^2} \frac{1}{\pi} P.V. \int_{-1}^1 \frac{1}{\sqrt{1-\xi^2}} \frac{\tilde{\phi}(t, \xi, 0)}{x'-\xi} d\xi \right). \quad (30) \quad \boxed{\text{ex3}}$$

Expression (28) was used in our previous article [\[8\]](#), while expression (30) will be more convenient in the present work. We present next an alternative deduction of (30). We introduce the function  $\chi$  defined by

$$\tilde{\phi} = \chi_{x'}$$

so that

$$\chi = \int_{-\infty}^{x'} \tilde{\phi}(s, y') ds,$$

(note that integrability at  $-\infty$  is guaranteed by the decay of  $\tilde{\phi}$  from (18) after integration by parts and use that the integral of the normal derivative of  $\tilde{\phi}$  vanishes by mass conservation (20)) and

$$\Delta \chi = 0.$$

We have then

$$\chi_{y'} = \frac{\partial}{\partial y'} \int_{-\infty}^{x'} \tilde{\phi} dx' = \int_{-\infty}^{x'} (\chi_{y'})_{x'} dx' = \begin{cases} 0, & x' \leq -1, \\ \chi_{y'}, & -1 < x' \leq 1, \\ 0, & x' > 1. \end{cases}$$

Since  $\Delta \chi = 0$ , by (23) we have

$$\chi_{x'}(x') = -\frac{1}{\pi} P.V. \int_{-1}^1 \frac{\chi_{y'}(\xi)}{x'-\xi} d\xi. \quad (31) \quad \boxed{\text{f1}}$$

Therefore, inverting Hilbert transform

$$\chi_{y'}(x') = \sqrt{1-x'^2} \frac{1}{\pi} P.V. \int_{-1}^1 \frac{1}{\sqrt{1-\xi^2}} \frac{\chi_{x'}(\xi)}{x' - \xi} d\xi, \quad (32) \quad \text{f2}$$

and taking  $x'$  derivative

$$\tilde{\phi}_{y'}(x') = \partial_{x'} \left( \sqrt{1-x'^2} \frac{1}{\pi} P.V. \int_{-1}^1 \frac{1}{\sqrt{1-\xi^2}} \frac{\tilde{\phi}(\xi)}{x' - \xi} d\xi \right) \quad (33) \quad \text{f3}$$

which, by (13), yields the following evolution problem

$$\tilde{\phi}_{tt} = -|f'(x')| \partial_{x'} \left( \sqrt{1-x'^2} \frac{1}{\pi} P.V. \int_{-1}^1 \frac{\tilde{\phi}(t, \xi)}{\sqrt{1-\xi^2}(x' - \xi)} d\xi \right). \quad (34) \quad \text{integr}$$

The equation (34) may be rewritten as an identical equation for  $\tilde{\zeta}$  after taking an additional time derivative and writing  $\tilde{\phi}_t = -\tilde{\zeta}$ . We remark again that the formulation (34) is equivalent to the formulation given in [8] (see (A)).

We need to complement (34) with suitable initial and boundary conditions, namely

$$\begin{aligned} \tilde{\phi}(0, x') &= \tilde{\phi}_0(x'), \\ \tilde{\phi}_t(0, x') &= \tilde{\phi}_1(x'), \end{aligned}$$

where, by mass conservation (eq. (21))

$$\int_{-1}^1 \frac{\tilde{\phi}_1(x')}{|f'(x')|} dx' = \int_{-1}^1 \frac{\tilde{\zeta}(x')}{|f'(x')|} dx' = 0. \quad (35) \quad \text{condi}$$

Moreover, from (25), (26), we impose on the initial data:

$$\int_{-1}^1 \frac{\tilde{\phi}_{0,x'}(x')}{\sqrt{1-x'^2}} dx' = 0, \quad (36) \quad \text{condi}$$

$$\int_{-1}^1 \frac{x' \tilde{\phi}_{0,x'}(x')}{\sqrt{1-x'^2}} dx' = 0, \quad (37) \quad \text{condi}$$

which are automatically fulfilled by defining an initial normal derivative  $\tilde{\phi}_{0,y'}(x')$  with vanishing mean value in  $[-1, 1]$  and the corresponding tangential derivative  $\tilde{\phi}_{0,x'}(x')$  defined by (23).

In the case when  $|f'(x')|$  is a symmetric function (corresponding to a symmetric domain  $\Omega$ ) it is useful to think of  $\tilde{\phi}(t, x')$  as decomposed into symmetric and antisymmetric part; that is

$$\tilde{\phi}(t, x') = S(t, x') + N(t, x')$$

with

$$S(t, x) = \frac{\tilde{\phi}(t, x) + \tilde{\phi}(t, -x)}{2}, \quad N(t, x) = \frac{\tilde{\phi}(t, x) - \tilde{\phi}(t, -x)}{2},$$

and the initial data decomposed accordingly

$$\begin{aligned} S_0(x) &= \frac{\tilde{\phi}_0(x) + \tilde{\phi}_0(-x)}{2}, & N_0(x) &= \frac{\tilde{\phi}_0(x) - \tilde{\phi}_0(-x)}{2}, \\ S_1(x) &= \frac{\tilde{\phi}_1(x) + \tilde{\phi}_1(-x)}{2}, & N_1(x) &= \frac{\tilde{\phi}_1(x) - \tilde{\phi}_1(-x)}{2}. \end{aligned}$$

Then, one can consider the evolution problem for symmetric and antisymmetric functions separately. For symmetric functions, the initial data need to satisfy (35), (37) while for antisymmetric functions the conditions are (35), (36). Finally, we remark that (35), (36), (37) do not only hold initially, but for any time by replacing  $(\tilde{\phi}_0(x'), \tilde{\phi}_1(x'))$  by  $(\tilde{\phi}(t, x'), \tilde{\phi}_t(t, x'))$ .

We discuss now on boundary conditions for (34). Two kind of conditions are customary: pinned end and free end boundary conditions. In pinned end conditions one imposes  $\tilde{\zeta}(t, \pm 1) = 0$ , and since  $\tilde{\phi}_t = -\tilde{\zeta}$ , this implies

$$\tilde{\phi}(t, \pm 1) = 0 \quad (\text{pinned-end boundary condition}). \quad (38) \quad \text{pinne}$$

In the case of free-end boundary conditions one imposes  $\tilde{\zeta}_{x'}(t, \pm 1) = 0$  and since  $\tilde{\phi}_{x't} = \tilde{\zeta}_{x'}$  this translates into the condition

$$\tilde{\phi}_{x'}(t, \pm 1) = 0 \quad (\text{free-end boundary condition}). \quad (39) \quad \boxed{\text{free}}$$

We discuss now the case with fluid injection, i.e. with the condition (14) where the flux  $j$  is such that  $\int \frac{\tilde{j}(t, x')}{|f'(x')|} dx' = 0$  (in order to preserve the total fluid mass). This extra contribution to the normal derivative leads to

$$\begin{aligned} \tilde{\phi}_{y'} \Big|_{y'=0} &= \partial_{x'} \left( \sqrt{1-x'^2} \frac{1}{\pi} P.V. \int_{-1}^1 \frac{1}{\sqrt{1-\xi^2}} \frac{\tilde{\phi}(t, \xi, 0)}{x' - \xi} d\xi \right) \\ &\quad - \frac{1}{\sqrt{1-x'^2}} \frac{1}{\pi^2} P.V. \int_{-1}^1 \frac{\sqrt{1-\xi^2}}{(x' - \xi)} \int_{\mathbb{R} \setminus [-1, 1]} \frac{\tilde{j}(t, z)}{|f'(z)|(\xi - z)} dz d\xi, \end{aligned} \quad (40)$$

and hence to the evolution equation

$$\tilde{\phi}_{tt} = -|f'(x')| \partial_{x'} \left( \sqrt{1-x'^2} \frac{1}{\pi} P.V. \int_{-1}^1 \frac{\tilde{\phi}(t, \xi)}{\sqrt{1-\xi^2}(x' - \xi)} d\xi \right) + h(t, x'), \quad (41) \quad \boxed{\text{integ}}$$

where

$$h(t, x') = \frac{|f'(x')|}{\sqrt{1-x'^2}} \frac{1}{\pi^2} P.V. \int_{-1}^1 \frac{\sqrt{1-\xi^2}}{(x' - \xi)} \int_{\mathbb{R} \setminus [-1, 1]} \frac{\tilde{j}(t, z)}{|f'(z)|(\xi - z)} dz d\xi. \quad (42) \quad \boxed{\text{hxt}}$$

Notice that  $\chi_{y'} = \int_{-\infty}^{x'} \tilde{\phi}_{y'} dx'$  implies expression (42) can be obtained by replacing (31) with

$$\chi_{x'}(x') = -\frac{1}{\pi} P.V. \int_{-1}^1 \frac{\chi_{y'}(\xi)}{x' - \xi} d\xi - \frac{1}{\pi} P.V. \int_{-\infty}^{\infty} \frac{J(\xi)}{x' - \xi} d\xi$$

where  $J$  is the primitive of  $\tilde{j}/|f'|$ , inverting the Hilbert transform (restricted to  $[-1, 1]$ ) in the first term at the right hand side as in (32), (33) and using finally (79) in the Appendix.

#### 4. SPECTRUM OF THE SLOSHING PROBLEM

Following [\[15\]](#), to study the interior controllability problem (41), we need to consider the homogeneous (backward in time) adjoint version in  $(0, T)$  as that given in (34). Then, let us consider first the eigenvalue problem

$$\lambda \frac{\tilde{\phi}}{|f'(x')|} = \partial_{x'} \left( \sqrt{1-x'^2} P.V. \frac{1}{\pi} \int_{-1}^1 \frac{\tilde{\phi}(\xi)}{\sqrt{1-\xi^2}(x' - \xi)} d\xi \right) \quad (43) \quad \boxed{\text{eigen}}$$

with the additional condition that  $\tilde{\phi}(\pm 1) = 0$  (pinned contact line). Let  $T_r, U_r$ , with  $r \in \mathbb{N} \cup \{0\}$ , be the Tchebyshev polynomials of the first and second kind respectively. Then, for  $r \geq 1$ ,

$$\begin{aligned} \frac{1}{\pi} P.V. \int_{-1}^1 \frac{T_r(\xi)}{\sqrt{1-\xi^2}(x' - \xi)} d\xi &= -U_{r-1}(x'), \\ \frac{d}{dx'} \left( \sqrt{1-x'^2} U_{r-1}(x') \right) &= -r \frac{T_r(x')}{\sqrt{1-x'^2}}. \end{aligned}$$

By writing

$$\tilde{\phi}(\xi) = \sum_{n=0}^{\infty} a_n T_n(\xi), \quad (44) \quad \boxed{\text{a-1}}$$

and using the identities above, we find

$$\mathcal{A}\tilde{\phi} \equiv \partial_{x'} \left( \sqrt{1-x'^2} \frac{1}{\pi} P.V. \int_{-1}^1 \frac{\tilde{\phi}(\xi)}{\sqrt{1-\xi^2}(x' - \xi)} d\xi \right) = \sum_{n=1}^{\infty} n a_n \frac{T_n(x')}{\sqrt{1-x'^2}}.$$

We use now the fact that  $\int_{-1}^1 \tilde{\phi}(x') \frac{dx'}{|f'(x')|} = 0$  (see (22)), to deduce

$$a_0 \int_{-1}^1 T_0(x') \frac{dx'}{|f'(x')|} + \sum_{n=1}^{\infty} a_n \int_{-1}^1 T_n(x') \frac{dx'}{|f'(x')|} = 0.$$



So that

$$a_0 = -\frac{1}{\int_{-1}^1 T_0(x') \frac{dx'}{|f'(x')|}} \sum_{n=1}^{\infty} a_n \int_{-1}^1 T_n(x') \frac{dx'}{|f'(x')|},$$

and conclude the estimate

$$a_0^2 \leq \frac{\sum_{n=1}^{\infty} \left( \int_{-1}^1 T_n(x') \frac{dx'}{|f'(x')|} \right)^2}{\left( \int_{-1}^1 T_0(x') \frac{dx'}{|f'(x')|} \right)^2} \sum_{n=1}^{\infty} a_n^2.$$

Now, let  $c_n$  such that

$$\frac{\sqrt{1-x'^2}}{|f'(x')|} = \sum_{n=0}^{\infty} c_n T_n(x').$$

Then, by the orthogonality of the Tchebyshev polynomials,

$$c_0 = \frac{1}{\pi} \int_{-1}^1 \frac{T_0(x')}{|f'(x')|} dx', \quad c_n = \frac{2}{\pi} \int_{-1}^1 \frac{T_n(x')}{|f'(x')|} dx'.$$

Therefore

$$\int_{-1}^1 \frac{1-x'^2}{|f'(x')|^2} \frac{1}{\sqrt{1-x'^2}} dx' = \sum_{n,m=0}^{\infty} c_n c_m \int_{-1}^1 \frac{T_n T_m}{\sqrt{1-x'^2}} dx' = \pi c_0^2 + \frac{\pi}{2} \sum_{n=1}^{\infty} c_n^2,$$

namely

$$\begin{aligned} \frac{1}{\pi} \left( \int_{-1}^1 T_0(x') \frac{dx'}{|f'(x')|} \right)^2 + \frac{2}{\pi} \sum_{n=1}^{\infty} \left( \int_{-1}^1 T_n(x') \frac{dx'}{|f'(x')|} \right)^2 \\ = \int_{-1}^1 \sqrt{1-x'^2} \frac{dx'}{|f'(x')|^2} < \infty, \end{aligned}$$

which implies

$$a_0^2 \leq C \sum_{n=1}^{\infty} a_n^2. \quad (45) \quad \boxed{a_0}$$

We consider now the scalar product

$$(\mathcal{A}\tilde{\phi}, \tilde{\psi})_{L^2} = \int_{-1}^1 \tilde{\psi} \frac{\partial \tilde{\phi}}{\partial \tilde{n}} dx',$$

and find

$$(\mathcal{A}\tilde{\phi}, \tilde{\psi})_{L^2} = \int_{-1}^1 \left( \sum_{m=0}^{\infty} b_m T_m(x') \right) \left( \sum_{n=1}^{\infty} n a_n \frac{T_n(x')}{\sqrt{1-x'^2}} \right) dx' = \frac{\pi}{2} \sum_{n=1}^{\infty} n a_n b_n, \quad (46) \quad \boxed{a_1}$$

so that,  $(\tilde{\psi}, \mathcal{A}\tilde{\phi})_{L^2} = (\mathcal{A}\tilde{\psi}, \tilde{\phi})_{L^2}$ , implying the selfadjoint character of the operator  $\mathcal{A}$ , a fact already shown in a previous section.

We study the problem

$$\mathcal{A}\tilde{\phi} = u, \quad (47) \quad \boxed{\text{phiu}}$$

where  $u \in L_w^2$  with  $w = \sqrt{1-x'^2}$  and

$$L_w^2 \equiv \left\{ u : \int_{-1}^1 \sqrt{1-x'^2} |u|^2 dx' < \infty \right\}.$$

Since  $\left\{ T_n(x')/\sqrt{1-x'^2} \right\}_{n=0}^{\infty}$  form an orthogonal base for  $L_w^2$ , given  $u \in L_w^2$  we can expand

$$u = \sum_{n=0}^{\infty} u_n \frac{T_n(x')}{\sqrt{1-x'^2}},$$

and hence write the following weak version of (47)

$$(\mathcal{A}\tilde{\phi}, \tilde{\psi})_{L^2} = (u, \tilde{\psi})_{L^2}, \quad (48) \quad \boxed{\text{weak1}}$$

to be satisfied for any  $\tilde{\psi}$  in

$$H_{w^{-1}}^{\frac{1}{2}} \equiv \left\{ \tilde{\psi} : \|\tilde{\psi}\|_{H_{w^{-1}}^{\frac{1}{2}}} = \|\tilde{\psi}\|_{L_{w^{-1}}^2} + \left( \sum_{n=1}^{\infty} n \left( \int_{-1}^1 \frac{T_n(x')}{\sqrt{1-x'^2}} \tilde{\psi}(x') dx' \right)^2 \right)^{1/2} < \infty \right\} \quad (49) \quad \text{psi1}$$

such that, in addition,  $\tilde{\psi}(\pm 1) = 0$  and

$$\int_{-1}^1 \tilde{\psi}(x') \frac{dx'}{|f'(x')|} = 0. \quad (50) \quad \text{psi2}$$

We denote the space of functions vanishing at  $\pm 1$  and satisfying both (49), (50) as  $H_{w^{-1},0}^{\frac{1}{2}}$ . Expanding  $\tilde{\psi}$  in the form

$$\tilde{\psi} = \sum_{n=0}^{\infty} b_n T_n(x'), \quad (51) \quad \text{psi3}$$

we can write the equation (48) in the form

$$\sum_{n=1}^{\infty} n a_n b_n = 2u_0 b_0 + \sum_{n=1}^{\infty} u_n b_n, \quad (52) \quad \text{weak2}$$

making it clear that  $(\mathcal{A}\tilde{\phi}, \tilde{\psi})_{L^2}$  defines a continuous bilinear form on  $H_{w^{-1},0}^{\frac{1}{2}}$ . Indeed,

**Remark 1.** *By the orthogonality of the Tchebyshev polynomials, from (51)*

$$\int_{-1}^1 \frac{T_n(x')}{\sqrt{1-x'^2}} \tilde{\psi}(x') dx' = \begin{cases} 0, & n \neq m, \\ \pi b_n, & n = m = 0, \\ \frac{\pi}{2} b_n, & n = m \neq 0. \end{cases}$$

Then, for  $n \geq 1$ ,

$$\sqrt{n} \int_{-1}^1 \frac{T_n(x')}{\sqrt{1-x'^2}} \tilde{\psi}(x') dx' = \frac{\pi}{2} \sqrt{n} b_n,$$

or, equivalently,

$$\left( \sum_{n=1}^{\infty} n \left( \int_{-1}^1 \frac{T_n(x')}{\sqrt{1-x'^2}} \tilde{\psi}(x') dx' \right)^2 \right)^{1/2} = \frac{\pi}{2} \left( \sum_{n=1}^{\infty} n |b_n|^2 \right)^{1/2}.$$

Thus from (46) and the last equality, we have

$$\begin{aligned} |(\mathcal{A}\tilde{\phi}, \tilde{\psi})_{L^2}| &= \frac{\pi}{2} \left| \sum_{n \geq 1} n a_n b_n \right| \leq \frac{\pi}{2} \left( \sum_{n \geq 1} n |a_n|^2 \right)^{1/2} \left( \sum_{n \geq 1} n |b_n|^2 \right)^{1/2} \\ &\leq \frac{2}{\pi} \|\tilde{\phi}\|_{H_{w^{-1},0}^{1/2}} \|\tilde{\psi}\|_{H_{w^{-1},0}^{1/2}}. \end{aligned}$$

$(\mathcal{A}\tilde{\phi}, \tilde{\psi})_{L^2}$  is also coercive (the  $L_{w^{-1}}^2$  part of the norm of  $\tilde{\phi}$  is trivially bounded by  $\sum_{n=1}^{\infty} n a_n^2$ , except for  $a_0^2$  that is also bounded by (45)):

**Remark 2.** *By the orthogonality of the Tchebyshev polynomials, from (44)*

$$\|\tilde{\phi}\|_{L_{w^{-1}}^2}^2 = \int_{-1}^1 \frac{|\tilde{\phi}|^2}{\sqrt{1-x'^2}} dx' = \pi a_0^2 + \frac{\pi}{2} \sum_{n=1}^{\infty} a_n^2.$$

Moreover,

$$a_n = \begin{cases} \frac{1}{\pi} \int_{-1}^1 \frac{T_n(x')}{\sqrt{1-x'^2}} \tilde{\phi}(x') dx', & n = 0, \\ \frac{2}{\pi} \int_{-1}^1 \frac{T_n(x')}{\sqrt{1-x'^2}} \tilde{\phi}(x') dx', & n \geq 1. \end{cases}$$

Then, from the last remark and (45), coerciveness holds as follows:

$$\begin{aligned}
(\mathcal{A}\tilde{\phi}, \tilde{\phi})_{L^2} &= \frac{\pi}{2} \sum_{n \geq 1} n |a_n|^2 \geq C \left[ \pi \sum_{n=1}^{\infty} a_n^2 + \frac{\pi}{2} \sum_{n=1}^{\infty} a_n^2 + \frac{\pi^2}{4} \sum_{n=1}^{\infty} n a_n^2 \right] \\
&\geq C \left[ \pi a_0^2 + \frac{\pi}{2} \sum_{n=1}^{\infty} a_n^2 + \frac{\pi^2}{4} \sum_{n=1}^{\infty} n a_n^2 \right] \\
&= C \left[ \|\tilde{\phi}\|_{L^2_{w^{-1}}}^2 + \sum_{n=1}^{\infty} n \left( \int_{-1}^1 \frac{T_n}{\sqrt{1-x'^2}} \tilde{\phi} dx' \right)^2 \right] \\
&= C \|\tilde{\phi}\|_{H^{1/2}_{w^{-1},0}}^2,
\end{aligned} \tag{53}$$

for a constant  $C > 0$ .

Finally,  $u \in L^2_w$  defines a linear continuous functional on  $H^{1/2}_{w^{-1},0}$ :

$$\begin{aligned}
\left| \int_{-1}^1 u \tilde{\phi} dx' \right| &\leq \int_{-1}^1 \sqrt[4]{1-x'^2} |u| \frac{\tilde{\phi}}{\sqrt[4]{1-x'^2}} dx' \\
&\leq \|u\|_{L^2_w} \|\tilde{\phi}\|_{L^2_{w^{-1}}} \\
&\leq \|u\|_{L^2_w} \|\tilde{\phi}\|_{H^{1/2}_{w^{-1},0}}.
\end{aligned}$$

Hence, by Lax-Milgram's theorem, there exists a unique weak solution  $\tilde{\phi}$  to (47) that belongs to  $H^{1/2}_{w^{-1},0}$ . If we consider now  $u = \frac{v}{|f'(x')|}$  then the problem

$$\mathcal{A}\tilde{\phi} = \frac{v}{|f'(x')|}$$

where  $v$  is such that

$$\int_{-1}^1 \sqrt{1-x'^2} \frac{v^2(x')}{|f'(x')|^2} dx' < \infty \tag{54}$$

has also a unique solution  $\tilde{\phi} \in H^{1/2}_{w^{-1},0}$ . Note that (54) is satisfied if  $v \in L^2_{|f'(x')|^{-1}}$  provided there exists a constant  $C$  such that

$$|f'(x')| \geq C \sqrt{1-x'^2}, \quad x' \in [-1, 1], \tag{55}$$

because

$$\int_{-1}^1 \sqrt{1-x'^2} \frac{v^2(x')}{|f'(x')|^2} dx' \leq \frac{1}{C} \int_{-1}^1 \frac{v^2(x')}{|f'(x')|} dx' \leq \frac{1}{C^2} \int_{-1}^1 \frac{v^2(x')}{\sqrt{1-x'^2}} dx',$$

or

$$\|\varphi\|_{L^2_{|f'|^{-1}}}^2 \leq \frac{1}{C} \|\varphi\|_{L^2_{w^{-1}}}^2.$$

We define now the operator  $\mathcal{T}$  such that  $\tilde{\phi} = \mathcal{T}v$ . If we view  $\mathcal{T}$  as an operator from  $L^2_{|f'(x')|^{-1}}$  to  $L^2_{|f'(x')|^{-1}}$ , since  $H^{1/2}_{w^{-1},0} \subset L^2_{w^{-1}} \subset L^2_{|f'(x')|^{-1}}$ , we can see that  $\mathcal{T}$  is a selfadjoint operator (due to the selfadjoint character of  $\mathcal{A}$ ) and also a compact operator (due to the compact embedding  $H^{1/2}_{w^{-1},0} \subset L^2_{w^{-1}}$  (see (B)) and the continuous inclusion  $L^2_{w^{-1}} \subset L^2_{|f'(x')|^{-1}}$ ). Moreover,  $\ker(\mathcal{T}) = \{0\}$  and

$$(\mathcal{T}v, v)_{L^2_{|f'|^{-1}}} = (\tilde{\phi}, |f'| \mathcal{A}\tilde{\phi})_{L^2_{|f'|^{-1}}} = (\tilde{\phi}, \mathcal{A}\tilde{\phi})_{L^2} \geq 0,$$

for any  $v \in L^2_{|f'|^{-1}}$ . Therefore, by the spectral decomposition theorem,  $L^2_{|f'|^{-1}}$  admits a Hilbert basis  $\{e_n\}$  formed by eigenvectors of  $\mathcal{T}$ , with eigenvalues  $\mu_n$  such that  $\mu_n > 0$  and  $\mu_n \rightarrow 0$  as  $n \rightarrow \infty$ . We have then  $\mathcal{T}e_n = \mu_n e_n$ ,  $e_n \in H^{1/2}_{w^{-1},0}$ , and thus

$$(\mathcal{A}e_n, \tilde{\psi})_{L^2} = \frac{1}{\mu_n} (\mathcal{A}(\mathcal{T}e_n), \tilde{\psi})_{L^2} = \frac{1}{\mu_n} \left( \frac{e_n}{|f'|}, \tilde{\psi} \right)_{L^2}, \quad \text{for any } \tilde{\psi} \in H^{1/2}_{w^{-1},0}. \tag{56}$$

This implies  $e_n$  is a weak solution to the eigenvalue problem (43) with  $\lambda = \lambda_n = \mu_n^{-1}$ . The base can be made orthonormal, i.e.

$$\int_{-1}^1 \frac{e_i e_j}{|f'(x')|} dx' = \delta_{ij}.$$

Let us remark that condition (55) is satisfied provided the angles  $\alpha_{1,2}$  between the fluid interface and the solid container satisfy

$$\alpha_{1,2} \geq \frac{\pi}{2},$$

since  $|f'(x')|$  has a singularity weaker (i.e. with larger exponent) than  $|x \pm 1|^{1/2}$  otherwise [2]. asmr2002applied

We summarize the results above in the following theorem:

**Theorem 1.** *Let  $\Omega$  be a domain such that the interior angles between the free liquid interface and the solid are greater or equal than  $\pi/2$ . There exist a Hilbert basis  $\{e_n\}_{n \geq 1}$  of  $L^2_{|f'|^{-1}}$  and a sequence  $\{\lambda_n\}_{n \geq 1}$  of real numbers with  $\lambda_n > 0 \forall n$  and  $\lambda_n \rightarrow +\infty$  such that*

$$e_n \in H^{\frac{1}{2}}_{w^{-1},0},$$

$$\mathcal{A}e_n = \lambda_n \frac{e_n}{|f'(x')|}.$$

As a final remark, notice that for bounded symmetric  $|f'(x')|$ , one has  $L^2_{|f'(x')|^{-1}} \subset L^2$  and hence vectors in  $L^2_{|f'|^{-1}}$  may be expanded in trigonometric basis of  $L^2$ . Using the base  $\{\sin(n\pi x')\}_{n=1}^{\infty}$  one can approximate antisymmetric eigenfunctions (solutions to (43)) with pinned contact lines while the set  $\{\cos((n + \frac{1}{2})\pi x')\}_{n=0}^{\infty}$  (with the extra mass conservation condition also imposed) allows to approximate symmetric eigenfunctions. Likewise, the sets  $\{\sin((n + \frac{1}{2})\pi x')\}_{n=0}^{\infty}$  and  $\{\cos(n\pi x')\}_{n=1}^{\infty}$  allow to find antisymmetric and symmetric eigenfunctions with free-end condition. This approach was followed in [8], in order to compute eigenvalues and eigenfunctions numerically for various domains. fontelos2020gravity

## 5. OBSERVABILITY AND ENERGY ESTIMATES

S5

We will consider next the control problem consisting of finding  $h(t, x')$  in equation (41) such that for given initial data the solution vanishes at some time  $T$ . As it is customary in controllability theory, we need to consider first the homogeneous adjoint problem and obtain observability estimates.

The analysis presented above allows us to consider the solution in  $(0, T)$  of the following homogeneous adjoint problem (see (34))

$$\frac{\tilde{\phi}_{tt}}{|f'|} + \mathcal{A}\tilde{\phi} = 0, \tag{57} \quad \text{o1}$$

with suitable initial conditions  $\tilde{\phi}(0) = \tilde{\phi}_0$ ,  $\tilde{\phi}'(0) = \tilde{\phi}_1$ , whose solution we can write as

$$\tilde{\phi} = \sum_{n=1}^{\infty} (A_n \cos(\theta_n t) + B_n \sin(\theta_n t)) e_n, \tag{58} \quad \text{o2}$$

where

$$\tilde{\phi}_0 = \sum_{n=1}^{\infty} A_n e_n, \quad \tilde{\phi}_1 = \sum_{n=1}^{\infty} \theta_n B_n e_n,$$

with

$$A_n = \int_{-1}^1 \frac{\tilde{\phi}_0 e_n}{|f'(x')|} dx' = (\tilde{\phi}_0, e_n)_{|f'|^{-1}}, \quad B_n = \frac{1}{\theta_n} \int_{-1}^1 \frac{\tilde{\phi}_1 e_n}{|f'(x')|} dx' = \frac{1}{\theta_n} (\tilde{\phi}_1, e_n)_{|f'|^{-1}}.$$

Hence  $\|\tilde{\phi}\|_{L^2(L^2_{|f'|^{-1}})}^2 = \int_0^T \int_{-1}^1 \frac{\tilde{\phi}^2}{|f'(x')|} dx' dt$  can be bounded from below by

$$\begin{aligned}
& \int_0^T \sum_n (A_n^2 \cos^2(\theta_n t) + B_n^2 \sin^2(\theta_n t) + 2A_n B_n \sin(\theta_n t) \cos(\theta_n t)) dt \\
&= \sum_n \left[ \frac{A_n^2}{2} \left( T + \frac{\sin(2\theta_n T)}{2\theta_n} \right) + \frac{B_n^2}{2} \left( T - \frac{\sin(2\theta_n T)}{2\theta_n} \right) + A_n B_n \frac{1 - \cos(2\theta_n T)}{2\theta_n} \right] \\
&\geq \sum_n \left[ \frac{A_n^2}{2} \left( T + \frac{\sin(2\theta_n T)}{2\theta_n} \right) + \frac{B_n^2}{2} \left( T - \frac{\sin(2\theta_n T)}{2\theta_n} \right) - \frac{A_n^2 + B_n^2}{2} \frac{1 - \cos(2\theta_n T)}{2\theta_n} \right] \\
&= \frac{T}{2} \sum_n \left[ A_n^2 \left( 1 - \frac{1}{2\theta_n T} + \frac{\sin(2\theta_n T)}{2\theta_n T} + \frac{\cos(2\theta_n T)}{2\theta_n T} \right) \right. \\
&\quad \left. + B_n^2 \left( 1 - \frac{1}{2\theta_n T} - \frac{\sin(2\theta_n T)}{2\theta_n T} + \frac{\cos(2\theta_n T)}{2\theta_n T} \right) \right] \\
&\geq C \frac{T}{2} \sum_n [A_n^2 + B_n^2],
\end{aligned} \tag{59} \quad \square_6$$

for some positive  $C > 0$  and  $T > 2.42/2 \min(\theta_n) = 2.42/2\theta_1$ .

From the orthogonality of the Hilbert basis  $\{e_n\}$ ,

$$\|\tilde{\phi}_0\|_{L^2_{|f'|^{-1}}}^2 = \int_{-1}^1 \frac{\tilde{\phi}_0^2}{|f'(x')|} dx' = \sum_{n=1}^{\infty} A_n^2. \tag{60} \quad \square_7$$

Moreover, from (56), the decomposition given in (58), and (57) we conclude  $\lambda_n = \theta_n^2$  and thus

$$\sum_{n=1}^{\infty} B_n^2 = \sum_n \left( \frac{1}{\theta_n} \int_{-1}^1 \frac{\tilde{\phi}_1 e_n}{|f'(x')|} dx' \right)^2 = \sum_n \frac{1}{\lambda_n} \left( \int_{-1}^1 \frac{\tilde{\phi}_1 e_n}{|f'(x')|} dx' \right)^2. \tag{61} \quad \square_8$$

Let us define, accordingly, the space

$$\tilde{H}^{-\frac{1}{2}} = \left\{ \tilde{\phi} : \sum_n \frac{1}{\lambda_n} \left( \int_{-1}^1 \frac{\tilde{\phi} e_n}{|f'(x')|} dx' \right)^2 < \infty \right\},$$

and its dual space

$$\tilde{H}^{\frac{1}{2}} = \left\{ \tilde{\phi} : \sum_n \lambda_n \left( \int_{-1}^1 \frac{\tilde{\phi} e_n}{|f'(x')|} dx' \right)^2 < \infty \right\}.$$

Note that, by (46) and definition of  $\lambda_n$  and  $e_n$ ,

$$\|\tilde{\phi}\|_{\tilde{H}^{\frac{1}{2}}}^2 = \sum \left( \int \frac{\tilde{\phi} \lambda_n e_n}{|f'(x)|} \right) \left( \int \frac{\tilde{\phi} e_n}{|f'(x)|} \right) = \int \tilde{\phi} \mathcal{A} \tilde{\phi} = \frac{\pi}{2} \sum_{n \geq 1} n a_n^2 \sim \|\tilde{\phi}\|_{H^{\frac{1}{2}}_{w^{-1}}}^2.$$

From (59)–(61), we conclude then for some  $C > 0$  and  $T$  sufficiently large,

$$\begin{aligned}
C^{-1} T \left( \|\tilde{\phi}_1\|_{\tilde{H}^{-\frac{1}{2}}}^2 + \|\tilde{\phi}_0\|_{L^2_{|f'|^{-1}}}^2 \right) &\geq \\
&\int_0^T \int_{-1}^1 \frac{\tilde{\phi}^2}{|f'(x')|} dx' dt \geq CT \left( \|\tilde{\phi}_1\|_{\tilde{H}^{-\frac{1}{2}}}^2 + \|\tilde{\phi}_0\|_{L^2_{|f'|^{-1}}}^2 \right).
\end{aligned}$$

This proves the following theorem:

**Theorem 2.** *The equation (57) is observable in time  $T > 0$ . That is, there exists a positive constant  $C > 0$  such that for  $T$  sufficiently large*

$$\int_0^T \int_{-1}^1 \frac{\tilde{\phi}^2}{|f'(x')|} dx' dt \geq CT \left( \|\tilde{\phi}_1\|_{\tilde{H}^{-\frac{1}{2}}}^2 + \|\tilde{\phi}_0\|_{L^2_{|f'|^{-1}}}^2 \right).$$

We consider next the nonhomogeneous problem (as in (41))

$$\frac{\tilde{\psi}_{tt}}{|f'|} + \mathcal{A}\tilde{\psi} = \frac{\tilde{h}}{|f'|}. \quad (62) \quad \boxed{\text{o12}}$$

It will be convenient the following energy estimate for the nonhomogeneous problem:

$$\frac{1}{2} \frac{d}{dt} \int_{-1}^1 \frac{\tilde{\psi}_t^2}{|f'(x')|} dx' + \int_{-1}^1 \tilde{\psi}_t \mathcal{A}\tilde{\psi} dx' = \int_{-1}^1 \frac{\tilde{h}\tilde{\psi}_t}{|f'(x')|} dx'. \quad (63) \quad \boxed{\text{o13}}$$

If we let

$$\tilde{\psi} = \sum_{n=1}^{\infty} c_n e_n,$$

then

$$c_n = \int_{-1}^1 \frac{\tilde{\psi} e_n}{|f'(x')|} dx',$$

$$\tilde{\psi}_t = \sum_{n=1}^{\infty} c_{n,t} e_n \implies \int_{-1}^1 \frac{\tilde{\psi}_t^2}{|f'(x')|} dx' = \sum_{n=1}^{\infty} c_{n,t}^2.$$

Noticing that

$$\begin{aligned} \int_{-1}^1 \tilde{\psi}_t \mathcal{A}\tilde{\psi} dx' &= \int_{-1}^1 \left( \sum_n c_{n,t} e_n \right) \left( \sum_m \lambda_m c_m \frac{e_m}{|f'|} \right) dx' \\ &= \sum_n \lambda_n c_{n,t} c_n \\ &= \frac{1}{2} \frac{d}{dt} \sum_n \lambda_n c_n^2, \end{aligned}$$

(63) is equivalent to the following inequality

$$\frac{dE}{dt} \leq \|\tilde{h}\|_{L^2_{|f'|^{-1}}} E^{\frac{1}{2}},$$

for the energy defined as

$$E = \frac{1}{2} \sum_n c_{n,t}^2 + \frac{1}{2} \sum_n \lambda_n c_n^2.$$

Hence, the natural initial data for which the problem is well-posed is  $(\tilde{\psi}_0, \tilde{\psi}_1) \in \tilde{H}^{\frac{1}{2}} \times L^2_{|f'|^{-1}}$  and, the energy is bounded provided  $\int_0^T \|\tilde{h}\|_{L^2_{|f'|^{-1}}}^2 dt < \infty$ , i.e.  $\tilde{h} \in L^2(0, T; L^2_{|f'|^{-1}})$ . Namely, we have the following:

**Theorem 3.** *For any  $\tilde{h} \in L^2(0, T; L^2_{|f'|^{-1}})$  and  $(\tilde{\psi}_0, \tilde{\psi}_1) \in \tilde{H}^{\frac{1}{2}} \times L^2_{|f'|^{-1}}$  equation (62) has a unique weak solution*

$$(\tilde{\psi}, \tilde{\psi}') \in C([0, T]; H^{\frac{1}{2}}_{w^{-1}, 0} \times L^2_{|f'|^{-1}}).$$

## 6. CONTROLLABILITY

Once we have studied in the previous sections the forward evolution equation and the backward homogeneous system, given by (62) and (57) respectively, we are in position of comparing them to get the controllability condition on the system.

That is, if we assume the control drives the initial data of system (62) to zero, by multiplying the source term in (62) by the solution to the adjoint problem (57), we obtain

$$\begin{aligned}
\int_0^T \int_{-1}^1 \frac{\tilde{h}\tilde{\phi}}{|f'|} dx' dt &= \int_0^T \int_{-1}^1 \left[ \frac{\tilde{\psi}_{tt}}{|f'|} + \mathcal{A}\tilde{\psi} \right] \tilde{\phi} dx' dt \\
&= \int_{-1}^1 \frac{\tilde{\psi}_t \tilde{\phi}|_0^T}{|f'|} dx' - \int_0^T \int_{-1}^1 \frac{\tilde{\psi}_t \tilde{\phi}_t}{|f'|} dx' dt + \int_0^T \int_{-1}^1 \tilde{\psi} \mathcal{A}\tilde{\phi} dx' dt \\
&= \int_{-1}^1 \frac{\tilde{\psi}_t \tilde{\phi}|_0^T}{|f'|} dx' - \int_{-1}^1 \frac{\tilde{\psi} \tilde{\phi}_t|_0^T}{|f'|} dx' + \int_0^T \int_{-1}^1 \tilde{\psi} \left[ \frac{\tilde{\phi}_{tt}}{|f'|} + \mathcal{A}\tilde{\phi} \right] dx' dt \\
&= - \int_{-1}^1 \frac{\tilde{\psi}_1 \tilde{\phi}_0}{|f'|} dx' + \int_{-1}^1 \frac{\tilde{\psi}_0 \tilde{\phi}_1}{|f'|} dx',
\end{aligned}$$

where  $(\tilde{\psi}_0, \tilde{\psi}_1) = (\tilde{\psi}(0), \tilde{\psi}_t(0)) \in \tilde{H}^{\frac{1}{2}} \times L^2_{|f'|^{-1}}$ . So that  $\tilde{h}$  can be chosen as the minimizer of the functional

$$J[\tilde{\phi}_0, \tilde{\phi}_1] = \frac{1}{2} \int_0^T \int_{-1}^1 \frac{\tilde{\phi}^2}{|f'|} dx' dt + \int_{-1}^1 \frac{\tilde{\psi}_1 \tilde{\phi}_0}{|f'|} dx' - \int_{-1}^1 \frac{\tilde{\psi}_0 \tilde{\phi}_1}{|f'|} dx'.$$

Notice that

$$\left| \int_{-1}^1 \frac{\tilde{\psi}_1 \tilde{\phi}_0}{|f'|} dx' - \int_{-1}^1 \frac{\tilde{\psi}_0 \tilde{\phi}_1}{|f'|} dx' \right| \leq \|\tilde{\phi}_0\|_{L^2_{|f'|^{-1}}} \|\tilde{\psi}_1\|_{L^2_{|f'|^{-1}}} + \|\tilde{\phi}_1\|_{\tilde{H}^{-\frac{1}{2}}} \|\tilde{\psi}_0\|_{\tilde{H}^{\frac{1}{2}}}.$$

As it is customary in controllability theory, coercivity of the functional  $J$  is guaranteed if the adjoint problem is observable in time  $T$ , that is:

$$\int_0^T dt \int_{-1}^1 \frac{\tilde{\phi}^2}{|f'|} dx' \geq C \left( \|\tilde{\phi}(0)\|_{L^2_{|f'|^{-1}}}^2 + \|\tilde{\phi}_t(0)\|_{\tilde{H}^{-\frac{1}{2}}}^2 \right), \quad (64) \quad \text{obs}$$

a fact that was proved in the previous section. Hence, we have proved the following controllability theorem:

**Theorem 4.** *The system (62) is exactly controllable in time  $T$ . That is, for any initial data  $(\tilde{\psi}_0, \tilde{\psi}_1) \in \tilde{H}^{\frac{1}{2}} \times L^2_{|f'|^{-1}}$ , there exist  $\tilde{h} \in L^2(0, T; L^2_{|f'|^{-1}})$  and  $T > 0$  such that*

$$\|\tilde{\psi}\|_{\tilde{H}^{\frac{1}{2}}_{w^{-1}, 0}} = 0, \quad \text{for } t > T.$$

The fact that the control  $\tilde{h} \in L^2((0, T); L^2_{|f'|^{-1}})$  implies that one can write

$$\tilde{h}(t, x') = \sum_n h_n(t) e_n,$$

with

$$\int_0^T \sum_n |h_n(t)|^2 dt < \infty.$$

We are going to discuss next on how to approach the control function  $\tilde{h}$ , defined at the fluid interface by means of a function  $j$  defined at the solid boundaries. This function  $j$  will represent a fluid injection at certain points  $\{x'_1, x'_2, \dots, x'_N\}$  of the solid boundary with a flow rate  $\{j_1(t), j_2(t), \dots, j_N(t)\}$  and under the mass conservation condition

$$\sum_{i=1}^N j_i(t) = 0.$$

More precisely,

$$\frac{\partial \tilde{\phi}}{\partial \tilde{n}} = \sum_i j_i(t) \delta(x' - x'_i). \quad (65) \quad \text{flux}$$

We replace the expression for  $\tilde{j}(t, z, 0)$  in (42) by the right hand side of (65) at the solid side walls and obtain

$$\begin{aligned}
& \frac{|f'(x')|}{\sqrt{1-x'^2}} \frac{1}{\pi^2} P.V. \int_{-1}^1 \frac{\sqrt{1-\xi^2}}{(x'-\xi)} \int_{\mathbb{R} \setminus [-1,1]} \frac{\tilde{j}(t, z, 0)}{|f'(z)|(\xi-z)} dz d\xi \\
&= \sum_i \frac{j_i(t)}{|f'(x'_i)|} \frac{|f'(x')|}{\sqrt{1-x'^2}} \frac{1}{\pi^2} P.V. \int_{-1}^1 \frac{\sqrt{1-\xi^2}}{(x'-\xi)(\xi-x'_i)} d\xi \\
&= \sum_i \frac{j_i(t)}{|f'(x'_i)|} \frac{|f'(x')|}{\sqrt{1-x'^2}} \frac{1}{x'-x'_i} \frac{1}{\pi^2} P.V. \int_{-1}^1 \sqrt{1-\xi^2} \left[ \frac{1}{x'-\xi} + \frac{1}{\xi-x'_i} \right] d\xi \\
&= \frac{|f'(x')|}{\sqrt{1-x'^2}} \sum_i j_i(t) \frac{G(x'_i) - G(x)}{\pi^2 |f'(x'_i)|} \frac{1}{x'-x'_i}, \tag{66}
\end{aligned}$$

with

$$G(x'_i) = P.V. \int_{-1}^1 \frac{\sqrt{1-\xi^2}}{(\xi-x'_i)} d\xi.$$

Hence, the challenge is to approximate the control function  $\tilde{h}$  by (66). One possibility is to approximate the first  $N-1$  modes so that

$$\int_{-1}^1 \frac{\tilde{h} e_i}{|f'(x')|} dx' = h_i(t), \quad i = 1, \dots, N-1,$$

or, denoting  $m_{ij} = \frac{1}{\pi^2 |f'(x'_j)|} \int_{-1}^1 \frac{G(x'_j) - G(x)}{\sqrt{1-x'^2}(x'-x'_j)} e_i dx'$  and including the mass conservation condition, solving the system

$$\begin{aligned}
& \sum_{j=1}^N m_{ij} j_j(t) = h_i(t), \quad i = 1, \dots, N-1, \\
& \sum_{i=1}^N j_i(t) = 0.
\end{aligned}$$

The fact that  $h_i(t) \in L^2(0, T)$  implies  $j_i(t) \in L^2(0, T)$ .

#### APPENDIX A. EQUIVALENT EXPRESSIONS OF THE OPERATOR $\mathcal{A}$

In the case when  $\tilde{\phi}$  is harmonic in the lower half plane,  $\tilde{\phi}$  satisfies (pinned contact line)

$$|\tilde{\phi}|/|x' \pm 1|^{\frac{1}{2}} \rightarrow 0 \quad \text{as } x' \rightarrow \mp 1$$



and  $\int_{-1}^1 \frac{\partial \tilde{\phi}}{\partial n} dx' = 0$  (see (20)). Then we have

$$\begin{aligned}
\pi \mathcal{A} \tilde{\phi} &\equiv \partial_{x'} \left( \sqrt{1-x'^2} P.V. \int_{-1}^1 \frac{1}{\sqrt{1-\xi^2}} \frac{1}{x'-\xi} \tilde{\phi}(\xi) d\xi \right) \\
&= -\frac{x'}{\sqrt{1-x'^2}} \int_{-1}^1 \frac{1}{\sqrt{1-\xi^2}} \frac{1}{x'-\xi} \tilde{\phi} - \sqrt{1-x'^2} \int_{-1}^1 \frac{1}{\sqrt{1-\xi^2}} \frac{1}{x'^2} \tilde{\phi} \\
&= -\frac{x'}{\sqrt{1-x'^2}} \int_{-1}^1 \frac{1}{\sqrt{1-\xi^2}} \frac{1}{x'-\xi} \tilde{\phi} - \sqrt{1-x'^2} \int_{-1}^1 \partial_\xi \left( \frac{1}{x'-\xi} \frac{1}{\sqrt{1-\xi^2}} \right) \tilde{\phi} \\
&\quad + \sqrt{1-x'^2} \int_{-1}^1 \frac{1}{x'-\xi} \partial_\xi \left( \frac{1}{\sqrt{1-\xi^2}} \right) \tilde{\phi} \\
&= -\frac{x'}{\sqrt{1-x'^2}} \int_{-1}^1 \frac{1}{\sqrt{1-\xi^2}} \frac{1}{x'-\xi} \tilde{\phi} + \sqrt{1-x'^2} \int_{-1}^1 \frac{1}{x'-\xi} \frac{1}{\sqrt{1-\xi^2}} \tilde{\phi}_\xi \\
&\quad + \sqrt{1-x'^2} \int_{-1}^1 \frac{1}{x'-\xi} \frac{\xi}{(1-\xi^2)^{\frac{3}{2}}} \tilde{\phi} \\
&= -\frac{1}{\sqrt{1-x'^2}} \int_{-1}^1 \frac{1}{(1-\xi^2)^{\frac{3}{2}}} \frac{x'(1-\xi^2) - \xi(1-x'^2)}{x'-\xi} \tilde{\phi} \\
&\quad + \sqrt{1-x'^2} \int_{-1}^1 \frac{1}{x'-\xi} \frac{1}{\sqrt{1-\xi^2}} \tilde{\phi}_\xi.
\end{aligned}$$

We now turn our attention to

$$\begin{aligned}
&-\frac{1}{\sqrt{1-x'^2}} \int_{-1}^1 \frac{1}{(1-\xi^2)^{\frac{3}{2}}} \frac{x'(1-\xi^2) - \xi(1-x'^2)}{x'-\xi} \tilde{\phi} \\
&= -\frac{1}{\sqrt{1-x'^2}} \int_{-1}^1 \frac{1}{(1-\xi^2)^{\frac{3}{2}}} \tilde{\phi} - \frac{x'}{\sqrt{1-x'^2}} \int_{-1}^1 \frac{\xi}{(1-\xi^2)^{\frac{3}{2}}} \tilde{\phi} \\
&= \frac{1}{\sqrt{1-x'^2}} \int_{-1}^1 \frac{\xi}{\sqrt{1-\xi^2}} \tilde{\phi}_\xi + \frac{x'}{\sqrt{1-x'^2}} \int_{-1}^1 \frac{1}{\sqrt{1-\xi^2}} \tilde{\phi}_\xi \\
&= 0,
\end{aligned}$$

where we have integrated by parts and used (36), (37), together with

$$\frac{d}{d\xi} \left( \frac{\xi}{\sqrt{1-\xi^2}} \right) = \frac{1}{(1-\xi^2)^{\frac{3}{2}}}, \quad \frac{d}{d\xi} \left( \frac{1}{\sqrt{1-\xi^2}} \right) = \frac{\xi}{(1-\xi^2)^{\frac{3}{2}}}.$$

Hence,

$$\mathcal{A} \tilde{\phi} = \sqrt{1-x'^2} \frac{1}{\pi} P.V. \int_{-1}^1 \frac{1}{x'-\xi} \frac{1}{\sqrt{1-\xi^2}} \tilde{\phi}_\xi(\xi) d\xi, \tag{67} \quad \boxed{\text{afi}}$$

for all  $\tilde{\phi}$  satisfying (36), (37). We can also show formula (67) by a second method that also extends to free-end boundary conditions. Let

$$\tilde{\phi} = \sum_{n=0}^{\infty} a_n T_n, \tag{68} \quad \boxed{\text{expan}}$$

and observe the following well-known identities

$$\frac{dT_n}{dx} = nU_{n-1}, \quad (69) \quad \boxed{\text{i1}}$$

$$-U_{n-1}(x) = \frac{1}{\pi} P.V. \int_{-1}^1 \frac{T_n(\xi)}{\sqrt{1-\xi^2}(x-\xi)} d\xi, \quad (70) \quad \boxed{\text{i2}}$$

$$T_{n+1} = xT_n - (1-x^2)U_{n-1}, \quad (71) \quad \boxed{\text{i3}}$$

$$T_n = U_n - xU_{n-1}, \quad (72) \quad \boxed{\text{i4}}$$

$$U_n = 2 \sum_{\text{odd } j}^n T_j, \quad n \text{ odd}, \quad (73) \quad \boxed{\text{i5}}$$

$$U_n = 2 \sum_{\text{even } j}^n T_j - 1, \quad n \text{ even}. \quad (74) \quad \boxed{\text{i6}}$$

Then, for each  $T_n$ , with  $n > 0$ , and using (69), (70), (73), (74), we have

$$\begin{aligned} & \sqrt{1-x'^2} \frac{1}{\pi} P.V. \int_{-1}^1 \frac{1}{x'-\xi} \frac{1}{\sqrt{1-\xi^2}} T'_n(\xi) d\xi \\ &= \sqrt{1-x'^2} \frac{n}{\pi} P.V. \int_{-1}^1 \frac{1}{x'-\xi} \frac{1}{\sqrt{1-\xi^2}} U_{n-1}(\xi) d\xi \\ &= \sqrt{1-x'^2} \frac{n}{\pi} P.V. \int_{-1}^1 \frac{1}{x'-\xi} \frac{1}{\sqrt{1-\xi^2}} \left\{ \begin{array}{l} 2 \sum_{\text{odd } j}^{n-1} T_j, \quad n \text{ even} \\ 2 \sum_{\text{even } j}^{n-1} T_j - 1, \quad n \text{ odd} \end{array} \right\} d\xi \\ &= -\sqrt{1-x'^2} n \left\{ \begin{array}{l} 2 \sum_{\text{odd } j}^{n-1} U_{j-1}, \quad n \text{ even} \\ 2 \sum_{\text{even } j}^{n-1} U_{j-1}, \quad n \text{ odd} \end{array} \right\}. \end{aligned}$$

Using (71) and adding in  $n$ :

$$\begin{aligned} \sum_{\text{odd } j}^{n-1} T_{j+1} &= x \sum_{\text{odd } j}^{n-1} T_j - (1-x^2) \sum_{\text{odd } j}^{n-1} U_{j-1}, \quad n \text{ even}, \\ \sum_{\text{even } j}^{n-1} T_{j+1} &= x \sum_{\text{even } j}^{n-1} T_j - (1-x^2) \sum_{\text{even } j}^{n-1} U_{j-1}, \quad n \text{ odd}, \end{aligned}$$

and using (73), (74) we get

$$\begin{aligned} \frac{U_n}{2} - 1 &= x' \frac{U_{n-1}}{2} - (1-x'^2) \sum_{\text{odd } j}^{n-1} U_{j-1}, \quad n \text{ even}, \\ \frac{U_n}{2} &= x' \frac{U_{n-1} + 1}{2} - (1-x'^2) \sum_{\text{even } j}^{n-1} U_{j-1}, \quad n \text{ odd}. \end{aligned}$$

Finally, using (72) we get

$$\begin{aligned} \frac{T_n}{2} - 1 &= -(1-x'^2) \sum_{\text{odd } j}^{n-1} U_{j-1}, \quad n \text{ even}, \\ \frac{T_n}{2} &= \frac{x'}{2} - (1-x'^2) \sum_{\text{even } j}^{n-1} U_{j-1}, \quad n \text{ odd}, \end{aligned}$$

and hence,

$$\sqrt{1-x'^2} \frac{1}{\pi} P.V. \int_{-1}^1 \frac{1}{x'-\xi} \frac{1}{\sqrt{1-\xi^2}} T'_n(\xi) d\xi = n \left\{ \begin{array}{l} \frac{T_n}{\sqrt{1-x'^2}} - \frac{2}{\sqrt{1-x'^2}}, \quad n \text{ even} \\ \frac{T_n}{\sqrt{1-x'^2}} - \frac{x'}{\sqrt{1-x'^2}}, \quad n \text{ odd} \end{array} \right\}. \quad (75) \quad \boxed{\text{i7}}$$

Then, from the well known formula

$$\frac{d}{dx} \left( \sqrt{1-x^2} U_{n-1}(x) \right) = -n \frac{T_n(x)}{\sqrt{1-x^2}},$$

and using (70) we can rewrite (75) as

$$\begin{aligned} & \sqrt{1-x'^2} \frac{1}{\pi} P.V. \int_{-1}^1 \frac{1}{x'-\xi} \frac{1}{\sqrt{1-\xi^2}} T'_n(\xi) d\xi \\ &= \frac{d}{dx'} \left( \sqrt{1-x'^2} \frac{1}{\pi} P.V. \int_{-1}^1 \frac{T_n(\xi)}{\sqrt{1-\xi^2}(x'-\xi)} d\xi \right) - n \left\{ \begin{array}{l} \frac{2}{\sqrt{1-x'^2}}, \quad n \text{ even} \\ \frac{x'}{\sqrt{1-x'^2}}, \quad n \text{ odd} \end{array} \right\}. \end{aligned} \quad (76) \quad \text{i8}$$

Using the identities (73), (74), the fact that  $T_0(\xi) = 1$ ,  $T_1(\xi) = \xi$ , and orthogonality of polynomials  $T_n$ , we find

$$\begin{aligned} n \int_{-1}^1 \frac{\xi}{\sqrt{1-\xi^2}} U_{n-1}(\xi) d\xi &= 2n, \quad n \text{ even}, \\ n \int_{-1}^1 \frac{1}{\sqrt{1-\xi^2}} U_{n-1}(\xi) d\xi &= n, \quad n \text{ odd}. \end{aligned}$$

Thus, from this two identities and (69), we can rewrite the last term at the right hand side of (76) as

$$n \left\{ \begin{array}{l} \frac{2}{\sqrt{1-x'^2}}, \quad n \text{ even} \\ \frac{x'}{\sqrt{1-x'^2}}, \quad n \text{ odd} \end{array} \right\} = \left\{ \begin{array}{l} \frac{1}{\sqrt{1-x'^2}} \int_{-1}^1 \frac{\xi}{\sqrt{1-\xi^2}} \frac{dT_n(\xi)}{d\xi} d\xi, \quad n \text{ even} \\ \frac{x'}{\sqrt{1-x'^2}} \int_{-1}^1 \frac{1}{\sqrt{1-\xi^2}} \frac{dT_n(\xi)}{d\xi} d\xi, \quad n \text{ odd} \end{array} \right\}. \quad (77) \quad \text{i9}$$

Finally, given  $\tilde{\phi}$  by (68) and using (76), (77), we conclude

$$\begin{aligned} & \sqrt{1-x'^2} \frac{1}{\pi} P.V. \int_{-1}^1 \frac{1}{x'-\xi} \frac{1}{\sqrt{1-\xi^2}} \tilde{\phi}_\xi(\xi) d\xi \\ &= \partial_{x'} \left( \sqrt{1-x'^2} \frac{1}{\pi} P.V. \int_{-1}^1 \frac{1}{\sqrt{1-\xi^2}} \frac{1}{x'-\xi} \tilde{\phi}(\xi) d\xi \right) \\ &\quad - \frac{1}{\sqrt{1-x'^2}} \frac{1}{\pi} \int_{-1}^1 \frac{\xi}{\sqrt{1-\xi^2}} \tilde{\phi}_\xi(\xi) d\xi - \frac{x'}{\sqrt{1-x'^2}} \frac{1}{\pi} \int_{-1}^1 \frac{1}{\sqrt{1-\xi^2}} \tilde{\phi}_\xi(\xi) d\xi, \end{aligned} \quad (78) \quad \text{algo}$$

which establishes again formula (67) under conditions (36), (37) for  $\tilde{\phi}$ . From (78) it also follows, after some straightforward algebra, the formula

$$\begin{aligned} & \partial_{x'} \left( \sqrt{1-x'^2} \frac{1}{\pi} P.V. \int_{-1}^1 \frac{1}{\sqrt{1-\xi^2}} \frac{1}{x'-\xi} \tilde{\phi}(\xi) d\xi \right) \\ &= \frac{1}{\sqrt{1-x'^2}} \frac{1}{\pi} P.V. \int_{-1}^1 \frac{\sqrt{1-\xi^2}}{x'-\xi} \tilde{\phi}_\xi(\xi) d\xi. \end{aligned} \quad (79) \quad \text{algo2}$$

## APPENDIX B. THE COMPACT EMBEDDING OF $H_{w-1,0}^{\frac{1}{2}}$ INTO $L_{w-1}^2$

Since the spaces  $H_{w-1,0}^{\frac{1}{2}}$  and  $L_{w-1}^2$  can be defined in terms of an orthogonal basis, following [4], let us prove the more general result  $h^{1/2} \xrightarrow{c} \ell^2$  for sequence spaces

$$h^{1/2} := \left\{ (a_n) : \sum_{n=1}^{\infty} n |a_n|^2 < \infty \right\}, \quad \ell^2 := \left\{ (a_n) : \sum_{n=1}^{\infty} |a_n|^2 < \infty \right\}.$$

Consider  $M \subseteq h^{1/2}$  a bounded set in  $h^{1/2}$ . Then, given  $\mathbf{a} \in \ell^2$ ,

$$\|\mathbf{a}\|_{\ell^2}^2 = \sum_{n=1}^{\infty} a_n^2 \leq \sum_{n=1}^{\infty} n a_n^2 = \|\mathbf{a}\|_{h^{\frac{1}{2}}}^2 < K^2,$$

where  $K$  is such that  $\|\mathbf{a}\|_{h^{\frac{1}{2}}} < K, \forall \mathbf{a} \in M$ . We take now  $Y_\varepsilon = \text{span}\{\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_N\}$  where  $N$  will be chosen later as a function of  $\varepsilon$ , and  $\mathbf{e}_i$  is the vector in  $\ell^2$  with all components zero except the  $i$ -th component that is 1. We write  $\mathbf{x} = a_1\mathbf{e}_1 + a_2\mathbf{e}_2 + \dots + a_N\mathbf{e}_N$  so that  $\mathbf{x} \in Y_\varepsilon \subset \ell^2$ ,  $\mathbf{x} = (a_1, a_2, \dots, a_{N-1}, 0, 0, \dots)$  and

$$\|\mathbf{x} - \mathbf{a}\|_{\ell^2}^2 = \sum_{n=N}^{\infty} a_n^2 = \sum_{n=N}^{\infty} \frac{1}{n} n a_n^2 \leq \frac{1}{N} \sum_{n=N}^{\infty} n a_n^2 \leq \frac{K^2}{N} < \frac{\varepsilon}{2},$$

which implies  $N = O(1/\varepsilon)$ . Let  $\mathbf{x} \in \overline{M}$ . There exists  $\mathbf{x}_n \in M$  such that  $\mathbf{x}_n \rightarrow \mathbf{x}$  in the  $\ell^2$  topology. For  $n$  sufficiently large,  $\|\mathbf{x}_n - \mathbf{x}\|_{\ell^2} < \frac{\varepsilon}{2}$  and all  $\mathbf{x}_n$  are at distance  $\frac{\varepsilon}{2}$  of  $Y_\varepsilon$ ; then by the triangle inequality  $\mathbf{x}$  is within  $\varepsilon$  of  $Y_\varepsilon$  and the closure of  $M$  in  $\ell^2$  is compact by Proposition 7.4 in [6] or Proposition 2.1 in [4]. Hence  $h^{1/2}$  is compactly embedded in  $\ell^2$ .

## REFERENCES

- [1] T. Alazard, P. Baldi, and D. Han-Kwan. Control of water waves. *Journal of the European Mathematical Society*, 20(3):657–745, 2018.
- [2] N. Asmar and G. Jones. *Applied Complex Analysis with Partial Differential Equations*. Prentice Hall, 2002.
- [3] T. B. Benjamin and J. C. Scott. Gravity-capillary waves with edge constraints. *Journal of Fluid Mechanics*, 92(2):241–267, 1979.
- [4] J. Bisgard et al. A compact embedding for sequence spaces. *Missouri Journal of Mathematical Sciences*, 24(2):182–189, 2012.
- [5] J.-M. Coron. *Control and nonlinearity*. Number 136. American Mathematical Soc., 2007.
- [6] K. Deimling. *Nonlinear functional analysis*. Dover Publications, Mineola, NY, 2010.
- [7] M. Fontelos and F. De La Hoz. Singularities in water waves and the rayleigh-taylor problem. *Journal of fluid mechanics*, 651:211, 2010.
- [8] M. Fontelos and J. López-Ríos. Gravity waves oscillations at semicircular and general 2d containers: an efficient computational approach to 2d sloshing problem. *Zeitschrift für angewandte Mathematik und Physik*, 71(3):1–24, 2020.
- [9] E. Godoy, A. Osses, J. H. Ortega, and A. Valencia. Modeling and control of surface gravity waves in a model of a copper converter. *Applied Mathematical Modelling*, 32(9):1696–1710, 2008.
- [10] J. Graham-Eagle. A new method for calculating eigenvalues with applications to gravity-capillary waves with edge constraints. In *Mathematical Proceedings of the Cambridge Philosophical Society*, volume 94, pages 553–564. Cambridge University Press, 1983.
- [11] H. Hochstadt. *Integral equations*, volume 91. John Wiley & Sons, 2011.
- [12] H. J. Kim, M. A. Fontelos, and H. J. Hwang. Capillary oscillations at the exit of a nozzle. *IMA Journal of Applied Mathematics*, 80(4):931–962, 2015.
- [13] D. Lannes. *The water waves problem: mathematical analysis and asymptotics*, volume 188. American Mathematical Soc., 2013.
- [14] R. Lecaros, J. López-Ríos, J. Ortega, and S. Zamorano. The stability for an inverse problem of bottom recovering in water-waves. *Inverse Problems*, 36(11):115002, 2020.
- [15] S. Micu and E. Zuazua. An introduction to the controllability of partial differential equations. *Quelques questions de théorie du contrôle. Sari, T., ed., Collection Travaux en Cours Hermann*, pages 69–157.
- [16] H. Nersisyan, D. Dutykh, and E. Zuazua. Generation of 2d water waves by moving bottom disturbances. *IMA Journal of Applied Mathematics*, 80(4):1235–1253, 2014.
- [17] E. Zuazua. *An Introduction to Exact Controllability for Distributed Systems*. Lecture notes, University of Lisbon, Lisbon, Portugal, 1990.
- [18] E. Zuazua. Controllability and observability of partial differential equations: some results and open problems. In *Handbook of differential equations: evolutionary equations*, volume 3, pages 527–621. Elsevier, 2007.

(M. A. Fontelos) (CORRESPONDING AUTHOR) ICMAT-CSIC, C/NICOLÁS CABRERA, NO 13-15 CAMPUS DE CANTOBLANCO, UAM, 28049 MADRID, SPAIN.

Email address: marco.fontelos@icmat.es

(J. López-Ríos) ESCUELA DE CIENCIAS MATEMÁTICAS Y COMPUTACIONALES, YACHAY TECH, SAN MIGUEL DE URQUQUÍ, HACIENDA SAN JOSÉ S/N, ECUADOR

Email address: jlopez@yachaytech.edu.ec