arXiv:2106.00427v3 [math.FA] 1 Jun 2022

WEIGHTED COMPOSITION OPERATORS ON THE FOCK SPACE: ITERATION AND SEMIGROUPS

I. CHALENDAR AND J.R. PARTINGTON

Abstract. This paper considers discrete and continuous semigroups of (weighted) composition operators on the Fock space. For discrete semigroups consisting of powers of a single operator, the asymptotic behaviour of the semigroups is analysed. For continuous semigroups and groups, a full classification of possible semigroups is given, and the generator is calculated.

1. Introduction and notation

For $1 \leq \nu < \infty$ the Fock space $\mathcal{F}^\nu$ consists of all entire functions $f : \mathbb{C} \to \mathbb{C}$ such that the norm

$$\|f\|_\nu := \left(\frac{\nu}{2\pi} \int_{\mathbb{C}} |f(z)|^\nu e^{-\nu|z|^2/2} \, dm(z) \right)^{1/\nu}$$

is finite.

Note that for $k \geq 0$ we have

$$\int_0^\infty r^k e^{-r^2} dr = \frac{1}{2}\Gamma\left(\frac{k+1}{2}\right),$$

from which we see that the functions e_n, $n = 0, 1, 2, \ldots$ such that

$$e_n(z) = z^n/\sqrt{n!} \tag{1}$$

form an orthonormal basis for the Hilbert space $\mathcal{F}^2$.

For entire functions w, φ, f we define the composition operator C_φ by $C_\varphi f = f \circ \varphi$ and the weighted composition operator $W_{w,\varphi}$ by $W_{w,\varphi} f = wC_\varphi f$.

Carswell, MacCluer and Schuster [8] characterised the bounded composition operators C_φ on $\mathcal{F}^2$ as follows:

2010 *Mathematics Subject Classification.* 47B33, 30H10, 30H20, 30D05.

Key words and phrases. Fock space, composition operator, weighted composition operator, iteration, strongly continuous semigroup.

Proposition 1.1. *The operator C_φ is bounded on $\mathcal{F}^2$ if and only if one of the following conditions holds:*

- *$\varphi(z) = az + b$ with $|a| < 1$, $b \in \mathbb{C}$;*
- *$\varphi(z) = az$ with $|a| = 1$.*

In the case $|a| < 1$ the operator C_φ is compact.

For weighted composition operators $W_{w,\varphi}$ we have the result of Le [15], extended by Hai and Khoi [13] as follows:

Proposition 1.2. *The weighted composition operator $W_{w,\varphi}$ with w not identically zero is bounded on $\mathcal{F}^\nu$ if and only if both the following conditions hold:*

- *$w \in \mathcal{F}^\nu$;*
- *$M(w,\varphi) := \sup_{z\in\mathbb{C}} |w(z)|^2 e^{(|\varphi(z)|^2-|z|^2)} < \infty$.*

Moreover, in this case $\varphi(z) = az + b$ with $|a| \leq 1$. If $|a| = 1$ then we also have

$$w(z) = w(0)e^{-\overline{b}az} \tag{2}$$

for $z \in \mathbb{C}$.

See also [7, 16, 18] for a recent discussion of these results.

Zhao [19] gave a criterion for the invertibility of a weighted composition operator on the Fock space; using the more recent results above we can simplify this as follows.

Proposition 1.3. *A weighted composition $W_{w,\varphi}$ is invertible on $\mathcal{F}^\nu$ if and only if $\varphi(z) = az + b$ with $|a| = 1$, and $w(z) = w(0)e^{-\overline{b}az}$ with $w(0) \neq 0$.*

Proof. Suppose that $W_{w,\varphi}$ is invertible. Then w must never vanish as if it did then any zeros of w would also be zeros of $W_{w,\varphi}f$ for every $f \in \mathcal{F}^\nu$ and $W_{w,\varphi}$ would not be surjective. For $f \in \mathcal{F}^\nu$ set $g(z) = w(z)f(az+b)$ with $0 < |a| \leq 1$. Then $f(\zeta)w(a^{-1}\zeta - a^{-1}b) = g(a^{-1}\zeta - a^{-1}b)$ for $\zeta \in \mathbb{C}$. Hence the inverse of $W_{w,\varphi}$ is also a weighted composition operator $W_{v,\psi}$

with $\psi(\zeta) = \varphi^{-1}(\zeta) = a^{-1}\zeta - a^{-1}b$, and thus $|a| = 1$ if $W_{v,\psi}$ is bounded. We also have a formula for w, namely (2). Similarly,

$$v(z) = v(0)e^{a\overline{b}a^{-1}z} = v(0)e^{\overline{b}z}$$

and since

$$1/w(a^{-1}z - a^{-1}b) = w(0)^{-1}e^{\overline{b}a(a^{-1}z - a^{-1}b)}$$

we find that $v(z) = 1/w(a^{-1}z - a^{-1}b)$ if $v(0) = w(0)^{-1}e^{-|b|^2}$. □

This result will be of further use to us when we discuss groups of weighted composition operators.

The paper is organised as follows. In Section 2 we study the asymptotics of iteration of bounded composition operators on Fock spaces. Theorem 2.2 is a complete answer to this problem. Then we consider C_0 semigroups of composition operators $(C_{\varphi_t})_{t\geq 0}$, providing explicit expressions of the semiflow $(\varphi_t)_{t\geq 0}$ in Theorem 2.3. We also prove that every C_0 semigroup $(T_t)_{t\geq 0}$ on $\mathcal{F}^2$ with generator $Af = Gf'$ is a semigroup of composition operators, which implies that $G(z) = az + b$ with $\mathrm{Re}(a) \leq 0$ and $b \in \mathbb{C}$. Moreover the convergence in norm as $t \to \infty$ of such $(T_t)_{t\geq 0}$ is equivalent to the condition $\mathrm{Re}(a) < 0$. We end this section with Theorem 2.10, which is a criterion, on the space $\mathcal{F}^2$ and for a generator A, for generating an analytic semigroup of composition operators.

Section 3 is concerned with weighted composition operators $W_{w,\varphi}$. We first characterize, in Theorem 3.1, the weights for which $W_{w,\varphi}$ is power-bounded on $\mathcal{F}^\nu$ in the particular case where $\varphi(z) = az + b$ with $|a| = 1$. The case $|a| < 1$ is much more delicate. We restrict our study to the case when the weight w does not vanish, which is the relevant situation for semigroups of weighted composition operators, as shown in Theorem 3.8. We consider two cases and split our results into Theorem 3.3 and Theorem 3.4. We also derive a result on the converegnce on the asymptotic behaviour of the iterates of $W_{w,\varphi}$ in Corollary 3.6, thanks to the description of its eigenvalues and eigenspaces described

in Theorem 3.5. The last part of the paper is a complete description of C_0 semigroups and C_0 groups of weighted composition operators on $\mathcal{F}^\nu$.

2. Composition operators

We begin with (unweighted) composition operators, where some fairly complete results will be derived.

2.1. **Iteration.** Note first that for a composition operator C_φ with $\varphi(z) = az + b$, and $a \neq 1$ we have

$$C_\varphi^n f(z) = f\left(a^n z + \frac{1-a^n}{1-a} b\right) \qquad \text{for} \quad n = 1, 2, 3, \ldots,$$

as is easily seen by induction. Also, for $\varphi(z) = z + b$ we clearly have

$$C_\varphi^n f(z) = f(z + nb) \qquad \text{for} \quad n = 1, 2, 3, \ldots.$$

There is a general convergence criterion for powers of an operator $T \in \mathcal{L}(X)$, where X is a Banach space, which may be found in [3, A-III, Sec. 3.7]; it was applied to composition operators in [2], and to weighted composition operators in [9].

An operator $T \in \mathcal{L}(X)$ is power-bounded if $\sup_{n\geq 1} \|T^n\| < \infty$.

Proposition 2.1. *Let $T \in \mathcal{L}(X)$ be power-bounded. Then the following assertions are equivalent:*
(i) $P := \lim T^n$ exists in $\mathcal{L}(X)$ and is of finite rank.
(ii) (a) $r_e(T) < 1$;
(b) $\sigma_p(T) \cap \mathbb{T} \subseteq \{1\}$;
(c) if 1 is in $\sigma(T)$ then it is a pole of the resolvent of order 1.
In that case, P is the residue at 1.

The following formula for $\|C_\varphi\|$ is given in [8] (see also [10]). If $\varphi(z) = az + b$ with $|a| < 1$, then

$$\|C_\varphi\| = \exp\left(\frac{1}{4}\frac{|b|^2}{1-|a|^2}\right). \tag{3}$$

Using these results we may prove the following.

Theorem 2.2. *The asymptotics of iteration of bounded composition operators on the Fock space $\mathcal{F}^\nu$ are as follows.*

(1) If $\varphi(z) = az$ with $|a| = 1$ and $a \neq 1$ then the sequence (C_φ^n) consists of unitary operators; it does not converge, even weakly.
(2) If $\varphi(z) = az + b$ with $|a| < 1$ then the sequence (C_φ^n) converges in norm to the operator mapping a function f to the constant function with value $f(b/(1-a))$.

Proof. In the first case the operators C_φ^n are clearly unitary; by considering the function $f(z) = z$ and $C_\varphi^n f(z) = f(a^n z)$ we see that weak convergence does not hold.

In the second case, the operator C_φ is power-bounded by (3) since if $a_n = a^n$ and $b_n = \frac{1-a^n}{1-a} b$, then $|b_n|^2/(1-|a_n|^2) \to |b|^2/|1-a|^2$. Now we can use Proposition 2.1. Since C_φ is compact, we have $r_e(C_\varphi) = 0$. For the point spectrum we suppose there is an eigenvalue λ of modulus 1 and eigenfunction f satisfying $f(az+b) = \lambda f(z)$. Iterating, we have

$$f(a^n z + b(1-a^n)/(1-a)) = \lambda^n f(z). \tag{4}$$

That is, $\lambda^n f(z) \to f(b/(1-a))$; but z was arbitrary, so f is identically 0 if $|\lambda| = 1$ but $\lambda \neq 1$.

Finally, if $1 \in \sigma(C_\varphi)$ then it is an eigenvalue, since C_φ is compact. Then (4) still holds for an eigenfunction f, and $f(z) = f(b/(1-a))$ so f is constant.

Since $\ker(\mathrm{Id} - C_\varphi)$ is 1-dimensional, we see that 1 is a pole of the resolvent of order at most 1. Indeed, assume that the pole order of 1 is larger than 1. Looking at the Jordan normal form of $T_0 := C_{\varphi|X_0}$, where $X_0 = P\mathcal{F}^\nu$ and P the residue, we see that there exists $f \in \mathcal{F}^\nu$ such that $(\mathrm{Id} - C_\varphi)f = 1_{\mathbb{C}}$. Evaluating at $z = \frac{b}{1-a}$ we obtain a contradiction. Thus the pole order is 1. It follows that P is the projection onto $\ker(C_\varphi - \mathrm{Id}) = \mathbb{C}1_{\mathbb{C}}$ along $\{f \in \mathcal{F}^\nu : f(b/(1-a)) = 0\}$. Thus $Pf = f(b/(1-a))1_{\mathbb{C}}$. This completes the proof that $C_\varphi^n \to P$ in norm thanks to Proposition 2.1. Thus (C_φ^n) converges in norm, and the

limit is the same as the pointwise limit, namely the rank-1 operator T with $Tf(z) = f(b/(1-a))$ for all f. □

2.2. **Semigroups.** The theory of C_0 semigroups (strongly continuous semigroups) is well described in many books, for example [1, 11, 17].

Recall that a C_0 semigroup $(T_t)_{t\geq 0}$ of bounded operators on a Banach space X satisfies the conditions

(1) $T_0 = \mathrm{Id}$;
(2) $T_{s+t} = T_s T_t$ for all $s, t \geq 0$;
(3) the mapping $t \mapsto T_t x$ is continuous for all $x \in X$.

The infinitesimal generator A of such a semigroup $(T_t)_{t\geq 0}$ is defined by

$$Ax = \lim_{t\to 0} \frac{T_t x - x}{t} \qquad (x \in \mathrm{dom}(A)). \tag{5}$$

In the situation when $T_t = W_{w_t,\varphi_t}$ for all $t \geq 0$ we therefore have

$$W_{w_{s+t},\varphi_{s+t}} = W_{w_s,\varphi_s} W_{w_t,\varphi_t} = W_{w_s(w_t\circ\varphi_s),\varphi_t\circ\varphi_s},$$

so that

$$w_{s+t} = w_s(w_t \circ \varphi_s) \text{ and } \varphi_{s+t} = \varphi_t \circ \varphi_s. \tag{6}$$

We write $\mathbb{C}_- := \{z \in \mathbb{C} : \mathrm{Re}\, z < 0\}$.

From now on all semigroups will be assumed to be strongly continuous.

Theorem 2.3. *A semigroup of bounded composition operators $(T_t)_{t\geq 0} = (C_{\varphi_t})_{t\geq 0}$ on $\mathcal{F}^\nu$ satisfies one of the following for $t > 0$:*

(1) $\varphi_t(z) = \exp(\lambda t)z + C(\exp(\lambda t) - 1)$ for some $\lambda \in \mathbb{C}_-$ and $C \in \mathbb{C}$;
(2) $\varphi_t(z) = \exp(\lambda t)z$ for some $\lambda \in i\mathbb{R}$.

The infinitesimal generator is given by $Af(z) = \lambda(z + C)f'(z)$, where $C = 0$ in case (2). Moreover $(T_t)_{t\geq 0}$ converges in norm as $t \to \infty$ if and only if $\lambda \in \mathbb{C}_-$, and it converges to the operator mapping a function f to the constant function with value $f(-C)$.

Proof. From Proposition 1.1 we may write $\varphi_t(z) = a_t z + b_t$; we have $a_{t+s} = a_t a_s$, so $a_t = \exp(\lambda t)$ for some $\lambda \in \mathbb{C}_- := \{z \in \mathbb{C} : \mathrm{Re}\, z < 0\}$ (if

a_t is identically zero then the semigroup is not strongly continuous at 0 since it would require $f(b_t) \to f(z)$ as $t \to 0$ for all functions f and for all $z \in \mathbb{C}$). Also, $b_{s+t} = a_t b_s + b_t = a_s b_t + b_s$. Thus we have

$$b_s(\exp(\lambda t) - 1) = b_t(\exp(\lambda s) - 1),$$

i.e., $b_t = C(\exp(\lambda t) - 1)$ for some constant $C \in \mathbb{C}$. If $\operatorname{Re}\lambda = 0$, then by Proposition 1.2 we have boundedness only for $C = 0$.

Then we have

$$Af(z) = \lim_{t\to 0} \frac{f(e^{\lambda t}z + C(e^{\lambda t} - 1)) - f(z)}{t} = \lambda(z + C)f'(z).$$

The last assertion is an obvious corollary of Theorem 2.2. □

A semigroup $(T_t)_{t\geq 0}$ is said to be quasicontractive if for some $\omega \in \mathbb{R}$ $\|T_t\| \leq e^{\omega t}$ for all $t \geq 0$.

Remark 2.4. *If $\varphi_t(z) = \exp(\lambda t) + C(\exp(\lambda t) - 1)$ with $\operatorname{Re}\lambda < 0$ then by (3), on the Fock space $\mathcal{F}^\nu$, we have*

$$\|T_t\| = \exp\left(\frac{1}{4}\frac{|C^2(\exp(\lambda t) - 1)|^2}{1 - |e^{2\lambda t}|}\right). \tag{7}$$

A routine calculation shows that

$$\frac{|\exp(\lambda t) - 1|^2}{1 - |e^{2\lambda t}|} = O(t)$$

as $t \to 0$. Hence (T_t) is a quasicontractive semigroup.

In [5, Thm. 3.9] It was shown that the following conditions are equivalent for a semigroup with generator $A : f \mapsto G(z)f'(z)$ on the Hardy space H^2, where $G \in H^2$:

(1) A generates a C_0 semigroup of composition operators on $H^2(\mathbb{D})$;
(2) A generates a quasicontractive semigroup on $H^2(\mathbb{D})$;
(3) $\operatorname{ess\,sup}_{z\in\mathbb{T}} \operatorname{Re}(\overline{z}G(z)) \leq 0$.

In the light of Theorem 2.3 and the following proposition we see that similar condition on G applies in the case of the Fock space, namely

$$\limsup_{|z|\to\infty} \operatorname{Re}(\overline{z}G(z)) \leq 0. \tag{8}$$

Proposition 2.5. *For an entire function G, condition* (8) *holds if and only if $G(z) = az + b$ with either (i) $a \in \mathbb{C}_-$ and $b \in \mathbb{C}$, or else (ii) $a \in i\mathbb{R}$ and $b = 0$.*

Proof. Since $\overline{z}(az + b) = a|z|^2 + b\overline{z}$, it is clear that functions G of the form above do satisfy (8).

Conversely, for an entire function G let $F(z) = (G(z) - G(0))/z$. Thus $\overline{z}G(z) = |z|^2F(z) + \overline{z}G(0)$. For G to satisfy (8) we must have $\mathrm{Re}(|z|^2F(z) + \overline{z}G(0)) < 1$, and hence $\mathrm{Re}(F(z) + G(0)/z) < 1$, for sufficiently large $|z|$. This is clearly impossible if F is a nonconstant polynomial, and also impossible if F has an essential singularity at ∞ (for example, by the Weierstrass–Casorati theorem). So F must be constant, and the conditions on a and b are now immediate. □

Next we consider the numerical range of a semigroup generator $A : f \mapsto G(z)f'(z)$ on the Fock space $\mathcal{F}^2$, namely

$$W(A) = \{\langle Af, f\rangle : f \in \mathcal{D}(A), \|f\| = 1\},$$

where $\mathcal{D}(A)$ denotes the domain of A. Recall that a semigroup is quasicontractive if and only if $W(A)$ is contained in some left half-plane.

Proposition 2.6. *If an entire function G is such that the operator $A : f \mapsto Gf'$ is the generator of a quasicontractive C_0 semigroup on $\mathcal{F}^2$, then G is a polynomial of degree at most* 1*, say $G(z) = az + b$, with* $\mathrm{Re}\, a \leq 0$.

Proof. We write $G(z) = \sum_{k=0}^{\infty} g_k z^k$ and examine the coefficients g_k. First, we calculate $\langle Af, f\rangle$ for $f(z) = e_n(z)$ as defined in (1).

$$\langle Af, f\rangle = \langle Gf', f\rangle = g_1 n\langle z^n, z^n\rangle/n! = ng_1.$$

Now if A generates a quasicontractive C_0 semigroup, there must be an upper bound on $\mathrm{Re}\, W(A)$, and hence $\mathrm{Re}\, g_1 \leq 0$.

Next, consider the function $f(z) = az^n + bz^{n+m}$ where $m \geq 1$ is fixed. We see that $\|f\|^2 = |a|^2 n! + |b|^2(n + m)!$.

Also

$$\begin{aligned}\langle Af, f\rangle &= \langle Gf', f\rangle \\ &= g_1(n|a|^2 n! + (n+m)|b|^2(n+m)!) + g_{m+1}(n+m)a\overline{b}(n+m)!.\end{aligned}$$

Now take $a = n^{m/2}\omega$ where $|\omega| = 1$, and $b = 1$, so that $\|f\|^2 = O(n!n^m)$ and

$$\langle Af, f\rangle = g_1(n^{m+1}n! + (n+m)(n+m)!) + g_{m+1}(n+m)(n^{m/2}\omega)(n+m)!.$$

That is, if $g_{m+1} \neq 0$, then for a suitable choice of ω, $\mathrm{Re}\langle Af, f\rangle/\|f\|^2$ grows as $|g_{m+1}|n^{m/2+1}$ as $n \to \infty$; hence $g_{m+1} = 0$ for all $m \geq 1$ if A is the generator of a quasicontractive C_0 semigroup. □

We can now go further and describe all semigroups with generators $A : f \mapsto Gf'$. In [4] there is a result applying to function spaces, i.e., Banach spaces X of holomorphic functions on a domain Ω for which point evaluations are continuous functionals. The result applies to such spaces satisfying the following supplementary condition:

(E) If (z_n) is a sequence in Ω such that $z_n \to z \in \overline{\Omega} \cup \{\infty\}$ and $\lim f(z_n)$ exists in $\mathbb{C}$ for all $f \in X$ then $z \in \Omega$.

Theorem 2.7. *Suppose that $(T_t)_{t\geq 0}$ is a C_0 semigroup on a function space X with property (E) such that the generator A is the operator $f \mapsto Gf'$ for all $f \in \mathrm{dom}(A)$ for some $G \in \mathrm{Hol}(\Omega)$. Then (T_t) is a semigroup of composition operators.*

Similar results are given in [12], for function spaces on the disc $\mathbb{D}$.

Theorem 2.7 clearly applies to the space $X = \mathcal{F}^2$ with $\Omega = \mathbb{C}$ and so we have the following corollary.

Corollary 2.8. *Every C_0 semigroup on $\mathcal{F}^2$ with generator $A : f \mapsto Gf'$ for some $G \in \mathrm{Hol}(\mathbb{C})$ consists entirely of composition operators.*

We may also consider the case of analytic semigroups, analogously to [6]. For $0 < \theta < \pi/2$ we define the sector Σ_θ by

$$\Sigma_\theta = \{re^{i\alpha} : r > 0, |\alpha| < \theta\},$$

and recall that an analytic (or holomorphic) semigroup is one for which there is an analytic mapping $T : \Sigma_\theta \to \mathcal{L}(X)$ satisfying the semigroup property, and which, on defining $T(0) = \mathrm{Id}$, is strongly continuous at 0, and satisfies $\sup_{\xi\in\Sigma_\theta\cap\mathbb{D}} \|T_\xi\| < \infty$.

We may test the analytic property by using Corollary 2.3 of [6], which asserts that if A is a Hilbert-space operator and A and $e^{\pm i\theta}A$ generate quasicontractive semigroups for some $0 < \theta < \pi/2$, then A generates an analytic semigroup on the sector Σ_θ.

As a variation on Corollary 2.5 of [6], we also have the following result.

Lemma 2.9. *Let $(T_t)_{t\geq 0}$ be a semigroup of composition operators on $\mathcal{F}^2$ and suppose that it has an analytic extension to Σ_θ for some $0 < \theta < \pi/2$. Then for every $\xi \in \Sigma_\theta$ the operator T_ξ is a composition operator.*

Proof. We note first that, writing $e_n(z) = z^n$ for $z \in \mathbb{C}$ we have $T_t(e_n) = (T_t(e_1))^n$ for all $n \in \mathbb{N}$ and $t \geq 0$. Now defining $F(\xi) = T_\xi e_n - (T_\xi e_1)^n$ we have that $\xi \mapsto F(\xi)(z_0)$ is an analytic function on Σ_θ for each $z_0 \in \mathbb{C}$; it vanishes on $\mathbb{R}_+$, and hence is identically zero. Thus by linearity and continuity $T_\xi = C_{\varphi_\xi}$ for each ξ, where $\varphi_\xi = T_\xi e_1$. □

We thus have the following result.

Theorem 2.10. *For an entire function G the operator $A : f \mapsto Gf'$ is the generator of an analytic semigroup of operators on $\mathcal{F}^2$ if and only if there is $\theta \in (0, \pi/2)$ such that*

$$\limsup_{|z|\to\infty} \mathrm{Re}(e^{\pm i\theta}\overline{z}G(z)) \leq 0. \tag{9}$$

In this case, the semigroup consists of composition operators and is quasicontractive.

3. Weighted composition operators

As is the case for weighted composition operators on other function spaces, the results that one can prove are necessarily less complete

than the results for (unweighted) composition operators. Nevertheless we may make some progress here.

3.1. **Iteration.** For the weighted composition operator $W_{w,\varphi}$ with $\varphi(z) = az + b$ and $a \neq 1$ we have

$$W^n_{w,\varphi} f(z) = w(z)w(\varphi(z))w(\varphi^2(z)) \dots w(\varphi^{n-1}(z)) f\left(a^n z + \frac{1-a^n}{1-a} b\right)$$

for $n = 1, 2, 3, \dots$, where we denote the kth iterate of φ by φ^k.

For $a = 1$ the corresponding formula is

$$W^n_{w,\varphi} f(z) = w(z)w(\varphi(z))w(\varphi^2(z)) \dots w(\varphi^{n-1}(z)) f\left(z + nb\right).$$

A necessary and sufficient condition for boundedness of $W_{w,\varphi}$ on $\mathcal{F}^\nu$ is given above in Proposition 1.2.

We do not have an exact expression for the norm of a weighted composition operator, but from [13] we know that for $\varphi(z) = az + b$ with $|a| \leq 1$ we have

$$\sqrt{M(w,\varphi)} \leq \|W_{w,\varphi}\| \leq \sqrt{M(w,\varphi)}/|a|. \tag{10}$$

Writing $M_z(w,\varphi) = |w(z)|^2 e^{(|\varphi(z)|^2 - |z|^2)}$ we have that $W_{w,\varphi}$ is compact if and only if $|a| < 1$ and $M_z(w,\varphi) \to 0$ as $|z| \to \infty$.

For power-boundedness, some results are given in [18], with applications to mean ergodicity. We proceed independently, and our main concern is with discrete and continuous semigroups.

First, it is possible to give an exact answer in the case $|a| = 1$.

Theorem 3.1. *Suppose that $\varphi(z) = az + b$ with $|a| = 1$. Then $W_{w,\varphi}$ is power-bounded on $\mathcal{F}^\nu$ if and only if $|w(0)| \leq e^{-|b|^2/2}$.*

Proof. (i) We begin with the case $a = 1$, and with $w(z) = w(0)e^{-\overline{b}az}$ we have

$$W^n_{w,\varphi} f(z) = w(z)w(z+b) \dots w(z+(n-1)b) f(z+nb).$$

That is, it is a weighted composition operator of the form W_{w_n,φ_n}, and we have

$$\begin{aligned} M_z(w_n,\varphi_n) &= |w_n(z)|^2 e^{|\varphi_n(z)|^2-|z|^2} \\ &= |w(0)|^{2n}|e^{-2\overline{b}z}|\,|e^{-2\overline{b}(z+b)}|\dots|e^{-2\overline{b}(z+(n-1)b)}|e^{|z+nb|^2-|z|^2} \\ &= |w(0)|^{2n}|e^{-2n\overline{b}z}|e^{-n(n-1)|b|^2}e^{2\operatorname{Re} nz\overline{b}}e^{n^2|b|^2} \\ &= |w(0)|^{2n}e^{n|b|^2} \end{aligned}$$

and this is uniformly bounded in n (and z) if and only if $|w(0)| \leq e^{-|b|^2/2}$.

(ii) For $a \in \mathbb{T}\setminus\{1\}$ we have

$$\begin{aligned} W^n_{w,\varphi}f(z) &= w(z)w(az+b)w\left(a^2z+\frac{1-a^2}{1-a}b\right)\dots w\left(a^{n-1}z+\frac{1-a^{n-1}}{1-a}b\right) \\ &\quad \times f\left(a^nz+\frac{1-a^n}{1-a}b\right), \end{aligned}$$

which again we write as W_{w_n,φ_n}. We now have

$$w_n(z) = w(0)^n e^{-\overline{b}az}e^{-\overline{b}a(az+b)}\dots e^{-\overline{b}a(a^{n-1}z+\frac{1-a^{n-1}}{1-a}b)}$$

and

$$\begin{aligned} M_z(w_n,\varphi_n) &= |w_n(z)|^2 e^{|\varphi_n(z)|^2-|z|^2} \\ &= |w(0)|^{2n}e^{-2|b|^2\operatorname{Re}(a(1+\frac{1-a^2}{1-a}+\dots+\frac{1-a^{n-1}}{1-a}))}e^{-2\operatorname{Re}(\overline{b}az\frac{1-a^n}{1-a})} \\ &\quad \times e^{|a^nz+\frac{1-a^n}{1-a}b|^2-|z|^2} \end{aligned} \tag{11}$$

and

$$e^{|a^nz+\frac{1-a^n}{1-a}b|^2-|z|^2} = e^{|\frac{1-a^n}{1-a}b|^2+2\operatorname{Re}(a^nz\overline{b}\frac{1-\overline{a}^n}{1-\overline{a}})} = e^{|\frac{1-a^n}{1-a}b|^2+2\operatorname{Re}(az\overline{b}\frac{1-a^n}{1-a})}$$

since $|a|=1$.

This gives

$$M_z(w_n,\varphi_n) = |w(0)|^{2n}e^{|\frac{1-a^n}{1-a}b|^2}e^{-2|b|^2\operatorname{Re}[(n-1)a/(1-a)-a^2(1-a^{n-1})/(1-a)^2]}$$

Note that for $|a|=1$ and $a\neq 1$ we have

$$\operatorname{Re}(a/(1-a)) = \frac{1}{2}\left(\frac{a}{1-a}+\frac{\overline{a}}{1-\overline{a}}\right) = \frac{1}{2}\left(\frac{a}{1-a}+\frac{1}{a-1}\right) = -\frac{1}{2},$$

so that

$$\sup_{z\in\mathbb{C}} M_z(w_n,\varphi_n) \leq K|w(0)|^{2n}e^{|b|^2(n-1)},$$

where K is independent of n; hence uniform boundedness of the powers $W^n_{w,\varphi}$ on $\mathcal{F}^\nu$ is again equivalent to the condition $|w(0)| \le e^{-|b|^2/2}$. $\square$

The analysis is apparently intractable in the case $|a| < 1$, however we may consider the case when the weight w does not vanish, which will be of importance when we come to consider semigroups. In this case, writing $\beta = 1 - |a|^2$ we have the following result from [7].

Theorem 3.2. *For $\varphi(z) = az + b$ with $|a| < 1$ and w nonvanishing the operator $W_{w,\varphi}$ is bounded on $\mathcal{F}^\nu$ if and only if*

$$w(z) = e^{p+qz+rz^2}$$

and either:
(i) $|r| < \beta/2$, in which case $W_{w,\varphi}$ is compact on $\mathcal{F}^\nu$, or
(ii) $|r| = \beta/2$ and, with $t = q + \overline{b}a$, one has either $t = 0$ or else $t \neq 0$ and $r = -\frac{\beta}{2}\frac{t^2}{|t|^2}$. In case (ii) $W_{w,\varphi}$ is not compact on $\mathcal{F}^\nu$.

The analysis splits into two cases:

Case 1. We take $\varphi(z) = az + b$ with $0 < |a| < 1$, and $w(z) = e^{p+qz}$. That is, $r = 0$.

Here we once again have (11) with $w(0) = e^p$ but to check boundedness all we have is (10).

Case 2. We take $\varphi(z) = az + b$ with $0 < |a| < 1$, and $w(z) = e^{p+qz+rz^2}$ with $0 < |r| \le \beta/2$.

In Case 1, we now have

$$\begin{aligned} M_z(w_n, \varphi_n) &= |w_n(z)|^2 e^{|\varphi_n(z)|^2 - |z|^2} \\ &= |w(0)|^{2n} |e^{2q(z+az+\ldots+a^{n-1}z)}| |e^{2qb(1+\frac{1-a^2}{1-a}+\ldots\frac{1-a^{n-1}}{1-a})}| e^{|\varphi_n(z)|^2-|z|^2} \\ &= e^{2n\,\mathrm{Re}\,p} e^{2\,\mathrm{Re}[qz(1-a^n)/(1-a)]} e^{2\,\mathrm{Re}\,qb[n/(1-a) - a(1-a^{n-1})/(1-a)^2]} \\ &\quad \times e^{|a^n z + \frac{1-a^n}{1-a} b|^2 - |z|^2} \qquad (12) \end{aligned}$$

Taking logarithms, we have the following:

Theorem 3.3. *In Case 1:*
if $\mathrm{Re}\,p + \mathrm{Re}\left(\frac{qb}{1-a}\right) > 0$ then $W_{w,\varphi}$ is not power-bounded on $\mathcal{F}^\nu$;
if $\mathrm{Re}\,p + \mathrm{Re}\left(\frac{qb}{1-a}\right) - \ln|a| < 0$ then $W_{w,\varphi}$ is power-bounded on $\mathcal{F}^\nu$.

Proof. In (12) we have that $a^n \to 0$, and the asymptotic behaviour of $\sup_z M_z(w_n, \varphi_n)$ depends only on the first and third factors. Now use (10). □

Note that this matches the condition of Theorem 3.1 if $|a| = 1$ and $q = -\overline{b}a$.

In Case 2 we have the extra factor $v(z) = e^{rz^2}$ in the weight, and Then

$$\begin{aligned} v(z)v(\varphi(z))\dots v(\varphi_{n-1}(z)) &= e^{rz^2}e^{r(az+b)^2}\dots e^{r(a^{n-1}z+b\frac{1-a^{n-1}}{1-a})^2} \\ &= e^{A+Bz+Cz^2}, \end{aligned}$$

where

$$\begin{aligned} A &= rb^2\left(\left(\frac{1-a}{1-a}\right)^2+\left(\frac{1-a^2}{1-a}\right)^2+\ldots+\left(\frac{1-a^{n-1}}{1-a}\right)^2\right) \\ &= \frac{rb^2}{(1-a)^2}\left(n-1-2a\frac{1-a^{n-1}}{1-a}+a^2\frac{1-a^{2n-2}}{1-a^2}\right), \\ B &= 2rb\left(a\frac{1-a}{1-a}+\ldots+a^{n-1}\frac{1-a^{n-1}}{1-a}\right) \\ &= \frac{2rb}{1-a}\left(a\frac{1-a^{n-1}}{1-a}-a^2\frac{1-a^{2n-2}}{1-a^2}\right) \end{aligned}$$

and

$$C = r + ra^2 + \ldots ra^{2n-2} = r\frac{1-a^{2n}}{1-a^2}.$$

Recall that $|r| \leq (1-|a|^2)/2$.

We therefore have the following result:

Theorem 3.4. *In Case 2:*
if $\operatorname{Re} p + \operatorname{Re}\left(\frac{qb}{1-a}\right) + \operatorname{Re}\left(\frac{rb^2}{(1-a)^2}\right) > 0$ *then* $W_{w,\varphi}$ *is not power-bounded on* $\mathcal{F}^\nu$*;*
if $\operatorname{Re} p + \operatorname{Re}\left(\frac{qb}{1-a}\right) + \operatorname{Re}\left(\frac{rb^2}{(1-a)^2}\right) - \ln|a| < 0$ *then* $W_{w,\varphi}$ *is power-bounded on* $\mathcal{F}^\nu$*.*

Proof. We again have that $a^n \to 0$, and the asymptotic behaviour of $\sup_z M_z(w_n, \varphi_n)$ is as in Case 1, with the additional terms from A and C. Once more we use (10). □

Coming to the question of convergence of iterates, we note that if $|r| < (1 - |a|^2)/2$, then $W_{w,\varphi}$ is compact, and so its essential spectrum is $\{0\}$. We now consider the point spectrum.

Theorem 3.5. *For a compact weighted composition operator $W_{w,\varphi}$ on $\mathcal{F}^\nu$ with w not the zero function, $\varphi(z) = az + b$ and $0 < |a| < 1$, the point spectrum consists of simple eigenvalues.*

Proof. The mapping φ has a Denjoy–Wolff point $\alpha \in \mathbb{C}$, i.e., a solution to $\varphi(\alpha) = \alpha$, namely $\alpha = b/(1 - a)$, to which all its iterates converge pointwise.

Factorizing out any possible zeros at $z = 0$, we may write

$$w(\alpha + z) = z^N \exp(\sum_{n=0}^{\infty} d_n z^n),$$

and let us suppose that $W_{w,\varphi} f = \lambda f$ with

$$f(\alpha + z) = z^M \exp(\sum_{n=0}^{\infty} c_n z^n).$$

Note that these expansions are valid in a disc centered at $z = 0$.

Now $\varphi(\alpha + z) = a(\alpha + z) + b = \alpha + az$, so we have

$$z^N \exp(\sum_{n=0}^{\infty} d_n z^n) a^M z^M \exp(\sum_{n=0}^{\infty} c_n a^n z^n) = \lambda z^M \exp(\sum_{n=0}^{\infty} c_n z^n).$$

We thus have $N = 0$, and $\lambda = a^M e^{d_0} = a^M w(\alpha)$; moreover, for some $k \in \mathbb{Z}$,

$$\sum_{n=1}^{\infty} (d_n + c_n a^n) z^n = \sum_{n=1}^{\infty} c_n z^n + 2\pi i k$$

at least on some disc $\{|z| < r\}$. So $k = 0$ and $c_n = d_n/(1 - a^n)$.

Thus the eigenvector (if it is in the space) is unique up to a constant factor.

□

Putting all these ideas together along with Proposition 2.1 we arrive at the following result.

Corollary 3.6. *For a nontrivial compact and power-bounded weighted composition $W_{w,\varphi}$ on $\mathcal{F}^\nu$ with $\varphi(z) = az + b$ and $0 < |a| < 1$, let $\alpha = b/(1-a)$. Then, provided that $a^M w(\alpha) \notin \mathbb{T}\setminus\{1\}$ for all $M = 0, 1, 2, \ldots$, the sequence $(W^n_{w,\varphi})$ converges in norm to the operator mapping f either to the constant function $f(\alpha)$ if $w(\alpha) = 1$, otherwise to the zero function.*

Remark 3.7. *Alternative approaches to Theorems 3.1, 3.3, 3.4 and 3.5 have been suggested by a referee. Namely, in Theorem 3.1 one can show that the operator $W_{w,\varphi}$ can be written as $w(0)e^{|b|^2/2}$ times a surjective isometry on $\mathcal{F}^\nu$. In Theorems 3.3, 3.4 and 3.5 the operator $W_{w,\varphi}$ is similar to an operator of the form $W_{\tilde{w},\tilde{\varphi}}$ where $\tilde{\varphi}(z) = az$: this approach may simplify the calculations.*

3.2. **Semigroups and groups.** For semigroups of weighted composition operators, say, $T_t f(z) = w_t(z) f(\varphi_t(z))$, we have the relation (6) so if $\varphi_t(z) = a_t z + b_t$ we obtain

$$w_{t+s}(z) = w_s(z) w_t(a_s z + b_s) = w_t(z) w_s(a_t z + b_t) \tag{13}$$

while $a_{t+s} = a_t a_s$ and $b_{s+t} = a_t b_s + b_t = a_s b_t + b_s$.

Theorem 3.8. *A C_0 semigroup $T_t f(z) = w_t(z) f(\varphi_t(z))$ $(t \geq 0)$ of weighted composition operators on the Fock space $\mathcal{F}^\nu$ has one of the following two expressions;*

1. *$\varphi_t(z) = \exp(\lambda t)z + C(\exp(\lambda t) - 1)$ for some $\lambda \in \mathbb{C}_-$ and $C \in \mathbb{C}$; in which case $w_t = e^{p_t + q_t z + r_t z^2}$ with p_t, q_t, r_t given respectively by (16), (15) and (14) below.*
2. *$\varphi_t(z) = \exp(\lambda t)z + C(\exp(\lambda t) - 1)$ for some $\lambda \in i\mathbb{R}$ and $C \in \mathbb{C}$, in which case $w_t(z) = w_t(0) \exp(\overline{C}(\exp(\lambda t) - 1)z)$ and moreover $w_t(0) = e^{\mu t} e^{|C|^2(e^{\lambda t} - 1)}$ for some $\mu \in \mathbb{C}$.*

Proof. The formula for φ_t follows as in Theorem 2.3. We now have two cases to analyse:

(1). The proof of [14, Lem. 2.1(b)], can be adapted and simplified to show that $w_t(z) \neq 0$ for all $t \geq 0$ and $z \in \mathbb{C}$.

For if there are a $z_0 \in \mathbb{C}$ and $t_0 \geq 0$ with $w_{t_0}(z_0) = 0$, then let $I = \{t \geq 0 : w_t(z_0) = 0\}$ and $\tau = \inf I$. Since the semigroup is strongly continuous we have $\tau \in I$ by considering the effect of the semigroup on a constant function; also $\tau > 0$ since $w_0(z) = 1$ for all z. By (13) we have $I = [\tau, \infty)$ and, taking $s = \tau/2$, we have

$$w_t(z_0)w_s(a_t z_0 + b_t) = w_{t+s}(z_0) = 0$$

for all $t \in (\tau/2, \tau)$ and hence $w_s(a_t z_0 + b_t) = 0$ for all such t. Hence w_s is identically zero, as it is zero on a nontrivial continuous path.

Now, by Theorem 3.2, $w_t(z)$ takes the form

$$w_t(z) = \exp(p_t + q_t z + r_t z^2).$$

From (13) we have

$$p_{t+s} + q_{t+s}z + r_{t+s}z^2 = p_s + q_s z + r_s z^2 + p_t + q_t(a_s z + b_s) + r_t(a_s z + b_s)^2,$$

For extra legibility, write $x = \exp(\lambda s)$ and $y = \exp(\lambda t)$; then $a_s = x$ and $b_s = C(x - 1)$; so the equations we have are

$$p_{t+s} + q_{t+s}z + r_{t+s}z^2 = p_s + q_s z + r_s z^2 + p_t + q_t(a_s z + b_s) + r_t(a_s z + b_s)^2,$$

giving

$$\begin{aligned} p_{t+s} &= p_s + p_t + q_t b_s + r_t b_s^2, \\ q_{t+s} &= q_s + q_t a_s + 2r_t a_s b_s, \\ r_{t+s} &= r_s + a_s^2 r_t. \end{aligned}$$

Thus $r_{t+s} = r_s + r_t x^2$, for which the general solution is $r_s = \alpha(1 - x^2)$ or

$$r_t = \alpha(1 - e^{2\lambda t}) \tag{14}$$

for some $\alpha \in \mathbb{C}$. (It is easy to check that $r_t/(1 - e^{2\lambda t})$ is necessarily constant.) The constant α must also be such that $|r_t| \leq (1 - |a_t|^2)/2$ by virtue of Theorem 3.2.

Next $q_{t+s} = q_s + q_t x + 2\alpha C(1-y^2)x(x-1)$, which has solution $q_s = 2\alpha C x(1-x) + \gamma(1-x)$ or

$$q_t = 2\alpha C e^{\lambda t}(1-e^{\lambda t}) + \gamma(1-e^{\lambda t}) \tag{15}$$

for some $\gamma \in \mathbb{C}$.

Finally, $p_{t+s} = p_s + p_t + 2\alpha C^2 y(1-y)(x-1) + C\gamma(1-y)(x-1) + \alpha C^2(1-y^2)(x-1)^2$, giving $p_s = -\alpha C^2(x-1)^2 - \gamma C(x-1) + \delta s$ or

$$p_t = -\alpha C^2(e^{\lambda t}-1)^2 - \gamma C(e^{\lambda t}-1) + \delta t \tag{16}$$

for some $\delta \in \mathbb{C}$.

(2). We require $w_t(z) = w_t(0)e^{-\overline{b_t}a_t z}$ to satisfy (13), that is,

$$w_{t+s}(0)e^{-\overline{b_{s+t}}a_{s+t}z} = w_s(0)e^{-\overline{b_s}a_s z}w_t(0)e^{-\overline{b_t}a_t(a_s z+b_s)}.$$

Observe that

$$\overline{b_{s+t}}a_{s+t} = \overline{b_s}a_s + \overline{b_t}a_t a_s,$$

that is,

$$\overline{C}(e^{-\lambda(s+t)}-1)e^{\lambda(s+t)} = \overline{C}(e^{-\lambda s}-1)e^{\lambda s} + \overline{C}(e^{-\lambda t}-1)e^{\lambda t}e^{\lambda s},$$

since

$$1 - e^{\lambda(s+t)} = 1 - e^{\lambda s} + (1-e^{\lambda t})e^{\lambda s}$$

holds for all s and t.

We also have

$$w_{t+s}(0) = w_s(0)w_t(0)e^{-\overline{b_t}a_t b_s} = w_t(0)w_s(0)e^{-\overline{b_s}a_s b_t}$$

since for this to happen we require

$$|C|^2(1-e^{\lambda t})(e^{\lambda s}-1) = |C|^2(1-e^{\lambda s})(e^{\lambda t}-1)$$

which holds for all s and t. Finally,

$$w_{t+s}(0) = w_s(0)w_t(0)e^{|C|^2(e^{\lambda t}-1)(e^{\lambda s}-1)}$$

or, with $v_t = w_t(0)e^{-|C|^2(e^{\lambda t}-1)}$,

$$v_{t+s} = v_s v_t e^{|C|^2((e^{\lambda t}-1)(e^{\lambda s}-1)-(e^{\lambda(t+s)}-1)+(e^{\lambda t}-1)+(e^{\lambda s}-1)} = v_s v_t$$

so that finally $w_t(0) = e^{\mu t}e^{|C|^2(e^{\lambda t}-1)}$ for some $\mu \in \mathbb{C}$. □

Remark 3.9. *We may also calculate infinitesimal generators using* (5)*, so that for* $T_t(f)(z) = w_t(z)f(\varphi_t(z))$ *with*

$$w_t(z) = \exp(p_t + q_t z + r_t z^2) \text{ and } \varphi_t(z) = a_t z + b_t$$

we have

$$\begin{aligned} Af(z) &= \lim_{t\to 0} w_t(z)\frac{f\circ\varphi_t(z) - f(z)}{t} + \lim_{t\to 0}\frac{w_t f(z) - f(z)}{t} \\ &= \left[\frac{\partial}{\partial t}(a_t z + b_t)f'(z) + \frac{\partial}{\partial t}(p_t + q_t z + r_t z^2)f(z)\right]_{t=0}. \end{aligned}$$

Finally, we may characterise one-parameter groups of weighted composition operators, using Proposition 1.3 and Theorem 3.8.

Corollary 3.10. *A* C_0 *group* $T_t f(z) = w_t(z)f(\varphi_t(z))$ $(t \in \mathbb{R})$ *of weighted composition operators on the Fock space* $\mathcal{F}^\nu$ *satisfies*

$$\varphi_t(z) = \exp(\lambda t)z + C(\exp(\lambda t) - 1)$$

for some $\lambda \in i\mathbb{R}$ *and* $C \in \mathbb{C}$*, in which case*

$$w_t(z) = w_t(0)\exp(\overline{C}(\exp(\lambda t) - 1)z)$$

and moreover $w_t(0) = e^{\mu t}e^{|C|^2(e^{\lambda t}-1)}$ *for some* $\mu \in \mathbb{C}$*.*

Proof. The first case of Theorem 3.8 does not give bounded inverses, but in the second case where $a_t = e^{\lambda t}$ and $b_t = C(e^{\lambda t} - 1)$ with $\lambda \in i\mathbb{R}$, we do have $a_{-t} = a_t^{-1}$ and $b_{-t} = -a_t^{-1}b_t$; also $w_{-t}(0) = w_t(0)^{-1}e^{-|b_t|^2}$, so that $T_{-t} = T_t^{-1}$ by Proposition 1.3. □

Remark 3.11. *It is also possible to consider the question of when a weighted composition semigroup has an analytic extension, analogously to Theorem 2.10. However, the question is simply whether* $e^{\pm i\theta}A$ *also generates a semigroup for some* $0 < \theta < \pi/2$*, which reduces to considering the semigroups in Theorem 3.8 with a complex parameter. Accordingly, we omit the details.*

Acknowledgment. The authors are grateful to a referee for a very careful of the reading of the paper and the correction of numerous errors.

References

[1] David Applebaum. *Semigroups of linear operators*, volume 93 of *London Mathematical Society Student Texts*. Cambridge University Press, Cambridge, 2019. With applications to analysis, probability and physics.

[2] W. Arendt, I. Chalendar, M. Kumar, and S. Srivastava. Asymptotic behaviour of the powers of composition operators on Banach spaces of holomorphic functions. *Indiana Univ. Math. J.*, 67(4):1571–1595, 2018.

[3] W. Arendt, A. Grabosch, G. Greiner, U. Groh, H. P. Lotz, U. Moustakas, R. Nagel, F. Neubrander, and U. Schlotterbeck. *One-parameter semigroups of positive operators*, volume 1184 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1986.

[4] Wolfgang Arendt and Isabelle Chalendar. Generators of semigroups on Banach spaces inducing holomorphic semiflows. *Israel J. Math.*, 229(1):165–179, 2019.

[5] C. Avicou, I. Chalendar, and J. R. Partington. A class of quasicontractive semigroups acting on Hardy and Dirichlet space. *J. Evol. Equ.*, 15(3):647–665, 2015.

[6] C. Avicou, I. Chalendar, and J. R. Partington. Analyticity and compactness of semigroups of composition operators. *J. Math. Anal. Appl.*, 437(1):545–560, 2016.

[7] Tom Carroll and Clifford Gilmore. Weighted composition operators on Fock spaces and their dynamics. *J. Math. Anal. Appl.*, 502(1):125234, 2021.

[8] Brent J. Carswell, Barbara D. MacCluer, and Alex Schuster. Composition operators on the Fock space. *Acta Sci. Math. (Szeged)*, 69(3-4):871–887, 2003.

[9] Isabelle Chalendar and Jonathan R. Partington. Weighted composition operators: isometries and asymptotic behaviour. *J. Operator Theory*, 86(1):189–201, 2021.

[10] Jineng Dai. The norm of composition operators on the Fock space. *Complex Var. Elliptic Equ.*, 64(9):1608–1616, 2019.

[11] Klaus-Jochen Engel and Rainer Nagel. *One-parameter semigroups for linear evolution equations*, volume 194 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 2000. With contributions by S. Brendle, M. Campiti, T. Hahn, G. Metafune, G. Nickel, D. Pallara, C. Perazzoli, A. Rhandi, S. Romanelli and R. Schnaubelt.

[12] Eva A. Gallardo-Gutiérrez and Dmitry V. Yakubovich. On generators of C_0-semigroups of composition operators. *Israel J. Math.*, 229(1):487–500, 2019.

[13] Pham Viet Hai and Le Hai Khoi. Boundedness and compactness of weighted composition operators on Fock spaces $\mathcal{F}^p(\mathbb{C})$. *Acta Math. Vietnam.*, 41(3):531–537, 2016.

[14] Wolfgang König. Semicocycles and weighted composition semigroups on H^p. *Michigan Math. J.*, 37(3):469–476, 1990.

[15] Trieu Le. Normal and isometric weighted composition operators on the Fock space. *Bull. Lond. Math. Soc.*, 46(4):847–856, 2014.

[16] Tesfa Mengestie and Werkaferahu Seyoum. Spectra of composition operators on Fock-type spaces. *Quaest. Math.*, 44(3):335–350, 2021.

[17] A. Pazy. *Semigroups of linear operators and applications to partial differential equations*, volume 44 of *Applied Mathematical Sciences*. Springer-Verlag, New York, 1983.

[18] Werkaferahu Seyoum and Tesfa Mengestie. Spectrums and uniform mean ergodicity of weighted composition operators on Fock spaces. *Bull. Malays. Math. Sci. Soc. (2)*, 45(1):455–481, 2022.
[19] Liankuo Zhao. Invertible weighted composition operators on the Fock space of $\mathbb{C}^N$. *J. Funct. Spaces*, pages Art. ID 250358, 5, 2015.

Isabelle Chalendar, Université Gustave Eiffel, LAMA, (UMR 8050), UPEM, UPEC, CNRS, F-77454, Marne-la-Vallée (France)
Email address: isabelle.chalendar@univ-eiffel.fr

Jonathan R. Partington, School of Mathematics, University of Leeds, Leeds LS2 9JT, UK
Email address: J.R.Partington@leeds.ac.uk