

Rényi Divergence in General Hidden Markov Models

Cheng-Der Fuh

Fanhai International School of Finance, Fudan University

Su-Chi Fuh

Department of Computer Science and Information Engineering

National Taipei University of Technology

Yuan-Chen Liu

Department of Computer Science, National Taipei University of Education

Chuan-Ju Wang

Research Center for Information Technology Innovation, Academia Sinica

Abstract

In this paper, we examine the existence of the Rényi divergence between two time invariant general hidden Markov models with arbitrary positive initial distributions. By making use of a Markov chain representation of the probability distribution for the general hidden Markov model and eigenvalue for the associated Markovian operator, we obtain, under some regularity conditions, convergence of the Rényi divergence. By using this device, we also characterize the Rényi divergence, and obtain the Kullback–Leibler divergence as $\alpha \rightarrow 1$ of the Rényi divergence. Several examples, including the classical finite state hidden Markov models, Markov switching models, and recurrent neural networks, are given for illustration. Moreover, we develop a non-Monte Carlo method that computes the Rényi divergence of two-state Markov switching models via the underlying invariant probability measure, which is characterized by the Fredholm integral equation.

Key words and phrases. Fredholm integration equation, Kullback–Leibler divergence, Markov switching models, recurrent neural network.

I. INTRODUCTION

Motivated by study of the information divergence in hidden Markov models (HMM), Markov switching models, and recurrent neural networks, we here investigate Rényi divergences in general hidden Markov models. A general hidden Markov model is, loosely speaking, a sequence $\mathbf{Y} = \{Y_n, n \geq 0\}$ of random variables obtained in the following way. First, a realization of a finite state Markov chain $\mathbf{X} = \{X_n, n \geq 0\}$ is created. This chain is sometimes called the regime and is not observed. Then, conditioned on \mathbf{X} , the \mathbf{Y} -variables are generated. Usually, the dependency of Y_n on \mathbf{X} is more or less local, as when $Y_n = g(X_n, \eta_n)$ or $Y_n = g(X_n, Y_{n-1}, \eta_n)$ for some function g and random sequence $\{\eta_n\}$, independent of \mathbf{X} . Y_n itself is generally not Markov and may in fact have a complicated dependency structure.

HMMs have been studied extensively, and have extraordinary applications in fields as varied as speech recognition, cf. Rabiner and Juang (1993), Rabiner (1989); handwritten recognition, cf. Hu et al. (1996); Kunda et al. (1989); human activity recognition, cf. Yamato et al. (1992); target detection and tracking, cf. Blanding et al. (2009); Tuğaç and Efe (2010); Vasuhi and Vaidehi (2014); computational molecular biology and bioinformatics, including DNA and protein modeling, cf. Churchill (1989); modeling, rapid detection, and tracking of malicious activity of terrorist groups, cf. Raghavan et al. (2013), Raghavan et al. (2014); and others, with just a small sample of references given for each application. A comprehensive survey of HMM research and applications can be found in Ephraim and Merhav (2002), Cappe et al. (2005), and Zucchini and MacDonald (2009), including an extensive bibliography.

A natural extension of the celebrated HMM is the following Markov switching models. We start with a simple real-valued first-order autoregression around one of two constants μ_1 or μ_2 :

$$Y_n = \mu_{X_n} + \psi Y_{n-1} + \varepsilon_n, \quad (1.1)$$

where $\varepsilon_n \sim N(0, \sigma^2)$, $|\psi| < 1$, and $\{X_n, n \geq 0\}$ is a 2-state Markov chain. When $\psi = 0$, (1.1) reduces to the classical Gaussian HMM.

Another interesting example is the recurrent neural network (RNN) in machine learning. Note that the RNN can take as input a variable-length sequence $y = (y_1, \dots, y_n)$ by recursively processing each symbol while maintaining its internal hidden state h . At each time step n , the RNN reads the symbol $Y_n \in \mathbf{R}^q$ and updates its hidden state $h_n \in \mathbf{R}^p$ by

$$h_n = f_\theta(Y_n, h_{n-1}), \quad (1.2)$$

where f_θ is a deterministic non-linear transition function, and θ is the parameter of f_θ . The transition function f_θ can be implemented with gated activation functions such as long short-term memory (LSTM), cf. Hochreiter and Schmidhuber (1997), or gated recurrent unit (GRU), cf. Cho et al. (2014). Although the hidden unit h_n is a general state random variable, a transformation in Section IV-B shows that the RNN in (1.2) can be formulated as a general HMM.

The Rényi divergence rate, cf. Rényi et al. (1961), and the Kullback–Leibler divergence in particular, have played a significant role in certain hypothesis-testing questions, cf. Koopmans (1960), Nemetz (1974). Furthermore, the Rényi entropy and the Rényi entropy rate have revealed several operational characterizations in the problem of fixed-length source coding, cf. Csiszar (1995), Chen and Alajaji (2001); unsupervised learning, cf. Jenssen et al. (2003); variable-length source coding, cf. Blumer and McEliece (1988), Campbell (1965), Jelinek (1968), and Rached et al. (2001); error exponent calculations, cf. Erez and Zamir (2001); policy optimization in reinforcement learning, cf. Metelli et al. (2018), Papini et al. (2019), and other areas such as Arikan (1996), Bassat and Raviv (1978), and Pronzato et al. (1997).

It is known that the root of the Kullback–Leibler divergence and the Rényi divergence is the celebrated Shannon entropy. The question of computing the Shannon entropy (or, simply, entropy) of a HMM was studied in an early paper by Blackwell (1959), in which the analysis suggests the intrinsic complexity of expressing the HMM entropy as a function of the process parameters. The author also presents an expression of the entropy in terms of a measure Q , which solves an integral equation dependent on the parameters of the process. In general, the measure is hard to extract from the equation in any explicit way. Fuh and Mei (2015) provide a numerical method to approximate the invariant measure and the Kullback–Leibler divergence for a two-state HMM. The problem of determining the residual noise of the best filter for a HMM was studied in Khasminskii and Zeitouni (1996), Ordentlich and Weissman (2004), Ordentlich and Weissman (2006), and Jacquet et al. (2008) investigate the asymptotic estimates of the HMM entropy rate. Furthermore, Zuk et al. (2005) present formulas for higher-order coefficients of the Taylor expansion in the symmetric case. Han and Marcus (2006a) and Han and Marcus (2006b) characterize the analyticity of the HMM entropy rate, and obtain a broad generalization of the results of Zuk et al. (2005).

For an explicit computation of the Rényi entropy of HMMs over finite alphabets in both finite-length and asymptotic regimes, Wu et al. (2017) discuss some convergence properties with no explicit formulas. For Shannon entropy the problem has been found hard and solvable only for specific cases, being related to an

intractable task in random matrix products—finding top Lapunov exponents, cf. Jacquet et al. (2008). In the case of finite state Markov chains, Rached et al. (2001) apply substochastic matrices in the asymptotic regime powers, which can be approximated by spectral analysis, to yield formulas on entropy rates.

Although there are some interesting papers on computing the Rényi divergence in a special HMM, systematic study for a general HMM is still lacking. To fill this gap, we investigate the Rényi divergence for a general HMM in this paper. We make three contributions. First, we note that a major difficulty for analyzing the Rényi divergence in general HMMs is that the joint probability can be expressed only in summation form; see equations (2.3) and (2.4) in Section II for instance. The contribution in this paper is that we provide a device which represents the joint probability as the L_1 -norm of a product of random matrices and treat it as a Markov chain in an enlarged state space. This representation enables us to apply results of the strong law of large numbers, and spectral theory for the associated Markovian operator of Markov random walks, to yield an explicit characterization of the Rényi divergence, and hence the Kullback–Leibler divergence. Second, our formulation of the HMM in a general sense covers several interesting examples, including finite state HMMs, Markov switching models, and RNNs. Third, we develop a non-Monte Carlo method that computes the Rényi divergence of a two-state Markov switching model via the underlying invariant probability measure, which is characterized by the Fredholm integral equation. For this purpose, we also provide an approximated Rényi divergence. Our numerical study shows that this approximated Rényi divergence is reasonably accurate in some simple cases.

The remainder of this paper is organized as follows. In Section II, we define the HMM as a Markov chain in a Markovian random environment, and represent the probability as the L_1 -norm of a product of Markovian random matrices. Then, we give a brief summary of eigenvalue and eigenfunction for Markovian operators. In Section III, we study the limiting behavior of the probability and characterize the Rényi divergence. The Kullback–Leibler divergence is defined and can be regarded as the limit of $\alpha \rightarrow 1$ in the Rényi divergence. In Section IV, we consider a few examples, including finite state Markov switching models and RNNs, which are commonly used in machine learning. In Section V, we give a numerical computation of the Rényi divergence by applying the Fredholm integral equation for a two-state Markov switching model. Section VI concludes.

II. HIDDEN MARKOV MODELS

A. Hidden Markov Models

In this section, we first provide a probability framework for a general HMM under which it can be regarded as a Markov chain in an enlarged state space. That is, there are two Markov chains associated with the general HMM to be described as follows. First, a general HMM is defined as a parameterized Markov chain in a Markovian random environment with the underlying environmental Markov chain viewed as missing data. Specifically, let $\mathbf{X} = \{X_n, n \geq 0\}$ be a Markov chain on a finite state space $\mathcal{X} = \{1, \dots, d\}$, with transition probability $p_{ij} = P\{X_1 = j | X_0 = i\}$ for $i, j = 1, \dots, d$, and stationary probability π_j . Suppose that a random sequence $\{Y_n\}_{n=0}^\infty$, taking values in \mathbf{R}^q , is adjoined to the chain such that $\{(X_n, Y_n), n \geq 0\}$ is a Markov chain on $\mathcal{X} \times \mathbf{R}^q$ satisfying $P\{X_1 \in A | X_0 = i, Y_0 = y\} = P\{X_1 \in A | X_0 = i\}$ for $A \in \mathcal{B}(\mathcal{X})$, the σ -algebra of \mathcal{X} . Conditioning on the full \mathbf{X} sequence, Y_n is a Markov chain with probability

$$P\{Y_{n+1} \in B | X_0, X_1, \dots; Y_0, Y_1, \dots, Y_n\} = P\{Y_{n+1} \in B | X_{n+1}, Y_n\} \quad a.s. \quad (2.1)$$

for each n and $B \in \mathcal{B}(\mathbf{R}^q)$, the Borel σ -algebra of \mathbf{R}^q . Note that in (2.1) the conditional probability of Y_{n+1} depends on X_{n+1} and Y_n only. Furthermore, we assume the existence of the conditional probability density $f(Y_k | X_k, Y_{k-1})$ of Y_k given X_k and Y_{k-1} with respect to a σ -finite measure \mathcal{L} on \mathbf{R}^q such that

$$P\{X_1 \in A, Y_1 \in B | X_0 = i, Y_0 = y_0\} = \sum_{j \in A} \int_{y \in B} p_{ij} f(y | j, y_0) \mathcal{L}(dy). \quad (2.2)$$

We also assume that the Markov chain $\{(X_n, Y_n), n \geq 0\}$ has a stationary probability with probability density function $\pi_j f(\cdot | j)$ with respect to \mathcal{L} . Now we give a formal definition as follows.

Definition II.1. $\{Y_n, n \geq 0\}$ is called a general hidden Markov model if there is a Markov chain $\{X_n, n \geq 0\}$ such that the process $\{(X_n, Y_n), n \geq 0\}$ is a Markov chain satisfying (2.1). A non-invertible function $g(Y_n)$ of Y_n is also called a general hidden Markov model.

Note that this HMM setting is defined in a general sense, which includes several interesting examples of Markov-switching Gaussian autoregression, cf. Hamilton (1989), and RNNs in machine learning, cf. Goodfellow et al. (2016). When Y_n are conditionally independent given \mathbf{X} , denote $S_n = \sum_{t=1}^n Y_t$. Then the Markov chain $\{(X_n, S_n), n \geq 0\}$ is called a Markov additive process, cf. Ney and Nummelin (1987), and $\{Y_n, n \geq 1\}$ is the celebrated HMM considered in engineering literature.

Next we follow a similar idea in Fuh (2004b) and Fuh and Tartakovsky (2019), to have a Markov chain representation of the probability. Note that the joint probability of the general HMM $\{Y_n, n \geq 0\}$ is

$$\begin{aligned} & P\{Y_0 \in B_0, Y_1 \in B_1, \dots, Y_n \in B_n\} \\ &= \int_{y_1 \in B_1} \cdots \int_{y_n \in B_n} p_n(y_1, \dots, y_n) \mathcal{L}(dy_n) \cdots \mathcal{L}(dy_1), \end{aligned} \quad (2.3)$$

where

$$p_n(y_0, y_1, \dots, y_n) = \sum_{x_0=1}^d \cdots \sum_{x_n=1}^d \nu_{x_0} f(y_0|x_0) \prod_{k=1}^n p_{x_{k-1}x_k} f(y_k|x_k, y_{k-1}), \quad (2.4)$$

where $\nu = (\nu_1, \dots, \nu_d)^t$ is an initial distribution of $\{X_n, n \geq 0\}$, which is positive P -a.s. Here t denotes the transpose of the underlying vector in \mathbf{R}^d

For a given column vector $a = (a_1, \dots, a_d)^t \in \mathbf{R}^d$, define the L_1 -norm of a as $\|a\| = \sum_{k=1}^d |a_k|$. The probability (2.4) can be represented as

$$p_n(y_0, y_1, \dots, y_n) = \|M_n \cdots M_1 M_0 \nu\|, \quad (2.5)$$

where y_0 is given from $f(\cdot|x_0)$ and

$$M_0 = \begin{bmatrix} f(y_0|x_0=1) & 0 & \cdots & 0 \\ 0 & f(y_0|x_0=2) & \cdots & 0 \\ \vdots & 0 & \ddots & \vdots \\ 0 & \cdots & 0 & f(y_0|x_0=d) \end{bmatrix}, \quad (2.6)$$

$$M_k = \begin{bmatrix} p_{11}f(y_k|x_k=1, y_{k-1}) & \cdots & p_{d1}f(y_k|x_k=1, y_{k-1}) \\ \vdots & \ddots & \vdots \\ p_{1d}f(y_k|x_k=d, y_{k-1}) & \cdots & p_{dd}f(y_k|x_k=d, y_{k-1}) \end{bmatrix}, \quad (2.7)$$

for $k = 1, \dots, n$.

Note that, for $k = 1, \dots, n$, the quantity $p_{ij}f(Y_k|X_k = j, Y_{k-1})$ in (2.7) represents $X_{k-1} = i$ and $X_k = j$, and Y_k is a Markov chain with transition probability density $f(y_k|x_k = j, y_{k-1})$ for given \mathbf{X} . By definition (2.1), $\{(X_n, Y_n), n \geq 0\}$ is a Markov chain, which implies that M_k is a sequence of Markovian random matrices. Therefore, by representation (2.5), $p_n(Y_0, Y_1, \dots, Y_n)$ is the L_1 -norm of a product of Markovian random matrices. Furthermore, let

$$T_n = M_n \cdots M_1 M_0. \quad (2.8)$$

Denote $P(\mathbf{R}^d)$ as the projection space on \mathbf{R}^d , $Gl(d, \mathbf{R})$ as the space of $d \times d$ matrices and S^{d-1} as the d -dimensional sphere. For $\bar{u} \in P(\mathbf{R}^d)$, $M \in Gl(d, \mathbf{R})$, let $M \cdot \bar{u} = \overline{Mu}$, and $\nu = \nu(\theta) = (\nu_1, \dots, \nu_d)^t \in S^{d-1}$, the unit sphere with respect to the L_1 -norm $\|\cdot\|$ in \mathbf{R}^d ; we have

$$\log \|T_n \nu\| = \log \frac{\|T_n \nu\|}{\|T_{n-1} \nu\|} + \dots + \log \frac{\|T_0 \nu\|}{\|\nu\|}. \quad (2.9)$$

Let $Z_n = (X_n, Y_n)$; define

$$W_0 = (Z_0, \overline{T_0 \nu}), W_1 = (Z_1, \overline{T_1 \nu}), \dots, W_n = (Z_n, \overline{T_n \nu}). \quad (2.10)$$

Then, W_0, W_1, \dots, W_n is a Markov chain on the state space $\mathcal{S} := \mathcal{X} \times \mathbf{R}^q \times P(\mathbf{R}^d)$ with the transition kernel

$$\mathbb{P}((z, \bar{u}), A \times B) := \mathbb{E}_z(I_{A \times B}(Z_1, \overline{M_1 u})) \quad (2.11)$$

for all $z \in \mathcal{X} \times \mathbf{R}^q$, $\bar{u} \in P(\mathbf{R}^d)$, $A \in \mathcal{X} \times \mathcal{B}(\mathbf{R}^q)$, and $B \in \mathcal{B}(P(\mathbf{R}^d))$, the Borel σ -algebra of $P(\mathbf{R}^d)$.

Note that the initial distribution of W_0 depends on Z_0 only, and Z_0 has the distribution $\nu_i f(y|i, y_0)$ as its initial distribution. We note that $\mathbb{P}_z := \mathbb{P}(\cdot, \cdot)$ in (2.11) depends only on z . Let $\mathbb{E}_z := \mathbb{E}_{(z, \bar{u})}$ denote the expectation under \mathbb{P}_z . By (2.2), the Markov chain $\{(X_n, Y_n), n \geq 0\}$ has transition density $p_{ij} f(y|j, y_0)$ with respect to \mathcal{L} . Therefore, the induced transition probability $\mathbb{P}(\cdot, \cdot)$ has a probability density $p(\cdot, \cdot)$ with respect to \mathcal{L} . Under condition C in Section II-B, it follows from Proposition 2 of Fuh (2004a) or Proposition 1 of Fuh and Tartakovsky (2019) that the Markov chain W_n has an invariant probability measure Π on \mathcal{S} . Note that \mathcal{L} is a product measure on $\mathcal{X} \times \mathbf{R}^q \times P(\mathbf{R}^d)$, and the first component has probability density $\nu_i f(y|i, y_0)$ with respect to \mathcal{L} . Now, for $M \in Gl(d, \mathbf{R})$, let $g : \mathcal{S} \times \mathcal{S} \rightarrow \mathbf{R}$ be $g((z_0, \bar{u}), (z_1, \overline{M u})) = \log \frac{\|M u\|}{\|u\|}$; then for ν defined in (2.4),

$$\log \|T_n \nu\| = g(W_{n-1}, W_n) + \dots + g(W_0, W_1) + g(W_0, W_0) \quad (2.12)$$

is an additive functional of the Markov chain $\{W_n, n \geq 0\}$, where $g(W_0, W_0) = \log \frac{\|T_0 \nu\|}{\|\nu\|}$.

B. Nonnegative transition probability kernel for Markov operator

To study the Kullback–Leibler divergence and Rényi divergence in general HMMs, we must consider a nonnegative transition probability kernel for a Markov operator of the induced Markov chain $\{W_n, n \geq 0\}$ defined in (2.10) on the state space $\mathcal{S} := \mathcal{X} \times \mathbf{R}^q \times P(\mathbf{R}^d)$, with the transition kernel \mathbb{P} in (2.11). Before that we need the following notation.

Note that $\{(X_n, Y_n), n \geq 0\}$ defined in (2.1) and (2.2) is a Markov chain on the state space $\mathcal{X} \times \mathbf{R}^q$. Below, we abuse the notation a bit to consider $\{Y_n, n \geq 0\}$ as a Markov chain on a general state space \mathbf{R}^q .

Definition II.2. A Markov chain $\{Y_n, n \geq 0\}$ on a general state space \mathbf{R}^q is said to be V -uniformly ergodic if there exists a measurable function $V : \mathbf{R}^q \rightarrow [1, \infty)$, with $\int V(y)\mathcal{L}(dy) < \infty$, such that

$$\lim_{n \rightarrow \infty} \sup_{y \in \mathbf{R}^q} \left\{ \frac{|E[h(Y_n)|Y_0 = y] - \int h(z)\mathcal{L}(dz)|}{V(y)} : |h| \leq V \right\} = 0. \quad (2.13)$$

Definition II.3. A Markov chain $\{Y_n, n \geq 0\}$ on a state space \mathbf{R}^q is said to be Harris recurrent if there exists a recurrent set $\mathcal{R} \in \mathcal{B}(\mathbf{R}^q)$, a probability measure φ on \mathcal{R} , a $\lambda > 0$, and an integer n_0 such that

$$P\{Y_n \in \mathcal{R} \text{ for some } n \geq 1 | Y_0 = y\} = 1, \quad (2.14)$$

$$P\{Y_{n_0} \in A | Y_0 = y\} \geq \lambda\varphi(A),$$

for all $y \in \mathcal{R}$ and $A \subset \mathcal{R}$.

It is known that under the irreducibility and aperiodicity assumption, V -uniform ergodicity implies that $\{X_n, n \geq 0\}$ is Harris recurrent, cf. Theorem 9.18 of Meyn and Tweedie (2009).

The following assumptions will be used throughout this paper.

Condition C:

C1. The Markov chain $\{(X_n, Y_n), n \geq 0\}$ defined in (2.1) and (2.2) is aperiodic and irreducible on $\mathcal{X} \times \mathbf{R}^q$. For each $j \in \mathcal{X}$, the conditional Markov chain $\{Y_n | X_n, n \geq 0\}$ is $V_j(\cdot)$ -uniformly ergodic for some $V_j(\cdot)$ on \mathbf{R}^q , such that there exists $p \geq 1$,

$$\sup_{y \in \mathbf{R}^q} E_y^\theta \left\{ \frac{V_j(Y_p)}{V_j(y)} \right\} < \infty \text{ for all } j \in \mathcal{X}. \quad (2.15)$$

C2. Assume $0 < \sup_{j \in \mathcal{X}} f(y|j, y_0) < \infty$, for all $y \in \mathbf{R}^q$. Denote $h(Y_1) = \max_{i \in \mathcal{X}} \sup_{y_0 \in \mathbf{R}^q} \sum_{j=1}^d p_{ij} f(Y_1|j, y_0)$. Assume there exists $p \geq 1$ as in C1 such that for all $i \in \mathcal{X}$,

$$\sup_{j \in \mathcal{X}, y \in \mathbf{R}^q} E_i^\theta \left\{ \log \left(h(Y_1)^p \frac{V_j(Y_p)}{V_j(y)} \right) \right\} < 0, \quad (2.16)$$

$$\sup_{j \in \mathcal{X}, y \in \mathbf{R}^q} E_i^\theta \left\{ h(Y_1) \frac{V_j(Y_1)}{V_j(y)} \right\} < \infty. \quad (2.17)$$

C3. Recall that \mathcal{L} is a σ -finite measure on \mathbf{R}^q defined in (2.2). Assume

$$\max_{i \in \mathcal{X}} \sup_{y_0 \in \mathbf{R}^q} \left| \sum_{j \in \mathcal{X}} \int_{y \in \mathbf{R}^q} \pi_i p_{ij} f(y|j, y_0) \mathcal{L}(dy) \right| < \infty.$$

Remark II.4. *C1 is an ergodic condition for the underlying Markov chain. The weighted mean contraction property (2.16) and the finite weighted mean average property (2.17), which appear in C2, guarantee that the induced Markov chain $\{W_n, n \geq 0\}$ is \tilde{V} -uniformly ergodic for a given function \tilde{V} , and hence to be Harris recurrent. In Section IV, we show that several interesting models satisfy these conditions. C3 is a constraint of the Rényi divergence (Kullback–Leibler divergence) and is a standard moment condition. The finiteness condition is quite natural and holds in most cases.*

The following proposition is a generalization of Theorem 3 in Fuh (2021b). Since the proof is the same as those in Lemmas 3 and 4 of Fuh (2006), it is omitted.

Proposition II.5. *Let $\{(X_n, Y_n), n \geq 0\}$ be the hidden Markov model given in (2.1) and (2.2), satisfying C1–C3. Then the induced Markov chain $\{W_n, n \geq 0\}$ is an aperiodic, irreducible, and Harris recurrent Markov chain, with the invariant probability Π . Furthermore there exist $a, C > 0$, such that $\mathbb{E}_w(\exp\{ag(W_0, W_1)\}) \leq C < \infty$ for all $w \in \mathcal{W}$.*

Under the Harris recurrent condition (2.14), it is known that, cf. Meyn and Tweedie (2009), W_n admits a regenerative scheme with i.i.d. inter-regeneration times for an augmented Markov chain, which is called the “split chain”. Heuristic speaking, let $\tau_\Delta(0) = \tau_\Delta$, and let $\{\tau_\Delta(j), j \geq 1\}$ denote the times of consecutive visits to a recurrent state $\Delta \in \mathcal{S}$. For a function $f : \mathcal{S} \rightarrow \mathbf{R}$, let $S_n = \sum_{i=0}^n f(W_i)$, $S_j(f) = \sum_{i=\tau_\Delta(j)+1}^{\tau_\Delta(j+1)} f(W_i)$. By the strong Markov property, the random variables $\{S_j(f), j \geq 0\}$ are independent and identically distributed random variables. Note that here we only consider $f(W_i)$; the case of $f(W_i, W_{i+1})$ is similar.

Let $\tau = \tau_\Delta$ be the first time (> 0) reaches the recurrent state Δ of the split chain. Let ν be an initial distribution on \mathcal{S} , and define

$$u(\vartheta, \zeta) = E_\nu e^{\vartheta S_\tau - \zeta \tau} \text{ for } \zeta \in \mathbf{R}. \quad (2.18)$$

Assume that

$$\Gamma := \{(\vartheta, \zeta) : u(\vartheta, \zeta) < \infty\} \text{ is an open subset on } \mathbf{R}^2. \quad (2.19)$$

Denote $\zeta_1 := S_1$. Ney and Nummelin (1987) shows that $\mathcal{D} = \{\vartheta : u(\vartheta, \zeta) < \infty \text{ for some } \zeta\}$ is an open set and that for $\vartheta \in \mathcal{D}$, the transition kernel

$$\hat{\mathbf{P}}_\vartheta(w, A) = \mathbb{E}_w \{e^{\vartheta \zeta_1} I_{\{W_1 \in A\}}\} \quad (2.20)$$

has a unique maximal simple real eigenvalue $e^{\Lambda(\vartheta)}$, where $\Lambda(\vartheta)$ is the unique solution of the equation $u(\vartheta, \Lambda(\vartheta)) = 1$, with corresponding right eigenfunctions $r(\cdot; \vartheta)$ and left eigenmeasures $l_\nu(\cdot; \vartheta)$ defined by

$$r(w; \vartheta) := \mathbb{E}_w \exp\{\vartheta S_\tau - \tau \Lambda(\vartheta)\}. \quad (2.21)$$

For a measurable subset $A \in \mathcal{B}(\mathcal{S})$, any initial distribution ν on \mathcal{S} and $w \in \mathcal{S}$, define

$$l_\nu(A; \vartheta) = \mathbb{E}_\nu \left[\sum_{n=0}^{\tau-1} e^{\vartheta S_n - n \Lambda(\vartheta)} I_{\{W_n \in A\}} \right], \quad (2.22)$$

$$l_w(A; \vartheta) = \mathbb{E}_w \left[\sum_{n=0}^{\tau-1} e^{\vartheta S_n - n \Lambda(\vartheta)} I_{\{W_n \in A\}} \right]. \quad (2.23)$$

To analyze $\hat{\mathbf{P}}_\vartheta$ in (2.20), for completeness, we state the following proposition, which is taken from Theorem 4.1 in Ney and Nummelin (1987). Note that by Proposition II.5, the induced Markov chain $\{W_n, n \geq 0\}$ is an aperiodic, irreducible, and Harris recurrent Markov chain, which implies that condition M1 in Theorem 4.1 of Ney and Nummelin (1987) holds.

Proposition II.6. *Let $\{(X_n, Y_n), n \geq 0\}$ be the hidden Markov model given in (2.1) and (2.2), satisfying C1–C3. Let $\hat{\mathbf{P}}_\vartheta(\cdot, \cdot)$ be the operator defined in (2.20), and $\Lambda(\cdot)$ be defined by the characteristic equation (2.18). Then*

(i) $\mathcal{D} = \{\vartheta : u(\vartheta, \zeta) < \infty \text{ for some } \zeta\}$ is an open set. Λ is analytic, strictly convex, and essentially smooth on \mathcal{D} .

(ii) For $\vartheta \in \mathcal{D}$, $\lambda(\cdot) = e^{\Lambda(\cdot)}$ is the largest eigenvalue of $\hat{\mathbf{P}}_\vartheta$ with (right) eigenfunction $\{r(w; \vartheta) : w \in \mathcal{S}\}$ and (left) eigenmeasure $\{l_\nu(A; \vartheta) : A \in \mathcal{S}\}$ having the representation (2.21) and (2.22).

(iii) There is a set $B \subset \mathcal{S}$ with $\varphi(B^c) = 0$, such that for each $w \in B$, $0 < r(w; \cdot) < \infty$ and is analytic on \mathcal{D} . If B is a small set, then $0 < l_\nu(B; \vartheta) < \infty$ and $0 < l_w(B; \vartheta) < \infty$ for all $w \in \mathcal{S}$ and is analytic on \mathcal{D} .

(iv) There exists a partition $\mathcal{S} = \cup_{i=1}^\infty \mathcal{S}_i$ and a sequence of functions $f_i : \mathbf{R} \rightarrow (0, \infty), i = 1, 2, \dots$, such that

$$r(w; \vartheta) \geq f_i(\vartheta) I_{\mathcal{S}_i}(w), \quad w \in \mathcal{S}, \vartheta \in \mathcal{D}, i = 1, 2, \dots$$

Remark II.7. *We will use Proposition II.6 (i)–(iv) in Section III and the rest of this paper, the reader is referred to Ney and Nummelin (1987) Theorem 4.1 and Lemma 4.5 for details. (iv) states that there is a countable partition of the state space $\mathcal{S} = \cup_{i=1}^\infty \mathcal{S}_i$, independent of ϑ , such that $r(w; \vartheta)$ is uniformly*

positive on each \mathcal{S}_i . However in order to apply (iii), one needs to extend to $0 < l_\nu(\mathcal{S}; \vartheta) < \infty$ and the uniform boundness of $r(w; \cdot)$ over the whole space \mathcal{S} . To this end, one needs extra condition and apply Theorem 4 of Chan and Lai (2003) under this additional assumption. For completeness, we include it as follows.

Note that $\{W_n, n \geq 0\}$ is \tilde{V} -uniformly ergodic as stated in Remark II.4.

C4. Assume (2.19) hold. Let C be a measurable subset of \mathcal{S} such that for any given initial distribution ν on \mathcal{S} ,

$$\mathcal{L}_\nu(C; \vartheta) < \infty \text{ and } \mathcal{L}_w(C; \vartheta) < \infty \text{ for all } w \in \mathcal{S}. \quad (2.24)$$

Let $\tilde{V} : \mathcal{S} \rightarrow [1, \infty)$ be a measurable function. Assume for some $0 < \beta < 1$ and $K > 0$, we have

$$E_w[e^{\vartheta W_1 - \Lambda(\vartheta)} \tilde{V}(W_1)] \leq (1 - \beta) \tilde{V}(w) \quad \forall w \notin C, \quad (2.25)$$

$$\sup_{w \in C} E_w[e^{\vartheta W_1 - \Lambda(\vartheta)} \tilde{V}(W_1)] = K < \infty \text{ and } \int \tilde{V}(w) \varphi(dw) < \infty, \quad (2.26)$$

where φ is defined in (2.14).

Remark II.8. Note that under condition C1–C4, we have $0 < l_\nu(\mathcal{S}; \vartheta) < \infty$ and $r(w; \cdot)$ is uniform boundness over the whole space \mathcal{S} . Although condition C4 is under the induced Markov chain $\{W_n, n \geq 0\}$, by using the results in Fuh (2021a), this condition holds for some interesting examples, see Section IV. Moreover, if the state space is finite (compact), which is commonly used in engineering, the above results hold.

III. RÉNYI DIVERGENCE

We state our main results in this section. Section III-A presents the convergence of the Rényi divergence. Section III-B defines the Kullback–Leibler divergence, and shows that the Kullback–Leibler divergence is the limit of the Rényi divergence as $\alpha \rightarrow 1$.

A. Rényi divergence

Let $\{Y_n, n \geq 0\}$ be the general HMM defined in (2.1). Denote $Y_{0:n} = \{Y_0, Y_1, \dots, Y_n\}$ and $y_{0:n} = \{y_0, y_1, \dots, y_n\}$. With the same notation used in Section II, denote $P^{(n)}(\cdot)$ and $Q^{(n)}(\cdot)$ as two proba-

bilities on $Y_{0:n}$. By (2.4), the probability density functions $p^{(n)}(\cdot)$ and $q^{(n)}(\cdot)$ of the random variables $\{Y_0, Y_1, \dots, Y_n\}$ under $P^{(n)}$ and $Q^{(n)}$ are given, respectively, by

$$\begin{aligned} p^{(n)}(y_{0:n}) &= p_n(y_0, y_1, \dots, y_n) \\ &= \sum_{x_0=1}^d \cdots \sum_{x_n=1}^d \nu_{x_0} f(y_0|x_0) \prod_{k=1}^n p_{x_{k-1}x_k} f(y_k; \theta|x_k, y_{k-1}), \end{aligned} \quad (3.1)$$

$$\begin{aligned} q^{(n)}(y_{0:n}) &= q_n(y_0, y_1, \dots, y_n) \\ &= \sum_{x_0=1}^d \cdots \sum_{x_n=1}^d \nu_{x_0} g(y_0|x_0) \prod_{k=1}^n p_{x_{k-1}x_k} g(y_k; \theta|x_k, y_{k-1}), \end{aligned} \quad (3.2)$$

where $g(y_k|x_k, y_{k-1})$ is the probability density of Q with respect to \mathcal{L} .

Recall the definition of the Rényi Divergence for independent and identically distributed random variables (i.i.d.) $\{\xi_n, n \geq 0\}$ as follows: for given $\alpha \in (0, 1) \cup (1, \infty)$, let

$$D_\alpha(f||g) = \frac{1}{\alpha - 1} \log E_f \left[\left(\frac{f(\xi_1)}{g(\xi_1)} \right)^{\alpha-1} \right]. \quad (3.3)$$

Now for given a hidden Markov model $\{Y_n, n \geq 0\}$ with transition probability density p and q , let

$$D_\alpha^n(p^{(n)}||q^{(n)}) = \frac{1}{\alpha - 1} \log E_p \left[\left(\frac{p_n(Y_0, Y_1, \dots, Y_n)}{q_n(Y_0, Y_1, \dots, Y_n)} \right)^{\alpha-1} \right]. \quad (3.4)$$

Note that here E_f (E_p) denotes the expectation under probability distribution f (p). We will use the same type of notation without specification here and afterward.

By (2.5) and (2.9), we have

$$(3.4) = \frac{1}{\alpha - 1} \log \mathbb{E}_p \left[\left(\frac{\|M_n^p \cdots M_1^p M_0^p \nu_p\|}{\|M_n^q \cdots M_1^q M_0^q \nu_q\|} \right)^{\alpha-1} \right], \quad (3.5)$$

where M_k^p is defined in (2.6) and (2.7) under the probability from $p^{(n)}$, and M_k^q is defined in (2.6) and (2.7) under the probability from $q^{(n)}$, for $k = 0, 1, \dots, n$. Here \mathbb{E} is defined as the expectation under the probability \mathbb{P} defined in (2.11).

Denote $g(W_0, W_0) = \log \frac{\|T_0^p \nu_p\|/\|\nu_p\|}{\|T_0^q \nu_q\|/\|\nu_q\|}$, and $g(W_{k-1}, W_k) = \log \frac{\|T_k^p \nu_p\|/\|T_{k-1}^p \nu_p\|}{\|T_k^q \nu_q\|/\|T_{k-1}^q \nu_q\|}$, for $k = 1, \dots, n$. Let $S_0 = g(W_0, W_0)$ and $S_n = S_0 + \sum_{k=1}^n g(W_{k-1}, W_k)$. Then by (2.12), we have

$$\begin{aligned}
& \mathbb{E}_p \left[\left(\frac{\|M_n^p \cdots M_1^p M_0^p \nu_p\|}{\|M_n^q \cdots M_1^q M_0^q \nu_q\|} \right)^{\alpha-1} \right] \\
&= \mathbb{E}_p \left[\exp \left\{ \log \left(\frac{\|M_n^p \cdots M_1^p M_0^p \nu_p\|}{\|M_n^q \cdots M_1^q M_0^q \nu_q\|} \right)^{\alpha-1} \right\} \right] \\
&= \mathbb{E}_p \left[\exp \left\{ (\alpha-1) \left(\log \frac{\|T_n^p \nu_p\|/\|T_{n-1}^p \nu_p\|}{\|T_n^q \nu_q\|/\|T_{n-1}^q \nu_q\|} + \cdots + \log \frac{\|T_1^p \nu_p\|/\|T_0^p \nu_p\|}{\|T_1^q \nu_q\|/\|T_0^q \nu_q\|} + \log \frac{\|T_0^p \nu_p\|/\|\nu_p\|}{\|T_0^q \nu_q\|/\|\nu_q\|} \right) \right\} \right] \\
&= \mathbb{E}_p \left[\exp \left\{ (\alpha-1) \left(g(W_{n-1}, W_n) + \cdots + g(W_0, W_1) + g(W_0, W_0) \right) \right\} \right] \\
&= \mathbb{E}_p \left[\exp \{ (\alpha-1) S_n \} \right].
\end{aligned} \tag{3.6}$$

Then, using $\hat{\mathbf{P}}_\alpha$ defined (2.20) in Section II-B with $\vartheta = \alpha - 1$, let ν_p (ν_q) be the initial distribution of the $\{W_n, n \geq 0\}$ under P (Q). Then we have

$$\mathbb{E}_\nu e^{(\alpha-1)S_\tau - \Lambda(\alpha)\tau} = u(\alpha, \Lambda(\alpha)) = 1 \tag{3.7}$$

$$\mathbb{E}_w e^{(\alpha-1)S_\tau - \Lambda(\alpha)\tau} = r(w, \alpha). \tag{3.8}$$

Let $\lambda(\alpha)$ be the largest eigenvalue of $\hat{\mathbf{P}}_\alpha$. Denote $r(w, \alpha)$ as the right eigenfunctions associated with $\lambda(\alpha)$ defined in (3.8). Define

$$\underline{r}(\alpha) = \inf_w r(w, \alpha), \quad \bar{r}(\alpha) = \sup_w r(w, \alpha), \tag{3.9}$$

Under conditions C1–C4, by Proposition II.6 (iv), the uniform positivity property, we have $0 < \underline{r}(\alpha) \leq \bar{r}(\alpha) < \infty$.

Theorem III.1. *Under conditions C1–C4, then the Rényi divergence rate between $p^{(n)}$ and $q^{(n)}$ is*

$$D_\alpha(p||q) := \lim_{n \rightarrow \infty} \frac{1}{n} D_\alpha^n(p^{(n)}||q^{(n)}) = \frac{1}{\alpha-1} \log \lambda(\alpha), \tag{3.10}$$

where $\lambda(\alpha)$ is the largest positive real eigenvalue of $\hat{\mathbf{P}}_\alpha$, and $0 < \alpha < 1$. Furthermore, the same result holds for $\alpha > 1$ if $P > 0$ and $Q > 0$.

Proof. Let $\lambda(\alpha)$ be the largest positive real eigenvalue of $\hat{\mathbf{P}}_\alpha$ defined in (2.20), with associated positive right eigenfunction $r(w, \alpha) > 0$ uniformly on \mathcal{S} . Then by (3.8), we have

$$\nu_\alpha \hat{\mathbf{P}}_\alpha^{n-1} r(w; \alpha) = \lambda^{n-1}(\alpha) r(w; \alpha). \tag{3.11}$$

Let $\underline{r}(\alpha)$ and $\bar{r}(\alpha)$ be defined in (3.9). Then $0 < \underline{r}(\alpha) \leq r(w, \alpha) \leq \bar{r}(\alpha) < \infty$, for all $w \in \mathcal{S}$. Let $\nu_\alpha \hat{\mathbf{P}}_\alpha^{n-1} \mathbf{1} = b(\alpha)$. Then by (3.9), we have

$$\lambda^{n-1}(\alpha) r(w; \alpha) = \nu_\alpha \hat{\mathbf{P}}_\alpha^{n-1} r(w; \alpha) \leq \bar{r}(\alpha) b(\alpha).$$

Similarly we have $\lambda^{n-1}(\alpha) r(w; \alpha) \geq \underline{r}(\alpha) b(\alpha)$.

Therefore,

$$\frac{r(w; \alpha)}{\bar{r}(\alpha)} \leq \frac{b(\alpha)}{\lambda^{n-1}(\alpha)} \leq \frac{r(w; \alpha)}{\underline{r}(\alpha)}.$$

Since

$$\frac{\int_w r(w; \alpha) \Pi(dw)}{\bar{r}(\alpha)} \leq \frac{b(\alpha)}{\lambda^{n-1}(\alpha)} \leq \frac{\int_w r(w; \alpha) \Pi(dw)}{\underline{r}(\alpha)},$$

we have

$$\frac{1}{n} \log \frac{\int_w r(w; \alpha) \Pi(dw)}{\bar{r}(\alpha)} \leq \frac{1}{n} \log \frac{b(\alpha)}{\lambda^{n-1}(\alpha)} \leq \frac{1}{n} \log \frac{\int_w r(w; \alpha) \Pi(dw)}{\underline{r}(\alpha)}. \quad (3.12)$$

Note that the constant terms in the upper- and lower-bound in (3.12) are bounded and independent of n , which approach 0 as $n \rightarrow \infty$. Therefore, we have

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log \frac{\nu_\alpha \hat{\mathbf{P}}_\alpha^{n-1} \mathbf{1}}{\lambda^{n-1}(\alpha)} = 0. \quad (3.13)$$

Hence

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log \nu_\alpha \hat{\mathbf{P}}_\alpha^{n-1} \mathbf{1} = \lim_{n \rightarrow \infty} \frac{1}{n} \log \lambda^{n-1}(\alpha) + \lim_{n \rightarrow \infty} \frac{1}{n} \log \frac{\nu_\alpha \hat{\mathbf{P}}_\alpha^{n-1} \mathbf{1}}{\lambda^{n-1}(\alpha)} = \log \lambda(\alpha). \quad (3.14)$$

Note that (3.14) holds for both p and q . Thus

$$\lim_{n \rightarrow \infty} \frac{1}{n} D_\alpha^n(p^{(n)} || q^{(n)}) = \frac{1}{\alpha - 1} \log \lambda(\alpha). \quad (3.15)$$

The proof is complete. \square

B. Kullback–Leibler divergence

By making use of Theorem III.1, we herein show that the Rényi divergence reduces to the Kullback–Leibler divergence as $\alpha \rightarrow 1$. Let us first note the following result about the computation of the Kullback–Leibler divergence rate between two general HMMs. The convergence rate of the Kullback–Leibler divergence has been investigated by Fuh (2004b), and Fuh and Mei (2015) for the parametric case. In

the following proposition, we then show that the Kullback–Leibler divergence for general HMMs can also be written in a form similar to that in the i.i.d. case.

Recall that $\{Y_0, Y_1, \dots\}$ is a general HMM. Let $p^{(n)}$ and $q^{(n)}$ be two probability distributions. Let P and Q be the probabilities associated with $p^{(n)}$ and $q^{(n)}$, respectively. Let ν_p and ν_q be two initial distributions with respect to $p^{(n)}$ and $q^{(n)}$, respectively. If $Q > 0$, then $q > 0$. Denote

$$D^n(p^{(n)}||q^{(n)}) = \log \left(\frac{p_n(Y_0, Y_1, \dots, Y_n)}{q_n(Y_0, Y_1, \dots, Y_n)} \right). \quad (3.16)$$

Proposition III.2. *Under conditions C1–C4, the Kullback–Leibler divergence rate between $p^{(n)}$ and $q^{(n)}$ is well-defined with*

$$K(p, q) = \lim_{n \rightarrow \infty} \frac{1}{n} D^n(p^{(n)}||q^{(n)}) = \mathbb{E}_{\Pi, p} [\log p_1(Y_0, Y_1)] - \mathbb{E}_{\Pi, q} [\log q_1(Y_0, Y_1)], \quad (3.17)$$

where $\mathbb{E}_{\Pi, p}$ ($\mathbb{E}_{\Pi, q}$) is the expectation of \mathbb{P}_p (\mathbb{P}_q) defined in (2.11) of Section 2.1 under the invariant probability Π of $\{W_n, n \geq 0\}$. Here \mathbb{P}_p denotes the probability under p .

Proof. Under conditions C1–C4, by Proposition II.6, the invariant probability Π of the induced Markov chain $\{W_n, n \geq 0\}$ exists. Recall $T_n = M_n \cdots M_1 M_0$ defined in (2.8). Now let M_n^p (M_n^q) be M_n defined in (2.6) and (2.7) when the probability is under P (Q). We can define T_n^p and T_n^q similarly. First, it is easy to see from (2.12) that

$$\begin{aligned} & \frac{1}{n} [\log p_n(Y_0, Y_1, \dots, Y_n) - \log q_n(Y_0, Y_1, \dots, Y_n)] = \frac{1}{n} [\log \|T_n^p\| - \log \|T_n^q\|] \\ & = \frac{1}{n} \sum_{i=1}^n g_p(W_i, W_{i-1}) - g_q(W_i, W_{i-1}). \end{aligned} \quad (3.18)$$

Taking $n \rightarrow \infty$ on both sides of (3.18), then by Proposition II.5 and the SLLN for Markov random walks in Meyn and Tweedie (2009), we have

$$\begin{aligned} K(p, q) &= \mathbb{E}_{\Pi, p} [g_p(W_1, W_0)] - \mathbb{E}_{\Pi, q} [g_q(W_1, W_0)] \\ &= \mathbb{E}_{\Pi, p} [\log p_1(Y_0, Y_1)] - \mathbb{E}_{\Pi, q} [\log q_1(Y_0, Y_1)], \end{aligned}$$

which completes the proof. \square

Theorem III.3. *Let $\alpha \in (0, 1) \cup (1, \infty)$. Assume conditions C1–C4 hold; then*

$$\begin{aligned} & \lim_{\alpha \rightarrow 1} \lim_{n \rightarrow \infty} \frac{1}{n} D_\alpha^n(p^{(n)}||q^{(n)}) = \lim_{n \rightarrow \infty} \lim_{\alpha \rightarrow 1} \frac{1}{n} D_\alpha^n(p^{(n)}||q^{(n)}) \\ & = \int_{w_0} \int_{w_1} \Pi_{w_0} \mathbb{P}(w_0, w_1) \log \frac{\mathbb{P}(w_0, w_1)}{\mathbb{Q}(w_0, w_1)} dw_1 dw_0 = K(p, q), \end{aligned} \quad (3.19)$$

which is the Kullback–Leibler divergence defined in (3.17).

Proof. To prove (3.19), we first consider the case of $\lim_{n \rightarrow \infty} \lim_{\alpha \rightarrow 1} \frac{1}{n} D_\alpha^n(p^{(n)} || q^{(n)})$. By (3.4), (3.5), and (3.6), we have

$$\begin{aligned} D_\alpha^n(p^{(n)} || q^{(n)}) &= \frac{1}{\alpha - 1} \log E_p \left[\left(\frac{p_n(Y_0, Y_1, \dots, Y_n)}{q_n(Y_0, Y_1, \dots, Y_n)} \right)^{\alpha-1} \right] \\ &= \frac{1}{\alpha - 1} \log \mathbb{E}_p \left[\left(\frac{\|M_n^p \cdots M_1^p M_0^p \nu_p\|}{\|M_n^q \cdots M_1^q M_0^q \nu_q\|} \right)^{\alpha-1} \right] \\ &= \frac{1}{\alpha - 1} \log \mathbb{E}_p \left[\exp \{ (\alpha - 1) S_n \} \right]. \end{aligned} \quad (3.20)$$

Then

$$\begin{aligned} &\lim_{n \rightarrow \infty} \lim_{\alpha \rightarrow 1} \frac{1}{n} D_\alpha^n(p^{(n)} || q^{(n)}) \\ &= \lim_{n \rightarrow \infty} \frac{1}{n} \lim_{\alpha \rightarrow 1} \frac{1}{\alpha - 1} \log \mathbb{E}_p \left[\exp \{ (\alpha - 1) S_n \} \right]. \\ &= \lim_{n \rightarrow \infty} \frac{1}{n} \mathbb{E}_p [S_n] = \int_{w_0} \int_{w_1} \Pi_{w_0} \mathbb{P}(w_0, w_1) \log \frac{\mathbb{P}(w_0, w_1)}{\mathbb{Q}(w_0, w_1)} dw_1 dw_0. \end{aligned} \quad (3.21)$$

Note that the second identity comes from L'Hospital's Rule and the last identity in (3.21) comes from (3.17) in Proposition III.2.

Next, we consider the case of $\lim_{\alpha \rightarrow 1} \lim_{n \rightarrow \infty} \frac{1}{n} D_\alpha^n(p^{(n)} || q^{(n)})$.

By (3.10) in Theorem III.1, we have

$$\lim_{n \rightarrow \infty} \frac{1}{n} D_\alpha^n(p^{(n)} || q^{(n)}) = \frac{1}{\alpha - 1} \log \lambda(\alpha).$$

To evaluate $\lim_{\alpha \rightarrow 1} \frac{1}{\alpha - 1} \log \lambda(\alpha)$, note by Proposition II.6 (i) that the eigenvalue $\lambda(\alpha)$ of $\hat{\mathbf{P}}_\alpha$ is a continuous differentiable function of α . Note that since $Q > 0$, we have

$$\lim_{\alpha \rightarrow 1} \lambda(\alpha) = 1.$$

Let a denote an arbitrary base of the logarithm. Then, by L'Hopital's rule, we find that

$$\lim_{\alpha \rightarrow 1} \frac{\log \lambda(\alpha)}{\alpha - 1} = \frac{1}{\ln a} \lambda'(1) := \frac{1}{\ln a} \frac{\partial \lambda(\alpha)}{\partial \alpha} \Big|_{\alpha=1}, \quad (3.22)$$

which is well defined by Proposition II.6 since the algebraic multiplicity of $\lambda(\alpha)$ is 1 by (2.20). The equation defining the largest positive eigenvalue $\lambda(\alpha) = 1$ of $\hat{\mathbf{P}}(w, A) := \mathbb{E}_w \{ I_{W_1 \in A} \} = \mathbb{P}_w(A)$. By Proposition II.6 (i), $\hat{\mathbf{P}}_\alpha$ is analytic for $\alpha \in \mathcal{D}$; therefore by (2.20), it is straightforward to check that $\hat{\mathbf{P}}_\alpha \rightarrow \hat{\mathbf{P}}$ as $\alpha \rightarrow 1$.

Note that for $\zeta \in \mathbf{R}$, $u(\alpha, \zeta) = \mathbb{E}_{\Pi} e^{(\alpha-1)S_{\tau} - \zeta\tau}$ defined in (2.18). Then the transition kernel $\hat{\mathbf{P}}_{\alpha}(w, A) = \mathbb{E}_w \{e^{(\alpha-1)\zeta_1} I_{\{W_1 \in A\}}\}$ has a maximal simple real eigenvalue $\lambda(\alpha) = e^{\Lambda(\alpha)}$, where $\Lambda(\alpha)$ is the unique solution of the equation $u(\alpha, \Lambda(\alpha)) = 1$. Then using $\Lambda(1) = \log \lambda(1) = \log 1 = 0$, we have

$$\begin{aligned} u(\alpha, \Lambda(\alpha)) = 1 &\implies \frac{\partial u(\alpha, \Lambda(\alpha))}{\partial \alpha} \Big|_{\alpha=1} = 0 \implies \mathbb{E}_{\Pi} \{(S_{\tau} - \Lambda'(\alpha)\tau) e^{(\alpha-1)S_{\tau} - \Lambda(\alpha)\tau}\} \Big|_{\alpha=1} = 0 \\ &\implies \mathbb{E}_{\Pi} \{(S_{\tau} - \Lambda'(\alpha)\tau) e^{(\alpha-1)S_{\tau} - \Lambda(\alpha)\tau}\} \Big|_{\alpha=1} = 0 \implies \mathbb{E}_{\Pi} \{(S_{\tau} - \Lambda'(1)\tau)\} = 0 \\ &\implies \lambda'(1) = \frac{\partial \lambda(\alpha)}{\partial \alpha} \Big|_{\alpha=1} = \frac{\mathbb{E}_{\Pi} S_{\tau}}{\mathbb{E}_{\Pi} \tau} = \mathbb{E}_{\Pi} S_1. \end{aligned} \quad (3.23)$$

The last identity in (3.23) comes from Lemma 5.2 of Ney and Nummelin (1987).

By using (3.22) and (3.23), we obtain

$$\lim_{\alpha \rightarrow 1} \frac{1}{\alpha - 1} \log \lambda(\alpha) = \int_{w_0} \int_{w_1} \Pi_{w_0} \mathbb{P}(w_0, w_1) \log \frac{\mathbb{P}(w_0, w_1)}{\mathbb{Q}(w_0, w_1)} dw_1 dw_0 = K(p, q), \quad (3.24)$$

which completes the proof. \square

IV. EXAMPLES

We present two examples of general HMMs in this section. Section IV-A considers the Markov switching models, whereas Section IV-B studies the RNN.

A. Markov switching models

We start with a simple real valued q -order autoregression around one of d constants μ_1, \dots, μ_d :

$$Y_n - \mu_{X_n} = \sum_{k=1}^q \psi_k (Y_{n-k} - \mu_{X_{n-k}}) + \varepsilon_n, \quad (4.1)$$

where $\varepsilon_n \sim N(0, \sigma^2)$, $|\psi_k| < 1$ for $k = 1, \dots, q$, and $\{X_n, n \geq 0\}$ is a d -state ergodic Markov chain. When $q = 4$ and $d = 2$, this model was studied by Hamilton (1989) in order to analyze the behavior of the U.S. real GNP. Note that the Markov switching model (4.1) includes the classical HMM by letting $\psi_k = 0$ for $k = 1, \dots, q$. To apply our theory in the form of (4.1), we consider a simple case of order 1 in (4.1) with $d = 2$. The extension to the general case is straightforward. In this case, the conditional probability given $X_n = x_n$ and $Y_{n-1} = y_{n-1}$, $n \geq 1$, is

$$f(y_n | x_n, y_{n-1}; \theta) = \frac{1}{\sqrt{2\pi}\sigma} \exp \left(- \frac{[(y_n - \mu_{x_n}) - \psi_1 (y_{n-1} - \mu_{x_{n-1}})]^2}{2\sigma^2} \right). \quad (4.2)$$

Denote $[p_{ij}]_{i,j=1,2}$ as the transition probability of the underlying Markov chain $\{X_n, n \geq 0\}$ and let $\theta = (p_{11}, p_{21}, \psi_1, \mu_1, \mu_2, \sigma^2)$ be the given parameter. Assume that $|\psi_1| < 1$ for the stability property, and

that there exists a constant $c > 0$ such that $\sigma^2 > c$. Moreover, we assume that $\mu_1 \neq \mu_2$. Since the state space of X_n is finite, we consider $0 < p_{ij} < 1$ for all $i, j = 1, 2$, and for $j = 1, 2$ let $V_j(y) = |y| + 1$ (cf. page 394 of Meyn and Tweedie (2009)) such that the condition C1 holds. Under the normal distribution assumption, it is easy to see that (2.15) in conditions C1 and C3 holds.

Next we check that the mean contraction property (2.16) in C2 for a simple Markov switching model with general innovation holds. Given $p \geq 1$ as in C2, and $|\psi| < 1$, let X_n be a two-state Markov chain, and $Y_n = \mu_{X_n} + \psi Y_{n-1} + \varepsilon_n$, where ε_n are i.i.d. random variables with $E|\varepsilon_1| = a < \infty$. Further, we assume both ε_1 have a positive probability density function with respect to the Lebesgue measure. Denote $h(Y_1) = C < 1$, $b = (1 - |\psi|^p)/(1 - |\psi|)$ and choose p such that $C^p(ab + 1) < 1$. Let $d(u, v) = |u - v|$. Then we have

$$\begin{aligned} & \sup_y \left\{ E_j \left(\log \frac{h(Y_1)^p V_j(Y_p)}{V_j(y)} \middle| Y_0 = y \right) \right\} \\ & < \sup_y \left\{ E_j \left(\log \frac{C^p(|\psi^p y + \sum_{k=0}^{p-1} \psi^k \varepsilon_{p-k}| + 1)}{|y| + 1} \middle| Y_0 = y \right) \right\} \\ & < \log \sup_y \left\{ \frac{C^p(|\psi^p y| + E|\sum_{k=0}^{p-1} \psi^k \varepsilon_{p-k}| + 1)}{|y| + 1} \right\} = \log \sup_y \left\{ \frac{C^p(|\psi^p y| + ab + 1)}{|y| + 1} \right\} < 0. \end{aligned} \quad (4.3)$$

By using the same argument, it is straightforward to check that (2.17) in C2 holds. By making use the same argument as that in Example 2 of Chan and Lai (2003), we can check C4 hold. In Section VI, we will present a numerical computation method of the Rényi divergence under model (4.1) with $q = 1$ and $d = 2$.

B. Recurrent neural network

Note that at the outset, RNN is a non-linear dynamical system commonly trained to fit sequence data via some variant of gradient descent. In this subsection, we treat RNN as a stochastic model as usual, cf. Goodfellow et al. (2016). Fuh (2021a) considers the example of RNN from a Markovian-iterated function system point of view, to study its stability. Here we apply a general HMM point of view to investigate RNN. Although some notation overlaps between these two parts, we include it here for completeness. A connection between the classical finite state HMM and RNN is in Buys et al. (2018).

An RNN can take as input a variable-length sequence $y = (y_1, \dots, y_n)$ by recursively processing each symbol while maintaining its internal hidden state h . At each time step n , the RNN reads the symbol

$Y_n \in \mathbf{R}^q$ and updates its hidden state $h_n \in \mathbf{R}^p$ by

$$h_n = f_\theta(Y_n, h_{n-1}), \quad (4.4)$$

where f_θ is a deterministic non-linear transition function, and θ is the parameter of f_θ .

The transition function f_θ can be implemented with gated activation functions such as long short-term memory (LSTM) or the gated recurrent unit (GRU). The joint probability of the RNN model sequence can be written as a product of conditional probabilities such that

$$P(Y_1, \dots, Y_n) = \prod_{k=1}^n P(Y_k | Y_1, \dots, Y_{k-1}) = \prod_{k=1}^n g_\gamma(h_{k-1}), \quad (4.5)$$

where g_γ is a function that maps the RNN hidden state h_{t-1} to a probability distribution over possible outputs, and γ is the parameter of g_γ .

To analyze (4.4) and (4.5), we provide a Markov chain framework as follows. Specifically, let $\mathbf{H} = \{h_n, n \geq 0\}$ be a sequence of random variables on $(\mathbf{R}^p, \mathcal{B}(\mathbf{R}^p))$, and suppose that a random sequence $\{Y_n\}_{n=0}^\infty$ taking values in \mathbf{R}^q is adjoined to \mathbf{H} such that $\{Z_n := (h_{n-1}, h_n, Y_n), n \geq 0\}$ is a Markov chain on $\mathbf{R}^p \times \mathbf{R}^p \times \mathbf{R}^q$ satisfying

$$\begin{aligned} & P\{(h_{n-1}, h_n) \in A, Y_n \in B | h_0, h_1, \dots, h_{n-1}; Y_0, Y_1, \dots, Y_{n-1}\} \\ &= \int_{y \in B} P\{(h_{n-1}, h_n) \in A | h_0, h_1, \dots, h_{n-1}; Y_0, Y_1, \dots, Y_{n-1}, Y_n \in dy\} \\ & \quad \times P\{Y_n \in dy | h_0, h_1, \dots, h_{n-1}; Y_0, Y_1, \dots, Y_{n-1}\} Q(dy) \\ &= \int_{y \in B} P\{(h_{n-1}, h_n) \in A | (h_{n-2}, h_{n-1}); Y_n \in dy\} P\{Y_n \in dy | (h_{n-2}, h_{n-1}); Y_{n-1}\} Q(dy) \\ &= \int_{y \in B} I_{\{(h_{n-1}, h_n) \in A | (h_{n-2}, h_{n-1}); Y_n \in dy\}} P\{Y_n \in dy | (h_{n-2}, h_{n-1}); Y_{n-1}\} Q(dy) \end{aligned} \quad (4.6)$$

for $A \in \mathcal{B}(\mathbf{R}^p \times \mathbf{R}^p)$, $B \in \mathcal{B}(\mathbf{R}^q)$ and each $n = 1, 2, \dots$.

By considering X_n to be degenerate and $\{(h_{n-1}, h_n, Y_n), n \geq 0\}$ to be a general-state Markov chain, we have $Y_n = g((h_{n-1}, h_n, Y_n))$ as a general HMM. Next, we consider the simple case, cf. Chung et al. (2015), in which the generating distribution is conditioned on h_{n-1} such that

$$Y_n = \mu_{y,n} + \sigma_{y,n} \varepsilon_n, \quad (4.7)$$

where $\sigma_{y,n} > 0$ P-a.s., $\varepsilon_n \sim N(0, 1)$ is a sequence of i.i.d. random variables, and ε_n is independent of $\{Y_{n-k}, k \geq 1\}$ for all n . Here, we assume that $\mu_{y,n}$ and $\sigma_{y,n}^2$ are the parameters of the generating

distribution such that $(\mu_{y,n}, \sigma_{y,n}^2) \sim g_\gamma(h_{n-1})$, with g_γ any highly flexible function such as a neural network.

To illustrate the general HMM approach, we consider two examples: linear RNN and LSTM. To start with, we consider the linear RNN. Let Y_n be the output model in (4.7); the linear RNN updates its hidden state using the following recurrence equation:

$$h_n = f_\theta(Y_n, h_{n-1}) = \delta_0 + \delta_1 h_{n-1} + \delta_2 Y_n, \quad (4.8)$$

where $\theta = (\delta_0, \delta_1, \delta_2)$ with $\delta_0 > 0$, $\delta_1 > 0$, and $\delta_2 > 0$ constants.

For an explicit representation, we analyze the linear RNN (4.7) and (4.8) as follows: let $Z_n = (h_{n-1}, h_n, Y_n)$ be the Markov chain on $\mathcal{X} := (\mathbf{R} \times \mathbf{R} \times \mathbf{R})$. Denote $\eta_n = h_{n-1}^{-1} Y_n$ and let $\tau_n = (\delta_1 + \delta_2 \eta_n) \in \mathbf{R}$. Let A_n be a 3×3 matrix written as

$$A_n = \begin{bmatrix} 0 & 1 & 0 \\ 0 & \tau_n & 0 \\ 0 & \eta_n & 0 \end{bmatrix}. \quad (4.9)$$

Note that although $\{A_n, n \geq 0\}$ are random matrices driven by the Markov chain $\{Z_n, n \geq 0\}$, since the randomness of Y_n comes only from the i.i.d. random variables ε_n , and since Y_n is independent of \mathcal{F}_{n-1} , the σ -algebra generated by Y_1, \dots, Y_{n-1} , $\{A_n, n \geq 0\}$ are i.i.d. random matrices.

Let $\xi_n = (0, \delta_0, 0)^t \in \mathbf{R}^3$. Then we have the following linear state space representation of the linear RNN (4.7) and (4.8): Z_n is a Markov chain governed by

$$Z_n = A_n Z_{n-1} + \xi_n, \quad (4.10)$$

and $Y_n := g(Z_n)$, the observed random quantity, is a non-invertible function of Z_n .

It is easy to check that the stability condition holds if $\mathbb{E}_\Pi \frac{Y_1}{h_1} < 1$ and $\delta_1 + \delta_2 \mathbb{E}_\Pi \frac{Y_1}{h_1} < 1$, where Π is the stationary distribution of the Markov chain $\{(Z_n, A_n \cdots A_1), n \geq 0\}$. The moment conditions hold under the normality assumption in (4.7). Note that Z_n defined in (4.10) is a V -uniformly ergodic Markov chain with $V(z) = \|z\|^2$, cf. Theorem 16.5.1 of Meyn and Tweedie (2009). By using the results in Bougerol and Picard (1992), it is straightforward to check that the stability condition and conditions C1 and C3 hold. By an argument similar to that in (4.3), C2 holds.

Next, we consider the LSTM network, cf. Hochreiter and Schmidhuber (1997). By using (4.7) as the output model for Y_n , we consider the hidden unit as follows. The state is a pair of vectors $s = (c, h) \in \mathbf{R}^{2d}$,

and the model is parameterized by eight matrices, $W_\Delta \in \mathbf{R}^{2d}$ and $U_\Delta \in \mathbf{R}^{d \times n}$, for $\Delta \in \{i, f, \sigma, z\}$. The state-transition map ϕ_{LSTM} for f_θ in (4.8) is defined as

$$\begin{aligned} f_t &= \sigma(W_f h_{t-1} + U_f y_t), & i_t &= \sigma(W_i h_{t-1} + U_i y_t), & o_t &= \sigma(W_o h_{t-1} + U_o y_t), \\ z_t &= \tanh(W_z h_{t-1} + U_z y_t), & c_t &= i_t \circ z_t + f_t \circ c_{t-1}, & h_t &= o_t \cdot \tanh(c_t), \end{aligned} \quad (4.11)$$

where \circ denotes elementwise multiplication, and σ is the logistic function.

Let $Z_n = (h_{n-1}, h_n, Y_n)$ be the Markov chain defined in (4.7) and (4.8). To provide conditions under which the r -step iterated system $\phi_{LSTM}^r = \phi_{LSTM} \circ \cdots \circ \phi_{LSTM}$ is stable, we denote $\|W\|_\infty$ as the induced ℓ_∞ matrix norm, which corresponds to the maximum absolute row sum $\max_i \sum_j |W_{ij}|$, and let $E\|f\|_\infty = \sup_t E\|f_t\|_\infty$. Since $\sigma < 1$ for given any weights W_f ; U_f and inputs y_t , we have $E\|f\|_\infty < 1$. This means the next state c_t must “forget” a non-trivial portion of c_{t-1} . We leverage this phenomenon to give sufficient conditions for ϕ_{LSTM} to be contractive in the ℓ_∞ norm, which in turn implies the system ϕ_{LSTM}^r is contractive in the ℓ_2 norm for $r = O(\log d)$.

By using the mean contraction under the normal distribution defined in (4.7), the following result is taken from Proposition 1 of Fuh (2021a), in which he shows that the iterated function system ϕ_{LSTM}^r is stable; see also Proposition 2 in Miller and Hardt (2018) for deterministic LSTM.

Proposition IV.1. *$\|W_f\|_\infty < B_w < \infty$, $\|U_f\|_\infty < B_u < \infty$, $\|Y_t\|_\infty \leq B_Y$, P -a.s for some random variable B_Y with $EB_Y < \infty$. Moreover, assume $\|W_i\|_\infty < (1 - E\|f\|_\infty)$, $\|W_o\|_\infty < (1 - E\|f\|_\infty)$, $\|W_z\|_\infty < (1/4)(1 - E\|f\|_\infty)$, $\|W_f\|_\infty < (1 - E\|f\|_\infty)^2$, and $r = O(\log d)$; then the iterated function system ϕ_{LSTM}^r is stable.*

Under this assumption, the state space of the Markov chain $\{Z_n = (h_{n-1}, h_n, Y_n), n \geq 0\}$ defined in the LSTM model (4.11) is compact, and hence is Harris recurrent and satisfies the V -uniformly ergodic assumptions C1. Under the normality assumption in (4.7), it is easy to see that $E|\varepsilon_1|^p < \infty$ for any $p > 0$. Therefore the moment conditions of C1 and C3 hold. The contraction property C2 holds due to the definition of the activation functions. C4 holds as the state space of the Markov chain is compact. By Theorem III.1 and Proposition III.2, we prove the existence of the Rényi divergence and Kullback–Leibler divergence, and provide a characterization. As numerical computations seem difficult, we will rely on Monte-Carlo simulations.

V. COMPUTATIONAL ISSUES IN GENERAL HMM

Since the Rényi divergence in general HMM involves an eigenvalue which is difficult to compute, we provide an approximated Rényi divergence in Section V-A. Next we present a theoretical background of the invariant measure for J_p^α in Section V-B, and report numerical computation of the Rényi divergence in Section V-C.

A. Approximated Rényi Divergence

Under conditions C1–C4, by Theorem III.1 the Rényi divergence between P and Q is

$$D_\alpha(p||q) := \lim_{n \rightarrow \infty} \frac{1}{n} D_\alpha^n(p^{(n)}||q^{(n)}) = \frac{1}{\alpha - 1} \log \lambda(\alpha). \quad (5.1)$$

Note that the computation of the Rényi divergence based on (5.1) involves the computation of the largest eigenvalue $\lambda(\alpha)$, which is not an easy task. Hence, instead of using (5.1), we will provide an alternative approach based on the recursive formula as follows. Recall that from (3.4), $D_\alpha(p||q)$ is defined as

$$D_\alpha(p||q) := \lim_{n \rightarrow \infty} \frac{1}{n} D_\alpha^n(p^{(n)}||q^{(n)}) = \frac{1}{\alpha - 1} \lim_{n \rightarrow \infty} \frac{1}{n} \log \mathbb{E}_p \left[\left(\frac{p_n(Y_0, Y_1, \dots, Y_n)}{q_n(Y_0, Y_1, \dots, Y_n)} \right)^{\alpha-1} \right], \quad (5.2)$$

where \mathbb{E}_p denotes the expectation according to \mathbb{P}_p , the probability of the Markov chain $\{W_n, \geq 0\}$ when the probability of $\{Y_0, Y_1, \dots, Y_n\}$ is under p_n .

Now, we seek to show that (5.2) can be computed via iterations. In particular, we have

$$p_n(y_0, y_1, \dots, y_n) = C_{n,1}(p) + \dots + C_{n,d}(p), \quad (5.3)$$

where

$$C_{t,j}(p) = f(y_t|x_{t-1} = j, y_{t-1}) \sum_{s=1}^d p_{sj} C_{t-1,s}(p), \quad (5.4)$$

with initial values $C_{1,j}(p) = \pi_j(p) f(y_1|x_0 = j)$, for $j = 1, \dots, d$. The following lemma gives an iteration method to compute the probability.

Lemma V.1. *Under conditions C1–C4, the probability can be calculated by*

$$(p_n(y_0, y_1, \dots, y_n))^{\alpha-1} = \prod_{t=1}^n (C_{t,1}^*(p) + \dots + C_{t,d}^*(p))^{\alpha-1}, \quad (5.5)$$

where

$$C_{t,j}^*(p) = \frac{f(y_t|x_{t-1} = j, y_{t-1}) \sum_{s=1}^d p_{sj} C_{t-1,s}(p)}{\sum_{s=1}^d C_{t-1,s}(p)},$$

with initial values $C_{1,j}^*(p) = \pi_j(p)f(y_1|x_0 = j)$, for $j = 1, \dots, d$.

Proof. By (5.3) and (5.4), and using induction for $n = 1, 2, \dots$, it is easy to show that

$$\begin{aligned}
& \prod_{t=1}^n (C_{t,1}^*(p) + \dots + C_{t,d}^*(p))^{\alpha-1} \\
&= \exp \left\{ (\alpha - 1) \sum_{t=1}^n \log(C_{t,1}^*(p) + \dots + C_{t,d}^*(p)) \right\} \\
&= \exp \left\{ (\alpha - 1) \left[\sum_{t=1}^n \log \left(\sum_{j=1}^d C_{t,j}(p) \right) - \sum_{t=1}^n \log \left(\sum_{s=1}^d C_{t-1,s}(p) \right) \right] \right\} \\
&= \exp \left\{ (\alpha - 1) \log \left(\sum_{j=1}^d C_{n,j}(p) \right) \right\} = \left(\sum_{j=1}^d C_{n,j}(p) \right)^{\alpha-1} = (p_n(y_0, y_1, \dots, y_n))^{\alpha-1}.
\end{aligned}$$

□

Remark V.2. Note that $C_{t,j}^*(p)$ in (5.5) generally have finite means, and thus $C_{n,j}^*(p)$ increase or decrease linearly in n . Hence, Lemma V.1 provides an algorithm that is computationally feasible when the time step is large. In addition, it is easy to see from (5.5) that the “normalization” of (5.3) reflects the idea of using the projection space $P(\mathbf{R}^d)$ defined in (2.10) and (2.11).

Denote

$$J^\alpha = \mathbb{E}_\Pi \left[\mathbb{E}_\Pi \left(\frac{C_{t,0}^*(p) + C_{t,1}^*(p)}{C_{t,0}^*(q) + C_{t,1}^*(q)} \right)^{\alpha-1} \middle| X_{t-1}, Y_{t-2}, W_{t-1} \right]. \quad (5.6)$$

By (5.2), (5.5), and (5.6) we observe that the Rényi divergence can be approximated as

$$\begin{aligned}
D_\alpha(p||q) &= \frac{1}{\alpha - 1} \lim_{n \rightarrow \infty} \frac{1}{n} \log \mathbb{E}_p \left\{ \frac{\exp \left\{ (\alpha - 1) \sum_{t=0}^n \log \left(\sum_{s=1}^d C_{t,s}^*(p) \right) \right\}}{\exp \left\{ (\alpha - 1) \sum_{t=0}^n \log \left(\sum_{s=1}^d C_{t,s}^*(q) \right) \right\}} \right\}, \quad (5.7) \\
&= \frac{1}{\alpha - 1} \frac{1}{n} \log \lim_{n \rightarrow \infty} \mathbb{E}_p \left\{ \exp \left\{ \sum_{t=0}^n \log \left(\frac{\sum_{s=1}^d C_{t,s}^*(p)}{\sum_{s=1}^d C_{t,s}^*(q)} \right)^{\alpha-1} \right\} \right\} \\
&\approx \frac{1}{\alpha - 1} \frac{1}{n} \log (J^\alpha)^n = \frac{1}{\alpha - 1} \log J^\alpha.
\end{aligned}$$

Remark V.3. Note that here in (5.7), we apply the following approximation

$$\mathbb{E}_p \left\{ \exp \left\{ \frac{1}{n} \sum_{t=0}^n \log \left(\frac{\sum_{s=1}^d C_{t,s}^*(p)}{\sum_{s=1}^d C_{t,s}^*(q)} \right)^{\alpha-1} \right\} \right\} \approx J^\alpha. \quad (5.8)$$

In other words, we approximate the largest eigenvalue $\lambda(\alpha)$ via $\lambda^n(\alpha) \approx (J^\alpha)^n$, which can be explained as follows. By (2.18), we have for given $\tau = n$,

$$E_\nu \left[e^{(\alpha-1)S_n} | \tau = n \right] = E_\nu \left[e^{\Lambda(\alpha)^n} | \tau = n \right] = E_\nu \left[\lambda^n(\alpha) | \tau = n \right].$$

Let T_1^α be defined as (2.8) with the form in (5.2). Denote A_τ as the number of epochs by the regeneration time τ ; then use $E\tau_1 \cdot A_\tau \sim n$ to approximate

$$(E(T_1^\alpha))^n \approx E e^{(\alpha-1)S_n} \approx E[e^{(\alpha-1)\sum_{j=1}^{\tau_n} S_{\tau_j}}] \approx E\left[\prod_{j=1}^{\tau_n} e^{(\alpha-1)S_{\tau_j}}\right] \approx E[e^{(\alpha-1)S_{\tau_1}}]^{\tau_n} = \lambda^{E\tau_1 \cdot A_\tau}(\alpha) \approx \lambda^n(\alpha).$$

In summary, we have more accurate approximation when $\alpha \approx 1$ or in the ‘almost i.i.d.’ case. In other words, we approximate $\lambda(\alpha)$ by $E(T_1^\alpha)$ via the idea of approximating the transition probability by the invariant probability.

B. Theoretical Background of the Invariant Measure for J^α

Since the Rényi divergence in general HMM in Theorem III.1 involves the largest eigenvalue of the operator defined in the induced Markov chain $\{W_n, n \geq 0\}$, it is not easy to compute in general. One standard way to calculate the Rényi divergence in general HMM is via Monte Carlo simulation. Specifically, we first generate $\{Y_t\}_{t=1}^n$ from the model \mathbb{P}_p . Second, we compute

$$\frac{1}{n} \log \mathbb{E}_p \left[\left(\frac{p_n(Y_0, Y_1, \dots, Y_n)}{q_n(Y_0, Y_1, \dots, Y_n)} \right)^{\alpha-1} \right]$$

via (5.5). Then, the Rényi divergence can be estimated by repeating the above procedure several times and averaging its results. Needless to say, Monte Carlo is time-consuming especially when repeated calculations are needed. In this section, we propose a faster algorithm to compute the Rényi divergence in a two-state Markov switching model. By (5.7), the Rényi divergence can be computed as $D_\alpha(p||q) = \frac{1}{\alpha-1} \log J^\alpha$, where J^α is defined in (5.6). For the case $\mathcal{X} = \{0, 1\}$, J^α can be computed numerically. Before stating the method, we first note that the invariant measure of J^α depends only on $W_t = C_{t,0}^*(p)/(C_{t,0}^*(p) + C_{t,1}^*(p)) \in [0, 1]$. The other important fact is that W_t depends to X_t and Y_{t-1} due to the fact that $\{(X_t, Y_t, Y_{t-1}, W_t), t \geq 0\}$ is a Markov chain ($Y_{-1} := 0$). To find the stationary distribution of W_t , we define $m_j(\cdot, \cdot)$ as the stationary density function satisfying

$$\Pr(X_t = j, Y_{t-1} = u, W_t \leq x) = \int_0^x m_j(u, w) dw.$$

By extending the argument as in Fuh and Mei (2015), the following proposition characterizes $m_j(\cdot, \cdot)$ via Fredholm integral equations. Before that, we require the following notation.

For ease of presentation, we will use p_{θ_1} for probability under \mathbb{P}_p and p_{θ} for a probability under \mathbb{P} with a parameter θ . We will denote $\varphi_j(\theta)$ as the parameter of the underlying Markov chain $X_t = j$ with parameter θ .

Define $z(w, x)$ as

$$z(w, x) = \frac{x}{1-x} \cdot \frac{p_{01}(\theta)w + p_{11}(\theta)(1-w)}{p_{00}(\theta)w + p_{10}(\theta)(1-w)},$$

for $j = 0, 1$,

$$Q_j(u, z) = P_{\theta_1} \left(\frac{g(Y_t | \varphi_0(\theta), u)}{g(Y_t | \varphi_1(\theta), u)} \leq z \middle| X_t = j, Y_{t-1} = u \right).$$

By using an argument similar to Theorem 3 of Fuh and Mei (2015), we have

Proposition V.4. *Under conditions C1–C4, for all $0 < x < 1$,*

$$\begin{aligned} m_0(u, x) &= p_{00}(\theta_1) \int_0^1 \int_{-\infty}^{\infty} g(u | \varphi_0(\theta_1), v) \frac{\partial}{\partial x} Q_0(u, z(w, x)) m_0(v, w) dv dw \\ &\quad + p_{10}(\theta_1) \int_0^1 \int_{-\infty}^{\infty} g(u | \varphi_1(\theta_1), v) \frac{\partial}{\partial x} Q_0(u, z(w, x)) m_1(v, w) dv dw, \\ m_1(u, x) &= p_{01}(\theta_1) \int_0^1 \int_{-\infty}^{\infty} g(u | \varphi_0(\theta_1), v) \frac{\partial}{\partial x} Q_1(u, z(w, x)) m_0(v, w) dv dw \\ &\quad + p_{11}(\theta_1) \int_0^1 \int_{-\infty}^{\infty} g(u | \varphi_1(\theta_1), v) \frac{\partial}{\partial x} Q_1(u, z(w, x)) m_1(v, w) dv dw. \end{aligned} \quad (5.9)$$

Remark V.5. *The key observation is that for a Markov switching model with a finite number of d -hidden states, the invariant measure can essentially be defined by d -functions whose ranges are in the $(d-1)$ -dimensional space. This is computationally challenging for $d \geq 3$, and numerically computationally feasible for $d = 2$, as the corresponding two-dimensional real-valued functions can be characterized by a two-dimensional Fredholm integral equation. Note that the Fredholm integral equation is well studied in mathematics, and the two-dimensional case can be numerically solved by discretizing and then finding the eigenvector of a (large) square matrix with respect to the eigenvalue.*

To illustrate the usefulness of Proposition V.4, we consider the following Markov switching regression model, in which the mean depends on X_t . A more general case of the means μ_{X_t} and $\mu_{X_{t-1}}$ is in Section V-C for the numerical computation.

Example V.6. Let $\{X_n, n \geq 0\}$ be a two-state ergodic (aperiodic, irreducible, and positive recurrent) Markov chain with transition probability matrix $P_{\theta} = \begin{pmatrix} p_{00} & p_{01} \\ p_{10} & p_{11} \end{pmatrix}$. Denote

$$Y_t = \mu_{X_t} + \psi_{X_t} Y_{t-1} + \sigma_{X_t} \epsilon_t, \quad (5.10)$$

where $\epsilon_t \sim N(0, 1)$. Denote $\theta = (p_{00}, p_{11}, \mu_0, \mu_1, \psi_0, \psi_1, \sigma_0, \sigma_1)$. To compute $Q_j(u, z)$ in (5.10), note that

$$\frac{g(Y_t | \varphi_0(\theta), u)}{g(Y_t | \varphi_1(\theta), u)} = \frac{\sigma_1}{\sigma_0} \exp \left\{ \zeta \left[\left(Y_t + \frac{\eta}{\zeta} \right)^2 - \frac{\eta^2}{\zeta^2} + \frac{\nu}{\zeta} \right] \right\},$$

where

$$\zeta = \frac{1}{2\sigma_1^2} - \frac{1}{2\sigma_0^2}, \quad \eta = \frac{\mu_0 + \psi_0 u}{2\sigma_0^2} - \frac{\mu_1 + \psi_1 u}{2\sigma_1^2} \quad \text{and} \quad \nu = -\frac{(\mu_0 + \psi_0 u)^2}{2\sigma_0^2} + \frac{(\mu_1 + \psi_1 u)^2}{2\sigma_1^2}.$$

It follows that $Q_j(u, z)$ can be computed as

$$Q_j(u, z) = \Pr \left(\Xi \leq \frac{1}{\zeta} \log \left(\frac{\sigma_0}{\sigma_1} z \right) + \frac{\eta^2}{\zeta^2} - \frac{\nu}{\zeta} \right),$$

where $\Xi \sim \sigma_j^2 \chi_1^2((\mu_j + \psi_j u + \eta/\zeta)^2)$.

Once we can approximate the stationary density $m_j(\cdot, \cdot)$ in Proposition V.4, J^α can be computed as follows and the Rényi divergence can be estimated.

$$\begin{aligned} J^\alpha &= \mathbb{E}_\Pi \left[\mathbb{E}_\Pi \left(\exp \left(\log \left(\frac{C_{t,0}^*(\theta_1) + C_{t,1}^*(\theta_1)}{C_{t,0}^*(\theta) + C_{t,1}^*(\theta)} \right) \right)^{\alpha-1} \middle| X_{t-1}, Y_{t-2}, W_{t-1} \right) \right] \quad (5.11) \\ &= \mathbb{E}_\Pi \left[\mathbb{E}_\Pi \left(\frac{C_{t,0}^*(\theta_1) + C_{t,1}^*(\theta_1)}{C_{t,0}^*(\theta) + C_{t,1}^*(\theta)} \right)^{\alpha-1} \middle| X_{t-1}, Y_{t-2}, W_{t-1} \right] \\ &= \sum_{j=0}^1 \int_0^1 \int_{-\infty}^{\infty} \sum_{i=0}^1 \mathbb{P}_{inv}(X_t = j | X_{t-1} = i) \\ &\quad \cdot \mathbb{E}_\Pi \left(\frac{C_{t,0}^*(\theta_1) + C_{t,1}^*(\theta_1)}{C_{t,0}^*(\theta) + C_{t,1}^*(\theta)} \right)^{\alpha-1} \middle| X_t = j, X_{t-1} = i, Y_{t-2} = v, W_{t-1} = w) m_i(v, w) dv dw \\ &= \int_0^1 \int_{-\infty}^{\infty} \left[p_{00}(\theta_1) G_{00}^\alpha(v, w) + p_{01}(\theta_1) G_{01}^\alpha(v, w) \right] m_0(v, w) dv dw \\ &\quad + \int_0^1 \int_{-\infty}^{\infty} \left[p_{10}(\theta_1) G_{10}^\alpha(v, w) + p_{11}(\theta_1) G_{11}^\alpha(v, w) \right] m_1(v, w) dv dw, \end{aligned}$$

where

$$\begin{aligned}
G_{ij}^\alpha(v, w) &= E_\Pi \left(\frac{C_{t,0}^*(\boldsymbol{\theta}_1) + C_{t,1}^*(\boldsymbol{\theta}_1)}{C_{t,0}^*(\boldsymbol{\theta}) + C_{t,1}^*(\boldsymbol{\theta})} \right)^{\alpha-1} | X_t = j, X_{t-1} = i, Y_{t-2} = v, W_{t-1} = w \\
&= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \left(\frac{[p_{00}(\boldsymbol{\theta}_1)w + p_{10}(\boldsymbol{\theta}_1)(1-w)]g(y|\varphi_0(\boldsymbol{\theta}_1), u) + [p_{01}(\boldsymbol{\theta}_1)w + p_{11}(\boldsymbol{\theta}_1)(1-w)]g(y|\varphi_1(\boldsymbol{\theta}_1), u)}{[p_{00}(\boldsymbol{\theta})w + p_{10}(\boldsymbol{\theta})(1-w)]g(y|\varphi_0(\boldsymbol{\theta}), u) + [p_{01}(\boldsymbol{\theta})w + p_{11}(\boldsymbol{\theta})(1-w)]g(y|\varphi_1(\boldsymbol{\theta}), u)} \right)^{\alpha-1} \\
&\quad \cdot g(u|\varphi_i(\boldsymbol{\theta}_1), v)g(y|\varphi_j(\boldsymbol{\theta}_1), u) du dy.
\end{aligned} \tag{5.12}$$

Then

$$D_\alpha(p||q) = \frac{1}{\alpha-1} \log J^\alpha = \frac{1}{\alpha-1} \log J^\alpha. \tag{5.13}$$

Remark V.7. We can see that this method involves two parts. The first part is solving the eigenvalue to approximate m_j , and the second part is using double integration to approximate J^α . If we carefully design our algorithm, we need only a few seconds to calculate the Rényi divergence.

C. Numerical Computation of the Rényi Divergence

To illustrate our method, in this subsection, we consider the following Markov switching model:

$$Y_t = \psi_1 \mu_{X_t} + \psi_2 \mu_{X_{t-1}} + \phi Y_{t-1} + \epsilon_t, \tag{5.14}$$

where $\epsilon_t \sim N(0, \sigma^2)$ and $X_t \in \mathcal{X} = \{0, 1\}$ is a Markov chain with transition probability matrix $P_\theta = \begin{pmatrix} p_{00} & p_{01} \\ p_{10} & p_{11} \end{pmatrix}$. Denote $\boldsymbol{\theta} = (p_{01}, p_{10}, \boldsymbol{\mu}, \phi, \psi_1, \psi_2, \sigma)$. The derivation of the Markov switching model (5.14) for numerical study will be given in the Appendix.

We give a summary of our numerical approximation of the invariant measure Π in model (5.11) and (5.12). Even though there are several ways to solve the two-dimensional Fredholm integral equation, the algorithm for solving the above integral equation should be carefully designed because $\mathbf{0}$ is one of its solutions. To resolve this problem, we vectorize m_j and take grid points on $(0, 1) \times (-\infty, \infty)$. Note that the standard techniques for the general Fredholm integral equation can be used to estimate $m_0(\cdot, \cdot), m_1(\cdot, \cdot)$ numerically from equation (5.9) in Proposition V.4. The key step is to consider two discrete approximations: $\int_0^1 \int_{-\infty}^{\infty} g(u|\phi(\boldsymbol{\theta}), v) \frac{\partial}{\partial x} Q(u, z(w, x)) m(v, w) dv dw$ and $\frac{\partial}{\partial x} Q(u, z(w, x))$. To this end, we first approximate $\int_{-\infty}^{\infty}$ by \int_{-a}^a , and consider the case that $-a = v_0 < v_1 < \dots < v_{N-1} < v_N = a$, $0 = x_0 < x_1 < \dots <$

$x_{N-1} < x_N = 1$ and $0 = w_0 < w_1 < \dots < w_{N-1} < w_N = 1$. One choice is to simply set $v_i = \frac{2ai}{N}$ and $x_i = w_i = \frac{i}{N}$ for $i = 0, 1, \dots, N$. Assume $m(a, 0) = m(a, 1) = m(b, 0) = m(b, 1) = 0$. Then

$$\begin{aligned} & \int_0^1 \int_{-a}^a g(u|\phi(\boldsymbol{\theta}), v) \frac{\partial}{\partial x} Q(u, z(w, x)) m(v, w) dv dw \\ & \approx \left[\frac{1}{N} g(u|\phi(\boldsymbol{\theta}), v_1) \frac{\partial}{\partial x} Q(u, z(w_1, x)) \right] m(v_1, w_1) \\ & + \sum_{j=2}^{N-1} \frac{1}{2N} \left[g(u|\phi(\boldsymbol{\theta}), v_j) \frac{\partial}{\partial x} Q(u, z(w_j, x)) + g(u|\phi(\boldsymbol{\theta}), v_j) \frac{\partial}{\partial x} Q(u, z(w_{j+1}, x)) \right] m(v_j, w_j) \end{aligned}$$

and

$$\frac{\partial}{\partial x} Q(u, z(u, x)) \approx \frac{Q(u, z(u, x + \Delta)) - Q(u, z(u, x - \Delta))}{2\Delta},$$

for $x = x_1, \dots, x_i, \dots, x_{N-1}$, where $\Delta = 1/(2N)$.

Now we apply $(u, x) = (u_1, x_1), \dots, (u_{N-1}, x_{N-1})$ to equation (5.9) to obtain $2(N-1)$ equations, and use the above approximations to discretize the right-hand side of equation (5.9). Write these $2(N-1)$ equations in a matrix form to yield $\mathbf{w} = \mathbf{M}\mathbf{w}$, where $\mathbf{w} = (m_0(u_1, x_1), \dots, m_0(u_{N-1}, x_{N-1}), m_1(u_1, x_1), \dots, m_1(u_{N-1}, x_{N-1}))^t$ and \mathbf{M} is a $2(N-1) \times 2(N-1)$ matrix whose entries values depend on $Q_0(u_i, z(w_j, x_i \pm \Delta))$ and $Q_1(u_i, z(w_j, x_i \pm \Delta))$, both of which can be computed for given values of (u_i, x_i) and w_j . Since $Q_i(u, z(w, 1)) = 1$ and $Q_i(u, z(w, 0)) = 0$ for all w and $i = 0, 1$, a nice property of the matrix \mathbf{M} is that the sum of each column equals 1, which is the main reason why we use the invariant joint density notation. This property ensures that the matrix \mathbf{M} has an eigenvalue (the largest) equal to 1 and thus the corresponding eigenvector \mathbf{w} is an efficient approximation to $m_0(\cdot, \cdot)$ and $m_1(\cdot, \cdot)$.

With the discretized approximation $\hat{m}_0(\cdot, \cdot)$ and $\hat{m}_1(\cdot, \cdot)$, we can estimate J_θ by approximating the integration in (5.11), thereby yielding an alternative way to compute the Rényi divergence. Now we show this alternative way is valuable, as the only method so far in the literature to estimate the Rényi divergence in general HMM is the Monte Carlo simulation of $\frac{1}{n} \log S_n$ for large value of n . It is expected that our proposed alternative non-Monte-Carlo method will allow one to check the accuracy and correctness of both methods.

We focus on the following 8 cases listed in Table I, and Table II presents the Rényi divergence for various $\alpha = 0.5, 0.8, 0.99, 0.999, 1.001, 1.01, 1.5, 2$, and the Kullback–Leibler divergence ($\alpha \rightarrow 1$) for the above 8 cases with two methods: simulation and numerical approximation. For simulation, we use the sample size as 2000 and the replication number as 100; for numerical approximation, we use the discretized lattice number as 16 and the lower and upper integral bounds as 15.

TABLE I
PARAMETERS FOR THE EIGHT EXAMPLES.

(1)	$\theta = (0.41, 0.6, (1, 0), 0, 1, 0, 2)$	$\theta_1 = (0.41, 0.6, (2, 1), 0, 1, 0, 1.5)$
(2)	$\theta = (0.41, 0.59, (1, 0), 0, 1, 0, 2)$	$\theta_1 = (0.41, 0.59, (2, 1), 0, 1, 0, 1.6)$
(3)	$\theta = (0.4, 0.59, (1, 0), 0, 1, 0, 1)$	$\theta_1 = (0.4, 0.59, [2, 1], 0, 1, 0, 0.9)$
(4)	$\theta = (0.4, 0.599, (1, 0), 0, 1, 0, 1)$	$\theta_1 = (0.4, 0.599, (2, 1), 0, 1, 0, 0.9)$
(5)	$\theta = (0.59, 0.4, (1, 0), 0, 1, 0, 1)$	$\theta_1 = (0.59, 0.4, (2, 1), 0, 1, 0, 0.9)$
(6)	$\theta = (0.599, 0.4, (1, 0), 0.2, 1, 0, 1)$	$\theta_1 = (0.599, 0.4, (2, 1), 0.3, 1, 0, 1.1)$
(7)	$\theta = (0.4, 0.59, (1, 0), 0.2, 1, 0.2, 1.1)$	$\theta_1 = (0.4, 0.59, (2, 1), 0.1, 1, 0.1, 1)$
(8)	$\theta = (0.4, 0.59, (1, 1), 0, 1, 0, 1)$	$\theta_1 = (0.4, 0.59, (2, 2), 0, 1, 0, 0.9)$

Next, we also report the value of the Rényi divergence and Kullback–Leibler divergence in this context. We consider two ways to estimate the Rényi divergence and Kullback–Leibler divergence. One is based on Monte Carlo simulations with the time step for convenience. The other is based on the invariant probability measure based on 500 discretizations over the interval $[0,1]$. The corresponding results are summarized in Table II, in which we report Rényi divergence based on numerical and Monte Carlo methods. The number in the parentheses indicates the standard deviation of the Monte Carlo estimator. We also present relative errors between these two methods and their corresponding running time in seconds.

From Table II, these two ways yield similar numerical results, and thus the two different methods validate each other. In particular, we feel confident that the sample size and the replication number for the above examples are large enough in the Monte Carlo simulation to estimate the Rényi divergence and Kullback–Leibler divergence. Note that the Rényi divergence increases as α increases, as that in the i.i.d. case. Furthermore, the Rényi divergence and Kullback–Leibler divergence get closer when $\alpha \rightarrow 1$. We also observe that our numerical method is stable for the Kullback–Leibler divergence, whereas it is sensitive to the underlying parameter change for the Rényi divergence. This may be due to the use of the approximated Rényi divergence.

VI. CONCLUSION

In this paper, we study the Rényi divergence for a general HMM, to cover the Markov switching model and RNN, including the classical HMM as a special case. The Kullback–Leibler divergence can be regarded as the limit of $\alpha \rightarrow 1$ of the Rényi divergence. Moreover, we express the Kullback–Leibler

TABLE II
RESULTS FOR CASES WITH VARIOUS α

α		Cases								Average
		1	2	3	4	5	6	7	8	time (sec)
0.5	Numerical	0.1091	0.0921	0.2196	0.2211	0.2225	0.2723	0.1227	0.2818	11.60
	Simulation	0.1097	0.0927	0.2211	0.2220	0.2239	0.2733	0.1293	0.2826	239.45
	R.E. (%)	(0.0145)	(0.0133)	(0.0214)	(0.0214)	(0.0217)	(0.026)	(0.015)	(0.0247)	
0.8	Numerical	0.1533	0.1324	0.3372	0.3387	0.3366	0.4566	0.1939	0.4243	11.68
	Simulation	0.1538	0.1329	0.3382	0.3395	0.3374	0.4575	0.1979	0.4250	240.43
	R.E. (%)	(0.0114)	(0.0109)	(0.0191)	(0.0192)	(0.0189)	(0.0275)	(0.0131)	(0.0212)	
0.99	Numerical	0.1762	0.1541	0.4072	0.4087	0.4032	0.5850	0.2363	0.5062	12.39
	Simulation	0.1767	0.1546	0.4079	0.4094	0.4036	0.5857	0.2388	0.5068	238.79
	R.E. (%)	(0.0101)	(0.0099)	(0.0184)	(0.0185)	(0.018)	(0.0295)	(0.0124)	(0.0199)	
0.999	Numerical	0.1772	0.1550	0.4104	0.4120	0.4063	0.5913	0.2382	0.5099	12.31
	Simulation	0.1777	0.1555	0.4111	0.4127	0.4067	0.5921	0.2407	0.5105	239.49
	R.E. (%)	(0.0101)	(0.0099)	(0.0184)	(0.0184)	(0.018)	(0.0296)	(0.0123)	(0.0198)	
KL ($\alpha \rightarrow 1$)	Numerical	0.1773	0.1552	0.4108	0.4123	0.4066	0.5920	0.2386	0.5104	13.34
	Simulation	0.1780	0.1558	0.4114	0.4129	0.4070	0.5928	0.2407	0.5106	235.01
	R.E. (%)	(0.0101)	(0.0099)	(0.0184)	(0.0184)	(0.0179)	(0.0296)	(0.0123)	(0.0198)	
1.001	Numerical	0.1774	0.1553	0.4112	0.4127	0.4069	0.5927	0.2387	0.5108	12.63
	Simulation	0.1779	0.1557	0.4118	0.4134	0.4073	0.5935	0.2411	0.5113	238.03
	R.E. (%)	(0.0101)	(0.0099)	(0.0184)	(0.0184)	(0.0179)	(0.0296)	(0.0123)	(0.0198)	
1.01	Numerical	0.1784	0.1562	0.4144	0.4159	0.4100	0.5991	0.2406	0.5145	13.39
	Simulation	0.1789	0.1567	0.4150	0.4166	0.4104	0.5999	0.2430	0.5151	235.90
	R.E. (%)	(0.01)	(0.0098)	(0.0184)	(0.0184)	(0.0179)	(0.0298)	(0.0123)	(0.0198)	
1.5	Numerical	0.2248	0.2014	0.5807	0.5823	0.5650	0.9971	0.3418	0.6995	13.23
	Simulation	0.2253	0.2019	0.5806	0.5828	0.5645	0.9967	0.3411	0.7000	232.73
	R.E. (%)	(0.0081)	(0.0082)	(0.0178)	(0.0178)	(0.0171)	(0.0437)	(0.0113)	(0.0181)	
2	Numerical	0.2601	0.2370	0.7330	0.7345	0.7054	1.5699	0.4364	0.8587	12.49
	Simulation	0.2606	0.2374	0.7321	0.7348	0.7041	1.5548	0.4335	0.8590	232.84
	R.E. (%)	(0.0069)	(0.0071)	(0.0181)	(0.0181)	(0.0174)	(0.1445)	(0.0114)	(0.0176)	

divergence of the general HMM as a top Lyapunov exponent of a well-defined product of Markovian random matrices. Since the Rényi divergence involves the largest eigenvalue of the associated Markov operator, which is notoriously difficult to compute, we turn our attention to asymptotic expansions, and derive an approximated Rényi divergence, which can be used for numerical approximation based on the Fredholm integral equation.

There are further studies along this line. First, it would be interesting to approximate the largest eigenvalue $\lambda(\alpha)$ to yield a more accurate numerical approximation. Second, we will study the case in which $\{X_n, n \geq 0\}$ is a general state Markov chain; or a more general HMM to cover regime switching state space models and regime switching GARCH(p, q) (stochastic volatility) models. Last, applications of the Rényi divergence and Kullback–Leibler divergence in general HMMs such as model selection, regularization, and variational inference are also interesting and merit further investigation.

REFERENCES

- Arikan, E. (1996). An inequality on guessing and its application to sequential decoding. *IEEE Trans. Inform. Theory* 42, 99–105.
- Bassat, M. B. and J. Raviv (1978). Rényi’s entropy and the probability of error. *IEEE Trans. Inform. Theory* 24, 324–330.
- Blackwell, D. H. (1959). The entropy of functions of finite-state Markov chains. *Matematika* 3(5), 143–150.
- Blanding, W. R., P. K. Willett, Y. Bar-Shalom, and S. Coraluppi (2009). Multisensor track management for targets with fluctuating SNR. *IEEE Trans. Aerosp. Electron. Syst.* 45, 1275–1292.
- Blumer, A. C. and R. J. McEliece (1988). The Rényi redundancy of generalized Huffman codes. *IEEE Trans. Inform. Theory* 34, 1242–1249.
- Bougerol, P. and N. Picard (1992). Stationarity of GARCH processes and some nonnegative time series. *J. Econom.* 52, 115–127.
- Buys, J., Y. Bisk, and Y. Choi (2018). Bridging HMMs and RNNs through architectural transformations. In *Proceedings of the 32nd Conference on Neural Information Processing Systems*, pp. IRASL workshop, Montréal, Canada.
- Campbell, L. L. (1965). A coding theorem and Rényi’s entropy. *Inform. Contr.* 8, 423–429.

- Cappe, O., E. Moulines, and Ryden (2005). *Inference in Hidden Markov Models*. Springer-Verlag, New York.
- Chan, H. P. and T. L. Lai (2003). Saddlepoint approximations and nonlinear boundary crossing probabilities of Markov random walks. *Ann. Appl. Probab.* 13, 395–429.
- Chen, P. N. and F. Alajaji (2001). Csiszár’s cutoff rates for arbitrary discrete sources. *IEEE Trans. Inform. Theory* 47, 330–338.
- Cho, K., B. van Merriënboer, C. Gulcehre, D. Bahdanau, F. Bougares, H. Schwenk, and Y. Bengio (2014). Learning phrase representations using RNN encoder-decoder for statistical machine translation. In *Proceedings of the 2014 Conference on Empirical Methods in Natural Language Processing*.
- Chung, J., Kastner, K., L. Dinh, K. Goel, A. C. Courville, and Y. A. Bengio (2015). Recurrent latent variable model for sequential data. *Advances in Neural Information Processing Systems*, 2980–2988.
- Churchill, G. A. (1989). Stochastic models for heterogeneous DNA sequences. *Bull. Math. Biol.* 51, 79–94.
- Csiszar, I. (1995). Generalized cutoff rates and Rényi’s information measures. *IEEE Trans. Inform. Theory* 41, 26–34.
- Ephraim, Y. and N. Merhav (2002). Hidden Markov processes. *IEEE Trans. Inform. Theory* 48, 1518–1569.
- Erez, U. and R. Zamir (2001). Error exponents of modulo-additive noise channels with side information at the transmitter. *IEEE Trans. Inform. Theory* 47, 210–218.
- Fuh, C. D. (2004a). Asymptotic operating characteristics of an optimal change point detection in hidden Markov models. *Ann. Stat.* 32, 2305–2339.
- Fuh, C. D. (2004b). On Bahadur efficiency of the maximum likelihood estimator in hidden Markov models. *Stat. Sin.* 14, 127–144.
- Fuh, C. D. (2006). Efficient likelihood estimation in state space models. *Ann. Statist.* 34, 2026–2068. Corrigendum in 38, 1279–1285, (2010).
- Fuh, C. D. (2021a). Asymptotic behavior for Markovian iterated function systems. *Stoch. Process. Their Appl.* 138, 186–211.
- Fuh, C. D. (2021b). Asymptotically optimal change point detection for composite hypothesis in state space models. *IEEE Trans. Inform. Theory* 67, 485–505.
- Fuh, C. D. and Y. Mei (2015). Quickest change detection and Kullback-Leibler divergence for two-state hidden Markov models. *IEEE Trans. Signal Process.* 63, 4866–4878.

- Fuh, C. D. and A. G. Tartakovsky (2019). Asymptotic Bayesian theory of quickest change detection for hidden Markov models. *IEEE Trans. Inform. Theory* 65, 511–529.
- Goodfellow, I., Y. Bengio, and A. Courville (2016). *Deep Learning*. MIT Press.
- Hamilton, J. D. (1989). A new approach to the economic analysis of nonstationary time series and the business cycle. *Econometrica* 57, 357–384.
- Han, G. and B. Marcus (2006a). Analyticity of entropy rate in families of hidden Markov chains (ii). In *Proceedings of the 2006 IEEE International Symposium on Information Theory*, pp. 103–107. IEEE.
- Han, G. and B. Marcus (2006b). Analyticity of entropy rate of hidden Markov chains. *IEEE Trans. Inform. Theory* 52, 5251–5266.
- Hochreiter, S. and J. Schmidhuber (1997). Long short-term memory. *Neural Comput.* 9, 1735–1780.
- Hu, J., M. Brown, and W. Turin (1996). HMM based on-line handwriting recognition. *IEEE Trans. Pattern Anal. Mach. Intell.* 18, 1039–1045.
- Jacquet, P., G. Seroussi, and W. Szpankowski (2008). On the entropy of a hidden Markov process. *Theor. Comput. Sci.* 395, 203–219.
- Jelinek, F. (1968). Buffer overflow in variable length coding of fixed rate sources. *IEEE Trans. Inform. Theory* 14, 490–501.
- Jenssen, R., K. Hild, D. Erdogmus, J. C. Principe, and T. Eltoft (2003). Clustering using renyi’s entropy. In *Proceedings of the International Joint Conference on Neural Networks, 2003.*, Volume 1, pp. 523–528. IEEE.
- Khasminskii, R. and O. Zeitouni (1996). Asymptotic filtering for finite state Markov chains. *Stoch. Process. Their Appl.* 63, 1–10.
- Koopmans, L. H. (1960). Asymptotic rate of discrimination for Markov processes. *Ann. Math. Stat.* 31, 982–994.
- Kunda, A., Y. He, and P. Bahl (1989). Recognition of handwritten word: First and second order hidden Markov model based approach. *Pattern Recognit.* 18, 283–297.
- Metelli, A. M., M. Papini, F. Faccio, and M. Restelli (2018). Policy optimization via importance sampling. *Advances in Neural Information Processing Systems* 31, 5442–5454.
- Meyn, S. P. and R. L. Tweedie (2009). *Markov Chains and Stochastic Stability*. Cambridge University Press.
- Miller, J. and M. Hardt (2018). Stable recurrent models. In *Proceedings of International Conference on*

Learning Representations.

- Nemetz, T. (1974). On the α -divergence rate for Markov-dependent hypotheses. *Probl. Contr. Inform. Theory* 3, 147–155.
- Ney, P. and E. Nummelin (1987). Markov additive processes I. Eigenvalue properties and limit theorems. *Ann. Probab.* 15, 561–592.
- Ordentlich, E. and T. Weissman (2004). New bounds on the entropy rate of hidden Markov processes. In *Proceedings of Information Theory Workshop*, pp. 117–122. IEEE.
- Ordentlich, E. and T. Weissman (2006). On the optimality of symbol by symbol filtering and denoising. *IEEE Trans. Inform. Theory* 52, 19–40.
- Papini, M., A. M. Metelli, L. Lupo, and M. Restelli (2019). Optimistic policy optimization via multiple importance sampling. In *Proceedings of International Conference on Machine Learning*, pp. 4989–4999. PMLR.
- Prinzato, L., H. P. Wynn, and A. A. Zhigljavsky (1997). Using Rényi entropies to measure uncertainty in search problems. *Lectures Appl. Math.* 33, 253–268.
- Rabiner, L. R. (1989). A tutorial on hidden Markov models and selected applications in speech recognition. *Proceedings of the IEEE* 77(2), 257–286.
- Rabiner, L. R. and B. H. Juang (1993). *Fundamentals of Speech Recognition*. Prentice Hall, New Jersey.
- Rached, Z., F. Alajaji, and L. L. Campbell (2001). Rényi’s divergence and entropy rates for finite alphabet Markov sources. *IEEE Trans. Inform. Theory* 47, 1553–1561.
- Raghavan, V., A. Galstyan, and A. G. Tartakovsky (2013). Hidden Markov models for the activity profile of terrorist groups. *Ann. Appl. Stat.* 7, 2402–2430.
- Raghavan, V., G. Steeg, A. Galstyan, and A. G. Tartakovsky (2014). Modeling temporal activity patterns in dynamic social networks. *IEEE Trans. Computat. Social Syst.* 1, 89–107.
- Rényi, A. et al. (1961). On measures of entropy and information. In *Proceedings of the Fourth Berkeley Symposium on Mathematical Statistics and Probability, Volume 1: Contributions to the Theory of Statistics*. The Regents of the University of California.
- Tuğaç, S. and M. Efe (2010). Hidden Markov model based target detection. In *2010 13th international conference on information fusion*, pp. 1–7. IEEE.
- Vasuhi, S. and V. Vaidehi (2014). Target detection and tracking for video surveillance. *WSEAS Trans. Signal Process.* 10, 168–177.

Wu, C., E. L. Xu, and G. Han (2017). Rényi entropy rate of hidden Markov processes. In *2017 IEEE International Symposium on Information Theory (ISIT)*, pp. 2970–2974. IEEE.

Yamato, J., J. Ohya, and K. Ishii (1992). Recognizing human action in time-sequential images using hidden Markov model. In *Proceedings of the IEEE Computer Society Conference on Computer Vision and Pattern Recognition*, pp. 379–385.

Zucchini, W. and I. MacDonald (2009). *Hidden Markov Models for Time Series: An Introduction Using R*. Chapman & Hall, New York.

Zuk, O., I. Kanter, and E. Domany (2005). Asymptotics of the entropy rate for a hidden Markov process. In *Proceedings of the Data Compression Conference*, pp. 173–182. IEEE.

VII. APPENDIX

A. Theoretical Study of the Invariant Probability in Model (5.14)

We consider the Markov switching model in (5.14),

$$Y_t = \psi_1 \mu_{X_t} + \psi_2 \mu_{X_{t-1}} + \phi Y_{t-1} + \epsilon_t,$$

where $\{X_t, t \geq 0\}$ is a Markov chain on a state space $\mathcal{X} = \{0, 1\}$, with transition probability matrix $P_\theta = \begin{pmatrix} p_{00} & p_{01} \\ p_{10} & p_{11} \end{pmatrix}$. Here we omit θ in p_{ij} to simplify the notation. We will add θ_1 in p_{ij} as $p_{ij}^{\theta_1}$ when the probability is $P = P_{\theta_1}$.

We approach this problem by reformulating model (5.14) as a 1-order four-state Markov switch model.

To do so, we first create a variable Z_n such that

- $Z_t = 0$ if $(X_{t-1}, X_t) = (0, 0)$, $Z_t = 1$ if $(X_{t-1}, X_t) = (0, 1)$,
- $Z_t = 2$ if $(X_{t-1}, X_t) = (1, 0)$, $Z_t = 3$ if $(X_{t-1}, X_t) = (1, 1)$.

Then $\{Z_t, t \geq 0\}$ is a Markov chain on state space $\mathcal{X} = \{0, 1, 2, 3\}$, with transition probability matrix

$$P_\theta = \begin{pmatrix} p_{00} & p_{01} & 0 & 0 \\ 0 & 0 & p_{10} & p_{11} \\ p_{00} & p_{01} & 0 & 0 \\ 0 & 0 & p_{10} & p_{11} \end{pmatrix}.$$

Denote the conditional density function of Y_t given Y_{t-1} and $(X_{t-1}, X_t) = (i, j)$ as $f_{ij, \theta}(Y_t|Y_{t-1})$. Then, we can represent the joint probability of Y_1, \dots, Y_n as

$$p_{\theta}(Y_1, \dots, Y_n) = \|\mathbf{M}_n \dots \mathbf{M}_1 \boldsymbol{\pi}_{\theta}\|, \quad (6.1)$$

where

$$\boldsymbol{\pi}_{\theta} = \begin{pmatrix} p_{00}p_{10}/(p_{01} + p_{10}) \\ p_{01}p_{10}/(p_{01} + p_{10}) \\ p_{10}p_{01}/(p_{01} + p_{10}) \\ p_{11}p_{01}/(p_{01} + p_{10}) \end{pmatrix}, \quad \mathbf{M}_1 = \begin{pmatrix} f_{00, \theta}(Y_1) & 0 & 0 & 0 \\ 0 & f_{01, \theta}(Y_1) & 0 & 0 \\ 0 & 0 & f_{10, \theta}(Y_1) & 0 \\ 0 & 0 & 0 & f_{11, \theta}(Y_1) \end{pmatrix},$$

and for $t \geq 2$,

$$\mathbf{M}_t = \begin{pmatrix} p_{00}f_{00, \theta}(Y_t|Y_{t-1}) & 0 & p_{00}f_{00, \theta}(Y_t|Y_{t-1}) & 0 \\ p_{01}f_{01, \theta}(Y_t|Y_{t-1}) & 0 & p_{01}f_{01, \theta}(Y_t|Y_{t-1}) & 0 \\ 0 & p_{10}f_{10, \theta}(Y_t|Y_{t-1}) & 0 & p_{10}f_{10, \theta}(Y_t|Y_{t-1}) \\ 0 & p_{11}f_{11, \theta}(Y_t|Y_{t-1}) & 0 & p_{11}f_{11, \theta}(Y_t|Y_{t-1}) \end{pmatrix}.$$

Let

$$A_{t, \theta} = p_{00}(A_{t-1, \theta} + C_{t-1, \theta})f_{00, \theta}(Y_t|Y_{t-1}), \quad B_{t, \theta} = p_{01}(A_{t-1, \theta} + C_{t-1, \theta})f_{01, \theta}(Y_t|Y_{t-1}), \\ C_{t, \theta} = p_{10}(B_{t-1, \theta} + D_{t-1, \theta})f_{10, \theta}(Y_t|Y_{t-1}), \quad D_{t, \theta} = p_{11}(B_{t-1, \theta} + D_{t-1, \theta})f_{11, \theta}(Y_t|Y_{t-1}),$$

with initial value $(A_{1, \theta}, B_{1, \theta}, C_{1, \theta}, D_{1, \theta}) = \mathbf{M}_1 \boldsymbol{\pi}_{\theta}$. Then we can calculate $p_{\theta}(Y_1, \dots, Y_n)$ recursively by

$$p_{\theta}(Y_1, \dots, Y_n) = A_{n, \theta} + B_{n, \theta} + C_{n, \theta} + D_{n, \theta}.$$

The log-joint probability of Y_1, \dots, Y_n can be computed as

$$\log p_{\theta}(Y_1, \dots, Y_n) = \sum_{t=1}^n \log(A_{t, \theta}^* + B_{t, \theta}^* + C_{t, \theta}^* + D_{t, \theta}^*),$$

where

$$A_{t, \theta}^* = \frac{p_{00}(A_{t-1, \theta}^* + C_{t-1, \theta}^*)}{A_{t-1, \theta}^* + B_{t-1, \theta}^* + C_{t-1, \theta}^* + D_{t-1, \theta}^*} f_{00, \theta}(Y_t|Y_{t-1}), \\ B_{t, \theta}^* = \frac{p_{01}(A_{t-1, \theta}^* + C_{t-1, \theta}^*)}{A_{t-1, \theta}^* + B_{t-1, \theta}^* + C_{t-1, \theta}^* + D_{t-1, \theta}^*} f_{01, \theta}(Y_t|Y_{t-1}), \\ C_{t, \theta}^* = \frac{p_{10}(B_{t-1, \theta}^* + D_{t-1, \theta}^*)}{A_{t-1, \theta}^* + B_{t-1, \theta}^* + C_{t-1, \theta}^* + D_{t-1, \theta}^*} f_{10, \theta}(Y_t|Y_{t-1}), \\ D_{t, \theta}^* = \frac{p_{11}(B_{t-1, \theta}^* + D_{t-1, \theta}^*)}{A_{t-1, \theta}^* + B_{t-1, \theta}^* + C_{t-1, \theta}^* + D_{t-1, \theta}^*} f_{11, \theta}(Y_t|Y_{t-1}),$$

with initial value $(A_{1,\theta}^*, B_{1,\theta}^*, C_{1,\theta}^*, D_{1,\theta}^*) = \mathbf{M}_1 \boldsymbol{\pi}_\theta$.

Then the Kullback–Leibler divergence and Rényi divergence can be computed as follows. Here we compute only the Kullback–Leibler divergence; the computation of the Rényi divergence can be done as that in (5.12). To start with, the Kullback–Leibler divergence can be computed by $J_{\theta_1} - J_\theta$, where $J_\theta = \mathbb{E}_\Pi \left[\log(A_{t,\theta}^* + B_{t,\theta}^* + C_{t,\theta}^* + D_{t,\theta}^*) \right]$.

Define $W_t = (A_{t,\theta}^* + C_{t,\theta}^*) / (A_{t,\theta}^* + B_{t,\theta}^* + C_{t,\theta}^* + D_{t,\theta}^*)$, and the stationary density function satisfying

$$\mathbb{P}(X_{t-1} = j, X_t = k, Y_{t-1} = u, W_t \leq x) = \int_0^x m_{jk}(u, w) dw.$$

Then we can express J_θ as

$$\begin{aligned} J_\theta &= \mathbb{E}_\Pi \left[\log(A_{t,\theta}^* + B_{t,\theta}^* + C_{t,\theta}^* + D_{t,\theta}^*) \right] \\ &= \mathbb{E}_\Pi \left[\mathbb{E}_\Pi \left(\log(A_{t,\theta}^* + B_{t,\theta}^* + C_{t,\theta}^* + D_{t,\theta}^*) \middle| X_{t-2}, X_{t-1}, Y_{t-2}, W_{t-1} \right) \right] \\ &= \sum_{i=0}^1 \sum_{j=0}^1 \sum_{k=0}^1 \int_0^1 \int_{-\infty}^{\infty} \mathbb{P}_\Pi(X_t = k | X_{t-1} = j) \\ &\quad \cdot \mathbb{E}_\Pi \left(\log(A_{t,\theta}^* + B_{t,\theta}^* + C_{t,\theta}^* + D_{t,\theta}^*) \middle| X_{t-2} = i, X_{t-1} = j, X_t = k, Y_{t-2} = v, W_{t-1} = w \right) m_{ij}(v, w) dv dw \\ &= \int_0^1 \int_{-\infty}^{\infty} \left[p_{00}^{\theta_1} G_{000}(v, w) + p_{01}^{\theta_1} G_{001}(v, w) \right] m_{00}(v, w) dv dw \\ &\quad + \int_0^1 \int_{-\infty}^{\infty} \left[p_{10}^{\theta_1} G_{010}(v, w) + p_{11}^{\theta_1} G_{011}(v, w) \right] m_{01}(v, w) dv dw \\ &\quad + \int_0^1 \int_{-\infty}^{\infty} \left[p_{00}^{\theta_1} G_{100}(v, w) + p_{01}^{\theta_1} G_{101}(v, w) \right] m_{10}(v, w) dv dw \\ &\quad + \int_0^1 \int_{-\infty}^{\infty} \left[p_{10}^{\theta_1} G_{110}(v, w) + p_{11}^{\theta_1} G_{111}(v, w) \right] m_{11}(v, w) dv dw, \end{aligned}$$

where

$$\begin{aligned} G_{ijk}(v, w) &= \mathbb{E}_\Pi \left(\log(A_{t,\theta}^* + B_{t,\theta}^* + C_{t,\theta}^* + D_{t,\theta}^*) \middle| X_{t-2} = i, X_{t-1} = j, X_t = k, Y_{t-2} = v, W_{t-1} = w \right) \\ &= \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \log \left[p_{00} w f_{00,\theta}(y|u) + p_{01} w f_{01,\theta}(y|u) \right. \\ &\quad \left. + p_{10} (1-w) f_{10,\theta}(y|u) + p_{11} (1-w) f_{11,\theta}(y|u) \right] f_{ij,\theta_1}(u|v) f_{jk,\theta_1}(y|u) du dy. \end{aligned}$$

It remains to approximate $\hat{m}_{ij}(\cdot)$. Note that

$$\begin{aligned}
& \mathbb{P}_{\Pi}(X_{t-1} = j, X_t = k, Y_{t-1} = u, W_t \leq x) \\
&= \sum_{i=0}^1 \sum_{l=0}^1 \int_0^1 \int_{-\infty}^{\infty} \mathbb{P}_{\Pi}(X_{t-1} = j, X_t = k, Y_{t-1} = u, W_t \leq x | X_{t-2} = i, X_{t-1} = l, Y_{t-2} = v, W_{t-1} = w) \\
&\quad \cdot m_{il}(v, w) dv dw \\
&= \sum_{i=0}^1 \int_0^1 \int_{-\infty}^{\infty} \mathbb{P}_{\Pi}(X_{t-1} = j, X_t = k, Y_{t-1} = u, W_t \leq x | X_{t-2} = i, X_{t-1} = j, Y_{t-2} = v, W_{t-1} = w) \\
&\quad \cdot m_{ij}(v, w) dv dw \\
&= \sum_{i=0}^1 \int_0^1 \int_{-\infty}^{\infty} \mathbb{P}_{\Pi}(X_t = k | X_{t-1} = j) f_{ij, \theta_1}(u|v) \\
&\quad \cdot \mathbb{P}_{\Pi}(W_t \leq x | X_{t-2} = i, X_{t-1} = j, X_t = k, Y_{t-1} = u, W_{t-1} = w) m_{ij}(v, w) dv dw \\
&= \sum_{i=0}^1 \int_0^1 \int_{-\infty}^{\infty} \mathbb{P}_{\Pi}(X_t = k | X_{t-1} = j) f_{ij, \theta_1}(u|v) Q_{jk}(x; u, w) m_{ij}(v, w) dv dw,
\end{aligned}$$

where

$$\begin{aligned}
Q_{jk}(x; u, w) &= \mathbb{P}_{\Pi}(W_t \leq x | X_{t-2} = i, X_{t-1} = j, X_t = k, Y_{t-1} = u, W_{t-1} = w) \\
&= \mathbb{P}_{\Pi} \left(\frac{A_{t, \theta}^* + C_{t, \theta}^*}{A_{t, \theta}^* + B_{t, \theta}^* + C_{t, \theta}^* + D_{t, \theta}^*} \leq x \middle| X_{t-2} = i, X_{t-1} = j, X_t = k, Y_{t-1} = u, W_{t-1} = w \right) \\
&= \mathbb{P}_{\Pi} \left((1-x)p_{00}wf_{00, \theta}(Y_t|u) - xp_{01}wf_{01, \theta}(Y_t|u) + (1-x)p_{10}(1-w)f_{10, \theta}(Y_t|u) \right. \\
&\quad \left. - xp_{11}(1-w)f_{11, \theta}(Y_t|u) \leq 0 \middle| X_{t-1} = j, X_t = k, Y_{t-1} = u \right).
\end{aligned}$$

It follows that

$$\begin{aligned}
m_{00}(u, x) &= \int_0^1 \int_{-\infty}^{\infty} p_{00}^{\theta_1} f_{00, \theta_1}(u|v) \frac{\partial}{\partial x} Q_{00}(x; u, w) m_{00}(v, w) dv dw \\
&\quad + \int_0^1 \int_{-\infty}^{\infty} p_{00}^{\theta_1} f_{10, \theta_1}(u|v) \frac{\partial}{\partial x} Q_{00}(x; u, w) m_{10}(v, w) dv dw \\
m_{01}(u, x) &= \int_0^1 \int_{-\infty}^{\infty} p_{01}^{\theta_1} f_{00, \theta_1}(u|v) \frac{\partial}{\partial x} Q_{01}(x; u, w) m_{00}(v, w) dv dw \\
&\quad + \int_0^1 \int_{-\infty}^{\infty} p_{01}^{\theta_1} f_{10, \theta_1}(u|v) \frac{\partial}{\partial x} Q_{01}(x; u, w) m_{10}(v, w) dv dw \\
m_{10}(u, x) &= \int_0^1 \int_{-\infty}^{\infty} p_{10}^{\theta_1} f_{01, \theta_1}(u|v) \frac{\partial}{\partial x} Q_{10}(x; u, w) m_{01}(v, w) dv dw \\
&\quad + \int_0^1 \int_{-\infty}^{\infty} p_{10}^{\theta_1} f_{11, \theta_1}(u|v) \frac{\partial}{\partial x} Q_{10}(x; u, w) m_{11}(v, w) dv dw \\
m_{11}(u, x) &= \int_0^1 \int_{-\infty}^{\infty} p_{11}^{\theta_1} f_{01, \theta_1}(u|v) \frac{\partial}{\partial x} Q_{11}(x; u, w) m_{01}(v, w) dv dw \\
&\quad + \int_0^1 \int_{-\infty}^{\infty} p_{11}^{\theta_1} f_{11, \theta_1}(u|v) \frac{\partial}{\partial x} Q_{11}(x; u, w) m_{11}(v, w) dv dw.
\end{aligned}$$

Note that in the model we consider in (5.14), the error term ϵ_t is a normal distribution with probability density function

$$\begin{aligned}
f_{ij, \theta}(Y_t|u) &= \frac{1}{\sqrt{\sigma^2 2\pi}} \exp \left[-\frac{[Y_t - (\psi_2 \mu_i + \psi_1 \mu_j + \phi u)]^2}{2\sigma^2} \right] \\
&= \frac{1}{\sqrt{\sigma^2 2\pi}} e^{-Y_t^2/(2\sigma^2)} \exp \left[\frac{2Y_t(\psi_2 \mu_i + \psi_1 \mu_j + \phi u) - (\psi_2 \mu_i + \psi_1 \mu_j + \phi u)^2}{2\sigma^2} \right].
\end{aligned}$$

It follows that

$$\begin{aligned}
Q_{jk}(x; u, w) &= \mathbb{P}_{\Pi} \left((1-x)p_{00}w \exp \left[\frac{2Y_t(\psi_2 \mu_0 + \psi_1 \mu_0 + \phi u) - (\psi_2 \mu_0 + \psi_1 \mu_0 + \phi u)^2}{2\sigma^2} \right] \right. \\
&\quad - xp_{01}w \exp \left[\frac{2Y_t(\psi_2 \mu_0 + \psi_1 \mu_1 + \phi u) - (\psi_2 \mu_0 + \psi_1 \mu_1 + \phi u)^2}{2\sigma^2} \right] \\
&\quad + (1-x)p_{10}(1-w) \exp \left[\frac{2Y_t(\psi_2 \mu_1 + \psi_1 \mu_0 + \phi u) - (\psi_2 \mu_1 + \psi_1 \mu_0 + \phi u)^2}{2\sigma^2} \right] \\
&\quad \left. - xp_{11}(1-w) \exp \left[\frac{2Y_t(\psi_2 \mu_1 + \psi_1 \mu_1 + \phi u) - (\psi_2 \mu_1 + \psi_1 \mu_1 + \phi u)^2}{2\sigma^2} \right] \right) \\
&\leq 0 \left| X_{t-1} = j, X_t = k, Y_{t-1} = u \right).
\end{aligned}$$