THE ALMOST PERIODIC GAUGE TRANSFORM — AN ABSTRACT SCHEME WITH APPLICATIONS TO DIRAC OPERATORS

JEAN LAGACÉ, SERGEY MOROZOV, LEONID PARNOVSKI, BERNHARD PFIRSCH, AND ROMAN SHTERENBERG

ABSTRACT. One of the main tools used to understand both qualitative and quantitative spectral behaviour of periodic and almost periodic Schrödinger operators is the method of gauge transform. In this paper, we extend this method to an abstract setting, thus allowing for greater flexibility in its applications that include, among others, matrix-valued operators. In particular, we obtain asymptotic expansions for the density of states of certain almost periodic systems of elliptic operators, including systems of Dirac type. We also prove that a range of periodic systems including the two-dimensional Dirac operators satisfy the Bethe–Sommerfeld property, that the spectrum contains a semi-axis — or indeed two semi-axes in the case of operators that are not semi-bounded.

CONTENTS

1. Introduction	Z
1.1. A Gauge transform.	2
1.2. Description of the results for elliptic systems and the Dirac operator	5
1.3. Description of the main results and plan of the paper	7
Acknowledgements	9
Part I : An abstract gauge transform scheme	9
2. Generalised almost-periodic operators	9
2.1. Generalised Sobolev spaces	9
2.2. An algebra of operators	10
3. Elliptic and diagonal operators	16
4. The Density of States Measure and von Neumann Algebras	20
4.1. Representations of the operator algebra	20
4.2. The density of states measure	23
5. Gauge Transform	28
5.1. The commutator equation	30
5.2. Weak gauge transform	32

Date: October 24, 2021.

J. LAGACÉ, S. MOROZOV, L. PARNOVSKI, B. PFIRSCH, AND R. SHTERENBERG	
5.3. Parallel weak gauge transform	34
5.4. Strong gauge transform	36
6. Systems of Almost Periodic Operators	38
6.1. Symbol formalism for systems of almost periodic operators	38
6.2. Gauge transform in \mathbf{S}_m^{∞} : the reduction to uncoupled operators	40
PART II : APPLICATIONS TO ASYMPTOTIC PROPERTIES OF SYSTEMS	44
7. Besicovitch space and systems of operators	46
7.1. Concrete systems of operators	47
7.2. Conditions on the perturbation and its frequency set	48
8. Asymptotic expansions for the IDS	50
8.1. IDS for uncoupled operators	51
8.2. Reduction to a finite interval	52
8.3. Reduction to a quasiperiodic operator	54
8.4. Reduction to uncoupled operators	56
9. The structure of periodic operators	58
9.1. Description of periodic operators	58
9.2. The overlap function	60
10. Systems of periodic operators – The Bethe-Sommerfeld property	63
10.1. The Bethe-Sommerfeld property	63
11. Bethe–Sommerfeld for uncoupled operators	65
11.1. A description of the spectrum	67
11.2. Description of the resonant sets	69
11.3. Volumes of intersections	71
11.4. Estimates on the overlap function	72
12. The Dirac Operator	77
12.1. Clifford algebras	78
12.2. Dirac operators	79
References	84

1. INTRODUCTION

1.1. **A Gauge transform.** During the last fifteen years, substantial progress has been made in the spectral theory of periodic and almost periodic scalar operators. An important tool that was developed during this period and was used to obtain asymptotic spectral results was the method of gauge transform (see, e.g., [Sob05, Sob06, PS10, PS12, MPS14, PS16, Ivr19, PS19]). This

2

method, which heavily uses commutator estimates, was originally created for classical pseudodifferential operators (see e.g. [Wei77, Roz78]) but was then modified to the periodic case by Sobolev [Sob05, Sob06] and to the almost periodic setting by Parnovski and Shterenberg [PS12]. The aim of this paper is to describe the method of gauge transform on an abstract level and then apply this abstract scheme to a concrete example — elliptic systems of operators (including Dirac operators).

Here is the basic setting: suppose that we are given an operator

$$A = A_0 + B_2$$

where A_0 is a diagonal operator in a given basis and B is a perturbation, which is assumed to be small in some sense. The standard example which the reader may want to keep in mind is

$$A_0 = \operatorname{diag}(a_1(-\Delta)^{\alpha/2}, \dots, a_m(-\Delta)^{\alpha/2}),$$

where $\alpha > 0$, $0 \neq a_j \in \mathbb{R}$, and *B* is a pseudo-differential perturbation of order smaller than α with periodic or almost periodic coefficients. For instance, a Dirac operator with an almost periodic potential can be brought to such a form by a unitary transformation. In many applications we will furthermore require *A* to be self-adjoint, even though our general scheme may not always require it.

We want to find an operator A' that is unitarily equivalent to A and is simpler – either diagonal or, failing this, has a form

$$A' = UAU^{-1} = A'_0 + B', (1.1)$$

where A'_0 is diagonal, U is unitary, and B' is a perturbation that is smaller than B. The notion of 'smallness' assumes that we have a small parameter, and B' has this small parameter entering in a higher power than B. The most common example of application to PDEs assumes that the order of B' is smaller than the order of B (so the role of the small parameter is played by the inverse of the energy), but in some cases the small parameter can be chosen to be a coupling constant, see [PS19]. The operators A and A' have the same spectrum and the hope is that it is easier to describe the spectrum of A', both quantitatively and qualitatively. As an example of the spectral properties we want to study, we list the following two types of problems:

- (1) Obtaining asymptotic expansions for the so-called *integrated density of states* $N(A; \lambda)$ as the spectral parameter λ tends to $\pm \infty$;
- (2) If *B* has periodic coefficients, to prove that whenever *A* is unbounded above (resp. below), its spectrum contains a semi-axis $[\lambda_0, \infty)$ (resp. $(-\infty, \lambda_0]$). Such an operator *A* is said to satisfy the *Bethe–Sommerfeld property*.

If we seek the unitary operator *U* in (1.1) in the form $U = \exp(i\Psi)$, then we have

$$A' = A_0 + B + i[A_0, \Psi] + i[B, \Psi] - \frac{1}{2}[[A_0, \Psi], \Psi] - \frac{1}{2}[[B, \Psi], \Psi] + R,$$
(1.2)

where *R* consists of further terms given by formally expanding the series for the exponentials $\exp(i\Psi)$. Our hope is to solve the equation

$$B + i[A_0, \Psi] = 0 \tag{1.3}$$

for Ψ , so that the second and third terms of (1.2) cancel each other. Ideally, the rest of the terms (starting from the fourth one) would indeed be smaller than *B*. In most cases, however, these two wishes turn out to be infeasible.

The main obstacle is that solutions Ψ to equation (1.3) involve a denominator that could be small for some *B* (for example, to have any hope of solving (1.3), the diagonal part of *B* has to be absent). Therefore, we usually have to modify our procedure and divide the perturbation *B* into two parts – good (or non-resonant) part $B^{\mathcal{NR}}$ for which the equation

$$B^{\mathcal{N}\mathcal{R}} + i[A_0, \Psi] = 0 \tag{1.4}$$

has a nice solution $\Psi^{\mathcal{N}\mathcal{R}}$ and bad (or resonant) part $B^{\mathcal{R}} = B - B^{\mathcal{N}\mathcal{R}}$ which we will be unable to destroy using our procedure. Thus, at the end we will have

$$A' = A'_0 + B^R + B', (1.5)$$

where B' is smaller (in order, say) than B. Of course, we also hope that the resonant part B^R is better in some sense than the initial perturbation B; in many applications, the operator B^R acts in subspaces of our Hilbert space generated by 'specially designated and geometrically defined' areas of the phase space.

After we have reduced our operator to the improved form (1.5), in principle we can repeat the same procedure — finitely, or even infinitely many times. The latter process is much more difficult to realise, and we will not give examples of it in this paper. However, in many settings we indeed have to run this procedure several times (more than once) in order to achieve the desired 'smallness' of the remainder. In other words, we construct the 'improved' operator in the form

$$A_n = \exp(i\Psi_n) \dots \exp(i\Psi_2) \exp(i\Psi_1) A \exp(-i\Psi_1) \exp(-i\Psi_2) \dots \exp(-i\Psi_n)$$

We call this method *the consecutive gauge transform*. Sometimes, it is more convenient to look for the improved operator in the form

$$A^{(n)} = \exp(i(\Psi_n + \dots + \Psi_2 + \Psi_1))A\exp(-i(\Psi_1 + \Psi_2 + \dots + \Psi_n)),$$

which we call *the parallel gauge transform*. In both situations, the operators Ψ_j are solutions of equations similar in form to (1.4).

Another important distinction between different variations of the gauge transform is as follows. In order to prove that the order of the remainder B' is smaller than the order of B, we have to estimate the orders of various commutators. Sometimes, it is enough to have the basic estimate: the order of the commutator is not greater than the sum of the orders of its entries. This estimate holds without any restrictions, but for it to be effective we need to have some a priori inequalities between the orders of the principal term A_0 and the perturbation B; we call this approach the weak gauge transform. On the other hand, quite often we can improve our estimate on commutators: for example, in the classical scalar pseudo-differential calculus, the order of the commutator can be estimated by the sum of the orders of the entries minus one. If we have such an estimate, we can guarantee that the order of B' is indeed smaller than the order of B, assuming nothing other than that the order of A_0 is larger than the order of B. This approach is called the strong gauge transform. In this paper, we will define the weak and strong gauge transforms rigorously and give a general abstract setting in which they can be applied. We discuss the advantages and drawbacks of both types of gauge transforms and finish with a couple of concrete applications.

The first application is to obtain asymptotic expansions for the density of states of elliptic almost periodic operator systems. Under some technical conditions described later, we may either obtain complete or limited expansions as the spectral parameter goes to $\pm\infty$. The other application is to prove that some elliptic periodic systems have the Bethe–Sommerfeld property. This will be done under the same conditions that allow us to obtain a complete asymptotic expansion for the density of states. In either of these cases, some Dirac operators are examples of those to which we can apply our results.

1.2. **Description of the results for elliptic systems and the Dirac operator.** While describing the precise class of operators *A* for which we obtain spectral asymptotics requires definitions that are made later, we can make these results explicit for Dirac operators in dimension 2 and 3 perturbed by classical pseudo-differential almost periodic operators right away. The two-dimensional Dirac operator with mass *M* acts in $L^2(\mathbb{R}^2; \mathbb{C}^2)$ and is given by

$$\mathbf{A}_{2,M} := -i(\sigma_1 \partial_{x_1} + \sigma_2 \partial_{x_2}) + \sigma_3 M,$$

where $\sigma_1, \sigma_2, \sigma_3$ are the Pauli matrices

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \text{and} \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

The three-dimensional Dirac operator with mass M acts in $L^2(\mathbb{R}^3; \mathbb{C}^4)$ and is given by

$$\mathbf{A}_{3,M} := -i\left(\gamma_1\partial_{x_1} + \gamma_2\partial_{x_2} + \gamma_3\partial_{x_3}\right) + \Gamma M,$$

where the matrices γ_i , Γ are the Dirac matrices (see [Upm02])¹

$$\gamma_j = \begin{pmatrix} \mathbf{0} & \sigma_j \\ \sigma_j & \mathbf{0} \end{pmatrix}$$
, and $\Gamma = \begin{pmatrix} \mathrm{Id}_2 & \mathbf{0} \\ \mathbf{0} & -\mathrm{Id}_2 \end{pmatrix}$.

We obtain asymptotic expansions for the density of states of operators of the type $\mathbf{A} = \mathbf{A}_{d,M} + \mathbf{B}$ under the assumption that **B** is a 'generic' almost periodic pseudo-differential perturbation. The precise meaning of generic is given in Section 7. The density of states for elliptic differential operators *A* that are not semi-bounded can be defined by the formula

$$N(\lambda; A) := \lim_{L \to \infty} \frac{N(\lambda; A_D^{(L)})}{(2L)^d}.$$

Here, $A_D^{(L)}$ is the restriction of A to the cube $[-L, L]^d$ with Dirichlet boundary condition, and $N(\lambda; A_D^{(L)})$ is the counting function for the discrete eigenvalues of $A_D^{(L)}$ in the interval $[0, \lambda)$ when $\lambda > 0$ and $(\lambda, 0]$ when $\lambda < 0$. Later, we will give several equivalent definitions of $N(\lambda)$ which are more convenient to work with and allow pseudo-differential perturbations.

Theorem 1.1. Let $\mathbf{A} = \mathbf{A}_{2,M} + \mathbf{B}$, where \mathbf{B} is a generic symmetric pseudo-differential operator with almost periodic coefficients of order $\beta < 1$ acting in $L^2(\mathbb{R}^2; \mathbb{C}^2)$. Then, there is a complete asymptotic expansion for the density of states of \mathbf{A} in the sense that for every K > -2, there is a finite set $L \subset (0, 2 + K)$ and constants C_j^{\pm} , $C_{i,\log}^{\pm}$, $j \in L \cup \{0\}$ such that

$$N(\pm\lambda;\mathbf{A}) = C_0^{\pm}\lambda^2 + \sum_{j \in L} \left(C_j^{\pm}\lambda^{2-j} + C_{j,\log}^{\pm}\lambda^j \log\lambda \right) + O\left(\lambda^{-K}\right)$$

as $\lambda \to \infty$.

Theorem 8.2 is a more general version of Theorem 1.1. It is applicable to elliptic systems of pseudodifferential operators whose principal symbol has only simple eigenvalues.

We obtain a restricted expansion for the three-dimensional case.

Theorem 1.2. Let $\mathbf{A} = \mathbf{A}_{3,M} + \mathbf{B}$, where **B** is a generic operator of the form

$$\mathbf{B} = B_1 \gamma_1 + B_2 \gamma_2 + B_3 \gamma_3 + B_{\Gamma} \Gamma + B_{\mathrm{Id}} \mathrm{Id}_4,$$

where each B_j , $j \in \{1,2,3,\Gamma, \text{Id}\}$ is a scalar symmetric pseudo-differential operator with almost periodic coefficients of order β , $0 \le \beta \le 1/2$. Then, writing $\gamma^* = \max \{\beta - 1, 2\beta - 1\}$ there is a finite

¹Many authors would write α_j for γ_j and β for Γ , see e.g. [Tha92]. We keep our convention in line with higherdimensional generalisations and to avoid some notational conflicts later on.

set $L \subset (0, 1 - \gamma^*)$ and constants C_j^{\pm} , $C_{j,\log}^{\pm}$, $j \in L \cup \{0\}$ such that

$$N(\pm\lambda;A) = C_0^{\pm}\lambda^3 + \sum_{j\in L} \left(C_j^{\pm}\lambda^{3-j} + C_{j,\log}^{\pm}\lambda^{3-j}\log\lambda \right) + O\left(\lambda^{2+\gamma^*}\right)$$

as $\lambda \to \infty$.

This time, it is Theorem 8.1 which is a more general version of Theorem 1.2. It is applicable to elliptic systems of pseudodifferential operators whose principal symbol has multiple eigenvalues under some more restrictive conditions on the perturbation.

We also obtain that two-dimensional Dirac operators satisfy the Bethe-Sommerfeld property.

Theorem 1.3. Let $\mathbf{A} = \mathbf{A}_{2,M} + \mathbf{B}$, where \mathbf{B} is a symmetric pseudo-differential operator of order $\beta < 1$ with periodic coefficients. Then, there exists $\lambda_0 > 0$ such that the spectrum of \mathbf{A} contains intervals $(-\infty, -\lambda_0]$ and $[\lambda_0, \infty)$.

This theorem also has a more general version in Theorem 10.1. It is applicable to systems whose principal symbol has only simple eigenvalues.

In Section 12, we also describe generalisations of these results to higher dimensional Dirac operators, and give some technical conditions under which we can get complete asymptotic expansions or the Bethe–Sommerfeld property for the three-dimensional Dirac operator.

1.3. **Description of the main results and plan of the paper.** In the first half of our paper, we discuss the gauge transform in an abstract setting. The setting is developed while keeping in mind particular applications to almost periodic operators. As such, the space on which the operators act looks like an abstract version of a Besicovitch space. In the second half, we will discuss the specific applications of the results obtained in the first half to elliptic systems of pseudo-differential almost periodic operators; in particular, in the last section we will show that Dirac operators are a specific example of them. An interesting part of the application of our methods to systems is that we need to intertwine and alternate the use of the weak and strong gauge transforms, whereas in the past only one type was used at a time. In order to help the reader familiar with previous literature on the method of gauge transform, we have kept the notation as close as possible to the one used in [MPS14, PS10].

Plan and results of Part I. In Section 2 we define an algebra of operators S^{∞} acting on a non separable Hilbert space which should be thought of as an abstract version of a Besicovitch space. For some set Ξ this algebra will be concretely realised on $\ell^2(\Xi)$ through a group action on its basis elements. This algebra is filtered as an algebra of pseudo-differential operators on $\ell^2(\Xi)$, and it has similar properties to those of classical pseudo-differential operators in the PDE sense. Their natural domains are generalisations of Sobolev spaces. This section contains many technical but

very useful lemmas describing boundedness properties, adjoints, compositions and commutators of operators in S^{∞} . One of the main differences with classical pseudo-differential operators is illustrated in Proposition 2.16, which plays the role of the Calderon–Vaillancourt theorem in our setting. It essentially says that we can directly correlate symbol norms of operators with the norms of individual summands in Paley-Wiener type decompositions.

In Section 3, we turn our attention to some natural subspaces of S^{∞} – operators that are either elliptic or diagonal. Just as in the classical setting, our definition of elliptic operators allows us to characterise natural domains of self-adjointness for operators in the algebra S^{∞} . The three main results of this section illustrate the three most important properties of elliptic operators. In Proposition 3.5, they are shown to admit a parametrix, and are therefore invertible up to a controllable error. Lemma 3.6 is used repeatedly throughout the paper and shows that lower order perturbations of elliptic operators are relatively bounded, with explicit bounds. Finally, in Proposition 3.7, we show that elliptic operators are closed and self-adjoint if symmetric.

In Section 4, we consider the situation where operators in S^{∞} are affiliated to a I_{∞} or II_{∞} factor. This is common in the study of almost-periodic operators and their generalisation. We define a general notion of density of states measures (DSM) in S^{∞} as traces in the affiliated I_{∞} or II_{∞} factor. We give a variational description of the DSM of an interval *J* even in situations where the operator is not bounded below. This is used to show the principal results of this section: small perturbations of elliptic self-adjoint operators do not change their density of states much. The definition of 'smallness' of the perturbation is made clear in that section. In Lemma 4.11, we control to what extent perturbations of smaller order can affect the DSM, whereas in Lemma 4.12 it is perturbations that are spectrally supported away from the interval *J* that are shown to have a small effect.

In Section 5, we describe the abstract gauge transform scheme, which is split into two cases: the weak and strong gauge transforms. In both cases, we describe the resonant regions geometrically as subsets of the index set Ξ . The consecutive scheme for the weak gauge transform is described in Lemmas 5.5 and 5.6 and Corollary 5.7, whereas the parallel scheme is described in Proposition 5.9. In both cases, only trivial estimates on the commutator are used. In Lemma 5.10, we describe conditions under which a stronger scheme can be used. Since conditions for the strong transform to be applicable are varying in nature, we do not attempt at completely classifying them.

Finally, in Section 6, we describe the case where the symbols are functions into $Mat^{m \times m}(\mathbb{C})$ rather than \mathbb{C} . We describe how this can be reduced to the abstract scalar case and introduce a new class of operator systems: uncoupled operators. Our goal is to show that under some specific conditions, elliptic systems are unitarily equivalent to uncoupled operators up to a remainder which we can control. In that light, the main results of this section are Theorems 6.4

and 6.6 which give explicit conditions under which one can use the weak gauge transform to conjugate elliptic symmetric operators into almost uncoupled ones. The remainders are small (in the sense of Section 4) perturbations.

Plan and results of Part II. In the second part, we apply the results of Part I to concrete systems of elliptic pseudo-differential operators with periodic and almost periodic perturbations. More specifically, we study operator systems of the form $\mathbf{A} = \mathbf{A}_0 + \mathbf{B}$, defined on a dense domain in $L^2(\mathbb{R}^d;\mathbb{C}^m)$ where \mathbf{A}_0 is defined as in (1.1), and \mathbf{B} is a pseudo-differential perturbation of order $\beta < \alpha$. In Section 7, we give a description of these operators in term of Besicovitch space, and we make the relevant definitions concerning periodic operators. In Sections 7.2 and 8, we obtain asymptotic expansions for the IDS. In Sections 10 and 11 we prove that some elliptic systems of operators have the Bethe–Sommerfeld property using some combinatorial geometric arguments. Finally in Section 12 we expose how Dirac operators may fit in our setting.

Since the precise description of the results requires some notations and language defined in Part I, we postpone their description to the beginning of Part II.

Acknowledgements. The research of JL and LP was supported by EPSRC grant EP/P024793/1. The research of JL was also partially supported by NSERC's postdoctoral fellowship. The research of SM was supported by RSF grant 18-11-00032. The research of RS was supported by NSF grant DMS-1814664.

PART I : AN ABSTRACT GAUGE TRANSFORM SCHEME

2. GENERALISED ALMOST-PERIODIC OPERATORS

In this section, we define an algebra of generalised almost-periodic operators. We start by defining the space on which those operators are defined. We also define generalised Sobolev spaces which are their natural domains. We then describe the algebraic properties of the generalised almost-periodic operators, and obtain version of the Calderon–Vaillancourt theorem in our context in Proposition 2.16.

2.1. **Generalised Sobolev spaces.** Let Ξ be an infinite, possibly uncountable set equipped with a weight function $\langle \cdot \rangle : \Xi \to [1, \infty)$. We will often call Ξ *the index set*. For $\gamma \in \mathbb{R}$ we define the spaces

$$\mathrm{H}^{\gamma}(\Xi) := \begin{cases} x : \Xi \to \mathbb{C} \\ x : \xi \mapsto x_{\xi} \end{cases} : \sum_{\xi \in \Xi} \langle \xi \rangle^{2\gamma} |x_{\xi}|^{2} < \infty \end{cases}$$

and

$$\mathrm{G}^\infty(\Xi):=\bigcap_{\gamma\in\mathbb{R}}\mathrm{H}^\gamma(\Xi).$$

In particular, every $x \in H^{\gamma}(\Xi)$ vanishes at all but countably many $\xi \in \Xi$. Every $H^{\gamma}(\Xi)$ is a Hilbert space with inner product

$$(x, y)_{\mathrm{H}^{\gamma}(\Xi)} := \sum_{\xi \in \Xi} \langle \xi \rangle^{2\gamma} x_{\xi} \overline{y_{\xi}}.$$
(2.1)

It is easy to see that $H^0(\Xi) = \ell^2(\Xi)$ with the standard orthonormal basis indexed bijectively from Ξ as

$$\mathscr{E} := \left\{ \mathbf{e}_{\xi} : \xi \in \Xi \right\}, \qquad \mathbf{e}_{\xi} : \eta \in \Xi \mapsto \begin{cases} 1 & \text{if } \eta = \xi, \\ 0 & \text{if } \eta \neq \xi \end{cases}$$

and that $H^{\gamma_1}(\Xi) \subset H^{\gamma_2}(\Xi)$ for all $\gamma_1 > \gamma_2 \in \mathbb{R}$. When there is no risk of confusion, we will write $H^{\gamma} := H^{\gamma}(\Xi)$.

2.2. An algebra of operators. Let *G* be a group that acts from the left on Ξ , so that the action is free, i.e. only the identity of *G* has fixed points. We denote by $g \triangleright \xi$ the action of $g \in G$ on $\xi \in \Xi$. Starting from the weight function $\langle \cdot \rangle$ on Ξ we define one on *G* by

$$\langle g \rangle := 1 + \sup_{\xi \in \Xi} \left| \langle g \triangleright \xi \rangle - \langle \xi \rangle \right|.$$
(2.2)

We assume that *G* has a *bounded range of action*, which means that $\langle g \rangle$ is finite for all $g \in G$.

It will be useful for future convenience to observe the following properties of the weight function:

Lemma 2.1. For all $f, g \in G, \xi \in \Xi$ and $t \in \mathbb{R}$ the following relations hold:

(1)

$$\langle g \rangle = \langle g^{-1} \rangle; \tag{2.3}$$

(2) Peetre-type inequalities:

$$\langle g \rangle^{-1} \langle \xi \rangle \le \langle g \triangleright \xi \rangle \le \langle g \rangle \langle \xi \rangle \tag{2.4}$$

and

$$\langle fg \rangle^{t} \le \min\left\{ \langle f \rangle^{t} \langle g \rangle^{|t|}, \langle f \rangle^{|t|} \langle g \rangle^{t} \right\}.$$
(2.5)

Proof. For all $g \in G$, $\xi \in \Xi$ the definition (2.2) implies (2.3) and the estimates

$$\max\left\{1, 1 + \langle \xi \rangle - \langle g \rangle\right\} \le \langle g \triangleright \xi \rangle \le \langle \xi \rangle + \langle g \rangle - 1.$$
(2.6)

Note the relations

$$a+1-b = ((b-1)(a-b)+a)/b \ge a/b$$
, for all $a \ge b \ge 1$.
 $a+b-1 \le a+b-1+(a-1)(b-1) = ab$, for all $a, b \ge 1$,

The first estimate in (2.4) follows from (2.6) and (2.7), the second from (2.6) and (2.8). Now by (2.2) and (2.7) for all $f, g \in G$ we obtain

$$\langle fg \rangle \leq 1 + \sup_{\xi \in \Xi} \left| \langle fg \triangleright \xi \rangle - \langle g \triangleright \xi \rangle \right| + \sup_{\xi \in \Xi} \left| \langle g \triangleright \xi \rangle - \langle \xi \rangle \right|$$

$$= \langle f \rangle + \langle g \rangle - 1 \leq \langle f \rangle \langle g \rangle,$$

$$(2.9)$$

which implies (2.5) for *t* > 0. Now (2.9) and (2.3) imply

$$\langle g \rangle = \langle f^{-1}fg \rangle \le \langle f \rangle \langle fg \rangle$$
 and $\langle f \rangle = \langle fgg^{-1} \rangle \le \langle fg \rangle \langle g \rangle$

which delivers (2.5) for t < 0. The case t = 0 is trivial.

Definition 2.2. We call a function $b: G \times \Xi \to \mathbb{C}$, $(g, \xi) \mapsto b_g(\xi)$ an almost periodic *symbol* if there exists a countable set $\Theta \subset G$, closed under inversion and containing the identity id_G , such that for all $g \in G \setminus \Theta$, $b_g(\xi) \equiv 0$. Whenever there is no risk of confusion, we will write $\mathrm{id} := \mathrm{id}_G$. We call Θ a *frequency set* for *b* and the functions $\{b_\theta(\cdot)\}_{\theta \in \Theta}$ the *Fourier coefficients* of *b*. For every symbol *b* and every $\gamma \in \mathbb{R}$, $l \ge 0$, we define the family of norms

$$\|b\|_{l}^{(\gamma)} := \sum_{\theta \in \Theta} \langle \theta \rangle^{l} \sup_{\xi \in \Xi} \left(\langle \xi \rangle^{-\gamma} | b_{\theta}(\xi) | \right).$$
(2.10)

The class of *symbols of order* γ is defined as

$$\mathbf{S}^{\gamma} := \mathbf{S}^{\gamma}(G, \Xi) := \left\{ b : G \times \Xi \to \mathbb{C} : \|b\|_{l}^{(\gamma)} < \infty \text{ for all } l \ge 0 \right\}.$$
 (2.11)

The space of symbols is naturally a linear space. It is clear that if Θ is a frequency set for a symbol, then any $\Gamma \supset \Theta$ is also one. It is obvious from the definition that $\|\cdot\|_l^{(\gamma)}$ is a decreasing function of γ and an increasing function of l, thus

$$\mathbf{S}^{\gamma_1} \subset \mathbf{S}^{\gamma_2}$$
, for all $\gamma_1 \leq \gamma_2$.

We introduce

$$\mathbf{S}^{\infty} := \bigcup_{\gamma \in \mathbb{R}} \mathbf{S}^{\gamma}$$
 and $\mathbf{S}^{-\infty} := \bigcap_{\gamma \in \mathbb{R}} \mathbf{S}^{\gamma}$

Lemma 2.3. For every $\gamma \in \mathbb{R}$, the space \mathbf{S}^{γ} equipped with the family of norms $\{\|\cdot\|_{l}^{(\gamma)}\}_{l\geq 0}$ is a Fréchet space.

Proof. Consider a sequence

 $(b_n)_{n\geq 1} \subset \mathbf{S}^{\gamma}$

that is Cauchy with respect to $\|\cdot\|_l^{(\gamma)}$ for every $l \ge 0$, and denote by $\Theta(n)$ a frequency set for each b_n . Then, for all $\theta \in G$, we observe that $b_{\theta}(\xi) := \lim_{n \to \infty} (b_n)_{\theta}(\xi)$ exists and vanishes outside the countable set $\Theta = \bigcup_n \Theta(n)$. It is a simple computation to see that $b \in \mathbf{S}^{\gamma}$ with $\|b_n - b\|_l^{(\gamma)} \to 0$, as $n \to \infty$, for all $l \ge 0$. Hence, the claim follows.

Definition 2.4. Let $b: G \times \Xi \to \mathbb{C}$ be a symbol with frequency set $\Theta \subset G$ and

$$(b_{\theta}(\xi))_{\theta \in \Theta} \in \ell^2(\Theta)$$
, for all $\xi \in \Xi$.

Then the *almost periodic linear operator associated to b* is

$$B := \operatorname{Op}(b) : \operatorname{span}(\mathscr{E}) \to \ell^2(\Xi)$$

defined by

$$B\mathbf{e}_{\xi} := \sum_{\theta \in \Theta} b_{\theta}(\xi) \mathbf{e}_{\theta \succ \xi}, \quad \text{for all } \xi \in \Xi.$$
(2.12)

Remark 2.5. If $b \in S^{\infty}$, then, in view of (2.10) and (2.11), $(b_{\theta}(\xi))_{\theta \in \Theta} \in \ell^{1}(\Theta) \subset \ell^{2}(\Theta)$ holds for all $\xi \in \Xi$. This means that we can associate an almost periodic operator to every symbol in S^{∞} . On the other hand, since the group action of *G* on Ξ is free, *b* can be recovered from *B* via the identity

$$b_g(\xi) = (\mathbf{e}_{g \triangleright \xi}, B\mathbf{e}_{\xi})_{\ell^2(\Xi)}, \quad \text{for all } g \in G, \, \xi \in \Xi.$$

$$(2.13)$$

Thus, there is a one-to-one correspondence between almost periodic symbols and almost periodic operators. This correspondence is in contrast to the case of classical pseudo-differential operators where this correspondence is only modulo smoothing operators. Hence, we allow ourselves to overload the notation and write $B = \operatorname{Op}(b) \in \mathbf{S}^{\gamma}$ if $b \in \mathbf{S}^{\gamma}$, $\gamma \in \mathbb{R} \cup \{\pm \infty\}$, and let $\|B\|_{l}^{(\gamma)} := \|b\|_{l}^{(\gamma)}$ for all $l \ge 0$, $\gamma \in \mathbb{R}$. Note that this correspondence gets lost if one does not require the group action of G on Ξ to be free. Our construction can be generalised to such non-free group actions, but for simplicity of the exposition we do not do it in this paper.

We call *B* quasi-periodic if *b* admits a finite frequency set. A simple example of a quasiperiodic operator of class \mathbf{S}^{γ} , $\gamma \in \mathbb{R}$, is Op(h) with

$$h_g(\xi) := \begin{cases} \widetilde{h}(\xi) & \text{if } g = \text{id,} \\ 0 & \text{otherwise.} \end{cases}$$

Here, \tilde{h} is a function on Ξ satisfying $|\tilde{h}(\xi)| \leq \langle \xi \rangle^{\gamma}$ for all $\xi \in \Xi$.

Remark 2.6. Our terminology is justified by the following example. Suppose that *G* is a locally compact abelian (LCA) group and G_B is its Bohr compactification, see [Shu78, §1]. Index by Ξ the set of characters $\tilde{\mathscr{E}} := \{\tilde{\mathbf{e}}_{\xi} : \xi \in \Xi\}$ of *G* or, equivalently, G_B . On CAP(*G*), the continuous almost periodic functions on *G*, we can define an inner product $(f, g) = \mathcal{M}(f\overline{g})$, where $\mathcal{M}(f)$ is the mean of *f* with respect to the normalised Haar measure on G_B . The Besicovitch space B²(*G*) is defined as the closure of CAP(*G*) with respect to the norm induced by this inner product. By [Shu78, Proposition 1.5], the map

$$\mathscr{E} \to \widetilde{\mathscr{E}} \quad \mathbf{e}_{\xi} \mapsto \widetilde{\mathbf{e}}_{\xi},$$

extends to an isometric isomorphism $\ell^2(\Xi) \to B^2(G)$. In particular, for $G = (\mathbb{R}^d, +)$, one has $\widetilde{\mathscr{E}} = \{\mathbf{x} \mapsto \exp(i\mathbf{x} \cdot \boldsymbol{\xi}), \ \boldsymbol{\xi} \in \mathbb{R}^d\}$ and the operators in $\mathbf{S}^{\infty}(\mathbb{R}^d, \mathbb{R}^d)$ correspond to almost periodic pseudo-differential operators in $B^2(\mathbb{R}^d)$ or $L^2(\mathbb{R}^d)$, see [Shu78, §3–4] and [PS12, Equation (8.8)]. The present work can be applied to more general settings, for example, when the underlying group *G* is non-abelian. Note that the Bohr compactification construction is inadequate in that situation, e.g., for $G = SL(2; \mathbb{R})$ we have $G_B = \{id\}$, see [Shu78, p. 4].

From Lemma 2.3 we obtain the following corollary.

Corollary 2.7. Let $(B_n)_{n\geq 1} \subset \mathbf{S}^{\gamma}$ be such that

$$\sum_{n\geq 1} \|B_n\|_l^{(\gamma)} < \infty$$

for all $l \ge 0$. Then the sum

$$B := \sum_{n \ge 1} B_n$$

converges in \mathbf{S}^{γ} with

$$||B||_l^{(\gamma)} \le \sum_{n\ge 1} ||B_n||_l^{(\gamma)}.$$

Up until now, operators from S^{∞} were only defined on span(\mathscr{E}). We now show that they can be extended in a natural way.

Lemma 2.8. For every $\beta, \gamma \in \mathbb{R}$ the operator $B \in S^{\gamma}$ can be uniquely extended to a bounded linear operator $B : H^{\beta} \to H^{\beta-\gamma}$. Moreover, we have the bound

$$\|B\|_{\mathrm{H}^{\beta} \to \mathrm{H}^{\beta-\gamma}} \le \|B\|_{|\beta-\gamma|}^{(\gamma)}$$

Proof. Let $x, y \in \text{span}(\mathcal{E})$, i.e. $x_{\xi} = y_{\xi} = 0$ for all but finitely many ξ . Then, the Cauchy–Schwarz and Peetre inequalities (2.4) imply

.

$$\begin{split} \left| (x, By)_{\mathbf{H}^{\beta-\gamma}} \right| &= \left| \sum_{\theta \in \Theta} \sum_{\xi \in \Xi} \langle \theta \succ \xi \rangle^{2(\beta-\gamma)} b_{\theta}(\xi) \overline{x_{\theta \succ \xi}} \, y_{\xi} \right| \\ &\leq \sum_{\theta \in \Theta} \langle \theta \rangle^{|\beta-\gamma|} \sup_{\xi \in \Xi} \left(\langle \zeta \rangle^{-\gamma} | b_{\theta}(\zeta) | \right) \times \\ &\times \left(\sum_{\xi \in \Xi} \langle \theta \succ \xi \rangle^{2(\beta-\gamma)} \left| x_{\theta \succ \xi} \right|^2 \right)^{1/2} \left(\sum_{\xi \in \Xi} \langle \xi \rangle^{2\beta} \left| y_{\xi} \right|^2 \right)^{1/2} \\ &\leq \|B\|_{|\beta-\gamma|}^{(\gamma)} \|x\|_{\mathbf{H}^{\beta-\gamma}} \|y\|_{\mathbf{H}^{\beta}}. \end{split}$$

The claim follows by density of span \mathscr{E} in H^{α} for all $\alpha \in \mathbb{R}$.

We obtain the following immediate corollary.

Corollary 2.9. *Every* $B \in S^0$ *extends to a bounded operator on* $\ell^2 = H^0$ *such that*

$$\|B\|_{\ell^2 \to \ell^2} \le \|B\|_0^{(0)}.$$

Definition 2.10. For $b \in \mathbf{S}^{\infty}$, we define

$$b_{\theta}^{\dagger}(\xi) := \begin{cases} \overline{b_{\theta^{-1}}(\theta \triangleright \xi)} & \text{if } \theta \in \Theta, \\ 0 & \text{if } \theta \in G \setminus \Theta \end{cases}$$
(2.14)

for all $\xi \in \Xi$, where Θ is a frequency set for *b*.

Lemma 2.11. If $b \in \mathbf{S}^{\gamma}$, then $b^{\dagger} \in \mathbf{S}^{\gamma}$. Moreover, for all $x, y \in \mathbf{H}^{\gamma}$, one has

$$(x, By)_{\ell^2(\Xi)} = (B^{\dagger}x, y)_{\ell^2(\Xi)}, \quad i.e. \ B^{\dagger} \subset B^*.$$
 (2.15)

In particular, B is symmetric on H^{γ} if and only if $B = B^{\dagger}$.

Proof. Every frequency set Θ for $b \in \mathbf{S}^{\gamma}$ is also one for b^{\dagger} . Moreover, since $\Theta = \Theta^{-1}$ holds by convention, (2.3) and (2.4) imply that for all $l \ge 0$,

$$\begin{split} \left\| b^{\dagger} \right\|_{l}^{(\gamma)} &= \sum_{\theta \in \Theta_{B}} \langle \theta \rangle^{l} \sup_{\xi \in \Xi} \left[\langle \xi \rangle^{-\gamma} | b_{\theta^{-1}}(\theta \triangleright \xi) | \right] \\ &= \sum_{\theta \in \Theta_{B}} \langle \theta \rangle^{l} \sup_{\xi \in \Xi} \left[\langle \theta^{-1} \triangleright \xi \rangle^{-\gamma} | b_{\theta^{-1}}(\xi) | \right] \\ &\leq \sum_{\theta \in \Theta_{B}} \langle \theta \rangle^{l+|\gamma|} \sup_{\xi \in \Xi} \left[\langle \xi \rangle^{-\gamma} | b_{\theta^{-1}}(\xi) | \right] \\ &= \sum_{\theta \in \Theta_{B}} \langle \theta \rangle^{l+|\gamma|} \sup_{\xi \in \Xi} \left[\langle \xi \rangle^{-\gamma} | b_{\theta}(\xi) | \right] \\ &= \| b \|_{l+|\gamma|}^{(\gamma)}, \end{split}$$

thus $b^{\dagger} \in \mathbf{S}^{\gamma}$ holds. Moreover, (2.13) and (2.14) yield

$$(\mathbf{e}_{\eta}, B\mathbf{e}_{\xi}) = (B^{\dagger}\mathbf{e}_{\eta}, \mathbf{e}_{\xi}), \quad \text{for all } \eta, \xi \in \Xi.$$
 (2.16)

In view of Lemma 2.8 and the density of span \mathscr{E} in H^{γ}, (2.16) extends to (2.15). This finishes the proof of the lemma.

Definition 2.12. Let $a, b \in S^{\infty}$ be symbols with frequency sets Θ_a and Θ_b . The *composed symbol* $a \circ b$ with frequency set

$$\Theta_{a \circ b} := \Theta_a \Theta_b := \{ \theta_a \theta_b : \theta_a \in \Theta_a, \theta_b \in \Theta_b \}$$

is defined as

$$(a \circ b)_{\theta}(\xi) := \sum_{\theta_a \theta_b = \theta} a_{\theta_a}(\theta_b \triangleright \xi) b_{\theta_b}(\xi) \quad \text{for all } \theta \in \Theta_{a \circ b}, \, \xi \in \Xi.$$

Lemma 2.13. For $\alpha, \beta \in \mathbb{R}$ let $A = Op(a) \in S^{\alpha}$ and $B = Op(b) \in S^{\beta}$. Then $AB \in S^{\alpha+\beta}$ and $AB = Op(a \circ b)$. Moreover, for all $l \ge 0$ we have the bound

$$\|AB\|_{l}^{(\alpha+\beta)} \le \|A\|_{l}^{(\alpha)} \|B\|_{l+|\alpha|}^{(\beta)}.$$
(2.17)

Proof. The frequency set $\Theta_{a \circ b}$ is, clearly, a countable set. For any $l \ge 0$, we have

$$\begin{split} \|a \circ b\|_{l}^{(\alpha+\beta)} &= \sum_{\theta \in \Theta_{a \circ b}} \sum_{\theta_{a} \theta_{b} = \theta} \langle \theta \rangle^{l} \sup_{\xi \in \Xi} \left(\langle \xi \rangle^{-\alpha-\beta} \left| a_{\theta_{a}}(\theta_{b} \triangleright \xi) \right| \left| b_{\theta_{b}}(\xi) \right| \right) \\ &\leq \sum_{\theta_{b} \in \Theta_{b}} \langle \theta_{b} \rangle^{l+|\alpha|} \sup_{\xi \in \Xi} \left(\langle \xi \rangle^{-\beta} \left| b_{\theta_{b}}(\xi) \right| \right) \times \\ &\times \sum_{\theta_{a} \in \Theta_{a}} \langle \theta_{a} \rangle^{l} \sup_{\zeta \in \Xi} \left(\langle \theta_{b} \triangleright \zeta \rangle^{-\alpha} \left| a_{\theta_{a}}(\theta_{b} \triangleright \zeta) \right| \right) \\ &\leq \|a\|_{l}^{(\alpha)} \|b\|_{l+|\alpha|}^{(\beta)}. \end{split}$$

Thus $a \circ b \in \mathbf{S}^{\alpha+\beta}$ and (2.12) implies $AB = \operatorname{Op}(a \circ b)$.

It is natural to consider operators from S^{∞} on the common domain H^{∞} . Then Lemmata 2.8, 2.11, and 2.13 yield the following corollary.

Corollary 2.14. $\mathbf{S}^{\infty} = \bigcup_{\gamma \in \mathbb{R}} \mathbf{S}^{\gamma}$ is a *-algebra of operators on H^{∞} , filtered by \mathbb{R} , with involution \dagger . The subalgebra of regularising operators $\mathbf{S}^{-\infty}$ forms a two-sided ideal of \mathbf{S}^{∞} .

We also consider the adjoint actions ad(A, B) := i(AB - BA) with the frequency set $\Theta_{ad(a,b)} = \Theta_{a\circ b} \cup \Theta_{b\circ a}$. The Fourier coefficients of ad(A, B) are

$$\mathrm{ad}(a,b)_{\theta}(\xi) = \mathrm{i}\left(\sum_{\theta_{a}\theta_{b}=\theta} a_{\theta_{a}}(\theta_{b} \triangleright \xi) b_{\theta_{b}}(\xi) - \sum_{\theta_{b}\theta_{a}=\theta} b_{\theta_{b}}(\theta_{a} \triangleright \xi) a_{\theta_{a}}(\xi)\right),\tag{2.18}$$

for all $\theta \in \Theta_{\mathrm{ad}(a,b)}$. If *G* is commutative, (2.18) simplifies to

$$\mathrm{ad}(a,b)_{\theta}(\xi) = \mathrm{i} \sum_{\theta_a \theta_b = \theta} \left(a_{\theta_a}(\theta_b \triangleright \xi) b_{\theta_b}(\xi) - b_{\theta_b}(\theta_a \triangleright \xi) a_{\theta_a}(\xi) \right).$$

For k = 1, 2, 3, ... and $A, B, B_1, ..., B_k \in S^{\infty}$, we define recursively

$$ad(A; B_1, \dots, B_k) := ad (ad(A; B_1, \dots, B_{k-1}), B_k),$$

$$ad^0(A, B) := A,$$

$$ad^k(A; B) := ad (ad^{k-1}(A; B); B).$$

(2.19)

The following lemma is a direct consequence of Lemma 2.13.

Lemma 2.15. Let $k \in \mathbb{N}$ and assume that $A_j \in \mathbf{S}^{\gamma_j}$ for $0 \le j \le k$. Put

$$\gamma = \sum_{j=0}^{k} \gamma_j, \qquad \hat{\gamma} = \sum_{j=0}^{k} |\gamma_j|.$$

Then $\operatorname{ad}(A_0; A_1, \dots, A_k) \in \mathbf{S}^{\gamma}$. Furthermore, if for all $0 \le j \le k$ we have $A_j = A_j^{\dagger}$, then $\operatorname{ad}(A_0; A_1, \dots, A_k) = \operatorname{ad}(A_0; A_1, \dots, A_k)^{\dagger}$. Moreover, for all $l \ge 0$ we have

$$\|\mathrm{ad}(A_0; A_1, \dots, A_k)\|_l^{(\gamma)} \le 2^k \prod_{j=0}^k \|A_j\|_{l+\hat{\gamma}-|\gamma_j|}^{(\gamma_j)}.$$
(2.20)

In particular, for any $A \in \mathbf{S}^{\alpha}$, $B \in \mathbf{S}^{0}$ and $k \in \mathbb{N}$ we obtain the estimate

$$\left\| \operatorname{ad}^{k}(A;B) \right\|_{l}^{(\alpha)} \leq 2^{k} \left\| A \right\|_{l}^{(\alpha)} \left(\left\| B \right\|_{l+|\alpha|}^{(0)} \right)^{k}.$$
(2.21)

For some Ξ and *G* it may be possible to improve this lemma and show that $ad(A, B) \in \mathbf{S}^{\gamma}$ holds with $\gamma < \alpha + \beta$ for all $A \in \mathbf{S}^{\alpha}$, $B \in \mathbf{S}^{\beta}$. This will be discussed in Section 5.4.

The following proposition provides bounds on norms of operators restricted to 'annuli' in Ξ .

Proposition 2.16. For $1 \le m \le M \le \infty$, let $\Upsilon \subset \{\xi \in \Xi : m \le \langle \xi \rangle \le M\}$ and denote by P_{Υ} the orthogonal projection in $\ell^2(\Xi)$ onto the closure of span $\{\mathbf{e}_{\xi} : \xi \in \Upsilon\}$. Then, for any $A \in \mathbf{S}^{\gamma}$ with $\gamma \ge 0$, the norm inequality

$$\|AP_{\Upsilon}\|_{\ell^2 \to \ell^2} \le M^{\gamma} \|A\|_0^{(\gamma)} \tag{2.22}$$

holds. For any $A \in \mathbf{S}^{\gamma}$ with $\gamma \leq 0$, we get the inequality

$$||AP_{\Upsilon}||_{\ell^2 \to \ell^2} \le m^{\gamma} ||A||_0^{(\gamma)}.$$

Proof. Observe that P_{Υ} is a quasi-periodic operator with a frequency set $\Theta = \{id\}$ and the symbol $(p_{\Upsilon})_{id} = \mathbf{1}_{\Upsilon}$ (the indicator function of Υ). Thus, for all $\gamma \in \mathbb{R}$ and $l \ge 0$,

$$\|P_{\Upsilon}\|_{l}^{(-\gamma)} = \sup_{\xi \in \Upsilon} \langle \xi \rangle^{\gamma} \le \begin{cases} m^{\gamma} & \text{if } \gamma \le 0, \\ M^{\gamma} & \text{if } \gamma \ge 0. \end{cases}$$
(2.23)

If $M < \infty$ or $\gamma \le 0$, then Corollary 2.9 and Lemma 2.13 imply the bound

$$\|AP_{\Upsilon}\|_{\ell^{2} \to \ell^{2}} \le \|A\|_{0}^{(\gamma)} \|P_{\Upsilon}\|_{|\gamma|}^{(-\gamma)},$$

and the statement of the lemma follows from (2.23). On the other hand, the inequality (2.22) is trivial for $M = \infty$ and $\gamma > 0$.

3. Elliptic and diagonal operators

In this section, we introduce particular classes of operators from S^{∞} and study their properties. Some of these classes do depend on the specific choice of orthonormal basis \mathscr{E} for $\ell^2(\Xi)$. However, the class of operators on which our main theorems depend, that of elliptic operators, is invariant under change of basis.

$$\mathbf{DS}^{\infty} := \{A = \operatorname{Op}(a) \in \mathbf{S}^{\infty} : \{\mathrm{id}\} \text{ is a frequency set for } a\}$$

For symbols of operators from \mathbf{DS}^{∞} we can suppress the subscript id, i.e. we let $a(\xi) := a_{id}(\xi)$ for all $A = Op(a) \in \mathbf{DS}^{\infty}$, $\xi \in \Xi$. For $\alpha \in \mathbb{R} \cup \{-\infty\}$ we define $\mathbf{DS}^{\alpha} := \mathbf{DS}^{\infty} \cap \mathbf{S}^{\alpha}$. Introduce the map $\mathcal{D}: \mathbf{S}^{\infty} \to \mathbf{DS}^{\infty}$, $A \mapsto A^{\mathcal{D}}$, that projects A = Op(a) onto its diagonal part $A^{\mathcal{D}} := Op(a^{\mathcal{D}})$ where

$$a^{\mathscr{D}}(\xi) := a_{\mathrm{id}}(\xi),$$

i.e.

$$A^{\mathscr{D}}\mathbf{e}_{\xi} = \langle \mathbf{e}_{\xi}, A\mathbf{e}_{\xi} \rangle \mathbf{e}_{\xi}$$

holds for all $\xi \in \Xi$. We also define the *off-diagonal part* as $A^{\mathcal{OD}} := \operatorname{Op}(a^{\mathcal{OD}})$ with $a^{\mathcal{OD}} := a - a^{\mathcal{D}}$.

Note that for any $A \in \mathbf{S}^{\alpha}$ with $\alpha \in \mathbb{R}$ and all $l \ge 0$,

$$\left\|A^{\mathscr{D}}\right\|_{l}^{(\alpha)} + \left\|A^{\mathscr{O}}\right\|_{l}^{(\alpha)} = \left\|A\right\|_{l}^{(\alpha)}$$
(3.1)

and

$$\left\|A^{\mathscr{D}}\right\|_{l}^{(\alpha)} = \left\|A^{\mathscr{D}}\right\|_{0}^{(\alpha)}.$$
(3.2)

Definition 3.2. The set **DES**^{α} of *diagonal elliptic operators of order* $\alpha \in \mathbb{R}$ is defined as the set of operators $A = Op(a) \in \mathbf{DS}^{\alpha}$ for which there exist *ellipticity parameters* $\kappa > 0$ and $r \ge 1$ such that

$$|a(\xi)| \ge \kappa \langle \xi \rangle^{\alpha}$$
 for all $\xi \in \Xi$ such that $\langle \xi \rangle \ge r$. (3.3)

Let the set of ellipticity parameters (κ, r) of *A* be denoted by EP(*A*). Note that $(\kappa, r) \in \text{EP}(A)$ implies $(\tilde{\kappa}, \tilde{r}) \in \text{EP}(A)$ for all $0 < \tilde{\kappa} \le \kappa$, and $\tilde{r} \ge r$.

Definition 3.3. The set SES^{α} of *strongly elliptic operators of order* $\alpha \in \mathbb{R}$ consists of operators $A \in S^{\alpha}$ such that $A^{\mathcal{D}} \in DES^{\alpha}$ and $A^{\mathcal{D}\mathcal{D}} \in S^{\gamma}$ for some $\gamma < \alpha$. For $(\kappa, r) \in EP(A^{\mathcal{D}})$ we define P_r as the diagonal operator with symbol $\mathbf{1}_{\{\xi; \langle \xi \rangle \leq r\}}$. We also define P_r^c as $Id - P_r$, and

$$\widetilde{A}_{\kappa,r} := A^{\mathscr{D}} P_r^c + \kappa r^{\alpha} P_r.$$
(3.4)

Definition 3.4. The set \mathbf{ES}^{α} of *elliptic operators of order* $\alpha \in \mathbb{R}$ consists of operators $A \in \mathbf{S}^{\alpha}$ for which there exists a unitary $U \in \mathbf{S}^{0}$ with $UAU^{\dagger} \in \mathbf{SES}^{\alpha}$.

As we did with diagonal operators, we set

$$\mathbf{T}^{\infty} := \bigcup_{\gamma \in \mathbb{R}} \mathbf{T}^{\gamma}, \ \mathbf{T}^{-\infty} := \bigcap_{\gamma \in \mathbb{R}} \mathbf{T}^{\gamma}, \text{ for } \mathbf{T} \in \{\mathbf{DES}, \mathbf{SES}, \mathbf{ES}\}.$$

Clearly, both **SES**^{α} and **ES**^{α} are closed under addition of operators in **S**^{β}, $\beta < \alpha$.

Proposition 3.5. Let $A \in \mathbf{S}^{\infty}$ and $\alpha > 0$ such that $A^{\mathcal{D}} \in \mathbf{DES}^{\alpha}$. For any $(\kappa, r) \in \mathrm{EP}(A^{\mathcal{D}})$ the operator $\widetilde{A}_{\kappa,r}$ is invertible with $\widetilde{A}_{\kappa,r}^{-1} \in \mathbf{DS}^{-\alpha}$ and for all $l \ge 0$ we have

$$\left\|\widetilde{A}_{\kappa,r}^{-1}\right\|_{l}^{(\gamma)} = \left\|\widetilde{A}_{\kappa,r}^{-1}\right\|_{0}^{(\gamma)} \le \kappa^{-1} \begin{cases} r^{-\alpha} & \text{for } \gamma \ge 0, \\ r^{-\alpha-\gamma} & \text{for } -\alpha \le \gamma < 0. \end{cases}$$
(3.5)

Moreover, the following estimates hold for all $\gamma \in \mathbb{R}$ *, l* \geq 0*:*

$$\left\|A\widetilde{A}_{\kappa,r}^{-1} - \mathrm{Id}\right\|_{l}^{(\gamma-\alpha)} \le r^{\alpha-\gamma} + \frac{1}{\kappa} \left(r^{\alpha-\gamma} \left\|A^{\mathscr{D}}\right\|_{0}^{(\alpha)} + \left\|A^{\mathscr{O}}\right\|_{l}^{(\gamma)}\right)$$
(3.6)

and

$$\left\|\widetilde{A}_{\kappa,r}^{-1}A - \mathrm{Id}\right\|_{l}^{(\gamma-\alpha)} \leq r^{\alpha-\gamma} + \frac{1}{\kappa} \left(r^{\alpha-\gamma} \left\|A^{\mathcal{D}}\right\|_{0}^{(\alpha)} + \left\|A^{\mathcal{O}\mathcal{D}}\right\|_{l+\alpha}^{(\gamma)}\right).$$
(3.7)

Proof. We have that $\widetilde{A}_{\kappa,r} - A^{\mathcal{D}} = (\kappa r^{\alpha} - A^{\mathcal{D}})P_r$, and since $P_r \in \mathbf{S}^{-\infty}$, which is an ideal of \mathbf{S}^{∞} , we observe that $\widetilde{A}_{\kappa,r} \equiv A^{\mathcal{D}} \mod \mathbf{S}^{-\infty}$. By (3.3) and (3.4), $\widetilde{A}_{\kappa,r} \in \mathbf{DS}^{\infty}$ and its symbol satisfies

$$\left|\tilde{a}_{\kappa,r}(\xi)\right| = |a_{\mathrm{id}}(\xi)| \mathbf{1}_{\langle\xi\rangle>r} + \kappa r^{\alpha} \mathbf{1}_{\langle\xi\rangle\leq r} \ge \kappa \langle\xi\rangle^{\alpha} \mathbf{1}_{\langle\xi\rangle>r} + \kappa r^{\alpha} \mathbf{1}_{\langle\xi\rangle\leq r}$$

for all $\xi \in \Xi$. Hence $\widetilde{A}_{\kappa,r}^{-1} = \operatorname{Op}\left(\widetilde{a}_{\kappa,r}^{-1}\right) \in \mathbf{DS}^{-\alpha}$ and (3.5) holds.

The estimates (3.6) and (3.7) follow by applying (2.17) term-wise to the right hand sides of the identities

$$A\widetilde{A}_{\kappa,r}^{-1} - \mathrm{Id} = -P_r + \kappa^{-1}r^{-\alpha}A^{\mathscr{D}}P_r + A^{\mathscr{O}}\widetilde{A}_{\kappa,r}^{-1},$$
$$\widetilde{A}_{\kappa,r}^{-1}A - \mathrm{Id} = -P_r + \kappa^{-1}r^{-\alpha}A^{\mathscr{D}}P_r + \widetilde{A}_{\kappa,r}^{-1}A^{\mathscr{O}}\mathscr{D}$$

and taking (3.2) into account.

Lemma 3.6. Let $\beta \in \mathbb{R}$, $\alpha > \max(\beta, 0)$, $0 < \gamma < \alpha$, and assume that $A \in SES^{\alpha}$ with $A^{\emptyset \mathcal{D}} \in S^{\gamma}$ and $B \in S^{\beta}$. Then for $\beta \leq 0$ the operator *B* is bounded, and, in the case of $\alpha > \beta > 0$, for every $x \in H^{\alpha}$ and

$$(\kappa, r) \in \operatorname{EP}(A^{\mathcal{D}}) \cap \left\{ r \ge \left(\|A^{\mathcal{O}\mathcal{D}}\|_{0}^{(\gamma)} / \kappa \right)^{1/(\alpha - \gamma)} \right\},$$

we have

$$\|Bx\| \le \frac{r^{\beta-\alpha} \|B\|_{0}^{(\beta)}}{\kappa - r^{\gamma-\alpha} \|A^{\mathcal{O}}\mathscr{D}\|_{0}^{(\gamma)}} \Big(\|Ax\| + \kappa r^{\alpha} \big(1 + \kappa^{-1} \|A^{\mathcal{D}}\|_{0}^{(\alpha)}\big) \|x\| \Big).$$
(3.8)

In particular, B is infinitesimally A-bounded in $\ell^2(\Xi)$.

Proof. The only non-trivial case is $\alpha > \beta > 0$. For every $x \in H^{\alpha}$ we have

$$||Bx|| \le ||B\widetilde{A}_{\kappa,r}^{-1}|| ||A^{\mathscr{D}}x|| + ||B(\widetilde{A}_{\kappa,r}^{-1}A^{\mathscr{D}} - \mathrm{Id})|| ||x||,$$

$$\begin{split} \|B(\widetilde{A}_{\kappa,r}^{-1}A^{\mathscr{D}} - \mathrm{Id})\| &\leq \|B\|_{0}^{(\beta)} \|\widetilde{A}_{\kappa,r}^{-1}A^{\mathscr{D}} - \mathrm{Id}\|_{|\beta|}^{(-\beta)} \\ &\leq r^{\beta} \|B\|_{0}^{(\beta)} \left(1 + \kappa^{-1} \|A^{\mathscr{D}}\|_{0}^{(\alpha)}\right) \end{split}$$

and

$$\begin{split} \|B\widetilde{A}_{\kappa,r}^{-1}\| &\leq \|B\|_{0}^{(\beta)} \|\widetilde{A}_{\kappa,r}^{-1}\|_{0}^{(-\beta)} \\ &\leq \kappa^{-1} \|B\|_{0}^{(\beta)} r^{\beta-\alpha}, \end{split}$$

and we obtain

$$\|Bx\| \le \kappa^{-1} r^{\beta-\alpha} \|B\|_0^{(\beta)} \Big(\|A^{\mathscr{D}}x\| + \kappa r^{\alpha} \big(1 + \kappa^{-1} \|A^{\mathscr{D}}\|_0^{(\alpha)}\big) \|x\| \Big),$$
(3.9)

which is (3.8) with $A^{\mathcal{D}} \in \mathbf{DES}^{\alpha}$ replacing A. Applying (3.9) with $B = A^{\mathcal{OD}}$, we arrive at

$$\begin{split} \|A^{\mathcal{O}\mathcal{D}}x\| &\leq \kappa^{-1} r^{\gamma-\alpha} \|A^{\mathcal{O}\mathcal{D}}\|_{0}^{(\gamma)} \|A^{\mathcal{D}}x\| \\ &+ r^{\gamma} \|A^{\mathcal{O}\mathcal{D}}\|_{0}^{(\gamma)} (1+\kappa^{-1} \|A^{\mathcal{D}}\|_{0}^{(\alpha)}) \|x\| \end{split}$$

Hence we have

$$\begin{split} \|Ax\| &\geq \|A^{\mathcal{D}}x\| - \|A^{\mathcal{O}\mathcal{D}}x\| \\ &\geq \left(1 - \kappa^{-1}r^{\gamma-\alpha} \|A^{\mathcal{O}\mathcal{D}}\|_{0}^{(\gamma)}\right) \|A^{\mathcal{D}}x\| - r^{\gamma} \|A^{\mathcal{O}\mathcal{D}}\|_{0}^{(\gamma)} \left(1 + \kappa^{-1} \|A^{\mathcal{D}}\|_{0}^{(\alpha)}\right) \|x\|, \end{split}$$

which implies

$$\|A^{\mathscr{D}}x\| \le \left(1 - \kappa^{-1} r^{\gamma - \alpha} \|A^{\mathscr{O}}\|_{0}^{(\gamma)}\right)^{-1} \left(\|Ax\| + r^{\gamma} \|A^{\mathscr{O}}\|_{0}^{(\gamma)} \left(1 + \kappa^{-1} \|A^{\mathscr{D}}\|_{0}^{(\alpha)}\right) \|x\|\right).$$
(3.10)

Substituting (3.10) into (3.9) we obtain (3.8).

We conclude the section with the following proposition.

Proposition 3.7. For $\alpha \in \mathbb{R}$, every operator from \mathbf{ES}^{α} is closed on $\mathrm{H}^{\max\{\alpha,0\}}$ in the Hilbert space $\ell^{2}(\Xi)$. Every symmetric operator from \mathbf{ES}^{α} defined on $\mathrm{H}^{\max\{\alpha,0\}}$ is self-adjoint.

Proof. For $\alpha \leq 0$ we have $\mathbf{ES}^{\alpha} \subset \mathbf{S}^{0}$, and the statements follow from Corollary 2.9. Now assume $A \in \mathbf{SES}^{\alpha}$ with $\alpha > 0$. By (2.1), Definition 3.3 and Lemma 2.8, for any $(\kappa, r) \in EP(A^{\textcircled{D}})$ we have the estimates

$$\kappa^2 \|x\|_{\mathrm{H}^{\alpha}}^2 \leq \|\widetilde{A}_{\kappa,r}x\|^2 \leq \left(\|\widetilde{A}_{\kappa,r}\|_{\alpha}^{(\alpha)}\right)^2 \|x\|_{\mathrm{H}^{\alpha}}^2$$

for all $x \in H^{\alpha}$. Hence the graph norm of $\widetilde{A}_{\kappa,r}$ is equivalent to the norm of H^{α} , and $\widetilde{A}_{\kappa,r}$ is closed on H^{α} . If $\widetilde{A}_{\kappa,r}$ is symmetric, then for every $x \in \text{dom}(\widetilde{A}^*_{\kappa,r})$ there exists $C_x \ge 0$ such that for all $y \in H^{\alpha}$

$$|(x, \widetilde{A}_{\kappa,r} y)| \le C_x \|y\|_{\ell^2(\Xi)}.$$

In particular, with $(y_n)_{\xi} := \overline{\widetilde{a}_{\kappa,r,id}(\xi)} \mathbf{1}_{\langle \xi \rangle \leq n} x_{\xi}$ for $n \geq r, \xi \in \Xi$, we obtain by (3.3) and (3.4) that

$$\sum_{\substack{\xi \in \Xi \\ \langle \xi \rangle \le n}} \langle \xi \rangle^{2\alpha} |x_{\xi}|^{2} \le \kappa^{-1}(x, \widetilde{A}_{\kappa, r} y_{n}) \le \kappa^{-1} C_{x} ||y_{n}||_{\ell^{2}(\Xi)}$$
$$\le \kappa^{-1} C_{x} \left\| \widetilde{A}_{\kappa, r} \right\|_{0}^{(\alpha)} \left(\sum_{\substack{\xi \in \Xi \\ \langle \xi \rangle \le n}} \langle \xi \rangle^{2\alpha} |x_{\xi}|^{2} \right)^{1/2}.$$

Passing to the limit $n \to \infty$, it follows from (2.1) that $||x||_{H^{\alpha}} \le \kappa^{-1} C_x ||\widetilde{A}_{\kappa,r}||_0^{(\alpha)}$, i.e.

 $\operatorname{dom}(\widetilde{A}_{\kappa,r}^*) \subset \operatorname{H}^{\alpha} = \operatorname{dom}(\widetilde{A}_{\kappa,r}),$

hence $\widetilde{A}_{\kappa,r}$ is self-adjoint. By Lemma 3.6 $A - \widetilde{A}_{\kappa,r}$ is infinitesimally $\widetilde{A}_{\kappa,r}$ -bounded, so that A is also self-adjoint (see, e.g., Theorems 3.4.2 and 4.1.9 in [BS87]).

For $A \in \mathbf{ES}^{\alpha}$, by Definition 3.4 there exist a unitary $U \in \mathbf{S}^{0}$ and $H \in \mathbf{SES}^{\alpha}$ such that $A = UHU^{\dagger}$. Moreover, it follows from Lemma 2.8 that $UH^{\alpha} = U^{\dagger}H^{\alpha} = H^{\alpha}$. Now, let $(x_{n})_{n \in \mathbb{N}} \subset H^{\alpha}$ with $x_{n} \to x$ and $U^{\dagger}HUx_{n} \to z$ in ℓ^{2} , as $n \to \infty$. Since U is bounded, $Ux_{n} \to Ux$ and $HUx_{n} \to Uz$, thus the closedness of H implies that $Ux \in H^{\alpha}$ and HUx = Uz, i.e. $x \in H^{\alpha}$ and $UHU^{\dagger}x = z$. Hence, A is closed on H^{α} and self-adjoint if symmetric.

4. The Density of States Measure and von Neumann Algebras

In this section, following [Shu79a], we consider a representation of S^{∞} into another operator algebra, affiliated with an infinite factor (accounting for the almost periodicity), and define the density of states measure (DSM) for self-adjoint operators in ES^{∞} with respect to this representation. For a suitable representation, this DSM will coincide with the classically defined DSM on elliptic differential operators with almost periodic coefficients. We follow the construction and terminology of [Shu79a, §1], generalising Shubin's symbol classes to the ones defined in Section 2.

4.1. **Representations of the operator algebra.** Let \mathfrak{H} be a Hilbert space and \mathfrak{A} a factor of either type I_{∞} or II_{∞} in $\mathscr{B}(\mathfrak{H})$, the algebra of bounded linear operators in \mathfrak{H} . Let \widetilde{H}^{∞} be a dense subspace of \mathfrak{H} and

$$\widetilde{\mathbf{S}}^{\infty} = \bigcup_{\gamma \in \mathbb{R}} \widetilde{\mathbf{S}}^{\gamma}$$

a *-algebra of unbounded linear operators in \mathfrak{H} defined on \widetilde{H}^{∞} , filtered by \mathbb{R} . We assume that $\widetilde{S}^{\infty}\widetilde{H}^{\infty} \subset \widetilde{H}^{\infty}$, and that \widetilde{S}^{∞} is invariant under the involution $\widetilde{A} \mapsto \widetilde{A}^{\dagger} := \widetilde{A}^{*}|_{\widetilde{H}^{\infty}}$, where \widetilde{A}^{*} is the adjoint to $\widetilde{A} \in \widetilde{S}^{\infty}$. We also suppose that \widetilde{S}^{∞} is affiliated with the factor \mathfrak{A} , denoted $\widetilde{S}^{\infty}\eta\mathfrak{A}$. Finally, we consider a representation $\rho : S^{\infty} \to \widetilde{S}^{\infty}$ having the following properties:

(i) ρ is a homomorphism of filtered *-algebras with $\rho(\mathbf{S}^{\gamma}) \subset \widetilde{\mathbf{S}}^{\gamma}$, for all $\gamma \in \mathbb{R}$.

(ii) For every $A \in S^0$, $\rho(A)$ extends to a bounded linear operator on \mathfrak{H} with

$$\|\rho(A)\|_{\mathfrak{H}\to\mathfrak{H}} = \|A\|_{\ell^{2}(\Xi)\to\ell^{2}(\Xi)}.$$
(4.1)

- (iii) For all $A \in \mathbf{S}^{\infty}$, $\rho(A)$ is closable in \mathfrak{H} with the closure $A^{\sharp} := \overline{\rho(A)}$. For every $\alpha > 0$ there exists a dense subspace $\widetilde{H}^{\alpha} \supset \widetilde{H}^{\infty}$ such that
 - (a) $\widetilde{H}^{\alpha} \subset \widetilde{H}^{\gamma}$ if $0 < \gamma \le \alpha$,
 - (b) $A \in \mathbf{DES}^{\alpha}$ implies dom $(A^{\sharp}) = \widetilde{H}^{\alpha}$,
 - (c) for all $B \in \mathbf{S}^0$, $B^{\sharp} \widetilde{H}^{\alpha} \subset \widetilde{H}^{\alpha}$.
- (iv) If $A \in \mathbf{DES}^{\alpha}$, $\alpha > 0$, is self-adjoint on H^{α} , then A^{\sharp} is self-adjoint.

Remark 4.1. When \mathfrak{A} is a I_{∞} factor some of the statements in this section become rather trivial. However, we include this case for applications in Section 10.

Remark 4.2. In [Shu79a], Shubin considers $G = \mathbb{R}^d$ acting on itself by translation, with almost periodic operators acting both in Besicovitch space $B^2(\mathbb{R}^d) \cong \ell^2(\mathbb{R}^d)$ and in $L^2(\mathbb{R}^d)$ through the Fourier integral representation of pseudo-differential operators. The appropriate Hilbert space is then

$$\mathfrak{H} = \mathrm{B}^2(\mathbb{R}^d) \otimes \mathrm{L}^2(\mathbb{R}^d),$$

and the II_{∞} factor \mathfrak{A} is generated by the two families of operators

$$\left\{ \mathbf{e}_{\boldsymbol{\xi}} \otimes \mathbf{e}_{\boldsymbol{\xi}} : \boldsymbol{\xi} \in \mathbb{R}^d \right\}$$
 and $\left\{ I \otimes T_{\boldsymbol{\xi}} : \boldsymbol{\xi} \in \mathbb{R}^d \right\}$,

where $\mathbf{e}_{\boldsymbol{\xi}}$ is multiplication by the character $\mathbf{e}_{\boldsymbol{\xi}}(\mathbf{x}) = e^{i\boldsymbol{\xi}\cdot\mathbf{x}}$ and $T_{\boldsymbol{\xi}}$ is the translation operator $T_{\boldsymbol{\xi}}f(\mathbf{x}) = f(\mathbf{x}-\boldsymbol{\xi})$. The representation ρ is given on $A = Op(a) \in \mathbf{S}^{\infty}$ by the linear operator $\rho(A) = a(\mathbf{x}+\mathbf{y}; D_{\mathbf{y}})$ acting on

$$\widetilde{\mathbf{H}}^{\infty} := \mathbf{B}^2(\mathbb{R}^d) \otimes \widehat{\mathbf{H}}^{\infty}(\mathbb{R}^d).$$

Here, **x** is the variable of functions in B²(\mathbb{R}^d), **y** is the variable of functions in L²(\mathbb{R}^d), $D_y = -i\nabla_y$, and $\hat{H}^{\infty}(\mathbb{R}^d) := \{f \in C^{\infty}(\mathbb{R}^d) : \partial^{\alpha} f \in L^2(\mathbb{R}^d) \text{ for all } \alpha \in \mathbb{N}_0^d\}.$

Properties (i) and (ii) of the representation ρ imply the following lemma.

Lemma 4.3. If $A \in \mathbf{S}^0$, then A^{\sharp} is defined on \mathfrak{H} and satisfies $(A^{\sharp})^* = (A^{\dagger})^{\sharp} = (A^*)^{\sharp}$ and $||A^{\sharp}||_{\mathfrak{H} \to \mathfrak{H}} = ||A||_{\ell^2(\Xi) \to \ell^2(\Xi)}$. In particular, the map $\mathbf{S}^0 \to \mathfrak{B}(\mathfrak{H})$, $A \mapsto A^{\sharp}$ is an injective homomorphism of *-algebras. If $U \in \mathbf{S}^0$ is unitary, then so is U^{\sharp} .

We will now carry over Lemma 3.6 to images under \sharp . This provides us with some information on the domains of operators from $(\mathbf{S}^{\infty})^{\sharp}$.

Lemma 4.4. Let $\beta \in \mathbb{R}$, $B \in S^{\beta}$ and $A \in SES^{\alpha}$ for some $\alpha > 0$. Then (1) $\bigcup_{\zeta > \max\{\beta, 0\}} \widetilde{H}^{\zeta} \subset \operatorname{dom}(B^{\sharp})$,

- (2) dom $(A^{\sharp}) = \widetilde{H}^{\alpha}$.
- (3) Suppose $\beta < \alpha$ and $0 < \gamma < \alpha$ with $A^{\emptyset \mathscr{D}} \in \mathbf{S}^{\gamma}$. Then for $\beta \leq 0$ the operator B^{\sharp} is bounded, and, otherwise, for every $\varphi \in \widetilde{H}^{\alpha}$ and

$$(\kappa,r)\in \operatorname{EP}(A^{\mathcal{D}})\cap \left\{r\geq \left(\|A^{\mathcal{OD}}\|_0^{(\gamma)}/\kappa\right)^{1/(\alpha-\gamma)}\right\},$$

we have

$$\|B^{\sharp}\varphi\|_{\mathfrak{H}} \leq \frac{r^{\beta-\alpha} \|B\|_{0}^{(\beta)}}{\kappa - r^{\gamma-\alpha} \|A^{\mathscr{O}\mathscr{D}}\|_{0}^{(\gamma)}} \Big(\|A^{\sharp}\varphi\|_{\mathfrak{H}} + \kappa r^{\alpha} \big(1 + \kappa^{-1} \|A^{\mathscr{D}}\|_{0}^{(\alpha)}\big) \|\varphi\|_{\mathfrak{H}} \Big).$$
(4.2)

In particular, B^{\sharp} is infinitesimally A^{\sharp} -bounded in \mathfrak{H} .

Proof. For $\beta \leq 0$ the statements (1) and (3) follow from (4.1). Let now $\beta > 0$ and assume that $0 < \gamma < \alpha$ with $A^{\oslash \mathscr{D}} \in \mathbf{S}^{\gamma}$. Following the proof of Lemma 3.6 and applying properties (i) and (ii) of the representation ρ where necessary, we derive (4.2) for $\varphi \in \widetilde{H}^{\infty}$. Consequently, the graph norm of $\rho(B)$ is dominated by the graph norm of $\rho(A)$, thus dom $(B^{\sharp}) \supset \text{dom}(A^{\sharp})$. Applying (4.2) for $A^{\oslash \mathscr{D}}$ instead of *B* and $A^{\mathscr{D}}$ instead of *A*, we conclude that the graph norms of $\rho(A)$ and $\rho(A^{\mathscr{D}})$ are equivalent, thus (iiib) implies dom $(A^{\sharp}) = \text{dom}((A^{\mathscr{D}})^{\sharp}) = \widetilde{H}^{\alpha}$, which is (2). Now (1) follows by varying $A \in \mathbf{SES}^{\alpha}$ with $\alpha > \beta$. Finally, we can extend (4.2) from \widetilde{H}^{∞} to \widetilde{H}^{α} by density with respect to the graph norm of A^{\sharp} .

Properties (iiib) and (iv) of the map ρ can also be extended to operators from the classes **ES**^{α}, $\alpha > 0$.

Lemma 4.5. Let
$$\alpha > 0$$
 and $A \in \mathbf{ES}^{\alpha}$. Then $\operatorname{dom}(A^{\sharp}) = \widetilde{H}^{\alpha}$ and for all unitary $U \in \mathbf{S}^{0}$

$$U^{\sharp}A^{\sharp}(U^{\sharp})^{*} = (UAU^{\dagger})^{\sharp} \qquad holds \ on \ \widetilde{H}^{\alpha}.$$

$$(4.3)$$

Moreover, if A is self-adjoint on H^{α} , then A^{\sharp} is self-adjoint.

Proof. Assume first that $A \in \mathbf{SES}^{\alpha}$, so that $A^{\mathcal{D}} \in \mathbf{DES}^{\alpha}$ and $A^{\mathcal{D}\mathcal{D}} \in \mathbf{S}^{\gamma}$ for some $0 < \gamma < \alpha$. According to Lemma 4.4(1,2) we have that dom $(A^{\sharp}) = \widetilde{\mathbf{H}}^{\alpha} \subset \operatorname{dom}((A^{\mathcal{D}\mathcal{D}})^{\sharp})$. Moreover, if A is self-adjoint, then $(A^{\mathcal{D}})^{\sharp}$ is self-adjoint on $\widetilde{\mathbf{H}}^{\alpha}$ and $\rho(A^{\mathcal{D}\mathcal{D}})$ is symmetric on $\widetilde{\mathbf{H}}^{\infty}$, as follows from properties (iv) and (i) of ρ , respectively. Since dom $((A^{\mathcal{D}\mathcal{D}})^{\sharp}) \supset \widetilde{\mathbf{H}}^{\alpha}$ is the closure of $\widetilde{\mathbf{H}}^{\infty}$ with respect to the graph norm of $(A^{\mathcal{D}\mathcal{D}})^{\sharp}$, the operator $(A^{\mathcal{D}\mathcal{D}})^{\sharp}$ is also symmetric on $\widetilde{\mathbf{H}}^{\alpha}$. Moreover, by Lemma 4.4(3) it is infinitesimally A^{\sharp} -bounded. Thus, [BS87, Theorem 4.1.9] implies that $A^{\sharp} = (A^{\mathcal{D}} + A^{\mathcal{D}\mathcal{D}})^{\sharp}$ is self-adjoint on $\widetilde{\mathbf{H}}^{\alpha}$.

Let now $A \in \mathbf{ES}^{\alpha}$. By definition, there exist $H \in \mathbf{SES}^{\alpha}$ and $V \in \mathbf{S}^{0}$ unitary such that $A = V^{\dagger}HV$ on H^{∞} . Since ρ is a *-homomorphism,

$$\rho(A) = \rho(V)^{\dagger} \rho(H) \rho(V) \tag{4.4}$$

holds on \widetilde{H}^{∞} . By Lemma 4.3 the operator V^{\sharp} is unitary, and property (iiic) implies that

$$V^{\sharp}\widetilde{H}^{\alpha} = \left(V^{\sharp}\right)^{*}\widetilde{H}^{\alpha} = \widetilde{H}^{\alpha}.$$
(4.5)

We have already proved in Lemma 4.4(2) that dom $(H^{\sharp}) = \tilde{H}^{\alpha}$, thus the argument at the end of the proof of Proposition 3.7 implies that $(V^{\sharp})^* H^{\sharp} V^{\sharp}$ is closed on \tilde{H}^{α} . As by (4.4) and Lemma 4.3 this operator is an extension of $\rho(A)$, it follows that dom $(A^{\sharp}) \subset \tilde{H}^{\alpha}$. Similarly, we have on \tilde{H}^{∞} that

$$\rho(H) = \rho(V)\rho(A)\rho(V)^{\dagger}$$

and $V^{\sharp}A^{\sharp}(V^{\sharp})^*$ is a closed operator on $V^{\sharp} \operatorname{dom}(A^{\sharp}) \subset \widetilde{H}^{\infty}$. Thus,

$$\widetilde{\mathrm{H}}^{\alpha} = \mathrm{dom}(H^{\sharp}) \subset V^{\sharp} \mathrm{dom}(A^{\sharp}),$$

and (4.5) yields $\widetilde{H}^{\alpha} \subset \operatorname{dom}(A^{\sharp})$. Hence $\operatorname{dom}(A^{\sharp}) = \widetilde{H}^{\alpha}$. Finally, let $U \in \mathbf{S}^{0}$ be unitary. Then

$$\rho(U)\rho(A)\rho(U)^{\dagger} = \rho(UAU^{\dagger}) \subset (UAU^{\dagger})^{\sharp}$$

and $U^{\sharp}A^{\sharp}(U^{\sharp})^{*}$ is a closed extension of $\rho(U)\rho(A)\rho(U)^{\dagger}$ on the domain $\widetilde{H}^{\alpha} = \operatorname{dom}((UAU^{\dagger})^{\sharp})$, i.e. (4.3) holds. If *A* is self-adjoint on H^{α} , then so is *H*, thus H^{\sharp} by the first part of the proof. Hence, the self-adjointness of A^{\sharp} follows from (4.3) with U = V and *H* instead of *A*.

4.2. The density of states measure. Since \mathfrak{A} is a factor of type I_{∞} or II_{∞} , there exists, by definition, a semi-finite faithful normal trace \mathfrak{T} on \mathfrak{A} , see [Dix81, I.6 and I.8.4]. Moreover, due to [Dix81, I.6.4, Corollary], this trace is unique up to multiplication by a positive number. Following the notation of [Naĭ72], we write $\mathfrak{L}\eta\mathfrak{A}$ to denote that $\mathfrak{L} \subset \mathfrak{H}$ is a closed linear subspace adjoint to \mathfrak{A} , i.e. $P_{\mathfrak{L}} \in \mathfrak{A}$, where $P_{\mathfrak{L}}$ is the projection onto \mathfrak{L} . If $\mathfrak{L}\eta\mathfrak{A}$, the relative dimension of \mathfrak{L} is defined by

$$\mathfrak{D}(\mathfrak{L}) := \mathfrak{T}(P_{\mathfrak{L}}) \in [0,\infty]$$

If \mathfrak{A} is a I_{∞} -factor, the range of the relative dimension is $c\mathbb{N}_0 \cup \{\infty\}$, for some c > 0. It is $[0, \infty]$ if \mathfrak{A} is a II_{∞} -factor.

Definition 4.6. Let $A \in \mathbf{S}^0 \cup \mathbf{ES}^\infty$ be symmetric and $J \subset \mathbb{R}$ be a Borel measurable set. Denote by $E_J(A^{\sharp})$ the spectral projection of A^{\sharp} for J. We define the *density of states measure* (DSM) of A on J, relative to the representation ρ , by

$$N(J;A) := \mathfrak{T}(E_I(A^{\sharp})) = \mathfrak{D}(E_I(A^{\sharp})\mathfrak{H}).$$

Remark 4.7. Usually, the dependence on the representation ρ and the factor \mathfrak{A} is unambiguous and is thus not reflected in the notation.

The following corollary generalises [PS12, Lemma 4.4]. It follows directly from Lemma 4.5 (or Lemma 4.3 for $A \in \mathbf{S}^0$) and the invariance of \mathfrak{T} under unitary transformations in \mathfrak{A} . We remark at

this point that, since $\tilde{\mathbf{S}}^{\infty}\eta\mathfrak{A}$, one has $U^{\sharp} \in \mathfrak{A}$ for every unitary $U \in \mathbf{S}^{0}$, see Lemma 4.3 and [Naĭ72, §35.1].

Corollary 4.8. Let $U \in S^0$ be unitary and let $A \in S^0 \cup ES^\infty$ be symmetric. Then one has $N(J; A) = N(J; UAU^{\dagger})$ for any Borel measurable set $J \subset \mathbb{R}$.

In the remainder of this section, we investigate the behaviour of the DSM for elliptic operators of positive order under perturbations. In [PS12, MPS14, PS16] such an analysis was conducted for operators that are bounded from below and the particular case $J = (-\infty, \lambda), \lambda \in \mathbb{R}$.

Before continuing, let us introduce the following notation. For any interval $J = [s, t] \subset \mathbb{R}$, s < t and $\varepsilon \in \mathbb{R}$, we define

$$J_{\varepsilon} := \begin{cases} \varnothing & \text{for } \varepsilon < \frac{s-t}{2} \\ [s-\varepsilon, t+\varepsilon] & \text{otherwise.} \end{cases}$$

The following lemma gives us a variational characterisation of the DSM (cf. [PS12, Lemma 4.1]).

Lemma 4.9. Let $A \in S^0 \cup ES^\infty$ be symmetric. Then, for any interval J = [q - r, q + r] with $q \in \mathbb{R}$ and r > 0, we have

$$N(J;A) = \sup \left\{ \mathfrak{D}(\mathfrak{L}) : \mathfrak{L} \subset \operatorname{dom}(A^{\sharp}), \ \mathfrak{L}\eta\mathfrak{A}, \\ \operatorname{and} \| (A^{\sharp} - q)\varphi \|_{\mathfrak{H}} \le r \|\varphi\|_{\mathfrak{H}} \ \forall \varphi \in \mathfrak{L} \right\}.$$

$$(4.6)$$

The analogous statement holds for the open interval J = (q - r, q + r) with strict inequality in (4.6).

Remark 4.10. Usually, variational characterisations such as (4.6) are given in terms of quadratic forms rather than norms. The reason why we cannot do so is because we do not assume the operator *A* to be semi-bounded, *J* a semi-infinite interval. One can interpret Lemma 4.9 in terms of quadratic forms as usual for the nonnegative operator $(A - q)^2$.

Proof. Choosing $\mathfrak{L} := E_J(A^{\sharp})\mathfrak{H}$, we observe that N(J; A) is at most the right hand side of (4.6). Suppose that there exists a subspace \mathfrak{L} that satisfies the assumptions on the righthand side of (4.6) and $\mathfrak{D}(\mathfrak{L}) > \mathfrak{D}(E_J(A^{\sharp})\mathfrak{H})$. Then [Naĭ72, §37.1, Lemma] implies that \mathfrak{L} contains an element φ orthogonal to $E_J(A^{\sharp})\mathfrak{H}$, implying that $\|(A^{\sharp} - q)\varphi\|_{\mathfrak{H}}^2 > r^2 \|\varphi\|_{\mathfrak{H}}^2$, which is a contradiction. \Box

The following lemma generalises [PS12, Corollary 4.3] to operators that are not necessarily bounded below and unbounded perturbations.

Lemma 4.11. Let $A \in SES^{\alpha}$, $\alpha > 0$, and $B \in S^{\beta}$, $\beta < \alpha$, symmetric operators. Let $J := [q-r, q+r] \subset \mathbb{R}$ be the interval of length 2r > 0 centred at $q \in \mathbb{R}$. Then there exists a constant $C \ge 0$ depending only

on A and β such that, for

$$\varepsilon := \varepsilon_{J,A,B} := \begin{cases} \|B\| & \text{if } \beta \le 0, \\ \frac{\|B\|_{0}^{(\beta)}}{2 + \|B\|_{0}^{(\beta)}} \left(r + |q| + C\left(1 + \|B\|_{0}^{(\beta)}\right)^{\frac{\alpha}{\alpha - \beta}}\right) & \text{if } \beta > 0, \end{cases}$$
(4.7)

the inequality

$$N(J_{-\varepsilon}; A) \le N(J; A + B) \le N(J_{\varepsilon}; A)$$
(4.8)

holds.

Proof. In view of Lemma 4.4(1,3) and property (i) of the representation ρ , one has that $(A+B)^{\sharp} = A^{\sharp} + B^{\sharp}$ on dom $(A^{\sharp}) \subset \text{dom}(B^{\sharp})$. Fix $\varphi \in \mathfrak{L} := E_J((A+B)^{\sharp})\mathfrak{H} \subset \text{dom}((A+B)^{\sharp}) = \text{dom}(A^{\sharp})$, so that

$$\left\| (A^{\sharp} + B^{\sharp} - q)\varphi \right\|_{\mathfrak{H}} \le r \left\| \varphi \right\|_{\mathfrak{H}}.$$

$$(4.9)$$

We will show that

$$\left\| (A^{\sharp} - q)\varphi \right\|_{\mathfrak{H}} \le (r + \varepsilon) \left\| \varphi \right\|_{\mathfrak{H}}$$
(4.10)

holds, which in view of (4.6) implies the second inequality in (4.8). Since

$$\begin{split} \left\| (A^{\sharp} - q)\varphi \right\|_{\mathfrak{H}} &\leq \left\| (A^{\sharp} + B^{\sharp} - q)\varphi \right\|_{\mathfrak{H}} + \left\| B^{\sharp}\varphi \right\|_{\mathfrak{H}} \\ &\leq r \left\| \varphi \right\|_{\mathfrak{H}} + \left\| B^{\sharp}\varphi \right\|_{\mathfrak{H}}, \end{split}$$
(4.11)

it is sufficient to estimate $\|B^{\sharp}\varphi\|_{\mathfrak{H}}$. For $\beta \leq 0$, Lemma 4.3 and Corollary 2.9 imply

$$\left\|B^{\sharp}\varphi\right\|_{\mathfrak{H}} \leq \left\|B^{\sharp}\right\| \left\|\varphi\right\|_{\mathfrak{H}} = \left\|B\right\| \left\|\varphi\right\|_{\mathfrak{H}} = \varepsilon \left\|\varphi\right\|_{\mathfrak{H}},$$

and (4.10) follows from (4.9) and (4.11).

From now on, we consider $\beta > 0$. By assumption we can choose $\gamma \in (\beta, \alpha)$ such that $A^{\mathcal{OD}} \in \mathbf{S}^{\gamma}$. Let $(\kappa, r) \in EP(A^{\mathcal{D}})$ with

$$r \ge \max\left\{\left(\frac{4\left(1+\|B\|_{0}^{(\beta)}\right)}{\kappa}\right)^{1/(\alpha-\beta)}; \quad \left(\frac{2\|A^{\mathcal{OD}}\|_{0}^{(\gamma)}}{\kappa}\right)^{1/(\alpha-\gamma)}\right\}$$

As $\varphi \in \text{dom}(A^{\sharp})$, Lemma 4.4(3) yields

$$\begin{split} \left\| B^{\sharp} \varphi \right\|_{\mathfrak{H}} &\leq 2 \left\| B \right\|_{0}^{(\beta)} \left[r^{\beta - \alpha} \kappa^{-1} \left\| A^{\sharp} \varphi \right\|_{\mathfrak{H}} + r^{\beta} \left(1 + \kappa^{-1} \left\| A^{\mathscr{D}} \right\|_{0}^{(\alpha)} \right) \left\| \varphi \right\|_{\mathfrak{H}} \right] \\ &\leq \frac{\left\| B \right\|_{0}^{(\beta)}}{2} \left[\frac{\left\| (A^{\sharp} - q) \varphi \right\|_{\mathfrak{H}} + |q| \| \varphi \|_{\mathfrak{H}}}{1 + \| B \|_{0}^{(\beta)}} + C \left(1 + \| B \|_{0}^{(\beta)} \right)^{\frac{\beta}{\alpha - \beta}} \left\| \varphi \right\|_{\mathfrak{H}} \right], \end{split}$$
(4.12)

where *C* is a constant only depending on *A* and β . Combining (4.11) and (4.12), we get

$$\frac{2 + \|B\|_{0}^{(\beta)}}{2\left(1 + \|B\|_{0}^{(\beta)}\right)} \|(A^{\sharp} - q)\varphi\|_{\mathfrak{H}} \leq \left[r + \frac{\|B\|_{0}^{(\beta)}}{2\left(1 + \|B\|_{0}^{(\beta)}\right)} \left(|q| + C\left(1 + \|B\|_{0}^{(\beta)}\right)^{\frac{\alpha}{\alpha - \beta}}\right)\right] \|\varphi\|_{\mathfrak{H}}.$$

Hence, we arrive at (4.10) with ε as in (4.7).

For the first inequality in (4.8) the only non-trivial case is $\varepsilon \leq r$. For all $\varphi \in E_{J-\varepsilon}(A^{\sharp})\mathfrak{H} \subset \operatorname{dom}(A^{\sharp})$ we have

$$\left\| (A^{\sharp} - q)\varphi \right\|_{\mathfrak{H}} \leq (r - \varepsilon) \left\| \varphi \right\|_{\mathfrak{H}}$$

This implies

$$\left\| (A^{\sharp} + B^{\sharp} - q)\varphi \right\|_{\mathfrak{H}} \le (r - \varepsilon) \left\| \varphi \right\|_{\mathfrak{H}} + \left\| B^{\sharp}\varphi \right\|_{\mathfrak{H}},$$

where in view of (4.12) and (4.7)

$$\left\| B^{\sharp} \varphi \right\|_{\mathfrak{H}} \leq \frac{\|B\|_{0}^{(\beta)}}{2+2 \|B\|_{0}^{(\beta)}} \left(r + \left| q \right| + C \left(1 + \|B\|_{0}^{(\beta)} \right)^{\frac{\alpha}{\alpha-\beta}} \right) \left\| \varphi \right\|_{\mathfrak{H}} \leq \varepsilon \left\| \varphi \right\|_{\mathfrak{H}}.$$

Thus, the first inequality in (4.8) follows and the lemma is proved.

The next lemma deals with perturbations that are 'spectrally far' from a given interval. It is a generalisation of [MPS14, Lemma 11.1] for operators that are not necessarily bounded below.

Lemma 4.12. For $\alpha > 0$, $\beta < \alpha$ let $H_0 \in \mathbf{DES}^{\alpha}$, $B \in \mathbf{S}^{\beta}$, and $A \in \mathbf{S}^0$ be symmetric operators and set $H := H_0 + B \in \mathbf{SES}^{\alpha}$. Suppose that there exists a family of orthogonal projections $\{P_l\}_{l=0}^L$ with $P_l \in \mathbf{S}^{-\alpha}$, $0 \le l \le L-1$, and $P_L \in \mathbf{S}^0$ that all commute with H_0 and satisfy

$$\sum_{l=0}^{L} P_l = I, \quad and \quad A = AP_0, \quad B_{n,l} := P_n BP_l = 0, \quad for \ |n-l| > 1.$$
(4.13)

Moreover, let J = (q - r, q + r) *be an interval such that*

$$D_l := \operatorname{dist}\left(J, \sigma\left((P_l H P_l)^{\sharp}\right)\right) > 0, \quad \text{for all } 0 \le l < L.$$

$$(4.14)$$

Finally, assume that

$$3^L r \ge d_L := \min_{1 \le l < L} D_l \tag{4.15}$$

and

$$\max_{0 \le l < L} (\|B_{l,l-1}\| + \|B_{l,l+1}\|) / D_l \le 1/4,$$
(4.16)

where we use the convention $B_{0,-1} := 0$.

т

Then for

$$\varepsilon := 3^{2 - \frac{L}{2}} \left(\frac{r}{d_L} \right)^{1/2} \|A\|$$
(4.17)

we have that

$$N(J_{-\varepsilon}; H) \le N(J; H + A) \le N(J_{\varepsilon}; H).$$

Proof. We only prove the first inequality; the second inequality follows analogously. It suffices to show that for any $\varphi \in E_{J_{-\epsilon}}(H^{\sharp})\mathfrak{H} \subset \operatorname{dom}(H^{\sharp}) = \widetilde{H}^{\alpha}$, one has

$$\left\| (H^{\sharp} + A^{\sharp} - q)\varphi \right\|_{\mathfrak{H}} \le r \left\| \varphi \right\|_{\mathfrak{H}}.$$

For any $K \in \mathbb{N}$, we split the interval $J_{-\varepsilon}$ into 2K + 1 subintervals of equal width: for $-K \le k \le K - 1$, set

$$I_k := \left(q + (2k-1)\frac{(r-\varepsilon)}{2K+1}, q + (2k+1)\frac{(r-\varepsilon)}{2K+1} \right]$$

and

$$I_K := \left(q + (2K-1)\frac{(r-\varepsilon)}{2K+1}, q+r-\varepsilon\right).$$

For $\varphi \in E_{J_{-\varepsilon}}(H^{\sharp})\mathfrak{H}$ and $-K \leq k \leq K$ define $\varphi^k := E_{I_k}(H^{\sharp})\varphi \in \widetilde{H}^{\alpha}$ and

$$\eta^{k} := H^{\sharp} \varphi^{k} - \left(q + 2k \frac{(r-\varepsilon)}{2K+1} \right) \varphi^{k}, \qquad (4.18)$$

so that

$$\left\|\eta^{k}\right\|_{\mathfrak{H}} \leq \frac{r}{(2K+1)} \left\|\varphi^{k}\right\|_{\mathfrak{H}}$$

$$(4.19)$$

holds. We also introduce

$$\varphi_l^k := P_l^{\sharp} \varphi_k$$
 and $\eta_l^k := P_l^{\sharp} \eta_k$, for $-K \le k \le K$ and $0 \le l \le L$.

For $0 \le l < L$, we clearly have $P_l^{\sharp} H^{\sharp} = (P_l H)^{\sharp}$ on \widetilde{H}^{∞} and, since $P_l H \in \mathbf{S}^0$, this identity extends to \widetilde{H}^{α} . Moreover, P_l commutes with H_0 so that (4.13) implies that on \widetilde{H}^{α}

$$P_l^{\sharp} H^{\sharp} = (P_l H)^{\sharp} = (P_l H P_l)^{\sharp} + B_{l,l-1}^{\sharp} + B_{l,l+1}^{\sharp}$$
$$= (P_l H P_l)^{\sharp} P_l^{\sharp} + B_{l,l-1}^{\sharp} P_{l-1}^{\sharp} + B_{l,l+1}^{\sharp} P_{l+1}^{\sharp}$$

where we use the convention $P_{-1} := 0$. Thus, applying P_l^{\sharp} to (4.18), we arrive at

$$\eta_{l}^{k} = B_{l,l-1}^{\sharp} \varphi_{l-1}^{k} + \left((P_{l} H P_{l})^{\sharp} - \left(q + 2k \frac{(r-\varepsilon)}{2K+1} \right) \right) \varphi_{l}^{k} + B_{l,l+1}^{\sharp} \varphi_{l+1}^{k}$$

for $0 \le l < L$, and Lemma 4.3 together with (4.14) and (4.19) gives for $0 \le l < L$,

$$\begin{split} \left\| \varphi_{l}^{k} \right\|_{\mathfrak{H}} &\leq D_{l}^{-1} \Big(\left\| \eta_{l}^{k} \right\|_{\mathfrak{H}} + \left\| B_{l,l-1} \right\| \left\| \varphi_{l-1}^{k} \right\|_{\mathfrak{H}} + \left\| B_{l,l+1} \right\| \left\| \varphi_{l+1}^{k} \right\|_{\mathfrak{H}} \Big) \\ &\leq \frac{r}{(2K+1)d_{L}} \left\| \varphi^{k} \right\|_{\mathfrak{H}} + \frac{\left\| \varphi_{l-1}^{k} \right\|_{\mathfrak{H}} + \left\| \varphi_{l+1}^{k} \right\|_{\mathfrak{H}}}{4}. \end{split}$$

Recursively for $0 \le l < L$ we deduce that

$$\left\|\varphi_{l}^{k}\right\|_{\mathfrak{H}} \leq \frac{2r}{(2K+1)d_{L}}\left\|\varphi^{k}\right\|_{\mathfrak{H}} + \frac{1}{3}\left\|\varphi_{l+1}^{k}\right\|_{\mathfrak{H}}.$$

Hence, employing the trivial bound $\|\varphi_L^k\|_{\mathfrak{H}} \leq \|\varphi^k\|_{\mathfrak{H}}$, we get that

$$\left\|\varphi_0^k\right\|_{\mathfrak{H}} \leq \left(\frac{3r}{(2K+1)d_L} + 3^{-L}\right) \left\|\varphi^k\right\|_{\mathfrak{H}}.$$

In view of Lemma 4.3, it follows that for all $-K \le k \le K$,

$$\begin{aligned} A^{\sharp} \varphi^{k} \Big\|_{\mathfrak{H}} &= \left\| (AP_{0})^{\sharp} \varphi^{k} \right\|_{\mathfrak{H}} \\ &= \left\| A^{\sharp} \varphi_{0}^{k} \right\|_{\mathfrak{H}} \\ &\leq \left(\frac{3r}{(2K+1)d_{L}} + 3^{-L} \right) \|A\| \left\| \varphi^{k} \right\|_{\mathfrak{H}}, \end{aligned}$$

whence the Cauchy-Schwarz inequality and the Pythagorean theorem yield

$$\begin{split} \left\| A^{\sharp} \varphi \right\|_{\mathfrak{H}} &\leq \sum_{-K \leq k \leq K} \left\| A^{\sharp} \varphi^{k} \right\|_{\mathfrak{H}} \\ &\leq \left(\frac{3r}{(2K+1)d_{L}} + 3^{-L} \right) \sqrt{2K+1} \left\| A \right\| \left\| \varphi \right\|_{\mathfrak{H}}. \end{split}$$

We choose

$$K = \left\lfloor \frac{3^{L+1}r}{2d_L} - \frac{1}{2} \right\rfloor + 1,$$

so that $\frac{3^{L+1}r}{d_L} \le 2K + 1 \le \frac{3^{L+2}r}{d_L}$. Then, by (4.15), we have

$$\left(\frac{3r}{(2K+1)d_L} + 3^{-L}\right)\sqrt{2K+1} \le 3^{2-\frac{L}{2}} \left(\frac{r}{d_L}\right)^{1/2}.$$

Consequently, we arrive at

$$\begin{split} \left\| (H^{\sharp} + A^{\sharp} - q)\varphi \right\|_{\mathfrak{H}} &\leq \left\| (H^{\sharp} - q)\varphi \right\|_{\mathfrak{H}} + \left\| A^{\sharp}\varphi \right\|_{\mathfrak{H}} \\ &\leq \left((r - \varepsilon) + 3^{2 - \frac{L}{2}} \left(\frac{r}{d_L} \right)^{1/2} \|A\| \right) \|\varphi\|_{\mathfrak{H}} \\ &= r \left\| \varphi \right\|_{\mathfrak{H}}, \end{split}$$

where we used that $\varphi \in E_{J_{-\varepsilon}}(H^{\sharp})$ and the value of ε given in (4.17).

5. GAUGE TRANSFORM

Let $\alpha \in \mathbb{R}$ and $A = Op(\alpha) \in SES^{\alpha}$ be symmetric, thus extends to a self-adjoint linear operator on H^{α} by Proposition 3.7.

Definition 5.1. For every symmetric $\Psi \in \mathbf{S}^0$, the unitary transformation of *A* into

$$[A] := [A]_{\Psi} := \exp(-i\Psi)A\exp(i\Psi)$$

is called a gauge transform.

We remark here that, due to Lemma 2.13 and Corollary 2.9, the series

$$\exp(i\Psi) = \sum_{k=0}^{\infty} \frac{\left(i\Psi\right)^k}{k!}$$
(5.1)

converges both in \mathbf{S}^0 and in the operator norm. In particular, Lemma 2.3 implies that $\exp(i\Psi) \in \mathbf{S}^0$, whence $\exp(i\Psi)$ is unitary and $[A]_{\Psi} \in \mathbf{ES}^{\alpha}$ is symmetric. The following lemma provides an expansion of $[A]_{\Psi}$ into a series of multiple commutators of A with Ψ , see (2.19) for the definition of ad^k .

Lemma 5.2. We have

$$[A]_{\Psi} = \sum_{k=0}^{\infty} \frac{1}{k!} \operatorname{ad}^{k}(A; \Psi),$$
 (5.2)

where the series converges absolutely in S^{α} .

Proof. Lemma 2.13 yields the bounds

$$\left\| \Psi^{j} A \Psi^{m} \right\|_{l}^{(\alpha)} \leq \left(\left\| \Psi \right\|_{l}^{(0)} \right)^{j} \left\| A \right\|_{l}^{(\alpha)} \left(\left\| \Psi \right\|_{l+|\alpha|}^{(0)} \right)^{m}, \text{ for all } l \geq 0.$$

Thus, the double series

$$[A]_{\Psi} = \sum_{j=0}^{\infty} \frac{(-i\Psi)^{j}}{j!} A \sum_{m=0}^{\infty} \frac{(i\Psi)^{m}}{m!} = \sum_{j,m=0}^{\infty} \frac{(-i\Psi)^{j}}{j!} A \frac{(i\Psi)^{m}}{m!}$$

converges absolutely in S^{α} . Recursively, we obtain

$$\operatorname{ad}^{k}(A; \Psi) = k! \sum_{j+m=k} \frac{(-i\Psi)^{j}}{j!} A \frac{(i\Psi)^{m}}{m!}, \text{ for all } k \ge 0.$$

In the remainder of this section, we look at gauge transforms that result in an operator $[A]_{\Psi}$
that is closer to a diagonal operator (i.e. an operator in DES ^{α}) than A. More precisely, we con-
struct Ψ in such a way that the gauge transform removes as much of the off-diagonal part A^{OD}
from A as possible. Let $\beta < \alpha$ such that $A^{\mathcal{O}\mathcal{D}} \in \mathbf{S}^{\beta}$. Then we aim at

$$[A]_{\Psi} = A^{\mathcal{D}} + A^{\mathcal{R}} + R, \tag{5.3}$$

where $A^{\mathcal{R}} \in \mathbf{S}^{\beta}$ is an off-diagonal *resonant* part (which our transformation cannot eliminate) and $R \in \mathbf{S}^{\gamma}$ for some $\gamma < \beta$. The exact form of the operators $A^{\mathcal{R}}$ and R depends on the choice of Ψ .

As a first step towards (5.3), let us rewrite the series (5.2) as

$$[A]_{\Psi} = A^{\mathscr{D}} + A^{\mathscr{O}\mathscr{D}} + \operatorname{ad}(A^{\mathscr{D}}; \Psi) + R,$$
(5.4)

with

$$R := \operatorname{ad}(A^{\mathscr{OD}}; \Psi) + \sum_{k=2}^{\infty} \frac{1}{k!} \operatorname{ad}^{k}(A; \Psi).$$
(5.5)

Suppose that $\Psi \in \mathbf{S}^{\zeta}$ with $\zeta \in \mathbb{R}$ and let $\kappa \in \mathbb{R}$. In order to achieve that $R \in \mathbf{S}^{\gamma}$ for some $\gamma < \beta$, we can use the following estimates:

- (1) If $\zeta < 0$, then by Lemma 2.15 we get $ad(X; \Psi) \in \mathbf{S}^{\kappa+\zeta}$ for all $X \in \mathbf{S}^{\kappa}$. We call a gauge transform that only uses these trivial bounds on the commutator norms *weak*.
- (2) Sometimes the structure of the commutators allows us to prove $ad(X; \Psi) \in \mathbf{S}^{\kappa+\zeta-\varepsilon}$ for some $\varepsilon > 0$ and appropriate $X \in \mathbf{S}^{\kappa}$. A gauge transform exploiting this improvement shall be called *strong*.

As we will see, the main issue with the strong gauge transform is that some conditions under which it can be used may not be formally invariant under the use of the gauge transform, which is in general an iterative scheme. Furthermore, due to combinatorial issues it may be harder to verify that those conditions are still satisfied as the number of steps increase. However, as we will see, in many situations it is sufficient to make one step of the strong gauge transform, and proceed from there with the weak one.

5.1. **The commutator equation.** We recall that after the gauge transform we would like to arrive at the operator $[A]_{\Psi}$ as in (5.3), in the best possible case with $A^{\mathscr{R}} = 0$. Comparing (5.3) with (5.4) we obtain that $A^{\mathscr{R}} = 0$ is equivalent to the commutator equation

$$\mathrm{ad}(A^{\mathscr{D}};\Psi) + A^{\mathscr{O}\mathscr{D}} = 0 \tag{5.6}$$

for $\Psi = Op(\psi)$.

Let A = Op(a) and Θ be a frequency set for *a*. By (2.18), equation (5.6) is solved if Θ is a frequency set for ψ and

$$a^{\mathscr{D}}(\theta \triangleright \xi)\psi_{\theta}(\xi) - \psi_{\theta}(\xi)a^{\mathscr{D}}(\xi) = \mathrm{i}a_{\theta}^{\mathscr{O}\mathscr{D}}(\xi)$$

holds for all $\theta \in \Theta' := \Theta \setminus \{id\}$ and $\xi \in \Xi$. This leads to

$$\psi_{\theta}(\xi) = \frac{\mathrm{i}a_{\theta}^{\mathcal{O}\mathcal{D}}(\xi)}{a^{\mathcal{D}}(\theta \triangleright \xi) - a^{\mathcal{D}}(\xi)},\tag{5.7}$$

for $\theta \in \Theta'$ and $\xi \in \Xi$. However, the problem of small denominators $a^{\mathscr{D}}(\theta \triangleright \xi) - a^{\mathscr{D}}(\xi)$ for some pairs (θ, ξ) generally prevents such choice of ψ . This motivates the following definition.

Definition 5.3. For $\delta \in \mathbb{R}$, s > 0, and $\theta \in G$, we call a set $\Lambda_{\theta}^{\delta,s} \subset \Xi$ a δ -resonant region generated by θ for $A^{\mathcal{D}}$ if it satisfies

$$\Lambda_{\theta}^{\delta,s} \supset \left\{ \xi \in \Xi : \left| a^{\mathscr{D}}(\theta \triangleright \xi) - a^{\mathscr{D}}(\xi) \right| \le s \langle \xi \rangle^{\delta} \right\}.$$
(5.8)

A corresponding *resonance cut-off* is a function $\chi := \chi^{\delta,s} : G \times \Xi \to \mathbb{R}$, mapping $(\theta, \xi) \mapsto \chi^{\delta,s}_{\theta}(\xi)$, such that for all $\theta \in G$, we have

$$0 \le \chi \le 1,$$

$$\chi_{\theta}^{\delta,s}(\xi) = 0, \text{ for all } \xi \in \Lambda_{\theta}^{\delta,s},$$

$$\overline{\chi_{\theta^{-1}}(\theta \triangleright \xi)} = \chi_{\theta}(\xi), \text{ for all } \xi \in \Xi.$$
(5.9)

For a fixed resonance cut-off, we define the *resonant* part $B^{\mathscr{R}} := \operatorname{Op}(b^{\mathscr{R}})$ and the *non-resonant* part $B^{\mathscr{NR}} := \operatorname{Op}(b^{\mathscr{NR}})$ of any operator $B = \operatorname{Op}(b) \in \mathbf{S}^{\infty}$ via their symbols

$$b^{\mathscr{R}} := b^{\mathscr{O}\mathscr{D}}(1 - \chi^{\delta,s}),$$

$$b^{\mathscr{N}\mathscr{R}} := b^{\mathscr{O}\mathscr{D}}\chi^{\delta,s}.$$
 (5.10)

Remark 5.4.

- (i) For any $\delta \in \mathbb{R}$ and s > 0, the only δ -resonant region generated by id is $\Lambda_{id}^{\delta,s} = \Xi$. Hence, every resonance cut-off χ satisfies $\chi_{id} \equiv 0$.
- (ii) If $\Lambda_{\theta}^{\delta,s}$ satisfies

$$\Lambda_{\theta^{-1}}^{\delta,s} = \theta \triangleright \Lambda_{\theta}^{\delta,s}, \quad \text{for all } \theta \in G,$$

then the resonance cut-off χ can be chosen as

$$\chi_{\theta}(\xi) := \mathbf{1}_{\Xi \setminus \Lambda_{\alpha}^{\delta,s}}(\xi), \text{ for all } (\theta, \xi) \in G \times \Xi.$$

(iii) If $B^{\mathcal{OD}} \in \mathbf{S}^{\gamma}$, $\gamma \in \mathbb{R}$, then

$$B^{\mathscr{O}\mathscr{D}} = B^{\mathscr{N}\mathscr{R}} + B^{\mathscr{R}},$$

and

$$\left\| B^{\mathcal{N}\mathcal{R}} \right\|_{l}^{(\gamma)} \leq \left\| B^{\mathcal{O}\mathcal{D}} \right\|_{l}^{(\gamma)}, \qquad \left\| B^{\mathcal{R}} \right\|_{l}^{(\gamma)} \leq \left\| B^{\mathcal{O}\mathcal{D}} \right\|_{l}^{(\gamma)}$$
(5.11)
If *B* is symmetric, then so are $B^{\mathcal{D}}, B^{\mathcal{N}\mathcal{R}}$, and $B^{\mathcal{R}}$.

hold for all $l \ge 0$. If *B* is symmetric, then so are $B^{\mathcal{D}}$, $B^{\mathcal{NR}}$, and $B^{\mathcal{R}}$.

With the help of Definition 5.3, the problem of small denominators in (5.7) can be circumvented. Let $\delta \in \mathbb{R}$, s > 0, and fix a resonance cut-off χ corresponding to δ -resonant regions $\Lambda_{\theta}^{\delta,s}$, $\theta \in G$, for $A^{\mathcal{D}}$. Using (5.10), we define

$$\psi_{\theta}^{\delta,s}(\xi) := \begin{cases} \frac{i a_{\theta}^{\mathcal{NR}}(\xi)}{a^{\mathscr{D}}(\theta \triangleright \xi) - a^{\mathscr{D}}(\xi)} & \text{if } \theta \in \Theta', \\ 0 & \text{otherwise.} \end{cases}$$
(5.12)

Recall that $A^{\mathcal{OD}} \in \mathbf{S}^{\beta}$ so that, in view of Remark 5.4(iii), $A^{\mathcal{NR}} \in \mathbf{S}^{\gamma}$ for some $\gamma \leq \beta$.

Lemma 5.5. Let $\gamma \leq \beta$ with $A^{\mathcal{NR}} \in \mathbf{S}^{\gamma}$. Then (5.12) defines a symbol $\psi^{\delta,s} \in \mathbf{S}^{\gamma-\delta}$. The operator $\Psi := \operatorname{Op}(\psi^{\delta,s})$ is symmetric with

$$\|\Psi\|_{l}^{(\gamma-\delta)} \leq \frac{1}{s} \left\| A^{\mathscr{N}\mathscr{R}} \right\|_{l}^{(\gamma)}, \text{ for all } l \geq 0.$$
(5.13)

It satisfies

$$\mathrm{ad}(A^{\mathscr{D}};\Psi) + A^{\mathscr{N}\mathscr{R}} = 0. \tag{5.14}$$

Proof. The bounds (5.13) follow directly from (5.12) and (5.8)–(5.10). The equation (5.14) follows as in (5.6)–(5.7) with \mathcal{NR} replacing \mathcal{OD} .

In view of (5.14), (5.4) takes the form

$$[A]_{\Psi} = A^{\mathscr{D}} + A^{\mathscr{R}} + R, \qquad (5.15)$$

with R defined in (5.5).

5.2. Weak gauge transform. Let $\gamma \leq \beta$ such that $A^{\mathcal{NR}} \in \mathbf{S}^{\gamma}$, $A^{\mathcal{OD}} \in \mathbf{S}^{\beta}$. We choose $\delta > \gamma$, so that $\gamma - \delta < 0$ in Lemma 5.5. Note that δ determines the size of the resonant regions and thus the efficiency of the gauge transform.

Lemma 5.6. Let $\Psi = Op(\psi^{\delta,s})$ be the operator defined in (5.12) and R be as in (5.5). Then $ad(A^{OD}; \Psi)$, $R \in \mathbf{S}^{\beta+\gamma-\delta}$ are symmetric and, for all $l \ge 0$,

$$\left\|\operatorname{ad}(A^{\mathscr{OD}};\Psi)\right\|_{l}^{(\beta+\gamma-\delta)} \leq \frac{2}{s} \left\|A^{\mathscr{OD}}\right\|_{l+|\gamma-\delta|}^{(\beta)} \left\|A^{\mathscr{NR}}\right\|_{l+|\beta|}^{(\gamma)},\tag{5.16}$$

and

$$\|R\|_{l}^{(\beta+\gamma-\delta)} \leq \frac{3}{s} \left\| A^{\mathscr{O}} \right\|_{l+|\gamma-\delta|}^{(\beta)} \left\| A^{\mathscr{N}} \right\|_{l+|\beta|+|\gamma-\delta|}^{(\gamma)} \exp\left(\frac{2}{s} \left\| A^{\mathscr{N}} \right\|_{l+|\beta|+|\gamma-\delta|}^{(\gamma)}\right).$$
(5.17)

Proof. The estimates (2.20) and (2.21) together with $\delta > \gamma$ and Lemma 5.5 imply that, for all $k \in \mathbb{N}$,

$$\begin{split} \left\| \operatorname{ad}^{k}(A^{\mathcal{OD}}; \Psi) \right\|_{l}^{(\beta+\gamma-\delta)} &\leq 2 \left\| \operatorname{ad}^{k-1}(A^{\mathcal{OD}}; \Psi) \right\|_{l+|\gamma-\delta|}^{(\beta)} \|\Psi\|_{l+|\beta|}^{(\gamma-\delta)} \\ &\leq \frac{2^{k}}{s} \left(\|\Psi\|_{l+|\beta|+|\gamma-\delta|}^{(0)} \right)^{k-1} \left\| A^{\mathcal{OD}} \right\|_{l+|\gamma-\delta|}^{(\beta)} \left\| A^{\mathcal{NR}} \right\|_{l+|\beta|}^{(\gamma)} \end{split}$$

Thus, (5.16) follows by choosing k = 1. Moreover, (5.13) implies that for all $k \in \mathbb{N}$,

$$\left\|\operatorname{ad}^{k}(A^{\mathscr{O}\mathscr{D}};\Psi)\right\|_{l}^{(\beta+\gamma-\delta)} \leq \left(\frac{2}{s}\left\|A^{\mathscr{N}\mathscr{R}}\right\|_{l+|\beta|+|\gamma-\delta|}^{(\gamma)}\right)^{k}\left\|A^{\mathscr{O}\mathscr{D}}\right\|_{l+|\gamma-\delta|}^{(\beta)}.$$
(5.18)

Similarly, we get from (5.14) that for all $k \ge 2$,

$$\begin{aligned} \left\| \operatorname{ad}^{k}(A^{\mathscr{D}}; \Psi) \right\|_{l}^{(\beta+\gamma-\delta)} &= \left\| \operatorname{ad}^{k-1}(A^{\mathscr{NR}}; \Psi) \right\|_{l}^{(\beta+\gamma-\delta)} \\ &\leq \left(\frac{2}{s} \left\| A^{\mathscr{NR}} \right\|_{l+|\beta|+|\gamma-\delta|}^{(\gamma)} \right)^{k-1} \left\| A^{\mathscr{NR}} \right\|_{l+|\gamma-\delta|}^{(\beta)}. \end{aligned}$$

$$(5.19)$$

Hence, the bounds (5.17) follow from (5.5), (5.18), (5.19), and (5.11).

Lemmata 5.5 and 5.6 have the following immediate corollary, which follows by choosing $\gamma = \beta$ and applying (5.11).

Corollary 5.7. Let $\Psi = Op(\psi^{\delta,s})$ be the operator defined in (5.12) and *R* be as in (5.5). Assume that $\delta > \beta$. Then $\Psi \in \mathbf{S}^{\beta-\delta}$ is symmetric with

$$\|\Psi\|_{l}^{(\beta-\delta)} \leq \frac{1}{s} \left\| A^{\mathcal{O}\mathcal{D}} \right\|_{l}^{(\beta)}, \text{ for all } l \geq 0.$$

Moreover, $\operatorname{ad}(A^{\mathcal{OD}}; \Psi)$, $R \in \mathbf{S}^{2\beta-\delta}$ are symmetric and for all $l \ge 0$,

$$\left\|\operatorname{ad}(A^{\mathscr{OD}};\Psi)\right\|_{l}^{(2\beta-\delta)} \leq \frac{2}{s} \left(\left\|A^{\mathscr{OD}}\right\|_{l+|\beta|+|\beta-\delta|}^{(\beta)}\right)^{2},$$

and

$$\|R\|_{l}^{(2\beta-\delta)} \leq \frac{3}{s} \left(\left\| A^{\mathscr{O}}\right\|_{l+|\beta|+|\beta-\delta|}^{(\beta)} \right)^{2} \exp\left(\frac{2}{s} \left\| A^{\mathscr{O}}\right\|_{l+|\beta|+|\beta-\delta|}^{(\beta)} \right).$$

As a consequence of Lemma 5.6 we have arrived at (5.3) with $R \in \mathbf{S}^{\beta+\gamma-\delta}$ and $\beta+\gamma-\delta < \beta$ as desired. One may now iterate the gauge transform to further reduce the order of the error term, starting from $[A]_{\Psi}$ in the next step. We call such an iterative scheme *consecutive gauge transform*. A few remarks on this iterative scheme are in order.

Remark 5.8. (i) At each step of the consecutive gauge transform, the resonant regions can be chosen differently.

(ii) Let us consider a consecutive gauge transform consisting of k steps, starting with the operator $A_0 := A$, and transforming into the operator

$$A_j := \exp(-i\Psi_j)A_{j-1}\exp(i\Psi_j) = \left[\dots\left[[A]_{\Psi_1}\right]_{\Psi_2}\dots\right]_{\Psi_j}$$

at step j = 1, 2, ..., k. Moreover, suppose for simplicity that $\delta > \beta$ and that $\Lambda_{\theta}^{\delta,s}$, $\theta \in G$, are δ -resonant regions for all $A_j^{\mathcal{D}}$, j = 0, 1, ..., k - 1, simultaneously (so that the resonant cutoff $\chi = \chi^{\delta,s}$ can be chosen at all steps as $\chi_{\theta}(\xi) = \mathbf{1}_{\Xi \setminus \Lambda_{\theta}^{\delta,s}}(\xi)$, $(\theta, \xi) \in G \times \Xi$). Then (5.15) and Corollary 5.7 imply that

$$A_1 := [A]_{\Psi_1} = A_0^{\mathscr{D}} + A_0^{\mathscr{R}} + R_1$$

with $R_1 \in \mathbf{S}^{2\beta-\delta}$. Repeating the procedure, we obtain

$$A_j = A_{j-1}^{\mathscr{D}} + A_{j-1}^{\mathscr{R}} + R_j, \quad j = 1, 2, \dots, k,$$

with $R_i \in \mathbf{S}^{\beta+j(\beta-\delta)}$.

(iii) A disadvantage of the consecutive gauge transform lies in the fact that already the operator $A_1 = [A]_{\Psi_1}$ may have a frequency set as large as

$$Z(\Theta) := \bigcup_{n \in \mathbb{N}} \Theta^n, \tag{5.20}$$

where

$$\Theta^n := \underbrace{\Theta \cdot \ldots \cdot \Theta}_{\text{product taken } n \text{ times}}$$

This set $Z(\Theta)$ is usually infinite, even when Θ is finite. Thus, the same holds for $A_1^{\mathscr{R}}$, which arises after the second step of gauge transform. This might be inconvenient since one generally likes to keep the structure of the resonant operators as simple as possible. However, one can resolve this issue by excluding the terms belonging to $\mathbf{S}^{\beta+j(\beta-\delta)}$ from $A_j^{\mathscr{NR}}$ at the *j*th step (by moving them to the remainder). Then Θ^{j+1} will be the frequency set for $A_j^{\mathscr{R}}$.

In the next sub-section, we describe a different iterative gauge transform scheme that we call the *parallel gauge transform*. This is often more convenient to work with than the consecutive gauge transform.

5.3. **Parallel weak gauge transform.** Here, we perform several steps of the gauge transform at the same time, i.e.

$$A^{(k)} = [A]_{\Psi^{(\widetilde{k})}},$$

where

$$\Psi^{(\widetilde{k})} = \sum_{j=1}^{\widetilde{k}} \Psi_j$$

for some $\tilde{k} \in \mathbb{N}$. Fix again $\delta \in \mathbb{R}$, s > 0 and a resonant cut-off $\chi^{\delta,s}$ satisfying (5.9) corresponding to δ -resonant regions $\Lambda_{\theta}^{\delta,s}$, $\theta \in G$, for $A^{\mathcal{D}}$, see Definition 5.3. Following [PS12, Section 9], the operators Ψ_l , B_l , and T_l are recursively defined by

$$B_{1} := A^{\otimes \mathscr{D}},$$

$$B_{l} := \sum_{j=1}^{l-1} \frac{1}{j!} \sum_{k_{1}+k_{2}+\dots+k_{j}=l-1} \operatorname{ad}(A^{\otimes \mathscr{D}}; \Psi_{k_{1}}, \Psi_{k_{2}}, \dots, \Psi_{k_{j}}), \ l \ge 2,$$

$$T_{l} := \sum_{j=2}^{l} \frac{1}{j!} \sum_{k_{1}+k_{2}+\dots+k_{j}=l} \operatorname{ad}(A^{\otimes}; \Psi_{k_{1}}, \Psi_{k_{2}}, \dots, \Psi_{k_{j}}), \ l \ge 2,$$

RO

and the relations

$$ad(A^{\mathcal{D}}, \Psi_1) + B_1^{\mathcal{N}\mathscr{R}} = 0,$$

$$ad(A^{\mathcal{D}}, \Psi_l) + B_l^{\mathcal{N}\mathscr{R}} + T_l^{\mathcal{N}\mathscr{R}} = 0, \ l \ge 2.$$

(5.21)

35

More precisely, let Θ be a frequency set for A and for all $l \ge 1$, let b_l and t_l be the symbols of B_l and T_l , respectively. Analogously to (5.12), we solve (5.21) by choosing $\Psi_l := \operatorname{Op}(\psi_l)$ with

$$(\psi_1)_{\theta}(\xi) := \begin{cases} \frac{i(b_1^{\mathcal{N}\mathcal{R}})_{\theta}(\xi)}{a^{\mathcal{D}}(\theta \triangleright \xi) - a^{\mathcal{D}}(\xi)} & \text{if } \theta \in \Theta', \\ 0 & \text{otherwise} \end{cases}$$

and

$$(\psi_l)_{\theta}(\xi) := \begin{cases} \frac{\mathrm{i}(b_l^{\mathcal{N}\mathcal{R}})_{\theta}(\xi) + \mathrm{i}(t_l^{\mathcal{N}\mathcal{R}})_{\theta}(\xi)}{a^{\mathscr{D}}(\theta \triangleright \xi) - a^{\mathscr{D}}(\xi)} & \text{if } \theta \in (\Theta^l)', \\ 0 & \text{otherwise} \end{cases}$$

for $l \ge 2$. Note that for all $l \ge 1$, Θ^l is a frequency set for B_l , T_l , and Ψ_l . Finally, put

$$Y_{\widetilde{k}} := \sum_{l=1}^{\widetilde{k}} B_l + \sum_{l=2}^{\widetilde{k}} T_l,$$

and

$$R_{\tilde{k}+1} := B_{\tilde{k}+1} + R_{\tilde{k}+1}^{(1)} + R_{\tilde{k}+1}^{(2)},$$

with

$$R_{\widetilde{k}+1}^{(1)} := \sum_{j \ge \widetilde{k}+1} \frac{1}{j!} \operatorname{ad}^{j}(A; \Psi^{(\widetilde{k})}),$$

$$R_{\widetilde{k}+1}^{(2)} := \sum_{j=1}^{\widetilde{k}} \frac{1}{j!} \sum_{k_{1}+k_{2}+\dots+k_{j} \ge \widetilde{k}+1} \operatorname{ad}(A; \Psi_{k_{1}}, \Psi_{k_{2}}, \dots, \Psi_{k_{j}}).$$

Then we arrive at

$$A^{(\tilde{k})} = A^{\mathcal{D}} + Y^{\mathcal{D}}_{\tilde{k}} + Y^{\mathcal{R}}_{\tilde{k}} + R_{\tilde{k}+1},$$
(5.22)

see Lemma 5.2, where $Y_{\tilde{k}}^{\mathscr{R}}$ is an operator with frequency set $\Theta_{\tilde{k}}$. The following Proposition provides norm estimates for the operators after the parallel gauge transform. In particular, it shows that if $\delta > \beta$, then we can assure that the error term $R_{\tilde{k}}$ belongs to classes of arbitrarily small order by choosing \tilde{k} sufficiently large.

Proposition 5.9. Let $A^{\mathcal{OD}} \in \mathbf{S}^{\beta}$ with $\delta > \beta$. Then we have for all $l \ge 0$,

$$\|\Psi_k\|_l^{(k(\beta-\delta))} \ll \left(\left\|A^{\mathscr{O}}\right\|_{l+n_k}^{(\beta)}\right)^k, \ k \ge 1$$

$$\|B_k\|_l^{(k(\beta-\delta)+\delta)} + \|T_k\|_l^{(k(\beta-\delta)+\delta)} \ll \left(\left\|A^{\mathscr{O}}\right\|_{l+n_k}^{(\beta)}\right)^k, \ k \ge 2,$$
(5.23)

where n_k is an increasing function of k, depending on k, β and δ , and the implied constants depend only on k, β , δ , and s in (5.8). Moreover, the operators $\Psi^{(\tilde{k})} \in \mathbf{S}^{\beta-\delta}$, $Y_{\tilde{k}} \in \mathbf{S}^{\beta}$, $R_{\tilde{k}+1} \in \mathbf{S}^{\tilde{k}(\beta-\delta)+\beta}$

are symmetric and satisfy the bounds

$$\begin{split} \|\Psi^{(\widetilde{k})}\|_{l}^{(\beta-\delta)} + \|Y_{\widetilde{k}}\|_{l}^{(\beta)} \ll \left(1 + \left\|A^{\mathcal{O}\mathcal{D}}\right\|_{l+n_{\widetilde{k}}}^{(\beta)}\right)^{\widetilde{k}} \\ \|R_{\widetilde{k}+1}\|_{l}^{(\widetilde{k}(\beta-\delta)+\beta)} \leq C_{A,\widetilde{k},\beta,\delta,l}, \end{split}$$

for all $l \ge 0$, where the implied constants only depend on \tilde{k}, β, δ , and s; and $C_{A,\tilde{k},\beta,\delta,l}$ is a bounded function of the symbol norms $\left\{ \|A^{OD}\|_{l}^{(\beta)} \right\}_{l \ge 0}, \tilde{k}, \beta, \delta, \text{ and } l.$

Proof. The bounds (5.23) are easily deduced from Corollary 5.7 by induction in k, estimating all involved commutators using (2.20). The estimates on the symbol norms of $\Psi^{(\tilde{k})}$ and $Y_{\tilde{k}}$ follow readily.

Let us prove the estimates on the norms of $R_{\tilde{k}+1}$. Starting with $R_{\tilde{k}+1}^{(1)}$ we note that, for $m \ge \tilde{k}+1$ and $\Psi := \Psi^{(\tilde{k})}$

$$\begin{split} & \left\|\operatorname{ad}^{m}(A;\Psi)\right\|_{l}^{(k(\beta-\delta)+\beta)} \\ & \leq \left\|\operatorname{ad}^{m}(A^{\mathcal{D}};\Psi)\right\|_{l}^{(\widetilde{k}(\beta-\delta)+\beta)} + \left\|\operatorname{ad}^{m}(A^{\mathcal{O}\mathcal{D}};\Psi)\right\|_{l}^{(\widetilde{k}(\beta-\delta)+\beta)} \\ & = \left\|\operatorname{ad}^{m-1}(Y_{\widetilde{k}}^{\mathcal{N}\mathcal{R}};\Psi)\right\|_{l}^{(\widetilde{k}(\beta-\delta)+\beta)} + \left\|\operatorname{ad}^{m}(A^{\mathcal{O}\mathcal{D}};\Psi)\right\|_{l}^{(\widetilde{k}(\beta-\delta)+\beta)} \\ & \leq 2^{m-\widetilde{k}-1} \left\|\operatorname{ad}^{\widetilde{k}}(Y_{\widetilde{k}}^{\mathcal{N}\mathcal{R}};\Psi)\right\|_{l}^{(\widetilde{k}(\beta-\delta)+\beta)} \left(\left\|\Psi\right\|_{l+|\widetilde{k}(\beta-\delta)+\beta|}^{(0)}\right)^{m-\widetilde{k}-1} \\ & + 2^{m-\widetilde{k}} \left\|\operatorname{ad}^{\widetilde{k}}(A^{\mathcal{O}\mathcal{D}};\Psi)\right\|_{l}^{(\widetilde{k}(\beta-\delta)+\beta)} \left(\left\|\Psi\right\|_{l+|\widetilde{k}(\beta-\delta)+\beta|}^{(0)}\right)^{m-\widetilde{k}}, \end{split}$$

where we apply (2.21) in the second inequality. Dividing by m! and summing over $m \ge \tilde{k} + 1$ we obtain a convergent sum, for which we use the estimates on the norms of $Y_{\tilde{k}}$ and $\Psi^{(\tilde{k})}$. Estimating $\|R_{\tilde{k}+1}^{(2)}\|_{l}^{(\tilde{k}(\beta-\delta)+\beta)}$ is somewhat easier since there are no convergence issues. This finishes the proof of the proposition.

5.4. **Strong gauge transform.** The aim of any (iterative) gauge transform scheme is to force the error term *R* after the gauge transform, see e.g. (5.3) or (5.22), into a class of (relatively) small order. For instance, in (5.4) and (5.5), we aim at $R \in \mathbf{S}^{\gamma}$ for some $\gamma < \beta$. If Ψ belongs to a class of negative order, as is the case if one can choose $\delta > \beta$ in Definition 5.3, for example, it can be trivially satisfied for some $\eta < \beta$, as was seen to be the case with the weak gauge transform. In some cases, however, one can not guarantee more than $\Psi \in \mathbf{S}^{0}$, whether it be by choosing $\delta = \beta$, or by introducing additional cut-offs in the definition of Ψ . This is notably the case for Schrödinger-type operators, whenever the perturbation is not in \mathbf{S}^{β} for $\beta < \alpha - 1$. In such a case, one can no longer rely on the trivial product estimates for ad($A^{\emptyset \mathcal{D}}, \Psi$) to get that $\mathrm{ad}(A^{\emptyset \mathcal{D}}, \Psi) \in \mathbf{S}^{\eta}$

for some $\eta < \beta$. In the next lemma, we give a sufficient condition that, nevertheless, yields the required improvement through commuting with Ψ .

Lemma 5.10. Suppose that $A \in \mathbf{ES}^{\alpha}$ and that $\Psi \in \mathbf{S}^{0}$ is defined as in (5.12). If both $\mathrm{ad}(A^{\mathscr{OD}}; \Psi) \in \mathbf{S}^{\gamma}$ and $\mathrm{ad}(A^{\mathscr{NR}}; \Psi) \in \mathbf{S}^{\gamma}$, then, with

$$[A]_{\Psi} = \exp(-i\Psi)A\exp(i\Psi),$$

we have that

$$R := [A]_{\Psi} - A^{\mathscr{D}} - A^{\mathscr{R}} \in \mathbf{S}^{\gamma}.$$

Remark 5.11. If, in addition to the assumptions of this lemma, we have $A^{@\mathscr{D}} \notin \mathbf{S}^{\gamma}$, this would mean that commuting with Ψ has improved order and we can therefore call this gauge transform strong. Note that we do not require improvements in order to happen at every iteration of the commutator, but only at the first step.

Proof. It follows from Lemma 5.2 and equation (5.14) that

$$R = \sum_{k\geq 1} \frac{1}{k!} \operatorname{ad}^{k}(A^{\mathcal{O}\mathcal{D}}; \Psi) + \sum_{k\geq 2} \frac{1}{k!} \operatorname{ad}^{k}(A^{\mathcal{D}}, \Psi)$$
$$= \sum_{k\geq 1} \frac{1}{k!} \operatorname{ad}^{k-1}(\operatorname{ad}(A^{\mathcal{O}\mathcal{D}}; \Psi); \Psi) + \sum_{k\geq 2} \frac{1}{k!} \operatorname{ad}^{k-1}(\operatorname{ad}(A^{\mathcal{N}\mathcal{R}}; \Psi), \Psi).$$

As in the proof of Lemma 5.2, both series converge absolutely in \mathbf{S}^{γ} .

Remark 5.12. The hypotheses of the previous lemma can be achieved in many ways. The most common one does not depend on the operator *A*, but only on the algebraic structure of \mathbf{S}^{∞} : it is when commutators naturally improve order. The principal example is pseudo-differential operators in $L^2(\mathbb{R}^d)$ that are almost periodic with respect to the translation group \mathbb{R}^d . To obtain the commutator estimates in this case one requires some limited smoothness in ξ . We refer to [Sob05, Lemma 3.4] for a proof, and [Sob06, PS10, PS12, MPS14, PS16] for examples of further applications. In all of these cases, the smooth structure of functions on \mathbb{R}^d was used, and the resonance cut-off functions were taken to be smooth approximations to indicator functions of the non-resonant regions rather the indicators themselves.

It is also possible that one cannot reach $\Psi \in \mathbf{S}^0$ through only non-resonant cut-offs. In such a case, in order to achieve convergence of the series for $\exp(i\Psi)$ and $[A]_{\Psi}$ achieved in (5.1) and Lemma 5.2 we will need energy cut-offs, i.e. cutting off large ξ . See [PS10, MPS14] where this idea is being used.

6. SYSTEMS OF ALMOST PERIODIC OPERATORS

In this section, we provide a construction suitable to describe almost periodic operators with matrix-valued symbols within the framework of Sections 2 – 5.

6.1. Symbol formalism for systems of almost periodic operators. Let the index set Ξ and the group *G* be as in Section 2. Let $m \in \mathbb{N}$ and $\mathbf{b} : G \times \Xi \to \mathscr{L}(\mathbb{C}^m)$ be a function such that there exists a countable frequency set $\Theta = \Theta^{-1} \subset G$ with $\mathbf{b}_{\theta}(\xi) = 0$ for all $\theta \in G \setminus \Theta$ and $\xi \in \Xi$. Furthermore, assume that

$$\sum_{\theta \in \Theta} \|\mathbf{b}_{\theta}(\xi)\|^2 < \infty, \text{ for all } \xi \in \Xi,$$

where $\|\cdot\|$ is the operator norm on $\mathscr{L}(\mathbb{C}^m)$. For every $\xi \in \Xi$, let $\{v_j(\xi) : j \in \mathbb{Z}/m\mathbb{Z}\}$ be an orthonormal basis for \mathbb{C}^m so that $\{\mathbf{e}_{\xi} \otimes v_j(\xi)\}_{\xi \in \Xi, j \in \mathbb{Z}/m\mathbb{Z}}$ is an orthonormal basis for $\ell^2(\Xi; \mathbb{C}^m) = \ell^2(\Xi) \otimes \mathbb{C}^m$. In analogy to (2.12), an almost periodic operator **B** in $\ell^2(\Xi; \mathbb{C}^m)$ with symbol **b** is defined by

$$\mathbf{B}(\mathbf{e}_{\xi} \otimes \nu_{j}(\xi)) := \sum_{\theta \in \Theta} \mathbf{e}_{\theta \succ \xi} \otimes [\mathbf{b}_{\theta}(\xi) \nu_{j}(\xi)].$$
(6.1)

We introduce the index set $\Xi_m := \Xi \times \mathbb{Z} / m\mathbb{Z}$ equipped with the weight function

$$\langle (\xi, j) \rangle_m := \langle \xi \rangle \tag{6.2}$$

and define the group $G_m := G \times \mathbb{Z}/m\mathbb{Z}$, and its (free) action on Ξ_m by

$$(g,k) \triangleright (\xi,j) := (g \triangleright \xi, k+j).$$
 (6.3)

Applying the unitary map $T_m: \ell^2(\Xi_m) \to \ell^2(\Xi; \mathbb{C}^m)$ defined by

$$T_m(\mathbf{e}_{(\xi,j)}) := \mathbf{e}_{\xi} \otimes \nu_j(\xi), \ (\xi,j) \in \Xi_m,$$
(6.4)

we can relate the operator **B** to the operator

$$B := T_m^* \mathbf{B} T_m \tag{6.5}$$

in $\ell^2(\Xi_m)$. For every $g \in G$ and $\xi \in \Xi$, let $[\mathbf{b}_g(\xi)]$ be the matrix representation of $\mathbf{b}_g(\xi) : \mathbb{C}^m \to \mathbb{C}^m$ with respect to the pair of bases $\{v_j(\xi)\}_j$ and $\{v_j(g \triangleright \xi)\}_j$ on the domain and codomain, respectively, i.e.

$$\mathbf{b}_{g}(\xi)v_{j}(\xi) = \sum_{k \in \mathbb{Z}/m\mathbb{Z}} \left[\mathbf{b}_{g}(\xi)\right]_{kj} v_{k}(g \succ \xi), \quad \text{for all } j \in \mathbb{Z}/m\mathbb{Z}.$$
(6.6)

Define the scalar symbol $b: G_m \times \Xi_m \to \mathbb{C}$ by

$$b_{(g,k)}(\xi,j) := \left[\mathbf{b}_g(\xi)\right]_{k+j\,j} \tag{6.7}$$

for $(g, k) \in G_m$ and $(\xi, j) \in \Xi_m$, and note that $\Theta \times \mathbb{Z}/m\mathbb{Z}$ is a frequency set for *b*. In view of (2.12), (6.3), and (6.7), we have

$$\operatorname{Op}(b)\mathbf{e}_{(\xi,j)} = \sum_{(\theta,k)\in\Theta\times\mathbb{Z}/m\mathbb{Z}} [\mathbf{b}_{\theta}(\xi)]_{kj} \mathbf{e}_{(\theta\succ\xi,k)}, \text{ for } (\xi,j)\in\Xi_m.$$
(6.8)

Hence, (6.1), (6.4), (6.5), (6.6), and (6.8) yield $T_m^* \mathbf{B} T_m = B = \operatorname{Op}(b)$. This justifies calling the operators from $\mathbf{S}^{\infty}(G_m, \Xi_m)$ systems of almost periodic operators, also known as matrix-valued operators. We shall use the notation

$$\mathbf{T}_{m}^{\gamma} := \mathbf{T}^{\gamma}(G_{m}, \Xi_{m}), \ \mathbf{T} \in \{\mathbf{S}, \mathbf{DS}, \mathbf{DES}, \mathbf{SES}, \mathbf{ES}\}, \ \gamma \in \mathbb{R} \cup \{\pm \infty\},$$
(6.9)

and

$$\mathbf{H}_{m}^{\gamma} := \mathbf{H}^{\gamma}(\Xi_{m}), \ \gamma \in \mathbb{R} \cup \{\pm \infty\}.$$

Since the map of symbols $\mathbf{b} \to b$ is one-to-one, for those \mathbf{b} which are mapped to $b \in \mathbf{S}_m^{\infty}$ we write $\operatorname{Op}(\mathbf{b}) := \operatorname{Op}(b)$. We use this identification to apply the results of Sections 2–5 without always making explicit the conjugation by the operators T_m . Note that $\langle (g, k) \rangle_m = \langle g \rangle$, for all $(g, k) \in G_m$, see (2.2) and (6.2). Hence, for $b \in \mathbf{S}_m^{\beta}$, we have the equivalence of norms,

$$\mathfrak{c}_{m}^{-1} \|b\|_{l}^{(\beta)} \leq \sum_{\theta \in \Theta} \langle \theta \rangle^{l} \sup_{\xi \in \Xi} \langle \xi \rangle^{-\beta} \|\mathbf{b}_{\theta}(\xi)\| \leq \mathfrak{c}_{m} \|b\|_{l}^{(\beta)},$$
(6.10)

where the constant $c_m > 0$ only depends on m. Two sub-algebras of \mathbf{S}_m^{∞} will be of particular interest in the sequel: uncoupled operators and diagonal operators.

Definition 6.1. The *uncoupled operators* in \mathbf{S}_m^{β} , $\beta \in \mathbb{R} \cup \{\pm \infty\}$, are defined by

 $\mathbf{US}_{m}^{\beta} := \{ \mathbf{B} \in \mathbf{S}_{m}^{\beta} : \text{the matrix } [\mathbf{b}_{g}(\xi)], \text{ see (6.6), is diagonal for all } g \in G, \ \xi \in \Xi \}$

 $= \{ \mathbf{B} \in \mathbf{S}_{m}^{\beta} : \text{the frequency set for } b \text{ can be chosen as a subset of } G \times \{0\} \}.$

For any operator $\mathbf{A} = \operatorname{Op}(\mathbf{a}) \in \mathbf{S}_m^{\alpha}$, $\alpha \in \mathbb{R} \cup \{\pm \infty\}$, we define the symbol

$$\left[\mathbf{a}_{g}^{\mathcal{U}}(\xi)\right]_{kj} := \begin{cases} \left[\mathbf{a}_{g}(\xi)\right]_{kj} & \text{if } k = j \\ 0 & \text{if } k \neq j \end{cases}, \text{ for all } g \in G, \xi \in \Xi, k, j \in \mathbb{Z}/m\mathbb{Z}.$$

We write $\mathbf{A}^{\mathscr{U}} := \operatorname{Op}(\mathbf{a}^{\mathscr{U}})$ for the projection of \mathbf{A} onto \mathbf{US}_m^{α} which we call the uncoupled part. We also denote $\mathbf{A}^{\mathscr{C}} := \mathbf{A} - \mathbf{A}^{\mathscr{U}}$ its coupled part. It can easily be seen that if $\mathbf{A} \in \mathbf{S}_m^{\gamma}$, $\gamma \in \mathbb{R}$ then for all $l \ge 0$

$$\left\|\mathbf{A}^{\mathscr{U}}\right\|_{l}^{(\gamma)} \leq \left\|\mathbf{A}\right\|_{l}^{(\gamma)}, \quad \left\|\mathbf{A}^{\mathscr{C}}\right\|_{l}^{(\gamma)} \leq \left\|\mathbf{A}\right\|_{l}^{(\gamma)}$$

$$(6.11)$$

and that if **A** is symmetric so are $\mathbf{A}^{\mathscr{U}}$ and $\mathbf{A}^{\mathscr{C}}$.

The second sub-algebra is \mathbf{DS}_m^{∞} , see (6.9) and Definition 3.1. Noting that $\mathrm{id}_{G_m} = (\mathrm{id}_G, 0)$, we infer from (6.7) that

$$\mathbf{DS}_m^{\infty} = \left\{ B \in \mathbf{US}_m^{\infty} : \mathbf{b}_g(\xi) = 0 \text{ for all } g \in G \setminus \{ \mathrm{id}_G \} \right\}$$

so that $\mathbf{DS}_m^{\infty} \subset \mathbf{US}_m^{\infty} \subset \mathbf{S}_m^{\infty}$. As in the scalar case, for any operator $\mathbf{A} = \operatorname{Op}(\mathbf{a}) \in \mathbf{S}_m^{\alpha}$, $\alpha \in \mathbb{R} \cup \{\pm \infty\}$, we denote by

$$\mathbf{A}^{\mathcal{D}} := \operatorname{Op}(\mathbf{a}^{\mathcal{D}})$$
$$\left[\mathbf{a}_{g}^{\mathcal{U}}(\xi)\right]_{kj} := \begin{cases} \left[\mathbf{a}_{g}(\xi)\right]_{kj} & \text{if } k = j \text{ and } g = \text{id} \\ 0 & \text{otherwise.} \end{cases}$$

and $\mathbf{A}^{\mathscr{O}\mathscr{D}} = \mathbf{A} - \mathbf{A}^{\mathscr{D}}$. This definition makes it so that $\mathbf{A}^{\mathscr{D}} = T_m A^{\mathscr{D}} T_m^*$. Similarly, if resonant and non-resonant regions are defined in terms of Ξ_m , we set $\mathbf{A}^{\mathscr{R}} = T_m A^{\mathscr{R}} T_m^*$ and $\mathbf{A}^{\mathscr{N}\mathscr{R}} = T_m A^{\mathscr{N}\mathscr{R}} T_m^*$. We can also combine notions of coupling and resonance; we set for instance $\mathbf{A}^{\mathscr{R},\mathscr{U}} = (\mathbf{A}^{\mathscr{R}})^{\mathscr{U}}$, and proceed similarly for other combinations of the indices.

The following lemma is useful when changing the reference orthonormal basis of \mathbb{C}^m . Nevertheless, for the rest of this section the reference basis of $\ell^2(\Xi;\mathbb{C}^m)$ will remain fixed as $\{\mathbf{e}_{\xi} \otimes v_j(\xi)\}_{(\xi,j)\in \Xi_m}$.

Lemma 6.2. Assume that, for every $\xi \in \Xi$, the set $\{u_j(\xi) : j \in \mathbb{Z}/m\mathbb{Z}\}$ is an orthonormal basis for \mathbb{C}^m . Then the unitary operator

$$\mathbf{U}: \ell^2(\Xi; \mathbb{C}^m) \to \ell^2(\Xi; \mathbb{C}^m), \quad \mathbf{e}_{\xi} \otimes \nu_j(\xi) \mapsto \mathbf{e}_{\xi} \otimes u_j(\xi)$$
(6.12)

satisfies $T_m^* \mathbf{U} T_m \in \mathbf{S}_m^0$.

Proof. It is clear from (6.12) that $T_m^* \mathbf{U} T_m = \operatorname{Op}(\mathbf{u})$ where $\mathbf{u}_g(\xi) \in U(m)$ is unitary for all $g \in G, \xi \in \Xi$, and $\{\operatorname{id}_G\}$ is a frequency set for \mathbf{u} . Thus the equivalence of norms (6.10) implies that $T_m^* \mathbf{U} T_m \in \mathbf{S}_m^0$.

The previous lemma has the following corollary, justifying our terminology of uncoupled operators.

Corollary 6.3. Let $\{v_j : j \in \mathbb{Z}/m\mathbb{Z}\}$ be a fixed basis for \mathbb{C}^m . Then any operator $A \in \mathbf{US}_m^\infty$ is unitarily equivalent to an orthogonal sum $\bigoplus_{j \in \mathbb{Z}/m\mathbb{Z}} A_j$ where for every $j \in \mathbb{Z}/m\mathbb{Z}$, A_j acts in $\ell^2(\Xi) \otimes \text{span}\{v_j\}$.

6.2. Gauge transform in S_m^{∞} : the reduction to uncoupled operators. We would like to apply a weak gauge transform to an operator in the class SES_m^{∞} — cf. (6.9) — in order to obtain an operator of the same ordrer with an uncoupled principal symbol. In this section, we give two sufficient conditions that allow us to do this. The first one is more restrictive on the off-diagonal part and

gives a non-trivial remainder, but allows for a principal symbol with multiple eigenvalue. The second one requires the principal symbol to have only simple eigenvalues, in which case the procedure is more efficient and the restrictions on the off-diagonal symbol are much milder.

Theorem 6.4. Let $\mathbf{A} = \operatorname{Op}(\mathbf{a}) \in \operatorname{SES}_m^{\alpha}$ be symmetric and let $\beta < \alpha$ be such that $\mathbf{A}^{\mathscr{C}} := \operatorname{Op}(\mathbf{a}^{\mathscr{C}}) \in \mathbf{S}_m^{\beta}$. Assume that

$$\left[\mathbf{a}_{\mathrm{id}}\right](\xi) = \langle \xi \rangle^{\alpha} \operatorname{diag}(a_1(\xi), \dots, a_m(\xi))$$

and that Θ is a frequency set for $\mathbf{a}^{\mathcal{OD}}$. Here, for $j \in \mathbb{Z}/m\mathbb{Z}$, $a_j : \Xi \to \mathbb{R}$ are bounded functions such that for all $\theta \in Z(\Theta) = \bigcup_{k=1}^{\infty} \Theta^k$,

$$\lim_{\langle \xi \rangle \to \infty} \frac{a_j(\theta \triangleright \xi)}{a_j(\xi)} = 1.$$
(6.13)

Suppose finally that there exists C, c > 0 such that for every $j \in \mathbb{Z}/m\mathbb{Z}$ and $k \in \mathbb{Z}/m\mathbb{Z} \setminus \{0\}$, either

$$\inf_{\langle \xi \rangle > C} \left| a_j(\xi) - a_{j+k}(\xi) \right| \ge c > 0, \tag{6.14}$$

or

$$\mathbf{a}^{\mathscr{O}\mathscr{D}}]_{j,j+k} \in \mathbf{S}^{2\beta-\alpha}.$$
(6.15)

Then, for all $\varepsilon > 0$ and $N \in \mathbb{N}$ there exists a symmetric operator $\Psi \in \mathbf{S}_m^{\beta - \alpha}$ such that

$$[\mathbf{A}]_{\Psi} = \exp(-i\Psi)\mathbf{A}\exp(i\Psi) = \mathbf{A}^{\mathscr{D}} + \mathbf{Y} + \mathbf{R}_1 + \mathbf{R}_2$$
(6.16)

where $\mathbf{Y} \in \mathbf{US}_m^{\beta}$, $\mathbf{R}_1 \in \mathbf{S}^{2\beta-\alpha}$, $\|\mathbf{R}_2\|_{\mathbf{H}_m^{\beta} \to \mathbf{H}_m^0} < \varepsilon$ and $\mathbf{Y}, \mathbf{R}_1, \mathbf{R}_2$ are symmetric. If \mathbf{A} is quasi-periodic, one can choose $\mathbf{R}_2 = 0$.

Remark 6.5. The conditions (6.14) and (6.13) are satisfied in the simple case of constant functions $a_i(\xi) = a_i \in \mathbb{R} \setminus \{0\}$.

Proof. Fix $\varepsilon' > 0$. We first eliminate the long-range coupling. Since $\|\mathbf{A}^{\oslash}\|_{0}^{(\beta)} < \infty$, there exists a finite subset $\widetilde{\Theta} \subset \Theta$, closed under inversion and containing the identity, such that

$$\sum_{\theta \in \Theta \setminus \widetilde{\Theta}} \sup_{\xi \in \Xi} \langle \xi \rangle^{-\beta} \left\| \mathbf{a}_{\theta}^{\mathscr{O}\mathscr{D}}(\xi) \right\| < \varepsilon'.$$
(6.17)

Let $\mathbf{B} := \operatorname{Op}(\mathbf{b})$ with the symbol

$$\mathbf{b}_{\theta}(\xi) := \begin{cases} \mathbf{a}_{\theta}^{\mathscr{O}\mathscr{D}}(\xi) & \text{if } \theta \in \widetilde{\Theta}, \\ 0 & \text{otherwise} \end{cases}$$

For $\widetilde{\mathbf{R}} := \mathbf{A}^{\mathcal{OD}} - \mathbf{B}$, (6.10) implies

$$\|\widetilde{\mathbf{R}}\|_{0}^{(\beta)} < \mathfrak{c}_{m}\varepsilon', \tag{6.18}$$

and we write $\widetilde{\mathbf{A}} := \mathbf{A}^{\mathcal{D}} + \mathbf{B}$ so that $\mathbf{A} = \widetilde{\mathbf{A}} + \widetilde{\mathbf{R}}$ and $\widetilde{\mathbf{A}}^{\mathcal{D}} = \mathbf{A}^{\mathcal{D}}$. For every $j \in \mathbb{Z}/m\mathbb{Z}$, define the set

$$I_i := \{k \in \mathbb{Z} / m\mathbb{Z} : (6.14) \text{ holds} \}.$$

Finiteness of $\tilde{\Theta}$ and bounded range of action imply that

$$\lim_{\langle \xi \rangle \to \infty} \sup_{\theta \in \widetilde{\Theta}} \left| \frac{\langle \theta \triangleright \xi \rangle}{\langle \xi \rangle} - 1 \right| = 0$$

Combining this with (6.13) and (6.14), as well as boundedness of the functions a_j implies the existence of s' depending on ε' and the constants c, C in (6.14) such that

$$\inf_{j\in\mathbb{Z}/m\mathbb{Z}}\inf_{k\in I_j}\inf_{\theta\in\widetilde{\Theta}}\inf_{\langle\xi\rangle>s'}\left|a_{j+k}(\theta\triangleright\xi)\langle\theta\triangleright\xi\rangle^{\alpha}-a_j(\xi)\langle\xi\rangle^{\alpha}\right|>\frac{c}{2}\langle\xi\rangle^{\alpha}.$$

Thus, for $(g, k) \in G_m$, the sets

$$\Lambda_{(g,k)}^{\alpha,c/2} := \begin{cases} \{(\xi,j) \in \Xi_m : \min\{\langle\xi\rangle, \langle g \triangleright \xi\rangle\} \le s'\}, & \text{if } g \in \widetilde{\Theta} \text{ and } k \in I_j, \\ \Xi_m, & \text{otherwise} \end{cases}$$

are α -resonant regions for the operator $\tilde{A} = T_m^* \mathbf{A} T_m$, cf. (5.8), and we choose the corresponding (scalar) resonance cut-off function

$$\chi_{(g,k)}(\xi,j) := \mathbf{1}_{\Xi_m \setminus \Lambda^{\alpha,c/2}_{(\sigma,k)}(\xi,j)},$$

see Remark 5.4(ii). Thus, taking Ψ as in Lemma 5.5, we have that $\Psi = T_m \Psi T_m^* \in \mathbf{S}_m^{\beta-\alpha}$. In view of (5.15), we deduce that

$$[\widetilde{\mathbf{A}}]_{\Psi} = \exp(-i\Psi)\widetilde{\mathbf{A}}\exp(i\Psi) = \mathbf{A}^{\mathscr{D}} + \widetilde{\mathbf{A}}^{\mathscr{R}} + \mathbf{R},$$

where Corollary 5.7 and conjugation by T_m give $\mathbf{R} \in \mathbf{S}_m^{2\beta-\alpha}$. We turn our attention to $\widetilde{\mathbf{A}}^{\mathscr{R}}$. We decompose it as $\widetilde{\mathbf{A}}^{\mathscr{R}} = \widetilde{\mathbf{A}}^{\mathscr{R},\mathscr{U}} + \widetilde{\mathbf{A}}^{\mathscr{R},\mathscr{C}}$. By definition of the resonant region $\Lambda_{(g,k)}^{\alpha,c/2}$, we have that

$$[\widetilde{\mathbf{a}}^{\mathcal{R},\mathcal{C}}]_{j,j+k} = [\widetilde{\mathbf{a}}^{\mathcal{OD},\mathcal{C}}]_{j,j+k} \quad \text{if } k \not\in I_j$$

and

$$\operatorname{supp}([\mathbf{a}^{\mathscr{R},\mathscr{C}}]_{j,j+k}) \subset \left\{ \xi \in \Xi : \min_{\theta \in \widetilde{\Theta}} \langle \theta \triangleright \xi \rangle \le s' \right\} \quad \text{if } k \in I_j.$$

By (6.15), for every $k \notin I_j$ we have $[\tilde{\mathbf{a}}^{\mathcal{R},\mathcal{C}}]_{j,j+k} \in \mathbf{S}^{2\beta-\alpha}$. Finiteness of $\tilde{\Theta}$ and bounded range of action imply that the support of $[\mathbf{a}^{\mathcal{R},\mathcal{C}}]_{j,j+k}$ is bounded for $k \in I_j$. Together, along with Proposition 2.16, this gives $\tilde{\mathbf{A}}^{\mathcal{R},\mathcal{C}} \in \mathbf{S}_m^{2\beta-\alpha}$.

All of this implies

$$[\mathbf{A}]_{\Psi} = \mathbf{A}^{\mathscr{D}} + \widetilde{\mathbf{A}}^{\mathscr{R},\mathscr{U}} + \widetilde{\mathbf{A}}^{\mathscr{R},\mathscr{C}} + \mathbf{R} + \exp(-i\Psi)\widetilde{\mathbf{R}}\exp(i\Psi).$$
(6.19)

We claim that this has the desired form (6.16) with

$$\mathbf{Y} = \widetilde{\mathbf{A}}^{\mathscr{R},\mathscr{U}}, \qquad \mathbf{R}_1 = \widetilde{\mathbf{A}}^{\mathscr{R},\mathscr{C}} + \mathbf{R}, \qquad \text{and} \qquad \mathbf{R}_2 = \exp(-i\Psi)\widetilde{\mathbf{R}}\exp(i\Psi).$$
(6.20)

Indeed, it follows from (5.11) and (6.11) that $\tilde{\mathbf{A}}^{\mathcal{R},\mathcal{U}} \in \mathbf{US}_m^{\beta}$. We have that $\tilde{\mathbf{A}}^{\mathcal{R},\mathcal{C}}$, $\mathbf{R} \in \mathbf{S}_m^{2\beta-\alpha}$ so that their sum \mathbf{R}_1 also is.

Finally, recall that $\exp(i\Psi) \in \mathbf{S}_m^0$ by Corollary 2.7. In particular, we have

$$\left\|\exp(i\Psi)\right\|_{|\beta|}^{(0)} \le \sum_{k=0}^{\infty} \frac{\left(\left\|\Psi\right\|_{|\beta|}^{(0)}\right)^k}{k!},$$

where by Corollary 5.7 and conjugation with T_m we have

$$\|\Psi\|_{|\beta|}^{(0)} \le \|\Psi\|_{|\beta|}^{(\beta-\alpha)} \le \frac{4}{c} \|\mathbf{B}\|_{|\beta|}^{(\beta)} \le \frac{4}{c} \left\|\mathbf{A}^{\mathcal{OD}}\right\|_{|\beta|}^{(\beta)},$$

with *c* the constant in (6.14). Consequently, $\|\exp(i\Psi)\|_{|\beta|}^{(0)}$ is bounded uniformly in $\varepsilon' \searrow 0$. Hence, for any $\varepsilon > 0$, choosing

$$0 < \varepsilon' < \frac{\varepsilon}{\left(\left\|\exp(\mathrm{i}\Psi)\right\|_{|\beta|}^{(0)}\right)^2} \mathfrak{c}_m,$$

we obtain by Lemma 2.8 and (6.18) that

$$\begin{split} \|\mathbf{R}_{2}\|_{\mathbf{H}_{m}^{\beta} \to \mathbf{H}_{m}^{0}} &\leq \left\|\exp(i\Psi)\right\|_{\mathbf{H}_{m}^{0} \to \mathbf{H}_{m}^{0}} \|\widetilde{\mathbf{R}}\|_{\mathbf{H}_{m}^{\beta} \to \mathbf{H}_{m}^{0}} \left\|\exp(i\Psi)\right\|_{\mathbf{H}_{m}^{\beta} \to \mathbf{H}_{m}^{\beta}} \\ &\leq \|\widetilde{\mathbf{R}}\|_{0}^{(\beta)} \left(\left\|\exp(i\Psi)\right\|_{|\beta|}^{(0)}\right)^{2} < \varepsilon. \end{split}$$

This finishes the proof.

Theorem 6.6. Let $\mathbf{A} = \operatorname{Op}(\mathbf{a}) \in \operatorname{SES}_m^{\alpha}$ be symmetric and let $\beta < \alpha$ such that $\mathbf{A}^{\mathscr{OD}} := \operatorname{Op}(\mathbf{a}^{\mathscr{OD}}) \in \mathbf{S}_m^{\beta}$ with frequency set $\Theta \subset G$. Assume that

$$[\mathbf{a}_{id}(\xi)] = \langle \xi \rangle^{\alpha} \operatorname{diag}(a_1(\xi), \dots, a_m(\xi))$$

for some bounded functions $a_i: \Xi \to \mathbb{R}$. Moreover, suppose that there exist C, c > 0 such that

$$\inf_{\langle \xi \rangle > C} \min_{j \neq k} |a_j(\xi) - a_k(\xi)| \ge c > 0, \tag{6.21}$$

and that, for all j = 1, 2, ..., m, and $\theta \in Z(\Theta) = \bigcup_{k=1}^{\infty} \Theta^k$,

$$\lim_{\langle \xi \rangle \to \infty} \frac{a_j(\theta \triangleright \xi)}{a_j(\xi)} = 1.$$

Then, for all $\varepsilon > 0$ and $N \in \mathbb{N}$ there exists a symmetric operator $\Psi \in \mathbf{S}_m^{\beta-\alpha}$ such that

$$[\mathbf{A}]_{\Psi} = \exp(i\Psi)\mathbf{A}\exp(i\Psi) = \mathbf{A}^{\mathcal{D}} + \mathbf{Y}^{\mathcal{U}} + \mathbf{R}_1 + \mathbf{R}_2$$

with $\mathbf{Y} \in \mathbf{S}_{m}^{\beta}$, $\mathbf{R}_{1} \in \mathbf{S}_{m}^{-N}$, $\|\mathbf{R}_{2}\|_{\mathbf{H}_{m}^{\beta} \to \mathbf{H}_{m}^{0}} < \varepsilon$, and \mathbf{Y} , \mathbf{R}_{1} , \mathbf{R}_{2} symmetric. If $\mathbf{A}^{\mathcal{OD}}$ is quasi-periodic, then one can choose $\mathbf{R}_{2} = \mathbf{0}$.

Proof. The proof essentially follows the scheme of the proof of Theorem 6.4. We first eliminate long-range coupling and find $\mathbf{B} \in \mathbf{S}_m^\beta$ and $\widetilde{\mathbf{R}}$ such that $\mathbf{A} = \mathbf{A}^{\mathcal{D}} + \mathbf{B} + \widetilde{\mathbf{R}}$ and

$$\|\widetilde{\mathbf{R}}\|_0^{(\beta)} < \varepsilon'$$

Assumption (6.21) leads this time to α -resonant regions

$$\Lambda_{(g,k)}^{\alpha,c/2} = \begin{cases} \{(\xi,j) \in \Xi_m : \min\{\langle\xi\rangle, \langle g \succ \xi\rangle\} \le s'\}, & \text{if } k \ne 0, \\ \Xi_m, & \text{if } k = 0 \end{cases}$$
(6.22)

for some *s'* depending on ε' . Put

$$K := \frac{N+\beta}{\alpha-\beta}$$

We apply a parallel weak gauge transform according to (5.22). We have from Proposition 5.9 and conjugating by T_m that there exists symmetric operators $\Psi \in \mathbf{S}_m^{\beta-\alpha}$, $\mathbf{Y} \in \mathbf{S}_m^{\beta}$ and $\mathbf{R} \in \mathbf{S}_m^{-N}$ such that

$$[\widetilde{\mathbf{A}}]_{\Psi} = \exp(-i\Psi)\widetilde{A}\exp(i\Psi) = \mathbf{A}^{\mathscr{D}} + \mathbf{Y}^{\mathscr{D}} + \mathbf{Y}^{\mathscr{R}} + \mathbf{R}$$

and

$$\|\Psi\|_{|\beta|}^{(\beta-\alpha)} \ll \left(1 + \left\|\mathbf{A}^{\mathscr{O}\mathscr{D}}\right\|_{n_{K}}^{(\beta)}\right)^{K}.$$
(6.23)

where the implicit constant depends only on *c* and *K*. Inequality (6.23) implies that $\|\exp(i\Psi)\|_{|\beta|}^{(0)}$ is uniformly bounded as $\varepsilon' \searrow 0$.

With the resonant region as in (6.22), for every $\mathbf{Y} \in \mathbf{S}_m^{\gamma}$, $\gamma < \alpha$ we have that $\mathbf{Y}^{\mathcal{D}} + \mathbf{Y}^{\mathcal{R}} = \mathbf{Y}^{\mathcal{U}} + \mathbf{R}_{\mathbf{Y}}$, where the symbol of $\mathbf{R}_{\mathbf{Y}}$ has bounded support, implying $\mathbf{R}_{\mathbf{Y}} \in \mathbf{S}_m^{-\infty}$. We put $\mathbf{R}_1 = \mathbf{R} + \mathbf{R}_{\mathbf{Y}} \in \mathbf{S}_m^{-N}$ and $\mathbf{R}_2 = \exp(-i\Psi)\widetilde{\mathbf{R}}\exp(i\Psi)$ gives us

$$\|\mathbf{R}_2\|_{\mathbf{H}_m^\beta \to \mathbf{H}_m^0} \le \varepsilon' \left(\|\exp(i\Psi)\|_0^{(|\beta|)} \right)^2.$$

Therefore for any $\varepsilon > 0$ by choosing $0 < \varepsilon' < \varepsilon \left(\left\| \exp(i\Psi) \right\|_{0}^{(|\beta|)} \right)^{-2}$ we obtain

$$[\mathbf{A}]_{\Psi} = \mathbf{A}^{\mathcal{D}} + \mathbf{Y}^{\mathcal{U}} + \mathbf{R}_1 + \mathbf{R}_2$$

with the claimed properties.

PART II : APPLICATIONS TO ASYMPTOTIC PROPERTIES OF SYSTEMS

In this second part, we consider some specific examples where the methods and results developed in the first half are applicable. As was mentioned earlier, these methods work very well for operators

$$H = H_0 + B$$

of Schrödinger type acting on $L^2(\mathbb{R}^d)$. Here, $H_0 = (-\Delta)^{\alpha/2}$ and *B* is a pseudo-differential perturbation of order $\beta < \alpha$. In particulars, the gauge transform method allows us to solve the following two types of problems, see [BP09, MPS14, Par08, PS16, PS12, PS10, Sob06] :

- obtain a complete asymptotic expansion for the integrated density of states of almost periodic operators, as the spectral parameter goes to infinity;
- Prove that some elliptic periodic operators have the Bethe–Sommerfeld property, which asserts that the spectrum of such operators contains a half-line [λ;∞) for some λ ∈ ℝ.

We now consider these questions in the setting of elliptic systems of operators. We establish answers to both of these problems in the case where symbols are periodic, for the Bethe-Sommerfeld property, and almost periodic, for the integrated density of states. We will do so by using the tools developed in Part I of this paper to reduce these operators to uncoupled operators. We will show that such a reduction cannot change the integrated density of states too much, and we will show that it cannot open infinitely many gaps in the spectrum. Since elliptic systems of operators do not have to be semi-bounded, we will obtain these results as the spectral parameter goes to $\pm\infty$. In order to do this, we will establish quantitative estimates based upon the results of Sections 4, 5 and 6 under generic assumptions about the perturbations.

In Section 7, we describe the Besicovitch space of almost periodic functions, and the operators acting on it. We also describe the structure of operators that are periodic rather than simply almost periodic, interpreting the Floquet-Bloch decomposition through the lens of almost periodic functions.

In Section 7.2, we describe the approach and the conditions required to prove the existence of complete asymptotics for the integrated density of states (IDS), and we state Theorem 8.2, which describes the asymptotic behaviour of the IDS for elliptic systems of operators. We reduce the problem to obtaining asymptotics for the IDS in families of intervals, which is the statement of Theorem 8.5.

In Section 8, we prove Theorem 8.5 by showing that it holds for uncoupled operators. We then use the gauge transform to show that it is sufficient to obtain a complete asymptotic expansion for uncoupled operators to get one for general systems.

In Section 10, we change perspective and we study periodic operators. In Theorem 10.1, we give conditions under which elliptic systems of periodic operators enjoy the Bethe–Sommerfeld property. We then use the reduction to uncoupled operators and bounds for the density of states obtained in Section 8 to show that it is sufficient to prove that the spectral overlap function is sufficiently bounded away from 0 for uncoupled operators. This will be done by reusing the results of Section 4, but interpreting fibrewise eigenvalue counting functions as instances of the IDS.

We prove those lower bounds in Section 11 by refining arguments based on combinatorial geometry that were previously used in proving the Bethe-Sommerfeld conjecture for Schrödingertype operators.

Finally in Section 12, we spend a few words to show that periodic and almost periodic perturbations of the Dirac operator fit in the framework that we have described in this part.

7. BESICOVITCH SPACE AND SYSTEMS OF OPERATORS

In this section, we turn back to the space $B^2(\mathbb{R}^d; \mathbb{C}^m)$ of almost periodic vector-valued functions, corresponding to the case where $G = \Xi = \mathbb{R}^d$ and

$$\mathbf{g}_1\mathbf{g}_2:=\mathbf{g}_1+\mathbf{g}_2, \quad \mathbf{g}\triangleright\boldsymbol{\xi}:=\mathbf{g}+\boldsymbol{\xi},$$

for all $\mathbf{g}_1, \mathbf{g}_2, \boldsymbol{\xi} \in \mathbb{R}^d$. The weight function is $\langle \boldsymbol{\xi} \rangle = 1 + |\boldsymbol{\xi}|$. From (2.2) we also get that $\langle \mathbf{g} \rangle = 1 + |\mathbf{g}|$ and that *G* has bounded range of action. The case m = 1 corresponds to the usual Besicovitch space. We now offer a concrete description of this space, along with a few results relating the properties of operators acting on L² and B². These results can be found in [CMS73, Shu78, Shu79b].

Let $\{v_1, ..., v_m\}$ be an orthonormal basis for \mathbb{C}^m and for $1 \le j \le m$ let

$$\mathbf{e}_{\boldsymbol{\xi},i}(\mathbf{x}) := \exp(i\boldsymbol{\xi}\cdot\mathbf{x}) \otimes \boldsymbol{v}_i,$$

The space $B^2(\mathbb{R}^d; \mathbb{C}^m)$ is the closure of

span
$$\left\{ \mathbf{e}_{\boldsymbol{\xi},j} : \boldsymbol{\xi} \in \mathbb{R}^d, j = 1, \dots, m \right\},\$$

taken with respect to the inner product

$$(f,g)_{\mathrm{B}} = \lim_{L \to \infty} \frac{1}{(2L)^d} \int_{[-L,L]^d} f \cdot \bar{g} \,\mathrm{d}\mathbf{x}$$

For the remainder of this article, we will use \mathbf{S}_m^{∞} , \mathbf{DS}_m^{∞} , etc. to refer to the spaces of almost periodic operators acting on $B^2(\mathbb{R}^d; \mathbb{C}^m)$. Let *A* be an operator in \mathbf{S}_m^{α} with symbol $\mathbf{a}(\mathbf{x}, \boldsymbol{\xi})$. The action of **A** in $L^2(\mathbb{R}^d; \mathbb{C}^m)$ as an operator in the Hörmander class $\Psi^{\alpha}(\mathbb{R}^d; \mathbb{C}^m)$ with almost periodic symbol is defined through the usual Fourier integral representation of pseudo-differential operators (see e.g. [Hör07]) as

$$\widetilde{\operatorname{Op}}(\mathbf{a})f(\mathbf{x}) = \frac{1}{(2\pi)^d} \iint_{\mathbb{R}^d \times \mathbb{R}^d} \exp(i\boldsymbol{\xi} \cdot (\mathbf{x} - \mathbf{y}))\mathbf{a}(\mathbf{x}, \boldsymbol{\xi})f(\mathbf{y}) \, \mathrm{d}\mathbf{y} \, \mathrm{d}\boldsymbol{\xi}.$$

The following proposition links its properties as an operator in L² and B², respectively.

Proposition 7.1. If $\mathbf{A} \in \mathbf{S}_m^{\infty}$ is bounded or elliptic, then

$$\operatorname{spec}_{\mathrm{B}^2}(\mathbf{A}) = \operatorname{spec}_{\mathrm{L}^2}(\mathbf{A})$$

as a set. In particular, if **A** is bounded, its norm in L^2 and B^2 coincide.

46

The proof of this proposition is exactly the same as the one in [Shu78] for the case m = 1. Indeed, it relies on some facts about function approximation proven in [Shu78, Lemmata 4.1 and 4.2] which remain true as m > 1 since they apply coordinatewise. Boundedness or ellipticity then implies Proposition 7.1. When we refer to the norm of an operator, we will not distinguish whether that operator is acting in $L^2(\mathbb{R}^d; \mathbb{C}^m)$ or $B^2(\mathbb{R}^d; \mathbb{C}^m)$ since those norms are the same.

As mentioned in Remark 4.2, there is a faithful, norm-preserving *-representation $\mathbf{A} \mapsto \mathbf{A}^{\sharp}$ of almost periodic operators given by $\mathbf{A}^{\sharp} := \mathbf{a}(\mathbf{x} + \mathbf{y}, D_{\mathbf{y}})$ acting in

$$\mathfrak{H}_m := \mathrm{B}^2(\mathbb{R}^d) \otimes \mathrm{L}^2(\mathbb{R}^d) \otimes \mathbb{C}^m.$$

Here **x** is a variable of functions in $B^2(\mathbb{R}^d; \mathbb{C}^m)$ and **y** is a variable of functions in $L^2(\mathbb{R}^d; \mathbb{C}^m)$. The operator \mathbf{A}^{\sharp} is interpreted as a direct integral over **x** of operators acting in $L^2(\mathbb{R}^d; \mathbb{C}^m)$. We denote by $e_I(x, y)$ the Schwartz kernel of the spectral projection $E_I(\mathbf{A})$. Note that in view of Proposition 7.1 and [CMS73], if $\mathbf{A} \leq \mathbf{B}$ as operators, then $\mathbf{A}^{\sharp} \leq \mathbf{B}^{\sharp}$ and $\|\mathbf{A}\| = \|\mathbf{A}^{\sharp}\|$.

Finally, the operator A^{\sharp} is affiliated to the ${\rm II}_{\infty}$ factor ${\mathfrak A}$ generated by the two families of operators

$$\left\{ \mathbf{e}_{\boldsymbol{\xi}} \otimes \mathbf{e}_{\boldsymbol{\xi}} \otimes M : \boldsymbol{\xi} \in \mathbb{R}^{d}, \ M \in \mathcal{M}_{m} \right\} \quad \text{and} \quad \left\{ I \otimes T_{\boldsymbol{\xi}} \otimes M : \boldsymbol{\xi} \in \mathbb{R}^{d}, \ M \in \mathcal{M}_{m} \right\}$$

where $\mathbf{e}_{\boldsymbol{\xi}}$ is the operator of multiplication by $e^{i\boldsymbol{\xi}\cdot\mathbf{x}}$, $T_{\boldsymbol{\xi}}$ is the operator of translation $T_{\boldsymbol{\xi}}f(\mathbf{x}) = f(\mathbf{x} - \boldsymbol{\xi})$ and \mathcal{M}_m is the algebra of $m \times m$ matrices with complex entries. This means that the results of Section 4–6 on the density of states measure (DSM) also called the integrated density of states (IDS) apply to this algebra of operators and this representation.

In the classical setting, the IDS is defined for differential operators using the large box limit and for pseudo-differential operators as the trace of the Schwartz kernel

$$N(J;\mathbf{A}) = M_{\mathbf{x}}(\operatorname{tr} e_{I}(\mathbf{x}, \mathbf{x})),$$

where *M* is the almost periodic mean. Note that this kernel is actually a smooth integral kernel whenever *J* is a bounded interval, see [PS16].

Our terminology for the IDS is justified in [Shu79b, Remark 3.1], where it is shown that the IDS as defined in Section 4 is the same as the one obtained from the classical definition for either differential or pseudo-differential operators.

7.1. **Concrete systems of operators.** From now on, we turn our attention to almost periodic pseudo-differential operators whose principal symbol is diagonal and nondegenerate.

Definition 7.2. A *uncoupleable* operator is an operator $\mathbf{A} \in \mathbf{ES}_m^{\alpha}$ for which there exists an unitary operator $\mathbf{U} \in \mathbf{S}_m^0$ so that $\mathbf{U}^* \mathbf{AU} = \mathbf{A}_0 + \mathbf{B} \in \mathbf{SES}_m^{\alpha}$ has the following properties.

• The *principal part* $\mathbf{A}_0 \in \mathbf{DES}_m^{\alpha}$, with symbol

$$\mathbf{a}_0(\boldsymbol{\xi}) = \operatorname{diag}\left(a_1 \,|\boldsymbol{\xi}|^{\alpha}, \dots, a_m \,|\boldsymbol{\xi}|^{\alpha}\right),\tag{7.1}$$

with $a_j \neq 0$ and without loss of generality $a_1 \ge ... \ge a_m$. We set $m^+ = \max j : a_j > 0$, where by convention $m_+ = 0$ if $a_1 < 0$.

The *subprincipal part* B ∈ S^β_m for β < α and has frequency set Θ. We also suppose that B is formally self-adjoint, i.e. that its symbol satisfies

$$\mathbf{b}_{\boldsymbol{\theta}}(\boldsymbol{\xi}) = \mathbf{b}_{-\boldsymbol{\theta}}(\boldsymbol{\xi} + \boldsymbol{\theta})^*$$

for all $\xi \in D$ and $\theta \in \Theta$, where for a matrix **a**, **a**^{*} is its conjugate transpose.

If $a_j \neq a_k$ for $j \neq k$, we say that **A** is a *competely uncoupleable* operator.

Remark 7.3. Since we are interested only in spectral properties of elliptic operators, for the remainder of this paper we can always assume that the operators are already in SES_m^{α} .

Without loss of generality, we assume that the frequency set Θ spans \mathbb{R}^d , contains **0**, and is symmetric about **0**. Recall from (5.20) that, using sum rather than product notations for the group of shifts in \mathbb{R}^d , that we also put

$$\Theta^{k} = \Theta + \dots + \Theta,$$

$$Z(\Theta) = \bigcup_{k \in \mathbb{N}} \Theta^{k}$$
(7.2)

The set $Z(\Theta)$ is countable and non-discrete, unless Θ generates a lattice.

where the sum is taken k times, and

7.2. Conditions on the perturbation and its frequency set. In this section, we state the exact conditions under which we can obtain asymptotics for the integrated density of states for a system of operators acting in $B^2(\mathbb{R}^d; \mathbb{C}^m)$. We also show how we can reduce the problem to computing the IDS solely on some intervals contained in a large enough range of energies.

We are interested in the asymptotics for the positive energy and negative energy integrated densities of states for a uncoupleable operator $\mathbf{A} = \mathbf{A}_0 + \mathbf{B}$, defined as

$$N^{+}(\lambda) := N^{+}(\lambda; \mathbf{A}) := N([0, \lambda); \mathbf{A}),$$

and

$$N^{-}(\lambda) := N^{-}(\lambda; \mathbf{A}) := N((-\lambda, 0]; \mathbf{A}),$$

as $\lambda \to \infty$. For this, we will need some conditions on the frequency set of the perturbation **B**. In Section 8, we reduce the operator **A** to a direct sum of operators of the type appearing in [MPS14]. In that paper, the perturbations are required to satisfy some conditions, which we describe for completeness. Conditions I and IV correspond to Conditions **A** and **C** in [MPS14] and we do not use them explicitly. Condition II addresses [MPS14, equation 2.4], while Condition III addresses [MPS14, Condition **B**]. We refer the reader to [MPS14], as well as [PS12] for a more detailed discussion around these conditions and their genericity.

We first need the following generic condition on the set $Z(\Theta)$ defined in (7.2).

Condition I. Suppose that $\theta_1, \ldots, \theta_d \in Z(\Theta)$. Then, $Z(\{\theta_1, \ldots, \theta_d\})$ is discrete.

This condition is clearly satisfied for periodic **B**, but for quasi-periodic or almost periodic **B** it is meaningful. The next two conditions describe how well **B** is approximated by finite sums of homogeneous functions of $\boldsymbol{\xi}$, and by quasi-periodic operators.

Condition II. There exists a constant $C_0 > 1$ and a discrete subset $J \subset (-\infty, \beta]$ such that for all $\theta \in \mathbb{R}^d$ and $|\xi| \ge C_0$,

$$(1 - \mathbf{1}_{C_0}(\boldsymbol{\xi}))\mathbf{b}_{\boldsymbol{\theta}}(\boldsymbol{\xi}) = \sum_{l \in J} |\boldsymbol{\xi}|^l \mathbf{b}_{\boldsymbol{\theta}}^{(l)} \left(\frac{\boldsymbol{\xi}}{|\boldsymbol{\xi}|}\right)$$

where $\mathbf{b}_{\theta}^{(l)} \in \mathbf{S}_{m}^{0}$ is positively homogeneous of degree 0. We also suppose that for $\boldsymbol{\eta} \in \mathbb{S}^{d-1}$, $\mathbf{b}_{\theta}^{(l)}(\boldsymbol{\eta})$ has a series representation (written in multi-index notation)

$$\mathbf{b}_{\boldsymbol{\theta}}^{(l)}(\boldsymbol{\eta}) = \sum_{\mathbf{n}\in\mathbb{N}_0^d} \mathbf{b}_{\boldsymbol{\theta}}^{(l,\mathbf{n})} \boldsymbol{\eta}^{\mathbf{n}},$$

which converges absolutely in a ball of radius greater than one of \mathbb{R}^d .

If **B** is quasi-periodic and J_0 is finite, these are the only conditions that we need. Otherwise, we need to find a quasi-periodic approximation of **B**. In view of (6.17), such an approximation will always exist, but we need a quantitative version of it.

Condition III. For every $k \in \mathbb{N}$, there exists $C_k > C_0$ such that for each $\rho > C_k$, there exists a finite symmetric $\tilde{\Theta} \subset (\Theta \cap \mathscr{B}(\rho^{1/k}))$ and a finite subset $J_k \subset (-\infty, \beta]$ with

$$#J_k \leq \rho^{1/k}$$

such that the symbol

$$\mathbf{r}_{\boldsymbol{\theta}}^{(k)}(\boldsymbol{\xi}) := \begin{cases} \mathbf{b}_{\boldsymbol{\theta}(\boldsymbol{\xi})} & \text{if } \boldsymbol{\theta} \notin \widetilde{\Theta}, \\ \mathbf{b}_{\boldsymbol{\theta}(\boldsymbol{\xi})} - \sum_{\iota \in J_k} |\boldsymbol{\xi}|^{\iota} \mathbf{b}_{\boldsymbol{\theta}}^{(\iota)} \left(\frac{\boldsymbol{\xi}}{|\boldsymbol{\xi}|}\right) & \text{if } \boldsymbol{\theta} \in \widetilde{\Theta}, \end{cases}$$
(7.3)

satisfies, for all $\ell \in \mathbb{N}$,

$$\left\|\mathbf{r}^{(k)}\right\|_{\ell}^{(\beta)} \le c_{\ell,k}\rho^{-k},\tag{7.4}$$

for some $c_{\ell,k} > 0$.

Finally, we need a Diophantine condition on the frequencies of *B*, for which we need some definitions. Fix $\tilde{k} \in \mathbb{N}$ (which will depend on the order of the remainder in the asymptotic expansion, but not on *k* as in Condition III). We say that \mathfrak{V} is a quasi-lattice subspace of dimension *q* if there are linearly independent $\theta_1, \ldots, \theta_q \in \widetilde{\Theta}^{\tilde{k}}$ such that $\mathfrak{V} = \operatorname{span}(\theta_1, \ldots, \theta_q)$. We denote by \mathcal{V} the collection of all quasi-lattice subspaces. For $\mathfrak{U}, \mathfrak{V} \in \mathcal{V}$, we write $\varphi(\mathfrak{U}, \mathfrak{V}) \in [0, \pi/2]$ to denote the angle between them, that is the angle between $\mathfrak{U} \ominus (\mathfrak{U} \cap \mathfrak{V})$ and $\mathfrak{V} \ominus (\mathfrak{U} \cap \mathfrak{V})$, where for a subspace $\mathfrak{W} \subset \mathfrak{V}, \mathfrak{V} \ominus \mathfrak{W}$ is the orthogonal complement of \mathfrak{W} in \mathfrak{V} . This angle is non-zero if and only if \mathfrak{U} and \mathfrak{V} are strongly distinct, *i.e.* if neither of them is a subspace of the other. Recalling that for any *k* the choice of $\widetilde{\Theta}$ depends on ρ , we put

$$R(\rho) = \sup_{\boldsymbol{\theta} \in \widetilde{\Theta}^{\widetilde{k}}} |\boldsymbol{\theta}|, \qquad r(\rho) = \inf_{\boldsymbol{\theta} \in \widetilde{(\Theta^{\widetilde{k}})'}} |\boldsymbol{\theta}|,$$

as well as

$$s := s(\rho) = s(\widetilde{\Theta}^k) := \inf \sin(\varphi(\mathfrak{U}, \mathfrak{V}))$$

where the infimum is over all strongly distinct pairs of subspaces in \mathcal{V} . It is clear that

$$R(\rho) = O\left(\rho^{1/k}\right),\tag{7.5}$$

where the implicit constant might depend on *k* and \tilde{k} ; however, we need the following condition for *r* and *s*.

Condition IV. For each fixed k and \tilde{k} , the sets $\tilde{\Theta}$ can be chosen in such way that for sufficiently large ρ , depending on k and \tilde{k} , the number of elements of $\tilde{\Theta}^{\tilde{k}}$ satisfies $\#\tilde{\Theta}^{\tilde{k}} \leq \rho^{1/k}$ and we have that

 $s(\rho) \ge \rho^{-1/k}$

and

$$r(\rho) \ge \rho^{-1/k}$$

Remark 7.4. Condition IV is automatically satisfied for quasi-periodic and smooth periodic **B**. See [PS12] for further discussion of this condition.

8. Asymptotic expansions for the IDS

We now suppose that the perturbation **B** satisfies Conditions I–IV and we set $\rho = \lambda^{1/\alpha}$, where α is the order of **A**₀. We prove the two following theorems, depending on whether all the a_j in (7.1) are distinct or not. Recall that $m_+ = \max\{j : a_j > 0\}$, with $m_+ = 0$ if $a_j < 0$ for all j.

Theorem 8.1. Let **A** be a uncoupleable operator with subprincipal part $\mathbf{B} \in \mathbf{S}_m^\beta$, $\beta \le \alpha/2$ satisfying Conditions I–IV. Suppose that there exists $\gamma \le 0$ such that whenever $a_j = a_k$ for some $1 \le j \ne k \le m$,

then $[\mathbf{B}]_{j,k} \in \mathbf{S}^{\gamma}$ and put $\gamma^* = \max(2\beta - \alpha, \gamma)$. Then, there exists a discrete set $L \subset (0, 1 - \gamma^*)$ and constants C_0^{\pm} and $C_{a,j}^{\pm}$, $0 \le q \le d - 1$, $j \in L$ such that

$$N^{\pm}(\mathbf{A};\rho^{\alpha}) = C_0^{\pm}\rho^d + \sum_{j \in L} \sum_{q=0}^{d-1} C_{j,q}^{\pm}\rho^{d-j}\log^q(\rho) + O(\rho^{d-1+\gamma^*}),$$
(8.1)

as $\rho \to \infty$. If $m^+ = m$ (resp. if $m^+ = 0$), then $C_0^- = C_{j,q}^- = 0$ (resp. $C_0^+ = C_{j,q}^+ = 0$) except for (j,q) = (d,0).

Theorem 8.2. Let **A** be a competely uncoupleable operator satisfying Conditions I–IV. Then, for every $K \in \mathbb{R}$ there exists a discrete set $L \subset (0, d + K)$ and constants $C_0^{\pm}, C_{q,j}^{\pm}, 0 \le q \le d - 1, j \in L$, such that

$$N^{\pm}(\mathbf{A};\rho^{\alpha}) = C_0^{\pm}\rho^d + \sum_{j\in L}\sum_{q=0}^{d-1} C_{j,q}^{\pm}\rho^{d-j}\log^q(\rho) + O(\rho^{-K}),$$
(8.2)

as $\rho \to \infty$. If $m^+ = m$ (resp. if $m^+ = 0$), then $C_0^- = C_{j,q}^- = 0$ (resp. $C_0^+ = C_{j,q}^+ = 0$) except for (j,q) = (d,0).

Remark 8.3. Note that the statement for $m^+ \in \{0, m\}$ follows from the operator being semibounded either above or below, respectively.

Note as well that if $J \subset \mathbb{Z}$, i.e. if the symbol of **A** is a classical symbol, see [Tay11, Chapter 7], then $L = \{0, ..., K + d - 1\}$.

The set *L* of allowable exponents can be made explicit, depending on *J* and *K*, see [MPS14, Remark 2.7].

The proof of Theorems 8.1 and 8.2 are obtained after many reductions to simpler cases. Recall that they are the general versions of Theorems 1.1 and 1.2 in the introduction.

8.1. **IDS for uncoupled operators.** In this subsection, we prove that the conclusion of Theorem 8.2 holds in the special case where $\mathbf{A} \in \mathbf{US}_m^{\alpha}$, regardless of whether an operator is uncoupleable or competely uncoupleable. This means that in addition of satisfying the conditions of Section 7.2, its symbol is given by

$$\mathbf{a}(\mathbf{x},\boldsymbol{\xi}) = \mathbf{a}_0(\boldsymbol{\xi}) + \mathbf{b}(\mathbf{x},\boldsymbol{\xi}),$$

where $\mathbf{b}(\mathbf{x}, \boldsymbol{\xi})$ is a diagonal matrix.

Proposition 8.4. Let $\mathbf{A} \in \mathbf{US}_m^{\alpha}$ be an uncoupleable operator satisfying conditions I–IV. Then, for every $K \in \mathbb{R}$ there exists a discrete set $L \subset (0, d + K)$ and constants $C_0^{\pm}, C_{q,j}^{\pm}, 0 \le q \le d - 1, j \in L$, such that

$$N^{\pm}(\mathbf{A};\rho^{\alpha}) = C_0^{\pm}\rho^d + \sum_{j \in L} \sum_{q=0}^{d-1} C_{j,q}^{\pm} \rho^{d-j} \log^q(\rho) + O(\rho^{-K}),$$

as $\rho \to \infty$. If $m^+ = m$ (resp. if $m^+ = 0$), then $C_0^- = C_{j,q}^- = 0$ (resp. $C_0^+ = C_{j,q}^+ = 0$) except for (j,q) = (d,0).

Proof. Since $\mathbf{A} \in \mathbf{US}_m^{\alpha}$, it can be split as a direct sum of operators $A_1 \oplus \ldots \oplus A_m$ acting in the mutually orthogonal subspaces $B^2(\mathbb{R}^d) \otimes v_j$. As such, we have that on any interval J,

$$N(J;\mathbf{A}) = \sum_{j=1}^{m} N(J;A_j).$$

This means that, for $j \le m^+$, A_j is semi-bounded below and acts invariantly on $B^2(\mathbb{R}^d) \otimes v_j$ as the operator considered in [MPS14]. For $j > m^+$, it is the operator $-\mathbf{A}_j$ that acts in such a way. From [MPS14, Theorem 2.5], this means that $N((-\infty, \lambda); \mathbf{A}_j)$ (resp. $N((\lambda, \infty); \mathbf{A}_j))$ enjoys an asymptotic expansion of the form (8.2) for $1 \le j \le m^+$ (resp. $m^+ < j \le m$). Observe that we have

$$N^{+}(\rho^{\alpha}; \mathbf{A}) = \sum_{j=1}^{m^{+}} N((-\infty, \rho^{\alpha}); A_{j}) - \sum_{j=1}^{m^{+}} N((-\infty, 0]; A_{j}) + \sum_{j=m^{+}+1}^{m} N((0, \rho^{\alpha}); A_{j}).$$

The terms in the first sum have the required asymptotic expansion. The terms in the second sum do not depend on ρ , hence they might only change the constant term in (8.2). Finally, the operators in the third sum are semi-bounded above, hence for ρ large enough the terms are constant and once again only affect the constant term. This proves the existence of the asymptotic expansion (8.2) for N^+ . The proof for N^- is the same, interchanging the role of the semi-bounded below and above operators.

8.2. **Reduction to a finite interval.** The strategy in this subsection is an adaptation of the one found in [MPS14, PS12]. It consists in showing that an asymptotic expansion holds in overlapping dyadic intervals I_n .

For K > -d, we choose ρ_0 sufficiently large, to be fixed later. For every $n \in \mathbb{N}$, we put $\rho_n := 2\rho_{n-1} = 2^n \rho_0$. We also define the intervals $I_n := [\rho_{n-1}, \rho_{n+1}]$. We prove the following theorem, which implies Theorems 8.1 and 8.2 as a corollary.

Theorem 8.5. Let **A** be an operator satisfying the conditions of either Theorem 8.1 or 8.2. Then, for either $K = -d + 1 - \gamma^*$ in the former case or any $K \in \mathbb{R}$ in the latter, there exists ρ_0 large enough, a discrete set $L \subset (0, d + K)$ and constants $C_0^{\pm}, C_{i,a}^{\pm}$ for every $j \in L$ and $0 \le q \le d - 1$ such that for every

 $n \in \mathbb{N}$ and every $0 < \mu < \nu$ with $\mu, \nu \in I_n$,

$$N((\mu^{\alpha}, \nu^{\alpha}); \mathbf{A}) = C_{0}^{+}(\nu^{d} - \mu^{d}) + \sum_{j \in L} \sum_{q=0}^{d-1} C_{j,q}^{+} \left(\nu^{d-j} \log^{q}(\nu) - \mu^{d-j} \log^{q}(\mu) \right) + O\left(\rho_{n}^{-K}\right),$$
(8.3)

where the implicit constants might depend on K, but not on n. Similarly,

$$\begin{split} N((-v^{\alpha},-\mu^{\alpha});\mathbf{A}) &= C_0^-(v^d-\mu^d) + \\ &+ \sum_{j \in L} \sum_{q=0}^{d-1} C_{j,q}^- \left(v^{d-j} \log^q(v) - \mu^{d-j} \log^q(\mu) \right) + O\left(\rho_n^{-K}\right). \end{split}$$

Remark 8.6. The reader familiar with previous works on the integrated density of states for almost periodic operators can notice that the roles of the dyadic decomposition in intervals I_n is slightly different here. In previous work, this decomposition was necessary because the resonant zones were significantly different for different spectral intervals were different. This yielded coefficients C^{\pm} depending possibly on *n*. It was however shown that the asymptotics had to match if the coefficients didn't grow too fast.

In our case, we need this decomposition in order to apply Theorem 8.9 when the perturbation is unbounded. Indeed, it relies on Lemma 4.12 which can only be applied for some interval with control on how far away the endpoints can be. We will therefore obtain asymptotics when both endpoints belong to a specific dyadic interval, then glue the intervals together. We end up comparing the density of states with the one obtained in [MPS14] for operators acting on scalar functions, i.e. the case m = 1. In such a case, the dependence on n of the coefficients has already been removed.

Proof of Theorems 8.1 and 8.2 assuming Theorem 8.5. We prove the theorem for N^+ , the proof for N^- is the same. For $K \in \mathbb{R}$, suppose that ρ_0 is large enough for Theorem 8.5 to hold. Suppose without loss of generality that for all n, ρ_n is a point of continuity of N^+ . For $\rho \in I_n$, we have that

$$N^{+}(\rho^{\alpha}) = N^{+}(\rho_{0}^{\alpha}) + \sum_{j=1}^{n-1} N((\rho_{j-1}^{\alpha}, \rho_{j}^{\alpha}); \mathbf{A}) + N((\rho_{n-1}^{\alpha}, \rho^{\alpha}); \mathbf{A})$$
$$= N^{+}(\rho_{0}^{\alpha}) + \sum_{j \in L} \sum_{q=0}^{d-1} C_{j,q}^{+} \left(\rho^{d-j} \log^{q}(\rho) - \rho_{0}^{d-j} \log^{q}(\rho_{0})\right) + \sum_{j=1}^{n} S_{j}$$

where $S_j = O(\rho_j^{-K})$. This implies that

$$\sum_{j=1}^{n} S_{j} \ll \rho_{0}^{-K} \sum_{j=1}^{n} 2^{-Kj} \ll \rho_{n}^{-K} \ll \rho^{-K}$$

since $\rho \in I_n$. One can see that the term depending on ρ_0 is O(1), so that it can be included either in the error term $O(\rho^{-K})$ when $K \le 0$ or in the constant term in (8.2) and (8.1) otherwise.

8.3. **Reduction to a quasiperiodic operator.** We now show in the following lemma that it is sufficient to prove Theorem 8.5 for quasiperiodic operators.

Lemma 8.7. Let $\mathbf{A} \in \mathbf{SES}_m^{\alpha}$ be an uncoupleable operator with subprincipal part $\mathbf{B} \in \mathbf{S}_m^{\beta}$ satisfying Condition III and $k \ge 2$. There exists $\rho_0 > 0$ and $0 < c_0 < 1$ so that for every $n \in \mathbb{N}$ there exists a quasi-periodic uncoupleable operator $\mathbf{A}' \in \mathbf{SES}_m^{\alpha}$ with frequency set $\widetilde{\Theta} \subset \mathcal{B}(\rho_n^{1/k})$ such that

• $\mathbf{A} - \mathbf{A}' \in \mathbf{S}_m^\beta$;

• supp
$$(\mathbf{a}^{\prime \mathscr{O} \mathscr{D}}) \subset \{ |\boldsymbol{\xi}| > c_0 \rho_n \};$$

• there is $\varepsilon \ll \rho_n^{\alpha-k}$ such that for all $J \subset I_n^{\alpha}$,

$$N((\pm J)_{-\varepsilon}; \mathbf{A}') \le N(\pm J; \mathbf{A}) \le N((\pm J)_{\varepsilon}; \mathbf{A}').$$
(8.4)

Proof. For $k \in \mathbb{N}$ let $\widetilde{\Theta} \subset \Theta \cap \mathscr{B}(\rho_n^{1/k})$ be the frequency set given by Condition III with $\rho = \rho_n$, and $\mathbf{R} \in \mathbf{S}_m^{\beta}$ be the operator with symbol given in (7.3), which by (7.4) satisfies $\|\mathbf{R}\|_0^{(\beta)} \ll \rho_n^{-k}$. Setting $\mathbf{A}'' = \mathbf{A} - \mathbf{R}$ we have that $\mathbf{A} - \mathbf{A}'' \in \mathbf{S}_m^{\beta}$, and that $\widetilde{\Theta}$ is a frequency set for \mathbf{A}'' and that as long as ρ_0 is large enough,

$$\left\|\mathbf{A}^{\prime\prime}\right\|_{0}^{(\gamma)} \le 2 \left\|\mathbf{A}\right\|_{0}^{(\gamma)} \tag{8.5}$$

for all $\beta \leq \gamma \leq \alpha$.

Writing any interval $J \subset \pm I_n^{\alpha}$ in the form [M - r, M + r], it is easy to see that $|M| + r \leq (2\rho_n)^{\alpha}$. Put $\beta_0 = \max{\{\beta, 0\}}$. By Lemma 4.11, estimate (8.4) holds with **A**'' instead of **A**' and

$$\varepsilon_{1} = \frac{\|\mathbf{R}\|_{0}^{(\beta_{0})}}{2 + \|\mathbf{R}\|_{0}^{(\beta_{0})}} \left(|M| + r + C(1 + \|\mathbf{R}\|_{0}^{(\beta_{0})})^{\frac{\alpha}{\alpha - \beta_{0}}} \right) \ll \rho_{n}^{\alpha - k}$$

instead of ε . Let us now define

$$\mathbf{A}' = \mathbf{A}'' - \mathbf{A}''^{\mathscr{O}\mathscr{D}} P_{c_0 \rho_n} = (\mathbf{A}'')^{\mathscr{D}} + \mathbf{A}''^{\mathscr{O}\mathscr{D}} (1 - P_{c_0 \rho_n}),$$

where $0 < c_0 < 1$ is to be determined later. By (8.5) and (3.1)

$$\left\|\mathbf{A}'\right\|_{0}^{(\gamma)} \le 4 \left\|\mathbf{A}\right\|_{0}^{(\gamma)}$$

for all $\beta \leq \gamma \leq \alpha$. We apply Lemma 4.12 with

$$H_0 = (\mathbf{A}'')^{\mathcal{D}}, \qquad B = \mathbf{A}''^{\mathcal{O}\mathcal{D}} (1 - P_{c_0 \rho_n}), \qquad A = \mathbf{A}''^{\mathcal{O}\mathcal{D}} P_{c_0 \rho_n}, \qquad H = \mathbf{A}'.$$

By Proposition 2.16,

$$\left\|\mathbf{A}^{\prime\prime\mathscr{O}\mathscr{D}}P_{c_0\rho_n}\right\| \leq (c_0\rho_n)^{\beta_0} \left\|\mathbf{A}^{\prime\prime\mathscr{O}\mathscr{D}}\right\|_0^{(\beta_0)}.$$

Set $X = \lfloor (2 - \alpha + k + \beta_0) \log_3 \rho_n \rfloor$, and let

$$Z_l := c_0 \rho_n + l \rho_n^{2/3}, \qquad 0 \le l \le X - 1, \tag{8.6}$$

so that if ρ_0 is large enough, $Z_{X-1} \leq 2c_0\rho_n$. For $0 \leq l \leq X$ introduce the family of projections

$$P_l := \begin{cases} P_{Z_0} & \text{for } l = 0, \\ P_{Z_l} - P_{Z_{l-1}} & \text{for } 0 < l < X, \\ 1 - P_{Z_{X-1}} & \text{for } l = X. \end{cases}$$

We now verify that the conditions of Lemma 4.12 are satisfied. It is clear that $\mathbf{B}P_{c_0\rho_n}P_{Z_0} = BP_{c_0\rho_n}$, and relations (4.13) follow from (7.5) and (8.6) as long as $k \ge 2$ and ρ_0 is large enough. By Proposition 2.16, for $0 \le l < X$,

$$\|P_{l}\mathbf{A}'P_{l}\| \leq Z_{X-1}^{\alpha} \|\mathbf{A}'\|_{0}^{(\alpha)} \leq 4(2c_{0}\rho_{n})^{\alpha} \|\mathbf{A}\|_{0}^{(\alpha)}.$$
(8.7)

We also have that

$$\left\| P_{l} \mathbf{A}^{\prime \prime \mathscr{O} \mathscr{D}} P_{l-1} \right\| + \left\| P_{l} \mathbf{A}^{\prime \prime \mathscr{O} \mathscr{D}} P_{l+1} \right\| \le 2Z_{X-1}^{\beta} \left\| \mathbf{A}^{\prime \prime} \right\|_{0}^{(\beta_{0})} \le 4(2c_{0}\rho_{n})^{\beta_{0}} \left\| \mathbf{A} \right\|_{0}^{(\beta_{0})}$$
(8.8)

For $0 \le l < X$, set

$$D_l = \operatorname{dist}(J, \operatorname{spec}(P_l \mathbf{A}' P_l)^{\sharp}).$$

By (8.7) and Lemma 4.3 for $l \le X - 1$

spec(
$$(P_l \mathbf{A}' P_l)^{\sharp}$$
) $\subset \left[-4(2c_0 \rho_n)^{\alpha} \|\mathbf{A}\|_0^{(\alpha)}, 4(2c_0 \rho_n)^{\alpha} \|A\|_0^{(\alpha)} \right],$

so that setting $c_0^{-\alpha} = 2^{\alpha+3} \|\mathbf{A}\|_0^{(\alpha)}$ gives

$$D_l \ge \frac{\rho_n^{\alpha}}{2},$$

in particular (4.14) holds. Combining with (8.8) we have that

$$\max_{0 \le l < X} \left\{ \frac{\left\| P_l \mathbf{A}^{\prime \prime \mathscr{O} \mathscr{D}} P_{l-1} \right\| + \left\| P_l \mathbf{A}^{\prime \prime \mathscr{O} \mathscr{D}} P_{l+1} \right\|}{D_l} \right\} \le 2(2c_0)^{\beta} \rho_n^{\beta - \alpha} \left\| \mathbf{A} \right\|_0^{(\beta_0)},$$

so that for ρ_0 large enough, (4.16) is satisfied.

Since the conditions of Lemma 4.12 are satisfied, for

$$\varepsilon_2 = 3^{2-X} \left\| \mathbf{A}^{\prime\prime \mathscr{O}\mathscr{D}} P_{c_0 \rho_n} \right\| \le 2\rho_n^{\alpha-k} \left\| \mathbf{A} \right\|_0^{(\beta_0)},$$

we have that

$$N(I_{-\varepsilon_1-\varepsilon_2};\mathbf{A}') \le N(I_{-\varepsilon_1};\mathbf{A}'') \le N(I;\mathbf{A});$$

and

$$N(I_{+\varepsilon_1+\varepsilon_2};\mathbf{A}') \ge N(I_{+\varepsilon_1};\mathbf{A}'') \ge N(I;\mathbf{A}).$$

Our claim therefore holds with $\varepsilon = \varepsilon_1 + \varepsilon_2$.

8.4. **Reduction to uncoupled operators.** In this subsection, we show that it is sufficient to prove Theorem 8.5 for quasi-periodic operators, it is sufficient to do so for uncoupled operators. At the end of the section, we finally prove Theorem 8.5 after all those reductions, which completes the proof of Theorems 8.1 and 8.2.

Theorem 8.8. Let **A** be an operator satisfying the conditions of Theorem 8.1. Then, for every $n \in \mathbb{N}$ there is an operator $A' \in \mathbf{US}_m^{\alpha}$ and $\varepsilon \ll \rho_n^{\gamma^*}$ such that for all $\mu, \nu \in I_n$ and $I = (\mu^{\alpha}, \nu^{\alpha})$,

$$N(\pm I_{-\varepsilon}; \mathbf{A}') \le N(\pm I; \mathbf{A}) \le N(\pm I_{\varepsilon}; \mathbf{A}').$$
(8.9)

In particular,

$$N(\pm I; \mathbf{A}) = N(\pm I; \mathbf{A}') + O\left(\rho_n^{d-1+\gamma^*}\right).$$
(8.10)

Proof. We only prove this theorem for *I*, the proof for -I follows from the same argument, making the relevant sign changes. By Proposition 8.4, we see that (8.10) follows from (8.9). We therefore only prove the latter.

We keep a quantitative track of the estimates found in Section 6.2. By Lemma 8.7, we can without loss of generality assume for some fixed $C_0 > 0$ that **A** is a quasi-periodic operator whose frequency set Θ lies in the ball $B(\rho_n^{1/k})$ for some $k \in \mathbb{N}$ and such that the support of $\mathbf{a}^{\mathcal{O}\mathcal{D}}$ lies in $\{|\xi| > C_0 \rho_n\}$. In particular, we can assume that there is s > 0 such that for all $\xi \in \text{supp}(\mathbf{a}^{\mathcal{O}\mathcal{D}})$, all $\theta \in \Theta$ and all j, k such that $a_j \neq a_k$,

$$\left|a_{j}|\boldsymbol{\theta}+\boldsymbol{\xi}|^{\alpha}-a_{k}|\boldsymbol{\xi}|^{\alpha}\right|>s|\boldsymbol{\xi}|^{\alpha}.$$

By Theorem 6.4, since **A** is quasi-periodic there are symmetric operators $\mathbf{Y} \in \mathbf{US}_m^{\beta}$, $\mathbf{R} \in \mathbf{S}_m^{\gamma^*}$ and $\Psi \in \mathbf{S}_m^{\beta-\alpha}$ such that **A** is unitarily equivalent through conjugation with $\exp(i\Psi)$ to

$$\mathbf{A}' + \mathbf{R} = \mathbf{A}^{\mathscr{D}} + \mathbf{Y} + \mathbf{R}.$$

Here, the symbol of Ψ is given by

$$[\boldsymbol{\psi}_{\boldsymbol{\theta}}(\boldsymbol{\xi})]_{j,k} = \frac{i[\mathbf{b}^{\mathscr{C}}_{\boldsymbol{\theta}}(\boldsymbol{\xi})]_{j,k}\chi_{j,k}}{a_{j}|\boldsymbol{\xi}+\boldsymbol{\theta}|^{\alpha}-a_{k}|\boldsymbol{\xi}|^{\alpha}},$$

where $\chi_{i,k} = 1$ if $a_i \neq a_k$ and 0 otherwise. Using the fact that

$$\mathrm{ad}(\mathbf{A}; \Psi) = \mathrm{ad}(\mathbf{A}^{\mathscr{O}\mathscr{D}}; \Psi) + \mathrm{ad}(\mathbf{A}^{\mathscr{D}}; \Psi) = \mathrm{ad}(\mathbf{A}^{\mathscr{O}\mathscr{D}}; \Psi) - \mathbf{A}^{\mathscr{N}\mathscr{R}},$$

the operator **R** is obtained from equations (6.19), (6.20) with $\tilde{\mathbf{R}} = 0$, and (5.5) by

$$\mathbf{R} = \mathbf{B}^{\mathcal{R},\mathcal{C}} + \mathrm{ad}(\mathbf{A}^{\mathcal{O}\mathcal{D}}; \mathbf{\Psi}) + \sum_{k=2}^{\infty} \frac{1}{k!} \mathrm{ad}^{k}(\mathbf{A}^{\mathcal{O}\mathcal{D}}; \mathbf{\Psi}) - \sum_{k=2}^{\infty} \frac{1}{k!} \mathrm{ad}^{k-1}(\mathbf{A}^{\mathcal{N}\mathcal{R}}; \mathbf{\Psi}).$$
(8.11)

By Lemma 4.11,

$$N(I_{-\varepsilon}; \mathbf{A}') \le N(I; \mathbf{A}' + \mathbf{R}) \le N(I_{\varepsilon}; \mathbf{A}')$$

for $\varepsilon = ||\mathbf{R}||$. Since Ψ has order $\beta - \alpha$ and is supported on $\{|\xi| > c\rho\}$, by Corollary 2.9 and Lemma 2.13 we have as in Proposition 2.16 that

$$\left\|\operatorname{ad}(\mathbf{A}^{\mathscr{O}\mathscr{D}}; \Psi)\right\| \ll \rho_n^{2\beta-\alpha} \left\|\mathbf{A}^{\mathscr{O}\mathscr{D}}\right\|_0^{(\beta)} \left\|\Psi\right\|_{|\beta|}^{(-\beta)},$$

so this gives the contribution from the second term in (8.11). The third and fourth terms uses the same estimate and the fact that this sum is absolutely convergent. Finally, for the first term we supposed that $\mathbf{B}^{\mathcal{R},\mathcal{C}} \in \mathbf{S}_m^{\gamma}$, and it is also supported on $\{|\xi| > c\rho\}$ so that by Proposition 2.16,

$$\left\|\mathbf{B}^{\mathscr{R},\mathscr{C}}\right\| \ll \rho_n^{\gamma}.$$

Together, this completes the proof.

When $a_j \neq a_k$ whenever $j \neq k$, we get the following stronger statement.

Theorem 8.9. Let **A** be an operator satisfying the hypotheses of Theorem 8.2. There is a decreasing sequence $\{\gamma_K\}_{K \in \mathbb{N}}, \gamma_K \to -\infty$ such that for all $K, n \in \mathbb{N}$ there is an operator $\mathbf{A}_K \in \mathbf{US}_m^{\alpha}$ and some $\varepsilon \ll \rho_n^{-\alpha - d - K}$ such that for all $\mu, \nu \in I_n$, and $I = (\mu^{\alpha}, \nu^{\alpha})$,

$$N(I_{-\varepsilon}; \mathbf{A}_K) \le N(I; \mathbf{A}) \le N(I_{\varepsilon}; \mathbf{A}_K)$$

and such that if $K_1 < K_2$, then

$$\mathbf{A}_{K_1} \equiv \mathbf{A}_{K_2} \mod \mathbf{S}_m^{\gamma_{K_1}}.$$

In particular,

$$N(\pm I;\mathbf{A}) = N(\pm I;\mathbf{A}_K) + O(\rho_n^{-K})$$

Proof. This statement is proven in the same way as the previous one, replacing the use of Theorem 6.4 with the parallel gauge transform Theorem 6.6, with a number of steps depending on *K*. This is possible because the condition on the terms coupling $a_j = a_k$ for $j \neq k$ is vacuously verified, so that it is assuredly preserved after each step of gauge transform. This yields a remainder $\mathbf{R} \in \mathbf{S}_m^{-N}$ for any *N*, allowing for the arbitrary precision in the approximation for the density of states.

Remark 8.10. Note that after this reduction, Conditions II and III, corresponding to [MPS14, Equation 2.4 and Condition B], do not hold anymore. However, the reason why these conditions are needed is to have a specific form for the functions $\mathbf{b}_{\theta_1}(\boldsymbol{\xi} + \boldsymbol{\theta}_2) \boldsymbol{\theta}_1, \boldsymbol{\theta}_2 \in \widetilde{\Theta}$, see [MPS14, Equation 10.5]. This expansion still holds if these conditions are imposed on the symbol prior to reduction to uncoupled operators.

Proof of Theorem 8.5. By Theorems 8.8 and 8.9, there is ρ_0 large enough so that for any *n* there is an operator $\mathbf{A}_K \in \mathbf{US}_m^{\alpha}$ such that for $\mu, \nu \in I_n$,

$$N((\mu^{\alpha}, \nu^{\alpha}); \mathbf{A}) = N((\mu^{\alpha}, \nu^{\alpha}); \mathbf{A}_{K}) + O(\rho_{n}^{-K}),$$

where $K = 1 - d - \gamma^*$ if **A** satisfies the hypotheses of Theorem 8.1 and $K \in \mathbb{R}$ if **A** satisfies the hypotheses of Theorem 8.2. Equation (8.3) (with coefficients $C_{j,q}^{\pm}$ depending on *n*) then follows by Proposition 8.4 for **A**_K and the fact that

$$N((\mu^{\alpha}, \nu^{\alpha}); \mathbf{A}_{K}) = N^{+}(\nu^{\alpha}; \mathbf{A}_{K}) - N^{+}(\mu^{\alpha}; \mathbf{A}_{K}).$$

$$(8.12)$$

In order to remove the dependence on *n* of the coefficients, it is sufficient for every $K \in \mathbb{N}$ to prove that they must agree for all *n* large enough. Since the coefficients obtained in Proposition 8.2 do not depend on *n*, this means that as soon as $\mu, \nu \in I_n \cap I_{n+1}$, (8.12) gives the same coefficients for the asymptotic expansion up to terms of order ρ_n^{-K} , which means that the coefficients need to agree for all *n* large enough.

9. The structure of periodic operators

We now turn our attention to periodic operators. In this section, we describe the structure of operators that are periodic with respect to some lattice Λ , and we give a quantitative approach to the study of the Bethe–Sommerfeld property. We realise the usual Bloch–Floquet decomposition through Besicovitch spaces.

9.1. **Description of periodic operators.** For periodic operators we assume that *G* is not \mathbb{R}^d but rather the dual lattice $\Theta := \Lambda^{\dagger} \subset \mathbb{R}^d$. We note that in this case $Z(\Theta) = \Theta$.

Invariance of **A** under the action of Λ means that for all $\mathbf{k} \in \mathbb{R}^d$, the subspace

$$\ell^{2}_{\mathbf{k}}(\Theta; \mathbb{C}^{m}) := \overline{\operatorname{span}\left\{\mathbf{e}_{\boldsymbol{\xi}, j} : 1 \le j \le m, \boldsymbol{\xi} \in \Theta + \mathbf{k}\right\}} \subset \mathrm{B}^{2}(\mathbb{R}^{d}; \mathbb{C}^{m})$$
(9.1)

is an invariant subspace for **A**, and we denote by $\mathbf{A}(\mathbf{k})$ the restriction of **A** to this subspace. It is clear from the definition that we can restrict ourselves to $\mathbf{k} \in \mathcal{O} = \mathbb{R}^d / \Theta$, and we call **k** a *quasimomentum*. For any $\boldsymbol{\xi} \in \mathbb{R}^d$, its fractional part { $\boldsymbol{\xi}$ } $\in \mathcal{O}$ is the image of $\boldsymbol{\xi}$ under the quotient map. The spectrum of **A** can be obtained as

$$\operatorname{spec}(\mathbf{A}) = \bigcup_{\mathbf{k} \in \mathcal{O}} \operatorname{spec}(\mathbf{A}(\mathbf{k})),$$

see [Kuc93, Theorem 4.5.1]. For every $\mathbf{k} \in \mathcal{O}$ the spectrum of $\mathbf{A}(\mathbf{k})$ in $\ell^2_{\mathbf{k}}(\Theta; \mathbb{C}^m)$ is discrete.

The usual approach to studying the $L^2(\mathbb{R}^d;\mathbb{C}^m)$ theory of periodic operators is through the Floquet-Bloch decomposition, see *e.g.* [Kuc93], where we represent **A** as a direct integral over

 \mathcal{O} of the fibre operators $\mathbf{A}(\mathbf{k})$. This would require us to introduce a considerable amount of machinery. However, the Bethe-Sommerfeld property is strictly about the spectrum as a set, and to every elliptic periodic operator acting in $L^2(\mathbb{R}^d; \mathbb{C}^m)$ there corresponds an elliptic operator acting in $B^2(\mathbb{R}^d; \mathbb{C}^m)$ with the same spectrum. We therefore consider periodic operators as operators on Besicovitch space where we decompose them according to the invariant subspaces (9.1). This makes our statements and proofs more direct.

Remark 9.1. The subspaces $\ell^2_{\mathbf{k}}(\Theta; \mathbb{C}^m)$ can be realised as

$$\mathrm{H}^{0}(\Theta + \mathbf{k}; \mathbb{C}^{m}) = \overline{\mathrm{span}\left(\left\{\mathbf{e}_{\boldsymbol{\theta}+\mathbf{k}, j}, \boldsymbol{\theta} \in \Theta, 1 \leq j \leq m\right\}\right)}.$$

The group *G* is Θ acting on Θ + **k** by translation. In this case, consider the I_{∞} factor \mathfrak{A} generated by { $\mathbf{e}_{\boldsymbol{\theta}} \otimes M : \boldsymbol{\theta} \in \Theta, M \in \mathcal{M}_m$ }.

It is clear that the restriction of the subalgebra of periodic operators to $\ell^2_{\mathbf{k}}(\Theta; \mathbb{C}^m)$ is affiliated to \mathfrak{A} , and that it respects the conditions described at the beginning of Section 4. The associated trace of the spectral projection over an interval *J* is simply $N(J; \mathbf{A}(\mathbf{k})) := \#\{j : \lambda_j(\mathbf{A}(\mathbf{k})) \in J\}$, the number of eigenvalues of $\mathbf{A}(\mathbf{k})$ in that interval.

We also make the observation that for a bounded, periodic self-adjoint operator Ψ , the restriction to $\ell^2_{\mathbf{k}}(\Theta; \mathbb{C}^m)$ of the unitary operator $\exp(i\Psi)$ is still unitary, since $\ell^2_{\mathbf{k}}(\Theta; \mathbb{C}^m)$ is an invariant subspace. This means that we can simultaneously use the gauge transform on each of the fibre operators and that the estimates from Section 4 hold uniformly for the counting function of the fibre operators.

Let us now describe the structure of the spectrum of A in terms of the spectra of the fibre operators $A(\mathbf{k})$. Since A is self-adjoint, it has Fredholm index 0, this implies that the Bloch variety

$$\{(\mathbf{k},\lambda)\subset \mathcal{O}\times\mathbb{R}:\lambda\in\operatorname{spec}(\mathbf{A}(\mathbf{k}))\}$$

is a principal analytic set [Kuc93, Corollary 3.1.6 and Section 3.4.C]. As such, if **A** is semi-bounded below, we can naturally label the eigenvalues of $\mathbf{A}(\mathbf{k})$ in non-decreasing order, counting multiplicity. Then, the functions $\lambda_j(\mathbf{k}) := \lambda_j(\mathbf{A}(\mathbf{k}))$ are piecewise analytic functions of **k**. If **A** is not semi-bounded, we can label the eigenvalues in non-decreasing order by $j \in \mathbb{Z}$ and it is possible to choose the labelling so that the functions $\lambda_j(\mathbf{k})$ are piecewise analytic. This requirement determines the labelling uniquely up to a uniform shift of the indices. Note that continuity in **k** of the functions λ_j and discreteness of the spectrum imply that labelling the eigenvalues at one quasimomentum **k** induces a labelling everywhere in \mathcal{O} . The interval

$$\iota_j := \iota_j(\mathbf{A}) := \bigcup_{\mathbf{k} \in \mathcal{O}^\dagger} \lambda_j(\mathbf{A}(\mathbf{k}))$$

is called the *j*th spectral band of **A**.

9.2. **The overlap function.** In order to prove that an operator has the Bethe-Sommerfeld property we study the band overlap, characterized by the *overlap function* $\zeta(\lambda; \mathbf{A})$, $\lambda \in \mathbb{R}$, introduced by M. Skriganov [Skr85]. The overlap function is defined as the maximal number *t* such that the symmetric interval $[\lambda - t, \lambda + t]$ is entirely contained in one band, *i.e.*

$$\zeta(\lambda; \mathbf{A}) := \begin{cases} \max_{j} \max\left\{t \ge 0 : [\lambda - t, \lambda + t] \subset \iota_{j}\right\} & \text{if } \lambda \in \operatorname{spec}(\mathbf{A}) \\ 0 & \text{if } \lambda \notin \operatorname{spec}(\mathbf{A}). \end{cases}$$
(9.2)

It is not hard to see that ζ is a continuous function of λ . In order to use our machinery we will relate the overlap function to the eigenvalue counting functions of the operators $\mathbf{A}(\mathbf{k})$. This type of idea has been used in the past but crucially relied on the fact that \mathbf{A} was semi-bounded below. In the following proposition we find an equivalent formulation that is robust under perturbations yet works for operators that are not semi-bounded. Recall that for an interval $I = [s, t] \subset \mathbb{R}$ and $\varepsilon \in \mathbb{R}$, we define

$$I_{\varepsilon} := \begin{cases} \varnothing & \text{for } \varepsilon < \frac{s-t}{2}, \\ [s-\varepsilon, t+\varepsilon] & \text{otherwise.} \end{cases}$$

Lemma 9.2. Suppose that \mathbf{A}_1 , \mathbf{A}_2 are self-adjoint periodic operators. Suppose that for all $p \in \{1, 2\}$, $\mathbf{k} \in \mathcal{O}$, $\mathbf{A}_p(\mathbf{k})$ has discrete spectrum. For $\lambda \in \mathbb{R}$ and t > 0, let

$$\delta = \min_{\mathbf{k} \in \mathcal{O}} \max\left\{ \operatorname{dist}(\mu; \operatorname{spec}(\mathbf{A}_1(\mathbf{k}))) : \mu \in [\lambda - t, \lambda + t] \right\}.$$
(9.3)

Suppose that there is $0 \le \varepsilon \le \delta/4$ such that for all $\mathbf{k} \in \mathcal{O}$ and any interval $I \subset [\lambda - t, \lambda + t]$

$$N(I;\mathbf{A}_{2}(\mathbf{k})) \leq N(I_{\varepsilon};\mathbf{A}_{1}(\mathbf{k})) \quad and \quad N(I;\mathbf{A}_{1}(\mathbf{k})) \leq N(I_{\varepsilon};\mathbf{A}_{2}(\mathbf{k})).$$

$$(9.4)$$

Then, for $p \in \{1,2\}$ *there exist sets of consecutive integers* $J_p \subset \mathbb{Z}$ *and surjective maps*

$$\lambda(\mathbf{A}_p(\mathbf{k})): J_p \to \operatorname{spec}(\mathbf{A}_p(\mathbf{k}))$$

such that for all $j \in J_p$, $\lambda_j(\mathbf{A}_p(\mathbf{k}))$ are continuous in \mathbf{k} and such that

$$\left|\lambda_{j}(\mathbf{A}_{1}(\mathbf{k})) - \lambda_{j}(\mathbf{A}_{2}(\mathbf{k}))\right| \leq \varepsilon \tag{9.5}$$

for all **k** and *j* such that $\lambda_j(\mathbf{A}_p(\mathbf{k})) \in [\lambda - t, \lambda + t]$.

Remark 9.3. We do not ask in the previous lemma that both operators share the properties of being either bounded, semi-bounded above or below, or unbounded in both directions.

Proof. For any $\mathbf{k} \in \mathcal{O}$, $0 < \eta \le \delta$, we say that $\mu \in [\lambda - t, \lambda + t]$ is η -distant (from the spectrum of $\mathbf{A}_1(\mathbf{k})$) at \mathbf{k} if

dist(
$$\mu$$
, spec($\mathbf{A}_1(\mathbf{k})$)) $\geq \eta$.

By (9.3), for every $\mathbf{k} \in \mathcal{O}^{\dagger}$ there exists $\mu \in [\lambda - t, \lambda + t]$ which is δ -distant at \mathbf{k} . By the second inequality in (9.4), if μ is η -distant at \mathbf{k} for some $\eta > \delta/4$, then for $p \in \{1, 2\}$

$$(\mu - \varepsilon, \mu + \varepsilon) \cap \operatorname{spec}(\mathbf{A}_p(\mathbf{k})) = \emptyset.$$
(9.6)

Choose $\mathbf{k}_0 \in \mathcal{O}$ and μ_0 a point δ -distant at \mathbf{k}_0 . Maps $j \mapsto \lambda_j(\mathbf{A}_p(\mathbf{k}))$ can be uniquely defined from the properties that they are nondecreasing, mapping to continuous functions in \mathbf{k} , and that $\lambda_0(\mathbf{A}_p(\mathbf{k}_0))$ is the smallest eigenvalue larger than μ_0 . Note that the sets J_1 and J_2 are both defined uniquely from these properties, in particular if \mathbf{A}_p is unbounded both above and below then $J_p = \mathbb{Z}$.

We now prove that, for all $\mathbf{k} \in \mathcal{O}$, if μ is $\delta/2$ -distant at \mathbf{k} , then for all $j \in J_1 \cap J_2$, then

$$\left(\lambda_{j}(\mathbf{A}_{1}(\mathbf{k}))-\mu\right)\left(\lambda_{j}(\mathbf{A}_{2}(\mathbf{k}))-\mu\right)>0,\tag{9.7}$$

in other words, for $p \in \{1,2\}$, $\lambda_j(\mathbf{A}_p(\mathbf{k}))$ are both on the same side of μ . The functions $\lambda_j(\mathbf{A}_p(\mathbf{k}))$ were constructed specifically so that (9.7) holds at \mathbf{k}_0 and μ_0 , our goal is to show that this property propagates to other μ and \mathbf{k} .

We first prove that if (9.7) holds for some $\mu \delta/2$ -distant at **k**, then it holds for all other $\nu \delta/2$ distant at **k**. This is a direct consequence of (9.4) and (9.6), which imply that

$$N([\mu,\nu];\mathbf{A}_1(\mathbf{k})) = N([\mu,\nu];\mathbf{A}_2(\mathbf{k}))$$

By continuity, for every $\mathbf{k} \in \mathcal{O}$ there is $s_{\mathbf{k}} > 0$ so that whenever μ is δ -distant at \mathbf{k} , μ is also $\delta/2$ distant at every $\mathbf{k}' \in \mathcal{B}(\mathbf{k}, s_{\mathbf{k}})$. This also implies that if (9.7) holds at \mathbf{k} for one of those μ , it also holds for that μ at every $\mathbf{k}' \in \mathcal{B}(\mathbf{k}, s_{\mathbf{k}})$, and therefore at every $v \delta/2$ -distant at \mathbf{k}' .

By compactness of \mathcal{O} , there are $\mathbf{k}_1, \dots, \mathbf{k}_\ell$ such that \mathcal{O} is covered by the balls $U_j = \mathscr{B}(\mathbf{k}_j, s_{\mathbf{k}_j}/2)$, with $0 \le j \le \ell$. If $U_j \cap U_{j'} \ne \emptyset$, we have that $\mathbf{k}_{j'} \in \mathscr{B}(\mathbf{k}_j, s_{\mathbf{k}_j})$, so that if (9.7) holds for some μ δ -distant at \mathbf{k}_j then it also holds for all $v \delta/2$ -distant at $\mathbf{k}_{j'}$, and therefore also at any $\mathbf{k}' \in U_{j'}$. By connectedness of \mathcal{O} , this means that (9.7) only needs to be verified for some \mathbf{k}_j , $0 \le j \le \ell$ and one $\mu \delta/2$ -distant at \mathbf{k}_j . Choosing \mathbf{k}_0 and μ_0 , this means that (9.7) holds everywhere.

Suppose now that for some $p \in \{1,2\}$ there is some $j \in J_p$ and $\mathbf{k} \in \mathcal{O}$ such that $\lambda_j(\mathbf{A}_p(\mathbf{k})) \in [\lambda - t, \lambda + t]$ and

$$\lambda_{j}(\mathbf{A}_{1}(\mathbf{k})) - \lambda_{j}(\mathbf{A}_{2}(\mathbf{k})) > \varepsilon.$$

Let μ be δ -distant at **k**. Without loss of generality assume that $[\mu, \infty) \cap \operatorname{spec}(\mathbf{A}_p) \neq \emptyset$ and let

$$j' = \min\left\{\ell : \lambda_{\ell}(\mathbf{A}_p)(\mathbf{k}) > \mu\right\}$$

Supposing that $j \ge j'$, and using (9.6) we obtain

$$N\left(\left[\mu,\lambda_{j}(\mathbf{A}_{2}(\mathbf{k}))\right];\mathbf{A}_{2}(\mathbf{k})\right) \geq j+1-j' > N\left(\left[\mu,\lambda_{j}(\mathbf{A}_{2}(\mathbf{k}))\right]_{\varepsilon};\mathbf{A}_{1}(\mathbf{k})\right),$$

which contradicts the second inequality in (9.4). Similarly, supposing that $\lambda_j(\mathbf{A}_2(\mathbf{k})) - \lambda_j(\mathbf{A}_1(\mathbf{k})) > \varepsilon$ contradicts the first inequality in (9.4). The case j < j' is treated analogously. We can therefore deduce that (9.5) holds at every $\mathbf{k} \in \mathcal{O}^{\dagger}$.

The previous lemma admits the following corollary in the situation where the difference $A_1 - A_2$ is a bounded operator. The reader interested solely in this case might notice that the proofs of the statement could have been more direct on its own.

Corollary 9.4. Let $\mathbf{A} = \mathbf{A}_0 + \mathbf{B}$ be a self-adjoint unbounded periodic operator such that $\mathbf{A}(\mathbf{k})$ has discrete spectrum for all $\mathbf{k} \in \mathcal{O}$ and such that \mathbf{B} is bounded. Then, there exist labelings $\lambda_j(\mathbf{A}_0(\mathbf{k}))$ and $\lambda_j(\mathbf{A}(\mathbf{k}))$ of the eigenvalues of the fibre operators such that the functions $\lambda_j(\mathbf{A}_0(\cdot))$ and $\lambda_j(\mathbf{A}(\cdot))$ are both continuous on \mathcal{O} and such that for every $\mathbf{k} \in \mathcal{O}$

$$\left|\lambda_{j}(\mathbf{A}_{0}(\mathbf{k}) - \lambda_{j}(\mathbf{A}(\mathbf{k}))\right| \leq \|\mathbf{B}\|.$$

Proof. It suffices to observe that $\|\mathbf{B}(\mathbf{k})\| \le \|\mathbf{B}\|$ for all $\mathbf{k} \in \mathcal{O}$. Defining the continuous family of operators $\mathbf{A}_t = \mathbf{A}_0 + t\mathbf{B}$, it is easy to see that $\mathbf{A}_1 = \mathbf{A}$ and $\|\mathbf{A}_t - \mathbf{A}_s\| = |t - s| \|\mathbf{B}\|$. From Lemma 4.11, we know that for all $I \subset \mathbb{R}$, (9.4) holds for $\mathbf{A}_s, \mathbf{A}_t$ with $\varepsilon = |t - s| \|\mathbf{B}\|$. It is also clear that δ defined in (9.3) is continuous in the parameter *t*. Setting

$$N = \left\lceil \frac{\|\mathbf{B}\|}{\min_{0 \le t \le 1} \varepsilon_t} \right\rceil,$$

and applying recursively Lemma 9.2 to the operators $\mathbf{A}_{j/N}$ and $\mathbf{A}_{(j+1)/N}$, with $0 \le j < N$ yields the result we seek.

The previous lemma and corollary provide us with an explicit way to compare the overlap function. This is made precise in the following proposition.

Proposition 9.5. Suppose that $\mathbf{A}_1, \mathbf{A}_2$ are self-adjoint, periodic operators such that for all $p \in \{1, 2\}$, $\mathbf{k} \in \mathcal{O} \mathbf{A}_p(\mathbf{k})$ has discrete spectrum. Suppose that for $\varepsilon > 0$ there is a non-decreasing labelling of their eigenvalues so that whenever $\lambda_j(\mathbf{A}_p(\mathbf{k})) \in [\lambda - 4\zeta(\lambda; \mathbf{A}_1), \lambda + 4\zeta(\lambda; \mathbf{A}_1)]$ we have

$$\left|\lambda_{j}(\mathbf{A}_{1}(\mathbf{k})) - \lambda_{j}(\mathbf{A}_{2}(\mathbf{k}))\right| \leq \varepsilon.$$
(9.8)

Then,

$$\zeta(\lambda; \mathbf{A}_2) \ge \zeta(\lambda; \mathbf{A}_1) - 2\varepsilon.$$

Proof. If $2\varepsilon > \zeta(\lambda; \mathbf{A}_1)$, the result follows trivially from nonnegativity of the overlap function. Otherwise, choose $j \in \mathbb{Z}$ such that

$$[\lambda - \zeta(\lambda; \mathbf{A}_1), \lambda + \zeta(\lambda; \mathbf{A}_1)] \subset \iota_i(\mathbf{A}_1).$$

Then (9.8) implies

$$[\lambda - \zeta(\lambda; \mathbf{A}_1) + \varepsilon, \lambda + \zeta(\lambda; \mathbf{A}_1) - \varepsilon] \subset \iota_i(\mathbf{A}_1)_{-\varepsilon} \subset \iota_i(\mathbf{A}_2).$$

The claim then follows from inspection of the definition (9.2) of the overlap function. \Box

10. Systems of periodic operators – The Bethe-Sommerfeld property

In this section we prove that certain systems of periodic operators enjoy the Bethe-Sommerfeld property in a quantitative way. This will imply that the spectrum of an elliptic periodic operator **A** in some classes contains a half-line. Our proof is again based on a reduction of the problem to uncoupled operators.

It is clear that if we show that the overlap function (9.2) is bounded away from 0 at sufficiently large λ for some operator **A**, then **A** has the Bethe–Sommerfeld property. This is the strategy employed in [PS10], where the self-adjoint operators of the form

$$A = (-\Delta)^{\alpha} + B,$$

with $B \in \mathbf{S}^{\beta}$, $\beta < 2\alpha$, and *B* is Λ -periodic are studied. It is shown in [PS10] that there are *S*, *c* and λ_0 , depending only on Θ and the symbol norms of *B*, such that for all $\lambda \ge \lambda_0$, $\zeta(\lambda; H) \ge c\lambda^S$. However, unlike in the situation of Proposition 8.4, it is no longer possible to simply extend the result to uncoupled operators, i.e. direct sums of operators acting on scalar functions. While it is certainly true that such operators enjoy the Bethe–Sommerfeld property, it could be possible *a priori* that the overlap function does not stay bounded away from 0 when direct sums are taken. This would imply that the reduction to uncoupled operators would be able to open gaps. Our aim is to show that such a situation is impossible for our class of operators.

10.1. **The Bethe-Sommerfeld property.** Our main theorem concerning systems of periodic operators is the following.

Theorem 10.1. Suppose that $\mathbf{A} \in \mathbf{ES}_m^{\alpha}$, $\alpha > 0$ is periodic, self-adjoint competely uncoupleable operator with $\mathbf{A}^{\mathcal{D}}$ of the form (7.1) and $a_j \neq a_k$ whenever $j \neq k$. Then, there exist positive $\tilde{\lambda}$, S, c such that

(1) *if* **A** *is unbounded above,* $[\tilde{\lambda}, \infty) \subset \operatorname{spec}(\mathbf{A})$ *and for every* $\lambda \geq \tilde{\lambda}$ *,* $\zeta(\lambda; \mathbf{A}) \geq c\lambda^{-S}$ *;*

(2) *if A is unbounded below*, $(-\infty, -\tilde{\lambda}] \subset \operatorname{spec}(\mathbf{A})$ *and for every* $\lambda \geq \tilde{\lambda}$, $\zeta(-\lambda; \mathbf{A}) \geq c\lambda^{-S}$.

The overlap exponent S depends only on α and the dimension d. The parameters $\tilde{\lambda}$ and c can be chosen uniformly in the symbol norms of **A** and $\mathbf{A}^{\mathscr{OD}}$.

Remark 10.2. Saying that the parameters are chosen uniformly in the symbol norms means that if **A** and **A**' are operators satisfying the conditions of Theorem 10.1 and for all $s, \ell, \|\mathbf{A}'\|_{\ell}^{(s)} \le \|\mathbf{A}\|_{\ell}^{(s)}$ and $\|\mathbf{A}'^{\mathcal{OD}}\|_{\ell}^{(s)} \le \|\mathbf{A}^{\mathcal{OD}}\|_{\ell}^{(s)}$ then the parameters obtained for **A** also work for **A**'.

Let us first observe that, as was the case in Sections 7.2 and 8, it is sufficient to prove case (1), case (2) will then follow from the former applied to the operator -A.

To prove Theorem 10.1, we proceed in two steps. The first one is the following proposition, the proof of which is delayed until Section 11.

Proposition 10.3. Let $\mathbf{A} \in \mathbf{US}_m^{\alpha} \cap \mathbf{ES}_m^{\alpha}$, $\alpha > 0$ be periodic, essentially self-adjoint, with $\mathbf{A}^{\mathcal{D}}$ of the form (7.1). Then, there exist $\tilde{\lambda}, S, c > 0$ such that

- (1) *if* **A** *is unbounded above, the interval* $[\lambda, \infty) \subset \operatorname{spec}(\mathbf{A})$ *and for every* $\lambda \geq \lambda, \zeta(\lambda; \mathbf{A}) \geq c\lambda^{-S}$ *;*
- (2) if **A** is unbounded below, the interval $(-\infty, -\tilde{\lambda}] \subset \operatorname{spec}(\mathbf{A})$ and for every $\lambda \geq \tilde{\lambda}$, $\zeta(-\lambda; \mathbf{A}) \geq c\lambda^{-S}$.

The overlap exponent S depends only on α and d. The parameters $\tilde{\lambda}$ and c can be chosen uniformly in the symbol norms of **A** and \mathbf{A}^{OD} .

The second step on the way to proving Theorem 10.1 is to show that small perturbations of an operator (in the sense of perturbations that do not change the integrated density of states much) cannot open a gap if the overlap function is large enough.

Lemma 10.4. Suppose that $\mathbf{A} \in \mathbf{ES}_m^{\alpha}$ satisfies the hypotheses of Theorem 10.1. Then, for every $K \in \mathbb{R}$, there exists an operator $\mathbf{A}_K \in \mathbf{US}_m^{\alpha} \cap \mathbf{SES}_m^{\alpha}$ satisfying the hypothesis of Proposition 10.3 such that for every λ large enough, we have that

$$\zeta(\lambda; \mathbf{A}) \ge \zeta(\lambda; \mathbf{A}_K) + O\left(\lambda^{-K}\right). \tag{10.1}$$

Proof. From Remark 9.1 and Theorem 8.9, it is possible for any *K* to find an operator $\mathbf{A}_K \in \mathbf{US}_m^{\alpha}$ such that for $\varepsilon = \lambda^{-\alpha-K}$ and any interval $I \subset [\lambda - 2\zeta(\lambda; \mathbf{A}), \lambda + 2\zeta(\lambda; \mathbf{A})]$ we have that for every $\mathbf{k} \in \mathcal{O}^{\dagger}$,

$$N(I;\mathbf{A}(\mathbf{k})) \le N(I_{\varepsilon};\mathbf{A}_{K}(\mathbf{k})) \quad \text{and} \quad N(I;\mathbf{A}_{K}(\mathbf{k})) \le N(I_{\varepsilon};\mathbf{A}(\mathbf{k})).$$
(10.2)

As in (9.3), put

$$\begin{split} \delta &= \min_{\mathbf{k} \in \mathcal{O}} \max \left\{ \operatorname{dist}(\mu; \operatorname{spec}(\mathbf{A}(\mathbf{k}))) : \mu \in [\lambda - 2\zeta(\lambda; \mathbf{A}), \lambda + 2\zeta(\lambda; \mathbf{A})] \right\} \\ &\geq C \frac{\zeta(\lambda; \mathbf{A})}{\max_{\mathbf{k}} N([\lambda - 2\zeta(\lambda; \mathbf{A}), \lambda + 2\zeta(\lambda; \mathbf{A})]; \mathbf{A}(\mathbf{k}))} \end{split}$$

for some C > 0. By Lemma 9.2 and Proposition 9.5, if $\varepsilon < \delta/4$, then (10.2) implies (10.1). It follows from Proposition 10.3 that $\zeta(\lambda; \mathbf{A}) \ge \lambda^{-S}$ for some *S*. Weyl's law implies that

$$\max_{\mathbf{k}} N([\lambda - 2\zeta(\lambda; \mathbf{A}), \lambda + 2\zeta(\lambda; \mathbf{A})]; \mathbf{A}(\mathbf{k})) = O\left(\lambda^{d/\alpha}\right).$$

It follows that by choosing $K > S + \frac{d}{\alpha}$, we have $\varepsilon < \delta/4$ for λ large enough, finishing the proof.

64

Before making our constructions explicit in the next section, we prove the following lemma which indicates that for uncoupled operators, we may suppose without loss of generality that they are semi-bounded below.

Lemma 10.5. Let $\mathbf{A} \in \mathbf{US}_m^{\alpha} \cap \mathbf{SES}_m^{\alpha}$ is self-adjoint and periodic, and suppose that

$$\mathbf{A} = \mathbf{A}_+ \oplus \mathbf{A}_-$$

with A_+ semi-bounded below and A_- semi-bounded above. Then, Theorem 10.1, part (1) holds for A if and only if it holds for A_+ . Similarly, Theorem 10.1, part (2) holds if and only if it holds for A_- .

Proof. We prove it for the implication of Theorem 10.1 part (1), the other one follows by replacing **A** by $-\mathbf{A}$. Since \mathbf{A}_{-} is semi-bounded above, there is some M > 0 such that $\operatorname{spec}(\mathbf{A}_{-}) \cap (M, \infty) = \emptyset$. It therefore follows that for $\lambda > M$,

$$\operatorname{spec}(\mathbf{A}_+) \cap [\lambda, \infty) = \operatorname{spec}(\mathbf{A}) \cap [\lambda, \infty)$$

and our claim follows.

11. Bethe-Sommerfeld for uncoupled operators

In this section we prove Proposition 10.3. Remark 9.1, Lemmas 9.2 and 10.5; and Theorem 8.9 tell us that it is sufficient to prove Proposition 10.3 after a few simplifying assumptions. First, from now on we assume that $\mathbf{A} \in \mathbf{US}_m^{\alpha} \cap \mathbf{SES}_m^{\alpha}$ is semi-bounded below, in other words $a_j > 0$ for all $1 \le j \le m$. Then, we also suppose that we have already applied the gauge transform in each component, and that we have removed the remainder that changes the IDS by $O(\rho^{-K})$ for some large *K*. In other words, we suppose that the symbol of **A** is given by

$$\mathbf{a}(\mathbf{x},\boldsymbol{\xi}) = \mathbf{a}^{\mathscr{D}}(\boldsymbol{\xi}) + \mathbf{b}^{\mathscr{R}}(\mathbf{x},\boldsymbol{\xi}),$$

where we have abused notation and used the same symbol for **A** and the operator after gauge transform.

The study of the Bethe-Sommerfeld property in high dimensions [BP09, Par08, PS10] relies on variations of a certain combinatorial geometric argument. The strategy goes as follows, using the notation from [PS10]. First, find a function $g : \mathbb{R}^d \to \operatorname{spec}(\mathbf{A})$ such that on fibres of the quotient map $\mathbb{R}^d \to \mathbb{R}^d / \Theta$, g is a bijection onto $\operatorname{spec}(\mathbf{A}(\mathbf{k}))$. Such a function can be chosen in such a way that

$$g(\boldsymbol{\xi}) - |\boldsymbol{\xi}|^{\alpha} = o(|\boldsymbol{\xi}|^{\alpha}).$$

The goal is then to show that for any large energy ρ^{α} the pre-image of some small interval $[\rho^{\alpha} - \delta, \rho^{\alpha} + \delta]$ contains a curve connecting $g^{-1}(\rho^{\alpha} - \delta)$ and $g^{-1}(\rho^{\alpha} + \delta)$, and that *g* is continuous along

that curve. Here, δ is a parameter that depends on ρ and corresponds to the overlap length at energy level ρ^{α} .

In order to show that such a curve exists we split the pre-image $g^{-1}([\rho^{\alpha} - \delta, \rho^{\alpha} + \delta])$ into bad (resonant) and good (non-resonant) regions. The latter are defined in such a way that *g* is continuous and radially increasing within them. The idea in [BP09, Par08, PS10] was to find a small radial interval in the non-resonant region where the eigenvalues of **A**(**k**) are all simple along that small interval, which in turn gives a lower bound for the overlap function.

This is done by showing that not only is the angular measure of the resonant region small, but that the total measure of intersections of translates of the resonant regions by elements of Θ with the non-resonant region is also small, in comparison to the volume of the non-resonant region. This implies the existence of a radial interval along which translations by $\theta \in \Theta$ stay inside the non-resonant region, which gives us the desired interval where the eigenvalues of $A(\mathbf{k})$ are simple and increasing.

In the case of an operator $\mathbf{A} \in \mathbf{US}_m^{\alpha}$ acting in $L^2(\mathbb{R}^d; \mathbb{C}^m)$, hoping for the eigenvalues of $\mathbf{A}(\mathbf{k})$ to be simple is unrealistic. However, such a requirement is not necessary. Indeed, suppose that the eigenvalues of operator H acting in $L^2(\mathbb{R}^d; \mathbb{C})$ are simple. Then, since the spectrum of $H \oplus H$ is the same as for H, $H \oplus H$ enjoys the Bethe–Sommerfeld property if and only if it H also does. However, the eigenvalues of $(H \oplus H)(\mathbf{k})$ are always at least double. This means that we need a new way of reasoning. We obtain the following conditions, which are sufficient to obtain lower bounds on the overlap function.

Proposition 11.1. Suppose that there is an interval $[\mathbf{k}_1, \mathbf{k}_2] \subset \mathcal{O}^{\dagger}$ and three families of real-valued continuous functions $\boldsymbol{\mu}, \boldsymbol{\nu}$ and $\boldsymbol{\tau}$ on $[\mathbf{k}_1, \mathbf{k}_2]$, satisfying the following properties.

• The equality of multisets (i.e. taking multiplicity into account)

$$\{\lambda_{p}(\mathbf{A}(\mathbf{k})); p \in \mathbb{N}\} = \{\mu(\mathbf{k}) : \mu \in \boldsymbol{\mu}\} \cup \{\nu(\mathbf{k}) : \nu \in \boldsymbol{\nu}\} \cup \cup \{\tau(\mathbf{k}) : \tau \in \boldsymbol{\tau}\}$$
(11.1)

holds for all $\mathbf{k} \in [\mathbf{k}_1, \mathbf{k}_2]$.

• For every $\mu \in \mu$, μ is increasing as **k** goes from \mathbf{k}_1 to \mathbf{k}_2 . Furthermore, either

 $\mu(\mathbf{k}_1) < \rho^{\alpha} - \delta \qquad or \qquad \mu(\mathbf{k}_2) > \rho^{\alpha} + \delta. \tag{11.2}$

• We have that $\#\mathbf{v} < \infty$. Furthermore,

$$\#\left\{\mu \in \boldsymbol{\mu} : [\rho^{\alpha} - \delta, \rho^{\alpha} + \delta] \subset \mu([\mathbf{k}_1, \mathbf{k}_2])\right\} > \#\boldsymbol{\nu}.$$
(11.3)

• For all $\tau \in \boldsymbol{\tau}$, we have that

$$\tau([\mathbf{k}_1, \mathbf{k}_2]) \cap [\rho^{\alpha} - \delta, \rho^{\alpha} + \delta] = \emptyset.$$
(11.4)

Then, $\zeta(\rho^{\alpha}; \mathbf{A}) \geq \delta$.

Proof. Since **A** is semi-bounded below, we can use Skriganov's characterisation [Skr85] of the overlap function as

$$\zeta(\rho^{\alpha}, \mathbf{A}) = \sup_{\mathbf{k}} \left\{ t : \min_{\mathbf{k}} N((-\infty, \rho^{\alpha} + t]; \mathbf{A}(\mathbf{k})) < \max_{\mathbf{k}} N((-\infty, \rho^{\alpha} - t); \mathbf{A}(\mathbf{k})) \right\}.$$
(11.5)

It then follows that if

$$N(\rho^{\alpha} - \delta; \mathbf{A}(\mathbf{k}_1)) - N(\rho^{\alpha} + \delta; \mathbf{A}(\mathbf{k}_2)) \ge 1,$$
(11.6)

then $\zeta(\rho^{\alpha}; \mathbf{A}) \ge \delta$. Equation (11.1) implies that we can separate the differences of each family $\boldsymbol{\mu}$, $\boldsymbol{\nu}$ and $\boldsymbol{\tau}$ in (11.6). Equation (11.4) ensures that

$$\#\left\{\tau \in \boldsymbol{\tau} : \tau(\mathbf{k}_1) \le \rho^{\alpha} - \delta\right\} - \#\left\{\tau \in \boldsymbol{\tau} : \tau(\mathbf{k}_2) \le \rho^{\alpha} + \delta\right\} = 0.$$
(11.7)

We also have that

$$\left| \# \left\{ v \in \boldsymbol{v} : v(\mathbf{k}_1) < \rho^{\alpha} - \delta \right\} - \# \left\{ v \in \boldsymbol{v} : v(\mathbf{k}_2) \le \rho^{\alpha} + \delta \right\} \right| \le \# \boldsymbol{v}.$$

Finally, from (11.2), (11.3) and the fact that all functions $\mu \in \mu$ are increasing, we have that

$$\#\left\{\boldsymbol{\mu} \in \boldsymbol{\mu} : \boldsymbol{\mu}(\mathbf{k}_1) \le \rho^{\alpha} - \delta\right\} - \#\left\{\boldsymbol{\mu} \in \boldsymbol{\mu} : \boldsymbol{\mu}(\mathbf{k}_2) \le \rho^{\alpha} + \delta\right\} > \#\boldsymbol{\nu}.$$
(11.8)

Indeed, assumption (11.2) ensures that every $\mu \in \mu$ contributes at least 0 to the lefthand side of (11.8), whereas (11.3) ensures that there at least $\#\nu + 1$ functions in μ contributing 1. Combining (11.7)–(11.8) yields (11.6).

In the remainder of this section, we therefore set out to prove the existence of the interval $[\mathbf{k}_1, \mathbf{k}_2]$ as well as of the families of functions $\boldsymbol{\mu}$, $\boldsymbol{\nu}$ and $\boldsymbol{\tau}$ satisfying the hypotheses of Proposition 11.1. We will, as we go along, use relevant results from [PS10], mostly about properties of the function *g* and volume of various sets associated with the preimage.

11.1. A description of the spectrum. In [PS10], Parnovski and Sobolev have constructed functions $g_j : \mathbb{R}^d \to \operatorname{spec}(A_j)$. When restricted to a coset $\{\xi\} + \Theta$, g_j is a bijection (respecting multiplicity) with $\operatorname{spec}(A_j(\{\xi\}))$. It is useful to use g_j to define a family of functions on the fundamental cell \mathcal{O} indexed by $\mathbf{p} := (j, \boldsymbol{\theta}) \in \{1, ..., m\} \times \Theta =: \widetilde{\Theta}$ as

$$g_{\mathbf{p}}(\mathbf{k}) := g_j(\mathbf{k} + \boldsymbol{\theta})$$

Since we assume in this section that

$$\mathbf{A} = A_1 \oplus \ldots \oplus A_m \in \mathbf{US}_m^{\alpha},$$

and that is is a semi-bounded below operator, we have by construction that $\{\lambda_p(\mathbf{A}(\mathbf{k})) : p \in \mathbb{N}\}\$ is a relabeling of $\{g_{(j,\theta)}(\mathbf{k}) : (j,\theta) \in \widetilde{\Theta}\}\$ in increasing order. In particular, the latter set is discrete, bounded from below and accumulates only at infinity.

To prove Proposition 10.3, it remains to show that there exists $S \in \mathbb{R}$ such that the hypotheses of Proposition 11.1 hold for some $\delta \gg \rho^S$. The families μ , ν and τ will be realised as a partition of the set $\{g_{\mathbf{p}} : \mathbf{p} \in \widetilde{\Theta}\}$.

Proposition 11.2. There exist $\rho_0 > 0$ and $S \in \mathbb{R}$ (depending on $\{a_1, ..., a_m\}$, α and the implicit constants in (11.12)) such that for all $\rho \ge \rho_0$, there exists an interval $[\mathbf{k}_1, \mathbf{k}_2]$ and three families of functions $\boldsymbol{\mu}, \boldsymbol{\nu}, \boldsymbol{\tau}$ on $[\mathbf{k}_1, \mathbf{k}_2]$ satisfying the hypotheses (11.1)–(11.4) of Proposition 11.1.

The remainder of this section is dedicated to the proof of this proposition. We give some useful properties of the functions g_i . For $1 \le j \le m$, there are $a_i > 0$ and $G_i : \mathbb{R}^d \to \mathbb{R}$ such that

$$g_{j}(\boldsymbol{\xi}) = a_{j} |\boldsymbol{\xi}|^{\alpha} + G_{j}(\boldsymbol{\xi}), \qquad a_{j} > 0.$$
 (11.9)

Furthermore, whenever

$$|\boldsymbol{\xi}| \approx \rho, \tag{11.10}$$

we have

 $\left|G_{j}(\boldsymbol{\xi})\right| \ll \rho^{\beta}$

for some $\beta < \alpha$, the implicit constant depending on the ones in (11.10).

We need to control the first and second derivatives of the functions G_j on a large enough region. To this end, for $\delta \in (0, \rho^{\alpha}/4)$, define the "annular" regions

$$\mathscr{A}_j := \mathscr{A}_j(\rho; \delta) := g_j^{-1} \left(\left[\rho^{\alpha} - \delta, \rho^{\alpha} + \delta \right] \right).$$

We suppose that for every $0 < \delta < \rho^{\alpha}/4$, \mathcal{A}_i can be decomposed further into a non-resonant set

$$\mathscr{B}_j := \mathscr{B}_j(\rho, \delta) \subset \mathscr{A}_j$$

and a resonant set

$$\mathscr{R}_j := \mathscr{A}_j \setminus \mathscr{B}_j.$$

A precise description of those sets is given in [PS10]; the non-resonant sets \mathscr{B}_j correspond to simple eigenvalues of the operators $A_j(\{\xi\})$, whereas the resonant sets \mathscr{R}_j correspond to clusters of eigenvalues. In particular, when $\xi \in \mathscr{B}_j$,

$$\mathbf{A}\boldsymbol{e}_{\boldsymbol{\xi}} \otimes \boldsymbol{v}_{j} = \boldsymbol{g}_{j}(\boldsymbol{\xi})\boldsymbol{e}_{\boldsymbol{\xi}} \otimes \boldsymbol{v}_{j}. \tag{11.11}$$

An important property of the non-resonant set is that the functions g_j behave well there. We suppose that the restriction of G_j to \mathscr{B}_j is of class C^2 and that for all $\boldsymbol{\xi} \in \mathscr{B}'_j$, the bounds

$$\begin{aligned} \left| \nabla G_j(\boldsymbol{\xi}) \right| \ll \rho^{\gamma}, \\ \left| \nabla^2 G_j(\boldsymbol{\xi}) \right| \ll \rho^{\sigma} \end{aligned} \tag{11.12}$$

hold, where $\gamma < \alpha - 1$ and $\sigma < \alpha - 2$, and the implicit constants are once again allowed to depend on the implicit constants in (11.10). In order to prove Proposition 11.2, it will be sufficient to show that the sets \mathscr{B}_j are, in a sense, large enough. In subsection 11.2 we constrain the radial projection on the sphere of those resonant and non-resonant sets.

11.2. **Description of the resonant sets.** The description of the resonant sets in this section expands on the results in [PS10, Sections 5,7,8,9]. We first introduce some notation. Most of the sets we define depend on the parameters ρ and δ ; however, we will be quick to drop the (explicit) dependence on these parameters to make notation lighter. For every $\boldsymbol{\xi} \in \mathbb{R}^d \setminus \{0\}$, let $\mathbf{u}_{\boldsymbol{\xi}} := |\boldsymbol{\xi}|^{-1} \boldsymbol{\xi}$ be the unit vector in the direction of $\boldsymbol{\xi}$. For $\nu > 0$ and $\boldsymbol{\theta} \in \Theta \setminus \{0\}$, we define spherical resonant regions as

$$\mathscr{S}(\boldsymbol{\theta};T) := \left\{ \boldsymbol{\zeta} \in \mathbb{S}^{d-1} : |\boldsymbol{\zeta} \cdot \mathbf{u}_{\boldsymbol{\theta}}| < T \right\}.$$

For any subset \mathscr{U} of the sphere \mathbb{S}^{d-1} , we denote its radial extension by

$$\mathscr{U}_{rd} := \left\{ \boldsymbol{\xi} \in \mathbb{R}^d : \mathbf{u}_{\boldsymbol{\xi}} \in \mathscr{U} \right\}$$

The following lemma is proved in [PS10, Section 5].

Lemma 11.3. There exists c (depending on the coefficients a_j) such that for every ρ large enough, and $\delta \in (0, c\rho^{\alpha})$, there are κ , and ν such that

$$0 < \kappa < d^{-2}, \qquad \Theta_{\kappa} := (\Theta \setminus \{0\}) \cap B(\rho^{\kappa}), \qquad \kappa d < \nu < 1$$

and such that for all $1 \le j \le m$,

$$\mathscr{R}_{j}(\rho;\delta) \subset \mathscr{A}_{j} \cap \bigcup_{\boldsymbol{\theta} \in \Theta_{\kappa}} \mathscr{S}(\boldsymbol{\theta};\rho^{-\nu})_{rd}.$$
(11.13)

Denoting

$$\mathcal{T}(\rho) := \mathbb{S}^{d-1} \setminus \bigcup_{\boldsymbol{\theta} \in \Theta_{\kappa}} \mathscr{S}(\boldsymbol{\theta}; \rho^{-\nu})$$

and

$$\widetilde{\mathscr{B}}_{j}(\rho) := \mathscr{A}_{j} \cap \mathscr{T}_{rd},$$

we also have that $\widetilde{\mathscr{B}}_j \subset \mathscr{B}_j$.

Remark 11.4. The inclusion (11.13) means that the radial projection to the sphere of the resonant region is included in the union of the sets $\mathscr{S}(\theta; \rho^{-\nu})$, justifying naming them spherical resonant regions. Consequently, we name \mathscr{T} the spherical non-resonant region.

The object of the next lemma is that small enough neighborhoods of the spherical resonant regions have small volume.

Lemma 11.5. Let κ , Θ_{κ} and ν be obtained in Lemma 11.3 and define

$$\widetilde{\mathscr{T}}(\rho) := \mathbb{S}^{d-1} \setminus \left(\bigcup_{\boldsymbol{\theta} \in \Theta_{\kappa}} \mathscr{S}(\boldsymbol{\theta}; 2\rho^{-\nu}) \right),$$
$$\mathscr{Z}_{j}(\rho; \delta) := \mathscr{A}_{j} \cap \bigcup_{\boldsymbol{\theta} \in \Theta_{\kappa}} \mathscr{S}_{rd}(\boldsymbol{\theta}; 2\rho^{-\nu}),$$
(11.14)

and

$$\mathscr{G}_{j}(\rho;\delta) := \mathscr{A}_{j} \cap \widetilde{\mathscr{T}}_{rd} = \mathscr{A}_{j} \setminus \mathscr{Z}_{j}.$$
(11.15)

Then, $\mathscr{G}_j \subset \mathscr{B}_j$, and for all $\boldsymbol{\xi} \in \mathscr{G}_j$, dist $(\boldsymbol{\xi}, \mathscr{R}_j) \gg \rho^{1-\nu}$. *Furthermore, for* $\varepsilon_0 = \nu - d\kappa > 0$,

$$\operatorname{vol}(\mathcal{Z}_{j}(\rho;\delta)) \ll \delta \rho^{d-\alpha-\varepsilon_{0}}.$$
 (11.16)

and

$$\operatorname{vol}(\mathscr{G}_{i}(\rho;\delta)) \asymp \delta \rho^{d-\alpha}.$$
 (11.17)

Proof. It is clear from the definition that $\mathscr{G}_j \subset \widetilde{\mathscr{B}}_j \subset \mathscr{B}_j$. Define the sets

$$\mathscr{I}_{i}(\boldsymbol{\zeta}) := \mathscr{A}_{i} \cap \{\boldsymbol{\zeta}\}_{rd}$$

It follows from [PS10, pp.518–519] that for all ζ , $\mathscr{I}_j(\zeta)$ is an interval of length $|\mathscr{I}_j| \ll \delta \rho^{1-\alpha}$ (uniformly in ζ), and

$$\mathscr{I}_{j} \subset \left\{ \boldsymbol{\xi} : |\boldsymbol{\xi}| \asymp \rho \right\}.$$

Furthermore, by definition

dist
$$\left(\widetilde{\mathcal{T}}(\rho), \bigcup_{\boldsymbol{\theta}\in\Theta_{\kappa}}\mathscr{S}(\boldsymbol{\theta}; \rho^{-\nu})\right) > \rho^{-\nu}.$$

It therefore follows from (11.13), (11.15) and basic trigonometry that $dist(\mathscr{G}_j, \mathscr{R}_j) \gg \rho^{1-\nu}$. For the volume estimate for \mathscr{Z}_j , we compute

$$\operatorname{vol}(\mathcal{Z}_{j}) \leq \sum_{\boldsymbol{\theta} \in \Theta_{\kappa}} \int_{\widetilde{\mathscr{S}}(\boldsymbol{\theta}; \rho^{-\nu})} \int_{\mathscr{I}_{j}(\boldsymbol{\zeta})} t^{d-1} dt d\boldsymbol{\zeta}$$

$$\ll \#(\Theta_{\kappa}) \max_{\boldsymbol{\theta}} \operatorname{vol}_{d-1}(\mathscr{S}(\boldsymbol{\theta}; 2\rho^{-\nu})) \delta \rho^{d-\alpha}.$$
(11.18)

Uniformly in $\boldsymbol{\theta}$ we have that, $\operatorname{vol}_{d-1}(\mathscr{S}(\boldsymbol{\theta}; 2\rho^{-\nu}) \ll \rho^{-\nu})$. We also have that $\#\Theta_{\kappa} \ll \rho^{d\kappa}$. Putting these two estimates in (11.18) yields (11.16). For the estimate on $\operatorname{vol}(\mathscr{G}_{i}(\rho; \delta))$, we observe that

 $\operatorname{vol}(\mathscr{G}_i) = \operatorname{vol}(\mathscr{A}_i) - \operatorname{vol}(\mathscr{Z}_i)$ and that by (11.9),

$$\operatorname{vol}(\mathscr{A}_i) \cong \delta \rho^{d-\alpha}$$

Estimate (11.17) then follows from the fact that $\operatorname{vol}(\mathcal{Z}_i) = o(\operatorname{vol}(\mathcal{A}_i))$.

11.3. Volumes of intersections. For $\mathbf{b}_1, \mathbf{b}_2 \in \mathbb{R}^d$ and $i, j \in \{1, ..., m\}$ we define the *crossing sets*

$$\mathscr{X}_{ij}(\rho,\delta,\mathbf{b}_1,\mathbf{b}_2) := (\mathscr{A}_i(\rho;\delta) + \mathbf{b}_1) \cap (\mathscr{A}_j(\rho;\delta) + \mathbf{b}_2).$$

We are interested in volume estimates, and since

$$\operatorname{vol}(\mathscr{X}_{i\,i}(\rho,\delta,\mathbf{b}_1,\mathbf{b}_2)) = \operatorname{vol}(\mathscr{X}_{i\,i}(\rho,\delta,\mathbf{0},\mathbf{b}_2-\mathbf{b}_1)),$$

we restrict ourselves to sets of the form

$$\mathscr{X}_{ii}(\mathbf{b}) := \mathscr{X}_{ii}(\rho, \delta, \mathbf{b}) := \mathscr{X}_{ii}(\rho, \delta, \mathbf{0}, \mathbf{b}).$$

Denote by $\varphi(\mathbf{a}, \mathbf{b})$ the (smaller) angle between **a** and **b**. For any angle $\omega \in [0, \pi]$, we define the set

$$\mathscr{X}_{i\,j,\omega}(\mathbf{b}) := \left\{ \boldsymbol{\xi} \in \mathscr{X}_{i\,j}(\mathbf{b}) : \varphi(\boldsymbol{\xi}, \boldsymbol{\xi} - \mathbf{b}) > \omega \right\}.$$

From [PS10, Section 9] since g_i, g_j are defined as in (11.9), so that the conditions (11.12) are respected, the following holds: for any $\omega \in (0, \pi)$, $\varepsilon > 0$, if $\delta \rho^{2-\alpha+2\varepsilon} \to 0$ as $\rho \to \infty$, then

$$\operatorname{vol}(\mathscr{X}_{ij,\omega}(\rho,\delta,\mathbf{b})) \ll \delta^2 \rho^{4-2\alpha+d+6\varepsilon} + \delta \rho^{1-\alpha-\varepsilon(d-1)}, \tag{11.19}$$

uniformly in **b**.

Proposition 11.6. There are constants c and C depending on δ , ω , and the numbers a_j such that $\mathscr{X}_{ij,\omega}(\mathbf{b}) \neq \emptyset$ implies that $c\rho \leq |\mathbf{b}| \leq C\rho$.

Proof. We first make the observation that there exists C > 0, depending on α and the numbers a_j such that if $|\mathbf{b}| > C\rho$, then for ρ large enough $\mathscr{A}_i \cap (\mathscr{A}_j + \mathbf{b}) = \emptyset$. On the other hand, it follows from basic planar trigonometry that for every ω , there exists c, depending on the constants in $|\boldsymbol{\xi}| \simeq \rho$, such that if $|\mathbf{b}| < c\rho$, then $\varphi(\boldsymbol{\xi}, \boldsymbol{\xi} - \mathbf{b}) \le \omega$.

Remark 11.7. The estimates in (11.19) were established under stronger hypotheses in [PS10, Theorem 9.1]. This was because of the fact that their methods required control of the volume of $\mathscr{X}_{ij}(\mathbf{b})$, for all $1 \ll |\mathbf{b}| \ll \rho$, while we only need to control the volume of $\mathscr{X}_{ij,\omega}(\mathbf{b})$ for $|\mathbf{b}| \asymp \rho$. The stronger conditions were needed to treat the case of small angles, and small \mathbf{b} (that is, such that $|\mathbf{b}| = o(\rho)$). Our approach does not require any such estimates.

Before going on, let us make the following notational convention.

Convention. For any family of subsets $\mathscr{E}(\delta) \subset \mathbb{R}^d$ depending on the parameter $\delta > 0$, we denote

$$\mathscr{E}'(\delta) := \mathscr{E}(Z\delta),$$

where *Z* is some large constant to be determined later and depending only on the dimension *d*, the order α and the numbers $\{a_1, \ldots, a_m\}$.

We now define crossing sets for the non-resonant sets \mathscr{G}_i . For **b** $\in \mathbb{R}^d$, let

$$\mathscr{Y}_{i\,i}(\mathbf{b}) := \mathscr{G}_i \cap (\mathscr{G}_i + \mathbf{b})$$

and for any angle $\omega \in (0, \pi)$,

$$\mathscr{Y}_{ij,\omega}(\mathbf{b}) := \left\{ \boldsymbol{\xi} \in \mathscr{Y}_{ij}(\mathbf{b}) : \varphi(\boldsymbol{\xi}, \boldsymbol{\xi} - \mathbf{b}) > \omega \right\} = \mathscr{X}_{ij,\omega}(\mathbf{b}) \cap \mathscr{Y}_{ij}(\mathbf{b}).$$

We need the following lemma.

Lemma 11.8. For any $\omega \in (0, \pi)$ and $\varepsilon > 0$, the condition $\delta \rho^{2-\alpha+2\varepsilon} \to 0$ as $\rho \to \infty$ implies

$$\operatorname{vol}\left(\bigcup_{i,j=1}^{m}\bigcup_{\boldsymbol{\theta}\in\Theta}\mathscr{Y}'_{ij,\omega}(\boldsymbol{\theta})\right)\ll\delta^{2}\rho^{4-2\alpha+2d+6\varepsilon}+\delta\rho^{1-\alpha+d-\varepsilon(d-1)},$$

the implicit constants depending only on δ , ω , Z, and the coefficients a_i .

Proof. It is sufficient to prove the result for a single pair *i*, *j*, then sum the estimates over all m^2 of those pairs. From Proposition 11.6, there are constants *c* and *C* depending only on ω , δ , *T* and the numbers a_i such that

$$\operatorname{vol}\left(\bigcup_{\boldsymbol{\theta}\in\Theta}\mathscr{Y}'_{ij,\omega}(\boldsymbol{\theta})\right) \leq \sum_{\substack{\boldsymbol{\theta}\in\Theta\\c\rho\leq|\boldsymbol{\theta}|\leq C\rho}}\operatorname{vol}\left(\mathscr{Y}_{ij,\omega}(\boldsymbol{\theta})\right) \\ \ll \delta^2 \rho^{4-2\alpha+2d+6\varepsilon} + \delta \rho^{1-\alpha+d-\varepsilon(d-1)\varepsilon}$$

where the last line comes from $\mathscr{Y}_{ij,\omega}(\boldsymbol{\theta}) \subset \mathscr{X}_{ij,\omega}(\boldsymbol{\theta})$, estimate (11.19) and the fact that

$$\#\left\{\boldsymbol{\theta}\in\Theta^{\dagger}:c\rho\leq|\boldsymbol{\theta}|\leq C\rho\right\}\ll\rho^{d}.$$

11.4. Estimates on the overlap function. Define

$$\mathcal{Z} := \bigcup_{1 \le j \le m} \mathcal{Z}_j, \quad \text{and} \quad \mathcal{G} := \bigcup_{1 \le j \le m} \mathcal{G}_j.$$

It follows directly from the definitions of \mathcal{Z}_j (11.14) and \mathcal{G}_j (11.15) that $\mathcal{G} \cap \mathcal{Z} = \emptyset$. Furthermore, for every $\boldsymbol{\xi} \in \mathcal{G}$ and every $1 \le j \le m$, $\boldsymbol{\xi} \in \mathcal{G}_j$ or $\boldsymbol{\xi} \notin \mathcal{A}_j$.

For every $\boldsymbol{\xi} \in \mathbb{R}^d$, and any subset $E \subset \mathbb{R}^d$, we define

$$n(\boldsymbol{\xi}, E) = \# \{ \boldsymbol{\theta} \in \Theta : \boldsymbol{\xi} + \boldsymbol{\theta} \in E \}.$$

Let us take a step back to see what is needed in order to prove Proposition 11.2. The estimates on the functions G_j in the non-resonant regions \mathscr{B}_j ensure that g_j are radially increasing in those regions.

We need three ingredients in order to prove the existence of a path (of the form, before projection down to \mathcal{O} , $[t_1, t_2]\boldsymbol{\zeta}$ for some $\boldsymbol{\zeta} \in \mathbb{S}^{d-1}$) satisfying hypotheses (11.1) – (11.4) in Proposition 11.2.

- That for $\xi \in \mathcal{G}_j$, the functions g_j are increasing not only in the radial direction but also along directions deviating from the radial one by angles smaller than $\pi/4$, at least for some controllable distance.
- That there are points $\xi \in \mathcal{G}$ such that $n(\xi; \mathcal{G}) > n(\xi; \mathcal{Z}')$.
- That there are some of those $\boldsymbol{\xi}$ such that for all $\boldsymbol{\theta} \in \Theta' = \Theta \setminus \{0\}$ satisfying $\boldsymbol{\xi} + \boldsymbol{\theta} \in \mathcal{G}'$, we have that $\varphi(\boldsymbol{\xi}, \boldsymbol{\xi} + \boldsymbol{\theta}) < \pi/4$.

The first of those statements is proved directly. For the last two, we will show that the proportion of $\boldsymbol{\xi} \in \mathcal{G}$ such that either $n(\boldsymbol{\xi}; \mathcal{Z}') \ge n(\boldsymbol{\xi}; \mathcal{G})$ or for which there exists $\boldsymbol{\theta} \in \Theta'$ such that $\boldsymbol{\xi} + \boldsymbol{\theta} \in \mathcal{G}'$ and $\varphi(\boldsymbol{\xi}, \boldsymbol{\xi} + \boldsymbol{\theta}) > \pi/4$ is smaller than 1. Actually, for some appropriate choice of the free parameter δ , we will in fact show that this proportion can be made arbitrarily small.

The following lemma addresses the first bullet point.

Lemma 11.9. Let v be as obtained in Lemma 11.3. For all $\xi \in \mathcal{G}_j$, and all \mathbf{b} such that $|\xi + \mathbf{b}| = \rho$ and $\varphi(\xi, \xi + \mathbf{b}) \leq \pi/4$ there is a $t_0 \gg \rho^{-\nu}$ such that for all $t \in [-t_0, t_0]$ the function $t \mapsto g_j(\xi + t(\xi + \mathbf{b}))$ is increasing and

$$\frac{\mathrm{d}}{\mathrm{d}t}g_j(\boldsymbol{\xi}+t(\boldsymbol{\xi}+\mathbf{b}))\gg\rho^{\alpha}.$$

The implicit constant in $t_0 \gg \rho^{-\nu}$ depends only on the functions g_j and the implicit constants in $|\boldsymbol{\xi} + \mathbf{b}| \simeq \rho$.

Proof. Since $\boldsymbol{\xi} \in \mathcal{G}_j$, we have that not only $\boldsymbol{\xi} \in \mathcal{B}_j$ for some *j*, but also, by Lemma 11.5, that there exists r > 0 such that for all j', dist $(\boldsymbol{\xi}; \mathcal{R}_{j'}) > r\rho^{1-\nu}$. Therefore, for $|t| \le t_0 := r\rho^{-\nu}$, we have that $\boldsymbol{\xi} + t(\boldsymbol{\xi} + \mathbf{b}) \in \mathcal{B}'_j$. By (11.12), we have that

$$\left|\frac{\mathrm{d}}{\mathrm{d}t}G_j(\boldsymbol{\xi}+t(\boldsymbol{\xi}+\mathbf{b}))\right|\ll\rho^{\gamma+1}=o\left(\rho^{\alpha}\right).$$

On the other hand,

$$\frac{\mathrm{d}}{\mathrm{d}t} |\boldsymbol{\xi} + t(\boldsymbol{\xi} + \mathbf{b})|^{\alpha} = \alpha |\boldsymbol{\xi} + t(\boldsymbol{\xi} + \mathbf{b})|^{\alpha - 2} \times \\ \times \left(|\boldsymbol{\xi}| |\boldsymbol{\xi} + \mathbf{b}| \cos\left(\varphi(\boldsymbol{\xi}, \boldsymbol{\xi} + \mathbf{b})\right) + t |\boldsymbol{\xi} + \mathbf{b}|^2 \right) \\ \gg \rho^{\alpha},$$

where the last line holds from the fact that $\cos \varphi(\xi, \xi + \mathbf{b}) > \sqrt{2}/2$.

Lemma 11.10. Let

$$\mathcal{N} := \left\{ \boldsymbol{\xi} \in \mathbb{R}^d : n(\boldsymbol{\xi}; \mathcal{G}) \leq mn(\boldsymbol{\xi}; \mathcal{Z}') \right\}$$

and $\mathcal{N}_{\mathcal{G}} = \mathcal{N} \cap \mathcal{G}$. Then, we have that

$$\operatorname{vol}(\mathcal{N}_{\mathcal{G}}) \ll \operatorname{vol}(\mathcal{Z}') \ll \delta \rho^{d-\alpha-\varepsilon_0},$$

where ε_0 is as in Lemma 11.5.

Proof. Observe first that for any $E \subset \mathbb{R}^d$, the function $n(\xi; E)$ is constant on the fibres $\xi \mod \Theta$, as such it is well defined on \mathcal{O} and \mathcal{N} is invariant under the action of Θ . We therefore have that

$$\operatorname{vol}(\mathscr{N}_{\mathscr{G}}) = \int_{\mathscr{N}/\Theta} n(\boldsymbol{\xi}; \mathscr{G}) \, \mathrm{d}\boldsymbol{\xi}$$
$$\leq m \int_{\mathscr{N}/\Theta} n(\boldsymbol{\xi}; \mathscr{Z}') \, \mathrm{d}\boldsymbol{\xi}$$
$$= m \operatorname{vol}(\mathscr{N} \cap \mathscr{Z}')$$
$$\leq m \operatorname{vol}(\mathscr{Z}').$$

The claim now follows from Lemma 11.5.

Lemma 11.11. Let

$$\mathcal{U} := \{ \boldsymbol{\xi} \in \mathcal{G} \setminus \mathcal{N} : \boldsymbol{\xi} + \boldsymbol{\theta}_1 \in \mathcal{Y}'_{ij,\pi/4}(\boldsymbol{\theta}_1 - \boldsymbol{\theta}_2) \\ for some \ 1 \le i, j \le m \ and \ \boldsymbol{\theta}_1, \boldsymbol{\theta}_2 \in \Theta \}.$$

Then, if ε *and* δ *are such that* $\delta \rho^{d-\alpha+2\varepsilon} \rightarrow 0$ *, we have*

$$\operatorname{vol}(\mathscr{U}) \ll \delta^2 \rho^{4-2\alpha+2d+6\varepsilon} + \delta \rho^{1-\alpha+d-\varepsilon(d-1)}.$$

Proof. Suppose that $\boldsymbol{\xi} \in \mathcal{U} \subset \mathcal{G}$, so that $\boldsymbol{\xi} \in \mathcal{G}_k$ for some $1 \le k \le m$. Consider the lattice elements $\boldsymbol{\theta}_1, \boldsymbol{\theta}_2 \in \Theta^{\dagger}$ such that $\boldsymbol{\xi} + \boldsymbol{\theta}_1 \in \mathcal{G}'_{ij,\pi/4}(\boldsymbol{\theta}_1 - \boldsymbol{\theta}_2)$. By definition of $\mathcal{G}'_{ij,\pi/4}$ and translation, this means that

$$\boldsymbol{\xi} \in (\mathscr{G}'_i - \boldsymbol{\theta}_1) \cap (\mathscr{G}'_i - \boldsymbol{\theta}_2),$$

and therefore that

$$\boldsymbol{\xi} \in \mathscr{Y}_{ki}'(-\boldsymbol{\theta}_1) \cap \mathscr{Y}_{kj}'(-\boldsymbol{\theta}_2).$$

74

Furthermore, $\varphi(\boldsymbol{\xi} + \boldsymbol{\theta}_1, \boldsymbol{\xi} + \boldsymbol{\theta}_2) > \pi/4$. As such,

$$\max\{\varphi(\boldsymbol{\xi},\boldsymbol{\xi}+\boldsymbol{\theta}_1),\varphi(\boldsymbol{\xi},\boldsymbol{\xi}+\boldsymbol{\theta}_2)\} > \pi/8.$$

Combining the previous two displays yields

$$\boldsymbol{\xi} \in \mathscr{Y}'_{ki,\pi/8}(-\boldsymbol{\theta}_1) \cup \mathscr{Y}_{kj,\pi/8}(-\boldsymbol{\theta}_2).$$

Therefore,

$$\operatorname{vol}(\mathscr{U}) \leq \operatorname{vol}\left(\bigcup_{i,j=1}^{m} \bigcup_{\boldsymbol{\theta} \in \Theta} \mathscr{Y}'_{ij,\pi/8}(\boldsymbol{\theta})\right)$$
$$\ll \delta^{2} \rho^{4-2\alpha+2d+6\varepsilon} + \delta \rho^{1-\alpha+d-\varepsilon(d-1)},$$

the last line holding by virtue of Lemma 11.8.

Proposition 11.12. Let

$$s := \min\left\{\frac{\alpha d - d^2 - 3d - \alpha - 2}{2(d+2)}, \alpha - d + \frac{\alpha - d - 2}{2(d+2)}\right\}.$$
(11.20)

For ρ large enough and $\delta = o(\rho^s)$, the set

$$\mathcal{K} := \mathcal{G} \setminus (\mathcal{N}_{\mathcal{G}} \cup \mathcal{U})$$

is non empty.

Proof. Recall from Lemma 11.5 that $vol(\mathcal{G}) \approx \delta \rho^{d-\alpha}$. On the other hand we have from Lemma 11.10 that there is $\varepsilon_0 > 0$ such that

$$\operatorname{vol}(\mathcal{N}_{\mathscr{G}}) \ll \delta \rho^{d-\alpha-\varepsilon_0}$$

and from Lemma 11.11 that as soon as $\delta \rho^{d-\alpha+2\varepsilon} \to 0$ we have that

$$\operatorname{vol}(\mathscr{U}) \ll \delta^2 \rho^{4-2\alpha+2d+6\varepsilon} + \delta \rho^{1-\alpha+d-\varepsilon(d-1)}.$$

Take

$$\varepsilon = \frac{\alpha - d - 2}{2(d+2)}.$$

Observe that indeed when $\delta = o(\rho^s)$ we have $\delta \rho^{2-\alpha+2\varepsilon} \to 0$ as $\rho \to \infty$. We also observe that with that choice of parameters $vol(\mathcal{U}) + vol(\mathcal{N}_{\mathcal{G}}) = o(vol(\mathcal{G}))$ and hence, for large enough ρ , \mathcal{K} is not empty.

We now have all the necessary ingredients to prove Proposition 11.2.

Proof of Proposition 11.2. Let *s* be defined as in (11.20). For any $\varepsilon > 0$, set $s' = \min\{s - \varepsilon, \alpha - \nu\}$ where *v* is obtained in Lemma 11.3, and put $\delta = \rho^{s'}$. Then, by Proposition 11.12 \mathcal{K} is nonempty

for ρ large enough; choose $\xi_0 \in \mathcal{K}$. For $1 \le j \le m$, let $\Gamma_j, \Gamma'_j \subset \Theta$ be defined as

$$\Gamma_j := \left\{ \boldsymbol{\theta} \in \Theta : \boldsymbol{\xi}_0 + \boldsymbol{\theta} \in \mathcal{G}_j \right\}, \qquad \Gamma'_j := \left\{ \boldsymbol{\theta} \in \Theta : \boldsymbol{\xi}_0 + \boldsymbol{\theta} \in \mathcal{G}'_j \right\}.$$

It follows from the definition of ${\mathcal K}$ that

$$Q := \sum_{j=1}^{m} \# \Gamma_j \ge n(\boldsymbol{\xi}_0; \mathcal{G}).$$

Since $\xi_0 \notin \mathcal{U}$, we have that $\varphi(\xi_0, \xi_0 + \theta) \le \pi/4$ for all $\theta \in \Gamma_j$. From equations (11.9)–(11.12), Lemma 11.9, since $\delta \ll \rho^{\alpha-\nu}$ there are $t \ll \rho^{-\nu}$ and Z_0 independent of ρ such that for all $1 \le j \le m$ and $\theta \in \Gamma_j$,

$$\rho^{\alpha} - Z_0 \delta \le g_j \left((1-t) \,\boldsymbol{\xi}_0 + \boldsymbol{\theta} \right) \le \rho^{\alpha} - \delta, \tag{11.21}$$

and

$$\rho^{\alpha} + \delta \le g_j \left((1+t) \,\boldsymbol{\xi}_0 + \boldsymbol{\theta} \right) \le \rho^{\alpha} + Z_0 \delta. \tag{11.22}$$

It is clear that if (11.21) and (11.22) hold, they also hold replacing Z_0 with any $Z > Z_0$. The precise value we assign to Z might change as the proof goes along, but it will remain independent of ρ and δ . We denote by \mathcal{J} the radial interval

$$\mathscr{J} := [(1-t)\xi_0, (1+t)\xi_0].$$

We now restrict ourselves to consider in (11.5) only those $\mathbf{k} \in \mathscr{J} \mod \Theta$. This corresponds, at the level of Besicovitch space, to the study of the operators

$$\mathbf{A}^{\mathscr{G}} := \mathbf{A} P_{\mathscr{G} + \Theta},$$

where the projection $P_{\mathscr{J}+\Theta}$ is defined in Proposition 2.16. We also define the projection P_{good} acting on the basis elements $e_{\xi} \otimes v_j$ as

$$P_{\text{good}} e_{\boldsymbol{\xi}} \otimes v_j := (P_{\mathcal{J}+\Gamma_i} e_{\boldsymbol{\xi}}) \otimes v_j.$$

From Lemma 11.9 and its proof, for all $\boldsymbol{\xi} \in \mathcal{J}$, and all $\boldsymbol{\theta} \in \Gamma_j$, we have that $\boldsymbol{\xi} + \boldsymbol{\theta} \in \mathcal{B}'_j$. It therefore follows from (11.11) and that P_{good} commutes with **A** and **A** \mathcal{I} . We decompose

$$\mathbf{A}^{\mathscr{F}} = \mathbf{A}_{\text{good}} + \mathbf{A}_{\text{bad}},$$

where

$$\mathbf{A}_{\text{good}} := \mathbf{A}^{\mathscr{J}} P_{\text{good}}$$
 and $\mathbf{A}_{\text{bad}} = \mathbf{A}^{\mathscr{J}} (\text{Id} - P_{\text{good}})$

For $\xi \in \mathcal{J}$, denote by $\boldsymbol{\mu} := \{\mu_p(\boldsymbol{\xi}) : 1 \le p \le Q\}$ the eigenvalues of the operators $\mathbf{A}_{good}(\{\boldsymbol{\xi}\})$, and $\hat{\boldsymbol{\nu}} := \{\hat{\boldsymbol{\nu}}_p(\boldsymbol{\xi}) : p \in \mathbb{N}\}$ the eigenvalues of $\mathbf{A}_{bad}(\{\boldsymbol{\xi}\})$, each ordered nondecreasingly pointwise. It follows that for every $\boldsymbol{\xi} \in \mathcal{J}$, we have the equality of multisets

$$\{\lambda_p(\mathbf{A}(\{\boldsymbol{\xi}\})): p \in \mathbb{N}\} = \boldsymbol{\mu} \cup \hat{\boldsymbol{\nu}}\}$$

Furthermore, for all $v \in \hat{v}$ corresponds (j, θ) such that either $\xi_0 + \theta \in \mathcal{Z}'_j$ or $\xi_0 + \theta \notin \mathcal{A}'_j$. By Lemma 11.9, for $1 \le j \le m$ and $\theta \in \Gamma_j$, the functions $g_j(\xi + \theta)$ are continuous and increasing for $\xi \in \mathcal{J}$. Since the functions μ_p are obtained simply by ordering the functions $g_j(\xi + \theta)$ for $1 \le j \le m$ and $\theta \in \Gamma_j$ in nondecreasing order at every point $\{\xi\}$, they are therefore themselves continuous and increasing on \mathcal{J} .

Let us now consider the functions $\hat{v}_{\mathbf{p}}$. Notice that they are formed by removing $\#\Gamma$ continuous branches from a family of continuous functions. They are therefore all continuous on \mathscr{J} , and [PS10, Theorem 3.6] also applies to them. In particular, one can choose $Z \ge Z_0$ large enough so that all functions $v_{\mathbf{p}}$ such that $v_{\mathbf{p}}(\boldsymbol{\xi}_0) \le \rho^{\alpha} - Z\delta/2$, respect the bound $v_{\mathbf{p}}(\boldsymbol{\xi}) < \rho^{\alpha} - \delta$ for all $\boldsymbol{\xi} \in \mathscr{J}$. Similarly, all functions $v_{\mathbf{p}}$ such that $v_{\mathbf{p}}(\boldsymbol{\xi}_0) > \rho^{\alpha} + Z\delta/2$ respect $v_{\mathbf{p}}(\boldsymbol{\xi}) > \rho^{\alpha} + \delta$ for all $\boldsymbol{\xi} \in \mathscr{J}$. We therefore define the sets

$$\boldsymbol{\tau} := \left\{ \boldsymbol{\nu} \in \hat{\boldsymbol{\nu}} : \left| \boldsymbol{\nu}(\boldsymbol{\xi}_0) - \boldsymbol{\rho}^{\alpha} \right| \ge \frac{Z\delta}{2} \right\}$$

and $\mathbf{v} = \hat{\mathbf{v}} \setminus \mathbf{\tau}$. In particular, for all $v \in \mathbf{v}$ there corresponds $(j, \boldsymbol{\theta})$ such that in $\boldsymbol{\xi}_0 + \boldsymbol{\theta} \in \mathscr{A}'_j$ and therefore in \mathscr{Z}'_i .

By construction, we have indeed that at every $\mathbf{k} \in [\mathbf{k}_1, \mathbf{k}_2]$ the images of the families $\boldsymbol{\mu}, \boldsymbol{\nu}, \boldsymbol{\tau}$ are the eigenvalues of $\mathbf{A}(\mathbf{k})$ counted with multiplicity, satisfying therefore (11.1). Hypothesis (11.2) is also satisfied since all $\boldsymbol{\mu} \in \boldsymbol{\mu}$ are increasing, and for all $\boldsymbol{\mu} \in \boldsymbol{\mu}$, either

$$\mu((1-t)\xi_0) < \rho^{\alpha} - \delta, \quad \text{or} \quad \mu((1+t)\xi_0) > \rho^{\alpha} + \delta.$$
 (11.23)

It follows from our choice of t in (11.21)–(11.22) that there are at least $Q \ge n(\boldsymbol{\xi}_0; \mathcal{G})$ of them so that both inequalities in (11.23). On the other hand, all $v \in \boldsymbol{v}$ correspond to some $(j, \boldsymbol{\theta}) \in \widetilde{\Theta}$ such that $\boldsymbol{\xi}_0 + \boldsymbol{\theta} \in \mathcal{Z}'$, hence there's at most $mn(\boldsymbol{\xi}_0; \mathcal{Z}') < n(\boldsymbol{\xi}_0; \mathcal{G})$ of them, hence hypothesis (11.3) holds. Finally, we constructed $\boldsymbol{\tau}$ explicitly so that for all $\tau \in \boldsymbol{\tau}$, hypothesis (11.4) holds.

Proposition 11.2 is therefore true, and we conclude that we have at least $\zeta(\rho^{\alpha}; \mathbf{A}) > \delta$. In view of our choice of δ this also gives us an overlap exponent at least

$$S \ge \min\{\alpha - \nu, s - \varepsilon\}$$

where $\varepsilon > 0$ is arbitrary and *s* is given in (11.20).

12. THE DIRAC OPERATOR

In this section, we aim to get conditions on perturbations of the Dirac operator so that the gauge transform and, more importantly, all the theorems from Part II can be applied. Basic facts and theorem on the Dirac operator are found in [GM91, Tha92]. We consider Dirac operators built through Clifford algebras, of which the usual two- and three-dimensional cases are

examples. We are then able to explicitly describe perturbations to which we can apply the gauge transform method and recover the results of Sections 7–11.

12.1. **Clifford algebras.** We give here basic facts about Clifford algebras used to construct the Dirac operator in the flat setting. They can be found in [GM91, Section 7]. Let $\mathbb{R}^{p,q}$ be the euclidean space of dimension p + q equipped with the canonical quadratic form η of signature (p, q). In our applications, we consider only the cases $\mathbb{R}^{0,d}$ (Euclidean) and $\mathbb{R}^{1,d}$ (Minkowski). We denote their orthonormal bases respectively $\{\mathbf{v}_1, \dots, \mathbf{v}_d\}$ and $\{\mathbf{v}_0, \mathbf{v}_1, \dots, \mathbf{v}_d\}$. Consider the algebra $\mathfrak{A}_{p,q}$ generated by $\{1, \mathbf{v}_1, \dots, \mathbf{v}_d\}$ or $\{1, \mathbf{v}_0, \dots, \mathbf{v}_d\}$ with the relations

$$\mathbf{v}_j \mathbf{v}_k + \mathbf{v}_k \mathbf{v}_j = -2\eta_{jk}.\tag{12.1}$$

It is easy to see that $\mathfrak{A}_{p,q}$ has dimension 2^{p+q} . For any subset $S := \{s_1, \ldots, s_k\} \subset \{0, \ldots, d\}$ (or of $\{1, \ldots, d\}$ in the euclidean setting), we denote by \mathbf{v}_S the element $\mathbf{v}_{s_1} \cdots \mathbf{v}_{s_k} \in \mathfrak{A}_{p,q}$, where by convention $\mathbf{v}_{\varnothing} = 1$. The Clifford algebra on $\mathbb{R}^{p,q}$ is isomorphic to the exterior algebra $\Lambda^*(\mathbb{R}^d)$.

From the anticommutation relation (12.1), we deduce that each pair of the 2^{p+q} generators of $\mathfrak{A}_{p,q}$ either commutes or anticommutes, according to the rule

When p + q is even, there is a faithful representation of $\mathfrak{A}_{p,q}$ acting on the spinor space $\mathbb{C}^{2^{(p+q)/2}}$. A specific representation by matrices constructed recursively is given in [Upm02] in the Euclidean and Minkowski cases. This representation γ has the property that for all $1 \le j \le d$, the matrix $\gamma_j := \gamma(\mathbf{v}_j)$ is skew-hermitian and squares to $-\operatorname{Id}_{p+q}$, $\gamma_0 := \gamma(\mathbf{v}_0)$ is hermitian and squares to the identity, and there is some |c| = 1 so that the grading operator $\Gamma := c \prod_j \gamma_j$ is a diagonal matrix of the form

$$\Gamma = \begin{pmatrix} \mathrm{Id}_{(p+q)/2} & 0\\ 0 & -\mathrm{Id}_{(p+q)/2} \end{pmatrix}.$$

We can observe that for all j, $\Gamma \gamma_j = -\gamma_j \Gamma$. The operator Γ is called "grading" because it induces a \mathbb{Z}_2 grading on $\mathfrak{A}_{p,q}$. The even subalgebra of $\mathfrak{A}_{p,q}$ consists of all the elements commuting with Γ , while the odd subspace consists of all the anti-commuting elements. In particular, all the γ_j are in the odd subspace, which is characterised as a product of an odd number of generators, while the even subalgebra is characterised as products of even number of generators.

Lemma 12.1. Let γ be an element of the odd subspace. Then, as a matrix it has the form

$$\gamma := \begin{pmatrix} \mathbf{0} & \mathbf{X} \\ \mathbf{Y} & \mathbf{0} \end{pmatrix},$$

where each of the blocks is a $m/2 \times m/2$ matrix.

Proof. This follows from a simple computation of the relation $\gamma\Gamma + \Gamma\gamma = 0$ on the matrix elements.

The representation γ also allows us to see that as a C^* -algebra, $\mathfrak{A}_{p,q}$ is naturally isomorphic to an algebra of operators on a Hilbert space \mathfrak{S}_{p+q} , called the *spinor space*. When *m* is even, we have that $\mathfrak{S}_{p+q} \cong \mathbb{C}^{2^{(p+q)/2}}$. Therefore, setting $m = 2^{(p+q)/2}$, we can use this representation to obtain operators in \mathbf{S}_m^{∞} .

12.2. **Dirac operators.** We define (spatial) Dirac operators differently depending on whether the number of spatial dimensions is even or odd.

Definition 12.2. Let *d* be odd. The *d*-dimensional free Dirac operator \mathbf{A}_d is the first order system acting on spinors in $L^2(\mathbb{R}^d; \mathbb{C}^m)$, for $m = 2^{\frac{d+1}{2}}$ given by

$$\mathbf{A}_d = \sum_{j=1}^d \gamma_j \partial_j,$$

where the γ_i are given by the representation of $\mathfrak{A}_{1,d-1}$ in $\mathscr{L}(\mathbb{C}^m)$.

Definition 12.3. Let *d* be even. The *d*-dimensional free Dirac operator \mathbf{A}_d is the first order system acting on spinors $L^2(\mathbb{R}^d; \mathbb{C}^m)$, for $m = 2^{\frac{d}{2}}$ given by

$$\mathbf{A}_d = \sum_{j=1}^d \gamma_j \boldsymbol{\partial}_j,$$

where the γ_i are given by the representation of $\mathfrak{A}_{0,d}$ in $\mathscr{L}(\mathbb{C}^m)$.

It is easy to see in both cases that $\mathbf{A}_d^2 = -\Delta \operatorname{Id}_m$.

Example 12.4. The two-dimensional Dirac operator with mass *M* is given in [Tha92, Equation 1.14] as

$$\mathbf{A}_{2,M} = -i\left(\sigma_1\partial_{x_1} + \sigma_2\partial_{x_2}\right) + \sigma_3M,$$

where $\sigma_1, \sigma_2, \sigma_3$ are the Pauli matrices

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \ \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \ \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

It is a perturbation of order 0 of the free Dirac operator. Indeed, the Pauli matrices can be used for a representation of the Clifford algebra $\mathfrak{A}_{0,2}$, and σ_3 corresponds to the grading operator Γ .

Example 12.5. The three-dimensional Dirac operator with mass *M* from [Tha92, Equation 1.11] given by

$$\mathbf{A}_{3,M} = -i\left(\gamma_1\partial_{x_1} + \gamma_2\partial_{x_2} + \gamma_3\partial_{x_3}\right) + \Gamma M$$

is also a perturbation of order 0 of the free Dirac operator. Here, the matrices γ_j are the Dirac γ -matrices used as a representation of $\mathfrak{A}_{1,3}$, and our notation generalises this notion, following [Upm02].

We now show that the operators A_d are elliptic in the sense of Section 3.

Proposition 12.6. Let m := m(d) be the dimension of the spinor space on which \mathbf{A}_d acts. The operator $\mathbf{U} \in \mathbf{S}_m^0$ with symbol

$$\mathbf{u}(\mathbf{x},\boldsymbol{\xi}) := \frac{\mathbf{1}_{\{|\boldsymbol{\xi}|\geq 1\}}(\boldsymbol{\xi})}{\sqrt{2}} \left(\mathrm{Id}_m + \frac{i}{|\boldsymbol{\xi}|} \Gamma \sum_{j=1}^d \boldsymbol{\xi}_j \gamma_j \right) + \mathbf{1}_{\{|\boldsymbol{\xi}|<1\}}(\boldsymbol{\xi}) \, \mathrm{Id}_m \tag{12.2}$$

is unitary. Furthermore, $\mathbf{U}\mathbf{A}_d\mathbf{U}^* \in \mathbf{D}\mathbf{E}\mathbf{S}_m^1$ and there is $\mathbf{R} \in \mathbf{S}_m^{-\infty}$ such that the symbol of $\mathbf{U}\mathbf{A}_d\mathbf{U}^* - \mathbf{R}$ is $|\boldsymbol{\xi}|\Gamma$.

Proof. The symbol of the adjoint of **U** is given, following (2.14), by

$$\mathbf{u}^{\dagger}(\mathbf{x},\boldsymbol{\xi}) = \frac{\mathbf{1}_{\{|\boldsymbol{\xi}|\geq 1\}}(\boldsymbol{\xi})}{\sqrt{2}} \left(\mathrm{Id}_{m} - \frac{i}{|\boldsymbol{\xi}|} \Gamma \sum_{j=1}^{d} \boldsymbol{\xi}_{j} \gamma_{j} \right) + \mathbf{1}_{\{|\boldsymbol{\xi}|<1\}}(\boldsymbol{\xi}) \, \mathrm{Id}_{m}$$

and we can compute that

$$\begin{bmatrix} \mathbf{u} \circ \mathbf{u}^{\dagger} \end{bmatrix} (\boldsymbol{\xi}) = \frac{\mathbf{1}_{\{|\boldsymbol{\xi}| \ge 1\}}(\boldsymbol{\xi})}{2} \left(\mathrm{Id}_{m} - \frac{1}{|\boldsymbol{\xi}|^{2}} \sum_{j,k=1}^{d} \Gamma^{2} \gamma_{j} \gamma_{k} \boldsymbol{\xi}_{j} \boldsymbol{\xi}_{k} \right) + \mathbf{1}_{\{|\boldsymbol{\xi}| < 1\}}(\boldsymbol{\xi}) \mathrm{Id}_{m}$$
$$= \frac{\mathbf{1}_{\{|\boldsymbol{\xi}| \ge 1\}}(\boldsymbol{\xi})}{2} \left(\mathrm{Id}_{m} - \frac{1}{|\boldsymbol{\xi}|^{2}} \sum_{j} \gamma_{j}^{2} \boldsymbol{\xi}_{j}^{2} \right) + \mathbf{1}_{\{|\boldsymbol{\xi}| < 1\}}(\boldsymbol{\xi}) \mathrm{Id}_{m}$$
$$= \mathrm{Id}_{m}.$$

In a very similar fashion, we see that the symbol of $\mathbf{U}\mathbf{A}_d\mathbf{U}^*$ is given by

$$[\mathbf{u} \circ \mathbf{a}_d \circ \mathbf{u}^{\mathsf{T}}](\boldsymbol{\xi}) = \mathbf{1}_{\{|\boldsymbol{\xi}| \ge 1\}}(\boldsymbol{\xi}) |\boldsymbol{\xi}| \Gamma + \mathbf{1}_{\{|\boldsymbol{\xi}| < 1\}}(\boldsymbol{\xi}) \mathbf{a}_d(\boldsymbol{\xi}).$$

This proves our claim where $\mathbf{R} \in \mathbf{S}_m^{-\infty}$ has symbol

$$\mathbf{r}(\boldsymbol{\xi}) = \mathbf{1}_{\{|\boldsymbol{\xi}| < 1\}}(\boldsymbol{\xi})(\mathbf{a}_d(\boldsymbol{\xi}) - |\boldsymbol{\xi}| \Gamma).$$

We now see that for d = m = 2, the operators $\mathbf{A}_2 + \mathbf{B}$, $\mathbf{B} \in \mathbf{S}_m^\beta$, $\beta < 1$ are unitarily equivalent to an operator satisfying the hypotheses of Theorems 8.2 and Theorem 10.1, which proves that we generically have a complete asymptotic expansion for the density of states, and that if **B** is periodic then **A** has the Bethe–Sommerfeld property. In other words, the following two theorems are proved, which are more precise reformulations of Theorems 1.1 and 1.3. **Theorem 12.7.** Let $\beta < 1$ and $\mathbf{A} = \mathbf{A}_2 + \mathbf{B}$, where $\mathbf{B} \in \mathbf{S}_2^{\beta}$ satisfies the generic conditions \mathbf{A} , \mathbf{B} and \mathbf{C} . Then, for every K > -2 there is a finite set $L \subset (0, 2 + K)$ so that for every $j \in L \cup \{0\}$ there are constants C_j^{\pm} , $C_{j,\log}^{\pm}$ such that

$$N^{\pm}(\mathbf{A};\lambda) = C_0^{\pm}\lambda^2 + \sum_{j \in L} \left(C_j^{\pm}\lambda^{2-j} + C_{j,\log}^{\pm}\lambda^{2-j}\log\lambda \right) + O\left(\lambda^{-K}\right)$$

as $\lambda \to \infty$.

Theorem 12.8. Let $\beta < 1$ and $\mathbf{A} = \mathbf{A}_2 + \mathbf{B}$, where $\mathbf{B} \in \mathbf{S}_2^{\beta}$ is periodic. Then, \mathbf{A} has the Bethe-Sommerfeld property, i.e. there exists $\lambda_0 > 0$ such that the spectrum of \mathbf{A} contains intervals $(-\infty, -\lambda_0]$ and $[\lambda_0, \infty)$.

We now want to address the question of the perturbations that are allowed whenever $d \ge 3$.

Proposition 12.9. For $\beta < 1$, and $0 \le j \le d$ (with 0 omitted when d is even), let $B_{\text{Id}}, B_{\Gamma}, B_j \in \mathbf{S}_1^{\beta}$ be scalar pseudo-differential operators of order β , and put

$$\mathbf{B} = B_{\mathrm{Id}} \, \mathrm{Id}_m + B_{\Gamma} \Gamma + \sum_j B_j \gamma_j.$$

Then, there are operators $\mathbf{B}' \in \mathbf{US}_m^\beta$, $\mathbf{R} \in \mathbf{S}_m^{\beta-1}$ and $\widetilde{\mathbf{B}} \in \mathbf{S}_m^\beta$ whose symbol has image in the odd subspace of $\mathfrak{A}_{p,q}$ such that

$$\mathbf{U}(\mathbf{A}_d + \mathbf{B})\mathbf{U}^* = \operatorname{Op}(|\boldsymbol{\xi}|)\Gamma + \mathbf{B}' + \mathbf{B} + \mathbf{R}$$
(12.3)

Proof. The unitary operator U from (12.2) can be written as

$$\mathbf{U} = \frac{1}{\sqrt{2}} \left(\mathrm{Id}_m + \sum_{j=1}^d U_j \Gamma \gamma_j \right) \mod \mathbf{S}_m^{-\infty}.$$

Here, $U_j \in \mathbf{S}_1^0$ are scalar pseudo-differential operators given by

$$U_j = \operatorname{Op}\left(\frac{i\boldsymbol{\xi}_j\boldsymbol{\chi}(\boldsymbol{\xi})}{|\boldsymbol{\xi}|}\right),$$

where χ is a smooth function supported in $\{|\boldsymbol{\xi}| \ge 1/2\}$ and $\chi(\boldsymbol{\xi}) \equiv 1$ for all $|\boldsymbol{\xi}| \ge 3/4$. We now compute $\mathbf{U}B_{\gamma}\gamma\mathbf{U}^*$ for different values of γ . All the sums range from 1 to d with additional restrictions, we have only written the restrictions to make notation lighter. For $1 \le j \le d$ we have

$$\mathbf{U}B_{j}\gamma_{j}\mathbf{U}^{*} = \frac{1}{2} \left(B_{j}\gamma_{j} + \sum_{k\neq j} [U_{k};B_{j}]\Gamma\gamma_{k}\gamma_{j} - (U_{j}B_{j} + B_{j}U_{j})\Gamma + \sum_{k} (U_{k}B_{j}U_{j} + U_{j}B_{j}U_{k})\gamma_{k} - \sum_{k} U_{k}B_{j}U_{k}\gamma_{j} - \sum_{\substack{\ell\neq j \\ k\neq j \\ k<\ell}} ([U_{\ell};B_{j}U_{k}] + [B_{j};U_{k}]U_{\ell})\gamma_{k}\gamma_{\ell}\gamma_{j} \right) \mod \mathbf{S}_{m}^{-\infty}.$$

$$(12.4)$$

Let us have a careful look at each of the six terms in Equation (12.4). The second and the last terms involve commutators of operators with scalar-valued symbols, they are in $\mathbf{S}_m^{\beta-1}$ and we put \mathbf{R}_j as their sum. The third term is in \mathbf{US}_m^β , and we denote it \mathbf{B}'_j . Finally, the first, fourth and fifth term are readily seen to have symbols in the odd subspace, we put $\widetilde{\mathbf{B}}_j$ as their sum.

The operator $\mathbf{U}B_0\gamma_0\mathbf{U}^*$ is computed similarly as in (12.4) with some of the terms vanishing. It is given by

$$\mathbf{U}B_{0}\gamma_{0}\mathbf{U}^{*} = \frac{1}{2} \left(B_{0}\gamma_{0} + \sum_{k} [U_{k}; B_{0}]\Gamma\gamma_{k}\gamma_{0} - \sum_{k} U_{k}B_{0}U_{k}\gamma_{0} - \sum_{k<\ell} ([U_{\ell}; B_{0}U_{k}] + [B_{0}; U_{k}]U_{\ell})\gamma_{k}\gamma_{\ell}\gamma_{0} \right) \mod \mathbf{S}_{m}^{-\infty}$$

The first and third term have image in the odd subspace, we put $\tilde{\mathbf{B}}_0$ as their sum. The second and last terms involve commutators of operators with scalar-valued symbols, as such they are in $\mathbf{S}_m^{\beta-1}$ and we put \mathbf{R}_0 as their sum. We note that there are no uncoupled terms.

The operator $\mathbf{U}B_{\Gamma}\Gamma\mathbf{U}^*$ is given by

$$\mathbf{U}B_{\Gamma}\Gamma\mathbf{U}^{*} = \frac{1}{2} \left(B_{\Gamma}\Gamma - \sum_{k} (U_{k}B_{\Gamma} + B_{\Gamma}U_{k})\gamma_{k} - \sum_{k} U_{k}B_{\Gamma}U_{k}\Gamma + \sum_{k<\ell} ([U_{\ell}; B_{\Gamma}U_{k}] + [B_{\Gamma}; U_{k}]U_{\ell})\Gamma\gamma_{\ell}\gamma_{k} \right) \mod \mathbf{S}_{m}^{-\infty}$$

This time, the first and third terms are seen to be in \mathbf{US}_m^β and we put their sum as \mathbf{B}'_{Γ} . The second term has symbol in the odd subspace and we denote it by $\widetilde{\mathbf{B}}_{\Gamma}$. The last term can be seen to be in $\mathbf{S}_m^{\beta-1}$ and we denote it by \mathbf{R}_{Γ} .

Finally, the operator $\mathbf{U}B_{\mathrm{Id}} \mathrm{Id}_m \mathbf{U}^*$ is given by

$$\mathbf{U}B_{\mathrm{Id}} \operatorname{Id}_{m} \mathbf{U}^{*} = \frac{1}{2} \Big(B_{\mathrm{Id}} \operatorname{Id}_{m} + \sum_{k} [U_{k}; B_{\mathrm{Id}}] \Gamma \gamma_{k} + \sum_{k} U_{k} B_{\mathrm{Id}} U_{k} \operatorname{Id}_{m} \\ + \sum_{k < \ell} ([U_{\ell}; B_{\mathrm{Id}} U_{k}] + [B_{\mathrm{Id}}; U_{k}] U_{\ell}) \gamma_{\ell} \gamma_{k} \Big) \mod \mathbf{S}_{m}^{-\infty}$$

This time, we see that the first and third terms are in \mathbf{US}_m^{β} , we put their sum as $\mathbf{B}'_{\mathrm{Id}}$, while the second and last terms are in $\mathbf{S}_m^{\beta-1}$ and we put their sum as \mathbf{R}_{Id} .

Finally, put $\widetilde{\mathbf{R}} \in \mathbf{S}_m^{-\infty}$ as the sum of the remainders mod $\mathbf{S}_m^{-\infty}$ obtained at every step. Combining all our computations and Proposition 12.6 gives us that (12.3) holds with

$$\mathbf{B}' = \mathbf{B}'_{\mathrm{Id}} + \mathbf{B}'_{\Gamma} + \sum_{j=1}^{d} \mathbf{B}'_{j}$$
$$\widetilde{\mathbf{B}} = \widetilde{\mathbf{B}}_{\mathrm{Id}} + \widetilde{\mathbf{B}}_{\Gamma} + \sum_{j=0}^{d} \widetilde{\mathbf{B}}_{j}$$
$$\mathbf{R} = \widetilde{\mathbf{R}} + \mathbf{R}_{\mathrm{Id}} + \mathbf{R}_{\Gamma} + \sum_{j=0}^{d} \mathbf{R}_{j}.$$

The next theorem follows and includes Theorem 1.2 as a special case when d = 3.

Theorem 12.10. Let m(d) be the dimension of the spinor space on which \mathbf{A}_d acts. For $\beta \le 1/2$ and $0 \le j \le d$ (with 0 omitted when d is even) let B_{Γ} , B_j , $B_{\mathrm{Id}} \in \mathbf{S}^{\beta}$ be scalar pseudo-differential operators satisfying Conditions I–IV, and put

$$\mathbf{B} = B_{\mathrm{Id}} \, \mathrm{Id}_m + B_{\Gamma} \Gamma + \sum_{j=0}^d B_j \gamma_j,$$

and $\mathbf{A} = \mathbf{A}_d + \mathbf{B}$. Then, putting $\gamma^* = \max\{\beta - 1, 2\beta - 1\}$, there exists a finite set $L \subset (0, 1 - \gamma^*)$ and constants C_0^{\pm} and $C_{i,a}^{\pm}$, $0 \le q \le d - 1$, $j \in L$ such that

$$N^{\pm}(\mathbf{A};\lambda) = C_0^{\pm}\lambda^d + \sum_{j \in L} \sum_{q=0}^{d-1} C_{j,q}^{\pm}\lambda^{d-j} \log^q \lambda + O\left(\lambda^{d-1+\gamma^*}\right)$$

as $\lambda \to \infty$.

Proof. It follows from Proposition 12.9 that **UAU**^{*} satisfies the hypotheses of Theorem 8.1 with $\gamma^* = \max \{\beta - 1, 2\beta - 1\}$. In particular, the restricted asymptotics of the IDS given in that theorem are true for such operators with $\alpha = 1$.

Finally, in some highly non-generic cases we can get complete asymptotic expansions and the Bethe–Sommerfeld property for *d*-dimensional Dirac operators with $d \ge 3$. We state both results and observe that they follow directly from the fact that after conjugation by **U**, these operators are uncoupled.

Theorem 12.11. Let m(d) be the dimension of the spinor space on which \mathbf{A}_d acts, $\beta < 1$ and $\mathbf{B} \in \mathbf{US}_m^\beta$ satisfying Conditions I–IV. Put $\mathbf{A} = \mathbf{A}_d + \mathbf{U}^*\mathbf{BU}$. Then, $N^{\pm}(\mathbf{A};\lambda)$ satisfies the complete asymptotic expansion (8.2) with $\alpha = 1$.

Theorem 12.12. Let m(d) be the dimension of the spinor space on which \mathbf{A}_d acts, $\beta < 1$ and $\mathbf{B} \in \mathbf{US}_m^{\beta}$ be periodic. Put $\mathbf{A} = \mathbf{A}_d + \mathbf{U}^* \mathbf{BU}$. Then, \mathbf{A} has the Bethe–Sommerfeld property.

REFERENCES

- [BP09] G. Barbatis and L. Parnovski. Bethe-Sommerfeld conjecture for pseudodifferential perturbation. *Comm. Partial Differential Equations*, 34(4-6):383–418, 2009.
- [BS87] M. Birman and M. Solomjak. *Spectral Theory of Self-Adjoint Operators in Hilbert Space*. Mathematics and its Applications. Springer Netherlands, 1987.
- [CMS73] L. A. Coburn, R. D. Moyer, and I. M. Singer. C*-algebras of almost periodic pseudodifferential operators. Acta Math., 130:279–307, 1973.
- [Dix81] J. Dixmier. Von Neumann Algebras, volume 27 of North-Holland Mathematical Library.
 North-Holland Publishing Co., Amsterdam-New York, 1981. With a preface by E. C.
 Lance, Translated from the second French edition by F. Jellett.
- [GM91] J. E. Gilbert and M. A. M. Murray. Clifford algebras and Dirac operators in harmonic analysis, volume 26 of Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 1991.
- [Hör07] L. Hörmander. The analysis of linear partial differential operators. III. Classics in Mathematics. Springer, Berlin, 2007. Pseudo-differential operators, Reprint of the 1994 edition.
- [Ivr19] V. Ivrii. Complete Semiclassical Spectral Asymptotics for Periodic and Almost Periodic Perturbations of Constant Operators, pages 583–606. Springer, 2019.
- [Kuc93] P. Kuchment. *Floquet theory for partial differential equations*, volume 60 of *Operator Theory: Advances and Applications*. Birkhäuser Verlag, Basel, 1993.
- [MPS14] S. Morozov, L. Parnovski, and R. Shterenberg. Complete asymptotic expansion of the integrated density of states of multidimensional almost-periodic pseudo-differential operators. Ann. Henri Poincaré, 15(2):263–312, 2014.
- [Naĭ72] M. A. Naïmark. Normed algebras. Wolters-Noordhoff Publishing, Groningen, third edition, 1972. Translated from the second Russian edition by Leo F. Boron, Wolters-Noordhoff Series of Monographs and Textbooks on Pure and Applied Mathematics.
- [Par08] L. Parnovski. Bethe-Sommerfeld conjecture. Ann. Henri Poincaré, 9(3):457–508, 2008.
- [PS10] L. Parnovski and A. V. Sobolev. Bethe-Sommerfeld conjecture for periodic operators with strong perturbations. *Invent. Math.*, 181(3):467–540, 2010.
- [PS12] L. Parnovski and R. Shterenberg. Complete asymptotic expansion of the integrated density of states of multidimensional almost-periodic Schrödinger operators. *Ann. of Math.* (2), 176(2):1039–1096, 2012.

- [PS16] L. Parnovski and R. Shterenberg. Complete asymptotic expansion of the spectral function of multidimensional almost-periodic Schrödinger operators. *Duke Math. J.*, 165(3):509–561, 2016.
- [PS19] L. Parnovski and R. Shterenberg. Perturbation theory for almost-periodic potentials I: one-dimensional case. *Comm. Math. Phys.*, 366(3):1229–1257, 2019.
- [Roz78] G. V. Rozenbljum. Near-similarity of operators and the spectral asymptotic behavior of pseudodifferential operators on the circle. *Trudy Moskov. Mat. Obshch.*, 36:59–84, 294, 1978.
- [Shu78] M. A. Shubin. Almost periodic functions and partial differential operators. *Uspehi Mat. Nauk*, 33(2):3–47, 247, 1978.
- [Shu79a] M. A. Shubin. Pseudodifferential almost-periodic operators and von Neumann algebras. *Trans. Mosc. Math. Soc.*, 35:103–166, 1979.
- [Shu79b] M. A. Shubin. Spectral theory and the index of elliptic operators with almost-periodic coefficients. *Uspekhi Mat. Nauk*, 34(2):95–135, 1979.
 - [Skr85] M. M. Skriganov. Geometric and arithmetic methods in the spectral theory of multidimensional periodic operators. *Trudy Mat. Inst. Steklov.*, 171:122, 1985.
- [Sob05] A. V. Sobolev. Integrated density of states for the periodic Schrödinger operator in dimension two. *Ann. Henri Poincaré*, 6(1):31–84, 2005.
- [Sob06] A. V. Sobolev. Asymptotics of the integrated density of states for periodic elliptic pseudo-differential operators in dimension one. *Rev. Mat. Iberoam.*, 22(1):55–92, 2006.
- [Tay11] M. E. Taylor. Partial differential equations II. Qualitative studies of linear equations, volume 116 of Applied Mathematical Sciences. Springer, New York, second edition, 2011.
- [Tha92] B. Thaller. *The Dirac equation*. Texts and Monographs in Physics. Springer-Verlag, Berlin, 1992.
- [Upm02] H. Upmeier. Dirac operator and real structure on Euclidean and Minkowski spacetime. In Noncommutative geometry and the standard model of elementary particle physics (Hesselberg, 1999), volume 596 of Lecture Notes in Phys., pages 136–151. Springer, Berlin, 2002.
- [Wei77] A. Weinstein. Asymptotics of eigenvalue clusters for the Laplacian plus a potential. *Duke Math. J.*, 44(4):883–892, 1977.

J. LAGACÉ, S. MOROZOV, L. PARNOVSKI, B. PFIRSCH, AND R. SHTERENBERG

DEPARTMENT OF MATHEMATICS, UNIVERSITY COLLEGE LONDON, GOWER STREET, LONDON, WC1E 6BT, UK *Email address*: j.lagace@ucl.ac.uk *Email address*: l.parnovski@ucl.ac.uk *Email address*: bernhard.pfirsch.15@alumni.ucl.ac.uk

MATHEMATISCHES INSTITUT DER UNIVERSITÄT MÜNCHEN, THERESIENSTR. 39, D-80333 MÜNCHEN, GERMANY *Email address*: morozov@math.lmu.de

UNIVERSITY OF ALABAMA AT BIRMINGHAM, 1300 UNIVERSITY BLVD, BIRMINGHAM, AL 35294. USA *Email address*: shterenb@math.uab.edu