

ON THE BOUNDEDNESS OF THE MAXIMAL AND FRACTIONAL MAXIMAL, POTENTIAL OPERATORS IN THE GLOBAL MORREY-TYPE SPACES WITH VARIABLE EXPONENTS

N.A. Bokayev, Zh.M. Onerbek

Key words: boundedness, Riesz potential, fractional maximal operator, global Morrey-type spaces with variable exponent.

AMS Mathematics Subject Classification: 31C99, 31B99.

Abstract. We consider the global Morrey-type spaces $GM_{p(\cdot),\theta(\cdot),w(\cdot)}(\Omega)$ with variable exponents $p(x)$, $\theta(x)$ and general function $w(x, r)$ defining these spaces. In the case of unbounded sets $\Omega \subset \mathbb{R}^n$, we prove boundedness of the Hardy–Littlewood maximal operator, potential type operator in these spaces.

1 Introduction

In this paper we consider the global Morrey-type spaces $GM_{p(\cdot),\theta(\cdot),w(\cdot)}(\Omega)$ with variable exponents $p(\cdot)$, $\theta(\cdot)$ and a general function $w(x, r)$ defining a Morrey-type norm. The Morrey spaces $M_{p,\lambda}$ are introduced in [1] in relation to the study of partial differential equations. Many classical operators of harmonic analysis (for example, maximal, fractional maximal, potential operators) were studied in the Morrey-type spaces with constant exponents p , θ [2, 3, 4]. The Morrey spaces also attracted attention of researchers in the area of variable exponent analysis; see [5, 6, 7, 8, 9, 10]. The Morrey spaces $\mathcal{L}_{p(\cdot),\lambda(\cdot)}$ with variable exponent $p(\cdot)$, $\lambda(\cdot)$ were introduced and studied in [5]. The general version $M_{p(\cdot),w(\cdot)}(\Omega)$, $\Omega \subset \mathbb{R}^n$ were introduced and studied in [11] in the case of bounded sets $\Omega \subset \mathbb{R}^n$, and in [12] in the case of unbounded sets $\Omega \subset \mathbb{R}^n$. The boundedness of maximal and potential type operators in the generalized Morrey-type spaces with a variable exponent were considered in [11] in the case of bounded sets $\Omega \subset \mathbb{R}^n$, in [12] in the case of unbounded sets $\Omega \subset \mathbb{R}^n$.

Let $f \in L_{loc}(\mathbb{R}^n)$. The Hardy–Littlewood maximal operator is defined as

$$Mf(x) = \sup_{r>0} \frac{1}{|B(x, r)|} \int_{\tilde{B}(x, r)} |f(y)| dy$$

where $B(x, r)$ is the ball in \mathbb{R}^n centered at the point $x \in \mathbb{R}^n$ and of the radius r , $\tilde{B}(x, r) = B(x, r) \cap \Omega$, $\Omega \subset \mathbb{R}^n$.

The fractional maximal operator of variable order $\alpha(x)$ is defined as

$$M^{\alpha(\cdot)}f(x) = \sup_{r>0} |B(x, r)|^{-1+\frac{\alpha(x)}{n}} \int_{\tilde{B}(x, r)} |f(y)| dy, 0 \leq \alpha(x) < n.$$

In the case of $\alpha(x) = \alpha = \text{const}$, this operator coincides with the classical fractional maximal operator M^α . If $\alpha(x) = 0$, then $M^{\alpha(\cdot)}$ coincides with the operator M .

The Riesz potential $I^{\alpha(x)}$ of variable order $\alpha(x)$ is defined by the following equality:

$$I^{\alpha(x)}f(x) = \int_{\mathbb{R}^n} \frac{f(y)}{|x-y|^{n-\alpha(x)}} dy, 0 < \alpha(x) < n.$$

In the case of $\alpha(x) = \alpha = \text{const}$, this operator coincides with the classical Riesz potential I^α .

2 Preliminaries. Variable Exponent Lebesgue Spaces $L_{p(\cdot)}$. Generalized variable exponent Morrey spaces $M_{p(\cdot),w(\cdot)}$

Let $p(x)$ be a measurable function on an open set $\Omega \subset \mathbb{R}^n$ with values $(1, \infty)$. Let

$$1 < p_- \leq p(x) \leq p_+ < \infty \quad (2.1)$$

where $p_- = p_-(\Omega) = \text{essinf}_{x \in \Omega} p(x)$, $p_+ = p_+(\Omega) = \text{esssup}_{x \in \Omega} p(x)$. We denote by $L_{p(\cdot)}(\Omega)$ the space of all measurable functions $f(x)$ on Ω such that

$$J_{p(\cdot)}(f) = \int_{\Omega} [f(x)]^{p(x)} dx < \infty,$$

where the norm is defined as follows

$$\|f\|_{p(\cdot)} = \inf \left\{ \eta > 0, J_{p(\cdot)} \left(\frac{f}{\eta} \right) \leq 1 \right\}.$$

This is a Banach space. The conjugate exponent p' is defined by the formula

$$p'(x) = \frac{p(x)}{p(x) - 1}$$

Holder inequality for the variable exponents $p(\cdot)$, $p'(\cdot)$ is

$$\int_{\Omega} f(x)g(x)dx \leq C(p) \|f\|_{L_{p(\cdot)}(\Omega)} \|g\|_{L_{p'(\cdot)}(\Omega)},$$

where $C(p) = \frac{1}{p_-} + \frac{1}{p'_-}$.

The variable exponent Lebesgue spaces $L_{p(\cdot)}$ were introduced in [13], and were investigated in [14, 15].

$\mathcal{P}(\Omega)$ is the set of measurable functions $p : \Omega \rightarrow [1, \infty)$, $\mathcal{P}^{\log}(\Omega)$ is the set of measurable functions $p(x)$ satisfying the local log-condition

$$|p(x) - p(y)| \leq \frac{A_p}{-\ln|x-y|}, \quad |x-y| \leq \frac{1}{2}, x, y \in \Omega$$

where A_p is independent of x и y . $\mathcal{P}^{\log}(\Omega)$ is the set of measurable functions $p(x)$ satisfying (2.1) and the log-condition. In the case of Ω is an unbounded set, we denote

by $\mathbb{P}_\infty^{\log}(\Omega)$ the set of exponents which is a subset of the set of $\mathbb{P}^{\log}(\Omega)$ and satisfying the decay condition

$$|p(x) - p(\infty)| \leq A_\infty \ln(2 + |x|), x \in \mathbb{R}^n.$$

Let $\mathbb{A}^{\log}(\Omega)$ be the set of bounded exponents $\alpha : \Omega \rightarrow \mathbb{R}$ satisfying the log-condition.

Let Ω be an open bounded set, $p \in \mathbb{P}^{\log}(\Omega)$ and $\lambda(x)$ be a measurable function on Ω with values in $[0, n]$. The variable Morrey space $\mathcal{L}_{p(\cdot), \lambda(\cdot)}(\Omega)$ is introduced in [3] with the norm

$$\|f\|_{\mathcal{L}_{p(\cdot), \lambda(\cdot)}(\Omega)} = \sup_{x \in \Omega, t > 0} t^{-\frac{\lambda(x)}{p(x)}} \|f\|_{L_{p(\cdot)}(\tilde{B}(x, t))}.$$

Let $w(x, r)$ be nonnegative measurable function on Ω , where $\Omega \subset \mathbb{R}^n$ is a open bounded set. The generalized Morrey type space $M_{p(\cdot), w(\cdot)}(\Omega)$ with variable exponent is defined in [11] with the norm

$$\|f\|_{M_{p(\cdot), w(\cdot)}(\Omega)} = \sup_{x \in \Omega, r > 0} \frac{r^{-\frac{n}{p(x)}}}{w(x, r)} \|f\|_{L_{p(\cdot)}(\tilde{B}(x, r))}.$$

Let $w(x, r)$ be nonnegative measurable function on Ω , where $\Omega \subset \mathbb{R}^n$ is a open unbounded set. The generalized Morrey type space $M_{p(\cdot), w(\cdot)}(\Omega)$ with variable exponent is defined in [12] with the norm

$$\|f\|_{M_{p(\cdot), w(\cdot)}(\Omega)} = \sup_{x \in \Omega, r > 0} \frac{\|f\|_{L_{p(\cdot)}(\tilde{B}(x, r))}}{w(x, r)}.$$

Let

$$\eta_p(x, r) = \begin{cases} \frac{n}{p(x)}, & \text{if } r \leq 1; \\ \frac{n}{p(\infty)}, & \text{if } r > 1. \end{cases}$$

Definition 1. Let $p \in P^{\log}(\Omega)$, $w(x, r)$ be a positive function on $\Omega \times [0, \infty]$, where $\Omega \in \mathbb{R}^n$. Global Morrey-type space with variable exponent $GM_{p(\cdot), \theta(\cdot), w(\cdot)}(\Omega)$ is defined as the set of functions $f \in L_{p(\cdot)}^{\text{loc}}(\Omega)$ with finite norm

$$\|f\|_{GM_{p(\cdot), \theta(\cdot), w(\cdot)}(\Omega)} = \sup_{x \in \Omega} \|w(x, r) r^{-\eta_p(x, r)} \|f\|_{L_{p(\cdot)}(\tilde{B}(x, r))}\|_{L_{\theta(\cdot)}(0, \infty)}.$$

We assume that the positive measurable function $w(x, r)$ satisfies the condition

$$\sup_{x \in \Omega} \|w(x, r)\|_{L_{\theta(\cdot)}(0, \infty)} < \infty.$$

Then the space contains bounded functions and thereby is nonempty. In the case of $w(x, r) = r^{-\frac{\lambda(x)}{p(x)} + \eta_p(x, r)}$, the corresponding space is denoted by $GM_{p(\cdot), \theta(\cdot)}^{\lambda(\cdot)}$:

$$GM_{p(\cdot), \theta(\cdot)}^{\lambda(\cdot)}(\Omega) = GM_{p(\cdot), w(\cdot), \theta} \Big|_{w(x, r) = r^{-\frac{\lambda(x)}{p(x)} + \eta_p(x, r)}},$$

$$\|f\|_{GM_{p(\cdot), \theta(\cdot)}^{\lambda(\cdot)}(\Omega)} = \sup_{x \in \Omega} \|w(x, r) r^{-\frac{\lambda(x)}{p(x)}} \|f\|_{L_{p(\cdot)}(\tilde{B}(x, r))}\|_{L_{\theta(\cdot)}(0, \infty)}.$$

In the case of $\theta = \infty$, the space $GM_{p(\cdot),\infty,w(\cdot)}(\Omega)$ coincides the generalized Morrey space with variable exponent $M_{p(\cdot),w(\cdot)}(\Omega)$ with finite quasi-norm

$$\|f\|_{M_{p(\cdot),w(\cdot)}(\Omega)} = \sup_{x \in \Omega} w(x, r) r^{-\eta_p(x,r)} \|f\|_{L_{p(\cdot)}(\tilde{B}(x,r))}.$$

If $p(\cdot) = p = \text{const}$, $\theta(x) = \theta = \text{const}$ then the space $GM_{p(\cdot),\theta(\cdot),w(\cdot)}(\Omega)$ coincides with the ordinary global Morrey space $GM_{p,\theta,w}(\Omega)$, considered in the works by V.I. Burdenkov, V. Guliev and others [2, 3, 4].

The Spanne and The Adams type theorems were proved in [5] for bounded sets Ω .

Theorem 2.1. Suppose that $p \in \mathbb{P}_\infty^{\log}(\Omega)$ and

$$\sup_{t>r} \frac{\text{essinf}_{t \leq s < \infty} w_1(x, s)}{t^{\eta_p(x,t)}} \leq C \frac{w_2(x, r)}{r^{\eta_p(x,r)}}$$

where C is independent of x and r . Then the maximal operator M from $M_{p(\cdot),w_1(\cdot)}(\Omega)$ to $M_{p(\cdot),w_2(\cdot)}(\Omega)$ is bounded.

Theorem 2.2 (Spanne type result with $\alpha = \text{const}$). Let $p \in \mathbb{P}_\infty^{\log}(\Omega)$, α and $p(\cdot), q(\cdot)$ satisfy $0 < \alpha < n$, $\frac{1}{q(x)} = \frac{1}{p(x)} - \frac{\alpha}{n}$, the functions w_1 and w_2 satisfy the condition

$$\int_r^\infty \frac{\text{essinf}_{t \leq s < \infty} w_1(x, s)}{t^{1+\eta_p(x,t)}} dt \leq C \frac{w_2(x, r)}{r^{\eta_q(x,r)}}$$

where C is independent of x and r . Then the operators M_α and I_α from $M_{p(\cdot),w_1(\cdot)}$ to $M_{q(\cdot),w_2(\cdot)}$ are bounded.

The next theorem was proved in [16].

Theorem 2.3. Suppose that $p \in \mathbb{P}_\infty^{\log}(\Omega)$, $\alpha \in \mathbb{A}^{\log}(\Omega)$ and $\alpha_- = \inf_{x \in \Omega} \alpha(x) > 0$, $(\alpha p)_+ = \sup_{x \in \Omega} \alpha(x)p(x) < n$. Then

$$\left\| \frac{1}{(1 + |x|)^{\gamma(x)}} I^{\alpha(\cdot)} f \right\|_{L_{q(\cdot)}(R^n)} \leq C \|f\|_{L_{p(\cdot)}(R^n)},$$

where

$$\frac{1}{q(x)} = \frac{1}{p(x)} - \frac{\alpha(x)}{n}, \gamma(x) = A_\infty \alpha(x) \left[1 - \frac{\alpha(x)}{n} \right] \leq \frac{n}{4} A_\infty$$

with A_∞ comes from (2.1).

The following results were obtained in [12].

Theorem 2.4. Suppose that $p \in \mathcal{P}^{\log}(\Omega)$ satisfies (2.1), and let $1 < \theta_1^- \leq \theta_1(t) \leq \theta_1^+ < \infty$, $0 < t < l$, $1 < \theta_2^- \leq \theta_2(t) \leq \theta_2^+ < \infty$, $0 < t < l$. Assume that there exists $\delta > 0$ such that $\theta_1(t) \leq \theta_2(t)$, $t \in (0, \delta)$, $(\theta_1, w_1) \in \mathcal{W}(\delta, l)$. If

$$\sup_{x \in \Omega, 0 < t < \delta} \int_0^t (w_2(x, \xi))^{\theta_2(\xi)} \left(\int_t^\delta \left(\frac{1}{r w_1(x, r)} \right)^{[\tilde{\theta}_1(\xi)]'} dr \right)^{\frac{\theta_2(\xi)}{[\tilde{\theta}_1(\xi)]'}} d\xi < \infty,$$

then the maximal operator M from $M_{p(\cdot),\theta_1(\cdot),w_1(\cdot)}(\Omega)$ to $M_{p(\cdot),\theta_2(\cdot),w_2(\cdot)}(\Omega)$ is bounded.

Theorem 2.5. Suppose that $p, \alpha \in \mathcal{P}^{\log}(\Omega)$ satisfy (2.1), and let $\alpha > 0$, $(\alpha p(\cdot))_+ = \sup_{x \in \Omega} \alpha p(x) < n$, $\frac{1}{p_2(x)} = \frac{1}{p_1(x)} - \frac{\alpha}{n}$, $1 < \theta_1^- \leq \theta_1(t) \leq \theta_1^+ < \infty$, $0 < t < l$, $1 < \theta_2^- \leq \theta_2(t) \leq \theta_2^+ < \infty$, $0 < t < l$. Assume that there exists $\delta > 0$ such that $\theta_1(t) \leq \theta_2(t)$, $t \in (0, \delta)$, $(\theta_1, w_1) \in \mathcal{W}(\delta, l)$. If

$$\sup_{x \in \Omega, 0 < t < \delta} \int_0^t (w_2(x, \xi))^{\theta_2(\xi)} \left(\int_t^\delta \left(\frac{r^{\alpha(x)-1}}{w_1(x, r)} \right)^{[\tilde{\theta}_1(\xi)]'} \frac{\frac{\theta_2(\xi)}{[\tilde{\theta}_1(\xi)]'}}{dr} d\xi < \infty,$$

then the operators I_α and M_α from $M_{p_1(\cdot), \theta_1(\cdot), w_1(\cdot)}(\Omega)$ to $M_{p_2(\cdot), \theta_2(\cdot), w_2(\cdot)}(\Omega)$ are bounded.

In [12] it was proved that, if $\alpha(x)$ is a variable, under the conditions of Theorem 2.2 the operators $\frac{1}{(1+|x|)^{\gamma(x)}} M^{\alpha(\cdot)}$ and $\frac{1}{(1+|x|)^{\gamma(x)}} I^{\alpha(\cdot)}$ from $M_{p(\cdot), w_1(\cdot)}$ to $M_{q(\cdot), w_2(\cdot)}$ are bounded, where $\gamma(x) = A_\infty \alpha(x) [1 - \frac{\alpha(x)}{n}] \leq \frac{n}{4} A_\infty$.

The next lemma was proved in [12].

Lemma 2.1. Assume that $p \in \mathbb{P}_\infty^{\log}(\Omega)$ and $f \in L_{loc}^{p(\cdot)}(\mathbb{R}^n)$. Then

$$\|f\|_{L_{p(\cdot)}(B(x,t))} \leq C t^{\eta_p(x,t)} \int_t^\infty r^{-\eta_p(x,r)-1} \|f\|_{L_{p(\cdot)}(B(x,r))} dr.$$

In the same paper, the next theorem was proved.

Theorem 2.6. Suppose that $p \in \mathbb{P}_\infty^{\log}(\Omega)$. Then

$$\|Mf\|_{L_{p(\cdot)}(\tilde{B}(x,t))} \leq C t^{\eta_p(x,t)} \sup_{r>2t} r^{-\eta_p(x,r)} \|f\|_{L_{p(\cdot)}(\tilde{B}(x,r))},$$

for every $f \in L_{p(\cdot)}(\Omega)$, where C is independent of $f, x \in \Omega$ and $t > 0$.

We prove the next necessary inequality.

Theorem 2.7. Let $p \in \mathbb{P}_\infty^{\log}(\Omega)$. Then

$$\|Mf\|_{L_{p(\cdot)}(\tilde{B}(x,t))} \leq C t^{\eta_p(x,t)} \int_t^\infty s^{-\eta_p(x,s)-1} \|f\|_{L_{p(\cdot)}(\tilde{B}(x,s))} ds, \quad (2.2)$$

where C is independent of f, x, t .

Proof. Using Theorem 2.6 and Lemma 2.1, we have

$$\begin{aligned} \|Mf\|_{L_{p(\cdot)}(\tilde{B}(x,t))} &\leq C t^{\eta_p(x,t)} \sup_{r>2t} r^{-\eta_p(x,r)} \|f\|_{L_{p(\cdot)}(\tilde{B}(x,r))} \leq C t^{\eta_p(x,t)} \sup_{r>t} r^{-\eta_p(x,r)} \|f\|_{L_{p(\cdot)}(\tilde{B}(x,r))} \\ &\leq C t^{\eta_p(x,t)} \sup_{r>t} \int_r^\infty s^{-\eta_p(x,s)-1} \|f\|_{L_{p(\cdot)}(B(x,s))} ds = C t^{\eta_p(x,t)} \int_t^\infty s^{-\eta_p(x,s)-1} \|f\|_{L_{p(\cdot)}(\tilde{B}(x,s))} ds. \end{aligned}$$

□

The next inequality was proved in [12].

Theorem 2.8. Let $p \in \mathbb{P}_\infty^{\log}(\Omega)$ and α, q satisfy conditions $0 < \alpha < n$, $\frac{1}{q(x)} = \frac{1}{p(x)} - \frac{\alpha}{n}$. Then the next inequality holds

$$\|I^\alpha f\|_{L_{q(\cdot)}(\tilde{B}(x,t))} \leq C t^{\eta_q(x,t)} \int_t^\infty r^{-\eta_q(x,r)-1} \|f\|_{L_{p(\cdot)}(\tilde{B}(x,r))} dr, t > 0 \quad (2.3)$$

where C is independent of x and t .

The inequality (2.3) holds, if we put $\frac{1}{(1+|y|)^{\gamma(y)}} I^{\alpha(\cdot)} f(y)$ instead of $I^\alpha f(x)$. Namely, the following is true.

Theorem 2.9. Let $p \in \mathbb{P}_\infty^{\log}(\Omega)$ and the function $\alpha(x), q(x)$ satisfy the condition $\frac{1}{q(x)} = \frac{1}{p(x)} - \frac{\alpha(x)}{n}$. Then for each $x \in \mathbb{R}^n, t > 0$ the following inequality holds

$$\left\| \frac{1}{(1+|y|)^{\gamma(y)}} I^{\alpha(\cdot)} f \right\|_{L_{q(\cdot)}(\tilde{B}(x,t))} \leq C t^{\eta_q(x,t)} \int_t^\infty r^{-\eta_q(x,r)-1} \|f\|_{L_{p(\cdot)}(\tilde{B}(x,r))} dr \quad (2.4)$$

Proof. We represent the function f as $f(x) = f_1(x) + f_2(x)$, $f_1(x) = f(x)\chi_{\tilde{B}(x,2t)}$, $f_2(x) = f(x)\chi_{\Omega \setminus \tilde{B}(x,2t)}$. Then

$$\frac{1}{(1+|y|)^{\gamma(y)}} I^{\alpha(\cdot)} f(y) = \frac{1}{(1+|y|)^{\gamma(y)}} I^{\alpha(\cdot)} f_1(y) + \frac{1}{(1+|y|)^{\gamma(y)}} I^{\alpha(\cdot)} f_2(y).$$

By Theorem 2.3,

$$\left\| \frac{1}{(1+|y|)^{\gamma(y)}} I^{\alpha(\cdot)} f_1 \right\|_{L_{q(\cdot)}(\tilde{B}(x,t))} \leq \left\| \frac{1}{(1+|y|)^{\gamma(y)}} I^{\alpha(\cdot)} f_1 \right\|_{L_{q(\cdot)}(\mathbb{R}^n)} \leq C \|f_1\|_{L_{p(\cdot)}(\mathbb{R}^n)} = C \|f\|_{L_{p(\cdot)}(B(x,2t))}.$$

By Lemma 2.1,

$$\left\| \frac{1}{(1+|y|)^{\gamma(y)}} I^{\alpha(\cdot)} f_1 \right\|_{L_{q(\cdot)}(\tilde{B}(x,t))} \leq C t^{\eta_q(x,t)} \int_{2t}^\infty r^{-\eta_q(x,r)-1} \|f\|_{L_{p(\cdot)}(\tilde{B}(x,r))} dr. \quad (2.5)$$

If $|x - z| \leq t$ and $|z - y| \geq 2t$, we have $\frac{1}{2}|z - y| \leq |x - y| \leq \frac{3}{2}|z - y|$. Using the inequality $\frac{1}{(1+|y|)^{\gamma(y)}} \leq 1$, we infer

$$\begin{aligned} \left\| \frac{1}{(1+|y|)^{\gamma(y)}} I^{\alpha(\cdot)} f_2 \right\|_{L_{q(\cdot)}(\tilde{B}(x,t))} &\leq \left\| \int_{\mathbb{R}^n \setminus B(x,2t)} |z - y|^{\alpha-n} f(y) dy \right\|_{L_{q(\cdot)}(\tilde{B}(x,t))} \\ &\leq C \int_{\mathbb{R}^n \setminus B(x,2t)} |x - y|^{\alpha-n} |f(y)| dy \|\chi_{B(x,t)}\|_{L_{q(\cdot)}(\mathbb{R}^n)}. \end{aligned}$$

Choosing $\beta > \frac{n}{q-}$, we obtain

$$\int_{\mathbb{R}^n \setminus B(x,2t)} |x - y|^{\alpha-n} |f(y)| dy = \beta \int_{\mathbb{R}^n \setminus B(x,2t)} |x - y|^{\alpha-n+\beta} |f(y)| \left(\int_{|x-y|}^\infty s^{-\beta-1} ds \right) dy$$

=

$$\begin{aligned}
&= \beta \int_{2t}^{\infty} s^{-\beta-1} \left(\int_{y \in \mathbb{R}^n : 2t \leq |x-y| \leq s} |x-y|^{\alpha-n+\beta} |f(y)| dy \right) ds \\
&\leq C \int_{2t}^{\infty} s^{-\beta-1} \|f\|_{L_{p(\cdot)}(B(x,s))} \| |x-y|^{\alpha-n+\beta} \|_{L_{p'(\cdot)}(B(x,s))} ds \\
&\leq C \int_{2t}^{\infty} s^{\alpha-\eta_p(x,s)-1} \|f\|_{L_{p(\cdot)}(B(x,s))} ds.
\end{aligned}$$

Therefore

$$\left\| \frac{1}{(1+|y|)^{\gamma(y)}} I^{\alpha(\cdot)} f_2 \right\|_{L_{q(\cdot)}(\tilde{B}(x,t))} \leq C t^{\eta_p(x,t)} \int_{2t}^{\infty} s^{-\eta_q(x,s)-1} \|f\|_{L_{p(\cdot)}(B(x,s))} ds$$

which, together with (2.5), yields (2.4). \square

Let u and v be a positive measurable functions. The dual Hardy operator is defined by the identity

$$\tilde{H}_{v,u} f(x) = v(x) \int_x^{\infty} f(t) u(t) dt, \quad x \in \mathbb{R}^n.$$

Suppose that a is a positive fixed number. Let $\theta_{1,a}(x) = \text{essinf}_{y \in [x,a]} \theta_1(y)$,

$$\tilde{\theta}_1(x) = \begin{cases} \theta_{1,a}(x) & \text{if } x \in [0, a]; \\ \bar{\theta}_1 = \text{const} & \text{if } x \in [a, \infty); \end{cases}, \quad \theta_1 = \text{essinf}_{x \in \mathbb{R}_+} \theta_1(x), \quad \Theta_2 = \text{esssup}_{x \in \mathbb{R}_+} \theta_2(x)$$

The next theorem was proved in [17].

Theorem 2.10. *Let $\theta_1(x)$ and $\theta_2(x)$ measurable functions on \mathbb{R}_+ . Suppose that there exists a positive number a for all $x > a$ holds $\theta_1(x) = \bar{\theta}_1 = \text{const}$, $\theta_2(x) = \bar{\theta}_2 = \text{const}$ and $1 < \theta_1 \leq \tilde{\theta}_1(x) \leq \theta_2(x) \leq \Theta_2 < \infty$ for a.a. If*

$$G = \sup_{t>0} \int_0^t [v(x)]^{\theta_2(x)} \left(\int_t^{\infty} u^{\tilde{\theta}_1(x)}(\tau) d\tau \right)^{\frac{\theta_2(x)}{(\tilde{\theta}_1)'(x)}} dx < \infty$$

then the operator $\tilde{H}_{v,u}$ is bounded from $L_{\theta_1(\cdot)}(\mathbb{R}^+)$ to $L_{\theta_2(\cdot)}(\mathbb{R}^+)$.

3 The Main Results

Theorem 3.1. *Assume that $p(\cdot) \in \mathbb{P}_{\infty}^{\log}(\Omega)$, and $\theta_1(x)$ and $\theta_2(x)$ are measurable functions on \mathbb{R}_+ . Suppose that there exists a positive number a such that for all $t > a$ we have $\theta_1(x) = \bar{\theta}_1 = \text{const}$, $\theta_2(x) = \bar{\theta}_2 = \text{const}$ and $1 < \theta_1 \leq \tilde{\theta}_1(x) \leq \theta_2(x) \leq \Theta_2 < \infty$ for a.a., the positive measurable functions w_1 u w_2 satisfy the condition*

$$A = \sup_{x \in \Omega, t>0} \int_0^t (w_2(x, r))^{\theta_2(r)} \left(\int_t^{\infty} \left(\frac{1}{w_1(x, s)s} \right)^{[\tilde{\theta}_1(r)]'} ds \right)^{\frac{\theta_2(r)}{[\tilde{\theta}_1(r)]'}} dr < \infty \quad (3.1)$$

Then the maximal operator M from $GM_{p(\cdot), \theta_1(\cdot), w_1(\cdot)}(\Omega)$ to $GM_{p(\cdot), \theta_2(\cdot), w_2(\cdot)}(\Omega)$ is bounded.

Corollary 3.1. Let $p(\cdot) \in \mathbb{P}_\infty^{\log}(\Omega)$, $w_1(x, r) = w_2(x, r) = r^{\beta(x)}$. If

$$\inf_{x \in \Omega, r > 0} (\beta(x) + 1)[\tilde{\theta}_1(r)]' > 1, \quad (3.2)$$

$$\sup_{x \in \Omega, t > 0} \int_0^t r^{\theta_2(r)\beta(x)} \frac{t^{[-(\beta(x)+1)[\tilde{\theta}_1(r)]' + 1] \frac{\theta_2(r)}{[\tilde{\theta}_1(r)]'}}}{[(\beta(x) + 1)[\tilde{\theta}_1(r)] - 1] \frac{\theta_2(r)}{[\tilde{\theta}_1(r)]'}} dr < \infty. \quad (3.3)$$

Then the maximal operator M from $GM_{p(\cdot), \theta_1(\cdot), r^{\beta(\cdot)}}(\Omega)$ to $GM_{p(\cdot), \theta_2(\cdot), r^{\beta(\cdot)}}(\Omega)$ is bounded.

The following theorems give Spanne-type results on the boundedness of the Riesz potential I^α in global Morrey-type spaces with variable exponent $GM_{p(\cdot), \theta(\cdot), w(\cdot)}(\Omega)$. In the following theorem $\alpha = \text{const}$.

Theorem 3.2. Assume that $p(\cdot) \in \mathbb{P}_\infty^{\log}(\Omega)$, the constant number α satisfies the condition $\alpha > 0$, $(\alpha p(\cdot))_+ = \sup_{x \in \Omega} \alpha p(x) < n$. Let $\theta_1(x)$ and $\theta_2(x)$ be measurable functions on R_+ . Suppose that there exists a positive number a such that for all $x > a$ we have $\theta_1(x) = \bar{\theta}_1 = \text{const}$, $\theta_2(x) = \bar{\theta}_2 = \text{const}$ and $1 < \theta_1 \leq \tilde{\theta}_1(x) \leq \theta_2(x) \leq \Theta_2 < \infty$ for a.a., the functions $p_1(x)$ u $p_2(x)$ satisfy $\frac{1}{p_2(x)} = \frac{1}{p_1(x)} - \frac{\alpha(x)}{n}$, and the functions w_1 u w_2 satisfy the condition

$$T = \sup_{x \in \Omega, t > 0} \int_0^t (w_2(x, r))^{\theta_2(r)} \left(\int_t^\infty \left(\frac{s^{\alpha-1}}{w_1(x, s)} \right)^{[\tilde{\theta}_1(r)]'} ds \right)^{\frac{\theta_2(r)}{[\tilde{\theta}_1(r)]'}} dr < \infty. \quad (3.4)$$

Then the operators I_α u M_α from $GM_{p_1(\cdot), \theta_1(\cdot), w_1(\cdot)}(\Omega)$ to $GM_{p_2(\cdot), \theta_2(\cdot), w_2(\cdot)}(\Omega)$ are bounded.

Corollary 3.2. Let $p(\cdot) \in P_\infty^{\log}(\Omega)$, $w_1(x, r) = w_2(x, r) = r^{\beta(x)}$. If

$$\sup_{x \in \Omega, r > 0} (\alpha - \beta(x) - 1)[\tilde{\theta}_1(r)]' < -1, \quad (3.5)$$

$$\sup_{x \in \Omega, t > 0} \int_0^t r^{\theta_2(r)\beta(x)} \frac{t^{[[\alpha - \beta(x) - 1][\tilde{\theta}_1(r)]' + 1] \frac{\theta_2(r)}{[\tilde{\theta}_1(r)]'}}}{[\beta(x) + 1 - \alpha][\tilde{\theta}_1(r)]' - 1} dr < \infty, \quad (3.6)$$

then the operators I_α u M_α from $GM_{p(\cdot), \theta_1(\cdot), r^{\beta(\cdot)}}(\Omega)$ to $GM_{p(\cdot), \theta_2(\cdot), r^{\beta(\cdot)}}(\Omega)$ are bounded.

In the following theorem $\alpha(x)$ is a variable exponent.

Theorem 3.3. Assume that $p(\cdot) \in \mathbb{P}_\infty^{\log}(\Omega)$, the function $\alpha(x)$ satisfies the condition $\alpha(x) > 0$, $(\alpha(\cdot)p(\cdot))_+ = \sup_{x \in \Omega} \alpha(x)p(x) < n$. Let $\theta_1(x)$ and $\theta_2(x)$ be measurable functions on R_+ . Suppose that there exists a positive number a such that for all $x > a$ we have $\theta_1(x) = \bar{\theta}_1 = \text{const}$, $\theta_2(x) = \bar{\theta}_2 = \text{const}$ and $1 < \theta_1 \leq \tilde{\theta}_1(x) \leq \theta_2(x) \leq \Theta_2 < \infty$ for a.a., the functions $p_1(x)$ u $p_2(x)$ satisfy $\frac{1}{p_2(x)} = \frac{1}{p_1(x)} - \frac{\alpha(x)}{n}$, the functions w_1 u w_2 satisfy the condition

$$T = \sup_{x \in \Omega, t > 0} \int_0^t (w_2(x, r))^{\theta_2(r)} \left(\int_t^\infty \left(\frac{s^{\alpha(x)-1}}{w_1(x, s)} \right)^{[\tilde{\theta}_1(r)]'} ds \right)^{\frac{\theta_2(r)}{[\tilde{\theta}_1(r)]'}} dr < \infty.$$

Then the operators $\frac{1}{(1+|x|)^{\gamma(x)}}I^{\alpha(\cdot)}$ u $\frac{1}{(1+|x|)^{\gamma(x)}}M^{\alpha(\cdot)}$ from $GM_{p_1(\cdot),\theta_1(\cdot),w_1(\cdot)}(\Omega)$ to $GM_{p_2(\cdot),\theta_2(\cdot),w_2(\cdot)}(\Omega)$ are bounded.

In the following theorem $\alpha = \text{const}$.

Theorem 3.4. Assume that $p(\cdot) \in P_{\infty}^{\log}(\Omega)$, bounded measurable functions $\theta_1(x)$, $\theta_2(x)$ satisfy $1 < \theta_{1-}(\cdot) \leq \theta_1(\cdot) \leq \theta_{1+}(\cdot)$, $1 < \theta_{2-}(\cdot) \leq \theta_2(\cdot) \leq \theta_{2+}(\cdot)$, and there exists a such that for all $t > a$ we have $\theta_1(t) = \text{const}$, $\theta_2(t) = \text{const}$. Let the positive measurable functions $w_1(x, r)$, $w_2(x, r)$ satisfy the condition

$$\sup_{x \in \Omega} \left\| w_2(x, r) \left\| \frac{1}{tw_1(x, t)} \right\|_{L_{\theta'_1(\cdot)}(r, \infty)} \right\|_{L_{\theta_2(\cdot)}(0, \infty)} < \infty.$$

Then the singular integral operator T from $GM_{p(\cdot),\theta_1(\cdot),w_1(\cdot)}$ to $GM_{p(\cdot),\theta_2(\cdot),w_2(\cdot)}$ is bounded.

Proof of Theorem 3.1. According to the definition and to Theorem 2.10, using the Holder inequality with variable exponents θ , θ' , we infer

$$\begin{aligned} \|Mf\|_{GM_{p(\cdot),\theta_2(\cdot),w_2(\cdot)}(\Omega)} &= \sup_{x \in \Omega} \left\| \frac{w_2(x, r)}{r^{\eta_p(x, r)}} \|Mf\|_{L_{p(\cdot)}(B(x, r))} \right\|_{L_{\theta_2(\cdot)}(0, \infty)} \\ &\leq C \sup_{x \in \Omega} \left\| w_2(x, r) \int_r^{\infty} t^{-\eta_p(x, t)-1} \|f\|_{L_{p(\cdot)}(B(x, t))} dt \right\|_{L_{\theta_2(\cdot)}(0, \infty)}, \end{aligned}$$

we denote

$$\tilde{H}_{v,u}f(r) = v(r) \int_r^{\infty} g(t)u(t)dt,$$

here $v(r) = w_2(x, r)$, $g(t) = \frac{w_1(x, t)}{t^{\eta_p(x, t)}} \|f\|_{L_{p(\cdot)}(B(x, t))}$, $u(t) = \frac{1}{w_1(x, t)t}$ for every fixed $x \in \Omega$. Then the condition (2.2) has the form (3.1), from which boundedness of the operator $\tilde{H}_{v,w}f(r)$ from $L_{\theta_1(\cdot)}(0, \infty)$ to $L_{\theta_2(\cdot)}(0, \infty)$ follows. Consequently, we have

$$\begin{aligned} \|Mf\|_{GM_{q(\cdot),\theta_2(\cdot),w_2(\cdot)}(\Omega)} &\leq A \cdot \sup_{x \in \Omega} \left\| w_1(x, t) t^{-\eta_p(x, t)} \|f\|_{L_{p(\cdot)}(B(x, t))} \right\|_{L_{\theta_1(\cdot)}(0, \infty)} = \\ &= A \cdot \|f\|_{GM_{p(\cdot),\theta_1(\cdot),w_1(\cdot)}}, \end{aligned}$$

this means that the operator M from $GM_{p(\cdot),\theta_1(\cdot),w_1(\cdot)}$ to $GM_{q(\cdot),\theta_2(\cdot),w_2(\cdot)}$ is bounded. \square

Proof of Corollary 3.1. The condition (3.1) has the form

$$\sup_{x \in \Omega, t > 0} \int_0^t r^{\theta_2(r)\beta(x)} \left(\int_t^{\infty} s^{-[\beta(x)+1][\tilde{\theta}_1(r)]'} ds \right)^{\frac{\theta_2(r)}{[\tilde{\theta}_1(r)]'}} dr < \infty.$$

By the convergence of the inner integral, we obtain the conditions (3.2) and (3.3). \square

Proof of Theorem 3.2. Using the definition and the Theorem 2.8, we have

$$\begin{aligned} \|I^\alpha\|_{GM_{q(\cdot),\theta_2(\cdot),w_2(\cdot)}(\Omega)} &= \sup_{x \in \Omega} \|w_2(x, r) r^{-\eta_q(x,r)} \|I_\alpha f\|_{L_{q(\cdot)}(B(x,r))}\|_{L_{\theta_2(\cdot)}(0,\infty)} \leq \\ &\leq C \sup_{x \in \Omega} \|w_2(x, r) \int_r^\infty t^{-\eta_q(x,t)-1} \|f\|_{L_{p(\cdot)}(B(x,t))} dt\|_{L_{\theta_2(\cdot)}(0,\infty)} \end{aligned}$$

We denote

$$\tilde{H}_{v,u} f(r) = v(r) \int_r^\infty g(t) u(t) dt,$$

we denote by $v(r) = w_2(x, r)$, $g(t) = \frac{w_1(x,t)}{t^{\eta_p(x,t)}} \|f\|_{L_{p(\cdot)}(B(x,t))}$, $u(t) = \frac{t^{\eta_p(x,t)-\eta_q(x,t)-1}}{w_1(x,t)}$ for every fixed $x \in \Omega$. Then the condition (2.2) has the form (3.4), from which boundedness of the operator $\tilde{H}_{v,u} f(r)$ from $L_{\theta_1(\cdot)}(0, \infty)$ to $L_{\theta_2(\cdot)}(0, \infty)$ follows. Consequently, we have

$$\begin{aligned} \|I^\alpha f\|_{GM_{q(\cdot),\theta_2(\cdot),w_2(\cdot)}(\Omega)} &\leq T \cdot \sup_{x \in \Omega} \|w_1(x, t) t^{-\eta_p(x,t)} \|f\|_{L_{p(\cdot)}(B(x,t))}\|_{L_{\theta_1(\cdot)}(0,\infty)} = \\ &= T \cdot \|f\|_{GM_{p(\cdot),\theta_1(\cdot),w_1(\cdot)}}, \end{aligned}$$

this means that the operator I^α from $GM_{p_1(\cdot),\theta_1(\cdot),w_1(\cdot)}$ to $GM_{p_2(\cdot),\theta_2(\cdot),w_2(\cdot)}$ is bounded. \square

Proof of Corollary 3.2. The condition (3.4) takes the form

$$\sup_{x \in \Omega, t > 0} \int_0^t r^{\theta_2(r)\beta(x)} \left(\int_t^\infty s^{(\alpha-1-\beta(x))[\tilde{\theta}_1(r)]'} ds \right)^{\frac{\theta_2(r)}{[\tilde{\theta}_1(r)]'}} dr < \infty.$$

By the convergence of the inner integral, we deduce the conditions (3.5) and (3.6). \square

Proof of Theorem 3.3. The proof of this theorem the same as the proof of Theorem 3.2, it is sufficient to put $\frac{1}{(1+|x|)^{\gamma(x)}} I^{\alpha(\cdot)} f(x)$ instead of $I^\alpha f(x)$. \square

References

- [1] C.B. Morrey, *On the solutions of quasi-linear elliptic partial differential equations*. Trans.Am.Math.Soc.43 (1938), 126-166.
- [2] V. Burenkov, H.Guliev *Necessary and sufficient conditions for boundedness of the maximal operator in the local Morrey-type spaces* . Studia Math.163 (2004),157-176.
- [3] V. Burenkov, V.Guliev *Necessary and sufficient conditions for boundedness of the Riesz potential in the local Morrey-type spaces* . Potential Anal.31 (2009),1-39
- [4] V. Burenkov, H.Guliev, V.Guliev *Necessary and sufficient conditions for boundedness of the Riesz potential in the local Morrey-type spaces* . Doklady Ross.Akad.Nauk.Matematika,412, no.5(2007), 585-589 (in Russian).English transl. in Acad.Sci.Dokl.Math.,76 (2007).
- [5] A. Almeida, J.Hasanov, S.Samko, *Maximal and potential operators in variable exponent Morrey spaces* . Georgian Mathematical Journal , Vol.15 (2008), no. 2., 195-208

- [6] A. Almeida, P.Hasto, *Besov spaces with variable smoothness and integrability*. Journal of functional anal. , Vol.258 (2010), 1628-1655
- [7] A. Almeida, S.Samko, *Characterization of Riesz and Bessel potentials on variable Lebesgue spaces*. Journal of function spaces and applications. , Vol.4 (2006)(2), 113-144
- [8] A. Almeida, S.Samko, *Embeddings of variable Hajlasz-Sobolev spaces into Holder spaces of variable order*. J.Math.Anal. and Appl., 353(2009), 489-496.
- [9] A. Almeida, S.Samko, *Fractional and hypersingular operators in variable exponent spaces on metric measure spaces*. Meditter.J.Math.,6(2009), 215-232.
- [10] J. Alvarez, C.Perez, *Estimates with A_∞ weights for various singular integral operators*. Boll.Un.Mat.Ital.A(7)8(1994),no.1, 123-133.
- [11] V. Guliyev, S.Samko, *Boundedness of maximal, potential type, and singular integral operators in the generalized variable exponent Morrey spaces*. J.Math.Sciences, Vol.170(2010),423-441.
- [12] V. Guliyev, S.Samko, *Maximal, potential, and singular operators in the generalized variable exponent Morrey spaces on unbounded sets*. J.Math.Sciences, Vol.193(2013),228-247.
- [13] I. Sharafutdinov *The topology of spaces $L_p(\cdot)$* . Math.Notes.1(1978), no.2
- [14] L. Diening, *Maximal function on generalized Lebesgue spaces $L_p(\cdot)$* . Math.Inequal.Appl,7(2004),no.2, 245-253
- [15] L. Diening, *Riesz potential and Sobolev embeddings on generalized Lebesgue and Sobolev spaces $L_{p(\cdot)}$ and $W_{k,p(\cdot)}$* . Math.Nachr.268(2004),31-43.
- [16] V. Kokilashvili, S.Samko, *On Sobolev theorem for the Riesz type potentials in Lebesgue spaces with variable exponent*. Z.Anal.Anwend.,22.N.4(2003), 899-910
- [17] D. E. Edmunds, V. Kokilashvili, and A. Meskhi *On the boundedness and compactness of weight Hardy operators in $L_p(x)$ spaces*. Georgian Math. J., Vol. 12 (2005), No. 1, 27-44.

Nurzhan Bokayev, Zhomart Onerbek
 Department of Mechanics and Mathematics
 L.N. Gumilyov Eurasian National University
 5 Munaitpasov St,
 010008 Astana, Nur-Sultan, Kazakhstan
 E-mail: onerbek.93@mail.ru