

CONVOLUTION WITH THE KERNEL $e^{s\langle x \rangle^q}$, $q \geq 1, s > 0$ WITHIN ULTRADISTRIBUTION SPACES

STEVAN PILIPOVIĆ, BOJAN PRANGOSKI, AND ĐORĐE VUČKOVIĆ

ABSTRACT. We consider the existence of convolution of Roumieu type ultradistribution with the kernel $e^{s(1+|x|^2)^{q/2}}$, $q \geq 1, s \in \mathbb{R} \setminus \{0\}$.

0. INTRODUCTION

In time frequency analysis, when one needs to analyse modulation or Wiener amalgam spaces with super-exponential weights, or more generally, to study ultradistributions of quasi-analytical classes, kernels of the form $e^{s(1+|x|^2)^{q/2}}$, $q \geq 1, s \in \mathbb{R} \setminus \{0\}$ are involved. Our aim is to show that it is possible to study convolution with such kernel beyond a very general theory we have developed in [17].

The problem of when a distribution in $\mathcal{D}'(\mathbb{R}^d)$ admits a \mathcal{D}' -convolution with the Gaussian $e^{s|x|^2}$, $s \in \mathbb{R} \setminus \{0\}$, was studied in [21] and the distributions that do satisfy this property were characterised by their growth behaviour. Later, the same problem was also solved in the setting of non-quasianalytic ultradistributions in [15] and applied in extending the Anti-Wick quantisation to more general symbol classes. Our aim in this article is to find conditions on the Gelfand-Shilov ultradistributions in $\mathcal{S}'_{\{p^{1/q_1}\}}^{\{M_p\}}(\mathbb{R}^d)$ which are sufficient for the existence of convolution with the function $x \mapsto e^{s\langle x \rangle^q}$, $s \in \mathbb{R} \setminus \{0\}$, $q_1 > q \geq 1$, where, as standard, $\langle x \rangle = (1 + |x|^2)^{1/2}$ (of course, when $q = 2$, $e^{s\langle x \rangle^2}$ is just a constant multiple of the Gaussian). Here $\{M_p\}_{p \in \mathbb{N}}$ is a Gevrey type sequence, bounded from below by $\{p!\}_{p \in \mathbb{N}}$, which controls the growth of the derivatives of the test functions.

The definition of the convolution of $S \in \mathcal{S}'_{\{p^{1/q_1}\}}^{\{M_p\}}(\mathbb{R}^d)$ with $e^{s\langle \cdot \rangle^q}$ we employ is the following: we say S is convolvable with $e^{s\langle \cdot \rangle^q}$ if for every test function $\varphi \in \mathcal{S}_{\{p^{1/q_1}\}}^{\{M_p\}}(\mathbb{R}^d)$, $(\varphi * e^{s\langle \cdot \rangle^q})S \in \mathcal{D}'_{L^1}^{\{M_p\}}(\mathbb{R}^d)$ (the ultradistributional analogue of the Schwartz space $\mathcal{D}'_{L^1}(\mathbb{R}^d)$) and $S * e^{s\langle \cdot \rangle^q} \in \mathcal{S}'_{\{p^{1/q_1}\}}^{\{M_p\}}(\mathbb{R}^d)$ is defined as

$$\langle S * e^{s\langle \cdot \rangle^q}, \varphi \rangle = \langle (\varphi * e^{s\langle \cdot \rangle^q})S, 1 \rangle$$

2010 *Mathematics Subject Classification.* 46F10 46F05.

Key words and phrases. Convolution with exponentials, Gelfand-Shilov spaces, ultradistributions.

B. Prangoski was partially supported by the bilateral project “Microlocal analysis and applications” funded by the Macedonian and Serbian academies of sciences and arts.

The work presented in this paper is partially supported by Ministry of Education and Science, Republic of Serbia, project no. 174024.

(see Subsection 1.1 for the reasons why this is indeed a well defined element of $\mathcal{S}'_{\{p!^{1/q_1}\}}^{\{M_p\}}(\mathbb{R}^d)$).

When $q_1 \leq 1$, this reduces to the definition of $\mathcal{S}'_{\{p!^{1/q_1}\}}^{\{M_p\}}$ -convolution of ultradistributions, see [17, Definition 5.7 and Theorem 5.8] (see also [6, 14, 9, 4]) and is analogous to one of the equivalent formulations of \mathcal{S}' -convolution in the distributional setting (see [19, 20]; see also [1, 5, 12, 13]).

When studying the convolution with the Gaussian, one of the key ingredients employed in [21] and [15] is the fact that $e^{s|x-t|^2}$ splits into a product of two parts: $e^{s|x|^2}$ and $e^{-2sxt+s|t|^2}$. However, when $q \neq 2$ this is no longer the case, and thus we can not apply the same ideas from [21] and [15]. In fact, a significant part of Section 2 is devoted to finding good bounds on the growth of the derivatives of $e^{s\langle x \rangle^q}$ before we provide sufficient conditions in Theorem 2.6. It is important to emphasise that, when $q = 2$, these are analogous to the ones given in the distributional setting as well as in the non-quasianalytic ultradistributional setting (see the remark at the very end of Section 2).

In Section 3 we give also necessary conditions for the existence of the convolution with $e^{s\langle \cdot \rangle^q}$, when $s > 0$. When $q = 1$ they turn out to be the same as the sufficient conditions, and thus providing a characterisation of the ultradistributions in $\mathcal{S}'_{\{p!^{1/q_1}\}}^{\{M_p\}}(\mathbb{R}^d)$ which are convolvable with $e^{s\langle \cdot \rangle}$, $s > 0$.

The main results of the paper can be summarised as follows (see Theorem 2.6 and Corollaries 3.5, 3.6 and 3.7):

Let $q_1 > q \geq 1$, $s \in \mathbb{R} \setminus \{0\}$ and $S \in \mathcal{S}'_{\{p!^{1/q_1}\}}^{\{p!\}}(\mathbb{R}^d)$.

1. If

$$e^{s\langle \cdot \rangle^q} e^{k\langle \cdot \rangle^{(q-1)q_1/(q_1-1)}} S \in \mathcal{D}'_{L^1}^{\{p!\}}(\mathbb{R}^d), \quad \text{for all } k \geq 0,$$

then the $\mathcal{S}'_{\{p!^{1/q_1}\}}^{\{p!\}}$ -convolution of S and $e^{s\langle \cdot \rangle^q}$ exists.

2. When $q = 1$, the $\mathcal{S}'_{\{p!^{1/q_1}\}}^{\{p!\}}$ -convolution of S and $e^{s\langle \cdot \rangle}$, $s > 0$, exists if and only if $e^{s\langle \cdot \rangle} S \in \mathcal{D}'_{L^1}^{\{p!\}}(\mathbb{R}^d)$.

3. Let $S \in \mathcal{S}'_{\{p!^{1/q_1}\}}^{\{p!^{2-1/q}\}}(\mathbb{R}^d)$. If the $\mathcal{S}'_{\{p!^{1/q_1}\}}^{\{p!^{2-1/q}\}}$ -convolution of S and $e^{s\langle \cdot \rangle^q}$, $s > 0$, exists then the $\mathcal{S}'_{\{p!^{1/q_1}\}}^{\{p!^{2-1/q}\}}$ -convolution of S and $e^{s'\langle \cdot \rangle^q}$ also exists for all $s' < s$, $s' \neq 0$.

Note that the Beurling ultradistribution spaces give even more flexibility in the analysis of these problems, so most of the results (with appropriate changes) can be easily transferred to the ultradistribution spaces of Beurling class.

1. PRELIMINARIES

Let $\{M_p\}_{p \in \mathbb{N}}$ be a sequence of positive numbers satisfying $M_0 = M_1 = 1$. Unless stated otherwise, we will always assume the following conditions on $\{M_p\}_{p \in \mathbb{N}}$ (cf. [10]):

$$(M.1) \quad M_p^2 \leq M_{p-1} M_{p+1}, \quad p \in \mathbb{Z}_+;$$

$$(M.2) \quad M_p \leq c_0 H^p \min_{0 \leq q \leq p} \{M_{p-q} M_q\}, \quad p, q \in \mathbb{N}, \text{ for some } c_0, H \geq 1;$$

(M.5) there exists $q > 0$ such that M_p^q is strongly non-quasianalytic (see [10]), i.e., there exists $c_0 \geq 1$ such that $\sum_{j=p+1}^{\infty} M_{j-1}^q / M_j^q \leq c_0 p M_p^q / M_{p+1}^q$, $\forall p \in \mathbb{Z}_+$;

(M.6) $p! \subset M_p$, i.e. there exist $C, L > 0$ such that $p! \leq CL^p M_p$, $\forall p \in \mathbb{N}$.

For each $\alpha \in \mathbb{N}^d$, we set $M_\alpha = M_{|\alpha|}$. Note that the Gevrey sequences $\{p!^\kappa\}_{p \in \mathbb{N}}$, with $\kappa \geq 1$, satisfy all of the above conditions.

Let $q > 0$. For every $h > 0$, we denote by $\mathcal{S}_{p^{1/q}, h}^{M_p}(\mathbb{R}^d)$ the Banach space of all $\varphi \in \mathcal{C}^\infty(\mathbb{R}^d)$ for which the following norm is finite

$$\sup_{\alpha \in \mathbb{N}^d} h^{|\alpha|} \|e^{h|\cdot|^q} \partial^\alpha \varphi\|_{L^\infty(\mathbb{R}^d)} / M_\alpha < \infty.$$

Recall, the Gelfand-Shilov test space $\mathcal{S}_{\{p^{1/q}\}}^{\{M_p\}}(\mathbb{R}^d)$ is defined as

$$\mathcal{S}_{\{p^{1/q}\}}^{\{M_p\}}(\mathbb{R}^d) = \varinjlim_{h \rightarrow 0^+} \mathcal{S}_{p^{1/q}, h}^{M_p}(\mathbb{R}^d).$$

Its strong dual $\mathcal{S}'_{\{p^{1/q}\}}^{\{M_p\}}(\mathbb{R}^d)$ is the space of Gelfand-Shilov ultradistributions; we refer to [2, 8, 17] for its properties. Next, for each $h > 0$, let $\mathcal{D}_{L^\infty}^{M_p, h}(\mathbb{R}^d)$ be the Banach space of all $\varphi \in \mathcal{C}^\infty(\mathbb{R}^d)$ such that $\sup_{\alpha \in \mathbb{N}^d} h^{|\alpha|} \|\partial^\alpha \varphi\|_{L^\infty(\mathbb{R}^d)} / M_\alpha < \infty$, and define

$$\mathcal{D}_{L^\infty}^{\{M_p\}}(\mathbb{R}^d) = \varinjlim_{h \rightarrow 0^+} \mathcal{D}_{L^\infty}^{M_p, h}(\mathbb{R}^d);$$

it is a complete barrelled (DF) -space (see [7, Section 4.3]). Fix $q > 0$ and denote by $\dot{\mathcal{B}}^{\{M_p\}}(\mathbb{R}^d)$ the closure of $\mathcal{S}_{\{p^{1/q}\}}^{\{M_p\}}(\mathbb{R}^d)$ in $\mathcal{D}_{L^\infty}^{\{M_p\}}(\mathbb{R}^d)$ and let $\mathcal{D}'_{L^1}^{\{M_p\}}(\mathbb{R}^d)$ be the strong dual of $\dot{\mathcal{B}}^{\{M_p\}}(\mathbb{R}^d)$. As $\mathcal{S}_{\{p^{1/q'}\}}^{\{M_p\}}(\mathbb{R}^d)$, with $q' > q$, is continuously and densely injected into $\mathcal{S}_{\{p^{1/q}\}}^{\{M_p\}}(\mathbb{R}^d)$, $\dot{\mathcal{B}}^{\{M_p\}}(\mathbb{R}^d)$ is the closure of $\mathcal{S}_{\{p^{1/q'}\}}^{\{M_p\}}(\mathbb{R}^d)$ in $\mathcal{D}_{L^\infty}^{\{M_p\}}(\mathbb{R}^d)$, for any $q' > 0$. We mention several important properties of the space $\dot{\mathcal{B}}^{\{M_p\}}(\mathbb{R}^d)$, $\mathcal{D}'_{L^1}^{\{M_p\}}(\mathbb{R}^d)$ and $\mathcal{D}_{L^\infty}^{\{M_p\}}(\mathbb{R}^d)$ and refer to [7, Section 4.3] for the complete account. First, $\dot{\mathcal{B}}^{\{M_p\}}(\mathbb{R}^d)$ is a complete barrelled (DF) -space and thus $\mathcal{D}'_{L^1}^{\{M_p\}}(\mathbb{R}^d)$ is an (F) -space. The strong dual of the latter is topologically isomorphic to $\mathcal{D}_{L^\infty}^{\{M_p\}}(\mathbb{R}^d)$ (i.e. $\mathcal{D}_{L^\infty}^{\{M_p\}}(\mathbb{R}^d)$ is the strong bidual of $\dot{\mathcal{B}}^{\{M_p\}}(\mathbb{R}^d)$). Denote by $\mathcal{D}_{L^\infty, c}^{\{M_p\}}(\mathbb{R}^d)$ the space $\mathcal{D}_{L^\infty}^{\{M_p\}}(\mathbb{R}^d)$ equipped with the topology of compact convex circled convergence from the duality $\langle \mathcal{D}'_{L^1}^{\{M_p\}}(\mathbb{R}^d), \mathcal{D}_{L^\infty}^{\{M_p\}}(\mathbb{R}^d) \rangle$. Then $\mathcal{D}_{L^\infty, c}^{\{M_p\}}(\mathbb{R}^d)$ is a complete space whose topology is weaker than the original one, $\dot{\mathcal{B}}^{\{M_p\}}(\mathbb{R}^d)$ is continuously and densely injected into $\mathcal{D}_{L^\infty, c}^{\{M_p\}}(\mathbb{R}^d)$ and the strong dual of $\mathcal{D}_{L^\infty, c}^{\{M_p\}}(\mathbb{R}^d)$ is topologically isomorphic to $\mathcal{D}'_{L^1}^{\{M_p\}}(\mathbb{R}^d)$ (see [17, Section 5]).

1.1. Convolution of Gelfand-Shilov ultradistributions. Let $0 < q \leq 1$ and $S, T \in \mathcal{S}'_{\{p^{1/q}\}}^{\{M_p\}}(\mathbb{R}^d)$. The $\mathcal{S}'_{\{p^{1/q}\}}^{\{M_p\}}$ -convolution of S and T is said to exist if $(S \otimes T)\varphi^\Delta \in \mathcal{D}'_{L^1}^{\{M_p\}}(\mathbb{R}^{2d})$, for all $\varphi \in \mathcal{S}_{\{p^{1/q}\}}^{\{M_p\}}(\mathbb{R}^d)$, where $\varphi^\Delta(x, y) = \varphi(x+y)$. When this is the case, $S * T \in \mathcal{S}'_{\{p^{1/q}\}}^{\{M_p\}}(\mathbb{R}^d)$

is defined as

$$\langle S * T, \varphi \rangle = {}_{\mathcal{D}'_{L^1} \{M_p\}(\mathbb{R}^{2d})} \langle (S \otimes T) \varphi^\Delta, 1 \rangle_{{}_{\mathcal{D}'_{L^\infty, c} \{M_p\}(\mathbb{R}^{2d})}}, \quad \varphi \in \mathcal{S}'_{\{p^{1/q}\}} \{M_p\}(\mathbb{R}^d),$$

where 1 denotes the constant one function; see the comments after [17, Definition 5.7] for the reason why $S * T$ is indeed a continuous functional on $\mathcal{S}'_{\{p^{1/q}\}} \{M_p\}(\mathbb{R}^d)$. Furthermore, we have the following result.

Theorem 1.1. [17, Theorem 5.8 and Remark 5.9] *Let $0 < q \leq 1$ and $S, T \in \mathcal{S}'_{\{p^{1/q}\}} \{M_p\}(\mathbb{R}^d)$. The following statements are equivalent:*

- (i) *the $\mathcal{S}'_{\{p^{1/q}\}} \{M_p\}$ -convolution of S and T exists;*
- (ii) *for all $\varphi \in \mathcal{S}'_{\{p^{1/q}\}} \{M_p\}(\mathbb{R}^d)$, $(\varphi * \check{T})S \in \mathcal{D}'_{L^1} \{M_p\}(\mathbb{R}^d)$;*
- (iii) *for all $\varphi \in \mathcal{S}'_{\{p^{1/q}\}} \{M_p\}(\mathbb{R}^d)$, $(\varphi * \check{S})T \in \mathcal{D}'_{L^1} \{M_p\}(\mathbb{R}^d)$.*

Furthermore, when the $\mathcal{S}'_{\{p^{1/q}\}} \{M_p\}$ -convolution of S and T exists,

$$\langle S * T, \varphi \rangle = {}_{\mathcal{D}'_{L^1} \{M_p\}(\mathbb{R}^d)} \langle (\varphi * \check{T})S, 1 \rangle_{{}_{\mathcal{D}'_{L^\infty, c} \{M_p\}(\mathbb{R}^d)}} = {}_{\mathcal{D}'_{L^1} \{M_p\}(\mathbb{R}^d)} \langle (\varphi * \check{S})T, 1 \rangle_{{}_{\mathcal{D}'_{L^\infty, c} \{M_p\}(\mathbb{R}^d)}}. \quad (1.1)$$

Our goal is to study the convolvability with $e^{s\langle \cdot \rangle^q}$, $q \geq 1$. We will adopt Theorem 1.1 (ii) as a definition for the existence of the convolution with $e^{s\langle \cdot \rangle^q}$. More precisely:

Definition 1.2. Let $q_1 > q \geq 1$ and $s \in \mathbb{R} \setminus \{0\}$. The ultradistribution $S \in \mathcal{S}'_{\{p^{1/q_1}\}} \{M_p\}(\mathbb{R}^d)$ admits a $\mathcal{S}'_{\{p^{1/q_1}\}} \{M_p\}$ -convolution with the function $e^{s\langle \cdot \rangle^q}$ if $(\varphi * e^{s\langle \cdot \rangle^q})S \in \mathcal{D}'_{L^1} \{M_p\}(\mathbb{R}^d)$, for all $\varphi \in \mathcal{S}'_{\{p^{1/q_1}\}} \{M_p\}(\mathbb{R}^d)$. When this is the case, $S * e^{s\langle \cdot \rangle^q} \in \mathcal{S}'_{\{p^{1/q_1}\}} \{M_p\}(\mathbb{R}^d)$ is defined by

$$\langle S * e^{s\langle \cdot \rangle^q}, \varphi \rangle := {}_{\mathcal{D}'_{L^1} \{M_p\}(\mathbb{R}^d)} \langle (\varphi * e^{s\langle \cdot \rangle^q})S, 1 \rangle_{{}_{\mathcal{D}'_{L^\infty, c} \{M_p\}(\mathbb{R}^d)}}, \quad \varphi \in \mathcal{S}'_{\{p^{1/q_1}\}} \{M_p\}(\mathbb{R}^d). \quad (1.2)$$

To see that (1.2) indeed defines a continuous functional on $\mathcal{S}'_{\{p^{1/q_1}\}} \{M_p\}(\mathbb{R}^d)$, it is enough to prove that the mapping

$$\varphi \mapsto (\varphi * e^{s\langle \cdot \rangle^q})S, \quad \mathcal{S}'_{\{p^{1/q_1}\}} \{M_p\}(\mathbb{R}^d) \rightarrow \mathcal{D}'_{L^1} \{M_p\}(\mathbb{R}^d), \quad (1.3)$$

is continuous (as $\mathcal{D}'_{L^1} \{M_p\}(\mathbb{R}^d)$ is the strong dual of $\mathcal{D}_{L^\infty, c} \{M_p\}(\mathbb{R}^d)$; see the comments before this section). Clearly $\varphi \mapsto (\varphi * e^{s\langle \cdot \rangle^q})S, \mathcal{S}'_{\{p^{1/q_1}\}} \{M_p\}(\mathbb{R}^d) \rightarrow \mathcal{S}'_{\{p^{1/q_1}\}} \{M_p\}(\mathbb{R}^d)$, is continuous and thus has a closed graph. As $(\varphi * e^{s\langle \cdot \rangle^q})S \in \mathcal{D}'_{L^1} \{M_p\}(\mathbb{R}^d)$, $\forall \varphi \in \mathcal{S}'_{\{p^{1/q_1}\}} \{M_p\}(\mathbb{R}^d)$, its graph is closed in $\mathcal{S}'_{\{p^{1/q_1}\}} \{M_p\}(\mathbb{R}^d) \times \mathcal{D}'_{L^1} \{M_p\}(\mathbb{R}^d)$. Now the Ptak closed graph theorem [18, Theorem 8.5, p. 166] implies that (1.3) is continuous ($\mathcal{D}'_{L^1} \{M_p\}(\mathbb{R}^d)$ is a Ptak space since it is an (F) -space; see [18, Section 4.8, p. 162]).

Remark 1.3. Let $q_1 > q \geq 1$ and $s > 0$. If S is a measurable function which satisfies $e^{k\langle \cdot \rangle^q} S \in L^\infty(\mathbb{R}^d)$, $\forall k > 0$, then the ordinary convolution of $e^{k\langle \cdot \rangle^q}$ and S exists. It is straightforward to verify that the $\mathcal{S}'_{\{p^{1/q_1}\}} \{M_p\}$ -convolution also exists and they are the same.

Similarly, when $s < 0$, if S is a measurable function such that $e^{-\varepsilon\langle \cdot \rangle^q} S \in L^\infty(\mathbb{R}^d)$, $\forall \varepsilon > 0$, then both the ordinary and the $\mathcal{S}'_{\{p^{1/q_1}\}}^{\{M_p\}}$ -convolution of S and $e^{s\langle \cdot \rangle^q}$ exist and they coincide.

2. SUFFICIENT CONDITIONS FOR THE CONVOLUTION WITH $e^{s\langle \cdot \rangle^q}$, $q \geq 1$, $s \in \mathbb{R} \setminus \{0\}$

The aim of this section is to give a sufficient condition for an ultradistribution $S \in \mathcal{S}'_{\{p^{1/q_1}\}}^{\{M_p\}}(\mathbb{R}^d)$ to be $\mathcal{S}'_{\{p^{1/q_1}\}}^{\{M_p\}}$ -convolvable with $e^{s\langle \cdot \rangle^q}$, where $q_1 > q \geq 1$ and $s \in \mathbb{R} \setminus \{0\}$.

Before we start, we recall the following multidimensional variant of the Faà di Bruno formula.

Proposition 2.1. ([3, Corollary 2.10]) *Let $|\alpha| = n \geq 1$ and $h(x_1, \dots, x_d) = f(g(x_1, \dots, x_d))$ with $g \in \mathcal{C}^n$ in a neighbourhood of x^0 and $f \in \mathcal{C}^n$ in a neighbourhood of $y^0 = g(x^0)$. Then*

$$\partial^\alpha h(x^0) = \sum_{r=1}^n f^{(r)}(y^0) \sum_{p(\alpha, r)} \alpha! \prod_{j=1}^n \frac{(\partial^{\alpha^{(j)}} g(x^0))^{k_j}}{k_j! (\alpha^{(j)}!)^{k_j}},$$

where

$$\begin{aligned} p(\alpha, r) = & \left\{ (k_1, \dots, k_n; \alpha^{(1)}, \dots, \alpha^{(n)}) \mid \text{for some } 1 \leq s \leq n, k_j = 0 \text{ and } \alpha^{(j)} = 0 \right. \\ & \text{for } 1 \leq j \leq n-s; k_j > 0 \text{ for } n-s+1 \leq j \leq n; \text{ and} \\ & 0 \prec \alpha^{(n-s+1)} \prec \dots \prec \alpha^{(n)} \text{ are such that} \\ & \left. \sum_{j=1}^n k_j = r, \sum_{j=1}^n k_j \alpha^{(j)} = \alpha \right\}. \end{aligned}$$

In the above formula, the convention $0^0 = 1$ is used. The relation \prec used in this proposition is defined in the following way (cf. [3]). We say that $\beta \prec \alpha$ when one of the following holds:

- (i) $|\beta| < |\alpha|$;
- (ii) $|\beta| = |\alpha|$ and $\beta_1 < \alpha_1$;
- (iii) $|\beta| = |\alpha|$, $\beta_1 = \alpha_1, \dots, \beta_k = \alpha_k$ and $\beta_{k+1} < \alpha_{k+1}$ for some $1 \leq k < d$.

We need precise estimates for the growth of the derivatives of $e^{s\langle \cdot \rangle^q}$. For this purpose we first give a bound for the derivatives of $\langle \cdot \rangle^q$.

Lemma 2.2. *For every $q \in \mathbb{R}$ there exists $C \geq 1$ such that*

$$|\partial^\alpha \langle x \rangle^q| \leq C^{|\alpha|} \alpha! \langle x \rangle^{q-|\alpha|}, \quad \text{for all } x \in \mathbb{R}^d, \alpha \in \mathbb{N}^d. \quad (2.1)$$

Proof. We apply the Faà di Bruno formula (Proposition 2.1) with $f(\rho) = \rho^q$ and $g(x) = \langle x \rangle$ to infer, for $\alpha \in \mathbb{N}^d \setminus \{0\}$,

$$|\partial^\alpha \langle x \rangle^q| \leq \sum_{m=1}^{|\alpha|} \langle x \rangle^{q-m} \left(\prod_{j=0}^{m-1} |q-j| \right) \alpha! \sum_{p(\alpha, m)} \prod_{j=1}^{|\alpha|} \frac{|\partial^{\alpha^{(j)}} \langle x \rangle|^{k_j}}{k_j! (\alpha^{(j)}!)^{k_j}}.$$

Notice that

$$\prod_{j=0}^{m-1} |q - j| \leq \prod_{j=0}^{m-1} (|q| + j) \leq (|q| + m)! \leq 2^{|q|+m} |q|! m!.$$

Recall the bound (see [11, (0.1.1), p. 10])

$$|\partial^\beta \langle x \rangle| \leq 2^{|\beta|+1} |\beta|! \langle x \rangle^{1-|\beta|} \leq 2(2d)^{|\beta|} |\beta|! \langle x \rangle^{1-|\beta|}, \quad \text{for all } x \in \mathbb{R}^d, \beta \in \mathbb{N}^d. \quad (2.2)$$

Hence, we deduce

$$|\partial^\alpha \langle x \rangle^q| \leq 2^{|q|} (2d)^{|\alpha|} |q|! \alpha! \langle x \rangle^{q-|\alpha|} \sum_{m=1}^{|\alpha|} 4^m m! \sum_{p(\alpha, m)} \prod_{j=1}^{|\alpha|} \frac{1}{k_j!}.$$

Now, [16, Lemma 7.4] gives

$$\sum_{m=1}^{|\alpha|} m! \sum_{p(\alpha, m)} \prod_{j=1}^{|\alpha|} \frac{1}{k_j!} \leq 2^{|\alpha|(d+1)}. \quad (2.3)$$

Plugging this in the above estimate, we deduce (2.1). \square

Lemma 2.3. *Let $s \in \mathbb{R} \setminus \{0\}$. Then for every $q > 0$ there exists $C \geq 1$ such that*

$$|\partial^\alpha e^{s\langle x \rangle^q}| \leq C^{|\alpha|} \alpha! e^{s\langle x \rangle^q} \sum_{m=1}^{|\alpha|} \frac{\langle x \rangle^{qm-|\alpha|}}{m!}, \quad \text{for all } x \in \mathbb{R}^d, \alpha \in \mathbb{N}^d \setminus \{0\}.$$

Proof. Let $\alpha \in \mathbb{N}^d \setminus \{0\}$. We apply the Faà di Bruno formula (Proposition 2.1) with $f(\rho) = e^{s\rho}$ and $g(x) = \langle x \rangle^q$ together with Lemma 2.2 to infer ($C'_q \geq 1$ is the constant from this lemma)

$$\begin{aligned} |\partial^\alpha e^{s\langle x \rangle^q}| &\leq \alpha! \sum_{m=1}^{|\alpha|} |s|^m e^{s\langle x \rangle^q} \sum_{p(\alpha, m)} \prod_{j=1}^{|\alpha|} \frac{C'_q |\alpha^{(j)}| k_j}{k_j!} \langle x \rangle^{qk_j - |\alpha^{(j)}| k_j} \\ &\leq \alpha! (|s| + 1) C'_q |\alpha| e^{s\langle x \rangle^q} \sum_{m=1}^{|\alpha|} \langle x \rangle^{qm-|\alpha|} \sum_{p(\alpha, m)} \prod_{j=1}^{|\alpha|} \frac{1}{k_j!}. \end{aligned}$$

Now, the claim in the lemma follows from (2.3). \square

As an immediate consequence we have the following result.

Corollary 2.4. *Let $s \in \mathbb{R} \setminus \{0\}$. Then for every $q > 0$ there exists $C \geq 1$ such that*

$$|\partial^\alpha e^{s\langle x \rangle^q}| \leq C^{|\alpha|} \alpha! e^{s\langle x \rangle^q + \langle x \rangle^{q-1}}, \quad \text{for all } x \in \mathbb{R}^d, \alpha \in \mathbb{N}^d.$$

Before we prove the main result of this section, we need a technical lemma.

Lemma 2.5. *For all $x, t \in \mathbb{R}^d$, $q \geq 1$, $s \in \mathbb{R} \setminus \{0\}$, the following inequality holds true:*

$$s\langle x - t \rangle^q - s\langle x \rangle^q \leq q2^{q-1} |s| |t| \langle x \rangle^{q-1} + q2^{q-1} |s| |t| \langle t \rangle^{q-1}. \quad (2.4)$$

Proof. Let $\lambda_1, \lambda_2 \geq 0$. By the mean value theorem, there exists λ' between λ_1 and λ_2 such that

$$s\langle \lambda_1 \rangle^q - s\langle \lambda_2 \rangle^q = sq(\lambda_1 - \lambda_2)\lambda'\langle \lambda' \rangle^{q-2} \quad (2.5)$$

$$\leq q|s||\lambda_1 - \lambda_2|\langle \max\{\lambda_1, \lambda_2\} \rangle^{q-1}. \quad (2.6)$$

We apply (2.6) with $\lambda_1 = |x - t|$ and $\lambda_2 = |x|$. If $|x| \geq |x - t|$, we infer

$$s\langle x - t \rangle^q - s\langle x \rangle^q \leq q|s||t|\langle x \rangle^{q-1}.$$

If $|x| \leq |x - t|$, then (2.6) together with the inequality $\langle x - t \rangle^{q-1} \leq 2^{q-1}\langle x \rangle^{q-1} + 2^{q-1}\langle t \rangle^{q-1}$ gives (2.4). \square

Theorem 2.6. Let $q_1 > q \geq 1$, $s \in \mathbb{R} \setminus \{0\}$ and $S \in \mathcal{S}'_{\{p^{1/q_1}\}}^{\{M_p\}}(\mathbb{R}^d)$. If

$$e^{s\langle \cdot \rangle^q} e^{k\langle \cdot \rangle^{(q-1)q_1/(q_1-1)}} S \in \mathcal{D}'_{L^1}^{\{M_p\}}(\mathbb{R}^d), \quad \text{for all } k \geq 0, \quad (2.7)$$

then the $\mathcal{S}'_{\{p^{1/q_1}\}}^{\{M_p\}}$ -convolution of S and $e^{s\langle \cdot \rangle^q}$ exists.

Remark 2.7. Notice that $e^{k\langle \cdot \rangle^{(q-1)q_1/(q_1-1)}}$ is a multiplier for $\mathcal{S}'_{\{p^{1/q_1}\}}^{\{M_p\}}(\mathbb{R}^d)$ because $(q-1)q_1/(q_1-1) < q < q_1$ and Corollary 2.4.

Remark 2.8. When $q > 1$, (2.7) is equivalent to the following seemingly weaker condition

$$e^{s\langle \cdot \rangle^q} e^{k\langle \cdot \rangle^{(q-1)q_1/(q_1-1)}} S \in \mathcal{S}'_{\{p^{(q_1-1)/(q_1(q-1))}\}}^{\{M_p\}}(\mathbb{R}^d), \quad \text{for all } k \geq 0. \quad (2.8)$$

To see this, denote $q' = (q-1)q_1/(q_1-1) < q_1$ and assume that (2.8) holds. As

$$e^{s\langle \cdot \rangle^q} e^{k\langle \cdot \rangle^{q'}} S = e^{-\langle \cdot \rangle^{q'}} (e^{s\langle \cdot \rangle^q} e^{(k+1)\langle \cdot \rangle^{q'}} S),$$

and $e^{-\langle \cdot \rangle^{q'}} \in \mathcal{S}'_{\{p^{1/q'}\}}^{\{M_p\}}(\mathbb{R}^d)$ (because of Corollary 2.4) it is enough to prove that for every $\psi \in \mathcal{S}'_{\{p^{1/q'}\}}^{\{M_p\}}(\mathbb{R}^d)$ and $T \in \mathcal{S}'_{\{p^{1/q'}\}}^{\{M_p\}}(\mathbb{R}^d)$, $\psi T \in \mathcal{D}'_{L^1}^{\{M_p\}}(\mathbb{R}^d)$. Since $\varphi \mapsto \psi\varphi$, $\mathcal{D}'_{L^\infty}^{\{M_p\}}(\mathbb{R}^d) \rightarrow \mathcal{S}'_{\{p^{1/q'}\}}^{\{M_p\}}(\mathbb{R}^d)$, is continuous, the mapping $\varphi \mapsto \langle T, \psi\varphi \rangle$, $\dot{\mathcal{B}}^{\{M_p\}}(\mathbb{R}^d) \rightarrow \mathbb{C}$, is continuous, and ψT coincides with it on the dense subspace $\mathcal{S}'_{\{p^{1/q'}\}}^{\{M_p\}}(\mathbb{R}^d)$ of $\dot{\mathcal{B}}^{\{M_p\}}(\mathbb{R}^d)$. Consequently, $\psi T \in \mathcal{D}'_{L^1}^{\{M_p\}}(\mathbb{R}^d)$.

Proof. Let $\varphi \in \mathcal{S}'_{\{p^{1/q_1}\}}^{\{M_p\}}(\mathbb{R}^d)$ be arbitrary but fixed. There exists $C, \eta > 0$ such that $|\varphi(x)| \leq Ce^{-\eta\langle x \rangle^{q_1}}$, $\forall x \in \mathbb{R}^d$. Corollary 2.4 implies

$$|D^\alpha \varphi * e^{s\langle \cdot \rangle^q}(x)| \leq CC_1^{|\alpha|} \alpha! e^{2^q \langle x \rangle^{q-1}} \int_{\mathbb{R}^d} e^{-\eta \langle t \rangle^{q_1}} e^{s\langle x-t \rangle^q} e^{2^q \langle t \rangle^{q-1}} dt.$$

Denote $\lambda = (\eta/(4^q q |s|))^{1/(q-1)}$ when $q > 1$ and $\lambda = 1$ when $q = 1$. Lemma 2.5 gives

$$|D^\alpha \varphi * e^{s\langle \cdot \rangle^q}(x)| \leq CC_1^{|\alpha|} \alpha! e^{2^q \langle x \rangle^{q-1}} e^{s\langle x \rangle^q} (I_1 + I_2),$$

where

$$I_1 = \int_{|t| \leq (|x|/\lambda)^{(q-1)/(q_1-1)}} e^{-\eta \langle t \rangle^{q_1}} e^{q2^{q-1}|s||t|\langle x \rangle^{q-1}} e^{q2^{q-1}|s|\langle t \rangle^q} e^{2^q \langle t \rangle^{q-1}} dt,$$

$$I_2 = \int_{|t| \geq (|x|/\lambda)^{(q-1)/(q_1-1)}} e^{-\eta \langle t \rangle^{q_1}} e^{q2^{q-1}|s||t| \langle x \rangle^{q-1}} e^{q2^{q-1}|s| \langle t \rangle^q} e^{2^q \langle t \rangle^{q-1}} dt.$$

Notice that

$$I_1 \leq e^{q2^{q-1}|s|\lambda^{-(q-1)/(q_1-1)} \langle x \rangle^{(q-1)q_1/(q_1-1)}} \|e^{-\eta \langle \cdot \rangle^{q_1}} e^{q2^{q-1}|s| \langle \cdot \rangle^q} e^{2^q \langle \cdot \rangle^{q-1}}\|_{L^1(\mathbb{R}^d)}.$$

To estimate I_2 , let $|t| \geq (|x|/\lambda)^{(q-1)/(q_1-1)}$. When $q > 1$, we infer

$$|t| \langle x \rangle^{q-1} \leq 2^q |t| + 2^q |t| |x|^{q-1} \leq 2^q |t| + 2^q \lambda^{q-1} |t|^{q_1},$$

which implies

$$I_2 \leq \|e^{-(\eta/2) \langle \cdot \rangle^{q_1}} e^{q2^{2q-1}|s| \langle \cdot \rangle} e^{q2^{q-1}|s| \langle \cdot \rangle^q} e^{2^q \langle \cdot \rangle^{q-1}}\|_{L^1(\mathbb{R}^d)}.$$

When $q = 1$, one immediately deduces $I_2 \leq e^2 \|e^{-\eta \langle \cdot \rangle^{q_1}} e^{2|s| \langle \cdot \rangle}\|_{L^1(\mathbb{R}^d)}$. Consequently, there exists $c_1 > 0$ such that

$$|D^\alpha \varphi * e^{s \langle \cdot \rangle^q}(x)| \leq C_2 C_1^{|\alpha|} \alpha! e^{s \langle x \rangle^q} e^{c_1 \langle x \rangle^{(q-1)q_1/(q_1-1)}}, \quad \text{for all } x \in \mathbb{R}^d, \alpha \in \mathbb{N}^d.$$

Corollary 2.4 now verifies that $e^{-s \langle \cdot \rangle^q} e^{-(c_1+1) \langle \cdot \rangle^{(q-1)q_1/(q_1-1)}} (\varphi * e^{s \langle \cdot \rangle^q}) \in \mathcal{D}_{L^\infty}^{\{M_p\}}(\mathbb{R}^d)$, which, in turn, gives

$$(\varphi * e^{s \langle \cdot \rangle^q}) S = \left(e^{-s \langle \cdot \rangle^q} e^{-(c_1+1) \langle \cdot \rangle^{(q-1)q_1/(q_1-1)}} (\varphi * e^{s \langle \cdot \rangle^q}) \right) \left(e^{s \langle \cdot \rangle^q} e^{(c_1+1) \langle \cdot \rangle^{(q-1)q_1/(q_1-1)}} S \right) \in \mathcal{D}_{L^1}^{\{M_p\}}(\mathbb{R}^d).$$

□

Remark 2.9. Notice that

$$\mathcal{S}_{\{p^{1/q_1}\}}^{\{M_p\}}(\mathbb{R}^d) = \lim_{h \rightarrow 0^+} \tilde{\mathcal{S}}_{p^{1/q_1}, h}^{M_p, h}(\mathbb{R}^d),$$

where $\tilde{\mathcal{S}}_{p^{1/q_1}, h}^{M_p, h}(\mathbb{R}^d)$ is the Banach space of all $\varphi \in \mathcal{C}^\infty(\mathbb{R}^d)$ such that

$$\sup_{\alpha, \beta \in \mathbb{N}^d} h^{|\alpha|+|\beta|} \|\langle \cdot \rangle^{|\beta|} \partial^\alpha \varphi\|_{L^\infty(\mathbb{R}^d)} / (M_\alpha \beta!^{1/q_1}) < \infty.$$

Thus, if $\{M_p\}_{p \in \mathbb{N}}$ satisfies the non-quasianalyticity condition $\sum_{p=1}^\infty M_p^{-1/p} < \infty$, one can consider the case when $q_1 = \infty$ as well; in fact $\mathcal{S}_{\{p^{1/0}\}}^{\{M_p\}}(\mathbb{R}^d)$ is exactly $\mathcal{D}^{\{M_p\}}(\mathbb{R}^d)$, i.e. the space of compactly supported ultradifferentiable functions of class $\{M_p\}$ (see [10]). In this case (and when $\{M_p\}_{p \in \mathbb{N}}$ additionally satisfies the condition (M.3) from [10]), when $q = 2$, the condition (2.7), which boils down to $e^{s \langle \cdot \rangle^2} e^{k \langle \cdot \rangle} S \in \mathcal{D}_{L^1}^{\{M_p\}}(\mathbb{R}^d)$, $\forall k \geq 0$, is known to be both necessary and sufficient for the existence of the $\mathcal{D}^{\{M_p\}}$ -convolution with the Gaussian $e^{s \langle \cdot \rangle^2}$; see [15, Theorem 4.3 (a)]. (See also [21] for the necessary and sufficient conditions for existence of convolution with the Gaussian $e^{s \langle \cdot \rangle^2}$ in the setting of Schwartz distributions.)

3. NECESSARY CONDITIONS FOR THE CONVOLUTION WITH $e^{s\langle \cdot \rangle^q}$, $q \geq 1, s > 0$

We now turn our attention to finding necessary conditions for the existence of the $\mathcal{S}'_{\{p^{1/q_1}\}}\{M_p\}$ -convolution with $e^{s\langle \cdot \rangle^q}$. We start with two technical results.

Lemma 3.1. *Let $q_1 > q \geq 1$, $s \in \mathbb{R} \setminus \{0\}$ and g be a measurable function on \mathbb{R}^d which satisfies the following: there exist $\eta_1, \eta_2, C'_1, C'_2 > 0$ such that*

$$C'_1 e^{-\eta_1 \langle x \rangle^{q_1}} \leq g(x) \leq C'_2 e^{-\eta_2 \langle x \rangle^{q_1}}, \quad a.e. \quad (3.1)$$

Then for every $k > 0$ there exists $c, \varepsilon \in (0, 1)$ such that

$$(g(\varepsilon \cdot) * e^{s\langle \cdot \rangle^q})(x) \geq c e^{s\langle x \rangle^q} e^{k\langle x \rangle^{(q-1)q_1/(q_1-1)}}, \quad \forall x \in \mathbb{R}^d.$$

Proof. Notice first that when $q = 1$ the claim immediately follows from

$$\langle x \rangle - \langle t \rangle \leq \langle x - t \rangle \leq \langle x \rangle + \langle t \rangle, \quad \text{for all } x, t \in \mathbb{R}^d.$$

So, throughout the rest of the proof we assume $q > 1$; additionally, we denote by $B(x, r)$ the closed ball with centre at $x \in \mathbb{R}^d$ and radius $r > 0$. Assume first $s > 0$ and let $k > 0$ be arbitrary but fixed. Pick $k_1 \geq 1 + 4(k+1)/(qs)$ and set $\varepsilon = 1/(4^{q_1}(\eta_1 + 1)k_1)$. Let $x \in \mathbb{R}^d$, $|x| \geq 2$, be arbitrary but fixed. For every $t \in B(-k_1|x|^{-1+(q-1)/(q_1-1)}x, 1)$, we infer

$$\begin{aligned} |x - t| &\geq |x + k_1|x|^{-1+(q-1)/(q_1-1)}x| - |k_1|x|^{-1+(q-1)/(q_1-1)}x + t| \\ &\geq |x| + k_1|x|^{(q-1)/(q_1-1)} - 1 \geq |x| + (k_1 - 1)|x|^{(q-1)/(q_1-1)}. \end{aligned}$$

We apply the equality (2.5) with $\lambda_1 = |x - t|$, $\lambda_2 = |x|$. Because of the above, $\lambda_1 > \lambda_2$ and thus $\lambda' \langle \lambda' \rangle^{q-2} \geq \langle x \rangle^{q-1}/2$. We infer

$$s\langle x - t \rangle^q \geq s\langle x \rangle^q + (qs(k_1 - 1)/2)|x|^{(q-1)/(q_1-1)}\langle x \rangle^{q-1} \geq s\langle x \rangle^q + (k+1)\langle x \rangle^{(q-1)q_1/(q_1-1)} \quad (3.2)$$

for all $t \in B(-k_1|x|^{-1+(q-1)/(q_1-1)}x, 1)$. On the other hand, with t as before, we have

$$\eta_1 \langle \varepsilon t \rangle^{q_1} \leq 2^{q_1} \eta_1 \langle \varepsilon \rangle^{q_1} + 2^{q_1} \eta_1 \langle \varepsilon k_1 |x|^{(q-1)/(q_1-1)} \rangle^{q_1} \leq 8^{q_1} \eta_1 + \langle x \rangle^{(q-1)q_1/(q_1-1)}.$$

Thus, denoting by ω_d the measure of the unit ball in \mathbb{R}^d , we deduce

$$\begin{aligned} (g(\varepsilon \cdot) * e^{s\langle \cdot \rangle^q})(x) &\geq C'_1 \int_{B(-k_1|x|^{-1+(q-1)/(q_1-1)}x, 1)} e^{-\eta_1 \langle \varepsilon t \rangle^{q_1}} e^{s\langle x-t \rangle^q} dt \\ &\geq \omega_d C'_1 e^{-8^{q_1} \eta_1} e^{s\langle x \rangle^q} e^{k\langle x \rangle^{(q-1)q_1/(q_1-1)}}. \end{aligned} \quad (3.3)$$

Hence, the above holds for all $x \in \mathbb{R}^d$, $|x| \geq 2$. As $g(\varepsilon \cdot) * e^{s\langle \cdot \rangle^q}$ is positive and continuous, this holds on all of \mathbb{R}^d as well possibly with a different constant in place of $\omega_d C'_1 e^{-8^{q_1} \eta_1}$.

Assume now $s < 0$. Let $k > 0$ be arbitrary but fixed. Pick $k_1 \geq 1 + 4^q(k+1)/(q|s|)$ and set $\varepsilon = 1/(4^{q_1}(\eta_1 + 1)k_1)$. There exists $c' \geq 4$ such that $k_1|x|^{-1+(q-1)/(q_1-1)} \leq 1/2$, for all $|x| \geq c'$. Let $x \in \mathbb{R}^d$, $|x| \geq c'$, be arbitrary but fixed. For $t \in B(k_1|x|^{-1+(q-1)/(q_1-1)}x, 1)$ we infer

$$|x - t| \leq |x|(1 - k_1|x|^{-1+(q-1)/(q_1-1)}) + 1 = |x| - k_1|x|^{(q-1)/(q_1-1)} + 1,$$

and thus $|x| \geq |x - t| + (k_1 - 1)|x|^{(q-1)/(q_1-1)}$. Furthermore

$$|x - t| \geq |x|(1 - k_1|x|^{-1+(q-1)/(q_1-1)}) - 1 \geq |x|/2 - 1 \geq |x|/4$$

(the very last inequality follows from $|x| \geq c' \geq 4$). We apply (2.5) with $\lambda_1 = |x - t|$ and $\lambda_2 = |x|$. Because of the above, $\lambda' \langle \lambda' \rangle^{q-2} \geq \langle x - t \rangle^{q-1}/2$ and consequently

$$\begin{aligned} s \langle x - t \rangle^q - s \langle x \rangle^q &= q|s|(|x| - |x - t|)\lambda' \langle \lambda' \rangle^{q-2} \geq q|s|((k_1 - 1)/2)|x|^{(q-1)/(q_1-1)} \langle x/4 \rangle^{q-1} \\ &\geq (k + 1) \langle x \rangle^{(q-1)q_1/(q_1-1)}, \end{aligned}$$

for all $t \in B(k_1|x|^{-1+(q-1)/(q_1-1)}x, 1)$. Now, similarly as above, one deduces the validity of (3.3) for all $|x| \geq c'$. As $g(\varepsilon \cdot) * e^{s \langle \cdot \rangle^q}$ is positive and continuous this implies the claim in the lemma. \square

Lemma 3.2. *Let $q_1 > q \geq 1$, $s > 0$ and g be a measurable function on \mathbb{R}^d which satisfies the following:*

$$\text{there exists } C', \eta' > 0 \text{ such that } 0 \leq g(x) \leq C' e^{-\eta' \langle x \rangle^{q_1}}, \quad a.e.$$

$$\text{measure}(\{x \in \mathbb{R}^d \mid g(x) > 0\}) > 0.$$

(i) *There exists $C > 1$ such that*

$$|D^\alpha(g * e^{s \langle \cdot \rangle^q})(x)| \leq C^{|\alpha|+1} \alpha! \langle x \rangle^{(q-1)|\alpha|} (g * e^{s \langle \cdot \rangle^q})(x), \quad \text{for all } x \in \mathbb{R}^d, \alpha \in \mathbb{N}^d.$$

(ii) *There exists $C > 1$ such that*

$$|D^\alpha(1/(g * e^{s \langle \cdot \rangle^q})(x))| \leq C^{|\alpha|+1} \alpha! \langle x \rangle^{(q-1)|\alpha|} / (g * e^{s \langle \cdot \rangle^q})(x), \quad \text{for all } x \in \mathbb{R}^d, \alpha \in \mathbb{N}^d.$$

Proof. The condition on g , as well as the fact that $s > 0$, imply

$$(g * e^{s \langle \cdot \rangle^q})(x) = \int_{\mathbb{R}^d} g(t) e^{s \langle x-t \rangle^q} dt \geq \|g\|_{L^1(\mathbb{R}^d)} > 0, \quad \forall x \in \mathbb{R}^d.$$

To prove (i) we apply Lemma 2.3 with $\alpha \in \mathbb{N}^d \setminus \{0\}$ to infer

$$|D^\alpha(g * e^{s \langle \cdot \rangle^q})(x)| \leq C^{|\alpha|} \alpha! (I_1 + I_2), \quad \text{where}$$

$$I_1 = \int_{|t| \leq |x|} g(t) e^{s \langle x-t \rangle^q} \sum_{m=1}^{|\alpha|} \frac{\langle x-t \rangle^{qm-|\alpha|}}{m!} dt, \quad I_2 = \int_{|t| \geq |x|} g(t) e^{s \langle x-t \rangle^q} \sum_{m=1}^{|\alpha|} \frac{\langle x-t \rangle^{qm-|\alpha|}}{m!} dt.$$

When $|t| \leq |x|$,

$$\langle x-t \rangle^{qm-|\alpha|} \leq \langle x-t \rangle^{(q-1)|\alpha|} \leq 2^{(q-1)|\alpha|} \langle x \rangle^{(q-1)|\alpha|},$$

whence $I_1 \leq e 2^{(q-1)|\alpha|} \langle x \rangle^{(q-1)|\alpha|} (g * e^{s \langle \cdot \rangle^q})(x)$. To estimate I_2 , we proceed as follows

$$I_2 \leq \int_{|t| \geq |x|} g(t) e^{s \langle x-t \rangle^q} e^{\langle x-t \rangle^{q-1}} dt \leq \int_{|t| \geq |x|} g(t) e^{s \langle 2t \rangle^q} e^{\langle 2t \rangle^{q-1}} dt \leq \|g\| e^{s 2^q \langle \cdot \rangle^q} e^{2^{q-1} \langle \cdot \rangle^{q-1}} \|1\|_{L^1(\mathbb{R}^d)}.$$

Now, (i) follows from the fact that $g * e^{s \langle \cdot \rangle^q}$ is bounded from below by a positive constant.

To prove (ii) we apply the Faà di Bruno formula (Proposition 2.1) to the composition of the functions $\rho \mapsto 1/\rho$ with $x \mapsto (g * e^{s \langle \cdot \rangle^q})(x)$. The first part of the lemma gives

$$|D^\alpha(1/(g * e^{s \langle \cdot \rangle^q})(x))| \leq \alpha! \sum_{m=1}^{|\alpha|} \frac{C^{|\alpha|+m} m! \langle x \rangle^{(q-1)|\alpha|}}{(g * e^{s \langle \cdot \rangle^q})(x)} \sum_{p(\alpha, m)} \prod_{j=1}^{|\alpha|} \frac{1}{k_j!}.$$

Now, the bound in (ii) follows from (2.3). \square

Remark 3.3. Under the assumptions of the above lemma, by applying Corollary 2.4, it is straightforward to verify that

$$|D^\alpha(g * e^{s\langle \cdot \rangle^q})(x)| \leq C^{|\alpha|+1} \alpha! e^{s'\langle x \rangle^q}, \quad \text{for all } x \in \mathbb{R}^d, \alpha \in \mathbb{N}^d,$$

for some $C > 1, s' > s$; i.e. $g * e^{s\langle \cdot \rangle^q}$ is a multiplier for $\mathcal{S}'_{\{p^{1/q_1}\}}(\mathbb{R}^d)$.

Now we are ready to prove the main results of this section; recall that $p!^\kappa \subset M_p$ means: there exist $C, L > 0$ such that $p!^\kappa \leq CL^p M_p, \forall p \in \mathbb{N}$.

Theorem 3.4. *Let $q_1 > q \geq 1, s > 0$ and g a measurable function on \mathbb{R}^d which satisfies (3.1).*

- (i) *Assume $p!^{2-1/q} \subset M_p$. If $S \in \mathcal{S}'_{\{p^{1/q_1}\}}(\mathbb{R}^d)$ satisfies $(g(\varepsilon \cdot) * e^{s\langle \cdot \rangle^q})S \in \mathcal{D}'_{L^1}(\mathbb{R}^d)$, $\forall \varepsilon \in (0, 1)$, then*

$$e^{s'\langle \cdot \rangle^q} S \in \mathcal{D}'_{L^1}(\mathbb{R}^d), \quad \text{for all } s' < s. \quad (3.4)$$

If $q = 1$, then (3.4) holds true even for $s' = s$.

- (ii) *Assume $p!^{2-1/q_1} \subset M_p$. If $S \in \mathcal{S}'_{\{p^{1/q_1}\}}(\mathbb{R}^d)$ satisfies $(g(\varepsilon \cdot) * e^{s\langle \cdot \rangle^q})S \in \mathcal{D}'_{L^1}(\mathbb{R}^d)$, $\forall \varepsilon \in (0, 1)$, then*

$$e^{s\langle \cdot \rangle^q} e^{k\langle \cdot \rangle^{(q-1)q_1/(q_1-1)}} S \in \mathcal{D}'_{L^1}(\mathbb{R}^d), \quad \text{for all } k \geq 0. \quad (3.5)$$

Proof. First we prove (ii). Let $k \geq 0$ be arbitrary but fixed. By Lemma 3.1 there exists $c, \varepsilon \in (0, 1)$ such that

$$(g(\varepsilon \cdot) * e^{s\langle \cdot \rangle^q})(x) \geq c e^{s\langle x \rangle^q} e^{(k+2)\langle x \rangle^{(q-1)q_1/(q_1-1)}}, \quad \forall x \in \mathbb{R}^d. \quad (3.6)$$

Lemma 3.2 (ii) implies

$$|D^\alpha(1/(g(\varepsilon \cdot) * e^{s\langle \cdot \rangle^q})(x))| \leq C_1 C^{|\alpha|} \alpha! \langle x \rangle^{(q-1)|\alpha|} e^{-s\langle x \rangle^q} e^{-(k+2)\langle x \rangle^{(q-1)q_1/(q_1-1)}}. \quad (3.7)$$

Since $t^N/N! \leq e^t$ for all $N \in \mathbb{N}$ and $t \geq 0$, we deduce

$$\langle x \rangle^{(q-1)|\alpha|} = |\alpha|!^{1-1/q_1} \left(\frac{\langle x \rangle^{|\alpha|(q-1)q_1/(q_1-1)}}{|\alpha|!} \right)^{1-1/q_1} \leq |\alpha|!^{1-1/q_1} e^{(1-1/q_1)\langle x \rangle^{(q-1)q_1/(q_1-1)}}$$

and then

$$|D^\alpha(1/(g(\varepsilon \cdot) * e^{s\langle \cdot \rangle^q})(x))| \leq C_1 C^{|\alpha|} |\alpha|!^{2-1/q_1} e^{-s\langle x \rangle^q} e^{-(k+1)\langle x \rangle^{(q-1)q_1/(q_1-1)}}.$$

Now, Corollary 2.4 yields $e^{s\langle \cdot \rangle^q} e^{k\langle \cdot \rangle^{(q-1)q_1/(q_1-1)}}/g(\varepsilon \cdot) * e^{s\langle \cdot \rangle^q} \in \mathcal{D}'_{L^\infty}(\mathbb{R}^d)$ and consequently

$$e^{s\langle \cdot \rangle^q} e^{k\langle \cdot \rangle^{(q-1)q_1/(q_1-1)}} S = \frac{e^{s\langle \cdot \rangle^q} e^{k\langle \cdot \rangle^{(q-1)q_1/(q_1-1)}}}{g(\varepsilon \cdot) * e^{s\langle \cdot \rangle^q}} ((g(\varepsilon \cdot) * e^{s\langle \cdot \rangle^q})S) \in \mathcal{D}'_{L^1}(\mathbb{R}^d).$$

To prove (i), assume first $q > 1$. Let $s' < s$ be arbitrary but fixed and denote $\kappa = s - s' > 0$. Pick $\varepsilon > 0$ such that (3.6) holds true with $k = 0$. By employing the bound

$$\langle x \rangle^{(q-1)|\alpha|} \leq \kappa^{-|\alpha|(1-1/q)} |\alpha|!^{1-1/q} e^{(1-1/q)\kappa \langle x \rangle^q},$$

similarly as above we deduce

$$|D^\alpha(1/(g(\varepsilon \cdot) * e^{s(\cdot)^q})(x))| \leq C_2 C^{|\alpha|} |\alpha|!^{2-1/q} e^{-s' \langle x \rangle^q} e^{-2 \langle x \rangle^{(q-1)q_1/(q_1-1)}}.$$

Now, one can apply the same technique as before to conclude the validity of (i). When $q = 1$, notice that (3.7) boils down to $|D^\alpha(1/(g(\varepsilon \cdot) * e^{s(\cdot)})(x))| \leq C_1 C^{|\alpha|} |\alpha|! e^{-s \langle x \rangle}$, which, in view of Corollary 2.4, immediately implies $e^{s(\cdot)}/g(\varepsilon \cdot) * e^{s(\cdot)} \in \mathcal{D}_{L^\infty}^{\{p\}}(\mathbb{R}^d)$; the rest of the proof is the same as before. \square

Applying the above theorem for $g = e^{-\langle \cdot \rangle^{q_1}}$ we immediately deduce the following result.

Corollary 3.5. *Let $q_1 > q \geq 1$ and $s > 0$.*

- (i) *Assume $p!^{2-1/q} \subset M_p$. If the $\mathcal{S}'_{\{p!^{1/q_1}\}}^{\{M_p\}}$ -convolution of $S \in \mathcal{S}'_{\{p!^{1/q_1}\}}^{\{M_p\}}(\mathbb{R}^d)$ and $e^{s(\cdot)^q}$ exists then (3.4) holds true. When $q = 1$, (3.4) holds true even for $s' = s$.*
- (ii) *Assume $p!^{2-1/q_1} \subset M_p$. If the $\mathcal{S}'_{\{p!^{1/q_1}\}}^{\{M_p\}}$ -convolution of $S \in \mathcal{S}'_{\{p!^{1/q_1}\}}^{\{M_p\}}(\mathbb{R}^d)$ and $e^{s(\cdot)^q}$ exists then (3.5) holds true.*

As a consequence of the above result together with Theorem 2.6 we have the following corollary.

Corollary 3.6. *Let $q_1 > 1$, $s > 0$ and $S \in \mathcal{S}'_{\{p!^{1/q_1}\}}^{\{M_p\}}(\mathbb{R}^d)$. The $\mathcal{S}'_{\{p!^{1/q_1}\}}^{\{M_p\}}$ -convolution of S and $e^{s(\cdot)}$ exists if and only if $e^{s(\cdot)} S \in \mathcal{D}_{L^1}^{\{M_p\}}(\mathbb{R}^d)$.*

Since $(q-1)q_1/(q_1-1) < q$, Corollary 3.5 together with Theorem 2.6 also imply the following result.

Corollary 3.7. *Let $q_1 > q \geq 1$, $s > 0$, $p!^{2-1/q} \subset M_p$ and $S \in \mathcal{S}'_{\{p!^{1/q_1}\}}^{\{M_p\}}(\mathbb{R}^d)$. If the $\mathcal{S}'_{\{p!^{1/q_1}\}}^{\{M_p\}}$ -convolution of S and $e^{s(\cdot)^q}$ exists then the $\mathcal{S}'_{\{p!^{1/q_1}\}}^{\{M_p\}}$ -convolution of S and $e^{s'(\cdot)^q}$ also exists for all $s' < s$, $s' \neq 0$.*

REFERENCES

- [1] C. Bargetz, N. Ortner, *Convolution of vector-valued distributions: a survey and comparison*, Diss. Math, 495 (2013) 1-51.
- [2] R. Carmichael, A. Kamiński, S. Pilipović, *Boundary Values and Convolution in Ultradistribution Spaces*, World Scientific Publishing Co. Pte. Ltd., 2007.
- [3] G. M. Constantine, T. H. Savits, *A multivariate Faà di Bruno formula with applications*, Transactions of the American Mathematical Society, **348** (1996), 503–520.
- [4] A. Debrouwere, J. Vindas, *On weighted inductive limits of spaces of ultradifferentiable functions and their duals*, Math. Nachr, 292(3) (2019), 573-602
- [5] P. Dierolf, J. Voigt, *Convolution and \mathcal{S}' -convolution of distributions*, Collect. Math. **29** (1978) 185–196.
- [6] P. Dimovski, S. Pilipović, B. Prangoski, J. Vindas, *Convolution of ultradistributions and ultradistribution spaces associated to translation-invariant Banach spaces*, Kyoto J. Math, **56** (2016), 401-440.
- [7] P. Dimovski, B. Prangoski, J. Vindas, *On a class of translation-invariant spaces of quasianalytic ultradistributions*, Novi Sad J. Math, **45** (2015), 143-175.
- [8] I. M. Gelfand, G. E. Shilov, *Generalized Functions, Volume 2: Spaces of Fundamental and Generalized Functions*, Acad. Press, New York, 1968.

- [9] A. Kamiński, D. Kovačević, S. Pilipović, *The equivalence of various definitions of the convolution of ultradistributions*, Trudy Mat. Inst. Steklov, 203 (1994), 307-322.
- [10] H. Komatsu, *Ultradistributions, I: Structure theorems and a characterization*, J. Fac. Sci. Univ. Tokyo, Sect. IA Math, 20 (1973), 25-105.
- [11] F. Nicola, L. Rodino, *Global Pseudo-Differential Calculus on Euclidean Spaces*, Vol. 4. Birkhäuser Basel, 2010.
- [12] N. Ortner, *On convolvability conditions for distributions*, Monatsh. Math, 160 (2010) 313-335.
- [13] N. Ortner, P. Wagner, *Distribution-Valued Analytic Functions - Theory and Applications*, edition swk, Hamburg, 2013.
- [14] S. Pilipović, B. Prangoski, *On the convolution of Roumieu ultradistributions through the ϵ tensor product*, Monatsh. Math, 173 (2014), 83-105.
- [15] S. Pilipović, B. Prangoski, *Anti-Wick and Weyl quantization on ultradistribution spaces*, J. Math. Pures Appl, 103 (2015), 472-503.
- [16] S. Pilipović, B. Prangoski, *Complex powers for a class of infinite order hypoelliptic operators*, Dissert. Math, 529 (2018), 1-58.
- [17] S. Pilipović, B. Prangoski, J. Vindas, *On quasianalytic classes of Gelfand Shilov type. Parametrix and convolution*, J. Math. Pures Appl, 116 (2018), 174-210.
- [18] H. H. Schaefer, *Topological vector spaces*, Springer-Verlag, New York Heidelberg Berlin, 1970.
- [19] L. Schwartz, *Théorie des distributions à valeurs vectorielles. I*, Ann. Inst. Fourier, 7 (1957), 1-141.
- [20] R. Shiraishi, *On the definition of convolution for distributions*, J. Sci. Hiroshima Univ. Ser. A 23 (1959) 19-32.
- [21] P. Wagner, *Zur Faltung von Distributionen*, Mathematische Annalen, **276**(3) (1987), 467-485

STEVAN PILIPOVIĆ, DEPARTMENT OF MATHEMATICS AND INFORMATICS, UNIVERSITY OF NOVI SAD, TRG DOSITEJA OBRADOVIĆA 4, 21000 NOVI SAD, SERBIA
Email address: `stevan.pilipovic@dmi.uns.ac.rs`

BOJAN PRANGOSKI, FACULTY OF MECHANICAL ENGINEERING, SS. CYRIL AND METHODIUS UNIVERSITY IN SKOPJE, KARPOS II BB, 1000 SKOPJE, MACEDONIA
Email address: `bprangoski@yahoo.com`

DORĐE VUČKOVIĆ, MATHEMATICAL INSTITUTE OF THE SERBIAN ACADEMY OF SCIENCES AND ARTS, KNEZ MIHAILOVA 36, 11000 BELGRADE, SERBIA
Email address: `djordjeplusja@gmail.com`