A NOTE ON EXTRAPOLATION OF COMPACTNESS

SHENYU LIU, HUOXIONG WU AND DONGYONG YANG

ABSTRACT. This note is devoted to the study of Hytönen's extrapolation theorem of compactness on weighted Lebesgue spaces. Two criteria of compactness of linear operators in the two-weight setting are obtained. As applications, we obtain two-weight compactness of commutators of Calderón–Zygmund operators, fractional integrals and bilinear Calderón–Zygmund operators.

1. INTRODUCTION

By a weight on \mathbb{R}^d we mean a locally integrable function $w \in L^1_{loc}(\mathbb{R}^d)$ that is positive almost everywhere. For $p \in (1, \infty)$, we define the weighted Lebesgue spaces

$$L^{p}(w) := \left\{ f : \mathbb{R}^{d} \to \mathbb{C} \text{ measurable } \middle| \|f\|_{L^{p}(w)} := \left(\int_{\mathbb{R}^{d}} |f(x)|^{p} w(x) dx \right)^{\frac{1}{p}} < \infty \right\}.$$

A weight w is called a Muckenhoupt A_p weight (write $w \in A_p$) if

$$[w]_{A_p} := \sup_{Q \subset \mathbb{R}^d} \langle w \rangle_Q \langle w^{1-p'} \rangle_Q^{p-1} < \infty_q$$

where the supremum is taken over all cubes $Q \subset \mathbb{R}^d$, $\langle w \rangle_Q := |Q|^{-1} \int_Q w$ and $p' := \frac{p}{p-1}$. We define $A_{\infty} := \bigcup_{p>1} A_p$ and

$$[w]_{A_{\infty}} := \sup_{Q \subset \mathbb{R}^d} \frac{1}{w(Q)} \int_Q M(w\chi_Q)(x) dx < \infty,$$

where M denotes the Hardy–Littlewood maximal operator. A weight w is called an $A_{p,q}$ weight (write $w \in A_{p,q}$) if

$$w]_{A_{p,q}} := \sup_{Q \subset \mathbb{R}^d} \langle w^q \rangle_Q^{\frac{1}{q}} \langle w^{-p'} \rangle_Q^{\frac{1}{p'}} < \infty, \quad 1 < p \le q < \infty.$$

It readily follows that

$$w \in A_{p,q} \iff w^q \in A_{1+q/p'} \subset A_q \iff w^{-p'} \in A_{1+p'/q} \subset A_{p'} \Rightarrow w^p \in A_p.$$

In this note, we show that the corresponding results of Hytönen [12], Hytönen and Lappas [13] are still valid in the two-weight setting. To be precise, we establish the following two extrapolation theorems of compactness:

Theorem 1.1. Let T be a linear operator simultaneously defined and bounded from $L^p(\sigma)$ to $L^p(\lambda)$ for all $p \in (1, \infty)$ and all $\sigma, \lambda \in A_p$. Suppose moreover that T is compact on $L^q(w)$ for some $q \in (1, \infty)$ and some $w \in A_q$. Then T is compact from $L^p(\sigma)$ to $L^p(\lambda)$ for all $p \in (1, \infty)$ and all $\sigma, \lambda \in A_p$.

Date: January 9, 2022.

²⁰²⁰ Mathematics Subject Classification. 47B38, 46B70, 42B35, 42B20.

Key words and phrases. Weighted extrapolation, compact operator, two-weight, commutator, Calderón-Zygmund operator, fractional integral, bilinear operator.

Theorem 1.1 says that one-weight compactness (or even unweighted compactness when we take $w \equiv 1$) bootstraps to two-weight compactness, if the two-weight boundedness is already known. We also consider the counterpart off-diagonal case in the two-weight setting as follows:

Theorem 1.2. Let T be a linear operator simultaneously defined and compact from $L^{p_1}(w_1^{p_1})$ to $L^{q_1}(w_1^{q_1})$ for some $1 < p_1 \le q_1 < \infty$ and some $w_1 \in A_{p_1,q_1}$. Suppose moreover that T is simultaneously defined and bounded from $L^p(\sigma^p)$ to $L^q(\lambda^q)$ for all $p, q \in (1, \infty)$ with $\frac{1}{p} - \frac{1}{q} = \frac{1}{p_1} - \frac{1}{q_1}$ and all $\sigma, \lambda \in A_{p,q}$. Then T is compact from $L^p(\sigma^p)$ to $L^q(\lambda^q)$ for all $p, q \in (1, \infty)$ all $p, q \in (1, \infty)$ with $\frac{1}{p} - \frac{1}{q} = \frac{1}{p_1} - \frac{1}{q_1} = \frac{1}{p_1} - \frac{1}{q_1}$ and all $\sigma, \lambda \in A_{p,q}$.

The rest of this note is organized as follows. In Section 2, we recover Hytönen's extrapolation theorem by extending the extrapolation theorem of the A_p weights in [12]. In Section 3, we obtain some generalizations of extrapolation of compactness in the two-weight setting. Section 4 is devoted to providing two applications to the two-weight compactness of linear operators, including commutators of Calderón–Zygmund operators and fractional integrals. We also give an application to the two-weight compactness of bilinear Calderón–Zygmund operators in Section 5.

Throughout this note, we write $A \leq B$ to denote $A \leq cB$ for some positive constant c which is independent of the main parameters. We write $A \approx B$ if $A \leq B$ and $B \leq A$.

2. Hytönen's extrapolation theorem: a revisit

Hytönen [12] first established a compact version of Rubio de Francia's weighted extrapolation theorem (see Theorem 2.6 below), whose approach relies on interpolation of compactness, see also [13, 14, 4]. In this section, we recover Hytönen's extrapolation theorem by applying an abstract tool about extrapolation of compact operators due to Cwikel [7, Theorem 2.1], in which there is no additional assumption on Banach couples in comparison to [12, Theorem 3.1] (see also Theorem 3.1 below).

Theorem 2.1 ([7]). Let (X_0, X_1) and (Y_0, Y_1) be Banach couples and T be a linear operator such that $T : X_0 + X_1 \to Y_0 + Y_1$ and $T : X_j \to Y_j$ boundedly for j = 0, 1. Let $[,]_{\theta}$ be the complex interpolation functor of Calderón. Suppose moreover that $T : [X_0, X_1]_{\theta} \to [Y_0, Y_1]_{\theta}$ is compact for at least one value θ_* of θ in (0, 1). Then $T : [X_0, X_1]_{\theta} \to [Y_0, Y_1]_{\theta}$ is compact for all values of θ in (0, 1).

To apply Theorem 2.1, we need a well-known result of Stein and Weiss [20] as follows, see also [2, Theorem 5.5.3].

Theorem 2.2 ([20, 2]). Let $q, t \in [1, \infty)$ and w_0, w_1 be two weights. Then for all $\theta \in (0, 1)$,

$$L^{p}(w) = [L^{t}(w_{0}), L^{q}(w_{1})]_{\theta},$$

where

(2.1)
$$\frac{1}{p} = \frac{1-\theta}{t} + \frac{\theta}{q}, \quad w^{\frac{1}{p}} = w_0^{\frac{1-\theta}{t}} w_1^{\frac{\theta}{q}}.$$

We also need the following powerful result of Hytönen [12, Lemma 4.3] that can be understood as an extrapolation of the A_p weights.

Theorem 2.3 ([12]). Let $p, q \in (1, \infty)$, $w \in A_p$ and $w_1 \in A_q$. Then there exist $\theta \in (0, 1)$, $t \in (1, \infty)$ and $w_0 \in A_t$ such that (2.1) holds.

To adapt Theorem 2.2 to our case, we need a further result described below:

Lemma 2.4. Let $p, q \in (1, \infty)$, $u \in A_p$ and $v \in A_q$. Then there exist $\theta_0, \theta_1 \in (0, 1)$, $r, s \in (1, \infty)$, $w_0 \in A_r$ and $w_1 \in A_s$ such that

(2.2)
$$\frac{1}{p} = \frac{1-\theta_0}{r} + \frac{\theta_0}{s}, \quad u^{\frac{1}{p}} = w_0^{\frac{1-\theta_0}{r}} w_1^{\frac{\theta_0}{s}}$$

(2.3)
$$\frac{1}{q} = \frac{1-\theta_1}{r} + \frac{\theta_1}{s}, \quad v^{\frac{1}{q}} = w_0^{\frac{1-\theta_1}{r}} w_1^{\frac{\theta_1}{s}}$$

Proof. For $v \in A_q$ and $u \in A_p$, from Theorem 2.3, there exist $\tilde{\theta}_0 \in (0,1)$, $r \in (1,\infty)$ and $w_0 \in A_r$ such that

(2.4)
$$\frac{1}{q} = \frac{1 - \hat{\theta}_0}{r} + \frac{\hat{\theta}_0}{p}, \quad v^{\frac{1}{q}} = w_0^{\frac{1 - \tilde{\theta}_0}{r}} u^{\frac{\tilde{\theta}_0}{p}}.$$

Then for $u \in A_p$ and $w_0 \in A_r$, again by Theorem 2.3, there exist $\theta_0 \in (0, 1)$, $s \in (1, \infty)$ and $w_1 \in A_s$ such that (2.2) holds. Combined with (2.2) and (2.4), we obtain (2.3) by taking $\theta_1 = \theta_0 \tilde{\theta}_0$ which completes the proof.

By combining with Theorem 2.2 and Lemma 2.4, we immediately obtain the following interpolation result.

Corollary 2.5. Let $\lambda \in [1, \infty)$, $p, q \in (\lambda, \infty)$, $u \in A_{p/\lambda}$ and $v \in A_{q/\lambda}$. Then

$$L^{p}(u) = [L^{r}(w_{0}), L^{s}(w_{1})]_{\theta_{0}}, \quad L^{q}(v) = [L^{r}(w_{0}), L^{s}(w_{1})]_{\theta_{1}}$$

for some $\theta_0, \theta_1 \in (0, 1), r, s \in (\lambda, \infty), w_0 \in A_{r/\lambda}$ and $w_1 \in A_{s/\lambda}$.

Now we are ready to recover Hytönen's extrapolation theorem in [12, Theorem 1.2]; see also [21] for the weighted compactness of Calderón-Zygmund operators.

Theorem 2.6 ([12]). Let $\lambda \in [1, \infty)$, $p_1 \in (\lambda, \infty)$ and T be a linear operator simultaneously defined and bounded on $L^{p_1}(\tilde{w})$ for all $\tilde{w} \in A_{p_1/\lambda}$, with the operator norm dominated by some increasing function of $[\tilde{w}]_{A_{p_1/\lambda}}$. Suppose in addition that T is compact on $L^{p_1}(w_1)$ for some $w_1 \in A_{p_1/\lambda}$. Then T is compact on $L^p(w)$ for all $p \in (\lambda, \infty)$ and all $w \in A_{p/\lambda}$.

Proof. First, by a rescaling version of the Rubio de Francia weighted extrapolation theorem (see, for example, [12, Theorem 1.1]), we have that T is a bounded linear operator on $L^p(w)$ for all $p \in (\lambda, \infty)$ and all $w \in A_{p/\lambda}$. Next, for any fixed $p \in (\lambda, \infty)$ and $w \in A_{p/\lambda}$, since T is compact on $L^{p_1}(w_1)$ for $p_1 \in (\lambda, \infty)$ and some $w_1 \in A_{p_1/\lambda}$, by Corollary 2.5,

$$L^{p}(w) = [L^{r}(w_{0}), L^{s}(w_{2})]_{\theta_{0}}, \quad L^{p_{1}}(w_{1}) = [L^{r}(w_{0}), L^{s}(w_{2})]_{\theta_{1}}$$

for some $\theta_0, \theta_1 \in (0, 1), r, s \in (\lambda, \infty), w_0 \in A_{r/\lambda}$ and $w_2 \in A_{s/\lambda}$. Finally, by Theorem 2.1 with $X_0 = Y_0 = L^r(w_0)$ and $X_1 = Y_1 = L^s(w_2)$, we obtain that T is compact on $L^p(w)$. This completes the proof.

Note that Theorem 2.6 only deals with linear operators and the classical Muckenhoupt A_p weight class. It would be interesting to extend it to the case of sublinear operators (e.g., see [25]), or more general weight classes (e.g., see [21]).

3. EXTENSIONS IN THE TWO-WEIGHT SETTING

In this section, we present the proofs of Theorems 1.1 and 1.2. Our abstract tool is the following complex interpolation theorem of compactness due to Cwikel and Kalton [8].

Theorem 3.1 ([8]). Let (X_0, X_1) and (Y_0, Y_1) be Banach couples and T be a linear operator such that $T : X_0 + X_1 \to Y_0 + Y_1$ and $T : X_j \to Y_j$ boundedly for j = 0, 1. Suppose moreover that $T : X_1 \to Y_1$ is compact. Then $T : [X_0, X_1]_{\theta} \to [Y_0, Y_1]_{\theta}$ is compact for $\theta \in (0, 1)$ under any of the following four side conditions:

- (a) X_1 has the UMD (unconditional martingale differences) property;
- (b) X_1 is reflexive, and $X_1 = [X_0, E]_{\alpha}$ for some Banach space E and $\alpha \in (0, 1)$;
- (c) $Y_1 = [Y_0, F]_\beta$ for some Banach space F and $\beta \in (0, 1)$;
- (d) X_0 and X_1 are both complexified Banach lattices of measurable functions on a common measure space.

We have swapped the roles of the indices 0 and 1 in comparison to [8]. It is known that all four side conditions hold for the weighted Lebesgue spaces, see [12, Corollary 3.3]. To obtain our extrapolation theorems of two-weight compactness, we need the following key result which is essentially an extension of the reverse Hölder inequality.

Lemma 3.2. Let $p, q \in (1, \infty)$, $\sigma, \lambda \in A_p$ and $w \in A_q$. Then there exist $\theta \in (0, 1)$, $t \in (1, \infty)$ and $\sigma_0, \lambda_0 \in A_t$ such that

(3.1)
$$\frac{1}{p} = \frac{1-\theta}{t} + \frac{\theta}{q}, \quad \sigma^{\frac{1}{p}} = \sigma_0^{\frac{1-\theta}{t}} w^{\frac{\theta}{q}}, \quad \lambda^{\frac{1}{p}} = \lambda_0^{\frac{1-\theta}{t}} w^{\frac{\theta}{q}}.$$

We recall that a weight w satisfies the reverse Hölder inequality for some $\eta \in (1, \infty)$ (write $w \in RH_{\eta}$) if

$$[w]_{RH_{\eta}} := \sup_{Q \subset \mathbb{R}^d} \langle w^{\eta} \rangle_Q^{\frac{1}{\eta}} \langle w \rangle_Q^{-1} < \infty.$$

Notice that the reverse Hölder inequality can simultaneously hold for a finite number of the Muckenhoupt weights. From this fact, one may see that the proof of Lemma 3.2 below is similar to that in [12, Lemma 4.3] with some minor changes.

Proof of Lemma 3.2. By a simple calculation of (3.1), we have that

$$t = t(\theta) = \frac{1-\theta}{\frac{1}{p} - \frac{\theta}{q}}, \quad \sigma_0 = \sigma_0(\theta) = \sigma^{\frac{t}{p(1-\theta)}} w^{-\frac{t\theta}{q(1-\theta)}}, \quad \lambda_0 = \lambda_0(\theta) = \lambda^{\frac{t}{p(1-\theta)}} w^{-\frac{t\theta}{q(1-\theta)}},$$

from which it suffices to show that there exists $\theta \in (0,1)$ such that $t \in (1,\infty)$ and $\sigma_0, \lambda_0 \in A_t$. Note that $t(0) = p \in (1,\infty)$. Then by continuity, we have $t \in (1,\infty)$ for small enough $\theta > 0$.

We now show that $\sigma_0, \lambda_0 \in A_t$. Observe that the weights σ and λ are independent of each other, so we deal with these two weights separately in the same way as [12, Lemma 4.3]. Let

$$\epsilon := \frac{\theta p}{q'}, \quad \delta := \frac{\theta p'}{q}, \quad r(\theta) := \frac{t(1+\epsilon)}{p(1-\theta)} = \frac{t(q'+\theta p)}{pq'(1-\theta)}, \quad s(\theta) := \frac{t'(1+\delta)}{p'(1-\theta)} = \frac{t'(q+\theta p')}{p'q(1-\theta)}.$$

A direct calculation yields that for any cube $Q \subset \mathbb{R}^d$,

$$(3.2) \quad \langle \sigma_0 \rangle_Q \langle \sigma_0^{1-t'} \rangle_Q^{t-1} = \langle \sigma^{\frac{t}{p(1-\theta)}} w^{-\frac{t\theta}{q(1-\theta)}} \rangle_Q \langle \sigma^{-\frac{t'}{p(1-\theta)}} w^{\frac{t'\theta}{q(1-\theta)}} \rangle_Q^{t-1} \\ \leq \langle \sigma^{\frac{t(1+\epsilon)}{p(1-\theta)}} \rangle_Q^{\frac{1}{1+\epsilon}} \langle w^{-\frac{t\theta}{q(1-\theta)}} \frac{q'+\theta p}{\theta p} \rangle_Q^{\frac{\epsilon}{1+\epsilon}} \langle \sigma^{-\frac{t'(1+\delta)}{p(1-\theta)}} \rangle_Q^{\frac{t-1}{1+\delta}} \langle w^{\frac{t'\theta}{q(1-\theta)}} \frac{q+\theta p'}{\theta p'} \rangle_Q^{\frac{\delta(t-1)}{1+\delta}}$$

$$= \langle \sigma^{r(\theta)} \rangle_Q^{\frac{1}{1+\epsilon}} \langle w^{(1-q')\frac{t(q'+\theta p)}{pq'(1-\theta)}} \rangle_Q^{\frac{\epsilon}{1+\epsilon}} \langle \sigma^{(1-p')\frac{t'(1+\delta)}{p'(1-\theta)}} \rangle_Q^{\frac{t-1}{1+\delta}} \langle w^{\frac{t'(q+\theta p')}{p'q(1-\theta)}} \rangle_Q^{\frac{\delta(t-1)}{1+\delta}} \\ = \langle \sigma^{r(\theta)} \rangle_Q^{\frac{1}{1+\epsilon}} \langle (w^{1-q'})^{r(\theta)} \rangle_Q^{\frac{\epsilon}{1+\epsilon}} \langle (\sigma^{1-p'})^{s(\theta)} \rangle_Q^{\frac{t-1}{1+\delta}} \langle w^{s(\theta)} \rangle_Q^{\frac{\delta(t-1)}{1+\delta}},$$

and similarly,

$$(3.3) \qquad \langle \lambda_0 \rangle_Q \langle \lambda_0^{1-t'} \rangle_Q^{t-1} \le \langle \lambda^{r(\theta)} \rangle_Q^{\frac{1}{1+\epsilon}} \langle (w^{1-q'})^{r(\theta)} \rangle_Q^{\frac{\epsilon}{1+\epsilon}} \langle (\lambda^{1-p'})^{s(\theta)} \rangle_Q^{\frac{t-1}{1+\delta}} \langle w^{s(\theta)} \rangle_Q^{\frac{\delta(t-1)}{1+\delta}}.$$

Observe that r(0) = s(0) = 1. Another application of continuity yields that for any fixed $\eta > 1$, $\max(r(\theta), s(\theta)) \leq \eta$ for small enough $\theta > 0$. Moreover, since $w \in A_q$, $w^{1-q'} \in A_{q'}, \sigma, \lambda \in A_p$ and $\sigma^{1-p'}, \lambda^{1-p'} \in A_{p'}$, then there exists $\eta > 1$ such that each of the above weights satisfies the reverse Hölder inequality. Thus, for small enough $\theta > 0$, using the reverse Hölder inequality, it follows from (3.2) and (3.3) that

$$\langle \sigma_0 \rangle_Q \langle \sigma_0^{1-t'} \rangle_Q^{t-1} \lesssim \langle \sigma \rangle_Q^{r(\theta)\frac{1}{1+\epsilon}} \langle w^{1-q'} \rangle_Q^{r(\theta)\frac{\epsilon}{1+\epsilon}} \langle \sigma^{1-p'} \rangle_Q^{s(\theta)\frac{t-1}{1+\delta}} \langle w \rangle_Q^{s(\theta)\frac{\delta(t-1)}{1+\delta}},$$

$$\langle \lambda_0 \rangle_Q \langle \lambda_0^{1-t'} \rangle_Q^{t-1} \lesssim \langle \lambda \rangle_Q^{r(\theta)\frac{1}{1+\epsilon}} \langle w^{1-q'} \rangle_Q^{r(\theta)\frac{\epsilon}{1+\epsilon}} \langle \lambda^{1-p'} \rangle_Q^{s(\theta)\frac{t-1}{1+\delta}} \langle w \rangle_Q^{s(\theta)\frac{\delta(t-1)}{1+\delta}}.$$

Observe that

$$\langle \sigma \rangle_Q^{r(\theta)\frac{1}{1+\epsilon}} \langle \sigma^{1-p'} \rangle_Q^{s(\theta)\frac{t-1}{1+\delta}} = \langle \sigma \rangle_Q^{\frac{t}{p(1-\theta)}} \langle \sigma^{1-p'} \rangle_Q^{(p-1)\frac{t}{p(1-\theta)}} \leq [\sigma]_{A_p}^{\frac{t}{p(1-\theta)}} = [\sigma]_{A_p}^{\frac{q}{q-p\theta}},$$

$$\langle w^{1-q'} \rangle_Q^{r(\theta)\frac{\epsilon}{1+\epsilon}} \langle w \rangle_Q^{s(\theta)\frac{\delta(t-1)}{1+\delta}} = \langle w^{1-q'} \rangle_Q^{(q-1)\frac{t\theta}{q(1-\theta)}} \langle w \rangle_Q^{\frac{t\theta}{q(1-\theta)}} \leq [w]_{A_q}^{\frac{t\theta}{q(1-\theta)}} = [w]_{A_q}^{\frac{\theta}{q-p\theta}}$$

Then we have

$$[\sigma_0]_{A_t} \lesssim [\sigma]_{A_p}^{\frac{q}{q-\theta_p}} [w]_{A_q}^{\frac{\theta_p}{q-\theta_p}} < \infty.$$

Similarly,

$$[\lambda_0]_{A_t} \lesssim [\lambda]_{A_p}^{\frac{q}{q-\theta_p}} [w]_{A_q}^{\frac{\theta_p}{q-\theta_p}} < \infty$$

This completes the proof.

By combining with Theorem 2.2 and Lemma 3.2, we immediately obtain the following interpolation result in the two-weight setting.

Corollary 3.3. Let $p, q \in (1, \infty)$, $\sigma, \lambda \in A_p$ and $w \in A_q$. Then

(3.4)
$$L^{p}(\sigma) = [L^{t}(\sigma_{0}), L^{q}(w)]_{\theta}, \quad L^{p}(\lambda) = [L^{t}(\lambda_{0}), L^{q}(w)]_{\theta}$$

for some $\theta \in (0, 1)$, $t \in (1, \infty)$ and $\sigma_0, \lambda_0 \in A_t$.

By Theorem 2.2 and using the same method as Lemma 3.2 (see also [13, Lemma 3.4]) with some minor changes, we further conclude the following interpolation result. The details are omitted here.

Lemma 3.4. Let $1 , <math>1 < p_1 \leq q_1 < \infty$, $\sigma, \lambda \in A_{p,q}$ and $w_1 \in A_{p_1,q_1}$. Then there exist $\theta \in (0, 1)$, $1 < p_0 \leq q_0 < \infty$ and $\sigma_0, \lambda_0 \in A_{p_0,q_0}$ such that

(3.5)
$$L^{p}(\sigma^{p}) = [L^{p_{0}}(\sigma^{p_{0}}_{0}), L^{p_{1}}(w^{p_{1}}_{1})]_{\theta}, \quad L^{q}(\lambda^{q}) = [L^{q_{0}}(\lambda^{q_{0}}_{0}), L^{q_{1}}(w^{q_{1}}_{1})]_{\theta},$$

where

$$\frac{1}{p} = \frac{1-\theta}{p_0} + \frac{\theta}{p_1}, \quad \frac{1}{q} = \frac{1-\theta}{q_0} + \frac{\theta}{q_1}, \quad \sigma = \sigma_0^{1-\theta} w_1^{\theta}, \quad \lambda = \lambda_0^{1-\theta} w_1^{\theta}.$$

We now turn to verify our main results.

Proof of Theorem 1.1. For any fixed $p \in (1, \infty)$ and $\sigma, \lambda \in A_p$, it suffices to show that T is compact from $L^p(\sigma)$ to $L^p(\lambda)$. Recall that T is compact on $L^q(w)$ for some $q \in (1, \infty)$ and some $w \in A_q$. By Corollary 3.3, we have that (3.4) holds for some $\theta \in (0, 1)$, $t \in (1, \infty)$ and $\sigma_0, \lambda_0 \in A_t$. Since T is bounded from $L^t(\sigma_0)$ to $L^t(\lambda_0)$, by Theorem 3.1, we obtain that T is compact from $L^p(\sigma)$ to $L^p(\lambda)$ by taking $X_0 = L^t(\sigma_0), Y_0 = L^t(\lambda_0)$ and $X_1 = Y_1 = L^q(w)$. This completes the proof.

Proof of Theorem 1.2. For any fixed $p, q \in (1, \infty)$ with $\frac{1}{p} - \frac{1}{q} = \frac{1}{p_1} - \frac{1}{q_1}$ and $\sigma, \lambda \in A_{p,q}$, it suffices to show that T is compact from $L^p(\sigma^p)$ to $L^q(\lambda^q)$. Recall that for $1 < p_1 \leq q_1 < \infty$ and $w_1 \in A_{p_1,q_1}$, T is compact from $L^{p_1}(w_1^{p_1})$ to $L^{q_1}(w_1^{q_1})$. By Lemma 3.4, we have that (3.5) holds for some $\theta \in (0, 1), p_0, q_0 \in (1, \infty)$ with $\frac{1}{p_0} - \frac{1}{q_0} = \frac{1}{p_1} - \frac{1}{q_1}$ and $\sigma_0, \lambda_0 \in A_{p_0,q_0}$. Since T is bounded from $L^{p_0}(\sigma_0^{p_0})$ to $L^{q_0}(\lambda_0^{q_0})$, by Theorem 3.1, we obtain that T is compact from $L^p(\sigma^p)$ to $L^q(\lambda^q)$ by taking $X_0 = L^{p_0}(\sigma_0^{p_0}), Y_0 = L^{q_0}(\lambda_0^{q_0}), X_1 = L^{p_1}(w_1^{p_1})$ and $Y_1 = L^{q_1}(w_1^{q_1})$. This completes the proof.

4. Applications to the two-weight compactness of linear operators

Notice that the unweighted compactness result is probably the most available and relevant case for most applications. Based on this fact, we point out that the nature of extrapolation of two-weight compactness (including the one-weight case) is essentially a combination of the reverse Hölder inequality and interpolation Theorem 3.1. From this point of view, we will give two applications below to the two-weight compactness of linear operators.

Both of our two applications in this section deal with commutators of the form

$$[b,T](f) := bT(f) - T(bf),$$

where the pointwise multiplier b is a locally integrable function. We say that a locally integrable function b belongs to $\text{BMO}_w(\mathbb{R}^d)$ with $w \in A_\infty$ if

$$\|b\|_{\mathrm{BMO}_w(\mathbb{R}^d)} := \sup_{Q \subset \mathbb{R}^d} w(Q)^{-1} \int_Q |b(x) - \langle b \rangle_Q | dx < \infty,$$

where the supremum is taken over all cubes $Q \subset \mathbb{R}^d$ and $w(Q) := \int_Q w$. When $w \equiv 1$, we simply write $BMO_w(\mathbb{R}^d)$ as $BMO(\mathbb{R}^d)$. We then define

$$\operatorname{CMO}(\mathbb{R}^d) := \overline{C_c^{\infty}(\mathbb{R}^d)}^{\operatorname{BMO}(\mathbb{R}^d)}$$

where $C_c^{\infty}(\mathbb{R}^d)$ denotes the space of smooth functions on \mathbb{R}^d with compact support and the closure is in the BMO norm. We say that a function $b \in BMO_w(\mathbb{R}^d)$ belongs to $CMO_w(\mathbb{R}^d)$ with $w \in A_{\infty}$ if the following three conditions hold:

- (a) $\lim_{a \to 0^+} \sup_{|Q|=a} w(Q)^{-1} \int_Q |b(x) \langle b \rangle_Q | dx = 0;$
- (b) $\lim_{a \to \infty} \sup_{|Q|=a} w(Q)^{-1} \int_Q |b(x) \langle b \rangle_Q | dx = 0;$
- (c) $\lim_{a \to \infty} \sup_{Q \cap Q(0,a) = \emptyset} w(Q)^{-1} \int_Q |b(x) \langle b \rangle_Q | dx = 0;$

where Q(0, a) denotes the cube centered at 0 with the side length 2a in \mathbb{R}^d . When $w \equiv 1$, it turns to be an equivalent characterization of $\text{CMO}(\mathbb{R}^d)$ due to Uchiyama [23, Lemma 3].

4.A. Commutators of Calderón–Zygmund operators. In our first application, we study the commutators of Calderón–Zygmund operators. Recall that a linear operator

T is a Calderón–Zygmund operator if it is an integral operator defined initially on $f \in$ $C_c^{\infty}(\mathbb{R}^d)$:

$$T(f)(x) := \int_{\mathbb{R}^d} K(x, y) f(y) dy, \quad x \notin \text{supp}(f),$$

and it extends to a bounded operator on $L^2(\mathbb{R}^d)$, where the kernel K satisfies the standard estimates

$$|K(x,y)| \lesssim \frac{1}{|x-y|^d}, \quad x \neq y,$$

and

$$|K(x+h,y) - K(x,y)| + |K(x,y+h) - K(x,y)| \lesssim \frac{|h|^o}{|x-y|^{d+\delta}}$$

for all $|x - y| \ge 2|h| > 0$ and some fixed $\delta \in (0, 1]$.

The two-weight boundedness of Bloom type for commutators of Calderón–Zygmund operators were first considered by Bloom [3] for the Hilbert transform in one dimension. His result was then improved by Holmes, Lacey and Wick [10] as follows:

Theorem 4.1 ([10]). Let T be a Calderón–Zygmund operator, $p \in (1, \infty)$, and $\sigma, \lambda \in A_p$. Suppose $b \in BMO_{v}(\mathbb{R}^{d})$ with $v = \sigma^{\frac{1}{p}} \lambda^{-\frac{1}{p}}$. Then

$$\|[b,T]\|_{L^p(\sigma)\to L^p(\lambda)} \lesssim \|b\|_{\mathrm{BMO}_v(\mathbb{R}^d)}.$$

For the application of Theorem 3.1, we need the following unweighted compactness result of Uchiyama [23] (see also [5, Theorem 2]).

Theorem 4.2 ([23]). Let T be a Calderón–Zygmund operator and $b \in CMO(\mathbb{R}^d)$. Then the commutator [b,T] is compact on $L^p(\mathbb{R}^d)$ for all $p \in (1,\infty)$.

The following sharp reverse Hölder inequality of the A_p weights for $p \in (1, \infty]$ is useful. **Theorem 4.3** ([19]). (a) Let $p \in (1, \infty]$, $w \in A_p$ and

$$r_w = \begin{cases} 1 + \frac{1}{2^{2p+d+1}[w]_{A_p}}, & p \in (1,\infty); \\ 1 + \frac{1}{2^{d+11}[w]_{A_\infty}}, & p = \infty. \end{cases}$$

Then for any cube $Q \subset \mathbb{R}^d$, $\langle w^{r_w} \rangle_Q^{\frac{1}{r_w}} \leq 2 \langle w \rangle_Q$. (b) If a weight $w \in RH_r$ for some $r \in (1, \infty)$, then $[w]_{A_\infty} \lesssim [w]_{RH_r} r'$.

For simplicity, we write

$$\tilde{r}_{p,w} := \min(r_w, r_{w^{1-p'}}) = 1 + \frac{1}{2^{2\max(p,p')+d+1}[w]_{A_p}^{\max(1,p'-1)}}$$

for any $p \in (1,\infty)$ and $w \in A_p$. Note that if $w \in A_\infty$, then from Theorem 4.3 (a), $w \in RH_{r_w}$. It follows that for any fixed $\eta \in (1, r_w)$ and any cube $Q \subset \mathbb{R}^d$,

$$\langle w^{\eta \frac{r_w}{\eta}} \rangle_Q^{\frac{\eta}{r_w}} = \langle w^{r_w} \rangle_Q^{\frac{\eta}{r_w}} \le 2^{\eta} \langle w \rangle_Q^{\eta} \le 2^{\eta} \langle w^{\eta} \rangle_Q,$$

which implies $w^{\eta} \in RH_{r_w/\eta}$. Thus, from Theorem 4.3 (b), $[w^{\eta}]_{A_{\infty}} \leq 2^{\eta} (r_w/\eta)' < \infty$ and hence $w^{\eta} \in A_{\infty}$.

Very recently, Lacey and Li [16] established the $L^p(\sigma) \to L^p(\lambda)$ compactness for commutators of Calderón–Zygmund operators under the assumption $b \in \text{CMO}_v(\mathbb{R}^d)$ with $v = \sigma^{\frac{1}{p}} \lambda^{-\frac{1}{p}}$, whose method relies on a fine decomposition of the Calderón–Zygmund kernel and uses the idea of the approximation of compact operators. We note that a combination of Theorems 4.1 and 4.2, together with Theorems 3.1 and 4.3, readily gives another version of the two-weight compactness of commutators of Calderón–Zygmund operators.

Corollary 4.4. Let T be a Calderón–Zygmund operator, $p \in (1, \infty)$, two weights $\sigma, \lambda \in A_p$. Suppose

$$b \in \overline{\bigcup_{\eta \in (1, r_{\sigma, \lambda}]} \operatorname{BMO}_{v^{\eta}}(\mathbb{R}^d) \cap \operatorname{CMO}(\mathbb{R}^d)}^{\operatorname{BMO}_v(\mathbb{R}^d)}$$

where

$$v = \sigma^{\frac{1}{p}} \lambda^{-\frac{1}{p}} \in A_2, \quad r_{\sigma,\lambda} = \min(\tilde{r}_{p,\sigma}, \tilde{r}_{p,\lambda})$$

= $1 + \frac{1}{2^{2\max(p,p')+d+1}\max\left([\sigma]_{A_p}^{\max(1,p'-1)}, [\lambda]_{A_p}^{\max(1,p'-1)}\right)}$

Then the commutator [b,T] is compact from $L^p(\sigma)$ to $L^p(\lambda)$.

Before giving our proof, we discuss the relationship between $BMO_{v^{\eta}}(\mathbb{R}^d) \cap CMO(\mathbb{R}^d)$ and $BMO_v(\mathbb{R}^d)$. Indeed, inspired by the work of Lerner, Ombrosi and Rivera-Ríos [17, Lemma 4.8], we have the following result.

Lemma 4.5. Let $v \in A_2$ and $r_v = 1 + \frac{1}{2^{5+d}[v]_{A_2}}$. Then for any $\eta \in (1, r_v]$, BMO_{v^η}(\mathbb{R}^d) \cap CMO(\mathbb{R}^d) \subset CMO_v(\mathbb{R}^d).

Proof. From Theorem 4.3 and the Hölder inequality, it readily follows that when $\eta \in (1, r_v]$, for any cube $Q \subset \mathbb{R}^d$,

$$\frac{1}{2} \langle v^{\eta} \rangle_Q^{\frac{1}{\eta}} \le \langle v \rangle_Q \le \langle v^{\eta} \rangle_Q^{\frac{1}{\eta}},$$

which further implies that

$$\begin{split} \frac{1}{v(Q)} \int_{Q} |b(x) - \langle b \rangle_{Q} |dx &\approx \frac{1}{v^{\eta}(Q)^{\frac{1}{\eta}} |Q|^{\frac{1}{\eta'}}} \int_{Q} |b(x) - \langle b \rangle_{Q} |dx \\ &\approx \left(\frac{1}{v^{\eta}(Q)} \int_{Q} |b(x) - \langle b \rangle_{Q} |dx\right)^{\frac{1}{\eta}} \left(\frac{1}{|Q|} \int_{Q} |b(x) - \langle b \rangle_{Q} |dx\right)^{\frac{1}{\eta'}} \\ &\lesssim \|b\|_{\mathrm{BMO}_{v^{\eta}}(\mathbb{R}^{d})}^{\frac{1}{\eta}} \left(\frac{1}{|Q|} \int_{Q} |b(x) - \langle b \rangle_{Q} |dx\right)^{\frac{1}{\eta'}}. \end{split}$$

Then from this and the definition of $CMO_v(\mathbb{R}^d)$, we conclude that

 $BMO_{v^{\eta}}(\mathbb{R}^d) \cap CMO(\mathbb{R}^d) \subset CMO_v(\mathbb{R}^d).$

This completes the proof.

Similarly, we conclude that if $v \in A_{\infty}$, let $r_v = 1 + \frac{1}{2^{d+11}[w]_{A_{\infty}}}$, then for any $\eta \in (1, r_v)$, $v^{\eta} \in A_{\infty}$ and

$$BMO_{v^{\eta}}(\mathbb{R}^d) \cap CMO(\mathbb{R}^d) \subset CMO_v(\mathbb{R}^d).$$

Proof of Corollary 4.4. First, by Theorem 4.1, we have that

$$||[b_0, T] - [b_1, T]||_{L^p(\sigma) \to L^p(\lambda)} = ||[b_0 - b_1, T]||_{L^p(\sigma) \to L^p(\lambda)} \lesssim ||b_0 - b_1||_{BMO_v(\mathbb{R}^d)},$$

provided that $b_0, b_1 \in BMO_v(\mathbb{R}^d)$. Therefore it suffices to show the two-weight compactness of the commutator [b, T] with $b \in BMO_{v^{\eta}}(\mathbb{R}^d) \cap CMO(\mathbb{R}^d)$ for $\eta \in (1, r_{\sigma, \lambda}]$.

From $b \in \text{CMO}(\mathbb{R}^d)$ and Theorem 4.2, the commutator [b, T] is compact on $L^p(\mathbb{R}^d)$. Moreover, for $\sigma, \lambda \in A_p$, by Theorem 4.3, it follows that $\sigma^{\eta}, \lambda^{\eta} \in A_p$ for $\eta \in (1, r_{\sigma,\lambda}]$. Then from $b \in \text{BMO}_{v^{\eta}}(\mathbb{R}^d)$ and Theorem 4.1, [b, T] is bounded from $L^p(\sigma^{\eta})$ to $L^p(\lambda^{\eta})$. Thus, from Theorem 3.1, we obtain that [b, T] is compact from $L^p(\sigma)$ to $L^p(\lambda)$ by taking $X_0 = L^p(\sigma^{\eta}), Y_0 = L^p(\lambda^{\eta}), X_1 = Y_1 = L^p(\mathbb{R}^d)$ and $\theta = 1 - 1/\eta$. This completes the proof.

Remark 4.6. Since our approach relies on the interpolation of compactness, the assumption of b in Corollary 4.4 seems to be slightly stronger than $b \in \text{CMO}_v(\mathbb{R}^d)$ in [16, Theorem 1.2] by Lemma 4.5. However, when d = 1, for any $v \in A_2$ and $\eta \in (1, r_v]$, we claim that

(4.1)
$$\overline{\mathrm{BMO}_{v^{\eta}}(\mathbb{R}) \cap \mathrm{CMO}(\mathbb{R})}^{\mathrm{BMO}_{v}(\mathbb{R})} = \mathrm{CMO}_{v}(\mathbb{R}).$$

To see this, note that by Theorem 4.3, we have $v^{\eta} \in A_2$. Then from [16, Theorem 4.1], we obtain that

$$C_c^{\infty}(\mathbb{R}) \subset \mathrm{BMO}_{v^{\eta}}(\mathbb{R}), \ \overline{C_c^{\infty}(\mathbb{R})}^{\mathrm{BMO}_v(\mathbb{R})} = \mathrm{CMO}_v(\mathbb{R}),$$

from which (4.1) follows. But it is still unknown whether (4.1) holds when d > 1, so recovering [16, Theorem 1.2] by our approach might deserve further study.

4.B. Commutators of fractional integral operators. In our second application, we study the commutators of fractional integral operators. Recall the classical fractional integral operator (or Riesz potential): given $\alpha \in (0, d)$, for a Schwartz function $f \in \mathcal{S}(\mathbb{R}^d)$, define the fractional integral operator I_{α} by

$$I_{\alpha}(f)(x) := \int_{\mathbb{R}^d} \frac{f(y)}{|x-y|^{d-\alpha}} dy.$$

The two-weight norm inequalities of Bloom type for commutators of fractional integrals have been considered in many literatures. For instance, Ding and Lu [9, Theorem 1] established the two-weight boundedness of commutators of fractional integrals $T_{\Omega,\alpha}$ with rough kernels $\Omega \in L^{\infty}(S^{d-1})$. However, to obtain our two-weight compactness result, we need a refined upper bounded estimate established recently by Holmes, Rahm and Spencer [11, Theorem 1.1] as follows:

Theorem 4.7 ([11]). Let $\alpha \in (0, d)$, $p, q \in (1, \infty)$ with $\frac{1}{p} = \frac{1}{q} + \frac{\alpha}{d}$ and $\sigma, \lambda \in A_{p,q}$. Suppose $b \in BMO_v(\mathbb{R}^d)$ with $v = \sigma \lambda^{-1}$. Then

$$\|[b, I_{\alpha}]\|_{L^{p}(\sigma^{p}) \to L^{q}(\lambda^{q})} \approx \|b\|_{\mathrm{BMO}_{v}(\mathbb{R}^{d})}.$$

For the application of Theorem 3.1, we need the following unweighted compactness result of Wang [24].

Theorem 4.8 ([24]). Let $\alpha \in (0, d)$, $p, q \in (1, \infty)$ with $\frac{1}{p} = \frac{1}{q} + \frac{\alpha}{d}$. Suppose $b \in \text{CMO}(\mathbb{R}^d)$, then $[b, I_{\alpha}] : L^p(\mathbb{R}^d) \to L^q(\mathbb{R}^d)$ is a compact operator.

A combination of Theorems 4.7 and 4.8, together with Theorems 3.1 and 4.3, gives the following two-weight compactness of commutators of fractional integral operators which is a new result to our best knowledge.

Corollary 4.9. Let $\alpha \in (0, d)$, $p, q \in (1, \infty)$ with $\frac{1}{p} = \frac{1}{q} + \frac{\alpha}{d}$ and $\sigma, \lambda \in A_{p,q}$. Suppose BMO_v(\mathbb{R}^d)

$$b \in \bigcup_{\eta \in (1, r_{\sigma, \lambda}]} BMO_{v^{\eta}}(\mathbb{R}^d) \cap CMO(\mathbb{R}^d)^{DHO_v(\mathbb{R}^d)},$$

where

$$v = \sigma \lambda^{-1} \in A_2, \quad r_{\sigma,\lambda} = \min(\tilde{r}_{1+p'/q,\sigma^{-p'}}, \tilde{r}_{1+q/p',\lambda^q})$$
$$= 1 + \frac{1}{2^{2\max(p'/q,q/p')+d+3}\max\left([\sigma]_{A_{p,q}}^{\max(p',q)}, [\lambda]_{A_{p,q}}^{\max(p',q)}\right)}$$

Then the commutator $[b, I_{\alpha}]$ is compact from $L^{p}(\sigma^{p})$ to $L^{q}(\lambda^{q})$.

Proof. First, by Theorem 4.7, we have that

$$\|[b_0, I_\alpha] - [b_1, I_\alpha]\|_{L^p(\sigma^p) \to L^q(\lambda^q)} = \|[b_0 - b_1, I_\alpha]\|_{L^p(\sigma^p) \to L^q(\lambda^q)} \approx \|b_0 - b_1\|_{BMO_v(\mathbb{R}^d)},$$

provided that $b_0, b_1 \in BMO_v(\mathbb{R}^d)$. Therefore it suffices to show the two-weight compactness of the commutator $[b, I_\alpha]$ with $b \in BMO_{v^\eta}(\mathbb{R}^d) \cap CMO(\mathbb{R}^d)$ for $\eta \in (1, r_{\sigma,\lambda}]$.

From $b \in \text{CMO}(\mathbb{R}^d)$ and Theorem 4.8, we deduce that the commutator $[b, I_\alpha]$: $L^p(\mathbb{R}^d) \to L^q(\mathbb{R}^d)$ is a compact operator. Observe that for any $w \in A_{p,q}$,

$$[w]_{A_{p,q}} = [w^q]_{A_{1+q/p'}}^{\frac{1}{q}} = [w^{-p'}]_{A_{1+p'/q}}^{\frac{1}{p'}}$$

It follows that for $\sigma, \lambda \in A_{p,q}$, we have $\sigma^{-p'} \in A_{1+p'/q}$ and $\lambda^q \in A_{1+q/p'}$. Then by Theorem 4.3, when $\eta \in (1, r_{\sigma,\lambda}]$, we see that $\sigma^{-p'\eta} \in A_{1+p'/q}$ and $\lambda^{q\eta} \in A_{1+q/p'}$. From this, we obtain $\sigma^{\eta}, \lambda^{\eta} \in A_{p,q}$. Moreover, from $b \in BMO_{v^{\eta}}(\mathbb{R}^d)$ and Theorem 4.7, $[b, I_{\alpha}]$ is bounded from $L^p(\sigma^{p\eta})$ to $L^q(\lambda^{q\eta})$. Thus, from Theorem 3.1, we obtain that $[b, I_{\alpha}]$ is compact from $L^p(\sigma^p)$ to $L^q(\lambda^q)$ by taking $X_0 = L^p(\sigma^{p\eta}), Y_0 = L^q(\lambda^{q\eta}), X_1 = L^p(\mathbb{R}^d), Y_1 = L^q(\mathbb{R}^d)$ and $\theta = 1 - 1/\eta$. This completes the proof.

5. Applications to the two-weight compactness of bilinear operators

In this section, we show that our approach applied to linear operators still works in the multilinear case. In the one-weight setting, this topic has been discussed carefully in [14, 4, 25]. For simplicity, we only study the two-weight compactness of bilinear operators. To be precise, we consider the first order bilinear commutators of bilinear Calderón–Zygmund operators defined as follows.

Let $\Delta := \{(x, y_0, y_1) \in (\mathbb{R}^d)^3 : x = y_0 = y_1\}$. We say that $K : (\mathbb{R}^d)^3 \setminus \Delta \to \mathbb{C}$ is a bilinear Calderón–Zygmund kernel if

$$|K(x, y_0, y_1)| \lesssim \frac{1}{(|x - y_0| + |x - y_1|)^{2d}},$$

and

$$|K(x+h, y_0, y_1) - K(x, y_0, y_1)| + |K(x, y_0 + h, y_1) - K(x, y_0, y_1)| + |K(x, y_0, y_1 + h) - K(x, y_0, y_1)| \lesssim \frac{|h|^{\delta}}{(|x - y_0| + |x - y_1|)^{2d + \delta}},$$

for all $\max(|x-y_0|, |x-y_1|) \geq 2|h|$ and some fixed $\delta \in (0, 1]$. Let T be a bilinear operator initially defined from $(\mathcal{S}(\mathbb{R}^d))^2$ to $\mathcal{S}'(\mathbb{R}^d)$. Then, we say that T is a bilinear Calderón– Zygmund operator if it extends to be bounded from $L^{p_0}(\mathbb{R}^d) \times L^{p_1}(\mathbb{R}^d)$ to $L^p(\mathbb{R}^d)$ for some $p_0, p_1 \in (1, \infty)$, where $\frac{1}{p} = \frac{1}{p_0} + \frac{1}{p_1}$, and there exists a bilinear Calderón–Zygmund kernel K such that for all $f_0, f_1 \in C_c^{\infty}(\mathbb{R}^d)$,

$$T(f_0, f_1)(x) := \int_{(\mathbb{R}^d)^2} K(x, y_0, y_1) f_0(y_0) f_1(y_1) dy_0 dy_1, \quad x \notin \operatorname{supp}(f_0) \cap \operatorname{supp}(f_1).$$

Given a bilinear Calderón–Zygmund operator T, a pair of locally integrable functions $\mathbf{b} = (b_0, b_1)$ and $\mathcal{I}(\neq \emptyset) \subset \{0, 1\}$, we define the first order bilinear commutators as follows:

(5.1)
$$[\mathbf{b}, T]_{\mathcal{I}}(f_0, f_1) = \begin{cases} b_0 T(f_0, f_1) - T(b_0 f_0, f_1), & \mathcal{I} = \{0\}; \\ b_1 T(f_0, f_1) - T(f_0, b_1 f_1), & \mathcal{I} = \{1\}; \\ [\mathbf{b}, [\mathbf{b}, T]_{\{0\}}]_{\{1\}}(f_0, f_1) = [\mathbf{b}, [\mathbf{b}, T]_{\{1\}}]_{\{0\}}(f_0, f_1), & \mathcal{I} = \{0, 1\} \end{cases}$$

In the bilinear case, we need a bilinear version of the interpolation theorem of compactness due to Cobos, Fernández-Cabrera and Martínez [6, Theorem 3.2]. Before this, we need to introduce some notations. For a Banach couple $\bar{X} = (X_0, X_1)$, we write X_i° for the closure of $X_0 \cap X_1$ in the norm of X_i for i = 0, 1. We denote by $\mathcal{B}(\bar{X} \times \bar{Y}, \bar{Z})$ the space of all bilinear operators T which are defined from $(X_0 \cap X_1) \times (Y_0 \cap Y_1)$ to $Z_0 \cap Z_1$ and satisfy

$$||T(f_0, f_1)||_{Z_i} \lesssim ||f_0||_{X_i} ||f_1||_{Y_i}, f_0 \in X_0 \cap X_1, f_1 \in Y_0 \cap Y_1, i = 0, 1,$$

where $\overline{Y} := (Y_0, Y_1)$ and $\overline{Z} := (Z_0, Z_1)$ are Banach couples.

Theorem 5.1 ([6]). Let $\bar{X} = (X_0, X_1)$ and $\bar{Y} = (Y_0, Y_1)$ be Banach couples. Assume that (Ω, μ) is a σ -finite measure space. Let $\bar{Z} = (Z_0, Z_1)$ be a couple of Banach function spaces on Ω , $\theta \in (0, 1)$ and $T \in \mathcal{B}(\bar{X} \times \bar{Y}, \bar{Z})$. If $T : X_0^{\circ} \times Y_0^{\circ} \to Z_0$ compactly and Z_0 has absolutely continuous norm, then T can be uniquely extended to a compact bilinear operator from $[X_0, X_1]_{\theta} \times [Y_0, Y_1]_{\theta}$ to $[Z_0, Z_1]_{\theta}$.

For the definitions of the Banach function space and the absolutely continuous norm, see, for example, [6, 14]. It is known that the assumptions in Theorem 5.1 hold for the weighted Lebesgue spaces, see [14, Lemma 3.3], which is sufficient for our applications of Theorem 5.1.

We recall that a weight vector $\mathbf{w} = (w_0, w_1)$ is called a (multiple) $A_{\mathbf{p}}$ weight (write $\mathbf{w} \in A_{\mathbf{p}}$) if

$$[\mathbf{w}]_{A_{\mathbf{p}}} := \sup_{Q \subset \mathbb{R}^d} \langle w_{\mathbf{p}} \rangle_Q^{\frac{1}{p}} \langle w_0^{1-p_0'} \rangle_Q^{\frac{1}{p_0'}} \langle w_1^{1-p_1'} \rangle_Q^{\frac{1}{p_1'}} < \infty,$$

where $\mathbf{p} = (p_0, p_1) \in (1, \infty)^2$, $\frac{1}{p} = \frac{1}{p_0} + \frac{1}{p_1}$ and $w_{\mathbf{p}} = w_0^{\frac{p}{p_0}} w_1^{\frac{p}{p_1}}$. The following is a characterization of $A_{\mathbf{p}}$ weights which is quite useful in applications:

(5.2)
$$\mathbf{w} \in A_{\mathbf{p}} \iff w_{\mathbf{p}} \in A_{2p}, \ w_i^{1-p_i'} \in A_{2p_i'}, \ i = 0, 1.$$

The two-weight boundedness of Bloom type of multilinear commutators of multilinear Calderón–Zygmund operators were first studied by Kunwar and Ou in [15]. Their result was then improved by Li [18] in terms of the genuinely multilinear weights. The following is a bilinear version of [18, Theorem 3.5].

Theorem 5.2 ([18]). Let T be a bilinear Calderón–Zygmund operator and $\mathcal{I}(\neq \emptyset) \subset \{0,1\}$. Let $[\mathbf{b},T]_{\mathcal{I}}$ be defined as in (5.1), $p_0, p_1 \in (1,\infty)$ and $\frac{1}{p} = \frac{1}{p_0} + \frac{1}{p_1}$. Suppose that $(w_0, w_1) \in A_{\mathbf{p}}$, where

$$w_i = \begin{cases} \sigma_i, & i \notin \mathcal{I}; \\ \sigma_i \text{ or } \lambda_i, & i \in \mathcal{I}. \end{cases}$$

 $If v_i := \sigma_i^{\frac{1}{p_i}} \lambda_i^{-\frac{1}{p_i}} \in A_{\infty} \text{ and } b_i \in BMO_{v_i}(\mathbb{R}^d) \text{ for each } i \in \mathcal{I}, \text{ then} \\ \|[\mathbf{b}, T]_{\mathcal{I}}\|_{L^{p_0}(\sigma_0) \times L^{p_1}(\sigma_1) \to L^p(\prod_{i \in \mathcal{I}} \lambda_i^{\frac{p}{p_i}} \prod_{j \notin \mathcal{I}} \sigma_j^{\frac{p}{p_j}})} \lesssim \prod_{i \in \mathcal{I}} \|b_i\|_{BMO_{v_i}(\mathbb{R}^d)}.$

We would like to mention that different from the linear case, in the multilinear case, by (5.2), we only have $\sigma_i^{1-p'_i}$, $\lambda_i^{1-p'_i} \in A_{2p'_i}$ for each $i \in \mathcal{I}$, which implies $v_i^{\frac{1}{2}} \in A_2$, but may not ensure $v_i \in A_{\infty}$, see [18, Example 2.12] for more details.

For the application of Theorem 5.1, we need the following unweighted compactness result of Bényi and Torres [1, Theorem 1], see also [22] for some developments in the one-weight setting.

Theorem 5.3 ([1]). Let T be a bilinear Calderón–Zygmund operator and $\mathcal{I}(\neq \emptyset) \subset \{0, 1\}$. Let $[\mathbf{b}, T]_{\mathcal{I}}$ be defined as in (5.1), $p_0, p_1, p \in (1, \infty)$ such that $\frac{1}{p} = \frac{1}{p_0} + \frac{1}{p_1}$. If $b_i \in \text{CMO}(\mathbb{R}^d)$ for each $i \in \mathcal{I}$, then $[\mathbf{b}, T]_{\mathcal{I}}$ is compact from $L^{p_0}(\mathbb{R}^d) \times L^{p_1}(\mathbb{R}^d)$ to $L^p(\mathbb{R}^d)$.

A combination of Theorems 5.2 and 5.3, together with Theorems 5.1 and 4.3, gives the following two-weight compactness of bilinear commutators of bilinear Calderón–Zygmund operators which is a new result to our best knowledge.

Corollary 5.4. Let T be a bilinear Calderón–Zygmund operator and $\mathcal{I}(\neq \emptyset) \subset \{0, 1\}$. Let $[\mathbf{b}, T]_{\mathcal{I}}$ be defined as in (5.1), $p_0, p_1, p \in (1, \infty)$ such that $\frac{1}{p} = \frac{1}{p_0} + \frac{1}{p_1}$. Suppose that $(w_0, w_1) \in A_{\mathbf{p}}$, where

$$w_i = \begin{cases} \sigma_i, & i \notin \mathcal{I}; \\ \sigma_i \text{ or } \lambda_i, & i \in \mathcal{I}. \end{cases}$$

If $v_i := \sigma_i^{\frac{1}{p_i}} \lambda_i^{-\frac{1}{p_i}} \in A_\infty$ and

$$b_i \in \overline{\bigcup_{\eta_{\mathcal{I}} \in (1, r_{\sigma_{\mathcal{I}}, \lambda_{\mathcal{I}}})} \mathrm{BMO}_{v_i^{\eta_{\mathcal{I}}}}(\mathbb{R}^d) \cap \mathrm{CMO}(\mathbb{R}^d)}^{\mathrm{BMO}_{v_i}(\mathbb{R}^d)}$$

for each $i \in \mathcal{I}$, where

$$\begin{aligned} r_{\sigma_{\mathcal{I}},\lambda_{\mathcal{I}}} &:= \min\left(\tilde{r}_{2p,\sigma_{0}^{p}\sigma_{1}^{p}}, \tilde{r}_{2p,\prod_{i\in\mathcal{I}}\lambda_{i}^{p}\prod_{j\notin\mathcal{I}}\sigma_{j}^{p}}, \tilde{r}_{2p_{0}',\sigma_{0}^{1-p_{0}'}}, \tilde{r}_{2p_{1}',\sigma_{1}^{1-p_{1}'}}, \min_{i\in\mathcal{I}}\tilde{r}_{2p_{i}',\lambda_{i}^{1-p_{i}'}}, \min_{i\in\mathcal{I}}r_{v_{i}}\right). \end{aligned}$$

$$Then \ [\mathbf{b},T]_{\mathcal{I}} \ is \ compact \ from \ L^{p_{0}}(\sigma_{0}) \times L^{p_{1}}(\sigma_{1}) \ to \ L^{p}(\prod_{i\in\mathcal{I}}\lambda_{i}^{p}\prod_{j\notin\mathcal{I}}\sigma_{j}^{p}). \end{aligned}$$

Proof. We only give the detailed proof of the corollary in the case $\mathcal{I} = \{0, 1\}$, since the proofs of the other two cases are similar and easier. First, by Theorem 5.2, we have that

$$\begin{split} \|[\mathbf{b}, T]_{\mathcal{I}} - [\mathbf{b}^*, T]_{\mathcal{I}} \| \\ &= \|[\mathbf{b}, [\mathbf{b}, T]_{\{0\}}]_{\{1\}} - [\mathbf{b}^*, [\mathbf{b}, T]_{\{0\}}]_{\{1\}} + [\mathbf{b}^*, [\mathbf{b}, T]_{\{0\}}]_{\{1\}} - [\mathbf{b}^*, [\mathbf{b}^*, T]_{\{0\}}]_{\{1\}} \| \\ &\leq \|[\mathbf{b} - \mathbf{b}^*, [\mathbf{b}, T]_{\{0\}}]_{\{1\}} \| + \|[\mathbf{b}^*, [\mathbf{b} - \mathbf{b}^*, T]_{\{0\}}]_{\{1\}} \| \end{split}$$

 $\lesssim \|b_0\|_{\mathrm{BMO}_{v_0}(\mathbb{R}^d)} \|b_1 - b_1^*\|_{\mathrm{BMO}_{v_1}(\mathbb{R}^d)} + \|b_1^*\|_{\mathrm{BMO}_{v_1}(\mathbb{R}^d)} \|b_0 - b_0^*\|_{\mathrm{BMO}_{v_0}(\mathbb{R}^d)},$

provided that $\mathbf{b} = (b_0, b_1), \mathbf{b}^* = (b_0^*, b_1^*) \in BMO_{v_0}(\mathbb{R}^d) \times BMO_{v_1}(\mathbb{R}^d)$, where

$$\|\cdot\| := \|\cdot\|_{L^{p_0}(\sigma_0) \times L^{p_1}(\sigma_1) \to L^p(\lambda_0^{\frac{p}{p_0}} \lambda_1^{\frac{p}{p_1}})}$$

Therefore, it suffices to show the two-weight compactness of the bilinear commutator $[\mathbf{b}, T]_{\mathcal{I}}$ with $b_i \in \text{BMO}_{v_i^{\eta_{\mathcal{I}}}}(\mathbb{R}^d) \cap \text{CMO}(\mathbb{R}^d)$ for each $i \in \mathcal{I}$, where $\eta_{\mathcal{I}} \in (1, r_{\sigma_{\mathcal{I}}, \lambda_{\mathcal{I}}})$.

From $b_i \in \text{CMO}(\mathbb{R}^d)$ for each $i \in \mathcal{I}$ and Theorem 5.3, we deduce that $[\mathbf{b}, T]_{\mathcal{I}}$ is compact from $L^{p_0}(\mathbb{R}^d) \times L^{p_1}(\mathbb{R}^d)$ to $L^p(\mathbb{R}^d)$. Next, for $(\sigma_0, \sigma_1), (\lambda_0, \lambda_1) \in A_{\mathbf{p}}$, from (5.2),

$$\sigma_0^{\frac{p}{p_0}} \sigma_1^{\frac{p}{p_1}}, \lambda_0^{\frac{p}{p_0}} \lambda_1^{\frac{p}{p_1}} \in A_{2p}, \ \sigma_i^{1-p_i'}, \lambda_i^{1-p_i'} \in A_{2p_i'}, \ i = 0, 1.$$

Then by Theorem 4.3, when $\eta_{\mathcal{I}} \in (1, r_{\sigma_{\mathcal{I}}, \lambda_{\mathcal{I}}})$, we see that

$$\sigma_0^{\frac{\eta_{\mathcal{I}}p}{p_0}}\sigma_1^{\frac{\eta_{\mathcal{I}}p}{p_1}}, \lambda_0^{\frac{\eta_{\mathcal{I}}p}{p_0}}\lambda_1^{\frac{\eta_{\mathcal{I}}p}{p_1}} \in A_{2p}, \ \sigma_i^{\eta_{\mathcal{I}}(1-p_i')}, \lambda_i^{\eta_{\mathcal{I}}(1-p_i')} \in A_{2p_i'}, \ v_i^{\eta_{\mathcal{I}}} \in A_{\infty}, \ i = 0, 1.$$

From this, again by (5.2), we have $(\sigma_0^{\eta_{\mathcal{I}}}, \sigma_1^{\eta_{\mathcal{I}}}), (\lambda_0^{\eta_{\mathcal{I}}}, \lambda_1^{\eta_{\mathcal{I}}}) \in A_{\mathbf{p}}$. Note that from $b_i \in BMO_{v_i^{\eta_{\mathcal{I}}}}(\mathbb{R}^d)$ for each $i \in \mathcal{I}$ and Theorem 5.2, we deduce that $[\mathbf{b}, T]_{\mathcal{I}}$ is bounded from $L^{p_0}(\sigma_0^{\eta_{\mathcal{I}}}) \times L^{p_1}(\sigma_1^{\eta_{\mathcal{I}}})$ to $L^p(\lambda_0^{\frac{\eta_{\mathcal{I}}p}{p_0}} \lambda_1^{\frac{\eta_{\mathcal{I}}p}{p_1}})$. Thus, by Theorem 5.1 with

$$(X_0, X_1) = (L^{p_0}(\mathbb{R}^d), L^{p_0}(\sigma_0^{\eta_{\mathcal{I}}})), \quad (Y_0, Y_1) = (L^{p_1}(\mathbb{R}^d), L^{p_1}(\sigma_1^{\eta_{\mathcal{I}}})),$$
$$(Z_0, Z_1) = (L^p(\mathbb{R}^d), L^p(\lambda_0^{\frac{\eta_{\mathcal{I}}p}{p_0}}\lambda_1^{\frac{\eta_{\mathcal{I}}p}{p_1}})), \quad \theta = \frac{1}{\eta_{\mathcal{I}}},$$

we obtain that $[\mathbf{b}, T]_{\mathcal{I}}$ is compact from $L^{p_0}(\sigma_0) \times L^{p_1}(\sigma_1)$ to $L^p(\lambda_0^{\frac{p}{p_0}} \lambda_1^{\frac{p}{p_1}})$. This completes the proof.

Remark 5.5. From our applications in Sections 4 and 5, we see that the two-weight compactness on weighted Lebesgue spaces of certain (multi-)linear operators is naturally obtained from the unweighted compactness and the two-weight boundedness without dealing with any concrete forms and characteristics of these operators. Based on this fact, our results may still valid for more general (multi-)linear integral operators, e.g., ω -Calderón–Zygmund operators with ω satisfying the Dini type estimates studied in [17].

Acknowledgments: Wu is supported by the NNSF of China (Grant Nos. 11871101 and 11771358). Yang is supported by the NNSF of China (Grant Nos. 11971402 and 11871254).

References

- A. Bényi and R. H. Torres. Compact bilinear operators and commutators. Proc. Amer. Math. Soc., 141(10):3609–3621, 2013.
- [2] J. Bergh and J. Löfström. Interpolation spaces. An introduction. Springer-Verlag, Berlin-New York, 1976. Grundlehren der Mathematischen Wissenschaften, No. 223.
- [3] S. Bloom. A commutator theorem and weighted BMO. Trans. Amer. Math. Soc., 292(1):103–122, 1985.
- [4] M. Cao, A. Olivo, and K. Yabuta. Extrapolation for multilinear compact operators and applications. arXiv:2011.13191, 2020.
- [5] A. Clop and V. Cruz. Weighted estimates for Beltrami equations. Ann. Acad. Sci. Fenn. Math., 38(1):91–113, 2013.
- [6] F. Cobos, L. M. Fernández-Cabrera, and A. Martínez. On compactness results of Lions-Peetre type for bilinear operators. *Nonlinear Anal.*, 199:111951, 9, 2020.

- [7] M. Cwikel. Real and complex interpolation and extrapolation of compact operators. Duke Math. J., 65(2):333–343, 1992.
- [8] M. Cwikel and N. J. Kalton. Interpolation of compact operators by the methods of Calderón and Gustavsson-Peetre. Proc. Edinburgh Math. Soc. (2), 38(2):261–276, 1995.
- [9] Y. Ding and S. Lu. Higher order commutators for a class of rough operators. Ark. Mat., 37(1):33–44, 1999.
- [10] I. Holmes, M. T. Lacey, and B. D. Wick. Commutators in the two-weight setting. Math. Ann., 367(1-2):51-80, 2017.
- I. Holmes, R. Rahm, and S. Spencer. Commutators with fractional integral operators. *Studia Math.*, 233(3):279–291, 2016.
- [12] T. Hytönen. Extrapolation of compactness on weighted spaces. arXiv:2003.01606, 2020.
- [13] T. Hytönen and S. Lappas. Extrapolation of compactness on weighted spaces II: Off-diagonal and limited range estimates. arXiv:2006.15858, 2020.
- [14] T. Hytönen and S. Lappas. Extrapolation of compactness on weighted spaces III: Bilinear operators. arXiv:2012.10407, 2020.
- [15] I. Kunwar and Y. Ou. Two-weight inequalities for multilinear commutators. New York J. Math., 24:980–1003, 2018.
- [16] M. Lacey and J. Li. Compactness of commutator of Riesz transforms in the two weight setting. arXiv:2010.15451, 2020.
- [17] A. K. Lerner, S. Ombrosi, and I. P. Rivera-Ríos. Commutators of singular integrals revisited. Bull. Lond. Math. Soc., 51(1):107–119, 2019.
- [18] K. Li. Multilinear commutators in the two-weight setting. arXiv:2006.09071, 2020.
- [19] C. Pérez. A course on singular integrals and weights. Advanced courses in Mathematics at the CRM-Barcelona series, Birkhaüser, 2013.
- [20] E. M. Stein and G. Weiss. Interpolation of operators with change of measures. Trans. Amer. Math. Soc., 87(1):159–172, 1958.
- [21] C. B. Stockdale, P. Villarroya, and B. D. Wick. Sparse domination results for compactness on weighted spaces. arXiv:1912.10290, 2019.
- [22] J. Tao, Q. Xue, D. Yang, and W. Yuan. XMO and weighted compact bilinear commutators. arXiv:1909.03173, 2019.
- [23] A. Uchiyama. On the compactness of operators of Hankel type. Tohoku Math. J. (2), 30(1):163–171, 1978.
- [24] S. Wang. The compactness of the commutator of fractional integral operator. Chin. Ann. Math, 8:475–482, 1987.
- [25] S. Wang and Q. Xue. On weighted compactness of commutators of bilinear maximal Calderón-Zygmund singular integral operators. arXiv:2012.12747, 2020.

Shenyu Liu, School of Mathematical Sciences, Xiamen University, Xiamen 361005, China

Email address: shenyuliu@stu.xmu.edu.cn

HUOXIONG WU, SCHOOL OF MATHEMATICAL SCIENCES, XIAMEN UNIVERSITY, XIAMEN 361005, CHINA

Email address: huoxwu@xmu.edu.cn

Dongyong Yang (Corresponding Author), School of Mathematical Sciences, Xiamen University, Xiamen 361005, China

Email address: dyyang@xmu.edu.cn