

Semi-Supervised Statistical Inference for High-Dimensional Linear Regression with Blockwise Missing Data

Fei Xue

FEI.XUE@PENNMEDICINE.UPENN.EDU

*Department of Biostatistics, Epidemiology and Informatics
University of Pennsylvania
Philadelphia, PA 19104, USA*

Rong Ma

RONGM@PENNMEDICINE.UPENN.EDU

*Department of Biostatistics, Epidemiology and Informatics
University of Pennsylvania
Philadelphia, PA 19104, USA*

Hongzhe Li

HONGZHE@PENNMEDICINE.UPENN.EDU

*Department of Biostatistics, Epidemiology and Informatics
University of Pennsylvania
Philadelphia, PA 19104, USA*

Editor:

Abstract

Blockwise missing data occurs frequently when we integrate multisource or multimodality data where different sources or modalities contain complementary information. In this paper, we consider a high-dimensional linear regression model with blockwise missing covariates and a partially observed response variable. Under this semi-supervised framework, we propose a computationally efficient estimator for the regression coefficient vector based on carefully constructed unbiased estimating equations and a multiple blockwise imputation procedure, and obtain its rates of convergence. Furthermore, building upon an innovative semi-supervised projected estimating equation technique that intrinsically achieves bias-correction of the initial estimator, we propose nearly unbiased estimators for the individual regression coefficients that are asymptotically normally distributed under mild conditions. By carefully analyzing these debiased estimators, asymptotically valid confidence intervals and statistical tests about each regression coefficient are constructed. Numerical studies and application analysis of the Alzheimer's Disease Neuroimaging Initiative data show that the proposed method performs better and benefits more from unsupervised samples than existing methods.

Keywords: Confidence intervals; Data integration; Hypothesis testing; Semi-supervised learning; Structural missing.

1. Introduction

Blockwise missing data arises when we integrate data from multiple modalities, sources, or studies. For instance, the Alzheimer's Disease Neuroimaging Initiative (ADNI) study collects magnetic resonance imaging (MRI), positron emission tomography (PET) imaging, genetics, cerebrospinal fluid (CSF), cognitive tests, and demographic information of patients (Mueller et al., 2005). However, due to the unavailability of the MRI or PET images for




Figure 1: White areas represent missing blocks, while colored areas represent observed blocks. (a) Blockwise missing structure for ADNI data. (b) One blockwise missing example.

some subjects, the biomarkers related to the images can be completely missing for these subjects. As a consequence, when we integrate data from multiple sources and group patients based on their missing patterns, blocks of values could be missing as illustrated in Figure 1 (a), where white areas represent the missing blocks. Multimodality data also appear in modern genomic studies of complex diseases. For example, the Genotype-Tissue Expression (GTEx) study contains RNA-seq gene expression data from over 45 tissues of more than 800 donors (Lonsdale et al., 2013). The gene expression data in the GTEx are also blockwise missing if an tissue sample was unavailable.

Many important scientific questions can be answered through an association or regression analysis. In this case, it is common that, for data sets with blockwise missing covariates, the response variable is also partially missing across the samples. For example, this situation could occur when the outcomes are expensive to collect, such as in electronic health records databases where labeling outcome of each individual is costly and time-consuming (Kohane, 2011; Gronsbell and Cai, 2018). In the GTEx study, samples are collected from only non-diseased tissue samples across individuals (Consortium et al., 2017), implying that the response is only partially observed when we predict gene expression in one tissue using gene expression levels in other tissues. As a consequence, to make the most use of such semi-supervised data sets, it is essential to develop methods that are well adaptive and effectively utilize the extra unsupervised samples for inferring the underlying models.

In this study, we consider a linear regression model

$$y = X^T \beta + \epsilon, \quad (1)$$

where $\beta \in \mathbb{R}^p$ is the regression coefficient vector, which is assumed to be sparse, $X \sim P_X$ is a random vector of regression covariates, and ϵ is a centered sub-Gaussian random variable with variance σ^2 and independent of X . We let the set of observed samples be $\mathcal{D} = \mathcal{D}_1 \cup \mathcal{D}_2$,

where \mathcal{D}_1 is the set of the unsupervised samples, whose response variable is not observable, and \mathcal{D}_2 the set of the supervised samples. All the samples in \mathcal{D} are independently drawn from the regression model (1), before going through certain missing mechanisms. In addition, we assume the covariates of the samples in \mathcal{D} are blockwise missing, and the missing patterns, including the variables associated with each missing block and the numbers of missing blocks, are shared between \mathcal{D}_1 and \mathcal{D}_2 . Our goal is to study the problems of statistical inference for the high-dimensional regression vector β in (1) based on such semi-supervised and blockwise missing observations.

1.1 Related Works

Several methods have been developed recently on blockwise missing data. Yuan et al. (2012) proposed an incomplete multi-source feature learning method that divides blockwise missing data into several learning tasks and builds models for all tasks simultaneously. Xiang et al. (2014) improved the incomplete multi-source feature learning method and propose a unified bi-level learning model that considers not only source-level analysis but also feature-level analysis. Moreover, Yu et al. (2020) proposed to use covariance information from the blockwise missing data to predict a continuous response under the linear regression framework. These methods perform analysis without imputation. There are also methods considering imputation for blockwise missing data. A structured matrix completion proposed by Cai et al. (2016) imputes blockwise missing values via the singular value decomposition. Xue and Qu (2020) proposed a multiple blockwise imputation approach and integrate informative estimating functions based on all the imputations for model selection and prediction. However, none of these existing methods incorporates a partially observed response, or aims for confidence intervals or hypotheses testing of regression models with blockwise missing data.

Concerning statistical inference for high-dimensional regression models under the supervised setting, there are a number of developments based on bias correction of regularized estimators, including Javanmard and Montanari (2014), van de Geer et al. (2014), Zhang and Zhang (2014), Ning and Liu (2017), Javanmard and Montanari (2018), and Neykov et al. (2018), among many others. For inference under the semi-supervised setting, Bellec et al. (2018) proposed a modified Lasso estimator for high-dimensional regression and carefully studied its prediction risk. Zhang and Bradic (2019) studied the semi-supervised inference of the mean and variance of the response with high-dimensional covariates. More recently, Cai and Guo (2020) proposed optimal semi-supervised inference procedures for the quadratic functionals in high-dimensional linear regression, and Deng et al. (2020) studied optimal high-dimensional semi-supervised inference that allows for potential model misspecification. Besides, there are many interesting works focusing on the semi-supervised inference in the classical fixed dimensional setting; see, for example, Wasserman and Lafferty (2008), Azriel et al. (2016), Chakraborty and Cai (2018), Zhang et al. (2019), and the references therein. Nevertheless, none of these methods addresses missing covariates, and, to the best of our knowledge, there are no existing methods that focus on the statistical inference for blockwise missing data in the context of semi-supervised learning.

1.2 Main Contributions

In this study, building upon carefully constructed unbiased estimating equations and a multiple blockwise imputation procedure that account for the structural missing covariates, we propose a computationally efficient semi-supervised sparse estimator for the high-dimensional regression coefficient vector based on a novel multiple blockwise imputed Dantzig selector, and obtain its theoretical properties under mild regularity conditions. Importantly, unlike the existing methods, our method does not require any complete observations across the available samples. In addition, we further develop an innovative semi-supervised projected estimating equation technique that intrinsically achieves bias-correction of the initial sparse estimator, leading to nearly unbiased estimators for the individual regression coefficients that are asymptotically normally distributed under mild conditions. By carefully analyzing these nearly unbiased estimators, asymptotically valid confidence intervals and statistical tests about each regression coefficient can be constructed accordingly. In particular, our theoretical analysis provide important insights on the benefit of the unsupervised samples on the proposed inference procedures, yielding their scalability and adaptivity to the semi-supervised data (see also the discussions after Theorems 1 and 3 for details).

1.3 Notation and Organization

Throughout, for a vector $a = (a_1, \dots, a_n)^\top \in \mathbb{R}^n$, we define the ℓ_p norm $\|a\|_p = (\sum_{i=1}^n |a_i|^p)^{1/p}$, the ℓ_0 norm $\|a\|_0 = \sum_{i=1}^n 1\{a_i \neq 0\}$, and the ℓ_∞ norm $\|a\|_\infty = \max_{1 \leq j \leq n} |a_j|$. For an index set $S \subset \{1, \dots, n\}$, we denote a_S as the subvector of a consisting all the components a_j where $j \in S$. In addition, we let $a_{-j} \in \mathbb{R}^{n-1}$ stand for the subvector of a without the j -th component. For a matrix $A \in \mathbb{R}^{p \times q}$, $\lambda_i(A)$ stands for the i -th largest singular value of A and $\lambda_{\max}(A) = \lambda_1(A)$, $\lambda_{\min}(A) = \lambda_{\min(p,q)}(A)$. For index sets $S_1 \subseteq [1 : p]$ and $S_2 \subseteq [1 : q]$, we denote $A_{S_1 S_2}$ as the submatrix of A consisting of its entries in the rows indexed by S_1 and the columns indexed by S_2 . We denote $\|A\|_\infty = \max_{i,j} |A_{ij}|$. For any positive integer n , we denote the set $\{1, 2, \dots, n\}$ as $[1 : n]$. For sequences $\{a_n\}$ and $\{b_n\}$, we write $a_n = o(b_n)$, $a_n \ll b_n$ or $b_n \gg a_n$ if $\lim_n a_n/b_n = 0$, and write $a_n = O(b_n)$, $a_n \lesssim b_n$ or $b_n \gtrsim a_n$ if there exists a constant C such that $a_n \leq Cb_n$ for all n . We write $a_n \asymp b_n$ if $a_n \lesssim b_n$ and $a_n \gtrsim b_n$. For a set A , we denote $|A|$ as its cardinality. Lastly, C, C_0, C_1, \dots are absolute constants.

The rest of the paper is organized as follows. In Section 2, we introduce our proposed semi-supervised inference procedures for parameter estimation, constructing confidence intervals, and hypotheses testing based on multiple blockwise imputation and a novel bias-correction technique. In Section 3, the theoretical justifications of the proposed inference procedures are provided. The empirical performance of the proposed methods as well as its advantages over alternative methods are assessed in Section 4. In Section 5, we apply the proposed methods to the Alzheimer's Disease Neuroimaging Initiative data sets to identify important biomarkers associated with the Alzheimer's disease. We conclude this article with a discussion in Section 6.

2. Semi-Supervised Inference with Multiple Blockwise Imputation

2.1 Estimation based on Multiple Blockwise Imputed Dantzig Selector

We start with a description of the multiple blockwise imputation procedure. This procedure is able to use more information from incomplete samples or cases than the traditional single regression imputation, which imputes missing values via regression models using all the observed variables as the predictors (Baraldi and Enders, 2010; Zhang, 2016; Campos et al., 2015). Specifically, we assume there are R groups of samples, whose missing covariates are the same within each group, and for any $r \in [1 : R]$, we let $\mathcal{H}(r) \subseteq \mathcal{D}$ be an index set of the samples in Group r . We also let $\mathcal{G}(r) \subseteq [1 : R]$ be the index set of the groups where missing variables of Group r and variables in at least one of the other sources are observed. For example, when there are three sources of data with $R = 5$ missing groups as shown in Figure 1 (b), then $\mathcal{G}(2) = \{1, 3, 4\}$. Group 5 is not in $\mathcal{G}(2)$, since it does not contain any information for variables in Source 3 which are missing in Group 2. We assume that $|\mathcal{G}(r)| \geq 1$ for each $r \in [1 : R]$, implying that each covariate is observed in at least one group.

Intuitively, we can predict the expected values of the missing variables for the samples in $\mathcal{H}(r)$ using the dependence structure between the missing variables and the non-missing variables as yielded by the samples of the groups in $\mathcal{G}(r)$ where both types of variables are observed. In other words, for any $r \in [1 : R]$, $i \in \mathcal{H}(r)$ and any $k \in \mathcal{G}(r)$, we can impute each of the missing variables X_{ij} by $E(X_{ij}|X_{iJ(r,k)})$ where $J(r,k) \subseteq [1 : p]$ is an index set of covariates that are observed in both Groups r and k . Throughout, for each $r \in [1 : R]$ and $i \in \mathcal{H}(r)$, we define $X_i^{(k)} = (X_{i1}^{(k)}, \dots, X_{ip}^{(k)})^T$ as the imputed random vector for sample i according to Group $k \in \mathcal{G}(r)$, so that $X_{ij}^{(k)} = E(X_{ij}|X_{iJ(r,k)})$ if the j -th covariate is missing in the i -th sample X_i , otherwise $X_{ij}^{(k)} = X_{ij}$.

Oftentimes, the conditional expectation $E(X_{ij}|X_{iJ(r,k)})$ can be achieved by fitting a linear regression model between X_{ij} and the random vector $X_{iJ(r,k)}$ based on the samples in Group k . To account for high-dimensionality, we consider the Lasso estimator (Tibshirani, 1996) defined as

$$\hat{\gamma}_{j,J(r,k)} = \arg \min_{\gamma \in \mathbb{R}^{|J(r,k)|}} \left\{ \frac{1}{|\mathcal{H}(k)|} \sum_{i \in \mathcal{H}(k)} (X_{ij} - X_{iJ(r,k)}^T \gamma)^2 + \tau \|\beta\|_1 \right\}, \quad (2)$$

where $\tau > 0$ is a tuning parameter. Then we can approximate the imputed variable $X_{ij}^{(k)} = E(X_{ij}|X_{iJ(r,k)})$ by $\hat{\gamma}_{j,J(r,k)}^T X_{iJ(r,k)}$. Note that $\mathcal{G}(r)$ could contain multiple elements, such as $|\mathcal{G}(2)| = 3$ under the example in Figure 1 (b). Consequently, for each missing variable X_{ij} associated to Group r , there could be multiple imputed values $\hat{\gamma}_{j,J(r,k)}^T X_{iJ(r,k)}$ associated with different $k \in \mathcal{G}(r)$. In the following, for each $r \in [1 : R]$ and $i \in \mathcal{H}(r)$, we define $\hat{X}_i^{(k)} = (\hat{X}_{i1}^{(k)}, \dots, \hat{X}_{ip}^{(k)})^T$ as the actual imputed observations of sample i based on Group $k \in \mathcal{G}(r)$, where $\hat{X}_{ij}^{(k)} = \hat{\gamma}_{j,J(r,k)}^T X_{iJ(r,k)}$ if the j -th covariate is missing in the i -th sample X_i , and otherwise $\hat{X}_{ij}^{(k)} = X_{ij}$.

With the above preparations, we are ready to introduce our multiple-imputed Dantzig selector estimator of the coefficient vector β , whose construction is based on the imputed observations of the supervised samples in \mathcal{D}_2 . The intuition of the construction of our

proposed estimator comes from the following observation. For any $r \in [1 : R]$, $k \in \mathcal{G}(r)$ and $i \in \mathcal{D}_2$, we consider

$$h_{irk}(\beta) = I(\xi_i = r)\{y_i - (X_i^{(k)})^T \beta\} \cdot X_{ia(k)}^{(k)}, \quad (3)$$

where ξ_i is a random group label taking values in $[1 : R]$, $a(k) \subseteq [1 : p]$ is the index set of the observed covariates in Group k , $X_{ia(k)}^{(k)}$ is a sub-vector of $X_i^{(k)}$ consisting of all the covariates observed in Group k . Under the linear regression model, whenever ξ_i is independent of all the covariates (missing completely at random), or only depends on the observed covariates (missing at random), it can be shown that $E\{h_{irk}(\beta)\} = 0$ (Xue and Qu, 2020). Hence, a system of unbiased estimating equations can be defined as

$$g(\beta) = \frac{1}{|\mathcal{D}_2|} \sum_{i \in \mathcal{D}_2} \begin{bmatrix} \hat{\theta}_1^{-1} h_{i1}(\beta) \\ \vdots \\ \hat{\theta}_R^{-1} h_{iR}(\beta) \end{bmatrix} = 0, \quad (4)$$

where $\hat{\theta}_r = |\mathcal{D}_2 \cap \mathcal{H}(r)| / |\mathcal{D}_2|$ is an estimate of observed rate for the r -th group among \mathcal{D}_2 , and $h_{ir}(\beta) = (h_{ir1}(\beta)^T, \dots, h_{ir|\mathcal{G}(r)}(\beta)^T)^T$ is a vector combining the components of the vectors in $\{h_{irk}(\beta)\}_{k \in \mathcal{G}(r)}$. In particular, $g(\beta)$ is a vector of dimension $M_g = \sum_{r=1}^R \sum_{k \in \mathcal{G}(r)} |a(k)|$.

However, the random vectors $X_i^{(k)}$ required by (3) and (4) are not fully observed. Instead, we use the imputed observations $\hat{X}_i^{(k)}$ as an approximation. Specifically, we define the imputed counterpart of $h_{irk}(\beta)$ as

$$\hat{h}_{irk}(\beta) = I(\xi_i = r)\{y_i - (\hat{X}_i^{(k)})^T \beta\} \cdot \hat{X}_{ia(k)}^{(k)}, \quad (5)$$

and define the imputed estimating function as

$$g_n(\beta) = \frac{1}{|\mathcal{D}_2|} \sum_{i \in \mathcal{D}_2} \begin{bmatrix} \hat{\theta}_1^{-1} \hat{h}_{i1}(\beta) \\ \vdots \\ \hat{\theta}_R^{-1} \hat{h}_{iR}(\beta) \end{bmatrix}, \quad (6)$$

where $n = |\mathcal{D}_2|$ and $\hat{h}_{ir}(\beta) = (\hat{h}_{ir1}(\beta)^T, \dots, \hat{h}_{ir|\mathcal{G}(r)}(\beta)^T)^T$ for each $r \in [1 : R]$. Finally, respecting the underlying sparsity of the coefficient vector β , we define the proposed estimator as

$$\hat{\beta} = \arg \min_{\beta \in \mathbb{R}^p} \|\beta\|_1, \quad \text{subject to } \|g_n(\beta)\|_\infty \leq \lambda, \quad (7)$$

where $\lambda > 0$ is a tuning parameter. In Section 3, we show that $\hat{\beta}$ is a consistent estimator and obtain its rate of convergence under mild regularity conditions.

2.2 Projected Estimating Equations and the Bias-Corrected Estimator

Although the proposed estimator $\hat{\beta}$ performs well in terms of point estimation, it is biased and cannot be directly adopted for developing powerful inference procedures such as confidence intervals and statistical tests. In this section, based on the idea of projected

estimation equations, we propose a novel bias-corrected estimator, which is asymptotically normally distributed around the true coefficients.

From the imputed estimating function $g_n(\beta)$, we define $g_n^*(\beta)$ be the subvector of $g_n(\beta)$ where we replace each $\widehat{h}_{irk}(\beta)$ in (5) by its subvector

$$\widehat{h}_{irk}^*(\beta) = I(\xi_i = r)\{y_i - (\widehat{X}_i^{(k)})^T \beta\} \cdot X_{iJ(r,k)}^{(k)} = I(\xi_i = r)\{y_i - (\widehat{X}_i^{(k)})^T \beta\} \cdot X_{iJ(r,k)}. \quad (8)$$

By considering the partial estimating function $g_n^*(\beta)$ rather than the complete function $g_n(\beta)$, the bias of the resulting estimators due to the imputation step is significantly reduced. In addition, we define

$$W_n = \text{diag} \left\{ \left(\frac{|\mathcal{D}_2|^2}{|\mathcal{D}_2 \cap \mathcal{H}(r)|^2} \sum_{i \in \mathcal{D}_2} I\{\xi_i = r\} X_{iJ(r,k)} X_{iJ(r,k)}^T \right)_{1 \leq r \leq R, k \in \mathcal{G}(r)} \right\}, \quad (9)$$

which is a block-diagonal matrix consisting of the block sub-matrices

$$\frac{|\mathcal{D}_2|^2}{|\mathcal{D}_2 \cap \mathcal{H}(r)|^2} \sum_{i \in \mathcal{D}_2} I\{\xi_i = r\} X_{iJ(r,k)} X_{iJ(r,k)}^T,$$

ordered first by $r \in [1 : R]$ and then by $k \in \mathcal{G}(r)$. Additionally, we define

$$G_n = \frac{\partial}{\partial \beta} g_n^*(\beta) = \frac{1}{|\mathcal{D}|} \sum_{i \in \mathcal{D}} \begin{bmatrix} \widehat{\theta}_1^{-1} \partial \widehat{h}_{i1}^*(\beta) / \partial \beta \\ \vdots \\ \widehat{\theta}_R^{-1} \partial \widehat{h}_{iR}^*(\beta) / \partial \beta \end{bmatrix}, \quad (10)$$

where for each $r \in [1 : R]$, we have $\partial \widehat{h}_{ir}^*(\beta) / \partial \beta \in \mathbb{R}^{m'_r \times p}$ with $m'_r = \sum_{k \in \mathcal{G}(r)} |J(r, k)|$, which consists of submatrices $\{I(\xi_i = r) X_{iJ(r,k)} (\widehat{X}_i^{(k)})^T\}_{k \in \mathcal{G}(r)}$ combined by row. Then, for any given $j \in [1 : p]$, we define the projected estimating function as

$$\widehat{S}_j(\beta) = \widehat{v}_j^T g_n^*(\beta), \quad (11)$$

with the projection vector \widehat{v}_j being the solution to the following optimization problem

$$\widehat{v}_j = \arg \min v^T W_n v, \quad \text{subject to } \|v^T G_n - e_j^T\|_\infty \leq \lambda', \quad (12)$$

where $\lambda' > 0$ is a tuning parameter, $e_j \in \mathbb{R}^p$ has 1 as its j -th element and 0 otherwise.

We set $\widehat{\beta}_j^* = (\widehat{\beta}_1, \dots, \widehat{\beta}_{j-1}, \beta_j, \widehat{\beta}_{j+1}, \dots, \widehat{\beta}_p)^T$, then, the bias-corrected estimator $\widetilde{\beta}_j$ of β_j is defined as the root of

$$\widehat{S}_j(\widehat{\beta}_j^*) = 0, \quad (13)$$

treated as a univariate equation of the scalar variable β_j . As a result, if we denote $\widetilde{\beta}_j^* = (\widehat{\beta}_1, \dots, \widehat{\beta}_{j-1}, \widetilde{\beta}_j, \widehat{\beta}_{j+1}, \dots, \widehat{\beta}_p)^T$, by Taylor expansion, we have

$$\begin{aligned} 0 &= \widehat{S}_j(\widetilde{\beta}_j^*) = \widehat{v}_j^T g_n^*(\widetilde{\beta}_j^*) + \widehat{v}_j^T G_n e_j \cdot (\widetilde{\beta}_j - \beta_j) \\ &= \widehat{v}_j^T g_n^*(\beta) + \widehat{v}_j^T G_n (\widehat{\beta}_j^* - \beta) + \widehat{v}_j^T G_n e_j \cdot (\widetilde{\beta}_j - \beta_j), \end{aligned} \quad (14)$$

which can be rewritten as

$$\tilde{\beta}_j - \beta_j = - \underbrace{\frac{\hat{v}_j^T g_n^*(\beta)}{\hat{v}_j^T G_n e_j}}_{\text{Stochastic Error}} - \underbrace{\frac{\hat{v}_j^T G_n (\hat{\beta}_j^* - \beta)}{\hat{v}_j^T G_n e_j}}_{\text{Bias}}. \quad (15)$$

The estimation error $\tilde{\beta}_j - \beta_j$ is decomposed into two parts. In particular, it can be shown that, the first term in (26) is the stochastic error, which is asymptotically normal with variance $\hat{v}_j^T W_n \hat{v}_j$, while the second term is the bias term, which can be bounded by

$$\left| \frac{\hat{v}_j^T G_n (\hat{\beta}_j^* - \beta)}{\hat{v}_j^T G_n e_j} \right| \leq \frac{\|(\hat{v}_j^T G_n)_{-j}\|_\infty \|(\hat{\beta}_j^* - \beta)_{-j}\|_1}{1 - |\hat{v}_j^T G_n e_j - 1|}. \quad (16)$$

Intuitively, the projected estimating function (11) is carefully constructed such that the bias of $\tilde{\beta}_j$ is dominated by the stochastic error. In other words, the constrained optimization problem (12) is rooted in the bias–variance trade-off: it aims to find a projection vector \hat{v}_j that simultaneously controls $\|(\hat{v}_j^T G_n)_{-j}\|_\infty$ and $|\hat{v}_j^T G_n e_j - 1|$ in (16), so that the bias term in (26) is negligible with respect to the stochastic error, while minimizing the variance of the stochastic error for the sake of statistical efficiency.

We emphasize that, although the imputed estimating function $g_n^*(\beta)$, like $g_n(\beta)$ defined in (6), is based on the imputed observations of the supervised data \mathcal{D}_2 , the projection vector \hat{v}_j defined in (12) is based on all the available samples \mathcal{D} , to achieve better accuracy.

2.3 Confidence Intervals and Statistical Tests

Concerning the above bias-corrected estimator $\tilde{\beta}_j$, we show in Section 3 that, conditional on the imputed covariates, under mild regularity conditions, it is asymptotically normally distributed, whose variance is $s_j^2/|\mathcal{D}_2|^2$, where

$$s_j^2 = \sigma^2 \sum_{i \in \mathcal{D}_2} \sum_{k \in \mathcal{G}(r), 1 \leq r \leq R} \frac{|\mathcal{D}_2|^2}{|\mathcal{D}_2 \cap \mathcal{H}(r)|^2} I\{\xi_i = r\} (\hat{v}_{j,rk}^T X_{iJ(r,k)})^2, \quad (17)$$

and for each $r \in [1 : R]$ and $k \in \mathcal{G}(r)$, $\hat{v}_{j,rk} \in \mathbb{R}^{|\mathcal{J}(r,k)|}$ is the subvector of the projection vector \hat{v}_j corresponding to the estimating functions in $g_n^*(\beta)$ associated to Group $k \in \mathcal{G}(r)$. As a consequence, for given $j \in [1 : p]$, suppose $\hat{\sigma}^2$ is a reasonably good estimator of the noise level σ^2 , then, an asymptotically $(1 - \alpha)$ -level confidence interval for β_j can be constructed as

$$\text{CI}_\alpha(\beta_j) = \left[\tilde{\beta}_j - \frac{z_{\alpha/2} \hat{s}_j}{|\mathcal{D}_2|}, \tilde{\beta}_j + \frac{z_{\alpha/2} \hat{s}_j}{|\mathcal{D}_2|} \right], \quad (18)$$

where

$$\hat{s}_j^2 = \hat{\sigma}^2 \sum_{i \in \mathcal{D}_2} \sum_{k \in \mathcal{G}(r), 1 \leq r \leq R} \frac{|\mathcal{D}_2|^2}{|\mathcal{D}_2 \cap \mathcal{H}(r)|^2} I\{\xi_i = r\} [\hat{v}_{j,rk}^T X_{iJ(r,k)}]^2, \quad (19)$$

and $z_{\alpha/2} = \Phi^{-1}(1 - \alpha/2)$ is the upper $\alpha/2$ -quantile of the standard normal distribution. In particular, a simple estimator for σ^2 can be defined as

$$\hat{\sigma}^2 = \frac{1}{\sum_{k \in \mathcal{G}(r), 1 \leq r \leq R} |\mathcal{D}_2 \cap \mathcal{H}(r)|} \sum_{i \in \mathcal{D}_2} \sum_{k \in \mathcal{G}(r), 1 \leq r \leq R} \left[I(\xi_i = r) \{y_i - (\hat{X}_i^{(k)})^T \hat{\beta}\} \right]^2, \quad (20)$$

which is used in the numerical analysis in this paper.

In parallel with the above confidence interval, we can also construct an asymptotically valid statistical test for the null hypothesis $H_0 : \beta_j = b_j$ for any $b_j \in \mathbb{R}$. Specifically, define the test statistic

$$T_j = |\mathcal{D}_2|(\tilde{\beta}_j - b_j)/\hat{s}_j. \quad (21)$$

Then an asymptotically α -level test is that we reject H_0 whenever $|T_j| > z_{\alpha/2}$.

With these component-wise test statistics, in light of the existing works on the supervised inference for high-dimensional regression such as Javanmard and Javadi (2019), Ma et al. (2020) and Zhang et al. (2020), among many others, one can also construct tests for the global null hypothesis $H_0 : \beta = 0$, and the multiple simultaneous hypotheses $H_{0j} : \beta_j = 0, j = 1, \dots, p$. For example, to test the global null hypothesis, one could adopt the maximum-type test statistic $M = \max_{1 \leq j \leq p} T_j^2$, and compare its empirical values to the quantile of a Gumbel distribution given in Theorem 1 of Ma et al. (2020); to test the simultaneous null hypotheses while controlling for false discovery rates, one could apply the modified Benjamini–Hochberg procedure described in Javanmard and Javadi (2019) and Ma et al. (2020) when the design covariates are weakly correlated, or the Benjamini–Yekutieli procedure (Benjamini and Yekutieli, 2001) if the design covariates are arbitrarily correlated.

3. Theoretical Justifications

This section provides theoretical justifications of the proposed inference procedures by studying the properties of the proposed estimator $\hat{\beta}$ and its bias-corrected counterpart $\tilde{\beta}$. Throughout, we assume the missing patterns, or the missing variables in each group $r \in [1 : R]$, are given, but allow the random group label ξ_i to be dependent on the covariates. For technical reasons, we assume for simplicity that the multiple blockwise imputation step (2) is performed using the unsupervised data \mathcal{D}_1 and a fixed portion of the supervised data \mathcal{D}_2 , whereas the construction of estimating functions $g_n(\beta)$ and $g_n^*(\beta)$ is based on the imputed observations of the other portion of the supervised data. In practice, however, this sample splitting procedure is not needed, and the proposed method works well empirically when the multiple blockwise imputation step is based on all the available samples; see our numerical results in Sections 4 and 5, as well as discussion in Section 6.

For any $r \in [1 : R]$, $k \in \mathcal{G}(r)$ and $i \in \mathcal{D}$, we define $\Sigma^{(r,k)} = E[I\{\xi_i = r\}X_i^{(k)}(X_i^{(k)})^T] \in \mathbb{R}^{p \times p}$ and denote $m(k) \subseteq [1 : p]$ as the index set of the missing covariates associated to Group k . Recall that $a(k)$ is the index set of the observed covariates in Group k , and $J(r, k)$ is the index set of the covariates observed in both Groups r and k . We also denote $N_r = |\mathcal{D}_1 \cap \mathcal{H}(r)|$, $n_r = |\mathcal{D}_2 \cap \mathcal{H}(r)|$, and $N = |\mathcal{D}_1|$. Regarding the missing patterns, we assume

(A1) R is a finite integer, and for all $r \in [1 : R]$ and $k \in \mathcal{G}(r)$, we have $|a(r)|/p, J(r, k)/p \in [C_1, C_2]$ for some constants $0 < C_1 < C_2 < 1$; and

(A2) For all $r \in [1 : R]$, we have $P(n_r/n, N_r/N \in [c_1, c_2]) \geq 1 - p^{-c}$ for some constants $0 < c_1 < c_2 < 1$ and $c > 0$.

These assumptions are mild as they essentially ensure the missing patterns are finite and balanced. For the design covariates, and the regression coefficient vector β , we assume

(A3) For $i \in \mathcal{D}$, $X_i \sim_{i.i.d.} N(0, \Sigma)$, where $C^{-1} \leq \lambda_{\min}(\Sigma) \leq \lambda_{\max}(\Sigma) \leq C$ for some absolute constant $C > 1$, and each row of $\Sigma^{-1} = (\omega_{ij})$ is s -sparse, in the sense that

$\max_{1 \leq i \leq p} \sum_{j=1}^p 1\{\omega_{ij} \neq 0\} \leq s$; and

(A4) β is s -sparse in the sense that $\|\beta\|_0 \leq s$, and $\|\beta\|_2$ is bounded by an absolute constant.

The following theorem concerns the convergence rates of the estimator $\hat{\beta}$ defined by (7).

Theorem 1 *Suppose (A1) to (A4) hold, $p \lesssim \{\min(N, n)\}^\eta$ for some constant $\eta > 0$, $s \ll \min\{\sqrt{n/\log p}, (n + N)/\log p\}$, and there exists some $r \in [1 : R]$, $k_1, k_2 \in \mathcal{G}(r)$ and some constant $c_0 > 0$, such that $m(k_1) \subseteq a(k_2)$, $m(k_2) \subseteq a(k_1)$, and $\lambda_{\min}(\Sigma_{a(k), a(k)}^{(r, k)}) \geq 7c_0 > c_0 \geq \lambda_{\max}(\Sigma_{a(k), m(k)}^{(r, k)})$ for $k = k_1, k_2$. Then, for $\lambda \asymp \sqrt{\log p/n} + s\sqrt{\log p/(n + N)}$, and sufficiently large (n, p) , it holds that, $\|\hat{\beta} - \beta\|_1 \lesssim s\lambda$ and $\|\hat{\beta} - \beta\|_2 \lesssim s^{1/2}\lambda$, with probability at least $1 - p^{-c}$, for some absolute constant $c > 0$.*

Some remarks about Theorem 1 are in order. Firstly, as indicated by assumptions (A1) and (A2), unlike many existing methods such as Cai et al. (2016), Kundu et al. (2019), and Yu et al. (2020), our method does not require availability of samples with complete observations. In other words, each sample in the data set could be partially missing, as long as the overall missing patterns are finite and asymptotically balanced. Secondly, our theorem shows that the rate of convergence of $\hat{\beta}$ under the ℓ_2 -norm is bounded by $\sqrt{s \log p/n} + s^{3/2} \sqrt{\log p/(n + N)}$. Compared to the minimax optimal rates $\sqrt{s \log p/n}$ for estimating β with complete observations of n samples (Verzelen, 2012), the above rate of $\hat{\beta}$ is sharp up to an additional $s^{3/2} \sqrt{\log p/(n + N)}$, which reflects the extra approximation error due to the multiple blockwise imputation step. Thirdly, since the above approximation error is partially driven by the number of the unsupervised samples N , the estimation of β using $\hat{\beta}$ could potentially benefit from additional unsupervised data or the improved imputation quality. In particular, when the number of unsupervised samples is sufficiently large, namely, $N \gtrsim s^2 n$, the approximation error is negligible comparing to $\sqrt{s \log p/n}$, and the estimator $\hat{\beta}$ achieves the minimax optimal rate when complete observations of n samples are available.

Moreover, one theoretical condition in Theorem 1 is the existence of two groups $\{k_1, k_2\} \subseteq \mathcal{G}(r)$ whose missing patterns are complementary, that is, $m(k_1) \subseteq a(k_2)$ and $m(k_2) \subseteq a(k_1)$, which is rather mild. The eigenvalue condition $\lambda_{\min}(\Sigma_{a(k), a(k)}^{(r, k)}) \geq 7c_0 > c_0 \geq \lambda_{\max}(\Sigma_{a(k), m(k)}^{(r, k)})$ requires the existence of a pair of groups $(r, k) \in [1 : R] \times \mathcal{G}(r)$ whose imputed random vector $X_i^{(k)}$ with $i \in \mathcal{H}(r)$ having its observed variables in $X_{ia(k)}^{(k)} = X_{ia(k)}$ individually informative in the sense that they are not too collinear with each other or the imputed variables $X_{ij}^{(k)}$ for $j \in m(r)$. As an example, as shown in Appendix E, this condition is satisfied in a simple case when the missing is completely at random and $\|(\Sigma^{-1})_{J(r, k), m(r)}\| \rightarrow 0$, that is, when there exists a pair $(r, k) \in [1 : R] \times \mathcal{G}(r)$ such that the missing variables in $m(r)$ are nearly conditionally independent of the observed variables in $J(r, k)$. Finally, for the selection of the tuning parameter λ , in practice, we recommend the cross validation, whose empirical performance is evaluated in Section 4.

The proof of Theorem 1, however, is involved, and very different from the existing results concerning the standard Dantzig selector or the Lasso estimator for the linear regression model (Candes and Tao, 2007; Bickel et al., 2009; Negahban et al., 2010). The complete proof can be found in Appendix B. In particular, as a key component of our theoretical analysis, a novel restricted singular value inequality is developed, which accounts for the

multiple blockwise-imputed samples, and plays a similar role as the restricted eigenvalue condition (Bickel et al., 2009; Raskutti et al., 2010), or the restricted strong convexity property (Negahban et al., 2010; Negahban and Wainwright, 2012) in the previous literature. The inequality, summarized as the following proposition, could be of independent interest. The proof is provided in Appendix D.4.

Proposition 2 *Under the conditions of Theorem 1, there exists some $r \in [1 : R]$ and $k \in \mathcal{G}(r)$, such that, with probability at least $1 - p^{-c}$ for some absolute constant $c > 0$,*

$$\inf_{\substack{\|u\|_2=1, u \in E_s \\ \|u_{a(k)}\|_2 \geq 1/2}} \left| n_r^{-1} \sum_{i=1}^n I\{\xi_i = r\} (u_{a(k)} / \|u_{a(k)}\|_2)^T \widehat{X}_{ia(k)}^{(k)} (\widehat{X}_i^{(k)})^T u \right| \geq c_0 \quad (22)$$

for some constant $c_0 > 0$ and $E_s(p) = \{\delta \in \mathbb{R}^p : \|\delta\|_2 = 1, \|\delta_{S^c}\|_1 \leq \|\delta_S\|_1, |S| \leq s\}$.

Our next theorem establishes the asymptotic normality of the bias-corrected estimator $\tilde{\beta}_j$, which in turn implies the asymptotic validity of the confidence intervals and the statistical tests proposed in Section 2.3.

Theorem 3 *Suppose the conditions of Theorem 1 hold. If, in addition, for $G(\beta) = \partial g^*(\beta) / \partial \beta$ with $g^*(\beta)$ being the population counterpart of $g_n^*(\beta)$, we have $\lambda_{\min}(E\{G(\beta)\}) \geq c$ for some absolute constant $c > 0$, $\mathcal{X} \asymp \sqrt{s \log p / (n + N)}$ and $s \ll \min\{(n + N)^{1/3} / \log^{2/3} p, (n + N)^{2/5} / (n \log^2 p)^{1/5}\}$, then, for each $j \in [1 : p]$, it holds that*

$$n(\tilde{\beta}_j - \beta_j) / s_j = AB + D, \quad (23)$$

where $A \rightarrow 1$ and $D \rightarrow 0$ in probability, and $B|\widehat{X} \rightarrow N(0, 1)$ in distribution, in which $\widehat{X} = \{\widehat{X}_i^{(k)}\}_{i \in \mathcal{D}_2}$ is the set of all the imputed observations.

In the above theorem, a regularity condition on the smallest singular value of the expected gradient $E\{G(\beta)\}$ is assumed, which ensures the existence of a projection vector \hat{v}_j satisfying the constraint in (12) with high probability. The additional condition on the sparsity $s \ll \min\{(n + N)^{1/3} / \log^{2/3} p, (n + N)^{2/5} / (n \log^2 p)^{1/5}\}$ essentially ensures that the remaining bias D in (23) is diminishing; it becomes weaker when the number of the unsupervised samples increases. This suggests that the debiased estimators $\tilde{\beta}_j$ can benefit from the extra unsupervised data not only in terms of improved imputation quality as discussed earlier, but also the accuracy of the asymptotic normality. Similar phenomena have also been observed by Cai and Guo (2020) and Deng et al. (2020) in the context of semi-supervised inference for the quadratic and the linear functionals without missing covariates.

4. Simulation

In this section, we provide simulation studies to compare the proposed method with existing methods, including the debiased Lasso method (Javanmard and Montanari, 2014) with complete cases, the Lasso projection method (van de Geer et al., 2014; Zhang and Zhang, 2014) with complete cases, the debiased Lasso method with single regression imputation,

	Source 1	Source 2	Source 3
Group 1			
Group 2			
Group 3			
Group 4			

Figure 2: Blockwise missing structure used for simulation.

and the Lasso projection method with the single regression imputation. Here, “single regression imputation” refers to predicting missing values via linear regressions with observed variables as predictors (Baraldi and Enders, 2010; Zhang, 2016; Campos et al., 2015).

For implementation of the proposed method, we use the “Rglpk” package (<https://cran.r-project.org/web/packages/Rglpk/index.html>) and the “osqp” package (<https://cran.r-project.org/web/packages/osqp/index.html>) in R to solve the linear programming problem in (7) and the quadratic programming problem in (12), respectively. The parameter λ is determined by 10-fold cross validation. We let $\lambda' = 0.1(\log p/n)^{1/2}$ and scale it up if there exists no solution to the the quadratic programming problem in (12). We use the R codes in <https://web.stanford.edu/~montanar/sslasso/> to implement the debiased Lasso method. For the Lasso projection method, we apply the “hdi” package (<https://cran.r-project.org/web/packages/hdi/index.html>) in R. We simulate $X_i, i = 1, \dots, n$, independent from a multivariate Gaussian distribution with mean 0 and a block-diagonal covariance matrix Σ , and generate $y_i, i = 1, \dots, n$, from (1) where $\epsilon_i \sim_{i.i.d.} N(0, 1)$. The relevant covariates share the same signal strength β_s .

We compare all the methods in the following three settings. In Settings 1 and 2, we assume informative missing with data from three sources and four missing pattern groups as shown in Figure 2. In Setting 3, we assume missing at random, and add one more data source where variables are all observed for each subject. In each setting, we construct confidence intervals for a relevant covariate with confidence level 95%, and evaluate each method using the coverage rate and average length of the confidence intervals based on 250 replications. Let S denote the number of data sources, p_i denote the number of total covariates in the i -th data sources, and q_i denote the number of relevant covariates in the i -th data sources for $1 \leq i \leq S$.

Setting 1 Let $n = 150$, $p = 200$, $q = 9$, $R = 4$, $S = 3$, $N = 300$, $\beta_s = 0.2$, $n_1 = 30$, $n_2 = 70$, $n_3 = n_4 = 25$, $p_1 = 115$, $p_2 = 45$, $p_3 = 40$, $q_1 = 5$, $q_2 = q_3 = 2$, and $\Sigma = \text{diag}\{I_{p_1}, A\}$, where A is a $(p_2 + p_3) \times (p_2 + p_3)$ exchangeable matrix with diagonal elements 1 and off-diagonal elements ρ . We let $\rho = 0.1$ or 0.3. Samples are sequentially randomly assigned into the complete case group with probabilities proportional to $\exp(-10y_i)$

for $1 \leq i \leq n + N$. Otherwise, they are uniformly assigned to the other three groups. The response is observed for uniformly selected one-third of samples in each missing group.

Setting 2 The same as Setting 1 except that $p = 700$ and $p_1 = 615$.

Setting 3 The same as Setting 1 except that $n = 120$, $S = 4$, $N = 750$, $n_1 = 15$, $n_2 = 65$, $n_3 = n_4 = 20$, $p_2 = 40$, $p_4 = 5$, $q_1 = 4$, $q_4 = 1$, and $\Sigma = \text{diag}\{I_{p_1}, A, I_{p_4}\}$. Samples are sequentially randomly assigned into the complete case group with probabilities proportional to $\exp(-10d_i)$ for $1 \leq i \leq n + N$, where d_i is the sum of the i -th samples of covariates in the fourth source of data. The missing pattern in Sources 1–3 are the same as that in Figure 2, while covariates in Source 4 are all observed. The response is observed for uniformly selected one-sixth of samples in each missing group.

The results of Settings 1–3 are provided in Table 1. The proposed method outperforms existing methods across all the settings in terms of coverage rate. In Setting 1, 80% observations are missing and the missingness is informative. As shown in Table 1, the coverage of the proposed method is at least 42.0% more than that of others methods when $\rho = 0.1$. In Setting 2, we consider more potential predictors to mimic the the Alzheimer’s Disease Neuroimaging Initiative data in Section 5. The proposed method still produces the largest coverage rate than existing methods. Moreover, when $\rho = 0.1$, the coverage rate of the proposed method is 94.4% which is close to 95%. In Setting 3, we simulate more unsupervised samples and consider missing at random mechanism. As shown in Table 1, the proposed method and the Lasso projection method with complete cases both achieve the 95% coverage. However, the average length of confidence intervals of the proposed method is much smaller than that of the Lasso projection method with complete cases, indicating that confidence intervals of the proposed method are more accurate.

5. Real data application

In this section, we apply the proposed method to the Alzheimer’s Disease Neuroimaging Initiative (ADNI) data set which contains multisource measurements: magnetic resonance imaging (MRI), positron emission tomography (PET) imaging, gene expression, and cognitive tests (Mueller et al., 2005). Among these measurements, the mini-mental state examination is often used for diagnosis of the Alzheimer’s Disease (Chapman et al., 2016). It is therefore important to identify the imaging and gene expression features that are associative and are predictive to the score of the mini-mental state examination. To identify biomarkers associated with the Alzheimer’s Disease, we use the score of the mini-mental state examination as our response variable, and treat MRI, PET, and gene expression variables as predictors. Specifically, the MRI variables contain volumes, surface areas, average cortical thickness, and standard deviation in cortical thickness of regions of interest in brain, which are extracted from the MRIs by the Center for Imaging of Neurodegenerative Diseases at the University of California, San Francisco. To mitigate bias due to different head sizes, we normalize the MRI variables via dividing region volumes, surface areas and cortical thicknesses by the whole brain volume, the total surface area, and the mean cortical thickness of each subject, respectively (Zhou et al., 2014; Kang et al., 2019). The PET variables are standard uptake value ratios of brain regions of interest, which represent metabolic activity

Method	$\rho = 0.1$		$\rho = 0.3$	
	Coverage rate	Average length	Coverage rate	Average length
Setting 1				
Proposed	0.920	0.581	0.876	0.560
DL-CC	0.264	0.274	0.248	0.291
LP-CC	0.636	0.423	0.644	0.429
DL-SI	0.036	0.140	0.036	0.135
LP-SI	0.648	0.326	0.732	0.362
Setting 2				
Proposed	0.944	0.931	0.908	0.881
DL-CC	0.000	0.004	0.000	0.008
LP-CC	0.628	0.428	0.668	0.443
DL-SI	0.036	0.146	0.016	0.140
LP-SI	0.804	0.375	0.800	0.380
Setting 3				
Proposed	0.956	0.864	0.968	0.824
DL-CC	0.264	0.209	0.356	0.271
LP-CC	0.952	1.201	0.968	1.255
DL-SI	0.388	0.301	0.376	0.283
LP-SI	0.740	0.387	0.804	0.420

Table 1: Simulation results of Settings 1–3. DL-CC:the debiased Lasso method with complete cases. LP-CC: the Lasso projection method with complete cases. DL-SI: the debiased Lasso method with single regression imputation. LP-SI: the Lasso projection method with single regression imputation.

and are provided by the Jagust Lab at the University of California, Berkeley. Gene expression levels at different probes are contributed by Bristol-Myers Squibb laboratories from blood samples of ADNI participants.

Although the ADNI is a longitudinal study, we focus on data collected in the second phase of the ADNI study (ADNI-2) at month 48 in this real data application. In total, there are 212 samples, 267 MRI variable, 113 PET variables, and 49386 gene expression variables. The blockwise missingness emerges when we combine data from MRI, PET and gene expression. The missing pattern structure is the same as that in Figure 2 with four groups and 69 complete observations. Due to relatively small sample sizes, we first screen gene expression variables via marginal correlations according to the sure independence screening (Fan and Lv, 2008) and retain 300 gene expression variables. Due to the missingness, we compute the marginal correlation between the response variable and each gene expression variable based on all available pairs of observations of the two variables.

We first apply the proposed method to all the $n = 212$ samples in order to identify the various biomarkers that are associated with the score of the mini-mental state examination. We test the simultaneous hypotheses $H_{0j} : \beta_j = 0, 1 \leq j \leq p = 680$, while controlling the false discovery rate, using the modified Benjamini–Hochberg procedure of Ma et al. (2020) with the proposed estimators $\tilde{\beta}_j$ and their variance estimators \tilde{s}_j^2 . The detailed description of the multiple testing procedure can be found in Appendix F.

The identified biomarkers by all the methods at significance level $\alpha = 0.01$ are provided in Table 3 of the Appendix. For the gene expression probes, we provide the corresponding gene names in the table. The proposed method identifies 36 biomarkers, including 19, 2, and 15 variables from the MRI, PET, and gene expression, respectively. Some of these biomarkers are also selected by other methods. Although debiased Lasso using complete cases or using single regression imputation seems to identify many more markers, based on our simulation results, many of the identified markers may be false positive since the corresponding confidence intervals do not provide the correct coverage probabilities.

Among the gene expressions, SFRP1 gene is selected by all the methods, which is a crucial player in AD pathogenesis (Esteve et al., 2019). PJA2 gene is only identified by the proposed method and has reduced expressions in AD patients than on normal controls. PJA2 gene has been shown to regulate AD marker genes in mouse hippocampal neuronal cells, indicating its the potential relevance to the pathophysiology of AD (Gong et al., 2020). Among the MRI related markers, “ST30SV” is identified by not only the proposed method but also DL-SI and LP-SI. It represents the volume of left inferior lateral ventricle, which is related to the AD (Bartos et al., 2019; Ledig et al., 2018). Yet, “ST101SV” and “ST35TA”, representing the volume of the right pallidum and the average cortical thickness of the left lateral occipital, respectively, are only identified by the proposed method. Both were shown to be associated with AD by other studies (Kautzky et al., 2018; Yang et al., 2019). Finally, the PET biomarker “CTX_RH_TEMPORALPOLE”, the standardized uptake value of the right temporal pole, is only identified by the proposed method. This agrees with the observation that hypometabolism in temporal lobe often appears in AD patients (Sanabria-Diaz et al., 2013).

To illustrate that the multiple sources in the ADNI study contain complementary information, we compare the proposed method with the Lasso using only the MRI, PET, or gene expression variables in terms of prediction. We also compare the proposed method

Method	Prediction mean squared error (SD)	Improvement rate
Proposed ($\hat{\beta}$)	13.898 (4.427)	—
MRI Lasso	15.546 (5.715)	10.6%
PET Lasso	16.975 (7.009)	18.1%
Gene Lasso	19.946 (8.909)	30.3%
CC Lasso	19.956 (9.724)	30.4%
Naive mean	21.018 (10.410)	33.9%

Table 2: Averages of prediction mean squared errors based on 150 replications. Proposed ($\hat{\beta}$): the proposed method with the estimator $\hat{\beta}$. MRI Lasso, PET Lasso, and Gene Lasso: Lasso method using only MRI, PET, and gene expression variables, respectively. CC Lasso: the Lasso method using only complete cases. Naive mean: using the sample mean of the response variable in the training sets for prediction. SD: standard deviation of prediction mean squared errors calculated based on the 150 replications.

with the naive mean prediction method and Lasso using only the complete observations, where the naive mean prediction method uses the sample mean of the response variable calculated based on training sets for prediction. Specifically, we randomly hide 10% of all the values of the response variable as testing responses 150 times, and apply all the methods to the remaining data. In each replication, we calculate the prediction mean squared error $\sum_{1 \leq i \leq T} (\hat{y}_i - y_i)^2 / T$, where y_i is a testing response, \hat{y}_i is the corresponding predicted value, and T is the number of testing responses. We also compute improvement rates of the proposed method relative to other methods in terms of the prediction mean squared error, which is defined as $(PE_{\mathcal{M}} - PE_{\mathcal{P}}) / PE_{\mathcal{P}}$, where $PE_{\mathcal{P}}$ and $PE_{\mathcal{M}}$ denote averages of the prediction mean squared errors of the proposed method and the method \mathcal{M} , respectively, based on the 150 replications.

As shown in Table 2, the proposed estimator $\hat{\beta}$ produces smaller prediction mean squared errors than other estimators, which indicates that the proposed method can achieve higher prediction accuracy than using data from only one source or using only complete cases. Specifically, the proposed method reduces the prediction mean squared errors of other methods by at least 10.6%. In particular, the improvement rate with respect to the Lasso method using only gene expression variables or using only complete cases is over 30%. Moreover, the standard deviation of the prediction mean squared errors of the proposed method is smaller than that of other methods, indicating that the proposed method is more stable.

In summary, the proposed estimator produces smaller prediction mean squared errors than using only one source data or using only complete observations, implying that integration of data from multiple sources and usage of incomplete observations are critical. Additionally, the proposed method identifies meaningful and important biomarkers that are not selected by other methods, indicating that the proposed method is more powerful in integrating multi-modality data.

6. Discussion

In this paper, we propose a computational efficient estimator of regression coefficients for blockwise missing covariates based on the multiple blockwise imputation. In addition,

we propose a new bias-corrected estimator via projected estimating functions, and develop statistical inference for each regression coefficient. The proposed estimator produces smaller prediction errors and the proposed method provides confidence intervals closer to nominal level than other methods.

In Section 3, sample splitting was used to obtain the theoretical properties of the proposed estimators. In particular, this can be directly extended to the case where the size of the unsupervised samples is much larger than that of the supervised samples, since the multiple blockwise imputation step can be performed based on only the unsupervised data. In this case, Theorems 1 and 3 continue to hold only with $n + N$ replaced by N . However, to best exploit the available information in the data, we still suggest using all the samples for the multiple imputation, and all the supervised samples for the inference, for better empirical performance. To this end, it is unclear to us whether the same theoretical result can be obtained without imposing additional stringent technical conditions. Nevertheless, given the fact that the proposed methods perform well numerically without sample splitting, it is of interest to develop novel technical tools to facilitate the theoretical analysis.

Although the missing not at random mechanism is not covered in our theoretical justifications, simulation studies in Section 4 show that the proposed method still outperforms other methods under some informative missing settings. This could possibly due to that the proposed method incorporates more groups in the imputation of each missing block via the multiple blockwise imputation. In this way, the proposed method could aggregate information from various groups to reduce the selection bias in different groups caused by the informative missing. In the future research, we can handle the missing not at random situations through modeling the missingness or using instrumental variables.

A few other extensions are also worth exploring in the future. Since Alzheimer’s disease is a progressive brain disease, we could improve the proposed method through incorporating longitudinal information in the estimating functions to increase estimation and inference. In addition, although the proposed method only considers linear regression with continuous responses, it is feasible to extend the proposed method for categorical responses via generalized linear models.

Appendix A. Biomarkers selected by various methods for ADNI data set

Method	Biomarkers selected
Proposed	ST101SV, ST104TS, ST107TS, ST110TS, ST121TS, ST124SV, ST26TA, ST30SV, ST35TA, ST57TS, ST58TA, ST60TA, ST68SV, ST6SV, ST83CV, ST84CV, ST84SA, ST84TA, ST94CV, CTX_RH_TEMPORALPOLE, LEFT_CHOROID_PLEXUS, PJA2, SFRP1, P4HA3, ABCG1, IKZF5, NLRP10, ACOXL, DNAH2, EIF4ENIF1, TEKT4, PPIL2, TRPM8, SIGMAR1, ANKRD13C, MAGI2

Method	Biomarkers selected
DL-CC	ST129TS, ST15CV, ST15SA, ST147SV, ST24TA, ST26TA, ST29SV, ST32CV, ST34CV, ST40TA, ST45CV, ST48TA, ST60TA, ST60TS, ST72TA, ST73CV, ST76SV, ST77SV, ST84SA, ST85CV, ST93CV, ST97TA, ST97TS, CC_ANTERIOR, CTX_LH_CUNEUS, CTX_LH_LINGUAL, CTX_LH_PERICALCARINE, CTX_RH_CUNEUS, SUMMARYSUVR_WHOLECEREBNORM_1.11CUTOFF, SUMMARYSUVR_COMPOSITE_REFNORM, PHF1, SFRP1, GOLGA8A GOLGA8B, IKZF5, MECR, NLRP10, PROKR2, TAS2R4, DAZAP2, SERPINH1, LDLR, TTYH1, PSMB2, MAGI2, PCDH9, POSTN
LP-CC	ST147SV, ST32CV, SFRP1
DL-SI	ST129TA, ST130TS, ST18SV, ST24TA, ST26TA, ST29SV, ST30SV, ST31CV, ST31TA, ST32CV, ST40CV, ST40TA, ST46TA, ST48TA, ST58CV, ST58TA, ST59CV, ST80SV, ST83CV, ST85CV, ST89SV, ST90CV, ST90TA, CC_POSTERIOR, CTX_LH_FUSIFORM, LEFT_LATERAL_VENTRICLE, SUMMARYSUVR_WHOLECEREBNORM_1.11CUTOFF, SUMMARYSUVR_COMPOSITE_REFNORM, CTX_LH_FRONTALPOLE, COL4A1, PHF1, SFRP1, SCFD2, ABCG1, CDH2, NAALAD2, TRIM6-TRIM34 TRIM34, IKZF5, MECR, OXT, NLRP10, PROKR2, ACOXL, TAS2R4, BRDT, CACYBP, PCSK6, DNAH2, SERPINH1, EIF4ENIF1, TEK4, STXBP1, PPIL2, ABHD14B, LDLR, TRPM8, PRMT6, SIGMAR1, ZNF195, PSMB2, CALD1, MAGI2, PYCR2
LP-SI	ST107TS, ST30SV, ST39SA, ST46TA, CTX_RH_PARAHIPPOCAMPAL, SUMMARYSUVR_COMPOSITE_REFNORM, SFRP1, DNAH2, EIF4ENIF1, TEK4, TRPM8, SIGMAR1, MAGI2

Table 3: Biomarkers identified by all the methods.

Appendix B. Proof of Theorem 1

The proof of the following theorem can be separated into two parts. In the first part, we show that the constraint $\|g_n(\beta)\|_\infty \leq \lambda$ is feasible with high probability. In the second part, we obtain the rate of convergence of the proposed estimator $\hat{\beta}$. For simplicity, and without loss of generality, we assume that there are $2n$ samples in \mathcal{D}_2 , where half of the samples along with the N samples in \mathcal{D}_1 are used for imputation, whereas the other half of the samples are used for constructing the estimating function.

Throughout, we adopt the following notations. Recall that $\hat{\gamma}_{j,J(r,k)}$ is the Lasso estimator defined in the main paper, we define the matrix $\hat{\Gamma}_{J(r,k),m(r)} \in \mathbb{R}^{|J(r,k)| \times |m(r)|}$ whose columns are $\hat{\gamma}_{j,J(r,k)}$ with $j \in m(r)$. In addition, let $\gamma_{j,J(r,k)}$ be such that $\mathbb{E}(X_j | X_{J(r,k)}) = \gamma_{j,J(r,k)}^\top X_{J(r,k)}$. In other words, by the properties of Gaussian distribution, we have $\gamma_{j,J(r,k)} \equiv -\Sigma_{-j,j}^{-1} \Sigma_{jj}$ where $\Sigma_{-j,j}^{-1}$ is the j -th column of Σ^{-1} with its j -th component removed. Sim-

ilarly, we define $\mathbf{\Gamma}_{J(r,k),m(r)} \in \mathbb{R}^{|J(r,k)| \times |m(r)|}$ as a matrix whose columns are $\gamma_{j,J(r,k)}$ with $j \in J(r,k)$. As a consequence, by (A1), (A2) and the standard estimation bound for the Lasso estimator (Bickel et al., 2009; Negahban et al., 2010), we have

$$\max_{j \in m(r)} \|\widehat{\gamma}_{j,J(r,k)} - \gamma_{j,J(r,k)}\|_2 \lesssim \sqrt{\frac{s \log p}{N+n}}, \quad (24)$$

with probability at least $1 - p^{-c}$. In addition, for simplicity, we may also write $\mathbf{\Gamma}_{J(r,k),m(r)}$ as $\mathbf{\Gamma}_{r,k}$ and write $\widehat{\mathbf{\Gamma}}_{J(r,k),m(r)}$ as $\widehat{\mathbf{\Gamma}}_{r,k}$.

Part I. Feasibility. We start with the following lemmas.

Lemma 4 *Under conditions (A1) (A2) and (A3), with probability at least $1 - p^{-c}$, it holds that $\|g(\beta)\|_\infty \lesssim \sqrt{\frac{\log p}{n}}$.*

Lemma 5 *Under conditions (A1) (A2) and (A3), if $s \lesssim \frac{N+n}{\log p}$, then with probability at least $1 - p^{-c}$, it holds that $\|g_n(\beta) - g(\beta)\|_\infty \lesssim s \sqrt{\frac{\log p}{N+n}} (1 + s \sqrt{\frac{\log p}{n}})$*

Combining the above two lemmas, we have

$$\|g_n(\beta)\|_\infty \leq \|g_n(\beta) - g(\beta)\|_\infty + \|g(\beta)\|_\infty \lesssim \sqrt{\frac{\log p}{n}} + s \sqrt{\frac{\log p}{N+n}} \left(1 + s \sqrt{\frac{\log p}{n}}\right) \quad (25)$$

with probability at least $1 - p^{-c}$. Whenever $s \ll \sqrt{n/\log p}$, the RHS of (25) can be bounded by $\lambda \asymp \sqrt{\log p/n} + s \sqrt{\log p/(n+N)}$, which shows that the constraint $\|g_n(\beta)\|_\infty \leq \lambda$ is feasible with high probability.

Part II. Rate of Convergence. Firstly, if we denote $g_n(\beta) = \frac{1}{n} \sum_{i=1}^n \widehat{\mathbf{h}}_i(\beta)$, where $\widehat{\mathbf{h}}_i(\beta)$ has its components $\frac{n}{n_r} I\{\xi_i = r\} [y_i - (\widehat{X}_i^{(k)})^\top \beta] \widehat{X}_{ia(k)}^{(k)}$, it then follows that $g_n(\widehat{\beta}) = \frac{1}{n} \sum_{i=1}^n \widehat{\mathbf{h}}_i(\widehat{\beta})$ where $\widehat{\mathbf{h}}_i(\widehat{\beta})$ has its components

$$\begin{aligned} & \frac{n}{n_r} I\{\xi_i = r\} [y_i - (\widehat{X}_i^{(k)})^\top \widehat{\beta}] \widehat{X}_{ia(k)}^{(k)} \\ &= \frac{n}{n_r} I\{\xi_i = r\} [X_i^\top \beta - (\widehat{X}_i^{(k)})^\top \widehat{\beta} + \epsilon_i] \widehat{X}_{ia(k)}^{(k)} \\ &= \frac{n}{n_r} I\{\xi_i = r\} [(\widehat{X}_i^{(k)})^\top (\beta - \widehat{\beta}) + (X_i - \widehat{X}_i^{(k)})^\top \beta + \epsilon_i] \widehat{X}_{ia(k)}^{(k)}. \end{aligned} \quad (26)$$

Now, we will show that

$$\left\| \frac{1}{n} \sum_{i=1}^n \frac{n}{n_r} I\{\xi_i = r\} \widehat{X}_{ia(k)}^{(k)} (\widehat{X}_i^{(k)})^\top (\beta - \widehat{\beta}) \right\|_\infty \lesssim \lambda. \quad (27)$$

To see this, by the definition of $\widehat{\beta}$ and Part I, we have $\left\| \frac{1}{n} \sum_{i=1}^n \widehat{\mathbf{h}}_{irk}(\widehat{\beta}) \right\|_\infty \leq \|g_n(\widehat{\beta})\|_\infty \lesssim \lambda$, with probability at least $1 - p^{-c}$. Then, in light of the decomposition (26), it suffices to show that, with probability at least $1 - p^{-c}$,

$$\left\| \frac{1}{n} \sum_{i=1}^n I\{\xi_i = r\} \epsilon_i \widehat{X}_{ia(k)}^{(k)} \right\|_\infty \leq \lambda, \quad (28)$$

and

$$\left\| \frac{1}{n} \sum_{i=1}^n I\{\xi_i = r\} (X_i - \widehat{X}_i^{(k)})^\top \beta \widehat{X}_{ia(k)}^{(k)} \right\|_\infty \leq \lambda. \quad (29)$$

This is done by the following lemma. Thus (27) holds.

Lemma 6 *Under the conditions of Theorem 1, (28) and (29) hold with probability at least $1 - p^{-c}$.*

By (27), we have

$$\left| \frac{1}{n} \sum_{i=1}^n \frac{n}{n_r} I\{\xi_i = r\} [\beta - \widehat{\beta}]_{a(k)}^\top \widehat{X}_{ia(k)}^{(k)} (\widehat{X}_i^{(k)})^\top (\beta - \widehat{\beta}) \right| \lesssim \lambda \|\beta - \widehat{\beta}\|_{a(k)} \leq \lambda s^{1/2} \|\beta - \widehat{\beta}\|_2. \quad (30)$$

Next, we would like to prove the lower bound

$$\left| \frac{1}{n} \sum_{i=1}^n \frac{n}{n_r} I\{\xi_i = r\} [\beta - \widehat{\beta}]_{a(k)}^\top \widehat{X}_{ia(k)}^{(k)} (\widehat{X}_i^{(k)})^\top (\beta - \widehat{\beta}) \right| \gtrsim \|\beta - \widehat{\beta}\|_{a(k)} \|\beta - \widehat{\beta}\|_2, \quad (31)$$

which along with the upper bound (30) implies the final results $\|\beta - \widehat{\beta}\|_2 \leq s^{1/2} \lambda$ and $\|\beta - \widehat{\beta}\|_1 \leq s \lambda$. To show (31), we need the following key proposition concerning a restricted singular value condition.

Now by the definition of $\widehat{\beta}$, it follows from the same argument of Candès and Tao (2007) that $\|\widehat{\beta} - \beta\|_{S^c} \leq \|\widehat{\beta} - \beta\|_S$, where $S = \text{supp}(\beta)$. If in addition

$$\|\beta - \widehat{\beta}\|_{a(k)} \geq \|\beta - \widehat{\beta}\|_{m(k)}, \quad (32)$$

we have $\frac{\|\beta - \widehat{\beta}\|_{a(k)}}{\|\beta - \widehat{\beta}\|_2} \geq 1/2$ so that by (22), the lower bound (31) hold. The rest of the proof is devoted to (32).

Proof of (32). The existence of r and $k \in \mathcal{G}(r)$ such that (32) holds follows directly from the assumption that, there exists an $r \in \{1, \dots, R\}$ and $k, k' \in \mathcal{G}(r)$ such that $m(k) \subset a(k')$ and $m(k') \subset a(k)$. In this case, if $\|\beta - \widehat{\beta}\|_{a(k)} < \|\beta - \widehat{\beta}\|_{m(k)}$, we have $\|\beta - \widehat{\beta}\|_{m(k')} < \|\beta - \widehat{\beta}\|_{a(k')}$. Thus, there exists some $k \in \mathcal{G}(r)$ such that (32) holds.

Appendix C. Proof of Theorem 3

Again, without loss of generality, we assume that there are $2n$ samples in \mathcal{D}_2 , where half of the samples along with the N samples in \mathcal{D}_1 are used for imputation, whereas the other half of the samples are used for constructing the estimating function. We need the following lemmas.

Lemma 7 *Suppose the smallest singular value of $\mathbb{E}G \in \mathbb{R}^{M'_g \times p}$, where $M'_g = \sum_{r=1}^R \sum_{k \in \mathcal{G}(r)} |J(r, k)| > p$, is bounded away from zero. Then there exists infinite numbers of $v \in \mathbb{R}^{M'_g}$ such that $\mathbb{E}G^\top v = e_j$ and $\|v\|_2 \leq C$ for some constant $C > 0$.*

Lemma 8 Let $v_j^* \in \mathbb{R}^{M'_g}$ be any vector such that $\mathbb{E}G^\top v_j^* = e_j$. Suppose the smallest singular value of $\mathbb{E}G$ is bounded away from zero, and $R \asymp 1$. Then with probability at least $1 - p^{-c}$, it holds that $\|(v_j^*)^\top G_n - e_j^\top\|_\infty \lesssim \sqrt{\frac{s \log p}{N+n}}$.

Lemma 9 Let $\hat{S}_j(\beta) = \hat{v}_j^\top g_n^*(\beta)$. If $s\lambda\lambda'\sqrt{n} = o(1)$, it holds that

$$\sqrt{n}\hat{S}_j(\hat{\beta}_j^*) = \sqrt{n}\hat{S}_j(\beta) + o_P(1). \quad (33)$$

Lemma 10 Conditional on $\hat{X} = \{\hat{X}_i^{(k)}\}_{i \in \mathcal{D}_2}$, it holds that $\frac{n\hat{S}_j(\beta) - \mu_j}{s_j} | \hat{X} \rightarrow_d N(0, 1)$, where $\mu_j = \sum_{i=1}^n \sum_{k \in \mathcal{G}(r), 1 \leq r \leq R} \frac{n}{n_r} I\{\xi_i = r\} (X_i - \hat{X}_i^{(k)})^\top \beta \hat{v}_{rk}^\top X_{iJ(r,k)}$, and

$$s_j^2 = \sigma^2 \sum_{i=1}^n \sum_{k \in \mathcal{G}(r), 1 \leq r \leq R} \frac{n^2}{n_r^2} I\{\xi_i = r\} [\hat{v}_{rk}^\top X_{iJ(r,k)}]^2.$$

Let $\tilde{\beta}_j^*$ be such that $\tilde{\beta}_j^*$ is $\hat{\beta}_j^*$ by replacing its j -th component by $\tilde{\beta}_j$. By the mean value theorem, we have

$$0 = \hat{S}_j(\tilde{\beta}_j^*) = \hat{S}_j(\hat{\beta}_j^*) + \hat{v}_j^\top [G_n]_{\cdot j} (\tilde{\beta}_j - \beta_j). \quad (34)$$

By Lemma 9, we have $0 = \sqrt{n}\hat{S}_j(\beta) + \sqrt{n}\hat{v}_j^\top [G_n]_{\cdot j} (\tilde{\beta}_j - \beta_j) + o_P(1)$, or

$$\frac{\sqrt{n}(\tilde{\beta}_j - \beta_j)}{s_j/\sqrt{n}} = -\frac{n\hat{S}_j(\beta)}{s_j\hat{v}_j^\top [G_n]_{\cdot j}} + o_P(1) = -\frac{n\hat{S}_j(\beta) - \mu_j}{s_j\hat{v}_j^\top [G_n]_{\cdot j}} - \frac{\mu_j}{s_j\hat{v}_j^\top [G_n]_{\cdot j}} + o_P(1).$$

In the above equation, by the definition of \hat{v}_j and Lemma 8, it holds that, with probability at least $1 - p^{-c}$, $|\hat{v}_j^\top [G_n(\hat{\beta}_j^*)]_{\cdot j} - 1| \lesssim \sqrt{\frac{s \log p}{N}}$, or $\hat{v}_j^\top [G_n(\hat{\beta}_j^*)]_{\cdot j} \rightarrow_P 1$. Hence, it suffices to show

$$\mu_j/\sqrt{n} \rightarrow_P 0 \quad (35)$$

and

$$s_j^2/n \geq c > 0 \quad (36)$$

with probability at least $1 - p^{-c}$. With these, it follows that $\frac{n(\tilde{\beta}_j - \beta_j)}{s_j} = AB + D$, where $A \rightarrow 1$, $D \rightarrow 0$ in probability and $B | \hat{X} \rightarrow_d N(0, 1)$. The rest of the proof is devoted to (35) and (36).

Proof of (35) and (36). The proof of (35) can be established as soon as we prove

$$\left| \sum_{k \in \mathcal{G}(r), 1 \leq r \leq R} \frac{n}{n_r} \frac{1}{n} \sum_{i=1}^n I\{\xi_i = r\} (X_i - \hat{X}_i^{(k)})^\top \beta \hat{v}_{j,rk}^\top X_{iJ(r,k)} \right| = o_P(n^{-1/2}).$$

Now since

$$\begin{aligned} & \left| \sum_{k \in \mathcal{G}(r), 1 \leq r \leq R} \frac{1}{n} \sum_{i=1}^n \frac{n}{n_r} I\{\xi_i = r\} \beta_{m(r)}^\top (\hat{\Gamma}_{r,k} - \Gamma_{r,k})^\top X_{iJ(r,k)} (X_{iJ(r,k)})^\top \hat{v}_{j,rk} \right| \\ & \lesssim \sum_{k \in \mathcal{G}(r), 1 \leq r \leq R} \sqrt{\frac{1}{n} \sum_{i=1}^n [\beta_{m(r)}^\top (\hat{\Gamma}_{r,k} - \Gamma_{r,k})^\top X_{iJ(r,k)}]^2} \sqrt{\frac{1}{n} \sum_{i=1}^n \frac{n^2}{n_r^2} I\{\xi_i = r\} [(X_{iJ(r,k)})^\top \hat{v}_{j,rk}]^2}. \end{aligned} \quad (37)$$

By (41) we have $\max_{k,r} \|(\hat{\Gamma}_{r,k} - \Gamma_{r,k})\beta_{m(r)}\|_2 \lesssim s\sqrt{\frac{\log p}{N+n}}$ with probability at least $1 - p^{-c}$. By the definition of \hat{v}_j , we have

$$\sqrt{\frac{1}{n} \sum_{i=1}^n \frac{n^2}{n_r^2} I\{\xi_i = r\} [(X_{iJ(r,k)})^\top \hat{v}_{j,rk}]^2} \leq \sqrt{\frac{1}{n} \sum_{i=1}^n \frac{n^2}{n_r^2} I\{\xi_i = r\} [(X_{iJ(r,k)})^\top v_{j,rk}^*]^2},$$

where by concentration inequality of sub-exponential random variables, the right hand side of the above inequality is bounded by a constant with probability at least $1 - p^{-c}$. Now to show (36), it suffices to show

$$\frac{\sigma^2}{n} \sum_{i=1}^n \sum_{k \in \mathcal{G}(r), 1 \leq r \leq R} \frac{n^2}{n_r^2} I\{\xi_i = r\} [\hat{v}_{j,rk}^\top X_{iJ(r,k)}]^2 \geq c > 0. \quad (38)$$

To see this, note that by definition of \hat{v}_j , we have

$$\left| \sum_{k \in \mathcal{G}(r), 1 \leq r \leq R} \frac{1}{n} \sum_{i=1}^n \frac{n}{n_r} I\{\xi_i = r\} X_{ij}^{(k)} (X_{iJ(r,k)})^\top \hat{v}_{j,rk} - 1 \right| \leq \lambda',$$

or

$$\sum_{k \in \mathcal{G}(r), 1 \leq r \leq R} \frac{1}{n} \sum_{i=1}^n \frac{n}{n_r} I\{\xi_i = r\} X_{ij}^{(k)} (X_{iJ(r,k)})^\top \hat{v}_{j,rk} \geq 1 - \lambda'.$$

By Cauchy-Schwartz inequality, we have

$$\sum_{k \in \mathcal{G}(r), 1 \leq r \leq R} \sqrt{\frac{1}{n} \sum_{i=1}^n [X_{ij}^{(k)}]^2} \sqrt{\frac{1}{n} \sum_{i=1}^n \frac{n^2}{n_r^2} I\{\xi_i = r\} [(X_{iJ(r,k)})^\top \hat{v}_{j,rk}]^2} \geq 1 - \lambda',$$

or

$$\sqrt{\sum_{k \in \mathcal{G}(r), 1 \leq r \leq R} \frac{1}{n} \sum_{i=1}^n \frac{n^2}{n_r^2} I\{\xi_i = r\} [(X_{iJ(r,k)})^\top \hat{v}_{j,rk}]^2} \gtrsim \frac{1 - \lambda'}{\max_{k \in \mathcal{G}(r), 1 \leq r \leq R} \sqrt{\frac{1}{n} \sum_{i=1}^n [X_{ij}^{(k)}]^2}}.$$

Now since by concentration inequality of subexponential random variables with probability at least $1 - p^{-c}$, $\max_{k \in \mathcal{G}(r), 1 \leq r \leq R} \sqrt{\frac{1}{n} \sum_{i=1}^n [X_{ij}^{(k)}]^2} \geq c > 0$, it follows that under the same event (38) holds.

Appendix D. Proof of Auxiliary Lemmas

D.1 Proof of Lemma 4

By definition, $g(\beta)$ has its subvectors of the form

$$\begin{aligned} \frac{1}{n_r} \sum_{i=1}^n I\{\xi_i = r\} [y_i - (X_i^{(k)})^\top \beta] X_{ia(k)}^{(k)} &= \frac{1}{n_r} \sum_{i=1}^n I\{\xi_i = r\} [X_i^\top \beta + \epsilon_i - (X_i^{(k)})^\top \beta] X_{ia(k)}^{(k)} \\ &= \frac{1}{n_r} \sum_{i=1}^n I\{\xi_i = r\} [(X_i - X_i^{(k)})^\top \beta + \epsilon_i] X_{ia(k)}^{(k)}. \end{aligned}$$

By assumptions (A3) and $\|\beta\|_2 \leq C$, each of $(X_i - X_i^{(k)})^\top \beta$, ϵ_i and $X_i a(k)^{(k)}$ are sub-Gaussian random variables and are mutually independent across different samples $i \in \{1, \dots, n\}$. Since the product of two sub-Gaussian random variables are sub-exponential, and by the properties of conditional expectations $\mathbb{E}[(X_i - X_i^{(k)})^\top \beta + \epsilon_i] X_{ia(k)}^{(k)} = 0, i = 1, \dots, n$, then it follows that, as long as $0 < c_1 < n/n_r < c_2 < 1$ for each $1 \leq r \leq R$ (A2), each component of $g(\beta)$ is a centered sub-exponential variable. Applying the concentration inequality for independent sub-exponential random variables (see, for example, Proposition 5.16 of Vershynin (2010)), we have $P(\|g(\beta)\|_\infty \leq C\sqrt{\frac{\log M_g}{n}}) \geq 1 - M_g^{-c}$, for some constants $C, c > 0$. The final results follows from (A1) so that $p \lesssim M_g \leq R^2 p \lesssim p$.

D.2 Proof of Lemma 6

To prove (28), we note that $\widehat{X}_{ia(k)}^{(k)}$ can be partitioned into two subvectors, namely, $X_{iJ(r,k)}$ and $\widehat{X}_{im(r)}$. On the one hand, by the concentration inequality for independent centered sub-exponential random random variables, we have

$$P\left(\left\|\frac{1}{n} \sum_{i=1}^n I\{\xi_i = r\} \epsilon_i X_{iJ(r,k)}\right\|_\infty \leq C\sqrt{\frac{\log p}{n}}\right) \geq 1 - p^{-c}. \quad (39)$$

On the other hand, for any $j \in m(r)$,

$$[\widehat{X}_{im(r)} \epsilon_i]_j = \widehat{\gamma}_{j,J(r,k)}^\top X_{iJ(r,k)} \epsilon_i = (\widehat{\gamma}_{j,J(r,k)} - \gamma_{j,J(r,k)})^\top X_{iJ(r,k)} \epsilon_i + \gamma_{j,J(r,k)}^\top X_{iJ(r,k)} \epsilon_i.$$

By the concentration inequality for independent centered sub-exponential random random variables, we have $P(\max_{j \in m(r)} |\frac{1}{n} \sum_{i=1}^n I\{\xi_i = r\} \gamma_{j,J(r,k)}^\top X_{iJ(r,k)} \epsilon_i| \leq C\sqrt{\frac{\log p}{n}}) \geq 1 - p^{-c}$. In addition, by (24), with probability at least $1 - p^{-c}$, we have

$$\begin{aligned} & \max_{j \in m(r)} \left| \frac{1}{n} \sum_{i=1}^n I\{\xi_i = r\} (\widehat{\gamma}_{j,J(r,k)} - \gamma_{j,J(r,k)})^\top X_{iJ(r,k)} \epsilon_i \right|_\infty \\ & \leq \max_{j \in m(r)} \|\widehat{\gamma}_{j,J(r,k)} - \gamma_{j,J(r,k)}\|_1 \left\| \frac{1}{n} \sum_{i=1}^n I\{\xi_i = r\} X_{iJ(r,k)} \epsilon_i \right\|_\infty \lesssim s \sqrt{\frac{\log p}{N+n}} \sqrt{\frac{\log p}{n}}. \end{aligned}$$

Therefore,

$$P\left(\max_{j \in m(r)} \left| \frac{1}{n} \sum_{i=1}^n I\{\xi_i = r\} \widehat{\gamma}_{j,J(r,k)}^\top X_{iJ(r,k)} \epsilon_i \right| \leq C \left(1 + s \sqrt{\frac{\log p}{N+n}}\right) \sqrt{\frac{\log p}{n}}\right) \geq 1 - p^{-c}. \quad (40)$$

Inequality (28) then follows from (39) and (40).

To prove (29), note that $(X_i - \widehat{X}_i^{(k)})^\top \beta \widehat{X}_{ia(k)}^{(k)} = X_{iJ(r,k)}^\top (\widehat{\Gamma}_{r,k} - \Gamma_{r,k}) \beta_{m(r)} \widehat{X}_{ia(k)}^{(k)}$. Again, the above vector consists of two parts, corresponding to the above partition of $\widehat{X}_{ia(k)}^{(k)}$. For

the first part, it follows that

$$\begin{aligned}
 & \left\| \frac{1}{n} \sum_{i=1}^n I\{\xi_i = r\} X_{iJ(r,k)}^\top (\hat{\mathbf{\Gamma}}_{r,k} - \mathbf{\Gamma}_{r,k}) \beta_{m(r)} X_{iJ(r,k)} \right\|_\infty \\
 &= \max_{k \in J(r,k)} \left| \frac{1}{n} \sum_{i=1}^n I\{\xi_i = r\} \beta_{m(r)}^\top (\hat{\mathbf{\Gamma}}_{r,k} - \mathbf{\Gamma}_{r,k})^\top X_{iJ(r,k)} X_{ik} \right| \\
 &\leq \|\beta_{m(r)}^\top (\hat{\mathbf{\Gamma}}_{r,k} - \mathbf{\Gamma}_{r,k})^\top\|_2 \cdot \max_{k \in J(r,k)} \left| \frac{1}{n} \sum_{i=1}^n I\{\xi_i = r\} \mathbf{u}_{r,k}^\top X_{iJ(r,k)} X_{ik} \right|,
 \end{aligned}$$

where $\mathbf{u}_{r,k}^\top = \beta_{m(r)}^\top (\hat{\mathbf{\Gamma}}_{r,k} - \mathbf{\Gamma}_{r,k})^\top / \|\beta_{m(r)}^\top (\hat{\mathbf{\Gamma}}_{r,k} - \mathbf{\Gamma}_{r,k})^\top\|_2$. Note that $\|\beta_{m(r)}\|_1 \leq \sqrt{s} \|\beta\|_2 \lesssim \sqrt{s}$. Then we have

$$\begin{aligned}
 \|\hat{\mathbf{\Gamma}}_{r,k} - \mathbf{\Gamma}_{r,k}\|_2 &\leq \sum_{j \in m(r)} |\beta_j| \cdot \|\hat{\gamma}_{j,J(r,k)} - \gamma_{j,J(r,k)}\|_2 \\
 &\leq \|\beta_{m(r)}\|_1 \cdot \max_{1 \leq j \leq m(r)} \|\hat{\gamma}_{j,J(r,k)} - \gamma_{j,J(r,k)}\|_2 \lesssim s \sqrt{\frac{\log p}{N+n}},
 \end{aligned} \tag{41}$$

and, by the concentration inequality of sub-exponential random variables and the independence between $\hat{\mathbf{\Gamma}}_{r,k}$ and X_i ,

$$\max_{k \in J(r,k)} \left| \frac{1}{n} \sum_{i=1}^n I\{\xi_i = r\} \mathbf{u}_{r,k}^\top X_{iJ(r,k)} X_{ik} \right| \leq C, \tag{42}$$

with probability at least $1 - p^{-c}$.

For the second part, it follows that

$$\begin{aligned}
 & \left\| \frac{1}{n} \sum_{i=1}^n I\{\xi_i = r\} X_{iJ(r,k)}^\top (\hat{\mathbf{\Gamma}}_{r,k} - \mathbf{\Gamma}_{r,k}) \beta_{m(r)} X_{iJ(r,k)}^\top \hat{\mathbf{\Gamma}}_{r,k} \right\|_\infty \\
 &= \max_{k \in m(r)} \left| \frac{1}{n} \sum_{i=1}^n I\{\xi_i = r\} \beta_{m(r)}^\top (\hat{\mathbf{\Gamma}}_{r,k} - \mathbf{\Gamma}_{r,k})^\top X_{iJ(r,k)} X_{iJ(r,k)}^\top \hat{\gamma}_{j,J(r,k)} \right| \\
 &\leq \|\beta_{m(r)}^\top (\hat{\mathbf{\Gamma}}_{r,k} - \mathbf{\Gamma}_{r,k})^\top\|_2 \cdot \max_{k \in m(r)} \left| \frac{1}{n} \sum_{i=1}^n I\{\xi_i = r\} \mathbf{u}_{r,k}^\top X_{iJ(r,k)} X_{iJ(r,k)}^\top \gamma_{j,J(r,k)} \right| \\
 &\quad + \|\beta_{m(r)}^\top (\hat{\mathbf{\Gamma}}_{r,k} - \mathbf{\Gamma}_{r,k})^\top\|_2 \cdot \max_{k \in m(r)} \left| \frac{1}{n} \sum_{i=1}^n I\{\xi_i = r\} \mathbf{u}_{r,k}^\top X_{iJ(r,k)} X_{iJ(r,k)}^\top (\hat{\gamma}_{j,J(r,k)} - \gamma_{j,J(r,k)}) \right|,
 \end{aligned}$$

where by the concentration inequality for sub-exponential random variables, we have with probability at least $1 - p^{-c}$,

$$\max_{k \in m(r)} \left| \frac{1}{n} \sum_{i=1}^n I\{\xi_i = r\} \mathbf{u}_{r,k}^\top X_{iJ(r,k)} X_{iJ(r,k)}^\top \gamma_{j,J(r,k)} \right| \leq C \tag{43}$$

and

$$\begin{aligned} & \max_{k \in m(r)} \left| \frac{1}{n} \sum_{i=1}^n I\{\xi_i = r\} \mathbf{u}_{r,k}^\top X_{iJ(r,k)} X_{iJ(r,k)}^\top (\widehat{\gamma}_{j,J(r,k)} - \gamma_{j,J(r,k)}) \right| \\ & \leq \max_{k \in m(r)} \|\widehat{\gamma}_{j,J(r,k)} - \gamma_{j,J(r,k)}\|_2 \cdot \left\| \frac{1}{n} \sum_{i=1}^n I\{\xi_i = r\} \mathbf{u}_{r,k}^\top X_{iJ(r,k)} X_{iJ(r,k)}^\top \mathbf{v}_j \right\|_2 \lesssim \sqrt{\frac{s \log p}{N+n}}, \end{aligned}$$

where in the second last inequality $\mathbf{v}_j = (\widehat{\gamma}_{j,J(r,k)} - \gamma_{j,J(r,k)}) / \|\widehat{\gamma}_{j,J(r,k)} - \gamma_{j,J(r,k)}\|_2$ and the last inequality follows from the concentration inequality for the sub-exponential random variables. In particular, in (43), we used the fact that $\sup_{j,r,k} \|\mathbf{\Gamma}_{r,k} \cdot j\|_2 \leq C$, which is implied by the assumption that $\Omega = \Sigma^{-1}$ has bounded eigenvalues from both above and below. The above inequalities imply (29).

D.3 Proof of Lemma 5

For each $r \in [1 : R]$, $i \in \mathcal{H}(r)$ and $k \in \mathcal{G}(r)$, we have

$$\widehat{h}_{irk}(\beta) = I(\xi_i = r) \{X_i - \widehat{X}_i^{(k)}\}^\top \beta + \epsilon_i \cdot \widehat{X}_{ia(k)}^{(k)} = I(\xi_i = r) \left[\{X_{im(r)} - \widehat{X}_{im(r)}^{(k)}\}^\top \beta_{m(r)} + \epsilon_i \right] \cdot \widehat{X}_{ia(k)}^{(k)},$$

and therefore $g(\beta) - g_n(\beta) = \frac{1}{n} \sum_{i=1}^n \{\mathbf{h}_i(\beta) - \widehat{\mathbf{h}}_i(\beta)\}$, where $\widehat{\mathbf{h}}_i(\beta) = (\widehat{\theta}_1^{-1} \widehat{h}_{i1}(\beta)^\top, \dots, \widehat{\theta}_R^{-1} \widehat{h}_{iR}(\beta)^\top)^\top$ and $\mathbf{h}_i(\beta) = (\widehat{\theta}_1^{-1} h_{i1}(\beta)^\top, \dots, \widehat{\theta}_R^{-1} h_{iR}(\beta)^\top)^\top$ which consist of all the (imputed) estimating functions for the i -th sample. In particular, for $k \in \mathcal{G}(r)$, we have

$$\begin{aligned} & \frac{1}{n} \sum_{i=1}^n \widehat{\theta}_r^{-1} \{h_{irk}(\beta) - \widehat{h}_{irk}(\beta)\} \\ & = \frac{1}{n} \sum_{i=1}^n \widehat{\theta}_r^{-1} I(\xi_i = r) \left([y_i - \{X_i^{(k)}\}^\top \beta] \cdot X_{ia(k)}^{(k)} - [y_i - \{\widehat{X}_i^{(k)}\}^\top \beta] \widehat{X}_{ia(k)}^{(k)} \right) \\ & = \frac{1}{n \widehat{\theta}_r} \sum_{i=1}^n I(\xi_i = r) \left(\left[\{X_{im(r)} - X_{im(r)}^{(k)}\}^\top \beta_{m(r)} + \epsilon_i \right] X_{ia(k)}^{(k)} - \left[\{X_{im(r)} - \widehat{X}_{im(r)}^{(k)}\}^\top \beta_{m(r)} + \epsilon_i \right] \widehat{X}_{ia(k)}^{(k)} \right) \\ & = \frac{1}{n \widehat{\theta}_r} \sum_{i=1}^n I(\xi_i = r) \left(X_{ia(k)}^{(k)} \{X_{im(r)} - X_{im(r)}^{(k)}\}^\top \beta_{m(r)} - \widehat{X}_{ia(k)}^{(k)} \{X_{im(r)} - \widehat{X}_{im(r)}^{(k)}\}^\top \beta_{m(r)} \right. \\ & \quad \left. + \epsilon_i \{X_{ia(k)}^{(k)} - \widehat{X}_{ia(k)}^{(k)}\} \right). \end{aligned}$$

Then,

$$\begin{aligned}
 & \left\| \frac{1}{n} \sum_{i=1}^n \hat{\theta}_r^{-1} \left\{ h_{irk}(\beta) - \hat{h}_{irk}(\beta) \right\} \right\|_{\infty} \\
 = & \hat{\theta}_r^{-1} \max_{1 \leq r \leq R, k \in \mathcal{G}(r), j \in a(k)} \left| \frac{1}{n} \sum_{i=1}^n I(\xi_i = r) \left(X_{ij}^{(k)} \{ X_{im(r)} - X_{im(r)}^{(k)} \}^{\top} \beta_{m(r)} \right. \right. \\
 & \left. \left. - \hat{X}_{ij}^{(k)} \{ X_{im(r)} - \hat{X}_{im(r)}^{(k)} \}^{\top} \beta_{m(r)} + \epsilon_i \cdot \{ X_{ij}^{(k)} - \hat{X}_{ij}^{(k)} \} \right) \right| \\
 = & \hat{\theta}_r^{-1} \max_{1 \leq r \leq R, k \in \mathcal{G}(r), j \in a(k)} \left| \frac{1}{n} \sum_{i=1}^n I(\xi_i = r) \left(X_{ij}^{(k)} X_{iJ(r,k)}^{\top} \left\{ \hat{\mathbf{\Gamma}}_{J(r,k),m(r)} - \mathbf{\Gamma}_{J(r,k),m(r)} \right\} \beta_{m(r)} \right. \right. \\
 & \left. \left. + \left\{ \hat{X}_{ij}^{(k)} - X_{ij}^{(k)} \right\} \left[X_{iJ(r,k)}^{\top} \left\{ \hat{\mathbf{\Gamma}}_{J(r,k),m(r)} - \mathbf{\Gamma}_{J(r,k),m(r)} \right\} - \left\{ \boldsymbol{\epsilon}_{im(r)}^{(k)} \right\}^{\top} \right] \beta_{m(r)} + \epsilon_i \cdot \left\{ X_{ij}^{(k)} - \hat{X}_{ij}^{(k)} \right\} \right) \right|,
 \end{aligned}$$

where $\boldsymbol{\epsilon}_{im(r)}^{(k)}$ is the residual term of the i -th sample in the regression model of $X_{im(r)}$ with $X_{iJ(r,k)}$ as covariates. In particular, $\boldsymbol{\epsilon}_{im(r)}^{(k)}$ is centered and uncorrelated with $X_{iJ(r,k)}$.

If $j \in J(r, k)$, then

$$\begin{aligned}
 & \left| \frac{1}{n_r} \sum_{i=1}^n I(\xi_i = r) \left(X_{ij}^{(k)} X_{iJ(r,k)}^{\top} \left\{ \hat{\mathbf{\Gamma}}_{J(r,k),m(r)} - \mathbf{\Gamma}_{J(r,k),m(r)} \right\} \beta_{m(r)} \right. \right. \\
 & \left. \left. + \left\{ \hat{X}_{ij}^{(k)} - X_{ij}^{(k)} \right\} \left[X_{iJ(r,k)}^{\top} \left\{ \hat{\mathbf{\Gamma}}_{J(r,k),m(r)} - \mathbf{\Gamma}_{J(r,k),m(r)} \right\} - \left\{ \boldsymbol{\epsilon}_{im(r)}^{(k)} \right\}^{\top} \right] \beta_{m(r)} + \epsilon_i \cdot \left\{ X_{ij}^{(k)} - \hat{X}_{ij}^{(k)} \right\} \right) \right| \\
 = & \left| \frac{1}{n_r} \sum_{i=1}^n I(\xi_i = r) X_{ij} X_{iJ(r,k)}^{\top} \mathbf{u}_{r,k} \right| \cdot \| (\hat{\mathbf{\Gamma}}_{J(r,k),m(r)} - \mathbf{\Gamma}_{J(r,k),m(r)}) \beta_{m(r)} \|_2 \lesssim s \sqrt{\frac{\log p}{N+n}}, \quad (44)
 \end{aligned}$$

where the last inequality follows from (41) and (42). If $j \in a(k) \setminus J(r, k)$, then

$$\begin{aligned}
 & \frac{1}{n_r} \sum_{i=1}^n I(\xi_i = r) \left(X_{ij}^{(k)} \mathbf{X}_{iJ(r,k)}^{\top} \left\{ \hat{\mathbf{\Gamma}}_{J(r,k),m(r)} - \mathbf{\Gamma}_{J(r,k),m(r)} \right\} \beta_{m(r)} \right. \\
 & \left. + \left\{ \hat{X}_{ij}^{(k)} - X_{ij}^{(k)} \right\} \left[X_{iJ(r,k)}^{\top} \left\{ \hat{\mathbf{\Gamma}}_{J(r,k),m(r)} - \mathbf{\Gamma}_{J(r,k),m(r)} \right\} - \left\{ \boldsymbol{\epsilon}_{im(r)}^{(k)} \right\}^{\top} \right] \beta_{m(r)} + \epsilon_i \cdot \left\{ X_{ij}^{(k)} - \hat{X}_{ij}^{(k)} \right\} \right) \\
 = & \left\{ \frac{1}{n_r} \sum_{i=1}^n I(\xi_i = r) X_{ij}^{(k)} \mathbf{X}_{iJ(r,k)}^{\top} \right\} \left\{ \hat{\mathbf{\Gamma}}_{J(r,k),m(r)} - \mathbf{\Gamma}_{J(r,k),m(r)} \right\} \beta_{m(r)} \\
 & + \left\{ \gamma_{j,J(r,k)} - \hat{\gamma}_{j,J(r,k)} \right\}^{\top} \left\{ \frac{1}{n_r} \sum_{i=1}^n I(\xi_i = r) X_{iJ(r,k)} X_{iJ(r,k)}^{\top} \right\} \left\{ \hat{\mathbf{\Gamma}}_{J(r,k),m(r)} - \mathbf{\Gamma}_{J(r,k),m(r)} \right\} \beta_{m(r)} \\
 & + \left\{ \gamma_{j,J(r,k)} - \hat{\gamma}_{j,J(r,k)} \right\}^{\top} \left\{ \frac{1}{n_r} \sum_{i=1}^n I(\xi_i = r) X_{iJ(r,k)} \left\{ \boldsymbol{\epsilon}_{im(r)}^{(k)} \right\}^{\top} \right\} \beta_{m(r)} \\
 & + \left\{ \gamma_{j,J(r,k)} - \hat{\gamma}_{j,J(r,k)} \right\}^{\top} \frac{1}{n_r} \sum_{i=1}^n I(\xi_i = r) \epsilon_i X_{iJ(r,k)} \\
 = & \gamma_{j,J(r,k)}^{\top} \left\{ \frac{1}{n_r} \sum_{i=1}^n I(\xi_i = r) \mathbf{X}_{iJ(r,k)} X_{iJ(r,k)}^{\top} \right\} \left\{ \hat{\mathbf{\Gamma}}_{J(r,k),m(r)} - \mathbf{\Gamma}_{J(r,k),m(r)} \right\} \beta_{m(r)}
 \end{aligned}$$

$$\begin{aligned}
 & + \{\gamma_{j,J(r,k)} - \widehat{\gamma}_{j,J(r,k)}\}^\top \left\{ \frac{1}{n_r} \sum_{i=1}^n I(\xi_i = r) X_{iJ(r,k)} X_{iJ(r,k)}^\top \right\} \left\{ \widehat{\mathbf{\Gamma}}_{J(r,k),m(r)} - \mathbf{\Gamma}_{J(r,k),m(r)} \right\} \beta_{m(r)} \\
 & + \{\gamma_{j,J(r,k)} - \widehat{\gamma}_{j,J(r,k)}\}^\top \left\{ \frac{1}{n_r} \sum_{i=1}^n I(\xi_i = r) X_{iJ(r,k)} \left\{ \boldsymbol{\varepsilon}_{im(r)}^{(k)} \right\}^\top \right\} \beta_{m(r)} \\
 & + \{\gamma_{j,J(r,k)} - \widehat{\gamma}_{j,J(r,k)}\}^\top \frac{1}{n_r} \sum_{i=1}^n I(\xi_i = r) \epsilon_i X_{iJ(r,k)} \\
 = & \widehat{\theta}_r^{-1} \gamma_{j,J(r,k)}^\top \left\{ \frac{1}{n} \sum_{i=1}^n I(\xi_i = r) X_{iJ(r,k)} X_{iJ(r,k)}^\top \right\} \left\{ \widehat{\mathbf{\Gamma}}_{J(r,k),m(r)} - \mathbf{\Gamma}_{J(r,k),m(r)} \right\} \beta_{m(r)} \\
 & + \widehat{\theta}_r^{-1} \{\gamma_{j,J(r,k)} - \widehat{\gamma}_{j,J(r,k)}\}^\top \left\{ \frac{1}{n} \sum_{i=1}^n I(\xi_i = r) X_{iJ(r,k)} X_{iJ(r,k)}^\top \right\} \left\{ \widehat{\mathbf{\Gamma}}_{J(r,k),m(r)} - \mathbf{\Gamma}_{J(r,k),m(r)} \right\} \beta_{m(r)} \\
 & + \widehat{\theta}_r^{-1} \{\gamma_{j,J(r,k)} - \widehat{\gamma}_{j,J(r,k)}\}^\top \left\{ \frac{1}{n} \sum_{i=1}^n I(\xi_i = r) X_{iJ(r,k)} \left\{ \boldsymbol{\varepsilon}_{im(r)}^{(k)} \right\}^\top \right\} \beta_{m(r)} \\
 & + \widehat{\theta}_r^{-1} \{\gamma_{j,J(r,k)} - \widehat{\gamma}_{j,J(r,k)}\}^\top \frac{1}{n} \sum_{i=1}^n I(\xi_i = r) \epsilon_i X_{iJ(r,k)} \\
 = & \widehat{\theta}_r^{-1} \gamma_{j,J(r,k)}^\top \mathbf{S}_n \left\{ \widehat{\mathbf{\Gamma}}_{J(r,k),m(r)} - \mathbf{\Gamma}_{J(r,k),m(r)} \right\} \beta_{m(r)} \\
 & + \widehat{\theta}_r^{-1} \gamma_{j,J(r,k)}^\top \mathbf{C}_{J(r,k),J(r,k)}^* \left\{ \widehat{\mathbf{\Gamma}}_{J(r,k),m(r)} - \mathbf{\Gamma}_{J(r,k),m(r)} \right\} \beta_{m(r)} \\
 & + \widehat{\theta}_r^{-1} \{\gamma_{j,J(r,k)} - \widehat{\gamma}_{j,J(r,k)}\}^\top \mathbf{S}_n \left\{ \widehat{\mathbf{\Gamma}}_{J(r,k),m(r)} - \mathbf{\Gamma}_{J(r,k),m(r)} \right\} \beta_{m(r)} \\
 & + \widehat{\theta}_r^{-1} \{\gamma_{j,J(r,k)} - \widehat{\gamma}_{j,J(r,k)}\}^\top \mathbf{C}_{J(r,k),J(r,k)}^* \left\{ \widehat{\mathbf{\Gamma}}_{J(r,k),m(r)} - \mathbf{\Gamma}_{J(r,k),m(r)} \right\} \beta_{m(r)} \\
 & + \widehat{\theta}_r^{-1} \{\gamma_{j,J(r,k)} - \widehat{\gamma}_{j,J(r,k)}\}^\top \left\{ \frac{1}{n} \sum_{i=1}^n I(\xi_i = r) X_{iJ(r,k)} \left\{ \boldsymbol{\varepsilon}_{im(r)}^{(k)} \right\}^\top \right\} \beta_{m(r)} \\
 & + \widehat{\theta}_r^{-1} \{\gamma_{j,J(r,k)} - \widehat{\gamma}_{j,J(r,k)}\}^\top \frac{1}{n} \sum_{i=1}^n I(\xi_i = r) \epsilon_i X_{iJ(r,k)}, \tag{45}
 \end{aligned}$$

where $\mathbf{C}_{J(r,k),J(r,k)}^* = \mathbb{E} \left\{ I(\xi_i = r) X_{iJ(r,k)} X_{iJ(r,k)}^\top \right\}$ and $\mathbf{S}_n = \frac{1}{n} \sum_{i=1}^n I(\xi_i = r) X_{iJ(r,k)} X_{iJ(r,k)}^\top - \mathbf{C}_{J(r,k),J(r,k)}^*$. For the first term in (45), we have, with probability at least $1 - p^{-c}$

$$\begin{aligned}
 & \left| \gamma_{j,J(r,k)}^\top \mathbf{S}_n \left\{ \widehat{\mathbf{\Gamma}}_{J(r,k),m(r)} - \mathbf{\Gamma}_{J(r,k),m(r)} \right\} \beta_{m(r)} \right| \\
 & \leq \|\gamma_{j,J(r,k)}\|_1 \|\mathbf{S}_n\|_\infty \left\| \left(\widehat{\mathbf{\Gamma}}_{J(r,k),m(r)} - \mathbf{\Gamma}_{J(r,k),m(r)} \right) \beta_{m(r)} \right\|_1 \\
 & \leq C \sqrt{s} \sqrt{\frac{\log p}{n}} s^{3/2} \sqrt{\frac{\log p}{N+n}} \lesssim \frac{s^2 \log p}{\sqrt{(N+n)n}} \tag{46}
 \end{aligned}$$

where the second last inequality follows from

$$\left\| \left(\widehat{\mathbf{\Gamma}}_{J(r,k),m(r)} - \mathbf{\Gamma}_{J(r,k),m(r)} \right) \beta_{m(r)} \right\|_1 \leq \max_{j \in m(r)} \|\widehat{\gamma}_{j,J(r,k)} - \gamma_{j,J(r,k)}\|_1 \|\beta_{m(r)}\|_1 \lesssim s^{3/2} \sqrt{\frac{\log p}{N+n}}. \tag{47}$$

and $P(\|\mathbf{S}_n\|_\infty \geq C\sqrt{\frac{\log p}{n}}) \leq p^{-c}$, which is a direct consequence of the concentration inequality of sub-exponential random variables. For the second term in (45), we have with probability at least $1 - p^{-c}$

$$\begin{aligned} & \left| \gamma_{j,J(r,k)}^\top \mathbf{C}_{J(r,k),J(r,k)}^* \left\{ \hat{\mathbf{\Gamma}}_{J(r,k),m(r)} - \mathbf{\Gamma}_{J(r,k),m(r)} \right\} \beta_{m(r)} \right| \\ & \leq \|\gamma_{j,J(r,k)}\|_2 \left\| \mathbf{C}_{J(r,k),J(r,k)}^* \right\|_2 \left\| \left\{ \hat{\mathbf{\Gamma}}_{J(r,k),m(r)} - \mathbf{\Gamma}_{J(r,k),m(r)} \right\} \beta_{m(r)} \right\|_2 \lesssim s \sqrt{\frac{\log p}{N+n}}, \end{aligned} \quad (48)$$

where the last inequality follows from (41). For the third term, we have, with probability at least $1 - p^{-c}$,

$$\begin{aligned} & \left| \left\{ \gamma_{j,J(r,k)} - \hat{\gamma}_{j,J(r,k)} \right\}^\top \mathbf{S}_n \left\{ \hat{\mathbf{\Gamma}}_{J(r,k),m(r)} - \mathbf{\Gamma}_{J(r,k),m(r)} \right\} \beta_{m(r)} \right| \\ & \leq \|\gamma_{j,J(r,k)} - \hat{\gamma}_{j,J(r,k)}\|_1 \|\mathbf{S}_n\|_\infty \left\| \hat{\mathbf{\Gamma}}_{J(r,k),m(r)} - \mathbf{\Gamma}_{J(r,k),m(r)} \right\|_1 \|\beta_{m(r)}\|_1 \\ & \leq Cs \sqrt{\frac{\log p}{N+n}} \sqrt{\frac{\log p}{n}} s \sqrt{\frac{\log p}{N+n}} \|\beta_{m(r)}\|_1 \lesssim \frac{s^2 \log^{3/2} p}{(N+n)\sqrt{n}}. \end{aligned} \quad (49)$$

The fourth term satisfies

$$\begin{aligned} & \left| \left\{ \gamma_{j,J(r,k)} - \hat{\gamma}_{j,J(r,k)} \right\}^\top \mathbf{C}_{J(r,k),J(r,k)}^* \left\{ \hat{\mathbf{\Gamma}}_{J(r,k),m(r)} - \mathbf{\Gamma}_{J(r,k),m(r)} \right\} \beta_{m(r)} \right| \\ & \leq \|\gamma_{j,J(r,k)} - \hat{\gamma}_{j,J(r,k)}\|_2 \left\| \mathbf{C}_{J(r,k),J(r,k)}^* \right\|_2 \left\| \left\{ \hat{\mathbf{\Gamma}}_{J(r,k),m(r)} - \mathbf{\Gamma}_{J(r,k),m(r)} \right\} \beta_{m(r)} \right\|_2 \\ & \lesssim \sqrt{\frac{s \log p}{N+n}} \cdot s \sqrt{\frac{\log p}{N+n}} \lesssim \frac{s^{3/2} \log p}{N+n}, \end{aligned} \quad (50)$$

with probability at least $1 - p^{-c}$. The fifth term satisfies

$$\begin{aligned} & \left| \left\{ \gamma_{j,J(r,k)} - \hat{\gamma}_{j,J(r,k)} \right\}^\top \left\{ \frac{1}{n} \sum_{i=1}^n I(\xi_i = r) X_{iJ(r,k)} \left\{ \boldsymbol{\varepsilon}_{im(r)}^{(k)} \right\}^\top \right\} \beta_{m(r)} \right| \\ & \leq \|\gamma_{j,J(r,k)} - \hat{\gamma}_{j,J(r,k)}\|_1 \left\| \frac{1}{n} \sum_{i=1}^n I(\xi_i = r) X_{iJ(r,k)} \left\{ \boldsymbol{\varepsilon}_{im(r)}^{(k)} \right\}^\top \right\|_\infty \|\beta_{m(r)}\|_1 \\ & \lesssim s \sqrt{\frac{\log p}{N+n}} \sqrt{\frac{\log p}{n}} \|\beta_{m(r)}\|_1 \lesssim \frac{s^{3/2} \log p}{\sqrt{(N+n)n}}, \end{aligned} \quad (51)$$

with probability at least $1 - p^{-c}$. Lastly, the sixth term satisfies

$$\begin{aligned} & \left| \left\{ \gamma_{j,J(r,k)} - \hat{\gamma}_{j,J(r,k)} \right\}^\top \frac{1}{n} \sum_{i=1}^n I(\xi_i = r) \epsilon_i X_{iJ(r,k)} \right| \\ & \leq \|\gamma_{j,J(r,k)} - \hat{\gamma}_{j,J(r,k)}\|_1 \left\| \frac{1}{n} \sum_{i=1}^n I(\xi_i = r) \epsilon_i X_{iJ(r,k)} \right\|_\infty \lesssim s \sqrt{\frac{\log p}{N+n}} \sqrt{\frac{\log p}{n}}, \end{aligned} \quad (52)$$

with probability at least $1 - p^{-c}$. Combining all the pieces together, it follows that with probability at least $1 - p^{-c}$,

$$\|g_n(\beta) - g(\beta)\|_\infty \lesssim s \sqrt{\frac{\log p}{N+n}} + s \sqrt{\frac{\log p}{N+n}} \cdot s \sqrt{\frac{\log p}{n}} \asymp s \sqrt{\frac{\log p}{N+n}} \left(1 + s \sqrt{\frac{\log p}{n}} \right).$$

where the last inequality follow from $s \lesssim (n+N)/\log p$.

D.4 Proof of Proposition 2

The proof is separated into two parts. For simplicity, we write $E_s(p)$ as E_s when there is no risk of confusion. In the first part, we show that with probability at least $1 - p^{-c}$,

$$\sup_{\substack{\|\mathbf{u}\|_2=1, \mathbf{u} \in E_s \\ \|\mathbf{u}_{a(k)}\|_2 \geq 1/2}} \left| \frac{\mathbf{u}_{a(k)}^\top}{\|\mathbf{u}_{a(k)}\|_2} \frac{1}{n} \sum_{i=1}^n \frac{n}{n_r} I\{\xi_i = r\} [\widehat{X}_{ia(k)}^{(k)} (\widehat{X}_i^{(k)})^\top - X_{ia(k)}^{(k)} (X_i^{(k)})^\top] \mathbf{u} \right| \leq c/4. \quad (53)$$

In the second part, we show that with probability at least $1 - p^{-c}$,

$$\inf_{\substack{\|\mathbf{u}\|_2=1, \mathbf{u} \in E_s \\ \|\mathbf{u}_{a(k)}\|_2 \geq 1/2}} \left| \frac{\mathbf{u}_{a(k)}^\top}{\|\mathbf{u}_{a(k)}\|_2} \frac{1}{n} \sum_{i=1}^n \frac{n}{n_r} I\{\xi_i = r\} X_{ia(k)}^{(k)} (X_i^{(k)})^\top \mathbf{u} \right| \geq c. \quad (54)$$

The above inequalities implies (22).

Part I. Note that by Hölder's inequality

$$\begin{aligned} & \sup_{\substack{\|\mathbf{u}\|_2=1, \mathbf{u} \in E_s \\ \|\mathbf{u}_{a(k)}\|_2 \geq 1/2}} \left| \frac{\mathbf{u}_{a(k)}^\top}{\|\mathbf{u}_{a(k)}\|_2} \frac{1}{n} \sum_{i=1}^n \frac{n}{n_r} I\{\xi_i = r\} [\widehat{X}_{ia(k)}^{(k)} (\widehat{X}_i^{(k)})^\top - X_{ia(k)}^{(k)} (X_i^{(k)})^\top] \mathbf{u} \right| \\ & \lesssim s \left\| \frac{1}{n} \sum_{i=1}^n \frac{n}{n_r} I\{\xi_i = r\} [\widehat{X}_{ia(k)}^{(k)} (\widehat{X}_i^{(k)})^\top - X_{ia(k)}^{(k)} (X_i^{(k)})^\top] \right\|_\infty. \end{aligned}$$

By definition, it holds that, after proper permutation of coordinates $\widehat{X}_i^{(k)} = ((X_{ia(r)}^{(k)})^\top, (\widehat{X}_{im(r)}^{(k)})^\top)^\top =$

$$(X_{ia(r)}^\top, X_{iJ(r,k)}^\top \widehat{\mathbf{\Gamma}}_{r,k})^\top = \begin{bmatrix} \mathbf{I}_{|a(r)|} \\ [\widehat{\mathbf{\Gamma}}_{r,k}^\top, \mathbf{0}] \end{bmatrix} X_{ia(r)}, \text{ and}$$

$$\widehat{X}_{ia(k)}^{(k)} = ((X_{i,a(k) \cap a(r)}^{(k)})^\top, (\widehat{X}_{im(r)}^{(k)})^\top)^\top = (X_{iJ(r,k)}^\top, X_{iJ(r,k)}^\top \widehat{\mathbf{\Gamma}}_{r,k})^\top = \begin{bmatrix} \mathbf{I}_{|J(r,k)|} \\ \widehat{\mathbf{\Gamma}}_{r,k}^\top \end{bmatrix} X_{iJ(r,k)}.$$

Note that

$$\begin{aligned} & \left\| \frac{1}{n} \sum_{i=1}^n \frac{n}{n_r} I\{\xi_i = r\} [\widehat{X}_{ia(k)}^{(k)} (\widehat{X}_i^{(k)})^\top - X_{ia(k)}^{(k)} (X_i^{(k)})^\top] \right\|_\infty \\ & \leq \left\| \frac{1}{n} \sum_{i=1}^n \frac{n}{n_r} I\{\xi_i = r\} [\widehat{X}_{ia(k)}^{(k)} - X_{ia(k)}^{(k)}] (X_i^{(k)})^\top \right\|_\infty + \left\| \frac{1}{n} \sum_{i=1}^n \frac{n}{n_r} I\{\xi_i = r\} X_{ia(k)}^{(k)} (X_i^{(k)} - \widehat{X}_i^{(k)})^\top \right\|_\infty \\ & \quad + \left\| \frac{1}{n} \sum_{i=1}^n \frac{n}{n_r} I\{\xi_i = r\} (\widehat{X}_{ia(k)}^{(k)} - X_{ia(k)}^{(k)}) (X_i^{(k)} - \widehat{X}_i^{(k)})^\top \right\|_\infty \end{aligned}$$

To control the first term, we have

$$\begin{aligned}
 & \left\| \frac{1}{n} \sum_{i=1}^n \frac{n}{n_r} I\{\xi_i = r\} [\widehat{X}_{ia(k)}^{(k)} - X_{ia(k)}^{(k)}] (X_i^{(k)})^\top \right\|_\infty \\
 &= \left\| \frac{1}{n} \sum_{i=1}^n \frac{n}{n_r} I\{\xi_i = r\} \begin{bmatrix} 0 \\ \widehat{\mathbf{\Gamma}}_{r,k}^\top - \mathbf{\Gamma}_{r,k}^\top \end{bmatrix} X_{iJ(r,k)} X_{ia(r)}^\top [\mathbf{I}_{|a(r)|} \quad [\mathbf{\Gamma}_{r,k}, 0]^\top] \right\|_\infty \\
 &= \max_{\substack{1 \leq j \leq |a(k)| \\ 1 \leq k \leq p}} \left| \frac{1}{n} \sum_{i=1}^n \frac{n}{n_r} I\{\xi_i = r\} \mathbf{u}_j^\top X_{iJ(r,k)} X_{ia(r)}^\top \mathbf{v}_k \right|
 \end{aligned}$$

where \mathbf{u}_j is the j -th row of $\begin{bmatrix} 0 \\ \widehat{\mathbf{\Gamma}}_{r,k}^\top - \mathbf{\Gamma}_{r,k}^\top \end{bmatrix}$ and \mathbf{v}_k is the k -th row of $\begin{bmatrix} \mathbf{I}_{|a(r)|} \\ [\mathbf{\Gamma}_{r,k}, 0]^\top \end{bmatrix}$. Now since for any $j \in m(r)$, $\|[\mathbf{\Gamma}_{r,k}]_{\cdot j}\|_2 \leq C$, and for all $1 \leq k \leq p$, $\|\mathbf{v}_k\|_2 \leq C$, then $\mathbf{X}_{ia(r)}^\top \mathbf{v}_k$ is sub-Gaussian random variable with parameter bounded by some absolute constant. On the other hand, by the Lasso bound (24) for the columns of $\widehat{\mathbf{\Gamma}}_{r,k}$, we have $\|\mathbf{u}_j\|_2 \leq \sqrt{s \log p / (n + N)}$, so that conditional on $\widehat{\mathbf{\Gamma}}_{r,k}$, $\mathbf{u}_j^\top \mathbf{X}_{iJ(r,k)}$ is a sub-Gaussian random variable with parameter asymptotically bounded by $\sqrt{s \log p / (n + N)}$. In addition, with probability at least $1 - p^{-c}$, the mean value satisfies $|\frac{n}{n_r} \mathbf{u}_j^\top \mathbb{E} I\{\xi_i = r\} X_{iJ(r,k)} X_{ia(r)}^\top \mathbf{v}_k| \lesssim \sqrt{\frac{s \log p}{N+n}} \lambda_{\max}(\Sigma)$, where the last inequality follows from $\lambda_{\max}(\mathbb{E} I\{\xi_i = r\} X_i X_i^\top) = \sup_{\|u\|_2=1} \mathbb{E} I\{\xi_i = r\} (X_i^\top u)^2 \leq \lambda_{\max}(\Sigma)$. Thus, by concentration inequality for sub-exponential random variables,

$$P\left(\max_{\substack{1 \leq j \leq |a(k)| \\ 1 \leq k \leq p}} \left| \frac{1}{n} \sum_{i=1}^n \frac{n}{n_r} I\{\xi_i = r\} \mathbf{u}_j^\top X_{iJ(r,k)} X_{ia(r)}^\top \mathbf{v}_k \right| \leq C_1 \sqrt{\frac{s \log p}{N+n}} + C_2 \frac{\sqrt{s \log p}}{\sqrt{(n+N)n}} \right) \geq 1 - p^{-c}.$$

Similarly, one can also show that $P(\|\frac{1}{n} \sum_{i=1}^n \frac{n}{n_r} I\{\xi_i = r\} X_{ia(k)}^{(k)} (X_i^{(k)} - \widehat{X}_i^{(k)})^\top\|_\infty \leq C \sqrt{\frac{s \log p}{N+n}}) \geq 1 - p^{-c}$, and $P(\|\frac{1}{n} \sum_{i=1}^n \frac{n}{n_r} I\{\xi_i = r\} [\widehat{X}_{ia(k)}^{(k)} - X_{ia(k)}^{(k)}] (X_i^{(k)} - \widehat{X}_i^{(k)})^\top\|_\infty \leq C \sqrt{\frac{s \log p}{N+n}}) \geq 1 - p^{-c}$. Since $s \lesssim \frac{N+n}{\log p}$, we have shown (53) holds with high probability.

Part II. If we denote $\mathbf{H}_n = \frac{1}{n} \sum_{i=1}^n \frac{n}{n_r} I\{\xi_i = r\} X_{ia(k)}^{(k)} (X_i^{(k)})^\top$, we have

$$\begin{aligned}
 \inf_{\substack{\mathbf{u} \in E_s \\ \|\mathbf{u}_{a(k)}\|_2 \geq 1/2}} \left| \frac{\mathbf{u}_{a(k)}^\top \mathbf{H}_n \mathbf{u}}{\|\mathbf{u}_{a(k)}\|_2} \right| &\geq \inf_{\substack{\mathbf{u} \in E_s \\ \|\mathbf{u}_{a(k)}\|_2 \geq 1/2}} \frac{\mathbf{u}_{a(k)}^\top [\mathbf{H}_n]_{\cdot a(k)} \mathbf{u}_{a(k)}}{\|\mathbf{u}_{a(k)}\|_2} - \sup_{\substack{\mathbf{u} \in E_s \\ \|\mathbf{u}_{a(k)}\|_2 \geq 1/2}} \frac{|\mathbf{u}_{a(k)}^\top [\mathbf{H}_n]_{\cdot m(k)} \mathbf{u}_{m(k)}|}{\|\mathbf{u}_{a(k)}\|_2} \\
 &\geq \frac{1}{2} \inf_{\|\mathbf{v}\|_2=1, \|\mathbf{v}\|_1 \leq 4\sqrt{s}} |\mathbf{v}^\top [\mathbf{H}_n]_{\cdot a(k)} \mathbf{v}| - \sup_{\substack{\|\mathbf{v}\|_1 \leq 4\sqrt{s}, \|\mathbf{u}\|_1 \leq 2\sqrt{s} \\ \|\mathbf{v}\|_2=1, \|\mathbf{u}\|_2=1}} |\mathbf{v}^\top [\mathbf{H}_n]_{\cdot m(k)} \mathbf{u}|,
 \end{aligned}$$

where the last inequality follows that, for any $\mathbf{u} \in E_s$ such that $\|\mathbf{u}_{a(k)}\|_2 \geq 1/2$, $\|\frac{\mathbf{u}_{a(k)}}{\|\mathbf{u}_{a(k)}\|_2}\|_1 \leq 2\|\mathbf{u}_{a(k)}\|_1 \leq 2\|\mathbf{u}\|_1 \leq 4\sqrt{s}$, and $\|\mathbf{u}_{m(k)}\|_1 \leq \|\mathbf{u}\|_1 \leq 2\sqrt{s}$. If we can show that, with probability at least $1 - p^{-c}$,

$$\frac{1}{2} \inf_{\|\mathbf{v}\|_2=1, \|\mathbf{v}\|_1 \leq 4\sqrt{s}} |\mathbf{v}^\top [\mathbf{H}_n]_{\cdot a(k)} \mathbf{v}| > 3c, \tag{55}$$

and

$$\sup_{\substack{\|\mathbf{v}\|_1 \leq 4\sqrt{s}, \|\mathbf{u}\|_1 \leq 2\sqrt{s} \\ \|\mathbf{v}\|_2=1, \|\mathbf{u}\|_2=1}} |\mathbf{v}^\top [\mathbf{H}_n]_{.m(k)} \mathbf{u}| < 2c, \quad (56)$$

it follows that under the same event (22) holds. Suppose $\Sigma^{(k)} = \mathbb{E} \frac{n}{n_r} I\{\xi_i = r\} X_i^{(k)} (X_i^{(k)})^\top \in \mathbb{R}^{p \times p}$. In the following, we will show that, under the condition

$$\lambda_{\min}(\Sigma_{a(k), a(k)}^{(k)}) \geq 7c > c \geq \lambda_{\max}(\Sigma_{a(k), m(k)}^{(k)}), \quad (57)$$

inequalities (55) and (56) holds with the stated probability.

In fact, under condition (57), it can be shown through Theorem 2.5 of Zhou (2009) (see also Corollary 2.7 of Mendelson et al. (2007) or Theorem 2.1 of Mendelson et al. (2008)) that, for $s \ll \frac{n}{\log p}$ and $n \gg \log p$, with probability at least $1 - e^{-cn}$, (55) holds. Specifically, by Theorem 2.5 of Zhou (2009), the proof of (55) reduces to show that

$$\mathbb{E} \sup_{\|\mathbf{v}\|_2=1, \|\mathbf{v}\|_1 \leq 4\sqrt{s}} |\langle \mathbf{g}, [\Sigma_{a(k), a(k)}^{(k)}]^{1/2} \mathbf{v} \rangle| \lesssim \sqrt{s \log p} \ll \sqrt{n}, \quad (58)$$

where $\mathbf{g} \sim N(0, \mathbf{I}_p)$. To see this, simply note that

$$\mathbb{E} \sup_{\|\mathbf{v}\|_2=1, \|\mathbf{v}\|_1 \leq 4\sqrt{s}} |\langle \mathbf{g}, [\Sigma_{a(k), a(k)}^{(k)}]^{1/2} \mathbf{v} \rangle| \leq \sup_{\|\mathbf{v}\|_2=1, \|\mathbf{v}\|_1 \leq 4\sqrt{s}} \|\mathbf{v}\|_1 \cdot \mathbb{E} \| [\Sigma_{a(k), a(k)}^{(k)}]^{1/2} \mathbf{g} \|_\infty \lesssim \sqrt{s \log p},$$

where the last inequality follows from the following property of the maxima of sub-Gaussian random variables, $\mathbb{E} \max_{1 \leq i \leq n} g_i \leq C \sqrt{\sigma^2 \log n}$, where g_1, \dots, g_n are centered sub-Gaussian random variables with parameter σ^2 , not necessarily independent.

It remains to show (56). Note that

$$\begin{aligned} & \sup_{\substack{\|\mathbf{v}\|_1 \leq 4\sqrt{s}, \|\mathbf{u}\|_1 \leq 2\sqrt{s} \\ \|\mathbf{v}\|_2=1, \|\mathbf{u}\|_2=1}} |\mathbf{v}^\top [\mathbf{H}_n]_{.m(k)} \mathbf{u}| \\ & \leq \sup_{\substack{\|\mathbf{v}\|_1 \leq 4\sqrt{s}, \|\mathbf{u}\|_1 \leq 2\sqrt{s} \\ \|\mathbf{v}\|_2=1, \|\mathbf{u}\|_2=1}} |\mathbf{v}^\top [\mathbf{H}_n]_{.m(k)} \mathbf{u} - \mathbb{E} \mathbf{v}^\top [\mathbf{H}_n]_{.m(k)} \mathbf{u}| + \sup_{\substack{\|\mathbf{v}\|_1 \leq 4\sqrt{s}, \|\mathbf{u}\|_1 \leq 2\sqrt{s} \\ \|\mathbf{v}\|_2=1, \|\mathbf{u}\|_2=1}} \mathbb{E} \mathbf{v}^\top [\mathbf{H}_n]_{.m(k)} \mathbf{u} \\ & \leq \sup_{\substack{\|\mathbf{v}\|_1 \leq 4\sqrt{s}, \|\mathbf{u}\|_1 \leq 2\sqrt{s} \\ \|\mathbf{v}\|_2=1, \|\mathbf{u}\|_2=1}} |\mathbf{v}^\top [\mathbf{H}_n]_{.m(k)} \mathbf{u} - \mathbb{E} \mathbf{v}^\top [\mathbf{H}_n]_{.m(k)} \mathbf{u}| + \lambda_{\max}(\Sigma_{a(k), m(k)}^{(k)}). \end{aligned}$$

It suffices to show that, with probability at least $1 - p^{-c}$,

$$\sup_{\substack{\|\mathbf{v}\|_1 \leq 4\sqrt{s}, \|\mathbf{u}\|_1 \leq 2\sqrt{s} \\ \|\mathbf{v}\|_2=1, \|\mathbf{u}\|_2=1}} |\mathbf{v}^\top [\mathbf{H}_n]_{.m(k)} \mathbf{u} - \mathbb{E} \mathbf{v}^\top [\mathbf{H}_n]_{.m(k)} \mathbf{u}| \leq c. \quad (59)$$

To see this, note that

$$\begin{aligned} & \sup_{\substack{\|\mathbf{v}\|_1 \leq 4\sqrt{s}, \|\mathbf{u}\|_1 \leq 2\sqrt{s} \\ \|\mathbf{v}\|_2=1, \|\mathbf{u}\|_2=1}} |\mathbf{v}^\top [\mathbf{H}_n]_{.m(k)} \mathbf{u} - \mathbb{E} \mathbf{v}^\top [\mathbf{H}_n]_{.m(k)} \mathbf{u}| \\ & \leq \sup_{\substack{\|\mathbf{v}\|_1 \leq 4\sqrt{s}, \|\mathbf{u}\|_1 \leq 2\sqrt{s} \\ \|\mathbf{v}\|_2=1, \|\mathbf{u}\|_2=1}} \|\mathbf{v}\|_1 \|\mathbf{u}\|_1 \cdot \| [\mathbf{H}_n]_{.m(k)} - \mathbb{E} [\mathbf{H}_n]_{.m(k)} \|_\infty, \end{aligned}$$

and $\sup_{\substack{\|\mathbf{v}\|_1 \leq 4\sqrt{s}, \|\mathbf{u}\|_1 \leq 2\sqrt{s} \\ \|\mathbf{v}\|_2=1, \|\mathbf{u}\|_2=1}} \|\mathbf{u}\|_1 \|\mathbf{v}\|_1 \leq 8s$. Then by the concentration inequality for independent sub-exponential random variables, we have, for any $(i, j) \in \{1, \dots, |a(k)|\} \times m(k)$, $P(|[\mathbf{H}_n]_{ij} - [\mathbb{E}\mathbf{H}_n]_{ij}| \leq C\sqrt{\frac{\log p}{n}}) \geq 1 - p^{-c}$, so that $P(\|\mathbf{H}_n - \mathbb{E}\mathbf{H}_n\|_\infty \leq C\sqrt{\frac{\log p}{n}}) \geq 1 - p^{-c}$. Then as long as $s \lesssim \sqrt{n/\log p}$, we have (59) hold with probability at least $1 - p^{-c}$. This completes the proof of the proposition.

D.5 Proof of Lemma 7

By the theory of inhomogeneous system of linear equations, $\mathbb{E}G^\top v_j = e_j$ has infinitely numbers of solutions if $\text{rank}(\mathbb{E}G) = \text{rank}([\mathbb{E}G, e_j]) < M'_g$. When the smallest singular value of $\mathbb{E}G$ is nonnegative, we have $\text{rank}(\mathbb{E}G) = p$, and the matrix $[\mathbb{E}G, e_j] \in \mathbb{R}^{p \times (M'_g+1)}$ is also of rank p , which by assumption is smaller than M'_g . Secondly, note that $1 = \|e_j\| = \|\mathbb{E}G^\top v_j\| \geq \lambda_{\min}(\mathbb{E}G)\|v_j\|_2$. Then as long as $\lambda_{\min}(\mathbb{E}G) \geq c > 0$, we have $\|v_j\|_2 \leq C$.

D.6 Proof of Lemma 8

Since $\|(v_j^*)^\top G_n - e_j^\top\|_\infty \leq \|(v_j^*)^\top (G_n - G)\|_\infty + \|(v_j^*)^\top G - e_j^\top\|_\infty$, it suffices to show that

$$\|(v_j^*)^\top G - e_j^\top\|_\infty \lesssim \sqrt{\frac{\log p}{n + N}}, \quad (60)$$

and

$$\|(v_j^*)^\top (G_n - G)\|_\infty = \max_{1 \leq j \leq p} |(v_j^*)^\top [G_n - G]_{\cdot j}| \lesssim R^2 \sqrt{\frac{s \log p}{N + n}}, \quad (61)$$

holds with probability at least $1 - p^{-c}$.

To prove (60), we use the similar argument as in the proof of Lemma 4. We notice that each component of $(v_j^*)^\top G$ is an average of n independent sub-exponential random variables and $\mathbb{E}(v_j^*)^\top G = e_j^\top$. Then by concentration inequality for sub-exponential random variables (Vershynin, 2010), inequality (60) holds with probability at least $1 - p^{-c}$.

The rest of the proof is devoted to (61). Let $h_{irk}(\beta) = (h_{irk\ell}(\beta))_{\ell \in a(k)}$. We can write $G_{\cdot j} = \left[\frac{1}{N_r + n_r} \sum_{i=1}^{N+n} \frac{\partial h'_{irk\ell}(\beta)}{\partial \beta_j} \right]_{1 \leq r \leq R, k \in \mathcal{G}(r), \ell \in a(k)}$, where the M_g elements are ordered first by the index r , and then k , and finally ℓ . Now for each (r, k, ℓ) , we have $\frac{\partial h'_{irk\ell}(\beta)}{\partial \beta_j} = -I(\xi_i = r) X_{ij}^{(k)} X_{i\ell}$. If j -th variable is not observed in i -th sample, we have $\frac{\partial h'_{irk\ell}(\beta)}{\partial \beta_j} = -I(\xi_i = r) \gamma_{j, J(r, k)}^\top X_{iJ(r, k)} X_{i\ell}$, where we used $X_{ij}^{(k)} = \gamma_{j, J(r, k)}^\top X_{iJ(r, k)}$. If j -th variable is observed in i -th sample, we have $\frac{\partial h'_{irk\ell}(\beta)}{\partial \beta_j} = -I(\xi_i = r) X_{ij} X_{i\ell}$. Similarly, if we write $[G_n]_{\cdot j} = \left[\frac{1}{N_r + n_r} \sum_{i=1}^{N+n} \frac{\partial \hat{h}'_{irk\ell}(\beta)}{\partial \beta_j} \right]_{1 \leq r \leq R, k \in \mathcal{G}(r), \ell \in a(k)}$, where $\frac{\partial \hat{h}'_{irk\ell}(\beta)}{\partial \beta_j} = -I(\xi_i = r) \hat{\gamma}_{j, J(r, k)}^\top X_{iJ(r, k)} X_{i\ell}$, if j -th variable is not observed in i -th sample, and $\frac{\partial h_{irk\ell}(\beta)}{\partial \beta_j} = -I(\xi_i = r) X_{ij} X_{i\ell}$, if j -th variable is observed in i -th sample. Thus, we have

$$[G - G_n]_{\cdot j} = \left[\frac{1}{N_r + n_r} \sum_{i=1}^{N+n} \left(\frac{\partial h'_{irk\ell}(\beta)}{\partial \beta_j} - \frac{\partial \hat{h}'_{irk\ell}(\beta)}{\partial \beta_j} \right) \right]_{1 \leq r \leq R, k \in \mathcal{G}(r), \ell \in a(k)}. \quad (62)$$

If the j -th variable is not observed in i -th sample, we have

$$\frac{\partial \hat{h}'_{irk}(\beta)}{\partial \beta_j} - \frac{\partial h'_{irk}(\beta)}{\partial \beta_j} = -I(\xi_i = r)(\hat{\gamma}_{j,J(r,k)} - \gamma_{j,J(r,k)})^\top X_{iJ(r,k)} X_{iJ(r,k)} \quad (63)$$

In the following we control the left hand side of (61) for j not observed in Group r . Specifically, let $v_{j,rk}^*$ be the subvector of v_j^* associated to Group $k \in \mathcal{G}(r)$. We have, with probability at least $1 - p^{-c}$,

$$\begin{aligned} & \sum_{r=1}^R \sum_{k \in \mathcal{G}(r)} \frac{1}{N_r + n_r} \sum_{i=1}^{N+n} I(\xi_i = r) (\hat{\gamma}_{j,J(r,k)} - \gamma_{j,J(r,k)})^\top X_{iJ(r,k)} X_{iJ(r,k)}^\top v_{j,rk}^* \\ & \leq \sum_{r=1}^R \sum_{k \in \mathcal{G}(r)} \sqrt{\frac{1}{N+n} \sum_{i=1}^{N+n} \left[\frac{N+n}{N_r + n_r} I(\xi_i = r) X_{iJ(r,k)}^\top (\hat{\gamma}_{j,J(r,k)} - \gamma_{j,J(r,k)}) \right]^2} \\ & \quad \times \sqrt{\frac{1}{N+n} \sum_{i=1}^{N+n} I(\xi_i = r) (X_{iJ(r,k)}^\top v_{j,rk}^*)^2} \\ & \lesssim \sum_{r=1}^R \sum_{k \in \mathcal{G}(r)} \sqrt{\frac{1}{n+N} \sum_{i=1}^{n+N} \left[X_{iJ(r,k)}^\top (\hat{\gamma}_{j,J(r,k)} - \gamma_{j,J(r,k)}) \right]^2} \sqrt{\frac{1}{n+N} \sum_{i=1}^{n+N} I(\xi_i = r) (X_{iJ(r,k)}^\top v_{j,rk}^*)^2} \\ & \lesssim R^2 \sqrt{\frac{s \log p}{N+n}} \end{aligned}$$

where the last inequality follows from the standard Lasso bound

$$P\left(\frac{1}{n+N} \sum_{i=1}^{n+N} \left[X_{iJ(r,k)}^\top (\hat{\gamma}_{j,J(r,k)} - \gamma_{j,J(r,k)}) \right]^2 \leq C \frac{s \log p}{N+n}\right) \geq 1 - p^{-c},$$

and the inequality

$$P\left(\max_{1 \leq r \leq R, k \in \mathcal{G}(r)} \frac{1}{n+N} \sum_{i=1}^{n+N} I(\xi_i = r) ((X_{ia(k)}^{(k)})^\top v_{j,rk}^*)^2 \leq C\right) \geq 1 - p^{-c}, \quad (64)$$

which can be established by

$$P\left(\max_{1 \leq r \leq R, k \in \mathcal{G}(r)} \left| \frac{1}{n+N} \sum_{i=1}^{n+N} (I(\xi_i = r) X_{iJ(r,k)}^\top v_{j,rk}^*)^2 - \mathbb{E} I(\xi_i = r) (X_{iJ(r,k)}^\top v_{j,rk}^*)^2 \right| \leq C \sqrt{\frac{\log p}{n+N}}\right) \geq 1 - p^{-c},$$

as a consequence of Proposition 5.16 of Vershynin (2010), and $\mathbb{E} I(\xi_i = r) (X_{iJ(r,k)}^\top v_{j,rk}^*)^2 \leq \|v_j^*\|_2^2 \lambda_{\max}(\Sigma) = O(1)$, where the last inequality follows from Lemma 7 and condition (A3).

If the j -th variable observed in i -th sample, we have

$$\frac{\partial h'_{irk\ell}(\beta)}{\partial \beta_j} - \frac{\partial \hat{h}'_{irk\ell}(\beta)}{\partial \beta_j} = 0 \quad (65)$$

Then combining the above two pieces, we have shown that (61) holds.

D.7 Proof of Lemma 9

By mean value theorem, we have

$$\hat{S}_j(\hat{\beta}_j^*) = \hat{v}_j^\top g_n^*(\hat{\beta}_j^*) = \hat{v}_j^\top g_n^*(\beta) + \hat{v}_j^\top G_n(\hat{\beta}_j^* - \beta). \quad (66)$$

To control the second term in (66), we note that, by Theorem 1, with probability at least $1 - p^{-c}$,

$$|\hat{v}_j^\top G_n(\hat{\beta}_j^* - \beta)| \leq \|[\hat{v}_j^\top G_n]_{-j}\|_\infty \|\hat{\beta}_j^* - \beta\|_1 \lesssim s\lambda'\lambda.$$

Thus, (33) is proved if $s\lambda'\lambda\sqrt{n} \rightarrow 0$, or, under the specific choices of the tuning parameters λ and λ' , we have $\frac{s^{3/2}\log p}{\sqrt{N+n}} + \frac{s^{5/2}\sqrt{n}\log p}{N+n} \rightarrow 0$, or $s \ll \min\left\{\left(\frac{N+n}{\log^2 p}\right)^{1/3}, \left(\frac{(N+n)^2}{n\log^2 p}\right)^{1/5}\right\}$.

D.8 Proof of Lemma 10

By definition

$$\begin{aligned} \hat{S}_j(\beta) &= \sum_{k \in \mathcal{G}(r), 1 \leq r \leq R} \frac{1}{n} \sum_{i=1}^n \hat{v}_{rk}^\top \hat{h}'_{irk}(\beta) \\ &= \frac{1}{n} \sum_{i=1}^n \sum_{k \in \mathcal{G}(r), 1 \leq r \leq R} \frac{n}{n_r} I\{\xi_i = r\} [(X_i - \hat{X}_i^{(k)})^\top \beta + \epsilon_i] \hat{v}_{rk}^\top X_{iJ(r,k)}. \end{aligned}$$

Let

$$\eta_i = \sum_{k \in \mathcal{G}(r), 1 \leq r \leq R} \frac{n}{n_r} I\{\xi_i = r\} [(X_i - \hat{X}_i^{(k)})^\top \beta + \epsilon_i] \hat{v}_{rk}^\top X_{iJ(r,k)}.$$

We have

$$\mathbb{E}[\eta_i | \hat{X}] = \sum_{k \in \mathcal{G}(r), 1 \leq r \leq R} \frac{n}{n_r} I\{\xi_i = r\} (X_i - \hat{X}_i^{(k)})^\top \beta \hat{v}_{rk}^\top X_{iJ(r,k)} \equiv \mu_{ji},$$

and

$$\text{Var}(\eta_i | \hat{X}) = \sigma^2 \sum_{k \in \mathcal{G}(r), 1 \leq r \leq R} \frac{n^2}{n_r^2} I\{\xi_i = r\} [\hat{v}_{rk}^\top X_{iJ(r,k)}]^2 \quad (67)$$

By the central limit theorem, we have the desired result.

Appendix E. Discussion of the Eigenvalue Condition

Note that

$$X_i^{(k)} = ((X_{ia(r)}^{(k)})^\top, (X_{im(r)}^{(k)})^\top)^\top = (X_{ia(r)}^\top, X_{iJ(r,k)}^\top \mathbf{\Gamma}_{r,k})^\top = \begin{bmatrix} \mathbf{I}_{|a(r)|} \\ [\mathbf{\Gamma}_{r,k}^\top, 0] \end{bmatrix} X_{ia(r)}. \quad (68)$$

It follows that

$$\begin{aligned}
 \Sigma^{(r,k)} &= \mathbb{E}I\{\xi_i = r\} X_i^{(k)} (X_i^{(k)})^\top \\
 &= \mathbb{E}I\{\xi_i = r\} \begin{bmatrix} \mathbf{I}_{|a(r)|} \\ \mathbf{\Gamma}_{r,k}^\top, 0 \end{bmatrix} X_{ia(r)} X_{ia(r)}^\top \begin{bmatrix} \mathbf{I}_{|a(r)|} & \\ & \mathbf{\Gamma}_{r,k}^\top, 0 \end{bmatrix}^\top \\
 &= \begin{bmatrix} \mathbf{I}_{|a(r)|} \\ \mathbf{\Gamma}_{r,k}^\top, 0 \end{bmatrix} \mathbf{C}_{a(r),a(r)}^* \begin{bmatrix} \mathbf{I}_{|a(r)|} & \\ & \mathbf{\Gamma}_{r,k}^\top, 0 \end{bmatrix}^\top \\
 &= \begin{bmatrix} \mathbf{C}_{a(r),a(r)}^* & \mathbf{C}_{a(r),J(r,k)}^* \mathbf{\Gamma}_{r,k} \\ \mathbf{\Gamma}_{r,k}^\top \mathbf{C}_{J(r,k),a(r)}^* & \mathbf{\Gamma}_{r,k}^\top \mathbf{C}_{J(r,k),J(r,k)}^* \mathbf{\Gamma}_{r,k} \end{bmatrix} \tag{69}
 \end{aligned}$$

It can be seen that condition (57) essentially reduces to that on $\mathbf{C}^* = \mathbb{E}I\{\xi_i = r\} X_i X_i^\top$ and $\mathbf{\Gamma}_{r,k}$. To see that there exists situation where the condition (57) holds, it suffices to consider the case where $\|\mathbf{\Gamma}_{r,k}\| \rightarrow 0$ or $\|[\Sigma^{-1}]_{m(r),J(r,k)}\| \rightarrow 0$ as $(n, p) \rightarrow \infty$, and the missing is completely at random with missing proportion $\mathbb{E}I\{\xi_i = r\} \geq c_0 > 0$. In this case, there exists some $c > 0$ such that $\lambda_{\min}(\Sigma_{a(k),a(k)}^{(r,k)}) \geq 7c > c \geq \lambda_{\max}(\Sigma_{a(k),m(k)}^{(r,k)})$.

Appendix F. The FDR Control Procedure

To test the simultaneous hypotheses

$$H_{0j} : \beta_j = 0 \quad \text{vs} \quad H_{aj} : \beta_j \neq 0, \quad 1 \leq j \leq p,$$

we apply the modified BH procedure of Ma et al. (2020). Specifically, we reject the null hypothesis H_{0j} at significance level α if $|T_{0j}| \geq \hat{t}$, where

$$\hat{t} = \inf \left\{ 0 \leq t \leq b_p : \frac{p\{2 - 2\Phi(t)\}}{\max\{\sum_{j=1}^p I(|T_{0j}| \geq t), 1\}} \leq \alpha \right\}, \tag{70}$$

$b_p = \sqrt{2 \log p - 2 \log \log p}$, $T_{0j} = n\tilde{\beta}_j/\hat{s}_j$, and $\Phi(t)$ is the cumulative distribution function of the standard normal distribution.

References

- David Azriel, Lawrence D Brown, Michael Sklar, Richard Berk, Andreas Buja, and Linda Zhao. Semi-supervised linear regression. *arXiv preprint arXiv:1612.02391*, 2016.
- Amanda N Baraldi and Craig K Enders. An introduction to modern missing data analyses. *Journal of School Psychology*, 48(1):5–37, 2010.
- Ales Bartos, David Gregus, Ibrahim Ibrahim, and Jaroslav Tintera. Brain volumes and their ratios in alzheimer s disease on magnetic resonance imaging segmented using freesurfer 6.0. *Psychiatry Research: Neuroimaging*, 287:70–74, 2019.
- Pierre C Bellec, Arnak S Dalalyan, Edwin Grappin, and Quentin Paris. On the prediction loss of the lasso in the partially labeled setting. *Electronic Journal of Statistics*, 12(2): 3443–3472, 2018.

- Yoav Benjamini and Daniel Yekutieli. The control of the false discovery rate in multiple testing under dependency. *Annals of Statistics*, pages 1165–1188, 2001.
- Peter J Bickel, Ya'acov Ritov, and Alexandre B Tsybakov. Simultaneous analysis of Lasso and Dantzig selector. *The Annals of Statistics*, 37(4):1705–1732, 2009.
- T. Tony Cai and Zijian Guo. Semi-supervised inference for explained variance in high-dimensional regression and its applications. *Journal of the Royal Statistical Society: Series B*, 82:391–419, 2020.
- Tianxi Cai, T Tony Cai, and Anru Zhang. Structured matrix completion with applications to genomic data integration. *Journal of the American Statistical Association*, 111(514):621–633, 2016.
- Sergio Campos, Luis Pizarro, Carlos Valle, Katherine R. Gray, Daniel Rueckert, and Héctor Allende. Evaluating imputation techniques for missing data in ADNI: A patient classification study. In *Progress in Pattern Recognition, Image Analysis, Computer Vision, and Applications*, pages 3–10. Springer International Publishing, 2015. ISBN 978-3-319-25751-8.
- Emmanuel Candes and Terence Tao. The dantzig selector: Statistical estimation when p is much larger than n . *The Annals of Statistics*, 35(6):2313–2351, 2007.
- Abhishek Chakraborty and Tianxi Cai. Efficient and adaptive linear regression in semi-supervised settings. *The Annals of Statistics*, 46(4):1541–1572, 2018.
- Kimberly R Chapman, Hanaan Bing-Canar, Michael L Alosco, Eric G Steinberg, Brett Martin, Christine Chaisson, Neil Kowall, Yorghos Tripodis, and Robert A Stern. Mini mental state examination and logical memory scores for entry into alzheimer's disease trials. *Alzheimer's research & therapy*, 8(1):9, 2016.
- GTEX Consortium et al. Genetic effects on gene expression across human tissues. *Nature*, 550(7675):204–213, 2017.
- Siyi Deng, Yang Ning, Jiwei Zhao, and Heping Zhang. Optimal semi-supervised estimation and inference for high-dimensional linear regression. *arXiv preprint arXiv:2011.14185*, 2020.
- Pilar Esteve, Javier Rueda-Carrasco, María Inés Mateo, María Jesús Martín-Bermejo, Jonathan Draffin, Guadalupe Pereyra, África Sandonís, Inmaculada Crespo, Inmaculada Moreno, Ester Aso, et al. Elevated levels of secreted-frizzled-related-protein 1 contribute to alzheimer's disease pathogenesis. *Nature Neuroscience*, 22(8):1258–1268, 2019.
- Jianqing Fan and Jinchi Lv. Sure independence screening for ultrahigh dimensional feature space. *Journal of the Royal Statistical Society: Series B*, 70(5):849–911, 2008.
- Mengting Gong, Shoudong Ye, Wen-Xing Li, Jian Zhang, Yanjun Liu, Jie Zhu, Wenwen Lv, Hui Zhang, Jing Wang, Aiping Lu, et al. Regulatory function of praja ring finger ubiquitin ligase 2 mediated by the p2rx3/p2rx7 axis in mouse hippocampal neuronal cells. *American Journal of Physiology-Cell Physiology*, 318(6):C1123–C1135, 2020.

- Jessica L Gronsbell and Tianxi Cai. Semi-supervised approaches to efficient evaluation of model prediction performance. *Journal of the Royal Statistical Society: Series B*, 80(3): 579–594, 2018.
- Adel Javanmard and Hamid Javadi. False discovery rate control via debiased lasso. *Electronic Journal of Statistics*, 13(1):1212–1253, 2019.
- Adel Javanmard and Andrea Montanari. Confidence intervals and hypothesis testing for high-dimensional regression. *Journal of Machine Learning Research*, 15(1):2869–2909, 2014.
- Adel Javanmard and Andrea Montanari. Debiasing the lasso: Optimal sample size for gaussian designs. *The Annals of Statistics*, 46(6A):2593–2622, 2018.
- Kai Kang, Jingheng Cai, Xinyuan Song, and Hongtu Zhu. Bayesian hidden markov models for delineating the pathology of alzheimer’s disease. *Statistical Methods in Medical Research*, 28(7):2112–2124, 2019.
- Alexander Kautzky, Rene Seiger, Andreas Hahn, Peter Fischer, Wolfgang Krampla, Siegfried Kasper, Gabor G Kovacs, and Rupert Lanzenberger. Prediction of autopsy verified neuropathological change of alzheimer’s disease using machine learning and mri. *Frontiers in Aging Neuroscience*, 10:406, 2018.
- Isaac S Kohane. Using electronic health records to drive discovery in disease genomics. *Nature Reviews Genetics*, 12(6):417–428, 2011.
- Prosenjit Kundu, Runlong Tang, and Nilanjan Chatterjee. Generalized meta-analysis for multiple regression models across studies with disparate covariate information. *Biometrika*, 106(3):567–585, 2019.
- Christian Ledig, Andreas Schuh, Ricardo Guerrero, Rolf A Heckemann, and Daniel Rueckert. Structural brain imaging in alzheimer’s disease and mild cognitive impairment: biomarker analysis and shared morphometry database. *Scientific Reports*, 8(1):1–16, 2018.
- John Lonsdale, Jeffrey Thomas, Mike Salvatore, Rebecca Phillips, Edmund Lo, Saboor Shad, Richard Hasz, Gary Walters, Fernando Garcia, Nancy Young, et al. The genotype-tissue expression (gtex) project. *Nature genetics*, 45(6):580–585, 2013.
- Rong Ma, T. Tony Cai, and Hongzhe Li. Global and simultaneous hypothesis testing for high-dimensional logistic regression models. *Journal of the American Statistical Association*, pages 1–15, 2020.
- Shahar Mendelson, Alain Pajor, and Nicole Tomczak-Jaegermann. Reconstruction and subgaussian operators in asymptotic geometric analysis. *Geometric and Functional Analysis*, 17(4):1248–1282, 2007.
- Shahar Mendelson, Alain Pajor, and Nicole Tomczak-Jaegermann. Uniform uncertainty principle for bernoulli and subgaussian ensembles. *Constructive Approximation*, 28(3): 277–289, 2008.

- Susanne G Mueller, Michael W Weiner, Leon J Thal, Ronald C Petersen, Clifford Jack, William Jagust, John Q Trojanowski, Arthur W Toga, and Laurel Beckett. The alzheimer’s disease neuroimaging initiative. *Neuroimaging Clinics*, 15(4):869–877, 2005.
- Sahand Negahban and Martin J Wainwright. Restricted strong convexity and weighted matrix completion: Optimal bounds with noise. *The Journal of Machine Learning Research*, 13(1):1665–1697, 2012.
- Sahand Negahban, Pradeep Ravikumar, Martin J Wainwright, and Bin Yu. A unified framework for high-dimensional analysis of M-estimators with decomposable regularizers. *Technical Report Number 979*, 2010.
- Matey Neykov, Yang Ning, Jun S Liu, and Han Liu. A unified theory of confidence regions and testing for high-dimensional estimating equations. *Statistical Science*, 33(3):427–443, 2018.
- Yang Ning and Han Liu. A general theory of hypothesis tests and confidence regions for sparse high dimensional models. *The Annals of Statistics*, 45(1):158–195, 2017.
- Garvesh Raskutti, Martin J Wainwright, and Bin Yu. Restricted eigenvalue properties for correlated gaussian designs. *The Journal of Machine Learning Research*, 11:2241–2259, 2010.
- Gretel Sanabria-Diaz, Eduardo Martínez-Montes, and Lester Melie-Garcia. Glucose metabolism during resting state reveals abnormal brain networks organization in the alzheimer’s disease and mild cognitive impairment. *PloS one*, 8(7):e68860, 2013.
- Robert Tibshirani. Regression shrinkage and selection via the lasso. *Journal of the Royal Statistical Society: Series B*, 58(1):267–288, 1996.
- Sara van de Geer, Peter Bühlmann, YaŌacov Ritov, and Ruben Dezeure. On asymptotically optimal confidence regions and tests for high-dimensional models. *The Annals of Statistics*, 42(3):1166–1202, 2014.
- Roman Vershynin. Introduction to the non-asymptotic analysis of random matrices. *arXiv preprint arXiv:1011.3027*, 2010.
- Nicolas Verzelen. Minimax risks for sparse regressions: Ultra-high dimensional phenomena. *Electronic Journal of Statistics*, 6:38–90, 2012.
- Larry Wasserman and John D Lafferty. Statistical analysis of semi-supervised regression. In *Advances in Neural Information Processing Systems*, pages 801–808, 2008.
- Shuo Xiang, Lei Yuan, Wei Fan, Yalin Wang, Paul M Thompson, Jieping Ye, Alzheimer’s Disease Neuroimaging Initiative, et al. Bi-level multi-source learning for heterogeneous block-wise missing data. *NeuroImage*, 102:192–206, 2014.
- Fei Xue and Annie Qu. Integrating multisource block-wise missing data in model selection. *Journal of the American Statistical Association*, pages 1–14, 2020.

- Huanqing Yang, Hua Xu, Qingfeng Li, Yan Jin, Weixiong Jiang, Jinghua Wang, Yina Wu, Wei Li, Cece Yang, Xia Li, et al. Study of brain morphology change in alzheimer’s disease and amnesic mild cognitive impairment compared with normal controls. *General Psychiatry*, 32(2), 2019.
- Guan Yu, Qufeng Li, Dinggang Shen, and Yufeng Liu. Optimal sparse linear prediction for block-missing multi-modality data without imputation. *Journal of the American Statistical Association*, 115(531):1406–1419, 2020.
- Lei Yuan, Yalin Wang, Paul M Thompson, Vaibhav A Narayan, Jieping Ye, and Alzheimer’s Disease Neuroimaging Initiative. Multi-source feature learning for joint analysis of incomplete multiple heterogeneous neuroimaging data. *NeuroImage*, 61(3):622–632, 2012.
- Anru Zhang, Lawrence D Brown, and T Tony Cai. Semi-supervised inference: General theory and estimation of means. *The Annals of Statistics*, 47(5):2538–2566, 2019.
- Cun-Hui Zhang and Stephanie S Zhang. Confidence intervals for low dimensional parameters in high dimensional linear models. *Journal of the Royal Statistical Society: Series B*, 76(1):217–242, 2014.
- Linjun Zhang, Rong Ma, T Tony Cai, and Hongzhe Li. Estimation, confidence intervals, and large-scale hypotheses testing for high-dimensional mixed linear regression. *arXiv preprint arXiv:2011.03598*, 2020.
- Yuqian Zhang and Jelena Bradic. High-dimensional semi-supervised learning: in search for optimal inference of the mean. *arXiv preprint arXiv:1902.00772*, 2019.
- Zhongheng Zhang. Missing data imputation: Focusing on single imputation. *Annals of Translational Medicine*, 4(1):9, 2016.
- Qi Zhou, Mohammed Goryawala, Mercedes Cabrerizo, Warren Barker, Ranjan Duara, and Malek Adjouadi. Significance of normalization on anatomical mri measures in predicting alzheimer’s disease. *The Scientific World Journal*, 2014, 2014.
- Shuheng Zhou. Restricted eigenvalue conditions on subgaussian random matrices. *arXiv preprint arXiv:0912.4045*, 2009.