

ON THE EMERGENCE OF QUANTUM BOLTZMANN FLUCTUATION DYNAMICS NEAR A BOSE-EINSTEIN CONDENSATE

THOMAS CHEN AND MICHAEL HOTT

ABSTRACT. In this work, we study the quantum fluctuation dynamics in a Bose gas on a torus $\Lambda = (L\mathbb{T})^3$ that exhibits Bose-Einstein condensation, beyond the leading order Hartree-Fock-Bogoliubov (HFB) theory. Given a Bose-Einstein condensate (BEC) with density $N \gg 1$ surrounded by thermal fluctuations with density 1, we assume that the system dynamics is generated by a Hamiltonian with mean-field scaling. We derive a quantum Boltzmann type dynamics from a second-order Duhamel expansion upon subtracting both the BEC dynamics and the HFB dynamics, with rigorous error control. Given a quasifree initial state, we determine the time evolution of the centered correlation functions $\langle a \rangle$, $\langle aa \rangle - \langle a \rangle^2$, $\langle a^+ a \rangle - |\langle a \rangle|^2$ at mesoscopic time scales $t \sim \lambda^{-2}$, where $0 < \lambda \ll 1$ is the coupling constant determining the HFB interaction, and a, a^+ denote annihilation and creation operators. While the BEC and the HFB fluctuations both evolve at a microscopic time scale $t \sim 1$, the Boltzmann dynamics is much slower, by a factor λ^2 . For large but finite N , we consider both the case of fixed system size $L \sim 1$, and the case $L \sim \lambda^{-2-}$. In the case $L \sim 1$, we show that the Boltzmann collision operator contains subleading terms that can become dominant, depending on time-dependent coefficients assuming particular values in \mathbb{Q} ; this phenomenon is reminiscent of the Talbot effect. For the case $L \sim \lambda^{-2-}$, we prove that the collision operator is well approximated by the expression predicted in the literature. In either of those cases, we have $\lambda \sim \left(\frac{\log \log N}{\log N} \right)^\alpha$, for different values of $\alpha > 0$.

CONTENTS

1. Introduction	2
1.1. Quantum dynamics and Boltzmann equations	2
1.2. Collisions of Fluctuations about a Bose-Einstein Condensate	5
1.3. Definition of the mathematical model	6
1.4. Leading order condensate dynamics: Hartree equation	8
1.5. Leading order fluctuation dynamics: HFB equations	9
1.6. Fourier transform	10
1.7. Lower order fluctuations: Emergence of Boltzmann dynamics	11
1.8. Fluctuations beyond HFB	16
1.9. Scaling of λ and N	16
2. Main Results	17
2.1. Notation	17
2.2. Assumptions	19
2.3. Statement of results	20
2.4. Sketch of the proof	24
3. Preliminaries	25

4. Trace estimates	28
4.1. Computation of the partition function	29
4.2. Operator estimates w.r.t. \mathcal{N}_b	32
5. Control of error terms in the expansion	39
5.1. Fixed, N -independent lattice $\Lambda^* \cong \mathbb{Z}^3$	44
5.2. Continuum approximation $\Lambda^* \rightarrow \mathbb{R}^3$	59
5.3. Centered expectations	70
6. Main order terms in the evolution of (Φ, f, g)	75
6.1. Discrete case	75
6.2. Continuum approximation	77
7. Effective equations	80
7.1. Discrete case	80
7.2. Continuum approximation	83
Appendix A. Calculus for creation and annihilation operators	85
Appendix B. Propagation of approximate restricted quasifreeness	91
Appendix C. Proof of convergence to mean field equations	98
C.1. Error bounds due to HFB evolution	103
C.2. Mollified energy conservation	104
References	108

1. INTRODUCTION

1.1. Quantum dynamics and Boltzmann equations. The main question we set out to answer in this work is:

Is there a scaling regime in an interacting quantum field theory, for which the emergence of collisional processes described by a Boltzmann equation can be rigorously established?

1.1.1. *Emergence of a quantum Boltzmann equation.* In analogy to Maxwell's and Boltzmann's theory of collisions in classical systems, Nordheim [251] in 1928 was the first to propose a Boltzmann equation for Bose and Fermi gases given by

$$\begin{aligned} \partial_t f(p) = Q_4(f) := & \int d\mathbf{p}_4 \delta(p_1 + p_2 - p_3 - p_4) \delta(E(p_1) + E(p_2) - E(p_3) - E(p_4)) \\ & |\mathcal{M}_{22}(\mathbf{p}_4)|^2 (\delta(p - p_1) + \delta(p - p_2) - \delta(p - p_3) - \delta(p - p_4)) \\ & ((1 \pm f(p_1))(1 \pm f(p_2))f(p_3)f(p_4) - f(p_1)f(p_2)(1 \pm f(p_3))(1 \pm f(p_4))). \end{aligned} \quad (1.1)$$

Here, f denotes the particle density in the spatially homogeneous case; '+' refers to the bosonic, and '-' refers to the fermionic equation, and $\mathbf{p}_4 = (p_1, p_2, p_3, p_4)$. In addition, $E(p) = \frac{1}{2}|p|^2$ denotes the free dispersion, and \mathcal{M}_{22} is the (microscopic) scattering cross section for $2 \leftrightarrow 2$ processes describing two thermal fluctuation scattering off of each other. As Nordheim already argues, the distribution of the outgoing particles needs to be taken into account, resulting in a quartic collision operator, in contrast to classical particles that are described by a quadratic collision operator. It is shown in [251] that the equilibrium is given by the Bose-Einstein, respectively the Fermi-Dirac statistics, and that an H -theorem holds true. In 1933, Uehling and Uhlenbeck [297] studied the linearization about the equilibrium, in

order to determine the associated hydrodynamics, and to compute the heat conductivity and the viscosity coefficient.

Subsequently, physicists have given formal derivations of the above quantum Boltzmann equation, using diagrammatic techniques from quantum field theory, see, e.g., [1, 188, 300]. This has given rise to interesting fundamental effective theories, such as the *Kadanoff-Baym equations*, see, e.g., [181, 277]. We also mention the important contributions by Bogoliubov and collaborators [63] and Prigogine and collaborators [262].

The first mathematically rigorous works on the derivation of the classical Boltzmann equation, a billiards model for a classical gas, go back to Cercignani [79] in 1972 and Lanford [202] in 1975, where they studied the Grad-limit [161] of a hard-sphere model. These works were later revisited and completed through works by Uchiyama [296], Cercignani-Illner-Pulvirenti [81], Spohn [285], Cercignani-Gerasimenko-Petrina [80], and Gallagher-Saint-Raymond-Texier [149].

In 1983, Hugenholtz [184] considered the commutator perturbation expansion with respect to the weak coupling constant λ . Implementing the kinetic time scale $t = T\lambda^{-2}$ used by van Hove [299], it was shown that, in the translation-invariant case, terms of order $O(\lambda)$ vanish as $\lambda \rightarrow 0$, and that terms of order $O(\lambda^2)$ are proportional to T . Using a selection rule, it is conjectured in [184] that only two-point correlations of higher orders in λ survive, motivating the assumption of *quasifreeness*. Quasifreeness, as we will see below, is, in a sense, a quantum analogue to the 'Stoßzahlansatz', also known as 'molecular chaos'. Hence, Hugenholtz argues, at leading order, the Boltzmann equation should arise for the evolution of the two-point function. Ho and Landau [179] later proved that, to second order in λ , this holds true.

In 2004, Erdős, Salmhofer, and Yau [129] extended the results by Hugenholtz, and by Ho and Landau, by introducing the concept of *restricted quasi-freeness*, i.e., quasi-freeness only up to six- or eight-point correlations. Assuming propagation of restricted quasi-freeness, they showed that a (time-dependent) Boltzmann equation arises from the second-order Duhamel expansion, under certain assumptions. Around the same time, Benedetto, Castella, Esposito and Pulvirenti, began a series of works that used Bogoliubov-Born-Green-Kirkwood-Yvon (BBGKY) hierarchies to derive a quadratic Boltzmann equation with a quantum collision kernel for a system obtained by iterating Duhamel's formula, ignoring the tail, and truncating the obtained hierarchy in the small-coupling limit [42, 45] respectively in the low-density regime [44]. In [43], they went on to show that the contributions to second order in the coupling recover (1.1) to leading order, evaluated at initial time with f_0 instead of f_t . Further works studying the BBGKY hierarchy in the context of the quantum Boltzmann equation include works by Gerasimenko and collaborators [152, 153, 155], see also references therein. A recent review in this direction can be found in [154]. For works using a second quantization approach, we refer to Spohn and collaborators [148, 234, 287, 288, 289], see also references therein.

In 2015, X. Chen and Y. Guo [94] showed that, if the marginals of the BBGKY hierarchy converge in the weak-coupling limit in a strong sense, and if the $W^{4,1}$ -regularity per particle remains uniformly bounded for some positive time, then the limiting hierarchy is that associated to the quadratic Boltzmann equation with a quantum collision kernel, instead of the Boltzmann equation derived in [129].

In a different line of work addressing the weakly disordered Anderson model, scattering of electrons at impurities in a lattice have been investigated extensively. For works studying the derivation of the linear Boltzmann equation in this context, we refer to [82, 90, 91, 134, 178, 180, 283]. For a more recent treatment, we refer to [165].

Lukkarinen-Spohn [234, 235] showed that the nonlinear Schrödinger equation (NLS) with random initial data leads to the wave kinetic equation: They present this system as a simplified model to gain insights into the emergence of Boltzmann-type dynamics in a quantum Bose gas. The study of wave-turbulence in the context of the long-time behavior of the NLS is a very active area of research, see, e.g., [18, 75, 107, 108, 113, 114, 143, 156, 191, 270].

The derivation of Boltzmann equations from a system of classical interacting particles is an extraordinarily active research field, see [20, 21, 58, 59, 60, 61, 80, 81, 115, 149, 193, 202, 253, 264, 265, 266, 278, 284, 296] and references therein. The methods differ vastly from the quantum field theoretic approach developed in the work at hand. For works studying well-posedness and other analytical properties of the classical Boltzmann equation, we refer to [4, 5, 6, 7, 8, 10, 13, 19, 25, 53, 54, 41, 81, 83, 84, 116, 118, 119, 123, 125, 158, 160, 163, 170, 172, 173, 185, 189, 238, 292, 291, 293, 298, 301, 302, 303] and references therein.

1.1.2. *Well-posedness.* The first well-posedness results for (1.1) go back to Dolbeaut [124] and Lions [219] for the fermionic case. X. Lu and collaborators have made significant progress on the fermionic Boltzmann-Uhlenbeck-Uehling (BUU) equation, see, e.g., [222, 226, 231]. For a recent work on the fermionic problem, we refer to [213].

A unified treatment of bosons and fermions can be found in [305, 306], and more recently [252]. In addition, generalized statistics such as anyons [28], and Haldane statistics [26, 30] have also been included in this line of study.

Works studying the relativistic quantum Boltzmann equation include [36, 136, 137] and references therein.

The quantum Landau equation, which can be viewed as a limit of the Boltzmann equation accounting for long-range interactions, has been studied in [11, 12, 37, 38, 205, 220]. Recently, He-Lu-Pulvirenti [175] showed that it can be obtained as a weak semi-classical limit from the quantum Boltzmann equation.

In 1924, Bose [64], and, independently in 1925, Einstein [127] predicted that, below a critical temperature, the ground state becomes gradually more populated, forming a macroscopic state called the *Bose-Einstein condensate* (BEC). In 1995, this phenomenon was independently experimentally verified by groups around Cornell and Wiemann [24] and Ketterle [111]. Both groups were awarded the 2001 Physics Nobel Prize.

In the mathematically rigorous PDE literature, the bosonic problem has been investigated first by X. Lu [221]. The long-time behavior for radial initial data was studied, obtaining global existence, local stability, conservation of energy, and estimates on moment production. Moreover, at low temperatures, it is shown in [221] that a solution concentrates at $p = 0$ for large time, and that, at high temperatures, the solution converges weakly to the Bose-Einstein distribution. Escobedo-Mischler-Valle [137] showed that bosonic entropy maximizers are given by

$$f_{eq}(p) = \frac{1}{e^{\beta(E(p)-\mu)} - 1} + m_0\delta(p), \quad (1.2)$$

where $m_0 \cdot \mu = 0$, see, e.g., [141]. The BEC density m_0 , the chemical potential μ and the inverse temperature β are uniquely determined by the moments $\int dp f_{eq}(p)$ and $\int dp E(p) f_{eq}(p)$, see, e.g., [223] and references therein. One has that m_0 vanishes above a critical temperature T_c , and is non-zero below T_c .

Well-posedness results have been formulated to account for solutions that form a Dirac mass at temperatures below T_c , and that stay bounded for temperatures above T_c . We refer to [26, 76, 141, 142, 223, 224, 227, 228, 229, 232] for the isotropic and space-homogeneous case, to [28, 29, 272, 252, 305, 306] for the space-dependent case, and to [73, 77, 212, 252] for the anisotropic case. Further works studying the blow-up behavior related to condensation include [40, 140, 290].

1.2. Collisions of Fluctuations about a Bose-Einstein Condensate. Pioneering works by Kirkpatrick and Dorfmann [195, 196] and Eckern [126] started analyzing the interplay of the BEC with thermal excitation cloud surrounding the condensate. They formally obtained a Boltzmann equation of the form

$$\partial_t f^{(ex)}(p) = n_c Q_3(f^{(ex)}) + Q_4(f^{(ex)}), \quad (1.3)$$

where Q_4 is given by (1.1) with E replaced by the Bogoliubov dispersion $\Omega = \sqrt{E(E + 2n_c\lambda)}$, and

$$\begin{aligned} Q_3(f) := & \int d\mathbf{p}_3 \delta(p_1 + p_2 - p_3) \delta(\Omega(p_1) + \Omega(p_2) - \Omega(p_3)) \\ & |\mathcal{M}_{21}(\mathbf{p}_3)|^2 (\delta(p - p_1) + \delta(p - p_2) - \delta(p - p_3)) \\ & ((1 + f(p_1))(1 + f(p_2))f(p_3) - f(p_1)f(p_2)(1 + f(p_3))). \end{aligned} \quad (1.4)$$

Here n_c denotes the BEC density, $f^{(ex)}$ the density of thermal fluctuation particles, $\bar{\lambda}$ is the coupling strength of the hard-sphere pair interaction $\bar{\lambda}\delta$, \mathcal{M}_{21} is the cross section for $2 \leftrightarrow 1$ processes describing collisions of 2 thermal particles, where one is either being absorbed into or emitted from the BEC, and $\mathbf{p}_3 = (p_1, p_2, p_3)$. Zaremba, Niguni and Griffin [304], see also [164], later extended their approach to include the dynamics of the condensate. In the *Hartree-Fock-Bogoliubov-Popov* approximation, and for a translation-invariant initial state, they formally argue that the condensate wave function Φ satisfies

$$\begin{aligned} i\partial_t \hat{\Phi}(p) = & (h_{Har} + 2\bar{\lambda}f^{(ex)}(p) - iQ_3(f^{(ex)}))\hat{\Phi}(p) \\ & + \bar{\lambda}g^{(ex)}(p)\bar{\Phi}(-p), \end{aligned} \quad (1.5)$$

and it is linked to the density via $n_c(t, x) = |\Phi_t(x)|^2$. Here h_{Har} denotes the Hartree Hamiltonian in momentum representation, see Section 1.4 below, and $g^{(ex)}$ denotes the rate of pair absorption into the condensate, which they discarded as a lower-order contribution. One of the motivations to study the coupled system between the condensate and the thermal cloud is to understand the nucleation process of the BEC, see, e.g., [52, 151, 187, 201, 257, 275]. For a review, we also refer to [263, 268].

Observe that, for large values of the condensate density n_c , we expect that Q_4 is of subleading order.

For mathematical works studying the system describing the two-component gas consisting of the condensate and the excitation cloud, we refer to [14, 27, 109, 110, 139, 250, 258, 259, 260, 279, 280].

1.3. Definition of the mathematical model. As a starting point for our analysis, we choose a single-species Bose gas at positive temperature trapped in a periodic box $\Lambda = (L\mathbb{T})^3$ of linear length L . We assume that the gas consists of two phases:

- (1) A Bose-Einstein condensate (BEC) with density N ,
- (2) thermal fluctuations with density ~ 1 .

We note that a significant part of this paper addresses the case of a fixed volume, where we may think of N as the number of bosons when $L = 1$. On the other hand, we will also consider the limit of large volume, where $N > 0$ will denote the number of bosons per unit volume (that is, the density); for convenience, we are not changing the notation. We assume mean-field interactions for which the kinetic energy of the condensate and the total (pair) interaction potential among particles are balanced, and analyze the interplay between the condensate dynamics and the dynamics of the fluctuation particles.

1.3.1. Definition of the model. Let $\mathcal{F} = \mathbb{C} \oplus \bigoplus_{n \geq 1} (L^2(\Lambda))^{\otimes_{sym} n}$ denote the bosonic Fock space, equipped with the inner product

$$\langle \Psi, \Phi \rangle_{\mathcal{F}} = \sum_{n \in \mathbb{N}_0} \langle \Psi_n, \Phi_n \rangle_{L^2(\mathbb{R}^{3n})} \quad (1.6)$$

for all $\Psi = (\Psi_n)_{n \in \mathbb{N}_0}$, $\Phi = (\Phi_n)_{n \in \mathbb{N}_0} \in \mathcal{F}$. For $n \in \mathbb{N}$, $f \in L^2(\mathbb{R}^3)$, $\Psi \in (L^2(\Lambda))^{\otimes_{sym} n}$, and $\mathbf{x}_{n-1} \in \mathbb{R}^{3(n-1)}$, we define

$$(a(f)\Psi)(\mathbf{x}_{n-1}) := \sqrt{n} \int_{\Lambda} dx \bar{f}(x) \Psi(x, \mathbf{x}_{n-1}). \quad (1.7)$$

For any $n \in \mathbb{N}$, let \mathcal{S}_n denote the permutation group of $\{1, \dots, n\}$. For $n \in \mathbb{N}_0$, $f \in L^2(\mathbb{R}^3)$, $\Psi \in (L^2(\Lambda))^{\otimes_{sym} n}$, and $\mathbf{x}_{n+1} = (x_1, \dots, x_{n+1}) \in \mathbb{R}^{3(n+1)}$, we define

$$(a^+(f)\Psi)(\mathbf{x}_{n+1}) := \frac{\sqrt{n+1}}{(n+1)!} \sum_{\pi \in \mathcal{S}_{n+1}} f(x_{\pi(1)}) \Psi(x_{\pi(2)}, \dots, x_{\pi(n+1)}). \quad (1.8)$$

Then we have that for all $f \in L^2(\Lambda)$, $\Phi, \Psi \in \mathcal{F}_{fin}$

$$\langle \Phi, a(f)\Psi \rangle_{\mathcal{F}} = \langle a^+(f)\Phi, \Psi \rangle_{\mathcal{F}}. \quad (1.9)$$

In addition, a and a^+ satisfy the canonical commutation relations (CCR) for any $f, g \in L^2(\Lambda)$

$$[a(f), a^+(g)] = \langle f, g \rangle, \quad [a(f), a(g)] = [a^+(f), a^+(g)] = 0. \quad (1.10)$$

Moreover, we introduce the operator-valued distributions a_x, a_y^+ by requiring

$$a(f) = \int_{\Lambda} dx \bar{f}(x) a_x, \quad (1.11)$$

$$a^+(g) = \int_{\Lambda} dx f(x) a_x^+ \quad (1.12)$$

for all $f, g \in L^2(\Lambda)$. These satisfy the CCR

$$[a_x, a_y^+] = \delta_{\Lambda}(x - y), \quad [a_x, a_y] = [a_x^+, a_y^+] = 0. \quad (1.13)$$

We call $\Omega_0 := (1, 0, 0, \dots) \in \mathcal{F}$ the *Fock vacuum*. Then we have that $a_x \Omega_0 = 0$ for all $x \in \Lambda$. In addition, for $\Psi \in \mathcal{F}_{fin}$, $x \in \Lambda$, $n \in \mathbb{N}$, and $\mathbf{x}_n \in \mathbb{R}^{3n}$, we have that

$$(a_x \Psi)^{(n)}(\mathbf{x}_n) = \sqrt{n+1} \Psi^{(n+1)}(x, \mathbf{x}_n). \quad (1.14)$$

We introduce the number operator

$$\mathcal{N}_b := \int_{\Lambda} dx a_x^\dagger a_x, \quad (1.15)$$

which satisfies

$$(\mathcal{N}_b \Psi)^{(n)} = n \Psi^{(n)}, \quad (1.16)$$

see, e.g., [50].

The Hamiltonian of the system studied in this paper has the following form. Let v be a sufficiently regular pair potential, see section 2.2, and $\lambda > 0$ a small coupling constant. Given $N > 0$, we define

$$\mathcal{H}_N := \frac{1}{2} \int_{\Lambda} dx a_x^\dagger (-\Delta_x) a_x + \frac{\lambda}{2N} \int_{\Lambda^2} dx dy v(x-y) a_x^\dagger a_y^\dagger a_y a_x. \quad (1.17)$$

Our aim is to study the evolution associated with the Hamiltonian \mathcal{H}_N given in (1.17). The condensate is accounted for by a coherent state $\mathcal{W}[\sqrt{N|\Lambda|}\phi_0]\Omega_0$, where

$$\mathcal{W}[f] := \exp(a^+(f) - a(f)) = \exp\left(\int_{\Lambda} dx (f(x)a_x^\dagger - \bar{f}(x)a_x)\right) \quad (1.18)$$

denotes the Weyl operator, and ϕ_0 is normalized, $\|\phi_0\|_{L^2(\Lambda)} = 1$. By the Baker-Campbell-Hausdorff formula,

$$\mathcal{W}[f]\Omega = e^{-\frac{1}{2}\|f\|_{L^2(\Lambda)}^2} \left(\frac{f^{\otimes n}}{\sqrt{n!}}\right)_{n \in \mathbb{N}_0}, \quad (1.19)$$

see, e.g., [50]. That is, in each fixed particle sector, the wave function is the product state determined by a single wave function f .

We assume that, at initial time $t = 0$, the system is described by a state

$$\nu_0(A) := \frac{\text{Tr}(e^{-\mathcal{K}}A)}{\text{Tr}(e^{-\mathcal{K}})} \quad (1.20)$$

for all observables $A \in \mathfrak{A}$, where \mathfrak{A} denotes the Weyl algebra generated by $\mathcal{W}[f]$, where $f \in \mathcal{S}(\Lambda)$ is a Schwartz function, see [70], section 5.2.3. In addition, the Baker-Campbell-Hausdorff formula yields

$$\mathcal{W}^*[f]a_x\mathcal{W}[f] = a_x + f(x). \quad (1.21)$$

Definition 1.1 (Quasifree state). *Let ν be a state and*

$$\nu^{(cen)}(A) := \nu(\mathcal{W}[\nu(a)]A\mathcal{W}^*[\nu(a)]) \quad (1.22)$$

denote its centering. We say ν is quasi-free iff

$$\begin{cases} \nu^{(cen)}(a^{\#1}a^{\#2} \dots a^{\#2n}) &= \overline{a^{\#1}a^{\#2} \dots a^{\#2n}} \\ &+ \text{all pair contractions}, \\ \nu^{(cen)}(a^{\#1}a^{\#2} \dots a^{\#2n-1}) &= 0 \end{cases}, \quad (1.23)$$

where $\overline{a^{\#1}a^{\#2}} := \nu^{(cen)}(a^{\#1}a^{\#2})$. (1.23) is referred to as Wick's Theorem.

Definition 1.2 (Number conserving state). *A state ν is called number conserving iff $\nu([A, \mathcal{N}_b]) = 0$ for every observable $A \in \mathfrak{A}$.*

We assume that the initial state ν_0 is number conserving, quasifree, and translation-invariant. In particular, we assume that the translation-invariant generator \mathcal{K} is given by

$$\mathcal{K} := \int_{\Lambda^*} dp K(p) a_p^+ a_p, \quad (1.24)$$

where $K(p) \geq \kappa_0$ for some $\kappa_0 > 0$. Observe that, by being number conserving, ν_0 is already centered.

The state describing the two-phase Bose gas is then given by

$$\rho_0(A) := \frac{1}{\text{Tr}(e^{-\mathcal{K}})} \text{Tr} \left(\mathcal{W}[\sqrt{N|\Lambda}]\phi_0 e^{-\mathcal{K}} \mathcal{W}^*[\sqrt{N|\Lambda}]\phi_0 A \right) \quad (1.25)$$

for all $A \in \mathfrak{A}$. The initial value problem (IVP) associated with the Hamiltonian \mathcal{H}_N and the initial state ρ_0 is then given by the von Neumann equation

$$i\partial_t \rho_t(A) = \rho_t([A, \mathcal{H}_N]) \quad (1.26)$$

for all observables $A \in \mathfrak{A}$. Below, we impose assumptions on v ensuring that \mathcal{H}_N is self-adjoint and that it induces a unitary evolution $e^{-it\mathcal{H}_N}$. Then, the solution of (1.26) is given by

$$\rho_t(A) = \rho_0(e^{it\mathcal{H}_N} A e^{-it\mathcal{H}_N}). \quad (1.27)$$

By making specific choices for A , we will study effective equations of key correlation functions characterized by the Bose gas.

1.4. Leading order condensate dynamics: Hartree equation. We expect the leading order dynamics of (1.26) to be described by the leading order condensate dynamics, as the BEC describes the bulk component of the Bose gas. Indeed, for instance, [166, 167, 168, 169, 269], based on Hepp's method and using coherent states, have shown, that, in a precise sense, $e^{-it\mathcal{H}_N} W[\sqrt{N|\Lambda}]\phi_0 \Omega_0$ is well approximated by $W[\sqrt{N|\Lambda}]\phi_t \Omega_0$ for $N \gg 1$, with approximation errors $o_N(1)$, where ϕ_t satisfies the Hartree equation

$$i\partial_t \phi_t = -\frac{1}{2} \Delta \phi_t + \lambda |\Lambda| (v * |\phi_t|^2) \phi_t. \quad (1.28)$$

The volume factor in the nonlinear interaction term accounts for our assumption that the L^2 -mass of the condensate is

$$\|\sqrt{N|\Lambda}]\phi_t\|_{L^2(\Lambda)}^2 = N|\Lambda|. \quad (1.29)$$

In the case $v = \delta$, (1.28) yields the nonlinear Schrödinger equation (NLS). Analogous statements have been proved for different choices of v with alternative approaches involving the corresponding BBGKY hierarchy, see, e.g., [2, 85, 89, 93, 95, 96, 97, 98, 99, 100, 130, 131, 132, 133, 135, 162, 177, 182, 194, 197, 214, 286], and other approaches, see [9, 15, 16, 17, 22, 23, 39, 65, 69, 120, 121, 122, 145, 146, 147, 157, 174, 183, 186, 192, 198, 203, 204, 207, 208, 245, 215, 216, 217, 218, 233, 236, 237, 255, 256, 271, 276, 281, 282]. For more background on the derivation of Hartree theory, we refer to [50, 159, 171, 206, 216, 249, 273].

Stationary, translation-invariant condensate. For simplicity, we choose to consider a stationary and translation-invariant solution of (1.28). Due to the normalization constraint $\|\phi_0\|_2 = 1$, we have

$$\phi_t = \phi_0 = |\Lambda|^{-\frac{1}{2}} \in \mathbb{R}_+. \quad (1.30)$$

Substituting this into (1.28) yields

$$0 = \lambda |\Lambda| |\phi_0|^2 \phi_0 \int_{\Lambda} dx v(x) = \frac{\lambda}{\sqrt{|\Lambda|}} \int_{\Lambda} dx v(x). \quad (1.31)$$

In particular, we assume

$$\int_{\Lambda} dx v(x) = 0, \quad (1.32)$$

with additional regularity properties introduced below. Henceforth, we assume that ϕ is stationary, translation-invariant and satisfies (1.30).

1.5. Leading order fluctuation dynamics: HFB equations. We next turn to the leading order corrections of the full dynamics past the leading order BEC Hartree dynamics. For this purpose, we consider the fluctuation dynamics described by $W^*[\sqrt{N|\Lambda}|\phi_0]e^{-it\mathcal{H}_N}W[\sqrt{N|\Lambda}|\phi_0]$. We show in Lemma B.1

$$W^*[\sqrt{N|\Lambda}|\phi_0]\mathcal{H}_N W[\sqrt{N|\Lambda}|\phi_0] = \mathcal{H}_{HFB} + \mathcal{H}_{cub} + \mathcal{H}_{quart}, \quad (1.33)$$

where

$$\begin{aligned} \mathcal{H}_{HFB} := & \int_{\Lambda^2} dx dy \left(a_x^+ \left(-\frac{1}{2} \delta(x-y) \Delta_x + \lambda v(x-y) \right) a_y \right. \\ & \left. + \frac{\lambda v(x-y)}{2} (a_x^+ a_y^+ + a_x a_y) \right), \end{aligned} \quad (1.34)$$

$$\mathcal{H}_{cub} := \frac{\lambda}{\sqrt{N}} \int_{\Lambda^2} dx dy v(x-y) a_x^+ (a_y + a_y^+) a_y, \quad (1.35)$$

$$\mathcal{H}_{quart} := \frac{\lambda}{2N} \int_{\Lambda^2} dx dy v(x-y) a_x^+ a_y^+ a_y a_x, \quad (1.36)$$

noting that (1.30) has been used to obtain these expressions. In particular, (1.33) implies that the fluctuation dynamics is determined by the unitary operator

$$\tilde{\mathcal{U}}_N(t) := W^*[\sqrt{N|\Lambda}|\phi_0]e^{-it\mathcal{H}_N}W[\sqrt{N|\Lambda}|\phi_0], \quad (1.37)$$

where

$$i\partial_t \tilde{\mathcal{U}}_N(t) = (\mathcal{H}_{HFB} + \mathcal{H}_{cub} + \mathcal{H}_{quart}) \tilde{\mathcal{U}}_N(t). \quad (1.38)$$

In the unitary evolution relative to the Hartree dynamics, the dynamics of thermal bosons is determined by two types of processes:

- (1) Emission and absorption of thermal bosons from and into the BEC, respectively.
- (2) Collisions between thermal bosons.

In particular, the Hamiltonian $\mathcal{H}_{HFB} + \mathcal{H}_{cub} + \mathcal{H}_{quart}$ describing this relative dynamics is not number conserving, as opposed to the original Hamiltonian \mathcal{H}_N .

Observe that conjugation by the Weyl operator $\mathcal{W}[\sqrt{N|\Lambda}|\phi_0]$ 'subtracts' the condensate dynamics, thereby revealing the relative dynamics. As a consequence, our focus will be on the IVP

$$\begin{cases} i\partial_t \rho_t^{(rel.BEC)}(A) &= \rho_t^{(rel.BEC)}([A, \mathcal{H}_{HFB} + \mathcal{H}_{cub} + \mathcal{H}_{quart}]), \\ \rho_0^{(rel.BEC)}(A) &= \frac{1}{\text{Tr}(e^{-\mathcal{K}})} \text{Tr}(e^{-\mathcal{K}} A). \end{cases} \quad (1.39)$$

Notice that the initial state $\rho_0^{(rel.BEC)}$ is chosen to be particle number conserving, in contrast to ρ_0 .

Observe that $\mathcal{H}_{HFB} = O(1)$, while $\mathcal{H}_{cub} = O(\frac{\lambda}{\sqrt{N}})$ and $\mathcal{H}_{quart} = O(\frac{\lambda}{N})$ are of lower order as $N \gg 1$. We thus expect the leading order fluctuation dynamics to be determined by the Hartree-Fock-Bogoliubov (HFB) dynamics described by the Hamiltonian \mathcal{H}_{HFB} . Bogoliubov [62] observed that a Hamiltonian of the form of \mathcal{H}_{HFB} can be diagonalized in terms of rotated creation and annihilation operators $b_x = a(u_x) + a^+(v_x)$. Based on this idea, and considering the more general case for a non-stationary, non-translation invariant condensate wave function ϕ_t , there are many works analyzing the emergence of the HFB dynamics, including [3, 31, 32, 55, 56, 57, 66, 67, 68, 71, 72, 78, 101, 102, 103, 105, 112, 166, 167, 168, 199, 200, 209, 239, 240, 241, 242, 243, 244, 246, 247, 254] and references therein. See also [206, 249] for more details.

Let $(\mathcal{V}_{HFB}(t))_{t \in \mathbb{R}}$ denote the unitary group associated with the generator \mathcal{H}_{HFB} , see (3.7) below. Then the density of HFB fluctuation particles, we have that

$$\mathcal{V}_{HFB}^*(t) \mathcal{N}_b \mathcal{V}_{HFB}(t) \lesssim \mathcal{N}_b + |\Lambda|, \quad (1.40)$$

see Remark 4.9 for details. In particular, the HFB fluctuation density has the order of magnitude

$$\frac{1}{|\Lambda|} \rho_0^{(rel.BEC)}(\mathcal{V}_{HFB}^*(t) \mathcal{N}_b \mathcal{V}_{HFB}(t)) \lesssim 1, \quad (1.41)$$

compared to the BEC density N , see also Lemma 4.1.

As we verify in Lemma 3.3, the HFB evolution captures oscillations between absorption into and emission from the BEC with frequency

$$\Omega = \sqrt{E(E + 2\lambda\hat{v})}, \quad (1.42)$$

which is the Bogoliubov dispersion relation. Here \hat{v} is the Fourier transform of v , see definition (1.43), and $E(p) = |p|^2/2$ denotes the free kinetic energy. We will assume $\hat{v} \geq 0$ to be non-negative. The Bogoliubov dispersion corresponds to the propagation of acoustic excitations, see, e.g., [112, 274] for more details.

1.6. Fourier transform. Before moving on to the next order corrections, we fix our conventions for the Fourier transform. Let the Fourier transform be given by

$$\hat{f}(p) := \int_{\Lambda} dx e^{ip \cdot x} f(x), \quad (1.43)$$

for all $p \in \Lambda^* = (\frac{2\pi}{L}\mathbb{Z})^3$, where $\Lambda = (L\mathbb{T})^3$. We denote by

$$a_p := a(e^{-ip \cdot (\cdot)}) = \int_{\Lambda} dx e^{ip \cdot x} a_x, \quad (1.44)$$

$$a_p^+ := a^+(e^{-ip \cdot (\cdot)}) = \int_{\Lambda} dx e^{-ip \cdot x} a_x^+ \quad (1.45)$$

the Fourier transforms of the operator-valued distributions a_x, a_y^+ . These satisfy the discrete CCR

$$[a_p, a_q^+] = |\Lambda| \delta_{p,q} =: \delta_{\Lambda^*}(p - q), \quad (1.46)$$

$$[a_p, a_q] = [a_p^+, a_q^+] = 0. \quad (1.47)$$

When the context is clear, we will omit the subscript ' Λ^* ' in δ_{Λ^*} .

Recalling (1.15), the number operator is given by

$$\mathcal{N}_b = \int_{\Lambda^*} dp a_p^+ a_p. \quad (1.48)$$

Then a_p, a_q^+ satisfy the bounds

$$\|a_p \Psi\|_{\mathcal{F}} \leq \|e^{-ip \cdot (\cdot)}\|_{L^2(\Lambda)} \|\mathcal{N}_b^{\frac{1}{2}} \Psi\|_{\mathcal{F}} = |\Lambda|^{\frac{1}{2}} \|\mathcal{N}_b^{\frac{1}{2}} \Psi\|_{\mathcal{F}}, \quad (1.49)$$

$$\|a_p^+ \Psi\|_{\mathcal{F}} = \sqrt{\|a_p \Psi\|_{\mathcal{F}}^2 + |\Lambda|} \leq |\Lambda|^{\frac{1}{2}} \|(\mathcal{N}_b + 1)^{\frac{1}{2}} \Psi\|_{\mathcal{F}}, \quad (1.50)$$

see, e.g., [50].

For convenience, we will use the notation

$$\int_{\Lambda^*} dp f(p) \equiv \frac{1}{|\Lambda|} \sum_{p \in \Lambda^*} f(p). \quad (1.51)$$

In contrast, we will denote the Lebesgue integral by $\int_{\mathbb{R}^3} dq f(q)$. Moreover, we will denote an n -tuple (q_1, q_2, \dots, q_n) , $n \in \mathbb{N}$, of vectors $q_j \in \mathbb{R}^d$ for some $d \in \mathbb{N}$, as

$$\mathbf{q}_n := (q_1, q_2, \dots, q_n). \quad (1.52)$$

1.7. Lower order fluctuations: Emergence of Boltzmann dynamics. In order to study corrections to the HFB dynamics, we subtract it from the dynamics (1.39) relative to the BEC Hartree dynamics by conjugation with $\mathcal{V}_{HFB}(t)$, the unitary group induced by \mathcal{H}_{HFB} . The resulting relative dynamics is determined by

$$\begin{cases} i\partial_t \nu_t(A) &= \nu_t([A, \mathcal{H}_{cub}(t) + \mathcal{H}_{quart}(t)]), \\ \nu_0(A) &= \frac{1}{\text{Tr}(e^{-\mathcal{K}})} \text{Tr}(e^{-\mathcal{K}} A), \end{cases} \quad (1.53)$$

where

$$\begin{aligned} \mathcal{H}_{cub}(t) &:= \mathcal{V}_{HFB}^*(t) \mathcal{H}_{cub} \mathcal{V}_{HFB}(t) \\ &= \frac{\lambda}{\sqrt{N}} \int_{(\Lambda^*)^3} d\mathbf{p}_3 \delta(p_1 + p_2 - p_3) \hat{v}(p_2) \\ &\quad \left(e^{i(\Omega(p_1) + \Omega(p_2) - \Omega(p_3))t} a_{p_1}^+ a_{p_2}^+ a_{p_3} + h.c. \right) + O\left(\frac{\lambda^2}{\sqrt{N}}\right) \end{aligned} \quad (1.54)$$

$$\begin{aligned} \mathcal{H}_{quart}(t) &:= \mathcal{V}_{HFB}^*(t) \mathcal{H}_{quart} \mathcal{V}_{HFB}(t) \\ &= \frac{\lambda}{2N} \int_{(\Lambda^*)^4} d\mathbf{p}_4 \delta(p_1 + p_2 - p_3 - p_4) \hat{v}(p_1 - p_3) \\ &\quad e^{i(\Omega(p_1) + \Omega(p_2) - \Omega(p_3) - \Omega(p_4))t} a_{p_1}^+ a_{p_2}^+ a_{p_3} a_{p_4} + O\left(\frac{\lambda^2}{N}\right), \end{aligned} \quad (1.55)$$

see Corollary 3.3. For a derivation of (1.53), we refer to Lemma B.1.

We are interested in the evolution of the density of fluctuation particles

$$f_t(p) := \frac{\nu_t(a_p^+ a_p)}{|\Lambda|}. \quad (1.56)$$

We have that

$$f_0(p) = \frac{1}{e^{K(p)} - 1}, \quad (1.57)$$

see Remark 4.2. The case $K(p) = \beta(E(p) - \mu)$, with inverse temperature $\beta > 0$, and chemical potential $\mu < 0$, corresponds to the Bose-Einstein distribution of the ideal Bose gas.

In order to study f_t , we need to extend (1.53) to hold for a more general class of operators. We prove in Lemma 4.1 a quantitative version of

$$\nu_0(\mathcal{N}_b^k) < \infty, \quad (1.58)$$

and in Corollary 4.10 a quantitative version of

$$\nu_t(\mathcal{N}_b^k) < \infty \quad (1.59)$$

for all $k \in \mathbb{N}$. Thus, we may extend the IVP (1.53) to observables

$$A = a_{p_1}^{\sharp_1} a_{p_2}^{\sharp_2} \dots a_{p_k}^{\sharp_k}. \quad (1.60)$$

In order to expand f_t , we follow [184, 179, 129] and apply Duhamel's formula three times

$$\begin{aligned} f_t(p) - f_0(p) = & \\ & - \frac{i}{|\Lambda|} \int_0^t ds \nu_0([a_p^+ a_p, \mathcal{H}_I(s)]) \end{aligned} \quad (1.61)$$

$$- \frac{1}{|\Lambda|} \int_{[0,t]^2} ds_2 \mathbf{1}_{s_1 \geq s_2} \nu_0([a_p^+ a_p, \mathcal{H}_I(s_1)], \mathcal{H}_I(s_2)) \quad (1.62)$$

$$+ \text{Rem}_t(p), \quad (1.63)$$

where we abbreviated

$$\mathcal{H}_I(t) := \mathcal{H}_{cub}(t) + \mathcal{H}_{quart}(t), \quad (1.64)$$

$$\text{Rem}_t(p) := \frac{i}{|\Lambda|} \int_{[0,t]^3} ds_3 \mathbf{1}_{s_1 \geq s_2 \geq s_3} \nu_{s_3}([a_p^+ a_p, \mathcal{H}_I(s_1)], \mathcal{H}_I(s_2), \mathcal{H}_I(s_3)) \quad (1.65)$$

and $\mathbf{s}_j = (s_1, \dots, s_j)$, $j \geq 2$.

Because $a_p^+ a_p$ commutes with $e^{-\mathcal{K}}$, the transport term (1.61) vanishes. Due to translation-invariance and ν_0 being number conserving, the transport term (1.61) vanishes.

For (1.62), observe that $\mathcal{H}_{cub}(t) \sim \frac{\lambda}{\sqrt{N}}$ is much larger than $\mathcal{H}_{quart}(t) \sim \frac{\lambda}{N}$. We thus expect the main contribution in (1.62) to stem from the terms involving \mathcal{H}_{cub} .

It is key to our analysis that we exploit the fact that the HFB dynamics happens on a much *shorter* time scale than the corrections coming from \mathcal{H}_{cub} and \mathcal{H}_{quart} . Observing that λ defines the coupling strength at the level of the HFB evolution, we consider the kinetic time scale defined by $t \sim \lambda^{-2} \gg 1$. In order to separate the corrections in (1.54) and (1.55) from the HFB oscillations, we choose $0 < \lambda \ll 1$.

Using quasifreeness of ν_0 , we thus expect the main contributions in (1.62) to be given by

$$\frac{1}{N} \int_0^T dS Q_S^{(mol)}(f_0)(p) \quad (1.66)$$

$$+ \left| - \frac{i}{|\Lambda|} \int_0^{T\lambda^{-2}} ds \nu_0([a_0, \mathcal{H}_{cub}(s)]) \right|^2 \delta(p), \quad (1.67)$$

where

$$\begin{aligned}
 & Q_S^{(mol)}(h)(p) \\
 &= \int_{(\Lambda^*)^3} d\mathbf{p}_3 \frac{\sin\left(\frac{T-S}{\lambda^2}(\Omega(p_1) + \Omega(p_2) - \Omega(p_3))\right)}{\Omega(p_1) + \Omega(p_2) - \Omega(p_3)} \delta(p_1 + p_2 - p_3) \\
 &\quad (\hat{v}(p_1) + \hat{v}(p_2))^2 (\delta(p - p_1) + \delta(p - p_2) - \delta(p - p_3)) \\
 &\quad ((1 + h(p_1))(1 + h(p_2))h(p_3) - h(p_1)h(p_2)(1 + h(p_3))) \tag{1.68}
 \end{aligned}$$

is a mollification of the cubic Boltzmann operator given in (1.4). The fact that energy conservation only holds approximately up to an error of order $\lambda^2 = O(t^{-1})$ is consistent with the time-energy Heisenberg uncertainty principle, see Remark 2.5 for more details. $Q_S^{(mol)}$ describes collisions between fluctuation particles, one of which is being absorbed into or emitted from the BEC. We expect that (1.67) is of size $\frac{T^2}{N\lambda^2}$ and that it dominates the Boltzmann collision term $\frac{1}{N} \int_0^T dS Q_S^{(mol)}$. It turns out that the presence of (1.67) owes to the fact that, above, we only subtracted the leading-order condensate dynamics. Thus, in order to resolve Boltzmann dynamics collisions, we need to pass to centered moments according to $a_p \rightarrow a_p - \nu_t(a_p)$. Denoting

$$\begin{aligned}
 F_T(p) &:= \frac{\nu_{T\lambda^{-2}}((a_p^+ - \nu_{T\lambda^{-2}}(a_p^+))(a_p - \nu_{T\lambda^{-2}}(a_p)))}{|\Lambda|} \\
 &= \frac{\nu_{T\lambda^{-2}}(a_p^+ a_p) - |\nu_{T\lambda^{-2}}(a_p)|^2}{|\Lambda|}, \tag{1.69}
 \end{aligned}$$

and using that ν_0 is number conserving, we expect – and will indeed prove – to have

$$F_T(p) - F_0(p) = \frac{1}{N} \int_0^T dS Q_S^{(mol)}(f_0)(p) + \text{Rem}_{T\lambda^{-2}}(p) + l.o.t., \tag{1.70}$$

where "l.o.t." abbreviates "lower order terms".

Before moving on, we would like to reflect on the validity of this identity.

1.7.1. *Fixed, N -independent lattice $\Lambda^* \cong \mathbb{Z}^3$.* Recall that the fluctuation particles, at leading order, propagate with the Bogoliubov dispersion Ω . These acoustic waves have the phase velocity

$$v_P(p) := \frac{\Omega(p)}{|p|} = \sqrt{\frac{E(p)}{2}} + \lambda \hat{v}(p). \tag{1.71}$$

Averaging this over all particles yields

$$\langle v_P \rangle_0 := \int_{\Lambda^*} dp f_0(p) v_P(p) \sim 1, \tag{1.72}$$

where we assume that f_0 is sufficiently regular for this argument. During the time $t \sim \lambda^{-2}$, the corresponding acoustic waves propagate a distance $\sim \lambda^{-2}$. In particular, we have

$$\lambda^{-2} \gtrsim L \Leftrightarrow \text{acoustic waves interfere with themselves.}$$

Thus, when λ is small enough, lower-order terms in (1.54) and (1.55) can constructively interfere to an extent as to contribute to leading order terms of F in (1.70). As we will see below, the effect of these contributions is large, depending on whether certain time-dependent expressions, coming from HFB oscillations, have particular

values in \mathbb{Q} . This phenomenon is slightly reminiscent of the *Talbot effect* [128, 190]. The absence of this effect has been discussed in the context of the kinetic wave equation, see, e.g., [75, 107, 108, 113, 114]. To the best of our knowledge, this phenomenon has not previously been discussed in the literature in the context of the quantum field theoretic emergence of Boltzmann equations.

1.7.2. *Continuum approximation* $\Lambda^* \rightarrow \mathbb{R}^3$. In order to elucidate the link with the expression for the BUU collision operator that is widely discussed in the literature, we present a continuum limit, that is derived here. However, we emphasize that this limit applies to the kernel itself, but not to the dynamics because sharp energy conservation cannot hold for finite times, due to the Heisenberg uncertainty.

As observed above, we expect that, for $L \gg \lambda^{-2}$, the self-interactions of the HFB waves to be negligible. In fact, we show that in this limit, we can approximate sums $\int_{\Lambda^*} dp \equiv \frac{1}{|\Lambda|} \sum_p$ by Lebesgue-integrals $\frac{1}{(2\pi)^3} \int_{\mathbb{R}^3}$, with control of errors that include oscillatory contributions, see Lemma 5.5. In this case, we establish that

$$F_T(p) - F_0(p) = \frac{T}{N} Q(f_0)(p) + \text{Rem}_{T\lambda^{-2}}(p) + O\left(\frac{1}{N^{1+\delta_N}}\right), \quad (1.73)$$

where

$$\begin{aligned} & Q(f_0)[J] \\ & := \frac{\pi}{(2\pi)^6} \int_{\mathbb{R}^9} dp_1 dp_2 dp_3 \delta(E(p_1) + E(p_2) - E(p_3)) \delta(p_1 + p_2 - p_3) \\ & \quad (\hat{v}(p_1) + \hat{v}(p_2))^2 (J(p_1) + J(p_2) - J(p_3)) \\ & \quad ((1 + f_0(p_1))(1 + f_0(p_2))f_0(p_3) - f_0(p_1)f_0(p_2)(1 + f_0(p_3))) \end{aligned} \quad (1.74)$$

is the energy-conserving cubic Boltzmann operator found in the above mentioned literature.

1.7.3. *Propagation of quasifreeness*. If we can show that $\text{Rem}_{T\lambda^{-2}}(p)$ is, in fact, of lower order compared to $\frac{1}{N} \int_0^T dS Q_S^{(mol)}$, then, by rearranging (1.70), we find that

$$F_T(p) - F_0(p) = \frac{1}{N} \int_0^T dS Q_S^{(mol)}(F)(p) + l.o.t., \quad (1.75)$$

which would prove that the next-to-leading order correction to the HFB dynamics of the particle density is described by a cubic Boltzmann equation. Comparing with the analysis of collisions for classical systems, bounding $\text{Rem}_t(p)$ in the present context includes controlling recollisions, as is necessary in the context of classical systems. For more details on the role of recollisions in the classical case, we refer, e.g., to [81].

While drawing comparisons to the classical case, we address the role of quasifreeness. With the given choice of a number conserving and translation-invariant initial state ν_0 , we have that the joint distribution function of n particles satisfies

$$f_{joint}(\mathbf{p}_n) := \frac{\nu_0(a_{p_1}^+ a_{p_2}^+ \dots a_{p_n}^+ a_{p_n} \dots a_{p_2} a_{p_1})}{|\Lambda|^n} \quad (1.76)$$

$$= f_0^{\otimes n}(\mathbf{p}_n) + O(|\Lambda|^{-1}), \quad (1.77)$$

where $\mathbf{p}_n = (p_1, \dots, p_n)$. In particular, we have that asymptotically, for large $L \gg 1$, quasifreeness implies *molecular chaos* in the classical sense, which refers to

factorization of the joint distribution. Propagation of the factorized form is called *propagation of chaos* in the classical context.

In the present quantum field theoretic context, it is an important task to understand in what sense and in which scaling regime propagation of quasifreeness can be observed. An important property of the HFB evolution is that it *preserves* quasifreeness. In particular, if a state $\langle \cdot \rangle_0$ is quasifree, then so is $\langle \mathcal{V}_{HFB}^*(t)(\cdot)\mathcal{V}_{HFB}(t) \rangle_0$. This is a natural consequence of the fact that the HFB dynamics arises as a quasifree reduction of the full evolution. In the recent literature, this is explained on the basis of the Dirac-Frenkel principle, see, for instance, [34] and [51] for more details.

Notice that the full evolution clearly does not preserve quasifreeness, as is expected for an interacting gas. In order to study propagation of quasifreeness, it is crucial to control the Duhamel term $\text{Rem}_t(p)$ accounting for all quantum 'recollisions'. Indeed, if we expand the evolution of arbitrary expectations

$$\nu_{T\lambda^{-2}}(a_{p_1}^{\#1} a_{p_2}^{\#2} \dots a_{p_k}^{\#k}) = \nu_0(a_{p_1}^{\#1} a_{p_2}^{\#2} \dots a_{p_k}^{\#k}) + \text{Rem}_{T\lambda^{-2}}(\mathbf{p}_k), \quad (1.78)$$

controlling $\text{Rem}_{T\lambda^{-2}}(\mathbf{p}_k)$ implies that $\nu_{T\lambda^{-2}}$ is quasifree to k^{th} order. Similar to [129], we do not need to propagate quasifreeness to arbitrary orders to derive a Boltzmann equation; instead, adopting their notion of *restricted quasifreeness*, it is sufficient to show that $\nu_{T\lambda^{-2}}$ is approximately restricted quasifree up to eight-point correlation functions, see (1.62). However, we choose not to explicitly prove such a statement, and, instead, calculate the evolution of f explicitly. We leave the proof of a more general result of the form (1.78) to the interested reader, which will be straightforward using the tools developed in this work.

It remains to understand how we can control $\text{Rem}_t(p)$. For that, we use the fact that \hat{v} is bounded to show that

$$\mathcal{H}_{cub}(t) \lesssim \frac{\lambda}{\sqrt{N}} (\mathcal{N}_b + |\Lambda|)^{\frac{3}{2}}, \quad (1.79)$$

$$\mathcal{H}_{quart}(t) \lesssim \frac{\lambda}{N} (\mathcal{N}_b + |\Lambda|)^2, \quad (1.80)$$

see Lemmata 4.4 and 4.5. When bounding products of non number conserving operators, one needs to take into account the growth of the particle number, see Lemma 4.3. This is consistent with the fact that fluctuation particles are being absorbed into and emitted from the BEC. It turns out that in order to control $\text{Rem}_t(p)$, it suffices to only consider three, and thus a fixed number of Duhamel iterations. Hence, we apply the bounds (1.49), (1.50) on $a_p^+ a_p$ with the bound (1.79) and (1.80) on $\mathcal{H}_I(t)$ to obtain

$$|\text{Rem}_{T\lambda^{-2}}(p)| \lesssim \frac{T^3}{N^{\frac{3}{2}} \lambda^3} \sup_{t \in [0, T\lambda^{-2}]} \nu_t \left((\mathcal{N}_b + |\Lambda|)^{\frac{11}{2}} \left(1 + \frac{\sqrt{\mathcal{N}_b + |\Lambda|}}{\sqrt{N}} \right)^3 \right). \quad (1.81)$$

In order to bound $\nu_t(\mathcal{N}_b^k)$, we employ a result by Rodnianski and Schlein [269], see also [74], to obtain that

$$\nu_t((\mathcal{N}_b + |\Lambda|)^{\frac{k}{2}}) \lesssim_{|\Lambda|, \ell} e^{K_\ell \lambda |\Lambda| t}. \quad (1.82)$$

In particular, (1.81) then yields the constraint

$$\frac{|\Lambda|}{\lambda} \lesssim \frac{\log N}{\log \log N}, \quad (1.83)$$

in order to suppress $\text{Rem}_{T\lambda^{-2}}(p)$ compared to the leading order Boltzmann term in the evolution of F , see (1.70). In particular, this implies the scaling

- (1) $L \sim 1$ fixed: $\lambda \sim \frac{\log \log N}{\log N}$,
- (2) $L \sim \lambda^{-2-}$: $\lambda \sim \left(\frac{\log \log N}{\log N}\right)^{\frac{1}{7}-}$.

1.8. Fluctuations beyond HFB. Now that we understand that approximate restricted quasifreeness can be propagated, we would like to comment on the fact that the fluctuation dynamics does not preserve number conservation. We have that, for a non number conserving quasifree state μ , a more general version of the Wick-Theorem holds: Let $b^\# := a^\# - \mu(a^\#)$. Then μ is quasifree if and only if it satisfies (1.23) with the operators $a^\#$ replaced by their centered counterparts $b^\#$.

Since quasifreeness implies that n -point correlation functions $\langle a^{\#1} \dots a^{\#n} \rangle$ are determined by the one- and two-point correlation functions, and since by (1.78), $\nu_{T\lambda^{-2}}$ is approximately restricted quasifree, we need to include the dynamics of

$$\Phi_t := \frac{\nu_t(a_0)}{|\Lambda|}, \quad (1.84)$$

$$g_t(p) := \frac{\nu_t(a_p a_{-p})}{|\Lambda|} \quad (1.85)$$

in our analysis. We have that Φ_t captures the corrections to the condensate dynamics. g_t is the rate of absorption into or emission from the BEC of pairs of thermal bosons.

Recall that, due to translation-invariance of the condensate, ν_t is also translation-invariant, which is why we have that $\nu_t(a_p) = \Phi_t \delta(p)$, see Lemma A.1. Due to ν_0 being number conserving, we have that $\Phi_0 = g_0 = 0$. Arguing as in the case of f above, we are also interested in the dynamics of the centered, mesoscopic counterparts

$$\Psi_T := \frac{\nu_{T\lambda^{-2}}(a_0)}{|\Lambda|}, \quad (1.86)$$

$$G_T(p) := \frac{\nu_{T\lambda^{-2}}(a_p a_{-p}) - \nu_{T\lambda^{-2}}(a_p) \nu_{T\lambda^{-2}}(a_{-p})}{|\Lambda|}. \quad (1.87)$$

We show that

$$\Psi_T = \frac{-ic_1(f_0)T}{N^{\frac{1}{2}}\lambda} + O\left(\frac{1}{N^{\frac{1}{2}+\delta_N}\lambda}\right), \quad (1.88)$$

$$G_T = (T + T^2)O\left(\frac{1}{N\lambda^2}\right), \quad (1.89)$$

where, assuming that \hat{v} is real-valued, $c_1(f_0)$ is real-valued. At leading order, we show that the dynamics of G is completely determined by F . As a consequence of (1.89), and recalling (1.61) and (1.62), we have that G merely contributes lower-order corrections to the Boltzmann dynamics for F .

1.9. Scaling of λ and N . We emphasize that the parameter $N \gg 1$ accounts for the L^2 mass of the BEC per unit volume, and is unrelated to the ~ 1 density of fluctuation particles around the BEC. Hence $1/N$ yields a small perturbation parameter in the expansion of the full dynamics, in addition to the coupling constant $0 < \lambda \ll 1$ characterizing the HFB dynamics. Boßmann et al. [67] give an entire expansion of the fluctuation dynamics in powers of λ/N , in terms of effective

Hamiltonians for a given order of precision. Our analysis differs in that we choose $\lambda \ll 1$ and $N \gg 1$ suitably in order to be able to extract effective equations for the moments F, G, Ψ , while keeping the error sufficiently small. In particular, our time scale is $O((\log N / \log \log N)^\alpha)$, $\alpha > 0$, instead of $O(1)$. In the latter case, the Boltzmann dynamics cannot be observed.

In the derivation of Boltzmann equations in classical collisional systems along the lines of Lanford's approach, $O(1)$ many collisions take place during the relevant time scale (which is inversely proportional to the mean free path). In our result, we encounter a similar situation; the error of order $O(N^{-1-})$, in (1.75) dominates after $O(1)$ collisions have taken place; this can be easily seen by iterating the Duhamel formula (1.75) twice (as the k -th order terms in the Duhamel expansion, of size $O(N^{-k})$, account for k collisions). We also note that, compliant with a kinetic scaling regime, particle velocities do not scale in our problem.

Our results are limited to the parameter regime $\lambda \sim (\log \log N / \log N)^{1/7-}$ when $L \sim \lambda^{-2-}$, similar for $\lambda \sim \log \log N / \log N$ when $L \sim 1$, and $N \gg 1$. This is, in part, due to technical reasons, but we do not expect the fluctuation dynamics to remain of the form (1.75), (1.88), (1.89) for longer time scales, even if our approach is extended to the next order of magnitude. We expect the analogous to hold for the parameter regime $\lambda \sim \frac{\log \log N}{\log N}$ when $L \sim 1$.

Remark 1.3. *For works studying the perturbation expansion for a fermionic gas, we refer to [33, 46, 47, 48, 49, 51, 86, 87, 90, 91, 104, 106, 210, 211, 248, 261] and references therein.*

2. MAIN RESULTS

2.1. Notation . We introduce the rescaled $L^a(\Lambda^*)$ norms

$$\|f\|_{L^a(\Lambda^*)} := |\Lambda|^{-\frac{1}{a}} \|f\|_{\ell^a(\Lambda^*)} \quad \text{if } 1 \leq a < \infty, \quad (2.1)$$

and accordingly

$$\|f\|_{a \cap \infty, d} := \|f\|_{L^a(\Lambda^*)} + \|f\|_{\ell^\infty(\Lambda^*)}, \quad (2.2)$$

$$\|f\|_d := \|f\|_{1 \cap \infty, d}, \quad (2.3)$$

$$\|f\|_{m, c} := \sum_{n=0}^m \left\| \langle |\cdot| \rangle^n D^{m-n} f \right\|_{L^1(\mathbb{R}^3)} \quad \text{if } m \in \mathbb{N}. \quad (2.4)$$

Moreover, we introduce the weight

$$w(p) := 1 + \frac{1}{|p|^2}. \quad (2.5)$$

Whenever $wf \in L^\infty(B_R(0) \setminus \{0\})$ for some $R > 0$, we define the weighted norms

$$\|f\|_{w, d} := \|wf\|_d, \quad (2.6)$$

$$\|f\|_{m, w, c} := \|wf\|_{m, c}. \quad (2.7)$$

Recall that

$$\begin{aligned} \mathcal{H}_{cub}(t) &= \frac{\lambda}{\sqrt{N}} \int_{(\Lambda^*)^3} d\mathbf{p}_3 \hat{v}(p_2) \delta(p_1 + p_2 - p_3) \\ &\quad e^{it\mathcal{H}_{HFB}} (a_{p_1}^+ a_{p_2}^+ a_{p_3} + h.c.) e^{-it\mathcal{H}_{HFB}}, \quad (2.8) \\ \mathcal{H}_{quart}(t) &= \frac{\lambda}{2N} \int_{(\Lambda^*)^3} d\mathbf{p}_3 \hat{v}(p_1 - p_3) \delta(p_1 + p_2 - p_3 - p_4) \end{aligned}$$

$$e^{it\mathcal{H}_{HFB}} a_{p_1}^+ a_{p_2}^+ a_{p_3} a_{p_4} e^{-it\mathcal{H}_{HFB}}, \quad (2.9)$$

where

$$\mathcal{H}_{HFB} = \int_{\Lambda^*} dp (E(p) + \lambda \hat{v}(p)) a_p^+ a_p + \frac{\lambda}{2} \int_{\Lambda^*} dp \hat{v}(p) (a_p^+ a_{-p}^+ + a_p a_{-p}), \quad (2.10)$$

$E(p) = \frac{|p|^2}{2}$, and $\int_{\Lambda^*} \equiv \frac{1}{|\Lambda|} \sum_{\Lambda^*}$. In this context, we also recall the Bogoliubov dispersion relation

$$\Omega(p) = \sqrt{E(p)(E(p) + 2\lambda \hat{v}(p))}. \quad (2.11)$$

We are interested in the evolution of the correlation functions

$$\Psi_T = \frac{\nu_{T\lambda^{-2}}(a_0)}{|\Lambda|}, \quad (2.12)$$

$$F_T(p) = \frac{\nu_{T\lambda^{-2}}(a_p^+ a_p) - |\nu_{T\lambda^{-2}}(a_p)|^2}{|\Lambda|}, \quad (2.13)$$

$$G_T(p) = \frac{\nu_{T\lambda^{-2}}(a_p a_{-p}) - \nu_{T\lambda^{-2}}(a_p) \nu_{T\lambda^{-2}}(a_{-p})}{|\Lambda|}. \quad (2.14)$$

As explained in 1.7.1, there are additional dominant terms in the Boltzmann collision terms for F , and also G in the case of $L \sim 1$ fixed. Thus, we introduce the collision operators

$$\mathcal{Q}_{d,G;T,\lambda}(h)[J] := \frac{1}{\lambda^2} \int_{[0,T]^2} \mathbf{1}_{S_1 \geq S_2} d\mathbf{S}_2 \text{col}_d(h, \lambda^2)[J](\mathbf{S}_2/\lambda^2), \quad (2.15)$$

$$q_{d,F;S_2,\lambda}^{(j)}(H_{S_2})[J] := \text{bol}^{(j)}(H, \lambda^2)[J](\mathbf{S}_2/\lambda^2), \quad j \in \{1, 2\}, \quad (2.16)$$

and the pair absorption operator

$$\begin{aligned} \mathcal{A}_{d;T,\lambda}(h)[J] &:= \int_0^T dS \text{abs}_{\text{quart},d}(h, \lambda^2)[J](S/\lambda^2) \\ &+ \frac{1}{\lambda^2} \int_{[0,T]^2} \mathbf{1}_{S_1 \geq S_2} d\mathbf{S}_2 \text{abs}_{\text{cub},d}(h, \lambda^2)[J](\mathbf{S}_2/\lambda^2), \end{aligned} \quad (2.17)$$

for any test function J , where $\mathbf{S}_2 = (S_1, S_2)$. The expressions for $\text{bol}^{(j)}$, $j \in \{1, 2\}$, col_d , $\text{abs}_{\text{quart},d}$, and $\text{abs}_{\text{cub},d}$ are lengthy, and we refer the reader to Section 5.1.1 for their definition. In the case $L \sim \lambda^{-2-}$, we also define the continuous counterparts $\mathcal{Q}_{c,G;T,\lambda}$, $\mathcal{A}_{c;T,\lambda}$ by replacing sums $\int_{\Lambda^*} dp$ over momenta by Lebesgue integrals $\frac{1}{(2\pi)^3} \int_{\mathbb{R}^3}$.

The subscript 'd' in the notation refers to the fact that momentum is summed over the discrete set Λ^* ; 'c' on the other hand refers to the continuum approximation. Henceforth, we refer to the case with $L \sim 1$ fixed as the 'discrete case', and $L \sim \lambda^{-2-}$ as the 'continuum approximation'.

$\mathcal{Q}_{d,G;T,\lambda}(h)$ resp. $\mathcal{Q}_{c,G;T,\lambda}(h)$ is a Boltzmann collision operator for G , and it is, after cancellations, quadratic in h , and $\mathcal{A}_{d;T,\lambda}(h)$ resp. $\mathcal{A}_{c;T,\lambda}(h)$ corresponds to the leading order of the expected rate of absorption of a pair of fluctuation bosons into the BEC, and it is linear in h .

We denote the mollified cubic Boltzmann operator in the evolution of F by

$$Q_{d;T-S,\lambda}(F_S)[J]$$

$$\begin{aligned}
 &:= \int_{(\Lambda^*)^3} d\mathbf{p}_3 \frac{\sin\left(\frac{\Omega(p_1)+\Omega(p_2)-\Omega(p_3)}{\lambda^2}(T-S)\right)}{\Omega(p_1)+\Omega(p_2)-\Omega(p_3)} \delta(p_1+p_2-p_3) \\
 &\quad (\hat{v}(p_1)+\hat{v}(p_2))^2 (J(p_1)+J(p_2)-J(p_3)) \\
 &\quad \left((1+F_S(p_1))(1+F_S(p_2))F_S(p_3) - F_S(p_1)F_S(p_2)(1+F_S(p_3)) \right). \quad (2.18)
 \end{aligned}$$

Again, in the continuum approximation, $Q_{c;T-S,\lambda}$ is defined by replacing the lattice sum \int_{Λ^*} by the Lebesgue integral $\frac{1}{(2\pi)^3} \int_{\mathbb{R}^3}$.

In addition, $q_{d,F;\mathbf{s}_2,\lambda}^{(j)}$ are higher order Boltzmann type collision terms for the equations governing F_T , where j accounts for the order λ^j from which they are derived. As explained above, they are of lower order when $L \sim \lambda^{-2-}$.

2.2. Assumptions. We summarize all the assumptions described in the previous section. We also add the following restrictions required in our results.

(1)

$$\nu_0(A) = \frac{1}{\text{Tr} e^{-\mathcal{K}}} \text{Tr} \left(e^{-\mathcal{K}} A \right) \quad (2.19)$$

for all observables A , is a quasifree, translation-invariant state that is number conserving, with

$$\mathcal{K} = \int_{\Lambda^*} dp K(p) a_p^+ a_p, \quad (2.20)$$

$$f_0(p) = \frac{\nu_0(a_p^+ a_p)}{|\Lambda|} = \frac{1}{e^{K(p)} - 1}, \quad (2.21)$$

and $K(p) \geq \kappa_0 > 0$.

(2) ν_t satisfies

$$\begin{cases}
 i\partial_t \nu_t(A) &= \nu_t([A, \mathcal{H}_{cub}(t) + \mathcal{H}_{quart}(t)]), \\
 \nu_0(A) &= \frac{1}{\text{Tr}(e^{-\mathcal{K}})} \text{Tr} (e^{-\mathcal{K}} A)
 \end{cases} \quad (2.22)$$

for all observables A .

(3) The Fourier transform \hat{v} of v , see Section 1.6, is a non-negative, radial function.

(4) If $L \sim 1$ is fixed, assume $\|\hat{v}\|_{w,d}, \|f_0\|_d < \infty$.

(5) If $L \sim \lambda^{-2-}$, assume $\|\hat{v}\|_{2(\lfloor \frac{\varepsilon}{2} \rfloor + 1), w, c}, \|f_0\|_{2(\lfloor \frac{\varepsilon}{2} \rfloor + 1), c} < \infty$.

In either case, we assume that \hat{v} satisfies

$$\int_{\Lambda} dx v(x) = \hat{v}(0) = 0. \quad (2.23)$$

Moreover, this implies that the leading order condensate wave function $\phi_0 \equiv |\Lambda|^{-1/2}$ can be chosen to be a stationary, translation-invariant solution of the Hartree equation

$$i\partial_t \phi_t = -\Delta \phi_t + \lambda |\Lambda| v * |\phi_t|^2 \phi_t. \quad (2.24)$$

As described above, N denotes the BEC density, and $\lambda > 0$ is an (additional) coupling constant, defining the HFB coupling, see (1.34).

We are now ready to formulate our main results.

2.3. Statement of results.

Theorem 2.1 (Discrete case). *Let $T > 0$, $L \geq 1$, and let $N > 0$ denote the BEC density. Choose*

$$\lambda = \frac{\log \log N}{\log N}. \quad (2.25)$$

Then, under the assumptions stated in Section 2.2 and with the notations in Section 2.1, there exist constants

$$C_0 = C_0(\|\hat{v}\|_{w,d}, \|f_0\|_d, |\Lambda|, T), \quad (2.26)$$

$$N_0 = N_0(\|\hat{v}\|_{w,d}, |\Lambda|, T), \quad (2.27)$$

such that, for all $N \geq N_0$ we have that

$$\delta_1 := \frac{\log \frac{1}{\lambda}}{\log N} > 0, \quad \delta_2 := \frac{1}{2} - \frac{4 \log \frac{1}{\lambda}}{\log N} > 0, \quad (2.28)$$

and that

$$\begin{aligned} & \left| \Psi_T + \frac{i}{N^{\frac{1}{2}\lambda}} \int_0^T dS \int_{\Lambda^*} dp \hat{v}(p) F_S(p) \right| \\ & \leq \frac{C_0}{N^{\frac{1}{2} + \delta_1 \lambda}}, \end{aligned} \quad (2.29)$$

$$\begin{aligned} & \left| \int_{\Lambda^*} dp (F_T(p) - F_0(p)) J(p) - \frac{1}{N} \left(\int_0^T dS Q_{d;T-S,\lambda}(F_S)[J] \right. \right. \\ & \quad \left. \left. + \int_{[0,T]^2} d\mathbf{S}_2 \mathbb{1}_{s_1 \geq s_2} \sum_{j=1}^2 \lambda^j q_{d,F;\mathbf{S}_2,\lambda}^{(j)}(F_{S_2})[J] \right) \right| \\ & \leq \frac{C_0 \|J\|_{\ell^\infty(\Lambda^*)}}{N^{1+\delta_1}}, \end{aligned} \quad (2.30)$$

$$\begin{aligned} & \left| \int_{\Lambda^*} dp G_T(p) J(p) - \frac{1}{N} \left(\mathcal{A}_{d;T,\lambda}(F)[J] + \mathcal{Q}_{d,G;T,\lambda}(F)[J] \right) \right| \\ & \leq \frac{C_0 \|J\|_{2 \cap \infty, d}}{N^{1+\delta_2}} \end{aligned} \quad (2.31)$$

for all test functions J . The error terms on the right-hand sides of (2.29)–(2.31) are subleading in N with respect to the main terms appearing on the respective left-hand sides.

Theorem 2.2 (Continuum approximation). *Let $T > 0$, $r > 6$, $\varepsilon > 0$, and let $N > 0$ denote the BEC density. Fix*

$$\lambda = \left(\frac{\log \log N}{\log N} \right)^{\frac{r}{(7+\varepsilon)r+6}}, \quad (2.32)$$

$$L = \lambda^{-2 - \frac{2}{r} - \varepsilon}. \quad (2.33)$$

Then, under the assumptions stated in Section 2.2 and with the notations in Section 2.1, there exist constants

$$C_0 = C_0(\|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c}, \|f_0\|_{2(\lfloor \frac{r}{2} \rfloor + 1), c}, r, \varepsilon, T), \quad (2.34)$$

$$C_1 = C_1(r, \varepsilon), \quad (2.35)$$

$$N_0 = N_0(\|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c}, r, \varepsilon, T), \quad (2.36)$$

such that, for all $N \geq N_0$, and for

$$\delta := \frac{C_1 \log \frac{1}{\lambda}}{\log N} > 0, \quad (2.37)$$

we have that

$$\begin{aligned} \left| \Psi_T + \frac{i}{(2\pi)^3 N^{\frac{1}{2}} \lambda} \int_0^T dS \int_{\mathbb{R}^3} dp \hat{v}(p) F_S(p) \right| \\ \leq \frac{C_0}{N^{\frac{1}{2} + \delta} \lambda}, \end{aligned} \quad (2.38)$$

$$\begin{aligned} \left| \int_{\Lambda^*} dp (F_T(p) - F_0(p)) J(p) - \frac{1}{N} \int_0^T dS Q_{c;T-S,\lambda}(F_S) \right| \\ \leq \frac{C_0 \|J\|_{W^2[\frac{1}{2}] + 2, \infty}}{N^{1+\delta}}, \end{aligned} \quad (2.39)$$

$$\begin{aligned} \left| \int_{\Lambda^*} dp G_T(p) J(p) - \frac{1}{N} \left(\mathcal{A}_{c;T,\lambda}(F)[J] + \mathcal{Q}_{c,G;T,\lambda}(F)[J] \right) \right| \\ \leq \frac{C_0 \|J\|_{2([\frac{T}{2}] + 1), c}}{N^{1+\delta}} \end{aligned} \quad (2.40)$$

for all test functions J . The error terms on the right-hand sides of (2.38)–(2.39) are subleading in N with respect to the main terms appearing on the respective left-hand sides. The main order term in the evolution of F is given by $\frac{T}{N} Q(f_0)[J]$, where

$$\begin{aligned} Q(f_0)[J] \\ := \frac{\pi}{(2\pi)^6} \int_{\mathbb{R}^9} dp_1 dp_2 dp_3 \delta(E(p_1) + E(p_2) - E(p_3)) \delta(p_1 + p_2 - p_3) \\ (\hat{v}(p_1) + \hat{v}(p_2))^2 (J(p_1) + J(p_2) - J(p_3)) \\ ((1 + f_0(p_1))(1 + f_0(p_2))f_0(p_3) - f_0(p_1)f_0(p_2)(1 + f_0(p_3))) \end{aligned} \quad (2.41)$$

denotes the (energy conserving) quantum Boltzmann collision operator.

Remark 2.3. Theorem 2.2 is not the continuum limit for the dynamics. Instead, it quantifies how the collision operator can be approximated by its continuous counterpart. In particular, energy conservation cannot hold precisely for finite times, in consistence with the Heisenberg uncertainty principle, see Remark 2.5.

Remark 2.4. In this work, we are not attempting to analyze G_T in more detail beyond (2.40). We expect a more detailed analysis to yield

$$\mathcal{A}_{c;T,\lambda}(F)[J] + \mathcal{Q}_{c,G;T,\lambda}(F)[J] \sim T + T^2, \quad (2.42)$$

based on similar arguments as we present to control the Boltzmann collision term for F .

Remark 2.5. We point out that $Q_{d;T-S,\lambda}$ resp. $Q_{c;T-S,\lambda}$ contain the Bogoliubov dispersion Ω for sound waves propagating as fluctuations around the BEC. As stated in Theorem 2.2, the collision operator Q emerges in the limit $\lambda \searrow 0$. In the limit $\lambda \searrow 0$, we have that $\Omega(p) \rightarrow E(p)$, whereby we retrieve the Dirac- δ on the

hypersurface $\{(p_1, p_2) \in \mathbb{R}^6 \mid E(p_1) + E(p_2) = E(p_1 + p_2)\}$ of energy conservation. Let

$$\delta_\varepsilon(x) := \frac{1}{\varepsilon\pi} \frac{\sin^2(x/\varepsilon)}{(x/\varepsilon)^2} = \frac{\varepsilon}{\pi} \frac{\sin^2(x/\varepsilon)}{x^2}, \quad (2.43)$$

which defines a Dirac sequence, $\int_{\mathbb{R}} dx \delta_\varepsilon(x) = 1$. Observing that

$$\frac{\sin\left(\Delta_{\text{cub}}\Omega \frac{T-S}{\lambda^2}\right)}{\Delta_{\text{cub}}\Omega} = -\partial_S \left((T-S) \delta_{\frac{2\lambda^2}{T-S}}(\Delta_{\text{cub}}\Omega) \right), \quad (2.44)$$

where $\Delta_{\text{cub}}\Omega(\mathbf{p}_2) = \Omega(p_1) + \Omega(p_2) - \Omega(p_1 + p_2)$, this implies that the mollified quantum Boltzmann collision terms in (2.30) and (2.39) have the form

$$\begin{aligned} & \int_0^T dS Q_{c/d; T-S, \lambda}(F_S)[J] \\ &= \int_0^T dS \int_{\mathcal{D}} d\mathbf{p}_3 \frac{\sin\left(\Delta_{\text{cub}}\Omega(\mathbf{p}_2) \frac{T-S}{\lambda^2}\right)}{\Delta_{\text{cub}}\Omega(\mathbf{p}_2)} H_S(\mathbf{p}_3) \\ &= \pi T \int_{\mathcal{D}} d\mathbf{p}_3 \delta_{\frac{2\lambda^2}{T}}(\Delta_{\text{cub}}\Omega(\mathbf{p}_2)) H_0(\mathbf{p}_3; J) \\ & \quad + \pi \int_0^T dS \int_{\mathcal{D}} d\mathbf{p}_3 \delta_{\frac{2\lambda^2}{T-S}}(\Delta_{\text{cub}}\Omega(\mathbf{p}_2)) (T-S) \partial_S H_S(\mathbf{p}_3; J), \end{aligned} \quad (2.45)$$

where $\mathbf{p}_3 = (p_1, p_2, p_3)$ and $\mathcal{D} = (\Lambda^*)^3$ in the discrete case, and $\mathcal{D} = \mathbb{R}^9$ for the continuum approximation. Here,

$$\begin{aligned} & H_S(\mathbf{p}_3; J) \\ &:= |\hat{v}(p_1) + \hat{v}(p_2)|^2 (J(p_1) + J(p_2) - J(p_3)) \delta(p_1 + p_2 - p_3) \\ & \quad \left((1 + F_S(p_1))(1 + F_S(p_2))F_S(p_3) - F_S(p_1)F_S(p_2)(1 + F_S(p_3)) \right), \end{aligned} \quad (2.46)$$

where, in the continuum approximation, H contains an additional factor $1/(2\pi)^6$. In particular, this representation makes manifest that the mollification of the quantum Boltzmann collision operator corresponds to approximate energy conservation up to an error of order $O(t^{-1}) = O(\lambda^2/T)$, in compliance with the time-energy Heisenberg uncertainty principle.

Remark 2.6. Q in (2.41) can be evaluated via the Coarea Formula, yielding

$$\begin{aligned} & Q(f_0)[J] \\ &= \frac{\pi}{(2\pi)^6} \int_{E(p_1)+E(p_2)=E(p_1+p_2)} \frac{d\mathcal{H}^5(\mathbf{p}_2)}{|\mathbf{p}_2|} (\hat{v}(p_1) + \hat{v}(p_2))^2 \\ & \quad (J(p_1) + J(p_2) - J(p_1 + p_2)) \\ & \quad \left((1 + f_0(p_1))(1 + f_0(p_2))f_0(p_1 + p_2) - f_0(p_1)f_0(p_2)(1 + f_0(p_1 + p_2)) \right) \end{aligned} \quad (2.47)$$

where $d\mathcal{H}^5$ is the induced Hausdorff measure on the hypersurface $\{(p_1, p_2) \in \mathbb{R}^6 \mid E(p_1) + E(p_2) = E(p_1 + p_2)\}$.

Remark 2.7. Theorem 2.2 implies that, for high regularity $r \gg 6$, for the maximal time scale $t_{\max} \sim \lambda^{-2} \sim (\log N / \log \log N)^{\frac{2}{7}-}$ and length $L \sim \lambda^{-2-} \ll t_{\max}$, we obtain that

$$\Psi_T \sim \frac{T}{N^{\frac{1}{2}-}} + O(N^{-\frac{1}{2}}), \quad (2.48)$$

$$F_T - F_0 \sim \frac{T}{N} + O(N^{-1-}) \quad (2.49)$$

for $N \gg 1$, where $F_0 \sim 1$. We point out that one of the difficulties in extracting a quantum Boltzmann dynamics for F stems from the fact that F is centered with $|\Psi|^2$, which is at least an order $O(N^{0+})$ larger than the Boltzmann collision term for F itself.

There are five characteristic length scales involved in our derivation:

- (1) a BEC with large density N ,
- (2) thermal fluctuations with density ~ 1 ,
- (3) the HFB coupling of size λ ,
- (4) the linear system size L ,
- (5) the time scale t .

A major difficulty that is overcome in this work is to identify a parameter regime which allows the Boltzmann dynamics to dominate over error terms.

Remark 2.8. The collision term for G is a functional only of F , due to our choice of initial data with $G_0 = 0$. Therefore, solving the Boltzmann equation for F , and substituting into

$$\mathcal{I}_G(F)[J] := \frac{1}{N} \mathcal{A}_{c;T,\lambda}(F)[J] + \frac{1}{N\lambda^2} \mathcal{Q}_{c;G;T,\lambda}(F)[J], \quad (2.50)$$

integration time, yields G_T . A key reason for which the extraction of the Boltzmann equation for F is a difficult problem, is the fact that we expect $\mathcal{I}_G(F)[J]$ to be of the same order of magnitude as the collision operator for F ,

$$\mathcal{I}_F(F)[J] := \frac{1}{N} \int_0^T dS \mathcal{Q}_{c;T-S,\lambda}(F_S)[J]. \quad (2.51)$$

In our case, $\mathcal{I}_G(F)[J]$ does not depend on G because the initial state ν_0 is chosen to conserve the particle number \mathcal{N}_b . The explicit expression for $\mathcal{I}_G(F)$ is somewhat lengthy and not sufficiently enlightening to be presented here. We obtain a closed system of equations for (Ψ, F, G) because of the approximate persistence of (restricted) quasifreeness of ν_0 in the scaling regime of this problem.

If we expand the dynamics of (Ψ, F, G) to lower orders, we expect $\mathcal{I}_F(\Psi, F, G)$ and $\mathcal{I}_G(\Psi, F, G)$ to be coupled non-trivially.

Remark 2.9. Our purpose of introducing a condensate of large density N is due to the fact that its subleading order interactions with the fluctuation field are of Boltzmann type (the leading order is determined by the HFB dynamics); the latter are not drowned out by the error term, due to the largeness of N . On the other hand, a quartic Boltzmann collision term emerges, as expected, from our analysis, but it is a lower order term that is buried in the error terms because it does not couple to the condensate. We refer to Proposition 5.1 for more details. We do expect an analysis to the next subleading order, conjecturally involving appropriate quantum corrections to the BEC and HFB dynamics, to reveal this fourth order collision term of Boltzmann-Uhlenbeck-Uehling type, separated from the error term.

Remark 2.10. If it is assumed that the BEC is non-stationary and that both the BEC and the thermal fluctuation field are not translationally invariant, the first term in the Duhamel expansion for $\nu_t(a_p^+ a_q)$, expressed with a single time integral, does not vanish, contrary to the situation considered in the paper at hand. This additional term couples nontrivially to both the HFB and the Boltzmann dynamics,

and will lead to a different system of PDEs than the one derived here. We expect the rigorous analysis of this system to be considerably more involved, and leave it for future work.

2.4. Sketch of the proof. The strategy of our proof is to start by calculating one collision, corresponding to calculating the Duhamel expansion to second order in the coupling λ . The tail in the Duhamel expansion corresponds to recollisions. Using bounds available in the second-quantization formulation, we are able to bound this tail. For that, it is crucial to exploit the fact that the initial condensate density N provides a large perturbation parameter. Accordingly, the resulting leading order cubic Boltzmann collision operator appears with an overall factor $1/N$. This allows us to compare the density of thermal fluctuation particles for positive times with the initial density. We can observe that the superposition of HFB oscillations results in corrections to the leading order $2 \leftrightarrow 1$ processes coming from the collision of two thermal particles and one of those being absorbed into or emitted from the BEC. In addition, if we consider the continuous approximation, we show that the collision operator can be approximated by its continuous counterpart.

We start by applying Duhamel's principle to (2.22), using one recursion for Φ , and two for f and g . The main term is recovered by evaluating the terms involving ν_0 . In Section 4, in order to control higher-order terms involving the full dynamics ν_t , we establish uniform-in-time bounds on $f[J]$, $g[J]$, $\mathcal{H}_{cub}(t)$, and $\mathcal{H}_{quart}(t)$ with respect to the number operator \mathcal{N}_b . We use an a-priori bound on the growth of $\nu_t(\mathcal{N}_b^k)$ established in [74, 269]. This will yield the restriction $|\Lambda|/\lambda = O(\log N / \log \log N)$. Moreover, we show closeness of ν_t to ν_0 in a suitable sense, which allows us to exploit that ν_0 is approximately quasifree.

In Section 5, we control the tail term in the Duhamel expansion using the previously established operator bounds and quantify propagation of moments of the number operator, using [74, 269]. The terms in the Duhamel expansion are expressed by way of $a(t) = \mathcal{V}_{HFB}^*(t)a\mathcal{V}_{HFB}(t)$ and $a^+(t)$. \mathcal{H}_{HFB} is quadratic in a and a^+ , thus $a(t)$ and $a^+(t)$ are linear combinations of a and a^+ . That enables us to control the proximity of ν_t to ν_0 , which is quasifree, in order to evaluate the main terms in the Duhamel expansion by means of Wick's Theorem.

In the discrete case, we observe interference phenomena in our scaling regime $L \leq t \sim \lambda^{-2}$. The Boltzmann collision terms are lattice sums in momentum space, and their magnitudes depend on whether terms that vary with time have particular values in \mathbb{Q} . This effect becomes negligible for box length $L \sim \lambda^{-2-} \gg t$ and $\lambda = \lambda(N) > 0$ chosen small enough. The latter means that we will ultimately set $\lambda = \lambda(N)$, and choose N large enough.

We prove a discretization Lemma 5.5 that allows us to improve the rate of convergence dependent on the regularity of f_0 and \hat{v} , beyond the trivial bound. As we noticed after completing this work, our approach here appears to be related to the numerical error estimates for the trapezoid rule in numerical mathematics via Poisson summation, see [294].

We use the Duhamel expansion and the approximate quasifreeness of ν_t to relate f_0 back to f_s . In order to control the large magnitude of condensate terms in the expansion, we need to rewrite the equations for the centered expectations $f_s(p) - |\Phi_s|^2 \delta(p)$ and $g_s(p) - \Phi_s^2 \delta(p)$.

In Section 6, we collect all results to compute the main order terms in Theorem 2.1 and 2.2, and in section 7, we compute the effective equations.

Remark 2.11. *We note that the Boltzmann-type equations in Theorem 2.1 and 2.2 are presented in their integral and weak forms. No smallness assumption on f is necessary for our result to hold. In this work, we will not further investigate questions regarding the well-posedness of the corresponding Cauchy problem of kinetic equations in the context of nonlinear PDE. Some works in this direction are referenced in Section 1.1.2, see in particular [27, 29, 140, 141, 142].*

3. PRELIMINARIES

Some of our estimates will be formulated for finite number subspaces of \mathcal{F} . We introduce the projectors

$$P_n := P_{\mathcal{F}_n}. \quad (3.1)$$

We will consider \mathcal{F}_n embedded into \mathcal{F} and we identify P_n as maps $\mathcal{F} \rightarrow \mathcal{F}$. By the spectral theorem,

$$P_n = \frac{1}{2\pi i} \oint_{\partial B_{1/2}(n)} \frac{dz}{\mathcal{N}_b - z}. \quad (3.2)$$

Observe that $a_x|_{F_n}$ defines a map

$$a_x : F_n \rightarrow F_{n-1} \quad (3.3)$$

for all $n \in \mathbb{N}$, with formal adjoint $a_x^+|_{F_{n-1}} : F_{n-1} \rightarrow F_n$. To study the weak formulation of the effective equations, we introduce

$$\begin{aligned} f[J] &:= \int dp J(p) a_p^+ a_p, \\ g[J] &:= \int dp J(p) a_{-p} a_p, \\ g^*[J] &:= g[J]^+ = \int dp J(p) a_p^+ a_{-p}^+. \end{aligned} \quad (3.4)$$

As a convenient notation for iterating Duhamel's formula, let

$$\Delta(t, j) := \{\mathbf{s}_j \in [0, t]^j \mid s_1 \geq \dots \geq s_j\} \quad (3.5)$$

be a j -simplex, where $t > 0$ and $j \in \mathbb{N}$. We also recall that

$$\mathcal{H}_I(t) = \mathcal{H}_{cub}(t) + \mathcal{H}_{quart}(t). \quad (3.6)$$

Let $(\mathcal{V}_{HFB}(t))_{t \in \mathbb{R}}$ be the solution of

$$\begin{cases} i\partial_t \mathcal{V}_{HFB}(t) &= \mathcal{H}_{HFB} \mathcal{V}_{HFB}(t), \\ \mathcal{V}_{HFB}(0) &= \mathbf{1}. \end{cases} \quad (3.7)$$

We denote the $d - 1$ -dimensional Hausdorff measure of a smooth, embedded hypersurface in \mathbb{R}^d by $d\mathcal{H}^{d-1}$.

If needed, we will keep track of the dependence of constants on parameters by adding the respective parameters in a subscript. Whenever the constants have no explicit dependence, they are assumed to be universal.

The following result is a direct consequence from iterating (2.22).

Lemma 3.1. *Assume that $A \in \mathfrak{A}$ is an observable, and that ν_t is defined as in (2.22). Then, for any $k \in \mathbb{N}$, we have that*

$$\begin{aligned} \nu_t(A) &= \sum_{\ell=0}^{k-1} (-i)^\ell \int_{\Delta[t,\ell]} ds_\ell \nu_0([\cdot \cdots [A, \mathcal{H}_I(s_1)], \cdots], \mathcal{H}_I(s_\ell)] \\ &\quad + (-i)^k \int_{\Delta[t,k]} ds_k \nu_{s_k}([\cdot \cdots [A, \mathcal{H}_I(s_1)], \cdots], \mathcal{H}_I(s_k)]. \end{aligned} \quad (3.8)$$

We will be particularly interested in the cases $A \in \{a_0, f[J], g[J]\}$. To study the expansion, we derive the following useful identity.

Lemma 3.2. *Let*

$$\Omega(p) := \sqrt{E(p)(E(p) + 2\lambda\hat{v}(p))} \quad (3.9)$$

denote the dispersion function for acoustic excitations, and let

$$a_p^+(t) := \mathcal{V}_{HFB}^*(t) a_p^+ \mathcal{V}_{HFB}(t), \quad (3.10)$$

$$a_p(t) = \mathcal{V}_{HFB}^*(t) a_p \mathcal{V}_{HFB}(t). \quad (3.11)$$

Moreover, let

$$\mathcal{M}(p) := \begin{pmatrix} -E(p) - \lambda\hat{v}(p) & -\lambda\hat{v}(p) \\ \lambda\hat{v}(p) & E(p) + \lambda\hat{v}(p) \end{pmatrix}. \quad (3.12)$$

Then,

$$\begin{pmatrix} a_p^+(t) \\ a_{-p}(t) \end{pmatrix} = \left[\cos(t\Omega(p)) \mathbf{1} - i \frac{\sin(t\Omega(p))}{\Omega(p)} \mathcal{M}(p) \right] \begin{pmatrix} a_p^+ \\ a_{-p} \end{pmatrix}. \quad (3.13)$$

Proof. We obtain the following system of ODEs,

$$\begin{aligned} i\partial_t a_p^+(t) &= \mathcal{V}_{HFB}^*(t) [a_p^+, \mathcal{H}_{HFB}^{\phi_0}] \mathcal{V}_{HFB}(t) \\ &= -(E(p) + \lambda\hat{v}(p)) a_p^+(t) - \lambda\hat{v}(p) a_{-p}(t), \end{aligned} \quad (3.14)$$

$$\begin{aligned} i\partial_t a_{-p}(t) &= \mathcal{V}_{HFB}^*(t) [a_{-p}, \mathcal{H}_{HFB}^{\phi_0}] \mathcal{V}_{HFB}(t) \\ &= (E(p) + \lambda\hat{v}(p)) a_{-p}(t) + \lambda\hat{v}(p) a_p^+(t), \end{aligned} \quad (3.15)$$

where $E(p) = \frac{v^2}{2}$, so that

$$i\partial_t \begin{pmatrix} a_p^+(t) \\ a_{-p}(t) \end{pmatrix} = \mathcal{M}(p) \begin{pmatrix} a_p^+(t) \\ a_{-p}(t) \end{pmatrix} \quad (3.16)$$

with

$$\mathcal{M}(p) := \begin{pmatrix} -E(p) - \lambda\hat{v}(p) & -\lambda\hat{v}(p) \\ \lambda\hat{v}(p) & E(p) + \lambda\hat{v}(p) \end{pmatrix}. \quad (3.17)$$

We observe that $\mathcal{M}(p)$ satisfies

$$\mathcal{M}^2(p) = \Omega^2(p) \mathbf{1}, \quad (3.18)$$

where

$$\Omega(p) := \sqrt{E(p)(E(p) + 2\lambda\hat{v}(p))} \quad (3.19)$$

denotes the dispersion function of acoustic excitations; see, for instance, [112, 274]. In particular, we have $\mathcal{M}^{-1}(p) = \frac{1}{\Omega^2(p)} \mathcal{M}(p)$.

Hence,

$$\begin{aligned} \begin{pmatrix} a_p^+(t) \\ a_{-p}(t) \end{pmatrix} &= \exp \left(-it \begin{pmatrix} -E(p) - \lambda \hat{v}(p) & -\lambda \hat{v}(p) \\ \lambda \hat{v}(p) & E(p) + \lambda \hat{v}(p) \end{pmatrix} \right) \begin{pmatrix} a_p^+ \\ a_{-p} \end{pmatrix} \\ &= \left(\cos(t\Omega(p)) \mathbf{1} - i \sin(t\Omega(p)) \frac{\mathcal{M}(p)}{\Omega(p)} \right) \begin{pmatrix} a_p^+ \\ a_{-p} \end{pmatrix}. \end{aligned} \quad (3.20)$$

This finishes the proof. \square

Corollary 3.3. *Assume $q \in [1, \infty]$ and $n \in \{0, 1, 2, 3\}$. Moreover, recall the assumptions $\hat{v} \geq 0$ with $\hat{v}(0) = 0$ from Section 2.2. V_1, V_2 , defined by*

$$a_p^+(t) = (e^{i\Omega(p)t} + i\lambda^2 \sin(\Omega(p)t) V_1(p)^2) a_p^+ + i\lambda \sin(\Omega(p)t) V_2(p) a_{-p}, \quad (3.21)$$

are given by

$$V_1(p)^2 := \frac{\hat{v}(p)^2}{\Omega(p)(\Omega(p) + E(p) + \lambda \hat{v}(p))}, \quad (3.22)$$

$$V_2(p) := \frac{\hat{v}(p)}{\Omega(p)}. \quad (3.23)$$

They satisfy the bounds

$$\|D^n V_1^2\|_q \leq C \|\hat{v}\|_{2(\lfloor \frac{n}{2} \rfloor + 1), w, c}^{n+2}, \quad (3.24)$$

$$\|D^n V_2\|_q \leq C \|\hat{v}\|_{2(\lfloor \frac{n}{2} \rfloor + 1), w, c}^{n+1}. \quad (3.25)$$

Similarly,

$$u_\lambda(t, p) := e^{i\Omega(p)t} + i\lambda^2 \sin(\Omega(p)t) V_1(p)^2 \quad (3.26)$$

$$v_\lambda(t, p) := i\lambda \sin(\Omega(p)t) V_2(p) \quad (3.27)$$

satisfy

$$\|u_\lambda(t, \cdot)\|_{\ell^\infty(\Lambda^*)} \leq C(1 + \lambda^2 \|\hat{v}\|_{w, d}^2), \quad (3.28)$$

$$\|v_\lambda(t, \cdot)\|_d \leq C\lambda \|\hat{v}\|_{w, d} \quad (3.29)$$

for all $q \in [1, \infty]$, all $\lambda \in (0, 1]$, and all $t > 0$.

Proof. Lemma 3.2 implies

$$a_p^+(t) = \cos(t\Omega(p)) a_p^+ - i \sin(t\Omega(p)) \left(\frac{-E(p) - \lambda \hat{v}(p)}{\Omega(p)} a_p^+ - \lambda \frac{\hat{v}(p)}{\Omega(p)} a_{-p} \right). \quad (3.30)$$

Define $a := E(p)$ and $b := E(p) + 2\lambda \hat{v}(p)$. Setting

$$V_1(p)^2 = \frac{1}{\lambda^2} \left(\frac{a+b}{2\sqrt{ab}} - 1 \right), \quad (3.31)$$

$$V_2(p) = \frac{1}{\lambda} \frac{b-a}{2\sqrt{ab}} = \frac{\hat{v}(p)}{\Omega(p)}, \quad (3.32)$$

we have that

$$a_p^+(t) = (e^{i\Omega(p)t} + i\lambda^2 \sin(\Omega(p)t) V_1(p)^2) a_p^+ + i\lambda \sin(\Omega(p)t) V_2(p) a_{-p}. \quad (3.33)$$

We start by establishing pointwise bounds. We have

$$\frac{a+b}{2\sqrt{ab}} - 1 = \frac{(\sqrt{a} - \sqrt{b})^2}{2\sqrt{ab}} = \frac{(b-a)^2}{2\sqrt{ab}(\sqrt{a} + \sqrt{b})^2}. \quad (3.34)$$

In particular, we get

$$\begin{aligned} V_1(p)^2 &= \frac{2\hat{v}(p)^2}{\Omega(p)(\sqrt{E(p)} + \sqrt{E(p) + 2\lambda\hat{v}(p)})^2} \\ &= \frac{\hat{v}(p)^2}{\Omega(p)(\Omega(p) + E(p) + \lambda\hat{v}(p))}. \end{aligned} \quad (3.35)$$

Using $\hat{v} \geq 0$, we have the pointwise bounds

$$|V_1^2| \leq 2\frac{\hat{v}^2}{E^2}, \quad (3.36)$$

$$|V_2| \leq \frac{\hat{v}}{E}. \quad (3.37)$$

We thus have that

$$\begin{aligned} \|V_1^2\|_d &\leq C\|\hat{v}\|_{w,d}^2, \\ \|V_2\|_d &\leq C\|\hat{v}\|_{w,d}. \end{aligned} \quad (3.38)$$

Finally, set

$$h_1(x) := \frac{2x^2}{\sqrt{1 + 2\lambda x}(1 + \sqrt{1 + 2\lambda x})^2}, \quad (3.39)$$

$$h_2(x) := \frac{x}{\sqrt{1 + 2\lambda x}}. \quad (3.40)$$

We have that

$$V_1(p)^2 = h_1\left(\frac{\hat{v}(p)}{E(p)}\right), \quad (3.41)$$

$$V_2(p) = h_2\left(\frac{\hat{v}(p)}{E(p)}\right). \quad (3.42)$$

Using Lemma C.3 with $n \in \{1, 2, 3\}$, we obtain

$$\|D^n V_1^2\|_q \leq C\|h_1\|_{C^n([0, \|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c}]})} \|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c}^n, \quad (3.43)$$

$$\|D^n V_2\|_q \leq C\|h_2\|_{C^n([0, \|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c}]})} \|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c}^n. \quad (3.44)$$

A straight-forward calculation yields

$$\|h_1\|_{C^n([0, \|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c}]})} \leq C\|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c}^2, \quad (3.45)$$

$$\|h_2\|_{C^n([0, \|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c}]})} \leq C\|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c}. \quad (3.46)$$

Combining (3.43), (3.44), (3.45), and (3.46) finishes the proof. \square

4. TRACE ESTIMATES

In this chapter, we omit the subscript Λ^* from \int_{Λ^*} in our notation, as we will only consider $\int \equiv \int_{\Lambda^*}$ as sums over Λ^* in the sense of (1.51).

4.1. Computation of the partition function. We establish estimates on the partition function on finite particle number subspaces of the Fock space \mathcal{F} . The estimates derived here are needed to control the error terms coming from the expansion in Lemma 3.1.

For the next lemma, let us introduce some notation. Let us denote

$$a_p^{(1)} := a_p^+ \quad \text{and} \quad a_p^{(-1)} := a_p. \quad (4.1)$$

Let

$$\begin{aligned} \mathcal{P}[a, a^+] := & \left\{ \int d\mathbf{p}_k h(\mathbf{p}_k, \vec{\sigma}_k) \prod_{j=1}^k a_{p_j}^{(\sigma_j)} \mid k \in \mathbb{N}, h(\cdot, \vec{\sigma}_k) \in \mathcal{S}'(\mathbb{R}^3), \right. \\ & \left. \vec{\sigma}_k = (\sigma_1, \dots, \sigma_k) \in \{\pm 1\}^k \right\} \end{aligned} \quad (4.2)$$

denote all monomials in a, a^+ . Here and henceforth, we define ordered operator products via multiplication from the right, by

$$\prod_{j=1}^{n+1} A_j := \left(\prod_{j=1}^n A_j \right) A_{n+1}, \quad n \in \mathbb{N}_0, \quad (4.3)$$

$$\prod_{j=1}^0 A_j := \mathbf{1}. \quad (4.4)$$

Let us denote

$$\text{sign}(A) := \sum_{j=1}^k \sigma_j \quad (4.5)$$

whenever $A = \int d\mathbf{p}_k h(\mathbf{p}_k, \vec{\sigma}_k) \prod_{j=1}^k a_{p_j}^{(\sigma_j)}$. Note that we have

$$[\mathcal{N}_b, \prod_{j=1}^k a_{p_j}^{(\sigma_j)}] = \sum_{j=1}^k \sigma_j \prod_{j=1}^k a_{p_j}^{(\sigma_j)}, \quad (4.6)$$

which, in turn, gives

$$[\mathcal{N}_b, A] = \text{sign}(A)A \quad (4.7)$$

whenever $A = \int d\mathbf{p}_k h(\mathbf{p}_k, \sigma_k) \prod_{j=1}^k a_{p_j}^{(\sigma_j)}$. Using the spectral decomposition

$$P_n = \frac{1}{2\pi i} \oint_{\partial B_{\frac{1}{2}}(n)} \frac{dz}{\mathcal{N}_b - z}, \quad (4.8)$$

with counter-clockwise contour, it follows that

$$P_n A = A P_{n - \text{sign}(A)} \quad (4.9)$$

for all $A \in \mathcal{P}[a, a^+]$. We will refer to (4.9) as the *Pull-Through Formula* for projectors, see [35].

Moreover, if $\text{sign}(A) \neq 0$, (4.7) implies that

$$\nu(A) = \frac{1}{\text{sign}(A)} \nu([\mathcal{N}_b, A]) = 0. \quad (4.10)$$

Lemma 4.1 (Moments of the number operator). *We have for all $\ell \in \mathbb{N}_0$ that*

$$\nu_0((\mathcal{N}_b + 1)^{\frac{\ell}{2}}) \leq C_{\ell, \|f_0\|_d} |\Lambda|^{\frac{\ell}{2}}. \quad (4.11)$$

Proof. Let $\mathcal{K}_\mu := \int_{\Lambda^*} dp (K(p) - \mu) a_p^+ a_p$. Then we obtain that

$$\begin{aligned} Z_0(\mu) &:= \text{Tr}(e^{-\mathcal{K}_\mu}) \\ &= \prod_{p \in \Lambda^*} \sum_{n=0}^{\infty} e^{-n(K(p) - \mu)} \\ &= \prod_{p \in \Lambda^*} \frac{1}{1 - e^{-(K(p) - \mu)}} \\ &= e^{-|\Lambda| \int_{\Lambda^*} dp \log(1 - e^{-(K(p) - \mu)})} \end{aligned} \quad (4.12)$$

for any $\mu < \kappa_0$. Let

$$\kappa_n := -(-\partial_\mu)^n \Big|_{\mu=0} \int_{\Lambda^*} dp \log(1 - e^{-(K(p) - \mu)}) \quad (4.13)$$

denote the n^{th} cumulant for $n \in \mathbb{N}$. Let $\binom{n}{\mathbf{r}_\ell}$ denote the multinomial coefficient, and $R(\ell) := \{\mathbf{r}_\ell \in \mathbb{N}_0^\ell \mid \sum_{n=1}^\ell n r_n = \ell\}$. Then the Faà di Bruno formula, see [295], yields

$$\begin{aligned} \nu_0(\mathcal{N}_b^\ell) &= \frac{(-\partial_\mu)^\ell \Big|_{\mu=0} Z_0(\mu)}{Z_0(0)} \\ &= \ell! \sum_{\mathbf{r}_\ell \in R(\ell)} \prod_{n=1}^\ell \frac{1}{r_n!} \left(\frac{|\Lambda| \kappa_n}{n!} \right)^{r_n}. \end{aligned} \quad (4.14)$$

Observe that we have

$$\kappa_1 = \int_{\Lambda^*} dp \frac{1}{e^{K(p)} - 1} = \int_{\Lambda^*} dp f_0(p), \quad (4.15)$$

recalling f_0 from (1.57). More generally, Lemma A.2 implies

$$\kappa_n \leq C_{n, \|f_0\|_a} \quad (4.16)$$

for all $n \in \mathbb{N}$. As a consequence of (4.14) and (4.16), we find that

$$\nu_0((\mathcal{N}_b + 1)^\ell) \leq C_{\ell, \|f_0\|_a} |\Lambda|^\ell, \quad (4.17)$$

where we used that $|\Lambda| \geq 1$. Using

$$\nu_0((\mathcal{N}_b + 1)^{\frac{\ell}{2}}) \leq \nu_0((\mathcal{N}_b + 1)^{\ell-1})^{\frac{1}{2}} \nu_0((\mathcal{N}_b + 1)^{\ell+1})^{\frac{1}{2}} \quad (4.18)$$

yields the half-integer case. This concludes the proof. \square

Remark 4.2. We use an argument from [117]. Using the fact that

$$e^{-\mathcal{K}} a_p^+ e^{\mathcal{K}} = e^{-K(p)} a_p^+, \quad (4.19)$$

cyclicity of the trace implies

$$\begin{aligned} f_0(p) &= \frac{\text{Tr}(e^{-\mathcal{K}} a_p^+ a_p)}{\text{Tr} e^{-\mathcal{K}} |\Lambda|} \\ &= e^{-K(p)} \frac{\text{Tr}(a_p^+ e^{-\mathcal{K}} a_p)}{\text{Tr} e^{-\mathcal{K}} |\Lambda|} \\ &= e^{-K(p)} \frac{\text{Tr}(e^{-\mathcal{K}} a_p a_p^+)}{\text{Tr} e^{-\mathcal{K}} |\Lambda|} \\ &= e^{-K(p)} (f_0(p) + 1), \end{aligned} \quad (4.20)$$

Thus, we have

$$f_0(p) = \frac{1}{e^{K(p)} - 1}. \quad (4.21)$$

Lemma 4.3 (Operator product bound). *Let $A_j \in \mathcal{P}[a, a^+]$ be monomials in a, a^+ , $\gamma_j > 0$, and $k_j \in \mathbb{N}$ be such that*

$$\|P_m A_j P_{m-\text{sign}(A_j)}\| \leq \gamma_j (m + |\Lambda|)^{k_j/2} \quad (4.22)$$

for all $j \in \{1, \dots, \ell\}$ and all $m \in \mathbb{N}_0$. Then we have that

$$|\nu(\prod_{j=1}^{\ell} A_j)| \leq \left(\prod_{j=1}^{\ell} \gamma_j \right) \nu \left((\mathcal{N} + \sum_{m=1}^{\ell} |\text{sign}(A_m)| + |\Lambda|)^{\sum_{j=1}^{\ell} k_j/2} \right) \quad (4.23)$$

for any state ν .

Proof. Observe that

$$\text{sign}(A) = \sum_{j=1}^{\ell} \text{sign}(A_j). \quad (4.24)$$

In addition,

$$\sum_{n=0}^{\infty} P_n = \mathbf{1}, \quad (4.25)$$

$$P_n f(\mathcal{N}_b) = P_n f(n) \quad (4.26)$$

holds for any measurable function f . Hence and by applying (4.9), we have that

$$\begin{aligned} |\nu(A)| &\leq \sum_{n=0}^{\infty} |\nu(P_n A P_{n-\text{sign}(A)})| \\ &= \sum_{n=0}^{\infty} |\nu(P_n \left(\prod_{j=1}^{\ell} P_{n-\sum_{m=1}^{j-1} \text{sign}(A_m)} A_j P_{n-\sum_{m=1}^j \text{sign}(A_m)} \right) P_{n-\text{sign}(A)})| \end{aligned} \quad (4.27)$$

Denoting $B_n := \prod_{j=1}^{\ell} P_{n-\sum_{m=1}^{j-1} \text{sign}(A_m)} A_j P_{n-\sum_{m=1}^j \text{sign}(A_m)}$, we can apply Cauchy-Schwarz to ν to obtain the upper bound

$$\begin{aligned} |\nu(A)| &\leq \sum_{n=0}^{\infty} \nu(P_n)^{\frac{1}{2}} \nu(P_{n-\text{sign}(A)} B_n^+ B_n P_{n-\text{sign}(A)})^{\frac{1}{2}} \\ &\leq \sum_{n=0}^{\infty} \|B_n\| \nu(P_n)^{\frac{1}{2}} \nu(P_{n-\text{sign}(A)})^{\frac{1}{2}} \\ &\leq \left(\sum_{n=0}^{\infty} \|B_n\| \nu(P_n) \right)^{\frac{1}{2}} \left(\sum_{n=0}^{\infty} \|B_n\| \nu(P_{n-\text{sign}(A)}) \right)^{\frac{1}{2}} \\ &= \left(\sum_{n=0}^{\infty} \|B_n\| \nu(P_n) \right)^{\frac{1}{2}} \left(\sum_{n=\text{sign}(A)-}^{\infty} \|B_{n+\text{sign}(A)}\| \nu(P_n) \right)^{\frac{1}{2}}. \end{aligned} \quad (4.28)$$

Here, we used Cauchy-Schwarz to $\sum_{n=0}^{\infty}$, followed by the fact that $P_{n-\text{sign}(A)} \equiv 0$ for all $n < \text{sign}(A)_+$, as $\mathcal{N}_b \geq 0$ in the sense of quadratic forms. Notice that we

have

$$\begin{aligned}
\|B_n\| &\leq \prod_{j=1}^{\ell} \|P_{n-\sum_{m=1}^{j-1} \text{sign}(A_m)} A_j P_{n-\sum_{m=1}^j \text{sign}(A_m)}\| \\
&\leq \prod_{j=1}^{\ell} \gamma_j \left(\left(n - \sum_{m=1}^{j-1} \text{sign}(A_m) \right)_+ + |\Lambda| \right)^{k_j/2} \\
&\leq \left(\prod_{j=1}^{\ell} \gamma_j \right) \left(n + \sum_{m=1}^{\ell} |\text{sign}(A_m)| + |\Lambda| \right)^{\sum_{j=1}^{\ell} k_j/2} \tag{4.29}
\end{aligned}$$

by assumption (4.22). Similarly, we have that

$$\begin{aligned}
\|B_{n+\text{sign}(A)}\| &\leq \prod_{j=1}^{\ell} \gamma_j \left(\left(n + \text{sign}(A) - \sum_{m=1}^{j-1} \text{sign}(A_m) \right)_+ + |\Lambda| \right)^{k_j/2} \\
&= \prod_{j=1}^{\ell} \gamma_j \left(\left(n + \sum_{m=j}^{\ell} \text{sign}(A_m) \right)_+ + |\Lambda| \right)^{k_j/2} \\
&\leq \left(\prod_{j=1}^{\ell} \gamma_j \right) \left(n + \sum_{m=1}^{\ell} |\text{sign}(A_m)| + |\Lambda| \right)^{\sum_{j=1}^{\ell} k_j/2} \tag{4.30}
\end{aligned}$$

due to (4.24). Collecting (4.28)–(4.30) and applying (4.26) again, we then obtain

$$\begin{aligned}
|\nu(A)| &\leq \left(\prod_{j=1}^{\ell} \gamma_j \right) \sum_{n=0}^{\infty} \left(n + \sum_{m=1}^{\ell} |\text{sign}(A_m)| + \max_{1 \leq j \leq \ell} k_j \right)^{\sum_{j=1}^{\ell} k_j/2} \nu(P_n) \\
&= \left(\prod_{j=1}^{\ell} \gamma_j \right) \nu \left(\mathcal{N} + \sum_{m=1}^{\ell} |\text{sign}(A_m)| + |\Lambda| \right)^{\sum_{j=1}^{\ell} k_j/2} \tag{4.31}
\end{aligned}$$

□

4.2. Operator estimates w.r.t. \mathcal{N}_b . In our analysis, we will bound various operators relative to powers of the particle number operator \mathcal{N}_b , in conjunction with previously established estimates on the partition function.

Lemma 4.4 ($\mathcal{H}_{cub}(t)$ bound). *We have that*

$$\mathcal{H}_{cub}(t) = \sum_{k=1}^{16} A_{cub}^{(k)}(t) \tag{4.32}$$

with monomials $A_{cub}^{(k)}(t) \in \mathcal{P}[a, a^+]$ such that

$$\|A_{cub}^{(k)}(t) P_M\| \leq C \|\hat{v}\|_{w,d} (1 + \lambda \|\hat{v}\|_{w,d})^6 \frac{\lambda(M + |\Lambda|)^{\frac{3}{2}}}{\sqrt{N}} \tag{4.33}$$

for any $M \in \mathbb{N}_0$, and $|\text{sign}(A_{cub}^{(k)}(t))| \leq 3$.

Proof. Corollary 3.3 and evenness of \hat{v} imply that

$$\begin{aligned}
&\mathcal{H}_{cub}(t) \\
&= \frac{\lambda}{\sqrt{N}} \int_{(\Lambda^*)^3} d\mathbf{p}_3 \hat{v}(p_2) \delta(p_1 + p_2 - p_3) a_{p_1}^+(t) a_{p_2}^+(t) a_{p_3}(t) + h.c.
\end{aligned}$$

$$= \frac{\lambda}{\sqrt{N}} \sum_{\vec{\sigma}_3 \in \{\pm 1\}^3} \int_{(\Lambda^*)^3} d\mathbf{p}_3 \delta\left(\sum_{j=1}^3 \sigma_j p_j\right) \hat{v}(p_2) \prod_{j=1}^3 (g_j(p_j, \sigma_j) a_{p_j}^{(\sigma_j)}) + h.c., \quad (4.34)$$

where $\vec{\sigma}_3 = (\sigma_1, \sigma_2, \sigma_3)$, and

$$g_j(p_j, \sigma_j) := \delta_{\sigma_j, 1} u_\lambda(t, p_j) + \delta_{\sigma_j, -1} v_\lambda(t, p_j), \quad j \in \{1, 2\}, \quad (4.35)$$

$$g_3(p_3, \sigma_3) := \delta_{\sigma_3, -1} \overline{u_\lambda}(t, p_3) + \delta_{\sigma_3, 1} \overline{v_\lambda}(t, p_3). \quad (4.36)$$

Abbreviating

$$A_{cub}(\vec{\sigma}_3) := \int d\mathbf{p}_3 \delta\left(\sum_{j=1}^3 \sigma_j p_j\right) \hat{v}(p_2) \prod_{j=1}^3 (g_j(p_j, \sigma_j) a_{p_j}^{(\sigma_j)}), \quad (4.37)$$

it is sufficient to prove (4.33) for $A_{cub}(\vec{\sigma}_3)$, since the adjoint satisfies

$$\begin{aligned} \|A_{cub}(\vec{\sigma}_3)^* P_M\| &= \|P_{M - \sum_{j=1}^3 \sigma_j} A_{cub}(\vec{\sigma}_3)^* P_M\| \\ &= \|A_{cub}(\vec{\sigma}_3) P_{M - \sum_{j=1}^3 \sigma_j}\| \end{aligned} \quad (4.38)$$

due to the Pull-Through formula (4.9).

Using Wick's Theorem in Appendix A, we have that

$$\begin{aligned} a_{p_1}^{(\sigma_1)} a_{p_2}^{(\sigma_2)} a_{p_3}^{(\sigma_3)} \delta\left(\sum_{j=1}^3 \sigma_j p_j\right) &= : a_{p_1}^{(\sigma_1)} a_{p_2}^{(\sigma_2)} a_{p_3}^{(\sigma_3)} : \delta\left(\sum_{j=1}^3 \sigma_j p_j\right) \\ &\quad + \delta(p_1 - p_2) \delta(p_3) \delta_{\sigma_1, -1} \delta_{\sigma_2, 1} a_0^{(\sigma_3)} \\ &\quad + \delta(p_1 - p_3) \delta(p_2) \delta_{\sigma_1, -1} \delta_{\sigma_3, 1} a_0^{(\sigma_2)} \\ &\quad + \delta(p_2 - p_3) \delta(p_1) \delta_{\sigma_2, -1} \delta_{\sigma_3, 1} a_0^{(\sigma_1)}. \end{aligned} \quad (4.39)$$

We want to apply Lemma A.4 to each of the terms associated with the terms in (4.39). For that, we need to ensure integrability for each of those terms. The terms involving $a_0^{(\sigma)}$ contain 2 momentum- δ where the only free momentum comes with a coefficient v_λ resp. $\overline{v_\lambda}$. Using

$$\|a_0^+ P_M\| = \|a_0 P_{M+1}\|, \quad (4.40)$$

Lemma A.4 then implies

$$\begin{aligned} \|a_0^{(\sigma)} P_M\| &= \delta_{\sigma, 1} \|a_0^+ P_M\| + \delta_{\sigma, -1} \|a_0 P_M\| \\ &\leq \sqrt{|\Lambda|(M+1)}. \end{aligned} \quad (4.41)$$

For the cubic term in (4.39) and if not all $\sigma_j = 1$ or all $\sigma_j = -1$, Lemma A.4 implies

$$\begin{aligned} &\| : A_{cub}(\vec{\sigma}_3) : P_M \| \\ &\leq \|\hat{v}(p_2)\|^{1/2} \prod_{j=1}^2 |v_\lambda(p_j)|^{\delta_{\sigma_j, -1}} |u_\lambda(p_3)|^{\delta_{\sigma_3, -1}} \delta\left(\sum_{j=1}^3 p_j \sigma_j\right)^{1/2} \|L_{\mathbf{p}_{J_+}}^\infty L_{\mathbf{p}_{J_-}}^2\| \\ &\quad \|\hat{v}(p_2)\|^{1/2} \prod_{j=1}^2 |u_\lambda(p_j)|^{\delta_{\sigma_j, 1}} |v_\lambda(p_3)|^{\delta_{\sigma_3, 1}} \delta\left(\sum_{j=1}^3 p_j \sigma_j\right)^{1/2} \|L_{\mathbf{p}_{J_-}}^\infty L_{\mathbf{p}_{J_+}}^2\| \\ &= (M + \sum_{j=1}^3 \sigma_j)_{|J_+|}^{1/2} (M)_{|J_-|}^{1/2}, \end{aligned} \quad (4.42)$$

using the notation in Lemma A.4 with $n = 3$. Integrability of $\hat{v}(p_2)$ is sufficient to yield that the RHS is, indeed, finite. If $\sigma_j = 1$ for all j , we have that

$$\begin{aligned} & \| : A_{cub}(1, 1, 1) : P_M \| \\ & \leq \left(\|\hat{v}(p_2)u_\lambda(t, p_1)u_\lambda(t, p_2)\overline{v_\lambda}(t, p_3)\|_{L_{p_2}^2 L_{p_1, p_3}^\infty}^2 (M - 2) \right. \\ & \quad \left. + \|\hat{v}(p_2)u_\lambda(t, p_1)u_\lambda(t, p_2)\overline{v_\lambda}(t, p_1 + p_2)\|_{L_{p_2}^2}^2 |\Lambda| \right)^{\frac{1}{2}} \sqrt{M(M - 1)}. \end{aligned} \quad (4.43)$$

Similarly, we find that

$$\begin{aligned} & \| : A_{cub}(-1, -1, -1) : P_M \| \\ & \leq \left(\|\hat{v}(p_2)v_\lambda(t, p_1)v_\lambda(t, p_2)\overline{u_\lambda}(t, p_3)\|_{L_{p_2}^2 L_{p_1, p_3}^\infty}^2 (M + 1) \right. \\ & \quad \left. + \|\hat{v}(p_2)v_\lambda(t, p_1)v_\lambda(t, p_2)\overline{u_\lambda}(t, p_1 + p_2)\|_{L_{p_2}^2 L_{p_1}^\infty}^2 |\Lambda| \right)^{\frac{1}{2}} \sqrt{(M + 3)(M + 2)}. \end{aligned} \quad (4.44)$$

Observe that

$$\begin{aligned} \|H\|_{L^2(\Lambda^*)} &= \frac{1}{\sqrt{|\Lambda|}} \|H\|_{\ell^2(\Lambda^*)} \\ &\leq \|H\|_d. \end{aligned} \quad (4.45)$$

Since we are summing over a lattice, there are terms for which momentum requires the summation over $\hat{v}(2p)$ or $v_\lambda(2p)$ resp. $\overline{v_\lambda}(2p)$. In order to use the upper bound $\|\hat{v}\|_{w,d}$, we employ the fact that

$$\left| \sum_{p \in \Lambda^*} H(2p) \right| \leq \|H\|_{\ell^1(\Lambda^*)}. \quad (4.46)$$

Collecting (4.38), (4.41), (4.42), (4.43), (4.44), (4.45), and employing Corollary 3.3, we have shown that

$$\|A_{cub}(\vec{\sigma}_3)P_M\| \leq C\|\hat{v}\|_{w,d}(1 + \lambda\|\hat{v}\|_{w,d})^6(M + |\Lambda|)^{\frac{3}{2}}, \quad (4.47)$$

where we also used the fact that $|\Lambda| \geq 1$. Together with (4.34), this finishes the proof. \square

Lemma 4.5 ($\mathcal{H}_{quart}(t)$ bound). *We have that*

$$\mathcal{H}_{quart}(t) = \sum_{k=1}^{16} A_{quart}^{(k)}(t) \quad (4.48)$$

with monomials $A_{quart}^{(k)}(t) \in \mathcal{P}[a, a^+]$ such that

$$\|A_{quart}^{(k)}(t)P_M\| \leq C\|\hat{v}\|_{w,d}(1 + \lambda\|\hat{v}\|_{w,d})^8 \frac{\lambda(M + |\Lambda|)^2}{N} \quad (4.49)$$

for any $M \in \mathbb{N}_0$, and $|\text{sign}(A_{quart}^{(k)}(t))| \leq 4$.

Proof. We follow the steps of the proof of Lemma 4.4. We have that

$$\begin{aligned} & \mathcal{H}_{quart}(t) \\ &= \frac{\lambda}{2N} \int d\mathbf{p}_4 \hat{v}(p_1 - p_3) \delta(p_1 + p_2 - p_3 - p_4) a_{p_1}^+(t) a_{p_2}^+(t) a_{p_3}(t) a_{p_4}(t) \\ &= \frac{\lambda}{2N} \sum_{\vec{\sigma}_4 \in \{\pm 1\}^4} \int d\mathbf{p}_4 \delta\left(\sum_{j=1}^4 \sigma_j p_j\right) \hat{v}(\sigma_1 p_1 + \sigma_3 p_3) \prod_{j=1}^4 (g_j(p_j, \sigma_j) a_{p_j}^{(\sigma_j)}), \end{aligned} \quad (4.50)$$

where $\vec{\sigma}_4 = (\sigma_1, \sigma_2, \sigma_3, \sigma_4)$, and

$$g_j(p_j, \sigma_j) := \begin{cases} u_\lambda(t, p_j)^{\delta_{\sigma_j, 1}} v_\lambda(t, p_j)^{\delta_{\sigma_j, -1}}, & j \in \{1, 2\}, \\ \overline{u_\lambda}(t, p_j)^{\delta_{\sigma_j, -1}} \overline{v_\lambda}(t, p_j)^{\delta_{\sigma_j, 1}}, & j \in \{3, 4\}. \end{cases} \quad (4.51)$$

Analogously to above, we denote

$$A_{quart}(\vec{\sigma}_4) := \int d\mathbf{p}_4 \delta\left(\sum_{j=1}^4 \sigma_j p_j\right) \hat{v}(\sigma_1 p_1 + \sigma_3 p_3) \prod_{j=1}^4 (g_j(p_j, \sigma_j) a_{p_j}^{(\sigma_j)}). \quad (4.52)$$

Wick-ordering using Lemma A.3 yields

$$\begin{aligned} & \delta\left(\sum_{j=1}^4 \sigma_j p_j\right) \prod_{j=1}^4 a_{p_j}^{(\sigma_j)} \\ &= \delta\left(\sum_{j=1}^4 \sigma_j p_j\right) : \prod_{j=1}^4 a_{p_j}^{(\sigma_j)} : + \sum_{\substack{j_1 < j_2, \\ j_3 < j_4}} \delta(p_{j_1} - p_{j_2}) \delta_{\sigma_{j_1}, -1} \delta_{\sigma_{j_2}, 1} \\ & \quad (\delta(\sigma_{j_3} p_{j_3} + \sigma_{j_4} p_{j_4}) : a_{p_{j_3}}^{(\sigma_{j_3})} a_{p_{j_4}}^{(\sigma_{j_4})} : + |\Lambda| \delta(p_{j_3} - p_{j_4}) \delta_{\sigma_{j_3}, -1} \delta_{\sigma_{j_4}, 1}), \end{aligned} \quad (4.53)$$

where $\{j_1, \dots, j_4\} = \{1, \dots, 4\}$.

We start with the 0-order terms. These occur whenever two pairs aa^+ exist, each with an annihilation operator to the left of a creation operator. This implies an integrable coefficient v_λ resp. $\overline{v_\lambda}$ for each of the corresponding momenta. Together with the deltas coming from the contraction, the resulting terms are integrable and are bounded by

$$\begin{aligned} & \|\Lambda\| \sum_{\substack{j_1 < j_2, \\ j_3 < j_4}} \int d\mathbf{p}_4 \hat{v}(\sigma_1 p_1 + \sigma_3 p_3) \prod_{j=1}^4 g_j(p_j, \sigma_j) \\ & \quad \delta(p_{j_1} - p_{j_2}) \delta(p_{j_3} - p_{j_4}) \delta_{\sigma_{j_1}, -1} \delta_{\sigma_{j_2}, 1} \delta_{\sigma_{j_3}, -1} \delta_{\sigma_{j_4}, 1} P_M \\ & \leq C \lambda^2 |\Lambda| \|\hat{v}\|_{w,d}^3 (1 + \lambda^2 \|\hat{v}\|_{w,d}^2)^2, \end{aligned} \quad (4.54)$$

where, as above $\{j_1, \dots, j_4\} = \{1, \dots, 4\}$.

For terms quadratic in a, a^+ , there again remain two independent momenta in the integration after integrating out the δ 's coming from momentum conservation and the single contraction. Observe that one of these δ 's comes from a contraction with an annihilation operator left to a creation operator. Thus, the corresponding momentum comes with an integrable factor $v_\lambda(t, p_{j_1})$ resp. $\overline{v_\lambda}(t, p_{j_1})$. The other momentum comes with a coefficient $v_\lambda(t, p_{j_3})^{\delta_{\sigma_{j_3}, -1}} : a_{p_{j_3}}^{(\sigma_{j_3})} a_{-\sigma_{j_3} p_{j_4}}^{(\sigma_{j_4})} :$, unless $j_3 = 3$. In that case, we have a coefficient $v(\sigma_1 p_1 + \sigma_3 p_3) : a_{p_3}^{(\sigma_3)} a_{-\sigma_3 p_4}^{(\sigma_4)} :$. Then, integrating first over p_{j_2}, p_{j_4} and then p_{j_3} , Lemma A.4 implies

$$\begin{aligned} & \left\| \sum_{\substack{j_1 < j_2, \\ j_3 < j_4}} \int d\mathbf{p}_4 \hat{v}(\sigma_1 p_1 + \sigma_3 p_3) \prod_{j=1}^4 g_j(p_j, \sigma_j) \right. \\ & \quad \left. \delta(p_{j_1} - p_{j_2}) \delta(\sigma_{j_3} p_{j_3} + \sigma_{j_4} p_{j_4}) \delta_{\sigma_{j_1}, -1} \delta_{\sigma_{j_2}, 1} : a_{p_{j_3}}^{(\sigma_{j_3})} a_{p_{j_4}}^{(\sigma_{j_4})} : P_M \right\| \\ & \leq C \|\hat{v}\|_{w,d} (1 + \lambda^2 \|\hat{v}\|_{w,d})^4 (M + |\Lambda|). \end{aligned} \quad (4.55)$$

Finally, if not all $\sigma_j = 1$ or $\sigma_j = -1$, Lemma A.4 implies

$$\begin{aligned} & \| : A_{quart}(\vec{\sigma}_4) : P_M \| \\ & \leq C \| |\hat{v}(\sigma_1 p_1 + \sigma_3 p_3)|^{\frac{1}{2}} \prod_{j=1}^2 |v_\lambda(t, p_j)|^{\delta_{\sigma_j, -1}} |u_\lambda(t, p_{j+2})|^{\delta_{\sigma_{j+2}, -1}} \delta(\sum_{j=1}^4 \sigma_j p_j)^{\frac{1}{2}} \|_{L_{p_{J_+} L_{p_{J_-}}}^\infty} \\ & \| |\hat{v}(\sigma_1 p_1 + \sigma_3 p_3)|^{\frac{1}{2}} \prod_{j=1}^2 |u_\lambda(t, p_j)|^{\delta_{\sigma_j, 1}} |v_\lambda(t, p_{j+2})|^{\delta_{\sigma_{j+2}, 1}} \delta(\sum_{j=1}^4 \sigma_j p_j)^{\frac{1}{2}} \|_{L_{p_{J_-} L_{p_{J_+}}}^\infty} \\ & (M+1)^2. \end{aligned} \quad (4.56)$$

By flipping the roles of (1, 2) and (3, 4), it is sufficient to show boundedness of only one of the norms. If $1 \in J_-$, we can use integrability of $|v_\lambda(t, p_1)|$, if $2 \in J_-$, there is a factor $|v_\lambda(t, p_2)|$. Integrability w.r.t. to p_3 and p_4 is always given due to the factor $|\hat{v}(\sigma_1 p_1 + \sigma_3 p_3)| \delta(\sum_{j=1}^4 \sigma_j p_j)$. In case $J_- = \emptyset$, Lemma A.4 implies

$$\begin{aligned} & \| : A_{quart}(1, 1, 1, 1) : P_M \| \\ & \leq (\| \hat{v}(p_1 + p_3) v_\lambda(t, p_1) v_\lambda(t, p_2) \overline{u_\lambda}(t, p_3) \overline{u_\lambda}(t, p_4) \|_{L_{p_2}^2 L_{p_3, p_4}^\infty}^2 (M-3) \\ & + \| \hat{v}(p_1 + p_3) v_\lambda(t, p_1) v_\lambda(t, p_2) \overline{u_\lambda}(t, p_3) \overline{u_\lambda}(t, p_1 + p_2 - p_3) \|_{L_{p_3}^2}^2 |\Lambda|)^{\frac{1}{2}} \\ & \sqrt{M(M-1)(M-2)}. \end{aligned} \quad (4.57)$$

$\| : A_{quart}(-1, -1, -1, -1) : P_M \|$ can be similarly bounded. Collecting (4.54), (4.55), (4.56) and (4.57), and using $|\Lambda| \geq 1$, we have shown that

$$\| A_{quart}(\vec{\sigma}_4) P_M \| \leq C \| \hat{v} \|_{w,d} (1 + \lambda \| \hat{v} \|_{w,d})^8 (M + |\Lambda|)^2. \quad (4.58)$$

Using (4.50), this concludes the proof. \square

Proposition 4.6. *Let $A \in \mathcal{P}[a, a^+]$ be a monomial in a, a^+ , $\gamma > 0$ and $\ell \in \mathbb{N}$ be such that*

$$\| AP_M \| \leq \gamma (M + |\Lambda|)^{\frac{\ell}{2}}, \quad (4.59)$$

and that $|\text{sign}(A)| \leq \ell$. Let $\mathbf{s}_{j+k} \in \Delta[t, j+k]$ and $t \geq 0$. For the moment, we abbreviate by $A \mathcal{H}_{cub}^j \mathcal{H}_{quart}^k(\mathbf{s}_{j+k})$ all terms that contain one factor A , j factors \mathcal{H}_{cub} , k factors \mathcal{H}_{quart} , all at possibly different times s_m . Then we have that

$$\begin{aligned} & |\nu(A \mathcal{H}_{cub}^j \mathcal{H}_{quart}^k(\mathbf{s}_{j+k}))| \\ & \leq \frac{\gamma (C \lambda \| \hat{v} \|_{w,d})^{j+k} (1 + \lambda \| \hat{v} \|_{w,d})^{6j+8k}}{N^{\frac{j}{2}+k}} \\ & \nu \left((\mathcal{N}_b + 3j + 4k + \ell + |\Lambda|)^{\frac{3j+4k+\ell}{2}} \right). \end{aligned} \quad (4.60)$$

for any state ν and all $\lambda > 0$ small enough.

Proof. Decomposing \mathcal{H}_{cub} and \mathcal{H}_{quart} into monomials as in Lemmata 4.4 and 4.5, we apply Lemmata 4.3, 4.4, 4.5 together with assumption (4.59) on A , we obtain that

$$\begin{aligned} & |\nu(A A_{cub}^j A_{quart}^k(\mathbf{s}_{j+k}))| \leq \frac{\gamma (C \lambda \| \hat{v} \|_{w,d})^{j+k} (1 + \lambda \| \hat{v} \|_{w,d})^{6j+8k}}{N^{\frac{j}{2}+k}} \\ & \nu \left((\mathcal{N}_b + 3j + 4k + \ell + |\Lambda|)^{\frac{3j+4k+\ell}{2}} \right). \end{aligned} \quad (4.61)$$

As a consequence, we obtain that

$$|\nu(A\mathcal{H}_{cub}^j \mathcal{H}_{quart}^k(\mathbf{s}_{j+k}))| \leq \frac{\gamma(C\lambda\|\hat{v}\|_{w,d})^{j+k}(1+\lambda\|\hat{v}\|_{w,d})^{6j+8k}}{N^{\frac{j}{2}+k}} \nu\left((\mathcal{N}_b + 3j + 4k + \ell + |\Lambda|)^{\frac{3j+4k+\ell}{2}}\right). \quad (4.62)$$

This finishes the proof. \square

Lemma 4.7. *Given a test function $J \in \ell^2(\Lambda^*) \cap \ell^\infty(\Lambda^*)$, we recall that*

$$f[J] = \int_{\Lambda^*} dp J(p) a_p^+ a_p, \quad (4.63)$$

$$g[J] = \int_{\Lambda^*} dp J(p) a_{-p} a_p. \quad (4.64)$$

Then we have

$$\|P_m f[J] P_n\| \leq \delta_{m,n} \|J\|_{\ell^\infty(\Lambda^*)} m, \quad (4.65)$$

$$\|P_m g[J] P_n\| \leq \delta_{n,m+2} \|J\|_{2 \cap \infty, d} (m+1 + |\Lambda|). \quad (4.66)$$

Proof. Due to $[\mathcal{N}, f[J]] = 0$, we have that

$$P_m f[J] P_n = \delta_{m,n} \int d\mathbf{p}_2 J(p_1) \delta(p_1 - p_2) a_{p_1}^+ a_{p_2} P_n. \quad (4.67)$$

Then Lemma A.4 in Appendix A implies

$$\|P_m f[J] P_n\| \leq \delta_{m,n} \|J\|_\infty m \quad (4.68)$$

For $g[J]$, we have that

$$P_m g[J] P_n = \delta_{n,m+2} \int d\mathbf{p}_2 J(p_1) \delta(p_1 + p_2) a_{p_1} a_{p_2} \quad (4.69)$$

Then Lemma A.4 implies

$$\begin{aligned} \|P_m g[J] P_n\| &\leq \delta_{n,m+2} \sqrt{\|J\|_\infty^2 (m+1) + |\Lambda| \|J\|_2^2} \sqrt{m+2} \\ &\leq \delta_{n,m+2} (\|J\|_{L^2(\Lambda^*)} + \|J\|_\infty) (m+1 + |\Lambda|) \end{aligned} \quad (4.70)$$

This finishes the proof. \square

Lemma 4.8 (Propagation of moments in HFB evolution). *The HFB evolution $\mathcal{V}_{HFB}(t)$ satisfies*

$$\|(\mathcal{N}_b + |\Lambda|)^{\frac{\ell}{2}} \mathcal{V}_{HFB}(t) (\mathcal{N}_b + |\Lambda|)^{-\frac{\ell}{2}}\| \leq e^{K_\ell \|\hat{v}\|_{w,d} \lambda t} \quad (4.71)$$

for all $\ell \in \mathbb{N}$ and some positive constants $K_\ell > 0$.

Proof. Let $\psi \in \mathcal{F}$. We have that

$$\begin{aligned} &i\partial_t \langle \mathcal{V}_{HFB}(t) \psi, (\mathcal{N}_b + |\Lambda|)^\ell \mathcal{V}_{HFB}(t) \psi \rangle \\ &= \langle \mathcal{V}_{HFB}(t) \psi, [(\mathcal{N}_b + |\Lambda|)^\ell, \mathcal{H}_{HFB}^{\phi_0}] \mathcal{V}_{HFB}(t) \psi \rangle \\ &= \sum_{n=0}^{\infty} (n + |\Lambda|)^\ell \langle \mathcal{V}_{HFB}(t) \psi, [P_n, \mathcal{H}_{cor}^{\phi_0}] \mathcal{V}_{HFB}(t) \psi \rangle. \end{aligned} \quad (4.72)$$

Employing (4.9) and recalling (1.34), we have that

$$[P_n, \mathcal{H}_{HFB}^{\phi_0}] = \lambda(P_n g[\hat{v}] P_{n+2} - P_{n-2} g[\hat{v}] P_n - h.c.). \quad (4.73)$$

As a consequence, we have that

$$\begin{aligned}
& \left| \partial_t \langle \mathcal{V}_{HFB}(t)\psi, (\mathcal{N}_b + |\Lambda|)^\ell \mathcal{V}_{HFB}(t)\psi \rangle \right| \\
&= 2\lambda \left| \sum_{n=0}^{\infty} (n + |\Lambda|)^\ell \operatorname{Im} \langle \mathcal{V}_{HFB}(t)\psi, (P_n g[\hat{v}] P_{n+2} - P_{n-2} g[\hat{v}] P_n) \mathcal{V}_{HFB}(t)\psi \rangle \right| \\
&= 2\lambda \left| \sum_{n=0}^{\infty} [(n + |\Lambda|)^\ell - (n + 2 + |\Lambda|)^\ell] \operatorname{Im} \langle \mathcal{V}_{HFB}(t)\psi, P_n g[\hat{v}] P_{n+2} \mathcal{V}_{HFB}(t)\psi \rangle \right|. \tag{4.74}
\end{aligned}$$

Using the Mean-Value Theorem and Cauchy-Schwarz, the last inequality implies

$$\begin{aligned}
& \left| \partial_t \langle \mathcal{V}_{HFB}(t)\psi, (\mathcal{N}_b + |\Lambda|)^\ell \mathcal{V}_{HFB}(t)\psi \rangle \right| \\
&\leq K_\ell \lambda \sum_{n=0}^{\infty} (n + 2 + |\Lambda|)^{\ell-1} \|P_n \mathcal{V}_{HFB}(t)\psi\| \|P_n g[\hat{v}] P_{n+2} \mathcal{V}_{HFB}(t)\psi\|. \tag{4.75}
\end{aligned}$$

Applying Lemma 4.7 followed by Young's inequality, we find

$$\begin{aligned}
& \left| \partial_t \langle \mathcal{V}_{HFB}(t)\psi, (\mathcal{N}_b + |\Lambda|)^\ell \mathcal{V}_{HFB}(t)\psi \rangle \right| \\
&\leq K_\ell \lambda \|\hat{v}\|_{w,d} \sum_{n=0}^{\infty} (n + 2 + |\Lambda|)^{\ell-1} (n + 1 + |\Lambda|) \\
&\quad \left(\langle \mathcal{V}_{HFB}(t)\psi, P_n \mathcal{V}_{HFB}(t)\psi \rangle + \langle \mathcal{V}_{HFB}(t)\psi, P_{n+2} \mathcal{V}_{HFB}(t)\psi \rangle \right) \\
&\leq K_\ell \lambda \|\hat{v}\|_{w,d} \langle \mathcal{V}_{HFB}(t)\psi, (\mathcal{N}_b + |\Lambda|)^\ell \mathcal{V}_{HFB}(t)\psi \rangle. \tag{4.76}
\end{aligned}$$

Employing Gronwall's Lemma concludes the proof. \square

Remark 4.9. *Using Corollary 3.3, we find that*

$$\begin{aligned}
\mathcal{V}_{HFB}^*(t) \mathcal{N}_b \mathcal{V}_{HFB}(t) &= \int dp (|u_\lambda(t,p)|^2 a_p^+ a_p + |v_\lambda(t,p)|^2 a_{-p} a_{-p}^+ \\
&\quad + u_\lambda(t,p) \overline{v_\lambda}(t,p) a_p^+ a_{-p}^+ + \overline{u_\lambda}(t,p) v_\lambda(t,p) a_p a_{-p}) \\
&= \int dp ((|u_\lambda(t,p)|^2 + |v_\lambda(t,p)|^2) a_p^+ a_p + |\Lambda| |v_\lambda(t,p)|^2 \\
&\quad + u_\lambda(t,p) \overline{v_\lambda}(t,p) a_p^+ a_{-p}^+ + \overline{u_\lambda}(t,p) v_\lambda(t,p) a_p a_{-p}), \tag{4.77}
\end{aligned}$$

where we used the CCR together with the fact that v_λ is even in p . Lemma 4.7 then implies

$$\left\| \mathcal{V}_{HFB}^*(t) (\mathcal{N}_b + |\Lambda|)^{\frac{1}{2}} \mathcal{V}_{HFB}(t) (\mathcal{N}_b + |\Lambda|)^{-\frac{1}{2}} \right\| \leq C_{\|\hat{v}\|_{w,d}}. \tag{4.78}$$

As an immediate consequence of Lemmata 4.8 and B.2, we obtain the following statement.

Corollary 4.10 (Propagation of moments). *Let $|\Lambda| \geq 1$. There exist constants $C_{\ell,\mu_0}, K_\ell > 0$ such that for any $\ell \in \mathbb{N}$, we have that*

$$\nu_t \left((\mathcal{N}_b + |\Lambda|)^{\frac{\ell}{2}} \right) \leq C_{\ell,\|\hat{v}\|_{w,d}} e^{K_\ell \|\hat{v}\|_{w,d} \lambda |\Lambda| t} |\Lambda|^{\frac{\ell}{2}}. \tag{4.79}$$

Proof. Assume that $\frac{\ell}{2} \in \mathbb{N}$. Define

$$\tilde{\mathcal{U}}_N(t) := \mathcal{W}^*[\sqrt{N|\Lambda}]\phi_0 e^{-i\mathcal{H}_N t} \mathcal{W}[\sqrt{N|\Lambda}]\phi_0, \tag{4.80}$$

and let

$$\mu_t(A) := \nu_0(\tilde{\mathcal{U}}_N^*(t) A \tilde{\mathcal{U}}_N(t)). \tag{4.81}$$

There exist $\lambda_k \geq 0$, $\sum_{k=1}^{\infty} \lambda_k = 1$, and $\Psi_k \in \mathcal{F}$ such that

$$\mu_t(A) = \sum_{k=1}^{\infty} \lambda_k \langle \Psi_k, A \Psi_k \rangle. \quad (4.82)$$

Then we have that

$$\begin{aligned} \nu_t((\mathcal{N}_b + 1)^{\frac{\ell}{2}}) &= \mu_t\left(\mathcal{V}_{HFB}^*(t)(\mathcal{N}_b + 1)^{\frac{\ell}{2}}\mathcal{V}_{HFB}(t)\right) \\ &= \sum_{k=1}^{\infty} \lambda_k \langle \Psi_k, \mathcal{V}_{HFB}^*(t)(\mathcal{N}_b + 1)^{\frac{\ell}{2}}\mathcal{V}_{HFB}(t)\Psi_k \rangle \\ &\leq e^{K_{\ell}\|\hat{v}\|_{w,d}\lambda t} \sum_{k=1}^{\infty} \lambda_k \langle \Psi_k, (\mathcal{N}_b + 1)^{\frac{\ell}{2}}\Psi_k \rangle \\ &= e^{K_{\ell}\|\hat{v}\|_{w,d}\lambda t} \mu_t((\mathcal{N}_b + 1)^{\frac{\ell}{2}}). \end{aligned} \quad (4.83)$$

using Lemma 4.8. Similarly, we write

$$\nu_0(A) = \sum_{k=1}^{\infty} \alpha_k \langle \Phi_k, A \Phi_k \rangle \quad (4.84)$$

for some $\alpha_k \geq 0$, $\sum_{k=1}^{\infty} \alpha_k = 1$, $\Phi_k \in \mathcal{F}$, and we obtain that

$$\begin{aligned} \mu_t((\mathcal{N}_b + 1)^{\frac{\ell}{2}}) &= \sum_{k=1}^{\infty} \alpha_k \langle \Phi_k, \tilde{\mathcal{U}}_N^*(t)(\mathcal{N}_b + 1)^{\frac{\ell}{2}}\tilde{\mathcal{U}}_N(t)\Phi_k \rangle \\ &\leq C_{\ell} e^{K_{\ell}\|\hat{v}\|_{w,d}\lambda|\Lambda|t} \sum_{k=1}^{\infty} \mu_k \langle \Phi_k, (\mathcal{N}_b + 1)^{\frac{\ell}{2}} \left(1 + \frac{\mathcal{N}_b}{N|\Lambda|}\right) \Phi_k \rangle \\ &= C_{\ell} e^{K_{\ell}\|\hat{v}\|_{w,d}\lambda|\Lambda|t} \nu_0\left((\mathcal{N}_b + 1)^{\frac{\ell}{2}} \left(1 + \frac{\mathcal{N}_b}{N|\Lambda|}\right)\right) \\ &\leq C_{\ell, \|f_0\|_d} e^{K_{\ell}\|\hat{v}\|_{w,d}\lambda|\Lambda|t} |\Lambda|^{\frac{\ell}{2}}, \end{aligned} \quad (4.85)$$

where we used Lemma B.2 followed by Lemma 4.1. Collecting (4.83) and (4.85) and using $|\Lambda| \geq 1$, we obtain that

$$\nu_t((\mathcal{N}_b + 1)^{\frac{\ell}{2}}) \leq C_{\ell, \|f_0\|_d} e^{K_{\ell}\|\hat{v}\|_{w,d}\lambda|\Lambda|t} |\Lambda|^{\frac{\ell}{2}}. \quad (4.86)$$

The half-integer case follows analogously to (4.18). This concludes the proof. \square

5. CONTROL OF ERROR TERMS IN THE EXPANSION

Again, we write $\int \equiv \int_{\Lambda^*}$ for brevity, to account for lattice sums over Λ^* in the sense of (1.51).

Proposition 5.1 (Tail estimates). *Let $T > 0$ and $t \leq T\lambda^{-2}$, $\lambda \in (0, 1)$, $|\Lambda| \geq 1$, and $J \in \ell^1(\Lambda^*) \cap \ell^\infty(\Lambda^*)$. Then the following holds true.*

(1)

$$\Phi_t = -i \int_0^t ds \frac{\nu_0([a_0, \mathcal{H}_{cub}(s)])}{|\Lambda|} + \text{Rem}_2(t; \Phi) \quad (5.1)$$

with

$$|\text{Rem}_2(t; \Phi)| \leq C_{\|\hat{v}\|_{w,d}, \|f_0\|_d} T^2 e^{C\|\hat{v}\|_{w,d}|\Lambda|/\lambda T} \frac{|\Lambda|^{\frac{3}{2}}}{N\lambda^2} \left(1 + \frac{|\Lambda|}{N}\right). \quad (5.2)$$

$$\begin{aligned}
(2) \quad \int dp J(p)(f_t(p) - f_0(p)) &= -i \int_0^t ds \frac{\nu_0([f[J], \mathcal{H}_{quart}(s)])}{|\Lambda|} \\
&\quad - \int_{\Delta[t,2]} ds_2 \frac{\nu_0([[f[J], \mathcal{H}_{cub}(s_1)], \mathcal{H}_{cub}(s_2)])}{|\Lambda|} \\
&\quad + \text{Rem}_2(t; f[J])
\end{aligned} \tag{5.3}$$

with

$$\begin{aligned}
&|\text{Rem}_2(t; f[J])| \\
&\leq C_{\|\hat{v}\|_{w,d}, \|f_0\|_d} \langle T \rangle^4 \|J\|_{\ell^\infty(\Lambda^*)} \frac{e^{C\|\hat{v}\|_{w,d}|\Lambda|/\lambda T} |\Lambda|^6}{\lambda^4 N^2} \left(1 + \frac{|\Lambda|}{N}\right)^2.
\end{aligned} \tag{5.4}$$

$$\begin{aligned}
(3) \quad \int dp J(p)g_t(p) &= -i \int_0^t ds \frac{\nu_0([g[J], \mathcal{H}_{quart}(s)])}{|\Lambda|} \\
&\quad - \int_{\Delta[t,2]} ds_2 \frac{\nu_0([[g[J], \mathcal{H}_{cub}(s_1)], \mathcal{H}_{cub}(s_2)])}{|\Lambda|} \\
&\quad + \text{Rem}_2(t; g[J])
\end{aligned} \tag{5.5}$$

with

$$\begin{aligned}
&|\text{Rem}_2(t; g[J])| \\
&\leq C_{\|\hat{v}\|_{w,d}, \|f_0\|_d} \langle T \rangle^4 \|J\|_{2 \cap \infty, d} \frac{e^{C\|\hat{v}\|_{w,d}|\Lambda|/\lambda T} |\Lambda|^6}{\lambda^4 N^2} \left(1 + \frac{|\Lambda|}{N}\right)^2.
\end{aligned} \tag{5.6}$$

Proof. Recall that ν_0 being number conserving implies

$$\nu_0(a_0) = \nu_0(a_p a_q) = 0. \tag{5.7}$$

Let $A \in \mathcal{P}[a, a^+]$, be a monomial with data such that

$$\|P_n A P_{n-\text{sign}(A)}\| \leq \gamma(n + |\Lambda|)^{\frac{\ell}{2}} \tag{5.8}$$

with $\ell, |\text{sign}(A)| \leq 2$. Lemma 3.1 implies

$$\begin{aligned}
\nu_t(A) &= \nu_0(A) - i \int_0^t ds \nu_0([A, \mathcal{H}_I(s)]) \\
&\quad - \int_{\Delta[t,2]} ds_2 \nu_{s_2}([[A, \mathcal{H}_I(s_1)], \mathcal{H}_I(s_2)]) \\
&= \nu_0(A) - i \int_0^t ds \nu_0([A, \mathcal{H}_I(s)]) \\
&\quad - \int_{\Delta[t,2]} ds_2 \nu_0([[A, \mathcal{H}_I(s_1)], \mathcal{H}_I(s_2)]) \\
&\quad + i \int_{\Delta[t,3]} ds_3 \nu_0([[[A, \mathcal{H}_I(s_1)], \mathcal{H}_I(s_2)], \mathcal{H}_I(s_3)]) \\
&\quad + \int_{\Delta[t,4]} ds_4 \nu_{s_4}([[[[A, \mathcal{H}_I(s_1)], \mathcal{H}_I(s_2)], \mathcal{H}_I(s_3)], \mathcal{H}_I(s_4)]),
\end{aligned} \tag{5.10}$$

where

$$\mathcal{H}_I(s) = \mathcal{H}_{cub}(s) + \mathcal{H}_{quart}(s), \tag{5.11}$$

see (3.6). In order to simplify notation, we shall abbreviate by $A\mathcal{H}_{cub}^j\mathcal{H}_{quart}^k(\mathbf{s}_{j+k})$ all terms that contain one factor A , j factors $\mathcal{H}_{cub}(s_\ell)$, k factors $\mathcal{H}_{quart}(s_m)$, all at possibly different times. Let $\mathbf{s}_{j+k} \in \Delta[t, j+k]$, $t \leq T\lambda^{-2}$, $\lambda \in (0, 1)$. Proposition 4.6 then implies

$$\begin{aligned} & |\nu_{\mathbf{s}_{j+k}}(A\mathcal{H}_{cub}^j\mathcal{H}_{quart}^k(\mathbf{s}_{j+k}))| \\ & \leq \frac{C_{j,k}\gamma\|\hat{v}\|_{w,d}^{j+k}\lambda^{j+k}(1+\lambda\|\hat{v}\|_{w,d})^{6j+8k}}{N^{\frac{j}{2}+k}}\nu_{\mathbf{s}_{j+k}}\left((\mathcal{N}_b+|\Lambda|)^{\frac{3j+4k+\ell}{2}}\right) \end{aligned} \quad (5.12)$$

for all $N > 0$ large enough. By Corollary 4.10, we obtain

$$\begin{aligned} & |\nu_{\mathbf{s}_{j+k}}(A\mathcal{H}_{cub}^j\mathcal{H}_{quart}^k(\mathbf{s}_{j+k}))| \\ & \leq C_{\|\hat{v}\|_{w,d},\|f_0\|_{d,j,k}}\gamma e^{K_{j,k}\|\hat{v}\|_{w,d}|\Lambda|/\lambda T}\frac{\lambda^{j+k}|\Lambda|^{\frac{3j+4k+\ell}{2}}}{N^{\frac{j}{2}+k}}. \end{aligned} \quad (5.13)$$

Similarly to (5.12), we have that Lemma 4.1 implies

$$\begin{aligned} & |\nu_0(A\mathcal{H}_{cub}^j\mathcal{H}_{quart}^k(\mathbf{s}_{j+k}))| \\ & \leq C_{\|\hat{v}\|_{w,d},\|f_0\|_{d,j,k}}\gamma\frac{\lambda^{j+k}|\Lambda|^{\frac{3j+4k+\ell}{2}}}{N^{\frac{j}{2}+k}}. \end{aligned} \quad (5.14)$$

Due to ν_0 being number conserving, see (4.10), we have that

$$|\nu_0(A\mathcal{H}_{cub}^j\mathcal{H}_{quart}^k(\mathbf{s}_{j+k}))| = 0, \quad (5.15)$$

whenever the total number of creation and annihilation operators in $A\mathcal{H}_{cub}^j\mathcal{H}_{quart}^k(\mathbf{s}_{j+k})$ is odd.

In the case $A = a_0$, due to Corollary 3.3 and using $\hat{v}(0) = 0$, we have that

$$a_0 = a_0(s) \quad (5.16)$$

for all $s \in \mathbb{R}$. With that, an easy computation yields

$$\begin{aligned} [a_0, \mathcal{H}_{cub}(s)] &= [a_0, \mathcal{H}_{cub}^{\phi_0}(s)] \\ &= \frac{\lambda}{\sqrt{N}} \int dp \hat{v}(p) (a_p^+(s)a_p(s) + a_p(s)a_{-p}(s)). \end{aligned} \quad (5.17)$$

Using Corollary 3.3, we obtain

$$[a_0, \mathcal{H}_{cub}(s)] = f[J_1(s)] + g[J_2(s)] + g^*[J_3(s)] + J_4(s), \quad (5.18)$$

where, with the notation of Corollary 3.3,

$$J_1(s, p) := \frac{\lambda\hat{v}(p)}{\sqrt{N}} (|u_\lambda(s, p)|^2 + |v_\lambda(s, p)|^2 + 2\overline{v_\lambda}(s, p)\overline{u_\lambda}(s, p)), \quad (5.19)$$

$$J_2(s, p) := \frac{\lambda\hat{v}(p)}{\sqrt{N}} (v_\lambda(s, p) + \overline{u_\lambda}(s, p))\overline{u_\lambda}(s, p), \quad (5.20)$$

$$J_3(s, p) := \frac{\lambda\hat{v}(p)}{\sqrt{N}} (u_\lambda(s, p) + \overline{v_\lambda}(s, p))\overline{v_\lambda}(s, p), \quad (5.21)$$

$$J_4(s) := \frac{\lambda}{\sqrt{N}} \int dp \hat{v}(p) (|v_\lambda(s, p)|^2 + \overline{u_\lambda}(s, p)\overline{v_\lambda}(s, p)). \quad (5.22)$$

Note, that in (5.18), we used the notation

$$g^*[J] = \int dp J(p)a_p^+a_{-p}^-. \quad (5.23)$$

Using Corollary 3.3 again, we find that

$$\|J_i(s)\|_d \leq C_{\|\hat{v}\|_{w,d}} \frac{\lambda}{\sqrt{N}} \quad \text{for } i \in \{1, 2, 3\}, \quad (5.24)$$

$$|J_4(s)| \leq C_{\|\hat{v}\|_{w,d}} \frac{\lambda}{\sqrt{N}} \quad (5.25)$$

for all $s \leq T\lambda^{-2}$.

Similarly, we compute

$$\begin{aligned} [a_0, \mathcal{H}_{quart}(s)] &= [a_0, \mathcal{H}_{quart}](s) \\ &= \frac{\lambda}{N} \int dp \hat{v}(p_2) (a_{p_1+p_2}^+ a_{p_2} a_{p_1})(s) \\ &= \frac{1}{\sqrt{N}} \mathcal{H}_{cub}^{(2)}(s). \end{aligned} \quad (5.26)$$

Here, we used the notation

$$\begin{aligned} \mathcal{H}_{cub}(s) &= \frac{\lambda}{\sqrt{N}} \int dp \hat{v}(p_2) ((a_{p_1}^+ a_{p_2}^+ a_{p_1+p_2})(s) + (a_{p_1+p_2}^+ a_{p_2} a_{p_1})(s)) \\ &=: \mathcal{H}_{cub}^{(1)}(s) + \mathcal{H}_{cub}^{(2)}(s). \end{aligned} \quad (5.27)$$

Notice that bounds of the form (5.12)–(5.14) hold with \mathcal{H}_{cub} replaced by $\mathcal{H}_{cub}^{(2)}$, due to the proof of Proposition 4.6.

Using (5.9) and (5.26), we have that

$$\Phi_t = -\frac{i}{|\Lambda|} \int_0^t ds \nu_0([a_0, \mathcal{H}_{cub}(s)]) + \text{Rem}_2(t; \Phi), \quad (5.28)$$

where

$$\begin{aligned} \text{Rem}_2(t; \Phi) &:= -\frac{i}{2\sqrt{N}} \int_0^t ds \frac{\nu_0(\mathcal{H}_{cub}^{(2)}(s))}{|\Lambda|} \\ &\quad - \int_{\Delta[t,2]} ds_2 \frac{\nu_{s_2}([f[J_1(s_1)] + g[J_2(s_1)] + g^*[J_3(s_1)], \mathcal{H}_I(s_2)])}{|\Lambda|} \\ &\quad + \frac{1}{2\sqrt{N}} \int_{\Delta[t,2]} ds_2 \frac{\nu_{s_2}([\mathcal{H}_{cub}^{(2)}(s_1), \mathcal{H}_I(s_2)])}{|\Lambda|}. \end{aligned} \quad (5.29)$$

Recall that, by definition (5.23),

$$|\nu_{s_2}([g^*[J_3(s_1)], \mathcal{H}_I(s_2)])| = |\nu_{s_2}([g[\overline{J_3}(s_1)], \mathcal{H}_I(s_2)])|. \quad (5.30)$$

Then Lemma 4.7 together with the bounds (5.13) and (5.24) imply

$$\begin{aligned} &\frac{|\nu_{s_2}([f[J_1(s_1)], \mathcal{H}_I(s_2)])|}{|\Lambda|}, \frac{|\nu_{s_2}([g[J_2(s_1)], \mathcal{H}_I(s_2)])|}{|\Lambda|}, \\ &\frac{|\nu_{s_2}([g^*[J_3(s_1)], \mathcal{H}_I(s_2)])|}{|\Lambda|} \\ &\leq \frac{C_{\|\hat{v}\|_{w,d}, \|f_0\|_d} e^{C\|\hat{v}\|_{w,d}|\Lambda|/\lambda T} \lambda^2 |\Lambda|^{\frac{3}{2}}}{N} \left(1 + \sqrt{\frac{|\Lambda|}{N}}\right). \end{aligned} \quad (5.31)$$

(5.15) with $A = \mathbf{1}$ implies

$$\frac{|\nu_0(\mathcal{H}_{cub}^{(2)}(s))|}{|\Lambda|\sqrt{N}} = 0. \quad (5.32)$$

(5.13) with $A = \mathbf{1}$ implies

$$\begin{aligned} & \frac{|\nu_{s_2}([\mathcal{H}_{cub}^{(2)}(s_1), \mathcal{H}_I(s_2)])|}{|\Lambda|\sqrt{N}} \\ & \leq \frac{C_{\|\hat{v}\|_{w,d}, \|\hat{v}\|_{w,d}} e^{C\|\hat{v}\|_{w,d}|\Lambda|/\lambda T} \lambda^2 |\Lambda|^2}{N^{\frac{3}{2}}} \left(1 + \sqrt{\frac{|\Lambda|}{N}}\right). \end{aligned} \quad (5.33)$$

Collecting (5.29), (5.31), (5.32), and (5.33), yields the bound

$$\begin{aligned} & |\text{Rem}_2(t; \Phi)| \\ & \leq C_{\|\hat{v}\|_{w,d}, \|f_0\|_d} e^{C\|\hat{v}\|_{w,d}|\Lambda|/\lambda T} \frac{\lambda^2 t^2 |\Lambda|^{\frac{3}{2}}}{N} \left(1 + \sqrt{\frac{|\Lambda|}{N}}\right)^2 \\ & \leq C_{\|\hat{v}\|_{w,d}, \|f_0\|_d} T^2 e^{C\|\hat{v}\|_{w,d}|\Lambda|/\lambda T} \frac{|\Lambda|^{\frac{3}{2}}}{N\lambda^2} \left(1 + \frac{|\Lambda|}{N}\right) \end{aligned} \quad (5.34)$$

for all $t \leq T\lambda^{-2}$, $\lambda \in (0, 1)$.

In the cases $A = f[J]$ and $A = g[J]$, we have that

$$\begin{aligned} \frac{\nu_t(A)}{|\Lambda|} &= \frac{\nu_0(f[J])}{|\Lambda|} \mathbb{1}_{A=f[J]} - i \int_0^t ds \frac{\nu_0([A, \mathcal{H}_{quart}(s)])}{|\Lambda|} \\ & \quad - \int_{\Delta[t,2]} ds_2 \frac{\nu_0([[A, \mathcal{H}_{cub}(s_1)], \mathcal{H}_{cub}(s_2)])}{|\Lambda|} \\ & \quad + \text{Rem}_2(t; A), \end{aligned} \quad (5.35)$$

where, using (5.7) and (5.10),

$$\begin{aligned} \text{Rem}_2(t; A) &:= -i \int_0^t ds \frac{\nu_0([A, \mathcal{H}_{cub}(s)])}{|\Lambda|} \\ & \quad - \int_{\Delta[t,2]} ds_2 \frac{\nu_0([[A, \mathcal{H}_{cub}(s_1)], \mathcal{H}_{quart}(s_2)])}{|\Lambda|} \\ & \quad - \int_{\Delta[t,2]} ds_2 \frac{\nu_0([[A, \mathcal{H}_{quart}(s_1)], \mathcal{H}_I(s_2)])}{|\Lambda|} \\ & \quad + i \int_{\Delta[t,3]} ds_3 \frac{\nu_0([[[A, \mathcal{H}_I(s_1)], \mathcal{H}_I(s_2)], \mathcal{H}_I(s_3)])}{|\Lambda|} \\ & \quad + \int_{\Delta[t,4]} ds_4 \frac{\nu_{s_4}([[[[A, \mathcal{H}_I(s_1)], \mathcal{H}_I(s_2)], \mathcal{H}_I(s_3)], \mathcal{H}_I(s_4)])}{|\Lambda|} \end{aligned} \quad (5.36)$$

Employing (5.14), Lemma 4.7 yields

$$\frac{|\nu_0(A\mathcal{H}_{quart}^2(\mathbf{s}_2))|}{|\Lambda|} \leq C_{\|\hat{v}\|_{w,d}, \|f_0\|_d} \frac{\|J\|_A \lambda^2 |\Lambda|^4}{N^2}, \quad (5.37)$$

$$\frac{|\nu_0(A\mathcal{H}_{cub}^2 \mathcal{H}_{quart}(\mathbf{s}_3))|}{|\Lambda|} \leq C_{\|\hat{v}\|_{w,d}, \|f_0\|_d} \frac{\|J\|_A \lambda^3 |\Lambda|^5}{N^2}, \quad (5.38)$$

$$\frac{|\nu_0(A\mathcal{H}_{quart}^3(\mathbf{s}_3))|}{|\Lambda|} \leq C_{\|\hat{v}\|_{w,d},\|f_0\|_d} \frac{\|J\|_A \lambda^3 |\Lambda|^6}{N^3}, \quad (5.39)$$

where we abbreviated

$$\|J\|_A := \|J\|_{\ell^\infty(\Lambda^*)} \mathbf{1}_{A=f[J]} + \|J\|_{2 \cap \infty, d} \mathbf{1}_{A=g[J]}. \quad (5.40)$$

Using (5.15), we have that

$$\begin{aligned} \frac{|\nu_0([A, \mathcal{H}_{cub}(s)])|}{|\Lambda|} &= \frac{|\nu_0(A\mathcal{H}_{cub}\mathcal{H}_{quart}(\mathbf{s}_2))|}{|\Lambda|} \\ &= \frac{|\nu_0(A\mathcal{H}_{cub}^3(\mathbf{s}_3))|}{|\Lambda|} = \frac{|\nu_0(A\mathcal{H}_{cub}\mathcal{H}_{quart}^2(\mathbf{s}_3))|}{|\Lambda|} = 0. \end{aligned} \quad (5.41)$$

(5.12) together with Lemma 4.7 implies

$$\begin{aligned} \frac{|\nu_{s_2}(A\mathcal{H}_T^4(\mathbf{s}_4))|}{|\Lambda|} \\ \leq C_{\|\hat{v}\|_{w,d},\|f_0\|_d} e^{C\|\hat{v}\|_{w,d}|\Lambda|/\lambda T} \frac{\|J\|_A \lambda^4 |\Lambda|^6}{N^2} \left(1 + \sqrt{\frac{|\Lambda|}{N}}\right)^4. \end{aligned} \quad (5.42)$$

Collecting (5.37)-(5.42), we arrive at

$$\begin{aligned} |\text{Rem}_2(t; A)| \\ \leq \frac{C_{\|\hat{v}\|_{w,d},\|f_0\|_d} e^{C\|\hat{v}\|_{w,d}|\Lambda|/\lambda T} \|J\|_A \lambda^2 t^2 |\Lambda|^4}{N^2} \left[1 + \lambda t |\Lambda| \left(1 + \frac{|\Lambda|}{N}\right)\right. \\ \left. + \lambda^2 t^2 |\Lambda|^2 \left(1 + \sqrt{\frac{|\Lambda|}{N}}\right)^4\right] \\ \leq C_{\|\hat{v}\|_{w,d},\|f_0\|_d} \langle T \rangle^4 \|J\|_A \frac{e^{C\|\hat{v}\|_{w,d}|\Lambda|/\lambda T} |\Lambda|^6}{\lambda^4 N^2} \left(\left(1 + \frac{|\Lambda|}{N}\right)^2 + \frac{\lambda^2}{|\Lambda|^2}\right) \\ \leq C_{\|\hat{v}\|_{w,d},\|f_0\|_d} \langle T \rangle^4 \|J\|_A \frac{e^{C\|\hat{v}\|_{w,d}|\Lambda|/\lambda T} |\Lambda|^6}{\lambda^4 N^2} \left(1 + \frac{|\Lambda|}{N}\right)^2 \end{aligned} \quad (5.43)$$

for all $t \leq T\lambda^{-2}$, $\lambda \in (0, 1)$, $|\Lambda| \geq 1$. This finishes the proof. \square

Up until now, all calculations did not further distinguish the cases of fixed $|\Lambda|$ and $|\Lambda|$ growing with N . In the latter situation, we will approximate lattice sums over Λ^* with integrals over \mathbb{R}^3 . It is crucial to note that oscillatory and dispersive properties differ fundamentally in these two cases.

5.1. Fixed, N -independent lattice $\Lambda^* \cong \mathbb{Z}^3$.

5.1.1. *Notation*. For the next result, we define

$$\Delta_{cub}H(\mathbf{p}_2) := H(p_1) + H(p_2) - H(p_1 + p_2), \quad (5.44)$$

$$\tilde{h} := 1 + h, \quad (5.45)$$

$$p_{12} := p_1 + p_2. \quad (5.46)$$

Recall from Corollary 3.3 that

$$V_1(p)^2 = \frac{\hat{v}(p)^2}{\Omega(p)(\Omega(p) + E(p) + \lambda\hat{v}(p))}, \quad (5.47)$$

$$V_2(p) = \frac{\hat{v}(p)}{\Omega(p)}. \quad (5.48)$$

In order to describe the dynamics of Φ , we introduce the condensate operator

$$\text{Con}_d(h)(T; \lambda) := -i \int_0^T dS \int_{\Lambda^*} dp \hat{v}(p) h_{\frac{S}{\lambda^2}}(p). \quad (5.49)$$

For the dynamics of f , we define the generalized Boltzmann operators

$$\begin{aligned} & \text{Bol}_d(h)[J](T; \lambda) \\ & := \frac{1}{\lambda^2} \int_{\Delta[T,2]} d\mathbf{S}_2 \int_{(\Lambda^*)^2} d\mathbf{p}_2 \cos\left(\frac{\Delta_{cub}\Omega(\mathbf{p}_2)(S_1 - S_2)}{\lambda^2}\right) \\ & \quad (\hat{v}(p_1) + \hat{v}(p_2))^2 \Delta_{cub} J(\mathbf{p}_2) \left(\tilde{h}_{\frac{S_2}{\lambda^2}}(p_1) \tilde{h}_{\frac{S_2}{\lambda^2}}(p_2) h_{\frac{S_2}{\lambda^2}}(p_1 + p_2) \right. \\ & \quad \left. - h_{\frac{S_2}{\lambda^2}}(p_1) h_{\frac{S_2}{\lambda^2}}(p_2) \tilde{h}_{\frac{S_2}{\lambda^2}}(p_1 + p_2) \right), \end{aligned} \quad (5.50)$$

$$\begin{aligned} & \text{bol}^{(1)}(h)[J](\mathbf{S}_2/\lambda^2) \\ & := \frac{2}{\lambda^2} \text{Im} \int_{(\Lambda^*)^2} d\mathbf{p}_2 (\hat{v}(p_1) + \hat{v}(p_2)) (\hat{v}(p_1) + \hat{v}(p_{12})) \\ & \quad V_2(p_1) e^{-i(\Omega(p_2) - \Omega(p_{12}))(S_1 - S_2)/\lambda^2} \\ & \quad \left((-J(-p_1) + J(p_2) - J(p_{12})) e^{i\Omega(p_1)S_2/\lambda^2} \sin(\Omega(p_1)S_1/\lambda^2) \right. \\ & \quad \left(\tilde{h}_{\frac{S_2}{\lambda^2}}(-p_1) h_{\frac{S_2}{\lambda^2}}(p_2) \tilde{h}_{\frac{S_2}{\lambda^2}}(p_{12}) - h_{\frac{S_2}{\lambda^2}}(-p_1) \tilde{h}_{\frac{S_2}{\lambda^2}}(p_2) h_{\frac{S_2}{\lambda^2}}(p_{12}) \right) \\ & \quad \left. + \Delta_{cub} J(\mathbf{p}_2) e^{i\Omega(p_1)S_1/\lambda^2} \sin(\Omega(p_1)S_2/\lambda^2) \right. \\ & \quad \left. \left(h_{\frac{S_2}{\lambda^2}}(p_1) h_{\frac{S_2}{\lambda^2}}(p_2) \tilde{h}_{\frac{S_2}{\lambda^2}}(p_{12}) - \tilde{h}_{\frac{S_2}{\lambda^2}}(p_1) \tilde{h}_{\frac{S_2}{\lambda^2}}(p_2) h_{\frac{S_2}{\lambda^2}}(p_{12}) \right) \right), \end{aligned} \quad (5.51)$$

$$\begin{aligned} & \text{Bol}_d^{(1)}(h)[J](T; \lambda) \\ & := \int_{\Delta[T,2]} d\mathbf{S}_2 \text{bol}^{(1)}(h)[J](\mathbf{S}_2/\lambda^2), \end{aligned} \quad (5.52)$$

$$\begin{aligned} & \text{bol}^{(2,1)}(h)[J](\mathbf{S}_2; \lambda) \\ & := \frac{1}{\lambda^2} \text{Im} \int_{(\Lambda^*)^2} d\mathbf{p}_2 (\hat{v}(p_1) + \hat{v}(p_2))^2 \Delta_{cub} J(\mathbf{p}_2) e^{i\Omega(\mathbf{p}_2)(S_1 - S_2)/\lambda^2} \\ & \quad \left(V_1(p_1)^2 (\sin(\Omega(p_1)S_1/\lambda^2) e^{-i\Omega(p_1)S_1/\lambda^2} - \sin(\Omega(p_1)S_2/\lambda^2) e^{i\Omega(p_1)S_2/\lambda^2}) \right. \\ & \quad + V_1(p_2)^2 (\sin(\Omega(p_2)S_1/\lambda^2) e^{-i\Omega(p_2)S_1/\lambda^2} - \sin(\Omega(p_2)S_2/\lambda^2) e^{i\Omega(p_2)S_2/\lambda^2}) \\ & \quad \left. - V_1(p_{12})^2 (\sin(\Omega(p_{12})S_1/\lambda^2) e^{i\Omega(p_{12})S_1/\lambda^2} - \sin(\Omega(p_{12})S_2/\lambda^2) e^{i\Omega(p_{12})S_2/\lambda^2}) \right) \\ & \quad \left(h_{\frac{S_2}{\lambda^2}}(p_1) h_{\frac{S_2}{\lambda^2}}(p_2) \tilde{h}_{\frac{S_2}{\lambda^2}}(p_{12}) - \tilde{h}_{\frac{S_2}{\lambda^2}}(p_1) \tilde{h}_{\frac{S_2}{\lambda^2}}(p_2) h_{\frac{S_2}{\lambda^2}}(p_{12}) \right), \end{aligned} \quad (5.53)$$

$$\begin{aligned} & \text{bol}^{(2,2)}(h)[J](\mathbf{S}_2/\lambda^2) \\ & := \frac{2}{\lambda^2} \text{Re} \int_{(\Lambda^*)^2} d\mathbf{p}_2 (\hat{v}(p_1) + \hat{v}(p_2)) \\ & \quad \left[\sin(\Omega(p_1)S_1/\lambda^2) \sin(\Omega(p_{12})S_1/\lambda^2) \right. \\ & \quad \left. e^{i\Omega(p_2)(S_1 - S_2)/\lambda^2 + i(\Omega(p_1) - \Omega(p_{12}))S_2/\lambda^2} \right. \\ & \quad \left. V_2(p_1) V_2(p_{12}) (\hat{v}(p_2) + \hat{v}(p_{12})) \right] \end{aligned}$$

$$\begin{aligned}
& (-J(-p_1) + J(p_2) + J(-p_{12})) \\
& \left(h \frac{s_2}{\lambda^2} (-p_1) \tilde{h} \frac{s_2}{\lambda^2} (p_2) \tilde{h} \frac{s_2}{\lambda^2} (-p_{12}) - \tilde{h} \frac{s_2}{\lambda^2} (-p_1) h \frac{s_2}{\lambda^2} (p_2) f(-p_{12}) \right) \\
& + \sin(\Omega(p_1) S_1 / \lambda^2) e^{i\Omega(p_2)(S_1 - S_2) / \lambda^2 - i\Omega(p_{12}) S_1 / \lambda^2} V_2(p_1) \\
& (-J(-p_1) + J(p_2) - J(p_{12})) \\
& \left((\hat{v}(p_1) + \hat{v}(p_2)) V_2(p_1) \sin(\Omega(p_1) S_2 / \lambda^2) e^{i\Omega(p_{12}) S_2 / \lambda^2} \right. \\
& \left. + (\hat{v}(p_2) + \hat{v}(p_{12})) V_2(p_{12}) \sin(\Omega(p_{12}) S_2 / \lambda^2) e^{-i\Omega(p_1) S_2 / \lambda^2} \right) \\
& \left(h \frac{s_2}{\lambda^2} (-p_1) \tilde{h} \frac{s_2}{\lambda^2} (p_2) h \frac{s_2}{\lambda^2} (p_{12}) - \tilde{h} \frac{s_2}{\lambda^2} (-p_1) h \frac{s_2}{\lambda^2} (p_2) \tilde{h} \frac{s_2}{\lambda^2} (p_{12}) \right) \\
& + \left(\frac{1}{2} e^{-i(\Omega(p_1) + \Omega(p_2)) S_2 / \lambda^2} \sin(\Omega(p_{12}) S_2 / \lambda^2) V_2(p_{12}) (\hat{v}(p_1) + \hat{v}(p_2)) \right. \\
& \left. + e^{-i(\Omega(p_1) + \Omega(p_{12})) S_2} \sin(\Omega(p_2) S_2 / \lambda^2) V_2(p_2) (\hat{v}(p_1) + \hat{v}(p_{12})) \right) \\
& \left. \left(\tilde{h} \frac{s_2}{\lambda^2} (p_1) \tilde{h} \frac{s_2}{\lambda^2} (p_2) \tilde{h} \frac{s_2}{\lambda^2} (-p_{12}) - h \frac{s_2}{\lambda^2} (p_1) h \frac{s_2}{\lambda^2} (p_2) h \frac{s_2}{\lambda^2} (-p_{12}) \right) \right], \tag{5.54}
\end{aligned}$$

$$\begin{aligned}
& \text{bol}^{(2)}(h)[J](\mathbf{S}_2 / \lambda^2) \\
& := \text{bol}^{(2,1)}(h)[J](\mathbf{S}_2; \lambda) + \text{bol}^{(2,2)}(h)[J](\mathbf{S}_2 / \lambda^2), \tag{5.55}
\end{aligned}$$

$$\begin{aligned}
& \text{Bol}_d^{(2)}(h)[J](T; \lambda) \\
& := \int_{\Delta[T, 2]} d\mathbf{S}_2 \text{bol}^{(2)}(h)[J](\mathbf{S}_2 / \lambda^2). \tag{5.56}
\end{aligned}$$

5.1.2. Results.

Proposition 5.2. *Let $T > 0$, $\lambda \in (0, 1)$, $|\Lambda| \geq 1$, and $J \in L^\infty(\mathbb{R}^3; \mathbb{R})$. Then the following holds.*

$$\begin{aligned}
(1) \quad & -i \int_0^{T\lambda^{-2}} ds \frac{\nu_0([a_0, \mathcal{H}_{cub}(s)])}{|\Lambda|} = \frac{1}{N^{\frac{1}{2}} \lambda} \text{Con}_d(f_0)(T; \lambda) + \text{err}_{1,d}^{(Bog)}\left(\frac{T}{\lambda^2}; \Phi\right) \tag{5.57}
\end{aligned}$$

where

$$|\text{err}_{1,d}^{(Bog)}\left(\frac{T}{\lambda^2}; \Phi\right)| \leq \frac{C \|\hat{v}\|_{w,d} \|f_0\|_d T}{N^{\frac{1}{2}}}, \tag{5.58}$$

$$\begin{aligned}
(2) \quad & \nu_0([f[J], \mathcal{H}_{quart}(s)]) = 0, \tag{5.59}
\end{aligned}$$

$$\begin{aligned}
(3) \quad & - \int_{\Delta[T\lambda^{-2}, 2]} d\mathbf{S}_2 \frac{\nu_0([f[J], \mathcal{H}_{cub}(s_1)], \mathcal{H}_{cub}(s_2)])}{|\Lambda|} \\
& = \frac{1}{N} (\text{Bol}_d(f_0)[J](T; \lambda) + \lambda \text{Bol}_d^{(1)}(f_0)[J](T; \lambda) + \lambda^2 \text{Bol}_d^{(2)}(f_0)[J](T; \lambda)) \\
& + J(0) |\Phi \frac{T}{\lambda^2}|^2 + \text{err}_{2,d}^{(Bog, Bol)}\left(\frac{T}{\lambda^2}; f[J]\right) + J(0) \text{err}_{2,d}^{(Bog, Con)}\left(\frac{T}{\lambda^2}; f\right), \tag{5.60}
\end{aligned}$$

where

$$|\text{err}_{2,d}^{(Bog, Bol)}\left(\frac{T}{\lambda^2}; f[J]\right)| \leq C \|\hat{v}\|_{w,d} \|f_0\|_d \|J\|_{\ell^\infty(\Lambda^*)} T^2 \frac{\lambda}{N}, \tag{5.61}$$

$$|\text{err}_{2,d}^{(Bog,Con)}(t; f)| \leq C_{\|\hat{v}\|_{w,d}, \|f_0\|_d} e^{C\|\hat{v}\|_{w,d}|\Lambda|/\lambda T} \frac{\langle T \rangle^4 |\Lambda|^{\frac{3}{2}}}{\lambda^3 N^{\frac{3}{2}}} \left(1 + \frac{|\Lambda|}{N}\right)^2 \left(1 + \frac{|\Lambda|^{\frac{3}{2}}}{\lambda N^{\frac{1}{2}}}\right), \quad (5.62)$$

$$(4) \quad -i \int_0^{T\lambda^{-2}} ds \frac{\nu_0(g[J], \mathcal{H}_{quart}(s))}{|\Lambda|} = \frac{1}{N} \int_0^T dS \text{abs}_{quart,d}(f_0)[J](S/\lambda^2), \quad (5.63)$$

The absorption operator $\text{abs}_{quart,d}(f_0)[J](S/\lambda^2)$ consists of terms of the form

$$(-i)^\ell \lambda^j \int dp dk e^{-i(m_1\Omega(p)+m_2\Omega(k))S/\lambda^2} (1 + f(p) + f(-p))J(p) V_1(p)^{2\alpha_1} V_2(p)^{\alpha_2} \hat{v}(p-k) V_1(k)^{2\alpha_3} V_2(k)^{\alpha_4} (f_0(k) + \iota), \quad (5.64)$$

with $\ell \in \mathbb{N}_0$, $\ell \leq 3$, $j \in \mathbb{N}_0$, $j \leq 7$, $m_1, m_2 \in \{0, \pm 2\}$, $\alpha_j \in \{0, 1, 2\}$ and $\iota \in \{0, 1\}$. The integrand contains a factor $f_0(k)$ or a factor $V_2(k)$.

$$(5) \quad - \int_{\Delta[T\lambda^{-2}, 2]} d\mathbf{s}_2 \frac{\nu_0([\![g[J], \mathcal{H}_{cub}(s_1)], \mathcal{H}_{cub}(s_2)]])}{|\Lambda|} \\ = \frac{1}{N\lambda^2} \int_{\Delta[T, 2]} d\mathbf{s}_2 (\text{col}_d(f_0)[J](\mathbf{s}_2/\lambda^2) + \text{abs}_{cub,d}(f_0)[J](\mathbf{s}_2/\lambda^2)) \\ + J(0) \left(\Phi_{T\lambda^{-2}}\right)^2 + J(0) \text{err}_2^{(Bog,Con)}\left(\frac{T}{\lambda^2}; g\right). \quad (5.65)$$

Here, the collision operator $\text{col}_d(f_0)[J](\mathbf{s}_2)$ consists of terms of the form

$$(-i)^{\ell_0} \lambda^{j_0} \int_{(\Lambda^*)^3} d\mathbf{p}_3 e^{-i\sum_{\ell=1}^2 s_\ell \sum_{k=1}^3 \sigma_{k,\ell} \Omega(p_k)} \delta(p_1 + p_2 - p_3) \\ J(p_{j_1}) \hat{v}(p_2) \hat{v}(p_{j_2}) \prod_{k=1}^3 V_1(p_k)^{2\alpha_{\ell_k}} V_2(p_k)^{\beta_{\ell_k}} \\ \left(\prod_{k=1}^3 (f_0(\tau_{k,1} p_k) + \frac{1 + \tau_{k,2}}{2}) - \prod_{k=1}^3 (f_0(-\tau_{k,1} p_k) + \frac{1 - \tau_{k,2}}{2}) \right), \quad (5.66)$$

where $\ell_0 \in \mathbb{N}_0$, $\ell_0 \leq 3$, $j_0 \in \mathbb{N}_0$, $j_0 \leq 12$, $\sigma_{k,\ell}, \tau_{k,\ell} \in \{\pm 1\}$, $j_1, j_2 \in \{1, 2, 3\}$ and $\alpha_{\ell_k}, \beta_{\ell_k} \in \{0, 1, 2\}$. Any term contains a product of at least two of the functions \hat{v} , f_0 , V_1 , and V_2 depending on at least two of the momenta $\{p_1, p_2, p_3\}$, which implies that the integrand in (5.66) exhibits the necessary regularity properties used in Propositions 5.7 and 5.9 below. The absorption operator $\text{abs}_{cub,d}(f_0)[J](\mathbf{s}_2)$ consists of terms of the form

$$(-i)^\ell \lambda^j \int_{(\Lambda^*)^2} dp dk e^{-is_1 m_1 \Omega(p) - is_2 m_2 \Omega(k)} J(p) (1 + f_0(p) + f_0(-p)) \\ \hat{v}(p) V_1(p)^{2\alpha_1} V_2(p)^{\alpha_2} \hat{v}(k) V_1(k)^{2\alpha_3} V_2(k)^{\alpha_4} (f_0(k) + \iota), \quad (5.67)$$

where $\ell \in \mathbb{N}_0$, $\ell \leq 3$, $j \in \mathbb{N}_0$, $j \leq 12$, $m_1, m_2 \in \{0, \pm 2\}$, $\alpha_j \in \{0, 1, 2\}$ and $\iota \in \{0, 1\}$. $|\text{err}_2^{(Bog,Con)}(\frac{T}{\lambda^2}; g)|$ satisfies the same bound as $|\text{err}_2^{(Bog,Con)}(\frac{T}{\lambda^2}; f)|$.

Proof. Set $t = T\lambda^{-2}$. Since here all integrals range over Λ^* in the sense of (1.51), we will omit the domain of integration in this proof. Using that $\nu_0(Q) = 0$ for

$[Q, \mathcal{N}_b] \neq 0$, (5.18), and recalling J_j from (5.19) and (5.22), we start by observing that

$$-i \int_0^t ds \frac{\nu_0([a_0, \mathcal{H}_{cub}(s)])}{|\Lambda|} = -i \int_0^t ds \left(\int dp f_0(p) J_1(s, p) + \frac{J_4(s)}{|\Lambda|} \right). \quad (5.68)$$

Here, we also used the fact that translation-invariance implies

$$\overline{a_p^{(\sigma)}} a_q^{(-\sigma)} = \delta(p - q)(f_0(p) + \delta_{\sigma, -1}). \quad (5.69)$$

Recall that, by Corollary 3.3,

$$\begin{aligned} J_1(s, p) &= \frac{\lambda \hat{v}(p)}{\sqrt{N}} (|u_\lambda(s, p)|^2 + |v_\lambda(s, p)|^2 + 2\overline{v_\lambda}(s, p)\overline{u_\lambda}(s, p)) \\ &= \frac{\lambda \hat{v}(p)}{\sqrt{N}} \left(1 + 2\lambda^2 \operatorname{Re}(e^{-i\Omega(p)s} i \sin(\Omega(p)s) V_1(p)) \right. \\ &\quad \left. + \lambda^4 \sin^2(\Omega(p)s) V_1(p)^4 + |v_\lambda(s, p)|^2 \right. \\ &\quad \left. + 2\overline{v_\lambda}(s, p)\overline{u_\lambda}(s, p) \right), \end{aligned} \quad (5.70)$$

$$J_4(s) = \frac{\lambda}{\sqrt{N}} \int dp \hat{v}(p) (|v_\lambda(s, p)|^2 + \overline{u_\lambda}(s, p)\overline{v_\lambda}(s, p)). \quad (5.71)$$

Thus, again by Corollary 3.3, we obtain

$$\begin{aligned} \|J_1(s) + \frac{\lambda \hat{v}}{\sqrt{N}}\|_\infty &\leq C \frac{\lambda \|\hat{v}\|_\infty}{\sqrt{N}} \left(\lambda^2 \|V_1\|_\infty^2 + \lambda^4 \|V_1\|_\infty^4 + \|v_\lambda(s, \cdot)\|_\infty^2 \right. \\ &\quad \left. + \|v_\lambda(s, \cdot)\|_\infty \|u_\lambda(t, \cdot)\|_\infty \right) \\ &\leq C \frac{\lambda^2 \|\hat{v}\|_{w,d}^2 (1 + \lambda \|\hat{v}\|_{w,d})^3}{\sqrt{N}}, \end{aligned} \quad (5.72)$$

$$\begin{aligned} |J_4(s)| &\leq C \frac{\lambda \|\hat{v}\|_1}{\sqrt{N}} (\|v_\lambda(s, \cdot)\|_\infty^2 + \|v_\lambda(s, \cdot)\|_\infty \|u_\lambda(t, \cdot)\|_\infty) \\ &\leq C \frac{\lambda^2 \|\hat{v}\|_{w,d}^2 (1 + \lambda \|\hat{v}\|_{w,d})^3}{\sqrt{N}} \end{aligned} \quad (5.73)$$

for all $s \geq 0$. Then (5.68), Lemma 4.7, and the definition (5.57) of $\operatorname{err}_{1,d}^{(Bog)}(t; \Phi)$ yield

$$\begin{aligned} |\operatorname{err}_{1,d}^{(Bog)}(t; \Phi)| &\leq C \frac{\lambda^2 t \|\hat{v}\|_{w,d}^2 (1 + \lambda \|\hat{v}\|_{w,d})^3 (|\Lambda|^{-1} + \|f_0\|_1)}{N^{\frac{1}{2}}} \\ &\leq \frac{C \|\hat{v}\|_{w,d} \|f_0\|_d T}{N^{\frac{1}{2}}}. \end{aligned} \quad (5.74)$$

Collecting (5.68), (5.72), (5.73), and (5.74), we have proved (5.57).

Next, we compute the dynamics of f . We use the fact that ν_0 is quasifree to obtain that

$$\begin{aligned} \nu_0([a_p^+ a_p, \mathcal{H}_{quart}(s)]) &= \overline{[a_p^+ a_p, \mathcal{H}_{quart}(s)]} + [a_p^+ a_p, \mathcal{H}_{quart}(s)] \\ &\propto f_0(p) \tilde{f}_0(p) - \tilde{f}_0(p) f_0(p) + 0 = 0, \end{aligned} \quad (5.75)$$

since, due to Corollary (3.3), $\mathcal{H}_{quart}(s)$ is a quartic polynomial.

We observe that for self-adjoint operators A, B, C and any state ν we have that

$$\nu([[A, B], C]) = \nu(\overline{[[A, B], C]}) = \text{Re}(\nu([[A, B], C])). \quad (5.76)$$

Similarly to the previous case, we have

$$\begin{aligned} \overline{[[f[J], \mathcal{H}_{cub}(s_1)], \mathcal{H}_{cub}(s_2)]} &= \overline{[[f[J], \mathcal{H}_{cub}(s_1)], \mathcal{H}_{cub}(s_2)]} \\ &= \overline{[[f[J], \mathcal{H}_{cub}(s_1)], \mathcal{H}_{cub}(s_2)]} \\ &= 0. \end{aligned} \quad (5.77)$$

This leaves us with exactly two types of possible contractions. These are

$$(-i)^2 \overline{[[f[J], \mathcal{H}_{cub}(s_1)], \mathcal{H}_{cub}(s_2)]} \quad (5.78)$$

and, using translation invariance and employing Lemma A.1,

$$\begin{aligned} (-i)^2 \overline{[[f[J], \mathcal{H}_{cub}(s_1)], \mathcal{H}_{cub}(s_2)]} &= -\frac{2J(0)}{|\Lambda|} \text{Re} \left(\overline{[a_0, \mathcal{H}_{cub}(s_1)]} \overline{[a_0^+, \mathcal{H}_{cub}(s_2)]} \right). \end{aligned} \quad (5.79)$$

(5.78) corresponds to a scattering or Boltzmann term, while (5.79) describes corrections to the evolution of the condensate.

We start by analyzing the condensate term. We have that

$$\begin{aligned} &\frac{-2J(0)}{|\Lambda|^2} \text{Re} \int_{\Delta[t,2]} ds_2 \overline{[a_0, \mathcal{H}_{cub}(s_1)]} \overline{[a_0^+, \mathcal{H}_{cub}(s_2)]} \\ &= \frac{-J(0)}{|\Lambda|^2} \int_{\Delta[t,2]} ds_2 \left(\overline{[a_0, \mathcal{H}_{cub}(s_1)]} \overline{[a_0^+, \mathcal{H}_{cub}(s_2)]} \right. \\ &\quad \left. + \overline{[a_0, \mathcal{H}_{cub}(s_2)]} \overline{[a_0^+, \mathcal{H}_{cub}(s_1)]} \right) \\ &= -\frac{J(0)}{2|\Lambda|^2} \int ds_2 \left(\overline{[a_0, \mathcal{H}_{cub}(s_1)]} \overline{[a_0^+, \mathcal{H}_{cub}(s_2)]} \right. \\ &\quad \left. + \overline{[a_0, \mathcal{H}_{cub}(s_2)]} \overline{[a_0^+, \mathcal{H}_{cub}(s_1)]} \right) \\ &= J(0) \left| -i \int_0^t \frac{ds}{|\Lambda|} \overline{[a_0, \mathcal{H}_{cub}(s)]} \right|^2. \end{aligned} \quad (5.80)$$

Here, we used the fact that

$$\overline{[a_0, \mathcal{H}_{cub}(s)]} = -\overline{[a_0^+, \mathcal{H}_{cub}(s)]}. \quad (5.81)$$

Next, we apply quasifreeness of ν_0 , followed by Proposition 5.1, to get that

$$\begin{aligned} -i \int_0^t ds \frac{\overline{[a_0, \mathcal{H}_{cub}(s)]}}{|\Lambda|} &= -i \int_0^t ds \frac{\nu_0([a_0, \mathcal{H}_{cub}(s)])}{|\Lambda|} \\ &= \Phi_t - \text{Rem}_2(t; \Phi). \end{aligned} \quad (5.82)$$

In particular, we have that the condensate term is given by

$$J(0)|\Phi_t|^2 + J(0) \text{err}_{2,d}^{(Bog, Con)}(t; f), \quad (5.83)$$

where

$$\begin{aligned} \text{err}_{2,d}^{(Bog,Con)}(t; f) &= -2\text{Re}\left((-i) \int_0^t ds \frac{\nu_0([a_0, \mathcal{H}_{cub}(s)])}{|\Lambda|} \overline{\text{Rem}_2(t; \Phi)}\right) \\ &\quad - |\text{Rem}_2(t; \Phi)|^2. \end{aligned} \quad (5.84)$$

Using (5.68), (5.72), and (5.73), we have that

$$\begin{aligned} \left| \int_0^t ds \frac{\nu_0([a_0, \mathcal{H}_{cub}(s)])}{|\Lambda|} \right| &\leq C_{\|\hat{v}\|_{w,d}} \frac{\lambda t(1 + \|f_0\|_d)}{\sqrt{N}} \\ &\leq \frac{C_{\|\hat{v}\|_{w,d}, \|f_0\|_d} T}{N^{\frac{1}{2}} \lambda} \end{aligned} \quad (5.85)$$

for all $\lambda \in (0, 1)$. Thus, Proposition 5.1 and (5.85) yield

$$\begin{aligned} &|\text{err}_{2,d}^{(Bog,Con)}(t; f)| \\ &\leq C_{\|\hat{v}\|_{w,d}, \|f_0\|_d} \left[\frac{T}{N^{\frac{1}{2}} \lambda} T^2 e^{C_{\|\hat{v}\|_{w,d}} |\Lambda| / \lambda T} \frac{|\Lambda|^{\frac{3}{2}}}{N \lambda^2} \left(1 + \frac{|\Lambda|}{N}\right) \right. \\ &\quad \left. + T^4 e^{C_{\|\hat{v}\|_{w,d}} |\Lambda| / \lambda T} \frac{|\Lambda|^3}{N^2 \lambda^4} \left(1 + \frac{|\Lambda|}{N}\right)^2 \right] \\ &\leq C_{\|\hat{v}\|_{w,d}, \|f_0\|_d} e^{C_{\|\hat{v}\|_{w,d}} |\Lambda| / \lambda T} \frac{\langle T \rangle^4 |\Lambda|^{\frac{3}{2}}}{\lambda^3 N^{\frac{3}{2}}} \left(1 + \frac{|\Lambda|}{N}\right)^2 \left(1 + \frac{|\Lambda|^{\frac{3}{2}}}{\lambda N^{\frac{1}{2}}}\right). \end{aligned} \quad (5.86)$$

Next, we compute the Boltzmann term for f . Observe that with the decomposition (5.27), we have that

$$\mathcal{H}_{cub}(t) = \mathcal{H}_{cub}^{(1)}(t) + \mathcal{H}_{cub}^{(2)}(t). \quad (5.87)$$

Notice that $\mathcal{H}_{cub}^{(1)}(t)$ and $\mathcal{H}_{cub}^{(2)}(t)$ are formal adjoints. In particular, we have that

$$\begin{aligned} &\nu_0([f[J], \mathcal{H}_{cub}(s_1)], \mathcal{H}_{cub}(s_2)) \\ &= 2\text{Re}\nu_0([f[J], \mathcal{H}_{cub}^{(1)}(s_1)], \mathcal{H}_{cub}^{(1)}(s_2)) \\ &\quad + 2\text{Re}\nu_0([f[J], \mathcal{H}_{cub}^{(1)}(s_1)], \mathcal{H}_{cub}^{(2)}(s_2)) \end{aligned} \quad (5.88)$$

With that, we sort the Boltzmann contractions of $\nu_0([f[J], \mathcal{H}_{cub}(s_1)], \mathcal{H}_{cub}(s_2))$ by powers of λ , i.e.,

$$\begin{aligned} -\frac{1}{|\Lambda|} \overbrace{[f[J], \mathcal{H}_{cub}(s_1)], \mathcal{H}_{cub}(s_2)]} &= \frac{\lambda^2}{N} (\text{bol}^{(0)}(f_0)[J](s_1, s_2) \\ &\quad + \lambda \text{bol}^{(1)}(f_0)[J](s_1, s_2) \\ &\quad + \lambda^2 \text{bol}^{(2)}(f_0)[J](s_1, s_2) \\ &\quad + \lambda^3 \text{bol}^{(3)}(f_0)[J](s_1, s_2)), \end{aligned} \quad (5.89)$$

Notice that the CCR imply

$$[f[J], a_{\sigma_1 p_1}^{(\sigma_1)} a_{\sigma_2 p_2}^{(\sigma_2)} a_{\sigma_3 p_{12}}^{(-\sigma_3)}] = \sum_{j=1}^3 \sigma_j (-1)^{\delta_{j,3}} J(p_j) a_{\sigma_1 p_1}^{(\sigma_1)} a_{\sigma_2 p_2}^{(\sigma_2)} a_{\sigma_3 p_{12}}^{(-\sigma_3)}. \quad (5.90)$$

We have that

$$\begin{aligned} \text{bol}^{(0)}(f_0)[J](\mathbf{s}_2) &= -2\text{Re} \int d\mathbf{p}_2 d\mathbf{k}_2 e^{i\Delta_{cub}\Omega(\mathbf{p}_2) s_1 - i\Delta_{cub}\Omega(\mathbf{k}_2) s_2} \\ &\quad \hat{v}(p_2) \hat{v}(k_2) \Delta_{cub} J(\mathbf{p}_2) \end{aligned}$$

$$\begin{aligned}
 & \frac{1}{|\Lambda|} \left(\overbrace{[a_{p_1}^+ a_{p_2}^+ a_{p_{12}}^+, a_{k_{12}}^+ a_{k_2} a_{k_1}]} + \overbrace{[a_{p_1}^+ a_{p_2}^+ a_{p_{12}}^+, a_{k_{12}}^+ a_{k_2} a_{k_1}]} \right) \\
 &= -2\text{Re} \int d\mathbf{p}_2 e^{i\Delta_{cub}\Omega(\mathbf{p}_2)(s_1-s_2)} \hat{v}(p_2) \\
 & \quad (\hat{v}(p_1) + \hat{v}(p_2)) \Delta_{cub} J(\mathbf{p}_2) \\
 & \quad (f_0(p_1) f_0(p_2) \tilde{f}_0(p_{12}) - \tilde{f}_0(p_1) \tilde{f}_0(p_2) f_0(p_{12})) \\
 &= \text{Re} \int d\mathbf{p}_2 e^{i\Delta_{cub}\Omega(\mathbf{p}_2)(s_1-s_2)} (\hat{v}(p_1) + \hat{v}(p_2))^2 \Delta_{cub} J(\mathbf{p}_2) \\
 & \quad (\tilde{f}_0(p_1) \tilde{f}_0(p_2) f_0(p_{12}) - f_0(p_1) f_0(p_2) \tilde{f}_0(p_{12})). \tag{5.91}
 \end{aligned}$$

Here, we used (5.69) followed by symmetry $p_1 \leftrightarrow p_2$.

Recalling Corollary 3.3, $\text{bol}^{(1)}(f_0)[J](s_1, s_2)$ is given by the Boltzmann contractions of

$$\begin{aligned}
 & -\frac{2}{|\Lambda|} \text{Re} i \int d\mathbf{p}_2 d\mathbf{k}_2 \hat{v}(p_2) \hat{v}(k_2) e^{i\Delta_{cub}\Omega(\mathbf{p}_2)s_1 + i\Delta_{cub}\Omega(\mathbf{k}_2)s_2} \\
 & \quad \left(e^{-i\Omega(p_1)s_1} \sin(\Omega(p_1)s_1) V_2(p_1) \nu_0([f[J], a_{-p_1} a_{p_2}^+ a_{p_{12}}^+], a_{k_1}^+ a_{k_2}^+ a_{k_{12}}]) \right. \\
 & \quad + e^{-i\Omega(p_2)s_1} \sin(\Omega(p_2)s_1) V_2(p_2) \nu_0([f[J], a_{p_1}^+ a_{-p_2} a_{p_{12}}^+], a_{k_1}^+ a_{k_2}^+ a_{k_{12}}]) \\
 & \quad + e^{-i\Omega(k_1)s_2} \sin(\Omega(k_1)s_2) V_2(k_1) \nu_0([f[J], a_{p_1}^+ a_{p_2}^+ a_{p_{12}}^+], a_{-k_1} a_{k_2}^+ a_{k_{12}}]) \\
 & \quad \left. + e^{-i\Omega(k_2)s_2} \sin(\Omega(k_2)s_2) V_2(k_2) \nu_0([f[J], a_{p_1}^+ a_{p_2}^+ a_{p_{12}}^+], a_{k_1}^+ a_{-k_2} a_{k_{12}}]) \right) \tag{5.92}
 \end{aligned}$$

Using (5.90), obtain, similarly to above,

$$\begin{aligned}
 & 2\text{Im} \int d\mathbf{p}_2 \hat{v}(p_2) \left[(-J(-p_1) + J(p_2) - J(p_{12})) \right. \\
 & \quad (\hat{v}(p_1) + \hat{v}(p_{12})) e^{i(\Omega(p_2) - \Omega(p_{12}))(s_1-s_2) + i\Omega(p_1)s_2} \sin(\Omega(p_1)s_1) V_2(p_1) \\
 & \quad (\tilde{f}_0(-p_1) f_0(p_2) \tilde{f}_0(p_{12}) - f_0(-p_1) \tilde{f}_0(p_2) f_0(p_{12})) \\
 & \quad + (J(p_1) - J(-p_2) - J(p_{12})) (\hat{v}(p_2) + \hat{v}(p_{12})) \\
 & \quad e^{i(\Omega(p_1) - \Omega(p_{12}))(s_1-s_2) + i\Omega(p_2)s_2} \sin(\Omega(p_2)s_1) V_2(p_2) \\
 & \quad (f_0(p_1) \tilde{f}_0(-p_2) \tilde{f}_0(p_{12}) - \tilde{f}_0(p_1) f_0(-p_2) f_0(p_{12})) \\
 & \quad + \left(e^{i(\Omega(p_1) - \Omega(p_{12}))(s_1-s_2) + i\Omega(p_2)s_1} \sin(\Omega(p_2)s_2) V_2(p_2) (\hat{v}(p_2) + \hat{v}(p_{12})) \right. \\
 & \quad \left. + e^{i(\Omega(p_2) - \Omega(p_{12}))(s_1-s_2) + i\Omega(p_1)s_1} \sin(\Omega(p_1)s_2) V_2(p_1) (\hat{v}(p_1) + \hat{v}(p_{12})) \right) \\
 & \quad \left. \Delta_{cub} J(\mathbf{p}_2) (f_0(p_1) f_0(p_2) \tilde{f}_0(p_{12}) - \tilde{f}_0(p_1) \tilde{f}_0(p_2) f_0(p_{12})) \right]. \tag{5.93}
 \end{aligned}$$

Here, we used the fact that \hat{v} , and thus Ω , are even functions. Using symmetry in $p_1 \leftrightarrow p_2$, we can further simplify this expression to

$$\begin{aligned}
 & \text{bol}^{(1)}(f_0)[J](s_2) \\
 & := 2\text{Im} \int d\mathbf{p}_2 (\hat{v}(p_1) + \hat{v}(p_2)) (\hat{v}(p_1) + \hat{v}(p_{12})) \\
 & \quad V_2(p_1) e^{i(\Omega(p_2) - \Omega(p_{12}))(s_1-s_2)} \\
 & \quad \left((-J(-p_1) + J(p_2) - J(p_{12})) e^{i\Omega(p_1)s_2} \sin(\Omega(p_1)s_1) \right.
 \end{aligned}$$

$$\begin{aligned}
& (\tilde{f}_0(-p_1)f_0(p_2)\tilde{f}_0(p_{12}) - f_0(-p_1)\tilde{f}_0(p_2)f_0(p_{12})) \\
& + \Delta_{cub}J(\mathbf{p}_2)e^{i\Omega(p_1)s_1}\sin(\Omega(p_1)s_2) \\
& (f_0(p_1)f_0(p_2)\tilde{f}_0(p_{12}) - \tilde{f}_0(p_1)\tilde{f}_0(p_2)f_0(p_{12})). \tag{5.94}
\end{aligned}$$

Recalling (5.52), we have that

$$\int_{\Delta_{[T\lambda^{-2}, 2]}} ds_2 \text{bol}^{(1)}(f_0)[J](s_2) = \text{Bol}_d^{(1)}(f_0)[J](T; \lambda). \tag{5.95}$$

Next, we compute the corrections with coefficient $\frac{\lambda^4}{N}$. These come either from a correction to the same momentum, $a_p^+ \rightarrow i\lambda^2 V_1(p)^2 \sin(\Omega(p)t)a_p^+$, or from two momentum flips, $a_p^+ \rightarrow i\lambda V_2(p) \sin(\Omega(p)t)a_{-p}$. In the first case, computations analogous to (5.91) yield

$$\begin{aligned}
& -2\text{Re}i \int d\mathbf{p}_2 \hat{v}(p_2)(\hat{v}(p_1) + \hat{v}(p_2))\Delta_{cub}J(\mathbf{p}_2)e^{i\Delta_{cub}\Omega(\mathbf{p}_2)(s_1-s_2)} \\
& \left(V_1(p_1)^2 (\sin(\Omega(p_1)s_1)e^{-i\Omega(p_1)s_1} - \sin(\Omega(p_1)s_2)e^{i\Omega(p_1)s_2}) \right. \\
& + V_1(p_2)^2 (\sin(\Omega(p_2)s_1)e^{-i\Omega(p_2)s_1} - \sin(\Omega(p_2)s_2)e^{i\Omega(p_2)s_2}) \\
& \left. - V_1(p_{12})^2 (\sin(\Omega(p_{12})s_1)e^{i\Omega(p_{12})s_1} - \sin(\Omega(p_{12})s_2)e^{i\Omega(p_{12})s_2}) \right) \\
& (f_0(p_1)f_0(p_2)\tilde{f}_0(p_{12}) - \tilde{f}_0(p_1)\tilde{f}_0(p_2)f_0(p_{12})). \tag{5.96}
\end{aligned}$$

Using symmetry in $p_1 \leftrightarrow p_2$ again, we obtain

$$\begin{aligned}
& \text{bol}_1^{(2)}(f_0)[J](s_2) \\
& := \text{Im} \int d\mathbf{p}_2 (\hat{v}(p_1) + \hat{v}(p_2))^2 \Delta_{cub}J(\mathbf{p}_2)e^{i\Delta_{cub}\Omega(\mathbf{p}_2)(s_1-s_2)} \\
& \left(V_1(p_1)^2 (\sin(\Omega(p_1)s_1)e^{-i\Omega(p_1)s_1} - \sin(\Omega(p_1)s_2)e^{i\Omega(p_1)s_2}) \right. \\
& + V_1(p_2)^2 (\sin(\Omega(p_2)s_1)e^{-i\Omega(p_2)s_1} - \sin(\Omega(p_2)s_2)e^{i\Omega(p_2)s_2}) \\
& \left. - V_1(p_{12})^2 (\sin(\Omega(p_{12})s_1)e^{i\Omega(p_{12})s_1} - \sin(\Omega(p_{12})s_2)e^{i\Omega(p_{12})s_2}) \right) \\
& (f_0(p_1)f_0(p_2)\tilde{f}_0(p_{12}) - \tilde{f}_0(p_1)\tilde{f}_0(p_2)f_0(p_{12})). \tag{5.97}
\end{aligned}$$

In the case of two momentum flips, the correction terms are given by the Boltzmann contractions of

$$\begin{aligned}
& -\frac{2}{|\Lambda|}\text{Re} \int d\mathbf{p}_2 d\mathbf{k}_2 \hat{v}(p_2)\hat{v}(k_2)e^{i\Delta_{cub}\Omega(\mathbf{p}_2)s_1 - i\Delta_{cub}\Omega(\mathbf{k}_2)s_2} \\
& \left(e^{-i(\Omega(p_1) - \Omega(p_{12}))s_1} i \sin(\Omega(p_1)s_1)(-i) \sin(\Omega(p_{12})s_1)V_2(p_1)V_2(p_{12}) \right. \\
& \quad \left. (-J(-p_1) + J(p_2) + J(-p_{12}))\nu_0([a_{-p_1}a_{p_2}^+ a_{-p_{12}}^+, a_{k_{12}}^+ a_{k_2} a_{k_1}]) \right) \\
& + e^{-i(\Omega(p_2) - \Omega(p_{12}))s_1} i \sin(\Omega(p_2)s_1)(-i) \sin(\Omega(p_{12})s_1)V_2(p_2)V_2(p_{12}) \\
& \quad \left. (J(p_1) - J(-p_2) + J(-p_{12}))\nu_0([a_{p_1}^+ a_{-p_2} a_{-p_{12}}^+, a_{k_{12}}^+ a_{k_2} a_{k_1}]) \right) \\
& + e^{-i\Omega(p_1)s_1 + i\Omega(k_2)s_2} i \sin(\Omega(p_1)s_1)(-i) \sin(\Omega(k_2)s_2)V_2(p_1)V_2(k_2) \\
& \quad \left. (-J(-p_1) + J(p_2) - J(p_{12}))\nu_0([a_{-p_1}a_{p_2}^+ a_{p_{12}}, a_{k_{12}}^+ a_{-k_2}^+ a_{k_1}]) \right) \\
& + e^{-i\Omega(p_2)s_1 + i\Omega(k_2)s_2} i \sin(\Omega(p_2)s_1)(-i) \sin(\Omega(k_2)s_2)V_2(p_2)V_2(k_2)
\end{aligned}$$

$$\begin{aligned}
& (J(p_1) - J(-p_2) - J(p_{12}))\nu_0([a_{p_1}^+ a_{-p_2} a_{p_{12}}, a_{k_{12}}^+ a_{-k_2}^+ a_{k_1}]) \\
& + e^{-i\Omega(p_1)s_1 + i\Omega(k_1)s_2} i \sin(\Omega(p_1)s_1)(-i) \sin(\Omega(k_1)s_2) V_2(p_1) V_2(k_1) \\
& (-J(-p_1) + J(p_2) - J(p_{12}))\nu_0([a_{-p_1} a_{p_2}^+ a_{p_{12}}, a_{k_{12}}^+ a_{k_2} a_{-k_1}^+]) \\
& + e^{-i\Omega(p_2)s_1 + i\Omega(k_1)s_2} i \sin(\Omega(p_2)s_1)(-i) \sin(\Omega(k_1)s_2) V_2(p_2) V_2(k_1) \\
& (J(p_1) - J(-p_2) - J(p_{12}))\nu_0([a_{p_1}^+ a_{-p_2} a_{p_{12}}, a_{k_{12}}^+ a_{k_2} a_{-k_1}^+]) \\
& + e^{i\Omega(p_{12})s_1 - i\Omega(k_{12})s_2} (-i) \sin(\Omega(p_{12})s_1) i \sin(\Omega(k_{12})s_2) V_2(p_{12}) V_2(k_{12}) \\
& (J(p_1) + J(p_2) + J(-p_{12}))\nu_0([a_{p_1}^+ a_{p_2}^+ a_{-p_{12}}^+, a_{-k_{12}} a_{k_2} a_{k_1}]) \Big). \quad (5.98)
\end{aligned}$$

This expression equals

$$\begin{aligned}
& -2\text{Re} \int d\mathbf{p}_2 \hat{v}(p_2) \Big[\sin(\Omega(p_1)s_1) \sin(\Omega(p_{12})s_1) e^{i\Omega(p_2)(s_1-s_2) + i(\Omega(p_1) - \Omega(p_{12}))s_2} \\
& V_2(p_1) V_2(p_{12}) (\hat{v}(p_2) + \hat{v}(p_{12})) (-J(-p_1) + J(p_2) + J(-p_{12})) \\
& (\tilde{f}_0(-p_1) f_0(p_2) f(-p_{12}) - f_0(-p_1) \tilde{f}_0(p_2) \tilde{f}_0(-p_{12})) \\
& + \sin(\Omega(p_2)s_1) \sin(\Omega(p_{12})s_1) e^{i\Omega(p_1)(s_1-s_2) + i(\Omega(p_2) - \Omega(p_{12}))s_2} \\
& V_2(p_2) V_2(p_{12}) (\hat{v}(p_1) + \hat{v}(p_{12})) (J(p_1) - J(-p_2) + J(-p_{12})) \\
& (f_0(p_1) \tilde{f}_0(-p_2) f(-p_{12}) - \tilde{f}_0(p_1) f_0(-p_2) \tilde{f}_0(-p_{12})) \\
& + \sin(\Omega(p_1)s_1) e^{i\Omega(p_2)(s_1-s_2) - i\Omega(p_{12})s_1} V_2(p_1) (-J(-p_1) + J(p_2) - J(p_{12})) \\
& (\hat{v}(p_1) V_2(p_1) \sin(\Omega(p_1)s_2) e^{i\Omega(p_{12})s_2} + \hat{v}(p_{12}) V_2(p_{12}) \sin(\Omega(p_{12})s_2) e^{-i\Omega(p_1)s_2}) \\
& (\tilde{f}_0(-p_1) f_0(p_2) \tilde{f}_0(p_{12}) - f_0(-p_1) \tilde{f}_0(p_2) f_0(p_{12})) \\
& + \sin(\Omega(p_2)s_1) e^{i\Omega(p_1)(s_1-s_2) - i\Omega(p_{12})s_1} V_2(p_2) (J(p_1) - J(-p_2) - J(p_{12})) \\
& (\hat{v}(p_2) V_2(p_2) \sin(\Omega(p_2)s_2) e^{i\Omega(p_{12})s_2} + \hat{v}(p_{12}) V_2(p_{12}) \sin(\Omega(p_{12})s_2) e^{-i\Omega(p_2)s_2}) \\
& (f_0(p_1) \tilde{f}_0(-p_2) \tilde{f}_0(p_{12}) - \tilde{f}_0(p_1) f_0(-p_2) f_0(p_{12})) \\
& + \sin(\Omega(p_1)s_1) e^{i\Omega(p_2)(s_1-s_2) - i\Omega(p_{12})s_1} V_2(p_1) \hat{v}(p_2) (-J(-p_1) + J(p_2) - J(p_{12})) \\
& (V_2(p_1) \sin(\Omega(p_1)s_2) e^{i\Omega(p_{12})s_2} + V_2(p_{12}) \sin(\Omega(p_{12})s_2) e^{i\Omega(p_1)s_2}) \\
& (\tilde{f}_0(-p_1) f_0(p_2) \tilde{f}_0(p_{12}) - f_0(-p_1) \tilde{f}_0(p_2) f_0(p_{12})) \\
& + \sin(\Omega(p_2)s_1) e^{i\Omega(p_1)(s_1-s_2) - i\Omega(p_{12})s_1} V_2(p_2) \hat{v}(p_1) (J(p_1) - J(-p_2) - J(p_{12})) \\
& (V_2(p_2) \sin(\Omega(p_2)s_2) e^{i\Omega(p_{12})s_2} + V_2(p_{12}) \sin(\Omega(p_{12})s_2) e^{i\Omega(p_2)s_2}) \\
& (f_0(p_1) \tilde{f}_0(-p_2) \tilde{f}_0(p_{12}) - \tilde{f}_0(p_1) f_0(-p_2) f_0(p_{12})) \\
& + V_2(p_2) e^{i(\Omega(p_1) + \Omega(p_2))s_1} \sin(\Omega(p_{12})s_1) (J(p_1) + J(p_2) + J(-p_{12})) \\
& + \left(e^{-i(\Omega(p_1) + \Omega(p_2))s_2} \sin(\Omega(p_{12})s_2) V_2(p_{12}) (\hat{v}(p_1) + \hat{v}(p_2)) \right. \\
& + e^{-i(\Omega(p_1) + \Omega(p_{12}))s_2} \sin(\Omega(p_2)s_2) V_2(p_2) (\hat{v}(p_1) + \hat{v}(p_{12})) \\
& \left. + \sin(\Omega(p_1)s_2) e^{-i(\Omega(p_2) + \Omega(p_{12}))s_2} V_2(p_1) (\hat{v}(p_2) + \hat{v}(p_{12})) \right) \\
& (f_0(p_1) f_0(p_2) f_0(-p_{12}) - \tilde{f}_0(p_1) \tilde{f}_0(p_2) \tilde{f}_0(-p_{12})) \Big]. \quad (5.99)
\end{aligned}$$

Using symmetry in $p_1 \leftrightarrow p_2$, this can be reduced to

$$\begin{aligned}
& -2\operatorname{Re} \int d\mathbf{p}_2 (\hat{v}(p_1) + \hat{v}(p_2)) \\
& \left[\sin(\Omega(p_1)s_1) \sin(\Omega(p_{12})s_1) e^{i\Omega(p_2)(s_1-s_2)+i(\Omega(p_1)-\Omega(p_{12}))s_2} \right. \\
& V_2(p_1)V_2(p_{12})(\hat{v}(p_2) + \hat{v}(p_{12}))(-J(-p_1) + J(p_2) + J(-p_{12})) \\
& (\tilde{f}_0(-p_1)f_0(p_2)f(-p_{12}) - f_0(-p_1)\tilde{f}_0(p_2)\tilde{f}_0(-p_{12})) \\
& + \sin(\Omega(p_1)s_1) e^{i\Omega(p_2)(s_1-s_2)-i\Omega(p_{12})s_1} V_2(p_1)(-J(-p_1) + J(p_2) - J(p_{12})) \\
& (\hat{v}(p_1)V_2(p_1) \sin(\Omega(p_1)s_2) e^{i\Omega(p_{12})s_2} + \hat{v}(p_{12})V_2(p_{12}) \sin(\Omega(p_{12})s_2) e^{-i\Omega(p_1)s_2}) \\
& (\tilde{f}_0(-p_1)f_0(p_2)\tilde{f}_0(p_{12}) - f_0(-p_1)\tilde{f}_0(p_2)f_0(p_{12})) \\
& + \sin(\Omega(p_1)s_1) e^{i\Omega(p_2)(s_1-s_2)-i\Omega(p_{12})s_1} V_2(p_1)\hat{v}(p_2)(-J(-p_1) + J(p_2) - J(p_{12})) \\
& (V_2(p_1) \sin(\Omega(p_1)s_2) e^{i\Omega(p_{12})s_2} + V_2(p_{12}) \sin(\Omega(p_{12})s_2) e^{i\Omega(p_1)s_2}) \\
& (\tilde{f}_0(-p_1)f_0(p_2)\tilde{f}_0(p_{12}) - f_0(-p_1)\tilde{f}_0(p_2)f_0(p_{12})) \\
& \left. \left(\frac{1}{2} e^{-i(\Omega(p_1)-\Omega(p_2))s_2} \sin(\Omega(p_{12})s_2) V_2(p_{12})(\hat{v}(p_1) + \hat{v}(p_2)) \right. \right. \\
& \left. \left. + e^{-i(\Omega(p_1)-\Omega(p_{12}))s_2} \sin(\Omega(p_2)s_2) V_2(p_2)(\hat{v}(p_1) + \hat{v}(p_{12})) \right) \right. \\
& \left. (f_0(p_1)f_0(p_2)f_0(-p_{12}) - \tilde{f}_0(p_1)\tilde{f}_0(p_2)\tilde{f}_0(-p_{12})) \right]. \tag{5.100}
\end{aligned}$$

This can be further be simplified to

$$\begin{aligned}
& \operatorname{bol}_2^{(2)}(f_0)[J](\mathbf{s}_2) \\
& := 2\operatorname{Re} \int d\mathbf{p}_2 (\hat{v}(p_1) + \hat{v}(p_2)) \\
& \left[\sin(\Omega(p_1)s_1) \sin(\Omega(p_{12})s_1) e^{i\Omega(p_2)(s_1-s_2)+i(\Omega(p_1)-\Omega(p_{12}))s_2} \right. \\
& V_2(p_1)V_2(p_{12})(\hat{v}(p_2) + \hat{v}(p_{12}))(-J(-p_1) + J(p_2) + J(-p_{12})) \\
& (f_0(-p_1)\tilde{f}_0(p_2)\tilde{f}_0(-p_{12}) - \tilde{f}_0(-p_1)f_0(p_2)f(-p_{12})) \\
& + \sin(\Omega(p_1)s_1) e^{i\Omega(p_2)(s_1-s_2)-i\Omega(p_{12})s_1} V_2(p_1) \\
& (-J(-p_1) + J(p_2) - J(p_{12})) \\
& \left((\hat{v}(p_1) + \hat{v}(p_2))V_2(p_1) \sin(\Omega(p_1)s_2) e^{-i\Omega(p_{12})s_2} \right. \\
& \left. + (\hat{v}(p_2) + \hat{v}(p_{12}))V_2(p_{12}) \sin(\Omega(p_{12})s_2) e^{-i\Omega(p_1)s_2} \right) \\
& (f_0(-p_1)\tilde{f}_0(p_2)f_0(p_{12}) - \tilde{f}_0(-p_1)f_0(p_2)\tilde{f}_0(p_{12})) \\
& \left(\frac{1}{2} e^{-i(\Omega(p_1)+\Omega(p_2))s_2} \sin(\Omega(p_{12})s_2) V_2(p_{12})(\hat{v}(p_1) + \hat{v}(p_2)) \right. \\
& \left. + e^{-i(\Omega(p_1)+\Omega(p_{12}))s_2} \sin(\Omega(p_2)s_2) V_2(p_2)(\hat{v}(p_1) + \hat{v}(p_{12})) \right) \\
& \left. (\tilde{f}_0(p_1)\tilde{f}_0(p_2)\tilde{f}_0(-p_{12}) - f_0(p_1)f_0(p_2)f_0(-p_{12})) \right]. \tag{5.101}
\end{aligned}$$

In addition, let

$$\operatorname{bol}^{(2)}(f_0)[J](\mathbf{s}_2) := \operatorname{bol}_1^{(2)}(f_0)[J](\mathbf{s}_2) + \operatorname{bol}_2^{(2)}(f_0)[J](\mathbf{s}_2). \tag{5.102}$$

Recalling definitions (5.53), (5.54), and (5.56), we obtain that

$$\int_{\Delta[T\lambda^{-2}, 2]} ds_2 \text{bol}^{(2)}(f_0)[J](s_2) = \text{Bol}_d^{(2)}(f_0)[J](T; \lambda). \quad (5.103)$$

In order to bound $\text{bol}^{(3)}(f_0)[J](s_2)$, we need to, first, look at integrability of the terms. Notice that, for Boltzmann contractions, the order of the creation/annihilation operators within a single argument of a commutator do not matter. In particular, we are interested in evaluating expressions involving the Boltzmann contractions of

$$[a_{\sigma_1 p_1}^{(\sigma_1)} a_{\sigma_2 p_2}^{(\sigma_2)} a_{\sigma_3 p_{12}}^{(-\sigma_3)}, a_{\tau_1 k_1}^{(\tau_1)} a_{\tau_2 k_2}^{(\tau_2)} a_{\tau_3 k_{12}}^{(-\tau_3)}], \quad (5.104)$$

which contain a factor

$$\begin{aligned} & (f_0(\sigma_1 p_1) + \frac{1 - \sigma_1}{2})(f_0(\sigma_2 p_2) + \frac{1 - \sigma_2}{2})(f_0(\sigma_3 p_{12}) + \frac{1 + \sigma_3}{2}) \\ & - (f_0(\sigma_1 p_1) + \frac{1 + \sigma_1}{2})(f_0(\sigma_2 p_2) + \frac{1 + \sigma_2}{2})(f_0(\sigma_3 p_{12}) + \frac{1 - \sigma_3}{2}) \\ & = \sigma_3 f_0(\sigma_1 p_1) f_0(\sigma_2 p_2) - \sigma_1 f_0(\sigma_2 p_2) f_0(\sigma_3 p_{12}) - \sigma_2 f_0(\sigma_1 p_1) f_0(\sigma_3 p_{12}) \\ & + (\sigma_3 - \sigma_2) f_0(\sigma_1 p_1) + (\sigma_3 - \sigma_1) f_0(\sigma_2 p_2) - (\sigma_1 + \sigma_2) f_0(\sigma_3 p_{12}). \end{aligned} \quad (5.105)$$

Observe that there is a global coefficient $\hat{v}(p_2)$. After evaluating all δ coming from contractions between p and k momenta, we want to verify integrability w.r.t. $dp_1 dp_2$. We are thus left with verifying integrability for the terms involving $(\sigma_3 - \sigma_1) f_0(\sigma_2 p_2)$. This term occurs only if $\sigma_3 = -\sigma_1$. Another global factor then is

$$V_2(p_1)^{\frac{1-\sigma_1}{2}} V_2(p_{12})^{\frac{1-\sigma_3}{2}} = V_2(p_1)^{\frac{1-\sigma_1}{2}} V_2(p_{12})^{\frac{1+\sigma_1}{2}}, \quad (5.106)$$

where we recall from (5.88) that we only need to consider $\mathcal{H}_{cub}^{(1)}(s_1)$ in the first argument of the commutator. In particular, there is an integrable factor w.r.t. dp_1 . With that, we have the estimate

$$|\text{bol}^{(3)}(f_0)[J](s_2)| \leq C_{\|\hat{v}\|_{w,d}, \|f_0\|_d} \|J\|_{\ell^\infty(\Lambda^*)}. \quad (5.107)$$

As a consequence, we have that

$$\begin{aligned} |\text{err}_{2,d}^{(Bog, Bol)}(t; f[J])| &= \left| \frac{\lambda^5}{N} \int_{\Delta[t, 2]} ds_2 \text{bol}^{(3)}(f_0)[J](s_2) \right| \\ &\leq C_{\|\hat{v}\|_{w,d}, \|f_0\|_d} \|J\|_{\ell^\infty(\Lambda^*)} \frac{\lambda^5 t^2}{N} \\ &\leq C_{\|\hat{v}\|_{w,d}, \|f_0\|_d} \|J\|_{\ell^\infty(\Lambda^*)} T^2 \frac{\lambda}{N}. \end{aligned} \quad (5.108)$$

This concludes the proof of (5.60).

Finally, we compute the dynamics of g . We write

$$-i \int_0^t \frac{ds}{|\Lambda|} [g[J], \mathcal{H}_{quart}(s)] = \frac{\lambda^2}{N} \int_0^t ds \text{abs}_{quart,d}(f_0)[J](s), \quad (5.109)$$

where $\text{abs}_{quart,d}(f_0)[J](s)$ consists of terms of the form

$$\begin{aligned} & (-i)^\ell \lambda^j \int dp dk e^{-is(m_1 \Omega(p) + m_2 \Omega(k))} J(p) \\ & [(1 + f_0(p))(1 + f_0(-p)) - f_0(p)f_0(-p)] \\ & V_1(p)^{2\alpha_1} V_2(p)^{\alpha_2} \hat{v}(p \pm k) V_1(k)^{2\alpha_3} V_2(k)^{\alpha_4} (f_0(k) + \iota), \end{aligned} \quad (5.110)$$

where $\ell \in \mathbb{N}_0$, $\ell \leq 3$, $j \in \mathbb{N}_0$, $j \leq 7$, $m_1, m_2 \in \{0, \pm 2\}$, $\alpha_j \in \{0, 1, 2\}$ and $\iota \in \{0, 1\}$. Here, we already employed the fact that $\hat{v}(0) = 0$. Using symmetry of the integrand w.r.t. $p \leftrightarrow -p$, we can further simplify the expression (5.110) to

$$\begin{aligned} & (-i)^\ell \lambda^j \int dp dk e^{-is(m_1\Omega(p)+m_2\Omega(k))} (1 + f_0(p) + f_0(-p)) J(p) \\ & V_1(p)^{2\alpha_1} V_2(p)^{\alpha_2} \hat{v}(p-k) V_1(k)^{2\alpha_3} V_2(k)^{\alpha_4} (f_0(k) + \iota). \end{aligned} \quad (5.111)$$

Observe that the only terms contributing to (5.109) are of the form $[aa, (a^+)^3a]$ with any permutation of $(a^+)^3a$, which is why the terms contain at least one momentum flip $a_p \rightarrow i\lambda V_2(p) \sin(\Omega(p)s) a_{-p}$. This justifies the extra factor λ on the RHS of (5.109). In the case $\iota = 1$, we need to have at least one annihilation operator left of a creation operator in the second argument of the commutator. This yields an additional factor $V_2(k)$. In particular, the integrand in (5.110) contains $\hat{v}(p \pm k) f_0(k)$ or $\hat{v}(p \pm k) V_2(k)$.

Next, we compute

$$\begin{aligned} & \frac{(-i)^2}{|\Lambda|} \int_{\Delta[t,2]} ds_2 \nu_0([g[J], \mathcal{H}_{cub}(s_1)], \mathcal{H}_{cub}(s_2)) \\ & = -\frac{1}{|\Lambda|} \int_{\Delta[t,2]} ds_2 \left(\overbrace{[[g[J], \mathcal{H}_{cub}(s_1)], \mathcal{H}_{cub}(s_2)]} \right) \end{aligned} \quad (5.112)$$

$$+ \overbrace{[[g[J], \mathcal{H}_{cub}(s_1)], \mathcal{H}_{cub}(s_2)]} \quad (5.113)$$

$$+ \overbrace{[[g[J], \mathcal{H}_{cub}(s_1)], \mathcal{H}_{cub}(s_2)]} \quad (5.114)$$

Similarly to above, we will refer to (5.112) as condensate contraction and to (5.113) as Boltzmann contraction. In addition, we call (5.114) pair absorption contractions.

We start again with the condensate term. We obtain that

$$\begin{aligned} & -\frac{1}{|\Lambda|} \int_{\Delta[t,2]} ds_2 \overbrace{[[g(J), \mathcal{H}_{cub}(s_1)], \mathcal{H}_{cub}(s_2)]} \\ & = -2J(0) \int_{\Delta[t,2]} \frac{ds_2}{|\Lambda|} \left(\overbrace{[a_0, \mathcal{H}_{cub}(s_1)]} \overbrace{[a_0, \mathcal{H}_{cub}(s_2)]} \right) \\ & = J(0) \Phi_t^2 + J(0) \text{err}_{2,d}^{(Bog,Con)}(t; g). \end{aligned} \quad (5.115)$$

where, analogously to (5.84),

$$\begin{aligned} \text{err}_{2,d}^{(Bog,Con)}(t; g) & := -2 \left(-i \int_0^t ds \frac{\nu_0([a_0, \mathcal{H}_{cub}(s)])}{|\Lambda|} \text{Rem}_2(t; \Phi) \right) \\ & \quad - (\text{Rem}_2(t; \Phi))^2. \end{aligned} \quad (5.116)$$

Using analogous estimates as for (5.86), we find that $|\text{err}_{2,d}^{(Bog,Con)}(t; g)|$ satisfies the same bounds as $|\text{err}_{2,d}^{(Bog,Con)}(t; f)|$.

For the Boltzmann contraction, we write

$$-\int_{\Delta[t,2]} \frac{ds_2}{|\Lambda|} \overbrace{[[g[J], \mathcal{H}_{cub}(s_1)], \mathcal{H}_{cub}(s_2)]}$$

$$= \frac{\lambda^2}{N} \int_{\Delta[t,2]} d\mathbf{s}_2 \text{col}_d(f_0)[J](\mathbf{s}_2). \quad (5.117)$$

The expressions in $\text{col}_d(f_0)[J](\mathbf{s}_2)$ are of the form

$$\begin{aligned} & (-i)^{\ell_0} \lambda^{j_0} \int d\mathbf{p}_3 e^{-i \sum_{\ell=1}^2 s_\ell \sum_{k=1}^3 \sigma_{k,\ell} \Omega(p_k)} \delta(p_1 + p_2 - p_3) \\ & J(p_{j_1}) \hat{v}(p_2) \hat{v}(p_{j_2}) \prod_{k=1}^3 V_1(p_k)^{2\alpha_{\ell_k}} V_2(p_k)^{\beta_{\ell_k}} \\ & \left(\prod_{k=1}^3 \left(f_0(\tau_{k,1} p_k) + \frac{1 + \tau_{k,2}}{2} \right) - \prod_{k=1}^3 \left(f_0(-\tau_{k,1} p_k) + \frac{1 - \tau_{k,2}}{2} \right) \right), \end{aligned} \quad (5.118)$$

where $\ell_0 \in \mathbb{N}_0^{\leq 3}$, $j_0 \in \mathbb{N}_0$, $j_0 \leq 12$, $\sigma_{k,\ell}, \tau_{k,\ell} \in \{\pm 1\}$, $j_1, j_2 \in \{1, 2, 3\}$ and $\alpha_{\ell_k}, \beta_{\ell_k} \in \{0, 1, 2\}$. We need to ensure integrability of each of these terms. More precisely, we will show that any term contains a product of at least two of the functions \hat{v} , f_0 , V_1 , and V_2 depending on at least two of the momenta p_1 , p_2 , or p_3 . We have that

$$\int dp J(p) a_p a_{-p} = \int dp J(p) a_{-p} a_p = \int dp J(-p) a_p a_{-p}, \quad (5.119)$$

where we used the CCR followed by substitution. In particular, we may assume without loss of generality that J is even. Then the CCR imply

$$\begin{aligned} [g[J], a_{\sigma_1 p_1}^{(\sigma_1)} a_{\sigma_2 p_2}^{(\sigma_2)} a_{\sigma_3 p_3}^{(-\sigma_3)}] &= \delta_{\sigma_1, 1} J(p_1) a_{-p_1} a_{\sigma_2 p_2}^{(\sigma_2)} a_{\sigma_3 p_3}^{(-\sigma_3)} \\ &+ \delta_{\sigma_2, 1} J(p_2) a_{\sigma_1 p_1}^{(\sigma_1)} a_{-p_2} a_{\sigma_3 p_3}^{(-\sigma_3)} \\ &+ \delta_{\sigma_3, -1} J(p_3) a_{\sigma_1 p_1}^{(\sigma_1)} a_{\sigma_2 p_2}^{(\sigma_2)} a_{p_3}. \end{aligned} \quad (5.120)$$

We will discuss the expressions related to one these three terms in detail; the remaining follow with analogous computations. Consider the Boltzmann contractions of

$$\nu_0 \left([a_{-p_1} a_{\sigma_2 p_2}^{(\sigma_2)} a_{\sigma_3 p_3}^{(-\sigma_3)}, a_{\tau_1 k_1}^{(\tau_1)} a_{\tau_2 k_2}^{(\tau_2)} a_{\tau_3 k_3}^{(-\tau_3)}] \right), \quad (5.121)$$

which is why, again, the order of the operators a and a^+ does not matter. Observe that it is sufficient to have a factor \hat{v} , f_0 , V_1 , or V_2 with momentum p_1 or p_{12} , since, due to (5.118), we always have a coefficient $\hat{v}(p_2)$. The Boltzmann contractions in (5.121) yield a factor

$$\begin{aligned} & (f_0(-p_1) + 1) \left(f_0(\sigma_2 p_2) + \frac{1 - \sigma_2}{2} \right) \left(f_0(\sigma_3 p_{12}) + \frac{1 + \sigma_3}{2} \right) \\ & - f_0(-p_1) \left(f_0(\sigma_2 p_2) + \frac{1 + \sigma_2}{2} \right) \left(f_0(\sigma_3 p_{12}) + \frac{1 - \sigma_3}{2} \right) \\ & = \sigma_3 f_0(-p_1) f_0(\sigma_2 p_2) + f_0(\sigma_2 p_2) f_0(\sigma_3 p_{12}) - \sigma_2 f_0(-p_1) f_0(\sigma_3 p_{12}) \\ & + (\sigma_3 - \sigma_2) f_0(-p_1) + (1 + \sigma_3) f_0(\sigma_2 p_2) + (1 - \sigma_2) f_0(\sigma_3 p_{12}). \end{aligned} \quad (5.122)$$

The only term in (5.122) that does not already involve a factor depending on a momentum other than p_2 , see also (5.118), is $(1 + \sigma_3) f_0(\sigma_2 p_2)$. This term only appears if $\sigma_3 = 1$, which we now want to consider.

Due to momentum flips in p_j , the corresponding term associated with $\mathcal{H}_{cub}^{(2)}(s_1)$ in the first argument of the commutator has a coefficient $V_2(p_1) V_2(p_3)$, yielding the remaining integrability w.r.t. dp_1 .

Thus, let us consider the terms associated with $\mathcal{H}_{cub}^{(2)}(s_1)$. If we contract $a_{\tau_2 k_2}^{(\tau_2)}$ in with $a_{-\sigma_1 p_1}^{(-\sigma_1)}$ or $a_{\sigma_3 p_3}^{(\sigma_3)}$, we obtain a factor $\hat{v}(p_1)$ or $\hat{v}(p_{12})$, yielding integrability w.r.t. dp_1 .

So, it remains to consider the case when $a_{\tau_2 k_2}^{(\tau_2)}$ is contracted with $a_{\sigma_2 p_2}^{(\sigma_2)}$. The remaining contractions yield either $(\tau_1, -\tau_3) = (1, 1)$ or $(-\tau_3, \tau_1) = (1, 1)$. In those cases, we obtain an additional factor $V_2(p_{12})$ or $V_2(p_1)$. This concludes the argument.

We are left with evaluating the pair absorption terms

$$\begin{aligned} & -\frac{1}{|\Lambda|} \int_{\Delta[t,2]} ds_2 [[g[J], \overbrace{\mathcal{H}_{cub}(s_1)}^{\text{---}}, \overbrace{\mathcal{H}_{cub}(s_2)}^{\text{---}}]] \\ & = \frac{\lambda^2}{N} \int_{\Delta[t,2]} ds_2 \text{abs}_{cub,d}(f_0)[J](s_2). \end{aligned} \quad (5.123)$$

$\text{abs}_{cub,d}(f_0)[J](s_2)$ consists of terms of the form

$$\begin{aligned} & (-i)^\ell \lambda^j \int dp dk e^{-is_1 m_1 \Omega(p) - is_2 m_2 \Omega(k)} J(p) (1 + f_0(p) + f_0(-p)) \hat{v}(p) \\ & V_1(p)^{2\alpha_1} V_2(p)^{\alpha_2} \hat{v}(k) V_1(k)^{2\alpha_3} V_2(k)^{\alpha_4} (f_0(k) + \iota), \end{aligned} \quad (5.124)$$

where $\ell \in \mathbb{N}_0$, $\ell \leq 3$, $j \in \mathbb{N}_0$, $j \leq 12$, $m_1, m_2 \in \{0, \pm 2\}$, $\alpha_j \in \{0, 1, 2\}$ and $\iota \in \{0, 1\}$. Here again, we take into account that terms involving $\hat{v}(0) = 0$ vanish. This concludes the proof. \square

Lemma 5.3. *We have the following expansions.*

(1) *Identifying the RHS with its continuous extension, we have that*

$$\begin{aligned} \int_{\Delta[t,2]} ds_2 e^{i(\omega_1 s_1 + \omega_2 s_2)} & = \frac{2}{\omega_2} \left(\frac{\sin^2((\omega_1 + \omega_2)t/2)}{\omega_1 + \omega_2} - \frac{\sin^2(\omega_1 t/2)}{\omega_1} \right) \\ & - \frac{i}{\omega_2} \left(\frac{\sin((\omega_1 + \omega_2)t)}{\omega_1 + \omega_2} - \frac{\sin(\omega_1 t)}{\omega_1} \right) \end{aligned} \quad (5.125)$$

for all $\omega_1, \omega_2 \in \mathbb{R}$.

(2) *The Bogoliubov dispersion Ω in Lemma 3.2 satisfies*

$$\begin{aligned} \Omega & = E + \lambda \hat{v}_{Bog} \\ & = E + \lambda \hat{v} - \lambda^2 \frac{\hat{v}^2}{2E} + \lambda^3 \text{err}_{Bog}, \end{aligned} \quad (5.126)$$

where $\hat{v}_{Bog} := \frac{2\hat{v}}{1 + \sqrt{1 + 2\lambda \frac{\hat{v}}{E}}}$ and err_{Bog} satisfies

$$\|\text{err}_{Bog}\|_\infty \leq C \|\hat{v}\|_{w,d}^3 \quad (5.127)$$

for all $\lambda > 0$.

Proof. For the first part, let $\omega_1, \omega_2, \omega_1 + \omega_2 \neq 0$. Then

$$\begin{aligned} \int_{\Delta[t,2]} ds_2 e^{i(\omega_1 s_1 + \omega_2 s_2)} & = \int_0^t ds \frac{e^{i\omega_1 s}}{i\omega_2} (e^{i\omega_2 s} - 1) \\ & = -\frac{1}{\omega_2} \left(\frac{e^{i(\omega_1 + \omega_2)t} - 1}{\omega_1 + \omega_2} - \frac{e^{i\omega_1 t} - 1}{\omega_1} \right). \end{aligned} \quad (5.128)$$

Using $1 - \cos(x) = 2 \sin(x/2)$, we have shown the first statement.

For the second part, we expand Ω using

$$\begin{aligned}\Omega &= E + \frac{\Omega^2 - E^2}{E + \Omega} \\ &= E + \frac{2\lambda\hat{v}}{1 + \sqrt{1 + 2\lambda\frac{\hat{v}}{E}}}.\end{aligned}\quad (5.129)$$

We Taylor expand

$$\frac{1}{1 + \sqrt{1 + x}} = \frac{1}{2} - \frac{x}{8} + R(x) \quad (5.130)$$

for some $R(x)$. An easy computation yields

$$\frac{|R(x)|}{x^2} \leq C \quad (5.131)$$

for all $x > 0$, and in the limit $x \searrow 0$. This concludes the proof. \square

Remark 5.4 (Talbot effect). *Lemma 5.3 shows that the leading term in $\text{Bol}_d(f_0)[J]$ is a sum of the form*

$$\frac{T}{\lambda^2} \int_{(\Lambda^*)^2} d\mathbf{p}_2 \frac{\sin^2\left(\frac{T(\Delta_{cub}E + \lambda\Delta_{cub}\hat{v}_{Bog})}{2\lambda^2}\right)}{\left(\frac{T(\Delta_{cub}E + \lambda\Delta_{cub}\hat{v}_{Bog})}{2\lambda^2}\right)^2} \Delta_{cub} J(\mathbf{p}_2) H(\mathbf{p}_2) \quad (5.132)$$

for some $H \sim 1$ with \hat{v}_{Bog} as defined in Lemma 5.3. Thus, as a function of T , the modulus of this sum oscillates between 0 and $O(\lambda^{-2})$. The size of these oscillations is dependent on the interaction profile \hat{v} , the lattice Λ , and the chosen sequence $\lambda = \lambda(N)$. A similar phenomenon occurs for $\text{Bol}_d^{(j)}$, which thereby also oscillate in modulus between 0 and $O(\lambda^{j-2})$, depending on T , and we also observe it for col_d , $\text{abs}_{quart,d}$, and $\text{abs}_{cub,d}$. Therefore, $\text{Bol}_d^{(j)}$ can dominate Bol_d , depending on whether $\frac{T}{4\pi\lambda^2} \Delta_{cub} \Omega(\mathbf{p}_2)$ lies in a small vicinity of $\frac{1}{4} + \mathbb{Z}$ for some $\mathbf{p}_2 \in (\Lambda^*)^2$. This is reminiscent of the Talbot effect, see [128, 190]. As heuristically explained above, $\text{Bol}_d^{(j)}$ are negligible in the case of large system length $L \sim \lambda^{-2-}$. Next, we will present a proof of this fact.

5.2. Continuum approximation $\Lambda^* \rightarrow \mathbb{R}^3$. In this section, we will write out the summation over the lattice Λ^* explicitly. In contrast, all integrals will be understood as integrals over \mathbb{R}^d in the respective dimension d .

Lemma 5.5. *Let $F_1(\mathbf{p}_2), F_2(\mathbf{p}_2) \in \{\pm\Omega(p_1) \pm \Omega(p_2) \pm \Omega(p_1 + p_2), 0\}$, $H \in C_c^\infty(\mathbb{R}^6)$, $\chi \in C_c^\infty(\mathbb{R}^3)$, and $\tau_1, \tau_2 \in \mathbb{R}$. Let $\langle x \rangle := (1 + x^2)^{\frac{1}{2}}$. Abbreviate*

$$\begin{aligned}\text{err}_{disc}^{(1)} &:= \lambda^2 \int_0^{T\lambda^{-2}} ds \left(\frac{1}{|\Lambda|^2} \sum_{\mathbf{p}_2 \in (\Lambda^*)^2} e^{i(\tau_1\Omega(p_1) + \tau_2\Omega(p_2))s} H(\mathbf{p}_2) \right. \\ &\quad \left. - \frac{1}{(2\pi)^6} \int d\mathbf{p}_2 e^{i(\tau_1\Omega(p_1) + \tau_2\Omega(p_2))s} H(\mathbf{p}_2) \right), \\ \text{err}_{disc}^{(2)} &:= \lambda^2 \int_{\Delta[T\lambda^{-2}, 2]} ds_2 \left(\frac{1}{|\Lambda|^2} \sum_{\mathbf{p}_2 \in (\Lambda^*)^2} e^{iF_1(\mathbf{p}_2)s_1 + iF_2(\mathbf{p}_2)s_2} H(\mathbf{p}_2) \right. \\ &\quad \left. - \frac{1}{(2\pi)^6} \int d\mathbf{p}_2 e^{iF_1(\mathbf{p}_2)s_1 + iF_2(\mathbf{p}_2)s_2} H(\mathbf{p}_2) \right).\end{aligned}\quad (5.133)$$

Then we have for any $r > 6$ and $\lambda > 0$ the following.

$$(1) \quad \left| \frac{1}{|\Lambda|} \sum_{p \in \Lambda^*} \chi(p) - \frac{1}{(2\pi)^3} \int_{\mathbb{R}^3} dp \chi(p) \right| \leq C_r \frac{\|\nabla|\frac{r}{2}\chi\|_1}{|\Lambda|^{\frac{r}{6}}}, \quad (5.134)$$

$$(2) \quad |\text{err}_{disc}^{(1)}| \leq C_{\|\hat{v}\|_2(\lfloor \frac{r}{2} \rfloor + 1), w, c, r} \frac{\langle T \rangle^{r+1}}{\lambda^{2r} |\Lambda|^{\frac{r}{3}}} \sum_{n=0}^{2(\lfloor \frac{r}{2} \rfloor + 1)} \left\| \langle |\mathbf{p}_2| \rangle^n D^{2(\lfloor \frac{r}{2} \rfloor + 1) - n} H \right\|_1, \quad (5.135)$$

(3) if $F_1 = F_2 \equiv 0$,

$$|\text{err}_{disc}^{(2)}| \leq C_{\|\hat{v}\|_2(\lfloor \frac{r}{2} \rfloor + 1), w, c, r} \frac{T^2}{\lambda^2 |\Lambda|^{\frac{r}{3}}} \|\nabla|^r H\|_1, \quad (5.136)$$

(4) if $(F_1, F_2) \neq (0, 0)$,

$$|\text{err}_{disc}^{(2)}| \leq C_{\|\hat{v}\|_2(\lfloor \frac{r}{2} \rfloor + 1), w, c, r} \frac{\langle T \rangle^{r+2}}{\lambda^{2r+2} |\Lambda|^{\frac{r}{3}}} \sum_{n=0}^{2(\lfloor \frac{r}{2} \rfloor + 1)} \left\| \langle |\mathbf{p}_2| \rangle^n D^{2(\lfloor \frac{r}{2} \rfloor + 1) - n} H \right\|_1 \quad (5.137)$$

Proof. Let

$$\tau := \frac{2\pi}{|\Lambda|^{\frac{1}{3}}}, \quad (5.138)$$

so that

$$\int_{\Lambda^*} dp \tilde{H}(p) = \frac{\tau^3}{(2\pi)^3} \sum_{p \in \Lambda^*} \tilde{H}(p). \quad (5.139)$$

Denote

$$G_\lambda(\mathbf{p}_2) := \lambda^2 \int_{\Delta[T\lambda^{-2}, 2]} d\mathbf{s}_2 e^{iF_1(\mathbf{p}_2)s_1 + iF_2(\mathbf{p}_2)s_2}. \quad (5.140)$$

Poisson summation implies

$$\begin{aligned} \tau^6 \sum_{\mathbf{p}_2 \in (\Lambda^*)^2} G_\lambda(\mathbf{p}_2) H(\mathbf{p}_2) &= \tau^6 \sum_{\mathbf{P}_2 \in \mathbb{Z}^6} G_\lambda(\tau \mathbf{P}_2) H(\tau \mathbf{P}_2) \\ &= \sum_{\mathbf{X}_2 \in \mathbb{Z}^6} \tau^6 \int_{\mathbb{R}^6} d\mathbf{P}_2 e^{2\pi i \mathbf{P}_2 \cdot \mathbf{X}_2} G_\lambda(\tau \mathbf{P}_2) H(\tau \mathbf{P}_2) \\ &= \sum_{\mathbf{X}_2 \in \mathbb{Z}^6} \int_{\mathbb{R}^6} d\mathbf{p}_2 e^{2\pi i \mathbf{p}_2 \cdot \mathbf{X}_2 / \tau} G_\lambda(\mathbf{p}_2) H(\mathbf{p}_2). \end{aligned} \quad (5.141)$$

As a consequence of (5.141), we obtain that

$$\text{err}_{disc}^{(2)} = \frac{1}{(2\pi)^6} \sum_{\mathbf{X}_2 \in \mathbb{Z}^6 \setminus \{0\}} \int_{\mathbb{R}^6} d\mathbf{p}_2 e^{2\pi i \mathbf{p}_2 \cdot \mathbf{X}_2 / \tau} G_\lambda(\mathbf{p}_2) H(\mathbf{p}_2). \quad (5.142)$$

Next, we have that

$$\begin{aligned} &|\mathbf{X}_2|^r \int_{\mathbb{R}^6} d\mathbf{p}_2 e^{2\pi i \mathbf{p}_2 \cdot \mathbf{X}_2 / \tau} G_\lambda(\mathbf{p}_2) H(\mathbf{p}_2) \\ &= \frac{\tau^r}{(2\pi)^r} \int_{\mathbb{R}^6} d\mathbf{p}_2 e^{2\pi i \mathbf{p}_2 \cdot \mathbf{X}_2 / \tau} |\nabla_{\mathbf{p}_2}|^r (G_\lambda(\mathbf{p}_2) H(\mathbf{p}_2)). \end{aligned} \quad (5.143)$$

In particular, we have that

$$\begin{aligned} |\text{err}_{disc}^{(2)}| &\leq C_r \tau^r \sum_{\mathbf{x}_2 \in \mathbb{Z}^6 \setminus \{0\}} \frac{1}{|\mathbf{x}_2|^r} \|\nabla|^r (G_\lambda H)\|_1 \\ &\leq \frac{C_r}{|\Lambda|^{\frac{r}{3}}} \|\nabla|^r (G_\lambda H)\|_1 \end{aligned} \quad (5.144)$$

due to $r > 6$. With analogous steps, we obtain

$$\begin{aligned} \left| \tau^3 \sum_{p \in \Lambda^*} \chi(p) - \int_{\mathbb{R}^3} dp \chi(p) \right| &= \frac{\tau^{\frac{r}{2}}}{(2\pi)^{\frac{r}{2}}} \left| \sum_{X \in \mathbb{Z}^3 \setminus \{0\}} \frac{1}{|X|^{\frac{r}{2}}} \int_{\mathbb{R}^3} dp e^{2\pi i p \cdot X / \tau} |\nabla|^{\frac{r}{2}} \chi(p) \right| \\ &\leq \frac{C_r}{|\Lambda|^{\frac{r}{6}}} \|\nabla|^{\frac{r}{2}} \chi\|_1. \end{aligned} \quad (5.145)$$

Interpolation implies

$$\|\nabla|^r (G_\lambda H)\|_1 \leq C \|(-\Delta)^{\lfloor \frac{r}{2} \rfloor + 1} (G_\lambda H)\|_1^{\frac{r}{2}} \|G_\lambda H\|_1^{\frac{1 + \lfloor \frac{r}{2} \rfloor - \frac{r}{2}}{\lfloor \frac{r}{2} \rfloor + 1}}, \quad (5.146)$$

where $t_0 > 0$ will be fixed below. The Leibniz rule implies for $k \in \mathbb{N}_0$ that

$$|(-\Delta)^k G_\lambda H| \leq C_k \sum_{n=0}^{2k} |D^n G_\lambda| |D^{2k-n} H|. \quad (5.147)$$

We start with the case $F_1 = F_2 \equiv 0$. Lemma 5.3 yields $G_\lambda = C \frac{T^2}{2} \lambda^{-2}$, which, employing (5.144), implies

$$\begin{aligned} |\text{err}_{disc}^{(2)}| &\leq C_r T^2 \lambda^{-2} \tau^r \|\nabla|^r H\|_1 \\ &\leq C_r \frac{T^2}{\lambda^2 |\Lambda|^{\frac{r}{3}}} \|\nabla|^r H\|_1. \end{aligned} \quad (5.148)$$

Next, let $F_1 \equiv 0 \wedge F_2 \neq 0$, $F_1 \neq 0 \wedge F_2 \equiv 0$, or $F_1 = -F_2 \neq 0$. Let $F_j \neq 0$ for some $j \in \{1, 2, 3\}$. Then, by Lemma 5.3, we obtain that

$$G_\lambda(\mathbf{p}_2) = \frac{T^2}{\lambda^2} G(F_j(\mathbf{p}_2) T / \lambda^2) \quad (5.149)$$

for some continuous function $G \in C_b^\infty(\mathbb{R})$. Moreover, the Faà di Bruno formula, see [295], implies

$$|D^n G(F_j T / \lambda^2)| \leq C_n \sum_{\substack{\mathbf{r}_n \in \\ R(n)}} |G^{(S(\mathbf{r}_n))}| (F_j T / \lambda^2) \prod_{\ell=1}^n \left[\frac{T}{\lambda^2} |D^\ell F_j| \right]^{r_\ell}, \quad (5.150)$$

where

$$R(n) := \left\{ \mathbf{r}_n \in \mathbb{N}_0^n \mid \sum_{\ell=1}^n \ell r_\ell = n \right\}, \quad (5.151)$$

$$S(\mathbf{r}_n) := \sum_{\ell=1}^n r_\ell. \quad (5.152)$$

Observe that due to Lemma 5.3, we have that

$$|\nabla F_j| \leq C_{\|\hat{v}\|_2, \lfloor \frac{r}{2} \rfloor + 1, w, c} \langle |\mathbf{p}_2| \rangle, \quad (5.153)$$

$$|D^\ell F_j| \leq C_{\|\hat{v}\|_2, \lfloor \frac{r}{2} \rfloor + 1, w, c} \quad (5.154)$$

for all $\lambda \in (0, 1)$, and all $\ell \geq 2$. Then (5.150) together with (5.153), (5.154), and the fact that $1 \leq S(\mathbf{r}_n) \leq n$ imply

$$\begin{aligned} |D^n G(F_j T / \lambda^2)| &\leq C_n \frac{T \langle T \rangle^{n-1} \langle |\nabla F_j| \rangle^n}{\lambda^{2n}} \left(\sum_{\ell=1}^n |G^{(\ell)}| \right) (F_j T / \lambda^2) \\ &\leq C_{\|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c}, n} \frac{T \langle T \rangle^{n-1} \langle |\mathbf{p}_2| \rangle^n}{\lambda^{2n}} \left(\sum_{\ell=1}^n |G^{(\ell)}| \right) (F_j T / \lambda^2) \end{aligned} \quad (5.155)$$

for all $\lambda \in (0, 1)$. Collecting (5.147), 5.155, we have that

$$\begin{aligned} &\|(-\Delta)^k (G_\lambda H)\|_1 \\ &\leq C_{\|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c}, r} \sum_{n=0}^{2k} \frac{T^3 \langle T \rangle^{n-1}}{\lambda^{2n+2}} \left\| \left(\sum_{\ell=1}^n |G^{(\ell)}| \right) (F_j T / \lambda^2) \langle |\mathbf{p}_2| \rangle^n D^{2k-n} H \right\|_1 \\ &\leq C_{\|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c}, r} \frac{\langle T \rangle^{2k+2}}{\lambda^{4k+2}} \left\| \left(\sum_{\ell=0}^{2k} |G^{(\ell)}| \right) \right\|_\infty \sum_{n=0}^{2k} \left\| \langle |\mathbf{p}_2| \rangle^n D^{2k-n} H \right\|_1 \end{aligned} \quad (5.156)$$

for all $k \in \{0, 1, 2, \dots, \lfloor \frac{r}{2} \rfloor + 1\}$. Then (5.146) and (5.156) yield

$$\begin{aligned} &\|\nabla^r (G_\lambda H)\|_1 \\ &\leq C_{\|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c}, r} \frac{\langle T \rangle^{r+2}}{\lambda^{2r+2}} \sum_{n=0}^{2(\lfloor \frac{r}{2} \rfloor + 1)} \left\| \langle |\mathbf{p}_2| \rangle^n D^{2(\lfloor \frac{r}{2} \rfloor + 1) - n} H \right\|_1, \end{aligned} \quad (5.157)$$

where we also used the fact that $\|H\|_1 \leq \|\langle |\mathbf{p}_2| \rangle^{2(\lfloor \frac{r}{2} \rfloor + 1)} H\|_1$. (5.144) and (5.157) yield

$$|\text{err}_{disc}^{(2)}| \leq C_{\|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c}, r} \frac{\langle T \rangle^{r+2}}{\lambda^{2r+2} |\Lambda|^{\frac{r}{3}}} \sum_{n=0}^{2(\lfloor \frac{r}{2} \rfloor + 1)} \left\| \langle |\mathbf{p}_2| \rangle^n D^{2(\lfloor \frac{r}{2} \rfloor + 1) - n} H \right\|_1 \quad (5.158)$$

Now, let $F_j \neq 0$ for all $j \in \{1, 2, 3\}$. In this case, Lemma 5.3 implies

$$G_\lambda(\mathbf{p}_2) = \frac{T^2}{\lambda^2} G\left(\frac{T}{\lambda^2}(F_1(\mathbf{p}_2), F_2(\mathbf{p}_2))\right) \quad (5.159)$$

for some smooth function $G \in C_b^\infty(\mathbb{R}^2)$. The Faà di Bruno formula implies

$$\begin{aligned} |D^n G\left(\frac{T}{\lambda^2}(F_1(\mathbf{p}_2), F_2(\mathbf{p}_2))\right)| &\leq C_n \sum_{\substack{\mathbf{r}_n \in \\ R(n)}} |D^{S(\mathbf{r}_n)} G|\left(\frac{T}{\lambda^2}(F_1(\mathbf{p}_2), F_2(\mathbf{p}_2))\right) \\ &\quad \prod_{\ell=1}^n \left[\frac{T}{\lambda^2} \sum_{j=1}^2 |D^\ell F_j| \right]^{r_\ell} \\ &\leq C_n \frac{T \langle T \rangle^{n-1} \langle \sum_{j=1}^2 |\nabla F_j| \rangle^n}{\lambda^{2n}} \\ &\quad \left(\sum_{\ell=1}^n |D^\ell G| \right) \left(\frac{T}{\lambda^2}(F_1(\mathbf{p}_2), F_2(\mathbf{p}_2)) \right). \end{aligned} \quad (5.160)$$

Collecting (5.147), (5.153), (5.154), and 5.160, we have that

$$\|(-\Delta)^k (G_\lambda H)\|_1$$

$$\begin{aligned}
 &\leq C_{\|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c, r}} \sum_{n=0}^{2k} \frac{T^3 \langle T \rangle^{n-1}}{\lambda^{2n+2}} \\
 &\quad \left\| \left(\sum_{\ell=1}^n |D^\ell G| \right) \left(\frac{T}{\lambda^2} (F_1, F_2) \right) \left\langle \sum_{j=1}^2 |\nabla F_j| \right\rangle^n D^{2k-n} H \right\|_1 \\
 &\leq C_{\|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c, r}} \frac{\langle T \rangle^{2k+2}}{\lambda^{4k+2}} \left\| \left(\sum_{\ell=0}^{2k} |D^\ell G| \right) \right\|_\infty \sum_{n=0}^{2k} \left\| \langle |\mathbf{p}_2| \rangle^n D^{2k-n} H \right\|_1 \quad (5.161)
 \end{aligned}$$

for all $k \in \{0, 1, 2, \dots, \lfloor \frac{r}{2} \rfloor + 1\}$. As a consequence of (5.146), we thus obtain

$$\|\nabla|^r(G_\lambda H)\|_1 \leq C_{\|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c, r}} \frac{\langle T \rangle^{r+2}}{\lambda^{2r+2}} \sum_{n=0}^{2(\lfloor \frac{r}{2} \rfloor + 1)} \left\| \langle |\mathbf{p}_2| \rangle^n D^{2(\lfloor \frac{r}{2} \rfloor + 1) - n} H \right\|_1 \quad (5.162)$$

(5.144) and (5.162) imply

$$|\text{err}_{disc}^{(2)}| \leq C_{\|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c, r}} \frac{\langle T \rangle^{r+2}}{\lambda^{2r+2} |\Lambda|^{\frac{r}{3}}} \sum_{n=0}^{2(\lfloor \frac{r}{2} \rfloor + 1)} \left\| \langle |\mathbf{p}_2| \rangle^n D^{2(\lfloor \frac{r}{2} \rfloor + 1) - n} H \right\|_1 \quad (5.163)$$

Finally, let

$$\mathring{G}_\lambda(\mathbf{p}_2) := \lambda^2 \int_0^{T\lambda^{-2}} ds e^{i(\tau_1 \Omega(p_1) + \tau_2 \Omega(p_2))s}. \quad (5.164)$$

A simple computation shows

$$\mathring{G}_\lambda(\mathbf{p}_2) = T \mathring{G}((\tau_1 \Omega(p_1) + \tau_2 \Omega(p_2)) \frac{T}{\lambda^2}) \quad (5.165)$$

for some smooth $\mathring{G} \in C_b^\infty(\mathbb{R})$. If $\tau_1 = \tau_2 = 0$, we have $\mathring{G} \equiv C$. If $(\tau_1, \tau_2) \neq (0, 0)$, a computation analogous to (5.155) yields

$$\begin{aligned}
 &|D^n \mathring{G}((\tau_1 \Omega(p_1) + \tau_2 \Omega(p_2)) T / \lambda^2)| \\
 &\leq C_{\|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c, n}} \frac{T \langle T \rangle^{n-1} \langle |\mathbf{p}_2| \rangle^n}{\lambda^{2n}} \left(\sum_{\ell=1}^n |\mathring{G}^{(\ell)}| \right) ((\tau_1 \Omega(p_1) + \tau_2 \Omega(p_2)) T / \lambda^2) \quad (5.166)
 \end{aligned}$$

for all $n \in \{0, 1, 2, \dots, 2(\lfloor \frac{r}{2} \rfloor + 1)\}$. In particular, with analogous steps that led to (5.158), we obtain

$$|\text{err}_{disc}^{(1)}| \leq C_{\|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c, r}} \frac{\langle T \rangle^{r+1}}{\lambda^{2r} |\Lambda|^{\frac{r}{3}}} \sum_{n=0}^{2(\lfloor \frac{r}{2} \rfloor + 1)} \left\| \langle |\mathbf{p}_2| \rangle^n D^{2(\lfloor \frac{r}{2} \rfloor + 1) - n} H \right\|_1. \quad (5.167)$$

This concludes the proof. \square

Remark 5.6. Observe that Lemma 5.5 implies that

$$\begin{aligned}
 \int_{\Lambda^*} dp f_0(p) &\leq \|f_0\|_1 + C_r \frac{\|\nabla|^{\frac{r}{2}} f_0\|_1}{|\Lambda|^{\frac{r}{6}}} \\
 &\leq C_r \|f_0\|_{2(\lfloor \frac{r}{2} \rfloor + 1), c} \quad (5.168)
 \end{aligned}$$

for any $r > 6$ as in Theorem 2.2, where C_r is independent of $|\Lambda| \geq 1$. In particular, we have that

$$\|f_0\|_d \leq C_r \|f_0\|_{2(\lfloor \frac{r}{2} \rfloor + 1), c}. \quad (5.169)$$

This allows us to use the previous estimates to prove Theorem 2.2 as well. Analogously, we have that

$$\|\hat{v}\|_{w,d} \leq C_r \|\hat{v}\|_{2(\lfloor \frac{T}{2} \rfloor + 1), w, c}. \quad (5.170)$$

in the assumption of Theorem 2.1, where, again C_r is independent of $|\Lambda| \geq 1$.

Likewise, we have that

$$\|J\|_{2 \cap \infty, d} \leq C_r \|J\|_{2(\lfloor \frac{T}{2} \rfloor + 1), c}. \quad (5.171)$$

For the next statement, define the continuous analogues Con_c , Bol_c , etc. of Con_d , Bol_d , etc. in (5.49)–(5.56), and (5.63), (5.65) by replacing the lattice sums \int_{Λ^*} over Λ^* in the sense of (1.51) by Lebesgue integrals $\frac{1}{(2\pi)^3} \int_{\mathbb{R}^3}$ over \mathbb{R}^3 .

Proposition 5.7. *Let $T > 0$, $J \in L^\infty(\mathbb{R}^3; \mathbb{R})$ and $r > 6$. Then the following holds for all $\lambda > 0$ small enough, dependent on $\|\hat{v}\|_{2(\lfloor \frac{T}{2} \rfloor + 1), w, c}$.*

$$(1) \quad -i \int_0^{T\lambda^{-2}} ds \frac{\nu_0([a_0, \mathcal{H}_{cub}(s)])}{|\Lambda|} = \frac{1}{N^{\frac{1}{2}} \lambda} \text{Con}_c(f_0)(T; \lambda) + \text{err}_{1,c}^{(Bog)}\left(\frac{T}{\lambda^2}; \Phi\right), \quad (5.172)$$

where

$$|\text{err}_{1,c}^{(Bog)}\left(\frac{T}{\lambda^2}; \Phi\right)| \leq \frac{C_{\|\hat{v}\|_{2(\lfloor \frac{T}{2} \rfloor + 1), w, c}, \|f_0\|_{2(\lfloor \frac{T}{2} \rfloor + 1), c}, r} T}{N^{\frac{1}{2}} \lambda} \left(\lambda + \frac{1}{|\Lambda|^{\frac{r}{6}}} \right), \quad (5.173)$$

$$(2) \quad \begin{aligned} & - \int_{\Delta[t, 2]} d\mathbf{s}_2 \frac{\nu_0([f[J], \mathcal{H}_{cub}(s_1)], \mathcal{H}_{cub}(s_2)])}{|\Lambda|} \\ &= \frac{1}{N} \text{Bol}_c(f_0)[J](T; \lambda) + J(0) |\Phi_{\frac{T}{\lambda^2}}|^2 \\ &+ \text{err}_{2,c}^{(Bog, Bol)}\left(\frac{T}{\lambda^2}; f[J]\right) + J(0) \text{err}_{2,c}^{(Bog, Con)}\left(\frac{T}{\lambda^2}; f\right), \end{aligned} \quad (5.174)$$

where

$$\begin{aligned} |\text{err}_{2,c}^{(Bog, Bol)}\left(\frac{T}{\lambda^2}; f[J]\right)| &\leq \frac{C_{\|\hat{v}\|_{2(\lfloor \frac{T}{2} \rfloor + 1), w, c}, \|f_0\|_{2(\lfloor \frac{T}{2} \rfloor + 1), c}, r} \langle T \rangle^{r+2} \|J\|_{W^2[\lfloor \frac{T}{2} \rfloor + 2, \infty}}{N} \\ &\quad \left(\frac{1}{\lambda^{2(r+1)} |\Lambda|^{\frac{r}{3}}} + \lambda |\log(\lambda)| \right), \\ |\text{err}_{2,c}^{(Bog, Con)}\left(\frac{T}{\lambda^2}; f[J]\right)| &\leq C_{\|\hat{v}\|_{2(\lfloor \frac{T}{2} \rfloor + 1), w, c}, \|f_0\|_{2(\lfloor \frac{T}{2} \rfloor + 1), c}, r} e^{C_r \|\hat{v}\|_{2(\lfloor \frac{T}{2} \rfloor + 1), w, c} |\Lambda| / \lambda T} \\ &\quad \frac{\langle T \rangle^4 |\Lambda|^{\frac{3}{2}}}{\lambda^3 N^{\frac{3}{2}}} \left(1 + \frac{|\Lambda|}{N} \right)^2 \left(1 + \frac{|\Lambda|^{\frac{3}{2}}}{\lambda N^{\frac{1}{2}}} \right), \end{aligned} \quad (5.175)$$

$$(3) \quad \begin{aligned} & -i \int_0^{T\lambda^{-2}} ds \frac{\nu_0(g[J], \mathcal{H}_{quart}(s))}{|\Lambda|} \\ &= \frac{1}{N} \int_0^T dS \text{abs}_{quart, c}(f_0)[J](S/\lambda^2) + \text{err}_{1,c}^{(dis)}\left(\frac{T}{\lambda^2}; g[J]\right) \end{aligned} \quad (5.176)$$

with

$$|\text{err}_{1,c}^{(dis)}(\frac{T}{\lambda^2}; g[J])| \leq \frac{C_{\|\hat{v}\|_{2(\lfloor \frac{T}{2} \rfloor + 1), w, c}, \|f_0\|_{2(\lfloor \frac{T}{2} \rfloor + 1), c}, r} \langle T \rangle^{r+1} \|J\|_{W^2[\frac{T}{2}] + 2, \infty}}{N\lambda^{2r} |\Lambda|^{\frac{r}{3}}} \quad (5.177)$$

(4)

$$\begin{aligned} & - \int_{\Delta[T\lambda^{-2}, 2]} d\mathbf{s}_2 \frac{\nu_0([\![g[J], \mathcal{H}_{cub}(s_1)], \mathcal{H}_{cub}(s_2)]])}{|\Lambda|} \\ & = \frac{1}{N\lambda^2} \int_{\Delta[T, 2]} d\mathbf{S}_2 (\text{col}_c(f_0)[J](\mathbf{S}_2/\lambda^2) + \text{abs}_{cub,c}(f_0)[J](\mathbf{S}_2/\lambda^2)) \\ & + \text{err}_{2,c}^{(dis)}(\frac{T}{\lambda^2}; g[J]) + J(0) (\Phi_{T\lambda^{-2}})^2 + J(0) \text{err}_{2,c}^{(Bog, Con)}(\frac{T}{\lambda^2}; g). \end{aligned} \quad (5.178)$$

where

$$|\text{err}_{2,c}^{(dis)}(\frac{T}{\lambda^2}; g[J])| \leq \frac{C_{\|\hat{v}\|_{2(\lfloor \frac{T}{2} \rfloor + 1), w, c}, \|f_0\|_{2(\lfloor \frac{T}{2} \rfloor + 1), c}, r} \langle T \rangle^{r+2} \|J\|_{W^2[\frac{T}{2}] + 2, \infty}}{N\lambda^{2(r+1)} |\Lambda|^{\frac{r}{3}}} \quad (5.179)$$

and $\text{err}_{2,c}^{(Bog, Con)}(\frac{T}{\lambda^2}; g)$ satisfies the same bound as $\text{err}_{2,c}^{(Bog, Con)}(\frac{T}{\lambda^2}; f)$.

Proof. Recall from Proposition 5.2 that

$$\begin{aligned} & -i \int_0^{T\lambda^{-2}} ds \frac{\nu_0([a_0, \mathcal{H}_{cub}(s)])}{|\Lambda|} \\ & = \frac{1}{N^{\frac{1}{2}}\lambda} \text{Con}_d(f_0)(T; \lambda) + \text{err}_{1,d}^{(Bog)}(\frac{T}{\lambda^2}; \Phi) \end{aligned} \quad (5.180)$$

with

$$\begin{aligned} |\text{err}_{1,d}^{(Bog)}(\frac{T}{\lambda^2}; \Phi)| & \leq \frac{C_{\|\hat{v}\|_{w,d}, \|f_0\|_d} T}{N^{\frac{1}{2}}} \\ & \leq \frac{C_{\|\hat{v}\|_{2(\lfloor \frac{T}{2} \rfloor + 1), w, c}, \|f_0\|_{2(\lfloor \frac{T}{2} \rfloor + 1), c}, r} T}{N^{\frac{1}{2}}} \end{aligned} \quad (5.181)$$

due to Remark 5.6. Let

$$\text{err}_{1,c}^{(dis)}(\frac{T}{\lambda^2}; \Phi) := \frac{1}{N^{\frac{1}{2}}\lambda} (\text{Con}_d(f_0)(T; \lambda) - \text{Con}_c(f_0)(T; \lambda)). \quad (5.182)$$

Then Lemma 5.5 implies

$$\begin{aligned} |\text{err}_{1,c}^{(dis)}(\frac{T}{\lambda^2}; \Phi)| & \leq \frac{T}{N^{\frac{1}{2}}\lambda} \left| \frac{(2\pi)^3}{|\Lambda|} \sum_{p \in \Lambda^*} \hat{v}(p) f_0(p) - \int_{\mathbb{R}^3} dp \hat{v}(p) f_0(p) \right| \\ & \leq C_r \frac{T}{N^{\frac{1}{2}}\lambda |\Lambda|^{\frac{r}{6}}} \|\nabla|^{\frac{r}{2}} \hat{v} f_0\|_1 \\ & \leq \frac{C_{\|\hat{v}\|_{2(\lfloor \frac{T}{2} \rfloor + 1), w, c}, \|f_0\|_{2(\lfloor \frac{T}{2} \rfloor + 1), c}, r} T}{N^{\frac{1}{2}}\lambda |\Lambda|^{\frac{r}{6}}}. \end{aligned} \quad (5.183)$$

Here, we used interpolation as in (5.146), together with the Leibniz rule. Applying (5.181), (5.183) implies

$$|\text{err}_{1,c}^{(Bog)}(\frac{T}{\lambda^2}; \Phi)| \leq |\text{err}_{1,d}^{(Bog)}(\frac{T}{\lambda^2}; \Phi)| + |\text{err}_{1,c}^{(dis)}(\frac{T}{\lambda^2}; \Phi)|$$

$$\leq \frac{C \|\hat{v}\|_{2(\lfloor \frac{T}{2} \rfloor + 1), w, c} \|f_0\|_{2(\lfloor \frac{T}{2} \rfloor + 1), c} r^T}{N^{\frac{1}{2}} \lambda} \left(\lambda + \frac{1}{|\Lambda|^{\frac{r}{6}}} \right). \quad (5.184)$$

Next, Proposition 5.2 together with Remark 5.6 yield that

$$\begin{aligned} & - \int_{\Delta[T\lambda^{-2}, 2]} ds_2 \frac{\nu_0([f[J], \mathcal{H}_{cub}(s_1)], \mathcal{H}_{cub}(s_2))}{|\Lambda|} \\ &= \frac{1}{N} (\text{Bol}_d(f_0)[J](T; \lambda) + \lambda \text{Bol}_d^{(1)}(f_0)[J](T; \lambda) + \lambda^2 \text{Bol}_d^{(2)}(f_0)[J](T; \lambda)) \\ & \quad + J(0) |\Phi_{\frac{T}{\lambda^2}}|^2 + \text{err}_{2,d}^{(Bog, Bol)}\left(\frac{T}{\lambda^2}; f[J]\right) + J(0) \text{err}_{2,d}^{(Bog, Con)}\left(\frac{T}{\lambda^2}; f\right), \end{aligned} \quad (5.185)$$

where

$$\begin{aligned} |\text{err}_{2,d}^{(Bog, Bol)}\left(\frac{T}{\lambda^2}; f[J]\right)| &\leq C \|\hat{v}\|_{2(\lfloor \frac{T}{2} \rfloor + 1), w, c} \|f_0\|_{2(\lfloor \frac{T}{2} \rfloor + 1), c} r^T \|J\|_{\ell^\infty(\Lambda^*)} \frac{\lambda}{N}, \quad (5.186) \\ |\text{err}_{2,d}^{(Bog, Con)}(t; f)| &\leq C \|\hat{v}\|_{2(\lfloor \frac{T}{2} \rfloor + 1), w, c} \|f_0\|_{2(\lfloor \frac{T}{2} \rfloor + 1), c} r^T e^{C_r \|\hat{v}\|_{2(\lfloor \frac{T}{2} \rfloor + 1), w, c} |\Lambda| \lambda T} \\ & \quad \frac{\langle T \rangle^4 |\Lambda|^{\frac{3}{2}}}{\lambda^3 N^{\frac{3}{2}}} \left(1 + \frac{|\Lambda|}{N}\right)^2 \left(1 + \frac{|\Lambda|^{\frac{3}{2}}}{\lambda N^{\frac{1}{2}}}\right). \end{aligned} \quad (5.187)$$

Let

$$\text{err}_{2,0}^{(dis)}\left(\frac{T}{\lambda^2}; f[J]\right) := \frac{1}{N} (\text{Bol}_d(f_0)[J](T; \lambda) - \text{Bol}_c(f_0)[J](T; \lambda)), \quad (5.188)$$

$$\text{err}_{2,1}^{(dis)}\left(\frac{T}{\lambda^2}; f[J]\right) := \frac{\lambda}{N} (\text{Bol}_d^{(1)}(f_0)[J](T; \lambda) - \text{Bol}_c^{(1)}(f_0)[J](T; \lambda)), \quad (5.189)$$

$$\text{err}_{2,2}^{(dis)}\left(\frac{T}{\lambda^2}; f[J]\right) := \frac{\lambda^2}{N} (\text{Bol}_d^{(2)}(f_0)[J](T; \lambda) - \text{Bol}_c^{(2)}(f_0)[J](T; \lambda)). \quad (5.190)$$

Recalling (5.50), (5.52), (5.56), (5.53), and (5.54), Lemma 5.5 then implies

$$\begin{aligned} \text{err}_{2,j}^{(dis)}\left(\frac{T}{\lambda^2}; f[J]\right) &\leq \frac{\lambda^j}{N \lambda^{2r+2} |\Lambda|^{\frac{r}{3}}} \\ &\leq \frac{C \|\hat{v}\|_{2(\lfloor \frac{T}{2} \rfloor + 1), w, c} \|f_0\|_{2(\lfloor \frac{T}{2} \rfloor + 1), c} r^{\langle T \rangle^{r+2}} \|J\|_{W^2[\lfloor \frac{T}{2} \rfloor + 2, \infty]}}{N \lambda^{2r+2} |\Lambda|^{\frac{r}{3}}} \end{aligned} \quad (5.191)$$

for $j \in \{0, 1, 2\}$. Lemma C.5 yields

$$\begin{aligned} & \frac{\lambda^j}{N} |\text{Bol}_c^{(j)}(f_0)[J](T; \lambda)| \\ & \leq C \|\hat{v}\|_{2(\lfloor \frac{T}{2} \rfloor + 1), w, c} \|f_0\|_{2(\lfloor \frac{T}{2} \rfloor + 1), c} r^T (1 + \log(T)) \|J\|_{C^1} \frac{\lambda |\log(\lambda)|}{N} \end{aligned} \quad (5.192)$$

for $j \in \{1, 2\}$, and all $\lambda > 0$ small enough, dependent on $\|\hat{v}\|_{2(\lfloor \frac{T}{2} \rfloor + 1), w, c}$. Let

$$\begin{aligned} & \frac{1}{N} \text{Bol}_c(f_0)[J](T; \lambda) + \text{err}_{2,c}^{(Bog, Bol)}\left(\frac{T}{\lambda^2}; f[J]\right) \\ & := \frac{1}{N} (\text{Bol}_d(f_0)[J](T; \lambda) + \lambda \text{Bol}_d^{(1)}(f_0)[J](T; \lambda) \\ & \quad + \lambda^2 \text{Bol}_d^{(2)}(f_0)[J](T; \lambda)) + \text{err}_{2,d}^{(Bog, Bol)}\left(\frac{T}{\lambda^2}; f[J]\right). \end{aligned} \quad (5.193)$$

Collecting (5.186), (5.191), and (5.192), we thus proved that

$$\begin{aligned}
 & |\text{err}_{2,c}^{(Bog,Bol)}\left(\frac{T}{\lambda^2}; f[J]\right)| \\
 & \leq C \|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c} \|f_0\|_{2(\lfloor \frac{r}{2} \rfloor + 1), c, r} \left(\frac{T^2 \|J\|_{\ell^\infty(\Lambda^*)} \lambda}{N} \right. \\
 & \quad + \frac{\langle T \rangle^{r+2} \|J\|_{W^2 \lfloor \frac{r}{2} \rfloor + 2, \infty}}{N \lambda^{2(r+1)} |\Lambda|^{\frac{r}{3}}} \\
 & \quad \left. + \frac{\lambda |\log(\lambda)| T (1 + \log(T)) \|J\|_{W^{1, \infty}}}{N} \right) \\
 & \leq \frac{C \|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c} \|f_0\|_{2(\lfloor \frac{r}{2} \rfloor + 1), c, r} \langle T \rangle^{r+2} \|J\|_{W^2 \lfloor \frac{r}{2} \rfloor + 2, \infty}}{N} \\
 & \quad \left(\frac{1}{\lambda^{2(r+1)} |\Lambda|^{\frac{r}{3}}} + \lambda |\log(\lambda)| \right). \tag{5.194}
 \end{aligned}$$

Finally, we compute the discretization error in the dynamics of g . Let

$$\begin{aligned}
 & \text{err}_{1,c}^{(dis)}\left(\frac{T}{\lambda^2}; g[J]\right) \\
 & := \frac{1}{N} \int_0^T dS (\text{abs}_{\text{quart}, d}(f_0)[J](S/\lambda^2) - \text{abs}_{\text{quart}, c}(f_0)[J](S/\lambda^2)). \tag{5.195}
 \end{aligned}$$

Lemma 5.5 together with Proposition 5.2 implies

$$|\text{err}_{1,c}^{(dis)}\left(\frac{T}{\lambda^2}; g[J]\right)| \leq \frac{C \|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c} \|f_0\|_{2(\lfloor \frac{r}{2} \rfloor + 1), c, r} \langle T \rangle^{r+1} \|J\|_{W^2 \lfloor \frac{r}{2} \rfloor + 2, \infty}}{N \lambda^{2r} |\Lambda|^{\frac{r}{3}}} \tag{5.196}$$

Moreover, let

$$\begin{aligned}
 & \text{err}_{2,c}^{(dis)}\left(\frac{T}{\lambda^2}; g[J]\right) \\
 & := \frac{1}{N \lambda^2} \int_{\Delta[T, 2]} d\mathbf{S}_2 (\text{col}_d(f_0)[J](\mathbf{S}_2/\lambda^2) - \text{col}_c(f_0)[J](\mathbf{S}_2/\lambda^2)) \\
 & \quad + \text{abs}_{\text{cub}, d}(f_0)[J](\mathbf{S}_2/\lambda^2) - \text{abs}_{\text{cub}, c}(f_0)[J](\mathbf{S}_2/\lambda^2)) \tag{5.197}
 \end{aligned}$$

Again, applying Lemma 5.5 together with Proposition 5.2, yields

$$|\text{err}_{2,c}^{(dis)}\left(\frac{T}{\lambda^2}; g[J]\right)| \leq \frac{C \|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c} \|f_0\|_{2(\lfloor \frac{r}{2} \rfloor + 1), c, r} \langle T \rangle^{r+2} \|J\|_{W^2 \lfloor \frac{r}{2} \rfloor + 2, \infty}}{N \lambda^{2r+2} |\Lambda|^{\frac{r}{3}}} \tag{5.198}$$

for all $\lambda > 0$ small enough, dependent on $\|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c}$. This concludes the proof. \square

For the next result, observe that

$$\Delta_{\text{cub}} E = E(p_1) + E(p_2) - E(p_1 + p_2) = -p_1 \cdot p_2. \tag{5.199}$$

Let

$$\begin{aligned}
 & \text{Bol}_{\text{fec}}(f_0)[J](T) \\
 & := \frac{\pi T}{(2\pi)^6} \int_{p_1 \perp p_2} d\mathcal{H}^5(\mathbf{p}_2) \Delta_{\text{cub}} J(\mathbf{p}_2) \frac{(\hat{v}(p_1) + \hat{v}(p_2))^2}{|\mathbf{p}_2|} \\
 & \quad (\tilde{f}_0(p_1) \tilde{f}_0(p_2) f_0(p_1 + p_2) - f_0(p_1) f_0(p_2) \tilde{f}_0(p_1 + p_2)) \tag{5.200}
 \end{aligned}$$

be the Boltzmann operator with energy conserving collision kernel for the free energy dispersion $E(p) = \frac{|p|^2}{2}$.

Proposition 5.8. *We have that*

$$\begin{aligned} & \text{Bol}_c(f_0)[J](T; \lambda) \\ &= \frac{\pi T}{(2\pi)^6} \int d\mathbf{p}_2 \delta_{\frac{2\lambda^2}{T}}(\Delta_{cub}\Omega(\mathbf{p}_2)) \Delta_{cub}J(\mathbf{p}_2) (\hat{v}(p_1) + \hat{v}(p_2))^2 \\ & \quad (\tilde{f}_0(p_1)\tilde{f}_0(p_2)f_0(p_1+p_2) - f_0(p_1)f_0(p_2)\tilde{f}_0(p_1+p_2)), \end{aligned} \quad (5.201)$$

where $\delta_\varepsilon(x) := \frac{\varepsilon \sin^2(x/\varepsilon)}{\pi x^2}$. Moreover, approximating $\text{Bol}(f_0)[J](T\lambda^{-2})$ with its continuum counterpart, i.e., replacing $\frac{1}{|\Lambda|} \sum_{\Lambda^*}$ by the (Lebesgue-) integral $\frac{1}{(2\pi)^3} \int_{\mathbb{R}^3}$, we find that

$$\frac{1}{N} \text{Bol}_c(f_0)[J](T; \lambda) = \frac{1}{N} \text{Bol}_{fec}(f_0)[J](T) + \text{err}_{2,c}^{(fec)}\left(\frac{T}{\lambda^2}; f[J]\right), \quad (5.202)$$

where

$$|\text{err}_{2,c}^{(fec)}\left(\frac{T}{\lambda^2}; f[J]\right)| \leq C \|\hat{v}\|_{2([\frac{T}{2}]_{+1}), w, c} \|f_0\|_{2([\frac{T}{2}]_{+1}), c} \langle T \rangle \|J\|_{W^{2,\infty}} \frac{\lambda}{N} \quad (5.203)$$

for all $\lambda > 0$ small enough, dependent on $\|\hat{v}\|_{2([\frac{T}{2}]_{+1}), w, c}$.

Proof. Observe that $\Delta_{cub}E = 0$ is equivalent to $p_1 \perp p_2$, see (5.199), and that

$$|\nabla \Delta_{cub}E(\mathbf{p}_2)| = \left| \begin{pmatrix} -p_2 \\ -p_1 \end{pmatrix} \right| = |\mathbf{p}_2|. \quad (5.204)$$

Moreover, notice that for any $\omega \in \mathbb{R} \setminus \{0\}$, we have

$$\begin{aligned} \text{Re} \int_{\Delta[t,2]} ds_2 e^{-i\omega(s_1-s_2)} &= \frac{1 - \cos(\omega t)}{\omega^2} \\ &= 2 \frac{\sin^2(\frac{\omega t}{2})}{\omega^2}. \end{aligned} \quad (5.205)$$

Observe that $\delta_a(x) = \frac{a \sin^2(x/a)}{\pi x^2}$ defines an approximate identity in the sense that we have

$$\delta_a(x) \geq 0 \quad \forall a > 0, \quad (5.206)$$

$$\int dx \delta_a(x) = 1 \quad \forall a > 0, \quad (5.207)$$

$$\int_{|x|>\rho} dx \delta_a(x) \leq \frac{Ca}{\rho} \quad \forall a, \rho > 0. \quad (5.208)$$

With that, we obtain

$$\begin{aligned} & \text{Bol}(f_0)[J](T; \lambda) \\ &= \frac{\pi T}{(2\pi)^6} \int d\mathbf{p}_2 \delta_{\frac{2\lambda^2}{T}}(\Delta_{cub}\Omega(\mathbf{p}_2)) \Delta_{cub}J(\mathbf{p}_2) (\hat{v}(p_1) + \hat{v}(p_2))^2 \\ & \quad (f_0(p_1+p_2) + f_0(p_1)f_0(p_1+p_2) + f_0(p_1+p_2)f_0(p_2) \\ & \quad - f_0(p_1)f_0(p_2)) \end{aligned} \quad (5.209)$$

This proves the first part of the statement.

For the second part, assume \mathbf{p}_2 ranges over \mathbb{R}^6 . Emphasizing the dependence of $\mathcal{E}(\lambda, \mathbf{p}_2) := \Delta_{cub} \Omega_\lambda(\mathbf{p}_2)$ on $\lambda > 0$, we abbreviate (5.209) as

$$\lambda^2 \text{Bol}(f_0)[J](T\lambda^{-2}) =: \int d\mathbf{p}_2 \delta_{\frac{2\lambda^2}{T}}(\mathcal{E}(\lambda, \mathbf{p}_2)) H(\mathbf{p}_2), \quad (5.210)$$

where we emphasize. Observe that $\mathcal{E}(0, \mathbf{p}_2) = \Delta_{cub} E(\mathbf{p}_2)$. Using the Fundamental Theorem of Calculus by the Coarea Formula, we obtain that

$$\begin{aligned} N \text{err}_2^{(fd)}\left(\frac{T}{\lambda^2}; f[J]\right) &:= \int d\mathbf{p}_2 \left[\delta_{\frac{2\lambda^2}{T}}(\mathcal{E}(\lambda, \mathbf{p}_2)) - \delta_{\frac{2\lambda^2}{T}}(\mathcal{E}(0, \mathbf{p}_2)) \right] H(\mathbf{p}_2) \\ &= \int_0^\lambda d\tau \delta'_{\frac{2\lambda^2}{T}}(\mathcal{E}(\tau, \mathbf{p}_2)) \partial_\tau \mathcal{E}(\tau, \mathbf{p}_2) H(\mathbf{p}_2) \\ &= \int_0^\lambda d\tau \int d\omega \delta'_{\frac{2\lambda^2}{T}}(\omega) \\ &\quad \int_{\mathcal{E}(\tau, \mathbf{p}_2)=\omega} d\mathcal{H}^5 \frac{\partial_\tau \mathcal{E}(\tau, \mathbf{p}_2) H(\mathbf{p}_2)}{|\nabla_{\mathbf{p}_2} \mathcal{E}(\tau, \mathbf{p}_2)|}. \end{aligned} \quad (5.211)$$

We prove in Lemma C.6 that

$$|\text{err}_2^{(fd)}(t; f[J])| \leq C \|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c} \|f_0\|_{2(\lfloor \frac{r}{2} \rfloor + 1), c} r^T \|J\|_{W^{2, \infty}} \frac{\lambda}{N} \quad (5.212)$$

for all $\lambda > 0$ small enough, dependent on $\|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c}$.

Next, using (5.207) together with the Coarea Formula, we have that

$$\begin{aligned} N \text{err}_2^{(ec)}\left(\frac{T}{\lambda^2}; f[J]\right) &:= \int d\mathbf{p}_2 \delta_{\frac{2\lambda^2}{T}}(\Delta_{cub} E(\mathbf{p}_2)) H(\mathbf{p}_2) \\ &\quad - \int_{\Delta_{cub} E=0} d\mathcal{H}^5 \frac{H}{|\nabla \Delta_{cub} E|} \\ &= \int d\omega \delta_{\frac{2\lambda^2}{T}}(\omega) \left(\int_{\Delta_{cub} E=\omega} d\mathcal{H}^5 \frac{H}{|\nabla \Delta_{cub} E|} \right. \\ &\quad \left. + \int_{\Delta_{cub} E=0} d\mathcal{H}^5 \frac{H}{|\nabla \Delta_{cub} E|} \right). \end{aligned} \quad (5.213)$$

We prove in Lemma C.7 that

$$|\text{err}_2^{(ec)}\left(\frac{T}{\lambda^2}; f[J]\right)| \leq C \|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c} \|f_0\|_{2(\lfloor \frac{r}{2} \rfloor + 1), c} r \sqrt{T} \|J\|_{W^{2, \infty}} \frac{\lambda}{N} \quad (5.214)$$

for all $\lambda > 0$ small enough, dependent on $\|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c}$. In particular, we have that

$$\text{err}_{2,c}^{(fec)}\left(\frac{T}{\lambda^2}; f[J]\right) = \text{err}_2^{(fd)}\left(\frac{T}{\lambda^2}; f[J]\right) + \text{err}_2^{(ec)}\left(\frac{T}{\lambda^2}; f[J]\right) \quad (5.215)$$

satisfies

$$|\text{err}_{2,c}^{(fec)}\left(\frac{T}{\lambda^2}; f[J]\right)| \leq C \|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c} \|f_0\|_{2(\lfloor \frac{r}{2} \rfloor + 1), c} r \langle T \rangle \|J\|_{W^{2, \infty}} \frac{\lambda}{N}. \quad (5.216)$$

This concludes the proof. \square

5.3. Centered expectations. In order to resolve the fluctuations around the HFB dynamics, we have to consider the dynamics relative to the condensate term, i.e.,

$$f_t^{(\Phi)}(p) := f_t(p) - |\Phi_t|^2 \delta(p). \quad (5.217)$$

Proposition 5.9. *Let $T > 0$, $\lambda \in (0, 1)$ and $|\Lambda| \geq 1$. Let $J \in L^\infty(\mathbb{R}^3; \mathbb{R})$. Then the following holds.*

$$(1) \quad \frac{1}{N^{\frac{1}{2}} \lambda} \text{Con}_d(f_0)(T; \lambda) = \frac{1}{N^{\frac{1}{2}} \lambda} \text{Con}_d(f^{(\Phi)})(T; \lambda) + \text{err}_{1,d}^{(cen)}\left(\frac{T}{\lambda^2}; \Phi\right), \quad (5.218)$$

where

$$\begin{aligned} & |\text{err}_{1,d}^{(cen)}\left(\frac{T}{\lambda^2}; \Phi\right)| \\ & \leq C_{\|\hat{v}\|_{w,d}, \|f_0\|_d} \langle T \rangle^3 \frac{|\Lambda|^{\frac{3}{2}} e^{C\|\hat{v}\|_{w,d}|\Lambda|/\lambda T}}{N\lambda^2} \\ & \quad \left(1 + \frac{|\Lambda|}{N}\right) \left(1 + \frac{1}{N^{\frac{1}{2}} \lambda |\Lambda|^{\frac{3}{2}}}\right), \end{aligned} \quad (5.219)$$

$$\begin{aligned} (2) \quad (a) \quad & \frac{1}{N} (\text{Bol}_d(f_0)[J](T; \lambda) + \lambda \text{Bol}_d^{(1)}(f_0)[J](T; \lambda) + \lambda^2 \text{Bol}_d^{(2)}(f_0)[J](T; \lambda)) \\ & = \frac{1}{N} (\text{Bol}_d(f^{(\Phi)})[J](T; \lambda) + \lambda \text{Bol}_d^{(1)}(f^{(\Phi)})[J](T; \lambda) \\ & \quad + \lambda^2 \text{Bol}_d^{(2)}(f^{(\Phi)})[J](T; \lambda)) + \text{err}_{2,d}^{(cen)}\left(\frac{T}{\lambda^2}; f[J]\right), \end{aligned} \quad (5.220)$$

with

$$\begin{aligned} & |\text{err}_{2,d}^{(cen)}\left(\frac{T}{\lambda^2}; f[J]\right)| \\ & \leq C_{\|\hat{v}\|_{w,d}, \|f_0\|_d} \|J\|_{\ell^\infty(\Lambda^*)} \langle T \rangle^4 \frac{|\Lambda|^{\frac{3}{2}} e^{C\|\hat{v}\|_{w,d}|\Lambda|/\lambda T}}{N^{\frac{3}{2}} \lambda^3} \\ & \quad \left(1 + \frac{|\Lambda|}{N}\right) \left(1 + \frac{1}{N^{\frac{1}{2}} \lambda |\Lambda|^{\frac{3}{2}}}\right) \\ & \quad \left[1 + \langle T \rangle^2 \frac{|\Lambda|^{\frac{3}{2}} e^{C\|\hat{v}\|_{w,d}|\Lambda|/\lambda T}}{N^{\frac{1}{2}} \lambda} \left(1 + \frac{|\Lambda|}{N}\right) \left(1 + \frac{1}{N^{\frac{1}{2}} \lambda |\Lambda|^{\frac{3}{2}}}\right)\right], \end{aligned} \quad (5.221)$$

$$\begin{aligned} (b) \quad & \frac{1}{N} \text{Bol}_c(f_0)[J](T; \lambda) \\ & = \frac{1}{N} \text{Bol}_c(f^{(\Phi)})[J](T; \lambda) + \text{err}_{2,c}^{(cen)}\left(\frac{T}{\lambda^2}; f[J]\right), \end{aligned} \quad (5.222)$$

where $\text{err}_{2,c}^{(cen)}\left(\frac{T}{\lambda^2}; f[J]\right)$ satisfies the same bound as $\text{err}_{2,d}^{(cen)}\left(\frac{T}{\lambda^2}; f[J]\right)$,

$$\begin{aligned} (3) \quad (a) \quad & \frac{1}{N} \int_0^T dS \text{abs}_{\text{quart},d}(f_0)[J](S/\lambda^2) \\ & = \frac{1}{N} \int_0^T dS \text{abs}_{\text{quart},d}(f^{(\Phi)})[J](S/\lambda^2) + \text{err}_{1,d}^{(fd)}\left(\frac{T}{\lambda^2}; g[J]\right) \end{aligned} \quad (5.223)$$

with

$$\begin{aligned}
& |\text{err}_{1,d}^{(cen)}(\frac{T}{\lambda^2}; g[J])| \\
& \leq C_{\|\hat{v}\|_{w,d}, \|f_0\|_d} \|J\|_{\ell^\infty(\Lambda^*)} \langle T \rangle^3 \frac{|\Lambda|^{\frac{3}{2}} e^{C\|\hat{v}\|_{w,d}|\Lambda|/\lambda T}}{N^{\frac{3}{2}}\lambda} \\
& \quad \left(1 + \frac{|\Lambda|}{N}\right) \left(1 + \frac{1}{N^{\frac{1}{2}}\lambda|\Lambda|^{\frac{3}{2}}}\right) \\
& \quad \left[1 + \langle T \rangle^2 \frac{|\Lambda|^{\frac{3}{2}} e^{C\|\hat{v}\|_{w,d}|\Lambda|/\lambda T}}{N^{\frac{1}{2}}\lambda} \left(1 + \frac{|\Lambda|}{N}\right) \left(1 + \frac{1}{N^{\frac{1}{2}}\lambda|\Lambda|^{\frac{3}{2}}}\right)\right], \quad (5.224)
\end{aligned}$$

(b)

$$\begin{aligned}
& \frac{1}{N\lambda^2} \int_{\Delta_{[T,2]}} d\mathbf{S}_2 (\text{col}_d(f_0)[J](\mathbf{S}_2/\lambda^2) + \text{abs}_{cub,d}(f_0)[J](\mathbf{S}_2/\lambda^2)) \\
& = \frac{1}{N\lambda^2} \int_{\Delta_{[T,2]}} d\mathbf{S}_2 (\text{col}_d(f^{(\Phi)})[J](\mathbf{S}_2/\lambda^2) \\
& \quad + \text{abs}_{cub,d}(f^{(\Phi)})[J](\mathbf{S}_2/\lambda^2)) + \text{err}_{2,d}^{(cen)}(\frac{T}{\lambda^2}; g[J]) \quad (5.225)
\end{aligned}$$

with

$$\begin{aligned}
& |\text{err}_{2,d}^{(cen)}(\frac{T}{\lambda^2}; g[J])| \\
& \leq C_{\|\hat{v}\|_{w,d}, \|f_0\|_d} \|J\|_{\ell^\infty(\Lambda^*)} \langle T \rangle^3 \frac{|\Lambda|^{\frac{3}{2}} e^{C\|\hat{v}\|_{w,d}|\Lambda|/\lambda T}}{N^{\frac{3}{2}}\lambda^3} \\
& \quad \left(1 + \frac{|\Lambda|}{N}\right) \left(1 + \frac{1}{N^{\frac{1}{2}}\lambda|\Lambda|^{\frac{3}{2}}}\right) \\
& \quad \left[1 + \langle T \rangle^2 \frac{|\Lambda|^{\frac{3}{2}} e^{C\|\hat{v}\|_{w,d}|\Lambda|/\lambda T}}{N^{\frac{1}{2}}\lambda} \left(1 + \frac{|\Lambda|}{N}\right) \left(1 + \frac{1}{N^{\frac{1}{2}}\lambda|\Lambda|^{\frac{3}{2}}}\right)\right]. \quad (5.226)
\end{aligned}$$

The analogous statements hold true for the continuum approximation if one replaces $(\|\hat{v}\|_{w,d}, \|f_0\|_d)$ on the RHS of the inequalities by $(C_r\|\hat{v}\|_{2(\lfloor \frac{T}{2} \rfloor + 1), w, c}, C_r\|f_0\|_{2(\lfloor \frac{T}{2} \rfloor + 1), c})$.

Proof. In the notation of the proof, we will focus on the discrete case. The bounds for the continuum approximation follow by replacing $(\|\hat{v}\|_{w,d}, \|f_0\|_d)$ by

$$(C_r\|\hat{v}\|_{2(\lfloor \frac{T}{2} \rfloor + 1), w, c}, C_r\|f_0\|_{2(\lfloor \frac{T}{2} \rfloor + 1), c}), \quad (5.227)$$

see Remark 5.6.

Let $t = T\lambda^{-2}$. Next, Lemma 3.1 and the definition (5.217) of $f^{(\Phi)}$ imply that

$$\begin{aligned}
\int dp J(p)(f_0(p) - f_t^{(\Phi)}(p)) & = \int dp J(p)(f_0(p) - f_t(p)) + J(0)|\Phi_t|^2 \\
& = -i \int_0^t ds \frac{\nu_s([f[J], \mathcal{H}_I(s)])}{|\Lambda|} + J(0)|\Phi_t|^2 \quad (5.228)
\end{aligned}$$

Using (5.13), we have the estimate

$$\begin{aligned}
& \left| \frac{\nu_s([f[J], \mathcal{H}_I(s)])}{|\Lambda|} \right| \\
& \leq C_{\|\hat{v}\|_{w,d}, \|f_0\|_d} e^{C\|f_0\|_d|\Lambda|/\lambda T} \|J\|_{\ell^\infty(\Lambda^*)} \frac{\lambda|\Lambda|^{\frac{3}{2}}}{N^{\frac{1}{2}}} \left(1 + \frac{|\Lambda|^{\frac{1}{2}}}{N^{\frac{1}{2}}}\right). \quad (5.229)
\end{aligned}$$

We have that

$$\Phi_t = -i \int_0^t ds \frac{\nu_s([a_0, \mathcal{H}_I(s)])}{|\Lambda|}. \quad (5.230)$$

Recall from (5.18) and (5.26) that

$$[a_0, \mathcal{H}_{cub}(s)] = f[J_1(s)] + g[J_2(s)] + g^*[J_3(s)] + J_4(s), \quad (5.231)$$

$$[a_0, \mathcal{H}_{quart}(s)] = \frac{1}{N^{\frac{1}{2}}} \mathcal{H}_{cub}^{(1)}(s). \quad (5.232)$$

Collecting (5.24), (5.25), and using Lemma 4.4, (5.13) yields

$$|\Phi_t| \leq C_{\|\hat{v}\|_{w,d}, \|f_0\|_d} T e^{C\|f_0\|_d |\Lambda|/\lambda T} \frac{1}{N^{\frac{1}{2}} \lambda} \left(1 + \frac{|\Lambda|^{\frac{1}{2}}}{N^{\frac{1}{2}}}\right). \quad (5.233)$$

Then (5.228), (5.229), and (5.233) yield that

$$\begin{aligned} & \left| \int dp J(p) (f_0(p) - f_t^{(\Phi)}(p)) \right| \\ & \leq C_{\|\hat{v}\|_{w,d}, \|f_0\|_d} \langle T \rangle^2 \|J\|_{\ell^\infty(\Lambda^*)} \frac{|\Lambda|^{\frac{3}{2}} e^{C\|\hat{v}\|_{w,d} |\Lambda|/\lambda T}}{N^{\frac{1}{2}} \lambda} \\ & \quad \left(1 + \frac{|\Lambda|}{N}\right) \left(1 + \frac{1}{N^{\frac{1}{2}} \lambda |\Lambda|^{\frac{3}{2}}}\right). \end{aligned} \quad (5.234)$$

As a consequence of (5.234) and recalling definition (5.49) of $\text{Con}(f)$, we find that

$$\begin{aligned} & |\text{err}_1^{(cen)}(\frac{T}{\lambda^2}; \Phi)| \\ & \leq C_{\|\hat{v}\|_{w,d}, \|f_0\|_d} \langle T \rangle^3 \frac{|\Lambda|^{\frac{3}{2}} e^{C\|\hat{v}\|_{w,d} |\Lambda|/\lambda T}}{N \lambda^2} \\ & \quad \left(1 + \frac{|\Lambda|}{N}\right) \left(1 + \frac{1}{N^{\frac{1}{2}} \lambda |\Lambda|^{\frac{3}{2}}}\right). \end{aligned} \quad (5.235)$$

After substitution and using the notation in the proof of Proposition 5.1, the terms in $\text{bol}^{(j)}(f_0)[J](t) - \text{bol}^{(j)}(f^{(\Phi)})[J](t)$ and $\text{col}^{(k)}(f_0)[J](t) - \text{col}^{(k)}(f^{(\Phi)})[J](t)$ are of the form

$$\int d\mathbf{p}_2 H_1(p_1) H_2(p_2) H_3(p_1 \pm p_2) f_{s_2}^{(\Phi)}(p_1) (f_0(p_2) - f_{s_2}^{(\Phi)}(p_2)), \quad (5.236)$$

$$\int d\mathbf{p}_2 H_1(p_1) H_2(p_2) H_3(p_1 \pm p_2) f_0(p_1) (f_0(p_2) - f_{s_2}^{(\Phi)}(p_2)), \quad (5.237)$$

$$\int d\mathbf{p}_2 H_1(p_1) H_2(p_2) H_3(p_{12}) (f_0(p_{12}) - f_{s_2}^{(\Phi)}(p_{12})), \quad (5.238)$$

where $H_j \in L_{s_2}^\infty(\mathbb{R}^2; L_p^\infty(\mathbb{R}^3))$ can differ in every line and in the last line, we require $H_1 \in L_{s_2}^\infty(\mathbb{R}^2; L_p^1(\mathbb{R}^3))$. We have that $s_2 \leq T\lambda^{-2}$ with $\lambda \in (0, 1)$.

Observe that

$$\begin{aligned} \int dp |H_1(p)| f_0(p) &= \frac{\nu_0(f(|H_1|))}{|\Lambda|} \\ &\leq \frac{\|H_1\|_\infty \nu_0(\mathcal{N}_b)}{|\Lambda|} \\ &\leq \|f_0\|_d \|H_1\|_\infty \end{aligned} \quad (5.239)$$

due to Lemma 4.7 and

$$\frac{\nu_0(\mathcal{N}_b)}{|\Lambda|} = \int dp f_0(p) \leq \|f_0\|_d. \quad (5.240)$$

(5.234) and (5.239) imply that the terms of the form (5.237) satisfy

$$\begin{aligned} & \left| \int d\mathbf{p}_2 H_1(p_1) H_2(p_2) H_3(p_1 \pm p_2) f_0(p_1) (f_0(p_2) - f_{s_2}^{(\Phi)}(p_2)) \right| \\ & \leq \int dp_1 |H_1(p_1)| |f_0(p_1)| \left| \int dp_2 H_2(p_2) H_3(p_1 \pm p_2) (f_0(p_2) - f_{s_2}^{(\Phi)}(p_2)) \right| \\ & \leq C_{\|\hat{v}\|_{w,d}, \|f_0\|_d} \|H_1\|_\infty \|H_2\|_\infty \|H_3\|_\infty \langle T \rangle^2 \frac{|\Lambda|^{\frac{3}{2}} e^{C\|\hat{v}\|_{w,d}|\Lambda|/\lambda T}}{N^{\frac{1}{2}}\lambda} \\ & \quad \left(1 + \frac{|\Lambda|}{N}\right) \left(1 + \frac{1}{N^{\frac{1}{2}}\lambda|\Lambda|^{\frac{3}{2}}}\right). \end{aligned} \quad (5.241)$$

For (5.236), (5.234) and (5.241) yield that

$$\begin{aligned} & \left| \int d\mathbf{p}_2 H_1(p_1) H_2(p_2) H_3(p_1 \pm p_2) f_{s_2}^{(\Phi)}(p_1) (f_0(p_2) - f_{s_2}^{(\Phi)}(p_2)) \right| \\ & \leq \left| \int d\mathbf{p}_2 H_1(p_1) H_2(p_2) H_3(p_1 \pm p_2) f_0(p_1) (f_0(p_2) - f_{s_2}^{(\Phi)}(p_2)) \right| \\ & \quad + \left| \int d\mathbf{p}_2 H_1(p_1) H_2(p_2) H_3(p_1 \pm p_2) (f_{s_2}^{(\Phi)}(p_1) - f_0(p_1)) (f_0(p_2) - f_{s_2}^{(\Phi)}(p_2)) \right| \\ & \leq C_{\|\hat{v}\|_{w,d}, \|f_0\|_d} \|H_1\|_\infty \langle T \rangle^2 \frac{|\Lambda|^{\frac{3}{2}} e^{C\|\hat{v}\|_{w,d}|\Lambda|/\lambda T}}{N^{\frac{1}{2}}\lambda} \\ & \quad \left(1 + \frac{|\Lambda|}{N}\right) \left(1 + \frac{1}{N^{\frac{1}{2}}\lambda|\Lambda|^{\frac{3}{2}}}\right) \\ & \quad \left(\|H_2\|_\infty \|H_3\|_\infty + \sup_{p_1} \left| \int dp_2 H_2(p_2) H_3(p_1 \pm p_2) (f_0(p_2) - f_{s_2}^{(\Phi)}(p_2)) \right| \right) \\ & \leq C_{\|\hat{v}\|_{w,d}, \|f_0\|_d} \|H_1\|_\infty \|H_2\|_\infty \|H_3\|_\infty \langle T \rangle^2 \frac{|\Lambda|^{\frac{3}{2}} e^{C\|\hat{v}\|_{w,d}|\Lambda|/\lambda T}}{N^{\frac{1}{2}}\lambda} \\ & \quad \left(1 + \frac{|\Lambda|}{N}\right) \left(1 + \frac{1}{N^{\frac{1}{2}}\lambda|\Lambda|^{\frac{3}{2}}}\right) \\ & \quad \left[1 + \langle T \rangle^2 \frac{|\Lambda|^{\frac{3}{2}} e^{C\|\hat{v}\|_{w,d}|\Lambda|/\lambda T}}{N^{\frac{1}{2}}\lambda} \left(1 + \frac{|\Lambda|}{N}\right) \left(1 + \frac{1}{N^{\frac{1}{2}}\lambda|\Lambda|^{\frac{3}{2}}}\right) \right]. \end{aligned} \quad (5.242)$$

Employing (5.234) again, we estimate (5.238) by

$$\begin{aligned} & \left| \int d\mathbf{p}_2 H_1(p_1) H_2(p_2) H_3(p_1 + p_2) (f_0(p_1 + p_2) - f_{s_2}^{(\Phi)}(p_1 + p_2)) \right| \\ & \leq C_{\|\hat{v}\|_{w,d}, \|f_0\|_d} \|H_1\|_1 \|H_2\|_\infty \|H_3\|_\infty \langle T \rangle^2 \frac{|\Lambda|^{\frac{3}{2}} e^{C\|\hat{v}\|_{w,d}|\Lambda|/\lambda T}}{N^{\frac{1}{2}}\lambda} \\ & \quad \left(1 + \frac{|\Lambda|}{N}\right) \left(1 + \frac{1}{N^{\frac{1}{2}}\lambda|\Lambda|^{\frac{3}{2}}}\right). \end{aligned} \quad (5.243)$$

Collecting (5.241)–(5.243), recalling definitions (5.50)–(5.56), and denoting $\text{Bol}^{(0)} := \text{Bol}$, we find the upper bound

$$\begin{aligned}
& \frac{\lambda^j}{N} |\text{Bol}^{(j)}(f_0)[J](T; \lambda) - \text{Bol}^{(j)}(f^{(\Phi)})[J](T; \lambda)| \\
& \leq C_{\|\hat{v}\|_w, d, \|f_0\|_d} \frac{\lambda^{2+j} t^2 \|J\|_{\ell^\infty(\Lambda^*)} \langle T \rangle^2 \frac{|\Lambda|^{\frac{3}{2}} e^{C\|\hat{v}\|_w, d|\Lambda|/\lambda T}}{N^{\frac{1}{2}} \lambda}}{N} \\
& \quad \left(1 + \frac{|\Lambda|}{N}\right) \left(1 + \frac{1}{N^{\frac{1}{2}} \lambda |\Lambda|^{\frac{3}{2}}}\right) \\
& \quad \left[1 + \langle T \rangle^2 \frac{|\Lambda|^{\frac{3}{2}} e^{C\|\hat{v}\|_w, d|\Lambda|/\lambda T}}{N^{\frac{1}{2}} \lambda} \left(1 + \frac{|\Lambda|}{N}\right) \left(1 + \frac{1}{N^{\frac{1}{2}} \lambda |\Lambda|^{\frac{3}{2}}}\right)\right]. \quad (5.244)
\end{aligned}$$

In particular, we have that

$$\begin{aligned}
& |\text{err}_{2,d}^{(cen)}(t; f[J])|, |\text{err}_{2,c}^{(cen)}(t; f[J])| \\
& \leq C_{\|\hat{v}\|_w, d, \|f_0\|_d} \|J\|_{\ell^\infty(\Lambda^*)} \langle T \rangle^4 \frac{|\Lambda|^{\frac{3}{2}} e^{C\|\hat{v}\|_w, d|\Lambda|/\lambda T}}{N^{\frac{3}{2}} \lambda^3} \\
& \quad \left(1 + \frac{|\Lambda|}{N}\right) \left(1 + \frac{1}{N^{\frac{1}{2}} \lambda |\Lambda|^{\frac{3}{2}}}\right) \\
& \quad \left[1 + \langle T \rangle^2 \frac{|\Lambda|^{\frac{3}{2}} e^{C\|\hat{v}\|_w, d|\Lambda|/\lambda T}}{N^{\frac{1}{2}} \lambda} \left(1 + \frac{|\Lambda|}{N}\right) \left(1 + \frac{1}{N^{\frac{1}{2}} \lambda |\Lambda|^{\frac{3}{2}}}\right)\right]. \quad (5.245)
\end{aligned}$$

Finally, it remains to estimate $\text{err}_j^{(cen)}(\frac{T}{\lambda^2}; g[J])$, $j \in \{1, 2\}$. Following analogous steps that lead to (5.245), Proposition 5.2 implies

$$\begin{aligned}
& |\text{err}_1^{(cen)}(\frac{T}{\lambda^2}; g[J])| \\
& \leq C_{\|\hat{v}\|_w, d, \|f_0\|_d} \|J\|_{\ell^\infty(\Lambda^*)} \langle T \rangle^3 \frac{|\Lambda|^{\frac{3}{2}} e^{C\|\hat{v}\|_w, d|\Lambda|/\lambda T}}{N^{\frac{3}{2}} \lambda} \\
& \quad \left(1 + \frac{|\Lambda|}{N}\right) \left(1 + \frac{1}{N^{\frac{1}{2}} \lambda |\Lambda|^{\frac{3}{2}}}\right) \\
& \quad \left[1 + \langle T \rangle^2 \frac{|\Lambda|^{\frac{3}{2}} e^{C\|\hat{v}\|_w, d|\Lambda|/\lambda T}}{N^{\frac{1}{2}} \lambda} \left(1 + \frac{|\Lambda|}{N}\right) \left(1 + \frac{1}{N^{\frac{1}{2}} \lambda |\Lambda|^{\frac{3}{2}}}\right)\right], \quad (5.246)
\end{aligned}$$

$$\begin{aligned}
& |\text{err}_2^{(cen)}(\frac{T}{\lambda^2}; g[J])| \\
& \leq C_{\|\hat{v}\|_w, d, \|f_0\|_d} \|J\|_{\ell^\infty(\Lambda^*)} \langle T \rangle^3 \frac{|\Lambda|^{\frac{3}{2}} e^{C\|\hat{v}\|_w, d|\Lambda|/\lambda T}}{N^{\frac{3}{2}} \lambda^3} \\
& \quad \left(1 + \frac{|\Lambda|}{N}\right) \left(1 + \frac{1}{N^{\frac{1}{2}} \lambda |\Lambda|^{\frac{3}{2}}}\right) \\
& \quad \left[1 + \langle T \rangle^2 \frac{|\Lambda|^{\frac{3}{2}} e^{C\|\hat{v}\|_w, d|\Lambda|/\lambda T}}{N^{\frac{1}{2}} \lambda} \left(1 + \frac{|\Lambda|}{N}\right) \left(1 + \frac{1}{N^{\frac{1}{2}} \lambda |\Lambda|^{\frac{3}{2}}}\right)\right]. \quad (5.247)
\end{aligned}$$

This concludes the proof. \square

6. MAIN ORDER TERMS IN THE EVOLUTION OF (Φ, f, g)

In this section, we will prove Theorem 2.2. We will always assume that $\lambda \in (0, 1)$ is small enough to comply with the estimates proven in all preceding steps.

6.1. Discrete case. Fix $L \geq 1$. Recall that we refer to the case of fixed Λ as the 'discrete case'.

6.1.1. *Main order term of Φ .* Using Propositions 5.1, and 5.2, we have that

$$\Phi_{\frac{T}{\lambda^2}} = \frac{1}{N^{\frac{1}{2}}\lambda} \text{Con}_d(f_0)(T; \lambda) + \text{Rem}_2\left(\frac{T}{\lambda^2}; \Phi\right) + \text{err}_{1,d}^{(Bog)}\left(\frac{T}{\lambda^2}; \Phi\right) \quad (6.1)$$

with

$$|\text{Rem}_2\left(\frac{T}{\lambda^2}; \Phi\right)| \leq C_{\|\hat{v}\|_{w,d}, \|f_0\|_d} T^2 e^{C\|\hat{v}\|_{w,d}|\Lambda|/\lambda T} \frac{|\Lambda|^{\frac{3}{2}}}{N\lambda^2}, \quad (6.2)$$

$$|\text{err}_{1,d}^{(Bog)}\left(\frac{T}{\lambda^2}; \Phi\right)| \leq \frac{C_{\|\hat{v}\|_{w,d}, \|f_0\|_d} T}{N^{\frac{1}{2}}}. \quad (6.3)$$

In particular, the main term is given by

$$\frac{1}{N^{\frac{1}{2}}\lambda} \text{Con}_d(f_0)(T; \lambda) = -\frac{iT}{N^{\frac{1}{2}}\lambda} \int_{\Lambda^*} dp \hat{v}(p) f_0(p), \quad (6.4)$$

see (5.49), and thus it is of size $N^{-\frac{1}{2}}\lambda^{-1}$. In order to suppress $\text{Rem}_2\left(\frac{T}{\lambda^2}; \Phi\right)$, we choose $\lambda = \log \log N / \log N$.

6.1.2. *Main order term of f .* For f , we apply Propositions 5.1, and 5.2 to obtain that

$$\begin{aligned} & \int_{\Lambda^*} dp \left(f_{\frac{T}{\lambda^2}}^{(\Phi)} - f_0^{(\Phi)} \right) J(p) \\ &= \frac{1}{N} \left(\text{Bol}_d(f_0)[J](T; \lambda) + \lambda \text{Bol}_d^{(1)}(f_0)[J](T; \lambda) + \lambda^2 \text{Bol}_d^{(2)}(f_0)[J](T; \lambda) \right) \\ &+ \text{Rem}_2\left(\frac{T}{\lambda^2}; f[J]\right) + \text{err}_{2,d}^{(Bog, Bol)}\left(\frac{T}{\lambda^2}; f[J]\right) + J(0) \text{err}_2^{(Bog, Con)}\left(\frac{T}{\lambda^2}; f\right), \end{aligned} \quad (6.5)$$

where

$$|\text{Rem}_2\left(\frac{T}{\lambda^2}; f[J]\right)| \leq C_{\|\hat{v}\|_{w,d}, \|f_0\|_d} \langle T \rangle^4 \|J\|_{\ell^\infty(\Lambda^*)} \frac{e^{C\|\hat{v}\|_{w,d}|\Lambda|/\lambda T} |\Lambda|^6}{\lambda^4 N^2}, \quad (6.6)$$

$$|\text{err}_{2,d}^{(Bog, Bol)}\left(\frac{T}{\lambda^2}; f[J]\right)| \leq C_{\|\hat{v}\|_{w,d}, \|f_0\|_d} \|J\|_{\ell^\infty(\Lambda^*)} T^2 \frac{\lambda}{N}, \quad (6.7)$$

$$|\text{err}_{2,d}^{(Bog, Con)}(t; f)| \leq C_{\|\hat{v}\|_{w,d}, \|f_0\|_d} e^{C\|\hat{v}\|_{w,d}|\Lambda|/\lambda T} \frac{\langle T \rangle^4 |\Lambda|^{\frac{3}{2}}}{\lambda^3 N^{\frac{3}{2}}} \left(1 + \frac{|\Lambda|^{\frac{3}{2}}}{\lambda N^{\frac{1}{2}}} \right). \quad (6.8)$$

Due to Lemma 5.3, we can expand the oscillatory factors in $\text{Bol}_d^{(j)}$ in powers of λ . Then the main order term depends on \hat{v} , $\lambda(N)$, and T . We are interested in terms up to $O(N^{-1})$ which, as we will see below, is the size of the main order term in the continuum approximation. For the same reasons as above, we choose $\lambda = \log \log N / \log N$.

6.1.3. *Main order term of g .* Similar to the case of f , we need to subtract the dynamics of the condensate in order to resolve the fluctuations defined by g . For that, we introduce

$$g_t^{(\Phi)}(p) := g_t(p) - \Phi_t^2 \delta(p). \quad (6.9)$$

Propositions 5.1 and 5.2 then yield

$$\begin{aligned} & \int_{\Lambda^*} dp g \frac{(\Phi)}{\lambda^2}(p) J(p) \\ &= \frac{1}{N} \int_0^T dS \text{abs}_{\text{quart},d}(f_0)[J](S/\lambda^2) \\ & \quad + \frac{1}{N\lambda^2} \int_{\Delta[T,2]} d\mathbf{S}_2 (\text{col}_d(f_0)[J](\mathbf{S}_2/\lambda^2) + \text{abs}_{\text{cub},d}(f_0)[J](\mathbf{S}_2/\lambda^2)) \\ & \quad + \text{Rem}_2\left(\frac{T}{\lambda^2}; g[J]\right) + J(0) \text{err}_2^{(\text{Bog}, \text{Con})}\left(\frac{T}{\lambda^2}; g\right) \end{aligned} \quad (6.10)$$

with

$$|\text{Rem}_2\left(\frac{T}{\lambda^2}; g[J]\right)| \leq C_{\|\hat{v}\|_{w,d}, \|f_0\|_d} \langle T \rangle^4 \|J\|_{2\cap\infty, d} \frac{e^{C\|\hat{v}\|_{w,d}|\Lambda|/\lambda T} |\Lambda|^6}{\lambda^4 N^2}, \quad (6.11)$$

$$|\text{err}_2^{(\text{Bog}, \text{Con})}\left(\frac{T}{\lambda^2}; g\right)| \leq C_{\|\hat{v}\|_{w,d}, \|f_0\|_d} e^{C\|\hat{v}\|_{w,d}|\Lambda|/\lambda T} \frac{\langle T \rangle^4 |\Lambda|^{\frac{3}{2}}}{\lambda^3 N^{\frac{3}{2}}} \left(1 + \frac{|\Lambda|^{\frac{3}{2}}}{\lambda N^{\frac{1}{2}}}\right). \quad (6.12)$$

As in the case of f , we again observe a phenomenon similar to the Talbot effect on $\text{abs}_{\text{quart},d}$, col_d , and $\text{abs}_{\text{cub},d}$. We choose N^{-1} as reference order.

6.1.4. *Conclusion for $L \sim 1$.* As described above, we choose $\lambda = \log \log(N)/\log(N)$. Let

$$\delta_{\Psi,d} = \min \left\{ \frac{\log \frac{1}{\lambda}}{\log N}, \frac{1}{2} - \frac{\log \frac{1}{\lambda}}{\log N} - \frac{C\|\hat{v}\|_{w,d}|\Lambda|T}{\log \log N} \right\}, \quad (6.13)$$

$$\delta_{F,d} = \min \left\{ \frac{\log \frac{1}{\lambda}}{\log N}, \frac{1}{2} - \frac{3 \log \frac{1}{\lambda}}{\log N} - \frac{C\|\hat{v}\|_{w,d}|\Lambda|T}{\log \log N} \right\}, \quad (6.14)$$

$$\delta_{G,d} = \frac{1}{2} - \frac{3 \log \frac{1}{\lambda}}{\log N} - \frac{C\|\hat{v}\|_{w,d}|\Lambda|T}{\log \log N}. \quad (6.15)$$

Then we obtain

$$\begin{aligned} & \left| \Phi \frac{(\Phi)}{\lambda^2} - \frac{1}{N^{\frac{1}{2}} \lambda} \text{Con}_d(f_0)(T; \lambda) \right| \\ & \leq \frac{C_{\|\hat{v}\|_{w,d}, \|f_0\|_d, |\Lambda|, T}}{N^{\frac{1}{2} + \delta_{\Psi,d}} \lambda}, \end{aligned} \quad (6.16)$$

$$\begin{aligned} & \left| \int_{\Lambda^*} dp \left(f \frac{(\Phi)}{\lambda^2}(p) - f_0^{(\Phi)}(p) \right) J(p) \right. \\ & \quad \left. - \frac{1}{N} \left(\text{Bol}_d(f_0)[J](T; \lambda) + \sum_{j=1}^2 \lambda^j \text{Bol}_d^{(j)}(f_0)[J](T; \lambda) \right) \right| \\ & \leq \frac{C_{\|\hat{v}\|_{w,d}, \|f_0\|_d, |\Lambda|, T} \|J\|_{\ell^\infty(\Lambda^*)}}{N^{1 + \delta_{F,d}}}, \end{aligned} \quad (6.17)$$

$$\left| \int_{\Lambda^*} dp g \frac{(\Phi)}{\lambda^2}(p) J(p) - \frac{1}{N} \int_0^T dS \text{abs}_{\text{quart},d}(f_0)[J](S/\lambda^2) \right|$$

$$\begin{aligned}
 & - \frac{1}{N\lambda^2} \int_{\Delta[T,2]} d\mathbf{S}_2 \left(\text{col}_d(f_0)[J](\mathbf{S}_2/\lambda^2) + \text{abs}_{\text{scub},d}(f_0)[J](\mathbf{S}_2/\lambda^2) \right) \\
 & \leq \frac{C_{\|\hat{v}\|_{w,d},\|f_0\|_d,|\Lambda|,T}\|J\|_{2\cap\infty,d}}{N^{1+\delta_{G,d}}}
 \end{aligned} \tag{6.18}$$

for all N larger than a universal constant. Notice that for all N large enough, we have that $\delta_{j,d} = \delta_{j,d}(N, |\Lambda|, T) > 0$ for $j \in \{\Psi, F, G\}$.

6.2. Continuum approximation. We have that errors coming from the tail in the Duhamel expansion grow like $\exp(C_r \|\hat{v}\|_{2(\lfloor \frac{T}{2} \rfloor + 1), w, c} T |\Lambda|/\lambda)$. As a consequence, we require

$$\frac{|\Lambda|}{\lambda} = O\left(\frac{\log(N)}{\log \log(N)}\right) \tag{6.19}$$

as $N \rightarrow \infty$. In order to use bounds established for general values of $|\Lambda|$ to the limit $|\Lambda| \rightarrow \infty$, for specific expressions of interest, we employ Remark 5.6.

6.2.1. Main order term of Φ . Using Propositions 5.1 and 5.7, we have that

$$\Phi_{\frac{T}{\lambda^2}} = \frac{1}{N^{\frac{1}{2}}\lambda} \text{Con}_c(f_0)(T; \lambda) + \text{Rem}_2\left(\frac{T}{\lambda^2}; \Phi\right) + \text{err}_{1,c}^{(Bog)}\left(\frac{T}{\lambda^2}; \Phi\right) \tag{6.20}$$

with

$$\begin{aligned}
 |\text{Rem}_2\left(\frac{T}{\lambda^2}; \Phi\right)| & \leq C_{\|\hat{v}\|_{2(\lfloor \frac{T}{2} \rfloor + 1), w, c}, \|f_0\|_{2(\lfloor \frac{T}{2} \rfloor + 1), c}, r} T^2 e^{C_r \|\hat{v}\|_{2(\lfloor \frac{T}{2} \rfloor + 1), w, c} |\Lambda|/\lambda T} \frac{|\Lambda|^{\frac{3}{2}}}{N\lambda^2} \\
 & \quad \left(1 + \frac{|\Lambda|}{N}\right), \\
 |\text{err}_{1,c}^{(Bog)}(t; \Phi)| & \leq \frac{C_{\|\hat{v}\|_{2(\lfloor \frac{T}{2} \rfloor + 1), w, c}, \|f_0\|_{2(\lfloor \frac{T}{2} \rfloor + 1), c}, r} T}{N^{\frac{1}{2}}\lambda} \left(\lambda + \frac{1}{|\Lambda|^{\frac{r}{6}}}\right)
 \end{aligned} \tag{6.21}$$

for some $r > 6$. The main order term is given by

$$\frac{1}{N^{\frac{1}{2}}\lambda} \text{Con}_c(f_0)(T; \lambda) = -\frac{iT}{(2\pi)^3 N^{\frac{1}{2}}\lambda} \int_{\mathbb{R}^3} dp \hat{v}(p) f_0(p), \tag{6.22}$$

and it is of size $N^{\frac{1}{2}}\lambda^{-1}$.

6.2.2. Main order term of f . For f , we apply Propositions 5.1, 5.7, and 5.8 to obtain that

$$\begin{aligned}
 & \int_{\Lambda^*} dp \left(f_{\frac{T}{\lambda^2}}^{(\Phi)}(p) - f_0^{(\Phi)}(p) \right) J(p) \\
 & = \frac{1}{N} \text{Bol}_{fec}(f_0)[J](T) \\
 & \quad + \text{Rem}_2\left(\frac{T}{\lambda^2}; f[J]\right) + \text{err}_{2,c}^{(Bog, Bol)}\left(\frac{T}{\lambda^2}; f[J]\right) \\
 & \quad + J(0) \text{err}_{2,c}^{(Bog, Con)}\left(\frac{T}{\lambda^2}; f\right) + \text{err}_{2,c}^{(fec)}(T\lambda^{-2}; f[J]),
 \end{aligned} \tag{6.23}$$

where

$$\begin{aligned}
 |\text{Rem}_2\left(\frac{T}{\lambda^2}; f[J]\right)| & \leq C_{\|\hat{v}\|_{2(\lfloor \frac{T}{2} \rfloor + 1), w, c}, \|f_0\|_{2(\lfloor \frac{T}{2} \rfloor + 1), c}, r} \langle T \rangle^4 \|J\|_{\infty} \\
 & \quad \frac{e^{C_r \|\hat{v}\|_{2(\lfloor \frac{T}{2} \rfloor + 1), w, c} |\Lambda|/\lambda T} |\Lambda|^6}{\lambda^4 N^2} \left(1 + \frac{|\Lambda|}{N}\right)^2,
 \end{aligned} \tag{6.24}$$

$$|\text{err}_{2,c}^{(Bog,Bol)}\left(\frac{T}{\lambda^2}; f[J]\right)| \leq \frac{C_{\|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c}, \|f_0\|_{2(\lfloor \frac{r}{2} \rfloor + 1), c}, r} \langle T \rangle^{r+2} \|J\|_{W^2(\lfloor \frac{r}{2} \rfloor + 2, \infty)}}{N} \left(\frac{1}{\lambda^{2(r+1)} |\Lambda|^{\frac{r}{3}}} + \lambda |\log(\lambda)| \right), \quad (6.25)$$

$$|\text{err}_{2,c}^{(Bog,Con)}\left(\frac{T}{\lambda^2}; f[J]\right)| \leq C_{\|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c}, \|f_0\|_{2(\lfloor \frac{r}{2} \rfloor + 1), c}, r} e^{C_r \|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c} |\Lambda| / \lambda T} \frac{\langle T \rangle^4 |\Lambda|^{\frac{3}{2}}}{\lambda^3 N^{\frac{3}{2}}} \left(1 + \frac{|\Lambda|}{N}\right)^2 \left(1 + \frac{|\Lambda|^{\frac{3}{2}}}{\lambda N^{\frac{1}{2}}}\right), \quad (6.26)$$

$$|\text{err}_{2,c}^{(fec)}\left(\frac{T}{\lambda^2}; f[J]\right)| \leq C_{\|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c}, \|f_0\|_{2(\lfloor \frac{r}{2} \rfloor + 1), c}, r} \langle T \rangle \|J\|_{W^{2,\infty}} \frac{\lambda}{N} \quad (6.27)$$

for all $\lambda > 0$ small enough, dependent on $\|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c}$. Recalling definition (5.200) of Bol_{fec} , the main term is given by

$$\frac{\pi T}{(2\pi)^6 N} \int_{p_1 \perp p_2} d\mathcal{H}^5(\mathbf{p}_2) \Delta_{cub} J(\mathbf{p}_2) \frac{(\hat{v}(p_1) + \hat{v}(p_2))^2}{|\mathbf{p}_2|} \left(\tilde{f}_0(p_1) \tilde{f}_0(p_2) f_0(p_1 + p_2) - f_0(p_1) f_0(p_2) \tilde{f}_0(p_1 + p_2) \right), \quad (6.28)$$

and it is of size N^{-1} . In order to have that the discretization error is negligible, we need to impose $\frac{1}{\lambda} = o(|\Lambda|^{\frac{r}{6(r+1)}})$ as $|\Lambda| \rightarrow \infty$. In particular, we may choose

$$L = \lambda^{-2 - \frac{2}{r} - \varepsilon} \quad (6.29)$$

for any arbitrary $\varepsilon > 0$. Recall that we required

$$\frac{|\Lambda|}{\lambda} = O\left(\frac{\log N}{\log \log N}\right) \quad (6.30)$$

as $N \rightarrow \infty$ to suppress the tail in the Duhamel expansion. We may thus choose

$$\lambda = \left(\frac{\log \log N}{\log N}\right)^{\frac{r}{(7+\varepsilon)r+6}}, \quad (6.31)$$

$$L = \left(\frac{\log N}{\log \log N}\right)^{\frac{(2+\varepsilon)r+2}{(7+\varepsilon)r+6}}. \quad (6.32)$$

6.2.3. *Main order term of g .* Propositions 5.1 and 5.7 imply

$$\begin{aligned} & \int_{\Lambda^*} dp g \frac{\langle \Phi \rangle}{\lambda^2}(p) J(p) \\ &= \frac{1}{N} \int_0^T dS \text{abs}_{quart,c}(f_0)[J](S/\lambda^2) \\ & \quad + \frac{1}{N\lambda^2} \int_{\Delta[T,2]} d\mathbf{S}_2 \left(\text{col}_c(f_0)[J](\mathbf{S}_2/\lambda^2) + \text{abs}_{cub,c}(f_0)[J](\mathbf{S}_2/\lambda^2) \right) \\ & \quad + \text{Rem}_2\left(\frac{T}{\lambda^2}; g[J]\right) + \text{err}_{1,c}^{(dis)}\left(\frac{T}{\lambda^2}; g[J]\right) \\ & \quad + \text{err}_{2,c}^{(dis)}\left(\frac{T}{\lambda^2}; g[J]\right) + J(0) \text{err}_{2,c}^{(Bog,Con)}\left(\frac{T}{\lambda^2}; g\right) \end{aligned} \quad (6.33)$$

with

$$|\text{Rem}_2\left(\frac{T}{\lambda^2}; g[J]\right)| \leq C_{\|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c}, \|f_0\|_{2(\lfloor \frac{r}{2} \rfloor + 1), c}, r} \langle T \rangle^4 \|J\|_{2(\lfloor \frac{r}{2} \rfloor + 1), c}$$

$$\frac{e^{C_r \|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c} |\Lambda| / \lambda T} |\Lambda|^6 \left(1 + \frac{|\Lambda|}{N}\right)^2}{\lambda^4 N^2}, \quad (6.34)$$

$$|\text{err}_{1,c}^{(dis)}(\frac{T}{\lambda^2}; g[J])| \leq \frac{C_{\|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c}, \|f_0\|_{2(\lfloor \frac{r}{2} \rfloor + 1), c}, r} \langle T \rangle^{r+1} \|J\|_{W^2[\frac{r}{2}] + 2, \infty}}{N \lambda^{2r} |\Lambda|^{\frac{r}{2}}} \quad (6.35)$$

$$|\text{err}_{2,c}^{(dis)}(\frac{T}{\lambda^2}; g[J])| \leq \frac{C_{\|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c}, \|f_0\|_{2(\lfloor \frac{r}{2} \rfloor + 1), c}, r} \langle T \rangle^{r+2} \|J\|_{W^2[\frac{r}{2}] + 2, \infty}}{N \lambda^{2(r+1)} |\Lambda|^{\frac{r}{2}}} \quad (6.36)$$

$$\begin{aligned} |\text{err}_{2,c}^{(Bog, Con)}(\frac{T}{\lambda^2}; g[J])| &\leq C_{\|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c}, \|f_0\|_{2(\lfloor \frac{r}{2} \rfloor + 1), c}, r} e^{C_r \|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c} |\Lambda| / \lambda T} \\ &\quad \frac{\langle T \rangle^4 |\Lambda|^{\frac{3}{2}}}{\lambda^3 N^{\frac{3}{2}}} \left(1 + \frac{|\Lambda|}{N}\right)^2 \left(1 + \frac{|\Lambda|^{\frac{3}{2}}}{\lambda N^{\frac{1}{2}}}\right) \end{aligned} \quad (6.37)$$

for all $\lambda > 0$ small enough, dependent on $\|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c}$. All the errors are suppressed for the choices of $|\Lambda|$ and λ as in the case of f above.

6.2.4. *Conclusion for $L \sim \lambda^{-2-}$.* Recall that we impose

$$\lambda = \left(\frac{\log \log N}{\log N}\right)^{\frac{r}{(7+\varepsilon)r+6}}, \quad (6.38)$$

$$L = \lambda^{-2-\frac{2}{r}-\varepsilon} = \left(\frac{\log N}{\log \log N}\right)^{\frac{(2+\varepsilon)r+2}{(7+\varepsilon)r+6}}. \quad (6.39)$$

Let

$$\delta_{\Psi, c} := \min \left\{ \frac{1}{2} - C_{r, \varepsilon} \frac{\log \frac{1}{\lambda}}{\log N} - \frac{C_r \|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c} T}{\log \log N}, \frac{\log \frac{1}{\lambda}}{\log N} \right\}, \quad (6.40)$$

$$\begin{aligned} \delta_{F, c} := \min \left\{ \frac{1}{2} - C_{r, \varepsilon} \frac{\log \frac{1}{\lambda}}{\log N} - \frac{C_r \|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c} T}{\log \log N}, \frac{\varepsilon r \log \frac{1}{\lambda}}{\log N}, \right. \\ \left. \frac{\log \frac{1}{\lambda} - \log \log \frac{1}{\lambda}}{\log N} \right\}, \end{aligned} \quad (6.41)$$

$$\delta_{G, c} := \min \left\{ \frac{1}{2} - C_{r, \varepsilon} \frac{\log \frac{1}{\lambda}}{\log N} - \frac{C_r \|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c} T}{\log \log N}, \frac{\varepsilon r \log \frac{1}{\lambda}}{\log N} \right\}. \quad (6.42)$$

Observe that we have

$$\|J\|_{W^{2, \infty}(\mathbb{R}^3)} \leq \|J\|_{2(\lfloor \frac{r}{2} \rfloor + 1), c}. \quad (6.43)$$

Then we have proved that, for some $N_0 = N_0(\|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c})$,

$$\begin{aligned} \left| \Phi_{\frac{T}{\lambda^2}} - \frac{1}{N^{\frac{1}{2}} \lambda} \text{Con}_c(f_0)(T; \lambda) \right| \\ \leq \frac{C_{\|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c}, \|f_0\|_{2(\lfloor \frac{r}{2} \rfloor + 1), c}, r, \varepsilon, T}}{N^{\frac{1}{2} + \delta_{\Psi, c}} \lambda}, \end{aligned} \quad (6.44)$$

$$\begin{aligned} \left| \int_{\Lambda^*} dp (f_{\frac{T}{\lambda^2}}^{(\Phi)}(p) - f_0^{(\Phi)}(p)) J(p) - \frac{1}{N} \text{Bol}_{f_{ec}}(f_0)[J](T) \right| \\ \leq \frac{C_{\|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c}, \|f_0\|_{2(\lfloor \frac{r}{2} \rfloor + 1), c}, r, \varepsilon, T} \|J\|_{W^2[\frac{r}{2}] + 2, \infty}}{N^{1 + \delta_{F, c}}}, \end{aligned} \quad (6.45)$$

$$\begin{aligned}
& \left| \int_{\Lambda^*} dp g_{\frac{T}{\lambda^2}}^{(\Phi)}(p) J(p) - \frac{1}{N} \int_0^T dS \text{abs}_{\text{quart},c}(f_0)[J](S/\lambda^2) \right. \\
& \quad - \frac{1}{N\lambda^2} \int_{\Delta[T,2]} d\mathbf{S}_2 (\text{col}_c(f_0)[J](\mathbf{S}_2/\lambda^2) \\
& \quad \left. + \text{abs}_{\text{cub},c}(f_0)[J](\mathbf{S}_2/\lambda^2)) \right| \\
& \leq \frac{C_{\|\hat{v}\|_{2(\lfloor \frac{T}{2} \rfloor + 1), w, c}, \|f_0\|_{2(\lfloor \frac{T}{2} \rfloor + 1), c}, r, \varepsilon, T} \|J\|_{2(\lfloor \frac{T}{2} \rfloor + 1), c}}{N^{1+\delta_{G,c}}} \quad (6.46)
\end{aligned}$$

for all $N \geq N_0(\|\hat{v}\|_{2(\lfloor \frac{T}{2} \rfloor + 1), w, c})$. Observe that for all N large enough, we have that $\delta_{j,c} = \delta_{j,c}(N, T) > 0$ for $j \in \{\Psi, F, G\}$.

7. EFFECTIVE EQUATIONS

After establishing the sizes of the leading order terms, we can derive the effective equations. This will prove Theorem 2.1.

7.1. Discrete case. Let $L \geq 1$ be fixed. Choose $\lambda = \log \log(N)/\log(N)$ as explained in section 6.1.4.

7.1.1. Evolution of Φ . We have that

$$\begin{aligned}
\Phi_{\frac{T}{\lambda^2}} &= \frac{1}{N^{\frac{1}{2}}\lambda} \text{Con}_d(f^{(\Phi)})(T; \lambda) \\
& \quad + \text{Rem}_2\left(\frac{T}{\lambda^2}; \Phi\right) + \text{err}_{1,d}^{(\text{Bog})}\left(\frac{T}{\lambda^2}; \Phi\right) + \text{err}_{1,d}^{(\text{cen})}\left(\frac{T}{\lambda^2}; \Phi\right), \quad (7.1)
\end{aligned}$$

see (6.1). Proposition 5.9 implies

$$\begin{aligned}
|\text{err}_{1,d}^{(\text{cen})}\left(\frac{T}{\lambda^2}; \Phi\right)| &\leq C_{\|\hat{v}\|_{w,d}, \|f_0\|_d} \langle T \rangle^3 \frac{|\Lambda|^{\frac{3}{2}} e^{C\|\hat{v}\|_{w,d}|\Lambda|/\lambda T}}{N\lambda^2} \\
& \quad \left(1 + \frac{|\Lambda|}{N}\right) \left(1 + \frac{1}{N^{\frac{1}{2}}\lambda|\Lambda|^{\frac{3}{2}}}\right) \\
& \leq \frac{C_{\|\hat{v}\|_{w,d}, \|f_0\|_{2(\lfloor \frac{T}{2} \rfloor + 1), c}, |\Lambda|, T}}{N^{\frac{1}{2}+\delta_{\Psi,d}}\lambda}. \quad (7.2)
\end{aligned}$$

for all N large enough. Observe that

$$\frac{1}{N^{\frac{1}{2}}\lambda} \text{Con}_d(f^{(\Phi)})(T; \lambda) = -\frac{i}{N^{\frac{1}{2}}\lambda} \int_0^T dS \int_{\Lambda^*} dp \hat{v}(p) f_{\frac{S}{\lambda^2}}^{(\Phi)}(p). \quad (7.3)$$

Then, analogously to (6.16) and employing (7.2), we obtain that

$$\left| \Phi_{\frac{T}{\lambda^2}} - \frac{1}{N^{\frac{1}{2}}\lambda} \text{Con}_d(f^{(\Phi)})(T; \lambda) \right| \leq \frac{C_{\|\hat{v}\|_{w,d}, \|f_0\|_d, |\Lambda|, T}}{N^{\frac{1}{2}+\delta_{\Psi,d}}\lambda}. \quad (7.4)$$

7.1.2. Evolution of f . (6.5) and Proposition 5.9 imply

$$\begin{aligned}
& \int_{\Lambda^*} dp \left(f_{\frac{T}{\lambda^2}}^{(\Phi)}(p) - f_0^{(\Phi)}(p) \right) J(p) \\
&= \frac{1}{N} \left(\text{Bol}_d(f^{(\Phi)})[J](T; \lambda) + \sum_{j=1}^2 \lambda^j \text{Bol}_d^{(1)}(f^{(\Phi)})[J](T; \lambda) \right) \\
& \quad + \text{Rem}_2\left(\frac{T}{\lambda^2}; f[J]\right) + \text{err}_{2,d}^{(\text{Bog}, \text{Bol})}\left(\frac{T}{\lambda^2}; f[J]\right)
\end{aligned}$$

$$+ J(0) \text{err}_{2,d}^{(Bog,Con)}\left(\frac{T}{\lambda^2}; f\right) + \text{err}_{2,d}^{(cen)}\left(\frac{T}{\lambda^2}; f[J]\right). \quad (7.5)$$

Recall that, due to Proposition 5.9,

$$\begin{aligned} |\text{err}_{2,d}^{(cen)}\left(\frac{T}{\lambda^2}; f[J]\right)| &\leq C_{\|\hat{v}\|_{w,d}, \|f_0\|_d} \|J\|_{\ell^\infty(\Lambda^*)} \langle T \rangle^4 \frac{|\Lambda|^{\frac{3}{2}} e^{C\|\hat{v}\|_{w,d}|\Lambda|/\lambda T}}{N^{\frac{3}{2}} \lambda^3} \\ &\quad \left(1 + \frac{|\Lambda|}{N}\right) \left(1 + \frac{1}{N^{\frac{1}{2}} \lambda |\Lambda|^{\frac{3}{2}}}\right) \\ &\quad \left[1 + \langle T \rangle^2 \frac{|\Lambda|^{\frac{3}{2}} e^{C\|\hat{v}\|_{w,d}|\Lambda|/\lambda T}}{N^{\frac{1}{2}} \lambda} \left(1 + \frac{|\Lambda|}{N}\right) \left(1 + \frac{1}{N^{\frac{1}{2}} \lambda |\Lambda|^{\frac{3}{2}}}\right)\right] \\ &\leq \frac{C_{\|\hat{v}\|_{w,d}, \|f_0\|_d, |\Lambda|, T} \|J\|_{\ell^\infty(\Lambda^*)}}{N^{1+\delta_{F,d}(N)}}, \end{aligned} \quad (7.6)$$

where we possibly enlarge the constant in the definition of $\delta_{F,d}$. In particular, similar to (6.17) we obtain that

$$\begin{aligned} &\left| \int_{\Lambda^*} dp \left(f \frac{(\Phi)}{\lambda^2}(p) - f_0^{(\Phi)}(p) \right) J(p) - \frac{1}{N} \left(\text{Bol}_d(f^{(\Phi)})[J](T; \lambda) \right. \right. \\ &\quad \left. \left. + \sum_{j=1}^2 \lambda^j \text{Bol}_d^{(j)}(f^{(\Phi)})[J](T; \lambda) \right) \right| \\ &\leq \frac{C_{\|\hat{v}\|_{w,d}, \|f_0\|_d, |\Lambda|, T} \|J\|_{\ell^\infty(\Lambda^*)}}{N^{1+\delta_{F,d}}}. \end{aligned} \quad (7.7)$$

7.1.3. *Evolution of g .* (6.10) yields

$$\begin{aligned} &\int_{\Lambda^*} dp g \frac{(\Phi)}{\lambda^2}(p) J(p) \\ &= \frac{1}{N} \int_0^T dS \text{abs}_{\text{quart},d}(f^{(\Phi)})[J](S/\lambda^2) \\ &\quad + \frac{1}{N\lambda^2} \int_{\Delta[T,2]} d\mathbf{S}_2 \left(\text{col}_d(f^{(\Phi)})[J](\mathbf{S}_2/\lambda^2) + \text{abs}_{\text{cub},d}(f^{(\Phi)})[J](\mathbf{S}_2/\lambda^2) \right) \\ &\quad + \text{Rem}_2\left(\frac{T}{\lambda^2}; g[J]\right) + J(0) \text{err}_{2,d}^{(Bog,Con)}\left(\frac{T}{\lambda^2}; g\right) \\ &\quad + \text{err}_1^{(cen)}\left(\frac{T}{\lambda^2}; g[J]\right) + \text{err}_2^{(cen)}\left(\frac{T}{\lambda^2}; g[J]\right), \end{aligned} \quad (7.8)$$

where, due to Proposition 5.9 and analogously to (7.6),

$$\begin{aligned} &|\text{err}_1^{(cen)}\left(\frac{T}{\lambda^2}; g[J]\right)|, |\text{err}_2^{(cen)}\left(\frac{T}{\lambda^2}; g[J]\right)| \\ &\leq \frac{C_{\|\hat{v}\|_{w,d}, \|f_0\|_d, |\Lambda|, T} \|J\|_{\ell^\infty(\Lambda^*)}}{N^{1+\delta_{G,d}}}, \end{aligned} \quad (7.9)$$

where we possibly enlarge the constant in the definition of $\delta_{G,d}$. As a consequence of (6.18), we thus obtain that

$$\begin{aligned} &\left| \int_{\Lambda^*} dp g \frac{(\Phi)}{\lambda^2}(p) J(p) - \frac{1}{N} \int_0^T dS \text{abs}_{\text{quart},d}(f^{(\Phi)})[J](S/\lambda^2) \right. \\ &\quad \left. - \frac{1}{N\lambda^2} \int_{\Delta[T,2]} d\mathbf{S}_2 \left(\text{col}_d(f^{(\Phi)})[J](\mathbf{S}_2/\lambda^2) + \text{abs}_{\text{cub},d}(f^{(\Phi)})[J](\mathbf{S}_2/\lambda^2) \right) \right| \end{aligned}$$

$$\leq \frac{C_{\|\hat{v}\|_{w,d}, \|f_0\|_d, |\Lambda|, T} \|J\|_{2 \cap \infty, d}}{N^{1+\delta_{G,d}(N)}}. \quad (7.10)$$

7.1.4. *Conclusion for $L \sim 1$.* Recall that $\lambda = \log \log(N)/\log(N)$. We consider the effective quantities

$$\Psi_T = \Phi_{\frac{T}{\lambda^2}}, \quad (7.11)$$

$$F_T = f_{\frac{T}{\lambda^2}}^{(\Phi)}, \quad (7.12)$$

$$G_T = g_{\frac{T}{\lambda^2}}^{(\Phi)}. \quad (7.13)$$

We have that

$$\begin{aligned} \lambda^2 \operatorname{Re} \int_{[\Delta[T\lambda^{-2}, 2]]} d\mathbf{S}_2 e^{-i\omega(s_1-s_2)} h(s_2) &= \lambda^2 \int_0^{\frac{T}{\lambda^2}} ds \frac{\sin(\omega(\frac{T}{\lambda^2} - s))}{\omega} h(s) \\ &= \int_0^T dS \frac{\sin\left(\frac{\omega}{\lambda^2}(T - S)\right)}{\omega} h\left(\frac{S}{\lambda^2}\right). \end{aligned} \quad (7.14)$$

For a function $H_t(p)$ and $j \in \{1, 2\}$, denote

$$\begin{aligned} Q_{d;T-S,\lambda}(H_S)[J] &= \int_{(\Lambda^*)^2} d\mathbf{p}_2 \frac{\sin\left(\frac{\Omega(p_1)+\Omega(p_2)-\Omega(p_1+p_2)}{\lambda^2}(T-S)\right)}{\Omega(p_1) + \Omega(p_2) - \Omega(p_1+p_2)} \\ &\quad (\hat{v}(p_1) + \hat{v}(p_2))^2 (J(p_1) + J(p_2) - J(p_1+p_2)) \\ &\quad (\widetilde{H}_S(p_1)\widetilde{H}_S(p_2)H_S(p_1+p_2) \\ &\quad - H_S(p_1)H_S(p_2)\widetilde{H}_S(p_1+p_2)), \end{aligned} \quad (7.15)$$

$$q_{d;F;\mathbf{S}_2,\lambda}^{(j)}(H_{S_2})[J] = \operatorname{bol}^{(j)}(H_{\cdot,\lambda^2})[J](\mathbf{S}_2/\lambda^2), \quad (7.16)$$

$$Q_{d;G;T,\lambda}(H)[J] = \frac{1}{\lambda^2} \int_{\Delta[T,2]} d\mathbf{S}_2 \operatorname{col}_d(H_{\cdot,\lambda^2})[J](\mathbf{S}_2/\lambda^2), \quad (7.17)$$

$$\begin{aligned} \mathcal{A}_{d;T,\lambda}(H)[J] &= \int_0^T dS \operatorname{abs}_{\text{quart},d}(H_{\cdot,\lambda^2})[J](S/\lambda^2) \\ &\quad + \frac{1}{\lambda^2} \int_{\Delta[T,2]} d\mathbf{S}_2 \operatorname{abs}_{\text{cub},d}(H_{\cdot,\lambda^2})[J](\mathbf{S}_2/\lambda^2), \end{aligned} \quad (7.18)$$

where we used Proposition 5.2. $Q_{d;G;T,\lambda}$ denotes a generalized collision operator and $\mathcal{A}_{d;T,\lambda}$ a generalized absorption operator. We have proved that

$$\begin{aligned} \left| \Psi_T + \frac{i}{N^{\frac{1}{2}}\lambda} \int_0^T dS \int_{\Lambda^*} dp \hat{v}(p) F_S(p) \right| \\ \leq \frac{C_{\|\hat{v}\|_{w,d}, \|f_0\|_d, |\Lambda|, T}}{N^{\frac{1}{2}+\delta_{\Psi,d}}\lambda}, \end{aligned} \quad (7.19)$$

$$\begin{aligned} \left| \int_{\Lambda^*} dp (F_T(p) - F_0(p)) J(p) - \frac{1}{N} \left(\int_0^T dS Q_{d;T-S,\lambda}(F_S)[J] \right. \right. \\ \left. \left. + \int_{\Delta[T,2]} d\mathbf{S}_2 \sum_{j=1}^2 \lambda^j q_{d;F;\mathbf{S}_2,\lambda}^{(j)}(F_{S_2})[J] \right) \right| \\ \leq \frac{C_{\|\hat{v}\|_{w,d}, \|f_0\|_d, |\Lambda|, T} \|J\|_{\ell^\infty(\Lambda^*)}}{N^{1+\delta_{F,d}}}, \end{aligned} \quad (7.20)$$

$$\begin{aligned} & \left| \int_{\Lambda^*} dp G_T(p) J(p) - \frac{1}{N} \left(\mathcal{A}_{d;T,\lambda}(F)[J] + \mathcal{Q}_{d,G;T,\lambda}(F)[J] \right) \right| \\ & \leq \frac{C_{\|\hat{v}\|_{w,d}, \|f_0\|_d, |\Lambda|, T} \|J\|_{2 \cap \infty, d}}{N^{1+\delta_{G,d}}} \end{aligned} \quad (7.21)$$

for all $N > 0$ larger than a universal constant.

7.2. Continuum approximation. Recall that

$$\lambda = \left(\frac{\log \log N}{\log N} \right)^{\frac{r}{(7+\varepsilon)r+6}}, \quad (7.22)$$

$$L = \lambda^{-2-\frac{2}{r}-\varepsilon} = \left(\frac{\log N}{\log \log N} \right)^{\frac{(2+\varepsilon)r+2}{(7+\varepsilon)r+6}}. \quad (7.23)$$

7.2.1. *Evolution of Φ .* (6.20) and Proposition 5.9 imply that

$$\begin{aligned} \Phi_{\frac{T}{\lambda^2}} &= \frac{1}{N^{\frac{1}{2}} \lambda} \text{Con}_c(f^{(\Phi)})(T; \lambda) \\ &+ \text{Rem}_2\left(\frac{T}{\lambda^2}; \Phi\right) + \text{err}_{1,c}^{(Bog)}\left(\frac{T}{\lambda^2}; \Phi\right) + \text{err}_1^{(cen)}(t; \Phi). \end{aligned} \quad (7.24)$$

We have that

$$\begin{aligned} |\text{err}_1^{(cen)}(t; \Phi)| &\leq C_{\|\hat{v}\|_{2(\lfloor \frac{T}{\lambda^2} \rfloor + 1), w, c}, \|f_0\|_{2(\lfloor \frac{T}{\lambda^2} \rfloor + 1), c}, r} \langle T \rangle^3 \frac{|\Lambda|^{\frac{3}{2}} e^{C_r \|\hat{v}\|_{2(\lfloor \frac{T}{\lambda^2} \rfloor + 1), w, c} |\Lambda| / \lambda T}}{N \lambda^2} \\ &\quad \left(1 + \frac{|\Lambda|}{N} \right) \left(1 + \frac{1}{N^{\frac{1}{2}} \lambda |\Lambda|^{\frac{3}{2}}} \right) \\ &\leq \frac{C_{\|\hat{v}\|_{2(\lfloor \frac{T}{\lambda^2} \rfloor + 1), w, c}, \|f_0\|_{2(\lfloor \frac{T}{\lambda^2} \rfloor + 1), c}, r, T}}{N^{\frac{1}{2} + \delta_{\Psi, c}} \lambda} \end{aligned} \quad (7.25)$$

by possibly enlarging the constant C in the definition (6.40) of $\delta_{\Psi, c}$. Combining this inequality with (6.44), we find that,

$$\left| \Phi_{\frac{T}{\lambda^2}} - \frac{1}{N^{\frac{1}{2}} \lambda} \text{Con}_c(f^{(\Phi)})(T; \lambda) \right| \leq \frac{C_{\|\hat{v}\|_{2(\lfloor \frac{T}{\lambda^2} \rfloor + 1), w, c}, \|f_0\|_{2(\lfloor \frac{T}{\lambda^2} \rfloor + 1), c}, r, \varepsilon, T}}{N^{\frac{1}{2} + \delta_{\Psi, c}} \lambda}. \quad (7.26)$$

7.2.2. *Evolution of f .* As a consequence of (6.23), and Proposition 5.8, 5.9, we have that

$$\begin{aligned} & \int_{\Lambda^*} dp \left(f_{\frac{T}{\lambda^2}}^{(\Phi)}(p) - f_0^{(\Phi)}(p) \right) J(p) \\ &= \frac{1}{N} \text{Bol}_c(f^{(\Phi)})(J)(T; \lambda) \\ &+ \text{Rem}_2\left(\frac{T}{\lambda^2}; f[J]\right) + \text{err}_{2,c}^{(Bog, Bol)}\left(\frac{T}{\lambda^2}; f[J]\right) \\ &+ J(0) \text{err}_2^{(Bog, Con)}\left(\frac{T}{\lambda^2}; f\right) + \text{err}_{2,c}^{(cen)}\left(\frac{T}{\lambda^2}; f[J]\right). \end{aligned} \quad (7.27)$$

Observe that

$$\begin{aligned} |\text{err}_{2,c}^{(cen)}\left(\frac{T}{\lambda^2}; f[J]\right)| &\leq C_{\|\hat{v}\|_{2(\lfloor \frac{T}{\lambda^2} \rfloor + 1), w, c}, \|f_0\|_{2(\lfloor \frac{T}{\lambda^2} \rfloor + 1), c}, r} \|J\|_{\infty} \langle T \rangle^4 \frac{|\Lambda|^{\frac{3}{2}} e^{C_r \|\hat{v}\|_{2(\lfloor \frac{T}{\lambda^2} \rfloor + 1), w, c} |\Lambda| / \lambda T}}{N^{\frac{3}{2}} \lambda^3} \\ &\quad \left(1 + \frac{|\Lambda|}{N} \right) \left(1 + \frac{1}{N^{\frac{1}{2}} \lambda |\Lambda|^{\frac{3}{2}}} \right) \end{aligned}$$

$$\begin{aligned}
& \left[1 + \langle T \rangle^2 \frac{|\Lambda|^{\frac{3}{2}} e^{C_r \|\hat{v}\|_{2(\lfloor \frac{T}{2} \rfloor + 1), w, c} |\Lambda| / \lambda T}}{N^{\frac{1}{2}} \lambda} \left(1 + \frac{|\Lambda|}{N} \right) \left(1 + \frac{1}{N^{\frac{1}{2}} \lambda |\Lambda|^{\frac{3}{2}}} \right) \right] \\
& \leq \frac{C_{\|\hat{v}\|_{2(\lfloor \frac{T}{2} \rfloor + 1), w, c}, \|f_0\|_{2(\lfloor \frac{T}{2} \rfloor + 1), c}, r, T} \|J\|_{\infty}}{N^{1 + \delta_{F, c}}} \tag{7.28}
\end{aligned}$$

by possibly enlarging the constants in the definition (6.41) of $\delta_{F, c}(N, \lambda)$. Then, (6.45) yields

$$\begin{aligned}
& \left| \int_{\Lambda^*} dp \left(f_{\frac{T}{\lambda^2}}^{(\Phi)}(p) - f_0^{(\Phi)}(p) \right) J(p) - \frac{1}{N} \text{Bol}_c(f^{(\Phi)})[J](T; \lambda) \right| \\
& \leq \frac{C_{\|\hat{v}\|_{2(\lfloor \frac{T}{2} \rfloor + 1), w, c}, \|f_0\|_{2(\lfloor \frac{T}{2} \rfloor + 1), c}, r, \varepsilon, T} \|J\|_{W^2[\frac{T}{2}] + 2, \infty}}{N^{1 + \delta_{F, c}}}. \tag{7.29}
\end{aligned}$$

7.2.3. *Evolution of g .* (6.33) and Proposition 5.9 imply

$$\begin{aligned}
& \int_{\Lambda^*} dp g_{\frac{T}{\lambda^2}}^{(\Phi)}(p) J(p) \\
& = \frac{1}{N} \int_0^T dS \text{abs}_{\text{quart}, c}(f^{(\Phi)})[J](S/\lambda^2) \\
& \quad + \frac{1}{N\lambda^2} \int_{\Delta[T, 2]} d\mathbf{S}_2 \left(\text{col}_c(f^{(\Phi)})[J](\mathbf{S}_2/\lambda^2) + \text{abs}_{\text{cub}, c}(f^{(\Phi)})[J](\mathbf{S}_2/\lambda^2) \right) \\
& \quad + \text{Rem}_2\left(\frac{T}{\lambda^2}; g[J]\right) + \text{err}_1^{(\text{dis})}\left(\frac{T}{\lambda^2}; g[J]\right) + \text{err}_2^{(\text{dis})}\left(\frac{T}{\lambda^2}; g[J]\right) \\
& \quad + J(0) \text{err}_2^{(\text{Bog}, \text{Con})}\left(\frac{T}{\lambda^2}; g\right) + \text{err}_1^{(\text{cen})}\left(\frac{T}{\lambda^2}; g[J]\right) + \text{err}_2^{(\text{cen})}\left(\frac{T}{\lambda^2}; g[J]\right). \tag{7.30}
\end{aligned}$$

Analogously to (7.28), we have that

$$\begin{aligned}
& \left| \text{err}_1^{(\text{cen})}\left(\frac{T}{\lambda^2}; g[J]\right) \right|, \left| \text{err}_2^{(\text{cen})}\left(\frac{T}{\lambda^2}; g[J]\right) \right| \\
& \leq \frac{C_{\|\hat{v}\|_{2(\lfloor \frac{T}{2} \rfloor + 1), w, c}, \|f_0\|_{2(\lfloor \frac{T}{2} \rfloor + 1), c}, r, \varepsilon, T} \|J\|_{\infty}}{N^{1 + \delta_{G, c}}}, \tag{7.31}
\end{aligned}$$

by again possibly enlarging the constants in the definition (6.42) of $\delta_{G, c}$. Applying (6.46), we obtain that

$$\begin{aligned}
& \left| \int_{\Lambda^*} dp g_{\frac{T}{\lambda^2}}^{(\Phi)}(p) J(p) - \frac{1}{N} \int_0^T dS \text{abs}_{\text{quart}, c}(f^{(\Phi)})[J](S/\lambda^2) \right. \\
& \quad \left. - \frac{1}{N\lambda^2} \int_{\Delta[T, 2]} d\mathbf{S}_2 \left(\text{col}_c(f^{(\Phi)})[J](\mathbf{S}_2/\lambda^2) \right) \right. \\
& \quad \left. + \text{abs}_{\text{cub}, c}(f^{(\Phi)})[J](\mathbf{S}_2/\lambda^2) \right| \\
& \leq \frac{C_{\|\hat{v}\|_{2(\lfloor \frac{T}{2} \rfloor + 1), w, c}, \|f_0\|_{2(\lfloor \frac{T}{2} \rfloor + 1), c}, r, \varepsilon, T} \|J\|_{2(\lfloor \frac{T}{2} \rfloor + 1), c}}{N^{1 + \delta_{G, c}}}. \tag{7.32}
\end{aligned}$$

7.2.4. *Conclusion for $L \sim \lambda^{-2-}$.* Recall that we impose

$$\lambda = \left(\frac{\log \log N}{\log N} \right)^{\frac{r}{(7+\varepsilon)r+6}}, \tag{7.33}$$

$$L = \lambda^{-2 - \frac{2}{r} - \varepsilon} = \left(\frac{\log N}{\log \log N} \right)^{\frac{(2+\varepsilon)r+2}{(7+\varepsilon)r+6}}. \tag{7.34}$$

Again, we consider the effective quantities

$$\Psi_T = \Phi_{\frac{T}{\lambda^2}}, \quad (7.35)$$

$$F_T = f_{\frac{T}{\lambda^2}}^{(\Phi)}, \quad (7.36)$$

$$G_T = g_{\frac{T}{\lambda^2}}^{(\Phi)}. \quad (7.37)$$

Let $Q_{c;T-S,\lambda}$, $Q_{c;G;T,\lambda}$ and $\mathcal{A}_{c;T,\lambda}$ be defined analogously to (2.18), (2.15) respectively (2.17) with sums over Λ^* replaced by integrals $\frac{1}{(2\pi)^3} \int_{\mathbb{R}^3}$ over \mathbb{R}^3 . Collecting (7.26), (7.29), and (7.32), we have proved that, for some possibly larger constant $N_0 = N_0(\|\hat{v}\|_{2(\lfloor \frac{T}{2} \rfloor + 1), w, c})$,

$$\begin{aligned} \left| \Psi_T + \frac{i}{(2\pi)^3 N^{\frac{1}{2}} \lambda} \int_0^T dS \int_{\mathbb{R}^3} dp \hat{v}(p) F_S(p) \right| \\ \leq \frac{C_{\|\hat{v}\|_{2(\lfloor \frac{T}{2} \rfloor + 1), w, c}, \|f_0\|_{2(\lfloor \frac{T}{2} \rfloor + 1), c, r, \varepsilon, T}}}{N^{\frac{1}{2} + \delta_{\Psi, c}} \lambda}, \end{aligned} \quad (7.38)$$

$$\begin{aligned} \left| \int_{\Lambda^*} dp (F_T(p) - F_0(p)) J(p) - \frac{1}{N} \int_0^T dS Q_{c;T-S,\lambda}(F_S) \right| \\ \leq \frac{C_{\|\hat{v}\|_{2(\lfloor \frac{T}{2} \rfloor + 1), w, c}, \|f_0\|_{2(\lfloor \frac{T}{2} \rfloor + 1), c, r, \varepsilon, T} \|J\|_{W^2[\lfloor \frac{T}{2} \rfloor + 2, \infty]}}}{N^{1 + \delta_{F, c}}}, \end{aligned} \quad (7.39)$$

$$\begin{aligned} \left| \int_{\Lambda^*} dp G_T(p) J(p) - \frac{1}{N} (\mathcal{A}_{c;T,\lambda}(F)[J] + \mathcal{Q}_{c;G;T,\lambda}(F)[J]) \right| \\ \leq \frac{C_{\|\hat{v}\|_{2(\lfloor \frac{T}{2} \rfloor + 1), w, c}, \|f_0\|_{2(\lfloor \frac{T}{2} \rfloor + 1), c, r, \varepsilon, T} \|J\|_{2(\lfloor \frac{T}{2} \rfloor + 1), c}}}{N^{1 + \delta_{G, c}}} \end{aligned} \quad (7.40)$$

for all $N \geq N_0$.

APPENDIX A. CALCULUS FOR CREATION AND ANNIHILATION OPERATORS

Lemma A.1. *Let ν be a translation invariant state, i.e.,*

$$\nu(A) = \nu(e^{ix \cdot \mathcal{P}} A e^{-ix \cdot \mathcal{P}}) \quad (A.1)$$

for all $x \in \mathbb{R}^3$ and all observables A . Then we have

$$\nu\left(\prod_{i=1}^m a_{p_i}^{(\sigma_i)}\right) = \frac{\delta(\sum_{i=1}^m \sigma_i p_i)}{|\Lambda|} \nu\left(\prod_{i=1}^m a_{p_i}^{(\sigma_i)}\right). \quad (A.2)$$

Proof. By translation invariance, we have that

$$\begin{aligned} \nu\left(\prod_{i=1}^m a_{p_i}^{(\sigma_i)}\right) &= \nu\left(e^{ix \cdot \mathcal{P}} \prod_{i=1}^m a_{p_i}^{(\sigma_i)} e^{-ix \cdot \mathcal{P}}\right) \\ &= e^{ix \cdot (\sum_{i=1}^m \sigma_i p_i)} \nu\left(\prod_{i=1}^m a_{p_i}^{(\sigma_i)}\right) \end{aligned} \quad (A.3)$$

for all $x \in \mathbb{R}^3$. Integrating both sides $\int_{\Lambda} dx$, we obtain

$$|\Lambda| \nu\left(\prod_{i=1}^m a_{p_i}^{(\sigma_i)}\right) = \delta\left(\sum_{i=1}^m \sigma_i p_i\right) \nu\left(\prod_{i=1}^m a_{p_i}^{(\sigma_i)}\right), \quad (A.4)$$

which yields the statement. \square

Lemma A.2 (Cumulant Formula). *Recall from (4.13) that*

$$\kappa_n = (-\partial_\mu)^n \Big|_{\mu=0} \int_{\Lambda^*} dp \log(1 - e^{-(K(p)-\mu)}). \quad (\text{A.5})$$

Then there are constants $a_{n,k} \in \mathbb{R}$ such that

$$\kappa_n = \sum_{k=1}^{n-1} a_{n,k} \int_{\Lambda^*} dp f_0(p)^k (1 + f_0(p))^{n-k} + \delta_{n,1} \int_{\Lambda^*} dp f_0(p), \quad (\text{A.6})$$

where $f_0(p) = (e^{K(p)} - 1)^{-1}$.

Proof. We will show, in more generality that

$$\kappa_n(\mu) := (-\partial_\mu)^n \int_{\Lambda^*} dp \log(1 - e^{-(K(p)-\mu)}) \quad (\text{A.7})$$

$$= \sum_{k=1}^{n-1} a_{n,k} \int_{\Lambda^*} dp f_\mu(p)^k (1 + f_\mu(p))^{n-k} + \delta_{n,1} \int_{\Lambda^*} dp f_\mu(p) \quad (\text{A.8})$$

for all $\mu \leq 0$, where $f_\mu(p) = (e^{K(p)-\mu} - 1)^{-1}$. A straightforward calculation yields

$$\kappa_1(\mu) = \int_{\Lambda^*} dp f_\mu(p), \quad (\text{A.9})$$

$$\kappa_2(\mu) = \int_{\Lambda^*} dp f_\mu(p)(1 + f_\mu(p)). \quad (\text{A.10})$$

Observe that

$$-\partial_\mu f_\mu = f_\mu(1 + f_\mu). \quad (\text{A.11})$$

Now assume that (A.8) for some fixed $n \in \mathbb{N}$, $n \geq 2$. By definition, we have that

$$\begin{aligned} \kappa_{n+1}(\mu) &= (-\partial_\mu) \kappa_n(\mu) \\ &= \int_{\Lambda^*} dp \sum_{k=1}^{n-1} a_{n,k} (k f_\mu(p)^k (1 + f_\mu(p))^{n+1-k} \\ &\quad + (n-k) f_\mu(p)^{k+1} (1 + f_\mu(p))^{n-k}). \end{aligned} \quad (\text{A.12})$$

After an index shift, we can further simplify this to

$$\begin{aligned} \kappa_{n+1}(\mu) &= a_{n,1} \int_{\Lambda^*} dp f_\mu(p)(1 + f_\mu(p))^n \\ &\quad + a_{n,n-1} \int_{\Lambda^*} dp f_\mu(p)^n (1 + f_\mu(p)) \\ &\quad + \sum_{k=2}^{n-1} (k a_{n,k} + (n+1-k) a_{n,k-1}) \\ &\quad \int_{\Lambda^*} dp f_\mu(p)^k (1 + f_\mu(p))^{n+1-k} \\ &= \sum_{k=1}^n a_{n+1,k} \int_{\Lambda^*} dp f_\mu(p)^k (1 + f_\mu(p))^{n+1-k}, \end{aligned} \quad (\text{A.13})$$

for some $a_{n+1,k} \in \mathbb{R}$. This finishes the proof. \square

For the following standard result, we need to introduce some notation. For a proof of the statement, we refer, e.g., to [34, 35]. Given a finite ordered subset $J = \{j_1 < j_2 < \dots < j_r\} \subset \mathbb{N}$ and $\sigma_{j_k} \in \{\pm 1\}$, we define the ordered product

$$\prod_{j \in J} a_{p_j}^{(\sigma_j)} := a_{p_{j_1}}^{(\sigma_{j_1})} \dots a_{p_{j_r}}^{(\sigma_{j_r})}. \quad (\text{A.14})$$

In addition, we abbreviate

$$\mathbf{p}_J := (p_{j_k})_{k=1}^r, \quad (\text{A.15})$$

as well as

$$a^{(\sigma)}(\mathbf{p}_J) := \prod_{j \in J} a_{p_j}^{(\sigma)}. \quad (\text{A.16})$$

Furthermore, we define the sets

$$J_{\pm} := \{j \in J \mid \sigma_j = \pm 1\} \quad (\text{A.17})$$

and the Wick-ordered product

$$: \prod_{j \in J} a_{p_j}^{(\sigma_j)} : := a^+(\mathbf{p}_{J_+}) a(\mathbf{p}_{J_-}) \quad (\text{A.18})$$

with all creation operators to the left, and all annihilation operators to the right.

Finally, in order to keep track of the correct scaling, it is useful to work with the rescaled $\ell^2(\Lambda^*)$ -norm

$$\|H\|_{L^2(\Lambda^*)} = \frac{1}{\sqrt{|\Lambda|}} \|H\|_{\ell^2(\Lambda^*)}, \quad (\text{A.19})$$

see (2.1). More generally, we also define

$$\|H\|_{L_{\mathbf{p}_m}^{\infty} L_{\mathbf{k}_n}^2((\Lambda^*)^{m+n})} := \sup_{\mathbf{p}_m \in (\Lambda^*)^m} \left(\int_{(\Lambda^*)^n} d\mathbf{k}_n |H(\mathbf{p}_m, \mathbf{k}_n)|^2 \right)^{\frac{1}{2}}, \quad (\text{A.20})$$

$$\|H\|_{L_{\mathbf{k}_n}^2 L_{\mathbf{p}_m}^{\infty}((\Lambda^*)^{m+n})} := \left(\int_{(\Lambda^*)^n} d\mathbf{k}_n \sup_{\mathbf{p}_m \in (\Lambda^*)^m} |H(\mathbf{p}_m, \mathbf{k}_n)|^2 \right)^{\frac{1}{2}}, \quad (\text{A.21})$$

where in the case $n = 0$, this norm reduces to $\|H\|_{L_{\mathbf{p}_m}^{\infty}((\Lambda^*)^m)}$, and in the case $m = 0$, to $\|H\|_{L_{\mathbf{k}_n}^2((\Lambda^*)^n)}$.

Lemma A.3 (Wick's Theorem). *Let $\sigma_j \in \{\pm 1\}$, $p_j \in \mathbb{R}^3$ for all $j \in \{1, \dots, n\}$, $n \in \mathbb{N}$. Then we have that*

$$\prod_{j=1}^n a_{p_j}^{(\sigma_j)} = \sum_{J \subseteq \{1, \dots, n\}} \langle \Omega_0, \prod_{j \in J} a_{p_j}^{(\sigma_j)} \Omega_0 \rangle : \prod_{j \in J} a_{p_j}^{(\sigma_j)} :. \quad (\text{A.22})$$

Lemma A.4 (Wick-ordered operator bound). *Let $M \in \mathbb{N}_0$, $n \in \mathbb{N}$, $J := \{1, \dots, n\}$, $\sigma_j \in \{\pm 1\}$ for all $j \in J$. Let $H : (\Lambda^*)^n \rightarrow \mathbb{C}$, and $g_j : \Lambda^* \rightarrow \mathbb{C}$ be given functions. Then the following holds true*

(1) *If $J_{\pm} \neq \emptyset$, we have that*

$$\begin{aligned} & \left\| \int_{(\Lambda^*)^n} d\mathbf{p}_n H(\mathbf{p}_n) \delta\left(\sum_{j=1}^n p_j \sigma_j\right) \prod_{j=1}^n g_j(p_j) : \prod_{j=1}^n a_{p_j}^{(\sigma_j)} : P_M \right\| \\ & \leq \| |H|^{\frac{1}{2}} \prod_{j \in J_-} g_j(p_j) \delta\left(\sum_{j=1}^n \sigma_j p_j\right)^{\frac{1}{2}} \|_{L_{\mathbf{p}_{J_+}}^{\infty} L_{\mathbf{p}_{J_-}}^2} \| |H|^{\frac{1}{2}} \prod_{j \in J_+} g_j(p_j) \delta\left(\sum_{j=1}^n \sigma_j p_j\right)^{\frac{1}{2}} \|_{L_{\mathbf{p}_{J_-}}^{\infty} L_{\mathbf{p}_{J_+}}^2} \end{aligned}$$

$$(M + \sum_{j=1}^n \sigma_j)_{|J_+|}^{\frac{1}{2}} (M)_{|J_-|}^{\frac{1}{2}} \mathbf{1}_{M \geq |J_-|},$$

where

$$(x)_m := \prod_{k=0}^{m-1} (x - k) \quad (\text{A.23})$$

denotes the falling factorial.

(2) If $J_+ = \emptyset$ and $n \geq 2$, we find that

$$\begin{aligned} & \left\| \int_{(\Lambda^*)^n} d\mathbf{p}_n H(\mathbf{p}_n) \delta\left(\sum_{j=1}^n p_j\right) \prod_{j=1}^n a_{p_j} P_M \right\| \\ & \leq \left(\|H\|_{L^2_{\mathbf{p}_{n-2} L^{\infty}_{\mathbf{p}_{n-1}, \mathbf{p}_n}}}^2 (M - n + 1) + |\Lambda| \left\| \delta\left(\sum_{j=1}^n p_j\right)^{\frac{1}{2}} H \right\|_{L^2_{\mathbf{p}_n}}^2 \right)^{\frac{1}{2}} (M)_{n-1}^{\frac{1}{2}}. \end{aligned} \quad (\text{A.24})$$

Similarly, in the case $J_- = \emptyset$, we have that

$$\left\| \int_{(\Lambda^*)^n} d\mathbf{p}_n H(\mathbf{p}_n) \delta\left(\sum_{j=1}^n p_j\right) a^+(\mathbf{p}_n) P_M \right\| \quad (\text{A.25})$$

$$\leq \left(\|H\|_{L^2_{\mathbf{p}_{n-2} L^{\infty}_{\mathbf{p}_{n-1}, \mathbf{p}_n}}}^2 (M + 1) + |\Lambda| \left\| \delta\left(\sum_{j=1}^n p_j\right)^{\frac{1}{2}} H \right\|_{L^2_{\mathbf{p}_n}}^2 \right)^{\frac{1}{2}} (M + n)_{n-1}^{\frac{1}{2}}. \quad (\text{A.26})$$

(3) If $n = 1$, we obtain

$$\|a_0 P_M\| = \sqrt{M|\Lambda|}. \quad (\text{A.27})$$

Proof. Let $\Phi_+ \in \mathcal{F}_{M+\sum_{j=1}^n \sigma_j}$ and $\Phi_- \in \mathcal{F}_M$ be two normalized test functions.

In the case $J_{\pm} \neq \emptyset$, we have that

$$\begin{aligned} & \left| \left\langle \Phi_+, \int_{(\Lambda^*)^n} d\mathbf{p}_n H(\mathbf{p}_n) \delta\left(\sum_{j=1}^n p_j \sigma_j\right) \prod_{j=1}^n g_j(p_j) : \prod_{j=1}^n a_{p_j}^{(\sigma_j)} : \Phi_- \right\rangle \right| \\ & \leq \int_{(\Lambda^*)^n} d\mathbf{p}_n |H(\mathbf{p}_n)| \delta\left(\sum_{j=1}^n p_j \sigma_j\right) \prod_{j=1}^n |g_j(p_j)| \|a(\mathbf{p}_{J_+}) \Phi_+\| \|a(\mathbf{p}_{J_-}) \Phi_-\|, \end{aligned} \quad (\text{A.28})$$

where we applied Cauchy-Schwarz w.r.t. the inner product on \mathcal{F} . Let $\alpha \in [0, 1]$ be arbitrary. Then Cauchy-Schwarz w.r.t. $d\mathbf{p}_n$ implies that we can estimate the last expression by

$$\begin{aligned} & \left(\int_{(\Lambda^*)^n} d\mathbf{p}_n \delta\left(\sum_{j=1}^n p_j \sigma_j\right) |H(\mathbf{p}_n)| \prod_{j \in J_-} |g_j(p_j)|^2 \|a(\mathbf{p}_{J_+}) \Phi_+\|^2 \right)^{\frac{1}{2}} \\ & \left(\int_{(\Lambda^*)^n} d\mathbf{p}_n \delta\left(\sum_{j=1}^n p_j \sigma_j\right) |H(\mathbf{p}_n)| \prod_{j \in J_+} |g_j(p_j)|^2 \|a(\mathbf{p}_{J_-}) \Phi_-\|^2 \right)^{\frac{1}{2}}. \end{aligned} \quad (\text{A.29})$$

Observe that the Pull-Through Formula implies that

$$\begin{aligned} \int d\mathbf{p}_m a^+(\mathbf{p}_m) a(\mathbf{p}_m) &= \int d\mathbf{p}_{m-1} a^+(\mathbf{p}_{m-1}) a(\mathbf{p}_{m-1}) (\mathcal{N}_b - m + 1)_+ \\ &= (\mathcal{N}_b)_m, \end{aligned} \quad (\text{A.30})$$

where we define $(\mathcal{N}_b)_m|_{\mathcal{F}_j} := 0$ for all $j \in \{0, 1, \dots, m-1\}$, in accordance with (3.3).

In particular, we have that $(\mathcal{N}_b)_m \geq 0$ as a quadratic form on \mathcal{F} .

Then, integrating the first factor first w.r.t. the momenta p_{J_-} and then p_{J_+} , and opposite for the second factor, (A.29) has the upper bound

$$\begin{aligned} & \| |H|^{\frac{1}{2}} \prod_{j \in J_-} g_j(p_j) \delta(\sum_{j=1}^n \sigma_j p_j)^{\frac{1}{2}} \|_{L_{\mathbf{p}_{J_+}}^\infty L_{\mathbf{p}_{J_-}}^2} \| (\mathcal{N}_b)_{|J_+|}^{\frac{1}{2}} \Phi_+ \| \\ & \| |H|^{\frac{1}{2}} \prod_{j \in J_+} g_j(p_j) \delta(\sum_{j=1}^n \sigma_j p_j)^{\frac{1}{2}} \|_{L_{\mathbf{p}_{J_-}}^\infty L_{\mathbf{p}_{J_+}}^2} \| (\mathcal{N}_b)_{|J_-|}^{\frac{1}{2}} \Phi_- \| . \end{aligned} \quad (\text{A.31})$$

Using the fact that $\Phi_+ \in \mathcal{F}_{M+\sum_{j=1}^n \sigma_j}$, $\Phi_- \in \mathcal{F}_M$ and that both are normalized with norm 1, we have that

$$\| (\mathcal{N}_b)_{|J_+|}^{\frac{1}{2}} \Phi_+ \| \leq (M + \sum_{j=1}^n \sigma_j)_{|J_+|}^{\frac{1}{2}} \mathbf{1}_{M+\sum_{j=1}^n \sigma_j \geq |J_+|} , \quad (\text{A.32})$$

$$\| (\mathcal{N}_b)_{|J_-|}^{\frac{1}{2}} \Phi_- \| \leq (M)_{|J_-|}^{\frac{1}{2}} \mathbf{1}_{M \geq |J_-|} . \quad (\text{A.33})$$

Observe that $\sum_{j=1}^n \sigma_j = |J_+| - |J_-|$. Collecting (A.28)–(A.33), we have proved that

$$\begin{aligned} & \left| \left\langle \Phi_+ , \int_{(\Lambda^*)^n} d\mathbf{p}_n H(\mathbf{p}_n) \delta(\sum_{j=1}^n p_j \sigma_j) \prod_{j=1}^n g_j(p_j) : \prod_{j=1}^n a_{p_j}^{(\sigma_j)} : \Phi_- \right\rangle \right| \\ & \leq \| |H|^{\frac{1}{2}} \prod_{j \in J_-} g_j(p_j) \delta(\sum_{j=1}^n \sigma_j p_j)^{\frac{1}{2}} \|_{L_{\mathbf{p}_{J_+}}^\infty L_{\mathbf{p}_{J_-}}^2} \| |H|^{\frac{1}{2}} \prod_{j \in J_+} g_j(p_j) \delta(\sum_{j=1}^n \sigma_j p_j)^{\frac{1}{2}} \|_{L_{\mathbf{p}_{J_-}}^\infty L_{\mathbf{p}_{J_+}}^2} \\ & (M + \sum_{j=1}^n \sigma_j)_{|J_+|}^{\frac{1}{2}} (M)_{|J_-|}^{\frac{1}{2}} \mathbf{1}_{M \geq |J_-|} . \end{aligned} \quad (\text{A.34})$$

In the case $J_- = \emptyset$, we have that

$$\left\| \int_{(\Lambda^*)^n} d\mathbf{p}_n H(\mathbf{p}_n) \delta(\sum_{j=1}^n p_j) a^+(\mathbf{p}_n) P_M \right\| \quad (\text{A.35})$$

$$= \left\| \int_{(\Lambda^*)^n} d\mathbf{p}_n \bar{H}(\mathbf{p}_n) \delta(\sum_{j=1}^n p_j) a(\mathbf{p}_n) P_{M+n} \right\| , \quad (\text{A.36})$$

which reduces to the case $J_+ = \emptyset$.

In the case $J_+ = \emptyset$ and $n \geq 2$, we find that

$$\begin{aligned} & \left\| \int_{(\Lambda^*)^n} d\mathbf{p}_n H(\mathbf{p}_n) \delta(\sum_{j=1}^n p_j) a(\mathbf{p}_n) \Phi_- \right\|^2 \\ & = \int_{(\Lambda^*)^{2n}} d\mathbf{p}_n d\mathbf{q}_n \bar{H}(\mathbf{p}_n) H(\mathbf{q}_n) \delta(\sum_{j=1}^n p_j) \delta(\sum_{j=1}^n q_j) \langle \Phi_- , a^+(\mathbf{p}_n) a(\mathbf{q}_n) \Phi_- \rangle \\ & = \int_{(\Lambda^*)^{2n}} d\mathbf{p}_n d\mathbf{q}_n \bar{H}(\mathbf{p}_n) H(\mathbf{q}_n) \delta(\sum_{j=1}^n p_j) \delta(\sum_{j=1}^n q_j) \\ & \quad \langle \Phi_- , a^+(\mathbf{p}_{n-1}) a_{q_n} a_{p_n}^+ a(\mathbf{q}_{n-1}) \Phi_- \rangle \\ & \quad - \int_{(\Lambda^*)} dp_n \| A_{p_n} \Phi_- \|^2 , \end{aligned} \quad (\text{A.37})$$

where

$$A_{p_n} := \int_{(\Lambda^*)^{n-1}} d\mathbf{p}_{n-1} H(\mathbf{p}_n) \delta\left(\sum_{j=1}^n p_j\right) a(\mathbf{p}_{n-1}). \quad (\text{A.38})$$

Using Cauchy-Schwarz first on the inner product on \mathcal{F} , and then w.r.t. $d\mathbf{p}_n d\mathbf{q}_n$, we find the upper bound

$$\begin{aligned} & \left\| \int_{(\Lambda^*)^n} d\mathbf{p}_n H(\mathbf{p}_n) \delta\left(\sum_{j=1}^n p_j\right) a(\mathbf{p}_n) \Phi_- \right\|^2 \\ & \leq \int_{(\Lambda^*)^{2n}} d\mathbf{p}_n d\mathbf{q}_n |H(\mathbf{p}_n)| |H(\mathbf{q}_n)| \delta\left(\sum_{j=1}^n p_j\right) \delta\left(\sum_{j=1}^n q_j\right) \\ & \quad \|a_{q_n}^+ a(\mathbf{p}_{n-1}) \Phi_- \| \|a_{p_n}^+ a(\mathbf{q}_{n-1}) \Phi_- \| \\ & \leq \left(\int_{(\Lambda^*)^{2n}} d\mathbf{p}_n d\mathbf{q}_n \delta\left(\sum_{j=1}^n p_j\right) \delta\left(\sum_{j=1}^n q_j\right) |H(\mathbf{q}_n)|^2 \|a_{q_n}^+ a(\mathbf{p}_{n-1}) \Phi_- \|^2 \right)^{\frac{1}{2}} \\ & \quad \left(\int_{(\Lambda^*)^{2n}} d\mathbf{p}_n d\mathbf{q}_n \delta\left(\sum_{j=1}^n p_j\right) \delta\left(\sum_{j=1}^n q_j\right) |H(\mathbf{p}_n)|^2 \|a_{p_n}^+ a(\mathbf{q}_{n-1}) \Phi_- \|^2 \right)^{\frac{1}{2}} \\ & = \int_{(\Lambda^*)^{2n-1}} d\mathbf{p}_{n-1} d\mathbf{q}_n \delta\left(\sum_{j=1}^n q_j\right) |H(\mathbf{q}_n)|^2 \|a_{q_n} a(\mathbf{p}_{n-1}) \Phi_- \|^2 \\ & \quad + |\Lambda| \int_{(\Lambda^*)^n} d\mathbf{q}_n \delta\left(\sum_{j=1}^n q_j\right) |H(\mathbf{q}_n)|^2 \|(\mathcal{N}_b)_{n-1}^{\frac{1}{2}} \Phi_- \|^2, \end{aligned} \quad (\text{A.39})$$

where we used that $[a_{q_n}, a_{q_n}^+] = |\Lambda|$ together with (A.30). Using (A.30) again, we conclude

$$\begin{aligned} & \left\| \int_{(\Lambda^*)^n} d\mathbf{p}_n H(\mathbf{p}_n) \delta\left(\sum_{j=1}^n p_j\right) a(\mathbf{p}_n) \Phi_- \right\|^2 \\ & \leq \|H\|_{L_{\mathbf{p}_{n-2}}^2 L_{\mathbf{p}_{n-1}, p_n}^\infty}^2 \|(\mathcal{N}_b)_{n-1}^{\frac{1}{2}} \Phi_- \|^2 + |\Lambda| \left\| \delta\left(\sum_{j=1}^n p_j\right)^{\frac{1}{2}} H \right\|_{L_{\mathbf{p}_n}^2}^2 \|(\mathcal{N}_b)_{n-1}^{\frac{1}{2}} \Phi_- \|^2 \\ & \leq \left(\|H\|_{L_{\mathbf{p}_{n-2}}^2 L_{\mathbf{p}_{n-1}, p_n}^\infty}^2 (M-n+1) + |\Lambda| \left\| \delta\left(\sum_{j=1}^n p_j\right)^{\frac{1}{2}} H \right\|_{L_{\mathbf{p}_n}^2}^2 \right) (M)_{n-1}. \end{aligned} \quad (\text{A.40})$$

Finally, we have

$$\begin{aligned} \|a_0 P_M\|^2 &= \sup_{\|\Phi\|=1} \langle P_M \Phi, a_0^+ a_0 P_M \Phi \rangle \\ &= \langle P_M \Phi, a_0 a_0^+ P_M \Phi \rangle - |\Lambda| \|P_M\| \\ &= \|a_0^+ P_M\|^2 - |\Lambda|, \end{aligned} \quad (\text{A.41})$$

where we used that $\|P_M\| = 1$. In addition, we have that

$$\|a_0^+ P_M\| = \|P_{M+1} a_0^+\| = \|a_0 P_{M+1}\|. \quad (\text{A.42})$$

(A.41) and (A.42) imply

$$\|a_0 P_M\|^2 = M|\Lambda|. \quad (\text{A.43})$$

This finishes the proof. \square

APPENDIX B. PROPAGATION OF APPROXIMATE RESTRICTED QUASIFREENESS

Lemma B.1. *Let*

$$\mathcal{U}_N(t) := \mathcal{V}_{HFB}^*(t) \mathcal{W}^*[\sqrt{N|\Lambda|}\phi_0] e^{-it\mathcal{H}_N} \mathcal{W}[\sqrt{N|\Lambda|}\phi_0], \quad (\text{B.1})$$

where for the definition of \mathcal{V}_{HFB} , we refer to (3.7). Then fluctuation dynamics \mathcal{U}_N obeys

$$\begin{cases} i\partial_t \mathcal{U}_N(t) &= (\mathcal{H}_{cub}(t) + \mathcal{H}_{quart}(t)) \mathcal{U}_N(t), \\ \mathcal{U}_N(0) &= \mathbf{1}, \end{cases} \quad (\text{B.2})$$

where $\mathcal{H}_{cub}(t)$ is defined in (2.8), $\mathcal{H}_{quart}(t)$ in (2.9).

Proof. We start by defining the auxiliary dynamics

$$\tilde{\mathcal{U}}_N(t) := \mathcal{W}^*[\sqrt{N|\Lambda|}\phi_0] e^{-i\mathcal{H}_N t} \mathcal{W}[\sqrt{N|\Lambda|}\phi_0]. \quad (\text{B.3})$$

We have that

$$i\partial_t \tilde{\mathcal{U}}_N(t) = \mathcal{W}^*[\sqrt{N|\Lambda|}\phi_0] \mathcal{H}_N \mathcal{W}[\sqrt{N|\Lambda|}\phi_0] \tilde{\mathcal{U}}_N(t). \quad (\text{B.4})$$

Using $\phi_0 = |\Lambda|^{-1/2}$, the explicit expressions for the terms on the right hand side are given by

$$\begin{aligned} & \mathcal{W}^*[\sqrt{N|\Lambda|}\phi_0] \mathcal{H}_N \mathcal{W}[\sqrt{N|\Lambda|}\phi_0] \\ &= \frac{N|\Lambda|\lambda}{2} \int_{\Lambda} dx v(x) + \frac{1}{2} \int_{\Lambda} dx a_x^+ (-\Delta_x) a_x \\ & \quad + \lambda \sqrt{N} \int_{\Lambda^2} dx dy v(x-y) (a_y^+ + a_y) + \lambda \int_{\Lambda^2} dx dy v(x-y) a_y^+ a_y \\ & \quad + \lambda \int_{\Lambda^2} dx dy v(x-y) (a_x^+ a_y + \frac{1}{2} (a_x^+ a_y^+ + a_y a_x)) \\ & \quad + \frac{\lambda}{\sqrt{N}} \int_{\Lambda^2} dx dy v(x-y) a_x^+ (a_y + a_y^+) a_y \\ & \quad + \frac{\lambda}{2N} \int_{\Lambda^2} dx dy v(x-y) a_x^+ a_y^+ a_y a_x. \end{aligned} \quad (\text{B.5})$$

Using the fact that $\hat{v}(0) = 0$, and recalling definitions (1.34)–(1.35), we thus obtain

$$\begin{cases} i\partial_t \tilde{\mathcal{U}}_N(t) &= (\mathcal{H}_{HFB} + \mathcal{H}_{cub} + \mathcal{H}_{quart}) \tilde{\mathcal{U}}_N(t), \\ \tilde{\mathcal{U}}_N(0) &= \mathbf{1}. \end{cases} \quad (\text{B.6})$$

In particular, using

$$\mathcal{U}_N(t) = \mathcal{V}_{HFB}^*(t) \tilde{\mathcal{U}}_N(t), \quad (\text{B.7})$$

a straight-forward calculation yields that \mathcal{U}_N satisfies (B.2). This concludes the proof. \square

Next, we adjust Proposition 3.1 in [74] to our present context.

Lemma B.2. *Assume $|\Lambda| \geq 1$. Let $\tilde{\mathcal{U}}_N$ be defined as in (B.6). Then there are $C_\ell, K_\ell > 0$ such that*

$$\left\| (\mathcal{N}_b + |\Lambda|)^{\frac{\ell}{2}} \tilde{\mathcal{U}}_N(t) (\mathcal{N}_b + |\Lambda|)^{-\frac{\ell}{2}} \left(1 + \frac{\mathcal{N}_b}{N|\Lambda|}\right)^{-\frac{1}{2}} \right\| \leq C_\ell e^{K_\ell \|\hat{v}\|_{w,d} \lambda |\Lambda| t} \quad (\text{B.8})$$

for all $\ell \in \mathbb{N}$.

Proof. We follow the steps of the proof in [74] and point out the differences. We show the statement by induction. Let $\psi \in \mathcal{F}$ be arbitrary.

Step 1: (B.6) implies that

$$\begin{aligned} \partial_t \langle \tilde{\mathcal{U}}_N(t)\psi, (\mathcal{N}_b + |\Lambda|)\tilde{\mathcal{U}}_N(t)\psi \rangle &= -i \langle \tilde{\mathcal{U}}_N(t)\psi, [\mathcal{N}_b, \mathcal{H}_{HFB} + \mathcal{H}_{cub} + \mathcal{H}_{quart}]\tilde{\mathcal{U}}_N(t)\psi \rangle \\ &= -i \langle \tilde{\mathcal{U}}_N(t)\psi, [\mathcal{N}_b, \mathcal{H}_{HFB} + \mathcal{H}_{cub}]\tilde{\mathcal{U}}_N(t)\psi \rangle. \end{aligned} \quad (\text{B.9})$$

Recalling (1.34) and (1.35), we have that

$$[\mathcal{N}_b, \mathcal{H}_{HFB}] = \lambda \int d\mathbf{p}_2 \hat{v}(p_1) \delta(p_1 - p_2) (a_{p_1}^+ a_{-p_2}^+ - a_{p_1} a_{-p_2}), \quad (\text{B.10})$$

$$[\mathcal{N}_b, \mathcal{H}_{cub}] = \frac{\lambda}{\sqrt{N}} \int d\mathbf{p}_3 \hat{v}(p_2) \delta(p_1 + p_2 - p_3) (a_{p_1}^+ a_{p_2}^+ a_{p_3} - a_{p_3}^+ a_{p_2} a_{p_1}) \quad (\text{B.11})$$

Employing Lemma A.4, (B.9) thus yields

$$\begin{aligned} &|\partial_t \langle \tilde{\mathcal{U}}_N(t)\psi, (\mathcal{N}_b + |\Lambda|)\tilde{\mathcal{U}}_N(t)\psi \rangle| \\ &\leq C \|\hat{v}\|_{w,d} \lambda \left(\langle \tilde{\mathcal{U}}_N(t)\psi, (\mathcal{N}_b + |\Lambda|)\tilde{\mathcal{U}}_N(t)\psi \rangle \right. \\ &\quad \left. + \frac{1}{\sqrt{N}} \langle \tilde{\mathcal{U}}_N(t)\psi, \mathcal{N}_b^{\frac{3}{2}} \tilde{\mathcal{U}}_N(t)\psi \rangle \right) \\ &\leq C \|\hat{v}\|_{w,d} \lambda \left(\langle \tilde{\mathcal{U}}_N(t)\psi, (\mathcal{N}_b + |\Lambda|)\tilde{\mathcal{U}}_N(t)\psi \rangle \right. \\ &\quad \left. + \frac{1}{N} \langle \tilde{\mathcal{U}}_N(t)\psi, \mathcal{N}_b^2 \tilde{\mathcal{U}}_N(t)\psi \rangle \right). \end{aligned} \quad (\text{B.12})$$

Using

$$\mathcal{W}[\sqrt{N|\Lambda}|\phi_0] a_p \mathcal{W}^*[\sqrt{N|\Lambda}|\phi_0] = a_p - \sqrt{N}\delta(p), \quad (\text{B.13})$$

we derive that

$$\begin{aligned} [\mathcal{N}_b, \mathcal{W}^*[\sqrt{N|\Lambda}|\phi_0]] &= -\sqrt{N}\mathcal{W}^*[\sqrt{N|\Lambda}|\phi_0] (a_0 + a_0^+) + N|\Lambda|\mathcal{W}^*[\sqrt{N|\Lambda}|\phi_0] \\ &= -(\sqrt{N}(a_0 + a_0^+) + N|\Lambda|)\mathcal{W}^*[\sqrt{N|\Lambda}|\phi_0], \end{aligned} \quad (\text{B.14})$$

$$[\mathcal{N}_b, \mathcal{W}[\sqrt{N|\Lambda}|\phi_0]] = \mathcal{W}[\sqrt{N|\Lambda}|\phi_0] (\sqrt{N}(a_0 + a_0^+) + N|\Lambda|). \quad (\text{B.15})$$

From these identities and using $[\mathcal{N}_b, \mathcal{H}_N] = 0$, we obtain that

$$\begin{aligned} [\mathcal{N}_b, \tilde{\mathcal{U}}_N(t)] &= [\mathcal{N}_b, \mathcal{W}^*[\sqrt{N|\Lambda}|\phi_0]] e^{-it\mathcal{H}_N} \mathcal{W}[\sqrt{N|\Lambda}|\phi_0] \\ &\quad + \mathcal{W}^*[\sqrt{N|\Lambda}|\phi_0] e^{-it\mathcal{H}_N} [\mathcal{N}_b, \mathcal{W}[\sqrt{N|\Lambda}|\phi_0]] \\ &= -\sqrt{N}(a_0 + a_0^+) \tilde{\mathcal{U}}_N(t) + \sqrt{N} \tilde{\mathcal{U}}_N(t) (a_0 + a_0^+). \end{aligned} \quad (\text{B.16})$$

As a consequence, we have that

$$\begin{aligned} &\langle \tilde{\mathcal{U}}_N(t)\psi, \mathcal{N}_b^2 \tilde{\mathcal{U}}_N(t)\psi \rangle \\ &= \langle \mathcal{N}_b \tilde{\mathcal{U}}_N(t)\psi, \tilde{\mathcal{U}}_N(t) \mathcal{N}_b \psi \rangle - \sqrt{N} \langle \mathcal{N}_b \tilde{\mathcal{U}}_N(t)\psi, (a_0 + a_0^+) \tilde{\mathcal{U}}_N(t)\psi \rangle \\ &\quad + \sqrt{N} \langle \mathcal{N}_b \tilde{\mathcal{U}}_N(t)\psi, \tilde{\mathcal{U}}_N(t) (a_0 + a_0^+) \psi \rangle. \end{aligned} \quad (\text{B.17})$$

Using Cauchy-Schwarz, we thus obtain

$$\begin{aligned} &\langle \tilde{\mathcal{U}}_N(t)\psi, \mathcal{N}_b^2 \tilde{\mathcal{U}}_N(t)\psi \rangle \\ &\leq \|\mathcal{N}_b \tilde{\mathcal{U}}_N(t)\psi\| \left(\|\tilde{\mathcal{U}}_N(t) \mathcal{N}_b \psi\| + \sqrt{N} (\|(a_0 + a_0^+) \tilde{\mathcal{U}}_N(t)\psi\| + \|\tilde{\mathcal{U}}_N(t) (a_0 + a_0^+) \psi\|) \right) \end{aligned}$$

$$\begin{aligned} &\leq \|\mathcal{N}_b \tilde{\mathcal{U}}_N(t)\psi\| \left(\|\mathcal{N}_b \psi\| + \sqrt{N} (\|a_0 \tilde{\mathcal{U}}_N(t)\psi\| + \|a_0^+ \tilde{\mathcal{U}}_N(t)\psi\| \right. \\ &\quad \left. + \|a_0 \psi\| + \|a_0^+ \psi\| \right), \end{aligned} \quad (\text{B.18})$$

where we also applied that $\tilde{\mathcal{U}}_N(t) : \mathcal{F} \rightarrow \mathcal{F}$ is a unitary transformation. Lemma A.4 implies

$$\begin{aligned} \|a_0 \psi\|^2 &= \sum_{M=0} \|P_M a_0 \psi\|^2 \\ &= \sum_{M=0} \|a_0 P_{M-1} \psi\|^2 \\ &\leq \sum_{M=0} (M-1) |\Lambda| \|P_{M-1} \psi\|^2 \\ &= |\Lambda| \|\sqrt{\mathcal{N}_b} \psi\|^2. \end{aligned} \quad (\text{B.19})$$

Similarly, we have that

$$\|a_0^+ \psi\| \leq \sqrt{|\Lambda|} \|\sqrt{\mathcal{N}_b + 1} \psi\| \quad (\text{B.20})$$

Employing (B.19) and (B.20), (B.18) implies

$$\begin{aligned} &\langle \tilde{\mathcal{U}}_N(t)\psi, \mathcal{N}_b^2 \tilde{\mathcal{U}}_N(t)\psi \rangle \\ &\leq \|\mathcal{N}_b \tilde{\mathcal{U}}_N(t)\psi\| (\|\mathcal{N}_b \psi\| + 2\sqrt{N|\Lambda|} (\|\sqrt{\mathcal{N}_b + 1} \tilde{\mathcal{U}}_N(t)\psi\| + \|\sqrt{\mathcal{N}_b + 1} \psi\|)) \end{aligned} \quad (\text{B.21})$$

Using Young's inequality implies

$$\begin{aligned} &\langle \tilde{\mathcal{U}}_N(t)\psi, \mathcal{N}_b^2 \tilde{\mathcal{U}}_N(t)\psi \rangle \\ &\leq \frac{1}{2} \langle \tilde{\mathcal{U}}_N(t)\psi, \mathcal{N}_b^2 \tilde{\mathcal{U}}_N(t)\psi \rangle + C \left(\langle \psi, (\mathcal{N}_b^2 + N|\Lambda|(\mathcal{N}_b + 1))\psi \rangle \right. \\ &\quad \left. N|\Lambda| \langle \tilde{\mathcal{U}}_N(t)\psi, (\mathcal{N}_b + 1) \tilde{\mathcal{U}}_N(t)\psi \rangle \right) \end{aligned} \quad (\text{B.22})$$

As a consequence, we find that

$$\begin{aligned} &\frac{1}{N} \langle \tilde{\mathcal{U}}_N(t)\psi, \mathcal{N}_b^2 \tilde{\mathcal{U}}_N(t)\psi \rangle \\ &\leq C|\Lambda| \left(\langle \tilde{\mathcal{U}}_N(t)\psi, (\mathcal{N}_b + 1) \tilde{\mathcal{U}}_N(t)\psi \rangle + \langle \psi, (\mathcal{N}_b + 1 + \frac{\mathcal{N}_b^2}{N|\Lambda|})\psi \rangle \right). \end{aligned} \quad (\text{B.23})$$

Plugging this into (B.12) and using $|\Lambda| \geq 1$, we obtain that

$$\begin{aligned} &|\partial_t \langle \tilde{\mathcal{U}}_N(t)\psi, (\mathcal{N}_b + |\Lambda|) \tilde{\mathcal{U}}_N(t)\psi \rangle| \\ &\leq C \|\hat{v}\|_{w,d} \lambda |\Lambda| \left(\langle \tilde{\mathcal{U}}_N(t)\psi, (\mathcal{N}_b + |\Lambda|) \tilde{\mathcal{U}}_N(t)\psi \rangle + \langle \psi, (\mathcal{N}_b + 1 + \frac{\mathcal{N}_b^2}{N|\Lambda|})\psi \rangle \right) \end{aligned} \quad (\text{B.24})$$

Gronwall's Lemma then implies

$$\langle \tilde{\mathcal{U}}_N(t)\psi, (\mathcal{N}_b + |\Lambda|) \tilde{\mathcal{U}}_N(t)\psi \rangle \leq e^{C \|\hat{v}\|_{w,d} \lambda |\Lambda| t} \langle \psi, (\mathcal{N}_b + |\Lambda| + \frac{\mathcal{N}_b^2}{N|\Lambda|})\psi \rangle. \quad (\text{B.25})$$

Step 2: Assume that

$$\begin{aligned} & \langle \tilde{\mathcal{U}}_N(t)\psi, (\mathcal{N}_b + |\Lambda|)^j \tilde{\mathcal{U}}_N(t)\psi \rangle \\ & \leq C_j e^{K_j \|\hat{v}\|_{w,d\lambda} |\Lambda|^t} \langle \psi, (\mathcal{N}_b + |\Lambda|)^j \left(1 + \frac{\mathcal{N}_b}{N|\Lambda|}\right) \psi \rangle \end{aligned} \quad (\text{B.26})$$

for all $1 \leq j \leq \ell$ and some constants C_j, K_j , and any $\psi \in \mathcal{F}$. We compute

$$\begin{aligned} & i\partial_t \langle \tilde{\mathcal{U}}_N(t)\psi, (\mathcal{N}_b + |\Lambda|)^{\ell+1} \tilde{\mathcal{U}}_N(t)\psi \rangle \\ & = \langle \tilde{\mathcal{U}}_N(t)\psi, [(\mathcal{N}_b + |\Lambda|)^{\ell+1}, \mathcal{H}_{cor}^{\phi_0} + \mathcal{H}_{cub}^{\phi_0}] \tilde{\mathcal{U}}_N(t)\psi \rangle \\ & = \sum_{j=1}^{\ell+1} \langle \tilde{\mathcal{U}}_N(t)\psi, (\mathcal{N}_b + |\Lambda|)^{j-1} [\mathcal{N}_b, \mathcal{H}_{cor}^{\phi_0} + \mathcal{H}_{cub}^{\phi_0}] (\mathcal{N}_b + |\Lambda|)^{\ell+1-j} \tilde{\mathcal{U}}_N(t)\psi \rangle \end{aligned} \quad (\text{B.27})$$

Let

$$A_{cub}[\hat{v}] := \int d\mathbf{p}_3 \hat{v}(p_2) \delta(p_1 + p_2 - p_3) a_{p_1}^+ a_{p_2}^+ a_{p_3}. \quad (\text{B.28})$$

Applying (B.10) and (B.11), (B.27) yields

$$\begin{aligned} & i\partial_t \langle \tilde{\mathcal{U}}_N(t)\psi, (\mathcal{N}_b + |\Lambda|)^{\ell+1} \tilde{\mathcal{U}}_N(t)\psi \rangle \\ & = 2 \sum_{j=1}^{\ell+1} \text{Im} \langle \tilde{\mathcal{U}}_N(t)\psi, (\mathcal{N}_b + |\Lambda|)^{j-1} (g[\hat{v}] - A_{cub}[\hat{v}]) (\mathcal{N}_b + |\Lambda|)^{\ell+1-j} \tilde{\mathcal{U}}_N(t)\psi \rangle \\ & = 2 \sum_{j=1}^{\ell+1} \text{Im} \sum_{m,n=0}^{\infty} (m + |\Lambda|)^{j-1} (n + |\Lambda|)^{\ell+1-j} \\ & \quad \langle \tilde{\mathcal{U}}_N(t)\psi, P_m (\lambda g[\hat{v}] - \frac{\lambda}{\sqrt{N}} A_{cub}[\hat{v}]) P_n \tilde{\mathcal{U}}_N(t)\psi \rangle. \end{aligned} \quad (\text{B.29})$$

Observe that we have

$$P_m g[\hat{v}] P_n = P_m g[\hat{v}] P_{m+2} \delta_{n,m+2}, \quad (\text{B.30})$$

$$P_m A_{cub}[\hat{v}] P_n = P_m A_{cub}[\hat{v}] P_{m-1} \delta_{n,m-1}. \quad (\text{B.31})$$

Lemma A.4 and (B.29) then imply

$$\begin{aligned} & |\partial_t \langle \tilde{\mathcal{U}}_N(t)\psi, (\mathcal{N}_b + |\Lambda|)^{\ell+1} \tilde{\mathcal{U}}_N(t)\psi \rangle| \\ & \leq C\lambda \sum_{j=1}^{\ell+1} \sum_{m,n=0}^{\infty} (m + |\Lambda|)^{j-1} (n + |\Lambda|)^{\ell+1-j} \|P_m \tilde{\mathcal{U}}_N(t)\psi\| \|P_n \tilde{\mathcal{U}}_N(t)\psi\| \\ & \quad \left(\|g[\hat{v}] P_{m+2}\| \delta_{n,m+2} + \frac{\|A_{cub}[\hat{v}] P_{m-1}\|}{\sqrt{N}} \delta_{n,m-1} \right) \\ & \leq C \|\hat{v}\|_{w,d\lambda} \sum_{j=1}^{\ell+1} \sum_{m,n=0}^{\infty} (m + |\Lambda|)^{j-1} (n + |\Lambda|)^{\ell+1-j} \\ & \quad \left(\|P_m \tilde{\mathcal{U}}_N(t)\psi\|^2 + \|P_n \tilde{\mathcal{U}}_N(t)\psi\|^2 \right) \left((m+2 + |\Lambda|) \delta_{n,m+2} + \frac{m^{\frac{3}{2}}}{\sqrt{N}} \delta_{n,m-1} \right) \\ & \leq C \|\hat{v}\|_{w,d\lambda} \sum_{j=1}^{\ell+1} \sum_{m=0}^{\infty} (m + |\Lambda|)^{j-1} (m+2 + |\Lambda|)^{\ell+1-j} \|P_m \tilde{\mathcal{U}}_N(t)\psi\|^2 \end{aligned}$$

$$(m + 2 + |\Lambda| + \frac{(m + 1)^{\frac{3}{2}}}{\sqrt{N}}). \quad (\text{B.32})$$

We can further estimate this by

$$\begin{aligned} & |\partial_t \langle \tilde{\mathcal{U}}_N(t) \psi, (\mathcal{N}_b + |\Lambda|)^{\ell+1} \tilde{\mathcal{U}}_N(t) \psi \rangle| \\ & \leq C(\ell + 1) \|\hat{v}\|_{w,d} \lambda \langle \tilde{\mathcal{U}}_N(t) \psi, (\mathcal{N}_b + |\Lambda| + 2)^{\ell+1} \tilde{\mathcal{U}}_N(t) \psi \rangle \\ & \quad + \langle \tilde{\mathcal{U}}_N(t) \psi, \frac{(\mathcal{N}_b + |\Lambda| + 2)^{\ell+\frac{3}{2}}}{\sqrt{N}} \tilde{\mathcal{U}}_N(t) \psi \rangle \\ & \leq C_\ell \|\hat{v}\|_{w,d} \lambda \langle \tilde{\mathcal{U}}_N(t) \psi, (\mathcal{N}_b + |\Lambda|)^{\ell+1} \tilde{\mathcal{U}}_N(t) \psi \rangle \\ & \quad + \frac{1}{N} \langle \tilde{\mathcal{U}}_N(t) \psi, (\mathcal{N}_b + |\Lambda|)^{\ell+2} \tilde{\mathcal{U}}_N(t) \psi \rangle \end{aligned} \quad (\text{B.33})$$

We claim that

$$\begin{aligned} & \frac{1}{N|\Lambda|} \langle \tilde{\mathcal{U}}_N(t) \psi, (\mathcal{N}_b + |\Lambda|)^{j+1} \tilde{\mathcal{U}}_N(t) \psi \rangle \\ & \leq C_j \left(e^{K_j \|\hat{v}\|_{w,d} \lambda |\Lambda| t} \langle \psi, (\mathcal{N}_b + |\Lambda|)^j \left(1 + \frac{\mathcal{N}_b}{N|\Lambda|} \right) \psi \rangle \right. \\ & \quad \left. + \langle \tilde{\mathcal{U}}_N(t) \psi, (\mathcal{N}_b + |\Lambda|)^j \tilde{\mathcal{U}}_N(t) \psi \rangle \right) \end{aligned} \quad (\text{B.34})$$

for all $1 \leq j \leq \ell + 1$ and all $\psi \in \mathcal{F}$. (B.34) for $j = \ell + 1$ together with (B.33) implies

$$\begin{aligned} & |\partial_t \langle \tilde{\mathcal{U}}_N(t) \psi, (\mathcal{N}_b + |\Lambda|)^{\ell+1} \tilde{\mathcal{U}}_N(t) \psi \rangle| \\ & \leq C_\ell \|\hat{v}\|_{w,d} \lambda |\Lambda| \left(\langle \tilde{\mathcal{U}}_N(t) \psi, (\mathcal{N}_b + |\Lambda|)^{\ell+1} \tilde{\mathcal{U}}_N(t) \psi \rangle \right. \\ & \quad \left. + e^{K_\ell \|\hat{v}\|_{w,d} \lambda |\Lambda| t} \langle \psi, (\mathcal{N}_b + |\Lambda|)^{\ell+1} \left(1 + \frac{\mathcal{N}_b}{N|\Lambda|} \right) \psi \rangle \right). \end{aligned} \quad (\text{B.35})$$

Gronwall's Lemma then implies

$$\begin{aligned} & |\langle \tilde{\mathcal{U}}_N(t) \psi, (\mathcal{N}_b + |\Lambda|)^{\ell+1} \tilde{\mathcal{U}}_N(t) \psi \rangle| \\ & \leq e^{K_\ell \|\hat{v}\|_{w,d} \lambda |\Lambda| t} \langle \psi, (\mathcal{N}_b + |\Lambda|)^{\ell+1} \psi \rangle \\ & \quad + C_\ell \|\hat{v}\|_{w,d} \lambda |\Lambda| t e^{K_\ell \|\hat{v}\|_{w,d} \lambda |\Lambda| t} \langle \psi, (\mathcal{N}_b + |\Lambda|)^{\ell+1} \left(1 + \frac{\mathcal{N}_b}{N|\Lambda|} \right) \psi \rangle \\ & \leq C_\ell e^{K_\ell \|\hat{v}\|_{w,d} \lambda |\Lambda| t} \langle \psi, (\mathcal{N}_b + |\Lambda|)^{\ell+1} \left(1 + \frac{\mathcal{N}_b}{N|\Lambda|} \right) \psi \rangle. \end{aligned} \quad (\text{B.36})$$

Thus, proving (B.34) for $j = \ell + 1$ concludes the proof. We have proved (B.34) for $j = 1$ in Step 1, (B.23). We have that (B.34) also holds for $j = 0$, observing that

$$\begin{aligned} & \mathcal{W}[\sqrt{N|\Lambda|}\phi_0] \mathcal{N}_b \mathcal{W}^*[\sqrt{N|\Lambda|}\phi_0] \\ & = \mathcal{N}_b - \sqrt{N}(a_0 + a_0^\dagger) + N|\Lambda| \\ & \leq 2(\mathcal{N}_b + 1 + N|\Lambda|), \end{aligned} \quad (\text{B.37})$$

which then commutes with $e^{-it\mathcal{H}_N}$.

Suppose (B.34) holds up to some $1 \leq j \leq \ell - 1$. Applying (B.16), we have that

$$\begin{aligned} & \frac{1}{N|\Lambda|} \langle \tilde{\mathcal{U}}_N(t) \psi, (\mathcal{N}_b + |\Lambda|)^{j+2} \tilde{\mathcal{U}}_N(t) \psi \rangle \\ & = \frac{1}{N|\Lambda|} \langle (\mathcal{N}_b + |\Lambda|)^{j+1} \tilde{\mathcal{U}}_N(t) \psi, \tilde{\mathcal{U}}_N(t) (\mathcal{N}_b + |\Lambda|) \psi \rangle \end{aligned}$$

$$\begin{aligned}
& -\frac{1}{\sqrt{N}|\Lambda|} \langle (\mathcal{N}_b + |\Lambda|)^{j+1} \tilde{\mathcal{U}}_N(t)\psi, (a_0 + a_0^+) \tilde{\mathcal{U}}_N(t)\psi \rangle \\
& + \frac{1}{\sqrt{N}|\Lambda|} \langle (\mathcal{N}_b + |\Lambda|)^{j+1} \tilde{\mathcal{U}}_N(t)\psi, \tilde{\mathcal{U}}_N(t)(a_0 + a_0^+)\psi \rangle. \tag{B.38}
\end{aligned}$$

We can bound the second term by

$$\begin{aligned}
& \frac{1}{\sqrt{N}|\Lambda|} |\langle (\mathcal{N}_b + |\Lambda|)^{j+1} \tilde{\mathcal{U}}_N(t)\psi, (a_0 + a_0^+) \tilde{\mathcal{U}}_N(t)\psi \rangle| \\
& \leq \alpha \langle \tilde{\mathcal{U}}_N(t)\psi, (\mathcal{N}_b + |\Lambda|)^{j+1} \tilde{\mathcal{U}}_N(t)\psi \rangle \\
& + \frac{1}{\alpha N |\Lambda|^2} \langle \tilde{\mathcal{U}}_N(t)\psi, (a_0 + a_0^+) (\mathcal{N}_b + |\Lambda|)^{j+1} (a_0 + a_0^+) \tilde{\mathcal{U}}_N(t)\psi \rangle. \tag{B.39}
\end{aligned}$$

Employing (B.19) and (B.20), we find that, for any $\tilde{\psi} \in \mathcal{F}$,

$$\begin{aligned}
& \langle \tilde{\psi}, (a_0 + a_0^+) (\mathcal{N}_b + |\Lambda|)^k (a_0 + a_0^+) \tilde{\psi} \rangle \\
& \leq (\|(\mathcal{N}_b + |\Lambda|)^{\frac{k}{2}} a_0 \tilde{\psi}\| + \|(\mathcal{N}_b + |\Lambda|)^{\frac{k}{2}} a_0^+ \tilde{\psi}\|)^2 \\
& = (\|a_0 (\mathcal{N}_b + |\Lambda| - 1)^{\frac{k}{2}} \tilde{\psi}\| + \|a_0^+ (\mathcal{N}_b + |\Lambda| + 1)^{\frac{k}{2}} \tilde{\psi}\|)^2 \\
& \leq C_k |\Lambda| \langle \tilde{\psi}, (\mathcal{N}_b + |\Lambda|)^{k+1} \tilde{\psi} \rangle, \tag{B.40}
\end{aligned}$$

i.e.,

$$(a_0 + a_0^+) (\mathcal{N}_b + |\Lambda|)^k (a_0 + a_0^+) \leq C_k |\Lambda| (\mathcal{N}_b + |\Lambda|)^{k+1}. \tag{B.41}$$

Employing (B.41) and choosing $\alpha > 0$ sufficiently large, (B.39) implies

$$\begin{aligned}
& \frac{1}{\sqrt{N}|\Lambda|} |\langle (\mathcal{N}_b + |\Lambda|)^{j+1} \tilde{\mathcal{U}}_N(t)\psi, (a_0 + a_0^+) \tilde{\mathcal{U}}_N(t)\psi \rangle| \\
& \leq C_j \langle \tilde{\mathcal{U}}_N(t)\psi, (\mathcal{N}_b + |\Lambda|)^{j+1} \tilde{\mathcal{U}}_N(t)\psi \rangle \\
& + \frac{1}{4N|\Lambda|} \langle \tilde{\mathcal{U}}_N(t)\psi, (\mathcal{N}_b + |\Lambda|)^{j+2} \tilde{\mathcal{U}}_N(t)\psi \rangle. \tag{B.42}
\end{aligned}$$

We bound the third term in (B.38) by

$$\begin{aligned}
& \frac{1}{\sqrt{N}|\Lambda|} |\langle (\mathcal{N}_b + |\Lambda|)^{j+1} \tilde{\mathcal{U}}_N(t)\psi, \tilde{\mathcal{U}}_N(t)(a_0 + a_0^+)\psi \rangle| \\
& \leq \frac{4}{|\Lambda|} \langle \tilde{\mathcal{U}}_N(t)(a_0 + a_0^+)\psi, (\mathcal{N}_b + |\Lambda|)^j \tilde{\mathcal{U}}_N(t)(a_0 + a_0^+)\psi \rangle \\
& + \frac{1}{4N|\Lambda|} \langle \tilde{\mathcal{U}}_N(t)\psi, (\mathcal{N}_b + |\Lambda|)^{j+2} \tilde{\mathcal{U}}_N(t)\psi \rangle. \tag{B.43}
\end{aligned}$$

The induction hypothesis (B.26) and (B.41) hence imply

$$\begin{aligned}
& \frac{1}{\sqrt{N}|\Lambda|} |\langle (\mathcal{N}_b + |\Lambda|)^{j+1} \tilde{\mathcal{U}}_N(t)\psi, \tilde{\mathcal{U}}_N(t)(a_0 + a_0^+)\psi \rangle| \\
& \leq \frac{C_j}{|\Lambda|} e^{K_j \|\hat{v}\|_{w, a\lambda|\Lambda|t}} \langle (a_0 + a_0^+)\psi, (\mathcal{N}_b + |\Lambda|)^j (1 + \frac{\mathcal{N}_b}{N|\Lambda|}) (a_0 + a_0^+)\psi \rangle \\
& + \frac{1}{4N|\Lambda|} \langle \tilde{\mathcal{U}}_N(t)\psi, (\mathcal{N}_b + |\Lambda|)^{j+2} \tilde{\mathcal{U}}_N(t)\psi \rangle \\
& \leq C_j e^{K_j \|\hat{v}\|_{w, a\lambda|\Lambda|t}} \langle \psi, (\mathcal{N}_b + |\Lambda|)^{j+1} (1 + \frac{\mathcal{N}_b}{N|\Lambda|}) \psi \rangle
\end{aligned}$$

$$+ \frac{1}{4N|\Lambda|} \langle \tilde{\mathcal{U}}_N(t)\psi, (\mathcal{N}_b + |\Lambda|)^{j+2} \tilde{\mathcal{U}}_N(t)\psi \rangle \quad (\text{B.44})$$

For the first term in (B.38), we apply (B.16) to the left and obtain

$$\begin{aligned} & \frac{1}{N|\Lambda|} \langle (\mathcal{N}_b + |\Lambda|)^{j+1} \tilde{\mathcal{U}}_N(t)\psi, \tilde{\mathcal{U}}_N(t)(\mathcal{N}_b + |\Lambda|)\psi \rangle \\ &= \frac{1}{N|\Lambda|} \langle \tilde{\mathcal{U}}_N(t)(\mathcal{N}_b + |\Lambda|)\psi, (\mathcal{N}_b + |\Lambda|)^j \tilde{\mathcal{U}}_N(t)(\mathcal{N}_b + |\Lambda|)\psi \rangle \\ & \quad - \frac{1}{\sqrt{N}|\Lambda|} \langle (\mathcal{N}_b + |\Lambda|)^j (a_0 + a_0^+) \tilde{\mathcal{U}}_N(t)\psi, \tilde{\mathcal{U}}_N(t)(\mathcal{N}_b + |\Lambda|)\psi \rangle \\ & \quad + \frac{1}{\sqrt{N}|\Lambda|} \langle (\mathcal{N}_b + |\Lambda|)^j \tilde{\mathcal{U}}_N(t)(a_0 + a_0^+)\psi, \tilde{\mathcal{U}}_N(t)(\mathcal{N}_b + |\Lambda|)\psi \rangle. \end{aligned} \quad (\text{B.45})$$

For the first term in (B.45), we use the induction hypothesis (B.34), and obtain

$$\begin{aligned} & \frac{1}{N|\Lambda|} \langle \tilde{\mathcal{U}}_N(t)(\mathcal{N}_b + |\Lambda|)\psi, (\mathcal{N}_b + |\Lambda|)^j \tilde{\mathcal{U}}_N(t)(\mathcal{N}_b + |\Lambda|)\psi \rangle \\ & \leq C_j \left(e^{K_j \|\hat{v}\|_{w, a\lambda|\Lambda|t}} \langle \psi, (\mathcal{N}_b + |\Lambda|)^{j+1} \left(1 + \frac{\mathcal{N}_b}{N|\Lambda|}\right) \psi \rangle \right. \\ & \quad \left. + \langle \tilde{\mathcal{U}}_N(t)(\mathcal{N}_b + |\Lambda|)\psi, (\mathcal{N}_b + |\Lambda|)^{j-1} \tilde{\mathcal{U}}_N(t)(\mathcal{N}_b + |\Lambda|)\psi \rangle \right) \\ & \leq C_j e^{K_j \|\hat{v}\|_{w, a\lambda|\Lambda|t}} \langle \psi, (\mathcal{N}_b + |\Lambda|)^{j+1} \left(1 + \frac{\mathcal{N}_b}{N|\Lambda|}\right) \psi \rangle, \end{aligned} \quad (\text{B.46})$$

where in the last step we employed (B.26). Using Cauchy-Schwarz, followed by Young's inequality, the second term in (B.45) can be bounded by

$$\begin{aligned} & \frac{1}{\sqrt{N}|\Lambda|} \left| \langle (\mathcal{N}_b + |\Lambda|)^j (a_0 + a_0^+) \tilde{\mathcal{U}}_N(t)\psi, \tilde{\mathcal{U}}_N(t)(\mathcal{N}_b + |\Lambda|)\psi \rangle \right| \\ & \leq \frac{1}{\sqrt{N}|\Lambda|} \| (\mathcal{N}_b + |\Lambda|)^{\frac{j}{2}} (a_0 + a_0^+) \tilde{\mathcal{U}}_N(t)\psi \| \| (\mathcal{N}_b + |\Lambda|)^{\frac{j}{2}} \tilde{\mathcal{U}}_N(t)(\mathcal{N}_b + |\Lambda|)\psi \| \\ & \leq \frac{1}{|\Lambda|} \left(\langle \tilde{\mathcal{U}}_N(t)\psi, (a_0 + a_0^+) (\mathcal{N}_b + |\Lambda|)^j (a_0 + a_0^+) \tilde{\mathcal{U}}_N(t)\psi \rangle \right. \\ & \quad \left. + \frac{1}{N} \langle \tilde{\mathcal{U}}_N(t)(\mathcal{N}_b + |\Lambda|)\psi, (\mathcal{N}_b + |\Lambda|)^j \tilde{\mathcal{U}}_N(t)(\mathcal{N}_b + |\Lambda|)\psi \rangle \right). \end{aligned} \quad (\text{B.47})$$

Employing (B.41) and then (B.46), we thus obtain

$$\begin{aligned} & \frac{1}{\sqrt{N}|\Lambda|} \left| \langle (\mathcal{N}_b + |\Lambda|)^j (a_0 + a_0^+) \tilde{\mathcal{U}}_N(t)\psi, \tilde{\mathcal{U}}_N(t)(\mathcal{N}_b + |\Lambda|)\psi \rangle \right| \\ & \leq C_j \langle \tilde{\mathcal{U}}_N(t)\psi, (\mathcal{N}_b + |\Lambda|)^{j+1} \tilde{\mathcal{U}}_N(t)\psi \rangle \\ & \quad + \frac{1}{N|\Lambda|} \langle \tilde{\mathcal{U}}_N(t)(\mathcal{N}_b + |\Lambda|)\psi, (\mathcal{N}_b + |\Lambda|)^j \tilde{\mathcal{U}}_N(t)(\mathcal{N}_b + |\Lambda|)\psi \rangle \\ & \leq C_j \langle \tilde{\mathcal{U}}_N(t)\psi, (\mathcal{N}_b + |\Lambda|)^{j+1} \tilde{\mathcal{U}}_N(t)\psi \rangle \\ & \quad + C_j e^{K_j \|\hat{v}\|_{w, a\lambda|\Lambda|t}} \langle \psi, (\mathcal{N}_b + |\Lambda|)^{j+1} \left(1 + \frac{\mathcal{N}_b}{N|\Lambda|}\right) \psi \rangle. \end{aligned} \quad (\text{B.48})$$

Similarly, the third term in (B.45) can be estimated using

$$\frac{1}{\sqrt{N}|\Lambda|} \left| \langle (\mathcal{N}_b + |\Lambda|)^j \tilde{\mathcal{U}}_N(t)(a_0 + a_0^+)\psi, \tilde{\mathcal{U}}_N(t)(\mathcal{N}_b + |\Lambda|)\psi \rangle \right|$$

$$\begin{aligned}
&\leq \frac{1}{|\Lambda|} \langle \tilde{\mathcal{U}}_N(t)(a_0 + a_0^+) \psi, (\mathcal{N}_b + |\Lambda|)^j \tilde{\mathcal{U}}_N(t)(a_0 + a_0^+) \psi \rangle \\
&\quad + \frac{1}{N|\Lambda|} \langle \tilde{\mathcal{U}}_N(t)(\mathcal{N}_b + |\Lambda|) \psi, (\mathcal{N}_b + |\Lambda|)^j \tilde{\mathcal{U}}_N(t)(\mathcal{N}_b + |\Lambda|) \psi \rangle \\
&\leq \frac{C_j}{|\Lambda|} e^{K_j \|\hat{v}\|_w, d\lambda|\Lambda|t} \langle \psi, (a_0 + a_0^+) (\mathcal{N}_b + |\Lambda|)^j (1 + \frac{\mathcal{N}_b}{N|\Lambda|}) (a_0 + a_0^+) \psi \rangle \\
&\quad + C_j e^{K_j \|\hat{v}\|_w, d\lambda|\Lambda|t} \langle \psi, (\mathcal{N}_b + |\Lambda|)^{j+1} (1 + \frac{\mathcal{N}_b}{N|\Lambda|}) \psi \rangle \\
&\leq C_j e^{K_j \|\hat{v}\|_w, d\lambda|\Lambda|t} \langle \psi, (\mathcal{N}_b + |\Lambda|)^{j+1} (1 + \frac{\mathcal{N}_b}{N|\Lambda|}) \psi \rangle, \tag{B.49}
\end{aligned}$$

where we used (B.26) and (B.46), followed by (B.41). Inserting (B.46), (B.48), and (B.49) into (B.45), we obtain that

$$\begin{aligned}
&\frac{1}{N|\Lambda|} \langle (\mathcal{N}_b + |\Lambda|)^{j+1} \tilde{\mathcal{U}}_N(t) \psi, \tilde{\mathcal{U}}_N(t) (\mathcal{N}_b + |\Lambda|) \psi \rangle \\
&\leq C_j e^{K_j \|\hat{v}\|_w, d\lambda|\Lambda|t} \langle \psi, (\mathcal{N}_b + |\Lambda|)^{j+1} (1 + \frac{\mathcal{N}_b}{N|\Lambda|}) \psi \rangle \\
&\quad + C_j \langle \tilde{\mathcal{U}}_N(t) \psi, (\mathcal{N}_b + |\Lambda|)^{j+1} \tilde{\mathcal{U}}_N(t) \psi \rangle. \tag{B.50}
\end{aligned}$$

Plugging (B.42), (B.44), and (B.50) into (B.38), we find that

$$\begin{aligned}
&\frac{1}{N|\Lambda|} \langle \tilde{\mathcal{U}}_N(t) \psi, (\mathcal{N}_b + |\Lambda|)^{j+2} \tilde{\mathcal{U}}_N(t) \psi \rangle \\
&\leq C_j e^{K_j \|\hat{v}\|_w, d\lambda|\Lambda|t} \langle \psi, (\mathcal{N}_b + |\Lambda|)^{j+1} (1 + \frac{\mathcal{N}_b}{N|\Lambda|}) \psi \rangle \\
&\quad + C_j \langle \tilde{\mathcal{U}}_N(t) \psi, (\mathcal{N}_b + |\Lambda|)^{j+1} \tilde{\mathcal{U}}_N(t) \psi \rangle. \tag{B.51}
\end{aligned}$$

This concludes the proof. \square

APPENDIX C. PROOF OF CONVERGENCE TO MEAN FIELD EQUATIONS

Lemma C.1. *Let $H, F \in C^1(\mathbb{R}^n)$ be such that*

$$\left\| \nabla \cdot \left(\frac{H \nabla F}{|\nabla F|} \right) \right\|_1 < \infty. \tag{C.1}$$

Then the following holds true for all $\omega \in \mathbb{R}$ and $g \in C_0^1(\mathbb{R})$:

- (1) $\int_{F=\omega} \frac{d\mathcal{H}^{n-1}}{|\nabla F|} H = \int_{F<\omega} dp \nabla \cdot \left(\frac{H \nabla F}{|\nabla F|^2} \right),$
- (2) $\int d\omega g(\omega) \partial_\omega \int_{F=\omega} \frac{d\mathcal{H}^{n-1}}{|\nabla F|} H = \int d\omega g(\omega) \int_{F=\omega} \frac{d\mathcal{H}^{n-1}}{|\nabla F|} \nabla \cdot \left(\frac{H \nabla F}{|\nabla F|^2} \right).$

Proof. The first statement is a direct consequence of the Divergence Theorem together with the fact that $\nabla F/|\nabla F|$ is the outer normal for $\{F < \omega\}$. For the second statement, we have that the divergence theorem implies that

$$\int d\omega g(\omega) \partial_\omega \int_{F=\omega} \frac{d\mathcal{H}^{n-1}}{|\nabla F|} H = - \int d\omega g'(\omega) \int_{F=\omega} \frac{d\mathcal{H}^{n-1}}{|\nabla F|} H, \tag{C.2}$$

where we used the assumptions $g \in C_0^1(\mathbb{R})$, (C.1), and the first statement. Employing the Coarea Formula, we have that

$$\int d\omega g(\omega) \partial_\omega \int_{F=\omega} \frac{d\mathcal{H}^{n-1}}{|\nabla F|} H = - \int dp g'(F) H. \tag{C.3}$$

Expanding with the factor $\frac{\nabla F \cdot \nabla F}{|\nabla F|^2}$, the Divergence Theorem, together with $g \in C_0^1(\mathbb{R})$, (C.1), then implies

$$\int d\omega g(\omega) \partial_\omega \int_{F=\omega} \frac{d\mathcal{H}^{n-1}}{|\nabla F|} H = \int dp g(F) \nabla \cdot \left(\frac{H \nabla F}{|\nabla F|^2} \right). \quad (\text{C.4})$$

Finally, the Coarea Formula implies

$$\int d\omega g(\omega) \partial_\omega \int_{F=\omega} \frac{d\mathcal{H}^{n-1}}{|\nabla F|} H = \int d\omega g(\omega) \int_{F=\omega} \frac{d\mathcal{H}^{n-1}}{|\nabla F|} \nabla \cdot \left(\frac{H \nabla F}{|\nabla F|^2} \right). \quad (\text{C.5})$$

This finishes the proof. \square

Lemma C.2. *Let $H, F_1, F_2 \in C^2(\mathbb{R}^6)$ be s.t.*

$$H, \nabla \cdot \frac{\nabla F_2 |H|}{|\nabla F_2|^2} \in L^1(\mathbb{R}^6). \quad (\text{C.6})$$

Then, for all $t \geq 0$, we have that

$$\begin{aligned} & \left| \int d\mathbf{p}_2 \int_{\Delta[t,2]} ds_2 e^{-i(F_1(\mathbf{p}_2)s_1 + F_2(\mathbf{p}_2)s_2)} H(\mathbf{p}_2) \right| \\ & \leq Ct \left(\log(t+1) \left\| \nabla \cdot \frac{\nabla F_2 |H|}{|\nabla F_2|^2} \right\|_1 + \|H\|_1 \right). \end{aligned} \quad (\text{C.7})$$

Proof. Observe that

$$\begin{aligned} \left| \int_{\Delta[t,2]} ds_2 e^{-i(\omega_1 s_1 + \omega_2 s_2)} \right| & \leq \int_0^t ds_1 \left| \int_0^{s_1} ds_2 e^{-i\omega_2 s_2} \right| \\ & \leq \int_0^t ds_1 \frac{2}{|\omega_2| + \frac{1}{s_1}} \\ & \leq \frac{2t}{|\omega_2| + \frac{1}{t}} \end{aligned} \quad (\text{C.8})$$

for all $\omega_1, \omega_2 \in \mathbb{R}$, $t \geq 0$, due to the fact that

$$\left| \int_{x_1}^{x_2} dy e^{-iay} \right| \leq \min\{x_2 - x_1; \frac{2}{|a|}\} \leq \frac{2}{|a| + \frac{1}{x_2 - x_1}} \quad (\text{C.9})$$

for all $x_1, x_2, a \in \mathbb{R}$, $x_2 \geq x_1$.

Then (C.8) followed by the Coarea Formula implies

$$\begin{aligned} & \left| \int d\mathbf{p}_2 \int_{\Delta[t,2]} ds_2 e^{-i(F_1(\mathbf{p}_2)s_1 + F_2(\mathbf{p}_2)s_2)} H(\mathbf{p}_2) \right| \\ & \leq Ct \int d\mathbf{p}_2 \frac{|H(\mathbf{p}_2)|}{|F_2(\mathbf{p}_2)| + \frac{1}{t}} \\ & \leq Ct \int d\omega \frac{1}{|\omega| + \frac{1}{t}} \int_{F_2=\omega} d\mathcal{H}^5(\mathbf{p}_2) \frac{|H(\mathbf{p}_2)|}{|\nabla F_2(\mathbf{p}_2)|}. \end{aligned} \quad (\text{C.10})$$

Lemma C.1 yields

$$\begin{aligned} \left| \int_{F_2=\omega} d\mathcal{H}^5(\mathbf{p}_2) \frac{|H(\mathbf{p}_2)|}{|\nabla F_2(\mathbf{p}_2)|} \right| & \leq \int_{F_2>\omega} d\mathbf{p}_2 \left| \nabla \cdot \frac{\nabla F_2(\mathbf{p}_2) |H(\mathbf{p}_2)|}{|\nabla F_2(\mathbf{p}_2)|^2} \right| \\ & \leq \left\| \nabla \cdot \frac{\nabla F_2 |H|}{|\nabla F_2|^2} \right\|_1. \end{aligned} \quad (\text{C.11})$$

Then, after splitting the domain of integration w.r.t. $d\omega$ into $(-1, 1) \cup (-1, 1)^c$, (C.10) can be bounded by

$$\begin{aligned} & \int_{-1}^1 d\omega \frac{1}{|\omega| + \frac{1}{t}} \sup_{\omega} \left| \int_{F_2=\omega} d\mathcal{H}^5(\mathbf{p}_2) \frac{|H(\mathbf{p}_2)|}{|\nabla F_2(\mathbf{p}_2)|} \right| \\ & \quad + \int d\mathbf{p}_2 \mathbf{1}_{|F_2(\mathbf{p}_2)| \geq 1} |H(\mathbf{p}_2)| \\ & \leq Ct \left(\log(t+1) \left\| \nabla \cdot \frac{\nabla F_2 H}{|\nabla F_2|^2} \right\|_1 + \|H\|_1 \right), \end{aligned} \quad (\text{C.12})$$

where, again, we used the Coarea Formula. \square

Lemma C.3. *Let $h \in C_b^n([0, \|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c}])$, $n \in \mathbb{N}$, $n \leq 3$. Then the following holds for all $\lambda \in (0, 1]$. We have that*

$$\left\| D^n \left[h \left(\frac{\lambda \hat{v}}{E} \right) \right] \right\|_{\infty} \leq C_n \lambda \|h\|_{C^n[0, \|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c} \lambda]} \|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c}^n. \quad (\text{C.13})$$

Moreover, Ω satisfies

$$\nabla \Omega(p) = (1 + \lambda m_{\Omega}(p))p \quad (\text{C.14})$$

$$\|m_{\Omega}\|_{\infty} \leq C \|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c}, \quad (\text{C.15})$$

$$\|D^2 \Omega - I\|_{\infty} \leq C \lambda \|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c}^2,$$

$$\|D^3 \Omega\|_{\infty} \leq C \lambda \|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c}^3. \quad (\text{C.16})$$

Proof. By the Faà di Bruno formula, we have that

$$\left| D^n \left[h \left(\frac{\lambda \hat{v}}{E} \right) \right] \right| \leq C_n \sum_{\mathbf{r}_n \in \hat{R}(n)} \left| h^{(S(\mathbf{r}_n))} \left(\frac{\lambda \hat{v}}{E} \right) \right| \lambda^{S(\mathbf{r}_n)} \prod_{j=1}^n \left[|D^j \left(\frac{\hat{v}}{E} \right)| \right]^{r_j}, \quad (\text{C.17})$$

where, $R(n)$ is defined as in (5.151) and

$$S(\mathbf{r}_n) := \sum_{k=1}^n r_k, \quad (\text{C.18})$$

see (5.152). Notice that $S(\mathbf{r}_n)$ satisfies

$$1 \leq S(\mathbf{r}_n) \leq n \quad (\text{C.19})$$

due to the summation condition

$$\sum_{j=1}^n j r_j = n. \quad (\text{C.20})$$

(C.19) allows us to extract a factor λ in (C.17) since it appears within the sum with power $S(\mathbf{r}_n)$. Then (C.17) implies

$$\left\| D^n \left[h \left(\frac{\lambda \hat{v}}{E} \right) \right] \right\|_{\infty} \leq C_n \lambda \|h\|_{C^n[0, \|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c} \lambda]} \|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c}^n. \quad (\text{C.21})$$

With analogous steps, we have that

$$\left\| \partial_{|p|}^n \left[h \left(\frac{\lambda \hat{v}}{E} \right) \right] \right\|_{\infty} \leq C_n \lambda \|h\|_{C^n[0, \|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c} \lambda]} \|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c}^n. \quad (\text{C.22})$$

We have that $\Omega = E\sqrt{1 + \frac{2\lambda\hat{v}}{E}}$, see (3.19), is radial since \hat{v} is radial. Thus we have that

$$\nabla\Omega(p) = p\left[\sqrt{1 + \frac{2\lambda\hat{v}(p)}{E(p)}} + \frac{|p|}{2}\left(\sqrt{1 + \frac{2\lambda\hat{v}}{E}}\right)'(p)\right]. \quad (\text{C.23})$$

Observe that

$$\begin{aligned} \left|\sqrt{1 + \frac{2\lambda\hat{v}}{E}} - 1\right| &\leq C \frac{\lambda\|\hat{v}\|_{2(\lfloor\frac{\alpha}{2}\rfloor+1),w,c}}{\sqrt{1 + \frac{2\lambda\hat{v}}{E}} + 1} \\ &\leq C\lambda\|\hat{v}\|_{2(\lfloor\frac{\alpha}{2}\rfloor+1),w,c}. \end{aligned} \quad (\text{C.24})$$

Then (C.22), and (C.24) imply that

$$\nabla\Omega(p) = (1 + \lambda m_\Omega(p))p \quad (\text{C.25})$$

$$\|m_\Omega\|_\infty \leq C\|\hat{v}\|_{2(\lfloor\frac{\alpha}{2}\rfloor+1),w,c}. \quad (\text{C.26})$$

Next, using the multivariate Leibniz rule together with (C.24), we then find that

$$\begin{aligned} D^2\Omega &= \sqrt{1 + \frac{2\lambda\hat{v}}{E}}I + 2p \vee D\sqrt{1 + \frac{2\lambda\hat{v}}{E}} + ED^2\sqrt{1 + \frac{2\lambda\hat{v}}{E}} \\ &= I + \lambda A \end{aligned} \quad (\text{C.27})$$

for some bounded matrix A . Using (C.21) with $h(x) = \sqrt{1 + 2x}$, and (C.24), we have that

$$\|A\|_\infty \leq C\|\hat{v}\|_{2(\lfloor\frac{\alpha}{2}\rfloor+1),w,c}^2 \quad (\text{C.28})$$

for all $\lambda \in (0, 1]$. Here, we used the fact that $\|h\|_{C^2([0, \|\hat{v}\|_{2(\lfloor\frac{\alpha}{2}\rfloor+1),w,c}]})} \leq C$.

Finally, the multivariate Leibniz rule implies

$$\begin{aligned} \|D^3\Omega\|_\infty &\leq C \sum_{k=0}^3 \left\| D^k(|p|^2) D^{3-k} \sqrt{1 + \frac{2\lambda\hat{v}}{E}} \right\|_\infty \\ &\leq C \sum_{k=0}^2 \left\| |p|^{2-k} D^{3-k} \sqrt{1 + \frac{2\lambda\hat{v}}{E}} \right\|_\infty. \end{aligned} \quad (\text{C.29})$$

Using (C.17) and the fact that

$$\left\| |p|^{2-k} D^j \left(\frac{\hat{v}}{E} \right) \right\|_\infty \leq C\|\hat{v}\|_{2(\lfloor\frac{\alpha}{2}\rfloor+1),w,c} \quad (\text{C.30})$$

for $j \in \{1, 2, 3\}$, we find that

$$\|D^3\Omega\|_\infty \leq C\lambda\|\hat{v}\|_{2(\lfloor\frac{\alpha}{2}\rfloor+1),w,c}^3. \quad (\text{C.31})$$

This concludes the proof. \square

Lemma C.4. *Let*

$$F_2(\mathbf{p}_2) := \sigma_1\Omega(p_1) + \sigma_2\Omega(p_2) + \sigma_{12}\Omega(p_{12}) \quad (\text{C.32})$$

and

$$T := \begin{pmatrix} (\sigma_1 + \sigma_{12})I & \sigma_{12}I \\ \sigma_{12}I & (\sigma_2 + \sigma_{12})I \end{pmatrix}. \quad (\text{C.33})$$

F_2 satisfies

$$\nabla F_2(0) = 0, \quad (\text{C.34})$$

$$\|\det(D^2 F_2) - \sigma_{12}(\sigma_{12}\sigma_1\sigma_2 + \sigma_1 + \sigma_2)^3\|_\infty \leq C\lambda\|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c}^2, \quad (\text{C.35})$$

$$\|D^2 F_2\|_\infty \leq C \quad (\text{C.36})$$

For all $\mathbf{p}_2, \mathbf{k}_2 \in \mathbb{R}^6$, and all $\lambda > 0$ small enough, dependent on $\|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c}$, we have that

$$|D^2 F_2(\mathbf{k}_2)\mathbf{p}_2| \in \left[\frac{|T\mathbf{p}_2|}{2}, \frac{3|T\mathbf{p}_2|}{2} \right]. \quad (\text{C.37})$$

In particular, in this case, we have

$$|D^2 F_2(\mathbf{k}_2)\mathbf{p}_2| \geq \begin{cases} |\mathbf{p}_2|/2, & \text{if } \sigma_1 = \sigma_2 = \pm\sigma_{12}, \\ |p_1|/2, & \text{if } \sigma_1 = -\sigma_2 = \sigma_{12}, \\ |p_2|/2, & \text{if } -\sigma_1 = \sigma_2 = \sigma_{12}. \end{cases}$$

Proof. Thanks to Lemma C.3, we have that

$$\nabla \Omega(p) = p(1 + \lambda m_\Omega(p)) \quad (\text{C.38})$$

with

$$\|m_\Omega\|_\infty \leq C\|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c}. \quad (\text{C.39})$$

This immediately implies (C.34).

Next, again due to Lemma C.3, we have that

$$D^2 \Omega = I + \lambda A \quad (\text{C.40})$$

for some matrix $A \in \mathbb{R}^{3 \times 3}$ with

$$\|A\|_\infty \leq C\|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c}^2 \quad (\text{C.41})$$

for all $\lambda \in (0, 1)$. Denoting $A_j := A(p_j)$, we obtain that

$$\begin{aligned} D^2 F_2 &= \begin{pmatrix} \sigma_1(I + \lambda A_1) + \sigma_{12}(I + \lambda A_{12}) & \sigma_{12}(I + \lambda A_{12}) \\ \sigma_{12}(I + \lambda A_{12}) & \sigma_2(I + \lambda A_2) + \sigma_{12}(I + \lambda A_{12}) \end{pmatrix} \\ &= \begin{pmatrix} (\sigma_1 + \sigma_{12})I & \sigma_{12}I \\ \sigma_{12}I & (\sigma_2 + \sigma_{12})I \end{pmatrix} + \lambda \begin{pmatrix} \sigma_1 A_1 + \sigma_{12} A_{12} & \sigma_{12} A_{12} \\ \sigma_{12} A_{12} & \sigma_2 A_2 + \sigma_{12} A_{12} \end{pmatrix} \\ &=: T + \lambda \tilde{B}. \end{aligned} \quad (\text{C.42})$$

This identity together with (C.41) immediately implies (C.36). Notice that due to (C.41), we have that

$$\|\tilde{B}\|_\infty \leq C\|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c}^2. \quad (\text{C.43})$$

Now, observe that

$$\begin{aligned} \det(T) &= \det \left(\begin{pmatrix} 0 & -(\sigma_{12}\sigma_1\sigma_2 + \sigma_1 + \sigma_2)I \\ \sigma_{12}I & (\sigma_2 + \sigma_{12})I \end{pmatrix} \right) \\ &= -\det \left(\begin{pmatrix} \sigma_{12}I & (\sigma_2 + \sigma_{12})I \\ 0 & -(\sigma_{12}\sigma_1\sigma_2 + \sigma_1 + \sigma_2)I \end{pmatrix} \right) \\ &= \sigma_{12}(\sigma_{12}\sigma_1\sigma_2 + \sigma_1 + \sigma_2)^3 \neq 0. \end{aligned} \quad (\text{C.44})$$

Thus we may rewrite $T + \lambda \tilde{B} =: (I + \lambda B)T$ with

$$\|B\|_\infty \leq C \|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c}^2. \quad (\text{C.45})$$

due to (C.43) and (C.44). Then, we have that

$$\|\det(I + \lambda B) - 1\|_\infty \leq C\lambda \|B\|_\infty \quad (\text{C.46})$$

for all $\lambda \in (0, 1)$.

Finally, Gershgorin's Circle Theorem implies that

$$\begin{aligned} \sigma(|I + \lambda B|^2) &\subseteq [1 - 2\lambda \|B\|_\infty - \lambda^2 \|B\|_\infty^2, 1 + 2\lambda \|B\|_\infty + \lambda^2 \|B\|_\infty^2] \\ &\subseteq \left[\frac{1}{4}, \frac{9}{4} \right] \end{aligned} \quad (\text{C.47})$$

for all $\lambda \in (0, 1)$ small enough, dependent on $\|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c}$. Then we have that

$$|D^2 F_2(\mathbf{k}_2) \mathbf{p}_2|^2 = (T \mathbf{p}_2)^T |I + \lambda B(\mathbf{k}_2)|^2 T \mathbf{p}_2, \quad (\text{C.48})$$

which together with (C.47) implies that

$$|D^2 F_2(\mathbf{k}_2) \mathbf{p}_2| \in \left[\frac{|T \mathbf{p}_2|}{2}, \frac{3|T \mathbf{p}_2|}{2} \right]. \quad (\text{C.49})$$

This finishes the proof. \square

C.1. Error bounds due to HFB evolution.

Lemma C.5. *We have that*

$$\begin{aligned} &\frac{\lambda^j}{N} |\text{BoI}_c^{(j)}(f_0)[J](T; \lambda)| \\ &\leq C \|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c} \|f_0\|_{2(\lfloor \frac{r}{2} \rfloor + 1), c}^r T(1 + \log(T)) \|J\|_{W^{1, \infty}} \frac{\lambda |\log(\lambda)|}{N} \end{aligned} \quad (\text{C.50})$$

for $j \in \{1, 2\}$, and all $\lambda > 0$ small enough, dependent on $\|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c}$.

Proof. $\frac{\lambda^j}{N} |\text{BoI}_c^{(j)}(f_0)[J](T; \lambda)|$ are of the form

$$\frac{\lambda^{j-2}}{N} \int_{\mathbb{R}^6} d\mathbf{p}_2 \int_{\Delta[T, 2]} d\mathbf{S}_2 e^{-i(F_1(\mathbf{p}_2)S_1 + F_2(\mathbf{p}_2)S_2)/\lambda^2} H(\mathbf{p}_2) \quad (\text{C.51})$$

with

$$F_2(\mathbf{p}_2) = \sigma_1 \Omega(p_1) + \sigma_2 \Omega(p_2) + \sigma_{12} \Omega(p_{12}) \quad (\text{C.52})$$

for some $\sigma_j \in \{\pm 1\}$ and $H : \mathbb{R}^6 \rightarrow \mathbb{R}$. Employing the bound on $D^2 F_2$ given in Lemma C.4 and the diamagnetic inequality $|\nabla|h| \leq |\nabla h|$, we find the estimate

$$\begin{aligned} \left\| \nabla \cdot \frac{\nabla F_2 |H|}{|\nabla F_2|^2} \right\|_1 &\leq \left\| \frac{\Delta F_2 H}{|\nabla F_2|^2} \right\|_1 + \left\| \frac{\nabla F_2^T D^2 F_2 \nabla F_2 H}{|\nabla F_2|^4} \right\|_1 + \left\| \frac{\nabla |H|}{|\nabla F_2|} \right\|_1 \\ &\leq C \left(\left\| \frac{H}{|\nabla F_2|^2} \right\|_1 + \left\| \frac{\nabla H}{|\nabla F_2|} \right\|_1 \right). \end{aligned} \quad (\text{C.53})$$

Let $B_{F_2} := \{|\nabla F_2| \leq 1\}$ denote the unit ball induced by ∇F_2 , and let $J_{\nabla F_2} := |\det(D^2 F_2)|$. Notice that for any $m \in [0, 4)$ and test function h , we have that

$$\left\| \frac{h}{|\nabla F_2|^m} \right\|_1 \leq \left\| \frac{J_{\nabla F_2}^{\frac{12-m}{24-3m}}}{|\nabla F_2|^m} \right\|_{L^{\frac{24-3m}{12-m}}(B_{F_2})} \left\| \frac{h}{J_{\nabla F_2}^{\frac{12-m}{24-3m}}} \right\|_{L^{\frac{24-3m}{12-2m}}(B_{F_2})}$$

$$\begin{aligned}
& + \left\| \frac{h}{|\nabla F_2|^m} \right\|_{L^1(B_{F_2}^c)} \\
& \leq C \left(\|h\|_{\frac{24-3m}{12-2m}} + \|h\|_1 \right)
\end{aligned} \tag{C.54}$$

for all $\lambda > 0$ small enough, dependent on $\|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c}$. Here, we used Lemma C.4 to bound $J_{\nabla F_2} \geq C$ for $\lambda \in (0, 1)$ small enough, substitution, and the fact that

$$\int_{B_1} \frac{d\mathbf{x}_6}{|\mathbf{x}_6|^m \frac{24-3m}{12-m}} \leq C \int_0^1 \frac{d|\mathbf{x}_6|}{|\mathbf{x}_6|^m \frac{24-3m}{12-m} - 5} < \infty \tag{C.55}$$

since, due to $m < 4$,

$$m \frac{24-3m}{12-m} - 5 = -3 \frac{(m-5)^2 - 1}{12-m} + 1 < 1. \tag{C.56}$$

In particular, (C.53) and (C.54) imply

$$\begin{aligned}
\left\| \nabla \cdot \frac{\nabla F_2 |H|}{|\nabla F_2|^2} \right\|_1 & \leq C \left(\|H\|_{W^{1,1}} + \|H\|_{\frac{9}{4}} + \|\nabla H\|_{\frac{21}{10}} \right) \\
& \leq C \left(\|H\|_{W^{1,1}} + \|H\|_{W^{1, \frac{9}{4}}} \right),
\end{aligned} \tag{C.57}$$

where we employed the fact that $m \mapsto \frac{24-3m}{12-2m}$ is an increasing function, and interpolation. With that, we apply Lemma C.2 to obtain

$$\begin{aligned}
& \left| \int d\mathbf{p}_2 \int_{\Delta[t, 2]} ds_2 e^{-i(F_1(\mathbf{p}_2)s_1 + F_2(\mathbf{p}_2)s_2)} H(\mathbf{p}_2) \right| \\
& \leq Ct \left(\log(1+t) \left\| \nabla \cdot \frac{\nabla F_2 |H|}{|\nabla F_2|^2} \right\|_1 + \|H\|_1 \right) \\
& \leq Ct(1 + \log(1+t)) \left(\|H\|_{W^{1,1}} + \|H\|_{W^{1, \frac{9}{4}}} \right).
\end{aligned} \tag{C.58}$$

As a consequence of (C.58) and using the chain rule, we obtain that

$$\begin{aligned}
& \frac{\lambda^j}{N} |\text{Bol}_c^{(j)}(f_0)[J](T; \lambda)| \\
& \leq C \|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c} \|f_0\|_{2(\lfloor \frac{r}{2} \rfloor + 1), c, r} T(1 + \log(T)) \|J\|_{W^{1, \infty}} \frac{\lambda |\log(\lambda)|}{N}
\end{aligned} \tag{C.59}$$

for $j \in \{1, 2\}$, and all $\lambda > 0$ small enough, dependent on $\|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c}$. This finishes the proof. \square

C.2. Mollified energy conservation.

Lemma C.6. *Let $\text{err}_2^{(fd)}(t; f[J])$ be defined as in (5.211). Then*

$$|\text{err}_2^{(fd)}(t; f[J])| \leq C \|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c} \|f_0\|_{2(\lfloor \frac{r}{2} \rfloor + 1), c, r} T \|J\|_{W^{2, \infty}} \frac{\lambda}{N} \tag{C.60}$$

for all $\lambda > 0$ small enough, dependent on $\|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c}$.

Proof. We want to apply integration by parts to

$$\begin{aligned}
\text{err}_2^{(fd)}(t; f[J]) & = \frac{1}{N} \int_0^\lambda d\tau \int d\omega \delta'_{\frac{2\lambda^2}{T}}(\omega) \\
& \int_{\mathcal{E}(\tau, \mathbf{p}_2) = \omega} d\mathcal{H}^5 \frac{\partial_\tau \mathcal{E}(\tau, \mathbf{p}_2) H(\mathbf{p}_2)}{|\nabla_{\mathbf{p}_2} \mathcal{E}(\tau, \mathbf{p}_2)|}.
\end{aligned} \tag{C.61}$$

For that, we need to establish a uniform bound on

$$\int_{\mathcal{E}(\tau, \mathbf{p}_2) = \omega} d\mathcal{H}^5 \frac{\partial_\tau \mathcal{E}(\tau, \mathbf{p}_2) H(\mathbf{p}_2)}{|\nabla_{\mathbf{p}_2} \mathcal{E}(\tau, \mathbf{p}_2)|}. \quad (\text{C.62})$$

The divergence theorem implies that

$$\left| \int_{\mathcal{E}(\tau, \mathbf{p}_2) = \omega} d\mathcal{H}^5 \frac{\partial_\tau \mathcal{E}(\tau, \mathbf{p}_2) H(\mathbf{p}_2)}{|\nabla_{\mathbf{p}_2} \mathcal{E}(\tau, \mathbf{p}_2)|} \right| \leq \left\| \nabla_{\mathbf{p}_2} \cdot \left(\frac{\nabla_{\mathbf{p}_2} \mathcal{E}(\tau) \partial_\tau \mathcal{E}(\tau) H}{|\nabla_{\mathbf{p}_2} \mathcal{E}(\tau)|^2} \right) \right\|_1. \quad (\text{C.63})$$

We compute

$$\partial_\tau \Omega_\tau = \frac{2\hat{v}}{\sqrt{1 + \frac{2\tau\hat{v}}{E}}}. \quad (\text{C.64})$$

Then $\partial_\tau \Omega_\tau$ satisfies

$$\|\partial_\tau \Omega_\tau\|_\infty \leq C \|\hat{v}\|_{2([\frac{\tau}{2}] + 1), w, c}, \quad (\text{C.65})$$

$$\begin{aligned} \|\nabla_p \partial_\tau \Omega_\tau\|_\infty &\leq C \left(\|\hat{v}\|_{2([\frac{\tau}{2}] + 1), w, c} + \|\hat{v}\|_\infty \left\| D \left(1 + \frac{2\tau\hat{v}}{E} \right)^{-\frac{1}{2}} \right\|_\infty \right) \\ &\leq C \|\hat{v}\|_{2([\frac{\tau}{2}] + 1), w, c}, \end{aligned} \quad (\text{C.66})$$

$$\begin{aligned} \|D_p^2 \partial_\tau \Omega_\tau\|_\infty &\leq C \left(\|D^2 \hat{v}\|_\infty + \|D \hat{v}\|_\infty \left\| D \left(1 + \frac{2\tau\hat{v}}{E} \right)^{-\frac{1}{2}} \right\|_\infty \right. \\ &\quad \left. + \|\hat{v}\|_\infty \left\| D^2 \left(1 + \frac{2\tau\hat{v}}{E} \right)^{-\frac{1}{2}} \right\|_\infty \right) \\ &\leq C \|\hat{v}\|_{2([\frac{\tau}{2}] + 1), w, c} \end{aligned} \quad (\text{C.67})$$

for all $\tau \in [0, \lambda]$, $\lambda \in (0, 1]$. Here we applied the Leibniz Formula, Lemma C.3. Then (C.65)-(C.67) imply

$$\|\partial_\tau \mathcal{E}(\tau, \mathbf{p}_2)\|_{W^{2, \infty}} \leq C \|\hat{v}\|_{2([\frac{\tau}{2}] + 1), w, c} \quad (\text{C.68})$$

for all $\tau \in [0, \lambda]$, $\lambda \in (0, 1)$. Following the steps of (C.53) in Lemma C.5, we find that

$$\begin{aligned} \left\| \nabla_{\mathbf{p}_2} \cdot \left(\frac{\nabla_{\mathbf{p}_2} \mathcal{E}(\tau) \partial_\tau \mathcal{E}(\tau) H}{|\nabla_{\mathbf{p}_2} \mathcal{E}(\tau)|^2} \right) \right\|_1 &\leq C \left(\left\| \frac{\partial_\tau \mathcal{E}(\tau) H}{|\nabla_{\mathbf{p}_2} \mathcal{E}(\tau)|^2} \right\|_1 \right. \\ &\quad \left. + \left\| \frac{\nabla_{\mathbf{p}_2} (\partial_\tau \mathcal{E}(\tau) H)}{|\nabla_{\mathbf{p}_2} \mathcal{E}(\tau)|} \right\|_1 \right) \end{aligned} \quad (\text{C.69})$$

due to Lemma C.4. Using the Mean-Value Theorem together with $\nabla_{\tau, \mathbf{p}_2} \mathcal{E}(\tau, 0) = 0$, see Lemma C.4, we find that, for some $\zeta_{\tau, \mathbf{p}_2} \in [0, \mathbf{p}_2]$,

$$\nabla_{\mathbf{p}_2} \mathcal{E}(\tau, \mathbf{p}_2) = D_{\mathbf{p}_2}^2 \mathcal{E}(\tau, \zeta_{\tau, \mathbf{p}_2}) \mathbf{p}_2. \quad (\text{C.70})$$

Using C.4 again, we conclude that

$$|\nabla_{\mathbf{p}_2} \mathcal{E}(\tau, \mathbf{p}_2)| \geq \frac{1}{2} \left| \begin{pmatrix} p_2 \\ p_1 \end{pmatrix} \right| = \frac{|\mathbf{p}_2|}{2} \quad (\text{C.71})$$

for all $\lambda > 0$ small enough, dependent on $\|\hat{v}\|_{2([\frac{\tau}{2}] + 1), w, c}$. Thus, we obtain that

$$\left| \frac{D^\ell [(\hat{v}(p_1) + \hat{v}(p_2))^2 f_0(p_{12})]}{|\nabla_{\mathbf{p}_2} \mathcal{E}(\tau, \mathbf{p}_2)|^m} \right|$$

$$\begin{aligned} &\leq C_{\|\hat{v}\|_{2(\lfloor \frac{\ell}{2} \rfloor + 1), w, c}, \|f_0\|_{2(\lfloor \frac{\ell}{2} \rfloor + 1), c}} \left(\frac{1}{|\mathbf{p}_2|^m} \mathbb{1}_{B_1}(\mathbf{p}_2) \right. \\ &\quad \left. + D^\ell [(\hat{v}(p_1) + \hat{v}(p_2))^2 f_0(p_{12})] \mathbb{1}_{B_1^c}(\mathbf{p}_2) \right) \end{aligned} \quad (\text{C.72})$$

for all $\ell, m \in \mathbb{N}_0$, $\ell \leq 2$. Hence,

$$\left\| \frac{D^\ell [(\hat{v}(p_1) + \hat{v}(p_2))^2 f_0(p_{12})]}{|\nabla_{\mathbf{p}_2} \mathcal{E}(\tau)|^m} \right\|_1 \leq C_{\|\hat{v}\|_{2(\lfloor \frac{\ell}{2} \rfloor + 1), w, c}, \|f_0\|_{2(\lfloor \frac{\ell}{2} \rfloor + 1), c}} \quad (\text{C.73})$$

for all $\ell, m \in \mathbb{N}_0$, $\ell \leq 2$, $m \leq 5$, and all $\lambda > 0$ small enough, dependent on $\|\hat{v}\|_{2(\lfloor \frac{\ell}{2} \rfloor + 1), w, c}$. With analogous arguments, we may replace $f_0(p_{12})$ in (C.73) by $f_0(p_{12})f_0(p_1)$, $f_0(p_{12})f_0(p_2)$, or $f_0(p_1)f_0(p_2)$ and obtain an analogous inequality. Then, (C.63), (C.68), (C.69), (C.73), and the definition (5.210) of H imply

$$\left| \int_{\mathcal{E}(\tau, \mathbf{p}_2) = \omega} d\mathcal{H}^5 \frac{\partial_\tau \mathcal{E}(\tau, \mathbf{p}_2) H(\mathbf{p}_2)}{|\nabla_{\mathbf{p}_2} \mathcal{E}(\tau, \mathbf{p}_2)|} \right| \leq C_{\|\hat{v}\|_{2(\lfloor \frac{\ell}{2} \rfloor + 1), w, c}, \|f_0\|_{2(\lfloor \frac{\ell}{2} \rfloor + 1), c}} T \|J\|_{W^1} \quad (\text{C.74})$$

for all $\omega \in \mathbb{R}$, $\tau \in [0, \lambda]$, λ small enough, dependent on $\|\hat{v}\|_{2(\lfloor \frac{\ell}{2} \rfloor + 1), w, c}$.

Next, by integration by parts w.r.t. $d\omega$, $\delta'_{\frac{2\lambda^2}{T}}(\omega) \rightarrow 0$ as $|\omega| \rightarrow \infty$, (C.74), and employing Lemma C.1, (C.61) implies that

$$\begin{aligned} \text{err}_2^{(fd)}(t; f[J]) &= -\frac{1}{N} \int_0^\lambda d\tau \int d\omega \delta_{\frac{2\lambda^2}{T}}(\omega) \int_{\mathcal{E}(\tau, \mathbf{p}_2) = \omega} \frac{d\mathcal{H}^5}{|\nabla_{\mathbf{p}_2} \mathcal{E}(\tau, \mathbf{p}_2)|} \\ &\quad \nabla_{\mathbf{p}_2} \cdot \left(\frac{\nabla_{\mathbf{p}_2} \mathcal{E}(\tau, \mathbf{p}_2) \partial_\tau \mathcal{E}(\tau, \mathbf{p}_2) H(\mathbf{p}_2)}{|\nabla_{\mathbf{p}_2} \mathcal{E}(\tau, \mathbf{p}_2)|^2} \right) \end{aligned} \quad (\text{C.75})$$

Lemma C.1 implies

$$\begin{aligned} &\left| \int_{\mathcal{E}(\tau, \mathbf{p}_2) = \omega} \frac{d\mathcal{H}^5}{|\nabla_{\mathbf{p}_2} \mathcal{E}(\tau, \mathbf{p}_2)|} \nabla_{\mathbf{p}_2} \cdot \left(\frac{\nabla_{\mathbf{p}_2} \mathcal{E}(\tau, \mathbf{p}_2) \partial_\tau \mathcal{E}(\tau, \mathbf{p}_2) H(\mathbf{p}_2)}{|\nabla_{\mathbf{p}_2} \mathcal{E}(\tau, \mathbf{p}_2)|^2} \right) \right| \\ &\leq \left\| \nabla_{\mathbf{p}_2} \cdot \left[\frac{\nabla_{\mathbf{p}_2} \mathcal{E}(\tau, \mathbf{p}_2)}{|\nabla_{\mathbf{p}_2} \mathcal{E}(\tau, \mathbf{p}_2)|^2} \nabla_{\mathbf{p}_2} \cdot \left(\frac{\nabla_{\mathbf{p}_2} \mathcal{E}(\tau, \mathbf{p}_2) \partial_\tau \mathcal{E}(\tau, \mathbf{p}_2) H(\mathbf{p}_2)}{|\nabla_{\mathbf{p}_2} \mathcal{E}(\tau, \mathbf{p}_2)|^2} \right) \right] \right\|_1 \\ &\leq C \left(\|D_{\mathbf{p}_2}^2 \mathcal{E}(\tau)\|_\infty^2 \left\| \frac{\partial_\tau \mathcal{E}(\tau) H}{|\nabla_{\mathbf{p}_2} \mathcal{E}(\tau, \mathbf{p}_2)|^4} \right\|_1 + \|D_{\mathbf{p}_2}^3 \mathcal{E}(\tau)\|_\infty \left\| \frac{\partial_\tau \mathcal{E}(\tau) H}{|\nabla_{\mathbf{p}_2} \mathcal{E}(\tau, \mathbf{p}_2)|^3} \right\|_1 \right. \\ &\quad \left. + \|D_{\mathbf{p}_2}^2 \mathcal{E}(\tau)\|_\infty \left\| \frac{D_{\mathbf{p}_2}(\partial_\tau \mathcal{E}(\tau) H)}{|\nabla_{\mathbf{p}_2} \mathcal{E}(\tau, \mathbf{p}_2)|^3} \right\|_1 + \left\| \frac{D_{\mathbf{p}_2}^2(\partial_\tau \mathcal{E}(\tau) H)}{|\nabla_{\mathbf{p}_2} \mathcal{E}(\tau, \mathbf{p}_2)|^2} \right\|_1 \right). \end{aligned} \quad (\text{C.76})$$

Lemma C.3 implies

$$\|D_{\mathbf{p}_2}^2 \mathcal{E}(\tau)\|_\infty \leq 1 + C \|\hat{v}\|_{2(\lfloor \frac{\ell}{2} \rfloor + 1), w, c}^\tau, \quad (\text{C.77})$$

$$\|D_{\mathbf{p}_2}^3 \mathcal{E}(\tau)\|_\infty \leq C \|\hat{v}\|_{2(\lfloor \frac{\ell}{2} \rfloor + 1), w, c}^3. \quad (\text{C.78})$$

Collecting (C.68), (C.73), (C.76), (C.77), and (C.78), we find that

$$\begin{aligned} &\left| \int_{\mathcal{E}(\tau, \mathbf{p}_2) = \omega} \frac{d\mathcal{H}^5}{|\nabla_{\mathbf{p}_2} \mathcal{E}(\tau, \mathbf{p}_2)|} \nabla_{\mathbf{p}_2} \cdot \left(\frac{\nabla_{\mathbf{p}_2} \mathcal{E}(\tau, \mathbf{p}_2) \partial_\tau \mathcal{E}(\tau, \mathbf{p}_2) H(\mathbf{p}_2)}{|\nabla_{\mathbf{p}_2} \mathcal{E}(\tau, \mathbf{p}_2)|^2} \right) \right| \\ &\leq C_{\|\hat{v}\|_{2(\lfloor \frac{\ell}{2} \rfloor + 1), w, c}, \|f_0\|_{2(\lfloor \frac{\ell}{2} \rfloor + 1), c}} T \|J\|_{W^{2, \infty}}. \end{aligned} \quad (\text{C.79})$$

Employing (C.73), (C.75), and (C.79), we have proved that

$$|\text{err}_2^{(fd)}(t; f[J])| \leq C_{\|\hat{v}\|_{2(\lfloor \frac{\ell}{2} \rfloor + 1), w, c}, \|f_0\|_{2(\lfloor \frac{\ell}{2} \rfloor + 1), c}} \frac{T \|J\|_{W^{2, \infty}}}{N} \int_0^\lambda d\tau \int d\omega \delta_{\frac{2\lambda^2}{T}}(\omega)$$

$$\leq C \|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c} \|f_0\|_{2(\lfloor \frac{r}{2} \rfloor + 1), c} \frac{T\lambda \|J\|_{W^{2, \infty}}}{N}, \quad (\text{C.80})$$

where in the last step, we applied the normalization (5.207) of $\delta_{\frac{2\lambda^2}{T}}$. This finishes the proof. \square

Lemma C.7. *Recall that, due to (5.213),*

$$\begin{aligned} \text{err}_2^{(ec)}(t; f[J]) &= \frac{1}{N} \int d\omega \delta_{\frac{2\lambda^2}{T}}(\omega) \left(\int_{\Delta_{cub} E = \omega} d\mathcal{H}^5 \frac{H}{|\nabla \Delta_{cub} E|} \right. \\ &\quad \left. + \int_{\Delta_{cub} E = 0} d\mathcal{H}^5 \frac{H}{|\nabla \Delta_{cub} E|} \right) \end{aligned} \quad (\text{C.81})$$

with H as defined in (5.210). Then we have that

$$|\text{err}_2^{(ec)}(t; f[J])| \leq C \|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c} \|f_0\|_{2(\lfloor \frac{r}{2} \rfloor + 1), c} \sqrt{T} \|J\|_{W^{2, \infty}} \frac{\lambda}{N} \quad (\text{C.82})$$

for all $\lambda > 0$ small enough, dependent on $\|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c}$.

Proof. We start by writing

$$\text{err}_2^{(ec)}(t; f[J]) =: \frac{1}{N} \int d\omega \delta_{\frac{2\lambda^2}{T}}(\omega) \mathcal{I}(\omega). \quad (\text{C.83})$$

Analogously to (C.74), we have that

$$|\mathcal{I}(\omega)| \leq C \|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c} \|f_0\|_{2(\lfloor \frac{r}{2} \rfloor + 1), c} T \|J\|_{W^{1, \infty}} \frac{1}{N}. \quad (\text{C.84})$$

Notice that, due to Lemma C.1, we have that

$$\mathcal{I}(\omega) = \int_0^\omega d\tau \int_{\Delta_{cub} E = \tau} \frac{d\mathcal{H}^5}{|\nabla \Delta_{cub} E|} \nabla \cdot \left(\frac{\nabla \Delta_{cub} E H}{|\nabla \Delta_{cub} E|^2} \right), \quad (\text{C.85})$$

where the integral respects the orientation of $[0, \omega]$ resp. $[\omega, 0]$. Using Lemma C.1, we thus obtain

$$\begin{aligned} |\mathcal{I}(\omega)| &\leq |\omega| \left\| \nabla \cdot \left[\frac{\nabla \Delta_{cub} E}{|\nabla \Delta_{cub} E|^2} \nabla \cdot \left(\frac{\nabla \Delta_{cub} E H}{|\nabla \Delta_{cub} E|^2} \right) \right] \right\|_1 \\ &\leq C |\omega| \left(\|D^2 \Delta_{cub} E\|_\infty^2 \left\| \frac{H}{|\nabla \Delta_{cub} E|^4} \right\|_1 + \left\| \frac{D^3 \Delta_{cub} E H}{|\nabla \Delta_{cub} E|^3} \right\|_1 \right. \\ &\quad \left. + \|D^2 \Delta_{cub} E\|_\infty \left\| \frac{DH}{|\nabla \Delta_{cub} E|^3} \right\|_1 + \left\| \frac{D^2 H}{|\nabla \Delta_{cub} E|^2} \right\|_1 \right), \end{aligned} \quad (\text{C.86})$$

for all $\lambda > 0$ small enough, dependent on $\|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c}$, analogously to (C.76).

Notice that $D^3 \Delta_{cub} E = 0$ and $\|D^2 \Delta_{cub} E\|_\infty \leq C$. Analogously to (C.80), we thus obtain the upper bound

$$|\mathcal{I}(\omega)| \leq C \|\hat{v}\|_{2(\lfloor \frac{r}{2} \rfloor + 1), w, c} \|f_0\|_{2(\lfloor \frac{r}{2} \rfloor + 1), c} |\omega| T \|J\|_{W^{2, \infty}}. \quad (\text{C.87})$$

We split the integral in (C.83) into the regions $(-\omega_0, \omega_0) \cup (-\omega_0, \omega_0)^c$ with ω_0 to be determined below. Then (C.84) and (C.87) yield

$$\begin{aligned} &|\text{err}_2^{(ec)}(t; f[J])| \\ &\leq \frac{1}{N} \int_{(-\omega_0, \omega_0)} d\omega \delta_{\frac{2\lambda^2}{T}}(\omega) |\mathcal{I}(\omega)| + \int_{(-\omega_0, \omega_0)^c} d\omega \delta_{\frac{2\lambda^2}{T}}(\omega) |\mathcal{I}(\omega)| \end{aligned}$$

$$\begin{aligned} &\leq \frac{C_{\|\hat{v}\|_{2(\lfloor \frac{x}{2} \rfloor + 1), w, c}, \|f_0\|_{2(\lfloor \frac{x}{2} \rfloor + 1), c}} T}{N} \left(\|J\|_{W^{2, \infty}} \int_{(-\omega_0, \omega_0)} d\omega \delta_{\frac{2\lambda^2}{T}}(\omega) |\omega| \right. \\ &\quad \left. + \|J\|_{W^{1, \infty}} \int_{(-\omega_0, \omega_0)^c} d\omega \delta_{\frac{2\lambda^2}{T}}(\omega) \right). \end{aligned} \quad (\text{C.88})$$

Employing the normalization condition (5.207) and the decay condition (5.208) on $\delta_{\frac{2\lambda^2}{T}}$, we thus obtain

$$|\text{err}_2^{(ec)}(t; f[J])| \leq \frac{C_{\|\hat{v}\|_{2(\lfloor \frac{x}{2} \rfloor + 1), w, c}, \|f_0\|_{2(\lfloor \frac{x}{2} \rfloor + 1), c}} T \|J\|_{W^{2, \infty}}}{N} \left(\omega_0 + \frac{\lambda^2}{\omega_0 T} \right). \quad (\text{C.89})$$

By now choosing $\omega_0 = \frac{\lambda}{\sqrt{T}}$, we have hence proved that

$$|\text{err}_2^{(ec)}(t; f[J])| \leq \frac{C_{\|\hat{v}\|_{2(\lfloor \frac{x}{2} \rfloor + 1), w, c}, \|f_0\|_{2(\lfloor \frac{x}{2} \rfloor + 1), c}} \sqrt{T} \|J\|_{W^{2, \infty}} \lambda}{N}. \quad (\text{C.90})$$

□

Acknowledgments: T.C. gratefully acknowledges support by the NSF through grants DMS-1151414 (CAREER), DMS-1716198, DMS-2009800, and the RTG Grant DMS-1840314 *Analysis of PDE*. M.H. was supported by a University of Texas at Austin Provost Graduate Excellence Fellowship, and by NSF grants DMS-1716198 and DMS-2009800 through T.C. M.H. would like to thank Darren King and Daniel Weser for helpful discussions and comments on the use of Geometric Measure Theory for some calculations. We also thank Esteban Cárdenas, Jacky Chong, Ryan Denlinger, Jürg Fröhlich, Irene Gamba, Laurent Lafleche, Nataša Pavlović, Israel Michael Sigal, and Avy Soffer for inspiring discussions, and some references. We thank the anonymous referees for valuable comments.

Conflict of Interest Statement: No conflicts of interests are connected to this article.

Data availability: Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

REFERENCES

- [1] A.A. Abrikosov, L.P. Gorkov, I.E. Dzyaloshinski, *Methods of quantum field theory in statistical physics*. Republication, with corrections, of the revised English edition (1963). Translated from the Russian and edited by Richard A. Silverman. Dover Publications, Inc., New York, xv+352 (1975).
- [2] R. Adami, F. Golse, A. Teta, *Rigorous derivation of the cubic NLS in dimension one*, J. Stat. Phys., **127**(6), pp. 1193-1220, (2007).
- [3] A. Adhikari, C. Brennecke, B. Schlein, *Bose-Einstein condensation beyond the Gross-Pitaevskii regime*, In Annales Henri Poincaré, pp. 1–71, Springer International Publishing (2020).
- [4] S. Akopian, *Global L^p solutions of the Boltzmann equation with an angle-potential concentrated collision kernel and convergence to a Landau solution*, Doctoral dissertation, UT Austin (2017).
- [5] S. Akopian, I.M. Gamba, *Existence and uniqueness of L^p solutions to the Boltzmann equation with an angle-potential concentrated collision kernel*, arXiv preprint arXiv:1611.06316 (2016).

- [6] R. Alexandre, Y. Morimoto, S. Ukai, C.-J. Xu, T. Yang, *The Boltzmann equation without angular cutoff in the whole space: II, Global existence for hard potential*, Anal. Appl. (Singap.), **9**(2):113–134, 2011.
- [7] R. Alexandre, Y. Morimoto, S. Ukai, C.-J. Xu, T. Yang, *Smoothing effect of weak solutions for the spatially homogeneous Boltzmann equation without angular cutoff*, Kyoto J. Math., **52**(3):433–463, 2012.
- [8] R. Alexandre, C. Villani, *On the Boltzmann equation for long-range interactions*, Comm. Pure Appl. Math., **55**(1):30–70, 2002.
- [9] O.E. Alon, A.I. Streltsov, L.S. Cederbaum, *Multiconfigurational time-dependent Hartree method for bosons: Many-body dynamics of bosonic systems*, Physical Review A, **77**(3), p.033613 (2008).
- [10] R. Alonso, *Existence of global solutions to the Cauchy problem for the inelastic Boltzmann equation with near-vacuum data*, Indiana University Mathematics Journal Vol. **58**, No. 3 (2009), pp. 999–1022.
- [11] R. Alonso, V. Bagland, L. Desvillettes, B. Lods, *About the Use of Entropy Production for the Landau-Fermi-Dirac Equation*, J. Stat. Phys., **183**(1), pp. 1–27 (2021).
- [12] R. Alonso, V. Bagland, B. Lods, *Long time dynamics for the Landau-Fermi-Dirac equation with hard potentials*, Journal of Differential Equations, **270**, pp. 596–663 (2021).
- [13] R. Alonso, I.M. Gamba, *Distributional and Classical Solutions to the Cauchy Boltzmann Problem for Soft Potentials with Integrable Angular Cross Section*, J. Stat. Phys. **137**: 11471165 (2009).
- [14] R. Alonso, I.M. Gamba, M.B. Tran, *The Cauchy problem for the quantum Boltzmann equation for bosons at very low temperature*, arXiv preprint arXiv:1609.07467 (2018).
- [15] Z. Ammari, M. Falconi, B. Pawilowski, *On the rate of convergence for the mean field approximation of many-body quantum dynamics*, Commun. Math. Sci. **14**, no. 5, pp. 1417–1442 (2016).
- [16] Z. Ammari, Q. Liard, C. Rouffort, *On Well-Posedness for General Hierarchy Equations of Gross-Pitaevskii and Hartree Type*, Archive for Rational Mechanics and Analysis, **238**(2), pp.845–900 (2020).
- [17] Z. Ammari, F. Nier, *Mean field limit for bosons and infinite dimensional phase-space analysis*, Annales Henri Poincaré. Vol. **9** No. 8, pp. 1503–1574 (2008).
- [18] I. Ampatzoglou, C. Collot, P. Germain, *Derivation of the kinetic wave equation for quadratic dispersive problems in the inhomogeneous setting*, arXiv preprint arXiv:2107.11819 (2021).
- [19] I. Ampatzoglou, I.M. Gamba, N. Pavlović, M. Tasković, *Global well-posedness of a binary-ternary Boltzmann equation*, arXiv preprint arXiv:1910.14476 (2019).
- [20] I. Ampatzoglou, J. Miller, N. Pavlović, *A Rigorous Derivation of a Boltzmann System for a Mixture of Hard-Sphere Gases*, arXiv preprint arXiv:2104.14480 (2021).
- [21] I. Ampatzoglou, N. Pavlović, *A rigorous derivation of a ternary Boltzmann equation for a classical system of particles*, Comm. Math. Phys. **387**, no. 2, 793–863 (2021).
- [22] I. Anapolitanos, M. Hott, *A simple proof of convergence to the Hartree dynamics in Sobolev trace norms*, Journal of Mathematical Physics **57**.12: 122108 (2016).
- [23] I. Anapolitanos, M. Hott, D. Hundertmark, *Derivation of the Hartree equation for compound Bose gases in the mean field limit*, Reviews in Mathematical Physics **29**.07: 1750022 (2017).
- [24] M.H. Anderson, J.R. Ensher, M.R. Matthews, C.E. Wieman, E.A. Cornell, *Observation of Bose-Einstein condensation in a dilute atomic vapor*, science, **269**(5221), 198–201 (1995).
- [25] L. Arkeryd, *Intermolecular forces of infinite range and the Boltzmann equation*, Archive for Rational Mechanics and Analysis, **77**(1), pp.11–21 (1981).
- [26] L. Arkeryd, *A quantum Boltzmann equation for Haldane statistics and hard forces; the space-homogeneous initial value problem*, Comm. Math. Phys., **298**, pp. 573–583 (2010).
- [27] L. Arkeryd, A. Nouri, *Bose condensates in interaction with excitations: a two-component space-dependent model close to equilibrium*, J. Stat. Phys. **160**, no. 1, pp. 209–238 (2015).
- [28] L. Arkeryd and A. Nouri, *Well-posedness of the Cauchy problem for a space-dependent anyon Boltzmann equation*, SIAM J. Math. Anal., **47**, pp. 4720–4742 (2015).
- [29] L. Arkeryd, A. Nouri, *On the Cauchy problem with large data for a space-dependent Boltzmann-Nordheim boson equation*, Commun. Math. Sci. **15**, no. 5, pp. 1247–1264 (2017).
- [30] L. Arkeryd, A. Nouri, *On a Boltzmann equation for Haldane statistics*, (English summary), Kinet. Relat. Models **12**, no. 2, pp. 323–346 (2019).

- [31] V. Bach, S. Breteaux, T. Chen, J. Fröhlich, I.M. Sigal, *The time-dependent Hartree-Fock-Bogoliubov equations for bosons*, arXiv preprint arXiv.org/abs/1602.05171 (2016)
- [32] V. Bach, S. Breteaux, T. Chen, J. Fröhlich, I.M. Sigal, *On the Hartree-Fock-Bogoliubov equations*, arXiv preprint arXiv:1805.04689 (2018)
- [33] V. Bach, S. Breteaux, S.P. Petrat, P. Pickl, T. Tzaneteas *Kinetic energy estimates for the accuracy of the time-dependent Hartree-Fock approximation with Coulomb interaction*, Journal de Mathématiques Pures et Appliquées **105**(1): pp. 1–30 (2016).
- [34] V. Bach, T. Chen, J. Fröhlich, I.M. Sigal *Smooth Feshbach map and operator-theoretic renormalization group methods*, J. Funct. Anal., **203**(1), pp. 44–92 (2003).
- [35] V. Bach, J. Fröhlich, I.M. Sigal, *Renormalization group analysis of spectral problems in quantumfield theory*, Adv. Math. **137**, 205–298, (1998).
- [36] G.C. Bae, J.W. Jang, S.B. Yun, *The relativistic quantum Boltzmann equation near equilibrium*, Archive for Rational Mechanics and Analysis, **240**(3), pp.1593–1644 (2021).
- [37] V. Bagland, *Well-posedness for the spatially homogeneous Landau-Fermi-Dirac equation for hard potentials*, Proceedings of the Royal Society of Edinburgh Section A: Mathematics, **134**(3), pp.415–447 (2004).
- [38] V. Bagland, M. Lemou, *Equilibrium states for the Landau-Fermi-Dirac equation*, Banach Center Publ, **66**, pp.29–37 (2004).
- [39] C. Bardos, F. Golse, N. J. Mauser, *Weak coupling limit of the N -particle Schrödinger equation*, Methods and Applications of Analysis, **7**(2), pp. 275–294, (2000).
- [40] J. Bandyopadhyay, J.J.L. Velázquez, *Blow-up rate estimates for the solutions of the bosonic Boltzmann-Nordheim equation*, J. Math. Phys. **56**, 063302 (2015).
- [41] N. Bellomo, G. Toscani, *On the Cauchy problem for the nonlinear Boltzmann equation: global existence, unique-ness and asymptotic stability*, J. Math. Phys. **26** (1985), no. 2, 334338.
- [42] D. Benedetto, F. Castella, R. Esposito, M. Pulvirenti, *Some considerations on the derivation of the nonlinear quantum Boltzmann equation*, J. Stat. Phys. **116**(1–4), pp. 381–410 (2004).
- [43] D. Benedetto, F. Castella, R. Esposito, M. Pulvirenti, *On the weak-coupling limit for bosons and fermions*, Mathematical Models and Methods in Applied Sciences, **15**(12), pp.1811–1843 (2005).
- [44] D. Benedetto, F. Castella, R. Esposito, M. Pulvirenti, *Some considerations on the derivation of the nonlinear quantum Boltzmann equation II: the low density regime*, Journal of statistical physics, **124**(2), pp.951–996 (2006).
- [45] D. Benedetto, F. Castella, R. Esposito, M. Pulvirenti, *From the N -body Schrödinger equation to the quantum Boltzmann equation: a term-by-term convergence result in the weak coupling regime*, Communications in mathematical physics **277**(1), pp.1–44 (2008).
- [46] N. Benedikter, P.T. Nam, M. Porta, B. Schlein, R. Seiringer, *Optimal upper bound for the correlation energy of a Fermi gas in the mean-field regime*, Communications in Mathematical Physics **374**(3), pp.2097–2150, (2020).
- [47] N. Benedikter, P.T. Nam, M. Porta, B. Schlein, R. Seiringer, *Correlation energy of a weakly interacting Fermi gas*, Inventiones mathematicae, pp.1–95 (2021).
- [48] N. Benedikter, P.T. Nam, M. Porta, B. Schlein, R. Seiringer, *Bosonization of fermionic many-body dynamics*, In Annales Henri Poincaré, pp. 1–40. Springer International Publishing, (2021).
- [49] N. Benedikter, M. Porta, B. Schlein, *Mean-field evolution of fermionic systems*. Communications in Mathematical Physics, **331**(3), pp.1087–1131 (2014).
- [50] N. Benedikter, M. Porta, B. Schlein, *Effective evolution equations from quantum dynamics*, (Vol. **7**). Berlin: Springer (2016).
- [51] N. Benedikter, J. Sok, J.P. Solovej, *The Dirac-Frenkel principle for reduced density matrices, and the Bogoliubov-de Gennes equations*, Annales Henri Poincaré Vol. **19**, No. 4, pp. 1167–1214 (2018).
- [52] M.J. Bijlsma, E. Zaremba, H.T.C. Stoof, *Condensate growth in trapped bose gases*, Physical Review A, **62**(6):063609 (2000).
- [53] A.V. Bobylev *Exact solutions of the nonlinear Boltzmann equation and the theory of relaxation of a Maxwell gas*, Teoret. Mat. Fiz., **60**(2):280–310 (1984).
- [54] A.V. Bobylev, I.M. Gamba, V. A. Panferov, *Moment inequalities and high-energy tails for Boltzmann equations with inelastic interactions*, J. Statist. Phys., **116**(5–6):1651–1682 (2004).

- [55] C. Boccato, *The excitation spectrum of the Bose gas in the Gross-Pitaevskii regime*, Reviews in Mathematical Physics, **33**(01), p.2060006 (2021).
- [56] C. Boccato, C. Brennecke, S. Cenatiempo, B. Schlein, *Complete Bose-Einstein condensation in the Gross-Pitaevskii regime*, Communications in Mathematical Physics, **359**(3), 975–1026 (2018).
- [57] C. Boccato, S. Cenatiempo, B. Schlein, *quantum many-body fluctuations around nonlinear Schrödinger dynamics*, Annales Henri Poincaré Vol. **18**, No. 1, pp. 113–191, Springer International Publishing (2017).
- [58] T. Bodineau, I. Gallagher, L. Saint-Raymond, *From hard sphere dynamics to the Stokes-Fourier equations: An analysis of the Boltzmann-Grad limit*, Annals of PDE, **3**(1), p.2 (2017).
- [59] T. Bodineau, I. Gallagher, L. Saint-Raymond, S. Simonella, *Fluctuation theory in the Boltzmann-Grad limit*, Journal of Statistical Physics, **180**(1), pp.873–895 (2020).
- [60] T. Bodineau, I. Gallagher, L. Saint-Raymond, S. Simonella, *Long-time correlations for a hard-sphere gas at equilibrium*, arXiv preprint arXiv:2012.03813 (2020), to appear in Communications in Pure and Applied Mathematics (2021).
- [61] T. Bodineau, I. Gallagher, L. Saint-Raymond, S. Simonella, *Long-time derivation at equilibrium of the fluctuating Boltzmann equation*, arXiv preprint arXiv:2201.04514 (2022).
- [62] N.N. Bogoliubov, *On the theory of superfluidity*, Izv. Akad. Nauk USSR, **11**, 77 (1947). Eng. Trans. J. Phys. (USSR), **11**, 23 (1947). See also *Lectures on quantum statistics*, vol. 1, Gordon and Breach (1967)
- [63] N.N. Bogoliubov, *Problems of dynamic theory in statistical physics; problemy dinamicheskoi teorii v statisticheskoi fiziki* (1946), in Studies in Statistical Mechanics, Vol. 1, J. de Boer and G. E. Uhlenbeck, eds. (North Holland, Amsterdam, 1962) .
- [64] S.N. Bose, *Plancks Gesetz und Lichtquantenhypothese*, Zeitschrift für Physik. **26**: 178–181 (1924).
- [65] L. Boßmann, *Derivation of the 1d nonlinear Schrödinger equation from the 3d quantum many-body dynamics of strongly confined bosons*, Journal of Mathematical Physics, **60**(3), p.031902 (2019).
- [66] L. Boßmann, N. Pavlović, P. Pickl, A. Soffer, *Higher order corrections to the mean-field description of the dynamics of interacting bosons*, Journal of Statistical Physics **178**(6), pp. 1362–1396 (2020).
- [67] L. Boßmann, S. Petrat, P. Pickl, A. Soffer, *Beyond Bogoliubov Dynamics*, Pure and Applied Analysis, **3**(4), pp. 677–726. (2022).
- [68] L. Boßmann, S. Petrat, R. Seiringer, *Asymptotic expansion of the low-energy excitation spectrum for weakly interacting bosons*, Forum of Mathematics, Sigma. **9**, e28 (2021).
- [69] L. Boßmann, S. Teufel, *Derivation of the 1d Gross-Pitaevskii equation from the 3d quantum many-body dynamics of strongly confined bosons*, In Annales Henri Poincaré Vol. **20**, No. 3, pp. 1003–1049. Springer International Publishing (2019).
- [70] O. Bratteli, D. W. Robinson, *Operator Algebras and quantum Statistical Mechanics 2. Equilibrium States. Models in quantum Statistical Mechanics*, Text and Monographs in Physics. Springer-Verlag, Berlin, 2 edition, (1996).
- [71] C. Brennecke, M. Caporaletti, B. Schlein, *Excitation Spectrum for Bose Gases beyond the Gross-Pitaevskii Regime*, arXiv preprint arXiv:2104.13003 (2021).
- [72] C. Brennecke, B. Schlein, S. Schraven, *Bose-Einstein Condensation with Optimal Rate for Trapped Bosons in the Gross-Pitaevskii Regime*, arXiv preprint arXiv:2102.11052 (2021)
- [73] M. Briant, A. Einav, *On the Cauchy problem for the homogeneous Boltzmann-Nordheim equation for bosons: local existence, uniqueness and creation of moments*, Journal of Statistical Physics, **163**(5), pp.1108–1156 (2016).
- [74] S. Buchholz, C. Saffirio, B. Schlein, *Multivariate central limit theorem in quantum dynamics*, Journal of Statistical Physics, **154**(1), pp.113–152 (2014).
- [75] T. Buckmaster, P. Germain, Z. Hani, J. Shatah, *Onset of the wave turbulence description of the longtime behavior of the nonlinear Schrödinger equation*, Inventiones mathematicae **225**(3), pp. 787–855 (2021).
- [76] S. Cai, *On the spatially homogeneous Boltzmann equation for Bose-Einstein particles with balanced potentials*, arXiv preprint arXiv:2101.00144 (2021).

- [77] S. Cai, X. Lu, *The spatially homogeneous Boltzmann equation for Bose-Einstein particles: rate of strong convergence to equilibrium*, (English summary) *J. Stat. Phys.* **175**, no. 2, 289–350 (2019).
- [78] S. Cenatiempo, *Bogoliubov theory for dilute Bose gases: The Gross-Pitaevskii regime*, *Journal of Mathematical Physics* **60**(8), p.081901 (2019).
- [79] C. Cercignani, *On the Boltzmann equation for rigid spheres*, *Transport Theory Statist. Phys.* **2**, no. 3, p. 211–225 (1972).
- [80] C. Cercignani, V. I. Gerasimenko, D. Y. Petrina, *Many-particle dynamics and kinetic equations*, Vol. **420**, Springer Science & Business Media (1997).
- [81] C. Cercignani, R. Illner, M. Pulvirenti, *The Mathematical Theory of Dilute Gases*, Springer, Berlin (1994)
- [82] T. Chen, *Localization lengths and Boltzmann limit for the Anderson model at small disorder in dimension 3*, *J. Stat. Phys.* **120**(1), pp. 279–337 (2005).
- [83] T. Chen, R. Denlinger, N. and Pavlović, *Local well-posedness for Boltzmann’s equation and the Boltzmann hierarchy via Wigner transform*, *Communications in Mathematical Physics* **368**(1), pp. 427–465 (2019).
- [84] T. Chen, R. Denlinger, N. and Pavlović, *Small data global well-posedness for a Boltzmann equation via bilinear spacetime estimates*, *Archive for Rational Mechanics and Analysis*, **240**(1), pp.327–381 (2021).
- [85] T. Chen, C. Hainzl, N. Pavlović, R. Seiringer, *Unconditional uniqueness for the cubic Gross-Pitaevskii hierarchy via quantum de Finetti*, *Commun. Pure Appl. Math.*, **68** (10), 1845–1884, 2015.
- [86] T. Chen, Y. Hong, N. Pavlović, *On the scattering problem for infinitely many fermions in dimensions $d \geq 3$ at positive temperature*, *Ann. Inst. H. Poincaré Anal. Non Linéaire* **35**, no. 2, pp. 393–416 (2018).
- [87] T. Chen, Y. Hong, N. Pavlović, *Global well-posedness of the NLS system for infinitely many fermions*, *Arch. Ration. Mech. Anal.* **224**, no. 1, pp. 91–123 (2017).
- [88] T. Chen, N. Pavlović, *On the Cauchy problem for focusing and defocusing Gross-Pitaevskii hierarchies*, *Discr. Contin. Dyn. Syst.*, **27** (2), 715–739 (2010).
- [89] T. Chen, N. Pavlović, *Derivation of the cubic NLS and Gross-Pitaevskii hierarchy from manybody dynamics in $d=3$ based on spacetime norms*, *Ann. H. Poincaré*, vol. **15**, no. 3, pp. 543–588. Springer Basel (2014).
- [90] T. Chen, I. Rodnianski, *Boltzmann limit for a homogeneous Fermi gas with dynamical Hartree-Fock interactions in a random medium*, *J. Stat. Phys.* **142**, no. 5, p. 1000–1051 (2011).
- [91] T. Chen, I. Sasaki, *Boltzmann limit and quasifreeness for a homogenous Fermi gas in a weakly disordered random medium*, *Journal of Statistical Physics*, **132**(2), pp. 329–353 (2008).
- [92] T. Chen, A. Soffer, *Mean field dynamics of a quantum tracer particle interacting with a boson gas*, *J. Funct. Anal.*, **276** (3), 971–1006 (2019).
- [93] T. Chen, K. Taliaferro, *Derivation in strong topology and global well-posedness of solutions to the Gross-Pitaevskii hierarchy*, *Communications in Partial Differential Equations*, **39**(9), pp.1658–1693 (2014).
- [94] X. Chen, Y. Guo *On the weak coupling limit of quantum many-body dynamics and the quantum Boltzmann equation*. *Kinetic & related models* 8.3: 443–465 (2015).
- [95] X. Chen, J. Holmer, *On the rigorous derivation of the 2D cubic nonlinear Schrödinger equation from 3D quantum many-body dynamics*. *Archive for Rational Mechanics and Analysis*, **210**(3), pp. 909–954 (2013).
- [96] X. Chen, J. Holmer, *On the Klainerman-Machedon conjecture for the quantum BBGKY hierarchy with self-interaction*, *J. Eur. Math. Soc. (JEMS)* **18** (6), 1161–1200 (2016).
- [97] X. Chen, J. Holmer, *Correlation structures, many-body scattering processes, and the derivation of the Gross-Pitaevskii hierarchy*, *Int. Math. Res. Not. IMRN* **2016**: **10**, 3051–3110 (2016).
- [98] X. Chen, J. Holmer, *Focusing quantum many-body dynamics: the rigorous derivation of the 1D focusing cubic nonlinear Schrödinger equation*. *Archive for Rational Mechanics and Analysis*, **221**(2), pp. 631–676 (2016).
- [99] X. Chen, J. Holmer, *The rigorous derivation of the 2D cubic focusing NLS from quantum many-body evolution*, *International Mathematics Research Notices*, **2017**(14), pp. 4173–4216, (2017).

- [100] X. Chen, J. Holmer, *The derivation of the \mathbb{T}^3 energy-critical NLS from quantum many-body dynamics*. *Inventiones mathematicae*, **217**(2), 433–547 (2019).
- [101] J.J.W. Chong, *Uniform in N global well-posedness of the time-dependent Hartree-Fock-Bogoliubov equations in \mathbb{R}^{1+1}* , *Letters in Mathematical Physics*, **108**(10), pp.2255–2283 (2018).
- [102] J.J.W. Chong, *Dynamics of large boson systems with attractive interaction and a derivation of the cubic focusing NLS equation in \mathbb{R}^3* , *Journal of Mathematical Physics* **62**, no. 4, 042106 (2021).
- [103] J.J.W. Chong, M.G. Grillakis, M. Machedon, Z. Zhao, *Global estimates for the Hartree-Fock-Bogoliubov equations*, *Communications in Partial Differential Equations*, pp.1–50 (2021).
- [104] J.J.W. Chong, L. Lafleche, C. Saffirio, *From many-body quantum dynamics to the Hartree-Fock and Vlasov equations with singular potentials*, arXiv preprint arXiv:2103.10946 (2021).
- [105] J.J.W. Chong, Z. Zhao, *Dynamical Hartree-Fock-Bogoliubov approximation of interacting bosons*, In *Annales Henri Poincaré* (pp. 1–59). Springer International Publishing (2021).
- [106] M.R. Christiansen, C. Hainzl, P.T. Nam, *The Random Phase Approximation for Interacting Fermi Gases in the Mean-Field Regime*, Preprint arXiv:2106.11161 (2021).
- [107] C. Collot, P. Germain, *On the derivation of the homogeneous kinetic wave equation*, Preprint arXiv:1912.10368 (2019).
- [108] C. Collot, P. Germain, *Derivation of the homogeneous kinetic wave equation: longer time scales*, Preprint, arXiv:2007.03508 (2020).
- [109] E. Cortés, M. Escobedo, *On a system of equations for the normal fluid-condensate interaction in a Bose gas*, *J. Func. An.*, **278**(2), p.108315 (2020).
- [110] G. Craciun, M.B. Tran, *A reaction network approach to the convergence to equilibrium of quantum Boltzmann equations for Bose gases*, *ESAIM: Control, Optimisation and Calculus of Variations* **27**: 83 (2021).
- [111] K.B. Davis, M.O. Mewes, M.R. Andrews, N.J. Van Druten, D.S. Durfee, D.M. Kurn, W. Ketterle, *Bose-Einstein condensation in a gas of sodium atoms*, *Physical review letters*, **75**(22), 3969 (1995).
- [112] D.A. Deckert, J. Fröhlich, P. Pickl, A. Pizzo, *Dynamics of sound waves in an interacting Bose gas*, *Advances in Mathematics*, **293**, pp.275–323 (2016).
- [113] Y. Deng, Z. Hani, *On the derivation of the wave kinetic equation for NLS*, In *Forum of Mathematics, Pi* (Vol. 9). Cambridge University Press (2021).
- [114] Y. Deng, Z. Hani, *Full derivation of the wave kinetic equation*, Preprint arXiv:2104.11204 (2021).
- [115] R. Denlinger, *The propagation of chaos for a rarefied gas of hard spheres in the whole space*, *Archive for Rational Mechanics and Analysis*, **229**(2), pp.885–952 (2018).
- [116] E. de la Canal, I.M. Gamba, M. Pavić-Čolić, *Propagation of L^p_β -norm, $1 < p \leq \infty$, for the system of Boltzmann equations for monatomic gas mixtures*, arXiv preprint arXiv:2001.09204 (2020).
- [117] P.J.H. Denteneer, *Second Quantization. Lecture notes with course quantum Theory [lecture notes] (Fall 2008)*, retrieved from <http://wwwhome.lorentz.leidenuniv.nl/~pjhdent/SecQuant08.pdf>
- [118] L. Desvillettes, F. Golse, *On a model Boltzmann equation without angular cutoff*, *Differential Integral Equations* **13**(4–6):567–594, (2000).
- [119] L. Desvillettes, B. Wennberg, *Smoothness of the solution of the spatially homogeneous Boltzmann equation without cutoff*, *Comm. Partial Differential Equations* **29**(1–2):133–155, (2004).
- [120] A. Deuchert, R. Seiringer, *Gross-Pitaevskii limit of a homogeneous Bose gas at positive temperature*, *Archive for Rational Mechanics and Analysis* **236**(3), pp.1217–1271, (2020).
- [121] A. Deuchert, R. Seiringer, *Semiclassical approximation and critical temperature shift for weakly interacting trapped bosons*, *Journal of Functional Analysis* **281**(6), p. 109096 (2021).
- [122] A. Deuchert, R. Seiringer, J. Yngvason, *Bose-Einstein condensation in a dilute, trapped gas at positive temperature*, *Communications in Mathematical Physics* **368**, no. 2, pp. 723–776 (2019).
- [123] R.J. DiPerna, P.-L. Lions, *On the Cauchy problem for Boltzmann equations: global existence and weak stability*, *Ann. Math.* **130**(2), 321–366 (1989).
- [124] J. Dolbeault, *Kinetic models and quantum effects: a modified Boltzmann equation for Fermi-Dirac particles*, *Arch. Ration. Mech. Anal.* **127**, pp. 101–131 (1994) .

- [125] R. Duan, *On the Cauchy problem for the Boltzmann equation in the whole space: global existence and uniform stability in $L^2_\xi(H^N_x)$* , J. Differ. Equ. **244**(12), pp. 3204–3234 (2008)
- [126] U. Eckern, *Relaxation processes in a condensed bose gas*, J. Low Temp. Phys. **54**, pp. 333–359, (1984).
- [127] A. Einstein, *Quantentheorie des einatomigen idealen Gases*, Sitzungsberichte der Preussischen Akademie der Wissenschaften. **1**: 3 (1925).
- [128] M.B. Erdoğan, N. Tzirakis, *Talbot effect for the cubic non-linear Schrödinger equation on the torus*. (English summary) Math. Res. Lett. **20** (2013), no. 6, 1081–1090.
- [129] L. Erdős, M. Salmhofer, H.-T. Yau, *On the quantum Boltzmann equation*. Journal of statistical physics, 116(1–4), 367–380 (2004).
- [130] L. Erdős, B. Schlein, H.-T. Yau, *Derivation of the Gross-Pitaevskii hierarchy for the dynamics of Bose-Einstein condensate*, Comm. Pure Appl. Math. **59** (12), 1659–1741 (2006).
- [131] L. Erdős, B. Schlein, H.-T. Yau, *Derivation of the cubic non-linear Schrödinger equation from quantum dynamics of many-body systems*, Invent. Math. **167**, 515–614 (2007).
- [132] L. Erdős, B. Schlein, H.-T. Yau, *Rigorous derivation of the Gross-Pitaevskii equation with a large interaction potential*. J. Amer. Math. Soc. **22** (4), 1099–1156 (2009).
- [133] L. Erdős, B. Schlein, H.-T. Yau *Derivation of the Gross-Pitaevskii equation for the dynamics of Bose-Einstein condensates*. Ann. of Math. (2) **172** (1), 291–370 (2010).
- [134] L. Erdős, H.-T. Yau, *Linear Boltzmann equation as the weak coupling limit of a random Schrödinger equation*, Comm. Pure Appl. Math. **53**(6), pp. 667–753, (2000).
- [135] L. Erdős, H.-T. Yau, *Derivation of the nonlinear Schrödinger equation from a many body Coulomb system*. Adv. Theor. Math. Phys. **5**, no. 6: pp. 1169–1205 (2001).
- [136] M. Escobedo, S. Mischler, M. Valle, *Homogeneous Boltzmann equation in quantum relativistic kinetic theory*, Department of Mathematics, Texas State University-San Marcos (2003).
- [137] M. Escobedo, S. Mischler, M. Valle, *Entropy maximisation problem for quantum relativistic particles*, Bull. Soc. Math. Fr. **133**, pp. 87–120 (2005).
- [138] M. Escobedo, S. Mischler, J.J.L. Velázquez, *Singular solutions for the Uehling-Uhlenbeck equation*, Proc. Roy. Soc. Edinburgh **138A**, pp. 67–107 (2008).
- [139] M. Escobedo, F. Pezzotti, and M. Valle, *Analytical approach to relaxation dynamics of condensed Bose gases*, Ann. Physics, **326** (4), pp. 808–827 (2011).
- [140] M. Escobedo, J.J.L. Velázquez, *A derivation of a new set of equations at the onset of the Bose-Einstein condensation*, Journal of Physics A: Mathematical and Theoretical, **41**(39), p.395208 (2008).
- [141] M. Escobedo, J.J.L. Velázquez, *On the blow up and condensation of supercritical solutions of the Nordheim equation for bosons*, Comm. Math. Phys. **330**, no. 1, 331–365 (2014).
- [142] M. Escobedo, J.J.L. Velázquez, *Finite time blow-up and condensation for the bosonic Nordheim equation*, Invent. Math., **200**, pp. 761–847 (2015).
- [143] M. Escobedo, J.J.L. Velázquez, *On the theory of weak turbulence for the nonlinear Schrödinger equation*, Mem. Amer. Math. Soc. **238** (2015), no. 1124, v+107 pp.
- [144] J. Fröhlich, S. Graffi, S. Schwarz, *Mean-field- and classical limit of many-body Schrödinger dynamics for bosons*, Comm. Math. Phys. **271**, no. 3, 681–697 (2007).
- [145] J. Fröhlich, A. Knowles, A. Pizzo, *Atomism and quantization*, J. Phys. A **40**, no. 12, 3033–3045 (2007).
- [146] J. Fröhlich, A. Knowles, S. Schwarz *On the Mean-Field Limit of Bosons with Coulomb Two-Body Interaction*, Comm. Math. Phys. **288** (3), 1023–1059 (2009).
- [147] J. Fröhlich, T.-P. Tsai, H.-T. Yau, *On a classical limit of quantum theory and the non-linear Hartree equation*, GAFA 2000 (Tel Aviv, 1999). Geom. Funct. Anal., Special Volume, Part I, 57–78 (2000).
- [148] M.L. Fürst, J. Lukkarinen, P. Mei, H. Spohn, *Derivation of a matrix-valued Boltzmann equation for the Hubbard model*, Journal of Physics A: Mathematical and Theoretical, **46**(48), p.485002 (2013).
- [149] I. Gallagher, L. Saint-Raymond, B. Texier, *From Newton to Boltzmann: hard spheres and short-range potentials*, European Mathematical Society (2013).
- [150] I.M. Gamba, M. Pavić-Čolić, *On existence and uniqueness to homogeneous Boltzmann flows of monatomic gas mixtures*, Archive for Rational Mechanics and Analysis, **235**(1), pp.723–781 (2020).
- [151] C.W. Gardiner, P. Zoller, R.J. Ballagh, M.J. Davis, *Kinetics of Bose-Einstein condensation in a trap*, Physical review letters, **79**(10):1793 (1997).

- [152] V.I. Gerasimenko *Heisenberg picture of quantum kinetic evolution in mean-field limit*, Kinetic & Related Models 4, no. 1: 385 (2011).
- [153] V.I. Gerasimenko, *Hierarchies of quantum evolution equations and dynamics of many-particle correlations*, International Journal of Evolution Equations, **7**(2), p.109 (2012).
- [154] V.I. Gerasimenko, *Kinetic equations and hierarchies of evolution equations of quantum systems*, arXiv preprint arXiv:2107.10872 (2021).
- [155] V.I. Gerasimenko, Z.A. Tsvir, *A description of the evolution of quantum states by means of the kinetic equation*, Journal of Physics A: Mathematical and theoretical **43**(48), p.485203 (2010).
- [156] P. Germain, A.D. Ionescu, M.B. Tran, *Optimal local well-posedness theory for the kinetic wave equation*, Journal of Functional Analysis, **279**(4), p.108570 (2020).
- [157] J. Ginibre, G. Velo, *The classical field limit of scattering theory for non-relativistic many-boson systems. I+II*, Communications in Mathematical Physics **66.1**: pp. 37–76 (1979), and **68**: pp. 45–68 (1979).
- [158] R. T. Glassey, *Global solutions to the Cauchy problem for the relativistic Boltzmann equation with near-vacuum data*, Comm. Math. Phys. **264** (2006), no. 3, 705724.
- [159] F. Golse, *On the dynamics of large particle systems in the mean field limit*, In Macroscopic and large scale phenomena: coarse graining, mean field limits and ergodicity (pp. 1–144). Springer, Cham (2016).
- [160] T. Goudon *On Boltzmann equations and Fokker-Planck asymptotics: influence of grazing collisions*, J. Statist. Phys., **89**(3–4):751–776, 1997.
- [161] H. Grad, *Principles of the kinetic theory of gases*, Handbuch der Physik **12**, Thermodynamik der Gase pp. 205–294 Springer-Verlag, Berlin-Göttingen-Heidelberg (1958).
- [162] P. Gressman, V. Sohinger, G. Staffilani, *On the uniqueness of solutions to the periodic 3D Gross-Pitaevskii hierarchy*, J. Funct. Anal. **266** (7), 4705–4764 (2014).
- [163] P.T. Gressman, R.M. Strain, *Global classical solutions of the Boltzmann equation without angular cutoff*, J. Am. Math. Soc. **24**(3), 771–847 (2011)
- [164] A. Griffin, T. Nikuni, E. Zaremba, *Bose-condensed gases at finite temperatures*, Cambridge University Press, Cambridge (2009).
- [165] J. Griffin, *Derivation of the linear Boltzmann equation from the damped quantum Lorentz gas with a general scatterer configuration*, arXiv preprint: arXiv:2201.08229 (2022).
- [166] M. Grillakis, M. Machedon, *Beyond mean field: On the role of pair excitations in the evolution of condensates*, Journal of Fixed Point Theory and Applications **14**, no. 1, pp. 91–111 (2013).
- [167] M. Grillakis, M. Machedon, *Pair excitations and the mean field approximation of interacting bosons, I*. Communications in Mathematical Physics **324.2**, pp. 601–636 (2013).
- [168] M. Grillakis, M. Machedon, *Pair excitations and the mean field approximation of interacting Bosons, II*. Communications in Partial Differential Equations **42.1**: pp. 24–67 (2017).
- [169] M. Grillakis, M. Machedon, A. Margetis, *Second-order corrections to mean field evolution for weakly interacting Bosons. I*, Comm. Math. Phys. **294** (1), 273–301 (2010).
- [170] Y. Guo, *Classical solutions to the Boltzmann equation for molecules with an angular cutoff*, Arch. Ration. Mech. Anal. **169**(4), 305–353 (2003).
- [171] S.J. Gustafson, I.M. Sigal, *Mathematical concepts of quantum mechanics*, Springer Science & Business Media (2011).
- [172] S.-Y. Ha, S.-E. Noh, *Global existence and stability of mild solutions to the inelastic Boltzmann system for gas mixtures*, Quart. Appl. Math. **68** (2010), no. 4, 671699.
- [173] S.-Y. Ha, S.-E. Noh, S.-B. Yun, *Global existence and stability of mild solutions to the Boltzmann system for gas mixtures*, Quart. Appl. Math. **65** (2007), no. 4, 757779.
- [174] C. Hainzl, *Another proof of BEC in the GP-limit*, Journal of Mathematical Physics **62**(5), p.051901 (2021).
- [175] L.B. He, X. Lu, M. Pulvirenti, *On semi-classical limit of spatially homogeneous quantum Boltzmann equation: weak convergence*, Communications in Mathematical Physics, **386**(1), pp. 143–223 (2021).
- [176] K. Hepp, *The classical limit for quantum mechanical correlation functions*, Comm. Math. Phys. **35**, pp. 265–277 (1974).
- [177] S. Herr, V. Sohinger, *The Gross-Pitaevskii hierarchy on general rectangular tori*, Archive for Rational Mechanics and Analysis **220**(3), pp.1119–1158 (2016).

- [178] P.D. Hislop, K. Kirkpatrick, S. Olla, J. Schenker, *Transport of a quantum particle in a time-dependent white-noise potential*, J. Math. Phys. **60**:8: 083303 (2019).
- [179] T.G. Ho, L.J. Landau, *Fermi gas on a lattice in the van Hove limit*, J. Statist. Phys. **87**, pp. 821–845 (1997).
- [180] T.G. Ho, L.J. Landau, A.J. Wilkins, *On the weak coupling limit for a Fermi gas in a random potential*, Rev. Math. Phys. **5**, pp. 209–298 (1992).
- [181] A. Hohenegger, A. Kartavtsev, M. Lindner, *Deriving Boltzmann equations from Kadanoff-Baym equations in curved space-time*, Physical Review D, **78**(8), p.085027 (2008).
- [182] Y. Hong, K. Taliaferro, Z. Xie, *Unconditional Uniqueness of the cubic Gross-Pitaevskii Hierarchy with Low Regularity*, SIAM Journal on Mathematical Analysis, **47**(5), pp.3314–3341 (2015).
- [183] M. Hott, *Convergence rate towards the fractional Hartree-equation with singular potentials in higher Sobolev norms*, Reviews in Mathematical Physics, Vol. **33**, No. 09, 2150029 (2021).
- [184] N.M. Hugenholtz, *Derivation of the Boltzmann equation for a Fermi gas*, J. Stat. Phys. **32**, pp. 231–254 (1983).
- [185] R. Illner, M. Shinbrot, *The Boltzmann Equation: Global Existence for a Rare Gas in an Infinite Vacuum*, Commun. Math. Phys. **95**, 217–226 (1984).
- [186] M. Jeblick, N. Leopold, P. Pickl, *Derivation of the time dependent Gross-Pitaevskii equation in two dimensions*, Communications in Mathematical Physics, **372**(1), pp.1–69 (2019).
- [187] C. Josserand and Y. Pomeau, *Nonlinear aspects of the theory of Bose-Einstein condensates*, Nonlinearity **14**, R25-R62 (2001).
- [188] L. P. Kadanoff, G. Baym, *quantum statistical mechanics. Green's function methods in equilibrium and nonequilibrium problems*, W. A. Benjamin, Inc., New York xi+203 (1962).
- [189] S. Kaniel, M. Shinbrot, *The Boltzmann Equation: Uniqueness and Local Existence*, Commun. math. Phys. **58**, 65–84 (1978).
- [190] L. Kapitanski, I. Rodnianski, *Does a quantum particle know the time?* Emerging applications of number theory (Minneapolis, MN, 1996), 355–371, IMA Vol. Math. Appl., **109**, Springer, New York, 1999.
- [191] A.H.M. Kierkels, J.J.L. Velázquez, *On the transfer of energy towards infinity in the theory of weak turbulence for the nonlinear Schrödinger equation*, J. Stat. Phys. **159**, no. 3, pp. 668–712 (2015).
- [192] M.K.H. Kiessling, *The Hartree limit of Born's ensemble for the ground state of a bosonic atom or ion*, Journal of mathematical physics, 53(9), p.095223 (2012).
- [193] F. King, *BBGKY hierarchy for positive potentials*, Ph.D. Thesis (1975)
- [194] K. Kirkpatrick, B. Schlein, G. Staffilani, *Derivation of the two dimensional nonlinear Schrödinger equation from many body quantum dynamics*, Amer. J. Math. **133** (1), 91–130 (2011).
- [195] T. R. Kirkpatrick, J. R. Dorfman, *Transport theory for a weakly interacting condensed Bose gas*, Phys. Rev. A (3), **28**(4), pp. 2576–2579 (1983).
- [196] T. R. Kirkpatrick, J. R. Dorfman, *Transport in a dilute but condensed nonideal bose gas: Kinetic equations*, J. Low Temp. Phys., **58**, pp. 301–331, (1985).
- [197] S. Klainerman, M. Machedon, *On the uniqueness of solutions to the Gross-Pitaevskii hierarchy*, Commun. Math. Phys. **279**, no. 1, 169–185 (2008).
- [198] A. Knowles, P. Pickl *mean-field Dynamics: Singular Potentials and Rate of Convergence*, Communications in Mathematical Physics. Volume **298**, Issue 1: 101–138 (2010).
- [199] E. Kuz, *Rate of convergence to mean field for interacting bosons*, Communications in Partial Differential Equations **40.10**: 1831–1854 (2015).
- [200] E. Kuz, *Exact evolution versus mean field with second-order correction for Bosons interacting via short-range two-body potential*, Differential and Integral Equations **30**, no. 7/8, 587–630 (2017).
- [201] R. Lacaze, P. Lallemand, Y. Pomeau, S. Rica, *Dynamical formation of a Bose-Einstein condensate*, Physica D 152-153, pp. 779–786 (2001).
- [202] O.E. Lanford, *On a derivation of the Boltzmann equation*, Astérisque, **40**, pp.117–137 (1976).
- [203] J. Lee, *On the time dependence of the rate of convergence towards Hartree dynamics for interacting Bosons*, Journal of Statistical Physics, 176(2), 358–381, (2019).
- [204] J.O. Lee, *Rate of Convergence Towards Semi-Relativistic Hartree Dynamics*, Annales Henri Poincaré. Vol. **14**. No. 2. SP Birkhäuser Verlag Basel, (2013).

- [205] M. Lemou, *Linearized quantum and relativistic Fokker-Planck-Landau equations*, Math. Methods Appl. Sci. **23**, pp. 1093–1119 (2000) .
- [206] M. Lewin, *mean-field limit of Bose systems: rigorous results*, Proceedings of the International Congress of Mathematical Physics (2015).
- [207] M. Lewin, P.T. Nam, N. Rougerie, *Derivation of Hartree’s theory for generic mean-field Bose systems*, Adv. Math. **254**, 570–621 (2014).
- [208] M. Lewin, P.T. Nam, B. Schlein, *Fluctuations around Hartree states in the mean-field regime*, Amer. J. Math. **137** (6), 1613–1650 (2015).
- [209] M. Lewin, P.T. Nam, S. Serfaty, J.P. Solovej, *Bogoliubov spectrum of interacting Bose gases*, Communications on Pure and Applied Mathematics **68**, no. 3, 413–471 (2015).
- [210] M. Lewin, J. Sabin, *The Hartree equation for infinitely many particles I. Wellposedness theory*, Commun. Math. Phys. **334**(1), p. 117–170, (2015).
- [211] M. Lewin, J. Sabin, *The Hartree equation for infinitely many particles, II: Dispersion and scattering in 2D*, Anal. PDE. **7**(6), pp. 1339–1363 (2014).
- [212] W. Li, X. Lu, *Global existence of solutions of the Boltzmann equation for Bose-Einstein particles with anisotropic initial data*, Journal of Functional Analysis, **276**(1), pp.231–283 (2019).
- [213] Z. Li, *Existence and uniqueness of solutions to the quantum Boltzmann equation for soft potentials*, arXiv preprint arXiv:2112.10357 (2021).
- [214] Q. Liard *On the mean-field approximation of many-boson dynamics*, J. Func. An., **273**(4): 1397–1442 (2017).
- [215] E.H. Lieb, R. Seiringer, *Proof of Bose-Einstein condensation for dilute trapped gases*, Phys. Rev. Lett. **88**, 170409 (2002).
- [216] E.H. Lieb, R. Seiringer, J.P. Solovej, J. Yngvason, *The mathematics of the Bose gas and its condensation*, Birkhäuser (2005).
- [217] E.H. Lieb, R. Seiringer, J. Yngvason, *Bosons in a trap: A rigorous derivation of the Gross-Pitaevskii energy functional*, Phys. Rev. A **61**, 043602–1–13 (2000).
- [218] E.H. Lieb, R. Seiringer, J. Yngvason, *A rigorous derivation of the Gross-Pitaevskii energy functional for a two-dimensional Bose gas*, Comm. Math. Phys. **224** (2001).
- [219] P.L. Lions, *Compactness in Boltzmann’s equation via Fourier integral operators and applications, III*, J. Math. Kyoto Univ. **34**(3), pp. 539–584 (1994).
- [220] S. Liu, *Smoothing effects for the classical solutions to the Landau-Fermi-Dirac equation*, Chinese Annals of Mathematics, Series B, **33**(6), pp. 857–876 (2012).
- [221] X. Lu, *A modified Boltzmann equation for Bose-Einstein particles: isotropic solutions and long-time behavior*, J. Stat. Phys. **98**, pp. 1335–1394 (2000).
- [222] X. Lu, *On spatially homogeneous solutions of a modified Boltzmann equation for Fermi-Dirac particles*, J. Stat. Phys. **105**, no. 1-2, 353–388 (2001).
- [223] X. Lu, *On isotropic distributional solutions to the Boltzmann equation for Bose-Einstein particles*, J. Stat. Phys. **116**, pp. 1597–1649 (2004).
- [224] X. Lu, *The Boltzmann equation for Bose-Einstein particles: velocity concentration and convergence to equilibrium*, J. Stat. Phys. **119**, 1027–1067 (2005)
- [225] X. Lu, *On the Boltzmann equation for Fermi-Dirac particles with very soft potentials: averaging compactness of weak solutions*, J. Stat. Phys. **124**, pp. 517–547 (2006) .
- [226] X. Lu, *On the Boltzmann equation for Fermi-Dirac particles with very soft potentials: Global existence of weak solutions*, J. Diff. Eq., **245**(7), pp.1705–1761 (2008).
- [227] X. Lu, *The Boltzmann equation for Bose-Einstein particles: condensation in finite time*, Journal of Statistical Physics, **150**(6), pp.1138–1176 (2013).
- [228] X. Lu, *The Boltzmann equation for Bose-Einstein particles: regularity and condensation*, J. Stat. Phys., **156**(3), pp.493–545 (2014).
- [229] X. Lu, *Long time convergence of the Bose-Einstein condensation*, J. Stat. Phys. **162**, pp. 652–670 (2016).
- [230] X. Lu, C. Mouhot, *On measure solutions of the Boltzmann equation, part I: moment production and stability estimates*, J. Diff. Eq. **252**(4):3305–3363 (2012).
- [231] X. Lu, B. Wennberg, *On stability and strong convergence for the spatially homogeneous Boltzmann equation for Fermi-Dirac particles*, Archive for rational mechanics and analysis, **168**(1), pp.1–34 (2003).
- [232] X. Lu, X. Zhang, *On the Boltzmann equation for 2D Bose-Einstein particles*. J. Stat. Phys., **143**(5), pp.990–1019 (2011).

- [233] J. Lührmann, *mean-field quantum dynamics with magnetic fields*, Journal of Mathematical Physics **53**:2: 022105 (2012).
- [234] J. Lukkarinen, H. Spohn, *Not to normal order-notes on the kinetic limit for weakly interacting quantum fluid*, Journal of Statistical Physics, **134**(5), pp. 1133–1172 (2009).
- [235] J. Lukkarinen, H. Spohn, *Weakly nonlinear Schrödinger equation with random initial data*, Inventiones mathematicae, **183**(1), pp.79–188 (2011).
- [236] A. Michelangeli, A. Olgiami, *Mean-field quantum dynamics for a mixture of Bose-Einstein condensates*, Anal.Math.Phys. 1–40 (2016).
- [237] A. Michelangeli, B. Schlein, *Dynamical collapse of boson stars*, Communications in Mathematical Physics **311**:3: 645–687 (2012).
- [238] S. Mischler, B. Perthame, *Boltzmann equation with infinite energy: renormalized solutions and distributional solutions for small initial data and initial data close to a Maxwellian*, SIAM J. Math. Anal. **28**, no. 5, 10151027 (1997).
- [239] D. Mitrouskas, S. Petrat, P. Pickl, *Bogoliubov corrections and trace norm convergence for the Hartree dynamics*, Reviews in Mathematical Physics, **31**(08), 1950024 (2019).
- [240] P.T. Nam, M. Napiórkowski, *Bogoliubov correction to the mean-field dynamics of interacting bosons*, Advances in Theoretical and Mathematical Physics 21, no. 3, 683–738 (2017).
- [241] P.T. Nam, M. Napiórkowski, *A note on the validity of Bogoliubov correction to mean-field dynamics*, Journal de Mathématiques Pures et Appliquées, 108(5), 662–688 (2017).
- [242] P.T. Nam, M. Napiórkowski, *Norm approximation for many-body quantum dynamics: focusing case in low dimensions*, Advances in Mathematics, **350**, 547–587 (2019).
- [243] P.T. Nam, M. Napiórkowski, *Two-term expansion of the ground state one-body density matrix of a mean-field Bose gas*, Calculus of Variations and Partial Differential Equations, 60(3), pp.1–30. (2021).
- [244] P.T. Nam, M. Napiórkowski, J. Ricaud, A. Triay, *Optimal rate of condensation for trapped bosons in the Gross-Pitaevskii regime*, arXiv preprint arXiv:2001.04364 (2020).
- [245] P.T. Nam, N. Rougerie, R. Seiringer, *Ground states of large bosonic systems: the Gross-Pitaevskii limit revisited*, Analysis & PDE **9**(2), pp.459–485 (2016).
- [246] P.T. Nam, R. Salzmänn, *Derivation of 3D Energy-Critical Nonlinear Schrödinger Equation and Bogoliubov Excitations for Bose Gases*, Communications in Mathematical Physics, 1–77 (2019).
- [247] P.T. Nam, R. Seiringer, *Collective excitations of Bose gases in the mean-field regime*, Archive for Rational Mechanics and Analysis, **215**(2), pp.381–417 (2015).
- [248] H. Narnhofer, G. Sewell, *Vlasov hydrodynamics of a quantum mechanical model*, Commun. Math. Phys. **79**(1), pp. 9–24 (1981).
- [249] M. Napiórkowski, *Dynamics of interacting bosons: a compact review*, arXiv preprint arXiv:2101.04594, (2021).
- [250] T.T. Nguyen, M.B. Tran, *Uniform in time lower bound for solutions to a quantum Boltzmann equation of bosons*, Arch. Ration. Mech. Anal. **231**, pp. 63–89 (2019).
- [251] L.W. Nordheim, *On the kinetic method in the new statistics and its application in the electron theory of conductivity*, Proc. Roy. Soc. 119, 689–698 (1928)
- [252] Z. Ouyang, L. Wu, *On the quantum Boltzmann Equation near Maxwellian and Vacuum*, Journal of Differential Equations, **316**, pp.471–551 (2022).
- [253] T. Paul, M. Pulvirenti, S. Simonella, *On the size of chaos in the mean field dynamics*, Archive for Rational Mechanics and Analysis, **231**(1), pp.285–317 (2019).
- [254] S. Petrat, P. Pickl, A. Soffer, *Derivation of the Bogoliubov time evolution for gases with finite speed of sound*, In Annales Henri Poincaré **21**, pp. 461–498 (2020).
- [255] P. Pickl, *Derivation of the time dependent Gross-Pitaevskii equation without positivity condition on the interaction*, Journal of Statistical Physics, 140(1), 76–89, (2010).
- [256] P. Pickl, *A simple derivation of mean field limits for quantum systems*, Lett. Math. Phys., **97** (2), pp. 151–164 (2011).
- [257] Y. Pomeau, M. A. Brachet, S. Métens, S. Rica. *Théorie cinétique d’un gaz de bose dilué avec condensat*, C. R. Acad. Sci. Paris Sér. IIb Méc. Phys. Astr., **327**, pp. 791–798 (1999).
- [258] Y. Pomeau, M.B. Tran, *Statistical physics of non equilibrium quantum phenomena*, Berlin: Springer (2019).
- [259] Y. Pomeau, M.B. Tran, *Boltzmann-type collision operators for Bogoliubov excitations of Bose-Einstein condensates: A unified framework*, Physical Review E, **101**(3), p.032119 (2020).

- [260] Y. Pomeau, M.B. Tran, *On a thermal cloud–Bose–Einstein condensate coupling system*, The European Physical Journal Plus, **136**(5), pp.1–11 (2021).
- [261] M. Porta, S. Rademacher, C. Saffirio, B. Schlein, *Mean field evolution of fermions with Coulomb interaction*, Journal of Statistical Physics **166**(6), pp.1345–1364 (2017).
- [262] I. Prigogine, *Nonequilibrium Statistical Mechanics*, Wiley-Interscience, New York 1962.
- [263] N. Proukakis, S. Gardiner, M. Davis, M. Szymańska, *quantum Gases: Finite temperature and non-equilibrium dynamics*, volume 1. World Scientific (2013).
- [264] M. Pulvirenti, *The weak-coupling limit of large classical and quantum systems*, In International Congress of Mathematicians Vol. **3**, pp. 229–256 (2006)
- [265] M. Pulvirenti, S. Simonella, *The Boltzmann-Grad limit of a hard sphere system: analysis of the correlation error*, Invent. Math. **207**(3), 1135–1237 (2017)
- [266] M. Pulvirenti, C. Saffirio, S. Simonella, *On the validity of the Boltzmann equation for short range potentials*, Reviews in Mathematical Physics, **26.02**, p.1450001, (2014).
- [267] L.E. Reichl, *A modern course in statistical physics*, (1999).
- [268] L.E. Reichl, M.B. Tran, *A kinetic model for very low temperature dilute Bose gases*, Journal of Physics A: Mathematical and Theoretical, Volume **52**, Number 6, 063001 (2019).
- [269] I. Rodnianski, B. Schlein, *quantum fluctuations and rate of convergence towards mean field dynamics*, Comm. Math. Phys. **291** (1), 31–61 (2009).
- [270] M. Rosenzweig, G. Staffilani, *Uniqueness of solutions to the spectral hierarchy in kinetic wave turbulence theory*, Physica D: Nonlinear Phenomena, p.133148 (2022).
- [271] N. Rougerie, *De Finetti theorems, mean-field limits and Bose-Einstein condensation*, Lecture notes, arXiv preprint arXiv:1506.05263 (2015).
- [272] G. Royat, *Etude de l'équation d'Uehling-Uhlenbeck: existence de solutions proches de Planckiennes et étude numérique*, PhD, Marseille (2010).
- [273] B. Schlein: *Derivation of effective evolution equations from microscopic quantum dynamics*, In: Lecture Notes for the Minicourse held at the 2008 CMI Summer School in Zurich, arXiv preprint arXiv:0807.4307 (2008).
- [274] R. Seiringer, *The excitation spectrum for weakly interacting bosons*, Commun. Math. Phys., **306**, 565–578 (2011).
- [275] D.V. Semikoz and J.I. Tkachev, *Condensation of bosons in the kinetic regime*, Phys. Rev. D **55**, pp. 489–502 (1997).
- [276] S. Shen, *The rigorous derivation of the T^2 focusing cubic NLS from 3D*, Journal of Functional Analysis, **280**(8), p.108934 (2021).
- [277] X.L. Sheng, N. Weickgenannt, E. Speranza, D.H. Rischke, Q. Wang, *From Kadanoff-Baym to Boltzmann equations for massive spin-1/2 fermions*, Physical Review D **104**, no. 1: 016029 (2021).
- [278] S. Simonella, H. Spohn, *Book review: from Newton to Boltzmann: hard spheres and short-range potentials, by Isabelle Gallagher, Laure Saint-Raymond and Benjamin Texier*. Bull. Am. Math. Soc. **52**, pp.533–538 (2015)
- [279] A. Soffer, M.B. Tran, *On the dynamics of finite temperature trapped Bose gases*, Advances in Mathematics **325**, pp.533–607 (2018).
- [280] A. Soffer, M.B. Tran, *On coupling kinetic and Schrödinger equations*, Journal of Differential Equations **265**(5), pp.2243–2279 (2018).
- [281] V. Sohinger, *A microscopic derivation of Gibbs measures for nonlinear Schrödinger equations with unbounded interaction potentials*, arXiv preprint arXiv:1904.08137 (2019).
- [282] V. Sohinger, *Gibbs Measures of Nonlinear Schrödinger Equations as Limits of quantum many-body States in Dimension $d \leq 3$* , In Frontiers in Analysis and Probability, pp. 371–382, Springer, Cham (2020).
- [283] H. Spohn, *Derivation of the transport equation for electrons moving through random impurities*, J. Stat. Phys., **17**, pp. 385–412 (1977).
- [284] H. Spohn, *Kinetic Equations from Hamiltonian Dynamics*, Rev. Mod. Phys. **52**, no. 3, 569–615 (1980).
- [285] H. Spohn, *Boltzmann hierarchy and Boltzmann equation*, in Kinetic theories and the Boltzmann equation (Montecatini), pp. 207–220 (1981).
- [286] H. Spohn, H. Neunzert, *On the Vlasov Hierarchy*, Math. Meth. in the Appl. Sci., **3**, 445–455 (1981) .
- [287] H. Spohn, *Quantum kinetic equations*, in: On Three Levels, pp. 1–10. Springer, Boston, MA, (1994).

- [288] H. Spohn, *The phonon Boltzmann equation, properties and link to weakly anharmonic lattice dynamics*, J. Stat. Phys. **124**(2–4), pp. 1041–1104 (2006).
- [289] H. Spohn, *Kinetic equations for quantum many-particle systems*, in Encyclopedia of Mathematical Physics (Springer, Berlin) (2007).
- [290] H. Spohn, *Kinetics of the Bose-Einstein condensation*, Physica D, **239**, pp. 627–634 (2010).
- [291] M. Tasković, R.J. Alonso, I.M. Gamba, N. Pavlović, *On Mittag-Leffler moments for the Boltzmann equation for hard potentials without cutoff* SIAM Journal on Mathematical Analysis, **50**(1), pp.834–869 (2018).
- [292] G. Toscani, *On the nonlinear Boltzmann equation in unbounded domains*, Arch. Rational Mech. Anal. **95**, no. 1, 3749 (1986).
- [293] G. Toscani, *Global solution of the initial value problem for the Boltzmann equation near a local Maxwellian*, Arch. Rational Mech. Anal. **102**, no. 3, 231241 (1988).
- [294] L. N. Trefethen, J.A.C. Weideman, *The exponentially convergent trapezoidal rule*. SIAM rev., **56**(3), pp. 385–458, (2014).
- [295] F. Turcu, C. Bonchiş, M. Najim, *Vector partitions, multi-dimensional Faà di Bruno formulae and generating algorithms*. Discrete Applied Mathematics, **272**, 90–99, (2020).
- [296] K. Uchiyama, *Derivation of the Boltzmann equation from particle dynamics*, Hiroshima mathematical journal, **18**(2), pp.245–297 (1988).
- [297] E.A. Uehling, G.E. Uhlenbeck, *Transport phenomena in Einstein-Bose and Fermi-Dirac gases*, Phys. Rev. **43**, 552–561 (1933).
- [298] S. Ukai, *On the existence of global solutions of mixed problem for the non-linear Boltzmann equation*, Proc. Jpn. Acad. **50**(3), 179–184 (1974)
- [299] L. van Hove, *quantum-mechanical perturbations giving rise to a statistical transport equation*, Physica **21**, 517–540 (1955).
- [300] R. van Leeuwen, N.E. Dahlen, *An introduction to nonequilibrium green functions*, Lecture notes, unpublished (2005)
- [301] C. Villani *On a new class of weak solutions to the spatially homogeneous Boltzmann and Landau equations*, Arch. Rational Mech. Anal., **143**(3): pp. 273–307 (1998).
- [302] C. Villani *A review of mathematical topics in collisional kinetic theory*, In Handbook of mathematical fluid dynamics, Vol. I, pp. 71–305. North-Holland, Amsterdam, (2002).
- [303] B. Wennberg *On moments and uniqueness for solutions to the space homogeneous Boltzmann equation*, Transport Theory Statist. Phys. **23**(4):533–539, (1994).
- [304] E. Zaremba, T. Nikuni, A. Griffin, *Dynamics of trapped Bose gases at finite temperatures*, Journal of Low Temperature Physics, **116**(3), pp.277–345 (1999).
- [305] Y. Zhang, X. Lu, *Boltzmann equations with quantum effects. I. Long time behavior of spatial decay solutions*, Tsinghua Sci. Technol. **7** (2002) 215–218.
- [306] Y. Zhang and X. Lu, *Boltzmann equations with quantum effects. II. entropy identity, existence and uniqueness of spatial decay solutions*, Tsinghua Sci. Technol. **7**, no. 3, 219–222 (2002).

(T. Chen) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TEXAS AT AUSTIN, AUSTIN TX 78712, USA

Email address: tc@math.utexas.edu

(M. Hott) SCHOOL OF MATHEMATICS, UNIVERSITY OF MINNESOTA TWIN CITIES, MINNEAPOLIS MN 55455, USA

Email address: mhott@umn.edu