

GLOBAL SOLUTIONS TO AN INITIAL BOUNDARY PROBLEM FOR THE COMPRESSIBLE 3-D MHD EQUATIONS WITH NAVIER-SLIP AND PERFECTLY CONDUCTING BOUNDARY CONDITIONS IN EXTERIOR DOMAINS

HAIRONG LIU¹, TAO LUO², AND HUA ZHONG³

ABSTRACT. An initial boundary value problem for compressible Magnetohydrodynamics (MHD) is considered on an exterior domain (with the first Betti number vanishes) in $\mathbb{R}^3$ in this paper. The global existence of smooth solutions near a given constant state for compressible MHD with the boundary conditions of Navier-slip for the velocity field and perfect conduction for the magnetic field is established. Moreover the explicit decay rate is given. In particular, the results obtained in this paper also imply the global existence of classical solutions for the full compressible Navier-Stokes equations with Navier-slip boundary conditions on exterior domains in three dimensions, which is not available in literature, to the best of knowledge of the authors¹.

Keywords: Exterior domain, Navier-slip boundary conditions, Perfectly conducting condition, Global regularity near boundaries, Compressible full MHD.

AMS Subject Classifications. 76W05, 76N10

1. INTRODUCTION AND MAIN THEOREMS

Magnetohydrodynamics (MHD) mainly investigates the dynamics of compressible quasineutrally ionized fluids under the influence of electromagnetic fields. It is well-known that the applications of Magnetohydrodynamics cover a very wide range of physical objects, from liquid metals to cosmic plasmas. In this paper, the 3-dimensional compressible full magnetohydrodynamic equations will be considered, which take the following form ([17, 22]):

$$(1.1) \quad \begin{cases} \rho_t + \operatorname{div}(\rho u) = 0, \\ \rho(u_t + u \cdot \nabla u) - \mu \Delta u - (\mu + \lambda) \nabla \operatorname{div} u + \nabla p = \operatorname{curl} H \times H, \\ \partial_t(\rho e) + \operatorname{div}(\rho e u) + p \operatorname{div} u = \Psi : \nabla u + \operatorname{div}(\kappa \nabla \mathcal{T}) + \eta |\operatorname{curl} H|^2, \\ H_t - \eta \Delta H = \operatorname{curl}(u \times H), \quad \operatorname{div} H = 0, \end{cases}$$

where ρ , $u = (u_1, u_2, u_3)$, $H = (H_1, H_2, H_3)$, e , $\mathcal{T}$ denote the density, the velocity, the magnetic field, the internal energy and temperature, respectively. The viscosity coefficients μ and λ of the fluid should satisfy

$$2\mu + 3\lambda > 0, \quad \mu > 0,$$

due to physical realities. $\kappa > 0$ is the heat conductivity, and $\eta > 0$ is the magnetic diffusivity which describe a magnetic diffusion coefficient of the magnetic field. Ψ is the viscous stress tensor, given by

$$\Psi = \mu(\nabla u + \nabla u^T) + \lambda \operatorname{div} u I.$$

¹School of Science, Nanjing Forestry University, Nanjing 210037, P.R.China. E-mail : hrliu@njfu.edu.cn.

²Department of Mathematics, City University of Hong Kong, 83 Tat Chee Avenue, Kowloon Tong, Hong Kong. E-mail: taoluo@cityu.edu.hk.

³School of Mathematics, Southwest Jiaotong University, Chengdu 611756, P.R. China. E-mail: huazhong@swjtu.edu.cn.

and $\Psi : \nabla u$ denotes the scalar product of two matrices. A calculation gives that

$$\Psi : \nabla u = \lambda(\operatorname{div} u)^2 + \sum_{i,j=1}^3 \frac{\mu}{2} \left(\frac{\partial u^i}{\partial x_j} + \frac{\partial u^j}{\partial x_i} \right)^2 = \lambda(\operatorname{div} u)^2 + 2\mu|S(u)|^2.$$

For simplicity, we study the case of polytropic ideal gas type with the equations of state:

$$e = c_v \mathcal{T}, \quad p = R\rho \mathcal{T},$$

where c_v and R are positive constants. In this case, we may write (1.1)₃ as

$$c_v \rho (\mathcal{T}_t + u \cdot \nabla \mathcal{T}) + p \operatorname{div} u = \kappa \Delta \mathcal{T} + \Psi : \nabla u + \eta |\operatorname{curl} H|^2.$$

Let U be a simply connected bounded smooth domain in $\mathbb{R}^3$, and $\Omega \equiv \mathbb{R}^3 \setminus \bar{U}$ be the exterior domain. In this paper, we study the initial-boundary value problem of (1.1) in Ω with the initial condition

$$(\rho, u, \mathcal{T}, H)|_{t=0} = (\rho_0, u_0, \mathcal{T}_0, H_0), \quad (1.2)$$

and the boundary conditions:

$$u \cdot n = 0, \quad \operatorname{curl} u \times n = 0, \quad \text{on } x \in \partial\Omega, \quad (1.3)$$

$$H \cdot n = 0, \quad \operatorname{curl} H \times n = 0, \quad \frac{\partial \mathcal{T}}{\partial n} = 0, \quad \text{on } x \in \partial\Omega. \quad (1.4)$$

Moreover, besides the compatibility condition of the initial data with the boundary condition, we also assume that the initial data satisfy the compatibility conditions,

$$(\rho_0, u_0, \mathcal{T}_0, H_0)(x) \rightarrow (1, 0, 1, 0), \quad \text{as } x \rightarrow \infty.$$

The boundary condition (1.3) means the Navier-slip boundary conditions for the velocity field, (1.3) is the perfectly conducting boundary condition for the magnetic field, and Neumann boundary condition for temperature.

The Navier-slip boundary conditions for the velocity field were first introduced by Navier in [28] and have been used in many applications, which are usually used in the large eddy simulations of turbulent flows, to compute the large eddies of a turbulent flow accurately by neglecting small flow structure. For this, the slip boundary conditions are more suitable than the Dirichlet boundary conditions [14].

The system of compressible MHD equations have been studied extensively by physicists and mathematicians because of its physical importance, rich and complex phenomena and mathematical challenges. We mainly review the results on the global existence of solutions related to the main theme of this paper, for which one may refer to [7, 20, 32] for one-dimensional case and [13, 17, 18, 20, 23, 24, 29, 36] for higher dimensions for example. The results for the global existence and large time behavior of solutions in 3-D for the Cauchy problem, can be found, for instance, the classical solutions in [20, 23, 24, 29, 36], the variational weak solutions in [13, 17, 18].

In the presence of physical boundaries, there have been extensive studies on the global existence and large time behavior of solutions to the initial boundary value problems for compressible MHD equations in 3-D for the homogenous Dirichlet condition for both velocity field u and the magnetic field H , i.e., $u = 0$ and $H = 0$ on the boundary. It is more physically relevant to consider the case that the magnetic field is non-zero on the boundary. In this direction, the global existence of classical solutions for 3-D compressible isentropic MHD was proved in [4] very recently in a bounded domain (see also [12] for the related results in 2-D and [3] for the related results for compressible isentropic Navier-Stokes equations in 3-D). For the compressible Navier-Stokes equations in 3-D with Navier-slip boundary conditions, there are only few results on the global solutions for the initial boundary value problem for the Navier-slip boundary conditions in 3-D, though results are available for some small physical parameter limits for local time, for instance, see the related zero viscosity limit [2, 27, 33–35] and [8, 16] for low Mach number limit.

Compared with the problem for 3-D compressible isentropic MHD equations studied in [4] in a bounded domain, the difficulty in proving the global existence of classical solution for the initial boundary value problem of (1.1)-(1.4) in an exterior domain in $\mathbb{R}^3$ is due to the unboundedness of the exterior domain, for which the Poincare type inequality to use the L^2 -norm of the derivatives to control the L^2 -norm of the solution itself is not available. For example, in a bounded domain, the dissipation estimates (the $L^2(\Omega \times [0, \infty))$ space-time estimates) of the velocity and magnetic fields can be obtained via the corresponding estimates of the derivatives which follow from the dissipation of the viscosity and magnetic diffusion in the basic energy estimates. This is an important decay mechanism for the problem on a bounded domain for which various exponential decay estimates are obtained in [4]. However, for the problem on an exterior domain studied in the present paper, only algebraic decay can be expected in general. Another difference from the case studied in [4] is that we can handle the case of the full compressible MHD system with variable entropy, while only isentropic case is studied in [4] for which possibly large oscillations and vacuum states (with small energy) are allowed. It should be noted that, for non-isentropic flows, even in the case without magnetic fields ($H = 0$), there have been no global-in-time theory available for the problems with physical boundaries under the Navier-slip boundary conditions.

For the boundary conditions of Navier-slip for the velocity field and the perfect conduction of magnetic field, it is more suitable to use the L^2 -norms of div and $curl$ to control that of the derivatives of the velocity and magnetic fields. This control usually depends on the topology of the domain, which relates to the first Betti number. A key observation is that first Betti number of the exterior domain $\Omega = \mathbb{R}^3 \setminus \bar{U}$ vanishes when U is a bounded simply connected open set. In this case, one can apply the refined div-curl estimate (see Proposition 2.3) to obtain the dissipation estimate of $\|\nabla u\|$ with the help of the dissipation estimates for $\|curl u\|$ and $\|div u\|$. This observation is important to obtain the various dissipation estimates.

The basic approach to obtain the global existence and decay estimates of classical solutions of the initial boundary value problem is based on the energy estimates, for which the boundary conditions (1.3)-(1.4) make the problem rather interesting and challenging. In fact, unlike the homogeneous Dirichlet boundary conditions $u = H = 0$ on the boundary, various estimates on $div u$, $curl u$ and $curl H$ are crucial for the problem investigated in this paper.

The main results of this paper are stated as follows.

Theorem 1.1. *Suppose that the initial data satisfy the compatibility condition with the boundary conditions. There exists a constant $\delta_1 > 0$ such that if*

$$\|\rho_0 - 1\|_2 + \|(u_0, \mathcal{T}_0 - 1, H_0)\|_3 \leq \delta_1,$$

then the initial boundary value problem (1.1)-(1.4) admits a unique strong solution $(\rho, u, \mathcal{T}, H)$ globally in time satisfying

$$\rho - 1 \in C^0(0, \infty, H^2(\Omega)) \cap C^1(0, \infty, H^1(\Omega)),$$

$$(u, \mathcal{T} - 1, H) \in C^0(0, \infty, H^3(\Omega)) \cap C^1(0, \infty, H^1(\Omega)),$$

and

$$\begin{aligned} & \left(\|(u, \mathcal{T} - 1, H)\|_3^2 + \|\rho - 1\|_2^2 + \|(\rho_t, u_t, H_t, \mathcal{T}_t)\|_1^2 \right)(t) \\ & + C \int_0^t \left(\|(\nabla u, \nabla \mathcal{T}, \nabla H)\|_2^2 + \|\nabla \rho\|_1^2 + \|(u_t, \mathcal{T}_t, H_t)\|_2^2 + \|\rho_t\|_1^2 \right) ds \\ & \leq \left(\|(u, \mathcal{T} - 1, H)\|_3^2 + \|\rho - 1\|_2^2 + \|(\rho_t, u_t, H_t, \mathcal{T}_t)\|_1^2 \right)(0). \end{aligned}$$

where $C > 0$ is a positive constant independent of t .

Remark 1.2. *The compatibility condition of the initial data with the boundary conditions is standard to ensure the regularity of the solutions. The data*

$$\|(\rho_t, u_t, H_t, \mathcal{T}_t)\|_1^2(0)$$

is determined by the the initial data $(\rho_0, u_0, \mathcal{T}_0, H_0)$ through the equations (1.1).

For the initial data close to the constant state $(1, 0, 1, 0)$ in higher-order Sobolev norm, we can improve the regularity of solutions in Theorem 1.1. Precisely,

Theorem 1.3. *Suppose that the initial data satisfy the compatibility condition with the boundary conditions. There exists a constant $\delta > 0$ such that if*

$$\|\rho_0 - 1\|_3 + \|(u_0, \mathcal{T}_0 - 1, H_0)\|_4 \leq \delta,$$

then the initial boundary value problem (1.1)- (1.4) admits a unique smooth solution $(\rho, u, \mathcal{T}, H)$ globally in time satisfying

$$\begin{aligned} \rho - 1 &\in C^0(0, \infty, H^3(\Omega)) \cap C^1(0, \infty, H^2(\Omega)), \\ (u, \mathcal{T} - 1, H) &\in C^0(0, \infty, H^4(\Omega)) \cap C^1(0, \infty, H^2(\Omega)), \end{aligned}$$

and

$$\begin{aligned} &\left(\|\rho - 1\|_3^2 + \|(u, \mathcal{T} - 1, H)\|_4^2 + \|(\rho_t, u_t, \mathcal{T}_t, H_t)\|_2^2 + \|(\rho_{tt}, u_{tt}, \mathcal{T}_{tt}, H_{tt})\|^2 \right)(t) \\ &+ C \int_0^t \left(\|(\nabla u, \nabla \mathcal{T}, \nabla H)\|_3^2 + \|\nabla \rho\|_2^2 + \|(\rho_t, u_t, H_t)\|_2^2 + \|\mathcal{T}_t\|_3^2 + \|(u_{tt}, \mathcal{T}_{tt}, H_{tt})\|_1^2 + \|\rho_{tt}\|^2 \right) ds \\ &\leq \|\rho - 1\|_3^2 + \|(u, \mathcal{T} - 1, H)\|_4^2 + \|(\rho_t, u_t, \mathcal{T}_t, H_t)\|_2^2 + \|(\rho_{tt}, u_{tt}, \mathcal{T}_{tt}, H_{tt})\|^2(0), \end{aligned}$$

where $C > 0$ is a positive constant independent of t .

In addition, we shall show that the solution in Theorem 1.3 approaches the stationary state as $t \rightarrow \infty$, and give the explicit decay rate. More precisely,

Theorem 1.4. *Let $(\rho, u, \mathcal{T}, H)$ be the global solution in Theorem 1.3, and $(1, u_s, 1, H_s)$ with $u_s = H_s = (0, 0, 0)$ be a stationary state. Then it holds:*

$$\begin{aligned} \|(\rho_t, u_t, \mathcal{T}_t, H_t)(t)\| &= O(t^{-1/2}) \quad \text{as } t \rightarrow \infty \\ \|(\nabla \rho, \nabla u, \nabla \mathcal{T}, \nabla H)(t)\| &= O(t^{-1/4}), \quad \text{as } t \rightarrow \infty \\ \|(\nabla^2 u, \nabla^2 \theta, \nabla^2 H)(t)\| &= O(t^{-1/4}), \quad \text{as } t \rightarrow \infty, \\ \|(u, \mathcal{T} - 1, H)(t)\|_{C^0(\Omega)} &= O(t^{-1/4}) \quad \text{as } t \rightarrow \infty \\ \|(\rho - 1)(t)\|_{C^0(\Omega)} &= O(t^{-1/8}) \quad \text{as } t \rightarrow \infty. \end{aligned}$$

Remark 1.5. *Note that for the IBVP of the Navier-Stokes equations with Dirichlet condition $u|_{\partial\Omega} = 0$ on exterior domain, the decay estimates of solutions to stationary state could be found in [9, 10, 19, 21, 26]. By contrast, the big difference in this paper is that we consider the Navier-slip boundary condition for u , instead of the homogeneous Dirichlet condition. Inspired by [9], we would like to form the inequalities as in lemma 2.7. Different from [9], Lemma 5.3 is used to obtain the estimate for $\|\text{curl curl } u\|$, which is important for us to derive the decay estimates of $\|(\nabla^2 u, \nabla \rho)\|$ (Lemma 5.4). Moreover, some classical elliptic estimates will be applied for H to find the corresponding decay rate.*

Remark 1.6. *The algebraic decay for Cauchy problem of MHD equations was shown in [6, 23, 36] by careful analysis of the linearized equation by using the Fourier transformation. However, due to the presence of the physical boundary, the approach of obtaining the decay for the Cauchy problem used in [6, 23, 36] cannot be applied to the problem studied in this paper. Moreover, the conditions required for initial data are lesser in the present paper.*

This paper is organized as follows. Section 2 contains some notations and basic lemmas for later use. In Section 3 we prove the global existence of strong solutions. The improved regularity of the strong solutions in Theorem 1.1 will be proved in Sec. 4. Finally, Section 5 is devoted to obtaining the decay rates. Through out of this paper, we introduce some notations for later use.

The L^p -norm on $x \in \Omega$ for $\phi(x, t)$ is given by

$$\|\phi\|_{L^p} \equiv \|\phi\|_{L^p(\Omega)} = \left(\int_{\Omega} |\phi|^p(x, t) dx \right)^{1/p}, \quad \|\phi\| \equiv \|\phi\|_{L^2},$$

$1 \leq p < +\infty$ and H^m is used to denote the standard Sobolev space with the following norm

$$\|\phi\|_m \equiv \|\phi\|_{H^m(\Omega)} = \left(\sum_{l=0}^m \|\nabla^l \phi\|^2 \right)^{1/2}.$$

$H^{m,p}$ is applied to express the standard Sobolev space with the following norm

$$\|\phi\|_{H^{m,p}} \equiv \left(\sum_{l=0}^m \|\nabla^l \phi\|_p^p \right)^{1/p}.$$

Besides, C will be used as a generic constant independent of time t .

2. NOTATIONS AND BASIC LEMMAS

In this section, we list some lemmas which will be frequently used throughout this paper.

Firstly, we recall some inequalities of Sobolev type.

Lemma 2.1. *Let $\Omega \subset \mathbb{R}^3$ be a domain with smooth boundary. Then*

- (i) $\|f\|_{C^0(\bar{\Omega})} \leq C\|f\|_{H^{m,p}}$ for $f \in H^{m,p}(\Omega)$, $mp > 3 > (m-1)p$.
- (ii) $\|f\|_{L^p} \leq C\|f\|_1$ for $f \in H^1(\Omega)$, $2 \leq p \leq 6$.
- (iii) $\|f\|_{L^6} \leq C\|\nabla f\|$ for $f \in H^1(\Omega)$.

The following lemma allows one to control the H^m -norm of a vector valued function v by its H^{m-1} -norm of $\text{curl} v$ and $\text{div} v$ (see [34]).

Lemma 2.2. *Let Ω be a domain in $\mathbb{R}^N$ with smooth boundary $\partial\Omega$ and outward normal n . Then there exists a constant $C > 0$, such that*

$$\|v\|_{H^s(\Omega)} \leq C(\|\text{div} v\|_{H^{s-1}(\Omega)} + \|\text{curl} v\|_{H^{s-1}(\Omega)} + |v \cdot n|_{H^{s-1/2}(\partial\Omega)} + \|v\|), \quad (2.1)$$

and

$$\|v\|_{H^s(\Omega)} \leq C(\|\text{div} v\|_{H^{s-1}(\Omega)} + \|\text{curl} v\|_{H^{s-1}(\Omega)} + |v \times n|_{H^{s-1/2}(\partial\Omega)} + \|v\|), \quad (2.2)$$

for any $v \in H^s(\Omega)$, $s \geq 1$.

In addition, since the first Betti number of the exterior domain Ω in our paper vanishes, we can apply Theorem 3.2 in [31] to obtain the following refined estimate of $\|\nabla v\|$ which is crucial in the case of exterior domain. We also note that the topological property of Ω is necessary for the following Proposition.

Proposition 2.3. ([31]) Let $\nabla v \in L^2(\Omega)$, $v \in L^6(\Omega)$ and $v \cdot n|_{\partial\Omega} = 0$. The estimate (C independent of v)

$$\|\nabla v\| \leq C_\Omega(\|div v\| + \|\operatorname{curl} v\|) \quad (2.3)$$

is true for all v as above if and only if Ω has a first Betti number of zero.

Next, we list some elliptic estimates for elliptic equations with Navier-slip or Neumann boundary conditions in the smooth exterior domain Ω .

Lemma 2.4. ([30]) Let s be an integer ≥ 0 . Suppose that u satisfy the Lamé equation:

$$\begin{cases} -\mu\Delta u - (\mu + \lambda)\nabla \operatorname{div} u = f, & \text{in } \Omega, \\ u \cdot n = 0, \quad \operatorname{curl} u \times n = 0 & \text{on } \partial\Omega \end{cases}$$

Then, it holds

$$\|\nabla^2 u\|_{H^s(\Omega)} \leq C(\|f\|_{H^s(\Omega)} + \|\nabla u\|_{L^2(\Omega)}).$$

In particular, the same conclusion also holds for Laplace operator, that is, for

$$v \cdot n = 0, \quad \operatorname{curl} v \times n = 0 \quad \text{on } \partial\Omega,$$

it holds

$$\|\nabla^2 v\|_{H^s(\Omega)} \leq C(\|\Delta v\|_{H^s(\Omega)} + \|\nabla v\|_{L^2(\Omega)}).$$

Lemma 2.5. ([11]) Let s be an integer ≥ 0 . Suppose that (v, p) is the solution of the Stokes problem:

$$\begin{cases} \operatorname{div} v = g, & \text{in } \Omega, \\ -\Delta v + \nabla p = f, & \text{in } \Omega, \\ v \cdot n = 0, \quad \operatorname{curl} v \times n = 0 & \text{on } \partial\Omega \end{cases}$$

Then, it holds

$$\|\nabla^2 v\|_{H^s(\Omega)} + \|\nabla p\|_{H^s(\Omega)} \leq C(\|f\|_{H^s(\Omega)} + \|g\|_{H^{s+1}(\Omega)} + \|\nabla v\|_{L^2(\Omega)}).$$

Lemma 2.6. ([1]) For any function $\theta \in H^s(\Omega)$ with $\frac{\partial \theta}{\partial n}|_{\partial\Omega} = 0$, then it holds

$$\|\nabla^2 \theta\|_{H^s(\Omega)} \leq C(\|\Delta \theta\|_{H^s(\Omega)} + \|\nabla \theta\|_{L^2(\Omega)}).$$

Finally, we recall the following differential inequalities which is the main argument in proving Theorem 1.3. The reader is referred to [25] for a proof.

Lemma 2.7. (i) Let $g \in C^1([t_0, \infty))$ such that $g \geq 0$, $E = \int_{t_0}^\infty g(t)dt < \infty$ and

$$g'(t) \leq a(t)g(t) \quad \text{for all } t \geq t_0$$

where $a \geq 0$, $M = \int_{t_0}^\infty a(t)dt < \infty$. Then

$$g(t) \leq \left((t_0 g(t_0) + 1) \exp(E + M) - 1 \right) t^{-1} \quad \forall t \geq t_0.$$

(ii) Let $g \in C^1([t_0, \infty))$ such that $g \geq 0$ and

$$g'(t) + c_0 g(t) \leq c_1 t^{-\alpha} \quad \text{for all } t \geq t_0$$

with positive constants c_0, c_1 and α . Then there exists a $t_1 = t_1(c_0, \alpha) \geq t_0$ such that

$$g(t) \leq C t^{-\alpha}, \quad \text{for } t \geq t_1.$$

3. GLOBAL EXISTENCE OF STRONG SOLUTIONS

Let

$$\rho = 1 + q, \mathcal{T} = 1 + \theta,$$

from the initial boundary value problem (1.1), then (q, u, θ, H) satisfy the following system

$$\begin{cases} q_t + \operatorname{div} u = -\operatorname{div}(qu), \\ \rho(u_t + u \cdot \nabla u) - \mu \Delta u - (\mu + \lambda) \nabla \operatorname{div} u + R \nabla q + R \nabla \theta = -R \nabla(q\theta) + \operatorname{curl} H \times H, \\ c_v \rho(\theta_t + u \cdot \nabla \theta) - \kappa \Delta \theta + R \operatorname{div} u = \lambda(\operatorname{div} u)^2 + 2\mu |S(u)|^2 - R(\rho\theta + q) \operatorname{div} u + \eta |\operatorname{curl} H|^2, \\ H_t - \eta \Delta H = \operatorname{curl}(u \times H), \quad \operatorname{div} H = 0, \end{cases} \quad (3.1)$$

with the initial data

$$(q, u, \theta, H)|_{t=0} = (q_0, u_0, \theta_0, H_0) \quad (3.2)$$

and the boundary conditions

$$u \cdot n = 0, \quad \operatorname{curl} u \times n = 0, \quad H \cdot n = 0, \quad \operatorname{curl} H \times n = 0, \quad \frac{\partial \theta}{\partial n} = 0, \quad \text{on } x \in \partial \Omega. \quad (3.3)$$

Theorem 1.1 will be proved in this section. The local-in-time well-posedness in the smooth norm is quite standard, therefore to prove Theorem 1.1, it suffices to prove the following *a priori* estimates. In order to describe clearly, we set $\mathcal{E}_1(t)$ and $\mathcal{D}_1(t)$ as follows:

$$\mathcal{E}_1(t) = \|(u, \theta, H)\|_3 + \|q\|_2 + \|(q_t, u_t, H_t, \theta_t)\|_1 \quad (3.4)$$

and

$$\mathcal{D}_1(t) = \|(\nabla u, \nabla \theta, \nabla H)\|_2 + \|\nabla q\|_1 + \|(u_t, \theta_t, H_t)\|_2 + \|q_t\|_1 + \|(q_{tt}, u_{tt}, \theta_{tt}, H_{tt})\|. \quad (3.5)$$

Proposition 3.1. (*a priori estimates*) Let (q, u, θ, H) be a solution to the initial boundary value problem (3.1), (3.2) and (3.3) in $t \in [0, T]$. Then there exists positive constants C and δ_1 which are independent of t , such that if

$$\sup_{0 \leq t \leq T} \mathcal{E}_1(t) \leq \delta_1,$$

then there holds, for any $t \in [0, T]$,

$$\mathcal{E}_1^2(t) + C \int_0^t \mathcal{D}_1^2(s) ds \leq C \mathcal{E}_1^2(0).$$

Now, we shall divide the proof of Proposition 3.1 into several lemmas.

3.1. Estimates for lower-order derivatives. To begin with, we have the following basic energy estimate.

Lemma 3.2.

$$\|(q, u, \theta, H)\|^2 + c_0 \int_0^t \|(\nabla u, \nabla \theta, \nabla H)\|^2 ds \leq \|(q, u, \theta, H)\|^2(0) + C \delta_1 \int_0^t \mathcal{D}_1^2(s) ds,$$

where $c_0 > 0$, $C > 0$ are positive constants independent of t .

Proof. Computing the following integral

$$\int_{\Omega} \{(3.1)_1 Rq + (3.1)_2 \cdot u + (3.1)_3 \theta + (3.1)_4 \cdot H\} dx,$$

and noting that $\Delta u = -\operatorname{curl}^2 u + \nabla \operatorname{div} u$, after integrating by parts, one has

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \int_{\Omega} \left(Rq^2 + \rho|u|^2 + c_v \rho \theta^2 + |H|^2 \right) dx + \mu \|\operatorname{curl} u\|^2 + (2\mu + \lambda) \|\operatorname{div} u\|^2 + \kappa \|\nabla \theta\|^2 + \eta \|\operatorname{curl} H\|^2 \\ &= \int_{\Omega} \left(\left(-\frac{1}{2} Rq^2 - R\rho \theta^2 \right) \operatorname{div} u + \left(\lambda (\operatorname{div} u)^2 + 2\mu |S(u)|^2 + \eta |\operatorname{curl} H|^2 \right) \theta \right) dx \\ &\leq C\mathcal{E}(t)\mathcal{D}_1^2(t). \end{aligned} \tag{3.6}$$

Applying Proposition 2.3 to u and H , it holds that

$$\|\nabla u\|^2 \leq C(\|\operatorname{curl} u\|^2 + \|\operatorname{div} u\|^2), \tag{3.7}$$

and

$$\|\nabla H\|^2 \leq C\|\operatorname{curl} H\|^2. \tag{3.8}$$

Therefore, integrating (3.6) over $[0, t]$, and noting that $2\mu + \lambda > 0$, we prove the basic energy estimate. $\square$

Next, we prove estimates of the first-order derivatives $(\nabla u, \nabla \theta, \nabla H)$.

Lemma 3.3.

$$\|(\nabla u, \nabla \theta, \nabla H)\|^2 + c \int_0^t \|(u_t, q_t, \theta_t, H_t)\|^2 ds \leq C\mathcal{E}_1^2(0) + C\delta_1 \int_0^t \mathcal{D}_1^2(s) ds,$$

where $c > 0$, $C > 0$ are positive constants independent of t .

Proof. Computing the following integral

$$\int_{\Omega} \{ (3.1)_2 \cdot u_t + (3.1)_1 \cdot q_t + (3.1)_3 \cdot \theta_t + (3.1)_4 \cdot H_t \} dx$$

integrating by parts, one has

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \left\{ \mu \|\operatorname{curl} u\|^2 + (2\mu + \lambda) \|\operatorname{div} u\|^2 + \kappa \|\nabla \theta\|^2 + \eta \|\operatorname{curl} H\|^2 \right\} - \frac{d}{dt} \int_{\Omega} R(q + \theta) \operatorname{div} u dx \\ &+ \|\sqrt{\rho} u_t\|^2 + \|q_t\|^2 + c_v \|\sqrt{\rho} \theta_t\| + \|H_t\|^2 \\ &= \int_{\Omega} \left((R+1)q_t + 2R\theta_t \right) \operatorname{div} u dx + \int_{\Omega} \operatorname{div}(qu) q_t dx + \int_{\Omega} \left(\rho u \cdot \nabla u - R\nabla(q\theta) + \operatorname{curl} H \times H \right) \cdot u_t dx \\ &+ \int_{\Omega} \left(-c_v \rho u \cdot \nabla \theta + \lambda (\operatorname{div} u)^2 + 2\mu |Su|^2 - R(\rho\theta + q) \operatorname{div} u + \eta |\operatorname{curl} H|^2 \right) \theta_t dx + \int_{\Omega} \operatorname{curl}(u \times H) \cdot H_t dx \\ &\leq \frac{1}{2} \left(\|\sqrt{\rho} u_t\|^2 + \|q_t\|^2 + c_v \|\sqrt{\rho} \theta_t\| + \|H_t\|^2 \right) + C \|\operatorname{div} u\|^2 + \mathcal{E}_1(t) \mathcal{D}_1^2(t), \end{aligned}$$

which gives

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \left\{ \mu \|\operatorname{curl} u\|^2 + (2\mu + \lambda) \|\operatorname{div} u\|^2 + \kappa \|\nabla \theta\|^2 + \eta \|\operatorname{curl} H\|^2 \right\} - \frac{d}{dt} \int_{\Omega} R(q + \theta) \operatorname{div} u dx \\ &+ \frac{1}{2} \left(\|\sqrt{\rho} u_t\|^2 + \|q_t\|^2 + c_v \|\sqrt{\rho} \theta_t\| + \|H_t\|^2 \right) \\ &\leq C \|\operatorname{div} u\|^2 + \delta_1 \mathcal{D}_1^2(t). \end{aligned} \tag{3.9}$$

Then, integrating (3.9) over $[0, t]$, then Lemma 3.2, (3.7) and (3.8) implies Lemma 3.3. $\square$

Now, we prove estimates of the first-order derivatives ∇q .

Lemma 3.4.

$$\|(\nabla q, \operatorname{div} u, \nabla \theta)\|^2 + c \int_0^t \|(\nabla \operatorname{div} u, \Delta \theta)\|^2 ds \leq \|(\nabla q, \operatorname{div} u, \nabla \theta)\|^2(0) + C\delta_1 \int_0^t \mathcal{D}_1^2(s) ds, \quad (3.10)$$

where $c > 0$, $C > 0$ are positive constants independent of t . Moreover,

$$\int_0^t \|\nabla^2 \theta\|^2 ds \leq C\mathcal{E}_1^2(0) + C\delta_1 \int_0^t \mathcal{D}_1^2(s) ds. \quad (3.11)$$

Proof. Computing the following integral

$$\int_{\Omega} \{ (3.1)_2 \cdot \nabla \operatorname{div} u + \nabla (3.1)_1 \cdot R \nabla q + \nabla (3.1)_3 \cdot \nabla \theta \} dx,$$

and integrating by parts, noting that $\int_{\Omega} \operatorname{curl}^2 u \cdot \nabla \operatorname{div} u dx = 0$ thanks to $\operatorname{curl} u \times n|_{\partial\Omega} = 0$, one has

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \int_{\Omega} \left(\rho |\operatorname{div} u|^2 + R |\nabla q|^2 + \rho |\nabla \theta|^2 \right) dx + (2\mu + \lambda) \|\nabla \operatorname{div} u\|^2 + \kappa \|\Delta \theta\|^2 \\ &= \frac{1}{2} \int_{\Omega} q_t |\operatorname{div} u|^2 dx + \frac{c_v}{2} \int_{\Omega} q_t |\nabla \theta|^2 dx - \int_{\Omega} u_t \cdot \nabla q \operatorname{div} u dx \\ &+ \int_{\Omega} \left(\rho u \cdot \nabla u + R \nabla(q\theta) - \operatorname{curl} H \times H \right) \cdot \nabla \operatorname{div} u dx \\ &- \int_{\Omega} \nabla \operatorname{div}(qu) \cdot R \nabla q dx - c_v \int_{\Omega} \left(\nabla q(\theta_t + u \cdot \nabla \theta) + \rho \nabla(u \cdot \nabla \theta) \right) \cdot \nabla \theta dx \\ &- \int_{\Omega} \nabla \left(R(\rho\theta + q) \operatorname{div} u \right) \cdot \nabla \theta dx + \int_{\Omega} \nabla \left(\lambda |\operatorname{div} u|^2 + 2\mu |Su|^2 + \eta |\operatorname{curl} H|^2 \right) \cdot \nabla \theta dx. \end{aligned}$$

The nonlinear linear terms on the right hand-side can be easily controlled by $\mathcal{E}_1(t)\mathcal{D}_1^2(t)$. Therefore, integrating over $[0, t]$ implies the desired estimate (3.10). Furthermore, applying Lemma 2.6 to θ , we get

$$\begin{aligned} \int_0^t \|\nabla^2 \theta\|^2 ds &\leq C \int_0^t \left(\|\Delta \theta\|^2 + \|\nabla \theta\|^2 \right) ds \\ &\leq C\mathcal{E}_1^2(0) + C\delta_1 \int_0^t \mathcal{D}_1^2(s) ds, \end{aligned}$$

due to Lemma 3.2 and (3.10). Thus the proof of Lemma 3.4 is completed. $\square$

In order to obtain the dissipation estimates of $\|\nabla^2 u\|^2$ and $\|\nabla^2 H\|^2$, in view of Lemma 2.4, we need to control $\|\operatorname{curl}^2 u\|_{L_t^2(L^2)}$ and $\|\operatorname{curl}^2 H\|_{L_t^2(L^2)}$.

Lemma 3.5.

$$\|(\sqrt{\rho} \operatorname{curl} u, \operatorname{curl} H)\|^2 + c \int_0^t \|(\operatorname{curl}^2 u, \operatorname{curl}^2 H)\|^2 ds \leq C\|(\sqrt{\rho} \operatorname{curl} u, \operatorname{curl} H)\|^2(0) + C\delta_1 \int_0^t \mathcal{D}_1^2(s) ds, \quad (3.12)$$

where $c > 0$, $C > 0$ are positive constants independent of t . Moreover,

$$\int_0^t \left(\|\nabla^2 u\|^2 + \|\nabla^2 H\|^2 \right) ds \leq C\mathcal{E}_1^2(0) + C\delta_1 \int_0^t \mathcal{D}_1^2(s) ds. \quad (3.13)$$

Proof. Let $\operatorname{curl} u = w$, then one has the identity

$$\operatorname{curl}(u \cdot \nabla) u = u \cdot \nabla w - w \cdot \nabla u + w \operatorname{div} u. \quad (3.14)$$

Taking operator curl to equation (3.1)₂, we easily derive the following equations for w :

$$\rho(w_t + u \cdot \nabla w) - \mu \Delta w = K + \operatorname{curl}(\operatorname{curl} H \times H), \quad (3.15)$$

where

$$K = -\rho w \cdot \nabla u - \rho w \operatorname{div} u - \nabla q \times (u_t + u \cdot \nabla u).$$

Multiplying (3.15) by w , noting that $\Delta w = -\operatorname{curl} \operatorname{curl} w$, integrating by parts with $w \times n|_{\partial\Omega} = 0$, we obtain

$$\frac{1}{2} \frac{d}{dt} \int_{\Omega} \rho |w|^2 dx + \mu \|\operatorname{curl} w\|^2 = \int_{\Omega} K \cdot w dx + \int_{\Omega} \operatorname{curl}(\operatorname{curl} H \times H) \cdot w dx. \quad (3.16)$$

We estimate the right-hand side as follows:

$$\begin{aligned} \int_{\Omega} K \cdot w dx &\leq C \left(\|w\|_{L^6} \|w\| \|\nabla u\|_{L^3} + \|\nabla q\|_{L^3} \|u_t\|_{L^6} \|w\| + \|\nabla q\|_{L^6} \|w\|_{L^6} \|u\|_{L^6} \|\nabla u\| \right) \\ &\leq \frac{\mu}{4} \|\operatorname{curl} w\|^2 + C \delta_1 \left(\|\nabla u_t\|^2 + \|\nabla u\|^2 \right), \end{aligned}$$

where we have used the estimate $\|w\|_{L^6} \leq C \|\nabla w\| \leq C(\|\operatorname{curl} w\|^2 + \|w\|^2)$ thanks to Lemma 2.2 with $w \times n|_{\partial\Omega} = 0$. Integrating by parts with the boundary condition $w \times n|_{\partial\Omega} = 0$ again implies

$$\begin{aligned} \int_{\Omega} \operatorname{curl}(\operatorname{curl} H \times H) \cdot w dx &= \int_{\Omega} (\operatorname{curl} H \times H) \cdot \operatorname{curl} w dx \\ &\leq \|H\|_{L^\infty} \|\operatorname{curl} H\| \|\operatorname{curl} w\| \\ &\leq \frac{\mu}{4} \|\operatorname{curl} w\|^2 + C \delta_1 \|\operatorname{curl} H\|^2. \end{aligned}$$

Putting these into (3.16) yields

$$\frac{1}{2} \frac{d}{dt} \int_{\Omega} \rho |w|^2 dx + \frac{\mu}{2} \|\operatorname{curl} w\|^2 \leq C \delta_1 \left(\|\nabla u_t\|^2 + \|\nabla u\|^2 + \|\nabla H\|^2 \right). \quad (3.17)$$

Similarly, letting $\operatorname{curl} H = \phi$, taking curl to equation (3.1)₄, we obtain

$$\phi_t + \eta \operatorname{curl}^2 \phi = \operatorname{curl}^2(u \times H). \quad (3.18)$$

Multiplying (3.18) by ϕ , integrating by parts with $\phi \times n|_{\partial\Omega} = 0$, one has

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \int_{\Omega} |\phi|^2 dx + \eta \|\operatorname{curl} \phi\|^2 &= \int_{\Omega} \operatorname{curl}^2(u \times H) \cdot \phi dx \\ &= \int_{\Omega} \operatorname{curl}(u \times H) \cdot \operatorname{curl} \phi dx \\ &\leq \frac{\eta}{2} \|\operatorname{curl} \phi\|^2 + C \|\operatorname{curl}(u \times H)\|^2 \\ &\leq \frac{\eta}{2} \|\operatorname{curl} \phi\|^2 + C \delta_1 \left(\|\nabla u\|^2 + \|\nabla H\|^2 \right). \end{aligned}$$

Therefore,

$$\frac{1}{2} \frac{d}{dt} \int_{\Omega} |\phi|^2 dx + \frac{\eta}{2} \|\operatorname{curl} \phi\|^2 \leq C \delta_1 \left(\|\nabla u\|^2 + \|\nabla H\|^2 \right). \quad (3.19)$$

Summing up (3.17) and (3.19), integrating over $[0, t]$ gives (3.12).

Furthermore, applying Lemma 2.4 with $v = H$, and noting that $\Delta H = -\text{curl}^2 H$, we get

$$\begin{aligned} \int_0^t \|\nabla^2 H\|^2 ds &\leq C \int_0^t \left(\|\Delta H\|^2 + \|\nabla H\|^2 \right) ds \\ &\leq C \int_0^t \left(\|\text{curl}^2 H\|^2 + \|\nabla H\|^2 \right) ds \\ &\leq C\mathcal{E}_1^2(0) + C\delta_1 \int_0^t \mathcal{D}_1^2(s) ds, \end{aligned}$$

thanks to the basic energy estimate Lemma 3.2 and (3.12).

Similarly, applying Lemma 2.4 with $v = u$, we obtain

$$\begin{aligned} \int_0^t \|\nabla^2 u\|^2 ds &\leq C \int_0^t \left(\|\Delta u\|^2 + \|\nabla u\|^2 \right) ds \\ &\leq C \int_0^t \left(\|\text{curl}^2 u\|^2 + \|\nabla \text{div} u\|^2 + \|\nabla u\|^2 \right) ds \\ &\leq C\mathcal{E}_1^2(0) + C\delta_1 \int_0^t \mathcal{D}_1^2(s) ds, \end{aligned}$$

thanks to Lemma 3.2, Lemma 3.4 and (3.12). □

Next, we prove estimates of the first-order temporal derivatives $(q_t, u_t, \theta_t, H_t)$.

Lemma 3.6.

$$\|(q_t, u_t, \theta_t, H_t)\|^2 + c \int_0^t \|(\nabla u_t, \nabla \theta_t, \nabla H_t)\|^2 ds \leq C \|(q_t, u_t, \theta_t, H_t)(0)\|^2 + C\delta_1 \int_0^t \mathcal{D}_1^2(s) ds,$$

where $c > 0$, $C > 0$ are positive constants independent of t .

Proof. Computing the following integral

$$\int_{\Omega} \{ \partial_t(3.1)_1 R q_t + \partial_t(3.1)_2 \cdot u_t + \partial_t(3.1)_3 \theta_t + \partial_t(3.1)_4 \cdot H_t \} dx,$$

and noting that differentiation of the system (3.1) with respect to t will keep the boundary conditions (3.3), it is easy to prove Lemma 3.6, we omit it. □

3.2. Estimates for the second-order derivatives. Firstly, we prove estimates of the second-order derivatives $(\nabla u_t, \nabla \theta_t, \nabla H_t)$.

Lemma 3.7.

$$\|(\nabla u_t, \nabla \theta_t, \nabla H_t)\|^2 + c \int_0^t \|(u_{tt}, q_{tt}, \theta_{tt}, H_{tt})\|^2 ds \leq C\mathcal{E}_1^2(0) + C\delta_1 \int_0^t \mathcal{D}_1^2(s) ds,$$

where $c > 0$, $C > 0$ are positive constants independent of t .

Proof. Computing the following integral

$$\int_{\Omega} \{ \partial_t(3.1)_2 \cdot u_{tt} + \partial_t(3.1)_1 \cdot q_{tt} + \partial_t(3.1)_3 \cdot \theta_{tt} + \partial_t(3.1)_4 \cdot H_{tt} \} dx$$

integrating by parts, one has

$$\begin{aligned}
& \frac{1}{2} \frac{d}{dt} \left\{ \mu \|\operatorname{curl} u_t\|^2 + (2\mu + \lambda) \|\operatorname{div} u_t\|^2 + \kappa \|\nabla \theta_t\|^2 + \eta \|\operatorname{curl} H_t\|^2 \right\} - \frac{d}{dt} \int_{\Omega} R(q_t + \theta_t) \operatorname{div} u_t dx \\
& + \|\sqrt{\rho} u_{tt}\|^2 + \|q_{tt}\|^2 + c_v \|\sqrt{\rho} \theta_{tt}\|^2 + \|H_{tt}\|^2 \\
& = \int_{\Omega} \left((R+1)q_{tt} + 2R\theta_{tt} \right) \operatorname{div} u_t dx + \int_{\Omega} \operatorname{div}(qu)_t q_{tt} dx \\
& + \int_{\Omega} \left(-q_t(u_t + u \cdot \nabla u) + \rho(u \cdot \nabla u)_t - R\nabla(q\theta)_t + (\operatorname{curl} H \times H)_t \right) \cdot u_{tt} dx \\
& + \int_{\Omega} \left(-q_t(\theta_t + u \cdot \nabla \theta) - c_v \rho(u \cdot \nabla \theta)_t \right) \theta_{tt} dx \\
& + \int_{\Omega} \left(\lambda(\operatorname{div} u)^2 + 2\mu|Su|^2 - R(\rho\theta + q)\operatorname{div} u + \eta|\operatorname{curl} H|^2 \right)_t \theta_{tt} dx + \int_{\Omega} \operatorname{curl}(u \times H)_t \cdot H_{tt} dx \\
& \leq \frac{1}{2} \left(\|\sqrt{\rho} u_{tt}\|^2 + \|q_{tt}\|^2 + c_v \|\sqrt{\rho} \theta_{tt}\|^2 + \|H_{tt}\|^2 \right) + C \|\operatorname{div} u_t\|^2 + C \mathcal{E}_1(t) \mathcal{D}_1^2(t).
\end{aligned} \tag{3.20}$$

On the other hand, applying Proposition 2.3 to u_t and H_t with $u_t \cdot n|_{\partial\Omega} = 0$, $H_t \cdot n|_{\partial\Omega} = 0$, it holds

$$\|\nabla u_t\|^2 \leq C(\|\operatorname{curl} u_t\|^2 + \|\operatorname{div} u_t\|^2), \quad \text{and} \quad \|\nabla H_t\|^2 \leq C\|\operatorname{curl} H_t\|^2. \tag{3.21}$$

Then, integrating (3.20) over $[0, t]$, recalling Lemma 3.6, and using (3.21) we prove Lemma 3.7. $\square$

Next, we prove estimates of the second-order derivatives ∇q_t .

Lemma 3.8.

$$\|(\nabla q_t, \sqrt{\rho} \operatorname{div} u_t, \sqrt{\rho} \nabla \theta_t)\|^2 + c \int_0^t \|(\nabla \operatorname{div} u_t, \Delta \theta_t)\|^2 ds \leq \|(\nabla q_t, \sqrt{\rho} \operatorname{div} u_t, \sqrt{\rho} \nabla \theta_t)\|^2(0) + C\delta_1 \int_0^t \mathcal{D}_1^2(s) ds, \tag{3.22}$$

where $c > 0$, $C > 0$ are positive constants independent of t . Moreover,

$$\int_0^t \|\nabla^2 \theta_t\|^2 ds \leq C\mathcal{E}_1^2(0) + C\delta_1 \int_0^t \mathcal{D}_1^2(s) ds. \tag{3.23}$$

Proof. Taking ∂_t to (3.1)₂, multiplying the resulted identity by $\nabla \operatorname{div} u_t$, and integrating over Ω , one has

$$\begin{aligned}
& \frac{1}{2} \frac{d}{dt} \int_{\Omega} \rho |\operatorname{div} u_t|^2 dx + (2\mu + \lambda) \|\nabla \operatorname{div} u_t\|^2 - R \int_{\Omega} (\nabla q_t + \nabla \theta_t) \cdot \nabla \operatorname{div} u_t dx \\
& = \frac{1}{2} \int_{\Omega} q_t |\operatorname{div} u_t|^2 dx + \int_{\Omega} u_{tt} \cdot \nabla q \operatorname{div} u_t dx \\
& + \int_{\Omega} \left(\rho(u \cdot \nabla u)_t + q_t(u_t + u \cdot \nabla u) + R\nabla(q\theta)_t - (\operatorname{curl} H \times H)_t \right) \cdot \nabla \operatorname{div} u_t dx \\
& \leq \|q_t\|_{L^\infty} \|\operatorname{div} u_t\|^2 + \|u_{tt}\| \|\nabla q\|_{L^3} \|\operatorname{div} u_t\|_{L^6} \\
& + \left(\|(u, q, \theta, H)\|_2 + \|q_t\|_1 + \|u\|_2 \|q_t\|_1 \right) \cdot \left(\|\nabla \operatorname{div} u_t\|^2 + \|\nabla u_t\|^2 + \|\nabla^2 u\|^2 + \|\nabla q_t\|^2 + \|\nabla \theta_t\|^2 + \|\nabla H_t\|^2 \right),
\end{aligned}$$

which yields

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \int_{\Omega} \rho |\operatorname{div} u_t|^2 dx + \frac{2\mu + \lambda}{2} \|\nabla \operatorname{div} u_t\|^2 - R \int_{\Omega} (\nabla q_t + \nabla \theta_t) \cdot \nabla \operatorname{div} u_t dx \\ & \leq \delta_1 \left(\|\operatorname{div} u_t\|^2 + \|u_{tt}\|^2 + \|\nabla \operatorname{div} u_t\|^2 + \|\nabla u_t\|^2 + \|\nabla^2 u\|^2 + \|\nabla q_t\|^2 + \|\nabla \theta_t\|^2 + \|\nabla H_t\|^2 \right). \end{aligned} \quad (3.24)$$

To eliminate the singular terms on the left-hand side of (3.24), we take ∂_t to (3.1)₃, multiply the resulted identity by $\Delta \theta_t$, and integrate over Ω , then it holds

$$\begin{aligned} & \frac{c_v}{2} \frac{d}{dt} \int_{\Omega} \rho |\nabla \theta_t|^2 dx + \kappa \|\Delta \theta_t\|^2 + R \int_{\Omega} \nabla \theta_t \cdot \nabla \operatorname{div} u_t dx \\ & = \frac{c_v}{2} \int_{\Omega} q_t |\nabla \theta_t|^2 dx + c_v \int_{\Omega} \theta_{tt} \nabla q \cdot \nabla \theta_t dx + \int_{\Omega} \left(\rho(u \cdot \nabla \theta)_t + q_t(\theta_t + u \cdot \nabla \theta) \right) \Delta \theta_t dx \\ & - \int_{\Omega} \partial_t \left(\lambda (\operatorname{div} u)^2 + 2\mu |S(u)|^2 - R(\rho \theta + q) \operatorname{div} u + \eta |\operatorname{curl} H|^2 \right) \Delta \theta_t dx \\ & \leq \|q_t\|_{L^3} \|\nabla \theta_t\|_{L^6} \|\nabla \theta_t\| + \|\nabla q\|_{L^3} \|\nabla \theta_t\|_{L^6} \|\theta_{tt}\| \\ & + \left(\|(u, \theta, H)\|_2 + \|(q, q_t)\|_1 \right) \cdot \left(\|\Delta \theta_t\|^2 + \|\nabla u_t\|^2 + \|\nabla \theta_t\|^2 + \|\nabla q_t\|^2 + \|\nabla \operatorname{div} u_t\|^2 + \|\nabla q_t\|^2 + \|\nabla^2 H_t\|^2 \right). \end{aligned}$$

Therefore, by using the elliptic estimate $\|\nabla^2 \theta_t\|^2 \leq C(\|\Delta \theta_t\|^2 + \|\nabla \theta_t\|^2)$, one has

$$\begin{aligned} & \frac{c_v}{2} \frac{d}{dt} \int_{\Omega} \rho |\nabla \theta_t|^2 dx + \frac{\kappa}{2} \|\Delta \theta_t\|^2 + R \int_{\Omega} \nabla \theta_t \cdot \nabla \operatorname{div} u_t dx \\ & \leq C \delta_1 \left(\|\theta_{tt}\|^2 + \|\nabla u_t\|^2 + \|\nabla \theta_t\|^2 + \|\nabla q_t\|^2 + \|\nabla \operatorname{div} u_t\|^2 + \|\nabla q_t\|^2 + \|\nabla^2 H_t\|^2 \right). \end{aligned} \quad (3.25)$$

Similarly, taking $\partial_t \nabla$ to (3.1)₁, multiplying the resulted identity by $R \nabla q_t$, and integrating over Ω , we get

$$\frac{R}{2} \frac{d}{dt} \int_{\Omega} |\nabla q_t|^2 dx + R \int_{\Omega} \nabla q_t \cdot \nabla \operatorname{div} u_t dx = R \int_{\Omega} \partial_t \nabla \operatorname{div}(qu) \cdot \nabla q_t dx. \quad (3.26)$$

The integration on the right hand side will appear term $\nabla^2 q_t$ which is not included in $\mathcal{D}_1(t)$, we shall work on it carefully.

$$\begin{aligned} & \int_{\Omega} \partial_t \nabla \operatorname{div}(qu) \cdot \nabla q_t dx = \int_{\Omega} u \cdot \nabla^2 q_t \cdot \nabla q_t dx \\ & + \int_{\Omega} \left(\operatorname{div} u \nabla q_t + q_t \nabla \operatorname{div} u + \operatorname{div} u_t \nabla q + q \nabla \operatorname{div} u_t + \nabla u_t \cdot \nabla q + u_t \cdot \nabla^2 q + \nabla u \cdot \nabla q_t \right) \cdot \nabla q_t dx \\ & \leq \int_{\Omega} u \cdot \nabla^2 q_t \cdot \nabla q_t dx + \left(\|\nabla u\|_{L^\infty} + \|\nabla \operatorname{div} u\|_{L^3} + \|\nabla q\|_{L^3} + \|q\|_{L^\infty} + \|\nabla^2 q\| \right) \cdot \left(\|\nabla q_t\|^2 + \|u_t\|_2^2 \right) \\ & \leq \int_{\Omega} u \cdot \nabla^2 q_t \cdot \nabla q_t dx + \delta_1 \left(\|\nabla q_t\|^2 + \|u_t\|_2^2 \right). \end{aligned}$$

In addition, integrating by parts with $u \cdot n|_{\partial\Omega} = 0$ gives

$$\int_{\Omega} u \cdot \nabla^2 q_t \cdot \nabla q_t dx = -\frac{1}{2} \int_{\Omega} \operatorname{div} u |\nabla q_t|^2 dx \leq \delta_1 \|\nabla q_t\|^2.$$

Therefore, plugging into (3.26), we obtain

$$\frac{R}{2} \frac{d}{dt} \int_{\Omega} |\nabla q_t|^2 dx + R \int_{\Omega} \nabla q_t \cdot \nabla \operatorname{div} u_t dx \leq \delta_1 \left(\|\nabla q_t\|^2 + \|u_t\|_2^2 \right). \quad (3.27)$$

Finally, summing (3.24), (3.25) and (3.27), and integrating the resulting estimates over $[0, t]$, we obtain the desired estimate (3.22).

In addition, applying Lemma 2.6 to θ_t , that is $\int_0^t \|\nabla^2 \theta_t\|^2 ds \leq C \int_0^t (\|\Delta \theta_t\|^2 + \|\nabla \theta_t\|^2) ds$, so together with Lemma 3.6 and (3.22), we obtain the estimate (3.23). This completes the proof of Lemma 3.8. $\square$

Lemma 3.9.

$$\|(\sqrt{\rho} \operatorname{curl} u_t, \operatorname{curl} H_t)\|^2 + c \int_0^t \|(\operatorname{curl}^2 u_t, \operatorname{curl}^2 H_t)\|^2 ds \leq \|(\sqrt{\rho} \operatorname{curl} u_t, \operatorname{curl} H_t)\|^2(0) + C\delta_1 \int_0^t \mathcal{D}_1^2(s) ds, \quad (3.28)$$

where $c > 0$, $C > 0$ are positive constants independent of t . Moreover, it holds

$$\int_0^t (\|\nabla^2 H_t\|^2 + \|\nabla^2 u_t\|^2) ds \leq C\mathcal{E}_1^2(0) + C\delta_1 \int_0^t \mathcal{D}_1^2(s) ds. \quad (3.29)$$

Proof. Notice that differentiation of the equations (3.15) and (3.18) with respect to t will keep the boundary conditions $w_t \times n|_{\partial\Omega} = 0$ and $\phi_t \times n|_{\partial\Omega} = 0$. Similar to the proof on Lemma 3.5, computing the integral

$$\int_{\Omega} \partial_t(3.15) \cdot w_t dx, \quad \text{and} \quad \int_{\Omega} \partial_t(3.18) \cdot \phi_t dx$$

gives

$$\frac{1}{2} \frac{d}{dt} \int_{\Omega} \rho |w_t|^2 dx + \frac{\mu}{2} \|\operatorname{curl} w_t\|^2 \leq C\mathcal{E}_1(t) \mathcal{D}_1^2(t), \quad (3.30)$$

and

$$\frac{1}{2} \frac{d}{dt} \int_{\Omega} |\phi_t|^2 dx + \frac{\eta}{2} \|\operatorname{curl} \phi_t\|^2 \leq C\mathcal{E}_1(t) \mathcal{D}_1^2(t). \quad (3.31)$$

Then, summing up (3.30) and (3.31), integrating over $[0, t]$ gives (3.28).

Furthermore, applying Lemma 2.4 with $v = H_t$, and noting that $\Delta H_t = -\operatorname{curl}^2 H_t$, we get

$$\begin{aligned} \int_0^t \|\nabla^2 H_t\|^2 ds &\leq C \int_0^t (\|\Delta H_t\|^2 + \|\nabla H_t\|^2) ds \\ &\leq C \int_0^t (\|\operatorname{curl}^2 H_t\|^2 + \|\nabla H_t\|^2) ds \\ &\leq C\mathcal{E}_1^2(0) + C\delta_1 \int_0^t \mathcal{D}_1^2(s) ds, \end{aligned}$$

thanks to Lemma 3.6 and (3.28).

Similarly, applying Lemma 2.4 with $v = u_t$, we get

$$\begin{aligned} \int_0^t \|\nabla^2 u_t\|^2 ds &\leq C \int_0^t (\|\Delta u_t\|^2 + \|\nabla u_t\|^2) ds \\ &\leq C \int_0^t (\|\operatorname{curl}^2 u_t\|^2 + \|\nabla \operatorname{div} u_t\|^2 + \|\nabla u_t\|^2) ds \\ &\leq C\mathcal{E}_1^2(0) + C\delta_1 \int_0^t \mathcal{D}_1^2(s) ds, \end{aligned}$$

thanks to Lemma 3.8, Lemma 3.6 and (3.28). $\square$

Next, we shall derive the estimate of $(\nabla^2 u, \nabla^2 H, \nabla^2 \theta)$ into the following two lemmas.

Lemma 3.10.

$$\|(\nabla \operatorname{div} u, \Delta \theta)\|^2 + c \int_0^t \|(\nabla q_t, \nabla \theta_t)\|^2 ds \leq C \|(\nabla \operatorname{div} u, \Delta \theta)\|^2(0) + C \delta_1 \int_0^t \mathcal{D}_1^2(s) ds, \quad (3.32)$$

where $c > 0$, $C > 0$ are positive constants independent of t .

Moreover,

$$\|\nabla^2 \theta\|^2 \leq C \mathcal{E}_1^2(0) + C \delta_1 \int_0^t \mathcal{D}_1^2(s) ds. \quad (3.33)$$

Proof. Taking ∂_t to equation (3.1)₂, multiplying the resulted equality by $\nabla \operatorname{div} u$ in $L^2(\Omega)$, and using the fact $\int_\Omega \operatorname{curl}^2 u_t \cdot \nabla \operatorname{div} u dx = 0$, we get

$$\begin{aligned} & \frac{2\mu + \lambda}{2} \frac{d}{dt} \int_\Omega |\nabla \operatorname{div} u|^2 dx - R \int_\Omega (\nabla q_t + \nabla \theta_t) \cdot \nabla \operatorname{div} u dx \\ &= \int_\Omega \rho(u_{tt} + u \cdot \nabla u_t) \cdot \nabla \operatorname{div} u dx + \int_\Omega \left(q_t u_t + (\rho u)_t \cdot \nabla u + R \nabla(q\theta)_t - (\operatorname{curl} H \times H)_t \right) \cdot \nabla \operatorname{div} u dx \\ &\leq \mathcal{E}_1(t) \mathcal{D}_1^2(t). \end{aligned} \quad (3.34)$$

To eliminate the singular terms on the left-hand side of (3.34), we apply operator ∇ to equations (3.1)₁ and (3.1)₃, then multiply the resulting equality by $R \nabla q_t$ and $\nabla \theta_t$ in $L^2(\Omega)$ to get

$$\begin{aligned} & R \|\nabla q_t\|^2 + R \int_\Omega \nabla \operatorname{div} u \cdot \nabla q_t dx \\ &= -R \int_\Omega \nabla(\operatorname{div}(qu)) \cdot \nabla q_t dx \\ &\leq \mathcal{E}_1(t) \mathcal{D}_1^2(t), \end{aligned} \quad (3.35)$$

and

$$\begin{aligned} & \frac{\kappa}{2} \frac{d}{dt} \int_\Omega |\Delta \theta|^2 dx + c_v \|\sqrt{\rho} \nabla \theta_t\|^2 + R \int_\Omega \nabla \operatorname{div} u \cdot \nabla \theta_t dx \\ &= \int_\Omega \left(c_v \nabla q \cdot (\theta_t + u \cdot \theta) + c_v \rho \nabla(u \cdot \nabla \theta) \right) \cdot \nabla \theta_t dx \\ &+ \int_\Omega \nabla \left(\lambda(\operatorname{div} u)^2 + 2\mu |S(u)|^2 - R(\rho\theta + q) \operatorname{div} u + \eta |\operatorname{curl} H|^2 \right) \cdot \nabla \theta_t dx \\ &\leq \mathcal{E}_1(t) \mathcal{D}_1^2(t). \end{aligned} \quad (3.36)$$

Then, from (3.34), combining with (3.35) and (3.36), we obtain (3.32).

Moreover, applying Lemma 2.6 to θ , that is $\|\nabla^2 \theta\|^2 \leq C(\|\Delta \theta\|^2 + \|\nabla \theta\|^2)$, so combining with Lemma 3.4 and (3.32), we obtain the estimate (3.33). The proof is completed. $\square$

Lemma 3.11.

$$\|(\sqrt{\rho} \operatorname{curl}^2 u, \operatorname{curl}^2 H)\|^2 + c \int_0^t \|(\operatorname{curl}^3 u, \operatorname{curl}^3 H)\|^2 ds \leq \|(\sqrt{\rho} \operatorname{curl}^2 u, \operatorname{curl}^2 H)\|^2(0) + C \delta_1 \int_0^t \mathcal{D}_1^2(s) ds, \quad (3.37)$$

where $c > 0$, $C > 0$ are positive constants independent of t . Moreover,

$$\|\nabla^2 H\|^2 + \int_0^t \|\nabla^3 H\|^2 \leq C \mathcal{E}_1^2(0) + C \delta_1 \int_0^t \mathcal{D}_1^2(s) ds. \quad (3.38)$$

and

$$\|\nabla^2 u\|^2 + \int_0^t \|\nabla \operatorname{curl}^2 u\|^2 \leq C\mathcal{E}_1^2(0) + C\delta_1 \int_0^t \mathcal{D}_1^2(s)ds. \quad (3.39)$$

Proof. Firstly, integrating by parts with $w_t \times n|_{\partial\Omega} = 0$, it holds

$$\begin{aligned} \int_{\Omega} \rho w_t \cdot \operatorname{curl}^2 w dx &= \int_{\Omega} \operatorname{curl}(\rho w_t) \cdot \operatorname{curl} w dx \\ &= \frac{1}{2} \frac{d}{dt} \int_{\Omega} \rho |\operatorname{curl} w|^2 dx - \frac{1}{2} \int_{\Omega} q_t |\operatorname{curl} w|^2 dx + \int_{\Omega} (\nabla q \times w_t) \cdot \operatorname{curl} w dx. \end{aligned}$$

Taking (3.15) inner with $\operatorname{curl}^2 w$, and integrating over Ω , we obtain

$$\begin{aligned} &\frac{1}{2} \frac{d}{dt} \int_{\Omega} \rho |\operatorname{curl} w|^2 dx + \mu \|\operatorname{curl}^2 w\|^2 \\ &= \frac{1}{2} \int_{\Omega} q_t |\operatorname{curl} w|^2 dx - \int_{\Omega} (\nabla q \times w_t) \cdot \operatorname{curl} w dx + \int_{\Omega} \left(K + \operatorname{curl}(\operatorname{curl} H \times H) - \rho u \cdot w \right) \cdot \operatorname{curl}^2 w dx \\ &\leq \mathcal{E}_1(t) \mathcal{D}_1^2(t). \end{aligned} \quad (3.40)$$

Similarly, taking (3.18) inner with $\operatorname{curl}^2 \phi$ in $L^2(\Omega)$, one has

$$\begin{aligned} &\frac{1}{2} \frac{d}{dt} \int_{\Omega} |\operatorname{curl} \phi|^2 dx + \eta \|\operatorname{curl}^2 \phi\|^2 = \int_{\Omega} \operatorname{curl}^2(u \times H) \cdot \operatorname{curl}^2 \phi dx \\ &\leq \mathcal{E}_1(t) \mathcal{D}_1^2(t). \end{aligned} \quad (3.41)$$

Therefore, integrating (3.40) and (3.41) over $[0, t]$ gives (3.37).

Moreover, since $\operatorname{curl}^2 u \cdot n|_{\partial\Omega} = 0$, applying Lemma 2.2, it holds

$$\begin{aligned} \int_0^t \|\nabla \operatorname{curl}^2 u\|^2 ds &\leq C \int_0^t \left(\|\operatorname{curl}^3 u\|^2 + \|\operatorname{curl}^2 u\|^2 \right) ds \\ &\leq C\mathcal{E}_1^2(0) + C\delta_1 \int_0^t \mathcal{D}_1^2(s)ds, \end{aligned}$$

due to Lemma 3.5 and (3.37).

Noting that $\operatorname{curl}^2 H \cdot n|_{\partial\Omega} = 0$, applying elliptic estimate and Lemma 2.2 again, we has

$$\begin{aligned} \int_0^t \|\nabla^3 H\|^2 ds &\leq C \int_0^t \left(\|\Delta H\|_1^2 + \|\nabla H\|^2 \right) ds \\ &\leq C \int_0^t \left(\|\operatorname{curl}^2 H\|_1^2 + \|\nabla H\|^2 \right) ds \\ &\leq C \int_0^t \left(\|\operatorname{curl}^3 H\|^2 + \|\operatorname{curl}^2 H\|^2 + \|\nabla H\|^2 \right) ds \\ &\leq C\mathcal{E}_1^2(0) + C\delta_1 \int_0^t \mathcal{D}_1^2(s)ds, \end{aligned}$$

thanks to Lemma 3.2, Lemma 3.5 and (3.37).

Furthermore, applying elliptic estimate and the fact $\operatorname{div} H = 0$, it holds

$$\begin{aligned}\|\nabla^2 H\|^2 &\leq C\left(\|\Delta H\|^2 + \|\nabla H\|^2\right) \\ &\leq C\left(\|\operatorname{curl}^2 H\|^2 + \|\nabla H\|^2\right) \\ &\leq C\mathcal{E}_1^2(0) + C\delta_1 \int_0^t \mathcal{D}_1^2(s)ds,\end{aligned}$$

thanks to Lemma 3.3 and (3.37). Applying elliptic estimate again, it holds

$$\begin{aligned}\|\nabla^2 u\|^2 &\leq C(\|\Delta u\|^2 + \|\nabla u\|^2) \\ &\leq C(\|\operatorname{curl}^2 u\|^2 + \|\nabla \operatorname{div} u\|^2 + \|\nabla u\|^2) \\ &\leq C\mathcal{E}_1^2(0) + C\delta_1 \int_0^t \mathcal{D}_1^2(s)ds,\end{aligned}$$

thanks to Lemma 3.3, Lemma 3.10 and (3.37). □

Now, we prove the estimate of $\nabla^2 q$.

Lemma 3.12.

$$\|\nabla^2 q\|^2 + c \int_0^t \|\nabla^2 \operatorname{div} u\|^2 ds \leq C\mathcal{E}_1^2(0) + \varepsilon \int_0^t \|\nabla^2 q\|^2 ds + C\delta_1 \int_0^t \mathcal{D}_1^2(s)ds. \quad (3.42)$$

Proof. Taking ∂_{x_i} to (3.1)₂, multiplying the resulting identities by $\partial_{x_i} \nabla \operatorname{div} u$, integrating over Ω , we obtain

$$\begin{aligned}(2\mu + \lambda)\|\nabla^2 \operatorname{div} u\|^2 - R \int_{\Omega} (\nabla^2 q + \nabla^2 \theta) \cdot \nabla^2 \operatorname{div} u dx \\ = -\mu \int_{\Omega} \nabla \operatorname{curl}^2 u \cdot \nabla^2 \operatorname{div} u dx + \int_{\Omega} \left(\nabla(\rho(u_t + u \cdot \nabla u)) + R\nabla^2(q\theta) - \nabla(\operatorname{curl} H \times H) \right) \cdot \nabla^2 \operatorname{div} u dx.\end{aligned}$$

To eliminate the singular terms on the left-hand side of the above, we compute the integral

$$\int_{\Omega} \{ \partial_{x_i} \nabla(3.1)_1 \cdot R(\partial_{x_i} \nabla q + \partial_{x_i} \nabla \theta) \} dx$$

to obtain

$$\begin{aligned}\frac{R}{2} \frac{d}{dt} \int_{\Omega} |\nabla^2 q|^2 dx + R \frac{d}{dt} \int_{\Omega} \nabla^2 q \cdot \nabla^2 \theta dx + R \int_{\Omega} (\nabla^2 q + \nabla^2 \theta) \cdot \nabla^2 \operatorname{div} u dx \\ = R \int_{\Omega} \nabla^2 q \cdot \nabla^2 \theta_t dx - R \int_{\Omega} \nabla^2 \operatorname{div}(qu) \cdot (\nabla^2 q + \nabla^2 \theta) dx,\end{aligned}$$

Summing up the above yields

$$\begin{aligned}
& \frac{R}{2} \frac{d}{dt} \int_{\Omega} |\nabla^2 q|^2 dx + R \frac{d}{dt} \int_{\Omega} \nabla^2 q \cdot \nabla^2 \theta dx + (2\mu + \lambda) \|\nabla^2 \operatorname{div} u\|^2 \\
&= -\mu \int_{\Omega} \nabla \operatorname{curl}^2 u \cdot \nabla^2 \operatorname{div} u dx + \int_{\Omega} \left(\nabla(\rho(u_t + u \cdot \nabla u)) + R \nabla^2(q\theta) - \nabla(\operatorname{curl} H \times H) \right) \cdot \nabla^2 \operatorname{div} u dx \\
&+ R \int_{\Omega} \nabla^2 q \cdot \nabla^2 \theta_t dx - R \int_{\Omega} (\nabla^2 q + \nabla^2 \theta) \cdot \nabla^2 \operatorname{div}(qu) dx \\
&\equiv K_1 + K_2 + K_3 + K_4.
\end{aligned} \tag{3.43}$$

Using Cauchy's inequality,

$$K_1 \leq \frac{2\mu + \lambda}{4} \|\nabla^2 \operatorname{div} u\|^2 + C \|\nabla \operatorname{curl}^2 u\|^2, \tag{3.44}$$

$$\begin{aligned}
K_2 &\leq \frac{2\mu + \lambda}{4} \|\nabla^2 \operatorname{div} u\|^2 + C \left(\|\nabla(\rho(u_t + u \cdot \nabla u))\|^2 + \|\nabla^2(q\theta)\|^2 + \|\nabla(\operatorname{curl} H \times H)\|^2 \right) \\
&\leq \frac{2\mu + \lambda}{4} \|\nabla^2 \operatorname{div} u\|^2 + \left(\|(u, q, \theta, H)\|_2^2 \right) \left(\|\nabla^2 u\|^2 + \|\nabla u_t\|^2 + \|\nabla^2 q\|^2 + \|\nabla^2 \theta\|^2 + \|\nabla \operatorname{curl} H\|^2 \right) \\
&\leq \frac{2\mu + \lambda}{4} \|\nabla^2 \operatorname{div} u\|^2 + \delta \left(\|\nabla^2 u\|^2 + \|\nabla u_t\|^2 + \|\nabla^2 q\|^2 + \|\nabla^2 \theta\|^2 + \|\nabla \operatorname{curl} H\|^2 \right),
\end{aligned} \tag{3.45}$$

and

$$K_3 \leq \varepsilon \|\nabla^2 q\|^2 + C_\varepsilon \|\nabla^2 \theta_t\|^2. \tag{3.46}$$

The term K_4 becomes

$$K_4 = R \int_{\Omega} (\nabla^2 q + \nabla^2 \theta) \cdot \left(\nabla^2 q \operatorname{div} u + 2 \nabla q \cdot \nabla \operatorname{div} u + q \nabla^2 \operatorname{div} u + \nabla^2 u \cdot \nabla q + u \cdot \nabla^3 q + 2 \nabla u \cdot \nabla^2 q \right) dx$$

The term $\nabla^3 q$ including on K_4 , by using integration by parts with $u \cdot n|_{\partial\Omega} = 0$, we obtain

$$\begin{aligned}
& R \int_{\Omega} (\nabla^2 q + \nabla^2 \theta) u \cdot \nabla^3 q dx \\
&= -\frac{R}{2} \int_{\Omega} \operatorname{div} u |\nabla^2 q|^2 dx - R \int_{\Omega} \nabla^3 \theta u \cdot \nabla^2 q dx - R \int_{\Omega} \operatorname{div} u \nabla^2 \theta \cdot \nabla^2 q dx \\
&\leq \left(\|\operatorname{div} u\|_{L^\infty} + \|u\|_{L^\infty} \right) \cdot \left(\|\nabla^2 q\|^2 + \|\nabla^2 \theta\|^2 + \|\nabla^3 \theta\|^2 \right) \\
&\leq \delta \left(\|\nabla^2 q\|^2 + \|\nabla^2 \theta\|^2 + \|\nabla^3 \theta\|^2 \right).
\end{aligned}$$

While the other terms on K_4 is controlled by $\delta \left(\|\nabla^2 q\|^2 + \|\nabla^2 \theta\|^2 + \|\nabla^3 u\|^2 \right)$. So

$$K_4 \leq \delta \left(\|\nabla^2 q\|^2 + \|\nabla^2 \theta\|^2 + \|\nabla^3 \theta\|^2 + \|\nabla^3 u\|^2 \right). \tag{3.47}$$

Putting estimates (3.44)-(3.47) into (3.43) implies

$$\begin{aligned}
& \frac{R}{2} \frac{d}{dt} \int_{\Omega} |\nabla^2 q|^2 dx + R \frac{d}{dt} \int_{\Omega} \nabla^2 q \cdot \nabla^2 \theta dx + \frac{2\mu + \lambda}{2} \|\nabla^2 \operatorname{div} u\|^2 \\
&\leq C \|\nabla \operatorname{curl}^2 u\|^2 + \varepsilon \|\nabla^2 q\|^2 + C_\varepsilon \|\nabla^2 \theta_t\|^2 + \delta_1 \left(\|\nabla^2 q\|^2 + \|\nabla^2 \theta\|^2 + \|\nabla^3 \theta\|^2 + \|\nabla^3 u\|^2 \right).
\end{aligned} \tag{3.48}$$

Integrating the above over $[0, t]$ yields

$$\begin{aligned} \frac{R}{4} \|\nabla^2 q\|^2 + \frac{2\mu + \lambda}{2} \int_0^t \|\nabla^2 \operatorname{div} u\|^2 ds &\leq C \|\nabla^2 \theta\|^2 + \varepsilon \int_0^t \|\nabla^2 q\|^2 ds \\ &+ C \int_0^t \left(\|\nabla \operatorname{curl}^2 u\|^2 + \|\nabla^2 \theta_t\|^2 \right) ds + \delta_1 \int_0^t \left(\|\nabla^2 q\|^2 + \|\nabla^2 \theta\|^2 + \|\nabla^3 \theta\|^2 + \|\nabla^3 u\|^2 \right) ds, \end{aligned}$$

which together (3.33), (3.23) and (3.39) implies the desired result (3.42). $\square$

At last, with Lemma 3.2-Lemma 3.12 at hand, we can use Lemma 2.4 to obtain the energy estimates of $\|(\nabla^3 u, \nabla^3 \theta, \nabla^3 H)\|$. Moreover, we complete the proof of Proposition 3.1 by adding Lemma 3.2-Lemma 3.12, and using equation (3.1)₂ to obtain the dissipation estimates of $\|\nabla^2 u\|_1^2 + \|\nabla q\|_1^2$ by choosing ε small in Lemma 3.12. Here, we omit the detail.

4. SMOOTH SOLUTION

In this section, we shall prove Theorem 1.3, that is to improve the regularity of the strong solutions in Theorem 1.1. Let

$$\mathcal{E}(t) = \|(u, \theta, H)\|_4 + \|q\|_3 + \|(q_t, u_t, H_t, \theta_t)\|_2 + \|(q_{tt}, u_{tt}, \theta_{tt}, H_{tt})\| \quad (4.1)$$

and

$$\mathcal{D}(t) = \|(\nabla u, \nabla \theta, \nabla H)\|_3 + \|\nabla q\|_2 + \|(q_t, u_t, H_t)\|_2 + \|\theta_t\|_3 + \|(u_{tt}, \theta_{tt}, H_{tt})\|_1 + \|q_{tt}\|. \quad (4.2)$$

We show a priori estimates of H^4 norm in the following.

Proposition 4.1. (a priori estimates of H^4 norm) *Let (q, u, θ, H) be a solution to the initial boundary value problem (3.1), (3.2) and (3.3) in $t \in [0, T]$. Then there exists positive constants C and δ which are independent of t , such that if*

$$\sup_{0 \leq t \leq T} \mathcal{E}(t) \leq \delta,$$

then there holds, for any $t \in [0, T]$,

$$\mathcal{E}^2(t) + C \int_0^t \mathcal{D}^2(s) ds \leq C \mathcal{E}^2(0).$$

We shall divide the proof of Proposition 4.1 into several lemmas. Firstly, with the results in Proposition 3.1 and using elliptic estimates, one can obtain the following dissipation estimates.

Lemma 4.2.

$$\int_0^t \|\nabla^2 \operatorname{curl}^2 u\|^2 ds \leq C \mathcal{E}_1^2(0), \quad (4.3)$$

$$\int_0^t \|\nabla^4 \theta\|^2 ds \leq C \mathcal{E}_1^2(0), \quad (4.4)$$

and

$$\int_0^t \|\nabla^4 H\|^2 ds \leq C \mathcal{E}_1^2(0). \quad (4.5)$$

Proof. Applying Lemma 2.2 with $v = \operatorname{curl}^2 u$,

$$\begin{aligned} \|\nabla^2 \operatorname{curl}^2 u\|^2 &\leq \|\operatorname{curl}^3 u\|_1^2 + \|\operatorname{curl}^2 u\|^2 + |\operatorname{curl}^2 u \cdot n|_{H^{3/2}(\partial\Omega)}^2 \\ &\leq \|\operatorname{curl}^3 u\|_1^2 + \|\operatorname{curl}^2 u\|^2, \end{aligned} \quad (4.6)$$

thanks to $\operatorname{curl}^2 u \cdot n|_{\partial\Omega} = 0$. Moreover, noting that

$$\|\operatorname{curl}^3 u\|_1^2 = \|\operatorname{curl}^2 w\|_1^2 = \|\Delta w\|_1^2,$$

and using the equation of w (3.15), we obtain

$$\begin{aligned} \|\Delta w\|_1^2 &\leq \|\rho w_t + \rho(u \cdot \nabla w - w \cdot \nabla u) + \nabla q \times (u_t + u \cdot \nabla u) + \rho w \operatorname{div} u - H \cdot \nabla \operatorname{curl} H + \operatorname{curl} H \cdot \nabla H\|_1^2 \\ &\leq C\|w_t\|_1^2 + \left(\|\nabla q\|_1^2 + \|u\|_3^2 + \|H\|_2^2\right) \cdot \left(\|\nabla w_t\|^2 + \|\nabla w\|_1^2 + \|u_t\|_2^2 + \|\nabla u\|_2^2 + \|\nabla H\|_2^2\right). \end{aligned}$$

Plugging the above back in (4.6), one gets

$$\begin{aligned} \int_0^t \|\nabla^2 \operatorname{curl}^2 u\|^2 ds &\leq C \int_0^t \left(\|w_t\|_1^2 + \|\operatorname{curl}^2 u\|^2\right) ds + \delta \int_0^t \left(\|\nabla w_t\|^2 + \|\nabla w\|_1^2 + \|u_t\|_2^2 + \|\nabla u\|_2^2 + \|\nabla H\|_2^2\right) ds \\ &\leq (C + \delta) \int_0^t \mathcal{D}_1^2(s) ds \leq C\mathcal{E}_1^2(0). \end{aligned}$$

For magnetic field H , applying Lemma 2.4 and using equation (3.1)₄, we get

$$\begin{aligned} \int_0^t \|\nabla^4 H\|^2 ds &\leq \int_0^t \left(\|\Delta H\|_2^2 + \|\nabla H\|^2\right) ds \\ &\leq \int_0^t \left(\|H_t\|_2^2 + \|\operatorname{curl}(u \times H)\|_2^2 + \|\nabla H\|^2\right) ds \\ &\leq \int_0^t \left(\|H_t\|_2^2 + \|H\|_2^2 \|\nabla u\|_2^2 + \|u\|_2^2 \|\nabla H\|_2^2 + \|\nabla H\|^2\right) ds \\ &\leq (C + \delta_1) \int_0^t \mathcal{D}_1^2(s) ds \leq C\mathcal{E}_1^2(0). \end{aligned}$$

Finally, we estimate the four-order derivatives of θ . Rewrite equation (3.1)₂ as

$$\kappa \Delta \theta = c_v \rho(\theta_t + u \cdot \nabla \theta) + R \operatorname{div} u - \lambda(\operatorname{div} u)^2 - 2\mu|S(u)|^2 + R(\rho\theta + q) \operatorname{div} u - \eta|\operatorname{curl} H|^2 \equiv N, \quad (4.7)$$

then, applying the elliptic estimates of θ , θ_t (Lemma 2.6), it is easy to obtain the estimates of $\|\nabla^4 \theta\|_{L_t^2(L^2)}$ and $\|\nabla^3 \theta_t\|_{L_t^2(L^2)}$. Precisely,

$$\begin{aligned} \int_0^t \|\nabla^4 \theta\|^2 ds &\leq \int_0^t \|N\|_2^2 ds + \int_0^t \|\nabla \theta\|^2 ds \\ &\leq (C + \delta) \int_0^t \mathcal{D}_1^2(s) ds \leq C\mathcal{E}_1^2(0). \end{aligned}$$

The proof of Lemma 4.2 is completed. □

Next, we prove estimates of the second-order temporal derivatives $(q_{tt}, u_{tt}, \theta_{tt}, H_{tt})$.

Lemma 4.3.

$$\|(q_{tt}, u_{tt}, \theta_{tt}, H_{tt})\|^2 + c \int_0^t \|(\nabla u_{tt}, \nabla \theta_{tt}, \nabla H_{tt})\|^2 ds \leq C\mathcal{E}^2(0). \quad (4.8)$$

where $c > 0$, $C > 0$ are positive constants independent of t . Moreover,

$$\|(u_t, \theta_t, H_t)\|_2^2 \leq C\mathcal{E}^2(0), \quad (4.9)$$

and

$$\int_0^t \|\nabla^3 \theta_t\|^2 ds \leq C\mathcal{E}^2(0). \quad (4.10)$$

Proof. Computing the following integral

$$\int_{\Omega} \{ \partial_{tt}(3.1)_1 R q_{tt} + \partial_{tt}(3.1)_2 \cdot u_{tt} + \partial_{tt}(3.1)_3 \theta_{tt} + \partial_{tt}(3.1)_4 \cdot H_{tt} \} dx,$$

and noting that differentiation of the system (3.1) with respect to t will keep the boundary conditions (3.3), after integrating by parts, one has

$$\begin{aligned} & \frac{1}{2} \frac{d}{dt} \int_{\Omega} \left(R q_{tt}^2 + \rho |u_{tt}|^2 + c_v \rho \theta_{tt}^2 + |H_{tt}|^2 \right) dx + \mu \|\operatorname{curl} u_{tt}\|^2 + (2\mu + \lambda) \|\operatorname{div} u_{tt}\|^2 + \kappa \|\nabla \theta_{tt}\|^2 + \eta \|\operatorname{curl} H_{tt}\|^2 \\ &= -R \int_{\Omega} (\operatorname{div}(qu))_{tt} q_{tt} dx - \int_{\Omega} \left(q_{tt}(u_t + u \cdot \nabla u) + 2q_t(u_t + u \cdot \nabla u)_t - \rho u_{tt} \cdot \nabla u + 2\rho u_t \cdot \nabla u_t \right) \cdot u_{tt} dx \\ &- R \int_{\Omega} (\nabla(q\theta))_{tt} \cdot u_{tt} dx + \int_{\Omega} (\operatorname{curl} H \times H)_{tt} \cdot u_{tt} dx \\ &- c_v \int_{\Omega} \left(q_{tt}(\theta_t + u \cdot \nabla \theta) + 2q_t(\theta_t + u \cdot \nabla \theta)_t - \rho u_{tt} \cdot \nabla \theta + 2\rho \theta_t \cdot \nabla \theta_t \right) \theta_{tt} dx \\ &+ \int_{\Omega} \left(\lambda (\operatorname{div} u)^2 + 2\mu |S(u)|^2 - R(\rho \theta + q) \operatorname{div} u + \eta |\operatorname{curl} H|^2 \right)_{tt} \theta_{tt} dx \\ &+ \int_{\Omega} (\operatorname{curl}(u \times H))_{tt} \cdot H_{tt} dx. \end{aligned} \tag{4.11}$$

The term involving third-order derivations of q can be estimated as follows:

$$\begin{aligned} & -R \int_{\Omega} (\nabla(q\theta))_{tt} \cdot u_{tt} dx = R \int_{\Omega} (q\theta)_{tt} \cdot \operatorname{div} u_{tt} dx \\ & \leq \frac{2\mu + \lambda}{4} \|\operatorname{div} u_{tt}\|^2 + C \|(q\theta)_{tt}\|^2 \\ & \leq \frac{2\mu + \lambda}{4} \|\operatorname{div} u_{tt}\|^2 + \left(\|(\theta, q)\|_2 + \|q_t\|_1 \right) \left(\|q_{tt}\|^2 + \|\theta_{tt}\|^2 + \|\nabla \theta_t\|^2 \right) \\ & \leq \frac{2\mu + \lambda}{4} \|\operatorname{div} u_{tt}\|^2 + C \delta_1 \mathcal{D}_1^2(t), \end{aligned} \tag{4.12}$$

and

$$\begin{aligned} & \int_{\Omega} (\operatorname{div}(qu))_{tt} q_{tt} dx = \int_{\Omega} u \cdot \nabla q_{tt} q_{tt} dx + \int_{\Omega} (\operatorname{div} u_{tt} q + \operatorname{div} u_t q_t + \operatorname{div} u q_{tt}) q_{tt} dx \\ &= -\frac{1}{2} \int_{\Omega} \operatorname{div} u q_{tt} dx + \int_{\Omega} (\operatorname{div} u_{tt} q + \operatorname{div} u_t q_t + \operatorname{div} u q_{tt}) q_{tt} dx \\ &\leq \frac{2\mu + \lambda}{4} \|\operatorname{div} u_{tt}\|^2 + \left(\|\operatorname{div} u\|_{L^\infty} + \|q\|_{L^\infty} + \|q_t\|_{L^3} \right) \left(\|\operatorname{div} u_{tt}\|^2 + \|\nabla \operatorname{div} u_t\|^2 + \|q_{tt}\|^2 \right) \\ &\leq \frac{2\mu + \lambda}{4} \|\operatorname{div} u_{tt}\|^2 + C \delta_1 \mathcal{D}_1^2(t). \end{aligned} \tag{4.13}$$

Noting that

$$(u \times H) \times n|_{\partial\Omega} = 0,$$

due to the vector identity $(u \times H) \times n = (u \cdot n)H - (H \cdot n)u$ and $u \cdot n|_{\partial\Omega} = 0$, $H \times n|_{\partial\Omega} = 0$, and moreover, differentiation with respect t will keep the boundary conditions, it holds

$$(u \times H)_{tt} \times n|_{\partial\Omega} = 0.$$

Then the last term on the right hand-side on (4.11) becomes

$$\begin{aligned}
\int_{\Omega} (\operatorname{curl}(u \times H))_{tt} \cdot H_{tt} dx &= \int_{\Omega} (u \times H)_{tt} \cdot \operatorname{curl} H_{tt} dx \\
&\leq \frac{\eta}{4} \|\operatorname{curl} H_{tt}\|^2 + C \|(u \times H)_{tt}\|^2 \\
&\leq \frac{\eta}{4} \|\operatorname{curl} H_{tt}\|^2 + C \delta_1 \mathcal{D}_1^2(t).
\end{aligned} \tag{4.14}$$

The other terms on the right hand-side on (4.11) can be estimated by $\delta_1 \mathcal{D}_1^2(t)$ by using Holder inequality and Sobolev inequalities directly. Therefore, (4.11) yields

$$\begin{aligned}
&\frac{1}{2} \frac{d}{dt} \int_{\Omega} \left(R q_{tt}^2 + \rho |u_{tt}|^2 + c_v \rho \theta_{tt}^2 + |H_{tt}|^2 \right) dx \\
&+ \mu \|\operatorname{curl} u_{tt}\|^2 + \frac{2\mu + \lambda}{2} \|\operatorname{div} u_{tt}\|^2 + \kappa \|\nabla \theta_{tt}\|^2 + \frac{\eta}{2} \|\operatorname{curl} H_{tt}\|^2 \\
&\leq C \delta_1 \mathcal{D}_1^2(t).
\end{aligned} \tag{4.15}$$

Finally, integrating (4.15) over $[0, t]$ and applying Proposition 2.3 to u_{tt} and H_{tt} , we obtain

$$\begin{aligned}
&\|(q_{tt}, u_{tt}, \theta_{tt}, H_{tt})\|^2 + c_{11} \int_0^t \|(\nabla u_{tt}, \nabla \theta_{tt}, \nabla H_{tt})\|^2 ds \\
&\leq C \|(q_{tt}, u_{tt}, \theta_{tt}, H_{tt})(0)\|^2 + C \delta_1 \int_0^t \mathcal{D}_1^2(s) ds \\
&\leq C \mathcal{E}^2(0),
\end{aligned}$$

where we have used Proposition 3.1 in the last inequality.

Furthermore, applying Lemma 2.4 to u_t , θ_t and H_t , it is easy to see (4.9).

Moreover, recalling (4.7), we see that

$$\begin{aligned}
\int_0^t \|\nabla^3 \theta_t\|^2 ds &\leq \int_0^t \|N_t\|_1^2 ds + \int_0^t \|\nabla \theta_t\|^2 ds \\
&\leq \int_0^t \|\nabla \theta_{tt}\|^2 ds + (C + \delta) \int_0^t \mathcal{D}_1^2(s) ds \\
&\leq C \mathcal{E}^2(0).
\end{aligned}$$

This complete the proof of Lemma 4.3. □

In order to obtain the estimate of $\|\nabla^2 q_t\|_{L_t^2(L^2)}$, we use the equation $q_t + \operatorname{div} u + \operatorname{div}(qu) = 0$, we shall prove $\|\nabla^3 q\|_{L_t^2(L^2)}$ and $\|\nabla^2 \operatorname{div} u\|_{L_t^2(L^2)}$ (see Lemma 3.12). To obtain $\|\nabla^3 q\|_{L_t^2(L^2)}$, by using the elliptic estimate of Stokes-type equation (3.1)₂. However, applying Lemma 2.5 to control $\|\nabla^3 q\|_{L_t^2(L^2)}$, we shall estimate $\|\operatorname{div} u\|_{L_t^2(H^3)}$ firstly.

Lemma 4.4.

$$\frac{R}{4} \|\nabla^3 q\|^2 + \frac{2\mu + \lambda}{2} \int_0^t \|\nabla^3 \operatorname{div} u\|^2 ds \leq C \mathcal{E}^2(0) + \varepsilon \int_0^t \|\nabla^3 q\|^2 ds + \delta \int_0^t \left(\|\nabla^3 q\|^2 + \|\nabla^4 u\|^2 \right) ds. \tag{4.16}$$

Proof. Taking $\partial_{x_i}\partial_{x_j}$ to (3.1)₂, multiplying the resulting identities by $\partial_{x_i}\partial_{x_j}\nabla\text{div}u$, integrating over Ω , we obtain

$$\begin{aligned} & (2\mu + \lambda)\|\nabla^3\text{div}u\|^2 - R \int_{\Omega} (\nabla^3q + \nabla^3\theta) \cdot \nabla^3\text{div}u dx \\ &= -\mu \int_{\Omega} \nabla^2\text{curl}^2u \cdot \nabla^3\text{div}u dx + \int_{\Omega} \left(\nabla^2(\rho(u_t + u \cdot \nabla u)) + R\nabla^2(q\theta) - \nabla(H \cdot \nabla H - \frac{1}{2}|\nabla H|^2) \right) \cdot \nabla^3\text{div}u dx \end{aligned}$$

Computing the integral $\int_{\Omega} \{ \partial_{x_i}\partial_{x_j}\nabla(3.1)_1 \cdot R(\partial_{x_i}\partial_{x_j}\nabla q + \partial_{x_i}\partial_{x_j}\nabla\theta) \} dx$, one has

$$\begin{aligned} & \frac{R}{2} \frac{d}{dt} \int_{\Omega} |\nabla^3q|^2 dx + R \frac{d}{dt} \int_{\Omega} \nabla^3q \cdot \nabla^3\theta dx + R \int_{\Omega} (\nabla^3q + \nabla^3\theta) \cdot \nabla^3\text{div}u dx \\ &= R \int_{\Omega} \nabla^3q \cdot \nabla^3\theta_t dx - R \int_{\Omega} \nabla^3\text{div}(qu) \cdot (\nabla^3q + \nabla^3\theta) dx, \end{aligned}$$

Summing up the above yields

$$\begin{aligned} & \frac{R}{2} \frac{d}{dt} \int_{\Omega} |\nabla^3q|^2 dx + R \frac{d}{dt} \int_{\Omega} \nabla^3q \cdot \nabla^3\theta dx + (2\mu + \lambda)\|\nabla^3\text{div}u\|^2 \\ &= -\mu \int_{\Omega} \nabla^2\text{curl}^2u \cdot \nabla^3\text{div}u dx + \int_{\Omega} \left(\nabla^2(\rho(u_t + u \cdot \nabla u)) + R\nabla^3(q\theta) - \nabla^2(\text{curl}H \times H) \right) \cdot \nabla^3\text{div}u dx \\ &+ R \int_{\Omega} \nabla^3q \cdot \nabla^3\theta_t dx - R \int_{\Omega} (\nabla^3q + \nabla^3\theta) \cdot \nabla^3\text{div}(qu) dx \equiv J_1 + J_2 + J_3 + J_4. \end{aligned} \tag{4.17}$$

Using Cauchy's inequality,

$$J_1 \leq \frac{2\mu + \lambda}{4} \|\nabla^3\text{div}u\|^2 + C \|\nabla^2\text{curl}^2u\|^2, \tag{4.18}$$

$$\begin{aligned} J_2 &\leq \frac{2\mu + \lambda}{4} \|\nabla^3\text{div}u\|^2 + C \left(\|\nabla^2(\rho(u_t + u \cdot \nabla u))\|^2 + \|\nabla^3(q\theta)\|^2 + \|\nabla^2(\text{curl}H \times H)\|^2 \right) \\ &\leq \frac{2\mu + \lambda}{4} \|\nabla^3\text{div}u\|^2 + \delta \left(\|\nabla q\|_1^2 + \|\nabla^2u_t\|^2 + \|\nabla^2u\|_1^2 + \|\nabla^3q\|^2 + \|\nabla^3\theta\|^2 + \|\nabla^3H\|^2 \right) \end{aligned} \tag{4.19}$$

and

$$J_3 \leq \varepsilon \|\nabla^3q\|^2 + C_\varepsilon \|\nabla^3\theta_t\|^2. \tag{4.20}$$

The term J_4 becomes

$$\begin{aligned} J_4 &= R \int_{\Omega} (\nabla^3q + \nabla^3\theta) \cdot \left(\nabla^3q\text{div}u + 3\nabla^2q \cdot \nabla\text{div}u + 3\nabla q \cdot \nabla^2\text{div}u + q\nabla^3\text{div}u \right. \\ &\quad \left. + \nabla^3u \cdot \nabla q + 3\nabla^2u \cdot \nabla^2q + 3\nabla u \cdot \nabla^3q + u \cdot \nabla^4q \right) dx \end{aligned}$$

The term $\nabla^4 q$ including on J_4 , by using integration by parts with $u \cdot n|_{\partial\Omega} = 0$, we obtain

$$\begin{aligned}
& R \int_{\Omega} (\nabla^3 q + \nabla^3 \theta) u \cdot \nabla^4 q dx \\
&= -\frac{R}{2} \int_{\Omega} \operatorname{div} u |\nabla^3 q|^2 dx - R \int_{\Omega} \nabla^4 \theta u \cdot \nabla^3 q dx - R \int_{\Omega} \operatorname{div} u \nabla^3 \theta \cdot \nabla^3 q dx \\
&\leq \left(\|\operatorname{div} u\|_{L^\infty} + \|u\|_{L^\infty} \right) \cdot \left(\|\nabla^3 q\|^2 + \|\nabla^3 \theta\|^2 + \|\nabla^4 \theta\|^2 \right) \\
&\leq \delta \left(\|\nabla^3 q\|^2 + \|\nabla^3 \theta\|^2 + \|\nabla^4 \theta\|^2 \right).
\end{aligned}$$

While the other terms on J_4 is controlled by $\delta \left(\|\nabla^3 q\|^2 + \|\nabla^3 \theta\|^2 + \|\nabla^4 u\|^2 \right)$. So

$$J_4 \leq \delta \left(\|\nabla^3 q\|^2 + \|\nabla^3 \theta\|^2 + \|\nabla^4 \theta\|^2 + \|\nabla^4 u\|^2 \right). \quad (4.21)$$

Putting estimates (4.18)-(4.21) into (4.17) implies

$$\begin{aligned}
& \frac{R}{2} \frac{d}{dt} \int_{\Omega} |\nabla^3 q|^2 dx + R \frac{d}{dt} \int_{\Omega} \nabla^3 q \cdot \nabla^3 \theta dx + \frac{2\mu + \lambda}{2} \|\nabla^3 \operatorname{div} u\|^2 \\
&\leq C \|\nabla^2 \operatorname{curl}^2 u\|^2 + \varepsilon \|\nabla^3 q\|^2 + C_\varepsilon \|\nabla^3 \theta_t\|^2 \\
&\quad + \delta \left(\|\nabla^3 q\|^2 + \|\nabla q\|_1^2 + \|\nabla^2 u\|_1^2 + \|\nabla^3 \theta\|^2 + \|\nabla^4 \theta\|^2 + \|\nabla^4 u\|^2 + \|\nabla^2 u_t\|^2 + \|\nabla^3 H\|^2 \right).
\end{aligned}$$

Then, integrating over $[0, t]$, and using Cauchy inequality and Proposition 3.1, Lemma 4.2 implies the desired estimate (4.16). $\square$

Finally, we use Lemma 2.4 to obtain the energy estimates $\|(\nabla^4 u, \nabla^4 \theta, \nabla^4 H)\|$. On the other hand, we apply the elliptic estimates of Stokes-type systems (Lemma 2.5) to obtain the estimates of $(\|\nabla^3 q\| + \|\nabla^4 u\|)_{L_t^2(L^2)}$. Therefore, the high-order energy is closed, i.e. Proposition 4.1 is proved.

Proof of Proposition 4.1 Applying stand elliptic estimates (i.e. Lemma 2.4) to u , θ and H , it is easy to get the energy estimates $\|(\nabla^4 u, \nabla^4 \theta, \nabla^4 H)\|$.

Next, rewriting (3.1)₂ as the following Stokes-type systems

$$\begin{cases} \operatorname{div} u = \operatorname{div} u, \\ -\mu \Delta u + R \nabla q = -\rho(u_t + u \cdot \nabla u) + (\mu + \lambda) \nabla \operatorname{div} u - R \nabla \theta - R \nabla(q\theta) + \operatorname{curl} H \times H \equiv f \end{cases}$$

Applying Lemma 2.5 to the above Stokes-type systems, one has

$$\begin{aligned}
\int_0^t \left(\|\nabla^2 u\|_2^2 + \|\nabla q\|_2^2 \right) ds &\leq C \left(\int_0^t \|f\|_2^2 ds + \int_0^t \|\operatorname{div} u\|_3^2 ds \right) \\
&\leq (C + \delta) \int_0^t \mathcal{D}_1^2(s) ds + C_* \int_0^t \|\nabla^3 \operatorname{div} u\|^2 ds \\
&\leq C \mathcal{E}_1^2(0) + C_* \int_0^t \|\nabla^3 \operatorname{div} u\|^2 ds,
\end{aligned} \quad (4.22)$$

where we have used the result of Proposition 3.1. Choosing ε small in Lemma 4.4, such that $\varepsilon C_* \leq \frac{2\mu + \lambda}{4}$, then, form (4.22), we have

$$\frac{R}{4} \|\nabla^3 q\|^2 + \frac{2\mu + \lambda}{4} \int_0^t \|\nabla^3 \operatorname{div} u\|^2 ds \leq C \mathcal{E}^2(0) + \delta \int_0^t \left(\|\nabla^3 q\|^2 + \|\nabla^4 u\|^2 \right) ds. \quad (4.23)$$

Therefore, together (4.22) with (4.23), for δ small,

$$\int_0^t \left(\|\nabla^2 u\|_2^2 + \|\nabla q\|_2^2 \right) ds \leq C\mathcal{E}^2(0).$$

So, we close the energy estimates. This complete the proof of Proposition 4.1. $\square$

5. DECAY RATE

For the solution (q, u, θ, H) of Theorem 1.3, one has

$$\int_0^\infty \|(\nabla q, \nabla u, \nabla \theta, \nabla H)\|_1^2 dt < \infty, \quad \int_0^\infty \left| \frac{d}{dt} \|(\nabla q, \nabla u, \nabla \theta, \nabla H)\|_1^2 \right| dt < \infty$$

where $q = \rho - 1$, $\theta = \mathcal{T} - 1$, it yields that

$$\|(\nabla q, \nabla u, \nabla \theta, \text{curl} H)\|_1 \rightarrow 0, \quad \text{as } t \rightarrow \infty. \quad (5.1)$$

In this section, we will prove Theorem 1.4 by the the following lemmas. Let

$$X(0, \infty; \mathcal{E}_0) := \{(q, u, \theta, H) \mid q \in C^0(0, \infty, H^3(\Omega)) \cap C^1(0, \infty, H^2(\Omega)),$$

$$(u, \theta, H) \in C^0(0, \infty, H^4(\Omega)) \cap C^1(0, \infty, H^2(\Omega)), \text{ and}$$

$$\mathcal{E}^2(t) + \int_0^t \mathcal{D}^2(s) ds \leq C\mathcal{E}_0^2\}$$

where $\mathcal{E}$ and $\mathcal{D}$ is same as (4.1) and (4.2). It is observed that the solution of IBVP (3.1)-(3.3) is sought in the space $X(0, \infty, \mathcal{E}_0)$, where $0 \leq \mathcal{E}_0 \leq C\delta$ and δ is given in Theorem 1.3.

The first and most important key point to prove Theorem 1.4 is to obtain the large-time behavior of the time derivatives $(q_t, u_t, \theta_t, H_t)$.

Lemma 5.1. *For every solution $(q, u, \theta, H) \in X(0, \infty; \mathcal{E}_0)$ of the problem (3.1)-(3.3), it exists a time $T_1 > 0$, depending on (q, u, θ, H) , such that*

$$\|(q_t, u_t, \theta_t, H_t)\| \leq Ct^{-1/2} \quad \forall t \geq T_1. \quad (5.2)$$

Proof. Differentiate (3.1)₁ with respect to t , multiply the resulting identity by Rq_t and integrate over Ω , it follows

$$\begin{aligned} & \frac{R}{2} \frac{d}{dt} \int_\Omega q_t^2 dx + R \int_\Omega \text{div} u_t q_t dx = -R \int_\Omega (\text{div}(qu))_t q_t dx \\ & = -R \int_\Omega q_t u_t \cdot \nabla q dx - R \int_\Omega q_t u \cdot \nabla q_t dx - R \int_\Omega q_t^2 \text{div} u dx - R \int_\Omega q q_t \text{div} u_t dx \\ & \equiv \sum_{j=1}^4 J_j. \end{aligned} \quad (5.3)$$

Utilizing Lemma 2.1 and Hölder's inequality, for $\epsilon > 0$ we deduce that

$$|J_1| \leq \|u_t\|_{L^6} \|\nabla q\|_{L^3} \|q_t\| \leq \epsilon \|\nabla u_t\|^2 + C_\epsilon \|\nabla q\|_1^2 \|q_t\|^2,$$

$$|J_4| \leq \|\text{div} u_t\| \|q\|_{L^\infty} \|q_t\| \leq \epsilon \|\nabla u_t\|^2 + C_\epsilon \|\nabla q\|_1^2 \|q_t\|^2.$$

Integration by parts with boundary condition $u \cdot n|_{\partial\Omega} = 0$ yields

$$J_2 = \frac{R}{2} \int_\Omega q_t^2 \text{div} u dx,$$

which implies that

$$J_2 + J_3 = -\frac{R}{2} \int_\Omega q_t^2 \text{div} u dx.$$

In view of equation (3.1)₁, it holds out that

$$\begin{aligned}
J_2 + J_3 &= \frac{R}{2} \int_{\Omega} q_t^2 (q_t + q \operatorname{div} u + u \cdot \nabla q) dx \\
&= \frac{R}{2} \frac{d}{dt} \int_{\Omega} q q_t^2 dx - R \int_{\Omega} q q_t q_{tt} dx + \frac{R}{2} \int_{\Omega} q_t^2 q \operatorname{div} u dx + \frac{R}{2} \int_{\Omega} q_t^2 u \cdot \nabla q dx \\
&\leq \frac{R}{2} \frac{d}{dt} \int_{\Omega} q q_t^2 dx - R \int_{\Omega} q q_t q_{tt} dx + \|(\nabla u, \nabla q)\|_2^2 \|q_t\|^2.
\end{aligned}$$

For the term $\int_{\Omega} q q_t q_{tt} dx$, using (3.1)₁ again, one has

$$\begin{aligned}
-R \int_{\Omega} q q_t q_{tt} dx &= R \int_{\Omega} q q_t (\operatorname{div} u + \operatorname{div}(qu))_t dx \\
&\leq \varepsilon \|\nabla u_t\|^2 + C_{\varepsilon} \|(\nabla u, \nabla q)\|_2^2 \|q_t\|^2.
\end{aligned}$$

Plugging all above inequalities into (5.3), we obtain

$$\frac{R}{2} \frac{d}{dt} \int_{\Omega} q_t^2 dx - \frac{R}{2} \frac{d}{dt} \int_{\Omega} q q_t^2 dx + R \int_{\Omega} \operatorname{div} u_t q_t dx \leq \varepsilon \|\nabla u_t\|^2 + C_{\varepsilon} \|(\nabla q, \nabla u)\|_2^2 \|q_t\|^2. \quad (5.4)$$

In a similar way we get the following inequalities. From (3.1)₂, we have

$$\begin{aligned}
&\frac{1}{2} \frac{d}{dt} \int_{\Omega} \rho |u_t|^2 dx + R \int_{\Omega} u_t \cdot \nabla q_t dx + R \int_{\Omega} u_t \cdot \nabla \theta_t dx + \mu \|\operatorname{curl} u_t\|^2 + (2\mu + \lambda) \|\operatorname{div} u_t\|^2 \\
&\leq (\varepsilon + \|\nabla u\|_1^2) (\|\nabla u_t\|^2 + \|\operatorname{curl} H_t\|^2) \\
&\quad + C_{\varepsilon} (\|u_t\|_1^2 + \|(\nabla q, \nabla u, \nabla \theta)\|_1^2 + \|\operatorname{curl} H\|_2^2) \cdot (\|q_t\|^2 + \|u_t\|^2 + \|\theta_t\|^2 + \|H_t\|^2).
\end{aligned} \quad (5.5)$$

From (3.1)₃, one has

$$\begin{aligned}
&\frac{c_v}{2} \frac{d}{dt} \int_{\Omega} \rho |\theta_t|^2 dx + R \int_{\Omega} \theta_t \operatorname{div} u_t dx + \kappa \|\nabla \theta_t\|^2 \\
&\leq (\varepsilon + \|\nabla u\|_1^2) (\|\nabla \theta_t\|^2 + \|\nabla u_t\|^2 + \|\operatorname{curl} H_t\|^2) \\
&\quad + C_{\varepsilon} (\|\theta_t\|_1^2 + \|(\nabla q, \nabla \theta)\|_1^2 + \|(\nabla u, \operatorname{curl} H)\|_2^2) \cdot (\|q_t\|^2 + \|u_t\|^2 + \|\theta_t\|^2).
\end{aligned} \quad (5.6)$$

And from (3.1)₄, it gives that

$$\frac{1}{2} \frac{d}{dt} \int_{\Omega} |H_t|^2 dx + \eta \|\operatorname{curl} H_t\|^2 \leq \varepsilon (\|\nabla u_t\|^2 + \|\operatorname{curl} H_t\|^2) + C_{\varepsilon} \|(\nabla u, \operatorname{curl} H)\|_1^2 \|H_t\|^2. \quad (5.7)$$

Summing up the estimates (5.4)-(5.7), and using Proposition 2.3, it implies that

$$\begin{aligned}
&\frac{1}{2} \frac{d}{dt} \{R \|q_t\|^2 + \|\sqrt{\rho} u_t\|^2 + c_v \|\sqrt{\rho} \theta_t\|^2 + \|H_t\|^2\} - \frac{R}{2} \frac{d}{dt} \int_{\Omega} q q_t^2 dx \\
&\quad + c_{*} \|\nabla u_t\|^2 + \kappa \|\nabla \theta_t\|^2 + \eta \|\operatorname{curl} H_t\|^2 \\
&\leq (\varepsilon + \|\nabla u\|_1^2) (\|\nabla \theta_t\|^2 + \|\nabla u_t\|^2 + \|\operatorname{curl} H_t\|^2) \\
&\quad + C_{\varepsilon} (\|u_t\|_1^2 + \|\theta_t\|_1^2 + \|(\nabla q, \nabla u, \operatorname{curl} H)\|_2^2 + \|\nabla \theta\|_1^2) \cdot (\|q_t\|^2 + \|u_t\|^2 + \|\theta_t\|^2 + \|H_t\|^2),
\end{aligned} \quad (5.8)$$

where $c_* > 0$ is some positive constant depending on μ , λ and C_Ω as in Proposition 2.3. Define

$$\begin{aligned}\phi(t) &\equiv \frac{1}{2} \{ R \|q_t\|^2 + \|\sqrt{\rho}u_t\|^2 + c_v \|\sqrt{\rho}\theta_t\|^2 + \|H_t\|^2 \} - \frac{R}{2} \int_\Omega q q_t^2 dx \\ &= \frac{R}{2} \int_\Omega (1-q) q_t^2 dx + \frac{1}{2} \{ \|\sqrt{\rho}u_t\|^2 + c_v \|\sqrt{\rho}\theta_t\|^2 + \|H_t\|^2 \},\end{aligned}$$

then it is obvious that

$$\|(q_t, u_t, \theta_t, H_t)\|^2 \leq C\phi(t).$$

Moreover, because the fact (5.1) is true, there exists a positive constant $T_1 > 0$ such that

$$\varepsilon + \|\nabla u\|_1^2 \leq \frac{1}{2} \min(c_*, \kappa, \eta), \quad \forall t \geq T_1.$$

Therefore,

$$\phi'(t) \leq a(t)\phi(t), \quad \forall t \geq T_1,$$

where $a(t) = C_\epsilon (\|u_t\|_1^2 + \|\theta_t\|_1^2 + \|(\nabla q, \nabla u, \text{curl} H)\|_2^2 + \|\nabla \theta\|_1^2)$. We have

$$\int_0^\infty \phi(t) dt < \infty, \quad \int_0^\infty a(t) dt < \infty.$$

due to $(q, u, \theta, H) \in X(0, \infty; \mathcal{E}_0)$, Then Lemma 2.7 (i) implies the assertion of Lemma 5.1. $\square$

Lemma 5.2. *For every solution $(q, u, \theta, H) \in X(0, \infty; \mathcal{E}_0)$ of the problem (3.1)-(3.3), it exists a time $T_2 = T_2(q, u, \theta, H) \geq T_1 > 0$ such that*

$$\|(\nabla u, \nabla \theta, \nabla H)\|^2 \leq C \|(q_t, u_t, \theta_t, H_t)\| \quad \forall t \geq T_2, \quad (5.9)$$

$$\|(\nabla^2 \theta, \nabla^2 H)\|^2 \leq C \|(q_t, u_t, \theta_t, H_t)\| \quad \forall t \geq T_2. \quad (5.10)$$

Proof. Multiplying

$$q_t + \text{div}(\rho u) = 0$$

by $R(\theta+1)\ln(q+1)$ and integrating over Ω , after integration by parts with the boundary condition $u \cdot n|_{\partial\Omega} = 0$, we have

$$\begin{aligned}- \int_\Omega R(\theta+1)u \cdot \nabla q dx &= - \int_\Omega Rq_t(\theta+1)\ln(q+1)dx + \int_\Omega R\rho \ln(q+1)u \cdot \nabla \theta dx \\ &\leq C\|q_t\| + C\|q\|_{L^3}\|u\|_{L^6}\|\nabla \theta\|,\end{aligned} \quad (5.11)$$

where we have used the fact

$$|\ln(q+1) - \ln 1| \leq C|q|.$$

Let us rewrite (3.1)₂ in the following form

$$\rho(u_t + u \cdot \nabla u) - \mu \Delta u - (\mu + \lambda) \nabla \text{div} u + R(\theta+1) \nabla q + R \nabla \theta = -Rq \nabla \theta + \text{curl} H \times H,$$

multiplication this equation by u , and integration by parts indicate that

$$\begin{aligned}&\mu \|\text{curl} u\|^2 + (2\mu + \lambda) \|\text{div} u\|^2 + \int_\Omega R(\theta+1)u \cdot \nabla q dx - \int_\Omega R \text{div} u \theta dx - \int_\Omega (\text{curl} H \times H) \cdot u dx \\ &= - \int_\Omega \rho u_t \cdot u dx - \int_\Omega \rho (u \cdot \nabla) u \cdot u dx - \int_\Omega Rqu \cdot \nabla \theta dx,\end{aligned}$$

then by means of Hölder's inequality, it leads to

$$\begin{aligned}&\mu \|\text{curl} u\|^2 + (2\mu + \lambda) \|\text{div} u\|^2 + \int_\Omega R(\theta+1)u \cdot \nabla q - \int_\Omega R \text{div} u \theta - \int_\Omega (\text{curl} H \times H) \cdot u \\ &\leq C\|u_t\| + C\|(u, q)\|_{L^3} \|(\nabla u, \nabla \theta)\|^2.\end{aligned} \quad (5.12)$$

Similarly, multiplying (3.1)₃ by θ , we have

$$\begin{aligned} & \kappa \|\nabla \theta\|^2 + R \int_{\Omega} \operatorname{div} u \theta dx \\ &= \int_{\Omega} (-c_v \rho (\theta_t + u \cdot \nabla \theta) - R(\rho \theta + q) \operatorname{div} u + \lambda (\operatorname{div} u)^2 + 2\mu |S(u)|^2 + \eta |\operatorname{curl} H|^2) \theta dx, \end{aligned}$$

which implies

$$\kappa \|\nabla \theta\|^2 + R \int_{\Omega} \operatorname{div} u \theta dx \leq C \|\theta_t\| + C(\|\theta\|_{L^\infty} + \|(u, q, \theta)\|_{L^3}) \|(\nabla u, \nabla \theta, \operatorname{curl} H)\|^2. \quad (5.13)$$

And, multiplying (3.1)₄ by H , and integrating by parts with the boundary condition $\operatorname{curl} H \times n|_{\partial\Omega} = 0$, it shows

$$\eta \|\operatorname{curl} H\|^2 - \int_{\Omega} \operatorname{curl}(u \times H) \cdot H = - \int_{\Omega} H_t \cdot H \leq C \|H_t\|. \quad (5.14)$$

Putting (5.11)-(5.14) together, it could be arrived at

$$\begin{aligned} & \mu \|\operatorname{curl} u\|^2 + (2\mu + \lambda) \|\operatorname{div} u\|^2 + \kappa \|\nabla \theta\|^2 + \eta \|\operatorname{curl} H\|^2 \\ & \leq C \|(q_t, u_t, \theta_t, H_t)\| + C(\|\theta\|_{L^\infty} + \|(u, q, \theta)\|_{L^3}) \|(\nabla u, \nabla \theta, \operatorname{curl} H)\|^2, \end{aligned}$$

where we have used the fact

$$\int_{\Omega} (\operatorname{curl} H \times H) \cdot u dx + \int_{\Omega} \operatorname{curl}(u \times H) \cdot H dx = 0.$$

Taking the same argument as (5.8) of Lemma 5.1, the desired results (5.9) is received.

In order to prove (5.10), we employ (3.1)₃ to get the estimate for $\Delta \theta$ firstly:

$$\kappa \Delta \theta = c_v \rho (\theta_t + u \cdot \nabla \theta) + R \operatorname{div} u - \lambda (\operatorname{div} u)^2 - 2\mu |S(u)|^2 + R(\rho \theta + q) \operatorname{div} u - \eta |\operatorname{curl} H|^2$$

which implies

$$\|\Delta \theta\|^2 \leq C(\|\theta_t\| + \|(\nabla u, \nabla \theta, \operatorname{curl} H)\|^2). \quad (5.15)$$

Recalling the stand elliptic estimate in Lemma 2.6:

$$\|\nabla^2 \theta\|^2 \leq C \left(\|\Delta \theta\|^2 + \|\nabla \theta\|^2 \right) \quad \text{for } \theta \in H^2(\Omega) \quad \text{with } \frac{\partial \theta}{\partial n} \Big|_{\partial\Omega} = 0, \quad (5.16)$$

which together with (5.9) and (5.15) induces that

$$\|\nabla^2 \theta\|^2 \leq C \|(q_t, u_t, \theta_t, H_t)\| \quad \forall t \geq T_2. \quad (5.17)$$

Similarly, we have the following equation from (3.1)₄:

$$\eta \Delta H = H_t - \operatorname{curl}(u \times H),$$

which implies that

$$\begin{aligned} \|\Delta H\|^2 & \leq \|H_t\|^2 + \|u\|_{L^\infty}^2 \|\nabla H\|^2 + \|H\|_{L^\infty} \|\nabla u\|^2 \\ & \leq C \|(q_t, u_t, \theta_t, H_t)\|. \end{aligned}$$

Furthermore, using the standard elliptic estimate, it holds

$$\|\nabla^2 H\|^2 \leq C \left(\|\Delta H\|^2 + \|\nabla H\|^2 \right) \leq C \|(q_t, u_t, \theta_t, H_t)\|.$$

Therefore, the proof of Lemma 5.2 is completed. $\square$

Lemma 5.3. *For every solution $(q, u, \theta, H) \in X(0, \infty; \mathcal{E}_0)$ of the problem (3.1)-(3.3), it exists a time $T_3 = T_3(q, u, \theta, H) \geq T_2$ such that*

$$\|\operatorname{curl}^2 u\| \leq C t^{-1/4} \quad \forall t \geq T_3 \quad (5.18)$$

Proof. Recalling the vorticity equation:

$$\rho(w_t + u \cdot \nabla w) - \mu \Delta w = K + \operatorname{curl}(\operatorname{curl} H \times H), \quad (5.19)$$

where

$$K = -\nabla q \times (u_t + u \cdot \nabla u) - \rho(w \cdot \nabla)u - \rho w \operatorname{div} u.$$

Taking the L^2 inner product of (5.19) with w , using integration by parts with $w \times n|_{\partial\Omega} = 0$, we obtain

$$\frac{1}{2} \frac{d}{dt} \|\sqrt{\rho} w\|^2 + \mu \|\operatorname{curl} w\|^2 = \int_{\Omega} K \cdot w dx + \int_{\Omega} (\operatorname{curl} H \times H) \cdot \operatorname{curl} w dx. \quad (5.20)$$

It is easy to check

$$\int_{\Omega} K \cdot w dx \leq C \|(q_t, u_t, H_t, \theta_t)\|$$

and

$$\begin{aligned} \int_{\Omega} (\operatorname{curl} H \times H) \cdot \operatorname{curl} w dx &\leq \frac{\mu}{2} \|\operatorname{curl} w\|^2 + C \|\operatorname{curl} H \times H\|^2 \\ &\leq \frac{\mu}{2} \|\operatorname{curl} w\|^2 + C \|(q_t, u_t, H_t, \theta_t)\|. \end{aligned}$$

after utilizing Sobolev's inequality and Lemma 5.2. Hence, (5.20) becomes

$$\frac{1}{2} \frac{d}{dt} \|\sqrt{\rho} w\|^2 + \frac{\mu}{2} \|\operatorname{curl} w\|^2 \leq C \|(q_t, u_t, H_t, \theta_t)\|. \quad (5.21)$$

On the other hand, multiplication (5.19) by w_t followed by integration over Ω gives that

$$\frac{\mu}{2} \frac{d}{dt} \|\operatorname{curl} w\|^2 + \|\sqrt{\rho} w_t\|^2 = - \int_{\Omega} \rho u \cdot \nabla w \cdot w_t dx + \int_{\Omega} K \cdot w_t dx + \int_{\Omega} \operatorname{curl}(\operatorname{curl} H \times H) \cdot w_t dx \quad (5.22)$$

Applying Sobolev's inequality and Lemma 5.2, we obtain

$$\begin{aligned} \int_{\Omega} \rho u \cdot \nabla w \cdot w_t dx &\leq C \|u\|_{L^\infty} \|\nabla w\| \|\sqrt{\rho} w_t\| \\ &\leq \frac{1}{8} \|\sqrt{\rho} w_t\|^2 + C \|\nabla u\|_1^2 \|\nabla w\|^2 \\ &\leq \frac{1}{8} \|\sqrt{\rho} w_t\|^2 + C \|\nabla u\|_1^2 \|\operatorname{curl} w\|^2 + C \|\nabla u\|_1^2 \|w\|^2 \\ &\leq \frac{1}{8} \|\sqrt{\rho} w_t\|^2 + C \|\nabla u\|_1^2 \|\operatorname{curl} w\|^2 + C \|(q_t, u_t, H_t, \theta_t)\|, \end{aligned} \quad (5.23)$$

$$\int_{\Omega} K \cdot w_t dx \leq \frac{1}{8} \|\sqrt{\rho} w_t\|^2 + C \|(q_t, u_t, H_t, \theta_t)\|, \quad (5.24)$$

and

$$\begin{aligned} \int_{\Omega} \operatorname{curl}(\operatorname{curl} H \times H) \cdot w_t dx &\leq \frac{1}{8} \|\sqrt{\rho} w_t\|^2 + C \|\nabla H\|_1^2 \|\nabla^2 H\|^2 \\ &\leq \frac{1}{8} \|\sqrt{\rho} w_t\|^2 + C \|(q_t, u_t, H_t, \theta_t)\|, \end{aligned} \quad (5.25)$$

then substituting these into (5.22), we derive that

$$\begin{aligned} \frac{\mu}{2} \frac{d}{dt} \|\operatorname{curl} w\|^2 + \frac{5}{8} \|\sqrt{\rho} w_t\|^2 \\ \leq C \|\nabla u\|_1^2 \|\operatorname{curl} w\|^2 + C \|(q_t, u_t, H_t, \theta_t)\|. \end{aligned} \quad (5.26)$$

Summing (5.21) and (5.26) up, one gets

$$\begin{aligned} & \frac{d}{dt} \{ \|\sqrt{\rho}w\|^2 + \mu \|\operatorname{curl} w\|^2 \} + c_0 \{ \|\operatorname{curl} w\|^2 + \|\sqrt{\rho}w_t\|^2 \} \\ & \leq C \|\nabla u\|_1^2 \|\operatorname{curl} w\|^2 + C \|(q_t, u_t, H_t, \theta_t)\|, \end{aligned} \quad (5.27)$$

for some positive constant $c_0 > 0$. From (5.1), there exists T'_3 such that if $t \geq T'_3$, then

$$C \|\nabla u\|_1^2 \leq \frac{c_0}{2},$$

which yields

$$\frac{d}{dt} \{ \|\sqrt{\rho}w\|^2 + \mu \|\operatorname{curl} w\|^2 \} + \frac{c_0}{2} \{ \|\operatorname{curl} w\|^2 + \|\sqrt{\rho}w_t\|^2 \} \leq C \|(q_t, u_t, H_t, \theta_t)\|.$$

So,

$$\frac{d}{dt} \{ \|\sqrt{\rho}w\|^2 + \mu \|\operatorname{curl} w\|^2 \} + \frac{c_0}{2} \|\operatorname{curl} w\|^2 \leq C \|(q_t, u_t, H_t, \theta_t)\|.$$

Putting (5.9) together with the above inequality induce that

$$\frac{d}{dt} \{ \|\sqrt{\rho}w\|^2 + \mu \|\operatorname{curl} w\|^2 \} + \frac{c_0}{2} \{ \|\operatorname{curl} w\|^2 + \|\sqrt{\rho}w\|^2 \} \leq C \|(q_t, u_t, H_t, \theta_t)\|, \quad \forall t > T_3 = \max\{T'_3, T_2\}$$

then from Lemma 5.1, we have

$$g'(t) + \frac{c_0}{2} g(t) \leq C t^{-1/2}, \quad \forall t > T_3$$

where $g(t)$ is defined as

$$g(t) \equiv \|\sqrt{\rho}w\|^2 + \mu \|\operatorname{curl} w\|^2.$$

An application of Lemma 2.7 (ii) implies

$$g(t) \leq C t^{-1/2} \quad \text{for } t \geq T_3.$$

Therefore, this completes the proof of Lemma 5.3. $\square$

Lemma 5.4. *For every solution $(q, u, \theta, H) \in X(0, \infty; \mathcal{E}_0)$ of the problem (3.1)-(3.3), it exists a time $T_4 \geq T_3$, such that*

$$\|\nabla^2 u\| + \|\nabla q\| \leq C t^{-1/4} \quad \forall t \geq T_4. \quad (5.28)$$

Proof. Consider (q, u) as the solution of the following Stokes problem:

$$\begin{cases} -\mu \Delta u + R \nabla q = G & \text{in } \Omega \\ \operatorname{div} u = \operatorname{div} u & \text{in } \Omega \\ u \cdot n|_{\partial\Omega} = 0, \quad \operatorname{curl} u \times n|_{\partial\Omega} = 0 \end{cases} \quad (5.29)$$

where

$$\begin{aligned} G &= (\mu + \lambda) \nabla \operatorname{div} u - \rho(u_t + u \cdot \nabla u) - R \nabla \theta - R \nabla(q\theta) + H \cdot \nabla H - \frac{1}{2} \nabla |H|^2 \\ &\equiv (\mu + \lambda) \nabla \operatorname{div} u + G_1 \end{aligned}$$

In view of Hölder's inequality, Sobolev's inequality and Lemma 5.2, it is easy to see that

$$\|G_1\|^2 \leq C \|(q_t, u_t, \theta_t, H_t)\| + \|\theta\|_{L^\infty}^2 \|\nabla q\|^2, \quad (5.30)$$

which implies that

$$\|G\|^2 \leq C \|\nabla \operatorname{div} u\|^2 + C \|(q_t, u_t, \theta_t, H_t)\| + \|\theta\|_{L^\infty}^2 \|\nabla q\|^2.$$

Then elliptic estimate (see Lemma 2.5) for the Stoke problem yields

$$\begin{aligned}\|\nabla^2 u\|^2 + \|\nabla q\|^2 &\leq C(\|G\|^2 + \|\operatorname{div} u\|_1^2 + \|\nabla u\|^2) \\ &\leq C\|\nabla \operatorname{div} u\|^2 + C\|(q_t, u_t, \theta_t, H_t)\| + \|\theta\|_{L^\infty}^2 \|\nabla q\|^2.\end{aligned}$$

By (5.1) and Lemma 2.1, there exists a time $T_4 > T_3$ such that

$$\|\theta\|_{L^\infty}^2 \leq C\|\nabla \theta\|_1^2 \leq \frac{1}{4}. \quad (5.31)$$

Therefore, we obtain the following inequality

$$\|\nabla^2 u\|^2 + \frac{3}{4}\|\nabla q\|^2 \leq C\|\nabla \operatorname{div} u\|^2 + C\|(q_t, u_t, \theta_t, H_t)\| \quad (5.32)$$

for $t \geq T_4$. Next, we shall estimate the term $\|\nabla \operatorname{div} u\|$. For this purpose, we take the derivative of the continuity equation with respect to x_i , $1 \leq i \leq 3$ as follows

$$\begin{aligned}(\operatorname{div} u)_{x_i} &= -\frac{1}{\rho}(q_{t,x_i} + q_{x_i} \operatorname{div} u + u \cdot \nabla q_{x_i} + u_{x_i} \cdot \nabla q) \\ &\equiv -\frac{1}{\rho}q_{t,x_i} - \frac{1}{\rho}F^i.\end{aligned} \quad (5.33)$$

It is clear that

$$\|F\|^2 \leq C\|\nabla q\|_2^2 \|\nabla u\|^2 \leq C\|(q_t, u_t, H_t, \theta_t)\|, \quad (5.34)$$

which together with (5.33) implies that

$$\|\nabla \operatorname{div} u\|^2 \leq C \int_{\Omega} |\nabla q_t|^2 dx + C\|(q_t, u_t, H_t, \theta_t)\|. \quad (5.35)$$

In order to deal with $\|\nabla q_t\|^2$, we follow the method given in [9]. Let $d(x) := \operatorname{dist}(x, \partial\Omega)$ satisfies

$$d \in C^2(\bar{\Gamma}_h), \Gamma_h = \{x \in \Omega | 0 \leq d(x) < h\}$$

and $\psi \in C^\infty(\mathbb{R}^+)$ be non-increasing with $\psi(d) = 1$ for $0 \leq d \leq \frac{1}{2}h$, and $\psi(d) = 0$ for $d \geq \frac{3}{4}h$. For $1 \leq i, k \leq 3$, we define

$$e_{ik}(x) := \delta_{ik} - \psi(d(x))^2 d_{x_i}(x) d_{x_k}(x).$$

For any smooth function $f : \Omega \rightarrow \mathbb{R}$, one has

$$\begin{aligned}f_{x_i} &= e_{ik} f_{x_k} + \psi(d)^2 d_{x_i} d_{x_k} f_{x_k} \quad 1 \leq i \leq 3, \\ |\nabla f|^2 &= e_{ik} f_{x_i} f_{x_k} + \psi(d)^2 (\nabla f \cdot \nabla d)^2.\end{aligned} \quad (5.36)$$

Since $\psi(0) = 1$ and $\nabla d(x) = n(x)$ for any $x \in \partial\Omega$, where n denotes the outer unit normal, we have

$$\begin{aligned}e_{ik} n_k|_{\partial\Omega} &= 0, \quad 1 \leq i \leq 3, \\ e_{ik} f_{x_k} n_i|_{\partial\Omega} &= 0.\end{aligned} \quad (5.37)$$

Therefore, we have the following equivalent form

$$\int_{\Omega} |\nabla q_t|^2 dx = \int_{\Omega} e_{ik} q_{t,x_i} q_{t,x_k} dx + \int_{\Omega} \psi(d)^2 (\nabla q_t \cdot \nabla d)^2 dx, \quad (5.38)$$

The first term on the right-hand side of (5.38) could be controlled as

$$\begin{aligned}\int_{\Omega} e_{ik} q_{t,x_i} q_{t,x_k} dx &= - \int_{\Omega} (e_{ik})_{x_i} q_t q_{t,x_k} dx - \int_{\Omega} e_{ik} q_t q_{t,x_i x_k} dx + \int_{\partial\Omega} e_{ik} n_i q_t q_{t,x_k} dx \\ &= - \int_{\Omega} (e_{ik})_{x_i} q_t q_{t,x_k} dx - \int_{\Omega} e_{ik} q_t q_{t,x_i x_k} dx \\ &\leq C\|q_t\| \|\nabla q_t\|_1 \leq C\|q_t\|.\end{aligned} \quad (5.39)$$

by integrating by parts with the boundary condition (5.37). Therefore, combining (5.32), (5.35) with (5.38)-(5.39) gives that

$$\|\nabla^2 u\|^2 + \|\nabla q\|^2 \leq C\|\psi(d)(\nabla q_t \cdot \nabla d)\|^2 + C\|(q_t, u_t, \theta_t, H_t)\|, \quad \forall t \geq T_3. \quad (5.40)$$

Multiply (5.33) by $(2\mu + \lambda)\psi(d)d_{x_i}$, and sum up all these equations for $i = 1, 2, 3$, then we get

$$\frac{2\mu + \lambda}{\rho}\psi(d)\nabla q_t \cdot \nabla d + (2\mu + \lambda)\psi(d)\nabla \operatorname{div} u \cdot \nabla d = \frac{2\mu + \lambda}{\rho}\psi(d)F \cdot \nabla d \quad (5.41)$$

In order to cancel the second term on the left-hand side on (5.41), we rewrite (5.29)₁ as

$$-(2\mu + \lambda)\nabla \operatorname{div} u + R\nabla q = -\mu \operatorname{curl} \operatorname{curl} u + G_1$$

then multiplying this identity by $\psi(d)\nabla d$, and adding the result to (5.41), we infer that

$$\begin{aligned} & \frac{2\mu + \lambda}{\rho}\psi(d)\nabla q_t \cdot \nabla d + R\psi(d)\nabla q \cdot \nabla d \\ &= \frac{2\mu + \lambda}{\rho}\psi(d)F \cdot \nabla d - \mu\psi(d)\operatorname{curl} \operatorname{curl} u \cdot \nabla d + \psi(d)G_1 \cdot \nabla d \equiv Q. \end{aligned} \quad (5.42)$$

Moreover, (5.34), (5.30), Lemma 5.1 and Lemma 5.3 imply that

$$\begin{aligned} \|Q\|^2 &\leq C\|\operatorname{curl} \operatorname{curl} u\|^2 + \|F\|^2 + \|G_1\|^2 \\ &\leq C\|\theta\|_{L^\infty}^2 \|\nabla q\|^2 + Ct^{-1/2}, \end{aligned} \quad (5.43)$$

where we have used the definition of $\psi(d)$ and $|\nabla n| = 1$. Therefore, (5.31) and (5.40) induce

$$\|\nabla^2 u\|^2 + \frac{1}{2}\|\nabla q\|^2 \leq C\|\psi(d)\nabla q \cdot \nabla d\|^2 + Ct^{-1/2}, \quad \text{for } t \geq \widetilde{T}_4. \quad (5.44)$$

On the other side, multiplying (5.42) by $\psi(d)\nabla q \cdot \nabla d$, it indicates

$$\begin{aligned} & \frac{d}{dt} \int_{\Omega} \frac{2\mu + \lambda}{\rho} (\psi(d)\nabla q \cdot \nabla d)^2 dx + R\|\psi(d)\nabla q \cdot \nabla d\|^2 \\ &= \int_{\Omega} (\psi(d)\nabla q \cdot \nabla d)Q dx + \int_{\Omega} \left(\frac{2\mu + \lambda}{\rho}\right)_t \psi(d)\nabla q \cdot \nabla d dx \\ &\leq \varepsilon\|\psi(d)\nabla q \cdot \nabla d\|^2 + C_\varepsilon(\|Q\|^2 + \|q_t\|^2). \end{aligned}$$

Therefore, from Lemma 5.1, (5.1), (5.43) and (5.44), we conclude that

$$\frac{d}{dt} \int_{\Omega} \frac{2\mu + \lambda}{\rho} (\psi(d)\nabla q \cdot \nabla d)^2 dx + \frac{R}{2}\|\psi(d)\nabla q \cdot \nabla d\|^2 \leq Ct^{-1/2}, \quad \text{for } t \geq T_4 \geq \widetilde{T}_4.$$

One can apply (ii) of Lemma 2.7 to gain

$$\int_{\Omega} \frac{2\mu + \lambda}{\rho} (\psi(d)\nabla q \cdot \nabla d)^2 dx \leq Ct^{-1/2},$$

which together with (5.44) finishes the proof of Lemma 5.4. $\square$

Now, we are ready to prove Theorem 1.3.

Proof of Theorem 1.3 Lemma 5.2 and Lemma 5.4 yield

$$\|(u, \theta, H)\|_{C^0(\bar{\Omega})} \leq C\|(\nabla u, \nabla \theta, \operatorname{curl} H)\|_1 \leq Ct^{-1/4}.$$

Because

$$\|q\|_{L^4}^4 \leq \|q\| \|q\|_{L^6}^3 \leq C\|\nabla q\|^3 \leq Ct^{-3/4},$$

and

$$\|\nabla q\|_{L^4}^4 \leq \|\nabla q\|_{L^\infty}^2 \|\nabla q\|^2 \leq Ct^{-1/2}.$$

Then we have

$$\|\rho - 1\|_{C^0(\bar{\Omega})} = \|q\|_{C^0(\bar{\Omega})} \leq C\|q\|_{H^{1,4}} \leq C\left(t^{-3/16} + t^{-1/8}\right) \leq Ct^{-1/8}.$$

Hence Theorem 1.3 is proved. $\square$

Acknowledgements

Liu's research is supported by National Natural Science Foundation of China (No.12071219, 12026432). Luo's research is supported by a grant from the Research Grants Council of the Hong Kong Special Administrative Region, China (Project No. 11307420). Zhong is supported by the Fundamental Research Funds for the Central Universities (No. A0920502052101-224). Luo would like to thank Professor Chongchun Zeng for helpful discussions on the Hodge type estimates.

REFERENCES

- [1] C. Amrouche, V. Girault, J. Giroire, Dirichlet and Neumann exterior problems for the n-dimensional Laplace operator: an approach in weighted Sobolev spaces, *J. Math. Pures Appl.* 76(1) (1997) 55-81.
- [2] L. C. Berselli, S. Spirito, On the Vanishing Viscosity Limit of 3D Navier-Stokes Equations under Slip Boundary Conditions in General Domains, *Commun. Math. Phys.*, 316 (2012) 171-198.
- [3] G.C. Cai, J.Li, Existence and exponential growth of global classical solutions to the compressible Navier-Stokes equations with slip boundary conditions in 3D bounded domains, arXiv:2102.06348.
- [4] Y. Chen, B. Huang, and X. Shi. Global strong solutions to the compressible Magnetohydrodynamic equations with slip boundary conditions in 3D bounded domains, arXiv:2102.07341.
- [5] Y. Chen, B. Huang, and X. Shi. Global Strong and Weak Solutions to the Initial-boundary-value Problem of 2D Compressible MHD System with Large Initial Data and Vacuum, arXiv:2105.11291.
- [6] Q. Chen, Z. Tan, Global existence and convergence rates of smooth solutions for the compressible magnetohydrodynamic equations, *Nonlinear Anal.* 72 (2010) 4438-4451.
- [7] G. Q. Chen, D.H. Wang, Global solution of nonlinear magnetohydrodynamics with large initial data, *J. Differential Equations* 182 (2002) 344-376.
- [8] W.Q. Cui, Y.B. Ou, D.D. Ren, Incompressible limit of full compressible magnetohydrodynamic equations with well-prepared data in 3-D bounded domains, *J. Math. Anal. Appl.* 427 (2015) 263-288.
- [9] K. Deckelnick, Decay estimates for the compressible Navier-Stokes equations in unbounded domain. *Math. Z.* 209 (1992) 115-130.
- [10] K. Deckelnick, L^2 -decay for the compressible Navier-Stokes equations in unbounded domains, *Commun. in Partial Differential Equations* 18 (1993) 1445-1476.
- [11] A. Dhifaoui, M. Meslameni, U. Razafison, Weighted Hilbert spaces for the stationary exterior Stokes problem with Navier slip boundary conditions, *J. Math. Anal. Appl.* 472(2) (2019) 1846-1871.
- [12] C. Dou, S. Jiang, and Q. Ju. Global existence and the low mach number limit for the compressible magnetohydrodynamic equations in a bounded domain with perfectly conducting boundary. *Z. Angew. Math. Phys.*, 64(6) (2013) 1661-1678.
- [13] B. Ducomet, E. Feireisl, The equations of Magnetohydrodynamics: On the interaction between matter and radiation in the evolution of gaseous stars, *Comm. Math. Phys.* 226 (2006) 595-629.
- [14] G.P. Galdi, W.J. Layton, Approximation of the larger eddies in fluid motion. II. A model for space-filtered flow, *Math. Models Methods Appl. Sci.* 10 (2000) 343-350.
- [15] J.C. Gao, Y.H. Chen, Z.A. Yao, Long-time behavior of solution to the compressible magnetohydrodynamic equations, *Nonlinear Anal.* 128 (2015) 122-135.
- [16] L. Guo, F.C. Li, F. Xie, Asymptotic limits of the isentropic compressible viscous magnetohydrodynamic equations with Navier-slip boundary conditions, *J. Differ. Equ.* 267 (2019) 6910-6957.
- [17] X.P. Hu, D.H. Wang, Global solutions to the three-dimensional full compressible magnetohydrodynamics flows, *Comm. Math. Phys.* 283 (2008) 253-284.
- [18] X.P. Hu, D.H. Wang, Global existence and large-time behavior of solutions to the three-dimensional equations of compressible magnetohydrodynamic flows, *Arch. Ration. Mech. Anal.* 202 (2011) 787-827.
- [19] S. Jiang, Global spherically symmetric solutions to the equations of a viscous polytropic ideal gas in an exterior domain, *Comm. Math. Phys.* 178 (1996) 339-374.
- [20] S. Kawashima, M. Okada, Smooth global solutions for the one-dimensional equations in magnetohydrodynamics, *Proc. Japan Acad. Ser. A* 58 (1982) 384-387.
- [21] T. Kobayashi, Y. Shibata, Decay estimates of solutions for the equations of motion of compressible viscous and heat conductive gases in an exterior domain in $\mathbb{R}^3$, *Comm. Math. Phys.* 200 (1999) 621-659.
- [22] L.D. Landau, E.M. Lifshitz, *Electrodynamics of Continuous Media*, 2nd ed., New York: Pergamon, 1984.

- [23] F.C. Li, H.J. Yu, Optimal decay rate of classical solutions to the compressible magnetohydrodynamic equations, *Proc. R. Soc. Edinb. Sect. A* 141 (2011) 109-126.
- [24] H.L. Li, X.Y. Xu, J.W. Zhang, Global classical solutions to the 3D compressible magnetohydrodynamic equations with large oscillations and vacuum, *SIAM J. Math. Anal.* 45 (2013) 1356-1387.
- [25] P. Maremonti, Asymptotic stability theorems for viscous fluid motions in exterior domains, *Rend. Semin. Mat. Univ. Padova* 71 (1984) 35-72.
- [26] A. Matsumura, T. Nishida, Initial boundary value problem for equations of motion of compressible viscous and heat conductive fluids, *Comm. Math. Phys.* 89 (1983) 445-464.
- [27] N. Masmoudi, F. Rousset, Uniform regularity for the Navier-Stokes equation with Navier boundary condition, *Arch. Ration. Mech. Anal.* 203 (2012) 529-575.
- [28] C.L.M.H. Navier, Sur les lois de l'équilibre et du mouvement des corps élastiques, *Mem. Acad. R. Sci. Inst. France* 6 (1827) 369.
- [29] X.K. Pu, B.L. Guo, Global existence and convergence rates of smooth solutions for the full compressible MHD equations, *Z. Angew. Math. Phys.* 64 (2013) 519-538.
- [30] Y. Sun, C. Wang, F. Zhang, A Beale-Kato-Majda blow-up criterion to the compressible Navier-Stokes equation, *J. Math. Pures Appl.* 95 (2011) 36-47.
- [31] W. Von Wahl, Estimating ∇u by $\operatorname{div} u$ and $\operatorname{curl} u$, *Mathematical Methods in the Applied Sciences*, 15 (1992) 123-143.
- [32] D.H. Wang, Large solutions to the initial-boundary value problem for planar magnetohydrodynamics, *SIAM J. Appl. Math.* 63 (2003) 1424-1441.
- [33] Y. Wang, Uniform regularity and vanishing dissipation limit for the full compressible Navier-Stokes system in three dimensional bounded domain, *Arch. Ration. Mech. Anal.* 221 (2016) 1345-1415.
- [34] Y. Xiao, Z. Xin, On the vanishing viscosity limit for the 3D Navier-Stokes equations with a slip boundary condition, *Commun. Pure Appl. Math.* 60 (2007) 1027-1055.
- [35] Y. Xiao, Z. Xin, J. Wu, Vanishing viscosity limit for the 3D magnetohydrodynamic system with a slip boundary condition, *J. Funct. Anal.* 257 (2009) 3375-3394.
- [36] J.W. Zhang, J.N. Zhao, Some decay estimates of solutions for the 3-D compressible isentropic magnetohydrodynamics, *Commun. Math. Sci.* 8 (2010) 835-850.