

SEMICLASSICAL STATES FOR A MAGNETIC NONLINEAR SCHRÖDINGER EQUATION WITH EXPONENTIAL CRITICAL GROWTH IN \mathbb{R}^2

PIETRO D'AVENIA AND CHAO JI

ABSTRACT. This paper is devoted to the magnetic nonlinear Schrödinger equation

$$\left(\frac{\varepsilon}{i}\nabla - A(x)\right)^2 u + V(x)u = f(|u|^2)u \text{ in } \mathbb{R}^2,$$

where $\varepsilon > 0$ is a parameter, $V : \mathbb{R}^2 \rightarrow \mathbb{R}$ and $A : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ are continuous functions and $f : \mathbb{R} \rightarrow \mathbb{R}$ is a C^1 function having exponential critical growth. Under a global assumption on the potential V , we use variational methods and Ljusternick-Schnirelmann theory to prove existence, multiplicity, concentration, and decay of nontrivial solutions for $\varepsilon > 0$ small.

CONTENTS

1. Introduction and main results	1
Notation	3
2. The variational framework and some preliminaries	4
3. The limit problem	11
4. Proof of Theorem 1.1	16
5. Proof of Theorem 1.2	29
6. Proof of Theorem 1.3	33
Acknowledgments	34
References	34

1. INTRODUCTION AND MAIN RESULTS

In this paper, we consider the following nonlinear Schrödinger equation

$$(1.1) \quad \left(\frac{\varepsilon}{i}\nabla - A(x)\right)^2 u + V(x)u = f(|u|^2)u \quad \text{in } \mathbb{R}^2,$$

where $\varepsilon > 0$ is a parameter, $u : \mathbb{R}^2 \rightarrow \mathbb{C}$, $A \in C(\mathbb{R}^2, \mathbb{R}^2)$, $V \in C(\mathbb{R}^2, \mathbb{R})$ satisfies

$$(V) \quad V_\infty = \liminf_{|x| \rightarrow +\infty} V(x) > V_0 = \inf_{x \in \mathbb{R}^2} V(x) > 0,$$

where V_∞ can be also $+\infty$, and $f \in C^1(\mathbb{R}, \mathbb{R})$ satisfies the following assumptions:

- (f₁) $f(t) = 0$ if $t \leq 0$;
- (f₂) there holds

$$\lim_{t \rightarrow +\infty} \frac{f(t^2)t}{e^{\alpha t^2}} = \begin{cases} 0, & \text{for } \alpha > 4\pi, \\ +\infty, & \text{for } 0 < \alpha < 4\pi, \end{cases}$$

(exponential critical growth);

- (f₃) there is a positive constant $\theta > 2$ such that, for every $t > 0$,

$$0 < \frac{\theta}{2}F(t) \leq tf(t),$$

Date: November 15, 2021.

2010 Mathematics Subject Classification. 35J20, 35J60, 35B33.

Key words and phrases. Nonlinear Schrödinger equation, Magnetic field, Exponential critical growth, Trudinger-Moser inequality, Variational methods.

where $F(t) = \int_0^t f(s)ds$;

(f₄) there exist two constants $p > 2$ and

$$C_p > \left(\frac{8\theta}{\theta-2}\right)^{\frac{p-2}{2}} \left(\frac{p-2}{p}\right)^{\frac{p-2}{2}} V^* S_p^{p/2} > 0$$

such that

$$f'(t) \geq \frac{p-2}{2} C_p t^{(p-4)/2} \quad \text{for all } t > 0,$$

where

$$(1.2) \quad S_p = \inf_{u \in H^1(\mathbb{R}^2, \mathbb{R}) \setminus \{0\}} \frac{\|\nabla u\|_2^2 + \|u\|_2^2}{\|u\|_p^2} \quad \text{and } V^* = \begin{cases} V_\infty, & \text{if } V_\infty < +\infty, \\ V_0, & \text{if } V_\infty = +\infty; \end{cases}$$

(f₅) $f'(t) \leq (e^{4\pi t} - 1)$ for any $t \geq 0$.

Equation (1.1) appears in the study of standing wave solutions $\psi(x, t) := e^{-i\omega/\hbar} u(x)$, with $\omega \in \mathbb{R}$, of the planar nonlinear Schrödinger equation

$$i\hbar \frac{\partial \psi}{\partial t} = \left(\frac{\hbar}{i} \nabla - A(x)\right)^2 \psi + U(x)\psi - f(|\psi|^2)\psi \quad \text{in } \mathbb{R}^2 \times \mathbb{R}.$$

In particular, the operator

$$\psi \mapsto \left(i\hbar \frac{\partial}{\partial t} - U(x)\right)\psi - \left(\frac{\hbar}{i} \nabla - A(x)\right)^2 \psi$$

appears in the study of the interaction of a matter field ψ with an external electromagnetic field whose electric potential is U and whose magnetic potential is A (here we are considering an electromagnetic potential which does not depend on the time variable) though the minimal coupling rule. Moreover, from a physical point of view, the study of the existence of solutions of (1.1), in particular of its ground states, and of their behavior as $\varepsilon \rightarrow 0^+$ (which is equivalent to send the Planck constant \hbar to zero and it is known as *semiclassical limit*) is of the greatest importance in the theory of superconductivity (see [23, 26] and references therein) and since the transition from Quantum Mechanics to Classical Mechanics can be formally performed by this limit.

If the magnetic potential $A \equiv 0$ and \mathbb{R}^2 is replaced by \mathbb{R}^N , then (1.1) reduces to a following nonlinear Schrödinger equation

$$(1.3) \quad -\varepsilon^2 \Delta u + V(x)u = f(|u|^2)u \quad \text{in } \mathbb{R}^N.$$

A vast literature is dedicated to (1.3), in particular concerning existence and multiplicity of solutions, by several methods. If the potential V satisfies the global assumption (V), introduced by Rabinowitz in [32], Cingolani and Lazzo in [17] considered multiplicity and concentration of positive solutions of (1.3) with $f(s) = s^{(p-2)/2}$ with $p > 2$ if $N = 1, 2$ and $2 < p < 2N/(N-2)$ if $N \geq 3$. Then, Alves and Figueiredo in [2] generalized the results in [17] to the p -Laplacian operator for a class of nonlinearities including power growth and in [3] considered multiplicity and concentration of positive solutions for N -Laplacian equation with exponential critical growth in \mathbb{R}^N for $N \geq 2$. Finally, let us mention [1, 5], that are devoted to critical cases in \mathbb{R}^2 .

Moreover, after the seminal paper [25], also the magnetic nonlinear Schrödinger equation (1.1) has been extensively investigated by variational and topological methods (see [9, 15, 16, 19, 21, 29, 30] and references therein). In particular, in [4], Alves, Figueiredo, and Furtado, using the penalization method and Ljusternik-Schnirelmann category theory, obtained a multiplicity result for (1.1) with a subcritical right hand side; in [10], Barile and Figueiredo considered the existence of a solution of (1.1) with an exponentially critically growing nonlinearity in \mathbb{R}^2 ; in [20], we got multiplicity and concentration for (1.1) when the potential V satisfies a local assumption due to del Pino and Felmer [22]. Let us mention also the recent contributions [7] concerning a multiplicity result for a nonlinear fractional magnetic Schrödinger equation with exponential critical growth in the one-dimensional case and [8], where existence and

multiplicity for a nonlinear fractional magnetic Schrödinger equation in \mathbb{R}^N , $N \geq 3$, in the presence of a subcritical nonlinearity and of a potential V satisfying a global assumption, has been proved.

The main result of this paper is the following.

Theorem 1.1. *Assume that (V) and (f_1) – (f_5) hold. Then, there exists $\varepsilon_0 > 0$ such that problem (1.1) admits a ground state solution for any $\varepsilon \in (0, \varepsilon_0)$.*

Moreover, introducing the sets

$$(1.4) \quad M := \{x \in \mathbb{R}^2 : V(x) = V_0\}$$

and

$$M_\delta := \{x \in \mathbb{R}^2 : \text{dist}(x, M) \leq \delta\}, \quad \delta > 0,$$

we are able to show, for small $\varepsilon > 0$, a multiplicity result for problem (1.1) and concentration phenomena of such solutions. More precisely, we have

Theorem 1.2. *Assume that (V) and (f_1) – (f_5) hold. Then, for any $\delta > 0$, there exists $\varepsilon_\delta > 0$ such that problem (1.1) has at least $\text{cat}_{M_\delta}(M)$ nontrivial solutions, for any $0 < \varepsilon < \varepsilon_\delta$.*

Finally, for our solutions we are able to prove their concentration and decay. Indeed we have

Theorem 1.3. *Let $\{\varepsilon_n\}$ be a sequence in $(0, +\infty)$ such that $\varepsilon_n \rightarrow 0^+$, and, for n large enough, let $\{u_{\varepsilon_n}\}$ be a sequence of solutions found in Theorem 1.1 or in Theorem 1.2. If $\eta_n \in \mathbb{R}^2$ is a global maximum point of $|u_{\varepsilon_n}|$, then*

$$\lim_{n \rightarrow \infty} V(\eta_n) = V_0.$$

Moreover, if, furtherly, $A \in C^1(\mathbb{R}^2, \mathbb{R}^2)$, then there exist $C, c > 0$ such that

$$(1.5) \quad |u_{\varepsilon_n}(x)| \leq Ce^{-c \frac{|x - \eta_n|}{\varepsilon_n}} \text{ in } \mathbb{R}^2.$$

We get our results combining variational methods and the Ljusternik-Schnirelmann theory.

This paper is a completion of our joint research project started with [20], where we assumed a local condition on V . Here the presence of the global condition (V) and the weaker assumption $A \in C(\mathbb{R}^2, \mathbb{R}^2)$, require more delicate estimates and a careful and deeper analysis of some technical aspects. Moreover, comparing our case with known results obtained for $A \equiv 0$, the presence of the magnetic field leads to consider a complex valued problem and this implies to get more delicate estimates. We are also able to weaken the assumptions of some technical lemmas (see, for instance, Lemma 2.6) and to study the behavior of the maximum points of the modulus of solutions and their decay both for the ground states obtained in Theorem 1.1 and for solutions found in Theorem 1.2.

The paper is organized as follows. In Section 2 we introduce the functional setting and we give some preliminaries. We study the limit problem in Section 3. Then, in Section 4, we prove Theorem 1.1. Section 5 is devoted to prove Theorem 1.2. Finally, in the last section, we prove Theorem 1.3.

Notation.

- C, C_1, C_2, \dots denote positive constants whose exact values can change from line to line;
- $B_R(y)$ denotes the open disk centered at $y \in \mathbb{R}^2$ with radius $R > 0$ and $B_R^c(y)$ denotes its complement in \mathbb{R}^2 ;
- $\|\cdot\|$, $\|\cdot\|_q$, and $\|\cdot\|_\infty$ denote the usual norms of the spaces $H^1(\mathbb{R}^2, \mathbb{R})$, $L^q(\mathbb{R}^2, \mathbb{K})$, and $L^\infty(\mathbb{R}^2, \mathbb{K})$, respectively, where $\mathbb{K} = \mathbb{R}$ or $\mathbb{K} = \mathbb{C}$.

2. THE VARIATIONAL FRAMEWORK AND SOME PRELIMINARIES

In this section we introduce the functional spaces that we use and a *classical* equivalent version of (1.1). We also present some results for a *limit problem* which will be useful for our arguments.

From now on, in the whole paper, we will assume that (V) holds.

For a function $u : \mathbb{R}^2 \rightarrow \mathbb{C}$, let us denote by

$$\nabla_A u := \left(\frac{\nabla}{i} - A \right) u$$

and consider the space

$$H_A^1(\mathbb{R}^2, \mathbb{C}) := \{u \in L^2(\mathbb{R}^2, \mathbb{C}) : |\nabla_A u| \in L^2(\mathbb{R}^2, \mathbb{R})\},$$

which is an Hilbert space endowed with the scalar product

$$\langle u, v \rangle := \operatorname{Re} \int_{\mathbb{R}^2} \left(\nabla_A u \overline{\nabla_A v} + u \bar{v} \right) dx, \quad \text{for any } u, v \in H_A^1(\mathbb{R}^2, \mathbb{C}),$$

where Re and the bar denote the real part of a complex number and the complex conjugation, respectively. Hence, let $\|\cdot\|_A$ be the norm induced by this inner product.

Observe that a useful tool when we work on $H_A^1(\mathbb{R}^2, \mathbb{C})$ is the diamagnetic inequality

$$(2.1) \quad |\nabla_A u(x)| \geq |\nabla |u|(x)|$$

(see e.g. [31, Theorem 7.21]).

Moreover, a simple change of variables allows us to write equation (1.1) as

$$(2.2) \quad \left(\frac{1}{i} \nabla - A_\varepsilon(x) \right)^2 u + V_\varepsilon(x) u = f(|u|^2) u \quad \text{in } \mathbb{R}^2,$$

where $A_\varepsilon(x) = A(\varepsilon x)$ and $V_\varepsilon(x) = V(\varepsilon x)$.

Thus, we introduce the Hilbert space

$$H_\varepsilon := \left\{ u \in H_{A_\varepsilon}^1(\mathbb{R}^2, \mathbb{C}) : \int_{\mathbb{R}^2} V_\varepsilon(x) |u|^2 dx < \infty \right\}$$

endowed with the scalar product

$$\langle u, v \rangle_\varepsilon := \operatorname{Re} \int_{\mathbb{R}^2} \left(\nabla_{A_\varepsilon} u \overline{\nabla_{A_\varepsilon} v} + V_\varepsilon(x) u \bar{v} \right) dx$$

and $\|\cdot\|_\varepsilon$ is the norm induced by $\langle \cdot, \cdot \rangle_\varepsilon$.

For every $\varepsilon > 0$, the functional space H_ε satisfies the following embeddings. For the sake of completeness here we give some details about their proof.

Lemma 2.1. *The space H_ε is continuously embedded in $L^r(\mathbb{R}^2, \mathbb{C})$ for $r \geq 2$ and compactly embedded in $L_{\text{loc}}^r(\mathbb{R}^2, \mathbb{C})$ for $r \geq 1$. Moreover, if $V_\infty = +\infty$, then H_ε is compactly embedded in $L^r(\mathbb{R}^2, \mathbb{C})$ for $r \geq 2$.*

Proof. According to the diamagnetic inequality (2.1), if $u \in H_\varepsilon$, then $|u| \in H^1(\mathbb{R}^2, \mathbb{R})$ and

$$(2.3) \quad \| |u| \| \leq C \| u \|_\varepsilon.$$

Thus, the continuous embedding $H_\varepsilon \hookrightarrow L^r(\mathbb{R}^2, \mathbb{C})$ is a consequence of (2.3) and of the continuous embedding $H^1(\mathbb{R}^2, \mathbb{R}) \hookrightarrow L^r(\mathbb{R}^2, \mathbb{R})$ for $r \geq 2$.

To get the compact embedding of H_ε in $L_{\text{loc}}^r(\mathbb{R}^2, \mathbb{C})$ for $r \geq 1$ we can argue in the following way.

Let K be an open subset of \mathbb{R}^2 with compact closure.

First of all, we prove that H_ε is continuously embedded in $H^1(K, \mathbb{C})$.

Indeed we have

$$\begin{aligned} \|u\|_{H^1(K, \mathbb{C})}^2 &= \int_K |\nabla u|^2 dx + \int_K |u|^2 dx = \int_K \left| \frac{\nabla}{i} u \right|^2 dx + \int_K |u|^2 dx \\ &\leq C \left(\int_K |\nabla_{A_\varepsilon} u|^2 dx + \int_K |A_\varepsilon u|^2 dx + \int_K V_\varepsilon(x) |u|^2 dx \right) \\ &\leq C \left(\int_K |\nabla_{A_\varepsilon} u|^2 dx + \int_K V_\varepsilon(x) |u|^2 dx \right) \leq C \|u\|_\varepsilon^2. \end{aligned}$$

Let now \mathcal{F} be a bounded subset of H_ε . We want to prove that $\mathcal{F}|_K$, which is a subset of $H^1(K, \mathbb{C})$ due to the continuous embedding proved before, has compact closure in $L^r(K, \mathbb{C})$, $r \geq 1$.

Let $\tilde{\mathcal{P}}$ be an extension operator found in [12, Theorem 9.7] for $H^1(K, \mathbb{R})$ and $\mathcal{P} := \tilde{\mathcal{P}}(\operatorname{Re}(u)) + i\tilde{\mathcal{P}}(\operatorname{Im}(u))$. We know that, for each $v \in H^1(K, \mathbb{R})$,

$$\tilde{\mathcal{P}}v|_K = v, \quad \|\tilde{\mathcal{P}}v\|_{L^2(\mathbb{R}^2, \mathbb{R})} \leq C\|v\|_{L^2(K, \mathbb{R})} \quad \text{and} \quad \|\tilde{\mathcal{P}}v\|_{H^1(\mathbb{R}^2, \mathbb{R})} \leq C\|v\|_{H^1(K, \mathbb{R})}.$$

Thus, we get

$$\mathcal{P}u|_K = \tilde{\mathcal{P}}(\operatorname{Re}(u))|_K + i\tilde{\mathcal{P}}(\operatorname{Im}(u))|_K = \operatorname{Re}(u) + i\operatorname{Im}(u) = u,$$

$$\begin{aligned} \|\mathcal{P}u\|_{L^2(\mathbb{R}^2, \mathbb{C})}^2 &= \|\tilde{\mathcal{P}}(\operatorname{Re}(u)) + i\tilde{\mathcal{P}}(\operatorname{Im}(u))\|_{L^2(\mathbb{R}^2, \mathbb{C})}^2 \leq C(\|\operatorname{Re}(u)\|_{L^2(K, \mathbb{R})}^2 + \|\operatorname{Im}(u)\|_{L^2(K, \mathbb{R})}^2) \\ &= C\|u\|_{L^2(K, \mathbb{C})}^2 \end{aligned}$$

and, analogously,

$$\begin{aligned} \|\mathcal{P}u\|_{H^1(\mathbb{R}^2, \mathbb{C})}^2 &= \|\tilde{\mathcal{P}}(\operatorname{Re}(u)) + i\tilde{\mathcal{P}}(\operatorname{Im}(u))\|_{H^1(\mathbb{R}^2, \mathbb{C})}^2 \leq C(\|\operatorname{Re}(u)\|_{H^1(K, \mathbb{R})}^2 + \|\operatorname{Im}(u)\|_{H^1(K, \mathbb{R})}^2) \\ &= C\|u\|_{H^1(K, \mathbb{C})}^2. \end{aligned}$$

Let now $\mathcal{H} := \mathcal{P}(\mathcal{F}|_K)$.

Since H_ε is continuously embedded in $H^1(K, \mathbb{C})$, we have that \mathcal{H} is bounded in $H^1(\mathbb{R}^2, \mathbb{C})$ and $\mathcal{H}|_K = \mathcal{F}|_K$.

Thus we can proceed in a classical way, see e.g. [12, Theorem 9.16], using [12, Theorem 4.26].

Indeed, first observe that, since K is bounded, we may assume $r \geq 2$.

Then, let $u \in C_c^\infty(\mathbb{R}^2, \mathbb{C})$ and $\tau_h u := u(\cdot + h)$, with $h \in \mathbb{R}^2$.

For every $x \in \mathbb{R}^2$ we have

$$\begin{aligned} |\tau_h u(x) - u(x)| &= \sqrt{[\operatorname{Re}(u)(x+h) - \operatorname{Re}(u)(x)]^2 + [\operatorname{Im}(u)(x+h) - \operatorname{Im}(u)(x)]^2} \\ &\leq |\operatorname{Re}(u)(x+h) - \operatorname{Re}(u)(x)| + |\operatorname{Im}(u)(x+h) - \operatorname{Im}(u)(x)| \\ &\leq \int_0^1 |v'_R(t)| dt + \int_0^1 |v'_I(t)| dt \\ &= \int_0^1 |h \cdot \nabla \operatorname{Re}(u)(x+th)| dt + \int_0^1 |h \cdot \nabla \operatorname{Im}(u)(x+th)| dt \\ &\leq |h| \left(\int_0^1 (|\nabla \operatorname{Re}(u)(x+th)| + |\nabla \operatorname{Im}(u)(x+th)|) dt \right) \\ &\leq \sqrt{2}|h| \int_0^1 |\nabla u(x+th)| dt \end{aligned}$$

where $v_R(t) := \operatorname{Re}(u)(x+th)$ and $v_I(t) := \operatorname{Im}(u)(x+th)$, and so, using the Hölder inequality,

$$|\tau_h u(x) - u(x)|^2 \leq 2|h|^2 \int_0^1 |\nabla u(x+th)|^2 dt.$$

Hence, using the density of $C_c^\infty(\mathbb{R}^2, \mathbb{C})$ in $H^1(\mathbb{R}^2, \mathbb{C})$, see [25, Proposition 2.1],

$$\|\tau_h u - u\|_2 \leq C|h| \|\nabla u\|_2 \leq C|h|$$

uniformly with respect to $u \in \mathcal{H}$.

For $r > 2$, if we take $q > r$, there exists $\alpha \in (0, 1)$ such that

$$\frac{1}{r} = \frac{\alpha}{2} + \frac{1-\alpha}{q},$$

so that, for every $u \in \mathcal{H}$,

$$\|\tau_h u - u\|_r \leq \|\tau_h u - u\|_2^\alpha \|\tau_h u - u\|_q^{1-\alpha} \leq C|h|^\alpha$$

with C independent of the elements of \mathcal{H} .

Thus, [12, Theorem 4.26] implies that the closure of $\mathcal{H}|_K = \mathcal{F}|_K$ is compact in $L^r(K, \mathbb{C})$.

Finally, if $V_\infty = +\infty$, let $\{u_n\} \subset H_\varepsilon$ be a sequence weakly convergent to u in H_ε and $v_n := u_n - u$.

We have that $v_n \rightarrow 0$ in $L^2(\mathbb{R}^2, \mathbb{C})$.

Indeed, for every $\eta > 0$, there exists $\sigma > 0$ large enough such that

$$\frac{1}{V(x)} < \eta \text{ in } B_\sigma^c(0).$$

Moreover, since $v_n \rightarrow 0$ in H_ε , then $v_n \rightarrow 0$ in $L_{\text{loc}}^2(\mathbb{R}^2, \mathbb{C})$. Thus, for n large enough,

$$\|v_n\|_2^2 < \eta + \frac{\eta}{V_0} \int_{B_{\sigma/\varepsilon}^c(0)} V_\varepsilon(x) |v_n|^2 dx < C\eta.$$

Hence, using the Gagliardo-Nirenberg inequality applied to $|v_n|$, for every $r \geq 2$ we have

$$\|v_n\|_r \leq C \|\nabla |v_n|\|_2^{1-2/r} \|v_n\|_2^{2/r},$$

and, by the diamagnetic inequality (2.1) and the boundedness of $\{v_n\}$ in H_ε , we conclude. \square

For compact supported functions in $H^1(\mathbb{R}^2, \mathbb{R})$, we have the following result which will be very important for some estimates below.

Lemma 2.2. *If $u \in H^1(\mathbb{R}^2, \mathbb{R})$ and u has a compact support, then $\omega := e^{iA(0) \cdot x} u \in H_\varepsilon$.*

Proof. Assume that $\text{supp}(u) \subset B_R(0)$. Since V is continuous, it is clear that

$$\int_{\mathbb{R}^2} V_\varepsilon(x) |\omega|^2 dx = \int_{B_R(0)} V_\varepsilon(x) |\omega|^2 dx \leq C \|u\|_2^2 < +\infty.$$

Moreover, since V and A are continuous, we have

$$\begin{aligned} \int_{\mathbb{R}^2} |\nabla_{A_\varepsilon} \omega|^2 dx &= \int_{\mathbb{R}^2} |\nabla \omega|^2 dx + \int_{\mathbb{R}^2} |A_\varepsilon(x)|^2 |\omega|^2 dx + 2\text{Re} \int_{\mathbb{R}^2} i A_\varepsilon(x) \bar{\omega} \nabla \omega dx \\ &\leq 2 \int_{\mathbb{R}^2} |\nabla \omega|^2 dx + 2 \int_{\mathbb{R}^2} |A_\varepsilon(x)|^2 |\omega|^2 dx \\ &\leq C \left[\int_{\mathbb{R}^2} |\nabla u|^2 dx + \int_{\mathbb{R}^2} |u|^2 dx \right] < +\infty \end{aligned}$$

and we conclude. \square

A useful tool when we work with nonlinearities with exponential critical growth is the following version of Trudinger-Moser inequality as stated e.g. in [1, Lemma 1.2] (see also [13, Lemma 2.1]).

Lemma 2.3. *If $\alpha > 0$ and $u \in H^1(\mathbb{R}^2, \mathbb{R})$, then*

$$\int_{\mathbb{R}^2} (e^{\alpha|u|^2} - 1) dx < +\infty.$$

Moreover, if $\|\nabla u\|_2^2 \leq 1$, $\|u\|_2 \leq M < +\infty$, and $0 < \alpha < 4\pi$, then there exists a positive constant $C(M, \alpha)$, which depends only on M and α , such that

$$\int_{\mathbb{R}^2} (e^{\alpha|u|^2} - 1) dx \leq C(M, \alpha).$$

Observe also that that (f_1) and (f_2) imply that, fixed $q > 2$, for any $\zeta > 0$ and $\alpha > 4\pi$, there exists a constant $C > 0$, which depends on q , α , ζ , such that

$$(2.4) \quad f(t) \leq \zeta + Ct^{(q-2)/2}(e^{\alpha t} - 1), \quad \text{for all } t \geq 0$$

and, using (f_3) ,

$$(2.5) \quad F(t) \leq \zeta t + Ct^{q/2}(e^{\alpha t} - 1), \quad \text{for all } t \geq 0.$$

Then, by (2.4) and (2.5), we have

$$(2.6) \quad f(t^2)t^2 \leq \zeta t^2 + C|t|^q(e^{\alpha t^2} - 1), \quad \text{for all } t \in \mathbb{R}$$

and

$$(2.7) \quad F(t^2) \leq \zeta t^2 + C|t|^q(e^{\alpha t^2} - 1), \quad \text{for all } t \in \mathbb{R}.$$

Moreover (f_4) and (f_5) imply that there exist two positive constants C_1 and C_2 such that

$$C_1 t^{(p-4)/2} \leq f'(t) \leq C_2 t, \quad \text{as } t \rightarrow 0^+$$

and so $p > 6$.

For $u \in H_\varepsilon$, let

$$(2.8) \quad \hat{u}_j := \varphi_j u,$$

where $j \in \mathbb{N}^*$ and $\varphi_j(x) = \varphi(2x/j)$ with $\varphi \in C_c^\infty(\mathbb{R}^2, \mathbb{R})$, $0 \leq \varphi \leq 1$, $\varphi(x) = 1$ if $|x| \leq 1$, and $\varphi(x) = 0$ if $|x| \geq 2$.

Note that \hat{u}_j belongs to H_ε and has compact support.

We have that any $u \in H_\varepsilon$ can be *approximated* by \hat{u}_j in the following sense. This property is classical, but, for completeness, we give here some details.

Lemma 2.4. *Let $\varepsilon > 0$. For any $u \in H_\varepsilon$, $\|\hat{u}_j - u\|_\varepsilon \rightarrow 0$ as $j \rightarrow +\infty$.*

Proof. Since $u \in H_\varepsilon$, by the Lebesgue Dominated Convergence Theorem, it is easy to see that

$$\int_{\mathbb{R}^2} V_\varepsilon(x) |\hat{u}_j - u|^2 dx = o_j(1).$$

Moreover, since

$$\begin{aligned} \int_{\mathbb{R}^2} |\nabla_{A_\varepsilon}(\hat{u}_j - u)|^2 dx &= \int_{\mathbb{R}^2} |\nabla(\hat{u}_j - u)|^2 dx + \int_{\mathbb{R}^2} |A_\varepsilon(x)|^2 |\hat{u}_j - u|^2 dx \\ &\quad + 2\operatorname{Re} \int_{\mathbb{R}^2} i \nabla(\hat{u}_j - u) A_\varepsilon(x) \overline{(\hat{u}_j - u)} dx \\ &\leq 2 \int_{\mathbb{R}^2} |\nabla(\hat{u}_j - u)|^2 dx + 2 \int_{\mathbb{R}^2} |A_\varepsilon(x)|^2 |\hat{u}_j - u|^2 dx, \end{aligned}$$

applying again the Lebesgue Dominated Convergence Theorem, we conclude. \square

In the spirit of [24, Lemma 3.2], we have the following result for bounded sequences in H_ε .

Lemma 2.5. *Let $s \geq 2$ and $\{u_n\} \subset H_\varepsilon$ be a bounded sequence. Then there exists a subsequence $\{u_{n_j}\}$ such that for any $\sigma > 0$ there is $r_{\sigma,s} > 0$ such that, for every $r \geq r_{\sigma,s}$,*

$$\limsup_j \int_{B_j(0) \setminus B_r(0)} |u_{n_j}|^s dx \leq \sigma.$$

Proof. By Lemma 2.1, for each $j \in \mathbb{N}$,

$$\int_{B_j(0)} |u_n|^s dx \rightarrow \int_{B_j(0)} |u|^s dx \text{ as } n \rightarrow +\infty.$$

Thus there exists $\nu_j \in \mathbb{N}$ such that

$$\left| \int_{B_j(0)} (|u_n|^s - |u|^s) dx \right| < \frac{1}{j} \quad \text{for all } n > \nu_j$$

and, without loss of generality, we can assume that $\nu_{j+1} \geq \nu_j$.

In particular, for $n_j := \nu_j + j$ we have

$$\int_{B_j(0)} (|u_{n_j}|^s - |u|^s) dx < \frac{1}{j}.$$

On the other hand, there exists $r_{\sigma,s}$ such that, for every $r \geq r_{\sigma,s}$,

$$\int_{\mathbb{R}^2 \setminus B_r(0)} |u|^s \leq \frac{\sigma}{3}.$$

Hence, for j large enough,

$$\begin{aligned} \int_{B_j(0) \setminus B_r(0)} |u_{n_j}|^s dx &= \int_{B_j(0)} (|u_{n_j}|^s - |u|^s) dx + \int_{B_j(0) \setminus B_r(0)} |u|^s dx + \int_{B_r(0)} (|u|^s - |u_{n_j}|^s) dx \\ &\leq \frac{1}{j} + \frac{\sigma}{3} + \frac{1}{j} \leq \sigma. \end{aligned}$$

□

In the proofs of our results, the following properties will be useful.

Lemma 2.6. *Let $\{u_n\} \subset H_\varepsilon$ be a bounded sequence and $\{u_{n_j}\}$ be a subsequence of $\{u_n\}$ such that $u_{n_j} \rightharpoonup u$ in H_ε and $\nabla|u_{n_j}| \rightarrow \nabla|u|$ a.e. in \mathbb{R}^2 , as $j \rightarrow +\infty$. Moreover let $\ell^2 := \limsup_j \|\nabla|u_{n_j}|\|_2^2$ and \hat{u}_j be defined as in (2.8).*

Then, up to a subsequence still denoted by $\{u_{n_j}\}$, we have that, if $\ell^2 < 1/4$,

$$(2.9) \quad \int_{\mathbb{R}^2} \left(F(|u_{n_j}|^2) - F(|u_{n_j} - \hat{u}_j|^2) - F(|\hat{u}_j|^2) \right) dx = o_j(1)$$

and, for every $\phi \in H_\varepsilon$, if $\ell^2 < 1/8$,

$$(2.10) \quad \int_{\mathbb{R}^2} \left| f(|u_{n_j}|^2)u_{n_j} - f(|u_{n_j} - \hat{u}_j|^2)(u_{n_j} - \hat{u}_j) - f(|\hat{u}_j|^2)\hat{u}_j \right|^2 = o_j(1).$$

Proof. Let us prove (2.9).

Using (2.4) and the Young inequality, we have that, fixed $q > 2$, for any $\zeta > 0$ and $\alpha > 4\pi$, there exists

a constant $C > 0$, which depends on q, α, ζ , such that

$$\begin{aligned} |F(|u_{n_j}|^2) - F(|u_{n_j} - \hat{u}_j|^2)| &\leq 2 \int_0^1 |f(|u_{n_j} - t\hat{u}_j|^2)| |u_{n_j} - t\hat{u}_j| |\hat{u}_j| dt \\ &\leq \zeta \int_0^1 |u_{n_j} - t\hat{u}_j| |\hat{u}_j| dt + C \int_0^1 |u_{n_j} - t\hat{u}_j|^{q-1} (e^{\alpha|u_{n_j} - t\hat{u}_j|^2} - 1) |\hat{u}_j| dt \\ &\leq \zeta(|u_{n_j}| + |u|)|u| + C(|u_{n_j}| + |u|)^{q-1} (e^{\alpha(|u_{n_j}| + |u|)^2} - 1)|u| \\ &\leq \zeta(|u_{n_j}|^2 + |u_{n_j}|^q (e^{\alpha(|u_{n_j}| + |u|)^2} - 1)) + C(|u|^2 + |u|^q (e^{\alpha(|u_{n_j}| + |u|)^2} - 1)). \end{aligned}$$

Then, by (2.7) we have

$$(2.11) \quad \begin{aligned} |F(|u_{n_j}|^2) - F(|u_{n_j} - \hat{u}_j|^2) - F(|\hat{u}_j|^2)| &\leq \zeta(|u_{n_j}|^2 + |u_{n_j}|^q (e^{\alpha(|u_{n_j}| + |u|)^2} - 1)) \\ &\quad + C(|u|^2 + |u|^q (e^{\alpha(|u_{n_j}| + |u|)^2} - 1)). \end{aligned}$$

Now, let

$$G_j^\zeta := \max \left\{ |F(|u_{n_j}|^2) - F(|u_{n_j} - \hat{u}_j|^2) - F(|\hat{u}_j|^2)| - \zeta(|u_{n_j}|^2 + |u_{n_j}|^q (e^{\alpha(|u_{n_j}| + |u|)^2} - 1)), 0 \right\}.$$

Note that, by (2.11), we have

$$(2.12) \quad G_j^\zeta \leq C(|u|^2 + |u|^q (e^{\alpha(|u_{n_j}| + |u|)^2} - 1))$$

and, by Lemma 2.4 and Lemma 2.1,

$$(2.13) \quad G_j^\zeta \rightarrow 0 \text{ a.e. in } \mathbb{R}^2$$

and

$$(2.14) \quad e^{\alpha(|u_{n_j}| + |u|)^2} \rightarrow e^{4\alpha|u|^2} \text{ a.e. in } \mathbb{R}^2$$

as $j \rightarrow +\infty$.

Let us show that

$$(2.15) \quad |u|^q (e^{\alpha(|u_{n_j}| + |u|)^2} - 1) \rightarrow |u|^q (e^{4\alpha|u|^2} - 1) \text{ in } L^1(\mathbb{R}^2, \mathbb{R}).$$

Since $\nabla|u_{n_j}| \rightarrow \nabla|u|$ a.e. in \mathbb{R}^2 as $j \rightarrow +\infty$, by the Fatou's Lemma, we deduce that

$$\|\nabla|u|\|_2^2 \leq \liminf_j \|\nabla|u_{n_j}|\|_2^2 \leq \ell^2$$

and so, for any fixed $\kappa \in (0, 1 - 4\ell^2)$, we have that for j large enough,

$$(2.16) \quad \|\nabla(|u_{n_j}| + |u|)\|_2^2 \leq 2\|\nabla|u_{n_j}|\|_2^2 + 2\|\nabla|u|\|_2^2 < 4\ell^2 + \kappa.$$

Let

$$p^* \in \left(1, \frac{1}{4\ell^2 + \kappa}\right), \quad \alpha \in \left(4\pi, \frac{4\pi}{p^*(4\ell^2 + \kappa)}\right).$$

By the inequality

$$(2.17) \quad (e^t - 1)^s \leq e^{ts} - 1, \text{ for } s > 1 \text{ and } t \geq 0,$$

inequality (2.16), and Lemma 2.3, we get

$$(2.18) \quad \int_{\mathbb{R}^2} (e^{\alpha(|u_{n_j}| + |u|)^2} - 1)^{p^*} dx \leq \int_{\mathbb{R}^2} (e^{(4\ell^2 + \kappa)p^*\alpha \left(\frac{|u_{n_j}| + |u|}{\sqrt{4\ell^2 + \kappa}}\right)^2} - 1) dx \leq C.$$

Thus, up to a subsequence, and applying the Severini-Egoroff Theorem [12, Theorem 4.29],

$$(e^{\alpha(|u_{n_j}| + |u|)^2} - 1) \rightharpoonup (e^{4\alpha|u|^2} - 1) \text{ in } L^{p^*}(\mathbb{R}^2, \mathbb{R})$$

and, since $|u|^q \in L^{q^*}(\mathbb{R}^2)$, where q^* is the conjugate exponent of p^* , we get (2.15).

Hence, by (2.12), (2.13), (2.14), and (2.15), applying a variant of the Lebesgue Dominated Convergence Theorem, we deduce that

$$\int_{\mathbb{R}^2} G_j^\zeta dx \rightarrow 0 \quad \text{as } j \rightarrow +\infty.$$

On the other hand, by the definition of G_j^ζ ,

$$|F(|u_{n_j}|^2) - F(|u_{n_j} - \hat{u}_j|^2) - F(|\hat{u}_j|^2)| \leq \zeta(|u_{n_j}|^2 + |u_{n_j}|^q(e^{\alpha(|u_{n_j}|+|u|)^2} - 1)) + G_j^\zeta.$$

Using the Hölder inequality, the boundedness of the sequence $\{u_n\}$, Lemma 2.1, and (2.18), we have

$$\int_{\mathbb{R}^2} \left(|u_{n_j}|^q(e^{\alpha(|u_{n_j}|+|u|)^2} - 1) \right) dx \leq \|u_{n_j}\|_{qq^*}^q \left[\int_{\mathbb{R}^2} |(e^{\alpha(|u_{n_j}|+|u|)^2} - 1)^{p^*} dx \right]^{\frac{1}{p^*}} \leq C.$$

Thus, the arbitrariness of ζ allows to conclude the proof of (2.9).

Now we prove (2.10).

First observe that, by Lemma 2.1 and Lemma 2.4, we have that

$$\left| f(|u_{n_j}|^2)u_{n_j} - f(|u_{n_j} - \hat{u}_j|^2)(u_{n_j} - \hat{u}_j) - f(|\hat{u}_j|^2)\hat{u}_j \right| \rightarrow 0 \text{ a.e. in } \mathbb{R}^2.$$

Moreover, using also (2.4), we have that fixed $q > 2$, for any $\zeta > 0$ and $\alpha > 4\pi$, there exists a constant $C > 0$, which depends on q, α, ζ , such that

$$\begin{aligned} & \left| f(|u_{n_j}|^2)u_{n_j} - f(|u_{n_j} - \hat{u}_j|^2)(u_{n_j} - \hat{u}_j) - f(|\hat{u}_j|^2)\hat{u}_j \right| \\ & \leq \zeta|u_{n_j}| + C|u_{n_j}|^{q-1}(e^{\alpha|u_{n_j}|^2} - 1) + \zeta|u_{n_j} - \hat{u}_j| + C|u_{n_j} - \hat{u}_j|^{q-1}(e^{\alpha|u_{n_j} - \hat{u}_j|^2} - 1) \\ & \quad + \zeta|\hat{u}_j| + C|\hat{u}_j|^{q-1}(e^{\alpha|\hat{u}_j|^2} - 1) \\ & \leq C \left[|u_{n_j}| + |u| + (|u_{n_j}| + |u|)^{q-1}(e^{\alpha(|u_{n_j}|+|u|)^2} - 1) \right] \end{aligned}$$

and so

$$\begin{aligned} (2.19) \quad & \left| f(|u_{n_j}|^2)u_{n_j} - f(|u_{n_j} - \hat{u}_j|^2)(u_{n_j} - \hat{u}_j) - f(|\hat{u}_j|^2)\hat{u}_j \right|^2 \\ & \leq C \left[|u_{n_j}|^2 + |u|^2 + (|u_{n_j}| + |u|)^{2(q-1)}(e^{\alpha(|u_{n_j}|+|u|)^2} - 1)^2 \right]. \end{aligned}$$

Arguing as before, for any fixed $\kappa \in (0, \frac{1}{2} - 4\ell^2)$, if

$$p^* \in \left(1, \frac{1}{2(4\ell^2 + \kappa)}\right) \text{ and } \alpha \in \left(4\pi, \frac{2\pi}{p^*(4\ell^2 + \kappa)}\right),$$

we have that

$$(2.20) \quad \int_{\mathbb{R}^2} (e^{\alpha(|u_{n_j}|+|u|)^2} - 1)^{2p^*} dx \leq \int_{\mathbb{R}^2} (e^{2(4\ell^2 + \kappa)p^* \alpha \left(\frac{|u_{n_j}|+|u|}{\sqrt{4\ell^2 + \kappa}}\right)^2} - 1) dx \leq C.$$

By Lemma 2.5 for $s = 2$, from $\{u_{n_j}\}$ we can extract a first subsequence, still denoted by $\{u_{n_j}\}$, such that for any $\sigma > 0$ there is $r_{\sigma,2} > 0$ such that, for every $r \geq r_{\sigma,2}$,

$$(2.21) \quad \limsup_j \int_{B_j(0) \setminus B_r(0)} |u_{n_j}|^2 dx \leq \sigma.$$

Applying again Lemma 2.5 for $s = 2(q-1)q^*$, where q^* is the conjugate exponent of p^* , from $\{u_{n_j}\}$ we can extract a further subsequence, still denoted by $\{u_{n_j}\}$, such that for any $\sigma > 0$ there is $r_{\sigma,2(q-1)q^*} > 0$

such that, for every $r \geq r_{\sigma, 2(q-1)q^*}$,

$$(2.22) \quad \limsup_j \int_{B_j(0) \setminus B_r(0)} |u_{n_j}|^{2(q-1)q^*} dx \leq \sigma.$$

Let now $\sigma > 0$ be arbitrary and $r \geq \max\{r_{\sigma, 2}, r_{\sigma, 2(q-1)q^*}\}$ large enough such that

$$(2.23) \quad \int_{B_r^c(0)} |u|^2 dx \leq \sigma \text{ and } \int_{B_r^c(0)} |u|^{2(q-1)q^*} dx \leq \sigma.$$

Since, $2(q-1)q^* > 2$ and by Lemma 2.1, there exists $w \in L^{2(q-1)q^*}(B_r(0))$ such that $|u_{n_j}| \leq w$ a.e. in $B_r(0)$.

Thus, by (2.19), for every $s \geq 1$ and a.e. $x \in B_r(0)$,

$$\begin{aligned} & \left| f(|u_{n_j}|^2)u_{n_j} - f(|u_{n_j} - \hat{u}_j|^2)(u_{n_j} - \hat{u}_j) - f(|\hat{u}_j|^2)\hat{u}_j \right|^2 \\ & \leq C \left[w^2 + |u|^2 + (w + |u|)^{2(q-1)}(e^{\alpha(|u_{n_j}| + |u|)^2} - 1)^2 \right]. \end{aligned}$$

Then, by (2.20), applying the Hölder inequality,

$$\int_{B_r(0)} (w + |u|)^{2(q-1)}(e^{\alpha(|u_{n_j}| + |u|)^2} - 1)^2 dx \leq C \left(\int_{B_r(0)} (w + |u|)^{2(q-1)q^*} dx \right)^{1/q^*},$$

and so

$$w^2 + |u|^2 + (w + |u|)^{2(q-1)}(e^{\alpha(|u_{n_j}| + |u|)^2} - 1)^2 \in L^1(B_r(0)).$$

Hence, applying the Lebesgue Dominated Convergence Theorem, we deduce that

$$\int_{B_r(0)} \left| f(|u_{n_j}|^2)u_{n_j} - f(|u_{n_j} - \hat{u}_j|^2)(u_{n_j} - \hat{u}_j) - f(|\hat{u}_j|^2)\hat{u}_j \right|^2 dx \rightarrow 0 \text{ as } j \rightarrow +\infty.$$

On the other hand, by (2.19), (2.20), (2.21), (2.22), and (2.23), for j large enough,

$$\begin{aligned} & \int_{B_j(0) \setminus B_r(0)} \left| f(|u_{n_j}|^2)u_{n_j} - f(|u_{n_j} - \hat{u}_j|^2)(u_{n_j} - \hat{u}_j) - f(|\hat{u}_j|^2)\hat{u}_j \right|^2 dx \\ & \leq C \int_{B_j(0) \setminus B_r(0)} \left(|u_{n_j}|^2 + |u|^2 + (|u_{n_j}| + |u|)^{2(q-1)}(e^{2\alpha(|u_{n_j}| + |u|)^2} - 1) \right) dx \\ & \leq C \left[\sigma + \left(\int_{B_j(0) \setminus B_r(0)} |u_{n_j}|^{2(q-1)q^*} dx \right)^{1/q^*} + \left(\int_{B_j(0) \setminus B_r(0)} |u|^{2(q-1)q^*} dx \right)^{1/q^*} \right] \\ & \leq C(\sigma + \sigma^{1/q^*}). \end{aligned}$$

Hence, since $\hat{u}_j = 0$ in $B_j^c(0)$ for any $j \geq 1$, the arbitrariness of $\sigma > 0$ allows to conclude. \square

3. THE LIMIT PROBLEM

As it is usual for this type of questions, we consider the *limit* problem

$$(3.1) \quad -\Delta u + \mu u = f(u^2)u, \quad u : \mathbb{R}^2 \rightarrow \mathbb{R}, \quad \mu > 0.$$

We have already considered this problem in [20, Section 2] for a particular value of μ , namely for $\mu = V_0$. Here, due to the assumptions on V , we need some generalizations.

First of all we observe that solutions of (3.1) can be found as critical points of the C^1 functional

$$I_\mu(u) := \frac{1}{2} \int_{\mathbb{R}^2} (|\nabla u|^2 + \mu u^2) dx - \frac{1}{2} \int_{\mathbb{R}^2} F(u^2) dx$$

defined in $H^1(\mathbb{R}^2, \mathbb{R})$. Moreover, if

$$\mathcal{N}_\mu := \{u \in H^1(\mathbb{R}^2, \mathbb{R}) \setminus \{0\} : I'_\mu(u)[u] = 0\} \quad \text{and} \quad c_\mu := \inf_{u \in \mathcal{N}_\mu} I_\mu(u),$$

using (f₁) and (f₄), we have that for each fixed $u \in H^1(\mathbb{R}^2, \mathbb{R}) \setminus \{0\}$, there is a unique $t(u) > 0$ such that

$$(3.2) \quad I_\mu(t(u)u) = \max_{t \geq 0} I_\mu(tu) \quad \text{and} \quad t(u)u \in \mathcal{N}_\mu.$$

Then, arguing as in [33, Lemma 4.1 and Theorem 4.2], we also have that

$$(3.3) \quad 0 < c_\mu = \inf_{u \in H^1(\mathbb{R}^2, \mathbb{R}) \setminus \{0\}} \max_{t \geq 0} I_\mu(tu).$$

Observe finally that, arguing as in [20, Lemma 2.3] where we consider only the case $\mu = V_0$, we can get that there exists a constant $K > 0$ such that, for all $u \in \mathcal{N}_\mu$, $\|u\|_\mu \geq K$.

In particular, if $\mu \in (0, V^*]$, the assumptions on f allow to get the existence of a positive *ground state* solution ω_μ of (3.1), namely such that $I_\mu(\omega_\mu) \leq I_\mu(v)$ for all positive solutions $v \in H^1(\mathbb{R}^2, \mathbb{R})$ of (3.1).

Lemma 3.1. *For every $\mu \in (0, V^*]$, there exists a positive and radially symmetric ground state solution $\omega_\mu \in H^1(\mathbb{R}^2, \mathbb{R})$ for problem (3.1).*

Proof. We can argue exactly as in [20, Lemma 2.2], applying [6, Corollary 1.5] to the function $h(t) := f(t^2)t/\mu$. Here we observe only that the condition

$$\text{there exist } \lambda > 0 \text{ and } p > 2 \text{ such that } h(t) \geq \lambda t^{p-1} \text{ for } t \geq 0 \text{ and } \lambda > \left(\frac{p-2}{p}\right)^{\frac{p-2}{2}} S_p^{p/2}$$

follows by (f₄), since

$$h(t) \geq \frac{C_p}{\mu} t^{p-1} \quad \text{for all } t \geq 0$$

and

$$\frac{C_p}{\mu} > \left(\frac{8\theta}{\theta-2}\right)^{\frac{p-2}{2}} \left(\frac{p-2}{p}\right)^{\frac{p-2}{2}} S_p^{p/2} > 8^{\frac{p-2}{2}} \left(\frac{p-2}{p}\right)^{\frac{p-2}{2}} S_p^{p/2} > \left(\frac{p-2}{p}\right)^{\frac{p-2}{2}} S_p^{p/2}.$$

By standard arguments we can assume ω_μ positive. We refer to [20, Lemma 2.2] for the remaining details. \square

Moreover the ground state ω_μ decays exponentially at infinity with its gradient and belongs to $C^2(\mathbb{R}^2, \mathbb{R}) \cap L^\infty(\mathbb{R}^2, \mathbb{R})$ (see also [34, Proposition 2.1]) and, for any $\mu_1, \mu_2 \in (0, V^*]$ with $0 < \mu_1 < \mu_2$, we have

$$(3.4) \quad c_{\mu_1} = I_{\mu_1}(\omega_{\mu_1}) \leq \max_{t \geq 0} I_{\mu_1}(t\omega_{\mu_2}) < \max_{t \geq 0} I_{\mu_2}(t\omega_{\mu_2}) = I_{\mu_2}(\omega_{\mu_2}) = c_{\mu_2}.$$

Let now $S_p > 0$ be defined as in (f₄) and, for every $\mu > 0$,

$$S_p^\mu := \inf_{u \in H^1(\mathbb{R}^2, \mathbb{R}) \setminus \{0\}} \frac{\|\nabla u\|_2^2 + \mu \|u\|_2^2}{\|u\|_p^2}.$$

We have

Lemma 3.2. $S_p^\mu = \mu^{2/p} S_p$.

Proof. Let $u \in H^1(\mathbb{R}^2, \mathbb{R}) \setminus \{0\}$.

If $\tilde{u} := u(\mu^{-1/2} \cdot)$, we have

$$\frac{\|\nabla u\|_2^2 + \mu \|u\|_2^2}{\|u\|_p^2} = \mu^{2/p} \frac{\|\nabla \tilde{u}\|_2^2 + \|\tilde{u}\|_2^2}{\|\tilde{u}\|_p^2} \geq \mu^{2/p} S_p$$

and, passing to the infimum, we get $S_p^\mu \geq \mu^{2/p} S_p$.

On the other hand, if $\tilde{u} := u(\mu^{1/2} \cdot)$, we have

$$\frac{\|\nabla u\|_2^2 + \|u\|_2^2}{\|u\|_p^2} = \mu^{-2/p} \frac{\|\nabla \tilde{u}\|_2^2 + \mu \|\tilde{u}\|_2^2}{\|\tilde{u}\|_p^2} \geq \mu^{-2/p} S_p^\mu$$

and, passing to the infimum, we get $S_p^\mu \leq \mu^{2/p} S_p$, concluding the proof. \square

Remark 3.3. Observe that, by [33, Theorem 1.35], S_p is achieved by a C^2 positive and radially symmetric function, and, arguing as in the proof of Lemma 3.2, this holds for S_p^μ too.

Now we give an upper estimate of the ground state energy for (3.1) with $\mu = V^*$, with V^* defined in (1.2).

Lemma 3.4. The minimax level c_{V^*} belongs to the interval $(0, \frac{\theta - 2}{16\theta})$.

Proof. As observed in Remark 3.3, there exists $\omega^* \in H^1(\mathbb{R}^2, \mathbb{R}) \setminus \{0\}$ such that

$$S_p^{V^*} = \frac{\|\nabla \omega^*\|_2^2 + V^* \|\omega^*\|_2^2}{\|\omega^*\|_p^2}.$$

Thus, using (3.3), (f₄), and Lemma 3.2, we have

$$\begin{aligned} 0 < c_{V^*} &\leq \max_{t \geq 0} I_{V^*}(t\omega^*) \leq \max_{t \geq 0} \left\{ \frac{t^2}{2} (\|\nabla \omega^*\|_2^2 + V^* \|\omega^*\|_2^2) - \frac{C_p t^p}{p} \|\omega^*\|_p^p \right\} \\ &= \frac{p-2}{2p} \left[\frac{(S_p^{V^*})^p}{C_p^2} \right]^{\frac{1}{p-2}} = \frac{p-2}{2p} \left[\frac{(V^*)^2 S_p^p}{C_p^2} \right]^{\frac{1}{p-2}} < \frac{\theta - 2}{16\theta}. \end{aligned}$$

\square

Finally we give the following result.

Lemma 3.5. Let $\mu \in (0, V^*]$, and $\{\omega_n\} \subset \mathcal{N}_\mu$ be a sequence of positive functions satisfying $I_\mu(\omega_n) \rightarrow c_\mu$. Then $\{\omega_n\}$ is bounded in $H^1(\mathbb{R}^2, \mathbb{R})$ and, up to a subsequence, $\omega_n \rightharpoonup \omega$ in $H^1(\mathbb{R}^2, \mathbb{R})$.

Moreover we have that:

- (1) if $\omega \neq 0$, then $\omega_n \rightarrow \omega \in \mathcal{N}_\mu$ in $H^1(\mathbb{R}^2, \mathbb{R})$ and ω is a ground state for (3.1);
- (2) if $\omega = 0$, then there exists $\{\tilde{y}_n\} \subset \mathbb{R}^2$ with $|\tilde{y}_n| \rightarrow +\infty$ and $\tilde{\omega} \in \mathcal{N}_\mu$ such that, up to a subsequence, $\omega_n(\cdot + \tilde{y}_n) \rightarrow \tilde{\omega}$ in $H^1(\mathbb{R}^2, \mathbb{R})$ and $\tilde{\omega}$ is a ground state for (3.1).

Proof. By (f₃) and Lemma 3.4, it follows that

$$\begin{aligned} \frac{\theta - 2}{2\theta} \limsup_n (\|\nabla \omega_n\|_2^2 + \mu \|\omega_n\|_2^2) &\leq \limsup_n \left\{ \left(\frac{1}{2} - \frac{1}{\theta} \right) (\|\nabla \omega_n\|_2^2 + \mu \|\omega_n\|_2^2) \right. \\ &\quad \left. + \int_{\mathbb{R}^2} \left(\frac{1}{\theta} f(\omega_n^2) \omega_n^2 - \frac{1}{2} F(\omega_n^2) \right) dx \right\} \\ &= \limsup_n \left\{ I_\mu(\omega_n) - \frac{1}{\theta} I'_\mu(\omega_n)[\omega_n] \right\} = c_\mu \leq c_{V^*} < \frac{\theta - 2}{16\theta}. \end{aligned}$$

Thus,

$$(3.5) \quad \ell^2 := \limsup_n (\|\nabla \omega_n\|_2^2 + \mu \|\omega_n\|_2^2) < \frac{1}{8}$$

and so there exists $\omega \in H^1(\mathbb{R}^2, \mathbb{R})$ such that, up to a subsequence, $\omega_n \rightharpoonup \omega$ in $H^1(\mathbb{R}^2, \mathbb{R})$. Moreover $\omega_n \rightarrow \omega$ in $L_{\text{loc}}^r(\mathbb{R}^2, \mathbb{R})$, for any $r \geq 1$ and $\omega_n \rightarrow \omega$ a.e. in $x \in \mathbb{R}^2$.

Now let us consider two different cases.

Case 1: $\omega \neq 0$.

Observe that, for every $\phi \in C_c^\infty(\mathbb{R}^2, \mathbb{R})$,

$$f(\omega_n^2)\omega_n\phi \rightarrow f(\omega^2)\omega\phi \text{ a.e. in } \mathbb{R}^2 \text{ as } n \rightarrow +\infty$$

and that, by (2.4), we have that for any $\zeta > 0$, $q > 2$, and $\alpha > 4\pi$, there exists $C > 0$ such that

$$(3.6) \quad |f(\omega_n^2)\omega_n\phi| \leq \zeta\omega_n|\phi| + C\omega_n^{q-1}(e^{\alpha\omega_n^2} - 1)|\phi| \leq \zeta\omega_n|\phi| + C(\omega_n^q + |\phi|^q)(e^{\alpha\omega_n^2} - 1),$$

with

$$\zeta\omega_n|\phi| + C(\omega_n^q + |\phi|^q)(e^{\alpha\omega_n^2} - 1) \rightarrow \zeta\omega|\phi| + C(\omega^q + |\phi|^q)(e^{\alpha\omega^2} - 1) \text{ a.e. in } \mathbb{R}^2 \text{ as } n \rightarrow +\infty.$$

Now, taking into account (3.5), let us fix $\kappa \in (0, 1 - 8\ell^2)$, and let

$$p^* \in \left(1, \frac{1}{8\ell^2 + \kappa}\right), \quad \alpha \in \left(4\pi, \frac{4\pi}{p^*(8\ell^2 + \kappa)}\right).$$

If $\Omega := \text{supp}(\phi)$, by [33, Lemma A.1], there exists $v \in L^{qq^*}(\Omega, \mathbb{R})$ such that $\omega_n \leq v$ a.e. in Ω , where $qq^* > 2$ and q^* is the conjugate exponent of p^* and, by (3.6), we have

$$|f(\omega_n^2)\omega_n\phi| \leq \zeta\omega_n|\phi| + C(v^q + |\phi|^q)(e^{\alpha\omega_n^2} - 1) \rightarrow \zeta\omega|\phi| + C(v^q + |\phi|^q)(e^{\alpha\omega^2} - 1) \text{ a.e. in } \Omega.$$

Arguing as in the proof of Lemma 2.6, we have

$$(e^{\alpha\omega_n^2} - 1) \rightharpoonup (e^{\alpha\omega^2} - 1) \text{ in } L^{p^*}(\mathbb{R}^2, \mathbb{R}).$$

Moreover, since $\omega_n \rightharpoonup \omega$ in $L^2(\mathbb{R}^2, \mathbb{R})$ we have that

$$\zeta \int_{\mathbb{R}^2} \omega_n |\phi| dx + C \int_{\mathbb{R}^2} (v^q + |\phi|^q)(e^{\alpha\omega_n^2} - 1) dx \rightarrow \zeta \int_{\mathbb{R}^2} \omega |\phi| dx + C \int_{\mathbb{R}^2} (v^q + |\phi|^q)(e^{\alpha\omega^2} - 1) dx$$

as $n \rightarrow +\infty$.

Hence, a variant of the Lebesgue Dominated Convergence Theorem implies that

$$\int_{\mathbb{R}^2} f(\omega_n^2)\omega_n\phi dx \rightarrow \int_{\mathbb{R}^2} f(\omega^2)\omega\phi dx,$$

and so ω is a nontrivial critical point for I_μ .

Since, by (f3) and the Fatou's Lemma,

$$\int_{\mathbb{R}^2} \left(\frac{1}{\theta} f(\omega^2)\omega^2 - \frac{1}{2} F(\omega^2) \right) dx \leq \liminf_n \int_{\mathbb{R}^2} \left(\frac{1}{\theta} f(\omega_n^2)\omega_n^2 - \frac{1}{2} F(\omega_n^2) \right) dx,$$

we have

$$\begin{aligned} c_\mu &\leq I_\mu(\omega) = I_\mu(\omega) - \frac{1}{\theta} I'_\mu(\omega)[\omega] \\ &= \left(\frac{1}{2} - \frac{1}{\theta} \right) (\|\nabla \omega\|_2^2 + \mu \|\omega\|_2^2) + \int_{\mathbb{R}^2} \left(\frac{1}{\theta} f(\omega^2)\omega^2 - \frac{1}{2} F(\omega^2) \right) dx \\ &\leq \liminf_n \left\{ \left(\frac{1}{2} - \frac{1}{\theta} \right) (\|\nabla \omega_n\|_2^2 + \mu \|\omega_n\|_2^2) + \int_{\mathbb{R}^2} \left(\frac{1}{\theta} f(\omega_n^2)\omega_n^2 - \frac{1}{2} F(\omega_n^2) \right) dx \right\} \\ &= \liminf_n \left\{ I_\mu(\omega_n) - \frac{1}{\theta} I'_\mu(\omega_n)[\omega_n] \right\} = c_\mu \end{aligned}$$

and, by the assumptions,

$$\begin{aligned}
0 &\leq \liminf_n \left[\left(\frac{1}{2} - \frac{1}{\theta} \right) (\|\nabla \omega_n\|_2^2 + \mu \|\omega_n\|_2^2 - \|\nabla \omega\|_2^2 - \mu \|\omega\|_2^2) \right] \\
&\leq \limsup_n \left[\left(\frac{1}{2} - \frac{1}{\theta} \right) (\|\nabla \omega_n\|_2^2 + \mu \|\omega_n\|_2^2 - \|\nabla \omega\|_2^2 - \mu \|\omega\|_2^2) \right] \\
&= \limsup_n \left[I_\mu(\omega_n) - c_\mu + \int_{\mathbb{R}^2} \left(\frac{1}{\theta} f(\omega^2) \omega^2 - \frac{1}{2} F(\omega^2) \right) dx - \int_{\mathbb{R}^2} \left(\frac{1}{\theta} f(\omega_n^2) \omega_n^2 - \frac{1}{2} F(\omega_n^2) \right) dx \right] \\
&= \int_{\mathbb{R}^2} \left(\frac{1}{\theta} f(\omega^2) \omega^2 - \frac{1}{2} F(\omega^2) \right) dx - \liminf_n \left[\int_{\mathbb{R}^2} \left(\frac{1}{\theta} f(\omega_n^2) \omega_n^2 - \frac{1}{2} F(\omega_n^2) \right) dx \right] \leq 0.
\end{aligned}$$

Case 2: $\omega = 0$.

In this case, there exist $R, \eta > 0$, and $(\tilde{y}_n) \subset \mathbb{R}^2$ such that

$$(3.7) \quad \lim_n \int_{B_R(\tilde{y}_n)} \omega_n^2 dx \geq \eta.$$

Indeed, if this does not hold, for any $R > 0$, one has

$$\lim_n \sup_{y \in \mathbb{R}^2} \int_{B_R(y)} \omega_n^2 dx = 0.$$

Thus, by [28, Chapter 6, Lemma 8.4], for every $\tau > 2$,

$$\lim_n \|\omega_n\|_\tau = 0.$$

Moreover, by (2.6), (2.17), Lemma 2.3, and (3.5), and arguing again as in Lemma 2.6, for $0 < \zeta < \mu/2$, $q > 2$ and $\alpha > 4\pi$, there exists $C > 0$ such that

$$\begin{aligned}
\|\nabla \omega_n\|_2^2 + \mu \|\omega_n\|_2^2 &\leq \zeta \|\omega_n\|_2^2 + C \int_{\mathbb{R}^2} \omega_n^q (e^{\alpha \omega_n^2} - 1) dx \\
&\leq \zeta \|\omega_n\|_2^2 + C \|\omega_n\|_{qq^*}^q \left(\int_{\mathbb{R}^2} (e^{p^* \alpha \omega_n^2} - 1) dx \right)^{1/p^*} \\
&\leq \zeta \|\omega_n\|_2^2 + C \|\omega_n\|_{qq^*}^q = \zeta \|\omega_n\|_2^2 + o_n(1).
\end{aligned}$$

Hence $\omega_n \rightarrow 0$ in $H^1(\mathbb{R}^2, \mathbb{R})$ as $n \rightarrow +\infty$, and so $I_\mu(\omega_n) \rightarrow 0$ as $n \rightarrow +\infty$. But this is in contradiction with $I_\mu(\omega_n) \rightarrow c_\mu > 0$ as $n \rightarrow +\infty$.

Now we claim that $|\tilde{y}_n| \rightarrow +\infty$.

Otherwise, there exists $\bar{R} > 0$ such that

$$\lim_n \int_{B_{\bar{R}}(0)} \omega_n^2 dx \geq \eta$$

and so $\omega \neq 0$.

Moreover $\omega_n(\cdot + \tilde{y}_n) \in \mathcal{N}_\mu$ and, the invariance of I_μ and $\|\nabla \cdot\|_2^2 + \mu \|\cdot\|_2^2$ by translations implies

$$I_\mu(\omega_n(\cdot + \tilde{y}_n)) \rightarrow c_\mu \text{ and } \limsup_n (\|\nabla \omega_n(\cdot + \tilde{y}_n)\|_2^2 + \mu \|\omega_n(\cdot + \tilde{y}_n)\|_2^2) < \frac{1}{8}.$$

Thus, arguing as before, there exists $\tilde{\omega} \in H^1(\mathbb{R}^2, \mathbb{R})$ which, by (3.7), is nontrivial, such that

$$\omega_n(\cdot + \tilde{y}_n) \rightharpoonup \tilde{\omega} \quad \text{in } H^1(\mathbb{R}^2, \mathbb{R}).$$

Hence, repeating the same arguments of Case 1, we conclude. \square

4. PROOF OF THEOREM 1.1

The goal of this section is to prove Theorem 1.1.

We say that $u \in H_\varepsilon$ is a weak solution of (2.2), if for any $\phi \in H_\varepsilon$

$$\operatorname{Re} \int_{\mathbb{R}^2} (\nabla_{A_\varepsilon} u \overline{\nabla_{A_\varepsilon} \phi} + V_\varepsilon(x) u \bar{\phi}) dx = \operatorname{Re} \int_{\mathbb{R}^2} f(|u|^2) u \bar{\phi} dx.$$

Such solutions can be found as the critical points of the C^1 functional $J_\varepsilon : H_\varepsilon \rightarrow \mathbb{R}$ given by

$$J_\varepsilon(u) = \frac{1}{2} \int_{\mathbb{R}^2} (|\nabla_{A_\varepsilon} u|^2 + V_\varepsilon(x) |u|^2) dx - \frac{1}{2} \int_{\mathbb{R}^2} F(|u|^2) dx.$$

The functional J_ε satisfies the Mountain Pass geometry, namely we have

Lemma 4.1. *For any $\varepsilon > 0$,*

- (i) *there exist $\beta, r > 0$ such that $J_\varepsilon(u) \geq \beta$ if $\|u\|_\varepsilon = r$;*
- (ii) *there exists $e \in H_\varepsilon$ with $\|e\|_\varepsilon > r$ such that $J_\varepsilon(e) < 0$.*

Proof. By (2.7), the Hölder and Sobolev inequalities, and (2.17), fixed $q > 2$, for any $\zeta > 0$ and $\alpha > 4\pi$, there exists a constant $C > 0$, which depends on q, α, ζ , such that

$$\begin{aligned} \int_{\mathbb{R}^2} F(|u|^2) dx &\leq \zeta \|u\|_2^2 + C \int_{\mathbb{R}^2} |u|^q (e^{\alpha|u|^2} - 1) dx \\ (4.1) \quad &\leq \zeta \|u\|_2^2 + C \|u\|_{2q}^q \left(\int_{\mathbb{R}^2} (e^{\alpha|u|^2} - 1)^2 dx \right)^{1/2} \\ &\leq \zeta \|u\|_2^2 + C \|u\|_\varepsilon^q \left(\int_{\mathbb{R}^2} (e^{2\alpha|u|^2} - 1) dx \right)^{1/2}. \end{aligned}$$

Moreover, by the diamagnetic inequality (2.1) and (V), for any $u \in H_\varepsilon \setminus \{0\}$,

$$\frac{|u|}{\|u\|_\varepsilon} \in H^1(\mathbb{R}^2, \mathbb{R}), \quad \left\| \frac{|u|}{\|u\|_\varepsilon} \right\|_2^2 \leq \frac{1}{V_0}, \quad \left\| \nabla \frac{|u|}{\|u\|_\varepsilon} \right\|_2^2 \leq 1.$$

Thus, for every $u \in H_\varepsilon$ with $\|u\|_\varepsilon = r > 0$ and $r^2 < \pi/\alpha$, using Lemma 2.3, we get that there exists $C > 0$ independent of r such that

$$(4.2) \quad \int_{\mathbb{R}^2} (e^{2\alpha|u|^2} - 1) dx = \int_{\mathbb{R}^2} (e^{2\alpha r^2 \left(\frac{|u|}{\|u\|_\varepsilon}\right)^2} - 1) dx < \int_{\mathbb{R}^2} (e^{2\pi \left(\frac{|u|}{\|u\|_\varepsilon}\right)^2} - 1) dx \leq C.$$

Hence, by (4.1) and (4.2), we get that for any $\zeta > 0$, there exists $C > 0$ independent of r such that

$$J_\varepsilon(u) \geq \frac{1}{2} r^2 - \frac{\zeta}{V_0} \int_{\mathbb{R}^2} V_\varepsilon(x) |u|^2 dx - C r^q \geq \frac{1}{2} \left(1 - \frac{\zeta}{V_0}\right) r^2 - C r^q$$

and so, since $q > 2$, for r small enough we obtain (i).

Finally, property (ii) is an easy consequence of (f₄), since, fixed $\varphi \in C_c^\infty(\mathbb{R}^2, \mathbb{C}) \setminus \{0\}$, we have that

$$J_\varepsilon(t\varphi) \leq \frac{t^2}{2} \|\varphi\|_\varepsilon^2 - \frac{C_p}{p} t^p \|\varphi\|_p^p.$$

□

Let now

$$(4.3) \quad c_\varepsilon := \inf_{\gamma \in \Gamma_\varepsilon} \max_{t \in [0,1]} J_\varepsilon(\gamma(t)),$$

where

$$\Gamma_\varepsilon = \{\gamma \in C([0,1], H_\varepsilon) : \gamma(0) = 0 \text{ and } J_\varepsilon(\gamma(1)) < 0\}.$$

To find ground states and multiple solutions for (1.1), as usual, we consider the Nehari manifold

$$\mathcal{N}_\varepsilon := \{u \in H_\varepsilon \setminus \{0\} : J'_\varepsilon(u)[u] = 0\}.$$

In virtue of (f₄), we can show that for any $u \in H_\varepsilon \setminus \{0\}$ there exists a unique $t_\varepsilon > 0$ such that $t_\varepsilon u \in \mathcal{N}_\varepsilon$ and

$$(4.4) \quad J_\varepsilon(t_\varepsilon u) = \max_{t \geq 0} J_\varepsilon(tu)$$

and, using classical arguments (see e.g. [33, Lemma 4.1 and Theorem 4.2]),

$$(4.5) \quad c_\varepsilon = \inf_{u \in H_\varepsilon \setminus \{0\}} \sup_{t \geq 0} J_\varepsilon(tu) = \inf_{u \in \mathcal{N}_\varepsilon} J_\varepsilon(u).$$

Moreover we have the following useful preliminary property.

Lemma 4.2. *There exist $K_1, K_2 > 0$ such that for every $\varepsilon > 0$ and for every $u \in \mathcal{N}_\varepsilon$*

$$\|u\|_\varepsilon \geq K_1 \text{ and } J_\varepsilon(u) \geq K_2.$$

Proof. Observe that, for every $\varepsilon > 0$, if $u \in \mathcal{N}_\varepsilon$, by (2.6), fixed $q > 2$, for any $\zeta > 0$ and $\alpha > 4\pi$, there exists a constant $C > 0$, which depends on q, α, ζ , such that

$$(4.6) \quad \|u\|_\varepsilon^2 = \int_{\mathbb{R}^2} f(|u|^2)|u|^2 \leq \zeta \|u\|_2^2 + C \int_{\mathbb{R}^2} |u|^q (e^{\alpha|u|^2} - 1) dx.$$

In particular, for $\zeta < V_0/2$, by (V), we have that

$$(4.7) \quad \|u\|_\varepsilon^2 - \zeta \|u\|_2^2 \geq \frac{1}{2} \|u\|_\varepsilon^2 + \frac{1}{2} \int_{\mathbb{R}^2} [V_\varepsilon(x) - V_0] |u|^2 dx \geq \frac{1}{2} \|u\|_\varepsilon^2.$$

By the Hölder inequality and (2.17), we infer that

$$(4.8) \quad \int_{\mathbb{R}^2} |u|^q (e^{\alpha|u|^2} - 1) dx \leq \|u\|_{2q}^q \left(\int_{\mathbb{R}^2} (e^{\alpha|u|^2} - 1)^2 dx \right)^{1/2} \leq C \|u\|_\varepsilon^q \left(\int_{\mathbb{R}^2} (e^{2\alpha|u|^2} - 1) dx \right)^{1/2}.$$

Now, assume by contradiction that for every $n \in \mathbb{N}$ there exist $\varepsilon_n > 0$ and $u_n \in \mathcal{N}_{\varepsilon_n}$ such that $\|u_n\|_{\varepsilon_n} \rightarrow 0$ as $n \rightarrow +\infty$. Then, using the diamagnetic inequality (2.1) and (V), also $\|\nabla|u_n|\|_2^2 + V_0\|u_n\|_2^2 \rightarrow 0$ as $n \rightarrow +\infty$ and so, for n large enough, using Lemma 2.3, for some $\bar{\alpha} \in (0, 4\pi)$,

$$(4.9) \quad \begin{aligned} \int_{\mathbb{R}^2} (e^{2\alpha|u_n|^2} - 1) dx &= \int_{\mathbb{R}^2} (e^{2\alpha(\|\nabla|u_n|\|_2^2 + V_0\|u_n\|_2^2)} \left(\frac{|u_n|}{\sqrt{\|\nabla|u_n|\|_2^2 + V_0\|u_n\|_2^2}} \right)^2 - 1) dx \\ &\leq \int_{\mathbb{R}^2} (e^{\bar{\alpha} \left(\frac{|u_n|}{\sqrt{\|\nabla|u_n|\|_2^2 + V_0\|u_n\|_2^2}} \right)^2} - 1) dx \leq C. \end{aligned}$$

Thus, by (4.6), (4.7), (4.8), and (4.9), we get

$$\frac{1}{2} \|u_n\|_{\varepsilon_n}^2 \leq C \|u_n\|_{\varepsilon_n}^q$$

reach a contradiction.

The second part is an immediate consequence of the first one observing that, if $u \in \mathcal{N}_\varepsilon$, by (f₃),

$$J_\varepsilon(u) = J_\varepsilon(u) - \frac{1}{\theta} J'_\varepsilon(u)[u] = \frac{\theta - 2}{2\theta} \|u\|_\varepsilon^2 + \frac{1}{\theta} \int_{\mathbb{R}^2} \left[f(|u|^2)|u|^2 - \frac{\theta}{2} F(|u|^2) \right] dx \geq \frac{\theta - 2}{2\theta} \|u\|_\varepsilon^2.$$

□

The functional J_ε satisfies the following fundamental property.

Lemma 4.3. *If $\{u_n\} \subset H_\varepsilon$ is a (PS) sequence for J_ε , then it is bounded in H_ε .*

Proof. Let $d := \lim_n J_\varepsilon(u_n) \in \mathbb{R}$. Using (f₃) we have that

$$\begin{aligned} \frac{\theta-2}{2\theta} \|u_n\|_\varepsilon^2 &\leq \left(\frac{1}{2} - \frac{1}{\theta}\right) \|u_n\|_\varepsilon^2 + \int_{\mathbb{R}^2} \left[\frac{1}{\theta} f(|u_n|^2) |u_n|^2 - \frac{1}{2} F(|u_n|^2) \right] dx \\ &= J_\varepsilon(u_n) - \frac{1}{\theta} J'_\varepsilon(u_n)[u_n] \\ &\leq d + o_n(1) + o_n(1) \|u_n\|_\varepsilon \end{aligned}$$

and we can conclude. \square

The following lemma is very important for the proof of Theorem 1.1.

Lemma 4.4. *There exists $\bar{\varepsilon} > 0$ such that, for all $\varepsilon \in (0, \bar{\varepsilon})$, $c_\varepsilon \leq c_{V_0}$.*

Proof. Assume for simplicity that $V(0) = V_0$ and let $\omega \in H^1(\mathbb{R}^2, \mathbb{R})$ be a positive ground state solution of the limit problem (3.1) with $\mu = V_0$ (see Lemma 3.1) and $\eta \in C_c^\infty(\mathbb{R}^2, [0, 1])$ be a cut-off function such that $\eta = 1$ in $B_1(0)$ and $\eta = 0$ in $B_2^c(0)$. Moreover, let

$$\omega_r(x) := \eta_r(x) \omega(x) e^{iA(0) \cdot x}, \quad \eta_r(x) = \eta(x/r) \text{ for } r > 0.$$

Observe that $|\omega_r| = \eta_r \omega$ and $\omega_r \in H_\varepsilon$ by Lemma 2.2.

Let $\varepsilon_n \rightarrow 0^+$ as $n \rightarrow +\infty$. Taking $t_{\varepsilon_n, r} > 0$ such that $t_{\varepsilon_n, r} \omega_r \in \mathcal{N}_{\varepsilon_n}$, by (4.5) we have

$$(4.10) \quad c_{\varepsilon_n} \leq J_{\varepsilon_n}(t_{\varepsilon_n, r} \omega_r) = \frac{t_{\varepsilon_n, r}^2}{2} \int_{\mathbb{R}^2} |\nabla_{A_{\varepsilon_n}} \omega_r|^2 dx + \frac{t_{\varepsilon_n, r}^2}{2} \int_{\mathbb{R}^2} V_{\varepsilon_n}(x) |\omega_r|^2 dx - \frac{1}{2} \int_{\mathbb{R}^2} F(t_{\varepsilon_n, r}^2 |\omega_r|^2) dx.$$

Observe that, for any fixed $r > 0$,

$$(4.11) \quad \lim_n \int_{\mathbb{R}^2} |\nabla_{A_{\varepsilon_n}} \omega_r|^2 dx = \int_{\mathbb{R}^2} |\nabla(\eta_r \omega)|^2 dx.$$

Indeed,

$$\begin{aligned} \int_{\mathbb{R}^2} |\nabla_{A_{\varepsilon_n}} \omega_r|^2 dx &= \int_{\mathbb{R}^2} \left| \frac{\nabla}{i}(\eta_r \omega) - [A_{\varepsilon_n}(x) - A(0)] \eta_r \omega \right|^2 dx \\ &= \int_{\mathbb{R}^2} |\nabla(\eta_r \omega)|^2 dx + \int_{\mathbb{R}^2} |A_{\varepsilon_n}(x) - A(0)|^2 \eta_r^2 \omega^2 dx \end{aligned}$$

and

$$\int_{\mathbb{R}^2} |A_{\varepsilon_n}(x) - A(0)|^2 \eta_r^2 \omega^2 dx \leq \int_{B_{2r}(0)} |A_{\varepsilon_n}(x) - A(0)|^2 \omega^2 dx = o_n(1).$$

Moreover

$$(4.12) \quad \lim_n \int_{\mathbb{R}^2} V_{\varepsilon_n}(x) |\omega_r|^2 dx = V_0 \int_{\mathbb{R}^2} \eta_r^2 \omega^2 dx$$

since

$$\left| \int_{\mathbb{R}^2} V_{\varepsilon_n}(x) |\omega_r|^2 dx - V_0 \int_{\mathbb{R}^2} \eta_r^2 \omega^2 dx \right| \leq \int_{\mathbb{R}^2} |V_{\varepsilon_n}(x) - V_0| \eta_r^2 \omega^2 dx \leq \int_{B_{2r}(0)} |V_{\varepsilon_n}(x) - V_0| \omega^2 dx = o_n(1).$$

We claim that, if $r > 0$ is fixed, $\{t_{\varepsilon_n, r}\}$ is bounded.

Indeed, let us assume by contradiction that there exists a subsequence, still denoted by $t_{\varepsilon_n, r}$, such that $t_{\varepsilon_n, r} \rightarrow +\infty$ as $n \rightarrow +\infty$.

Since $J'_{\varepsilon_n}(t_{\varepsilon_n, r} \omega_r)[t_{\varepsilon_n, r} \omega_r] = 0$, by (f₁) and (f₄) we have

$$\int_{\mathbb{R}^2} |\nabla_{A_{\varepsilon_n}} \omega_r|^2 dx + \int_{\mathbb{R}^2} V_{\varepsilon_n}(x) |\omega_r|^2 dx = \int_{\mathbb{R}^2} f(t_{\varepsilon_n, r}^2 |\omega_r|^2) |\omega_r|^2 dx \geq \int_{B_r(0)} f(t_{\varepsilon_n, r}^2 |\omega|^2) |\omega|^2 dx \geq C t_{\varepsilon_n, r}^{p-2} \rightarrow +\infty$$

as $n \rightarrow +\infty$, which is a contradiction with (4.11) and (4.12).

Thus, up to a subsequence, there exists $t_r \geq 0$ such that $t_{\varepsilon_n, r} \rightarrow t_r$ as $n \rightarrow +\infty$ for r fixed.

Therefore, by (4.10),

$$(4.13) \quad \limsup_n c_{\varepsilon_n} \leq \frac{t_r^2}{2} \int_{\mathbb{R}^2} |\nabla(\eta_r \omega)|^2 dx + \frac{t_r^2}{2} V_0 \int_{\mathbb{R}^2} |\eta_r \omega|^2 dx - \frac{1}{2} \int_{\mathbb{R}^2} F(t_r^2 |\eta_r \omega|^2) dx = I_{V_0}(t_r \eta_r \omega).$$

Let us consider now $r_m \rightarrow +\infty$ as $m \rightarrow +\infty$.

Since

$$\begin{aligned} \int_{\mathbb{R}^2} |\nabla(\eta_{r_m} \omega - \omega)|^2 dx &= \int_{\mathbb{R}^2} |\nabla \eta_{r_m}|^2 \omega^2 dx + 2 \int_{\mathbb{R}^2} (\eta_{r_m} - 1) \omega \nabla \eta_{r_m} \nabla \omega dx + \int_{\mathbb{R}^2} (\eta_{r_m} - 1)^2 |\nabla \omega|^2 dx, \\ \int_{\mathbb{R}^2} |\nabla \eta_{r_m}|^2 \omega^2 dx &\leq \frac{C}{r_m^2} \int_{B_{r_m}^c} \omega^2 dx = o_m(1), \\ \left| \int_{\mathbb{R}^2} (\eta_{r_m} - 1) \omega \nabla \eta_{r_m} \nabla \omega dx \right| &\leq \frac{C}{r_m} \left(\int_{B_{r_m}^c} \omega^2 dx \right)^{1/2} \left(\int_{B_{r_m}^c} |\nabla \omega|^2 dx \right)^{1/2} = o_m(1), \\ \int_{\mathbb{R}^2} (\eta_{r_m} - 1)^2 |\nabla \omega|^2 dx &\leq \int_{B_{r_m}^c} |\nabla \omega|^2 dx = o_m(1), \\ \int_{\mathbb{R}^2} (\eta_{r_m} - 1)^2 \omega^2 dx &\leq \int_{B_{r_m}^c} \omega^2 dx = o_m(1), \end{aligned}$$

then $\eta_{r_m} \omega \rightarrow \omega$ in $H^1(\mathbb{R}^2, \mathbb{R})$ as $m \rightarrow +\infty$.

Now, let $\tau_{r_m} > 0$ such that $\tau_{r_m} \eta_{r_m} \omega \in \mathcal{N}_{V_0}$.

Arguing as before, since

$$(4.14) \quad \int_{\mathbb{R}^2} |\nabla(\eta_{r_m} \omega)|^2 dx + V_0 \int_{\mathbb{R}^2} \eta_{r_m}^2 \omega^2 dx = \int_{\mathbb{R}^2} f(\tau_{r_m}^2 |\eta_{r_m} \omega|^2) \eta_{r_m}^2 \omega^2 dx,$$

we have that $\{\tau_{r_m}\}$ is bounded.

Thus, up to a subsequence, $\tau_{r_m} \rightarrow \bar{\tau} \geq 0$ as $m \rightarrow +\infty$ and, by (4.14), the uniqueness of the projection in \mathcal{N}_{V_0} , and since $\omega \in \mathcal{N}_{V_0}$ we get that $\bar{\tau} = 1$.

Hence, by (4.13) and (3.2),

$$\limsup_n c_{\varepsilon_n} \leq I_{V_0}(t_{r_m} \eta_{r_m} \omega) \leq I_{V_0}(\tau_{r_m} \eta_{r_m} \omega)$$

and so, passing to the limit as $m \rightarrow +\infty$, we get

$$\limsup_n c_{\varepsilon_n} \leq I_{V_0}(\omega) = c_{V_0}.$$

□

In our arguments we will use the following result that is a version of the celebrated Lions Lemma (see e.g. [33] and [20, Lemma 3.3]).

Lemma 4.5. *Let $\{u_n\} \subset H_\varepsilon$ satisfy $J'_\varepsilon(u_n)[u_n] \rightarrow 0$, $u_n \rightharpoonup 0$ in H_ε , as $n \rightarrow +\infty$, and $\ell^2 := \limsup_n \|\nabla |u_n|\|_2^2 < 1$. Then, one of the following alternatives occurs:*

- (i) $u_n \rightarrow 0$ in H_ε as $n \rightarrow +\infty$;
- (ii) there are a sequence $\{y_n\} \subset \mathbb{R}^2$ and constants $R, \beta > 0$ such that

$$\liminf_n \int_{B_R(y_n)} |u_n|^2 dx \geq \beta.$$

Proof. If (ii) does not hold, then, for every $R > 0$,

$$\lim_n \sup_{y \in \mathbb{R}^2} \int_{B_R(y)} |u_n|^2 dx = 0.$$

Since $\{|u_n|\}$ is bounded in $H^1(\mathbb{R}^2)$, [28, Chapter 6, Lemma 8.4] implies that

$$(4.15) \quad \|u_n\|_\tau \rightarrow 0 \text{ as } n \rightarrow +\infty \text{ for all } \tau > 2.$$

Moreover, using (2.6), we have that, fixed $q > 2$, for any $\zeta > 0$ and $\alpha > 4\pi$, there exists $C > 0$ such that

$$(4.16) \quad \begin{aligned} 0 \leq \|u_n\|_\varepsilon^2 &= \int_{\mathbb{R}^2} f(|u_n|^2) |u_n|^2 dx + o_n(1) \\ &\leq \zeta \|u_n\|_2^2 + C \int_{\mathbb{R}^2} |u_n|^q (e^{\alpha|u_n|^2} - 1) dx + o_n(1) \\ &\leq \frac{\zeta}{V_0} \|u_n\|_\varepsilon^2 + C \int_{\mathbb{R}^2} |u_n|^q (e^{\alpha|u_n|^2} - 1) dx + o_n(1). \end{aligned}$$

Let us fix $\kappa \in (0, 1 - \ell^2)$ and take

$$r \in \left(1, \frac{1}{\ell^2 + \kappa}\right), \quad \alpha \in \left(4\pi, \frac{4\pi}{r(\ell^2 + \kappa)}\right).$$

Thus, the Hölder inequality, (2.17), and Lemma 2.3, imply that, for n large enough (such that $\|\nabla|u_n|\|_2^2 < \ell^2 + \kappa$),

$$\int_{\mathbb{R}^2} |u_n|^q (e^{\alpha|u_n|^2} - 1) dx \leq \|u_n\|_{qr'}^q \left(\int_{\mathbb{R}^2} (e^{r\alpha(\ell^2 + \kappa)} \left(\frac{|u_n|}{\sqrt{\ell^2 + \kappa}}\right)^2 - 1) dx \right)^{1/r} \leq C \|u_n\|_{qr'}^q$$

and so, using (4.15) and (4.16) we can conclude. \square

Now we prove a fundamental result on the $(PS)_d$ sequences for J_ε in the case $V_\infty < \infty$.

Lemma 4.6. *Let $d \in \mathbb{R}$, $V_\infty < +\infty$ and $\{u_n\} \subset H_\varepsilon$ be a $(PS)_d$ sequence for J_ε such that $\ell^2 := \limsup_n \|\nabla|u_n|\|_2^2 < 1$ and $u_n \rightharpoonup 0$ in H_ε . If $u_n \not\rightarrow 0$ in H_ε , then $d \geq c_{V_\infty}$.*

Proof. For every $n \in \mathbb{N}$, let $t_n := t(|u_n|) \in (0, \infty)$ be such that $t_n|u_n| \in \mathcal{N}_{V_\infty}$ (see the first part of Section 3).

Firstly we show that $\limsup_n t_n \leq 1$.

Assume by contradiction that there exist $\delta > 0$ and a subsequence, still denoted by $\{t_n\}$, such that, for every $n \in \mathbb{N}$,

$$(4.17) \quad t_n \geq 1 + \delta.$$

Since $\{u_n\}$ is a $(PS)_d$ sequence for J_ε , we have

$$(4.18) \quad \int_{\mathbb{R}^2} (|\nabla_{A_\varepsilon} u_n|^2 + V_\varepsilon(x) |u_n|^2) dx = \int_{\mathbb{R}^2} f(|u_n|^2) |u_n|^2 dx + o_n(1).$$

On the other hand, $t_n|u_n| \in \mathcal{N}_{V_\infty}$, then one has

$$(4.19) \quad \int_{\mathbb{R}^2} (|\nabla|u_n||^2 + V_\infty |u_n|^2) dx = \int_{\mathbb{R}^2} f(t_n^2 |u_n|^2) |u_n|^2 dx.$$

Putting together (4.18), (4.19) and using the diamagnetic inequality (2.1) we obtain

$$\int_{\mathbb{R}^2} (f(t_n^2 |u_n|^2) - f(|u_n|^2)) |u_n|^2 dx \leq \int_{\mathbb{R}^2} (V_\infty - V_\varepsilon(x)) |u_n|^2 dx + o_n(1).$$

Now, by the assumption (V), we can see that for every $\eta > 0$ there exists $R = R(\eta) > 0$ such that, for every $|x| \geq R/\varepsilon$,

$$V_\infty - V_\varepsilon(x) \leq \eta.$$

Then, using the boundedness of $\{u_n\}$ in H_ε and that $u_n \rightarrow 0$ in $L^2(B_{R/\varepsilon}(0), \mathbb{C})$ by Lemma 2.1,

$$\int_{\mathbb{R}^2} (V_\infty - V(\varepsilon x)) |u_n|^2 dx \leq V_\infty \int_{B_{R/\varepsilon}(0)} |u_n|^2 dx + \eta \int_{B_R^c(0)} |u_n|^2 dx \leq o_n(1) + C\eta.$$

Thus,

$$(4.20) \quad \int_{\mathbb{R}^2} \left(f(t_n^2 |u_n|^2) - f(|u_n|^2) \right) |u_n|^2 dx \leq o_n(1) + C\eta.$$

Moreover, since $u_n \not\rightarrow 0$ in H_ε , by Lemma 4.5, there exist a sequence $\{y_n\} \subset \mathbb{R}^2$, and two positive numbers R, β such that

$$(4.21) \quad \liminf_{n \rightarrow \infty} \int_{B_R(y_n)} |u_n|^2 dx \geq \beta.$$

Now, let us consider $v_n := |u_n|(\cdot + y_n)$. Taking into account (V), the diamagnetic inequality (2.1), and the boundedness of $\{u_n\}$ in H_ε , we can see that

$$\|\nabla v_n\|_2^2 + V_0 \|v_n\|_2^2 = \|\nabla |u_n|\|_2^2 + V_0 \|u_n\|_2^2 \leq \|u_n\|_\varepsilon^2 \leq C.$$

Therefore $v_n \rightharpoonup v$ in $H^1(\mathbb{R}^2, \mathbb{R})$, $v_n \rightarrow v$ in $L_{\text{loc}}^r(\mathbb{R}^2, \mathbb{R})$ for all $r \geq 1$, and, by (4.21),

$$\int_{B_R(0)} |v|^2 dx = \lim_{n \rightarrow \infty} \int_{B_R(0)} |v_n|^2 dx \geq \beta.$$

Thus there exists $\Omega \subset B_R(0)$ with positive measure such that $v \not\equiv 0$ in Ω .

Moreover, by (4.17), (f₄), and (4.20), we obtain

$$\int_{\mathbb{R}^2} \left(f((1+\delta)^2 v_n^2) - f(v_n^2) \right) v_n^2 dx \leq o_n(1) + C\eta,$$

and so, (f₄) and the Fatou's Lemma imply

$$0 < \int_{\Omega} \left(f((1+\delta)^2 v^2) - f(v^2) \right) v^2 dx \leq C\eta.$$

Thus, by the arbitrariness of $\eta > 0$, we get a contradiction.

Now, two cases can occur.

Case 1: $\limsup_n t_n = 1$.

In this case there exists a subsequence still denoted by $\{t_n\}$ such that $t_n \rightarrow 1$.

Taking into account that $\{u_n\}$ is a $(PS)_d$ sequence for J_ε , (3.2), (3.3), and the diamagnetic inequality (2.1), we have

$$(4.22) \quad \begin{aligned} d + o_n(1) &= J_\varepsilon(u_n) \geq J_\varepsilon(u_n) - I_{V_\infty}(t_n |u_n|) + c_{V_\infty} \\ &\geq \frac{1-t_n^2}{2} \int_{\mathbb{R}^2} |\nabla |u_n||^2 dx + \frac{1}{2} \int_{\mathbb{R}^2} (V_\varepsilon(x) - t_n^2 V_\infty) |u_n|^2 dx \\ &\quad + \frac{1}{2} \int_{\mathbb{R}^2} (F(t_n^2 |u_n|^2) - F(|u_n|^2)) dx + c_{V_\infty}. \end{aligned}$$

Since $\{|u_n|\}$ is bounded in $H^1(\mathbb{R}^2, \mathbb{R})$ and $t_n \rightarrow 1$, it is easy to see that

$$(4.23) \quad \frac{1-t_n^2}{2} \int_{\mathbb{R}^2} |\nabla |u_n||^2 dx = o_n(1).$$

Moreover, by (V), we have that for every $\eta > 0$ there exists $R = R(\zeta) > 0$ such that for any $x \in \mathbb{R}^2$ with $|x| > R/\varepsilon$, it holds

$$V_\varepsilon(x) - t_n^2 V_\infty = (V_\varepsilon(x) - V_\infty) + (1 - t_n^2) V_\infty \geq -\eta + (1 - t_n^2) V_\infty.$$

Thus, using as before the boundedness of $\{u_n\}$ in H_ε , that $|u_n| \rightarrow 0$ in $L^r(B_{R/\varepsilon})$ for $r \geq 1$, and $t_n \rightarrow 1$, for n large enough we get

$$(4.24) \quad \begin{aligned} \int_{\mathbb{R}^2} (V_\varepsilon(x) - t_n^2 V_\infty) |u_n|^2 dx &\geq (V_0 - t_n^2 V_\infty) \int_{B_{R/\varepsilon}(0)} |u_n|^2 dx \\ &\quad + ((1 - t_n^2) V_\infty - \eta) \int_{B_{R/\varepsilon}^c(0)} |u_n|^2 dx \\ &\geq o_n(1) - C\eta. \end{aligned}$$

Finally let $\vartheta, l > 1$ such that $l\vartheta\ell^2 < 1$. Since $t_n \rightarrow 1$, for n large enough, applying the Mean Value Theorem and using also (f4), we get

$$|F(t_n^2 |u_n|^2) - F(|u_n|^2)| \leq |t_n^2 - 1| f(\theta_n(x) |u_n|^2) |u_n|^2 \leq |t_n^2 - 1| f(\vartheta |u_n|^2) |u_n|^2.$$

Then

$$(4.25) \quad \left| \int_{\mathbb{R}^2} (F(t_n^2 |u_n|^2) - F(|u_n|^2)) dx \right| \leq |t_n^2 - 1| \int_{\mathbb{R}^2} f(\vartheta |u_n|^2) |u_n|^2 dx = o_n(1).$$

Indeed, $t_n \rightarrow 1$. Moreover, if we take

$$\kappa \in \left(0, \frac{1}{l\vartheta} - \ell^2\right) \text{ and } \alpha \in \left(4\pi, \frac{4\pi}{l\vartheta(\ell^2 + \kappa)}\right),$$

by (2.4), fixed $q > 2$, for any $\zeta > 0$, there exists a constant $C > 0$ such that

$$\int_{\mathbb{R}^2} f(\vartheta |u_n|^2) |u_n|^2 dx \leq \zeta \|u_n\|_2^2 + C \int_{\mathbb{R}^2} |u_n|^q (e^{\vartheta\alpha |u_n|^2} - 1) dx$$

and the right hand side is bounded due to the boundedness of $\{u_n\}$ and since the Hölder inequality, (2.17), Lemma 2.3, and

$$\|\nabla |u_n|\|_2^2 < \ell^2 + \kappa < \frac{1}{l\vartheta} < 1 \text{ for } n \text{ large enough,}$$

imply that

$$\begin{aligned} \int_{\mathbb{R}^2} |u_n|^q (e^{\vartheta\alpha |u_n|^2} - 1) dx &\leq \|u_n\|_{q^{l'}}^q \left(\int_{\mathbb{R}^2} (e^{l\vartheta\alpha |u_n|^2} - 1) dx \right)^{1/l} \\ &\leq \|u_n\|_{q^{l'}}^q \left(\int_{\mathbb{R}^2} (e^{l\vartheta\alpha(\ell^2 + \kappa) \left(\frac{|u_n|}{\sqrt{\ell^2 + \kappa}}\right)^2} - 1) dx \right)^{1/l} \leq C, \end{aligned}$$

where $l' > 1$ is the conjugate exponent of l .

Hence, putting together (4.22), (4.23), (4.24), and (4.25), we can obtain

$$d + C\eta + o_n(1) \geq c_{V_\infty},$$

and, due to the arbitrariness of η , taking the limit as $n \rightarrow \infty$, we conclude.

Case 2: $\limsup_n t_n = \bar{t} < 1$.

In this case there exists a subsequence still denoted by $\{t_n\}$, such that $t_n \rightarrow \bar{t}$ and $t_n < 1$ for n large enough.

Since $\{u_n\}$ is a bounded $(PS)_d$ sequence for J_ε in H_ε , we have

$$(4.26) \quad d + o_n(1) = J_\varepsilon(u_n) - \frac{1}{2} J'_\varepsilon(u_n)[u_n] = \frac{1}{2} \int_{\mathbb{R}^2} (f(|u_n|^2) |u_n|^2 - F(|u_n|^2)) dx.$$

Note that, by (f₃), the map $t \mapsto f(t)t - F(t)$ is increasing for $t > 0$. Hence, this fact combined with $t_n|u_n| \in \mathcal{N}_{V_\infty}$, $t_n < 1$, and (4.26), allows us to get

$$\begin{aligned} c_{V_\infty} &\leq I_{V_\infty}(t_n|u_n|) = I_{V_\infty}(t_n|u_n|) - \frac{t_n}{2} I'_{V_\infty}(t_n|u_n|)[|u_n|] \\ &= \frac{1}{2} \int_{\mathbb{R}^2} \left(f(t_n^2|u_n|^2) t_n^2|u_n|^2 - F(t_n^2|u_n|^2) \right) dx \\ &\leq \frac{1}{2} \int_{\mathbb{R}^2} \left(f(|u_n|^2) |u_n|^2 - F(|u_n|^2) \right) dx \\ &= d + o_n(1). \end{aligned}$$

Passing to the limit as $n \rightarrow +\infty$, we complete the proof. \square

We shall also use the next result about the (PS) sequences.

Lemma 4.7. *Let $\{u_n\} \subset H_\varepsilon$ be a (PS) sequence for J_ε such that $\ell^2 := \limsup_n \|u_n\|_\varepsilon^2 < 1$. Then, up to a subsequence, $\nabla|u_n| \rightarrow \nabla|u|$ a.e. in \mathbb{R}^2 .*

Proof. Let $\{u_n\} \subset H_\varepsilon$ be a (PS) sequence for J_ε with $\limsup_n \|u_n\|_\varepsilon^2 < 1$. By Lemma 4.3 we have that $\{u_n\}$ is bounded in H_ε and so, up to a subsequence, $u_n \rightharpoonup u$ in H_ε , $u_n \rightarrow u$ in $L^r_{\text{loc}}(\mathbb{R}^2, \mathbb{C})$ for any $r \geq 1$ and $u_n \rightarrow u$ a.e. in \mathbb{R}^2 . Since the norm is weakly lower semicontinuous, it follows that $\|u\|_\varepsilon^2 < 1$.

Now observe that, for every $\phi \in C_c^\infty(\mathbb{R}^2, \mathbb{C})$,

$$f(|u_n|^2)u_n\bar{\phi} \rightarrow f(|u|^2)u\bar{\phi} \text{ a.e. in } \mathbb{R}^2 \text{ as } n \rightarrow +\infty$$

and, by (2.4), fixed $q > 2$, for any $\zeta > 0$ and $\alpha > 4\pi$, there exists $C > 0$ such that, for every $\phi \in C_c^\infty(\mathbb{R}^2, \mathbb{R})$,

$$|\operatorname{Re}[f(|u_n|^2)u_n\bar{\phi}]| \leq |f(|u_n|^2)u_n\bar{\phi}| \leq \zeta|u_n||\phi| + C|u_n|^{q-1}(e^{\alpha|u_n|^2} - 1)|\phi|$$

with

$$\zeta|u_n||\phi| + C|u_n|^{q-1}(e^{\alpha|u_n|^2} - 1)|\phi| \rightarrow \zeta|u||\phi| + C|u|^{q-1}(e^{\alpha|u|^2} - 1)|\phi| \text{ a.e. in } \mathbb{R}^2 \text{ as } n \rightarrow +\infty.$$

Moreover, let $r^*, \tau > 1$ such that $r^*\tau\ell^2 < 1$. Using the Hölder inequality, (2.17), Sobolev inequality, Lemma 2.3, and taking

$$\kappa \in \left(0, \frac{1}{r^*\tau} - \ell^2\right) \text{ and } \alpha \in \left(4\pi, \frac{4\pi}{r^*\tau(\ell^2 + \kappa)}\right),$$

we have that, for n large enough, such that $\|\nabla|u_n|\|_2^2 \leq \ell^2 + \kappa$,

$$\begin{aligned} (4.27) \quad \int_{\mathbb{R}^2} \left[|u_n|^{q-1}(e^{\alpha|u_n|^2} - 1) \right]^{r^*} dx &\leq \|u_n\|_{\tau'r^*(q-1)}^{r^*(q-1)} \left(\int_{\mathbb{R}^2} (e^{\alpha r^*\tau|u_n|^2} - 1) dx \right)^{1/\tau} \\ &\leq C \|u_n\|_\varepsilon^{r^*(q-1)} \left(\int_{\mathbb{R}^2} (e^{\alpha r^*\tau(\ell^2 + \kappa)} \left(\frac{|u_n|}{\sqrt{\ell^2 + \kappa}} \right)^2 - 1) dx \right)^{1/\tau} \leq C, \end{aligned}$$

where τ' is the conjugate exponent of τ .

Then, up to a subsequence and applying the Severini-Egoroff Theorem, we get

$$|u_n|^{q-1}(e^{\alpha|u_n|^2} - 1) \rightharpoonup |u|^{q-1}(e^{\alpha|u|^2} - 1) \text{ in } L^r(\mathbb{R}^2, \mathbb{R}).$$

Thus, since, up to a subsequence, also $|u_n| \rightharpoonup |u|$ in $L^2(\mathbb{R}^2, \mathbb{R})$, we have, as $n \rightarrow +\infty$,

$$\zeta \int_{\mathbb{R}^2} |u_n||\phi| dx + C \int_{\mathbb{R}^2} |u_n|^{q-1}(e^{\alpha|u_n|^2} - 1)|\phi| dx \rightarrow \zeta \int_{\mathbb{R}^2} |u||\phi| dx + C \int_{\mathbb{R}^2} |u|^{q-1}(e^{\alpha|u|^2} - 1)|\phi| dx.$$

Hence, by a variant of the Lebesgue Dominated Convergence Theorem, we obtain that, for every $\phi \in C_c^\infty(\mathbb{R}^2, \mathbb{C})$

$$\operatorname{Re} \int_{\mathbb{R}^2} f(|u_n|^2)u_n\bar{\phi} dx \rightarrow \operatorname{Re} \int_{\mathbb{R}^2} f(|u|^2)u\bar{\phi} dx,$$

from which we get $J'_\varepsilon(u) = 0$.

Let now $R > 0$ be arbitrary and ψ be a $C_c^\infty(\mathbb{R}^2, [0, 1])$ function such that $\psi \equiv 1$ in $B_R(0)$ and $\psi \equiv 0$ in $B_{2R}^c(0)$.

First we observe that, since for every $v \in H_\varepsilon$,

$$\nabla_{A_\varepsilon}(v\psi) = \psi \nabla_{A_\varepsilon} v - iv \nabla \psi,$$

and then, by the Hölder inequality,

$$\|\nabla_{A_\varepsilon}(v\psi)\|_2^2 = \|\psi \nabla_{A_\varepsilon} v\|_2^2 + 2\operatorname{Re} \int_{\mathbb{R}^2} i\bar{v}\psi \nabla_{A_\varepsilon} v \nabla \psi dx + \|v\| \|\nabla \psi\|_2^2 \leq C(\|\nabla_{A_\varepsilon} v\|_2^2 + \|v\|_2^2),$$

we have that $v\psi \in H_\varepsilon$ and that the boundedness of $\{u_n\}$ in H_ε implies the boundedness of $\{u_n\psi\}$ in H_ε . By (2.4), the Hölder and diamagnetic inequalities, (2.17), Lemma 2.1, and Lemma 2.3, for $q > 2$ and $\alpha > 4\pi$,

$$\begin{aligned} \left| \operatorname{Re} \int_{\mathbb{R}^2} f(|u|^2) u \overline{(u - u_n)} \psi dx \right| &\leq \int_{\mathbb{R}^2} f(|u|^2) |u| |u - u_n| \psi dx \\ &\leq \zeta \int_{\mathbb{R}^2} |u| |u - u_n| \psi dx \\ &\quad + C \int_{\mathbb{R}^2} |u|^{q-1} (e^{\alpha|u|^2} - 1) |u - u_n| \psi dx \\ &\leq \zeta \|u\|_2 \left(\int_{B_{2R}(0)} |u - u_n|^2 dx \right)^{\frac{1}{2}} \\ &\quad + C \left(\int_{\mathbb{R}^2} |u|^{2(q-1)} (e^{2\alpha|u|^2} - 1) dx \right)^{\frac{1}{2}} \left(\int_{B_{2R}(0)} |u - u_n|^2 dx \right)^{\frac{1}{2}} \\ &\leq o_n(1) + C \|u\|_{4(q-1)}^{q-1} \left(\int_{\mathbb{R}^2} (e^{4\alpha|u|^2} - 1) dx \right)^{\frac{1}{4}} o_n(1) \\ &= o_n(1). \end{aligned}$$

Moreover, by Lemma 2.1 and the Hölder inequality,

$$\left| \operatorname{Re} \int_{\mathbb{R}^2} V_\varepsilon(x) u \overline{(u - u_n)} \psi dx \right| \leq C \|u\|_2 \left(\int_{B_{2R}(0)} |u - u_n|^2 dx \right)^{\frac{1}{2}} = o_n(1)$$

and

$$\left| \operatorname{Re} \int_{\mathbb{R}^2} i \overline{(u - u_n)} \nabla_{A_\varepsilon} u \nabla \psi dx \right| \leq C \|\nabla_{A_\varepsilon} u\|_2 \left(\int_{B_{2R}(0) \setminus B_R(0)} |u - u_n|^2 dx \right)^{\frac{1}{2}} = o_n(1).$$

Thus

$$\begin{aligned} \operatorname{Re} \int_{\mathbb{R}^2} \nabla_{A_\varepsilon} u \overline{\nabla_{A_\varepsilon} (u - u_n)} \psi dx &= J'_\varepsilon(u) [(u - u_n)\psi] - \operatorname{Re} \int_{\mathbb{R}^2} V_\varepsilon(x) u \overline{(u - u_n)} \psi dx \\ &\quad - \operatorname{Re} \int_{\mathbb{R}^2} i \overline{(u - u_n)} \nabla_{A_\varepsilon} u \nabla \psi dx + \operatorname{Re} \int_{\mathbb{R}^2} f(|u|^2) u \overline{(u - u_n)} \psi dx \\ &= o_n(1). \end{aligned} \tag{4.28}$$

Now, using (4.28), $J'_\varepsilon(u_n) \rightarrow 0$ as $n \rightarrow +\infty$, and the boundedness of the sequence $\{u_n\psi\}$ in H_ε , we obtain

$$\begin{aligned}
 0 &\leq \int_{B_R(0)} |\nabla_{A_\varepsilon} u_n - \nabla_{A_\varepsilon} u|^2 dx \leq \int_{\mathbb{R}^2} |\nabla_{A_\varepsilon} u_n - \nabla_{A_\varepsilon} u|^2 \psi dx \\
 (4.29) \quad &= \operatorname{Re} \int_{\mathbb{R}^2} \nabla_{A_\varepsilon} u_n \overline{\nabla_{A_\varepsilon} (u_n - u)} \psi dx + o_n(1) \\
 &= J'_\varepsilon(u_n)[(u_n - u)\psi] + K_{1,n} + K_{2,n} + K_{3,n} + o_n(1) \\
 &= K_{1,n} + K_{2,n} + K_{3,n} + o_n(1),
 \end{aligned}$$

where

$$\begin{aligned}
 K_{1,n} &= \operatorname{Re} \int_{\mathbb{R}^2} i \overline{(u - u_n)} \nabla_{A_\varepsilon} u_n \nabla \psi dx, \\
 K_{2,n} &= \operatorname{Re} \int_{\mathbb{R}^2} V_\varepsilon(x) u_n \overline{(u - u_n)} \psi dx, \\
 K_{3,n} &= \operatorname{Re} \int_{\mathbb{R}^2} f(|u_n|^2) u_n \overline{(u_n - u)} \psi dx.
 \end{aligned}$$

By Lemma 2.1, the Hölder inequality, and the boundedness of $\{u_n\}$ in H_ε , we get

$$|K_{1,n}| \leq \int_{B_{2R}(0) \setminus B_R(0)} |\nabla_{A_\varepsilon} u_n| |u - u_n| |\nabla \psi| dx \leq C \|u_n\|_\varepsilon \left(\int_{B_{2R}(0) \setminus B_R(0)} |u - u_n|^2 dx \right)^{1/2} = o_n(1)$$

and

$$|K_{2,n}| \leq \int_{B_{2R}(0)} V_\varepsilon(x) |u_n| |u - u_n| dx \leq C \|u_n\|_2 \left(\int_{B_{2R}(0)} |u - u_n|^2 dx \right)^{1/2} = o_n(1).$$

Moreover, by (2.4), the Hölder inequality and the boundedness of $\{u_n\}$ in H_ε , for $q > 2$ and suitable $\tau > 1$ and $\alpha > 4\pi$, if τ' is the conjugate exponent of τ ,

$$\begin{aligned}
 |K_{3,n}| &\leq \int_{B_{2R}(0)} |f(|u_n|^2)| |u_n| |u_n - u| dx \\
 &\leq \zeta \|u_n\|_2 \left(\int_{B_{2R}(0)} |u_n - u|^2 dx \right)^{1/2} + C \int_{B_{2R}(0)} |u_n - u| |u_n|^{q-1} (e^{\alpha|u_n|^2} - 1) dx \\
 &\leq o_n(1) + C \left(\int_{B_{2R}(0)} |u_n - u|^{\tau'} dx \right)^{1/\tau'} \left(\int_{\mathbb{R}^2} [|u_n|^{q-1} (e^{\alpha|u_n|^2} - 1)]^\tau dx \right)^{1/\tau} \\
 &= o_n(1),
 \end{aligned}$$

since, arguing as in (4.27), for n large enough,

$$\int_{\mathbb{R}^2} [|u_n|^{q-1} (e^{\alpha|u_n|^2} - 1)]^\tau dx \leq C.$$

Hence, by (4.29), we obtain that, for any $R > 0$,

$$\int_{B_R(0)} |\nabla_{A_\varepsilon} u_n - \nabla_{A_\varepsilon} u|^2 dx \rightarrow 0, \text{ as } n \rightarrow +\infty,$$

which implies that $\nabla_{A_\varepsilon} u_n \rightarrow \nabla_{A_\varepsilon} u$ a.e. in \mathbb{R}^2 . Thus, using also that $u_n \rightarrow u$ a.e. in \mathbb{R}^2 , we have

$$\nabla u_n \rightarrow \nabla u \text{ a.e. in } \mathbb{R}^2$$

and we can conclude. \square

Now we provide a range of levels in which the functional J_ε satisfies the (PS) condition.

Proposition 4.8. *Let $c \in \mathbb{R}$ and $\{u_n\}$ be a $(PS)_c$ sequence for J_ε such that $\limsup_{n \rightarrow \infty} \|u_n\|_\varepsilon^2 < 1/8$. Assume that $c < c_{V_\infty}$ if $V_\infty < +\infty$ or c is arbitrary if $V_\infty = +\infty$. Then, $\{u_n\}$ admits a convergent subsequence in H_ε .*

Proof. Since the sequence $\{u_n\}$ is bounded in H_ε , up to a subsequence, $u_n \rightharpoonup u$ in H_ε and $u_n \rightarrow u$ in $L_{\text{loc}}^r(\mathbb{R}^2, \mathbb{C})$ for any $r \geq 1$. Moreover, arguing as in Lemma 4.7, we also have that, for any $\phi \in H_\varepsilon$,

$$\operatorname{Re} \int_{\mathbb{R}^2} f(|u_n|^2) u_n \bar{\phi} dx \rightarrow \operatorname{Re} \int_{\mathbb{R}^2} f(|u|^2) u \bar{\phi} dx \text{ as } n \rightarrow +\infty.$$

Thus u is a critical point of J_ε and, using (f3),

$$(4.30) \quad J_\varepsilon(u) = J_\varepsilon(u) - \frac{1}{\theta} J'_\varepsilon(u)[u] = \frac{\theta-2}{2\theta} \|u\|_\varepsilon^2 + \frac{1}{\theta} \int_{\mathbb{R}^2} \left(f(|u|^2) |u|^2 - \frac{\theta}{2} F(|u|^2) \right) dx \geq 0.$$

Now, let $\{u_{n_j}\}$ be a subsequence of $\{u_n\}$ satisfying the assumptions of Lemma 4.7 and Lemma 2.6 and let $v_j := u_{n_j} - \hat{u}_j$ where $\hat{u}_j = \varphi_j u$, with $\varphi_j(x) = \varphi(2x/j)$, $\varphi \in C_c^\infty(\mathbb{R}^2, \mathbb{R})$, $0 \leq \varphi \leq 1$, $\varphi(x) = 1$ if $|x| \leq 1$, and $\varphi(x) = 0$ if $|x| \geq 2$. We claim that

$$(4.31) \quad J_\varepsilon(v_j) = c - J_\varepsilon(u) + o_j(1),$$

and

$$(4.32) \quad J'_\varepsilon(v_j) = o_j(1).$$

In order to prove (4.31), let us observe that

$$J_\varepsilon(v_j) - J_\varepsilon(u_{n_j}) + J_\varepsilon(\hat{u}_j) = \|\hat{u}_j\|_\varepsilon^2 - \langle u_{n_j}, \hat{u}_j \rangle_\varepsilon + \frac{1}{2} \int_{\mathbb{R}^2} (F(|u_{n_j}|^2) - F(|v_j|^2) - F(|\hat{u}_j|^2)) dx.$$

In view of the weak convergence of $\{u_{n_j}\}$ to u in H_ε and using Lemma 2.4, we can see that

$$\|\hat{u}_j\|_\varepsilon^2 - \langle u_{n_j}, \hat{u}_j \rangle_\varepsilon = o_j(1).$$

Moreover, by Lemma 4.7 and (2.9) in Lemma 2.6, we have that

$$\int_{\mathbb{R}^2} (F(|u_{n_j}|^2) - F(|v_j|^2) - F(|\hat{u}_j|^2)) dx = o_j(1),$$

getting (4.31).

To prove (4.32), we observe that, for any $\phi \in H_\varepsilon$,

$$\begin{aligned} |(J'_\varepsilon(v_j) - J'_\varepsilon(u_{n_j}) + J'_\varepsilon(\hat{u}_j))[\phi]| &= \left| \operatorname{Re} \int_{\mathbb{R}^2} (f(|u_{n_j}|^2) u_{n_j} - f(|v_j|^2) v_j - f(|\hat{u}_j|^2) \hat{u}_j) \bar{\phi} dx \right| \\ &\leq \int_{\mathbb{R}^2} |f(|u_{n_j}|^2) u_{n_j} - f(|v_j|^2) v_j - f(|\hat{u}_j|^2) \hat{u}_j| |\phi| dx \end{aligned}$$

and so, by Lemma 4.7 and (2.10) in Lemma 2.6, applying the diamagnetic inequality (2.1), we have that

$$\sup_{\|\phi\|_\varepsilon \leq 1} |(J'_\varepsilon(v_j) - J'_\varepsilon(u_{n_j}) + J'_\varepsilon(\hat{u}_j))[\phi]| \rightarrow 0.$$

Thus, since $J'_\varepsilon(u_{n_j}) \rightarrow 0$ and $J'_\varepsilon(\hat{u}_j) \rightarrow J'_\varepsilon(u) = 0$, (4.32) holds.

Now, if $V_\infty < +\infty$ and $c < c_{V_\infty}$, by (4.30), we have that

$$(4.33) \quad c - J_\varepsilon(u) \leq c < c_{V_\infty}.$$

Moreover, by (4.31) and (4.32), $\{v_j\}$ is a $(PS)_{c-J_\varepsilon(u)}$ sequence for J_ε with $v_j \rightharpoonup 0$ in H_ε .

Note also that, due to the weakly lower semicontinuity of the norm,

$$\|u\|_\varepsilon^2 \leq \liminf_j \|u_{n_j}\|_\varepsilon^2.$$

Then, using also the diamagnetic inequality and Lemma 2.4,

$$(4.34) \quad \limsup_j (\|\nabla|v_j|\|_2^2 + V_0\|v_j\|_2^2) \leq \limsup_j \|u_{n_j} - \hat{u}_j\|_\varepsilon^2 \leq 2(\limsup_j \|u_{n_j}\|_\varepsilon^2 + \limsup_j \|\hat{u}_j\|_\varepsilon^2) < \frac{1}{2}.$$

Hence, (4.33) and Lemma 4.6 imply that $v_j \rightarrow 0$ in H_ε and so, by Lemma 2.4, we get that $u_{n_j} \rightarrow u$ in H_ε as $j \rightarrow +\infty$.

If $V_\infty = +\infty$, then, by Lemma 2.4 and Lemma 2.1, $v_j \rightarrow 0$ in $L^r(\mathbb{R}^2, \mathbb{C})$ for any $r \geq 2$. Arguing as in (4.34), we also have

$$\limsup_j (\|\nabla|v_j|\|_2^2 + V_0\|v_j\|_2^2) < 1/2 < 1$$

and so, by (2.6), the Hölder inequality, (2.17), and Lemma 2.3, we deduce that, fixed $q > 2$, for suitable $\tau > 1$ and $\alpha > 4\pi$ such that $\alpha\tau < 8\pi$, for j large enough,

$$(4.35) \quad \begin{aligned} \|v_j\|_\varepsilon^2 &= \int_{\mathbb{R}^2} f(|v_j|^2)|v_j|^2 dx + o_j(1) \leq C \int_{\mathbb{R}^2} |v_j|^q (e^{\alpha|v_j|^2} - 1) dx + o_j(1) \\ &\leq C \|v_j\|_{q\tau'}^q \left(\int_{\mathbb{R}^2} (e^{\alpha|v_j|^2} - 1)^\tau dx \right)^{1/\tau} + o_j(1) \\ &\leq C \|v_j\|_{q\tau'}^q \left(\int_{\mathbb{R}^2} \left(e^{\frac{\alpha\tau}{2}(\sqrt{2}|v_j|)^2} - 1 \right) dx \right)^{1/\tau} + o_j(1) = o_j(1), \end{aligned}$$

where τ' is the conjugate exponent of τ .

Hence $u_{n_j} \rightarrow u$ in H_ε as $j \rightarrow +\infty$ and the proof is completed. \square

To show that \mathcal{N}_ε is a natural constraint, namely that the constrained critical points of the functional J_ε on \mathcal{N}_ε are critical points of J_ε in H_ε , we prove the following preliminary result.

Proposition 4.9. *Let $c \in \mathbb{R}$ and $\{u_n\}$ be a $(PS)_c$ sequence for J_ε restricted to \mathcal{N}_ε with $\limsup_{n \rightarrow \infty} \|u_n\|_\varepsilon^2 < 1/8$. Assume that $c < c_{V_\infty}$ if $V_\infty < +\infty$ or c is arbitrary if $V_\infty = +\infty$. Then, $\{u_n\}$ has a convergent subsequence in \mathcal{N}_ε .*

Proof. Let $\{u_n\} \subset \mathcal{N}_\varepsilon$ be a $(PS)_c$ sequence of J_ε restricted to \mathcal{N}_ε . Then, $J_\varepsilon(u_n) \rightarrow c$ as $n \rightarrow +\infty$ and there exists $\{\lambda_n\} \subset \mathbb{R}$ such that

$$(4.36) \quad J'_\varepsilon(u_n) = \lambda_n T'_\varepsilon(u_n) + o_n(1),$$

with

$$T_\varepsilon : H_\varepsilon \rightarrow \mathbb{R}, \quad T_\varepsilon(u) := \|u\|_\varepsilon^2 - \int_{\mathbb{R}^2} f(|u|^2)|u|^2 dx.$$

Using (f₄), we get

$$\begin{aligned} T'_\varepsilon(u_n)[u_n] &= 2\|u_n\|_\varepsilon^2 - 2 \int_{\mathbb{R}^2} f(|u_n|^2)|u_n|^2 dx - 2 \int_{\mathbb{R}^2} f'(|u_n|^2)|u_n|^4 dx \\ &= -2 \int_{\mathbb{R}^2} f'(|u_n|^2)|u_n|^4 dx \leq -(p-2)C_p \int_{\mathbb{R}^2} |u_n|^p dx \leq 0. \end{aligned}$$

Moreover $\{T'_\varepsilon(u_n)\}$ is bounded. Indeed, using the Hölder inequality, (2.4), and (f₅),

$$\begin{aligned} |T'_\varepsilon(u_n)[\phi]| &\leq 2\|u_n\|_\varepsilon \|\phi\|_\varepsilon + 2 \int_{\mathbb{R}^2} f(|u_n|^2)|u_n||\phi| dx + 2 \int_{\mathbb{R}^2} f'(|u_n|^2)|u_n|^3|\phi| dx \\ &\leq C \left[\|u_n\|_\varepsilon \|\phi\|_\varepsilon + \int_{\mathbb{R}^2} |u_n|^{q-1}|\phi|(e^{\alpha|u_n|^2} - 1) dx + \int_{\mathbb{R}^2} |u_n|^3|\phi|(e^{4\pi|u_n|^2} - 1) dx \right]. \end{aligned}$$

Then, using similar arguments as in (4.35) and taking $\alpha > 4\pi$ and $r > 1$ with $\alpha r < 32\pi$, for n large enough,

$$\begin{aligned} \int_{\mathbb{R}^2} |u_n|^{q-1} |\phi| (e^{\alpha|u_n|^2} - 1) dx &\leq \left(\int_{\mathbb{R}^2} |u_n|^{r'(q-1)} |\phi|^{r'} dx \right)^{1/r'} \left(\int_{\mathbb{R}^2} (e^{\frac{\alpha r}{8}(2\sqrt{2}|u_n|)^2} - 1) dx \right)^{1/r} \\ &\leq C \|u_n\|_\varepsilon^{q-1} \|\phi\|_\varepsilon, \end{aligned}$$

where r' is the conjugate exponent of r .

Analogously

$$\int_{\mathbb{R}^2} |u_n|^3 |\phi| (e^{4\pi|u_n|^2} - 1) dx \leq C \|u_n\|_\varepsilon^3 \|\phi\|_\varepsilon.$$

Thus, using the boundedness of $\{u_n\}$, for every $\phi \in H_\varepsilon$ we have that

$$|T'_\varepsilon(u_n)[\phi]| \leq C(\|u_n\|_\varepsilon + \|u_n\|_\varepsilon^{q-1} + \|u_n\|_\varepsilon^3) \|\phi\|_\varepsilon \leq C \|\phi\|_\varepsilon.$$

Thus, up to a subsequence, $T'_\varepsilon(u_n)[u_n] \rightarrow \varsigma \leq 0$.

We claim that $T'_\varepsilon(u_n)[u_n] \not\rightarrow 0$.

Indeed, if $T'_\varepsilon(u_n)[u_n] \rightarrow 0$, then also $u_n \rightarrow 0$ in $L^p(\mathbb{R}^2, \mathbb{C})$, since

$$o_n(1) = |T'_\varepsilon(u_n)[u_n]| \geq C \int_{\mathbb{R}^2} |u_n|^p dx.$$

Thus, by interpolation, $u_n \rightarrow 0$ in $L^\tau(\mathbb{R}^2, \mathbb{C})$, for all $\tau > 2$.

Moreover, using similar arguments as before and taking $\alpha > 4\pi$ and $r > 1$ with $\alpha r < 32\pi$, we get that, for every $\zeta > 0$ and for n large enough,

$$\|u_n\|_\varepsilon^2 = \int_{\mathbb{R}^2} f(|u_n|^2) |u_n|^2 dx \leq \frac{\zeta}{V_0} \|u_n\|_\varepsilon^2 + C \|u_n\|_{q^r}^q \left(\int_{\mathbb{R}^2} (e^{\frac{\alpha r}{8}(2\sqrt{2}|u_n|)^2} - 1) dx \right)^{1/r'} = \frac{\zeta}{V_0} \|u_n\|_\varepsilon^2 + o_n(1)$$

and so $u_n \rightarrow 0$ in H_ε , which is in contradiction with Lemma 4.2.

Hence, since $\{u_n\} \subset \mathcal{N}_\varepsilon$, by (4.36) we get $\lambda_n = o_n(1)$ and so $J'_\varepsilon(u_n) = o_n(1)$. Then, applying Proposition 4.8, we conclude. \square

Arguing as in the first part of the proof of Proposition 4.9, we have

Corollary 4.10. *The constrained critical points of the functional J_ε on \mathcal{N}_ε are critical points of J_ε in H_ε .*

Now we can complete the proof of Theorem 1.1.

Proof of Theorem 1.1. By Lemma 4.1, we know that the functional J_ε has the Mountain Pass geometry. Thus, there exists a $(PS)_{c_\varepsilon}$ sequence $\{u_n\} \subset H_\varepsilon$ for J_ε , with c_ε defined in (4.3). By Lemma 4.3 we have that $\{u_n\}$ is bounded in H_ε and, by (f3), Lemma 4.4, and (3.4), for ε small enough,

$$\begin{aligned} \frac{\theta - 2}{2\theta} \|u_n\|_\varepsilon^2 &\leq \left(\frac{1}{2} - \frac{1}{\theta} \right) \|u_n\|_\varepsilon^2 + \frac{1}{\theta} \int_{\mathbb{R}^2} \left(f(|u_n|^2) |u_n|^2 - \frac{\theta}{2} F(|u_n|^2) \right) dx = J_\varepsilon(u_n) - \frac{1}{\theta} J'_\varepsilon(u_n)[u_n] \\ &\leq c_\varepsilon + o_n(1) + o_n(1) \|u_n\|_\varepsilon \leq c_{V_0} + o_n(1) + o_n(1) \|u_n\|_\varepsilon \leq c_{V^*} + o_n(1) + o_n(1) \|u_n\|_\varepsilon, \end{aligned}$$

Thus, using Lemma 3.4, for n large enough we have

$$\frac{\theta - 2}{2\theta} \|u_n\|_\varepsilon^2 < \frac{\theta - 2}{16\theta}$$

and so $\limsup_n \|u_n\|_\varepsilon^2 < 1/8$. Hence, (3.4) and Proposition 4.8 allow us to conclude. \square

We conclude this section showing the behavior of the energy levels c_ε of the solutions found in Theorem 1.1.

Proposition 4.11. *Let $\{\varepsilon_n\} \subset (0, +\infty)$ be such that $\varepsilon_n \rightarrow 0^+$ as $n \rightarrow +\infty$ and, for n large enough, u_n be a respective ground state found in Theorem 1.1 for $\varepsilon = \varepsilon_n$ with energy $J_{\varepsilon_n}(u_n) = c_{\varepsilon_n}$. Up to a subsequence, we have that*

$$c_{\varepsilon_n} \rightarrow c_{V_0}.$$

Proof. By Lemma 4.2 and Lemma 4.4, $0 < K_2 \leq c_{\varepsilon_n} \leq c_{V_0}$. Thus, up to a subsequence, $c_{\varepsilon_n} \rightarrow \bar{c} > 0$. Since, using (f₃), Lemma 4.4, the monotonicity of c_μ 's, and Lemma 3.4,

$$\begin{aligned} \frac{\theta - 2}{2\theta} \limsup_n \|u_n\|_{\varepsilon_n}^2 &\leq \limsup_n \left[\left(\frac{1}{2} - \frac{1}{\theta} \right) \|u_n\|_{\varepsilon_n}^2 + \frac{1}{\theta} \int_{\mathbb{R}^2} \left(f(|u_n|^2) |u_n|^2 - \frac{\theta}{2} F(|u_n|^2) \right) dx \right] \\ &= \lim_n \left[J_{\varepsilon_n}(u_n) - \frac{1}{\theta} J'_{\varepsilon_n}(u_n)[u_n] \right] = \bar{c} \leq c_{V_0} < \frac{\theta - 2}{16\theta} \end{aligned}$$

and so, by the diamagnetic inequality (2.1),

$$(4.37) \quad \limsup_n \|\nabla |u_n|\|_2^2 \leq \limsup_n (\|\nabla |u_n|\|_2^2 + V_0 \|u_n\|_2^2) \leq \limsup_n \|u_n\|_{\varepsilon_n}^2 < \frac{1}{8}.$$

Moreover, since $\bar{c} > 0$, we have that $\|u_n\|_{\varepsilon_n} \not\rightarrow 0$. Thus, arguing as in Lemma 4.5, we get that there exist a sequence $(\tilde{y}_n) \subset \mathbb{R}^2$ and constants $R, \gamma > 0$ such that

$$(4.38) \quad \liminf_n \int_{B_R(\tilde{y}_n)} |u_n|^2 dx \geq \gamma.$$

Let $v_n := u_n(\cdot + \tilde{y}_n)$.

By (4.37) we have that $\{|v_n|\}$ is bounded in $H^1(\mathbb{R}^2, \mathbb{R})$ and so, up to a subsequence, $|v_n| \rightharpoonup v$ in $H^1(\mathbb{R}^2, \mathbb{R})$, with $v \neq 0$ by (4.38).

Let now $t_n > 0$ be such that $t_n |v_n| \in \mathcal{N}_{V_0}$ and $y_n := \varepsilon_n \tilde{y}_n$.

Using the diamagnetic inequality (2.1)

$$c_{V_0} \leq I_{V_0}(t_n |v_n|) = I_{V_0}(t_n |u_n|) \leq J_{\varepsilon_n}(t_n u_n) \leq \max_{t \geq 0} J_{\varepsilon_n}(t u_n) = J_{\varepsilon_n}(u_n) = c_{\varepsilon_n} \leq c_{V_0},$$

and we conclude. \square

5. PROOF OF THEOREM 1.2

In this section, by Ljusternik-Schnirelmann category theory, we prove a multiplicity result for problem (2.2).

First, let us provide some useful preliminaries.

Let us start with the following useful compactness result.

Proposition 5.1. *Let $\varepsilon_n \rightarrow 0^+$ and $\{u_n\} \subset \mathcal{N}_{\varepsilon_n}$ be such that $J_{\varepsilon_n}(u_n) \rightarrow c_{V_0}$. Then there exists $\{\tilde{y}_n\} \subset \mathbb{R}^2$ such that the sequence $\{|v_n|\} \subset H^1(\mathbb{R}^2, \mathbb{R})$, where $v_n(x) := u_n(x + \tilde{y}_n)$, has a convergent subsequence in $H^1(\mathbb{R}^2, \mathbb{R})$. Moreover, up to a subsequence, $y_n := \varepsilon_n \tilde{y}_n \rightarrow y \in M$ as $n \rightarrow +\infty$.*

Proof. Since $\{u_n\} \subset \mathcal{N}_{\varepsilon_n}$ and $J_{\varepsilon_n}(u_n) \rightarrow c_{V_0}$, arguing as in the first part of the proof of Lemma 3.5 and using the monotonicity of c_μ 's, we have $\limsup_n \|u_n\|_{\varepsilon_n}^2 < 1/8$ and so, by the diamagnetic inequality (2.1),

$$(5.1) \quad \limsup_n \|\nabla |u_n|\|_2^2 \leq \limsup_n (\|\nabla |u_n|\|_2^2 + V_0 \|u_n\|_2^2) < \frac{1}{8}.$$

Moreover, since $c_{V_0} > 0$, we have that $\|u_n\|_{\varepsilon_n} \not\rightarrow 0$. Thus, arguing as in Lemma 4.5, we get that there exist a sequence $(\tilde{y}_n) \subset \mathbb{R}^2$ and constants $R, \beta > 0$ such that

$$(5.2) \quad \liminf_n \int_{B_R(\tilde{y}_n)} |u_n|^2 dx \geq \beta.$$

Let $v_n(x) = u_n(x + \tilde{y}_n)$. By (5.1) we have that $\{|v_n|\}$ is bounded in $H^1(\mathbb{R}^2, \mathbb{R})$ and so, up to a subsequence, $|v_n| \rightharpoonup v$ in $H^1(\mathbb{R}^2, \mathbb{R})$, with $v \neq 0$ by (5.2).

Let now $t_n > 0$ be such that $\tilde{v}_n := t_n|v_n| \in \mathcal{N}_{V_0}$ and $y_n := \varepsilon_n \tilde{y}_n$.

By the diamagnetic inequality (2.1)

$$c_{V_0} \leq I_{V_0}(\tilde{v}_n) = I_{V_0}(t_n|u_n|) \leq J_{\varepsilon_n}(t_n u_n) \leq \max_{t \geq 0} J_{\varepsilon_n}(t u_n) = J_{\varepsilon_n}(u_n) = c_{V_0} + o_n(1),$$

and so $I_{V_0}(\tilde{v}_n) \rightarrow c_{V_0}$ as $n \rightarrow +\infty$.

Thus, by Lemma 3.5 for $\mu = V_0$, $\{\tilde{v}_n\}$ is bounded in $H^1(\mathbb{R}^2, \mathbb{R})$ and, since $|v_n| \not\rightarrow 0$ in $H^1(\mathbb{R}^2, \mathbb{R})$, then $\{t_n\}$ is bounded too and so, up to a subsequence, $t_n \rightarrow t_0 \geq 0$.

Actually we have that $t_0 > 0$.

Indeed, if $t_0 = 0$, then, due to the boundedness of $\{|v_n|\}$ in $H^1(\mathbb{R}^2, \mathbb{R})$, $\tilde{v}_n \rightarrow 0$ in $H^1(\mathbb{R}^2, \mathbb{R})$, so that $I_{V_0}(\tilde{v}_n) \rightarrow 0$, in contradiction with $c_{V_0} > 0$.

Hence, up to a subsequence, $\tilde{v}_n \rightharpoonup \tilde{v} := t_0 v \neq 0$ in $H^1(\mathbb{R}^2, \mathbb{R})$ and so, by Lemma 3.5, $\tilde{v}_n \rightarrow \tilde{v}$ in $H^1(\mathbb{R}^2, \mathbb{R})$, which gives $|v_n| \rightarrow v$ in $H^1(\mathbb{R}^2, \mathbb{R})$.

Let us show now that $\{y_n\}$ is bounded.

Assume by contradiction that $\{y_n\}$ is not bounded. Then there exists a subsequence such that $|y_n| \rightarrow +\infty$.

If $V_\infty = +\infty$, since, by (V),

$$\inf_{B_R(\tilde{y}_n)} V_{\varepsilon_n} = \inf_{B_{R\varepsilon_n}(y_n)} V \rightarrow +\infty,$$

then

$$\begin{aligned} \int_{\mathbb{R}^2} V(\varepsilon_n x + y_n) |v_n|^2 dx &\geq \int_{B_R(0)} V(\varepsilon_n x + y_n) |v_n|^2 dx = \int_{B_R(0)} V(\varepsilon_n x + y_n) |u_n(x + \tilde{y}_n)|^2 dx \\ &= \int_{B_R(\tilde{y}_n)} V(\varepsilon_n z) |u_n(z)|^2 dz \geq \inf_{B_R(\tilde{y}_n)} V_{\varepsilon_n} \int_{B_R(\tilde{y}_n)} |u_n(z)|^2 dz \rightarrow +\infty, \end{aligned}$$

in contradiction with

$$\int_{\mathbb{R}^2} V(\varepsilon_n x + y_n) |v_n|^2 dx \leq \|u_n\|_{\varepsilon_n}^2 \leq C.$$

If $V_\infty \in \mathbb{R}$, since $\tilde{v}_n \rightarrow \tilde{v}$ in $H^1(\mathbb{R}^2, \mathbb{R})$, by the Fatou's Lemma and since $H^1(\mathbb{R}^2, \mathbb{R})$ is continuously embedded in $L^2(\mathbb{R}^2, \mathbb{R})$,

$$V_\infty \lim_n \int_{\mathbb{R}^2} |\tilde{v}_n|^2 dx = V_\infty \int_{\mathbb{R}^2} |\tilde{v}|^2 dx \leq \liminf_n \int_{\mathbb{R}^2} V(\varepsilon_n x + y_n) |\tilde{v}_n|^2 dx,$$

and so, by the diamagnetic inequality (2.1) and (4.4),

$$\begin{aligned} c_{V_0} &= I_{V_0}(\tilde{v}) < I_{V_\infty}(\tilde{v}) = \lim_n I_{V_\infty}(\tilde{v}_n) \\ &\leq \liminf_n \frac{1}{2} \int_{\mathbb{R}^2} V(\varepsilon_n x + y_n) |\tilde{v}_n|^2 dx - \frac{V_\infty}{2} \int_{\mathbb{R}^2} |\tilde{v}|^2 dx + \lim_n I_{V_\infty}(\tilde{v}_n) \\ (5.3) \quad &\leq \liminf_n \left(\frac{1}{2} \int_{\mathbb{R}^2} (|\nabla \tilde{v}_n|^2 + V(\varepsilon_n x + y_n) |\tilde{v}_n|^2) dx - \frac{1}{2} \int_{\mathbb{R}^2} F(|\tilde{v}_n|^2) dx \right) \\ &= \liminf_n \left(\frac{t_n^2}{2} \int_{\mathbb{R}^2} (|\nabla |u_n||^2 + V_{\varepsilon_n}(z) |u_n|^2) dz - \frac{1}{2} \int_{\mathbb{R}^2} F(t_n^2 |u_n|^2) dz \right) \\ &\leq \liminf_n J_{\varepsilon_n}(t_n u_n) \leq \liminf_n J_{\varepsilon_n}(u_n) = c_{V_0}, \end{aligned}$$

which gives a contradiction.

Hence, up to a subsequence, $y_n \rightarrow y \in \mathbb{R}^2$.

If $y \notin M$, then $V(y) > V_0$ and we can argue as in (5.3) to get a contradiction and so the proof is complete. \square

Let now $\delta > 0$ be fixed, $\omega \in H^1(\mathbb{R}^2, \mathbb{R})$ be a positive ground state solution of problem (3.1) for $\mu = V_0$, and $\eta \in C^\infty([0, +\infty), [0, 1])$ be a non-increasing cut-off function such that $\eta(t) = 1$ if $0 \leq t \leq \delta/2$ and $\eta(t) = 0$ if $t \geq \delta$.

For any $y \in M$, let

$$\Psi_{\varepsilon,y}(x) := \eta(|\varepsilon x - y|) \omega\left(\frac{\varepsilon x - y}{\varepsilon}\right) \exp\left(i\tau_y\left(\frac{\varepsilon x - y}{\varepsilon}\right)\right),$$

where M is defined in (1.4) and $\tau_y(x) := A_1(y)x_1 + A_2(y)x_2$.

Let $t_{\varepsilon,y} > 0$ be the unique positive number such that $t_{\varepsilon,y}\Psi_{\varepsilon,y} \in \mathcal{N}_\varepsilon$ (and so $\max_{t \geq 0} J_\varepsilon(t\Psi_{\varepsilon,y}) = J_{\varepsilon,y}(t_{\varepsilon,y}\Psi_{\varepsilon,y})$) and let us consider

$$\Phi_\varepsilon := y \in M \mapsto t_{\varepsilon,y}\Psi_{\varepsilon,y} \in \mathcal{N}_\varepsilon.$$

Observe that, by the assumptions on V , M is compact and, for any $y \in M$, $\Phi_\varepsilon(y)$ has compact support. Moreover we have

Lemma 5.2. *As $\varepsilon \rightarrow 0^+$,*

$$\|\Psi_{\varepsilon,y}\|_\varepsilon^2 \rightarrow \int_{\mathbb{R}^2} (|\nabla \omega|^2 + V_0 \omega^2) dx$$

uniformly with respect to $y \in M$.

Proof. First we observe that

$$\sup_{y \in M} \left| \int_{\mathbb{R}^2} V_\varepsilon(x) |\Psi_{\varepsilon,y}|^2 dx - V_0 \int_{\mathbb{R}^2} \omega^2 dx \right| \rightarrow 0 \text{ as } \varepsilon \rightarrow 0^+.$$

Indeed, taking $z = (\varepsilon x - y)/\varepsilon$,

$$\sup_{y \in M} \left| \int_{\mathbb{R}^2} |\Psi_{\varepsilon,y}|^2 dx - \int_{\mathbb{R}^2} \omega^2 dx \right| \leq \int_{\mathbb{R}^2} |\eta^2(|\varepsilon z|) - 1| \omega^2(z) dz \leq \int_{B_{\delta/(2\varepsilon)}^c(0)} \omega^2(z) dz \rightarrow 0.$$

Moreover

$$\begin{aligned} \sup_{y \in M} \left| \int_{\mathbb{R}^2} [V_\varepsilon(x) - V_0] |\Psi_{\varepsilon,y}|^2 dx \right| &\leq \sup_{y \in M} \int_{\mathbb{R}^2} |V(\varepsilon z + y) - V_0| \eta^2(|\varepsilon z|) \omega^2(z) dz \\ &\leq \int_{B_{\delta/\varepsilon}(0)} \sup_{y \in M} |V(\varepsilon z + y) - V_0| \omega^2(z) dz \\ &= \int_{\mathbb{R}^2} \sup_{y \in M} |V(\varepsilon z + y) - V_0| \chi_{B_{\delta/\varepsilon}(0)}(z) \omega^2(z) dz \rightarrow 0, \end{aligned}$$

applying the Lebesgue Dominated Convergence Theorem, since, if $|\varepsilon z| < \delta$, then, due to the compactness of M , $|\varepsilon z + y|$ is bounded when $y \in M$,

$$\sup_{y \in M} |V(\varepsilon z + y) - V_0| \chi_{B_{\delta/\varepsilon}(0)}(z) \omega^2(z) \leq C \omega^2(z) \in L^1(\mathbb{R}^2, \mathbb{R}),$$

and $\sup_{y \in M} |V(\varepsilon z + y) - V_0| \chi_{B_{\delta/\varepsilon}(0)}(z) \omega^2(z)$ converges to 0 a.e. in \mathbb{R}^2 as $\varepsilon \rightarrow 0^+$.

Then, since, taking again $z = (\varepsilon x - y)/\varepsilon$,

$$\begin{aligned} \int_{\mathbb{R}^2} |\nabla_{A_\varepsilon} \Psi_{\varepsilon,y}|^2 dx &= \varepsilon^2 \int_{\mathbb{R}^2} |\eta'(|\varepsilon z|)|^2 \omega^2(z) dz + \int_{\mathbb{R}^2} \eta^2(|\varepsilon z|) |\nabla \omega(z)|^2 dz \\ &\quad + \int_{\mathbb{R}^2} |A(y) - A(\varepsilon z + y)|^2 \eta^2(|\varepsilon z|) \omega^2(z) dz \\ &\quad + 2\varepsilon \int_{\mathbb{R}^2} \eta(|\varepsilon z|) \eta'(|\varepsilon z|) \omega(z) \nabla \omega(z) \cdot \frac{z}{|z|} dz, \end{aligned}$$

and,

$$\begin{aligned}
\left| \int_{\mathbb{R}^2} \eta^2(|\varepsilon z|) |\nabla \omega(z)|^2 dz - \int_{\mathbb{R}^2} |\nabla \omega(z)|^2 dz \right| &\leq \int_{\mathbb{R}^2} |\eta^2(|\varepsilon z|) - 1| |\nabla \omega(z)|^2 dz \leq \int_{B_{\delta/(2\varepsilon)}^c(0)} |\nabla \omega(z)|^2 dz, \\
\int_{\mathbb{R}^2} |\eta'(|\varepsilon z|)|^2 \omega^2(z) dz &\leq C \int_{B_{\delta/(2\varepsilon)}^c(0)} \omega^2(z) dz, \\
\int_{\mathbb{R}^2} |\eta(|\varepsilon z|) \eta'(|\varepsilon z|) \omega(z) \nabla \omega(z)| dz &\leq C \int_{B_{\delta/(2\varepsilon)}^c(0)} \omega(z) |\nabla \omega| dz, \\
\sup_{y \in M} \int_{\mathbb{R}^2} |A(y) - A(\varepsilon z + y)|^2 \eta^2(|\varepsilon z|) \omega^2(z) dz &\leq \int_{B_{\delta/\varepsilon}(0)} \sup_{y \in M} |A(y) - A(\varepsilon z + y)|^2 \omega^2(z) dz \rightarrow 0,
\end{aligned}$$

due to the continuity of A and a similar argument as before.

Thus we infer

$$\sup_{y \in M} \left| \int_{\mathbb{R}^2} |\nabla_{A_\varepsilon} \Psi_{\varepsilon, y}|^2 dx - \int_{\mathbb{R}^2} |\nabla \omega|^2 dx \right| \rightarrow 0$$

as $\varepsilon \rightarrow 0^+$ and we conclude. \square

Hence we can conclude in the following way.

Proof of Theorem 1.2. Arguing as in [20, Lemma 4.1], and using Lemma 5.2, we can prove that $J_\varepsilon(\Phi_\varepsilon(y)) \rightarrow c_{V_0}$ uniformly with respect to $y \in M$ as $\varepsilon \rightarrow 0^+$.

Then, given $\delta > 0$, if $\rho > 0$ with $M_\delta \subset B_\rho$, let

$$\Upsilon : \mathbb{R}^2 \rightarrow \mathbb{R}^2, \quad \Upsilon(x) := \begin{cases} x, & \text{if } |x| < \rho, \\ \rho x/|x|, & \text{if } |x| \geq \rho \end{cases}$$

and then let us define the barycenter map as

$$\beta_\varepsilon := u \in \mathcal{N}_\varepsilon \mapsto \frac{1}{\|u\|_2^2} \int_{\mathbb{R}^2} \Upsilon(\varepsilon x) |u(x)|^2 dx \in \mathbb{R}^2.$$

By [20, Lemma 4.2] we have that $\beta_\varepsilon(\Phi_\varepsilon(y)) \rightarrow y$ as $\varepsilon \rightarrow 0^+$ uniformly in $y \in M$.

Let now $h : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ be any positive function such that $h(\varepsilon) \rightarrow 0$ as $\varepsilon \rightarrow 0^+$ and

$$\tilde{\mathcal{N}}_\varepsilon := \{u \in \mathcal{N}_\varepsilon : J_\varepsilon(u) \leq c_{V_0} + h(\varepsilon)\}.$$

Since $|J_\varepsilon(\Phi_\varepsilon(y)) - c_{V_0}| \rightarrow 0$ uniformly in $y \in M$ as $\varepsilon \rightarrow 0^+$, then $\tilde{\mathcal{N}}_\varepsilon \neq \emptyset$ for $\varepsilon > 0$ small enough.

Thus, arguing as in [20, Lemma 4.4],

$$\lim_{\varepsilon \rightarrow 0^+} \sup_{u \in \tilde{\mathcal{N}}_\varepsilon} \text{dist}(\beta_\varepsilon(u), M_\delta) = 0.$$

Hence, as in [17, Section 6], we can find $\tilde{\varepsilon}_\delta > 0$ such that for any $\varepsilon \in (0, \tilde{\varepsilon}_\delta)$, the following diagram

$$M \xrightarrow{\Phi_\varepsilon} \tilde{\mathcal{N}}_\varepsilon \xrightarrow{\beta_\varepsilon} M_\delta$$

is well defined and $\beta_\varepsilon \circ \Phi_\varepsilon$ is homotopically equivalent to the embedding $\iota : M \rightarrow M_\delta$. Thus, [11, Lemma 4.3] (see also [18, Lemma 2.2]) implies that

$$(5.4) \quad \text{cat}_{\tilde{\mathcal{N}}_\varepsilon}(\tilde{\mathcal{N}}_\varepsilon) \geq \text{cat}_{M_\delta}(M).$$

Now observe that J_ε is bounded from below on \mathcal{N}_ε (see the preliminary results in Section 4 and in particular (4.5)). Moreover, using, respectively, Lemma 4.4, Lemma 3.4, and (3.4), for ε small enough,

$$\begin{aligned}
c_\varepsilon &\leq c_{V_0} < c_{V_0} + h(\varepsilon), \\
c_{V^*} + h(\varepsilon) &< \frac{\theta - 2}{16\theta},
\end{aligned}$$

and, if $V_\infty < +\infty$,

$$(5.5) \quad c_{V_0} + h(\varepsilon) < c_{V_\infty}.$$

Thus, if $\{u_n\}$ is a $(PS)_c$ sequence for $J_\varepsilon|_{\mathcal{N}_\varepsilon}$ with $c \in [c_\varepsilon, c_{V_0} + h(\varepsilon)]$, then, by (f₃) and (3.4),

$$\begin{aligned} \frac{\theta-2}{2\theta} \|u_n\|_\varepsilon^2 &\leq \left(\frac{1}{2} - \frac{1}{\theta}\right) \|u_n\|_\varepsilon^2 + \frac{1}{\theta} \int_{\mathbb{R}^2} \left(f(|u_n|^2)|u_n|^2 - \frac{\theta}{2} F(|u_n|^2)\right) dx \\ &= J_\varepsilon(u_n) - \frac{1}{\theta} J'_\varepsilon(u_n)[u_n] = c + o_n(1) \\ &< c_{V_0} + h(\varepsilon) + o_n(1) \leq c_{V^*} + h(\varepsilon) + o_n(1). \end{aligned}$$

Then, passing to the lim sup on n we have

$$\frac{\theta-2}{2\theta} \limsup_n \|u_n\|_\varepsilon^2 \leq c_{V^*} + h(\varepsilon) < \frac{\theta-2}{16\theta}$$

for ε small enough and so

$$\limsup_n \|u_n\|_\varepsilon^2 < \frac{1}{8}.$$

Hence, using (5.5) if $V_\infty < +\infty$, we can apply Proposition 4.9, obtaining that $J_\varepsilon|_{\mathcal{N}_\varepsilon}$ satisfies the $(PS)_c$ condition for $c \in [c_\varepsilon, c_{V_0} + h(\varepsilon)]$. Then, by [33, Theorem 5.20], the Ljusternik-Schnirelmann theory for C^1 functionals implies that $\tilde{\mathcal{N}}_\varepsilon$ contains at least $\text{cat}_{\tilde{\mathcal{N}}_\varepsilon}(\tilde{\mathcal{N}}_\varepsilon)$ critical points of $J_\varepsilon|_{\mathcal{N}_\varepsilon}$. Observing that, due to Corollary 4.10, such critical points are critical points for J_ε and using (5.4), we conclude the proof of the multiplicity result. \square

6. PROOF OF THEOREM 1.3

This last section is dedicated to the study of the behavior of the maximum points of the modulus of the solutions obtained in Theorem 1.1 and in Theorem 1.2 and to their the decay property.

Proof of Theorem 1.3. Let $\varepsilon_n \rightarrow 0^+$ as $n \rightarrow +\infty$ and let u_{ε_n} be a nontrivial solution of (2.2) found in Theorem 1.1 or in Theorem 1.2. For ground states found in Theorem 1.1, by Proposition 4.11, up to a subsequence, we have that $J_{\varepsilon_n}(u_{\varepsilon_n}) = c_{\varepsilon_n} \rightarrow c_{V_0}$ and, for solutions found in Theorem 1.2 we have that $J_{\varepsilon_n}(u_{\varepsilon_n}) \leq c_{V_0} + h(\varepsilon_n)$. Thus, Proposition 5.1 applies and so we can consider $\{\tilde{y}_n\} \subset \mathbb{R}^2$, and $v_{\varepsilon_n} := u_{\varepsilon_n}(\cdot + \tilde{y}_n)$, which is a solution of

$$\left(\frac{1}{i}\nabla - A_{\varepsilon_n}(x + \tilde{y}_n)\right)^2 v_{\varepsilon_n} + V_{\varepsilon_n}(x + \tilde{y}_n)v_{\varepsilon_n} = f(|v_{\varepsilon_n}|^2)v_{\varepsilon_n} \quad \text{in } \mathbb{R}^2,$$

such that, up to a subsequence, $|v_{\varepsilon_n}| \rightarrow v$ in $H^1(\mathbb{R}^2, \mathbb{R})$ and $y_n = \varepsilon_n \tilde{y}_n \rightarrow y \in M$.

Arguing as in [20, Lemma 5.1], we have that $\{|v_{\varepsilon_n}|\}$ is bounded in $L^\infty(\mathbb{R}^2, \mathbb{R})$ and

$$(6.1) \quad \lim_{|x| \rightarrow +\infty} |v_{\varepsilon_n}(x)| = 0 \quad \text{uniformly in } n \in \mathbb{N}.$$

In addition, by the first part of the proof of Proposition 5.1, there exist $R, \beta > 0$ such that

$$\liminf_n \int_{B_R(\tilde{y}_n)} |u_{\varepsilon_n}|^2 dx \geq \beta.$$

Then, if $\kappa > 0$ such that $0 < \pi\kappa^2 R^2 < \beta$, for n large enough,

$$\pi\kappa^2 R^2 \leq \int_{B_R(\tilde{y}_n)} |u_{\varepsilon_n}|^2 dx \leq \pi R^2 \|v_{\varepsilon_n}\|_\infty^2,$$

and so

$$(6.2) \quad \|v_{\varepsilon_n}\|_\infty \geq \kappa.$$

Let us show the concentration of global maximum points p_n of $|v_{\varepsilon_n}|$.

If $z_n := p_n + \tilde{y}_n$, by the definition of v_{ε_n} , we have that each z_n is a global maximum point of $|u_{\varepsilon_n}|$. Moreover, since by (6.1) and (6.2), $p_n \in B_R(0)$ for some $R > 0$, $\eta_n := \varepsilon_n z_n = \varepsilon_n p_n + \varepsilon_n \tilde{y}_n \rightarrow y \in M$ and so

$$\lim_n V(\eta_n) = V_0.$$

Now let us assume that $A \in C^1(\mathbb{R}^2, \mathbb{R}^2)$ and let us show decay of the modulus of our solutions.

Arguing, for instance, as in [16, Remark 1.2], $v_{\varepsilon_n} \in C^1(\mathbb{R}^2, \mathbb{C})$. Moreover, by (6.1) and (f₁), there exists a $R_1 > 0$ such that

$$(6.3) \quad f(|v_{\varepsilon_n}|^2) \leq \frac{V_0}{2} \text{ for all } |x| \geq R_1 \text{ and uniformly in } n \in \mathbb{N}.$$

Since $v_{\varepsilon_n} \in C^1(\mathbb{R}^2, \mathbb{C})$ and (6.1) holds, there exist $c, C > 0$ with $c^2 < V_0/2$ such that

$$|v_{\varepsilon_n}(x)| \leq Ce^{-cR_1} \text{ for } |x| = R_1.$$

Let $\varphi(x) := Ce^{-c|x|}$ in \mathbb{R}^2 .

Observe that

$$(6.4) \quad \Delta\varphi \leq c^2\varphi \text{ in } \mathbb{R}^2 \setminus \{0\}.$$

Moreover, by the Kato's inequality (see [27, Lemma A]) and (V), $|v_{\varepsilon_n}|$ satisfies (in the weak sense)

$$-\Delta|v_{\varepsilon_n}| + V_0|v_{\varepsilon_n}| \leq f(|v_{\varepsilon_n}|^2)|v_{\varepsilon_n}| \text{ in } \mathbb{R}^2.$$

Thus, by (6.3), we have that, uniformly in $n \in \mathbb{N}$,

$$(6.5) \quad -\Delta|v_{\varepsilon_n}| + \frac{V_0}{2}|v_{\varepsilon_n}| \leq 0 \text{ for } |x| \geq R_1.$$

Let, now, $\varphi_n := \varphi - |v_{\varepsilon_n}|$.

By (6.4) and (6.5), we have

$$\begin{cases} -\Delta\varphi_n + \frac{V_0}{2}\varphi_n \geq 0 & \text{for } |x| \geq R_1, \\ \varphi_n \geq 0 & \text{for } |x| = R_1, \\ \lim_{|x| \rightarrow +\infty} \varphi_n(x) = 0. \end{cases}$$

The classical maximum principle (see e.g. [14, Theorem 7.4.2]) implies that $\varphi_n \geq 0$ for $|x| \geq R_1$ and so we obtain that

$$|u_{\varepsilon_n}(x)| = |v_{\varepsilon_n}(x - \tilde{y}_n)| \leq Ce^{-c|x - \tilde{y}_n|} \text{ for all } |x| \geq R_1 \text{ and uniformly in } n \in \mathbb{N}.$$

Observe that, due to the change of variable introduced in Section 2 to write (1.1) in the equivalent form (2.2), the points η_n are maximizers of solutions of (1.1) and (1.5) holds. \square

ACKNOWLEDGMENTS

P. d'Avenia is supported by PRIN project 2017JPCAPN *Qualitative and quantitative aspects of non-linear PDEs*. C. Ji is partially supported by Shanghai Natural Science Foundation (20ZR1413900).

REFERENCES

- [1] C.O. Alves, D. Cassani, C. Tarsi, M. Yang, Existence and concentration of ground state solutions for a critical nonlocal Schrödinger equation in \mathbb{R}^2 , J. Differential Equations 261 (2016), 1933–1972. 2, 6
- [2] C.O. Alves, G.M. Figueiredo, Existence and multiplicity of positive solutions to a p -Laplacian equations in \mathbb{R}^N , Differential Integral Equations 19 (2006), 143–162. 2
- [3] C.O. Alves, G.M. Figueiredo, On multiplicity and concentration of positive solutions for a class of quasilinear problems with critical exponential growth in \mathbb{R}^N , J. Differential Equations 246 (2009), 1288–1311. 2
- [4] C.O. Alves, G.M. Figueiredo, M.F. Furtado, Multiple solutions for a nonlinear Schrödinger equation with magnetic fields, Comm. Partial Differential Equations 36 (2011), 1565–1586. 2

- [5] C.O. Alves, M.A.S. Souto, Multiplicity of positive solutions for a class of problems with exponential critical growth in \mathbb{R}^2 , *J. Differential Equations* 244 (2008), 1502–1520. [2](#)
- [6] C.O. Alves, M.A.S. Souto, M. Montenegro, Existence of a ground state solution for a nonlinear scalar field equation with critical growth, *Calc. Var. Partial Differential Equations* 43 (2012), 537–554. [12](#)
- [7] V. Ambrosio, On a fractional magnetic Schrödinger equation in \mathbb{R} with exponential critical growth, *Nonlinear Anal.* 183 (2019), 117–148. [2](#)
- [8] V. Ambrosio, P. d’Avenia, Nonlinear fractional magnetic Schrödinger equation: existence and multiplicity, *J. Differential Equations* 264 (2018), 3336–3368. [2](#)
- [9] G. Arioli, A. Szulkin, A semilinear Schrödinger equation in the presence of a magnetic field, *Arch. Ration. Mech. Anal.* 170 (2003), 277–295. [2](#)
- [10] S. Barile, G.M. Figueiredo, An existence result for Schrödinger equations with magnetic fields and exponential critical growth, *J. Elliptic Parabol. Equ.* 3 (2017), 105–125. [2](#)
- [11] V. Benci, G. Cerami, Multiple positive solutions of some elliptic problems via the Morse theory and the domain topology, *Calc. Var. Partial Differential Equations* 2 (1994), 29–48. [32](#)
- [12] H. Brezis, *Functional analysis, Sobolev spaces and partial differential equations*, Universitext, Springer, New York, 2011. [5](#), [6](#), [9](#)
- [13] D.M. Cao, Nontrivial solution of semilinear elliptic equation with critical exponent in \mathbb{R}^2 , *Comm. Partial Differential Equations* 17 (1992), 407–435. [6](#)
- [14] W. Chen, C. Li, *Methods on nonlinear elliptic equations*, AIMS Series on Differential Equations & Dynamical Systems 4, American Institute of Mathematical Sciences (AIMS), Springfield, 2010. [34](#)
- [15] S. Cingolani, Semiclassical stationary states of nonlinear Schrödinger equations with an external magnetic field, *J. Differential Equations* 188 (2003), 52–79. [2](#)
- [16] S. Cingolani, L. Jeanjean, S. Secchi, Multi-peak solutions for magnetic NLS equations without non-degeneracy conditions, *ESAIM Control Optim. Calc. Var.* 15 (2009), 653–675. [2](#), [34](#)
- [17] S. Cingolani, M. Lazzo, Multiple semiclassical standing waves for a class of nonlinear Schrödinger equations, *Topol. Methods Nonlinear Anal.* 10 (1997), 1–13. [2](#), [32](#)
- [18] S. Cingolani, M. Lazzo, Multiple positive solutions to nonlinear Schrödinger equations with competing potential functions, *J. Differential Equations* 160 (2000), 118–138. [32](#)
- [19] S. Cingolani, S. Secchi, Semiclassical states for NLS equations with magnetic potentials having polynomial growths, *J. Math. Phys.* 46 (2005), 053503, 19pp. [2](#)
- [20] P. d’Avenia, C. Ji, Multiplicity and concentration results for a magnetic Schrödinger equation with exponential critical growth in \mathbb{R}^2 , to appear on *Int. Math. Res. Not. IMRN*, doi: [10.1093/imrn/rnaa074](#). [2](#), [3](#), [11](#), [12](#), [19](#), [32](#), [33](#)
- [21] P. d’Avenia, M. Squassina, Ground states for fractional magnetic operators, *ESAIM Control Optim. Calc. Var.* 24 (2018), 1–24. [2](#)
- [22] M. del Pino, P.L. Felmer, local mountain passes for semilinear elliptic problems in unbounded domains, *Calc. Var. Partial Differ. Equ.* 4 (1996) 121–137. [2](#)
- [23] M. Del Pino, P. Felmer, P. Sternberg, Boundary concentration for eigenvalue problems related to the onset of superconductivity, *Comm. Math. Phys.* 210 (2000), 413–446. [2](#)
- [24] Y. Ding, F. Lin, Solutions of perturbed Schrödinger equations with critical nonlinearity, *Calc. Var. Partial Differ. Equ.* 30 (2007), 231–249. [7](#)
- [25] M.J. Esteban, P.-L. Lions, Stationary solutions of nonlinear Schrödinger equations with an external magnetic field, *Partial differential equations and the calculus of variations I*, 401–449, *Progr. Nonlinear Differential Equations Appl.* 1, Birkhäuser Boston, Boston, 1989. [2](#), [6](#)
- [26] B. Helffer, A. Morame, Magnetic bottles in connection with superconductivity, *J. Funct. Anal.* 185 (2001), 604–680. [2](#)
- [27] T. Kato, Schrödinger operators with singular potentials, *Israel J. Math.* 13 (1972), 135–148. [34](#)
- [28] O. Kavian, *Introduction à la théorie des points critiques et applications aux problèmes elliptiques*, Mathématiques & Applications 13, Springer-Verlag, Paris, 1993. [15](#), [20](#)
- [29] K. Kurata, Existence and semi-classical limit of the least energy solution to a nonlinear Schrödinger equation with electromagnetic fields, *Nonlinear Anal.* 41 (2000), 763–778. [2](#)
- [30] S. Liang, S. Shi, On multi-bump solutions of nonlinear Schrödinger equation with electromagnetic fields and critical nonlinearity in \mathbb{R}^N , *Calc. Var. Partial Differential Equations* 56 (2017), 25, 29 pp. [2](#)
- [31] E.H. Lieb, M. Loss, *Analysis*, Second edition, Graduate Studies in Mathematics 14, American Mathematical Society, Providence, 2001. [4](#)
- [32] P.H. Rabinowitz, On a class of nonlinear Schrödinger equations, *Z. Angew. Math. Phys.* 43 (1992), 270–291. [2](#)
- [33] M. Willem, *Minimax theorems*, Progress in Nonlinear Differential Equations and their Applications 24, Birkhäuser Boston, Boston, 1996. [12](#), [13](#), [14](#), [17](#), [19](#), [33](#)
- [34] J. Zhang, J.M. do Ó, Standing waves for nonlinear Schrödinger equations involving critical growth of Trudinger-Moser type, *Z. Angew. Math. Phys.* 66 (2015), 3049–3060. [12](#)

(P. d'Avenia)

DIPARTIMENTO DI MECCANICA, MATEMATICA E MANAGEMENT
POLITECNICO DI BARI

VIA ORABONA 4, 70125 BARI, ITALY

Email address: pietro.davenia@poliba.it

(C. Ji)

DEPARTMENT OF MATHEMATICS

EAST CHINA UNIVERSITY OF SCIENCE AND TECHNOLOGY

SHANGHAI 200237, PR CHINA

Email address: jichao@ecust.edu.cn