

DG-LMC: A Turn-key and Scalable Synchronous Distributed MCMC Algorithm via Langevin Monte Carlo within Gibbs

Vincent Plassier^{1,2,*}, Maxime Vono^{1,*}, Alain Durmus^{3,*}, and Eric Moulines²

¹Lagrange Mathematics and Computing Research Center, Paris, France

²Ecole Polytechnique, Université Paris-Saclay, Palaiseau, France

³Ecole Normale Supérieure Paris-Saclay, Gif-sur-Yvette, France

^{*}Equal contribution

November 16, 2021

Abstract

Performing reliable Bayesian inference on a big data scale is becoming a keystone in the modern era of machine learning. A workhorse class of methods to achieve this task are Markov chain Monte Carlo (MCMC) algorithms and their design to handle distributed datasets has been the subject of many works. However, existing methods are not completely either reliable or computationally efficient. In this paper, we propose to fill this gap in the case where the dataset is partitioned and stored on computing nodes within a cluster under a master/slaves architecture. We derive a user-friendly centralised distributed MCMC algorithm with provable scaling in high-dimensional settings. We illustrate the relevance of the proposed methodology on both synthetic and real data experiments.

1 Introduction

In the current machine learning era, data acquisition has seen significant progress due to rapid technological advances which now allow for more accurate, cheaper and faster data storage and collection. This data quest is motivated by modern machine learning techniques and algorithms which are now well-proven and have become common tools for data analysis. In most cases, the empirical success of these methods are based on a very large sample size (Bardenet et al., 2017; Bottou et al., 2018). This need for data is also theoretically justified by data probabilistic modelling which asserts that under appropriate conditions, the more data can be processed, the more accurate the inference can be performed. However, in recent years, several challenges have emerged regarding the use and access to data in mainstream machine learning methods. Indeed, first the amount of data is now so large

that it has outpaced the increase in computation power of computing resources (Verbraeken et al., 2020). Second, in many modern applications, data storage and/or use are not on a single machine but shared across several units (Raicu et al., 2006; Bernstein and Newcomer, 2009). Third, life privacy is becoming a prime concern for many users of machine learning applications who are therefore asking for methods preserving data anonymity (Shokri and Shmatikov, 2015; Abadi et al., 2016). Distributed machine learning aims at tackling these issues. One of its popular paradigms, referred to as data-parallel approach, is to consider that the training data are divided across multiple machines. Each of these units constitutes a worker node of a computing network and can perform a *local* inference based on the data it has access. Regarding the choice of the network, several options and frameworks have been considered. We focus here on the master/slaves architecture where the worker nodes communicate with each other through a device called the *master* node.

Under this framework, we are interested in carrying Bayesian inference about a parameter $\theta \in \mathbb{R}^d$ based on observed data $\{\mathbf{y}_k\}_{k=1}^n \in \mathcal{Y}^n$ (Robert, 2001). The dataset is assumed to be partitioned into S *shards* and stored on S machines among a collection of b worker nodes. The subset of observations associated to worker $i \in [b]$ is denoted by \mathbf{y}_i , with potentially $\mathbf{y}_i = \{\emptyset\}$ if $i \in [S + 1 : b]$, $b > S$. The posterior distribution of interest is assumed to admit a density w.r.t. the d -dimensional Lebesgue measure which factorises across workers, *i.e.*,

$$\pi(\theta \mid \mathbf{y}_{1:n}) = Z_\pi^{-1} \prod_{i=1}^b e^{-U_{\mathbf{y}_i}(\mathbf{A}_i \theta)}, \quad (1)$$

where $Z_\pi = \int_{\mathbb{R}^d} \prod_{i=1}^b e^{-U_{\mathbf{y}_i}(\mathbf{A}_i \theta)} d\theta$ is a normalisation constant and $\mathbf{A}_i \in \mathbb{R}^{d_i \times d}$ are matrices that might act on the parameter of interest. For $i \in [b]$, the potential function $U_{\mathbf{y}_i} : \mathbb{R}^{d_i} \rightarrow \mathbb{R}$ is assumed to depend only on the subset of observations \mathbf{y}_i . Note that for $i \in [S + 1 : b]$, $b > S$, $U_{\mathbf{y}_i}$ does not depend on the data but only on the prior. For the sake of brevity, the dependency of π w.r.t. the observations $\{\mathbf{y}_i\}_{i=1}^b$ is notationally omitted and for $i \in [b]$, $U_{\mathbf{y}_i}$ is simply denoted by U_i .

To sample from π given by (1) in a distributed fashion, a large number of approximate methods have been proposed in the past ten years (Neiswanger et al., 2014; Ahn et al., 2014; Rabinovich et al., 2015; Scott et al., 2016; Nemeth and Sherlock, 2018; Chowdhury and Jermaine, 2018; Rendell et al., 2021). Despite multiple research lines, to the best of authors' knowledge, none of these proposals has been proven to be satisfactory. Indeed, the latter are not completely either computationally efficient in high-dimensional settings, reliable or theoretically grounded (Jordan et al., 2019).

This work is an attempt to fill this gap. To this purpose, we follow the data augmentation approach introduced in Vono et al. (2021) and referred to as asymptotically exact data augmentation (AXDA). Given a tolerance parameter ρ , the main idea behind this method-

ology is to consider a joint distribution $\Pi_{\boldsymbol{\rho}}$ on the extended state space $\mathbb{R}^d \times \prod_{i=1}^b \mathbb{R}^{d_i}$ such that $\Pi_{\boldsymbol{\rho}}$ has a density w.r.t. the Lebesgue measure of the form $(\boldsymbol{\theta}, \mathbf{z}_{1:b}) \mapsto \prod_{i=1}^b \Pi_{\boldsymbol{\rho}}^i(\boldsymbol{\theta}, \mathbf{z}_i)$, with $\boldsymbol{\theta} \in \mathbb{R}^d$ and $\mathbf{z}_i \in \mathbb{R}^{d_i}$, $i \in [b]$. $\Pi_{\boldsymbol{\rho}}$ is carefully designed so that its marginal w.r.t. $\boldsymbol{\theta}$, denoted by $\pi_{\boldsymbol{\rho}}$, is a proxy of (1) for which quantitative approximation bounds can be derived and are controlled by $\boldsymbol{\rho}$. In addition, for any $i \in [b]$, $\Pi_{\boldsymbol{\rho}}^i(\boldsymbol{\theta}, \mathbf{z}_i)$ only depends on the data \mathbf{y}_i , and therefore plays a role similar to the local posterior $\pi^i(\boldsymbol{\theta}) \propto e^{-U_i(\mathbf{A}_i \boldsymbol{\theta})}$ in popular embarrassingly parallel approaches (Neiswanger et al., 2014; Scott et al., 2016). However, compared to this class of methods, AXDA does not seek for each worker to sample from $\Pi_{\boldsymbol{\rho}}^i$. Following a data augmentation strategy based on Gibbs sampling, AXDA instead requires each worker to sample from the conditional distribution $\Pi_{\boldsymbol{\rho}}(\mathbf{z}_i | \boldsymbol{\theta})$ and to communicate its sample to the master. $\Pi_{\boldsymbol{\rho}}$ is generally chosen such that sampling from $\Pi_{\boldsymbol{\rho}}(\boldsymbol{\theta} | \mathbf{z}_{1:b})$ is easy and does not require access to the data. However, two main challenges remain: one has to sample efficiently from the conditional distribution $\Pi_{\boldsymbol{\rho}}(\mathbf{z}_i | \boldsymbol{\theta})$ for $i \in [b]$ and avoid too frequent communication rounds on the master. Existing AXDA-based approaches unfortunately do not fulfill these important requirements (Vono et al., 2019; Rendell et al., 2021). In this work, we leverage these issues by considering the use of the Langevin Monte Carlo (LMC) algorithm to approximately sample from $\Pi_{\boldsymbol{\rho}}(\mathbf{z}_i | \boldsymbol{\theta})$ (Rossky et al., 1978; Roberts and Tweedie, 1996).

Our contributions are summarised in what follows. (1) We introduce in Section 2 a new methodology called Distributed Gibbs using Langevin Monte Carlo (DG-LMC). (2) Importantly, we provide in Section 3 a detailed quantitative analysis of the induced bias and show explicit convergence results. This stands for our main contribution and to the best of authors’ knowledge, this theoretical study is one of the most complete among existing works which focused on distributed Bayesian machine learning with a master/slaves architecture. In particular, we discuss the complexity of our algorithm, the choice of hyperparameters, and provide practitioners with simple prescriptions to tune them. Further, we provide a thorough comparison of our method with existing approaches in Section 4. (3) Finally, in Section 5, we show the benefits of the proposed sampler over popular and recent distributed MCMC algorithms on several numerical experiments. All the proofs are postponed to the Appendices.

Notations and conventions. The Euclidean norm on \mathbb{R}^d is denoted by $\|\cdot\|$. For $n \geq 1$, we refer to $\{1, \dots, n\}$ with the notation $[n]$ and for $i_1, i_2 \in \mathbb{N}$, $i_1 \leq i_2$, $\{i_1, \dots, i_2\}$ with the notation $[i_1 : i_2]$. For $0 \leq i < j$ and $(\mathbf{u}_k; k \in \{i, \dots, j\})$, we use the notation $\mathbf{u}_{i:j}$ to refer to the vector $[\mathbf{u}_i^\top, \dots, \mathbf{u}_j^\top]^\top$. We denote by $N(\mathbf{m}, \boldsymbol{\Sigma})$ the Gaussian distribution with mean vector \mathbf{m} and covariance matrix $\boldsymbol{\Sigma}$. For a given matrix $\mathbf{M} \in \mathbb{R}^{d \times d}$, we denote its smallest eigenvalue by $\lambda_{\min}(\mathbf{M})$. We denote by $\mathcal{B}(\mathbb{R}^d)$ the Borel σ -field of \mathbb{R}^d . We define the Wasserstein distance of order 2 for any probability measures μ, ν on \mathbb{R}^d with finite

2-moment by $W_2(\mu, \nu) = (\inf_{\zeta \in \mathcal{T}(\mu, \nu)} \int_{\mathbb{R}^d \times \mathbb{R}^d} \|\theta - \theta'\|^2 d\zeta(\theta, \theta'))^{1/2}$, where $\mathcal{T}(\mu, \nu)$ is the set of transference plans of μ and ν .

2 Distributed Gibbs using Langevin Monte Carlo (DG-LMC)

In this section, we present the proposed methodology which is based on the AXDA statistical framework and the popular LMC algorithm.

AXDA relies on the decomposition of the target distribution π given in (1) to introduce an extended distribution which enjoys favorable properties for distributed computations. This distribution is defined on the state space $\mathbb{R}^d \times \mathbf{Z}$, $\mathbf{Z} = \prod_{i=1}^b \mathbb{R}^{d_i}$, and admits a density w.r.t. the Lebesgue measure given, for any $\theta \in \mathbb{R}^d$, $\mathbf{z}_{1:b} \in \mathbf{Z}$, by

$$\Pi_{\rho}(\theta, \mathbf{z}_{1:b}) \propto \prod_{i=1}^b \tilde{\Pi}_{\rho}^i(\theta, \mathbf{z}_i), \quad (2)$$

where $\tilde{\Pi}_{\rho}^i(\theta, \mathbf{z}_i) = \exp(-U_i(\mathbf{z}_i) - \|\mathbf{z}_i - \mathbf{A}_i \theta\|^2 / 2\rho_i)$ and $\rho = \{\rho_i\}_{i=1}^b \in \mathbb{R}_+^b$ is a sequence of positive tolerance parameters. Note that $\tilde{\Pi}_{\rho}$ is not necessarily a probability density function. Actually, for Π_{ρ} to define a proper probability density, *i.e.* $\int_{\mathbb{R}^d \times \mathbf{Z}} \prod_{i=1}^b \tilde{\Pi}_{\rho}^i(\theta, \mathbf{z}_i) d\theta d\mathbf{z}_{1:b} < \infty$, some conditions are required.

H1. *There exists $b' \in [b-1]$ such that the following conditions hold: $\min_{i \in [b']} \inf_{\mathbf{z}_i \in \mathbb{R}^{d_i}} U_i(\mathbf{z}_i) > -\infty$, and $\max_{i \in [b'+1:b]} \int_{\mathbb{R}^{d_i}} e^{-U_i(\mathbf{z}_i)} d\mathbf{z}_i < \infty$. In addition, $\sum_{j=b'+1}^b \mathbf{A}_j^{\top} \mathbf{A}_j$ is invertible.*

The next result shows that these mild assumptions are sufficient to guarantee that the extended model (2) is well-defined.

Proposition 1. *Assume H1. Then, for any $\rho \in \mathbb{R}_+^b$, Π_{ρ} in (2) is a proper density.*

The data augmentation scheme (2) is approximate in the sense that the θ -marginal defined by

$$\pi_{\rho}(\theta) = \int_{\mathbf{Z}} \Pi_{\rho}(\theta, \mathbf{z}_{1:b}) d\mathbf{z}_{1:b}, \quad (3)$$

coincides with (1) only in the limiting case $\max_{i \in [b]} \rho_i \downarrow 0$ (Scheffé, 1947). For a fixed ρ , quantitative results on the induced bias in total variation distance can be found in Vono et al. (2019). The main benefit of working with (2) is that conditionally upon θ , auxiliary variables $\{\mathbf{z}_i\}_{i=1}^b$ are independent. Therefore, they can be sampled in parallel within a Gibbs sampler. For $i \in [b]$, the conditional density of \mathbf{z}_i given θ writes

$$\Pi_{\rho}(\mathbf{z}_i | \theta) \propto \exp\left(-U_i(\mathbf{z}_i) - \frac{\|\mathbf{z}_i - \mathbf{A}_i \theta\|^2}{2\rho_i}\right). \quad (4)$$

On the other hand, the conditional distribution of θ given $\mathbf{z}_{1:b}$ is a Gaussian distribution

$$\Pi_{\rho}(\theta | \mathbf{z}_{1:b}) = \mathcal{N}(\mu(\mathbf{z}_{1:b}), \mathbf{Q}^{-1}), \quad (5)$$

with precision matrix $\mathbf{Q} = \sum_{i=1}^b \mathbf{A}_i^\top \mathbf{A}_i / \rho_i$ and mean vector $\boldsymbol{\mu}(\mathbf{z}_{1:b}) = \mathbf{Q}^{-1} \sum_{i=1}^b \mathbf{A}_i^\top \mathbf{z}_i / \rho_i$. Under **H1**, note that \mathbf{Q} is invertible and therefore this conditional Gaussian distribution is well-defined. Since sampling from high-dimensional Gaussian distributions can be performed efficiently (Vono et al., 2020), this Gibbs sampling scheme is interesting as long as sampling from (4) is cheap. Vono et al. (2019) proposed the use of a rejection sampling step requiring to set $\rho_i = \mathcal{O}(1/d_i)$. When $d_i \gg 1$, this condition unfortunately leads to prohibitive computational costs and hence prevents its practical use for general Bayesian inference problems. Instead of sampling exactly from (4), Rendell et al. (2021) rather proposed to use Metropolis-Hastings algorithms. However, it is not clear whether this choice indeed leads to efficient sampling schemes. To tackle these issues, we propose to build upon LMC to end

Algorithm 1 Distributed Gibbs using LMC (DG-LMC)

Input: burn-in T_{bi} ; for $i \in [b]$, tolerance parameters $\rho_i > 0$, step-sizes $\gamma_i \in (0, \rho_i / (1 + \rho_i M_i)]$, local LMC steps $N_i \geq 1$.
 Initialise $\boldsymbol{\theta}^{(0)}$ and $\mathbf{z}_{1:b}^{(0)}$.
for $t = 0$ **to** $T - 1$ **do**
 // Sampling from $\Pi_{\boldsymbol{\rho}}(\mathbf{z}_{1:b} | \boldsymbol{\theta})$
 for $i = 1$ **to** b // In parallel on the b workers **do**
 $\mathbf{u}_i^{(0)} = \mathbf{z}_i^{(t)}$
 for $k = 0$ **to** $N_i - 1$ // N_i local LMC steps **do**
 $\boldsymbol{\xi}_i^{(k,t)} \sim \mathcal{N}(\mathbf{0}_{d_i}, \mathbf{I}_{d_i})$
 $\mathbf{g}_i = (1 - \frac{\gamma_i}{\rho_i}) \mathbf{u}_i^{(k)} + \frac{\gamma_i}{\rho_i} \mathbf{A}_i \boldsymbol{\theta}^{(t)} - \gamma_i \nabla U_i(\mathbf{u}_i^{(k)})$
 $\mathbf{u}_i^{(k+1)} = \mathbf{g}_i + \sqrt{2\gamma_i} \boldsymbol{\xi}_i^{(k,t)}$ // See (4)
 end for
 $\mathbf{z}_i^{(t+1)} = \mathbf{u}_i^{(N_i)}$
 end for
 // Sampling from $\Pi_{\boldsymbol{\rho}}(\boldsymbol{\theta} | \mathbf{z}_{1:b})$
 $\boldsymbol{\theta}^{(t+1)} \sim \mathcal{N}(\boldsymbol{\mu}(\mathbf{z}_{1:b}^{(t+1)}), \mathbf{Q}^{-1})$ // See (5)
end for
Output: samples $\{\boldsymbol{\theta}^{(t)}\}_{t=T_{\text{bi}}-1}^T$.

up with a distributed MCMC algorithm which is both simple to implement, efficient and amenable to a theoretical study. LMC stands for a popular way to approximately generate samples from a given distribution based on the Euler-Maruyama discretisation scheme of the overdamped Langevin stochastic differential equation (Roberts and Tweedie, 1996). At iteration t of the considered Gibbs sampling scheme and given a current parameter $\boldsymbol{\theta}^{(t)}$, LMC applied to (4) considers, for $i \in [b]$, the recursion

$$\mathbf{z}_i^{(t+1)} = (1 - \frac{\gamma_i}{\rho_i}) \mathbf{z}_i^{(t)} + \frac{\gamma_i}{\rho_i} \mathbf{A}_i \boldsymbol{\theta}^{(t)} - \gamma_i \nabla U_i(\mathbf{z}_i^{(t)}) + \sqrt{2\gamma_i} \boldsymbol{\xi}_i^{(t)}$$




Figure 1: Illustration of one global iteration of Algorithm 1. For each worker, the width of the green box represents the amount of time required to perform one LMC step.

where $\gamma_i > 0$ is a fixed step-size and $(\xi_i^{(k)})_{k \in \mathbb{N}, i \in [b]}$ a sequence of independent and identically distributed (i.i.d.) d -dimensional standard Gaussian random variables. Only using a single step of LMC on each worker might incur important communication costs. To mitigate the latter while increasing the proportion of time spent on exploring the state-space, we instead allow each worker to perform $N_i \geq 1$ LMC steps (Dieuleveut and Patel, 2019; Rendell et al., 2021). Letting N_i varies across workers prevents Algorithm 1 to suffer from a significant block-by-the-slowest delay in cases where the response times of the workers are unbalanced (Ahn et al., 2014). The proposed algorithm, coined Distributed Gibbs using Langevin Monte Carlo (DG-LMC), is depicted in Algorithm 1 and illustrated in Figure 1.

3 Detailed analysis of DG-LMC

In this section, we derive quantitative bias and convergence results for DG-LMC and show that its mixing time only scales quadratically w.r.t. the dimension d . We also discuss the choice of hyperparameters and provide guidelines to tune them.

3.1 Non-Asymptotic Analysis

The scope of our analysis will focus on smooth and strongly log-concave target posterior distributions π . While these assumptions may be restrictive in practice, they allow for a detailed theoretical study of the proposed algorithm.

- H2.** (i) For any $i \in [b]$, U_i is twice continuously differentiable and $\sup_{\mathbf{z}_i \in \mathbb{R}^{d_i}} \|\nabla^2 U_i(\mathbf{z}_i)\| \leq M_i$.
(ii) For any $i \in [b]$, U_i is m_i -strongly convex: there exists $m_i > 0$ such that $m_i \mathbf{I}_{d_i} \preceq \nabla^2 U_i$.

Under these assumptions, it is shown in Lemma S16 in the Appendix that $-\log \pi$ is strongly convex with constant

$$m_U = \lambda_{\min}(\sum_{i=1}^b m_i \mathbf{A}_i^\top \mathbf{A}_i) . \quad (6)$$

Behind the use of LMC, the main motivation is to end up with a simple hybrid Gibbs sampler amenable to a non-asymptotic theoretical analysis based on previous works (Durmus and Moulines, 2019; Dalalyan and Karagulyan, 2019). In the following, this study is carried out using the Wasserstein distance of order 2.

3.1.1 Convergence Results

DG-LMC introduced in Algorithm 1 defines a homogeneous Markov chain $(V_t)_{t \in \mathbb{N}} = (\theta_t, Z_t)_{t \in \mathbb{N}}$ with realisations $(\theta^{(t)}, \mathbf{z}_{1:b}^{(t)})_{t \in \mathbb{N}}$. We denote by $P_{\rho, \gamma, \mathbf{N}}$ the Markov kernel associated with $(V_t)_{t \in \mathbb{N}}$. Since no Metropolis-Hastings step is used in combination with LMC, the proposed algorithm does not fall into the class of Metropolis-within-Gibbs samplers (Roberts and Rosenthal, 2006). Therefore, a first step is to show that $P_{\rho, \gamma, \mathbf{N}}$ admits a unique invariant distribution and is geometrically ergodic. We proceed via an appropriate synchronous coupling which reduces the convergence analysis of $(V_t)_{t \in \mathbb{N}}$ to that of the marginal process $(Z_t)_{t \in \mathbb{N}}$. While the proof of the convergence of $(Z_t)_{t \in \mathbb{N}}$ shares some similarities with LMC (Durmus and Moulines, 2019), the analysis of $(Z_t)_{t \in \mathbb{N}}$ is much more involved and especially in the case $\max_{i \in [b]} N_i > 1$. We believe that the proof techniques we developed to show the next result can be useful to the study of other MCMC approaches based on LMC.

Proposition 2. *Assume **H1-H2** and let $c > 0$ and $\gamma = \{\gamma_i\}_{i=1}^b$ $\mathbf{N} = \{N_i\}_{i=1}^b$ satisfying $\max_{i \in [b]} \gamma_i \leq \bar{\gamma}$, $\min_{i \in [b]} \{N_i \gamma_i\} / \max_{i \in [b]} \{N_i \gamma_i\} \geq c$ and $\max_{i \in [b]} \{N_i \gamma_i\} \leq C_1$ where $\bar{\gamma}, C_1$ are explicit constants only depending on $(m_i, M_i, \rho_i)_{i \in [b]}$ ¹². Then, there exists a probability measure $\Pi_{\rho, \gamma, \mathbf{N}}$ such that $\Pi_{\rho, \gamma, \mathbf{N}}$ is invariant for $P_{\rho, \gamma, \mathbf{N}}$. Moreover there exists $C_2 > 0$ such that for any integer $t \geq 0$ and $\mathbf{v} = (\theta, \mathbf{z}) \in \mathbb{R}^d \times \mathcal{Z}$, we have*

$$W_2(\delta_{\mathbf{v}} P_{\rho, \gamma, \mathbf{N}}^t, \Pi_{\rho, \gamma, \mathbf{N}}) \leq C_2 \cdot (1 - \min_{i \in [b]} \{N_i \gamma_i m_i\} / 2)^t \cdot W_2(\delta_{\mathbf{v}}, \Pi_{\rho, \gamma, \mathbf{N}}).$$

Explicit expressions for C_1 and C_2 are given in Proposition S13 in the Appendix. Finally, if $\mathbf{N} = N \mathbf{1}_b$ for $N \geq 1$, then $\Pi_{\rho, \gamma, \mathbf{N}} = \Pi_{\rho, \gamma, \mathbf{1}_b}$.

We now discuss Proposition 2. If we set, for any $i \in [b]$, $N_i = 1$, the convergence rate in Proposition 2 becomes equal to $1 - \min_{i \in [b]} \{\gamma_i m_i\} / 2$. In this specific case, we show in Proposition S5 in the Appendix that DG-LMC actually admits the tighter convergence rate $1 - \min_{i \in [b]} \{\gamma_i m_i\}$ which simply corresponds to the rate at which the slowest LMC conditional kernel converges. On the other hand, when $\max_{i \in [b]} N_i > 1$, the convergence of $P_{\rho, \gamma, \mathbf{N}}$ towards $\Pi_{\rho, \gamma, \mathbf{N}}$ only holds if $\max_{i \in [b]} \{N_i \gamma_i\}$ is sufficiently small. This condition is necessary to ensure a contraction in W_2 and can be understood intuitively as follows in the case where $\mathbf{N} = N \mathbf{1}_b$ and $\gamma = \gamma \mathbf{1}_b$. Given two vectors (θ_k, θ'_k) and an appropriate coupling (Z_{k+1}, Z'_{k+1}) , we can show that $Z_{k+1} - Z'_{k+1}$ involves two competing terms: one keeping

¹When $\mathbf{N} = \mathbf{1}_b$, $C_1 = \bar{\gamma} = 1 / \max_{i \in [b]} \{M_i + \rho_i^{-1}\}$.

²When $\max_{i \in [b]} N_i > 1$, C_1 is of order $\min_{i \in [b]} \rho_i^2$ when $\max_{i \in [b]} \rho_i \rightarrow 0$, see Lemma S12 in the Appendix.

$Z_{k+1} - Z'_{k+1}$ close to $Z_k - Z'_k$ and another one driving $Z_{k+1} - Z'_{k+1}$ away from $\theta_k - \theta'_k$ (and therefore of $Z_k - Z'_k$) as N increases. This implies that N stands for a trade-off and the product $N\gamma$ cannot be arbitrarily chosen. Finally, it is worth mentioning that the tolerance parameters $\{\rho_i\}_{i \in [b]}$ implicitly drive the convergence rate of DG-LMC. In the case $N_i = 1$, a sufficient condition on the step-sizes to ensure a contraction is $\gamma_i \leq 2/(M_i + m_i + 1/\rho_i)$. We can denote that the smaller ρ_i , the smaller γ_i and the slower the convergence.

Starting from the results of Proposition 2, we can analyse the convergence properties of DG-LMC. We specify our result to the case where we take for the specific initial distribution

$$\mu_{\boldsymbol{\rho}}^{\star} = \delta_{\mathbf{z}^{\star}} \otimes \Pi_{\boldsymbol{\rho}}(\cdot | \mathbf{z}^{\star}) , \quad (7)$$

where $\mathbf{z}^{\star} = ([\mathbf{A}_1 \boldsymbol{\theta}^{\star}]^{\top}, \dots, [\mathbf{A}_b \boldsymbol{\theta}^{\star}]^{\top})^{\top}$, $\boldsymbol{\theta}^{\star} = \arg \min \{-\log \pi\}$ and $\Pi_{\boldsymbol{\rho}}(\cdot | \mathbf{z}^{\star})$ is defined in (5). Note that sampling from $\mu_{\boldsymbol{\rho}}^{\star}$ is straightforward and simply consists in setting $\mathbf{z}^{(0)} = \mathbf{z}^{\star}$ and drawing $\boldsymbol{\theta}^{(0)}$ from $\Pi_{\boldsymbol{\rho}}(\cdot | \mathbf{z}^{\star})$. For $t \geq 1$, we consider the marginal law of θ_t initialised at \mathbf{v}^{\star} with distribution $\mu_{\boldsymbol{\rho}}^{\star}$ and denote it $\Gamma_{\mathbf{v}^{\star}}^t$. As mentioned previously, the proposed approach relies on two approximations which both come with some bias we need to control. This naturally brings us to consider the following inequality based on the triangular inequality and the definition of the Wasserstein distance:

$$W_2(\Gamma_{\mathbf{v}^{\star}}^t, \pi) \leq W_2(\mu_{\boldsymbol{\rho}}^{\star} P_{\boldsymbol{\rho}, \gamma, N}^t, \Pi_{\boldsymbol{\rho}, \gamma, N}) + W_2(\Pi_{\boldsymbol{\rho}, \gamma, N}, \Pi_{\boldsymbol{\rho}}) + W_2(\pi_{\boldsymbol{\rho}}, \pi) , \quad (8)$$

where $\Pi_{\boldsymbol{\rho}, \gamma, N}$, $\Pi_{\boldsymbol{\rho}}$ and $\pi_{\boldsymbol{\rho}}$ are defined in Proposition 2, (2) and (3), respectively. In Proposition S14 in the Appendix, we provide an upper bound on the first term on the right hand side based on Proposition 2. In the next section, we focus on controlling the last two terms on the right hand side.

3.1.2 Quantitative Bounds on the Bias

The error term $W_2(\pi_{\boldsymbol{\rho}}, \pi)$ in (8) is related to the underlying AXDA framework which induces an approximate posterior representation $\pi_{\boldsymbol{\rho}}$. It can be controlled by the sequence of positive tolerance parameters $\{\rho_i\}_{i=1}^b$. By denoting $\bar{\rho} = \max_{i \in [b]} \rho_i$, Proposition 3 shows that this error can be quantitatively assessed and is of order $\mathcal{O}(\bar{\rho})$ for sufficiently small values of this parameter.

Proposition 3. *Assume H 1, H 2. In addition, let $\mathbf{A} = [\mathbf{A}_1^{\top}, \dots, \mathbf{A}_b^{\top}]^{\top}$ and denote $\sigma_U^2 = \|\mathbf{A}^{\top} \mathbf{A}\| \max_{i \in [b]} \{M_i^2\} / m_U$, where m_U is defined in (6). Then, for any $\bar{\rho} \leq \sigma_U^2 / 12$,*

$$W_2(\pi_{\boldsymbol{\rho}}, \pi) \leq \sqrt{2/m_U} \max(A_{\boldsymbol{\rho}}, B_{\boldsymbol{\rho}}) ,$$

where $A_{\boldsymbol{\rho}} = d\mathcal{O}(\bar{\rho})$ and $B_{\boldsymbol{\rho}} = d^{1/2}\mathcal{O}(\bar{\rho})$ for $\bar{\rho} \downarrow 0$. Explicit expressions for $A_{\boldsymbol{\rho}}, B_{\boldsymbol{\rho}}$ are given in Appendix S3 in the Appendix.

Table 1: For the specific initialisation \mathbf{v}^* with distribution μ_{ρ}^* given in (7), dependencies w.r.t. d and ε of the parameters involved in Algorithm 1 and of $t_{\text{mix}}(\varepsilon; \mathbf{v}^*)$ to get a W_2 -error of at most ε .

Assumptions		ρ_{ε}	γ_{ε}	N_{ε}	$t_{\text{mix}}(\varepsilon; \mathbf{v}^*)$	Gradient evaluations
H1, H2	d	$\mathcal{O}(d^{-1})$	$\mathcal{O}(d^{-3})$	$\mathcal{O}(d)$	$\mathcal{O}(d^2 \log(d))$	$\mathcal{O}(d^3 \log(d))$
	ε	$\mathcal{O}(\varepsilon)$	$\mathcal{O}(\varepsilon^4)$	$\mathcal{O}(\varepsilon^{-2})$	$\mathcal{O}(\varepsilon^{-2} \log(\varepsilon))$	$\mathcal{O}(\varepsilon^{-4} \log(\varepsilon))$
H1, H2, H3	d	$\mathcal{O}(d^{-1})$	$\mathcal{O}(d^{-2})$	$\mathcal{O}(1)$	$\mathcal{O}(d^2 \log(d))$	$\mathcal{O}(d^2 \log(d))$
	ε	$\mathcal{O}(\varepsilon)$	$\mathcal{O}(\varepsilon^2)$	$\mathcal{O}(1)$	$\mathcal{O}(\varepsilon^{-2} \log(\varepsilon))$	$\mathcal{O}(\varepsilon^{-2} \log(\varepsilon))$

In the case where π is Gaussian, the approximate distribution π_{ρ} admits an explicit expression and is Gaussian as well (e.g. when $b = 1$, the mean is the same and the covariance matrix is inflated by a factor $\rho \mathbf{I}_d$), see for instance [Rendell et al. \(2021, Section S2\)](#) and [Vono et al. \(2021, Section 5.1\)](#). Hence, an explicit expression for $W_2(\pi_{\rho}, \pi)$ can be derived. Based on this result, we can check that the upper bound provided by Proposition 3 matches the same asymptotics as $\rho \rightarrow 0$ and $d \rightarrow \infty$.

The second source of approximation error is induced by the use of LMC within Algorithm 1 to target the conditional distribution $\Pi_{\rho}(\mathbf{z}_{1:b} | \boldsymbol{\theta})$ in (4). The stationary distribution of $P_{\rho, \gamma, \mathbf{N}}$ whose existence is ensured in Proposition 2 differs from Π_{ρ} . The associated bias is assessed quantitatively in Proposition 4.

Proposition 4. *Assume **H1-H2**. For any $i \in [b]$, define $\tilde{M}_i = M_i + 1/\rho_i$ and let $\gamma \in (\mathbb{R}_+^*)^b$, $\mathbf{N} \in (\mathbb{N}^*)^b$ such that for any $i \in [b]$,*

$$\gamma_i \leq \frac{m_i}{40\tilde{M}_i^2} \min_{i \in [b]} (m_i/\tilde{M}_i)^2 / \max_{i \in [b]} (m_i/\tilde{M}_i)^2, \quad (9)$$

$$N_i = \lfloor m_i \min_{i \in [b]} \{m_i/\tilde{M}_i\}^2 / (20\gamma_i \tilde{M}_i^2 \max_{i \in [b]} \{m_i/\tilde{M}_i\}^2) \rfloor. \quad (10)$$

Then, we have

$$W_2^2(\Pi_{\rho, \gamma, \mathbf{N}}, \Pi_{\rho}) \leq C_3 \sum_{i=1}^b d_i \gamma_i \tilde{M}_i^2,$$

where $C_3 > 0$ only depends of $(m_i, M_i, \mathbf{A}_i, \rho_i)_{i=1}^b$ and is explicitly given in Proposition S29 in the Appendix.

With the notation $\bar{\gamma} = \max_{i \in [b]} \gamma_i$, Proposition 4 implies that $W_2(\Pi_{\rho}, \Pi_{\rho, \gamma, \mathbf{N}}) \leq \mathcal{O}(\bar{\gamma}^{1/2})(\sum_{i=1}^b d_i)^{1/2}$ for $\bar{\gamma} \downarrow 0$. Note that this result is in line with [Durmus and Moulines](#)

(2019, Corollary 7) and can be improved under further regularity assumptions on U , as shown below.

H3. U is three times continuously differentiable and there exists $L_i > 0$ such that for all $\mathbf{z}_i, \mathbf{z}'_i \in \mathbb{R}^{d_i}$, $\|\nabla^2 U_i(\mathbf{z}_i) - \nabla^2 U_i(\mathbf{z}'_i)\| \leq L_i \|\mathbf{z}_i - \mathbf{z}'_i\|$.

Proposition 5. Assume **H1-H2-H3**. For any $i \in [b]$, define $\tilde{M}_i = M_i + 1/\rho_i$ and let $\gamma \in (\mathbb{R}_+^*)^b$, $\mathbf{N} \in (\mathbb{N}^*)^b$ such that for any $i \in [b]$, (9) and (10) hold. Then, we have

$$W_2^2(\Pi_{\rho, \gamma, \mathbf{N}}, \Pi_{\rho}) \leq C_4 \sum_{i \in [b]} d_i \gamma_i (1/\tilde{M}_i^2 + \gamma_i \tilde{M}_i^2),$$

where $C_4 > 0$ only depends on $(m_i, M_i, L_i, \mathbf{A}_i, \rho_i)_{i=1}^b$ and is explicitly given in Proposition S33 in the Appendix.

3.1.3 Mixing Time with Explicit Dependencies

Based on explicit non-asymptotic bounds shown in Propositions 2, 3 and 4 and the decomposition (8), we are now able to analyse the scaling of Algorithm 1 in high dimension. Given a prescribed precision $\varepsilon > 0$ and an initial condition \mathbf{v}^* with distribution μ_{ρ}^* given in (7), we define the ε -mixing time associated to $\Gamma_{\mathbf{v}^*}$ by

$$t_{\text{mix}}(\varepsilon; \mathbf{v}^*) = \min \{t \in \mathbb{N} : W_2(\Gamma_{\mathbf{v}^*}^t, \pi) \leq \varepsilon\}.$$

This quantity stands for the minimum number of DG-LMC iterations such that the θ -marginal distribution is at most at an ε W_2 -distance from the initial target π . Under the condition that $b \max_{i \in [b]} d_i = \mathcal{O}(d)$ and by assuming for simplicity that for any $i \in [b]$, $m_i = m, M_i = M, L_i = L, \rho_i = \rho, \gamma_i = \gamma$ and $N_i = N$, Table 1 gathers the dependencies w.r.t. d and ε of the parameters involved in Algorithm 1 and of $t_{\text{mix}}(\varepsilon; \mathbf{v}^*)$ to get a W_2 -error of at most ε . Note that the mixing time of Algorithm 1 scales at most quadratically (up to polylogarithmic factors) in the dimension. When **H3** holds, we can see that the number of local iterations becomes independent of d and ε which leads to a total number of gradient evaluations with better dependencies w.r.t. to these quantities. Up to the authors' knowledge, these explicit results are the first among the centralised distributed MCMC literature and in particular give the dependency w.r.t. d and ε of the number of local LMC iterations on each worker. Overall, the proposed approach appears as a scalable and reliable alternative for high-dimensional and distributed Bayesian inference.

3.2 DG-LMC in Practice: Guidelines for Practitioners

We now discuss practical guidelines for setting the values of hyperparameters involved in Algorithm 1. Based on Proposition 2, we theoretically show an optimal choice of order $N_i \gamma_i \asymp m_i \rho_i^2 / (\rho_i M_i + 1)^2$. Ideally, within the considered distributed setting, the

optimal value for $(N_i, \gamma_i)_{i \in [b]}$ would boil down to optimise the value of $\max_{i \in [b]} \{N_i \gamma_i\}$ under the constraints derived in Proposition 2 combined with communication considerations. In particular, this would imply a comprehensive modelling of the communication costs including I/O bandwidths constraints. These optimisation tasks fall outside the scope of the present paper and therefore we let the search of optimal values for future works. Since our aim here is to provide practitioners with simple prescriptions, we rather focus on general rules involving tractable quantities.

3.2.1 Selection of γ and ρ

From Durmus and Moulines (2017) and references therein, a simple sufficient condition on step-sizes $\gamma = \{\gamma_i\}_{i=1}^b$ to guarantee the stability of LMC is $\gamma_i \leq \rho_i / (\rho_i M_i + 1)$ for $i \in [b]$. Both the values of γ_i and ρ_i are subject to a bias-variance trade-off. More precisely, large values yield a Markov chain with small estimation variance but high asymptotic bias. Conversely, small values produce a Markov chain with small asymptotic bias but which requires a large number of iterations to obtain a stable estimator. We propose to mitigate this trade-off by setting γ_i to a reasonably large value, that is for $i \in [b]$, $\gamma_i \in [0.1\rho_i / (\rho_i M_i + 1), 0.5\rho_i / (\rho_i M_i + 1)]$. Since γ_i saturates to $1/M_i$ when $\rho_i \rightarrow \infty$, there is no computational advantage to choose very large values for ρ_i . Based on several numerical studies, we found that setting ρ_i of the order of $1/M_i$ was a good compromise between computational efficiency and asymptotic bias.

3.2.2 N : A Trade-Off between Asymptotic Bias and Communication Overhead

In a similar vein, the choice of $N = \{N_i\}_{i=1}^b$ also stands for a trade-off but here between asymptotic accuracy and communication costs. Indeed, a large number of local LMC iterations reduces the communication overhead but at the expense of a larger asymptotic bias since the master parameter is not updated enough. Ahn et al. (2014) proposed to tune the number of local iterations N_i on a given worker based on the amount of time needed to perform one local iteration, denoted here by τ_i . Given an average number of local iterations N_{avg} , the authors set $N_i = q_i N_{\text{avg}} b$ with $q_i = \tau_i^{-1} / \sum_{k=1}^b \tau_k^{-1}$ so that $b^{-1} \sum_{i=1}^b N_i = N_{\text{avg}}$. As mentioned by the aforementioned authors, this choice allows to keep the block-by-the-slowest delay small by letting fast workers perform more iterations in the same wall-clock time. Although they showed how to tune N_i w.r.t. communication considerations, they let the choice of N_{avg} to the practitioner. Here, we propose a simple guideline to set N_{avg} such that N_i stands for a good compromise between the amount of time spent on exploring the state-space and communication overhead. As highlighted in the discussion after Proposition 2, as γ_i becomes smaller, more local LMC iterations are required to sufficiently explore the latent space before the global consensus round on the master. Assuming for any $i \in [b]$ that γ_i has been chosen following our guidelines in

Table 2: Synthetic overview of the main existing distributed MCMC methods under a master-slave architecture. The column *Exact* means that the Markov chain defined by the MCMC sampler admits (1) as invariant distribution. The column *Comm.* reports the communication frequency. A value of 1 means that the sampler communicates after every iteration. T stands for the total number of iterations and $N < T$ is a tunable parameter to mitigate communication costs. The acronym D-SGLD stands for distributed stochastic gradient Langevin dynamics.

METHOD	TYPE	EXACT	COMM.	BIAS BOUNDS	SCALING
WANG AND DUNSON (2013)	ONE-SHOT	×	$1/T$	✓	$\mathcal{O}(e^d)$
NEISWANGER ET AL. (2014)	ONE-SHOT	×	$1/T$	×	$\mathcal{O}(e^d)$
MINSKER ET AL. (2014)	ONE-SHOT	×	$1/T$	✓	UNKNOWN
SRIVASTAVA ET AL. (2015)	ONE-SHOT	×	$1/T$	×	UNKNOWN
WANG ET AL. (2015)	ONE-SHOT	×	$1/T$	✓	$\mathcal{O}(e^d)$
SCOTT ET AL. (2016)	ONE-SHOT	×	$1/T$	×	UNKNOWN
NEMETH AND SHERLOCK (2018)	ONE-SHOT	×	$1/T$	×	UNKNOWN
JORDAN ET AL. (2019)	ONE-SHOT	×	$1/T$	✓	UNKNOWN
AHN ET AL. (2014)	D-SGLD	×	$1/N$	×	UNKNOWN
CHEN ET AL. (2016)	D-SGLD	×	1	✓	UNKNOWN
EL MEKKAoui ET AL. (2020)	D-SGLD	×	$1/N$	✓	UNKNOWN
RABINOVICH ET AL. (2015)	G. CONSENSUS	×	$1/N$	×	UNKNOWN
CHOWDHURY AND JERMAINE (2018)	G. CONSENSUS	✓	1	N/A	UNKNOWN
RENDELL ET AL. (2021)	G. CONSENSUS	×	$1/N$	✓	UNKNOWN
THIS PAPER	G. CONSENSUS	×	$1/N$	✓	$\mathcal{O}(d^2 \log(d))$

Section 3.2.1, this suggests to set $N_{\text{avg}} = \lceil (1/b) \sum_{i \in [b]} \rho_i / (\gamma_i [\rho_i M_i + 1]) \rceil$.

4 Related work

As already mentioned in Section 1, hosts of contributions have focused on deriving distributed MCMC algorithms to sample from (1). This section briefly reviews the main existing research lines and draws a detailed comparison with the proposed methodology.

4.1 Existing distributed MCMC methods

Existing methodologies are mostly approximate and can be loosely speaking divided into three groups: *one-shot*, distributed stochastic gradient MCMC and *global consensus* approaches. To ease the understanding, a synthetic overview of their main characteristics is presented in Table 2.

One-shot approaches stand for communication-efficient schemes where workers and master only exchange information at the very beginning and the end of the sampling task;

similarly to MapReduce schemes (Dean and Ghemawat, 2004). Most of these methods assume that the posterior density factorises into a product of local posteriors and launch independent Markov chains across workers to target them. The local posterior samples are then combined through the master node using a single final aggregation step. This step turns to be the milestone of one-shot approaches and was the topic of multiple contributions (Wang and Dunson, 2013; Neiswanger et al., 2014; Minsker et al., 2014; Srivastava et al., 2015; Scott et al., 2016; Nemeth and Sherlock, 2018). Unfortunately, the latter are either infeasible in high-dimensional settings or have been shown to yield inaccurate posterior representations empirically, if the posterior is not near-Gaussian, or if the local posteriors differ significantly (Wang et al., 2015; Dai et al., 2019; Rendell et al., 2021). Alternative schemes have been recently proposed to tackle these issues but their theoretical scaling w.r.t. the dimension d is currently unknown (Jordan et al., 2019; Mesquita et al., 2020).

Albeit popular in the machine learning community, distributed stochastic gradient MCMC methods (Ahn et al., 2014) suffer from high variance when the dataset is large because of the use of stochastic gradients (Brosse et al., 2018). Some surrogates have been recently proposed to reduce this variance such as the use of *stale* or *conductive* gradients (Chen et al., 2016; El Mekkaoui et al., 2020). However, these variance reduction methods require an increasing number of workers for the former and come at the price of a prohibitive pre-processing step for the latter. In addition, it is currently unclear whether these methods are able to generate efficiently accurate samples from a given target distribution.

Contrary to aforementioned distributed MCMC approaches, global consensus methods periodically share information between workers by performing a consensus round between the master and the workers (Rabinovich et al., 2015; Chowdhury and Jermaine, 2018; Vono et al., 2019; Rendell et al., 2021). Again, they have been shown to perform well in practice but their theoretical understanding is currently limited.

4.2 Comparison with the proposed methodology

Table 2 compares Algorithm 1 with existing approaches detailed previously. In addition to having a simple implementation and guidelines, it is worth noticing that DG-LMC appears to benefit from favorable convergence properties compared to the other considered methodologies.

We complement this comparison with an informal discussion on the computational and communication complexities of Algorithm 1. Recall that the dataset is assumed to be partitioned into S shards and stored on S workers among a collection of b computing nodes. Suppose that the s -th shard has size n_s , and let T be the number of total MCMC iterations and c_{com} the communication cost. In addition, denote by $c_{\text{eval}}^{(i)}$ the approximate wall-clock time required to evaluate U_i or its gradient. For the ease of exposition, we do not discuss the additional overhead due to bandwidth restrictions and assume similar computation costs, *i.e.*, $Nc_{\text{eval}} \simeq N_i c_{\text{eval}}^{(i)}$, to perform each local LMC step at each iteration of Algorithm 1. Under these assumptions, the total complexity of Algorithm 1 is $\mathcal{O}(T[2c_{\text{com}} + Nc_{\text{eval}}])$.

Following the same reasoning, distributed stochastic gradient Langevin dynamics (D-SGLD) and one-shot approaches admit complexities of the order $\mathcal{O}(T[2c_{\text{com}} + Nc_{\text{eval}}n_{\text{mb}}/n_s])$ and $\mathcal{O}(Tc_{\text{eval}} + 2c_{\text{com}})$, respectively. The integer n_{mb} stands for the mini-batch size used in D-SGLD. Despite their very low communication overhead, existing one-shot approaches are rarely reliable and therefore not necessarily efficient to sample from π given a prescribed computational budget, see [Rendell et al. \(2021\)](#) for a recent overview. D-SGLD seems to enjoy a lower complexity than Algorithm 1 when n_{mb} is small. Unfortunately, this choice comes with two main shortcomings: (i) a larger number of iterations T to achieve the same precision because of higher variance of gradient estimators, and (ii) a smaller amount of time spent on exploration compared to communication latency. By falling into the global consensus class of methods, the proposed methodology hence appears as a good compromise between one-shot and D-SGLD algorithms in terms of both computational complexity and accuracy. Section 5 will enhance the benefits of Algorithm 1 by showing experimentally better convergence properties and posterior approximation.

5 Experiments

This section compares numerically DG-LMC with the most popular and recent centralised distributed MCMC approaches namely D-SGLD and the global consensus Monte Carlo (GCMC) algorithm proposed in [Rendell et al. \(2021\)](#). Since all these approaches share the same communication latency, this feature is not discussed here.

5.1 Toy Gaussian Example

In this toy example, we first illustrate the behavior of DG-LMC w.r.t. the number of local iterations which drives the communication overhead. We consider the conjugate Gaussian model $\pi(\boldsymbol{\theta}|\mathbf{y}_{1:n}) \propto \mathcal{N}(\boldsymbol{\theta}|\mathbf{0}_d, \boldsymbol{\Sigma}_0) \prod_{i=1}^n \mathcal{N}(\mathbf{y}_i|\boldsymbol{\theta}, \boldsymbol{\Sigma}_1)$, with positive definite matrices $\boldsymbol{\Sigma}_0, \boldsymbol{\Sigma}_1$. We set $d = 2$, allocate $n = 20,000$ observations to a cluster made of $b = 10$ workers and compare DG-LMC with D-SGLD. Both MCMC algorithms have been run using the same number of local iterations N per worker and for a fixed budget of $T = 100,000$ iterations including a burn-in period equal to $T_{\text{bi}} = 10,000$. Regarding DG-LMC, we follow the guidelines in Section 3.2.1 and set for all $i \in [b]$, $\mathbf{A}_i = \mathbf{I}_d$, $\rho_i = 1/(5M_i)$ and $\gamma_i = 0.25\rho_i/(\rho_iM_i + 1)$. On the other hand, D-SGLD has been run with batch-size $n/(10b)$ and a step-size chosen such that the resulting posterior approximation is similar to that of DG-LMC for $N = 1$. Figure 2 depicts the results for $N = 1$ and $N = 10$ on the left and right columns, respectively. The top row (resp. middle row) shows the contours of the b local posteriors in dashed grey, the contours of the target posterior in red and the 2D histogram built with DG-LMC (resp. D-SGLD) samples in blue (resp. green). When required, a zoomed version of these figures is depicted at the top right corner. It can be noted that DG-LMC exhibits better mixing properties while achieving similar performances as shown by the autocorrelation function (ACF) on the bottom row.




Figure 2: Toy Gaussian experiment. (left) $N = 1$ local iterations and (right) $N = 10$. (top) DG-LMC, (middle) D-SGLD and (bottom) ACF comparison between DG-LMC and D-SGLD.

Furthermore, its posterior approximation is robust to the choice of N in contrast to D-SGLD, which needs further tuning of its step-size to yield an accurate posterior representation. This feature is particularly important for distributed computations since N is directly related to communication costs and might often change depending upon the hardware architecture.

5.2 Bayesian Logistic Regression

This second experiment considers a more challenging problem namely Bayesian logistic regression. We use the *covtype*³ dataset with $d = 54$ and containing $n = 581,012$ observations partitioned into $b = 16$ shards. We set $N = 10$, $T = 200,000$, $T_{bi} = T/10$ for all approaches, and again used the guidelines in Section 3.2.1 to tune DG-LMC. Under the Bayesian paradigm, we are interested in performing uncertainty quantification by estimating highest

³www.csie.ntu.edu.tw/~cjlin/libsvmtools/datasets




Figure 3: Logistic regression. From left to right: negative log-posterior, ACF, HPD relative error after and during the sampling procedure.




Figure 4: Bayesian neural network. (left) probability of the most probable label for 8 examples and (right) probability of each label for a single example.

posterior density (HPD) regions. For any $\alpha \in (0, 1)$, define $\mathcal{C}_\alpha = \{\boldsymbol{\theta} \in \mathbb{R}^d; -\log \pi(\boldsymbol{\theta}|\mathbf{y}_{1:n}) \leq \eta_\alpha\}$ where $\eta_\alpha \in \mathbb{R}$ is chosen such that $\int_{\mathcal{C}_\alpha} \pi(\boldsymbol{\theta}|\mathbf{y}_{1:n}) d\boldsymbol{\theta} = 1 - \alpha$. For the three approximate MCMC approaches, we computed the relative HPD error based on the scalar summary η_α , *i.e.* $|\eta_\alpha - \eta_\alpha^{\text{true}}|/\eta_\alpha^{\text{true}}$ where $\eta_\alpha^{\text{true}}$ has been estimated using the Metropolis adjusted Langevin algorithm. The parameters of GCMC and D-SGLD have been chosen such that all MCMC algorithms achieve similar HPD error. Figure 3 shows that this error is reasonable and of the order of 1%. Nonetheless, one can denote that DG-LMC achieves this precision level faster than GCMC and D-SGLD due to better mixing properties. This confirms that the proposed methodology is indeed efficient and reliable to perform Bayesian analyses compared to existing popular methodologies.

5.3 Bayesian Neural Network

Up to now, both our theoretical and experimental results focused on the strongly log-concave scenario and showed that even in this case, DG-LMC appeared as a competitive alternative. In this last experiment, we propose to end the study of DG-LMC on an open note without ground truth by tackling the challenging sampling problem associated to Bayesian neural networks. We consider the MNIST training dataset consisting of $n = 60,000$ observations partitioned into $b = 50$ shards and such that for any $i \in [n]$ and $k \in [10]$, $\mathbb{P}(y_i = k|\boldsymbol{\theta}, \mathbf{x}_i) = \beta_k$ where β_k is the k -th element of $\sigma(\sigma(\mathbf{x}_i^\top \mathbf{W}_1 + \mathbf{b}_1)\mathbf{W}_2 + \mathbf{b}_2)$, $\sigma(\cdot)$ is the sigmoid function, \mathbf{x}_i are covariates, and \mathbf{W}_1 , \mathbf{W}_2 , \mathbf{b}_1 and \mathbf{b}_2 are matrices of size 784×128 , 128×10 , 1×128 and 1×10 , respectively. We set normal priors for each weight matrix and bias vector, $N = 10$ and ran DG-LMC with constant hyperparameters across workers $(\rho, \gamma) = (0.02, 0.005)$ and D-SGLD using a step-size of 10^{-5} . Exact MCMC approaches are too computationally costly to launch for this experiment and therefore no ground truth about the true posterior distribution is available. To this purpose, Figure 4 only compares the credibility regions associated to the posterior predictive distribution. Similarly to previous experiments, we found that D-SGLD was highly sensitive to hyperparameters choices (step-size and mini-batch size). Except for a few testing examples, most of conclusions given by DG-LMC and D-SGLD regarding the predictive uncertainty coincide. In addition, posterior accuracies on the test set given by both algorithms are similar.

6 Conclusion

In this paper, a simple algorithm coined DG-LMC has been introduced for distributed MCMC sampling. In addition, it has been established that this method inherits favorable convergence properties and numerical illustrations support our claims.

Acknowledgements

The authors acknowledge support of the Lagrange Mathematics and Computing Research Center.

References

- Abadi, M., A. Chu, I. Goodfellow, H. B. McMahan, I. Mironov, K. Talwar, and L. Zhang (2016). Deep learning with differential privacy. In *Proceedings of the 2016 ACM SIGSAC Conference on Computer and Communications Security, CCS '16*, New York, NY, USA, pp. 308–318. Association for Computing Machinery.
- Ahn, S., B. Shahbaba, and M. Welling (2014). Distributed Stochastic Gradient MCMC. In E. P. Xing and T. Jebara (Eds.), *Proceedings of the 31st International Conference on Machine Learning*, Volume 32 of *Proceedings of Machine Learning Research*, pp. 1044–1052.
- Bakry, D., I. Gentil, and M. Ledoux (2013). *Analysis and geometry of Markov diffusion operators*, Volume 348. Springer Science & Business Media.
- Bardenet, R., A. Doucet, and C. Holmes (2017). On Markov chain Monte Carlo methods for tall data. *Journal of Machine Learning Research* 18(47), 1–43.
- Bernstein, P. A. and E. Newcomer (2009). *Principles of Transaction Processing* (2nd ed.). San Francisco, CA, USA: Morgan Kaufmann Publishers Inc.
- Bottou, L., F. E. Curtis, and J. Nocedal (2018). Optimization methods for large-scale machine learning. *SIAM Review* 60(2), 223–311.
- Brosse, N., E. Moulines, and A. Durmus (2018). The Promises and Pitfalls of Stochastic Gradient Langevin Dynamics. In *Neural Information Processing Systems*, pp. 8278–8288.
- Chen, C., N. Ding, C. Li, Y. Zhang, and L. Carin (2016). Stochastic Gradient MCMC with Stale Gradients. In D. Lee, M. Sugiyama, U. Luxburg, I. Guyon, and R. Garnett (Eds.), *Advances in Neural Information Processing Systems*, Volume 29, pp. 2937–2945. Curran Associates, Inc.
- Chowdhury, A. and C. Jermaine (2018). Parallel and Distributed MCMC via Shepherdng Distributions. In *Proceedings of the Twenty-First International Conference on Artificial Intelligence and Statistics*, Volume 84, pp. 1819–1827.
- Dai, H., M. Pollock, and G. Roberts (2019). Monte Carlo fusion. *Journal of Applied Probability* 56(1), 174–191.

- Dalalyan, A. S. and A. Karagulyan (2019). User-friendly guarantees for the langevin monte carlo with inaccurate gradient. *Stochastic Processes and Their Applications* 129(12), 5278–5311.
- Dean, J. and S. Ghemawat (2004). MapReduce: Simplified Data Processing on Large Clusters. In *OSDI’04: Sixth Symposium on Operating System Design and Implementation*, San Francisco, CA, pp. 137–150.
- Dieuleveut, A. and K. K. Patel (2019). Communication trade-offs for Local-SGD with large step size. In *Advances in Neural Information Processing Systems*, Volume 32, pp. 13601–13612.
- Douc, R., E. Moulines, P. Priouret, and P. Soulier (2018). *Markov chains*. Springer.
- Durmus, A. and E. Moulines (2017, 06). Nonasymptotic convergence analysis for the unadjusted Langevin algorithm. *The Annals of Applied Probability* 27(3), 1551–1587.
- Durmus, A. and E. Moulines (2019). High-dimensional Bayesian inference via the unadjusted Langevin algorithm. *Bernoulli* 25(4A), 2854–2882.
- El Mekkaoui, K., D. Mesquita, P. Blomstedt, and S. Kaski (2020). Distributed stochastic gradient MCMC for federated learning. *arXiv preprint arXiv:2004.11231*.
- Jordan, M. I., J. D. Lee, and Y. Yang (2019). Communication-Efficient Distributed Statistical Inference. *Journal of the American Statistical Association* 114(526), 668–681.
- Kent, J. (1978, 12). Time-reversible diffusions. *Advances in Applied Probability* 10, 819–835.
- Ledoux, M. (2001). *The Concentration of Measure Phenomenon*, Volume 89 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI.
- Mesquita, D., P. Blomstedt, and S. Kaski (2020). Embarrassingly Parallel MCMC using Deep Invertible Transformations. In R. P. Adams and V. Gogate (Eds.), *Proceedings of The 35th Uncertainty in Artificial Intelligence Conference*, Volume 115 of *Proceedings of Machine Learning Research*, pp. 1244–1252.
- Minsker, S., S. Srivastava, L. Lin, and D. Dunson (2014). Scalable and robust Bayesian inference via the median posterior. In *Proceedings of the 31st International Conference on Machine Learning*.
- Neiswanger, W., C. Wang, and E. P. Xing (2014). Asymptotically exact, embarrassingly parallel MCMC. In *Proceedings of the 30th Conference on Uncertainty in Artificial Intelligence*.
- Nemeth, C. and C. Sherlock (2018, 06). Merging MCMC Subposteriors through Gaussian-Process Approximations. *Bayesian Analysis* 13(2), 507–530.

- Otto, F. and C. Villani (2000). Generalization of an Inequality by Talagrand and Links with the Logarithmic Sobolev Inequality. *Journal of Functional Analysis* 173(2), 361 – 400.
- Rabinovich, M., E. Angelino, and M. I. Jordan (2015). Variational Consensus Monte Carlo. In *Advances in Neural Information Processing Systems*, Volume 28, pp. 1207–1215.
- Raicu, I., I. Foster, A. Szalay, and G. Turcu (2006). AstroPortal: A Science Gateway for Large-scale Astronomy Data Analysis. In *TeraGrid Conference*, pp. 12–15.
- Rendell, L. J., A. M. Johansen, A. Lee, and N. Whiteley (2021). Global consensus Monte Carlo. *Journal of Computational and Graphical Statistics* 30(2), 249–259.
- Revuz, D. and M. Yor (2013). *Continuous martingales and Brownian motion*, Volume 293. Springer Science & Business Media.
- Robert, C. P. (2001). *The Bayesian Choice: from decision-theoretic foundations to computational implementation* (2 ed.). New York: Springer.
- Roberts, G. O. and J. S. Rosenthal (2006, 11). Harris recurrence of Metropolis-within-Gibbs and trans-dimensional Markov chains. *Annals of Applied Probability* 16(4), 2123–2139.
- Roberts, G. O. and R. L. Tweedie (1996, 12). Exponential convergence of Langevin distributions and their discrete approximations. *Bernoulli* 2(4), 341–363.
- Rosky, P. J., J. D. Doll, and H. L. Friedman (1978). Brownian dynamics as smart Monte Carlo simulation. *The Journal of Chemical Physics* 69(10), 4628–4633.
- Scheffé, H. (1947). A useful convergence theorem for probability distributions. *The Annals of Mathematical Statistics* 18(3), 434–438.
- Scott, S. L., A. W. Blocker, F. V. Bonassi, H. A. Chipman, E. I. George, and R. E. McCulloch (2016). Bayes and Big Data: The Consensus Monte Carlo Algorithm. *International Journal of Management Science and Engineering Management* 11, 78–88.
- Shokri, R. and V. Shmatikov (2015). Privacy-preserving deep learning. In *2015 53rd Annual Allerton Conference on Communication, Control, and Computing (Allerton)*, pp. 909–910.
- Srivastava, S., V. Cevher, Q. Dinh, and D. Dunson (2015). WASP: Scalable Bayes via barycenters of subset posteriors. In *Proceedings of the 18th International Conference on Artificial Intelligence and Statistics*, Volume 38, pp. 912–920.
- Verbraeken, J., M. Wolting, J. Katzy, J. Kloppenburg, T. Verbelen, and J. S. Rellermeyer (2020, March). A survey on distributed machine learning. *ACM Comput. Surv.* 53(2).
- Villani, C. (2008). *Optimal Transport: Old and New*. Springer Berlin Heidelberg.

- Vono, M., N. Dobigeon, and P. Chainais (2019). Split-and-augmented Gibbs sampler - Application to large-scale inference problems. *IEEE Transactions on Signal Processing* 67(6), 1648–1661.
- Vono, M., N. Dobigeon, and P. Chainais (2020). High-dimensional Gaussian sampling: A review and a unifying approach based on a stochastic proximal point algorithm. *arXiv preprint arXiv:2010.01510*.
- Vono, M., N. Dobigeon, and P. Chainais (2021). Asymptotically exact data augmentation: models, properties and algorithms. *Journal of Computational and Graphical Statistics* 30(2), 335–348.
- Vono, M., D. Paulin, and A. Doucet (2019). Efficient MCMC sampling with dimension-free convergence rate using ADMM-type splitting. *arXiv preprint arXiv:1905.11937*.
- Wang, X. and D. B. Dunson (2013). Parallelizing MCMC via Weierstrass sampler. *arXiv preprint arXiv:1312.4605*.
- Wang, X., F. Guo, K. A. Heller, and D. B. Dunson (2015). Parallelizing MCMC with random partition trees. In *Advances in Neural Information Processing Systems*.

APPENDIX

DG-LMC: A Turn-key and Scalable Synchronous Distributed MCMC Algorithm via Langevin Monte Carlo within Gibbs

Notations and conventions. We denote by $\mathcal{B}(\mathbb{R}^d)$ the Borel σ -field of \mathbb{R}^d , $\mathbb{M}(\mathbb{R}^d)$ the set of all Borel measurable functions f on \mathbb{R}^d , $\|f\|_\infty = \sup_{\mathbf{x} \in \mathbb{R}^d} |f(\mathbf{x})|$ and $\|\cdot\|$ the Euclidean norm on \mathbb{R}^d . For μ a probability measure on $(\mathbb{R}^d, \mathcal{B}(\mathbb{R}^d))$ and $f \in \mathbb{M}(\mathbb{R}^d)$ a μ -integrable function, denote by $\mu(f)$ the integral of f with respect to (w.r.t.) μ . Let μ and ν be two sigma-finite measures on $(\mathbb{R}^d, \mathcal{B}(\mathbb{R}^d))$. Denote by $\mu \ll \nu$ if μ is absolutely continuous w.r.t. ν and $d\mu/d\nu$ the associated density. Let μ, ν be two probability measures on $(\mathbb{R}^d, \mathcal{B}(\mathbb{R}^d))$. Define the Kullback-Leibler (KL) divergence of μ from ν by

$$\text{KL}(\mu|\nu) = \begin{cases} \int_{\mathbb{R}^d} \frac{d\mu}{d\nu}(\mathbf{x}) \log \left(\frac{d\mu}{d\nu}(\mathbf{x}) \right) d\nu(\mathbf{x}), & \text{if } \mu \ll \nu \\ +\infty & \text{otherwise.} \end{cases}$$

In addition, define the Pearson χ^2 -divergence of μ from ν by

$$\chi^2(\mu|\nu) = \begin{cases} \int_{\mathbb{R}^d} \left(\frac{d\mu}{d\nu}(\mathbf{x}) - 1 \right)^2 d\nu(\mathbf{x}), & \text{if } \mu \ll \nu \\ +\infty & \text{otherwise.} \end{cases}$$

We say that ζ is a transference plan of μ and ν if it is a probability measure on $(\mathbb{R}^d \times \mathbb{R}^d, \mathcal{B}(\mathbb{R}^d \times \mathbb{R}^d))$ such that for all measurable set A of \mathbb{R}^d , $\zeta(A \times \mathbb{R}^d) = \mu(A)$ and $\zeta(\mathbb{R}^d \times A) = \nu(A)$. We denote by $\mathcal{T}(\mu, \nu)$ the set of transference plans of μ and ν . In addition, we say that a couple of \mathbb{R}^d -random variables (X, Y) is a coupling of μ and ν if there exists $\zeta \in \mathcal{T}(\mu, \nu)$ such that (X, Y) are distributed according to ζ . Let \mathbf{M} be a $d \times d$ symmetric positive definite matrix. Denote $\langle \cdot, \cdot \rangle_{\mathbf{M}}$ the scalar product corresponding to \mathbf{M} , defined for any $\mathbf{x}, \mathbf{y} \in \mathbb{R}^d$ by $\langle \mathbf{x}, \mathbf{y} \rangle_{\mathbf{M}} = \mathbf{x}^\top \mathbf{M} \mathbf{y}$. Denote $\|\cdot\|_{\mathbf{M}}$ the corresponding norm. We denote by $\mathcal{P}_2(\mathbb{R}^d)$ the set of probability measures with finite 2-moment: for all $\mu \in \mathcal{P}_2(\mathbb{R}^d)$, $\int_{\mathbb{R}^d} \|\mathbf{x}\|^2 d\mu(\mathbf{x}) < \infty$. We define the Wasserstein distance of order 2 associated with $\|\cdot\|_{\mathbf{M}}$ for any probability measures $\mu, \nu \in \mathcal{P}_2(\mathbb{R}^d)$ by

$$W_{\mathbf{M}}^2(\mu, \nu) = \inf_{\zeta \in \mathcal{T}(\mu, \nu)} \int_{\mathbb{R}^d \times \mathbb{R}^d} \|\mathbf{x} - \mathbf{y}\|_{\mathbf{M}}^2 d\zeta(\mathbf{x}, \mathbf{y}).$$

In the case when $\mathbf{M} = \mathbf{I}_d$, we will denote the Wasserstein distance of order 2 by W_2 . By Villani (2008, Theorem 4.1), for all μ, ν probability measures on \mathbb{R}^d , there exists a transference plan $\zeta^* \in \mathcal{T}(\mu, \nu)$ such that for any coupling (X, Y) distributed according to ζ^* , $W_{\mathbf{M}}(\mu, \nu) = \mathbb{E}[\|\mathbf{x} - \mathbf{y}\|_{\mathbf{M}}^2]^{1/2}$. This kind of transference plan (respectively coupling) will be called an optimal transference plan (respectively optimal coupling) associated with $W_{\mathbf{M}}$. By Villani (2008, Theorem 6.16), $\mathcal{P}_2(\mathbb{R}^d)$ equipped with the Wasserstein distance $W_{\mathbf{M}}$

is a complete separable metric space. The total variation norm between two probability measures μ and ν on $(\mathbb{R}^d, \mathcal{B}(\mathbb{R}^d))$ is defined by

$$\|\mu - \nu\|_{\text{TV}} = \sup_{f \in \mathcal{M}(\mathbb{R}^d), \|f\|_{\infty} \leq 1} \left| \int_{\mathbb{R}^d} f(\mathbf{x}) d\mu(\mathbf{x}) - \int_{\mathbb{R}^d} f(\mathbf{x}) d\nu(\mathbf{x}) \right|.$$

For the sake of simplicity, with little abuse, we shall use the same notations for a probability distribution and its associated probability density function. For a Markov chain with transition kernel P on \mathbb{R}^d and invariant distribution π , we define the ε -mixing time associated to a statistical distance D , precision $\varepsilon > 0$ and initial distribution ν , by

$$t_{\text{mix}}(\varepsilon; \nu) = \min \left\{ t \geq 0 \mid D(\nu P^t, \pi) \leq \varepsilon \right\},$$

which stands for the minimum number of steps of the Markov chain such that its distribution is at most at an ε D -distance from the invariant distribution π . For $n \geq 1$, we refer to the set of integers between 1 and n with the notation $[n]$. The d -multidimensional Gaussian probability distribution with mean $\boldsymbol{\mu}$ and covariance matrix $\boldsymbol{\Sigma}$ is denoted by $N(\boldsymbol{\mu}, \boldsymbol{\Sigma})$. When $\boldsymbol{\mu} = \mathbf{0}_d$ and $\boldsymbol{\Sigma} = \mathbf{I}_d$, the associated probability density function is denoted by ϕ_d . Let $F : \mathbb{R}^d \rightarrow \mathbb{R}$ be a twice continuously differentiable function, denote $\vec{\Delta}$ the vector Laplacian of F defined, for all $x \in \mathbb{R}^d$, by $\vec{\Delta}F(x) = \{\sum_{l=1}^d (\partial^2 F_k)(x) / \partial x_l^2\}_{k=1}^d$. For $0 \leq i < j$, we use the notation $\mathbf{u}_{i:j}$ to refer to the vector $[\mathbf{u}_i^\top, \dots, \mathbf{u}_j^\top]^\top$ built by stacking $j - i + 1$ vectors $(\mathbf{u}_k; k \in \{i, \dots, j\})$. For a given matrix $\mathbf{M} \in \mathbb{R}^{d \times d}$, we denote its smallest and largest eigenvalues by $\lambda_{\min}(\mathbf{M})$ and $\lambda_{\max}(\mathbf{M})$, respectively. Fix $b \in \mathbb{N}^*$ and let $\mathbf{M}_1, \dots, \mathbf{M}_b$ be d -dimensional matrices. We denote $\prod_{\ell=i}^j \mathbf{M}_\ell = \mathbf{M}_j \dots \mathbf{M}_i$ if $i \leq j$ and with the convention $\prod_{\ell=i}^j \mathbf{M}_\ell = \mathbf{I}_d$ if $i > j$. For any $b \in \mathbb{N}^*$, $(d_i)_{i \in [b]} \in (\mathbb{N}^*)^b$ and $(\mathbf{M}_i)_{i \in [b]} \in \otimes_{i \in [b]} \mathbb{R}^{d_i \times d_i}$, we denote $\text{diag}(\mathbf{M}_1, \dots, \mathbf{M}_b)$ the unique matrix $\mathbf{M} \in \mathbb{R}^{(\sum_i d_i) \times (\sum_i d_i)}$ satisfying for any $\mathbf{u} = (\mathbf{u}_1, \dots, \mathbf{u}_b) \in \mathbb{R}^{d_1} \times \dots \times \mathbb{R}^{d_b}$, $\mathbf{M}\mathbf{u} = \sum_{i=1}^b \mathbf{M}_i \mathbf{u}_i$ which corresponds to

$$\mathbf{M} = \begin{pmatrix} \mathbf{M}_1 & \mathbf{0}_{d_1, d_2} & \cdots & \mathbf{0}_{d_b, d_b} \\ \mathbf{0}_{d_2, d_1} & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & \mathbf{0}_{d_{b-1}, d_b} \\ \mathbf{0}_{d_b, d_1} & \cdots & \mathbf{0}_{d_b, d_{b-1}} & \mathbf{M}_b \end{pmatrix}.$$

For any $\mathbf{v} \in \mathbb{R}^b$, define the block diagonal matrix

$$\mathbf{D}_{\mathbf{v}} = \text{diag}(v_1 \cdot \mathbf{I}_{d_1}, \dots, v_b \cdot \mathbf{I}_{d_b}) \in \mathbb{R}^{p \times p}. \quad (\text{S1})$$

For any symmetric matrices $\mathbf{S}_1, \mathbf{S}_2 \in \mathbb{R}^{p \times p}$, we note $\mathbf{S}_1 \preceq \mathbf{S}_2$ if and only if, for any $\mathbf{u} \in \mathbb{R}^p$, we have $\mathbf{u}^\top (\mathbf{S}_2 - \mathbf{S}_1) \mathbf{u} \geq 0$. Let $(\mathbf{X}, \mathcal{X})$ and $(\mathbf{Y}, \mathcal{Y})$ be two measurable spaces, we say that a transition probability kernel on $(\mathbf{Y} \times \mathbf{X}) \times \mathcal{Y}$ is a conditional Markov kernel. One elementary step in most Gibbs samplers corresponds to a conditional Markov kernel.

Table of contents

S1 Proof of Proposition 1	24
S1.1 Technical lemma	24
S1.2 Proof of Proposition 1	25
S2 Proof of Proposition 2	25
S2.1 Single local LMC iteration	26
S2.1.1 Supporting lemmata	27
S2.1.2 Specific case of Proposition 2	29
S2.2 Multiple local LMC iterations	30
S2.2.1 Technical lemmata	31
S2.2.2 Proof of Proposition 2	39
S3 Proof of Proposition 3	43
S3.1 Technical lemmata	43
S3.2 Proof of Proposition 3	46
S4 Proof of Proposition 4 and Proposition 5	49
S4.1 Synchronous coupling and a first estimate	50
S4.2 Proof of Proposition 4	56
S4.3 Proof of Proposition 5	60
S5 Explicit mixing times	67
S5.1 Lower bound on the number of iterations n_ε	67
S5.2 Upper bound on the tolerance parameter ρ_ε	74
S5.3 Upper bound on the step-size γ_ε and number of local iteration N_ε	75
S5.4 Discussion	76

S1 Proof of Proposition 1

Let $b' \in [b - 1]$, $p' = \sum_{i=b'+1}^b d_i$ and consider

$$\mathbf{B}_{b'}^\top = [\mathbf{A}_{b'+1}^\top / \rho_{b'+1}^{1/2} \cdots \mathbf{A}_b^\top / \rho_b^{1/2}] \in \mathbb{R}^{d \times p'}, \quad \bar{\mathbf{B}}_{b'} = \mathbf{B}_{b'}^\top \mathbf{B}_{b'} = \sum_{i=b'+1}^b \{\mathbf{A}_i^\top \mathbf{A}_i / \rho_i\} \in \mathbb{R}^{d \times d}. \quad (\text{S2})$$

Note that under **H1**, $\bar{\mathbf{B}}_{b'}$ is invertible. Indeed, it is a symmetric positive definite matrix since for any $\boldsymbol{\theta} \in \mathbb{R}^d$, $\langle \bar{\mathbf{B}}_{b'} \boldsymbol{\theta}, \boldsymbol{\theta} \rangle \geq [\min_{i \in [b]} \rho_i^{-1}] \langle \sum_{i=b'+1}^b \mathbf{A}_i^\top \mathbf{A}_i \boldsymbol{\theta}, \boldsymbol{\theta} \rangle > 0$ using that $\sum_{i=b'+1}^b \mathbf{A}_i^\top \mathbf{A}_i$

is invertible. Define the orthogonal projection onto the range of $\mathbf{B}_{b'}$ and the diagonal matrix:

$$\mathbf{P}_{b'} = \mathbf{B}_{b'} \bar{\mathbf{B}}_{b'}^{-1} \mathbf{B}_{b'}^\top, \quad \tilde{\mathbf{D}}_{b'} = \text{diag}(\mathbf{I}_{d_{b'+1}}/\rho_{b'+1}, \dots, \mathbf{I}_{d_b}/\rho_b). \quad (\text{S3})$$

S1.1 Technical lemma

Lemma S1. Assume **H1**. For any $(\boldsymbol{\theta}, \mathbf{z}_{b'+1:b}) \in \mathbb{R}^d \times \mathbb{R}^{p'}$, setting $\mathbf{z} = \mathbf{z}_{b'+1:b}$, we have

$$\begin{aligned} \sum_{i=b'+1}^b \left\{ \|\mathbf{z}_i - \mathbf{A}_i \boldsymbol{\theta}\|^2 / \rho_i \right\} &= (\tilde{\mathbf{D}}_{b'}^{1/2} \mathbf{z})^\top \{ \mathbf{I}_{p'} - \mathbf{P}_{b'} \} (\tilde{\mathbf{D}}_{b'}^{1/2} \mathbf{z}) \\ &\quad + (\boldsymbol{\theta} - \bar{\mathbf{B}}_{b'}^{-1} \mathbf{B}_{b'}^\top \tilde{\mathbf{D}}_{b'}^{1/2} \mathbf{z})^\top \bar{\mathbf{B}}_{b'} (\boldsymbol{\theta} - \bar{\mathbf{B}}_{b'}^{-1} \mathbf{B}_{b'}^\top \tilde{\mathbf{D}}_{b'}^{1/2} \mathbf{z}). \end{aligned}$$

Proof. Setting $\mathbf{b} = \mathbf{B}_{b'}^\top \tilde{\mathbf{D}}_{b'}^{1/2} \mathbf{z}$ and using the fact that $\bar{\mathbf{B}}_{b'}$ is symmetric, we have

$$\begin{aligned} \sum_{i=b'+1}^b \left\{ \|\mathbf{z}_i - \mathbf{A}_i \boldsymbol{\theta}\|^2 / \rho_i \right\} &= \boldsymbol{\theta}^\top \bar{\mathbf{B}}_{b'} \boldsymbol{\theta} - 2 \boldsymbol{\theta}^\top \mathbf{b} + \sum_{i=b'+1}^b \|\mathbf{z}_i\|^2 / \rho_i \\ &= \sum_{i=b'+1}^b \|\mathbf{z}_i\|^2 / \rho_i - \mathbf{b}^\top \bar{\mathbf{B}}_{b'}^{-1} \mathbf{b} + (\boldsymbol{\theta} - \bar{\mathbf{B}}_{b'}^{-1} \mathbf{b})^\top \bar{\mathbf{B}}_{b'} (\boldsymbol{\theta} - \bar{\mathbf{B}}_{b'}^{-1} \mathbf{b}). \end{aligned}$$

Using that $\mathbf{b}^\top \bar{\mathbf{B}}_{b'}^{-1} \mathbf{b} = (\tilde{\mathbf{D}}_{b'}^{1/2} \mathbf{z})^\top \mathbf{P}_{b'} (\tilde{\mathbf{D}}_{b'}^{1/2} \mathbf{z})$ and $\mathbf{P}_{b'}$ is a projection, $\mathbf{P}_{b'}^2 = \mathbf{P}_{b'}$ completes the proof. \square

S1.2 Proof of Proposition 1

Proposition S2. Assume **H1**. Then, the function $\psi : (\boldsymbol{\theta}, \mathbf{z}_{1:b}) \mapsto \prod_{i=1}^b \exp\{-U_i(\mathbf{z}_i) - \|\mathbf{z}_i - \mathbf{A}_i \boldsymbol{\theta}\|^2 / (2\rho_i)\}$ is integrable on $\mathbb{R}^d \times \mathbb{R}^p$, where $p = \sum_{i=1}^b d_i$.

Proof. Using **H1** and the Fubini theorem, there exists $C_1 > 0$ such that:

$$\begin{aligned} &\int_{\mathbb{R}^d} \left[\prod_{i=1}^{b'} \int_{\mathbb{R}^{d_i}} e^{-U_i(\mathbf{z}_i)} e^{-\frac{\|\mathbf{z}_i - \mathbf{A}_i \boldsymbol{\theta}\|^2}{2\rho_i}} d\mathbf{z}_i \cdot \prod_{j=b'+1}^b \int_{\mathbb{R}^{d_j}} e^{-U_j(\mathbf{z}_j)} e^{-\frac{\|\mathbf{z}_j - \mathbf{A}_j \boldsymbol{\theta}\|^2}{2\rho_j}} d\mathbf{z}_j \right] d\boldsymbol{\theta} \\ &\leq C_1 \int_{\mathbb{R}^d} \left[\prod_{i=1}^{b'} \int_{\mathbb{R}^{d_i}} e^{-\frac{\|\mathbf{z}_i - \mathbf{A}_i \boldsymbol{\theta}\|^2}{2\rho_i}} d\mathbf{z}_i \cdot \prod_{j=b'+1}^b \int_{\mathbb{R}^{d_j}} e^{-U_j(\mathbf{z}_j)} e^{-\frac{\|\mathbf{z}_j - \mathbf{A}_j \boldsymbol{\theta}\|^2}{2\rho_j}} d\mathbf{z}_j \right] d\boldsymbol{\theta} \\ &\leq C_1 \prod_{i=1}^{b'} (2\pi\rho_i)^{d_i/2} \int_{\mathbb{R}^d} \left[\prod_{j=b'+1}^b \int_{\mathbb{R}^{d_j}} e^{-U_j(\mathbf{z}_j)} \exp\left(-\|\mathbf{z}_j - \mathbf{A}_j \boldsymbol{\theta}\|^2 / (2\rho_j)\right) d\mathbf{z}_j \right] d\boldsymbol{\theta} \\ &= C_1 \prod_{i=1}^{b'} (2\pi\rho_i)^{d_i/2} \int_{\mathbb{R}^{d_{b'+1}}} \cdots \int_{\mathbb{R}^{d_b}} \left[\prod_{j=b'+1}^b e^{-U_j(\mathbf{z}_j)} \right] \left[\int_{\mathbb{R}^d} \prod_{j=b'+1}^b e^{-\frac{\|\mathbf{z}_j - \mathbf{A}_j \boldsymbol{\theta}\|^2}{2\rho_j}} d\boldsymbol{\theta} \right] d\mathbf{z}_{b'+1:b}. \end{aligned} \quad (\text{S4})$$

Using Lemma S1 and the fact that $\mathbf{I}_{p'} - \mathbf{P}_{b'}$ is positive definite, we obtain

$$\begin{aligned}
& \int_{\mathbb{R}^d} \prod_{j=b'+1}^b \exp\left(-\|\mathbf{z}_j - \mathbf{A}_j \boldsymbol{\theta}\|^2 / (2\rho_j)\right) d\boldsymbol{\theta} \\
&= \exp\left(-(\tilde{\mathbf{D}}_{b'}^{1/2} \mathbf{z})^\top \{\mathbf{I}_{p'} - \mathbf{P}_{b'}\} (\tilde{\mathbf{D}}_{b'}^{1/2} \mathbf{z}) / 2\right) \\
&\times \int_{\mathbb{R}^d} \exp\left(-(\boldsymbol{\theta} - \bar{\mathbf{B}}_{b'}^{-1} \mathbf{B}_{b'}^\top \tilde{\mathbf{D}}_{b'}^{1/2} \mathbf{z})^\top \bar{\mathbf{B}}_{b'} (\boldsymbol{\theta} - \bar{\mathbf{B}}_{b'}^{-1} \mathbf{B}_{b'}^\top \tilde{\mathbf{D}}_{b'}^{1/2} \mathbf{z}) / 2\right) d\boldsymbol{\theta} \\
&\leq \det(\bar{\mathbf{B}}_{b'})^{-1/2} (2\pi)^{d/2}.
\end{aligned}$$

Then, the proof is completed by plugging this expression into (S4) and using from H1 that $\mathbf{z}_{b'+1:b} \mapsto \prod_{j=b'+1}^b e^{-U_j(\mathbf{z}_j)}$ is integrable. \square

S2 Proof of Proposition 2

This section aims at proving Proposition 2 in the main paper. To ease the understanding, we dissociate the scenarios where $\max_{i \in [b]} N_i = 1$ and $\max_{i \in [b]} N_i > 1$. In addition, in all this section $\boldsymbol{\rho} \in (\mathbb{R}_+^*)^b$ is assumed to be fixed.

S2.1 Single local LMC iteration

In this section, we assume that a single LMC step is performed locally on each worker, that is $\max_{i \in [b]} N_i = 1$. For this, we introduce the conditional Markov transition kernel defined for any $\boldsymbol{\gamma} = (\gamma_1, \dots, \gamma_b)$, $\boldsymbol{\theta} \in \mathbb{R}^d$, $\mathbf{z} = (\mathbf{z}_1, \dots, \mathbf{z}_b) \in \mathbb{R}^{d_1} \times \dots \times \mathbb{R}^{d_b}$, and for $i \in [b]$, $\mathbf{B}_i \in \mathcal{B}(\mathbb{R}^{d_i})$, by

$$Q_{\boldsymbol{\rho}, \boldsymbol{\gamma}}(\mathbf{z}, \mathbf{B}_1 \times \dots \times \mathbf{B}_b | \boldsymbol{\theta}) = \prod_{i=1}^b R_{\rho_i, \gamma_i}(\mathbf{z}_i, \mathbf{B}_i | \boldsymbol{\theta}), \quad (\text{S5})$$

where

$$R_{\rho_i, \gamma_i}(\mathbf{z}_i, \mathbf{B}_i | \boldsymbol{\theta}) = \int_{\mathbf{B}_i} \exp\left\{-\frac{1}{4\gamma_i} \left\| \tilde{\mathbf{z}}_i - \left(1 - \frac{\gamma_i}{\rho_i}\right) \mathbf{z}_i - \frac{\gamma_i}{\rho_i} \mathbf{A}_i \boldsymbol{\theta} + \gamma_i \nabla U_i(\mathbf{z}_i) \right\|^2\right\} \frac{d\tilde{\mathbf{z}}_i}{(4\pi\gamma_i)^{d_i/2}}. \quad (\text{S6})$$

Recall that $p = \sum_{i=1}^b d_i$. The considered Gibbs sampler in Algorithm 1 defines a homogeneous Markov chain $X_n^\top = (\boldsymbol{\theta}_n^\top, Z_n^\top)_{n \geq 1}$ where $Z_n^\top = ([Z_n^1]^\top, \dots, [Z_n^b]^\top)$. Indeed, it is easy to show that for any $n \in \mathbb{N}$ and measurable bounded function $f : \mathbb{R}^p \rightarrow \mathbb{R}_+$, $\mathbb{E}[f(Z_{n+1}) | X_n] = \int_{\mathbb{R}^p} f(\mathbf{z}) Q_{\boldsymbol{\rho}, \boldsymbol{\gamma}}(Z_n, d\mathbf{z} | \boldsymbol{\theta}_n)$ and therefore $(X_n)_{n \in \mathbb{N}}$ is associated with the Markov kernel defined, for any $\mathbf{x}^\top = (\boldsymbol{\theta}^\top, \mathbf{z}^\top) \in \mathbb{R}^d \times \mathbb{R}^p$ and $\mathbf{A} \in \mathcal{B}(\mathbb{R}^d)$, $\mathbf{B} \in \mathcal{B}(\mathbb{R}^p)$, by

$$P_{\boldsymbol{\rho}, \boldsymbol{\gamma}}(\mathbf{x}, \mathbf{A} \times \mathbf{B}) = \int_{\mathbf{B}} Q_{\boldsymbol{\rho}, \boldsymbol{\gamma}}(\mathbf{z}, d\tilde{\mathbf{z}} | \boldsymbol{\theta}) \int_{\mathbf{A}} \Pi_{\boldsymbol{\rho}}(d\tilde{\boldsymbol{\theta}} | \tilde{\mathbf{z}}), \quad (\text{S7})$$

where $\Pi_{\boldsymbol{\rho}}(\cdot|\tilde{\mathbf{z}})$ is defined in (5). Let $(\xi_n)_{n \geq 1}$ be a sequence of i.i.d. d -dimensional standard Gaussian random variables independent of the family of independent random variables $\{(\eta_n^i)_{n \geq 1} : i \in [b]\}$ where for any $i \in [b]$ and $n \geq 1$, η_n^i is a d_i -dimensional standard Gaussian random variable. We define the stochastic processes $(X_n, \tilde{X}_n)_{n \geq 0}$ on $\mathbb{R}^p \times \mathbb{R}^p$ starting from $(X_0, \tilde{X}_0) = (\mathbf{x}, \tilde{\mathbf{x}}) = ((\boldsymbol{\theta}^\top, \mathbf{z}^\top)^\top, (\tilde{\boldsymbol{\theta}}^\top, \tilde{\mathbf{z}}^\top)^\top)$ and following the recursion for $n \geq 0$,

$$X_{n+1} = (\theta_{n+1}^\top, Z_{n+1}^\top)^\top, \quad \tilde{X}_{n+1} = (\tilde{\theta}_{n+1}^\top, \tilde{Z}_{n+1}^\top)^\top, \quad (\text{S8})$$

where $Z_{n+1} = ([Z_{n+1}^1]^\top, \dots, [Z_{n+1}^b]^\top)^\top$, $\tilde{Z}_{n+1} = ([\tilde{Z}_{n+1}^1]^\top, \dots, [\tilde{Z}_{n+1}^b]^\top)^\top$ are defined, for any $i \in [b]$, by

$$\begin{aligned} Z_{n+1}^i &= (1 - \gamma_i/\rho_i) Z_n^i + (\gamma_i/\rho_i) \mathbf{A}_i \theta_n - \gamma_i \nabla U_i(Z_n^i) + \sqrt{2\gamma_i} \eta_{n+1}^i, \\ \tilde{Z}_{n+1}^i &= (1 - \gamma_i/\rho_i) \tilde{Z}_n^i + (\gamma_i/\rho_i) \mathbf{A}_i \tilde{\theta}_n - \gamma_i \nabla U_i(\tilde{Z}_n^i) + \sqrt{2\gamma_i} \eta_{n+1}^i, \end{aligned} \quad (\text{S9})$$

and $\theta_{n+1}, \tilde{\theta}_{n+1}$ by

$$\theta_{n+1} = \bar{\mathbf{B}}_0^{-1} \mathbf{B}_0^\top \tilde{\mathbf{D}}_0^{1/2} Z_{n+1} + \bar{\mathbf{B}}_0^{-1/2} \xi_{n+1}, \quad \tilde{\theta}_{n+1} = \bar{\mathbf{B}}_0^{-1} \mathbf{B}_0^\top \tilde{\mathbf{D}}_0^{1/2} \tilde{Z}_{n+1} + \bar{\mathbf{B}}_0^{-1/2} \xi_{n+1}, \quad (\text{S10})$$

where $\bar{\mathbf{B}}_0$, \mathbf{B}_0 and $\tilde{\mathbf{D}}_0$ are given in (S2) and (S3), respectively. Note that X_n and \tilde{X}_n are distributed according to $\delta_{\mathbf{x}} P_{\boldsymbol{\rho}, \gamma}^n$ and $\delta_{\tilde{\mathbf{x}}} P_{\boldsymbol{\rho}, \gamma}^n$, respectively. Hence, by definition of the Wasserstein distance of order 2, it follows that

$$W_2(\delta_{\mathbf{x}} P_{\boldsymbol{\rho}, \gamma}^n, \delta_{\tilde{\mathbf{x}}} P_{\boldsymbol{\rho}, \gamma}^n) \leq \mathbb{E} \left[\|X_n - \tilde{X}_n\|^2 \right]^{1/2}. \quad (\text{S11})$$

Thus, in this section we focus on upper bounding the squared norm $\|X_n - \tilde{X}_n\|$ from which we get an explicit bound on the Wasserstein distance thanks to the previous inequality.

S2.1.1 Supporting lemmata

Note that **H1** implies the invertibility of the matrix \mathbf{B}_0 defined in (S2) since we have the existence of $b' \in [b-1]$, such that $\sum_{i=b'+1}^b \lambda_{\min}(\mathbf{A}_i^\top \mathbf{A}_i)/\rho_i > 0$ and by the semi-positiveness of the symmetric matrices $\{\mathbf{A}_i^\top \mathbf{A}_i\}_{i \in [b]}$, we get that $\lambda_{\min}(\mathbf{B}_0) = \sum_{i=1}^b \lambda_{\min}(\mathbf{A}_i^\top \mathbf{A}_i)/\rho_i \geq \sum_{i=b'+1}^b \lambda_{\min}(\mathbf{A}_i^\top \mathbf{A}_i)/\rho_i$. To prove Proposition 2 in the case $\max_{i \in [b]} N_i = 1$, we first upper bound (S83) by building upon the following two technical lemmas.

Lemma S3. *Assume **H1** and consider $(X_n, \tilde{X}_n)_{n \in \mathbb{N}}$ defined in (S8). Then, for any $n \in \mathbb{N}$, it holds almost surely that*

$$\|X_{n+1} - \tilde{X}_{n+1}\|^2 \leq (1 + \|\bar{\mathbf{B}}_0^{-1} \mathbf{B}_0^\top \tilde{\mathbf{D}}_0^{1/2}\|^2) \|Z_{n+1} - \tilde{Z}_{n+1}\|^2.$$

Proof. Let $n \geq 0$. By (S10), we have $\theta_{n+1} - \tilde{\theta}_{n+1} = \bar{\mathbf{B}}_0^{-1} \mathbf{B}_0^\top \tilde{\mathbf{D}}_0^{1/2} (Z_{n+1} - \tilde{Z}_{n+1})$ which implies that

$$\|X_{n+1} - \tilde{X}_{n+1}\|^2 = \|\theta_{n+1} - \tilde{\theta}_{n+1}\|^2 + \|Z_{n+1} - \tilde{Z}_{n+1}\|^2 \leq (1 + \|\bar{\mathbf{B}}_0^{-1} \mathbf{B}_0^\top \tilde{\mathbf{D}}_0^{1/2}\|^2) \|Z_{n+1} - \tilde{Z}_{n+1}\|^2.$$

□

Define the contraction factor

$$\kappa_\gamma = \max_{i \in [b]} \{ |1 - \gamma_i m_i| \vee |1 - \gamma_i (M_i + 1/\rho_i)| \} . \quad (\text{S12})$$

Then, the following result holds.

Lemma S4. Assume **H1-H2** and let $\gamma \in (\mathbb{R}_+^*)^b$. Then for any $\mathbf{x} = (\mathbf{z}^\top, \boldsymbol{\theta}^\top)^\top$, $\tilde{\mathbf{x}} = (\tilde{\mathbf{z}}^\top, \tilde{\boldsymbol{\theta}}^\top)^\top$, with $(\boldsymbol{\theta}, \tilde{\boldsymbol{\theta}}) \in (\mathbb{R}^d)^2$ and $(\mathbf{z}, \tilde{\mathbf{z}}) \in (\mathbb{R}^p)^2$, for any $n \geq 1$, we have

$$\begin{aligned} W_2(\delta_{\mathbf{x}} P_{\rho, \gamma}^n, \delta_{\tilde{\mathbf{x}}} P_{\rho, \gamma}^n) &\leq \kappa_\gamma^{n-1} \cdot \left((1 + \|\bar{\mathbf{B}}_0^{-1} \mathbf{B}_0^\top \tilde{\mathbf{D}}_0^{1/2}\|^2) \cdot \frac{\max_{i \in [b]} \{\gamma_i\}}{\min_{i \in [b]} \{\gamma_i\}} \right)^{1/2} \\ &\quad \times \left[\kappa_\gamma \|\mathbf{z} - \tilde{\mathbf{z}}\| + \|\mathbf{D}_{\gamma/\sqrt{\rho}} \mathbf{B}_0\| \|\boldsymbol{\theta} - \tilde{\boldsymbol{\theta}}\| \right] , \end{aligned}$$

where $\mathbf{D}_{\gamma/\sqrt{\rho}}$ is defined as in (S1) with $\gamma/\sqrt{\rho} = (\gamma_1/\rho_1^{1/2}, \dots, \gamma_b/\rho_b^{1/2})$, $\bar{\mathbf{B}}_0$, \mathbf{B}_0 , $P_{\rho, \gamma}$ and κ_γ are given in (S2), (S7), (S12), respectively.

Proof. Consider $(X_k, \tilde{X}_k)_{k \in \mathbb{N}}$ defined in (S8). By (S83) and Lemma S3, we need to bound $(\|Z_k - \tilde{Z}_k\|)_{k \in \mathbb{N}}$. Let $n \in \mathbb{N}^*$. For any $i \in [b]$, we have by (S9), that

$$Z_{n+1}^i - \tilde{Z}_{n+1}^i = \left(1 - \frac{\gamma_i}{\rho_i}\right) (Z_n^i - \tilde{Z}_n^i) + \frac{\gamma_i}{\rho_i} \mathbf{A}_i (\theta_n - \tilde{\theta}_n) - \gamma_i \left(\nabla U_i(Z_n^i) - \nabla U_i(\tilde{Z}_n^i) \right) . \quad (\text{S13})$$

Since U_i is twice differentiable, we have

$$\nabla U_i(Z_n^i) - \nabla U_i(\tilde{Z}_n^i) = \int_0^1 \nabla^2 U_i(\tilde{Z}_n^i + t(Z_n^i - \tilde{Z}_n^i)) dt \cdot (Z_n^i - \tilde{Z}_n^i) .$$

Using $\theta_n - \tilde{\theta}_n = \bar{\mathbf{B}}_0^{-1} \mathbf{B}_0^\top \tilde{\mathbf{D}}_0^{1/2} (Z_n - \tilde{Z}_n)$, it follows that

$$\begin{aligned} Z_{n+1}^i - \tilde{Z}_{n+1}^i &= \left(\left[1 - \frac{\gamma_i}{\rho_i}\right] \mathbf{I}_{d_i} - \gamma_i \int_0^1 \nabla^2 U_i(\tilde{Z}_n^i + t(Z_n^i - \tilde{Z}_n^i)) dt \right) (Z_n^i - \tilde{Z}_n^i) \\ &\quad + \frac{\gamma_i}{\rho_i} \mathbf{A}_i \bar{\mathbf{B}}_0^{-1} \mathbf{B}_0^\top \tilde{\mathbf{D}}_0^{1/2} (Z_n - \tilde{Z}_n) . \end{aligned}$$

Consider the $p \times p$ block diagonal matrix defined by

$$\mathbf{D}_{U, n} = \text{diag} \left(\gamma_1 \int_0^1 \nabla^2 U_1(\tilde{Z}_n^1 + t(Z_n^1 - \tilde{Z}_n^1)) dt, \dots, \gamma_b \int_0^1 \nabla^2 U_b(\tilde{Z}_n^b + t(Z_n^b - \tilde{Z}_n^b)) dt \right) .$$

With the projection matrix \mathbf{P}_0 defined in (S3), the difference $Z_{n+1} - \tilde{Z}_{n+1}$ can be rewritten as

$$Z_{n+1} - \tilde{Z}_{n+1} = \left(\mathbf{I}_p - \mathbf{D}_{U, n} - \mathbf{D}_{\gamma}^{1/2} \mathbf{D}_{\gamma/\rho}^{1/2} (\mathbf{I}_p - \mathbf{P}_0) \tilde{\mathbf{D}}_0^{1/2} \right) (Z_n - \tilde{Z}_n) ,$$

where $\mathbf{D}_{\gamma/\rho}$ is defined as in (S1) with $\gamma/\rho = (\gamma_1/\rho_1, \dots, \gamma_b/\rho_b)$. Since $\mathbf{D}_{U,n}$ commutes with \mathbf{D}_γ and \mathbf{P}_0 is an orthogonal projection matrix, using **H2-(i)-(ii)**, we get

$$\begin{aligned} & \|Z_{n+1} - \tilde{Z}_{n+1}\|_{\mathbf{D}_\gamma^{-1}} \\ &= \|\mathbf{D}_\gamma^{-1/2}(\mathbf{D}_\gamma^{1/2}\mathbf{D}_\gamma^{-1/2} - \mathbf{D}_\gamma^{1/2}\mathbf{D}_{U,n}\mathbf{D}_\gamma^{-1/2} - \mathbf{D}_\gamma^{1/2}\mathbf{D}_{\gamma/\rho}^{1/2}(\mathbf{I}_p - \mathbf{P}_0)\mathbf{D}_{\gamma/\rho}^{1/2}\mathbf{D}_\gamma^{-1/2})(Z_n - \tilde{Z}_n)\| \\ &\leq \|\mathbf{I}_p - \mathbf{D}_{U,n} - \mathbf{D}_{\gamma/\rho}^{1/2}(\mathbf{I}_p - \mathbf{P}_0)\mathbf{D}_{\gamma/\rho}^{1/2}\| \|Z_n - \tilde{Z}_n\|_{\mathbf{D}_\gamma^{-1}}. \end{aligned}$$

Note that **H1** and **H2** and the fact that \mathbf{P}_0 is an orthogonal projector, so $\mathbf{0}_p \preceq \mathbf{I}_p - \mathbf{P}_0$, imply that

$$\begin{aligned} \text{diag}(\{1 - \gamma_1(M_1 + 1/\rho_1)\}\mathbf{I}_{d_1}, \dots, \{1 - \gamma_b(M_b + 1/\rho_b)\}\mathbf{I}_{d_b}) &\preceq \mathbf{I}_p - \mathbf{D}_{U,n} - \mathbf{D}_{\gamma/\rho}^{1/2}(\mathbf{I}_p - \mathbf{P}_0)\mathbf{D}_{\gamma/\rho}^{1/2} \\ &\preceq \text{diag}(\{1 - \gamma_1 m_1\}\mathbf{I}_{d_1}, \dots, \{1 - \gamma_b m_b\}\mathbf{I}_{d_b}). \end{aligned}$$

Therefore, we get

$$\begin{aligned} \|Z_{n+1} - \tilde{Z}_{n+1}\|_{\mathbf{D}_\gamma^{-1}} &\leq \max_{i \in [b]} \{\max(|1 - \gamma_i m_i|, |1 - \gamma_i(M_i + 1/\rho_i)|)\} \|Z_n - \tilde{Z}_n\|_{\mathbf{D}_\gamma^{-1}} \\ &= \kappa_\gamma \|Z_n - \tilde{Z}_n\|_{\mathbf{D}_\gamma^{-1}}. \end{aligned} \tag{S14}$$

An immediate induction shows, for any $n \geq 1$,

$$\|Z_n - \tilde{Z}_n\|_{\mathbf{D}_\gamma^{-1}} \leq \kappa_\gamma^{n-1} \|Z_1 - \tilde{Z}_1\|_{\mathbf{D}_\gamma^{-1}}. \tag{S15}$$

In addition, by (S13), we have for any $i \in [b]$,

$$Z_1^i - \tilde{Z}_1^i = \left(1 - \frac{\gamma_i}{\rho_i}\right)(\mathbf{z}_i - \tilde{\mathbf{z}}_i) + \frac{\gamma_i}{\rho_i} \mathbf{A}_i(\boldsymbol{\theta} - \tilde{\boldsymbol{\theta}}) - \gamma_i(\nabla U_i(\mathbf{z}_i) - \nabla U_i(\tilde{\mathbf{z}}_i)).$$

It follows that $Z_1 - \tilde{Z}_1 = (\mathbf{I}_p - \mathbf{D}_{\gamma/\rho} - \mathbf{D}_{U,0})(\mathbf{z} - \tilde{\mathbf{z}}) + \mathbf{D}_{\gamma/\rho} \tilde{\mathbf{D}}_0^{-1/2} \mathbf{B}_0(\boldsymbol{\theta} - \tilde{\boldsymbol{\theta}})$. Using the triangle inequality and **H2** gives

$$\begin{aligned} \|Z_1 - \tilde{Z}_1\|_{\mathbf{D}_\gamma^{-1}} &\leq (\min_{i \in [b]} \{\gamma_i\})^{-1/2} \|(\mathbf{I}_p - \mathbf{D}_{\gamma/\rho} - \mathbf{D}_{U,0})(\mathbf{z} - \tilde{\mathbf{z}}) + (\mathbf{D}_{\gamma/\sqrt{\rho}} \mathbf{B}_0(\boldsymbol{\theta} - \tilde{\boldsymbol{\theta}}))\| \\ &\leq (\min_{i \in [b]} \{\gamma_i\})^{-1/2} \left[\|\mathbf{I}_p - \mathbf{D}_{\gamma/\rho} - \mathbf{D}_{U,0}\| \|\mathbf{z} - \tilde{\mathbf{z}}\| + \|\mathbf{D}_{\gamma/\sqrt{\rho}} \mathbf{B}_0\| \|\boldsymbol{\theta} - \tilde{\boldsymbol{\theta}}\| \right] \\ &\leq (\min_{i \in [b]} \{\gamma_i\})^{-1/2} \left[\max_{i \in [b]} \{|1 - \gamma_i(m_i + 1/\rho_i)|, |1 - \gamma_i(M_i + 1/\rho_i)|\} \|\mathbf{z} - \tilde{\mathbf{z}}\| \right. \\ &\quad \left. + \|\mathbf{D}_{\gamma/\sqrt{\rho}} \mathbf{B}_0\| \|\boldsymbol{\theta} - \tilde{\boldsymbol{\theta}}\| \right] \\ &\leq (\min_{i \in [b]} \{\gamma_i\})^{-1/2} \left[\kappa_\gamma \|\mathbf{z} - \tilde{\mathbf{z}}\| + \|\mathbf{D}_{\gamma/\sqrt{\rho}} \mathbf{B}_0\| \|\boldsymbol{\theta} - \tilde{\boldsymbol{\theta}}\| \right]. \end{aligned}$$

Combining (S15) and the previous inequality and using Lemma S3, we get for $n \geq 1$,

$$\begin{aligned} \|X_n - \tilde{X}_n\|^2 &\leq \kappa_\gamma^{2(n-1)} \left(1 + \|\bar{\mathbf{B}}_0^{-1} \mathbf{B}_0^\top \tilde{\mathbf{D}}_0^{1/2}\|^2\right) \frac{\max_{i \in [b]} \{\gamma_i\}}{\min_{i \in [b]} \{\gamma_i\}} \\ &\quad \times \left[\kappa_\gamma \|\mathbf{z} - \tilde{\mathbf{z}}\| + \|\mathbf{D}_{\gamma/\sqrt{\rho}} \mathbf{B}_0\| \|\boldsymbol{\theta} - \tilde{\boldsymbol{\theta}}\| \right]^2. \end{aligned}$$

The proof is concluded by (S83). \square

S2.1.2 Specific case of Proposition 2

Based on the previous lemmata, we provide in what follows a specific instance of Proposition 2 in the scenario where $\max_{i \in [b]} N_i = 1$.

Proposition S5. *Assume **H1-H2** and let $\gamma \in (\mathbb{R}_+^*)^b$ such that, for any $i \in [b]$, $\gamma_i \leq 2(m_i + M_i + 1/\rho_i)^{-1}$. Then, $P_{\rho, \gamma}$ defined in (S7) admits a unique stationary distribution $\Pi_{\rho, \gamma}$ and for any $\mathbf{x} = (\mathbf{z}^\top, \boldsymbol{\theta}^\top)^\top$ with $\boldsymbol{\theta} \in \mathbb{R}^d$, $\mathbf{z} \in \mathbb{R}^p$ and any $n \in \mathbb{N}^*$, we have*

$$W_2^2(\delta_{\mathbf{x}} P_{\rho, \gamma}^n, \Pi_{\rho, \gamma}) \leq \left(1 - \min_{i \in [b]} \{\gamma_i m_i\}\right)^{2(n-1)} \left((1 + \|\bar{\mathbf{B}}_0^{-1} \mathbf{B}_0^\top \tilde{\mathbf{D}}_0^{1/2}\|^2) \cdot \frac{\max_{i \in [b]} \{\gamma_i\}}{\min_{i \in [b]} \{\gamma_i\}} \right) \\ \times \int_{\mathbb{R}^d \times \mathbb{R}^p} \left[(1 - \min_{i \in [b]} \{\gamma_i m_i\}) \|\mathbf{z} - \tilde{\mathbf{z}}\| + \|\mathbf{D}_{\gamma/\sqrt{\rho}} \mathbf{B}_0\| \|\boldsymbol{\theta} - \tilde{\boldsymbol{\theta}}\| \right]^2 d\Pi_{\rho, \gamma}(\tilde{\mathbf{x}}),$$

where $\bar{\mathbf{B}}_0, \mathbf{B}_0, \tilde{\mathbf{D}}_0, P_{\rho, \gamma}$ are defined in (S2) and (S3).

Proof. For any $i \in [b]$, note that the condition $0 < \gamma_i \leq 2(m_i + M_i + 1/\rho_i)^{-1}$ ensures that $\kappa_\gamma = 1 - \min_{i \in [b]} \{\gamma_i m_i\} \in (0, 1)$ and the proof follows from Lemma S4 combined with Douc et al. (2018, Lemma 20.3.2, Theorem 20.3.4). \square

S2.2 Multiple local LMC iterations

In this section, we consider the general case $\max_{i \in [b]} N_i \geq 1$. For this, we introduce the conditional Markov transition kernel defined for any $\gamma = (\gamma_1, \dots, \gamma_b)$, $\mathbf{N} = (N_1, \dots, N_b)$, $\boldsymbol{\theta} \in \mathbb{R}^d$, $\mathbf{z} = (\mathbf{z}_1, \dots, \mathbf{z}_b) \in \mathbb{R}^{d_1} \times \dots \times \mathbb{R}^{d_b}$, for $i \in [b]$ and $\mathbf{B}_i \in \mathcal{B}(\mathbb{R}^{d_i})$, by

$$Q_{\rho, \gamma, \mathbf{N}}(\mathbf{z}, \mathbf{B}_1 \times \dots \times \mathbf{B}_b | \boldsymbol{\theta}) = \prod_{i=1}^b R_{\rho_i, \gamma_i}^{N_i}(\mathbf{z}_i, \mathbf{B}_i | \boldsymbol{\theta}), \quad (\text{S16})$$

where R_{ρ_i, γ_i} is defined by (S6). Then, as in the case $\max_{i \in [b]} N_i = 1$, the Gibbs sampler presented in Algorithm 1 defines a homogeneous Markov chain $X_n^\top = (\theta_n^\top, Z_n^\top)_{n \geq 1}$ where $Z_n^\top = ([Z_n^1]^\top, \dots, [Z_n^b]^\top)$. Indeed, it is easy to show that for any $n \in \mathbb{N}$ and measurable function $f : \mathbb{R}^p \rightarrow \mathbb{R}_+$, $\mathbb{E}[f(Z_{n+1}) | X_n] = \int_{\mathbb{R}^p} f(\mathbf{z}) Q_{\rho, \gamma, \mathbf{N}}(Z_n, d\mathbf{z} | \theta_n)$. Therefore, $(X_n)_{n \in \mathbb{N}}$ is associated with the Markov kernel defined, for any $\mathbf{x}^\top = (\boldsymbol{\theta}^\top, \mathbf{z}^\top)$ and $\mathbf{A} \in \mathcal{B}(\mathbb{R}^d)$, $\mathbf{B} \in \mathcal{B}(\mathbb{R}^p)$, by

$$P_{\rho, \gamma, \mathbf{N}}(\mathbf{x}, \mathbf{A} \times \mathbf{B}) = \int_{\mathbf{B}} Q_{\rho, \gamma, \mathbf{N}}(\mathbf{z}, d\tilde{\mathbf{z}} | \boldsymbol{\theta}) \int_{\mathbf{A}} \Pi_{\rho}(d\tilde{\boldsymbol{\theta}} | \tilde{\mathbf{z}}), \quad (\text{S17})$$

where $\Pi_{\rho}(\cdot | \tilde{\mathbf{z}})$ is defined in (5). We now define a coupling between $\delta_{\mathbf{x}} P_{\rho, \gamma, \mathbf{N}}^n$ and $\delta_{\tilde{\mathbf{x}}} P_{\rho, \gamma, \mathbf{N}}^n$ for any $n \geq 1$ and $\mathbf{x}, \tilde{\mathbf{x}} \in \mathbb{R}^d \times \mathbb{R}^p$. Let $(\xi_n)_{n \geq 1}$ be a sequence of i.i.d. d -dimensional standard

Gaussian random variables independent of the family of independent random variables $\{(\eta_n^i)_{n \geq 1} : i \in [b]\}$ where for any $i \in [b]$ and $n \geq 1$, η_n^i is a d_i -dimensional standard Gaussian random variable. Define by induction the synchronous coupling $(\theta_n, Z_n)_{n \geq 0}, (\tilde{\theta}_n, \tilde{Z}_n)_{n \geq 0}$, for any $i \in [b]$ starting from $(\theta_0, Z_0) = \mathbf{x} = (\boldsymbol{\theta}, \mathbf{z})$, $(\tilde{\theta}_0, \tilde{Z}_0) = \tilde{\mathbf{x}} = (\tilde{\boldsymbol{\theta}}, \tilde{\mathbf{z}})$ and for any $n \geq 0$ by

$$\begin{aligned}\tilde{Z}_{n+1}^i &= \tilde{Y}_{N_i}^{(i,n)} , & \tilde{\theta}_{n+1} &= \bar{\mathbf{B}}_0^{-1} \mathbf{B}_0^\top \tilde{\mathbf{D}}_0^{1/2} \tilde{Z}_{n+1} + \bar{\mathbf{B}}_0^{-1/2} \xi_{n+1} , \\ Z_{n+1}^i &= Y_{N_i}^{(i,n)} , & \theta_{n+1} &= \bar{\mathbf{B}}_0^{-1} \mathbf{B}_0^\top \mathbf{D}_0^{1/2} Z_{n+1} + \bar{\mathbf{B}}_0^{-1/2} \xi_{n+1} ,\end{aligned}\tag{S18}$$

where $\bar{\mathbf{B}}_0, \mathbf{B}_0, \tilde{\mathbf{D}}_0$ are given by (S2)-(S3) and $\tilde{Y}_0^{(i,n)} = \tilde{Z}_n^i$, $Y_0^{(i,n)} = Z_n^i$, and for any $k \in \mathbb{N}$

$$\begin{aligned}\tilde{Y}_{k+1}^{(i,n)} &= \tilde{Y}_k^{(i,n)} - \gamma_i \nabla V_i(\tilde{Y}_k^{(i,n)}) + (\gamma_i / \rho_i) \mathbf{A}_i \tilde{\theta}_n + \sqrt{2\gamma_i} \eta_{k+1}^{(i,n)} , \\ Y_{k+1}^{(i,n)} &= Y_k^{(i,n)} - \gamma_i \nabla V_i(Y_k^{(i,n)}) + (\gamma_i / \rho_i) \mathbf{A}_i \theta_n + \sqrt{2\gamma_i} \eta_{k+1}^{(i,n)} ,\end{aligned}\tag{S19}$$

where, for any $\mathbf{z}_i \in \mathbb{R}^{d_i}$, V_i is defined by

$$V_i(\mathbf{z}_i) = U_i(\mathbf{z}_i) + (2\rho_i)^{-1} \|\mathbf{z}_i\|^2 .\tag{S20}$$

For any $n, k \in \mathbb{N}$ consider the $p \times p$ matrices defined by

$$\begin{aligned}\mathbf{H}_{U,k}^{(n)} &= \text{diag} \left(\gamma_1 \int_0^1 \nabla^2 U_1((1-s)Y_k^{(1,n)} + s\tilde{Y}_k^{(1,n)}) ds, \right. \\ &\quad \left. \dots, \gamma_b \int_0^1 \nabla^2 U_b((1-s)Y_k^{(b,n)} + s\tilde{Y}_k^{(b,n)}) ds \right) , \\ \mathbf{J}(k) &= \text{diag} \left(\mathbf{1}_{[N_1]}(k+1) \cdot \mathbf{I}_{d_1}, \dots, \mathbf{1}_{[N_b]}(k+1) \cdot \mathbf{I}_{d_b} \right) ,\end{aligned}\tag{S21}$$

$$\mathbf{C}_k^{(n)} = \mathbf{J}(k)(\mathbf{D}_{\gamma/\rho} + \mathbf{H}_{U,k}^{(n)}) ,\tag{S22}$$

$$\mathbf{M}_{k+1}^{(n)} = (\mathbf{I}_p - \mathbf{C}_0^{(n)})^{-1} \dots (\mathbf{I}_p - \mathbf{C}_k^{(n)})^{-1} , \quad \text{with } \mathbf{M}_0^{(n)} = \mathbf{I}_p .\tag{S23}$$

Under **H2**, we have $\|\mathbf{C}_k^{(n)}\| \leq \max_{i \in [b]} \{\gamma_i(M_i + 1/\rho_i)\}$, thus if we suppose that for any $i \in [b]$, $0 < \gamma_i < (M_i + 1/\rho_i)^{-1}$, the matrix $(\mathbf{I}_p - \mathbf{C}_k^{(n)})$ is invertible. In addition, for any $n \in \mathbb{N}, k \geq \max_{i \in [b]} \{N_i\}$, $\mathbf{C}_k^{(n)} = \mathbf{0}_{p \times p}$, hence the sequence $(\mathbf{M}_k^{(n)})_{k \in \mathbb{N}}$ is stationary and we denote its limit by $\mathbf{M}_\infty^{(n)}$ which is equal to $\mathbf{M}_{\max_{i \in [b]} \{N_i\}}^{(n)}$.

S2.2.1 Technical lemmata

Similarly to Lemma S3, the following result shows that it is enough to consider the marginal process $(Z_n, \tilde{Z}_n)_{n \geq 0}$ to control

$$W_2(\delta_{\mathbf{x}} P_{\rho, \gamma, N}^n, \delta_{\tilde{\mathbf{x}}} P_{\rho, \gamma, N}^n) \leq \mathbb{E} \left[\|X_n - \tilde{X}_n\|^2 \right]^{1/2} .\tag{S24}$$

Lemma S6. Assume **H1** and let $\mathbf{N} \in (\mathbb{N}^*)^b, \boldsymbol{\gamma} \in (\mathbb{R}_+^*)^b$. Then, for any $n \in \mathbb{N}$, the random variables $X_n = (\theta_n^\top, Z_n^\top)^\top, \tilde{X}_n = (\tilde{\theta}_n^\top, \tilde{Z}_n^\top)^\top$ defined in (S18) satisfy

$$\|\tilde{X}_{n+1} - X_{n+1}\|^2 \leq (1 + \|\bar{\mathbf{B}}_0^{-1} \mathbf{B}_0^\top \tilde{\mathbf{D}}_0^{1/2}\|^2) \|\tilde{Z}_{n+1} - Z_{n+1}\|^2,$$

where $\bar{\mathbf{B}}_0, \mathbf{B}_0, \tilde{\mathbf{D}}_0$ are defined in (S2)-(S3).

Proof. The proof is similar to the proof of Lemma S3 and is omitted. \square

To ease notation, for any $i \in [b]$, we consider all along this section the quantities

$$\tilde{m}_i = m_i + 1/\rho_i, \quad \tilde{M}_i = M_i + 1/\rho_i. \quad (\text{S25})$$

The following lemma provides an explicit expression for $\|\tilde{Z}_{n+1} - Z_{n+1}\|$ with respect to $\|\tilde{Z}_n - Z_n\|$.

Lemma S7. Assume **H1-H2** and let $\mathbf{N} \in (\mathbb{N}^*)^b, \boldsymbol{\gamma} \in (\mathbb{R}_+^*)^b$ such that, for any $i \in [b], \gamma_i < \tilde{M}_i^{-1}$. Then, for any $n \geq 1$, we have

$$\begin{aligned} \|\tilde{Z}_{n+1} - Z_{n+1}\|_{\mathbf{D}_{N\boldsymbol{\gamma}}^{-1}} &\leq \left\| [\mathbf{M}_\infty^{(n)}]^{-1} + \sum_{k=0}^{\infty} [\mathbf{M}_\infty^{(n)}]^{-1} \mathbf{M}_{k+1}^{(n)} \mathbf{J}(k) \mathbf{D}_N^{-1/2} \mathbf{D}_{\boldsymbol{\gamma}/\boldsymbol{\rho}}^{1/2} \mathbf{P}_0 \mathbf{D}_{\boldsymbol{\gamma}/\boldsymbol{\rho}}^{1/2} \mathbf{D}_N^{1/2} \right\| \\ &\quad \times \|\tilde{Z}_n - Z_n\|_{\mathbf{D}_{N\boldsymbol{\gamma}}^{-1}}, \end{aligned} \quad (\text{S26})$$

where $(\mathbf{M}_k^{(n)})_{k \in \mathbb{N}}$ is defined in (S23), $(\tilde{Z}_k, Z_k)_{k \in \mathbb{N}}$ in (S18), $N\boldsymbol{\gamma} = (\gamma_1 N_1, \dots, \gamma_b N_b)$ and $\boldsymbol{\gamma}/\boldsymbol{\rho} = (\gamma_1/\rho_1, \dots, \gamma_b/\rho_b)$.

Proof. Let $n \geq 1$. By (S19), for any $i \in [b], k \in \mathbb{N}$, we obtain

$$\begin{aligned} \tilde{Y}_{k+1}^{(i,n)} - Y_{k+1}^{(i,n)} &= \left(\mathbf{I}_{d_i} - \gamma_i \int_0^1 \nabla^2 V_i((1-s)Y_k^{(i,n)} + s\tilde{Y}_k^{(i,n)}) ds \right) (\tilde{Y}_k^{(i,n)} - Y_k^{(i,n)}) \\ &\quad + (\gamma_i/\rho_i) \mathbf{A}_i (\tilde{\theta}_n - \theta_n). \end{aligned}$$

Consider the process $(\tilde{Y}_k^{(n)}, Y_k^{(n)}) = \{\tilde{Y}_k^{(i,n)}, Y_k^{(i,n)}\}_{i=1}^b\}_{k \in \mathbb{N}}$ with values in $\mathbb{R}^p \times \mathbb{R}^p$ defined for any $i \in [b], k \geq 0$, by

$$\tilde{Y}_k^{(i,n)} = \tilde{Y}_{\min(k, N_i)}^{(i,n)}, \quad Y_k^{(i,n)} = Y_{\min(k, N_i)}^{(i,n)}. \quad (\text{S27})$$

By (S18), we have $\mathbf{A}_i(\tilde{\theta}_n - \theta_n) = \mathbf{A}_i \bar{\mathbf{B}}_0^{-1} \mathbf{B}_0^\top \tilde{\mathbf{D}}_0^{1/2} (\tilde{Z}_n - Z_n)$. Since $\mathbf{B}_0^\top = [\mathbf{A}_1^\top/\rho_1^{1/2} \cdots \mathbf{A}_b^\top/\rho_b^{1/2}]$ and $\mathbf{P}_0 = \mathbf{B}_0 \bar{\mathbf{B}}_0^{-1} \mathbf{B}_0^\top$ is the orthogonal projection matrix defined in (S3), it follows that

$$\tilde{Y}_{k+1}^{(n)} - Y_{k+1}^{(n)} = (\mathbf{I}_p - \mathbf{C}_k^{(n)}) (\tilde{Y}_k^{(n)} - Y_k^{(n)}) + \mathbf{J}(k) \mathbf{D}_{\boldsymbol{\gamma}/\sqrt{\boldsymbol{\rho}}} \mathbf{P}_0 \tilde{\mathbf{D}}_0^{1/2} (\tilde{Y}_0^{(n)} - Y_0^{(n)}). \quad (\text{S28})$$

Since $\mathbf{D}_{\mathbf{N}\gamma}$ commutes with $\mathbf{C}_k^{(n)}$ and $\mathbf{J}(k)$, multiplying (S28) by $\mathbf{M}_{k+1}^{(n)}\mathbf{D}_{\mathbf{N}\gamma}^{-1/2}$, yields

$$\begin{aligned} \mathbf{M}_{k+1}^{(n)}\mathbf{D}_{\mathbf{N}\gamma}^{-1/2}(\tilde{\mathbf{Y}}_{k+1}^{(n)} - \mathbf{Y}_{k+1}^{(n)}) &= \mathbf{M}_k^{(n)}\mathbf{D}_{\mathbf{N}\gamma}^{-1/2}(\tilde{\mathbf{Y}}_k^{(n)} - \mathbf{Y}_k^{(n)}) \\ &\quad + \mathbf{M}_{k+1}^{(n)}\mathbf{J}(k)\mathbf{D}_{\mathbf{N}}^{-1/2}\mathbf{D}_{\gamma/\rho}^{1/2}\mathbf{P}_0\tilde{\mathbf{D}}_0^{1/2}(\tilde{\mathbf{Y}}_0^{(n)} - \mathbf{Y}_0^{(n)}) . \end{aligned} \quad (\text{S29})$$

By definition of the processes in (S18)-(S19) and (S27), we have for $k \geq \max_{i \in [b]} \{N_i\}$, $(\tilde{\mathbf{Y}}_k^{(n)}, \mathbf{Y}_k^{(n)}) = (\tilde{Z}_{n+1}, Z_{n+1})$ and $\mathbf{J}(k) = \mathbf{0}_{p \times p}$. Therefore summing the previous equality (S29) yields

$$\begin{aligned} \mathbf{M}_{\infty}^{(n)}\mathbf{D}_{\mathbf{N}\gamma}^{-1/2}(\tilde{Z}_{n+1} - Z_{n+1}) &= [\mathbf{M}_0^{(n)} + \sum_{k=0}^{\infty} \mathbf{M}_{k+1}^{(n)}\mathbf{J}(k)\mathbf{D}_{\mathbf{N}}^{-1/2}\mathbf{D}_{\gamma/\rho}^{1/2}\mathbf{P}_0\mathbf{D}_{\gamma/\rho}^{1/2}\mathbf{D}_{\mathbf{N}}^{1/2}] \\ &\quad \times \mathbf{D}_{\mathbf{N}\gamma}^{-1/2}(\tilde{\mathbf{Y}}_0^{(n)} - \mathbf{Y}_0^{(n)}) . \end{aligned}$$

Multiplying this last equality by $[\mathbf{M}_{\infty}^{(n)}]^{-1}$ and applying the norm $\|\cdot\|_{\mathbf{D}_{\mathbf{N}\gamma}^{-1}}$ concludes the proof. \square

The three following lemmata aim at providing an explicit upper bound on (S26). To this end, for $n, k \in \mathbb{N}$ and $i \in [b]$, consider $\mathbf{C}_k^{(i,n)}$ corresponding to the i -th diagonal block of $\mathbf{C}_k^{(n)}$ defined in (S22), *i.e.*

$$\mathbf{C}_k^{(i,n)} = \mathbb{1}_{[N_i]}(k+1)\gamma_i \left\{ \rho_i^{-1} \mathbf{I}_{d_i} + \int_0^1 \nabla^2 U_i((1-s)\mathbf{Y}_k^{(i,n)} + s\tilde{\mathbf{Y}}_k^{(i,n)}) \mathrm{d}s \right\} \in \mathbb{R}^{d_i \times d_i} , \quad (\text{S30})$$

where, for any $n \in \mathbb{N}$ and $i \in [b]$, $(\mathbf{Y}_k^{(i,n)}, \tilde{\mathbf{Y}}_k^{(i,n)})_{k \in \mathbb{N}}$ is defined in (S19). Thus, using the definition (S23) of $\mathbf{M}_k^{(n)}$, we can write $[\mathbf{M}_{\infty}^{(n)}]^{-1}\mathbf{M}_k^{(n)}$ as a block-diagonal matrix $\text{diag}([\mathbf{M}_{\infty}^{(n)}]^{-1}\mathbf{M}_k^{(n)})^1, \dots, ([\mathbf{M}_{\infty}^{(n)}]^{-1}\mathbf{M}_k^{(n)})^b$ where for any $i \in [b]$, $([\mathbf{M}_{\infty}^{(n)}]^{-1}\mathbf{M}_k^{(n)})^i = \prod_{l=k}^{N_i-1} (\mathbf{I}_{d_i} - \mathbf{C}_l^{(i,n)}) \in \mathbb{R}^{d_i \times d_i}$.

Lemma S8. Assume **H1-H2** and let $\mathbf{N} \in (\mathbb{R}_+^*)^b$, $\gamma \in (\mathbb{R}_+^*)^b$ such that, for any $i \in [b]$, $\gamma_i < \tilde{M}_i^{-1}$. Then, for any $i \in [b]$, $n \in \mathbb{N}$ and $k \in [N_i]$, we have

$$\|([\mathbf{M}_{\infty}^{(n)}]^{-1}\mathbf{M}_k^{(n)})^i - \mathbf{I}_{d_i} - \sum_{l=k}^{\infty} \mathbf{C}_l^{(i,n)}\| \leq \exp\{(N_i - k)\gamma_i \tilde{M}_i\} - 1 - (N_i - k)\gamma_i \tilde{M}_i ,$$

where $\mathbf{M}_k^{(n)}, \tilde{M}_i$ are defined in (S23), (S25) respectively, and $\mathbf{M}_{\infty}^{(n)}$ is the limit of the stationary sequence $(\mathbf{M}_k^{(n)})_{k \in \mathbb{N}}$.

Proof. Let $n \in \mathbb{N}$, $i \in [b]$ and $k \in [N_i]$. The approximation error between $\prod_{l=k}^{\infty} (\mathbf{I}_{d_i} - \mathbf{C}_l^{(i,n)})$

and its linear approximation can be upper bounded as

$$\begin{aligned}
\left\| \prod_{l=k}^{\infty} (\mathbf{I}_{d_i} - \mathbf{C}_l^{(i,n)}) - \mathbf{I}_{d_i} - \sum_{l=k}^{\infty} \mathbf{C}_l^{(i,n)} \right\| &= \left\| \sum_{m=2}^{\infty} (-1)^m \sum_{k \leq l_1 < \dots < l_m} \mathbf{C}_{l_1}^{(i,n)} \dots \mathbf{C}_{l_m}^{(i,n)} \right\| \\
&\leq \sum_{m=2}^{\infty} \sum_{k \leq l_1 < \dots < l_m} \|\mathbf{C}_{l_1}^{(i,n)}\| \dots \|\mathbf{C}_{l_m}^{(i,n)}\| = \prod_{l=k}^{\infty} (1 + \|\mathbf{C}_l^{(i,n)}\|) - 1 - \sum_{l \geq k} \|\mathbf{C}_l^{(i,n)}\| \\
&\leq \exp \left(\sum_{l=k}^{\infty} \|\mathbf{C}_l^{(i,n)}\| \right) - 1 - \sum_{l=k}^{\infty} \|\mathbf{C}_l^{(i,n)}\| ,
\end{aligned}$$

where the products and the sums are well defined since for any $l \geq N_i$, we have $\mathbf{C}_l^{(i,n)} = \mathbf{0}_{d_i}$. Finally, the proof is concluded using that $x \mapsto \exp(x) - 1 - x$ is increasing on \mathbb{R} and for $l \in \mathbb{N}$, $\|\mathbf{C}_l^{(i,n)}\| \leq \gamma_i \tilde{M}_i \mathbf{1}_{[N_i]}(l+1)$ from **H2-(i)**. \square

For any $\mathbf{N} = (N_1, \dots, N_b) \in (\mathbb{N}^*)^b$, $\boldsymbol{\gamma} = (\gamma_1, \dots, \gamma_b) \in (\mathbb{R}_+^*)^b$, define the $p \times p$ block matrices

$$\begin{aligned}
\mathbf{S}_1 &= \text{diag}(\{1 - N_1 \gamma_1 \tilde{M}_1\} \mathbf{I}_{d_1}, \dots, \{1 - N_b \gamma_b \tilde{M}_b\} \mathbf{I}_{d_b}) , \\
\mathbf{S}_2 &= \mathbf{I}_p - \sum_{l=0}^{\infty} \mathbf{J}(l) \mathbf{H}_{U,l}^{(n)} - (\mathbf{D}_N \mathbf{D}_{\boldsymbol{\gamma}/\rho})^{1/2} (\mathbf{I}_p - \mathbf{P}_0) (\mathbf{D}_N \mathbf{D}_{\boldsymbol{\gamma}/\rho})^{1/2} , \\
\mathbf{S}_3 &= \text{diag}(\{1 - N_1 \gamma_1 m_1\} \mathbf{I}_{d_1}, \dots, \{1 - N_b \gamma_b m_b\} \mathbf{I}_{d_b}) ,
\end{aligned} \tag{S31}$$

where for any $i \in [b]$, \tilde{M}_i is defined in (S25) and $\mathbf{P}_0, \mathbf{J}(l), \mathbf{H}_{U,l}^{(n)}$ are defined in (S3), (S86), (S87), respectively.

Lemma S9. Assume **H1-H2**. Then, for any $\mathbf{N} \in (\mathbb{N}^*)^b$, $\boldsymbol{\gamma} \in (\mathbb{R}_+^*)^b$, we have

$$\mathbf{S}_1 \preceq \mathbf{S}_2 \preceq \mathbf{S}_3 .$$

As a result, under the additional assumption, for any $i \in [b]$, $\gamma_i N_i \leq 2/(m_i + M_i + 1/\rho_i)$, we get

$$\|\mathbf{S}_2\| \leq 1 - \min_{i \in [b]} \{N_i \gamma_i m_i\} . \tag{S32}$$

Proof. Since \mathbf{P}_0 is an orthogonal projection defined in (S3), we have $\mathbf{P}_0 \preceq \mathbf{I}_p$, therefore we easily get

$$\mathbf{0}_{p \times p} \preceq (\mathbf{D}_N \mathbf{D}_{\boldsymbol{\gamma}/\rho})^{1/2} (\mathbf{I}_p - \mathbf{P}_0) (\mathbf{D}_N \mathbf{D}_{\boldsymbol{\gamma}/\rho})^{1/2} \preceq \mathbf{D}_N \mathbf{D}_{\boldsymbol{\gamma}/\rho}$$

and **H2-(i)-(ii)** imply

$$\text{diag}(N_1 \gamma_1 m_1 \mathbf{I}_{d_1}, \dots, N_b \gamma_b m_b \mathbf{I}_{d_b}) \preceq \sum_{l=0}^{\infty} \mathbf{J}(l) \mathbf{H}_{U,l}^{(n)} \preceq \text{diag}(N_1 \gamma_1 M_1 \mathbf{I}_{d_1}, \dots, N_b \gamma_b M_b \mathbf{I}_{d_b}) .$$

Substracting these previous inequalities and adding \mathbf{I}_p complete the first part of the proof. The additional condition, for any $i \in [b]$, $\gamma_i N_i \leq 2/(m_i + M_i + 1/\rho_i)$, ensures that \mathbf{S}_1 is definite-positive. Since $\mathbf{S}_1 \preceq \mathbf{S}_2$, we deduce that \mathbf{S}_2 is symmetric positive-definite as well. Then, $\|\mathbf{S}_2\|$ is equal to the largest eigenvalue of \mathbf{S}_2 . The inequality $\mathbf{S}_2 \preceq \mathbf{S}_3$ concludes the second part of the proof. \square

For any $\mathbf{N} = (N_1, \dots, N_b) \in (\mathbb{N}^*)^b$, $\boldsymbol{\gamma} = (\gamma_1, \dots, \gamma_b) \in (\mathbb{R}_+^*)^b$, define

$$r_{\boldsymbol{\gamma}, \boldsymbol{\rho}, \mathbf{N}} = \max_{i \in [b]} \{N_i \gamma_i / \rho_i\} \max_{i \in [b]} \{N_i \gamma_i \tilde{M}_i\} \left(1/2 + \max_{i \in [b]} \{N_i \gamma_i \tilde{M}_i\}\right) + 4 \max_{i \in [b]} \{N_i \gamma_i \tilde{M}_i\}^2, \quad (\text{S33})$$

where \tilde{M}_i is defined in (S25).

Lemma S10. Assume **H 1-H 2**. Let $\mathbf{N} \in (\mathbb{N}^*)^b$, $\boldsymbol{\gamma} \in (\mathbb{R}_+^*)^b$ such that, for any $i \in [b]$, $N_i \gamma_i \leq 2/(m_i + \tilde{M}_i)$ and $\gamma_i < \tilde{M}_i^{-1}$. Then, for any $n \in \mathbb{N}$, we have

$$\|[\mathbf{M}_\infty^{(n)}]^{-1} + \sum_{k=0}^{\infty} [\mathbf{M}_\infty^{(n)}]^{-1} \mathbf{M}_{k+1}^{(n)} \mathbf{J}(k) \mathbf{D}_N^{-1/2} \mathbf{D}_{\boldsymbol{\gamma}/\boldsymbol{\rho}}^{1/2} \mathbf{P}_0 \mathbf{D}_{\boldsymbol{\gamma}/\boldsymbol{\rho}}^{1/2} \mathbf{D}_N^{1/2}\| \leq 1 - \min_{i \in [b]} \{N_i \gamma_i m_i\} + r_{\boldsymbol{\gamma}, \boldsymbol{\rho}, \mathbf{N}},$$

where \mathbf{P}_0 , $\mathbf{D}_{\boldsymbol{\gamma}/\boldsymbol{\rho}}$, $\mathbf{J}(k)$, $\mathbf{M}_k^{(n)}$ and $r_{\boldsymbol{\gamma}, \boldsymbol{\rho}, \mathbf{N}}$ are defined in (S3), (S21), (S23) and (S33), respectively.

Proof. Let $n \in \mathbb{N}$. For any $k \in \mathbb{N}$, define

$$\mathbf{R}_k^{(n)} = \prod_{l=k}^{\infty} (\mathbf{I}_p - \mathbf{C}_l^{(n)}) - \mathbf{I}_p + \sum_{l=k}^{\infty} \mathbf{C}_l^{(n)}, \quad \mathbf{R}_k^{(i,n)} = \prod_{l=k}^{\infty} (\mathbf{I}_{d_i} - \mathbf{C}_l^{(i,n)}) - \mathbf{I}_{d_i} + \sum_{l=k}^{\infty} \mathbf{C}_l^{(i,n)}, \quad i \in [b], \quad (\text{S34})$$

where $(\mathbf{C}_l^{(i,n)})_{l \in \mathbb{N}}$ is defined in (S30) and remark that the products and the sums are well defined since for any $l \geq N_i$, we have $\mathbf{C}_l^{(i,n)} = \mathbf{0}_{d_i}$. By noting, for any $k \in [\max_{i \in [b]} N_i]$, that $[\mathbf{M}_\infty^{(n)}]^{-1} \mathbf{M}_k^{(n)} = \prod_{l=k}^{\infty} (\mathbf{I}_p - \mathbf{C}_l^{(n)})$, it follows that $[\mathbf{M}_\infty^{(n)}]^{-1} \mathbf{M}_k^{(n)} = \mathbf{I}_p - \sum_{l=k}^{\infty} \mathbf{C}_l^{(n)} + \mathbf{R}_k^{(n)}$. Since for any $i \in [b]$, $l \geq N_i$, $\mathbf{R}_k^{(i,n)} = \mathbf{0}_{d_i}$, thus we have $\mathbf{J}(k) \mathbf{R}_{k+1}^{(n)} = \mathbf{R}_{k+1}^{(n)}$. In addition, using that $\mathbf{M}_0^{(n)} = \mathbf{I}_p$, $\mathbf{C}_l^{(n)} = \mathbf{J}(l)(\mathbf{D}_{\boldsymbol{\gamma}/\boldsymbol{\rho}} + \mathbf{H}_{U,l}^{(n)})$, $\mathbf{D}_N = \sum_{k=0}^{\infty} \mathbf{J}(k)$, $\mathbf{D}_N \mathbf{C}_l^{(n)} = \mathbf{C}_l^{(n)} \mathbf{D}_N$,

we get

$$\begin{aligned}
& [\mathbf{M}_\infty^{(n)}]^{-1} + \sum_{k=0}^{\infty} [\mathbf{M}_\infty^{(n)}]^{-1} \mathbf{M}_{k+1}^{(n)} \mathbf{J}(k) \mathbf{D}_N^{-1/2} \mathbf{D}_{\gamma/\rho}^{1/2} \mathbf{P}_0 \mathbf{D}_{\gamma/\rho}^{1/2} \mathbf{D}_N^{1/2} \\
&= \mathbf{I}_p - \sum_{l=0}^{\infty} \mathbf{C}_l^{(n)} + \sum_{k=0}^{\infty} \mathbf{J}(k) \mathbf{D}_N^{-1/2} \mathbf{D}_{\gamma/\rho}^{1/2} \mathbf{P}_0 \mathbf{D}_{\gamma/\rho}^{1/2} \mathbf{D}_N^{1/2} - \sum_{k=0}^{\infty} \sum_{l=k+1}^{\infty} \mathbf{J}(k) \mathbf{D}_N^{-1/2} \mathbf{C}_l^{(n)} \mathbf{D}_{\gamma/\rho}^{1/2} \mathbf{P}_0 \mathbf{D}_{\gamma/\rho}^{1/2} \mathbf{D}_N^{1/2} \\
&\quad + \mathbf{R}_0^{(n)} + \sum_{k=0}^{\infty} \mathbf{R}_{k+1}^{(n)} \mathbf{J}(k) \mathbf{D}_N^{-1/2} \mathbf{D}_{\gamma/\rho}^{1/2} \mathbf{P}_0 \mathbf{D}_{\gamma/\rho}^{1/2} \mathbf{D}_N^{1/2} \\
&= \mathbf{I}_p - \sum_{l=0}^{\infty} \mathbf{J}(l) \mathbf{H}_{U,l}^{(n)} - \left(\sum_{k=0}^{\infty} \mathbf{J}(k) \right) \mathbf{D}_N^{-1/2} \mathbf{D}_{\gamma/\rho}^{1/2} (\mathbf{I}_p - \mathbf{P}_0) \mathbf{D}_{\gamma/\rho}^{1/2} \mathbf{D}_N^{1/2} \\
&\quad - \sum_{l=1}^{\infty} \left(\sum_{k=0}^{l-1} \mathbf{J}(k) \right) \mathbf{D}_N^{-1/2} \mathbf{C}_l^{(n)} \mathbf{D}_{\gamma/\rho}^{1/2} \mathbf{P}_0 \mathbf{D}_{\gamma/\rho}^{1/2} \mathbf{D}_N^{1/2} + \mathbf{R}_0^{(n)} + \sum_{k=0}^{\infty} \mathbf{J}(k) \mathbf{D}_N^{-1/2} \mathbf{R}_{k+1}^{(n)} \mathbf{D}_{\gamma/\rho}^{1/2} \mathbf{P}_0 \mathbf{D}_{\gamma/\rho}^{1/2} \mathbf{D}_N^{1/2} \\
&= \mathbf{S}_2 - \sum_{l=1}^{\infty} \left(\sum_{k=0}^{l-1} \mathbf{J}(k) \right) \mathbf{D}_N^{-1} \mathbf{C}_l^{(n)} (\mathbf{D}_N \mathbf{D}_{\gamma/\rho})^{1/2} \mathbf{P}_0 (\mathbf{D}_{\gamma/\rho} \mathbf{D}_N)^{1/2} \\
&\quad + \mathbf{R}_0^{(n)} + \sum_{k=1}^{\infty} \mathbf{D}_N^{-1} \mathbf{R}_k^{(n)} (\mathbf{D}_N \mathbf{D}_{\gamma/\rho})^{1/2} \mathbf{P}_0 (\mathbf{D}_{\gamma/\rho} \mathbf{D}_N)^{1/2}, \tag{S35}
\end{aligned}$$

where \mathbf{S}_2 is defined in (S31). We now bound the different terms of (S35) separately. First, using (S32), we have

$$\|\mathbf{S}_2\| \leq 1 - \min_{i \in [b]} \{N_i \gamma_i m_i\}. \tag{S36}$$

By recalling $\mathbf{R}_0^{(n)}$ defined in (S34), Lemma S8 shows that

$$\|\mathbf{R}_0^{(n)}\| \leq \max_{i \in [b]} \|\mathbf{R}_0^{(i,n)}\| = \max_{i \in [b]} \left\{ \left\| \prod_{l=0}^{\infty} (\mathbf{I}_{d_i} - \mathbf{C}_l^{(i,n)}) - \mathbf{I}_{d_i} - \sum_{l=0}^{\infty} \mathbf{C}_l^{(i,n)} \right\| \right\} \tag{S37}$$

$$\leq \max_{i \in [b]} \left\{ \exp \left(\sum_{l=0}^{\infty} \|\mathbf{C}_l^{(i,n)}\| \right) - 1 - \sum_{l=0}^{\infty} \|\mathbf{C}_l^{(i,n)}\| \right\} \tag{S38}$$

$$\leq \max_{i \in [b]} \left\{ \exp\{(N_i - 1)\gamma_i \tilde{M}_i\} - 1 - (N_i - 1)\gamma_i \tilde{M}_i \right\} \tag{S39}$$

$$\leq \max_{i \in [b]} \left\{ ((N_i - 1)\gamma_i \tilde{M}_i)^2 e^{(N_i - 1)\gamma_i \tilde{M}_i} \right\} / 2 \tag{S40}$$

$$\leq 4 \max_{i \in [b]} \{(N_i - 1)\gamma_i \tilde{M}_i\}^2, \tag{S41}$$

where, in the penultimate line, we used for any $t \geq 0$, that $\exp(t) - 1 - t \leq t^2 \exp(t)/2$.

Regarding the second term of (S35), using that \mathbf{P}_0 is an orthogonal projector, we get

$$\left\| \sum_{l=1}^{\infty} \left(\sum_{k=0}^{l-1} \mathbf{J}(k) \right) \mathbf{D}_N^{-1} \mathbf{C}_l^{(n)} (\mathbf{D}_N \mathbf{D}_{\gamma/\rho})^{1/2} \mathbf{P}_0 (\mathbf{D}_N \mathbf{D}_{\gamma/\rho})^{1/2} \right\|$$

$$\leq \max_{i \in [b]} \left(\frac{N_i \gamma_i}{\rho_i} \right) \left\| \sum_{l=1}^{\infty} \left(\sum_{k=0}^{l-1} \mathbf{J}(k) \right) \mathbf{D}_N^{-1} \mathbf{C}_l^{(n)} \right\|.$$

Combining the following upper bound

$$\left\| \sum_{l=1}^{\infty} \left(\sum_{k=0}^{l-1} \mathbf{J}(k) \right) \mathbf{D}_N^{-1} \mathbf{C}_l^{(n)} \right\| \leq \max_{i \in [b]} \left\{ \frac{1}{N_i} \sum_{l=1}^{\infty} l \|\mathbf{C}_l^{(i,n)}\| \right\}$$

with the fact, for any $i \in [b]$, that $\|\mathbf{C}_l^{(i,n)}\| \leq \gamma_i \tilde{M}_i \mathbf{1}_{[N_i]}(l+1)$, we get that

$$\left\| \sum_{l=1}^{\infty} \left(\sum_{k=0}^{l-1} \mathbf{J}(k) \right) \mathbf{D}_N^{-1} \mathbf{C}_l^{(n)} (\mathbf{D}_N \mathbf{D}_{\gamma/\rho})^{1/2} \mathbf{P}_0 (\mathbf{D}_N \mathbf{D}_{\gamma/\rho})^{1/2} \right\|$$

$$\leq \max_{i \in [b]} \left(\frac{N_i \gamma_i}{\rho_i} \right) \max_{i \in [b]} \left\{ \frac{N_i \gamma_i \tilde{M}_i}{2} \right\}. \quad (\text{S42})$$

To upper bound the last term of (S35), we start from the following inequality

$$\left\| \sum_{k=1}^{\infty} \mathbf{D}_N^{-1} \mathbf{R}_k^{(n)} \right\| \leq \max_{i \in [b]} \left\{ \frac{1}{N_i} \sum_{k=1}^{N_i-1} \|\mathbf{R}_k^{(i,n)}\| \right\}.$$

Lemma S8 shows that for any $k \in [N_i - 1]$ and $i \in [b]$, $\|\mathbf{R}_k^{(i,n)}\| \leq \exp\{(N_i - k)\gamma_i \tilde{M}_i\} - 1 - (N_i - k)\gamma_i \tilde{M}_i$. Then, for any $i \in [b]$, we have

$$\frac{1}{N_i} \sum_{k=1}^{N_i-1} \|\mathbf{R}_k^{(i,n)}\| \leq \frac{1}{N_i} \sum_{k=1}^{N_i-1} [\exp\{(N_i - k)\gamma_i \tilde{M}_i\} - 1 - (N_i - k)\gamma_i \tilde{M}_i]$$

$$\leq (N_i \gamma_i \tilde{M}_i)^{-1} \int_0^{N_i \gamma_i \tilde{M}_i} (e^t - 1 - t) dt \leq \frac{(N_i \gamma_i \tilde{M}_i)^2}{12} (e^{N_i \gamma_i \tilde{M}_i} + 1)$$

$$\leq \max_{i \in [b]} \{(N_i \gamma_i \tilde{M}_i)^2\}, \quad (\text{S43})$$

where we have used $e^2 + 1 \leq 12$. Plugging (S43), (S42), (S41) into (S32), we get

$$\left\| [\mathbf{M}_{\infty}^{(n)}]^{-1} + \sum_{k \in \mathbb{N}} [\mathbf{M}_{\infty}^{(n)}]^{-1} \mathbf{M}_{k+1}^{(n)} \mathbf{J}(k) \mathbf{D}_N^{-1/2} \mathbf{D}_{\gamma/\rho}^{1/2} \mathbf{P}_0 \mathbf{D}_{\gamma/\rho}^{1/2} \mathbf{D}_N^{1/2} \right\| \leq 1 - \min_{i \in [b]} \{N_i \gamma_i m_i\} + r_{\gamma, \rho, N},$$

where $r_{\gamma, \rho, N}$ is defined in (S33). \square

Lemma S11. Assume **H1-H2**. Let $\mathbf{N} \in (\mathbb{N}^*)^b, \gamma \in (\mathbb{R}_+^*)^b$ such that, for any $i \in [b]$, $N_i \gamma_i \leq 2/(m_i + \tilde{M}_i)$ and $\gamma_i < \tilde{M}_i^{-1}$. Then, for any $\mathbf{x} = (\mathbf{z}^\top, \boldsymbol{\theta}^\top)^\top, \tilde{\mathbf{x}} = (\tilde{\mathbf{z}}^\top, \tilde{\boldsymbol{\theta}}^\top)^\top \in \mathbb{R}^{p+d}$, with $(\boldsymbol{\theta}, \tilde{\boldsymbol{\theta}}) \in (\mathbb{R}^d)^2, (\mathbf{z}, \tilde{\mathbf{z}}) \in (\mathbb{R}^p)^2$ and any $n \geq 1$ we have

$$W_2^2(\delta_{\tilde{\mathbf{x}}} P_{\rho, \gamma, \mathbf{N}}^n, \delta_{\mathbf{x}} P_{\rho, \gamma, \mathbf{N}}^n) \leq (1 - \min_{i \in [b]} \{N_i \gamma_i m_i\} + r_{\gamma, \rho, \mathbf{N}})^{2n-2} (1 + \|\bar{\mathbf{B}}_0^{-1} \mathbf{B}_0^\top \tilde{\mathbf{D}}_0^{1/2}\|^2) \\ \times \frac{\max_{i \in [b]} \{N_i \gamma_i\}}{\min_{i \in [b]} \{N_i \gamma_i\}} \left[\|\mathbf{M}_\infty^{(0)}\|^{-1} \|\tilde{\mathbf{z}} - \mathbf{z}\| + (\sum_{i \in [b]} \|\mathbf{A}_i\|/\rho_i) \|\tilde{\boldsymbol{\theta}} - \boldsymbol{\theta}\| \right]^2,$$

where $\mathbf{B}_0, \bar{\mathbf{B}}_0, \tilde{\mathbf{D}}_0, P_{\rho, \gamma, \mathbf{N}}, \mathbf{M}_\infty^{(0)}, r_{\gamma, \rho, \mathbf{N}}$ are defined in (S2), (S3), (S17), (S23), (S33), respectively.

Proof. Combining Lemma S7 and Lemma S10, we have for $n \geq 1$,

$$\|\tilde{Z}_{n+1} - Z_{n+1}\|_{\mathbf{D}_{\mathbf{N}\gamma}^{-1}} \leq (1 - \min_{i \in [b]} \{N_i \gamma_i m_i\} + r_{\gamma, \rho, \mathbf{N}}) \|\tilde{Z}_n - Z_n\|_{\mathbf{D}_{\mathbf{N}\gamma}^{-1}}.$$

Thereby, for any $n \geq 1$, we obtain by induction

$$\|\tilde{Z}_n - Z_n\|_{\mathbf{D}_{\mathbf{N}\gamma}^{-1}} \leq (1 - \min_{i \in [b]} \{N_i \gamma_i m_i\} + r_{\gamma, \rho, \mathbf{N}})^{n-1} \|\tilde{Z}_1 - Z_1\|_{\mathbf{D}_{\mathbf{N}\gamma}^{-1}}. \quad (\text{S44})$$

Define the process $((\tilde{Y}_k^{(0)}, Y_k^{(0)}) = \{\tilde{Y}_k^{(i,0)}, Y_k^{(i,0)}\}_{i=1}^b)_{k \in \mathbb{N}}$ with values in $\mathbb{R}^p \times \mathbb{R}^p$ defined for any $i \in [b], k \geq 0$ by

$$\tilde{Y}_k^{(i,0)} = \tilde{Y}_{\min(k, N_i)}^{(i,0)}, \quad Y_k^{(i,0)} = Y_{\min(k, N_i)}^{(i,0)}.$$

By (S18), it follows that for any $i \in [b]$, $(\tilde{Z}_1^i, Z_1^i) = (\tilde{Y}_{N_i}^{(i,0)}, Y_{N_i}^{(i,0)})$ where $(\tilde{Y}_0^{(i,0)}, Y_0^{(i,0)}) = (\tilde{Z}_0^i, Z_0^i)$. We get by (S19) for $k \geq 0$,

$$\tilde{Y}_{k+1}^{(0)} - Y_{k+1}^{(0)} = (\mathbf{I}_p - \mathbf{C}_k^{(0)}) (\tilde{Y}_k^{(0)} - Y_k^{(0)}) + \mathbf{J}(k) \mathbf{D}_{\gamma/\sqrt{\rho}} \mathbf{B}_0 (\tilde{\boldsymbol{\theta}}_0 - \boldsymbol{\theta}_0).$$

Hence, for $k \geq 0$, we obtain

$$\mathbf{M}_{k+1}^{(0)} \mathbf{D}_{\mathbf{N}\gamma}^{-1/2} (\tilde{Y}_{k+1}^{(0)} - Y_{k+1}^{(0)}) = \mathbf{M}_k^{(0)} \mathbf{D}_{\mathbf{N}\gamma}^{-1/2} (\tilde{Y}_k^{(0)} - Y_k^{(0)}) + \mathbf{M}_{k+1}^{(0)} \mathbf{J}(k) \mathbf{D}_{\mathbf{N}}^{-1/2} \mathbf{D}_{\gamma/\rho}^{1/2} \mathbf{B}_0 (\tilde{\boldsymbol{\theta}}_0 - \boldsymbol{\theta}_0).$$

Summing the previous equality gives

$$\mathbf{M}_\infty^{(0)} \mathbf{D}_{\mathbf{N}\gamma}^{-1/2} (\tilde{Y}_N^{(0)} - Y_N^{(0)}) = \mathbf{M}_0^{(0)} \mathbf{D}_{\mathbf{N}\gamma}^{-1/2} (\tilde{Y}_0^{(0)} - Y_0^{(0)}) + \sum_{k=0}^{\infty} \mathbf{M}_{k+1}^{(0)} \mathbf{J}(k) \mathbf{D}_{\mathbf{N}}^{-1/2} \mathbf{D}_{\gamma/\rho}^{1/2} \mathbf{B}_0 (\tilde{\boldsymbol{\theta}}_0 - \boldsymbol{\theta}_0).$$

Multiplying by $[\mathbf{M}_\infty^{(0)}]^{-1}$ and using the fact that $(\boldsymbol{\theta}_0, Y_0^{(0)}) = (\boldsymbol{\theta}, \mathbf{z})$, $(\tilde{\boldsymbol{\theta}}_0, \tilde{Y}_0^{(0)}) = (\tilde{\boldsymbol{\theta}}, \tilde{\mathbf{z}})$, we get

$$\mathbf{D}_{\mathbf{N}\gamma}^{-1/2} (\tilde{Z}_1 - Z_1) = [\mathbf{M}_\infty^{(0)}]^{-1} \mathbf{D}_{\mathbf{N}\gamma}^{-1/2} (\tilde{\mathbf{z}} - \mathbf{z}) + \sum_{k=0}^{\infty} [\mathbf{M}_\infty^{(0)}]^{-1} \mathbf{M}_{k+1}^{(0)} \mathbf{J}(k) \mathbf{D}_{\mathbf{N}}^{-1/2} \mathbf{D}_{\gamma/\rho}^{1/2} \mathbf{B}_0 (\tilde{\boldsymbol{\theta}} - \boldsymbol{\theta}).$$

Plugging the result in (S52) implies for any $n \geq 1$,

$$\begin{aligned} \|\tilde{Z}_n - Z_n\|_{\mathbf{D}_{N\gamma}^{-1}} &\leq (1 - \min_{i \in [b]} \{N_i \gamma_i m_i\} + r_{\gamma, \rho, N})^{n-1} \left[\|[\mathbf{M}_\infty^{(0)}]^{-1}\| \|\tilde{\mathbf{z}} - \mathbf{z}\|_{\mathbf{D}_{N\gamma}^{-1}} \right. \\ &\quad \left. + \left\| \sum_{k=0}^{\infty} [\mathbf{M}_\infty^{(0)}]^{-1} \mathbf{M}_{k+1}^{(0)} \mathbf{J}(k) \mathbf{D}_N^{-1/2} \mathbf{D}_{\gamma/\rho}^{1/2} \mathbf{B}_0 \right\| \|\tilde{\boldsymbol{\theta}} - \boldsymbol{\theta}\| \right]. \quad (\text{S45}) \end{aligned}$$

By **H2-(ii)** and the definitions of $\mathbf{C}_l^{(0)}, \mathbf{M}_k^{(0)}$ given in (S22), (S23), we have $\|\mathbf{I}_{d_i} - \mathbf{C}_l^{(i,0)}\| \leq 1 - \gamma_i \tilde{m}_i$. As a result and since $([\mathbf{M}_\infty^{(0)}]^{-1} \mathbf{M}_k^{(0)})^i = \prod_{l=0}^{k-1} (\mathbf{I}_{d_i} - \mathbf{C}_l^{(i,0)})$, the triangle inequality implies

$$\begin{aligned} \left\| \sum_{k=0}^{\infty} [\mathbf{M}_\infty^{(0)}]^{-1} \mathbf{M}_{k+1}^{(0)} \mathbf{J}(k) \mathbf{D}_N^{-1/2} \mathbf{D}_{\gamma/\rho}^{1/2} \mathbf{B}_0 \right\| &\leq \sum_{i \in [b]} \sqrt{\gamma_i / N_i} (\|\mathbf{A}_i\| / \rho_i) \sum_{k=1}^{N_i} \|([\mathbf{M}_\infty^{(0)}]^{-1} \mathbf{M}_k^{(0)})^i\| \\ &\leq \sum_{i \in [b]} \sqrt{\gamma_i / N_i} (\|\mathbf{A}_i\| / \rho_i) \sum_{k=0}^{N_i-1} (1 - \gamma_i \tilde{m}_i)^k \\ &\leq \sum_{i \in [b]} \|\mathbf{A}_i\| \sqrt{N_i \gamma_i / \rho_i}. \end{aligned}$$

Plugging this result in (S45), we get

$$\begin{aligned} \|\tilde{Z}_n - Z_n\|_{\mathbf{D}_{N\gamma}^{-1}} &\leq (1 - \min_{i \in [b]} \{N_i \gamma_i m_i\} + r_{\gamma, \rho, N})^{n-1} \left[\|[\mathbf{M}_\infty^{(0)}]^{-1}\| \|\tilde{\mathbf{z}} - \mathbf{z}\|_{\mathbf{D}_{N\gamma}^{-1}} \right. \\ &\quad \left. + \left(\sum_{i \in [b]} \|\mathbf{A}_i\| \sqrt{N_i \gamma_i / \rho_i} \right) \|\tilde{\boldsymbol{\theta}} - \boldsymbol{\theta}\| \right]. \end{aligned}$$

Finally, Lemma S6 gives

$$\begin{aligned} \|\tilde{X}_n - X_n\|^2 &\leq (1 - \min_{i \in [b]} \{N_i \gamma_i m_i\} + r_{\gamma, \rho, N})^{2n-2} \cdot (1 + \|\bar{\mathbf{B}}_0^{-1} \mathbf{B}_0^\top \tilde{\mathbf{D}}_0^{1/2}\|^2) \frac{\max_{i \in [b]} \{N_i \gamma_i\}}{\min_{i \in [b]} \{N_i \gamma_i\}} \\ &\quad \times \left[\|[\mathbf{M}_\infty^{(0)}]^{-1}\| \|\tilde{\mathbf{z}} - \mathbf{z}\| + \left(\sum_{i \in [b]} \|\mathbf{A}_i\| / \rho_i \right) \|\tilde{\boldsymbol{\theta}} - \boldsymbol{\theta}\| \right]^2. \end{aligned}$$

Plugging this result into (S24) concludes the proof. \square

The following result gives a condition on $\max_{i \in [b]} \{N_i \gamma_i\}$ to simplify the contracting term in Lemma S11 to $1 - \min_{i \in [b]} \{N_i \gamma_i m_i\} / 2$. To this end, define

$$\begin{aligned} A_0 &= \max_{i \in [b]} \{\tilde{M}_i\} \max_{i \in [b]} \{1 / \rho_i\} / 2 + 4 \max_{i \in [b]} \{\tilde{M}_i\}^2, \\ A_1 &= \max_{i \in [b]} \{\tilde{M}_i\}^2 \max_{i \in [b]} \{1 / \rho_i\}. \end{aligned}$$

Lemma S12. Assume **H1-H2** and let $c \in \mathbb{R}_+^*$, $\mathbf{N} \in (\mathbb{N}^*)^b$, $\gamma \in (\mathbb{R}_+^*)^b$ such that

$$\begin{aligned} \min_{i \in [b]} \{N_i \gamma_i\} / \max_{i \in [b]} \{N_i \gamma_i\} &\geq c, \\ \max_{i \in [b]} \{N_i \gamma_i\} &\leq \frac{c \min_{i \in [b]} \{m_i\}}{2A_0 + \sqrt{2A_1 c \min_{i \in [b]} \{m_i\}}} \wedge \frac{2}{\max_{i \in [b]} \{m_i + M_i + 1/\rho_i\}}. \end{aligned} \quad (\text{S46})$$

Then, $1 - \min_{i \in [b]} \{N_i \gamma_i m_i\} + r_{\gamma, \rho, \mathbf{N}} < 1 - \min_{i \in [b]} \{N_i \gamma_i m_i\} / 2 < 1$, where $r_{\gamma, \rho, \mathbf{N}}$ is defined in (S33).

Proof. The proof is straightforward solving a second order polynomial inequality and using for any $a, b \in \mathbb{R}_+^*$, $a + \frac{b^2}{2a+b} \leq \sqrt{a^2 + b^2}$. \square

S2.2.2 Proof of Proposition 2

The next proposition quantifies the convergence of $\delta_{\mathbf{x}} P_{\rho, \gamma, \mathbf{N}}^n$ towards $\Pi_{\rho, \gamma}$ in $(\mathcal{P}_2(\mathbb{R}^d), W_2)$, where $\Pi_{\rho, \gamma}$ is the stationnary distribution derived in Proposition S5. In addition, it generalises and gives a more formal statement than Proposition 2.

Proposition S13. Assume **H1-H2** and let $c > 0$ and $\gamma = \{\gamma_i\}_{i=1}^b$, $\mathbf{N} \in (\mathbb{N}^*)^b$ such that (S46) is satisfied, for any $i \in [b]$, $N_i \gamma_i < 2 / \max_{i \in [b]} \{m_i + \tilde{M}_i\}$ and $\gamma_i < \tilde{M}_i^{-1}$. Then, $P_{\rho, \gamma, \mathbf{N}}$ defined in (S17) admits a unique invariant probability measure $\Pi_{\rho, \gamma, \mathbf{N}}$. In addition, for any $\mathbf{x} = (\mathbf{z}^\top, \boldsymbol{\theta}^\top)^\top$ with $(\boldsymbol{\theta}, \mathbf{z}) \in \mathbb{R}^d \times \mathbb{R}^p$, any integer $n \geq 1$, we have

$$\begin{aligned} W_2^2(\delta_{\mathbf{x}} P_{\rho, \gamma, \mathbf{N}}^n, \Pi_{\rho, \gamma}) &\leq (1 - \min_{i \in [b]} \{N_i \gamma_i m_i\} / 2)^{2n-2} \cdot (1 + \|\bar{\mathbf{B}}_0^{-1} \mathbf{B}_0^\top \tilde{\mathbf{D}}_0^{1/2}\|^2) \frac{\max_{i \in [b]} \{N_i \gamma_i\}}{\min_{i \in [b]} \{N_i \gamma_i\}} \\ &\quad \times \int_{\mathbb{R}^d \times \mathbb{R}^p} \left[\|\mathbf{M}_\infty^{(0)}\|^{-1} \|\tilde{\mathbf{z}} - \mathbf{z}\| + \left(\sum_{i \in [b]} \|\mathbf{A}_i\| / \rho_i \right) \|\tilde{\boldsymbol{\theta}} - \boldsymbol{\theta}\| \right]^2 d\Pi_{\rho, \gamma}(\tilde{\mathbf{x}}), \end{aligned}$$

where $\mathbf{B}_0, \bar{\mathbf{B}}_0, \mathbf{M}_\infty^{(0)}$ are defined in (S2), (S23), respectively.

Finally, if $\mathbf{N} = N(1, \dots, 1) = N\mathbf{1}_b$ for $N \geq 1$, then $\Pi_{\rho, \gamma, \mathbf{N}} = \Pi_{\rho, \gamma, \mathbf{1}_b}$.

Proof. Note that under the conditions on γ and \mathbf{N} stated in Proposition S13, Lemma S12 ensures that $1 - \min_{i \in [b]} \{N_i \gamma_i m_i\} / 2 < 1$. Then, from Lemma S11 and Douc et al. (2018, Lemma 20.3.2, Theorem 20.3.4), we deduce the existence and unicity of a stationary distribution $\Pi_{\rho, \gamma, \mathbf{N}}$ for $P_{\rho, \gamma, \mathbf{N}}$. The proof is concluded by using the upper bound given in Lemma S11.

We now show the last statement and assume that $\mathbf{N} = N\mathbf{1}_b$, for $N \geq 1$. By Proposition S5, we have the existence and unicity of a stationary distribution $\Pi_{\rho, \gamma, \mathbf{1}_b}$ which is invariant for $P_{\rho, \gamma}$ defined in (S7). For ease of notation, we simply denote $\Pi_{\rho, \gamma, \mathbf{1}_b}$ by $\Pi_{\rho, \gamma}$. We now show that $\Pi_{\rho, \gamma}$ is also invariant for $P_{\rho, \gamma, \mathbf{N}}$ defined in (S17). Using the fact that

$P_{\rho, \gamma}$ defined in (S7) leaves $\Pi_{\rho, \gamma}$ invariant from Proposition S5 and Fubini's theorem, we get for any $\mathbf{A} \in \mathcal{B}(\mathbb{R}^d)$ and $\mathbf{B} \in \mathcal{B}(\mathbb{R}^p)$,

$$\begin{aligned}
& \Pi_{\rho, \gamma} P_{\rho, \gamma, \mathbf{N}}(\mathbf{A} \times \mathbf{B}) \tag{S47} \\
&= \int_{\mathbf{A} \times \mathbf{B}} \int_{\mathbb{R}^d \times \mathbb{R}^p} \Pi_{\rho, \gamma}(\mathrm{d}\tilde{\boldsymbol{\theta}}, \mathrm{d}\tilde{\mathbf{z}}) P_{\rho, \gamma, \mathbf{N}}((\tilde{\boldsymbol{\theta}}, \tilde{\mathbf{z}}), (\mathrm{d}\boldsymbol{\theta}, \mathrm{d}\mathbf{z})) \\
&= \int_{\mathbf{A} \times \mathbf{B}} \int_{\mathbb{R}^d \times \mathbb{R}^p} \Pi_{\rho, \gamma}(\mathrm{d}\tilde{\boldsymbol{\theta}}, \mathrm{d}\tilde{\mathbf{z}}) Q_{\rho, \gamma, \mathbf{N}}(\tilde{\mathbf{z}}, \mathrm{d}\mathbf{z} | \tilde{\boldsymbol{\theta}}) \Pi_{\rho}(\mathrm{d}\boldsymbol{\theta} | \mathbf{z}) \\
&= \int_{\mathbf{A} \times \mathbf{B}} \int_{\mathbb{R}^d \times \mathbb{R}^p} \Pi_{\rho, \gamma}(\mathrm{d}\tilde{\boldsymbol{\theta}}, \mathrm{d}\tilde{\mathbf{z}}) \left[\prod_{i=1}^b R_{\rho_i, \gamma_i}^{N_i}(\tilde{\mathbf{z}}_i, \mathrm{d}\mathbf{z}_i | \tilde{\boldsymbol{\theta}}) \right] \Pi_{\rho}(\mathrm{d}\boldsymbol{\theta} | \mathbf{z}) \\
&= \int_{\mathbf{A} \times \mathbf{B}} \int_{\mathbb{R}^d \times \mathbb{R}^p} \Pi_{\rho, \gamma}(\mathrm{d}\tilde{\boldsymbol{\theta}}, \mathrm{d}\tilde{\mathbf{z}}) \int_{\mathbb{R}^p} \left[\prod_{i=1}^b R_{\rho_i, \gamma_i}(\tilde{\mathbf{z}}_i, \mathrm{d}\tilde{\mathbf{z}}_i^{(1)} | \tilde{\boldsymbol{\theta}}) \right] \left[\prod_{i=1}^b R_{\rho_i, \gamma_i}^{N_i-1}(\tilde{\mathbf{z}}_i^{(1)}, \mathrm{d}\mathbf{z}_i | \tilde{\boldsymbol{\theta}}) \right] \Pi_{\rho}(\mathrm{d}\boldsymbol{\theta} | \mathbf{z}) \\
&= \int_{\mathbf{A} \times \mathbf{B}} \int_{\mathbb{R}^d \times \mathbb{R}^p} \left[\int_{\mathbb{R}^d \times \mathbb{R}^p} \Pi_{\rho, \gamma}(\mathrm{d}\tilde{\boldsymbol{\theta}}, \mathrm{d}\tilde{\mathbf{z}}) \left[\prod_{i=1}^b R_{\rho_i, \gamma_i}(\tilde{\mathbf{z}}_i, \mathrm{d}\tilde{\mathbf{z}}_i^{(1)} | \tilde{\boldsymbol{\theta}}) \right] \Pi_{\rho}(\mathrm{d}\tilde{\boldsymbol{\theta}}^{(1)} | \tilde{\mathbf{z}}_i^{(1)}) \right] \\
&\quad \times \left[\prod_{i=1}^b R_{\rho_i, \gamma_i}^{N_i-1}(\tilde{\mathbf{z}}_i^{(1)}, \mathrm{d}\mathbf{z}_i | \tilde{\boldsymbol{\theta}}) \right] \Pi_{\rho}(\mathrm{d}\boldsymbol{\theta} | \mathbf{z}) \\
&= \int_{\mathbf{A} \times \mathbf{B}} \int_{\mathbb{R}^d \times \mathbb{R}^p} \Pi_{\rho, \gamma}(\mathrm{d}\tilde{\boldsymbol{\theta}}^{(1)}, \mathrm{d}\tilde{\mathbf{z}}^{(1)}) \left[\prod_{i=1}^b R_{\rho_i, \gamma_i}^{N_i-1}(\tilde{\mathbf{z}}_i^{(1)}, \mathrm{d}\mathbf{z}_i | \tilde{\boldsymbol{\theta}}^{(1)}) \right] \Pi_{\rho}(\mathrm{d}\boldsymbol{\theta} | \mathbf{z}) .
\end{aligned}$$

Using a straightforward induction, we finally get

$$\int_{\mathbf{A} \times \mathbf{B}} \int_{\mathbb{R}^d \times \mathbb{R}^p} \Pi_{\rho, \gamma}(\mathrm{d}\tilde{\boldsymbol{\theta}}, \mathrm{d}\tilde{\mathbf{z}}) P_{\rho, \gamma, \mathbf{N}}((\tilde{\boldsymbol{\theta}}, \tilde{\mathbf{z}}), (\mathrm{d}\boldsymbol{\theta}, \mathrm{d}\mathbf{z})) = \int_{\mathbf{A} \times \mathbf{B}} \Pi_{\rho, \gamma}(\mathrm{d}\boldsymbol{\theta}, \mathrm{d}\mathbf{z}) ,$$

which shows that $P_{\rho, \gamma, \mathbf{N}}$ leaves $\Pi_{\rho, \gamma}$ invariant. Since this stationary distribution is unique, we conclude that $\Pi_{\rho, \gamma, \mathbf{N}} = \Pi_{\rho, \gamma}$. \square

We specify our result to the case where we take a specific initial distribution. To define it, consider

$$\mathbf{x}^* = ([\boldsymbol{\theta}^*]^\top, [\mathbf{z}^*]^\top)^\top, \text{ where } \boldsymbol{\theta}^* = \arg \min \{-\log \pi\} \text{ and } \mathbf{z}^* = ([\mathbf{A}_1 \boldsymbol{\theta}^*]^\top, \dots, [\mathbf{A}_b \boldsymbol{\theta}^*]^\top)^\top . \tag{S48}$$

We define the probability measure

$$\mu_{\rho}^* = \delta_{\mathbf{z}^*} \otimes \Pi_{\rho}(\cdot | \mathbf{z}^*) . \tag{S49}$$

Note that sampling from μ_ρ^\star is straightforward and simply consists in setting $\mathbf{z}_0 = \mathbf{z}^\star$ and $\boldsymbol{\theta}_0 = \bar{\mathbf{B}}_0^{-1} \mathbf{B}_0^\top \tilde{\mathbf{D}}_0^{1/2} \mathbf{z}_0 + \bar{\mathbf{B}}_0^{-1/2} \xi$, where ξ is a d -dimensional standard Gaussian random variable. We now specify our result when using μ_ρ^\star as an initial distribution. Define the \mathbf{z} -marginal under $\Pi_{\rho, \gamma}$ by

$$\pi_{\rho, \gamma}^{\mathbf{z}} = \int_{\mathbb{R}^d} \Pi_{\rho, \gamma}(\mathrm{d}\boldsymbol{\theta}, \mathbf{z}), \quad (\text{S50})$$

and the transition kernel of the Markov chain $\{Z_n\}_{n \geq 0}$, for all $\mathbf{z} \in \mathbb{R}^p$ and $\mathbf{B} \in \mathcal{B}(\mathbb{R}^p)$, by

$$P_{\rho, \gamma, N}^{\mathbf{z}}(\mathbf{z}, \mathbf{B}) = \int_{\mathbb{R}^d} Q_{\rho, \gamma, N}(\mathbf{z}, \mathbf{B} | \boldsymbol{\theta}) \Pi_\rho(\mathrm{d}\boldsymbol{\theta} | \mathbf{z}), \quad (\text{S51})$$

where $\Pi_\rho(\cdot | \cdot)$ and $Q_{\rho, \gamma, N}$ are defined in (5) and (S16), respectively.

Proposition S14. *Assume **H1-H2** and let $c > 0$ and $\gamma = \{\gamma_i\}_{i=1}^b$, $\mathbf{N} \in (\mathbb{N}^*)^b$ such that (S46) is satisfied, for any $i \in [b]$, $N_i \gamma_i < 2 / \max_{i \in [b]} \{m_i + \tilde{M}_i\}$ and $\gamma_i < \tilde{M}_i^{-1}$. Then, for any integer $n \geq 1$, we have*

$$\begin{aligned} W_2(\mu_\rho^\star P_{\rho, \gamma, N}^n, \Pi_{\rho, \gamma}) &\leq 2^{1/2} (1 - \min_{i \in [b]} \{N_i \gamma_i m_i\} / 2)^{n-1} \cdot (1 + \|\bar{\mathbf{B}}_0^{-1} \mathbf{B}_0^\top \tilde{\mathbf{D}}_0^{1/2}\|^2)^{1/2} \max_{i \in [b]} \{N_i \gamma_i\}^{1/2} \\ &\quad \times \left\{ \int_{\mathbb{R}^d} \|\mathbf{z}_1 - \mathbf{z}^\star\|_{\mathbf{D}_{N\gamma}^{-1}}^2 \pi_{\rho, \gamma}^{\mathbf{z}}(\mathrm{d}\mathbf{z}_1) + \int_{\mathbb{R}^d} \|\mathbf{z}_1 - \mathbf{z}^\star\|_{\mathbf{D}_{N\gamma}^{-1}}^2 P_{\rho, \gamma, N}^{\mathbf{z}}(\mathbf{z}^\star, \mathrm{d}\mathbf{z}_1) \right\}^{1/2}, \end{aligned}$$

where $\bar{\mathbf{B}}_0, \mathbf{B}_0, \tilde{\mathbf{D}}_0$ are defined in (S2)-(S3).

Proof. Consider for $n \in \mathbb{N}^*$, $X_n = (\theta_n^\top, Z_n^\top)^\top$, $\tilde{X}_n = (\tilde{\theta}_n^\top, \tilde{Z}_n^\top)^\top$ defined in (S18) with X_0 distributed according to μ_ρ^\star and \tilde{X}_0 distributed according to $\Pi_{\rho, \gamma}$. Combining Lemma S7, Lemma S10 and Lemma S12, we have for $n \geq 1$,

$$\|\tilde{Z}_{n+1} - Z_{n+1}\|_{\mathbf{D}_{N\gamma}^{-1}} \leq (1 - \min_{i \in [b]} \{N_i \gamma_i m_i\} / 2) \|\tilde{Z}_n - Z_n\|_{\mathbf{D}_{N\gamma}^{-1}}.$$

Thereby, for any $n \geq 1$, we obtain by induction

$$\|\tilde{Z}_n - Z_n\|_{\mathbf{D}_{N\gamma}^{-1}} \leq (1 - \min_{i \in [b]} \{N_i \gamma_i m_i\} / 2)^{n-1} \|\tilde{Z}_1 - Z_1\|_{\mathbf{D}_{N\gamma}^{-1}}. \quad (\text{S52})$$

Using $\|\tilde{Z}_1 - Z_1\|_{\mathbf{D}_{N\gamma}^{-1}}^2 \leq 2\|\tilde{Z}_1 - \mathbf{z}^\star\|_{\mathbf{D}_{N\gamma}^{-1/2}}^2 + 2\|Z_1 - \mathbf{z}^\star\|_{\mathbf{D}_{N\gamma}^{-1/2}}^2$ combined with the definition of the Wasserstein distance and Lemma S6 give

$$\begin{aligned} W_2(\mu_\rho^\star P_{\rho, \gamma, N}^n, \Pi_{\rho, \gamma}) &\leq \mathbb{E} \left[\|\tilde{X}_n - X_n\|^2 \right]^{1/2} \\ &\leq (1 + \|\bar{\mathbf{B}}_0^{-1} \mathbf{B}_0^\top \tilde{\mathbf{D}}_0^{1/2}\|^2)^{1/2} \max_{i \in [b]} \{N_i \gamma_i\}^{1/2} \mathbb{E} \left[\|\tilde{Z}_n - Z_n\|_{\mathbf{D}_{N\gamma}^{-1/2}}^2 \right]^{1/2} \\ &\leq 2^{1/2} (1 - \min_{i \in [b]} \{N_i \gamma_i m_i\} / 2)^{n-1} (1 + \|\bar{\mathbf{B}}_0^{-1} \mathbf{B}_0^\top \tilde{\mathbf{D}}_0^{1/2}\|^2)^{1/2} \max_{i \in [b]} \{N_i \gamma_i\}^{1/2} \\ &\quad \times \mathbb{E} \left[\|\tilde{Z}_1 - \mathbf{z}^\star\|_{\mathbf{D}_{N\gamma}^{-1/2}}^2 + \|Z_1 - \mathbf{z}^\star\|_{\mathbf{D}_{N\gamma}^{-1/2}}^2 \right]^{1/2}. \quad (\text{S53}) \end{aligned}$$

Since \tilde{X}_0 is distributed according to the stationary distribution $\Pi_{\rho, \gamma}$, \tilde{X}_1 also and therefore \tilde{Z}_1 is distributed according to $\pi_{\rho, \gamma}^z$. Finally, by definition Z_1 has distribution $P_{\rho, \gamma, N}^z(\mathbf{z}^*, \cdot)$, therefore (S53) completes the proof. \square

S3 Proof of Proposition 3

The proof of Proposition 3 stands for a generalization of Vono et al. (2019, Proposition 6) which only considered the specific case $\rho_i = \rho^2$ for $i \in [b]$. This section is divided into two parts, the first gathers lemmas which allow us to upper bound the ξ^2 -divergence between π_ρ and π . Then, in the second subsection, we combine these results to control the Wasserstein distance $W_2(\pi_\rho, \pi)$ by showing that it is smaller than $\chi^2(\pi_\rho | \pi)$. For any $\theta \in \mathbb{R}^d$ and $\rho \in (\mathbb{R}_+^*)^b$, define

$$U_i^{\rho_i}(\mathbf{A}_i \theta) = -\log \left(\int_{\mathbf{z}_i \in \mathbb{R}^d} \exp\{-U_i(\mathbf{z}_i) - \|\mathbf{z}_i - \mathbf{A}_i \theta\|^2 / (2\rho_i)\} d\mathbf{z}_i / (2\pi\rho_i)^{d_i/2} \right), \quad (\text{S54})$$

$$\overline{B}(\theta) = \sum_{i=1}^b \rho_i \|\nabla U_i(\mathbf{A}_i \theta)\|^2 / 2, \quad (\text{S55})$$

$$\underline{B}(\theta) = \sum_{i=1}^b \left\{ \rho_i \|\nabla U_i(\mathbf{A}_i \theta)\|^2 / [2(1 + \rho_i M_i)] - d_i \log(1 + \rho_i M_i) / 2 \right\} \quad (\text{S56})$$

and consider

$$U(\theta) = \sum_{i \in [b]} U_i(\mathbf{A}_i \theta), \quad U^\rho(\theta) = \sum_{i \in [b]} U_i^{\rho_i}(\mathbf{A}_i \theta).$$

S3.1 Technical lemmata

We start this subsection by Lemma S15 which allow us to bound the ratio between the integrals defined by $\int_{\mathbb{R}^d} \exp\{-\sum_{i \in [b]} U_i^{\rho_i}(\mathbf{A}_i \theta)\}$ and $\int_{\mathbb{R}^d} \exp\{-\sum_{i \in [b]} U_i(\mathbf{A}_i \theta)\} d\theta$.

Lemma S15. *Assume H1-H2-(i) and let $\rho \in (\mathbb{R}_+^*)^b$. Then, we have $\underline{B}(\theta) \leq U(\theta) - U^\rho(\theta)$, for any $\theta \in \mathbb{R}^d$. If we assume in addition that for any $i \in [b]$, U_i is convex, we have $U(\theta) - U^\rho(\theta) \leq \overline{B}(\theta)$, for any $\theta \in \mathbb{R}^d$.*

Proof. The proof follows from the same lines as in Vono et al. (2019, Lemma 14). In what follows, we give it for the sake of completeness. First, note for any $\theta \in \mathbb{R}^d$ and $i \in [b]$,

$$\exp\{U_i(\mathbf{A}_i \theta) - U_i^{\rho_i}(\mathbf{A}_i \theta)\} = \int_{\mathbb{R}^{d_i}} \exp\left(U_i(\mathbf{A}_i \theta) - U_i(\mathbf{z}_i) - \|\mathbf{z}_i - \mathbf{A}_i \theta\|^2 / (2\rho_i)\right) \frac{d\mathbf{z}_i}{(2\pi\rho_i)^{d_i/2}}. \quad (\text{S57})$$

Using **H2-(i)**, and a second order Taylor expansion, for any $\boldsymbol{\theta} \in \mathbb{R}^d, i \in [b], \mathbf{z}_i \in \mathbb{R}^{d_i}$, we have

$$U_i(\mathbf{A}_i \boldsymbol{\theta}) - U_i(\mathbf{z}_i) \geq \nabla U_i(\mathbf{A}_i \boldsymbol{\theta})^\top (\mathbf{A}_i \boldsymbol{\theta} - \mathbf{z}_i) - M_i \|\mathbf{A}_i \boldsymbol{\theta} - \mathbf{z}_i\|^2 / 2.$$

Hence, using (S57), we have for any $\boldsymbol{\theta} \in \mathbb{R}^d$ and $i \in [b]$,

$$\begin{aligned} \exp \left(\sum_{i=1}^b U_i(\mathbf{A}_i \boldsymbol{\theta}) - U_i^{\rho_i}(\mathbf{A}_i \boldsymbol{\theta}) \right) &\geq \prod_{i=1}^b \exp \left(\frac{\rho_i}{2(1 + \rho_i M_i)} \|\nabla U_i(\mathbf{A}_i \boldsymbol{\theta})\|^2 \right) (1 + \rho_i M_i)^{-d_i/2} \\ &= \exp(\underline{B}(\boldsymbol{\theta})). \end{aligned}$$

Similarly, under the assumption that for any $i \in [b]$, U_i is convex, the proof for the upper bound follows from the same lines using, for any $i \in [b]$, $\boldsymbol{\theta} \in \mathbb{R}^d$ and $\mathbf{z}_i \in \mathbb{R}^{d_i}$, that

$$U_i(\mathbf{A}_i \boldsymbol{\theta}) - U_i(\mathbf{z}_i) \leq \nabla U_i(\mathbf{A}_i \boldsymbol{\theta})^\top (\mathbf{A}_i \boldsymbol{\theta} - \mathbf{z}_i).$$

□

Lemma S16. Assume **H1-H2**. Then, U is m_U -strongly convex with $m_U = \lambda_{\min}(\sum_{i=1}^b m_i \mathbf{A}_i^\top \mathbf{A}_i)$.

Proof. Using by **H2-(i)** that for any $i \in [b]$, U_i is twice differentiable and by **H2-(ii)** the fact that for any $i \in [b]$, U_i is m_i -strongly convex, we have for any $\boldsymbol{\theta} \in \mathbb{R}^d$

$$\nabla^2 U(\boldsymbol{\theta}) = \sum_{i=1}^b \mathbf{A}_i^\top \nabla^2 U_i(\mathbf{A}_i \boldsymbol{\theta}) \mathbf{A}_i \succeq \sum_{i=1}^b m_i \mathbf{A}_i^\top \mathbf{A}_i \succeq \lambda_{\min} \left(\sum_{i=1}^b m_i \mathbf{A}_i^\top \mathbf{A}_i \right) \mathbf{I}_d = m_U \mathbf{I}_d.$$

□

For any $\boldsymbol{\theta} \in \mathbb{R}^d$, define

$$\beta(\boldsymbol{\theta}) = \left(\sum_{i=1}^b \rho_i \left\| \nabla U_i(\mathbf{A}_i \boldsymbol{\theta}) \right\|^2 \right)^{1/2}. \quad (\text{S58})$$

Lemma S17. Assume **H2-(i)** and let $\boldsymbol{\rho} \in (\mathbb{R}_+^*)^b$. Then β is a Lipschitz function w.r.t. $\|\cdot\|$, with Lipschitz constant

$$L_\beta = \lambda_{\max} \left(\sum_{i=1}^b \rho_i M_i^2 \mathbf{A}_i^\top \mathbf{A}_i \right)^{1/2}. \quad (\text{S59})$$

Proof. For any $\boldsymbol{\theta}_1, \boldsymbol{\theta}_2 \in \mathbb{R}^d$, we have using $|(\sum_{i=1}^b a_i^2)^{1/2} - (\sum_{i=1}^b b_i^2)^{1/2}| \leq (\sum_{i=1}^b (a_i - b_i)^2)^{1/2}$, that

$$|\beta(\boldsymbol{\theta}_1) - \beta(\boldsymbol{\theta}_2)| \leq \left(\sum_{i=1}^b \rho_i \|\nabla U_i(\mathbf{A}_i \boldsymbol{\theta}_1) - \nabla U_i(\mathbf{A}_i \boldsymbol{\theta}_2)\|^2 \right)^{1/2} \leq \left(\sum_{i=1}^b \rho_i M_i^2 \|\mathbf{A}_i(\boldsymbol{\theta}_1 - \boldsymbol{\theta}_2)\|^2 \right)^{1/2},$$

which completes the proof. □

Suppose **H2-(ii)** and for any $i \in [b]$, denote $\boldsymbol{\theta}_i^*$ a minimiser of $\boldsymbol{\theta} \mapsto U_i(\mathbf{A}_i \boldsymbol{\theta})$.

Lemma S18. Assume **H1-H2** and let $\boldsymbol{\rho} \in (\mathbb{R}_+^*)^b$. Then for any $s < m_U/(12L_\beta^2)$, where L_β is defined in (S59), we have

$$\log \pi[e^{s\{\beta^2 - \pi[\beta]^2\}}] \leq 8s^2 L_\beta^4 / m_U^2 + 4s^2 \{\pi[\beta]\}^2 L_\beta^2 / m_U. \quad (\text{S60})$$

In addition,

$$\pi(\beta^2) \leq 2dL_\beta^2 / m_U + 2 \sum_{i=1}^b \rho_i M_i^2 \|\mathbf{A}_i(\boldsymbol{\theta}^* - \boldsymbol{\theta}_i^*)\|^2. \quad (\text{S61})$$

Proof. Using the decomposition

$$\beta^2(\boldsymbol{\theta}) - \{\pi[\beta]\}^2 = (\beta(\boldsymbol{\theta}) - \pi[\beta])^2 + 2\pi[\beta](\beta(\boldsymbol{\theta}) - \pi[\beta])$$

and the Cauchy-Schwarz inequality imply, for any $s > 0$,

$$\pi[e^{s\{\beta^2 - \{\pi[\beta]\}^2\}}] \leq \{\pi[e^{2s\{\beta - \pi[\beta]\}^2}]\}^{1/2} \cdot \{\pi[e^{4s\pi[\beta]\{\beta - \pi[\beta]\}}]\}^{1/2}. \quad (\text{S62})$$

The proof consists in bounding the two terms in the right-hand sided. Since $\beta : \mathbb{R}^d \rightarrow \mathbb{R}$ is L_β -Lipschitz by Lemma S17, for any $0 \leq s \leq m_U/(12L_\beta^2)$, using Vono et al. (2019, Lemma 16) and Lemma S16 gives setting $\bar{\beta} = \beta - \pi[\beta]$, that

$$\pi[\exp\{2s(\bar{\beta}^2 - \pi[\bar{\beta}]^2)\}] \leq \exp(16s^2 L_\beta^4 / m_U^2). \quad (\text{S63})$$

In addition, using Bakry et al. (2013, Proposition 5.4.1), Lemma S17 and Lemma S16, we get for any $s \geq 0$,

$$\pi[e^{4s\pi[\beta](\beta - \pi[\beta])}] \leq e^{8s^2 \{\pi[\beta]\}^2 L_\beta^2 / m_U}.$$

Plugging this result and (S63) into (S62), we get

$$\pi[e^{s\{\beta^2 - \{\pi[\beta]\}^2\}}] \leq \exp(s\pi(\bar{\beta}^2) + 8s^2 L_\beta^4 / m_U^2 + 4s^2 \{\pi[\beta]\}^2 L_\beta^2 / m_U).$$

The proof of (S60) follows using $\pi(\bar{\beta}^2) = \pi(\beta^2) - [\pi(\beta)]^2$ and rearranging terms.

Using the Young inequality, **H2-(i)**, $\nabla U_i(\mathbf{A}_i \boldsymbol{\theta}_i^*) = 0$, $\nabla U(\boldsymbol{\theta}^*) = 0$, we have

$$\begin{aligned} \pi(\beta^2) &= \int_{\mathbb{R}^d} \left(\sum_{i=1}^b \rho_i \|\nabla U_i(\mathbf{A}_i \boldsymbol{\theta})\|^2 \right) \pi(\boldsymbol{\theta}) \, d\boldsymbol{\theta} \\ &\leq 2 \int_{\mathbb{R}^d} \left(\sum_{i=1}^b \rho_i M_i^2 \|\mathbf{A}_i(\boldsymbol{\theta} - \boldsymbol{\theta}^*)\|^2 \right) \pi(\boldsymbol{\theta}) \, d\boldsymbol{\theta} + 2 \sum_{i=1}^b \rho_i M_i^2 \|\mathbf{A}_i(\boldsymbol{\theta}^* - \boldsymbol{\theta}_i^*)\|^2 \\ &\leq 2\lambda_{\max} \left(\sum_{i=1}^b \rho_i M_i^2 \mathbf{A}_i^\top \mathbf{A}_i \right) \int_{\mathbb{R}^d} \|\boldsymbol{\theta} - \boldsymbol{\theta}^*\|^2 \pi(\boldsymbol{\theta}) \, d\boldsymbol{\theta} + 2 \sum_{i=1}^b \rho_i M_i^2 \|\mathbf{A}_i(\boldsymbol{\theta}^* - \boldsymbol{\theta}_i^*)\|^2 \\ &\leq 2dL_\beta^2 / m_U + 2 \sum_{i=1}^b \rho_i M_i^2 \|\mathbf{A}_i(\boldsymbol{\theta}^* - \boldsymbol{\theta}_i^*)\|^2, \end{aligned}$$

where we have used $\pi[\|\boldsymbol{\theta} - \boldsymbol{\theta}^*\|^2] \leq d/m_U$ by [Durmus and Moulines \(2019, Proposition 1 \(ii\)\)](#) and [Lemma S16](#). \square

[Proposition 1](#) shows that $\pi_{\boldsymbol{\rho}}(\cdot) = \int_{\mathbb{R}^p} \Pi_{\boldsymbol{\rho}}(\cdot, \mathbf{z}) d\mathbf{z}$ is well-defined and as such admits a finite normalising constant. These two quantities are defined by

$$Z_{\pi_{\boldsymbol{\rho}}} = \int_{\mathbb{R}^d} \exp \left\{ - \sum_{i \in [b]} U_i^{\rho_i}(\mathbf{A}_i \boldsymbol{\theta}) \right\} d\boldsymbol{\theta}, \quad \pi_{\boldsymbol{\rho}}(\cdot) = \exp \left\{ - \sum_{i \in [b]} U_i^{\rho_i}(\mathbf{A}_i \cdot) \right\} / Z_{\pi_{\boldsymbol{\rho}}}. \quad (\text{S64})$$

Finally, note that the following quantity Z_{π} is a normalising constant of π associated with the potential U , *i.e.* $\pi = e^{-U}/Z_{\pi}$,

$$Z_{\pi} = \int_{\mathbb{R}^d} \exp \left\{ - \sum_{i \in [b]} U_i(\mathbf{A}_i \boldsymbol{\theta}) \right\} d\boldsymbol{\theta}. \quad (\text{S65})$$

Lemma S19. *Assume [H1-H2](#) and let $\boldsymbol{\rho} \in (\mathbb{R}_+^*)^b$. Suppose in addition that $6L_{\beta}^2 \leq m_U$ where L_{β} is given in [\(S59\)](#). Then, we have*

$$\log(Z_{\pi_{\boldsymbol{\rho}}}/Z_{\pi}) \leq \left\{ dL_{\beta}^2/m_U + \sum_{i=1}^b \rho_i M_i^2 \|\mathbf{A}_i(\boldsymbol{\theta}^* - \boldsymbol{\theta}_i^*)\|^2 \right\} (1 + 2L_{\beta}^2/m_U) + 2L_{\beta}^4/m_U^2.$$

Proof. From the definitions [\(S64\)](#) and [\(S65\)](#), we have $Z_{\pi_{\boldsymbol{\rho}}}/Z_{\pi} = \int_{\mathbb{R}^d} \pi(\boldsymbol{\theta}) \exp\{\sum_{i=1}^b U_i(\mathbf{A}_i \boldsymbol{\theta}) - U_i^{\rho_i}(\mathbf{A}_i \boldsymbol{\theta})\} d\boldsymbol{\theta}$. By [Lemma S15](#), we obtain

$$Z_{\pi_{\boldsymbol{\rho}}}/Z_{\pi} \leq \int_{\mathbb{R}^d} \pi(\boldsymbol{\theta}) \exp(\overline{B}(\boldsymbol{\theta})) d\boldsymbol{\theta}.$$

Note that $\overline{B} = \beta^2/2$ by [\(S55\)-\(S58\)](#), hence using that $6L_{\beta}^2 \leq m_U$, [Lemma S18](#) applied with $s = 1/2$ shows that

$$\log \left(\int_{\mathbb{R}^d} \pi(\boldsymbol{\theta}) \exp(\overline{B}(\boldsymbol{\theta})) d\boldsymbol{\theta} \right) \leq \pi[\beta^2]/2 + 2L_{\beta}^4/m_U^2 + \{\pi[\beta]\}^2 L_{\beta}^2/m_U.$$

Using [Lemma S18-\(S61\)](#) and $\pi[\beta] \leq \pi[\beta^2]$ concludes the proof. \square

S3.2 Proof of Proposition 3

Based on the technical lemmas derived in [Appendix S3.1](#), we are now ready to bound the Wasserstein distance of order 2 between π and $\pi_{\boldsymbol{\rho}}$.

Proof of Proposition Proposition 3. Let $\boldsymbol{\rho} \in (\mathbb{R}_+^*)^b$ such that $\max_{i \in [b]} \rho_i = \bar{\rho} \leq \sigma_U^2/12$, where $\sigma_U^2 = \|\mathbf{A}^\top \mathbf{A}\| \max_{i \in [b]} \{M_i^2\}/m_U$. Then, by definition of L_β (S59), we get

$$12L_\beta^2 \leq m_U . \quad (\text{S66})$$

and Lemma S18 can be applied for $s = 1$ and Lemma S19 too. By Lemma S16, $U = -\log \pi$ is m_U -strongly convex therefore π satisfies a log-Sobolev inequality with constant m_U (Ledoux, 2001, Theorem 5.2). Finally, Otto and Villani (2000, Theorem 1) shows that π satisfies for any $\nu \in \mathcal{P}_2(\mathbb{R}^d)$:

$$W_2(\nu, \pi) \leq \sqrt{(2/m_U) \text{KL}(\nu|\pi)} \leq \sqrt{(2/m_U) \chi^2(\pi_\rho|\pi)} , \quad (\text{S67})$$

where χ^2 is the chi-square divergence and where we have used for the last inequality that $\text{KL}(\pi_\rho|\pi) \leq \chi^2(\pi_\rho|\pi)$ since for any $t > 0$, $\log(t) \leq t - 1$. We now bound $\chi^2(\pi_\rho|\pi)$. By (S64) and (S65), for any $\boldsymbol{\theta} \in \mathbb{R}^d$, consider the decomposition given by

$$\pi_\rho(\boldsymbol{\theta})/\pi(\boldsymbol{\theta}) - 1 = (Z_\pi/Z_{\pi_\rho}) \exp \left(\sum_{i=1}^b (U_i(\mathbf{A}_i \boldsymbol{\theta}) - U_i^{\rho_i}(\mathbf{A}_i \boldsymbol{\theta})) \right) - 1 . \quad (\text{S68})$$

In the sequel, we will both lower and upper bound (S68) in order to upper bound $|1 - \pi_\rho(\boldsymbol{\theta})/\pi(\boldsymbol{\theta})|$. Using the fact that for all $x \in \mathbb{R}$, $\exp(x) - 1 \geq x$, Lemmas S15 and S19 yield

$$\begin{aligned} \pi_\rho(\boldsymbol{\theta})/\pi(\boldsymbol{\theta}) - 1 &\geq \log(Z_\pi/Z_{\pi_\rho}) + \sum_{i=1}^b (U_i(\mathbf{A}_i \boldsymbol{\theta}) - U_i^{\rho_i}(\mathbf{A}_i \boldsymbol{\theta})) \\ &\geq - \left\{ dL_\beta^2/m_U + \sum_{i=1}^b \rho_i M_i^2 \|\mathbf{A}_i(\boldsymbol{\theta}^\star - \boldsymbol{\theta}_i^\star)\|^2 \right\} (1 + 2L_\beta^2/m_U) - 2L_\beta^4/m_U^2 + \underline{B}(\boldsymbol{\theta}) \geq -A_1 , \end{aligned} \quad (\text{S69})$$

where

$$\begin{aligned} A_1 &= \left\{ dL_\beta^2/m_U + \sum_{i=1}^b \rho_i M_i^2 \|\mathbf{A}_i(\boldsymbol{\theta}^\star - \boldsymbol{\theta}_i^\star)\|^2 \right\} (1 + 2L_\beta^2/m_U) \\ &\quad + 2L_\beta^4/m_U^2 + \sum_{i=1}^b (d_i/2) \log(1 + \rho_i M_i) , \end{aligned}$$

where we have used in the last inequality that $\underline{B}(\boldsymbol{\theta}) \geq -\sum_{i=1}^b (d_i/2) \log(1 + \rho_i M_i)$ by (S55). In addition, by (S64) and (S65) $Z_{\pi_\rho}/Z_\pi = \int_{\mathbb{R}^d} \pi(\boldsymbol{\theta}) \exp\{\sum_{i=1}^b U_i(\mathbf{A}_i \boldsymbol{\theta}) - U_i^{\rho_i}(\mathbf{A}_i \boldsymbol{\theta})\} d\boldsymbol{\theta}$, which implies by Lemma S15 and Jensen inequality

$$Z_{\pi_\rho}/Z_\pi \geq \int_{\mathbb{R}^d} \pi(\boldsymbol{\theta}) \exp(\underline{B}(\boldsymbol{\theta})) d\boldsymbol{\theta} \geq \exp(\pi[\underline{B}]) .$$

It follows by (S68) that $\pi_{\boldsymbol{\rho}}(\boldsymbol{\theta})/\pi(\boldsymbol{\theta}) - 1 \leq \exp(\bar{B}(\boldsymbol{\theta}) - \pi(\underline{B})) - 1$. Combining this result and (S69), it follows that the Pearson χ^2 -divergence between π and $\pi_{\boldsymbol{\rho}}$ can be upper bounded as where

$$\chi^2(\pi_{\boldsymbol{\rho}}|\pi) \leq \max(A_1^2, A_2), \quad A_2 = \int_{\mathbb{R}^d} (\exp(\bar{B}(\boldsymbol{\theta}) - \pi(\underline{B})) - 1)^2 \pi(\boldsymbol{\theta}) d\boldsymbol{\theta}.$$

We now provide an explicit bound for A_2 . First by Jensen inequality, we have $\pi(\exp(\bar{B})) \geq \exp(\pi(\bar{B}))$ which implies that $\exp(-\pi(\underline{B}))\pi[\exp(\bar{B})] \geq \prod_{i=1}^b (1 + \rho_i M_i)^{d_i/2}$ by (S55). Therefore, using that $\bar{B} = \beta^2/2$ by (S55)-(S58) and Lemma S18 with $s = 1$ since (S66) holds, we get by (S55),

$$\begin{aligned} A_2 &= \int_{\mathbb{R}^d} (\exp(\bar{B}(\boldsymbol{\theta}) - \pi(\underline{B})) - 1)^2 \pi(\boldsymbol{\theta}) d\boldsymbol{\theta} \\ &= \exp(-2\pi(\underline{B}))\pi[\exp(2\bar{B})] - 2\exp(-\pi(\underline{B}))\pi[\exp(\bar{B})] + 1 \\ &\leq \prod_{i=1}^b (1 + \rho_i M_i)^{d_i} \cdot \exp(-\pi\{\sum_{i=1}^b (\rho_i/(1 + \rho_i M_i))\|\nabla U_i(\mathbf{A}_i \cdot)\|^2\})\pi[\exp(\beta^2)] \\ &\quad - 2 \prod_{i=1}^b (1 + \rho_i M_i)^{d_i/2} + 1 \\ &\leq \prod_{i=1}^b (1 + \rho_i M_i)^{d_i} \cdot \exp(\pi\{\sum_{i=1}^b (\rho_i^2 M_i/(1 + \rho_i M_i))\|\nabla U_i(\mathbf{A}_i \cdot)\|^2\}) \\ &\quad \times \exp\left(8L_{\beta}^4/m_U^2 + 4\{2dL_{\beta}^2/m_U + 2\sum_{i=1}^b \rho_i M_i^2\|\mathbf{A}_i(\boldsymbol{\theta}^* - \boldsymbol{\theta}_i^*)\|^2\}L_{\beta}^2/m_U\right) \end{aligned} \quad (\text{S70})$$

$$- 2 \prod_{i=1}^b (1 + \rho_i M_i)^{d_i/2} + 1, \quad (\text{S71})$$

where we have used for the last inequality that for $\boldsymbol{\theta} \in \mathbb{R}^d$, $\beta(\boldsymbol{\theta})^2 - \sum_{i=1}^b (\rho_i/(1 + \rho_i M_i))\|\nabla U_i(\mathbf{A}_i \boldsymbol{\theta})\|^2 = \sum_{i=1}^b (\rho_i^2 M_i/(1 + \rho_i M_i))\|\nabla U_i(\mathbf{A}_i \boldsymbol{\theta})\|^2$, $\pi[\beta]^2 \leq \pi[\beta^2]$ by the Cauchy-Schwartz inequality and Lemma S18-(S61). Similarly to the proof of Lemma S18-(S61), by **H2-(i)**, $\nabla U_i(\mathbf{A}_i \boldsymbol{\theta}_i^*) = 0$, $\nabla U(\boldsymbol{\theta}^*) = 0$, Durmus and Moulines (2019, Proposition 1 (ii)) and Lemma S16, we have

$$\begin{aligned} \pi\left[\sum_{i=1}^b (\rho_i^2 M_i/(1 + \rho_i M_i))\|\nabla U_i(\mathbf{A}_i \cdot)\|^2\right] &\leq \pi\left[\sum_{i=1}^b \rho_i^2 M_i\|\nabla U_i(\mathbf{A}_i \cdot)\|^2\right] \\ &\leq 2d\lambda_{\max}\left(\sum_{i=1}^b \rho_i^2 M_i^3 \mathbf{A}_i^{\top} \mathbf{A}_i\right)/m_U + 2\sum_{i=1}^b \rho_i^2 M_i^3\|\mathbf{A}_i(\boldsymbol{\theta}^* - \boldsymbol{\theta}_i^*)\|^2. \end{aligned}$$

Therefore, we get by (S71)

$$\begin{aligned}
A_2 \leq A_3 &= \prod_{i=1}^b (1 + \rho_i M_i)^{d_i} \exp \left(2d\lambda_{\max} \left(\sum_{i=1}^b \rho_i^2 M_i^3 \mathbf{A}_i^\top \mathbf{A}_i \right) / m_U + 2 \sum_{i=1}^b \rho_i^2 M_i^3 \|\mathbf{A}_i(\boldsymbol{\theta}^\star - \boldsymbol{\theta}_i^\star)\|^2 \right) \\
&\exp \left(8L_\beta^4 / m_U^2 + 8 \left[dL_\beta^2 / m_U + \sum_{i=1}^b \rho_i M_i^2 \|\mathbf{A}_i(\boldsymbol{\theta}^\star - \boldsymbol{\theta}_i^\star)\|^2 \right] L_\beta^2 / m_U \right) - 2 \prod_{i=1}^b (1 + \rho_i M_i)^{d_i/2} + 1 .
\end{aligned} \tag{S72}$$

It follows by (S70) and (S67) that

$$W_2(\pi_\rho, \pi) \leq \sqrt{(2/m_U) \max(A_1^2, A_3)} , \tag{S73}$$

where A_1 and A_3 are given by (S69) and (S72) respectively. Using that $L_\beta^2 = \mathcal{O}(\bar{\rho})$ and an expansion of the bound as $\bar{\rho} \rightarrow 0$ completes the proof. \square

S4 Proof of Proposition 4 and Proposition 5

As in Appendix S2, we assume in all this section that $\boldsymbol{\rho} \in (\mathbb{R}_+^*)^b$ is fixed. For any $\boldsymbol{\gamma} = (\gamma_1, \dots, \gamma_b) \in (\mathbb{R}_+^*)^b$, we establish in this section explicit bounds on $W_2(\pi_{\boldsymbol{\rho}, \boldsymbol{\gamma}, \mathbf{N}}, \pi_\rho)$ where π_ρ is given in (1) and $\pi_{\boldsymbol{\rho}, \boldsymbol{\gamma}, \mathbf{N}}$ is the marginal distribution defined by

$$\pi_{\boldsymbol{\rho}, \boldsymbol{\gamma}, \mathbf{N}}(\mathbf{A}) = \Pi_{\boldsymbol{\rho}, \boldsymbol{\gamma}, \mathbf{N}}(\mathbf{A} \times \mathbb{R}^p) , \quad \mathbf{A} \in \mathcal{B}(\mathbb{R}^d) ,$$

of the stationary probability measure $\Pi_{\boldsymbol{\rho}, \boldsymbol{\gamma}, \mathbf{N}}$ associated with the Markov chain $(Z_n, \theta_n)_{n \geq 0}$ defined in Algorithm 1. Note that in the case $\mathbf{N} = N(1, \dots, 1)$, this distribution is independent of N , see Proposition S13. To this purpose, we define an “ideal” dynamics from which we cannot sample but which converges geometrically towards Π_ρ under appropriate conditions. The corresponding ideal process will play the same role as the Langevin dynamics for the study of the unadjusted Langevin algorithm (Durmus and Moulines, 2019). This dynamics is defined as follows. Consider first for any $\boldsymbol{\theta} \in \mathbb{R}^d$, $i \in [b]$, the stochastic differential equation (SDE) defined by

$$d\tilde{Y}_t^{i, \boldsymbol{\theta}} = -\nabla V_i(\tilde{Y}_t^{i, \boldsymbol{\theta}}) dt - \rho_i^{-1} \mathbf{A}_i \boldsymbol{\theta} + \sqrt{2} dB_t^i , \tag{S74}$$

where $(B_t^i)_{t \geq 0}$ is a d_i -dimensional Brownian motion and V_i is defined in (S20). Note that under H2-(i), this SDE admits a unique strong solution (Revuz and Yor, 2013, Theorem (2.1) in Chapter IX). Denote for any $i \in [b]$, the Markov semi-group associated to (S74) by $(\tilde{R}_{\rho_i, t}^i)_{t \geq 0}$ defined for any $\tilde{\mathbf{y}}_0^i \in \mathbb{R}^{d_i}$, $t \geq 0$ and $\mathbf{B}_i \in \mathcal{B}(\mathbb{R}^{d_i})$ by

$$\tilde{R}_{\rho_i, t}^i(\tilde{\mathbf{y}}_0^i, \mathbf{B}_i | \boldsymbol{\theta}) = \mathbb{P}(\tilde{Y}_t^{i, \boldsymbol{\theta}, \tilde{\mathbf{y}}_0^i} \in \mathbf{B}_i) ,$$

where $(\tilde{Y}_t^{i,\boldsymbol{\theta},\tilde{\mathbf{y}}_0^i})_{t \geq 0}$ is a solution of (S74) with $\tilde{Y}_0^{i,\boldsymbol{\theta},\tilde{\mathbf{y}}_0^i} = \tilde{\mathbf{y}}_0^i$. For any bounded measurable function $f_i : \mathbb{R}^{d_i} \rightarrow \mathbb{R}_+$, Lemma S20 shows the measurability of the function $(\boldsymbol{\theta}, \tilde{\mathbf{y}}_0^i) \mapsto \mathbb{E}[f_i(\tilde{Y}_t^{i,\boldsymbol{\theta},\tilde{\mathbf{y}}_0^i})]$ on $\mathbb{R}^d \times \mathbb{R}^{d_i}$ and therefore $\tilde{R}_{\rho_i,t}^i$ is a conditional Markov kernel.

Lemma S20. *For any bounded measurable function $f_i : \mathbb{R}^{d_i} \rightarrow \mathbb{R}_+$ and function f_i satisfying **H2-(i)**, the mapping $(\tilde{\boldsymbol{\theta}}_0, \tilde{\mathbf{y}}_0^i) \mapsto \mathbb{E}[f_i(\tilde{Y}_t^{i,\tilde{\boldsymbol{\theta}}_0,\tilde{\mathbf{y}}_0^i})]$ is Borel measurable.*

Proof. Consider the following stochastic differential equation

$$\begin{cases} d\tilde{\boldsymbol{\theta}}_t = \mathbf{0}_d, \\ d\tilde{Y}_t^i = -\nabla V_i(\tilde{Y}_t^i) dt - \rho_i^{-1} \mathbf{A}_i \tilde{\boldsymbol{\theta}}_t + \sqrt{2} dB_t^i. \end{cases}$$

Using Revuz and Yor (2013, Theorem (2.4) in Chapter IX), since U_i satisfies **H2-(i)**, there exists a unique solution $(\tilde{X}_t^{\tilde{\mathbf{x}}})_{t \geq 0} = (\tilde{\boldsymbol{\theta}}_t, \tilde{Y}_t^i)_{t \geq 0}$ with initial condition $\tilde{\mathbf{x}} = (\tilde{\boldsymbol{\theta}}_0^\top, (\tilde{\mathbf{y}}_0^i)^\top)^\top \in \mathbb{R}^p$. Then, the proof follows from Revuz and Yor (2013, Theorem (1.9) in Chapter IX) and the fact that \tilde{Y}_t^i is the unique solution of (S74) with $\boldsymbol{\theta} = \boldsymbol{\theta}_0$. \square

Define for any $\boldsymbol{\theta} \in \mathbb{R}^d$, $\mathbf{z} = (\mathbf{z}_1^\top, \dots, \mathbf{z}_b^\top)^\top \in \mathbb{R}^p$, and for $i \in [b]$, $B_i \in \mathcal{B}(\mathbb{R}^{d_i})$,

$$\tilde{Q}_{\rho,\gamma}(\mathbf{z}, B_1 \times \dots \times B_b | \boldsymbol{\theta}) = \prod_{i=1}^b \tilde{R}_{\rho_i, N_i \gamma_i}^i(\mathbf{z}_i, B_i | \boldsymbol{\theta}),$$

and consider the Markov kernel defined, for any $\mathbf{x}^\top = (\boldsymbol{\theta}^\top, \mathbf{z}^\top)$ and $A \in \mathcal{B}(\mathbb{R}^d)$, $B \in \mathcal{B}(\mathbb{R}^p)$, by

$$\tilde{P}_{\rho,\gamma}(\mathbf{x}, A \times B) = \int_B \tilde{Q}_{\rho,\gamma}(\mathbf{z}, d\tilde{\mathbf{z}} | \boldsymbol{\theta}) \int_A \Pi_\rho(d\tilde{\boldsymbol{\theta}} | \tilde{\mathbf{z}}), \quad (\text{S75})$$

where $\Pi_\rho(\cdot | \tilde{\mathbf{z}})$ is defined in (5). Note that $P_{\rho,\gamma,N}$ can be interpreted as a discretised version of $\tilde{P}_{\rho,\gamma}$ using the Euler-Maruyama scheme.

In the sequel, we first derive technical lemmata in Appendix S4.1 that are used to prove both Proposition 4 and Proposition 5. Based on these lemmata, we then prove each proposition in a dedicated section, namely Appendix S4.2 and Appendix S4.3.

S4.1 Synchronous coupling and a first estimate

The main idea to prove Proposition 4 and Proposition 5 is to define $(X_n, \tilde{X}_n)_{n \in \mathbb{N}}$ such that for any $n \in \mathbb{N}$, (X_n, \tilde{X}_n) is a coupling between $\delta_{\mathbf{x}} P_{\rho,\gamma,N}^n$ defined in (S17) and $\delta_{\tilde{\mathbf{x}}} \tilde{P}_{\rho,\gamma}^n$, and satisfies

$$\mathbb{E} \left[\left\| X_n - \tilde{X}_n \right\|^2 \right] \leq c_1(\mathbf{x}, \tilde{\mathbf{x}}) e^{-c_2 \min_{i \in [b]} \{\gamma_i m_i\}} + c_3 \gamma^\alpha,$$

where $c_2, c_3 > 0$ and $\alpha \in \{1, 2\}$ depending if **H3** holds or not. Conditioning with respect to (X_0, \tilde{X}_0) with distribution $\delta_{\mathbf{x}} \otimes \Pi_{\boldsymbol{\rho}}$, using the definition of the Wasserstein distance of order 2 and taking $n \rightarrow \infty$, we obtain

$$W_2(\pi_{\boldsymbol{\rho}}, \pi_{\boldsymbol{\rho}, \gamma, \mathbf{N}}) \leq W_2(\Pi_{\boldsymbol{\rho}}, \Pi_{\boldsymbol{\rho}, \gamma, \mathbf{N}}) \leq \tilde{c}_3 \gamma^\alpha ,$$

where $\tilde{c}_3 > 0$. We now provide the rigorous construction of $(X_n, \tilde{X}_n)_{n \in \mathbb{N}}$.

Let $\{(B_t^{(i,n)})_{t \geq 0} : i \in [b], n \in \mathbb{N}\}$ be independent random variables such that for any $i \in [b]$, the sequences $\{(B_t^{(i,n)})_{t \geq 0} : n \in \mathbb{N}\}$ are i.i.d. d_i -dimensional Brownian motions and let $(\xi_n)_{n \geq 0}$ be a sequence of i.i.d. standard d -dimensional Gaussian random variables independent of $\{(B_t^{(i,n)})_{t \geq 0} : i \in [b], n \in \mathbb{N}\}$. Consider the stochastic process $(\tilde{X}_n)_{n \geq 0}$ on $\mathbb{R}^d \times \mathbb{R}^p$ starting from \tilde{X}_0 distributed according to $\Pi_{\boldsymbol{\rho}}$ and defined by the recursion: for $n \in \mathbb{N}$, $i \in [b]$,

$$\tilde{X}_{n+1} = (\tilde{\theta}_{n+1}^\top, \tilde{Z}_{n+1}^\top)^\top, \quad \tilde{Z}_{n+1}^i = \tilde{Y}_{N_i \gamma_i}^{(i,n)}, \quad \tilde{\theta}_{n+1} = \bar{\mathbf{B}}_0^{-1} \mathbf{B}_0^\top \tilde{\mathbf{D}}_0^{1/2} \tilde{Z}_{n+1} + \bar{\mathbf{B}}_0^{-1/2} \xi_{n+1}, \quad (\text{S76})$$

where $(\tilde{Y}_t^{(i,n)})_{t \geq 0}$, is a solution of (S74) starting from \tilde{Z}_n^i with parameter $\boldsymbol{\theta} \leftarrow \boldsymbol{\theta}_n$. Similarly to the process $(X_n)_{n \in \mathbb{N}}$ defined in Algorithm 1, the process $(\tilde{X}_n)_{n \in \mathbb{N}}$ defines a homogeneous Markov chain. Indeed, it is easy to show that for any $n \in \mathbb{N}$ and measurable function $f : \mathbb{R}^p \rightarrow \mathbb{R}_+$, $\mathbb{E}[f(\tilde{Z}_{n+1}) | \tilde{X}_n] = \int_{\mathbb{R}^p} f(\tilde{\mathbf{z}}) \tilde{Q}_{\boldsymbol{\rho}, \gamma}(\tilde{Z}_n, d\tilde{\mathbf{z}} | \tilde{\theta}_n)$ and therefore $(\tilde{X}_n)_{n \in \mathbb{N}}$ is associated with (S75).

Proposition S21. *Assume **H1-H2-(i)**, and let $\mathbf{N} \in (\mathbb{N}^*)^b, \gamma \in (\mathbb{R}_+^*)^b$. Then, the Markov kernel $\tilde{P}_{\boldsymbol{\rho}, \gamma}$ defined in (S75) admits $\Pi_{\boldsymbol{\rho}}$ as an invariant probability measure.*

Proof. By property of the Langevin diffusion defined in (S74), for all $\boldsymbol{\theta}_0 \in \mathbb{R}^d$, the Markov kernel $\tilde{Q}_{\boldsymbol{\rho}, \gamma}(\cdot | \boldsymbol{\theta}_0)$ admits $\Pi_{\boldsymbol{\rho}}(\cdot | \boldsymbol{\theta}_0)$ as invariant measure, see e.g. Roberts and Tweedie (1996) or Kent (1978). Thus, for any $\boldsymbol{\theta}_0 \in \mathbb{R}^d$ and $\mathbf{B} \in \mathcal{B}(\mathbb{R}^p)$, we have

$$\int_{\mathbf{B}} \Pi_{\boldsymbol{\rho}}(\mathbf{z}_1 | \boldsymbol{\theta}_0) d\mathbf{z}_1 = \int_{\mathbf{z}_0 \in \mathbb{R}^p} \tilde{Q}_{\boldsymbol{\rho}, \gamma}(\mathbf{z}_0, \mathbf{B} | \boldsymbol{\theta}_0) \Pi_{\boldsymbol{\rho}}(\mathbf{z}_0 | \boldsymbol{\theta}_0) d\mathbf{z}_0. \quad (\text{S77})$$

Denote by $\pi_{\boldsymbol{\rho}}^\theta, \pi_{\boldsymbol{\rho}}^{\mathbf{z}}$ the marginals under $\Pi_{\boldsymbol{\rho}}$: $\pi_{\boldsymbol{\rho}}^\theta(\mathbf{A}) = \Pi_{\boldsymbol{\rho}}(\mathbf{A} \times \mathbb{R}^p)$, $\pi_{\boldsymbol{\rho}}^{\mathbf{z}}(\mathbf{B}) = \Pi_{\boldsymbol{\rho}}(\mathbb{R}^d \times \mathbf{B})$, for $\mathbf{A} \in \mathcal{B}(\mathbb{R}^d)$ and $\mathbf{B} \in \mathcal{B}(\mathbb{R}^p)$, and consider the Markov chain $(\tilde{X}_n)_{n \in \mathbb{N}}$ defined in (S76). For

any measurable function $f : \mathbb{R}^{d+p} \rightarrow \mathbb{R}_+$, the Fubini-Tonelli theorem gives

$$\begin{aligned}
\mathbb{E}[f(\tilde{X}_1)] &= \int_{\mathbb{R}^{d+p}} \int_{\mathbb{R}^{d+p}} f(\mathbf{x}_1) \Pi_{\rho}(\boldsymbol{\theta}_1 | \mathbf{z}_1) d\boldsymbol{\theta}_1 \tilde{Q}_{\rho, \gamma}(\mathbf{z}_0, d\mathbf{z}_1 | \boldsymbol{\theta}_0) \Pi_{\rho}(\boldsymbol{\theta}_0, \mathbf{z}_0) d\boldsymbol{\theta}_0 d\mathbf{z}_0 \\
&= \int_{\mathbb{R}^d} \int_{\mathbb{R}^p} f(\mathbf{x}_1) \Pi_{\rho}(\boldsymbol{\theta}_1 | \mathbf{z}_1) \int_{\mathbb{R}^d} \left[\int_{\mathbb{R}^p} \tilde{Q}_{\rho, \gamma}(\mathbf{z}_0, d\mathbf{z}_1 | \boldsymbol{\theta}_0) \Pi_{\rho}(\mathbf{z}_0 | \boldsymbol{\theta}_0) d\mathbf{z}_0 \right] \pi_{\rho}^{\boldsymbol{\theta}}(\boldsymbol{\theta}_0) d\boldsymbol{\theta}_0 d\boldsymbol{\theta}_1 \\
&= \int_{\mathbb{R}^d} \int_{\mathbb{R}^p} f(\mathbf{x}_1) \Pi_{\rho}(\boldsymbol{\theta}_1 | \mathbf{z}_1) \left[\int_{\boldsymbol{\theta}_0 \in \mathbb{R}^d} \Pi_{\rho}(\mathbf{z}_1 | \boldsymbol{\theta}_0) \pi_{\rho}^{\boldsymbol{\theta}}(\boldsymbol{\theta}_0) d\boldsymbol{\theta}_0 \right] d\mathbf{z}_1 d\boldsymbol{\theta}_1 \quad (\text{S78}) \\
&= \int_{\mathbb{R}^d} \int_{\mathbb{R}^p} f(\mathbf{x}_1) \Pi_{\rho}(\boldsymbol{\theta}_1 | \mathbf{z}_1) \pi_{\rho}^{\mathbf{z}}(\mathbf{z}_1) d\mathbf{z}_1 d\boldsymbol{\theta}_1 \\
&= \int_{\mathbb{R}^{d+p}} f(\mathbf{x}_1) \Pi_{\rho}(\boldsymbol{\theta}_1, \mathbf{z}_1) d\mathbf{z}_1 d\boldsymbol{\theta}_1 = \mathbb{E}[f(\tilde{X}_0)] ,
\end{aligned}$$

where we have used (S77) in (S78). Therefore, X_1 has distribution Π_{ρ} and the Markov kernel $\tilde{P}_{\rho, \gamma}$ admits Π_{ρ} as a stationary distribution, which completes the proof. \square

Define by induction the synchronous coupling $(X_n = (\theta_n, Z_n))_{n \geq 0}, (\tilde{X}_n = (\tilde{\theta}_n, \tilde{Z}_n))_{n \geq 0}$, starting from $(\theta_0, Z_0) = (\boldsymbol{\theta}, \mathbf{z}), (\tilde{\theta}_0, \tilde{Z}_0)$ distributed according to Π_{ρ} , for any $i \in [b]$ and $n \geq 0$, as

$$\begin{aligned}
\tilde{Z}_{n+1}^i &= \tilde{Y}_{N_i \gamma_i}^{(i, n)} , & \tilde{\theta}_{n+1} &= \bar{\mathbf{B}}_0^{-1} \mathbf{B}_0^{\top} \tilde{\mathbf{D}}_0^{1/2} \tilde{Z}_{n+1} + \bar{\mathbf{B}}_0^{-1/2} \xi_{n+1} , \\
Z_{n+1}^i &= Y_{N_i \gamma_i}^{(i, n)} , & \theta_{n+1} &= \bar{\mathbf{B}}_0^{-1} \mathbf{B}_0^{\top} \tilde{\mathbf{D}}_0^{1/2} Z_{n+1} + \bar{\mathbf{B}}_0^{-1/2} \xi_{n+1} ,
\end{aligned} \quad (\text{S79})$$

where we consider for any $i \in [b], k \in \mathbb{N}$, for $t \in [k\gamma_i, (k+1)\gamma_i)$

$$\begin{aligned}
\tilde{Y}_t^{(i, n)} &= \tilde{Y}_{k\gamma_i}^{(i, n)} - \int_{k\gamma_i}^t \nabla V_i(\tilde{Y}_l^{(i, n)}) dl + (t - k\gamma_i)(\rho_i)^{-1} \mathbf{A}_i \tilde{\theta}_n + 2^{1/2} (B_t^{(i, n)} - B_{k\gamma_i}^{(i, n)}) , \\
Y_t^{(i, n)} &= Y_{k\gamma_i}^{(i, n)} - (t - k\gamma_i) \nabla V_i(Y_{k\gamma_i}^{(i, n)}) + (t - k\gamma_i)(\rho_i)^{-1} \mathbf{A}_i \theta_n + 2^{1/2} (B_t^{(i, n)} - B_{k\gamma_i}^{(i, n)}) .
\end{aligned} \quad (\text{S80})$$

Let $\mathcal{G}_0 = \sigma(Z_0, \tilde{Z}_0, \theta_0, \tilde{\theta}_0)$, for any $n \in \mathbb{N}^*$, let

$$\mathcal{G}_n = \sigma\{(Z_0, \tilde{Z}_0, \theta_0, \tilde{\theta}_0), (B_t^{(i, k)})_{t \geq 0} : i \in [b], k \leq n\} , \quad (\text{S81})$$

and for any $t \geq 0$, let $\mathcal{H}_t^{(n)} = \sigma\{(B_s^{(i, n)})_{s \leq t} : i \in [b]\}$, and

$$\mathcal{F}_t^{(n)} \text{ the } \sigma\text{-field generated by } \mathcal{H}_t^{(n)} \text{ and } \mathcal{G}_{n-1} . \quad (\text{S82})$$

Note that X_n and \tilde{X}_n are distributed according to $\Pi_{\rho} \tilde{P}_{\rho}^n$ and $\delta_{\tilde{\mathbf{x}}} P_{\rho, \gamma, \mathbf{N}}^n$, respectively. Hence, by definition of the Wasserstein distance of order 2, it follows since $\Pi_{\rho} \tilde{P}_{\rho}^n = \Pi_{\rho}$ by Proposition S21 that

$$W_2(\Pi_{\rho}, \delta_{\tilde{\mathbf{x}}} P_{\rho, \gamma, \mathbf{N}}^n) \leq \mathbb{E} \left[\|X_n - \tilde{X}_n\|^2 \right]^{1/2} . \quad (\text{S83})$$

We start this section by a first estimate on $\mathbb{E}[\|X_n - \tilde{X}_n\|^2]^{1/2}$ and some technical results needed for the proof of Proposition 4 and Proposition 5. The following result holds regarding the process $(\tilde{Y}_t^{(i,n)})_{t \in \mathbb{R}_+}$ defined, for any $i \in [b]$ and $n \in \mathbb{N}$, in (S80).

Lemma S22. *Assume **H1-H2**. For $i \in [b], n \in \mathbb{N}$, denote by $\mathbf{z}_{n,\star}^i$ the unique minimiser of $\mathbf{z}_i \in \mathbb{R}^{d_i} \mapsto U_i(\mathbf{z}_i) + \|\mathbf{z}_i - \mathbf{A}_i \tilde{\theta}_n\|/(2\rho_i)$. Then, for any $i \in [b], k \in \mathbb{N}$ and $n \in \mathbb{N}$,*

$$\mathbb{E}^{\mathcal{G}_n} [\|\tilde{Y}_{k\gamma_i}^{(i,n)} - \mathbf{z}_{n,\star}^i\|^2] \leq d_i/\tilde{m}_i. \quad (\text{S84})$$

where \tilde{m}_i is defined in (S25).

Proof. Let $n \in \mathbb{N}$. By Durmus and Moulines (2019, Proposition 1), for $i \in [b]$ and $k \in \mathbb{N}$, we have

$$\mathbb{E}^{\mathcal{F}_{k\gamma_i}^{(n)}} \|\tilde{Y}_{k\gamma_i}^{(i,n)} - \mathbf{z}_{n,\star}^i\|^2 \leq \|\tilde{Z}_n^i - \mathbf{z}_{n,\star}^i\|^2 e^{-2k\gamma_i \tilde{m}_i} + (d_i/\tilde{m}_i)(1 - e^{-2k\gamma_i \tilde{m}_i}). \quad (\text{S85})$$

By (S80), using Proposition S21 we get that \tilde{X}_n has distribution $\Pi_{\boldsymbol{\rho}}$, therefore given $\tilde{\theta}_n$, \tilde{Z}_n has distribution $\Pi_{\boldsymbol{\rho}}(\cdot|\tilde{\theta}_n)$. Then, using (S85), Durmus and Moulines (2019, Proposition 1(ii)) combined with **H2**, and since $(\tilde{Z}_n^1, \dots, \tilde{Z}_n^b)$ are independent given $\tilde{\theta}_n$, we get the stated result. \square

Lemma S23. *Assume **H1** and let $\mathbf{N} \in (\mathbb{N}^*)^b, \boldsymbol{\gamma} \in (\mathbb{R}_+^*)^b$. Then, for any $n \in \mathbb{N}$, the random variables $X_n = (\theta_n^\top, Z_n^\top)^\top, \tilde{X}_n = (\tilde{\theta}_n^\top, \tilde{Z}_n^\top)^\top$ defined in (S79) satisfy*

$$\|\tilde{X}_{n+1} - X_{n+1}\|^2 \leq (1 + \|\bar{\mathbf{B}}_0^{-1} \mathbf{B}_0^\top \tilde{\mathbf{D}}_0^{1/2}\|^2) \|\tilde{Z}_{n+1} - Z_{n+1}\|^2,$$

where $\bar{\mathbf{B}}_0, \mathbf{B}_0, \tilde{\mathbf{D}}_0$ are defined in (S2)-(S3).

Proof. The proof is similar to the proof of Lemma S3 and is omitted. \square

For any $k, n \in \mathbb{N}, s \in \mathbb{R}_+$ consider the $p \times p$ matrices defined by

$$\mathbf{J}(k, s) = \text{diag} \left(\mathbb{1}_{[N_1]}(k+1) \mathbb{1}_{[0, \gamma_1]}(s) \cdot \mathbf{I}_{d_1}, \dots, \mathbb{1}_{[N_b]}(k+1) \mathbb{1}_{[0, \gamma_b]}(s) \cdot \mathbf{I}_{d_b} \right), \quad (\text{S86})$$

$$\mathbf{H}_{U,k}^{(n)} = \text{diag} \left(\gamma_1 \int_0^1 \nabla^2 U_1((1-s)Y_{k\gamma_1}^{(1,n)} + s\tilde{Y}_{k\gamma_1}^{(1,n)}) \, ds, \right. \quad (\text{S87})$$

$$\left. \dots, \gamma_b \int_0^1 \nabla^2 U_b((1-s)Y_{k\gamma_b}^{(b,n)} + s\tilde{Y}_{k\gamma_b}^{(b,n)}) \, ds \right),$$

$$\mathbf{C}_k^{(n)} = \mathbf{J}(k, 0)(\mathbf{D}_{\boldsymbol{\gamma}/\boldsymbol{\rho}} + \mathbf{H}_{U,k}^{(n)}), \quad (\text{S88})$$

$$\mathbf{M}_{k+1}^{(n)} = (\mathbf{I}_p - \mathbf{C}_0^{(n)})^{-1} \dots (\mathbf{I}_p - \mathbf{C}_k^{(n)})^{-1}, \quad \text{with } \mathbf{M}_0^{(n)} = \mathbf{I}_p. \quad (\text{S89})$$

Similarly to (S19), for $n, k \in \mathbb{N}$ and $i \in [b]$, consider $\mathbf{C}_k^{(i,n)}$ corresponding to the i -th diagonal block of $\mathbf{C}_k^{(n)}$ defined in (S88), *i.e.*

$$\mathbf{C}_k^{(i,n)} = \mathbb{1}_{[N_i]}(k+1)\gamma_i \left\{ \rho_i^{-1} \mathbf{I}_{d_i} + \int_0^1 \nabla^2 U_i((1-s)Y_{k\gamma_i}^{(i,n)} + s\tilde{Y}_{k\gamma_i}^{(i,n)}) ds \right\} \in \mathbb{R}^{d_i \times d_i}, \quad (\text{S90})$$

where, for any $n \in \mathbb{N}$ and $i \in [b]$, $(Y_{k\gamma_i}^{(i,n)}, \tilde{Y}_{k\gamma_i}^{(i,n)})_{k \in \mathbb{N}}$ is defined in (S80).

Lemma S24. Assume **H1-H2** and let $\gamma \in (\mathbb{R}_+^*)^b$ such that, for any $i \in [b]$, $\gamma_i < 1/\tilde{M}_i$. Then, for any $n, k \in \mathbb{N}$, the matrix $(\mathbf{I}_p - \mathbf{C}_k^{(n)})$ is invertible and in addition, for any $i \in [b]$, we have

$$\|\mathbf{I}_{d_i} - \mathbf{C}_k^{(i,n)}\| \leq 1 - \gamma_i \tilde{m}_i,$$

where $\mathbf{C}_k^{(i,n)}$ is defined in (S90).

Proof. Let $i \in [b], n, k \in \mathbb{N}$. By **H2**, we have $\|\nabla^2 U_i\| \leq M_i$ which implies by (S90) that $\|\mathbf{C}_k^{(i,n)}\| \leq \gamma_i \tilde{M}_i$. Since $\gamma_i < 1/\tilde{M}_i$, the matrix $\mathbf{I}_p - \mathbf{C}_k^{(i,n)}$ is invertible and so is $\mathbf{I}_p - \mathbf{C}_k^{(n)}$. In addition, following the same lines as the proof of Lemma S9 implies $\|\mathbf{I}_{d_i} - \mathbf{C}_k^{(i,n)}\| \leq \max\{|1 - \gamma_i \tilde{m}_i|, |1 - \gamma_i \tilde{M}_i|\} = 1 - \gamma_i \tilde{m}_i$. \square

For any $n, k \in \mathbb{N}, i \in [b]$, if $\gamma_i \in (0, 1/\tilde{M}_i)$, Lemma S24 shows the invertibility of the matrices $\mathbf{I}_p - \mathbf{C}_k^{(n)}$. Therefore, $\mathbf{M}_\infty^{(n)}$ is invertible and we can define

$$\mathbf{T}_1^{(n)} = [\mathbf{M}_\infty^{(n)}]^{-1} + \sum_{k=0}^{\infty} [\mathbf{M}_\infty^{(n)}]^{-1} \mathbf{M}_{k+1}^{(n)} \mathbf{J}(k, 0) \mathbf{D}_N^{-1/2} \mathbf{D}_{\gamma/\rho}^{1/2} \mathbf{P}_0 \mathbf{D}_{\gamma/\rho}^{1/2} \mathbf{D}_N^{1/2}, \quad (\text{S91})$$

$$\mathbf{T}_2^{(n)} = \sum_{k=0}^{\infty} \left\{ [\mathbf{M}_\infty^{(n)}]^{-1} \mathbf{M}_{k+1}^{(n)} \mathbf{D}_{N\gamma}^{-1/2} \int_0^{+\infty} \mathbf{J}(k, l) [\nabla V(\tilde{Y}_{k\gamma+l}^{(n)}) - \nabla V(\tilde{Y}_{k\gamma}^{(n)})] dl \right\}. \quad (\text{S92})$$

Using these matrices, we have the following result.

Lemma S25. Assume **H1-H2** and let $\mathbf{N} \in (\mathbb{N}^*)^b, \gamma \in (\mathbb{R}_+^*)^b$ such that, for any $i \in [b]$, $\gamma_i < 1/\tilde{M}_i$. Then, for any $n \geq 1$,

$$\mathbf{D}_{N\gamma}^{-1/2} (\tilde{Z}_{n+1} - Z_{n+1}) = \mathbf{T}_1^{(n)} (\tilde{Z}_n - Z_n) - \mathbf{T}_2^{(n)}, \quad (\text{S93})$$

where $(Z_n, \tilde{Z}_n)_{n \in \mathbb{N}}$ is defined in (S79) and $\mathbf{D}_{N\gamma} = \text{diag}(N_1 \gamma_1 \mathbf{I}_{d_1}, \dots, N_b \gamma_b \mathbf{I}_{d_b}) \in \mathbb{R}^{p \times p}$.

Proof. Let $i \in [b]$ and $n \geq 1$. Recall that V_i is defined in (S20) and for $\mathbf{z} \in \mathbb{R}^p$, denote $V(\mathbf{z}) = \sum_{i=1}^b V_i(\mathbf{z}_i)$. For any $k \in \mathbb{N}$, we have

$$\nabla V_i(\tilde{Y}_{k\gamma_i}^{(i,n)}) - \nabla V_i(Y_{k\gamma_i}^{(i,n)}) = \left[\int_0^1 \nabla^2 V_i((1-s)Y_{k\gamma_i}^{(i,n)} + s\tilde{Y}_{k\gamma_i}^{(i,n)}) ds \right] (\tilde{Y}_{k\gamma_i}^{(i,n)} - Y_{k\gamma_i}^{(i,n)}).$$

For $k \geq 0$, it follows from (S80) that

$$\begin{aligned} \tilde{Y}_{(k+1)\gamma_i}^{(i,n)} - Y_{(k+1)\gamma_i}^{(i,n)} &= \left(\mathbf{I}_{d_i} - \gamma_i \int_0^1 \nabla^2 V_i((1-s)Y_{k\gamma_i}^{(i,n)} + s\tilde{Y}_{k\gamma_i}^{(i,n)}) ds \right) (\tilde{Y}_{k\gamma_i}^{(i,n)} - Y_{k\gamma_i}^{(i,n)}) \\ &\quad - \int_0^{\gamma_i} [\nabla V_i(\tilde{Y}_{k\gamma_i+l}^{(i,n)}) - \nabla V_i(Y_{k\gamma_i+l}^{(i,n)})] dl + (\gamma_i/\rho_i) \mathbf{A}_i(\tilde{\theta}_n - \theta_n). \end{aligned} \quad (\text{S94})$$

Consider the process $(\tilde{Y}_t^{(n)}, Y_t^{(n)})_{t \in \mathbb{R}_+}$ valued in $\mathbb{R}^p \times \mathbb{R}^p$ and defined for any $t \geq 0$ by

$$\tilde{Y}_t^{(n)} = \tilde{Y}_{\min(t, N_i \gamma_i)}^{(n)}, \quad Y_t^{(n)} = Y_{\min(t, N_i \gamma_i)}^{(n)}. \quad (\text{S95})$$

The process (S95) is continuous with respect to t and defined so that its component $(\tilde{Y}_t^{(i,n)}, Y_t^{(i,n)})$ equals (\tilde{Y}_t^i, Y_t^i) for $t \leq N_i \gamma_i$ and is constant for $t > N_i \gamma_i$. For $l \geq 0$, we write $(\tilde{Y}_{k\gamma+l}^{(n)}, Y_{k\gamma+l}^{(n)}) = (\tilde{Y}_{k\gamma_i+l}^{(i,n)}, Y_{k\gamma_i+l}^{(i,n)})_{i \in [b]} \in \mathbb{R}^p \times \mathbb{R}^p$. Using the matrices defined in (S89), for $k \in \mathbb{N}$, we obtain

$$\begin{aligned} \tilde{Y}_{(k+1)\gamma}^{(n)} - Y_{(k+1)\gamma}^{(n)} &= (\mathbf{I}_p - \mathbf{C}_k^{(n)}) (\tilde{Y}_{k\gamma}^{(n)} - Y_{k\gamma}^{(n)}) - \int_0^\infty \mathbf{J}(k, l) [\nabla V(\tilde{Y}_{k\gamma+l}^{(n)}) - \nabla V(Y_{k\gamma+l}^{(n)})] dl \\ &\quad + \mathbf{J}(k, 0) \mathbf{D}_{\gamma/\sqrt{\rho}} \mathbf{P}_0 \tilde{\mathbf{D}}_0^{1/2} (\tilde{Y}_0^{(n)} - Y_0^{(n)}), \end{aligned} \quad (\text{S96})$$

where \mathbf{P}_0 is defined in (S3). Recall the matrix $\mathbf{M}_k^{(n)}$ defined in (S89) with $\mathbf{M}_0^{(n)} = \mathbf{I}_p$ and for $k \geq 1$, $\mathbf{M}_k^{(n)} = (\mathbf{I}_p - \mathbf{C}_0^{(n)})^{-1} \dots (\mathbf{I}_p - \mathbf{C}_{k-1}^{(n)})^{-1}$. By multiplying (S96) by $\mathbf{M}_{k+1}^{(n)} \mathbf{D}_{N\gamma}^{-1/2}$, we have

$$\begin{aligned} \mathbf{M}_{k+1}^{(n)} \mathbf{D}_{N\gamma}^{-1/2} (\tilde{Y}_{(k+1)\gamma}^{(n)} - Y_{(k+1)\gamma}^{(n)}) &= \mathbf{M}_k^{(n)} \mathbf{D}_{N\gamma}^{-1/2} (\tilde{Y}_{k\gamma}^{(n)} - Y_{k\gamma}^{(n)}) \\ &\quad - \mathbf{M}_{k+1}^{(n)} \mathbf{D}_{N\gamma}^{-1/2} \int_0^\infty \mathbf{J}(k, l) [\nabla V(\tilde{Y}_{k\gamma+l}^{(n)}) - \nabla V(Y_{k\gamma+l}^{(n)})] dl \\ &\quad + \mathbf{M}_{k+1}^{(n)} \mathbf{J}(k, 0) \mathbf{D}_N^{-1/2} \mathbf{D}_{\gamma/\rho}^{1/2} \mathbf{P}_0 \tilde{\mathbf{D}}_0^{1/2} (\tilde{Y}_0^{(n)} - Y_0^{(n)}). \end{aligned}$$

By (S95) and (S79), we have for $t \geq \max_{i \in [b]} \{\gamma_i N_i\}$, $(\tilde{Z}_{n+1}, Z_{n+1}) = (\tilde{Y}_t, Y_t)$. Therefore, summing the previous expression over k , we get

$$\begin{aligned} \mathbf{M}_\infty^{(n)} \mathbf{D}_{N\gamma}^{-1/2} (\tilde{Z}_{n+1} - Z_{n+1}) &= - \sum_{k=0}^\infty \mathbf{M}_{k+1}^{(n)} \mathbf{D}_{N\gamma}^{-1/2} \int_0^\infty \mathbf{J}(k, l) [\nabla V(\tilde{Y}_{k\gamma+l}^{(n)}) - \nabla V(Y_{k\gamma+l}^{(n)})] dl \\ &\quad + \left[\mathbf{M}_0^{(n)} + \sum_{k=0}^\infty \mathbf{M}_{k+1}^{(n)} \mathbf{J}(k, 0) \mathbf{D}_N^{-1/2} \mathbf{D}_{\gamma/\rho}^{1/2} \mathbf{P}_0 \mathbf{D}_{\gamma/\rho}^{1/2} \mathbf{D}_N^{1/2} \right] \mathbf{D}_{N\gamma}^{-1/2} \cdot (\tilde{Z}_n - Z_n). \end{aligned}$$

By Lemma S24, $\mathbf{M}_\infty^{(n)}$ is invertible and the proof is concluded by multiplying the previous equality by $[\mathbf{M}_\infty^{(n)}]^{-1}$. \square

Based on Lemma S25, we have the following relation between $\|\tilde{Z}_{n+1} - Z_{n+1}\|^2$ and $\|\tilde{Z}_n - Z_n\|^2$.

Lemma S26. Assume **H1-H2** and let $\mathbf{N} \in (\mathbb{N}^*)^b, \gamma \in (\mathbb{R}_+^*)^b$ such that, for any $i \in [b], \gamma_i < 1/\tilde{M}_i$. Then, for any $\epsilon > 0$ and $n \geq 1$,

$$\|\tilde{Z}_{n+1} - Z_{n+1}\|_{\mathbf{D}_{\mathbf{N}\gamma}^{-1}}^2 \leq (1 + 2\epsilon)\|\mathbf{T}_1^{(n)}\|^2\|\tilde{Z}_n - Z_n\|_{\mathbf{D}_{\mathbf{N}\gamma}^{-1}}^2 + (1 + 1/\{2\epsilon\})\|\mathbf{T}_2^{(n)}\|^2.$$

where $(Z_n, \tilde{Z}_n)_{n \in \mathbb{N}}$ is defined in (S79) and $\mathbf{D}_{\mathbf{N}\gamma} = \text{diag}(N_1\gamma_1\mathbf{I}_{d_1}, \dots, N_b\gamma_b\mathbf{I}_{d_b}) \in \mathbb{R}^{p \times p}$.

Proof. The proof follows from Lemma S25 and by using the fact that for $\mathbf{a}, \mathbf{b} \in \mathbb{R}^p, \epsilon > 0$ we have $2\langle \mathbf{a}, \mathbf{b} \rangle \leq 2\epsilon\|\mathbf{a}\|^2 + (1/\{2\epsilon\})\|\mathbf{b}\|^2$. \square

Similarly to Lemma S10, we have the following result regarding the contracting term.

Lemma S27. Assume **H1-H2** and let $\mathbf{N} \in (\mathbb{N}^*)^b, \gamma \in (\mathbb{R}_+^*)^b$ such that, for any $i \in [b], \gamma_i < 1/\tilde{M}_i$ and $N_i\gamma_i \leq 2/(m_i + \tilde{M}_i)$. Then, for any $n \geq 0$, we have

$$\|\mathbf{T}_1^{(n)}\| \leq 1 - \min_{i \in [b]} \{N_i\gamma_i m_i\} + r_{\gamma, \rho, \mathbf{N}},$$

where $\mathbf{T}_1^{(n)}$ and $r_{\gamma, \rho, \mathbf{N}}$ are defined in (S91) and (S33), respectively.

Proof. The proof is similar to the proof of Lemma S10 and therefore is omitted. \square

In the next lemma, we upper bound the coefficient $r_{\gamma, \rho, \mathbf{N}}$ defined in (S33). For this, we explicit a choice of \mathbf{N} that we denote $\mathbf{N}^* = (N_1^*(\gamma_1), \dots, N_b^*(\gamma_b)) \in (\mathbb{N}^*)^b$ defined for any $i \in [b]$, any $\gamma_i > 0$, by

$$N_i^*(\gamma_i) = \lfloor m_i \min_{i \in [b]} \{m_i/\tilde{M}_i\}^2 / (20\gamma_i \tilde{M}_i^2 \max_{i \in [b]} \{m_i/\tilde{M}_i\}^2) \rfloor, \quad (\text{S97})$$

where $\tilde{M}_i = M_i + 1/\rho_i$.

Lemma S28. Assume **H1-H2** and let $\gamma \in (\mathbb{R}_+^*)^b$ such that, for any $i \in [b]$,

$$\gamma_i \leq \frac{m_i}{40\tilde{M}_i^2} \left(\frac{\min_{i \in [b]} \{m_i/\tilde{M}_i\}}{\max_{i \in [b]} \{m_i/\tilde{M}_i\}} \right)^2.$$

Then, for any $i \in [b]$, we have $N_i^*(\gamma_i) \in \mathbb{N}^*$ and

$$r_{\gamma, \rho, \mathbf{N}^*} < \min_{i \in [b]} \{N_i^*(\gamma_i)\gamma_i m_i\} / 2,$$

where $r_{\gamma, \rho, \mathbf{N}^*}$ is defined in (S33).

Proof. The assumption on γ_i combined with the definition (S97) of $N_i^*(\gamma_i)$ imply $N_i^*(\gamma_i) \geq 2$, using in addition $m_i \leq M_i$, $\max_{i \in [b]} \{N_i^*(\gamma_i) \gamma_i \tilde{M}_i \mathbb{1}_{N_i^*(\gamma_i) > 1}\} \leq 1/20$ and

$$\begin{aligned} \frac{1}{20} \left(\frac{\min_{i \in [b]} \{m_i / \tilde{M}_i\}}{\max_{i \in [b]} \{m_i / \tilde{M}_i\}} \right)^2 &\geq \frac{N_i^*(\gamma_i) \gamma_i \tilde{M}_i^2}{m_i} > \frac{1}{20} \left(\frac{\min_{i \in [b]} \{m_i / \tilde{M}_i\}}{\max_{i \in [b]} \{m_i / \tilde{M}_i\}} \right)^2 - \frac{\gamma_i \tilde{M}_i^2}{m_i} \\ &\geq \frac{1}{40} \left(\frac{\min_{i \in [b]} \{m_i / \tilde{M}_i\}}{\max_{i \in [b]} \{m_i / \tilde{M}_i\}} \right)^2. \end{aligned} \quad (\text{S98})$$

Using the definition (S33) of $r_{\gamma, \rho, \mathbf{N}}$, we have $r_{\gamma, \rho, \mathbf{N}} < 5 \max_{i \in [b]} \{N_i^*(\gamma_i) \gamma_i \tilde{M}_i \mathbb{1}_{N_i^*(\gamma_i) > 1}\}^2$. Thus, plugging (S98) in the previous inequality gives

$$r_{\gamma, \rho, \mathbf{N}} \leq \max_{i \in [b]} \{m_i / \tilde{M}_i\}^2 \max_{i \in [b]} \left\{ \frac{N_i^*(\gamma_i) \gamma_i \tilde{M}_i^2}{m_i} \right\} < \frac{\min_{i \in [b]} \{m_i / \tilde{M}_i\}^4}{80 \max_{i \in [b]} \{m_i / \tilde{M}_i\}^2}. \quad (\text{S99})$$

In addition, (S98) also shows that

$$\frac{1}{40} \left(\frac{\min_{i \in [b]} \{m_i / \tilde{M}_i\}}{\max_{i \in [b]} \{m_i / \tilde{M}_i\}} \right)^2 \left(\frac{m_i}{\tilde{M}_i} \right)^2 \leq N_i^*(\gamma_i) \gamma_i m_i. \quad (\text{S100})$$

Therefore, combining (S99) and (S100) completes the proof. \square

S4.2 Proof of Proposition 4

We first give the formal statement of Proposition 4.

Proposition S29. Assume **H1-H2** and let $\gamma \in (\mathbb{R}_+^*)^b$, $\mathbf{N} \in (\mathbb{N}^*)^b$ such that for any $i \in [b]$, $\gamma_i \leq m_i / 40 \tilde{M}_i^2 (\min_{i \in [b]} \{m_i / \tilde{M}_i\} / \max_{i \in [b]} \{m_i / \tilde{M}_i\})^2$ and $N_i = \lfloor m_i \min_{i \in [b]} \{m_i / \tilde{M}_i\}^2 / (20 \gamma_i \tilde{M}_i^2 \max_{i \in [b]} \{m_i / \tilde{M}_i\}) \rfloor$. Then, we have

$$\begin{aligned} W_2^2(\Pi_{\rho, \gamma, \mathbf{N}}, \Pi_{\rho}) &\leq \frac{4(1 + \|\bar{\mathbf{B}}_0^{-1} \mathbf{B}_0^\top \tilde{\mathbf{D}}_0^{1/2}\|^2) \max_{i \in [b]} \{m_i / \tilde{M}_i^2\}}{5 \min_{i \in [b]} \{m_i / \tilde{M}_i\}^2 \max_{i \in [b]} \{m_i / \tilde{M}_i\}^2} \\ &\quad \times \sum_{i=1}^b d_i \gamma_i m_i (1 + \gamma_i^2 \tilde{M}_i^2 / 12 + \gamma_i \tilde{M}_i^2 / (2 \tilde{m}_i)), \end{aligned}$$

where $\bar{\mathbf{B}}_0, \mathbf{B}_0, \tilde{\mathbf{D}}_0$ are defined in (S2)-(S3), and for any $i \in [b]$, \tilde{m}_i, \tilde{M}_i are defined in (S25).

By Lemma S23 and Lemma S26, we can note that the proof of Proposition S29 boils down to derive an upper bound on $\|\mathbf{T}_2^{(n)}\|^2$ defined in (S92) for $n \in \mathbb{N}$. The following lemma provides such a bound.

Lemma S30. Assume **H1-H2** and let $\mathbf{N} \in (\mathbb{N}^*)^b, \gamma \in (\mathbb{R}_+^*)^b$ such that, for any $i \in [b]$, $\gamma_i < 1/\tilde{M}_i$. Then, for any $n \in \mathbb{N}$, we have

$$\mathbb{E} \left[\|\mathbf{T}_2^{(n)}\|^2 \right] \leq \sum_{i=1}^b d_i N_i \gamma_i^2 \tilde{M}_i^2 \left[1 + \gamma_i^2 \tilde{M}_i^2 / 12 + \gamma_i \tilde{M}_i^2 / (2\tilde{m}_i) \right],$$

where $\tilde{m}_i, \tilde{M}_i, \mathbf{T}_2^{(n)}$ are defined in (S25) and (S92), respectively.

Proof. Let $n \in \mathbb{N}$. Using (S86), we can write, for any $l \in \mathbb{R}_+$ and $k \in \mathbb{N}$, $\mathbf{J}(k, l)$ as a block-diagonal matrix $\text{diag}(\mathbf{J}^1(k, l), \dots, \mathbf{J}^b(k, l))$ with $\mathbf{J}^i(k, l) = \mathbf{1}_{[N_i]}(k+1) \mathbf{1}_{[0, \gamma_i]}(s) \cdot \mathbf{I}_{d_i}$ for any $i \in [b]$. By (S89) and using for any $k \in \mathbb{N}$, that $[\mathbf{M}_\infty^{(n)}]^{-1} \mathbf{M}_{k+1}^{(n)} = \prod_{l=k+1}^\infty (\mathbf{I}_{d_i} - \mathbf{C}_l^{(i,n)})$ is finite by (S88), we have

$$\begin{aligned} \|\mathbf{T}_2^{(n)}\|^2 &= \left\| \sum_{k=0}^\infty [\mathbf{M}_\infty^{(n)}]^{-1} \mathbf{M}_{k+1}^{(n)} \mathbf{D}_{\mathbf{N}\gamma}^{-1/2} \int_0^\infty \mathbf{J}(k, l) [\nabla V(\tilde{Y}_{k\gamma+l}^{(n)}) - \nabla V(\tilde{Y}_{k\gamma}^{(n)})] dl \right\|^2 \\ &= \sum_{i=1}^b \frac{1}{N_i \gamma_i} \left\| \sum_{k=0}^\infty \prod_{l=k+1}^\infty (\mathbf{I}_{d_i} - \mathbf{C}_l^{(i,n)}) \int_0^{\gamma_i} \mathbf{J}^i(k, 0) [\nabla V_i(\tilde{Y}_{k\gamma_i+l}^{(i,n)}) - \nabla V_i(\tilde{Y}_{k\gamma_i}^{(i,n)})] dl \right\|^2. \end{aligned} \quad (\text{S101})$$

Since for any $i \in [b]$, $k \geq N_i$ we have $\mathbf{J}^i(k, 0) = \mathbf{C}_l^{(i,n)} = \mathbf{0}_{d_i \times d_i}$, (S101) can be rewritten as

$$\|\mathbf{T}_2^{(n)}\|^2 = \sum_{i=1}^b \frac{1}{N_i \gamma_i} \left\| \sum_{k=0}^{N_i-1} \prod_{l=k+1}^{N_i-1} (\mathbf{I}_{d_i} - \mathbf{C}_l^{(i,n)}) \int_0^{\gamma_i} \mathbf{J}^i(k, 0) [\nabla V_i(\tilde{Y}_{k\gamma_i+l}^{(i,n)}) - \nabla V_i(\tilde{Y}_{k\gamma_i}^{(i,n)})] dl \right\|^2,$$

and the Cauchy-Schwarz inequality gives

$$\|\mathbf{T}_2^{(n)}\|^2 \leq \sum_{i=1}^b \frac{1}{\gamma_i} \left(\sum_{k=0}^{N_i-1} \left\| \prod_{l=k+1}^{N_i-1} (\mathbf{I}_{d_i} - \mathbf{C}_l^{(i,n)}) \right\|^2 \left\| \int_0^{\gamma_i} [\nabla V_i(\tilde{Y}_{k\gamma_i+l}^{(i,n)}) - \nabla V_i(\tilde{Y}_{k\gamma_i}^{(i,n)})] dl \right\|^2 \right). \quad (\text{S102})$$

Since, for any $i \in [b]$, $\gamma_i \tilde{M}_i < 1$, we get using Lemma S24,

$$\left\| \prod_{l=k+1}^{N_i-1} (\mathbf{I}_{d_i} - \mathbf{C}_l^{(i,n)}) \right\|^2 \leq \{1 - \gamma_i \tilde{m}_i\}^{2(N_i-k-1)}.$$

By combining (S102) with the previous result and the Jensen inequality, we have

$$\|\mathbf{T}_2^{(n)}\|^2 \leq \sum_{i=1}^b \sum_{k=0}^{N_i-1} \{1 - \gamma_i \tilde{m}_i\}^{2(N_i-k-1)} \int_0^{\gamma_i} \left\| \nabla V_i(\tilde{Y}_{k\gamma_i+l}^{(i,n)}) - \nabla V_i(\tilde{Y}_{k\gamma_i}^{(i,n)}) \right\|^2 dl. \quad (\text{S103})$$

For $i \in [b]$, using [Durmus and Moulines \(2019, Lemma 21\)](#) applied to the potential $V_i^\theta : \mathbf{y}^i \mapsto U_i(\mathbf{y}^i) + \|\mathbf{y}^i - \mathbf{A}_i \theta\|^2 / (2\rho_i)$ yields

$$\begin{aligned} \int_0^{\gamma_i} \mathbb{E}^{\mathcal{F}_{k\gamma_i}^{(n)}} \|\nabla V_i(\tilde{Y}_{k\gamma_i+l}^{(i,n)}) - \nabla V_i(\tilde{Y}_{k\gamma_i}^{(i,n)})\|^2 dl &= \int_0^{\gamma_i} \mathbb{E}^{\mathcal{F}_{k\gamma_i}^{(n)}} \|\nabla V_i^{\tilde{\theta}_n}(\tilde{Y}_{k\gamma_i+l}^{(i,n)}) - \nabla V_i^{\tilde{\theta}_n}(\tilde{Y}_{k\gamma_i}^{(i,n)})\|^2 dl \\ &\leq \gamma_i^2 \tilde{M}_i^2 \left[d_i + d_i \gamma_i^2 \tilde{M}_i^2 / 12 + (\gamma_i \tilde{M}_i^2 / 2) \|\tilde{Y}_{k\gamma_i}^{(i,n)} - \mathbf{z}_{n,\star}^i\|^2 \right], \end{aligned} \quad (\text{S104})$$

where $\mathbf{z}_{n,\star}^i = \arg \min_{\mathbf{z}_i \in \mathbb{R}^{d_i}} V_i^{\tilde{\theta}_n}(\mathbf{z}_i)$.

By (S104), (S84), Lemma S22 and since $\max_{i \in [b]} \gamma_i \tilde{m}_i < 1$, we get

$$\begin{aligned} \sum_{i=1}^b \sum_{k=0}^{N_i-1} \{1 - \gamma_i \tilde{m}_i\}^{2(N_i-k-1)} \int_0^{\gamma_i} \mathbb{E} \|\nabla V_i(\tilde{Y}_{k\gamma_i+l}^{(i,n)}) - \nabla V_i(\tilde{Y}_{k\gamma_i}^{(i,n)})\|^2 dl \\ \leq \sum_{i=1}^b d_i N_i \gamma_i^2 \tilde{M}_i^2 [1 + \gamma_i^2 \tilde{M}_i^2 / 12 + \gamma_i \tilde{M}_i^2 / (2\tilde{m}_i)]. \end{aligned}$$

Combining this result with (S103) completes the proof. \square

We can now combine Lemma S30 and Lemma S27 with Lemma S26 to get the following bound.

Lemma S31. *Assume **H1-H2** and let $\mathbf{N} \in (\mathbb{N}^*)^b$, $\boldsymbol{\gamma} \in (\mathbb{R}_+^*)^b$ such that, for any $i \in [b]$, $\gamma_i < 1/\tilde{M}_i$, $N_i \gamma_i \leq 2/(m_i + \tilde{M}_i)$. Suppose in addition $\kappa_{\boldsymbol{\gamma}, \boldsymbol{\rho}, \mathbf{N}} = \min_{i \in [b]} \{N_i \gamma_i m_i\} - r_{\boldsymbol{\gamma}, \boldsymbol{\rho}, \mathbf{N}} \in (0, 1)$, where $r_{\boldsymbol{\gamma}, \boldsymbol{\rho}, \mathbf{N}}$ is defined in (S33). Then, for $n \geq 1$, we have*

$$\begin{aligned} \mathbb{E} \left[\|\tilde{Z}_n - Z_n\|_{\mathbf{D}_{\mathbf{N}\boldsymbol{\gamma}}^{-1}}^2 \right] &\leq (1 - \kappa_{\boldsymbol{\gamma}, \boldsymbol{\rho}, \mathbf{N}} + \kappa_{\boldsymbol{\gamma}, \boldsymbol{\rho}, \mathbf{N}}^2 / 2)^{2(n-1)} \mathbb{E} \left[\|\tilde{Z}_1 - Z_1\|_{\mathbf{D}_{\mathbf{N}\boldsymbol{\gamma}}^{-1}}^2 \right] \\ &\quad + 2\kappa_{\boldsymbol{\gamma}, \boldsymbol{\rho}, \mathbf{N}}^{-2} \sum_{i=1}^b d_i N_i \gamma_i^2 \tilde{M}_i^2 \left(1 + \frac{\gamma_i^2 \tilde{M}_i^2}{12} + \frac{\gamma_i \tilde{M}_i^2}{2\tilde{m}_i} \right), \end{aligned}$$

where, for any $i \in [b]$, \tilde{M}_i and \tilde{m}_i are defined in (S25).

Proof. Taking expectation in Lemma S26, we get for any $n \in \mathbb{N}$, $\epsilon > 0$ that

$$\mathbb{E} \left[\left\| \tilde{Z}_{n+1} - Z_{n+1} \right\|_{\mathbf{D}_{\mathbf{N}\boldsymbol{\gamma}}^{-1}}^2 \right] \leq (1 + 2\epsilon) \mathbb{E} \left[\|\mathbf{T}_1^{(n)}\|^2 \|\tilde{Z}_n - Z_n\|_{\mathbf{D}_{\mathbf{N}\boldsymbol{\gamma}}^{-1}}^2 \right] + (1 + 1/\{2\epsilon\}) \mathbb{E} \left[\|\mathbf{T}_2^{(n)}\|^2 \right],$$

where $\mathbf{T}_1^{(n)}$ and $\mathbf{T}_2^{(n)}$ are defined in (S91) and (S92), respectively. To ease notation, denote $\mathbf{B} = \sum_{i=1}^b d_i N_i \gamma_i^2 \tilde{M}_i^2 (1 + \gamma_i^2 \tilde{M}_i^2 / 12 + \gamma_i \tilde{M}_i^2 / (2\tilde{m}_i))$. Using Lemma S30, we obtain for any $n \in \mathbb{N}$, $\epsilon > 0$

$$\mathbb{E} \left[\left\| \tilde{Z}_{n+1} - Z_{n+1} \right\|_{\mathbf{D}_{\mathbf{N}\boldsymbol{\gamma}}^{-1}}^2 \right] \leq (1 + 2\epsilon) \mathbb{E} \left[\|\mathbf{T}_1^{(n)}\|^2 \|\tilde{Z}_n - Z_n\|_{\mathbf{D}_{\mathbf{N}\boldsymbol{\gamma}}^{-1}}^2 \right] + (1 + 1/\{2\epsilon\}) \mathbf{B}. \quad (\text{S105})$$

In addition, Lemma S27 implies that $\|T_1^{(n)}\|^2 \leq (1 - \kappa_{\gamma, \rho, \mathbf{N}})^2$ almost surely. Therefore, taking $\epsilon = (1 - [1 - \kappa_{\gamma, \rho, \mathbf{N}}]^2)/(4[1 - \kappa_{\gamma, \rho, \mathbf{N}}]^2)$, (S105) yields for any $n \geq 0$,

$$\mathbb{E} \left[\left\| \tilde{Z}_{n+1} - Z_{n+1} \right\|_{\mathbf{D}_{N\gamma}^{-1}}^2 \right] \leq \frac{1 + (1 - \kappa_{\gamma, \rho, \mathbf{N}})^2}{2} \mathbb{E} \left[\left\| \tilde{Z}_n - Z_n \right\|_{\mathbf{D}_{N\gamma}^{-1}}^2 \right] + \frac{1 + (1 - \kappa_{\gamma, \rho, \mathbf{N}})^2}{1 - (1 - \kappa_{\gamma, \rho, \mathbf{N}})^2} \mathbf{B}.$$

An easy induction implies for any $n \geq 1$,

$$\mathbb{E} [\left\| \tilde{Z}_n - Z_n \right\|_{\mathbf{D}_{N\gamma}^{-1}}^2] \leq \left(\frac{1 + (1 - \kappa_{\gamma, \rho, \mathbf{N}})^2}{2} \right)^{n-1} \mathbb{E} [\left\| \tilde{Z}_1 - Z_1 \right\|_{\mathbf{D}_{N\gamma}^{-1}}^2] + 2 \frac{1 + (1 - \kappa_{\gamma, \rho, \mathbf{N}})^2}{(1 - (1 - \kappa_{\gamma, \rho, \mathbf{N}})^2)^2} \mathbf{B}. \quad (\text{S106})$$

Since $\kappa_{\gamma, \rho, \mathbf{N}}^2 = (\min_{i \in [b]} \{N_i \gamma_i m_i\} + r_{\gamma, \rho, \mathbf{N}})^2$ and using $\kappa_{\gamma, \rho, \mathbf{N}}^2 \leq 1$, we obtain

$$\begin{aligned} (1 + (1 - \kappa_{\gamma, \rho, \mathbf{N}})^2)/2 &= 1 - \kappa_{\gamma, \rho, \mathbf{N}} + \kappa_{\gamma, \rho, \mathbf{N}}^2/2, \\ (1 + (1 - \kappa_{\gamma, \rho, \mathbf{N}})^2)/(1 - (1 - \kappa_{\gamma, \rho, \mathbf{N}})^2)^2 &\leq \kappa_{\gamma, \rho, \mathbf{N}}^{-2}. \end{aligned}$$

Combining these inequalities with (S106) and (S105) completes the proof. \square

Lemma S32. Assume **H1-H2** and let $\mathbf{N} \in (\mathbb{N}^*)^b, \gamma \in (\mathbb{R}_+^*)^b$ such that, for any $i \in [b]$, $\gamma_i < 1/\tilde{M}_i, N_i \gamma_i \leq 2/(m_i + \tilde{M}_i)$ and $\kappa_{\gamma, \rho, \mathbf{N}} = \min_{i \in [b]} \{N_i \gamma_i m_i\} - r_{\gamma, \rho, \mathbf{N}} \in (0, 1)$, where $r_{\gamma, \rho, \mathbf{N}}$ is defined in (S33). Then, for any $\mathbf{x} \in \mathbb{R}^{d+p}$ and $n \geq 1$, we have

$$\begin{aligned} &W_2^2(\delta_{\mathbf{x}} P_{\rho, \gamma, \mathbf{N}}^n, \Pi_{\rho}) \\ &\leq (1 - \kappa_{\gamma, \rho, \mathbf{N}} + \kappa_{\gamma, \rho, \mathbf{N}}^2/2)^{2(n-1)} (1 + \|\bar{\mathbf{B}}_0^{-1} \mathbf{B}_0^{\top} \tilde{\mathbf{D}}_0^{1/2}\|^2) \max_{i \in [b]} \{N_i \gamma_i\} \mathbb{E} [\left\| \tilde{Z}_1 - Z_1 \right\|_{\mathbf{D}_{N\gamma}^{-1}}^2] \\ &\quad + \frac{2(1 + \|\bar{\mathbf{B}}_0^{-1} \mathbf{B}_0^{\top} \tilde{\mathbf{D}}_0^{1/2}\|^2) \max_{i \in [b]} \{N_i \gamma_i\}}{\kappa_{\gamma, \rho, \mathbf{N}}^2} \sum_{i=1}^b d_i N_i \gamma_i^2 \tilde{M}_i^2 [1 + \gamma_i^2 \tilde{M}_i^2/12 + \gamma_i \tilde{M}_i^2/(2\tilde{m}_i)], \end{aligned}$$

where $\bar{\mathbf{B}}_0, \mathbf{B}_0, \tilde{\mathbf{D}}_0$ are defined in (S2)-(S3), $P_{\rho, \gamma, \mathbf{N}}$ is defined in (S17), $(\tilde{Z}_n, Z_n)_{n \in \mathbb{N}}$ is defined in (S79) and for any $i \in [b]$, \tilde{M}_i, \tilde{m}_i are defined in (S25).

Proof. By Lemma S31, we have the following upper bound for $n \geq 1$,

$$\begin{aligned} \mathbb{E} [\left\| \tilde{Z}_n - Z_n \right\|_{\mathbf{D}_{N\gamma}^{-1}}^2] &\leq (1 - \kappa_{\gamma, \rho, \mathbf{N}} + \kappa_{\gamma, \rho, \mathbf{N}}^2/2)^{2(n-1)} \mathbb{E} [\left\| \tilde{Z}_1 - Z_1 \right\|_{\mathbf{D}_{N\gamma}^{-1}}^2] \\ &\quad + 2\kappa_{\gamma, \rho, \mathbf{N}}^{-2} \sum_{i=1}^b d_i N_i \gamma_i^2 \tilde{M}_i^2 \left(1 + \frac{\gamma_i^2 \tilde{M}_i^2}{12} + \frac{\gamma_i \tilde{M}_i^2}{2\tilde{m}_i} \right). \end{aligned}$$

Using (S79), Lemma S23, combined with the previous inequality, we get for any $n \geq 1, \mathbf{x} \in \mathbb{R}^{d+p}$,

$$\begin{aligned}
& W_2^2(\Pi_{\rho}, \delta_{\mathbf{x}} P_{\rho, \gamma, \mathbf{N}}^n) \\
& \leq (1 + \|\bar{\mathbf{B}}_0^{-1} \mathbf{B}_0^{\top} \tilde{\mathbf{D}}_0^{1/2}\|^2) \mathbb{E}[\|\tilde{Z}_n - Z_n\|^2] \\
& \leq (1 + \|\bar{\mathbf{B}}_0^{-1} \mathbf{B}_0^{\top} \tilde{\mathbf{D}}_0^{1/2}\|^2) \max_{i \in [b]} \{N_i \gamma_i\} \mathbb{E}[\|\tilde{Z}_n - Z_n\|_{\mathbf{D}_{\mathbf{N}\gamma}^{-1}}^2] \\
& \leq (1 - \kappa_{\gamma, \rho, \mathbf{N}} + \kappa_{\gamma, \rho, \mathbf{N}}^2/2)^{2(n-1)} (1 + \|\bar{\mathbf{B}}_0^{-1} \mathbf{B}_0^{\top} \tilde{\mathbf{D}}_0^{1/2}\|^2) \max_{i \in [b]} \{N_i \gamma_i\} \mathbb{E}[\|\tilde{Z}_1 - Z_1\|_{\mathbf{D}_{\mathbf{N}\gamma}^{-1}}^2] \\
& \quad + \frac{2(1 + \|\bar{\mathbf{B}}_0^{-1} \mathbf{B}_0^{\top} \tilde{\mathbf{D}}_0^{1/2}\|^2) \max_{i \in [b]} \{N_i \gamma_i\}}{\kappa_{\gamma, \rho, \mathbf{N}}^2} \sum_{i=1}^b d_i N_i \gamma_i^2 \tilde{M}_i^2 \left(1 + \frac{\gamma_i^2 \tilde{M}_i^2}{12} + \frac{\gamma_i \tilde{M}_i^2}{2\tilde{m}_i}\right).
\end{aligned}$$

Hence the stated result. \square

Proof of Proposition 4/Proposition S29.

Proof. Since for any $i \in [b]$, $\gamma_i \leq m_i/40\tilde{M}_i^2(\min_{i \in [b]} \{m_i/\tilde{M}_i\}/\max_{i \in [b]} \{m_i/\tilde{M}_i\})^2$, setting

$$N_i^*(\gamma_i) = \lfloor m_i \min_{i \in [b]} \{m_i/\tilde{M}_i\}^2 / (20\gamma_i \tilde{M}_i^2 \max_{i \in [b]} \{m_i/\tilde{M}_i\}^2) \rfloor$$

implies $\kappa_{\gamma, \rho, \mathbf{N}^*} \in (0, 1)$ by Lemma S28. Thereby, letting n tend towards infinity in Lemma S32 and using Proposition S13 conclude the proof. \square

S4.3 Proof of Proposition 5

We first give the formal statement of Proposition 5.

Proposition S33. Assume **H1-H2-H3** and let $\gamma \in (\mathbb{R}_+^*)^b$, $\mathbf{N} \in (\mathbb{N}^*)^b$ such that for any $i \in [b]$, $\gamma_i \leq m_i/40\tilde{M}_i^2(\min_{i \in [b]} \{m_i/\tilde{M}_i\}/\max_{i \in [b]} \{m_i/\tilde{M}_i\})^2$ and $N_i = \lfloor m_i \min_{i \in [b]} \{m_i/\tilde{M}_i\}^2 / (20\gamma_i \tilde{M}_i^2 \max_{i \in [b]} \{m_i/\tilde{M}_i\}^2) \rfloor$. Then, we have

$$W_2^2(\Pi_{\rho, \gamma, \mathbf{N}}, \Pi_{\rho}) \leq 4(1 + \|\bar{\mathbf{B}}_0^{-1} \mathbf{B}_0^{\top} \tilde{\mathbf{D}}_0^{1/2}\|^2) \frac{\max_{i \in [b]} \{m_i/\tilde{M}_i^2\}}{\min_{i \in [b]} \{m_i/\tilde{M}_i\}^2} \mathcal{R}^*(\gamma),$$

where setting $\mathfrak{f}_i = m_i/(20\tilde{M}_i)$,

$$\mathcal{R}^*(\gamma) = \sum_{i=1}^b \left\{ d_i \gamma_i^2 \tilde{M}_i^2 + \frac{d_i \gamma_i^2 \mathfrak{f}_i}{\tilde{M}_i} \left(d_i L_i^2 + \frac{\tilde{M}_i^4}{\tilde{m}_i} \right) + d_i \gamma_i \tilde{M}_i \mathfrak{f}_i^3 (1 + \mathfrak{f}_i + \mathfrak{f}_i^2) \right\}, \quad (\text{S107})$$

$\bar{\mathbf{B}}_0, \mathbf{B}_0, \tilde{\mathbf{D}}_0$ are defined in (S2)-(S3), and for any $i \in [b]$, \tilde{m}_i, \tilde{M}_i are defined in (S25).

We provide the proof of Proposition 5 in what follows. Similarly to Lemma S26 for the proof of Proposition 4, we derive an explicit relation between $\|\tilde{Z}_{n+1} - Z_{n+1}\|$ and $\|\tilde{Z}_n - Z_n\|$.

Lemma S34. *Assume H1-H2-H3 and let $\mathbf{N} \in (\mathbb{N}^*)^b, \gamma \in (\mathbb{R}_+^*)^b$ such that for any $i \in [b]$, $N_i \gamma_i \leq 2/(m_i + \tilde{M}_i)$ and $\gamma_i < 1/\tilde{M}_i$. Then, for $n \geq 1$, we have*

$$\mathbb{E}[\|\tilde{Z}_{n+1} - Z_{n+1}\|_{\mathbf{D}_{\mathbf{N}\gamma}^{-1}}^2]^{1/2} \leq (1 - \min_{i \in [b]} \{N_i \gamma_i m_i\} + r_{\gamma, \rho, \mathbf{N}}) \mathbb{E}[\|\tilde{Z}_n - Z_n\|_{\mathbf{D}_{\mathbf{N}\gamma}^{-1}}^2]^{1/2} + \mathcal{R}(\gamma, \mathbf{N})^{1/2},$$

where

$$\begin{aligned} \mathcal{R}(\gamma, \mathbf{N}) = & \sum_{i=1}^b d_i N_i \gamma_i^3 (d_i L_i^2 + \tilde{M}_i^4 / \tilde{m}_i) + \sum_{i=1}^b \left(d_i \gamma_i^2 \tilde{M}_i^2 + d_i N_i^3 \gamma_i^4 \tilde{M}_i^4 \right) \\ & + \sum_{i=1}^b d_i N_i^4 \gamma_i^5 \tilde{M}_i^5 (1 + N_i \gamma_i \tilde{M}_i), \end{aligned} \quad (\text{S108})$$

$(\tilde{Z}_n, Z_n)_{n \in \mathbb{N}}$ is defined in (S79), $r_{\gamma, \rho, \mathbf{N}}$ in (S33) and for any $i \in [b]$, \tilde{m}_i, \tilde{M}_i are defined in (S25).

Proof. Let $n \in \mathbb{N}$. For any $k \in \mathbb{N}$, recall that $\mathbf{M}_k^{(n)}$ is defined in (S89) and invertible by Lemma S24. Define

$$w_n = \mathbf{D}_{\mathbf{N}\gamma}^{-1/2} (\tilde{Z}_n - Z_n).$$

Under this notation, the result given in Lemma S25 can be rewritten as

$$w_{n+1} = \mathbf{T}_1^{(n)} w_n - \mathbf{T}_2^{(n)},$$

where $\mathbf{T}_1^{(n)}$ and $\mathbf{T}_2^{(n)}$ are defined in (S91) and (S92), respectively. By the Minkowsky inequality and using (S81), we have

$$\mathbb{E}^{\mathcal{G}_n} [\|w_{n+1}\|^2]^{1/2} \leq \mathbb{E}^{\mathcal{G}_n} [\|\mathbf{T}_1^{(n)} w_n\|^2]^{1/2} + \mathbb{E}^{\mathcal{G}_n} [\|\mathbf{T}_2^{(n)}\|^2]^{1/2}. \quad (\text{S109})$$

Since by Lemma S27,

$$\|\mathbf{T}_1^{(n)}\| \leq 1 - \min_{i \in [b]} \{N_i \gamma_i m_i\} + r_{\gamma, \rho, \mathbf{N}}, \quad (\text{S110})$$

it remains to bound $\mathbb{E}^{\mathcal{G}_n} [\|\mathbf{T}_2^{(n)}\|^2]$ to complete the proof.

For any $i \in [b]$, recall the function $V_i^{\theta_n} : \mathbb{R}^{d_i} \rightarrow \mathbb{R}$ defined for any $\mathbf{y}^i \in \mathbb{R}^{d_i}$ by $V_i^{\theta_n}(\mathbf{y}^i) = U_i(\mathbf{y}^i) + \|\mathbf{y}^i - \mathbf{A}_i \theta_n\|^2 / (2\rho_i)$. For any $i \in [b], k \in \mathbb{N}$, using the Itô formula, we

have for $l \in [k\gamma_i, (k+1)\gamma_i)$,

$$\begin{aligned} \nabla V_i(\tilde{Y}_{k\gamma_i+l}^{(i,n)}) - \nabla V_i(\tilde{Y}_{k\gamma_i}^{(i,n)}) &= \int_{k\gamma_i}^{k\gamma_i+l} \{ \nabla^2 V_i^{\theta_n}(\tilde{Y}_u^{(i,n)}) \nabla V_i^{\theta_n}(\tilde{Y}_u) + \vec{\Delta}(\nabla V_i^{\theta_n})(\tilde{Y}_u^{(i,n)}) \} du \\ &\quad + \sqrt{2} \int_{k\gamma_i}^{k\gamma_i+l} \nabla^2 V_i^{\theta_n}(\tilde{Y}_u^{(i,n)}) dB_u^i. \end{aligned} \quad (\text{S111})$$

For any $i \in [b]$, $k \in \mathbb{N}$, define

$$\begin{aligned} a_{1,k}^{(i,n)} &= \mathbb{1}_{[N_i]}(k+1) [\mathbf{M}_\infty^{(i,n)}]^{-1} \mathbf{M}_{k+1}^{(i,n)} \int_0^{\gamma_i} \int_{k\gamma_i}^{k\gamma_i+l} \nabla^2 V_i^{\theta_n}(\tilde{Y}_u^{(i,n)}) \nabla V_i^{\theta_n}(\tilde{Y}_u^{(i,n)}) du dl, \\ a_{2,k}^{(i,n)} &= \mathbb{1}_{[N_i]}(k+1) [\mathbf{M}_\infty^{(i,n)}]^{-1} \mathbf{M}_{k+1}^{(i,n)} \int_0^{\gamma_i} \int_{k\gamma_i}^{k\gamma_i+l} \vec{\Delta}(\nabla V_i^{\theta_n})(\tilde{Y}_u^{(i,n)}) du dl, \\ a_{3,k}^{(i,n)} &= \sqrt{2} \mathbb{1}_{[N_i]}(k+1) [\mathbf{M}_\infty^{(i,n)}]^{-1} \mathbf{M}_{k+1}^{(i,n)} \int_0^{\gamma_i} \int_{k\gamma_i}^{k\gamma_i+l} \nabla^2 V_i^{\theta_n}(\tilde{Y}_u^{(i,n)}) dB_u^i dl. \end{aligned}$$

With these notation and by (S111), we have

$$\begin{aligned} \|T_2^{(n)}\|^2 &= \sum_{i \in [b]} \frac{1}{N_i \gamma_i} \left\| \sum_{k \in \mathbb{N}} \{a_{1,k}^{(i,n)} + a_{2,k}^{(i,n)} + a_{3,k}^{(i,n)}\} \right\|^2 \\ &\leq E_1 + E_2 + E_3, \end{aligned} \quad (\text{S112})$$

where for any $j \in [3]$, $E_j = 3 \sum_{i \in [b]} \left\| \sum_{k=0}^{N_i-1} a_{j,k}^{(i,n)} \right\|^2 / (N_i \gamma_i)$. We now bound $\{E_j\}_{j \in [3]}$.

Upper bound on E_1 . For any $i \in [b]$, $k \in \mathbb{N}$, recall that we have $[\mathbf{M}_\infty^{(i,n)}]^{-1} \mathbf{M}_{k+1}^{(i,n)} = \prod_{l=k+1}^\infty (\mathbf{I}_{d_i} + \mathbf{C}_l^{(i,n)})$ where $\mathbf{C}_l^{(i,n)}$ is defined in (S88). In addition, since we suppose for any $i \in [b]$, that $\gamma_i \tilde{M}_i < 1$, Lemma S24 implies

$$\left\| \prod_{l=k+1}^{N_i-1} (\mathbf{I}_{d_i} - \mathbf{C}_l^{(i,n)}) \right\|^2 \leq \{1 - \gamma_i \tilde{m}_i\}^{2(N_i-k-1)}.$$

Combining this result with the Cauchy-Schwarz inequality, we obtain

$$\frac{1}{N_i} \left\| \sum_{k=0}^{N_i-1} a_{1,k}^{(i,n)} \right\|^2 \leq \sum_{k=0}^{N_i-1} \left\| \int_0^{\gamma_i} \int_{k\gamma_i}^{k\gamma_i+l} \nabla^2 V_i^{\theta_n}(\tilde{Y}_u^{(i,n)}) \nabla V_i^{\theta_n}(\tilde{Y}_u^{(i,n)}) du dl \right\|^2. \quad (\text{S113})$$

For $i \in [b]$, using the definition of $\mathbf{z}_{n,\star}^i = \arg \min_{\mathbf{y}^i \in \mathbb{R}^{d_i}} V_i^{\theta_n}(\mathbf{y}^i) \in \mathbb{R}^{d_i}$, we have $\nabla V_i^{\theta_n}(\mathbf{z}_{n,\star}^i) = \mathbf{0}_{d_i}$. Therefore, for $i \in [b]$, $k \in \mathbb{N}$, conditioning with respect to $\mathcal{F}_{k\gamma_i}^{(n)}$ defined in (S82) and

using the \tilde{M}_i -Lipschitz property of $V_i^{\theta_n}$ by **H2** gives

$$\begin{aligned} \mathbb{E}^{\mathcal{F}_{k\gamma_i}^{(n)}} [\|\nabla^2 V_i^{\theta_n}(\tilde{Y}_u^{(i,n)}) \nabla V_i^{\theta_n}(\tilde{Y}_u^{(i,n)})\|^2] &\leq \tilde{M}_i^2 \mathbb{E}^{\mathcal{F}_{k\gamma_i}^{(n)}} [\|\nabla V_i^{\theta_n}(\tilde{Y}_u^{(i,n)}) - \nabla V_i^{\theta_n}(\mathbf{z}_{n,\star}^i)\|^2] \\ &\leq \tilde{M}_i^4 \mathbb{E}^{\mathcal{F}_{k\gamma_i}^{(n)}} [\|\tilde{Y}_u^{(i,n)} - \mathbf{z}_{n,\star}^i\|^2] . \end{aligned}$$

For any $i \in [b]$, $k \in \mathbb{N}$, combining this result with the Jensen inequality yields

$$\begin{aligned} &\mathbb{E}^{\mathcal{F}_{k\gamma_i}^{(n)}} \left[\left\| \int_0^{\gamma_i} \int_{k\gamma_i}^{k\gamma_i+l} \nabla^2 V_i^{\theta_n}(\tilde{Y}_u^{(i,n)}) \nabla V_i^{\theta_n}(\tilde{Y}_u^{(i,n)}) \, du \, dl \right\|^2 \right] \\ &\leq \gamma_i \int_0^{\gamma_i} l \int_{k\gamma_i}^{k\gamma_i+l} \mathbb{E}^{\mathcal{F}_{k\gamma_i}^{(n)}} [\|\nabla^2 V_i^{\theta_n}(\tilde{Y}_u^{(i,n)}) \nabla V_i^{\theta_n}(\tilde{Y}_u^{(i,n)})\|^2] \, du \, dl \\ &\leq \gamma_i \tilde{M}_i^4 \int_0^{\gamma_i} l \int_{k\gamma_i}^{k\gamma_i+l} \mathbb{E}^{\mathcal{F}_{k\gamma_i}^{(n)}} [\|\tilde{Y}_u^{(i,n)} - \mathbf{z}_{n,\star}^i\|^2] \, du \, dl . \end{aligned} \quad (\text{S114})$$

By Lemma S22, we have for any $i \in [b]$, $u \in \mathbb{R}_+$,

$$\mathbb{E}^{\mathcal{G}_n} [\|\tilde{Y}_u^{(i,n)} - \mathbf{z}_{n,\star}^i\|^2] \leq d_i / \tilde{m}_i . \quad (\text{S115})$$

Injecting this result in (S114) yields

$$\mathbb{E} \left[\int_0^{\gamma_i} l \int_{k\gamma_i}^{k\gamma_i+l} \mathbb{E}^{\mathcal{F}_{k\gamma_i}^{(n)}} [\|\tilde{Y}_u^{(i,n)} - \mathbf{z}_{n,\star}^i\|^2] \, du \, dl \right] \leq d_i \gamma_i^3 / (3\tilde{m}_i) .$$

Finally, this inequality, (S114) and (S113), we get

$$\mathbb{E} [E_1] \leq \sum_{i=1}^b d_i N_i \gamma_i^3 \tilde{M}_i^4 / \tilde{m}_i . \quad (\text{S116})$$

Upper bound on E_2 . Using the Cauchy-Schwarz inequality, we have

$$\frac{1}{N_i} \left\| \sum_{k=0}^{N_i-1} a_{2,k}^{(i,n)} \right\|^2 \leq \sum_{k=0}^{N_i-1} \left\| \int_0^{\gamma_i} \int_{k\gamma_i}^{k\gamma_i+l} \vec{\Delta}(\nabla V_i^{\theta_n})(\tilde{Y}_u^{(i,n)}) \, du \, dl \right\|^2 .$$

By **H3**, we have for any $\mathbf{z}_i \in \mathbb{R}^{d_i}$, $\|\vec{\Delta}(\nabla V_i^{\theta_n})(\mathbf{z}_i)\|^2 \leq d_i^2 L_i^2$. Therefore, we obtain

$$\begin{aligned} \left\| \int_0^{\gamma_i} \int_{k\gamma_i}^{k\gamma_i+l} \vec{\Delta}(\nabla V_i^{\theta_n})(\tilde{Y}_u^{(i,n)}) \, du \, dl \right\|^2 &\leq \gamma_i \int_0^{\gamma_i} l \int_{k\gamma_i}^{k\gamma_i+l} \|\vec{\Delta}(\nabla V_i^{\theta_n})(\tilde{Y}_u^{(i,n)})\|^2 \, du \, dl \\ &\leq d_i^2 \gamma_i^4 L_i^2 / 3 . \end{aligned}$$

Thus, we get

$$\mathbb{E} [E_2] \leq \sum_{i=1}^b d_i^2 N_i \gamma_i^3 L_i^2 . \quad (\text{S117})$$

Upper bound on E_3 . For any $i \in [b], k \in \mathbb{N}$, define

$$\Delta_{3,k}^{(i,n)} = \int_0^{\gamma_i} \int_{k\gamma_i}^{k\gamma_i+l} \nabla^2 V_i^{\theta_n}(\tilde{Y}_u^{(i,n)}) dB_u^i dl.$$

Using for any $i \in [b], k \in \mathbb{N}$, $[\mathbf{M}_\infty^{(i,n)}]^{-1} \mathbf{M}_{k+1}^{(i,n)} = \mathbf{I}_{d_i} - \sum_{l=k+1}^\infty \mathbf{C}_l^{(i,n)} + \mathbf{R}_k^{(i,n)}$ where $\mathbf{R}_k^{(i,n)}$ is defined in (S34), we have, for any $i \in [b], k \in \mathbb{N}$,

$$\begin{aligned} \left\| \sum_{k=0}^{N_i-1} a_{3,k}^{(i,n)} \right\|^2 &= \left\| \sqrt{2} \sum_{k=0}^{N_i-1} \prod_{l=k+1}^{N_i} [\mathbf{I}_{d_i} - \mathbf{C}_l^{(i,n)}] \Delta_{3,k}^{(i,n)} \right\|^2 \\ &= 2 \sum_{k_1, k_2=0}^{N_i-1} \langle \mathbf{R}_{k_1}^{(i,n)} \Delta_{3,k_1}^{(i,n)}, \mathbf{R}_{k_2}^{(i,n)} \Delta_{3,k_2}^{(i,n)} \rangle + 2 \sum_{k_1, k_2=0}^{N_i-1} \langle \Delta_{3,k_1}^{(i,n)}, \Delta_{3,k_2}^{(i,n)} \rangle \\ &\quad + 2 \sum_{k_1, k_2=0}^{N_i-1} \left\langle \sum_{l=k_1+1}^{N_i} \mathbf{C}_l^{(i,n)} \Delta_{3,k_1}^{(i,n)}, \sum_{l=k_2+1}^{N_i} \mathbf{C}_l^{(i,n)} \Delta_{3,k_2}^{(i,n)} \right\rangle \\ &\quad - 4 \sum_{k_1, k_2=0}^{N_i-1} \left\langle \sum_{l=k_1+1}^{N_i} \mathbf{C}_l^{(i,n)} \Delta_{3,k_1}^{(i,n)}, \Delta_{3,k_2}^{(i,n)} \right\rangle + 4 \sum_{k_1, k_2=0}^{N_i-1} \langle \mathbf{R}_{k_1}^{(i,n)} \Delta_{3,k_1}^{(i,n)}, \Delta_{3,k_2}^{(i,n)} \rangle \\ &\quad - 4 \sum_{k_1, k_2=0}^{N_i-1} \langle \mathbf{R}_{k_1}^{(i,n)} \Delta_{3,k_1}^{(i,n)}, \sum_{l=k_2+1}^{N_i} \mathbf{C}_l^{(i,n)} \Delta_{3,k_2}^{(i,n)} \rangle. \end{aligned} \quad (\text{S118})$$

We now control the quantities which appear in (S118). First, by **H2**, for any $i \in [b], \mathbf{x}^i, \mathbf{y}^i \in \mathbb{R}^{d_i}$, note that we have

$$\|\nabla^2 V_i^{\theta_n}(\mathbf{x}^i) \mathbf{y}^i\| \leq \tilde{M}_i \|\mathbf{y}^i\|.$$

By the Jensen inequality and the Itô isometry, for any $k \in \mathbb{N}$, we get

$$\begin{aligned} \mathbb{E}^{\mathcal{F}_{k\gamma_i}^{(n)}} [\|\Delta_{3,k}^{(i,n)}\|^2] &= \mathbb{E}^{\mathcal{F}_{k\gamma_i}^{(n)}} \left[\left\| \int_0^{\gamma_i} \int_{k\gamma_i}^{k\gamma_i+l} \nabla^2 V_i^{\theta_n}(\tilde{Y}_u^{(i,n)}) dB_u^i dl \right\|^2 \right] \\ &\leq \gamma_i \tilde{M}_i^2 \int_0^{\gamma_i} \mathbb{E}^{\mathcal{F}_{k\gamma_i}^{(n)}} \left[\left\| \int_{k\gamma_i}^{k\gamma_i+l} dB_u^i \right\|^2 \right] dl = d_i \gamma_i^3 \tilde{M}_i^2 / 2. \end{aligned} \quad (\text{S119})$$

In addition, since for $i \in [b]$, $(\int_0^t \nabla^2 V_i^{\theta_n}(\tilde{Y}_u^{(i,n)}) dB_u^i)_{t \geq 0}$ is a $(\mathcal{F}_t^{(n)})_{t \geq 0}$ -martingale, for $(k_1, k_2) \in \{0, \dots, N_i - 1\}^2$ such that $k_1 < k_2$, we obtain

$$\mathbb{E}^{\mathcal{G}_n} [\Delta_{3,k_1}^{(i,n) \top} \Delta_{3,k_2}^{(i,n)}] = \mathbb{E}^{\mathcal{G}_n} [\mathbb{E}^{\mathcal{F}_{k_2\gamma_i}^{(n)}} [\Delta_{3,k_1}^{(i,n) \top} \Delta_{3,k_2}^{(i,n)}]] = 0.$$

Therefore,

$$\sum_{k_1, k_2=0}^{N_i-1} \mathbb{E}^{\mathcal{G}_n} [\langle \Delta_{3,k_1}^{(i,n)}, \Delta_{3,k_2}^{(i,n)} \rangle] = d_i N_i \gamma_i^3 \tilde{M}_i^2 / 2.$$

Second, since for any $i \in [b], l \in \mathbb{N}$, $\mathbf{C}_l^{(i,n)} \in \mathbb{R}^{d_i \times d_i}$ is symmetric positive semi-definite, we have

$$\sum_{k_1, k_2=0}^{N_i-1} \left\langle \sum_{l=k_1+1}^{N_i} \mathbf{C}_l^{(i,n)} \Delta_{3,k_1}^{(i,n)}, \Delta_{3,k_2}^{(i,n)} \right\rangle = \left\langle \left\{ \sum_{l=1}^N C_l \right\} \sum_{k_1=0}^{l-1} \Delta_{3,k_1}, \sum_{k_1=0}^{l-1} \Delta_{3,k_2} \right\rangle \geq 0.$$

Third, using for any $i \in [b], l \in \mathbb{N}$, using $\|\mathbf{C}_l^{(i,n)}\| \leq \gamma_i \tilde{M}_i$ by definition (S88) and **H 2** and combining the Cauchy-Schwarz inequality with (S119), for any $i \in [b], (k_1, k_2) \in \{0, \dots, N_i - 1\}^2$, we get

$$\sum_{k_1, k_2=0}^{N_i-1} \mathbb{E}^{\mathcal{G}_n} \left[\left\langle \sum_{l=k_1+1}^{N_i} \mathbf{C}_l^{(i,n)} \Delta_{3,k_1}^{(i,n)}, \sum_{l=k_2+1}^{N_i} \mathbf{C}_l^{(i,n)} \Delta_{3,k_2}^{(i,n)} \right\rangle \right] \leq d_i N_i^4 \gamma_i^5 \tilde{M}_i^4 / 8.$$

Using (S119) again and Lemma S8, for $i \in [b]$, we obtain

$$\begin{aligned} & \sum_{k_1, k_2=0}^{N_i-1} \mathbb{E}^{\mathcal{G}_n} \left[\langle \mathbf{R}_{k_1}^{(i,n)} \Delta_{3,k_1}^{(i,n)}, \mathbf{R}_{k_2}^{(i,n)} \Delta_{3,k_2}^{(i,n)} \rangle \right] \leq (d_i \gamma_i^3 \tilde{M}_i^2 / 2) \sum_{k_1, k_2=0}^{N_i-1} \mathbb{E} \left[\|\mathbf{R}_{k_1}^{(i,n)}\| \|\mathbf{R}_{k_2}^{(i,n)}\| \right] \\ & \leq (d_i \gamma_i^3 \tilde{M}_i^2 / 2) \left\{ \sum_{k=0}^{N_i-1} (\exp[(N_i - k) \gamma_i \tilde{M}_i] - 1 - [(N_i - k) \gamma_i \tilde{M}_i]) \right\}^2 \\ & \leq (d_i \gamma_i^3 \tilde{M}_i^2 / 2) \left\{ (\tilde{M}_i \gamma_i)^{-1} \int_0^{N_i \gamma_i \tilde{M}_i} \{e^t - 1 - t\} dt \right\}^2 \\ & \leq \frac{(e^{N_i \gamma_i \tilde{M}_i} + 1)^2}{288} d_i N_i^6 \gamma_i^7 \tilde{M}_i^6. \end{aligned}$$

Similarly, we get Moreover, using the Cauchy-Schwarz inequality, for any $i \in [b]$ we get

$$\begin{aligned} & \sum_{k_1, k_2=0}^{N_i-1} \mathbb{E} [\langle \Delta_{k_1}^{(i,n)}, \mathbf{R}_{k_2}^{(i,n)} \Delta_{k_2}^{(i,n)} \rangle] \leq \sum_{k_1, k_2=0}^{N_i-1} \mathbb{E} \left[\|\Delta_{k_1}^{(i,n)}\| \|\Delta_{k_2}^{(i,n)}\| \|\mathbf{R}_{k_2}^{(i,n)}\| \right] \\ & \leq \frac{d_i N_i \gamma_i^3 \tilde{M}_i^2}{24} (e^{N_i \gamma_i \tilde{M}_i} + 1) N_i^3 \gamma_i^2 \tilde{M}_i^2 \\ & \leq d_i N_i^4 \gamma_i^5 \tilde{M}_i^4 \frac{e^{N_i \gamma_i \tilde{M}_i} + 1}{24}. \end{aligned}$$

In addition, for any $i \in [b]$, we have also

$$\sum_{k_1, k_2=0}^{N_i-1} \mathbb{E} \left[\left\langle \mathbf{R}_{k_1}^{(i,n)} \Delta_{3,k_1}^{(i,n)}, \sum_{l=k_2+1}^{N_i} \mathbf{C}_l^{(i,n)} \Delta_{3,k_2}^{(i,n)} \right\rangle \right] \leq d_i N_i^5 \gamma_i^6 \tilde{M}_i^5 \frac{e^{N_i \gamma_i \tilde{M}_i} + 1}{24}. \quad (\text{S120})$$

For any $i \in [b], k \in \mathbb{N}$, regrouping the previous results and using that $N_i \gamma_i \tilde{M}_i \leq 2$ give

$$\mathbb{E}[E_3] \leq \sum_{i=1}^b \{d_i N_i \gamma_i^2 \tilde{M}_i^2 + d_i N_i^3 \gamma_i^4 \tilde{M}_i^4\} + \sum_{i=1}^b d_i N_i^4 \gamma_i^5 \tilde{M}_i^5 (1 + N_i \gamma_i \tilde{M}_i). \quad (\text{S121})$$

Combination of our previous results. Injecting the three upper bounds (S116), (S117), (S121) in (S112), we get

$$\begin{aligned} \mathbb{E} \left[\|T_2^{(n)}\|^2 \right] &\leq \sum_{i=1}^b d_i N_i \gamma_i^3 (d_i L_i^2 + \tilde{M}_i^4 / \tilde{m}_i) + \sum_{i=1}^b \{d_i \gamma_i^2 \tilde{M}_i^2 + d_i N_i^3 \gamma_i^4 \tilde{M}_i^4\} \\ &\quad + \sum_{i=1}^b d_i N_i^4 \gamma_i^5 \tilde{M}_i^5 (1 + N_i \gamma_i \tilde{M}_i). \end{aligned} \quad (\text{S122})$$

Using the recursion defined in (S109), and combining the upper bounds derived in (S110) and (S122) completes the proof. \square

Lemma S35. Assume **H1-H2-H3** and let $\mathbf{N} \in (\mathbb{N}^*)^b, \boldsymbol{\gamma} \in (\mathbb{R}_+^*)^b$ such that for any $i \in [b]$, $N_i \gamma_i \leq 2/(m_i + \tilde{M}_i)$, $\gamma_i < 1/\tilde{M}_i$ and $\kappa_{\boldsymbol{\gamma}, \boldsymbol{\rho}, \mathbf{N}} = \min_{i \in [b]} \{N_i \gamma_i m_i\} - r_{\boldsymbol{\gamma}, \boldsymbol{\rho}, \mathbf{N}} \in (0, 1)$, where $r_{\boldsymbol{\gamma}, \boldsymbol{\rho}, \mathbf{N}}$ is defined in (S33). Then, for $n \geq 1$, we have

$$\mathbb{E} \left[\|\tilde{Z}_{n+1} - Z_{n+1}\|_{\mathbf{D}_{N\boldsymbol{\gamma}}^{-1}}^2 \right]^{1/2} \leq (1 - \kappa_{\boldsymbol{\gamma}, \boldsymbol{\rho}, \mathbf{N}})^{n-1} \mathbb{E} \left[\|\tilde{Z}_1 - Z_1\|_{\mathbf{D}_{N\boldsymbol{\gamma}}^{-1}}^2 \right]^{1/2} + \{\kappa_{\boldsymbol{\gamma}, \boldsymbol{\rho}, \mathbf{N}}\}^{-1} \mathcal{R}(\boldsymbol{\gamma}, \mathbf{N}),$$

where $\mathcal{R}(\boldsymbol{\gamma}, \mathbf{N})$ is given in (S108).

Proof. The proof follows from Lemma S34 combined with a straightforward induction. \square

Proof of Proposition 5/Proposition S33.

Proof of Proposition 5/Proposition S33. For any $i \in [b]$, consider

$$N_i^*(\gamma_i) = \lfloor m_i \min_{i \in [b]} \{m_i / \tilde{M}_i\}^2 / (20 \gamma_i \tilde{M}_i^2 \max_{i \in [b]} \{m_i / \tilde{M}_i\}^2) \rfloor.$$

By Proposition S13 and Lemma S28, $P_{\boldsymbol{\rho}, \boldsymbol{\gamma}, \mathbf{N}}$ converges in W_2 to $\Pi_{\boldsymbol{\rho}, \boldsymbol{\gamma}}$. Therefore, using (S83), Lemma S23 and Lemma S35 and taking $n \rightarrow +\infty$, we obtain

$$W_2^2(\Pi_{\boldsymbol{\rho}, \boldsymbol{\gamma}, \mathbf{N}}, \Pi_{\boldsymbol{\rho}}) \leq 4(1 + \|\bar{\mathbf{B}}_0^{-1} \mathbf{B}_0^\top \tilde{\mathbf{D}}_0^{1/2}\|^2) \frac{\max_{i \in [b]} \{N_i^*(\gamma_i) \gamma_i\}}{\min_{i \in [b]} \{N_i^*(\gamma_i) \gamma_i m_i\}} \mathcal{R}(\boldsymbol{\gamma}, \mathbf{N}^*(\boldsymbol{\gamma})). \quad (\text{S123})$$

By definition of $N_i^*(\gamma_i)$, we have $\gamma_i \tilde{M}_i N_i^*(\gamma_i) \leq \mathfrak{f}_i = m_i / (20 \tilde{M}_i)$ which completes the proof upon using it in (S123). \square

S5 Explicit mixing times

This section aims at providing mixing times for DG-LMC with explicit dependencies w.r.t. the dimension d and the prescribed precision ε . We specify our result to the case where for any $i \in [b]$, $m_i = m$, $M_i = M$, $L_i = L$, $\rho_i = \rho$, $\gamma_i = \gamma$, $N_i = N$ and for the specific initial distribution

$$\mu_{\rho}^{\star} = \delta_{\mathbf{z}^{\star}} \otimes \Pi_{\rho}(\cdot | \mathbf{z}^{\star}), \quad (\text{S124})$$

where

$$\mathbf{x}^{\star} = ([\boldsymbol{\theta}^{\star}]^{\top}, [\mathbf{z}^{\star}]^{\top})^{\top}, \text{ where } \boldsymbol{\theta}^{\star} = \arg \min \{-\log \pi\} \text{ and } \mathbf{z}^{\star} = ([\mathbf{A}_1 \boldsymbol{\theta}^{\star}]^{\top}, \dots, [\mathbf{A}_b \boldsymbol{\theta}^{\star}]^{\top})^{\top}. \quad (\text{S125})$$

Note that sampling from μ_{ρ}^{\star} is straightforward and simply consists in setting $\mathbf{z}_0 = \mathbf{z}^{\star}$ and $\boldsymbol{\theta}_0 = \bar{\mathbf{B}}_0^{-1} \mathbf{B}_0^{\top} \tilde{\mathbf{D}}_0^{1/2} \mathbf{z}_0 + \bar{\mathbf{B}}_0^{-1/2} \xi$, where ξ is a d -dimensional standard Gaussian random variable. Starting from this initialisation, we consider the marginal law of θ_n for $n \geq 1$ and denote it $\Gamma_{\mathbf{x}^{\star}}^n$. By Proposition S13, since for any $i \in [b]$, $N_i = N$, the stationary distribution associated to $P_{\rho, \gamma, \mathbf{N}}$ is $\Pi_{\rho, \gamma} = \Pi_{\rho, \gamma, \mathbf{1}_b}$. We build upon the natural decomposition of the bias:

$$W_2(\Gamma_{\mathbf{x}^{\star}}^n, \pi) \leq W_2(\mu_{\rho}^{\star} P_{\rho, \gamma, \mathbf{N}}^n, \Pi_{\rho, \gamma}) + W_2(\Pi_{\rho, \gamma}, \Pi_{\rho}) + W_2(\pi_{\rho}, \pi),$$

where $\Pi_{\rho, \gamma}$, Π_{ρ} and π_{ρ} are defined in Proposition 2, (2) and (3), respectively. The following subsections focus on deriving conditions on n_{ε} , γ_{ε} , N_{ε} and ρ_{ε} to satisfy $W_2(\Gamma_{\mathbf{x}^{\star}}^{n_{\varepsilon}}, \pi) \leq \varepsilon$, where $\varepsilon > 0$.

S5.1 Lower bound on the number of iterations n_{ε}

In this section, we derive a lower bound on n_{ε} such that $W_2(\mu_{\rho}^{\star} P_{\rho, \gamma, \mathbf{N}}^{n_{\varepsilon}}, \Pi_{\rho, \gamma}) \leq \varepsilon/3$ following the result provided in Proposition S14. Recall that we define the \mathbf{z} -marginal under $\Pi_{\rho, \gamma}$ by

$$\pi_{\rho, \gamma}^{\mathbf{z}} = \int_{\mathbb{R}^d} \Pi_{\rho, \gamma}(\boldsymbol{\theta}, \mathbf{z}) d\boldsymbol{\theta}, \quad (\text{S126})$$

and the transition kernel of the Markov chain $\{Z_n\}_{n \geq 0}$, for all $\mathbf{z} \in \mathbb{R}^p$ and $\mathbf{B} \in \mathcal{B}(\mathbb{R}^p)$, by

$$P_{\rho, \gamma, \mathbf{N}}^{\mathbf{z}}(\mathbf{z}, \mathbf{B}) = \int_{\mathbb{R}^d} Q_{\rho, \gamma, \mathbf{N}}(\mathbf{z}, \mathbf{B} | \boldsymbol{\theta}) \Pi_{\rho}(\boldsymbol{\theta} | \mathbf{z}) d\boldsymbol{\theta}, \quad (\text{S127})$$

where $\Pi_{\rho}(\cdot | \mathbf{z})$ and $Q_{\rho, \gamma, \mathbf{N}}$ are defined in (5) and (S16), respectively. In the case $\mathbf{N} = \mathbf{1}_b$, we simply denote $P_{\rho, \gamma, \mathbf{N}}^{\mathbf{z}}$ by $P_{\rho, \gamma}^{\mathbf{z}}$. We need to bound in Proposition S14 the factor

$$\left\{ \int_{\mathbb{R}^d} \|\mathbf{z}_1 - \mathbf{z}^{\star}\|_{\mathbf{D}_{\mathbf{N}\gamma}^{-1}}^2 \pi_{\rho, \gamma}^{\mathbf{z}}(d\mathbf{z}_1) + \int_{\mathbb{R}^d} \|\mathbf{z}_1 - \mathbf{z}^{\star}\|_{\mathbf{D}_{\mathbf{N}\gamma}^{-1}}^2 P_{\rho, \gamma, \mathbf{N}}^{\mathbf{z}}(\mathbf{z}^{\star}, d\mathbf{z}_1) \right\}^{1/2}. \quad (\text{S128})$$

Our next results provide such bounds.

Lemma S36. Assume **H1**. Then, the transition kernel $P_{\rho, \gamma}^{\mathbf{z}}$ leaves $\pi_{\rho, \gamma}^{\mathbf{z}}$ invariant, that is $\pi_{\rho, \gamma}^{\mathbf{z}} P_{\rho, \gamma}^{\mathbf{z}} = \pi_{\rho, \gamma}^{\mathbf{z}}$, where $\pi_{\rho, \gamma}^{\mathbf{z}}$ is defined by (S50).

Proof. We have for any $\mathbf{B} \in \mathcal{B}(\mathbb{R}^p)$

$$\int_{\mathbf{B}} \pi_{\rho, \gamma}^{\mathbf{z}}(d\mathbf{z}) = \int_{\mathbf{B}} \int_{\mathbb{R}^d} \Pi_{\rho, \gamma}(d\boldsymbol{\theta}, d\mathbf{z}) = \int_{\mathbf{B}} \pi_{\rho, \gamma}^{\mathbf{z}}(d\mathbf{z}) \int_{\mathbb{R}^d} \Pi_{\rho, \gamma}(d\boldsymbol{\theta}|\mathbf{z}) .$$

Therefore, using the fact that $P_{\rho, \gamma}$ leaves $\Pi_{\rho, \gamma}$ invariant from Proposition S5 and Fubini's theorem, we get

$$\begin{aligned} \int_{\mathbf{B}} \pi_{\rho, \gamma}^{\mathbf{z}}(d\mathbf{z}) &= \int_{\mathbf{B}} \int_{\mathbb{R}^d} \Pi_{\rho, \gamma}(d\boldsymbol{\theta}, d\mathbf{z}) = \int_{\mathbf{B}} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d \times \mathbb{R}^p} \Pi_{\rho, \gamma}(d\tilde{\boldsymbol{\theta}}, d\tilde{\mathbf{z}}) P_{\rho, \gamma}((\tilde{\boldsymbol{\theta}}, \tilde{\mathbf{z}}), (d\boldsymbol{\theta}, d\mathbf{z})) \\ &= \int_{\mathbf{B}} \int_{\mathbb{R}^d} \int_{\mathbb{R}^d \times \mathbb{R}^p} \Pi_{\rho, \gamma}(d\tilde{\boldsymbol{\theta}}, d\tilde{\mathbf{z}}) Q_{\rho, \gamma}(\tilde{\mathbf{z}}, d\mathbf{z}|\tilde{\boldsymbol{\theta}}) \Pi_{\rho}(\boldsymbol{\theta}|\mathbf{z}) d\boldsymbol{\theta} \\ &= \int_{\mathbf{B}} \int_{\mathbb{R}^d \times \mathbb{R}^p} \Pi_{\rho, \gamma}(d\tilde{\boldsymbol{\theta}}, d\tilde{\mathbf{z}}) Q_{\rho, \gamma}(\tilde{\mathbf{z}}, d\mathbf{z}|\tilde{\boldsymbol{\theta}}) \int_{\mathbb{R}^d} \Pi_{\rho}(\boldsymbol{\theta}|\mathbf{z}) d\boldsymbol{\theta} \\ &= \int_{\mathbb{R}^d} \pi_{\rho, \gamma}^{\mathbf{z}}(d\tilde{\mathbf{z}}) P_{\rho, \gamma}^{\mathbf{z}}(\tilde{\mathbf{z}}, \mathbf{B}) . \end{aligned} \tag{S129}$$

□

For any $i \in [b]$, let $\boldsymbol{\theta}_i^*$ a minimiser of $\boldsymbol{\theta} \mapsto U_i(\mathbf{A}_i \boldsymbol{\theta})$, and define

$$\mathbf{u}^* = ([\mathbf{A}_1(\boldsymbol{\theta}^* - \boldsymbol{\theta}_1^*)]^\top, \dots, [\mathbf{A}_b(\boldsymbol{\theta}^* - \boldsymbol{\theta}_b^*)]^\top)^\top \tag{S130}$$

Lemma S37. Assume **H1-H2** and let $\mathbf{N} \in (\mathbb{N}^*)^b$, $\gamma, \rho \in (\mathbb{R}_+^*)^b$ such that, for any $i \in [b]$, $\gamma_i \leq 2/(m_i + M_i + 1/\rho_i)$ and denote $\mathbf{z}^* = ([\mathbf{A}_1 \boldsymbol{\theta}^*]^\top, \dots, [\mathbf{A}_b \boldsymbol{\theta}^*]^\top)^\top$. Then, for any $\mathbf{z} \in \mathbb{R}^p$ and $\varepsilon > 0$,

$$\begin{aligned} \int_{\mathbb{R}^p} \|\tilde{\mathbf{z}} - \mathbf{z}^*\|_{\mathbf{D}_{\mathbf{N}\gamma}^{-1}}^2 P_{\rho, \gamma}^{\mathbf{z}}(\mathbf{z}, d\tilde{\mathbf{z}}) &\leq \min_{i \in [b]} \{N_i\}^{-1} \left[\kappa_\gamma^2 (1 + 2\varepsilon) \|\mathbf{z} - \mathbf{z}^*\|_{\mathbf{D}_\gamma^{-1}}^2 \right. \\ &\quad \left. + (1 + 1/(2\varepsilon)) \max_{i \in [b]} \{\gamma_i M_i^2\} \|\mathbf{u}^*\|^2 + \text{Tr}(\mathbf{D}_{\gamma/\rho} \mathbf{P}_0) + 2 \sum_{i=1}^b d_i \right] , \end{aligned}$$

where the transition kernel $P_{\rho, \gamma}^{\mathbf{z}}$ is defined in (S51) with $\mathbf{N} = \mathbf{1}_b$.

Proof. Let $\gamma_i \leq 2/(m_i + M_i + 1/\rho_i)$ for any $i \in [b]$. Let ξ be a d -dimensional Gaussian random variable independent of $\{\eta^i : i \in [b]\}$ where for any $i \in [b]$, η^i is a d_i -dimensional Gaussian random variable. Let $\mathbf{z} \in \mathbb{R}^p$ and Z be the random variable distributed according to $\delta_{\mathbf{z}} P_{\rho, \gamma}^{\mathbf{z}}$, and defined by

$$\boldsymbol{\theta} = \bar{\mathbf{B}}_0^{-1} \mathbf{B}_0^\top \tilde{\mathbf{D}}_0^{1/2} \mathbf{z} + \bar{\mathbf{B}}_0^{-1/2} \xi ,$$

and for any $i \in [b]$,

$$\begin{aligned}
Z^i &= (1 - \gamma_i/\rho_i) \mathbf{z}_i - \gamma_i \nabla U_i(\mathbf{z}_i) + \frac{\gamma_i}{\rho_i} \mathbf{A}_i \theta + \sqrt{2\gamma_i} \eta^i \\
&= (1 - \gamma_i/\rho_i) \mathbf{z}_i - \gamma_i \nabla U_i(\mathbf{z}_i) + \frac{\gamma_i}{\rho_i} \mathbf{A}_i \bar{\mathbf{B}}_0^{-1} \mathbf{B}_0^\top \tilde{\mathbf{D}}_0^{1/2} \mathbf{z} + \frac{\gamma_i}{\rho_i} \mathbf{A}_i \bar{\mathbf{B}}_0^{-1/2} \xi + \sqrt{2\gamma_i} \eta^i \\
&= (1 - \gamma_i/\rho_i) \mathbf{z}_i - \gamma_i [\nabla U_i(\mathbf{z}_i) - \nabla U_i(\mathbf{A}_i \theta^*)] - \gamma_i [\nabla U_i(\mathbf{A}_i \theta^*) - \nabla U_i(\mathbf{A}_i \theta_i^*)] \\
&\quad + \frac{\gamma_i}{\rho_i} \mathbf{A}_i \bar{\mathbf{B}}_0^{-1} \mathbf{B}_0^\top \tilde{\mathbf{D}}_0^{1/2} \mathbf{z} + \frac{\gamma_i}{\rho_i} \mathbf{A}_i \bar{\mathbf{B}}_0^{-1/2} \xi + \sqrt{2\gamma_i} \eta^i .
\end{aligned}$$

Let

$$\begin{aligned}
\mathbf{D}_U^* &= \text{diag} \left(\gamma_1 \int_0^1 \nabla^2 U_1(\mathbf{z}_1 + t(\mathbf{A}_1 \theta^* - \mathbf{z}_1)) dt, \dots, \gamma_b \int_0^1 \nabla^2 U_b(\mathbf{z}_b + t(\mathbf{A}_b \theta^* - \mathbf{z}_b)) dt \right), \\
\tilde{\mathbf{D}}_U^* &= \text{diag} \left(\gamma_1 \int_0^1 \nabla^2 U_1(\mathbf{A}_1 \theta^* + t(\mathbf{A}_1 \theta_1^* - \mathbf{A}_1 \theta^*)) dt, \dots, \gamma_b \int_0^1 \nabla^2 U_b(\mathbf{A}_b \theta^* + t(\mathbf{A}_b \theta_b^* - \mathbf{A}_b \theta^*)) dt \right).
\end{aligned} \tag{S131}$$

Since $\mathbf{P}_0 \mathbf{D}_\rho^{-1/2} \mathbf{z}^* = \mathbf{D}_\rho^{-1/2} \mathbf{z}^*$, it follows that

$$\mathbf{Z} - \mathbf{z}^* = \left[\mathbf{I}_p - \mathbf{D}_U^* - \mathbf{D}_\gamma^{1/2} \mathbf{D}_{\gamma/\rho}^{1/2} (\mathbf{I}_p - \mathbf{P}_0) \mathbf{D}_\rho^{-1/2} \right] (\mathbf{z} - \mathbf{z}^*) - \tilde{\mathbf{D}}_U^* \mathbf{u}^* + \mathbf{D}_\gamma^{1/2} \mathbf{D}_{\gamma/\rho}^{1/2} \mathbf{B}_0 \bar{\mathbf{B}}_0^{-1/2} \xi + \mathbf{D}_{2\gamma}^{1/2} \eta .$$

With the notation $\mathbf{H} = \mathbf{I}_p - \mathbf{D}_U^* - \mathbf{D}_\gamma^{1/2} \mathbf{D}_{\gamma/\rho}^{1/2} (\mathbf{I}_p - \mathbf{P}_0) \mathbf{D}_\rho^{-1/2}$, (S14), and using the fact that for any $\varepsilon > 0$, $\mathbf{a}, \mathbf{b} \in \mathbb{R}^d$, $|\langle \mathbf{a}, \mathbf{b} \rangle| \leq \varepsilon \|\mathbf{a}\|^2 + (4\varepsilon)^{-1} \|\mathbf{b}\|^2$, it follows, for any $\mathbf{z} \in \mathbb{R}^p$, that

$$\begin{aligned}
&\int_{\mathbb{R}^p} \|\tilde{\mathbf{z}} - \mathbf{z}^*\|_{\mathbf{D}_\gamma^{-1}}^2 P_{\rho, \gamma}^{\mathbf{z}}(\mathbf{z}, d\tilde{\mathbf{z}}) \\
&= \int_{\mathbb{R}^p} \int_{\mathbb{R}^d} \|\mathbf{H}(\mathbf{z} - \mathbf{z}^*) - \tilde{\mathbf{D}}_U^* \mathbf{u}^* + \mathbf{D}_\gamma^{1/2} \mathbf{D}_{\gamma/\rho}^{1/2} \mathbf{B}_0 \bar{\mathbf{B}}_0^{-1/2} \xi + \mathbf{D}_{2\gamma}^{1/2} \eta\|_{\mathbf{D}_\gamma^{-1}}^2 \phi_d(\xi) d\xi \phi_p(\eta) d\eta \\
&= \|\mathbf{H}(\mathbf{z} - \mathbf{z}^*) - \tilde{\mathbf{D}}_U^* \mathbf{u}^*\|_{\mathbf{D}_\gamma^{-1}}^2 + \text{Tr}(\mathbf{D}_{\gamma/\rho} \mathbf{P}_0) + 2 \sum_{i=1}^b d_i \\
&\leq \kappa_\gamma^2 \|\mathbf{z} - \mathbf{z}^*\|_{\mathbf{D}_\gamma^{-1}}^2 - 2 \langle \mathbf{H}(\mathbf{z} - \mathbf{z}^*), \tilde{\mathbf{D}}_U^* \mathbf{u}^* \rangle_{\mathbf{D}_\gamma^{-1}} + \|\tilde{\mathbf{D}}_U^* \mathbf{u}^*\|_{\mathbf{D}_\gamma^{-1}}^2 + \text{Tr}(\mathbf{D}_{\gamma/\rho} \mathbf{P}_0) + 2 \sum_{i=1}^b d_i \\
&\leq \kappa_\gamma^2 (1 + 2\varepsilon) \|\mathbf{z} - \mathbf{z}^*\|_{\mathbf{D}_\gamma^{-1}}^2 + \left(1 + \frac{1}{2\varepsilon}\right) \max_{i \in [b]} \{\gamma_i M_i^2\} \|\mathbf{u}^*\|^2 + \text{Tr}(\mathbf{D}_{\gamma/\rho} \mathbf{P}_0) + 2 \sum_{i=1}^b d_i .
\end{aligned}$$

□

Proposition S38. Assume **H1-H2** and let $\mathbf{N} \in (\mathbb{N}^*)^b, \gamma, \rho \in (\mathbb{R}_+^*)^b$ such that, for any $i \in [b]$, $\gamma_i \leq 2/(m_i + M_i + 1/\rho_i)$. Then, we have

$$\begin{aligned} & \int_{\mathbb{R}^d} \|\mathbf{z}_1 - \mathbf{z}^*\|_{\mathbf{D}_{N\gamma}^{-1}}^2 \pi_{\rho, \gamma}^{\mathbf{z}}(d\mathbf{z}_1) \\ & \leq \min_{i \in [b]} \{N_i\}^{-1} \frac{2}{1 - \kappa_\gamma^2} \left(\frac{1 + \kappa_\gamma^2}{1 - \kappa_\gamma^2} \max_{i \in [b]} \{\gamma_i M_i^2\} \|\mathbf{u}^*\|^2 + \text{Tr}(\mathbf{D}_{\gamma/\rho} \mathbf{P}_0) + 2 \sum_{i=1}^b d_i \right), \end{aligned}$$

with κ_γ defined in (S12).

Proof. With the choice $\varepsilon = (1 - \kappa_\gamma^2)/(4\kappa_\gamma^2)$ in Lemma S37 and using Lemma S36, we have

$$\begin{aligned} & \int_{\mathbb{R}^p} \|\tilde{\mathbf{z}} - \mathbf{z}^*\|_{\mathbf{D}_\gamma^{-1}}^2 \pi_{\rho, \gamma}^{\mathbf{z}}(d\tilde{\mathbf{z}}) \leq \frac{\kappa_\gamma^2 + 1}{2} \int_{\mathbb{R}^p} \|\mathbf{z} - \mathbf{z}^*\|_{\mathbf{D}_\gamma^{-1}}^2 \pi_{\rho, \gamma}^{\mathbf{z}}(d\mathbf{z}) + \frac{1 + \kappa_\gamma^2}{1 - \kappa_\gamma^2} \max_{i \in [b]} \{\gamma_i M_i^2\} \|\mathbf{u}^*\|^2 \\ & + \text{Tr}(\mathbf{D}_{\gamma/\rho} \mathbf{P}_0) + 2 \sum_{i=1}^b d_i. \end{aligned}$$

Rearranging terms concludes the proof. \square

Lemma S39. Assume **H1-H2** and let $\mathbf{N} \in (\mathbb{N}^*)^b, \gamma, \rho \in (\mathbb{R}_+^*)^b$ such that, for any $i \in [b]$, $N_i \gamma_i \leq 2/(m_i + M_i + 1/\rho_i), \gamma_i \tilde{M}_i < 1$ and denote $\mathbf{z}^* = ([\mathbf{A}_1 \boldsymbol{\theta}^*]^\top, \dots, [\mathbf{A}_b \boldsymbol{\theta}^*]^\top)^\top$. Then, we have

$$\int_{\mathbb{R}^p} \|\tilde{\mathbf{z}} - \mathbf{z}^*\|_{\mathbf{D}_{N\gamma}^{-1}}^2 P_{\rho, \gamma, \mathbf{N}}^{\mathbf{z}}(\mathbf{z}^*, d\tilde{\mathbf{z}}) \leq 2 \sum_{i=1}^b \gamma_i N_i (1 + \text{Tr}(\mathbf{P}_0)/\rho_i) + 4 \sum_{i=1}^b d_i.$$

where the transition kernel $P_{\rho, \gamma, \mathbf{N}}^{\mathbf{z}}$ is defined in (S51).

Proof. Let $\{(\eta_k^i)_{k \geq 1} : i \in [b]\}$ be independent random variables such that for any $i \in [b]$, the sequences $\{(\eta_k^i)_{k \geq 1}\}$ are i.i.d. d_i -dimensional Brownian motions and let ξ a d -dimensional standard Gaussian random variable independent of $\{(\eta_k^i)_{k \geq 1} : i \in [b]\}$. Consider the stochastic process $(Y_k)_{k \in \mathbb{N}}$ initialised for any $i \in [b]$ at $Y_0^i = \mathbf{A}_i \boldsymbol{\theta}^*$ and defined, for any $i \in [b], k \in \mathbb{N}$, by

$$Y_{k+1}^i = Y_k^i - \gamma_i \nabla V_i(Y_k^i) + (\gamma_i/\rho_i) \mathbf{A}_i \boldsymbol{\theta} + \sqrt{2\gamma_i} \eta_{k+1}^i, \quad (\text{S132})$$

where the potential $V_i = \mathbf{y}^i \mapsto U_i(\mathbf{y}^i) + \|\mathbf{y}^i\|^2/(2\rho_i)$ and

$$\boldsymbol{\theta} = \bar{\mathbf{B}}_0^{-1} \mathbf{B}_0^\top \tilde{\mathbf{D}}_0^{1/2} \mathbf{z}^* + \bar{\mathbf{B}}_0^{-1/2} \xi. \quad (\text{S133})$$

In addition, we define the random variable $Z = (Z^1, \dots, Z^b)$, for any $i \in [b]$, as

$$Z^i = Y_{N_i}^i.$$

By definition, note that Z is distributed according to $P_{\rho, \gamma, N}^{\mathbf{z}^*}(\mathbf{z}^*, \cdot)$. Define the process $(Y_k = \{Y_k^i\}_{i=1}^b)_{k \in \mathbb{N}}$ valued in $\mathbb{R}^p \times \mathbb{R}^p$ defined for any $i \in [b]$, $k \geq 0$ by

$$Y_k^i = Y_{\min(k, N_i)}^i.$$

and consider the following matrices defined, for any $k \in \mathbb{N}$, by

$$\begin{aligned} \mathbf{H}_{U,k} &= \text{diag} \left(\gamma_1 \int_0^1 \nabla^2 U_1((1-s)Y_k^1 + s\mathbf{z}^*) ds, \right. \\ &\quad \left. \dots, \gamma_b \int_0^1 \nabla^2 U_b((1-s)Y_k^b + s\mathbf{z}^*) ds \right), \\ \mathbf{J}(k) &= \text{diag} \left(\mathbb{1}_{[N_1]}(k+1) \cdot \mathbf{I}_{d_1}, \dots, \mathbb{1}_{[N_b]}(k+1) \cdot \mathbf{I}_{d_b} \right), \end{aligned} \quad (\text{S134})$$

$$\mathbf{C}_k = \mathbf{J}(k)(\mathbf{D}_{\gamma/\rho} + \mathbf{H}_{U,k}), \quad (\text{S135})$$

$$\mathbf{M}_{k+1} = (\mathbf{I}_p - \mathbf{C}_0)^{-1} \dots (\mathbf{I}_p - \mathbf{C}_k)^{-1}, \quad \text{with } \mathbf{M}_0 = \mathbf{I}_p. \quad (\text{S136})$$

Using these notation and (S132), for any $k \in \mathbb{N}$, we get

$$Y_{k+1} - \mathbf{z}^* = (\mathbf{I}_p - \mathbf{C}_k)(Y_k - \mathbf{z}^*) + \mathbf{J}(k) \left(\mathbf{D}_{\gamma/\sqrt{\rho}} \mathbf{B}_0 \theta - \mathbf{D}_{\gamma} \nabla V(\mathbf{z}^*) + \mathbf{D}_{2\gamma}^{1/2} \eta_{k+1} \right).$$

Multiplying the previous equality by $\mathbf{M}_{k+1} \mathbf{D}_{N\gamma}^{-1/2}$, we obtain, for $k \geq 0$,

$$\begin{aligned} \mathbf{M}_{k+1} \mathbf{D}_{N\gamma}^{-1/2} (Y_{k+1} - \mathbf{z}^*) &= \mathbf{M}_k \mathbf{D}_{N\gamma}^{-1/2} (Y_k - \mathbf{z}^*) \\ &\quad + \mathbf{M}_{k+1} \mathbf{J}(k) \mathbf{D}_{N\gamma}^{-1/2} \left(\mathbf{D}_{\gamma/\sqrt{\rho}} \mathbf{B}_0 \theta - \mathbf{D}_{\gamma} \nabla V(\mathbf{z}^*) + \mathbf{D}_{2\gamma}^{1/2} \eta_{k+1} \right). \end{aligned}$$

Summing the previous equality over $k \in \mathbb{N}$ gives

$$\begin{aligned} \mathbf{M}_{\infty} \mathbf{D}_{N\gamma}^{-1/2} (Y_N - \mathbf{z}^*) &= \mathbf{M}_0 \mathbf{D}_{N\gamma}^{-1/2} (Y_0 - \mathbf{z}^*) \\ &\quad + \sum_{k=0}^{\infty} \mathbf{M}_{k+1} \mathbf{J}(k) \mathbf{D}_{N\gamma}^{-1/2} \left(\mathbf{D}_{\gamma/\sqrt{\rho}} \mathbf{B}_0 \theta - \mathbf{D}_{\gamma} \nabla V(\mathbf{z}^*) + \mathbf{D}_{2\gamma}^{1/2} \eta_{k+1} \right). \end{aligned}$$

Multiplying the last equality by $[\mathbf{M}_{\infty}]^{-1}$ and using the fact that $Y_0 = \mathbf{z}^*$, we get

$$\mathbf{D}_{N\gamma}^{-1/2} (Z - \mathbf{z}^*) = \sum_{k=0}^{\infty} [\mathbf{M}_{\infty}]^{-1} \mathbf{M}_{k+1} \mathbf{J}(k) \mathbf{D}_{N\gamma}^{-1/2} \left(\mathbf{D}_{\gamma/\sqrt{\rho}} \mathbf{B}_0 \theta - \mathbf{D}_{\gamma} \nabla V(\mathbf{z}^*) + \mathbf{D}_{2\gamma}^{1/2} \eta_{k+1} \right). \quad (\text{S137})$$

Recall that $\mathbf{P}_0 = \mathbf{B}_0 \bar{\mathbf{B}}_0^{-1} \mathbf{B}_0^{\top}$. Hence, by (S133) and using $\mathbf{P}_0 \mathbf{D}_{\rho}^{-1/2} \mathbf{z}^* = \mathbf{D}_{\rho}^{-1/2} \mathbf{z}^*$, we get

$$\mathbf{D}_{\gamma/\sqrt{\rho}} \mathbf{B}_0 \theta - \mathbf{D}_{\gamma} \nabla V(\mathbf{z}^*) = \mathbf{D}_{\gamma/\sqrt{\rho}} \mathbf{B}_0 \bar{\mathbf{B}}_0^{-1/2} \xi - \mathbf{D}_{\gamma} \nabla U(\mathbf{z}^*).$$

Plugging this equality into (S137) yields

$$\begin{aligned}
\mathbf{D}_{N\gamma}^{-1/2}(Z - \mathbf{z}^*) &= - \sum_{k=0}^{\infty} [\mathbf{M}_{\infty}]^{-1} \mathbf{M}_{k+1} \mathbf{J}(k) \mathbf{D}_{\gamma/N}^{1/2} \nabla U(\mathbf{z}^*) \\
&\quad + \sum_{k=0}^{\infty} [\mathbf{M}_{\infty}]^{-1} \mathbf{M}_{k+1} \mathbf{J}(k) \mathbf{D}_{\gamma/(N\rho)}^{1/2} \mathbf{B}_0 \bar{\mathbf{B}}_0^{-1/2} \xi \\
&\quad + \sqrt{2} \sum_{k=0}^{\infty} [\mathbf{M}_{\infty}]^{-1} \mathbf{M}_{k+1} \mathbf{J}(k) \mathbf{D}_N^{-1/2} \eta_{k+1} .
\end{aligned} \tag{S138}$$

Recall that $[\mathbf{M}_{\infty}]^{-1} \mathbf{M}_{k+1} = (([\mathbf{M}_{\infty}]^{-1} \mathbf{M}_{k+1})^1, \dots, ([\mathbf{M}_{\infty}]^{-1} \mathbf{M}_{k+1})^b)$ is a block-diagonal matrix where, for any $i \in [b]$, $([\mathbf{M}_{\infty}]^{-1} \mathbf{M}_{k+1})^i = \prod_{l=k+1}^{\infty} (\mathbf{I}_{d_i} - \mathbf{C}_l^i)$ where \mathbf{C}_l^i is defined in (S135). In addition, since we suppose for any $i \in [b]$, that $\gamma_i \tilde{M}_i < 1$, Lemma S24 implies

$$\left\| \prod_{l=k+1}^{N_i-1} (\mathbf{I}_{d_i} - \mathbf{C}_l^i) \right\|^2 \leq (1 - \gamma_i \tilde{m}_i)^{2(N_i-k-1)} .$$

We now upper bound separately each term on the right-hand side of (S138). First, using the Cauchy-Schwarz inequality, we have

$$\begin{aligned}
\left\| \sum_{k=0}^{\infty} [\mathbf{M}_{\infty}]^{-1} \mathbf{M}_{k+1} \mathbf{J}(k) \mathbf{D}_{\gamma/N}^{1/2} \right\|^2 &\leq \sum_{i=1}^b (\gamma_i / N_i) \left\| \sum_{k=0}^{\infty} ([\mathbf{M}_{\infty}]^{-1} \mathbf{M}_{k+1})^i \mathbf{J}^i(k) \right\|^2 \\
&\leq \sum_{i=1}^b (\gamma_i / N_i) \left\| \sum_{k=0}^{N_i-1} \prod_{l=k+1}^{N_i-1} (\mathbf{I}_{d_i} - \mathbf{C}_l^i) \right\|^2 \\
&\leq \sum_{i=1}^b \gamma_i \sum_{k=0}^{N_i-1} \left\| \prod_{l=k+1}^{N_i-1} (\mathbf{I}_{d_i} - \mathbf{C}_l^i) \right\|^2 \\
&\leq \sum_{i=1}^b \gamma_i \sum_{k=0}^{N_i-1} (1 - \gamma_i \tilde{m}_i)^{2(N_i-k-1)} \\
&\leq \sum_{i=1}^b N_i \gamma_i .
\end{aligned} \tag{S139}$$

Second, using the same techniques as for the above inequality, we obtain

$$\left\| \sum_{k=0}^{\infty} [\mathbf{M}_{\infty}]^{-1} \mathbf{M}_{k+1} \mathbf{J}(k) \mathbf{D}_{\gamma/(N\rho)}^{1/2} \mathbf{B}_0 \bar{\mathbf{B}}_0^{-1/2} \xi \right\|^2 \leq \sum_{i=1}^b \frac{N_i \gamma_i}{\rho_i} \left\| \mathbf{B}_0 \bar{\mathbf{B}}_0^{-1/2} \xi \right\|^2 \tag{S140}$$

Finally, the third term can be upper-bounded as

$$\mathbb{E} \left[\left\| \sqrt{2} \sum_{k=0}^{\infty} [\mathbf{M}_{\infty}]^{-1} \mathbf{M}_{k+1} \mathbf{J}(k) \mathbf{D}_{\mathbf{N}}^{-1/2} \eta_{k+1} \right\|^2 \right] \leq 2 \sum_{i=1}^b d_i. \quad (\text{S141})$$

Combining (S138), (S139), (S140) and (S141), we get

$$\int_{\mathbb{R}^p} \|\tilde{\mathbf{z}} - \mathbf{z}^*\|_{\mathbf{D}_{\mathbf{N}\gamma}^{-1}}^2 P_{\rho, \gamma, \mathbf{N}}^{\mathbf{z}}(\mathbf{z}^*, d\tilde{\mathbf{z}}) \leq \sum_{i=1}^b \gamma_i N_i (1 + \text{Tr}(\mathbf{P}_0)/\rho_i) + 2 \sum_{i=1}^b d_i.$$

□

Given $\varepsilon > 0$, we are now ready to provide a condition on the number of iterations n_{ε} to achieve $W_2(\mu_{\rho}^* P_{\rho, \gamma, \mathbf{N}}^{n_{\varepsilon}}, \Pi_{\rho, \gamma}) \leq \varepsilon/3$ in the case where for any $i \in [b]$, $m_i = m$, $M_i = M$, $\rho_i = \rho$, $\gamma_i = \gamma$ and $N_i = N$. Define

$$\begin{aligned} \mathbf{E}_0^2 = & 9(1 + \|\bar{\mathbf{B}}_0^{-1} \mathbf{B}_0^{\top} \tilde{\mathbf{D}}_0^{1/2}\|) 2N\gamma \left[\frac{2}{N(1 - \kappa_{\gamma}^2)} \left(\frac{1 + \kappa_{\gamma}^2}{1 - \kappa_{\gamma}^2} \cdot \gamma M^2 \|\mathbf{u}^*\|^2 \right. \right. \\ & \left. \left. + (\gamma/\rho) \text{Tr}(\mathbf{P}_0) + 2 \sum_{i=1}^b d_i \right) + 2b\gamma N (1 + \text{Tr}(\mathbf{P}_0)/\rho) + 4 \sum_{i=1}^b d_i \right]. \end{aligned}$$

Theorem S40. Assume **H1-H2** and assume that for any $i \in [b]$, $m_i = m$ and $M_i = M$. In addition, let $\mathbf{N} = N\mathbf{1}_b$, $\gamma = \gamma\mathbf{1}_b$, $\rho = \rho\mathbf{1}_b$, $\rho > 0, \gamma > 0, N \geq 1$, such that $\gamma < 1/\tilde{M}$, $N\gamma < 2/(m + \tilde{M})$, and (S46) is satisfied. Then, for any $\varepsilon > 0$, any

$$n_{\varepsilon} \geq 2 \log(\mathbf{E}_0/\varepsilon) / (N\gamma m),$$

we have, $W_2(\mu_{\rho}^* P_{\rho, \gamma, \mathbf{N}}^{n_{\varepsilon}}, \Pi_{\rho, \gamma}) \leq \varepsilon/3$.

Proof. By some algebra and using $1/\log(1/(1-x)) \leq 1/x$ for $0 < x < 1$, the proof directly follows from Proposition S14 combined with Proposition S38 and Lemma S39. □

S5.2 Upper bound on the tolerance parameter ρ_ε

Define

$$\begin{aligned}
R_0 &= 2\sigma_U^2 \left(d\sigma_U^2 + \sum_{i=1}^b M_i^2 \|\mathbf{A}_i(\boldsymbol{\theta}^* - \boldsymbol{\theta}_i^*)\|^2 \right) + 2\sigma_U^4, \\
R_1 &= d\sigma_U^2 + \sum_{i=1}^b M_i^2 \|\mathbf{A}_i(\boldsymbol{\theta}^* - \boldsymbol{\theta}_i^*)\|^2 + \sum_{i=1}^b d_i M_i / 2 \\
R_2 &= 2d \max_{i \in [b]} \{M_i\} \sigma_U^2 + 2 \sum_{i=1}^b M_i^3 \|\mathbf{A}_i(\boldsymbol{\theta}^* - \boldsymbol{\theta}_i^*)\|^2 + 8\sigma_U^4 + 8\sigma_U^2 \left[2d\sigma_U^2 \right. \\
&\quad \left. + 2 \sum_{i=1}^b M_i^2 \|\mathbf{A}_i(\boldsymbol{\theta}^* - \boldsymbol{\theta}_i^*)\|^2 \right].
\end{aligned}$$

Recall that $\bar{\rho} = \max_{i \in [b]} \{\rho_i\}$. Then, the following result holds.

Lemma S41. Assume **H1-H2**. For any $\varepsilon > 0$, let $\boldsymbol{\rho}_\varepsilon \in (\mathbb{R}_+^*)^b$ such that

$$\begin{aligned}
\bar{\rho}_\varepsilon &\leq \frac{-R_1 + \sqrt{R_1^2 + 4R_0\varepsilon m_U^{1/2}/(3\sqrt{2})}}{2R_0} \wedge \frac{\varepsilon\sqrt{m_U}}{3\sqrt{2}\sqrt{R_2 + [R_2/(12\sigma_U^2) + \sum_{i=1}^b d_i M_i]^2}} \\
&\wedge \frac{1}{12\sigma_U^2} \wedge \frac{-\sum_{i=1}^b d_i M_i + \sqrt{(\sum_{i=1}^b d_i M_i)^2 + 6R_2}}{2R_2}.
\end{aligned}$$

Then, $W_2(\pi_{\boldsymbol{\rho}_\varepsilon}, \pi) \leq \varepsilon/3$.

Proof. Let $\varepsilon > 0$. From (S73), for any $\bar{\rho} \leq 1/(12\sigma_U^2)$, $W_2(\pi_{\boldsymbol{\rho}}, \pi) \leq \sqrt{\frac{2}{m_U}} \max(A_1, A_3^{1/2})$, where A_1, A_3 are defined in (S69) and (S72) respectively. This implies that $W_2(\pi_{\boldsymbol{\rho}}, \pi) \leq \varepsilon/3$ is verified if $\max(A_1, A_3^{1/2}) \leq \varepsilon\sqrt{m_U}/(3\sqrt{2})$. First, $A_1 \leq \varepsilon\sqrt{m_U}/(3\sqrt{2})$ holds if

$$\bar{\rho} \leq \frac{-R_1 + \sqrt{R_1^2 + 4R_0\varepsilon m_U^{1/2}/(3\sqrt{2})}}{2R_0} \wedge \frac{1}{12\sigma_U^2}. \quad (\text{S142})$$

We now focus on A_3 . Using the fact that for any $x \in \mathbb{R}$, $e^x \geq x + 1$, we have $2 \prod_{i=1}^b (1 + \rho_i M_i)^{d_i} \geq 2 + \sum_{i=1}^b d_i \log(1 + \rho_i M_i)$ and therefore

$$A_3 \leq \exp \left(\bar{\rho}^2 R_2 + \sum_{i=1}^b d_i \log(1 + \rho_i M_i) \right) - 1 - \sum_{i=1}^b d_i \log(1 + \rho_i M_i).$$

Since $\sum_{i=1}^b d_i \log(1 + \rho_i M_i) \leq \bar{\rho} \sum_{i=1}^b d_i M_i$, $\bar{\rho}^2 R_2 + \sum_{i=1}^b d_i \log(1 + \rho_i M_i) \leq 3/2$ holds for

$$\bar{\rho} \leq \frac{-\sum_{i=1}^b d_i M_i + \sqrt{(\sum_{i=1}^b d_i M_i)^2 + 6R_2}}{2R_2}. \quad (\text{S143})$$

Since for any $x \leq 3/2$, $e^x \leq 1 + x + x^2$ and using the fact that $\bar{\rho} \leq 1/(12\sigma_U^2)$, it follows that

$$A_3 \leq \bar{\rho}^2 R_2 + \left(\bar{\rho}^2 R_2 + \bar{\rho} \sum_{i=1}^b d_i M_i \right)^2 \leq \bar{\rho}^2 \left[B_1 + \left(\frac{R_2}{12\sigma_U^2} + \sum_{i=1}^b d_i M_i \right)^2 \right].$$

Hence $A_3^{1/2} \leq \varepsilon \sqrt{m_U}/(3\sqrt{2})$ holds under (S143) and

$$\bar{\rho} \leq \frac{\varepsilon \sqrt{m_U}}{3\sqrt{2} \sqrt{R_2 + \left(\frac{R_2}{12\sigma_U^2} + \sum_{i=1}^b d_i M_i \right)^2}}. \quad (\text{S144})$$

The proof is concluded by combining (S142), (S143) and (S144). \square

S5.3 Upper bound on the step-size γ_ε and number of local iteration N_ε

Based on Proposition S29 or Proposition S33, we now determine an upper bound on γ_ε to ensure $W_2(\Pi_\rho, \Pi_{\rho, \gamma_\varepsilon}) \leq \varepsilon/3$ in the case $\mathbf{N} = N\mathbf{1}_b$, $\gamma = \gamma\mathbf{1}_b$, $\rho = \rho\mathbf{1}_b$ where $\rho > 0$, $\gamma > 0$, $N \geq 1$. The following results hold depending if H3 is considered. Define

$$C_\rho = \frac{4\tilde{M}^2(1 + \|\bar{\mathbf{B}}_0^{-1}\mathbf{B}_0^\top \tilde{\mathbf{D}}_0^{1/2}\|^2)}{5m}, \quad (\text{S145})$$

$$C_0 = (\tilde{M}^2/2) \left[\tilde{M}/\tilde{m} + 1/6 \right] \sum_{i=1}^b d_i, \quad C_1 = \sum_{i=1}^b d_i, \quad C_2 = \varepsilon^2/(9C_\rho).$$

Lemma S42. Assume H1-H2 and assume for any $i \in [b]$, $m_i = m$ and $M_i = M$. In addition, let $\rho, \gamma_\varepsilon > 0$ and $N_\varepsilon \geq 1$ such that $\rho = \rho\mathbf{1}_b$, $\gamma_\varepsilon = \gamma_\varepsilon\mathbf{1}_b$, $\mathbf{N}_\varepsilon = N_\varepsilon\mathbf{1}_b$ and $\varepsilon > 0$ satisfying

$$\gamma_\varepsilon \leq \frac{-C_1 + \sqrt{C_1^2 + 4C_0C_2}}{2C_0} \wedge \frac{m}{40\tilde{M}^2}. \quad (\text{S146})$$

Then $W_2(\Pi_\rho, \Pi_{\rho, \gamma_\varepsilon}) \leq \varepsilon/3$.

Proof. Let $\varepsilon > 0$. By Proposition S29, note that $W_2^2(\Pi_\rho, \Pi_{\rho, \gamma_\varepsilon}) \leq \varepsilon^2/9$ is satisfied if

$$C_0\gamma_\varepsilon^2 + C_1\gamma_\varepsilon \leq C_2.$$

This inequality is satisfied under the choice (S147). \square

We now provide a condition on \mathbf{N} and γ when **H3** is considered.

Lemma S43. *Assume **H1-H2** and assume for any $i \in [b]$, $m_i = m$, $M_i = M$ and $L_i = L$. In addition, let $\rho, \gamma_\varepsilon > 0$ and $N_\varepsilon \geq 1$ such that $\boldsymbol{\rho} = \rho \mathbf{1}_b$, $\boldsymbol{\gamma}_\varepsilon = \gamma_\varepsilon \mathbf{1}_b$, $\mathbf{N}_\varepsilon = N_\varepsilon \mathbf{1}_b$ and $\varepsilon > 0$ satisfying*

$$\gamma_\varepsilon \leq \frac{\varepsilon}{6b\sqrt{5 \max_{i \in [b]} \{d_i\} C_\rho \tilde{M}^2 [4 + (\max_{i \in [b]} \{d_i\} L^2 m)/(20\tilde{M}^4)]}} \wedge \frac{m}{40\tilde{M}^2} \quad (\text{S147})$$

$$\wedge \frac{\varepsilon}{6b(5C_\rho \max_{i \in [b]} \{d_i\} m^3 / \tilde{M}^2)} , \quad (\text{S148})$$

where C_ρ is defined in (S145). Then $W_2(\Pi_\rho, \Pi_{\rho, \gamma_\varepsilon}) \leq \varepsilon/3$.

Proof. In Proposition S33, we dissociate $R^*(\gamma)$ into two contributions and the conditions we impose on γ_ε ensure $W_2(\Pi_\rho, \Pi_{\rho, \gamma_\varepsilon}) \leq \varepsilon/3$. More precisely, we have $\sum_{i=1}^b d_i \gamma_i^2 \tilde{M}_i^2 + \frac{d_i \gamma_i^2 \mathfrak{f}_i}{M_i} (d_i L_i^2 + \frac{\tilde{M}_i^4}{m_i}) \leq 2\varepsilon^2/9$ and $\sum_{i=1}^b d_i \gamma_i \tilde{M}_i \mathfrak{f}_i^3 (1 + \mathfrak{f}_i + \mathfrak{f}_i^2) \leq 2\varepsilon^2/9$ where $\mathfrak{f}_i < 1$ for any $i \in [b]$. \square

S5.4 Discussion

Let $\boldsymbol{\rho}_\varepsilon = \rho_\varepsilon \mathbf{1}_b$ such that $W_2(\pi_{\boldsymbol{\rho}_\varepsilon}, \pi) \leq \varepsilon/3$. From Lemma S41, $\rho_\varepsilon = \mathcal{O}(\varepsilon/d)$ when $\varepsilon \rightarrow 0$ and $d \rightarrow \infty$. Similarly, let $\boldsymbol{\gamma}_\varepsilon = \gamma_\varepsilon \mathbf{1}_b$ such that $W_2(\Pi_{\boldsymbol{\rho}_\varepsilon}, \Pi_{\boldsymbol{\rho}_\varepsilon, \boldsymbol{\gamma}_\varepsilon}) < \varepsilon/3$. Under **H1-H2**, we obtain by Lemma S42 $\gamma_\varepsilon = \mathcal{O}(\varepsilon^4/d^3)$. On the other hand, when **H3** is additionally assumed, we get by Lemma S43 $\gamma_\varepsilon = \mathcal{O}(\varepsilon^2/d^2)$. Finally, to apply Theorem S40 for the previous choices γ_ε and ρ_ε , we obtain for $\mathbf{N}_\varepsilon = N_\varepsilon \mathbf{1}_b$ the conditions $N_\varepsilon = \mathcal{O}(d/\varepsilon^2)$ and $N_\varepsilon = \mathcal{O}(1)$ under **H1-H2** and **H1-H2-H3**, respectively. In both scenarios, Theorem S40 implies $n_\varepsilon = \mathcal{O}(d^2 \log(d)/(\varepsilon^2 |\log(\varepsilon)|))$. This concludes the results depicted in Table 1 in the main paper.