

REDUCTION COHOMOLOGY OF RIEMANN SURFACES

A. ZUEVSKY

ABSTRACT. We study the algebraic conditions leading to the chain property of complexes for vertex operator algebra n -point functions (with their convergence assumed) with differential being defined through reduction formulas. The notion of the reduction cohomology of Riemann surfaces is introduced. Algebraic, geometrical, and cohomological meanings of reduction formulas is clarified. A counterpart of the Bott-Segal theorem for Riemann surfaces in terms of the reductions cohomology is proven. It is shown that the reduction cohomology is given by the cohomology of n -point connections over the vertex operator algebra bundle defined on a genus g Riemann surface $\Sigma^{(g)}$. The reduction cohomology for a vertex operator algebra with formal parameters identified with local coordinates around marked points on $\Sigma^{(g)}$ is found in terms of the space of analytical continuations of solutions to Knizhnik-Zamolodchikov equations. For the reduction cohomology, the Euler-Poincare formula is derived. Examples for various genera and vertex operator cluster algebras are provided.

1. INTRODUCTION

The natural problem of computing continuous cohomologies for non-commutative structures on manifolds has proven to be a subject of great geometrical interest [BS, Kaw, PT, Fei, Fuks, Wag]. As it was demonstrated in [Fei, Wag], the ordinary Gelfand-Fuks cohomology of the Lie algebra of holomorphic vector fields on complex manifolds turns to be not the most effective and general one. For Riemann surfaces, and even for higher dimension complex manifolds, the classical cohomology of vector fields becomes trivial [Kaw]. The Lie algebra of holomorphic vector fields is not always an interesting Lie algebra. For example, it is zero for a compact Riemann surface of genus greater than one, and one looks for other algebraic objects having locally the same cohomology. In [Fei] Feigin obtained various results concerning (co)-homology of the Lie algebra cosimplicial objects of holomorphic vector fields $Lie(M)$. Despite results in previous approaches, it is desirable to find a way to enrich cohomological structures which motivates constructions of more refined cohomology description for non-commutative algebraic structures. In the seminal paper [BS], the authors proved that the Gelfand-Fuks cohomology $H^*(Vect(M))$ of vector fields on a smooth compact manifold M is isomorphic to the singular cohomology of the space of continuous cross sections of a certain fibre bundle over M . An important problem of revealing relations between non-commutative structures and geometrical objects on complex manifolds still remains underinvestigated in the literature [PT].

Key words and phrases. Cohomology; Vertex algebras; Riemann surfaces; Cluster algebras.

The main idea of this paper is to introduce and compute the reduction cohomology vertex operator algebras [B, FLM, FHL, K] with formal parameters considered as local coordinates on a genus g compact Riemann surface [FK, Bo, Gu, A] is to study cohomology of spaces of converging functions with respect to adding new sets of pairs of vertex operator algebra elements and corresponding formal parameters [BZF, DGM, DL, EO, FMS]. Due to structure of correlations functions [FMS] and reduction relations [Y, Zhu, MTZ, GT, TW] among them, one can form chain complexes of n -point functions (with their convergence assumed) that are fine enough to describe local geometry of Riemann surfaces. Another meaning of the reduction cohomology of Riemann surfaces is how sections of certain bundles with values in higher genus generalizations of elliptic functions change along growing of number of marked points on a surface. In addition to that we prove a version of the Bott-Segal [BS, Wag, PT] theorem for compact Riemann surfaces of arbitrary genus. It relates the reduction cohomology with cohomology of a space of sections of certain vertex operator algebra bundle [BZF]. Our new vertex operator algebra approach to cohomology of compact Riemann surfaces involves Lie-algebraic formal series with applications of techniques used in surgery of spheres [Huang]. In contrast to more geometrical methods, e.g., in ordinary cosimplicial cohomology for Lie algebras [Fei, Wag], the reduction cohomology pays more attention to the analytical structure of elements of chain complex spaces. Computational methods involving reduction formulas proved their effectiveness in conformal field theory, geometrical descriptions of intertwined modules for Lie algebras, and differential geometry of integrable models. In section 2 we give the definition of the reduction cohomology and prove a proposition relating it to cohomology of a vertex operator algebra bundle in terms of n -point connections. In section 3 the main proposition expressing the reduction cohomology in terms of spaces of auxiliary functions on Riemann surfaces is proven. Results of this paper are useful for cosimplicial cohomology theory of smooth manifolds, generalizations of the Bott-Segal theorem have their consequences in conformal field theory [Fei, Wag], deformation theory [O], non-commutative geometry, modular forms, and the theory of foliations.

2. CHAIN COMPLEX FOR VERTEX OPERATOR ALGEBRA n -POINT FUNCTIONS

2.1. Spaces of n -point correlation functions. In this section we recall definitions and some properties of correlation functions for vertex operator algebras on Riemann surfaces [FHL, Zhu, FMS, DGM]. Let us fix a vertex operator algebra V . Depending on its commutation relations and configuration of a genus g Riemann surface $\Sigma^{(g)}$, the space of all V n -point functions can contain various forms of complex functions defined on $\Sigma^{(g)}$.

We denote by $\mathbf{v}_n = (v_1, \dots, v_n) \in V^{\otimes n}$ a tuple of vertex operator algebra elements (see Appendix 5 for definition of a vertex operator algebra). Mark n points $\mathbf{p}_n = (p_1, \dots, p_n)$ on a Riemann surface of genus $\Sigma^{(g)}$. Denote by $\mathbf{z}_n = (z_1, \dots, z_n)$ local coordinates around $\mathbf{p}_n \in \Sigma^{(g)}$. Let us introduce the notation: $\mathbf{x}_n = (\mathbf{v}_n, \mathbf{z}_n)$. On a genus g Riemann surfaces an $n \geq 0$ -point correlation function $\mathcal{F}_n^{(g)}(\mathbf{x}_n, B^{(g)})$ (here $B^{(g)}$ denotes the set of its parameters) has certain specific form depending on g ,

construction of a Riemann surface $\Sigma^{(g)}$, type of conformal field theory model used for definitions of n -point functions, and the type of commutation relations for V -elements. In addition to that, it depends on a set of moduli parameters $B^{(g)} \in \mathcal{B}^{(g)}$ where we denote by $\mathcal{B}^{(g)}$ a subset of the moduli space of genus g Riemann surfaces $\Sigma^{(g)}$ obtained by specific ways of sewing of lower genus Riemann surfaces. In particular, for n -point functions considered on Riemann surfaces, $B^{(g)}$ is an element of the space $[\mathbf{Y}]$ $\mathcal{B}^{(g)} = (\Omega^{(i)}; (\epsilon_i); (\rho_i); (\mu_i))$, $0 \leq i \leq g$, where vectors (ϵ_i) , (ρ_i) , are Riemann surface sewing parameters $[\mathbf{Y}]$, (μ_i) are any further modular space parameters, and $\Omega^{(i)}$ is the period matrix of corresponding Riemann surface $\Sigma^{(g_i)}$ used in the procedure of $\Sigma^{(g)}$ construction.

Definition 1. For a Riemann surface Σ of genus g , a V -module W , and $n \geq 0$, \mathbf{x}_n on Σ , we consider the spaces of n -point correlation functions

$$C_{(g)}^n(W) = \left\{ \mathcal{F}_{W,n}^{(g)}(\mathbf{x}_n, B^{(g)}) \right\}.$$

Note that we choose the same V -module W for all \mathbf{x}_n . In what follows, we will omit where possible W and $B^{(g)}$. The co-boundary operator $\delta_{(g)}^n(v_{n+1})$ on $C_{(g)}^n(W)$ -space is defined according to the reduction formulas for V -module W on a genus g Riemann surface (cf. particular examples in subsections 4.1–4.5, [Zhu, MTZ, GT, TW]).

Definition 2. For $g \geq 0$, $n \geq 0$, and any $x_{n+1} \in V \times \mathbb{C}$, define

$$\begin{aligned} \delta_{(g)}^n : C_{(g)}^n(W) &\rightarrow C_{(g)}^{n+1}(W), \\ \delta_n^{(g)} = H^{(g)} &= H_1^{(g)} + H_2^{(g)}, \end{aligned} \quad (2.1)$$

with operators $H_1^{(g)}(x_{n+1})$, $H_2^{(g)}(x_{n+1})$ given by

$$\begin{aligned} H_1^{(g)}(x_{n+1}) \mathcal{F}_n^{(g)}(\mathbf{x}_n) &= \sum_{l=1}^{l(g)} f_1^{(g)}(\mathbf{x}_{n+1}, l) T_l^{(g)} \mathcal{F}_n^{(g)}(\mathbf{x}_n), \\ H_2^{(g)}(x_{n+1}) \mathcal{F}_n^{(g)}(\mathbf{x}_n) &= \sum_{k=1}^n \sum_{m \geq 0} f_2^{(g)}(\mathbf{x}_{n+1}, k, m) \\ &\quad \cdot T_k^{(g)}(v_{n+1}(m)) \mathcal{F}_{W,n}^{(g)}(\mathbf{x}_n), \end{aligned} \quad (2.2)$$

where $l(g) \geq 0$ is a constant depending on g , and the meaning of indexes $1 \leq k \leq n$, $1 \leq l \leq l(g)$, $m \geq 0$ explained below.

Operator-valued functions $f_1^{(g)}(\mathbf{x}_{n+1}, l)$, $f_2^{(g)}(\mathbf{x}_{n+1}, k, m)$, $T_k^{(g)}(v_{n+1}(m))$ depend on genus of a Riemann surface $\Sigma^{(g)}$. $T_l^{(g)}$ and $T_k^{(g)}(v(m))$ are operators of insertion of certain function of vertex operator algebra modes into $\mathcal{F}_{W,n}^{(g)}(\mathbf{x}_n)$ at the k -th entry:

$$T_{l,k}^{(g)}(v(m)) \mathcal{F}_{W,n}^{(g)}(\mathbf{x}_n) = \mathcal{F}_{W,n}^{(g)}\left(\dots, \left(T_l^{(g)}(v(m)).x_n\right)_k, \dots\right),$$

where we use the notation

$$(\gamma \cdot)_k x_n = (x_1, \dots, \gamma.x_k, \dots, x_n),$$

for an operator γ acting on k -th entry. Note that commutation properties of $H_1^{(g)}(x_{n+1})$ and $H_2^{(g)}(x_{n+1})$ depend on genus g . The reduction formulas have the form:

$$\mathcal{F}_{W,n+1}^{(g)}(\mathbf{x}_{n+1}) = H^{(g)}(x_{n+1}) \mathcal{F}_{W,n}^{(g)}(\mathbf{x}_n), \quad (2.3)$$

for $n \geq 0$.

Remark 1. The author's conjecture (based on [Y, MT, MTZ, TZ, GT, TW, BKT]) is that for all possible configurations of a genus g Riemann surface, the form of the reduction relations (2.2) is coinvariant and it is given by a sum of operators acting on the an n -point correlation function

$$\mathcal{F}_{W,n}^{(g)}(\mathbf{x}_{n+1}) = \sum_{\substack{1 \leq l \leq l^{(g)} \\ m \geq 0, k \geq 0}} f^{(g)}(\mathbf{x}_{n+1}, k, l, m) T^{(g)}(k, l, m) \mathcal{F}_{W,n}^{(g)}(\mathbf{x}_n).$$

Remark 2. The reductions formulas have an interpretation in terms of torsors [BZF] (Chapter 6). In such formulation \mathbf{x}_n is a torsor with respect to the group of transformation of the space $V^{\otimes n} \times \mathbb{C}^n$. In particular, from (2.3) we see that $T_k^{(g)}(u(m))$ -operators act on $V^{\otimes n}$ -entries of \mathbf{x}_n in $\mathcal{F}_{W,n}^{(g)}(\mathbf{x}_n, B^{(g)})$, while $T_{k,l}^{(g)}$ -operators act on \mathbf{z}_n of $\mathcal{F}_{W,n}^{(g)}(\mathbf{x}_n, B^{(g)})$ as a complex function.

For $n \geq 0$, let us denote by \mathfrak{V}_n the subsets of all $\mathbf{x}_{n+1} \in V^{\otimes(n+1)} \times \mathbb{C}^{n+1}$, such that the chain condition

$$H^{(g)}(x_{n+1}) H^{(g)}(x_n) \mathcal{F}_{W,n}^{(g)}(\mathbf{x}_n) = 0, \quad (2.4)$$

for the coboundary operators (2.2) for complexes $C_{(g)}^n(W)$ is satisfied.

Explicitly, the chain condition (2.4) leads to an infinite $n \geq 0$ set of equations involving functions $f_1^{(g)}(\mathbf{x}_{n+1}, l)$, $f_2^{(g)}(\mathbf{x}_{n+1}, k, m)$, and $\mathcal{F}_{W,n}^{(g)}(\mathbf{x}_n)$:

$$\begin{aligned} & \left(\sum_{l'=1, l=1}^{l^{(g)}} f_1^{(g)}(\mathbf{x}_{n+2}, l') f_1^{(g)}(\mathbf{x}_{n+1}, l; B^{(g)}) T_{l'}^{(g)} T_l^{(g)} \right. \\ & + \sum_{l'=1, k=1}^{l^{(g)}} \sum_{m \geq 0}^n f_1^{(g)}(\mathbf{x}_{n+2}, l') f_2^{(g)}(m, k) T_{l'}^{(g)} T_k^{(g)}(v_{n+1}(m)) \\ & + \sum_{k'=1}^{n+1} \sum_{m' \geq 0}^{l^{(g)}} f_2^{(g)}(k', m') f_1^{(g)}(\mathbf{x}_{n+2}, l) T_{k'}^{(g)}(v_{n+1}(m')) T_l^{(g)} \\ & \left. + \sum_{k'=1}^{n+1} \sum_{k=1}^n \sum_{\substack{m', \\ m \geq 0}} f_2^{(g)}(k', m') f_2^{(g)}(k, m) T_{k'}^{(g)}(v_{n+2}(m')) T_k^{(g)}(v_{n+1}(m)) \right) \\ & \cdot \mathcal{F}_{W,n}^{(g)}(\mathbf{x}_n) = 0. \end{aligned} \quad (2.5)$$

Remark 3. (2.5) contain finite series and narrows the space of compatible n -point functions. The subspaces of $C_{(g)}^n(W)$, $g \geq 0$, $n \geq 0$, of genus g n -point functions such that the condition (2.5) is fulfilled for reduction cohomology complexes are non-empty.

Indeed, for all g , the condition (2.5) represents an infinite $n \geq 0$ set of functional-differential equations (with finite number of summands) on converging complex functions $\mathcal{F}_{W,n}^{(g)}(\mathbf{x}_n)$ defined for n local complex variables on a Riemann surface of genus g with functional coefficients $f_1^{(g)}(v_{n+1}, l)$ and $f_2^{(g)}(k, m)$ (in our examples in subsection 4.1–4.5, these are generalizations of genus g elliptic functions) on $\Sigma^{(g)}$. Note that (see examples in Section 4), all vertex operator algebra elements of $\mathbf{v}_n \in V^{\otimes n}$, as non-commutative parameters are not present in final form of functional-differential equations since they incorporated into either matrix elements, traces, etc. According to the theory of such equations [FK, Gu], each equation in the infinite set of (2.5) always have a solution in domains they are defined. Thus, there always exist solutions of (2.5) defining $\mathcal{F}_{W,n}^{(g)} \in C_{(g)}^n(W)$, and they are not empty.

Definition 3. The spaces with conditions (2.5) constitute a semi-infinite chain complex

$$0 \longrightarrow C_{(g)}^0 \xrightarrow{\delta_{(g)}^0} C_{(g)}^1 \xrightarrow{\delta_{(g)}^1} \dots \longrightarrow C_{(g)}^{n-1} \xrightarrow{\delta_{(g)}^{n-1}} C_{(g)}^n \longrightarrow \dots \quad (2.6)$$

For $n \geq 1$, we call corresponding cohomology

$$H_{(g)}^n = \text{Ker } \delta_{(g)}^n / \text{Im } \delta_{(g)}^{n-1}, \quad (2.7)$$

the n -th reduction cohomology of a vertex operator algebra V -module W on a compact Riemann surface $\Sigma^{(g)}$ of genus g .

In particular, the operators $T_l^{(g)}$, $0 \leq l \leq l(g)$, $T_k^{(g)}(u(m))$, $m \geq 0$, $1 \leq k \leq n$, form a set of generators of an infinite-dimensional continual Lie algebra $\mathfrak{g}(V)$ endowed with a natural grading indexed by l and m .

Indeed, we set the space of functions $\mathcal{F}_{W,n}^{(g)}$ as the base algebra [Sav] for the continual Lie algebra $\mathfrak{g}(V)$, and the generators as

$$\begin{aligned} X_{0,l} \left(\mathcal{F}_{W,n}^{(g)}(\mathbf{x}_n) \right) &= T_l^{(g)} \left(\mathcal{F}_{W,n}^{(g)}(\mathbf{x}_n) \right), \\ X_{k,m} \left(\mathcal{F}_{W,n}^{(g)}(\mathbf{x}_n) \right) &= T_k^{(g)}(u(m)) \left(\mathcal{F}_{W,n}^{(g)}(\mathbf{x}_n) \right). \end{aligned} \quad (2.8)$$

for $0 \leq l \leq l(g)$, $m \geq 0$, $1 \leq k \leq n$. Then the commutation relations for vertex operator algebra modes $v_{n+1}(m)$ in the action of operators $T_l^{(g)}$ and $T_{k,l}^{(g)}$ on v_k , $1 \leq k \leq n$ inside $\mathcal{F}_{W,n}^{(g)}$ represent the commutation relations of the continual Lie algebra $\mathfrak{g}(V)$. Jacobi identities for $\mathfrak{g}(V)$ follow from Jacobi identities (5.4) for a vertex operator algebra V .

Remark 4. Recall that we consider genus g Riemann surfaces resulting from combinations of sewing procedures of [Y]. Accordingly, due to [MT, TZ, TZ1, TZ2, GT, TW], corresponding genus g n -point functions are obtained coherently by combining lower genus functions. Then, relations among n -point functions of various genera appear. One is able to consider a cohomology theory taking into account such relations. For

instance, for the ϵ -formalism of [Y] one has

$$\begin{aligned} C_n^{(g)}(W) &\rightarrow C_{n_1+1}^{(g_1)}(W_1) \times C_{n_2+1}^{(g_2)}(W_2), \\ \mathcal{F}_{W,n}^{(g)}(\mathbf{x}_n) &= \sum_{l \geq 0} \epsilon^l \mathcal{F}_{W,n+1}^{(g_1)}\left((\bar{u}, z), \mathbf{x}_n; B^{(g_1)}\right) \mathcal{F}_{w,n+1}^{(g_2)}\left((u, z), \mathbf{x}'_n; B^{(g_2)}\right), \end{aligned} \quad (2.9)$$

for $g = g_1 + g_2$, $n = n_1 + n_2$, and $W = W_1 \otimes W_2$. In the ρ -formalism [Y] one has

$$\begin{aligned} C_n^{(g)}(W) &\rightarrow C_{n_1+1}^{(g-1)}(W), \\ \mathcal{F}_{W,n}^{(g)}(\mathbf{x}_n) &= \sum_{l \geq 0} \rho^l \mathcal{F}_{W,n+2}^{(g-1)}\left((u, z), \mathbf{x}_n; (\bar{u}, z), \mathbf{x}'_n; B^{(g-1)}\right). \end{aligned} \quad (2.10)$$

2.2. Geometrical meaning of reduction formulas and conditions (2.5). In this section we show that the reduction formulas have the form of multipoint connections generalizing ordinary holomorphic connections on complex curves [BZF].

2.2.1. Holomorphic n -point connections. Let us define the notion of a multipoint connection which will be useful for identifying reduction cohomology in section 3. Motivated by the definition of a holomorphic connection for a vertex operator algebra bundle (cf. Section 6, [BZF] and [Gu]) over a smooth complex curve, we introduce the definition of the multiple point connection over $\Sigma^{(g)}$.

Definition 4. Let \mathcal{V} be a holomorphic vector bundle over $\Sigma^{(g)}$, and $\mathcal{X}_0 \subset \Sigma^{(g)}$ be its subdomain. Denote by \mathcal{SV} the space of sections of \mathcal{V} . A multi-point connection \mathcal{G} on \mathcal{V} is a \mathbb{C} -multi-linear map

$$\mathcal{G} : \left(\Sigma^{(g)}\right)^{\times n} \times V^{\otimes n} \rightarrow \mathbb{C},$$

such that for any holomorphic function f , and two sections $\phi(p)$ and $\psi(p')$ of \mathcal{V} at points p and p' on $\mathcal{X}_0 \subset \Sigma^{(g)}$ correspondingly, we have

$$\sum_{q, q' \in \mathcal{X}_0 \subset \Sigma^{(g)}} \mathcal{G}(f(\psi(q)).\phi(q')) = f(\psi(p')) \mathcal{G}(\phi(p)) + f(\phi(p)) \mathcal{G}(\psi(p')), \quad (2.11)$$

where the summation on left hand side is performed over locuses of points q, q' on \mathcal{X}_0 . We denote by Con_n the space of n -point connections defined over $\Sigma^{(g)}$.

Geometrically, for a vector bundle \mathcal{V} defined over $\Sigma^{(g)}$, a multi-point connection (2.11) relates two sections ϕ and ψ at points p and p' with a number of sections on $\mathcal{X}_0 \subset \Sigma^{(g)}$.

Definition 5. We call

$$G(\phi, \psi) = f(\phi(p)) \mathcal{G}(\psi(p')) + f(\psi(p')) \mathcal{G}(\phi(p)) - \sum_{q, q' \in \mathcal{X}_0 \subset \Sigma^{(g)}} \mathcal{G}(f(\psi(q')).\phi(q)), \quad (2.12)$$

the form of a n -point connection \mathcal{G} . The space of n -point connection forms will be denoted by G^n .

Here we prove the following

Proposition 1. *n -point correlation functions of the space $\left\{ \mathcal{F}_{W,n}^{(g)}(\mathbf{x}_n), n \geq 0 \right\}$ form n -point connections on the space of sections of the vertex operator algebra bundle \mathcal{V} associated to V . For $n \geq 0$, the reduction cohomology of a compact Riemann surface of genus g is*

$$H_{(g)}^n(W) = H_{(g)}^n(\mathcal{SV}) = \text{Con}^n / G^{n-1}, \quad (2.13)$$

is isomorphic to the cohomology of the space of \mathcal{V} -sections.

Remark 5. Proposition 1 is a vertex operator algebra version of the main proposition of [BS, Wag], i.e., the Bott-Segal theorem for Riemann surfaces.

Proof. In [BZF] (Chapter 6, subsection 6.5.3) the vertex operator bundle \mathcal{V} was explicitly constructed. It is easy to see that n -point connections are holomorphic connection on the bundle \mathcal{V} with the following identifications. For non-vanishing $f(\phi(p))$ let us write (2.3) as

$$\mathcal{G}(\psi(p')) = -\frac{f(\psi(p'))}{f(\phi(p))} \mathcal{G}(\phi(p)) + \frac{1}{f(\phi(p))} \sum_{q,q' \in \mathcal{X}_0 \subset \mathcal{X}} \mathcal{G}(f(\psi(q)).\phi(q')). \quad (2.14)$$

Let us set

$$\begin{aligned} \mathcal{G} &= \mathcal{F}_{W,n}^{(g)}(\mathbf{x}_n), \\ \psi(p') &= (\mathbf{x}_{n+1}), \\ \phi(p) &= (\mathbf{x}_n), \\ \mathcal{G}(f(\psi(q)).\phi(q')) &= T_k^{(g)}(v(m)) \mathcal{F}_{W,n}^{(g)}(\mathbf{x}_n), \\ -\frac{f(\psi(p'))}{f(\phi(p))} \mathcal{G}(\phi(p)) &= \sum_{l=1}^{l(g)} f_1^{(g)}(v_{n+1}, l) T_l^{(g)} \mathcal{F}_{W,n}^{(g)}(\mathbf{x}_n), \\ \frac{1}{f(\phi(p))} \sum_{\substack{q_n, q'_n \in \\ \mathcal{X}_0 \subset \Sigma^{(g)}}} \mathcal{G}(f(\psi(q)).\phi(q')) &= \sum_{k=1}^n \sum_{m \geq 0} f_2^{(g)}(k, m) \\ &\quad \cdot T_k^{(g)}(v(m)) \mathcal{F}_{W,n}^{(g)}(\mathbf{x}_n). \end{aligned} \quad (2.15)$$

Thus, the formula (2.15) gives (2.3). Recall [BZF] the construction of the vertex operator algebra bundle \mathcal{V} . According to Proposition 6.5.4 of [BZF], one canonically (i.e., coordinate independently) associates $\text{End} \mathcal{V}$ -valued sections \mathcal{Y}_p of \mathcal{V}^* (the bundle dual to \mathcal{V}) to matrix elements of a number of vertex operators on appropriate punctured disks around points with local coordinates \mathbf{z}_n on $\Sigma^{(g)}$. The spaces of such \mathcal{V} -sections for each n are described by identifications (2.15). Taking into account the construction of Section 6 (subsection 6.6.1, in particular, construction 6.6.4, and Proposition 6.6.7) of [BZF], we see that n -point functions are connections on the space of sections of \mathcal{V} , and the reduction cohomology (2.7) is represented by (2.13). \square

For the chain condition (2.5) we have

$$\begin{aligned}
0 = \mathcal{G}(\chi(p'')) &= -\frac{f(\chi(p''))}{f(\psi(p'))} \mathcal{G}(\psi(p')) + \frac{1}{f(\psi(p'))} \sum_{\tilde{q}_n, \tilde{q}'_n \in \mathcal{X}_0 \subset \Sigma^{(g)}} \mathcal{G}(f(\chi(\tilde{q})) \cdot \psi(\tilde{q}')), \\
0 &= \mathcal{G}(\chi(p'')) \\
&= \frac{f(\chi(p''))}{f(\phi(p))} \mathcal{G}(\phi(p)) - \frac{f(\chi(p''))}{f(\psi(p')) f(\phi(p))} \sum_{q_n, q'_n \in \mathcal{X}_0 \subset \Sigma^{(g)}} \mathcal{G}(f(\psi(q)) \cdot \phi(q')) \\
&+ \frac{1}{f(\psi(p'))} \sum_{\tilde{q}_n, \tilde{q}'_n \in \mathcal{X}_0 \subset \Sigma^{(g)}} \mathcal{G}(f(\chi(\tilde{q})) \cdot \psi(\tilde{q}')), \tag{2.16}
\end{aligned}$$

The geometrical meaning of (2.5) consists in the following. Since in (2.2) operators act on vertex operator algebra elements only, we can interpret it as a relation on modes of V with functional coefficients. In particular, all operators T change vertex operator algebra elements by action either of $o(v) = v_{\text{wt}v-1}$. or positive modes of $v(m)$., $m \geq 0$. Recall that for all g the n -point functions possess certain modular properties with respect to groups depending on genus g and modular space $\mathcal{B}^{(g)}$ parameters. Moreover, the reduction formulas (2.3) are used to prove modular invariance for n -point functions. Due to automorphic properties of n -point functions, (2.5) can be also interpreted as relations among modular forms. The condition (2.3) defines a complex variety in $\mathbf{z}_n \in \mathbb{C}^n$ with non-commutative parameters $\mathbf{v}_n \in V^{\otimes n}$. As most identities (e.g., trisecant identity [Fa, TZ1] and triple product identity [K, MTZ, TZ2]) for n -point functions (2.5) has its algebraic-geometrical meaning. The condition (2.5) relates finite series of vertex operator algebra correlations functions on a genus g Riemann surface with rational function coefficients (at genus $g = 0$) [FHL], or elliptic functions (at genus $g = 1$) [Zhu, MT, MTZ], or generalizations of classical elliptic functions (at genus $g \geq 2$) [GT, TW]. Since in all cases n -point functions possess certain modular properties, we treat (2.5) as a source of new identities on modular forms at corresponding genus of Riemann surfaces.

3. COHOMOLOGY

In this section we compute the reduction cohomology defined by (2.6)–(2.7). The main result of this paper is the following.

Proposition 2. *The n -th reduction cohomology of a vertex operator algebra V -module W is the space of analytical continuations of solutions $\mathcal{F}_{W,n}^{(g)}(\mathbf{x}_n)$ to a Knizhnik-Zamolodchikov equation, and provided by series of auxiliary functions recursively generated by reduction formulas (2.3) with $x_i \notin \mathfrak{V}_i$, for $1 \leq i \leq n$.*

Remark 6. The first cohomology is given by the space of transversal (i.e., with vanishing sum over q, q') one-point connections $\mathcal{F}_{W,1}^{(g)}(x_1)$ provided by coefficients in terms of series of generalizations of elliptic functions (cf. Appendix 6). The second cohomology is given by a space of generalized higher genus complex kernels corresponding to V and $\Sigma^{(g)}$.

Proof. By definition (2.7), the n -th reduction cohomology is defined by the subspace of $C_n^{(g)}(W)$ of functions $\mathcal{F}_{W,n}^{(g)}(\mathbf{x}_n)$ satisfying

$$\begin{aligned} & \left(\sum_{l=1}^{l(g)} f_1^{(g)}(\mathbf{x}_{n+1}, l) T_l^{(g)}(v_{n+1}) \right. \\ & \quad \left. + \sum_{k=1}^n \sum_{m \geq 0} f_2^{(g)}(k, m) T_k^{(g)}(v_{n+1}(m)) \right) \mathcal{F}_{W,n}^{(g)}(\mathbf{x}_n) = 0, \end{aligned} \quad (3.1)$$

modulo the subspace of $C_n^{(g)}(W)$ n -point functions $\mathcal{F}_{W,n}^{(g)}(\mathbf{x}'_n)$ resulting from:

$$\begin{aligned} \mathcal{F}_{W,n}^{(g)}(\mathbf{x}'_n) &= \left(\sum_{l=1}^{l(g)} f_1^{(g)}(\mathbf{x}'_n, l) T_l^{(g)} \right. \\ & \quad \left. + \sum_{k=1}^{n-1} \sum_{m \geq 0} f_2^{(g)}(k, m) T_k^{(g)}(v'_n(m)) \right) \mathcal{F}_{W,n-1}^{(g)}(\mathbf{x}'_{n-1}). \end{aligned} \quad (3.2)$$

We assume that, subject to other fixed parameters, n -point functions are completely determined by all choices $\mathbf{x}_n \in V^{\otimes n} \times \mathbb{C}^n$. Thus, the reduction cohomology can be treated as depending on set of \mathbf{x}_n only with appropriate action of endomorphisms generated by x_{n+1} . Consider a non-vanishing solution $\mathcal{F}_{W,n}^{(g)}(\mathbf{x}_n)$ to (3.1) for some \mathbf{x}_n . Let us use the reduction formulas (2.3) recursively for each x_i , $1 \leq i \leq n$ of \mathbf{x}_n in order to express $\mathcal{F}_{W,n}^{(g)}(\mathbf{x}_n)$ in terms of the partition function $\mathcal{F}_{W,0}^{(g)}(B^{(g)})$, i.e., we obtain

$$\mathcal{F}_{W,n}^{(g)}(\mathbf{x}_n) = \mathcal{D}^{(g)}(\mathbf{x}_n) \mathcal{F}_{W,0}^{(g)}(B^{(g)}), \quad (3.3)$$

as in [MT, MTZ, TZ, TZ1, TZ2]. It is clear that $x_i \notin \mathfrak{V}_i$ for $1 \leq i \leq n$, i.e., at each stage of the recursion procedure towards (3.3), otherwise $\mathcal{F}_{W,n}^{(g)}(\mathbf{x}_n)$ would be zero. Thus, $\mathcal{F}_{W,n}^{(g)}(\mathbf{x}_n)$ is explicitly known and is represented as a series of auxiliary functions $\mathcal{D}^{(g)}$ depending on V , genus g , and moduli space parameters $B^{(g)}$. Consider now $\mathcal{F}_{W,n}^{(g)}(\mathbf{x}'_n)$ given by (3.2). It either vanishes when $v_{n-i} \in \mathfrak{V}_{n-i}$, $2 \leq i \leq n$, or given by (3.3) with \mathbf{x}'_n arguments.

The way the reduction relations (2.3) were derived in [MT, MTZ, GT, TW, BKT] is exactly the same as for the vertex operator algebra derivation [KZ, TK] for the Knizhnik-Zamolodchikov equations. The general idea is consider the double integration of $\mathcal{F}_{W,n}^{(g)}(\mathbf{x}_n)$ along small circles around two auxiliary variables with the action of reproduction kernels inserted. Then, these procedure leads to recursion formulas relating $\mathcal{F}_{W,n+1}^{(g)}$ and $\mathcal{F}_{W,n}^{(g)}$ with functional coefficients depending on the nature of the vertex operator algebra V , and the way a Riemann surface $\Sigma^{(g)}$ was constructed. Thus, in our context, (3.1) is seen as a version of the Knizhnik-Zamolodchikov equation. In [Y, MT, MTZ] formulas to n -point functions in various specific examples of V and configuration of Riemann surfaces were explicitly and recursively obtained.

In terms of x_{n+1} , by using (5.1)–(5.6), we are able to transfer in (3.1) the action of v_{n+1} -modes into an analytical continuation of $\mathcal{F}_{W,n}^{(g)}(\mathbf{x}_n)$ multi-valued holomorphic functions to domains $D_n \subset \Sigma^{(g)}$ with $z_i \neq z_j$ for $i \neq j$. Namely, in (3.1), the operators $T_l^{(g)}$ and $T_k^{(g)}(v_n(m))$ act by certain modes $v_{n+1}(m)$ of a vertex operator algebra element v_{n+1} on $\mathbf{v}_n \in V^{\otimes n}$. Using vertex operator algebra associativity property (5.6) we express the action of operators $T_l^{(g)}$ and $T_k^{(g)}(v_n(m))$ in terms of modes $v_{n+1}(m)$ inside vertex operators in actions of V -modes on the whole vertex operator at expense of a shift of their formal parameters \mathbf{z}_n by z_{n+1} , i.e., $\mathbf{z}'_n = \mathbf{z}_n + z_{n+1}$. Note that under such associativity transformations v -part of \mathbf{x}_n , i.e., \mathbf{v}_n remains the untouched. Thus, the n -th reduction cohomology of a V -module W is given by the space of analytical continuations of n -point functions $\mathcal{F}_{W,n}^{(g)}(\mathbf{x}_n)$ with $\mathbf{x}_{n-1} \notin \mathfrak{B}_{n-1}$ that are solutions to the Knizhnik-Zamolodchikov equations (3.1). The above analytic extensions for the Knizhnik-Zamolodchikov equations generated by x_{n+1} and with coefficients provided by functions $f_1^{(g)}(\mathbf{x}_{n+1}, n)$ and $f_2^{(g)}(\mathbf{x}_{n+1}, k, m)$ on the Riemann surface $\Sigma^{(g)}$ \square

This result is coherent with considerations of [TUY]. We illustrate the proposition above in cases of zero, one, two, and higher genera in section 4. One can make connection with the first cohomology of grading-restricted vertex operator algebras in terms of derivations, and to the second cohomology in terms of square-zero extensions of V by W [Huang2].

Euler-Poincare formula. In [Fuks], for a Lie algebra, we find a celebrated formula relating sums over dimensions of chain complex spaces and dimensions of homologies for a graded infinite-dimensional Lie algebra. Suppose a V -module W is endowed with a complex grading $W = \bigcup_{m \in \mathbb{C}} W_{(m)}$, i.e., one could consider $m = m' + \beta(m)$, where $m' \in \mathbb{Z}$, and $\beta(m) \in \mathbb{C}$ [MTZ]. In our case all spaces $C_{(g)}^n(W)$ are infinite-dimensional, but, according to definition of a vertex operator algebra, $\dim W_{(m)} \leq \infty$, $m \in \mathbb{C}$. Thus, for a fixed g , each $C_{(g)}^n(W)$ can be endowed with separation with respect to m . Then the complex (2.6) decomposes into sum of complexes

$$0 \longrightarrow C_{(g)}^0(m) \xrightarrow{\delta_{(g)}^0} C_{(g)}^1(m) \xrightarrow{\delta_{(g)}^1} \dots \longrightarrow C_{(g)}^{n-1}(m) \xrightarrow{\delta_{(g)}^{n-1}} C_{(g)}^n(m) \longrightarrow \dots \quad (3.4)$$

For $g \geq 0$, $l \geq 0$, let

$$q_{n,m} = \dim C_{(g)}^n(V_{(m)}) = \dim \left\{ \mathcal{F}_{W,n}^{(g)}(\mathbf{x}_n) : \mathbf{x}_n \in V_{(m)}^{\otimes n}, m \in \mathbb{C} \right\}.$$

Let

$$p_{m,n} = \dim H^n(V_{(m)}) = \dim \left\{ \left(\text{Ker } \delta_{(g)}^n / \text{Im } \delta_{(g)}^{n-1} \right) |_{V_{(m)}}, v_{n+1} \in V_{(m)} \right\},$$

be dimensions of corresponding cohomology spaces. In our context we find that for fixed $m \in \mathbb{C}$, and $n \geq 0$,

$$\sum_{n \geq 0}^N (-1)^n (q_{n,m} - p_{n,m}) = 0, \quad (3.5)$$

Indeed, let us consider relations of chain complex spaces and cohomology spaces for vertex operator algebra elements that belong to fixed $V_{(m)}$ grading subspace

of V . Recall that according to the definition of a vertex operator algebra (see Appendix 5) subspaces $W_{(m)}$ are finite dimensional. Thus, as a functional space, $\{\mathcal{F}_{W,n}^{(g)}(\mathbf{x}_n), \mathbf{v}_n \in V_{(m)}^{\otimes n}\}$ is finite-dimensional. Consider now the cohomology spaces $\{H_{(g)}^n, \mathbf{v}_{n+1} \in V_{(m)}^{\otimes(n+1)}\}$. Let us fix $v_{n+1} \in V_{(m)}$. For the same reason as above, $\text{rank}(\text{Ker } \delta_{(g)}^n) < \infty$, $\text{rank}(\text{Im } \delta_{(g)}^{n-1}) < \infty$, as subspaces of $n+1$ - and n -point functions. Thus, $\text{rank} H_{(g)}^n|_{V_{(m)}} < \infty$. Using then the standard Euler-Poincaré formula [Fuks] considerations for rank we obtain (3.5). The formula (3.5) has deep number-theoretical meanings as equality of generating functions for series expansions for automorphic forms.

4. EXAMPLES

The reduction cohomology depends on actual coefficients of the Knizhnik-Zamolodchikov equations (3.1). Note that for $n = 0$, $\mathcal{F}_{W,0}^{(g)}(B^{(g)})$, $g \geq 0$, is the partition function (or graded dimension for $g = 1$), for a vertex operator algebra V module W . In this section we provide examples of vertex operator algebras considered on Riemann surfaces of various genus g .

4.1. Vertex operator algebra n -point functions on the sphere. For $u, \mathbf{v}_n \in V$, and a homogeneous $u \in V$, we find the formula [FHL, FLM] for the n -point functions

$$\mathcal{F}_{W,n}^{(0)}(\mathbf{x}_n) = \langle u', \mathbf{Y}(\mathbf{x}_n)u \rangle,$$

where

$$\mathbf{Y}(\mathbf{x}_n) = Y(x_1) \dots Y(x_n),$$

The partition function is given by

$$\mathcal{F}_{W,0}^{(0)} = \langle u'_{(a)}, u_{(b)} \rangle = \delta_{a,b}.$$

The reduction operators of (2.2) are [FHL]

$$\begin{aligned} H_1^{(0)}(\mathbf{x}_{n+1}) \mathcal{F}_{W,n}^{(0)}(\mathbf{x}_n) &= T_1(o(v)) \mathcal{F}_{W,n}^{(0)}(\mathbf{x}_n), \\ H_2^{(0)}(\mathbf{x}_{n+1}) \mathcal{F}_{W,n}^{(0)}(\mathbf{x}_n) &= \sum_{k=1}^n \sum_{m \geq 0} f_{wt(v_{n+1}),m}(z_{n+1}, z_r) T_k(v(m)) \mathcal{F}_{W,n}^{(0)}(\mathbf{x}_n), \end{aligned} \quad (4.1)$$

where we define $f_{wt(v,m)}^{(0)}(z, w)$ is a rational function defined by

$$\begin{aligned} f_{n,m}^{(0)}(z, w) &= \frac{z^{-n}}{m!} \left(\frac{d}{dw} \right)^m \frac{w^n}{z-w}, \\ \iota_{z,w} f_{n,m}^{(0)}(z, w) &= \sum_{j \in \mathbb{N}} \binom{n+j}{m} z^{-n-j-1} w^{n+j-1}. \end{aligned}$$

Let us take x_{n+1} as the variable of expansion. The n -th reduction cohomology $H_{(0)}^n(W)$ is given the space of solutions to the Knizhnik-Zamolodchikov equation (4.4)

of the n -point function $\mathcal{F}_{W,0}^{(0)}(\mathbf{x}_n)$ (not given by $\delta_0^{n-1} \mathcal{F}_{W,n-1}^{(0)}(\mathbf{x}_{n-1})$)

$$\begin{aligned} \mathbf{Y}(\mathbf{x}_n) &\mapsto o(v_{n+1}) \mathbf{Y}(\mathbf{x}_n), \\ \mathbf{x}_k &\mapsto v_{(m)} \cdot \mathbf{x}_k, \end{aligned} \quad (4.2)$$

(i.e., generated by x_{n+1} -endomorphisms) of solutions $\mathcal{F}_{V,n}^{(0)}$ of the Knizhnik-Zamolodchikov equation

$$0 = \mathcal{F}_{W,n}^{(0)}(o(v_{n+1}) \mathbf{x}_n) + \sum_{k=1}^n \sum_{m \geq 0} f_{wt(v_{n+1}),m}^{(0)}(z_{n+1}, z_k) \mathcal{F}_{W,n}^{(0)}((v_{n+1}(m))_k \cdot \mathbf{x}_n), \quad (4.3)$$

with rational function coefficients $f_{wt(v_{n+1}),m,k}^{(0)}(z_{n+1}, z_k)$, modulo the space of n -point functions obtained by the recursion procedure. Using (5.1)–(5.6) we obtain from (4.3), and the standard Virasoro algebra representation $L_V(0) = z_{n+1} \partial_{z_{n+1}}$, we obtain

$$\left(\partial_{z_{n+1}} + \sum_{k=1}^n \tilde{f}_{wt(v_{n+1}),m}^{(0)}(z_{n+1}, z_k) \right) \mathcal{F}_{W,n}^{(0)}(\mathbf{x}_n + (z_{n+1})_k) = 0, \quad (4.4)$$

which is the Knizhnik-Zamolodchikov equation on analytical continuation of $\mathcal{F}_{W,n}^{(0)}(\mathbf{x}_n + (z_{n+1})_k)$ with a different $\tilde{f}^{(0)}$. Using the reduction formulas (2.3) we obtain

$$\mathcal{F}_{W,n}^{(0)}(\mathbf{x}_n + (z_{n+1})_k) = \mathcal{D}^{(0)}(\mathbf{x}_{n+1}),$$

where $\mathcal{D}^{(0)}(\mathbf{x}_{n+1})$ is given by the series of rational-valued functions in $\mathbf{x}_{n+1} \notin \mathfrak{V}_n$ resulting from the recursive procedure starting from n -point function to the partition function. Thus, in this example, the n -th cohomology is the space of analytic extensions [FHL, FLM] of rational function solutions to the equation (4.3) with rational function coefficients.

4.2. Vertex operator algebra n -point functions on the torus. In order to consider modular-invariance of n -point functions at genus one, Zhu introduced [Zhu] a second square-bracket VOA $(V, Y[.,.], \mathbf{1}_V, \tilde{\omega})$ associated to a given VOA $(V, Y(.,.), \mathbf{1}_V, \omega)$. The new square bracket vertex operators are

$$Y[v, z] = \sum_{n \in \mathbb{Z}} v[n] z^{-n-1} = Y(q_z^{L(0)} v, q_z - 1),$$

with $q_z = e^z$, while the new conformal vector is

$$\tilde{\omega} = \omega - \frac{c_V}{24} \mathbf{1}_V.$$

For v of $L(0)$ weight $wt(v) \in \mathbb{R}$ and $m \geq 0$,

$$v[m] = m! \sum_{i \geq m} c(wt(v), i, m) v(i),$$

where

$$\sum_{m=0}^i c(wt(v), i, m) x^m = \binom{wt(v) - 1 + x}{i}.$$

For $\mathbf{v}_n \in V^{\otimes n}$ the genus one n -point function [Zhu] has the form

$$F_n^{(1)}(\mathbf{x}_n; \tau) = \text{Tr}_V \left(\mathbf{Y} \left(\mathbf{q}^{L(0)} \mathbf{v}_n, \mathbf{q}_n \right) q^{L(0) - c_V/24} \right),$$

for $q = e^{2\pi i \tau}$ and $q_i = e^{z_i}$, where τ is the torus modular parameter, and c_V is the central charge of V -Virasoro algebra. Then the genus one Zhu recursion formula is given by the following [Zhu]. For any v_{n+1} , $\mathbf{v}_n \in V^{\otimes n}$ we find for an $n+1$ -point function

$$\begin{aligned} H_1^{(1)}(\mathbf{x}_{n+1}) &= F_n^{(1)}(o(v_{n+1}) \mathbf{x}_n; \tau) \\ H_2^{(1)}(\mathbf{x}_{n+1}) &= \sum_{k=1}^n \sum_{m \geq 0} P_{m+1}(z_{n+1} - z_k, \tau) F_V^{(1)}((v[m])_k, \mathbf{x}_n; \tau). \end{aligned} \quad (4.5)$$

In this theorem $P_m(z, \tau)$ denote higher Weierstrass functions defined by

$$P_m(z, \tau) = \frac{(-1)^m}{(m-1)!} \sum_{n \in \mathbb{Z} \neq 0} \frac{n^{m-1} q_z^n}{1 - q^n}.$$

The partition function

$$\mathcal{F}_{W,0}^{(1)}(\tau) = \text{Tr}_V \left(q^{L_V(0) - c/24} \right), \quad (4.6)$$

is called the graded dimension for V . Let us fix $\mathbf{x}_{n+1} \in V^{\otimes n} \times \mathbb{C}^n$. The n -th reduction cohomology H^n is the space of extensions of the Knizhnik-Zamolodchikov equation solutions

$$0 = F_V^{(1)}(o(v_{n+1}) \mathbf{x}_n; \tau) + \sum_{k=1}^{n+1} \sum_{m \geq 0} P_{m+1}(z_{n+1} - z_k, \tau) F_n^{(1)}((v_{n+1}[m])_k, \mathbf{x}_n; \tau). \quad (4.7)$$

This space can be described by complex kernels (i.e., prime forms [Mu]) given by sums of elliptic functions.

4.3. Genus two n -point functions. In this subsection we recall [GT] the construction and reduction formulas for vertex operator algebra n -point functions on a genus two Riemann surface formed in the sewing procedure of two tori due to [Y]. Let us assume that a vertex operator algebra V is of strong-type. Thus, it possess a non-degenerate bilinear form. For a V -basis $\{u^{(a)}\}$ we define the dual basis $\{\bar{u}^{(a)}\}$ with respect to the bilinear form where

$$\langle \bar{u}^{(a)}, u^{(b)} \rangle_{\text{sq}} = \delta_{ab}.$$

Definition 6. The genus two partition function (zero-point function) for V is defined by

$$\mathcal{F}_{V,0}^{(2)}(B^{(2)}) = \sum_{r \geq 0} \epsilon^r \sum_{\bar{u} \in V_{[r]}} \mathcal{F}_{V,1}^{(1)}(u; \tau_1) \mathcal{F}_{V,1}^{(1)}(u; \tau_2), \quad (4.8)$$

where $B^{(g)} = (\tau_1, \tau_2, \epsilon)$ in the ϵ -sewing procedure for constructing a genus two Riemann surface [Y, MT, TZ1], and the internal sum is taken over any $V_{[n]}$ -basis, and \bar{u} is the dual of u with respect to the bilinear form on V .

We then recall [GT] a formal genus two reduction formulas for all n -point functions.

Definition 7. Let $v_{n+1} \in V$ be inserted at $x_{n+1} \in \widehat{\Sigma}_1^{(1)}$, $\mathbf{v}_k \in V^{\otimes k}$ be inserted at $\mathbf{y}_k \in \widehat{\Sigma}_1^{(1)}$ and $\mathbf{v}'_l \in V^{\otimes l}$ be inserted at $\mathbf{y}'_l \in \widehat{\Sigma}_2^{(1)}$, a punctured torus

$$\widehat{\Sigma}_a^{(1)} = \Sigma_a^{(1)} \setminus \{z_a, |z_a| \leq |\epsilon|/r_a\}, \quad (4.9)$$

where we here we use the convention

$$\bar{1} = 2, \quad \bar{2} = 1. \quad (4.10)$$

We consider the corresponding genus two n -point function

$$\begin{aligned} \mathcal{F}_{V,n}^{(2)}(x_{n+1}; \mathbf{x}_k; \mathbf{x}'_l; B^{(2)}) \\ = \sum_{r \geq 0} \epsilon^r \sum_{u \in V_{[r]}} \mathcal{F}_{V,k+1}^{(1)}(Y[x_{n+1}] \mathbf{Y}[\mathbf{x}_k]_k u; \tau_1) \mathcal{F}_{V,l}^{(1)}(\mathbf{Y}[\mathbf{x}'_l]_l \bar{u}; \tau_2). \end{aligned} \quad (4.11)$$

where the sum as in (4.8).

Remark 7. (4.8) is independent of the choice of V -basis. One could also write a similar expression by inserting the $Y[x]$ vertex operator at $z \in \widehat{\Sigma}_2^{(1)}$ on the right hand side of (4.11).

First, one defines the functions $\mathcal{F}_{V,n,a}^{(2)}$ for $a \in \{1, 2\}$, by

$$\begin{aligned} \mathcal{F}_{V,n,1}^{(2)}(\mathbf{x}_{n+1}; B^{(2)}) &= \sum_{r \geq 0} \epsilon^r \sum_{u \in V_{[r]}} \text{Tr}_V \left(o(v_{n+1}) \mathbf{Y}(\mathbf{q}_{\mathbf{z}_k}^{\mathbf{L}(0)} \mathbf{v}_k, \mathbf{q}_{\mathbf{z}_k}) q_1^{L_V(0) - c_V/24} \right) \\ &\quad \cdot \mathcal{F}_{V,n-k}^{(1)}(\mathbf{Y}[\mathbf{x}_{k+1,n}] \bar{u}; \tau_2), \end{aligned}$$

$$\begin{aligned} \mathcal{F}_{V,n,2}^{(2)}(\mathbf{x}_{n+1}; B^{(2)}) &= \sum_{r \geq 0} \epsilon^r \sum_{u \in V_{[r]}} \mathcal{F}_{V,k}^{(1)}(\mathbf{Y}[\mathbf{x}_k] u; \tau_1) \\ &\quad \cdot \text{Tr}_V \left(o(v_{n+1}) \mathbf{Y}(\mathbf{q}_{\mathbf{z}_{k+1,n}}^{\mathbf{L}(0)} \mathbf{v}_{k+1}, \mathbf{q}_{\mathbf{z}_{k+1}}) q_2^{L_V(0) - c_V/24} \right), \end{aligned}$$

and $\mathcal{F}_{V,n,3}^{(2)}(\mathbf{x}_{n+1}; B^{(2)}) = \mathbb{X}_1^\Pi$ of (6.7). We also define

Definition 8. Let $f_a^{(2)}(p; z_{n+1})$, for $p \geq 1$, and $a = 1, 2$ be given by

$$f_a^{(2)}(p; z_{n+1}) = 1^{\delta_{ba}} + (-1)^{p\delta_{b\bar{a}}} \epsilon^{1/2} \left(\mathbb{Q}(p; z_{n+1}) \left(\tilde{\Lambda}_{\bar{a}} \right)^{\delta_{ba}} \right) (1), \quad (4.12)$$

for $z_{n+1} \in \widehat{\Sigma}_b^{(1)}$. Let $f_3^{(2)}(p; z_{n+1})$, for $z_{n+1} \in \Sigma_a^{(1)}$, be an infinite row vector given by

$$f_3^{(2)}(p; z_{n+1}) = \left(\mathbb{R}(z_{n+1}) + \mathbb{Q}(p; z_{n+1}) \left(\tilde{\Lambda}_{\bar{a}} \Lambda_a + \Lambda_{\bar{a}} \Gamma \right) \right) \Pi. \quad (4.13)$$

In [GT] it is proven that the genus two $n = k + l$ -point function for a quasi-primary vector v_{n+1} of weight $p = \text{wt}[v_{n+1}]$ inserted at $x \in \widehat{\Sigma}_1^{(1)}$, and general vectors \mathbf{v}_k and

\mathbf{v}'_l inserted at $\mathbf{x}_k \in \widehat{\Sigma}_1^{(1)}$, respectively, has the following operators $H_1^{(2)}(\mathbf{x}_{n+1})$ and $H_2^{(2)}(\mathbf{x}_{n+1})$:

$$\begin{aligned} H_1^{(2)}(x_{n+1}) \mathcal{F}_{V,n}^{(2)}(\mathbf{x}_n; B^{(2)}) &= \sum_{l=1}^3 f_l^{(2)}(p; z_{n+1}) \mathcal{F}_{V,n,l}^{(2)}(\mathbf{x}_{n+1}; B^{(2)}), \\ H_2^{(2)}(x_{n+1}) \mathcal{F}_{V,n}^{(2)}(\mathbf{x}_n; B^{(2)}) &= \sum_{i=1}^n \sum_{j \geq 0} \mathcal{P}_{j+1}(p; z_{n+1}, z_i) \mathcal{F}_{V,n}^{(2)}((v_{n+1}[j])_i \cdot \mathbf{x}_n; B^{(2)}), \end{aligned} \quad (4.14)$$

with $\mathcal{P}_{j+1}(p; x, y)$ of (6.10). A similar expression corresponds to v_{n+1} inserted on $x \in \widehat{\Sigma}_2^{(1)}$.

Remark 8. The functions $\mathcal{P}_{j+1}(p; x, y)$ are holomorphic for $x \neq y$ on the sewing domain in the cases $p = 1, 2$.

4.4. Vertex operator algebra n -point functions and reduction formulas in genus g Schottky parameterization. In this section we recall [TZ, TW, T2] the construction and reduction relations for vertex operator algebra n -point functions defined on a genus g Riemann surface formed in the Schottky parameterization. In particular, the formal partition and n -point correlation functions for a vertex operator algebra associated to a genus g Riemann surface $\Sigma^{(g)}$ are introduced in the Schottky scheme. All expressions here are functions of formal variables $w_{\pm a}$, $\rho_a \in \mathbb{C}$ and vertex operator parameters. Then we recall the genus g reduction formula with universal coefficients that have a geometrical meaning and are meromorphic on a Riemann surface $\Sigma^{(g)}$. These coefficients are generalizations of the elliptic Weierstrass functions [L]. For a $2g$ vertex operator algebra V states

$$\mathbf{b} = (b_{-1}, b_1; \dots; b_{-g}, b_g),$$

and corresponding local coordinates

$$\mathbf{w} = (w_{-1}, w_1; \dots; w_{-g}, w_g),$$

of $2g$ points $(p_{-1}, p_1; \dots; p_{-g}, p_g)$ on the Riemann sphere $\Sigma^{(0)}$, consider the genus zero $2g$ -point correlation function

$$\begin{aligned} Z^{(0)}(\mathbf{b}, \mathbf{w}) &= Z^{(0)}(b_{-1}, w_{-1}; b_1, w_1; \dots; b_{-g}, w_{-g}; b_g, w_g) \\ &= \prod_{a \in \mathcal{I}_+} \rho_a^{\text{wt}(b_a)} Z^{(0)}(\bar{b}_1, w_{-1}; b_1, w_1; \dots; \bar{b}_g, w_{-g}; b_g, w_g). \end{aligned}$$

where $\mathcal{I}_+ = \{1, 2, \dots, g\}$. Let

$$\mathbf{b}_+ = (b_1, \dots, b_g),$$

denote an element of a V -tensor product $V^{\otimes g}$ -basis with dual basis

$$\mathbf{b}_- = (b_{-1}, \dots, b_{-g}),$$

with respect to the bilinear form $\langle \cdot, \cdot \rangle_{\rho_a}$ (cf. Appendix 5).

Let w_a for $a \in \mathcal{I}$ be $2g$ formal variables. One identify them with the canonical Schottky parameters (for details of the Schottky construction, see [TW, T2]). One can define the genus g partition function as

$$Z_V^{(g)} = Z_V^{(g)}(\mathbf{w}, \boldsymbol{\rho}) = \sum_{\mathbf{b}_+} Z^{(0)}(\mathbf{b}, \mathbf{w}), \quad (4.15)$$

for

$$(\mathbf{w}, \boldsymbol{\rho}) = (w_{\pm 1}, \rho_1; \dots; w_{\pm g}, \rho_g).$$

Now we recall the formal reduction formulas for all genus g Schottky n -point functions. One defines the genus g formal n -point function for n vectors $\mathbf{v}_n \in V^{\otimes n}$ inserted at \mathbf{y}_n by

$$Z_V^{(g)}(\mathbf{v}, \mathbf{y}) = Z_V^{(g)}(\mathbf{v}, \mathbf{y}; \mathbf{w}, \boldsymbol{\rho}) = \sum_{\mathbf{b}_+} Z^{(0)}(\mathbf{v}, \mathbf{y}; \mathbf{b}, \mathbf{w}), \quad (4.16)$$

where

$$Z^{(0)}(\mathbf{v}, \mathbf{y}; \mathbf{b}, \mathbf{w}) = Z^{(0)}(v_1, y_1; \dots; v_n, y_n; b_{-1}, w_{-1}; \dots; b_g, w_g).$$

Let U be a vertex operator subalgebra of V where V has a U -module decomposition

$$V = \bigoplus_{\alpha \in A} W_\alpha,$$

for U -modules W_α and some indexing set A . Let

$$W_\alpha = \bigotimes_{a=1}^g W_{\alpha_a},$$

denote a tensor product of g modules

$$Z_{W_\alpha}^{(g)}(\mathbf{v}, \mathbf{y}) = \sum_{\mathbf{b}_+ \in W_\alpha} Z^{(0)}(\mathbf{v}, \mathbf{y}; \mathbf{b}, \mathbf{w}), \quad (4.17)$$

where here the sum is over a basis $\{\mathbf{b}_+\}$ for W_α . It follows that

$$Z_V^{(g)}(\mathbf{v}, \mathbf{y}) = \sum_{\alpha \in A} Z_{W_\alpha}^{(g)}(\mathbf{v}, \mathbf{y}), \quad (4.18)$$

where the sum ranges over $\alpha = (\alpha_1, \dots, \alpha_g) \in A$, for $A = A^{\otimes g}$. Finally, one defines corresponding formal n -point correlation differential forms

$$\begin{aligned} \mathcal{F}_V^{(g)}(\mathbf{v}, \mathbf{y}) &= Z_V^{(g)}(\mathbf{v}, \mathbf{y}) d\mathbf{y}^{\text{wt}(\mathbf{v})}, \\ \mathcal{F}_{W_\alpha}^{(g)}(\mathbf{v}, \mathbf{y}) &= Z_{W_\alpha}^{(g)}(\mathbf{v}, \mathbf{y}) d\mathbf{y}^{\text{wt}(\mathbf{v})}, \end{aligned} \quad (4.19)$$

where

$$d\mathbf{y}^{\text{wt}(\mathbf{v})} = \prod_{k=1}^n dy_k^{\text{wt}(v_k)}.$$

Recall notations and identifications given in Appendix 6.3. In [TW] they prove that the genus g $(n+1)$ -point formal differential $\mathcal{F}_{W_\alpha}^{(g)}(x; \mathbf{v}, \mathbf{y})$, for $x_{n+1} = (v_{n+1}, y_{n+1})$, for a quasiprimary vector $v_{n+1} \in U$ of weight $\text{wt}(v_{n+1}) = p$ inserted at a point p_0 , with

the coordinate y_{n+1} , and general vectors \mathbf{v}_n inserted at points \mathbf{p}_n with coordinates \mathbf{y}_n satisfies the recursive identity for $\mathbf{x}_n = (\mathbf{v}, \mathbf{y})$

$$\begin{aligned} \mathcal{F}_{W,n+1,\alpha}^{(g)}(x_{n+1}; \mathbf{x}_n) &= \left(H_1^{(g)} + H_2^{(g)} \right) \mathcal{F}_{W,n}^{(g)}(\mathbf{x}_n), \\ H_1^{(g)}(\mathbf{x}_{n+1}) \mathcal{F}_{W,n}^{(g)}(\mathbf{x}_n) &= \sum_{a=1}^g \Theta_a(y_{n+1}) O_a^{W,\alpha}(v_{n+1}; \mathbf{x}_n), \\ H_2^{(g)}(\mathbf{x}_{n+1}) \mathcal{F}_{W,n+1}^{(g)}(x_{n+1}) &= \sum_{k=1}^n \sum_{j \geq 0} \partial^{(0,j)} \Psi_p(y_{n+1}, y_k) \\ &\quad \cdot \mathcal{F}_{W,n,\alpha}^{(g)}((u(j))_k, \mathbf{x}_n) dy_k^j \end{aligned} \quad (4.20)$$

Here $\partial^{(0,j)}$ is given by

$$\partial^{(i,j)} f(x, y) = \partial_x^{(i)} \partial_y^{(j)} f(x, y),$$

for a function $f(x, y)$, and $\partial^{(0,j)}$ denotes partial derivatives with respect to x and y_j . The forms $\Psi_p(y_{n+1}, y_k) dy_k^j$ given by (6.24), $\Theta_a(x)$ is of (6.26), and $O_a^{W,\alpha}(v_{n+1}; \mathbf{x}_n)$ of (6.27). Similar definitions as in previous subsections can be formulated for a vertex operator algebra module.

4.5. Vertex operator cluster algebras. Finally, we would like to describe the example of vertex operator cluster algebras introduced in [Zu] and reveal their cohomological nature. The condition (2.5) for mutation (4.26) to be involutive gives us a cohomological condition (vanishing square of the n -th coboundary operator) on modular forms. Cluster algebras introduced in [FZ1] have numerous applications in various areas of mathematics [FG1, FG2, FG3, FG4, GSV1, GSV2, GSV3, FST, KS, N1, N2, DFK, GLS, HL, Ke1, Ke2, Na, Sch, N1, N2, Sch]. In this subsection we recall [Zu] definition of a vertex operator cluster algebra and show its cohomological nature. Let us fix a vertex operator operator algebra V . Chose n -marked points $p_i, i = 1, \dots, n$ on a genus g compact Riemann surface formed by one of procedures [Y, TZ] mentioned in Introduction. In the vicinity of each marked point p_i define a local coordinate z_i with zero at p_i . For n -tuples of elements \mathbf{x}_n let us denote

$$\mathcal{Y}(\mathbf{x}_n) = (Y(x_1), \dots, Y(x_n)).$$

Definition 9. We define a vertex operator cluster algebra seed

$$\left(\mathbf{v}_n, \mathcal{Y}(\mathbf{x}_n), \mathcal{F}_{W,n}^{(g)}(\mathbf{x}_n) \right). \quad (4.21)$$

The mutation is defined as follows:

Definition 10. For \mathbf{v}_n , we define the mutation \mathbf{v}'_n of \mathbf{v}_n in the direction $k \in 1, \dots, n$ as

$$\mathbf{v}'_n = \mu_k(x_{n+1}, m) \mathbf{v}_n = ((F_k(v_{n+1}(m)))_k \mathbf{v}_n), \quad (4.22)$$

for some $m \geq 0$, and V -valued functions $F_k^{(g)}(v_{n+1}(m))$, depending on genus g of the Riemann surface. Note that due to the property (5.3) we get a finite number of terms

as a result of the action of $v_{n+1}(m)$ on v_k , $1 \leq k \leq n$. For the n -tuple of vertex operators we define

$$\mathcal{Y}(\mathbf{x}'_n) = \mu_k(x_{n+1}, m) \mathcal{Y}(\mathbf{x}_n) = \left(Y \left(G_k^{(g)}(v_{n+1}(m)) \cdot \mathbf{x}_n \right) \right), \quad (4.23)$$

where $G_k^{(g)}(v_{n+1}(m))$ are other V -valued functions. For $u \in V$, $w \in \mathbb{C}$, the mutation $\mu(x_{n+1})$ of $\mathcal{F}_n^{(g)}(\mathbf{x}_n)$,

$$\mathcal{F}_{W,n}^{(g)'}(\mathbf{x}'_n) = \mu(x_{n+1}) \mathcal{F}_{W,n}^{(g)}(\mathbf{x}_n), \quad (4.24)$$

where $\mu(x_{n+1})$ is given by the coboundary operator

$$\mu(x_{n+1}) = \delta_{(g)}^n,$$

defined by summation over mutations in all possible directions k , $1 \leq k \leq n$, with auxiliary functions $f_1^{(g)}(x_{n+1}, l)$ and $f_2^{(g)}(m, k)$, $1 \leq l \leq l(g)$, $1 \leq k \leq n$, $m \geq 1$,

$$\mathcal{F}_{W,n}^{(g)'}(\mathbf{x}') = \delta_{(g)}^n \mathcal{F}_{W,n}^{(g)}(\mathbf{x}_n). \quad (4.25)$$

We also require the involutivity condition

$$\mu(x_{n+1}) \mu(x_{n+1}) = \text{Id}, \quad (4.26)$$

define the mutation of the seed (4.21) in the direction $k \in 1, \dots, n$ of $x_{n+1} \in V$. We denote by $\mathcal{C}_{(G)}^n$ the space of seeds (4.21) for particular $n \geq 1$, satisfying condition (4.26).

The involutivity condition can be expressed in terms of coboundary operator $\delta_{(g)}^n$ as (2.5) with $v_{n+2} = v_{n+1}$ with extra term Id on the right hand side.

Definition 11. For a fixed g , definitions 9–10 and involutivity condition (4.26) for mutation determine the structure of a vertex operator cluster algebra \mathcal{CG}_n of dimension n . We call the full vertex operator cluster algebra the union $\bigcup_{n \geq 0} \mathcal{CG}_n$.

It is naturally graded by n . of \mathcal{CG}_n . The cohomology of vertex operator cluster algebras will be considered elsewhere. In [Zu] we have proven the following

Proposition 3. For a vertex operator algebra V such that $\dim V_k = 1$, $k \in \mathbb{Z}$, and with $u = \mathbf{1}_V$, $w \in \mathbb{C}$, and

$$F_k(u(m)).v = G_k(u(m)).v = \xi_{u,v} u[-1].v,$$

$$T_k^{(g)}(u(m)) = u[m],$$

for $m \geq 0$, and $\xi_{u,v} \in \mathbb{C}$, $\xi_{u,v}^2 = 1$, depending on u and v , in (4.22), (4.23), and (4.25), the mutation

$$\begin{aligned} \mu &= (\mu_k(\mathbf{1}_V, -1), \mu_k(\mathbf{1}_V, -1), \mu(\mathbf{1}_V, w)), \\ \left(\mathbf{v}'_n, \mathcal{Y}(\mathbf{x}'_n), \mathcal{F}_{W,n}^{(g)'}(\mathbf{x}'_n) \right) &= \mu \left(\mathbf{v}, \mathcal{Y}(\mathbf{x}_n), \mathcal{F}_{W,n}^{(g)}(\mathbf{x}_n) \right), \end{aligned} \quad (4.27)$$

defined by (4.22), (4.23), (4.25) is an involution, i.e.,

$$\mu \mu = \text{Id}.$$

□

5. APPENDIX: VERTEX OPERATOR ALGEBRAS

In this subsection we recall the notion of a vertex operator algebra [B, DL, FHL, FLM, K, LL, MN].

Definition 12. A vertex operator algebra is determined by a quadruple $(V, Y, \mathbf{1}_V, \omega)$, where V is a linear space endowed with a \mathbb{Z} -grading with

$$V = \bigoplus_{r \in \mathbb{Z}} V_r,$$

with $\dim V_r < \infty$. The state $\mathbf{1}_V \in V_0$, $\mathbf{1}_V \neq 0$, is the vacuum vector and $\omega \in V_2$ is the conformal vector with properties described below. The vertex operator Y is a linear map

$$Y : V \rightarrow \text{End}(V)[[z, z^{-1}]],$$

for formal variable z so that for any vector $u \in V$ we have a vertex operator

$$Y(u, z) = \sum_{n \in \mathbb{Z}} u(n) z^{-n-1}. \quad (5.1)$$

The linear operators (modes) $u(n) : V \rightarrow V$ satisfy creativity

$$Y(u, z)\mathbf{1}_V = u + O(z), \quad (5.2)$$

and lower truncation

$$u(n)v = 0, \quad (5.3)$$

conditions for each $u, v \in V$ and $n \gg 0$. Finally, the vertex operators satisfy the Jacobi identity

$$\begin{aligned} z_0^{-1} \delta \left(\frac{z_1 - z_2}{z_0} \right) Y(u, z_1) Y(v, z_2) - z_0^{-1} \delta \left(\frac{z_2 - z_1}{-z_0} \right) Y(v, z_2) Y(u, z_1) \\ = z_2^{-1} \delta \left(\frac{z_1 - z_0}{z_2} \right) Y(Y(u, z_0)v, z_2). \end{aligned} \quad (5.4)$$

These axioms imply locality, skew-symmetry, associativity and commutativity conditions:

$$(z_1 - z_2)^N Y(u, z_1) Y(v, z_2) = (z_1 - z_2)^N Y(v, z_2) Y(u, z_1), \quad (5.5)$$

$$Y(u, z)v = e^{zL(-1)}Y(v, -z)u,$$

$$(z_0 + z_2)^N Y(u, z_0 + z_2) Y(v, z_2) w = (z_0 + z_2)^N Y(Y(u, z_0)v, z_2) w, \quad (5.6)$$

$$u(k)Y(v, z) - Y(v, z)u(k) = \sum_{j \geq 0} \binom{k}{j} Y(u(j)v, z) z^{k-j}, \quad (5.7)$$

for $u, v, w \in V$ and integers $N \gg 0$.

For the conformal vector ω one has

$$Y(\omega, z) = \sum_{n \in \mathbb{Z}} L(n) z^{-n-2}, \quad (5.8)$$

where $L(n)$ satisfies the Virasoro algebra for some central charge C

$$[L(m), L(n)] = (m - n)L(m + n) + \frac{C}{12}(m^3 - m)\delta_{m, -n}\text{Id}_V, \quad (5.9)$$

where Id_V is identity operator on V . Each vertex operator satisfies the translation property

$$\partial_z Y(u, z) = Y(L(-1)u, z). \quad (5.10)$$

The Virasoro operator $L(0)$ provides the \mathbb{Z} -grading with

$$L(0)u = ru,$$

for $u \in V_r$, $r \in \mathbb{Z}$.

For $v = \mathbf{1}_V$ one has

$$Y(\mathbf{1}_V, z) = \text{Id}_V. \quad (5.11)$$

Note also that modes of homogeneous states are graded operators on V , i.e., for $v \in V_k$,

$$v(n) : V_m \rightarrow V_{m+k-n-1}. \quad (5.12)$$

In particular, let us define the zero mode $o(v)$ of a state of weight $\text{wt}(v) = k$, i.e., $v \in V_k$, as

$$o(v) = v(\text{wt}(v) - 1), \quad (5.13)$$

extending to V additively. Similar definition is given for a vertex operator algebra module [K].

Definition 13. Given a vertex operator algebra V , one defines the adjoint vertex operator with respect to $\alpha \in \mathbb{C}$, by

The adjoint operators defined by the mapping $z \mapsto 1/z$, i.e., with $\alpha = \epsilon$,

$$Y_\epsilon^\dagger[v, z] = Y \left[\exp \left(\frac{z}{\epsilon} L[1] \right) \left(-\frac{\epsilon}{z^2} \right)^{L[0]} v, \frac{\epsilon}{z} \right].$$

associated with the formal Möbius map [FHL]

$$z \mapsto \frac{\alpha}{z}.$$

Definition 14. An element $u \in V$ is called quasiprimary if

$$L(1)u = 0.$$

For quasiprimary u of weight $\text{wt}(u)$ one has

$$u^\dagger(n) = (-1)^{\text{wt}(u)} \alpha^{n+1-\text{wt}(u)} u(2\text{wt}(u) - n - 2).$$

Definition 15. A bilinear form

$$\langle \cdot, \cdot \rangle : V \times V \rightarrow \mathbb{C},$$

is called invariant if [FHL, Li]

$$\langle Y(u, z)a, b \rangle = \langle a, Y^\dagger(u, z)b \rangle, \quad (5.14)$$

for all $a, b, u \in V$.

Notice that the adjoint vertex operator $Y^\dagger(.,.)$ as well as the bilinear form $\langle ., . \rangle$, depend on α . In terms of modes, we have

$$\langle u(n)a, b \rangle = \langle a, u^\dagger(n)b \rangle. \quad (5.15)$$

Choosing $u = \omega$, and for $n = 1$ implies

$$\langle L(0)a, b \rangle = \langle a, L(0)b \rangle.$$

Thus,

$$\langle a, b \rangle = 0,$$

when $\text{wt}(a) \neq \text{wt}(b)$.

Definition 16. A vertex operator algebra is called of strong-type if

$$V_0 = \mathbb{C}\mathbf{1}_V,$$

and V is simple and self-dual, i.e., V is isomorphic to the dual module V' as a V -module.

It is proven in [Li] that a strong-type vertex operator algebra V has a unique invariant non-degenerate bilinear form up to normalization. This motivates

Definition 17. The form $\langle ., . \rangle$ on a strong-type vertex operator algebra V is the unique invariant bilinear form $\langle ., . \rangle$ normalized by

$$\langle \mathbf{1}_V, \mathbf{1}_V \rangle = 1.$$

Given a vertex operator algebra $(V, Y(.,.), \mathbf{1}_V, \omega)$, one can find an isomorphic vertex operator algebra $(V, Y[.,.], \mathbf{1}_V, \tilde{\omega})$ called [Zhu] the square-bracket vertex operator algebra. Both algebras have the same underlying vector space V , vacuum vector $\mathbf{1}_V$, and central charge. The vertex operator $Y[.,.]$ is determined by

$$Y[v, z] = \sum_{n \in \mathbb{Z}} v[n]z^{-n-1} = Y\left(q_z^{L(0)}v, q_z - 1\right).$$

The new square-bracket conformal vector is

$$\tilde{\omega} = \omega - \frac{c}{24}\mathbf{1}_V,$$

with the vertex operator

$$Y[\tilde{\omega}, z] = \sum_{n \in \mathbb{Z}} L[n]z^{-n-2}.$$

The square-bracket Virasoro operator mode $L[0]$ provides an alternative \mathbb{Z} -grading on V , i.e., $\text{wt}[v] = k$ if

$$L[0]v = kv,$$

where $\text{wt}[v] = \text{wt}(v)$ for primary v , and $L(n)v = 0$ for all $n > 0$. We can similarly define a square-bracket bilinear form $\langle ., . \rangle_{\text{sq}}$.

6. APPENDIX: GENUS g GENERALIZATIONS OF ELLIPTIC FUNCTIONS

In this Appendix we recall [T2] genus g generalizations of classical elliptic functions.

6.1. Classical elliptic functions. In this subsection we recall the classical elliptic functions and modular forms [Se, La].

Definition 18. For an integer $k \geq 2$, the Eisenstein series is given by

$$E_k(\tau) = E_k(q) = \delta_{n, \text{even}} \left(-\frac{B_k}{k!} + \frac{2}{(k-1)!} \sum_{n \geq 1} \sigma_{k-1}(n) q^n \right),$$

where $\tau \in \mathbb{H}$, $q = e^{2\pi i \tau}$,

$$\sigma_{k-1}(n) = \sum_{d|n} d^{k-1},$$

and k -th Bernoulli number B_k .

If $k \geq 4$ then $E_k(\tau)$ is a modular form of weight k on $\text{SL}(2, \mathbb{Z})$, while $E_2(\tau)$ is a quasi-modular form.

Definition 19. For integer $k \geq 1$, define elliptic functions $z \in \mathbb{C}$:

$$\begin{aligned} P_k(z, \tau) &= \frac{(-1)^{k-1}}{(k-1)!} \partial_z^{k-1} P_1(z, \tau), \\ P_1(z, \tau) &= \frac{1}{z} - \sum_{k \geq 2} E_k(\tau) z^{k-1}. \end{aligned}$$

In particular

$$P_2(z, \tau) = \wp(z, \tau) + E_2(\tau),$$

for Weierstrass function $\wp(z, \tau)$ with periods $2\pi i$ and $2\pi i \tau$. $P_1(z, \tau)$ is related to the quasi-periodic Weierstrass σ -function with

$$P_1(z + 2\pi i \tau, \tau) = P_1(z, \tau) - 1.$$

6.2. Genus two counterparts of Weierstrass functions. In this section we recall the definition of genus two Weierstrass functions [GT]. For $m, n \geq 1$, we first define a number of infinite matrices and row and column vectors:

$$\begin{aligned} \Gamma(m, n) &= \delta_{m, -n+2p-2}, \\ \Delta(m, n) &= \delta_{m, n+2p-2}. \end{aligned} \tag{6.1}$$

We also define the projection matrix

$$\Pi = \Gamma^2 = \begin{bmatrix} \mathbb{I}_{2p-1} & 0 \\ 0 & \ddots \end{bmatrix}, \tag{6.2}$$

where Id_{2p-3} denotes the $2p-3$ dimensional identity matrix and $\text{Id}_{-1} = 0$. Let Λ_a for $a \in \{1, 2\}$ be the matrix with components

$$\begin{aligned} \Lambda_a(m, n) &= \Lambda_a(m, n; \tau_a, \epsilon) \\ &= \epsilon^{(m+n)/2} (-1)^{n+1} \binom{m+n-1}{n} E_{m+n}(\tau_a). \end{aligned} \tag{6.3}$$

Note that

$$\Lambda_a = S A_a S^{-1}, \tag{6.4}$$

for A_a given by

$$A_a = A_a(k, l, \tau_a, \epsilon) = \frac{(-1)^{k+1} \epsilon^{(k+l)/2}}{\sqrt{kl}} \frac{(k+l-1)!}{(k-1)!(l-1)!} E_{k+l}(\tau_a).$$

introduce the infinite dimensional matrices for S a diagonal matrix with components

$$S(m, n) = \sqrt{m} \delta_{mn}. \quad (6.5)$$

Let $\mathbb{R}(x)$ for $x \in \widehat{\Sigma}_a^{(1)}$ be the row vector with components

$$\mathbb{R}(x; m) = \epsilon^{\frac{m}{2}} P_{m+1}(x, \tau_a). \quad (6.6)$$

Let \mathbb{X}_a for $a \in \{1, 2\}$ be the column vector with components

$$\begin{aligned} \mathbb{X}_1(m) &= \mathbb{X}_1 \left(m; v_{n+1}, \mathbf{x}_n; B^{(2)} \right) \\ &= \epsilon^{-m/2} \sum_{u \in V} \mathcal{F}_{V,k}^{(1)}(\mathbf{Y}[\mathbf{x}_k] v[m] u; \tau_1) \mathcal{F}_{V,n-k}^{(1)}(\mathbf{Y}[\mathbf{x}_{k+1,n}] \bar{u}; \tau_2), \\ \mathbb{X}_2(m) &= \mathbb{X}_2 \left(m; v_{n+1}, \mathbf{x}_n; B^{(2)} \right) \\ &= \epsilon^{-m/2} \sum_{u \in V} \mathcal{F}_{V,k}^{(1)}(\mathbf{Y}[\mathbf{x}_k] u; \tau_1) \mathcal{F}_{V,m-k}^{(1)}(\mathbf{Y}[\mathbf{x}_{n-k}] v[m] \bar{u}; \tau_2). \end{aligned} \quad (6.7)$$

Introduce also $\mathbb{Q}(p; x)$ an infinite row vector defined by

$$\mathbb{Q}(p; x) = \mathbb{R}(x) \Delta \left(\mathbb{1} - \tilde{\Lambda}_{\bar{a}} \tilde{\Lambda}_a \right)^{-1}, \quad (6.8)$$

for $x \in \widehat{\Sigma}_a^{(1)}$. Notice that

$$\tilde{\Lambda}_a = \Lambda_a \Delta.$$

On introduces

$$\mathbb{P}_{j+1}(x) = \frac{(-1)^j}{j!} \mathbb{P}_1(x),$$

for $x \in \widehat{\Sigma}_a^{(1)}$, and $j \geq 0$, is the column with components

$$\mathbb{P}_{j+1}(x; m) = \epsilon^{\frac{m}{2}} \binom{m+j-1}{j} (P_{j+m}(x, \tau_a) - \delta_{j0} E_m(\tau_a)). \quad (6.9)$$

Definition 20. One defines

$$\mathcal{P}_1(p; x, y) = \mathcal{P}_1(p; x, y; \tau_1, \tau_2, \epsilon),$$

for $p \geq 1$ by

$$\begin{aligned} \mathcal{P}_1(p; x, y) &= P_1(x - y, \tau_a) - P_1(x, \tau_a) \\ &\quad - \mathbb{Q}(p; x) \tilde{\Lambda}_{\bar{a}} \mathbb{P}_1(y) - (1 - \delta_{p1}) (\mathbb{Q}(p; x) \Lambda_{\bar{a}}) (2p - 2), \end{aligned}$$

for $x, y \in \widehat{\Sigma}_a^{(1)}$, and

$$\begin{aligned} \mathcal{P}_1(p; x, y) &= (-1)^{p+1} \left[\mathbb{Q}(p; x) \mathbb{P}_1(y) + (1 - \delta_{p1}) \epsilon^{p-1} P_{2p-1}(x) \right. \\ &\quad \left. + (1 - \delta_{p1}) \left(\mathbb{Q}(p; x) \tilde{\Lambda}_{\bar{a}} \Lambda_a \right) (2p - 2) \right], \end{aligned}$$

for $x \in \widehat{\Sigma}_a^{(1)}$, $y \in \widehat{\Sigma}_{\bar{a}}^{(1)}$. For $j > 0$, define

$$\mathcal{P}_{j+1}(p; x, y) = \frac{1}{j!} \partial_y^j (\mathcal{P}_1(p; x, y)),$$

i.e.,

$$\mathcal{P}_{j+1}(p; x, y) = \delta_{a, \bar{a}} P_{j+1}(x - y) + (-1)^{j+1} \cdot \mathbb{Q}(p; x) \left(\tilde{\Lambda}_{\bar{a}} \right)^{\delta_{a, \bar{a}}} \mathbb{P}_{j+1}(y), \quad (6.10)$$

for $x \in \widehat{\Sigma}_a^{(1)}$, $y \in \widehat{\Sigma}_{\bar{a}}^{(1)}$. One calls $\mathcal{P}_{j+1}(p; x, y)$ the genus two generalized Weierstrass functions.

6.3. Genus g generalizations of elliptic functions. For purposes of the formula (4.20) we recall here certain definitions [TW]. Define a column vector

$$X = (X_a(m)),$$

indexed by $m \geq 0$ and $a \in \mathcal{I}$ with components

$$X_a(m) = \rho_a^{-\frac{m}{2}} \sum_{b_+} Z^{(0)}(\dots; u(m)b_a, w_a; \dots), \quad (6.11)$$

and a row vector

$$p(x) = (p_a(x, m)),$$

for $m \geq 0, a \in \mathcal{I}$ with components

$$p_a(x, m) = \rho_a^{\frac{m}{2}} \partial^{(0, m)} \psi_p^{(0)}(x, w_a). \quad (6.12)$$

Introduce the column vector

$$G = (G_a(m)),$$

for $m \geq 0, a \in \mathcal{I}$, given by

$$G = \sum_{k=1}^n \sum_{j \geq 0} \partial_k^{(j)} q(y_k) Z_V^{(g)}((u(j))_k \mathbf{x}_n),$$

where $q(y) = (q_a(y; m))$, for $m \geq 0, a \in \mathcal{I}$, is a column vector with components

$$q_a(y; m) = (-1)^p \rho_a^{\frac{m+1}{2}} \partial^{(m, 0)} \psi_p^{(0)}(w_{-a}, y), \quad (6.13)$$

and

$$R = (R_{ab}(m, n)),$$

for $m, n \geq 0$ and $a, b \in \mathcal{I}$ is a doubly indexed matrix with components

$$R_{ab}(m, n) = \begin{cases} (-1)^p \rho_a^{\frac{m+1}{2}} \rho_b^{\frac{n}{2}} \partial^{(m, n)} \psi_p^{(0)}(w_{-a}, w_b), & a \neq -b, \\ (-1)^p \rho_a^{\frac{m+n+1}{2}} \mathcal{E}_m^n(w_{-a}), & a = -b, \end{cases} \quad (6.14)$$

where

$$\mathcal{E}_m^n(y) = \sum_{\ell=0}^{2p-2} \partial^{(m)} f_\ell(y) \partial^{(n)} y^\ell, \quad (6.15)$$

$$\psi_p^{(0)}(x, y) = \frac{1}{x - y} + \sum_{\ell=0}^{2p-2} f_\ell(x) y^\ell, \quad (6.16)$$

for any Laurent series $f_\ell(x)$ for $\ell = 0, \dots, 2p-2$. Define the doubly indexed matrix $\Delta = (\Delta_{ab}(m, n))$ by

$$\Delta_{ab}(m, n) = \delta_{m, n+2p-1} \delta_{ab}. \quad (6.17)$$

Denote by

$$\tilde{R} = R\Delta,$$

and the formal inverse $(I - \tilde{R})^{-1}$ is given by

$$(I - \tilde{R})^{-1} = \sum_{k \geq 0} \tilde{R}^k. \quad (6.18)$$

Define $\chi(x) = (\chi_a(x; \ell))$ and

$$o(u; \mathbf{v}, \mathbf{y}) = (o_a(u; \mathbf{v}, \mathbf{y}; \ell)),$$

are finite row and column vectors indexed by $a \in \mathcal{I}$, $0 \leq \ell \leq 2p-2$ with

$$\chi_a(x; \ell) = \rho_a^{-\frac{\ell}{2}} (p(x) + \tilde{p}(x)(I - \tilde{R})^{-1} R)_a(\ell), \quad (6.19)$$

$$o_a(\ell) = o_a(u; \mathbf{v}, \mathbf{y}; \ell) = \rho_a^{\frac{\ell}{2}} X_a(\ell), \quad (6.20)$$

and where

$$\tilde{p}(x) = p(x)\Delta.$$

$\psi_p(x, y)$ is defined by

$$\psi_p(x, y) = \psi_p^{(0)}(x, y) + \tilde{p}(x)(I - \tilde{R})^{-1} q(y). \quad (6.21)$$

For each $a \in \mathcal{I}_+$ we define a vector

$$\theta_a(x) = (\theta_a(x; \ell)),$$

indexed by $0 \leq \ell \leq 2p-2$ with components

$$\theta_a(x; \ell) = \chi_a(x; \ell) + (-1)^p \rho_a^{p-1-\ell} \chi_{-a}(x; 2p-2-\ell). \quad (6.22)$$

Now define the following vectors of formal differential forms

$$\begin{aligned} P(x) &= p(x) dx^p, \\ Q(y) &= q(y) dy^{1-p}, \end{aligned} \quad (6.23)$$

with

$$\tilde{P}(x) = P(x)\Delta.$$

Then with

$$\Psi_p(x, y) = \psi_p(x, y) dx^p dy^{1-p}, \quad (6.24)$$

we have

$$\Psi_p(x, y) = \Psi_p^{(0)}(x, y) + \tilde{P}(x)(I - \tilde{R})^{-1} Q(y). \quad (6.25)$$

Defining

$$\Theta_a(x; \ell) = \theta_a(x; \ell) dx^p, \quad (6.26)$$

and

$$O_a(u; \mathbf{v}, \mathbf{y}; \ell) = o_a(u; \mathbf{v}, \mathbf{y}; \ell) d\mathbf{y}^{\text{wt}(\mathbf{v})}, \quad (6.27)$$

Remark 9. The $\Theta_a(x)$, and $\Psi_p(x, y)$ coefficients depend on $p = \text{wt}(u)$ but are otherwise independent of the vertex operator algebra V .

REFERENCES

- [A] Ahlfors, L. Some remarks on Teichmüller's space of Riemann surfaces, *Ann.Math.* **74** (1961) 171–191.
- [Ba] Baker, H.F. *Abel's Theorem and the Allied Theory Including the Theory of Theta Functions*, Cambridge University Press (Cambridge, 1995).
- [BPZ] Belavin, A., Polyakov, A. and Zamolodchikov, A.: Infinite conformal symmetry in two-dimensional quantum field theory. *Nucl. Phys.* **B241** 333–380 (1984).
- [Be1] Bers, L. Inequalities for finitely generated Kleinian groups, *J.Anal.Math.* **18** 23–41 (1967).
- [Be2] Bers, L. Automorphic forms for Schottky groups, *Adv. Math.* **16** 332–361 (1975).
- [BKT] Bringmann, K., Krauel, M., Tuite, M. Zhu reduction for Jacobi n -point functions and applications. *Trans. Amer. Math. Soc.* 373 (2020), no. 5, 3261–3293.
- [Bo] Bobenko, A. Introduction to compact Riemann surfaces, in *Computational Approach to Riemann Surfaces*, edited Bobenko, A. and Klein, C., Springer-Verlag (Berlin, Heidelberg, 2011).
- [B] Borcherds, R.E.: Vertex algebras, Kac-Moody algebras and the monster. *Proc. Nat. Acad. Sci.* **83**, 3068–3071 (1986).
- [Bu] Burnside, W. On a class of automorphic functions, *Proc. L. Math.Soc.* **23** 49–88 (1891).
- [BS] R. Bott, G.Segal, The cohomology of the vector fields on a manifold, *Topology* Volume 16, Issue 4, 1977, Pages 285–298.
- [BZF] D. Ben-Zvi and E. Frenkel *Vertex algebras on algebraic curves*. American Mathematical Society, 2 edition, 2004.
- [Co] Codogni, G. Vertex algebras and Teichmüller modular forms, arXiv:1901.03079.
- [DFK] Ph. Di Francesco and R. Kedem, Q-systems as cluster algebras. II. *Lett. Math. Phys.* **89**, no. 3, (2009) 183.
- [DGM] L. Dolan, P. Goddard and P. Montague, Conformal field theories, representations and lattice constructions, *Commun. Math. Phys.* **179** (1996) p. 61.
- [DL] C. Dong and J. Lepowsky, *Generalized Vertex Algebras and Relative Vertex Operators* (Progress in Math. Vol. **112**, Birkhäuser, Boston, 1993).
- [DFK] Ph. Di Francesco and R. Kedem, Q-systems as cluster algebras. II. *Lett. Math. Phys.* **89**, no. 3, (2009) 183.
- [EO] Eguchi, T. and Ooguri, H. Conformal and current algebras on a general Riemann surface, *Nucl. Phys.* **B282** 308–328 (1987).
- [Fa] Fay, J.D. *Theta Functions on Riemann Surfaces*, Lecture Notes in Mathematics, Vol. 352. Springer-Verlag, (Berlin-New York, 1973).
- [FHL] I. Frenkel, Y.-Z. Huang and J. Lepowsky, *On Axiomatic Approaches to Vertex Operator Algebras and Modules*, *Mem. AMS.* **104** No. 494 (1993).
- [FLM] I. Frenkel, J. Lepowsky and A. Meurman, *Vertex Operator Algebras and the Monster, Pure and Appl. Math.* Vol. **134** (Academic Press, Boston, 1988).
- [FMS] Ph. Di Francesco, P., Mathieu, P. and Senechal, D.: *Conformal Field Theory*. Springer Graduate Texts in Contemporary Physics, Springer-Verlag, New York (1997).
- [FK] H.M. Farkas and I. Kra, *Riemann surfaces*, (Springer-Verlag, New York, 1980).
- [FST] S. Fomin, M. Shapiro, D. Thurston, Cluster algebras and *Acta Mathematica* **201** (2008) 83.
- [FZ1] S. Fomin, A. Zelevinsky, Cluster algebras. I. Foundations, *J. of the Amer. Math. Soc.* (2002) **15** (2) 49. 7–529 .
- [FZ2] S. Fomin, A. Zelevinsky. Cluster algebras. II. Finite type classification, *Inventiones Mathematicae* **154** (1): (2003) 63–121 (2003).
- [FZ3] S. Fomin, A. Zelevinsky. Cluster algebras. IV. Coefficients, *Compositio Mathematica* **143** (1): (2007) p. 112–164 (2007).

- [FG1] V. Fock and A. Goncharov, Moduli spaces of local systems and higher Teichmüller theory. *Publ. Math. Inst. Hautes Études Sci.* (2006) **103**, 1.
- [FG2] V. Fock and A. Goncharov, Cluster ensembles, quantization and the dilogarithm, *Ann. Sci. Éc. Norm. Supér.* (4) **42**, no. 6, (2009) 865.
- [FG3] V. Fock and A. Goncharov, Cluster ensembles, quantization and the dilogarithm. II, in *The intertwiner. Algebra, arithmetic, and geometry: in honor of Yu. I. Manin*. Vol. I, (Progr. Math., 269, Birkhäuser Boston, Inc., Boston, MA, 2009) p. 655.
- [FG4] V. Fock and A. Goncharov, Dual Teichmüller and lamination spaces, in *Handbook of Teichmüller theory*. Vol. I, IRMA Lect. Math. Theor. Phys., 11 (Eur. Math. Soc., Zürich, 2007), p. 647.
- [FS] Freidan, D. and Shenker, S.: The analytic geometry of two-dimensional conformal field theory. *Nucl. Phys.* **B281** 509–545 (1987).
- [Fo] Ford, L.R. *Automorphic Functions*, AMS-Chelsea, (Providence, 2004).
- [Fei] Feigin, B. L.: Conformal field theory and Cohomologies of the Lie algebra of holomorphic vector fields on a complex curve. *Proc. ICM, Kyoto, Japan*, 71–85 (1990)
- [Fuks] Fuks, D. B.: Cohomology of Infinite Dimensional Lie algebras. New York and London: Consultant Bureau 1986
- [FST] S. Fomin, M. Shapiro, D. Thurston, Cluster algebras and *Acta Mathematica* **201** (2008) 83.
- [GSV1] M. Gekhtman, M. Shapiro and A. Vainshtein, *Mosc. Math. J.* **3**, no. 3, (2003) 899.
- [GSV2] M. Gekhtman, M. Shapiro and A. Vainshtein, *Duke Math. J.* **127**, no. 2, (2005) 291.,
- [GSV3] M. Gekhtman, M. Shapiro, and A. Vainshtein. Cluster algebras and Poisson geometry, in *Mathematical Surveys and Monographs*, vol. 167 (American Mathematical Society, Providence, RI, 2010).
- [GT] T. Gilroy and M. Tuite, Genus Two Zhu Theory for Vertex Operator Algebras, arXiv:1511.07664.
- [Gu] R. C. Gunning. Lectures on Vector Bundles over Riemann Surfaces. (MN-6), Volume 6, Mathematical Notes, 105, Princeton University Press, 1968
- [GLS] Ch. Geiss, B. Leclerc and J. Schröer. Cluster structures on *Selecta Math.* (N.S.) **19**, no. 2, (2013) 337.
- [HL] D. Hernandez and B. Leclerc. Cluster algebras and quantum affine algebras, *Duke Math. J.* **154**, no. 2, (2010) 265.
- [Huang] Y.-Zh. Huang, *Two-dimensional conformal geometry and vertex operator algebras*, Progress in Mathematics, Vol. 148, Birkhäuser, Boston, 1997.
- [Huang1] Y.-Zh. Huang, A cohomology theory of grading-restricted vertex algebras, *Comm. Math. Phys.* **327** (2014), no. 1, 279–307.
- [Huang2] Y.-Zh. Huang, The first and second cohomologies of grading-restricted vertex algebras, *Communications in Mathematical Physics* volume 327, 261–278 (2014)
- [K] V. Kac, *Vertex Algebras for Beginners, Second Ed.* (Univ. Lect. Ser. **10**, AMS, 1998).
- [Ke1] B. Keller, Cluster algebras, quiver representations and triangulated categories, “Triangulated categories”, 76–160, in *London Math. Soc. Lecture Note Ser.*, (Cambridge Univ. Press, Cambridge, 2010) p. 375.
- [Ke2] B. Keller, Quantum loop algebras, quiver varieties, and cluster algebras, in *Representations of algebras and related topics*, (EMS Ser. Congr. Rep., Eur. Math. Soc., Zürich, 2011) p. 117.
- [KS] M. Kontsevich and Y. Soibelman, Stability structures, Donaldson- Thomas invariants and cluster transformations, arXiv:0811.2435, 2008.
- [Kn] Knizhnik, V.G.: Multiloop amplitudes in the theory of quantum strings and complex geometry. *Sov. Phys. Usp.* **32** 945–971 (1989).
- [Kaw] Kawazumi, N. On the complex analytic Gel’fand-Fuks cohomology of open Riemann surfaces. *Ann. Inst. Fourier (Grenoble)* **43** (1993), no. 3, 655–712.
- [KS] M. Kontsevich and Y. Soibelman, Stability structures, Donaldson- Thomas invariants and cluster transformations, arXiv:0811.2435, 2008.
- [KZ] Knizhnik, V.G.; Zamolodchikov, A.B. (1984), Current Algebra and Wess-Zumino Model in Two-Dimensions, *Nucl. Phys. B*, **247** (1): 83–103.

- [LL] J. Lepowsky and H. Li, *Introduction to Vertex Algebras*, (Progress in Math. Vol. **227**, Birkhäuser, Boston, 2004).
- [L] S. Lang, *Introduction to Modular Forms* (Springer, Berlin, 1976).
- [La] Lang, S.: Elliptic functions. Springer-Verlag, New York (1987).
- [Li] Li, H.: Symmetric invariant bilinear forms on vertex operator algebras. *J. Pure. Appl. Alg.* **96**, 279–297 (1994).
- [MT] G. Mason and M.P. Tuite, "Vertex operators and modular forms" in *A window into zeta and modular physics*, (Math. Sci. Res. Inst. Publ., 57, Cambridge Univ. Press, Cambridge, 2010) p. 183.
- [MT1] G. Mason and M.P. Tuite, Torus Chiral n -point Functions for Free Boson and Lattice Vertex Operator Algebras, *Commun. Math. Phys.* **235** (2003), 47–68 (2003).
- [MT2] G. Mason, and M.P. Tuite, On genus two Riemann surfaces formed from sewn tori, *Commun. Math. Phys.* **270** (2007) p. 587–634 (2007).
- [MT3] G. Mason and M.P. Tuite, Free bosonic vertex operator algebras on genus two Riemann surfaces I, *Commun. Math. Phys.* **300** (2010) p. 673–713 (2010).
- [MT31] Mason, G. and Tuite, M.P. Free bosonic vertex operator algebras on genus two Riemann surfaces II, in *Conformal Field Theory, Automorphic Forms and Related Topics*, Contributions in Mathematical and Computational Sciences **8** 183–225, Springer Verlag, (Berlin, Heidelberg, 2014).
- [MTZ] G. Mason, M. P. Tuite, A. Zuevsky, Torus n -point functions for R -graded vertex operator superalgebras and continuous fermion orbifolds. *Commun. Math. Phys.* 283 no. 2 (2008) p. 305–342 (2008).
- [McI] McIntyre, A. Analytic torsion and Faddeev-Popov ghosts, SUNY PhD thesis 2002, hdl.handle.net/11209/10688.
- [McIT] McIntyre, A. and Takhtajan, L.A. Holomorphic factorization of determinants of Laplacians on Riemann surfaces and a higher genus generalization of Kronecker's first limit formula, *GAFA, Geom.Funct.Anal.* **16** 1291–1323 (2006).
- [MN] A. Matsuo and K. Nagatomo, *Axioms for a Vertex Algebra and the Locality of Quantum Fields*, (Math. Soc. of Japan Memoirs Vol. **4**, Tokyo, 1999).
- [Miy1] M. Miyamoto, Modular invariance of vertex operator algebras satisfying C_2 -cofiniteness, *Duke Math. J. Vol.* **122** No. 1 (2004) p. 51.
- [Mu] Mumford, D. *Tata Lectures on Theta I and II*, Birkhäuser, (Boston, 1983).
- [N1] T. Nakanishi, Dilogarithm identities for conformal field theories and cluster algebras: simply laced case, *Nagoya Math. J.* **202**, (2011) 23–43 .
- [N2] T. Nakanishi, Periodicities in cluster algebras and dilogarithm identities, in *Representations of algebras and related topics*, (EMS Ser. Congr. Rep., Eur. Math. Soc., Zürich, 2011) p. 407.
- [Na] H. Nakajima, Quiver varieties and cluster algebras, *Kyoto J. Math.* **51** no. 1, (2011) p. 71–126 (2011).
- [O] Odesskii, A. Deformations of complex structures on Riemann surfaces and integrable structures of Whitham type hierarchies, [arXiv:1505.07779](https://arxiv.org/abs/1505.07779).
- [PT] Patras, F., Thomas, J.-C. Cochain algebras of mapping spaces and finite group actions. *Topology Appl.* 128 (2003), no. 2-3, 189–207.
- [Po] B. Ponsot Monodromy of solutions of the Knizhnik-Zamolodchikov equation: $SL(2, \mathbb{C})/SU(2)$ WZNW model, *Nuclear Physics B* Volume 642, Issues 1–2, 21 October 2002, Pages 114–138.
- [Sch] R. Schiffler, On cluster algebras arising from unpunctured surfaces. II. *Adv. Math.* **223** no. 6 (2010) 1885.
- [Sav] M. V. Saveliev, Integro-differential nonlinear equations and continual Lie algebras. *Comm. Math. Phys.* 121 (1989), no. 2, 283–290.
- [SV1] M. V. Saveliev, A. M. Vershik. Continuum analogues of contragredient Lie algebras. *Commun. Math. Phys.* 126, 367, 1989;
- [SV2] M. V. Saveliev, A. M. Vershik. New examples of continuum graded Lie algebras. *Phys. Lett. A*, 143, 121, 1990.
- [Se] J.-P. Serre, *A Course in Arithmetic* (Springer, New York, 1973).

- [TUY] A. Tsuchiya, K. Ueno and Y. Yamada, Conformal field theory on universal family of stable curves with gauge symmetries, in: *Advanced Studies in Pure Math.*, Vol. 19, Kinokuniya Company Ltd., Tokyo, 1989, 459–566.
- [TK] Tsuchiya, A.; Kanie, Y. (1988), Vertex operators in conformal field theory on $P(1)$ and monodromy representations of braid group, *Adv. Stud. Pure Math.*, 16, pp. 297–372 (Erratum in volume 19, pp. 675–682.)
- [T2] Tuite, M.P. The Heisenberg generalized vertex operator algebra on a Riemann surface. Lie groups, number theory, and vertex algebras, 321–342, *Contemp. Math.*, 768, Amer. Math. Soc., [Providence], RI, [2021]
- [TZ] Tuite, M.P. and Zuevsky, A. The bosonic vertex operator algebra on a genus g Riemann surface, *RIMS Kokyuroko* **1756** 81–93 (2011).
- [TZ1] Tuite, M.P. and Zuevsky, A.: The Szego kernel on a sewn Riemann surface. *Commun. Math. Phys.* **306**, 617–645 (2011).
- [TZ2] Tuite, M.P. and Zuevsky, A.: Genus two partition and correlation functions for fermionic vertex operator superalgebras I. *Commun. Math. Phys.* 306 419–447 (2011).
- [TZ3] Tuite, M.P. and Zuevsky, A.: Genus two partition and correlation functions for fermionic vertex operator superalgebras II. arXiv:1308.2441.
- [TW] Tuite, M.P., M. Welby. General Genus Zhu Recursion for Vertex Operator Algebras. arXiv:1911.06596
- [Vershik] Vershik, A. Lie algebras generated by dynamical systems. *Algebra i Analiz* 4 (1992), no. 6, 103–113; reprinted in *St. Petersburg Math. J.* 4 (1993), no. 6, 1143–1151.
- [Wag] Wagemann, F.: Differential graded cohomology and Lie algebras of holomorphic vector fields. *Comm. Math. Phys.* 208 (1999), no. 2, 521–540
- [Y] Yamada, A.: Precise variational formulas for abelian differentials. *Kodai. Math. J.* **3**, 114–143 (1980).
- [Zhu] Y. Zhu, Modular-invariance of characters of vertex operator algebras, *J. Amer. Math. Soc.* **9** No. 1, (1996) P. 237–302 (1996).
- [Zu] A. Zuevsky, Cluster algebras based on vertex operator algebras. *Internat. J. Modern Phys. B* 30 (2016), no. 28–29, 1640030, 13 pp. 17B69 (13F60)

INSTITUTE OF MATHEMATICS, CZECH ACADEMY OF SCIENCES, ŽITNA 25, PRAGUE, CZECH REPUBLIC

Email address: zuevsky@yahoo.com