

# The action of GT-shadows on child's drawings

Vasily A. Dolgushev

*To the memory of Andrey Loginov*

## Abstract

GT-shadows [7] are tantalizing objects that can be thought of as approximations of elements of the mysterious Grothendieck-Teichmüller group  $\widehat{\mathrm{GT}}$  introduced by V. Drinfeld in 1990. GT-shadows form a groupoid  $\mathrm{GTSh}$  whose objects are finite index subgroups of the pure braid group  $\mathrm{PB}_4$ , that are normal in  $\mathrm{B}_4$ . The goal of this paper is to describe the action of GT-shadows on Grothendieck's child's drawings and show that this action agrees with that of  $\widehat{\mathrm{GT}}$ . We discuss the hierarchy of orbits of child's drawings with respect to the actions of  $\mathrm{GTSh}$ ,  $\widehat{\mathrm{GT}}$ , and the absolute Galois group  $G_{\mathbb{Q}}$  of rationals. We prove that the monodromy group and the passport of a child's drawing are invariant with respect to the action of the subgroupoid  $\mathrm{GTSh}^{\heartsuit}$  of charming GT-shadows. We use the action of GT-shadows on child's drawings to prove that every Abelian child's drawing admits a Belyi pair defined over  $\mathbb{Q}$ . Finally, we describe selected examples of non-Abelian child's drawings.

## Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
1.1	Notational conventions . . . . .	5
<b>2</b>	<b>Background material</b>	<b>6</b>
2.1	Reminder of GT-shadows . . . . .	6
2.1.1	$\widehat{\mathrm{GT}}$ versus the groupoid $\mathrm{GTSh}$ . . . . .	9
2.2	Permutation pairs, permutation triples and child's drawings . . . . .	11
2.3	Representing child's drawings by group homomorphisms . . . . .	12
2.4	The action of $\widehat{\mathrm{GT}}$ on child's drawings and the functor $\mathcal{A} : \widehat{\mathrm{GT}}_{\mathrm{NFI}} \rightarrow \mathrm{Dessin}$ . .	14
<b>3</b>	<b>The action of GT-shadows on child's drawings</b>	<b>17</b>
3.1	The hierarchy of orbits . . . . .	21
3.2	The monodromy group and the passport are invariant with respect to the action of GT-shadows . . . . .	23
<b>4</b>	<b>Abelian child's drawings</b>	<b>25</b>
4.1	$\mathrm{GTSh}$ -orbit of an Abelian child's drawing is a singleton . . . . .	27
<b>5</b>	<b>Examples of <math>\mathrm{GTSh}^{\heartsuit}</math>-orbits for (non-Abelian) child's drawings</b>	<b>28</b>
5.1	An example of degree 6, genus 0 and orbit size 2 . . . . .	28
5.2	An example of degree 5, genus 0 and orbit size 2 . . . . .	29
5.3	Examples of child's drawing subordinate to $\mathrm{N}^{(5)}$ and $\mathrm{N}^{(29)}$ from list (3.19) . .	30
<b>A</b>	<b>Charming GT-shadows satisfy the simplified hexagon relations</b>	<b>32</b>

# 1 Introduction

The profinite version  $\widehat{\mathbf{GT}}$  of the Grothendieck-Teichmueller group [5, Section 4] [26], [28] connects the topology to number theory in a fascinating way.  $\widehat{\mathbf{GT}}$  receives an injective homomorphism [20], [28, Section 3.2] from the absolute Galois group  $G_{\mathbb{Q}}$  of rationals. It acts on Grothendieck's child's drawings and this action is compatible with the embedding  $G_{\mathbb{Q}} \hookrightarrow \widehat{\mathbf{GT}}$  and the standard action of  $G_{\mathbb{Q}}$ .

Just as  $G_{\mathbb{Q}}$ , the group  $\widehat{\mathbf{GT}}$  is a rather intractable object. For example, currently, we know explicitly only two elements of  $\widehat{\mathbf{GT}}$ : the identity element and the element that comes from the complex conjugation. The author also believes that, for an arbitrary child's drawing  $\mathcal{D}$ , tools of modern mathematics do not allow us to say much about the orbit  $\widehat{\mathbf{GT}}(\mathcal{D})$ .

Let us denote by  $B_n$  (resp.  $PB_n$ ) the Artin braid group (resp. the pure braid group) on  $n$  strands. We denote by  $x_{ij}$ ,  $1 \leq i < j \leq n$  the standard generators of  $PB_n$  and recall that the elements  $x_{12}, x_{23}$  generate a free subgroup of  $PB_3$ . In this paper, we tacitly identify the free group  $F_2$  on two generators with  $\langle x_{12}, x_{23} \rangle \leq PB_3$ .

$\widehat{\mathbf{GT}}$  can be defined as the group of continuous automorphisms of the profinite completion  $\widehat{\mathbf{PaB}}$  of the operad  $\mathbf{PaB}$  of parenthesized braids [3], [7, Appendix A], [11, Chapter 6], [33].  $\mathbf{PaB}$  is an operad in the category of groupoids and it is “assembled” from the braid groups  $(B_n)_{n \geq 2}$ . Objects of the groupoid  $\mathbf{PaB}(n)$  are completely parenthesized sequences of number  $1, 2, \dots, n$  in which each number appears exactly once. For example,  $\text{Ob}(\mathbf{PaB}(3))$  has 12 objects:  $(1, 2)3$ ,  $1(2, 3)$ ,  $(2, 1)3$ ,  $2(1, 3)$ ,  $\dots$

The isomorphisms  $\alpha \in \mathbf{PaB}((1, 2)3, 1(2, 3))$  and  $\beta \in \mathbf{PaB}((1, 2), (2, 1))$  shown in figure 1.1 play an important role for  $\mathbf{PaB}$ .




Fig. 1.1: The isomorphisms  $\alpha$  and  $\beta$

Due to [11, Theorem 6.2.4],  $\mathbf{PaB}$  is generated by  $\alpha$  and  $\beta$  (as the operad in the category of groupoids) and any relation involving  $\alpha$  and  $\beta$  is a consequence of the two hexagon relations and the pentagon relation (see (A.13), (A.14) and (A.15) in [7, Appendix A]).

Since  $\alpha$  and  $\beta$  are topological generators of  $\widehat{\mathbf{PaB}}$ , every  $\hat{T} \in \widehat{\mathbf{GT}}$  is uniquely determined by the values  $\hat{T}(\alpha) \in \widehat{\mathbf{PaB}}((1, 2)3, 1(2, 3))$  and  $\hat{T}(\beta) \in \widehat{\mathbf{PaB}}((1, 2), (2, 1))$ . In addition, since the automorphism group of  $(1, 2)3$  (resp.  $(1, 2)$ ) in  $\widehat{\mathbf{PaB}}$  is  $\widehat{PB}_3$  (resp.  $\widehat{PB}_2 \cong \widehat{\mathbb{Z}}$ ), every  $\hat{T} \in \widehat{\mathbf{GT}}$  is uniquely determined by a pair  $(\hat{m}, \hat{f}) \in \widehat{\mathbb{Z}} \times \widehat{PB}_3$ . Using the hexagon relations and the pentagon relation, one can show that  $\hat{f} \in \hat{F}_2$ . In fact, one can show that  $\hat{f}$  belongs to the topological closure of the commutator subgroup  $[\hat{F}_2, \hat{F}_2]$ .

It is easy to see that the action of  $\widehat{\mathbf{GT}}$  on  $\widehat{PB}_3 \cong \text{Aut}((1, 2)3)$  descends to the action on  $\hat{F}_2 \leq \widehat{PB}_3$  and it is given (on the topological generators of  $\hat{F}_2$ ) by the formulas

$$\hat{T}_{\hat{F}_2}(x) := x^{2\hat{m}+1}, \quad \hat{T}_{\hat{F}_2}(y) := \hat{f}^{-1}y^{2\hat{m}+1}\hat{f}, \quad (1.1)$$

where  $(\hat{m}, \hat{f})$  is the pair in  $\widehat{\mathbb{Z}} \times \widehat{\mathbb{F}}_2$  corresponding to  $\hat{T}$ .

Let  $S_d$  be the symmetric group of degree  $d$  and  $\text{Hom}_{\text{tran}}(\widehat{\mathbb{F}}_2, S_d)$  be the set of continuous group homomorphisms  $\psi : \widehat{\mathbb{F}}_2 \rightarrow S_d$  for which the subgroup  $\psi(\widehat{\mathbb{F}}_2)$  acts transitively on the set  $\{1, 2, \dots, d\}$ . Here  $S_d$  is considered with the discrete topology.

The set  $\text{Hom}_{\text{tran}}(\widehat{\mathbb{F}}_2, S_d)$  carries the obvious action of  $S_d$  by conjugation and a **child's drawing** of degree  $d$  can be defined as an orbit of the  $S_d$ -action on  $\text{Hom}_{\text{tran}}(\widehat{\mathbb{F}}_2, S_d)$ . In these terms, it is easy to define the (right) action of  $\widehat{\text{GT}}$  on child's drawings: given a child's drawing  $[\psi]$  represented by a continuous group homomorphism  $\psi : \widehat{\mathbb{F}}_2 \rightarrow S_d$  and  $\hat{T} \in \widehat{\text{GT}}$  the child's drawing  $[\psi]^{\hat{T}}$  is represented by the continuous group homomorphism

$$\psi \circ \hat{T}_{\widehat{\mathbb{F}}_2} : \widehat{\mathbb{F}}_2 \rightarrow S_d.$$

Let us denote by  $\text{NFI}_{\text{PB}_4}(\text{B}_4)$  the poset of finite index normal subgroups  $\text{N} \trianglelefteq \text{B}_4$  satisfying the condition  $\text{N} \leq \text{PB}_4$ . In paper [7], the authors introduced the concept of a **GT-shadow**. Loosely speaking, a GT-shadow is an onto morphism of (truncated) operads

$$\text{PaB}^{\leq 4} \rightarrow \text{PaB}^{\leq 4} / \sim_{\text{N}},$$

where  $\sim_{\text{N}}$  is an equivalence relation on the set of morphisms of  $\text{PaB}^{\leq 4}$  that comes from an element  $\text{N}$  of the poset  $\text{NFI}_{\text{PB}_4}(\text{B}_4)$ .

In paper [7], it was proved that GT-shadows from a groupoid  $\text{GTSh}$  whose objects are elements of  $\text{NFI}_{\text{PB}_4}(\text{B}_4)$ . For every  $\text{K}, \text{N} \in \text{NFI}_{\text{PB}_4}(\text{B}_4)$ , the set  $\text{GTSh}(\text{K}, \text{N})$  may be identified with the set of isomorphisms of (truncated) operads

$$\text{PaB}^{\leq 4} / \sim_{\text{K}} \xrightarrow{\cong} \text{PaB}^{\leq 4} / \sim_{\text{N}}.$$

In this paper, we use the poset  $\text{NFI}_{\text{PB}_4}(\text{B}_4)$  and the set of all child's drawings to form a category **Dessin**. We show that  $\widehat{\text{GT}}$  acts naturally on the poset  $\text{NFI}_{\text{PB}_4}(\text{B}_4)$  and the action of  $\widehat{\text{GT}}$  on child's drawings gives us a cofunctor

$$\mathcal{A} : \widehat{\text{GT}}_{\text{NFI}} \rightarrow \text{Dessin}$$

from the corresponding transformation groupoid  $\widehat{\text{GT}}_{\text{NFI}}$  to the category **Dessin**.

We show (see Theorem 3.1) that GT-shadows act on child's drawings in the sense that we have a natural cofunctor

$$\mathcal{A}^{sh} : \text{GTSh} \rightarrow \text{Dessin}. \quad (1.2)$$

Recall [7, Section 2.4] that, for every  $\hat{T} \in \widehat{\text{GT}}$  and  $\text{N} \in \text{NFI}_{\text{PB}_4}(\text{B}_4)$ , we can produce a GT-shadow  $T_{\text{N}}$  with the target  $\text{N}$ , and  $T_{\text{N}}$  may be viewed as an approximation of  $\hat{T}$ . Using this passage from elements of  $\widehat{\text{GT}}$  to GT-shadows we define a functor

$$\mathcal{PR} : \widehat{\text{GT}}_{\text{NFI}} \rightarrow \text{GTSh}. \quad (1.3)$$

We prove (see Theorem 3.2) that the action of GT-shadows on child's drawings is compatible with the action of  $\widehat{\text{GT}}$  in the sense that the functors

$$\mathcal{A} \quad \text{and} \quad \mathcal{A}^{sh} \circ \mathcal{PR}$$

are equal in the strict sense.

Using Theorem 3.2 and the compatibility of the  $G_{\mathbb{Q}}$ -action and the  $\widehat{\text{GT}}$ -action on child's drawings, we deduce several interesting corollaries:

- Corollary 3.9 describes the hierarchy of orbits of the  $\text{GTSh}$ -action,  $\widehat{\text{GT}}$ -action and  $G_{\mathbb{Q}}$ -action on the set of child's drawings.
- Corollary 3.12 may be viewed as a version of [17, Proposition 14]; this statement gives us a useful bound on the degree of the field of moduli of a child's drawing.
- Corollary 3.13 tells us that, using the poset  $\text{NFI}_{\text{PB}_4}(\text{B}_4)$ , we can produce many examples of Galois child's drawings that admit Belyi pairs defined over  $\mathbb{Q}$ .

Using consequences of the hexagon relations for  $\text{GT}$ -shadows, we show (see Theorem 3.16) that the passport of a child's drawing is invariant with respect to the action of (charming)  $\text{GT}$ -shadows.

In this paper, we also prove some basic facts about Abelian child's drawings and show (see Corollary 4.8) that every Abelian child's drawing admits a Belyi pair defined over  $\mathbb{Q}$ .

Finally, we describe selected examples of non-Abelian child's drawings whose  $\text{GTSh}$ -orbits were computed using software package [6]. Whenever possible, the results were compared to  $G_{\mathbb{Q}}$ -orbits from database [24]. It is amazing to see that, if a  $\text{GTSh}$ -orbit and the  $G_{\mathbb{Q}}$ -orbit of a child's drawing can be computed then these orbits coincide.

## Organization of the paper

Section 2 is devoted to the background material. It contains a brief reminder of the groupoid  $\text{GTSh}$  and its link to  $\widehat{\text{GT}}$ . In this section, we also recall child's drawings, introduce the category **Dessin** and define the action of  $\widehat{\text{GT}}$  on child's drawings in terms of a cofunctor from a certain transformation groupoid  $\widehat{\text{GT}}_{\text{NFI}}$  to **Dessin**.

In Section 3, we define the action of  $\text{GT}$ -shadows on child's drawings, describe the relationship between orbits of a child's drawing with respect to different actions, and prove that the monodromy group and the passport of a child's drawing are invariant with respect to the action of (charming)  $\text{GT}$ -shadows.

Section 4 is devoted to various properties of Abelian child's drawings. In this section, we prove that every Abelian child's drawing admits a Belyi pair defined over  $\mathbb{Q}$ .

In Section 5, we describe selected examples of non-Abelian child's drawings whose  $\text{GTSh}$ -orbits were computed.

In Appendix A, we prove that charming  $\text{GT}$ -shadows satisfy versions of relations (4.3) and (4.4) from [5, Section 4].

**Remark 1.1** We should remark that the notation  $\widehat{\text{GT}}$  is a bit misleading: the hat over “ $\text{GT}$ ” does not mean that  $\widehat{\text{GT}}$  is a profinite completion of some “well known group  $\text{GT}$ ”. However, please see [7, Theorem 3.8].

**Acknowledgement.** V.A.D. is thankful to John Voight for many illuminating discussions and for his patience with his (possibly naive) questions. V.A.D. is thankful to Leila Schneps for her unbounded enthusiasm about  $\text{GT}$ -shadows and everything related to  $\text{GT}$ -shadows. V.A.D. is thankful to Benjamin Enriquez for his suggestion to use the transformation groupoid corresponding to the action of  $\widehat{\text{GT}}$  on  $\text{NFI}_{\text{PB}_4}(\text{B}_4)$ . V.A.D. benefitted from discussions with David Harbater, Julia Hartmann and Florian Pop. V.A.D. is thankful to

Khanh Le, Aidan Lorenz, Chelsea Zackey for discussion and their contributions to the software package GT. V.A.D. is thankful to Pavol Severa for showing him the works of Pierre Guillot. V.A.D. acknowledges Temple University for 2021 Summer Research Award.

**Memorial note.** Andrey Loginov (1977-2016) was an experimental physicist, and his work helped to push frontiers of knowledge related to the fundamental questions about elementary particles. There were several moments in my life when he played for me a role of an older brother. Life connected us in bizarre ways by cities scattered around the globe: Berdychiv, Chicago, Dolgoprudny, Moscow and Tomsk. It is very sad that I did not have a chance to say a proper goodbye to him...

## 1.1 Notational conventions

For a set  $X$  with an equivalence relation and  $a \in X$  we will denote by  $[a]$  the equivalence class represented by the element  $a$ .

The notation  $B_n$  (resp.  $PB_n$ ) is reserved for the Artin braid group on  $n$  strands (resp. the pure braid group on  $n$  strands). The standard generators of  $B_n$  are denoted by  $\sigma_1, \dots, \sigma_{n-1}$  and the standard generators of  $PB_n$  are denoted by  $x_{ij}$  for  $1 \leq i < j \leq n$ . We set

$$c := x_{23}x_{12}x_{13} \in PB_3$$

and recall that  $c$  generates the center  $\mathcal{Z}(PB_3)$  of  $PB_3$  (as well as the center  $\mathcal{Z}(B_3)$  of  $B_3$ ). The free group  $F_2$  on two generators is tacitly identified with the subgroup  $\langle x_{12}, x_{23} \rangle \leq PB_3$ . Occasionally, we denote the standard generators of  $F_2$  by  $x$  and  $y$ , i.e.  $x := x_{12}$  and  $y := x_{23}$ .

$S_d$  denotes the group of bijections  $\{1, 2, \dots, d\} \xrightarrow{\cong} \{1, 2, \dots, d\}$  (i.e. the symmetric group of degree  $d$ ). A subgroup  $H \leq S_d$  is called **transitive** if its standard action on  $\{1, 2, \dots, d\}$  is transitive. Every finite group is tacitly considered with the discrete topology. For a group  $G$ , the notation  $[G, G]$  is reserved for the commutator subgroup of  $G$ . For a normal subgroup  $H \trianglelefteq G$  of finite index, we denote by  $\mathbf{NFI}_H(G)$  the poset of finite index normal subgroups  $N$  in  $G$  such that  $N \leq H$ . The notation  $\mathbf{NFI}(G)$  is reserved for the poset of finite index normal subgroups of  $G$ . For  $\mathbf{N} \in \mathbf{NFI}(G)$ ,  $\mathcal{P}_{\mathbf{N}}$  denotes the standard projection

$$\mathcal{P}_{\mathbf{N}} : G \rightarrow G/\mathbf{N}, \quad \mathcal{P}_{\mathbf{N}}(g) := g\mathbf{N}.$$

Moreover,  $\widehat{\mathcal{P}}_{\mathbf{N}}$  denotes the standard continuous group homomorphism from the profinite completion  $\widehat{G}$  of  $G$  to the finite group  $G/\mathbf{N}$ . For a group  $G$  and  $g \in G$ , the notation  $\mathcal{Z}_G(g)$  is reserved for the centralizer of  $g$  in  $G$ . If a group  $G$  is residually finite (e.g.  $G = F_n$ ,  $PB_n$  or  $G = B_n$ ), then we tacitly identify  $G$  with its image in the profinite completion  $\widehat{G}$ .

For objects  $a, b$  of a category  $\mathcal{C}$ ,  $\mathcal{C}(a, b)$  often denotes the set of morphisms from  $a$  to  $b$ . For a groupoid  $\mathcal{G}$ , the notation  $\gamma \in \mathcal{G}$  means that  $\gamma$  is a **morphism** of this groupoid. We assume that all functors are covariant and the word “cofunctor” means a contravariant functor.

We will freely use the language of operads [8, Section 3], [11, Chapter 1], [22], [23], [31]. In this paper, we encounter operads in the category of sets and in the category of (topological) groupoids. The category of topological groupoids is treated in the “strict sense”. For example, the associativity axioms for the *elementary insertions*<sup>1</sup>  $\circ_i$  (for operads in the category of groupoids) are satisfied “on the nose”.

---

<sup>1</sup>In the literature, elementary insertions are sometimes called *partial compositions*.

For an integer  $q \geq 1$ , a  $q$ -**truncated operad** in the category of groupoids is a collection of groupoids  $\{\mathcal{G}(n)\}_{1 \leq n \leq q}$  such that

- For every  $1 \leq n \leq q$ , the groupoid  $\mathcal{G}(n)$  is equipped with an action of  $S_n$ .
- For every triple of integers  $i, n, m$  such that  $1 \leq i \leq n$ ,  $n, m, n + m - 1 \leq q$  we have functors

$$\circ_i : \mathcal{G}(n) \times \mathcal{G}(m) \rightarrow \mathcal{G}(n + m - 1). \quad (1.4)$$

- The axioms of the operad for  $\{\mathcal{G}(n)\}_{1 \leq n \leq q}$  are satisfied in the cases where all the arities are  $\leq q$ .

For every operad  $\mathcal{O}$  and for every integer  $q \geq 1$ , the disjoint union

$$\mathcal{O}^{\leq q} := \bigsqcup_{n=0}^q \mathcal{O}(n)$$

is clearly a  $q$ -truncated operad.

In this paper, we mostly consider 4-truncated operads. So, we will simply call them **truncated operads**.

The operad  $\text{PaB}$  of parenthesized braids, its truncation  $\text{PaB}^{\leq 4}$  and its completion  $\widehat{\text{PaB}}^{\leq 4}$  play an important role in this paper. See [3], [7, Appendix A], [11, Chapter 6], [33] for more details about these objects.

## 2 Background material

### 2.1 Reminder of GT-shadows

In this section, we review the groupoid  $\text{GTSh}$  whose objects are elements of the poset  $\text{NFI}_{\text{PB}_4}(\text{B}_4)$  and whose morphisms are called **GT-shadows**.

Recall [7, Section 2.2] that every element  $\mathbf{N} \in \text{NFI}_{\text{PB}_4}(\text{B}_4)$  gives us a compatible equivalence relation  $\sim_{\mathbf{N}}$  on the truncation

$$\text{PaB}^{\leq 4} := \text{PaB}(1) \sqcup \text{PaB}(2) \sqcup \text{PaB}(3) \sqcup \text{PaB}(4) \quad (2.1)$$

of the operad  $\text{PaB}$ , i.e.

- $\sim_{\mathbf{N}}$  is an equivalence relation on the set of morphisms of  $\text{PaB}(1) \sqcup \text{PaB}(2) \sqcup \text{PaB}(3) \sqcup \text{PaB}(4)$ ;
- $\sim$  is compatible with the actions of the symmetric groups  $S_2, S_3, S_4$ ;
- $\sim$  is compatible with the composition of morphisms and the operadic insertions;
- finally, the quotient operad  $\text{PaB}^{\leq 4} / \sim$  is finite.

More precisely, given  $\mathbf{N} \in \text{NFI}_{\text{PB}_4}(\text{B}_4)$ , we produce  $\mathbf{N}_{\text{PB}_3} \in \text{NFI}_{\text{PB}_3}(\text{B}_3)$  and  $\mathbf{N}_{\text{PB}_2} \in \text{NFI}_{\text{PB}_2}(\text{B}_2)$ . (Since  $\text{PB}_2$  is the infinite cyclic group and  $\text{B}_2$  is Abelian,  $\mathbf{N}_{\text{PB}_2}$  is uniquely determined by its index  $N_{\text{ord}} := |\text{PB}_2 : \mathbf{N}_{\text{PB}_2}|$ ). Then  $\mathbf{N}$  (resp.  $\mathbf{N}_{\text{PB}_3}$ ,  $\mathbf{N}_{\text{PB}_2}$ ) gives us an equivalence relation on  $\text{PaB}(4)$  (resp.  $\text{PaB}(3)$ ,  $\text{PaB}(2)$ ). Due to [7, Proposition 2.4], these

equivalence relations on  $\mathbf{PaB}(4)$ ,  $\mathbf{PaB}(3)$  and  $\mathbf{PaB}(2)$  assemble into a compatible equivalence relation<sup>2</sup>  $\sim_{\mathbf{N}}$  on (2.1).

For  $\mathbf{N} \in \mathbf{NFI}_{\mathbf{PB}_4}(\mathbf{B}_4)$ , we denote by  $\mathcal{P}_{\mathbf{N}}$  the standard projection

$$\mathcal{P}_{\mathbf{N}} : \mathbf{PaB}^{\leq 4} \rightarrow \mathbf{PaB}^{\leq 4} / \sim_{\mathbf{N}} . \quad (2.2)$$

Similarly,  $\widehat{\mathcal{P}}_{\mathbf{N}}$  denotes the standard continuous (onto) map of truncated operads

$$\widehat{\mathcal{P}}_{\mathbf{N}} : \widehat{\mathbf{PaB}}^{\leq 4} \rightarrow \mathbf{PaB}^{\leq 4} / \sim_{\mathbf{N}} , \quad (2.3)$$

where  $\widehat{\mathbf{PaB}}^{\leq 4}$  is the profinite completion of  $\mathbf{PaB}^{\leq 4}$ .

We set

$$\mathbf{N}_{\mathbf{F}_2} := \mathbf{N}_{\mathbf{PB}_3} \cap \mathbf{F}_2 , \quad (2.4)$$

where  $\mathbf{F}_2$  is (tacitly) identified with the subgroup  $\langle x_{12}, x_{23} \rangle \leq \mathbf{PB}_3$ .

A *GT-pair* with the target  $\mathbf{N}$  is a morphism of truncated operads  $\mathbf{PaB}^{\leq 4} \rightarrow \mathbf{PaB}^{\leq 4} / \sim_{\mathbf{N}}$  and such morphisms are in bijection with pairs

$$(m, f\mathbf{N}_{\mathbf{PB}_3}) \in \{0, 1, \dots, N_{\text{ord}} - 1\} \times \mathbf{PB}_3 / \mathbf{N}_{\mathbf{PB}_3} \quad (2.5)$$

satisfying the hexagon relations

$$\sigma_1 x_{12}^m f^{-1} \sigma_2 x_{23}^m f \mathbf{N}_{\mathbf{PB}_3} = f^{-1} \sigma_1 \sigma_2 x_{12}^{-m} c^m \mathbf{N}_{\mathbf{PB}_3} , \quad (2.6)$$

$$f^{-1} \sigma_2 x_{23}^m f \sigma_1 x_{12}^m \mathbf{N}_{\mathbf{PB}_3} = \sigma_2 \sigma_1 x_{23}^{-m} c^m f \mathbf{N}_{\mathbf{PB}_3} \quad (2.7)$$

and the pentagon relation

$$\varphi_{234}(f) \varphi_{1,23,4}(f) \varphi_{123}(f) \mathbf{N} = \varphi_{1,2,34}(f) \varphi_{12,3,4}(f) \mathbf{N} . \quad (2.8)$$

Both sides of (2.6) and (2.7) are elements of  $\mathbf{B}_3 / \mathbf{N}_{\mathbf{PB}_3}$  and both sides of (2.8) are elements of  $\mathbf{PB}_4 / \mathbf{N}$ . Explicit formulas for the group homomorphisms  $\varphi_{234}, \varphi_{1,23,4}, \varphi_{123}, \varphi_{1,2,34}$  and  $\varphi_{12,3,4}$  from  $\mathbf{PB}_3$  to  $\mathbf{PB}_4$  are given in [7, Appendix A.4, (A.18)].

Just as in [7], we tacitly identify morphisms  $\mathbf{PaB}^{\leq 4} \rightarrow \mathbf{PaB}^{\leq 4} / \sim_{\mathbf{N}}$  with pairs (2.5) satisfying (2.6), (2.7) and (2.8).

It is convenient to represent GT-pairs by tuples  $(m, f) \in \mathbb{Z} \times \mathbf{PB}_3$  and we denote by  $[m, f]$  the GT-pair represented by the tuple  $(m, f)$ . For a GT-pair  $[m, f]$  with the target  $\mathbf{N}$ , we denote by

$$T_{m,f} : \mathbf{PaB}^{\leq 4} \rightarrow \mathbf{PaB}^{\leq 4} / \sim_{\mathbf{N}} \quad (2.9)$$

the corresponding morphism of (truncated) operads.

It is clear that, for every GT-pair  $[m, f]$  with the target  $\mathbf{N}$ ,  $T_{m,f}$  gives us the following group homomorphisms

$$T_{m,f}^{\mathbf{PB}_4} : \mathbf{PB}_4 \rightarrow \mathbf{PB}_4 / \mathbf{N}, \quad T_{m,f}^{\mathbf{PB}_3} : \mathbf{PB}_3 \rightarrow \mathbf{PB}_3 / \mathbf{N}_{\mathbf{PB}_3}, \quad T_{m,f}^{\mathbf{PB}_2} : \mathbf{PB}_2 \rightarrow \mathbf{PB}_2 / \mathbf{N}_{\mathbf{PB}_2} .$$

For more details, see [7, Corollary 2.7].

A *GT-shadow* with the target  $\mathbf{N}$  is an **onto** morphism of truncated operads  $\mathbf{PaB}^{\leq 4} \rightarrow \mathbf{PaB}^{\leq 4} / \sim_{\mathbf{N}}$  and such morphisms are in bijection with pairs (2.5) that satisfy the following conditions:

---

<sup>2</sup>Note that  $\mathbf{PaB}(1)$  is the groupoid with exactly one object and exactly one (identity) morphism.

- $(m, f)$  obeys relations (2.6), (2.7), (2.8),
- $2m + 1$  represents a unit in the ring  $\mathbb{Z}/N_{\text{ord}}\mathbb{Z}$ , and
- the group homomorphism  $T_{m,f}^{\text{PB}_3} : \text{PB}_3 \rightarrow \text{PB}_3/\mathbf{N}_{\text{PB}_3}$  is onto.

We will identify GT-shadows (with the target  $\mathbf{N}$ ) with pairs (2.5) satisfying the above conditions and denote by  $\text{GT}(\mathbf{N})$  the set of GT-shadows with the target  $\mathbf{N}$ .

Let  $\mathbf{N} \in \text{NFI}_{\text{PB}_4}(\text{B}_4)$  and  $[m, f] \in \text{GT}(\mathbf{N})$ . Due to [7, Proposition 2.11], the “kernel” of the morphism  $T_{m,f}$  (2.9) is a compatible equivalence relation  $\sim_{\mathbf{N}^s}$ , where

$$\mathbf{N}^s := \ker(T_{m,f}^{\text{PB}_4}).$$

Hence  $T_{m,f}$  induces an isomorphism of truncated operads

$$T_{m,f}^{\text{isom}} : \text{PaB}^{\leq 4} / \sim_{\mathbf{N}^s} \xrightarrow{\cong} \text{PaB}^{\leq 4} / \sim_{\mathbf{N}} \quad (2.10)$$

and factors as follows:  $T_{m,f} = T_{m,f}^{\text{isom}} \circ \mathcal{P}_{\mathbf{N}^s}$ .

Thus GT-shadows form a groupoid  $\text{GTSh}$ : objects of  $\text{GTSh}$  are elements of  $\text{NFI}_{\text{PB}_4}(\text{B}_4)$ ; for  $\mathbf{N}^{(1)}, \mathbf{N}^{(2)} \in \text{NFI}_{\text{PB}_4}(\text{B}_4)$ ,

$$\text{GTSh}(\mathbf{N}^{(2)}, \mathbf{N}^{(1)}) := \{[m, f] \in \text{GT}(\mathbf{N}^{(1)}) \mid \mathbf{N}^{(2)} = \ker(T_{m,f}^{\text{PB}_4})\} \quad (2.11)$$

or equivalently  $\text{GTSh}(\mathbf{N}^{(2)}, \mathbf{N}^{(1)})$  consists of isomorphisms of (truncated) operads

$$\text{PaB}^{\leq 4} / \sim_{\mathbf{N}^{(2)}} \xrightarrow{\cong} \text{PaB}^{\leq 4} / \sim_{\mathbf{N}^{(1)}}.$$

The composition of morphisms is defined via the identification of elements  $[m, f] \in \text{GT}(\mathbf{N})$  with isomorphisms (2.10). (See (2.21) below for the explicit formula of the composition of (practical) GT-shadows in terms of their representatives.)

Recall [7, Corollary 2.8] that, for every  $[m, f] \in \text{GT}(\mathbf{N})$ , the homomorphism  $T_{m,f}^{\text{PB}_3} : \text{PB}_3 \rightarrow \text{PB}_3/\mathbf{N}_{\text{PB}_3}$  is given by the explicit formulas:

$$T_{m,f}^{\text{PB}_3}(x_{12}) = x_{12}^{2m+1} \mathbf{N}_{\text{PB}_3}, \quad T_{m,f}^{\text{PB}_3}(x_{23}) = f^{-1} x_{23}^{2m+1} f \mathbf{N}_{\text{PB}_3}, \quad T_{m,f}^{\text{PB}_3}(c) = c^{2m+1} \mathbf{N}_{\text{PB}_3}. \quad (2.12)$$

Since  $\text{PB}_3 \cong \langle x_{12}, x_{23} \rangle \times \langle c \rangle$ , the restriction of  $T_{m,f}^{\text{PB}_3}$  to  $\text{F}_2 = \langle x_{12}, x_{23} \rangle$  gives us a group homomorphism

$$T_{m,f}^{\text{F}_2} : \text{F}_2 \rightarrow \text{F}_2/\mathbf{N}_{\text{F}_2} \quad (2.13)$$

with

$$T_{m,f}^{\text{F}_2}(x) := x^{2m+1} \mathbf{N}_{\text{F}_2}, \quad T_{m,f}^{\text{F}_2}(y) := f^{-1} y^{2m+1} f \mathbf{N}_{\text{F}_2}. \quad (2.14)$$

Let us prove the following useful statement:

**Proposition 2.1** *For every  $\mathbf{N} \in \text{NFI}_{\text{PB}_4}(\text{B}_4)$  and every  $[m, f] \in \text{GT}(\mathbf{N})$ , the group homomorphism (2.13) is onto.*

**Proof.** Due to [7, Proposition 2.3],  $N_{\text{ord}}$  is the least common multiple of the orders of the elements  $x_{12} \mathbf{N}_{\text{PB}_3}$ ,  $x_{23} \mathbf{N}_{\text{PB}_3}$  and  $c \mathbf{N}_{\text{PB}_3}$  in  $\text{PB}_3/\mathbf{N}_{\text{PB}_3}$ .

Therefore, since  $2m + 1$  is coprime with  $N_{\text{ord}}$ , there exist  $q_1, q_2 \in \mathbb{Z}$  such that

$$T_{m,f}^{\text{F}_2}(x_{12}^{q_1}) = x_{12} \mathbf{N}_{\text{F}_2}, \quad T_{m,f}^{\text{F}_2}(x_{23}^{q_2}) = f^{-1} x_{23} f \mathbf{N}_{\text{F}_2}. \quad (2.15)$$



Since the homomorphism  $T_{m,f}^{\text{PB}_3}$  is onto, there exists  $w \in \text{PB}_3$  such that

$$T_{m,f}^{\text{PB}_3}(w) = f\mathbf{N}_{\text{PB}_3} \quad (2.16)$$

Since  $\text{PB}_3 = \langle x_{12}, x_{23} \rangle \times \langle c \rangle$ ,

$$w = w_1 c^k,$$

where  $w_1 \in \text{F}_2$ . Therefore, using (2.16), we get

$$T_{m,f}^{\text{F}_2}(w_1 x_{23}^{q_2} w_1^{-1}) = T_{m,f}^{\text{PB}_3}(w_1 c^k x_{23}^{q_2} c^{-k} w_1^{-1}) =$$

$$T_{m,f}^{\text{PB}_3}(w x_{23}^{q_2} w^{-1}) = f(f^{-1} x_{23} f) f^{-1} \mathbf{N}_{\text{PB}_3} = x_{23} \mathbf{N}_{\text{PB}_3}.$$

Thus,  $T_{m,f}^{\text{F}_2}(w_1 x_{23}^{q_2} w_1^{-1}) = x_{23} \mathbf{N}_{\text{F}_2}$ .

Since both generators  $x_{12} \mathbf{N}_{\text{F}_2}$  and  $x_{23} \mathbf{N}_{\text{F}_2}$  of  $\text{F}_2/\mathbf{N}_{\text{F}_2}$  belong to  $T_{m,f}^{\text{F}_2}(\text{F}_2)$ , we proved that  $T_{m,f}^{\text{F}_2}$  is indeed onto.  $\square$

### 2.1.1 $\widehat{\text{GT}}$ versus the groupoid GTSh

Just as in [7], we denote by  $\mathcal{I}$  the standard morphism  $\text{PaB}^{\leq 4} \rightarrow \widehat{\text{PaB}}^{\leq 4}$ .

Let  $\hat{T} \in \widehat{\text{GT}}$  and  $\mathbf{N} \in \text{NFI}_{\text{PB}_4}(\text{B}_4)$ . It was shown in [7, Section 2.4] that the formula

$$T_{\mathbf{N}} := \widehat{\mathcal{P}}_{\mathbf{N}} \circ \hat{T} \circ \mathcal{I} : \text{PaB}^{\leq 4} \rightarrow \text{PaB}^{\leq 4} / \sim_{\mathbf{N}} \quad (2.17)$$

defines a GT-shadow with the target  $\mathbf{N}$ .

Let us prove that

**Proposition 2.2** *The assignment*

$$\mathbf{N}^{\hat{T}} := \ker \left( \text{PB}_4 \xrightarrow{T_{\mathbf{N}}^{\text{PB}_4}} \text{PB}_4 / \mathbf{N} \right) \quad (2.18)$$

*defines a right action of  $\widehat{\text{GT}}$  on the set  $\text{NFI}_{\text{PB}_4}(\text{B}_4)$ . Moreover, the assignment  $\hat{T} \mapsto T_{\mathbf{N}}$  defines a functor  $\mathcal{PR}$  from the corresponding transformation groupoid to GTSh.*

**Proof.** It is clear that, if  $\hat{T}$  is the identity element of  $\widehat{\text{GT}}$ , then

$$T_{\mathbf{N}} = \mathcal{P}_{\mathbf{N}} : \text{PaB}^{\leq 4} \rightarrow \text{PaB}^{\leq 4} / \sim_{\mathbf{N}}.$$

For every  $\hat{T} \in \widehat{\text{GT}}$ , we have the following commutative diagram of maps of truncated operads:

$$\begin{array}{ccc} \widehat{\text{PaB}}^{\leq 4} & \xrightarrow{\hat{T}} & \widehat{\text{PaB}}^{\leq 4} \\ \widehat{\mathcal{P}}_{\mathbf{N}^{\hat{T}}} \downarrow & & \downarrow \widehat{\mathcal{P}}_{\mathbf{N}} \\ \text{PaB}^{\leq 4} / \sim_{\mathbf{N}^{\hat{T}}} & \xrightarrow{T_{\mathbf{N}}^{\text{isom}}} & \text{PaB}^{\leq 4} / \sim_{\mathbf{N}} \end{array} \quad (2.19)$$

Let  $\hat{T}_1, \hat{T}_2 \in \widehat{\mathbf{GT}}$ ,  $\mathbf{N} \in \mathbf{NFI}_{\mathbf{PB}_4}(\mathbf{B}_4)$ ,  $\mathbf{N}^{(1)} := \mathbf{N}^{\hat{T}_1}$  and  $\mathbf{N}^s := (\mathbf{N}^{(1)})^{\hat{T}_2}$ . Combining the corresponding commutative diagrams for  $\hat{T}_1$  and  $\hat{T}_2$ , we get the commutative diagram:

$$\begin{array}{ccccc}
 & \widehat{\mathbf{PaB}}^{\leq 4} & \xrightarrow{\hat{T}_2} & \widehat{\mathbf{PaB}}^{\leq 4} & \xrightarrow{\hat{T}_1} & \widehat{\mathbf{PaB}}^{\leq 4} \\
 & \downarrow \hat{\mathcal{P}}_{\mathbf{N}^s} & & \downarrow \hat{\mathcal{P}}_{\mathbf{N}^{(1)}} & & \downarrow \hat{\mathcal{P}}_{\mathbf{N}} \\
 \mathbf{PaB}^{\leq 4} & \xrightarrow{\mathcal{P}_{\mathbf{N}^s}} & \mathbf{PaB}^{\leq 4} / \sim_{\mathbf{N}^s} & \xrightarrow{T_{2, \mathbf{N}^{(1)}}^{\text{isom}}} & \mathbf{PaB}^{\leq 4} / \sim_{\mathbf{N}^{(1)}} & \xrightarrow{T_{1, \mathbf{N}}^{\text{isom}}} & \mathbf{PaB}^{\leq 4} / \sim_{\mathbf{N}} \\
 & \searrow & & \searrow & & \searrow & \\
 & & & & & & T_{\mathbf{N}}
 \end{array} \tag{2.20}$$

where  $\hat{T} := \hat{T}_1 \circ \hat{T}_2$  and  $T_{\mathbf{N}}$  is the corresponding  $\mathbf{GT}$ -shadow with the target  $\mathbf{N}$ .

Thus

$$(\mathbf{N}^{\hat{T}_1})^{\hat{T}_2} = \mathbf{N}^{\hat{T}_1 \circ \hat{T}_2}.$$

We proved that formula (2.18) defines a (right) action of  $\widehat{\mathbf{GT}}$  on  $\mathbf{NFI}_{\mathbf{PB}_4}(\mathbf{B}_4)$ .

Let us denote by  $\widehat{\mathbf{GT}}_{\mathbf{NFI}}$  the corresponding transformation groupoid. The set of objects of  $\widehat{\mathbf{GT}}_{\mathbf{NFI}}$  is  $\mathbf{NFI}_{\mathbf{PB}_4}(\mathbf{B}_4)$  and, for  $\mathbf{N}^{(1)}, \mathbf{N}^{(2)} \in \mathbf{NFI}_{\mathbf{PB}_4}(\mathbf{B}_4)$ , the set of morphisms  $\widehat{\mathbf{GT}}_{\mathbf{NFI}}(\mathbf{N}^{(1)}, \mathbf{N}^{(2)})$  consists of elements  $\hat{T} \in \widehat{\mathbf{GT}}$  such that  $(\mathbf{N}^{(2)})^{\hat{T}} = \mathbf{N}^{(1)}$ .

The diagram in (2.20) tells us that, if  $\hat{T} := \hat{T}_1 \circ \hat{T}_2$ , then

$$T_{\mathbf{N}}^{\text{isom}} = T_{1, \mathbf{N}}^{\text{isom}} \circ T_{2, \mathbf{N}^{(1)}}^{\text{isom}}.$$

Thus the assignment  $\hat{T} \mapsto T_{\mathbf{N}}$  indeed defines a functor  $\mathcal{PR}$  from  $\widehat{\mathbf{GT}}_{\mathbf{NFI}}$  to  $\mathbf{GTSh}$ . (On the level of objects, the functor  $\mathcal{PR}$  operates as the identity map.)  $\square$

Recall [7, Section 2.6] that a  $\mathbf{GT}$ -shadow  $[m, f] \in \mathbf{GT}(\mathbf{N})$  is called **genuine** if there exists  $\hat{T} \in \widehat{\mathbf{GT}}$  such that  $T_{m, f} = T_{\mathbf{N}}$ , i.e.  $[m, f]$  comes from an element of  $\widehat{\mathbf{GT}}$ . If such  $\hat{T} \in \widehat{\mathbf{GT}}$  does not exist then the  $\mathbf{GT}$ -shadow  $[m, f]$  is called **fake**.

It was shown in [7, Section 2] that genuine  $\mathbf{GT}$ -shadows satisfy additional conditions.

A  $\mathbf{GT}$ -shadow is called **practical**, if it can be represented by a pair  $(m, f)$  where<sup>3</sup>  $f \in \mathbf{F}_2$ . Since every genuine  $\mathbf{GT}$ -shadow is practical<sup>4</sup>, in this paper, we assume that all  $\mathbf{GT}$ -shadows are practical. In particular,  $\mathbf{GT}(\mathbf{N})$  denotes the set of all practical  $\mathbf{GT}$ -shadows with the target  $\mathbf{N}$ . Furthermore, we will use the same notation  $\mathbf{GTSh}$  for the (sub)groupoid of a practical  $\mathbf{GT}$ -shadows.

If  $[m_1, f_1] \in \mathbf{GTSh}(\mathbf{N}^{(2)}, \mathbf{N}^{(1)})$ ,  $[m_2, f_2] \in \mathbf{GTSh}(\mathbf{N}^{(3)}, \mathbf{N}^{(2)})$  and

$$\begin{aligned}
 m &:= 2m_1m_2 + m_1 + m_2, \\
 f(x, y) &:= f_1(x, y) f_2(x^{2m_1+1}, f_1(x, y)^{-1} y^{2m_1+1} f_1(x, y)),
 \end{aligned} \tag{2.21}$$

then the pair  $(m, f)$  represents a  $\mathbf{GT}$ -shadow in  $\mathbf{GTSh}(\mathbf{N}^{(3)}, \mathbf{N}^{(1)})$  and  $[m, f] := [m_1, f_1] \circ [m_2, f_2]$ , i.e. formula (2.21) defines the composition of (practical)  $\mathbf{GT}$ -shadows. For more details, see [7, Remark 2.15].

<sup>3</sup>Recall that  $\mathbf{F}_2$  is identified with the subgroup  $\langle x_{12}, x_{23} \rangle \leq \mathbf{PB}_3$ .

<sup>4</sup>See, for example, [7, Proposition 2.20].

It was proved in [7, Proposition 2.20] that, for every genuine GT-shadow  $[m, f] \in \text{GT}(\mathbf{N})$ , the coset  $f\mathbf{N}_{F_2}$  belongs to the commutator subgroup  $[F_2/\mathbf{N}_{F_2}, F_2/\mathbf{N}_{F_2}]$

$$f\mathbf{N}_{F_2} \in [F_2/\mathbf{N}_{F_2}, F_2/\mathbf{N}_{F_2}]. \quad (2.22)$$

GT-shadows satisfying this additional property are called **charming**.

Just as in [7], the notation  $\text{GT}^\heartsuit(\mathbf{N})$  is reserved for the subset of charming GT-shadows in  $\text{GT}(\mathbf{N})$ . Due to [7, Proposition 2.22], charming GT-shadows form a subgroupoid  $\text{GTSh}^\heartsuit$  of  $\text{GTSh}$ .

**Remark 2.3** Due to Proposition 2.1, the condition about the homomorphism  $T_{m,f}^{F_2}$  in [7, Definition 2.19] is redundant. In other words, a GT-shadow  $[m, f] \in \text{GT}(\mathbf{N})$  is charming if and only if condition (2.22) is satisfied.

**Remark 2.4** Since, for every  $\hat{T} \in \widehat{\text{GT}}$  and  $\mathbf{N} \in \text{NFI}_{\text{PB}_4}(\mathbf{B}_4)$ , the GT-shadow  $T_{\mathbf{N}} \in \text{GT}(\mathbf{N})$  is charming, the functor  $\mathcal{PR} : \widehat{\text{GT}}_{\text{NFI}} \rightarrow \text{GTSh}$  from Proposition 2.2 lands in the subgroupoid  $\text{GTSh}^\heartsuit$ .

Recall that the groupoid  $\text{GTSh}^\heartsuit$  is highly disconnected. Indeed, if  $\mathbf{K}, \mathbf{N} \in \text{NFI}_{\text{PB}_4}(\mathbf{B}_4)$  have different indices in  $\text{PB}_4$ , then  $\text{GTSh}^\heartsuit(\mathbf{K}, \mathbf{N})$  is empty. Just as in [7],  $\text{GTSh}_{\text{conn}}^\heartsuit(\mathbf{N})$  denotes the connected component of  $\mathbf{N}$  in the groupoid  $\text{GTSh}^\heartsuit$ . Elements  $\mathbf{N} \in \text{NFI}_{\text{PB}_4}(\mathbf{B}_4)$  for which the connected component  $\text{GTSh}_{\text{conn}}^\heartsuit(\mathbf{N})$  has exactly one object are called **isolated** and they play a special role:

- For every isolated element  $\mathbf{N} \in \text{NFI}_{\text{PB}_4}(\mathbf{B}_4)$ ,  $\text{GT}^\heartsuit(\mathbf{N})$  is a finite group.
- Due to [7, Proposition 3.3], the subposet  $\text{NFI}_{\text{PB}_4}^{\text{isolated}}(\mathbf{B}_4)$  of isolated elements in  $\text{NFI}_{\text{PB}_4}(\mathbf{B}_4)$  is cofinal: for every  $\mathbf{N} \in \text{NFI}_{\text{PB}_4}(\mathbf{B}_4)$ , there exists  $\mathbf{K} \in \text{NFI}_{\text{PB}_4}^{\text{isolated}}(\mathbf{B}_4)$  such that  $\mathbf{K} \leq \mathbf{N}$ .
- Due to [7, Proposition 3.6],  $\text{NFI}_{\text{PB}_4}^{\text{isolated}}(\mathbf{B}_4)$  is closed under taking finite intersections.
- Due to [7, Proposition 3.7], the assignment  $\mathbf{N} \rightarrow \text{GT}^\heartsuit(\mathbf{N})$  upgrades to a functor  $\mathcal{ML}$  from the poset  $\text{NFI}_{\text{PB}_4}^{\text{isolated}}(\mathbf{B}_4)$  to the category of finite groups.
- Finally, due to [7, Theorem 3.8], the limit of  $\mathcal{ML}$  is isomorphic to  $\widehat{\text{GT}}$ .

## 2.2 Permutation pairs, permutation triples and child's drawings

A **permutation pair** of degree  $d$  is a pair  $c = (c_1, c_2) \in S_d \times S_d$  for which the subgroup  $\langle c_1, c_2 \rangle \leq S_d$  is transitive. We consider permutation pairs with the action of  $S_d$  by conjugation

$$h(c) := (hc_1h^{-1}, hc_2h^{-1}), \quad h \in S_d.$$

Recall that a **child's drawing** of degree  $d$  is a conjugacy class  $[c]$  of a permutation pair  $c = (c_1, c_2)$ . Let us denote by  $\mathbf{ct}$  the standard map from  $S_d$  to the set  $\mathfrak{P}_d$  of partitions of  $d$ : for  $h \in S_d$ ,  $\mathbf{ct}(h)$  is the cycle structure of  $h$ .

The (conjugacy class of the) permutation group  $\langle c_1, c_2 \rangle \leq S_d$  is called the **monodromy group** of the child's drawing  $[c]$ .

In the literature<sup>5</sup> [1], [2], [10], [12], [13], [14], [21], [27], [30], [34], child's drawings (of degree  $d$ ) are often represented by permutation triples, i.e. elements  $(g_1, g_2, g_3)$  in  $(S_d)^3$  such that

- $g_1 g_2 g_3 = \text{id}$  and
- the subgroup  $\langle g_1, g_2, g_3 \rangle \leq S_d$  is transitive.

The assignment

$$(c_1, c_2) \mapsto (c_1, c_2, c_2^{-1} c_1^{-1}) \quad (2.23)$$

gives us an obvious bijection from the set of permutation pairs (of degree  $d$ ) to the set of permutation triples (of degree  $d$ ).  $S_d$  acts on permutation triples by conjugation

$$h(g_1, g_2, g_3) := (h g_1 h^{-1}, h g_2 h^{-1}, h g_3 h^{-1})$$

and this action is compatible with bijection (2.23).

The triple of partitions  $(\mathbf{ct}(c_1), \mathbf{ct}(c_2), \mathbf{ct}(c_2^{-1} c_1^{-1}))$  is called the **passport** of a child's drawing  $[c]$ .

### 2.3 Representing child's drawings by group homomorphisms

It is convenient to represent a child's drawing  $[c]$  of degree  $d$  by the group homomorphism  $\psi : \mathbf{F}_2 \rightarrow S_d$ :

$$\psi(x) := c_1, \quad \psi(y) := c_2.$$

So we denote by

$$\text{Hom}_{\text{tran}}(\mathbf{F}_2, S_d) \quad (2.24)$$

the set of group homomorphisms  $\psi : \mathbf{F}_2 \rightarrow S_d$  for which  $\psi(\mathbf{F}_2)$  is transitive.

Two homomorphisms  $\psi, \tilde{\psi} \in \text{Hom}_{\text{tran}}(\mathbf{F}_2, S_d)$  represent the same child's drawing if and only if  $\exists h \in S_d$  such that

$$\tilde{\psi}(w) = h \psi(w) h^{-1}, \quad \forall w \in \mathbf{F}_2.$$

In other words, the set of child's drawing of degree  $d$  can be identified with the set of orbits

$$\text{Hom}_{\text{tran}}(\mathbf{F}_2, S_d)_{S_d} \quad (2.25)$$

of the  $S_d$ -action on  $\text{Hom}_{\text{tran}}(\mathbf{F}_2, S_d)$  by conjugation.

Note that  $\ker(\psi)$  depends only on the child's drawing  $[\psi]$  but not on a particular choice of a representative  $\psi \in \text{Hom}_{\text{tran}}(\mathbf{F}_2, S_d)$ .

Child's drawings can also be represented by continuous group homomorphisms from  $\widehat{\mathbf{F}}_2$  to  $S_d$ . The goal of the following proposition is to recall this equivalent description:

**Proposition 2.5** *Let  $\text{Hom}(\widehat{\mathbf{F}}_2, S_d)$  be the set of continuous group homomorphisms<sup>6</sup>  $\widehat{\mathbf{F}}_2 \rightarrow S_d$  and*

$$\text{Hom}(\widehat{\mathbf{F}}_2, S_d)_{\text{tran}} := \{ \hat{\psi} \in \text{Hom}(\widehat{\mathbf{F}}_2, S_d) \mid \hat{\psi}(\widehat{\mathbf{F}}_2) \text{ is a transitive subgroup of } S_d \}.$$

<sup>5</sup>This list of references is far from complete.

<sup>6</sup>Recall that every finite group (for example,  $S_d$ ) is considered with the discrete topology.

The assignment

$$\hat{\psi} \mapsto \hat{\psi}|_{F_2} \quad (2.26)$$

gives us a bijection from  $\text{Hom}(\widehat{F}_2, S_d)$  (resp. from  $\text{Hom}_{\text{tran}}(\widehat{F}_2, S_d)$ ) to  $\text{Hom}(F_2, S_d)$  (resp. to  $\text{Hom}_{\text{tran}}(F_2, S_d)$ ). This assignment is compatible with the action of  $S_d$  by conjugation and it gives us a bijection between the set

$$\text{Hom}_{\text{tran}}(\widehat{F}_2, S_d)_{S_d} \quad (2.27)$$

of orbits of the  $S_d$ -action and the set of child's drawings of degree  $d$ .

**Proof.** We will prove this proposition by constructing the inverse of the assignment in (2.26).

Let  $\psi \in \text{Hom}(F_2, S_d)$  and  $K := \ker(\psi)$ . The homomorphism  $\psi : F_2 \rightarrow S_d$  factors as follows

$$\psi = \psi_K \circ \mathcal{P}_K,$$

where the homomorphism  $\psi_K : F_2/K \rightarrow S_d$  is defined by the formula

$$\psi_K(wK) := \psi(w). \quad (2.28)$$

We denote by  $\hat{\psi}$  the continuous homomorphism  $\widehat{F}_2 \rightarrow S_d$  defined by the formula

$$\hat{\psi} := \psi_K \circ \widehat{\mathcal{P}}_K. \quad (2.29)$$

We claim that the assignment

$$\psi \mapsto \hat{\psi}$$

gives us the inverse of the map from  $\text{Hom}(\widehat{F}_2, S_d)$  to  $\text{Hom}(F_2, S_d)$  defined in (2.26).

Indeed,  $\hat{\psi}|_{F_2}$  clearly coincides with  $\psi$ . Thus it remains to show that, if

$$\psi = \hat{\varphi}|_{F_2} \quad (2.30)$$

for  $\hat{\varphi} \in \text{Hom}(\widehat{F}_2, S_d)$ , then  $\hat{\psi}$  coincides with  $\hat{\varphi}$ .

Let  $K := \ker(\psi)$  and  $\psi_K$  be the homomorphism  $F_2/K \rightarrow S_d$  defined by (2.28), where  $\psi$  is defined in (2.30). As above, we set  $\hat{\psi} := \psi_K \circ \widehat{\mathcal{P}}_K$  and observe that

$$\hat{\psi}|_{F_2} = \hat{\varphi}|_{F_2}.$$

Since  $F_2$  is dense in  $\widehat{F}_2$ ,  $S_d$  is Hausdorff and  $\hat{\psi}, \hat{\varphi}$  are continuous, we conclude that  $\hat{\psi} = \hat{\varphi}$ .

It is easy to see that, for every  $\hat{\psi} \in \text{Hom}(\widehat{F}_2, S_d)$ ,  $\psi(F_2) = \psi(\widehat{F}_2)$ . Moreover the assignments (2.26) and  $\psi \mapsto \hat{\psi}$  are compatible with the (adjoint) action of  $S_d$ . Thus the remaining statements of the proposition are obvious.  $\square$

**Remark 2.6** In the above proof, we showed that every group homomorphism  $\psi : F_2 \rightarrow S_d$  extends uniquely to a continuous group homomorphism  $\hat{\psi} : \widehat{F}_2 \rightarrow S_d$ . Equation (2.29) is an explicit definition of this extension. See also [25, Lemma 1.1.16, (b)].

**Remark 2.7** Depending on the context, we will represent a child's drawing of degree  $d$  by a group homomorphism  $\psi : \mathbf{F}_2 \rightarrow S_d$  or by a continuous group homomorphism  $\hat{\psi} : \widehat{\mathbf{F}}_2 \rightarrow S_d$ . Of course, due to Proposition 2.5, we have

$$[\hat{\psi}] = [\psi|_{\mathbf{F}_2}]$$

for every  $\hat{\psi} \in \text{Hom}_{\text{tran}}(\widehat{\mathbf{F}}_2, S_d)$ .

**Remark 2.8** There is a natural bijection between the set  $\text{Hom}_{\text{tran}}(\mathbf{F}_2, S_d)_{S_d}$  and the set of conjugacy classes of index  $d$  subgroups  $H$  in  $\mathbf{F}_2$ . This bijection sends  $[\psi] \in \text{Hom}_{\text{tran}}(\mathbf{F}_2, S_d)_{S_d}$  to the conjugacy class (in  $\mathbf{F}_2$ ) of the subgroup<sup>7</sup>

$$H_\psi := \{h \in \mathbf{F}_2 \mid \psi(h)(1) = 1\}.$$

The inverse of this correspondence operates as follows: given a subgroup  $H \leq \mathbf{F}_2$  of index  $d$ , we choose a bijection between the set  $\mathbf{F}_2/H$  of left cosets of  $H$  and the set  $\{1, 2, \dots, d\}$ ; then the canonical action of  $\mathbf{F}_2$  on  $\mathbf{F}_2/H$  gives us a desired group homomorphism  $\psi : \mathbf{F}_2 \rightarrow S_d$ . Thus every child's drawing of degree  $d$  can be represented by an index  $d$  subgroup in  $\mathbf{F}_2$ .

**Remark 2.9** Recall [18, Section 1.3] that conjugacy classes of index  $d$  subgroups in  $\mathbf{F}_2$  are in bijection with connected degree  $d$  covering spaces of  $\mathbb{CP}^1 \setminus \{0, 1, \infty\}$ . We say that a child's drawing  $[H]$  represented by a finite index subgroup  $H \leq \mathbf{F}_2$  is **Galois** if the corresponding covering of  $\mathbb{CP}^1 \setminus \{0, 1, \infty\}$  is Galois. Due to [18, Proposition 1.39], the child's drawing  $[H]$  is Galois if and only if  $H$  is a normal subgroup of  $\mathbf{F}_2$ .

Recall that a **Belyi pair** is a pair  $(X, \gamma)$ , where  $X$  is a smooth projective curve defined over  $\overline{\mathbb{Q}}$  and  $\gamma : X \rightarrow \mathbb{P}_{\overline{\mathbb{Q}}}^1$  is a morphism of curves unramified outside of the set  $\{0, 1, \infty\}$ . Using [32, Proposition 4.5.13] and [32, Theorem 4.6.10], one can show<sup>8</sup> that isomorphism classes of connected degree  $d$  coverings of  $\mathbb{CP}^1 \setminus \{0, 1, \infty\}$  are in bijection with isomorphism classes of degree  $d$  Belyi pairs. Thus child's drawings can be also represented by Belyi pairs. The standard action of  $G_{\mathbb{Q}}$  on child's drawings is often defined in terms of Belyi pairs.

For more details, we refer to the reader to books [12], [21], blog [13], and survey [30]. The curious reader may also try to “surf through” the database of Belyi pairs [24].

## 2.4 The action of $\widehat{\mathbf{GT}}$ on child's drawings and the functor $\mathcal{A} : \widehat{\mathbf{GT}}_{\text{NFI}} \rightarrow \text{Dessin}$

The goal of the following proposition is to recall the action of  $\widehat{\mathbf{GT}}$  on child's drawings.

**Proposition 2.10** *Let  $\hat{T} \in \widehat{\mathbf{GT}}$  and  $\hat{T}_{\widehat{\mathbf{F}}_2}$  be the corresponding continuous automorphism of  $\widehat{\mathbf{F}}_2 \leq \widehat{\text{PB}}_3 = \text{PaB}((12)3, 1(23))$ . Then the formula*

$$\hat{\psi}^{\hat{T}} := \hat{\psi} \circ \hat{T}_{\widehat{\mathbf{F}}_2} \tag{2.31}$$

*defines a right action of  $\widehat{\mathbf{GT}}$  on the set  $\text{Hom}(\widehat{\mathbf{F}}_2, S_d)$ . This action descends to the action of  $\widehat{\mathbf{GT}}$  on  $\text{Hom}_{\text{tran}}(\widehat{\mathbf{F}}_2, S_d)$  and on  $\text{Hom}_{\text{tran}}(\widehat{\mathbf{F}}_2, S_d)_{S_d}$ .*

<sup>7</sup>Since the subgroup  $\psi(\mathbf{F}_2) \leq S_d$  acts transitively on  $\{1, 2, \dots, d\}$ , the subgroup  $H_{\psi, i} := \{h \in \mathbf{F}_2 \mid \psi(h)(i) = i\} \leq \mathbf{F}_2$  is conjugate to  $H_\psi$  in  $\mathbf{F}_2$  for every  $i \in \{1, 2, \dots, d\}$ .

<sup>8</sup>Some mathematicians also like to cite [15].

**Proof.** It is obvious that, for  $\hat{T}^{(1)}, \hat{T}^{(2)} \in \widehat{\mathbf{GT}}$ ,

$$(\hat{T}^{(1)} \circ \hat{T}^{(2)})_{\hat{F}_2} = \hat{T}_{\hat{F}_2}^{(1)} \circ \hat{T}_{\hat{F}_2}^{(2)}.$$

Thus the first statement of the proposition is obvious.

Since  $\hat{T}_{\hat{F}_2}$  is an isomorphism  $\hat{F}_2 \xrightarrow{\sim} \hat{F}_2$ , the subgroups  $\hat{\psi} \circ \hat{T}_{\hat{F}_2}(\hat{F}_2) \leq S_d$  and  $\hat{\psi}(\hat{F}_2) \leq S_d$  coincide. Moreover, the resulting action of  $\widehat{\mathbf{GT}}$  on  $\text{Hom}_{\text{tran}}(\hat{F}_2, S_d)$  clearly commutes with the action of  $S_d$ . Hence the remaining two statements of the proposition are also obvious.  $\square$

**On the Ihara embedding.** Recall [20] that  $G_{\mathbb{Q}}$  injects into the group  $\widehat{\mathbf{GT}}$

$$G_{\mathbb{Q}} \hookrightarrow \widehat{\mathbf{GT}}, \quad (2.32)$$

and the standard action of  $G_{\mathbb{Q}}$  on child's drawings agrees with homomorphism (2.32) and the above action of  $\widehat{\mathbf{GT}}$  on child's drawings. For details, please see [20, Proposition 1.6], [20, Theorem 1.7] and [28, Section 3.2]. We will call homomorphism (2.32) the **Ihara embedding**.

For our purposes, it is convenient to describe the action of  $\widehat{\mathbf{GT}}$  on child's drawing in terms of a functor from the transformation groupoid  $\widehat{\mathbf{GT}}_{\text{NFI}}$  to certain category assembled from child's drawings and elements of the poset  $\text{NFI}_{\text{PB}_4}(\text{B}_4)$ .

**Definition 2.11** *Let  $\mathbf{N} \in \text{NFI}_{\text{PB}_4}(\text{B}_4)$  and  $[\psi]$  be a child's drawing of degree  $d$  represented by a homomorphism  $\psi : F_2 \rightarrow S_d$ . We say that a child's drawing  $[\psi]$  is **subordinate** to  $\mathbf{N}$  if*

$$\mathbf{N}_{F_2} \leq \ker(\psi). \quad (2.33)$$

*If  $[\psi]$  is subordinate to  $\mathbf{N}$ , then we say that  $\mathbf{N}$  **dominates**  $[\psi]$ .*

It is easy to see that condition (2.33) does not depend on the choice of representing homomorphism  $\psi$ . Moreover, if a child's drawing is represented by a continuous homomorphism  $\hat{\psi} : \hat{F}_2 \rightarrow S_d$  then  $[\hat{\psi}]$  is subordinate to  $\mathbf{N}$  if and only if  $\mathbf{N}_{F_2} \leq \ker(\hat{\psi}|_{F_2})$ . We denote by  $\text{Dessin}(\mathbf{N})$  the set of child's drawings subordinate to  $\mathbf{N}$ .

It is clear that, if  $\mathbf{K} \leq \mathbf{N}$  ( $\mathbf{K}, \mathbf{N} \in \text{NFI}_{\text{PB}_4}(\text{B}_4)$ ) and child's drawing  $[\psi]$  is subordinate to  $\mathbf{N}$  then  $[\psi]$  is also subordinate to  $\mathbf{K}$ . In other words, if  $\mathbf{K} \leq \mathbf{N}$ , then  $\text{Dessin}(\mathbf{N}) \subset \text{Dessin}(\mathbf{K})$ .

Let us introduce the category **Dessin** whose objects are elements of  $\text{NFI}_{\text{PB}_4}(\text{B}_4)$ . For  $\mathbf{N}^{(1)}, \mathbf{N}^{(2)} \in \text{NFI}_{\text{PB}_4}(\text{B}_4)$ , morphisms from  $\mathbf{N}^{(1)}$  to  $\mathbf{N}^{(2)}$  are functions from  $\text{Dessin}(\mathbf{N}^{(1)})$  to  $\text{Dessin}(\mathbf{N}^{(2)})$ .

**Proposition 2.12** *For every  $\hat{T} \in \widehat{\mathbf{GT}}$  and  $[\hat{\psi}] \in \text{Dessin}(\mathbf{N})$ , the child's drawing represented by the homomorphism*

$$\hat{\psi} \circ \hat{T}|_{F_2} : F_2 \rightarrow S_d$$

*is subordinate to  $\mathbf{N}^{\hat{T}}$ . Moreover, the formulas*

$$\mathcal{A}(\mathbf{N}) := \mathbf{N}, \quad \mathcal{A}(\hat{T})([\hat{\psi}]) := [\hat{\psi} \circ \hat{T}] \quad (2.34)$$

*define a cofunctor*

$$\mathcal{A} : \widehat{\mathbf{GT}}_{\text{NFI}} \rightarrow \text{Dessin}.$$

**Proof.** Identifying  $F_2$  and  $\widehat{F}_2$  with the corresponding subgroups of  $\widehat{PB}_3 = \text{PaB}((1, 2)3, (1, 2)3)$  and using (2.17) we see that  $N_{F_2}^{\widehat{T}}$  is the kernel of the homomorphism

$$\widehat{\mathcal{P}}_{N_{F_2}} \circ \widehat{T}|_{F_2} : F_2 \rightarrow F_2/N_{F_2}.$$

Let  $\psi := \widehat{\psi}|_{F_2}$ . Since  $N_{F_2} \leq \ker(\psi)$ ,  $\psi$  induces the group homomorphism

$$\tilde{\psi} : F_2/N_{F_2} \rightarrow S_d, \quad \tilde{\psi}(wN_{F_2}) := \psi(w).$$

Composing  $\tilde{\psi}$  with  $\widehat{\mathcal{P}}_{N_{F_2}}$  we get a continuous group homomorphism  $\tilde{\psi} \circ \widehat{\mathcal{P}}_{N_{F_2}} : \widehat{F}_2 \rightarrow S_d$ . Since  $\tilde{\psi} \circ \widehat{\mathcal{P}}_{N_{F_2}}|_{F_2} = \widehat{\psi}|_{F_2}$ ,  $F_2$  is dense in  $\widehat{F}_2$  and  $S_d$  is Hausdorff, we conclude that

$$\widehat{\psi} = \tilde{\psi} \circ \widehat{\mathcal{P}}_{N_{F_2}}.$$

Hence

$$\widehat{\psi} \circ \widehat{T}|_{F_2} = \tilde{\psi} \circ (\widehat{\mathcal{P}}_{N_{F_2}} \circ \widehat{T}|_{F_2}). \quad (2.35)$$

Since the kernel of  $\widehat{\mathcal{P}}_{N_{F_2}} \circ \widehat{T}|_{F_2}$  is  $N_{F_2}^{\widehat{T}}$ , (2.35) implies that

$$N_{F_2}^{\widehat{T}} \leq \ker(\widehat{\psi} \circ \widehat{T}|_{F_2}).$$

The first statement of the proposition is proved.

Using the statement we just proved and Proposition 2.10 we see that, for all  $N^{(1)}, N^{(2)} \in \text{NFI}_{PB_4}(B_4)$ , the second formula in (2.34) defines a map

$$\widehat{GT}_{\text{NFI}}(N^{(1)}, N^{(2)}) \rightarrow \text{Dessin}(N^{(2)}, N^{(1)}).$$

Moreover, since  $\widehat{\psi} \circ \text{id}_{\widehat{F}_2} = \widehat{\psi}$  and

$$\widehat{\psi} \circ (\widehat{T}^{(1)} \circ \widehat{T}^{(2)})_{\widehat{F}_2} = (\widehat{\psi} \circ \widehat{T}_{\widehat{F}_2}^{(1)}) \circ \widehat{T}_{\widehat{F}_2}^{(2)}, \quad \forall \widehat{T}^{(1)}, \widehat{T}^{(2)} \in \widehat{GT}, \quad \widehat{\psi} \in \text{Hom}(\widehat{F}_2, S_d),$$

we conclude that the formulas in (2.34) indeed define a cofunctor  $\mathcal{A}$  from the transformation groupoid  $\widehat{GT}_{\text{NFI}}$  to the category **Dessin**.  $\square$

Using the Ihara embedding  $G_{\mathbb{Q}} \hookrightarrow \widehat{GT}$  and the action of  $\widehat{GT}$  on  $\text{NFI}_{PB_4}(B_4)$ , we get the action of  $G_{\mathbb{Q}}$  on  $\text{NFI}_{PB_4}(B_4)$ . We denote by  $G_{\mathbb{Q}, \text{NFI}}$  the corresponding transformation groupoid and by  $\text{Ih}$  the natural functor

$$\text{Ih} : G_{\mathbb{Q}, \text{NFI}} \rightarrow \widehat{GT}_{\text{NFI}} \quad (2.36)$$

coming from the Ihara embedding.

Composing  $\text{Ih}$  with the cofunctor  $\mathcal{A} : \widehat{GT}_{\text{NFI}} \rightarrow \text{Dessin}$ , we get a cofunctor from the groupoid  $G_{\mathbb{Q}, \text{NFI}}$  to the category **Dessin**. We denote this cofunctor by  $\mathcal{A}^{\mathbb{Q}}$ :

$$\mathcal{A}^{\mathbb{Q}} := \mathcal{A} \circ \text{Ih} : G_{\mathbb{Q}, \text{NFI}} \rightarrow \text{Dessin}. \quad (2.37)$$



### 3 The action of GT-shadows on child's drawings

It is convenient to introduce the action of GT-shadows on child's drawings as a functor from the groupoid GTSh to the category Dessin:

**Theorem 3.1** *Let  $\mathbf{N}^{(1)}, \mathbf{N}^{(2)} \in \mathbf{NFI}_{\mathbb{P}B_4}(B_4)$ ,  $[m, f] \in \text{GTSh}(\mathbf{N}^{(2)}, \mathbf{N}^{(1)})$  and  $[\psi] \in \text{Dessin}(\mathbf{N}^{(1)})$  be a child's drawing represented by a homomorphism  $\psi : \mathbf{F}_2 \rightarrow S_d$ . Let  $\psi^{(m,f)} : \mathbf{F}_2 \rightarrow S_d$  be the homomorphism defined by the formulas*

$$\psi^{(m,f)}(x) := \psi(x^{2m+1}), \quad \psi^{(m,f)}(y) := \psi(f^{-1}y^{2m+1}f). \quad (3.1)$$

Then

- $\psi^{(m,f)}$  does not depend on the choice of the pair  $(m, f)$  representing the GT-shadow  $[m, f]$ ;
- $\psi^{(m,f)}$  represents a child's drawing of degree  $d$  subordinate to  $\mathbf{N}^{(2)}$ .

Moreover, the formulas

$$\mathcal{A}^{sh}(\mathbf{N}) := \mathbf{N}, \quad \mathcal{A}^{sh}([m, f])([\psi]) := [\psi^{(m,f)}] \quad (3.2)$$

define a cofunctor  $\mathcal{A}^{sh} : \text{GTSh} \rightarrow \text{Dessin}$ .

**Proof.** Since  $\mathbf{N}_{\mathbf{F}_2}^{(1)} \leq \ker(\psi)$  the formula

$$\tilde{\psi}(w\mathbf{N}_{\mathbf{F}_2}^{(1)}) := \psi(w)$$

defines a group homomorphism  $\tilde{\psi} : \mathbf{F}_2/\mathbf{N}_{\mathbf{F}_2}^{(1)} \rightarrow S_d$  and  $\tilde{\psi}(\mathbf{F}_2/\mathbf{N}_{\mathbf{F}_2}^{(1)}) = \psi(\mathbf{F}_2)$ . In particular, the subgroup  $\tilde{\psi}(\mathbf{F}_2/\mathbf{N}_{\mathbf{F}_2}^{(1)})$  is transitive.

Due to (2.14),

$$\psi^{(m,f)} = \tilde{\psi} \circ T_{m,f}^{\mathbf{F}_2}. \quad (3.3)$$

Since the homomorphism  $T_{m,f}^{\mathbf{F}_2} : \mathbf{F}_2 \rightarrow \mathbf{F}_2/\mathbf{N}_{\mathbf{F}_2}^{(1)}$  does not depend on the choice of the representative  $(m, f)$  of the GT-shadow  $[m, f]$ , the first statement of the proposition is proved.

Since  $T_{m,f}^{\mathbf{F}_2} : \mathbf{F}_2 \rightarrow \mathbf{F}_2/\mathbf{N}_{\mathbf{F}_2}^{(1)}$  is onto  $\psi^{(m,f)}(\mathbf{F}_2) = \tilde{\psi}(\mathbf{F}_2/\mathbf{N}_{\mathbf{F}_2}^{(1)})$ . Hence  $\psi^{(m,f)}(\mathbf{F}_2)$  is a transitive subgroup of  $S_d$ .

We know that  $\ker(T_{m,f}^{\mathbf{F}_2}) = \mathbf{N}_{\mathbf{F}_2}^{(2)}$ . Hence (3.3) implies that

$$\mathbf{N}_{\mathbf{F}_2}^{(2)} \leq \ker(\psi^{(m,f)}).$$

We proved the second statement of the proposition.

It is easy to see that  $[\psi^{(m,f)}]$  does not depend on the representative  $\psi$  of the child's drawing  $[\psi]$ . Thus the formula

$$\mathcal{A}^{sh}([m, f])([\psi]) := [\psi^{(m,f)}]$$

defines a map  $\text{GTSh}(\mathbf{N}^{(2)}, \mathbf{N}^{(1)}) \rightarrow \text{Dessin}(\mathbf{N}^{(1)}, \mathbf{N}^{(2)})$ .

It is also easy to see that, for every  $[\psi] \in \text{Dessin}(\mathbf{N}^{(1)})$ ,  $[\psi^{(0,1_{\mathbf{F}_2})}] = [\psi]$ .

Thus it remains to show that, for all  $[m_1, f_1] \in \text{GTSh}(\mathbf{N}^{(2)}, \mathbf{N}^{(1)})$ ,  $[m_2, f_2] \in \text{GT}(\mathbf{N}^{(2)})$ , and  $[\psi] \in \text{Dessin}(\mathbf{N}^{(1)})$ ,

$$\mathcal{A}^{sh}([m_1, f_1] \circ [m_2, f_2])([\psi]) = [(\psi^{(m_1, f_1)})^{(m_2, f_2)}]. \quad (3.4)$$

Equation (3.4) can be verified directly using (2.21).

Here is another way to prove (3.4). Let  $[m, f] := [m_1, f_1] \circ [m_2, f_2]$ . Since the composition of GT-shadows is defined via the identification of elements of  $\widehat{\text{GT}}(\mathbf{N})$  with isomorphisms (2.10), we have

$$T_{m,f} = T_{m_1,f_1}^{\text{isom}} \circ T_{m_2,f_2}.$$

Hence

$$T_{m,f}^{\mathbf{F}_2} = T_{m_1,f_1}^{\mathbf{F}_2, \text{isom}} \circ T_{m_2,f_2}^{\mathbf{F}_2}. \quad (3.5)$$

Since  $\psi^{(m,f)} = \tilde{\psi} \circ T_{m,f}^{\mathbf{F}_2}$ , identity (3.5) implies that

$$\psi^{(m,f)} = (\tilde{\psi} \circ T_{m_1,f_1}^{\mathbf{F}_2, \text{isom}}) \circ T_{m_2,f_2}^{\mathbf{F}_2}. \quad (3.6)$$

Since  $\tilde{\psi} \circ T_{m_1,f_1}^{\mathbf{F}_2, \text{isom}} \circ \mathcal{P}_{\mathbf{N}_{\mathbf{F}_2}}^{(2)} = \psi^{(m_1,f_1)}$ , identity (3.6) implies the desired equation in (3.4).  $\square$

The action of GT-shadows on child's drawings is compatible with the action of  $\widehat{\text{GT}}$  in the following sense:

**Theorem 3.2** *Let  $\mathcal{PR}$  be the natural functor from  $\widehat{\text{GT}}_{\text{NFI}}$  to  $\text{GTSh}^\heartsuit$  introduced in the proof of Proposition 2.2 and let  $\mathcal{A}$  be the cofunctor from  $\widehat{\text{GT}}_{\text{NFI}}$  to  $\text{Dessin}$  defined in Proposition 2.12. The diagram of (co)functors*

$$\begin{array}{ccc} \widehat{\text{GT}}_{\text{NFI}} & \xrightarrow{\mathcal{PR}} & \text{GTSh}^\heartsuit \\ & \searrow \mathcal{A} & \swarrow \mathcal{A}^{sh} \\ & \text{Dessin} & \end{array} \quad (3.7)$$

*commutes “on the nose”.*

**Proof.** Since all three functors operate as the identity map on the level of objects, we need to show that, for every  $\mathbf{N} \in \text{NFI}_{\text{PB}_4}(\mathbf{B}_4)$ ,  $\hat{T} \in \widehat{\text{GT}}$  and  $[\psi] \in \text{Dessin}(\mathbf{N})$  we have

$$\hat{\psi} \circ \hat{T}|_{\mathbf{F}_2} = \tilde{\psi} \circ T_{\mathbf{N}}^{\mathbf{F}_2}, \quad (3.8)$$

where  $\hat{\psi} : \hat{\mathbf{F}}_2 \rightarrow S_d$  is the continuous group homomorphism that extends  $\psi : \mathbf{F}_2 \rightarrow S_d$ , and  $\tilde{\psi}$  is the group homomorphism  $\mathbf{F}_2/\mathbf{N}_{\mathbf{F}_2} \rightarrow S_d$  defined by the formula

$$\tilde{\psi}(w \mathbf{N}_{\mathbf{F}_2}) := \psi(w).$$

Due to the relation defining the GT-shadow  $T_N : \text{PaB}^{\leq 4} \rightarrow \text{PaB}^{\leq 4} / \sim_N$  in terms of  $\hat{T} \in \widehat{\text{GT}}$  (see equation (2.17)), the diagram

$$\begin{array}{ccc} \hat{\mathbf{F}}_2 & \xrightarrow{\hat{T}|_{\hat{\mathbf{F}}_2}} & \hat{\mathbf{F}}_2 \\ \uparrow & \searrow \hat{\mathcal{P}}_{\mathbf{N}_{\mathbf{F}_2}} & \downarrow \\ \mathbf{F}_2 & \xrightarrow{T_{\mathbf{N}}^{\mathbf{F}_2}} & \mathbf{F}_2/\mathbf{N}_{\mathbf{F}_2} \end{array} \quad (3.9)$$

commutes.

Using equation (2.29) from the proof of Proposition 2.5 and the relation  $N_{F_2} \leq \ker(\psi)$ , it is easy to see that the diagram

$$\begin{array}{ccc} \widehat{F}_2 & & \\ \widehat{\mathcal{P}}_{N_{F_2}} \downarrow & \searrow \hat{\psi} & \\ F_2/N_{F_2} & \xrightarrow{\tilde{\psi}} & S_d \end{array} \quad (3.10)$$

commutes.

Putting (3.9) and (3.10) together, we get the following commutative diagram

$$\begin{array}{ccccc} \widehat{F}_2 & \xrightarrow{\hat{T}|_{\widehat{F}_2}} & \widehat{F}_2 & & \\ & & \downarrow \widehat{\mathcal{P}}_{N_{F_2}} & \searrow \hat{\psi} & \\ F_2 & \xrightarrow{T_N^{F_2}} & F_2/N_{F_2} & \xrightarrow{\tilde{\psi}} & S_d \end{array}$$

Thus equation (3.8) is proved and Theorem 3.2 follows.  $\square$

Let us prove that

**Proposition 3.3** *For every child's drawing  $\mathcal{D}$ , there exists  $N \in \text{NFI}_{\text{PB}_4}(\text{B}_4)$  that dominates  $\mathcal{D}$ , i.e.  $\mathcal{D} \in \text{Dessin}(N)$ . In fact, for every child's drawing  $\mathcal{D}$ , there are infinitely many elements  $K \in \text{NFI}_{\text{PB}_4}(\text{B}_4)$  such that  $\mathcal{D} \in \text{Dessin}(K)$ .*

**Proof.** Let  $\psi$  be a group homomorphism from  $F_2$  to  $S_d$  that represents a child's drawing  $\mathcal{D}$ . The following formulas

$$\begin{aligned} \varphi(x_{12}) &:= \psi(x), & \varphi(x_{23}) &:= \psi(y), & \varphi(x_{13}) &:= \psi(x^{-1}y^{-1}), \\ \varphi(x_{14}) &= \varphi(x_{24}) = \varphi(x_{34}) &:= 1_{S_d} \end{aligned} \quad (3.11)$$

define a group homomorphism  $\varphi : \text{PB}_4 \rightarrow S_d$ .

Indeed, since  $x_{13} = x_{12}^{-1}x_{23}^{-1}c$ , the first three equations in (3.11) define a group homomorphism from  $\text{PB}_3$  to  $S_d$ . Since the elements  $x_{14}, x_{24}, x_{34}$  generate a free subgroup of  $\text{PB}_4$  and  $\text{PB}_4$  is isomorphic to the semi-direct product  $\text{PB}_3 \ltimes \langle x_{14}, x_{24}, x_{34} \rangle$ , the formulas in (3.11) define a group homomorphism  $\varphi : \text{PB}_4 \rightarrow S_d$ .

It is clear that

$$\ker(\varphi|_{\text{PB}_3}) \cap F_2 = \ker(\psi). \quad (3.12)$$

Unfortunately, in general,  $\ker(\varphi)$  is not normal in  $\text{B}_4$ .

We denote by  $N$  the normal core of  $\ker(\varphi)$  in  $\text{B}_4$ . Since  $\ker(\varphi)$  has finite index in  $\text{B}_4$ , so does  $N$ . In addition,  $N \leq \text{PB}_4$ . Thus  $N \in \text{NFI}_{\text{PB}_4}(\text{B}_4)$ .

Using the definition of  $N_{\text{PB}_3}$  (see equation (2.4) in [7, Section 2.2]) and the inclusion  $N \leq \ker(\varphi)$ , we conclude that

$$N_{\text{PB}_3} \leq \ker(\varphi|_{\text{PB}_3}).$$

Combining this observation with identity (3.12), we conclude that

$$\mathbf{N}_{\mathbf{F}_2} \leq \ker(\psi).$$

Thus  $\mathbf{N}$  dominates the child's drawing  $[\psi]$ .

To prove the second statement, let us show that, for every  $\mathbf{K} \in \mathbf{NFI}_{\mathbf{PB}_4}(\mathbf{B}_4)$  that dominates  $[\psi]$ , there exists  $\tilde{\mathbf{K}} \in \mathbf{NFI}_{\mathbf{PB}_4}(\mathbf{B}_4)$  such that

- $\tilde{\mathbf{K}}$  dominates  $[\psi]$  and
- $\tilde{\mathbf{K}}$  is properly contained in  $\mathbf{K}$ .

Since  $\mathbf{K}$  is non-trivial, there exists a non-identity element  $w \in \mathbf{K}$ . Since  $\mathbf{B}_4$  is residually finite, there exists  $\mathbf{H} \in \mathbf{NFI}(\mathbf{B}_4)$  such that  $w \notin \mathbf{H}$ . We set

$$\tilde{\mathbf{K}} := \mathbf{H} \cap \mathbf{K}.$$

It is clear that  $\tilde{\mathbf{K}} \in \mathbf{NFI}_{\mathbf{PB}_4}(\mathbf{B}_4)$  and  $\tilde{\mathbf{K}}$  is properly contained in  $\mathbf{K}$ .

Since  $\tilde{\mathbf{K}} \leq \mathbf{K}$ ,  $\tilde{\mathbf{K}}$  also dominates  $[\psi]$ . □

**Remark 3.4** Although paper [17] uses a more restrictive assumption on the analogue of  $\mathbf{N} \in \mathbf{NFI}_{\mathbf{PB}_4}(\mathbf{B}_4)$ , a discussion that is very similar to the proof of Proposition 3.3 can be found on page 225 of [17].

**Remark 3.5** The relation of sub-ordinance from Definition 2.11 has a loose analogy with a natural relationship between child's drawings and finite Galois extensions of  $\mathbb{Q}$ . One can say that a child's drawing  $\mathcal{D}$  is **subordinate** to a finite Galois extension  $E \supset \mathbb{Q}$  if  $\mathcal{D}$  can be represented by a Belyi pair  $(X, \gamma)$  defined over an intermediate field of the extension  $E \supset \mathbb{Q}$ . The direct analogue of the first statement of Proposition 3.3 for this relation is obvious because every finite field extension  $K \supset \mathbb{Q}$  is contained in a finite Galois extension of  $\mathbb{Q}$ . The direct analogue of the second statement of Proposition 3.3 is also obvious since the field extension  $\overline{\mathbb{Q}} \supset \mathbb{Q}$  is infinite.

It is relatively easy to see that, if a child's drawing  $[\psi]$  is Galois, then so is  $[\psi]^g$  for every  $g \in G_{\mathbb{Q}}$ . The corresponding version of this statement for the action of GT-shadows requires a proof:

**Proposition 3.6** *Let  $\mathbf{N} \in \mathbf{NFI}_{\mathbf{PB}_4}(\mathbf{B}_4)$ ,  $[\psi] \in \mathbf{Dessin}(\mathbf{N})$  and  $[m, f] \in \mathbf{GT}(\mathbf{N})$ . If the child's drawing  $[\psi]$  is Galois then so is  $[\psi]^{[m, f]}$ .*

**Proof.** Let  $\psi : \mathbf{F}_2 \rightarrow S_d$  be a homomorphism that represents the child's drawing  $[\psi]$ . Note that  $[\psi]$  is Galois if and only if the order of the subgroup  $\psi(\mathbf{F}_2) \leq S_d$  is  $d$ . (This is equivalent to the statement that the stabilizer of 1 in  $\mathbf{F}_2$  coincides with its normal core.)

Let us denote by  $\tilde{\psi}$  the homomorphism from  $\mathbf{F}_2/\mathbf{N}_{\mathbf{F}_2} \rightarrow S_d$  defined by the formula

$$\tilde{\psi}(w\mathbf{N}_{\mathbf{F}_2}) = \psi(w).$$

The child's drawing  $[\psi]^{[m, f]}$  is represented by the homomorphism

$$\tilde{\psi} \circ T_{m, f}^{\mathbf{F}_2} : \mathbf{F}_2 \rightarrow S_d.$$

Since  $T_{m,f}^{F_2} : F_2 \rightarrow F_2/N_{F_2}$  is onto (see Proposition 2.1) and  $\tilde{\psi}(F_2/N_{F_2}) = \psi(F_2)$ ,

$$\tilde{\psi} \circ T_{m,f}^{F_2}(F_2) = \psi(F_2). \quad (3.13)$$

Thus the order of the subgroup  $\tilde{\psi} \circ T_{m,f}^{F_2}(F_2) \leq S_d$  also coincides with the degree  $d$ . Hence the child's drawing  $[\psi]^{[m,f]}$  is Galois.  $\square$

The following proposition shows that there is a large supply of Galois child's drawings whose  $\text{GTSh}^\heartsuit$ -orbits are singletons:

**Proposition 3.7** *For every  $N \in \text{NFI}_{\text{PB}_4}(B_4)$ , the Galois child's drawing  $\mathcal{D}_N$  represented by  $N_{F_2}$  is subordinate to  $N$ . Moreover, if  $N$  is isolated, then the orbit  $\text{GT}^\heartsuit(N)(\mathcal{D}_N)$  is a singleton.*

**Proof.** We set  $d := |F_2 : N_{F_2}|$  and choose a bijection between the set  $F_2/N_{F_2}$  of left cosets and  $\{1, 2, \dots, d\}$ . Then the standard (left) action of  $F_2$  on  $F_2/N_{F_2}$  gives us a group homomorphism

$$\psi : F_2 \rightarrow S_d$$

which represents the child's drawing  $\mathcal{D}_N$ .

Since  $N_{F_2}$  is normal in  $F_2$ ,  $\ker(\psi) = N_{F_2}$ . Thus  $\mathcal{D}_N$  is subordinate to  $N$ .

For the second statement, we assume that  $N$  is isolated.

Let  $\tilde{\psi}$  be the homomorphism  $F_2/N_{F_2} \rightarrow S_d$  defined by the formula

$$\tilde{\psi}(wN_{F_2}) := \psi(w)$$

and let  $[m, f] \in \text{GT}^\heartsuit(N)$ .

The (new) child's drawing  $\mathcal{D}_N^{[m,f]}$  is represented by the homomorphism

$$\psi^{(m,f)} := \tilde{\psi} \circ T_{m,f}^{F_2} : F_2 \rightarrow S_d.$$

Since  $\ker(\psi) = N_{F_2}$ , the homomorphism  $\tilde{\psi}$  is injective.

Since  $N$  is isolated, the kernel of the morphism  $T_{m,f} : \text{PaB}^{\leq 4} \rightarrow \text{PaB}^{\leq 4} / \sim_N$  is the compatible equivalence relation corresponding to  $N$ . Hence  $\ker(T_{m,f}^{\text{PB}_3}) = N_{\text{PB}_3}$  and therefore  $\ker(T_{m,f}^{F_2}) = N_{F_2}$ .

Combining this observation with the injectivity of  $\tilde{\psi}$ , we conclude that

$$\ker(\psi^{(m,f)}) = N_{F_2}.$$

Due to Proposition 3.6, the child's drawing  $[\psi^{(m,f)}]$  is Galois. Hence  $[\psi^{(m,f)}]$  is represented by  $\ker(\psi^{(m,f)}) = N_{F_2}$ .

The proposition is proved.  $\square$

### 3.1 The hierarchy of orbits

Combining Theorem 3.2 with the definition of the cofunctor  $\mathcal{A}^\mathbb{Q}$  (see (2.37)), we get the following statement:

**Corollary 3.8** *The diagram of (co)functors*

$$\begin{array}{ccccc}
 G_{\mathbb{Q}, \text{NFI}} & \xrightarrow{\text{Ih}} & \widehat{\text{GT}}_{\text{NFI}} & \xrightarrow{\mathcal{PR}} & \text{GTSh}^\heartsuit \\
 & \searrow \mathcal{A}^{\mathbb{Q}} & \downarrow \mathcal{A} & \swarrow \mathcal{A}^{sh} & \\
 & & \text{Dessin} & & 
 \end{array}
 \tag{3.14}$$

commutes “on the nose”.  $\square$

Let  $K, N \in \text{NFI}_{\text{PB}_4}(\text{B}_4)$  and  $K \leq N$ . Recall [7, Section 3.2] that, if a pair  $(m, f) \in \mathbb{Z} \times \text{F}_2$  represents a charming GT-shadow with the target  $K$ , then the same pair  $(m, f)$  also represents a charming GT-shadow with the target  $N$ . Hence, for every pair  $K, N \in \text{NFI}_{\text{PB}_4}(\text{B}_4)$  with  $K \leq N$ , we have a natural map

$$P_{K,N} : \text{GT}^\heartsuit(K) \rightarrow \text{GT}^\heartsuit(N). \tag{3.15}$$

Let  $\psi$  be a homomorphism  $\text{F}_2 \rightarrow S_d$  that represents a child’s drawing subordinate to  $N$ . Since  $K \leq N$ ,  $[\psi]$  is also subordinate to  $K$ .

Let  $[m, f] \in \text{GT}^\heartsuit(K)$ . Since  $[\psi]^{[m,f]}$  depends only on the residue class of  $m$  modulo  $\text{lcm}(\text{ord}(x \ker(\psi)), \text{ord}(y \ker(\psi)))$  and the coset  $f \ker(\psi) \in \text{F}_2 / \ker(\psi)$ , we have

$$[\psi]^{[m,f]} = [\psi]^{P_{K,N}([m,f])}. \tag{3.16}$$

Combining this observation with Corollary (3.8) we get the following statement:

**Corollary 3.9** *Let  $K, N$  be elements of  $\text{NFI}_{\text{PB}_4}(\text{B}_4)$  that dominate a child’s drawing  $\mathcal{D}$  and  $K \leq N$ . Then we have the following hierarchy of orbits:*

$$\text{GT}^\heartsuit(N)(\mathcal{D}) \supset \text{GT}^\heartsuit(K)(\mathcal{D}) \supset \widehat{\text{GT}}(\mathcal{D}) \supset G_{\mathbb{Q}}(\mathcal{D}). \tag{3.17}$$

$\square$

**Remark 3.10** There may be examples of pairs  $K, N \in \text{NFI}_{\text{PB}_4}(\text{B}_4)$  with  $K \leq N$  such that the natural map  $P_{K,N}$  (3.15) is not onto. So, in principle, there may be examples of  $K, N \in \text{NFI}_{\text{PB}_4}(\text{B}_4)$  with  $K \leq N$  and  $\mathcal{D} \in \text{Dessin}(N)$  for which the inclusion

$$\text{GT}^\heartsuit(N)(\mathcal{D}) \supset \text{GT}^\heartsuit(K)(\mathcal{D})$$

is proper. At the time of writing, the author did not find any examples of this kind. In fact, for all examples in which both orbits  $\text{GT}^\heartsuit(N)(\mathcal{D})$  and  $G_{\mathbb{Q}}(\mathcal{D})$  can be computed, we have  $\text{GT}^\heartsuit(N)(\mathcal{D}) = G_{\mathbb{Q}}(\mathcal{D})$ .

**Remark 3.11** Since the orbit  $\widehat{\text{GT}}(\mathcal{D})$  is finite and  $\widehat{\text{GT}}$  is the limit of the functor that sends  $N \in \text{NFI}_{\text{PB}_4}^{\text{isolated}}(\text{B}_4)$  to the finite group  $\text{GT}^\heartsuit(N)$  (see [7, Theorem 3.8]), for every child’s drawing  $\mathcal{D}$ , there exists  $N \in \text{NFI}_{\text{PB}_4}^{\text{isolated}}(\text{B}_4)$  such that

- $\mathcal{D} \in \text{Dessin}(N)$ , i.e.  $N$  dominates  $\mathcal{D}$ , and
- $\widehat{\text{GT}}(\mathcal{D}) = \text{GT}^\heartsuit(N)(\mathcal{D})$ .

See also, the Corollary and the Problem on page 227 of [17].

Let  $\mathcal{D}$  be a child's drawing and  $H_{\mathcal{D}}$  be the stabilizer of  $\mathcal{D}$  in  $G_{\mathbb{Q}}$ . Recall that the **field of moduli** of a child's drawing  $\mathcal{D}$  is the (finite) extension  $M_{\mathcal{D}} \supset \mathbb{Q}$  corresponding to the (closed) subgroup  $H_{\mathcal{D}} \leq G_{\mathbb{Q}}$  (via the Galois correspondence). Since  $[M_{\mathcal{D}} : \mathbb{Q}] = |G_{\mathbb{Q}} : H_{\mathcal{D}}|$ , the orbit-stabilizer theorem and Corollary 3.9 imply the following version of [17, Proposition 14]:

**Corollary 3.12** *For every child's drawing  $\mathcal{D}$  and every  $\mathbf{N} \in \text{NFI}_{\text{PB}_4}(\text{B}_4)$  that dominates  $\mathcal{D}$ , we have*

$$[M_{\mathcal{D}} : \mathbb{Q}] \leq |\text{GT}^{\heartsuit}(\mathbf{N})(\mathcal{D})|. \quad (3.18)$$

Let us also recall [4, Proposition 2.5], [29, Corollary on page 2] that, if the child's drawing  $\mathcal{D}$  is Galois then it admits a Belyi pair  $(X, \gamma)$  defined over its field of moduli  $M_{\mathcal{D}}$ . Combining this observation with Corollary (3.9), we conclude that if  $\mathcal{D}$  is a Galois child's drawing and  $\text{GT}^{\heartsuit}(\mathbf{N})(\mathcal{D})$  is a singleton (for some  $\mathbf{N}$  that dominates  $\mathcal{D}$ ), then  $\mathcal{D}$  admits a Belyi pair  $(X, \gamma)$  defined over  $\mathbb{Q}$ .

A large supply of examples of such Galois child's drawings comes from isolated elements of  $\text{NFI}_{\text{PB}_4}(\text{B}_4)$ . Indeed, combining [4, Proposition 2.5], [29, Corollary on page 2] with Proposition 3.7 and Corollary (3.9), we get the following statement:

**Corollary 3.13** *For every isolated element  $\mathbf{N} \in \text{NFI}_{\text{PB}_4}(\text{B}_4)$ , the child's drawing represented by the subgroup  $\mathbf{N}_{\mathbb{F}_2} \trianglelefteq \mathbb{F}_2$  admits a Belyi pair defined over  $\mathbb{Q}$ .  $\square$*

**Remark 3.14** Section 4 of [7] presents the basic information about 35 selected elements

$$\mathbf{N}^{(0)}, \mathbf{N}^{(1)}, \dots, \mathbf{N}^{(34)} \quad (3.19)$$

of the poset  $\text{NFI}_{\text{PB}_4}(\text{B}_4)$ . More information about these elements can be found in [6]. According to Table 1 (on page 40) in [7], 27 of these 35 elements are isolated. Due to Corollary 3.13, for every isolated element  $\mathbf{N}$  in list (3.19), the child's drawing represented by  $\mathbf{N}_{\mathbb{F}_2}$  admits a Belyi pair defined over  $\mathbb{Q}$ . Moreover, for many of these isolated elements, the degrees of the corresponding child's drawings are quite large. For example, the degree of the child's drawing represented by  $\mathbf{N}_{\mathbb{F}_2}^{(34)}$  is  $20575296 = 2^6 \cdot 3^8 \cdot 7^2$ .

**Remark 3.15** Proposition 3.7, Remark 3.14 and examples considered in Section 5 indicate that the size of the orbit  $\text{GT}^{\heartsuit}(\mathbf{N})(\mathcal{D})$  (for some  $\mathbf{N}$  dominating a child's drawing  $\mathcal{D}$ ) may be significantly smaller than the number of the child's drawings with the same passport as  $\mathcal{D}$ . Thus the bound on the degree of the field of moduli (3.18) has more practical value than the one given in [30, Proposition 7.1].

### 3.2 The monodromy group and the passport are invariant with respect to the action of GT-shadows

It is easy to see that the degree and the monodromy group of a child's drawing are invariant with respect to the action of GT-shadows. (See, for example, (3.13) in the proof of Proposition 3.6).

Let us prove that the passport of a child's drawing is invariant with respect to the action of charming GT-shadows.

**Theorem 3.16** *Let  $N \in \text{NFI}_{\text{PB}_4}(\text{B}_4)$  and  $\psi$  be a homomorphism  $\text{F}_2 \rightarrow S_d$  that represents  $[\psi] \in \text{Dessin}(N)$ . Then, for every charming GT-shadow  $[m, f]$  in  $\text{GT}(N)$ , the child's drawing  $[\psi]^{[m, f]}$  has the same passport as  $[\psi]$ .*

**Proof.** We set

$$x := x_{12}, \quad y := x_{23}, \quad z := (xy)^{-1}, \quad w := (yx)^{-1},$$

and

$$g_x := \psi(x), \quad g_y := \psi(y), \quad g_z := \psi(z).$$

Due to (3.1), the child's drawing  $[\psi]^{[m, f]}$  is represented by the triple:

$$(g_x^{2m+1}, \psi(f)^{-1} g_y^{2m+1} \psi(f), \psi(f)^{-1} g_y^{-2m-1} \psi(f) g_x^{-2m-1}). \quad (3.20)$$

The passport of the child's drawing  $[\psi]$  is the triple of partitions  $(\text{ct}(g_x), \text{ct}(g_y), \text{ct}(g_z))$ .

Thus our goal is to show that

$$\text{ct}(g_x^{2m+1}) = \text{ct}(g_x), \quad \text{ct}(\psi(f)^{-1} g_y^{2m+1} \psi(f)) = \text{ct}(g_y), \quad (3.21)$$

and

$$\text{ct}(\psi(f)^{-1} g_y^{-2m-1} \psi(f) g_x^{-2m-1}) = \text{ct}(g_z) \quad (3.22)$$

It is clear that the second equation in (3.21) is equivalent to  $\text{ct}(g_y^{2m+1}) = \text{ct}(g_y)$ . Thus equations (3.21) are consequences of the following simple fact about permutations: if

$$\gcd(q, \text{ord}(h)) = 1,$$

then the permutations  $h^q$  and  $h$  have the same cycle structure.

The integer  $2m + 1$  is coprime with the orders of  $g_x := \psi(x)$  and  $g_y := \psi(y)$  because  $2m + 1$  is coprime with the orders of  $x_{12}\text{N}_{\text{F}_2}$  and  $x_{23}\text{N}_{\text{F}_2}$ .

The proof of equation (3.22) requires more work.

Since the GT-shadow  $[m, f]$  is charming, we may assume, without loss of generality, that  $f \in [\text{F}_2, \text{F}_2]$ . Hence Proposition A.3 from Appendix A implies that the pair  $(m, f)$  satisfies relations (A.9) and (A.10).

Conjugating (A.10) by  $x^{-m}$ , we get

$$f(z, x) z^m f(y, z) y^m f(x, y) x^m \in \text{N}_{\text{F}_2}. \quad (3.23)$$

Furthermore, conjugating (3.23) by  $\theta := \sigma_1 \sigma_2 \sigma_1$ , we get

$$f(w, y) w^m f(x, w) x^m f(y, x) y^m \in \text{N}_{\text{F}_2}. \quad (3.24)$$

Equation (A.10) implies that

$$y^m f(x, y) \text{N}_{\text{F}_2} = f(y, z)^{-1} z^{-m} f(z, x)^{-1} x^{-m} \text{N}_{\text{F}_2} \quad (3.25)$$

Equation (3.24) implies that

$$f(y, x) y^m \text{N}_{\text{F}_2} = x^{-m} f(x, w)^{-1} w^{-m} f(w, y)^{-1} \text{N}_{\text{F}_2} \quad (3.26)$$

To prove equation (3.22), we need to show that the permutations

$$\psi(xy) \quad \text{and} \quad \psi(x^{2m+1} f^{-1} y^{2m+1} f) \quad (3.27)$$



have the same cycle structure.

We have

$$x^{2m+1}f^{-1}y^{2m+1}f \mathbf{N}_{\mathbf{F}_2} = x^{2m+1}f(y, x)y^{2m+1}f(x, y) \mathbf{N}_{\mathbf{F}_2} = x^{2m+1}(f(y, x)y^m)y(y^mf(x, y)) \mathbf{N}_{\mathbf{F}_2}.$$

Applying (3.25) to  $y^mf(x, y)$  and (3.26) to  $f(y, x)y^m$  we get<sup>9</sup>

$$\begin{aligned} x^{2m+1}f^{-1}y^{2m+1}f \mathbf{N}_{\mathbf{F}_2} &= x^{2m+1}x^{-m}f(x, w)^{-1}w^{-m}f(w, y)^{-1}yf(y, z)^{-1}z^{-m}f(z, x)^{-1}x^{-m} \mathbf{N}_{\mathbf{F}_2} \\ &\sim xf(x, w)^{-1}w^{-m}f(w, y)^{-1}yf(y, z)^{-1}z^{-m}f(z, x)^{-1} \mathbf{N}_{\mathbf{F}_2} \\ &= xf(w, x)w^{-m}f(y, w)yf(z, y)z^{-m}f(x, z) \mathbf{N}_{\mathbf{F}_2}. \end{aligned}$$

Applying the obvious relations  $xwx^{-1} = z$ ,  $zyz^{-1} = w$  to  $xf(w, x)w^{-m}$ ,  $f(y, w)y$ , respectively, and using (A.4) and (A.5) from Appendix A, we see that, up to conjugation in  $\mathbf{F}_2/\mathbf{N}_{\mathbf{F}_2}$ ,

$$\begin{aligned} x^{2m+1}f^{-1}y^{2m+1}f \mathbf{N}_{\mathbf{F}_2} &\sim f(z, x)z^{-m}xyf(y, z)f(z, y)z^{-m}f(x, z) \mathbf{N}_{\mathbf{F}_2} = \\ &f(z, x)z^{-m}xy z^{-m}f(x, z) \mathbf{N}_{\mathbf{F}_2} \sim (xy)^{2m+1} \mathbf{N}_{\mathbf{F}_2}. \end{aligned}$$

Hence the permutations  $\psi(x^{2m+1}f^{-1}y^{2m+1}f)$  and  $\psi(xy)^{2m+1}$  have the same cycle structure.

To prove that the permutations in (3.27) have the same cycle structure, it remains to show that the cycle structure of  $\psi(xy)^{2m+1}$  coincides with the cycle structure of  $\psi(xy)$ . It suffices to show that the order of  $xy\mathbf{N}_{\mathbf{PB}_3}$  divides  $N_{\text{ord}}$ .

Since  $xy\mathbf{N}_{\mathbf{PB}_3} = z^{-1}\mathbf{N}_{\mathbf{PB}_3}$ , we need to show that the order of  $z\mathbf{N}_{\mathbf{PB}_3}$  divides  $N_{\text{ord}}$ .

Since  $x_{23}^{N_{\text{ord}}} \in \mathbf{N}_{\mathbf{PB}_3}$  and  $\sigma_1 x_{23} \sigma_1^{-1} = c x_{23}^{-1} x_{12}^{-1} = cz$ , we have

$$c^{N_{\text{ord}}} z^{N_{\text{ord}}} = (cz)^{N_{\text{ord}}} \in \mathbf{N}_{\mathbf{PB}_3}.$$

Combining this observation with  $c^{N_{\text{ord}}} \in \mathbf{N}_{\mathbf{PB}_3}$ , we conclude that  $z^{N_{\text{ord}}}\mathbf{N}_{\mathbf{PB}_3} = 1$ . Hence the order of  $z\mathbf{N}_{\mathbf{PB}_3}$  divides  $N_{\text{ord}}$ .

Since the permutations in (3.27) have the same cycle structure, (3.22) is proved.  $\square$

## 4 Abelian child's drawings

Recall that the monodromy group of a child's drawing  $[\psi]$  of degree  $d$  is defined up to conjugation in  $S_d$ . It is clear that the following definition does not depend on the choice of the representative of the monodromy group of a child's drawing.

**Definition 4.1** *A child's drawing  $\mathcal{D}$  is called **Abelian** if its monodromy group is Abelian.*

For example, for every  $d \geq 1$ , the pair  $((1, 2, \dots, d), (1, 2, \dots, d))$  represents an Abelian child's drawing of degree  $d$ .

Let us prove that

**Proposition 4.2** *Every Abelian child's drawing  $[(g_1, g_2)]$  is Galois. In particular, the order of the monodromy group  $G := \langle g_1, g_2 \rangle$  of  $[(g_1, g_2)]$  coincides with the degree of  $[(g_1, g_2)]$ .*

---

<sup>9</sup>In these calculations,  $\sim$  means "conjugate in  $\mathbf{F}_2/\mathbf{N}_{\mathbf{F}_2}$ ".

**Proof.** Let  $\psi : F_2 \rightarrow S_d$  be the group homomorphism corresponding to  $(g_1, g_2)$ ,  $K := \ker(\psi)$  and  $H$  be the stabilizer of 1 in  $F_2$ . Note that, since the action of  $F_2$  on  $\{1, 2, \dots, d\}$  is transitive,  $d = |G : H|$ .

Since the quotient  $F_2/K \cong G$  is Abelian, every subgroup  $N \leq F_2$  that contains  $K$  is normal in  $F_2$ . In particular,  $H$  is normal in  $F_2$ . Thus the child's drawing  $[\psi]$  is Galois.

Since  $K$  is the normal core of  $H$  in  $F_2$  and  $H$  is normal, we conclude that  $H = K$ . Thus the order of the monodromy group of  $[(g_1, g_2)]$  is  $|G : H| = d$ .  $\square$

Using the basic properties of group actions, we can prove the following useful properties of Abelian child's drawings:

**Proposition 4.3** *Let  $[(g_1, g_2)]$  be an Abelian child's drawing of degree  $d$ . Then*

- a) *For every  $i \in \{1, 2\}$ , the permutation  $g_i$  is a product of disjoint cycles of the same length.*
- b) *If  $g_1$  (resp.  $g_2$ ) is a cycle of length  $d$  then  $g_2 \in \langle g_1 \rangle$  (resp.  $g_1 \in \langle g_2 \rangle$ ).*

**Proof.** As above, we set  $G := \langle g_1, g_2 \rangle$ .

For a), it suffices to prove that  $g_1$  is a product of disjoint cycles of the same length.

Let  $H_1 := \langle g_1 \rangle$  and

$$\bigsqcup_{t=1}^r O_t$$

be the partition of the set  $\{1, 2, \dots, d\}$  into the orbits of the action of  $H_1$ . If  $r = 1$ , i.e.  $g_1$  is a cycle of length  $d$ , then there is nothing to prove. So we consider the case when the number of orbits  $r$  is  $\geq 2$ . Our goal is to prove that all orbits have the same size.

Since  $g_2$  commutes with  $g_1$ , the subgroup  $H_2 := \langle g_2 \rangle$  acts on the set of orbits  $\{O_1, \dots, O_r\}$ . Furthermore, since the permutation group  $G$  is transitive,  $H_2$  acts on  $\{O_1, \dots, O_r\}$  transitively.

Therefore, for two distinct orbits  $O$  and  $\tilde{O}$ , there exists  $k \in \mathbb{Z}_{\geq 1}$  such that  $g_2^k(i) \in \tilde{O}$  for every  $i \in O$ , i.e.  $g_2^k|_O$  is a map

$$O \rightarrow \tilde{O}. \tag{4.1}$$

Since  $g_2^{-k}|_{\tilde{O}}$  is the inverse of (4.1), we conclude that  $O$  and  $\tilde{O}$  have the same size. Thus the first statement is proved.

The second statement is a particular case of one of the exercises in [9, Section 4.3].

Here is a proof of this statement.

Without loss of generality, we may assume that  $g_1 = (1, 2, \dots, d)$ .

If  $g_2(1) = 1$  then, for every  $k \in \{2, 3, \dots, d\}$ , we have

$$g_2(k) = g_2(g_1^{k-1}(1)) = g_1^{k-1}(g_2(1)) = g_1^{k-1}(1) = k.$$

Thus, in this case,  $g_2 = \text{id}$ .

If  $g_2(1) = k > 1$  then the permutation  $h := g_1^{1-k}g_2$  satisfies these two properties:

- $h$  commutes with  $g_1$  and
- $h(1) = 1$ .

Using the previous argument, we conclude that  $h = \text{id}$  and hence  $g_2 \in \langle g_1 \rangle$ .  $\square$

**Remark 4.4** Assume that we are in the set-up of Proposition 4.3 and  $\{1, 2, \dots, d\}$  partitions into  $r \geq 2$  orbits of the  $\langle g_1 \rangle$ -action. Although,  $\langle g_2 \rangle$  acts transitively on the set of these orbits, in general,  $g_2$  is not a product of cycles of length  $r$ . For example, the permutations

$$g_1 = (1, 2, 3, 4)(5, 6, 7, 8)(9, 10, 11, 12), \quad g_2 = (1, 6, 12, 3, 8, 10)(2, 7, 9, 4, 5, 11)$$

generate an Abelian and transitive subgroup of  $S_{12}$ .

**Remark 4.5** Since every Abelian child's drawing is Galois, Proposition 4.3 can be deduced from the properties of Galois branched coverings (see, for example, [32, Proposition 3.2.10]). Of course, it is strange to use Riemann's existence theorem [16] for a statement that can be proved using basic properties of group actions.

#### 4.1 GTSh-orbit of an Abelian child's drawing is a singleton

Let us prove the following auxiliary statement:

**Proposition 4.6** *Suppose that  $(g_1, g_2) \in S_d \times S_d$  generate an Abelian and transitive subgroup  $G \leq S_d$ . Then, for every  $r \in \mathbb{Z}$  such that  $\gcd(r, \text{ord}(g_1)) = \gcd(r, \text{ord}(g_2)) = 1$ , there exists  $h \in S_d$  such that*

$$g_1^r = hg_1h^{-1} \quad \text{and} \quad g_2^r = hg_2h^{-1}. \quad (4.2)$$

**Proof.** Recall that, due to Proposition 4.2, the group  $G$  has order  $d$ . The key fact that is used in the proof of the desired statement is that any  $G$ -set of size  $d$  with a transitive action of  $G$  is isomorphic to the  $G$ -set  $G$  with the standard  $G$ -action by multiplication.

Let us show that  $r$  is coprime to  $\text{ord}(w)$  for every  $w \in G$ .

Since  $r$  is coprime to  $\text{ord}(g_1)$  and  $\text{ord}(g_2)$ ,  $r$  is coprime to  $\text{ord}(g_1^{t_1})$  and  $\text{ord}(g_2^{t_2})$  for all integers  $t_1$  and  $t_2$ .

Since  $G = \langle g_1, g_2 \rangle$  is Abelian, every  $w \in G$  can be written in the form

$$w = g_1^{t_1} g_2^{t_2}.$$

Let  $k_i := \text{ord}(g_i^{t_i})$  and  $k$  be the least common multiple of  $k_1$  and  $k_2$ . Since  $r$  is coprime to  $k_1$  and  $k_2$ ,  $r$  is also coprime to  $k$ .

Since  $w^k = \text{id}$ , we conclude that  $\text{ord}(w) | k$ . Thus we proved that  $r$  is coprime to  $\text{ord}(w)$  for every  $w \in G$ .

Since the group  $G = \langle g_1, g_2 \rangle$  is Abelian, the formula

$$\theta(w) := w^r, \quad w \in G \quad (4.3)$$

defines a group endomorphism  $\theta : G \rightarrow G$ .

Moreover, since  $r$  is coprime to  $\text{ord}(w)$  for every  $w \in G$ ,  $w^r = \text{id}$  if and only if  $w = \text{id}$ . Hence the homomorphism  $\theta$  is injective. Since the group  $G$  is finite, the injectivity of  $\theta$  implies its surjectivity. Thus  $\theta$  is actually an automorphism of  $G$ .

Precomposing the inclusion homomorphism  $G \rightarrow S_d$  with  $\theta$ , we get the new action  $\mathbf{a}$  of  $G$  given by the formula

$$\mathbf{a}(w)(i) := w^r(i).$$

Since  $\theta$  is an automorphism, the new action of  $G$  on  $\{1, 2, \dots, d\}$  is also transitive. Since  $G$  has order  $d$  and the new action of  $G$  is transitive, we conclude that the new  $G$ -set is isomorphic to the original one.

The proposition is proved.  $\square$

Let us use Proposition 4.6 to prove that the orbit of every Abelian child's drawing (with respect to the action of GT-shadows) is a singleton:

**Corollary 4.7** *Let  $[(g_1, g_2)]$  be an Abelian child's drawing of degree  $d$  and  $\mathbf{N} \in \mathbf{NFI}_{\text{PB}_4}(\mathbf{B}_4)$  such that  $[(g_1, g_2)]$  is subordinate to  $\mathbf{N}$ . For every  $[m, f] \in \text{GT}(\mathbf{N})$ , we have*

$$[(g_1, g_2)]^{[m, f]} = [(g_1, g_2)].$$

**Proof.** Let  $\psi : \mathbf{F}_2 \rightarrow \langle g_1, g_2 \rangle$  be the homomorphism that sends  $x_{12}$  (resp.  $x_{23}$ ) to  $g_1$  (resp. to  $g_2$ ). Furthermore, let  $h := \psi(f)$ .

We know that  $[(g_1, g_2)]^{[m, f]}$  is represented by the pair

$$(g_1^{2m+1}, h^{-1}g_2^{2m+1}h).$$

Since the group  $\langle g_1, g_2 \rangle$  is Abelian,  $h^{-1}g_2^{2m+1}h = g_2^{2m+1}$ . Hence the child's drawing  $[(g_1, g_2)]^{[m, f]}$  is represented by the pair

$$(g_1^{2m+1}, g_2^{2m+1}).$$

Since  $g_1$  (resp.  $g_2$ ) is the image of  $x_{12}\mathbf{N}_{\mathbf{F}_2}$  (resp.  $x_{23}\mathbf{N}_{\mathbf{F}_2}$ ) and  $\text{ord}(x_{12}\mathbf{N}_{\mathbf{F}_2}) = \text{ord}(x_{12}\mathbf{N}_{\text{PB}_3})$ ,  $\text{ord}(x_{23}\mathbf{N}_{\mathbf{F}_2}) = \text{ord}(x_{23}\mathbf{N}_{\text{PB}_3})$ , we have  $\text{ord}(g_1) | \text{ord}(x_{12}\mathbf{N}_{\text{PB}_3})$  and  $\text{ord}(g_2) | \text{ord}(x_{23}\mathbf{N}_{\text{PB}_3})$ .

Hence, using the fact that  $N_{\text{ord}}$  is a multiple of the orders  $\text{ord}(x_{12}\mathbf{N}_{\text{PB}_3})$ ,  $\text{ord}(x_{23}\mathbf{N}_{\text{PB}_3})$  and  $2m+1$  is coprime with  $N_{\text{ord}}$ , we conclude that  $2m+1$  is coprime with the integers  $\text{ord}(g_1)$  and  $\text{ord}(g_2)$ .

Thus, applying Proposition 4.6, we conclude that the pairs  $(g_1^{2m+1}, g_2^{2m+1})$  and  $(g_1, g_2)$  represent the same child's drawing.  $\square$

Combining [4, Proposition 2.5], [29, Corollary on page 2] with Proposition 4.2 and Corollaries 3.9, 4.7, we deduce the following statement:

**Corollary 4.8** *Every Abelian child's drawing  $\mathcal{D}$  admits a Belyi pair  $(X, \gamma)$  defined over  $\mathbb{Q}$ .*

**Remark 4.9** As far as the author understands, the statement of the above corollary was proved directly in paper [19] by R.A. Hidalgo (see [19, Corollary 3.5]).

## 5 Examples of $\text{GTSh}^\heartsuit$ -orbits for (non-Abelian) child's drawings

### 5.1 An example of degree 6, genus 0 and orbit size 2

Let us denote by  $\mathcal{D}_{6,0}$  the degree 6 child's drawing represented by triple

$$((1, 4, 5, 2)(3, 6), (1, 6, 3, 2)(4, 5), ((1, 3), (2, 4))). \quad (5.1)$$

The passport of  $\mathcal{D}_{6,0}$  is

$$((4, 2), (4, 2), (2, 2, 1, 1))$$

and its genus is 0. A Belyi map representing  $\mathcal{D}_{6,0}$  can be found at <https://beta.lmfdb.org/Belyi/6T10/4>

The  $G_{\mathbb{Q}}$ -orbit of  $\mathcal{D}_{6,0}$  has size 2 and the Galois conjugate of  $\mathcal{D}_{6,0}$  is represented by the permutation triple

$$((1, 4, 5, 2)(3, 6), (1, 2, 5, 6)(3, 4), (3, 5)(4, 6)). \quad (5.2)$$

Using [6], we found an element  $\mathbf{N} \in \mathbf{NFI}_{\mathrm{PB}_4}(\mathbf{B}_4)$  that dominates  $\mathcal{D}_{6,0}$ .  $\mathbf{N}$  is the kernel of a group homomorphism  $\mathrm{PB}_4 \rightarrow S_{192}$  stored in the file *E\_dde6genus0* (see [6, Section 6]). Here are some basic facts about this element of  $\mathbf{NFI}_{\mathrm{PB}_4}(\mathbf{B}_4)$ :

- $|\mathrm{PB}_4 : \mathbf{N}| = 289, 207, 845, 356, 544 = 2^{10} \cdot 3^{24}$ ;
- $|\mathrm{PB}_3 : \mathbf{N}_{\mathrm{PB}_3}| = 46656 = 2^6 \cdot 3^6$ ;
- $|\mathrm{F}_2 : \mathbf{N}_{\mathrm{F}_2}| = 11664 = 2^4 \cdot 3^6$ ;
- the order of the commutator subgroup  $[\mathrm{F}_2/\mathbf{N}_{\mathrm{F}_2}, \mathrm{F}_2/\mathbf{N}_{\mathrm{F}_2}]$  is  $729 = 3^6$ ;
- $N_{\mathrm{ord}} := |\mathrm{PB}_2 : \mathbf{N}_{\mathrm{PB}_2}| = 4$ ;
- $\mathbf{N}$  is isolated and  $\mathrm{GT}^\heartsuit(\mathbf{N})$  is a non-Abelian group of order  $32 = 2^5$ .

The orbit  $\mathrm{GT}^\heartsuit(\mathbf{N})(\mathcal{D}_{6,0})$  coincides with the  $G_{\mathbb{Q}}$ -orbit of  $\mathcal{D}_{6,0}$ .

The interesting feature of  $\mathcal{D}_{6,0}$  is that the  $\mathrm{GT}$ -shadow  $[3, 1_{\mathrm{F}_2}] \in \mathrm{GT}^\heartsuit(\mathbf{N})$  corresponding to the complex conjugation acts trivially on  $\mathcal{D}_{6,0}$ . A  $\mathrm{GT}$ -shadow that transforms  $\mathcal{D}_{6,0}$  to its Galois conjugate is represented by the pair

$$(1, yxyx^2y^2x^{-3}y^{-4}) \in \mathbb{Z} \times [\mathrm{F}_2, \mathrm{F}_2].$$

In fact, there are  $\mathrm{GT}$ -shadows that transforms  $\mathcal{D}_{6,0}$  to its Galois conjugate and belong to the kernel of the virtual cyclotomic character. Here is an example of such a  $\mathrm{GT}$ -shadow

$$[0, yxy^3xy^2x^2y^{-6}x^{-4}] \in \mathrm{GT}^\heartsuit(\mathbf{N}).$$

## 5.2 An example of degree 5, genus 0 and orbit size 2

Let  $\mathcal{D}_{5,0}$  be the child's drawing of degree 5 represented by the permutation triple

$$((1, 4, 5, 2), (2, 3, 5, 4), (1, 4)(2, 3)). \quad (5.3)$$

The passport of  $\mathcal{D}_{5,0}$  is

$$((4, 1), (4, 1), (2, 2, 1))$$

and its genus is 0. A Belyi map representing  $\mathcal{D}_{5,0}$  can be found at <https://beta.lmfdb.org/Belyi/5T3/4>

The  $G_{\mathbb{Q}}$ -orbit of  $\mathcal{D}_{5,0}$  has size 2 and the Galois conjugate of  $\mathcal{D}_{5,0}$  is represented by the permutation triple

$$((1, 2, 5, 4), (1, 5, 2, 3), (1, 4)(2, 3)). \quad (5.4)$$

Using [6], we found an element  $\mathbf{N} \in \mathbf{NFI}_{\mathrm{PB}_4}(\mathbf{B}_4)$  that dominates  $\mathcal{D}_{5,0}$ .  $\mathbf{N}$  is the kernel of a group homomorphism  $\mathrm{PB}_4 \rightarrow S_{160}$  stored in the file *E\_dde5genus0* (see [6, Section 6]).

Here are some basic facts about this element of  $\mathbf{NFI}_{\mathrm{PB}_4}(\mathbf{B}_4)$ :

- $|\mathrm{PB}_4 : \mathbf{N}| = 250000000000 = 2^{10} \cdot 5^{12}$ ;
- $|\mathrm{PB}_3 : \mathbf{N}_{\mathrm{PB}_3}| = 8000 = 2^6 \cdot 5^3$ ;
- $|\mathrm{F}_2 : \mathbf{N}_{\mathrm{F}_2}| = 2000 = 2^4 \cdot 5^3$ ;
- the order of the commutator subgroup  $[\mathrm{F}_2/\mathbf{N}_{\mathrm{F}_2}, \mathrm{F}_2/\mathbf{N}_{\mathrm{F}_2}]$  is  $125 = 5^3$ ;

- $N_{\text{ord}} := |\text{PB}_2 : \text{N}_{\text{PB}_2}| = 4$ ;
- $\text{N}$  is not settled and the connected component of  $\text{N}$  in  $\text{GTSh}^\heartsuit$  has exactly two objects; the size of  $\text{GT}^\heartsuit(\text{N})$  is 16.

The orbits  $\text{GT}^\heartsuit(\text{N})(\mathcal{D}_{5,0})$  and  $G_{\mathbb{Q}}(\mathcal{D}_{5,0})$  coincide.

### 5.3 Examples of child's drawing subordinate to $\text{N}^{(5)}$ and $\text{N}^{(29)}$ from list (3.19)

Recall that [7, Section 4] presents basic information about 35 selected elements  $\text{N}^{(0)}, \dots, \text{N}^{(34)}$  of  $\text{NFI}_{\text{PB}_4}(\text{B}_4)$ . In this subsection, we describe child's drawings of degrees 7, 15 and 18 subordinate to  $\text{N}^{(29)}$  and a child's drawing of degree 8 subordinate to  $\text{N}^{(5)}$ .

Here is the basic information about  $\text{N}^{(29)}$ :

- $|\text{PB}_4 : \text{N}^{(29)}| = 2520 = 2^3 \cdot 3^2 \cdot 5 \cdot 7$ ;
- $|\text{PB}_3 : \text{N}_{\text{PB}_3}^{(29)}| = 136080 = 2^4 \cdot 3^5 \cdot 5 \cdot 7$ ;
- $|\text{F}_2 : \text{N}_{\text{F}_2}^{(29)}| = 45,360 = 2^4 \cdot 3^4 \cdot 5 \cdot 7$ ;
- $N_{\text{ord}}^{(29)} = 6$ ;
- $\text{N}^{(29)}$  is an isolated object of  $\text{NFI}_{\text{PB}_4}(\text{B}_4)$  and  $\text{GT}^\heartsuit(\text{N}^{(29)})$  is non-Abelian group of order  $48 = 2^4 \cdot 3$ .

The child's drawings represented by the following permutation triples

$$((1, 2, 3)(4, 5)(6, 7), (1, 5, 6)(2, 7)(3, 4), (1, 4)(2, 6)(3, 7, 5)), \quad (5.5)$$

$$\begin{aligned} &((1, 2, 3, 4, 5, 6)(7, 8, 9, 10, 11, 12)(13, 14, 15), \\ &(1, 2, 6, 12, 9, 15)(3, 7, 13)(4, 11, 14, 5, 8, 10), \\ &(1, 2, 15, 11, 8, 3)(4, 13, 9, 5, 10, 12)(6, 14, 7)) \end{aligned} \quad (5.6)$$

$$\begin{aligned} &((1, 10, 17, 2, 9, 18)(3, 12, 13, 4, 11, 14)(5, 8, 15, 6, 7, 16), \\ &(1, 16, 11, 2, 15, 12)(3, 18, 7, 4, 17, 8)(5, 14, 9, 6, 13, 10) \\ &(1, 3, 5)(2, 4, 6)(7, 9, 11)(8, 10, 12)(13, 15, 17)(14, 16, 18)) \end{aligned} \quad (5.7)$$

are subordinate to  $\text{N}^{(29)}$ . We denote by  $\mathcal{D}_{7,0}$  (resp.  $\mathcal{D}_{15,4}$ ,  $\mathcal{D}_{18,4}$ ) the child's drawing represented by the permutation triple in (5.5) (resp. in (5.6), (5.7)).

The child's drawings  $\mathcal{D}_{7,0}$ ,  $\mathcal{D}_{15,4}$  and  $\mathcal{D}_{18,4}$  have degrees 7, 15 and 18, respectively.

The passport of  $\mathcal{D}_{7,0}$  is  $((3, 2, 2), (3, 2, 2), (3, 2, 2))$  and its genus is zero. A Belyi map that represents  $\mathcal{D}_{7,0}$  can be found at <https://beta.lmfdb.org/Belyi/7T6/3.2.2/3.2.2/3.2.2/a>.

The passports of  $\mathcal{D}_{15,4}$  and  $\mathcal{D}_{18,4}$  are

$$((6, 6, 3), (6, 6, 3), (6, 6, 3)) \quad \text{and} \quad ((6, 6, 6), (6, 6, 6), (3, 3, 3, 3, 3, 3))$$

respectively. Both child's drawings  $\mathcal{D}_{15,4}$  and  $\mathcal{D}_{18,4}$  have genus 4.

The orbit  $\mathrm{GT}^\heartsuit(\mathbf{N}^{(29)})(\mathcal{D}_{7,0})$  coincides with the  $G_{\mathbb{Q}}$ -orbit of  $\mathcal{D}_{7,0}$ . In fact, using [6], one can show that there is only one child's drawing with the passport  $((3, 2, 2), (3, 2, 2), (3, 2, 2))$ . Hence the  $G_{\mathbb{Q}}$ -orbit of  $\mathcal{D}_{7,0}$  and the  $\mathrm{GT}^\heartsuit(\mathbf{N})$ -orbit of  $\mathcal{D}_{7,0}$  (for any  $\mathbf{N} \in \mathrm{NFI}_{\mathrm{PB}_4}(\mathbf{B}_4)$  that dominates  $\mathcal{D}_{7,0}$ ) must have size 1.

The orbit  $\mathrm{GT}^\heartsuit(\mathbf{N}^{(29)})(\mathcal{D}_{15,4})$  has two elements and the “conjugate” of  $\mathcal{D}_{15,4}$  is represented by the permutation triple

$$\begin{aligned} &((1, 6, 5, 4, 3, 2)(7, 12, 11, 10, 9, 8)(13, 15, 14), \\ &(1, 15, 9, 12, 6, 2)(3, 13, 7)(4, 10, 8, 5, 14, 11), \\ &(1, 6, 2, 7, 10, 14)(3, 11, 9, 4, 8, 15)(5, 12, 13)). \end{aligned} \tag{5.8}$$

Currently, database [24] does not contain Belyi maps of degree 15. However, we can prove that

**Claim 5.1** *The  $G_{\mathbb{Q}}$ -orbit of  $\mathcal{D}_{15,4}$  coincides with the orbit  $\mathrm{GT}^\heartsuit(\mathbf{N}^{(29)})(\mathcal{D}_{15,4}) = \{\mathcal{D}_{15,4}, \mathcal{D}_{15,4}^*\}$ , where  $\mathcal{D}_{15,4}^*$  is the child's drawing represented by the permutation triple in (5.8).*

**Proof.** The image of the complex conjugation in  $\mathrm{GT}^\heartsuit(\mathbf{N}^{(29)})$  is represented by the pair  $(-1, 1_{\mathbb{F}_2})$  and

$$\mathcal{D}_{15,4}^* = \mathcal{D}_{15,4}^{[-1, 1_{\mathbb{F}_2}]}$$

Hence  $\mathrm{GT}^\heartsuit(\mathbf{N}^{(29)}) \subset G_{\mathbb{Q}}(\mathcal{D}_{15,4})$ . The inclusion  $G_{\mathbb{Q}}(\mathcal{D}_{15,4}) \subset \mathrm{GT}^\heartsuit(\mathbf{N}^{(29)})$  is a consequence of Corollary 3.9.  $\square$

We should remark that the  $G_{\mathbb{Q}}$ -orbit of  $\mathcal{D}_{15,4}$  is significantly smaller than the number of child's drawings with the passport  $((6, 6, 3), (6, 6, 3), (6, 6, 3))$ . Using [6], one can show that the number of child's drawings with the passport  $((6, 6, 3), (6, 6, 3), (6, 6, 3))$  is  $\geq 260$ .

The orbit  $\mathrm{GT}^\heartsuit(\mathbf{N}^{(29)})(\mathcal{D}_{18,4})$  is a singleton. The child's drawing  $\mathcal{D}_{18,4}$  is special because the corresponding (degree 18) covering of  $\mathbb{CP}^1 - \{0, 1, \infty\}$  is Galois. In fact, the child's drawing  $\mathcal{D}_{18,4}$  can be represented by the subgroup of  $\mathbb{F}_2$  that corresponds to the commutator subgroup  $[\mathbb{F}_2/\mathbf{N}_{\mathbb{F}_2}, \mathbb{F}_2/\mathbf{N}_{\mathbb{F}_2}]$ . (See Remark 2.8.)

Here is the basic information about element  $\mathbf{N}^{(5)}$  from (3.19):

- $|\mathrm{PB}_4 : \mathbf{N}^{(5)}| = 24 = 2^3 \cdot 3$ ;
- $|\mathrm{PB}_3 : \mathbf{N}_{\mathrm{PB}_3}^{(5)}| = 864 = 2^5 \cdot 3^3$ ;
- $|\mathbb{F}_2 : \mathbf{N}_{\mathbb{F}_2}^{(5)}| = 288 = 2^5 \cdot 3^2$ ;
- $N_{\mathrm{ord}}^{(5)} = 6$ ;
- $\mathbf{N}^{(5)}$  is an isolated object of  $\mathrm{NFI}_{\mathrm{PB}_4}(\mathbf{B}_4)$  and the group  $\mathrm{GT}^\heartsuit(\mathbf{N}^{(5)})$  is isomorphic to  $D_6$  (the dihedral group of order 12).

The child's drawing  $\mathcal{D}_{8,0}$  represented by the permutation triple

$$((1, 2, 3)(4, 5, 6), (1, 8, 5)(2, 4, 7), (1, 3, 7, 4, 6, 8)(2, 5)). \tag{5.9}$$

is subordinate to  $\mathbf{N}^{(5)}$ .

$\mathcal{D}_{8,0}$  has genus 0 and its passport is

$$((3, 3, 1, 1), (3, 3, 1, 1), (6, 2)).$$

The number of child's drawings with this passport is 5.

The orbit  $\text{GT}^\heartsuit(\mathbf{N}^{(5)})(\mathcal{D}_{8,0})$  is a singleton. Hence the  $G_{\mathbb{Q}}$ -orbit of  $\mathcal{D}_{8,0}$  is also a singleton.

Since  $\mathbf{N}^{(19)} \leq \mathbf{N}^{(5)}$ ,  $\mathbf{N}^{(19)}$  (from the list in (3.19)) also dominates  $\mathcal{D}_{8,0}$ .

The author could not find  $\mathcal{D}_{8,0}$  in [24]. However,  $\mathcal{D}_{8,0}$  can be transformed, by the action of  $B_3$ , to the child's drawing represented by the Belyi map <https://beta.lmfdb.org/Belyi/8T12/6.2/3.3>.

## A Charming GT-shadows satisfy the simplified hexagon relations

Let us prove the following auxiliary statement:

**Proposition A.1** *Let  $\mathbf{N} \in \text{NFI}_{\text{PB}_4}(\mathbf{B}_4)$ . If a pair  $(m, f) \in \mathbb{Z} \times \mathbf{F}_2$  satisfies hexagon relations (2.6), (2.7) (modulo  $\mathbf{N}_{\text{PB}_3}$ ) then*

$$f(x, y)f(y, x) \in \mathbf{N}_{\mathbf{F}_2}, \quad (\text{A.1})$$

where  $\mathbf{N}_{\mathbf{F}_2} := \mathbf{N}_{\text{PB}_3} \cap \mathbf{F}_2$ .

**Proof.** Let us consider the element

$$\sigma_1^{2m+1} f^{-1} \sigma_2^{2m+1} f \sigma_1^{2m+1} \mathbf{N}_{\text{PB}_3} \in B_3 / \mathbf{N}_{\text{PB}_3} \quad (\text{A.2})$$

On the one hand, (2.6) implies that

$$\begin{aligned} (\sigma_1^{2m+1} f^{-1} \sigma_2^{2m+1} f) \sigma_1^{2m+1} \mathbf{N}_{\text{PB}_3} &= f^{-1} \sigma_1 \sigma_2 c^m \sigma_1^{-2m} \sigma_1^{2m+1} \mathbf{N}_{\text{PB}_3} = \\ f^{-1} \sigma_1 \sigma_2 c^m \sigma_1 \mathbf{N}_{\text{PB}_3} &= f^{-1} \theta c^m \mathbf{N}_{\text{PB}_3}, \end{aligned}$$

where  $\theta := \sigma_1 \sigma_2 \sigma_1$ .

On the other hand, using (2.7) and the relation  $\sigma_1 \theta = \theta \sigma_2$  we get

$$\begin{aligned} \sigma_1^{2m+1} (f^{-1} \sigma_2^{2m+1} f \sigma_1^{2m+1}) \mathbf{N}_{\text{PB}_3} &= \sigma_1^{2m+1} \sigma_2 \sigma_1 c^m \sigma_2^{-2m} f \mathbf{N}_{\text{PB}_3} = \\ \sigma_1^{2m} \theta c^m \sigma_2^{-2m} f \mathbf{N}_{\text{PB}_3} &= \theta \sigma_2^{2m} c^m \sigma_2^{-2m} f \mathbf{N}_{\text{PB}_3} = \theta f c^m \mathbf{N}_{\text{PB}_3}. \end{aligned}$$

Thus  $\theta f c^m \mathbf{N}_{\text{PB}_3} = f^{-1} \theta c^m \mathbf{N}_{\text{PB}_3}$  and hence

$$\theta f \theta^{-1} \mathbf{N}_{\text{PB}_3} = f^{-1} \mathbf{N}_{\text{PB}_3}. \quad (\text{A.3})$$

Since  $\theta x_{12} \theta^{-1} = x_{23}$  and  $\theta x_{23} \theta^{-1} = x_{12}$  and  $f \in \mathbf{F}_2$ , relation (A.3) implies (A.1).  $\square$

If we assume that  $f \in [\mathbf{F}_2, \mathbf{F}_2]$ , then (A.1) implies additional useful properties:

**Proposition A.2** *Let  $\mathbf{N} \in \text{NFI}_{\text{PB}_4}(\mathbf{B}_4)$  and  $\mathbf{N}_{\mathbf{F}_2} := \mathbf{N}_{\text{PB}_3} \cap \mathbf{F}_2$ . If  $f \in [\mathbf{F}_2, \mathbf{F}_2]$  satisfies (A.1), then*

$$f(y, z)f(z, y) \in \mathbf{N}_{\mathbf{F}_2}, \quad (\text{A.4})$$

and

$$f(z, x)f(x, z) \in \mathbf{N}_{\mathbf{F}_2}, \quad (\text{A.5})$$

where  $z := y^{-1}x^{-1}$ .



**Proof.** Relation (A.1) obviously implies that

$$f(x, y)f(y, x) \in \mathbf{N}_{\text{PB}_3}. \quad (\text{A.6})$$

Since

$$\sigma_1 \sigma_2 x_{12} (\sigma_1 \sigma_2)^{-1} = x_{23} \quad \text{and} \quad \sigma_1 \sigma_2 x_{23} (\sigma_1 \sigma_2)^{-1} = x_{23}^{-1} x_{12}^{-1} c,$$

conjugating (A.6) by  $\sigma_1 \sigma_2$  gives us

$$f(y, zc)f(zc, y) \in \mathbf{N}_{\text{PB}_3}. \quad (\text{A.7})$$

Since  $f \in [\mathbf{F}_2, \mathbf{F}_2]$  and  $c \in \mathcal{Z}(\text{PB}_3)$ , relation (A.7) implies (A.4).

Conjugating relation (A.7) by  $\sigma_1 \sigma_2$ , we get

$$f(zc, x)f(x, zc) \in \mathbf{N}_{\text{PB}_3}. \quad (\text{A.8})$$

Since  $f(z, x) \in [\mathbf{F}_2, \mathbf{F}_2]$  and  $c \in \mathcal{Z}(\text{PB}_3)$ , relation (A.8) implies (A.5).  $\square$

We will now get the versions of relations (4.3) and (4.4) from Section 4 of Drinfeld's foundational paper [5]:

**Proposition A.3** *Let  $\mathbf{N} \in \mathbf{NFI}_{\text{PB}_4}(\mathbf{B}_4)$  and  $(m, f) \in \mathbb{Z} \times [\mathbf{F}_2, \mathbf{F}_2]$ . The pair  $(m, f)$  satisfies hexagon relations (2.6), (2.7) (modulo  $\mathbf{N}_{\text{PB}_3}$ ) if and only if  $(m, f)$  satisfies the relations*

$$f(x, y)f(y, x) \in \mathbf{N}_{\mathbf{F}_2}, \quad (\text{A.9})$$

$$x^m f(z, x) z^m f(y, z) y^m f(x, y) \in \mathbf{N}_{\mathbf{F}_2}, \quad (\text{A.10})$$

where  $\mathbf{N}_{\mathbf{F}_2} := \mathbf{N}_{\text{PB}_3} \cap \mathbf{F}_2$  and  $z := y^{-1}x^{-1}$ .

**Proof.** Recall that

$$x := x_{12}, \quad y := x_{23}, \quad z := y^{-1}x^{-1}, \quad w := x^{-1}y^{-1}.$$

Using the relations

$$\sigma_2^{-1} x_{12} \sigma_2 = x_{23}^{-1} x_{12}^{-1} c = zc, \quad \sigma_2^{-1} x_{23} \sigma_2 = x_{23}$$

and the properties  $c \in \mathcal{Z}(\text{PB}_3)$ ,  $f^{-1} \in [\mathbf{F}_2, \mathbf{F}_2]$ , we rewrite the left hand side of (2.6) as follows:

$$\sigma_1 x_{12}^m f^{-1} \sigma_2 x_{23}^m f \mathbf{N}_{\text{PB}_3} = \sigma_1 \sigma_2 c^m z^m f^{-1}(z, y) y^m f \mathbf{N}_{\text{PB}_3}. \quad (\text{A.11})$$

Using the relations

$$\sigma_2^{-1} \sigma_1^{-1} x_{12} \sigma_1 \sigma_2 = x_{23}^{-1} x_{12}^{-1} c = zc, \quad \sigma_2^{-1} \sigma_1^{-1} x_{23} \sigma_1 \sigma_2 = x_{12} = x$$

and the properties  $c \in \mathcal{Z}(\text{PB}_3)$ ,  $f^{-1} \in [\mathbf{F}_2, \mathbf{F}_2]$ , we rewrite the right hand side of (2.6) as follows:

$$f^{-1} \sigma_1 \sigma_2 x_{12}^{-m} c^m \mathbf{N}_{\text{PB}_3} = \sigma_1 \sigma_2 c^m f^{-1}(z, x) x^{-m} \mathbf{N}_{\text{PB}_3}. \quad (\text{A.12})$$

Combining (A.11) and (A.12), we see that relation (2.6) is equivalent to

$$x^m f(z, x) z^m f^{-1}(z, y) y^m f(x, y) \in \mathbf{N}_{\mathbf{F}_2}. \quad (\text{A.13})$$

Using the relations

$$\sigma_2 \sigma_1 x_{12} \sigma_1^{-1} \sigma_2^{-1} = c x_{12}^{-1} x_{23}^{-1} = cw, \quad \sigma_2 \sigma_1 x_{23} \sigma_1^{-1} \sigma_2^{-1} = x_{12} = x$$

and the properties  $c \in \mathcal{Z}(\text{PB}_3)$ ,  $f \in [\mathbf{F}_2, \mathbf{F}_2]$ , we rewrite the right hand side of (2.7) as follows:

$$\sigma_2 \sigma_1 x_{23}^{-m} c^m f \mathbf{N}_{\text{PB}_3} = x^{-m} f(w, x) \sigma_2 \sigma_1 c^m \mathbf{N}_{\text{PB}_3} \quad (\text{A.14})$$

Using the relations

$$\sigma_2 x_{12} \sigma_2^{-1} = cw, \quad \sigma_2 x_{23} \sigma_2^{-1} = x_{23} = y$$

and the properties  $c \in \mathcal{Z}(\text{PB}_3)$ ,  $f \in [\mathbf{F}_2, \mathbf{F}_2]$ , we rewrite the left hand side of (2.7) as follows:

$$f^{-1} \sigma_2 x_{23}^m f \sigma_1 x_{12}^m \mathbf{N}_{\text{PB}_3} = f^{-1} y^m f(w, y) w^m \sigma_2 \sigma_1 c^m \mathbf{N}_{\text{PB}_3}. \quad (\text{A.15})$$

Combining (A.15) and (A.14), we see that relation (2.7) is equivalent to

$$y^m f(w, y) w^m f^{-1}(w, x) x^m f^{-1}(x, y) \in \mathbf{N}_{\text{PB}_3}. \quad (\text{A.16})$$

Conjugating (A.16) by  $\theta := \sigma_1 \sigma_2 \sigma_1$ , we see that relation (2.7) is equivalent to

$$x^m f(z, x) z^m f^{-1}(z, y) y^m f^{-1}(y, x) \in \mathbf{N}_{\mathbf{F}_2}. \quad (\text{A.17})$$

We can now prove the desired equivalence. Indeed, if a pair  $(m, f) \in \mathbb{Z} \times [\mathbf{F}_2, \mathbf{F}_2]$  satisfies (2.6) and (2.7) then, due to Proposition A.1, relation (A.9) holds.

Moreover, relation (A.4) from Proposition A.2 implies that

$$f^{-1}(z, y) \mathbf{N}_{\mathbf{F}_2} = f(y, z) \mathbf{N}_{\mathbf{F}_2}. \quad (\text{A.18})$$

Hence (A.13) implies (A.10).

We proved that relations (2.6), (2.7) imply relations (A.9), (A.10).

Let us now assume that a pair  $(m, f) \in \mathbb{Z} \times [\mathbf{F}_2, \mathbf{F}_2]$  satisfies relations (A.9) and (A.10). Due to Proposition A.2, relation (A.18) holds. Using (A.9), (A.10) and (A.18) we deduce relations (A.13) and (A.17). Since we showed above that relation (A.13) (resp. relation (A.17)) is equivalent to (2.6) (resp. (2.7)), we proved that relations (A.9), (A.10) imply hexagon relations (2.6), (2.7).  $\square$

## References

- [1] N.M. Adrianov, N.Y. Amburg, V.A. Dremov, Yu. Yu. Kochetkov, E.M. Kreines, Yu.A. Levitskaya, V.F. Nasretdinova, G.B. Shabat, A catalogue of Belyi functions of dessins d'enfants with at most four edges, (*in Russian*) Fundam. Prikl. Mat. **13**, 6 (2007) 35–112; *translation in* J. Math. Sci. (N.Y.) **158**, 1 (2007) 22–80.
- [2] J. Anderson, I. Bouw, O. Ejder, N. Girgin, V. Karemaker, M. Manes, Dynamical Belyi maps, Women in numbers Europe II, 57–82, Assoc. Women Math. Ser., **11**, Springer, Cham, 2018; arXiv:1703.08563.
- [3] D. Bar-Natan, On Associators and the Grothendieck-Teichmueller Group I, Selecta Math. (N.S.) **4**, 2 (1998) 183–212; arXiv:q-alg/9606021

- [4] K. Coombes and D. Harbater, Hurwitz families and arithmetic Galois groups, *Duke Math. J.* **52**, 4 (1985) 821–839.
- [5] V. Drinfeld, On quasitriangular quasi-Hopf algebras and on a group that is closely connected with  $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ , *Algebra i Analiz* **2**, 4 (1990) 149–181.
- [6] V. A. Dolgushev, Documentation for the package GT is available at <https://math.temple.edu/~vald/PackageGT/README.pdf>
- [7] V. A. Dolgushev, K.Q. Le and A.A. Lorenz, What are GT-shadows? arXiv:2008.00066.
- [8] V.A. Dolgushev and C.L. Rogers, Notes on algebraic operads, graph complexes, and Willwacher’s construction, *Mathematical aspects of quantization*, 25–145, *Contemp. Math.*, **583**, Amer. Math. Soc., Providence, RI, 2012.
- [9] D. S. Dummit and R. M. Foote, *Abstract Algebra*, 3rd Edition, John Wiley and Sons, 2003.
- [10] J. Ellenberg, Galois invariants of dessins d’enfants, *Arithmetic fundamental groups and noncommutative algebra* (Berkeley, CA, 1999), 27–42, *Proc. Sympos. Pure Math.*, **70**, Amer. Math. Soc., Providence, RI, 2002.
- [11] B. Fresse, Homotopy of operads and Grothendieck-Teichmueller groups. Part 1, The algebraic theory and its topological background, *Mathematical Surveys and Monographs*, **217**. AMS, Providence, RI, 2017. xlv+532 pp.
- [12] E. Gironde and G. González-Diez, Introduction to compact Riemann surfaces and dessins d’enfants, *London Mathematical Society Student Texts*, **79**. Cambridge University Press, Cambridge, 2012.
- [13] E.H. Goins, The blog on Dessins d’Enfants, <https://dessindenfants.wordpress.com/>
- [14] A. Grothendieck, Esquisse d’un programme, *London Math. Soc. Lecture Note Ser.*, **242**, Geometric Galois actions, 1, 5–48, Cambridge Univ. Press, Cambridge, 1997.
- [15] A. Grothendieck and M. Raynaud, Revêtements étales et groupe fondamental, (SGA1), *Lecture Notes in Mathematics*, **224**. Springer-Verlag, Berlin-New York, 1971.
- [16] D. Harbater, Riemann’s Existence Theorem, <https://www2.math.upenn.edu/~harbater/RETppr.pdf>
- [17] D. Harbater and L. Schneps, Approximating Galois orbits of dessins, *Geometric Galois actions*, 1, 205–230, *London Math. Soc. Lecture Note Ser.*, **242**, Cambridge Univ. Press, Cambridge, 1997.
- [18] A. Hatcher, Algebraic Topology, <https://pi.math.cornell.edu/~hatcher/AT/AT.pdf>
- [19] R. A. Hidalgo, Homology closed Riemann surfaces, *Q. J. Math.* **63**, 4 (2012) 931–952.
- [20] Y. Ihara, On the embedding of  $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$  into  $\widehat{\text{GT}}$ , *with an appendix by M. Emsalem and P. Lochak*, *London Math. Soc. Lecture Note Ser.*, **200**, The Grothendieck theory of dessins d’enfants (Luminy, 1993), 289–321, Cambridge Univ. Press, Cambridge, 1994.
- [21] S.K. Lando and A.K. Zvonkin, Graphs on surfaces and their applications. With an appendix by Don B. Zagier, *Encyclopaedia of Mathematical Sciences*, **141**. Low-Dimensional Topology, II. Springer-Verlag, Berlin, 2004.

- [22] J.-L. Loday and B. Vallette, Algebraic operads, *Grundlehren der Mathematischen Wissenschaften*, **346**. Springer, Heidelberg, 2012. xxiv+634 pp.
- [23] M. Markl, S. Shnider and J. Stasheff, Operads in algebra, topology and physics, *Mathematical Surveys and Monographs*, **96**. American Mathematical Society, Providence, RI, 2002. x+349 pp.
- [24] M. Musty, S. Schiavone, J. Sijsling and J. Voight, A database of Belyi maps, <https://beta.lmfdb.org/Belyi/>; a detailed description of its construction can be found in <https://arxiv.org/abs/1805.07751>
- [25] L. Ribes and P. Zalesskii, Profinite groups. Second edition. *A Series of Modern Surveys in Mathematics* **40**. Springer-Verlag, Berlin, 2010. xvi+464 pp.
- [26] L. Schneps, An introduction to Grothendieck-Teichmueller theory, Slides of the lectures are available at <https://webusers.imj-prg.fr/~leila.schneps/mit.html>
- [27] L. Schneps, Dessins d'enfants on the Riemann Sphere, in *The Grothendieck Theory of Dessins d'Enfants*, LMS Lecture Notes **200**, Cambridge U. Press, 1994.
- [28] L. Schneps, The Grothendieck-Teichmueller group: a survey, in *Geometric Galois Theory I*, LMS Lecture Notes **242**, Cambridge U. Press, 1997.
- [29] J. Sijsling and J. Voight, Errata and Addenda: on computing Belyi maps, <https://math.dartmouth.edu/~jvoight/articles/birch-descent-errata.pdf>
- [30] J. Sijsling and J. Voight, On computing Belyi maps, Publ. Math. Besançon: Algèbre Théorie Nr. 2014/1, Presses Univ. Franche-Comté, Besançon, 73–131.
- [31] J. Stasheff, What is ... an operad? Notices Amer. Math. Soc. **51**, 6 (2004) 630–631.
- [32] T. Szamuely, Galois groups and fundamental groups, *Cambridge Studies in Advanced Mathematics*, **117**. Cambridge University Press, Cambridge, 2009.
- [33] D.E. Tamarkin, Formality of chain operad of little discs, Lett. Math. Phys. **66**, 1-2 (2003) 65–72.
- [34] L. Zapponi, What is... a dessin d'enfant? Notices of AMS, **50**, 7 (2003) 788–789.

DEPARTMENT OF MATHEMATICS, TEMPLE UNIVERSITY,  
 WACHMAN HALL RM. 638  
 1805 N. BROAD ST.,  
 PHILADELPHIA PA, 19122 USA  
*E-mail address:* **vald@temple.edu**