

CONIVEAU FILTRATIONS AND MILNOR OPERATION Q_n

NOBUAKI YAGITA

ABSTRACT. Let BG be the classifying space of an algebraic group G over the field \mathbb{C} of complex numbers. We compute a new stable birational invariant defined by the difference of two coniveau filtrations of a smooth projective approximation of $BG \times \mathbb{P}^\infty$.

1. INTRODUCTION

Let p be a prime number and $A = \mathbb{Q}, \mathbb{Z}$ or \mathbb{Z}/p^i for $i \geq 1$. Let X be a smooth algebraic variety over $k = \mathbb{C}$. Let us recall the coniveau filtration of the cohomology with coefficients in A ,

$$N^c H^i(X; A) = \sum_{Z \subset X} \ker(j^* : H^i(X; A) \rightarrow H^i(X - Z, A))$$

where $Z \subset X$ runs through the closed subvarieties of codimension at least c of X , and $j : X - Z \subset X$ is the complementary open immersion.

Similarly, we can define the *strong* coniveau filtration by

$$\tilde{N}^c H^i(X; A) = \sum_{f: Y \rightarrow X} \operatorname{im}(f_* : H^{i-2r}(Y; A) \rightarrow H^i(X, A))$$

where the sum is over all proper morphism $f : Y \rightarrow X$ from a smooth complex variety Y of $\dim(Y) = \dim(X) - r$ with $r \geq c$, and f_* its transfer (Gysin map). It is immediate that $\tilde{N}^c H^*(X; A) \subset N^c H^*(X; A)$.

It was hoped that when X is proper, the strong coniveau filtration was just the coniveau filtration, i.e., $\tilde{N}^c H^i(X; A) = N^c H^i(X; A)$. In fact Deligne shows they are the same for $A = \mathbb{Q}$. However, Benoist and Ottem ([1]) recently show that they are not equal for $A = \mathbb{Z}$.

Let G be an algebraic group such that $H^*(BG; \mathbb{Z})$ has p -torsion for the classifying space BG defined by Totaro [17]. The purpose of this paper is to compute the $\operatorname{mod}(p)$ stable birational invariant of X (Proposition 2.4 in [1])

$$DH^*(X; A) = N^1 H^*(X; A) / (p, \tilde{N}^1 H^*(X; A))$$

for projective approximations X for BG ([4],[17],[10]). In fact, we see that $DH^*(X; \mathbb{Z}) \neq 0$ happen very frequently in the above cases.

Here an approximation (for $\text{degree} \leq N$) is the projective (smooth) variety $X = X(N)$ such that there is a map $g : X \rightarrow BG \times \mathbb{P}^\infty$ with

$$g^* : H^*(BG \times \mathbb{P}^\infty; A) \cong H^*(X; A) \quad \text{for } * < N.$$

(In this paper, we say X is an approximation for BG when it is that of $BG \times \mathbb{P}^\infty$ strictly speaking.) Let us write $DH^*(X; \mathbb{Z})$ by $DH^*(X)$ simply as usual.

2010 *Mathematics Subject Classification.* Primary 20G10, 55R35, 14C15 ; Seconary 57T25.
Key words and phrases. coneveau filtration, Milnor operation, classifying spaces.

Here we give an example that we can compute a nonzero $DH^*(X)$. For $G = (\mathbb{Z}/p)^3$ the elementary abelian p -group of $rank = 3$, we know (for p odd)

$$H^*(BG; \mathbb{Z}/p) \cong \mathbb{Z}/p[y_1, y_2, y_3] \otimes \Lambda(x_1, x_2, x_3), \quad |x_i| = 1, \quad Q_0(x_i) = y_i$$

$$H^*(BG)/p \cong \mathbb{Z}/p[y_1, y_2, y_3](1, Q_0(x_i x_j), Q_0(x_1 x_2 x_3) | 1 \leq i < j \leq 3)$$

where $Q_0 = \beta$ is the Bockstein operation, and the notation $R(a, \dots, b)$ (resp. $R\{a, \dots, b\}$) means the R -submodule (resp. the free R -module) generated by a, \dots, b .

Theorem 1.1. *Let $G = (\mathbb{Z}/p)^3$. Given $N > 2p + 3$, there are approximations $X = X(N)$ for BG such that we have for $\alpha_{ij} = Q_0(x_i x_j)$, $\alpha = Q_0(x_1 x_2 x_3)$*

$$DH^*(X) \cong DH^3(X) \oplus DH^4(X)$$

$$\cong \mathbb{Z}/p\{\alpha_{ij}, \alpha | 1 \leq i < j \leq 3\} \quad \text{when } * < N.$$

But we have $DH^*(X; \mathbb{Z}/p) = 0$ when $* < N$.

Benoist and Ottem give examples, where the above invariants are nonzero for $A = \mathbb{Z}_{(2)}$ by using the Steenrod squares. We note that some of their arguments can be extended for $A = \mathbb{Z}_{(p)}$ by using the Milnor operation Q_n for odd primes p . However it seems not so easy to give a nontrivial example for $A = \mathbb{Z}/p$ in the case X is an approximation for BG as the above examples show.

For connected groups we have

Theorem 1.2. *Let G be a simply connected group such that $H^*(BG)$ has p -torsion. Then there is an approximation X for BG such that $DH^4(X) \neq 0$.*

Theorem 1.3. *Let p be an odd prime number, and $G = PGL_p$. Then there is an approximation X for BG such that $DH^3(X) \neq 0$.*

Theorem 1.4. *Let $p = 2$ and $X_n = X_n(N)$ be approximations for BSO_n with $n \geq 3$. Then for sufficient large N*

$$DH^*(X_{2m+1}) \supset \mathbb{Z}/2\{w_3, w_5, \dots, w_{2m+1}\} \quad \text{for } * < N$$

where w_i is the i -th Stiefel-Whitney class.

2. TRANSFER AND Q_n

The Milnor operation (in $H^*(-; \mathbb{Z}/p)$) is defined by $Q_0 = \beta$ and for $n \geq 1$

$$Q_n = P^{\Delta_n} \beta - \beta P^{\Delta_n}, \quad \Delta_n = (0, \dots, 0, \overset{n}{1}, 0, \dots)$$

(For details see [8], §3.1 in [18]) where β is the Bockstein operation and P^α for $\alpha = (\alpha_1, \alpha_2, \dots)$ is the fundamental base of the module of finite sums of products of reduced powers. .

Lemma 2.1. *Let f_* be the transfer (Gysin) map (for proper smooth) $f : X \rightarrow Y$. Then $Q_n f_*(x) = f_* Q_n(x)$ for $x \in H^*(X; \mathbb{Z}/p)$.*

The above lemma is known (see the proof of Lemma 7.1 in [23]). The transfer f_* is expressed as $g^* f'_*$ such that

$$f'_*(x) = i^*(Th(1) \cdot x), \quad x \in H^*(X; \mathbb{Z}/p)$$

for some maps g, f', i and the Thom class $Th(1)$. Since $Q_n(Th(1)) = 0$ and Q_i is a derivation, we get the lemma. However, we give here the another computational proof.

Proof of Lemma 2.1. Recall the following Grothendieck formula (e.g., [Q1])

$$(1) \quad P_t(f_*(x)) = f_*(c_t \cdot P_t(x)).$$

Here the total reduced powers $P_t(x)$ are defined

$$P_t(x) = \sum_{\alpha} P^{\alpha}(x) t^{\alpha} \in H^*(X; \mathbb{Z}/p)[t_1, t_2, \dots] \quad \text{with } t^{\alpha} = t_1^{\alpha_1} t_2^{\alpha_2} \dots$$

where $\alpha = (\alpha_1, \alpha_2, \dots)$ and $\deg(t^{\alpha}) = \sum_i 2\alpha_i(p^i - 1)$ (each element in $H^*(X; \mathbb{Z}/p)$ is represented as a homogeneous part respective to the above degree). The total Chern class c_t is defined similarly, for the Chern classes of the normal bundle of the map f .

We consider the above equation with the assumption such that $t_n^2 = 0$ and $t_j = 0$ for $j \neq n$, i.e., $P_t(x) \in H^*(X; \mathbb{Z}/p) \otimes \Lambda(t_n)$. That means

$$(2) \quad P_t(f_*(x)) = (1 + P^{\Delta_n} t_n)(f_*(x))$$

$$(3) \quad f_*(c_t \cdot P_t(x)) = f_*((1 + c_{p^n-1} t_n)(x + P^{\Delta_n}(x) t_n)) = f_*(x + (c_{p^n-1} x + P^{\Delta_n}(x)) t_n).$$

From (1), we see (2) = (3) and we have

$$(4) \quad P^{\Delta_n}(f_*(x)) = f_*(c_{p^n-1} x + P^{\Delta_n}(x)).$$

By the definition, β commutes with f_* , and we have

$$(5) \quad P^{\Delta_n} \beta(f_*(x)) = P^{\Delta_n} f_*(\beta x) = f_*(c_{p^n-1} \beta x + P^{\Delta_n}(\beta x)).$$

On the other hand

$$(6) \quad \beta P^{\Delta_n} f_*(x) \stackrel{(4)}{=} \beta f_*(c_{p^n-1} x + P^{\Delta_n}(x)) = f_*(c_{p^n-1} \beta x + \beta P^{\Delta_n}(x)).$$

Then (5) – (6) gives that $(P^{\Delta_n} \beta - \beta P^{\Delta_n}) f_*(x) = f_*(P^{\Delta_n} \beta - \beta P^{\Delta_n})(x)$. \square

By the definition, each cohomology operation h (i.e., an element in the Steenrod algebra) is written (with $Q^B = Q_0^{b_0} Q_1^{b_1} \dots$) by

$$h = \sum_{A, B} P^A Q^B \quad \text{with } A = (a_1, \dots), \quad B = (b_0, \dots) \quad b_i = 0 \text{ or } 1.$$

Corollary 2.2. *We have $P_t Q^B(f_*(x)) = f_*(c_t \cdot P_t Q^B(x))$.*

Hence cohomology operations h (for $H^*(-; \mathbb{Z}/p)$) which commute with all transfer f_* are cases $c_t = 1$, i.e. $A = 0$ which are only products Q^B of Milnor operations Q_i .

3. CONIVEAU FILTRATIONS

Bloch-Ogus [2] give a spectral sequence such that its E_2 -term is given by

$$E(c)_2^{c, *-c} \cong H_{Zar}^c(X, \mathcal{H}_A^{*-c}) \implies H_{et}^*(X; A)$$

where \mathcal{H}_A^* is the Zariski sheaf induced from the presheaf given by $U \mapsto H_{et}^*(U; A)$ for an open $U \subset X$.

The filtration for this spectral sequence is defined as the coniveau filtration

$$N^c H_{et}^*(X; A) = F(c)^{c, *-c}$$

where the infinite term $E(c)_{\infty}^{c, *-c} \cong F(c)^{c, *-c} / F(c)^{c+1, *-c-1}$ and

$$N^c H_{et}^*(X; A) = \sum_{Z \subset X; \text{codim}_X(Z) \leq c} \ker(j^* : H_{et}^*(X; A) \rightarrow H_{et}^*(X - Z, A)).$$

Here we recall the motivic cohomology $H^{*,*'}(X; \mathbb{Z}/p)$ defined by Voevodsky and Suslin ([18],[20],[21]) so that

$$H^{i,i}(X; \mathbb{Z}/p) \cong H_{et}^i(X; \mathbb{Z}/p) \cong H^i(X; \mathbb{Z}/p).$$

Let us write $H_{et}^*(X; \mathbb{Z})$ simply by $H_{et}^*(X)$ as usual. Note that we do *not* assume $H_{et}^*(X) \cong H^*(X)$ while we have the natural map $H_{et}^*(X) \rightarrow H^*(X)$.

Let $0 \neq \tau \in H^{0,1}(Spec(\mathbb{C}); \mathbb{Z}/p)$. Then by the multiplying by τ , we can define a map $H^{*,*'}(X; \mathbb{Z}/p) \rightarrow H^{*,*'+1}(X; \mathbb{Z}/p)$. By Deligne (foot note (1) in Remark 6.4 in [2]) and Paranjape (Corollary 4.4 in [9]), it is proven that there is an isomorphism of the coniveau spectral sequence with the τ -Bockstein spectral sequence $E(\tau)_r^{*,*'} (see also [16], [22]).$

Theorem 3.1. (*Deligne*) *Let $A = \mathbb{Z}/p$. Then we have the isomorphism of spectral sequence*

$$E(c)_r^{c,*-c} \cong E(\tau)_{r-1}^{*,*'-c} \quad \text{for } r \geq 2.$$

Hence the filtrations are the same, i.e. $N^c H_{et}^(X; \mathbb{Z}/p) = F_\tau^{*,*'-c}$ where*

$$F_\tau^{*,*'-c} = Im(\times \tau^c : H^{*,*'-c}(X; \mathbb{Z}/p) \rightarrow H^{*,*}(X; \mathbb{Z}/p)).$$

Lemma 3.2. *Let $x \in H^{*,*}(X)$ and also in $N^c H^*(X; \mathbb{Z}/p)$. Then if the map $H^{*+1,*-c}(X) \rightarrow H^{*+1,*}(X)$ is injective, then $x \in N^c H^*(X)$.*

Proof. Consider the exact sequences

$$\begin{array}{ccccccc} \xrightarrow{p} & x'' \in H^{*,*'-c}(X) & \xrightarrow{r_1} & x' \in H^{*,*'-c}(X; \mathbb{Z}/p) & \xrightarrow{\delta_1} & H^{*+1,*-c}(X) \\ & \downarrow f_1 & & \downarrow f_2 & & \downarrow f_3 \\ \xrightarrow{p} & x \in H^{*,*}(X) & \xrightarrow{r_2} & H^{*,*}(X; \mathbb{Z}/p) & \xrightarrow{\delta_2} & H^{*+1,*}(X) \end{array}$$

By the assumption of this lemma, we can take $x' \in H^{*,*'-c}(X; \mathbb{Z}/p)$ such that $r_2(x) = f_2(x')$. So $\delta_2 f_2(x') = 0$. Since f_3 is injective, we see $\delta_1(x') = 0$. Hence there is $x'' \in H^{*,*'-c'}(X)$ such that $r_1(x'') = x'$. \square

Let $cl : CH^*(X) \otimes A \rightarrow H^{2*}(X; A)$ be the cycle map. and $Im(cl)^+$ is the positive degree parts of its image.

Lemma 3.3. *We see that $Im(cl)^+ \subset N^* H^{2*}(X; A)$.*

Proof. Recall that $H^{*,*'}(X; A) \subset N^{*-*'} H^*(X; A)$. We have $H^{2*,*}(X; A) \cong CH^*(X) \otimes A$. Since $2* > *$ for $* \geq 1$, we see $cl(y) \in N^1 H^{2*}(X; A)$. \square

Hence $y \in CH^*(X) \otimes A$ is represented by closed algebraic set supported Y , while Y may be singular.

On the other hand, by Totaro [17], we have the modified cycle map \bar{cl}

$$cl : CH^*(X) \otimes A \xrightarrow{\bar{cl}} MU^{2*}(X) \otimes_{MU^*} A \xrightarrow{\rho} H^{2*}(X; A)$$

for the complex cobordism theory $MU^*(X)$. It is known [11] that elements in $MU^{2*}(X)$ is written by a proper map to X from a stable almost complex manifold Y . while it does not mean complex manifold.

The following lemma is well known.

Lemma 3.4. *If $x \in Im(\rho)$ for $\rho : MU^*(X)/p \rightarrow H^*(X; \mathbb{Z}/p)$, then we have $Q_i(x) = 0$ for all $i \geq 0$.*

Proof. In fact, it is known that $d = Q_i$ for the first nonzero differential d of the Atiyah-Hirzebruch spectral sequence converging to the connected Morava K-theory $k(i)^*(X)$ with $k(i)^* = \mathbb{Z}/p[v_i]$. We have the natural maps

$$\rho : MU^*(X)/p \xrightarrow{\rho_1} k(i)^*(X) \xrightarrow{\rho_2} H^*(X; \mathbb{Z}/p).$$

Hence $Q_i \rho_2(x) = 0$ implies $Q_i \rho(x) = 0$. \square

Lemma 3.5. (*reciprocity law*) *If $a \in \tilde{N}^* H^{2*}(X; A)$, then for each $g \in DH^{*'}(X; A)$ we have $ag \in \tilde{N}^* H^{2*+*'}(X; A)$.*

Proof. Suppose we have $f : Y \rightarrow X$ with $f_*(a') = a$. Then

$$f_*(a' f^*(g)) = f_*(a')g = ag$$

by Frobenius reciprocity law. \square

Let G be an algebraic group (over \mathbb{C}) and r be a complex representation $r : G \rightarrow U_n$ the unitary group. Then we can define the Chern class $c_i = r^* c_i^U$. Here the Chern classes c_i^U in $H^*(BU_n) \cong \mathbb{Z}[c_1^U, \dots, c_n^U]$ are defined by using the Gysin map as $c_n^U = i_{n,*}(1)$ for

$$i_n^* : H^*(BU_n) \cong H_{U_n}^*(pt.) \xrightarrow{i_n^*} H_{U_n}^{*+2i}(\mathbb{C}^{\times i}) \cong H^{*+2i}(BU_n)$$

where $H_{U_n}(X) = H^*(EU_n \times_{U_n} X)$ is the U_n -equivariant cohomology.

Lemma 3.6. *For an approximation X of BG and for each $g \in H^{*'}(BG; A)$, we have $gc_i = 0 \in DH^*(X; A)$.*

The following lemma is proved by Colliot Thérène and Voisin [3] by using the affirmative answer of the Bloch-Kato conjecture by Voevodsky. ([20]. [21])

Lemma 3.7. ([3]) *Let X be a smooth complex variety. Then any torsion element in $H^*(X)$ is in $N^1 H^*(X)$.*

4. THE MAIN LEMMAS

The following lemma is the Q_i -version of one of results by Benoist and Ottem.

Lemma 4.1. *Let $\alpha \in N^1 H^s(X)$ for $s = 3$ or 4 . If $Q_i(\alpha) \neq 0 \in H^*(X; \mathbb{Z}/p)$ for some $i \geq 1$, then*

$$DH^s(X) \supset \mathbb{Z}/p\{\alpha\}, \quad DH^s(X; \mathbb{Z}/p^t) \supset \mathbb{Z}/p\{\alpha\} \quad \text{for } t \geq 2.$$

Proof. Suppose $\alpha \in \tilde{N}^1 H^s(X)$ for $s = 3$ or 4 , i.e. there is a smooth Y with $f : Y \rightarrow X$ such that the transfer $f_*(\alpha') = \alpha$ for $\alpha' \in H^*(Y)$. Then for $s = 4$

$$\begin{aligned} Q_i(\alpha') &= (P^{\Delta_i} \beta - \beta P^{\Delta_i})(\alpha') = (-\beta P^{\Delta_i})(\alpha') = -\beta(\alpha')^{p^i} \\ &= -p^i(\beta\alpha')(\alpha')^{p^i-1} = 0 \quad (\text{by Cartan formula}) \end{aligned}$$

since $\beta(\alpha') = 0$ and $P^{\Delta_i}(y) = y^{p^i}$ for $\dim(y) = 2$. (For $s = 3$, we get also $Q_i(\alpha') = 0$ since $P^{\Delta_i}(x) = 0$ for $\dim(x) = 1$.) This contradicts to the commutativity of Q_i and f_* .

The case $A = \mathbb{Z}/p^t$, $t \geq 2$ is proved similarly, since for $\alpha' \in H^*(X; A)$ we see $\beta\alpha' = 0 \in H^*(X; \mathbb{Z}/p)$. \square

We will extend Lemma 4.1, $s = 4$, for the larger s , by using MU -theory of Eilenberg-MacLane spaces.

Recall that $K = K(\mathbb{Z}, n)$ is the Eilenberg-MacLane space such that the homotopy group $[X, K] \cong H^n(X; \mathbb{Z})$, i.e., each element $x \in H^n(X; \mathbb{Z})$ is represented by a homotopy map $x; X \rightarrow K$. Let $\eta_n \in H^n(K; \mathbb{Z})$ corresponding the identity map. (For $K' = K(\mathbb{Z}/p, n)$ define $\eta'_n \in H^n(K'; \mathbb{Z}/p)$ by the identity element of K' .) We know the image $\rho(MU^*(K)) \subset H^*(K; \mathbb{Z})/p$.

Lemma 4.2. ([15], [13]) *We have the isomorphism*

$$\rho : MU^*(K) \otimes_{MU^*} \mathbb{Z}/p \cong \mathbb{Z}/p[Q_{i_1} \dots Q_{i_{n-2}} \eta_n | 0 < i_1 < \dots < i_{n-2}],$$

$$\rho : MU^*(K') \otimes_{MU^*} \mathbb{Z}/p \cong \mathbb{Z}/p[Q_{i_1} \dots Q_{i_{n-1}} Q_0 \eta'_n | 0 < i_1 < \dots < i_{n-1}],$$

where the notation $\mathbb{Z}/p[a, \dots]$ exactly means $\mathbb{Z}/p[a, \dots]/(a^2 | |a| = \text{odd})$.

The following lemma is an extension of Lemma 4.1 for $s \geq 4$.

Lemma 4.3. *Let $\alpha \in N^c H^{n+2c}(X)$, $n \geq 2$, $c \geq 1$. Suppose that there is a sequence $0 < i_1 < \dots < i_{n-1}$ with*

$$Q_{i_1} \dots Q_{i_{n-1}} \alpha \neq 0 \quad \text{in } H^*(X; \mathbb{Z}/p).$$

Then $D^c H^(X) = N^c H^*(X)/(p, \tilde{N}^c H^*(X)) \supset \mathbb{Z}/p\{\alpha\}$.*

Proof. Suppose $\alpha \in \tilde{N}^c H^{n+2c}(X)$, i.e. there is a smooth Y of $\dim(Y) = \dim(X) - c$ with $f : Y \rightarrow X$ such that the transfer $f_*(\alpha') = \alpha$ for $\alpha' \in H^n(Y)$.

Identify the map $\alpha' : Y \rightarrow K$ with $\alpha' = (\alpha')^* \eta_n$. We still see from Lemma 4.2,

$$Q(\alpha') = Q_{i_1} \dots Q_{i_{n-2}} ((\alpha')^* \eta_n) \in \text{Im}(MU^*(Y) \rightarrow H^*(Y; \mathbb{Z}/p)).$$

From Lemma 3.4, we see

$$Q_{i_{n-1}} Q(\alpha') = Q_{i_{n-1}} Q_{i_1} \dots Q_{i_{n-2}} (\alpha') = 0 \in H^*(Y; \mathbb{Z}/p).$$

Therefore $Q_{i_{n-1}} Q(\alpha)$ must be zero by the commutativity of f_* and Q_i . \square

Remark. For $\alpha \in N^c H^{n+2c}(X; \mathbb{Z}/p)$, we can give the $A = \mathbb{Z}/p$ version of the above lemma from the second isomorphism in Lemma 4.2. But we can see $Q_{i_1} \dots Q_{i_n} Q_0 \alpha = 0$ always, (even if $Q_{i_1} \dots Q_{i_{n-1}} \alpha \neq 0$). Hence we do not say about $D^c H^*(X; \mathbb{Z}/p)$.

5. CLASSIFYING SPACES FOR FINITE GROUPS

Let G be a finite group or an algebraic group, and BG its classifying space. For example, when $G = G_m$ is the multiple group, we see

$$BG_m = BS^1 \cong \mathbb{P}^\infty, \quad H^*(\mathbb{P}^\infty) \cong \mathbb{Z}[y] \quad \text{with } |y = c_1| = 2,$$

for the infinite (complex) projective space \mathbb{P}^∞ . Note that BG_m is a colimit of complex projective spaces. Though BG itself is not a colimit of complex projective varieties, we can take a complex projective variety $X(N)$ ([4]) for a given $N \geq 3$ such that there is a map $j : X(N) \rightarrow BG \times \mathbb{P}^\infty$ with

$$H^*(BG \times \mathbb{P}^\infty; A) \xrightarrow{j^*} H^*(X(N); A) \quad \text{for } * < N.$$

Note that the quotient

$$N^n H^*(X; A) / (\tilde{N}^n H^*(X; A))$$

is invariant under replacing X with $X \times \mathbb{P}^m$ for all n and all abelian groups A . In fact, from Künneth formula,

$$H^*(X \times \mathbb{P}^m; A) \cong H^*(X; A) \otimes \mathbb{Z}[y]/(y^{m+1}),$$

where $y \in CH^1(\mathbb{P}^m)$. Let $Ideal(y)$ be the ideal of $H^*(X \times \mathbb{P}^m; A)$ generated by y . Then $Ideal(y) \subset \tilde{N}^* H^*(X \times \mathbb{P}^m; A)$ by the Frobenius reciprocity law (Lemma 3.5). Moreover Benoist and Ottem show the above quotient when $n = 1$ is a stable birational invariant (Proposition 2.4 in [1]).

In this paper, we say the above $X(N)$ as a (*degree* $\leq N$) complex projective approximation *for* BG (which is an approximation *of* $BG \times \mathbb{P}^\infty$ strictly speaking), and let us write the (*mod*(p)) stable rational invariant by

$$DH^*(X; A) = N^1 H^*(X; A)/(p, \tilde{N}^1 H^*(X; A)).$$

Hereafter, we consider $DH^*(X; A)$ when $A = \mathbb{Z}$. Let p be an odd prime. (The case $p = 2$ is different but a similar argument works.) Let $G = (\mathbb{Z}/p)^3$ the *rank* $= 3$ elementary abelian p -group. Then the *mod*(p) cohomology is

$$H^*(BG; \mathbb{Z}/p) \cong H^*(B\mathbb{Z}/p; \mathbb{Z}/p)^{3\otimes} \cong \mathbb{Z}/p[y_1, y_2, y_3] \otimes \Lambda(x_1, x_2, x_3).$$

Here $|y_i| = 2, |x_i| = 1, \beta(x_i) = y_i$, and $\Lambda(a, \dots, b)$ is the \mathbb{Z}/p -exterior algebra generated by a, \dots, b .

The integral cohomology (modulo p) is isomorphic to

$$H^*(BG)/p \cong \text{Ker}(Q_0) \cong H(H^*(BG; \mathbb{Z}/p); Q_0) \oplus \text{Im}(Q_0)$$

where $H(-; Q_0) = \text{Ker}(Q_0)/\text{Im}(Q_0)$ is the homology with the differential Q_0 . It is immediate that $H(H^*(B\mathbb{Z}/p; \mathbb{Z}/p); Q_0) \cong \mathbb{Z}/p$. By the Künneth formula, we have $H(H^*(BG; \mathbb{Z}/p); Q_0) \cong (\mathbb{Z}/p)^{3\otimes} \cong \mathbb{Z}/p$. Hence we have

$$H^*(BG)/p \cong \mathbb{Z}/p\{1\} \oplus \text{Im}(Q_0)$$

$$\cong \mathbb{Z}/p[y_1, y_2, y_3](1, Q_0(x_i x_j), Q_0(x_1 x_2 x_3) | i < j)$$

where the notation $R(a, \dots, b)$ (resp. $R\{a, \dots, b\}$) means the R -submodule (resp. the free R -module) generated by a, \dots, b . Here we note $H^+(BG)$ is just p -torsion.

Also note that y_1, y_2, y_3 are represented by the Chern classes c_1 . From Lemma 3.6, we see

$$Ideal(y_1, y_2, y_3) = 0 \in DH^*(X).$$

We know $Q_i(y_j) = y_j^{p^i}$ and Q_j is a derivation. Let us write

$$\alpha = Q_0(x_1 x_2 x_3) = y_1 x_2 x_3 - y_2 x_1 x_3 + y_3 x_1 x_2.$$

Note $\alpha \in H^4(X)$, $p\alpha = 0$, and $\alpha \in N^1 H^*(X)$ from Lemma 3.7. Moreover

$$Q_1(\alpha) = Q_1(y_1 x_2 x_3) - \dots = y_1 y_2^p x_3 - y_1 y_3^p x_2 - \dots \neq 0 \in H^*(X; \mathbb{Z}/p).$$

Similarly, for $\alpha_{ij} = Q_0(x_i x_j)$, we see $Q_1(\alpha_{ij}) \neq 0$. Hence from Lemma 4.1 and Lemma 3.6, we have

Theorem 5.1. *Let $X = X(N)$ with $N > 2p+3$ be an approximation for $B(\mathbb{Z}/p)^3$. Then we have*

$$DH^*(X) \cong \mathbb{Z}/p\{\alpha_{ij}, \alpha | 1 \leq i < j \leq 3\} \quad \text{for } * < N.$$

Proof. We see $H^*(BG)/(p, y_1, y_2, y_3) \cong \mathbb{Z}/p\{1, \alpha_{ij}, \alpha\}$. Of course $1 \notin N^1 H^*(X)$, we have the theorem from Lemma 4.1. \square

Theorem 5.2. *Let $X = X(N)_n$ be an approximation for $(B\mathbb{Z}/p)^n$ with $N > |Q_0 Q_1 \dots Q_{n-1}(x_1 \dots x_n)|$. Then we have for $\alpha_{i_1, \dots, i_s} = Q_0(x_{i_1} \dots x_{i_s})$,*

$$DH^*(X) \supset \mathbb{Z}/p\{(\alpha_{i_1, \dots, i_s} | 2 \leq s, 0 < i_1 < i_2 \dots < i_s \leq n\} \quad \text{for } * < N.$$

Proof. We have the theorem from Lemma 4.3 and $Q_{i_1} \dots Q_{i_{s-2}}(\alpha_{i_1, \dots, i_s})$ is

$$Q_{i_1} \dots Q_{i_{s-2}} Q_0(x_{i_1} \dots x_{i_s}) = y_{i_1}^{p^{i_1}} \dots y_{i_{s-2}}^{p^{i_{s-2}}} y_{i_{s-1}} x_{i_s} + \dots \neq 0.$$

(Note the $n = |\alpha'|$ in Lemma 4.3 is written by $s - 1$ here.) \square

Corollary 5.3. *If $n \neq m \geq 3$, then $X(N)_n$ and $X(N)_m$ are not stable birational equivalent.*

Next we study small non abelian p -groups. Let G be a non abelian group of order p^3 (see §8, for details). Then $H^{even}(BG)$ is generated by Chern classes, and $H^{odd}(BG)$ is a (just) p -torsion. We can identify $H^{odd}(BG) \subset H^{odd}(BG; \mathbb{Z}/p)$. The operation Q_1 acts on $H^{odd}(X)$, and induces the injection

$$Q_1 : H^{odd}(BG) \hookrightarrow H^{even}(BG).$$

Such groups are four types (see §8 below), and they are called extraspecial p -groups $G = p_{\pm}^{1+2}$ of order p^3 . When $G = Q_8 = 2_+^{1+2}$ the quaternion group of order 8, we know $H^{odd}(X) = 0$. However when $G = D_8 = 2_+^{1+2}$ the dihedral group of order 8, the cohomology $H^{odd}(BG)$ is generated as an $H^{even}(BG)$ module by an element e of $\deg(e) = 3$. When $G = E = p_+^{1+2}$ for $p \geq 3$, $H^{odd}(BG)$ is generated by e_1, e_2 with $\deg(e_i) = 3$. When $G = M = p_-^{1+2}$ for $p \geq 3$, $H^{odd}(BG)$ is generated by e' but $\deg(e') = 2p + 1$.

From Lemma 3.5 (Frobenius reciprocity) and the main lemma (Lemma 4.1), we have the following theorem.

Theorem 5.4. *Let $X = X(N)$ with $N > 2p + 3$ be an approximation for an extraspecial p -group G of order p^3 . Then we have for $* < N$*

$$DH^*(X) \cong \begin{cases} 0 & \text{for } G = Q_8 \\ \mathbb{Z}/2\{e\} & \text{for } G = D_8 \\ 0 \text{ or } \mathbb{Z}/p\{e'\} & \text{for } G = M \\ \mathbb{Z}/p\{e_1, e_2\} & \text{for } G = E. \end{cases}$$

In particular, the above theorem implies that when $G = p_+^{1+2}$, there is an X with $DH^3(X) \neq 0$ but for all $X = X(N)$ we see $DH^*(X) = 0$ for $4 \leq * < N$.

6. CONNECTED GROUPS

At first, we consider when $G = U_n$, SU_n or Sp_{2n} for all p , where the cohomology $H^*(BG)$ has no torsion. Then $H^*(BG)$ is generated by Chern classes, e.g.,

$$H^*(BU_n) \cong CH^*(BU_n) \cong \mathbb{Z}_{(p)}[c_1, \dots, c_n],$$

$$H^*(BSp_{2n}) \cong CH^*(BSp_{2n}) \cong \mathbb{Z}_{(p)}[c_2, c_4, \dots, c_{2n}].$$

Hence $DH^*(X) = 0$ for the approximations X for these groups.

Next we consider the case $G = SO_3$ and $p = 2$. Then

$$H^*(BG; \mathbb{Z}/2) \cong \mathbb{Z}/2[w_1, w_2, w_3]/(w_1) \cong \mathbb{Z}/2[w_2, w_3],$$

where w_i is the i -th Stiefel-Whitney class for $SO_3 \subset O_3$ and $w_i^2 = c_i$ is the i -th Chern class for $SO_3 \subset U_3$. (Also it is the elementary symmetric polynomial in $\mathbb{Z}/2[y_1, \dots, y_i]$.) Here we know $Q_0(w_2) = w_3$, and $Q_1(w_3) = w_3^2 = c_3$. Therefore we have [22]

$$\begin{aligned} H^*(BG; \mathbb{Z}/2) &\cong \mathbb{Z}/2[c_2, c_3]\{1, w_2, w_3 = Q_0(w_2), w_2 w_3 = Q_1 w_2\} \\ &\cong \mathbb{Z}/2[c_2, c_3] \otimes \Lambda(Q_0, Q_1)\{w_2\} \oplus \mathbb{Z}/2[c_2]. \end{aligned}$$

In particular $H^*(BG)/2 \cong \text{Ker}(Q_0) \cong \mathbb{Z}/2[c_2, c_3]\{1, w_3\}$. Then from Lemma 4.1, we have

Theorem 6.1. *Let $G = SO_3$ and X be an approximation of BG for $6 < N$. Then $DH^*(X) \cong \mathbb{Z}/2\{w_3\}$ for $* < N$.*

Using Lemma 4.3, we have

Theorem 6.2. *Let $X_n = X_n(N)$ be approximations for BSO_n for $n \geq 3$. Moreover let $|Q_1 \dots Q_{2m-1}(w_{2m+1})| < N$. Then*

$$DH^*(X_{2m+1}) \supset \mathbb{Z}/2\{w_3, w_5, \dots, w_{2m+1}\} \quad \text{for } * < N.$$

Proof. Since $Q_0 w_{2i} = w_{2i+1}$, we see $w_{2i+1} \in N^1 H^{2i+1}(X)$. We have the theorem, from Lemma 4.3 and the restriction to $H^*(B(\mathbb{Z}/2)^{2i}; \mathbb{Z}/2)$,

$$Q_1 \dots Q_{2i-2}(w_{2i+1}) = Q_1 \dots Q_{2i-2} Q_0(w_{2i}) = y_1 y_2^2 \dots y_{2i-1}^{2^{2i-2}} x_{2i} + \dots \neq 0.$$

□

We next consider simply connected groups. Let us write by X an approximation of BG_2 for the exceptional simple group G_2 of $\text{rank} = 2$. The $\text{mod}(2)$ cohomology is generated by the Stiefel-Whitney classes w_i of the real representation $G_2 \rightarrow SO_7$

$$H^*(BG_2; \mathbb{Z}/2) \cong \mathbb{Z}/2[w_4, w_6, w_7], \quad P^1(w_4) = w_6, \quad Q_0(w_6) = w_7,$$

$$H^*(BG_2) \cong (D' \oplus D'/2[w_7]^+) \quad \text{where } D' = \mathbb{Z}[w_4, c_6].$$

Then we have $Q_1 w_4 = w_7, Q_2(w_7) = w_7^2 = c_7$ (the Chern class).

The Chow ring of BG_2 is also known

$$CH^*(BG_2) \cong (D\{1, 2w_4\} \oplus D/2[c_7]^+) \quad \text{where } D = \mathbb{Z}[c_4, c_6] \quad c_i = w_i^2.$$

In particular the cycle map $cl : CH^*(BG) \rightarrow H^*(BG)$ is injective.

It is known $w_4 \in N^1 H^*(X; \mathbb{Z}/2)$ ([22]) and from Lemma 3.2, we see $w_4 \in N^1 H^*(X)$. Since $Q_1(w_4) = w_7 \neq 0$, from Lemma 4.1, we have $DH^4(X) \neq 0$ ([1]). Moreover $H^*(BG)/(c_4, c_6, c_7) \cong \Lambda(w_4, w_7)$ implies

Proposition 6.3. *For X an approximation for BG_2 , we have the surjection*

$$\Lambda(w_4, w_7)^+ \twoheadrightarrow DH^*(X) \quad \text{for } * < N.$$

By Voevodsky [18], [19], we have the Q_i operation also in the motivic cohomology $H^{*,*'}(X; \mathbb{Z}/p)$ with $\deg(Q_i) = (2p^i - 1, p - 1)$. Then we can take

$$\deg(w_4) = (4, 3), \quad \deg(w_6) = (6, 4), \quad \deg(w_7) = (7, 4), \quad \deg(c_7) = (14, 7).$$

By Theorem 3.1, the above means

$$w_7 = Q_1 w_4 \in N^{7-4} H^*(X; \mathbb{Z}/2) = N^3 H^*(X; \mathbb{Z}/2).$$

We can not see here that $0 \neq w_7 \in DH^*(X)$, but see the following proposition.

Proposition 6.4. *For X the approximation for BG_2 , we have*

$$\mathbb{Z}/2\{w_7\} \subset D^3H^*(X) = N^3H^*(X)/(2, \tilde{N}^3H^*(X)).$$

Proof. Suppose $w_7 \in \tilde{N}^3H^*(X)$. That is, there is $x \in H^1(Y)$ with $f_*(x) = w_7$ for $f : Y \rightarrow X$. Act Q_2 on $H^*(Y; \mathbb{Z}/2)$, and

$$Q_2(x) = (P^{\Delta_2}\beta + \beta P^{\Delta_2})(x) = 0$$

since $\beta(x) = 0$ and $P^i(x) = Sq^{2i}(x) = 0$ for $i > 0$. But $Q_2w_7 = c_7 \neq 0$. This contradicts to the commutativity of f_* and Q_2 . \square

Theorem 6.5. *Let G be a simply connected group such that $H^*(BG)$ has p -torsion. Let $X = X(N)$ be an approximation for BG for $N \geq 2p + 3$. Then $DH^4(X) \neq 0$.*

Proof. It is only need to prove the theorem when G is a simple group having p torsion in $H^*(BG)$. Let $p = 2$. It is well known that there is an embedding $j : G_2 \subset G$ such that (see [10], [25] for details)

$$H^4(BG) \xrightarrow{j^*} H^4(BG_2) \cong \mathbb{Z}\{w_4\}.$$

Let $x = (j^*)^{-1}w_4 \in H^4(BG)$. From Lemma 3.1 in [25], we see that $2x$ is represented by Chern classes. Hence $2x$ is the image from $CH^*(X)$, and so $2x \in N^1H^4(X)$. This means there is an open set $U \subset X$ such that $2x = 0 \in H^*(U)$ that is. x is 2-torsion in $H^*(U)$. Hence from Lemma 3.5, we have $x \in N^1H^4(U)$, and so there is $U' \subset U$ such that $x = 0 \in H^4(U')$. This implies $x \in N^1H^4(X)$.

Since $j^*(Q_1x) = Q_1w_4 = w_7 \neq 0$, we see $Q_1x \neq 0$. From the main lemma (Lemma 4.1), we see $DH^4(X) \neq 0$ for G .

For the cases $p = 3, 5$, we consider the exceptional groups F_4, E_8 respectively. Each simply connected simple group G contains F_4 for $p = 3$, E_8 for $p = 5$. There is $x \in H^4(BG)$ such that px is a Chern class and $Q_1(x) \neq 0 \in H^*(BG; \mathbb{Z}/p)$. In fact, there is embedding $j : (\mathbb{Z}/p)^3 \subset G$ with $j^*(x) = Q_0(x_1x_2x_3)$. Hence we have the theorem. \square

Corollary 6.6. *Let X be an approximation for $BSpin_n$ with $n \geq 7$ or BG for an exceptional group G . Then X is not stable rational.*

Remark. Kordonskii [6], Merkurjev (Corollary 5.8 in [7]), and Reichstein-Scavia show [14] that $BSpin_n$ itself is stably rational when $n \leq 14$. These facts do not mean contradictions. The (Ekedahl) approximation X is not stable rationally equivalent to $GL_M/G = BG$ where M is a large number such that G acts freely on GL_M . On the other hand X is constructed from a quasi projective variety BG as taking intersections of subspaces of $\mathbb{P}^{M'}$ for a large M' . (The author thanks Federico Scavia who pointed out this remark.)

At last of this section, we consider the case $G = PGL_p$. We have (for example Theorem 1.5, 1.7 in [5]) additively

$$H^*(BG; \mathbb{Z}/p) \cong M \oplus N \quad \text{with } M \xrightarrow{add.} \mathbb{Z}/p[x_4, x_6, \dots, x_{2p}],$$

$$N = SD \otimes \Lambda(Q_0, Q_1)\{u_2\} \quad \text{with } SD = \mathbb{Z}/p[x_{2p+2}, x_{2p^2-2p}]$$

where $x_{2p+2} = Q_1Q_0u_2$ and suffix means its degree. The Chow ring is given as

$$CH^*(BG)/p \cong M \oplus SD\{Q_0Q_1(u_2)\}.$$

From Lemma 4.1, we have ;

Theorem 6.7. *Let p be odd. For an approximation X for $BPGL_p$, we see $\mathbb{Z}/p\{Q_0u_2\} \subset DH^*(X)$, and moreover there is a surjection*

$$\mathbb{Z}/p[x_{2p^2-2p}]\{Q_0u_2\} \twoheadrightarrow DH^*(X)/(Im(cl)) \quad \text{for } * < N$$

for the cycle map $cl : CH^*(X) \rightarrow H^{2*}(X)$.

7. \mathbb{Z}/p -COEFFICIENT COHOMOLOGY FOR ABELIAN GROUPS

In the preceding sections, we have seen that cases $DH^*(X; A) \neq 0$ are not so rare for $A = \mathbb{Z}_{(p)}$, \mathbb{Z}/p^i , $i \geq 2$. However currently it seems difficult to make such example for $A = \mathbb{Z}/p$. (Recall the remark in the last in §4.)

Question 7.1. $DH^*(X; \mathbb{Z}/p) = 0$ for each smooth projective variety X ?

At first, we consider the case $G = (\mathbb{Z}/p)^3$. We can take a *quasi* projective approximation $\bar{X}(N)$ of $B\mathbb{Z}/p$ explicitly by the quotient (the N -dimensional lens space)

$$\bar{X}(N) = \mathbb{C}^{N*}/(\mathbb{Z}/p) \quad \text{where } \mathbb{C}^{N*} = (\mathbb{C}^N - \{0\}).$$

Next we consider the projective approximation

$$X(N) \rightarrow \bar{X}(N) \times \mathbb{P}^N \rightarrow B\mathbb{Z}/p \times \mathbb{P}^\infty.$$

Let us write X_i (resp. X'_i) the above $\bar{X}(N)$ (resp. $\bar{X}(N-1)$) for a sufficient large number N . Let

$$i_1 : Y_1 = X'_1 \times X_2 \times X_3 \rightarrow X = X_1 \times X_2 \times X_3.$$

Similarly we define Y_2, Y_3 , and the disjoint union $Y = Y_1 \sqcup Y_2 \sqcup Y_3$.

Recall that for p : odd

$$H^*(X; \mathbb{Z}/p) \cong \mathbb{Z}/p[y_1, y_2, y_3]/(y_1^{N+1}, y_2^{N+1}, y_3^{N+1}) \otimes \Lambda(x_1, x_2, x_3),$$

and $H^*(Y_i; \mathbb{Z}/p) \cong H^*(X; \mathbb{Z}/p)/(y_i^N)$ for $i = 1, 2, 3$. For $p = 2$, some graded ring $grH^*(X; \mathbb{Z}/2)$ is isomorphic to the above ring (in fact $x_i^2 = y_i$).

For the embedding $f_i : X'_i \rightarrow X_i$, it is known $f_{i*}(1) = c_1(N_i)$ where N_i is the normal bundle for $X'_i \subset X_i$. Hence the transfer is given by

$$f_{1*}(1) = y_1, \quad f_{2*}(1) = y_2, \quad f_{3*}(1) = y_3.$$

Therefore we have for $x = (x_2x_3 + x_3x_1 + x_1x_2) \in H^*(Y_1 \sqcup Y_2 \sqcup Y_3; \mathbb{Z}/p)$,

$$f_*(x) = y_1x_2x_3 + y_2x_3x_1 + y_3x_1x_2 = Q_0(x_1x_2x_3) = \alpha.$$

(Note that the element $x = (x_1x_2 + x_2x_3 + x_3x_1)$ is not in the integral cohomology $H^*(Y)$.) Thus we see $\alpha \in \tilde{N}^c H^*(X; \mathbb{Z}/p)$. More generally, we see

Theorem 7.2. *We have the approximation $X = X(N)$ for $(B\mathbb{Z}/p)^n$ such that*

$$DH^*(X; \mathbb{Z}/p) = 0 \quad * < N.$$

We recall here the motivic cohomology. By Voevodsky [Vo 1], $H^{*,*'}(B\mathbb{Z}/p; \mathbb{Z}/p)$ satisfies the Künneth formula so that (for p odd)

$$H^{*,*'}(B(\mathbb{Z}/p)^n; \mathbb{Z}/p) \cong \mathbb{Z}/p[\tau, y_1, \dots, y_n]/(y_1^{N+1}, \dots, y_n^{N+1}) \otimes \Lambda(x_1, \dots, x_n).$$

Here $0 \neq \tau \in H^{0,1}(Spec(\mathbb{C}); \mathbb{Z}/p)$, and $deg(y_i) = (2, 1)$, $deg(x_i) = (1, 1)$.

From Theorem 3.1, we can identify $N^c H_{et}^*(X; \mathbb{Z}/p) = F_\tau^{*,*-c}$ where $F_\tau^{*,*-c} = Im(\times \tau^c : H^{*,*-c}(X; \mathbb{Z}/p) \rightarrow H^{*,*}(X; \mathbb{Z}/p))$.

Corollary 7.3. (Theorem 5.1 in [16]) Let $X = X(N)$ be the approximation for $(B\mathbb{Z}/p)^n$ for a sufficient large N . Then we have

$$H^*(X; \mathbb{Z}/p)/N^1 H^*(X; \mathbb{Z}/p) \cong \Lambda(x_1, \dots, x_n).$$

Proof. Let $x \in \text{Ideal}(y_1, \dots, y_n) \subset H^{*,*'}(X; \mathbb{Z}/p)$. Then $\deg(x) = (*, *')$ with $* > *'$, and x is a multiplying of τ . Hence $x \in N^1 H^*(X; \mathbb{Z}/p)$. \square

Proof of Theorem 7.2. Let $x \in N^1 H^*(X; \mathbb{Z}/p)$. From the above corollary, $x \in \text{Ideal}(y_1, \dots, y_n)$ which is in the image of the transfer (such as $Y_1 = X'_1 \times X_2 \times \dots \times X_n$). That is $x \in \tilde{N}^1 H^*(X; \mathbb{Z}/p)$. \square

We can extend Lemma 7.3, by using the following lemma. Let us write by XG an approximation for BG .

Lemma 7.4. Let G have a Sylow p -subgroup S . If $DH^*(XS; \mathbb{Z}/p) = 0$, then so for BG .

Proof. Let $j : BS \rightarrow BG$ and $i : Y \rightarrow XS$. We consider maps

$$H^*(Y; \mathbb{Z}/p) \xrightarrow{i_*} H^*(XS; \mathbb{Z}/p) \xrightarrow{j^*} H^*(XG; \mathbb{Z}/p).$$

Here $j_* = \text{cor}_S^G$ is the transfer for finite groups.

Note $j^* N^1 H^*(XG; \mathbb{Z}/p) \subset N^1 H^*(XS; \mathbb{Z}/p)$ by the naturality of j^* . Hence given $x \in N^1 H^*(XG; \mathbb{Z}/p)$, the element $y = j^*(x)$ is in $N^1 H^*(XS; \mathbb{Z}/p)$.

By the assumption in this lemma, there is $y' \in H^*(Y; \mathbb{Z}/p)$ with $i_*(y') = y$. Thus we have $j_* i_*(y') = j_* y = j_* j^*(x) = [G; S]x$. \square

Similarly, we can prove ;

Corollary 7.5. Let G have an abelian p Sylow subgroup, and $X = X(N)$ be an approximation for BG . Then $DH^*(X; \mathbb{Z}/p) = 0$ for $* < N$.

8. THE GROUPS Q_8 AND D_8

When $|G| = p^3$, we have the short exact sequence

$$0 \rightarrow C \rightarrow G \rightarrow V \rightarrow 0$$

where $C \cong \mathbb{Z}/p$ is in the center and $V \cong \mathbb{Z}/p \times \mathbb{Z}/p$. Let us take generators such that $C = \langle c \rangle$, $V = \langle a, b \rangle$. Moreover we can take $[a, b] = c$ when G is non-abelian.

There are two cases, when $p = 2$, the quaternion group Q_8 and the dihedral group D_8 . We will show here

Theorem 8.1. Let $X = X(N)$ be an approximation for Q_8 or D_8 . Then $DH^*(X; \mathbb{Z}/2) = 0$ for $* < N$.

8.1. The case $G = Q_8$. Then $a^2 = b^2 = c$. Its cohomologies are well known (see [12])

$$\begin{aligned} H^*(BG)/2 &\cong \mathbb{Z}/2[y_1, y_2, c_2]/(y_i^2, y_1 y_2) \quad |y_i| = 2, \\ H^*(BG; \mathbb{Z}/2) &\cong \mathbb{Z}/2[x_1, x_2, c_2]/(x_1 x_2 + y_1 + y_2, x_1 y_2 + x_2 y_1) \\ &\cong \mathbb{Z}/2\{1, x_1, y_1, x_2, y_2, w\} \otimes \mathbb{Z}/2[c_2] \end{aligned}$$

where $x_i^2 = y_i$, $|x_i| = 1$, and $w = y_1 x_2 = y_2 x_1$, $|w| = 3$.

Therefore, we see

$$H^*(BG; \mathbb{Z}/2)/(y_1, y_2, c_2) \cong \mathbb{Z}/2\{1, x_1, x_2\}.$$

Of course $\deg(x_i) = (1, 1)$ in $H^{*,*'}(BG; \mathbb{Z}/2)$ and they are not in $N^1 H^*(BG; \mathbb{Z}/2)$. Thus we have Theorem 8.1 for $G = Q_8$.

8.2. The case $G = D_8$. Then $a^2 = c, b^2 = 1$. It is well known

$$H^*(BG)/2 \cong \mathbb{Z}/2[y_1, y_2, c_2]/(y_1 y_2)\{1, e\} \quad \text{with } |e| = 3.$$

The mod 2 cohomology is written [12]

$$\begin{aligned} H^*(BG; \mathbb{Z}/2) &\cong \mathbb{Z}/2[x_1, x_2, u]/(x_1 x_2) \quad (\text{with } |u| = 2) \\ &\cong (\oplus_{j=1}^2 \mathbb{Z}/2[y_j]\{y_j, x_j, y_j u, x_j u\} \oplus \mathbb{Z}/2\{1, u\}) \otimes \mathbb{Z}/2[c_2]. \end{aligned}$$

Here $y_j = x_j^2, u^2 = c_2$ and $Q_0(u) = (x_1 + x_2)u = e, Q_1 Q_0(u) = (y_1 + y_2)c_2$.

We note $y_1, y_2, c_2 \in CH^*(BG)/2$ and

$$H^*(BG; \mathbb{Z}/2)/(y_1, y_2, c_2) \cong (\oplus_{j=1}^2 \mathbb{Z}/2\{x_j, x_j u\}) \oplus \mathbb{Z}/2\{1, u\}.$$

Moreover, $\deg(x_j) = (1, 1), \deg(u) = (2, 2)$ in the motivic cohomology $H^{*,*'}(BG; \mathbb{Z}/2)$ and they are not in $N^1 H^*(BG; \mathbb{Z}/2)$. Here we note $\deg(x_j u) = (3, 3)$, but there is $u'_j \in H^{3,2}(BG; \mathbb{Z}/2)$ with $x_j u = \tau u'_j$ from Lemma 6.2 in [24] (i.e., $x_j u \in N^1 H^*(X; \mathbb{Z}/2)$).

Hence for the proof of Lemma 8.1 (for $G = D_8$), it is only need to show

Lemma 8.2. *We have $x_i u \in \tilde{N}^1 H^*(BG; \mathbb{Z}/2)$.*

To prove the above lemma, for a G -variety H , we consider the equivariant cohomology

$$H_G^*(H; \mathbb{Z}/p) = H^*(E(N) \times_G H; \mathbb{Z}/p)$$

where $E(N)$ is an (approximation of) contractible free G -variety. Let us write

$$X_G H = \text{approx. of } E(N) \times_G H \text{ so that } H_G^*(H; \mathbb{Z}/p) \cong H^*(X_G H; \mathbb{Z}/p).$$

For a closed embedding $i : H \subset K$ of G -varieties, we can define the transfer

$$i_* : H_G^*(H; \mathbb{Z}/p) \rightarrow H_G^*(K; \mathbb{Z}/p) \quad \text{by } i : X_G H \xrightarrow{id \times_G i} X_G K.$$

Hereafter in this section, let $G = D_8$. We recall arguments in [24]. We define the 2-dimensional representation $\tilde{c} : G \rightarrow U_2$ such that $\tilde{c}(a) = \text{diag}(i, -i)$ and $\tilde{c}(b)$ is the permutation matrix $(1, 2)$. By this representation, we identify that $W = \mathbb{C}^{2*} = \mathbb{C}^2 - \{0\}$ is an G -variety. Note G acts freely on $W \times \mathbb{C}^*$ but it does not so on $W = \mathbb{C}^{2*}$.

The fixed points set on W under b is

$$W^{(b)} = \{(x, x) | x \in \mathbb{C}^*\} = \mathbb{C}^*\{e'\}, \quad e' = \text{diag}(1, 1) \in GL_2(\mathbb{C}).$$

Similarly $W^{(bc)} = \mathbb{C}^*\{a^{-1}e'\}$. Take

$$H_0 = \mathbb{C}^*\{e', ae'\}, \quad H_1 = \mathbb{C}^*\{g^{-1}e', q^{-1}ae'\}$$

where $g \in GL_2(\mathbb{C})$ with $g^{-1}bg = ab$ (note $(ab)^2 = 1$).

Let us write $H = H_0 \sqcup H_1$. Then G acts on H_i and acts freely on $\mathbb{C}^{2*} - H$. In fact it does not contain fixed points of non-trivial stabilizer groups.

We consider the transfer for some G -set H in \mathbb{C}^{2*} , and induced equivariant cohomology

$$i_* : H_G^*(H; \mathbb{Z}/2) \rightarrow H_G^*(\mathbb{C}^{2*}; \mathbb{Z}/2).$$

Lemma 8.3. *We have*

$$H_G^*(H_0; \mathbb{Z}/2) \cong \mathbb{Z}/2[y] \otimes \Lambda(x, z) \quad \text{with } y = x^2, |x| = |z| = 1.$$

Proof. We consider the group extension $0 \rightarrow \langle a \rangle \rightarrow G \rightarrow \langle b \rangle \rightarrow 0$ and the induced spectral sequence

$$E_2^{*,*'} = H^*(B\langle b \rangle; H_{\langle a \rangle}^{*'}(H_0; \mathbb{Z}/2)) \implies H_G^*(H_0; \mathbb{Z}/2).$$

Since $\langle a \rangle \cong \mathbb{Z}/4$ acts freely on H_0 , we see

$$H_0/\langle a \rangle \cong \mathbb{C}^*\{e', ae'\}/\langle a \rangle \cong \mathbb{C}^*/\langle a \rangle \cong \mathbb{C}^*.$$

Therefore we have

$$H_{\langle a \rangle}^*(H_0; \mathbb{Z}/2) \cong H^*(\mathbb{C}^*/\langle a \rangle; \mathbb{Z}/2) \cong H^*(\mathbb{C}^*; \mathbb{Z}/2) \cong \Lambda(z) \quad |z| = 1.$$

Since $\langle b \rangle$ acts trivially on $\Lambda(z)$ we have $H_G^*(H; \mathbb{Z}/2) \cong H^*(B\langle b \rangle; \mathbb{Z}/2) \otimes \Lambda(z)$. \square

Note $H_G^*(H_0; \mathbb{Z}/2) \cong H_G^*(H_1; \mathbb{Z}/2)$ and hence we see

$$H_G^*(H; \mathbb{Z}/2) \cong \oplus_{j=1}^2 \mathbb{Z}/2[y_j]\{1_j, y_j, x_j, x_j z_j, z_j\}.$$

We consider the long exact sequence

$$(*) \quad \dots \rightarrow H_G^*(\{0\}; \mathbb{Z}/2) \xrightarrow{i_* \cong c_2} H_G^{*+4}(\mathbb{C}^2; \mathbb{Z}/2) \rightarrow H_G^{*+4}(\mathbb{C}^{2*}; \mathbb{Z}/2) \rightarrow \dots$$

and we have $H_G^*(\mathbb{C}^{2*}; \mathbb{Z}/2) \cong H^*(BG; \mathbb{Z}/2)/(c_2)$. Hence, we get

$$H_G^*(\mathbb{C}^{2*}; \mathbb{Z}/2) \cong (\oplus_{j=1}^2 \mathbb{Z}/2[y_j]\{y_j, x_j, x_j u'_j, u'_j\}) \oplus \mathbb{Z}/2\{1, u\}$$

Now we consider the transfer $H_G^*(H; \mathbb{Z}/2) \xrightarrow{i_*} H_G^{*+2}(\mathbb{C}^{2*}; \mathbb{Z}/2)$. We have explicitly (page 527 in [24])

$$i_*(1_j) = y_j, \quad i_*(x_j) = y_j x_j, \quad i_*(x_j z_j) = x_j u'_j, \quad i_*(z_j) = u'_j.$$

Therefore we have Theorem 8.1 for $G = D_8$.

To see the above fact we recall the long exact sequence for $i : H \subset \mathbb{C}^{2*}$

$$(**) \quad \dots \rightarrow H_G^{*+1}(\mathbb{C}^{2*} - H; \mathbb{Z}/2) \xrightarrow{\delta} H_G^*(H; \mathbb{Z}/2) \xrightarrow{i_*} H_G^{*+2}(\mathbb{C}^{2*}; \mathbb{Z}/2) \\ \xrightarrow{j_*} H_G^{*+2}(\mathbb{C}^{2*} - H; \mathbb{Z}/2) \rightarrow \dots$$

The transfer i_* is determined by the following lemma.

Lemma 8.4. *In the above (**), we see $\delta = 0$, and hence i_* is injective.*

Proof. Since G acts freely on $\mathbb{C}^{2*} - H$, we have

$$H_G^*(\mathbb{C}^{2*} - H; \mathbb{Z}/2) \cong H^*((\mathbb{C}^{2*} - H)/G; \mathbb{Z}/2),$$

which is zero when $* > 4 = 2\dim((\mathbb{C}^{2*} - H)/G)$. Hence δ must be zero for $* > 4$, and i_* is injective for $* > 4$. In particular, $i_*(y_j^2 z_j) = y_j^2 u'_j$. Since $H_G^*(H; \mathbb{Z}/2)$ is $\mathbb{Z}/2[y_1]$ -free (or $\mathbb{Z}/2[y_2]$ -free,) we see $i_*(z_j) = u'_j$. \square

REFERENCES

- [1] O. Benoist and J. Ottem. Two coniveau filtrations. *Duke Math J.* **170** (2021) 2719-2715.
- [2] S. Bloch and A. Ogus. Gersten's conjecture and the homology of schemes. *Ann. Scient. Éc. Norm. Sup.* **7** (1974) 181-202.
- [3] J.L. Colliot Thérène and C. Voisin. Cohomologie non ramifiée conjecture de Hodge entière. *Duke Math. J.* **161** (2012), 735-801.
- [4] T. Ekedahl. Approximating classifying spaces by smooth projective varieties *preprint* **44** (2009) arXiv:0905.1538.
- [5] M. Kameko and N. Yagita. The Brown-Peterson cohomology of the classifying cohomologies of projective unitary group $PU(n)$ and exceptional Lie groups. *Trans. Amer. Math. Soc.* **360** (2008) 2265-2284.

- [6] E. Kordonskii. Stable rationality of the group $Spin_{10}$. *Russian Math. Soc.* **360** (2000) 178-179.
- [7] A. Merkurjev. Rationally problem for classifying spaces of Spinor groups. *Proc. Steklov Inst. Math.* **307** (2019) 115-124.
- [8] J. Milnor. The Steenrod algebra and its dual. *Ann. Math.* **67** (1958) 150-171.
- [9] K. Paranjape. Some spectral sequences for filtered complexes and applications. *J. Algebra* **186** (1996) 793-806.
- [10] A. Pirutka and N. Yagita. Note on the counterexamples for the integral integral Tate conjecture over finite fields. *Doc. Math.* (2015) Extra vol. : Alexander S. Merkurjev's sixties birthday. 501-511.
- [11] D. Quillen. Elementary proofs of some results of the cobordism theory using Steenrod operations. *Advances in Math.* **7** (1971), 29-56.
- [12] D. Quillen. The mod 2 cohomology rings of extra-special 2-groups and the spinor groups. *Math. Ann.* **194** (1971), 197-212.
- [13] D. Ravenel, S. Wilson and N. Yagita. Brown-Peterson cohomology from Morava K -theory. *K-theory*. **15** (1998), 147-199.
- [14] Z. Reichstein and F. Scavia. The Noether problem for spinor groups of small rank. *J. Algebra* **48** (2020), 134-152.
- [15] M. Tamanoi. The image of the BP-Thom map for Eilenberg-MacLane spaces. *Trans. AMS.* **249** (1997), 1209-1237.
- [16] M. Tezuka and N. Yagita. The image of the map from group cohomology to Galois cohomology. *Trans. Amer. Math. Soc.* **363** (2011), 4475-4503.
- [17] B. Totaro. The Chow ring of classifying spaces. *Proc. of Symposia in Pure Math. "Algebraic K-theory" (1997: University of Washington, Seattle)* **67** (1999), 248-281.
- [18] V. Voevodsky. The Milnor conjecture. www.math.uiuc.edu/K-theory/0170 (1996).
- [19] V. Voevodsky. Reduced power operations in motivic cohomology. *Publ. Math. IHES* **98** (2003), 1-57.
- [20] V. Voevodsky. Motivic cohomology with $\mathbb{Z}/2$ coefficient. *Publ. Math. IHES* **98** (2003), 59-104.
- [21] V. Voevodsky. On motivic cohomology with \mathbb{Z}/l -coefficients. *Ann. Math.* **174** (2011) 401-438.
- [22] N. Yagita. Coniveau filtration of cohomology of groups. *Proc. London Math. Soc.* **101** (2010) 179-206.
- [23] N. Yagita. Algebraic BP -theory and norm varieties. *Hokkaido Math. J.* **41** (2012), 275-316.
- [24] N. Yagita. Chow rings of nonabelian p -groups of order p^3 . *J. Math. Soc. Japan* **64** (2012), 507-531.
- [25] N. Yagita. Chern classes and the Rost cohomological invariant. *Kodai Math. J.* **36** (2013), 174-178.

[N. YAGITA] DEPARTMENT OF MATHEMATICS, FACULTY OF EDUCATION, IBARAKI UNIVERSITY, MITO, IBARAKI, JAPAN

Email address: nobuaki.yagita.math@vc.ibaraki.ac.jp