A test for partial correlation between repeatedly observed nonstationary nonlinear timeseries

Kenneth D. Harris^{1*}

¹UCL Institute of Neurology, Queen Square, London WC1N 3BG, UK. * Correspondence: <u>kenneth.harris@ucl.ac.uk</u>.

We describe a family of statistical tests to measure partial correlation in vectorial timeseries. The test measures whether an observed timeseries Y can be predicted from a second series X, even after accounting for a third series Z which may correlate with X. It does not make any assumptions on the nature of these timeseries, such as stationarity or linearity, but it does require that multiple statistically independent recordings of the 3 series are available. Intuitively, the test works by asking if the series Y recorded on one experiment can be better predicted from X recorded on the same experiment than on a different experiment, after accounting for the prediction from Z recorded on both experiments.

T N many fields of science, observations consist of **⊥** timeseries. These timeseries often autocorrelated, meaning that observations from a single timeseries at different times are not statistically independent. Such autocorrelations mean that standard statistical tests, which make an assumption of independent, identically distributed observations, cannot be applied (Box, 2008; Granger and Newbold, 1974; Haugh, 1976; Phillips, 1986; Yule, 1926). Most proposed solutions to the problem focus on removing autocorrelations for example by fitting linear autoregressive models. However, this approach requires that the series be accurately fit by stationary linear models, which is often untrue (or impossible to show) in practice.

This problem is greatly ameliorated if one has access to multiple repetitions of the experiment. For example, consider an experiment yielding two timeseries X and Y which we consider as vectors of length T. The timeseries are statistically independent if $\mathbb{P}(X,Y) = \mathbb{P}(X)\mathbb{P}(Y)$: in other words, if there is no correlation between the entire history of **X** with the entire history of **Y**. Given only a single observation of the vectors \mathbf{X} and \mathbf{Y} , we cannot test for violation of independence without making further assumptions such as stationarity: methods such as Pearson correlation, which assume independence of timepoints, givens erroneously significant "nonsense correlations" (Yule, 1926). However, if the experiment is repeated N times, and we assume that the histories X_i and Y_i (i =1...N) observed on each repeat are independent vectors, we can employ a "session permutation" method to test for independence (Harris, 2020), without any further assumptions. This method essentially asks whether the relationship between X_i and Y_i is larger than that between X_i and Y_j for $i \neq j$, using a permutation test that randomizes across experiments. The test can be based on any measure of similarity between timeseries, including but not limited to Pearson correlation.

Here, we describe a family of tests for partial correlation between repeatedly observed timeseries. We assume that each of N experiments yielded 3 vectorial timeseries X, Y and Z. We ask whether a correlation between X and Y exists beyond a common effect of Z. No assumptions of stationarity or linearity are made, in fact X, Y and Z can be arbitrary vectors rather than timeseries.

The test

We assume N statistically independent experiments, each of which gives an observation of three vectorial timeseries: \mathbf{X}_i , \mathbf{Y}_i and \mathbf{Z}_i , for $i = 1 \dots N$, considered as matrices of sizes $T \times p_i$, $T \times q_i$ and $T \times r_i$, respectively. Write \mathbf{X} for the collection of all \mathbf{X}_i and \mathbf{Z}_i . We would like to test the null hypothesis that

$$Y_i = Z_i W_i + E_i$$

where the E_i are timeseries independent of each other and of X, and W_i is an unknown deterministic $p_i \times r_i$ matrix. This null hypothesis models the idea that Y may be linearly predicted from Z but there is

no further effect of **X** on **Y** once the effect of **Z** is considered, even though **X** and **Z** may themselves be correlated.

Write P_i for the $T \times T$ projection matrix orthogonal to all columns of Z_i . Under the null, $P_iY_i = P_iE_i$. Let $\rho(X; Y)$ be a measure of the predictability of a vector timeseries Y from a second timeseries Y, for example the Pearson coefficient of multiple linear regression. For a statistical test, you might first think to ask if $\rho(X_i; P_iY_i)$ is bigger than $\rho(X_j; P_iY_i)$ for $i \neq j$, i.e. if the relationship of E and E is strongest when they come from the same experiment. But $\rho(X_i; P_iY_i)$ is not identically distributed to $\rho(X_j; P_iY_i)$ for $j \neq i$ since Y_i and P_i are correlated but Y_i and P_i are not.

Instead, let $P_{i,j}$ be the $T \times T$ projection matrix orthogonal to the columns of both Z_i and Z_j ; by definition $P_{j,i} = P_{i,j}$. We will compare $\rho(X_i; P_{i,j}Y_i)$ to $\rho(X_j; P_{i,j}Y_i)$, Specifically, define

$$G_i = \frac{1}{N} \sum_{i=1}^{N} \rho(\mathbf{X}_i; \mathbf{P}_{i,j} \mathbf{Y}_i) - \rho(\mathbf{X}_j; \mathbf{P}_{i,j} \mathbf{Y}_i)$$

Under the null, $P_{i,j}Y_i = P_{i,j}E_i$, so $G_i = \frac{1}{N}\sum_{j=1}^{N}\rho(X_i; P_{i,j}E_i) - \rho(X_j; P_{i,j}E_i)$. Because the E_i are independent, the G_i are conditionally independent given \mathcal{X} , although they need not be identically distributed.

Under the null, the expectation of $\sum_{i=1}^{N} G_i$ conditional on $\boldsymbol{\mathcal{X}}$ is zero. Indeed,

$$\sum_{i=1}^{N} G_i = \frac{1}{N} \sum_{i,j=1}^{N} \rho(\mathbf{X}_i; \mathbf{P}_{i,j} \mathbf{E}_i) - \rho(\mathbf{X}_j; \mathbf{P}_{i,j} \mathbf{E}_i)$$
$$= \frac{1}{N} \sum_{i,j=1}^{N} \rho(\mathbf{X}_i; \mathbf{P}_{i,j} \mathbf{E}_i) - \rho(\mathbf{X}_i; \mathbf{P}_{i,j} \mathbf{E}_j)$$

For the last equality we have used the fact that $P_{j,i} = P_{i,j}$. Now because E_i and E_j are identically distributed and independent of everything else,

$$\mathbb{E}\left[\sum_{i=1}^{N} G_{i} \mid \mathbf{X}\right]$$

$$= \frac{1}{N} \sum_{i,j=1}^{N} \mathbb{E}\left[\rho\left(\mathbf{X}_{i}; \; \mathbf{P}_{i,j} \mathbf{E}_{i}\right) - \rho\left(\mathbf{X}_{i}; \; \mathbf{P}_{i,j} \mathbf{E}_{j}\right) \middle| \mathbf{X}\right] = 0$$

Therefore, under the null, the statistics G_i are statistically independent conditional on X, with




Figure 1: detecting a genuine correlation. In this simulation X_i and Y_i are both equal to a sum of two step functions, one of which is constant across experiments. Z_i equals the constant step function, and after removing it from Y_i , the residual $P_{i,j}Y_i$ is reliably more correlated to X_i than to X_j , and statistical significance is detected.

expectations that sum to zero. They are marginally exchangeable by independence of experiments, and are thus marginally identically distributed. Nevertheless, they need not be either conditionally or marginally Gaussian. To test the null hypothesis in such circumstances, a t-test that the mean of G_i is zero is valid provided the distributions of the G_i are close to symmetric (Cressie, 1980; Efron, 1969); this can be checked by usual methods such as histograms or QQ plots.

Example

We illustrate the test with a simple example (Figs. 1, 2) in which X and Y are identical, and are given by a sum of two step functions. The first step function S_0 is the same for each observation i, being 0 for the first half of the timeseries and 1 for the second half. The second step function S_i is a short pulse which occurs at a different time for different experiments i, but always the same time for X and Y. We




Figure 2: correlations due to common effects of Z are not detected. X_i and Y_i are as in Fig. 1, but now Z_i is the step function that varies between experiments. Because both Z_1 and Z_2 are projected out, the similarity of $P_{1,2}Y_1$ to X_1 is the same as to X_2 , so no correlation is detected.

simulate $X_i = Y_i = S_0 + S_i$ for each i. We consider two cases: that $Z_i = S_0$ for all i, or that $Z_i = S_i$.

If $\mathbf{Z}_i = \mathbf{S}_0$, there is a partial correlation of \mathbf{X}_i and \mathbf{Y}_i given \mathbf{Z}_i , because the time of the pulse varies between experiments, but \mathbf{Z}_i is always the same. Thus, $\rho(\mathbf{X}_i; \mathbf{P}_{i,j}\mathbf{Y}_i)$ is reliably larger than $\rho(\mathbf{X}_j; \mathbf{P}_{i,j}\mathbf{Y}_i)$, and the test finds statistical significance (Figure 1).

However, if $\mathbf{Z}_i = \mathbf{S}_i$, there is no a partial correlation of \mathbf{X}_i and \mathbf{Y}_i given \mathbf{Z}_i , since the correlation between \mathbf{X}_i and \mathbf{Y}_i can be explained by a common dependence on \mathbf{Z}_i . Thus, $\rho(\mathbf{X}_i; \mathbf{P}_{i,j}\mathbf{Y}_i)$ and $\rho(\mathbf{X}_j; \mathbf{P}_{i,j}\mathbf{Y}_i)$ are comparable, and the test finds no statistical significance (Figure 2).

Finally, to show why it is essential to use $P_{i,j}Y_i$ rather than simply P_iY_i , we repeat the simulation with $Z_i = S_i$, but comparing $\rho(X_i; P_iY_i)$ with $\rho(X_j; P_iY_i)$. This invalid test erroneously finds a




Figure 3: both Z_i and Z_j must be projected out to avoid erroneous correlations. All data are as in Fig. 2 but only Z_i is projected from Y_i . Because Z_j is not projected from Y_i , the residual P_iY_i is more similar to X_j than X_i , leading to an erroneously significant negative partial correlation.

significant partial correlation where none exists (Figure 3).

Discussion

We have described a family of statistical tests for partial correlation between vectorial timeseries. The timeseries are not assumed stationary or linear, and in fact the test applies to any data consisting of vectors depending on an index t. The test requires multiple observations of these timeseries, and its statistical power will increase with the number of observations available. Intuitively, the test measures how much better one can predict \mathbf{Y}_i from \mathbf{X}_i measured in the same experiment, compared to predicting \mathbf{Y}_i from \mathbf{X}_j measured in a different experiment. To remove a possible common effect of the confounding variable \mathbf{Z}_i both \mathbf{Z}_i and \mathbf{Z}_j are projected out from the T-dimensional vector \mathbf{Y}_i .

The test depends on a user-supplied measure $\rho(X,Y)$ of the degree of association between two vectorial timeseries. It makes no assumptions about

this measure, so anything can be used. When dealing with 1-dimensional timeseries (as in the examples presented here), Pearson correlation is a natural choice. In higher dimensions one could for example use the fraction of variance explained by multiple linear regression or ridge regression. If **Y** is high dimensional, it may be advisable to use regularize further with methods such as reduced rank regression or canonical correlation analysis.

The test may find use in multiple scientific fields where time-dependent experiments are performed repeatedly. An example in neuroscience would be if one makes multivariate recordings from the brains of subjects performing a behavioral task, and would like to test if some behavioral variables correlate with brain activity even after the effect of other behavioral variables are controlled for. Unlike the "pseudosession method" (Harris, 2020), the current test requires multiple repeats of an experiment. However the matrices W_i can differ between experiments, and so it is possible for these experiments to be made from different subjects, or even include different numbers of observations. Furthermore, unlike the "pseudosession method" this method can be used when the predictor variables **X** and **Z** are not randomly generated, but produced by the subject themselves according to an unknown distribution.

REFERENCES

Box, G.E.P. (2008). Time series analysis forecasting and control / George E.P. Box, Gwilym M. Jenkins, Gregory C. Reinsel. (Hoboken, N.J.: John Wiley).

Cressie, N. (1980). Relaxing assumptions in the one sample ttest. Aust. J. Stat. 22, 143–153.

Efron, B. (1969). Student's t-Test Under Symmetry Conditions. J. Am. Stat. Assoc. *64*, 1278–1302.

Granger, C.W.J., and Newbold, P. (1974). Spurious regressions in econometrics. J. Econom. 2, 111–120.

Harris, K.D. (2020). Nonsense correlations in neuroscience. BioRxiv 2020.11.29.402719.

Haugh, L.D. (1976). Checking the independence of two covariance-stationary time series: a univariate residual cross-correlation approach. J. Am. Stat. Assoc. 71, 378–385.

Phillips, P.C.B. (1986). Understanding spurious regressions in econometrics. J. Econom. 33, 311–340.

Yule, G.U. (1926). Why do we Sometimes get Nonsense-Correlations between Time-Series?--A Study in Sampling and the Nature of Time-Series. J. R. Stat. Soc. 89, 1–63.