GLOBAL SCHAUDER THEORY FOR MINIMIZERS OF THE $H^{s}(\Omega)$ ENERGY

MOUHAMED MOUSTAPHA FALL AND XAVIER ROS-OTON

ABSTRACT. We study the regularity of minimizers of the functional $\mathcal{E}(u) := [u]_{H^s(\Omega)}^2 + \int_{\Omega} fu$. This corresponds to understanding solutions for the regional fractional Laplacian in $\Omega \subset \mathbb{R}^N$. More precisely, we are interested on the global (up to the boundary) regularity of solutions, both in the case of free minimizers in $H^s(\Omega)$ (i.e., Neumann problem), or in the case of Dirichlet condition $u \in H_0^s(\Omega)$ when $s > \frac{1}{2}$.

Our main result establishes the sharp regularity of solutions in both cases: $u \in C^{2s+\alpha}(\overline{\Omega})$ in the Neumann case, and $u/\delta^{2s-1} \in C^{1+\alpha}(\overline{\Omega})$ in the Dirichlet case. Here, δ is the distance to $\partial\Omega$, and $\alpha < \alpha_s$, with $\alpha_s \in (0, 1-s)$ and $2s + \alpha_s > 1$. We also show the optimality of our result: these estimates fail for $\alpha > \alpha_s$, even when f and $\partial\Omega$ are C^{∞} .

1. INTRODUCTION AND MAIN RESULTS

Given $s \in (0,1)$ and a bounded domain $\Omega \subset \mathbb{R}^N$, we study the regularity of minimizers of energy functionals of the type

$$\mathcal{E}(u) := [u]_{H^s(\Omega)}^2 + \int_{\Omega} f u \, dx, \qquad (1.1)$$

where $f \in L^2(\Omega)$,

$$[u]_{H^{s}(\Omega)}^{2} = \frac{c_{N,s}}{4} \int_{\Omega} \int_{\Omega} \frac{|u(x) - u(y)|^{2}}{|x - y|^{N + 2s}} \, dx \, dy,$$

and $c_{N,s} = \frac{s4^s \Gamma\left(\frac{N}{2}+s\right)}{\pi^{\frac{N}{2}} \Gamma(1-s)}.$

Notice that, as in the classical case s = 1, one can study two types of minimizers:

• Free minimizers $u \in H^s(\Omega)$ of (1.1) —i.e., minimizing among all functions in $H^s(\Omega)$ —, which corresponds to a **Neumann problem**.

• Minimizers with prescribed zero boundary data —i.e., minimizing among functions $u \in H_0^s(\Omega)$ —, which corresponds to a **Dirichlet problem**. Notice however that one needs $s > \frac{1}{2}$ in order to have a trace operator¹, thus the Dirichlet problem makes sense only for $s > \frac{1}{2}$.

The aim of this paper is to understand the global (up to the boundary) regularity of solutions in both cases:

If f and Ω are regular enough, what is the regularity of minimizers u?

This is the question that motivates our work.

²⁰¹⁰ Mathematics Subject Classification. 35B65, 35R11.

Key words and phrases. Regional fractional Laplacian, regularity, Schauder estimates, Censored processes. ¹When $s \leq \frac{1}{2}$ we have $H_0^s(\Omega) = H^s(\Omega)$ for any Lipschitz domain $\Omega \subset \mathbb{R}^N$, see e.g. [14, Theorem 1.4.2.4].

Minimizers of (1.1) solve the equation $(-\Delta)^s_{\Omega} u = f$ in Ω in the weak sense, where

$$(-\Delta)^{s}_{\Omega}u(x) = c_{N,s} \text{p.v.} \int_{\Omega} \frac{u(x) - u(y)}{|x - y|^{N+2s}} dy$$
 (1.2)

is the so-called regional fractional Laplacian. Recall that this operator, subjetcs to Dirichlet or Neumann boundary conditions are generated by Lèvy-type processes which are censored to leave Ω , see e.g. [3].

Problems involving the regional fractional Laplacian have been studied both from the point of view of Probability [3, 7, 8, 15, 16] and of Analysis [2, 6, 12, 17]. The best known regularity results in this direction are those in [12], where the first author established that for Ω sufficiently regular and $f \in L^{\infty}(\Omega)$, solutions u are $C^{2s-\varepsilon}(\overline{\Omega})$ in the Neumann case and that $u/\delta^{2s-1} \in C^{1-\varepsilon}(\overline{\Omega})$ for the Dirichlet problem, for every $\varepsilon > 0$.

Our main results here establish for the first time Schauder-type estimates for such problem, which we show to be optimal. They can be summarized as follows:

Theorem 1.1 (Neumann problem). Let $s \in (0,1)$, and let $\alpha_s \in (0,1-s)$ be given by Lemma 2.6, which satisfies $2s + \alpha_s > 1$. Let $\Omega \subset \mathbb{R}^N$ be any bounded $C^{\max(1,2s+\alpha)}$ domain, and $f \in C^{\alpha}(\overline{\Omega})$ with $\int_{\Omega} f \, dx = 0$. Let $\alpha < \alpha_s$ with $\alpha + 2s \neq 1$.

Let u be the minimizer of (1.1) in $H^{s}(\Omega)$. Then, $u \in C^{2s+\alpha}(\overline{\Omega})$ and

$$\|u\|_{C^{2s+\alpha}(\overline{\Omega})} \le C \|f\|_{C^{\alpha}(\overline{\Omega})},$$

with C depending only on N, s, α , and Ω . In addition, if α is such that $2s + \alpha > 1$ then

 $\partial_{\nu} u = 0$ on $\partial \Omega$.

Moreover, for $\alpha > \alpha_s$, solutions u are in general not $C^{2s+\alpha}(\overline{\Omega})$.

The critical exponent $\alpha_s > 0$ is given by an explicit equation in terms of Gamma functions; see Remark 1.4 below. For example, in the simplest case $s = \frac{1}{2}$, it is the first positive solution of $1 + \alpha = \frac{1}{\pi} \tan(\pi \alpha)$.

Next, we notice that Theorem 1.1 implies that the minimizer u satisfies $(-\Delta)^s_{\Omega} u \in C^{\alpha}(\overline{\Omega})$ and thus it is in fact a classical solution: $(-\Delta)^s_{\Omega} u(x) = f(x)$ for all $x \in \overline{\Omega}$. This will follow from the results in Section 4 below.

Remark 1.2. It is natural to wonder whether, for $s \in (0, \frac{1}{2}]$, solutions to the Neumann problem are C^1 up to the boundary or not and have zero normal derivative when f and Ω are regular enough.

Our result answers this question for the first time: solutions are always $C^1(\overline{\Omega})$, for every $s \in (0,1)$. Furthermore, since $\alpha_s < 1-s$, then solutions are in general not $C^2(\overline{\Omega})$.

We think it is quite surprising that the answer is the same for all $s \in (0, 1)$.

In the Dirichlet case our result reads as follows:

Theorem 1.3 (Dirichlet problem). Given $s \in (\frac{1}{2}, 1)$, let $\alpha_s \in (0, 1-s)$ be given by Lemma 2.6. Let $\Omega \subset \mathbb{R}^N$ be a $C^{2,\beta}$ domain and $f \in C^{\alpha}(\overline{\Omega})$, with $\alpha < \min\{\alpha_s, 2s - 1, \beta\}$ and $\alpha + 2s \neq 1$.

Let u be the minimizer of (1.1) in $H_0^s(\Omega)$. Then, $u/\delta^{2s-1} \in C^{1+\alpha}(\overline{\Omega})$ and

$$\|u/\delta^{2s-1}\|_{C^{1+\alpha}(\overline{\Omega})} \le C\|f\|_{C^{\alpha}(\overline{\Omega})},$$

with C depending only on N, s, α , and Ω . Here, $\delta \in C^{2,\beta}(\overline{\Omega})$ is a positive function in Ω that coincides with dist $(\cdot, \partial \Omega)$ near $\partial \Omega$.

In addition, if we denote $\psi := u/\delta^{2s-1}$, we have

$$\partial_{\nu}\psi = -(N-1)H_{\partial\Omega}\psi \quad on \quad \partial\Omega, \tag{1.3}$$

where $H_{\partial\Omega}$ is the mean curvature of $\partial\Omega$ and ν is the exterior normal of $\partial\Omega$.

Moreover, for $\alpha > \alpha_s$ the quotient u/δ^{2s-1} does not belong in general to $C^{1+\alpha}(\overline{\Omega})$.

We point out that (1.3) does not hold in the local case s = 1. Indeed, in that case we have $\psi = -\partial_{\nu}u$ and hence $\partial_{\nu}\psi = -(N-1)H_{\partial\Omega}\psi + f$ on $\partial\Omega$, where we used $\Delta u = \partial_{\nu}^{2}u + (N-1)H_{\partial\Omega}\partial_{\nu}u$ on $\partial\Omega$.

A difficulty that arises in the proof of Theorem 1.3 in dimension $N \geq 2$ is that $(-\Delta)^s_{\Omega} \delta^{2s-1}$ does not in general belong to $L^{\infty}(\Omega)$ even if Ω is C^{∞} . In fact near $\partial\Omega$, we have that $(-\Delta)^s_{\Omega} \delta^{2s-1}(x) \simeq g(x) \log(\delta)$, with $g|_{\partial\Omega} = (N-1)H_{\partial\Omega}$. This leads to the assumption Ω to be of class $C^{2,\beta}$ with $\beta > \alpha$. To overcome this difficulty we consider the correction $\delta^{2s-1} + b\delta^{2s}$, for some appropriately chosen function b defined on $\partial\Omega$. This yields also the identity (1.3).

Remark 1.4 (The 1D case). Let us briefly explain what happens in dimension 1.

Let us assume for example that $\Omega = \mathbb{R}_+$. Then, we will show that any solution to $(-\Delta)^s_{\Omega} u = 0$ in $(0,1) \subset \mathbb{R}$ has an expansion of the form

$$u(x) = c_0 + a_0 x^{2s-1} + a_1 x^{\beta_1} + \dots + a_k x^{\beta_k} + \dots \quad for \quad x \approx 0, \tag{1.4}$$

where all β_k 's can be characterized as the positive solutions of

$$\frac{\Gamma(\beta+1)}{\Gamma(\beta-2s+1)\Gamma(2s)} = \frac{1}{\pi} \frac{\sin\left(\pi(\beta-2s)\right)\sin(\pi s)}{\sin\left(\pi(\beta-s)\right)}.$$

By an appropriate analysis of such equation², we will show that

$$\beta_{k+1} > 1 + \beta_k$$
 for all $k \ge 1$.

and $\beta_1 = 2s + \alpha_s$. In addition, we will see that $\alpha_s \in (0, 1-s)$ and $2s + \alpha_s > 1$.

In case of zero Neumann boundary conditions on $\partial\Omega$, we must have $a_0 = 0$ in (1.4), while in case of zero Dirichlet conditions we must clearly have $c_0 = 0$. This already shows that the $C^{2s+\alpha}$ regularity result from Theorem 1.1 cannot hold for $\alpha > \alpha_s$, as stated above.

In higher dimensions several difficulties arise, and we will have an expansion of this type with some new terms, both coming from the tangential extra variables and from the curvature of the domain.

We expect the assumptions on Ω in both Theorems 1.1 and 1.3 to be optimal.

1.1. **Preliminaries and definitions.** Let us next give some important definitions that will be used throughout the paper.

First, recall that given any open set $\Omega \subset \mathbb{R}^N$, the regional fractional Laplacian in Ω is defined by (1.2), and its associated bilinear form is given by

$$D_{\Omega}(u,v) = \frac{c_{N,s}}{2} \int_{\Omega} \int_{\Omega} \frac{(u(x) - u(y))(v(x) - v(y))}{|x - y|^{N + 2s}} \, dx \, dy$$

The natural notion of weak solution is the following.

Definition 1.5. We say that u is a (weak) solution of

$$(-\Delta)^s_{\Omega} u = f \quad in \quad \Omega \cap B_1 \tag{1.5}$$

²Notice that when $s = \frac{1}{2}$ this equation becomes $\pi\beta = \tan(\pi\beta)$. In this case, it is easy to visualize and prove all properties of the exponents β_k .

with zero Neumann boundary condition on $\partial \Omega \cap B_1$ if $u \in H^s(\Omega)$ and

$$D_{\Omega}(u,\varphi) = \int_{\Omega} f\varphi \, dx \qquad \forall \varphi \in C_c^{\infty}(B_1).$$

On the other hand, when $s > \frac{1}{2}$, we say that u is a (weak) solution of (1.5) with zero Dirichlet boundary condition on $\partial \Omega \cap B_1$ if $u \in H^s_0(\Omega)$ and

$$D_{\Omega}(u,\varphi) = \int_{\Omega} f\varphi \, dx \qquad \forall \varphi \in C_c^{\infty}(B_1 \cap \Omega).$$

By approximation, we can of course take any $\varphi \in H_0^s(\Omega \cap B_1)$.

We notice that being a weak solution is equivalent to being a minimizer of the functional (1.1), for both the Neumann and the Dirichlet case.

1.2. Acknowledgements. The first author's work is supported by the Alexander von Humboldt foundation. The second author was supported by the European Research Council (ERC) under Grant Agreement No 801867.

1.3. Organization of the paper. The paper is organized as follows. In Section 2 we completely characterize all 1D homogeneous solutions, and prove (1.4). In Section 3 we use this to prove Liouville theorems in a half-space. Then, in Section 4 we provide some new estimates for the regional fractional Laplacian in domains. Finally, in Sections 5 and 6 we prove Theorems 1.1 and 1.3, respectively.

2. 1D homogeneities

The aim of this section is to classify all possible homogeneities in dimension 1 and, as a consequence, to prove (1.4).

Roughly speaking, we want to classify all solutions $u \in H^s_{loc}([0,\infty))$ of

$$(-\Delta)^s_{\mathbb{R}_+} u = 0 \quad \text{in} \quad \mathbb{R}_+.$$

To do so, we will define

$$v(x) := \begin{cases} u(x) - u(0) & \text{for } x \ge 0\\ 0 & \text{for } x \le 0, \end{cases}$$

and notice that we have³ $(-\Delta)^{s}v(x) = (-\Delta)^{s}_{\mathbb{R}_{+}}u(x) + u(x)c_{1,s}\int_{\mathbb{R}_{-}}|x-y|^{-1-2s} dy$. Then by a direct computation, we find

$$\begin{cases} (-\Delta)^s v - \frac{a_s}{x^{2s}} v = 0 & \text{in } \mathbb{R}_+ \\ v = 0 & \text{in } \mathbb{R}_-, \end{cases}$$
(2.2)

where $a_s := \frac{c_{1,s}}{2s}$. This allows us to use the Caffarelli-Silvestre extension (see e.g. [4]). Now note that, since our aim is to obtain $C^{2s+\alpha}$ regularity, we will need to consider

Now note that, since our aim is to obtain $C^{2s+\alpha}$ regularity, we will need to consider solutions to (2.2) that grow like $|x|^{2s+\alpha}$ at infinity. Thus, the equations in (2.2) needs to be understood in a generalized sense.

In the following of this paper we let $\chi_R \in C_c^{\infty}(B_{2R})$ be such that $\chi_R \equiv 1$ in B_R .

³Here, $(-\Delta)^s$ denotes the fractional Laplacian in \mathbb{R} , i.e., (1.2) with $\Omega = \mathbb{R}$.

Definition 2.1. As in⁴ [1, Section 3], given a function with polynomial growth

$$\int_{\mathbb{R}^N} \frac{|u(x)|}{1+|x|^{N+2s+k}} \, dx < \infty,$$

we say that

$$(-\Delta)^s u \stackrel{k}{=} f \quad in \quad \Omega \subset \mathbb{R}^N$$

if there exists a family of polynomials p_R of degree at most k-1 and a family of functions f_R such that $(-\Delta)^s(u\chi_R) = f_R + p_R$ in $\Omega \cap B_{R/2}$, with $f_R \to f$ uniformly in Ω as $R \to \infty$.

Thanks to [1, Lemma 3.3], this is equivalent to saying that there exists an extension $\tilde{u}(x,y)$ with polynomial growth in \mathbb{R}^{N+1}_+ , satisfying $\tilde{u}(x,0) = u(x)$ and $\operatorname{div}(t^{1-2s}\nabla \tilde{u}) = 0$ in $\{y > 0\}$, such that

$$-\lim_{t\to 0} t^{1-2s} \partial_t \tilde{u} = \overline{\kappa}_s f \quad on \quad \Omega \cap \{y=0\}.$$

It is easy to check that, if $\int_{\mathbb{R}^N} |D^k u(x)|/(1+|x|^{N+2s}) dx < \infty$, and $(-\Delta)^s D^k u = D^k f$ in Ω for some function f, then $(-\Delta)^s u \stackrel{k}{=} f$ in Ω .

In the extended variables $(x,t) \in \mathbb{R} \times \mathbb{R}_+$ —and denoting v(x,t) the extension of v(x)—, (2.2) reads as

$$\begin{cases} \operatorname{div}(t^{1-2s}\nabla v) = 0 & \text{in } \{t > 0\} \\ -\lim_{t \to 0} t^{1-2s} \partial_t v = \overline{\kappa}_s \frac{a_s}{x^{2s}} v & \text{on } \{t = 0, \ x > 0\} \\ v = 0 & \text{on } \{t = 0, \ x \le 0\}, \end{cases}$$
(2.3)

where $\overline{\kappa}_s = \frac{\Gamma(1-s)}{2^{2s-1}\Gamma(s)}$.

If we look for homogeneous solutions

$$v(r\cos\theta, r\sin\theta) = r^{\beta}\psi(\theta), \qquad (2.4)$$

with r > 0 and $\theta \in [0, \pi]$, we are led to the following eigenvalue problem for the function ψ in the space $H^1((0, \pi); \sin(\theta)^{1-2s} d\theta)$:

$$\begin{cases} -\left(\sin(\theta)^{1-2s}\psi'\right)' = \lambda\sin(\theta)^{1-2s}\psi & \text{for } \theta \in (0,\pi) \\ -\lim_{\theta \to 0}\sin(\theta)^{1-2s}\psi'(\theta) = \overline{\kappa}_s a_s \psi(0) & \\ \psi(\pi) = 0, \end{cases}$$
(2.5)

where

$$\lambda = \beta(\beta - 2s + 1).$$

Problem (2.5) possesses a sequence of increasing eigenvalues

$$\lambda_0(s) < \lambda_1(s) \leq \ldots,$$

with corresponding eigenfunctions $\psi_k \in L^2((0,\pi); \sin(\theta)^{1-2s})$, normalized as

$$\int_0^\pi \sin(\theta)^{1-2s} \psi_k^2(\theta) \, d\theta = 1.$$

Moreover, see e.g. [12], ⁵ we have $\lambda_0(s) = 0$. Using this, we can prove the following.

⁴This kind of definition was originally introduced in [10]. However, the definition in [10] requires only pointwise convergence of f_R , while here (as in [1]) we take uniform convergence.

⁵This follows from the Poincaré type inequality: $\lambda_0(s) = \inf_{\substack{\psi \in C^1([0,\pi])\\\psi(\pi)=0}} \frac{\int_0^{\pi} \sin(\theta)^{1-2s} (\psi'(\theta))^2 d\theta - \overline{\kappa}_s a_s \psi(0)^2}{\int_0^{\pi} \sin(\theta)^{1-2s} \psi_k^2(\theta) d\theta} = 0.$

Lemma 2.2. Let λ_k and ψ_k as above, and let $\beta_k = \frac{2s-1}{2} + \sqrt{\lambda_k(s) + (\frac{2s-1}{2})^2}$. Then, the functions

$$v_k(r,\theta) = r^{\beta_k} \psi_k(\theta)$$

are homogeneous solutions of (2.3).

Moreover, if $w \in H^1(B^+_{\rho}; t^{1-2s}dxdt) \cap C(\overline{B^+_{\rho}})$ is any weak solution of

$$\begin{cases} \operatorname{div}(t^{1-2s}\nabla w) = 0 & \text{in } B_{\rho}^{+} \\ -\lim_{t \to 0} t^{1-2s} \partial_{t} w = \overline{\kappa}_{s} a_{s} x^{-2s} w & \text{on } \{t = 0, \ 0 < x < \rho\} \\ w = 0 & \text{on } \{t = 0, \ -\rho < x \le 0\}, \end{cases}$$
(2.6)

then

$$w(r\cos\theta, r\sin\theta) = \sum_{k=0}^{\infty} \omega_k r^{\beta_k} \psi_k(\theta), \qquad (2.7)$$

with $\omega_k \in \mathbb{R}$.

Proof. The first part of the lemma holds by construction of λ_k and ψ_k .

For the second part, notice that since ψ_k are an orthonormal basis of $L^2((0,\pi); \sin(\theta)^{1-2s} d\theta)$, then we can write

$$w(\rho\cos\theta,\rho\sin\theta) = \sum_{k\geq 0} \omega_k \rho^{\beta_k} \psi_k(\theta)$$

for some constants $a_k \in \mathbb{R}$ (recall that here ρ is a fixed positive number).

But then, the function

$$\tilde{w}(r\cos\theta, r\sin\theta) := \sum_{k\geq 0} \omega_k r^{\beta_k} \psi_k(\theta)$$

is a solution of (2.6) that coincides with w on ∂B_{ρ} . Moreover, since $\lambda_0(s) = 0$, then $\beta_0 = \max(2s - 1, 0)$ so that by continuity, $\omega_0 = 0$ for $s \leq 1/2$. This implies that $\tilde{w} \in H^1(B_{\rho}; t^{1-2s} dx dt)$. Now by uniqueness of solutions, the two functions \tilde{w} and w must coincide.

As a consequence, we find:

Corollary 2.3. Let $u \in H^{s}([0,1]) \cap C([0,1])$ be a function satisfying

$$(-\Delta)^s_{\mathbb{R}_+} u = 0 \quad in \quad (0,1) \subset \mathbb{R},$$

in the sense that $(-\Delta)^s v \stackrel{k}{=} a_s x^{-2s} v$ in (0,1) for some $k \in \mathbb{N}$, with $v := (u-u(0))1_{\mathbb{R}_+}$. Then, near x = 0 we have

$$u(x) = \omega_0 x^{\beta_0} + \omega_1 x^{\beta_1} + \omega_2 x^{\beta_2} + \dots,$$

where $\omega_i \in \mathbb{R}$, and the exponents β_i are given by Lemma 2.2.

Proof. It follows from Lemma 2.2.

Our next goal is to find an explicit equation for these exponents β_k . We start with the following

Lemma 2.4. Let $s \in (0,1)$ and $\beta > -1$. Then, there exists a unique β -homogeneous function v_β satisfying

$$\begin{cases} \operatorname{div}(t^{1-2s}\nabla v_{\beta}) = 0 & \text{in } \mathbb{R}^{2}_{+} \\ v_{\beta}(x,0) = (x_{+})^{\beta} & \text{on } \{t=0\}, \end{cases}$$

Moreover, such function satisfies

$$-\lim_{t\to 0} t^{1-2s} \partial_t v_\beta = \overline{\kappa}_s \frac{\Gamma(\beta+1)}{\Gamma(\beta-2s+1)} \frac{\sin(\pi(\beta-s))}{\sin(\pi(\beta-2s))} (x_+)^{\beta-2s} ds$$

Equivalently, in \mathbb{R} we have

$$(-\Delta)^{s}(x_{+})^{\beta} \stackrel{k}{=} \frac{\Gamma(\beta+1)}{\Gamma(\beta-2s+1)} \frac{\sin(\pi(\beta-s))}{\sin(\pi(\beta-2s))} (x_{+})^{\beta-2s} \quad for \quad x > 0,$$
(2.8)

where the equality must be understood in the sense of Definition 2.1, and with $k > \beta - 2s$.

Proof. We will show⁶ (2.8). Assume first $\beta \in (-1, 2s)$. Then, at x = 1 we have:

$$(-\Delta)^{s}(x_{+})^{\beta} = c_{1,s} \int_{\mathbb{R}} \left(1 - (1+y)_{+}^{\beta} \right) \frac{dy}{|y|^{1+2s}} = c_{1,s} \frac{2\pi\Gamma(-2s)}{\Gamma(-\beta)\Gamma(1-2s+\beta)} \frac{\cos(\pi s)\sin(\pi(\beta-s))}{\sin(\pi\beta)\sin(\pi(\beta-2s))}$$

where we used [0, Preposition 4.2]

where we used [9, Proposition 4.3].

Moreover, by definition of $(-\Delta)^s$, we have

$$c_{1,s} = \frac{4^{s}\Gamma(s+\frac{1}{2})}{\sqrt{\pi}|\Gamma(-s)|} = \frac{2\Gamma(2s)}{|\Gamma(-s)|\Gamma(s)|} = \frac{2s}{\pi}\Gamma(2s)\sin(\pi s),$$

where we used

$$\Gamma(z+\frac{1}{2})\Gamma(z) = 2^{1-2z}\sqrt{\pi}\Gamma(2z), \qquad \Gamma(z+1) = z\Gamma(z), \qquad \text{and} \qquad \Gamma(1-z)\Gamma(z) = \frac{\pi}{\sin(\pi z)}.$$

Combining the previous expressions, we find that at x = 1

$$(-\Delta)^{s}(x_{+})^{\beta} = \frac{2s}{\pi} \Gamma(2s) \sin(\pi s) \frac{2\pi\Gamma(-2s)}{\Gamma(-\beta)\Gamma(1-2s+\beta)} \frac{\cos(\pi s) \sin(\pi(\beta-s))}{\sin(\pi\beta) \sin(\pi(\beta-2s))}$$
$$= \frac{\Gamma(\beta+1)}{\Gamma(\beta-2s+1)} \frac{\sin(\pi(\beta-s))}{\sin(\pi(\beta-2s))},$$

where we used again the properties of the Γ function and that $\sin(2\pi s) = 2\sin(\pi s)\cos(\pi s)$. Thus, by homogeneity, (2.8) follows in case $\beta < 2s$.

Once we have the result for $\beta \in (-1, 2s)$, the general result follows by induction on $\lfloor \beta \rfloor$, by noticing that we can take derivatives in (2.8).

As a consequence of the previous Lemma, we find:

Lemma 2.5. Let $\beta > -1$, and v_{β} be as in Lemma 2.4. Then, v_{β} solves (2.3) if and only if

$$\frac{\Gamma(\beta+1)}{\Gamma(\beta-2s+1)\Gamma(2s)} = \frac{1}{\pi} \frac{\sin(\pi(\beta-2s))\sin(\pi s)}{\sin(\pi(\beta-s))}.$$
(2.9)

In particular, the exponents β_k are characterized as solutions to this equation.

Proof. The statement is equivalent to (2.2) in the generalized sense. Thus, we need

$$(-\Delta)^s (x_+)^\beta = \frac{c_{1,s}}{2s} (x_+)^{\beta - 2s}$$

Thanks to the previous Lemma, this is equivalent to

$$\frac{\Gamma(\beta+1)}{\Gamma(\beta-2s+1)}\frac{\sin(\pi(\beta-s))}{\sin(\pi(\beta-2s))} = \frac{1}{\pi}\Gamma(2s)\sin(\pi s),$$

and the result follows.

 $^{^{6}\}mathrm{Equivalently},$ one could do these computations in the extended variables, see [19, Appendix A] for a related result.

Let us next analyze the equation (2.9).

Lemma 2.6. There exists a sequence $0 \leq \beta_0 < \beta_1 < \beta_2 < ... \rightarrow \infty$ of solutions of (2.9). Moreover, they satisfy

$$\beta_0 = \max\{2s - 1, 0\},\$$

$$\beta_k \in (k + \beta_0, k + s)$$
 and $\beta_{k+1} > 1 + \beta_k$ for $k \ge 1$.

In particular, we have $\beta_1 = 2s + \alpha_s$, with

$$\alpha_s \in (0, 1-s) \qquad and \qquad 2s + \alpha_s > 1$$

for all $s \in (0, 1)$.

Proof. Let us first notice that both $\beta = 2s - 1$ and $\beta = 0$ are solutions of (2.9). We let

$$\beta_0 := \max\{2s - 1, 0\},\$$

and look for solutions $\beta > \beta_0$.

We consider the two functions

$$h_1(\beta) = \frac{\Gamma(\beta - 2s + 1)\Gamma(2s)}{\Gamma(\beta + 1)}$$

and

$$h_2(\beta) = \frac{\pi \sin(\pi(\beta - s))}{\sin(\pi(\beta - 2s))\sin(\pi s)}$$

and we need to study the intersection of their graphs for $\beta \geq \beta_0$.

Let us first analyse the two functions separately, and then combine the informations in order to see their intersections.

First notice that, by definition of the Beta function and its relation to the Gamma function, for $\beta > 2s - 1$ we have

$$h_1(\beta) = B(\beta - 2s + 1, 2s) = \int_0^1 t^{\beta - 2s} (1 - t)^{2s - 1} dt.$$

On the other hand, using $\sin(a+b) = \sin a \cos b + \sin b \cos a$ we find

$$h_2(\beta) = \frac{\pi}{\tan(\pi(\beta - 2s))} + \frac{\pi}{\tan(\pi s)}.$$

The function h_1 satisfies:

$$\lim_{\beta \downarrow 2s-1} h_1(\beta) = +\infty, \qquad h_1(0) = \frac{\pi}{\sin(2\pi s)}, \qquad \lim_{\beta \to +\infty} h_1(\beta) = 0,$$

and

 $h_1(\beta)$ is positive, decreasing, and convex in $(2s - 1, \infty)$.

The function h_2 satisfies:

$$h_2(\beta)$$
 is 1-periodic, $\lim_{\beta \downarrow 2s-1} h_2(\beta) = +\infty, \qquad h_2(0) = \frac{\pi}{\sin(2\pi s)}$

 $h_2(\beta)$ is decreasing in the intervals $(2s-1,2s) + \mathbb{N}$, convex in $(2s-1,2s-\frac{1}{2}) + \mathbb{N}$,

it is positive in the intervals $(2s - 1, s) + \mathbb{N}$, and negative in $(s, 2s) + \mathbb{N}$.

We then split the analysis into three cases:

Case 1. Assume $s = \frac{1}{2}$. This is the easiest case, and we have $h_1(\beta) = 1/\beta$ and $h_2(\beta) = \pi/\tan(\pi(\beta - 1))$. It is easy to see that these two functions intersect exactly once at each interval (k, k + 1), with $k \in \mathbb{N}$, at a point $\beta_k \in (k, k + \frac{1}{2})$ for $k \ge 1$ satisfying $\beta_{k+1} > 1 + \beta_k$.

Case 2. Assume $s > \frac{1}{2}$. In this case, $\beta_0 = 2s - 1$. Then, we claim that the functions h_1 and h_2 intersect exactly once in each of the intervals $(2s - 1 + k, 2s + k), k \in \mathbb{N}, k \ge 1$.

Indeed, notice first that h_1 is continuous, positive, and decreasing in $(2s - 1, \infty)$, while h_2 is continuous, positive, and decreasing in (2s - 1 + k, s + k), $k \in \mathbb{N}$, with $h_2 \to +\infty$ as $\beta \downarrow 2s - 1 + k$ and $h_2(s + k) = 0$. Thus, these two functions intersect at least once in each of the intervals (2s - 1 + k, s + k), $k \ge 1$, and they never intersect in [s + k, 2s + k] (since $h_2 \le 0$ therein).

It remains to see that they cannot intersect twice in one of these intervals. For this, we notice that both h_1 and h_2 are differentiable in the intervals (2s - 1 + k, s + k). Moreover, for any $\beta \in (2s - 1 + k, s + k)$ and $k \ge 1$ we have

$$h_1'(\beta) \ge h_1'(2s) = -\int_0^1 |\log t| (1-t)^{2s-1} dt \ge -1,$$
$$h_2'(\beta) = -\frac{\pi^2}{\sin^2(-\pi(\beta - 2s))} \le -\pi^2.$$

This means that $h'_2 < h'_1$ in each of these intervals, and therefore they cannot intersect twice in (2s - 1 + k, s + k), for $k \ge 1$, thanks to Rolle's theorem.

It only remains to see that the two functions do not intersect in the first interval (2s-1, s). In that case, however, we are looking for solutions of $(-\Delta)^s_{\mathbb{R}_+} x^\beta = 0$ in \mathbb{R}_+ , with $\beta \in (2s-1, s)$, which we know they do not exist.

Case 3. Assume $s < \frac{1}{2}$. In this case, $\beta_0 = 0$. Then, we claim that the functions h_1 and h_2 intersect exactly once in each of the intervals $(2s - 1 + k, 2s + k), k \in \mathbb{N}, k \ge 1$.

Notice first that h_1 is continuous, positive, and decreasing in $[0, \infty)$, while h_2 is continuous, positive, and decreasing in (2s - 1 + k, s + k), $k \in \mathbb{N}$, with $h_2 \to +\infty$ as $\beta \downarrow 2s - 1 + k$ and $h_2(s + k) = 0$. Thus, these two functions intersect at least once in each of the intervals (2s - 1 + k, s + k), $k \ge 1$, and they never intersect in [s + k, 2s + k] (since $h_2 \le 0$ therein).

To see that they cannot intersect twice in one of these intervals, notice that for $\beta \in (k+s, k+2s)$ the two functions cannot intersect —since h_2 is negative therein—, while for $\beta \in (2s-1+k, k), k \geq 1$, they cannot intersect either —since $h_1 < h_1(0) = h_2(0) = h_2(k) < h_2$ therein.

Instead, for $\beta \in (k, k+s), k \ge 1$, we have⁷

$$h_1'(\beta) \ge h_1'(1) = -\int_0^1 t^{1-2s} |\log t| (1-t)^{2s-1} dt \ge -\frac{1}{2s},$$
$$h_2'(\beta) \le h_2'(1) = -\frac{\pi^2}{\sin^2(\pi(1-2s))} \le -\frac{1}{s^2}.$$

This means that in such intervals we have $h'_1 > h'_2$, and thus the functions cannot intersect twice in (k, k+s) for all $k \ge 1$ by Rolle's theorem.

Notice also that we have showed that $\beta_k \in (k, k+s)$ for all $k \ge 1$. Moreover, since h_1 is decreasing and h_2 is periodic, then $\beta_{k+1} > 1 + \beta_k$.

Finally, notice that in the interval (0, 2s) we know that there are no solutions to $(-\Delta)_{\mathbb{R}_+}^s x^\beta = 0$ in \mathbb{R}_+ for $\beta \in (0, 2s)$, and we are done.

⁷This can be seen by using that $t^{1-2s} < 1$ and that, since $|\log t|$ is decreasing and $(1-t)^{2s-1}$ is increasing, then $\int_0^1 |\log t| (1-t)^{2s-1} dt \le \int_0^1 |\log t| dt \int_0^1 (1-t)^{2s-1} dt = \frac{1}{2s}$.

3. LIOUVILLE THEOREMS ON THE HALF-SPACE

3.1. Solutions with growth larger than 2s. We denote $\mathbb{R}^N_+ = \{x = (x', x_N) \in \mathbb{R}^{N-1} \times \mathbb{R} : x_N > 0\}$ and $B'_r(z)$ the ball in \mathbb{R}^{N-1} with radius r centred at z, while $B^+_r := B_r \cap \mathbb{R}^N_+$.

Definition 3.1. ⁸ Let U be an open subset of \mathbb{R}^N with $0 \in \overline{U}$. Let $u \in H^s(U)$, $f \in L^1_{loc}(U)$ and $\alpha < 1$. We say that u is a solution to $(-\Delta)^s_{\mathbb{R}^N_+} u \stackrel{\alpha}{=} f$ in U if for all $R \ge 1$, there exist a constant $c_R \in \mathbb{R}$ and a function $\mathcal{R}_R \in C^{0,1}(B_{R/2})$, such that

$$(-\Delta)^s_{\mathbb{R}^N_+}(\chi_R u) = f + c_R + \mathcal{R}_R \qquad in \ U \cap B^+_{R/2}, \tag{3.1}$$

with $\mathcal{R}_R(0) = 0$ and $\|\nabla \mathcal{R}_R\|_{L^{\infty}(B_{R/2})} \leq C_0 R^{\alpha - 1}$.

Remark 3.2. (i) We observe that if $(-\Delta)_{\mathbb{R}^N^+}^s u \stackrel{\alpha}{=} 0$ in U and $|u(x)| \leq C(1+|x|^{2s-\varepsilon})$, with $\varepsilon > 0$, then $(-\Delta)_{\mathbb{R}^N}^s u=0$ in U.

(ii) If u satisfies (3.1) and $|\stackrel{+}{u}(x)| \leq C_0(1+|x|^{2s+\alpha})$. Then for any open set $U' \subset U$ with $U' \cap B_1^+ \neq \emptyset$, there exists $C = C(N, s, \alpha, U')$ such that

$$|c_R| \le C \left(C_0 R^{\alpha} + \|f\|_{L^1(U')} \right).$$
(3.2)

Indeed, consider $\phi \in C_c^{\infty}(U' \cap B_1^+)$ with $\int_{\mathbb{R}^N} \phi \, dx = 1$. Then multiply (3.1) by ϕ and integrate over \mathbb{R}^N_+ to get

$$c_{R} = \int_{\mathbb{R}^{N}} \chi_{R} u(-\Delta)^{s}_{\mathbb{R}^{N}_{+}} \phi \, dx - \int_{U'} \phi f \, dx - \int_{U'} \mathcal{R}_{R} \phi \, dx.$$

nce $|(-\Delta)^{s}_{\mathbb{R}^{N}_{+}} \phi(x)| \leq C(1+|x|)^{-N-2s}$, we get (3.2).

The following result is contained in [12].

Lemma 3.3. Let $v \in H^s(B_2^+) \cap L^{\infty}(\mathbb{R}^N)$ be a solution to

$$(-\Delta)^{s}_{\mathbb{R}^{N}_{+}}v = g \quad in \quad B^{+}_{2},$$
 (3.3)

with zero Dirichlet or Neumann boundary conditions on $\partial \mathbb{R}^N_+ \cap B_2$. Then

$$||v||_{C^{\beta}(\overline{B_{1}^{+}})} \leq C(N, s, \beta)(||v||_{L^{\infty}(\mathbb{R}^{N})} + ||g||_{L^{\infty}(\mathbb{R}^{N})})$$

with $\beta = 2s - 1$ if 2s > 1 and $\beta \in (0, 2s)$ if $2s \leq 1$. If moreover $g \in C^k(B_2)$, k = 1, 2, then

$$\|\nabla_{x'}^{k}v\|_{L^{\infty}(B_{1}^{+})} \leq C(N,s)\left(\|v\|_{L^{\infty}(\mathbb{R}^{N})} + \|g\|_{C^{k}(B_{2})}\right).$$

3.2. Liouville theorem.

Si

Lemma 3.4. Let $u \in H^s_{loc}(\overline{\mathbb{R}^N_+})$ satisfy, for $\alpha < 1$, with $2s + \alpha \neq 1$ and $2s + \alpha < 2$, $|u(x)| \leq C(1 + |x|)^{2s + \alpha}$

and

$$(-\Delta)^s_{\mathbb{R}^N_+} u \stackrel{\alpha}{=} 0 \qquad in \ \mathbb{R}^N_+,$$

⁸This definition, is slightly different from Definition 2.1. Here, we ask more regularity on the reminder \mathcal{R}_R to simplify the proofs.

with zero Dirichlet or Neumann boundary conditions on $\partial \mathbb{R}^N_+$. Then $u(x', x_N) = a \cdot x' + b(x_N)$ in the Neumann case, while in the Dirichlet case $u(x', x_N) = a \cdot x' x_N^{2s-1} + \omega(x_N)$. Here $a \in \mathbb{R}^{N-1}$ is a constant and ω solves

$$(-\Delta)^s_{\mathbb{R}_+}\omega \stackrel{\alpha}{=} 0 \qquad in \ \mathbb{R}_+$$

Proof. By definition, we have

$$(-\Delta)^{s}_{\mathbb{R}^{N}_{+}}(\chi_{R}u) = c_{R} + \mathcal{R}_{R} \qquad \text{in } B^{+}_{R/2}.$$
 (3.4)

Next, let $v(x) = (\chi_R u)(Mx)$, for M = R/2. Then

$$(-\Delta)^s_{\mathbb{R}^N_+} v = M^{2s} c_R + M^{2s} \mathcal{R}_R(M \cdot) \qquad \text{in } B_1^+.$$

Now by Lemma 3.3,

$$\|\nabla_{x'}v\|_{L^{\infty}(B_{1/2})} \leq C(N,s) \left(\|v\|_{L^{\infty}(\mathbb{R}^N)} + M^{2s}|c_R| + \|M^{2s}\nabla_{x'}\mathcal{R}_R(Mx)\|_{L^{\infty}(B_1)} \right).$$

Hence, recalling the definition of \mathcal{R}_R , and Remark 3.2(*ii*), we get

$$R \| \nabla_{x'}(\chi_R u) \|_{L^{\infty}(B_{R/4})} \le C(N, s) R^{2s + \alpha}.$$

Since $|u(x)\nabla_{x'}\chi_R(x)| \leq CR^{2s+\alpha-1}$, we deduce that, for all R > 1,

$$\|\nabla_{x'} u\|_{L^{\infty}(B_{R/4})} \le C(N, s) R^{2s + \alpha - 1}.$$

As a consequence

$$|\nabla_{x'}u(x)| \le C(N,s)(1+|x|^{2s+\alpha-1}) \qquad \text{for all } x \in \mathbb{R}^N_+.$$
(3.5)

Differentiating (3.4) in x' and using Lemma 3.3, we obtain $\nabla_{x'} u \in H^s_{loc}(\overline{\mathbb{R}^N_+})$ and

$$(-\Delta)^s_{\mathbb{R}^N_+}(\chi_R \nabla_{x'} u) = -(-\Delta)^s_{\mathbb{R}^N_+}(u \nabla_{x'} \chi_R) + \nabla_{x'} \mathcal{R}_R \quad \text{in } B^+_{R/2}.$$
(3.6)

Thanks to (3.5) and the fact that $|\nabla_{x'}\mathcal{R}_R| + |(-\Delta)^s_{\mathbb{R}^N_+}(u\nabla_{x'}\chi_R)| \leq CR^{\alpha-1}$ in $B^+_{R/2}$, we can send $R \to \infty$ to deduce that

$$(-\Delta)^s_{\mathbb{R}^N_+} \nabla_{x'} u = 0 \qquad \text{in } \mathbb{R}^N_+$$

with $\nabla_{x'} u \in H^s_{loc}(\overline{\mathbb{R}^N_+})$ satisfies the same boundary condition as u. In view of (3.5) and since $2s + \alpha - 1 < 2s$, by applying the Liouville theorems in [12], we deduce that $\nabla_{x'} u(x', x_N)$ is a constant vector on \mathbb{R}^N_+ in the Neumann case and is proportional to $x_N^{\max(2s-1,0)}$ in the Dirichlet case.

Lemma 3.5 (1D Liouville theorem). Let $s \in (0,1)$, and let α_s be given by Lemma 2.6. Let $\omega \in H^s_{loc}(\overline{\mathbb{R}_+})$ satisfy

$$|\omega(x)| \le C(1+|x|)^{2s+\alpha}, \quad \text{with} \quad \alpha < \alpha_s, \tag{3.7}$$

and

$$(-\Delta)^s_{\mathbb{R}_+}\omega \stackrel{\alpha}{=} 0 \qquad in \ \mathbb{R}_+$$

with zero Dirichlet or zero Neumann at $\partial \mathbb{R}_+$. Then, $\omega(x) = c_1 + c_2 x^{2s-1}$, for some $c_1, c_2 \in \mathbb{R}$.

Proof. By Lemma 3.3, $(\omega - \omega(0))1_{\mathbb{R}_+} \in H^s_{loc}([0,\infty)) \cap C([0,\infty))$. In addition, we have that

$$L_s\left[(\omega - \omega(0))\mathbf{1}_{\mathbb{R}_+}\right] \stackrel{\alpha}{=} 0 \qquad \text{in } \mathbb{R}_+$$

where

$$L_s := (-\Delta)^s - a_s x^{-2s}.$$

By [1, Lemma 3.3], there exists $W \in H^1_{loc}(\overline{\mathbb{R}^2_+}; t^{1-2s} dx dt) \cap C(\overline{\mathbb{R}^+_2})$ such that

$$|W(x,t)| \le C(1 + (t + |x|)^K), \tag{3.8}$$

for some constants $K \in \mathbb{N}, C > 0$, and

$$\begin{cases} \operatorname{div}(t^{1-2s}\nabla W) = 0 & \text{in } \mathbb{R} \times (0,\infty) \\ -t^{1-2s}\partial_t W = \overline{\kappa}_s a_s x^{-2s} W & \text{on } \mathbb{R}_+ \times \{0\} \\ W = (\omega - \omega(0)) \mathbb{1}_{\mathbb{R}_+} & \text{on } \mathbb{R} \times \{0\}. \end{cases}$$

Thanks to Lemma 2.2, for all r > 0,

$$W(r\cos\theta, r\sin\theta) = \sum_{k=0}^{\infty} b_k r^{\beta_k} \psi_k(\theta)$$

By the Parseval identity and (3.8), we get

$$\sum_{k=0}^{\infty} b_k^2 r^{2\beta_k} = \int_0^{\pi} W^2(r\cos\theta, r\sin\theta)\sin(\theta)^{1-2s} \, d\theta \le C(1+r)^{2K}$$

Hence, letting r to ∞ we find that $W(r\cos\theta, r\sin\theta) = \sum_{\beta_k \leq K} b_k r^{\beta_k} \psi_k(\theta)$. As a consequence

$$W(r,0) = \sum_{\beta_k \le K} b_k r^{\beta_k} \psi_k(0).$$

Recall that $2s + \alpha < \beta_1 < \beta_k$ for all k > 1. Therefore since $W(r, 0) = \omega(r) - \omega(0)$ we then get from (3.7) and the above identity that $b_k = 0$ for all k > 1, so that $\omega(r) = \omega(0) + b_0 r^{\beta_0} \psi_0(0)$.

Combining Lemma 3.5, Lemma 3.4, and the regularity theory in [2, 12], where have that $u \in C^{2s-1+\varepsilon}(\overline{B_1^+})$ in the zero Neuman case, we get the

Theorem 3.6 (Liouville theorem: Neumann and Dirichlet). Let $s \in (0,1)$ and let $\alpha_s \in$ (0, 1-s) be given by Lemma 2.6. Let $u \in H^s_{loc}(\overline{\mathbb{R}^N_+})$ satisfy,

$$|u(x)| \le C_0 (1+|x|)^{2s+\alpha}, \quad with \quad \alpha < \alpha_s,$$

and

$$(-\Delta)^s_{\mathbb{R}^N_+} u \stackrel{\alpha}{=} 0.$$

- (i) If u satisfies zero Neumann boundary condition on ∂ℝ^N₊, then u(x', x_N) = B + A ⋅ x' for all x ∈ ℝ^N₊, for some constants B ∈ ℝ and A ∈ ℝ^{N-1}.
 (ii) If s > ¹/₂ and u = 0 on ∂ℝ^N₊ then u(x) = (B + A ⋅ x')x^{2s-1}_N for all x = (x', x_N) ∈ ℝ^N₊, for some constants B ∈ ℝ and A ∈ ℝ^{N-1}.

In this Section we establish some new estimates related to the regional fractional Laplacian. To alleviate the notations, we assume in this section that $c_{N,s}$, the constant in (1.2), is equal to 1. We start with the following:

Lemma 4.1. Let $s \in (0,1)$, $\alpha \in (0,1)$ be such that $1 < 2s + \alpha < 2$, $\Omega \subset \mathbb{R}^N$ be a $C^{2s+\alpha}$ domain, and $u \in C^{2s+\alpha}(\overline{\Omega})$ satisfying $\partial_{\nu}u = 0$ on $\partial\Omega$. Then,

$$\|(-\Delta)^{s}_{\Omega}u\|_{C^{\alpha}(\overline{\Omega})} \leq C\|u\|_{C^{2s+\alpha}(\overline{\Omega})},$$

with C depending only on N, s, α , and Ω .

Proof. It suffices to see that, for any $x' \in B_r(x)$, where $B_{2r}(x) \subset \Omega$, we have

$$\left| (-\Delta)^s_{\Omega} u(x) - (-\Delta)^s_{\Omega} u(x') \right| \le Cr^{\alpha};$$

see e.g. [13, Appendix A].

For this, we write

$$(-\Delta)^{s}_{\Omega}u(x') = \int_{(-x'+\Omega)\cap B_r} \frac{u(x') - u(x'+z)}{|z|^{N+2s}} dz + \int_{(-x'+\Omega)\setminus B_r} \frac{u(x') - u(x'+z)}{|z|^{N+2s}} dz$$
$$=: I_1(x') + I_2(x'),$$

for any $x' \in B_r(x)$. Notice that r > 0 is fixed, and that $B_r(x') \subset \Omega$ for all such x'. Let us now treat the two terms I_1 and I_2 separately.

For I_1 we have

$$I_1(x') = \int_{B_r} \frac{2u(x') - u(x'+z) - u(x'-z)}{|z|^{N+2s}} dz = \int_{B_r} \int_0^1 [\nabla u(x'+tz) - \nabla u(x'-tz)] \cdot z \frac{dt \, dz}{|z|^{N+2s}}$$

Thus, since

$$\left|\nabla u(x+tz) - \nabla u(x-tz) - \nabla u(x'+tz) + \nabla u(x'-tz)\right| \le C|z|^{2s+\alpha-1},$$

we deduce that

$$|I_1(x) - I_1(x')| \le C \int_{B_r} |z|^{2s+\alpha-1} \frac{|z|dz}{|z|^{N+2s}} \le Cr^{\alpha}.$$

For I_2 we have

$$\nabla I_2(x') = \int_{\Omega \setminus B_r(x')} \frac{\nabla u(x') - \nabla u(y)}{|x' - y|^{N+2s}} dy + \int_{\partial \Omega} \frac{u(x') - u(y)}{|x' - y|^{N+2s}} \nu(y) \, d\sigma(y).$$

The first term can be bounded as follows.

$$\left| \int_{\Omega \setminus B_r(x')} \frac{\nabla u(x') - \nabla u(y)}{|x' - y|^{N+2s}} dy \right| \le C \int_{\Omega \setminus B_r(x')} \frac{|x' - y|^{2s+\alpha-1}}{|x' - y|^{N+2s}} dy \le Cr^{\alpha-1}.$$

For the second term in I_2 we have notice that, since $\partial_{\nu} u = 0$ on $\partial \Omega$, then

$$\left|u(x') - u(x^*)\right| \le Cd^{2s+\alpha},$$

where $x^* \in \partial \Omega$ is the projection of x' onto $\partial \Omega$, and $d = \operatorname{dist}(x', \partial \Omega) \ge r$. Then

$$|u(x') - u(y)| \le Cd^{2s+\alpha} + |u(x^*) - u(y)|,$$

and therefore

$$\left| \int_{\partial\Omega} \frac{u(x') - u(y)}{|x' - y|^{N+2s}} \, \nu(y) \, d\sigma(y) \right| \le C d^{\alpha - 1} + \left| \int_{\partial\Omega} \frac{u(x^*) - u(y)}{|x' - y|^{N+2s}} \, \nu(y) \, d\sigma(y) \right|.$$

Using that $\partial \Omega$ is locally a $C^{2s+\alpha}$ graph, we will show that

$$\left| \int_{\partial\Omega} \frac{u(x^*) - u(y)}{|x' - y|^{N+2s}} \,\nu(y) \,d\sigma(y) \right| \le C d^{\alpha - 1},\tag{4.1}$$

and hence, combining the previous inequalities and using that d > r, we find

$$|\nabla I_2(x')| \le Cr^{\alpha - 1}$$

Thus, since this holds for any $x' \in B_r(x)$, we deduce

$$|I_2(x) - I_2(x')| \le C|x - x'|r^{\alpha - 1} \le Cr^{\alpha},$$

and the result follows.

It only remains to prove (4.1). Clearly, it is enough to show

$$\left| \int_{\partial\Omega \cap B_{\rho_0}(x^*)} \frac{u(x^*) - u(y)}{|x' - y|^{N+2s}} \nu(y) \, d\sigma(y) \right| \le C d^{\alpha - 1},\tag{4.2}$$

for some small $\rho_{\circ} > 0$. For this, notice that

$$u(y) = u(x^*) + \nabla u(x^*) \cdot (y - x^*) + O(|y - x^*|^{2s + \alpha})$$

and

$$\nu(y) = \nu(x^*) + O(|y - x^*|^{2s + \alpha - 1}).$$

Therefore,

$$(u(y) - u(x^*))\nu(y) = \nabla u(x^*) \cdot (y - x^*)\nu(x^*) + O(|y - x^*|^{2s + \alpha}).$$

Since $\nu(x^*)$ and $\nabla u(x^*)$ are fixed (and bounded), then

$$\left| \int_{\partial\Omega \cap B_{\rho_{\circ}}(x^{*})} \frac{u(x^{*}) - u(y)}{|x' - y|^{N+2s}} \nu(y) \, d\sigma(y) \right| \leq C \left| \int_{\partial\Omega \cap B_{\rho_{\circ}}(x^{*})} (y - x^{*}) \, \frac{d\sigma(y)}{|x' - y|^{N+2s}} \right| + C d^{\alpha - 1}.$$

To bound the last integral, let us assume that $x^* = 0$ and that $\partial \Omega \cap B_{\rho_0}$ can be written as a graph in the $e_N = -\nu(x^*)$ direction, i.e., $\partial \Omega \cap B_{\rho_o}(x^*) = \{x_N = g(x_1, ..., x_{N-1})\}$. We denote $z = (x_1, ..., x_{N-1})$, and then

$$\int_{\partial\Omega\cap B_{\rho_{\circ}}} (y-x^{*}) \, \frac{d\sigma(y)}{|x'-y|^{N+2s}} = \int_{B_{\rho_{\circ}}'} \left(z,g(z)\right) \frac{J(z)dz}{\left(|d-g(z)|^{2}+|z|^{2}\right)^{\frac{N+2s}{2}}},$$

where J(z) is the Jacobian of the change of variables, and $B'_{\rho_o} \subset \mathbb{R}^{N-1}$. Since $g \in C^{2s+\alpha}$ and $g(0) = |\nabla g(0)| = 0$, then $|g(z)| \leq C|z|^{2s+\alpha}$, and the last component of such integral can be bounded by

$$\int_{B'_{\rho_{0}}} |g(z)| \frac{J(z)dz}{\left(|d-g(z)|^{2}+|z|^{2}\right)^{\frac{N+2s}{2}}} \leq C \int_{B'_{\rho_{0}}} |z|^{2s+\alpha} \frac{dz}{\left(d^{2}+|z|^{2}\right)^{\frac{N+2s}{2}}} \leq C d^{\alpha-1},$$

where we used⁹ that $|d - g(z)|^2 + |z|^2 \ge \frac{1}{2}d^2 - |g(z)|^2 + |z|^2 \ge \frac{1}{2}(d^2 + |z|^2)$ in B_{ρ_o} . To control the first n-1 components, we symmetrize the integral, and then we need to

bound 1

$$\frac{1}{2} \int_{B'_{\rho_0}} z \left\{ \frac{J(z)}{\left(|d - g(z)|^2 + |z|^2\right)^{\frac{N+2s}{2}}} - \frac{J(-z)}{\left(|d - g(-z)|^2 + |z|^2\right)^{\frac{N+2s}{2}}} \right\} dz.$$

⁹It is easy to see that $|a - b|^2 \ge \frac{1}{2}a^2 - b^2$ for all $a, b \in \mathbb{R}$.

As before, we have $|d - g(\pm z)|^2 + |z|^2 \approx d^2 + |z|^2$, and hence

 $\left| \left(|d - g(z)|^2 + |z|^2 \right)^{-\frac{N+2s}{2}} - \left(|d - g(-z)|^2 + |z|^2 \right)^{-\frac{N+2s}{2}} \right| \le C \left| g(z) - g(-z) \right| \left(d^2 + |z|^2 \right)^{-\frac{N+2s+1}{2}}.$ Moreover,

$$|J(z) - J(-z)| \le C|z|^{2s+\alpha-1}$$
 and $|g(z) - g(-z)| \le C|z|^{2s+\alpha}$.

Therefore,

$$\left| \int_{B'_{\rho_{\circ}}} z \frac{J(z)dz}{\left(|d-g(z)|^{2}+|z|^{2}\right)^{\frac{N+2s}{2}}} \right| \leq C \int_{B'_{\rho_{\circ}}} |z| \left\{ \frac{|z|^{2s+\alpha}}{\left(d^{2}+|z|^{2}\right)^{\frac{N+2s+1}{2}}} + \frac{|z|^{2s+\alpha-1}}{\left(d^{2}+|z|^{2}\right)^{\frac{N+2s}{2}}} \right\} dz$$
$$\leq C \int_{B'_{\rho_{\circ}}} \frac{|z|^{2s+\alpha}}{\left(d^{2}+|z|^{2}\right)^{\frac{N+2s}{2}}} dz \leq C d^{\alpha-1}.$$

Hence, (4.2) follows, and we are done.

In case $2s + \alpha < 1$ we have the following.

Lemma 4.2. Let $s, \alpha > 0$ be such that $2s + \alpha < 1$, let $\Omega \subset \mathbb{R}^N$ be any open set, and let $u \in C^{2s+\alpha}(\overline{\Omega})$. Then,

$$\|(-\Delta)^{s}_{\Omega}u\|_{C^{\alpha}(\overline{\Omega})} \leq C \|u\|_{C^{2s+\alpha}(\overline{\Omega})},$$

with C depending only on N, s, Ω and α .

Proof. It is easy to see that

$$\|(-\Delta)_{\Omega}^{s}u\|_{L^{\infty}(\Omega)} \leq \sup_{x\in\Omega} \int_{\Omega} \frac{|u(x)-u(y)|}{|x-y|^{N+2s}} \, dy \leq C \|u\|_{C^{2s+\alpha}(\overline{\Omega})}.$$

Now, for $x, x' \in \Omega$ let r = 2|x - x'|. We have

$$\begin{split} J(x,x') &:= (-\Delta)_{\Omega}^{s} u(x) - (-\Delta)_{\Omega}^{s} u(x') = \int_{|x-y| < r} \frac{u(x) - u(y)}{|x-y|^{N+2s}} \, dy - \int_{|x-y| < r} \frac{u(x') - u(y)}{|x'-y|^{N+2s}} \, dy \\ &+ (u(x) - u(x')) \int_{|x-y| > r} \frac{1}{|x'-y|^{N+2s}} \, dy \\ &- \int_{|x-y| > r} (u(x) - u(y)) \left(\frac{1}{|x-y|^{N+2s}} - \frac{1}{|x'-y|^{N+2s}} \right) \, dy. \end{split}$$

We observe that if |y - x| < r, then $|y - x'| \le 3r/2$ and if |y - x| > r, then $|y - x'| \ge r/2$. Moreover for all $t \in (0, 1)$, if $|x - y| \ge r$, then $|tx + (1 - t)x' - y| \ge |y - x|/2$. We then have

$$\begin{aligned} |J(x,x')| &\leq Cr^{\alpha} \\ &+ Cr^{2s+\alpha} \int_{|x'-y|>r/2} |x'-y|^{-N-2s} \, dy + C \int_0^1 \int_{|x-y|\ge r} \frac{|x-x'||x-y|^{2s+\alpha} dy}{|tx+(1-t)x'-y|^{N+2s+1}} dt \\ &\leq Cr^{\alpha} + C|x-x'| \int_{|x-y|\ge r} |x-y|^{2s+\alpha} |x-y|^{-N-2s-1} \leq Cr^{\alpha}. \end{aligned}$$

The proof is complete.

Finally, we provide a new formula for the regional fractional Laplacian on domains. This can be used to give a different proof of Lemma 4.1 in case $s > \frac{1}{2}$.

Lemma 4.3. Let Ω be a bounded $C^{\max(2s,1)}$ domain. Let $u \in C^{2s+\alpha}(\overline{\Omega})$, with $2s+\alpha > 1$ and $\partial_{\nu}u = 0$ on $\partial\Omega$. Then, for $x \in \overline{\Omega}$ we have

$$(-\Delta)^{s}_{\Omega}u(x) = \int_{\Omega} \frac{u(x) - u(y) - \nabla u(x) \cdot (x - y)}{|x - y|^{N + 2s}} \, dy + \frac{1}{N + 2s - 2} \int_{\partial\Omega} \frac{\left(\nabla u(x) - \nabla u(y)\right) \cdot \nu(y)}{|x - y|^{N + 2s - 2}} \, d\sigma(y).$$

where ν is the unit exterior normal of Ω . Moreover, all the above integrals converge absolutely.

Notice that, for simplicity, we establish the result in case of bounded domains. Still, an analogous result could be proved for unbounded domains.

Proof of Lemma 4.3. We have

$$(-\Delta)^{s}_{\Omega}u(x) = \int_{\Omega} \frac{u(x) - u(y) - \nabla u(x) \cdot (x - y)}{|x - y|^{N + 2s}} \, dy + p.v. \int_{\Omega} \frac{\nabla u(x) \cdot (x - y)}{|x - y|^{N + 2s}} \, dy$$

Define

$$I_{\varepsilon}(x) := \int_{\Omega \cap \partial B_{\varepsilon}(x)} \frac{\nabla u(x) \cdot \eta_{\varepsilon}(y)}{|x - y|^{N + 2s - 2}} d\sigma(y),$$

where η_{ε} is the unit exterior normal of $B_{\varepsilon}(x)$. Using that $\nabla_y |x - y|^{-N-2s+2} = (N + 2s - 2)(x - y)|x - y|^{-N-2s}$, the divergence theorem and the fact that $\nabla u(y) \cdot \nu(y) = 0$ for $y \in \partial\Omega$, we then get

$$\begin{split} (N+2s-2)p.v.\int_{\Omega} \frac{\nabla u(x) \cdot (x-y)}{|x-y|^{N+2s}} \, dy &= p.v. \int_{\partial \Omega} \frac{\nabla u(x) \cdot \nu(y)}{|x-y|^{N+2s-2}} d\sigma(y) \\ &= \int_{\partial \Omega} \frac{[\nabla u(x) - \nabla u(y)] \cdot \nu(y)}{|x-y|^{N+2s-2}} d\sigma(y) + \lim_{\varepsilon \to 0} I_{\varepsilon}(x). \end{split}$$

If $x \in \Omega$ then we immediately get $\lim_{\varepsilon \to 0} I_{\varepsilon}(x) = 0$ and the result follows. We now consider the case $x \in \partial \Omega$, then denote by $H_x := T_x \partial \Omega$ the tangent plane of $\partial \Omega$ at x and H_x^+ the upper half space containing $-\nu(x)$ with boundary H_x . The proof is now complete once we show that

$$I_{\varepsilon}(x) = -C(N)\varepsilon^{1-2s}\nabla u(x) \cdot \nu(x) + o_{\varepsilon}(1) = o_{\varepsilon}(1), \qquad (4.3)$$

where $C(N) = \int_{S^{N-1}_{+}} \theta_N \, d\sigma(\theta)$. We have

$$I_{\varepsilon}(x) = \int_{\Omega \cap \partial B_{\varepsilon}(x)} \frac{\nabla u(x) \cdot \eta_{\varepsilon}(y)}{|y - x|^{N+2s-2}} d\sigma(y) = \varepsilon^{-N-2s+1} \int_{\Omega \cap \partial B_{\varepsilon}(x)} \nabla u(x) \cdot (y - x) d\sigma(y)$$

$$= \varepsilon^{-N-2s+1} \int_{H_{x}^{+} \cap \partial B_{\varepsilon}(x)} \nabla u(x) \cdot (y - x) d\sigma(y) - \varepsilon^{-N-2s+1} \int_{\Omega \triangle H_{x}^{+} \cap \partial B_{\varepsilon}(x)} \nabla u(x) \cdot (y - x) d\sigma(y).$$

(4.4)

By oddness, we have $\int_{H_x^+ \cap \partial B_{\varepsilon}(x)} \nabla u(x) \cdot (y-x) d\sigma(y) = -\varepsilon^N \nabla u(x) \cdot \nu(x) \int_{S_+^{N-1}} \theta_N d\sigma(\theta)$. On the other hand, since Ω is of class C^{2s} , we can write a neighbourhood of x in $\partial\Omega$ as a graph of a function $y_N = \gamma(y') = o(|y'|^{2s})$ over the tangent plane H_x . We then observe that if $y \in \Omega \bigtriangleup H_x^+ \cap \partial B_{\varepsilon}(x)$ then $|y'-x| \le \varepsilon$ and $|y_N| \le \gamma(y')$, so that

$$|\Omega \bigtriangleup H_x^+ \cap \partial B_{\varepsilon}(x)| = \varepsilon^{N-2} o(\varepsilon^{2s}).$$

We thus conclude from this and (4.4) that (4.3) holds.

Remark 4.4. It follows from the above proof that the map

$$x\mapsto \int_{\Omega\cap |x-y|>\varepsilon}\frac{u(x)-u(y)}{|x-y|^{N+2s}}\,dy$$

converges in $C_{loc}(\overline{\Omega})$ to $(-\Delta)^s_{\Omega}u$, as $\varepsilon \to 0$. The proof also shows that for $(-\Delta)^s_{\Omega}u(x)$ to be defined at $x \in \partial\Omega$, then necessarily, $\partial_{\nu}u(x) = 0$. See also [16], where these type of results were obtained.

5. Regional Neumann problem

In this section, we consider Ω a domain of class $C^{\max(1,2s+\alpha)}$ with $0 \in \partial\Omega$ and $\nu(0) = -e_N$. For, $2 > 2s + \alpha > 1$, we suppose that there exists a global diffeomorphism $\psi \in C^{2s+\alpha}(\mathbb{R}^N;\mathbb{R}^N)$, satisfying

$$\partial \Omega \cap B_2 = \psi(B'_2 \times \{0\}), \qquad \Omega \cap B_2 = \psi(B^+_2), \qquad \psi(0) = 0$$
 (5.1)

and

$$-\frac{\partial_{y_N}\psi(y',0)}{|\partial_{y_N}\psi(y',0)|} = \nu(\psi(y',0)) \quad \text{for } y' \in B_2',$$
(5.2)

where ν is the exterior normal of $\partial\Omega$, see [1] for the construction of such map. In the case $2s + \alpha < 1$, we simply assume that $\partial\Omega \cap B_2$ is parameterized by a global diffeo-

morphism $\psi \in C^1(\mathbb{R}^N; \mathbb{R}^N)$ satisfying (5.1).

Next, we define the operator \mathcal{L}_{ψ} by

$$\mathcal{L}_{\psi}w(x) := c_{N,s} \int_{B_2^+} \frac{w(x) - w(y)}{|\psi(x) - \psi(y)|^{N+2s}} Jac_{\psi}(y) \, dy, \tag{5.3}$$

so that for $u(x) = w(\psi(x))$, we get

$$(-\Delta)^{s}_{\psi(B_{2}^{+})}u(\psi(x)) = \mathcal{L}_{\psi}w(x).$$
(5.4)

We also define the bilinear form $\mathcal{D}_{\psi}: H^s(\mathbb{R}^N_+) \times H^s(\mathbb{R}^N_+) \to \mathbb{R}$ corresponding to this operator as

$$\mathcal{D}_{\psi}(w,\varphi) = \frac{c_{N,s}}{2} \int_{B_2^+ \times B_2^+} \frac{(w(x) - w(y))(\varphi(x) - \varphi(y))}{|\psi(x) - \psi(y)|^{N+2s}} Jac_{\psi}(x) Jac_{\psi}(y) \, dy dx.$$

Let $A \in \mathbb{R}^{N-1} = \{y_N = 0\}$ and define

$$q_A(y) = A \cdot y = A \cdot y' \tag{5.5}$$

We now define

$$Q_A : \mathbb{R}^N \to \mathbb{R}^N, \qquad Q_A(x) = q_A(\psi^{-1}(x))$$

We have

$$Q_A(\psi(y)) = q_A(y) = A \cdot y',$$

and thus

$$0 = \partial_{y_N} q_A(y) = \nabla Q_A(\psi(y)) \cdot \partial_{y_N} \psi(y)$$

This, together with (5.2), imply that

$$\partial_{\nu}Q_A = 0 \qquad \text{on } \partial\Omega \cap B_2.$$

Hence, since $Q_A \in C^{2s+\alpha}(B_2 \cap \overline{\Omega})$ for $2s + \alpha > 1$, by Lemma 4.1 and (5.4) we get:

Lemma 5.1. Let \mathcal{L}_{ψ} and q_A be given by (5.4) and (5.5), respectively. Then,

$$\|\mathcal{L}_{\psi}q_A\|_{C^{\alpha}(\overline{B_1^+})} \le C|A|.$$

with C depending only on α , $\|\psi\|_{C^{2s+\alpha}(B_2)}$, N, s.

Proof. It follows from Lemma 4.1 and (5.4).

Let

$$\ell = \ell_{\alpha} = \begin{cases} 1 & \text{for } 2s + \alpha > 1 \\ 0 & \text{for } 2s + \alpha < 1. \end{cases}$$
(5.6)

For r > 0 and $w \in L^2_{loc}(\mathbb{R}^N)$, let $A_{r,w} \in \mathbb{R}^{N-1}$ such that

$$\inf_{A \in \mathbb{R}^N} \|w - \ell A \cdot y'\|_{L^2(B_r^+)}^2 = \|w - \ell A_{r,w} \cdot y'\|_{L^2(B_r^+)}^2.$$
(5.7)

Then

$$\ell A_{r,w} \cdot e_i = \frac{1}{\int_{B_r^+} z_i^2 \, dz} \int_{B_r^+} w(y) y_i \, dy.$$
(5.8)

In particular

$$\int_{B_r^+} (w(y) - \ell A_{r,w} \cdot y') y_i \, dy = 0 \qquad \text{for all } i = 1, \dots, N - 1.$$
(5.9)

Before stating the main result of the present section, we record first the following result from [12].

Lemma 5.2 ([12]). Let $u \in H^s(\mathbb{R}^N_+)$ and $f \in L^\infty(\mathbb{R}^N_+)$ be a solution to $\mathcal{D}_{\psi}(u,\varphi) = \int_{\mathbb{R}^N_+} u\varphi \, dx$ for all $\varphi \in C_c^\infty(B_2)$. Then, for all $\varrho \in (0, 2s)$, there exists $C = C(N, s, \varrho) > 0$ such that

$$\|u\|_{C^{2s-\varrho}(B_1)} \le C\left(\|u\|_{L^2(\mathbb{R}^N_+)} + \|f\|_{L^{\infty}(\mathbb{R}^N_+)}\right)$$

We next show the following.

Proposition 5.3. Let $s \in (0,1)$ and $\alpha_s \in (0,1-s)$ be given by Lemma 2.6. Let $\alpha \in (0,\alpha_s)$, with $2s + \alpha \neq 1$, ψ be as above, with $\|\psi\|_{C^{\max(1,2s+\alpha)}(\mathbb{R}^N)} \leq c_0$. Let $f \in C^{\alpha}(\mathbb{R}^N_+)$ and $w \in H^s(\mathbb{R}^N_+) \cap C(\mathbb{R}^N_+)$ such that w(0) = 0 and

$$\mathcal{D}_{\psi}(w,\varphi) = \int_{\mathbb{R}^N_+} f\varphi \, dx \qquad \text{for all } \varphi \in C^{\infty}_c(B_2).$$

Let $\rho < 2s + \alpha - 1$ if $2s + \alpha > 1$. Then, provided

$$\ell|A_{r,w}| \le c_0 r^{-\varrho},\tag{5.10}$$

there exists $C = C(N, s, \alpha, c_0, \varrho, \beta)$ such that

$$\sup_{r>0} r^{-2s-\beta} \|w - \ell A_{r,w} \cdot y'\|_{L^{\infty}(B_r^+)} \le C(\|w\|_{L^{\infty}(\mathbb{R}^N_+)} + \|f\|_{C^{\alpha}(\mathbb{R}^N_+)}),$$
(5.11)

where $\beta = \alpha - \varrho$, $\ell = \ell_{\alpha}$ and $A_{r,u}$ are given by (5.6) and (5.7) respectively.

Proof. Suppose that (5.11) does not hold. Then for all integer $n \geq 2$, there exist $\psi_n \in C^{\max(1,2s+\alpha)}(\mathbb{R}^N)$, $f_n \in C_c^{\alpha}(\mathbb{R}^N_+)$ and $w_n \in H^s(\mathbb{R}^N_+) \cap C(\mathbb{R}^N_+)$, with $w_n(0) = 0$, such that

$$\|\psi_n\|_{C^{\max(1,2s+\alpha)}(\mathbb{R}^N)} \le c_0, \tag{5.12}$$

$$\|w_n\|_{L^{\infty}(\mathbb{R}^N_+)} + \|f_n\|_{C^{\alpha}(\mathbb{R}^N_+)} \le 1$$
(5.13)

such that

$$\ell |A_{r,w_n}| \le c_0 r^{-\varrho},\tag{5.14}$$

$$\mathcal{D}_{\psi_n}(w_n,\varphi) = \int_{\mathbb{R}^N_+} f_n \varphi \, dx \qquad \text{for all } \varphi \in C_c^\infty(B_2).$$
(5.15)

and

$$\sup_{r>0} r^{-2s-\alpha} \|w_n - \ell A_{r,w_n} \cdot y'\|_{L^{\infty}(B_r^+)} > n.$$

Then, since $2s + \beta > 1$ for $2s + \alpha > 1$, by a well known (see e.g. [18]), we can find sequences $r_n \to 0$ and $\theta_n \to \infty$ such that the function

$$v_n(y) := \frac{1}{r_n^{2s+\alpha}\theta_n} \left(w_n(r_n y) - r_n \ell A_{r_n, w_n} \cdot y' \right)$$

satisfies

$$\|v_n\|_{L^{\infty}(B_1^+)} \ge \frac{1}{4} \tag{5.16}$$

and, in view of (5.14), for all $M \ge 1$ with $Mr_n \le 1$,

$$\|v_n\|_{L^{\infty}(B_M^+)} \le CM^{2s+\beta},\tag{5.17}$$

while for $2s + \alpha < 1$

$$||v_n||_{L^{\infty}(B^+_M)} \le CM^{2s+\beta}$$
 for all $M \ge 1$, (5.18)

Moreover, by construction (recalling (5.9)),

$$v_n(0) = 0, \qquad \int_{B_1^+} v_n(y) y_i \, dy = 0 \qquad \text{for all } i = 1, \dots, N-1.$$
 (5.19)

Now for $Mr_n > 1$ and $2s + \alpha > 1$, we have

$$\|v_n\|_{L^{\infty}(B_M^+)} \le \frac{1}{r_n^{2s+\alpha}\theta_n} (1+r_n^{-\varrho}) \le M^{2s+\beta} + M^{2s+\beta+\varrho} \le 2M^{2s+\alpha}.$$

In view of this, (5.17) and (5.18), we see that whenever $2s + \alpha \neq 1$,

$$||v_n||_{L^{\infty}(B_M^+)} \le CM^{2s+\alpha}$$
 for all $M \ge 1$. (5.20)

Define $\tilde{\psi}_n(x) = \frac{1}{r_n} \psi(r_n x)$ and let $1 < R < \frac{1}{r_n}$. Then by a change of variable in (5.15), we thus get

$$\mathcal{D}_{\widetilde{\psi}_n}(\chi_R v_n, \varphi) = \int_{\mathbb{R}^N_+} G_R^n \varphi \, dx \qquad \text{for all } \varphi \in C_c^\infty(B_{\frac{R}{2}}), \tag{5.21}$$

where

$$G_{R}^{n}(x) := \frac{1}{r_{n}^{\beta}\theta_{n}} f_{n}(r_{n}x) + c_{N,s} \int_{\mathbb{R}^{N}_{+}} \frac{(\chi_{R}(x) - \chi_{R}(y))v_{n}(y)}{|\tilde{\psi}_{n}(x) - \tilde{\psi}_{n}(y)|^{N+2s}} J_{n}(y) \, dy - \frac{\ell}{r_{n}^{\alpha}\theta_{n}} \mathcal{L}_{\psi_{n}}[A_{r_{n},w_{n}} \cdot y'](r_{n}x) + c_{N,s} \int_{\mathbb{R}^{N}_{+}} \frac{(\chi_{R}(x) - \chi_{R}(y))v_{n}(y)}{|\tilde{\psi}_{n}(x) - \tilde{\psi}_{n}(y)|^{N+2s}} J_{n}(y) \, dy - \frac{\ell}{r_{n}^{\alpha}\theta_{n}} \mathcal{L}_{\psi_{n}}[A_{r_{n},w_{n}} \cdot y'](r_{n}x) + c_{N,s} \int_{\mathbb{R}^{N}_{+}} \frac{(\chi_{R}(x) - \chi_{R}(y))v_{n}(y)}{|\tilde{\psi}_{n}(x) - \tilde{\psi}_{n}(y)|^{N+2s}} J_{n}(y) \, dy - \frac{\ell}{r_{n}^{\alpha}\theta_{n}} \mathcal{L}_{\psi_{n}}[A_{r_{n},w_{n}} \cdot y'](r_{n}x) + c_{N,s} \int_{\mathbb{R}^{N}_{+}} \frac{(\chi_{R}(x) - \chi_{R}(y))v_{n}(y)}{|\tilde{\psi}_{n}(x) - \tilde{\psi}_{n}(y)|^{N+2s}} J_{n}(y) \, dy - \frac{\ell}{r_{n}^{\alpha}\theta_{n}} \mathcal{L}_{\psi_{n}}[A_{r_{n},w_{n}} \cdot y'](r_{n}x) + c_{N,s} \int_{\mathbb{R}^{N}_{+}} \frac{(\chi_{R}(x) - \chi_{R}(y))v_{n}(y)}{|\tilde{\psi}_{n}(x) - \tilde{\psi}_{n}(y)|^{N+2s}} J_{n}(y) \, dy - \frac{\ell}{r_{n}^{\alpha}\theta_{n}} \mathcal{L}_{\psi_{n}}[A_{r_{n},w_{n}} \cdot y'](r_{n}x) + c_{N,s} \int_{\mathbb{R}^{N}_{+}} \frac{(\chi_{R}(x) - \chi_{R}(y))v_{n}(y)}{|\tilde{\psi}_{n}(x) - \tilde{\psi}_{n}(y)|^{N+2s}} J_{n}(y) \, dy - \frac{\ell}{r_{n}^{\alpha}\theta_{n}} \mathcal{L}_{\psi_{n}}[A_{r_{n},w_{n}} \cdot y'](r_{n}x) + c_{N,s} \int_{\mathbb{R}^{N}_{+}} \frac{(\chi_{R}(x) - \chi_{R}(y))v_{n}(y)}{|\tilde{\psi}_{n}(x) - \tilde{\psi}_{n}(y)|^{N+2s}} J_{n}(y) \, dy - \frac{\ell}{r_{n}^{\alpha}\theta_{n}} \mathcal{L}_{\psi_{n}}[A_{r_{n},w_{n}} \cdot y'](r_{n}x) + c_{N,s} \int_{\mathbb{R}^{N}_{+}} \frac{(\chi_{R}(x) - \chi_{R}(y))v_{n}(y)}{|\tilde{\psi}_{n}(x) - \tilde{\psi}_{n}(y)|^{N+2s}} J_{n}(y) \, dy - \frac{\ell}{r_{n}^{\alpha}\theta_{n}} \mathcal{L}_{\psi_{n}}[A_{r_{n},w_{n}} \cdot y'](r_{n}x) + c_{N,s} \int_{\mathbb{R}^{N}_{+}} \frac{(\chi_{R}(x) - \chi_{R}(y))v_{n}(y)}{|\tilde{\psi}_{n}(x) - \tilde{\psi}_{n}(y)|^{N+2s}} J_{n}(y) \, dy - \frac{\ell}{r_{n}^{\alpha}\theta_{n}} \mathcal{L}_{\psi_{n}}[A_{r_{n},w_{n}} \cdot y'](r_{n}x) + c_{N,s} \int_{\mathbb{R}^{N}_{+}} \frac{(\chi_{R}(x) - \chi_{R}(y))v_{n}(y)}{|\tilde{\psi}_{n}(x) - \chi_{R}(y)|^{N+2s}} J_{n}(y) \, dy - \frac{\ell}{r_{n}^{\alpha}\theta_{n}} \mathcal{L}_{\psi_{n}}[A_{r_{n},w_{n}} \cdot y'](r_{n}x) + c_{N,s} \int_{\mathbb{R}^{N}_{+}} \frac{(\chi_{R}(x) - \chi_{R}(y))v_{n}(y)}{|\tilde{\psi}_{n}(x) - \chi_{R}(y)|^{N+2s}} J_{n}(y)} \, dy - \frac{\ell}{r_{n}^{\alpha}\theta_{n}} \mathcal{L}_{\psi_{n}}[A_{r_{n},w_{n}} \cdot y'](r_{n}x) + c_{N,s} \int_{\mathbb{R}^{N}_{+}} \frac{(\chi_{R}(x) - \chi_{R}(y))v_{n}(y)}{|\tilde$$

where $J_n(y) := Jac_{\tilde{\psi}_n}(y) \mathbf{1}_{B^+_{\frac{1}{r_n}}}(y)$. Since ψ_n is a global diffeomorphism, by (6.15) and increasing c_0 if necessary, we obtain

 $c_0|x-y| \ge |\widetilde{\psi}_n(x) - \widetilde{\psi}_n(y)| \ge c_0^{-1}|x-y|, \qquad |D\widetilde{\psi}_n(x)| \le c_0 \qquad \text{for all } x, y \in \mathbb{R}^N.$ (5.22) Define, for $x \in B_{R/2}$,

$$F_{R}^{n}(x) := c_{N,s} \int_{\mathbb{R}^{N}_{+}} \frac{(\chi_{R}(x) - \chi_{R}(y))v_{n}(y)}{|\tilde{\psi}_{n}(x) - \tilde{\psi}_{n}(y)|^{N+2s}} J_{n}(y) \, dy$$

= $c_{N,s} \int_{\mathbb{R}^{N}_{+} \cap \{|y| \ge R\}} \frac{(1 - \chi_{R}(y))v_{n}(y)}{|\tilde{\psi}_{n}(x) - \tilde{\psi}_{n}(y)|^{N+2s}} J_{n}(y) \, dy.$

Since $\tilde{\psi}_n(0) = 0$, by the mean value theorem, we have $F_R^n(x) = F_R^n(0) + \mathcal{R}_R^n(x)$, where

$$\mathcal{R}_R^n(x) := -(N+2s)c_{N,s} \int_0^1 \int_{\mathbb{R}^N_+} \frac{(\widetilde{\psi}_n(x) - \widetilde{\psi}_n(y)) \cdot \widetilde{\psi}_n(x)}{|\widetilde{\psi}_n(x) - \widetilde{\psi}_n(y)|} \frac{(1 - \chi_R(y))v_n(y)}{|t\widetilde{\psi}_n(x) - \widetilde{\psi}_n(y)|^{N+2s+1}} J_n(y) \, dy \, dt$$

Note that by (5.22), $|t\tilde{\psi}_n(x) - \tilde{\psi}_n(y)| \ge \frac{c_0}{2}|y|$ whenever $|y| \ge R$ and $|x| \le \frac{R}{2c_0^2}$. Therefore, by (5.20) and (5.22), there exists $C, n_0 > 0$ such that

$$|\nabla \mathcal{R}_R^n(x)| \le CR^{\alpha - 1} \qquad \text{for all } R > 1, \ |x| \le \frac{R}{2c_0^2} \text{ and } n \ge n_0 \tag{5.23}$$

and by construction we have $\mathcal{R}_R^n(0) = 0$. We can thus write

$$G_R^n(x) = H_R^n(x) + c_R^n + \mathcal{R}_R^n(x), \qquad (5.24)$$

with

$$H_R^n(x) = \frac{1}{r_n^\beta \theta_n} \left\{ f_n(r_n x) - f_n(0) + \mathcal{L}_{\psi_n} [A_{r_n, w_n} \cdot y'](r_n x) - \mathcal{L}_{\psi_n} [A_{r_n, w_n} \cdot y'](0) \right\}$$

and

$$c_R^n = \frac{1}{r_n^\beta \theta_n} \left\{ f_n(0) + \mathcal{L}_{\psi_n} [A_{r_n, w_n} \cdot y'](0) \right\} + F_R^n(0).$$

Recalling that $||f_n||_{C^{\alpha}(\mathbb{R}^N_{+})} \leq 1$, we then get from (5.14) and Lemma 5.1 that

$$|H_R^n(x)| \le \frac{r_n^{\alpha}C}{r_n^{\beta}\theta_n} + \ell \frac{r_n^{\alpha-\varrho}}{r_n^{\beta}\theta_n} \le \frac{C(R)}{\theta_n} \qquad \text{for all } x \in B_{R/2}.$$
(5.25)

Let us now show that c_R^n is bounded. For this, we pick $\phi \in C_c^{\infty}(B_{\frac{1}{2c_0^2}}^+)$ with $\int_{\mathbb{R}^N} \phi \, dx = 1$. Then multiply (5.21) by ϕ and integrate over \mathbb{R}^N_+ to get

$$|c_{R}^{n}| \leq \int_{\mathbb{R}^{N}_{+}} |\chi_{R}(x)v_{n}(x)| |\mathcal{L}_{\widetilde{\psi}_{n}}\phi(x)| \, dx + \int_{\mathbb{R}^{N}_{+}} |H_{R}^{n}(x)\phi(x)| \, dx + \int_{\mathbb{R}^{N}_{+}} |\mathcal{R}_{R}^{n}(x)\phi(x)| \, dx \leq C(R).$$
(5.26)

In view of this, (5.21), (5.24), (5.25) and (5.23), we can apply Lemma 5.2, to deduce that $(v_n)_n$ is bounded in $C_{loc}^{s+\delta}(\overline{\mathbb{R}^N_+})$, for some $\delta > 0$, and converges to some $v \in C_{loc}^{s+\delta}(\overline{\mathbb{R}^N_+}) \subset H^s_{loc}(\overline{\mathbb{R}^N_+})$. In addition passing to the limit in (5.16), (5.20) and (5.19), we get

$$|v(x)| \le C(1+|x|^{2s+\alpha}), \tag{5.27}$$

$$\|v\|_{L^{\infty}(B_1^+)} \ge \frac{1}{4}, \quad v(0) = 0, \quad \int_{B_1^+} v(y)y_i \, dx = 0, \text{ for } i = 1, \dots, N-1.$$
 (5.28)

Passing to the limit in (5.21), (5.23) and using (5.26), we get, for all $\phi \in C_c^{\infty}(B_{\underline{Rc_0^2}})$,

$$c_{N,s} \int_{\mathbb{R}^{2N}_+} \frac{((\eta_R v)(x) - (\eta_R v)(y))(\phi(x) - \phi(y))}{|Bx - By|^{N+2s}} \det(B)^2 dy dx = \int_{\mathbb{R}^N_+} (c_R^\infty + \mathcal{R}_R^\infty(x))\phi(x) dx,$$

where $B = \lim_{n \to \infty} D\widetilde{\psi}_n(0) = \lim_{n \to \infty} D\psi_n(0)$ and

$$c_R^{\infty} \in \mathbb{R}, \qquad \mathcal{R}_R^{\infty}(0) = 0, \qquad |\nabla \mathcal{R}_R^{\infty}(x)| \le CR^{\alpha - 1} \qquad \text{for all } R \ge 2 \text{ and } x \in B_{\frac{R}{2c_0^2}}.$$
 (5.29)

Recall from the construction of ψ_n that $B = \text{diag}(1, \ldots, 1, \lambda)$ is a diagonal matrix, for some $\lambda \neq 0$. Hence by scaling and (6.39), we can assume that

$$(-\Delta)^s_{\mathbb{R}^N_+} v \stackrel{\alpha}{=} 0 \qquad \text{in } \mathbb{R}^N_+.$$

Applying Theorem 3.4 and using (6.37), we deduce that $v(x) = b + a \cdot x'$. This is in contradiction with (5.28).

As a consequence, we have :

Corollary 5.4. Let $s \in (0,1)$ and $\alpha_s \in (0,1-s)$ be given by Lemma 2.6. Let $\alpha \in (0,\alpha_s)$, ψ be as above, with $\|\psi\|_{C^{\max(1,2s+\alpha)}(\mathbb{R}^N)} \leq c_0$. Let $f \in C^{\alpha}(\mathbb{R}^N_+)$ and $w \in H^s(B_2^+) \cap L^{\infty}(\mathbb{R}^N_+)$ such that

$$\mathcal{D}_{\psi}(w,\varphi) = \int_{\mathbb{R}^N_+} f\varphi \, dx \qquad \text{for all } \varphi \in C^{\infty}_c(B_2).$$

Then, there exists $C = C(N, s, c_0, \alpha)$ such that for all $z \in B'_{1/2}$ there exists a vector $e = e_z \in \mathbb{R}^{N-1} =: \partial \mathbb{R}^N_+$, with $|e| \leq C$ and

$$\sup_{r>0} r^{-2s-\alpha} \|w - w(z) - e \cdot (y' - z)\|_{L^{\infty}(B^+_r(z))} \le C \left(\|w\|_{L^{\infty}(\mathbb{R}^N_+)} + \|f\|_{C^{\alpha}(\mathbb{R}^N_+)} \right).$$
(5.30)

In particular, if $2s + \alpha > 1$ then $\partial_{y_N} w = 0$ on $B'_{1/2}$.

Proof. Thanks to Lemma 5.2, the function $v_z(x) = \chi_2(x)(w(x+z) - w(z))$ satisfies the hypothesis of Proposition 5.3. We can thus prove (5.30) with z = 0. We assume for simplicity that $\|w\|_{L^{\infty}(\mathbb{R}^N_+)} + \|f\|_{C^{\alpha}(\mathbb{R}^N_+)} \leq 1$.

If $2s + \alpha_s < 1$, then the result clearly holds with e = 0. We start with (5.30) and we distinguish the two cases 2s > 1 and 2s < 1. We start with the case 2s > 1, where Lemma 5.2, yields $|w(y)| \leq C|y|$, so that (6.25) holds with $\rho = 0$. By (5.11) and an (algebraic) iteration argument (see e.g. [5,18]), we can find $e \in \mathbb{R}^{N-1}$ such that

$$||w - e \cdot y'||_{L^{\infty}(B^+_r)} \le Cr^{2s+\alpha}, \qquad |e| \le C.$$
 (5.31)

We finally consider the case 2s < 1. Here, (5.11) implies that $|w(y)| \leq C|y|^{2s+\alpha'}$ as soon as $2s + \alpha' < 1$ (i.e. $\alpha' < 1 - 2s < \alpha$). As a consequence w satisfies (6.25) with $\rho = 1 - 2s - \alpha'$. Therefore choosing α' close to 1 - 2s so that $2s + \alpha - \rho > 1$, we get from (5.11) and an iteration argument, as above, that

$$||w - e \cdot y'||_{L^{\infty}(B^+_r)} \le Cr^{2s + \alpha - \varrho}, \qquad |e| \le C,$$
 (5.32)

which again implies that $|w(y)| \leq C|y|$. Hence w satisfies (5.33) with $\rho = 0$ which, once again by an iteration argument, yields (5.32) with $\rho = 0$.

If 2s = 1 then Lemma 5.2 implies that $|w(y)| \leq C|y|^{1-\varrho}$, for all $\varrho \in (0, 1)$ so that (6.25) holds with $\varrho \in (0, \alpha_{\frac{1}{2}})$. We thus get, from (5.11) and an iteration argument, as above, a vector $e \in \mathbb{R}^{N-1}$ such that

$$\|w - e \cdot y'\|_{L^{\infty}(B_r^+)} \le Cr^{1+\alpha_{\frac{1}{2}}-\varrho}, \qquad |e| \le C.$$
(5.33)

This gives $|w(y)| \leq C|y|$, so that w satisfies (6.25) with $\rho = 0$. Hence as above, we get (5.33) with $\rho = 0$.

We will also need the following interior Schauder estimate.

Lemma 5.5. Let Ω be an open set containing B_2 and $\alpha \in (0,1)$. Let $f \in C^{\alpha}(B_2)$ and $v \in H^s(\Omega)$ be a solution to

$$(-\Delta)^s_{\Omega}v = f \qquad in \ B_2$$

satisfying $C_0 := \int_{\Omega} \frac{|v(y)| dy}{1 + |y|^{N+2s+1}} < \infty$. Then, provided $2s + \alpha \notin \mathbb{N}$,

$$\|v\|_{C^{2s+\alpha}(B_1)} \le C(N, s, \alpha)(C_0 + \|v\|_{L^2(B_2)} + \|f - f(0)\|_{C^{\alpha}(B_2)})$$

Proof. We have

$$(-\Delta)^s_{\Omega}(\chi_2 v) = f + g_v \quad \text{in } B_1,$$

where for $x \in B_1$

$$g_v(x) = c_{N,s} \int_{\Omega} (1 - \chi_2(y)) |x - y|^{-N - 2s} v(y) \, dy.$$

Letting $\widetilde{v} = \chi_2 v \in H^s(\mathbb{R}^N)$, we get

$$(-\Delta)^{s}_{\mathbb{R}^{N}}\widetilde{v} + \widetilde{v}V = f + g_{v} = c_{0} + (f - f(0)) + (g_{v} - g_{v}(0)) \quad \text{in } B_{1/2},$$

where $c_0 = f(0) + g_v(0), V(x) = c_{N,s} \int_{\mathbb{R}^N \setminus \Omega} |x - y|^{-N-2s} dy$. Clearly $\|V\|_{C^3(B_1)} \leq C(N,s)$. Moreover $\|g_v - g_v(0)\|_{C^1(B_{1/2})} \leq C(N,s)C_0$. Now as in Remark 3.2(*ii*), we can estimate

$$\begin{aligned} |c_0| &\leq C(N,s) \left(C_0 + \|\widetilde{v}V\|_{L^1(B_{1/4})} + \|g_v - g_v(0)\|_{L^1(B_{1/4})} + \|f - f(0)\|_{L^1(B_{1/4})} \right) \\ &\leq C(N,s) \left(C_0 + \|v\|_{L^2(B_2)} + \|f - f(0)\|_{L^{\infty}(B_2)} \right). \end{aligned}$$

By the interior regularity estimates in [11], we have that

$$\|\widetilde{v}\|_{C^{2s-\varepsilon}(B_{1/4})} \le C(N, s, \varepsilon)(C_0 + \|v\|_{L^2(B_2)} + \|f - f(0)\|_{L^{\infty}(B_2)}).$$

We can now apply the interior Schauder estimate in [18] and a standard bootstrap (only necessary for 2s < 1 because $\alpha < 1$) argument to get the desired estimate.

We can now prove the following.

Theorem 5.6. Let $s \in (0,1)$ and $\alpha_s \in (0,1-s)$ be given by Lemma 2.6, which satisfies $2s + \alpha_s > 1$. Let $\Omega \subset \mathbb{R}^N$ be a $C^{\max(1,2s+\alpha)}$ domain, with $0 \in \partial\Omega$, and let $f \in C^{\alpha}(\overline{\Omega} \cap B_1)$, with $\alpha < \alpha_s$ and with $\alpha + 2s \neq 1$. Let $u \in H^s(\Omega)$ be such that

$$D_{\Omega}(u,\varphi) = \int_{\Omega} f\varphi \, dx \qquad \forall \varphi \in C_c^{\infty}(B_1).$$

Then,

$$\|u\|_{C^{2s+\alpha}(\overline{\Omega}\cap B_{1/2})} \le C\left(\|u\|_{L^{2}(\Omega)} + \|f\|_{C^{\alpha}(\overline{\Omega}\cap B_{1})}\right).$$
(5.34)

with C depending only on N, s, α , and Ω .

Moreover, if $2s + \alpha > 1$ then $\partial_{\nu} u = 0$ on $\partial \Omega \cap B_1$, where ν is the exterior normal of $\partial \Omega$. *Proof.* We assume for simplicity that $||u||_{L^2(\Omega)} + ||f||_{C^{\alpha}(\overline{\Omega})} \leq 1$ and that Ω satisfies (5.1) and (5.2).

By Corollary 5.4 and the change of variable $w(y) = u(\psi(y))$ we have that, for all $\zeta \in \partial \Omega \cap B_{1/2}$, there exists $e = e_{\zeta} \in \mathbb{R}^{N-1} =: \partial \mathbb{R}^N_+$ with $|e_{\zeta}| \leq C$ and such that

$$\|u - u(\zeta) - \ell_{\alpha} e \cdot (\psi^{-1}(\cdot) - \zeta)\|_{L^{\infty}(B_r(\zeta) \cap \Omega)} \le Cr^{2s + \alpha}.$$
(5.35)

In addition, using also (5.2), we have $\partial_{\nu} u = 0$ on $\partial \Omega \cap B_{1/2}$ as soon as $2s + \alpha > 1$.

Let $x_0 \in \Omega \cap B_1$ and $\zeta \in \partial \Omega$ be such that $\operatorname{dist}(x_0, \partial \Omega) = |x_0 - \zeta|$. Put $\rho := \frac{|x_0 - \zeta|}{4}$ and $\overline{x} = \frac{x_0 - \zeta}{2} \in \Omega$ then $B_{2\rho}(\overline{x}) \subset \Omega$. We next, define

$$v_{\rho}(y) = (u(\overline{x} + \rho y) - u(\zeta)) - \ell_{\alpha} e \cdot (\psi^{-1}(\rho y + \overline{x}) - \zeta), \qquad f_{\rho}(y) = \rho^{2s} f(\overline{x} + \rho y)$$

and $\Omega_{\rho} = \frac{1}{\rho} (\Omega - \overline{x})$. Then

$$(-\Delta)^s_{\Omega_\rho} v_\rho = f_\rho + F_\rho =: G_\rho \qquad \text{in } B_2 \subset \Omega_\rho, \tag{5.36}$$

where $F_{\rho}(y) := \ell_{\alpha} \rho^{2s} [(-\Delta)^s_{\Omega}(e \cdot \psi^{-1})](\overline{x} + \rho y))$. Since $f \in C^{\alpha}(\Omega)$, using Lemma 4.1, we then get $[G_{\rho}]_{C^{\alpha}(B_1)} \leq C \rho^{2s+\alpha}$. By (5.35), we have that $\|v_{\rho}\|_{L^{\infty}(B_1)} \leq C \rho^{2s+\alpha}$ and

$$\|v_{\rho}(\cdot/\rho)\|_{L^{\infty}(B_{R\rho}\cap(\Omega-\overline{x}))} \le \|v_{\rho}(\cdot/\rho)\|_{L^{\infty}(B_{R\rho+2\rho}\cap(\Omega-\zeta))} \le C(R\rho)^{2s+\alpha} \quad \text{for all } R \ge 1.$$
(5.37)

Now applying Lemma 5.5 to (5.36), we obtain

$$\|v_{\rho}\|_{C^{2s+\alpha}(B_{1/4})} \le C(N,s,\alpha) \left(\int_{\Omega_{\rho} \setminus B_{1}} \frac{|v_{\rho}(y)| dy}{|y|^{N+2s+1}} + \|v_{\rho}\|_{L^{\infty}(B_{1})} + \|G_{\rho} - G_{\rho}(0)\|_{C^{\alpha}(B_{1})} \right).$$

Thanks to (5.37) and a change of variable we get

$$\int_{\Omega_{\rho} \setminus B_{1}} \frac{|v_{\rho}(y)| dy}{|y|^{N+2s+1}} = \rho^{2s+1} \int_{(\Omega-\overline{x}) \setminus B_{\rho}} \frac{|v_{\rho}(x/\rho)| dx}{|x|^{N+2s+1}}$$

$$\leq \sum_{i=0}^{\infty} (2^{i}\rho)^{-N-2s-1} \rho^{2s+1} \int_{(\Omega-\overline{x})\{|y|<2^{i+1}\rho\}} |v_{\rho}(x/\rho)|, dx$$

$$\leq C \sum_{i=0}^{\infty} (2^{i}\rho)^{-N-2s-1} \rho^{2s+1} (2^{i+1}\rho)^{N+2s+\alpha} \leq C \rho^{2s+\alpha}.$$

We conclude that $||v_{\rho}||_{C^{2s+\alpha}(B_{1/4})} \leq C\rho^{2s+\alpha}$. Therefore scaling and translating back, we get for $2s + \alpha < 1$ that

$$[u]_{C^{2s+\alpha}(B_{\rho}(\overline{x}))} \le C \tag{5.38}$$

and thus $[u]_{C^{2s+\alpha}(B_{1/2}\cap\Omega)} \leq C$ because \overline{x} is arbitrary.

If now $2s + \alpha > 1$, we obtain

$$[\nabla u - \nabla e \cdot (\psi^{-1}(\cdot) - \zeta)]_{C^{2s+\alpha-1}(B_{\rho}(\overline{x}))} \le C.$$

Hence, since $[\nabla e \cdot (\psi^{-1}(\cdot) - \zeta)]_{C^{2s+\alpha-1}(B_{\rho}(\overline{x}))} \leq C$, we get

$$[\nabla u]_{C^{2s+\alpha-1}(B_{\rho}(\overline{x}))} \le C.$$

This implies

$$[\nabla u]_{C^{2s+\alpha-1}(B_{1/2}\cap\overline{\Omega})} \le C.$$

Finally, we give the:

Proof of Theorem 1.1. The result follows from Theorem 5.6.

6. Regional Dirichlet problem

In this section, we consider $\phi \in C^{2,\beta}(\mathbb{R}^{N-1})$ satisfying $\|\phi\|_{C^{2,\beta}(\mathbb{R}^{N-1})} \leq \frac{1}{4}$ and $\phi(0) = |\nabla\phi(0)| = 0$. We define $\Phi(x', x_N) = (x', x_N + \phi(x'))$. We start with the following result.

Lemma 6.1. Let $\eta \in C_c^{2s-1}(\mathbb{R})$ with $\eta = 0$ on \mathbb{R}_- and $\ell(x') := a \cdot x' + b$. Let $f(x) = \eta(x_N)\ell(x')$. Then, for all $x \in \mathbb{R}^N_+$,

$$c_{N,s} \int_{\mathbb{R}^{N}_{+}} \frac{f(x) - f(y)}{|\Phi(x) - \Phi(y)|^{N+2s}} dy = \ell(x') T_{1}(x') (-\Delta)^{s}_{\mathbb{R}_{+}} \eta(x_{N}) + \ell(x') T_{3}(x') (-\Delta)^{s}_{\mathbb{R}_{+}} (x_{N}\eta) - T_{2}(x') \cdot a \int_{\mathbb{R}} \frac{(t - x_{N})(\eta(t) - \eta(x_{N}))}{|x_{N} - t|^{1+2s}} dt - T_{3}(x')\ell(x') \left(\eta(-\Delta)^{s}_{\mathbb{R}_{+}} x_{N} + x_{N}(-\Delta)^{s}_{\mathbb{R}_{+}} \eta\right) + h(x),$$

where $\|h\|_{C^{\min(\alpha,\beta)}(\overline{B_1^+})} \leq C \|\eta\|_{C^{2s-1}(\mathbb{R})}$ for all $\alpha < 2s - 1$, $T_1(0) = 1$, $T_2(0) = 0$ and $\|T_j\|_{C^{1,\beta}(B_1')} \leq C(N,s,\beta)$, for j = 1, 2. Moreover, $\|T_3\|_{C^{\beta}(B_1')} \leq C(N,s,\beta)$ and

$$T_3(0) = -D^2 \phi(0)[e_i, e_i] = -\Delta \phi(0).$$
(6.1)

Proof. We have

$$\begin{split} &\int_{\mathbb{R}^{N}_{+}} \frac{f(x) - f(y)}{|\Phi(x) - \Phi(y)|^{N+2s}} \, dy = \int_{\mathbb{R}^{N-1}} \int_{-x_{N}}^{\infty} \frac{f(x) - f(x+z)}{|\Phi(x) - \Phi(x+z)|^{N+2s}} \, dz \\ &= \int_{\mathbb{R}^{N-1}} \int_{-x_{N}}^{\infty} \frac{f(x) - f(x+z)}{|D\Phi(x')z|^{N+2s}} \, dz + \int_{\mathbb{R}^{N-1}} \int_{-x_{N}}^{\infty} \frac{f(x) - f(x+z)}{|z|^{N+2s}} B(x', |z'|, z'/|z'|, z/|z|) \, dz, \end{split}$$

where

$$B(x', r, \theta', \theta) = \frac{1}{|\int_0^1 D\Phi(x' + tr\theta')\theta dt|^{N+2s}} - \frac{1}{|D\Phi(x')\theta|^{N+2s}}.$$

Recall that for all $y \in \mathbb{R}^N$

$$D\Phi(x+r\theta')y = y + \left(\nabla\phi(x'+r\theta')\cdot y'\right)e_N.$$

We can thus write

$$B(x', r, \theta', \theta) = r\mu_1(x', \theta', \theta) + r\mathcal{O}(x', r, \theta', \theta), \qquad (6.2)$$

where

$$\mu_1(x',\theta',\theta) = \partial_r B(x',0,\theta',\theta) = -(N+2s) \frac{D^2 \Phi(x')[\theta',\theta-(\theta\cdot e_N)e_N] \cdot D\Phi(x')\theta}{|D\Phi(x')\theta|^{N+2s+2}},$$

and

$$\mathcal{O}(x', r, \theta', \theta) = \partial_r B(x', r, \theta', \theta) - \partial_r B(x', 0, \theta', \theta).$$

We then have $\mu_1, \mathcal{O} \in C_x^\beta(B_2), \ \mu_1(x', -\theta', -\theta) = -\mu_1(x', \theta', \theta), \ |\mathcal{O}(x', r, \theta', \theta)| \leq C \min(1, r^\beta)$ and

$$|\mathcal{O}(x',r,\theta',\theta)| \le C\min(1,r^{\beta}), \qquad |\mathcal{O}(x',r,\theta',\theta) - \mathcal{O}(\overline{x}',r,\theta',\theta)| \le C\min(|x'-\overline{x}'|,r^{\beta}).$$
(6.3)

We thus get from (6.2),

$$c_{N,s} \int_{\mathbb{R}^{N}_{+}} \frac{f(x) - f(y)}{|\Phi(x) - \Phi(y)|^{N+2s}} \, dy = c_{N,s} \int_{\mathbb{R}^{N-1}} \int_{-x_{N}}^{\infty} \frac{f(x) - f(x+z)}{|D\Phi(x')z|^{N+2s}} \, dz$$

+ $c_{N,s} \int_{\mathbb{R}^{N-1}} \int_{-x_{N}}^{\infty} \frac{f(x) - f(x+z))}{|z|^{N+2s}} |z'| \mu_{1}(x', z'/|z'|, z/|z|) \, dz$
+ $c_{N,s} \int_{\mathbb{R}^{N-1}} \int_{-x_{N}}^{\infty} \frac{f(x) - f(x+z))}{|z|^{N+2s}} |z'| \mathcal{O}(x', |z'|, z/|z|) \, dz$
= $I_{1}(x) + I_{2}(x) + I_{3}(x).$ (6.4)

Letting $T_1(x') := \frac{1}{c_{1,s}} \int_{\mathbb{R}^{N-1}} \frac{c_{N,s}}{|D\Phi(x')(z',1)|^{N+2s}} dz'$ and $T_2(x') \cdot e_i = c_{N,s} \int_{\mathbb{R}^{N-1}} \frac{z_i}{|D\Phi(x')(z',1)|^{N+2s}} dz'$, we then get

$$\begin{split} I_1(x) &= \ell(x') \int_{\mathbb{R}^{N-1}} \int_{-x_N}^\infty \frac{\eta(x_N) - \eta(x_N + z_N)}{|D\Phi(x')z|^{N+2s}} dz - \int_{\mathbb{R}^{N-1}} \int_{-x_N}^\infty \frac{a \cdot z' \eta(x_N + z_N)}{|D\Phi(x')z|^{N+2s}} dz \\ &= T_1(x')\ell(x')c_{1,s} \int_{-x_N}^\infty \frac{\eta(x_N) - \eta(x_N + z_N)}{|z_N|^{1+2s}} dz_N - T_2(x') \cdot a \int_{-x_N}^\infty z_N \frac{\eta(x_N + z_N)}{|z_N|^{1+2s}} dz_N \\ &= T_1(x')\ell(x')(-\Delta)_{\mathbb{R}_+}^s \eta(x_N) - T_2(x') \cdot a \int_{\mathbb{R}^+} \frac{(t - x_N)\eta(t)}{|x_N - t|^{1+2s}} dt. \end{split}$$

Since $\eta = 0$ on \mathbb{R}_{-} , then

$$\int_{\mathbb{R}^+} \frac{(t-x_N)\eta(t)}{|x_N-t|^{1+2s}} \, dt = \int_{\mathbb{R}} \frac{(t-x_N)\eta(t)}{|x_N-t|^{1+2s}} \, dt = \int_{\mathbb{R}} \frac{(t-x_N)(\eta(t)-\eta(x_N))}{|x_N-t|^{1+2s}} \, dt$$

We thus conclude that

$$I_1(x) = T_1(x')\ell(x')(-\Delta)^s_{\mathbb{R}_+}\eta(x_N) - T_2(x') \cdot a \int_{\mathbb{R}} \frac{(t-x_N)(\eta(t)-\eta(x_N))}{|x_N-t|^{1+2s}} dt.$$
(6.5)

Since $D\Phi(0) = id$, we see that $T_1(0) = \frac{1}{c_{1,s}} \int_{\mathbb{R}^{N-1}} \frac{c_{N,s}}{(1+|z'|^2)^{\frac{N+2s}{2}}} dz' = 1$ and by oddness we have $T_2(0) = 0.$

We consider next I_2 . Similarly, letting $T_3(x') = \frac{c_{N,s}}{c_{1,s}} \int_{\mathbb{R}^{N-1}} \frac{|z'|}{|(z',1)|^{N+2s}} \mu_1(x', \frac{z'}{|z'|}, \frac{(z',1)}{|(z',1)|}) dz'$, we obtain

$$I_{2}(x) = T_{3}(x')\ell(x')c_{1,s} \int_{-x_{N}}^{\infty} z_{N} \frac{\eta(x_{N}) - \eta(x_{N} + z_{N})}{|z_{N}|^{1+2s}} dz_{N} + h_{1}(x)$$

$$= T_{3}(x')\ell(x')c_{1,s} \int_{\mathbb{R}^{+}} \frac{(t - x_{N})(\eta(x_{N}) - \eta(t))}{|x_{N} - t|^{1+2s}} dt + h_{1}(x)$$

$$= T_{3}(x')\ell(x') \left((-\Delta)_{\mathbb{R}^{+}}^{s}(x_{N}\eta) - \eta(-\Delta)_{\mathbb{R}^{+}}^{s}x_{N} - x_{N}(-\Delta)_{\mathbb{R}^{+}}^{s}\eta \right) + h_{1}(x), \qquad (6.6)$$

where

$$h_1(x) := c_{N,s} \int_{\mathbb{R}^{N-1}} \int_{-x_N}^{\infty} \frac{|z'|a \cdot z'\eta(x_N + z_N)}{|z|^{N+2s}} \mu_1(x', z'/|z'|, z/|z|) dz$$

= $c_{N,s} \int_{\mathbb{R}^{N-1}} \frac{|z'|a \cdot z'}{|(z', 1)|^{N+2s}} \mu_1(x', \frac{z'}{|z'|}, \frac{(z', 1)}{|(z', 1)|}) dz' \int_{-x_N}^{\infty} \frac{z_N \eta(x_N + z_N)}{|z_N|^{2s}} dz_N.$

Since 2s > 1, we can easily see that

$$\|h_1\|_{C^{\min(\beta,2s-1)}(\overline{B_1^+})} \le C|a| \|\eta\|_{C^{2s-1}(\mathbb{R})}.$$
(6.7)

Using that $D\Phi(0) = Id$, a change of variable and integration by part, we get

$$c_{1,s}T_{3}(0) = -(N+2s)c_{N,s} \int_{\mathbb{R}^{N-1}} \frac{|z'|^{2}D^{2}\phi(0)[z'/|z'|,z'/|z'|]}{|(z',1)|^{N+2s+2}} dz'$$

$$= -(N+2s)c_{N,s} \int_{0}^{\infty} \frac{r^{N}}{(r^{2}+1)^{\frac{N+2s+2}{2}}} dr \sum_{i=1}^{N-1} D^{2}\phi(0)[e_{i},e_{i}] \int_{S^{N-2}} \theta_{1}^{2}d\sigma(\theta)$$

$$= -(N+2s)\frac{c_{N,s}|S^{N-2}|}{N-1} \int_{0}^{\infty} \frac{r^{N}}{(r^{2}+1)^{\frac{N+2s+2}{2}}} dr \sum_{i=1}^{N-1} D^{2}\phi(0)[e_{i},e_{i}]$$

$$= -c_{N,s}|S^{N-2}| \int_{0}^{\infty} \frac{r^{N-2}}{(r^{2}+1)^{\frac{N+2s}{2}}} dr \sum_{i=1}^{N-1} D^{2}\phi(0)[e_{i},e_{i}] = -c_{1,s} \sum_{i=1}^{N-1} D^{2}\phi(0)[e_{i},e_{i}]. \quad (6.8)$$

Here, we used that $c_{N,s} \int_{\mathbb{R}^{N-1}} \frac{1}{(1+|z'|^2)^{\frac{N+2s}{2}}} dz' = c_{1,s}$. Finally, using (6.3), we see that

$$||I_3||_{C^{\min(\alpha,\beta)}(\overline{B_1^+})} \le C ||f||_{C^{2s-1}(\mathbb{R})}$$
 for all $\alpha \in (0, 2s-1).$

From this together with (6.8), (6.7), (6.6), (6.5) and (6.4), we get the result. **Lemma 6.2.** Let $\overline{\eta} \in C_c^{\infty}(-3,3)$ with $\overline{\eta} = 1$ on (-2,2) and $\ell(x') := a \cdot x' + b$. Define $P(x) = \overline{\eta}(x_N)(x_N)_+^{2s-1}\ell(x') - bT_3(0)\overline{\eta}(x_N)(x_N)_+^{2s}$,

where $T_3(0)$ is given by Lemma 6.1. Then, for all $x \in B_1^+$,

$$c_{N,s} \int_{\mathbb{R}^N_+} \frac{P(x) - P(y)}{|\Phi(x) - \Phi(y)|^{N+2s}} \, dy = g_1(x) \log(x_N) + g_2(x),$$

where $||g_i||_{C^{\min(\beta,\alpha)}(\overline{B_1^+})} \leq C(N, s, \beta, \alpha, \overline{\eta})(|a| + |b|)$ for all $\alpha < 2s - 1$ and $g_1(0) = 0$. Proof. Recall from Lemma 6.1 that

$$c_{N,s} \int_{\mathbb{R}^N_+} \frac{P(x) - P(y)}{|\Phi(x) - \Phi(y)|^{N+2s}} \, dy = L_1(x) + L_2(x) + h(x), \tag{6.9}$$

where $h \in C^{\beta}(\overline{B_1^+})$,

$$L_1(x) = \ell(x')T_1(x')(-\Delta)^s_{\mathbb{R}_+}(x_N^{2s-1}\overline{\eta}) + \ell(x')T_3(x')(-\Delta)^s_{\mathbb{R}_+}(x_N^{2s}\overline{\eta}) - bT_3(0)T_1(x')(-\Delta)^s_{\mathbb{R}_+}(x_N^{2s}\overline{\eta}) - bT_3(0)T_3(x')(-\Delta)^s_{\mathbb{R}_+}(x_N^{2s+1}\overline{\eta})$$

and

$$L_{2}(x) = -T_{2}(x') \cdot a \int_{\mathbb{R}} \frac{(t - x_{N})(t_{+}^{2s-1}\overline{\eta}(t) - (x_{N})_{+}^{2s-1}\overline{\eta}(x_{N}))}{|x_{N} - t|^{1+2s}} dt$$

$$-T_{3}(x')\ell(x') \left(\overline{\eta}x_{N}^{2s-1}(-\Delta)_{\mathbb{R}_{+}}^{s}x_{N} + x_{N}(-\Delta)_{\mathbb{R}_{+}}^{s}(x_{N}^{2s-1}\overline{\eta})\right)$$

$$-bT_{3}(0)T_{2}(x') \cdot a \int_{\mathbb{R}} \frac{(t - x_{N})(t_{+}^{2s}\overline{\eta}(t) - (x_{N})_{+}^{2s}\overline{\eta}(x_{N}))}{|x_{N} - t|^{1+2s}} dt$$

$$-bT_{3}(0)T_{3}(x') \left(\overline{\eta}x_{N}^{2s}(-\Delta)_{\mathbb{R}_{+}}^{s}x_{N} + x_{N}(-\Delta)_{\mathbb{R}_{+}}^{s}(x_{N}^{2s}\overline{\eta})\right).$$
(6.10)

We now write

$$L_{1}(x) = -bT_{3}(0)(T_{1}(x') - 1)(-\Delta)^{s}_{\mathbb{R}_{+}}(x_{N}^{2s}\overline{\eta}) + b(T_{3}(x') - T_{3}(0))(-\Delta)^{s}_{\mathbb{R}_{+}}(x_{N}^{2s}\overline{\eta}) + a \cdot x'T_{3}(x')(-\Delta)^{s}_{\mathbb{R}_{+}}(x_{N}^{2s}\overline{\eta}) - bT_{3}(x')(-\Delta)^{s}_{\mathbb{R}_{+}}(x_{N}^{2s+1}\overline{\eta}) + \ell(x')T_{1}(x')(-\Delta)^{s}_{\mathbb{R}_{+}}(x_{N}^{2s-1}\overline{\eta}).$$

Moreover since $x_N \mapsto x_N^{2s+1}\overline{\eta}(x_N)$ has zero derivative at $x_N = 0$, we get $(-\Delta)_{\mathbb{R}_+}^s(x_N^{2s+1}\overline{\eta}) \in C^{\beta}([0,1])$ by Lemma 4.1. In addition, since $(-\Delta)_{\mathbb{R}_+}^s x^{2s-1} = 0$, we see that $(-\Delta)_{\mathbb{R}_+}^s(x^{2s-1})\overline{\eta} \in C^{\beta}([0,1])$. Therefore, recalling that $T_1(0) = 1$, we thus obtain

$$L_1(x) = g_1(x)H(x_N) + g_2(x), (6.11)$$

for some functions g_i as in the statement of the lemma and $H(x_N) = (-\Delta)^s_{\mathbb{R}_+}(x_N^{2s}\overline{\eta})(x_N)$. To estimate L_2 , we first observe that

$$c_{1,s} \int_{\mathbb{R}} \frac{(t-x_N)(t_+^{2s-1}\overline{\eta}(t)-(x_N)_+^{2s-1}\overline{\eta}(x_N))}{|x_N-t|^{1+2s}} dt$$

= $-(-\Delta)^s (x_N(x_N)_+^{2s-1}\overline{\eta}) + x_N(-\Delta)^s ((x_N)_+^{2s-1}\overline{\eta})$
= $-(-\Delta)^s_{\mathbb{R}_+} ((x_N)^{2s}\overline{\eta}) - a_s + x_N c_{N,s} \int_{\mathbb{R}_+} \frac{\overline{\eta}(x_N) - \overline{\eta}(t) t^{2s-1}}{|x_N-t|^{1+2s}} dt,$

where we used that $(-\Delta)_{\mathbb{R}_+}^s x_N^{2s-1} = 0$ and $(-\Delta)^s x_N = 0$ and the formula $(-\Delta)^s (\mathbb{1}_{\mathbb{R}_+} u)(x) = (-\Delta)_{\mathbb{R}_+}^s u + a_s x^{-2s} u$ in \mathbb{R}_+ . We have that

$$c_{1,s} \int_{\mathbb{R}} \frac{(t-x_N)(t_+^{2s}\overline{\eta}(t) - (x_N)_+^{2s}\overline{\eta}(x_N))}{|x_N - t|^{1+2s}} dt = -(-\Delta)^s ((x_N)_+^{2s+1}\overline{\eta}) + x_N (-\Delta)^s (x_N)_+^{2s}\overline{\eta}$$
$$= -(-\Delta)^s ((x_N)_+^{2s+1}\overline{\eta}) + x_N (-\Delta)_{\mathbb{R}_+}^s (x_N^{2s}\overline{\eta}) + a_s x_N.$$
(6.12)

Moreover $(-\Delta)_{\mathbb{R}_+}^s x_N = \kappa_s x_N^{1-2s}$, for some $\kappa_s > 0$. Next, using that $(-\Delta)_{\mathbb{R}_+}^s x^{2s-1} = 0$, we see that $(-\Delta)_{\mathbb{R}_+}^s (x^{2s-1}\overline{\eta}) \in C^{\beta}([0,1])$. Since $T_2(0) = 0$, we thus get from (6.12) and (6.13),

$$L_2(x) = g_1(x)H(x_N) + g_2(x), (6.13)$$

for some functions g_i as in the statement of the lemma and $H(x_N) = (-\Delta)^s_{\mathbb{R}_+}(x_N^{2s}\overline{\eta})(x_N)$. To conclude, we observe that

$$H'(x_N) = 2s(-\Delta)^s (x_N^{2s-1}\overline{\eta})(x_N) + (-\Delta)^s (x_N^{2s}\overline{\eta}')(x_N) - a_s\overline{\eta}'(x_N)$$

= $2s(-\Delta)^s_{\mathbb{R}_+} (x_N^{2s-1}\overline{\eta})(x_N) + a_s x_N^{-1}\overline{\eta}(x_N) + (-\Delta)^s (x_N^{2s}\overline{\eta}')(x_N) - a_s\overline{\eta}'(x_N).$

As a consequence, $x_N \mapsto H'(x_N) - a_s x_N^{-1} \in C^{\infty}([0,1])$. It follows that for $x_N \in (0,1)$, we have $H(x_N) = -\int_{x_N}^{1/2} H'(t) dt + H(1/2) = a_s \log x_N + \tilde{h}(x_N)$, with $\tilde{h} \in C^{\infty}([0,1])$. In view of this, (6.13), (6.11) and (6.9), we get the result.

6.1. Expansion near the boundary. In the remaining of this section, we define \mathcal{B}_{Φ} : $H^{s}(\mathbb{R}^{N}_{+}) \times H^{s}(\mathbb{R}^{N}_{+}) \to \mathbb{R}$ by

$$\mathcal{B}_{\Phi}(u,\varphi) := \frac{c_{N,s}}{2} \int_{\mathbb{R}^N_+ \times \mathbb{R}^N_+} \frac{(u(x) - u(y))(\varphi(x) - \varphi(y))}{|\Phi(x) - \Phi(y)|^{N+2s}} \, dx dy.$$

We start recalling the following result from [12].

Lemma 6.3 ([12]). Let 2s > 1. Let $f \in L^p(B_2^+)$ and $u \in H_0^s(\mathbb{R}^N_+)$ by a solution to

$$\mathcal{B}_{\Phi}(u,\varphi) = \int_{\mathbb{R}^N_+} f\varphi \, dx \qquad \text{for all } \varphi \in C^{\infty}_c(B^+_2).$$

Then for all p > N, there exists C = C(N, s, p) > 0 such that

$$\|u/x_N^{2s-1}\|_{C^{1-N/p}(B_1^+)} \le C\left(\|u\|_{L^2(\mathbb{R}^N_+)} + \|f\|_{L^p(B_2^+)}\right).$$

We state the next result which is crucial for the proof of Theorem 6.7.

Proposition 6.4. Let $s \in (\frac{1}{2}, 1)$, $\alpha_s \in (0, 1 - s)$ be given by Lemma 2.6 and $\beta \in (0, \alpha_s)$. Let $\alpha \in (0, \min(\beta, 2s - 1))$, with $2s + \alpha \neq 1$, $f \in C^{\alpha}(\mathbb{R}^N_+)$ and $w \in H^s_0(\mathbb{R}^N_+)$ such that

$$\mathcal{B}_{\Phi}(w,\varphi) = \int_{\mathbb{R}^N_+} f\varphi \, dx \qquad \text{for all } \varphi \in C^{\infty}_c(B^+_2)$$

and

$$\|w/x_N^{2s-1}\|_{L^{\infty}(\mathbb{R}^N_+)} + \|f\|_{C^{\alpha}(\mathbb{R}^N_+)} \le 1.$$

Let $\overline{\eta} \in C_c^{\infty}(-3,3)$ with $\overline{\eta} = 1$ on (-2,2). Then, there exists $C = C(N, s, \alpha, \beta, \overline{\eta})$ such that

$$\sup_{r>0} r^{-1-\alpha} \left\| \frac{w}{x_N^{2s-1}} - \left(a_{r,w} \cdot x' + b_{r,w} (1 - T_3(0)x_N) \right) \overline{\eta}(x_N) \right\|_{L^{\infty}(B_r^+)} \le C, \tag{6.14}$$

where $a_{r,w} := A_{r,w/x_N^{2s-1}}$ is given by (5.7),

$$b_{r,w} = \frac{1}{\int_{B_r^+} (1 - T_3(0)y_N)^2 dy} \int_{B_r^+} \frac{w}{x_N^{2s-1}}(y)(1 - T_3(0)y_N) dy,$$

and $T_3(0) = -\Delta \phi(0)$ is given by (6.1).

Proof. Suppose on the contrary that (6.14) does not hold. Then for all integer $n \ge 2$, there exist $\phi_n \in C^{2,\beta}(\mathbb{R}^{N-1}), f_n \in C^{\alpha}(\mathbb{R}^N_+)$ and $w_n \in H^s_0(\mathbb{R}^N_+)$, such that

$$\|\phi_n\|_{C^{2,\beta}(\mathbb{R}^{N-1})} \le \frac{1}{4}, \qquad \phi_n(0) = |\nabla\phi_n(0)| = 0, \tag{6.15}$$

$$\|w_n/x_N^{2s-1}\|_{L^{\infty}(\mathbb{R}^N_+)} + \|f_n\|_{C^{\alpha}(\mathbb{R}^N_+)} \le 1$$
(6.16)

with

$$\mathcal{B}_{\Phi}(w_n,\varphi) = \int_{\mathbb{R}^N_+} f_n \varphi \, dx \qquad \text{for all } \varphi \in C^{\infty}_c(B^+_2).$$
(6.17)

and

$$\sup_{r>0} r^{-1-\alpha} \left\| \frac{w_n}{x_N^{2s-1}} - \left(a_{r,w_n} \cdot x' + b_{r,w_n} (1 - T_3^n(0)x_N) \right) \overline{\eta}(x_N) \right\|_{L^{\infty}(B_r^+)} > n, \tag{6.18}$$

where (recalling (6.1))

$$T_3^n(0) := -\Delta \phi_n(0). \tag{6.19}$$

Define

$$Q_r(x) = a_{r,w_n} \cdot x' + b_{r,w_n} (1 - T_3^n(0)x_N).$$

By (6.16), we can define the monotone nonincrease sequence of function θ_n by

$$\theta_n(\rho) := \sup_{r > \rho} r^{-1-\alpha} \left\| \frac{w_n}{x_N^{2s-1}} - Q_r(x)\overline{\eta}(x_N) \right\|_{L^{\infty}(B_r^+)}.$$
(6.20)

Obviously, for $n \ge 2$, by (6.18), there exists $\rho_n > 0$ such that

$$\theta_n(\rho_n) > n/2 \ge 1. \tag{6.21}$$

Hence, provided $n \ge 2$, by definition of $\theta_n(\rho_n)$ as a supremum and the monotonicity of θ_n , there exists $r_n \ge \rho_n$ such that

$$\theta_{n}(r_{n}) \geq r_{n}^{-1-\alpha} \left\| \frac{w_{n}}{x_{N}^{2s-1}} - Q_{r_{n}}(x)\overline{\eta}(x_{N}) \right\|_{L^{\infty}(B_{r_{n}}^{+})}$$
$$\geq \theta_{n}(\rho_{n}) - 1/2 \geq (1 - 1/2)\theta_{n}(\rho_{n}) \geq \frac{1}{2}\theta_{n}(r_{n}), \tag{6.22}$$

Also by (6.21) and the above estimate, we have that $\theta_n(r_n) \geq \frac{n}{2} - \frac{1}{2} \to \infty$ as $n \to \infty$. We now define $v_n \in H_0^s(\mathbb{R}^N_+)$ by

$$v_n(x) := \frac{1}{r_n^{2s+\alpha} \theta_n(r_n)} \left(w_n(r_n x) - Q_{r_n}(r_n x) \overline{\eta}(r_n x_N) (r_n x_N)^{2s-1} \right).$$

In view of (6.22), it clearly satisfies

$$\|v_n/x_N^{2s-1}\|_{L^{\infty}(B_1^+)} \ge \frac{1}{2}.$$
(6.23)

Moreover, by construction (recalling (5.9)),

$$\int_{B_1^+} \frac{v_n}{x_N^{2s-1}}(y)(1 - T_3^n(0)y_N)dy = 0, \qquad \int_{B_1^+} \frac{v_n}{x_N^{2s-1}}(y)y_i\,dy = 0 \qquad \text{for all } i = 1, \dots, N-1.$$
(6.24)

Note that by Lemma 6.3, (6.17) and (6.16),

$$|a_{r,w_n}| \le c_0 r^{-\varrho}, \quad \text{for all } \varrho \in (0, \beta - \alpha).$$
 (6.25)

By (6.15) and (6.19), for $M \le 1$, we have

$$\|Q_{2r} - Q_r\|_{L^2(B_M)}^2 \asymp |B_M| \left(M^2 |a_{2r,w_n} - a_{r,w_n}|^2 + |b_{2r,w_n} - b_{r,w_n}|^2 \right).$$
(6.26)

Therefore by (6.20), for $2r \leq 1$,

$$\begin{aligned} 2r|a_{2r,w_n} - a_{r,w_n}| + |b_{2r,w_n} - b_{r,w_n}| \\ &\leq C \left\| w_n / x_N^{2s-1} - Q_{2r} \right\|_{L^{\infty}(B_{2r}^+)} + C \left\| w_n / x_N^{2s-1} - Q_r \right\|_{L^{\infty}(B_r^+)} \\ &\leq C \left(\theta_n (2r) (2r)^{1+\alpha} + \theta_n(r) r^{1+\alpha} \right) \leq C \theta_n(r) r^{1+\alpha}. \end{aligned}$$

Using the monotonicity of θ_n , we see that for all $m \in \mathbb{N}$ such that $2^m r \leq 1$, we get

$$\begin{aligned} \|Q_{2^{m}r} - Q_{r}\|_{L^{\infty}(B_{2^{m}r})} &\leq C \|Q_{2^{m}r} - Q_{r}\|_{L^{2}(B_{2^{m}r})} \leq C \sum_{i=1}^{m} \|Q_{2^{i}r} - Q_{2^{i-1}}\|_{L^{2}(B_{2^{i}r})} \\ &\leq C \sum_{i=1}^{m} 2^{i}r|a_{2^{i}r,w_{n}} - a_{2^{i-1}r,w_{n}}| + |b_{2^{i}r,w_{n}} - b_{2^{i-1}r,w_{n}}| \leq C\theta_{n}(r) \sum_{i=1}^{m} 2^{i(1+\beta)}r^{1+\alpha} \\ &\leq C\theta_{n}(r)(2^{m}r)^{1+\alpha}. \end{aligned}$$

Hence for all $M \ge 1$ such that $Mr \le 1$, we get

$$||Q_{Mr} - Q_r||_{L^{\infty}(B_{Mr})} \le \theta_n(r)(Mr)^{1+\alpha}.$$

From this, (6.20) and the monotonicity of θ_n , we then have, for $Mr \leq 1$,

$$\begin{split} \|v_n/x_N^{2s-1}\|_{L^{\infty}(B_M^+)} &= \frac{1}{r_n^{1+\alpha}\theta_n(r_n)} \|w_n/x_N^{2s-1} - Q_{r_n}\|_{L^{\infty}(B_{Mr_n}^+)} \\ &+ \frac{1}{r_n^{1+\alpha}\theta_n(r_n)} \|w_n/x_N^{2s-1} - Q_{Mr_n}\|_{L^{\infty}(B_{Mr_n}^+)} + \frac{1}{r_n^{1+\alpha}\theta_n(r_n)} \|Q_{Mr_n} - Q_{r_n}\|_{L^{\infty}(B_{Mr_n})} \\ &< CM^{1+\alpha}. \end{split}$$

As a consequence,

$$\|v_n\|_{L^{\infty}(B_M^+)} \le CM^{2s+\alpha} \qquad \text{whenever } Mr_n \le 1.$$
(6.27)

Now for $Mr_n > 1$ and using (6.15), we have (recall that $2s + \alpha > 1$)

$$\|v_n\|_{L^{\infty}(B_M^+)} \le \frac{C}{r_n^{2s+\alpha}\theta_n(r_n)} (1 + r_n^{-\varrho} + Mr_n + |T_3^n(0)|Mr_n) \le CM^{2s+\alpha+\varrho}.$$

In view of this and (6.27) we deduce that

$$||v_n||_{L^{\infty}(B_M^+)} \le CM^{2s+\beta}$$
 for all $M \ge 1$. (6.28)

Letting $\widetilde{\Phi}_n(x) = \frac{1}{r_n} \Phi(r_n x)$, we thus get

$$\mathcal{B}_{\tilde{\Phi}_n}(\chi_R v_n, \varphi) = \int_{\mathbb{R}^N_+} G_R^n \varphi \, dx \qquad \text{for all } \varphi \in C_c^\infty(B^+_{\frac{1}{r_n}}), \tag{6.29}$$

where

$$G_{R}^{n}(x) := \frac{1}{r_{n}^{\beta}\theta_{n}(r_{n})} f_{n}(r_{n}x) + c_{N,s} \int_{\mathbb{R}^{N}_{+}} \frac{(\chi_{R}(x) - \chi_{R}(y))v_{n}(y)}{|\tilde{\Phi}_{n}(x) - \tilde{\Phi}_{n}(y)|^{N+2s}} \, dy - \frac{1}{r_{n}^{\beta}\theta_{n}(r_{n})} (\mathcal{L}_{\Phi_{n}}P_{n})(r_{n}x),$$

$$P_{n}(x) = \overline{\eta}(x_{N}) \left(a_{r_{n},w_{n}} \cdot x' + b_{r_{n},w_{n}}(1 - T_{3}^{n}(0)x_{N})\right) x_{N}^{2s-1}$$
and $(\mathcal{L}_{\Phi_{n}}P_{n})(x) = c_{N,s} \int_{\mathbb{R}^{N}_{+}} \frac{P_{n}(x) - P_{n}(y)}{|\Phi_{n}(x) - \Phi_{n}(y)|^{N+2s}} \, dy.$ By Lemma 6.2,
 $(\mathcal{L}_{\Phi_{n}}P_{n})(r_{n}x) = g_{1,n}(r_{n}x) \log(r_{n}x_{N}) + g_{2,n}(r_{n}x)$ (6.30)

and thus using (6.25) we see that

$$\frac{1}{r_n^{\beta}\theta_n(r_n)} |(\mathcal{L}_{\Phi_n} P_n)(r_n x) - (g_{2,n}(r_n x) - g_{2,n}(0))| \le \frac{Cr_n^{\beta}|x|^{\beta}}{r_n^{\alpha}\theta_n(r_n)} \left(1 + |\log x_N| + |\log r_n|\right) (r_n^{-\varrho} + 1) \le \frac{C|x|^{\beta}}{\theta_n(r_n)} \left(1 + |\log x_N|\right).$$
(6.31)

Since $|\nabla \phi_n| \leq \frac{1}{4}$ and $\widetilde{\Phi}_n(x) = x + (0, r_n^{-1}\phi_n(r_n x))$, we obtain

$$2|x-y| \ge |\widetilde{\Phi}_n(x) - \widetilde{\Phi}_n(y)| \ge \frac{1}{2}|x-y|, \qquad |D\widetilde{\Phi}_n(x)| \le 2 \qquad \text{for all } x, y \in \mathbb{R}^N.$$
(6.32)

Define, for $x \in B_{R/2}$,

$$F_R^n(x) := c_{N,s} \int_{\mathbb{R}^N_+} \frac{(\chi_R(x) - \chi_R(y))v_n(y)}{|\tilde{\Phi}_n(x) - \tilde{\Phi}_n(y)|^{N+2s}} dy$$
$$= c_{N,s} \int_{\mathbb{R}^N_+ \cap \{|y| \ge R\}} \frac{(1 - \chi_R(y))v_n(y)}{|\tilde{\Phi}_n(x) - \tilde{\Phi}_n(y)|^{N+2s}} dy.$$

By the mean value theorem and the fact that $\tilde{\Phi}_n(0) = 0$, we have $F_R^n(x) = F_R^n(0) + \mathcal{R}_R^n(x)$, where

$$\mathcal{R}_R^n(x) := -(N+2s)c_{N,s} \int_0^1 \int_{\mathbb{R}^N_+} \frac{(\widetilde{\Phi}_n(x) - \widetilde{\Phi}_n(y)) \cdot \widetilde{\Phi}_n(x)}{|\widetilde{\Phi}_n(x) - \widetilde{\Phi}_n(y)|} \frac{(1 - \chi_R(y))v_n(y)}{|t\widetilde{\Phi}_n(x) - \widetilde{\Phi}_n(y)|^{N+2s+1}} \, dy \, dt.$$

Note that by (6.32), $|t\tilde{\Phi}_n(x) - \tilde{\Phi}_n(y)| \ge \frac{1}{4}|y|$ whenever $|y| \ge R$ and $|x| \le \frac{R}{4}$. Therefore, by (5.20) and (6.32), there exists C > 0 such that

$$|\nabla \mathcal{R}_R^n(x)| \le C R^{\beta - 1} \qquad \text{for all } R > 1, \ |x| \le \frac{R}{4} \text{ and } n \ge 2$$
(6.33)

and by construction we have $\mathcal{R}_R^n(0) = 0$. We can thus write

$$G_R^n(x) = H_R^n(x) + c_R^n + \mathcal{R}_R^n(x),$$
 (6.34)

with (recalling (6.30))

$$H_R^n(x) = \frac{1}{r_n^\alpha \theta_n(r_n)} \left\{ f_n(r_n x) - f_n(0) - (\mathcal{L}_{\Phi_n} P_n)(r_n x) + (g_{2,n}(r_n x) - g_{2,n}(0)) \right\}$$

and

$$c_R^n = \frac{1}{r_n^{\alpha} \theta_n(r_n)} \left(f_n(0) - g_{2,n}(0) \right) + F_R^n(0).$$

Recalling that $||f_n||_{C^{\alpha}(\mathbb{R}^N_{+})} \leq 1$, we then get from (6.31) that

$$|H_R^n(x)| \le \frac{C(R)}{\theta_n(r_n)} (1 + |\log x_N|) \qquad \text{for all } x \in B_{R/2}^+,$$

so that

$$\|H_R^n\|_{L^p(B_{R/2})} \le \frac{C(R)}{\theta_n(r_n)} \qquad \text{for all } p > 1.$$
(6.35)

Let us now show that $(c_R^n)_{n \in \mathbb{N}}$ is bounded. For this, we pick $\phi \in C_c^{\infty}(B_{1/4}^+)$ with $\int_{\mathbb{R}^N} \phi \, dx = 1$ and let $R \geq 2$. Then multiply (6.29) by ϕ and integrate over \mathbb{R}^N_+ to get

$$|c_{R}^{n}| \leq \int_{\mathbb{R}^{N}} |\chi_{R}(x)v_{n}(x)| |\mathcal{L}_{\widetilde{\Phi}_{n}}\phi(x)| \, dx + \int_{B_{R/2}} |H_{R}^{n}(x)\phi(x)| \, dx + \int_{B_{R/2}} |\mathcal{R}_{R}^{n}(x)\phi(x)| \, dx \leq C(R).$$
(6.36)

In view of this, (6.29), (6.34), (6.35) and (6.33), we can apply Lemma 6.3, to deduce that v_n is bounded in $C_{loc}^{2s-1}(\overline{\mathbb{R}^N_+}) \cap H^s_{loc}(\overline{\mathbb{R}^N_+})$ and converges to some $v \in C_{loc}^{2s-1}(\overline{\mathbb{R}^N_+}) \cap H^s_{loc}(\overline{\mathbb{R}^N_+})$. Moreover v_n/x_N^{2s-1} converges to v/x_N^{2s-1} in $C_{loc}^{1-\varepsilon}(\overline{\mathbb{R}^N_+})$, for all $\varepsilon \in (0, 1)$. In addition passing to the limit in (6.23), (6.28) and (6.24), we get

$$|v(x)| \le C|x|^{2s+\beta}$$
 for all $x \in \mathbb{R}^N$, (6.37)

$$\|v/x_N^{2s-1}\|_{L^{\infty}(B_1^+)} \ge \frac{1}{2}, \qquad \int_{B_1^+} \frac{v}{x_N^{2s-1}} (1 - T_3^{\infty}(0)x_N) \, dx = \int_{B_1^+} \frac{v}{x_N^{2s-1}} x_i \, dx = 0, \qquad (6.38)$$

for $i = 1, \ldots, N-1$, where $T_3^{\infty}(0)$ is the limit $T_3^n(0)$ as $n \to \infty$ and satisfies $|T_3^{\infty}(0)| \le 1/4$. Passing to the limit in (6.29), (6.33) and using (6.36), we get, for all $\phi \in C_c^{\infty}(B_{\frac{R}{4}})$,

$$\int_{\mathbb{R}^{2N}_+} \frac{((\eta_R v)(x) - (\eta_R v)(y))(\varphi(x) - \varphi(y))}{|x - y|^{N+2s}} dy dx = \int_{\mathbb{R}^N_+} (c_R^\infty + \mathcal{R}_R^\infty(x))\varphi(x) dx$$

and

$$c_R^{\infty} \in \mathbb{R}, \qquad \mathcal{R}_R^{\infty}(0) = 0, \qquad |\nabla \mathcal{R}_R^{\infty}(x)| \le CR^{\beta - 1} \qquad \text{for all } R \ge 2 \text{ and } x \in B_{\frac{R}{2}}.$$
 (6.39)

Consequently, since $\beta < \alpha_s < 1$,

$$(-\Delta)^s_{\mathbb{R}^N_+} v \stackrel{\beta}{=} 0 \qquad \text{in } \mathbb{R}^N_+.$$

Applying Theorem 3.4 and using (6.37), we deduce that $\frac{v}{x_N^{2s-1}}(x) = b + a \cdot x'$. This is in contradiction with (6.38).

Corollary 6.5. Under the assumptions of Proposition 6.4, there exist $a \in \mathbb{R}^{N-1}$ and $b \in \mathbb{R}$ with $|a| + |b| \leq C$ such that letting $\psi = \frac{w}{x_{N}^{2s-1}}$, we get

$$\left\|\psi - \left((a \cdot x' + b) - bT_3(0)x_N\right)\right\|_{L^{\infty}(B_r^+)} \le Cr^{1+\alpha} \quad \text{for all } r > 0.$$
(6.40)

Moreover $a = \nabla_{x'}\psi(0)$, $b = \psi(0)$ and $\partial_{x_N}\psi(0) = -bT_3(0)$.

Proof. Let $Q_r(x) = a_{r,w} \cdot x' + b_{r,w}(1 - T_3(0)x_N)$. Let $0 < \rho_2 \le \rho_1/4 \le 1/4$. Pick $k \in \mathbb{N}$ and $\sigma \in [1/4, 1/2]$ such that $\rho_2 = \sigma^k \rho_1$. Then provided $r \in (0, 1)$, by (6.26) and (6.14) we get

$$\begin{split} \rho_{1}|a_{\rho_{2},w} - a_{\rho_{1},w}| + |b_{\rho_{2},w} - b_{\rho_{1},w}| \\ &\leq \sum_{i=0}^{k-1} \sigma^{i+1}\rho_{1}|a_{\sigma^{i+1}\rho_{1},w} - a_{\sigma^{i}\rho_{1},w}| + \sum_{i=0}^{k-1} |b_{\sigma^{i+1}\rho_{1},w} - b_{\sigma^{i}\rho_{1},w}| \\ &\leq C \frac{1}{|B_{\sigma^{i+1}\rho_{1}}|^{\frac{1}{2}}} \|Q_{\sigma^{i+1}\rho_{1}} - Q_{\sigma^{i}\rho_{1}}\|_{L^{2}(B_{\sigma^{i+1}\rho_{1}})} \\ &\leq C \sum_{i=0}^{k-1} \left(\|\psi - Q_{\sigma^{i}\rho_{1}}\|_{L^{\infty}(B_{\sigma^{i}\rho_{1}})} + \|\psi - Q_{\sigma^{i+1}\rho_{1}}\|_{L^{\infty}(B_{\sigma^{i+1}\rho_{1}})} \right) \\ &\leq C \sum_{i=0}^{k-1} \left((\sigma^{i}\rho_{1})^{1+\alpha} + (\sigma^{i+1}\rho_{1})^{1+\alpha} \right) \leq C\rho_{1}^{1+\alpha}. \end{split}$$

This implies that the limits $a := \lim_{r\to 0} a_{r,w}$ and $b := \lim_{r\to 0} b_{r,w}$ exist. Moreover, letting $\rho_2 \to 0$ and $\rho_1 = 1/2$ in the above estimate, we get $|a| + |b| \le C$. Hence (6.40) follows. It is by now easy to see from (6.40), that $b = \psi(0)$ and $\nabla \psi(0) = (a, -bT_3(0))$.

Corollary 6.6. Under the hypothesis of Proposition 6.4, there exists a constant C depending only on N, s, α and β such that for all $\xi \in B_1^+ \cap \mathbb{R}e_N$,

$$[w/x_N^{2s-1}]_{C^{1+\alpha}(B_{\xi_N/2}(\xi))} \le C.$$

Proof. Let $\xi := (0, 2\rho) \in B_1^+$ with $\rho < 1/2$. We define

$$P(x) = \overline{\eta}(x_N)(x_N)_+^{2s-1}(a \cdot x' + b) - bT_3(0)\overline{\eta}(x_N)(x_N)_+^{2s},$$

where b and a are given by Corollary 6.5 and $T_3(0)$ is given by Lemma 6.2. Put $v_{\xi}(y) = w(\xi + \rho y) - P(\xi + \rho y)$, $f_{\xi}(y) = \rho^{2s} f(\xi + \rho y)$. We have

$$\mathcal{L}_{\Phi_{\rho}}v_{\xi} = f_{\xi}(x) + \rho^{2s}(\mathcal{L}_{\Phi}P)(\xi + \rho x) \qquad \text{for all } \varphi \in C_{c}^{\infty}(B_{1}^{+}), \tag{6.41}$$

where $\Phi_{\rho}(x) = \rho^{-1} \Phi(\rho x + \xi) = (x', x_N + \rho^{-1} \phi(\rho x'))$ and

$$\mathcal{L}_{\Phi_{\rho}}v(x) = c_{N,s} \int_{\{y_N > -2\}} \frac{v(x) - v(y)}{|\Phi_{\rho}(x) - \Phi_{\rho}(y)|^{N+2s}} \, dy.$$

By Lemma 6.2, we have

$$F(x) := (\mathcal{L}_{\Phi}P)(\xi + \rho x) = g_1(\xi + \rho x)\log(2\rho + \rho x_N) + g_2(\xi + \rho x)$$

32

from which it follows that $||F||_{C^{\beta}([0,1))} \leq C\rho^{\beta}|\log \rho| \leq C\rho^{\alpha}$. Note that $|u_{\xi}(x)| \leq C(1 + |x|^{2s+\alpha})$ and thus since $\Phi_{\rho} \in C^{2}(\mathbb{R}^{N}; \mathbb{R}^{N})$ is a global (volume preserving) diffeomorphism, by a change of variable and (6.41), we obtain

$$(-\Delta)^s_{\Omega_\rho} v_{\xi} \circ \Phi_\rho^{-1} = G_{\xi} \circ \Phi_\rho^{-1} \qquad \text{in } \Omega_\rho$$

where $G_{\xi}(x) := f_{\xi}(x) + \rho^{2s} F(\xi + \rho x)$ and $\Omega_{\rho} = \Phi_{\rho}^{-1}(\{y_N > -2\})$. We note that $[G_{\xi}]_{C^{\alpha}(B_{1/2})} \leq \rho^{2s+\alpha}$, $\|v_{\xi} \circ \Phi_{\rho}^{-1}\|_{L^{\infty}(B_{1/2})} \leq \rho^{2s+\alpha}$ and $B_1 \subset \Omega_{\rho}$. We then apply Lemma 5.5 to deduce that

$$\begin{split} \|v_{\xi} \circ \Phi_{\rho}^{-1}\|_{C^{2s+\alpha}(B_{1/4})} &\leq C\left(\int_{\Omega_{\rho} \setminus B_{1/8}} |y|^{-N-2s-1} |v_{\xi} \circ \Phi_{\rho}^{-1}(y)| dy + \rho^{2s+\alpha}\right) \\ &\leq C\left(\int_{\mathbb{R}^{N}_{+} \setminus B_{1/16}} |x|^{-N-2s-1} |v_{\xi}(x)| dx + \rho^{2s+\alpha}\right) \\ &\leq C\left(\sum_{i=0}^{\infty} \int_{2^{i-3} > |x| > 2^{i-4}} |x|^{-N-2s-1} |v_{\xi}(x)| dx + \rho^{2s+\alpha}\right) \\ &\leq C\left(\sum_{i=0}^{\infty} (2^{i}\rho)^{-N-2s-1} \rho^{2s+1} \int_{2^{i-3}\rho > |z| > 2^{i-4}\rho} v_{\xi}(z/\rho) |dz + \rho^{2s+\alpha}\right) \\ &\leq C\left(\sum_{i=0}^{\infty} (2^{i}\rho)^{-N-2s-1} \rho^{2s+1} \int_{2^{i}\rho > |x|} |w(x) - P(x)| dx + \rho^{2s+\alpha}\right) \\ &\leq C\left(\sum_{i=0}^{\infty} 2^{i(\alpha-1)} \rho^{2s+\alpha} + \rho^{2s+\alpha}\right) \leq C\rho^{2s+\alpha}, \end{split}$$

where we used (6.40) which yields $|w(x) - P(x)| \le C|x|^{2s+\alpha}$. We conclude that $\|v_{\xi}\|_{C^{2s+\alpha}(B_{1/4})} \le C\rho^{2s+\alpha}$.

Scaling and translating back, we then get

$$||w - P||_{L^{\infty}(B_{\rho/4}(\xi))} \le C\rho^{2s+\alpha}, \qquad [\nabla w - \nabla P]_{C^{\alpha}(B_{\rho/4}(\xi))} \le C\rho^{2s-1}.$$

Combining the above estimates with

$$\|1/x_N^{2s-1}\|_{L^{\infty}(B_{\rho/4}(\xi))} \le C\rho^{1-2s}, \qquad [\nabla(1/x_N^{2s-1})]_{C^{\alpha}(B_{\rho/4}(\xi))} \le C\rho^{-2s-\alpha}$$

we finally get

$$[\nabla(w/x^{2s-1}) - \nabla(P/x^{2s-1})]_{C^{\alpha}(B_{\rho/4}(\xi))} \le C.$$

Since $[\nabla(P/x^{2s-1})]_{C^{\alpha}(B_{\rho/4}(\xi))} = 0$, this yields

$$[\nabla(w/x^{2s-1})]_{C^{\alpha}(B_{\rho/4}(\xi))} \le C$$

and the proof is complete.

We now prove the following.

Theorem 6.7. Given $s \in (\frac{1}{2}, 1)$, let $\alpha_s \in (0, 1-s)$ be given by Lemma 2.6. Let $\Omega \subset \mathbb{R}^N$ be a $C^{2,\beta}$ domain, with $0 \in \partial\Omega$, and $f \in C^{\alpha}(\overline{\Omega} \cap B_1)$ with $\alpha < \min\{\alpha_s, 2s-1, \beta\}$ and $\alpha + 2s \neq 1$. Let $u \in H_0^s(\Omega)$ be such that

$$D_{\Omega}(u,\varphi) = \int_{\Omega} f\varphi \, dx \qquad \forall \varphi \in C_c^{\infty}(B_1 \cap \Omega),$$

and let $\delta = dist(x, \partial \Omega)$. Then,

$$\|u/\delta^{2s-1}\|_{C^{1+\alpha}(B_{\rho/2}\cap\overline{\Omega})} \le C\left(\|u\|_{L^2(\Omega)} + \|f\|_{C^{\alpha}(\overline{\Omega})}\right),\tag{6.42}$$

with $C, \rho > 0$ depending only on N, s, α , and Ω .

Moreover, letting $\psi = u/\delta^{2s-1}$, we have

$$\partial_{\nu}\psi(\sigma) = -(N-1)H_{\partial\Omega}(\sigma)\psi(\sigma) \qquad \text{for all } \sigma \in \partial\Omega \cap B_1, \tag{6.43}$$

where $H_{\partial\Omega}$ is the mean curvature of $\partial\Omega$.

Proof. We assume for simplicity that $||u||_{L^2(\Omega)} + ||f||_{C^{\alpha}(\overline{\Omega})} \leq 1$. Hence by Lemma 6.3, we get $||u/\delta^{2s-1}||_{L^{\infty}(\Omega \cap B_{1/2})} + ||u||_{C^{2s-1}(\overline{\Omega} \cap B_{1/2})} \leq C$.

Up to a scaling and a rotation, we may assume that $B_{1/4} \cap \partial \Omega$ is contained in the graph $\{x_N = \gamma(x')\}$ with $\gamma \in C^{2,\beta}(B'_{1/4})$. For $q \in \partial \Omega$, let $\nu(q)$ denote the unit exterior normal vector of Ω at q and denote $T_q \partial \Omega = \nu(q)^{\perp}$ the tangent plane of $\partial \Omega$ at q.

By the local inversion theorem, there exist $r_0 \in (0, 1/4]$ depending only on N, β and $\|\gamma\|_{C^{2,\beta}(B_1')}$ such that for any $q \in B_{r_0} \cap \partial\Omega$, there exist an orthonormal basis $(E_1(q) \dots, E_{N-1}(q))$ of $T_q \partial\Omega$ and $\tilde{\phi} \in C^{2,\beta}(B_{r_0}')$ such that

$$x' \mapsto q + \sum_{i=1}^{N-1} x_i E_i(q) - \widetilde{\phi}(x')\nu(q) : B'_{r_0} \to \partial\Omega$$

with $\|\widetilde{\phi}\|_{C^{2,\beta}(B'_{r_0})} \leq C(N,\beta,\gamma,r_0)$. It is clear that $\widetilde{\phi}(0) = |\nabla\widetilde{\phi}(0)| = 0$. We let $\eta \in C_c^{\infty}(B'_{r_0})$ with $\eta \equiv 1$ on $B'_{r_0/2}$ and put $\phi = \eta\widetilde{\phi}$. For all $q \in B_{r_0} \cap \partial\Omega$, we define

$$\Phi: \mathbb{R}^N \to \mathbb{R}^N, \qquad \Phi(x', x_N) = q + \sum_{i=1}^{N-1} x_i E_i(q) - (x_N + \phi(x')) \nu(q). \tag{6.44}$$

Next, decreasing r_0 if necessary, Φ is a global $C^{2,\beta}$ diffeomorphism and clearly $\text{Det}D\Phi(x) = 1$ for all $x \in \mathbb{R}^N$. We may assume that the (signed) distance function $d \in C^{2,\beta}(\{d < r_0/2\})$. We also recall that the mean curvature of $\partial\Omega$ at q is given by

$$H_{\partial\Omega}(q) := \frac{1}{N-1} \Delta\phi(0). \tag{6.45}$$

We define $Q_{\delta} := B'_{\delta} \times (0, \delta)$ and $\mathcal{Q}_{\delta} := \Phi(Q_{\delta}) \subset \Omega$. For any $x_0 \in \mathcal{Q}_{r_0/9}$, let $q \in \partial \Omega \cap B_{r_0}$ such that $|x_0 - q| = d(x_0)$. Letting $v = \chi_{r_0/4} u$, we have

$$(-\Delta)^s_{\Omega} v = f + c_{N,s} \int_{\Omega} \frac{(1 - \chi_{r_0/4}(y))u(y)}{|x - y|^{N+2s}} \, dy =: f_1.$$

Clearly $||f_1||_{C^{\alpha}(\mathcal{Q}_{r_{\alpha}/8})} \leq C$ for all $\alpha \in (0, \min(2s-1, \beta))$. Now we get

$$(-\Delta)^s_{\mathcal{Q}_{r_0/2}}v = f_1 - (-\Delta)^s_{\Omega \setminus \mathcal{Q}_{r_0/2}}v =: f_2,$$

and obviously $||f_2||_{C^{\alpha}(\mathcal{Q}_{r_0/8})} \leq C$ because $||v||_{C^{2s-1}(\mathcal{Q}_{r_0/8})} \leq C$. We define $w(x) = v(\Phi(x))$ and $g_2(x) = f_2(\Phi(x))$. Then by a change of variable, we get

$$c_{N,s} \int_{Q_{r_0/2}} \frac{w(x) - w(y)}{|\Phi(x) - \Phi(y)|^{N+2s}} \, dy = g_2(x).$$

Hence $w \in H^s(\mathbb{R}^N_+)$ satisfies $\|w/x_N^{2s-1}\|_{L^{\infty}(\mathbb{R}^N_+)} + \|w\|_{C^{2s-1}(\mathbb{R}^N_+)} \leq C$ and solves

$$c_{N,s} \int_{\mathbb{R}^N_+} \frac{w(x) - w(y)}{|\Phi(x) - \Phi(y)|^{N+2s}} \, dy = g_3(x), \tag{6.46}$$

with $g_3(x) = g_2(x) - c_{N,s} \int_{\mathbb{R}^N_+ \setminus Q_{r_0/2}} \frac{w(x) - w(y)}{|\Phi(x) - \Phi(y)|^{N+2s}} dy$. We have that $||g_3||_{C^{\alpha}(Q_{r_0/8})} \leq C$. Recall that $x_0 = q - |x_0 - q|\nu(q) = \Phi(0, |x_0 - q|)$. We can thus apply Corollary 6.6 to the

Recall that $x_0 = q - |x_0 - q|\nu(q) = \Phi(0, |x_0 - q|)$. We can thus apply Corollary 6.6 to the equation (6.46), to get

$$[w/x_N^{2s-1}]_{C^{1+\alpha}(B_{r/2}(\xi))} = [u \circ \Phi/x_N^{2s-1}]_{C^{1+\alpha}(B_{r/2}(\xi))} \le C, \quad \text{with } \xi = \Phi^{-1}(x_0) \in Q_{\frac{r_0}{9}} \cap \mathbb{R}e_N.$$
(6.47)

where $r = \xi_N/2 = |x_0 - q|/2$. In addition, Corollary 6.40, (6.1) and (6.45) yield

$$\partial_{x_N}[w/x_N^{2s-1}](0) = -T_3(0)[w/x_N^{2s-1}](0) = (N-1)H_{\partial\Omega}(q)[w/x_N^{2s-1}](0).$$
(6.48)

In view of (6.44) and (6.47), by changing variables, we thus obtained

 $[u/\delta^{2s-1}]_{C^{1+\alpha}(B_{r/8}(x_0))} \le C,$

where we used that $x \mapsto \frac{d(\Phi(x))}{x_N} \in C^{1+\alpha}(B_{r_0/2})$ and that $d = \delta$ in Ω . This implies that $[u/\delta^{2s-1}]_{C^{1+\alpha}(\mathcal{Q}_{r_0/9})} \leq C$ and (6.42) follows.

Finally using (6.48) and the fact that $d(\Phi(0, x_N)) = d(0, x_N) = x_N + O(x_N^2)$, we conclude that

$$\partial_{\nu}[u/\delta^{2s-1}](q) = -(N-1)H_{\partial\Omega}(q)[u/\delta^{2s-1}](q)$$

and the proof is complete.

We finally give the:

Proof of Theorem 1.3. The result follows from Theorem 6.7 and a covering argument. \Box

References

- N. Abatangelo and X. Ros-Oton, Obstacle problems for integro-differential operators: higher regularity of free boundaries, Adv. Math. 360 (2020), 106-931.
- [2] A. Audrito, J.-C. Felipe-Navarro and X. Ros-Oton, The Neumann problem for the fractional Laplacian: regularity up to the boundary. Preprint arXiv (2020), https://arxiv.org/abs/2006.10026v1
- [3] K. Bogdan, K. Burdzy, Z-Q. Chen, *Cencored stable processes*. Probab. Theory Relat. Fields. 127 (2003), 89-152.
- [4] L. Caffarelli, L. Silvestre, An extension problem related to the fractional Laplacian. Comm. Partial Differential Equations 32 (2007), 1245-1260.
- [5] S. Campanato, Proprieta di Holderianita di alcuni classi di funzioni. Ann. Scuolla, Norm. Sup. Pisa 17 (1963), 175-1118.
- [6] H. Chan, D. Gómez-Castro, J. L. Vázquez, Blow-up phenomena in nonlocal eigenvalue problems: When theories of L¹ and L² meet, J. Funct. Anal. 280 (2021), 108845.
- [7] Z.-Q. Chen, P. Kim, Green function estimate for censored stable processes. Probab. Theory Relat. Fields 124 (2002), 595-610.
- [8] Z.-Q. Chen, P. Kim, R. Song, Two-sided heat kernel estimates for censored stable-like processes. Probab. Theory Relat. Fields 146 (2010), 361.
- S. Dipierro, X. Ros-Oton, J. Serra, E. Valdinoci, Non-symmetric stable operators: regularity theory and integration by parts, preprint arXiv (2020) https://arxiv.org/abs/2012.04833.
- [10] S. Dipierro, O. Savin, E. Valdinoci, Definition of fractional Laplacian for functions with polynomial growth, Rev. Mat. Iberoam. 35 (2019), 1079-1122.
- [11] M. M. Fall, Regularity estimates for nonlocal Schrödinger equations. Discrete Contin. Dyn. Syst. 39 (2019), 1405–1456.
- [12] M. M. Fall, Regional fractional Laplacians: boundary regularity. Preprint arXiv (2020), https: //arxiv.org/abs/2007.04808v2

- [13] X. Fernández-Real, X. Ros-Oton, *Regularity Theory for Elliptic PDE*, forthcoming book (2021), available at the webpage of the authors.
- [14] P. Grisvard, *Elliptic Problems in Nonsmooth Domains*. SIAM Classics in Applied Mathematics, 2011.
- [15] Q. Y. Guan, Integration by Parts Formula for Regional Fractional Laplacian. Commun. Math. Phys. 266 (2006), 289-329.
- [16] Q. Y. Guan, Z. M. Ma, Reflected symmetric α-stable processes and regional fractional Laplacian. Probab. Theory Related Fields. 134 (2006), 649-694.
- [17] C. Mou, Y. Yi, Interior Regularity for Regional Fractional Laplacian. Commun. Math. Phys. (2015) 340: 233.
- [18] X. Ros-Oton and J. Serra, Regularity theory for general stable operators, J. Differential Equations 260 (2016), 8675–8715.
- [19] X. Ros-Oton and J. Serra, Boundary regularity for fully nonlinear integro-differential equations, Duke Math. J. 165 (2016), 2079–2154.

M.M.F.: AFRICAN INSTITUTE FOR MATHEMATICAL SCIENCES IN SENEGAL, KM 2, ROUTE DE JOAL, B.P. 14 18. MBOUR, SÉNÉGAL

Email address: mouhamed.m.fall@aims-senegal.org, mouhamed.m.fall@gmail.com

X.R.: ICREA, Pg. Lluis Companys 23, 08010 Barcelona, Spain; and Universitat de Barcelona, Departament de Matemàtiques i Informàtica, Gran Via de les Corts Catalanes 585, 08007 Barcelona, Spain.

Email address: xros@ub.edu