

Asymptotic Behavior of Common Connections in Sparse Random Networks

Bikramjit Das*

Tiandong Wang[†]

Gengling Dai[‡]

March 7, 2022

Abstract

Random network models generated using sparse exchangeable graphs have provided a mechanism to study a wide variety of complex real-life networks. In particular, these models help with investigating power-law properties of degree distributions, number of edges, and other relevant network metrics which support the scale-free structure of networks. Previous work on such graphs imposes a marginal assumption of univariate regular variation (e.g., power-law tail) on the bivariate generating graphex function. In this paper, we study sparse exchangeable graphs generated by graphex functions which are multivariate regularly varying. We also focus on a different metric for our study: the distribution of the number of common vertices (connections) shared by a pair of vertices. The number being high for a fixed pair is an indicator of the original pair of vertices being connected. We find that the distribution of number of common connections are regularly varying as well, where the tail indices of regular variation are governed by the type of graphex function used. Our results are verified on simulated graphs by estimating the relevant tail index parameters.

MSC 2010 subject classifications. 05C82, 60F15, 60G70

Key words. Random networks, common connections, power laws, multivariate regular variation.

1 Introduction

Modeling social, economic, and biological networks has become a principal area of interest for data scientists in the recent decades. The degree distribution provides an idea about the structure of the network, e.g., a power-law behavior here indicating the so-called “scale-free” structure; see [Voitalov et al. \[2019\]](#) for a recent discussion. Another quantity of potential interest to network scientists, especially in the context of social networking platforms like Facebook, Instagram, LinkedIn, and Twitter, is the number of common connections or common friends between two vertices. This can be thought of as a generalization of the degree of a vertex.

*Engineering Systems and Design, Singapore University of Technology and Design, 8 Somapah Road, 487372 Singapore, *E-mail:* bikram@sutd.edu.sg

[†]Department of Statistics, Texas A&M University, TX77843, United States, *E-mail:* twang@stat.tamu.edu

[‡]Engineering Systems and Design, Singapore University of Technology and Design, 8 Somapah Road, 487372 Singapore, *E-mail:* gengling.dai@mymail.sutd.edu.sg

Network recommendation systems use this number as one of the metrics to suggest potential new vertices (friends) to platform users [Gupta et al., 2013]. The connection between *link prediction* and the size of common connections has been explored in the literature; see Barabási et al. [2002], Liben-Nowell and Kleinberg [2007], Newman [2001] for further discussions.

Although of interest to network scientists, only few theoretical results exist in the literature characterizing the asymptotic distribution of common connections for growing network models. While studying clustering coefficients for random intersection graph, Bloznelis and Kurauskas [2016] find the asymptotic distribution of common connections for a randomly selected pair of vertices. In addition, for a certain class of linear preferential attachment models, Das and Ghosh [2021] identify the growth rate for common connections between two fixed vertices in the graph. In our paper, we study common connections in the context of the erstwhile popular and flexible framework of exchangeable random graph models called *graphex processes*.

Graphex models, or vertex-exchangeable random graphs have been used extensively to study statistical network models that are dense (Hoover [1979], Aldous [1981], Lovász and Szegeedy [2006], Diaconis and Janson [2008]). Recent work has successfully extended the classical graphex framework to sparse networks, which mimics many real-world networks; see Borgs et al. [2019], Caron and Fox [2017]. In particular, they point out that exchangeable random measures admit a representation theorem due to Kallenberg [1990], giving a general construction for such graph models. Moreover, Veitch and Roy [2015] and Borgs et al. [2018] show how such construction naturally generalizes the dense exchangeable graphex framework to the sparse regime, and analyze some of the properties of the associated class of random graphs, called graphex processes; also see for instance, Janson [2016, 2017], Veitch and Roy [2019] and Borgs et al. [2019]. Furthermore, Herlau et al. [2016] and Todeschini et al. [2020] have developed sparse graph models with (overlapping) community structure within the graphex framework.

In this paper, we work under the framework of sparse vertex-exchangeable random graphs generated by graphex functions which are *multivariate regularly varying*, a modest generalization of power-law tailed functions. The regular variation of the graphex function creates a sparse graph, and we show that the distribution of common connections in the generated graphs is also regularly varying, i.e., approximately power-law tailed. Moreover, we observe that depending on the type of the graphex function, i.e., whether it is a product of univariate functions (called *separable*), or not (called *non-separable*), we get different asymptotic behaviors of common connections in the random graph. We estimate necessary parameters in simulated random graphs to numerically verify and support the results obtained.

The remaining of the paper proceeds as follows. In Section 2, we introduce the random graph framework, as well as necessary definitions and assumptions of multivariate regularly varying functions in this context. We also characterize effects of the regular variation assumption on graphex functions to particular marginals of the graphex functions. The main results on the asymptotic distribution of common connections are collected in Section 3. In addition, Section 4 presents results from simulation studies, where we provide typical examples of sparse random networks and verify our results of Section 3. Finally, concluding remarks are given in Section 5.

2 Random graphs, regular variation and graphex functions

In this section, we create our framework for network models via graphex processes, discuss multivariate regular variation, and derive regular variation properties for functionals of graphex processes.

2.1 Random graph models

We now present a framework which models sparse random graphs using (infinite) Poisson point processes on $\mathbb{R}_+^2 = [0, \infty)^2$. Finite graphs are obtained by appropriately truncating the support of the point process. The model representation is based on the work of [Borgs et al. \[2018\]](#), [Caron and Fox \[2017\]](#), [Caron et al. \[2020\]](#), [Veitch and Roy \[2015\]](#). Here we follow their construction and notation. Such representation is also related to Kallenberg exchangeable graphs [\[Kallenberg, 1990\]](#), and can be characterized by a symmetric measurable function, $W : [0, \infty)^2 \mapsto [0, 1]$, often called a *graphex function*. A variety of sparse random graphs can be parametrized using the graphex function W (we ignore terms corresponding to isolated edges and stars).

Let $\Pi = (\theta_i, \eta_i)_{i=1,2,\dots}$ be a unit-rate Poisson process on \mathbb{R}_+^2 . In this representation, vertices are embedded at some location $\theta_i \in \mathbb{R}_+$, and there is a latent variable $\eta_i \in (0, \infty)$ which along with the graphex function W determines the edges of the graph. For this paper, we will define a graphex function as follows.

Definition 2.1. A symmetric measurable function $W : [0, \infty)^2 \mapsto [0, 1]$ is a *graphex function* if it satisfies the following criteria:

- (a) $0 < \overline{W} = \int_0^\infty \int_0^\infty W(x, y) dx dy < \infty$;
- (b) $\int_0^\infty W(x, x) dx < \infty$;
- (c) $\lim_{x \rightarrow 0} W(x, x)$ and $\lim_{x \rightarrow \infty} W(x, x)$ both exist.

Note that we have $\lim_{x \rightarrow 0} W(x, x) = 0$, since $\overline{W} < \infty$. Other necessary regularity conditions on W will be imposed later. Edges of the graph induced by Π and W are given by a point process:

$$Z = \sum_{i,j} Z_{ij} \delta_{(\theta_i, \theta_j)}$$

where $Z_{ij} = Z_{ji}$ is a binary variable which takes the value 1 if there is an edge between θ_i and θ_j and 0 otherwise. Given the graphex function W and $i \leq j$, we have

$$Z_{ij} | \Pi, W \sim \text{Bernoulli}(W(\eta_i, \eta_j))$$

and $\{Z_{ij} : i \leq j\}$ are independent random variables. We obtain a finite size random graph family $(\mathcal{G}_t)_{t \geq 0}$, often called a graphex process, by restricting the process $\sum_{i,j} Z_{ij} \delta_{(\theta_i, \theta_j)}$ to $[0, t]^2$. Here $\mathcal{G}_t = (\mathcal{E}_t, \mathcal{V}_t)$ denotes a graph of size $t \geq 0$ with edge set \mathcal{E}_t and vertex set \mathcal{V}_t given by

$$\begin{aligned} \mathcal{E}_t &= \{\theta_i | \theta_i \leq t \text{ and } Z_{ik} = 1 \text{ for some } k \text{ with } \theta_k \leq t\}, \\ \mathcal{V}_t &= \{(\theta_i, \theta_j) | \theta_i, \theta_j \leq t \text{ and } Z_{ij} = 1\}. \end{aligned}$$

The following quantities are of interest for studying properties of graphex processes: the univariate and bivariate graphex marginal functions $\mu_1 : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ and $\mu_2 : \mathbb{R}_+^2 \rightarrow \mathbb{R}_+$ are given by

$$\mu_1(x) = \int_0^\infty W(x, y) dy, \tag{2.1}$$

$$\mu_2(x, y) = \int_0^\infty W(x, z) W(y, z) dz. \tag{2.2}$$

The integrability of W in \mathbb{R}_+^2 guarantees that both μ_1 and μ_2 are well-defined and finite. We can check that $\mu_1(\eta_i)$ is proportional to the average number of vertices that a vertex θ_i is connected to, whereas $\mu_2(\eta_i, \eta_j)$ is proportional to the average number of common connections (vertices) between the two vertices at θ_i and θ_j . We also further define the d -variate graphex marginal for $d \geq 3$ as:

$$\mu_d(x_1, \dots, x_d) := \int_0^\infty \prod_{i=1}^d W(x_i, z) dz, \quad x_i \neq x_j, \quad i \neq j. \quad (2.3)$$

2.2 Regular variation

Studies involving the Kallenberg exchangeable *sparse* graphs have been occasionally carried out under the assumption that the univariate graphex marginal $\mu_1(x)$ has a power-law-like tail (regularly varying) as $x \rightarrow \infty$; see [Caron et al. \[2020\]](#), [Naulet et al. \[2021\]](#). Instead of imposing a condition on μ_1 , we start by imposing important conditions on the generating graphex function, W , which is the direct input for the generation of the underlying network. It turns out that the regularity conditions on W are also closely connected to the theory of multivariate regular variation [[Bingham et al., 1989](#), [Resnick, 2007](#)], which we now outline.

A function $f : (0, \infty) \rightarrow (0, \infty)$ is regularly varying (at infinity) if $\lim_{t \rightarrow \infty} f(tx)/f(t) = x^\beta$ for $x > 0$ and $\beta \in \mathbb{R}$. Here β is the tail index or index of regular variation and we write $f \in \mathcal{RV}_\beta$. In dimensions $d > 1$ this can be extended to multivariate regular variation, cf. [de Haan and Resnick \[1987\]](#), [Stam \[1977\]](#): suppose $\mathbb{C} \subset \mathbb{R}_+^d$ is a cone, that is, $\mathbf{x} \in \mathbb{C}$ if and only if $t\mathbf{x} \in \mathbb{C}$ for all $t > 0$. A function $f : \mathbb{C} \rightarrow (0, \infty)$ is (multivariate) regularly varying with limit function λ with $\lambda(\mathbf{x}) > 0$ for $\mathbf{x} \in \mathbb{C}$ and tail index β : if there exists a function $g : (0, \infty) \rightarrow (0, \infty)$ with $g \in \mathcal{RV}_\beta$ such that

$$\lim_{t \rightarrow \infty} \frac{f(t\mathbf{x})}{g(t)} = \lambda(\mathbf{x}), \mathbf{x} \in \mathbb{C}, \quad (2.4)$$

or, equivalently: if there exists a function $b : (0, \infty) \rightarrow (0, \infty)$ with $b \in \mathcal{RV}_{-1/\beta}$

$$\lim_{t \rightarrow \infty} tf(b(t)\mathbf{x}) = \lambda(\mathbf{x}), \mathbf{x} \in \mathbb{C}. \quad (2.5)$$

As a consequence, we have the homogeneity property that $\lambda(t\mathbf{x}) = t^\beta \lambda(\mathbf{x}), \mathbf{x} \in \mathbb{C}, t > 0$. Moreover, we can obtain (2.5) from (2.4) by defining $b(t) = g^\leftarrow(1/t)$ where f^\leftarrow denotes the generalized left inverse of a monotone function; since without loss of generality, g can be assumed to be monotone. We write $f \in \mathcal{MRV}(\beta, g, \lambda, \mathbb{C})$ when following (2.4) or $f \in \mathcal{MRV}(\beta, b, \lambda, \mathbb{C})$ when following (2.5); one or more of the parameters are also often dropped for convenience. Both b and g are referred to as *scaling* functions.

2.3 Tail behavior of the univariate and bivariate graphex marginals

The following result characterizes the asymptotic behaviour of μ_1 and μ_2 at infinity, given that W is multivariate regularly varying. The proof uses ideas from [[de Haan and Resnick, 1987](#), Theorem 2.1]. A uniformity condition in addition to regular variation is imposed to guarantee regular variation of the integrals μ_1, μ_2 , see [[de Haan and Resnick, 1987](#), Section 2] for further discussions. Let $\|\cdot\| : (x, y) \rightarrow \|(x, y)\|$ denote a norm on \mathbb{R}^2 .

Theorem 2.2. Suppose the graphex function $W : \mathbb{R}_+^2 \rightarrow [0, 1]$ is regularly varying on $\mathbb{R}_+^2 \setminus \{\mathbf{0}\}$ with limit function ω and scaling function $h \in \mathcal{RV}_{-\alpha}$ where $\alpha > 1$. Moreover assume that ω is bounded on the unit sphere $\mathbb{N} := \{(x, y) \in \mathbb{R}_+^2 \setminus \{\mathbf{0}\} : \|(x, y)\| = 1\}$ and satisfies the uniformity condition

$$\lim_{t \rightarrow \infty} \sup_{(x, y) \in \mathbb{N}} \left| \frac{W(tx, ty)}{h(t)} - \omega(x, y) \right| = 0. \quad (2.6)$$

Then the following holds:

(i) The univariate graphex marginal μ_1 is regularly varying with tail index $-(\alpha - 1)$ and

$$\lim_{t \rightarrow \infty} \frac{\mu_1(tx)}{th(t)} = \int_0^\infty \omega(x, y) dy \quad x > 0. \quad (2.7)$$

(ii) The bivariate graphex marginal μ_2 is regularly varying on $(0, \infty)^2$ with tail index $-(2\alpha - 1)$ and

$$\lim_{t \rightarrow \infty} \frac{\mu_2(tx, ty)}{th^2(t)} = \int_0^\infty \omega(x, z)\omega(y, z) dz \quad (x, y) \in (0, \infty)^2. \quad (2.8)$$

(iii) The convergence in (2.8) is uniform on sets bounded away from the axes, i.e., for any fixed $\delta > 0$:

$$\lim_{t \rightarrow \infty} \sup_{x \wedge y \geq \delta} \left| \frac{\mu_2(tx, ty)}{th^2(t)} - \int_0^\infty \omega(x, z)\omega(y, z) dz \right| = 0. \quad (2.9)$$

Proof. First, using a change of variable $v = tz$ we have

$$\frac{\mu_1(tx)}{th(t)} = \int_0^\infty \frac{W(tx, z)}{th(t)} dz = \int_0^\infty \frac{W(tx, tv)}{h(t)} dv, \quad x > 0, \quad (2.10)$$

$$\frac{\mu_2(tx, ty)}{th^2(t)} = \int_0^\infty \frac{W(tx, z)W(ty, z)}{th^2(t)} dz = \int_0^\infty \frac{W(tx, tv)}{h(t)} \frac{W(ty, tv)}{h(t)} dv, \quad (x, y) \in (0, \infty)^2. \quad (2.11)$$

By our assumptions, the integrands in (2.10) and (2.11) are non-negative and converge to $\omega(x, v)$ and $\omega(x, v)\omega(y, v)$ respectively as $t \rightarrow \infty$. Hence if we show that the integrands are bounded by an integrable function and the limit are integrable, then (2.7) and (2.8) holds using dominated convergence. Since all norms in \mathbb{R}^2 are equivalent, without loss of generality let us fix a particular norm in \mathbb{R}^2 given by $\|(x, y)\| = |x| \vee |y|$.

(i) For fixed $x > 0$, and any $v \geq 0$, with $t > 0$,

$$\frac{W(tx, tv)}{h(t)} = \frac{W(t\|(x, v)\| \cdot \frac{(x, v)}{\|(x, v)\|})}{h(t\|(x, v)\|)} \frac{h(t\|(x, v)\|)}{h(t)}.$$

Since $\|(x, v)\| \geq x > 0$, and $(x, v)/\|(x, v)\| \in \mathbb{N}$, given a fixed $\eta > 0$, we obtain from (2.6) that for $t > t_0 \equiv t_0(x, \eta)$,

$$\frac{W\left(t\|(x, v)\| \frac{(x, v)}{\|(x, v)\|}\right)}{h(t\|(x, v)\|)} \leq \sup_{\mathbf{a} \in \mathbb{N}} \omega(\mathbf{a}) + \eta < \infty, \quad (2.12)$$

since ω is bounded on \aleph . Using Potter's bounds [Resnick, 2007], for a large enough t , $h(t\|(x, v)\|)/h(t)$ is bounded by $c\|(x, v)\|^{-\alpha+\rho}$ for a constant $c > 0$ and ρ such that $0 < \rho < \alpha - 1$. Hence,

$$\frac{W(tx, tv)}{h(t)} \leq C\|(x, v)\|^{-\alpha+\rho}, \quad (2.13)$$

for some constant $C > 0$, and

$$\int_0^\infty C\|(x, v)\|^{-\alpha+\rho} dv = Cx^{-\alpha+\rho} \int_0^x dv + C \int_x^\infty u^{-\alpha+\rho} dv < \infty, \quad (2.14)$$

which shows that the integrand in (2.10) is bounded by an integrable function. By homogeneity property we also have $\omega(t(x, y)) = t^{-\alpha}\omega(x, y)$ for $(x, y) \in \mathbb{R}_+^2 \setminus \{\mathbf{0}\}$ and hence

$$\int_0^\infty \omega(x, y) dy = \int_0^\infty \|(x, y)\|^{-\alpha} \omega\left(\frac{(x, y)}{\|(x, y)\|}\right) dy \leq \sup_{\mathbf{a} \in \aleph} \omega(\mathbf{a}) \int_0^\infty (x \vee y)^{-\alpha} dy < \infty, \quad (2.15)$$

since ω is bounded in \aleph and $\alpha > 1$. Hence (2.7) is a consequence of $W \in \mathcal{MRV}(-\alpha, h, \omega)$, (2.13), (2.14), (2.15) and the dominated convergence theorem. Therefore $\mu_1 \in \mathcal{RV}_{-(\alpha-1)}$.

- (ii) Using the bounds in (2.13), (2.14) (and finding similar bounds for $W(ty, tv)/h(t)$), we can conclude that for any $(x, y) \in (0, \infty)^2$, the integrand in (2.11) is bounded by an integrable function. To see that the limit (as $t \rightarrow \infty$) of the integrand in (2.11) is integrable, observe that

$$\begin{aligned} \int_0^\infty \omega(x, z)\omega(y, z) dz &= \int_0^\infty \|(x, z)\|^{-\alpha} \omega\left(\frac{(x, z)}{\|(x, z)\|}\right) \|(y, z)\|^{-\alpha} \omega\left(\frac{(y, z)}{\|(y, z)\|}\right) dz \\ &\leq \left(\sup_{\mathbf{a} \in \aleph} \omega(\mathbf{a})\right)^2 \int_0^\infty (x \vee z)^{-\alpha} (y \vee z)^{-\alpha} dz < \infty, \end{aligned}$$

since ω is bounded in \aleph and $\alpha > 1$. Then (2.11) follows from the dominated convergence theorem, and we have

$$\mu_2 \in \mathcal{MRV}\left(-(2\alpha - 1), th^2(t), \int_0^\infty \omega(x, z)\omega(y, z) dz, (0, \infty)^2\right).$$

- (iii) From de Haan and Resnick [1987, Theorem 2.1] we know that the uniformity condition (2.6) also implies that for any $\delta > 0$,

$$\lim_{t \rightarrow \infty} \sup_{\|(x, y)\| > \delta} \left| \frac{W(tx, ty)}{h(t)} - \omega(x, y) \right| = 0. \quad (2.16)$$

Now for fixed $\delta > 0$ and $x \wedge y > \delta$,

$$\begin{aligned} \left| \frac{\mu_2(tx, ty)}{th^2(t)} - \int_0^\infty \omega(x, z)\omega(y, z) dz \right| &\leq \int_0^\infty \left| \frac{W(tx, tz)W(ty, tz)}{h^2(t)} - \omega(x, z)\omega(y, z) \right| dz \\ &\leq \int_0^\infty \frac{W(tx, tz)}{h(t)} \left| \frac{W(ty, tz)}{h(t)} - \omega(y, z) \right| dz + \int_0^\infty \omega(y, z) \left| \frac{W(tx, tz)}{h(t)} - \omega(x, z) \right| dz. \end{aligned}$$

Since $\{(x, y) \in \mathbb{R}_+^2 : x \wedge y > \delta\} \subset \{(x, y) \in \mathbb{R}_+^2 : x \vee y = \|(x, y)\| > \delta\}$, then by (2.16), given $\epsilon > 0$, there exists t_0 such that for $t > t_0$, both $|\frac{W(ty, tz)}{h(t)} - \omega(y, z)| < \epsilon$ and $|\frac{W(tx, tz)}{h(t)} - \omega(x, z)| < \epsilon$ for any (x, y) with $x \wedge y > \delta$. Therefore for $t > t_0$,

$$\begin{aligned} \sup_{x \wedge y > \delta} \left| \frac{\mu_2(tx, ty)}{th^2(t)} - \int_0^\infty \omega(x, z) \omega(y, z) \, dz \right| &\leq \epsilon \sup_{x > \delta} \int_0^\infty \frac{W(tx, tz)}{h(t)} \, dz + \epsilon \sup_{y > \delta} \int_0^\infty \omega(y, z) \, dz \\ &= \epsilon \sup_{x > \delta} A_{x,t} + \epsilon \sup_{y > \delta} B_y \quad (\text{say}). \end{aligned}$$

From (2.13) and (2.14), since $x > \delta > 0$, for large enough t , $\sup_{x > \delta} A_{x,t} \leq C_1$ for some constant $C_1 > 0$. Also using (2.15), we can check that $\sup_{y > \delta} B_y \leq C_2$ for some constant $C_2 > 0$. Therefore

$$\lim_{t \rightarrow \infty} \sup_{x \wedge y \geq \delta} \left| \frac{\mu_2(tx, ty)}{th^2(t)} - \int_0^\infty \omega(x, z) \omega(y, z) \, dz \right| \leq \epsilon(C_1 + C_2)$$

and since $\epsilon > 0$ is arbitrary, we have (2.9). □

Example 2.3. Consider the graphex function $W : \mathbb{R}_+^2 \rightarrow (0, \infty)$ with $\alpha > 1$:

$$W(x, y) = \frac{1}{1 + x^\alpha + y^\alpha}. \quad (2.17)$$

Here

$$W \in \mathcal{MRV} \left(-\alpha, h(t) = t^{-\alpha}, \omega(x, y) = \frac{1}{x^\alpha + y^\alpha}, (0, \infty)^2 \right),$$

and we can check that the uniformity condition (2.6) holds. Therefore the univariate marginal $\mu_1 \in \mathcal{RV}_{-(\alpha-1)}$ and for $x > 0$,

$$\lim_{t \rightarrow \infty} \frac{\mu_1(tx)}{th(t)} = \int_0^\infty \omega(x, y) \, dy = x^{-\alpha+1} \int_0^\infty \frac{1}{1 + z^\alpha} \, dz = \frac{\pi}{\alpha} \operatorname{cosec}(\pi/\alpha) x^{-\alpha+1}.$$

See Gradshteyn and Ryzhik [2007, §3.241.2]. Moreover, the bivariate marginal satisfies

$$\mu_2 \in \mathcal{MRV}(-(2\alpha - 1), th^2(t), \nu(\cdot), (0, \infty)^2),$$

where ν is hard to compute in closed form in general. For $\alpha = 2$, we can compute ν in closed form, which is given by

$$\nu(x, y) = \lim_{t \rightarrow \infty} \frac{\mu_2(tx, ty)}{th^2(t)} = \int_0^\infty \frac{1}{x^2 + z^2} \frac{1}{y^2 + z^2} \, dz = \frac{\pi}{2x^2y + 2xy^2}, \quad (x, y) \in (0, \infty)^2.$$

Remark 1. A classical example of a multivariate regularly varying W is when it is *separable*, i.e., $W(x, y) = U(x)U(y)$ with $U \in \mathcal{RV}_{-\alpha}$ for some $\alpha > 1$. Unfortunately, the uniformity condition (2.6) fails to hold in this case and we cannot apply Theorem 2.2. Nevertheless, we can still ascertain the tail behavior in this case easily because of the separable structure.

Lemma 2.4. Let $W : \mathbb{R}_+^2 \rightarrow [0, 1]$ be a graphex function such that $W(x, y) = U(x)U(y)$ where $U : \mathbb{R}_+ \rightarrow (0, \infty)$ satisfies $U \in \mathcal{RV}_{-\alpha}$ for some $\alpha > 1$. Then the following holds:

(i) $W \in \mathcal{MRV}(-2\alpha, U^2(t), \omega(x, y), (0, \infty)^2)$ where

$$\lim_{t \rightarrow \infty} \frac{W(tx, ty)}{U^2(t)} = x^{-\alpha} y^{-\alpha} =: \omega(x, y), \quad (x, y) \in (0, \infty)^2,$$

(ii) $\mu_1 \in \mathcal{RV}_{-\alpha}$ where

$$\lim_{t \rightarrow \infty} \frac{\mu_1(tx)}{U(t)} = x^{-\alpha} \int_0^\infty U(z) dz, \quad x > 0,$$

(iii) $\mu_2 \in \mathcal{MRV}(-2\alpha, U^2(t), C\omega(x, y), (0, \infty)^2)$ where $C = \int_0^\infty U^2(z) dz$.

Proof. The proof is an easy consequence of $U \in \mathcal{RV}_{-\alpha}$ and is omitted here. \square

Example 2.5. Consider the graphex function $W : \mathbb{R}_+^2 \rightarrow (0, \infty)$ with $\alpha > 1$:

$$W(x, y) = U(x)U(y) := \frac{1}{1+x^\alpha} \frac{1}{1+y^\alpha}.$$

Clearly,

$$W \in \mathcal{MRV}\left(-2\alpha, U^2(t), \omega(x, y) = \frac{1}{x^\alpha} \frac{1}{y^\alpha}, (0, \infty)^2\right).$$

Using Example 2.3, the univariate marginal $\mu_1 \in \mathcal{RV}_{-\alpha}$ and for $x > 0$,

$$\lim_{t \rightarrow \infty} \frac{\mu_1(tx)}{U(t)} = x^{-\alpha} \int_0^\infty U(z) dz = x^{-\alpha} \int_0^\infty \frac{1}{1+z^\alpha} dz = \frac{\pi}{\alpha} \operatorname{cosec}(\pi/\alpha) x^{-\alpha}.$$

Moreover, $\mu_2 \in \mathcal{MRV}(-2\alpha, U^2(t), C\omega(x, y), (0, \infty)^2)$, where $C = ((\pi/\alpha)\operatorname{cosec}(\pi/\alpha))^2$.

3 Common connections

In this section, our goal is to understand the behavior of the number of common connections between two vertices, first for a fixed pair of vertices and eventually for a randomly chosen pair in a graph driven by the (Kallenberg exchangeable) graphex process as described in Section 2.

3.1 Distribution of common connections in \mathcal{G}_t

Recall that from the unit-rate Poisson process $\Pi = (\theta_i, \eta_i)_{i=1,2,\dots}$, the finite-size graphex \mathcal{G}_t is created by restricting $\sum_{i,j} Z_{ij} \delta_{(\theta_i, \theta_j)}$ to $[0, t]^2$, where $Z_{ij} \sim \text{Bernoulli}(W(\eta_i, \eta_j))$, and $Z_{ij} = Z_{ji}$. The first co-ordinate in Π does not provide any relevant information other than picking the plausible vertices, so for ease of computation, we project Π to its second co-ordinate as

$$\Pi_t^\eta = \{\eta_i | (\theta_i, \eta_i) \in \Pi_t\},$$

so that Π_t^η is a one-dimensional Poisson process with rate t .

Let I be an index set. Then following [Veitch and Roy, 2015, Section 5], for a locally finite, simple sequence $\phi = (x_i)_{i \in I}$, and a sequence of values $\{u_{ij}\} \in [0, 1]$ with $u_{ij} = u_{ji}$, $i, j \in I$, we

define the *2nd order common connections function* or *2-c-degree function* at (vertices) $x_i, x_j \in \phi$ to be

$$C(x_i, x_j, \phi, (u_{ij})) := \sum_{x_k \in \Pi_t^\eta \setminus \{x_i, x_j\}} \mathbf{1}_{\{W(x_i, x_k) \geq u_{ik}\}} \mathbf{1}_{\{W(x_j, x_k) > u_{jk}\}}. \quad (3.1)$$

With this definition, for the graphex \mathcal{G}_t realized at $Z_{ij} = z_{ij}$ (which are realizations of $\text{Unif}(0, 1)$ random variables), the 2-c-degree function at $(\theta_1, \eta_1), (\theta_2, \eta_2) \in \Pi_t$ is

$$C(\eta_1, \eta_2, \Pi_t^\eta, (z_{ij})). \quad (3.2)$$

Taking η_1, η_2 to be the proxy for the points $(\theta_1, \eta_1), (\theta_2, \eta_2) \in \Pi_t$, we refer to $C(\eta_1, \eta_2, \Pi_t^\eta, (z_{ij}))$ as the number of common friends/connections of η_1, η_2 in \mathcal{G}_t . We can analogously define a *k-c-degree function*, for the number of common connections between $k \geq 2$ vertices as well.

We now explore how $C(\eta_1, \eta_2, \Pi_t^\eta, (Z_{ij}))$ behaves for fixed t , and later in Section 3.2, we study the asymptotic behavior of common friends for a pair of randomly chosen vertices in \mathcal{G}_t as $t \rightarrow \infty$. Since for a finite set of points $x_1, \dots, x_k \in \mathbb{R}_+$, the probability that $x_1, \dots, x_k \in \Pi_t^\eta$ is 0, it is traditional to use some tools from Palm theory. First note that the conditional distribution of Π_t^η given x_1, \dots, x_k is equivalent to the distribution of $\Pi_t^\eta \cup \{x_1, \dots, x_k\}$ which is the (extended) Slivnyak-Mecke theorem if Π_t^η is a Poisson process; see [Moller and Waagepetersen, 2003, Theorem 3.3]. To make the paper self-contained, we state the extended Slivnyak-Mecke theorem below.

Theorem 3.1. *Let $S \subset \mathbb{R}^d$ and define \mathcal{N} as the set of locally finite subsets of S . Let $X = (x_i)_{i \geq 1} \in \mathcal{N}$ be a Poisson point process on S with a diffuse and locally finite mean measure μ . Then, for any $n \geq 1$ and any measurable function $h : \mathcal{N} \times S^n \mapsto [0, \infty)$,*

$$\mathbb{E} \left[\sum_{\xi_1 \neq \dots \neq \xi_n \in X} h(X \setminus \{\xi_1, \dots, \xi_n\}) \right] = \int_{S^n} \mathbb{E} [h(X \setminus \{\xi_1, \dots, \xi_n\})] \mu(d\xi_1) \cdots \mu(d\xi_n).$$

With Theorem 3.1 available, we present in the following proposition that the 2-c-degree of two fixed vertices in a the graphex \mathcal{G}_t is Poisson distributed.

Proposition 3.2. *Let $x, y \in \mathbb{R}_+$, and $\zeta_{\{i,j\}}$ be a symmetric array of iid $\text{Uniform}[0, 1]$ random variables. Then $C(x, y, \Pi_t^\eta \cup \{x, y\}, (\zeta_{\{i,j\}}))$ follows a Poisson distribution with rate $t\mu_2(x, y)$.*

Proof. In the sequel, we suppress the notation as

$$C(x, y) \equiv C(x, y, \Pi_t^\eta \cup \{x, y\}, (\zeta_{\{i,j\}})),$$

and use $i(s) \equiv i(s, \Pi_t^\eta)$ to denote the index of the point $s \in \Pi_t^\eta$ with respect to the natural ordering on \mathbb{R}_+ . Then we have

$$C(x, y) = \sum_{z \in \Pi_t^\eta} \mathbf{1}_{\{W(x, z) \geq \zeta_{i(x)i(z)}\}} \cdot \mathbf{1}_{\{W(y, z) \geq \zeta_{i(y)i(z)}\}}.$$

Note also that

$$\int_{\mathbb{R}_+} \int_{[0,1]} \int_{[0,1]} \mathbf{1}_{\{u \leq W(x, z)\}} \mathbf{1}_{\{v \leq W(y, z)\}} t du dv dz = t \int_{\mathbb{R}_+} W(x, z) W(y, z) dz = t\mu_2(x, y) < \infty.$$

Then by Campbell's theorem (cf. [Kingman, 1993, Section 5.3]), the characteristic function of $C(x, y)$ is

$$\begin{aligned}
\mathbb{E}[\exp(itC(x, y))] &= \mathbb{E}[\exp(it \sum_{z \in \Pi_t^\eta} \mathbb{1}_{\{W(x, z) \geq \zeta_{i(x)i(z)}\}} \cdot \mathbb{1}_{\{W(y, z) \geq \zeta_{i(y)i(z)}\}})] \\
&= \exp \left\{ \int_{\mathbb{R}_+} \int_{[0,1]} \int_{[0,1]} (1 - e^{it \mathbb{1}_{\{u \leq W(x, z)\}} \mathbb{1}_{\{v \leq W(y, z)\}}}) t du dv dz \right\} \\
&= \exp \left\{ t \sum_{n=1}^{\infty} \frac{(it)^n}{n!} \int_{\mathbb{R}_+} \int_{[0,1]} \int_{[0,1]} \mathbb{1}_{\{u \leq W(x, z)\}} \mathbb{1}_{\{v \leq W(y, z)\}} du dv dz \right\} \\
&= \exp \left\{ t \mu_2(x, y) (e^{it} - 1) \right\}.
\end{aligned}$$

Hence, $C(x, y)$ is a Poisson random variable with rate $t \mu_2(x, y)$. \square

Proposition 3.2 characterizes the distribution for a fixed pair of (x, y) in \mathcal{G}_t with t also fixed. In the next section, we extend our analysis to the asymptotic behavior of the 2-c-degree distribution when t goes to infinity.

3.2 Asymptotic distribution of common connections




Figure 3.1: The empirical distribution of common connections in a simulated sparse network generated from the graphex process with $W(x, y) = (1 + x + y)^{-3}$, $x, y \geq 0$.

To analyze the asymptotic distribution, we start with a simulated example where $W(x, y) = (1 + x + y)^{-3}$, $x, y \geq 0$. We compute the empirical distribution of the number of common friends, i.e.,

$$\frac{\sum_{x, y \in \Pi_t^\eta} \mathbb{1}_{\{C(x, y) = k\}}}{\sum_{x, y \in \Pi_t^\eta} \mathbb{1}_{\{C(x, y) > 0\}}}, \quad k \geq 1,$$

and numerical results are plotted in Figure 3.1. We observe that the plot looks linear until a certain point and then it quickly tapers down, indicating a certain cut-off prior to which a power-law behavior is quite evident. Excluding data beyond the cut-off, we estimate the slope of the log-log plot and obtain an estimated slope between 1 and 2. From this we hypothesize that the 2-c-degree of a randomly chosen pair of vertices, D_c , is roughly following

$$\mathbb{P}(D_c = k) \approx k^{-1-a}, \quad a \in (0, 1),$$

for large k , which implies, for t large,

$$t\mathbb{P}\left(\frac{D_c}{t^{1/a}} > x\right) \approx x^{-a}. \quad (3.3)$$

Since $1/a > 1$, then for a randomly chosen pair of vertices (as long as we move away from the cutoff), the growth rate of the common friends is faster than $O(t)$. Note that by the construction of the model, for x, y small, the expected number of common friends between vertices with labels x, y is $t\mu_2(x, y)$, which is growing no faster than $O(t)$. This leads to the cut-off as observed in the figure. Later in Theorem 3.4, we stay away from the cut-off, and give the theoretical analysis on the asymptotic features for $x, y \geq b(t)\epsilon$, with $b(t) \rightarrow \infty$.

Definition 3.3. Define $N_t^\epsilon(k)$ as the number of vertex pairs with second co-ordinate value $\eta > b(t)\epsilon$ and who have $k \geq 1$ common friends in the graph \mathcal{G}_t , i.e.

$$N_t^\epsilon(k) := \sum_{x, y \in \Pi_t^\eta, x, y > b(t)\epsilon} \mathbb{1}_{\{C(x, y) = k\}}, \quad \epsilon > 0.$$

We study the behavior of $N_t^\epsilon(k)$ when W is multivariate regularly varying. When $W \in \mathcal{MRV}(-\alpha, h(t), \omega(x, y), \mathbb{R}_+^2 \setminus \{\mathbf{0}\})$ is non-separable, we choose the scaling function to be $b \in \mathcal{RV}_{1/(2\alpha-1)}$, $\alpha > 2$, such that

$$\lim_{t \rightarrow \infty} t\mu_2(b(t)x, b(t)y) = \int_0^\infty \omega(x, z)\omega(y, z)dz. \quad (3.4)$$

When W is separable, i.e., $W(x, y) = U(x)U(y)$, and $W \in \mathcal{MRV}(-2\alpha, U^2(t), \omega(x, y), (0, \infty)^2)$, we set $b(t) = (1/U)^\leftarrow(\sqrt{t})$, so that $b \in \mathcal{RV}_{1/(2\alpha)}$, $\alpha > 1$ and

$$\mu_2 \in \mathcal{MRV}\left(-2\alpha, b(t), \int_0^\infty \omega(x, z)\omega(y, z)dz, (0, \infty)^2\right).$$

Assume that for $q > 1/2$ and some constant $C_4 > 0$, we have

$$\mu_4(x_1, x_2, x_3, x_4) \leq C_4 (\mu_2(x_1, x_2)\mu_2(x_3, x_4))^q. \quad (3.5)$$

We now give the asymptotic behavior of $\mathbb{E}[N_t^\epsilon(k)]$ for both separable and non-separable cases.

Theorem 3.4. Suppose (i) $W \in \mathcal{MRV}(-\alpha, \mathbb{R}_+^2 \setminus \{\mathbf{0}\})$ and condition (2.6) is satisfied, or, (ii) $W(x, y) = U(x)U(y)$ with $U \in \mathcal{RV}_{-\alpha}$ for $\alpha > 0$. Also assume that (3.5) is satisfied, and that $N_t^\epsilon(k)$ is as defined in Definition 3.3. Let $b(t) \in \mathcal{RV}_{1/\gamma}$ for $\gamma > 1$ and $\lambda(x, y)$ be some functions satisfying

$$\lim_{t \rightarrow \infty} t\mu_2(b(t)x, b(t)y) = \lambda(x, y), \quad x, y > 0. \quad (3.6)$$

Then for $k \geq 1$, we have

$$\frac{N_t^\epsilon(k)}{(tb(t))^2} \xrightarrow{p} \frac{1}{k!} \int_\epsilon^\infty \int_\epsilon^\infty (\lambda(u, v))^k e^{-\lambda(u, v)} du dv. \quad (3.7)$$

Proof. First note that (3.6) is a consequence of either Theorem 2.2 where $\gamma = 2\alpha - 1$, or, Lemma 2.4 where $\gamma = 2\alpha$. From Proposition 3.2, we have for $x, y > 0$, $C(x, y)$ follows a Poisson distribution with rate $t\mu_2(x, y)$. Now by the extended Slivnyak-Mecke theorem given in Theorem 3.1, we see that

$$\begin{aligned}\mathbb{E}[N_t^\epsilon(k)] &= t^2 \int_{b(t)\epsilon}^{\infty} \int_{b(t)\epsilon}^{\infty} \mathbb{P}[C(x, y) = k] dx dy \\ &= t^2 \int_{b(t)\epsilon}^{\infty} \int_{b(t)\epsilon}^{\infty} \frac{(t\mu_2(x, y))^k}{k!} e^{-t\mu_2(x, y)} dx dy \\ &= t^2 b(t)^2 \int_{\epsilon}^{\infty} \int_{\epsilon}^{\infty} \frac{(t\mu_2(b(t)u, b(t)v))^k}{k!} e^{-t\mu_2(b(t)u, b(t)v)} du dv,\end{aligned}\quad (3.8)$$

which, either by using Theorem 2.2, or, Lemma 2.4, converges to the limit on the right hand side of (3.7) when divided by $t^2 b(t)^2$ (as $t \rightarrow \infty$). With (3.8) available, we then show the convergence in (3.7) by proving

$$\text{Var}[N_t^\epsilon(k)] = O\left(t^{-\kappa} \ell(t) \mathbb{E}[N_t^\epsilon(k)]^2\right), \quad (3.9)$$

for some $\kappa > 0$ and a slowly varying function $\ell(t)$. Note that

$$\begin{aligned}(N_t^\epsilon(k))^2 &= N_t^\epsilon(k) + \sum_{x_i \in \Pi_t^\eta, i=1,2,3} \mathbb{1}_{\{C(x_1, x_2)=k, C(x_1, x_3)=k\}} \mathbb{1}_{\{x_1 \neq x_2 \neq x_3\}} \mathbb{1}_{\{x_i > b(t)\epsilon, i=1,2,3\}} \\ &\quad + \sum_{x_i \in \Pi_t^\eta, i=1, \dots, 4} \mathbb{1}_{\{C(x_1, x_2)=k, C(x_3, x_4)=k\}} \mathbb{1}_{\{x_1 \neq x_2 \neq x_3 \neq x_4\}} \mathbb{1}_{\{x_i > b(t)\epsilon, i=1, \dots, 4\}} \\ &=: N_t^\epsilon(k) + A_t^\epsilon(k) + B_t^\epsilon(k).\end{aligned}$$

Since

$$\mu_4(x_1, x_2, x_3, x_4) \leq \frac{1}{2} (\mu_2(x_1, x_2) + \mu_2(x_3, x_4)),$$

then applying Theorem 3.1 and the assumption in (3.5) gives that

$$\begin{aligned}\mathbb{E}[B_t^\epsilon(k)] - (\mathbb{E}[N_t^\epsilon(k)])^2 &\leq t^4 \sum_{l=0}^{k-1} \int_{(b(t)\epsilon, \infty)^4} \frac{(t\mu_2(x_1, x_2))^l}{l!} \frac{(t\mu_2(x_3, x_4))^l}{l!} \frac{(t\mu_4(x_1, x_2, x_3, x_4))^{k-l}}{(k-l)!} \\ &\quad \times e^{-t(\mu_2(x_1, x_2) + \mu_2(x_3, x_4) - \mu_4(x_1, x_2, x_3, x_4))} dx_1 dx_2 dx_3 dx_4 \\ &\leq t^4 \sum_{l=0}^{k-1} \int_{(b(t)\epsilon, \infty)^4} \frac{(t\mu_2(x_1, x_2))^l}{l!} \frac{(t\mu_2(x_3, x_4))^l}{l!} \frac{(t\mu_4(x_1, x_2, x_3, x_4))^{k-l}}{(k-l)!} \\ &\quad \times e^{-t/2(\mu_2(x_1, x_2) + \mu_2(x_3, x_4))} dx_1 dx_2 dx_3 dx_4 \\ &\leq \sum_{l=0}^{k-1} \frac{C_4^{2l} t^{4+k+l}}{(l!)^2 (k-l)!} \left(\int_{(b(t)\epsilon, \infty)^2} (\mu_2(x_1, x_2))^{l+q(k-l)} e^{-t/2\mu_2(x_1, x_2)} dx_1 dx_2 \right)^2,\end{aligned}\quad (3.10)$$

and using the results in (3.8), the upper bound above is of order

$$= O\left(t^{4-(2q-1)}(b(t))^4\right) = O\left(t^{-(2q-1)}(tb(t))^4\right). \quad (3.11)$$

For μ_3 , we have that

$$\begin{aligned}
\mu_3(x_1, x_2, x_3) &= \int_0^\infty \prod_{i=1}^3 W(x_i, z) dz \\
&= \int_0^\infty \left(\sqrt{W(x_1, z)W(x_2, z)} \right) \left(\sqrt{W(x_1, z)W(x_3, z)} \right) dz \\
&\leq \left(\int_0^\infty W(x_1, z)W(x_2, z)^2 dz \right)^{1/2} \left(\int_0^\infty W(x_1, z)W(x_3, z)^2 dz \right)^{1/2} \\
&\leq \sqrt{\mu_2(x_1, x_2)\mu_2(x_1, x_3)}.
\end{aligned}$$

Then, similarly, for $\mathbb{E}[A_t^\epsilon(k)]$, the extended Slivnyak-Mecke theorem gives:

$$\begin{aligned}
\mathbb{E}[A_t^\epsilon(k)] &\leq t^3 \sum_{l=0}^k \int_{(b(t)\epsilon, \infty)^3} \frac{(t\mu_2(x_1, x_2))^l}{l!} \frac{(t\mu_2(x_1, x_3))^l}{l!} \frac{(t\mu_3(x_1, x_2, x_3))^{k-l}}{(k-l)!} \\
&\quad \times e^{-t(\mu_2(x_1, x_2) + \mu_2(x_1, x_3) - \mu_3(x_1, x_2, x_3))} dx_1 dx_2 dx_3 \\
&\leq \sum_{l=0}^k \frac{t^{3+k+l}}{(l!)^2 (k-l)!} \int_{(b(t)\epsilon, \infty)^3} (\mu_2(x_1, x_2))^{l+(k-l)/2} (\mu_2(x_1, x_3))^{l+(k-l)/2} \\
&\quad e^{-t/2(\mu_2(x_1, x_2) + \mu_2(x_1, x_3))} dx_1 dx_2 dx_3,
\end{aligned}$$

and applying the same argument as in (3.8), the bound is of order

$$= O\left(t^3 (b(t))^3\right) = O(t^{-(1+1/\gamma)} \ell(t) (tb(t))^4), \quad (3.12)$$

where $b(t) \in \mathcal{RV}_{1/\gamma}$ and we write $1/b(t) = t^{-(1+1/\gamma)} \ell(t)$ for some slowly varying function $\ell(t)$. Combining (3.12) with (3.11), (3.9) holds with $\kappa \equiv \min(2q-1, 1 + \frac{1}{\gamma}) > 0$. \square

Remark 2. When $W(\cdot, \cdot)$ is separable, the condition in (3.5) holds with $q = 1$. Then the upper bounds in (3.10) and (3.12) are of orders

$$O\left(t^{2/\alpha+3} (\ell_1(t))^4\right), \quad O\left(t^{3/(2\alpha)+3} (\ell_1(t))^3\right),$$

respectively, where $\ell_1(t)$ is a slowly varying function such that $b(t) = t^{1/(2\alpha)} \ell_1(t)$. Therefore, the convergence results in Theorem 3.4 hold for $W(\cdot, \cdot)$ separable.

3.2.1 Tail asymptotics for common connections

Note that the convergence results in Theorem 3.4 do not provide an explicit explanation for the power-law behavior as observed in Figure 3.1. Hence, we proceed with a detailed discussion on the tail distribution of 2-c-degrees. Consider the function

$$f(x) = x^k e^{-x}, \quad x \in A,$$

and we see that f is increasing in x if $k \geq \sup_{x \in A} x$.

Separable case. In the separable case, assume in addition that $U(x)U(y) \leq U(x+y)$, $x, y \geq 0$, then the integrand in (3.8) satisfies

$$(t\mu_2(b(t)x, b(t)y))^k e^{-t\mu_2(b(t)x, b(t)y)} \leq (C \cdot tU(b(t)(x+y)))^k e^{-C \cdot tU(b(t)(x+y))},$$

with $C = \int_0^\infty U^2(z)dz$, as long as $k \geq \sup_{x,y \geq b(t)\epsilon} t\mu_2(b(t)x, b(t)y) = \lambda(\epsilon, \epsilon)$. Therefore, for $k \geq \lambda(\epsilon, \epsilon)$, we have

$$\begin{aligned} & \frac{1}{k!} \int_\epsilon^\infty \int_\epsilon^\infty (t\mu_2(b(t)x, b(t)y))^k e^{-t\mu_2(b(t)x, b(t)y)} dx dy \\ & \leq \frac{1}{k!} \int_\epsilon^\infty \int_\epsilon^\infty (C \cdot tU(b(t)(x+y)))^k e^{-C \cdot tU(b(t)(x+y))} dx dy, \end{aligned} \quad (3.13)$$

and since $b(t) = (1/U)^\leftarrow(\sqrt{t})$ in the separable case, then for some slowly varying function ℓ , we have

$$tU(b(t)(x+y)) = t^{1/2}(x+y)^{-\alpha} \frac{\ell(b(t)(x+y))}{\ell(b(t))}.$$

Assume further that

$$\lim_{t \rightarrow \infty} \ell(t) = C_0 > 0, \quad (3.14)$$

then the upper bound in (3.13) is of order $O(t^{1/\alpha} k^{-1-2/\alpha})$. By (3.3), we may write $k = O(t^{1/a})$, $a \in (0, 1)$, then the upper bound is of order

$$k^{-\frac{1}{\alpha}(2-a)-1},$$

and $\frac{1}{\alpha}(2-a) + 1 \in (1 + 1/\alpha, 1 + 2/\alpha)$.

Next, we give a lower bound for the asymptotic power-law behavior of the c-degrees in the separable case. Since

$$\mu_2(x, y) = CU(x)U(y) \leq \frac{C}{2}(U^2(x) + U^2(y)),$$

then we have for $b \in \mathcal{RV}_{1/(2\alpha)}$,

$$\begin{aligned} & \frac{b^2(t)}{k!} \int_\epsilon^\infty \int_\epsilon^\infty (t\mu_2(x, y))^k e^{-t\mu_2(x, y)} dx dy \\ & \geq \frac{1}{k!} \int_\epsilon^\infty \int_\epsilon^\infty [tCU(b(t)x)U(b(t)y)]^k e^{-\frac{tC}{2}(U^2(b(t)x) + U^2(b(t)y))} dx dy \\ & = C^k \frac{t^k}{k!} \left[\int_0^\infty U^k(b(t)x) e^{-\frac{tC}{2}U^2(b(t)x)} dx - \int_0^\epsilon U^k(b(t)x) e^{-\frac{tC}{2}U^2(b(t)x)} dx \right]^2. \end{aligned}$$

Using [Caron et al., 2020, Lemma S3.5], since $U^2(x) \in \mathcal{RV}_{-2\alpha}$,

$$\int_0^\infty U^k(x) e^{-\frac{tC}{2}U^2(x)} dx \sim \frac{1}{2\alpha} \left(\frac{tC}{2} \right)^{\frac{1}{2\alpha} - \frac{k}{2}} \Gamma\left(\frac{k}{2} - \frac{1}{2\alpha}\right) \ell_1(t),$$

for some slowly varying function $\ell_1(t)$. Note that when ϵ is small,

$$\int_0^\epsilon \left(t^{1/2} U(b(t)x) \right)^k e^{-\frac{tC}{2}U^2(b(t)x)} dx \approx \int_0^\epsilon x^{-\alpha k} e^{-\frac{C}{2}x^{-2\alpha}} dx,$$

which is also small. Therefore, using Stirling's approximation,

$$\begin{aligned}
& \frac{1}{k!} \int_{b(t)\epsilon}^{\infty} \int_{b(t)\epsilon}^{\infty} (t\mu_2(x, y))^k e^{-t\mu_2(x, y)} dx dy \\
& \gtrsim C^k \frac{t^k}{k!} (2\alpha)^{-2} \left(\frac{tC}{2} \right)^{\frac{1}{\alpha}-k} \left(\Gamma \left(\frac{k}{2} - \frac{1}{2\alpha} \right) \right)^2 (\ell_1(t))^2 \\
& = \frac{C^{1/\alpha}}{2^{2+1/\alpha-k} \alpha^2} \cdot t^{\frac{1}{\alpha}} \cdot \left(\frac{\Gamma(\frac{k}{2} - \frac{1}{2\alpha})}{\Gamma(\frac{k}{2} + 1)} \right)^2 \cdot \left(\frac{\Gamma(\frac{k}{2} + 1)}{\Gamma(k+1)} \right)^2 (\ell_1(t))^2 \\
& \approx \frac{C^{1/\alpha}}{2^{2+1/\alpha-k} \alpha^2} \cdot t^{2+\frac{1}{\alpha}} \cdot \left(\frac{k}{2} \right)^{-\frac{1}{\alpha}-2} \cdot \frac{\sqrt{2\pi k}}{2^{k+1}} (\ell_1(t))^2 \\
& = O \left(t^{\frac{1}{\alpha}} k^{-\left(\frac{1}{\alpha} + \frac{3}{2}\right)} \right), \tag{3.15}
\end{aligned}$$

if we assume (3.14). Hence for small ϵ and k is large, (3.15) provides a lower bound for the asymptotic power-law behavior of the 2-c-degree distribution, i.e., for large t ,

$$\frac{N_t^\epsilon(k)}{t^2 b(t)^2} \gtrsim O \left(k^{-\left(\frac{1}{\alpha} + \frac{3}{2}\right)} \right) \quad (\text{in probability}).$$

Non-separable case. In the non-separable case, we need the following technical assumption in order to further study the limit behavior of $\mathbb{E}[N_t^\epsilon(k)]/(tb(t))^2$. For $(x, y) \in (0, \infty)^2$, assume there exist constants $C_0 > 0$ and $x_0 \geq 0$ such that for all $x, y \geq x_0$,

$$W(x, y) \leq C_0 (\mu_1(x))^\theta (\mu_1(y))^\theta, \quad \mu_1(x_0) > 0, \tag{3.16}$$

where $\theta > 1/2$ is a positive constant. As discussed in Caron et al. [2020], the assumption in (3.16) is satisfied with $\theta = 0$ when the function W is separable. Another example is

$$W(x, y) = (1 + x + y)^{-\alpha}, \quad \alpha > 2, \tag{3.17}$$

then we have

$$\mu_1(x) = \frac{1}{\alpha-1} (1+x)^{-(\alpha-1)}, \quad W(x, y) \leq (\alpha-1)^{\frac{\alpha}{\alpha-1}} (\mu_1(x)\mu_1(y))^{\frac{\alpha}{2(\alpha-1)}}.$$

In fact, the assumption in (3.16) implies for $x, y > 0$,

$$\begin{aligned}
\mu_2(x, y) &= \int_0^\infty W(x, z) W(y, z) dz \\
&\leq C_0^2 \int_0^\infty \mu_1(z)^{2\theta} dz \cdot (\mu_1(x)\mu_1(y))^\theta =: C' (\mu_1(x)\mu_1(y))^\theta.
\end{aligned}$$

Let $b \in \mathcal{RV}_{1/(2\alpha-1)}$ be some scaling function such that (3.4) holds. Assume further that $\mu_1(x)\mu_1(y) \leq \mu_1(x+y)$ then we have for $x, y \geq \epsilon > 0$,

$$t\mu_2(b(t)x, b(t)y) \leq t (\mu_1(b(t)(x+y)))^\theta = t^{1-\frac{\theta(\alpha-1)}{2\alpha-1}} (x+y)^{-(\alpha-1)} \frac{\ell_0(b(t)(x+y))}{(\ell_b(t))^{\alpha-1}},$$

where $\ell_0(\cdot)$, $\ell_b(\cdot)$ are two slowly varying functions satisfying

$$\mu_1(x) = x^{-(\alpha-1)} \ell_0(x), \quad b(t) = t^{1/(2\alpha-1)} \ell_b(t).$$

Suppose also both ℓ_0 and ℓ_b satisfy (3.14). Then repeating the reasoning in the separable case, we see that the upper bound for

$$\frac{1}{k!} \int_{\epsilon}^{\infty} \int_{\epsilon}^{\infty} (t\mu_2(b(t)x, b(t)y))^k e^{-t\mu_2(b(t)x, b(t)y)} dx dy$$

is of order

$$t^{\frac{2}{\theta(\alpha-1)} - \frac{2}{2\alpha-1}} k^{-1 - \frac{2}{\theta(\alpha-1)}} = t^{\frac{4}{\alpha} - \frac{2}{2\alpha-1}} k^{-1 - \frac{4}{\alpha}}.$$

Recall (3.3), and we may write $k = O(t^{1/a})$, with $a \in (0, 1)$, then the upper bound is of order

$$k^{-1 - \frac{4}{\alpha}(1-a) - \frac{2a}{2\alpha-1}},$$

and $1 + \frac{4}{\alpha}(1-a) + \frac{2a}{2\alpha-1} \in (1 + 2/(2\alpha-1), 1 + 4/\alpha)$. Note that for the example in (3.17), we have $\ell_0, \ell_b \equiv 1$ and $\theta = \frac{\alpha}{2(\alpha-1)}$, which satisfy all those conditions listed above.

4 Simulation Studies

To see how tight the bounds derived in the previous section are, especially when the entire network is considered (e.g. the scenario in Figure 3.1), we further examine the slope estimates through simulation studies. We first choose two different sets of values for α :

- (1) Separable case: $W(x, y) = (1+x)^{-\alpha}(1+y)^{-\alpha}$, $\alpha \in \{1.5, 2, 3, 4, 5\}$.
- (2) Non-separable case: $W(x, y) = (1+x+y)^{-\alpha}$, $\alpha \in \{2, 2.6, 3, 4, 5\}$.

For each combination of $W(\cdot, \cdot)$ and α , we simulate 500 replications with $t = 1000$. Here all simulated replications have a similar distribution shape as depicted in Figure 3.1. In other words, when plotted on a log-log scale, the empirical distribution shows a linearly decaying pattern as long as we stay away from the cutoff. Therefore, we estimate the power-law index by the absolute value of the slope estimate from a simple linear regression. To avoid the cutoff area, we choose a proportion of the empirical distribution such that the fitted regression returns an $R^2 \geq 99.5\%$. This is done by first fitting a linear regression to the entire empirical distribution, then leaving out observations in the cutoff area until the target R^2 is achieved.

Following the method described above, we collect all tail index estimates (slope estimates) for the distribution of common connections, and plot them as boxplots in Figure 4.1. The left panel contains all simulated separable examples, i.e. $W(x, y) = (1+x)^{-\alpha}(1+y)^{-\alpha}$, $\alpha \in \{1.5, 2, 3, 4, 5\}$, whereas all non-separable examples, i.e. $W(x, y) = (1+x+y)^{-\alpha}$, $\alpha \in \{2, 2.6, 3, 4, 5\}$, are given in the right panel of Figure 4.1. Both separable and non-separable cases reveal the same pattern that when $W(\cdot, \cdot)$ decays at a faster rate or α is large, the distribution of common connections will have a heavier tail. Similar pattern is also observed from the empirical degree distribution as given in Caron and Fox [2017], Caron et al. [2020], Veitch and Roy [2015].

To assess the tightness of bounds provided in the previous section, we summarize the mean, variance and range based on the 500 replications for each combination of $W(\cdot, \cdot)$ and α . For the separable case, we compute the *coverage proportion* for the derived upper and lower bounds for the tail index:

$$\left[1 + \frac{1}{\alpha}, \max\left(\frac{3}{2} + \frac{1}{\alpha}, 1 + \frac{2}{\alpha}\right)\right], \quad (4.1)$$




Figure 4.1: Boxplots of estimated tail indices for the distribution of common connections based on simulated sparse networks with graphex function $W(x, y)$. Left: $W(x, y) = (1 + x)^{-\alpha}(1 + y)^{-\alpha}$, $\alpha \in \{1.5, 2, 3, 4, 5\}$. Right: $W(x, y) = (1 + x + y)^{-\alpha}$, $\alpha \in \{2, 2.6, 3, 4, 5\}$.

i.e., the proportion of estimated tail indices that are within the interval in (4.1). Numerical results are given in Table 4.1. Overall, the mean and standard deviation are both decreasing as α increases, and the coverage proportion remains high, as long as α is neither too small nor too large. This indicates that the bounds indicated in (4.1) perform well in these examples, even though we now take all $(x, y) \in \Pi_t^\eta$ into consideration.

For the non-separable case, we examine the tightness of the derived upper bound for the tail index, i.e.,

$$\left(1 + \frac{2}{(2\alpha - 1)}, 1 + \frac{4}{\alpha}\right), \quad (4.2)$$

in Table 4.2. Similar to the separable case, both mean and standard deviation are decreasing as α increases. Except the $\alpha = 2$ case, all other coverage proportions reach 100%. The relative poor coverage proportion for $\alpha = 2$ may be due to the fact that $\int_0^\infty \mu_1(x)dx = \infty$ when $\alpha = 2$.

Comparing Tables 4.1 and 4.2, we see that for the same α , a separable $W(\cdot, \cdot)$ tends to generate a heavier tail for the common connection distribution than that in a non-separable $W(\cdot, \cdot)$ case. Note that although in both cases $W \in \mathcal{MRV}(-\alpha, \mathbb{R}_+^2 \setminus \{\mathbf{0}\})$, these two models tend to generate structures which are quite different, for example, we obtain different behavior of graphex marginals μ_1 and μ_2 in the two cases. Using Theorem 2.2, for the non-separable W we have $\mu_1 \in \mathcal{RV}_{-(\alpha-1)}$. On the other hand, for the separable case, $W \in \mathcal{MRV}(-\alpha, \mathbb{R}_+^2 \setminus \{\mathbf{0}\})$ and moreover $W \in \mathcal{MRV}(-2\alpha, (0, \infty)^2)$, and from Lemma 2.4 we have $\mu_1 \in \mathcal{RV}_{-\alpha}$. In order to compare models with similar average connection behaviors, we tabulate in Table 4.3 the means and ranges of the estimated slope, where the μ_1 tail indices are the same in the two models. Even here we observe that the common connection distribution tends to have heavier tails in the separable case. A similar comparison matching μ_2 indices in the separable ($\alpha = 1.5$ with μ_2 index $2\alpha = 3$) and non-separable ($\alpha = 2$ with μ_2 index $2\alpha - 1 = 3$) cases also leads to observing

a heavier tail behavior in the former.

α	Mean	Std Dev	Range	Coverage (%)
1.5	2.095	0.052	[1.980, 2.193]	88.8
2	1.769	0.056	[1.689, 1.879]	100
3	1.438	0.039	[1.366, 1.508]	100
4	1.322	0.041	[1.249, 1.394]	99
5	1.238	0.033	[1.182, 1.308]	84.8

Table 4.1: Mean, standard deviation, and range based on the 500 replications for each combination of $W(\cdot, \cdot)$ and α in the separable case. The coverage proportion of the upper and lower bounds in (4.1) is given in the last column.

α	Mean	Std Dev	Range	Coverage (%)
2	2.228	0.045	[2.140, 2.310]	81.4
2.6	2.121	0.049	[2.022, 2.210]	100
3	2.081	0.041	[1.992, 2.150]	100
4	1.878	0.040	[1.781, 1.950]	100
5	1.754	0.045	[1.670, 1.829]	100

Table 4.2: Mean, standard deviation, and range based on the 500 replications for each combination of $W(\cdot, \cdot)$ and α in the non-separable case. The coverage proportion of the upper bound is given in the last column.

Index of μ_1	Separable model			Non-separable model		
	α	Mean	Range	α	Mean	Range
2	2	1.769	[1.689, 1.879]	3	2.081	[1.992, 2.150]
3	3	1.438	[1.366, 1.508]	4	1.878	[1.781, 1.950]
4	4	1.322	[1.249, 1.394]	5	1.754	[1.670, 1.829]

Table 4.3: The α value, mean and range based on the 500 replications for each combination of $W(\cdot, \cdot)$ and α in both the non-separable case and separable cases where the tail indices of the univariate graphex marginal μ_1 are the same.

5 Conclusion

In this paper, we work under the paradigm of sparse exchangeable graphs generated by multivariate regularly varying graphex functions and investigate the distributional behavior of common connections between pairs of vertices. We conclude that the asymptotic distribution of number of common connections of randomly chosen pair of vertices, given that the vertices chosen are above an (increasing) threshold also has a power-law behavior under certain regularity conditions. We distinguish between two types of generating graphex functions here, separable and non-separable, each of which leads to different asymptotic tail behavior. We manage to derive bounds for the tail behavior but not the exact rate, and our simulation studies verify the bounds obtained.

One future direction would be to find appropriate minimal conditions on the graphex function, so that we obtain tighter bounds for the tail rate of the distribution of number of common connections of randomly chosen pair of vertices. Network scientists have found this measure a good indicator for link-prediction between vertices. We believe understanding 2-c-degree functions, and possibly k -c-degree functions will also lead us towards a generating mechanism for networks like Facebook and LinkedIn, where connections are often made through friendship recommendation.

Data Availability Statement

The datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request.

References

- D. J. Aldous. Representations for partially exchangeable arrays of random variables. *Journal of Multivariate Analysis*, 11(4):581–598, 1981.
- A. Barabási, H. Jeong, Z. Néda, E. Ravasz, A. Schubert, and T. Vicsek. Evolution of the social network of scientific collaborations. *Physica A: Statistical mechanics and its applications*, 311(3):590–614, 2002.
- N. H. Bingham, C. M. Goldie, and J. L. Teugels. *Regular variation*, volume 27 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1989.
- M. Bloznelis and V. Kurauskas. Clustering function: another view on clustering coefficient. *Journal of Complex Networks*, 4(1):61–86, 2016.
- C. Borgs, J. T. Chayes, H. Cohn, and N. Holden. Sparse exchangeable graphs and their limits via graphon processes. *Journal of Machine Learning Research*, 18(210):1–71, 2018.
- C. Borgs, J. T. Chayes, H. Cohn, and V. Veitch. Sampling perspectives on sparse exchangeable graphs. *The Annals of Probability*, 47(5):2754–2800, 2019.
- F. Caron and E. B. Fox. Sparse graphs using exchangeable random measures. *J. R. Stat. Soc. Ser. B. Stat. Methodol.*, 79(5):1295–1366, 2017.
- F. Caron, F. Panero, and J. Rousseau. On sparsity, power-law and clustering properties of graphex processes. *Arxiv preprint*, 2020. URL <https://arxiv.org/pdf/1708.03120>.
- B. Das and S. Ghosh. Growth of common friends in a preferential attachment model. *Stochastic Models*, forthcoming, 2021. doi: 10.1080/15326349.2021.1910521.
- L. de Haan and S. I. Resnick. On regular variation of probability densities. *Stochastic Processes and their Applications*, 25:83 – 93, 1987.
- P. Diaconis and S. Janson. Graph limits and exchangeable random graphs. *Rend. Mat. Appl. (7)*, 28(1):33–61, 2008.

- I. Gradshteyn and I. Ryzhik. *Table of Integrals, series, and products*. Academic Press, 7th edition, 2007.
- P. Gupta, A. Goel, J. Lin, A. Sharma, D. Wang, and R. Zadeh. WTF: The who to follow service at twitter. In *Proceedings of the 22Nd International Conference on World Wide Web, WWW '13*, pages 505–514, New York, NY, USA, 2013. ACM.
- T. Herlau, M. N. Schmidt, and M. Mørup. Completely random measures for modelling block-structured sparse networks. In D. Lee, M. Sugiyama, U. Luxburg, I. Guyon, and R. Garnett, editors, *Advances in Neural Information Processing Systems*, volume 29, pages 4260–4268. Curran Associates, Inc., 2016.
- D. N. Hoover. Relations on probability spaces and arrays of random variables. *preprint*, 1979. URL <https://www.stat.berkeley.edu/~aldous/Research/hoover.pdf>.
- S. Janson. Graphons and cut metric on sigma-finite measure spaces. *Arxiv preprint*, 2016. URL <https://arxiv.org/pdf/1608.01833.pdf>.
- S. Janson. On convergence for graphexes. *Arxiv preprint*, 2017. URL <https://arxiv.org/pdf/1702.06389.pdf>.
- O. Kallenberg. Exchangeable random measures in the plane. *Journal of Theoretical Probability*, 3(1):81–136, 1990.
- J. F. C. Kingman. *Poisson Processes*. Oxford Studies in Probability. Clarendon Press, 1 edition, 1993.
- D. Liben-Nowell and J. Kleinberg. The link-prediction problem for social networks. *Journal of the American Society for Information Science and Technology*, 58(7):1019–1031, 2007.
- L. Lovász and B. Szegedy. Limits of dense graph sequences. *J. Combin. Theory Ser. B*, 96(6):933–957, 2006.
- J. Møller and R. P. Waagepetersen. *Statistical inference and simulation for spatial point processes*. CRC Press, 2003.
- Z. Naulet, D. M. Roy, E. Sharma, and V. Veitch. Bootstrap estimators for the tail-index and for the count statistics of graphex processes. *Electronic Journal of Statistics*, 15(1):282 – 325, 2021.
- M. E. J. Newman. Clustering and preferential attachment in growing networks. *Physical review E*, 64(2):025102, 2001.
- S. Resnick. *Heavy Tail Phenomena: Probabilistic and Statistical Modeling*. Springer-Verlag, New York, 2007.
- A. Stam. Regular variation in \mathbb{R}_+^d and the Abel-Tauber theorem. Technical Report, unpublished, Mathematisch Instituut, Rijksuniversiteit Groningen, August 1977.
- A. Todeschini, X. Miskouridou, and F. Caron. Exchangeable random measures for sparse and modular graphs with overlapping communities. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 82(2):487–520, 2020.

- V. Veitch and D. M. Roy. The class of random graphs arising from exchangeable random measures. *Arxiv preprint*, 2015. URL <https://arxiv.org/pdf/1512.03099.pdf>.
- V. Veitch and D. M. Roy. Sampling and estimation for (sparse) exchangeable graphs. *The Annals of Statistics*, 47(6):3274–3299, 2019.
- I. Voitalov, P. van der Hoorn, R. van der Hofstad, and D. Krioukov. Scale-free networks well done. *Phys. Rev. Research*, 1(033034), Oct 2019.