

# Spectral dimensions of Kreĭn-Feller operators and $L^q$ -spectra

Marc Kesseböhmer<sup>1,\*</sup>, Aljoscha Niemann\*

*Fachbereich 3 – Mathematik und Informatik, University of Bremen, Bibliothekstr. 1, 28359 Bremen, Germany*

---

## Abstract

We study the spectral dimensions and spectral asymptotics of Kreĭn-Feller operators for arbitrary finite Borel measures on  $(0, 1)$ . Connections between the spectral dimension, the  $L^q$ -spectrum, the partition entropy and the optimized coarse multifractal dimension are established. In particular, we show that the upper spectral dimension always corresponds to the fixed point of the  $L^q$ -spectrum of the corresponding measure. Natural bounds reveal intrinsic connections to the Minkowski dimension of the support of the associated Borel measure. Further, we give a sufficient condition on the  $L^q$ -spectrum to guarantee the existence of the spectral dimension. As an application, we confirm the existence of the spectral dimension of self-conformal measures with or without overlap as well as of certain measures of pure point type. We construct a simple example for which the spectral dimension does not exist and determine explicitly its upper and lower spectral dimension.

*Keywords:* Kreĭn-Feller operator, spectral asymptotics,  $L^q$ -spectrum, Dirichlet forms, Minkowski dimension, coarse multifractal formalism

*2020 MSC:* 35P20; 35J05; 28A80; 42B35; 45D05

---

## Contents

<b>1</b>	<b>Introduction and statement of main results</b>	<b>2</b>
<b>2</b>	<b>Spectral theory for Kreĭn-Feller operators</b>	<b>9</b>
2.1	Form approach . . . . .	9
2.2	Spectral problem, min-max principle, and sub- and superadditivity . . .	13
2.3	Neumann boundary conditions . . . . .	18
<b>3</b>	<b>Optimal partitions and partition entropy</b>	<b>19</b>
3.1	Basic observations . . . . .	19

---

\*Corresponding authors

*Email addresses:* [mhk@uni-bremen.de](mailto:mhk@uni-bremen.de) (Marc Kesseböhmer), [niemann1@uni-bremen.de](mailto:niemann1@uni-bremen.de) (Aljoscha Niemann)

<sup>1</sup>This research was supported by the DFG grant Ke 1440/3-1.

3.2	Some global bounds . . . . .	20
<b>4</b>	<b><math>L^q</math>-spectrum and coarse multifractal formalism</b>	<b>22</b>
4.1	The $L^q$ -spectrum . . . . .	23
4.2	Upper bounds in terms of the $L^q$ -spectrum . . . . .	27
4.3	Lower bounds in terms of the optimize coarse multifractal dimension . . . . .	29
4.4	Proof of main results . . . . .	33
<b>5</b>	<b>Examples</b>	<b>35</b>
5.1	$C^1$ -self-conformal measures . . . . .	35
5.2	Homogeneous Cantor measures . . . . .	37
5.3	Pure point measures . . . . .	40

## 1. Introduction and statement of main results

This article is devoted to the subtle connections between Kreĭn-Feller operators, geometric measure theory, and concepts of fractality and complexity and thus gives new partial answers to Mark Kac’s famous question “*Can One Hear the Shape of a Drum?*”.

We investigate the spectral properties of the Kreĭn-Feller operator  $\Delta_{\nu,[a,b]}$  for a given finite Borel measure  $\nu$  on the interval  $(a, b)$ . The Kreĭn-Feller operator has been introduced in [Fel57, KK58, Kre51]. We will elaborate the abstract approach via the (Dirichlet) form  $\mathcal{E}_{\nu,[a,b]}(f, g) := \int_a^b \nabla_{\Lambda} f \nabla_{\Lambda} g \, d\Lambda$  (with  $\Lambda$  denoting the Lebesgue measure) densely defined on  $L^2_{\nu} = L^2_{\nu}([a, b])$  as proposed e. g. in [KKVW09, BNT03, HLN06]. This approach gives rise to an associated Laplace operator  $\Delta_{\nu,[a,b]}$  and an orthonormal basis of eigenfunctions with non-negative eigenvalues  $(\lambda^n(\mathcal{E}_{\nu,[a,b]}))_{n \in \mathbb{N}}$  tending to  $\infty$ . As a consequence of the Dirichlet/Neumann bracketing (Prop. 2.18) we see that the asymptotic behavior of the eigenvalues does not depend on the chosen boundary condition nor on the interval containing  $\text{supp } \nu$  and we state the corresponding results neither referring to boundary conditions nor to the ambient interval. We denote the number of eigenvalues of  $\mathcal{E}_{\nu,[a,b]}$  not exceeding  $x \geq 0$  by  $N_{\nu,[a,b]}(x)$  and refer to  $N_{\nu,[a,b]}$  as the *eigenvalue counting function*. If the ambient interval is clear from the context and  $\text{supp } \nu \subset [a, b]$ , we will also suppress the dependence on the interval, i. e. we write  $\mathcal{E}_{\nu} = \mathcal{E}_{\nu,[a,b]}$ ,  $N_{\nu} = N_{\nu,[a,b]}$ ,  $\Delta_{\nu} = \Delta_{\nu,[a,b]}$ ,  $\lambda_{\nu}^n = \lambda^n(\mathcal{E}_{\nu,[a,b]})$  etc. Using some invariance properties and bi-Lipschitz equivalence we can assume without loss of generality that the ambient interval is given by  $[0, 1]$  and we will assume this throughout the paper. In this setup we give rigorous proofs for the sub- and superadditivity of the eigenvalue counting function which is crucial in our analysis. We define the upper and lower exponent of divergence of  $N_{\nu}$  by

$$s_{\nu} := \liminf_{x \rightarrow \infty} \frac{\log(N_{\nu}(x))}{\log(x)} \quad \text{and} \quad \bar{s}_{\nu} := \limsup_{x \rightarrow \infty} \frac{\log(N_{\nu}(x))}{\log(x)}$$

and refer to these numbers as the *upper*, resp. *lower*, *spectral dimension* of  $\mathcal{E}_{\nu}$ ,  $\Delta_{\nu}$  or, just  $\nu$ , respectively. If the two values coincide we denote the common value by  $s_{\nu}$  and call it the *spectral dimension*. Note that we always have  $\bar{s}_{\nu} \leq 1/2$  which has been shown in [BS66, BS67]. The case for measures with non-trivial absolutely continuous part was

completely solved in an elegant way in [BS70] using a variational approach. In this case, for a Borel measure  $\nu$  on the unit interval with absolutely continuous non-vanishing part  $\sigma\Lambda$  and singular part  $\eta$ , we have

$$\lim_{x \rightarrow \infty} \frac{N_{\eta+\sigma\Lambda}(x)}{x^{1/2}} = \frac{1}{\pi} \int \sqrt{\sigma} d\Lambda$$

and particularly the spectral dimension exists and equals  $1/2$ . Besides this estimate, many partial results have been obtained showing that there is a subtle relation between spectral properties and geometric data of  $\nu$ , which has been a major line of investigation since the famous result by Hermann Weyl [Wey1]. Weyl's result includes the  $d$ -dimensional Lebesgue case with spectral dimension  $d/2$ ; for this reason, some authors prefer to set the spectral dimension to twice the value defined here.

We will make use of the following notation. For any two functions  $f, g : \mathbb{R}_{\geq 0} \rightarrow \mathbb{R}_{> 0}$  we write  $f \ll g$  if there exists a positive constant  $c$  such that  $cf(x) \leq g(x)$  for all  $x$  large; we write  $f \asymp g$ , if both  $f \ll g$  and  $g \ll f$  holds. The asymptotic behavior of  $N_\nu$  strongly depends on the measure  $\nu$ . In general, one can not expect that  $N_\nu$  obeys a power law with a positive exponent  $s > 0$ , i. e.  $N_\nu(x) \asymp x^s$  (for counter examples see e. g. [Arz14] or Examples 5.2 and 5.9). However, if we restrict our attention to the spectral dimension this connection becomes more feasible and surprisingly, we are able to treat arbitrary Borel measures on  $(0, 1)$  and determine the upper spectral dimension solely from the data provided by the measure-geometric information carried by the  $L^q$ -spectrum of  $\nu$ . Under mild regularity conditions on the measure we can guarantee the existence of the spectral dimension. Also with the help of the  $L^q$ -spectrum we are able to construct first examples for which the spectral dimension does not exist. In this way we give a partial answer to Kac's question in terms of the spectral dimension revealing how the measure theoretic properties of  $\nu$  and the topological properties of its "fractal" support are interwoven to determine the spectral dimension (see Theorem 1.1 and Corollary 1.7). This striking connection will be the main line of investigation in this paper.

Our approach lies in the construction of optimal partitions as introduced in Section 3 and to make use of the min-max principle. This idea goes back to the fundamental works of [BS67, BS74, BS70] which deal with a systematic treatment of Sobolev embedding theorems in tandem with variational principles. The following observation motivates the definition of optimal  $\nu$ -partitions. Due to Proposition 2.18, it is sufficient to consider  $\Delta_\nu$  under Dirichlet boundary conditions and we will therefore mainly restrict ourselves to this case. The first key property to determine the spectral dimension is the sub- and superadditivity of the eigenvalue counting function with respect to interval partitions. Let  $P$  be a  $\nu$ -partition of  $[0, 1]$ , that is a partition with half open intervals  $\{(c_k, c_{k+1}] : k = 1, \dots, n-1\}$ ,  $n \in \mathbb{N}_{\geq 2}$ , such that  $0 = c_1 < c_2 < \dots < c_n = 1$  and there are no  $\nu$ -atoms on the endpoints  $\nu(\{c_k\}) = 0$ . Then by Proposition 2.16, we have

$$\sum_{k=1}^n N_{\nu, [c_k, c_{k+1}]}(x) \leq N_\nu(x) \leq \sum_{k=1}^n N_{\nu, [c_k, c_{k+1}]}(x) + n - 2.$$

The second key observation is the Poincaré inequality, i. e. the first eigenvalue  $\lambda_{\nu, [a, b]}^1$  of  $\Delta_{\nu, [a, b]}$  can be bounded from below by  $(\Lambda([a, b])\nu([a, b]))^{-1}$  (see Lemma 3.5). In

particular, if  $\Lambda([a, b])\nu([a, b]) < 1/x$ , then  $N_{\nu, [a, b]}(x) = 0$ . For a  $\nu$ -partition  $P$  of  $[0, 1]$  such that  $\Lambda(C)\nu(C) < 1/x$  for all  $C \in P$  and some  $x > 0$ , the superadditivity implies

$$N_{\nu}(x) \leq \text{card}(P).$$

Similar ideas will be employed to derive appropriate lower bounds. These observations lead to the following definitions. Let us consider the set  $\Pi$  of all finite  $\nu$ -partitions of  $[0, 1]$  and set

$$\mathcal{N}^R(x) := \inf \left\{ \text{card}(P) : P \in \Pi : \max_{C \in P} \nu(C) \Lambda(C) < 1/x \right\}.$$

The following quantities

$$\bar{h}_{\nu} := \limsup_{x \rightarrow \infty} \frac{\log \mathcal{N}^R(x)}{\log x}, \quad \underline{h}_{\nu} := \liminf_{x \rightarrow \infty} \frac{\log \mathcal{N}^R(x)}{\log x},$$

will be called the *upper*, resp. *lower*,  $\nu$ -partition entropy. These quantities have interesting connections to the classical work of Birman and Solomjak [BS66, BS67, Bor70] as explained in Section 3.1. Note that ignoring intervals of measure zero in the definition of the  $\nu$ -partition will not change the value of the  $\nu$ -partition entropy.

For effective lower bounds, we need the following variant of this notion. Let  $\Pi_0$  denote the set of finite disjoint collections of subintervals of  $I$  such that there are no atoms on the endpoints of each interval, and for  $m > 1$ , set

$$\mathcal{N}_m^L(x) := \sup \left\{ \text{card}(P) : P \in \Pi_0 : \min_{C \in P} \nu(\langle I \rangle_m) \Lambda(\langle I \rangle_m) \geq \frac{4}{x(m-1)} \right\},$$

where  $\langle I \rangle_m \subset I$  denotes the interval of length  $\Lambda(I)/m$  centered in  $I$ . Then the *lower* and *upper  $m$ -reduced  $\nu$ -partition entropy* is given by

$$\underline{h}_m^L := \liminf_{x \rightarrow \infty} \frac{\log \mathcal{N}_m^L(x)}{\log x} \quad \text{and} \quad \bar{h}_m^L := \limsup_{x \rightarrow \infty} \frac{\log \mathcal{N}_m^L(x)}{\log x}.$$

The second important object depends on dyadic partitions of the unit interval and gives rise to the notion of the  $L^q$ -spectrum of  $\nu$  given by

$$\beta_{\nu}(q) := \limsup_{n \rightarrow \infty} \beta_n^{\nu}(q) \quad \text{with} \quad \beta_n^{\nu}(q) := \frac{1}{\log 2^n} \log \sum_{C \in \mathcal{D}_{\nu, n}} \nu(C)^q,$$

where  $\mathcal{D}_{\nu, n} := \{A_n^k : k \in \{1, \dots, 2^n\}, \nu(A_n^k) > 0\}$  and  $A_n^k := ((k-1)2^{-n}, k2^{-n}]$  and  $q \in \mathbb{R}$ . Note that this notion has recently gained a lot of attention in the context of fractal geometry in the work of [Hoc14, Shm19] – connections to our work on spectral dimensions will be provided in the forthcoming paper [KN21b]. Each  $\beta_n^{\nu}$  is differentiable and convex with  $\beta_n^{\nu}(1) = 0$  and there exists a unique fixed point  $q_n$ , i. e.  $\beta_n^{\nu}(q_n) = q_n$ . Also  $\beta_{\nu}(0) =: \delta_{\nu}^* = \overline{\dim}_M(\text{supp } \nu)$ , where  $\overline{\dim}_M(A)$  denotes the upper Minkowski dimension of  $A \subset \mathbb{R}$ . As a shorthand notation we will also write  $\overline{\dim}_M(\nu)$  instead of  $\overline{\dim}_M(\text{supp } \nu)$ . It will turn out (see Theorem 1.1) that

$$\bar{q}_{\nu} := \limsup q_n$$

completely determines the upper spectral dimension. We also need the following left and right derivatives of  $\beta_\nu$  in 1:

$$\bar{\delta}_\nu := \lim_{t \nearrow 1} \beta_\nu(t) / (1-t) \quad \text{and} \quad \underline{\delta}_\nu := \lim_{t \searrow 1} \beta_\nu(t) / (1-t).$$

Conceptually, for the lower bound we borrow ideas from the *coarse multifractal analysis* (see [Fal14, Ric95]). For all  $n \in \mathbb{N}$ , we define for  $\alpha > 0$

$$\bar{F}_\nu(\alpha) := \limsup_n \frac{\log^+(\mathcal{N}_\alpha(n))}{\log 2^n} \quad \text{and} \quad \underline{F}_\nu(\alpha) := \liminf_n \frac{\log^+(\mathcal{N}_\alpha(n))}{\log 2^n},$$

where

$$\mathcal{N}_\alpha(n) := \text{card } M_n(\alpha), \quad M_n(\alpha) := \{0 \leq k < 2^n : \nu(A_k^n) \geq 2^{-\alpha n}\}$$

and refer to the quantities

$$\bar{F}_\nu := \sup_{\alpha > 0} \frac{\bar{F}_\nu(\alpha)}{(1+\alpha)} \quad \text{and} \quad \underline{F}_\nu := \sup_{\alpha > 0} \frac{\underline{F}_\nu(\alpha)}{(1+\alpha)}$$

as the *upper*, resp. *lower, optimized coarse multifractal dimension*. We shall see later that the highly relevant connections between the optimal and the dyadic partition approach are strongly linked by ideas from the theory of large deviations.

The following list of results gives the main achievements of this paper. The proofs are postponed to Sections 3 and 4.

**Theorem 1.1.** *For all  $1 < m \leq 3$ , we have*

$$\underline{F}_\nu \leq \underline{h}_\nu^m \leq \underline{s}_\nu \leq \underline{h}_\nu \leq \bar{h}_\nu = \bar{h}_\nu^m = \bar{s}_\nu = \bar{q}_\nu = \bar{F}_\nu. \quad (1.1)$$

*In particular,  $\bar{q}_\nu \leq 1/2$  and the following necessary and sufficient conditions hold for the spectral dimension to exist:*

$$\underline{s}_\nu = \bar{s}_\nu \implies \underline{h}_\nu = \bar{h}_\nu = s_\nu \quad \text{and} \quad \sup_{m > 1} \underline{h}_\nu^m = \bar{h}_\nu \implies \underline{s}_\nu = \bar{s}_\nu = \bar{h}_\nu.$$

*We would like to emphasize that if the spectral dimension exists, then it is given by purely measure-geometric data, which is encoded in the  $\nu$ -partition entropy  $\bar{h}_\nu = \underline{h}_\nu$ . We call  $\nu$  regular, if  $\sup_{m > 1} \underline{h}_\nu^m = \bar{h}_\nu$ , in which case the spectral dimension exists. If  $\nu$  is MF-regular, that is  $\underline{F}_\nu = \bar{F}_\nu$ , then in the above chain of inequalities (1.1) we have everywhere equality and especially  $\nu$  is regular. Moreover, if for some  $m > 1$  we have  $\underline{h}_\nu^m \geq 1/2$ , then  $s_\nu = 1/2 = \bar{h}_\nu$ .*

The following theorem shows that the  $L^q$ -spectrum is a valuable auxiliary concept to determine the spectral behavior of a given measure  $\nu$ . We call the measure  $\nu$   $L^q$ -regular, if  $\beta_\nu(q) = \liminf_n \beta_n^\nu(q)$  for  $q \in (\bar{q}_\nu - \varepsilon, \bar{q}_\nu)$ , for some  $\varepsilon > 0$ , or  $\beta_\nu(\bar{q}_\nu) = \liminf_n \beta_n^\nu(\bar{q}_\nu)$  and  $\beta_\nu$  is differentiable in  $\bar{q}_\nu$ .

**Theorem 1.2.** *If  $\nu$  is  $L^q$ -regular, then it is MF-regular.*

This result is optimal in the sense that there is an easily accessible example of a measure  $\nu$  which is not  $L^q$ -regular and for which  $\bar{s}_\nu = \bar{h}_\nu > \underline{h}_\nu = \underline{s}_\nu$  and the spectral dimension does not exist (see Section 5.2). It should be noted that  $L^q$  regularity is much easier accessible than regularity or MF regularity and that, to the best of our knowledge, all measures examined in the literature for which the spectral dimension is known are  $L^q$ -regular.

As a first application of these considerations, we present the case of measures  $\nu$  with absolutely continuous part based solely on the  $L^q$ -spectrum.

**Corollary 1.3.** *If  $\nu$  has an absolutely continuous part with respect to  $\Lambda$ , then for  $q \in [0, 1]$ , the  $L^q$ -spectrum exists as a limit with*

$$\beta_\nu(q) = 1 - q.$$

*In particular,  $\nu$  is  $L^q$ -regular and the spectral dimension exists and equals  $1/2$ .*

Also for measures without an absolutely continuous part we have the following rigidity result in terms of reaching the maximum possible value  $1/2$  of the spectral dimension.

**Corollary 1.4.** *The following rigidity result holds:*

1. *If  $\bar{s}_\nu = 1/2$ , then  $\beta_\nu(q) = 1 - q$  for all  $q \in [0, 1]$ .*
2. *If  $\beta_\nu(q) = \lim \beta_n^\nu(q) = 1 - q$  for some  $q \in (0, 1)$ , then  $\beta_\nu(q) = 1 - q$  for all  $q \in [0, 1]$  and  $s_\nu = 1/2$ .*

In the following proposition we present lower bounds of the lower spectral dimension in terms of the subdifferential  $\partial\beta_\nu(q) := \{a \in \mathbb{R} : \forall t \in \mathbb{R} : \beta_\nu(t) \geq a(t - q) + \beta_\nu(q)\}$  of  $\beta_\nu$  in  $q$ .

**Proposition 1.5.** *If for  $q \in [0, 1]$  we have  $\beta_\nu(q) = \lim_n \beta_n^\nu(q)$  and  $-\partial\beta_\nu(q) = [a, b]$ , then*

$$\frac{aq + \beta_\nu(q)}{1 + b} \leq \underline{s}_\nu.$$

*Remark 1.6.* In the case that  $\beta_\nu(\bar{q}_\nu) = \lim \beta_n^\nu(\bar{q}_\nu)$  and  $\beta_\nu$  is differentiable in  $\bar{q}_\nu$ , we infer  $\bar{q}_\nu \leq \underline{s}_\nu$  and hence obtain a direct proof of the regularity statement, namely,  $\bar{q}_\nu = \underline{s}_\nu = \bar{s}_\nu$ .

As a corollary to Proposition 1.5 and Theorem 1.1 we improve the known general upper bound of the spectral dimension of  $1/2$  as obtained in [BS66] in terms of the upper Minkowski dimension. Furthermore, we obtain a general lower bound of  $\underline{s}_\nu$  in terms of the left and right-hand derivative of  $\beta_\nu$ .

**Corollary 1.7.** *For the upper and lower spectral dimension we have the following general upper and lower bounds depending on the topological support of  $\nu$ , namely  $\delta_\nu^* = \overline{\dim}_M(\nu)$ , and right and left derivative of  $\beta_\nu$  in 1:*

$$\frac{\underline{\delta}_\nu}{1 + \underline{\delta}_\nu} \leq \underline{s}_\nu \leq \bar{s}_\nu \leq \frac{\delta_\nu^*}{1 + \delta_\nu^*} \leq \frac{1}{2}$$

and

$$\bar{s}_\nu = \frac{\delta_\nu^*}{1 + \delta_\nu^*} \iff \bar{\delta}_\nu = \delta_\nu^*.$$

*Remark 1.8.* It is worth pointing out that these bounds have been first observed in the self-similar case under the open set condition in [SV95, p. 245] (in this case the Minkowski dimension and the Hausdorff of  $\text{supp } \nu$  coincide as well as  $\beta_\nu$  is differentiable in 1 and  $\beta'_\nu(1)$  coincides with the Hausdorff dimension of  $\nu$ ). Furthermore, note that in the case that  $\nu$  has an atomic part we always have  $\underline{\delta}_\nu = 0$  (see Fact 4.6). Hence, the lower bound  $\underline{\delta}_\nu/(1 + \bar{\delta}_\nu)$  is only meaningful in the case of atomless measures.

Regarding Kac's question, if  $\beta_\nu$  is differentiable in 1, then the spectral dimension is determined by fractal-geometric quantities as follows

$$\dim_H(\nu) = \dim_P(\nu) \leq \frac{\underline{s}_\nu}{1 - \underline{s}_\nu} \leq \frac{\bar{s}_\nu}{1 - \bar{s}_\nu} \leq \overline{\dim}_M(\nu).$$

Under the additionally regularity condition imposed on  $\nu$ , we have the following rigidity result:

**Corollary 1.9.** *If  $\nu$  is  $L^q$ -regular, then we have*

1.  $\underline{\delta}_\nu = \delta_\nu^* \iff s_\nu = \frac{\delta_\nu^*}{1 + \delta_\nu^*},$
2.  $\overline{\dim}_M(\nu) = \dim_H(\nu) \implies s_\nu = \frac{\delta_\nu^*}{1 + \delta_\nu^*},$
3.  $\beta_\nu$  is differentiable in 1 and  $s_\nu = \frac{\delta_\nu^*}{1 + \delta_\nu^*}$  implies

$$\overline{\dim}_M(\nu) = \dim_H(\nu) = \underline{\dim}_H(\nu) := \inf \{ \dim_H(E) : \nu(E) > 0 \} = \dim_P(\nu),$$

where  $\dim_H(E)$ ,  $\dim_P(E)$  denotes the Hausdorff dimension, packing dimension of  $E$ , respectively.

The second implication explains that in earlier publication (e. g. [Fuj87, UH59]) the spectral dimension has typically been connected to the Hausdorff dimension of  $\nu$ . The implication in (3) concerning the Hausdorff and packing dimension goes back to the work of Ngai and Heurteaux [Nga97, Heu98].

The last implication is optimal in the sense that we find a measure  $\nu$  (Example 5.11) with  $1 = \delta_\nu^* = \bar{\delta}_\nu$ ,  $\beta$  is not differentiable in 1 and  $0 = \dim_H(\nu) = \dim_P(\nu) < \bar{\delta}_\nu = 1$ . We note that the equality  $\delta_\nu^* = \dim_P(\nu)$  is implied by  $\delta_\nu^* = \dim_H(\nu)$ ,  $\delta_\nu^* = h^*(\nu) = \limsup_n (\log 2^n)^{-1} \sum_{C \in \mathcal{D}_n} \nu(C) \log \nu(C)$ , or by the identities  $\delta_\nu^* = h_*(\nu) = \liminf_n (\log 2^n)^{-1} \sum_{C \in \mathcal{D}_n} \nu(C) \log \nu(C)$ .

The following proposition complements the connection of the Minkowski dimension by establishing an upper bound of the lower spectral dimension in terms of the lower Minkowski dimension  $\underline{\dim}_M(\nu)$  of the support of  $\nu$ .

**Proposition 1.10.** *We always have*

$$\underline{s}_\nu \leq \underline{h}_\nu \leq \underline{\dim}_M(\nu).$$

Next we give a lower bound for the lower spectral dimension in terms of the Minkowski dimension (if it exists) and the derivative of the  $L^q$ -spectrum in 0.

**Corollary 1.11.** *If the Minkowski dimension of  $\text{supp } \nu$  exists, i. e.  $\overline{\dim}_M(\nu) = \underline{\dim}_M(\nu)$ , and  $-\partial\beta_\nu(0) = [a, b]$ , then*

$$\frac{\delta_\nu^*}{1+b} \leq \underline{s}_\nu.$$

The upper spectral dimension is stable in the following sense.

**Corollary 1.12.** *For two finite Borel measures  $\nu_1$  and  $\nu_2$  on  $(0, 1)$  with upper spectral dimension  $\overline{s}_{\nu_1}$  and  $\overline{s}_{\nu_2}$ , we have that the upper spectral dimension of  $\nu_1 + \nu_2$  is given by  $\max\{\overline{s}_{\nu_1}, \overline{s}_{\nu_2}\}$ . If additionally,  $\underline{s}_{\nu_1} = \overline{s}_{\nu_1} \geq \overline{s}_{\nu_2}$ , then its spectral dimension exists and equals  $\overline{s}_{\nu_1}$ . If  $\underline{s}_{\nu_1} = \overline{s}_{\nu_1} \geq \overline{s}_{\nu_2}$  and  $\nu_1$  is regular, then so is  $\nu_1 + \nu_2$ .*

Moreover, the spectral dimension (and in fact also finer spectral asymptotic properties) is stable under bi-Lipschitz mappings.

**Proposition 1.13.** *Let  $\nu$  be a Borel measure on  $(0, 1)$ . Further let  $g : [0, 1] \rightarrow [a, b]$  be a bi-Lipschitz mapping. Then uniformly for  $i \in \mathbb{N}$ , we have*

$$\lambda_{\nu, [0,1]}^i \asymp \lambda_{\nu \circ g^{-1}, [a,b]}^i.$$

In the last section we give a sequence of examples showing how our approach allows to analyze in detail particular classes of measures: First, we determine the spectral dimension of measures that stem from *non-linear iterated function systems* (with or without overlap), see Section 5.1.

**Theorem 1.14.** *For any  $C^1$ -self-conformal measure  $\nu$ , the spectral dimension of  $\Delta_\nu$  exists and equals  $\overline{q}_\nu$ .*

Second, we study *homogeneous Cantor measure*, for which we provide examples such that the spectral dimension does not exist with  $\overline{s}_\nu = \overline{h}_\nu > \underline{h}_\nu = \underline{s}_\nu$  and an other example with spectral dimension 0. Finally, we consider *pure point measure* such that its spectral dimension exists and attains any given value in  $[0, 1/2]$ .

We end this introduction with some concluding remarks. Our newly established connection to the  $L^q$ -spectrum puts the results of [UH59], [Fuj87], [SV95] and [Nga11, NTX18, NX19, NX20] from an abstract point of view in the right context; for the self-similar measure  $\nu$  under the OSC, the  $L^q$ -spectrum of  $\nu$  can be expressed in terms of the associated pressure function. In the case of self-conformal measures with overlaps the situation becomes much more involved; nevertheless, by a result in [BF20], the  $L^q$ -spectrum will also exist in this situation and some explicit formulae can be provided in the self-similar setting (see [KN21b]).

We remark that the operator  $\Delta_{\nu, [a,b]}$  has been generalized by considering the form  $(f, g) \mapsto \int_{[a,b]} \nabla_\mu f \nabla_\mu g \, d\mu$  where  $\mu$  is an atomless Borel measure on  $[a, b]$  with and  $\nabla_\mu f$  denotes the weak derivative with respect to  $\mu$ , i. e.  $f(x) = f(a) + \int_{[a,x]} \nabla_\mu f \, d\mu$ ,  $x \in [a, b]$ . The associated generalized Kreĭn-Feller operator  $\Delta_{\nu, \mu, [a,b]}$  and its spectral properties have been studied in [Küc86], [Vol05], [Fre11, Fre08, Fre05, FZ02]. In [KNSW19] it has been shown that by a simple transformation of measure spaces by virtue of the distribution function  $F_\mu$  of  $\mu$  the generalized Kreĭn-Feller operator can be reduced to the classical one, namely  $\Delta_{\nu \circ F_\mu^{-1}, \Lambda, [F_\mu(a), F_\mu(b)]}$ , which has the same spectral properties. For this reason we have restricted our attention to the case  $\mu = \Lambda$ .

Finally, we would like to note that the operator  $\Delta_\nu$  has also been generalized to higher dimensions in [Sol94, NS95, Tri97, FT99, NS01, HLN06, DN15, DN21, NX21]. In the forthcoming paper [KN22] we elaborate the ideas of the present paper also for higher dimensions.

Furthermore, the operator  $\Delta_\nu$  can be generalized to polyharmonic operators in higher dimensions which has been considered in [BS74, BS70]. Polyharmonic operators with respect to self-similar measures under the OSC have been studied in [Naz04] using the same approach as in [SV95]. We would like to point out that our variational approach can also be used to solve the spectral problem in these higher dimensions as well as for polyharmonic operators; this will be the content of the forthcoming paper [KN21a], in which we improve known bounds on the spectral asymptotics for polyharmonic operators with respect to any Borel measures on  $(0, 1)^d$ ,  $d \in \mathbb{N}$ .

## 2. Spectral theory for Kreĭn-Feller operators

In this chapter, we develop a form approach for the Kreĭn-Feller operator that allows us to rigorously prove the sub- and superadditivity of the eigenvalue counting function. In doing so, we follow various ideas as found, for example, in [KKVW09, Fag12, HLN06]. Crucially for our purposes, and in contrast to previous work, we consider ambient intervals for which the boundary points are not necessarily in the support of the underlying measure, which moreover may have a non-vanishing pure point part. As this setup is not covered in the literature, we decided to provide the details necessary for our purpose. In particular, we give as a key observation Proposition 2.13, which indeed seems to be used implicitly throughout the existing literature.

### 2.1. Form approach

For a fixed finite Borel measure  $\nu$  on  $(a, b)$  let us define

$$C_\nu([a, b]) := \{f \in C([a, b]) \mid f \text{ is affine linear on the components of } [a, b] \setminus \text{supp } \nu\},$$

where  $C([a, b])$  denotes the space of continuous real-valued functions on  $[a, b]$ . Note that  $(C_\nu([a, b]), \|\cdot\|_\infty)$  is a Banach space. Further, we define the *Sobolev space* of weakly differentiable functions

$$H^1(a, b) := \left\{ f : [a, b] \rightarrow \mathbb{R} \mid \exists g \in L^2_\Lambda([a, b]) : f(x) = f(a) + \int_{[a, x]} g \, d\Lambda, x \in [a, b] \right\}$$

together with its closed subspace

$$H^1_0(a, b) := \{u \in H^1(a, b) \mid u(a) = u(b) = 0\}.$$

The function  $g$  in the definition of  $H^1(a, b)$  is uniquely determined by  $f$  and we write  $\nabla_\Lambda f := g$ . The space  $H^1_0(a, b)$  with the bilinear form

$$(u, v)_{H^1_0} := \int_{(a, b)} uv \, d\Lambda + \int_{(a, b)} \nabla_\Lambda f \nabla_\Lambda g \, d\Lambda$$

defines a Hilbert space. We will see that

$$\mathcal{E}_{v,[a,b]}(f, g) := \int_{(a,b)} \nabla_{\Lambda} f \nabla_{\Lambda} g \, d\Lambda$$

with (Dirichlet) domain  $\text{dom}(\mathcal{E}_v) := H_0^1(a, b) \cap C_v([a, b])$  defines a form for  $L_v^2$ . First note that  $\mathcal{E}_{v,[a,b]}$  is well defined. Indeed, for  $f \in \text{dom}(\mathcal{E}_{v,[a,b]})$  with  $f = 0$   $v$ -a. e. we have that  $f$  vanishes on  $\text{supp } v$  by continuity. Moreover, by definition of  $\text{dom}(\mathcal{E}_{v,[a,b]})$ , we have that  $f$  also vanishes on the complement of  $\text{supp } v$ . Therefore, for all  $g \in \text{dom}(\mathcal{E}_{v,[a,b]})$ , it follows that  $\mathcal{E}_{v,[a,b]}(f, g) = 0$ . In the following we need the decomposition

$$[a, b] \setminus \text{supp } v = A_1 \cup A_2 \cup \bigcup_{i \in I} (a_i, b_i),$$

where  $I \subset \mathbb{N}$  and  $A_1 := [a, d_1]$  if  $a \notin \text{supp } v$  otherwise  $A_1 = \emptyset$ ,  $A_2 := (d_2, b]$  if  $b \notin \text{supp } v$  and the intervals  $[a, c_1)$ ,  $(c_2, b]$ ,  $(a_i, b_i)$ ,  $i \in I$ , are mutually disjoint.

**Lemma 2.1.** *The map  $\tau_{v,[a,b]} : H_0^1(a, b) \rightarrow \text{dom}(\mathcal{E}_v)$*

$$\tau_{v,[a,b]}(f)(x) := \begin{cases} f(a_i) + \frac{f(b_i) - f(a_i)}{b_i - a_i} (x - a_i), & x \in (a_i, b_i), i \in I, \\ \frac{f(d_1)}{d_1 - a} (x - a), & x \in [a, d_1), a \notin \text{supp } v, \\ \frac{f(d_2)}{b - d_2} (b - x), & x \in (d_2, b], b \notin \text{supp } v, \\ f(x), & x \in \text{supp } v \end{cases}$$

is surjective,  $\tau_{v,[a,b]}(f) = f$  as elements of  $L_v^2$ , and we have

$$\nabla_{\Lambda|_{[a,b]}} \tau_{v,[a,b]}(f)(x) = \begin{cases} \frac{f(b_i) - f(a_i)}{b_i - a_i}, & x \in (a_i, b_i), i \in I, \\ \frac{f(d_1)}{d_1 - a}, & x \in [a, d_1), a \notin \text{supp } v, \\ -\frac{f(d_2)}{b - d_2}, & x \in (d_2, b], b \notin \text{supp } v, \\ \nabla_{\Lambda} f(x), & x \in \text{supp } v. \end{cases}$$

*Proof.* We follow the arguments as outlined in [KKVW09]. For  $f \in H_0^1(a, b)$ , we set

$$\tau_n(f)(x) := \begin{cases} f(a_i) + \frac{f(b_i) - f(a_i)}{b_i - a_i} (x - a_i), & x \in (a_i, b_i), i \in I, i \leq n \\ \frac{f(d_1)}{d_1 - a} (x - a), & x \in [a, d_1), a \notin \text{supp } v, \\ \frac{f(d_2)}{b - d_2} (b - x), & x \in (d_2, b], b \notin \text{supp } v, \\ f(x), & \text{otherwise,} \end{cases}$$

and choose a  $L_{\Lambda}^2$  representative of  $\nabla_{\Lambda} f$  to define

$$g_n(x) := \begin{cases} \frac{f(b_i) - f(a_i)}{b_i - a_i}, & x \in (a_i, b_i), i \in I, i \leq n \\ \frac{f(d_1)}{d_1 - a}, & x \in [a, d_1), a \notin \text{supp } v, \\ -\frac{f(d_2)}{b - d_2}, & x \in (d_2, b], b \notin \text{supp } v, \\ \nabla_{\Lambda} f(x), & \text{otherwise.} \end{cases}$$

Then we have  $\tau_n(f) \in H_0^1(a, b)$  with  $\nabla_\Lambda \tau_n(f) = g_n$  a. e. and clearly we have  $\tau_n(f) = f$  in  $L_v^2$ . Since  $f$  is uniformly continuous on the compact set  $[a, b]$ , for  $n$  tending to infinity, we have  $\tau_n(f) \rightarrow \tau_{v,[a,b]}(f) \in C_v([a, b])$  uniformly on  $[a, b]$ . Now, for each  $x \in [a, b]$ , set  $g(x) := \lim_{n \rightarrow \infty} g_n(x)$ . Then by Cauchy-Schwarz inequality, we have

$$\begin{aligned} \int g^2 d\Lambda &= \sum_{i \in I} \int_{(a_i, b_i)} g^2 d\Lambda + \int_{\text{supp } v} g^2 d\Lambda = \frac{(f(b_i) - f(a_i))^2}{b_i - a_i} + \int_{\text{supp } v} (\nabla_\Lambda f)^2 d\Lambda \\ &= \sum_{i \in I} \frac{\left( \int_{(a_i, b_i)} \nabla_\Lambda f d\Lambda \right)^2}{(b_i - a_i)} + \int_{\text{supp } v} (\nabla_\Lambda f)^2 d\Lambda \\ &\leq \sum_{i \in I} \int_{(a_i, b_i)} (\nabla_\Lambda f)^2 d\Lambda + \int_{\text{supp } v} (\nabla_\Lambda f)^2 d\Lambda = \int (\nabla_\Lambda f)^2 d\Lambda < \infty \end{aligned}$$

implying  $g \in L_{\Lambda|_{(a,b)}}^2$ . Furthermore, for all  $n \in \mathbb{N}$ , we have

$$|g_n(x)|^2 \leq (|g_n(x)| + |\nabla_\Lambda f(x)|)^2 \leq 4(|g_n(x)|^2 + |\nabla_\Lambda f(x)|^2)$$

and by the dominated convergence theorem we conclude  $g_n \rightarrow g$  in  $L_{\Lambda|_{(a,b)}}^2$ . This implies  $\tau_{v,[a,b]}(f) \in \text{dom}(\mathcal{E}_v)$  and  $\nabla_\Lambda \tau_{v,[a,b]}(f) = g$ .  $\square$

**Proposition 2.2.** *The set  $\text{dom}(\mathcal{E}_{v,[a,b]})$  is dense in  $L_v^2$ .*

*Proof.* Here we follow the arguments similar to [KKVW09]. Let  $C_c(a, b)$  denote the space of continuous functions from  $(a, b)$  to  $\mathbb{R}$  with compact support and let  $C_c^1(a, b)$  be the space of differentiable functions from  $(a, b)$  to  $\mathbb{R}$  with compact support. By Lemma 2.1 every  $f \in C_c^1(a, b)$  possesses a representative  $\tau_{v,[a,b]}(f) \in \text{dom}(\mathcal{E}_{v,[a,b]})$ , using  $f(a) = f(b) = 0$ . It is well known that  $C_c^1(a, b)$  lies dense in  $C_c(a, b)$  with respect to the uniform norm and  $C_c(a, b)$  in  $L_v^2$ , hence we obtain that  $\text{dom}(\mathcal{E}_{v,[a,b]})$  is dense in  $L_v^2$ .  $\square$

**Proposition 2.3.** *The set  $\text{dom}(\mathcal{E}_v)$  equipped with the inner product*

$$(f, g)_{\mathcal{E}_v} := \langle f, g \rangle_v + \mathcal{E}_v(f, g),$$

*defines a Hilbert space, i. e.  $\mathcal{E}_v$  is closed with respect to  $L_v^2$ .*

*Proof.* Here, we follow ideas of [Fag12, Lemma 6.4.]. Let  $(u_n)_{n \in \mathbb{N}}$  be a Cauchy sequence in  $\text{dom}(\mathcal{E}_{v,[a,b]})$  with respect to the norm induced by  $(\cdot, \cdot)_{\mathcal{E}_{v,[a,b]}}$ , i. e.

$$\|u_n - u_m\|_{L_v^2}^2 + \|\nabla_\Lambda u_n - \nabla_\Lambda u_m\|_{L_\Lambda^2}^2 \rightarrow 0$$

for  $n, m \rightarrow \infty$ . Since  $L_v^2$  and  $L_\Lambda^2$  are Hilbert spaces, there exist  $u \in L_v^2$  and  $f \in L_\Lambda^2$  such that

$$\lim_{n \rightarrow \infty} \|u_n - u\|_{L_v^2} = 0 \text{ and } \lim_{n \rightarrow \infty} \|f - \nabla_\Lambda u_n\|_{L_\Lambda^2} = 0.$$

Using Cauchy-Schwarz inequality, we find uniformly for all  $x \in [a, b]$ ,

$$\begin{aligned} \left| u_n(x) - u_n(a) - \underbrace{\int_{[a,x]} f(z) \, d\Lambda(z)}_{=:g(x)} \right| &= \left| \int_{[a,x]} (f(z) - \nabla_\Lambda u_n(z)) \, d\Lambda(z) \right| \\ &\leq \sqrt{\Lambda([a, b])} \|\nabla_\Lambda u_n - f\|_{L_\Lambda^2} \rightarrow 0, \end{aligned}$$

for  $n \rightarrow \infty$ . In particular,  $u_n - u_n(a) \rightarrow g$  in  $L_v^1$  and therefore  $u_n(a) = (u_n(a) - u_n) + u_n \rightarrow u - g$  in  $L_v^1$ . But this shows that the sequence  $(u_n(a))_{n \in \mathbb{N}}$  converge to  $u_0 := \nu([a, b])^{-1} \int u - g \, d\nu$ . Hence, we conclude that  $u_n$  convergence uniformly to  $h := u_0 + g \in H^1(a, b)$  with  $\nabla_\Lambda h = f$ , in particular, uniform convergence guarantees that  $h(a) = h(b) = 0$  and  $h \in C_v([a, b])$ .  $\square$

**Lemma 2.4.** *There exists a constant  $C$  such that for every  $f \in H^1(a, b)$*

$$\|f\|_\infty \leq C \left( \|\nabla_\Lambda f\|_{L_\Lambda^2(a,b)}^2 + \|f\|_{L_v^2}^2 \right)^{1/2}$$

and, for all  $f \in H_0^1(a, b)$ ,

$$\int_{(a,b)} f^2 \, d\nu \leq \nu([a, b]) (b - a) \int_{(a,b)} (\nabla_\Lambda f)^2 \, d\Lambda.$$

*Proof.* The first inequality has been proved in [KKVW09, Lemma 1.4.]. Further, for fixed  $f \in H_0^1(a, b)$ , we have

$$\begin{aligned} \int_{(a,b)} f(x)^2 \, d\nu(x) &= \int \left( \int_{(a,x)} (\nabla_\Lambda f) \, d\Lambda \right)^2 \, d\nu(x) \\ &\leq \int \left( (x - a) \int_{(a,x)} (\nabla_\Lambda f)^2 \, d\Lambda \right) \, d\nu(x) \\ &\leq (b - a) \nu([a, b]) \int_{(a,b)} (\nabla_\Lambda f)^2 \, d\Lambda. \end{aligned}$$

$\square$

*Remark 2.5.* Due to the above lemma, we can equivalently consider  $\text{dom}(\mathcal{E}_\nu)$  as a Hilbert space whose inner product is given by  $\mathcal{E}_\nu$ .

**Proposition 2.6.** *The inclusion from the Hilbert space  $(\text{dom}(\mathcal{E}_\nu), \mathcal{E}_\nu)$  into  $L_v^2$  is compact.*

*Proof.* Let  $U \subset \text{dom}(\mathcal{E}_{\nu, [a,b]})$  be a bounded set with respect to  $(\cdot, \cdot)_{\mathcal{E}_{\nu, [a,b]}}$  which means there exists  $c > 0$  such that

$$\sup_{f \in U} \left( \int_{(a,b)} (\nabla_\Lambda f)^2 \, d\Lambda \right)^{1/2} \leq c.$$

We have to show  $U$  is pre-compact in  $L_v^2$ . This will be done by showing that  $U$  is bounded in  $C([a, b])$  with respect to  $\|\cdot\|_\infty$  and equicontinuous. The Arzelà–Ascoli Theorem then guarantees that  $U$  is pre-compact in  $C([a, b])$ . Hence  $C([a, b])$  is continuously embedded in  $L_v^2$  noting that  $\|f\|_{L_v^2} \leq \nu([a, b]) \|f\|_\infty$ , for all  $f \in C([a, b])$ . The set  $U$  is also pre-compact in  $L_v^2$ . Using the Cauchy-Schwarz inequality, for all  $x, y \in [a, b]$ , we have

$$|f(x) - f(y)| = \left| \int_{(x,y)} \nabla_\Lambda f \, d\Lambda \right| \leq |x - y|^{1/2} \left( \int_{(a,b)} (\nabla_\Lambda f)^2 \, d\Lambda \right)^{1/2} \leq c |x - y|^{1/2}.$$

Hence,  $U$  is equicontinuous. Lemma 2.4 implies that  $U$  is also bounded in  $(C([a, b]), \|\cdot\|_\infty)$ .  $\square$

If  $\nu(\{a, b\}) = 0$  then Proposition 2.3 and Proposition 2.2 show that the densely defined form  $\mathcal{E}_{\nu, [a, b]}$  is closed. Hence, by [Dav95, Theorem 4.4.2] or [Kat76, Theorem 2.1., p. 322] there exists a non-negative self-adjoint operator  $\Delta_{\nu, [a, b]}$  which is characterized as follows;  $f \in \text{dom}(\mathcal{E}_{\nu, [a, b]})$  lies in the domain  $\text{dom}(\Delta_{\nu, [a, b]})$  of the Laplacian  $\Delta_{\nu, [a, b]}$ , if and only if there exists  $h \in L_v^2$  such that

$$\mathcal{E}_{\nu, [a, b]}(g, f) = \langle g, h \rangle_\nu, \quad \text{for all } g \in \text{dom}(\mathcal{E}_{\nu, [a, b]})$$

with  $h =: \Delta_{\nu, [a, b]} f =: \Delta_\nu f$ .

Observe that  $H_0^1(a, b)$  is compactly embedded in  $L_v^2$ , consequently the closed subspace  $\text{dom}(\mathcal{E}_\nu)$  is also compactly embedded in  $L_v^2$ . Consequently, we conclude (see [Tri92, Theorem 4.5.1. and p. 258]) that there exists an orthonormal system of eigenfunctions of  $\Delta_\nu$  of  $L_v^2$  with non-negative eigenvalues  $(\lambda_\nu^n)_{n \in \mathbb{N}}$  tending to  $\infty$  given  $\text{supp } \nu$  is not finite.

## 2.2. Spectral problem, min-max principle, and sub- and superadditivity

**Definition 2.7.** An element  $f \in \text{dom}(\mathcal{E}_\nu) \setminus \{0\}$  is called *eigenfunction* of  $\mathcal{E}_\nu$  with *eigenvalue*  $\lambda$ , if for all  $g \in \text{dom}(\mathcal{E}_\nu)$ , we have

$$\mathcal{E}_\nu(f, g) = \lambda \cdot \langle f, g \rangle_\nu.$$

We denote the number of eigenvalues of  $\mathcal{E}_\nu$  not exceeding  $x$  by  $N_\nu(x)$  and refer to  $N_\nu = N_{\nu, [a, b]}$  as the *eigenvalue counting function*.

*Remark 2.8.* Note that for any fixed  $c > 0$ ,  $\lambda$  is eigenvalue of  $\mathcal{E}_\nu$  if and only if  $c\lambda$  is an eigenvalue of  $\mathcal{E}_{c\nu}$ . Therefore, the spectral dimension does not change after rescaling the measure and we may assume that  $\nu$  is Borel probability measure.

*Remark 2.9.* Let  $\mathcal{E}$  be a closed form with domain  $\text{dom}(\mathcal{E})$  densely defined on  $L_v^2$ , in particular,  $\text{dom}(\mathcal{E})$  defines a Hilbert space with respect to  $(f, g)_\mathcal{E} := \langle f, g \rangle_\nu + \mathcal{E}(f, g)$ , and assume that the inclusion from  $(\text{dom}(\mathcal{E}), (\cdot, \cdot)_\mathcal{E})$  into  $L_v^2$  is compact. Then the *Poincaré–Courant–Fischer–Weyl min-max principle* is applicable, that is for the  $i$ -th eigenvalue  $\lambda^i(\mathcal{E})$  of  $\mathcal{E}$ ,  $i \in \mathbb{N}$ , we have (see also [Dav95, KL93])

$$\lambda^i(\mathcal{E}) = \inf \left\{ \sup \left\{ \frac{\mathcal{E}(\psi, \psi)}{\langle \psi, \psi \rangle_\nu} : \psi \in G^\star \right\} : G \prec_i \text{dom}(\mathcal{E}) \right\},$$

where  $J \prec_i H$  means that  $J$  is a linear subspace of the Hilbert space  $H$  and the vector space dimension of  $J$  is equal to  $i \in \mathbb{N}$  and we set  $J^\star := J \setminus \{0\}$ .

Let  $P := \{(c_k, c_{k+1}] : k = 1, \dots, n-1\}$  a  $\nu$ -partition of  $(a, b]$ ,  $n \in \mathbb{N}_{\geq 2}$ ,  $a = c_1 < c_2 < \dots < c_n = b$ . For a subinterval  $(c_k, c_{k+1})$  we may define the restricted form  $\mathcal{E}_{\nu, [c_k, c_{k+1}]}$  on  $L^2_{\nu|_{[c_k, c_{k+1}]}}$  with domain  $\text{dom}(\mathcal{E}_{\nu, [c_k, c_{k+1}]}) := H_0^1(c_k, c_{k+1}) \cap C_\nu(c_k, c_{k+1})$  by

$$\mathcal{E}_{\nu, [c_k, c_{k+1}]}(f, g) = \int_{c_k}^{c_{k+1}} \nabla_\Lambda f \nabla_\Lambda g \, d\Lambda.$$

Consider the natural direct sum Hilbert space

$$L^2_{\nu, P} := \bigoplus_{k=1}^n L^2_{\nu|_{[c_k, c_{k+1}]}}$$

with inner product

$$\langle (f_k), (g_k) \rangle_{\nu, P} := \sum_{k=1}^n \langle f_k, g_k \rangle_{\nu|_{[c_k, c_{k+1}]}}$$

and corresponding Sobolev spaces

$$H_P^1 := \bigoplus_{k=1}^n H_0^1(c_k, c_{k+1}).$$

We define the *direct sum form*

$$\mathcal{E}_{\nu, P}((f_k), (g_k)) := \sum_{k=1}^n \mathcal{E}_{\nu, [c_k, c_{k+1}]}(f_k, g_k)$$

with domain

$$\text{dom}(\mathcal{E}_{\nu, P}) := \bigoplus_{k=1}^n \text{dom}(\mathcal{E}_{\nu, [c_k, c_{k+1}]}).$$

Both  $\text{dom}(\mathcal{E}_{\nu, P})$  and  $H_P^1$  are Hilbert spaces with respect to the inner product given by  $\mathcal{E}_{\nu, P}$ .

**Lemma 2.10.** *The form  $\mathcal{E}_{\nu, P}$  is densely defined and closed.*

*Proof.* This follows from Proposition 2.3 and 2.2 applied to each component.  $\square$

The eigenvalue counting function with respect to  $\mathcal{E}_{\nu, P}$  will be denoted by  $N_{\nu, P}$ .

**Lemma 2.11.** *For a  $\nu$ -partition  $P := \{(c_k, c_{k+1}] : k = 1, \dots, n-1\}$  and all  $x > 0$ , we have*

$$N_{\nu, P}(x) = \sum_{k=1}^n N_{\nu, [c_k, c_{k+1}]}(x).$$

*Proof.* Let  $(f_k)_k \in \text{dom}(\mathcal{E}_{\nu, P})$  be an eigenfunction with eigenvalue  $\lambda > 0$ , i. e. for  $(g_k)_k \in \text{dom}(\mathcal{E}_{\nu, P})$

$$\sum_{k=1}^n \mathcal{E}_{\nu, [c_k, c_{k+1}]}(f_k, g_k) = \lambda \sum_{k=1}^n \langle f_k, g_k \rangle_{\nu|_{[c_k, c_{k+1}]}}.$$

Hence, if  $f_k \neq 0$  it follows all  $g \in \text{dom}(\mathcal{E}_{\nu|_{[c_k, c_{k+1}]}})$  that  $(g_k)_k = (0, \dots, 0, g, 0, \dots, 0) \in \text{dom}(\mathcal{E}_{\nu, P})$

$$\mathcal{E}_{\nu, [c_k, c_{k+1}]}(f_k, g) = \lambda \langle f_k, g \rangle_{\nu|_{[c_k, c_{k+1}]}}.$$

This shows that  $f_k$  is an eigenfunction of  $\mathcal{E}_{\nu, [c_k, c_{k+1}]}$  with eigenvalue  $\lambda$ , which implies

$$N_{\nu, P}(x) \leq \sum_{k=1}^n N_{\nu, [c_k, c_{k+1}]}(x).$$

In a similar way it follows  $N_{\nu, P}(x) \geq \sum_{k=1}^n N_{\nu, [c_k, c_{k+1}]}(x)$ .  $\square$

**Lemma 2.12.** For a  $\nu$ -partition  $P := \{(c_k, c_{k+1}) : k = 1, \dots, n-1\}$  we have

$$\text{dom}(\mathcal{E}_{\nu, P})^\perp := \{f \in H_P^1 : \forall g \in \text{dom}(\mathcal{E}_{\nu, P}) : \mathcal{E}_{\nu, P}(f, g) = 0\} = \{f \in H_P^1 : \|f\|_{L_{\nu, P}^2} = 0\}.$$

*Proof.* Pick  $f := (f_j)_{j=1, \dots, n} \in \left(\bigoplus_{k=1}^n \text{dom}(\mathcal{E}_{\nu, [c_k, c_{k+1}]})\right)^\perp$  and  $i \in \{1, \dots, n\}$  with  $\nu([c_i, c_{i+1}]) > 0$ . Then define for  $x \in \text{supp } \nu \cap [c_i, c_{i+1}]$

$$g_x(y) := f_i(x) \frac{(y - c_i)}{x - c_i} \mathbb{1}_{[c_i, x]} + f_i(x) \frac{c_{i+1} - y}{c_{i+1} - x} \mathbb{1}_{(x, c_{i+1}]}$$

Hence,

$$\begin{aligned} 0 &= \mathcal{E}_{\nu, [c_i, c_{i+1}]}(f_i, g_x) = \int_{[c_i, x]} \nabla_\Lambda f_i \nabla_\Lambda g_x \, d\Lambda + \int_{(x, c_{i+1}]} \nabla_\Lambda f_i \nabla_\Lambda g_x \, d\Lambda \\ &= \frac{f_i(x)}{x - c_i} (f_i(x) - f_i(c_i)) + \frac{-f_i(x)}{c_{i+1} - x} (f_i(c_{i+1}) - f_i(x)) = \frac{f_i(x)^2}{x - c_i} + \frac{f_i(x)^2}{c_{i+1} - x}, \end{aligned}$$

and consequently, for all  $x \in \text{supp } \nu \cap (c_i, c_{i+1})$ , we have  $f_i(x) = 0$ ; in particular,  $\|f\|_{L_{\nu, P}^2} = 0$ . On the other hand, for  $f \in \{g \in H_P^1 : \|g\|_{L_{\nu, P}^2} = 0\}$  and each  $k \in \{1, \dots, n\}$ ,  $f_k$  vanishes  $\nu|_{(c_k, c_{k+1})}$ -a. e. and, using the continuity of  $f_k$ , we obtain  $f_k = 0$  on  $\text{supp } \nu|_{[c_i, c_{i+1}]}$ . To simplify notation, assume  $c_k, c_{k+1} \in \text{supp } \nu$ . Now, with  $\bigcup_{i \in I} (a_i, b_i) = [c_k, c_{k+1}] \setminus \text{supp } \nu|_{[c_k, c_{k+1}]}$ , one easily verifies that  $\nabla_\Lambda f_k = \sum_{i \in I} \mathbb{1}_{(a_i, b_i)} \nabla_\Lambda f_k \in L_\Lambda^2$  and we obtain for all  $g \in \text{dom}(\mathcal{E}_{\nu, [c_k, c_{k+1}]})$ ,

$$\begin{aligned} \mathcal{E}_{\nu, [c_k, c_{k+1}]}(f_k, g) &= \int_{[c_k, c_{k+1}]} \nabla_\Lambda f_k \nabla_\Lambda g \, d\Lambda \\ &= \sum_{i \in I} \int_{(a_i, b_i)} \nabla_\Lambda f_k \nabla_\Lambda g \, d\Lambda \\ &= \sum_{i \in I} \frac{g(b_i) - g(a_i)}{b_i - a_i} (f_k(b_i) - f_k(a_i)) = 0. \end{aligned}$$

Since  $k$  was arbitrary the definition of  $\mathcal{E}_{\nu, P}$  gives  $f \in \text{dom}(\mathcal{E}_{\nu, P})^\perp$ .  $\square$

The following extended min-max principle will be crucial for the proof of sub- and superadditivity, and it seems to have been used implicitly in earlier work along these lines.

**Proposition 2.13.** For a  $\nu$ -partition  $P := ((c_k, c_{k+1}]: k = 1, \dots, n)$  and all  $i \in \mathbb{N}$ , we have

$$\begin{aligned} \lambda_i(\mathcal{E}_{\nu, P}) &= \inf \left\{ \sup \left\{ \frac{\mathcal{E}_{\nu, P}(\psi, \psi)}{\langle \psi, \psi \rangle_{\nu, P}} : \psi \in G^\star \right\} : G \prec_i \text{dom}(\mathcal{E}_{\nu, P}) \right\} \\ &= \inf \left\{ \sup \left\{ \frac{\mathcal{E}_{\nu, P}(\psi, \psi)}{\langle \psi, \psi \rangle_{\nu, P}} : \psi \in G^\star \right\} : G \prec_i H_P^1 \right\}. \end{aligned}$$

*Proof.* The part ‘ $\geq$ ’ follows from the inclusion  $\mathcal{D} := \text{dom}(\mathcal{E}_{\nu, P}) = \bigoplus_{k=1}^n \text{dom} \mathcal{E}_{\nu, [c_k, c_{k+1}]} \subset \bigoplus_{k=1}^n H_0^1(c_k, c_{k+1})$ . For the reverse inequality we consider an  $i$ -dimensional subspace  $G = \text{span}(f_1, \dots, f_i) \subset \bigoplus_{k=1}^n H_0^1(c_k, c_{k+1})$ . There exists a unique decomposition  $f_j = f_{1,j} + f_{2,j}$  with  $f_{1,j} \in \mathcal{D}$  and  $f_{2,j} \in \mathcal{D}^\perp$ ,  $j = 1, \dots, i$ . Suppose that  $(f_{1,j})_j$  are not linearly independent, then there exists a non-zero element  $g \in G \cap \mathcal{D}^\perp$ . To see this fix  $(\lambda_1, \dots, \lambda_n) \neq (0, \dots, 0)$  with  $\lambda_1 f_{1,1} + \dots + \lambda_i f_{1,i} = 0$ . Then

$$\underbrace{\lambda_1 (f_{1,1} + f_{1,2}) + \dots + \lambda_i (f_{1,i} + f_{2,i})}_{\in G^\star} = \underbrace{\lambda_1 f_{2,1} + \dots + \lambda_i f_{2,i}}_{\in \text{dom}(\mathcal{E}_{\nu, P})^\perp} =: g.$$

Using  $\mathcal{D}^\perp \subset \{f \in H_P^1 : \|f\|_{L_{\nu, P}^2} = 0\}$  by Lemma 2.12 and  $\mathcal{E}_{\nu, P}(g, g) > 0$ , we get in this case

$$\sup \left\{ \frac{\mathcal{E}_{\nu, P}(\psi, \psi)}{\langle \psi, \psi \rangle_{\nu, P}} : \psi \in G^\star \right\} = \infty.$$

Otherwise, for every vector  $(a_j) \in \mathbb{R}^i \setminus \{0\}$ , we have

$$\begin{aligned} & \frac{\mathcal{E}_{\nu, P}(\sum_j a_j f_{1,j} + \sum_j a_j f_{2,j}, \sum_j a_j f_{1,j} + \sum_j a_j f_{2,j})}{\langle \sum_j a_j f_{1,j} + \sum_j a_j f_{2,j}, \sum_j a_j f_{1,j} + \sum_j a_j f_{2,j} \rangle_{\nu, P}} \\ &= \frac{\mathcal{E}_{\nu, P}(\sum_j a_j f_{1,j}, \sum_j a_j f_{1,j}) + \mathcal{E}_{\nu, P}(\sum_j a_j f_{2,j}, \sum_j a_j f_{2,j})}{\langle \sum_j a_j f_{1,j}, \sum_j a_j f_{1,j} \rangle_{\nu, P}} \\ &\geq \frac{\mathcal{E}_{\nu, P}(\sum_j a_j f_{1,j}, \sum_j a_j f_{1,j})}{\langle \sum_j a_j f_{1,j}, \sum_j a_j f_{1,j} \rangle_{\nu, P}}. \end{aligned}$$

Note that  $\text{span}(f_{1,1}, \dots, f_{1,i}) \subset \text{dom}(\mathcal{E}_{\nu, P})$  is also  $i$ -dimensional subspace in  $L_{\nu, P}^2$  by our definition of  $\text{dom}(\mathcal{E}_{\nu, P})$ . Hence, in any case the reverse inequality follows.  $\square$

**Lemma 2.14.** Let  $\theta, \vartheta$  Borel measures on  $(0, 1)$  with  $\text{card}(\text{supp } \vartheta) = \text{card}(\text{supp } \theta) = \infty$ . Then for each  $i \in \mathbb{N}$ ,

$$\lambda_{\vartheta+\theta}^i \leq \lambda_\theta^i,$$

and for all  $x \geq 0$

$$\max(N_\vartheta(x), N_\theta(x)) \leq N_{\vartheta+\theta}(x).$$

*Proof.* For all  $f \in H_0^1(a, b) \setminus \{0\}$ , we have

$$\frac{\int (\nabla_\Lambda f)^2 d\Lambda}{\int f^2 d(\theta + \vartheta)} \leq \frac{\int (\nabla_\Lambda f)^2 d\Lambda}{\int f^2 d\vartheta}, \quad \frac{\int (\nabla_\Lambda f)^2 d\Lambda}{\int f^2 d(\theta + \vartheta)} \leq \frac{\int (\nabla_\Lambda f)^2 d\Lambda}{\int f^2 d\theta}$$

and the claim follows from Proposition 2.13.  $\square$

*Remark 2.15.* We can reformulate the above lemma as follows. If we assume  $\theta - \vartheta$  is again a positive measure, then, for each  $i \in \mathbb{N}$ , we have

$$\lambda_{\theta}^i \leq \lambda_{\vartheta}^i,$$

and for all  $x \geq 0$ ,

$$N_{\theta}(x) \leq N_{\vartheta}(x).$$

Now we are in the position also to state the proof of Proposition 1.13.

*Proof of Proposition 1.13.* For  $\mathcal{E}_v(\varphi, \psi) = \int_{[0,1]} \nabla_{\Lambda} \varphi \nabla_{\Lambda} \psi \, d\Lambda$  using the bi-Lipschitz property, we have

$$\begin{aligned} \frac{\mathcal{E}_{v \circ g^{-1}, [a,b]}(\psi, \psi)}{\langle \psi, \psi \rangle_{v \circ g^{-1}}} &= \frac{\int_{[a,b]} (\nabla_{\Lambda} \psi)^2 \, d\Lambda}{\langle \psi \circ g, \psi \circ g \rangle_v} = \frac{\int_{[0,1]} ((\nabla_{\Lambda} \psi) \circ g)^2 |\nabla_{\Lambda} g| \, d\Lambda}{\langle \psi \circ g, \psi \circ g \rangle_v} \\ &= \frac{\int_{[0,1]} (\nabla_{\Lambda} (\psi \circ g))^2 (|\nabla_{\Lambda} g|)^{-1} \, d\Lambda}{\langle \psi \circ g, \psi \circ g \rangle_v} \asymp \frac{\mathcal{E}_{v, [0,1]}(\psi \circ g, \psi \circ g)}{\langle \psi \circ g, \psi \circ g \rangle_v}. \end{aligned}$$

Since  $\Phi : H_0^1(a, b) \rightarrow H_0^1(0, 1)$ ,  $\Phi(f) := f \circ g$  defines a linear bijection with  $\nabla_{\Lambda} \Phi(f) = (\nabla_{\Lambda} f) \circ g |\nabla_{\Lambda} g|$  and inverse  $\Phi^{-1}(f) = f \circ g^{-1}$ . We see that for any linear subspace  $G \subset H_0^1(a, b)$  with  $\dim(G) = i$  with respect to  $\mathcal{E}_{v, [0,1]}$  if, and only if

$$\dim(\Phi(G)) = i$$

with respect to  $\mathcal{E}_{v \circ g^{-1}, [a,b]}$ . The extended min-max principle from Proposition 2.13 proves the claim.  $\square$

**Proposition 2.16.** For a  $v$ -partition  $P := ((c_k, c_{k+1}]: k = 1, \dots, n)$  and all  $x \geq 0$ , we have

$$\sum_{k=1}^n N_{v, [c_k, c_{k+1}]}(x) \leq N_v(x) \leq \sum_{k=1}^n N_{v, [c_k, c_{k+1}]}(x) + n - 2.$$

*Proof.* For the  $i$ -th eigenvalue we have with Proposition 2.13

$$\begin{aligned} \lambda_i(\mathcal{E}_{v, P}) &= \inf \left\{ \sup \left\{ \frac{\sum_k \mathcal{E}_{v, [c_k, c_{k+1}]}(\psi_k, \psi_k)}{\sum_k \langle \psi_k, \psi_k \rangle_{v|_{(c_k, c_{k+1}]}}} : (\psi_k) \in G^{\star} \right\} : G \prec_i \text{dom}(\mathcal{E}_{v, P}) \right\} \\ &\geq \inf \left\{ \sup \left\{ \frac{\mathcal{E}_v(\psi, \psi)}{\langle \psi, \psi \rangle_v} : \psi \in G^{\star} \right\} : G \prec_i H_0^1(a, b) \right\} \\ &= \inf \left\{ \sup \left\{ \frac{\mathcal{E}_v(\psi, \psi)}{\langle \psi, \psi \rangle_v} : \psi \in G^{\star} \right\} : G \prec_i \text{dom}(\mathcal{E}_v) \right\} = \lambda_i(\mathcal{E}_v). \end{aligned}$$

This gives

$$\sum_{k=1}^n N_{v, [c_k, c_{k+1}]}(x) = N_{v, P}(x) \leq N_v(x).$$

For the second inequality we make use of  $H_{0,P}^1 := \{f \in H_0^1(a, b) : f(c_k) = 0, k = 1, \dots, n\} \simeq H_P^1$  and  $\dim(H_0^1(a, b)/H_{0,P}^1) = n - 2$ . We now estimate as in the proof of [Kig01, Theorem 4.1.7.]:

$$\begin{aligned}
\lambda_{i+n-2}(\mathcal{E}_\nu) &= \inf \left\{ \sup \left\{ \frac{\mathcal{E}_\nu(\psi, \psi)}{\langle \psi, \psi \rangle_\nu} : \psi \in G^\star \right\} : G \subset_{i+n-2} H_0^1(a, b) \right\} \\
&\geq \inf \left\{ \sup \left\{ \frac{\mathcal{E}_\nu(\psi, \psi)}{\langle \psi, \psi \rangle_\nu} : \psi \in G^\star \right\} : G \subset_i H_{0,P}^1 \right\} \\
&= \inf \left\{ \sup \left\{ \frac{\sum \mathcal{E}_{\nu, [c_k, c_{k+1}]}(\psi_k, \psi_k)}{\sum \langle \psi_k, \psi_k \rangle_{\nu|_{(c_k, c_{k+1})}}} : (\psi_k) \in G^\star \right\} : G \subset_i H_P^1 \right\} \\
&= \inf \left\{ \sup \left\{ \frac{\sum \mathcal{E}_{\nu, [c_k, c_{k+1}]}(\psi_k, \psi_k)}{\sum \langle \psi_k, \psi_k \rangle_{\nu|_{(c_k, c_{k+1})}}} : (\psi_k) \in G^\star \right\} : G \subset_i \text{dom}(\mathcal{E}_{\nu, P}) \right\} \\
&= \lambda_i(\mathcal{E}_{\nu, P}).
\end{aligned}$$

□

**Corollary 2.17.** *Let  $\nu$  be a Borel measure with  $\text{supp } \nu \subset (a, b) \subset [c, d]$ . Then*

$$N_{\nu, [a, b]}(x) \leq N_{\nu, [c, d]}(x) \leq N_{\nu, [a, b]}(x) + 2.$$

### 2.3. Neumann boundary conditions

Although we have shown that Dirichlet boundary conditions alone are sufficient to determine spectral asymptotics, for completeness we give the relationship to Neumann boundary conditions, known as Neumann-Dirichlet bracketing. Let  $I_{\nu, a}$  and  $I_{\nu, b}$  denote possibly empty components of  $[a, b] \setminus \text{supp } \nu$  containing  $a$ , respectively  $b$  and set

$$C_\nu^N([a, b]) := \{f \in C_\nu([a, b]) \mid f \text{ is const. on } I_{\nu, a} \text{ and } I_{\nu, b}\}.$$

Note that  $(C_\nu^N([a, b]), \|\cdot\|_\infty)$  is a Banach space. We define the form

$$\mathcal{E}_{\nu, [a, b]}^N(f, g) := \int_{[a, b]} \nabla_\Lambda f \nabla_\Lambda g \, d\Lambda$$

with Neumann domain  $\text{dom}(\mathcal{E}_{\nu, [a, b]}^N) := H^1(a, b) \cap C_\nu^N([a, b])$ . Again  $\mathcal{E}_\nu^N$  is well defined. The set  $\text{dom}(\mathcal{E}_{\nu, [a, b]}^N)$  is dense in  $L_\nu^2$  and equipped it with the inner product  $(f, g)_{\mathcal{E}_{\nu, [a, b]}^N} := \langle f, g \rangle_\nu + \mathcal{E}_{\nu, [a, b]}^N(f, g)$ , defines a Hilbert space.

Using [Dav95, Theorem 4.4.2] or [Kat76, Theorem 2.1., p. 322], there exists a non-negative self-adjoint operator  $\Delta_{\nu, [a, b]}^N$  which is characterized as follows;  $f \in \text{dom}(\mathcal{E}_{\nu, [a, b]}^N)$  lies in the domain  $\text{dom}(\Delta_{\nu, [a, b]}^N)$  of the Laplacian  $\Delta_{\nu, [a, b]}^N$  if and only if there exists  $h \in L_\nu^2$  such that

$$\mathcal{E}_{\nu, [a, b]}(g, f) = \langle g, h \rangle_\nu, \quad \text{for all } g \in \text{dom}(\mathcal{E}_{\nu, [a, b]}^N)$$

with  $h := \Delta_{\nu, [a, b]}^N f$ . Furthermore, let  $N_\nu^N$  denote the eigenvalue counting function of  $\Delta_{\nu, [a, b]}^N$ .

Also in our situation the min-max principle gives the usual Dirichlet-Neumann bracketing.

**Proposition 2.18.** *Let  $\nu$  be a Borel measure on  $(a, b)$ , then, for all  $x \geq 0$ , we have*

$$N_{\nu, [a, b]}(x) \leq N_{\nu, [a, b]}^N(x) \leq N_{\nu, [a, b]}(x) + 2.$$

*Proof.* For  $g \in \text{dom}(\mathcal{E}_\nu)$  we define

$$\widehat{g}(x) := g(x) + \mathbb{1}_{[a, a']}((a' - x)g(a') / (a' - a)) + \mathbb{1}_{(b', b]}((x - b')g(b') / (b - b')) \in \text{dom}(\mathcal{E}_\nu^N).$$

Since  $\mathcal{E}_\nu(g, g) \geq \mathcal{E}_\nu(\widehat{g}, \widehat{g})$  and  $\langle g, g \rangle_\nu = \langle \widehat{g}, \widehat{g} \rangle_\nu$  this gives

$$\begin{aligned} \lambda_i(\mathcal{E}_{\nu, [a, b]}) &= \inf \left\{ \sup \left\{ \frac{\mathcal{E}_{\nu, [a, b]}(g, g)}{\langle g, g \rangle_\nu} : g \in G^\star \right\} : G <_i \text{dom}(\mathcal{E}_{\nu, [a, b]}) \right\} \\ &\geq \inf \left\{ \sup \left\{ \frac{\mathcal{E}_{\nu, [a, b]}(\widehat{g}, \widehat{g})}{\langle \widehat{g}, \widehat{g} \rangle_\nu} : g \in G^\star \right\} : G <_i \text{dom}(\mathcal{E}_{\nu, [a, b]}) \right\} \\ &\geq \inf \left\{ \sup \left\{ \frac{\mathcal{E}_{\nu, [a, b]}(g, g)}{\langle g, g \rangle_\nu} : g \in G^\star \right\} : G <_i \text{dom}(\mathcal{E}_{\nu, [a, b]}^N) \right\} \\ &= \lambda_i(\mathcal{E}_{\nu, [a, b]}^N). \end{aligned}$$

For the reversed inequality we note that the dimension of the quotient vector space  $\text{dom}(\mathcal{E}_{\nu, [a, b]}^N) / (\text{dom}(\mathcal{E}_{\nu, [a, b]}) \cap \text{dom}(\mathcal{E}_{\nu, [a, b]}^N))$  is equal to two. Then using again the proof of [Kig01, Theorem 4.1.7.], we deduce

$$\begin{aligned} \lambda^{i+2}(\mathcal{E}_{\nu, [a, b]}^N) &= \inf \left\{ \sup \left\{ \frac{\mathcal{E}_{\nu, [a, b]}(g, g)}{\langle g, g \rangle_\nu} : g \in G^\star \right\} : G <_{i+2} \text{dom}(\mathcal{E}_{\nu, [a, b]}^N) \right\} \\ &\geq \inf \left\{ \sup \left\{ \frac{\mathcal{E}_{\nu, [a, b]}(g, g)}{\langle g, g \rangle_\nu} : g \in G^\star \right\} : G <_i \text{dom}(\mathcal{E}_{\nu, [a, b]}) \cap \text{dom}(\mathcal{E}_{\nu, [a, b]}^N) \right\} \\ &\geq \inf \left\{ \sup \left\{ \frac{\mathcal{E}_{\nu, [a, b]}(g, g)}{\langle g, g \rangle_\nu} : g \in G^\star \right\} : G <_i \text{dom}(\mathcal{E}_{\nu, [a, b]}) \right\} = \lambda^i(\mathcal{E}_{\nu, [a, b]}). \end{aligned}$$

□

### 3. Optimal partitions and partition entropy

Recall the definition of  $\Pi$ ,  $\Pi_0$ ,  $\mathcal{N}^R(x)$ ,  $\mathcal{N}_m^L(x)$ ,  $\bar{h}_\nu$ ,  $\underline{h}_\nu$ ,  $\underline{h}_\nu^m$ ,  $\bar{h}_\nu^m$  as provided in the introduction.

#### 3.1. Basic observations

In this section we establish a connection of our approach and the classical work of Birman and Solomjak [BS66, BS67, Bor70]. This connection will come into play again in Example 5.7.

Let  $\Upsilon_n$  denote the set of all finite  $\nu$ -partitions consisting of at most  $n \in \mathbb{N}$  half open intervals of  $(a, b]$  with  $\nu(\{a, b\}) = 0$  and let us define

$$\gamma_n := \gamma_{n, \nu} := \inf_{\Xi \in \Upsilon_n} \max_{I \in \Xi} \Lambda(I)\nu(I),$$

and

$$\underline{\alpha}_\nu := \liminf_{n \rightarrow \infty} \frac{\log(\gamma_n)}{-\log(n)}, \quad \bar{\alpha}_\nu := \limsup_{n \rightarrow \infty} \frac{\log(\gamma_n)}{-\log(n)}. \quad (3.1)$$

*Remark 3.1.* If  $\nu$  has an absolutely continuous part, then the limit in (3.1) is equal to 2 and in general, we have that  $\underline{\alpha}_\nu \in [2, \infty]$ .

Note also that in the definition of  $\gamma_n$  for general  $\nu$  (i. e. also allowing for a pure point part) one cannot replace the infimum with the minimum. Indeed, consider the Dirac measure  $\delta_{1/2}$  on  $[0, 1]$ . Then for every  $1/2 > \epsilon > 0$ ,  $\{(0, 1/2 - \epsilon], (1/2 - \epsilon, 1/2 + \epsilon], (1/2 + \epsilon, 1]\}$  is  $\nu$  partition, hence we obtain for  $n \geq 3$  that  $\gamma_n \leq 2\epsilon$ ,  $\gamma_n = 0$  and the minimum does not exist.

**Lemma 3.2.** *For all  $n \in \mathbb{N}$  we have  $\gamma_{2^{n+1}} \leq \frac{1}{2}\gamma_{2^n} < \gamma_{2^n}$ . If  $\nu$  is atomless, then  $\gamma_{2^{n+1}} \leq \frac{1}{4}\gamma_{2^n}$ .*

*Proof.* For  $\Xi \in \Upsilon$ ,  $\text{card } \Xi \leq 2^n$  we can divide each  $I \in \Xi$  into two sub-intervals  $I_1, I_2$  of same length. The new resulting partition denoted by  $\Xi'$  satisfies  $\text{card}(\Xi') \leq 2^{n+1}$  and

$$\max_{I \in \Xi'} \nu(I)\Lambda(I) \leq \frac{1}{2} \max_{I \in \Xi} \nu(I)\Lambda(I) < \max_{I \in \Xi} \nu(I)\Lambda(I).$$

This implies the first claim of the Lemma. For the second statement fix  $I \in \Xi$  and let  $F$  denote the (continuous) distribution function of the normalized restriction of  $\nu$  to  $I$ . Then choose  $t \in (1/4, 1)$  such that  $tF(t) = 1/4$ . An elementary calculation then shows that for such  $t$  we have  $(1-t)(1-F(t)) = 5/4 - t - 1/(4t) \leq 1/4$ . Now we subdivide  $I$  in a left interval  $I_1$  of length  $t\Lambda(I)$  and the remaining right interval  $I_2$ . The resulting partition now fulfills the inequality as above with  $1/2$  replaced by  $1/4$ .  $\square$

*Remark 3.3.* If the measure  $\nu$  is atomless, we deduce from Lemma 3.2 by induction  $\gamma_{2^n} \leq 4^{-n+1}\gamma_1 = 4^{-n+1}\nu([a, b])(b-a)$  and as a simple consequence we find  $\underline{\alpha}_\nu \geq 2$ . This will give rise to the famous upper bound  $1/2$  on the spectral dimension as obtained in [BS66, BS67] also for arbitrary measure  $\nu$ .

**Lemma 3.4.** *We have  $\bar{h}_\nu = 1/\underline{\alpha}_\nu$  and  $\underline{h}_\nu = 1/\bar{\alpha}_\nu$ .*

*Proof.* The proof follows along the same lines as the proof of the elementary Lemma [KZ04, Lemma 2.2]. First note that for  $0 < \epsilon < 1$  we have  $\mathcal{N}^R(1/\epsilon) = \inf \{n \in \mathbb{N} \mid \gamma_n < \epsilon\}$ . By Lemma 3.2 we have that  $(\gamma_{2^n})_n$  is a strictly decreasing null sequences or eventually constant zero. The latter case is immediate. For the first case the strict monotonicity gives as in [KZ04, Lemma 2.2] for  $B(\epsilon) := \inf \{n \in \mathbb{N} \mid \gamma_{2^n} < \epsilon\}$

$$\limsup_{n \rightarrow \infty} \frac{n \log 2}{-\log \gamma_{2^n}} = \limsup_{\epsilon \searrow 0} \frac{B(\epsilon) \log 2}{-\log \epsilon}.$$

The left hand side gives  $1/\underline{\alpha}_\nu$  by squeezing  $m \rightarrow \infty$  on subsequences  $2^{n-1} < m \leq 2^n$  and the right hand side  $\bar{h}_\nu$  by noting that  $2^{B(\epsilon)-1} \leq \mathcal{N}^R(1/\epsilon) \leq 2^{B(\epsilon)}$ . The same result holds for  $\liminf$ .  $\square$

### 3.2. Some global bounds

**Lemma 3.5** (Poincaré inequality). *Let  $\nu$  be a Borel measure on  $(a, b)$ . Then for the first eigenvalue of  $\mathcal{E}_{\nu, [a, b]}$ , we have*

$$\lambda_\nu^1 \geq \frac{1}{\nu([a, b])\Lambda([a, b])}.$$

*Proof.* Here, we follow [Min20, Lemma 4.6.]. Let  $f$  be the eigenfunction with eigenvalue  $\lambda_\nu^1$  with  $\|f\|_\nu = 1$ . Then by the Cauchy-Schwarz inequality, for all  $x \in [a, b]$ , we have

$$f(x)^2 = |f(x) - f(a)|^2 = \left| \int_{[a,x]} \nabla_\Lambda f \, d\Lambda \right|^2 \leq \Lambda([a, b]) \|\nabla_\Lambda f\|_\Lambda^2$$

Now, integrating with respect to  $\nu$  gives

$$1 \leq \nu([a, b]) \Lambda([a, b]) \|\nabla_\Lambda f\|_\Lambda^2 = \nu([a, b]) \Lambda([a, b]) \lambda_\nu^1.$$

□

In the following we will use the approximation theory of Birman and Solomjak developed in [BS67, Theorem 2.1, Theorem 2.2], see also [BS66, BB71]. Let  $\nu$  be a Borel measure on  $[a, b]$ . For every  $n \in \mathbb{N}$  there exists a partition of half open intervals of  $(a, b]$  denoted by  $\Xi$  such that  $\text{card}(\Xi) \leq n$  and

$$\max_{I \in \Xi} \Lambda(I) \nu(I) \leq C_1 n^{-2} \nu([a, b])(b-a),$$

where  $C_1 > 0$  is independent of  $\nu$ . Assuming now  $\nu$  to be atomless the constant is given by  $C_1 = 1$ . Putting  $n := \lceil \sqrt{x(b-a)\nu([a, b])} \rceil$  and note that by Lemma 3.5 for every  $C \in \Xi$  with  $\nu(C) > 0$ , we have

$$\lambda_{\nu, C}^1 \geq \frac{1}{\nu(C)\Lambda(C)} \geq \frac{1}{\max_{\Gamma \in \Xi} \Lambda(\Gamma)\nu(\Gamma)} \geq \frac{n^2}{\nu([a, b])(b-a)},$$

hence applying Proposition 2.16 delivers

$$N_\nu(x) \leq \sum_{C \in \Xi, \nu(C) > 0} N_{\nu, \Lambda, C}(x) + n \leq n \leq 1 + \sqrt{\nu([a, b])(b-a)x},$$

where we used that  $\lambda_{\nu, C}^1 \geq n^2 (C_1 \nu([a, b])(b-a))^{-1} > x$  by the choice of  $n$ .

The estimate can be improved in the case that  $\nu$  is singular with respect to  $\Lambda$ . To prove this we need the following refinement which is developed in [Bor71]:

*There is a constant  $C_1 > 0$  such that for every measure  $\nu$  singular with respect to  $\Lambda$  and every  $\varepsilon > 0$  there exists  $n_{\nu, \varepsilon} \in \mathbb{N}$  such that for every  $n > n_{\nu, \varepsilon}$  there is a partition of half open intervals of  $(a, b]$  denoted by  $\Xi$  such that  $\text{card}(\Xi) \leq n$  and*

$$\max_{I \in \Xi} \Lambda(I) \nu(I) \leq \varepsilon C_1 \frac{\nu([a, b])(b-a)}{(1 - n_{\nu, \varepsilon}/n)^2 n^2}.$$

Hence, for  $\nu$  atomless and singular with respect to  $\Lambda$ , and for every  $\varepsilon > 0$  and all  $x > 0$  large enough, we have

$$N_\nu(x) \leq \sqrt{\varepsilon x}.$$

In particular,  $\lim_{x \rightarrow \infty} x^{-1/2} N_\nu(x) = 0$ .

**Lemma 3.6.** For the subinterval  $C = [c, d]$  of  $[a, b]$  assume  $\nu(\{c, d\}) = 0$  and fix  $\alpha_1, \alpha_2 \in C$  such that  $c < \alpha_1 < \alpha_2 < d$  and  $\nu([\alpha_1, \alpha_2]) > 0$ . Then

$$\lambda_{\nu, C}^1 \leq \frac{\left(\frac{\alpha_2 - \alpha_1}{\alpha_1 - c} + \frac{\alpha_2 - \alpha_1}{d - \alpha_2}\right)}{\nu([\alpha_1, \alpha_2]) \Lambda([\alpha_1, \alpha_2])}.$$

In particular, if  $\nu(\langle C \rangle_m) > 0$  for some  $m > 1$ , then

$$\lambda_{\nu, C}^1 \leq \frac{4/(m-1)}{\nu(\langle C \rangle_m) \Lambda(\langle C \rangle_m)}.$$

*Proof.* Define

$$f(x) := \frac{x-c}{\alpha_1-c} \mathbb{1}_{[c, \alpha_1]} + \mathbb{1}_{[\alpha_1, \alpha_2]} + \frac{d-x}{d-\alpha_2} \mathbb{1}_{(\alpha_2, d]},$$

then  $f \in H_0^1(c, d)$  and the min-max principle gives

$$\lambda_{\nu, C}^1 \leq \frac{\int (f')^2 d\Lambda}{\int f^2 d\nu} = \frac{\left(\frac{1}{\alpha_1-c} + \frac{1}{d-\alpha_2}\right)}{\int f^2 d\nu} \leq \frac{(\alpha_2 - \alpha_1) \left(\frac{1}{\alpha_1-c} + \frac{1}{d-\alpha_2}\right)}{\nu([\alpha_1, \alpha_2]) \Lambda([\alpha_1, \alpha_2])}.$$

For  $\alpha_1 = \frac{1}{2} \left(c + d - \frac{d-c}{m}\right)$ ,  $\alpha_2 = \frac{1}{2} \left(c + d + \frac{d-c}{m}\right)$  and provided  $\nu(\langle C \rangle_m) > 0$ , one obtains  $\lambda_{\nu, \Lambda, C}^1 \leq 4/(m-1) (\nu(\langle C \rangle_m) \Lambda(\langle C \rangle_m))^{-1}$ .  $\square$

**Proposition 3.7.** Let  $\nu$  be a Borel measure on  $(a, b)$ . Then for each  $m > 1$  and all  $x > 0$ , we have

$$\mathcal{N}_m^L(x) \leq N_\nu(x) \leq \mathcal{N}^R(x).$$

In particular,

$$\underline{h}_\nu^m \leq \underline{s}_\nu \leq \bar{s}_\nu \leq \bar{h}_\nu \text{ and } \underline{s}_\nu \leq \underline{h}_\nu \text{ and } \bar{h}_\nu^m \leq \bar{s}_\nu$$

and if

$$\sup_{m>1} \underline{h}_\nu^m = \bar{h}_\nu,$$

then the spectral dimension exists and equals  $\bar{h}_\nu$ .

*Proof.* This follows immediately from the sub- and superadditivity of the eigenvalue counting functions (Proposition 2.16) in tandem with Lemma 3.5 and Lemma 3.6.  $\square$

#### 4. $L^q$ -spectrum and coarse multifractal formalism

In this section we establish a connection between the spectral dimension,  $L^q$ -spectrum and the coarse multifractal formalism.

4.1. The  $L^q$ -spectrum

In this section we collect some important facts about the  $L^q$ -spectrum providing proofs only for those which are not straight forward or have not been covered in the literature.

We let  $\nu$  be an arbitrary finite Borel measure with bounded support. For  $k \in \mathbb{Z}$  and  $n \in \mathbb{N}$  we let  $A_k^n := (k2^{-n}, (k+1)2^{-n}]$  and set  $\mathcal{D}_{\nu,n} := \{A_k^n : k \in \mathbb{Z}, \nu(A_k^n) > 0\}$ ,  $\mathcal{D}_\nu := \bigcup_n \mathcal{D}_{\nu,n}$ . For  $q \in \mathbb{R}$ , the  $L^q$ -spectrum of  $\nu$  is defined by

$$\beta_\nu(q) := \limsup_{n \rightarrow \infty} \beta_n(q) \quad \text{with} \quad \beta_n^\nu(q) := \beta_n(q) := \frac{1}{n \log 2} \log \sum_{C \in \mathcal{D}_{\nu,n}} \nu(C)^q.$$

Let us begin with some basic facts.

**Fact 4.1.** *The function  $\beta_\nu$  will not alter when we take  $d$ -adic intervals instead of dyadic ones (see e. g. [Rie95, Proposition 2 and Remarks, p. 466] or [Rie93, Proposition 1.6.] and note that the definition in [Rie93, Proposition 1.6.] coincides with our definition for  $q \geq 0$ ). More precisely, for fixed  $\delta > 0$ , set*

$$G_{\nu,\delta} := G_\delta := \{(\delta l, (l+1)\delta] \mid l \in \mathbb{N}, \nu((\delta l, (l+1)\delta]) > 0\}$$

and let  $(\delta_n)$  be an admissible sequence, i. e.  $\delta_n \in (0, 1)^\mathbb{N}$ ,  $\delta_n \rightarrow 0$  and there exists a constant  $C > 0$  such that  $C\delta_n \leq \delta_{n+1} \leq \delta_n$ , for all  $n \in \mathbb{N}$ . Then for  $q \geq 0$ ,

$$\limsup_{\delta \downarrow 0} \frac{1}{-\log(\delta)} \log \sum_{C \in G_\delta} \nu(C)^q = \limsup_{m \rightarrow \infty} \frac{1}{-\log(\delta_m)} \log \sum_{C \in G_{\delta_m}} \nu(C)^q.$$

In particular, for  $\delta_m := 2^{-m}$ , the above expression equals  $\beta_\nu(q)$ .

**Fact 4.2.** *For  $c > 0$  we have  $\beta_{c\nu} = \beta_\nu$ . Therefore, we can assume without loss of generality that  $\nu$  is a probability measure.*

**Fact 4.3.** *The function  $\beta_\nu$  is — as a pointwise limit superior of convex function — again convex and we have (see also [Rie95, Corollary ])*

$$\beta_\nu(0) = \overline{\dim}_M(\nu) =: \delta_\nu^* \quad \text{and} \quad \beta_\nu(1) = 0.$$

Hence, the left, respectively right, derivative in 1 exists and its negative value will be denoted by  $\bar{\delta}_\nu := \lim_{q \nearrow 1} -\beta_\nu(q) / (1 - q)$  and  $\underline{\delta}_\nu := \lim_{q \searrow 1} -\beta_\nu(q) / (1 - q)$ . For later use we also define the right derivative of  $\beta$  in 0 denoted by  $\varrho_\nu := \lim_{t \searrow 0} (\beta_\nu(t) - \delta_\nu^*) / t$ . Note that in many situations we have  $\bar{\delta}_\nu = \dim_H(\nu)$ . The function  $\beta_\nu$  is non-increasing and non-negative on  $\mathbb{R}_{<1}$  and  $\liminf_n \beta_n(q) \geq -q$  for all  $q \geq 0$ .

**Fact 4.4.** *Since  $\beta_\nu$  is a proper convex function, its Legendre transform exists, given by*

$$\widehat{\beta}_\nu(\alpha) := \inf_{q \in \mathbb{R}} \beta_\nu(q) + \alpha q.$$

**Fact 4.5.** *For any two finite Borel measures  $\mu_1, \mu_2$  with bounded support and for  $q \geq 0$ , we have*

$$\beta_{\mu_1 + \mu_2}(q) = \max(\beta_{\mu_1}, \beta_{\mu_2})(q).$$

If  $\beta_{\mu_1}(q) \geq \beta_{\mu_2}(q)$  and  $\beta_{\mu_1}(q)$  exists as a limit, then so does  $\beta_{\mu_1 + \mu_2}(q) = \beta_{\mu_1}(q)$ .

*Proof.* We readily check, for  $q \geq 0$ ,

$$\sum_{C \in \mathcal{D}_{\mu_1, n}} (\mu_1(C))^q \leq \sum_{C \in \mathcal{D}_{\mu_1 + \mu_2, n}} (\mu_1(C) + \mu_2(C))^q \leq 2^q \left( \sum_{C \in \mathcal{D}_{\mu_1, n}} \mu_1(C)^q + \sum_{C \in \mathcal{D}_{\mu_2, n}} \mu_2(C)^q \right)$$

implying our claim.  $\square$

**Fact 4.6.** *If  $\nu$  has an atomic part, then  $\beta_\nu(q) = 0$ , for all  $q \geq 1$ .*

*Proof.* Assume that  $\nu$  has an atom in  $x_0 \in (0, 1)$  and let  $q > 1$ . Then for every  $n \in \mathbb{N}$ , we have that  $0 < \nu(\{x_0\})^q \leq \sum_{C \in \mathcal{D}_{\nu, n}} \nu(C)^q$  and hence  $0 \leq \beta_\nu(q) \leq \beta_\nu(1) = 0$ .  $\square$

**Fact 4.7.** *If  $\delta_\nu^* = 0$ , then  $\beta_\nu(q) = 0$ , for all  $q \geq 0$  and  $\dim_H(\nu) = \dim_P(\nu) = \underline{\delta}_\nu = \bar{\delta}_\nu = 0$ .*

*Proof.* Note that  $\beta$  is a proper convex non-increasing function with  $\beta_\nu(0) = \beta_\nu(1) = 0$ . This implies the first claim as well as  $\underline{\delta}_\nu = \bar{\delta}_\nu = 0$ . Further, recall that

$$\dim_H(\nu) = \inf \left\{ \dim_H(A) : \nu(A^c) = 0 \right\} \text{ and } \dim_P(\nu) = \inf \left\{ \dim_P(A) : \nu(A^c) = 0 \right\}.$$

Since  $\nu((\text{supp } \nu)^c) = 0$  and  $\dim_H(\text{supp } \nu) \leq \dim_P(\text{supp } \nu) \leq \overline{\dim}_M(\text{supp } \nu) = 0$  the second claim follows.  $\square$

Recall from the introduction the definition of  $\bar{q}_\nu := \limsup_{n \rightarrow \infty} q_n^\nu$ , where  $q_n^\nu$  denotes the unique fixed point of  $\beta_n^\nu$ .

**Fact 4.8.** *We have*

$$\begin{aligned} \bar{q}_\nu &= \inf \{ q > 0 : \beta_\nu(q) - q \leq 0 \} \\ &= \inf \left\{ q > 0 : \limsup_{n \rightarrow \infty} \frac{1}{n} \log \sum_{C \in \mathcal{D}_{\nu, n}} (\nu(C) \wedge (C))^q \leq 0 \right\} \\ &= \inf \left\{ q > 0 : \sum_{C \in \mathcal{D}_\nu} (\nu(C) \wedge (C))^q < \infty \right\} \\ &= \sup_{\alpha \geq 0} \frac{\widehat{\beta}_\nu(\alpha)}{1 + \alpha}. \end{aligned}$$

*If  $\bar{q}_\nu > 0$ , then  $\bar{q}_\nu$  is the unique fixed point of  $\beta_\nu$  in  $(0, 1)$ , i. e.  $\beta_\nu(\bar{q}_\nu) = \bar{q}_\nu \in (0, 1)$ .*

*Proof.* In view of Fact 4.2 we assume without loss of generality that  $\nu$  is a probability measure. Let us begin with the first equality assuming  $\inf \{ q > 0 : \beta_\nu(q) - q \leq 0 \} > 0$ . Recall that  $\beta_\nu$  defines a convex and real valued function on  $(0, 1]$  and hence  $\beta_\nu$  is also continuous on  $(0, \infty)$  (see e. g. [Rie95, Lemma 5]). Since  $\beta_\nu$  is also decreasing with  $\lim_{t \searrow 0} \beta_\nu(t) > 0$  and  $\beta_\nu(1) = 0$  there exists a unique fixed point of  $\beta_\nu$  which coincides with  $\inf \{ q > 0 : \beta_\nu(q) - q \leq 0 \}$ . This fixed point also coincides with  $\bar{q}_\nu$ , which can be checked easily by contradiction. This proves also the last claim. If, on the other hand,




Figure 4.1: Moment generating function  $\beta_\nu$  for  $\nu$  with (dashed) tangent to  $\beta_\nu$  of slope  $-\alpha$  in  $(q, \beta_\nu(q))$  which intersects the  $y$ -axis in its Legendre transform  $\widehat{\beta}_\nu(\alpha)$  and the bisector of the first quadrant in  $\widehat{\beta}_\nu(\alpha)/(1+\alpha)$ . Here  $\nu$  is chosen to be the  $(0.05, 0.95)$ -Salem measure with full support  $\text{supp } \nu = [0, 1]$ . The intersection of  $\beta_\nu$  with the  $y$ -axis gives the Minkowski dimension of  $\text{supp } \nu$ , namely 1, and the intersection with the (dotted) tangent to  $\beta_\nu$  in  $(0, 1)$  gives the Hausdorff dimension  $\dim_H(\nu)$  of the measure which equals  $(0.05 \log(0.05) + 0.95 \log(0.95)) / \log(2)$ . The spectral dimension  $s_\nu$  is equal to the maximum over all  $\widehat{\beta}_\nu(\alpha)/(1+\alpha)$  given by  $\bar{q}_\nu$ .

$\inf \{q > 0 : \beta_\nu(q) - q \leq 0\} = 0$ , then  $\beta_\nu(q) = 0$  for all  $q > 0$  and again by contradiction  $\bar{q}_\nu = 0$ . This proves the first equality.

The fact that  $\limsup_{n \rightarrow \infty} (n \log(2))^{-1} \log \sum_{C \in \mathcal{D}_{\nu, n}} (\nu(C) \Lambda(C))^q = \beta_\nu(q) - q$  for all  $q \in \mathbb{R}$  gives the second equality.

The third equality follows similar to [JKL14]. Let us assume  $\bar{q}_\nu > 0$  and fix  $0 < q < \bar{q}_\nu$ . Then by the second characterization, we have  $\limsup_{n \rightarrow \infty} n^{-1} \log \sum_{C \in \mathcal{D}_{\nu, n}} (\nu(C) \Lambda(C))^q = \epsilon > 0$  and in particular, for infinitely many  $n$ , for  $(n_j)$  say, we have  $\sum_{C \in \mathcal{D}_{\nu, n_j}} (\nu(C) \Lambda(C))^q > e^{n_j \epsilon / 2}$ . Consequently,

$$\sum_{C \in \mathcal{D}_\nu} (\nu(C) \Lambda(C))^q \geq \sum_{j=0}^{\infty} \sum_{C \in \mathcal{D}_{\nu, n_j}} (\nu(C) \Lambda(C))^q \geq \sum_{j=0}^{\infty} e^{n_j \epsilon / 2} = \infty.$$

Hence,  $\bar{q}_\nu \leq \inf \{q > 0 : \sum_{C \in \mathcal{D}_\nu} (\nu(C) \Lambda(C))^q < \infty\}$ , which is trivially fulfilled if  $\bar{q}_\nu = 0$ . For the reverse inequality and  $\bar{q}_\nu \geq 0$  arbitrary, fix  $\epsilon > 0$  and  $q > \bar{q}_\nu$  such that for all

$n > N$ , we have  $\sum_{C \in \mathcal{D}_{v,n}} (\nu(C) \Lambda(C))^q < 2^{n\epsilon/2}$ . Then

$$\begin{aligned} \sum_{C \in \mathcal{D}_v} (\nu(C) \Lambda(C))^{q+\epsilon} &\leq \sum_{C \in \mathcal{D}} (\nu(C) \Lambda(C))^q \Lambda(C)^\epsilon = \sum_{n=0}^{\infty} \sum_{C \in \mathcal{D}_{v,n}} (\nu(C) \Lambda(C))^q 2^{-n\epsilon} \\ &\leq \left( \sum_{n=0}^N \sum_{C \in \mathcal{D}_{v,n}} (\nu(C) \Lambda(C))^q 2^{-n\epsilon} + \sum_{n=N}^{\infty} \sum_{C \in \mathcal{D}_{v,n}} (\nu(C) \Lambda(C))^q 2^{-n\epsilon} \right) \\ &\leq \left( \sum_{n=0}^N \sum_{C \in \mathcal{D}_{v,n}} (\nu(C) \Lambda(C))^q 2^{-n\epsilon} + \sum_{n=N}^{\infty} 2^{n\epsilon/2} 2^{-n\epsilon} \right) < \infty. \end{aligned}$$

Hence  $\inf \left\{ q > 0 : \sum_{C \in \mathcal{D}_v} (\nu(C) \Lambda(C))^q < \infty \right\} \leq \bar{q}_v$ .

Now we prove the last equality. Note that, for every  $q' > \bar{q}_v$ , we have  $\beta_v(q') \leq q'$  and therefore,

$$\sup_{\alpha \geq 0} \frac{\widehat{\beta}_v(\alpha)}{1 + \alpha} \leq \sup_{\alpha \geq 0} \frac{q' + q'\alpha}{1 + \alpha} = q'.$$

This shows  $\sup_{\alpha \geq 0} \widehat{\beta}_v(\alpha) / (1 + \alpha) \leq \bar{q}_v$  and equality holds if  $\bar{q}_v = 0$ . Assuming  $\bar{q}_v > 0$ , we already know that  $\bar{q}_v$  is the unique fixed point of  $\beta_v$ . If  $-a$  is in the subdifferential  $\partial\beta_v(\bar{q}_v)$  of  $\beta$  in  $\bar{q}_v = \beta_v(\bar{q}_v)$  with  $a \geq 0$ , then (see Fig. 4.1)

$$\sup_{\alpha \geq 0} \frac{\widehat{\beta}_v(\alpha)}{1 + \alpha} \geq \frac{\inf_{q \in \mathbb{R}} \beta_v(q) + aq}{1 + a} \geq \frac{\beta_v(\bar{q}_v) - a(q - \bar{q}_v) + aq}{1 + a} = \bar{q}_v.$$

□

**Proposition 4.9** ([Rie93],[Rie95, Prop. 2]). *Let  $\nu$  be a Borel measure on  $[0, 1]$  and  $g : [0, 1] \rightarrow [a, b]$  be a bi-Lipschitz mapping. Then*

$$\beta_\nu = \beta_{\nu \circ g^{-1}}.$$

*Proof.* For completeness we give a short proof. First, by our assumption we have that there exist  $c_1, c_2 > 0$  such that for all  $x, y \in [0, 1]$

$$c_1 |x - y| \leq |g(x) - g(y)| \leq c_2 |x - y|.$$

For  $C \in G_{\nu \circ g^{-1}, 2^{-n}}$ ,

$$\frac{1}{c_2} \text{diam}(C) \leq \text{diam}(g^{-1}(C)) \leq \frac{1}{c_1} \text{diam}(C) = \frac{1}{c_1 2^n}.$$

This shows that there are at most two elements  $C_1, C_2$  of  $G_{\nu, (2^n c_1)^{-1}}$  intersecting  $g^{-1}(C)$ . Furthermore, each element of  $G_{\nu, (2^n c_1)^{-1}}$  that intersects  $g^{-1}(C)$  can intersect at most  $\lceil c_2/c_1 \rceil$  other element  $g^{-1}(C')$  with  $C' \in G_{\nu \circ g^{-1}, 2^{-n}}$ . This implies

$$\sum_{C \in G_{\nu \circ g^{-1}, 2^{-n}}} \nu(g^{-1}(C))^q \leq \left\lceil \frac{c_2}{c_1} \right\rceil 2^q \sum_{C \in G_{\nu, (2^n c_1)^{-1}}} \nu(C)^q.$$

By Fact 4.1, we therefore have

$$\beta_{\nu \circ g^{-1}}(q) \leq \limsup_{n \rightarrow \infty} \frac{\log \left( \sum_{C \in G_{\nu, (2^n c_1)^{-1}}} \nu(C)^q \right)}{\log(c_1 2^n)} = \beta_\nu(q).$$

The same argument applied to  $\nu \circ g^{-1}$  instead of  $\nu$  and  $g^{-1}$  instead of  $g$  gives the reverse inequality.  $\square$

Next, we have to ensure that with out loss of generality we can assume that  $\nu$  has no atoms in the set of the set  $\mathcal{D}^* := \{2^{-n}k \mid k \in \mathbb{Z}, n \in \mathbb{N}\}$  of dyadic numbers.

**Lemma 4.10.** *Let  $\nu$  be a Borel measure on  $(0, 1)$ . Then there exists an arbitrary small constant  $\alpha > 0$  such that for  $g_\alpha : x \mapsto x + \alpha$ , we have  $\nu \circ g_\alpha^{-1}$  has no atoms in  $\mathcal{D}^*$ .*

*Proof.* Fix  $\epsilon > 0$ . By Fubini's Theorem and the fact that  $\mathcal{D}^*$  is countable, we have

$$\begin{aligned} \int_0^\epsilon \int_{(0,1)} \mathbb{1}_{\mathcal{D}^*}(y + \alpha) d\nu(y) d\Lambda(\alpha) &= \int_{(0,1)} \int_0^\epsilon \mathbb{1}_{\mathcal{D}^*}(y + \alpha) d\Lambda(\alpha) d\nu(y) \\ &= \int_{(0,1)} \Lambda|_{[0,\epsilon]}(\mathcal{D}^* - y) d\nu(y) = 0. \end{aligned}$$

Hence,  $\int_{(0,1)} \mathbb{1}_{\mathcal{D}^*}(y + \alpha) d\nu(y) = 0$ , for  $\Lambda$ -a. e.  $\alpha \in [0, \epsilon]$ . In particular, there exists  $\alpha \in [0, \epsilon]$  such that  $\int_{(0,1)} \mathbb{1}_{\mathcal{D}^*}(y + \alpha) d\nu(y) = \nu \circ g_\alpha^{-1}(\mathcal{D}^*) = \sum_{y \in \mathcal{D}^*} \nu \circ g_\alpha^{-1}(\{y\}) = 0$ .  $\square$

Let  $\nu$  be a bounded Borel probability measure with bounded support contained in  $(0, 1)$ . Then we find a bi-Lipschitz map  $f$  mapping the support of  $\nu$  into  $(\epsilon, 1 - \epsilon)$  for some  $0 < \epsilon < 1$  and after a small affine linear transformation via  $x \mapsto x + \alpha$  the support of  $\nu \circ f^{-1} \circ g_\alpha^{-1}$  is still subset of  $(0, 1)$  and  $\nu \circ f^{-1} \circ g_\alpha^{-1}$  has no atoms in the dyadic numbers  $\mathcal{D}^*$ . In virtue of Proposition 1.13, Lemma 4.10 and Corollary 2.17 these transformations and enlarging the domain also does not change the spectral properties and the  $L^q$ -spectrum Hence, we will from now on assume without loss of generality that  $\nu$  is a probability measure on  $(0, 1)$  such that  $\nu(\mathcal{D}^*) = 0$ .

#### 4.2. Upper bounds in terms of the $L^q$ -spectrum

In this section we let  $\nu$  be an arbitrary Borel probability measure on  $(0, 1)$  with  $\text{card}(\text{supp } \nu) = \infty$  and  $\nu(\mathcal{D}^*) = 0$ . The idea is now to approximate the quantity  $\mathcal{N}^R$  by constructing  $\nu$ -partitions of dyadic intervals. Let us define  $\mathcal{D} := \bigcup_n \mathcal{D}_n$  with  $\mathcal{D}_n := \{A_k^n : k = 0, \dots, 2^n - 1\}$ . The following lemma is obvious.

**Lemma 4.11.** *For every  $1 > t > 0$*

$$P_t := \left\{ C \in \mathcal{D} : \nu(C) \wedge (C) < t \ \& \ \exists C' \in \mathcal{D}_{\lfloor \log_2(\wedge(C)) \rfloor - 1} : C' \supset C \ \& \ \nu(C') \wedge (C') \geq t \right\}$$

*is a  $\nu$ -partition.*

**Proposition 4.12.** *We always have*

$$\bar{s}_\nu \leq \bar{h}_\nu \leq \bar{q}_\nu.$$

*Proof.* We will show  $\bar{q}_v \geq \bar{h}_v$  which together with Proposition 3.7 proves the claim. Let us choose  $q_1 > \bar{q}_v$ . By Lemma 4.11 for  $t \in (0, 1)$ , we have

$$P_t = \left\{ C \in \mathcal{D} : \nu(C) \Lambda(C) < t \ \& \ \exists C' \in \mathcal{D}_{|\log_2(\Lambda(C))|-1} : C' \supset C \ \& \ \nu(C') \Lambda(C') \geq t \right\}$$

is a  $\nu$ -partition of  $[0, 1]$  by dyadic intervals. We also need the set

$$Q_t := \{C \in \mathcal{D} : \nu(C) \Lambda(C) \geq t\}.$$

Note that for  $C \in P_t$  there is exactly one  $C' \in Q_t \cap \mathcal{D}_{|\log_2(\Lambda(C))|-1}$  with  $C \subset C'$  and for each  $C' \in Q_t \cap \mathcal{D}_{|\log_2(\Lambda(C))|-1}$  there are at most two elements of  $P_t \cap \mathcal{D}_{|\log_2(\Lambda(C))|}$  such that they are subsets of  $C'$ . This shows the following crucial inequality

$$\text{card } P_t \leq 2 \text{ card } Q_t.$$

This allows us to estimate

$$t^{q_1} \text{ card } P_t = \sum_{C \in P_t} t^{q_1} \leq 2 \sum_{C \in Q_t} t^{q_1} \leq 2 \sum_{C \in Q_t} (\nu(C) \Lambda(C))^{q_1} \leq 2 \sum_{C \in \mathcal{D}} (\nu(C) \Lambda(C))^{q_1}.$$

With this at hand and with the help of Fact 4.8 we find that the right hand side is finite and therefore we conclude  $\limsup_{t \searrow 0} -\log(\text{card } P_t) / \log(t) \leq q_1$ . By Proposition 3.7 we finally conclude

$$\bar{s}_v \leq \bar{h}_v \leq \limsup_{t \searrow 0} \frac{\log \text{card } P_t}{-\log t} \leq \bar{q}_v.$$

□

*Proof of Proposition 1.10.* Since

$$\frac{\log(\mathcal{N}^R(2^{n-1}))}{\log(2^{n-1})} \leq \frac{\log(2^{n\beta_n(0)+1} + 1)}{\log(2^{n-1})}$$

it follows that  $\underline{s} \leq \underline{h}_v \leq \underline{\dim}_M(\nu)$ . □

**Proposition 4.13.** *Under the assumption that there exists a subsequence  $(n_k)_{k \in \mathbb{N}} \in \mathbb{N}^{\mathbb{N}}$  and a constant  $K > 0$  such that for all  $k \in \mathbb{N}$*

$$\max_{C \in \mathcal{D}_{\nu, n_k}} \nu(C)^{q_{n_k}} \leq \frac{K}{2^{\beta_{n_k}(0)n_k}} \sum_{C \in \mathcal{D}_{\nu, n_k}} \nu(C)^{q_{n_k}}$$

and  $\lim_{k \rightarrow \infty} q_{n_k} = \liminf_{n \rightarrow \infty} q_n$ , where  $q_n \geq 0$  is the unique solution to  $\beta_n(q_n) = q_n$ , we have

$$\underline{s}_v \leq \underline{h}_v \leq \liminf_{n \rightarrow \infty} q_n \leq \liminf_{n \rightarrow \infty} \frac{\beta_n(0)}{1 + \beta_n(0)} = \frac{\underline{\dim}_M(\nu)}{1 + \underline{\dim}_M(\nu)}.$$

*Proof.* We only consider the case  $\underline{\dim}_M(\nu) > 0$ . Since for all  $k \in \mathbb{N}$ ,

$$2^{q_{n_k} n_k} = \sum_{C \in \mathcal{D}_{\nu, n_k}} \nu(C)^{q_{n_k}},$$

we have

$$\max_{C \in \mathcal{D}_{v, n_k}} \nu(C)^{q_{n_k}} \leq \frac{K}{2^{\beta_{n_k}(0)n_k}} \sum_{C \in \mathcal{D}_{v, n_k}} \nu(C)^{q_{n_k}} = K \frac{2^{n_k q_{n_k}}}{2^{n_k \beta_{n_k}(0)}}.$$

Hence,

$$\max_{C \in \mathcal{D}_{v, n_k}} \nu(C) \Lambda(C) \leq \frac{K^{1/q_{n_k}}}{2^{n_k \beta_{n_k}(0)/q_{n_k}}}.$$

Using  $\underline{\dim}_M(\nu) > 0$  we find  $n_k \beta_{n_k}(0) \rightarrow \infty, k \rightarrow \infty$ . Thus we obtain

$$\begin{aligned} \frac{\log \left( \mathcal{N}^R \left( \frac{2^{\beta_{n_k}(0)n_k/q_{n_k}}}{2K^{1/q_{n_k}}} \right) \right)}{\log \left( \frac{2^{\beta_{n_k}(0)n_k/q_{n_k}}}{2K^{1/q_{n_k}}} \right)} &\leq \frac{\log \left( 22^{n_k \beta_{n_k}(0)} + 1 \right)}{\log \left( 2^{n_k \beta_{n_k}(0)} / q_{n_k} - \left( \log(K^{1/q_{n_k}}) - \log(2) \right) \right)} \\ &\leq \frac{\log(4)}{\log \left( 2^{n_k \beta_{n_k}(0)} / q_{n_k} - \left( \log(K^{1/q_{n_k}}) - \log(2) \right) \right)} \\ &\quad + \frac{1}{1/q_{n_k} - \left( \log(K^{1/q_{n_k}}) - \log(2) \right) / \log \left( 2^{\beta_{n_k}(0)n_k/q_{n_k}} \right)}. \end{aligned}$$

Therefore,  $\underline{s}_\nu \leq \underline{h}_\nu \leq \liminf_{n \rightarrow \infty} q_n$ .  $\square$

#### 4.3. Lower bounds in terms of the optimize coarse multifractal dimension

Here we again assume that  $\nu$  is a finite Borel measure on  $(0, 1)$  with  $\text{card}(\text{supp } \nu) = \infty$  and  $\nu(\mathcal{D}^*) = 0$ . To find a good lower bound, we use the concept of the *upper* and *lower optimize coarse multifractal dimension*  $\underline{F}_\nu, \overline{F}_\nu$  as defined in the introduction.

**Lemma 4.14.** *For fixed  $\alpha > 0$  and with  $n_x^\alpha := \left\lceil \log \left( \frac{x}{2} \right) / (\log(2)(\alpha + 1)) \right\rceil$  we have for every  $x > 2^{3+\alpha}$*

$$N_\nu(x) \geq \mathcal{N}_3^L(x) \geq \frac{N_\alpha(n_x^\alpha)}{3} - 3,$$

or with  $x_n^\alpha := 2^{n(\alpha+1)+1}$  we have for every  $n \in \mathbb{N}$

$$N_\nu(x_n^\alpha) \geq \frac{N_\alpha(n)}{3} - 3.$$

*Proof.* For fixed  $n \in \mathbb{N}, \alpha > 0$  let  $(c_1, d_1], \dots, (c_{N_\alpha(n)}, d_{N_\alpha(n)})$  denote the intervals of  $M_n(\alpha)$  ordered in the natural way, i. e.  $d_i < c_{i+1}$ . For  $i = 0, \dots, \lfloor N_\alpha(n)/3 - 2 \rfloor - 1$  define

$$D_{n,i} := \left( c_{2+3i} - \frac{1}{2^n}, c_{2+3i} \right] \cup (c_{2+3i}, d_{2+3i}] \cup \left( d_{2+3i}, d_{2+3i} + \frac{1}{2^n} \right).$$

Note that we have

$$\nu \left( \langle D_{n,i} \rangle_3 \right) \Lambda \left( \langle D_{n,i} \rangle_3 \right) = \nu \left( (c_{2+3i}, d_{2+3i}] \right) \Lambda \left( (c_{2+3i}, d_{2+3i}] \right) \geq 2^{-n(\alpha+1)},$$

hence for  $n_x^\alpha := \left\lceil \log \left( \frac{x}{2} \right) / (\log(2)(\alpha + 1)) \right\rceil$  and  $x > 2^{3+\alpha}$ , we have

$$\nu \left( \langle D_{n_x^\alpha, i} \rangle_3 \right) \Lambda \left( \langle D_{n_x^\alpha, i} \rangle_3 \right) \geq \frac{2}{x}$$

which shows

$$\begin{aligned} \mathcal{N}_{3,v}^L(x) &= \sup \left\{ \text{card}(P) : P \in \Pi_0 : \min_{C \in P} v(\langle I \rangle_3) \wedge \langle I \rangle_3 \geq \frac{2}{x} \right\} \\ &\geq \left\lfloor \frac{\mathcal{N}_\alpha(n_x^\alpha)}{3} - 2 \right\rfloor \geq \frac{\mathcal{N}_\alpha(n_x^\alpha)}{3} - 3. \end{aligned}$$

□

Now we can give a lower bound on  $\underline{s}_v$  and  $\bar{s}_v$  in terms of the lower and upper optimize coarse multifractal dimension.

**Proposition 4.15.** *As a general lower (upper) bound for the lower (upper) spectral dimension for all  $m \in (1, 3]$ , we have*

$$\underline{F}_v \leq \underline{h}_v^m \leq \underline{s}_v \text{ and } \bar{F}_v \leq \bar{h}_v^m \leq \bar{s}_v.$$

*Proof.* First note that Proposition 3.7 gives  $\underline{h}_v^m \leq \underline{s}_v$  and  $\bar{h}_v^m \leq \bar{s}_v$  for all  $m > 1$ . With  $n_x^\alpha := \left\lfloor \frac{\log(\frac{x}{3})}{\log(2^{(2\alpha+1)})} \right\rfloor$  and  $x > 2^{3+\alpha}$  with Lemma 4.14 for every  $\alpha \geq 0$  and  $x > 2^{3+\alpha}$ , we have

$$\frac{\log^+(3\mathcal{N}_3^L(x) + 3)}{\log(x)} \geq \frac{\log^+(\mathcal{N}_\alpha(n_x^\alpha))}{\log x}.$$

Hence, for all  $1 < m \leq 3$ ,

$$\begin{aligned} \underline{h}_v^m &\geq \underline{h}_v^3 = \liminf_x \frac{\log^+(3\mathcal{N}_3^L(x) + 3)}{\log x} \geq \liminf_x \frac{\log^+(\mathcal{N}_\alpha(n_x^\alpha))}{\log x} \\ &= \liminf_x \frac{\log^+(\mathcal{N}_\alpha(n_x^\alpha))}{(1+\alpha)\log(2^{\log_2(\frac{x}{3})/(1+\alpha)}) + \log(2)} \\ &\geq \liminf_x \frac{\log^+(\mathcal{N}_\alpha(n_x^\alpha))}{(1+\alpha)\log 2^{n_x^\alpha} + \log(2)(2+\alpha)} \\ &= \liminf_x \frac{\log^+(\mathcal{N}_\alpha(n_x^\alpha))}{(1+\alpha)\log 2^{n_x^\alpha}} \geq \liminf_{n \rightarrow \infty} \frac{\log^+(\mathcal{N}_\alpha(n))}{(1+\alpha)\log 2^n}, \end{aligned}$$

which implies  $\underline{h}_v^m \geq \underline{F}_v$ . We also have for  $x_m^\eta := 2^{m(\eta+1)+1}$  with  $\eta > 0$ ,

$$\begin{aligned} \bar{h}_v^m &\geq \bar{h}_v^3 \geq \limsup_{m \rightarrow \infty} \frac{\log^+(3\mathcal{N}_3^L(x_m^\eta) + 3)}{\log(x_m^\eta)} \geq \limsup_{m \rightarrow \infty} \frac{\log^+(\mathcal{N}_\eta(n_{x_m^\eta}^\eta))}{\log x_m^\eta} \\ &\geq \limsup_{m \rightarrow \infty} \frac{\log^+(\mathcal{N}_\eta(n_{x_m^\eta}^\eta))}{(1+\eta)\log 2^{n_{x_m^\eta}^\eta}} = \limsup_{m \rightarrow \infty} \frac{\log^+(\mathcal{N}_\eta(m))}{(1+\eta)\log 2^m}, \end{aligned}$$

where we used  $n_{x_m^\eta}^\eta = m$ . Thus,  $\bar{h}_v^m \geq \bar{F}_v$ . □

We need the following elementary observation from large deviation theory which seems not to be standard in the relevant literature.

**Lemma 4.16.** *Suppose  $(X_n)_{n \in \mathbb{N}}$  are real-valued random variables on some probability spaces  $(\Omega_n, \mathcal{A}_n, \mu_n)$  such that the rate function  $\mathfrak{c}(t) := \limsup_{n \rightarrow \infty} c_n(t)$  is a proper convex function with  $c_n(t) := a_n^{-1} \log \int \exp tX_n d\mu_n$ ,  $t \in \mathbb{R}$ ,  $a_n \rightarrow \infty$  and such that  $0$  belongs to the interior of the domain of finiteness  $\{t \in \mathbb{R} : \mathfrak{c}(t) < \infty\}$ . Let  $I = (a, d)$  be an open interval containing the subdifferential  $\partial\mathfrak{c}(0) = [b, c]$  of  $\mathfrak{c}$  in  $0$ . Then there exists  $r > 0$  such that for all  $n$  sufficiently large,*

$$\mu_n(a_n^{-1}X_n \notin I) \leq 2 \exp(-ra_n).$$

*Proof.* We assume that  $\partial\mathfrak{c}(0) = [b, c]$  and  $I = (a, d)$  with  $a < b \leq c < d$ . First note that the assumptions ensure that  $-\infty < b \leq c < \infty$ . By Chebyshev inequality for all  $q > 0$ ,

$$\mu_n(a_n^{-1}X_n \geq d) = \mu_n(qX_n \geq qa_nd) \leq \exp(-qa_nd) \int \exp(qX_n) d\mu_n$$

implying

$$\limsup a_n^{-1} \log \mu_n(a_n^{-1}X_n \geq d) \leq \inf_{q>0} \mathfrak{c}(q) - qd = \inf_{q \in \mathbb{R}} \mathfrak{c}(q) - qd,$$

where the equality follows from the assumption  $d \notin \partial\mathfrak{c}(0)$  and  $\mathfrak{c}(q) - qd \geq (c - d)q \geq 0$  for all  $q \leq 0$ . Similarly, we find

$$\limsup a_n^{-1} \log \mu_n(a_n^{-1}X_n \leq a) \leq \inf_{q<0} \mathfrak{c}(q) - qa = \inf_{q \in \mathbb{R}} \mathfrak{c}(q) - qa.$$

We are left to show that both upper bounds are negative. We show the first case by contradiction – the other case follows in exactly the same way. Assuming  $\inf_{q \in \mathbb{R}} \mathfrak{c}(q) - qd = 0$  implies for all  $q \in \mathbb{R}$  that  $\mathfrak{c}(q) - qd \geq 0$ , or after rearranging,  $\mathfrak{c}(q) - \mathfrak{c}(0) \geq dq$ . This means, according to the definition of the subdifferential, that  $d \in \partial\mathfrak{c}(0)$ , contradicting our assumptions.  $\square$

**Proposition 4.17.** *Assume for some  $q \in [0, 1]$  that  $\beta_v(q) = \lim \beta_{n_k}^v(q)$  on a subsequence  $(n_k)$ . Then we have  $[a, b] = -\partial\beta_v(q) \supset [a', b'] = -\partial \limsup_k \beta_{n_k}(q)$  and*

$$\frac{aq + \beta_v(q)}{1 + b} \leq \frac{a'q + \beta_v(q)}{1 + b'} \leq \sup_{t>b} \liminf_{k \rightarrow \infty} \frac{\log \mathcal{N}_t(n_k)}{\log 2^{n_k}(1 + t)} \leq \sup_{t>b} \limsup_{k \rightarrow \infty} \frac{\log \mathcal{N}_t(n_k)}{\log 2^{n_k}(1 + t)}.$$

*Proof.* Without loss of generality we can assume  $b < \infty$ . The fact that  $-\partial\beta_v(q) \supset -\partial \limsup_k \beta_{n_k}(q)$  follows immediately from  $\limsup_k \beta_{n_k} \leq \beta_v$ . It suffices to prove the second inequality. Now, for all  $k \in \mathbb{N}$  and  $s < a \leq b < t$ , we have

$$\begin{aligned} \mathcal{N}_t(n_k) &\geq \text{card} \left\{ \underbrace{0 \leq \ell < 2^{n_k} : 2^{-s n_k} > \nu(A_{n_k}^\ell) > 2^{-t n_k}}_{:= L_{n_k}^{s,t}} \right\} \\ &\geq \sum_{\ell \in L_{n_k}^{s,t}} \nu(A_{n_k}^\ell)^q 2^{s n_k q} \geq 2^{s n_k q + n_k \beta_{n_k}(q)} \sum_{1 \leq \ell \leq 2^{n_k}} \mathbb{1}_{L_{n_k}^{s,t}}(\ell) \nu(A_{n_k}^\ell)^q 2^{-n_k \beta_{n_k}(q)} \\ &= 2^{s n_k q + n_k \beta_{n_k}(q)} \left( 1 - \sum_{1 \leq \ell \leq 2^{n_k}} \mathbb{1}_{(L_{n_k}^{s,t})^c}(\ell) \nu(A_{n_k}^\ell)^q 2^{-n_k \beta_{n_k}(q)} \right). \end{aligned}$$

We use the lower large deviation principle for the process  $X_k(\ell) := \log^+ v(A_{n_k}^\ell)$  with probability measure on  $\{1, \dots, 2^{n_k}\}$  given by  $\mu_k(\{\ell\}) := v(A_{n_k}^\ell)^q / 2^{-n_k \beta_{n_k}(q)}$ . We find for the free energy function

$$\begin{aligned} c(x) &:= \limsup_{n_k} \frac{1}{\log 2^{n_k}} \log \left( \mathbb{E}_{\mu_k} (\exp x X_k) \right) = \limsup_{n_k} \frac{1}{\log 2^{n_k}} \log \left( \sum_{C \in \mathcal{D}_{n_k}} v(C)^{x+q} / 2^{n_k \beta_{n_k}(q)} \right) \\ &= \limsup_{n_k} \beta_{n_k}(q+x) - \beta_v(q), \end{aligned}$$

with  $-\partial c(0) = [a', b'] \subset [a, b] \subset (s, t)$  and hence, by Lemma 4.16, there exists a constant  $r > 0$  depending on  $s, t$  and  $q$  such that for  $k$  large,

$$\sum_{1 \leq \ell \leq 2^{n_k}} \mathbb{1}_{(L_{n_k}^{s,t})^c}(\ell) v(A_{n_k}^\ell)^q / 2^{n_k \beta_{n_k}(q)} = \mu_k \left( \frac{X_k}{\log(2^{n_k})} \notin (-t, -s) \right) \leq 2 \exp(-rn_k).$$

Therefore,

$$\liminf_{k \rightarrow \infty} \frac{\log \mathcal{N}_t(n_k)}{\log 2^{n_k}} \geq sq + \beta_v(q)$$

for all  $s < a$  and  $t > b$  and therefore

$$\sup_{t > b} \liminf_{k \rightarrow \infty} \frac{\log \mathcal{N}_t(n_k)}{\log 2^{n_k}(1+t)} \geq \sup_{t > b} \frac{aq + \beta_v(q)}{1+t} = \frac{aq + \beta_v(q)}{1+b}.$$

□

**Proposition 4.18.** *For all  $m \in (1, 3]$ , we have*

$$\bar{q}_v = \bar{F}_v = \bar{h}_v^m = \bar{s}_v.$$

*Proof.* We only have to consider the case  $\bar{q}_v > 0$ . First note that the family of convex function  $(\beta_n)$  restricted to  $[0, 1]$  only takes values in  $[0, 1]$  and on any compact interval  $[c, d] \subset (0, 1)$  and for all  $c \leq x \leq y \leq d$ , we have

$$\frac{\beta_n(x) - \beta_n(0)}{x-0} \leq \frac{\beta_n(y) - \beta_n(x)}{y-x} \leq \frac{\beta_n(1) - \beta_n(y)}{1-y},$$

which implies

$$|\beta_n(y) - \beta_n(x)| \leq \max \left\{ \frac{1}{c}, \frac{1}{1-d} \right\} |x-y|$$

and hence  $(\beta_n)$  is uniformly Lipschitz and by Arzelà–Ascoli relatively compact. Using this fact we find a subsequence  $(n_k)$  such that  $\lim_k \beta_{n_k}(\bar{q}_v) = \limsup \beta_n(\bar{q}_v) = \bar{q}_v$  and  $\beta_{n_k}$  converges uniformly to the convex function  $\beta^\sharp$  on  $[\bar{q}_v - \delta, \bar{q}_v + \delta] \subset [0, 1]$  for  $\delta$  sufficiently small and put  $[a, b] = -\partial \beta^\sharp(\bar{q}_v)$ . Then by the convexity of  $\beta^\sharp$  we find for every  $\delta > \epsilon > 0$  an element  $q \in (\bar{q}_v - \epsilon, \bar{q}_v)$  such that  $\beta^\sharp$  is differentiable in  $q$  with  $-(\beta^\sharp)'(q) \in [b, b + \epsilon]$  since the points where  $\beta^\sharp$  is differentiable on  $(0, 1)$  lie dense in  $(0, 1)$  which follows from the fact that  $\beta^\sharp$  is a decreasing function and left-hand derivatives of convex functions are

left-continuous and non-decreasing. In the same way as in the proof Proposition 4.17, we obtain,

$$\begin{aligned} \sup_{t>b} \limsup_{n \rightarrow \infty} \frac{\log \mathcal{N}_t(n)}{\log 2^n(1+t)} &\geq \sup_{t>b} \limsup_{k \rightarrow \infty} \frac{\log \mathcal{N}_t(n_k)}{\log 2^{n_k}(1+t)} \\ &\geq \frac{-(\beta^\#)'(q)q + \beta^\#(q)}{1 - (\beta^\#)'(q)} \\ &\geq \frac{b(\bar{q}_v - \varepsilon) + \beta^\#(\bar{q}_v)}{1 + b + \varepsilon} \\ &= \frac{b(\bar{q}_v - \varepsilon) + \bar{q}_v}{1 + b + \varepsilon}. \end{aligned}$$

Taking  $\varepsilon \rightarrow 0$

$$\bar{F}_v = \sup_{\alpha \geq 0} \frac{\bar{F}_v(\alpha)}{1 + \alpha} \geq \bar{q}_v.$$

The second part of Proposition 4.15 and Proposition 4.12 give the claimed equalities.  $\square$

#### 4.4. Proof of main results

Now, we are able to prove the main results by combining the results of Section 4.3 and Section 4.2. Recall from the introduction that we call  $v$  *regular* if  $\underline{F}_v = \bar{F}_v$  and  $L^q$ -*regular*, if  $\beta_v(q) = \liminf_n \beta_n^v(q)$  for  $q \in (\bar{q}_v - \varepsilon, \bar{q}_v)$ , for some  $\varepsilon > 0$ , or  $\beta_v(\bar{q}_v) = \liminf_n \beta_n^v(\bar{q}_v)$  and  $\beta_v$  is differentiable in  $\bar{q}_v$ .

*Proof of Theorem 1.2.* Since by Proposition 4.18 we have  $\bar{F}_v = \bar{q}_v$ , it suffices to show that  $L^q$ -regularity implies

$$\underline{F}_v \geq \bar{q}_v.$$

Indeed, let us assume  $[a, b] = -\partial\beta_v(\bar{q}_v)$ . Then by the convexity of  $\beta_v$  we find for every  $\varepsilon \in (0, \bar{q}_v)$  an element  $q \in (\bar{q}_v - \varepsilon, \bar{q}_v)$  such that  $\beta_v$  is differentiable in  $q$  with  $-\beta'_v(q) \in [b, b + \varepsilon]$  since the points where  $\beta_v$  is differentiable on  $(0, 1)$  lie dense in  $(0, 1)$  which follows from the fact that  $\beta_v$  is a decreasing function and the left-hand continuity of left-hand derivative of the convex function  $\beta_v$ . Then by Proposition 4.17, we have

$$\begin{aligned} \sup_{t \geq 0} \liminf_{n \rightarrow \infty} \frac{\log^+(\mathcal{N}_t(n))}{\log 2^n(1+t)} &\geq \sup_{t > -\beta'_v(q)} \liminf_{n \rightarrow \infty} \frac{\log^+(\mathcal{N}_t(n))}{\log 2^n(1+t)} \\ &\geq \frac{-\beta'_v(q)q + \beta_v(q)}{1 - \beta'_v(q)} \geq \frac{b(\bar{q}_v - \varepsilon) + \bar{q}_v}{1 + b + \varepsilon}. \end{aligned}$$

Taking the limit  $\varepsilon \rightarrow 0$

$$\sup_{\alpha \geq 0} \liminf_{n \rightarrow \infty} \frac{\log^+(\mathcal{N}_\alpha(n))}{\log 2^n(1+\alpha)} \geq \frac{b\bar{q}_v + \bar{q}_v}{1 + b} = \bar{q}_v.$$

The case  $\beta_v$  exists as limit in  $\bar{q}_v$  and is differentiable in  $\bar{q}_v$  is covered by Proposition 4.17.  $\square$

*Proof of Theorem 1.1.* The first claim follows from Propositions 4.12, 4.15, 3.7 and 4.18.  $\square$

*Proof of Corollary 1.12.* The first part follows immediately from Theorem 1.1 in tandem with Fact 4.5. For the second part note that by Lemma 2.14

$$s_{\nu_1} \leq \max\{\underline{s}_{\nu_1}, \underline{s}_{\nu_1}\} \leq \underline{s}_{\nu_1+\nu_2} \leq \bar{s}_{\nu_1+\nu_2} = \bar{F}_{\nu_1+\nu_2} = \max\{\bar{s}_{\nu_1}, \bar{s}_{\nu_2}\} = s_{\nu_1}$$

and using monotonicity properties of the coarse optimized multifractal dimension

$$\bar{F}_{\nu_1} = \max\{\underline{F}_{\nu_1}, \underline{F}_{\nu_2}\} \leq \underline{F}_{\nu_1+\nu_2} \leq \bar{F}_{\nu_1+\nu_2} = s_{\nu_1} = \bar{F}_{\nu_1}.$$

$\square$

*Proof of Corollary 1.3.* First, we show that for an absolutely continuous measure  $\nu$  the  $L^q$ -spectrum always exists as limit on  $[0, 1]$ . By Hölder's inequality for all  $C \in \mathcal{D}_{\nu, n}$  and  $q \in (0, 1)$ , we have

$$2^n \int_C \varrho^q d\Lambda \leq \left( \int_C 2^{n/(1-q)} d\Lambda \right)^{1-q} \left( \int_C \varrho d\Lambda \right)^q = \left( 2^{n \frac{q}{1-q}} \right)^{1-q} \left( \int_C \varrho d\Lambda \right)^q = 2^{nq} \left( \int_C \varrho d\Lambda \right)^q.$$

This leads to

$$\beta_n^\nu(q) \geq \frac{\log \left( \int_{[0,1]} \varrho^q d\Lambda \right)}{\log(2^n)} + 1 - q,$$

implying  $1 - q \leq \liminf_n \beta_n^\nu(q) \leq \limsup_n \beta_n^\nu(q) \leq 1 - q$  for all  $q \in [0, 1]$ . Now, the remaining claims follow from Fact 4.5 and Corollary 1.4.  $\square$

*Proof of Proposition 1.5.* By Proposition 4.17,

$$\frac{aq + \beta_\nu(q)}{1 + b} \leq \sup_{t > b} \liminf_{n \rightarrow \infty} \frac{\log \mathcal{N}_t(n)}{\log 2^n(1 + t)} \leq \underline{F}_\nu \leq \underline{s}_\nu.$$

$\square$

*Proof of Corollary 1.7.* For the case  $\delta_\nu^* = 0$  we combine Fact 4.7 and the upper estimate from Proposition 4.12. If  $\delta_\nu^* > 0$ , this is a direct consequence of Proposition 4.12 and the general properties of  $\beta_\nu$  on  $[0, 1]$  as a convex non-increasing function not exceeding the line  $t \mapsto \delta_\nu^*(1 - t)$  which intersects the bisector in  $\delta_\nu^*/(1 + \delta_\nu^*)$ . The last claim follows from the fact that  $\beta_\nu$  is linear on  $[0, 1]$  if and only if  $\delta_\nu = \delta_\nu^*$  and in this case  $\bar{q}_\nu = \delta_\nu^*/(1 + \delta_\nu^*)$ . The lower bound follows from Proposition 1.5 using the fact that  $\beta_\nu$  always exists as a limit in 1.  $\square$

*Remark 4.19.* The case  $\delta_\nu^* = 0$  immediately gives  $\bar{s}_\nu = 0$ . If we use more information on  $\beta_\nu$  for the case  $\delta_\nu^* > 0$  we find a better upper bound; namely, with  $q_1 := \inf\{s : \beta_\nu(s) \leq 0\}$  we have

$$\bar{s}_\nu \leq \bar{q}_\nu \leq \frac{q_1 \delta_\nu^*}{q_1 + \delta_\nu^*}.$$

*Proof of Corollary 1.9.* By Theorem 1.1 we have  $\bar{q}_v = \bar{s}_v$  and if  $\nu$  is  $L^q$ -regular, the claim then follows from Corollary 1.7. In [Heu07, Theorem 3.1] it has been shown that  $\bar{\delta}_v \geq \dim_P(E) \geq \dim_H(\nu) \geq \underline{\dim}_H(\nu) \geq \underline{\delta}_v$  and this gives the second claim as a consequence of the first equivalence.

The last claim follows from the convexity of  $\beta_\nu$ ,  $\beta_\nu(0) \leq 1$  and  $\beta_\nu(1) = 0$  in tandem with  $\beta_\nu(\delta_v^*/(1 + \delta_v^*)) = \delta_v^*/(1 + \delta_v^*)$  which implies  $\beta_\nu(q) = \delta_v^*(1 - q)$  for  $q \in [0, 1]$ . Now, [Heu07, Theorem 3.1] implies  $\delta_v^* = \dim_H(\nu) = \dim_P(\nu)$ .  $\square$

*Proof of Corollary 1.4.* If  $\bar{s}_v = 1/2$  then it follows from Proposition 4.12 that  $1/2 = \bar{s}_v \leq \bar{q}_v \leq 1/2$ . The convexity of  $\beta_\nu$  and the fact that  $\beta_\nu(1) = 0$  and  $\beta_\nu(0) \leq 1$  forces  $\beta_\nu(q) = 1 - q$  for all  $q \in [0, 1]$ . The second statement is an immediate consequence of Proposition 4.17 by observing that, as in case (1), by convexity we have  $\beta_\nu(q) = 1 - q$  for all  $q \in [0, 1]$ . This implies the differentiability of  $\beta_\nu$  in the particular point  $q \in (0, 1)$ , where by assumption  $\beta_\nu(q) = \lim \beta_n^r(q)$ . Now, applying Proposition 4.17 gives

$$\frac{1}{2} = \frac{q + 1 - q}{1 + 1} \leq s_\nu.$$

Since we always have  $\bar{s}_v \leq 1/2$ , our claim follows.  $\square$

## 5. Examples

### 5.1. $C^1$ -self-conformal measures

For simplicity, let us start with the self-similar case.

**Example 5.1.** Here, we consider for fixed  $m \in \mathbb{N}$  self-similar probability measure  $\nu$  with probability vector  $(p_1, \dots, p_m) \in (0, 1)^m$  and contractions  $T_i(x) = \sigma_i x + b_i$  for  $i = 1, \dots, m$  with  $\sigma_i \in (0, 1)$  which maps from  $[0, 1]$  to  $[0, 1]$  and fulfills the open set condition (i. e.  $T_i((0, 1)) \cap T_j((0, 1)) = \emptyset$  for all  $j \neq i$ ). That is the unique Borel measure which satisfy

$$\nu(A) = \sum_{i=1}^m p_i \nu(T_i^{-1}(A)), A \in \mathfrak{B}([0, 1]).$$

Using [Rie95, Theorem 15] we obtain for  $q \geq 0$  that  $\beta_\nu(q)$  is given by

$$\sum_{i=1}^m p_i^q \sigma_i^{\beta_\nu(q)} = 1.$$

Thus, we can conclude that  $\bar{q}_v$  is determined by the equality  $\sum_{i=1}^m (p_i \sigma_i)^{\bar{q}_v} = 1$ . Moreover, let us consider the natural choice of weights given by  $p_i := \sigma_i^u$ , where  $u$  is the zero of the associated pressure function, i. e.  $\sum_{i=1}^m \sigma_i^u = 1$ . Then for  $q \geq 0$ , we have  $\beta_\nu(q) = u - uq$  and consequently  $u = \delta_\nu = \delta_\nu^*$ . Hence, Corollary 1.9 gives

$$s_\nu = \bar{q}_v = \underline{q}_v = \frac{\delta_\nu^*}{1 + \delta_\nu^*}.$$

Now, we consider  $C^1$ -self-conformal measures as in [Fen07] which do not have to satisfy any separation conditions. Let  $U \subset \mathbb{R}$  be an open set and  $S : U \rightarrow \mathbb{R}$  a  $C^1$ -map with  $|S'(x)| > 0$  for all  $x \in U$ . Moreover, assume that  $S$  is a contracting, i. e. there exists  $0 < c < 1$  such for all  $x, y \in U$  we have  $|S(x) - S(y)| \leq c|x - y|$ . We call  $\{S_i : [0, 1] \rightarrow [0, 1]\}_{i=1, \dots, \ell}$  a  $C^1$ -conformal iterated function system on  $[0, 1]$ , if each  $S_i$  extends to an injective contracting  $C^1$ -map  $S_i : U \rightarrow U$  on an open set  $U \supset [0, 1]$ .

Given a  $C^1$ -conformal iterated function system  $\{S_i : [0, 1] \rightarrow [0, 1]\}_{i=1, \dots, \ell}$  then there exists a non-empty compact set  $K \subset [0, 1]$  such that

$$K = \bigcup_{i=1}^{\ell} S_i(K).$$

For a probability vector  $(p_1, \dots, p_\ell)$  there is a unique Borel probability measure  $\nu$  satisfying

$$\nu = \sum_{i=1}^{\ell} p_i \nu \circ S_i^{-1}.$$

The measure  $\nu$  with support  $\text{supp } \nu = K$  is called *self-conformal measure*. Furthermore, we assume that  $S_i$  do not have all the same fixed point, then  $\nu$  is atomless ([PS00]). This can be seen as follows. Fix  $x \in [0, 1]$  such that  $\nu(\{x\}) = \max(\nu(\{y\}) : y \in [0, 1]) =: m$ . Since with  $I := \{1, \dots, \ell\}$

$$\nu(\{x\}) = \sum_{\omega \in I^n} p_\omega \nu(T_\omega^{-1}\{x\})$$

we find  $\nu(T_\omega^{-1}\{x\}) = m$  for every  $\omega \in I^n$  and  $n \in \mathbb{N}$ . Hence, if  $m > 0$ , we have  $x \in T_\omega([0, 1])$  for all  $\omega \in I^n$  and  $n \in \mathbb{N}$  and  $x$  is the common fixed point for all contractions. In [Fen07, Corollary 4.5.], building on the earlier work [PS00], it has been proved that the  $L^q$ -spectrum  $\beta_\nu$  of the self-conformal measure  $\nu$  exists as a limit.

**Proposition 5.2** ([Fen07, Corollary 4.5.], [PS00]). *For every  $C^1$ -self-conformal measure  $\nu$  we have that the  $L^q$ -spectrum  $\beta_\nu$  exists as a limit on  $\mathbb{R}_{>0}$ .*

*Proof of Theorem 1.14.* Combining Theorems 1.2 and 1.1 readily proves Theorem 1.14.  $\square$

Note, that every measure on  $[0, 1]$  which is absolutely continuous with respect to the Lebesgue measure has spectral dimension  $1/2$ . Hence, we obtain the following corollary.

**Corollary 5.3.** *If the fixed point  $\bar{q}_\nu$  of  $\beta_\nu$  is smaller than  $1/2$ , then  $\nu$  is singular with respect to the Lebesgue measure.*

*Remark 5.4.* Note that Example 5.11 shows that there exist singular measures with spectral dimension  $1/2$ . Hence, in general we can not expect the equivalence of  $\bar{q}_\nu < 1/2$  and  $\nu \perp \Lambda$ .

## 5.2. Homogeneous Cantor measures

Let us recall the construction of general homogeneous Cantor measures as in [Arz14, Min20, BH97] allowing us to construct examples for which the spectral dimension does not exist. Let  $J$  be finite or countably infinite subset of  $\mathbb{N}$ . For every  $j \in J$  we define an iterated function system  $\mathcal{S}^{(j)}$ . For  $i = 1, 2$  let  $S_i^{(j)} : [a, b] \rightarrow [a, b]$  be defined by

$$S_i^{(j)}(x) = r_i^{(j)}x + c_i^{(j)}$$

with  $r_i^{(j)} \in (0, 1)$  and  $c_i^{(j)} \in \mathbb{R}$  are chosen such that

$$a = S_1^{(j)}(a) < S_1^{(j)}(b) \leq S_2^{(j)}(a) < S_2^{(j)}(b) = b.$$

This ensures the open set condition. We define  $\mathcal{S}^{(j)} = (S_1^{(j)}, S_2^{(j)})$ . Moreover let us define an *environment sequence*  $\xi := (\xi_i)_{i \in \mathbb{N}} \in J^{\mathbb{N}}$ . Each  $\xi_i$  represent an iterated function system  $\mathcal{S}^{(\xi_i)}$ . To give a suitable coding we define a word space  $W_n := \{1, 2\}^n$  of words with length  $n$ . For  $n \in \mathbb{N}$  and  $\omega := (\omega_1, \dots, \omega_n) \in W_n$  we set

$$S_\omega^{(\xi)} := S_{\omega_1}^{(\xi_1)} \circ S_{\omega_2}^{(\xi_2)} \circ \dots \circ S_{\omega_n}^{(\xi_n)}.$$

Now, for any environment sequence  $\xi$ , we construct a probability measure  $\nu^{(\xi)}$  on  $[a, b]$  with support  $K^{(\xi)}$  defined by

$$K^{(\xi)} := \bigcap_{n=1}^{\infty} \bigcup_{\omega \in W_n} (S_\omega^{(\xi_1)} \circ S_\omega^{(\xi_2)} \circ \dots \circ S_\omega^{(\xi_n)})([a, b]).$$

For any  $j \in J$ , let  $(p_i^j)_{i=1,2} \in (0, 1)^2$  with  $\sum_{i=1}^2 p_i^j = 1$ . Moreover, let  $p_\omega^{(\xi)} := \prod_{i=1}^n p_{\omega_i}^{(\xi_i)}$  be for  $n \in \mathbb{N}$  and  $\omega := (\omega_1, \dots, \omega_n) \in W_n$ . Then define the following sequence of probability measures:  $\nu_0 := \frac{1}{b-a} \Lambda|_{[a,b]}$  and for  $n \in \mathbb{N}$  and  $A \in \mathfrak{B}([a, b])$

$$\nu_n(A) := \sum_{\omega \in W_n} p_\omega^{(\xi)} \nu_0 \left( (S_\omega^{(\xi)})^{-1}(A) \right)$$

then  $\nu^{(\xi)}$  is defined by

$$\nu^{(\xi)}(A) = \lim_{n \rightarrow \infty} \nu_n(A).$$

The theorem of Vitali-Hahn-Saks ensures that  $\nu^{(\xi)}$  is a probability measure on  $\mathfrak{B}([a, b])$  (see [Arz14, Lemma 3.1.2.]) and for every  $n \in \mathbb{N}$  and  $\omega \in W_n$ , we have  $\nu^{(\xi)}(S_\omega^{(\xi)}([a, b])) = p_\omega^{(\xi)}$ .

**Example 5.5** (Homogeneous Cantor measure with non-converging  $L^q$ -spectrum). Now, let us consider the following environment

$$\xi_i := \begin{cases} 1 & , \exists \ell \in \mathbb{N}_0 : 2^{2\ell} < i \leq 2^{2\ell+1} \\ 2 & , \exists \ell \in \mathbb{N}_0 : 2^{2\ell+1} < i \leq 2^{2\ell+2} \end{cases}$$

and  $S_1^{(1)}(x) := \frac{x}{4}, S_2^{(1)}(x) := \frac{x}{4} + \frac{3}{4}, S_1^{(2)}(x) := \frac{x}{16}, S_2^{(2)}(x) := \frac{x}{16} + \frac{15}{16}, x \in [0, 1]$ . Furthermore, let  $p_1, p_2 \in (0, 1)$  be with  $p_1 + p_2 = 1$  then define for every  $j \in \{1, 2\}$   $p_1^j = p_1$  and  $p_2^j = p_2$ . First, observe that for  $n \in \mathbb{N}$  and  $\omega \in W_n$ ,

$$\Lambda \left( S_\omega^{(\xi)}([0, 1]) \right) = \begin{cases} 2^{-2^{2l+1} \cdot 8/3 + 10/3 - 4(n-2^{2l+1})} & , 2^{2l+1} < n \leq 2^{2l+2} \\ 2^{-10/3 \cdot 2^{2l+2} + 10/3 - 2(n-2^{2l+2})} & , 2^{2l+2} < n \leq 2^{2l+3} \end{cases}.$$

Hence, we obtain for all  $n \in \mathbb{N}$  with  $2^{2l+1} < n \leq 2^{2l+2}, q \geq 0$  and  $\omega \in W_n$

$$\begin{aligned} & \frac{1}{-\log \left( \Lambda \left( S_\omega^{(\xi)}([0, 1]) \right) \right)} \log \left( \sum_{l=0, v(A_l^n) > 0}^{2^{-\log_2 \left( \Lambda \left( S_\omega^{(\xi)}([a, b]) \right) \right) - 1}} v^{(\xi)} \left( A_l^n \right)^q \right) \\ &= \frac{1}{-\log \left( \Lambda \left( S_\omega^{(\xi)}([0, 1]) \right) \right)} \log \left( \sum_{l \in W_n} v^{(\xi)} \left( S_l^{(\xi)}([a, b]) \right)^q \right) \\ &= \frac{\log_2 \left( p_1^q + p_2^q \right)}{-4/3 \cdot 2^{2l+1}/n + 4 - 10/(3n)} \end{aligned}$$

and, for  $2^{2l+2} < n \leq 2^{2l+3}$ ,

$$\begin{aligned} \frac{1}{-\log \left( \Lambda \left( S_\omega^{(\xi)}([0, 1]) \right) \right)} \log \left( \sum_{l \in W_n} v^{(\xi)} \left( S_l^{(\xi)}([a, b]) \right)^q \right) &= \frac{\log \left( p_1^q + p_2^q \right)}{-\log \left( \Lambda \left( S_\omega^{(\xi)}([0, 1]) \right) \right)} \\ &= \frac{\log_2 \left( p_1^q + p_2^q \right)}{4/3 \frac{2^{2l+2}}{n} + 2 - 10/(3n)}. \end{aligned}$$

Therefore, by Fact 4.1,

$$\beta_{v^{(\xi)}}(q) = \begin{cases} \frac{3}{8} \log_2 \left( p_1^q + p_2^q \right) & \text{for } 0 \leq q \leq 1 \\ \frac{3}{10} \log_2 \left( p_1^q + p_2^q \right) & \text{for } q > 1 \end{cases}$$

and

$$\beta_{\underline{v}^{(\xi)}}(q) := \liminf \beta_n(q) = \begin{cases} \frac{3}{10} \log_2 \left( p_1^q + p_2^q \right) & \text{for } 0 \leq q \leq 1 \\ \frac{3}{8} \log_2 \left( p_1^q + p_2^q \right) & \text{for } q > 1 \end{cases}.$$


Hence, by Theorem 1.1, the upper spectral dimension of  $\Delta_{v^{(\xi)}}$  is given by the unique solution of

$$p_1^{\bar{q}_{v^{(\xi)}}} + p_2^{\bar{q}_{v^{(\xi)}}} = 2^{\frac{8\bar{q}_{v^{(\xi)}}}{3}}.$$

Furthermore,  $q_{\underline{v}^{(\xi)}}$  is the unique solution of

$$\left( 2^{-\frac{10}{3}} p_1 \right)^{q_{\underline{v}^{(\xi)}}} + \left( 2^{-\frac{10}{3}} p_2 \right)^{q_{\underline{v}^{(\xi)}}} = 1.$$

See Figure 5.1 for the two graphs. For the special case  $p_1 = 1/2$  we obtain

Figure 5.1:  $\beta_{\nu^{(\xi)}}$  and  $\liminf \beta_n^{\nu^{(\xi)}}$  with  $p_1 = 0.25$ .

$$\beta_{\nu^{(\xi)}}(q) = \begin{cases} \frac{3}{8}(1-q) & \text{for } 0 \leq q \leq 1 \\ \frac{3}{10}(1-q) & \text{for } q > 1 \end{cases}$$

and

$$\underline{\beta}_{\nu^{(\xi)}}(q) := \liminf \beta_n(q) = \begin{cases} \frac{3}{10}(1-q) & \text{for } 0 \leq q \leq 1 \\ \frac{3}{8}(1-q) & \text{for } q > 1 \end{cases}.$$

In particular in this special case, we have,  $\underline{\dim}_M(\text{supp } \nu) = 3/10$  and  $\overline{\dim}_M(\nu) = 3/8$ . Now, applying Theorem 1.1 in tandem with Proposition 4.13, we conclude

$$\underline{s}_{\nu^{(\xi)}} \leq \frac{\underline{\dim}_M(\nu)}{1 + \underline{\dim}_M(\nu)} = \frac{3}{13} < \frac{3}{11} = \frac{\overline{\dim}_M(\nu)}{1 + \overline{\dim}_M(\nu)} = \overline{s}_{\nu^{(\xi)}}.$$

Now, let us prove that  $\underline{s}_{\nu^{(\xi)}} = \frac{3}{13}$ . Define for  $n \in \mathbb{N}$

$$R(n) = \begin{cases} -2^{2\ell+1} \cdot 8/3 + 10/3 - 4(n - 2^{2\ell+1}) & , 2^{2\ell+1} < n \leq 2^{2\ell+2}, \\ -10/3 \cdot 2^{2\ell+2} + 10/3 - 2(n - 2^{2\ell+2}) & , 2^{2\ell+2} < n \leq 2^{2\ell+3}. \end{cases}$$

First note that for  $m = 1 + \frac{1}{2^\ell}$  and all  $\omega \in W_n$ , there exists  $\eta \in W_2$  such that

$$S_{\omega\eta}^{(\xi)}([0, 1]) \subset \langle S_{\omega}^{(\xi)}([0, 1]) \rangle_m.$$

Hence, we obtain

$$\nu(\langle S_{\omega}^{(\xi)}([0, 1]) \rangle_m) \wedge (\langle S_{\omega}^{(\xi)}([0, 1]) \rangle_m) \geq \frac{2^{-n+R(n)-2}}{m}. \quad (5.1)$$

Recall that by Proposition 3.7, for  $x > 0$ ,

$$\mathcal{N}_m^L(x) = \sup \left\{ \text{card}(P) : P \in \Pi_0 : \min_{C \in P} \nu(\langle I \rangle_m) \wedge (\langle I \rangle_m) \geq \frac{4}{x(m-1)} \right\} \leq N_\nu(x).$$

Now, with  $x_n = m \frac{2^{n-R(n)+4}}{m-1}$  by Eq. (5.1) we have that  $\mathcal{N}_m^L(x_n) \geq 2^n$ , which implies

$$\begin{aligned} \frac{\log(N_V(x_n))}{\log(x_n)} &\geq \left(1 - \frac{R(n)}{n} + 4/n + \frac{\log(m/(m-1))}{n \log(2)}\right)^{-1} \\ &= \begin{cases} \left(1 + \frac{2^{2l+1} \cdot 8/3 - 10/3 + 4(n-2^{2l+1})}{n} + 4/n - \frac{\log(m/(m-1))}{n \log(2)}\right)^{-1} & , 2^{2l+1} < n \leq 2^{2l+2}, \\ \left(1 + \frac{10/3 \cdot 2^{2l+2} - 10/3 + 2(n-2^{2l+2})}{n} + 4/n - \frac{\log(m/(m-1))}{n \log(2)}\right)^{-1} & , 2^{2l+2} < n \leq 2^{2l+3}, \end{cases} \\ &\geq \begin{cases} \left(1 + \frac{8}{3} + (4 - 10/3)/n - \frac{\log(m/(m-1))}{n \log(2)}\right)^{-1} & , 2^{2l+1} < n \leq 2^{2l+2}, \\ \left(1 + \frac{10}{3} + (4 - 10/3)/n - \frac{\log(m/(m-1))}{n \log(2)}\right)^{-1} & , 2^{2l+2} < n \leq 2^{2l+3}. \end{cases} \end{aligned}$$

This shows

$$\liminf_{n \rightarrow \infty} \frac{\log(N_V(x_n))}{\log(x_n)} \geq \frac{1}{1 + 10/3}.$$

Since  $x_n \leq x_{n+1} \leq 2^4 x_n$ ,

$$\liminf_{x \rightarrow \infty} \frac{\log(N_V(x))}{\log(x)} \geq \liminf_{n \rightarrow \infty} \frac{\log(N_V(x_n))}{\log(x_n)} \geq \frac{1}{1 + 10/3}.$$

For  $x > 0$ , choose  $n \in \mathbb{N}$  such that  $x_n < x \leq x_{n+1}$  and therefore

$$\liminf_{x \rightarrow \infty} \frac{\log(N_V(x))}{\log(x)} \geq \liminf_{n \rightarrow \infty} \frac{\log(N_V(x_n))}{\log(x_n) + \log(x_{n+1}/x_n)} \geq \frac{1}{1 + 10/3}.$$

**Example 5.6** (Homogeneous Cantor measure with vanishing  $L^q$ -spectrum). Let us now construct an example for which the spectral dimension equals 0. Let us consider the following environment  $\xi_n := n$ ,  $n \in \mathbb{N}$ , and  $S_1^{(n)}(x) := \frac{x}{4^n}$ ,  $S_2^{(n)}(x) := \frac{x}{4^n} + \frac{4^n - 1}{4^n}$ ,  $x \in [0, 1]$ ,  $p_1^n = p_2^n = 1/2$ . It follows that  $\overline{\dim}_M(K^{(\xi)}) = 0$  with Corollary 1.7 that  $s_V = 0$ . Indeed, let us verify that  $\dim_M(K^{(\xi)}) = 0$ . For every  $\omega \in W_m$ , we have

$$\Lambda(S_\omega^{(\xi)}([0, 1])) = 2^{-(m+1)m}.$$

Now, for  $n \in \mathbb{N}$ , there exists  $m(n) = m \in \mathbb{N}$  such that  $2^{-(m+2)(m+1)} \leq 2^{-n} \leq 2^{-(m+1)m}$ . In particular,  $m(n) \ll \sqrt{n}$  and we conclude

$$\frac{\log(\beta_n(0))}{\log(2^n)} \leq \frac{\log(\beta_{(m+2)(m+1)}(0))}{\log(2^n)} = \frac{m(n) + 1}{n} \rightarrow 0 \text{ for } n \rightarrow \infty.$$

### 5.3. Pure point measures

In this section we give examples of singular measures  $\eta$  on  $(0, 1)$  of pure point type such that the spectral dimension attains any value in  $[0, 1/2]$ . To fix notation, throughout this section, we write  $\eta := \sum p_k \delta_{x_k}$  with  $(p_k) \in (\mathbb{R}_{>0})^{\mathbb{N}}$ ,  $\sum p_k < \infty$  and  $(x_k) \in (0, 1)^{\mathbb{N}}$ .

The first example shows that it is possible for the spectral dimension to be 0 even though the Minkowski dimension is 1.

**Example 5.7.** In this example we consider pure atomic measures  $\eta$  with  $(x_n)_{n \in \mathbb{N}} \in (\mathbb{Q} \cap (0, 1))^{\mathbb{N}}$  such that  $x_n \neq x_m$  for  $m \neq n$ , and  $p_n \leq C_1 e^{-n}$ , for all  $n \in \mathbb{N}$ . We will show that the spectral dimension exists and equals  $s_\eta = 0$ . Define for  $I_k^n := \left(x_k - \frac{b_n}{e^n}, x_k + \frac{b_n}{e^n}\right) \cap [0, 1]$  with  $b_n := \min\{|x_{l_1} - x_{l_2}| : l_1 \neq l_2, l_1, l_2 \leq n\}$  for all  $k = 1, \dots, n$ . Then we have  $x_k - \frac{b_n}{e^n}, x_k + \frac{b_n}{e^n} \notin \mathbb{Q}$  and

$$\max_{k=1, \dots, n} \Lambda(I_k^n) \nu(I_k^n) \leq \sum_{l=1}^{\infty} p_l \max_{k=1, \dots, n} \Lambda(I_k^n) \leq \frac{C_1}{e-1} \frac{2}{e^n}.$$

Let  $A_k^n$  denote the disjoint half open intervals such that  $(0, 1] \setminus \bigcup_{k=1}^n I_k^n = \bigcup_{k=1}^{m(n)} A_k^n$  such that  $m(n) \leq n+1$ . Then we conclude

$$\max_{k=1, \dots, m(n)} \Lambda(A_k^n) \nu(A_k^n) \leq \sum_{l=n+1}^{\infty} p_l \leq C_1 \frac{e^{-n}}{e-1},$$

implying  $\gamma_{2n+1} \leq \frac{2C_1}{e-1} e^{-n}$  and therefore, for every  $m \in \mathbb{N}$  with  $2n+1 \leq m \leq 2n+3$ ,

$$\gamma_m \leq \gamma_{2n+1} \ll e^{-(m-3)/2}.$$

This shows  $\underline{\alpha}_\nu = \infty$  implying by Lemma 3.4 that  $s_\eta = 0$ .

If  $(x_k) \in (0, 1)^{\mathbb{N}}$  is strictly decreasing, then in [FW17]  $\Delta_\eta$  is called Kreĭn-Feller operator of Stieltjes type. We start with a general observation which is a consequence of the Poincaré inequality as stated in Lemma 3.5.

**Lemma 5.8.** Assume that  $(x_k) \in (0, 1)^{\mathbb{N}}$  is strictly decreasing such that for an increasing function  $f : \mathbb{Z}_+ \rightarrow \mathbb{R}_+$  and all  $k \in \mathbb{N}_{>1}$ ,

$$\frac{x_k + x_{k-1}}{2} \sum_{l=k}^{\infty} p_l \leq 1/f(k).$$

Then, for all  $x \geq 0$ , we have  $N_\eta(x) - 2 \leq \check{f}^{-1}(x) := \inf\{n \in \mathbb{N}_{>0} : f(n) \geq x\}$ .

*Proof.* The Poincaré inequality (Lemma 3.5) guarantees for all  $k \in \mathbb{N}$

$$\lambda_{\eta, [0, \frac{x_k + x_{k-1}}{2}]}^1 \geq \frac{1}{\frac{x_k + x_{k-1}}{2} \sum_{l=k}^{\infty} p_l}.$$

Since

$$\frac{1}{\frac{x_k + x_{k-1}}{2} \sum_{l=k}^{\infty} p_l} \geq f(k) \geq x \implies k \geq \check{f}^{-1}(x),$$

using the superadditivity of the Neumann eigenvalue counting function with  $k_x := \check{f}^{-1}(x)$ , by Proposition 2.16, we have

$$N_\eta(x) \leq N_{\eta, [0, \frac{x_{k_x} + x_{k_x-1}}{2}]}(x) + N_{\eta, [\frac{x_{k_x} + x_{k_x-1}}{2}, 1]}(x) + 2 \leq 2 + \check{f}^{-1}(x).$$

□

**Example 5.9** (Dirac comb with exponential decay). If  $(x_n)$  or  $(\sum_{k \geq m} p_k)_m$  decays exponentially, then  $N_\eta(x) \ll \log(x)$ , hence the spectral dimension  $s_\eta$  equals 0. Further, the  $L^q$ -spectrum is given by

$$\beta_\eta(q) = \begin{cases} \overline{\dim}_M(\eta) & q = 0, \\ 0 & q > 0. \end{cases}$$

**Example 5.10** (Dirac comb with at most power law decay). Assume that  $(x_k) \in (0, 1)^{\mathbb{N}}$  is strictly decreasing and

$$p_n \gg n^{-u_1} f_1(n), \quad (x_n - x_{n+1}) \gg n^{-u_2} f_2(n),$$

with  $u_1, u_2 \geq 1$  and  $\lim_{n \rightarrow \infty} \log(f_i(n))/\log(n) = 0$  for  $i = 1, 2$ , then

$$\frac{1}{u_1 + u_2} \leq \underline{s}_\eta$$

and, in particular, if  $u_1 + u_2 = 2$ , we have  $s_\eta = \frac{1}{2}$ . To see this, define

$$I_k := \left[ x_k - \frac{\min\{x_k - x_{k+1}, x_{k-1} - x_k\}}{2}, x_k + \frac{\min\{x_k - x_{k+1}, x_{k-1} - x_k\}}{2} \right]$$

for  $k = 1, \dots, n$ . Then we have  $\nu(\langle I_k \rangle_2) = \nu(I_k)$  and for  $n$  large enough

$$Cn^{-(u_1+u_2)} f_1(n) f_2(n) \leq \min_{k=1, \dots, n} \nu(\langle I_k \rangle_2) \Lambda(\langle I_k \rangle_2)$$

with  $C > 0$  suitable, which implies for every  $\varepsilon > 0$

$$\mathcal{N}_2^L(x) \gg x^{\frac{1}{u_1+u_2+\varepsilon}}$$

proving the claim. Since, in the case  $u_1 + u_2 = 2$ , we always have  $\bar{s}_\eta \leq 1/2$ , it readily follows that  $s_\eta = 1/2$ .

**Example 5.11** (Dirac comb with at most geometric decay and full dimension). Assume that  $(x_k) \in (0, 1)^{\mathbb{N}}$  is strictly decreasing and

$$\frac{f_1(n)}{n} \ll p_n, \quad \frac{f_2(n)}{n} \ll (x_{n-1} - x_n)$$

with  $\lim_{n \rightarrow \infty} \log(f_i(n))/\log(n) = 0$ , for  $i = 1, 2$ , then spectral dimension exists and equals

$$s_\eta = 1/2$$

and, for  $q \in [0, 1]$ ,

$$\beta_\eta(q) = \begin{cases} 1 - q & , q \in [0, 1], \\ 0 & , q > 1. \end{cases}$$

Indeed, observe that Example 5.10 implies  $1/2 \leq \underline{s}_\eta$  and by Theorem 1.1 and Corollary 1.7 we also have  $\underline{s}_\eta \leq \bar{s}_\eta = \bar{q}_\eta \leq 1/2$  which shows  $s_\eta = 1/2$ . The second statement is then a direct consequence of the first part of Corollary 1.4.

The following example shows that the spectral dimension attains every value in  $(0, 1/2)$ .

**Example 5.12** (Dirac comb with power law decay). If

$$\lim_{n \rightarrow \infty} -\log p_n / \log n = u_1 > 1 \text{ and } x_k := (k+1)^{-u_2}, k \in \mathbb{N}, u_2 > 0,$$

then the  $L^q$ -spectrum exists as a limit on the positive half-line and we have

$$\beta_\eta(q) = \begin{cases} \frac{1}{u_2+1} - q^{-\frac{u_1}{u_2+1}} & \text{for } q \in [0, 1/u_1], \\ 0 & \text{for } q > 1/u_1. \end{cases}$$

Consequently,  $\eta$  is  $L^q$ -regular and the spectral dimension exists and equals

$$s_\eta = \frac{1}{u_1 + u_2 + 1}.$$

In particular, for  $u_1 = u_2 + 1$  we have

$$s_\eta = \frac{\dim_M(\eta)}{2}.$$

This can be seen as follows. For every  $\varepsilon > 0$  uniformly in  $n \in \mathbb{N}$ , we have  $n^{-(u_1+\varepsilon)} \ll p_n \ll n^{-u_1+\varepsilon}$ . For suitable  $C > 0$ ,

$$x_m - x_{m+1} = \frac{(m+1)^{u_2} - m^{u_2}}{m^{u_2}(m+1)^{u_2}} = \frac{1}{(m+1)^{u_2}m} \frac{\left(\frac{m+1}{m}\right)^{u_2} - 1}{1/m} \geq \frac{C}{m^{u_2+1}}.$$

If  $2^{-n} < C(m+1)^{-(u_2+1)}$ , then  $m < (2^n C)^{\frac{1}{u_2+1}}$ . Combining these observations, we obtain

$$\sum_{C \in \mathcal{D}_n} \eta(C)^q \geq \sum_{k=1}^{C^{1/(u_2+1)} 2^{n/(u_2+1)}} p_k^q \gg \sum_{k=1}^{C^{1/(u_2+1)} 2^{n/(u_2+1)}} k^{-(u_1+\varepsilon)q} \asymp 2^{n(-(u_1+\varepsilon)q+1)/(u_2+1)}.$$

For  $q \in [0, 1/(u_1 + \varepsilon))$ , this gives

$$\beta_\eta(q) \geq \liminf_n \beta_n^q(q) \geq \frac{1}{u_2+1} - q \frac{u_1 + \varepsilon}{u_2+1}.$$

Letting  $\varepsilon \rightarrow 0$  this shows for  $q \in [0, 1/u_1]$

$$\beta_\eta(q) \geq \liminf_n \beta_n^q(q) \geq \begin{cases} \frac{1}{u_2+1} - q \frac{u_1}{u_2+1} & \text{for } q \in [0, 1/u_1], \\ 0 & \text{for } q > 1/u_1. \end{cases}$$

Moreover, for  $m \geq (2^n C)^{\frac{1}{u_2+1}}$  and  $k 2^{-n} < x_m$ , we have  $k < 2^n (2^n C)^{-\frac{u_2}{u_2+1}}$ . From this

inequality, using the integral test for convergence, we obtain for  $q > 1/(u_1 - \varepsilon)$ ,

$$\begin{aligned} \sum_{k=0}^{2^n-1} \eta(A_k^n)^q &= \sum_{k=0}^{2^n-1} \left( \sum_{k2^{-n} < x_m \leq (k+1)2^{-n}} p_m \right)^q \\ &\ll \sum_{m < C^{1/u_2} 2^{n/u_2}} m^{q(-u_1+\varepsilon)} + C^{-\frac{u_2}{u_2+1}} 2^{\frac{n}{u_2+1}} \left( \sum_{k2^{-n} < x_m \leq (k+1)2^{-n}} p_m \right)^q \\ &\ll 1 + 2^{n \left( \frac{1}{u_2+1} + q \left( -\frac{(u_1-\varepsilon-1)}{u_2} + \frac{(u_1-\varepsilon-1)-u_2}{u_2(u_2+1)} \right) \right)} = 1 + 2^{n \left( \frac{1}{u_2+1} - q \frac{u_1-\varepsilon}{u_2+1} \right)} \ll 1. \end{aligned}$$

Hence,  $\beta_\eta(q) = 0$  for  $q \geq 1/u_1$ . Since  $\beta_\eta(0) = \delta_\eta^* = 1/(u_2 + 1)$ , by the convexity of  $\beta_\eta$ , it follows that for all  $q \in [0, 1/u_1]$ ,

$$\frac{1}{u_2+1} - q \frac{u_1}{u_2+1} \leq \liminf_n \beta_n^q(q) \leq \beta_\eta(q) \leq \frac{1}{u_2+1} - q \frac{u_1}{u_2+1}.$$

Theorem 1.1 in tandem with Theorem 1.2 then gives  $s_\eta = \bar{q}_\eta = (u_1 + u_2 + 1)^{-1}$ .

The last example demonstrates how one can improve Example 5.9, if one knows the exact exponential asymptotics of  $p_n$  and  $x_n$ , which in turn forces an exact logarithmic asymptotics for the eigenvalue counting function. The following elementary lemma is provided for preparation.

**Lemma 5.13.** *Let  $v := p\delta_z$  be with  $z \in (0, 1)$ ,  $p > 0$  and  $0 < a < z$ ,  $0 < b < 1 - z$ . Then the first eigenvalue of  $\Delta_{v, [z-a, z+b]}$  is given by*

$$\lambda_{v, [z-a, z+b]}^1 = \frac{a+b}{pab}.$$

*Proof.* Observe that each function  $f \in \text{dom}(\mathcal{E}_{v, [z-a, z+b]})$  is of triangular shape, i. e.  $f(x) = f_{a,b,c}(x) := c \frac{(x-(z-a))}{a} \mathbb{1}_{[z-a, z]} + c \frac{(z+b-x)}{b} \mathbb{1}_{(z, z+b]}$ . Therefore, using the min-max principle we get

$$\lambda_{\eta, [z-a, z+b]}^1 = \inf \left\{ \frac{\mathcal{E}(\psi, \psi)}{\langle \psi, \psi \rangle_\eta} : \psi \in \text{dom}(\mathcal{E}_{v, [z-a, z+b]})^* \right\} = \inf_{c>0} c^2 \left( \frac{1}{a} + \frac{1}{b} \right) / (c^2 p) = \frac{a+b}{pab}.$$

□

**Example 5.14** (Dirac comb with exponential decay – precise asymptotics). Let us consider the case  $\sum_{k=1}^n p_k = 1 - e^{-\alpha n}$ , and  $x_k := e^{-\beta k}$  for some  $\alpha, \beta > 0$ . Then

$$\lim_{x \rightarrow \infty} \frac{N_\eta(x)}{\log(x)} = \frac{1}{\alpha + \beta}.$$

To see this, we consider the intervals  $A_n := [x_n - (x_n - x_{n+1})/2, x_n + (x_{n-1} - x_n)/2]$ , for  $n > 1$ . Then by Lemma 5.13

$$\lambda_{\eta, A_n}^1 = \frac{e^{(\alpha+\beta)n} 2 (e^\beta - e^{-\beta})}{(1 - e^{-\beta})(e^\alpha - 1)(e^\beta - 1)}$$

implying

$$N_{\eta, [0,1]}(x) \geq \left\lfloor \frac{1}{(\alpha + \beta)} \log \left( x \frac{(1 - e^{-\beta})(e^\alpha - 1)(e^\beta - 1)}{2(e^\beta - e^{-\beta})} \right) \right\rfloor.$$

For the upper bound, we observe that for  $A^k := [0, \frac{x_k + x_{k-1}}{2}]$ , we have

$$\lambda_{\eta, A^k}^1 \geq \frac{1}{\frac{x_k + x_{k-1}}{2} \sum_{\ell=k}^{\infty} p^\ell} = \frac{2}{e^{-\beta k + (k-1)\alpha} (1 + e^{-\beta})} = 2 \frac{e^{(k-1)\alpha + k\beta}}{1 + e^{-\beta}}.$$

Then, for  $x > 4$ , we obtain

$$2 \frac{e^{(k-1)\alpha + k\beta}}{1 + e^{-\beta}} > x \iff k > \frac{1}{(\alpha + \beta)} \log \left( (1 + e^{-\beta}) \frac{e^\alpha x}{2} \right).$$

This gives for  $n_x = \left\lceil \frac{1}{(\alpha + \beta)} \log \left( (1 + e^{-\beta}) \frac{e^\alpha x}{2} \right) \right\rceil$

$$N_{\eta, [0,1]}(x) \leq N_{\eta, A^k}(x) + N_{\eta, [\frac{x_{n_x} + x_{n_x-1}}{2}, 1]}(x) + 2 \leq 3 + (n_x - 1) \leq 2 + \frac{1}{(\alpha + \beta)} \log \left( (1 + e^{-\beta}) \frac{e^\alpha x}{2} \right)$$

and the claim follows.

## References

- [Arz14] P. Arzt, *Eigenvalues of Measure Theoretic Laplacians on Cantor-like Sets*, PhD thesis, Universität Siegen, 2014.
- [BB71] M. Š. Birman and V. V. Borzov, *The asymptotic behavior of the discrete spectrum of certain singular differential operators*, Problems of mathematical physics, No. 5: spectral theory (Russian), 1971, pp. 24–38. MR 0301568
- [BF20] Julien Barral and De-Jun Feng, *On multifractal formalism for self-similar measures with overlaps*, Mathematische Zeitschrift (2020), 1432–1823.
- [BH97] M. T. Barlow and B. M. Hambly, *Transition density estimates for Brownian motion on scale irregular Sierpinski gaskets*, Ann. Inst. H. Poincaré Probab. Statist. **33** (1997), no. 5, 531–557. MR 1473565
- [BNT03] Erik J. Bird, Sze-Man Ngai, and Alexander Teplyaev, *Fractal Laplacians on the unit interval*, Ann. Sci. Math. Québec **27** (2003), no. 2, 135–168. MR 2103098
- [Bor70] V. V. Borzov, *The quantitative characteristics of singular measures*, Problems of Mathematical Physics, No. 4: Spectral Theory, Wave Processes, Izdat. Leningrad. Univ., Leningrad, 1970, pp. 42–47. MR 0281860
- [Bor71] ———, *Quantitative characteristics of singular measures*, pp. 37–42, Springer US, Boston, MA, 1971.

- [BS66] M. Š. Birman and M. Z. Solomjak, *Approximation of functions of the  $W_p^\alpha$ -classes by piece-wise-polynomial functions*, Dokl. Akad. Nauk SSSR **171** (1966), 1015–1018. MR 0209733
- [BS67] ———, *Piecewise polynomial approximations of functions of classes  $W_p^\alpha$* , Mat. Sb. (N.S.) **73** (115) (1967), 331–355. MR 0217487
- [BS70] ———, *The principal term of the spectral asymptotics for "non-smooth" elliptic problems*, Funkcional. Anal. i Priložen. **4** (1970), no. 4, 1–13. MR 0278126
- [BS74] ———, *Quantitative analysis in Sobolev's imbedding theorems and applications to spectral theory*, Tenth Mathematical School (Summer School, Kaciveli/Nalchik, 1972) (Russian), 1974, pp. 5–189. MR 0482138
- [Dav95] E. Brian Davies, *Spectral theory and differential operators*, Cambridge Studies in Advanced Mathematics, Cambridge University Press, 1995.
- [DN15] Da-Wen Deng and Sze-Man Ngai, *Eigenvalue estimates for Laplacians on measure spaces*, J. Funct. Anal. **268** (2015), no. 8, 2231–2260. MR 3318648
- [DN21] ———, *Estimates for sums and gaps of eigenvalues of Laplacians on measure spaces*, Proc. Roy. Soc. Edinburgh Sect. A **151** (2021), no. 2, 842–861. MR 4241300
- [Fag12] Alessandra Faggionato, *Spectral analysis of 1D nearest-neighbor random walks and applications to subdiffusive trap and barrier models*, Electron. J. Probab. **17** (2012), no. 15, 36. MR 2892328
- [Fal14] Kenneth Falconer, *Fractal geometry*, third ed., John Wiley & Sons, Ltd., Chichester, 2014, Mathematical foundations and applications. MR 3236784
- [Fel57] W. Feller, *Generalized second order differential operators and their lateral conditions*, Illinois J. Math. **1** (1957), 459–504. MR 0092046
- [Fen07] De-Jun Feng, *Gibbs properties of self-conformal measures and the multifractal formalism*, Ergodic Theory Dynam. Systems **27** (2007), no. 3, 787–812. MR 2322179
- [Fre05] Uta Freiberg, *Spectral asymptotics of generalized measure geometric Laplacians on Cantor like sets*, Forum Math. **17** (2005), no. 1, 87–104. MR 2110540
- [Fre08] ———, *Prüfer angle methods in spectral analysis of Krein-Feller operators*, Proceedings of RIMS Workshop on Stochastic Analysis and Applications, RIMS Kôkyûroku Bessatsu, B6, Res. Inst. Math. Sci. (RIMS), Kyoto, 2008, pp. 71–84. MR 2407555

- [Fre11] ———, *Refinement of the spectral asymptotics of generalized Krein Feller operators*, Forum Math. **23** (2011), no. 2, 427–445. MR 2787628
- [FT99] Walter Farkas and Hans Triebel, *The distribution of eigenfrequencies of anisotropic fractal drums*, J. London Math. Soc. (2) **60** (1999), no. 1, 224–236. MR 1721826
- [Fuj87] T. Fujita, *A fractional dimension, self-similarity and a generalized diffusion operator*, Probabilistic methods in mathematical physics (Katata/Kyoto, 1985), Academic Press, Boston, MA, 1987, pp. 83–90. MR 933819
- [FW17] Andreas Fleige and Henrik Winkler, *An indefinite inverse spectral problem of Stieltjes type*, Integral Equations Operator Theory **87** (2017), no. 4, 491–514. MR 3648085
- [FZ02] U. Freiberg and M. Zähle, *Harmonic calculus on fractals—a measure geometric approach. I*, Potential Anal. **16** (2002), no. 3, 265–277. MR 1885763
- [Heu98] Yanick Heurteaux, *Estimations de la dimension inférieure et de la dimension supérieure des mesures*, Ann. Inst. H. Poincaré Probab. Statist. **34** (1998), no. 3, 309–338. MR 1625871
- [Heu07] ———, *Dimension of measures: the probabilistic approach*, Publ. Mat. **51** (2007), no. 2, 243–290. MR 2334791
- [HLN06] Jiaxin Hu, Ka-Sing Lau, and Sze-Man Ngai, *Laplace operators related to self-similar measures on  $\mathbb{R}^d$* , J. Funct. Anal. **239** (2006), no. 2, 542–565. MR 2261337
- [Hoc14] Michael Hochman, *On self-similar sets with overlaps and inverse theorems for entropy*, Ann. of Math. (2) **180** (2014), no. 2, 773–822. MR 3224722
- [JKL14] Johannes Jaerisch, Marc Kesseböhmer, and Sanaz Lamei, *Induced topological pressure for countable state Markov shifts*, Stoch. Dyn. **14** (2014), no. 2, 1350016, 31. MR 3190211
- [Kat76] Tosio Kato, *Perturbation theory for linear operators*, second ed., Springer-Verlag, Berlin-New York, 1976, Grundlehren der Mathematischen Wissenschaften, Band 132. MR 0407617
- [Kig01] Jun Kigami, *Analysis on fractals*, Cambridge Tracts in Mathematics, Cambridge University Press, 2001.
- [KK58] I. S. Kac and M. G. Kreĭn, *Criteria for the discreteness of the spectrum of a singular string*, Izv. Vysš. Učebn. Zaved. Matematika **1958** (1958), no. 2 (3), 136–153. MR 0139804
- [KKVW09] Ulrike Kant, Tobias Klauss, Jürgen Voigt, and Matthias Weber, *Dirichlet forms for singular one-dimensional operators and on graphs*, J. Evol. Equ. **9** (2009), no. 4, 637–659. MR 2563669

- [KL93] Jun Kigami and Michel L. Lapidus, *Weyl's problem for the spectral distribution of Laplacians on p.c.f. self-similar fractals*, *Comm. Math. Phys.* **158** (1993), no. 1, 93–125. MR 1243717
- [KN21a] Marc Kesseböhmer and Aljoscha Niemann, *Approximation order of Kolmogorov diameters via  $L^q$ -spectra and applications to polyharmonic operators*, arXiv: [2107.07932](https://arxiv.org/abs/2107.07932) (2021).
- [KN21b] ———, *Spectral asymptotics of Kreĭn-Feller operators for weak Gibbs measures on self-conformal fractals with overlaps*, arXiv: [2107.02616](https://arxiv.org/abs/2107.02616) (2021).
- [KN22] ———, *Spectral dimensions of Kreĭn-Feller operators in higher dimensions*, arXiv: [2202.05247](https://arxiv.org/abs/2202.05247) (2022).
- [KNSW19] M. Kesseböhmer, A. Niemann, T. Samuel, and H. Weyer, *Generalised Kreĭn-Feller operators and gap diffusions under transformations of measure spaces*, arXiv: [1909.08832](https://arxiv.org/abs/1909.08832) (2019).
- [Kre51] M. G. Kreĭn, *Determination of the density of a nonhomogeneous symmetric cord by its frequency spectrum*, *Doklady Akad. Nauk SSSR (N.S.)* **76** (1951), 345–348. MR 0042045
- [Küc86] U. Küchler, *On sojourn times, excursions and spectral measures connected with quasidiffusions*, *J. Math. Kyoto Univ.* **26** (1986), no. 3, 403–421.
- [KZ04] Marc Kesseböhmer and Sanguo Zhu, *Quantization dimension via quantization numbers*, *Real Anal. Exchange* **29** (2003/04), no. 2, 857–866. MR 2083820
- [Min20] Lenon Alexander Minorics, *Spectral asymptotics for Krein-Feller operators with respect to  $V$ -variable Cantor measures*, *Forum Math.* **32** (2020), no. 1, 121–138. MR 4048458
- [Naz04] A. I. Nazarov, *Logarithmic asymptotics of small deviations for some Gaussian processes in the  $L_2$ -norm with respect to a self-similar measure*, *Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI)* **311** (2004), no. Veroyatn. i Stat. 7, 190–213, 301. MR 2092208
- [Nga97] Sze-Man Ngai, *A dimension result arising from the  $L^q$ -spectrum of a measure*, *Proc. Amer. Math. Soc.* **125** (1997), no. 10, 2943–2951. MR 1402878
- [Nga11] ———, *Spectral asymptotics of Laplacians associated with one-dimensional iterated function systems with overlaps*, *Canad. J. Math.* **63** (2011), no. 3, 648–688. MR 2828537
- [NS95] K. Naimark and M. Solomyak, *The eigenvalue behaviour for the boundary value problems related to self-similar measures on  $\mathbb{R}^d$* , *Math. Res. Lett.* **2** (1995), no. 3, 279–298. MR 1338787

- [NS01] ———, *Eigenvalue distribution of some fractal semi-elliptic differential operators: combinatorial approach*, Integral Equations Operator Theory **40** (2001), no. 4, 495–506. MR 1839473
- [NTX18] Sze-Man Ngai, Wei Tang, and Yuanyuan Xie, *Spectral asymptotics of one-dimensional fractal Laplacians in the absence of second-order identities*, Discrete Contin. Dyn. Syst. **38** (2018), no. 4, 1849–1887. MR 3809018
- [NX19] Sze-Man Ngai and Yuanyuan Xie,  *$L^q$ -spectrum of self-similar measures with overlaps in the absence of second-order identities*, J. Aust. Math. Soc. **106** (2019), no. 1, 56–103. MR 3897401
- [NX20] ———, *Spectral asymptotics of Laplacians related to one-dimensional graph-directed self-similar measures with overlaps*, Ark. Mat. **58** (2020), no. 2, 393–435. MR 4176086
- [NX21] ———, *Spectral asymptotics of Laplacians associated with a class of higher-dimensional graph-directed self-similar measures*, Nonlinearity **34** (2021), no. 8, 5375–5398.
- [PS00] Yuval Peres and Boris Solomyak, *Existence of  $L^q$  dimensions and entropy dimension for self-conformal measures*, Indiana Univ. Math. J. **49** (2000), no. 4, 1603–1621. MR 1838304
- [Rie93] Rolf Riedi, *An improved Multifractal Formalism and Self-Affine Measures*, Ph.D. thesis, ETH Zuerich, 1993.
- [Rie95] ———, *An improved multifractal formalism and self-similar measures*, J. Math. Anal. Appl. **189** (1995), no. 2, 462–490. MR 1312056
- [Shm19] Pablo Shmerkin, *On Furstenberg’s intersection conjecture, self-similar measures, and the  $L^q$  norms of convolutions*, Ann. of Math. (2) **189** (2019), no. 2, 319–391. MR 3919361
- [Sol94] Michael Solomyak, *Eigenvalue estimates for a class of operators related to self-similar measures*, Journées “Équations aux Dérivées Partielles” (Saint-Jean-de-Monts, 1994), École Polytech., Palaiseau, 1994, pp. Exp. No. XI, 6. MR 1298682
- [SV95] M. Solomyak and E. Verbitsky, *On a spectral problem related to self-similar measures*, Bull. London Math. Soc. **27** (1995), no. 3, 242–248. MR 1328700
- [Tri92] Hans Triebel, *Higher analysis*, Hochschulbücher für Mathematik. [University Books for Mathematics], Johann Ambrosius Barth Verlag GmbH, Leipzig, 1992, Translated from the German by Bernhardt Simon [Bernhard Simon] and revised by the author. MR 1193032
- [Tri97] ———, *Fractals and spectra*, Monographs in Mathematics, vol. 91, Birkhäuser Verlag, Basel, 1997, Related to Fourier analysis and function spaces. MR 1484417

- [UH59] Toshio Uno and Imsik Hong, *Some consideration of asymptotic distribution of eigenvalues for the equation  $d^2u/dx^2 + \lambda q(x)u = 0$* , Jpn. J. Math. **29** (1959), 152–164. MR 118891
- [Vol05] Hans Volkmer, *Eigenvalue problems of atkinson, feller and krein, and their mutual relationship*, Electronic Journal of Differential Equations **2005** (2005).
- [Wey11] H. Weyl, *Über die asymptotische Verteilung der Eigenwerte*, Nachr. Ges. Wiss. Göttingen, Math.-Phys. Kl. **1911** (1911), 110–117 (German).