ON THE PERRON SOLUTION OF THE CALORIC DIRICHLET PROBLEM: AN ELEMENTARY APPROACH

ALESSIA E. KOGOJ AND ERMANNO LANCONELLI

" I like easy tricks." Louis Nirenberg

ABSTRACT. By an easy trick taken from caloric polynomial theory we construct a family \mathscr{B} of *almost regular* domains for the caloric Dirichlet problem. \mathscr{B} is a basis of the Euclidean topology. This allows to build, with a basically elementary procedure, the Perron solution to the caloric Dirichlet problem on every bounded domain.

1. INTRODUCTION AND KEY THEOREM

As it is very well known today, the construction of the Perron solution to the Dirichlet problem for the heat equation on a general bounded domain only rests upon three basic principles: the *caloric maximum principle*; the *convergence principle*, i.e., the closure of the sheaf of the caloric functions with respect to the uniform convergence; the *solvability principle*, i.e., the solvability of the caloric first boundary value problem on the open sets of a basis of the Euclidean topology.

The first one of this principles is very elementary, and the second one is a simple consequence of some good properties of the Gauss–Weierstrass kernel, the fundamental solution of the heat equation. On the contrary, the proof of the solvability principle has been always considered in literature a difficult task, requiring Volterra integral equation theory, or double layer potential method, or an involved procedure based on a reflection principle (see e.g., respectively, Bauer [Bau66, Chapter 1, Section 2], Watson [Wat12, Chapter 2, Section 2.2], Constantinescu and Cornea [CC72, Chapter 3, Section 3.3]).

The aim of this note is to draw attention to an easy trick - borrowed from caloric polynomial theory - allowing to construct with a very elementary procedure, a basis of open sets on which the first boundary value problem for the heat equation is solvable. The crucial point of this procedure is Theorem 1.1 below, in which w denotes the polynomial

$$w(z) = w(x,t) := t - |x|^2.$$

Here and in what follows

$$z = (x, t), \quad x \in \mathbb{R}^N, \quad t \in \mathbb{R},$$

1

denotes the point of \mathbb{R}^{N+1} ; |x| is the Euclidean norm of x.

²⁰¹⁰ Mathematics Subject Classification. 35K05; 35K20.

Key words and phrases. Heat equation, caloric Dirichlet problem, Perron solution.

We will use the notation H to denote the *heat operator*

$$H := \Delta - \partial_t,$$

where $\Delta := \sum_{j=1}^{N} \partial_{x_j}^2$ is the Laplacian in \mathbb{R}^N . We call *caloric* the smooth functions solutions to Hu = 0. If $\Omega \subseteq \mathbb{R}^{N+1}$ is open, we will denote by $\mathscr{C}(\Omega)$ the linear space of the caloric functions in Ω .

To our aim it is convenient to fix some more notations. If $\alpha = (\alpha_1, \ldots, \alpha_N, \alpha_{N+1})$ is a multi-index with non negative integer components, we let

$$|\alpha|_c = caloric \ hight \ of \ \alpha := \alpha_1 + \ldots + \alpha_N + 2\alpha_{N+1}$$

A polynomial in \mathbb{R}^{N+1} is a function of the kind

$$p(z) = \sum_{|\alpha|_c \le m} a_{\alpha} z^{\alpha}, \quad a_{\alpha} \in \mathbb{R} \text{ for every } \alpha$$

where $m \in \mathbb{Z}$, $m \ge 0$. In this case we say that p has caloric degree $\le m$; we will say that p has caloric degree equal to m if $\sum_{|\alpha|_c=m} a_{\alpha} z^{\alpha}$ is not identically zero.

Here is the key theorem of our note.

Theorem 1.1. Le p be a polynomial in \mathbb{R}^{N+1} . Then, there exists a unique polynomial q in \mathbb{R}^{N+1} such that

(1.1)
$$H(wq) = -Hp$$

Proof. Let us denote by m the caloric degree of -Hp and by \mathcal{P}_m the linear space of the polynomials in \mathbb{R}^{N+1} having caloric degree less than or equal to m. Since w has caloric degree two, then $H(wq) \in \mathcal{P}_m$ if $q \in \mathcal{P}_m$; therefore

$$q \longmapsto T(q) := H(wq)$$

maps \mathcal{P}_m in \mathcal{P}_m . To prove that (1.1) has a unique solution q it is (more than) enough to show that T is surgetive and injective. To this end, since \mathcal{P}_m is a linear space of finite dimension and T linearly maps \mathcal{P}_m in \mathcal{P}_m , we only have to prove that T is injective. Let $q \in \mathcal{P}_m$ be such that T(q) = 0. Then,

$$u := wq$$

is caloric in \mathbb{R}^{N+1} hence, in particular, in the open region

(1.2)
$$P := \{ (x,t) \in \mathbb{R}^{N+1} : t > |x|^2 \}.$$

Moreover, since w = 0 on

(1.3)
$$\partial P = \{ (x,t) \in \mathbb{R}^{N+1} : t = |x|^2 \},\$$

 $u|_{\partial P} = wq|_{\partial P} = 0$. Then, by the caloric maximum principle (see e.g. [BB19, Theorem 8.2]) u = 0 in P. Since w > 0 in P, this implies q = 0 in P, hence in \mathbb{R}^{N+1} . We have so proved that q = 0 if T(q) = 0, that is the injectivity of T, completing the proof.

From the previous theorem one immediately obtains the following corollary, in which ∂P is the closed set in (1.3), boundary of the region P in (1.2).

Corollary 1.2. Let p be a polynomial in \mathbb{R}^{N+1} . Then, there exists a unique polynomial u_p in \mathbb{R}^{N+1} such that

(1.4)
$$\begin{cases} Hu_p = 0 & in \mathbb{R}^{N+1}, \\ u_p = p & on \partial P. \end{cases}$$

Proof. Let q be a polynomial satisfying (1.1). Then $u_p = wq + p$ solves (1.4). Moreover, if v is any polynomial solving (1.4), then $v - u_p$ is caloric in \mathbb{R}^{N+1} hence in P - and $v - u_p = 0$ on ∂P . The caloric maximum principle implies $v - u_p = 0$ in P, hence in \mathbb{R}^{N+1} . Then $v = u_p$, that is the uniqueness part of the Corollary.

2. A proof of the solvability principle: A basis of H-almost regular domains


Let $z_0 = (x_0, t_0) \in \mathbb{R}^{N+1}$ and let r > 0. We call:

(i) caloric bowl of bottom z_0 and opening r the open set

$$B(z_0, r) := \{ (x, t) \in \mathbb{R}^{N+1} : |x - x_0|^2 < t - t_0 < r^2 \};$$

(ii) normal or caloric boundary of
$$B(z_0, r)$$
 the subset of $\partial B(z_0, r)$

$$\partial_n B(z_0, r) := \{ (x, t) \in \mathbb{R}^{N+1} : |x - x_0|^2 = t - t_0, \ 0 \le t - t_0 \le r^2 \}.$$



$$\mathscr{B} := \{ B(z_0, r) : z_0 \in \mathbb{R}^{N+1}, r > 0 \}$$

is a basis of the Euclidean topology.

The aim of this section is to show that every caloric bowl is H-almost regular in the sense of the following theorem.

Theorem 2.1. Let $z_0 \in \mathbb{R}^{N+1}$ and r > 0 be arbitrarily fixed and let $B = B(z_0, r)$ be the caloric bowl of bottom z_0 and opening r. Then for every $\varphi \in C(\partial_n B, \mathbb{R})$ there exists a unique solution to the boundary value problem

(2.1)
$$\begin{cases} Hu = 0 \quad in B, \\ u = \varphi \quad on \ \partial_n B \end{cases}$$

Precisely: there exists a unique function u_{φ}^B caloric in B and continuous up to $B \cup \partial_n B$ such that

$$u_{\varphi}^{B}(z) = \varphi(z) \text{ for every } z \in \partial_{n}B.$$

Moreover $u_{\varphi}^B \geq 0$ if $\varphi \geq 0$.

Proof. The uniqueness of u_{φ}^{B} and its positivity when φ is positive is a direct consequence of the caloric maximum principle (see e.g. [BB19, Theorem 8.2]). To prove the existence of u_{φ}^{B} we may, and do, assume $z_{0} = (0, 0)$, since H is left translation invariant and

$$B(z_0, r) = z_0 + B(0, r), \quad 0 \in \mathbb{R}^{N+1}.$$

Let $\varphi \in C(\partial_n B, \mathbb{R})$ and let $(p_k)_{k \in \mathbb{N}}$ be a sequence of polynomials in \mathbb{R}^{N+1} uniformly convergent to φ on $\partial_n B$. By using the notation of Corollary 1.2, we let

$$u_k := u_{p_k}, \quad k \in \mathbb{N}.$$

Then, u_k is caloric in B since it is caloric in \mathbb{R}^{N+1} . Moreover $u_k = p_k$ on ∂P in (1.3), hence on $\partial_n B$. Therefore, by the caloric maximum principle,

$$\max_{\overline{B}} |u_k - u_h| = \max_{\partial_n B} |u_k - u_h| = \max_{\partial_n B} |p_k - p_h| \longrightarrow 0 \text{ as } k, h \longrightarrow \infty.$$

From this, one gets the existence of a function $u \in C(\overline{B}, \mathbb{R})$ which is caloric in B - thanks to the convergence principle - and such that

$$u|_{\partial_n B} = \lim_{k \to \infty} u_k|_{\partial B} = \lim_{k \to \infty} p_k|_{\partial B} = \varphi.$$

Thus, the function u is the requested function u^B_{ω} .

Remark 2.2. From the above proof it follows that the solution u_{φ} of (2.1) actually is continuous up to \overline{B} .

For completeness reasons, and also to stress its elementary character, in Appendix A we will give a simple proof of the convergence principle. In Appendix B we sketch how to construct the caloric Perron solutions starting from Theorem 2.1.

APPENDIX A. THE CONVERGENCE PRINCIPLE

• The fundamental solution of H. The Gauss-Weierstrass kernel, i.e., the function

$$\Gamma: \mathbb{R}^{N+1} \longrightarrow \mathbb{R}, \quad \Gamma(x,t) = \begin{cases} 0 & \text{if } t \le 0, \\ (4\pi t)^{-\frac{N}{2}} \exp\left(-\frac{|x|^2}{4t}\right) & \text{if } t > 0, \end{cases}$$

$$\Box$$

is the fundamental solution with pole at the origin of the heat operator in \mathbb{R}^{N+1} . Γ is smooth in $\mathbb{R}^{N+1} \setminus \{(0,0)\}$ and locally summable in \mathbb{R}^{N+1} . Its crucial property is the following one:

(A.1)
$$\varphi(z) = -\int_{\mathbb{R}^{N+1}} \Gamma(z-\zeta) H\varphi(\zeta) \, d\zeta$$

for every $z \in \mathbb{R}^{N+1}$ and for every $\varphi \in C_0^{\infty}(\mathbb{R}^{N+1}, \mathbb{R})$.

• Caloric norm and caloric disks. If $z = (x, t) \in \mathbb{R}^{N+1}$ we let

$$||z|| = caloric norm of z := (|x|^4 + t^2)^{\frac{1}{4}}$$

We call *caloric disk* of center $z_0 \in \mathbb{R}^{N+1}$ and radius r > 0 the open set

$$D(z_0, r) := \{ z \in \mathbb{R}^{N+1} : \| z - z_0 \| < r \}.$$

• A representation formula for caloric functions. An easy consequence of property (A.1) is the following representation theorem

Theorem A.1. Let $\Omega \subseteq \mathbb{R}^{N+1}$ be open. For every caloric disk $D = D(z_0, r)$ such that $\overline{2D} := \overline{D(z_0, 2r)} \subseteq \Omega$, there exists a function

$$(z,\zeta) \longmapsto K_D(z,\zeta)$$

of class C^{∞} in an open set containing $\overline{D \times 2D}$ such that

$$u(z) = \int_{2D} K_D(z,\zeta) u(\zeta) \ d\zeta \quad \text{for every } z \in D$$

and for every u caloric in Ω .

Proof. Let $\psi \in C_0^{\infty}(2D, \mathbb{R})$ be such that $\psi \equiv 1$ in a neighbourhood of \overline{D} . Then $\psi u \in C_0^{\infty}(2D, \mathbb{R})$ and $u = \psi u$ in D. As a consequence, by (A.1),

$$\begin{split} u(z) &= -\int_{2D} \Gamma(z-\zeta) H(\psi u)(\zeta) \ d\zeta \\ &= -\int_{D} \Gamma(z-\zeta) \left(u(\zeta) H\psi(\zeta) + 2\langle \nabla u(\zeta), \nabla \psi(\zeta) \rangle \right) \ d\zeta \quad \text{for every } z \in D, \end{split}$$

where ∇ and \langle , \rangle denote, respectively, the gradient and the inner product in \mathbb{R}^N . Integrating by parts the second summand at the last right hand side we find

$$u(z) = \int_{2D} K_D(z,\zeta) u(\zeta) \ d\zeta,$$

where, denoting by Δ the Laplacian with respect to the spatial variables,

$$K_D(z,\zeta) = -\Gamma(z-\zeta)H\psi(\zeta) - 2\langle\nabla\Gamma(z-\zeta),\nabla\psi(\zeta)\rangle + \Gamma(z-\zeta)\Delta\psi(\zeta).$$

Then K_D is a smooth function in a neighbourhood of $D \times 2D$ and we are done.

• The convergence principle.

As recalled in the Introduction the convergence principle is the statement of the following theorem.

Theorem A.2. Let (u_k) be a sequence of caloric functions in an open set $\Omega \subseteq \mathbb{R}^{N+1}$. Suppose (u_k) uniformly convergent to a function $u : \Omega \longrightarrow \mathbb{R}$ on every compact subset of Ω . Then,

$$u \in C^{\infty}(\Omega, \mathbb{R})$$
 and $Hu = 0$ in Ω .

Proof. It is enough to show that u is smooth and caloric in every caloric disk D such that $\overline{2D} \subseteq \Omega$. So, let D be such a disk. Then, by Theorem A.1,

(A.2)
$$u_k(z) = \int_{2D} K_D(z,\zeta) u_k(\zeta) \ d\zeta \text{ for every } z \in D$$

and for every $k \in \mathbb{N}$. Since (u_k) is uniformly convergent on $\overline{2D}$, letting k go to infinity in (A.2), we get

$$u(z) = \int_{2D} K_D(z,\zeta) u(\zeta) \ d\zeta$$
 for every $z \in D$.

The smoothness of the kernel K_D implies $u \in C^{\infty}(D, \mathbb{R})$. To show that u is caloric in D we argue as follows. Denoting

$$H^* := \Delta + \partial_t$$

the formal adjoint of H, for every $\varphi \in C_0^{\infty}(D, \mathbb{R})$ we have

$$\int_D (Hu)\varphi \ dz = \int_D u \ H^*\varphi \ dz = \lim_{k \to \infty} \int_D u_k \ H^*\varphi \ dz = \lim_{k \to \infty} \int_D (Hu_k) \ \varphi \ dz = 0,$$

since u_k is caloric for every $k \in \mathbb{N}$. Hence,

$$\int_{D} (Hu) \varphi \, dz = 0 \quad \text{for every } \varphi \in C_0^{\infty}(\Omega, \mathbb{R}).$$

This implies Hu = 0 in D, completing the proof.

• A consequence of convergence principle.

A family \mathscr{F} of real functions in an open set $\Omega \subseteq \mathbb{R}^{N+1}$ is said up directed if for every $u, v \in \mathscr{F}$ there exists $w \in \mathscr{F}$ such that

$$u \leq w, \quad v \leq w.$$

Then Theorem A.2 and a Real Analysis lemma imply the following result.

Theorem A.3. Let \mathscr{F} be an up directed family of caloric functions in an open set $\Omega \subseteq \mathbb{R}^{N+1}$. Let $u : \Omega \longrightarrow] - \infty, \infty]$,

$$u := \sup \mathscr{F}.$$

If u is bounded above on every compact subset of Ω , then

$$u \in C^{\infty}(\Omega, \mathbb{R})$$
 and $Hu = 0$ in Ω .

 $\mathbf{6}$

Proof. A Real Analysis Lemma (see e.g. [AG01, Lemma 3.7.1]) implies the existence of a monotone increasing sequence (u_k) of functions in \mathscr{F} such that

$$\lim_{k \to \infty} u_k = u \quad \text{pointwise in } \Omega.$$

Now, we can argue as in the proof of Theorem A.2. Let D be a caloric disk such that $\overline{2D} \subseteq \Omega$. Then identity (A.2) holds for every function $u_k, k \in \mathbb{N}$. Since

$$u_1 \le u_k$$
 and $\sup_{2D} u_k \le \sup_{2D} u < \infty$,

by Lebesgue Dominate Convergence Theorem, letting k go to infinity in (A.2), one obtains

$$u(z) = \int_{2D} K_D(z,\zeta) \ u(\zeta) \ d\zeta \qquad \forall z \in D.$$

The smoothness of the kernel K implies the smoothness of u in D so that, since (u_k) is increasing, by Dini Theorem (u_k) converges uniformly on every compact subset of D. As a consequence, by Theorem A.1, u is caloric in D. This completes the proof of Theorem A.3 since D is any caloric disk such that $\overline{2D} \subseteq \Omega$.

• The operator $f \mapsto h_f^B$.

Let B be a caloric bowl. As an application of Theorem A.3 we show how to extend the operator

$$C(\partial_n B, \mathbb{R}) \ni \varphi \longmapsto u_{\varphi}^B \in \mathscr{C}(B),$$

defined in Theorem 2.1, to the bounded above lover semicontinuous functions.

Let B be a caloric bowl and let

$$f:\partial_n B\longrightarrow]-\infty,\infty[$$

be a bounded above lover semicontinuous function. Define

$$\mathscr{F}(B,f) := \{ u^B_{\varphi} \ : \ \varphi \in C(\partial_n B, \mathbb{R}), \ \varphi \leq f \}$$

and

$$h_f^B := \sup \mathscr{F}(B, f).$$

Obviously, if the function f is continuous, then $h_f^B = u_f^B$, so that $f \mapsto h_f^B$ is an extension of $f \mapsto u_f^B$. By using the caloric maximum principle, it is easy to show that $\mathscr{F}(B, f)$ is up directed and that

$$h_f^B \le m$$
 if $m = \sup_{\partial_n B} f$.

Then, by Theorem A.3,

$$h_f^B \in \mathscr{C}(B)$$

APPENDIX B. THE CALORIC PERRON SOLUTION

• Mean Value Theorem for caloric functions. For every $z_0 \in \mathbb{R}^{N+1}$ and every r > 0 we let

$$\Omega_r(z_0) =$$
 Pini–Watson ball with pole at z_0 and radius r

$$:= \left\{ z \in \mathbb{R}^{N+1} : \Gamma(z_0 - z) > (4\pi r)^{-\frac{N}{2}} \right\}$$

and

$$\begin{split} W(z) &= W(x,t) &= & \text{Watson kernel} \\ &:= & \frac{1}{4} \left(\frac{|x|}{t} \right)^2. \end{split}$$

We also denote by $M_r(u)(z_0)$ the average operator

$$M_r(u)(z_0) = \left(\frac{1}{4\pi r}\right)^{\frac{N}{2}} \int_{\Omega_r(z_0)} u(\zeta) \ W(z-\zeta) \ d\zeta.$$

Then, the following theorem holds

Theorem B.1. Let $\Omega \subseteq \mathbb{R}^{N+1}$ be open and let $u \in C(\Omega, \mathbb{R})$. The following statements are equivalent:

(i) For every Pini–Watson ball $\Omega_r(z)$ with closure contained in Ω

$$u(z) = M_r(u)(z).$$

(ii) For every $z \in \Omega$ there exists r(z) > 0 such that

$$u(z) = M_r(u)(z)$$
 for $0 < r < r(z)$.

(iii) $u \in C^{\infty}(\Omega, \mathbb{R})$ and

$$Hu = 0 \ in \ \Omega.$$

Proof. See Watson [Wat12, Chapter 1 and Chapter 2].

• Supercaloric and subcaloric functions. Let $\Omega \subseteq \mathbb{R}^{N+1}$ be open and let

$$u:\Omega\longrightarrow\mathbb{R}$$

be a lower semicontinuous function. We say that u is supercaloric in Ω if for every $z \in \Omega$ there exists r(z) > 0 such that

$$u(z) \ge M_r(u)(z) \qquad \text{for } 0 < r < r(z).$$

We say that u is subcaloric in Ω if -u is supercaloric. With $\overline{\mathscr{C}}(\Omega)$ ($\underline{\mathscr{C}}(\Omega)$), we denote the family of the supercaloric (subcaloric) functions in Ω . It can be elementarily proved that a sufficiently smooth function u is supercaloric (subcaloric) in an open set Ω if and only if

$$Hu \leq 0$$
 in Ω $(Hu \geq 0$ in Ω).

• A caloric Perron-type regularitation. To begin with, we fix some notation. If B is a caloric bowl we denote

$$B := B \smallsetminus \partial_n B.$$

Equivalently

$$\hat{B} = B \cup \operatorname{top}(B),$$

where

$$top(B) := \partial B \smallsetminus \partial_n B$$

Let us consider a bounded above supercaloric function u in an open set $\Omega \subseteq \mathbb{R}^{N+1}$. If $B = B(z_0, r)$ is a caloric bowl such that $2B := B(z_0, 2r) \subseteq \Omega$, we define

$$u_B:\Omega\longrightarrow\mathbb{R}$$

as follows:

$$u_B(z) = u(z)$$
 if $z \notin \hat{B}$,

8

and

$$u_B(z) = h_f^{2B}(z) \quad \text{if} \ z \in \hat{B},$$

where $f = u|_{\partial_n 2B}$.

We want to explicitly remark that u_B is *caloric* in *B* and *continuous* up to \hat{B} . The function u_B is what we call *caloric Perron-type regularization of u in B*. It satisfies all the crucial properties of the classical harmonic Perron regularization. Precisely, the following theorem holds.

Theorem B.2. Let u be a bounded above supercaloric function in an open set $\Omega \subseteq \mathbb{R}^{N+1}$ and let B be a bowl such that $2B \subseteq \Omega$. Then,

- (i) $u_B \in \overline{\mathscr{C}}(\Omega);$
- (*ii*) $u_B \leq u$;
- (iii) u_B is caloric in B and continuous in \hat{B} ;
- (iv) if $v \in \overline{\mathscr{C}}(\Omega)$ and $v \leq u$ then $v_B \leq u_B$.

Proof. The proof of this theorem follows basic standard lines in harmonic and caloric Potential Theory. It uses in a crucial way the properties of the operator $f \mapsto u_f^{2B}$ and the minimum principle for supercaloric functions (for this principle we directly refer to Watson's monograph [Wat12, Theorem 3.11].

We close this appendix with the following point.

• The caloric Perron solution.

Let $\Omega \subseteq \mathbb{R}^{N+1}$ be open and bounded and let $\varphi \in C(\partial\Omega, \mathbb{R})$. We let

 $\overline{\mathcal{U}}^\Omega_\varphi := \{ u \in \overline{\mathscr{C}}(\Omega) \ : \ u \text{ bounded above, } \liminf_{x \to y} u(x) \geq \varphi(y) \ \forall y \in \partial \Omega \},$

and

$$\overline{H}^{\Omega}_{\varphi} := \inf \overline{\mathcal{U}}^{\Omega}_{\varphi}$$

We also let

$$\underline{H}^\Omega_\varphi:=-\overline{H}^\Omega_{-\varphi}$$

From the quoted above minimum principle for supercaloric functions one easily gets

$$m \leq \underline{H}^{\Omega}_{\varphi} \leq \overline{H}^{\Omega}_{\varphi} \leq M,$$

where

$$m = \min_{\partial \Omega} \varphi$$
 and $M = \max_{\partial \Omega} \varphi$.

Actually, a stronger result holds

Theorem B.3. For every $\varphi \in C(\partial\Omega, \mathbb{R})$ one has

(i)
$$\overline{H}^{\Omega}_{\varphi} \in \mathscr{C}(\Omega)$$

(ii) $\overline{H}^{\Omega}_{\varphi} = \underline{H}^{\Omega}_{\varphi}$.

This is the caloric version of the celebrated Perron–Wiener Theorem for the harmonic functions. It can be proved with a standard procedure in which the caloric Perron-type regularization plays the crucial rôle.

Acknowledgment

The first author has been partially supported by the Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM).

References

- [AG01] D. H. Armitage and S. J. Gardiner. Classical potential theory. Springer Monographs in Mathematics. Springer-Verlag London, Ltd., London, 2001.
- [Bau66] H. Bauer. Harmonische R\u00e5ume und ihre Potentialtheorie. Ausarbeitung einer im Sommersemester 1965 an der Universit\u00e5t Hamburg gehaltenen Vorlesung. Lecture Notes in Mathematics, No. 22. Springer-Verlag, Berlin-New York, 1966.
- [BB19] S. Biagi and A. Bonfiglioli. An introduction to the geometrical analysis of vector fields with applications to maximum principles and Lie groups. World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2019.
- [CC72] C. Constantinescu and A. Cornea. Potential theory on harmonic spaces. Springer-Verlag, New York-Heidelberg, 1972. With a preface by H. Bauer, Die Grundlehren der mathematischen Wissenschaften, Band 158.
- [Wat12] N. A. Watson. Introduction to Heat Potential Theory. Mathematical Surveys and Monographs vol.182, Amer. Math. Soc., Providence RI, 2012.

DIPARTIMENTO DI SCIENZE PURE E APPLICATE (DISPEA), UNIVERSITÀ DEGLI STUDI DI URBINO CARLO BO, PIAZZA DELLA REPUBBLICA 13, 61029 URBINO (PU), ITALY.

Email address: alessia.kogoj@uniurb.it

DIPARTIMENTO DI MATEMATICA, ALMA MATER STUDIORUM UNIVERSITÀ DI BOLOGNA, PIAZZA DI PORTA SAN DONATO 5, 40126 BOLOGNA, ITALY.

Email address: ermanno.lanconelli@uniurb.it