

# General Remarks on the One-loop Contributions to the Muon Anomalous Magnetic Moment

Bingrong Yu <sup>a, b \*</sup>, Shun Zhou <sup>a, b †</sup>

<sup>a</sup>Institute of High Energy Physics, Chinese Academy of Sciences, Beijing 100049, China

<sup>b</sup>School of Physical Sciences, University of Chinese Academy of Sciences, Beijing 100049, China

## Abstract

The latest measurement of the muon anomalous magnetic moment  $a_\mu \equiv (g_\mu - 2)/2$  at the Fermi Laboratory has found a  $4.2\sigma$  discrepancy with the theoretical prediction of the Standard Model (SM). Motivated by this exciting progress, we investigate in the present paper the general one-loop contributions to  $a_\mu$  within the SM and beyond. First, different from previous works, the analytical formulae of relevant loop functions after integration are now derived and put into compact forms with the help of the Passarino-Veltman functions. Second, given the interactions of muon with new particles running in the loop, we clarify when the one-loop contribution to  $a_\mu$  could take the correct positive sign as desired. Third, possible divergences in the zero- and infinite-mass limits are examined, and the absence of any divergences in the calculations leads to some consistency conditions for the construction of ultraviolet complete models. Applications of our general formulae to specific models, such as the SM, seesaw models,  $Z'$  and leptoquark models, are also discussed.

---

\*E-mail: yubr@ihep.ac.cn

†E-mail: zhoush@ihep.ac.cn (corresponding author)

# 1 Introduction

The Standard Model (SM) of particle physics, based on the relativistic quantum field theory with gauge symmetry  $SU(3)_C \otimes SU(2)_L \otimes U(1)_Y$ , has proved to be a very successful theory in describing strong, weak and electromagnetic interactions among all discovered elementary particles. Recently, the Muon  $(g - 2)$  Collaboration at Fermi National Accelerator Laboratory has announced their precise measurement of the muon anomalous magnetic moment  $a_\mu \equiv (g_\mu - 2)/2$  with  $g_\mu$  being the spin  $g$ -factor of muon [1], which combined with the previous E821 measurement at Brookhaven National Laboratory in 2006 [2] leads to a  $4.2\sigma$  discrepancy with the SM prediction [3–6], namely,

$$\Delta a_\mu \equiv a_\mu^{\text{Exp}} - a_\mu^{\text{SM}} = (251 \pm 59) \times 10^{-11} . \quad (1)$$

If confirmed by more precise measurements and theoretical predictions [3, 7], such a discrepancy will definitely signify new physics beyond the SM (BSM). There have already been a great number of theoretical works attempting to explain this discrepancy by new physics [8–85]. See, e.g., Refs. [86–89], for comprehensive reviews on this topic and also for more earlier references.

In order to account for the deviation  $\Delta a_\mu$  in Eq. (1) by new physics, one may just introduce the BSM particles and their interactions with the SM particles. In comparison with a specific model, such a phenomenological approach is less model-dependent and has been taken up in many previous works. As a matter of fact, the one-loop contributions to  $\Delta a_\mu$  in this approach have been systematically calculated in Ref. [90] and later verified in Ref. [91]. In these two papers, the final results of  $\Delta a_\mu$  have been given in terms of uncompleted integrals, since the Feynman parametrization has been used to deal with the one-loop calculations and the integration over the Feynman parameter is left intact. The approximate expressions of these one-loop contributions at the leading order of small particle-mass ratios calculated in the Feynman gauge are collected in Ref. [88], as well as in some earlier literature [92, 93].

Two shortcomings of the existing results in Refs. [90, 91] should be noticed. First, for a general interaction without assuming hierarchical particle masses, it is usually difficult to integrate the Feynman parameter out and derive the final analytical results. Second, even with some assumptions of particle masses, one must be extremely careful with the order of series expansions and the integration. For instance, the contribution to  $a_\mu$  from the Higgs boson in the SM is given by [90]

$$a_\mu^{\text{SM Higgs}} \propto \int_0^1 \frac{x^2 (2 - x)}{1 - x + x^2 (m_\mu^2/M_H^2)} dx , \quad (2)$$

where  $m_\mu \approx 106$  MeV is the muon mass and  $M_H \approx 125$  GeV is the Higgs-boson mass. Although  $m_\mu \ll M_H$  is perfectly satisfied, it is problematic to first expand the integrand in Eq. (2) as a series of the small mass ratio  $m_\mu/M_H$  and then integrate them out order by order. The reason is simply that the integral at each order is actually divergent, whereas the whole integral in Eq. (2) is finite as it should be.

In this work, we are thus motivated to revisit the one-loop contributions to  $a_\mu$  and derive the final analytical formulae. First, instead of the Feynman parametrization, we implement the Passarino-Veltman functions [94] and present the results in compact forms by using the publicly

available Package-X [95, 96]. Without any integrals in the final formulae, one can safely expand those general expressions as the series of the small mass ratios without any divergences, which should be helpful for phenomenological studies. Second, while there have been many attempts in the literature to find positive contributions to  $\Delta a_\mu$  (e.g., the two-Higgs-doublet model and the  $Z'$  model), some contributions emerge with a wrong sign (e.g., type-I and type-II seesaw models for neutrino masses). One immediate question is whether one can set up a simple criterion for positive contributions to  $\Delta a_\mu$  in an arbitrary new-physics model without repeating tedious loop calculations. Given all the interactions between the BSM particles and the SM particles, as well as the basic properties of those particles (e.g., masses, spins and electric charges), we find that with reasonable assumptions of particle masses it is indeed possible to make a quick judgment about the sign of  $\Delta a_\mu$ . For example, this goal can be achieved according to some simple criteria on the model parameters, such as the electric charges of the particles running in the loop and the relative sizes of the coupling constants.

The remaining part of the present paper is organized as follows. In Sec. 2, following the earlier works [90, 91], we classify the one-loop contributions to  $a_\mu$  by the particles running in the loop. More explicitly, there are three different categories: (1) a fermion and a scalar boson (i.e., the FS-type); (2) a fermion and a vector boson (i.e., the FV-type); and (3) a fermion, a scalar boson and a vector boson (i.e., the FSV-type). For each type of models, general expressions of  $a_\mu$  without any approximations are derived by using the Passarino-Veltman functions. Some simplifications of the formulae in various limits of hierarchical particle masses are discussed. The general formulae derived in Sec. 2 are then applied in Sec. 3 to some concrete models that have been extensively discussed in the literature. We also demonstrate how to make a quick judgment about the sign of  $\Delta a_\mu$  for a concrete model. Finally, our main conclusions are summarized in Sec. 4.

## 2 One-loop Contributions

Since we are interested in the extra one-loop contributions to  $a_\mu$  from the BSM particles, it is necessary to specify the particles running in the loop, given the external particles (i.e., the muon and the photon). Without resorting to a specific model, one can follow the phenomenological approach in Refs. [90, 91] and write down the interaction terms for the BSM and SM particles. As already mentioned above, there are three possible scenarios in which the basic symmetries, such as the Lorentz invariance and the  $U(1)$  gauge symmetry of quantum electrodynamics (QED), are required to be preserved.

### 2.1 The FS-type

First, we consider the case where there is a fermion  $F$  and a scalar boson  $S$  in the loops that contribute to  $a_\mu$ . The Feynman diagrams are shown in Fig. 1. These FS-type diagrams appear in many well-known models, such as the SM Higgs, the two-Higgs-doublet model, the type-II seesaw model, and the scalar leptoquark model.

After the spontaneous symmetry breaking of  $SU(2)_L \otimes U(1)_Y$  to  $U(1)_{\text{QED}}$ , the interactions




Figure 1: The Feynman diagrams for the FS-type contributions to the muon anomalous magnetic moment  $a_\mu$ , where the particle four-momenta have been explicitly specified and represented by arrows along the internal and external lines.

among the newly-introduced particles and the relevant SM particles can be written as

$$\mathcal{L}_{\text{int}}^{\text{FS}} = [\bar{F} (C_S + C_P \gamma_5) \mu S^\dagger + \text{h.c.}] + Q_F e \bar{F} \not{A} F + i Q_S e (S^\dagger \partial_\mu S - S \partial_\mu S^\dagger) A^\mu, \quad (3)$$

where  $C_S$  and  $C_P$  stand respectively for the scalar-type and pseudo-scalar-type coupling constants. The interaction between  $S$  and the photon is determined by the minimal coupling arising from the kinetic term of  $S$  via the replacement  $\partial_\mu \rightarrow \partial_\mu - i Q_S e A_\mu$  as in the ordinary QED, so is the interaction between the fermion  $F$  and the photon, where  $Q_S$  and  $Q_F$  denote the corresponding electric charges in units of the electric charge of positrons. Due to the conservation of electric charges, we always have  $Q_F + Q_S = -1$ , which will be implied in the subsequent discussions in this subsection.

Now it is straightforward to find out the amplitudes corresponding to the Feynman diagrams in Fig. 1(a) and Fig. 1(b), namely,

$$\begin{aligned} \bar{u}(p_2) \left( -ie \Gamma_{\text{FS}}^{\mu, (a)} \right) u(p_1) &= i Q_S e \int \frac{d^4 k}{(2\pi)^4} \left[ \bar{u}(p_2) i (C_S^* - C_P^* \gamma_5) \frac{i}{\not{k} - M_F} i (C_S + C_P \gamma_5) u(p_1) \right. \\ &\quad \left. \times \frac{i}{(p_2 - k)^2 - M_S^2} \frac{i}{(p_1 - k)^2 - M_S^2} (p_1 + p_2 - 2k)^\mu \right], \end{aligned} \quad (4)$$

and

$$\begin{aligned} \bar{u}(p_2) \left( -ie \Gamma_{\text{FS}}^{\mu, (b)} \right) u(p_1) &= i Q_F e \int \frac{d^4 k}{(2\pi)^4} \left[ \bar{u}(p_2) i (C_S^* - C_P^* \gamma_5) \frac{i}{\not{p}_2 - \not{k} - M_F} \gamma^\mu \right. \\ &\quad \left. \times \frac{i}{\not{p}_1 - \not{k} - M_F} i (C_S + C_P \gamma_5) u(p_1) \frac{i}{k^2 - M_S^2} \right]. \end{aligned} \quad (5)$$

From the overall amplitude, one can extract the FS-type contributions to the muon anomalous magnetic moment

$$a_\mu^{\text{FS}} = \frac{1}{16\pi^2} f_{\text{FS}} \left( \frac{m_\mu}{M_S}, \frac{M_F}{M_S}, C_S, C_P, Q_S \right), \quad (6)$$

where the exact relation  $Q_F = -1 - Q_S$  has been used and the loop function  $f_{\text{FS}}$  has been derived without making any approximations. For later convenience, we cast  $f_{\text{FS}}$  in the following form

$$f_{\text{FS}}(x, y, C_S, C_P, Q_S) = (|C_S|^2 - |C_P|^2) f_{\text{FS}}^-(x, y, Q_S) + (|C_S|^2 + |C_P|^2) f_{\text{FS}}^+(x, y, Q_S) , \quad (7)$$

where<sup>1</sup>

$$f_{\text{FS}}^-(x, y, Q_S) \equiv \frac{2y}{x} + \frac{2y \ln y}{x^3} [1 - y^2 + x^2(1 + Q_S)] + \frac{2y\Lambda(x^2, y, 1)}{x\lambda(x^2, y^2, 1)} \{x^4(1 + Q_S) + Q_S x^2 - [2 + x^2(2 + Q_S)]y^2 + y^4 + 1\} , \quad (8)$$

and

$$f_{\text{FS}}^+(x, y, Q_S) \equiv -2Q_S - \frac{1}{x^2} (2 + x^2 - 2y^2) - \frac{2 \ln y}{x^4} [1 - y^2(2 + x^2 - y^2) + Q_S x^2(1 - y^2)] + \frac{2\Lambda(x^2, y, 1)}{x^2 \lambda(x^2, y^2, 1)} \left\{ (y^2 - 1) [(y^2 - 1)^2 - x^2(1 - Q_S) - x^2 y^2(2 + Q_S)] + x^4 [Q_S + y^2(1 + Q_S)] \right\} , \quad (9)$$

with

$$\Lambda(a, b, c) \equiv \frac{\lambda^{1/2}(a, b^2, c^2)}{a} \ln \left( \frac{-a + b^2 + c^2 + \lambda^{1/2}(a, b^2, c^2)}{2bc} \right) \quad (10)$$

coming from the branch cut of the Passarino-Veltman  $B_0$  function and  $\lambda(a, b, c) \equiv a^2 + b^2 + c^2 - 2ab - 2ac - 2bc$  being the Källén kinematic function. Note that the terms in  $f_{\text{FS}}^-$  ( $f_{\text{FS}}^+$ ) are all proportional to the odd (even) powers of  $y$ , reflecting the anti-commutation relation between  $\gamma_5$  and  $\gamma^\mu$ . Some helpful comments on the loop function are in order.

- The general expression of  $f_{\text{FS}}$  in Eq. (7) has been written as the sum of two terms, one of which is proportional to  $(|C_S|^2 - |C_P|^2)$  and the other to  $(|C_S|^2 + |C_P|^2)$ . The former term corresponds to the chirality flip via the coupling between  $S$  and muon as well as the mass of  $F$ . This term would vanish if  $S$  is not coupled simultaneously to the left- and right-handed muon (i.e.,  $|C_S|^2 - |C_P|^2 = 0$ ) or if  $F$  is massless (i.e.,  $M_F = 0$ ). In contrast, the latter term corresponds to the chirality flip via the SM Yukawa coupling of muon, which is always nonzero since muon is massive. But it may be suppressed, in comparison with the former term, by the mass ratio  $m_\mu/M_F$ .
- For the electrically neutral  $F$  or  $S$ , only one diagram in Fig. 1(a) or Fig. 1(b) contributes to  $a_\mu$ . In this case, one can simply substitute  $Q_S = -1$  or  $Q_S = 0$  into Eqs. (6)-(9) to derive the final results.

The general formulae of the FS-type contribution to  $a_\mu$  in Eqs. (6)-(9) are exact, but it is difficult to get more information from these expressions. Therefore, we make some assumptions on the relevant particle masses and simplify the formulae. As will be shown below, some interesting results can be obtained.

---

<sup>1</sup>At first glance it seems that the first few terms with negative powers of  $x = m_\mu/M_S$  in  $f_{\text{FS}}^-$  and  $f_{\text{FS}}^+$  would be divergent as  $x$  approaches zero. However, this is not the case since these negative-power terms are exactly cancelled by the relevant terms in the  $\Lambda$  function and the final results exhibit only the terms with positive powers of  $x$ , which would vanish for  $m_\mu/M_S \rightarrow 0$ , as will be shown in Sec. 2.1.1 below.

### 2.1.1 $m_\mu \ll M_F, M_S$

The first scenario is to assume that both  $F$  and  $S$  are particles much heavier than muon, i.e.,  $m_\mu \ll M_F, M_S$ . Note that it is not necessary to specify the relative size of  $M_F$  and  $M_S$ , which can actually be comparable. In terms of the small mass ratio  $m_\mu/M_S$ , we can expand the functions  $f_{\text{FS}}^\pm$  in Eqs. (8) and (9) as follows

$$\begin{aligned} f_{\text{FS}} = & + (|C_S|^2 - |C_P|^2) \sum_{n=1}^{\infty} g_{\text{FS}}^{(2n-1)} \left( \frac{M_F}{M_S}, Q_S \right) \left( \frac{m_\mu}{M_S} \right)^{2n-1} \\ & + (|C_S|^2 + |C_P|^2) \sum_{n=1}^{\infty} g_{\text{FS}}^{(2n)} \left( \frac{M_F}{M_S}, Q_S \right) \left( \frac{m_\mu}{M_S} \right)^{2n}, \end{aligned} \quad (11)$$

where the terms of odd (even) powers come from  $f_{\text{FS}}^-$  ( $f_{\text{FS}}^+$ ). In light of  $m_\mu/M_S \ll 1$ , it is an excellent approximation to keep only the first two leading terms in Eq. (11), i.e., the first- and second-order loop functions


$$g_{\text{FS}}^{(1)}(y, Q_S) = \frac{y [y^4 (1 + 2Q_S) - 4y^2 (1 + Q_S + Q_S \ln y) + 4(1 + Q_S) \ln y + 3 + 2Q_S]}{(y^2 - 1)^3}, \quad (12)$$

and

$$\begin{aligned} g_{\text{FS}}^{(2)}(y, Q_S) = & \frac{1}{3(y^2 - 1)^4} \{ y^6 (1 + 3Q_S) - 3y^4 (2 + Q_S + 4Q_S \ln y) \\ & + 3y^2 [1 - Q_S + 4(1 + Q_S) \ln y] + 2 + 3Q_S \}. \end{aligned} \quad (13)$$

Based on the basic properties of the loop functions  $g_{\text{FS}}^{(n)}(y, Q_S)$  with  $n$  being positive integers, we can make a few interesting observations.

- First of all, one may wonder whether the loop functions  $g_{\text{FS}}^{(1)}(y, Q_S)$  and  $g_{\text{FS}}^{(2)}(y, Q_S)$  in Eqs. (12) and (13) become divergent at the point  $y = 1$ , i.e.,  $M_F = M_S$ . After a direct calculation, we verify that all the functions  $g_{\text{FS}}^{(n)}(y, Q_S)$  are actually finite and continuous at  $y = 1$ , implying the absence of a singularity at  $M_F = M_S$ .
- Second, all the loop functions  $g_{\text{FS}}^{(n)}(y, Q_S)$  tend to be zero as  $y$  is approaching infinity (i.e., in the limit  $M_F \gg M_S$ ). On the other hand, if  $y$  is approaching zero (i.e., in the limit  $M_F \ll M_S$ ), then we have  $g_{\text{FS}}^{(n)}(y, Q_S)$  with  $n$  being odd tends to be zero, as a consequence of the fact that these contributions come from the chirality flip via the mass of  $F$ . Moreover, the loop functions  $g_{\text{FS}}^{(n)}(y, Q_S)$  with  $n$  being even are finite in the limit of  $M_F \ll M_S$  or  $y = 0$ , and the finite value depends on  $n$  and  $Q_S$ .
- Third, we emphasize whether the loop functions  $g_{\text{FS}}^{(n)}(y, Q_S)$  at any values of  $y$  are positive or negative depends sensitively on the electric charge  $Q_S$ . In general, there exist both an upper bound  $Q_S^{\text{U}}$  and a lower bound  $Q_S^{\text{L}}$  for the electric charge  $Q_S$  such that  $g_{\text{FS}}^{(n)}(y, Q_S) \geq 0$  for  $Q_S \geq Q_S^{\text{U}}$  and  $g_{\text{FS}}^{(n)}(y, Q_S) \leq 0$  for  $Q_S \leq Q_S^{\text{L}}$  hold. However, for  $Q_S^{\text{L}} < Q_S < Q_S^{\text{U}}$ , the sign of the loop functions will be indefinite. The exact values of the corresponding  $Q_S^{\text{U}}$  and  $Q_S^{\text{L}}$  vary with  $n$ .



(a) The first-order loop function  $g_{\text{FS}}^{(1)}(y, Q_S)$  for  $Q_S = -1$  (red dashed line),  $Q_S = -2/3$  (gray solid line) and  $Q_S = -1/2$  (blue dash-dotted line). Note that for  $Q_S \geq -1/2$ ,  $g_{\text{FS}}^{(1)}(y, Q_S)$  is non-negative; for  $Q_S \leq -1$ ,  $g_{\text{FS}}^{(1)}(y, Q_S)$  is non-positive; for  $-1 < Q_S < -1/2$ , the sign of  $g_{\text{FS}}^{(1)}(y, Q_S)$  is indefinite.

(b) The second-order loop function  $g_{\text{FS}}^{(2)}(y, Q_S)$  for  $Q_S = -2/3$  (red dashed line),  $Q_S = -1/2$  (gray solid line) and  $Q_S = -1/3$  (blue dash-dotted line). Note that for  $Q_S \geq -1/3$ ,  $g_{\text{FS}}^{(2)}(y, Q_S)$  is non-negative; for  $Q_S \leq -2/3$ ,  $g_{\text{FS}}^{(2)}(y, Q_S)$  is non-positive; for  $-2/3 < Q_S < -1/3$ , the sign of  $g_{\text{FS}}^{(2)}(y, Q_S)$  is indefinite.

Figure 2: Numerical illustrations for the first- and second-order loop functions relevant for the FS-type contributions to  $a_\mu$  for different values of the electric charge  $Q_S$ .

Loop functions	$y \rightarrow 0$	$y \rightarrow \infty$	$y \rightarrow 1$	$Q_S^L$	$Q_S^U$
$g_{\text{FS}}^{(1)}(y, Q_S)$	0	0	$\frac{2}{3} + Q_S$	-1	$-\frac{1}{2}$
$g_{\text{FS}}^{(2)}(y, Q_S)$	$\frac{2}{3} + Q_S$	0	$\frac{1}{6}(1 + 2Q_S)$	$-\frac{2}{3}$	$-\frac{1}{3}$

Table 1: Summary of the asymptotic values of  $g_{\text{FS}}^{(1)}(y, Q_S)$  and  $g_{\text{FS}}^{(2)}(y, Q_S)$  in the limits of  $y \rightarrow 0$ ,  $y \rightarrow \infty$  and  $y \rightarrow 1$ , as well as the corresponding critical values  $Q_S^U$  and  $Q_S^L$ .

For illustration we focus on the first- and second-order loop functions  $g_{\text{FS}}^{(1)}(y, Q_S)$  and  $g_{\text{FS}}^{(2)}(y, Q_S)$  in Eqs. (12) and (13), and show them in Fig. 2 for different values of  $Q_S$ . The critical values  $Q_S^U = -1/2$  and  $Q_S^L = -1$  have been found for  $g_{\text{FS}}^{(1)}(y, Q_S)$ , while  $Q_S^U = -1/3$  and  $Q_S^L = -2/3$  for  $g_{\text{FS}}^{(2)}(y, Q_S)$ . In addition, the asymptotic values of  $g_{\text{FS}}^{(1)}(y, Q_S)$  and  $g_{\text{FS}}^{(2)}(y, Q_S)$  in the limits of  $y \rightarrow 0$ ,  $y \rightarrow \infty$  and  $y \rightarrow 1$ , together with the critical values  $Q_S^U$  and  $Q_S^L$ , have been summarized in Table 1. These observations are useful in judging whether the FS-type contribution to  $a_\mu$  is positive or negative. For instance, for the pure left- or right- handed Yukawa-type interactions, we have  $|C_S|^2 - |C_P|^2 = 0$  and thus the dominant contribution to  $a_\mu$  is given by  $g_{\text{FS}}^{(2)}(y, Q_S)$  in Eq. (11). In this case, one can conclude that  $\Delta a_\mu^{\text{FS}} > 0$  for  $Q_S \geq -1/3$  and  $\Delta a_\mu^{\text{FS}} < 0$  for  $Q_S \leq -2/3$ .

### 2.1.2 $m_\mu, M_F \ll M_S$

Then we proceed with another scenario where  $S$  is much heavier than  $F$  and muon, i.e.,  $m_\mu, M_F \ll M_S$ . To the order of  $\mathcal{O}(m_\mu^2/M_S^2)$ , one obtains

$$f_{\text{FS}} = \frac{1}{3} \left( \frac{m_\mu}{M_S} \right)^2 \left\{ (|C_S|^2 + |C_P|^2) (2 + 3Q_S) - 3(|C_S|^2 - |C_P|^2) \frac{M_F}{m_\mu} \right. \\ \left. \times \left[ 3 + 2Q_S + 4(1 + Q_S) \ln \left( \frac{M_F}{M_S} \right) \right] \right\}, \quad (14)$$

where the mass ratio  $M_F/m_\mu$  can take any positive values. In the limit of  $M_F/m_\mu \rightarrow 0$ , which is valid for  $F$  being neutrinos in the SM, the loop function in Eq. (14) is reduced to

$$f_{\text{FS}} = \frac{1}{3} \left( \frac{m_\mu}{M_S} \right)^2 (|C_S|^2 + |C_P|^2) (2 + 3Q_S), \quad (15)$$

where the term proportional to  $(|C_S|^2 - |C_P|^2)$  vanishes because of the fact that the chirality flip via the mass of  $F$  becomes impossible in the limit  $M_F/m_\mu \rightarrow 0$ . As indicated by Eq. (15), in this case, the contribution to  $a_\mu$  is positive only for  $Q_S > -2/3$ .

If the mass hierarchy  $m_\mu \ll M_F$  is further assumed, then Eq. (14) can be reduced to

$$f_{\text{FS}} = -\frac{m_\mu M_F}{M_S^2} (|C_S|^2 - |C_P|^2) \left[ 3 + 2Q_S + 4(1 + Q_S) \ln \left( \frac{M_F}{M_S} \right) \right]. \quad (16)$$

It is worthwhile to mention that the result in Eq. (16) can also be obtained by taking  $y \equiv M_F/M_S \ll 1$  in the function  $g_{\text{FS}}^{(1)}(y, Q_S)$  in Eq. (12).

### 2.1.3 $m_\mu, M_S \ll M_F$

Finally we consider the scenario where both muon and  $S$  are much lighter than  $F$ , i.e.,  $m_\mu, M_S \ll M_F$ . To the order of  $\mathcal{O}(m_\mu^2/M_F^2)$ , we have

$$f_{\text{FS}} = (|C_S|^2 - |C_P|^2) (1 + 2Q_S) \left( \frac{m_\mu}{M_F} \right) + \frac{1}{3} (|C_S|^2 + |C_P|^2) (1 + 3Q_S) \left( \frac{m_\mu}{M_F} \right)^2, \quad (17)$$

which can also be obtained by taking the limit of  $y = M_F/M_S \rightarrow \infty$  in Eqs. (12) and (13). Note that to the leading and next-to-leading orders of  $m_\mu/M_F$ , the contributions to  $a_\mu$  have nothing to do with the mass of  $S$ , as long as  $F$  is much heavier than muon and  $S$ . In particular, we shall have a finite result even for a massless  $S$ . This is very different from the FV-case where there is a vector boson instead of a scalar boson in the loop, as we will see in the next subsection, the loop function would be divergent as the mass of the vector boson were approaching zero.

## 2.2 The FV-type

Then we turn to the case where there is a fermion  $F$  and a vector boson  $V$  in the loops that contribute to  $a_\mu$ . The relevant Feynman diagrams are shown in Fig. 3. These FV-type diagrams actually appear in some realistic models, such as the  $W$ - and  $Z$ -boson contributions in the SM and the type-I seesaw model, the  $Z'$  model, and the vector leptoquark model.




Figure 3: The Feynman diagrams for the FV-type contributions to the muon anomalous magnetic moment  $a_\mu$ , where the particle four-momenta have been explicitly specified and represented by arrows along the internal and external lines.

After the spontaneous breaking of the  $SU(2)_L \otimes U(1)_Y$  gauge symmetry, the interactions among all the relevant particles can be written as

$$\begin{aligned} \mathcal{L}_{\text{int}}^{\text{FV}} = & \left[ \bar{F} (C_V \gamma^\mu + C_A \gamma^\mu \gamma_5) \mu V_\mu^\dagger + \text{h.c.} \right] + Q_F e \bar{F} \not{A} F + ie Q_V \left[ (\partial_\mu V_\nu^\dagger - \partial_\nu V_\mu^\dagger) A^\mu V^\nu \right. \\ & \left. - (\partial_\mu V_\nu - \partial_\nu V_\mu) A^\mu V^{\dagger\nu} + (V_\mu V_\nu^\dagger - V_\nu V_\mu^\dagger) \partial^\mu A^\nu \right], \end{aligned} \quad (18)$$

where  $C_V$  and  $C_A$  stand respectively for the vector-type and axial-vector-type coupling constants. The interaction between  $V$  (or  $F$ ) and photon is determined by the minimal coupling as in the ordinary QED. The last two terms  $(V_\mu V_\nu^\dagger - V_\nu V_\mu^\dagger) \partial^\mu A^\nu$  in Eq. (18) have been added to ensure the renormalizability of the theory [97]. In addition, the conservation of electric charges requires the relation  $Q_F + Q_V = -1$  to hold.

The amplitudes corresponding to the Feynman diagrams in Fig. 3(a) and Fig. 3(b) can be explicitly written down as below<sup>2</sup>

$$\begin{aligned} \bar{u}(p_2) \left( -ie \Gamma_{\text{FV}}^{\mu, (a)} \right) u(p_1) = & i Q_V e \int \frac{d^4 k}{(2\pi)^4} \left\{ \bar{u}(p_2) i (C_V^* \gamma^\sigma + C_A^* \gamma^\sigma \gamma_5) \frac{i}{\not{k} - M_F} i (C_V \gamma^\rho \gamma_5 + C_A \gamma^\rho \gamma_5) \right. \\ & \times u(p_1) \frac{-i}{(p_1 - k)^2 - M_V^2} \frac{-i}{(p_2 - k)^2 - M_V^2} \left[ \eta_{\rho\alpha} - \frac{(p_1 - k)_\rho (p_1 - k)_\alpha}{M_V^2} \right] \\ & \times \left[ \eta_{\sigma\beta} - \frac{(p_2 - k)_\sigma (p_2 - k)_\beta}{M_V^2} \right] \left[ \eta^{\beta\alpha} (2k - p_1 - p_2)^\mu \right. \\ & \left. \left. + \eta^{\alpha\mu} (2p_1 - p_2 - k)^\beta + \eta^{\mu\beta} (2p_2 - k - p_1)^\alpha \right] \right\}, \end{aligned} \quad (19)$$

<sup>2</sup>Throughout this paper, whenever the vector boson  $V$  is involved, we perform all the calculations in the unitary gauge so that  $V$  can be either a gauge boson or a vector-like matter particle.

and

$$\begin{aligned} \bar{u}(p_2) \left( -ie\Gamma_{\text{FV}}^{\mu,(\text{b})} \right) u(p_1) &= iQ_F e \int \frac{d^4 k}{(2\pi)^4} \left[ \bar{u}(p_2) i (C_V^* \gamma^\sigma + C_A^* \gamma^\sigma \gamma_5) \frac{i}{\not{p}_2 - \not{k} - M_F} \gamma^\mu \right. \\ &\quad \left. \times \frac{i}{\not{p}_1 - \not{k} - M_F} i (C_V \gamma^\rho + C_A \gamma^\rho \gamma_5) u(p_1) \frac{-i}{k^2 - M_V^2} \left( \eta_{\rho\sigma} - \frac{k_\rho k_\sigma}{M_V^2} \right) \right], \end{aligned} \quad (20)$$

where  $\eta_{\mu\nu} = \text{diag}(+, -, -, -)$  is the metric tensor of the Minkowski space-time. From Eqs. (19) and (20) one can extract the FV-type contributions to  $a_\mu$ , i.e.,

$$a_\mu^{\text{FV}} = \frac{1}{16\pi^2} f_{\text{FV}} \left( \frac{m_\mu}{M_V}, \frac{M_F}{M_V}, C_V, C_A, Q_V \right), \quad (21)$$

where the exact relation  $Q_F = -1 - Q_V$  has been used and the loop function  $f_{\text{FV}}$  has been derived without making any approximations. Similar to  $f_{\text{FS}}$  in the previous subsection,  $f_{\text{FV}}$  can also be divided into two parts

$$f_{\text{FV}}(x, y, C_V, C_A, Q_V) = (|C_V|^2 - |C_A|^2) f_{\text{FV}}^-(x, y, Q_V) + (|C_V|^2 + |C_A|^2) f_{\text{FV}}^+(x, y, Q_V), \quad (22)$$

where

$$\begin{aligned} f_{\text{FV}}^-(x, y, Q_V) &= 4Q_V xy - \frac{2y}{x} (2 + y^2 - 2x^2) - \frac{2y \ln y}{x^3} \{ 2 + x^2 (2Q_V - 1) - x^4 (1 + Q_V) \\ &\quad + y^2 [(2 + Q_V)x^2 - 1] - y^4 \} - \frac{2y\Lambda(x^2, y, 1)}{x\lambda(x^2, y^2, 1)} \{ 2 - x^2 [3 + x^4 + Q_V (-2 + 3x^2 \\ &\quad + x^4)] + y^2 [-3 - Q_V x^2 + x^4 (3 + 2Q_V)] - x^2 y^4 (3 + Q_V) + y^6 \}, \end{aligned} \quad (23)$$

and

$$\begin{aligned} f_{\text{FV}}^+(x, y, Q_V) &= -2Q_V (2 + x^2 + y^2) - \frac{1}{x^2} \left[ (x^2 - 2)^2 + 3x^2 y^2 - 2y^2 (1 + y^2) \right] - \frac{2 \ln y}{x^4} \\ &\quad \times \{ 2 + x^2 (2Q_V - 3 - 3Q_V x^2) + y^2 [-3 + x^2 (2 - Q_V) + x^4 (1 + Q_V)] \\ &\quad - x^2 y^4 (2 + Q_V) + y^6 \} - \frac{2\Lambda(x^2, y, 1)}{x^2 \lambda(x^2, y^2, 1)} \{ (1 + Q_V x^2) (2 - 5x^2 + 3x^4) \\ &\quad + y^2 [-5 + x^2 (4 + x^2 + x^4 + Q_V (-3 + x^2 + x^4))] \\ &\quad + y^4 [3 - 2x^2 - x^4 (3 + 2Q_V)] + y^6 [1 + x^2 (3 + Q_V)] - y^8 \}. \end{aligned} \quad (24)$$

Although the expressions of the loop functions in Eqs. (23) and (24) seem to be lengthy, the integration has been explicitly carried out and the results are exact in the sense that no approximations have been made. As in the case of FS-type contributions, the anti-commutation relation between  $\gamma_5$  and  $\gamma^\mu$  enforces the terms in  $f_{\text{FV}}^-$  (or  $f_{\text{FV}}^+$ ) to be proportional to the odd (or even) powers of  $y$ . Some comments on the loop functions are in order.

- The loop function  $f_{\text{FV}}^-$  corresponds to the chirality flips via the coupling between  $V$  and muon or via the mass of  $F$ , which would vanish if  $V$  were not coupling simultaneously to the left- and right- handed muon (i.e.,  $|C_V|^2 - |C_A|^2 = 0$ ) or if  $F$  were massless (i.e.,  $M_F = 0$ ). In contrast, the other loop function  $f_{\text{FV}}^+$  corresponds to the chirality flips via the SM Yukawa coupling of muon, which is always nonzero because of non-vanishing muon mass. But it may be suppressed, when compared to  $f_{\text{FV}}^-$ , by the mass ratio  $m_\mu/M_F$ .

- If  $F$  or  $V$  is electrically neutral, then only the diagram in Fig. 3(a) or Fig. 3(b) contributes to  $a_\mu$ . In this case, one can simply set  $Q_V = -1$  or  $Q_V = 0$  in Eqs. (21)-(24) to derive the final results.

In the remaining part of this subsection we shall impose some hierarchical conditions on the masses of the relevant particles to simplify the general expressions of Eqs. (23) and (24).

### 2.2.1 $m_\mu \ll M_F, M_V$

Suppose that both  $F$  and  $V$  are much heavier than muon, i.e.,  $m_\mu \ll M_F, M_V$ , whereas the relative sizes of  $M_F$  and  $M_V$  are not specified. Expanding the general functions  $f_{FV}^\mp$  in Eqs. (23) and (24) as the series of  $m_\mu/M_V$ , one obtains

$$\begin{aligned} f_{FV} = & + (|C_V|^2 - |C_A|^2) \sum_{n=1}^{\infty} g_{FV}^{(2n-1)} \left( \frac{M_F}{M_V}, Q_V \right) \left( \frac{m_\mu}{M_V} \right)^{2n-1} \\ & + (|C_V|^2 + |C_A|^2) \sum_{n=1}^{\infty} g_{FV}^{(2n)} \left( \frac{M_F}{M_V}, Q_V \right) \left( \frac{m_\mu}{M_V} \right)^{2n}, \end{aligned} \quad (25)$$

where the odd- and even-power terms come from  $f_{FV}^-$  and  $f_{FV}^+$ , respectively. In the assumption of  $m_\mu \ll M_V$ , it is a good approximation to keep the first two leading terms in Eq. (25), i.e., the first- and second-order loop functions


$$\begin{aligned} g_{FV}^{(1)}(y, Q_V) = & \frac{y}{(y^2 - 1)^3} \{ y^6 (1 + 2Q_V) - 12y^4 Q_V (1 - \ln y) + 3y^2 [1 + 6Q_V - 4(1 + Q_V) \ln y] \\ & - 4(1 + 2Q_V) \} , \end{aligned} \quad (26)$$

and

$$\begin{aligned} g_{FV}^{(2)}(y, Q_V) = & \frac{-1}{3(y^2 - 1)^4} \{ y^8 (5 + 9Q_V) - y^6 [7(2 + 9Q_V) - 36Q_V \ln y] \\ & + y^4 [39 + 117Q_V - 36(1 + Q_V) \ln y] - y^2 (38 + 81Q_V) + 2(4 + 9Q_V) \} . \end{aligned} \quad (27)$$

Some interesting observations on the basic properties of the loop functions  $g_{FV}^{(n)}(y, Q_V)$  (with  $n$  being positive integers) can be made.

- All the loop functions  $g_{FV}^{(n)}(y, Q_V)$  are finite and continuous at  $y = 1$ , implying the absence of a singularity at  $M_V = M_F$ . As  $y$  is approaching zero (i.e., in the limit  $M_F \ll M_V$ ),  $g_{FV}^{(n)}(y, Q_V)$  with  $n$  being odd tends to be zero, reflecting the fact that these contributions arise from the chirality flip via the mass of  $F$ , while  $g_{FV}^{(n)}(y, Q_V)$  with  $n$  being even has a finite limit that depends on both  $n$  and  $Q_V$ . When  $y$  is approaching infinity (i.e., in the limit  $M_F \gg M_V$ ),  $g_{FV}^{(n)}(y, Q_V)$  with  $n > 2$  becomes zero. However, in the same limit, we have  $g_{FV}^{(2)}(y, Q_V) \rightarrow -5/3 - 3Q_V$ , while  $g_{FV}^{(1)}(y, Q_V) \rightarrow \infty$  for  $Q_V \neq -1/2$  and  $g_{FV}^{(1)}(y, Q_V) \rightarrow 0$  for  $Q_V = -1/2$ .
- There exist both an upper bound  $Q_V^U$  and a lower bound  $Q_V^L$  for the electric charge  $Q_V$  such that  $g_{FV}^{(n)}(y, Q_V) \geq 0$  ( $\leq 0$ ) for  $Q_V \geq Q_V^U$  with  $n$  being odd (even) and  $g_{FV}^{(n)}(y, Q_V) \leq 0$  ( $\geq 0$ ) for  $Q_V \leq Q_V^L$  with  $n$  being odd (even). However, for  $Q_V^L < Q_V < Q_V^U$ , the sign of the loop



(a) The first-order loop function  $g_{\text{FV}}^{(1)}(y, Q_V)$  for  $Q_V = -1/2$  (red dashed line),  $Q_V = -2/5$  (gray solid line) and  $Q_V = -1/3$  (blue dash-dotted line). Note that for  $Q_V \geq -0.387$ ,  $g_{\text{FV}}^{(1)}(y, Q_V)$  is non-negative; for  $Q_V \leq -1/2$ ,  $g_{\text{FV}}^{(1)}(y, Q_V)$  is non-positive; for  $-1/2 < Q_V < -0.387$ , the sign of  $g_{\text{FV}}^{(1)}(y, Q_V)$  is indefinite.

(b) The second-order loop function  $g_{\text{FV}}^{(2)}(y, Q_V)$  for  $Q_V = -5/9$  (red dashed line),  $Q_V = -1/2$  (gray solid line) and  $Q_V = -1/3$  (blue dash-dotted line). Note that for  $Q_V \geq -0.433$ ,  $g_{\text{FV}}^{(2)}(y, Q_V)$  is non-positive; for  $Q_V \leq -5/9$ ,  $g_{\text{FV}}^{(2)}(y, Q_V)$  is non-negative; for  $-5/9 < Q_V < -0.433$ , the sign of  $g_{\text{FV}}^{(2)}(y, Q_V)$  is indefinite.

Figure 4: Numerical illustrations for the first- and second-order loop functions relevant for the FV-type contributions to  $a_\mu$  for different values of the electric charge  $Q_V$ .

functions will be indefinite. The exact values of the corresponding  $Q_V^{\text{U}}$  and  $Q_V^{\text{L}}$  vary with  $n$ . For a better illustration we have shown the first- and second-order loop functions  $g_{\text{FV}}^{(1)}(y, Q_V)$  and  $g_{\text{FV}}^{(2)}(y, Q_V)$  in Fig. 4 for different values of  $Q_V$ . The asymptotic behaviors together with the critical values  $Q_V^{\text{U}}$  and  $Q_V^{\text{L}}$  are also summarized in Table 2. This observation is useful in judging the sign of the FV-type contribution to  $a_\mu$ . For instance, for the pure left- or right-handed interaction, we have  $|C_V|^2 - |C_A|^2 = 0$  and thus the dominant contribution to  $a_\mu$  is given by  $g_{\text{FV}}^{(2)}(y, Q_V)$  in Eq. (25). In this case, one can conclude that  $\Delta a_\mu^{\text{FV}} > 0$  for  $Q_V \leq -5/9$  and  $\Delta a_\mu^{\text{FV}} < 0$  for  $Q_V \geq -0.433$ .

The loop functions of the FV-type are quite different from those of the FS-type. In particular, the divergence in the first-order loop function  $g_{\text{FV}}^{(1)}(y, Q_V)$  as  $M_F/M_V \rightarrow \infty$  stems from the fact that the power of  $y$  in the numerator of  $g_{\text{FV}}^{(1)}(y, Q_V)$  is higher than that in its denominator, while the power of  $y$  in the numerator of  $g_{\text{FS}}^{(1)}(y, Q_S)$  is lower than that in the denominator.<sup>3</sup> However, as a physical observable, the muon anomalous magnetic moment should not blow up as the mass of  $F$  approaches infinity. The only way for  $a_\mu$  to be finite in the limit of  $M_F \rightarrow \infty$  is to demand that the couplings between muon and  $F$  should be further suppressed by the inverse powers of  $M_F$ . Therefore, in the theory with a heavy fermion coupled to muon and a vector boson, we can draw the following model-independent constraint: *If the heavy fermion is coupled simultaneously to both left- and right-handed muon and the vector boson, then the couplings should decrease at*

<sup>3</sup>If one calculates the Feynman amplitudes in Fig. 3 in the Feynman gauge, then the highest power of  $y$  in the numerator will be reduced by two and there is no divergence in  $g_{\text{FV}}^{(1)}(y, Q_V)$  [88]. However, there will be contributions from extra diagrams corresponding to the couplings among muon,  $F$  and Goldstone bosons, leading to divergences as  $M_F \rightarrow \infty$ .

Loop functions	$y \rightarrow 0$	$y \rightarrow \infty$	$y \rightarrow 1$	$Q_V^L$	$Q_V^U$
$g_{\text{FV}}^{(1)}(y, Q_S)$	0	$\infty$ (for $Q_V \neq -1/2$ ) 0 (for $Q_V = -1/2$ )	$2 + 5Q_V$	$-\frac{1}{2}$	-0.387
$g_{\text{FV}}^{(2)}(y, Q_S)$	$-\frac{8}{3} - 6Q_V$	$-\frac{5}{3} - 3Q_V$	$-\frac{13}{6} - 5Q_V$	$-\frac{5}{9}$	-0.433

Table 2: Summary of the asymptotic values of  $g_{\text{FV}}^{(1)}(y, Q_V)$  and  $g_{\text{FV}}^{(2)}(y, Q_V)$  in the limit of  $y \rightarrow 0$ ,  $y \rightarrow \infty$  and  $y \rightarrow 1$ , as well as the corresponding critical values  $Q_V^U$  and  $Q_V^L$ . Note that the upper-critical charges  $Q_V^U$  are irrational numbers and thus three significant digits are maintained.

least as  $\mathcal{O}(1/\sqrt{M_F})$  in the limit of  $M_F \rightarrow \infty$ . For an ultraviolet-complete model, this is indeed satisfied since the couplings are usually proportional to the mixing parameters between muon and  $F$ , which are suppressed by the heavy fermion mass.

### 2.2.2 $m_\mu, M_F \ll M_V$

Then we assume that both muon and  $F$  are much lighter than  $V$ , i.e.,  $m_\mu, M_F \ll M_V$ . To the order of  $\mathcal{O}(m_\mu^2/M_V^2)$  one obtains

$$f_{\text{FV}} = -\frac{2}{3} \left( \frac{m_\mu}{M_V} \right)^2 \left[ (|C_V|^2 + |C_A|^2) (4 + 9Q_V) - 6 (|C_V|^2 - |C_A|^2) (1 + 2Q_V) \frac{M_F}{m_\mu} \right], \quad (28)$$

where the mass ratio  $M_F/m_\mu$  can take any positive values. In the limit of  $M_F/m_\mu \rightarrow 0$ , which is valid for  $F$  being neutrinos in the SM, the loop function in Eq. (28) is reduced to

$$f_{\text{FV}} = -\frac{2}{3} \left( \frac{m_\mu}{M_V} \right)^2 (|C_V|^2 + |C_A|^2) (4 + 9Q_V), \quad (29)$$

where the term proportional to  $(|C_V|^2 - |C_A|^2)$  vanishes. As indicated by Eq. (29), in this case, the contribution to  $a_\mu$  is positive only for  $Q_V < -4/9$ .

Moreover, if  $m_\mu \ll M_F$  holds, then Eq. (28) can be greatly simplified to

$$f_{\text{FV}} = 4 \frac{m_\mu M_F}{M_V^2} (|C_V|^2 - |C_A|^2) (1 + 2Q_V), \quad (30)$$

which can also be obtained by setting  $y \equiv M_F/M_V \ll 1$  in the function  $g_{\text{FV}}^{(1)}(y, Q_V)$  in Eq. (26).

### 2.2.3 $m_\mu, M_V \ll M_F$

Finally we consider the scenario where both muon and  $V$  are much lighter than  $F$ , i.e.,  $m_\mu, M_V \ll M_F$ . To the order of  $\mathcal{O}(m_\mu^2/M_F^2)$ , we have

$$f_{\text{FV}} = (|C_V|^2 - |C_A|^2) (1 + 2Q_V) \frac{m_\mu M_F}{M_V^2} - \frac{1}{3} (|C_V|^2 + |C_A|^2) (5 + 9Q_V) \left( \frac{m_\mu}{M_V} \right)^2, \quad (31)$$




Figure 5: The Feynman diagrams for the FSV-type contributions to the muon anomalous magnetic moment  $a_\mu$ , where the particle four-momenta have been explicitly specified and represented by arrows along the internal and external lines.

which can also be obtained by taking the limit of  $y = M_F/M_V \rightarrow \infty$  in Eqs. (26) and (27). Two interesting observations can be made.

First, as has been emphasized in the last paragraph of Sec. 2.2.1, for the theory with a heavy fermion coupled simultaneously to both left- and right-handed muon (i.e.,  $|C_V|^2 - |C_A|^2 \neq 0$ ), the first-order loop function, which scales as  $\mathcal{O}(M_F m_\mu/M_V^2)$ , will be divergent for  $M_F \rightarrow \infty$ . Hence a finite contribution to  $a_\mu$  requires  $C_V$  and  $C_A$  to decrease no more slowly than  $\mathcal{O}(1/\sqrt{M_F})$ .

Second, the right-hand side of Eq. (31) becomes singular as  $M_V$  approaches zero. In order to track this singularity one can take  $M_V = 0$  from the very beginning in Eqs. (19) and (20).<sup>4</sup> After doing so, one can find that the contribution from Fig. 3(b) is finite while that from Fig. 3(a) is infinite. This result can be understood by recalling the famous Weinberg-Witten theorem [98]: *For the theory with a Lorentz-covariant conserved current, massless particles of spin  $j > 1/2$  with a nonvanishing charge, corresponding to the conserved current, cannot exist.* In our case, the particles running in the loop are coupled to the external photon via the electromagnetic interaction, and the corresponding electromagnetic current is of course Lorentz-covariant and conserved. The Weinberg-Witten theorem excludes the existence of a massless vector boson with a nonvanishing electric charge. In other words, the vector boson in the Feynman diagram in Fig. 3(a) cannot be massless, implying the absence of the aforementioned singularity.

## 2.3 The FSV-type

In this subsection, we study the scenario where there is a fermion  $F$ , a scalar  $S$  and a vector boson  $V$  in the loops that contribute to  $a_\mu$ . The relevant Feynman diagrams are shown in Fig. (5). These FSV-type diagrams usually appear in the calculations in the  $R_\xi$  gauge, for which there are extra contributions from Goldstone bosons. However, since we perform all the calculation in the unitary

<sup>4</sup>Note that when taking  $M_V = 0$ , we have to modify the propagator of the vector boson in Eq. (20) to  $-i\eta_{\rho\sigma}/k^2$ .

gauge, there are no such diagrams in the SM. In the extensions of the SM, such diagrams indeed exist and  $F$ ,  $S$  and  $V$  can all be identified with physical particles. A typical example is the type-II seesaw model, which will be discussed in Sec. 3.3.

As in the previous cases, we write down the general Lagrangian for the relevant particles contributing to  $a_\mu$ , i.e.,<sup>5</sup>

$$\mathcal{L}_{\text{int}}^{\text{FSV}} = [\overline{F}(C_S + C_P\gamma_5)\mu S^\dagger + \overline{F}(C_V\gamma^\mu + C_A\gamma^\mu\gamma_5)\mu V_\mu^\dagger + \text{h.c.}] + \mu_0 e A^\mu (V_\mu^\dagger S + V_\mu S^\dagger) , \quad (32)$$

where  $\mu_0$  stands for a free parameter of mass dimension and will be fixed in a concrete model.

The amplitudes corresponding to the Feynman diagrams in Fig. 5(a) and Fig. 5(b) are given by

$$\begin{aligned} \overline{u}(p_2) \left( -ie\Gamma_{\text{FSV}}^{\mu,(a)} \right) u(p_1) &= i\mu_0 e \int \frac{d^4k}{(2\pi)^4} \left\{ \overline{u}(p_2) i (C_V^* \gamma^\alpha + C_A^* \gamma^\alpha \gamma_5) \frac{i}{\not{k} - M_F} \right. \\ &\quad \times (C_S + C_P \gamma_5) u(p_1) \frac{i}{(p_1 - k)^2 - M_S^2} \frac{-i\eta^{\mu\beta}}{(p_2 - k)^2 - M_V^2} \\ &\quad \left. \times \left[ \eta_{\alpha\beta} - \frac{(p_2 - k)_\alpha (p_2 - k)_\beta}{M_V^2} \right] \right\} , \end{aligned} \quad (33)$$

and

$$\begin{aligned} \overline{u}(p_2) \left( -ie\Gamma_{\text{FSV}}^{\mu,(b)} \right) u(p_1) &= i\mu_0 e \int \frac{d^4k}{(2\pi)^4} \left\{ \overline{u}(p_2) i (C_S^* - C_P^* \gamma_5) \frac{i}{\not{k} - M_F} \right. \\ &\quad \times (C_V \gamma^\alpha + C_A \gamma^\alpha \gamma_5) u(p_1) \frac{i}{(p_2 - k)^2 - M_S^2} \frac{-i\eta^{\mu\beta}}{(p_1 - k)^2 - M_V^2} \\ &\quad \left. \times \left[ \eta_{\alpha\beta} - \frac{(p_1 - k)_\alpha (p_1 - k)_\beta}{M_V^2} \right] \right\} . \end{aligned} \quad (34)$$

From these amplitudes, one can extract the FSV-type contributions to  $a_\mu$  as follows

$$a_\mu^{\text{FSV}} = \frac{1}{16\pi^2} \left( \frac{\mu_0}{M_V} \right) f_{\text{FSV}} \left( \frac{m_\mu}{M_V}, \frac{m_\mu}{M_S}, \frac{m_\mu}{M_F}, C_S, C_P, C_V, C_A \right) , \quad (35)$$

where the loop function can be decomposed into two parts<sup>6</sup>

$$\begin{aligned} f_{\text{FSV}}(x, y, z, C_S, C_P, C_V, C_A) &= + [\text{Re}(C_A^* C_P) - \text{Re}(C_S^* C_V)] f_{\text{FSV}}^-(x, y, z) \\ &\quad + [\text{Re}(C_A^* C_P) + \text{Re}(C_S^* C_V)] f_{\text{FSV}}^+(x, y, z) , \end{aligned} \quad (36)$$

---

<sup>5</sup>Here we only include the renormalizable interactions. The cases of the axion and axion-like particles [14, 18, 20, 26, 42, 99–103], which contribute to the muon  $g-2$  via the dimension-five non-renormalizable effective operator, are not included.

<sup>6</sup>In the special case, where both  $S$  and  $V$  are only coupled to either left-handed or right-handed muon, then the first part vanishes. If  $S$  is only coupled to left-handed (or right-handed) muon and  $V$  to right-handed (or left-handed) muon, then the second part vanishes.

with

$$\begin{aligned}
f_{\text{FSV}}^-(x, y, z) = & \frac{1}{3x^3y^4z^6(x^2-y^2)^2} \left\{ -2x^2y^2z^6(x^2-y^2)[x^4+y^4+x^2y^2(2y^2-x^2-1)] \right. \\
& + x^4y^4z^4(x^4-y^4) + y^6 \ln\left(\frac{x}{y}\right) [2y^2z^6 + x^2z^4(3(3x^2-y^2)(1+z^2) - 4z^2) \\
& + x^6(1-z^2)((x^2+y^2)(1-z^2)^2 - 6z^2(1+z^2))] + z^4(x^2-y^2)^3 \ln\left(\frac{y}{z}\right) \\
& \times [-2(x^2+y^2)z^2 + 3x^2y^2(1+z^2)] + x^6y^2z^2\Lambda\left(1, \frac{1}{y}, \frac{1}{z}\right) [2z^4(2y^2-x^2) \\
& + y^4(x^2+y^2)(1-z^2)^2 + y^2z^2(x^2-5y^2)(1+z^2)] - y^6x^2z^2\Lambda\left(1, \frac{1}{x}, \frac{1}{z}\right) \\
& \left. \times [2z^4(2x^2-y^2) + x^4(y^2+x^2)(1-z^2)^2 + x^2z^2(y^2-5x^2)(1+z^2)] \right\}, \quad (37)
\end{aligned}$$

and

$$\begin{aligned}
f_{\text{FSV}}^+(x, y, z) = & \frac{1}{3x^3y^4z^7(x^2-y^2)} \left\{ -x^2y^2z^4(x^2-y^2)[2y^2(z^2-2x^2) + x^2z^2(2+y^2)] \right. \\
& + 2y^6 \ln\left(\frac{x}{y}\right) [z^4(z^2-3x^2) + x^4(1-z^2)(3z^2-x^2(1-z^2)^2)] - 2z^2 \\
& \times (x^2-y^2) \ln\left(\frac{y}{z}\right) [3x^2y^2(x^2y^2(1-z^2) - z^2(x^2+y^2)) + z^4(x^4+x^2y^2+y^4)] \\
& + 2x^2y^6z^2\Lambda\left(1, \frac{1}{x}, \frac{1}{z}\right) [z^2(z^2-x^2) - x^2(1-z^2)(z^2-x^2+x^2z^2)] \\
& \left. - 2y^2x^6z^2\Lambda\left(1, \frac{1}{y}, \frac{1}{z}\right) [z^2(z^2-y^2) - y^2(1-z^2)(z^2-y^2+y^2z^2)] \right\}. \quad (38)
\end{aligned}$$

Notice that the last term in  $f_{\text{FSV}}^-(f_{\text{FSV}}^+)$  can be obtained by simply exchanging between  $x$  and  $y$  in the next-to-last term in  $f_{\text{FSV}}^-(f_{\text{FSV}}^+)$  up to a minus sign.

Although the general expressions in Eqs. (37) and (38) are more complicated than those in the previous cases, we have included these exact results for completeness. In the SM extensions, these formulae may find some practical applications. In fact, the explicit expressions can be greatly simplified under the conditions of strong mass hierarchies. For instance, we assume that muon is much lighter than  $V$  and  $S$ , i.e.,  $m_\mu \ll M_V, M_S$ . Hence, to the order of  $\mathcal{O}(m_\mu^3/M_V^3)$ , one has

$$\begin{aligned}
f_{\text{FSV}} = & + \left\{ [\text{Re}(C_A^* C_P) - \text{Re}(C_S^* C_V)] g_{\text{FSV}}^{(1)}\left(\frac{M_V}{M_S}\right) \right\} \left(\frac{m_\mu}{M_V}\right) \\
& + \left\{ [\text{Re}(C_A^* C_P) - \text{Re}(C_S^* C_V)] g_{\text{FSV}}^{(3-)}\left(\frac{M_V}{M_S}, \frac{M_F}{m_\mu}\right) \right. \\
& \left. + [\text{Re}(C_A^* C_P) + \text{Re}(C_S^* C_V)] g_{\text{FSV}}^{(3+)}\left(\frac{M_V}{M_S}, \frac{M_F}{m_\mu}\right) \right\} \left(\frac{m_\mu}{M_V}\right)^3, \quad (39)
\end{aligned}$$

where

$$g_{\text{FSV}}^{(1)}(x) = \frac{-2x^2}{(x^2-1)^2} (1-x^2+2\ln x), \quad (40)$$

$$g_{\text{FSV}}^{(3-)}(x, y) = \frac{-2x^2}{3(x^2-1)^2} [(1+x^2)\ln x + 1-x^2] (3y^2-1), \quad (41)$$

$$g_{\text{FSV}}^{(3+)}(x, y) = \frac{4x^2y}{3(x^2-1)} \ln x. \quad (42)$$

Note that the loop functions in Eqs. (40)-(42) are all continuous at  $x = 1$ . More explicitly, we have the finite limits

$$\lim_{x \rightarrow 1} g_{\text{FSV}}^{(1)}(x) = 1, \quad \lim_{x \rightarrow 1} g_{\text{FSV}}^{(3-)}(x, y) = 0, \quad \lim_{x \rightarrow 1} g_{\text{FSV}}^{(3+)}(x, y) = \frac{2y}{3},$$

indicating the absence of any singularities at  $M_V = M_S$ . In general, it is impossible to reach a simple conclusion on the sign of the loop functions or that of  $a_\mu^{\text{FSV}}$ . The situation is even worse for the exact loop functions in Eqs. (37) and (38). However, if  $\text{Re}(C_A^* C_P) = \text{Re}(C_S^* C_V)$  holds, then we can see from Eq. (39) that only the loop function  $g_{\text{FSV}}^{(3+)}(M_V/M_S, M_F/m_\mu)$  is involved. In this special case, one can immediately verify that  $g_{\text{FSV}}^{(3+)}(M_V/M_S, M_F/m_\mu)$  in Eq. (42) will always be positive for arbitrary mass ratios  $M_V/M_S$  and  $M_F/m_\mu$ . Thus the sign of  $a_\mu$  is just determined by the sign of  $\mu_0 [\text{Re}(C_A^* C_P) + \text{Re}(C_S^* C_V)]$ .

### 3 Practical Applications

In this section, we take some concrete models, which have been extensively discussed in the literature, as examples to illustrate how our general formulae derived in Sec. 2 can be practically applied to the calculations of muon  $g - 2$ . On the other hand, the latest measurement of the fine structure constant from the recoil velocity of rubidium atom brings about a  $1.6 \sigma$  discrepancy<sup>7</sup> with the theoretical prediction for the electron anomalous magnetic moment [104]. We emphasize that our general formulae derived in Sec. 2 can be directly used to calculate the one-loop contributions to electron  $g - 2$  by only replacing the mass and couplings of muon with those of electron. However, in the remaining part of this section, we shall focus on the one-loop contributions to muon anomalous magnetic moment for clarity.

In the subsequent discussions, we first apply our general formulae to the SM and rederive the well-known one-loop QED and electroweak (EW) contributions to muon  $g - 2$ . Then, we investigate several new-physics models, including the type-I seesaw model, the type-II seesaw model, the  $Z'$  model and the scalar leptoquark model. Special attention will be paid to a quick judgment about the sign of  $\Delta a_\mu$  in concrete models.

#### 3.1 SM contributions

Since Julian Schwinger first calculated the one-loop QED correction to the anomalous magnetic moment of electron in 1948 [106], great efforts have been made to the precise calculation of higher-order corrections to the anomalous magnetic moment of charged leptons. See, for example, Ref. [3], for a recent review. The one-loop EW corrections to muon  $g - 2$  were calculated more than fifty years ago [107–113]. In this subsection, we reproduce the well-known results of one-loop QED and EW contributions to  $a_\mu$  by using our general formulae derived in the last section.

- *QED.* The one-loop QED contribution to the muon anomalous magnetic moment belongs to the FV-type, corresponding to Fig. 6(a). Setting  $Q_V = 0$ ,  $M_F = m_\mu$ ,  $M_V = 0$ ,  $C_V = e$

---

<sup>7</sup>Notice that an earlier measurement using the caesium atom leads to a  $2.4 \sigma$  discrepancy with an opposite sign [105].




Figure 6: The Feynman diagrams for the one-loop QED and EW corrections to the muon  $g - 2$  in the SM: (a) QED; (b) SM Higgs boson; (c) SM charged gauge boson  $W^-$ ; (d) SM neutral gauge boson  $Z$ .

and  $C_A = 0$  in Eqs. (21)-(24), one obtains

$$a_\mu^{\text{QED}} = \frac{1}{16\pi^2} \lim_{x \rightarrow \infty} f_{\text{FV}}(x, x, e, 0, 0) = \frac{\alpha}{2\pi}, \quad (43)$$

where  $\alpha = e^2/(4\pi)$  is the fine structure constant. Note that the massless photon in the loop is not forbidden by Weinberg-Witten theorem since it is neutral and thus not coupled to the Lorentz-covariant conserved electromagnetic current. As a consequence, the contribution from Fig. 3(b) is actually finite in the limit of  $M_V \rightarrow 0$ .

- *Higgs boson.* The SM Higgs boson contributes to the muon anomalous magnetic moment at one-loop level via Fig. 6(b), belonging to the FS-type. Since the Higgs boson is much heavier than muon, it is safe to use the approximate expression in Eq. (14). Taking  $Q_S = 0$ ,  $M_F = m_\mu$ ,  $C_S = -\sqrt{2}\lambda m_\mu/M_H$  and  $C_P = 0$ , we arrive at

$$a_\mu^{\text{Higgs}} = -\frac{\lambda}{24\pi^2} \left[ 7 + 12 \ln \left( \frac{m_\mu}{M_H} \right) \right] \left( \frac{m_\mu}{M_H} \right)^4, \quad (44)$$

where  $\lambda$  and  $M_H$  are the quartic self-coupling and Higgs-boson mass, respectively. One can check that the result in Eq. (44) is equivalent to that of the Higgs contribution in Ref. [90] after completing the Feynman integral, which is proportional to the integral in Eq. (2), and

retaining the leading-order term of  $m_\mu/M_H$ . However, our method is much more convenient since the integral in Eq. (2) is rather tedious to perform analytically without imposing any conditions on mass hierarchies.

- *W and Z bosons.* The  $W$  and  $Z$  bosons contribute to the muon  $g - 2$  at the one-loop level via Fig. 6(c) and Fig. 6(d), belonging to the FV-type. Substituting  $Q_V = 1$ ,  $C_V = -C_A = g/(2\sqrt{2})$  into Eq. (29) and  $Q_V = 0$ ,  $C_V = g(-1 + 4\sin^2\theta_W)/(4\cos\theta_W)$ ,  $C_A = g/(4\cos\theta_W)$ ,  $M_F/m_\mu = 1$  into Eq. (28), respectively, we get

$$a_\mu^W = \frac{5G_F m_\mu^2}{12\sqrt{2}\pi^2}, \quad (45)$$

$$a_\mu^Z = -\frac{G_F m_\mu^2}{24\sqrt{2}\pi^2} \left[ 5 - (1 - 4\sin^2\theta_W)^2 \right], \quad (46)$$

where  $G_F$  is the Fermi constant,  $g$  is the  $SU(2)$  gauge coupling and  $\theta_W$  is the Weinberg angle. Note that the positive sign of  $a_\mu^W$  can be simply understood via the factor in Eq. (29), i.e.,  $-(2/3) \times (4 + 9Q_{W-}) = +10/3 > 0$ . As the Higgs contribution in Eq. (44) is highly suppressed by the Yukawa coupling of muon, the total one-loop EW contributions to  $a_\mu$  turn out to be

$$a_\mu^{\text{EW}} \approx a_\mu^W + a_\mu^Z = \frac{G_F m_\mu^2}{24\sqrt{2}\pi^2} \left[ 5 + (1 - 4\sin^2\theta_W)^2 \right], \quad (47)$$

which perfectly reproduces the well-known result in the literature [3, 114, 115].

Thus far we have rederived the one-loop QED and EW corrections to  $a_\mu$ . Although the results are not new at all, we have demonstrate that our formulae are advantageous in avoiding any tedious integrals stemming from the Feynman parametrization. More examples about new physics models contributing to muon  $g - 2$  will be discussed below.

### 3.2 Type-I seesaw model

The type-I seesaw model [116–120] extends the SM by adding three right-handed neutrinos  $N_R$ , which are the SM gauge singlets and possess a large Majorana mass term. In this model, tiny Majorana masses of three active neutrinos can be generated via the seesaw mechanism on the one hand; the CP-violating and out-of-equilibrium decays of heavy Majorana neutrinos in the early Universe could explain the cosmological matter-antimatter asymmetry via the leptogenesis mechanism on the other hand. The gauge-invariant Lagrangian of the type-I seesaw model reads

$$\mathcal{L}_{\text{Type-I}} = \mathcal{L}_{\text{SM}} - \left( \bar{L} Y_\nu \tilde{H} N_R + \frac{1}{2} \overline{N_R^C} M_R N_R + \text{h.c.} \right), \quad (48)$$

where  $L = (\nu_L, l_L)^T$  and  $\tilde{H} = i\sigma_2 H^*$  with  $H = (\varphi^+, \varphi^0)^T$  are the left-handed lepton doublet and the Higgs doublet,  $Y_\nu$  is the Dirac neutrino Yukawa coupling matrix while  $M_R$  is the Majorana mass matrix for right-handed neutrino singlets. After the SM gauge symmetry is spontaneously broken down, the Dirac mass matrix linking the left-handed and right-handed neutrinos is given




Figure 7: The Feynman diagram for the contributions from heavy and light Majorana neutrinos to the muon anomalous magnetic moment in the type-I seesaw model.

by  $M_D = Y_\nu v / \sqrt{2}$  with  $v \approx 246$  GeV the vacuum expectation value (VEV) of Higgs field. The overall  $6 \times 6$  neutrino mass matrix can be diagonalized by a  $6 \times 6$  unitary matrix via

$$\begin{pmatrix} \mathcal{V} & \mathcal{R} \\ \mathcal{S} & \mathcal{U} \end{pmatrix}^\dagger \begin{pmatrix} 0 & M_D \\ M_D^T & M_R \end{pmatrix} \begin{pmatrix} \mathcal{V} & \mathcal{R} \\ \mathcal{S} & \mathcal{U} \end{pmatrix}^* = \begin{pmatrix} \widehat{M}_\nu & 0 \\ 0 & \widehat{M}_R \end{pmatrix}, \quad (49)$$

where  $\widehat{M}_\nu = \text{diag}(m_1, m_2, m_3)$  and  $\widehat{M}_R = \text{diag}(M_1, M_2, M_3)$  with  $m_i (M_i)$  (for  $i = 1, 2, 3$ ) being light (heavy) neutrino masses, and  $\mathcal{V}, \mathcal{R}, \mathcal{S}, \mathcal{U}$  are all  $3 \times 3$  matrices satisfying unitarity conditions  $\mathcal{V}\mathcal{V}^\dagger + \mathcal{R}\mathcal{R}^\dagger = \mathcal{S}\mathcal{S}^\dagger + \mathcal{U}\mathcal{U}^\dagger = \mathbf{1}$  and  $\mathcal{V}\mathcal{S}^\dagger + \mathcal{R}\mathcal{U}^\dagger = \mathcal{S}\mathcal{V}^\dagger + \mathcal{U}\mathcal{R}^\dagger = \mathbf{0}$ . Note that in the limit of  $M_i \rightarrow \infty$ , we have  $\mathcal{R} \approx M_D M_i^{-1} \rightarrow 0$  and thus  $\mathcal{V}$  is reduced to the unitary PMNS matrix [121, 122]. The neutrino mass eigenstates  $(\widehat{\nu}_L, \widehat{N}_R^C)$  and the flavor eigenstates  $(\nu_L, N_R^C)$  are related by

$$\begin{pmatrix} \nu_L \\ N_R^C \end{pmatrix} = \begin{pmatrix} \mathcal{V} & \mathcal{R} \\ \mathcal{S} & \mathcal{U} \end{pmatrix} \begin{pmatrix} \widehat{\nu}_L \\ \widehat{N}_R^C \end{pmatrix}, \quad (50)$$

and the interaction term relevant to the muon anomalous magnetic moment is given by

$$\mathcal{L}_{\text{int}}^{\text{Type-I}} = \frac{g}{2\sqrt{2}} \bar{\mu} \gamma^\mu (1 - \gamma_5) \left( \mathcal{V}_{\mu i} \widehat{\nu}_i + \mathcal{R}_{\mu i} \widehat{N}_i \right) W_\mu^- + \text{h.c.}, \quad (51)$$

where  $\mathcal{V}_{\alpha i}$  and  $\mathcal{R}_{\alpha i}$  denotes the  $(\alpha, i)$ -element of  $\mathcal{V}$  and  $\mathcal{R}$  (for  $\alpha = e, \mu, \tau$  and  $i = 1, 2, 3$ ). As indicated by Eq. (51), due to the light-heavy neutrino mixing in Eq. (50), both light and heavy Majorana neutrinos are present in the charged-current interaction.

The contributions to the muon anomalous magnetic moment in the type-I seesaw model are of the FV-type and depicted in Fig. 7. For the heavy neutrinos, using Eq. (27) with the identification of  $Q_V = -1$ ,  $M_V = M_W$ ,  $M_F = M_i$  and  $C_V = -C_A = g\mathcal{R}_{\mu i}^*/(2\sqrt{2})$ , we obtain

$$a_\mu^N = \frac{G_F m_\mu^2}{4\sqrt{2}\pi^2} \sum_{i=1}^3 |\mathcal{R}_{\mu i}|^2 \mathcal{G}\left(\frac{M_i^2}{M_W^2}\right), \quad (52)$$

where the loop function  $\mathcal{G}(x)$  comes from the second-order loop function in Eq. (27), i.e.,

$$\mathcal{G}(x) = \frac{1}{2} g_{\text{FV}}^{(2)}(\sqrt{x}, -1) = \frac{10 - 43x + 78x^2 - 49x^3 + 4x^4 + 18x^3 \ln x}{6(1-x)^4}, \quad (53)$$

which is always positive and monotonically decreasing with the limits  $\mathcal{G}(0) = 5/3$  and  $\mathcal{G}(\infty) = 2/3$ . Thus the pure contributions from the heavy neutrinos to  $a_\mu$  are positive, as a consequence of the previous observation that  $g_{\text{FV}}^{(2)}(y, Q_V) > 0$  if the electric charge of  $W^-$  is smaller than the lower critical charge, i.e.,  $Q_{W^-} = -1 < Q_V^L = -5/9$  (cf. Table 2).

For the light neutrinos, making use of Eq. (28) with  $Q_V = -1$ ,  $M_V = M_W$ ,  $M_F = m_i$  and  $C_V = -C_A = g\mathcal{V}_{\mu i}^*/(2\sqrt{2})$ , one gets

$$a_\mu^\nu = \frac{G_F m_\mu^2}{4\sqrt{2}\pi^2} \times \frac{5}{3} \sum_{i=1}^3 |\mathcal{V}_{\mu i}|^2. \quad (54)$$

The contributions from the SM neutrinos can be obtained from Eq. (54) simply by imposing the unitarity condition  $|\mathcal{V}_{\mu 1}|^2 + |\mathcal{V}_{\mu 2}|^2 + |\mathcal{V}_{\mu 3}|^2 = 1$ , which is valid in the decoupling limit of  $M_i \rightarrow \infty$ . As a result, we have

$$a_\mu^{\nu, \text{SM}} = \frac{G_F m_\mu^2}{4\sqrt{2}\pi^2} \times \frac{5}{3}. \quad (55)$$

The final result for the extra contribution to muon  $g-2$  in the type-I seesaw model can be obtained by subtracting the SM neutrino contribution, namely,

$$\Delta a_\mu^{\text{Type-I}} = a_\mu^N + a_\mu^\nu - a_\mu^{\nu, \text{SM}} = \frac{G_F m_\mu^2}{4\sqrt{2}\pi^2} \sum_{i=1}^3 \left\{ |\mathcal{R}_{\mu i}|^2 \left[ \mathcal{G}\left(\frac{M_i^2}{M_W^2}\right) - \frac{5}{3} \right] \right\}, \quad (56)$$

which is obviously negative because of the upper bound on the loop function (i.e.,  $\mathcal{G}(x) < 5/3$  for  $x > 0$ ). Note that the unitarity condition  $\mathcal{V}\mathcal{V}^\dagger + \mathcal{R}\mathcal{R}^\dagger = \mathbf{1}$  has been used. The result in Eq. (56) is the same as that derived in Ref. [52], where the method of Feynman parametrization has been implemented instead.

### 3.3 Type-II seesaw model

Another intriguing way to generate tiny neutrino masses is the type-II seesaw model [123–128], which augments the SM with a triplet scalar with the hypercharge  $-1$ , i.e.,

$$\Delta = \sqrt{2} \begin{pmatrix} \xi^-/\sqrt{2} & -\xi^0 \\ \xi^{--} & -\xi^-/\sqrt{2} \end{pmatrix}. \quad (57)$$

The triplet has been written in the adjoint representation and the electric charge of each component has been explicitly indicated.

The neutral components of the triplet (i.e.,  $\xi^0$  and its complex conjugate) mix with those of the SM Higgs doublet, leading to two CP-even scalar bosons ( $H^0, h^0$ ) and one CP-odd scalar boson  $A^0$ . The Feynman diagrams for their contributions to  $a_\mu$  are shown in Fig. 8(f), which however are highly suppressed by the Yukawa coupling of muon. The leading contributions are made by the singly- and doubly-charged scalars and the relevant interaction reads

$$\begin{aligned} \mathcal{L}_{\text{int}}^{\text{Type-II}} = & \left[ -\frac{1}{2} \cos \beta \left( Y_\Delta^\dagger \mathcal{V} \right)_{\mu i} \bar{\nu}_i (1 - \gamma_5) \mu H^+ - \frac{1}{\sqrt{2}} (Y_\Delta)_{\alpha\mu}^* \bar{l}_\alpha^C (1 - \gamma_5) \mu H^{++} \right. \\ & \left. - \frac{g}{2\sqrt{2}} \mathcal{V}_{\mu i}^* \bar{\nu}_i \gamma^\mu (1 - \gamma_5) \mu W_\mu^+ + \text{h.c.} \right] - \sqrt{2} g e v_\Delta \cos \beta (H^- W_\mu^+ + H^+ W_\mu^-) A^\mu, \quad (58) \end{aligned}$$




Figure 8: The Feynman diagrams for the contributions from the charged and neutral scalars to the muon anomalous magnetic moment in the type-II seesaw model: (a) and (b) from the doubly-charged scalar  $H^{--}$  and the charged leptons  $l_\alpha^C$  (for  $\alpha = e, \mu, \tau$ ); (c), (d) and (e) from the singly-charged scalar  $H^-$  and the neutrinos  $\nu_i$  (for  $i = 1, 2, 3$ ); (f) from the neutral scalars, where  $H^0$ ,  $h^0$  and  $A^0$  are two CP-even and one CP-odd neutral scalars coming from the mixing between the SM Higgs boson and the neutral components of the triplet scalar.

where  $Y_\Delta$  represents the Yukawa coupling matrix between the scalar triplet and the lepton doublet, and  $\mathcal{V}$  is the unitary PMNS matrix. The flavor eigenstates  $(\xi^\pm, \xi^{\pm\pm})$  are related to the mass eigenstates  $(H^\pm, H^{\pm\pm})$  via

$$\xi^\pm = H^\pm \cos \beta - G^\pm \sin \beta, \quad \xi^{\pm\pm} = H^{\pm\pm}, \quad (59)$$

with  $\beta = \cos^{-1} \left( v / \sqrt{v^2 + 2v_\Delta^2} \right)$  and  $v_\Delta$  being the VEV of  $\Delta$ . The Feynman diagrams contributing to  $a_\mu$  are shown in Fig. 8. The diagrams involving doubly-charged scalars can be found in Fig. 8(a) and Fig. 8(b). Noticing  $m_\alpha \ll M_{H^{\pm\pm}}$  (for  $\alpha = e, \mu, \tau$ ) and utilizing Eq. (14) with  $C_S = -C_P = -(Y_\Delta)_{\alpha\mu}^* / \sqrt{2}$ ,  $Q_S = -2$ ,  $M_S = M_{H^{\pm\pm}}$  and  $M_F = m_\alpha$ , one can immediately obtain

$$\Delta a_\mu^{H^{\pm\pm}} = -\frac{1}{12\pi^2} \left( Y_\Delta^\dagger Y_\Delta \right)_{\mu\mu} \left( \frac{m_\mu}{M_{H^{\pm\pm}}} \right)^2. \quad (60)$$

In addition, Figs. 8(c), (d) and (e) represent the contributions from the singly-charged scalars. For Fig. 8(c), we can also use Eq. (14) but with  $C_S = -C_P = -\cos \beta \left( Y_\Delta^\dagger \mathcal{V} \right)_{\mu i} / 2$ ,  $Q_S = -1$ ,  $M_S = M_{H^\pm}$  and  $M_F = m_\alpha$  to derive

$$\Delta a_\mu^{H^\pm(c)} = -\frac{1}{96\pi^2} \left( Y_\Delta^\dagger Y_\Delta \right)_{\mu\mu} \left( \frac{m_\mu}{M_{H^\pm}} \right)^2, \quad (61)$$

where we have used the experimental bound  $v_\Delta \ll v$  coming from the precision measurements of the  $\rho$  parameter. For Fig. 8(d) and Fig. 8(e), which belong to the FSV-type diagrams, we notice




Figure 9: The Feynman diagram for the contribution from the new gauge boson  $Z'$  to the muon anomalous magnetic moment.

that  $m_\mu \ll M_{H^\pm}, M_W$  is perfectly satisfied and then make use of Eq. (39) with  $C_S = -C_P = -\cos\beta \left(Y_\Delta^\dagger \mathcal{V}\right)_{\mu i} / 2$ ,  $C_V = -C_A = g\mathcal{V}_{\mu i}^* / (2\sqrt{2})$ ,  $\mu_0 = -\sqrt{2}gv_\Delta \cos\beta$ ,  $M_S = M_{H^\pm}$ ,  $M_V = M_W$  and  $M_F = m_\alpha$ . The final result is

$$\Delta a_\mu^{H^\pm(\text{d})+(\text{e})} = \frac{G_F v_\Delta^2}{6\sqrt{2}\pi^2} \frac{m_\mu^2 \ln(M_W^2/M_{H^\pm}^2)}{M_W^2 - M_{H^\pm}^2} \left(Y_\Delta^\dagger Y_\Delta\right)_{\mu\mu}, \quad (62)$$

where is highly suppressed by the tiny neutrino masses due to  $G_F v_\Delta^2 \propto m_i^2/M_W^2$ . Therefore, the contribution in Eq. (62) can be safely neglected, and the overall result is given by

$$\Delta a_\mu^{H^\pm} \approx \Delta a_\mu^{H^\pm(\text{c})} = -\frac{1}{96\pi^2} \left(Y_\Delta^\dagger Y_\Delta\right)_{\mu\mu} \left(\frac{m_\mu}{M_{H^\pm}}\right)^2. \quad (63)$$

As shown in Eqs. (60) and (63), the dominant contributions to the muon anomalous magnetic moment in the type-II seesaw turn out to be negative in the whole parameter space of  $Y_\Delta$ . The same results have also been obtained in Ref. [129]. It is worthwhile to mention that this wrong sign can be simply observed from the second-order loop function in Eq. (13). Since the electric charges of the charged scalars in the type-II seesaw model are smaller than its lower critical charge  $Q_S^L = -2/3$  (cf. Table 1), we have

$$g_{\text{FS}}^{(2)}(y, Q_S = -1) < 0, \quad g_{\text{FS}}^{(2)}(y, Q_S = -2) < 0, \quad (64)$$

implying the overall negative sign of  $\Delta a_\mu$ . Note that  $|C_S|^2 - |C_P|^2 = 0$  holds for both doubly- and singly-charged scalars, so the first-order loop function in Eq. (12) is irrelevant.

### 3.4 $Z'$ model

The linear combinations  $B/3 - L_\alpha$ ,  $L_e - L_\mu$  and  $L_\mu - L_\tau$  of the baryon number  $B$  and the lepton number  $L_\alpha$  (for  $\alpha = e, \mu, \tau$ ) are anomaly-free, so any one of them can naturally be promoted to a gauge symmetry<sup>8</sup> [130–133]. The neutral gauge boson  $Z'$  corresponding to the new U(1) gauge

<sup>8</sup>Note that  $L_e - L_\mu$  and  $L_\mu - L_\tau$  can be gauged without including right-handed neutrinos, whereas  $B/3 - L_\alpha$  (for  $\alpha = e, \mu, \tau$ ) can be gauged only when at least one right-handed neutrino is introduced.

symmetry mediates extra interaction among charged leptons and can contribute to the muon  $g-2$ . For illustration, we assume that the coupling is flavor-diagonal and both  $(\nu_\mu, \mu)_L$  and  $\mu_R$  take the same charge under this new  $U(1)$  gauge group. The relevant Lagrangian is

$$\mathcal{L}_{\text{int}}^{Z'} = g_{Z'} \bar{\mu} \gamma^\mu \mu Z'_\mu, \quad (65)$$

with  $g_{Z'}$  the coupling constant. The Feynman diagram contributing to the muon  $g-2$  is shown in Fig. 9. By inserting  $C_V = g_{Z'}$ ,  $C_A = 0$ ,  $Q_V = 0$ ,  $M_V = M_{Z'}$  and  $M_F = m_\mu$  into Eqs. (21)-(24), one obtains

$$\Delta a_\mu^{Z'} = \frac{g_{Z'}^2}{16\pi^2} \mathcal{H}\left(\frac{m_\mu}{M_{Z'}}\right), \quad (66)$$

where the loop function

$$\mathcal{H}(x) = \frac{2}{x^4} \left[ x^2 (x^2 - 2) + 2 (2x^2 - 1) \ln x + 2x^2 (4x^2 - 2x^4 - 1) \frac{\Lambda(x^2, x, 1)}{\lambda(x^2, x^2, 1)} \right]. \quad (67)$$

It is easy to verify that the loop function in Eq. (67) is non-negative with the following finite limits

$$\lim_{x \rightarrow 0} \mathcal{H}(x) = 0, \quad \lim_{x \rightarrow \infty} \mathcal{H}(x) = 2.$$

Thus one can reach the conclusion that the neutral vector boson with a vector-like interaction with muon will always make a positive contribution to  $a_\mu$ . The formulae in Eqs. (66) and (67) are valid for the whole parameter space of  $M_{Z'}$ . In particular, in the hierarchical cases of  $m_\mu \ll M_{Z'}$  and  $M_{Z'} \ll m_\mu$ , we have

$$\Delta a_\mu^{Z'} = \begin{cases} \frac{g_{Z'}^2}{12\pi^2} \left(\frac{m_\mu}{M_{Z'}}\right)^2, & m_\mu \ll M_{Z'}; \\ \frac{g_{Z'}^2}{8\pi^2}, & M_{Z'} \ll m_\mu. \end{cases} \quad (68)$$

Moreover, in the limit of  $M_{Z'} \rightarrow m_\mu$  or equivalently  $x = m_\mu/M_{Z'} \rightarrow 1$ , we can obtain  $\mathcal{H}(x)|_{x \rightarrow 1} = 2(2\pi - 3\sqrt{3})/(3\sqrt{3})$  from Eq. (67). Note that the result in Eq. (68), which has previously been obtained in the literature [52, 134], is valid only in the assumption of a strong mass hierarchy. For a complete scan of the parameter space of  $M_{Z'}$  in the most general case, it is necessary to use the exact formulae in Eqs. (66)-(67).

### 3.5 Leptoquark

Leptoquarks are hypothetical scalar or vector bosons that are coupled simultaneously to leptons and quarks. In the first place, leptoquarks arise naturally from the theories of grand unification. At present, they have been extensively discussed in the literature as new-physics explanations for, e.g., tiny neutrino masses via radiative corrections,  $B$  anomalies and muon anomalous magnetic moment. See, for example, Ref. [135], for a recent review on leptoquarks. For illustration, we consider the scalar leptoquark  $S_1$ , whose quantum numbers are assigned as  $(\bar{\mathbf{3}}, \mathbf{1}, 1/3)$  under the  $SU(3)_C \otimes SU(2)_L \otimes U(1)_Y$  gauge group. The main contribution to the muon anomalous magnetic




Figure 10: The Feynman diagrams for the contribution from the scalar leptoquark  $S_1$  to the muon anomalous magnetic moment.

moment comes from the interaction among muon, leptoquark and top quark, as shown in Fig. 10. The relevant Lagrangian can be written as

$$\mathcal{L}_{\text{int}}^{S_1} = \bar{t}^C (C_S + C_P \gamma_5) \mu S_1 + \text{h.c.} . \quad (69)$$

Assuming  $S_1$  to be much heavier than muon and substituting  $Q_S = Q_{S_1^\dagger} = -1/3$ ,  $M_S = M_{S_1}$  and  $M_F = m_t$  into Eq. (12), we arrive at

$$\Delta a_\mu^{S_1} = \frac{1}{16\pi^2} (|C_S|^2 - |C_P|^2) \frac{m_\mu}{M_{S_1}} \mathcal{I} \left( \frac{m_t}{M_{S_1}} \right) , \quad (70)$$

where the loop function  $\mathcal{I}(x)$  is derived from the first-order loop function in Eq. (12), i.e.,

$$\mathcal{I}(x) = 3g_{\text{FS}}^{(1)} \left( x, -\frac{1}{3} \right) = \frac{x [x^4 - 8x^2 + 7 + 4(x^2 + 2) \ln x]}{(x^2 - 1)^3} , \quad (71)$$

with the color factor 3 being included. Notice that we have assumed  $S_1$  to be coupled to both left- and right-handed muons, implying that the contribution to  $a_\mu$  is dominated by the first-order loop function. Two helpful comments are in order. First, the sign of  $\Delta a_\mu^{S_1}$  in Eq. (70) depends on the relative sizes of  $|C_S|$  and  $|C_P|$ , since the function  $g_{\text{FS}}^{(1)}(x, -1/3)$  is always positive. The latter can be understood by noticing that the electric charge of  $S_1^\dagger$  is larger than the upper critical charge, i.e.,  $Q_{S_1^\dagger} = -1/3 > Q_S^{\text{U}} = -1/2$  (cf. Table 1). Second, if we further assume that  $S_1$  is also much heavier than top quark, then the expression in Eq. (70) will be reduced to

$$\Delta a_\mu^{S_1} = \frac{1}{16\pi^2} (|C_S|^2 - |C_P|^2) \frac{m_\mu m_t}{M_{S_1}^2} \left[ -7 - 8 \ln \left( \frac{m_t}{M_{S_1}} \right) \right] . \quad (72)$$

This result has previously been derived in the literature [50, 136, 137]. However, for a complete scan of the parameter space, it is necessary to utilize the general expression of the loop function in Eq. (71).

## 4 Summary

Motivated by the recent measurement of the muon anomalous magnetic moment  $a_\mu$ , which is at odds with the SM prediction at the  $4.2\sigma$  level, we examine possible new-physics contributions to  $a_\mu$  at the one-loop order. A model-independent analysis of the general loop functions has been carried out. The main results in the present paper are summarized as follows.

First, the contributions to  $a_\mu$  at the one-loop order can be classified into three categories by different types of particles running in the loop: (1) The FS-type with one fermion and one scalar boson; (2) The FV-type with one fermion and one vector boson; (3) The FSV-type with one fermion, one scalar boson and one vector boson. Although the one-loop Feynman diagrams for  $a_\mu$  have been studied for a long time, it is surprising that the general formulae have never been explicitly given in the literature. Instead of expressing the final results in terms of the integral over the Feynman parameter, we use the Passarino-Veltman functions to derive a general and compact formula for  $a_\mu$  without any integrals in the end. The final results can be found in Eqs. (6)-(9) for the FS-type, Eqs. (21)-(24) for the FV-type, and Eqs. (35)-(38) for the FSV-type.

Second, we attempt to clarify when the new-physics contribution to  $a_\mu$  is positive so as to reconcile the theoretical prediction with the experimental measurement. For this purpose, we have investigated the basic properties of the loop functions of the FS- and FV-type. In the assumption that the new particles are much heavier than muon, the loop functions can be series expanded in terms of the mass ratio  $m_\mu/M_S$  (for the FS-type) or  $m_\mu/M_V$  (for the FV-type). At different orders of  $m_\mu/M_S$  or  $m_\mu/M_V$ , the loop functions exhibit interesting behaviors depending on the electric charges of the particles running in the loop. Roughly speaking, there exists an upper and a lower critical value for the electric charge  $Q_S$  or  $Q_V$ , which can be used to claim whether the contribution to  $a_\mu$  is positive or negative. The properties of the first- and second-order loop functions are summarized in Table 1 and Table 2. These properties have also been applied to make a quick judgment about the sign of  $\Delta a_\mu$  in a concrete model.

Third, it is interesting to find that there are two extra constraints for the FV-type model. First, if there exists in the theory a heavy fermion that is coupled simultaneously to both left- and right-handed muon and the vector boson, then the couplings should decrease no more slowly than  $\mathcal{O}(1/\sqrt{M_F})$  in the limit  $M_F \rightarrow \infty$ . Second, according to the Weinberg-Witten theorem, there cannot be a massless vector boson with a nonvanishing electric charge, contributing to muon anomalous magnetic moment. These constraints may be automatically satisfied in a self-consistent theory, but should be carefully taken into account in phenomenological studies.

Finally, our general formulae have been applied to several well-motivated theoretical models. The correct results in all these models can be reproduced in a straightforward way. If the  $4.2\sigma$  discrepancy between the theoretical prediction and the experimental measurement of  $a_\mu$  is further confirmed, new physics will definitely be introduced to explain such a discrepancy. In this case, the main results presented in our paper will be helpful for both model building and phenomenological studies associated with the muon anomalous magnetic moment.

# Acknowledgements

This work was supported in part by the National Natural Science Foundation of China under grant No. 11775232 and No. 11835013, by the Key Research Program of the Chinese Academy of Sciences under grant No. XDPB15, and by the CAS Center for Excellence in Particle Physics.

# References

- [1] B. Abi *et al.* [Muon  $g-2$ ], “Measurement of the Positive Muon Anomalous Magnetic Moment to 0.46 ppm,” Phys. Rev. Lett. **126** (2021) no.14, 141801 [arXiv:2104.03281 [hep-ex]].
- [2] G. W. Bennett *et al.* [Muon  $g-2$  Collaboration], “Final Report of the Muon E821 Anomalous Magnetic Moment Measurement at BNL,” Phys. Rev. D **73** (2006) 072003 [hep-ex/0602035].
- [3] T. Aoyama, N. Asmussen, M. Benayoun, J. Bijnens, T. Blum, M. Bruno, I. Caprini, C. M. Carloni Calame, M. Cè and G. Colangelo, *et al.* “The anomalous magnetic moment of the muon in the Standard Model,” Phys. Rept. **887** (2020), 1-166 [arXiv:2006.04822 [hep-ph]].
- [4] M. Davier, A. Hoecker, B. Malaescu and Z. Zhang, “Reevaluation of the Hadronic Contributions to the Muon  $g-2$  and to  $\alpha(M_Z)$ ,” Eur. Phys. J. C **71**, 1515 (2011) [erratum: Eur. Phys. J. C **72**, 1874 (2012)] [arXiv:1010.4180 [hep-ph]].
- [5] M. Davier, A. Hoecker, B. Malaescu and Z. Zhang, “Reevaluation of the hadronic vacuum polarisation contributions to the Standard Model predictions of the muon  $g-2$  and  $\alpha(m_Z^2)$  using newest hadronic cross-section data,” Eur. Phys. J. C **77**, no.12, 827 (2017) [arXiv:1706.09436 [hep-ph]].
- [6] M. Davier, A. Hoecker, B. Malaescu and Z. Zhang, “A new evaluation of the hadronic vacuum polarisation contributions to the muon anomalous magnetic moment and to  $\alpha(m_Z^2)$ ,” Eur. Phys. J. C **80**, no.3, 241 (2020) [erratum: Eur. Phys. J. C **80**, no.5, 410 (2020)] [arXiv:1908.00921 [hep-ph]].
- [7] S. Borsanyi, Z. Fodor, J. N. Guenther, C. Hoelbling, S. D. Katz, L. Lellouch, T. Lippert, K. Miura, L. Parato and K. K. Szabo, *et al.* “Leading hadronic contribution to the muon magnetic moment from lattice QCD,” Nature **593**, no.7857, 51-55 (2021) [arXiv:2002.12347 [hep-lat]].
- [8] H. B. Zhang, C. X. Liu, J. L. Yang and T. F. Feng, “Muon anomalous magnetic dipole moment in the  $\mu\nu$ SSM,” [arXiv:2104.03489 [hep-ph]].
- [9] G. Arcadi, L. Calibbi, M. Fedele and F. Mescia, “Muon  $g-2$  and  $B$ -anomalies from Dark Matter,” [arXiv:2104.03228 [hep-ph]].
- [10] B. Zhu and X. Liu, “Probing light dark matter with scalar mediator: muon  $(g-2)$  deviation, the proton radius puzzle,” [arXiv:2104.03238 [hep-ph]].

- [11] T. Nomura and H. Okada, “Explanations for anomalies of muon anomalous magnetic dipole moment,  $b \rightarrow s\mu\bar{\mu}$  and radiative neutrino masses in a leptoquark model,” [arXiv:2104.03248 [hep-ph]].
- [12] S. Baum, M. Carena, N. R. Shah and C. E. M. Wagner, “The Tiny (g-2) Muon Wobble from Small- $\mu$  Supersymmetry,” [arXiv:2104.03302 [hep-ph]].
- [13] M. Endo, K. Hamaguchi, S. Iwamoto and T. Kitahara, “Supersymmetric Interpretation of the Muon  $g - 2$  Anomaly,” [arXiv:2104.03217 [hep-ph]].
- [14] S. F. Ge, X. D. Ma and P. Pasquini, “Probing the Dark Axion Portal with Muon Anomalous Magnetic Moment,” [arXiv:2104.03276 [hep-ph]].
- [15] W. Ahmed, I. Khan, J. Li, T. Li, S. Raza and W. Zhang, “The Natural Explanation of the Muon Anomalous Magnetic Moment via the Electroweak Supersymmetry from the Gm-SUGRA in the MSSM,” [arXiv:2104.03491 [hep-ph]].
- [16] P. Das, M. K. Das and N. Khan, “The FIMP-WIMP dark matter and Muon g-2 in the extended singlet scalar model,” [arXiv:2104.03271 [hep-ph]].
- [17] X. F. Han, T. Li, H. X. Wang, L. Wang and Y. Zhang, “Lepton-specific inert two-Higgs-doublet model confronted with the new results for muon and electron g-2 anomalies and multi-lepton searches at the LHC,” [arXiv:2104.03227 [hep-ph]].
- [18] W. Y. Keung, D. Marfatia and P. Y. Tseng, “Axion-like particles, two-Higgs-doublet models, leptoquarks, and the electron and muon  $g - 2$ ,” [arXiv:2104.03341 [hep-ph]].
- [19] Y. Bai and J. Berger, “Muon  $g - 2$  in Lepton Portal Dark Matter,” [arXiv:2104.03301 [hep-ph]].
- [20] V. Brdar, S. Jana, J. Kubo and M. Lindner, “Semi-secretly interacting ALP as an explanation of Fermilab muon  $g - 2$  measurement,” [arXiv:2104.03282 [hep-ph]].
- [21] L. Zu, X. Pan, L. Feng, Q. Yuan and Y. Z. Fan, “Constraining  $U(1)_{L_\mu-L_\tau}$  charged dark matter model for muon  $g - 2$  anomaly with AMS-02 electron and positron data,” [arXiv:2104.03340 [hep-ph]].
- [22] M. Ibe, S. Kobayashi, Y. Nakayama and S. Shirai, “Muon  $g - 2$  in Gauge Mediation without SUSY CP Problem,” [arXiv:2104.03289 [hep-ph]].
- [23] P. Cox, C. Han and T. T. Yanagida, “Muon  $g - 2$  and Co-annihilating Dark Matter in the MSSM,” [arXiv:2104.03290 [hep-ph]].
- [24] M. Abdughani, Y. Z. Fan, L. Feng, Y. L. Sming Tsai, L. Wu and Q. Yuan, “A common origin of muon g-2 anomaly, Galaxy Center GeV excess and AMS-02 anti-proton excess in the NMSSM,” [arXiv:2104.03274 [hep-ph]].

- [25] M. Van Beekveld, W. Beenakker, M. Schutten and J. De Wit, “Dark matter, fine-tuning and  $(g - 2)_\mu$  in the pMSSM,” [arXiv:2104.03245 [hep-ph]].
- [26] M. A. Buen-Abad, J. Fan, M. Reece and C. Sun, “Challenges for an axion explanation of the muon  $g - 2$  measurement,” [arXiv:2104.03267 [hep-ph]].
- [27] C. H. Chen, C. W. Chiang and T. Nomura, “Muon  $g - 2$  in two-Higgs-doublet model with type-II seesaw mechanism,” [arXiv:2104.03275 [hep-ph]].
- [28] K. S. Babu, S. Jana, M. Lindner and V. P. K, “Muon  $g - 2$  Anomaly and Neutrino Magnetic Moments,” [arXiv:2104.03291 [hep-ph]].
- [29] P. M. Ferreira, B. L. Gonçalves, F. R. Joaquim and M. Sher, “ $(g - 2)_\mu$  in the 2HDM and slightly beyond – an updated view,” [arXiv:2104.03367 [hep-ph]].
- [30] W. Yin, “Muon  $g - 2$  Anomaly in Anomaly Mediation,” [arXiv:2104.03259 [hep-ph]].
- [31] C. Han, “Muon  $g-2$  and CP violation in MSSM,” [arXiv:2104.03292 [hep-ph]].
- [32] Y. Gu, N. Liu, L. Su and D. Wang, “Heavy Bino and Slepton for Muon  $g-2$  Anomaly,” [arXiv:2104.03239 [hep-ph]].
- [33] F. Wang, L. Wu, Y. Xiao, J. M. Yang and Y. Zhang, “GUT-scale constrained SUSY in light of E989 muon  $g-2$  measurement,” [arXiv:2104.03262 [hep-ph]].
- [34] L. Calibbi, M. L. López-Ibañez, A. Melis and O. Vives, “Implications of the Muon  $g-2$  result on the flavour structure of the lepton mass matrix,” [arXiv:2104.03296 [hep-ph]].
- [35] H. X. Wang, L. Wang and Y. Zhang, “muon  $g - 2$  anomaly and  $\mu$ - $\tau$ -philic Higgs doublet with a light CP-even component,” [arXiv:2104.03242 [hep-ph]].
- [36] T. Li, J. Pei and W. Zhang, “Muon Anomalous Magnetic Moment and Higgs Potential Stability in the 331 Model from  $E_6$ ,” [arXiv:2104.03334 [hep-ph]].
- [37] M. Cadeddu, N. Cargioli, F. Dordei, C. Giunti and E. Picciau, “Muon and electron  $g-2$ , proton and cesium weak charges implications on dark  $\mathbf{Z_d}$  models,” [arXiv:2104.03280 [hep-ph]].
- [38] J. Chen, Q. Wen, F. Xu and M. Zhang, “Flavor Anomalies Accommodated in A Flavor Gauged Two Higgs Doublet Model,” [arXiv:2104.03699 [hep-ph]].
- [39] P. Escribano, J. Terol-Calvo and A. Vicente, “ $(g - 2)_{e,\mu}$  in an extended inverse type-III seesaw,” [arXiv:2104.03705 [hep-ph]].
- [40] J. C. Eung and T. Mondal, “Leptophilic bosons and muon  $g-2$  at lepton colliders,” [arXiv:2104.03701 [hep-ph]].
- [41] A. Aboubrahim, M. Klasen and P. Nath, “What Fermilab  $(g - 2)_\mu$  experiment tells us about discovering SUSY at HL-LHC and HE-LHC,” [arXiv:2104.03839 [hep-ph]].

- [42] B. Bhattacharya, A. Datta, D. Marfatia, S. Nandi and J. Waite, “Axion-like particles resolve the  $B \rightarrow \pi K$  and  $g-2$  anomalies,” [arXiv:2104.03947 [hep-ph]].
- [43] J. L. Yang, H. B. Zhang, C. X. Liu, X. X. Dong and T. F. Feng, “Muon  $(g - 2)$  in the B-LSSM,” [arXiv:2104.03542 [hep-ph]].
- [44] G. Arcadi, Á. S. De Jesus, T. B. De Melo, F. S. Queiroz and Y. S. Villamizar, “A 2HDM for the  $g-2$  and Dark Matter,” [arXiv:2104.04456 [hep-ph]].
- [45] C. T. Lu, R. Ramos and Y. L. Sming Tsai, “Shedding light on dark matter with recent muon  $(g - 2)$  and Higgs exotic decay measurements,” [arXiv:2104.04503 [hep-ph]].
- [46] M. Chakraborti, L. Roszkowski and S. Trojanowski, “GUT-constrained supersymmetry and dark matter in light of the new  $(g - 2)_\mu$  determination,” JHEP **05** (2021), 252 [arXiv:2104.04458 [hep-ph]].
- [47] T. Li, M. A. Schmidt, C. Y. Yao and M. Yuan, “Charged lepton flavor violation in light of the muon magnetic moment anomaly and colliders,” [arXiv:2104.04494 [hep-ph]].
- [48] J. Y. Cen, Y. Cheng, X. G. He and J. Sun, “Flavor Specific  $U(1)_{B_q-L_\mu}$  Gauge Model for Muon  $g - 2$  and  $b \rightarrow s\bar{\mu}\mu$  Anomalies,” [arXiv:2104.05006 [hep-ph]].
- [49] D. Borah, M. Dutta, S. Mahapatra and N. Sahu, “Muon  $(g - 2)$  and XENON1T Excess with Boosted Dark Matter in  $L_\mu - L_\tau$  Model,” [arXiv:2104.05656 [hep-ph]].
- [50] D. Marzocca and S. Trifinopoulos, “A Minimal Explanation of Flavour Anomalies: B-Meson Decays, Muon Magnetic Moment, and the Cabibbo Angle,” [arXiv:2104.05730 [hep-ph]].
- [51] M. Du, J. Liang, Z. Liu and V. Tran, “A vector leptoquark interpretation of the muon  $g - 2$  and  $B$  anomalies,” [arXiv:2104.05685 [hep-ph]].
- [52] S. Zhou, “Neutrino Masses, Leptonic Flavor Mixing and Muon  $(g - 2)$  in the Seesaw Model with the  $U(1)_{L_\mu-L_\tau}$  Gauge Symmetry,” [arXiv:2104.06858 [hep-ph]].
- [53] K. Ban, Y. Jho, Y. Kwon, S. C. Park, S. Park and P. Y. Tseng, “A comprehensive study of vector leptoquark on the  $B$ -meson and Muon  $g-2$  anomalies,” [arXiv:2104.06656 [hep-ph]].
- [54] L. A. Anchordoqui, I. Antoniadis, X. Huang, D. Lüst and T. R. Taylor, “Muon  $g - 2$  discrepancy within D-brane string compactifications,” [arXiv:2104.06854 [hep-ph]].
- [55] A. E. Cárcamo Hernández, S. Kovalenko, M. Maniatis and I. Schmidt, “Fermion mass hierarchy and  $g-2$  anomalies in an extended 3HDM Model,” [arXiv:2104.07047 [hep-ph]].
- [56] H. Baer, V. Barger and H. Serce, “muon anomalous magnetic moment, supersymmetry, naturalness, LHC search limits and the landscape,” [arXiv:2104.07597 [hep-ph]].
- [57] W. Altmannshofer, S. A. Gadam, S. Gori and N. Hamer, “Explaining  $(g - 2)_\mu$  with Multi-TeV Sleptons,” [arXiv:2104.08293 [hep-ph]].

- [58] R. Balkin, C. Delaunay, M. Geller, E. Kajomovitz, G. Perez, Y. Shpilman and Y. Soreq, “A Custodial Symmetry for Muon  $g-2$ ,” [arXiv:2104.08289 [hep-ph]].
- [59] G. Cacciapaglia, C. Cot and F. Sannino, “Naturalness of lepton non-universality and muon  $g-2$ ,” [arXiv:2104.08818 [hep-ph]].
- [60] A. Dasgupta, S. K. Kang and M. Park, “Neutrino mass and  $(g - 2)_\mu$  with dark  $U(1)_D$  symmetry,” [arXiv:2104.09205 [hep-ph]].
- [61] A. Aboubrahim, P. Nath and R. M. Syed, “Yukawa coupling unification in an  $SO(10)$  model consistent with Fermilab  $(g - 2)_\mu$  result,” [arXiv:2104.10114 [hep-ph]].
- [62] E. Ma, “Gauged lepton number, Dirac neutrinos, dark matter, and muon  $g - 2$ ,” Phys. Lett. B **819** (2021), 136402 [arXiv:2104.10324 [hep-ph]].
- [63] A. Jueid, J. Kim, S. Lee and J. Song, “Type-X two Higgs doublet model in light of the muon  $g - 2$ : confronting Higgs and collider data,” [arXiv:2104.10175 [hep-ph]].
- [64] P. Fileviez Pérez, C. Murgui and A. D. Plascencia, “Leptoquarks and Matter Unification: Flavor Anomalies and the Muon  $g - 2$ ,” [arXiv:2104.11229 [hep-ph]].
- [65] V. G. Baryshevsky and P. I. Porshnev, “Predicting outcomes of electric dipole and magnetic moment experiments,” [arXiv:2104.12230 [hep-ph]].
- [66] K. Ghorbani, “Light vector dark matter with scalar mediator and muon  $g-2$  anomaly,” [arXiv:2104.13810 [hep-ph]].
- [67] J. A. Carpio, K. Murase, I. M. Shoemaker and Z. Tabrizi, “High-energy cosmic neutrinos as a probe of the vector mediator scenario in light of the muon  $g - 2$  anomaly and Hubble tension,” [arXiv:2104.15136 [hep-ph]].
- [68] B. A. Arbuzov and I. V. Zaitsev, “Calculation of the contribution to muon  $g - 2$  due to the effective anomalous three boson interaction and the new experimental result,” [arXiv:2105.00903 [hep-ph]].
- [69] J. S. Alvarado, S. F. Mantilla, R. Martinez and F. Ochoa, “A non-universal  $U(1)_X$  extension to the Standard Model to study the  $B$  meson anomaly and muon  $g - 2$ ,” [arXiv:2105.04715 [hep-ph]].
- [70] M. Chakraborti, S. Heinemeyer and I. Saha, “Improved  $(g - 2)_\mu$  Measurements and Supersymmetry : Implications for  $e^+e^-$  colliders,” [arXiv:2105.06408 [hep-ph]].
- [71] W. F. Chang, “One colorful resolution to the neutrino mass generation, three lepton flavor universality anomalies, and the Cabibbo angle anomaly,” [arXiv:2105.06917 [hep-ph]].
- [72] M. D. Zheng and H. H. Zhang, “Studying the  $b \rightarrow s\ell^+\ell^-$  Anomalies and  $(g - 2)_\mu$  in RPV-MSSM Framework with Inverse Seesaw,” [arXiv:2105.06954 [hep-ph]].

- [73] B. Dutta, S. Ghosh, P. Huang and J. Kumar, “Explaining  $g_\mu - 2$  and  $R_{K^{(*)}}$  using the light mediators of  $U(1)_{T3R}$ ,” [arXiv:2105.07655 [hep-ph]].
- [74] D. Zhang, “Radiative neutrino masses, lepton flavor mixing and muon  $g - 2$  in a leptoquark model,” [arXiv:2105.08670 [hep-ph]].
- [75] W. S. Hou, R. Jain, C. Kao, G. Kumar and T. Modak, “Collider Prospects for Muon  $g - 2$  in General Two Higgs Doublet Model,” [arXiv:2105.11315 [hep-ph]].
- [76] V. Cirigliano, W. Dekens, J. de Vries, K. Fuyuto, E. Mereghetti and R. Ruiz, “Leptonic anomalous magnetic moments in  $\nu$ SMEFT,” [arXiv:2105.11462 [hep-ph]].
- [77] L. Allwicher, L. Di Luzio, M. Fedele, F. Mescia and M. Nardecchia, “What is the scale of new physics behind the muon  $g - 2$ ?,” [arXiv:2105.13981 [hep-ph]].
- [78] L. B. Jia, “Dark leptophilic scalar with the updated muon  $g - 2$  anomaly,” [arXiv:2105.13805 [hep-ph]].
- [79] B. De, D. Das, M. Mitra and N. Sahoo, “Magnetic Moments of Leptons, Charged Lepton Flavor Violations and Dark Matter Phenomenology of a Minimal Radiative Dirac Neutrino Mass Model,” [arXiv:2106.00979 [hep-ph]].
- [80] A. Dey, J. Lahiri and B. Mukhopadhyaya, “Muon  $g-2$  and a type-X two Higgs doublet scenario: some studies in high-scale validity,” [arXiv:2106.01449 [hep-ph]].
- [81] N. Arkani-Hamed and K. Harigaya, “Naturalness and the muon magnetic moment,” [arXiv:2106.01373 [hep-ph]].
- [82] K. S. Jeong, J. Kawamura and C. B. Park, “Mixed modulus and anomaly mediation in light of the muon  $g - 2$  anomaly,” [arXiv:2106.04238 [hep-ph]].
- [83] Z. Li, G. L. Liu, F. Wang, J. M. Yang and Y. Zhang, “Gluino-SUGRA scenarios in light of FNAL muon  $g-2$  anomaly,” [arXiv:2106.04466 [hep-ph]].
- [84] A. Crivellin and M. Hoferichter, “Consequences of chirally enhanced explanations of  $(g - 2)_\mu$  for  $h \rightarrow \mu\mu$  and  $Z \rightarrow \mu\mu$ ,” [arXiv:2104.03202 [hep-ph]].
- [85] J. Cao, J. Lian, Y. Pan, D. Zhang and P. Zhu, “Improved  $(g - 2)_\mu$  Measurement and Singlino dark matter in the general NMSSM,” [arXiv:2104.03284 [hep-ph]].
- [86] F. Jegerlehner and A. Nyffeler, “The Muon  $g-2$ ,” Phys. Rept. **477** (2009), 1-110 [arXiv:0902.3360 [hep-ph]].
- [87] M. Lindner, M. Platscher and F. S. Queiroz, “A Call for New Physics : The Muon Anomalous Magnetic Moment and Lepton Flavor Violation,” Phys. Rept. **731** (2018), 1-82 [arXiv:1610.06587 [hep-ph]].

- [88] P. Athron, C. Balázs, D. H. Jacob, W. Kotlarski, D. Stöckinger and H. Stöckinger-Kim, “New physics explanations of  $a_\mu$  in light of the FNAL muon  $g - 2$  measurement,” [arXiv:2104.03691 [hep-ph]].
- [89] A. Keshavarzi, K. S. Khaw and T. Yoshioka, “Muon  $g - 2$ : current status,” [arXiv:2106.06723 [hep-ex]].
- [90] J. P. Leveille, “The Second Order Weak Correction to  $(G-2)$  of the Muon in Arbitrary Gauge Models,” Nucl. Phys. B **137**, 63 (1978).
- [91] S. R. Moore, K. Whisnant and B. L. Young, “Second Order Corrections to the Muon Anomalous Magnetic Moment in Alternative Electroweak Models,” Phys. Rev. D **31** (1985), 105
- [92] F. S. Queiroz and W. Shepherd, “New Physics Contributions to the Muon Anomalous Magnetic Moment: A Numerical Code,” Phys. Rev. D **89**, no.9, 095024 (2014) [arXiv:1403.2309 [hep-ph]].
- [93] A. Crivellin, M. Hoferichter and P. Schmidt-Wellenburg, “Combined explanations of  $(g - 2)_{\mu,e}$  and implications for a large muon EDM,” Phys. Rev. D **98**, no.11, 113002 (2018) [arXiv:1807.11484 [hep-ph]].
- [94] G. Passarino and M. J. G. Veltman, “One Loop Corrections for  $e^+ e^-$  Annihilation Into  $\mu^+ \mu^-$  in the Weinberg Model,” Nucl. Phys. B **160** (1979), 151-207
- [95] H. H. Patel, “Package-X: A Mathematica package for the analytic calculation of one-loop integrals,” Comput. Phys. Commun. **197** (2015), 276-290 [arXiv:1503.01469 [hep-ph]].
- [96] H. H. Patel, “Package-X 2.0: A Mathematica package for the analytic calculation of one-loop integrals,” Comput. Phys. Commun. **218** (2017), 66-70 [arXiv:1612.00009 [hep-ph]].
- [97] C. Biggio, M. Bordone, L. Di Luzio and G. Ridolfi, “Massive vectors and loop observables: the  $g - 2$  case,” JHEP **10** (2016), 002 [arXiv:1607.07621 [hep-ph]].
- [98] S. Weinberg and E. Witten, “Limits on Massless Particles,” Phys. Lett. B **96** (1980), 59-62
- [99] S. M. Barr and A. Zee, “Electric Dipole Moment of the Electron and of the Neutron,” Phys. Rev. Lett. **65**, 21-24 (1990) [erratum: Phys. Rev. Lett. **65**, 2920 (1990)]
- [100] D. Chang, W. F. Chang, C. H. Chou and W. Y. Keung, “Large two loop contributions to  $g-2$  from a generic pseudoscalar boson,” Phys. Rev. D **63**, 091301 (2001) [arXiv:hep-ph/0009292 [hep-ph]].
- [101] W. J. Marciano, A. Masiero, P. Paradisi and M. Passera, “Contributions of axionlike particles to lepton dipole moments,” Phys. Rev. D **94**, no.11, 115033 (2016) [arXiv:1607.01022 [hep-ph]].
- [102] M. Bauer, M. Neubert, S. Renner, M. Schnubel and A. Thamm, “Axionlike Particles, Lepton-Flavor Violation, and a New Explanation of  $a_\mu$  and  $a_e$ ,” Phys. Rev. Lett. **124**, no.21, 211803 (2020) [arXiv:1908.00008 [hep-ph]].

- [103] C. Cornella, P. Paradisi and O. Sumensari, “Hunting for ALPs with Lepton Flavor Violation,” JHEP **2001**, 158 (2020) [arXiv:1911.06279 [hep-ph]].
- [104] L. Morel, Z. Yao, P. Cladé and S. Guellati-Khélifa, “Determination of the fine-structure constant with an accuracy of 81 parts per trillion,” Nature **588**, no.7836, 61-65 (2020)
- [105] R. H. Parker, C. Yu, W. Zhong, B. Estey and H. Müller, “Measurement of the fine-structure constant as a test of the Standard Model,” Science **360**, 191 (2018) [arXiv:1812.04130 [physics.atom-ph]].
- [106] J. S. Schwinger, “On Quantum electrodynamics and the magnetic moment of the electron,” Phys. Rev. **73**, 416 (1948).
- [107] S. J. Brodsky and J. D. Sullivan, “W Boson Contribution To The Anomalous Magnetic Moment Of The Muon,” Phys. Rev. **156**, 1644 (1967).
- [108] T. Burnett and M. J. Levine, “Intermediate vector boson contribution to the muon’s anomalous magnetic moment,” Phys. Lett. **24B**, 467 (1967).
- [109] R. Jackiw and S. Weinberg, “Weak interaction corrections to the muon magnetic moment and to muonic atom energy levels,” Phys. Rev. D **5** (1972), 2396-2398
- [110] I. Bars and M. Yoshimura, “Muon magnetic moment in a finite theory of weak and electromagnetic interaction,” Phys. Rev. D **6**, 374 (1972).
- [111] K. Fujikawa, B. W. Lee and A. I. Sanda, “Generalized Renormalizable Gauge Formulation of Spontaneously Broken Gauge Theories,” Phys. Rev. D **6**, 2923 (1972).
- [112] G. Altarelli, N. Cabibbo and L. Maiani, “The Drell-Hearn sum rule and the lepton magnetic moment in the Weinberg model of weak and electromagnetic interactions,” Phys. Lett. **40B**, 415 (1972).
- [113] W. A. Bardeen, R. Gastmans and B. E. Lautrup, “Static quantities in Weinberg’s model of weak and electromagnetic interactions,” Nucl. Phys. B **46**, 319 (1972).
- [114] A. Czarnecki, W. J. Marciano and A. Vainshtein, “Refinements in electroweak contributions to the muon anomalous magnetic moment,” Phys. Rev. D **67**, 073006 (2003) [erratum: Phys. Rev. D **73**, 119901 (2006)] [arXiv:hep-ph/0212229 [hep-ph]].
- [115] C. Gnendiger, D. Stöckinger and H. Stöckinger-Kim, “The electroweak contributions to  $(g - 2)_\mu$  after the Higgs boson mass measurement,” Phys. Rev. D **88**, 053005 (2013) [arXiv:1306.5546 [hep-ph]].
- [116] P. Minkowski, “ $\mu \rightarrow e\gamma$  at a Rate of One Out of  $10^9$  Muon Decays?,” Phys. Lett. B **67** (1977), 421-428
- [117] T. Yanagida, “Horizontal Symmetry and Masses of Neutrinos,” Prog. Theor. Phys. **64** (1980), 1103

- [118] M. Gell-Mann, P. Ramond and R. Slansky, “Complex Spinors and Unified Theories,” Conf. Proc. C **790927** (1979), 315-321 [arXiv:1306.4669 [hep-th]].
- [119] S. L. Glashow, “The Future of Elementary Particle Physics,” NATO Sci. Ser. B **61** (1980), 687
- [120] R. N. Mohapatra and G. Senjanovic, “Neutrino Mass and Spontaneous Parity Nonconservation,” Phys. Rev. Lett. **44** (1980), 912
- [121] B. Pontecorvo, “Mesonium and anti-mesonium,” Sov. Phys. JETP **6**, 429 (1957)
- [122] Z. Maki, M. Nakagawa and S. Sakata, “Remarks on the unified model of elementary particles,” Prog. Theor. Phys. **28**, 870 (1962).
- [123] W. Konetschny and W. Kummer, “Nonconservation of Total Lepton Number with Scalar Bosons,” Phys. Lett. B **70** (1977), 433-435
- [124] M. Magg and C. Wetterich, “Neutrino Mass Problem and Gauge Hierarchy,” Phys. Lett. B **94** (1980), 61-64
- [125] J. Schechter and J. W. F. Valle, “Neutrino Masses in  $SU(2) \times U(1)$  Theories,” Phys. Rev. D **22** (1980), 2227
- [126] T. P. Cheng and L. F. Li, “Neutrino Masses, Mixings and Oscillations in  $SU(2) \times U(1)$  Models of Electroweak Interactions,” Phys. Rev. D **22** (1980), 2860
- [127] R. N. Mohapatra and G. Senjanovic, “Neutrino Masses and Mixings in Gauge Models with Spontaneous Parity Violation,” Phys. Rev. D **23** (1981), 165
- [128] G. Lazarides, Q. Shafi and C. Wetterich, “Proton Lifetime and Fermion Masses in an  $SO(10)$  Model,” Nucl. Phys. B **181** (1981), 287-300
- [129] T. Fukuyama, H. Sugiyama and K. Tsumura, “Constraints from muon  $g-2$  and LFV processes in the Higgs Triplet Model,” JHEP **03** (2010), 044 [arXiv:0909.4943 [hep-ph]].
- [130] R. Foot, G. C. Joshi, H. Lew and R. R. Volkas, “Charge quantization in the standard model and some of its extensions,” Mod. Phys. Lett. A **5** (1990), 2721-2732
- [131] R. Foot, “New Physics From Electric Charge Quantization?,” Mod. Phys. Lett. A **6** (1991), 527-530
- [132] X. G. He, G. C. Joshi, H. Lew and R. R. Volkas, “NEW Z-prime PHENOMENOLOGY,” Phys. Rev. D **43** (1991), 22-24
- [133] X. G. He, G. C. Joshi, H. Lew and R. R. Volkas, “Simplest Z-prime model,” Phys. Rev. D **44** (1991), 2118-2132
- [134] A. Bodas, R. Coy and S. J. D. King, “Solving the electron and muon  $g - 2$  anomalies in  $Z'$  models,” Eur. Phys. J. C **81**, no.12, 1065 (2021) [arXiv:2102.07781 [hep-ph]].

- [135] I. Doršner, S. Fajfer, A. Greljo, J. F. Kamenik and N. Košnik, “Physics of leptoquarks in precision experiments and at particle colliders,” *Phys. Rept.* **641** (2016), 1-68 [arXiv:1603.04993 [hep-ph]].
- [136] E. Coluccio Leskow, G. D’Ambrosio, A. Crivellin and D. Müller, “ $(g - 2)\mu$ , lepton flavor violation, and  $Z$  decays with leptoquarks: Correlations and future prospects,” *Phys. Rev. D* **95**, no.5, 055018 (2017) [arXiv:1612.06858 [hep-ph]].
- [137] A. Crivellin, D. Mueller and F. Saturnino, “Correlating  $h \rightarrow \mu^+ \mu^-$  to the Anomalous Magnetic Moment of the Muon via Leptoquarks,” [arXiv:2008.02643 [hep-ph]].