NORM INFLATION WITH INFINITE LOSS OF REGULARITY AT GENERAL INITIAL DATA FOR NONLINEAR WAVE EQUATIONS IN WIENER AMALGAM AND FOURIER AMALGAM SPACES

DIVYANG G. BHIMANI AND SAIKATUL HAQUE

ABSTRACT. We study the strong ill-posedness (norm inflation with infinite loss of regularity) for the nonlinear wave equation at every initial data in Wiener amalgam and Fourier amalgam spaces with negative regularity. In particular these spaces contain Fourier-Lebesgue, Sobolev and some modulation spaces. The equations are posed on \mathbb{R}^d and on torus \mathbb{T}^d and involve a smooth power nonlinearity. Our results are sharp with respect to well-posedness results of Bényi and Okoudjou (2009) and Cordero and Nicola (2009) in the Wiener amalgam and modulation space cases. In particular, we also complement norm inflation result of Christ, Colliander and Tao (2003) and Forlano and Okamoto (2020) by establishing infinite loss of regularity in the aforesaid spaces.

1. INTRODUCTION

We study strong ill-posedness for nonlinear wave (NLW) equations of the form

$$\begin{cases} \hat{c}_t^2 u - \Delta u = \pm u^{\rho}(\overline{u})^{\sigma - \rho} \\ (u(0, \cdot), u_t(0, \cdot)) = (u_0, u_1) \end{cases}$$
(1.1)

 $(t,x) \in \mathbb{R} \times \mathcal{M}$, where $\mathcal{M} = \mathbb{R}^d$ or \mathbb{T}^d , $\sigma \in \mathbb{N}$, $\rho \in \mathbb{N} \cup \{0\}$ with $\sigma \ge \max(\rho, 2)$, in Fourier amalgam and Wiener amalgam spaces. We recall:

Definition 1.1 (Hadamard's well/ill posedness).

- The Cauchy problem (1.1) is called locally well-posed from $\mathcal{X} = X_0 \times X_1$ to Y if for every bounded set $B \subset \mathcal{X}$, there exist T > 0 and a Banach space $X_T \hookrightarrow C([0, T], Y)$ such that (i) for all $\vec{u}_0 := (u_0, u_1) \in B$, (1.1) has a unique solution $u \in X_T$ with $(u(0, \cdot), \partial_t u(0, \cdot)) = \vec{u}_0$ (ii) the solution map $\vec{u}_0 \mapsto u$ is continuous from $(B, \|\cdot\|_{\mathcal{X}})$ to C([0, T], Y). In particular, when $\mathcal{X} = Y_s \times Y_{s-1}$ (in our context, $Y_s = \hat{w}_s^{p,q}$ or $W_s^{2,q}$ to be defined below) and $Y = Y_s$, we say (1.1) is locally well-posed in Y_s .
- The Cauchy problem (1.1) is *ill-posed* if the solution map is not continuous.
- We say norm inflation (NI) occurs at $\vec{u}_0 \in \mathcal{X}$ in \mathcal{X} (in short $NI_{\mathcal{X}}(\vec{u}_0)$) if given $\epsilon > 0$, there exist $\vec{v}_0 \in \mathcal{X}, 0 < t < \epsilon$ with $\|\vec{u}_0 - \vec{v}_0\|_{\mathcal{X}} < \epsilon$ such that for the solution v to (1.1) corresponding to the data \vec{v}_0 one has $\|v(t)\|_{X_0} > \epsilon^{-1}$. If $\mathcal{X} = \mathcal{X}_s = Y_s \times Y_{s-1}$, we write $NI_s(\vec{u}_0)$ for $NI_{\mathcal{X}}(\vec{u}_0)$ if there is no confusion.
- If $NI_{\mathcal{X}}(0)$ occurs, we say (mere) norm inflation (at zero) occurs to (1.1) in \mathcal{X} .
- We say NI occurs with *infinite loss of regularity* at \vec{u}_0 in \mathcal{X}_s if for any given $\theta \in \mathbb{R}, \epsilon > 0$, there exist $\vec{v}_0 \in \mathcal{X}_s, 0 < t < \epsilon$ with $\|\vec{u}_0 \vec{v}_0\|_{\mathcal{X}_s} < \epsilon$ such that for the solution v to (1.1) corresponding to the data \vec{v}_0 one has $\|v(t)\|_{X_{\theta}} > \epsilon^{-1}$.

²⁰¹⁰ Mathematics Subject Classification. 35L05, 42B35 (primary), 35B30 (secondary).

Key words and phrases. Nonlinear wave equations; Norm inflation (strong ill-posedness); Wiener amalgam spaces, Fourier amalgam spaces, Fourier-Lebesgue spaces; modulation spaces,

• If the conditions in the preceding definition occurs for θ on a proper subset of \mathbb{R} , we say NI occurs with *finite loss of regularity* at \vec{u}_0 in \mathcal{X}_s .

Recently in [15], Oh and Forlano have introduced Fourier amalgam spaces $\widehat{w}_s^{p,q}(\mathcal{M})$ (with $1 \leq p, q \leq \infty, s \in \mathbb{R}$):

$$\widehat{w}_{s}^{p,q}(\mathcal{M}) = \left\{ f \in \mathcal{S}'(\mathcal{M}) : \|f\|_{\widehat{w}_{s}^{p,q}} = \left\| \|\chi_{n+Q_{1}}(\xi)\mathcal{F}f(\xi)\|_{L_{\xi}^{p}(\widehat{\mathcal{M}})} \langle n \rangle^{s} \right\|_{\ell_{n}^{q}(\mathbb{Z}^{d})} < \infty \right\},$$

where $Q_1 = (-\frac{1}{2}, \frac{1}{2}]^d$, \mathcal{F} denotes the Fourier transform and $\langle \cdot \rangle^s = (1 + |\cdot|^2)^{s/2}$. Here $\widehat{\mathcal{M}}$ denotes the Pontryagin dual of \mathcal{M} , i.e. $\widehat{\mathcal{M}} = \mathbb{R}^d$ if $\mathcal{M} = \mathbb{R}^d$ and $\widehat{\mathcal{M}} = \mathbb{Z}^d$ if $\mathcal{M} = \mathbb{T}^d$. See [27, Part II], among others, for details on space $\mathcal{S}'(\mathcal{M})$. The Fourier-Lebesgue spaces $\mathcal{F}L_s^q(\mathcal{M})$ is defined by

$$\mathcal{F}L^q_s(\mathcal{M}) = \left\{ f \in \mathcal{S}'(\mathcal{M}) : \mathcal{F}f \langle \cdot \rangle^s \in L^q(\widehat{\mathcal{M}}) \right\}$$

On the other hand, the modulation $M_s^{p,q}(\mathcal{M})$ and Wiener amalgam $W_s^{p,q}(\mathcal{M})$ spaces were introduced by Feichtinger in early 1980's in [14]. To recall their definitions, let $\rho \in \mathcal{S}(\mathbb{R}^d)$, $\rho : \mathbb{R}^d \to [0,1]$ be a smooth function satisfying $\rho(\xi) = 1$ if $|\xi|_{\infty} \leq \frac{1}{2}$ and $\rho(\xi) = 0$ if $|\xi|_{\infty} \geq 1$. Let ρ_n be a translation of ρ , that is, $\rho_n(\xi) = \rho(\xi - n), n \in \mathbb{Z}^d$ and denote $\sigma_n(\xi) = \frac{\rho_n(\xi)}{\sum_{\ell \in \mathbb{Z}^d} \rho_\ell(\xi)}, n \in \mathbb{Z}^d$. Then the frequency-uniform decomposition operators can be defined by

$$\Box_n = \mathcal{F}^{-1} \sigma_n \mathcal{F}.$$

Now the modulation $M_s^{p,q}(\mathcal{M})$ and and Wiener amalgam spaces $W_s^{p,q}(\mathcal{M})$, (with $1 \leq p, q \leq \infty, s \in \mathbb{R}$) are defined by the norms:

$$\|f\|_{M^{p,q}_s(\mathcal{M})} = \left\| \|\Box_n f\|_{L^p_x(\mathcal{M})} \langle n \rangle^s \right\|_{\ell^q_n(\mathbb{Z}^d)} \quad \text{and} \quad \|f\|_{W^{p,q}_s(\mathcal{M})} = \left\| \|\Box_n f \cdot \langle n \rangle^s \|_{\ell^q_n(\mathbb{Z}^d)} \right\|_{L^p_x(\mathcal{M})}$$

See Remark 1.6 for equivalent characterization via short-time Fourier transform (STFT) of these spaces. It is known that

$$\widehat{w}_{s}^{p,q}(\mathcal{M}) = \begin{cases} \mathcal{F}L_{s}^{q}(\mathcal{M}) & \text{if } p = q \\ M_{s}^{2,q}(\mathcal{M}) & \text{if } p = 2 \\ H^{s}(\mathcal{M}) = M_{s}^{2,2}(\mathcal{M}) = W_{s}^{2,2}(\mathcal{M}) & \text{if } p = q = 2 \\ \mathcal{F}L_{s}^{q}(\mathcal{M}) = M_{s}^{p,q}(\mathcal{M}) = W_{s}^{p,q}(\mathcal{M}) & \text{if } \mathcal{M} = \mathbb{T}^{d}. \end{cases}$$

(See [25, Section 5] and [17, Proposition 11.3.1] for details.) In last two decades, modulation $M_s^{p,q}$ and Wiener amalgam $W_s^{p,q}$ spaces have been extensively studied in PDEs, see e.g. [2, 3, 5–8, 13, 24, 26, 29–31]. In particular, we have at least good local well-posedness theory for NLW (1.1) in these spaces. We summarize them in the following:

Theorem A (well-posedness). Let $1 \leq p, q \leq \infty, q' > (\sigma - 1)d$ where q' is the Hölder conjugate of q, and let $Y_s = W_s^{p,q}(\mathbb{R}^d)$ or $M_s^{p,q}(\mathbb{R}^d)$ or $\mathcal{F}L^1(\mathbb{R}^d)$.

- (1) [3,4,8,13,16,24] NLW (1.1) is locally well-posed in Y_s for $s \ge 0$.
- (2) [22, 28] NLW (1.1) is locally well-posed in $H^s(\mathbb{R}^d)$ for $s \ge s(\sigma, d), d \ge 2$, some $s(\sigma, d) > 0$.

To the best of authors knowledge there is no local well-posedness results for (1.1) for in $\hat{w}_s^{p,q}$ spaces (except the cases p = 2 or p = q = 1, s = 0). On the other hand, there is extensive literature on ill-posedness for (1.1). We summarize some of them which are most suitable in our context.

Theorem B (ill-posedness).

- (1) [12, Theorem 6] Norm inflation at zero occurs for (1.1) with nonlinearity $\pm |u|^2 u$ in $H^s(\mathbb{R})$ for -1/2 < s < 0.
- (2) [16, Theorem 1.7] NI at general initial data for (1.1) with nonlinearity $\pm u^{\sigma}$ (the case $\rho = \sigma$ in our setting) occurs in $\widehat{w}_s^{p,q}(\mathbb{R}^d)$ $(1 \leq p, q \leq \infty)$ for s < 0.
- (3) [20] NI with finite loss of regularity for (1.1) with nonlinearity $-u^{\sigma}(\sigma > 3)$ in $H^{s}(\mathbb{R}^{d})$ with some positive s (precisely for $1 < s < d/2 2/(\sigma 1), d \ge 3$).

The aim of this note is to complement positive results (Theorem A) by establishing strong ill-posedness for (1.1) in these spaces with negative regularity s < 0. We also complement ill-posedness results (Theorem B) by exhibiting infinite loss of regularity at general initial data. We now state our main result.

Theorem 1.2. Assume that $1 \le p, q \le \infty, s < 0$ and let

$$X_s^{p,q}(\mathcal{M}) = \begin{cases} \widehat{w}_s^{p,q}(\mathbb{R}^d) \text{ or } W_s^{2,q}(\mathbb{R}^d) \text{ for } \mathcal{M} = \mathbb{R}^d \\ \mathcal{F}L_s^q(\mathbb{T}^d) \text{ for } \mathcal{M} = \mathbb{T}^d \end{cases} \text{ and } \mathcal{X}_s^{p,q}(\mathcal{M}) = X_s^{p,q}(\mathcal{M}) \times X_{s-1}^{p,q}(\mathcal{M}).$$

Then norm inflation with infinite loss of regularity occurs to (1.1) at each element in $\mathcal{X}_s^{p,q}(\mathcal{M})$: For any $\theta \in \mathbb{R}$, $\varepsilon > 0$ and $\vec{u}_0 \in \mathcal{X}_s^{p,q}(\mathcal{M})$ there exist $\vec{u}_{0,\varepsilon} \in \mathcal{X}_s^{p,q}(\mathcal{M})$ and T > 0 satisfying

$$\|\vec{u}_0 - \vec{u}_{0,\varepsilon}\|_{\mathcal{X}^{p,q}_s} < \varepsilon, \quad 0 < T < \varepsilon$$

such that the corresponding smooth solution u_{ε} to (1.1) with data $\vec{u}_{0,\varepsilon}$ exists on [0,T] and

$$\|u_{\varepsilon}(T)\|_{X^{p,q}_{\theta}} > \frac{1}{\varepsilon}$$

In particular, for any T > 0, the solution map $X_s^{p,q} \times X_{s-1}^{p,q} \ni (u_0, u_1) \mapsto u \in C([0,T], X_{\theta}^{p,q})$ for (1.1) is discontinuous everywhere in $X_s^{p,q} \times X_{s-1}^{p,q}$ for all $\theta \in \mathbb{R}$.

For $q' > (\sigma - 1)d$, Theorem 1.2 is sharp in the sense that (1.1) is strongly ill-posed in $W_s^{2,q}(\mathbb{R}^d)$, $M_s^{2,q}(\mathbb{R}^d)$ for s < 0 while by Theorem A it is locally well-posed for $s \ge 0$. In fact, even *mere* ill-posedness in $W_s^{2,q}(\mathbb{R}^d)$ is completely new as it is not cover by Theorem B. The particular case $X_s^{p,q} = \hat{w}_s^{p,q}$ of Theorem 1.2 recover Theorem B (2) and further reveal that even worse situation occurs as the infinite loss of regularity (everywhere in $\hat{w}_s^{p,q} \times \hat{w}_{s-1}^{p,q}$) is present in all dimensions. Theorem 1.2 also complements Theorem B (3) by taking $\rho = \sigma$ in (1.1) by establishing infinite loss of regularity.

The circle of ideas to establish ill-posedness, via Fourier analytic approach, is originated from the abstract argument [1, Theorem 2 and Section 3] of Bejenaru and Tao in the context of quadratic nonlinear Schrödinger equation (NLS). In fact, the idea is to rewrite the solution of (1.1) as a power series expansion (Lemma 2.7) in terms of Picard iterations. It is then sufficient to establish the discontinuity for one Picard iterate to get the discontinuity of solution map at zero. This method is further developed by Iwabuchi and Ogawa [18] to establish stronger phenomena of NI for NLS. Later Kishimoto [19] establish NI for NLS with more general nonlinearity¹ on the special domain $\mathbb{R}^{d_1} \times \mathbb{T}^{d_2}$, $d = d_1 + d_2$. The idea is to show that one term in the series exhibits instability and dominates all the other terms (Lemmata 3.3, 3.6 and 3.2, 3.5) after adding a perturbation $\vec{\phi}_{0,N}$ to the data \vec{u}_0 , see (3.3). We shall notice that the existence time T > 0 is allowed to shrink for the purpose of establishing norm inflation while in [1] it is fixed and uniform with respect to the initial data. See [19, Section 2]

¹Precisely this nonlinarity: $\sum_{j=1}^{n} \nu_j u^{\rho_j}(\bar{u})^{\sigma_j - \rho_j}$ where $\nu_j \in \mathbb{C}, \sigma_j \in \mathbb{N}, \rho_j \in \mathbb{N} \cup \{0\}$ with $\sigma_j \ge \max(\rho_j, 2)$.

for detail discussion on the approach. On the other hand, Oh [23] use power series expansion indexed by trees to establish NI at general initial data for cubic NLS on \mathcal{M} . Forlano and Okomoto [16] use this Fourier analytic approach, following presentation from [23], for NLW (1.1) (in the particular case $\rho = \sigma$) to establish norm inflation at general initial data.

In addition to these ideas, in the present paper, we fix the size of the support of perturbation $\vec{\phi}_{0,N}$ in (3.3) of the initial data on the frequency side. This simplifies our analysis. Moreover, we choose this perturbation to be real valued and symmetric on the Fourier side i.e. $\mathcal{F}\vec{\phi}_{0,N}(-\cdot) = \mathcal{F}\vec{\phi}_{0,N}$. This enables us to consider more general nonlinearity $u^{\rho}(\bar{u})^{\sigma-\rho}$ as compared to u^{σ} in [16]. In order to get infinite loss of regularity (specifically, while showing the $X^{p,q}_{\theta}$ -norm of the solution is arbitrarily large for all θ), we restrict the Fourier transform of the solution at a particular frequency (say at $n = e_1$). This will allow us to compare two discrete Lebesgue norms with different weights i.e.

$$\|\langle \cdot \rangle^{\theta} f\|_{\ell^{q}(n=e_{1})} = 2^{(s-\theta)/2} \|\langle \cdot \rangle^{s} f\|_{\ell^{q}(n=e_{1})} \quad \text{for all } \theta \in \mathbb{R}$$

which eventually leads to infinite loss of regularity. In addition to this, it also allows us to use Plancherel Theorem to achieve the lower bound estimate for the second nontrivial Picard iterate in $W_s^{2,q}$ -norm, see Lemma 3.6. We note that a similar idea was used by Kishimoto in [19, Appendix A] in the context of NLS to achieve norm inflation at zero with infinite loss of regularity in $H^s(\mathcal{M})$. Kishimoto [19] used modulation space $M^{2,1}$ to justify the convergence power series expansion while in the present paper we use Wiener algebra $\mathcal{F}L^1$ as in [16,23]. We employ these ideas together with the refinement of ideas used in [5,16,19,23] and properties of $\widehat{w}_s^{p,q}, W_s^{p,q}$ to prove Theorem 1.2.

Given $\lambda > 0$, if u solves (1.1), then scaling $u_{\lambda}(t, x) := \lambda^{\frac{2}{\sigma-1}} u(\lambda t, \lambda x)$ also solves (1.1) with rescaled initial data $\lambda^{\frac{2}{\sigma-1}}(u_0(\lambda x), u_1(\lambda x))$. This scaling leaves the homogeneous Sobolev \dot{H}^s space invariant when $s = s_c = d/2 - 2/(\sigma - 1)$. The ill-posedness below the scaling critical regularity s_c has been studied in [9, 20, 21]. We note that NI with finite loss of regularity at zero initial data for (1.1) is initiated by G. Lebeau in $[20]^2$, see Theorem B (3). While we initiate, to the best of the authors' knowledge, NI with infinite loss of regularity at general initial data for (1.1) in the present paper. Our study of infinite loss of regularity for (1.1)is inspired form known NLS case in [10, 11, 19] while the inspiration of NI at general initial data comes from [16, 23]. We also note that the details of proofs Lebeau in [20] does not seem to work for negative regularity s < 0. In [5, 10, 11], Carles and his collaborators have used geometric optics approach to establish infinite loss of regularity for NLS. Recently in [7], Bhimani and Haque have used the Fourier analytic approach to establish infinite loss of regularity for fractional Hartree and cubic NLS for some negative regularity (i.e for all $s < -\epsilon < 0$ for some $\epsilon > 0$). On the other hand Theorem 1.2 establishes infinite loss of regularity for NLW (1.1) for all s < 0. This is in strict contrast as compared to known NLS case results proved in [5, 7, 10] and Theorem 1.2 thus reveal new phenomena for NLW (1.1). We conclude our discussion with the following remarks.

Remark 1.3. In order to get the upper bound for each Picard iterate we must have $s \leq 0$ (Lemma 3.2) while we get the lower bound for one dominating Picard iterate for all $s \in \mathbb{R}$ (Lemma 3.3). The restriction s < 0 will be required at the final stage in the proof of Theorem 1.2. Our approach thus may not work for $s \geq 0$.

 $^{^{2}}$ This result is stronger than finite loss of regularity at zero as it is obatied with a single datum instead of a sequence of initial data converging to zero.

Remark 1.4. In [5, Theorem 1.6], Bhimani and Carles have established infinite loss of regularity, via geometric optics approach, for NLS in $M_s^{p,q}$ for all $1 \leq p, q \leq \infty$ and for some s < 0. This somewhat indicates that restriction on p = 2 in $M_s^{p,q}, W_s^{p,q}$ in Theorem 1.2 is just due to our approach and we believe that Theorem 1.2 is also true for any $1 \leq p \leq \infty$ in $M_s^{p,q}, W_s^{p,q}$. In fact, by taking p = 2, we can work on the Fourier side due to Plancherel theorem. This makes our analysis somewhat simple. On the other hand, for $p \neq 2$, we do not know how to deal with frequency-uniform decomposition operators \Box_n . Also taking Theorem A into account, the case s = 0 with $q' \leq (\sigma - 1)d$ in Theorem 1.2 remains open. We plan to address these issues in our future work.

Remark 1.5. Our method proof should be applicable to (1.1) on special domain $Z = \mathbb{R}^{d_1} \times \mathbb{T}^{d_2}$ $(d_1, d_2 \in \mathbb{N} \cup \{0\})$ with more general type non-linearity $\sum_{j=1}^n \nu_j u^{\rho_j}(\bar{u})^{\sigma_j - \rho_j}$ where $\nu_j \in \mathbb{C}$, $\sigma_j \in \mathbb{N}$, $\rho_j \in \mathbb{N} \cup \{0\}$ with $\sigma_j \ge \max(\rho_j, 2)$; as in the NLS case [19].

Remark 1.6. The STFT of a $f \in \mathcal{S}'(\mathcal{M})$ with respect to a window function $0 \neq g \in \mathcal{S}(\mathcal{M})$ is defined by

$$V_g f(x, y) = \int_{\mathcal{M}} f(t) \overline{T_x g(t)} e^{-2\pi i y \cdot t} dt, \ (x, y) \in \mathcal{M} \times \widehat{\mathcal{M}}$$

whenever the integral exists. Here, $T_x g(t) = g(tx^{-1})$ is the translation operator on \mathcal{M} . It is known [29, Proposition 2.1], [14] that

$$\|f\|_{M^{p,q}_s} \approx \left\| \|V_g f(x,y)\|_{L^p(\mathcal{M})} \langle y \rangle^s \right\|_{L^q(\widehat{\mathcal{M}})} \quad \text{and} \quad \|f\|_{W^{p,q}_s(\mathcal{M})} \approx \left\| \|V_g f(x,y) \langle y \rangle^s \|_{L^q(\widehat{\mathcal{M}})} \right\|_{L^p(\mathcal{M})}$$

The definition of the modulation space is independent of the choice of the particular window function, see [17, Proposition 11.3.2(c)].

2. Key Lemmas

For $u_0, u_1 \in \mathcal{S}(\mathbb{R}^d)$, (or $\in C^{\infty}(\mathbb{T}^d)$ in torus case) the wave propagator $S(t)(u_0, u_1)$ is given by $S(t)(u_0, u_1) = \cos(t|\nabla|)u_0 + \frac{\sin(t|\nabla|)}{|\nabla|}u_1$, in other words

$$\mathcal{F}S(t)(u_0, u_1)(\xi) = \cos(t|\xi|)\mathcal{F}u_0(\xi) + \frac{\sin(t|\xi|)}{|\xi|}\mathcal{F}u_1(\xi), \quad \xi \in \widehat{\mathcal{M}}, \ t \in \mathbb{R}.$$

Let $\mathcal{X}_s^{p,q}(\mathcal{M}) = \widehat{w}_s^{p,q}(\mathcal{M}) \times \widehat{w}_{s-1}^{p,q}(\mathcal{M})$ with the norm

$$\|\overrightarrow{u_0}\|_{\mathcal{X}^{p,q}_s} = \|u_0\|_{\widehat{w}^{p,q}_s} + \sqrt{2} \|u_1\|_{\widehat{w}^{p,q}_{s-1}} \quad (1 \le p,q \le \infty, s \in \mathbb{R}).$$

When s = 0, we write $\mathcal{X}^{p,q}(\mathcal{M}) = \mathcal{X}^{p,q}_0(\mathcal{M})$.

Lemma 2.1. Let $\mathcal{X}_s^{p,q}(\mathcal{M})$ be defined as above and $0 \leq t \leq 1$. Then $||S(t)(\vec{u}_0)||_{\hat{w}_s^{p,q}} \leq ||\vec{u}_0||_{\mathcal{X}_s^{p,q}}$. *Proof.* Since $|\cos(t|\xi|)| \leq 1$ and for $0 \leq t \leq 1$,

$$\frac{|\sin(t|\xi|)|}{|\xi|} (1+|\xi|^2)^{1/2} \leqslant \begin{cases} t(1+|\xi|^2)^{1/2} \leqslant \sqrt{2} & \text{if } |\xi| \leqslant 1\\ \frac{(1+|\xi|^2)^{1/2}}{|\xi|} \leqslant \sqrt{2} & \text{if } |\xi| \ge 1 \end{cases}$$

the result follows from the definition of $\widehat{w}_s^{p,q}(\mathcal{M})$.

For $f_1, \dots, f_{\sigma} \in \mathcal{S}(\mathbb{R}^d)$ (or $\in C^{\infty}(\mathbb{T}^d)$ in torus case), we define the multilinear operator \mathcal{H}_{σ} associated to the nonlinearity in (1.1) as follows

$$\mathcal{H}_{\sigma}(f_1,\ldots,f_{2\sigma+1}) = \prod_{\ell=1}^{\rho} f_{\ell} \prod_{m=\rho+1}^{\sigma} \bar{f}_m$$

When $f_1 = \cdots = f_{\sigma} = f$, we write $\mathcal{H}_{\sigma}(f_1, \ldots, f_{\sigma}) = \mathcal{H}_{\sigma}(f)$. We set

$$\mathcal{N}_{\sigma}(u_1,\ldots,u_{\sigma})(t) = \int_0^t \frac{\sin((t-\tau)|\nabla|)}{|\nabla|} \mathcal{H}_{\sigma}(u_1(\tau),\cdots,u_{\sigma}(\tau))d\tau$$

and write $\mathcal{N}_{\sigma}(u_1, \ldots, u_{\sigma}) = \mathcal{N}_{\sigma}(u)$ for $u = u_1 = \cdots = u_{\sigma}$. Recall that solution of (1.1) satisfies

$$u(t) = S(t)(u_0, u_1) \pm \int_0^t \frac{\sin((t-\tau)|\nabla|)}{|\nabla|} \mathcal{H}_{\sigma}(u(\tau)) d\tau = S(t)(u_0, u_1) \pm \mathcal{N}_{\sigma}(u)(t)$$
(2.1)

Definition 2.2 (Picard iteration). Let us set $S_1[\vec{u}_0](t) = S(t)(\vec{u}_0)$ and

$$S_{k}[\vec{u}_{0}](t) = \sum_{\substack{k_{1},...,k_{\sigma} \ge 1\\k_{1}+\dots+k_{\sigma}=k}} \int_{0}^{t} \frac{\sin((t-\tau)|\nabla|)}{|\nabla|} \mathcal{H}_{\sigma}\left(S_{k_{1}}[\vec{u}_{0}],...,S_{k_{\sigma}}[\vec{u}_{0}]\right)(\tau)d\tau \quad (k \ge 2).$$

Remark 2.3. The empty sums in Definition 2.2 are considered as zeros. In view of this one can see that $S_{(\sigma-1)\ell+2}[\vec{u}_0] = S_{(\sigma-1)\ell+3}[\vec{u}_0] = \cdots = S_{(\sigma-1)(\ell+1)}[\vec{u}_0] = 0$ for all $\ell \in \mathbb{N} \cup \{0\}$.

Lemma 2.4 (Algebra property). The spaces $\widehat{w}_s^{p,q}(\mathcal{M})$ is a pointwise $\mathcal{F}L^1$ -module with norm inequality

$$\|fg\|_{\widehat{w}_s^{p,q}} \leqslant \|f\|_{\mathcal{F}L^1} \|g\|_{\widehat{w}_s^{p,q}} \quad (1 \leqslant p,q \leqslant \infty, s \in \mathbb{R})$$

In particular, $\mathcal{F}L^1$ is an algebra under pointwise multiplication, i.e. $\|fg\|_{\mathcal{F}L^1} \leq \|f\|_{\mathcal{F}L^1} \|g\|_{\mathcal{F}L^1}$. *Proof.* Follows from Young's inequality.

Lemma 2.5 (See [19]). Let $\{b_k\}_{k=1}^{\infty}$ be a sequence of nonnegative real numbers such that

$$b_k \leqslant C \sum_{\substack{k_1, \dots, k_\sigma \ge 1\\k_1 + \dots + k_\sigma = k}} b_{k_1} \cdots b_{k_\sigma} \quad \forall \ k \ge 2.$$

Then we have $b_k \leq b_1 C_0^{k-1}$, for all $k \geq 1$, where $C_0 = \frac{\pi^2}{6} (C\sigma^2)^{1/(\sigma-1)} b_1$.

Lemma 2.6. For $0 \leq t \leq 1$, for $k \geq \sigma$ one has $\|S_k[\vec{u}_0](t)\|_{\hat{w}_s^{p,q}} \leq C^k t^{2\frac{k-1}{\sigma-1}} \|\vec{u}_0\|_{\mathcal{X}_s^{p,q}}^{k-1}$.

Proof. Let $\{a_k\}$ be a sequence of nonnegative real numbers such that

$$a_1 = 1, \quad a_k = \frac{\sigma - 1}{2k - \sigma - 1} \sum_{\substack{k_1, \dots, k_\sigma \ge 1\\k_1 + \dots + k_\sigma = k}} a_{k_1} \cdots a_{k_\sigma} \quad \forall \ k \ge 2$$

where C > 1 to be chosen later. By Lemma 2.5 (2.5), we have $a_k \leq c_0^k$ for some $c_0 = c_0(\sigma) > 0$. In view of this it is enough to prove: $\|S_k[\vec{u}_0](t)\|_{\hat{w}_s^{p,q}} \leq a_k t^{2\frac{k-1}{\sigma-1}} \|\vec{u}_0\|_{\mathcal{X}_s^{p,q}}^{k-1}$. By Definition 2.2 and the fact $|\sin \tau| \leq |\tau|$ together with Lemma 2.4, we have

$$\|S_k[\vec{u}_0](t)\|_{\hat{w}_s^{p,q}} \leqslant \sum_{\substack{k_1,\dots,k_\sigma \ge 1\\k_1+\dots+k_\sigma=k}} \int_0^t |t-\tau| \, \|S_{k_1}[\vec{u}_0](\tau)\|_{\hat{w}_s^{p,q}} \prod_{\ell=2}^\sigma \|S_{k_\ell}[\vec{u}_0](\tau)\|_{\mathcal{F}L^1} \, d\tau.$$

Therefore, by Lemma 2.1, we have

$$\|S_{\sigma}[\vec{u}_{0}](t)\|_{\hat{w}_{s}^{p,q}} \leq t \int_{0}^{t} \|S_{1}[\vec{u}_{0}](\tau)\|_{\hat{w}_{s}^{p,q}} \prod_{\ell=2}^{\sigma} \|S_{1}[\vec{u}_{0}](\tau)\|_{\mathcal{F}L^{1}} d\tau \leq t^{2} \|\vec{u}_{0}\|_{\mathcal{F}L^{1}}^{\sigma-1} \|\vec{u}_{0}\|_{\mathcal{X}_{s}^{p,q}}.$$

Since $a_{\sigma} = 1$, the claim is true for $k = \sigma$. Assume that the claim is true up to the level k - 1. Then

$$\begin{split} \|S_{k}[\vec{u}_{0}](t)\|_{\hat{w}_{s}^{p,q}} &\leq \sum_{\substack{k_{1},\dots,k_{\sigma}\geq 1\\k_{1}+\dots+k_{\sigma}=k}} \|\vec{u}_{0}\|_{\mathcal{F}L^{1}}^{k_{1}-1} \|\vec{u}_{0}\|_{\mathcal{X}_{s}^{p,q}} \prod_{\ell=2}^{\sigma} \|\vec{u}_{0}\|_{\mathcal{F}L^{1}}^{k_{\ell}} a_{k_{1}}\cdots a_{k_{\sigma}} t \int_{0}^{t} \tau^{2\frac{k-\sigma}{\sigma-1}} d\tau \\ &= a_{k} \|\vec{u}_{0}\|_{\mathcal{F}L^{1}}^{k-1} \|\vec{u}_{0}\|_{\mathcal{X}_{s}^{p,q}} t^{2\frac{k-1}{\sigma-1}}. \end{split}$$

Hence, the claim is true at the level k. This completes the proof.

Lemma 2.7. If $0 < T \ll \min(1, M^{-\frac{\sigma-1}{2}})$, then for any $\vec{u_0} \in \overrightarrow{\mathcal{FL}}^1$ with $\|\vec{u_0}\|_{\overrightarrow{\mathcal{FL}}^1} \leq M$, there exists a unique solution u to integral equation (2.1) given by

$$u = \sum_{k=1}^{\infty} S_k[\vec{u}_0] = \sum_{\ell=0}^{\infty} S_{2\sigma\ell+1}[\vec{u}_0]$$
(2.2)

which converges absolutely in $C([0,T], \mathcal{F}L^1)$.

Proof. The proof goes in a similar line as the proof of Lemma 2.4 in [16]. Since the nonlinearity is different in our case, we shall briefly present the proof for the convenience of reader. Define

$$\Psi(u)(t) = S(t)[\vec{u}_0] \pm \int_0^t \frac{\sin((t-\tau)|\nabla|)}{|\nabla|} \mathcal{H}_{\sigma}(u(\tau)) d\tau.$$

Let $0 < T \leq 1$. By Lemma 2.1 and following the proof of Lemma 2.6, we obtain

$$\begin{split} \|\Psi(u)\|_{C([0,T],\mathcal{F}L^{1})} &\lesssim \|\vec{u}_{0}\|_{\overrightarrow{\mathcal{F}L}^{1}} + T^{2} \|u\|_{C([0,T],\mathcal{F}L^{1})}^{\sigma}, \\ \|\Psi(u) - \Psi(v)\|_{C([0,T],\mathcal{F}L^{1})} &\lesssim T^{2} \max\left(\|u\|_{C([0,T],\mathcal{F}L^{1})}^{\sigma-1}, \|v\|_{C([0,T],\mathcal{F}L^{1})}^{\sigma-1}\right) \|u - v\|_{C([0,T],\mathcal{F}L^{1})}. \end{split}$$

Then considering the ball $B_{2M}^T = \left\{ \phi \in C([0,T], \mathcal{F}L^1) : \|\phi\|_{C([0,T], \mathcal{F}L^1)} \leq 2M \right\}$ with $T^2, T^2 M^{\sigma-1} \ll 1$, we find a unique fixed point of Ψ in B_{2M}^T . Hence, the solution to (2.1). This proves the existence of unique solution. It is not hard to show that this solution is given by (2.2) (see for e.g. [16, Lemma 2.4]). This completes the proof.

3. The proof of Theorem 1.2

We first prove NI with infinite loss of regularity at general data in $\overrightarrow{\mathcal{FL}^{1}}(\mathcal{M}) \cap \mathcal{X}_{s}^{p,q}(\mathcal{M})$. Subsequently, for general data in $\mathcal{X}_{s}^{p,q}(\mathcal{M})$ we use the density of $\overrightarrow{\mathcal{FL}^{1}}(\mathcal{M}) \cap \mathcal{X}_{s}^{p,q}(\mathcal{M})$ in $\mathcal{X}_{s}^{p,q}(\mathcal{M})$ (s < 0). So let us begin with $\vec{u}_{0} \in \overrightarrow{\mathcal{FL}^{1}}(\mathcal{M}) \cap \mathcal{X}_{s}^{p,q}(\mathcal{M})$. Let $N, R \gg 1, Q = [-1, 1]^{d}$ and $e_{1} = (1, 0, ..., 0) \in \mathbb{R}^{d}$. Set $\Sigma_{N} = \{\pm Ne_{1}, \pm 2Ne_{1}\}$ and

$$\mathcal{F}\phi_{0,N} = R\chi_{\Omega_N}$$
 with $\Omega_N = \bigcup_{\eta \in \Sigma_N} (\eta + Q).$ (3.1)

Note that $\mathcal{F}\phi_{0,N} = \mathcal{F}\phi_{0,N}(-\cdot)$ by the symmetry of the set Q and Σ and

$$\|\phi_{0,N}\|_{\hat{w}_{s}^{p,q}} = R\|\|\chi_{n+Q_{1}}\hat{\phi}_{0,N}\|_{L^{p}}\langle n\rangle^{s}\|_{\ell^{q}} = R\left(\sum_{n\in\mathbb{Z}^{d}}|(n+Q_{1})\cap\Omega_{N}|^{\frac{q}{p}}\langle n\rangle^{qs}\right)^{1/q} \sim RN^{s}.$$
 (3.2)

We take

$$\vec{\phi}_{0,N} = (\phi_{0,N}, 0) \text{ and } \vec{u}_{0,N} = \vec{u}_0 + \vec{\phi}_{0,N}.$$
 (3.3)

Lemma 3.1 (See Lemma 3.6 in [19]). Let $\vec{\phi}_{0,N}$ be given by (3.3). Then there exists C > 0 such that for all $k \in \mathbb{N}$, we have

$$\left|\operatorname{supp} \mathcal{F}S_k[\vec{\phi}_{0,N}](t)\right| \leq C^k, \quad \forall t \geq 0.$$

Proof. Note that $\operatorname{supp} \mathcal{F}S_1[\vec{\phi}_{0,N}](t) \subset \operatorname{supp} \mathcal{F}u_0$; which is contained in 4 cubes with volumes 2^d . Hence, $\left|\operatorname{supp} \mathcal{F}S_1[\vec{\phi}_{0,N}](t)\right| \leq 2^d 4$, $\forall t \geq 0$. Thus, it is enough to prove the following claim: $\operatorname{supp} \mathcal{F}S_k[\vec{\phi}_{0,N}](t)$ is contained in $4c_d^{k-1}$ number of cubes with volume 2^d . Clearly the claim is true for k = 1. Assume that the claim is true upto k - 1 stage. Then

$$\operatorname{supp} \mathcal{F}S_k[\vec{\phi}_{0,N}](t) \subset \sum_{\substack{k_1,\dots,k_{2\sigma+1} \ge 1\\k_1+\dots+k_{2\sigma+1}=k}} \operatorname{supp} v_{k_j}(t)$$

where $v_{k_{\ell}}$ is either $\mathcal{F}S_{k_{\ell}}[\vec{\phi}_{0,N}]$ or $\mathcal{F}S_{k_{\ell}}[\vec{\phi}_{0,N}]$. Using induction we conclude that the set in RHS is contained in

$$d^{2\sigma} \prod_{\substack{k_1, \dots, k_{2\sigma+1} \ge 1 \\ k_1 + \dots + k_{2\sigma+1} = k}} 4c_d^{k_j - 1} = 4^{2\sigma+1} d^{2\sigma} c_d^{k-2\sigma-1} = 4(4d)^{2\sigma} c_d^{k-2\sigma-1}$$

number of cubes with volume 2^d . Set $c_d = 4d$ and $C = c_d = 4d$ to conclude.

3.1. Estimates in $\widehat{w}_s^{p,q}$. The next result is the analogue of [19, Lemma 3.7].

Lemma 3.2. Let $\vec{u}_{0,N}$ be given by (3.3), $s \leq 0, 1 \leq p, q \leq \infty$ and $0 \leq t \leq 1$. Then there exists C > 0 independent of R, N, t such that followings hold:

 $(1) \|\vec{u}_{0,N} - \vec{u}_0\|_{\mathcal{X}^{p,q}_s} \lesssim RN^s$ $(2) \|S_1[\vec{u}_{0,N}](t)\|_{\hat{w}^{p,q}_s} \lesssim 1 + RN^s$ $(3) \|S_{2\sigma+1}[\vec{u}_{0,N}](t) - S_{2\sigma+1}[\vec{\phi}_{0,N}](t)\|_{\hat{w}^{p,q}_s} \lesssim t^2 R^{\sigma-1}$ $(4) \|S_k[\vec{u}_{0,N}](t)\|_{\hat{w}^{p,q}_s} \lesssim C^k R^k t^{2\frac{k-1}{\sigma-1}}.$

Proof. (1) follows from (3.2). By Lemma 2.1 and (3.2) we have $||S_1[\vec{\phi}_{0,N}](t)||_{\hat{w}_s^{p,q}} \leq RN^s$. Then (2) follows by triangular inequality. By Lemma 2.6 (with $p = q = \infty$, s = 0) and (3.2), we obtain

$$\begin{aligned} \|S_{k}[\vec{\phi}_{0,N}](t)\|_{\hat{w}_{s}^{p,q}} &\leq \sup_{\xi \in \widehat{\mathcal{M}}} |\mathcal{F}S_{k}[\vec{\phi}_{0,N}](t,\xi)| |\operatorname{supp} \mathcal{F}S_{k}[\vec{\phi}_{0,N}](t)|_{\mu_{\widehat{\mathcal{M}}}}^{1/p} \|\langle n \rangle^{s}\|_{\ell^{q}((n+Q_{1})\cap\operatorname{supp} \mathcal{F}S_{k}[\vec{u}_{0}](t))} \\ &\leq C^{k} R^{k} t^{2\frac{k-1}{\sigma-1}} \|\langle n \rangle^{s}\|_{\ell^{q}((n+Q_{1})\cap\operatorname{supp} \mathcal{F}S_{k}[\vec{u}_{0}](t))} \cdot \end{aligned}$$

where $|A|_{\mu_{\widehat{\mathcal{M}}}}$ denotes the $\widehat{\mathcal{M}}$ -measure of the set A. Since $s \leq 0$, for any bounded set $D \subset \mathbb{Z}^d$, we have $\|\langle n \rangle^s \|_{\ell^q(n \in D)} \leq \|\langle n \rangle^s \|_{\ell^q(n \in B_D)}$ where $B_D \subset \mathbb{R}^d$ is the minimal ball centred at the origin with $|D| \leq |B_D|$. In view of this and Lemma 3.1, we obtain

$$\langle n \rangle^s \|_{\ell^q(\operatorname{supp} \widehat{S}_k[\vec{u}_0](t))} \leq \| \langle n \rangle^s \|_{\ell^q(\{|n| \leq C^{k/d}\})} \lesssim C^{k/q}.$$

Therefore

$$|S_k[\vec{\phi}_{0,N}](t)||_{\widehat{w}_s^{p,q}} \leqslant C^k R^k t^{2\frac{k-1}{\sigma-1}}.$$
(3.4)

Now observe that

$$I_{k}(t) := S_{k}[\vec{u}_{0,N}](t) - S_{k}[\vec{\phi}_{0,N}](t)$$

$$= \sum_{\substack{k_{1},\dots,k_{\sigma} \geq 1\\k_{1}+\dots+k_{\sigma}=k}} \mathcal{N}(S_{k_{1}}[\vec{u}_{0} + \vec{\phi}_{0,N}], \cdots, S_{k_{\sigma}}[\vec{u}_{0} + \vec{\phi}_{0,N}]) - \mathcal{N}(S_{k_{1}}[\vec{\phi}_{0,N}], \cdots, S_{k_{\sigma}}[\vec{\phi}_{0,N}])$$

$$= \sum_{\substack{k_{1},\dots,k_{\sigma} \geq 1\\k_{1}+\dots+k_{\sigma}=k}} \sum_{(\vec{\psi}_{1},\dots,\vec{\psi}_{\sigma})\in\mathcal{C}} \mathcal{N}(S_{k_{1}}[\vec{\psi}_{1}], \cdots, S_{k_{\sigma}}[\vec{\psi}_{\sigma}])$$

where $\mathcal{C} = \{\vec{u}_0, \vec{\phi}_{0,N}\}^{\sigma} \setminus \{(\vec{\phi}_{0,N}, \cdots, \vec{\phi}_{0,N})\}$. Observe that \mathcal{C} has at least one coordinate as \vec{u}_0 . Using Lemma 2.6 it follows that

$$\begin{split} \|I_{k}(t)\|_{\hat{w}_{s}^{p,q}} &\lesssim \sum_{\substack{k_{1},\dots,k_{\sigma} \geq 1\\k_{1}+\dots+k_{\sigma}=k}} t \int_{0}^{t} \sum_{(\vec{v}_{1},\dots,\vec{v}_{\sigma})\in\mathcal{C}} \|S_{k_{1}}[\vec{v}_{1}]\|_{\hat{w}_{s}^{p,q}} \prod_{j=2}^{\sigma} \|S_{k_{j}}[\vec{v}_{j}]\|_{\mathcal{F}L^{1}} \\ &\leqslant (2^{\sigma}-1)2\|\vec{u}_{0}\|_{\mathcal{X}_{s}^{p,q}} (\|\vec{u}_{0}\|_{\mathcal{F}L^{1}}^{k-1} + \|\vec{\phi}_{0,N}\|_{\mathcal{F}L^{1}}^{k-1}) t \int_{0}^{t} \tau^{2\frac{k-\sigma}{\sigma-1}} d\tau \sum_{\substack{k_{1},\dots,k_{\sigma} \geq 1\\k_{1}+\dots+k_{\sigma}=k}} a_{k_{1}} \cdots a_{k_{\sigma}} \\ &\leqslant 2^{\sigma+2} a_{k} t^{2\frac{k-1}{\sigma-1}} R^{k-1} \|\vec{u}_{0}\|_{\mathcal{X}_{s}^{p,q}} \leqslant C^{k} t^{2\frac{k-1}{\sigma-1}} R^{k-1} \|\vec{u}_{0}\|_{\mathcal{X}_{s}^{p,q}} \end{split}$$

as $R \gg 1$. Note that (3) is the particular case $k = \sigma$ and (4) follows using the above and (3.4).

In the proof of next the lemma we follow the strategy of Proposition 3.4 in [16]. Although Proposition 3.4 in [16] considers a different nonlinearity (u^{σ}) , the symmetry about the origin of the real valued function $\mathcal{F}u_0$ allows us to cover our choice of nonlinearity $(u^{\rho}\bar{u}^{\sigma-\rho})$. We have presented the proof in detail as it will be used in the proof of the similar estimates in $W_s^{2,q}(\mathbb{R}^d)$ spaces (Lemma 3.6).

Lemma 3.3. Let $\vec{\phi}_{0,N}$ be given by (3.3), $s \in \mathbb{R}$, $1 \leq p, q \leq \infty$, $N^{-1/2} < T \ll 1$. Then we have

$$\|S_{\sigma}[\vec{\phi}_{0,N}](T)\|_{\hat{w}_{s}^{p,q}} \ge \left\| \left\| \chi_{n+Q_{1}}(\xi)\mathcal{F}S_{\sigma}[\vec{\phi}_{0,N}](T)(\xi) \right\|_{L^{p}_{\xi}} \langle n \rangle^{s} \right\|_{\ell^{q}(n=e_{1})} \gtrsim R^{\sigma}T^{2}.$$

Proof. We shall first briefly gives the guideline of the proof. In order to establish the required lower estimate for $||S_{\sigma}[\vec{\phi}_{0,N}](T)||_{\hat{w}_{s}^{p,q}}$, we first write $\mathcal{F}S_{\sigma}[\vec{\phi}_{0,N}](T)(\xi)$ (with $\xi \in Q_2$) in terms of a double sum (see (3.5)). We shall naturally arrive summation over \mathcal{A} by the choice of data (3.3) with large N. And then over \mathcal{B} by applying suitable trigonometric identities for cosine functions. Further we will divide the terms under summation into two category: one collection of good terms (say I_0) which helps us to get the lower bound and the other collection of bad terms (I_1) which none the less will have some upper bound. Subsequently we will choose the time T so that the good terms dominate over the bad terms to achieve the required estimate. We now produce the details of the proof. Note that $\overline{\mathcal{F}u_0}(-\cdot) = \mathcal{F}u_0$. Set

$$\Gamma_{\xi} = \left\{ (\xi_1, \cdots, \xi_{2\sigma+1}) \in \mathbb{R}^{\sigma d} : \sum_{\ell=1}^{\sigma} \xi_{\ell} = \xi \right\}$$

and $d\Gamma_{\xi}$ denote the $(\sigma - 1)$ -dimensional Lebesgue measure on the hyperplane Γ_{ξ} . Note that for $\xi \in Q_2$, using $N \gg 1$ we have

$$\mathcal{F}S_{\sigma}[\phi_{0,N}](T)(\xi)$$

$$= \int_{0}^{T} \frac{\sin((T-t)|\xi|)}{|\xi|} \left[\left(\bigwedge_{\ell=1}^{\rho} \cos(t|\cdot|)\mathcal{F}u_{0} \right) * \left(\bigwedge_{m=\rho+1}^{\sigma} \cos(t|\cdot|)\overline{\mathcal{F}u_{0}}(-\cdot) \right) \right](\xi)dt$$

$$= \int_{0}^{T} \frac{\sin((T-t)|\xi|)}{|\xi|} \int_{\Gamma_{\xi}} \prod_{\ell=1}^{\sigma} \cos(t|\xi_{\ell}|)\mathcal{F}u_{0}(\xi_{\ell})d\Gamma_{\xi}dt$$

$$= R^{\sigma} \sum_{\mathcal{A}} \int_{0}^{T} \frac{\sin((T-t)|\xi|)}{|\xi|} \int_{\Gamma_{\xi}} \prod_{\ell=1}^{\sigma} \cos(t|\xi_{\ell}|)\mathbf{1}_{\eta_{\ell}+Q_{2}}(\xi_{\ell})d\Gamma_{\xi}dt$$

$$= \frac{R^{\sigma}}{4^{\sigma}} \sum_{\mathcal{A}} \sum_{\mathcal{B}} \int_{0}^{T} \frac{\sin((T-t)|\xi|)}{|\xi|} \int_{\Gamma_{\xi}} \cos\left(t\sum_{\ell=1}^{\sigma} \epsilon_{\ell}|\xi_{\ell}|\right) \prod_{\ell=1}^{\sigma} \mathbf{1}_{\eta_{\ell}+Q_{2}}(\xi_{\ell})d\Gamma_{\xi}dt \qquad (3.5)$$

where the sums are taken over

$$\mathcal{A} = \left\{ (\eta_1, \cdots, \eta_\sigma) \in \Sigma^\sigma : \sum_{\ell=1}^\sigma \eta_\ell = 0 \right\}, \quad \mathcal{B} = \{ (\varepsilon_1, \cdots, \varepsilon_\sigma) \in \{\pm 1\}^\sigma \}$$

respectively. For $\eta = (\eta_1, \cdots, \eta_\sigma) \in \Sigma^{\sigma}$ set

$$\mathcal{B}_0(\eta) = \left\{ (\varepsilon_1, \cdots, \varepsilon_\sigma) \in \{\pm 1\}^\sigma : \sum_{\ell=1}^\sigma \varepsilon_\ell |\eta_\ell| = 0 \right\}, \quad \mathcal{B}_1(\eta) = \mathcal{B} \setminus \mathcal{B}_0(\eta).$$

Then splitting the inner sum in (3.5) over $\mathcal{B}_0(\eta)$ and $\mathcal{B}_1(\eta)$ we write

$$\mathcal{F}S_{\sigma}[\vec{\phi}_{0,N}](T)(\xi) = \frac{R^{\sigma}}{4^{\sigma}} \sum_{\mathcal{A}} \left(I_0(\eta, T, \xi) + I_1(\eta, T, \xi) \right).$$

Note that for each $\eta \in \mathcal{A}$ the set $\mathcal{B}_0(\eta)$ is non empty. This is because for $\eta \in \mathcal{A}$, $\sum_{\ell=1}^{\sigma} \eta_\ell = 0$ and then $\left(\frac{(\eta_1)_1}{|\eta_1|}, \cdots, \frac{(\eta_{\sigma})_1}{|\eta_{\sigma}|}\right) \in \mathcal{B}_0(\eta)$ (here $(\eta_\ell)_1$ denotes the first coordinate of η_ℓ). For a fixed $\varepsilon = (\varepsilon_1, \cdots, \varepsilon_{\sigma}) \in \mathcal{B}_0(\eta)$ and $\xi_\ell \in \eta_\ell + Q_2$ one has (using triangular inequality)

$$\left|\sum_{\ell=1}^{\sigma} \varepsilon_{\ell} |\xi_{\ell}|\right| \leq \left|\sum_{\ell=1}^{\sigma} \varepsilon_{\ell} |\eta_{\ell}|\right| + \left|\sum_{\ell=1}^{\sigma} |\xi_{\ell} - \eta_{\ell}|\right| \leq 1.$$

Therefore, for $0 \leq t \leq T \ll 1$, we have $\cos\left(t \sum_{\ell=1}^{\sigma} \varepsilon_{\ell} |\xi_{\ell}|\right) \geq \frac{1}{2}$. On the other hand, we have

$$\frac{\sin((T-t)|\xi|)}{|\xi|} \gtrsim T - t$$

provided $0 \leq t < T \ll 1$ and $\xi \in Q_2$. Hence, for $\xi \in Q_2$, we have

$$I_0(\eta, T, \xi) \gtrsim \sum_{\varepsilon \in \mathcal{B}_0} \int_0^T (T - t) \int_{\Gamma_{\xi}} \prod_{\ell=1}^{\sigma} \mathbf{1}_{\eta_\ell + Q_2}(\xi_\ell) d\Gamma_{\xi} dt \gtrsim T^2 \mathbf{1}_{Q_2}(\xi)$$

as $\mathbf{1}_{\alpha+Q_2} * \mathbf{1}_{\beta+Q_2} \ge c_d \mathbf{1}_{\alpha+\beta+Q_2}$ for $\alpha, \beta \in \mathbb{R}^d$. Therefore, for $\xi \in Q_2$, $N \gg 1$ and $0 < T \ll 1$, we have

$$\frac{R^{\sigma}}{4^{\sigma}} \sum_{\mathcal{A}} I_0(\eta, T, \xi) \gtrsim T^2 R^{\sigma} \mathbf{1}_{Q_2}(\xi).$$
(3.6)

Note that for $\varepsilon \in \mathcal{B}_1(\eta)$ and $\xi_\ell \in \eta_\ell + Q_2$ one has $|\sum_{\ell=1}^{\sigma} \varepsilon_\ell |\xi_\ell|| \leq N$, this together with

$$\sum_{\ell=1}^{\sigma} \varepsilon_{\ell} |\xi_{\ell}| = \sum_{\ell=1}^{\sigma} \varepsilon_{\ell} |\eta_{\ell}| + \sum_{\ell=1}^{\sigma} \varepsilon_{\ell} (|\xi_{\ell}| - |\eta_{\ell}|) = \sum_{\ell=1}^{\sigma} \varepsilon_{\ell} |\eta_{\ell}| + \mathcal{O}(1)$$

implies

$$\left|\sum_{\ell=1}^{\sigma} \varepsilon_{\ell} |\xi_{\ell}|\right| \sim N.$$
(3.7)

Therefore using this in

$$\begin{split} I_{1}(\eta, T, \xi) \\ &= \sum_{\varepsilon \in \mathcal{B}_{1}(\eta)} \frac{1}{2|\xi|} \int_{0}^{T} \int_{\Gamma_{\xi}} \left[\sin\left(T|\xi| - t\left(|\xi| + \sum_{\ell=1}^{\sigma} \varepsilon_{\ell}|\xi_{\ell}|\right)\right) \right) + \sin\left(T|\xi| - t\left(|\xi| - \sum_{\ell=1}^{\sigma} \varepsilon_{\ell}|\xi_{\ell}|\right)\right) \right] \\ &\times \prod_{\ell=1}^{\sigma} \mathbf{1}_{\eta_{\ell} + Q_{2}}(\xi_{\ell}) d\Gamma_{\xi} dt \\ &= \sum_{\varepsilon \in \mathcal{B}_{1}(\eta)} \frac{1}{2|\xi|} \int_{\Gamma_{\xi}} \left[\frac{\cos\left(T|\xi| - t\left(|\xi| + \sum_{\ell=1}^{\sigma} \varepsilon_{\ell}|\xi_{\ell}|\right)\right)}{|\xi| + \sum_{\ell=1}^{\sigma} \varepsilon_{\ell}|\xi_{\ell}|} + \frac{\cos\left(T|\xi| - t\left(|\xi| - \sum_{\ell=1}^{\sigma} \varepsilon_{\ell}|\xi_{\ell}|\right)\right)}{|\xi| - \sum_{\ell=1}^{\sigma} \varepsilon_{\ell}|\xi_{\ell}|} \right]_{t=0}^{T} \\ &\times \prod_{\ell=1}^{\sigma} \mathbf{1}_{\eta_{\ell} + Q_{2}}(\xi_{\ell}) d\Gamma_{\xi} \end{split}$$

we get $|I_1(\eta, T, \xi)| \leq N^{-1} \mathbf{1}_{2^{\sigma-1}Q_2}(\xi)$ for $\xi \in \frac{3}{4}e_1 + Q_{\frac{1}{2}}$ as $\mathbf{1}_{\alpha+Q_2} * \mathbf{1}_{\beta+Q_2} \leq c_d \mathbf{1}_{\alpha+\beta+2Q_2}$. Hence for $\xi \in \frac{3}{4}e_1 + Q_{\frac{1}{2}}$,

$$\frac{R^{\sigma}}{4^{\sigma}} \left| \sum_{\mathcal{A}} I_1(\eta, T, \xi) \right| \lesssim N^{-1} R^{\sigma} \mathbf{1}_{2^{\sigma-1}Q_2}(\xi).$$
(3.8)

Therefore using (3.6) we have for $\xi \in \frac{3}{4}e_1 + Q_{\frac{1}{2}} \subset Q_2$

$$|\mathcal{F}S_{\sigma}[\vec{\phi}_{0,N}](T,\xi)| \gtrsim T^2 R^{\sigma}$$
(3.9)

provided $T^2 \gg N^{-1}$ and $0 < T \ll 1$. Thus we conclude

$$\|S_{\sigma}[\vec{\phi}_{0,N}](T)\|_{\hat{w}_{s}^{p,q}} \ge \left\| \left\| \chi_{n+Q_{1}}(\xi)\mathcal{F}S_{\sigma}[\vec{\phi}_{0,N}](T)(\xi) \right\|_{L^{p}_{\xi}} \langle n \rangle^{s} \right\|_{\ell^{q}(n=e_{1})} \gtrsim T^{2}R^{\sigma}$$
(3.10)

if $N^{-1/2} < T \ll 1$.

3.2. Estimates in $W_s^{2,q}$.

Lemma 3.4 (inclusion). Let $p, q, q_1, q_2 \in [1, \infty]$ and $s \in \mathbb{R}$. Then (1) $||f||_{W_s^{2,q}} \leq ||f||_{\hat{w}_s^{2,q}}$ if $q \leq 2$ (2) $||f||_{W_s^{p,q_1}} \leq ||f||_{W_s^{p,q_2}}$ if $q_1 \geq q_2$.

Proof. (1) is a consequence of Minkowski inequality and Plancherel theorem whereas (2) follows from the fact that $\ell^{q_2} \hookrightarrow \ell^{q_1}$ if $q_1 \ge q_2$.

Lemma 3.5. Let $\vec{u}_{0,N}$ be given by (3.3), $s \leq 0$, $1 \leq q \leq \infty$ and $0 \leq t \leq 1$. Then there exists C > 0 independent of R, N, t such that followings hold:

- (1) $\|\vec{u}_{0,N} \vec{u}_0\|_{\mathcal{W}^{2,q}_s} \lesssim RN^s$, where $\mathcal{W}^{2,q}_s = W^{2,q}_s \times W^{2,q}_{s-1}$
- (2) $||S_1[\vec{u}_{0,N}](t)||_{W^{2,q}_s} \lesssim 1 + RN^s$
- (3) $\|S_{\sigma}[\vec{u}_{0,N}](t) S_{\sigma}[\vec{\phi}_{0,N}](t)\|_{W^{2,q}_{s}} \lesssim t^2 R^{\sigma-1}$
- (4) $\|S_k[\vec{u}_{0,N}](t)\|_{W^{2,q}_*} \lesssim C^k R^k t^{2\frac{k-1}{\sigma-1}}$.

Proof. By Lemma 3.4, we have

$$\|\vec{u}_{0,N} - \vec{u}_0\|_{W_s^{2,q}} \lesssim \begin{cases} \|\vec{u}_{0,N} - \vec{u}_0\|_{\hat{w}_s^{2,q}} \lesssim RN^s & \text{for } q \in [1,2] \\ \|\vec{u}_{0,N} - \vec{u}_0\|_{W_s^{2,2}} \lesssim RN^s & \text{for } q \in (2,\infty] \end{cases}$$

using Lemma 3.2 (1). Similarly the other estimates also follow from Lemma 3.2.

Lemma 3.6. Let
$$\vec{\phi}_{0,N}$$
 be given by (3.3), $s \in \mathbb{R}$, $1 \leq q \leq \infty$, $N^{-1/2} < T \ll 1$. Then we h
 $\|S_{\sigma}[\vec{\phi}_{0,N}](T)\|_{W^{2,q}_{s}} \ge \|\|\Box_{n}S_{\sigma}[\vec{\phi}_{0,N}](T)(\xi)(1+|n|)^{s}\|_{\ell^{q}(n=e_{1})}\|_{L^{2}_{\xi}} \gtrsim R^{\sigma}T^{2}.$

Proof. Note that using Plancherel theorem and (3.9) we have

$$\begin{aligned} \|S_{\sigma}[\vec{\phi}_{0,N}](T)\|_{W^{2,q}_{s}} &\geq \left\| \|\mathcal{F}^{-1}\sigma_{n}\mathcal{F}S_{\sigma}[\vec{\phi}_{0,N}](T)(x)(1+|n|)^{s}\|_{\ell^{q}(n=e_{1})} \right\|_{L^{2}_{x}} \\ &= 2^{s/2} \left\| \sigma_{e_{1}}\mathcal{F}S_{\sigma}[\vec{\phi}_{0,N}](T)(\xi) \right\|_{L^{2}_{\xi}} \gtrsim R^{\sigma}T^{2}. \end{aligned}$$

This completes the proof.

3.3. Proof of main result.

Proof of Theorem 1.2. We first consider the case when $X_s^{p,q}(\mathcal{M}) = \hat{w}_s^{p,q}(\mathcal{M})$. In view of the comment at the beginning of this section, it is enough to prove NI with infinite loss of regularity at $\vec{u}_0 \in \overline{\mathcal{FL}^1}(\mathcal{M}) \cap \mathcal{X}_s^{p,q}(\mathcal{M})$. By Lemma 2.7, we have the existence of a unique solution to (1.1) with initial condition given by (3.3) in $\mathcal{FL}^1(\mathcal{M})$ up to time T whenever $(\|\vec{u}_0\|_{\overline{\mathcal{FL}^1}} + R)^{\frac{\sigma-1}{2}}T \ll 1$ which is implied by (0) $R^{\frac{\sigma-1}{2}}T \ll 1$ as $R \gg 1$. In view of Lemma 3.2 and if $RT^{\frac{2}{\sigma-1}} \ll 1$ (\Leftrightarrow (0)), $\sum_{\ell=2}^{\infty} \|S_{(\sigma-1)\ell+1}[\vec{u}_{0,N}](T)\|_{\hat{w}_s^{p,q}}$ can be dominated by the sum of the geometric series. Specifically, we have

$$\sum_{\ell=2}^{\infty} \left\| S_{(\sigma-1)\ell+1}[\vec{u}_{0,N}](T) \right\|_{\hat{w}_{s}^{p,q}} \lesssim R \sum_{\ell=2}^{\infty} (CR)^{(\sigma-1)\ell} T^{2\ell} \lesssim R^{2\sigma-1} T^{4}.$$
(3.11)

Note that

$$\left\|u_{N}(T)\right\|_{\widehat{w}_{\theta}^{p,q}} \geq \left\|\left\|\chi_{n+Q_{1}}\mathcal{F}u_{N}(T)\right\|_{L^{p}}\left\langle n\right\rangle^{\theta}\right\|_{\ell^{q}(n=e_{1})} \sim_{\theta,s} \left\|\left\|\chi_{n+Q_{1}}\mathcal{F}u_{N}(T)\right\|_{L^{p}}\left\langle n\right\rangle^{s}\right\|_{\ell^{q}(n=e_{1})}$$

ave

therefore by Lemma 2.7 and triangle inequality, we obtain

$$\begin{aligned} \|u_{N}(T)\|_{\hat{w}_{\theta}^{p,q}} \\ \gtrsim \|\|\chi_{n+Q_{1}}\mathcal{F}S_{\sigma}[u_{0,N}](T)\|_{L^{p}}\langle n\rangle^{s}\|_{\ell^{q}(n=e_{1})} - c\left(\|\|\chi_{n+Q_{1}}\mathcal{F}S_{1}[u_{0,N}](T)\|_{L^{p}}\langle n\rangle^{s}\|_{\ell^{q}(n=e_{1})} \right. \\ \left. + \sum_{\ell=2}^{\infty} \|\|\chi_{n+Q_{1}}\mathcal{F}S_{2\sigma\ell}[u_{0,N}](T)\|_{L^{p}}\langle n\rangle^{s}\|_{\ell^{q}(n=e_{1})} \right) \\ \gtrsim \|\|\chi_{n+Q_{1}}\mathcal{F}S_{\sigma}[u_{0,N}](T)\|_{L^{p}}\langle n\rangle^{s}\|_{\ell^{q}(n=e_{1})} - c\|S_{1}[\vec{u}_{0,N}](T)\|_{\hat{w}_{s}^{p,q}} - c\sum_{\ell=2}^{\infty} \|S_{(\sigma-1)\ell+1}[\vec{u}_{0,N}](T)\|_{\hat{w}_{s}^{p,q}}. \end{aligned}$$

Assume $m \in \mathbb{N}$ be given. In order to ensure $||u(T)||_{\widehat{w}_{q}^{p,q}} \gtrsim ||S_{\sigma}[\vec{u}_{0,N}](T)||_{\widehat{w}_{s}^{p,q}} \gg m$ we rely on the conditions

$$\int \|S_1[\vec{u}_{0,N}](T)\|_{\hat{w}_s^{p,q}},$$
(3.12)

$$\left\| \left\| \chi_{n+Q_1} \mathcal{F} S_{\sigma} [u_{0,N}](T) \right\|_{L^p} \langle n \rangle^s \right\|_{\ell^q (n=e_1)} \gg \begin{cases} \sum_{\ell=2}^{\infty} \left\| S_{(\sigma-1)\ell+1} [\vec{u}_{0,N}](T) \right\|_{\hat{w}_s^{p,q}}, & (3.13) \\ m. & (3.14) \end{cases}$$

To achieve (3.12)-(3.14), we rely on Lemmata 3.2, 3.3. To use Lemma 3.3, we impose (i) $N^{-1/2} \ll T \ll 1$. In view of Lemmata 3.2, 3.3, and (3.11), to prove (3.13) it is sufficient to have: (ii) (a) $R^{\sigma}T^2 \gg R^{2\sigma-1}T^4 \iff R^{\sigma-1}T^2 \ll 1 \iff (0)$ and (ii) (b) $T^2R^{\sigma-1} \ll R^{\sigma}T^2 \iff R \gg 1$. To achieve (3.14) we impose (iii) $R^{\sigma}T^2 \gg m$ (along with (ii) (b)). To ensure $\|\vec{u}_{0,N} - \vec{u}_0\|_{\mathcal{X}^{p,q}_s} < 1/m$, in view of (3.2) we impose (iv) $RN^s < 1/m$. At the end (iii) (along with (ii) (b)) imply (3.12) using Lemma 3.2.

Case $-\frac{1}{\sigma-1} \leq s < 0$. Set $R = N^{-s-\delta}, T = N^{\frac{\sigma-1}{2}(s+\frac{\delta}{2})}$ with $0 < \delta \ll 1$ satisfying $\frac{\sigma+1}{2}\delta < -s$. Note that with $N \gg 1$ we have $R \gg 1$ and

$$\begin{split} N^{-\frac{1}{2}}T^{-1} &\sim N^{-\frac{1}{2} - \frac{\sigma-1}{2}(s+\frac{\delta}{2})} = N^{-\frac{\sigma-1}{2}(s+\frac{1}{\sigma-1}+\frac{\delta}{2})} \ll 1 & \text{as } s + \frac{1}{\sigma-1} + \frac{\delta}{2} > s + \frac{1}{\sigma-1} \geqslant 0 \\ T &\sim N^{\frac{\sigma-1}{2}(s+\delta/2)} \ll 1 & \text{as } s + \delta/2 < -\sigma\delta/2 < 0 \\ R^{\sigma-1}T^2 &\sim N^{-(\sigma-1)(s+\delta)+(\sigma-1)(s+\frac{\delta}{2})} = N^{-(\sigma-1)\frac{\delta}{2}} \ll 1 & \text{as } -(\sigma-1)\delta/2 < 0 \\ RN^s &\sim N^{-(s+\delta)+s} = N^{-\delta} \ll \frac{1}{m} & \text{as } -\delta < 0 \\ R^{\sigma}T^2 &\sim N^{-\sigma(s+\delta)+(\sigma-1)(s+\delta/2)} = N^{-s-(\sigma+1)\delta/2} \gg m & \text{as } -s - (\sigma+1)\delta/2 > 0. \end{split}$$

Case $s < -\frac{1}{\sigma-1}$. Set $R = N^{\frac{1}{\sigma-1}-\delta}, T = N^{-\frac{1}{2}+\frac{\sigma-1}{4}\delta}$ with $0 < \delta \ll 1$ satisfying $(\sigma+1)\delta < \frac{2}{\sigma-1}$ and $(\sigma-1)\frac{\delta}{4} < \frac{1}{2}$.

Note that with $N \gg 1$ we have $R \gg 1$ and

$$\begin{split} N^{-1/2}T^{-1} &\sim N^{-1/2+1/2-(\sigma-1)\delta/4} = N^{-(\sigma-1)\delta/4} \ll 1 & \text{as } (\sigma-1)\delta/4 > 0 \\ T &\sim N^{-1/2+(\sigma-1)\delta/4} \ll 1 & \text{as } -1/2 + (\sigma-1)\delta/4 < 0 \\ R^{\sigma-1}T^2 &\sim N^{1-(\sigma-1)\delta-1+(\sigma-1)\delta/2} = N^{-(\sigma-1)\delta/2} \ll 1 & \text{as } -(\sigma-1)\delta/2 < 0 \\ RN^s &\sim N^{1/(\sigma-1)-\delta+s} \ll \frac{1}{m} & \text{as } 1/(\sigma-1) + s < 0 \\ R^{\sigma}T^2 &\sim N^{\sigma(\frac{1}{\sigma-1}-\delta)-1+(\sigma-1)\frac{\delta}{2}} = N^{\frac{1}{\sigma-1}-(\sigma+1)\frac{\delta}{2}} \gg m & \text{as } 1/(\sigma-1) - (\sigma+1)\delta/2 > 0 \end{split}$$

Thus with both the cases the conditions (i)-(iv) are satisfied and hence we are done with the case $\mathcal{X}_s^{p,q} = \hat{w}_s^{p,q}$.

For the case $X_s^{p,q} = W_s^{2,q}$ we use same argument as above: Note that using Lemmata 3.5, 3.6.

$$\begin{split} \|u_{N}(T)\|_{W^{2,q}_{\theta}} & \geq \left\| \|\Box_{n}u_{N}(T)\langle n\rangle^{\sigma} \|_{\ell^{q}(n=e_{1})} \right\|_{L^{2}} \sim_{\theta,s} \left\| \|\Box_{n}u_{N}(T)\langle n\rangle^{s} \|_{\ell^{q}(n=e_{1})} \right\|_{L^{2}} \\ & \geq \left\| \|\Box_{n}S_{\sigma}[u_{0,N}](T)\langle n\rangle^{s} \|_{\ell^{q}(n=e_{1})} \right\|_{L^{2}} - c\|S_{1}[u_{0,N}](T)\|_{W^{2,q}_{s}} - c\sum_{\ell=2}^{\infty} \|S_{(\sigma-1)\ell+1}[\vec{u}_{0,N}](T)\|_{W^{2,q}_{s}} \\ & \geq \left\| \|\Box_{n}S_{\sigma}[u_{0,N}](T)\langle n\rangle^{s} \|_{\ell^{q}(n=e_{1})} \right\|_{L^{2}} \gg m. \end{split}$$

and $\|\vec{u}_{0,N} - \vec{u}_0\|_{\mathcal{X}^{2,q}_s} < 1/m$ provided we choose R, N, T as in the case of $\hat{w}_s^{p,q}$.

Acknowledgement: D.G. B is thankful to DST-INSPIRE (DST/INSPIRE/04/2016/001507) for the research grant. S. H. acknowledges Dept of Atomic Energy, Govt of India, for the financial support and Harish-Chandra Research Institute for the research facilities provided. Both authors are grateful to professor Rémi Carles for his thoughtful comments and fruitful discussion on the topic. D. G. B. is grateful to professor Tadahiro Oh and Justin Forlano for sending their preprint [15] and introducing to Fourier amalgam spaces.

References

- I. BEJENARU AND T. TAO, Sharp well-posedness and ill-posedness results for a quadratic non-linear Schrödinger equation, J. Funct. Anal., 233 (2006), pp. 228–259.
- [2] A. BÉNYI, K. GRÖCHENIG, K. A. OKOUDJOU, AND L. G. ROGERS, Unimodular Fourier multipliers for modulation spaces, J. Funct. Anal., 246 (2007), pp. 366–384.
- [3] A. BÉNYI AND K. A. OKOUDJOU, Local well-posedness of nonlinear dispersive equations on modulation spaces, Bull. Lond. Math. Soc., 41 (2009), pp. 549–558.
- [4] Á. BÉNYI AND K. A. OKOUDJOU, Modulation Spaces: With Applications to Pseudodifferential Operators and Nonlinear Schrödinger Equations, 2020.
- [5] D. G. BHIMANI AND R. CARLES, Norm inflation for nonlinear Schrodinger equations in Fourier-Lebesgue and modulation spaces of negative regularity, Journal of Fourier Analysis and Applications, 26 (2020).
- [6] D. G. BHIMANI, M. GRILLAKIS, AND K. A. OKOUDJOU, The Hartree–Fock equations in modulation spaces, Communications in Partial Differential Equations, (2020), pp. 1–30.
- [7] D. G. BHIMANI AND S. HAQUE, Strong ill-posedness for fractional Hartree and cubic NLS equations, preprint arXiv:2101.03991, (2021).
- [8] D. G. BHIMANI AND P. K. RATNAKUMAR, Functions operating on modulation spaces and nonlinear dispersive equations, J. Funct. Anal., 270 (2016), pp. 621–648.

...

- N. BURQ AND N. TZVETKOV, Random data Cauchy theory for supercritical wave equations. I. Local theory, Invent. Math., 173 (2008), pp. 449–475.
- [10] R. CARLES, E. DUMAS, AND C. SPARBER, Geometric optics and instability for NLS and Davey-Stewartson models, J. Eur. Math. Soc. (JEMS), 14 (2012), pp. 1885–1921.
- [11] R. CARLES AND T. KAPPELER, Norm-inflation with infinite loss of regularity for periodic NLS equations in negative Sobolev spaces, Bull. Soc. Math. France, 145 (2017), pp. 623–642.
- [12] M. CHRIST, J. COLLIANDER, AND T. TAO, Ill-posedness for nonlinear Schrödinger and wave equations. arXiv:0311048, 2003.
- [13] E. CORDERO AND F. NICOLA, Remarks on Fourier multipliers and applications to the wave equation, J. Math. Anal. Appl., 353 (2009), pp. 583–591.
- [14] H. G. FEICHTINGER, Modulation spaces on locally compact Abelian groups. Technical Report, University of Vienna, 1983, and in "Wavelets and Their Applications" (eds. M. Krishna, R. Radha and S. Thangavelu), 99-140, Allied Publishers, New Delhi, 2003., 1983. Available on researchgate.net.
- [15] J. FORLANO AND T. OH, Normal form approach to the one-dimensional cubic nonlinear Schrödinger equation in Fourier-amalgam spaces, preprint.
- [16] J. FORLANO AND M. OKAMOTO, A remark on norm inflation for nonlinear wave equations, Dynamics of Partial Differential Equations, (2020), pp. 361–381.
- [17] K. GRÖCHENIG, Foundations of time-frequency analysis, Applied and Numerical Harmonic Analysis, Birkhäuser Boston, Inc., Boston, MA, 2001.
- [18] T. IWABUCHI AND T. OGAWA, Ill-posedness for the nonlinear Schrödinger equation with quadratic non-linearity in low dimensions, Trans. Amer. Math. Soc., 367 (2015), pp. 2613–2630.
- [19] N. KISHIMOTO, A remark on norm inflation for nonlinear Schrödinger equations, Communications on Pure & Applied Analysis, 18 (2019), p. 1375.
- [20] G. LEBEAU, Perte de régularité pour les équations d'ondes sur-critiques, Bull. Soc. Math. France, 133 (2005), pp. 145–157.
- [21] H. LINDBLAD, A sharp counterexample to the local existence of low-regularity solutions to nonlinear wave equations, Duke Math. J., 72 (1993), pp. 503–539.
- [22] H. LINDBLAD AND C. D. SOGGE, On existence and scattering with minimal regularity for semilinear wave equations, J. Funct. Anal., 130 (1995), pp. 357–426.
- [23] T. OH, A remark on norm inflation with general initial data for the cubic nonlinear Schrödinger equations in negative Sobolev spaces, Funkcial. Ekvac., 60 (2017), pp. 259–277.
- [24] M. REICH AND M. REISSIG, Wave equations in modulation spaces—decay versus loss of regularity, in New tools for nonlinear PDEs and application, Trends Math., Birkhäuser/Springer, Cham, 2019, pp. 371–390.
- [25] M. RUZHANSKY, M. SUGIMOTO, J. TOFT, AND N. TOMITA, Changes of variables in modulation and Wiener amalgam spaces, Math. Nachr., 284 (2011), pp. 2078–2092.
- [26] M. RUZHANSKY, M. SUGIMOTO, AND B. WANG, Modulation spaces and nonlinear evolution equations, in Evolution equations of hyperbolic and Schrödinger type, vol. 301 of Progr. Math., Birkhäuser/Springer Basel AG, Basel, 2012, pp. 267–283.
- [27] M. RUZHANSKY AND V. TURUNEN, Pseudo-differential operators and symmetries, vol. 2 of Pseudo-Differential Operators. Theory and Applications, Birkhäuser Verlag, Basel, 2010. Background analysis and advanced topics.
- [28] T. TAO, Low regularity semi-linear wave equations, Comm. Partial Differential Equations, 24 (1999), pp. 599–629.
- [29] B. WANG AND H. HUDZIK, The global Cauchy problem for the NLS and NLKG with small rough data, J. Differential Equations, 232 (2007), pp. 36–73.
- [30] B. WANG, Z. HUO, C. HAO, AND Z. GUO, Harmonic analysis method for nonlinear evolution equations. I, World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2011.
- [31] B. WANG, Z. LIFENG, AND G. BOLING, Isometric decomposition operators, function spaces $E_{p,q}^{\lambda}$ and applications to nonlinear evolution equations, J. Funct. Anal., 233 (2006), pp. 1–39.

D. BHIMANI AND S. HAQUE

DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF SCIENCE EDUCATION AND RESEARCH, DR. HOMI BHABHA ROAD, PUNE 411008, INDIA

Email address: divyang.bhimani@iiserpune.ac.in

Harish-Chandra Research Institute, Chhatnag Road, Jhunsi, Prayagraj (Allahabad) 211019, India

Email address: saikatulhaque@hri.res.in