

Energy distribution of solutions to defocusing semi-linear wave equation in higher dimensional space *

Liang Li, Ruipeng Shen

Centre for Applied Mathematics
Tianjin University
Tianjin, China

May 9, 2021

Abstract

The topic of this paper is a semi-linear, defocusing wave equation $u_{tt} - \Delta u = -|u|^{p-1}u$ in sub-conformal case in the higher dimensional space whose initial data are radical and come with a finite energy. We prove some decay estimates of the the solutions if initial data decay at a certain rate as the spatial variable tends to infinity. A combination of this property with a method of characteristic lines give a scattering result if the initial data satisfy

$$E_{\kappa}(u_0, u_1) = \int_{\mathbb{R}^d} (|x|^{\kappa} + 1) \left(\frac{1}{2} |\nabla u_0(x)|^2 + \frac{1}{2} |u_1(x)|^2 + \frac{1}{p+1} |u_0(x)|^{p+1} \right) dx < +\infty.$$

Here $\kappa = \frac{(2-d)p+(d+2)}{p+1}$.

1 Introduction

1.1 Background

In this work we consider the defocusing nonlinear wave equation in dimensions $d \geq 3$.

$$\begin{cases} \partial_t^2 u - \Delta u = -|u|^{p-1}u, & (x, t) \in \mathbb{R}^d \times \mathbb{R}; \\ u(\cdot, 0) = u_0; \\ u_t(\cdot, 0) = u_1. \end{cases} \quad (CP1)$$

The conserved energy is defined by

$$E(u, u_t) = \int_{\mathbb{R}^d} \left(\frac{1}{2} |\nabla u(x, t)|^2 + \frac{1}{2} |u_t(x, t)|^2 + \frac{1}{p+1} |u(x, t)|^{p+1} \right) dx.$$

Local theory Defocusing nonlinear wave equations

$$\partial_t^2 u - \Delta u = -|u|^{p-1}u, \quad (x, t) \in \mathbb{R}^d \times \mathbb{R}$$

have been extensively studied, especially in the 3 or higher dimensional space. The existence and uniqueness of solutions to semi-linear wave equation like (CP1) follows a combination of suitable Strichartz estimates and a fixed-point argument, Kapitanski [1] and Lindblad-Sogge [2] give more details.

*MSC classes: 35L05, 35L71.

Conjecture 1.1. *Any solution to (CP1) with initial data $(u_0, u_1) \in \dot{H}^{s_p} \times \dot{H}^{s_p-1}$ must exist for all time $t \in \mathbb{R}$ and scatter in both two time directions.*

This is still an open problem. Although there are many related results by different methods.

Scattering results with a priori estimates It has been proved that if a solution u with a maximal lifespan I satisfies an a priori estimate

$$\sup_{t \in I} \|(u(\cdot, t), u_t(\cdot, t))\|_{\dot{H}^{s_p} \times \dot{H}^{s_p-1}(\mathbb{R}^d)} < +\infty,$$

then u is defined for all time t and scatters. In fact, there are many works for different ranges of d and p , sometimes with a radial assumption. Different methods were used for different range of d and p . The details can be found in Kenig-Merle [4], Killip-Visan [14] (3 dimension), Killip-Visan [15] (all dimensions) for energy supercritical case and R.Shen [5], Dodson-Lawrie [6] (3 dimension), Rodriguez [16] (dimension 4 and 5) for energy subcritical case.

Strong Assumption on initial data There are multiple scattering results if we assume that the initial data satisfy stronger regularity and/or decay conditions. These results are usually proved via a suitable global space-time integral estimate. In $d \geq 3$ case, if the initial data (u_0, u_1) satisfy

$$\int_{\mathbb{R}^d} (1 + |x|)^2 \left(\frac{1}{2} |\nabla u_0(x)|^2 + \frac{1}{2} |u_1(x)|^2 + \frac{1}{p+1} |u_0(x)|^{p+1} \right) < +\infty$$

the conformal conservation law (see Ginibre-Velo [18] and Hidano [19]) leads to the scattering of solutions for $1 + 4/(d-1) \leq p < 1 + 4/(d-2)$. In 3-dimension case, R.Shen [17] proved the scattering result for $3 \leq p \leq 5$ if initial data (u_0, u_1) are radial and satisfy

$$\int_{\mathbb{R}^3} (|x|^\kappa + 1) \left(\frac{1}{2} |\nabla u_0|^2 + \frac{1}{2} |u_1|^2 + \frac{1}{p+1} |u_0|^{p+1} \right) dx < +\infty,$$

here $\kappa > \kappa_0(p) = \frac{5-p}{p+1}$. In $d = 3, p = 3$ case, Dodson [10] gives a proof of the conjecture above for (CP1) with radial data.

But most of these results above are in the conformal case or super-conformal case.

1.2 Main tools

Next we introduce main tools of this paper. The first tool is still a Morawetz-type estimate, and the second tool is the method of characteristic lines.

Morawetz estimates This kind of estimates were first found by Morawetz [3] for wave/Klein-Gordon equations. Lin-Strauss [7] then generalized Morawetz estimates to Schrödinger equations. Colliander-Keel-Staffilani-Takaoka-Tao [8] introduced interaction Morawetz estimates for Schrödinger equations. Nowadays the Morawetz estimate has been one of the most important tools in the study of dispersive equations.

Method of characteristic lines R.Shen [9] generalizes the 3D method to higher dimensions. Let u be a radial solution to (CP1) with a finite energy, reduce the equation to a one-dimensional one by defining $w(r, t) = r^{\frac{d-1}{2}} u(r, t)$, and considering the equation that w satisfies

$$(\partial_t + \partial_r)(w_t - w_r) = \partial_t^2 w - \partial_r^2 w = -\frac{(d-1)(d-3)}{4} dr^{\frac{d-5}{2}} u - r^{\frac{d-1}{2}} |u|^{p-1} u.$$

This enable us to evaluate the variation of $w_t \pm w_r$ along characteristic lines $t \mp r = \text{Const}$ and obtain plentiful information about the asymptotic behaviour of solutions.

1.3 Main Results

Now we give the main results of this work. Throughout this paper we always assume $d \geq 3$ and $p < 1 + \frac{4}{d-1}$. In this case it is well known that if the initial data come with a finite energy, then the solution exist for all time. Please refer to Ginibre-Velo [21]. Our first result is about the energy distribution of solutions to (CP1).

Theorem 1.1. *Assume $d \geq 3$, $1 + \frac{4}{d-1} > p > 1 + \frac{2}{d-1}$. Let u be a solution to (CP1) with a finite energy. Then*

(a) *The following limits hold as time tends to infinity*

$$\lim_{t \rightarrow \pm\infty} \int_{|x| < |t|} \frac{|t| - |x|}{|t|} e(x, t) dx = 0.$$

(b) *The inward/outward part of energy vanishes as time tends positive/negative infinity.*

$$\lim_{t \rightarrow \pm\infty} \int_{\mathbb{R}^d} \left(|u_r \pm u_t|^2 + |\nabla u|^2 + |u|^{p+1} \right) dx = 0.$$

(c) *Furthermore, if the initial data satisfy $E_\kappa(u_0, u_1) < +\infty$ for a constant*

$$\kappa \in \left(0, \frac{(d-1)(p-1)-2}{2} \right), \quad \text{if } 1 + \frac{4}{d-1} > p > 1 + \frac{2}{d-1};$$

then we have the following decay estimates

$$\lim_{t \rightarrow \pm\infty} \int_{|x| < |t|} \frac{|t| - |x|}{|t|^{1-\kappa}} e(x, t) dx = 0;$$

$$\lim_{t \rightarrow \pm\infty} |t|^\kappa \int_{\mathbb{R}^d} \left(|u_r \pm u_t|^2 + |\nabla u|^2 + |u|^{p+1} \right) dx = 0.$$

As an application of the theory on energy distribution, we also prove the following scattering result.

Theorem 1.2. *Assume $d \geq 3$, $1 + \frac{4}{d-1} > p > \frac{3-d+2\sqrt{d^2-d+1}}{d-1}$. Let u be a radial solution to (CP1) with a finite energy and the initial data (u_0, u_1) satisfy $E_{\frac{(2-d)p+(d+2)}{p+1}}(u_0, u_1) < +\infty$, then the solution u scatters in both two time directions. More precisely, there exist two radial finite-energy free waves \tilde{u}^+, \tilde{u}^- , so that*

$$\lim_{t \rightarrow \pm\infty} \left\| (\tilde{u}^\pm(\cdot, t) - u(\cdot, t), \tilde{u}_t^\pm(\cdot, t) - u_t(\cdot, t)) \right\|_{\dot{H}^1 \times L^2(\mathbb{R}^d)} = 0.$$

Here $E_{\frac{(2-d)p+(d+2)}{p+1}}(u_0, u_1) = \int_{\mathbb{R}^d} \left(|x|^{\frac{(2-d)p+(d+2)}{p+1}} + 1 \right) \left(\frac{1}{2} |\nabla u_0(x)|^2 + \frac{1}{2} |u_1(x)|^2 + \frac{1}{p+1} |u_0(x)|^{p+1} \right) dx$.

1.4 The Structure of This Paper

This paper is organized as follows. We first give a few preliminary results in section 2, In section 3 we give a Morewetz identity and a Morawetz inequality, which is the main tool of this paper. Next in section 4 we prove the energy distribution properties of the solutions. Finally we prove the scattering theory of the solution u under an additional decay assumption in the last section.

2 Preliminary Results

Notations In this work we will use the notation $e(x, t)$ for the energy density

$$e(x, t) = \frac{1}{2} |\nabla u(x, t)|^2 + \frac{1}{2} |u_t(x, t)|^2 + \frac{1}{p+1} |u(x, t)|^{p+1}.$$

We use $u_r, \nabla u$ for the derivative in the radial direction and the covariant derivative on the sphere centred at the origin, respectively;

$$u_r(x, t) = \frac{x}{|x|} \cdot \nabla u(x, t); \quad \nabla u = \nabla u - u_r \frac{x}{|x|}; \quad |\nabla u|^2 = |u_r|^2 + |\nabla u|^2.$$

We also define the weighted energy

$$E_\kappa(u_0, u_1) = \int_{\mathbb{R}^d} (|x|^\kappa + 1) \left(\frac{1}{2} |\nabla u_0(x)|^2 + \frac{1}{2} |u_1(x)|^2 + \frac{1}{p+1} |u_0(x)|^{p+1} \right) dx.$$

In this work σ_R represents the regular measure of the sphere $\{x \in \mathbb{R}^d : |x| = R\}$. We also define c_d to be the area of the united sphere \mathbb{S}^{d-1} . Thus we have the following identities for any radial function $f(x)$

$$\int_{|x|=r} f(x) d\sigma_r(x) = c_d r^{d-1} f(r); \quad \int_{\mathbb{R}^d} f(x) dx = c_d \int_0^\infty f(r) r^{d-1} dr.$$

The notation $A \lesssim B$ means that there exists a constant c , so that the inequality $A \leq cB$ holds. We may also put subscript(s) to indicate that the constant c depends on the given subscript(s) but nothing else. In particular, the symbol \lesssim_1 is used if c is an absolute constant.

Lemma 2.1. (*Pointwise Estimate*) Assume $d \geq 3$. all radial $\dot{H}^1(\mathbb{R}^d)$ functions u satisfy

$$|u(r)| \lesssim_d r^{-\frac{d-2}{2}} \|u\|_{\dot{H}^1}, \quad r > 0.$$

If u also satisfy $u \in L^{p+1}(\mathbb{R}^d)$, then its decay is stronger as $r \rightarrow +\infty$.

$$|u(r)| \lesssim_d r^{-\frac{2(d-1)}{p+3}} \|u\|_{\dot{H}^1}^{\frac{2}{p+3}} \|u\|_{L^{p+1}}^{\frac{p+1}{p+3}}, \quad r > 0.$$

This lemma has been known for many years, more details can be found, for example, in C.E.Kenig and F.Merle [4] for $d = 3$ and R.Shen [9] for $d \geq 3$.

Lemma 2.2. (*Radiation Field*) Assume that $d \geq 3$ and let u be a solution to the free wave equation $\partial_t^2 u - \Delta u = 0$ with initial data $(u_0, u_1) \in H^1 \times L^2(\mathbb{R}^d)$. Then

$$\lim_{t \rightarrow +\infty} \int_{\mathbb{R}^d} \left(|\nabla u(x, t)|^2 - |u_r(x, t)|^2 + \frac{|u(x, t)|^2}{|x|^2} \right) dx = 0,$$

and there exists a unique function $G_+ \in L^2(\mathbb{R} \times \mathbb{S}^{d-1})$ such that

$$\lim_{t \rightarrow +\infty} \int_0^\infty \int_{\mathbb{S}^{d-1}} \left| r^{\frac{d-1}{2}} \partial_t u(r\theta, t) - G_+(r-t, \theta) \right|^2 d\theta dr = 0;$$

$$\lim_{t \rightarrow +\infty} \int_0^\infty \int_{\mathbb{S}^{d-1}} \left| r^{\frac{d-1}{2}} \partial_r u(r\theta, t) + G_+(r-t, \theta) \right|^2 d\theta dr = 0.$$

In addition, the map

$$(u_0, u_1) \rightarrow \sqrt{2} G_+ \\ \dot{H}^1 \times L^2(\mathbb{R}^d) \rightarrow L^2(\mathbb{R} \times \mathbb{S}^{d-1})$$

is a bijective isometry.

This result was known many years ago, please see Friedlander [11, 12]. Duyckaerts-Kenig-Merle [13] gives a proof for all dimensions $d \geq 3$.

3 Morawetz identity and Morewetz inequality

3.1 Morawetz identity

Proposition 3.1. (Morawetz identity) *Let u be a solution to (CP1) with a finite energy E . Then the following identity holds for any $R > 0$ and time $t_1 < t_2$.*

$$\begin{aligned} & \frac{1}{2R} \int_{t_1}^{t_2} \int_{|x| < R} \left(|\nabla u|^2 + |u_t|^2 + \frac{(d-1)(p-1)-2}{p+1} |u|^{p+1} \right) dx dt + \frac{d-1}{4R^2} \int_{t_1}^{t_2} \int_{|x|=R} |u|^2 d\sigma_R(x) dt \\ & + \int_{t_1}^{t_2} \int_{|x| > R} \left(\frac{|\nabla u|^2}{|x|} + \frac{(d-1)(p-1)}{2(p+1)} \cdot \frac{|u|^{p+1}}{|x|} + \frac{(d-3)(d-1)}{4} \cdot \frac{|u|^2}{|x|^3} \right) dx dt \\ & + \sum_{i=1,2} \int_{|x| < R} \left(\frac{R^2 - |x|^2}{2R^2} |u_r|^2 + \frac{1}{2} \left| \frac{|x|}{R} u_r + \frac{(d-1)u}{2R} + (-1)^i u_t \right|^2 + \frac{(d^2-1)|u|^2}{8R^2} + \frac{|\nabla u|^2}{2} + \frac{|u|^{p+1}}{p+1} \right) \Bigg|_{t=t_i} dx \\ & + \sum_{i=1,2} \int_{|x| > R} \left(\frac{1}{2} \left| u_r + \frac{d-1}{2} \frac{u}{|x|} + (-1)^i u_t \right|^2 + \frac{|\nabla u|^2}{2} + \frac{|u|^{p+1}}{p+1} + \frac{(d-1)(d-3)|u(x, t_i)|^2}{8|x|^2} \right) \Bigg|_{t=t_i} dx = 2E. \end{aligned}$$

Proof. we follow a similar argument to the given by Perthame and Vega in the final section of their work [20]. Let us first consider solutions with compact support. Given a positive constant R , we define two radial function Ψ and φ by

$$\nabla \Psi = \begin{cases} x, & \text{if } |x| \leq R; \\ Rx/|x|, & \text{if } |x| \geq R; \end{cases} \quad \varphi = \begin{cases} 1/2, & \text{if } |x| \leq R; \\ 0, & \text{if } |x| > R. \end{cases}$$

Since u is defined for all time t , we may also define a function on R

$$\mathcal{E}(t) = \int_{\mathbb{R}^d} u_t(x, t) \left(\nabla u(x, t) \cdot \nabla \Psi + u(x, t) \left(\frac{\Delta \Psi}{2} - \varphi \right) \right) dx. \quad (1)$$

We may differentiate \mathcal{E} , utilize the equation

$$u_{tt} - \Delta u = -|u|^{p-1}u,$$

apply integration by parts and obtain

$$\begin{aligned} -\mathcal{E}'(t) &= \int_{\mathbb{R}^d} \left(\sum_{i,j=1}^d u_i \Psi_{ij} u_j - \varphi |\nabla u|^2 + \varphi |u_t|^2 \right) dx + \frac{1}{4} \int_{\mathbb{R}^d} \nabla(|u|^2) \cdot \nabla(\Delta \Psi - 2\varphi) dx \\ &+ \int_{\mathbb{R}^d} |u|^{p+1} \left(\frac{p-1}{2(p+1)} \Delta \Psi - \varphi \right) dx \\ &= I_1 + I_2 + I_3. \end{aligned} \quad (2)$$

Here we have

$$\begin{aligned} \Psi_{ij} &= \begin{cases} \delta_{ij}, & \text{if } |x| < R; \\ \frac{R\delta_{ij}}{|x|} - \frac{Rx_i x_j}{|x|^3}, & \text{if } |x| > R; \end{cases} \quad \Delta \Psi = \begin{cases} d, & \text{if } |x| < R; \\ R(d-1)/|x|, & \text{if } |x| > R; \end{cases} \\ \Delta \Psi - 2\varphi &= \begin{cases} d-1, & \text{if } |x| \leq R; \\ R(d-1)/|x|, & \text{if } |x| \geq R; \end{cases} \in C(\mathbb{R}^d). \end{aligned}$$

When $|x| > R$, we may calculate

$$\sum_{i,j=1}^d u_i \Psi_{ij} u_j = \sum_{i,j=1}^d u_i \left(\frac{R\delta_{ij}}{|x|} - \frac{Rx_i x_j}{|x|^3} \right) u_j = \frac{R}{|x|} |\nabla u|^2 - \frac{R|\nabla u \cdot x|^2}{|x|^3} = \frac{R}{|x|} |\nabla u|^2. \quad (3)$$

Thus we have

$$I_1 = \frac{1}{2} \int_{|x| < R} (|\nabla u|^2 + |u_t|^2) dx + R \int_{|x| > R} \frac{|\nabla u|^2}{|x|} dx. \quad (4)$$

The last term in the equality above, a basic computation shows

$$I_3 = \frac{(d-1)(p-1)-2}{2(p+1)} \int_{|x| < R} |u|^{p+1} dx + \frac{(p-1)(d-1)R}{2(p+1)} \int_{|x| > R} \frac{|u|^{p+1}}{|x|} dx. \quad (5)$$

Let us calculate the left hand carefully

$$\begin{aligned} I_2 &= \frac{1}{4} \int_{\mathbb{R}^d} \nabla (|u|^2) \cdot \nabla (\Delta \Psi - 2\varphi) dx \\ &= \frac{1}{4} \int_{|x| > R} \nabla (|u|^2) \cdot \frac{-R(d-1)x}{|x|^3} dx \\ &= \frac{1}{4} \int_{|x| > R} \left[\operatorname{div} \left(|u|^2 \cdot \frac{-R(d-1)x}{|x|^3} \right) + (d-3)(d-1) \frac{R}{|x|^3} |u|^2 \right] dx \\ &= \frac{d-1}{4R} \int_{|x|=R} |u|^2 d\sigma_R(x) + \frac{(d-3)(d-1)}{4} \int_{|x| > R} \frac{|u|^2}{|x|^3} dx. \end{aligned} \quad (6)$$

Since $-\mathcal{E}'(t) = I_1 + I_2 + I_3$, we have

$$\int_{t_1}^{t_2} (I_1 + I_2 + I_3) dt = \mathcal{E}(t_1) - \mathcal{E}(t_2). \quad (7)$$

We rewrite in the form of

$$\begin{aligned} R\mathcal{E}(t_1) &= \int_{\mathbb{R}^d} Ru_t(x, t) \left(\nabla u(x, t) \cdot \nabla \Psi + u(x, t) \left(\frac{\Delta \Psi}{2} - \varphi \right) \right) dx \\ &= \frac{1}{2} \int_{\mathbb{R}^d} \left(R^2 |u_t(x, t_1)|^2 + \left| \nabla u(x, t_1) \cdot \nabla \Psi + u(x, t_1) \left(\frac{\Delta \Psi}{2} - \varphi \right) \right|^2 \right) dx \\ &\quad - \frac{1}{2} \int_{\mathbb{R}^d} \left| \nabla u(x, t_1) \cdot \nabla \Psi + u(x, t_1) \left(\frac{\Delta \Psi}{2} - \varphi \right) - Ru_t(x, t_1) \right|^2 dx \\ &= J_1 - J_2. \end{aligned} \quad (8)$$

Then we calculate J_1, J_2

$$\begin{aligned} J_1 &= \frac{1}{2} \int_{\mathbb{R}^d} \left(R^2 |u_t|^2 + |\nabla u \cdot \nabla \Psi|^2 + \left(\frac{\Delta \Psi}{2} - \varphi \right) \nabla (|u|^2) \cdot \nabla \Psi + \left(\frac{\Delta \Psi}{2} - \varphi \right)^2 |u|^2 \right) dx \\ &= \frac{1}{2} \int_{\mathbb{R}^d} \left[R^2 |u_t|^2 + |\nabla u \cdot \nabla \Psi|^2 - \operatorname{div} \left(\left(\frac{\Delta \Psi}{2} - \varphi \right) \nabla \Psi \right) |u|^2 + \left(\frac{\Delta \Psi}{2} - \varphi \right)^2 |u|^2 \right] dx. \end{aligned} \quad (9)$$

A basic calculation shows

$$\operatorname{div} \left(\left(\frac{\Delta \Psi}{2} - \varphi \right) \nabla \Psi \right) = \begin{cases} d(d-1)/2, & \text{if } |x| < R; \\ \frac{(d-1)(d-2)R^2}{2|x|^2}, & \text{if } |x| > R. \end{cases} \quad (10)$$

Thus we have

$$\begin{aligned}
J_1 &= \frac{1}{2} \int_{|x| < R} \left[R^2 |u_t|^2 + |x \cdot \nabla u|^2 + \frac{1-d^2}{4} |u|^2 \right] dx \\
&\quad + \frac{1}{2} \int_{|x| > R} \left[R^2 |u_t|^2 + R^2 |u_r|^2 + \frac{R^2(d-1)(3-d)|u|^2}{4|x|^2} \right] dx \\
&= R^2 E - R^2 \int_{|x| < R} \left[\frac{R^2 - |x|^2}{2R^2} |u_r|^2 + \frac{d^2 - 1}{8R^2} |u|^2 + \frac{1}{2} |\nabla u|^2 + \frac{1}{p+1} |u|^{p+1} \right] dx \\
&\quad + \frac{(d-1)(3-d)R^2}{8} \int_{|x| > R} \frac{|u|^2}{|x|^2} dx - R^2 \int_{|x| > R} \left(\frac{1}{2} |\nabla u|^2 + \frac{1}{p+1} |u|^{p+1} \right) dx.
\end{aligned} \tag{11}$$

In addition we have

$$J_2 = \frac{1}{2} \int_{|x| < R} \left| x \cdot \nabla u + \frac{d-1}{2} u - Ru_t \right|^2 dx + \frac{R^2}{2} \int_{|x| > R} \left| \frac{x}{|x|} \cdot \nabla u + \frac{d-1}{2} \frac{u}{|x|} - u_t \right|^2 dx. \tag{12}$$

Combining J_1, J_2 , we obtain

$$\begin{aligned}
R\mathcal{E}(t_1) &= R^2 E - R^2 \int_{|x| > R} \left(\frac{1}{2} \left| u_r + \frac{d-1}{2} \frac{u}{|x|} - u_t \right|^2 + \frac{|\nabla u|^2}{2} + \frac{|u|^{p+1}}{p+1} + \frac{(d-1)(d-3)}{8} \frac{|u|^2}{|x|^2} \right) dx \\
&\quad - R^2 \int_{|x| < R} \left[\frac{R^2 - |x|^2}{2R^2} |u_r|^2 + \frac{1}{2} \left| \frac{|x|}{R} u_r + \frac{d-1}{2R} u - u_t \right|^2 + \frac{|\nabla u|^2}{2} + \frac{|u|^{p+1}}{p+1} + \frac{(d^2-1)|u|^2}{8R^2} \right] dx.
\end{aligned}$$

Finally we find a similar expression of $-R\mathcal{E}(t_2)$

$$\begin{aligned}
-R\mathcal{E}(t_2) &= R^2 E - R^2 \int_{|x| > R} \left(\frac{1}{2} \left| u_r + \frac{(d-1)u}{2|x|} + u_t \right|^2 + \frac{|\nabla u|^2}{2} + \frac{|u|^{p+1}}{p+1} + \frac{(d-1)(d-3)|u|^2}{8|x|^2} \right) dx \\
&\quad - R^2 \int_{|x| < R} \left[\frac{R^2 - |x|^2}{2R^2} |u_r|^2 + \frac{1}{2} \left| \frac{|x|}{R} u_r + \frac{(d-1)u}{2R} + u_t \right|^2 + \frac{|\nabla u|^2}{2} + \frac{|u|^{p+1}}{p+1} + \frac{(d^2-1)|u|^2}{8R^2} \right] dx.
\end{aligned}$$

Then plug all the expressions of I_1, I_2, I_3 and $-R\mathcal{E}(t_2), -R\mathcal{E}(t_2)$ in to the integral identity to finish the proof if u is compactly supported. In order to deal with the general solution u , we fix a smooth radial cut-off function $\phi : \mathbb{R}^2 \rightarrow [0, 1]$ so that

$$\phi(x) = \begin{cases} 1, & \text{if } |x| \leq 1; \\ 0, & \text{if } |x| > 2; \end{cases}$$

define initial data $(u_{0,R'}(x), u_{1,R'}(x)) = \phi(x/R')(u(x, t_1), u_t(x, t_1))$ and consider the corresponding solution $u_{R'}$ to (CP1). The argument above shows that $u_{R'}$ satisfies Morawetz identity. We observe

- 1) The identity $u_{R'}(x, t) = u(x, t)$ holds if $|x| < R' + t_1 - t$ by finite speed of propagation;
- 2) $E(u_{0,R'}, u_{1,R'}) \rightarrow E$ as $R' \rightarrow \infty$.
- 3) The energies of $u_{R'}$ and u in the region where $u_{R'} \neq u$ both converge to zero as $R' \rightarrow +\infty$ by finite speed of propagation and energy conservation law.

These facts enable us to take the limit $R' \rightarrow +\infty$ and prove Morawetz identity for general solutions u . \square

3.2 Morawetz inequalities

A combination of Morawetz identity and finite speed of propagation gives a few useful inequalities, which is the main tool of this paper. The key observation here is that if R is large, the first term in the Morawetz identity is almost $2E$ when $t_1 \leq -R$ and $t_2 \leq R$, thus all other terms must be small.

Corollary 3.2. *Let u be a solution to (CP1) with initial data $(u_0, u_1) \in \left(\dot{H}^1(\mathbb{R}^d) \cap L^{p+1}(\mathbb{R}^d)\right) \times L^2(\mathbb{R}^d)$. Given any $R > 0, r \geq 0$ we have*

$$\sum_{j=1}^6 M_j \leq \int_{\mathbb{R}^d} \min\{|x|/R, 1\} \left(|\nabla u_0|^2 + |u_1|^2 + \frac{2}{p+1} |u_0|^{p+1} \right) dx.$$

The notations M_j are defined by

$$\begin{aligned} M_1 &= \frac{1}{2R} \int_{R < |t| < R+r} \int_{|x| < R} \left(|\nabla u|^2 + |u_t|^2 + \frac{(d-1)(p-1)-2}{p+1} |u|^{p+1} \right) dx dt; \\ M_2 &= \frac{(d-1)(p-1)-4}{2(p+1)R} \int_{-R}^R \int_{|x| < R} |u|^{p+1} dx dt; \\ M_3 &= \frac{d-1}{4R^2} \int_{-R-r}^{R+r} \int_{|x|=R} |u|^2 d\sigma_R(x) dt; \\ M_4 &= \int_{-R-r}^{R+r} \int_{|x| > R} \left(\frac{|\nabla u|^2}{|x|} + \frac{(d-1)(p-1)}{2(p+1)} \frac{|u|^{p+1}}{|x|} + \frac{(d-3)(d-1)}{4} \frac{|u|^2}{|x|^3} \right) dx dt; \\ M_5 &= \sum_{\pm} \int_{|x| < R} \left(\frac{R^2 - |x|^2}{2R^2} |u_r|^2 + \frac{1}{2} \left| \frac{|x|}{R} u_r + \frac{(d-1)u}{2R} \pm u_t \right|^2 + \frac{(d^2-1)|u|^2}{8R^2} + \frac{|\nabla u|^2}{2} + \frac{|u|^{p+1}}{p+1} \right) \Big|_{t=\pm(R+r)} dx; \\ M_6 &= \sum_{\pm} \int_{|x| > R} \left(\frac{1}{2} \left| u_r + \frac{(d-1)u}{2|x|} \pm u_t \right|^2 + \frac{|\nabla u|^2}{2} + \frac{1}{p+1} |u|^{p+1} + \frac{(d-1)(d-3)}{8} \frac{|u|^2}{|x|^2} \right) \Big|_{t=\pm(R+r)} dx. \end{aligned}$$

Proof. We first choose $t_1 = -R - r, t_2 = R + r$ in the Morawetz identity, the first term above can be written as a sum of three terms

$$\begin{aligned} & \frac{1}{2R} \int_{-R-r}^{R+r} \int_{|x| < R} \left(|\nabla u|^2 + |u_t|^2 + \frac{(d-1)(p-1)-2}{p+1} |u|^{p+1} \right) dx \\ &= M_1 + \frac{1}{2R} \int_{-R}^R \int_{|x| < R} \left(|\nabla u|^2 + |u_t|^2 + \frac{(d-1)(p-1)-2}{p+1} |u|^{p+1} \right) dx \\ &= M_1 + M_2 + \frac{1}{2R} \int_{-R}^R \int_{|x| < R} \left(|\nabla u|^2 + |u_t|^2 + \frac{2}{p+1} |u|^{p+1} \right) dx. \end{aligned}$$

Thus

$$\frac{1}{2R} \int_{-R}^R \int_{|x| < R} \left(|\nabla u|^2 + |u_t|^2 + \frac{2}{p+1} |u|^{p+1} \right) dx dt + \sum_{j=1}^6 M_j = 2E. \quad (13)$$

In order to prove the first inequality we only need to show

$$\begin{aligned} I &= 2E - \frac{1}{2R} \int_{-R}^R \int_{|x| < R} \left(|\nabla u|^2 + |u_t|^2 + \frac{2}{p+1} |u|^{p+1} \right) dx dt \\ &\leq \int_{\mathbb{R}^d} \min\{|x|/R, 1\} \left(|\nabla u_0|^2 + |u_1|^2 + \frac{2}{p+1} |u_0|^{p+1} \right) dx. \end{aligned}$$

This follows energy conservation law and finite speed of propagation of energy

$$\begin{aligned}
I &= \frac{1}{R} \int_{-R}^R \int_{|x|>R} \left(\frac{1}{2} |\nabla u|^2 + \frac{1}{2} |u_t|^2 + \frac{1}{p+1} |u|^{p+1} \right) dx dt \\
&\leq \frac{1}{R} \int_{-R}^R \int_{|x|>R-|t|} \left(\frac{1}{2} |\nabla u_0|^2 + \frac{1}{2} |u_1|^2 + \frac{1}{p+1} |u_0|^{p+1} \right) dx dt \\
&= \frac{1}{R} \int_{\mathbb{R}^d} \min\{|x|, R\} \left(|\nabla u_0|^2 + |u_1|^2 + \frac{2}{p+1} |u_0|^{p+1} \right) dx.
\end{aligned}$$

□

4 Energy Distribution

In this section we prove Theorem 1.1. It suffices to consider the positive time direction $t > 0$, since the wave equation is time-reversible.

We choose $R = t, r = 0$ in corollary 3.3. The following inequalities hold for large time $t > 0$.

$$\begin{aligned}
&\sum_{\pm} \int_{|x|<t} \left(\frac{t^2 - |x|^2}{2t^2} |u_r|^2 + \frac{1}{2} \left| \frac{|x|}{t} u_r + \frac{(d-1)u}{2t} \pm u_t \right|^2 + \frac{(d^2-1)|u|^2}{8t^2} + \frac{|\nabla u|^2}{2} + \frac{|u|^{p+1}}{p+1} \right) dx \\
&+ \sum_{\pm} \int_{|x|>t} \left(\frac{1}{2} \left| u_r + \frac{(d-1)u}{2|x|} \pm u_t \right|^2 + \frac{|\nabla u|^2}{2} + \frac{1}{p+1} |u|^{p+1} + \frac{(d-1)(d-3)}{8} \frac{|u|^2}{|x|^2} \right) dx \\
&\leq \frac{4 - (d-1)(p-1)}{2(p+1)t} \int_{-t}^t \int_{|x|<t} |u(x, t')|^{p+1} dx dt' + \int_{\mathbb{R}^d} \min\{|x|/R, 1\} \left(|\nabla u_0|^2 + |u_1|^2 + \frac{2}{p+1} |u_0|^{p+1} \right) dx.
\end{aligned} \tag{14}$$

We observe that

$$|\nabla u|^2 + |u_t|^2 \lesssim_1 |u_r|^2 + \left| \frac{|x|}{t} u_r + \frac{(d-1)u}{2t} + u_t \right|^2 + \frac{d^2-1}{8} \frac{|u|^2}{t^2} + |\nabla u|^2$$

and

$$\begin{aligned}
|u_r + u_t|^2 &\lesssim_1 \frac{(t - |x|)^2}{t^2} |u_r|^2 + \left| \frac{|x|}{t} u_r + \frac{(d-1)u}{2t} + u_t \right|^2 + \frac{d^2-1}{8} \frac{|u|^2}{t^2}, \quad \text{if } |x| < t; \\
|u_r + u_t|^2 &\lesssim_1 \left| u_r + \frac{(d-1)u}{2|x|} + u_t \right|^2 + \frac{(d-1)(d-3)}{8} \frac{|u|^2}{|x|^2}, \quad \text{if } |x| > t.
\end{aligned}$$

Then obtain

$$\begin{aligned}
&\frac{1}{p+1} \int_{\mathbb{R}^d} (|u(x, t)|^{p+1} + |u(x, -t)|^{p+1}) dx + c_1 \sum_{\pm} \int_{|x|<t} \frac{t - |x|}{t} (|\nabla u(x, \pm t)|^2 + |u_t(x, \pm t)|^2) dx \\
&+ c_2 \sum_{\pm} \int_{\mathbb{R}^d} (|(u_r \pm u_t)(x, \pm t)|^2 + |\nabla u(x, \pm t)|^2) dx \\
&\leq \frac{4 - (d-1)(p-1)}{2(p+1)t} \int_{-t}^t \int_{|x|<t} |u(x, t')|^{p+1} dx dt' + 2 \int_{\mathbb{R}^d} \min\{|x|/t, 1\} e(x, 0) dx.
\end{aligned}$$

Here $c_1, c_2 > 0$ are absolute constants. For convenience we introduce the notation

$$\begin{aligned}
Q(t) &= \frac{1}{p+1} \int_{\mathbb{R}^d} (|u(x, t)|^{p+1} + |u(x, -t)|^{p+1}) dx + c_1 \sum_{\pm} \int_{|x|<t} \frac{t - |x|}{t} (|\nabla u(x, \pm t)|^2 + |u_t(x, \pm t)|^2) dx \\
&+ c_2 \sum_{\pm} \int_{\mathbb{R}^d} (|(u_r \pm u_t)(x, \pm t)|^2 + |\nabla u(x, \pm t)|^2) dx.
\end{aligned}$$

Then the inequality above implies that $Q(t)$ satisfies the recurrence formula

$$Q(t) \leq \frac{\lambda}{t} \int_0^t Q(t') dt' + 2 \int_{\mathbb{R}^d} \min\{|x|/t, 1\} e(x, 0) dx.$$

Here

$$0 < \lambda = \frac{4 - (d-1)(p-1)}{2} < 1.$$

Proof of part (a)(b) We may rewrite the recurrence formula as

$$Q(t) \leq \frac{\lambda}{t} \int_0^t Q(t') dt' + o(1).$$

We may take upper limits of both sides and obtain an inequality

$$\limsup_{t \rightarrow +\infty} Q(t) \leq \limsup_{t \rightarrow +\infty} \frac{\lambda}{t} \int_0^t Q(t') dt' \leq \lambda \limsup_{t \rightarrow +\infty} Q(t).$$

We recall the fact $\lambda \in (0, 1)$ and observe that $Q(t) \lesssim E$ is uniformly bounded, therefore we have

$$\limsup_{t \rightarrow +\infty} Q(t) = 0.$$

This verifies (a)(b).

Proof of part (c) Now we assume that initial data satisfy additional decay assumption. We start by multiplying both sides by $t^{\kappa-1}$ and integrate from $t = 1$ to $t = T$, utilize our assumption on initial data, then obtain

$$\begin{aligned} \int_1^T t^{\kappa-1} Q(t) dt &\leq \int_1^T t^{\kappa-1} \left(\frac{\lambda}{t} \int_0^t Q(t') dt' \right) dt + C_\kappa \int_{\mathbb{R}^d} \min\{|x|, |x|^\kappa\} e(x, 0) dx \\ &\leq \frac{\lambda}{1-\kappa} \int_0^T \min\{(t')^{\kappa-1}, 1\} Q(t') dt' + C_\kappa \int_{\mathbb{R}^d} \min\{|x|, |x|^\kappa\} e(x, 0) dx \\ &\leq \frac{\lambda}{1-\kappa} \int_1^T (t')^{\kappa-1} Q(t') dt' + C_\kappa \int_{\mathbb{R}^d} \min\{|x|, |x|^\kappa\} e(x, 0) dx + C_k E. \end{aligned}$$

Here

$$\frac{\lambda}{1-\kappa} = \frac{4 - (d-1)(p-1)}{2(1-\kappa)} < 1,$$

since we have assumed $\kappa < \frac{(d-1)(p-1)-2}{2}$. Therefore we have

$$\int_1^T t^{\kappa-1} Q(t) dt \lesssim_{p,\kappa} \int_{\mathbb{R}^d} \min\{|x|, |x|^\kappa\} e(x, 0) dx + E.$$

Because neither the right hand side nor the implicit constant here depends on T , we make $T \rightarrow +\infty$ to conclude

$$\int_1^\infty t^{\kappa-1} Q(t) dt < +\infty.$$

Combining this with the fact $Q(t) \lesssim E$, we have

$$\int_0^\infty t^{\kappa-1} Q(t) dt < +\infty.$$

We may multiply both sides of the recurrence formula by t^κ :

$$t^\kappa Q(t) \leq \lambda \int_0^t t^{\kappa-1} Q(t') dt' + 2 \int_{\mathbb{R}^d} \min\{|x|/t^{\kappa-1}, t^\kappa\} e(x, 0) dx.$$

Finally we apply dominated convergence theorem to finish the proof of Theorem 1.1.

Remark 4.1. When $d > 3$, we have

$$|u_r + u_t|^2 \lesssim_1 \left| u_r + \frac{(d-1)u}{2|x|} + u_t \right|^2 + \frac{(d-1)(d-3)}{8} \frac{|u|^2}{|x|^2}, \quad \text{if } |x| > t.$$

But the term $\frac{(d-1)(d-3)}{8} \frac{|u|^2}{|x|^2} = 0$ when $d = 3$, so the inequality above does not hold. We can redefine

$$\begin{aligned} Q(t) = & \frac{1}{p+1} \int_{\mathbb{R}^d} (|u(x, t)|^{p+1} + |u(x, -t)|^{p+1}) dx + c_1 \sum_{\pm} \int_{|x| < t} \frac{t - |x|}{t} (|\nabla u(x, \pm t)|^2 + |u_t(x, \pm t)|^2) dx \\ & + c_2 \sum_{\pm} \int_{|x| < t} (|(u_r \pm u_t)(x, \pm t)|^2 + |\nabla u(x, \pm t)|^2) dx \\ & + c_2 \sum_{\pm} \int_{|x| > t} \left(\left| u_r + \frac{(d-1)u}{2|x|} \pm u_t \right|^2 + |\nabla u(x, \pm t)|^2 \right) dx. \end{aligned}$$

We also have

$$\limsup_{t \rightarrow +\infty} Q(t) = 0,$$

and

$$t^\kappa Q(t) \leq \lambda \int_0^t t^{\kappa-1} Q(t') dt' + 2 \int_{\mathbb{R}^d} \min\{|x|/t^{\kappa-1}, t^\kappa\} e(x, 0) dx.$$

Because

$$|u_r + u_t|^2 \lesssim_1 \left| u_r + \frac{(d-1)u}{2|x|} + u_t \right|^2 + c \frac{|u|^2}{|x|^2}, \quad \text{if } |x| > t.$$

We can finish the proof of Theorem 1.1 by an estimate.

$$\begin{aligned} \int_{|x| > t} \frac{|u|^2}{|x|^2} dx & \lesssim \left(\int_{|x| > t} (|u|^2)^{\frac{p+1}{2}} dx \right)^{\frac{2}{p+1}} \left(\int_{|x| > t} (|x|^{-2})^{\frac{p+1}{p-1}} dx \right)^{\frac{p-1}{p+1}} \\ & \lesssim_p t^{\frac{-2\kappa}{p+1} + \frac{p-5}{p+1}}. \end{aligned}$$

When $d = 3$, we have $p \in (2, 3)$ and the inequality

$$t^{\frac{-2\kappa}{p+1} + \frac{p-5}{p+1}} \ll t^{-\kappa},$$

thus

$$\int_{|x| > t} \frac{|u|^2}{|x|^2} dx \ll t^{-\kappa}.$$

5 Scattering Theory

5.1 Transformation to 1D

In order to take full advantage of our radial assumption, we use the following transformation: if u is a radial solution to (CP1), then $w(r, t) = r^{\frac{d-1}{2}} u(r, t)$, where $|x| = r$, is a solution to one-dimensional wave equation

$$(\partial_t + \partial_r)(w_t - w_r) = \partial_t^2 w - \partial_r^2 w = -\frac{(d-1)(d-3)}{4} r^{\frac{d-5}{2}} u - r^{\frac{d-1}{2}} |u|^{p-1} u.$$

We define

$$v_+(r, t) = w_t(r, t) - w_r(r, t); \quad v_-(r, t) = w_t(r, t) + w_r(r, t).$$

A simple calculation shows that v_{\pm} satisfy the equation

$$(\partial_t \pm \partial_r) v_{\pm}(r, t) = \partial_t^2 w - \partial_r^2 w = -\frac{(d-1)(d-3)}{4} r^{\frac{d-5}{2}} u - r^{\frac{d-1}{2}} |u|^{p-1} u.$$

This gives variation of v_{\pm} along characteristic lines $t \pm r = \text{Const.}$

$$v_+(t_2 - \eta, t_2) - v_+(t_1 - \eta, t_1) = \int_{t_1}^{t_2} f(t - \eta, t) dt, \quad t_2 > t_1 > \eta; \quad (15)$$

$$v_-(s - t_2, t_2) - v_-(s - t_1, t_1) = \int_{t_1}^{t_2} f(s - t, t) dt, \quad t_1 < t_2 < s. \quad (16)$$

Here the function $f(r, t)$ is defined by

$$f(r, t) = -\frac{(d-1)(d-3)}{4} r^{\frac{d-5}{2}} u(r, t) - r^{\frac{d-1}{2}} |u|^{p-1} u(r, t).$$

Then we give the upper bounds of the integral above. According to Lemma 2.1 we have

$$\begin{aligned} & \int_{t_1}^{t_2} (t - \eta)^{\frac{d-5}{2}} |u(t - \eta, t)| dt \\ & \leq \left\{ \int_{t_1}^{t_2} \left[(t - \eta)^{\frac{d-3}{2}} |u(t - \eta, t)| \right]^2 dt \right\}^{1/2} \left\{ \int_{t_1}^{t_2} [(t - \eta)^{-1}]^2 dt \right\}^{1/2} \\ & \leq \left\{ \int_{\eta}^{\infty} (t - \eta)^{d-3} |u(t - \eta, t)|^2 dt \right\}^{1/2} (t_1 - \eta)^{-1/2} \\ & \lesssim_d E^{1/2} (t_1 - \eta)^{-1/2}. \end{aligned}$$

In addition we have

$$\begin{aligned} & \int_{t_1}^{t_2} (t - \eta)^{\frac{d-1}{2}} |u(t - \eta, t)|^p dt \\ & \leq \left\{ \int_{t_1}^{t_2} \left[(t - \eta)^{\frac{(d-1)p}{p+1}} |u(t - \eta, t)|^p \right]^{\frac{p+1}{p}} dt \right\}^{\frac{p}{p+1}} \left\{ \int_{t_1}^{t_2} \left[(t - \eta)^{-\frac{(d-1)(p-1)}{2(p+1)}} \right]^{p+1} dt \right\}^{\frac{1}{p+1}} \\ & \leq \left\{ \int_{\eta}^{\infty} (t - \eta)^{d-1} |u(t - \eta, t)|^{p+1} dt \right\}^{\frac{p}{p+1}} \left\{ \int_{t_1}^{t_2} (t - \eta)^{-\frac{(d-1)(p-1)}{2}} dt \right\}^{\frac{1}{p+1}} \\ & \lesssim_d E^{\frac{p}{p+1}} (t_1 - \eta)^{-\frac{(d-1)(p-1)-2}{2(p+1)}}. \end{aligned}$$

we combine these estimates above with (15) and (16) to obtain

Lemma 5.1. *Let u be a radial solution wave equation with a finite energy E . Then we have*

$$\begin{aligned} |v_+(t_2 - \eta, t_2) - v_+(t_1 - \eta, t_1)| & \lesssim_d E^{1/2} (t_1 - \eta)^{-1/2} + E^{\frac{p}{p+1}} (t_1 - \eta)^{-\beta(d,p)/2}; \\ |v_-(s - t_2, t_2) - v_-(s - t_1, t_1)| & \lesssim_d E^{1/2} (s - t_2)^{-1/2} + E^{\frac{p}{p+1}} (s - t_2)^{-\beta(d,p)/2}. \end{aligned}$$

Here $\beta(d, p) = \frac{(d-1)(p-1)-2}{p+1}$.

5.2 Scattering by energy decay

In this section we prove Theorem 1.2. Let us recall the lemma 5.1, we obtain there exists a function $g_+(\eta) \in L^2(\mathbb{R})$ with $\|g_+\|_{L^2(\mathbb{R})}^2 \leq E/c_d$, so that

$$v_+(t - \eta, t) \rightarrow 2g_+(\eta) \quad \text{in } L_{loc}^2(\mathbb{R}), \quad \text{as } t \rightarrow +\infty.$$

The asymptotic behaviour of v_- is similar as $t \rightarrow -\infty$

$$v_-(s-t, t) \rightarrow 2g_-(s) \quad \text{in } L_{loc}^2(\mathbb{R}), \quad \text{as } t \rightarrow -\infty,$$

let $t_2 \rightarrow +\infty$ in the first inequality of lemma 5.1, we have

$$|2g_+(\eta) - v_+(t-\eta, t)| \lesssim_{d,p,E} (t-\eta)^{-\beta(d,p)/2}, \quad \eta < t-1.$$

We apply a change of variable $r = t - \eta$ and rewrite this in the form

$$|v_+(r, t) - 2g_+(t-r)| \lesssim_{d,p,E} r^{-\beta(d,p)/2}, \quad r > 1.$$

Similarly we have

$$|v_-(r, t) - 2g_-(t+r)| \lesssim_{d,p,E} r^{-\beta(d,p)/2}, \quad r > 1.$$

These gives the following upper limits

$$\limsup_{t \rightarrow +\infty} \int_{t-c \cdot t^{\beta(d,p)}}^{t+R} \left(|v_+(r, t) - 2g_+(t-r)|^2 + |v_-(r, t) - 2g_-(t+r)|^2 \right) dr \lesssim_{d,p,E} c. \quad (17)$$

We ignore $g_-(t+r)$ in the upper limits above because

$$\lim_{t \rightarrow +\infty} \int_0^\infty |g_-(t+r)|^2 dr = \lim_{t \rightarrow +\infty} \int_t^\infty |g_-(s)|^2 ds = 0.$$

We recall $v_\pm = w_t \mp w_r$ and rewrite the upper limits above in term of w

$$\limsup_{t \rightarrow +\infty} \int_{t-c \cdot t^{\beta(d,p)}}^{t+R} \left(|w_r(r, t) + g_+(t-r)|^2 + |w_t(r, t) - g_+(t-r)|^2 \right) dr \lesssim_{d,p,E} c.$$

Next we utilize the identities $r^{\frac{d-1}{2}} u_r = w_r - (d-1)r^{\frac{d-3}{2}} u/2$, $r^{\frac{d-1}{2}} u_t = w_t$ and a direct consequence of the pointwise estimate $|u(r, t)| \lesssim_{d,E} r^{-\frac{2(d-1)}{p+3}}$ (lemma 2.1) to conclude

$$\limsup_{t \rightarrow +\infty} \int_{t-c \cdot t^{\beta(d,p)}}^{t+R} \left(\left| r^{\frac{d-1}{2}} u_r(r, t) + g_+(t-r) \right|^2 + \left| r^{\frac{d-1}{2}} u_t(r, t) - g_+(t-r) \right|^2 \right) dr \lesssim_{d,p,E} c. \quad (18)$$

By lemma 1.2 (radiation fields), there exists a radial free wave \tilde{u}^+ , so that

$$\lim_{t \rightarrow +\infty} \int_0^\infty \left(\left| r^{\frac{d-1}{2}} \tilde{u}_r^+(r, t) + g_+(t-r) \right|^2 + \left| r^{\frac{d-1}{2}} \tilde{u}_t^+(r, t) - g_+(t-r) \right|^2 \right) dr = 0. \quad (19)$$

Therefore we have

$$\limsup_{t \rightarrow +\infty} \int_{t-c \cdot t^{\beta(d,p)}}^{t+R} r^{d-1} \left(|u_t(r, t) - \tilde{u}_t^+(r, t)|^2 + |u_r(r, t) - \tilde{u}_r^+(r, t)|^2 \right) dr \lesssim_{d,p,E} c. \quad (20)$$

Finite speed of propagation of energy implies

$$\limsup_{R \rightarrow \infty} \int_{t+R}^\infty r^{d-1} \left(|u_t(r, t) - \tilde{u}_t^+(r, t)|^2 + |u_r(r, t) - \tilde{u}_r^+(r, t)|^2 \right) dr = 0. \quad (21)$$

We combine (20) with (21) and obtain

$$\limsup_{t \rightarrow +\infty} \int_{t-c \cdot t^{\beta(d,p)}}^{+\infty} r^{d-1} \left(|u_t(r, t) - \tilde{u}_t^+(r, t)|^2 + |u_r(r, t) - \tilde{u}_r^+(r, t)|^2 \right) dr \lesssim_{d,p,E} c. \quad (22)$$

Finally we consider the region $\{x : |x| < t - c \cdot t^{\beta(p)}\}$. We utilize the conclusion of Theorem 1.1.(c), and obtain

$$\lim_{t \rightarrow +\infty} \int_{|x| < t - c \cdot t^{\beta(d,p)}} e(x, t) dx \lesssim_c \lim_{t \rightarrow +\infty} t^{\frac{(2-d)p+(d+2)}{p+1}} \int_{|x| < t} \frac{t - |x|}{t} e(x, t) dx = 0.$$

Please note that in the sub-conformal range our assumption $p > \frac{3-d+2\sqrt{d^2-d+1}}{d-1}$ guarantees that $\frac{(2-d)p+(d+2)}{p+1} < \frac{(d-1)(p-1)-2}{2}$.

We also have

$$\lim_{t \rightarrow +\infty} \int_{|x| < t - ct^{\beta(p)}} \left(|\nabla \tilde{u}^+(x, t)|^2 + |\tilde{u}_t^+(x, t)|^2 \right) dx = 0.$$

Combining these two limits we obtain

$$\lim_{t \rightarrow +\infty} \int_{|x| < t - ct^{\beta(d,p)}} \left(|\nabla \tilde{u}^\pm(x, t) - \nabla u(x, t)|^2 + |\tilde{u}_t^\pm(x, t) - u_t(x, t)|^2 \right) dx = 0.$$

We combine this with stronger exterior scattering to conclude

$$\limsup_{t \rightarrow +\infty} \int_{\mathbb{R}^d} \left(|\nabla \tilde{u}^\pm(x, t) - \nabla u(x, t)|^2 + |\tilde{u}_t^\pm(x, t) - u_t(x, t)|^2 \right) dx \lesssim_{d,p,E} c.$$

We make $c \rightarrow 0^+$ and finish the proof.

Acknowledgement

The second author is financially supported by National Natural Science Foundation of China Projects 12071339, 11771325.

References

- [1] L.Kapitanski. “Weak and yet weaker solutions of semilinear wave equations.” *Communications in Partial Differential Equations* 19(1994):1629-1676.
- [2] H.Lindblad, and C.Sogge. “On existence and scattering with minimal regularity for semilinear wave equations.” *Journal of Functional Analysis* 130(1995):357-426.
- [3] C.S.Morawetz. “Time decay for the nonlinear Kelein-Gordon equations.” *Proceedings of the Royal Society.London. Series A* 306(1968):291-296.
- [4] C.E.King, and F.Merle. “Nondispersive radial solutions to energy supercritical non-linear wave equations, with applications.” *American Journal of Mathematics* 133, No 4(2011):1029-1065.
- [5] R.Shen. “On the energy subcritical, nonlinear wave equation in R^3 with radial data.” *Analysis and PDE* 6(2013):1929-1987.
- [6] B.Dodson and A.Lawrie. “Scattering for the radial 3d cubic wave equation.” *Analysis and PDE* 8(2015):467-497.
- [7] J.Lin, and W.Strauss. “Decay and scattering of solutions of a nonlinear Schrödinger equation.” *Journal of Functional Analysis* 30(1987): 245-263.
- [8] J.Colliander, M.Keel, G.Staffilani, H.Takaoka, and T.Tao. “Global existence and scattering for rough solutions to a nonlinear Schrödinger equation in R^3 .” *Communications on Pure and Applied Mathematics* 57(2004):987-1014.

- [9] R.Shen. “Long time behaviour of finite-energy radial solutions to energy subcritical wave equation in higher dimensions.” *arXiv Preprint* 1912.12913.
- [10] B.Dodson. “Global well-posedness and scattering for the radial, defocusing, cubic nonlinear wave equation.” *arXiv Preprint* 1809.08284.
- [11] F.G.Friedlander. “On the radiation field of pulse solutions of the wave equations of the wave equation.” *Proceeding of the Royal Society A* 269(1962):53-65.
- [12] F.G.Friedlander. “Radiation fields and hyperbolic scattering theory.” *Mathematical Proceedings of Cambridge Philosophical Society* 88(1980):483-515.
- [13] T.Duyckaerts, C.E.Kening, and F.Merle. “Scattering profile for global solutions of the energy-critical wave equation.” *Journal of European Mathematical Society* 21(2019):2117-2162.
- [14] R.Killip, and M.Visan. “The defocusing energy-supercritical nonlinear wave equation in three space dimensions.” *Transactions of The American Mathematical Society* 363(2011):3893-3934.
- [15] R.Killip, and M.Visan. “The radial defocusing energy-supercritical nonlinear wave equation in all space dimensions.” *Proceedings of the American Mathematical Society* 139(2011):1805-1817.
- [16] C.Rodriguez. “Scattering for radial energy-subcritical wave equations in dimensions 4 and 5.” *Communications in Partial Differential Equations* 42(2017):852-894.
- [17] R.Shen. “Energy distribution of radial solutions to energy subcritical wave equation with an application on scattering theory.” *Transactions of The American Mathematical Society* 374 (2021), 3827-3857.
- [18] J.Ginibre, and G.Velo, “Conformal invariance and time decay for nonlinear wave equations.” *Annales de l’institut Henri Poincaré (A) Physique théorique* 47(1987):221-276.
- [19] K.Hidano. “Conformal conservation law, time decay and scattering for nonlinear wave equation.” *Journal D’analysis Mathématique* 91(2003):269-295.
- [20] B.Perthame, and L.vega. “Morrey-Campanato estimates for Helmholtz equation.” *Journal of Functional Analysis*. 164(1999):340-355.
- [21] J.Ginibre and G.Velo. “The global Cauchy problem for the non linear Klein-Gordon equation.” *Math Z.* 189 (1985):487-505.