

# More on verification of probability forecasts for football outcomes: score decompositions, reliability, and discrimination analyses.

Jean-Louis Foulley

IMAG, Université de Montpellier, France : [foulley-jl@gmail.com](mailto:foulley-jl@gmail.com)

## Abstract

Forecast of football outcomes in terms of Home Win, Draw and Away Win relies largely on ex ante probability elicitation of these events and ex post verification of them via computation of probability scoring rules (Brier, Ranked Probability, Logarithmic, Zero-One scores). Usually, appraisal of the quality of forecasting procedures is restricted to reporting mean score values. The purpose of this article is to propose additional tools of verification, such as score decompositions into several components of special interest. Graphical and numerical diagnoses of reliability and discrimination and kindred statistical methods are presented using different techniques of binning (fixed thresholds, quantiles, logistic and iso regression). These procedures are illustrated on probability forecasts for the outcomes of the UEFA Champions League (C1) at the end of the group stage based on typical Poisson regression models with reasonably good results in terms of reliability as compared to those obtained from bookmaker odds and whatever the technique used. Links with research in machine learning and different areas of application (meteorology, medicine) are discussed.

## 1 Introduction

The list of areas opening to forecast would be exceedingly long if one wishes to draw it up in details: from economic inflation, employment rates, weather and climate change, medical diagnosis and biological tests to media and entertainment market as well as gambling & sporting events among others. Forecasting consists first of producing forecasts from available data and methodologies, and second to assess their quality. Forecasts are basically of two types: pointwise or by means of probability. Quantify the uncertainty of a forthcoming event highlights the superiority of probability forecasts over categorical ones even economically (Winkler & Murphy, 1979; Dawid, 1986). As far as football matches are concerned, outcomes are in terms of either results such as Win, Draw or Loss (WDL)-also abbreviated as Home Win, Draw and Away Win (H,D,A respectively) for two-legged matches- i.e., categorical data, or in terms of scorelines  $\{Y(A), Y(B)\}$  goals in match (A vs B) i.e., pairs of integers: see the review by Read et al. (2021).

Here we will be concerned with probability forecasts of WDL (HAD) results. In this area, much effort has been devoted to statistical approaches to forecasting, especially by modelling outcomes of matches (Scarf and Selliti-Rangel, 2019). There is a growing demand for relevant and acute probabilistic forecasting due to the large audience for TV broadcasting of association football matches and the related huge betting markets. Traditionally in football, one summarizes HDA forecast performance by a synthetic criterium such

as the mean square error known as Brier's score (1950). Although this statistic brought useful information, it is only an average measure of the overall accuracy of such predictions. Analytical attributes of this accuracy can be brought out such as Reliability, Resolution, Discrimination, Refinement to shed more light on some qualities or deficiencies of issued forecasts. The purpose of this note is mostly pedagogical with the aim to review the main tools available to that respect, and to illustrate them on data of a well-known club competition with the UEFA champions League (also known in brief as C1) forecasted by simple Poisson regression models. Focus is on a distribution-oriented approach based on the joint distribution of elementary forecast, binary outcome pairs and its factorization. We will deal with several decompositions of the Brier score (BRS) applied to each binary outcome considered separately namely 1) the Murphy decomposition derived from the so-called Calibration-Refinement factorization (CR), and 2) the Likelihood-base (LB) and Yates's decompositions. In parallel to the CR decomposition, we will present a graphical device known as the reliability diagram which allows diagnostic of strength or deficiency in this component. In each case, comparison will be made between forecasts derived from a simple Poisson loglinear model (POI), and to Bookmaker Odds implied probabilities (ODD). Finally, in the last (4<sup>th</sup>) section, we discuss the main points raised by using these decompositions and possible implications for improving forecast efficiency.

## 2 Decomposition of the Brier score

### 2.1 Basic theory

Let  $X$  be the binary outcome of the event  $H$  with probability  $q$  and  $P$  the random variable probabilistic forecast of  $X$  taking values  $p$ . Taking as scoring rule, the quadratic or Half-Brier Score defined as the loss function  $S(P, X) = (P - X)^2$ , and using the conditioning de conditioning rule, the Murphy (1973) decomposition of its expectation can be written as

$$\mathbb{E}[S(P, X)] = \text{Var}(X) - \text{Var}_P[\mathbb{E}_X(X|P)] + \mathbb{E}_P\left\{\left[\mathbb{E}_X(X|P) - P\right]^2\right\}, \quad (1)$$

where expectation is taken with respect to the pairs of forecasts and outcomes  $P, X$ .

Examination of this formula immediately identifies the 3 components of this decomposition:

- 1) Uncertainty (UNC) equal to  $\text{Var}(X) = q(1-q)$ , the variance of the outcome that is out of control of the forecaster,
- 2) Resolution (RES) equal to  $\text{Var}_P[\mathbb{E}_X(X|P)]$  referring to the variability between the conditional expectations of the observed outcomes given their forecasts,
- 3) Reliability (REL) or (Mis) Calibration equal to  $\mathbb{E}_P\left\{\left[\mathbb{E}_X(X|P) - P\right]^2\right\}$  measuring the average squared differences between the conditional expectation of the outcome and its forecast.

Murphy and Winkler (1987) also gave the dual decomposition of (1)

$$\mathbb{E}[S(P, X)] = \text{Var}(P) - \text{Var}_X[\mathbb{E}_P(P|X)] + \mathbb{E}_X\left\{\left[\mathbb{E}_P(P|X) - X\right]^2\right\}. \quad (2)$$

This decomposition is known as the Likelihood-base factorization as opposed to the Calibration-Refinement factorization of formula (1) referring to the 2-way decomposition of the joint distribution of  $(P, X)$ . This formula leads to identify three components: i) Refinement (REF) equal to  $\text{Var}(P)$ , the variance of

probabilistic forecasts also known as Sharpness, ii) Discrimination (DIS) equal to  $\text{Var}_X [\mathbb{E}_P(P|X)]$  i.e., the variance between conditional distributions of forecasts given the outcomes  $X$ , iii) Conditional bias type 2 as called by Bradley et al, (2003) equal to  $\mathbb{E}_X \left\{ \left[ \mathbb{E}_P(P|X) - X \right]^2 \right\}$  which is the dual expression of resolution. A special case of interest is the one with  $P$  having a probability mass function (pmf) concentrated at  $p = \mathbb{E}(X)$ , the mean of the marginal distribution of the binary outcome (the so-called climatological forecast). Then the forecast has no refinement ( $\text{Var} P = 0$ ), no discrimination as well, necessarily no resolution, even though it is perfectly calibrated. In fact, it is the only forecast being both reliable with no discrimination (Bröcker, 2012). In that case,  $\mathbb{E}[S(P, X)]$  reduces to its uncertain component UNC as an upper reference value for the expected Brier score. That is the reason why the decomposition in (1) is often expressed as fractions of UNC and the complement to one of the scaled Brier Score (BS) as a Brier Skill Score

$$BSS = 1 - BS / BS_{ref} = (REL - RES) / UNC. \quad (3)$$

This formula clearly emphasizes the trade-off between these two components with the aim of increasing resolution without neglecting reliability.

There is another decomposition by Yates (1982) deserving as much attention as it defines influential components of the expected quadratic score which are easier to understand:

$$\begin{aligned} \mathbb{E}[S(P, X)] = & \text{Var}(X) - 2\text{Cov}(P, X) + \text{Var}_X [\mathbb{E}_P(P|X)] \\ & + \mathbb{E}_X [\text{Var}_P(P|X)] + [\mathbb{E}(P) - \mathbb{E}(X)]^2. \end{aligned} \quad (4)$$

This formula stems from the basic expression  $\mathbb{E}[S(P, X)] = \text{Var}(P - X) + [\mathbb{E}(P) - \mathbb{E}(X)]^2$  with  $\text{Var}(P) = \text{Var}_X [\mathbb{E}_P(P|X)] + \mathbb{E}_X [\text{Var}_P(P|X)]$  and  $\mathbb{E}(P|X)$  being the best predictor of  $P$  given  $X$ . As this predictor is a linear one:  $\mathbb{E}(P|X) = a + bX$  with  $a = \mathbb{E}(P|X = 0)$ ,  $b = \mathbb{E}(P|X = 1) - \mathbb{E}(P|X = 0)$ ,  $\text{Var}_X [\mathbb{E}_P(P|X)] = b^2 \text{Var}(X)$ ,  $\text{Cov}(P, X) = b \text{Var}(X)$ . These last two expressions highlight the key role of the regression coefficient  $b$  of  $P$  on  $X$  equal to the difference in expected value of forecasts pertaining to future positive outcomes from those out of negative ones.

Yates (1982) emphasized the different influence of the two components of  $\text{Var}(P)$ . The first one, which is the between class variance  $\text{Var}_X [\mathbb{E}_P(P|X)]$ , he qualified as “VarPmin”, is beneficial. The second one, which is the within class variance  $\mathbb{E}_X [\text{Var}_P(P|X)]$ , called “ΔVarP” or “Scattered Variance” representing the lack of sharpness of the distributions of  $P|X = x$ , is detrimental. The last term in (4) measures marginal bias and was called “Calibration or Reliability-in-Large” (CIL or RIL). In short, this decomposition is written as: UNC- 2COV+ VarPmin+ ΔVarP+ RIL. Actually, Yates’ and the LB decompositions are closely related although not subject to the same interpretation, with REF=VarPmin+ΔVarP, DIS= VarPmin and CB2=UNC-2COV+VarPmin+RIL.

Table 1: Calibration-Refinement decomposition of Brier's score pertaining to Home Win, Draw and Away Win under the Poisson regression model.

RESULT	BRS	SKI (%)	B-TEST	UNC	MET	REL	RES
HWIN	0.1849	24.8	1.035	0.2458	INT	0.0035 (1.4)	0.0644 (26.2)
			[0.309]		QUA	0.0030 (1.2)	0.0639 (26.0)
					ISO	<b>0.0116 (4.7)</b>	<b>0.0725 (29.5)</b>
DRAW	0.1849	1.4	3.995	0.1875	INT	0.0010 (0.5)	0.0036 (1.9)
			[0.045]		QUA	0.0031 (1.7)	0.0058 (3.1)
					ISO	<b>0.0099 (5.3)</b>	<b>0.0125 (6.7)</b>
AWIN	0.1700	21.2	0.001	0.2158	INT	0.0047 (2.2)	0.0505 (23.4)
			[0.975]		QUA	0.0029 (1.3)	0.0487 (22.5)
					ISO	<b>0.0078 (3.6)</b>	<b>0.0537 (24.9)</b>

BRS=REL-RES+UNC with REL: Reliability, RES: Resolution, UNC: Uncertainty according to different binning procedures (INT: Interval; QUA: Quantile; ISO-Regression).and expressed both in absolute value and p.100 of uncertainty (UNC).

Skill (SKI) defined as  $SKI=(BRS_{ref}-BRS)/BRS_{ref}$  where  $BRS_{ref}=UNC$  so that  $SKI=(RES-REL)/UNC$

B-TEST: Brier-Score Test for departure of its expectation from that induced by the null hypothesis of perfect forecast calibration expressed with its corresponding statistic and P-value within brackets.

## 2.2 Estimation from data

Practically, verification takes place from a data sample made of pairs  $\{(p_i, x_i), i=1, \dots, N\}$  of ex ante probabilistic forecasts  $p_i$  and ex post binary outcomes  $x_i$ . Most quantities introduced previously can be estimated by their regular moment estimators. The expected quadratic score  $\mathbb{E}[S(P, X)]$  is traditionally estimated by the empirical score:

$$\bar{S}(\mathbf{p}) = N^{-1} \sum_{i=1}^N S(p_i, x_i). \quad (5)$$

For the CR decomposition, REL and RES require estimations of  $\mathbb{E}(X | P)$ . If the forecasts take  $K$  distinct values  $\{p_k, k=1, \dots, K\}$  with  $n_k$  occurrences of binary outcomes  $X$ , then  $\mathbb{E}(X | P = p_k) = \bar{X}_k = X_{k+} / n_k$  with  $X_{k+} = \sum_{i=1}^N I(p_i = p_k) X_i$  and  $\bar{X} = (\sum_{i=1}^N X_i) / N$ . In such cases, the Murphy (1973) decomposition is fully applicable without restrictions:

$$REL = N^{-1} \sum_{k=1}^K n_k (\bar{x}_k - p_k)^2, \quad RES = N^{-1} \sum_{k=1}^K n_k (\bar{x}_k - \bar{x})^2, \quad UNC = \bar{x}(1 - \bar{x}). \quad (6)$$

In fact, in many applications as in forecasting Football match results, we stay in-between discrete and continuous distributions, facing many distinct forecast values. In such cases, forecasts have to be distributed into intervals named bins  $B_1, \dots, B_d, \dots, B_D$  and averaged within bins i.e., letting  $I_d = \{i : p_i \in B_d\}$ , pairs  $\{(p_d, x_d), d=1, \dots, D\}$  are computed as  $p_d = n_d^{-1} \sum_{i \in I_d} p_i$ ,  $\hat{x}_d = x_{d+} / n_d$  where  $n_d = \#I_d$ ,  $\hat{x}_{d+} = n_d^{-1} \sum_{i \in I_d} x_i$ . To avoid inconsistencies in the CR decomposition, two extra components of within bin variance and covariance must be added to those in (6). We skip such complications by adopting a simple procedure as advocated by Siegert (2017). Letting as in (5)  $\bar{S}(\hat{\mathbf{x}}) = N^{-1} \sum_{i=1}^N S(f_i, \hat{x}_i)$  and  $\bar{S}(\bar{x} \mathbf{1}_N) = N^{-1} \sum_{i=1}^N S(f_i, \bar{x})$  with  $\mathbf{1}_N$ , the unit vector of size N, the components of the mean score  $\bar{S}(\mathbf{p})$  reduce to

$$REL = \bar{S}(\mathbf{p}) - \bar{S}(\hat{\mathbf{x}}), \quad RES = \bar{S}(\bar{x} \mathbf{1}_N) - \bar{S}(\hat{\mathbf{x}}), \quad UNC = \bar{S}(\bar{x} \mathbf{1}_N). \quad (7)$$

This decomposition automatically satisfies the equality  $\bar{S}(\mathbf{p}) = REL - RES + UNC$ , and is equivalent to the original Murphy decomposition in the case of distinct discrete forecasts. It also ensures that i) resolution is nil when  $\mathbf{p}$  is perfectly calibrated ( $\mathbf{p} = \hat{\mathbf{x}}$ ), and ii) the constant climatological forecast  $\mathbf{p} = \bar{x}\mathbf{1}_N$  is the only forecast satisfying  $RES=REL=0$ . Finally, it is potentially applicable to other proper probabilistic scoring rules, as the ignorance score  $L(P, X) = -X \log(P) - (1 - X) \log(1 - P)$  (Dawid, 1986; Bröcker, 2012). Moreover, the statistic  $2N \left[ \bar{L}(\mathbf{p}) - \bar{L}(\hat{q}_p) \right]$  is an analog of the log-likelihood ratio statistic for a perfectly reliable forecast having an asymptotic Chi-square distribution with degrees of freedom equal to the number of parameters specifying the model for  $q_p = \Pr(X = 1 | P = p)$ .

Different binning techniques are available such as fixed threshold intervals and fixed quantile intervals with potential optimization of their number (Bröcker, 2012; Gweon and Yu, 2019). A promising one relies on the non-parametric isotonic regression implemented via the pool-adjacent-violators (PAV) algorithm with optimality properties (Dimitriadis et al., 2021).

It also provides a reliability diagram featuring graphically the CR decomposition of the joint distribution  $[P, X] = [P][X | P]$  of binned data by the marginal (refinement) distribution of forecasts and plots of (re) calibrated probabilities  $\hat{x}_d$  against automatically binned forecasts  $p_d$ .

Table 2: Calibration analysis via fitting a logistic model of the probability of Homewin, Draw and Awaywin (AWIN) on the logit of its probabilistic forecast under a Poisson regression model (POI)

Category	Criterion	Estimation	SE	T-Statistics	DF	P-value
Homewin	intercept	-0.259	0.119	4.700	1	0.030
	slope	1.113	0.129	0.765	1	0.382
	D0 vs D1	423.085 vs 417.489		5.596	2	0.061
Draw	intercept	0.153	0.466	0.108	1	0.742
	slope	0.932	0.346	0.039	1	0.843
	D0 vs D1	426.981 vs 422.981		4.000	2	0.135
Awaywin	intercept	0.076	0.149	0.261	1	0.610
	slope	1.053	0.134	0.156	1	0.693
	D0 vs D1	389.458 vs 389.176		0.282	2	0.870

Intercept ( $\alpha$ ) and slope ( $\beta$ ) of the logit regression model with their estimation and standard error (SE). Deviance  $D(k) = -2L(k)$  where  $L(k)$  is the loglikelihood of the null model (0:  $\alpha = 0; \beta = 1$ ) vs the unspecified parameter model (1:  $\alpha \neq 0; \beta \neq 0$ ); T-statistics: Wald for intercept=0 and slope=1; Deviance differences  $\Delta D = D0 - D1$  and their corresponding degrees of freedom (DF) and P-values

To that respect, another way to assess Reliability via the conditional distribution  $[X | P]$  of outcomes  $X$  given  $P$  is through a regression model, but in the framework of logistic instead of linear regression chosen by Reade et al. (2021). Following Cox (1958), the model relating  $X$  to  $P$  is written via a logit linear predictor:  $\text{logit}[\Pr(X_i = 1)] = \alpha + \beta \text{logit}(p_i)$  with  $\alpha = 0$  and  $\beta = 1$  for perfect reliability and typical patterns of reliability diagrams with i) ( $\alpha > 0, \beta = 1$ ) for concave under-forecasting profiles, ii) ( $\alpha < 0, \beta = 1$ ) for convex over-forecasting profiles as well as iii) ( $\alpha = 0, \beta > 1$ ) for sigmoid, and iv) ( $\alpha = 0, \beta < 1$ ) for inverse-sigmoid profiles. Statistical tests are available (Wald and likelihood ratio tests) for challenging the different hypotheses about such patterns.

### 3 Application

The purpose of this illustration is to assess the performance of probability forecasts of outcomes of the UEFA champions league (the so called C1) matches played during the group stage (GS). Four seasons were considered from 2017 to 2020. Forecast is based on a simple log linear Poisson regression model applied to score lines with intercept, home effect and two time-dependent ELO team covariates. The models was fitted to ex ante data, namely score lines of all the matches played during the 3 previous seasons e.g., 2017, 2018 and 2019 as training sample to forecasts of the 96 GS matches of 2020; the same applies to forecasts of the 2019 GS based on 2016, 2017 and 2018 seasons and so on. Inference about parameters of the Poisson loglinear model is based on posterior distributions and probability forecasts are obtained as expectations of predictive distributions. Computations are carried out via the Win/OpenBUGS software. As top reference, we considered the classical Bookmaker Odds (ODD) as 3-Way Odds implied Probabilities with probabilities derived as  $p_{m,j} = o_{m,j}^{-1} / \sum_{k=1}^3 o_{m,k}^{-1}$  where  $o_{m,j}$  is the betting odd for  $j=1,2,3$  (WDL) edited by OddsPortal (here an average of 10 to 12 odds from well-known betting companies).

Table 3: Characteristics of conditional distributions of probability forecasts given the outcomes under two Forecasting procedures: Poisson regression (POI) and Odds Probabilities (ODD)

Method		Home Win		Draw		Away Win	
		POI	ODD	POI	ODD	POI	ODD
Sample sizes		217-167		288-96		263-121	
Mean %	X=0	37.88	35.01	20.22	20.97	24.30	23.24
	X=1	63.08	62.73	22.34	23.89	44.84	48.62
	Dif 1-0	24.20	27.71	2.02	2.93	20.54	25.38
Wilcoxon	Z	9.93	10.76	3.59	3.55	9.09	10.13
	P-val	<0.0001	<0.0001	0.0002	0.0002	<0.0001	<0.0001
KS	D	0.473	0.511	0.236	0.236	0.447	0.521
	P-val	<0.0001	<0.0001	0.0007	0.0007	<0.0001	<0.0001
C-statistic	Estimation	0.795	0.820	0.622	0.624	0.789	0.820

Sample sizes of forecasts having X=0 vs X=1 respectively; Z: Normal approximation of the Wilcoxon-test with one sided P value; KS: Kolmogorov-Smirnoff two sample test on Max [F(X=0)-F(X=1)] C-statistic: Harrell's concordance index varying from 0.5 (no discrimination) to 1 (perfect discrimination) equal to AUC (area under the ROC curve)

#### 3.1 CR decomposition of Brier's score

Due to the large number of unique probability profiles (377 among N=384 matches), forecasts were binned in three different ways: i) Fixed threshold intervals: D=10 from 0.0 to 1.0 for Home Win; D=5 with bounds at 0.10,0.15,0.20,0.25 and 0.35 for Draw and D=8 for Away Win with the first 7 bins equally spaced from 0 to 0.7 and the last one from 0.7; ii) Quantile thresholds intervals: deciles for Home Win and Away Win and quintiles for Draws iii) Bins automatically determined by the pool-adjacent-violators (PAV) algorithm used to set up the nonparametric isotonic regression deployed to estimate the conditional  $q_p = \Pr(X = 1 | P = p)$  outcome probabilities by minimizing the regression MSE with respect to D:

$$\sum_{d=1}^D \sum_{i=1}^N I(p_i \in [b_d, b_{d+1}]) (q_d - p_i)^2, \quad (8)$$

under the constraints of isotonicity ( $q_d$  estimation is a non-decreasing function of the original  $p_i$ 's). Results are displayed in Table 1, for Home Win, Draw and Away Win categories considered separately in terms of absolute values of Mean Brier score and its components (REL, RES, ACC) and Skill. Lack of reliability turns out to be small (lower than 5.5% of UNC) with miscalibration estimated a little bit higher under iso-regression. These values are supported by statistics and P values of Brier's score tests of departure from zero miscalibration (Spiegelhalter, 1986; Sellier-Moiseiwitch and David, 1993).




Figure 1: Reliability Diagrams for Home Win Probability Forecasts with plots of the Conditional Probability Events (CEP) against the Forecast Probability Values via Iso Regression. On the left : Reliability with Point 95% Consistency Bands. On the right: Discrimination diagrams with marginal distributions of forecasts and calibrated values.

Skill values are also appreciable for Home win (25 to 30%) and Away win (20 to 30%) with a little advantage of Odds vs Poisson of around 5. On the opposite, skill remains quite poor for Draw: 1.4 to 3.0% for Poisson and Odds respectively. Corresponding reliability diagrams were produced with an example shown for Home Win under Poisson and forecasts on Fig 1. There is some evidence of over-forecasting for Home Win practically all over the forecast range for Poisson, and at least, below  $p < 0.60$  for Odds (not shown) resulting in under-forecasted Draws. These conclusions are confirmed by the results (table 2) of the logistic regression with intercept and slope calibration coefficients showing clearly over-forecasting of Home Wins ( $\hat{\alpha} = -0.26$ ,  $\hat{\beta} = 1.11$ ) with Pvalues equal to 0.03 and 0.38 respectively while Away Wins are very well calibrated ( $\hat{\alpha} = 0.076$ ,  $\hat{\beta} = 1.053$ ).

### 3.2 LB and Yates' decompositions of Brier's score

Likelihood-base as well as Yates' decomposition of Brier Score rely largely on the concept of discrimination between the conditional distributions of forecasts with positive outcomes vs forecast with negative

outcomes. Characteristics of these two distributions are given in Table 3 and Fig 2. Differences between the means of these two conditional distributions are much more marked for Home Win and Away Win than for Draw. Again, these differences are more pronounced with Odds than Poisson as also reported by the Harrell c-statistics around 0.8 for Home Win and Away Win, and only 0.6 for Draw. Graphically, the boxplots confirmed this situation showing a clear separation of the two distributions for Home Win and Away Win and a tiny one for Draw.

Detailed accounts of Yates' and LB decompositions are shown on Table 4. In short, what emerges from them lies in the large role and weight given to the covariance component: 41 to 48% for Home Win and Away Win under Poisson and 50 to 55% with Odds with nevertheless a non-negligible part devoted to "noise" variance of forecasts (15 to 18%). The same picture applies to Draw but with much more tiny components, especially discrimination and covariance.

Table 4: Yates's and LB decompositions of Brier's score pertaining to Home Win, Draw and Away Win under the Poisson regression model (POI)

Factors	Home Win		Draw		Away Win		All	
	value	%	value	%	value	%	value	%
UNC	0.2458	100.0	0.1875	100.0	0.2158	100.0	0.6490	100.0
(-2)COV	-0.1190	-48.4	-0.0076	-4.0	-0.0886	-41.1	-0.2150	-29.5
VPB	0.0144	5.8	0.0001	0.0	0.0091	4.2	0.0236	3.6
VPW	0.0413	16.8	0.0031	1.7	0.0336	15.6	0.0780	12.0
RIL	0.0024	1.0	0.0017	0.9	0.0000	0.0	0.0042	0.6
REF	0.0556	22.6	0.0032	1.7	0.0427	19.8	0.1016	15.6
-DIS	-0.0144	-5.8	-0.0001	-0.0	-0.0091	-4.2	-0.0236	-3.6
CB2	0.1436	58.4	0.1817	96.9	0.1363	63.2	0.5396	83.1
BRS	0.1849	75.2	0.1848	98.6	0.1700	78.8	0.5397	83.1

Yates's decomposition into 5 components as follows  $BRS=UNC-2COV+VPB+VPW+RIL$  with UNC: Uncertainty, COV: Covariance between forecast and outcome, VPB: Variance among means of probability forecasts with outcome=1 and outcome=0, VPW : Average of Within groups variance and RIL marginal bias squared between the two groups, according to forecasting procedures (POI and ODD models) . Likelihood base decomposition into 3 components  $BRS=REF-DIS+CB2$  with REF: Refinement or Sharpness of forecast variance, DIS=Discrimination same as VPB and CB2: Type 2 bias equal to  $VPW-2COV+RIL$ .

## 4 Discussion

This presentation was deliberately restricted to the most popular (strictly) proper scoring rules as this properness property is a cornerstone of decision theory based on minimizing expected loss (or maximizing utility) (Bernardo, 1979; Gneiting and Raftery, 2007). They provide an incentive for ex ante honesty and reward ex post accuracy. Little was said about verifying probability forecasts for multiple categories taken simultaneously. Probability scoring rules are extended easily to that situation as shown in Table 4 for Yates' decomposition. Unhappily, such an extension is not straightforward for the CR decomposition. One reason for that lies on how to define bins. Procedures have been proposed to that respect by Broecker (2012) based on some functions of the probability vector for the J multiple categories of interest. A simple way to handle

the multiclass setting is by treating the problem as  $J$  one-versus-all binary events via e.g., a logistic-type regression with standard normalization of outcome probabilities. More generally, such forecasting verification methods already gained much attention in other fields especially in machine learning especially due to miscalibration of neural networks and its applications to health and medicine (Guo et al, 2017).




Figure 2: Box-plots of the conditional distributions of probabilistic forecasts of binary events (Home Win, Draw and Away Win) given the observed outcomes ( $X=0$ ) and ( $X=1$ ) and according to the Poisson loglinear regression (poi)

Although much of the theory and applications originated from meteorological literature (Winkler and Murphy, 1979; Jolliffe and Stephenson, 2003; Wilks, 2011), there had been a few attempts to apply some of these analytical procedures to football match results, especially in the EPL (Forrest and Simmons, 2000; Selliti Rangel, 2018; Wheatcroft, 2019; Read et al, 2021) but not enough. This area would benefit from a more systematic utilization. Regarding our application to the UEFA Champions League, it turns out that forecasts of Away Wins is both well calibrated and refined with good discrimination properties. The same trend can be seen on Home Wins, but with a trade-off between that over-forecasted category and under-forecasted draws with little discrimination of draw forecasts. It looks as if this category stands apart from the two others.

## References

- Bradley, A. A., T. Hashino, and S. S. Schwartz. (2003), "Distributions oriented verification of probability forecasts for small data samples", *Weather Forecasting*, 18, 903–917.
- Brier, G. (1950), "Verification of forecasts expressed in terms of probability", *Monthly Weather Review*, 78, 1-3.

- Bröcker, J. (2012) Probability forecasts. In *Forecast Verification: A Practitioner's Guide in Atmospheric Science*, Second Edition. Edited by I. T. Jolliffe, and D.B. Stephenson. 2012 John Wiley & Sons, pp 119-139.
- Cox, D.R. (1958), "Two Further Applications of a Model for Binary Regression", *Biometrika*, 45, 562-565.
- Dawid, A. P. (1986), "Probability Forecasting", *Encyclopedia of Statistical Science*, 7, 210-218.
- Dimitriadis, T. Gneiting, T. and Jordan, A.I. (2021) *Stable reliability diagrams for probabilistic classifiers*. Proceedings of the National Academy of Sciences of the United States of America **118** (8) DOI: [10.1073/pnas.2016191118](https://doi.org/10.1073/pnas.2016191118)
- Forrest, D., and R. Simmons. (2000), "Forecasting Sport: The Behaviour and Performance of Football Tipsters", *International Journal of Forecasting*, 16, 317-331.
- Gneiting, T., and A.E. Raftery. (2007), "Strictly Proper Scoring Rules, Prediction, and Estimation", *Journal of the American Statistical Association*, 102, 359-378.
- Guo, C., Pleiss, G., and K.Q., Weinberger.(2017), "On Calibration of Modern Neural Networks", arXiv: 1706.04599v2
- Gweon, H., and H. Yu. (2019), "How reliable is your reliability diagram?" *Pattern Recognition Letters*, 125, 687-693.
- Jolliffe, I., and D. Stephenson. (2003), "Forecast Verification: A Practitioner's Guide in Atmospheric Science", John Wiley-Blackwell
- Murphy, A.H. (1973), "A new vector partition of the probability score", *Journal of Applied Meteorology*, 12, 595-600.
- Murphy, A. H., and R.L., Winkler. (1987), "A general framework for forecast verification". *Monthly Weather Review* ,155, 1330-1338.
- Reade, J. J., Singleton C., and A. Brown. A. (2020) Evaluating Strange Forecasts: The Curious Case of Football Match Scorelines. *Scottish Journal of Political Economy*, 68, 261-285.
- Scarf, P., and J. Selliti Rangel Jr. (2016), "Models for outcomes of soccer matches". In *Handbook of Statistical Methods and Analyses in Sports*, Eds by J. Albert, M.E., Glickman, T. B, Swartz ,and R.H. Koning, 341-354. CRC Press, Chapman pp. 341-354.
- Seillier-Moiseiwitsch, F., and Dawid, A.P. (1993), "On Testing the Validity of Sequential Probability Forecasts", *Journal of the American Statistical Association*, 88, 55-359.
- Selliti Rangel Jr, J. (2018), "Estimation and Forecasting Team Strength Dynamics in Football: Investigation into Structural Breaks", PhD thesis, University of Salford, UK.
- Siegert, S. (2017), "Simplifying and generalising Murphy's Brier score decomposition." *Quarterly Journal of the Royal Meteorological Society*, 143, 1178-1183.
- Spiegelhalter, D. J. (1986), "Probabilistic prediction in patient management and clinical trials", *Statistics in Medicine* **5**, 421-433.
- Wheatcroft, E. (2019), "Interpreting the skill score form of forecast performance metrics", *International Journal of Forecasting*, 35, 573-579.
- Wilks, D. S. (2011), "Statistical Methods in the Atmospheric Sciences", 3<sup>rd</sup> Edition Academic Press, Oxford.
- Winkler, R.L., and A.H. Murphy. (1979), "The Use of Probabilities in Forecasts of Maximum and Minimum Temperatures", *The Meteorological Magazine*, 108, 317-329.
- Yates, J.F. (1982), "External correspondence: Decompositions of the mean probability score", *Organizational Behavior and Human Performance*, 30, 132-156.