

# Propagation phenomena in periodic patchy landscapes with interface conditions

François Hamel\*, Frithjof Lutscher† and Mingmin Zhang‡

*In memory of Pavol Brunovský, with admiration for a great mathematician*

## Abstract

This paper is concerned with a model for the dynamics of a single species in a one-dimensional heterogeneous environment. The environment consists of two kinds of patches, which are periodically alternately arranged along the spatial axis. We first establish the well-posedness for the Cauchy problem. Next, we give existence and uniqueness results for the positive steady state and we analyze the long-time behavior of the solutions to the evolution problem. Afterwards, based on dynamical systems methods, we investigate the spreading properties and the existence of pulsating traveling waves in the positive and negative directions. It is shown that the asymptotic spreading speed,  $c^*$ , exists and coincides with the minimal wave speed of pulsating traveling waves in positive and negative directions. In particular, we give a variational formula for  $c^*$  by using the principal eigenvalues of certain linear periodic eigenvalue problems.

*Mathematics Subject Classification:* 35K57; 92D25; 35J60.

*Keywords:* Patchy landscapes; Interface conditions; Periodic media; Asymptotic spreading speeds; Pulsating traveling waves.

## 1 Introduction

Reaction-diffusion equations of the type

$$u_t = \Delta u + f(u), \quad t > 0, \quad x \in \mathbb{R}^N, \quad (1.1)$$

have been introduced in the pioneering works of Fisher [18] and Kolmogorov, Petrovsky and Piskunov [28]. They are motivated by population genetics and aim at throwing light on the spatial spread of advantageous genetic features. The nonlinear reaction term considered there is that of logistic growth. Archetypes of such nonlinearities are  $f(u) = u(1 - u)$  or extensions like  $f(u) = u(1 - u^2)$ , which are sometimes also called monostable since they have exactly one stable nonnegative steady state. Skellam [47] then investigated this type of models in order to study spatial propagation of species and proposed quantitative explanations for the spread of muskrats throughout Europe at the beginning of the 20th century.

Since these celebrated works, a vast mathematical literature has been devoted to the homogeneous equation (1.1). It is of particular interest to investigate the structure of traveling front

---

\*Aix-Marseille Univ, CNRS, Centrale Marseille, I2M, Marseille, France ([francois.hamel@univ-amu.fr](mailto:francois.hamel@univ-amu.fr)).

†University of Ottawa, Department of Mathematics and Statistics & Department of Biology, Ottawa, Canada ([flutsche@uottawa.ca](mailto:flutsche@uottawa.ca)).

‡Aix-Marseille Univ, CNRS, Centrale Marseille, I2M, Marseille, France, and School of Mathematical Sciences, University of Science and Technology of China, Hefei, Anhui 230026, China ([mingmin.zhang.math@gmail.com](mailto:mingmin.zhang.math@gmail.com)).

solutions and their stability, propagation or invasion and spreading properties. The former is related to the well-known result that this equation has a family of planar traveling fronts of the form  $u(t, x) = U(x \cdot e - ct)$  for any given vector of unit norm  $e \in \mathbb{S}^{N-1}$ , which is the direction of propagation. Here,  $c > 0$  is the constant speed of the front and  $U : \mathbb{R} \rightarrow (0, 1)$  is the wave profile. It was proved in [28] that, under KPP assumptions,  $f(0) = f(1) = 0$  and  $0 < f(s) \leq f'(0)s$  in  $s \in (0, 1)$ , there exists a threshold speed  $c^* = 2\sqrt{f'(0)} > 0$  such that no fronts exist for  $c < c^*$ , while for each  $c \geq c^*$ , there is a unique (up to shift in space or time variables) planar front of the type  $U(x \cdot e - ct)$ . Such fronts are stable with respect to some natural classes of perturbations; see, for instance, [10, 22, 28, 29, 42, 48]. Many papers were also dedicated to such planar fronts for other types of nonlinearities  $f(\cdot)$ , for example the bistable and combustion type; see, e.g., [3, 9, 16, 17, 27].

While traveling waves are interesting mathematical objects, they do not necessarily represent biologically realistic scenarios. For questions of biological invasions, it is more realistic to study how a locally introduced population would spread. Corresponding mathematical invasion and spreading properties for model (1.1) were established by Aronson and Weinberger [4]. Under the same assumptions on  $f$  as in the previous paragraph, they proved that, starting with a nonnegative, compactly supported, continuous function  $u_0 \geq 0$ ,  $u_0 \not\equiv 0$ , the solution  $u(t, x)$  of (1.1) spreads with speed  $c^*$  in all directions for large times. More precisely,  $\max_{|x| \leq ct} |u(t, x) - 1| \rightarrow 0$  as  $t \rightarrow +\infty$  for each  $c \in [0, c^*)$ , and  $\max_{|x| \geq ct} u(t, x) \rightarrow 0$  for each  $c > c^*$ .

Most landscapes are not homogeneous, as model (1.1) implicitly assumes. Several possible generalizations of the equation to heterogeneous landscapes exist, for example

$$u_t = \nabla \cdot (D(x) \nabla u) u + f(x, u), \quad t > 0, \quad x \in \mathbb{R}^N, \quad (1.2)$$

in periodic media (by periodic, we mean that  $D(\cdot + k) = D$  and  $f(\cdot + k, s) = f(\cdot, s)$  for all  $k \in L_1 \mathbb{Z} \times \dots \times L_N \mathbb{Z}$  and  $s \in [0, 1]$ , where  $L_1, \dots, L_N$  are given positive real numbers). For such equations, standard traveling fronts do not exist in general. Instead, the notion of traveling fronts is replaced by the more general concept of pulsating fronts [46]. If a (unique) periodic positive steady state  $p(x)$  of (1.2) exists, a pulsating traveling front connecting 0 and  $p(x)$  is a solution of the type  $u(t, x) = U(x \cdot e - ct, x)$  with  $c \neq 0$  and  $e$  a unit vector representing the direction of propagation, if the function  $U : \mathbb{R} \times \mathbb{R}^N \rightarrow \mathbb{R}$  satisfies

$$\begin{cases} U(-\infty, x) = p(x), \quad U(+\infty, x) = 0 & \text{uniformly in } x \in \mathbb{R}^N, \\ U(s, \cdot) \text{ is periodic in } \mathbb{R}^N & \text{for all } s \in \mathbb{R}. \end{cases}$$

Moreover, for every  $x \in \mathbb{R}^N$ , the function  $t \mapsto u(t, x + cte)$  is in general quasi-periodic.

Berestycki, Hamel and Roques [7] gave a complete and rigorous mathematical analysis of the periodic heterogeneous model (1.2) in any space dimension. They required the coefficient functions  $D$  and  $f$  to be sufficiently smooth, yet some of their results are valid under reduced regularity assumptions. Solutions in [7] are still at least of class  $C^{1,\rho}$  with respect to  $x$  for all  $\rho \in (0, 1)$ , while the solutions considered in the present paper are even not continuous at all points in general, or not of class  $C^1$  even when they are continuous after a rescaling. In [7], existence, uniqueness and stability results were established. A criterion for species persistence and the effects of fragmentation on species survival were studied. Furthermore, the same authors studied the question of biological invasion and existence of pulsating fronts and they proved a variational formula of the minimal speed of such pulsating fronts and then analyzed the dependency of this speed on the heterogeneity of the medium [8]. We refer the reader to [5, 6, 20, 23, 26, 30, 31, 49] for more results on the existence, uniqueness and qualitative results of pulsating traveling fronts. For some results in space-time periodic media, see, e.g., [37–39]. We also refer to [5, 6, 13–15, 19, 25, 44, 50–52] for the existence and qualitative results with other types of nonlinearities or various boundary conditions in periodic domains.

While equation (1.2) and the corresponding theory is mathematically elegant, it is very difficult if not impossible to apply the model to biological invasions since the data requirements of finding diffusion coefficients and growth rates for continuously changing landscape characteristics are too costly. Alternatively, landscape ecology views natural environments as patches of homogeneous habitats such as forests, grasslands, or marshes, possibly fragmented by natural or artificial barriers like rivers or roads. Each patch is relatively homogeneous within but significantly different from adjacent patches. Shigesada, Kawasaki and Teramoto [46] used this perspective of a patchy landscape and proposed a heterogeneous extension of (1.1) with piecewise constant coefficient functions; see also [45]. For simplicity, they considered only two types of patches, arranged in a periodically alternating way. More specifically, they introduced the following equation

$$u_t = (D(x)u_x)_x + u(\mu(x) - u), \quad t > 0, \quad x \in \mathbb{R} \setminus S, \quad (1.3)$$

where  $\mu(x)$  is interpreted as the intrinsic growth rate of the population,  $D(x)$  is the diffusion coefficient, and  $S$  is the set of all interfaces between all adjacent patches. These functions are piecewise constant, namely, for  $m \in \mathbb{Z}$ ,

$$\begin{cases} D(x) = d_1(> 0), & \mu(x) = \mu_1, & ml - l_1 < x < ml & \text{(in patches of type 1),} \\ D(x) = d_2(> 0), & \mu(x) = \mu_2, & ml < x < ml + l_2 & \text{(in patches of type 2),} \end{cases}$$

with  $l_1, l_2 > 0$  and  $l = l_1 + l_2$ . Without loss of generality, one can assume that  $\mu_1 \geq \mu_2$ . Furthermore, if one assumes that the medium is not unfavorable everywhere, then  $\mu_1 > 0$ , i.e., type-1 patches support population growth and are “source” patches in biological terms. However,  $\mu_2$  can be negative, so that patches of type 2 do not support population growth and are “sink” patches in biological terms.

The above model is not complete. At the boundaries or interfaces between two patches, matching conditions must be imposed. Shigesada, Kawasaki and Teramoto [46] required continuity of density and flux, i.e., for  $t > 0$ :

$$u(t, x^-) = u(t, x^+), \quad D(x^-)u_x(t, x^-) = D(x^+)u_x(t, x^+),$$

for all  $x = mL$  and  $x = mL + L_2$  ( $m = 0, \pm 1, \pm 2, \dots$ ). Superscripts  $\pm$  denote one-sided limits from the right and the left, respectively. When we take  $D(x) \equiv d$ ,  $\mu(x) \equiv \mu > 0$ , problem (1.3) is reduced to the Fisher-KPP equation with threshold speed  $2\sqrt{d\mu}$ . We shall refer to this kind of model, i.e., the differential equations combined with interface matching conditions as *patch model* or *patch problem*.

Shigesada and coauthors obtained the invasion conditions in terms of the sizes of patches, diffusivities and growth rates. They proved that the population spreads successfully if and only if the invasion condition is satisfied. Moreover, when invasion occurs, the distribution of the population initially localized in a bounded area always evolves into a pulsating front. The velocity was calculated with the aid of the dispersion relation, based on linearization at low density [46]. When diffusion is constant, the rigorous analytical results in [7] apply to this patch model, but the case of discontinuous diffusion and interface matching conditions is not covered.

Recently, Maciel and Lutscher [34] introduced novel interface matching conditions, based on the work by Ovaskainen and Cornell [40]. The population flux is still continuous at an interface, but the density is not. We will explain these conditions in detail below. These matching conditions not only allow us to include patch preference data, which are frequently collected in the field, into reaction-diffusion models, but also remove some biologically unrealistic behavior that the model with the continuous-density interface conditions above shows (see [34] for a thorough discussion). A number of recent studies use this new framework to study questions of persistence

and spread [2, 35] and apply it to marine reserve design [1]. All these studies show that the correct choice of interface conditions has a crucial effect on basic quantities such as population persistence conditions and spread rates in periodic environments. Recently, Maciel and Lutscher [36] showed how different movement strategies for competing species in patchy landscapes can lead to different outcomes of competition. Maciel and coauthors found evolutionarily steady movement strategies in a two-patch landscape [33].

Our paper is devoted to a rigorous analytical study of the periodic patch model with two types of alternately arranged patches in a one-dimensional habitat. The population may grow or decay, depending on patch type, and diffusivity may change between patches. The setting and assumptions will be made precise below. The aim of the present work is first to rigorously prove the well-posedness of this somewhat nonstandard patch model starting with nonnegative continuous and bounded initial data. Then we investigate the long-time behavior and spatial dynamics of this type of model in the framework of a periodic environment with monostable dynamics. We give a criterion for the existence and uniqueness of a positive and bounded steady state. Furthermore, under the hypothesis that the species can persist, we shall prove the existence of an asymptotic spreading speed  $c^*$  of the solution to the Cauchy problem and we show that this spreading speed coincides with the minimal speed for rightward and leftward pulsating fronts. Moreover, the asymptotic spreading speed  $c^*$  can be characterized using a family of eigenvalues. To the best of our knowledge, the results that had previously been discussed only formally or observed numerically in [34], are proved rigorously here for the first time.

## 2 Model presentation and statement of the main results

### 2.1 The model and some equivalent formulations

Our model is a joint generalization of the models in [46] and [34]. We consider a patchy periodically alternating landscape consisting of two types of patches (say, type 1 and 2); see Figure 1. Each patch is homogeneous within. We denote the length of patch type  $i$  ( $i = 1, 2$ ) by  $l_i$  so that the period is  $l = l_1 + l_2$ . Accordingly, the real line is divided into intervals of the form

$$I_n = [nl - l_1, nl + l_2], \quad n \in \mathbb{Z},$$

each consisting of two adjacent patches. Such intervals were called “tiles” in [12].




Figure 1: Schematic figure of the one-dimensional periodic patchy environment.

For  $n \in \mathbb{Z}$ , let  $I_{1n} = (nl - l_1, nl)$  be the patches of type 1 and  $I_{2n} = (nl, nl + l_2)$  be the patches of type 2. On each patch  $I_{in}$ , we denote by  $v_{in} = v|_{I_{in}}$  the density of the population, by  $d_i$  the constant diffusion coefficients, and by  $f_i$  the corresponding reaction nonlinearities. Our model then reads, for  $n \in \mathbb{Z}$ ,

$$\begin{cases} \frac{\partial v_{1n}}{\partial t} = d_1 \frac{\partial^2 v_{1n}}{\partial x^2} + f_1(v_{1n}), & t > 0, x \in (nl - l_1, nl), \\ \frac{\partial v_{2n}}{\partial t} = d_2 \frac{\partial^2 v_{2n}}{\partial x^2} + f_2(v_{2n}), & t > 0, x \in (nl, nl + l_2). \end{cases} \quad (2.1)$$

In (2.1), the equations for  $v_{in} = v|_{I_{in}}$  are set in the open intervals  $(nl - l_1, nl)$  and  $(nl, nl + l_2)$ , but it will eventually turn out that the constructed solutions are such that the functions  $v_{in}$

can be extended in  $(0, +\infty) \times [nl - l_1, nl]$  or  $(0, +\infty) \times [nl, nl + l_2]$  as  $C_{t;x}^{1;2}$  functions, so that equations (2.1) will be satisfied in the closed intervals  $[nl - l_1, nl]$  and  $[nl, nl + l_2]$ . The matching conditions for the population density and flux at the interfaces are given by

$$\begin{cases} v_{1n}(t, x^-) = kv_{2n}(t, x^+), & d_1(v_{1n})_x(t, x^-) = d_2(v_{2n})_x(t, x^+), & t > 0, x = nl, \\ kv_{2n}(t, x^-) = v_{1(n+1)}(t, x^+), & d_2(v_{2n})_x(t, x^-) = d_1(v_{1(n+1)})_x(t, x^+), & t > 0, x = nl + l_2, \end{cases} \quad (2.2)$$

with parameter

$$k = \frac{\alpha}{1 - \alpha} \times \frac{d_2}{d_1}. \quad (2.3)$$

Here,  $\alpha \in (0, 1)$  denotes the probability that an individual at the interface chooses to move to the adjacent patch of type 1, and  $1 - \alpha$  the probability that it moves to the patch of type 2. Individuals cannot stay at the interfaces. These interface conditions were derived in [40] and studied in more detail in [34]. They reflect the movement behavior of individuals when they come to the edge of a patch. With these interface conditions, the population density is discontinuous across a patch interface in the presence of patch preference and/or when the diffusion rates in these two kinds of patches are different. Throughout the paper, we assume that the reaction terms  $f_i$  ( $i = 1, 2$ ) have the properties:

$$f_i \in C^1(\mathbb{R}), \quad f_i(0) = 0, \quad \text{and there is } K_i > 0 \text{ such that } f_i \leq 0 \text{ in } [K_i, +\infty). \quad (2.4)$$

Without loss of generality, we will consider type-1 patches as more favorable than type-2 patches, that is,  $f'_1(0) \geq f'_2(0)$ . In some statements, we will also assume that type-1 patches are “source” patches, i.e., patches where the intrinsic growth rate of the population is positive ( $f'_1(0) > 0$ ), while type-2 patches may be source patches ( $f'_2(0) > 0$ ), or “sink” patches ( $f'_2(0) < 0$ ), or such that  $f'_2(0) = 0$ . In order to investigate the long-time behavior and spatial dynamics, we will further assume in some statements that the functions  $f_i$  satisfy the strong Fisher-KPP assumption:

$$\begin{cases} \text{the functions } s \mapsto \frac{f_i(s)}{s} \text{ are non-increasing in } s > 0 \text{ for } i = 1, 2, \\ \text{and decreasing in } s > 0 \text{ for at least one } i. \end{cases} \quad (2.5)$$

For instance,  $f_i$  satisfying hypotheses (2.4)–(2.5) can be functions of the type  $f_i(s) = s(\mu_i - s)$ .

Since the discontinuity in the densities at the interfaces makes the problem quite delicate to study, we rescale the densities in such a way that the matching conditions become continuous in the density. More precisely, we set  $u_{1n}(t, x) = v_{1n}(t, x)$  for  $t \geq 0$ ,  $x \in (nl - l_1, nl)$  and  $n \in \mathbb{Z}$ , and  $u_{2n}(t, x) = kv_{2n}(t, x)$  for  $t \geq 0$ ,  $x \in (nl, nl + l_2)$  and  $n \in \mathbb{Z}$ . Then  $u_{1n}$  satisfy the same equations as  $v_{1n}$  with  $\tilde{f}_1(s) = f_1(s)$ , while  $u_{2n}$  satisfy the equations of  $v_{2n}$  with  $f_2$  replaced by  $\tilde{f}_2(s) = kf_2(s/k)$ . We notice that  $\tilde{f}_i$  ( $i = 1, 2$ ) satisfy the same hypotheses as  $f_i$  with  $K_i$  replaced by  $\tilde{K}_i$  where  $\tilde{K}_1 = K_1$  and  $\tilde{K}_2 = kK_2$ . Thanks to the change of variables, the interface conditions for the densities are now continuous; however, the flux interface conditions become discontinuous, namely,

$$\begin{cases} u_{1n}(t, x^-) = u_{2n}(t, x^+), & d_1(u_{1n})_x(t, x^-) = \frac{d_2}{k}(u_{2n})_x(t, x^+), & t > 0, x = nl, \\ u_{2n}(t, x^-) = u_{1(n+1)}(t, x^+), & \frac{d_2}{k}(u_{2n})_x(t, x^-) = d_1(u_{1(n+1)})_x(t, x^+), & t > 0, x = nl + l_2. \end{cases}$$

We drop the tilde from hereon. Notice that the properties (2.4) and (2.5) are invariant under this change. Putting it all together, we are led to the following problem:

$$\begin{cases} \frac{\partial u_{1n}}{\partial t} = d_1 \frac{\partial^2 u_{1n}}{\partial x^2} + f_1(u_{1n}), & t > 0, x \in (nl - l_1, nl), \\ \frac{\partial u_{2n}}{\partial t} = d_2 \frac{\partial^2 u_{2n}}{\partial x^2} + f_2(u_{2n}), & t > 0, x \in (nl, nl + l_2), \end{cases} \quad (2.6)$$

with continuous density conditions and discontinuous flux interface conditions,

$$\begin{cases} u_{1n}(t, x^-) = u_{2n}(t, x^+), & (u_{1n})_x(t, x^-) = \sigma(u_{2n})_x(t, x^+), & t > 0, x = nl, \\ u_{2n}(t, x^-) = u_{1(n+1)}(t, x^+), & \sigma(u_{2n})_x(t, x^-) = (u_{1(n+1)})_x(t, x^+), & t > 0, x = nl + l_2, \end{cases} \quad (2.7)$$

in which, from (2.3), we have

$$\sigma = \frac{d_2}{kd_1} = \frac{1-\alpha}{\alpha} > 0. \quad (2.8)$$

When  $\sigma = 1$  (that is,  $\alpha = 1/2$ ), the model (2.6)–(2.7) is reduced to the one in [46].

From now on, we denote by

$$S_1 = l\mathbb{Z}$$

the interface points between  $(nl - l_1, nl)$  and  $(nl, nl + l_2)$ , and by

$$S_2 = \{s + l_2 : s \in l\mathbb{Z}\}$$

the interface points between  $(nl, nl + l_2)$  and  $(nl + l_2, (n+1)l)$ . Therefore,  $S = S_1 \cup S_2$  represents all the interface points in  $\mathbb{R}$ . For convenience of our analysis, by setting  $u(t, x) = u_{1n}(t, x)$  for  $t > 0$  and  $x \in (nl - l_1, nl)$ ,  $u(t, x) = u_{2n}(t, x)$  for  $t > 0$  and  $x \in (nl, nl + l_2)$ ,  $u(t, x) = u_{1n}(t, x^-) = u_{2n}(t, x^+)$  for  $t > 0$  and  $x = nl$ , and  $u(t, x) = u_{2n}(t, x^-) = u_{1(n+1)}(t, x^+)$  for  $t > 0$  and  $x = nl + l_2$ , we rewrite the above model (2.6)–(2.7) in the following form:

$$\begin{cases} u_t - d(x)u_{xx} = f(x, u), & t > 0, x \in \mathbb{R} \setminus S, \\ u(t, x^-) = u(t, x^+), \quad u_x(t, x^-) = \sigma u_x(t, x^+), & t > 0, x \in S_1, \\ u(t, x^-) = u(t, x^+), \quad \sigma u_x(t, x^-) = u_x(t, x^+), & t > 0, x \in S_2, \end{cases} \quad (2.9)$$

where the diffusivity  $d$  and nonlinearity  $f$  are given by

$$d(x) = \begin{cases} d_1, & x \in (nl - l_1, nl), \\ d_2, & x \in (nl, nl + l_2), \end{cases} \quad f(x, s) = \begin{cases} f_1(s), & x \in (nl - l_1, nl), \\ f_2(s), & x \in (nl, nl + l_2), \end{cases} \quad (2.10)$$

and the parameter  $\sigma > 0$  is defined as in (2.8). Conditions (2.4) and (2.5) on  $f_i$  are equivalent to the following ones:

$$\begin{cases} \forall x \in \mathbb{R} \setminus S, f(x, \cdot) \in C^1(\mathbb{R}), f(x, 0) = 0, \\ \exists M = \max(K_1, K_2) > 0, \forall x \in \mathbb{R} \setminus S, \forall s \geq M, f(x, s) \leq 0, \\ \forall x \in (nl - l_1, nl), f(x, \cdot) = f_1, \quad \forall x \in (nl, nl + l_2), f(x, \cdot) = f_2. \end{cases} \quad (2.11)$$

and

$$\begin{cases} \text{the functions } s \mapsto \frac{f(x, s)}{s} \text{ are non-increasing in } s > 0 \text{ in all patches,} \\ \text{and decreasing in } s > 0 \text{ in at least one type of patch.} \end{cases} \quad (2.12)$$

From now on, we always assume that (2.11) is satisfied. Throughout this paper, unless otherwise specified, we always write  $I$  for an arbitrary patch in  $\mathbb{R}$  of either type, i.e., either  $I = (nl - l_1, nl)$  or  $I = (nl, nl + l_2)$ .



## 2.2 Well-posedness of the Cauchy problem (2.9)–(2.10)

Since the patch model considered in this paper is not standard, we shall first establish the well-posedness of the Cauchy problem (2.9)–(2.10) with hypotheses (2.11) on  $f$  and with nonnegative bounded and continuous initial conditions  $u_0 : \mathbb{R} \rightarrow \mathbb{R}$ . Before proceeding with the analysis, we present here the definition of a classical solution to (2.9)–(2.10).

**Definition 2.1.** For  $T \in (0, +\infty]$ , we say that a continuous function  $u : [0, T) \times \mathbb{R} \rightarrow \mathbb{R}$  is a classical solution of the Cauchy problem (2.9)–(2.10) in  $[0, T) \times \mathbb{R}$  with an initial condition  $u_0$ , if  $u(0, \cdot) = u_0$ , if  $u|_{(0, T) \times \bar{I}} \in C_{t;x}^{1;2}((0, T) \times \bar{I})$  for each patch  $I = (nl - l_1, nl)$  or  $(nl, nl + l_2)$ , and if all identities in (2.9) are satisfied pointwise for  $0 < t < T$ .

**Theorem 2.2.** Under assumption (2.11), for any nonnegative bounded continuous initial condition  $u_0$ , there is a nonnegative bounded classical solution  $u$  in  $[0, +\infty) \times \mathbb{R}$  of the Cauchy problem (2.9)–(2.10) such that, for any  $\tau > 0$  and any patch  $I \subset \mathbb{R}$ ,

$$\|u\|_{[\tau, +\infty) \times \bar{I}} \|u\|_{C_{t;x}^{1;\gamma;2,\gamma}([\tau, +\infty) \times \bar{I})} \leq C,$$

with a positive constant  $C$  depending on  $\tau$ ,  $l_{1,2}$ ,  $d_{1,2}$ ,  $f_{1,2}$ ,  $\sigma$  and  $\|u_0\|_{L^\infty(\mathbb{R})}$ , and with a universal positive constant  $\gamma \in (0, 1)$ . Moreover,  $u(t, x) > 0$  for all  $(t, x) \in (0, +\infty) \times \mathbb{R}$  if  $u_0 \not\equiv 0$ , and  $u(t, x) = u(t, x + l)$  for all  $(t, x) \in [0, +\infty) \times \mathbb{R}$  if  $u_0(x) = u_0(x + l)$  for all  $x \in \mathbb{R}$ . Lastly, the solutions depend monotonically and continuously on the initial data, in the sense that if  $u_0 \leq v_0$  then the corresponding solutions satisfy  $u \leq v$  in  $[0, +\infty) \times \mathbb{R}$ , and for any  $T \in (0, +\infty)$  the map  $u_0 \mapsto u$  is continuous from  $C^+(\mathbb{R}) \cap L^\infty(\mathbb{R})$  to  $C([0, T] \times \mathbb{R}) \cap L^\infty([0, T] \times \mathbb{R})$  equipped with the sup norms, where  $C^+(\mathbb{R})$  denotes the set of nonnegative continuous functions in  $\mathbb{R}$ .

We remark that the existence and uniqueness of a global bounded periodic classical solution to such a patch model was considered in [33] for (2.1)–(2.2) with periodic and possibly discontinuous initial data. By contrast, our result is established for general continuous and bounded initial data. Moreover, we also discuss the continuous dependence of solutions on initial data and give a priori estimates, which will play a critical role in the monotone semiflow argument used in the sequel. The well-posedness proof here can also be adapted to other non-periodic patch problems.

## 2.3 Existence, uniqueness and attractiveness of a positive periodic steady state

To investigate the existence and uniqueness of a positive bounded steady state as well as the large-time behavior of solutions to the Cauchy problem, we first study the following eigenvalue problem. From [34, 46] (see also Lemma 6.2 below), there exists a principal eigenvalue  $\lambda_1$ , defined as the unique real number such that there exists a unique continuous function  $\phi : \mathbb{R} \rightarrow \mathbb{R}$  with  $\phi|_{\bar{I}} \in C^\infty(\bar{I})$  for each patch  $I$ , that satisfies

$$\begin{cases} \mathcal{L}_0 \phi := -d(x)\phi'' - f_s(x, 0)\phi = \lambda_1 \phi, & x \in \mathbb{R} \setminus S, \\ \phi(x^-) = \phi(x^+), \quad \phi'(x^-) = \sigma \phi'(x^+), & x \in S_1, \\ \phi(x^-) = \phi(x^+), \quad \sigma \phi'(x^-) = \phi'(x^+), & x \in S_2, \\ \phi(x) \text{ is periodic, } \phi > 0, \quad \|\phi\|_{L^\infty(\mathbb{R})} = 1. \end{cases} \quad (2.13)$$

By periodic, we mean that  $\phi(\cdot + l) = \phi$  in  $\mathbb{R}$ . In the sequel we say that 0 is an unstable steady state of (2.9) if  $\lambda_1 < 0$ , otherwise the state 0 is said to be stable (i.e.,  $\lambda_1 \geq 0$ ). These definitions will be seen to be natural in view of the results we prove here. By applying (2.13) at minimal

and maximal points of the positive continuous periodic function  $\phi$ , whether these points be in patches or on the interfaces, it easily follows that

$$-f'_1(0) \leq \lambda_1 \leq -f'_2(0)$$

(remember that  $f'_1(0) \geq f'_2(0)$  without loss of generality). In particular, if  $\lambda_1 < 0$ , then  $f'_1(0) > 0$ , that is,  $f_s(x, 0)$  is necessarily positive (at least) in the favorable patches.

We first state a criterion for the existence of a continuous solution  $p : \mathbb{R} \rightarrow \mathbb{R}$  (such that  $p|_{\bar{I}} \in C^2(\bar{I})$  for each patch  $I$ ) to the elliptic problem:

$$\begin{cases} -d(x)p''(x) - f(x, p(x)) = 0, & x \in \mathbb{R} \setminus S, \\ p(x^-) = p(x^+), \quad p'(x^-) = \sigma p'(x^+), & x \in S_1, \\ p(x^-) = p(x^+), \quad \sigma p'(x^-) = p'(x^+), & x \in S_2. \end{cases} \quad (2.14)$$

**Theorem 2.3.** (i) Assume that 0 is an unstable solution of (2.14) (i.e.,  $\lambda_1 < 0$ ) and that  $f$  satisfies (2.11). Then there exists a bounded positive and periodic solution  $p$  of (2.14).

(ii) Assume that 0 is a stable solution of (2.14) (i.e.,  $\lambda_1 \geq 0$ ) and that  $f$  satisfies (2.11)–(2.12). Then 0 is the only nonnegative bounded solution of (2.14).

For reaction-diffusion equations that describe population dynamics in general periodically fragmented landscapes but do not include movement behavior at interfaces, the criteria for existence (and uniqueness) of the stationary problem in arbitrary dimension can be found in [7]. It turns out that the approach there can be adapted to our periodic patch model with the additional nonstandard interface conditions.

Let us now provide an insight into the stability of the trivial solution of (2.14). Under certain reasonable hypotheses on the diffusivities, the sizes of favorable and unfavorable patches, as well as the nonlinearities, the principal eigenvalue  $\lambda_1$  of (2.13) can indeed be negative. For example, when all patches support population growth, namely  $f'_1(0) > 0$  and  $f'_2(0) > 0$ , then the zero state is unstable. When the landscape consists of source and sink patches, i.e., when  $f'_1(0) > 0 > f'_2(0)$ , the stability of the zero state depends on the relationships between patch size, patch preference, diffusivity and growth rates. In the case  $\sigma = 1$ , Shigesada and coworkers derived such a stability criterion [46]; the case for general  $\sigma > 0$  can be found in [34]. We here derive an even more general formula, when we only assume that  $f'_2(0) \leq f'_1(0)$ . To do so, we first observe that the continuous functions  $x \mapsto \phi(-l_1 - x)$  and  $x \mapsto \phi(l_2 - x)$  still solve (2.13) as  $\phi$  does, and by uniqueness we get that  $\phi(-l_1 - x) = \phi(l_2 - x) = \phi(x)$  for all  $x \in \mathbb{R}$ , hence  $\phi'(-l_1/2) = \phi'(l_2/2) = 0$ . Then, as in [34, 46], by solving (2.13) in  $[-l_1/2, 0]$  and in  $[0, l_2/2]$  with zero derivatives at  $-l_1/2$  and  $l_2/2$ , and by matching the interface conditions at 0, we find that  $\lambda_1$  is the smallest root in  $[-f'_1(0), -f'_2(0)]$  of the equation:

$$\sqrt{\frac{f'_1(0) + \lambda_1}{d_1}} \tan\left(\sqrt{\frac{f'_1(0) + \lambda_1}{d_1}} \times \frac{l_1}{2}\right) = \sigma \sqrt{-\frac{\lambda_1 + f'_2(0)}{d_2}} \tanh\left(\sqrt{-\frac{\lambda_1 + f'_2(0)}{d_2}} \times \frac{l_2}{2}\right). \quad (2.15)$$

When  $0 < f'_2(0) \leq f'_1(0)$  or when  $0 = f'_2(0) < f'_1(0)$  (irrespective of the other parameters), then the trivial solution of (2.14) is unstable (i.e.,  $\lambda_1 < 0$ ). When  $f'_2(0) \leq f'_1(0) \leq 0$ , then  $\lambda_1 \geq 0$ . When  $f'_2(0) < 0 < f'_1(0)$ , we then derive that the trivial solution of (2.14) is stable ( $\lambda_1 \geq 0$ ) if

$$l_1 \leq l_1^c := 2\sqrt{\frac{d_1}{f'_1(0)}} \tan^{-1}\left(\sigma \sqrt{\frac{-d_1 f'_2(0)}{d_2 f'_1(0)}} \tanh\left(\sqrt{\frac{-f'_2(0)}{d_2}} \times \frac{l_2}{2}\right)\right) \quad (2.16)$$



(notice that  $l_1^c > 0$ ), and unstable ( $\lambda_1 < 0$ ) if  $l_1 > l_1^c$ . The persistence threshold  $l_1^c$  is decreasing with  $f_1'(0) > 0$  and increasing with  $d_1$  and  $l_2$ . Passing to the limit  $l_2 \rightarrow +\infty$ , we find that

$$l_1^c \rightarrow L_1^c := 2\sqrt{\frac{d_1}{f_1'(0)}} \tan^{-1} \left( \sigma \sqrt{\frac{-d_1 f_2'(0)}{d_2 f_1'(0)}} \right).$$

Therefore, as long as  $l_1 > L_1^c$ , the trivial solution of (2.14) is unstable (i.e.,  $\lambda_1 < 0$ ), no matter how large the size of the unfavorable patches is. Similarly, there is a critical rate

$$(f_2'(0))^c = -\frac{d_2 f_1'(0)}{\sigma^2 d_1} \left[ \tan \left( \sqrt{\frac{f_1'(0)}{d_1}} \times \frac{l_1}{2} \right) \right]^2$$

such that, if  $0 > f_2'(0) > (f_2'(0))^c$ , then the trivial solution of (2.14) is unstable (i.e.,  $\lambda_1 < 0$ ), no matter how large the size of the unfavorable patch is.

It also follows from (2.15) that, provided  $f_2'(0) \neq f_1'(0)$ , the principal eigenvalue  $\lambda_1$  is increasing with respect to  $\sigma > 0$ , that is,  $\lambda_1$  is decreasing with respect to  $\alpha \in (0, 1)$ . When  $\alpha \in (0, 1)$  increases, then the individuals at the interfaces have more propensity to go to patches of type 1 rather than to patches of type 2. This means that the relative advantage of the more favorable patches becomes more prominent:  $\lambda_1$  decreases and the 0 solution has more chances to become unstable. It is also easy to see that  $\lambda_1 \rightarrow -f_1'(0)$  as  $\sigma \rightarrow 0^+$  (that is, as  $\alpha \rightarrow 1^-$ ), hence 0 is unstable if  $\alpha \approx 1$ , provided the patches of type 1 support population growth. On the other hand,  $\lambda_1 \rightarrow \min(d_1 \pi^2 / l_1^2 - f_1'(0), -f_2'(0))$  as  $\sigma \rightarrow +\infty$  (that is, as  $\alpha \rightarrow 0^+$ ). Therefore, if  $f_1'(0) \geq d_1 \pi^2 / l_1^2$ , and even if  $f_2'(0) < 0$ , then 0 is still unstable when  $\alpha$  is small (and actually whatever the value of  $\alpha \in (0, 1)$  and the other parameters may be).

Next, we state a Liouville type result for problem (2.14).

**Theorem 2.4.** *Assume that  $f$  satisfies (2.11)–(2.12) and that the zero solution of (2.14) is unstable (i.e.,  $\lambda_1 < 0$ ). Then there exists at most one positive and bounded solution  $p$  of (2.14). Furthermore, such a solution  $p$ , if any, is periodic and  $\inf_{\mathbb{R}} p = \min_{\mathbb{R}} p > 0$ .*

Under the assumptions of Theorem 2.3 (i) and Theorem 2.4, we now look at the global attractiveness of the unique positive and bounded stationary solution  $p$  of (2.14) for the solutions of the Cauchy problem (2.9)–(2.10).

**Theorem 2.5.** *Assume that  $f$  satisfies (2.11)–(2.12). Let  $u$  be the solution of the Cauchy problem (2.9)–(2.10) with a nonnegative bounded and continuous initial datum  $u_0 \neq 0$ .*

- (i) *If 0 is an unstable solution of (2.14) (i.e.,  $\lambda_1 < 0$ ), then  $u(t, \cdot)|_{\bar{I}} \rightarrow p|_{\bar{I}}$  in  $C^2(\bar{I})$  as  $t \rightarrow +\infty$  for each patch  $I$ , where  $p$  is the unique positive bounded and periodic solution of (2.14) given by Theorem 2.3 (i) and Theorem 2.4.<sup>1</sup>*
- (ii) *If 0 is a stable solution of (2.14) (i.e.,  $\lambda_1 \geq 0$ ), then  $u(t, \cdot) \rightarrow 0$  uniformly in  $\mathbb{R}$  as  $t \rightarrow +\infty$ .*

## 2.4 Spreading speeds and pulsating traveling waves

In this subsection, we assume that the zero solution of (2.14) is unstable (i.e.,  $\lambda_1 < 0$ ) and that  $f$  satisfies (2.11)–(2.12). Let  $p$  be the unique positive bounded and periodic solution of (2.14)

---

<sup>1</sup>This statement shows that the solution  $u$  converges as  $t \rightarrow +\infty$  locally uniformly in space to the space-periodic function  $p$ . For a convergence result to time-periodic solutions for time-periodic quasilinear parabolic equations, we refer to [11].

obtained from Theorem 2.3 (i) and Theorem 2.4. After showing in Theorem 2.5 (i) the attractiveness of  $p$ , we now want to describe the way the positive steady state  $p$  invades the whole domain.

Let  $\mathcal{C}$  be the space of all bounded and uniformly continuous functions from  $\mathbb{R}$  to  $\mathbb{R}$  equipped with the compact open topology, i.e., we say that  $u_n \rightarrow u$  as  $n \rightarrow +\infty$  in  $\mathcal{C}$  when  $u_n \rightarrow u$  locally uniformly in  $\mathbb{R}$ . For  $u, v \in \mathcal{C}$ , we write  $u \geq v$  when  $u(x) \geq v(x)$  for all  $x \in \mathbb{R}$ ,  $u > v$  when  $u \geq v$  and  $u \not\equiv v$ , and  $u \gg v$  when  $u(x) > v(x)$  for all  $x \in \mathbb{R}$ . Notice that  $p \in \mathcal{C}$  is periodic and satisfies  $p \gg 0$ . We define

$$\mathcal{C}_p = \{v \in \mathcal{C} : 0 \leq v \leq p\}. \quad (2.17)$$

Let  $\mathbb{P}$  be the set of all continuous and periodic functions from  $\mathbb{R}$  to  $\mathbb{R}$  equipped with the  $L^\infty$ -norm, and  $\mathbb{P}^+ = \{u \in \mathbb{P} : u \geq 0\}$ .

The first result of this section states the existence of a speed of invasion by the state  $p$ .

**Theorem 2.6.** *Assume that  $f$  satisfies (2.11)–(2.12) and that the zero solution of (2.14) is unstable (i.e.,  $\lambda_1 < 0$ ). Then there is an asymptotic spreading speed,  $c^* > 0$ , given explicitly by*

$$c^* = \inf_{\mu > 0} \frac{-\lambda(\mu)}{\mu},$$

where  $\lambda(\mu)$  is the principal eigenvalue of the operator

$$\mathcal{L}_\mu \psi(x) := -d(x)\psi''(x) + 2\mu d(x)\psi'(x) - (d(x)\mu^2 + f_s(x, 0))\psi(x) \quad \text{for } x \in \mathbb{R} \setminus S,$$

acting on the set

$$\begin{aligned} E_\mu = \{ \psi \in C(\mathbb{R}) : & \psi|_{\bar{I}} \in C^2(\bar{I}) \text{ for each patch } I, \psi \text{ is periodic in } \mathbb{R}, \\ & [-\mu\psi + \psi'](x^-) = \sigma[-\mu\psi + \psi'](x^+) \text{ for } x \in S_1, \\ & \sigma[-\mu\psi + \psi'](x^-) = [-\mu\psi + \psi'](x^+) \text{ for } x \in S_2 \}, \end{aligned}$$

such that the following statements are valid:

- (i) if  $u$  is the solution to problem (2.9)–(2.10) with a compactly supported initial condition  $u_0 \in \mathcal{C}_p$ , then  $\lim_{t \rightarrow +\infty} \sup_{|x| \geq ct} u(t, x) = 0$  for every  $c > c^*$ ;
- (ii) if  $u_0 \in \mathcal{C}_p$  with  $u_0 \not\equiv 0$ , then  $\lim_{t \rightarrow +\infty} \max_{|x| \leq ct} |u(t, x) - p(x)| = 0$  for every  $0 \leq c < c^*$ .

It finally turns out that the asymptotic spreading speed  $c^*$  is also related to some speeds of rightward or leftward periodic (also called pulsating) traveling waves, whose definition is recalled:

**Definition 2.7.** *A bounded continuous solution  $u : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$  of problem (2.9)–(2.10) is called a periodic rightward traveling wave connecting  $p(x)$  to 0 if it has the form  $u(t, x) = W(x - ct, x)$ , where  $c \in \mathbb{R}$  and the function  $W : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$  has the properties: for each  $s \in \mathbb{R}$  the map  $x \mapsto W(x + s, x)$  is continuous<sup>2</sup> and the map  $x \mapsto W(s, x)$  is periodic, and for each  $x \in \mathbb{R}$  the map  $s \mapsto W(s, x)$  is decreasing with  $W(-\infty, x) = p(x)$  and  $W(+\infty, x) = 0$ .*

*Similarly, a bounded continuous solution  $u : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$  of problem (2.9)–(2.10) is called a periodic leftward traveling wave connecting 0 to  $p(x)$  if it has the form  $u(t, x) = W(x + ct, x)$ , where  $c \in \mathbb{R}$  and the function  $W : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$  has the properties: for each  $s \in \mathbb{R}$  the map  $x \mapsto W(x + s, x)$  is continuous and the map  $x \mapsto W(s, x)$  is periodic, and for each  $x \in \mathbb{R}$  the map  $s \mapsto W(s, x)$  is increasing with  $W(-\infty, x) = 0$  and  $W(+\infty, x) = p(x)$ .*

<sup>2</sup>Notice that the continuity of  $x \mapsto W(x + s, x)$  is automatic if  $c \neq 0$ , since  $u$  is assumed to be continuous itself in  $\mathbb{R} \times \mathbb{R}$ .

The following result shows that the asymptotic spreading speed  $c^*$  given in Theorem 2.6 coincides with minimal speeds of periodic traveling waves in the positive and negative directions.

**Theorem 2.8.** *Assume that the zero solution of (2.14) is unstable (i.e.,  $\lambda_1 < 0$ ) and that  $f$  satisfies (2.11)–(2.12). Let  $c^*$  be the asymptotic spreading speed given in Theorem 2.6. Then the following statements are valid:*

- (i) *problem (2.9)–(2.10) has a periodic rightward traveling wave  $W(x - ct, x)$  connecting  $p(x)$  to 0, in the sense of Definition 2.7, if and only if  $c \geq c^*$ ;*
- (ii) *problem (2.9)–(2.10) has a periodic leftward traveling wave  $W(x + ct, x)$  connecting 0 to  $p(x)$ , in the sense of Definition 2.7, if and only if  $c \geq c^*$ .*

**Remark 2.9.** It is known that for the standard spatially periodic Fisher-KPP problem (1.2) with  $N = 1$ , the variational characterization of minimal speeds in terms of a family of principal eigenvalues implies that the minimal wave speeds of rightward and leftward pulsating waves are the same. Theorem 2.8 shows that this property still holds true for our one-dimensional patchy periodic habitat, with nonstandard movement behavior at interfaces.

**Outline of the paper.** The rest of the paper is organized as follows. In the next section, we give the proof of Theorem 2.2 on the well-posedness of the Cauchy problem (2.9)–(2.10). Section 4 is devoted to the study of the stationary problem (2.14) and we give the proofs of Theorems 2.3 and 2.4. In Section 5, we prove Theorem 2.5 on the large-time behavior of the evolution problem. Finally, Section 6 is devoted to the proofs of Theorems 2.6 and 2.8, based on the abstract monotone semiflow method developed in [30, 31, 49]. Lastly, the appendix is devoted to giving supplementary comparison results concerning finitely many patches, which play an essential role in the well-posedness argument in Section 3.

### 3 Well-posedness of the Cauchy problem (2.9)–(2.10): proof of Theorem 2.2

In this section, we establish the well-posedness of the Cauchy problem (2.9)–(2.10) with nonnegative, bounded and continuous initial data. We first show the existence of classical solutions based on a semigroup argument and an approximation approach. Then we prove that the solutions depend monotonically and continuously on the initial data.

#### 3.1 Truncated problem

Fix  $n \in \mathbb{N}$ . We consider the following truncated problem of (2.9)–(2.10) in the finite interval  $[-nl, nl]$ , which consists of  $4n$  disjoint patches (see Figure 2):

$$\begin{cases} \frac{\partial u_{1m}}{\partial t} = d_1 \frac{\partial^2 u_{1m}}{\partial x^2} + f_1(u_{1m}), & t > 0, \ x \in (ml - l_1, ml), \ m = 0, \pm 1, \dots, \pm(n-1), n, \\ \frac{\partial u_{2m}}{\partial t} = d_2 \frac{\partial^2 u_{2m}}{\partial x^2} + f_2(u_{2m}), & t > 0, \ x \in (ml, ml + l_2), \ m = 0, \pm 1, \dots, \pm(n-1), -n, \end{cases} \quad (3.1)$$

together with interface conditions

$$\begin{cases} u_{1m}(t, x^-) = u_{2m}(t, x^+), & (u_{1m})_x(t, x^-) = \sigma(u_{2m})_x(t, x^+), \\ & t > 0, \ x = ml, \ m = 0, \pm 1, \dots, \pm(n-1), \\ u_{2m}(t, x^-) = u_{1(m+1)}(t, x^+), & \sigma(u_{2m})_x(t, x^-) = (u_{1(m+1)})_x(t, x^+), \\ & t > 0, \ x = ml + l_2, \ m = 0, \pm 1, \dots, \pm(n-1), -n, \end{cases} \quad (3.2)$$

and boundary conditions at  $x = \pm nl$ :

$$u_{1n}(t, (-nl)^+) = u_{2(-n)}(t, (nl)^-) = 0, \quad t > 0. \quad (3.3)$$

For consistency of notations, we set

$$\begin{cases} I_{1m} = (ml - l_1, ml) & \text{for } m \in J_1 = \{0, \pm 1, \dots, \pm(n-1), n\}, \\ I_{2m} = (ml, ml + l_2) & \text{for } m \in J_2 = \{0, \pm 1, \dots, \pm(n-1), -n\}. \end{cases}$$

We number these  $4n$  patches from left to right by  $I_{2(-n)}, I_{1(-n+1)}, \dots, I_{10}, I_{20}, \dots, I_{2(n-1)}, I_{1n}$ , so that

$$[-nl, nl] = \left( \bigcup_{j \in J_1} \overline{I_{1j}} \right) \cup \left( \bigcup_{j \in J_2} \overline{I_{2j}} \right).$$




Figure 2: Truncated interval  $[-nl, nl]$ .

For a solution  $(u_{2(-n)}, \dots, u_{1n})$  of (3.1)–(3.3), we define  $u : (0, +\infty) \times [-nl, nl] \rightarrow \mathbb{R}$  such that, for  $t > 0$ ,

$$\begin{cases} u(t, x) = u_{1j}(t, x) & \text{if } x \in I_{1j} \text{ with } j \in J_1, \\ u(t, x) = u_{2j}(t, x) & \text{if } x \in I_{2j} \text{ with } j \in J_2, \\ u(t, \cdot) \text{ is extended by continuity} & \text{at the interior interfaces and by 0 at } \pm nl. \end{cases} \quad (3.4)$$

We finally set

$$\mathcal{C}_0 = \{\varphi \in C([-nl, nl]) : \varphi(-nl) = \varphi(nl) = 0\}, \quad (3.5)$$

equipped with the sup norm.

**Definition 3.1.** For  $T \in (0, +\infty]$ , we say that a continuous function  $u : [0, T] \times [-nl, nl] \rightarrow \mathbb{R}$  is a classical solution to the truncated problem (3.1)–(3.4) in  $[0, T] \times [-nl, nl]$  with an initial condition  $u_0 \in \mathcal{C}_0$ , if  $u(0, \cdot) = u_0$ , if  $u|_{(0, T) \times \bar{I}} \in C_{t; x}^{1; 2}((0, T) \times \bar{I})$  for each patch  $I \subset [-nl, nl]$ , and if all identities in (3.1)–(3.4) are satisfied pointwise for  $0 < t < T$ .

**Theorem 3.2.** Under the assumption (2.11), the Cauchy problem (3.1)–(3.4) with a nonnegative initial condition  $u_0 \in \mathcal{C}_0$  admits a unique bounded classical solution  $u : [0, +\infty) \times [-nl, nl] \rightarrow \mathbb{R}$ . Furthermore,

$$0 \leq u \leq \max(K_1, K_2, \|u_0\|_{L^\infty(-nl, nl)}) \quad \text{in } [0, +\infty) \times [-nl, nl]. \quad (3.6)$$

If  $0 \leq u_0 \leq v_0$  in  $[-nl, nl]$  with  $u_0, v_0$  in  $\mathcal{C}_0$ , then the solutions  $u$  and  $v$  with respective initial conditions  $u_0$  and  $v_0$  satisfy  $u \leq v$  in  $[0, +\infty) \times [-nl, nl]$ .

The uniqueness and comparison properties immediately follow from Proposition A.2 in the appendix. In what follows, we prove the existence of a bounded classical solution to (3.1)–(3.4), relying on semigroup theory. To do so, we first introduce some auxiliary spaces and recast the truncated Cauchy problem into the abstract form:

$$\begin{cases} \frac{dU}{dt} + AU = f(U), & t > 0, \\ U(0) = U_0, \end{cases} \quad (3.7)$$

where  $U = (u_{2(-n)}, \dots, u_{1n})^T$ ,  $U_0 = (u_0|_{(-nl, -nl+l_2)}, \dots, u_0|_{(nl-l_1, nl)})^T$ , and  $A$  and  $f(U)$  are defined by

$$A = \begin{pmatrix} -d_2 \partial_{xx} & & & & \\ & -d_1 \partial_{xx} & & & \\ & & \ddots & & \\ & & & -d_2 \partial_{xx} & \\ & & & & -d_1 \partial_{xx} \end{pmatrix}_{4n \times 4n} \quad \text{and} \quad f(U) = \begin{pmatrix} f_2(u_{2(-n)}) \\ f_1(u_{1(-n+1)}) \\ \vdots \\ f_2(u_{2(n-1)}) \\ f_1(u_{1n}) \end{pmatrix}. \quad (3.8)$$

Set

$$X = L^2(I_{2(-n)}) \times \dots \times L^2(I_{1n}),$$

with elements viewed as column vectors. With a slight abuse of notation,  $X$  can be identified with  $L^2(-nl, nl)$ . We then define an inner product in  $X$  as follows:

$$\langle U, V \rangle_X = \sum_{j \in J_1} \int_{I_{1j}} u_{1j} v_{1j} + \frac{1}{k} \sum_{j \in J_2} \int_{I_{2j}} u_{2j} v_{2j}, \quad (3.9)$$

which induces the norm  $U \mapsto \|U\|_X = \sqrt{\langle U, U \rangle_X}$  and makes  $X$  a Hilbert space.<sup>3</sup> We also introduce other Hilbert spaces

$$\begin{cases} \mathcal{H}^1 = \{(u_{2(-n)}, \dots, u_{1n})^T \in H^1(I_{2(-n)}) \times \dots \times H^1(I_{1n}), \\ \quad u_{1m}(x) = u_{2m}(x), \quad x = ml, \quad m = 0, \pm 1, \dots, \pm(n-1), \\ \quad u_{2m}(x) = u_{1(m+1)}(x), \quad x = ml + l_2, \quad m = 0, \pm 1, \dots, \pm(n-1), -n\}, \\ \mathcal{H}_0^1 = \{(u_{2(-n)}, \dots, u_{10}, u_{20}, \dots, u_{1n})^T \in \mathcal{H}^1 : u_{2(-n)}(-nl) = u_{1n}(nl) = 0\}, \end{cases}$$

with elements viewed as column vectors, equipped with the norm

$$\|U\|_{\mathcal{H}^1} = \|U\|_{\mathcal{H}_0^1} = \sqrt{\sum_{j \in J_1} \|u_{1j}\|_{H^1(I_{1j})}^2 + \frac{1}{k} \sum_{j \in J_2} \|u_{2j}\|_{H^1(I_{2j})}^2}.$$

From the Sobolev embeddings and with a slight abuse of notation,  $\mathcal{H}^1$  and  $\mathcal{H}_0^1$  can be identified with  $H^1(-nl, nl)$  and  $H_0^1(-nl, nl)$ , respectively, and viewed as subsets of  $C([-nl, nl])$  and  $\mathcal{C}_0$ , respectively, with definition (3.5). Furthermore, in  $\mathcal{H}_0^1$  the norms  $\|U\|_{\mathcal{H}_0^1}$  and  $\|U'\|_X$  are equivalent, from Poincaré's inequality. We finally define the Hilbert space

$$\mathcal{H}^2 = H^2(I_{2(-n)}) \times \dots \times H^2(I_{1n}),$$

with elements viewed as column vectors, equipped with the norm

$$\|U\|_{\mathcal{H}^2} = \sqrt{\sum_{j \in J_1} \|u_{1j}\|_{H^2(I_{1j})}^2 + \frac{1}{k} \sum_{j \in J_2} \|u_{2j}\|_{H^2(I_{2j})}^2},$$

and the subspace

$$\mathcal{D}(A) = \left\{ (u_{2(-n)}, \dots, u_{1n})^T \in \mathcal{H}^2 \cap \mathcal{H}_0^1 : \begin{aligned} & (u_{1m})'(x) = \sigma(u_{2m})'(x), \quad x = ml, \quad m = 0, \pm 1, \dots, \pm(n-1), \\ & \sigma(u_{2m})'(x) = (u_{1(m+1)})'(x), \quad x = ml + l_2, \quad m = 0, \pm 1, \dots, \pm(n-1), -n \end{aligned} \right\}. \quad (3.10)$$

---

<sup>3</sup>We recall that  $k > 0$  is given in (2.3). In all integrals, we integrate with respect to the one-dimensional Lebesgue measure.

From the Sobolev embeddings, the set  $\mathcal{H}^2$  can be viewed as a subset of  $C^1(\overline{I_{2(-n)}}) \times \cdots \times C^1(\overline{I_{1n}})$  and, with a slight abuse of notation, it can also be identified with the set of  $\varphi$  in  $L^2(-nl, nl)$  such that  $\varphi|_I \in H^2(I)$  for each patch  $I \subset [-nl, nl]$ . As for  $\mathcal{D}(A)$ , one has  $\mathcal{D}(A) \subset \mathcal{H}^2 \cap \mathcal{H}_0^1 \subset X$  and, from the Sobolev embeddings,  $\mathcal{D}(A)$  is a Banach space when endowed with the norm  $\|\cdot\|_{\mathcal{H}^2}$ . With a slight abuse of notation,  $\mathcal{D}(A)$  can also be identified with the set of  $\varphi$  in  $H_0^1(-nl, nl)$  such that  $\varphi|_I \in H^2(I)$  for each patch  $I \subset [-nl, nl]$  and  $\varphi$  satisfies the above flux conditions at the interior interfaces.

The proof of the well-posedness of the Cauchy problem (3.1)–(3.4) is based on the following auxiliary lemma.

**Lemma 3.3.** *The linear operator  $A : \mathcal{D}(A) \subset X \rightarrow X$  is symmetric maximal monotone, and  $-A$  is the infinitesimal generator of an analytic semigroup on  $X$ .*

*Proof.* For any  $U = (u_{2(-n)}, \dots, u_{1n})^T \in \mathcal{D}(A)$ , by using (2.8) and the interface and boundary conditions given in the definition of  $\mathcal{D}(A)$ , we have

$$\begin{aligned} \langle AU, U \rangle_X &= \sum_{j \in J_1} \int_{I_{1j}} (-d_1 u_{1j}'') u_{1j} + \frac{1}{k} \sum_{j \in J_2} \int_{I_{2j}} (-d_2 u_{2j}'') u_{2j} \\ &= \sum_{j \in J_1} \int_{I_{1j}} d_1 u_{1j}' u_{1j}' + \frac{1}{k} \sum_{j \in J_2} \int_{I_{2j}} d_2 u_{2j}' u_{2j}' \geq \beta \|U\|_{\mathcal{H}_0^1}^2 \geq 0, \end{aligned} \quad (3.11)$$

where  $\beta > 0$  is a positive constant independent of  $U \in \mathcal{D}(A) \subset \mathcal{H}_0^1$ , given by Poincaré's inequality. Therefore,  $A$  is monotone. The symmetry of  $A$  is also obvious from a similar calculation.

Next, we shall prove that, for every  $\lambda \geq 0$ , the range  $\mathcal{R}(\lambda I_X + A)$  of  $\lambda I_X + A$  is equal to  $X$  ( $I_X$  denotes the identity operator in  $X$ ), that is, for any  $F \in X$ , there exists  $U \in \mathcal{D}(A)$  such that  $\lambda U + AU = F$  (such a  $U$  is then unique from (3.11)). For any  $F = (f_{2(-n)}, \dots, f_{1n})^T \in X$ , we consider the following boundary value problem:

$$\begin{cases} -d_2 u_{2(-n)}'' + \lambda u_{2(-n)} = f_{2(-n)}, & \text{in } (-nl, -nl + l_2), \\ -d_1 u_{1m}'' + \lambda u_{1m} = f_{1m}, & \text{in } (ml - l_1, ml), \quad m = 0, \pm 1, \dots, \pm(n-1), \\ -d_2 u_{2m}'' + \lambda u_{2m} = f_{2m}, & \text{in } (ml, ml + l_2), \quad m = 0, \pm 1, \dots, \pm(n-1), \\ -d_1 u_{1n}'' + \lambda u_{1n} = f_{1n}, & \text{in } (nl - l_1, nl), \end{cases} \quad (3.12)$$

with interface conditions

$$\begin{cases} u_{1m}(x) = u_{2m}(x), & (u_{1m})'(x^-) = \sigma(u_{2m})'(x^+), \\ & x = ml, \quad m = 0, \pm 1, \dots, \pm(n-1), \\ u_{2m}(x) = u_{1(m+1)}(x), & \sigma(u_{2m})'(x^-) = (u_{1(m+1)})'(x^+), \\ & x = ml + l_2, \quad m = 0, \pm 1, \dots, \pm(n-1), -n, \end{cases} \quad (3.13)$$

and boundary conditions

$$u_{2(-n)}(-nl) = u_{1n}(nl) = 0. \quad (3.14)$$

Problem (3.12)–(3.14) is first converted into a weak problem, which consists in finding  $U \in \mathcal{H}_0^1$  such that

$$a(U, V) = \langle F, V \rangle_X \quad \text{for all } V \in \mathcal{H}_0^1, \quad (3.15)$$

where the bilinear form  $a$  is defined in  $\mathcal{H}_0^1 \times \mathcal{H}_0^1$  by

$$a(U, V) = \sum_{j \in J_1} \int_{I_{1j}} (d_1 u_{1j}' v_{1j}' + \lambda u_{1j} v_{1j}) + \frac{1}{k} \sum_{j \in J_2} \int_{I_{2j}} (d_2 u_{2j}' v_{2j}' + \lambda u_{2j} v_{2j}).$$



It is clear that the bilinear form defined in  $\mathcal{H}_0^1 \times \mathcal{H}_0^1$  is continuous and coercive (from Poincaré's inequality again). Then, by the Lax-Milgram theorem, problem (3.15) admits a unique solution  $U \in \mathcal{H}_0^1$ , and we have

$$\|U\|_{\mathcal{H}_0^1} \leq C\|F\|_X, \quad (3.16)$$

for some constant  $C > 0$  only depending on  $d_{1,2}$ ,  $k$ ,  $n$  and  $l_{1,2}$ . Furthermore, owing to the definition of  $a$ , the solution  $U$  belongs to  $\mathcal{D}(A)$  and satisfies (3.12)–(3.14). By rewriting the equations as  $u''_{ij} = (\lambda u_{ij} - f_{ij})/d_i$  for  $j \in J_i$  and  $i \in \{1, 2\}$ , and taking  $L^2$ -norms on both sides, we get  $\|u''_{ij}\|_{L^2(I_{ij})} \leq (1/d_i) \times (\lambda \|u_{ij}\|_{L^2(I_{ij})} + \|f_{ij}\|_{L^2(I_{ij})})$ . By (3.16), we finally obtain the overall  $H^2$  estimate  $\|U\|_{\mathcal{H}^2} \leq C'\|F\|_X$  for some constant  $C' > 0$  only depending on  $d_{1,2}$ ,  $k$ ,  $n$ ,  $l_{1,2}$  and  $\lambda$ . We then conclude that  $\mathcal{R}(\lambda I_X + A) = X$ , that  $\lambda I_X + A$  is invertible from  $\mathcal{D}(A)$  onto  $X$  and that  $(\lambda I_X + A)^{-1}$  is bounded from  $X$  onto  $\mathcal{D}(A)$ . In particular,  $\mathcal{R}(I_X + A) = X$  and the operator  $A$  is maximal monotone. It is then densely defined and closed, and  $\mathcal{D}(A)$  is also a Banach space if endowed with the graph norm  $\|U\|_{\mathcal{D}(A)}$  of  $A$ .

Lastly, let us show that  $-A$  is the infinitesimal generator of an analytic semigroup on  $X$ . First of all, since  $A$  is monotone, one has  $\lambda\|u\|_X \leq \|(\lambda I_X + A)u\|_X$  for every  $\lambda \geq 0$  and  $u \in \mathcal{D}(A)$ , hence

$$\|(\lambda I_X + A)^{-1}\|_{\mathcal{L}(X)} \leq \frac{1}{\lambda} \quad \text{for every } \lambda > 0.$$

Therefore, the Hille-Yosida theorem implies that the operator  $-A$  is the infinitesimal generator of a contraction semigroup on  $X$ . On the other hand, by viewing  $A$  in the complexified Hilbert spaces associated with  $X$  and  $\mathcal{H}^2$ , one sees from (3.11) and the symmetry of  $A$  that the numerical range  $\mathcal{S}(-A)$  of  $-A$  is included in  $\mathbb{R}$  and more precisely in an interval  $(-\infty, -\delta]$ , for some  $\delta > 0$ . Fix any  $\theta \in (0, \pi/2)$  and denote

$$\Sigma_\theta := \{\lambda \in \mathbb{C}^* : |\arg \lambda| < \pi - \theta\}.$$

Then there is  $C_\theta > 0$  such that  $\text{dist}(\lambda, \overline{\mathcal{S}(-A)}) \geq C_\theta|\lambda|$  for all  $\lambda \in \Sigma_\theta$ , where  $\text{dist}(\lambda, \overline{\mathcal{S}(-A)})$  represents the distance in  $\mathbb{C}$  between  $\lambda$  and  $\overline{\mathcal{S}(-A)}$ . We observe that  $\Sigma_\theta \cap \rho(-A) \neq \emptyset$ , since any  $\lambda > 0$  is in the resolvent set  $\rho(-A)$  of  $-A$  from the above analysis. Therefore, [41, Theorem 1.3.9] then states that  $\Sigma_\theta \subset \rho(-A)$  and

$$\|(\lambda I_X + A)^{-1}\|_{\mathcal{L}(X)} \leq \frac{1}{C_\theta|\lambda|} \quad \text{for all } \lambda \in \Sigma_\theta.$$

Since 0 belongs to  $\rho(-A)$  as well, we conclude by [41, Theorem 2.5.2] that  $-A$  is the infinitesimal generator of an analytic semigroup on  $X$ . The proof of Lemma 3.3 is thereby complete.  $\square$

With Lemma 3.3 in hand, we are now ready to prove Theorem 3.2 on the well-posedness of the Cauchy problem (3.1)–(3.4).

*Proof of Theorem 3.2.* The proof is divided into two main steps. The first one assumes an additional hypothesis on the functions  $f_i$  in (2.11), and the second one deals with the general case of  $f_i$  satisfying (2.11).

*Step 1: in addition to (2.11), assume that  $f_1$  and  $f_2$  are globally Lipschitz continuous from  $\mathbb{R}$  to  $\mathbb{R}$ .* The function  $f$  given in (3.8) is then Lipschitz continuous from  $X$  to  $X$ . Therefore, it follows from Lemma 3.3 and [53, Theorem 2.5.1] that, for each  $U_0 \in X$ , problem (3.7) has a unique global mild solution  $U \in C([0, +\infty), X)$ , satisfying

$$U(t) = e^{-tA}U_0 + \int_0^t e^{-(t-s)A}f(U(s)) \, ds \quad (3.17)$$

for all  $t \geq 0$ . Note that the function  $t \mapsto f(U(t))$  belongs to  $C([0, +\infty), X)$  as well. By [32, Lemma 7.1.1], the integral on the right-hand side of (3.17) belongs to  $C_{loc}^\gamma((0, +\infty), X)$  for any  $\gamma \in (0, 1)$ . Since  $t \mapsto e^{-tA}U_0$  is of class  $C^\infty((0, +\infty), \mathcal{D}(A))$ , we see that  $U \in C_{loc}^\gamma((0, +\infty), X)$  for any  $\gamma \in (0, 1)$  and that the function  $t \mapsto f(U(t))$  belongs to  $C_{loc}^\gamma((0, +\infty), X)$  too for any  $\gamma \in (0, 1)$ . It then follows from [32, Theorem 4.3.1] that  $U \in C_{loc}^\gamma((0, +\infty), \mathcal{D}(A)) \cap C_{loc}^{1,\gamma}((0, +\infty), X)$  for any  $\gamma \in (0, 1)$ . As a consequence,  $U$  is a classical solution of (3.7), with equalities in  $X$ .

Furthermore, by Lemma 3.3 and the fact that  $0 \in \rho(A)$ , we can define fractional powers  $A^\beta$  of  $A$ . For  $0 < \beta \leq 1$ ,  $A^\beta$  is a closed operator whose domain  $\mathcal{D}(A^\beta)$  is dense in  $X$  and  $\mathcal{D}(A) \hookrightarrow \mathcal{D}(A^\beta) \hookrightarrow X$  continuously. Endowed with the graph norm  $\|U\|_{\mathcal{D}(A^\beta)}$  of  $A^\beta$ ,  $\mathcal{D}(A^\beta)$  is a Banach space. Since  $A$  is sectorial and  $\inf\{\operatorname{Re}(\lambda) : \lambda \notin \rho(A)\} > 0$ , it follows that  $A^\beta$  is invertible with bounded inverse  $(A^\beta)^{-1} \in \mathcal{L}(X)$  and that the norm  $\|U\|_{\mathcal{D}(A^\beta)}$  is equivalent to  $\|A^\beta U\|_X$  in  $\mathcal{D}(A^\beta)$ . From [43, Lemma 37.8], one has, for each  $1/4 < \beta \leq 1$  and  $\delta \in (0, 2\beta - 1/2)$ , a continuous embedding  $\mathcal{D}(A^\beta) \hookrightarrow \mathcal{C}_0 \cap C^{0,\delta}([-nl, nl])$ .<sup>4</sup> From now on, we fix  $\beta \in (1/4, 1)$ . We also observe that  $f : \mathcal{D}(A^\beta) \rightarrow X$  is globally Lipschitz continuous: indeed, for any  $U, V \in \mathcal{D}(A^\beta)$ , there holds

$$\begin{aligned} \|f(U) - f(V)\|_X &\leq L\|U - V\|_X = L\|(A^\beta)^{-1}A^\beta U - (A^\beta)^{-1}A^\beta V\|_X \\ &\leq L\|(A^\beta)^{-1}\|_{\mathcal{L}(X)}\|A^\beta U - A^\beta V\|_X \\ &\leq L\|(A^\beta)^{-1}\|_{\mathcal{L}(X)}\|U - V\|_{\mathcal{D}(A^\beta)}, \end{aligned}$$

for some constant  $L \in [0, +\infty)$  independent of  $U, V \in \mathcal{D}(A^\beta)$ . Now, for any  $U_0 \in \mathcal{D}(A^\beta) (\subset X)$ , the unique global solution  $U \in C([0, +\infty), X) \cap C_{loc}^\gamma((0, +\infty), \mathcal{D}(A)) \cap C_{loc}^{1,\gamma}((0, +\infty), X)$  (for any  $\gamma \in (0, 1)$ ) of (3.7), given in the previous paragraph, also belongs to  $C([0, +\infty), \mathcal{D}(A^\beta))$  and then to  $C([0, +\infty), \mathcal{C}_0 \cap C^{0,\delta}([-nl, nl]))$  for any  $\delta \in (0, 2\beta - 1/2)$ . Since  $U$  satisfies (3.17) for all  $t \geq 0$ , we then get by [24, Theorem 3.5.2] and [43, Lemma 37.8] the existence of some  $\eta \in (0, 1)$ ,  $\theta \in (1/4, 1)$  and  $\omega \in (0, 2\theta - 1/2)$  such that  $U \in C_{loc}^{1,\eta}((0, +\infty), \mathcal{D}(A^\theta))$  and

$$U \in C_{loc}^{1,\eta}((0, +\infty), \mathcal{C}_0 \cap C^{0,\omega}([-nl, nl])).$$

Since  $\mathcal{D}(A) \hookrightarrow \mathcal{D}(A^\beta)$ , it follows from the previous two paragraphs that, for any  $U_0 \in X$ , the solution  $U \in C([0, +\infty), X) \cap C((0, +\infty), \mathcal{D}(A)) \cap C^1((0, +\infty), X)$  of (3.7) belongs to

$$C((0, +\infty), \mathcal{C}_0 \cap C^{0,\delta}([-nl, nl])) \cap C_{loc}^{1,\eta}((0, +\infty), \mathcal{C}_0 \cap C^{0,\omega}([-nl, nl])).$$

Moreover, if  $U_0 \in \mathcal{D}(A^\beta)$ , then  $U \in C([0, +\infty), \mathcal{C}_0 \cap C^{0,\delta}([-nl, nl]))$ . One infers that, for any  $U_0 \in X$ , the function  $u$  defined as in (3.4) (with similar definition at  $t = 0$ ) is continuous in  $(0, +\infty) \times [-nl, nl]$ , vanishes on  $(0, +\infty) \times \{\pm nl\}$ , is of class  $C^1$  with respect to  $t$  in  $(0, +\infty) \times [-nl, nl]$ , with  $u$  and  $\frac{\partial u}{\partial t}$  Hölder continuous in  $[\tau, \tau'] \times [-nl, nl]$  for every  $0 < \tau < \tau' < +\infty$ . Therefore, for each patch  $I \subset [-nl, nl]$  of type  $i \in \{1, 2\}$  and for each  $0 < \tau < \tau' < +\infty$ , the function  $f_i(u)$  is Hölder continuous in  $[\tau, \tau'] \times \bar{I}$ , hence equation (3.7) implies that  $u|_{[\tau, \tau'] \times \bar{I}}$  is of class  $C^2$  with respect to  $x$  and  $\frac{\partial^2 u}{\partial x^2}|_{[\tau, \tau'] \times \bar{I}}$  is Hölder continuous in  $[\tau, \tau'] \times \bar{I}$ . In particular,  $u$  is a classical solution of (3.1)–(3.4) for  $t > 0$ . Furthermore, if  $U_0 \in \mathcal{D}(A^\beta)$ , then  $u$  is also continuous in  $[0, +\infty) \times [-nl, nl]$ .

It remains to show, still in this step 1, that  $u$  is bounded and continuous up to  $t = 0$  when  $u_0 \in \mathcal{C}_0$ . To do so, we first prove a comparison principle for the solutions when the initial conditions are in  $X$ . Take any  $V_0, W_0 \in X$  such that  $v_0 \leq w_0$  almost everywhere in  $[-nl, nl]$ ,

<sup>4</sup>With a slight abuse of notation, the embedding  $\mathcal{D}(A^\beta) \hookrightarrow \mathcal{C}_0 \cap C^{0,\delta}([-nl, nl])$  means that the elements  $U = (u_{2(-n)}, \dots, u_{1n})^T$  of  $\mathcal{D}(A^\beta)$  have continuous components  $u_{ij}$  in each corresponding closed patch  $\bar{I}_{ij}$ , and that the function equal to each  $u_{ij}$  on each closed patch  $\bar{I}_{ij}$  is well defined, continuous in  $[-nl, nl]$ , vanishes at  $\pm nl$  and is Hölder continuous of exponent  $\delta$  in  $[-nl, nl]$ , with a sup norm and a Hölder norm controlled by  $\|U\|_{\mathcal{D}(A^\beta)}$ .

with obvious notations for  $v_0$  and  $w_0$ . There exist then two sequences  $(V_{0j})_{j \in \mathbb{N}}$  and  $(W_{0j})_{j \in \mathbb{N}}$  in  $\mathcal{D}(A) (\subset \mathcal{D}(A^\beta) \subset \mathcal{C}_0)$  such that  $v_{0j} \leq w_{0j}$  everywhere in  $[-nl, nl]$  (with obvious notations) for all  $j \in \mathbb{N}$ , and  $V_{0j} \rightarrow V_0$ ,  $W_{0j} \rightarrow W_0$  in  $X$  as  $j \rightarrow +\infty$ . For each  $j \in \mathbb{N}$ , with obvious notations, let  $v_j$  and  $w_j$  be the classical solutions of (3.1)–(3.4) with initial conditions  $v_{0j}$  and  $w_{0j}$ . The functions  $v_j$  and  $w_j$  are continuous in  $[0, +\infty) \times [-nl, nl]$ , from the previous paragraph. Therefore, the maximum principle of Proposition A.2 implies that

$$v_j \leq w_j \text{ in } [0, +\infty) \times [-nl, nl],$$

for all  $j \in \mathbb{N}$ . Since, for each  $t \geq 0$ , the map  $U_0 \mapsto U(t)$  given by (3.17) is continuous (and even Lipschitz continuous) from  $X$  to  $X$  by [53, Theorem 2.5.1], one infers that, for each  $t > 0$ ,  $v(t, \cdot) \leq w(t, \cdot)$  almost everywhere in  $(-nl, nl)$  and then everywhere in  $[-nl, nl]$  by continuity. To sum up,

$$v \leq w \text{ in } [0, +\infty) \times [-nl, nl]. \quad (3.18)$$

If  $u_0 \in \mathcal{C}_0$  with  $u_0 \geq 0$  in  $[-nl, nl]$ , one can also assume without loss of generality that  $0 \leq u_{0k} \leq \|u_0\|_{L^\infty(-nl, nl)}$  in  $[-nl, nl]$ . Remembering (2.11), the constant functions equal to 0 and  $\max(K_1, K_2, \|u_0\|_{L^\infty(-nl, nl)})$  are, respectively, a subsolution and a supersolution, in the sense of Definition A.1, of the problem (3.1)–(3.4) satisfied by the continuous and classical solution  $u_k$  in  $[0, +\infty) \times [-nl, nl]$ . The maximum principle of Proposition A.2 then yields  $0 \leq u_k \leq \max(K_1, K_2, \|u_0\|_{L^\infty(-nl, nl)})$  in  $[0, +\infty) \times [-nl, nl]$  for all  $k \in \mathbb{N}$ , hence

$$0 \leq u \leq \max(K_1, K_2, \|u_0\|_{L^\infty(-nl, nl)}) \text{ in } (0, +\infty) \times [-nl, nl], \quad (3.19)$$

by passing to the limit as  $k \rightarrow +\infty$  for each  $t > 0$ , as in the previous paragraph. Notice that (3.19) holds as well on  $\{0\} \times [-nl, nl]$  by assumption on  $u_0$ .

Lastly, consider again any  $u_0 \in \mathcal{C}_0 (\subset X)$  in  $[-nl, nl]$  and let us show that  $u$  is continuous up to time  $t = 0$ . Let  $\varepsilon > 0$  be arbitrary. Let  $\underline{u}_0$  and  $\bar{u}_0$  be two functions in  $\mathcal{D}(A) (\subset \mathcal{D}(A^\beta) \subset X)$  such that

$$u_0 - \varepsilon \leq \underline{u}_0 \leq u_0 \leq \bar{u}_0 \leq u_0 + \varepsilon \text{ in } [-nl, nl]$$

(with obvious notations for the functions  $\underline{u}_0$  and  $\bar{u}_0$ , which can be chosen in  $C^2([-nl, nl]) \cap \mathcal{C}_0$  with zero derivatives at the interior interfaces) and let  $\underline{u}$  and  $\bar{u}$  be the two classical solutions of (3.1)–(3.4) with initial conditions  $\underline{u}_0$  and  $\bar{u}_0$ . From the above arguments, the functions  $\underline{u}$  and  $\bar{u}$  are continuous in  $[0, +\infty) \times [-nl, nl]$ , and  $\underline{u} \leq u \leq \bar{u}$  in  $[0, +\infty) \times [-nl, nl]$  from (3.18). Finally, there is  $t_0 > 0$  such that

$$u_0 - 2\varepsilon \leq \underline{u}_0 - \varepsilon \leq \underline{u} \leq u \leq \bar{u} \leq \bar{u}_0 + \varepsilon \leq u_0 + 2\varepsilon \text{ in } [0, t_0] \times [-nl, nl],$$

from which it follows that the  $C((0, +\infty) \times [-nl, nl])$  function  $u$  is also continuous up to time  $t = 0$ . It is therefore a bounded classical solution of (3.1)–(3.4) in  $[0, +\infty) \times [-nl, nl]$  with initial condition  $u_0$ , in the sense of Definition 3.1.

*Step 2: the general case of assumption (2.11).* Consider a nonnegative initial condition  $u_0$  in  $\mathcal{C}_0$ . Denote  $K = \max(K_1, K_2, \|u_0\|_{L^\infty(-nl, nl)})$  and, for  $i = 1, 2$ , let  $\tilde{f}_i : \mathbb{R} \rightarrow \mathbb{R}$  be a globally Lipschitz continuous function of class  $C^1(\mathbb{R})$  such that  $\tilde{f}_i|_{[0, K_i]} = f_i|_{[0, K_i]}$  and  $\tilde{f}_i \leq 0$  in  $[K_i, +\infty)$ . From Step 1, there is a unique bounded classical solution  $u$  of (3.1)–(3.4) in  $[0, +\infty) \times [-nl, nl]$  with initial condition  $u_0$ , but with the nonlinearities  $\tilde{f}_i$  instead of  $f_i$ , and  $u$  satisfies (3.19) in  $[0, +\infty) \times [-nl, nl]$ . From (3.19) and the choice of  $\tilde{f}_i$ , the function  $u$  is then a bounded classical solution of the problem (3.1)–(3.4) in  $[0, +\infty) \times [-nl, nl]$  with initial condition  $u_0$  and with the original nonlinearities  $f_i$ .

Since the uniqueness and comparison properties in Theorem 3.2 directly follow from Proposition A.2, the proof of Theorem 3.2 is thereby complete.  $\square$

### 3.2 Proof of Theorem 2.2

We prove the existence of a solution to the problem (2.9)–(2.10) through a truncation and approximation argument. Set

$$\varepsilon := \min\left(\frac{l_1}{4}, \frac{l_2}{4}\right) > 0.$$

We first fix a sequence of cut-off functions  $(\delta^n)_{n \in \mathbb{N}}$  in  $C(\mathbb{R})$  such that, for each  $n \in \mathbb{N}$ ,

$$0 \leq \delta^n \leq 1 \text{ in } \mathbb{R}, \text{ and } \delta^n(x) := \begin{cases} 1 & \text{if } x \in [-nl - \varepsilon + l_2, nl - l_1 + \varepsilon], \\ 0 & \text{if } x \notin (-nl, nl). \end{cases} \quad (3.20)$$

We can for instance define uniquely  $\delta^n$  by also assuming that  $\delta^n$  is affine in  $[-nl, -nl - \varepsilon + l_2]$  and in  $[nl - l_1 + \varepsilon, nl]$ .

Let us now take any nonnegative bounded continuous function  $u_0 : \mathbb{R} \rightarrow \mathbb{R}$ . For each  $n \in \mathbb{N}$ , we consider the truncated problem (3.1)–(3.4) on  $[-nl, nl]$ , with initial condition  $\delta^n u_0|_{[-nl, nl]}$ . This problem involves  $4n$  patches still identified by  $I_{2(-n)}, I_{1(-n+1)}, \dots, I_{2(n-1)}, I_{1n}$  as before, and the nonnegative initial condition  $\delta^n u_0|_{[-nl, nl]}$  belongs to the space  $\mathcal{C}_0$  defined in (3.5). Therefore, by Theorem 3.2, there is a unique bounded classical solution  $u^n$  of (3.1)–(3.4) in  $[0, +\infty) \times [-nl, nl]$  with initial condition  $\delta^n u_0|_{[-nl, nl]}$ , and  $u^n$  also satisfies (3.6) in  $[0, +\infty) \times [-nl, nl]$ . Moreover, for every  $m < n \in \mathbb{N}$ , one has

$$0 \leq u^m(0, \cdot) = \delta^m u_0|_{[-ml, ml]} \leq u_0|_{[-ml, ml]} = \delta^n u_0|_{[-ml, ml]} = u^n(0, \cdot) \text{ in } [-nl, nl]$$

and  $u^n(t, \pm ml) \geq 0$  for all  $t \geq 0$  by (3.6), hence

$$u^m \leq u^n|_{[0, +\infty) \times [-ml, ml]} \text{ in } [0, +\infty) \times [-ml, ml]$$

by Proposition A.2. As a consequence, for each  $(t, x) \in [0, +\infty) \times \mathbb{R}$ , the sequence  $(u^n(t, x))_{n \geq |x|/l}$  is non-decreasing and ranges in  $[0, K]$  with

$$K := \max(K_1, K_2, \|u_0\|_{L^\infty(\mathbb{R})}), \quad (3.21)$$

hence the sequence  $(u^n(t, x))_{n \geq |x|/l}$  converges to a quantity  $u(t, x) \in [0, K]$ , that is,

$$u^n(t, x) \rightarrow u(t, x) \in [0, K] \text{ as } n \rightarrow +\infty. \quad (3.22)$$

Notice also that, if  $u_0 \not\equiv 0$ , then  $u^n(0, \cdot) \geq \not\equiv 0$  in  $[-nl, nl]$  for all  $n$  large enough, hence  $u^n > 0$  in  $(0, +\infty) \times (-nl, nl)$  for all  $n$  large enough by Proposition A.2, and finally  $u(t, x) > 0$  for all  $(t, x) \in (0, +\infty) \times \mathbb{R}$  because the sequence  $(u^n(t, x))_{n \geq |x|/l}$  is non-decreasing for each  $(t, x) \in [0, +\infty) \times \mathbb{R}$ .

In order to show that  $u$  is a classical solution of (2.9)–(2.10), we need further differential estimates on the sequence  $(u^n)_{n \in \mathbb{N}}$ . Consider any  $0 < \tau \leq \tau' < +\infty$  and any patch  $I \subset \mathbb{R}$ . Let us assume that  $I$  is of type 1, that is,  $I = I_{1m} = (ml - l_1, ml)$  for some  $m \in \mathbb{Z}$  (the case of a patch  $I$  of type 2 can be dealt with similarly). Let us fix an arbitrary  $\nu \in (0, 1/2)$ , say for instance  $\nu = 1/4$ . Since the solutions  $u^n$  (for  $n \geq |m| + 1$ ) of (3.1)–(3.4) are uniformly bounded in  $[0, +\infty) \times [-(|m| + 1)l, (|m| + 1)l]$ , it follows from standard interior parabolic estimates that

$$\sup_{n \geq |m|+1} \|u^n(\cdot, ml - l_1 - \varepsilon)\|_{C^{1,\nu}([\tau/2, +\infty))} + \sup_{n \geq |m|+1} \|u^n(\cdot, ml + \varepsilon)\|_{C^{1,\nu}([\tau/2, +\infty))} \leq C_0,$$

for some positive constant  $C_0$  only depending on  $\tau, l_{1,2}, d_2, f_2$  and  $K$  given in (3.21), hence on  $\tau, l_{1,2}, d_2, f_{1,2}$  and  $\|u_0\|_{L^\infty(\mathbb{R})}$ . Consider then two  $C^3([ml - l_1 - \varepsilon, ml + \varepsilon])$  functions  $g : [ml - l_1 - \varepsilon, ml + \varepsilon] \rightarrow [0, 1]$  and  $h : [ml - l_1 - \varepsilon, ml + \varepsilon] \rightarrow [0, 1]$  such that

$$\begin{cases} g(ml - l_1 - \varepsilon) = h(ml + \varepsilon) = 0, \\ g(ml + \varepsilon) = h(ml - l_1 - \varepsilon) = 1, \\ g'(ml - l_1) = h'(ml - l_1) = g'(ml) = h'(ml) = 0. \end{cases}$$

They can be chosen so that their  $C^3([ml - l_1 - \varepsilon, ml + \varepsilon])$  norms only depend on  $l_{1,2}$ . Consider now, for each  $n \geq |m| + 1$ , the function  $\tilde{u}^n$  defined in  $[\tau/2, +\infty) \times [ml - l_1 - \varepsilon, ml + \varepsilon]$  by

$$\tilde{u}^n(t, x) = u^n(t, x) - h(x)u^n(t, ml - l_1 - \varepsilon) - g(x)u^n(t, ml + \varepsilon). \quad (3.23)$$

Each such function  $\tilde{u}^n$  is continuous in  $[\tau/2, +\infty) \times [ml - l_1 - \varepsilon, ml + \varepsilon]$  and has restrictions of class  $C_{t,x}^{1,2}$  in  $[\tau/2, +\infty) \times [ml - l_1 - \varepsilon, ml - l_1]$ , in  $[\tau/2, +\infty) \times [ml - l_1, ml] = [\tau/2, +\infty) \times \bar{I}$  and in  $[\tau/2, +\infty) \times [ml, ml + \varepsilon]$ . Furthermore, from (3.1)–(3.4) and (3.23), one has

$$\left\{ \begin{array}{ll} \frac{\partial \tilde{u}^n}{\partial t} = d_2 \frac{\partial^2 \tilde{u}^n}{\partial x^2} + \tilde{f}_2^n(t, x, \tilde{u}^n(t, x)), & t \geq \tau/2, x \in (ml - l_1 - \varepsilon, ml - l_1), \\ \frac{\partial \tilde{u}^n}{\partial t} = d_1 \frac{\partial^2 \tilde{u}^n}{\partial x^2} + \tilde{f}_1^n(t, x, \tilde{u}^n(t, x)), & t \geq \tau/2, x \in (ml - l_1, ml), \\ \frac{\partial \tilde{u}^n}{\partial t} = d_2 \frac{\partial^2 \tilde{u}^n}{\partial x^2} + \tilde{f}_2^n(t, x, \tilde{u}^n(t, x)), & t \geq \tau/2, x \in (ml, ml + \varepsilon), \\ \tilde{u}^n(t, ml - l_1 - \varepsilon) = \tilde{u}^n(t, ml + \varepsilon) = 0, & t \geq \tau/2, \\ \tilde{u}^n(t, (ml - l_1)^-) = \tilde{u}^n(t, (ml - l_1)^+), & t \geq \tau/2, \\ \sigma \tilde{u}_x^n(t, (ml - l_1)^-) = \tilde{u}_x^n(t, (ml - l_1)^+), & t \geq \tau/2, \\ \tilde{u}^n(t, (ml)^-) = \tilde{u}^n(t, (ml)^+), & t \geq \tau/2, \\ \tilde{u}_x^n(t, (ml)^-) = \sigma \tilde{u}_x^n(t, (ml)^+), & t \geq \tau/2, \end{array} \right.$$

with

$$\begin{aligned} \tilde{f}_i^n(t, x, s) &= f_i(s + h(x)u^n(t, ml - l_1 - \varepsilon) + g(x)u^n(t, ml + \varepsilon)) \\ &\quad - h(x)u_t^n(t, ml - l_1 - \varepsilon) - g(x)u_t^n(t, ml + \varepsilon) \\ &\quad + d_i h''(x)u^n(t, ml - l_1 - \varepsilon) + d_i g''(x)u^n(t, ml + \varepsilon). \end{aligned}$$

In other words, each function  $\tilde{u}^n$  solves a truncated problem similar to (3.1)–(3.4), but this time on the interval  $[ml - l_1 - \varepsilon, ml + \varepsilon]$  (with only three patches) and with nonlinearities  $\tilde{f}_i^n(t, x, s)$  which are still of class  $C^1$  with respect to  $s$ , with partial derivatives equal to  $f'_i(s + h(x)u^n(t, ml - l_1 - \varepsilon) + g(x)u^n(t, ml + \varepsilon))$ , and are now Hölder continuous of any exponent  $\nu$  with respect to  $(t, x) \in [\tau/2, +\infty) \times [ml - l_1 - \varepsilon, ml + \varepsilon]$  uniformly with respect to  $s$  and  $n$ . Remember that  $\tau' \geq \tau$ , hence  $\tau' - \tau/2 \geq \tau/2$ . Since the sequence  $(\tilde{u}^n(\tau' - \tau/2, \cdot))_{n \geq |m|+1}$  is bounded in particular in  $L^2(ml - l_1 - \varepsilon, ml + \varepsilon)$ , it then follows with similar notations and arguments as in the proof of Theorem 3.2 that there is a universal constant  $\gamma \in (0, 1)$  such that the sequence

$$\left( \tilde{u}^n(\tau' - \tau/4, \cdot)|_{(ml-l_1-\varepsilon, ml-l_1)}, \tilde{u}^n(\tau' - \tau/4, \cdot)|_{(ml-l_1, ml)}, \tilde{u}^n(\tau' - \tau/4, \cdot)|_{(ml, ml+\varepsilon)} \right)_{n \geq |m|+1}$$

is bounded in the set  $\mathcal{D}(A)$  (defined as in (3.10), but with now only three patches) and the sequences  $(\tilde{u}^n)_{n \geq |m|+1}$  and  $(\tilde{u}_t^n)_{n \geq |m|+1}$  are bounded in  $C^\gamma([\tau', \tau' + 1] \times [ml - l_1 - \varepsilon, ml + \varepsilon])$ , with bounds depending only on  $\sup_{n \geq |m|+1} \|\tilde{u}^n(\tau' - \tau/2, \cdot)\|_{L^2(ml-l_1-\varepsilon, ml+\varepsilon)}$ ,  $\tau$ ,  $l_{1,2}$ ,  $d_{1,2}$ ,  $f_{1,2}$  and  $\sigma$ , hence only on  $\tau$ ,  $l_{1,2}$ ,  $d_{1,2}$ ,  $f_{1,2}$ ,  $\sigma$  and  $\|u_0\|_{L^\infty(\mathbb{R})}$  (notice that these bounds are independent of  $\tau' \in [\tau, +\infty)$ ). Owing to the definitions of  $\tilde{f}_i^n$  and  $\tilde{u}^n$ , one infers that the sequence  $(\tilde{f}_i^n(\cdot, \cdot, \tilde{u}^n(\cdot, \cdot)))_{n \geq |m|+1}$  is bounded in  $C^\gamma([\tau', \tau' + 1] \times [ml - l_1 - \varepsilon, ml + \varepsilon])$ , hence so is the sequence

$$\left( \frac{\partial^2 \tilde{u}^n|_{[\tau', \tau'+1] \times [ml-l_1-\varepsilon, ml-l_1]}}{\partial x^2}, \frac{\partial^2 \tilde{u}^n|_{[\tau', \tau'+1] \times [ml-l_1, ml]}}{\partial x^2}, \frac{\partial^2 \tilde{u}^n|_{[\tau', \tau'+1] \times [ml, ml+\varepsilon]}}{\partial x^2} \right)_{n \geq |m|+1}$$

in  $C^\gamma([\tau', \tau' + 1] \times [ml - l_1 - \varepsilon, ml - l_1]) \times C^\gamma([\tau', \tau' + 1] \times [ml - l_1, ml]) \times C^\gamma([\tau', \tau' + 1] \times [ml, ml + \varepsilon])$ , with bounds depending only on  $\tau$ ,  $l_{1,2}$ ,  $d_{1,2}$ ,  $f_{1,2}$ ,  $\sigma$  and  $\|u_0\|_{L^\infty(\mathbb{R})}$ . Finally, using (3.23) again, the

sequence  $(u^n|_{[\tau', \tau'+1] \times \bar{I}})_{n \geq m+1}$  is bounded in  $C_{t;x}^{1,\gamma;2,\gamma}([\tau', \tau'+1] \times \bar{I})$ , and, since the bound does not depend on  $\tau' \in [\tau, +\infty)$ , the sequence  $(u^n|_{[\tau, +\infty) \times \bar{I}})_{n \geq m+1}$  is bounded in  $C_{t;x}^{1,\gamma;2,\gamma}([\tau, +\infty) \times \bar{I})$  by a constant depending only on  $\tau, l_{1,2}, d_{1,2}, f_{1,2}, \sigma$  and  $\|u_0\|_{L^\infty(\mathbb{R})}$ .

From the Arzelà-Ascoli theorem and the uniqueness of the limit  $u$  in (3.22), it follows that  $u^n \rightarrow u$  as  $n \rightarrow +\infty$  in  $C_{t;x}^{1;2}([\tau_1, \tau_2] \times \bar{I})$  for every  $0 < \tau_1 \leq \tau_2$  and every patch  $I \subset \mathbb{R}$ , hence  $u$  is a bounded classical solution of (2.9)–(2.10) in  $(0, +\infty) \times \mathbb{R}$ . Furthermore, for every  $\tau > 0$  and every patch  $I \subset \mathbb{R}$ , there holds

$$\|u|_{[\tau, +\infty) \times \bar{I}}\|_{C_{t;x}^{1,\gamma;2,\gamma}([\tau, +\infty) \times \bar{I})} \leq C,$$

for some constant  $C$  only depending on  $\tau, l_{1,2}, d_{1,2}, f_{1,2}, \sigma$  and  $\|u_0\|_{L^\infty(\mathbb{R})}$ .

Next, we shall prove the continuity of the function  $u$  up to time  $t = 0$ . Fix any  $x_0 \in \mathbb{R}$ ,  $R > 0$ , and  $\eta > 0$ . With  $K = \max(K_1, K_2, \|u_0\|_{L^\infty(\mathbb{R})})$  as in (3.21), one can choose two nonnegative functions  $\underline{u}_0$  and  $\bar{u}_0$  in  $C^2(\mathbb{R}) \cap L^\infty(\mathbb{R})$  such that both  $\underline{u}_0$  and  $K - \bar{u}_0$  are supported in  $[x_0 - 2R, x_0 + 2R]$ , and such that

$$0 \leq \underline{u}_0(x) \leq u_0(x) \leq \bar{u}_0(x) \leq K \quad \text{for all } x \in [x_0 - 2R, x_0 + 2R], \quad (3.24)$$

and

$$u_0(x) - \eta \leq \underline{u}_0(x) \leq u_0(x) \leq \bar{u}_0(x) \leq u_0(x) + \eta \quad \text{for all } x \in [x_0 - R, x_0 + R]. \quad (3.25)$$

These functions  $\underline{u}_0$  and  $\bar{u}_0$  can also be chosen so that their derivatives vanish at all interface points in  $[x_0 - 2R, x_0 + 2R]$ . There are then  $B > 0$  large enough and  $t_0 > 0$  small enough such that  $Bt_0 \leq 1$  and the  $C_{t;x}^{1;2}([0, +\infty) \times \mathbb{R})$  functions  $(t, x) \mapsto \underline{u}_0(x) - Bt$  and  $(t, x) \mapsto \bar{u}_0(x) + Bt$  are, respectively, a sub- and a supersolution of truncated problem (3.1)–(3.4) for  $(t, x) \in [0, t_0] \times [-nl, nl]$  and for any  $n \in \mathbb{N}$  large enough so that  $[x_0 - 2R, x_0 + 2R] \subset [-nl, nl]$  and  $\delta^n u_0 = u_0$  in  $[x_0 - 2R, x_0 + 2R]$ , where  $\delta^n$  is the cut-off function defined in (3.20). Remembering (3.24), the inequality (3.6) satisfied by  $u^n$  and the fact that  $\underline{u}_0$  and  $K - \bar{u}_0$  are supported in  $[x_0 - 2R, x_0 + 2R]$  (hence,  $\underline{u}_0(\pm nl) - Bt \leq 0 \leq K \leq \bar{u}_0(\pm nl) + Bt$  for all  $t \geq 0$ ), it follows from Proposition A.2 that

$$\underline{u}_0(x) - Bt \leq u^n(t, x) \leq \bar{u}_0(x) + Bt \quad \text{for all } (t, x) \in [0, t_0] \times [-nl, nl]$$

and for all  $n$  large enough. By passing to the limit  $n \rightarrow +\infty$  for any  $(t, x) \in (0, t_0] \times \mathbb{R}$ , one gets that

$$\underline{u}_0(x) - Bt \leq u(t, x) \leq \bar{u}_0(x) + Bt \quad \text{for all } (t, x) \in (0, t_0] \times \mathbb{R}.$$

Together with (3.25), there is then  $t_1 > 0$  such that  $|u(t, x) - u_0(x)| \leq 2\eta$  for all  $(t, x) \in (0, t_1] \times [x_0 - R, x_0 + R]$ . Finally, since  $\eta > 0$  was arbitrary, this shows that  $u$  is continuous up to time  $t = 0$ , and that  $u(t, \cdot) \rightarrow u_0$  locally uniformly as  $t \rightarrow 0^+$ . To sum up,  $u$  is a nonnegative bounded classical solution of (2.9)–(2.10) in  $[0, +\infty) \times \mathbb{R}$  with initial condition  $u_0$ , in the sense of Definition 2.1.

We point out that the definition of the solution  $u$  does not depend on the choice of the sequence of cut-off functions  $(\delta^n)_{n \in \mathbb{N}}$  given in (3.20). So, let  $(\hat{\delta}^n)_{n \in \mathbb{N}}$  be another such sequence, and, for each  $n \in \mathbb{N}$ , let  $\hat{u}^n$  be the unique bounded classical solution of (3.1)–(3.4) with initial condition  $\hat{\delta}^n u_0|_{[-nl, nl]}$ . As above, the sequence  $(\hat{u}^n)_{n \in \mathbb{N}}$  converges monotonically pointwise in  $[0, +\infty) \times \mathbb{R}$  to a nonnegative bounded classical solution  $\hat{u}$  of (2.9)–(2.10) in  $[0, +\infty) \times \mathbb{R}$ . Furthermore, for every  $n \in \mathbb{N}$ , one has

$$u^n(0, \cdot) = \delta^n u_0|_{[-nl, nl]} \leq u_0|_{[-nl, nl]} = \hat{\delta}^{n+1} u_0|_{[-nl, nl]} = \hat{u}^{n+1}(0, \cdot)|_{[-nl, nl]} \quad \text{in } [-nl, nl].$$



Since  $\hat{u}^{n+1}$  is nonnegative in  $[0, +\infty) \times [-(n+1)l, (n+1)l]$  and in particular on  $[0, +\infty) \times \{\pm nl\}$ , Proposition A.2 implies that  $u^n \leq \hat{u}^{n+1}$  in  $[0, +\infty) \times [-nl, nl]$ . The passage to the limit as  $n \rightarrow +\infty$  pointwise in  $[0, +\infty) \times \mathbb{R}$  yields  $u \leq \hat{u}$  in  $[0, +\infty) \times \mathbb{R}$ . The reverse inequality also holds similarly, hence  $u \equiv \hat{u}$  in  $[0, +\infty) \times \mathbb{R}$ .

Consider now the case where  $u_0$  is periodic, that is,  $u_0(x) = u_0(x+l)$  for all  $x \in \mathbb{R}$ . With the same notations as above, for every  $n \in \mathbb{N}$ , one has

$$u^n(0, x) = \delta^n(x)u_0(x) \leq \delta^{n+2}(x+l)u_0(x+l) = u^{n+2}(0, x+l) \quad \text{for all } x \in [-nl, nl].$$

Since  $u^{n+2}(t, -nl+l) \geq 0$  and  $u^{n+2}(t, nl+l) \geq 0$  for all  $t \geq 0$ , Proposition A.2 implies that  $u^n(t, x) \leq u^{n+2}(t, x+l)$  for all  $(t, x) \in [0, +\infty) \times [-nl, nl]$ , hence  $u(t, x) \leq u(t, x+l)$  for all  $(t, x) \in [0, +\infty) \times \mathbb{R}$ . The reverse inequality can be proved similarly. Therefore,

$$u(t, x) = u(t, x+l) \quad \text{for all } (t, x) \in [0, +\infty) \times \mathbb{R}.$$

Let us then show a comparison principle for the solutions constructed above. Consider any two nonnegative bounded continuous functions  $u_0$  and  $v_0$  such that

$$0 \leq u_0 \leq v_0 \text{ in } \mathbb{R}.$$

For each  $n \in \mathbb{N}$ , let  $u^n$  and  $v^n$  be the unique bounded classical solution of (3.1)–(3.4) with respective initial conditions  $\delta^n u_0|_{[-nl, nl]}$  and  $\delta^n v_0|_{[-nl, nl]}$ . The sequences  $(u^n)_{n \in \mathbb{N}}$  and  $(v^n)_{n \in \mathbb{N}}$  converge monotonically pointwise in  $[0, +\infty) \times \mathbb{R}$  to nonnegative bounded classical solutions  $u$  and  $v$  of (2.9)–(2.10) in  $[0, +\infty) \times \mathbb{R}$ . Since for each  $n \in \mathbb{N}$ , one has  $u^n(0, \cdot) \leq v^n(0, \cdot)$  in  $[-nl, nl]$ , Proposition A.2 yields  $u^n \leq v^n$  in  $[0, +\infty) \times [-nl, nl]$ , hence

$$u \leq v \text{ in } [0, +\infty) \times \mathbb{R}$$

at the limit  $n \rightarrow +\infty$ .

Finally, let us show the local-in-time continuous dependence of the solutions  $u$  with respect to the initial condition. We actually show more, that is, for each  $T > 0$ , the map  $u_0 \mapsto u$  is Lipschitz continuous from  $C^+(\mathbb{R}) \cap L^\infty(\mathbb{R})$  to  $C([0, T] \times \mathbb{R}) \cap L^\infty([0, T] \times \mathbb{R})$  equipped with the sup norms. Consider two functions  $u_0, v_0$  in  $C^+(\mathbb{R}) \cap L^\infty(\mathbb{R})$ , and denote

$$\underline{u}_0 = \max(0, u_0 - \|u_0 - v_0\|_{L^\infty(\mathbb{R})}) \quad \text{and} \quad \bar{u}_0 = u_0 + \|u_0 - v_0\|_{L^\infty(\mathbb{R})}.$$

Then,  $\underline{u}_0$  and  $\bar{u}_0$  are in  $C^+(\mathbb{R}) \cap L^\infty(\mathbb{R})$ , and  $0 \leq \underline{u}_0 \leq \min(u_0, v_0) \leq \max(u_0, v_0) \leq \bar{u}_0$  in  $\mathbb{R}$ , with  $\|\bar{u}_0 - \underline{u}_0\|_{L^\infty(\mathbb{R})} \leq 2\|u_0 - v_0\|_{L^\infty(\mathbb{R})}$ . For each  $n \in \mathbb{N}$ , let  $u^n, v^n, \underline{u}^n$  and  $\bar{u}^n$  be the unique bounded classical solutions of (3.1)–(3.4) with respective initial conditions  $\delta^n u_0|_{[-nl, nl]}$ ,  $\delta^n v_0|_{[-nl, nl]}$ ,  $\delta^n \underline{u}_0|_{[-nl, nl]}$  and  $\delta^n \bar{u}_0|_{[-nl, nl]}$ , with the cut-off function  $\delta^n$  given in (3.20). The sequences  $(u^n)_{n \in \mathbb{N}}, (v^n)_{n \in \mathbb{N}}, (\underline{u}^n)_{n \in \mathbb{N}}$  and  $(\bar{u}^n)_{n \in \mathbb{N}}$  converge monotonically pointwise in  $[0, +\infty) \times \mathbb{R}$  to some nonnegative bounded classical solutions  $u, v, \underline{u}$  and  $\bar{u}$  of (2.9)–(2.10) in  $[0, +\infty) \times \mathbb{R}$ , with respective initial conditions  $u_0, v_0, \underline{u}_0$  and  $\bar{u}_0$ . Now, for each  $n \in \mathbb{N}$ , one has

$$\begin{aligned} 0 \leq \underline{u}^n(0, \cdot) = \delta^n \underline{u}_0|_{[-nl, nl]} &\leq \min(\delta^n u_0|_{[-nl, nl]}, \delta^n v_0|_{[-nl, nl]}) \leq \max(\delta^n u_0|_{[-nl, nl]}, \delta^n v_0|_{[-nl, nl]}) \\ &\leq \delta^n \bar{u}_0|_{[-nl, nl]} = \bar{u}^n(0, \cdot) \leq \bar{u}_0|_{[-nl, nl]} \leq \|\bar{u}_0\|_{L^\infty(\mathbb{R})} \end{aligned}$$

in  $[-nl, nl]$  and  $\|\bar{u}^n(0, \cdot) - \underline{u}^n(0, \cdot)\|_{L^\infty([-nl, nl])} \leq 2\|u_0 - v_0\|_{L^\infty(\mathbb{R})}$ . From the previous paragraphs, one has

$$0 \leq \underline{u}^n \leq \min(u^n, v^n) \leq \max(u^n, v^n) \leq \bar{u}^n \leq \bar{K} := \max(K_1, K_2, \|\bar{u}_0\|_{L^\infty(\mathbb{R})}) \quad (3.26)$$

in  $[0, +\infty) \times [-nl, nl]$ . The function  $w^n := \bar{u}^n - \underline{u}^n$  is continuous and nonnegative in  $[0, +\infty) \times [-nl, nl]$  and its restriction to  $(0, +\infty) \times \bar{I}$ , for a patch  $I$  of type  $i \in \{1, 2\}$  included in  $(-nl, nl)$ , is of class  $C_{t;x}^{1;2}((0, +\infty) \times \bar{I})$  and satisfies

$$\frac{\partial w^n}{\partial t} = d_i \frac{\partial^2 w^n}{\partial x^2} + f_i(\bar{u}^n(t, x)) - f_i(\underline{u}^n(t, x)) \leq d_i \frac{\partial^2 w^n}{\partial x^2} + Lw^n \quad \text{in } (0, +\infty) \times I,$$

with  $L := \max(\|f'_1\|_{L^\infty([0, \bar{K}])}, \|f'_2\|_{L^\infty([0, \bar{K}])}) \in [0, +\infty)$ . Furthermore, the function  $w^n$  satisfies the interface conditions (3.2) and it vanishes on  $[0, +\infty) \times \{\pm nl\}$ . On the other hand, the nonnegative  $C^\infty([0, +\infty) \times [-nl, nl])$  function  $(t, x) \mapsto \bar{w}(t, x) := 2\|u_0 - v_0\|_{L^\infty(\mathbb{R})} e^{Lt}$  satisfies the interface conditions (since it is independent of  $x$ ) in (3.2) and  $\frac{\partial \bar{w}}{\partial t} = L\bar{w} = d_i \frac{\partial^2 \bar{w}}{\partial x^2} + L\bar{w}$  in  $[0, +\infty) \times \bar{I}$ , for each patch  $I$  of type  $i \in \{1, 2\}$ . Lastly,

$$0 \leq w^n(0, \cdot) = \bar{u}^n(0, \cdot) - \underline{u}^n(0, \cdot) \leq 2\|u_0 - v_0\|_{L^\infty(\mathbb{R})} = \bar{w}(0, \cdot) \quad \text{in } [-nl, nl].$$

Proposition A.2 then implies that  $0 \leq \bar{u}^n(t, x) - \underline{u}^n(t, x) = w^n(t, x) \leq \bar{w}(t, x)$  for all  $(t, x) \in [0, +\infty) \times [-nl, nl]$ . By passing to the limit as  $n \rightarrow +\infty$  pointwise in  $[0, +\infty) \times \mathbb{R}$ , one gets that  $0 \leq \bar{u}(t, x) - \underline{u}(t, x) \leq 2\|u_0 - v_0\|_{L^\infty(\mathbb{R})} e^{Lt}$ , while  $0 \leq \underline{u} \leq \min(u, v) \leq \max(u, v) \leq \bar{u}$  in  $[0, +\infty) \times \mathbb{R}$  by (3.26). As a conclusion,

$$|u(t, x) - v(t, x)| \leq 2\|u_0 - v_0\|_{L^\infty(\mathbb{R})} e^{Lt} \quad \text{for all } (t, x) \in [0, +\infty) \times \mathbb{R}, \quad (3.27)$$

which yields the Lipschitz continuity of the map  $u_0 \mapsto u$  from  $C^+(\mathbb{R}) \cap L^\infty(\mathbb{R})$  to  $C([0, T] \times \mathbb{R}) \cap L^\infty([0, T] \times \mathbb{R})$  equipped with the sup norms, for each  $T > 0$ . As a conclusion, the proof of Theorem 2.2 is complete.  $\square$

## 4 Existence and uniqueness of a stationary solution

In this section, we focus on the stationary problem (2.14). In Section 4.1, we show Theorem 2.3 on the existence and non-existence of a positive periodic bounded solution of (2.14). Section 4.2 is devoted to the proof of Theorem 2.4 on the uniqueness of such a solution.

### 4.1 Existence of solutions: proof of Theorem 2.3

(i) Assume that (2.11) is fulfilled and that 0 is an unstable solution of (2.14), that is,  $\lambda_1 < 0$ , where  $\lambda_1$  is the principal eigenvalue of the eigenvalue problem (2.13), associated with the principal eigenfunction  $\phi$ . Since  $f(x, \cdot)|_I = f_i$  is of class  $C^1(\mathbb{R})$  for each  $x \in \mathbb{R} \setminus S$  belonging to a patch  $I$  of type  $i \in \{1, 2\}$ , there exists  $\kappa_0 > 0$  small enough such that, for all  $0 < \kappa \leq \kappa_0$ ,

$$f(x, \kappa\phi(x)) \geq \kappa\phi(x)f_s(x, 0) + \frac{\lambda_1}{2}\kappa\phi(x) \quad \text{for all } x \in \mathbb{R} \setminus S,$$

hence  $\kappa\phi$  then satisfies

$$-d(x)\kappa\phi''(x) - f(x, \kappa\phi(x)) \leq -d(x)\kappa\phi''(x) - \kappa\phi(x)f_s(x, 0) - \frac{\lambda_1}{2}\kappa\phi(x) = \frac{\lambda_1}{2}\kappa\phi(x) < 0 \quad (4.1)$$

for all  $x \in \mathbb{R} \setminus S$ , as well as the interface conditions in (2.14). With  $M > 0$  as in (2.11), we can then fix  $\kappa \in (0, \kappa_0]$  so that  $\kappa\phi \leq M$  in  $\mathbb{R}$ . Now, for each  $n \in \mathbb{N}$ , let  $u^n$  be the unique bounded classical solution of (3.1)–(3.4) with initial condition  $u^n(0, \cdot) = M\delta^n|_{[-nl, nl]}$ , with the cut-off function  $\delta^n$  given in (3.20). From the proof of Theorem 2.2, the sequence  $(u^n)_{n \in \mathbb{N}}$  converges monotonically pointwise in  $[0, +\infty) \times \mathbb{R}$  to a nonnegative bounded classical solution  $u$  of (2.9)–(2.10) in  $[0, +\infty) \times \mathbb{R}$ , with initial condition  $M$ , and  $u(t, x) = u(t, x+l)$  for all  $(t, x) \in [0, +\infty) \times \mathbb{R}$ .

Furthermore, by (2.11), the constant  $M$  is a supersolution of (3.1)–(3.4) in  $[0, +\infty) \times [-nl, nl]$  for each  $n \in \mathbb{N}$ , in the sense of Definition A.1. Proposition A.2 implies that

$$u^n(t, x) \leq M \quad \text{for all } t \geq 0 \text{ and } x \in [-nl, nl].$$

In particular, for each  $h \geq 0$  and  $n \in \mathbb{N}$ , one has  $u^n(h, x) \leq M = u^{n+1}(0, x)$  for all  $x \in [-nl, nl]$ , together with  $u^n(t + h, \pm nl) = 0 \leq u^{n+1}(t, \pm nl)$  for all  $t \geq 0$ . Hence  $u^n(t + h, x) \leq u^{n+1}(t, x)$  for all  $t \geq 0$  and  $x \in [-nl, nl]$ , by Proposition A.2 again. Therefore,

$$u(t + h, x) \leq u(t, x) \quad \text{for all } (t, x) \in [0, +\infty) \times \mathbb{R},$$

by passing to the limit as  $n \rightarrow +\infty$ . In other words, the nonnegative continuous function  $u$  is non-increasing in  $t$ , and, together with the periodicity in space and the Schauder estimates of Theorem 2.2, there is a continuous periodic solution  $p : \mathbb{R} \rightarrow [0, M]$  of (2.14) such that  $u(t, \cdot) \rightarrow p$  uniformly in  $\mathbb{R}$  as  $t \rightarrow +\infty$ , and  $u(t, \cdot)|_{\bar{I}} \rightarrow p|_{\bar{I}}$  in  $C^2(\bar{I})$  for each patch  $I \subset \mathbb{R}$ . Finally, since the periodic continuous function  $\kappa\phi$  has restrictions of class  $C^2(\bar{I})$  for each patch  $I \subset \mathbb{R}$  and satisfies (4.1) and the interface conditions in (2.14) (it is a subsolution of this problem), and since  $\kappa\phi \leq M = u(0, \cdot)$  in  $\mathbb{R}$ , a similar comparison result as in Proposition A.2 (but adapted here to the case of spatially periodic super- and subsolutions) yields  $\kappa\phi(x) \leq u(t, x)$  for all  $t \geq 0$  and  $x \in \mathbb{R}$ , hence  $\kappa\phi(x) \leq p(x)$  for all  $x \in \mathbb{R}$  at the limit  $t \rightarrow +\infty$ . As a conclusion, there exists a positive and periodic continuous solution  $p$  of (2.14) satisfying  $\kappa\phi \leq p \leq M$  in  $\mathbb{R}$ , that is, Theorem 2.3 (i) is proved.

(ii) Next, in addition to (2.11), we assume that (2.12) holds, that  $p$  is a nonnegative bounded continuous solution to the elliptic problem (2.14), and that 0 is a stable solution of (2.14), that is,  $\lambda_1 \geq 0$ . Let  $\phi$  be the unique positive solution of (2.13). In (2.12), let us assume that  $s \mapsto f_1(s)/s$  is decreasing with respect to  $s > 0$  (the case when  $s \mapsto f_2(s)/s$  is decreasing with respect to  $s > 0$  can be handled similarly). We infer that, for every  $\gamma > 0$ ,

$$f(x, \gamma\phi(x)) = f_1(\gamma\phi(x)) < f'_1(0)\gamma\phi(x) = f_s(x, 0)\gamma\phi(x) \quad \text{for all } x \in (nl - l_1, nl) \text{ and } n \in \mathbb{Z},$$

while

$$f(x, \gamma\phi(x)) = f_2(\gamma\phi(x)) \leq f'_2(0)\gamma\phi(x) = f_s(x, 0)\gamma\phi(x) \quad \text{for all } x \in (nl, nl + l_2) \text{ and } n \in \mathbb{Z}.$$

Hence, for all  $\gamma > 0$ ,

$$\begin{cases} -d_1\gamma\phi''(x) - f_1(\gamma\phi(x)) > -d_1\gamma\phi''(x) - f'_1(0)\gamma\phi(x) = \lambda_1\gamma\phi(x) \geq 0, & x \in (nl - l_1, nl), \\ -d_2\gamma\phi''(x) - f_2(\gamma\phi(x)) \geq -d_2\gamma\phi''(x) - f'_2(0)\gamma\phi(x) = \lambda_1\gamma\phi(x) \geq 0, & x \in (nl, nl + l_2). \end{cases} \quad (4.2)$$

Since  $\phi$  is bounded from below by a positive constant (because it is positive, periodic and continuous), and since  $p$  is bounded, one can define

$$\gamma^* = \inf \{ \gamma > 0, \gamma\phi > p \text{ in } \mathbb{R} \} \in [0, +\infty).$$

Our goal is to show that  $\gamma^* = 0$ . Assume by way of contradiction that  $\gamma^* > 0$ , and set  $z := \gamma^*\phi - p$ . Then  $z \geq 0$  in  $\mathbb{R}$  and there exists a sequence  $(x_m)_{m \in \mathbb{N}}$  in  $\mathbb{R}$  such that  $z(x_m) \rightarrow 0$  as  $m \rightarrow +\infty$ . Moreover,  $z$  satisfies

$$\begin{cases} -d_1 z''(x) - b(x)z(x) > 0, & x \in (nl - l_1, nl), \\ -d_2 z''(x) - b(x)z(x) \geq 0, & x \in (nl, nl + l_2), \\ z(x^-) = z(x^+), \quad z'(x^-) = \sigma z'(x^+), & x = nl, \\ z(x^-) = z(x^+), \quad \sigma z'(x^-) = z'(x^+), & x = nl + l_2, \end{cases} \quad (4.3)$$

for some bounded function  $b$  defined in  $\mathbb{R} \setminus S$ .

Assume at first that up to a subsequence,  $x_m \rightarrow \bar{x} \in \mathbb{R}$  as  $m \rightarrow +\infty$ . By continuity of  $\phi$  and  $p$ , one has  $z(\bar{x}) = 0$ . We distinguish two cases. Assume first that  $\bar{x} \in \mathbb{R} \setminus S$ . It is easily seen from the strong elliptic maximum principle and the Hopf lemma, applied by induction from one patch to an adjacent one, that  $z \equiv 0$  in  $\mathbb{R}$ . This is a contradiction with the strict inequality in the first line of (4.3). Thus,  $z > 0$  in  $\mathbb{R} \setminus S$  and  $\bar{x} \in S$ , hence the Hopf lemma yields  $z'(\bar{x}^-) < 0$  and  $z'(\bar{x}^+) > 0$ , contradicting the interface condition in (4.3).

In the general case, let  $\bar{x}_m \in (-l_1, l_2]$  be such that  $x_m - \bar{x}_m \in l\mathbb{Z}$ . Then up to some subsequence, one can assume that there is  $\bar{x}_\infty \in [-l_1, l_2]$  such that  $\bar{x}_m \rightarrow \bar{x}_\infty$  as  $m \rightarrow +\infty$ . Set  $z_m = \gamma^* \phi_m - p_m = \gamma^* \phi - p_m$ , where  $\phi_m(x) := \phi(x + x_m - \bar{x}_m) = \phi(x)$  and  $p_m(x) := p(x + x_m - \bar{x}_m)$ . Since  $d(x)$  and  $f(x, \cdot)$  are periodic in  $x$ , one then infers from (2.14) and (4.2) that each function  $z_m$  satisfies

$$\begin{cases} -d_1 z_m''(x) - f_1(\gamma^* \phi(x)) + f_1(p_m(x)) > 0, & x \in (nl - l_1, nl), \\ -d_2 z_m''(x) - f_2(\gamma^* \phi(x)) + f_2(p_m(x)) \geq 0, & x \in (nl, nl + l_2), \\ z_m(x^-) = z_m(x^+), \quad z_m'(x^-) = \sigma z_m'(x^+), & x = nl, \\ z_m(x^-) = z_m(x^+), \quad \sigma z_m'(x^-) = z_m'(x^+), & x = nl + l_2. \end{cases}$$

From standard elliptic estimates, it follows that up to some subsequence, the sequences  $(p_m)_{m \in \mathbb{N}}$  and  $(z_m)_{m \in \mathbb{N}}$  converge as  $m \rightarrow +\infty$  to some functions  $p_\infty$  and  $z_\infty$  locally uniformly in  $\mathbb{R}$ , and in  $C^2(\bar{I})$  for each patch  $I \subset \mathbb{R}$ , with  $z_\infty = \gamma^* \phi - p_\infty$  and

$$\begin{cases} -d_1 z_\infty''(x) - f_1(\gamma^* \phi(x)) + f_1(p_\infty(x)) \geq 0, & x \in (nl - l_1, nl), \\ -d_2 z_\infty''(x) - f_2(\gamma^* \phi(x)) + f_2(p_\infty(x)) \geq 0, & x \in (nl, nl + l_2), \\ z_\infty(x^-) = z_\infty(x^+), \quad z_\infty'(x^-) = \sigma z_\infty'(x^+), & x = nl, \\ z_\infty(x^-) = z_\infty(x^+), \quad \sigma z_\infty'(x^-) = z_\infty'(x^+), & x = nl + l_2. \end{cases} \quad (4.4)$$

Moreover,  $z_\infty \geq 0$  in  $\mathbb{R}$ ,  $z_\infty(\bar{x}_\infty) = 0$ , and the first inequality in (4.4) is actually strict from the strict sign in the first line of (4.2) applied with  $\gamma^* > 0$ , and from the periodicity of  $\phi$ . From similar lines as above, one reaches a contradiction by using the strong elliptic maximum principle and the Hopf lemma together with the interface conditions in (4.4).

Consequently,  $\gamma^* = 0$ , whence  $p \equiv 0$ . This completes the proof of Theorem 2.3.  $\square$

## 4.2 Uniqueness of solutions: proof of Theorem 2.4

In order to prove the uniqueness of the positive stationary solution, we show the following crucial property.

**Proposition 4.1.** *Assume (2.11) and that 0 is an unstable solution of (2.14) (i.e.,  $\lambda_1 < 0$ ). Let  $p$  be a bounded nonnegative continuous solution of the stationary problem (2.14). Then, either  $p \equiv 0$  in  $\mathbb{R}$ , or  $\inf_{\mathbb{R}} p > 0$ .*

Note that Proposition 4.1 also holds in particular in the class of periodic solutions. However, we look in Theorem 2.4 at the uniqueness within a more general class of functions which are not assumed to be *a priori* periodic. Proposition 4.1, which implies that any positive solution of (2.14) is in fact bounded from below by a positive constant, will be the essence in proving uniqueness under the additional assumption (2.12).

We prove Proposition 4.1 via a series of lemmas. First of all, for any  $R > 0$  and  $y \in \mathbb{R}$ , we claim that there exists a unique real number (principal eigenvalue)  $\lambda_R^y$  and a unique nonnegative

continuous and piecewise smooth function (principal eigenfunction)  $\varphi_R^y$  in  $[-R, R]$  satisfying

$$\begin{cases} -d(x+y)(\varphi_R^y)''(x) - f_s(x+y, 0)\varphi_R^y(x) = \lambda_R^y \varphi_R^y(x), & x \in (-R, R) \setminus (S-y), \\ \varphi_R^y(x^-) = \varphi_R^y(x^+), \quad (\varphi_R^y)'(x^-) = \sigma(\varphi_R^y)'(x^+), & x = nl - y \in (-R, R), \\ \varphi_R^y(x^-) = \varphi_R^y(x^+), \quad \sigma(\varphi_R^y)'(x^-) = (\varphi_R^y)'(x^+), & x = nl + l_2 - y \in (-R, R), \\ \varphi_R^y > 0 \text{ in } (-R, R), \quad \varphi_R^y(\pm R) = 0, \quad \|\varphi_R^y\|_{L^\infty(-R, R)} = 1. \end{cases} \quad (4.5)$$

We sketch the proof below. For convenience, we denote by  $J_s$  and  $K_r$  the finitely many shifted (by  $-y$ ) patches of type 1 and of type 2 in  $(-R, R)$  so that

$$(-R, R) \setminus (S-y) = \left( \bigcup_s J_s \right) \cup \left( \bigcup_r K_r \right).$$

The functions  $d(\cdot + y)$  and  $f_s(\cdot + y, 0)$  are now constant in each patch  $J_s$  or  $K_r$ . Consider the Hilbert space  $H = H_0^1(-R, R)$  and the Banach space

$$G = \{u \in C([-R, R]) : u|_{\overline{J_s}} \in C^1(\overline{J_s}), u|_{\overline{K_r}} \in C^1(\overline{K_r}), u(\pm R) = 0\},$$

equipped with the norms

$$\begin{cases} \|u\|_H &= \sqrt{\sum_s \|u\|_{H^1(J_s)}^2 + \sum_r \frac{1}{k} \|u\|_{H^1(K_r)}^2} \\ &= \sqrt{\sum_s \int_{J_s} (|u'|^2 + u^2) + \sum_r \frac{1}{k} \int_{K_r} (|u'|^2 + u^2)} \geq \min\left(1, \frac{1}{\sqrt{k}}\right) \|u\|_{H^1(-R, R)}, \\ \|u\|_G &= \sum_s \|u\|_{C^1(\overline{J_s})} + \sum_r \frac{1}{k} \|u\|_{C^1(\overline{K_r})}. \end{cases}$$

Set

$$\Lambda := \max(f_1'(0), f_2'(0)) + 1. \quad (4.6)$$

For  $g \in G$ , let us solve the following problem

$$\begin{cases} -d(\cdot + y)u'' - f_s(\cdot + y, 0)u + \Lambda u = g, & \text{in } (-R, R) \setminus (S-y), \\ u(x^-) = u(x^+), \quad u'(x^-) = \sigma u'(x^+), & x = nl - y \in (-R, R), \\ u(x^-) = u(x^+), \quad \sigma u'(x^-) = u'(x^+), & x = nl + l_2 - y \in (-R, R), \\ u(\pm R) = 0, \end{cases} \quad (4.7)$$

first in the weak sense: that is, we look for a solution  $u \in H$  such that

$$B(u, z) = \langle g, z \rangle \quad \text{for all } z \in H, \quad (4.8)$$

where the bilinear form  $B$  is defined by

$$B(u, z) = \sum_s \int_{J_s} (d_1 u' z' + (\Lambda - f_1'(0))uz) + \sum_r \frac{1}{k} \int_{K_r} (d_2 u' z' + (\Lambda - f_2'(0))uz), \quad (4.9)$$

and

$$\langle g, z \rangle = \sum_s \int_{J_s} gz + \sum_r \frac{1}{k} \int_{K_r} gz.$$

Clearly, the map  $z \mapsto \langle g, z \rangle$  is continuous in  $H$ , and  $B$  is continuous in  $H \times H$ . Moreover, it is easily seen that, for any  $u \in H$ ,

$$\begin{aligned} B(u, u) &= \sum_s \int_{J_s} (d_1(u')^2 + (\Lambda - f'_1(0))u^2) + \sum_r \frac{1}{k} \int_{K_r} (d_2(u')^2 + (\Lambda - f'_2(0))u^2) \\ &\geq \min(d_1, d_2, 1) \|u\|_H^2, \end{aligned}$$

whence  $B$  is coercive. The Lax-Milgram theorem yields the existence of a unique  $u \in H$  (hence,  $u$  is continuous in  $[-R, R]$ , by identifying  $u$  with its unique continuous representative, and  $u(\pm R) = 0$ ) satisfying (4.8), and  $\|u\|_{H^1(-R, R)} \leq C_1 \|g\|_{L^2(-R, R)}$  for a positive constant  $C_1$  only depending on  $d_{1,2}$  and  $k$ . Thus,

$$\max_{[-R, R]} |u| \leq C_2 \|g\|_{L^2(-R, R)} \leq C_3 \|g\|_G$$

for some positive constants  $C_2$  and  $C_3$  only depending on  $d_{1,2}$ ,  $k$ , and  $R$ . Furthermore, owing to the definitions of  $B$  in (4.9) and of  $\sigma$  in (2.8), the function  $u$  has restrictions in  $\mathcal{I}$  belonging to  $H^2(\mathcal{I})$  for each patch  $\mathcal{I}$  of the type  $J_s$  or  $K_r$  in  $(-R, R) \setminus (S - y)$  and  $u$  satisfies the equations and the interface conditions in (4.7). In particular,  $-d_i u'' + (\Lambda - f'_i(0))u = g$  in  $L^2(\mathcal{I})$  for each patch  $\mathcal{I}$  of the type  $J_s$  or  $K_r$  in  $(-R, R) \setminus (S - y)$ , hence  $\max_{\mathcal{I} \in (\cup_s J_s) \cup (\cup_r K_r)} \|u|_{\mathcal{I}}''\|_{L^2(\mathcal{I})} \leq C_4 \|g\|_{L^2(-R, R)}$  for a positive constant  $C_4$  only depending on  $d_{1,2}$ ,  $k$  and  $f'_{1,2}(0)$ , and then  $u|_{\overline{\mathcal{I}}} \in C^2(\overline{\mathcal{I}})$  and

$$\max_{\mathcal{I} \in (\cup_s J_s) \cup (\cup_r K_r)} \left( \max_{\overline{\mathcal{I}}} |u|_{\mathcal{I}}'' \right) \leq C_5 \|g\|_G$$

for a positive constant  $C_5$  only depending on  $d_{1,2}$ ,  $k$ ,  $R$  and  $f'_{1,2}(0)$ . Notice in particular that  $u$  then belongs to  $G$ . Using again the equations satisfied by  $u$  and the fact that  $u$  necessarily has an interior critical (with vanishing derivative) point in  $(-R, R)$  thanks to (4.7), it follows that  $\max_{\mathcal{I} \in (\cup_s J_s) \cup (\cup_r K_r)} \left( \max_{\overline{\mathcal{I}}} |u|_{\mathcal{I}}' \right) \leq C_6 \|g\|_{L^2(-R, R)} \leq C_7 \|g\|_G$  for some positive constants  $C_6$  and  $C_7$  only depending on  $d_{1,2}$ ,  $k$ ,  $R$  and  $f'_{1,2}(0)$ , and finally that  $u|_{\overline{\mathcal{I}}} \in C^3(\overline{\mathcal{I}})$  for each patch  $\mathcal{I}$  of the type  $J_s$  or  $K_r$  in  $(-R, R) \setminus (S - y)$  and

$$\max_{\mathcal{I} \in (\cup_s J_s) \cup (\cup_r K_r)} \|u|_{\mathcal{I}}\|_{C^3(\overline{\mathcal{I}})} \leq C_8 \|g\|_G$$

for a positive constant  $C_8$  only depending on  $d_{1,2}$ ,  $k$ ,  $R$  and  $f'_{1,2}(0)$

The mapping  $T: g \in G \mapsto Tg := u \in G$  is obviously linear. The previous estimates and the Arzelà-Ascoli theorem yield the compactness of  $T$ . Let now  $\mathcal{K}$  be the cone  $\mathcal{K} = \{u \in G : u \geq 0 \text{ in } [-R, R]\}$ . Its interior  $\mathring{\mathcal{K}}$  is not empty, and  $\mathcal{K} \cap (-\mathcal{K}) = \{0\}$ . We claim that, if  $g \in \mathcal{K} \setminus \{0\}$ , then  $u \in \mathring{\mathcal{K}}$ . Indeed, by using  $z := u^- = \max(-u, 0) \in H$  in (4.8) one has

$$-\sum_s \int_{J_s} (d_1((u^-)')^2 + (\Lambda - f'_1(0))(u^-)^2) - \sum_r \frac{1}{k} \int_{K_r} (d_2((u^-)')^2 + (\Lambda - f'_2(0))(u^-)^2) = \langle g, u^- \rangle \geq 0,$$

hence  $u^- \equiv 0$ , that is,  $u \geq 0$  in  $[-R, R]$ . By finitely many applications of the strong elliptic maximum principle and the Hopf lemma, together with the fact that  $g \not\equiv 0$ , one concludes that  $u > 0$  in  $(-R, R)$  and that  $u'(-R) > 0$  and  $u'(R) < 0$ . Therefore,  $T(\mathcal{K} \setminus \{0\}) \subset \mathring{\mathcal{K}}$ . From the Krein-Rutman theory, there exists a unique positive real number  $\tilde{\lambda}_R^y$  and a unique function  $\varphi_R^y \in \mathring{K}$  such that  $\tilde{\lambda}_R^y T\varphi_R^y = \varphi_R^y$  and, say,  $\|\varphi_R^y\|_{L^\infty(-R, R)} = 1$ . Hence,  $\varphi_R^y$  is continuous in  $[-R, R]$ , of class  $C^3(\overline{\mathcal{I}})$  (and then  $C^\infty(\overline{\mathcal{I}})$  by induction) for each patch  $\mathcal{I}$  of the type  $J_s$  or  $K_s$  in  $(-R, R) \cap (S - y)$ , and

$$\begin{cases} -d(x+y)(\varphi_R^y)''(x) - f_s(x+y, 0)\varphi_R^y(x) + \Lambda\varphi_R^y(x) = \tilde{\lambda}_R^y\varphi_R^y(x), & x \in (-R, R) \setminus (S - y), \\ \varphi_R^y(x^-) = \varphi_R^y(x^+), \quad (\varphi_R^y)'(x^-) = \sigma(\varphi_R^y)'(x^+), & x = nl - y \in (-R, R), \\ \varphi_R^y(x^-) = \varphi_R^y(x^+), \quad \sigma(\varphi_R^y)'(x^-) = (\varphi_R^y)'(x^+), & x = nl + l_2 - y \in (-R, R), \\ \varphi_R^y > 0 \text{ in } (-R, R), \quad \varphi_R^y(\pm R) = 0, \quad \|\varphi_R^y\|_{L^\infty(-R, R)} = 1. \end{cases}$$



Therefore,  $\lambda_R^y := \tilde{\lambda}_R^y - \Lambda > -\Lambda = -\max(f_1'(0), f_2'(0)) - 1$  is the first eigenvalue of (4.5) associated with a unique continuous function  $\varphi_R^y$  in  $[-R, R]$  that is positive in  $(-R, R)$  and of class  $C^\infty(\bar{\mathcal{I}})$  for each patch  $\mathcal{I}$  of the type  $J_s$  or  $K_s$  in  $(-R, R) \cap (S - y)$ . Furthermore, for each  $R > 0$ , the interval  $(-R, R)$  contains at least a patch  $\mathcal{I}$  of the type  $J_s$  or  $K_s$  of length larger than or equal to  $\ell_R := \min(l_1, l_2, R)$ , hence  $\lambda_R^y \leq \max(-f_1'(0) + d_1\pi^2/\ell_R^2, -f_2'(0) + d_2\pi^2/\ell_R^2)$  from the positivity of  $\varphi_R^y$  in  $\mathcal{I}$ . As a consequence, for each  $R > 0$ , there is a constant  $N_R$  such that

$$|\tilde{\lambda}_{R'}^y| + |\lambda_{R'}^y| \leq N_R \text{ for every } R' \geq R \text{ and } y \in \mathbb{R}.$$

Since both  $\lambda_R^y$  and  $\varphi_R^y$  are unique, the aforementioned estimates and compactness arguments imply that, for each  $R > 0$ , the maps  $y \mapsto \lambda_R^y$  and  $y \mapsto \varphi_R^y$  are continuous in  $\mathbb{R}$  (the continuity of  $y \mapsto \varphi_R^y$  is understood in the sense of the uniform topology in  $[-R, R]$ ). Note also that since  $d$  and  $f$  are periodic in  $x$ , it follows that  $\lambda_R^y$  and  $\varphi_R^y$  are periodic with respect to  $y$  as well.

Similarly, for each  $y \in \mathbb{R}$ , there exists a unique principal eigenvalue  $\lambda^y$  and a unique principal eigenfunction  $\phi^y$  of the periodic problem

$$\begin{cases} -d(x+y)(\phi^y)''(x) - f_s(x+y, 0)\phi^y(x) = \lambda^y\phi^y(x), & x \in \mathbb{R} \setminus (S-y), \\ \phi^y(x^-) = \phi^y(x^+), \quad (\phi^y)'(x^-) = \sigma(\phi^y)'(x^+), & x = nl - y, \\ \phi^y(x^-) = \phi^y(x^+), \quad \sigma(\phi^y)'(x^-) = (\phi^y)'(x^+), & x = nl + l_2 - y, \\ \phi^y \text{ is periodic, } \phi^y > 0 \text{ in } \mathbb{R}, \quad \|\phi^y\|_{L^\infty(\mathbb{R})} = 1, \end{cases} \quad (4.10)$$

where  $\phi^y$  is continuous in  $\mathbb{R}$  and  $\phi^y|_{\bar{\mathcal{I}}}$  is of class  $C^\infty(\bar{\mathcal{I}})$  for each shifted patch  $\mathcal{I} \subset \mathbb{R} \setminus (S - y)$ . First of all, it is straightforward to observe:

**Lemma 4.2.** *The principal eigenvalue  $\lambda^y$  of (4.10) does not depend on  $y$ , that is,  $\lambda^y = \lambda^0 = \lambda_1$  for all  $y \in \mathbb{R}$ , where  $\lambda_1$  is the principal eigenvalue of the eigenvalue problem (2.13).*

*Proof.* Setting  $\phi(x) := \phi^y(x - y)$  for  $x \in \mathbb{R}$ , the function  $\phi$  satisfies

$$\begin{cases} -d(x)\phi''(x) - f_s(x, 0)\phi(x) = \lambda^y\phi(x), & x \in \mathbb{R} \setminus S, \\ \phi(x^-) = \phi(x^+), \quad \phi'(x^-) = \sigma\phi'(x^+), & x = nl, \\ \phi(x^-) = \phi(x^+), \quad \sigma\phi'(x^-) = \phi'(x^+), & x = nl + l_2, \\ \phi \text{ is periodic, } \phi > 0 \text{ in } \mathbb{R}, \quad \|\phi\|_{L^\infty(\mathbb{R})} = 1. \end{cases}$$

By uniqueness of the principal eigenvalue, one then has  $\lambda^y = \lambda^0 = \lambda_1$ .  $\square$

The second lemma provides a comparison between  $\lambda_R^y$  and  $\lambda_1$ .

**Lemma 4.3.** *For all  $y \in \mathbb{R}$  and  $R > 0$ , one has  $\lambda_R^y > \lambda_1$ .*

*Proof.* Fix any  $y \in \mathbb{R}$  and  $R > 0$ , and assume by way of contradiction that  $\lambda_R^y \leq \lambda_1$ . Notice that the continuous function  $\varphi_R^y$  satisfies

$$\begin{cases} -d(x+y)(\varphi_R^y)'' - f_s(x+y, 0)\varphi_R^y - \lambda_1\varphi_R^y = (\lambda_R^y - \lambda_1)\varphi_R^y, & \text{in } (-R, R) \setminus (S-y), \\ \varphi_R^y(x^-) = \varphi_R^y(x^+), \quad (\varphi_R^y)'(x^-) = \sigma(\varphi_R^y)'(x^+), & x = nl - y \in (-R, R), \\ \varphi_R^y(x^-) = \varphi_R^y(x^+), \quad \sigma(\varphi_R^y)'(x^-) = (\varphi_R^y)'(x^+), & x = nl + l_2 - y \in (-R, R), \\ \varphi_R^y > 0 \text{ in } (-R, R), \quad \varphi_R^y(\pm R) = 0, \quad \|\varphi_R^y\|_{L^\infty(-R, R)} = 1, \end{cases} \quad (4.11)$$

while the continuous solution  $\phi^y = \phi(\cdot + y)$  of (4.10) satisfies

$$\begin{cases} -d(x+y)(\phi^y)'' - f_s(x+y, 0)\phi^y - \lambda_1\phi^y = 0, & \text{in } (-R, R) \setminus (S-y), \\ \phi^y(x^-) = \phi^y(x^+), \quad (\phi^y)'(x^-) = \sigma(\phi^y)'(x^+), & x = nl - y \in (-R, R), \\ \phi^y(x^-) = \phi^y(x^+), \quad \sigma(\phi^y)'(x^-) = (\phi^y)'(x^+), & x = nl + l_2 - y \in (-R, R), \\ \phi^y > 0, & \text{in } [-R, R], \end{cases}$$

where  $\phi$  is the principal eigenfunction of (2.13). Therefore,  $\phi^y > \kappa \varphi_R^y$  in  $[-R, R]$  for all  $\kappa > 0$  small enough. Define

$$\kappa^* = \sup \{ \kappa > 0 : \phi^y > \kappa \varphi_R^y \text{ in } [-R, R] \} \in (0, +\infty).$$

By continuity,  $\phi^y \geq \kappa^* \varphi_R^y$  in  $[-R, R]$  and there exists  $x_0 \in [-R, R]$  such that  $\phi^y(x_0) = \kappa^* \varphi_R^y(x_0)$ . But since  $\phi^y > 0$  in  $[-R, R]$  and  $\varphi_R^y(\pm R) = 0$ , one infers that  $x_0 \in (-R, R)$ . On the other hand, the function  $\kappa^* \varphi_R^y$  satisfies (4.11), hence

$$-d(x+y)(\kappa^* \varphi_R^y)'' - f_s(x+y, 0)\kappa^* \varphi_R^y - \lambda_1 \kappa^* \varphi_R^y \leq 0 \text{ in } (-R, R) \setminus (S-y),$$

thanks to the assumption  $\lambda_R^y \leq \lambda_1$ . It then follows from finitely many applications of the strong elliptic maximum principle and the Hopf lemma that  $\kappa^* \varphi_R^y \equiv \phi^y$  in  $(-R, R)$  and then in  $[-R, R]$  by continuity, a contradiction with the boundary conditions at  $x = \pm R$ . Consequently,  $\lambda_R^y > \lambda_1$  for all  $y \in \mathbb{R}$  and  $R > 0$ .  $\square$

For any two positive real numbers  $R_1 < R_2$ , by replacing  $\lambda_R^y$  with  $\lambda_{R_1}^y$  and  $\lambda_1$  with  $\lambda_{R_2}^y$  in the above proof, and by noticing that  $\varphi_{R_2}^y > 0$  in  $[-R_1, R_1]$ , a similar argument as that of Lemma 4.3 implies that  $\lambda_{R_1}^y > \lambda_{R_2}^y$ . That is,

**Lemma 4.4.** *For all  $y \in \mathbb{R}$ , the function  $R \mapsto \lambda_R^y$  is decreasing in  $R > 0$ .*

The last result before the proof of Proposition 4.1 is the following convergence result.

**Lemma 4.5.** *One has  $\lim_{R \rightarrow +\infty} \lambda_R^y = \lambda_1$  uniformly in  $y \in \mathbb{R}$ .*

*Proof.* First of all, from Lemma 4.4 and the periodicity and continuity with respect to  $y$ , it is sufficient to show that  $\lambda_R^y \rightarrow \lambda_1$  as  $R \rightarrow +\infty$  for each  $y \in \mathbb{R}$ , from Dini's theorem. So let us fix  $y \in \mathbb{R}$  in the proof. For  $R > 0$ , consider the elliptic operator  $\mathcal{L}_R^y u := -d(x+y)u'' - f_s(x+y, 0)u$  with domain

$$E_R = \{ \psi \in H_0^1(-R, R) : \psi|_{\mathcal{I}} \in H^2(\mathcal{I}) \text{ for all } \mathcal{I}, \text{ and } \psi \text{ satisfies the interface conditions in (4.5)} \},$$

where  $\mathcal{I}$  is any patch of the type  $J_s$  or  $K_r$  in  $(-R, R) \setminus (S-y)$ . Note in particular that the principal eigenfunction  $\varphi_R^y$  of (4.5) belongs to  $E_R \setminus \{0\}$ . Owing to (2.8), the operator  $\mathcal{L}_R^y$  is symmetric with respect to the inner product

$$\langle u, v \rangle_R = \sum_s \int_{J_s} uv + \sum_r \frac{1}{k} \int_{K_r} uv, \quad u, v \in L^2(-R, R).$$

Therefore, one has the following variational formula for  $\lambda_R^y$ :

$$\lambda_R^y = \min_{\psi \in E_R \setminus \{0\}} \frac{B(\psi, \psi) - \Lambda \langle \psi, \psi \rangle_R}{\langle \psi, \psi \rangle_R} = \min_{\psi \in E_R \setminus \{0\}} \frac{\langle \mathcal{L}_R^y \psi, \psi \rangle_R}{\langle \psi, \psi \rangle_R}, \quad (4.12)$$

with  $\Lambda$  and  $B$  as in (4.6) and (4.9). It is easy to see that one can choose a family of  $C^2(\mathbb{R})$  functions  $(\chi_R^y)_{R \geq 2}$ , such that  $\sup_{R \geq 2} \|\chi_R^y\|_{C^2(\mathbb{R})} < +\infty$  and, for each  $R \geq 2$ ,

$$\begin{cases} \chi_R^y(x) = 1 & \text{for all } x \in [-R+1, R-1], \\ \chi_R^y(x) = 0 & \text{for all } x \in (-\infty, -R] \cup [R, +\infty), \\ (\chi_R^y)'(x) = 0 & \text{for all } x \in S-y, \\ 0 \leq \chi_R^y \leq 1 & \text{in } \mathbb{R}. \end{cases}$$

Let  $\phi^y$  be the solution of (4.10). The function  $\psi_R^y := \phi^y \chi_R^y$  belongs to  $E_R \setminus \{0\}$  and

$$\frac{\langle \mathcal{L}_R^y \psi_R^y, \psi_R^y \rangle_R}{\langle \psi_R^y, \psi_R^y \rangle_R} = \frac{\langle \mathcal{L}_R^y \phi^y, \phi^y \rangle_{R-1} + D_R}{\langle \psi_R^y, \psi_R^y \rangle_R} = \frac{\lambda_1 \langle \phi^y, \phi^y \rangle_{R-1} + D_R}{\langle \psi_R^y, \psi_R^y \rangle_R}$$

from Lemma 4.2, where

$$\begin{aligned} D_R &= \langle \mathcal{L}_R^y \psi_R^y, \psi_R^y \rangle_R - \langle \mathcal{L}_R^y \phi^y, \phi^y \rangle_{R-1} \\ &= \sum_s \int_{\tilde{J}_s} -d(\cdot + y)(\phi^y \chi_R^y)''(\phi^y \chi_R^y) - f_s(\cdot + y, 0)(\phi^y \chi_R^y)^2 \\ &\quad + \sum_r \frac{1}{k} \int_{\tilde{K}_r} -d(\cdot + y)(\phi^y \chi_R^y)''(\phi^y \chi_R^y) - f_s(\cdot + y, 0)(\phi^y \chi_R^y)^2, \end{aligned}$$

and  $\tilde{J}_s$  and  $\tilde{K}_r$  stand for the patches of type 1 and 2 in  $((-R, -R+1) \cup (R-1, R)) \setminus (S-y)$ , that is,  $((-R, -R+1) \cup (R-1, R)) \setminus (S-y) = (\cup_s \tilde{J}_s) \cup (\cup_r \tilde{K}_r)$ . Since  $\phi^y$  is periodic and  $\phi^y|_{\bar{\mathcal{I}}} \in C^2(\bar{\mathcal{I}})$  (and even  $C^\infty(\bar{\mathcal{I}})$ ) for each patch  $\mathcal{I} \subset \mathbb{R} \setminus (S-y)$ , the quantities  $\|\phi^y|_{\bar{\mathcal{I}}}\|_{C^2(\bar{\mathcal{I}})}$  are bounded independently of  $\mathcal{I}$ . Since  $\sup_{R \geq 2} \|\chi_R^y\|_{C^2(\mathbb{R})} < +\infty$ , it follows that there exists  $C > 0$  such that  $|D_R| \leq C$  for all  $R \geq 2$ . Likewise, one also has

$$\sup_{R \geq 2} |\langle \psi_R^y, \psi_R^y \rangle_R - \langle \phi^y, \phi^y \rangle_{R-1}| < +\infty.$$

On the other hand, since  $\phi^y$  is continuous, positive and periodic, there exists  $\delta > 0$  such that  $\phi^y \geq \delta > 0$  in  $\mathbb{R}$ , hence  $\langle \phi^y, \phi^y \rangle_{R-1} \geq 2 \min(1, 1/k) \delta^2 (R-1)$  and

$$\frac{\langle \psi_R^y, \psi_R^y \rangle_R}{\langle \phi^y, \phi^y \rangle_{R-1}} \rightarrow 1 \quad \text{as } R \rightarrow +\infty.$$

Using the estimates above, one gets that

$$\frac{\langle \mathcal{L}_R^y \psi_R^y, \psi_R^y \rangle_R}{\langle \psi_R^y, \psi_R^y \rangle_R} \rightarrow \lambda_1 \quad \text{as } R \rightarrow +\infty,$$

and, together with (4.12) and Lemma 4.3, it follows that  $\lambda_R^y \rightarrow \lambda_1$  as  $R \rightarrow +\infty$ . As already emphasized, this provides the desired conclusion.  $\square$

Now we are in a position to give the proof of Proposition 4.1.

*Proof of Proposition 4.1.* Assume (2.11) and  $\lambda_1 < 0$ . Let  $p$  be a continuous nonnegative bounded solution of the stationary problem (2.14), with  $p|_{\bar{I}} \in C^2(\bar{I})$  for each patch  $I$  in  $\mathbb{R}$ . Assume that  $p \not\equiv 0$ . By an immediate induction, the strong maximum principle and the Hopf lemma then imply that  $p > 0$  in  $\mathbb{R}$ . Now, from Lemma 4.5, there is  $R > 0$  such that

$$\forall y \in \mathbb{R}, \quad \lambda_R^y < \frac{\lambda_1}{2} < 0, \quad (4.13)$$

where  $(\lambda_R^y, \varphi_R^y)$  denotes the eigenpair of (4.5). From (2.11), one can choose  $\kappa_0 > 0$  small enough such that, for all  $0 < \kappa \leq \kappa_0$  and  $y \in \mathbb{R}$ ,

$$f(x+y, \kappa \varphi_R^y(x)) \geq f_s(x+y, 0) \kappa \varphi_R^y(x) + \frac{\lambda_1}{2} \kappa \varphi_R^y(x) \quad \text{for all } x \in [-R, R] \setminus (S-y). \quad (4.14)$$

For each  $y \in \mathbb{R}$ , the function  $p^y := p(\cdot + y)$  satisfies

$$\begin{cases} -d(x+y)(p^y)''(x) = f(x+y, p^y(x)), & x \in \mathbb{R} \setminus (S-y), \\ p^y(x^-) = p^y(x^+), \quad (p^y)'(x^-) = \sigma(p^y)'(x^+), & x = nl - y, \\ p^y(x^-) = p^y(x^+), \quad \sigma(p^y)'(x^-) = (p^y)'(x^+), & x = nl + l_2 - y, \end{cases} \quad (4.15)$$

while, for each  $\kappa \in (0, \kappa_0]$ , the continuous function  $\kappa\varphi_R^y$  satisfies

$$\begin{cases} -d(x+y)\kappa(\varphi_R^y)''(x) - f_s(x+y, 0)\kappa\varphi_R^y(x) = \lambda_R^y\kappa\varphi_R^y(x), & \text{in } (-R, R) \setminus (S-y), \\ \kappa\varphi_R^y(x^-) = \kappa\varphi_R^y(x^+), \quad \kappa(\varphi_R^y)'(x^-) = \sigma\kappa(\varphi_R^y)'(x^+), & x = nl - y \in (-R, R), \\ \kappa\varphi_R^y(x^-) = \kappa\varphi_R^y(x^+), \quad \sigma\kappa(\varphi_R^y)'(x^-) = \kappa(\varphi_R^y)'(x^+), & x = nl + l_2 - y \in (-R, R), \\ \kappa\varphi_R^y > 0 \text{ in } (-R, R), \quad \kappa\varphi_R^y(\pm R) = 0. \end{cases} \quad (4.16)$$

It then follows from (4.13)–(4.14) that, for each  $\kappa \in (0, \kappa_0]$  and  $y \in \mathbb{R}$ ,

$$-d(x+y)\kappa(\varphi_R^y)''(x) - f(x+y, \kappa\varphi_R^y(x)) \leq \left(\lambda_R^y - \frac{\lambda_1}{2}\right)\kappa\varphi_R^y(x) < 0 \text{ for } x \in (-R, R) \setminus (S-y). \quad (4.17)$$

Let us finally consider any  $y \in \mathbb{R}$  and prove that  $p^y \geq \kappa_0\varphi_R^y$  in  $[-R, R]$ . Assuming not and using the continuity and positivity of  $p^y$  and the continuity of  $\varphi_R^y$ , one can then define

$$\kappa^* = \sup\{\kappa \in (0, \kappa_0] : p^y \geq \kappa\varphi_R^y \text{ in } [-R, R]\} \in (0, \kappa_0)$$

and one has  $p^y \geq \kappa^*\varphi_R^y$  in  $[-R, R]$  with equality at a point  $x_0 \in [-R, R]$ . Since  $p^y > 0$  in  $\mathbb{R}$  and  $\varphi_R^y(\pm R) = 0$ , there holds  $x_0 \in (-R, R)$ . From (4.15)–(4.17) and finitely many applications of the strong maximum principle and the Hopf lemma, one gets that  $p^y \equiv \kappa^*\varphi_R^y$  in  $(-R, R)$  and then in  $[-R, R]$  by continuity, a contradiction with the boundary conditions at  $x = \pm R$ . As a consequence,  $p^y \geq \kappa_0\varphi_R^y$  in  $[-R, R]$ .

Thus,  $p(y) = p^y(0) \geq \kappa_0\varphi_R^y(0)$  for all  $y \in \mathbb{R}$ . Since the function  $y \mapsto \kappa_0\varphi_R^y(0)$  is periodic, continuous and positive, one concludes that  $\inf_{\mathbb{R}} p > 0$ .  $\square$

*Proof of Theorem 2.4.* Assume that  $f$  satisfies (2.11)–(2.12) and that  $\lambda_1 < 0$ . Let  $q$  and  $p$  be two positive bounded solutions of (2.14) (in the sense that  $q$  and  $p$  are continuous in  $\mathbb{R}$  and have restrictions in  $\bar{I}$  of class  $C^2(\bar{I})$  for each patch  $I \subset \mathbb{R}$ ). Applying Proposition 4.1, there exists  $\varepsilon > 0$  such that  $q \geq \varepsilon$  and  $p \geq \varepsilon$  in  $\mathbb{R}$ . One can then define the positive real number

$$\gamma^* = \sup\{\gamma > 0 : q > \gamma p \text{ in } \mathbb{R}\} \in (0, +\infty).$$

We shall prove that  $\gamma^* \geq 1$ , which will easily yield the conclusion by interchanging the roles of  $p$  and  $q$ . Assume by way of contradiction that  $\gamma^* < 1$ , and set  $z := q - \gamma^*p \geq 0$ . From the definition of  $\gamma^*$ , there exists a sequence  $(x_m)_{m \in \mathbb{N}}$  such that  $z(x_m) \rightarrow 0$  as  $m \rightarrow +\infty$ . Moreover, the nonnegative function  $z$  is continuous in  $\mathbb{R}$ , has restrictions in  $\bar{I}$  of class  $C^2(\bar{I})$  for each patch  $I \subset \mathbb{R}$ , and it satisfies

$$\begin{cases} -d(x)z''(x) - f(x, q(x)) + \gamma^*f(x, p(x)) = 0, & x \in \mathbb{R} \setminus S, \\ z(x^-) = z(x^+), \quad z'(x^-) = \sigma z'(x^+), & x = nl, \\ z(x^-) = z(x^+), \quad \sigma z'(x^-) = z'(x^+), & x = nl + l_2. \end{cases} \quad (4.18)$$

As in the proof of Theorem 2.3, let us assume in (2.12) that  $s \mapsto f_1(s)/s$  is decreasing with respect to  $s > 0$  (the case when  $s \mapsto f_2(s)/s$  is decreasing with respect to  $s > 0$  can be handled similarly). Since  $\gamma^* \in (0, 1)$  and  $p$  is positive in  $\mathbb{R}$ , one then has  $\gamma^*f(x, p(x)) = \gamma^*f_1(p(x)) <$

$f_1(\gamma^*p(x)) = f(x, \gamma^*p(x))$  for  $x \in (nl - l_1, nl)$  and  $n \in \mathbb{Z}$ , while  $\gamma^*f(x, p(x)) = \gamma^*f_2(p(x)) \leq f_2(\gamma^*p(x)) = f(x, \gamma^*p(x))$  for  $x \in (nl, nl + l_2)$  and  $n \in \mathbb{Z}$ . Hence, (4.18) implies that

$$\begin{cases} -d_1 z''(x) - f_1(q(x)) + f_1(\gamma^*p(x)) > 0, & x \in (nl - l_1, nl), \\ -d_2 z''(x) - f_2(q(x)) + f_2(\gamma^*p(x)) \geq 0, & x \in (nl, nl + l_2). \end{cases}$$

Therefore,  $z$  satisfies a problem of the type (4.3) for some bounded function  $b$  defined in  $\mathbb{R} \setminus S$ . Hence, as in the proof of Theorem 2.3, if  $x_m \rightarrow \bar{x} \in \mathbb{R}$  up to extraction of a subsequence, one gets a contradiction by using the strong maximum principle and the Hopf lemma. In the general case, let  $\bar{x}_m \in (-l_1, l_2]$  be such that  $x_m - \bar{x}_m \in l\mathbb{Z}$ , and let  $\bar{x}_\infty \in [-l_1, l_2]$  be such that  $\bar{x}_m \rightarrow \bar{x}_\infty$  as  $m \rightarrow +\infty$ , up to extraction of some subsequence. Next, set  $z_m = q_m - \gamma^*p_m$ , where  $q_m(x) := q(x + x_m - \bar{x}_m)$  and  $p_m(x) := p(x + x_m - \bar{x}_m)$ . Since  $d(x)$  and  $f(x, u)$  are periodic with respect to  $x$ , it follows from (4.18) that the functions  $z_m$ 's satisfy

$$\begin{cases} -d(x)z_m''(x) - f(x, q_m(x)) + \gamma^*f(x, p_m(x)) = 0 & x \in \mathbb{R} \setminus S, \\ z_m(x^-) = z_m(x^+), \quad z_m'(x^-) = \sigma z_m'(x^+), & x = nl, \\ z_m(x^-) = z_m(x^+), \quad \sigma z_m'(x^-) = z_m'(x^+), & x = nl + l_2. \end{cases}$$

The sequences of continuous functions  $(q_m)_{m \in \mathbb{N}}$  and  $(p_m)_{m \in \mathbb{N}}$  are bounded in  $L^\infty(\mathbb{R})$ , and then in  $C^{2,\nu}(\bar{I})$  for each  $\nu \in (0, 1)$  and each patch  $I \subset \mathbb{R}$  from standard elliptic estimates. Thus, there exist three continuous functions  $q_\infty \geq \varepsilon$ ,  $p_\infty \geq \varepsilon$  and  $z_\infty = q_\infty - \gamma^*p_\infty \geq 0$  such that, up to extraction of some subsequence,  $(q_m|_{\bar{I}}, p_m|_{\bar{I}}, z_m|_{\bar{I}}) \rightarrow (q_\infty|_{\bar{I}}, p_\infty|_{\bar{I}}, z_\infty|_{\bar{I}})$  in  $C^2(\bar{I})$  as  $m \rightarrow +\infty$  for each patch  $I \subset \mathbb{R}$ . Furthermore,

$$\begin{cases} -d(x)z_\infty''(x) - f(x, q_\infty(x)) + \gamma^*f(x, p_\infty(x)) = 0, & x \in \mathbb{R} \setminus S, \\ z_\infty(x^-) = z_\infty(x^+), \quad z_\infty'(x^-) = \sigma z_\infty'(x^+), & x = nl, \\ z_\infty(x^-) = z_\infty(x^+), \quad \sigma z_\infty'(x^-) = z_\infty'(x^+), & x = nl + l_2, \end{cases}$$

and  $z_\infty \geq 0$  in  $\mathbb{R}$  with  $z_\infty(\bar{x}_\infty) = 0$ . Using the positivity of  $p_\infty$  and the same argument as for problem (4.18) above, one reaches a contradiction.

Consequently,  $\gamma^* \geq 1$ , whence  $q \geq p$  in  $\mathbb{R}$ . By interchanging the roles of  $q$  and  $p$ , one also gets  $p \geq q$  in  $\mathbb{R}$ . The uniqueness is therefore obtained. Furthermore, if  $p$  is a positive solution of (2.14), so is the function  $x \mapsto p(x + l)$ . This implies that  $p$  is periodic. The proof of Theorem 2.4 is thereby complete.  $\square$

## 5 Long-time behavior: proof of Theorem 2.5

This section is devoted to the proof of Theorem 2.5 on the large-time behavior of the solutions of the evolution problem (2.9)–(2.10).

*Proof of Theorem 2.5.* Assume that  $f$  satisfies (2.11)–(2.12). Let  $u$  be the solution, given in Theorem 2.2, of the Cauchy problem (2.9)–(2.10) with a nonnegative, bounded and continuous initial datum  $u_0 \not\equiv 0$ . We know from Theorem 2.2 that  $u$  is continuous in  $[0, +\infty) \times \mathbb{R}$  and positive in  $(0, +\infty) \times \mathbb{R}$ .

(i) Assume that 0 is an unstable solution of (2.14), that is,  $\lambda_1 < 0$ , and let  $p$  be the unique positive bounded and periodic solution of (2.14) given by Theorem 2.3 (i) and Theorem 2.4. The function  $p$  is continuous in  $\mathbb{R}$  and has restriction in  $\bar{I}$  of class  $C^2(\bar{I})$  for each patch  $I \subset \mathbb{R}$ . With the notations of Section 4.2 and from Lemma 4.5, one can fix  $R > 0$  large enough so

that  $\lambda_R^0 < \lambda_1/2 < 0$  and (4.16)–(4.17) hold with  $y = 0$  for all  $\kappa > 0$  small enough, where  $(\lambda_R^0, \varphi_R^0)$  denotes the unique eigenpair solving (4.5) with  $y = 0$ :

$$\begin{cases} -d(x)(\varphi_R^0)''(x) - f_s(x, 0)\varphi_R^0(x) = \lambda_R^0\varphi_R^0(x), & x \in (-R, R) \setminus S, \\ \varphi_R^0(x^-) = \varphi_R^0(x^+), \quad (\varphi_R^0)'(x^-) = \sigma(\varphi_R^0)'(x^+), & x = nl \in (-R, R), \\ \varphi_R^0(x^-) = \varphi_R^0(x^+), \quad \sigma(\varphi_R^0)'(x^-) = (\varphi_R^0)'(x^+), & x = nl + l_2 \in (-R, R), \\ \varphi_R^0 > 0 \text{ in } (-R, R), \quad \varphi_R^0(\pm R) = 0, \quad \|\varphi_R^0\|_{L^\infty(-R, R)} = 1. \end{cases}$$

From the continuity and positivity of  $p$  and  $u(1, \cdot)$  in  $\mathbb{R}$ , one can fix  $\kappa > 0$  small enough so that (4.16)–(4.17) hold with  $y = 0$ , together with  $\kappa\varphi_R^0 < p$  and  $\kappa\varphi_R^0 < u(1, \cdot)$  in  $[-R, R]$ .

Define now a function  $v_0$  in  $\mathbb{R}$  by

$$v_0(x) = \begin{cases} \kappa\varphi_R^0(x) & \text{for } x \in [-R, R], \\ 0 & \text{for } x \in \mathbb{R} \setminus [-R, R]. \end{cases}$$

The function  $v_0$  is nonnegative, continuous and bounded in  $\mathbb{R}$ , with  $v_0 \not\equiv 0$  in  $\mathbb{R}$ . Let  $v$  be the solution of the Cauchy problem (2.9)–(2.10) with initial datum  $v_0$ , given by Theorem 2.2. For each  $n \in \mathbb{N}$ , let  $v^n$  (respectively  $u^n$ ) be the unique bounded classical solution of (3.1)–(3.4) with initial condition  $v^n(0, \cdot) = \delta^n v_0|_{[-nl, nl]}$  (respectively  $u^n(0, \cdot) = \delta^n u_0|_{[-nl, nl]}$ ), with the cut-off function  $\delta^n$  given in (3.20). From Theorem 2.2, the sequence  $(v^n)_{n \in \mathbb{N}}$  (respectively  $(u^n)_{n \in \mathbb{N}}$ ) converges monotonically pointwise in  $[0, +\infty) \times \mathbb{R}$  to the function  $v$  (respectively  $u$ ). For each  $n \in \mathbb{N}$ , there holds  $0 \leq v^n(0, \cdot) \leq v_0 \leq p$  in  $[-nl, nl]$ , hence  $0 \leq v^n(t, x) \leq p(x)$  for all  $(t, x) \in [0, +\infty) \times [-nl, nl]$  by Proposition A.2, and

$$0 \leq v(t, x) \leq p(x) \quad \text{for all } (t, x) \in [0, +\infty) \times \mathbb{R}$$

by passing to the limit  $n \rightarrow +\infty$  (actually, we also know from Theorem 2.2 that  $v > 0$  in  $(0, +\infty) \times \mathbb{R}$ ). Furthermore, for each  $n \geq R/l + 1$ , since  $v^n(t, \pm R) \geq 0$  for all  $t \geq 0$  and since

$$v^n(0, \cdot)|_{[-R, R]} = \delta^n v_0|_{[-R, R]} = v_0|_{[-R, R]} = \kappa\varphi_R^0$$

with  $\kappa\varphi_R^0$  satisfying (4.16)–(4.17), Proposition A.2 again implies that

$$v^n(t, x) \geq \kappa\varphi_R^0(x) \quad \text{for all } (t, x) \in [0, +\infty) \times [-R, R].$$

Together with the nonnegativity of  $v^n$  in  $[0, +\infty) \times [-nl, nl]$  and the definition of  $v_0$ , this yields

$$v^n(h, x) \geq v_0(x) \geq v^n(0, x) \quad \text{for all } x \in [-nl, nl] \text{ and } h \geq 0,$$

hence  $v^n(t+h, x) \geq v^n(t, x)$  for all  $(t, x) \in [0, +\infty) \times [-nl, nl]$  and  $h \geq 0$ , and finally  $v(t+h, x) \geq v(t, x)$  for all  $(t, x) \in [0, +\infty) \times \mathbb{R}$  and  $h \geq 0$ . Therefore, the function  $v$  is non-decreasing in  $t$ . Since it is positive in  $(0, +\infty)$  and since  $v(t, x) \leq p(x)$  for all  $(t, x) \in [0, +\infty) \times \mathbb{R}$ , the Schauder estimates of Theorem 2.2 yield the existence of a positive bounded solution  $q$  of (2.14) such that  $v(t, \cdot) \rightarrow q$  as  $t \rightarrow +\infty$  locally uniformly in  $\mathbb{R}$  and  $v(t, \cdot)|_{\bar{I}} \rightarrow q|_{\bar{I}}$  as  $t \rightarrow +\infty$  in  $C^2(\bar{I})$  for each patch  $I \subset \mathbb{R}$ . It follows from Theorem 2.4 that  $q \equiv p$  in  $\mathbb{R}$ . In other words,

$$v(t, \cdot) \xrightarrow[t \rightarrow +\infty]{} p \text{ locally uniformly in } \mathbb{R} \text{ and } v(t, \cdot)|_{\bar{I}} \xrightarrow[t \rightarrow +\infty]{} p|_{\bar{I}} \text{ in } C^2(\bar{I}) \text{ for each patch } I \subset \mathbb{R}. \quad (5.1)$$

On the other hand, since the continuous functions  $\kappa\varphi_R^0$  and  $u(1, \cdot)$  satisfy  $\kappa\varphi_R^0 < u(1, \cdot)$  in  $[-R, R]$  and since  $u^n(1, \cdot) \rightarrow u(1, \cdot)$  locally uniformly in  $\mathbb{R}$  as  $n \rightarrow +\infty$ , there is  $n_0 \geq R/l$  such that  $\kappa\varphi_R^0(x) \leq u^n(1, x)$  for all  $x \in [-R, R]$  and for all  $n \geq n_0$ . Since  $u^n \geq 0$  in  $[0, +\infty) \times [-nl, nl]$ , it follows that  $v^n(0, x) \leq v_0(x) \leq u^n(1, x)$  for all  $x \in [-nl, nl]$  and for all  $n \geq n_0$ , hence



$v^n(t, x) \leq u^n(t + 1, x)$  for all  $(t, x) \in [0, +\infty) \times [-nl, nl]$  and for all  $n \geq n_0$ , by Proposition A.2. Therefore,

$$v(t, x) \leq u(t + 1, x) \quad \text{for all } (t, x) \in [0, +\infty) \times \mathbb{R}. \quad (5.2)$$

Lastly, define  $M_1 := \max(M, \|u_0\|_{L^\infty(\mathbb{R})})$  with  $M > 0$  as in (2.11). As in the proof of Theorem 2.3 (i), the solution  $w$  (given by Theorem 2.2) of the Cauchy problem (2.9)–(2.10) with initial datum  $M_1$ , is non-increasing in  $t$ , periodic in  $x$ , and converges as  $t \rightarrow +\infty$  uniformly in  $\mathbb{R}$  to a nonnegative periodic bounded solution  $\bar{p}$  of (2.14). Furthermore,

$$0 \leq u(t, x) \leq w(t, x) \quad \text{for all } (t, x) \in [0, +\infty) \times \mathbb{R} \quad (5.3)$$

by Theorem 2.2. Together with (5.1)–(5.2), one infers that  $\bar{p} \geq p (> 0)$  in  $\mathbb{R}$ , and then  $\bar{p} \equiv p$  by Theorem 2.4. Since  $v(t, x) \leq u(t + 1, x) \leq w(t + 1, x)$  for all  $(t, x) \in [0, +\infty) \times \mathbb{R}$ , one then concludes that  $u(t, \cdot) \rightarrow p$  as  $t \rightarrow +\infty$  locally uniformly in  $\mathbb{R}$ , and together with the Schauder estimates of Theorem 2.2, that  $u(t, \cdot)|_{\bar{I}} \rightarrow p|_{\bar{I}}$  as  $t \rightarrow +\infty$  in  $C^2(\bar{I})$  for each patch  $I \subset \mathbb{R}$ .

(ii) Let us now assume that 0 is a stable solution of (2.14), that is,  $\lambda_1 \geq 0$ . By defining  $w$  and  $\bar{p}$  as in the previous paragraph (the definitions of  $w$  and  $\bar{p}$  did not use the stability properties of 0), Theorem 2.3 (ii) then yields  $\bar{p} \equiv 0$  in  $\mathbb{R}$ . Together with (5.3) and the uniform convergence  $w(t, \cdot) \rightarrow \bar{p} = 0$  in  $\mathbb{R}$  as  $t \rightarrow +\infty$ , one concludes that  $u(t, \cdot) \rightarrow 0$  as  $t \rightarrow +\infty$  uniformly in  $\mathbb{R}$ . The proof of Theorem 2.5 is thereby complete.  $\square$

An immediate corollary of Theorem 2.5, which will be used in the proofs of Theorems 2.6 and 2.8 in Section 6, is the following result.

**Corollary 5.1.** *Assume that  $f$  satisfies (2.11)–(2.12) and that 0 is an unstable solution of (2.14) (i.e.,  $\lambda_1 < 0$ ). Let  $p$  be the unique positive bounded and periodic solution of (2.14) given by Theorem 2.3 (i) and Theorem 2.4. Let  $u$  denote the solution of the Cauchy problem (2.9)–(2.10) with a nonnegative bounded and continuous initial datum  $u_0 \not\equiv 0$ . If  $u_0$  is periodic, then  $u(t, \cdot) \rightarrow p$  as  $t \rightarrow +\infty$  uniformly in  $\mathbb{R}$ .*

*Proof.* We already know from Theorem 2.2 that, for each  $t \geq 0$ , the function  $x \mapsto u(t, x)$  is periodic. Since  $u(t, \cdot) \rightarrow p$  as  $t \rightarrow +\infty$  locally uniformly in  $\mathbb{R}$  by Theorem 2.5, the conclusion follows.  $\square$

## 6 Spreading speeds and periodic traveling waves: proofs of Theorems 2.6 and 2.8

This section is devoted to the study of the spatial dynamics of the problem (2.9)–(2.10). We will prove the existence of an asymptotic spreading speed  $c^*$ , which can be given explicitly by a variational formula using principal eigenvalues of certain linear operators. Moreover, the spreading speed coincides with the minimal speed for pulsating traveling waves. The main approach is based on the abstract dynamical systems theory for monostable evolution systems established in the seminal work in [49] and further developed in [30, 31].

Hereafter we assume that the 0 solution of (2.14) is unstable (i.e.,  $\lambda_1 < 0$ ) and that  $f$  satisfies (2.11)–(2.12). By Theorem 2.3 (i) and Theorem 2.4, there exists a unique positive bounded periodic solution  $p$  of (2.14). We point out that, with these hypotheses, populations starting with any bounded nonnegative and non-trivial initial condition always persist, by Theorem 2.5. Using the notations of [31, Section 5], we define  $\mathcal{H} = \mathbb{R}$ ,  $\tilde{\mathcal{H}} = l\mathbb{Z}$ ,  $X = Y = \mathbb{R}$ ,  $\beta = p$  and  $\mathcal{M} = \mathcal{C}_p$ , with  $\mathcal{C}_p$  given in (2.17). We also define a family of maps  $\{Q_t\}_{t \geq 0}$  in  $\mathcal{C}_p$  by

$$Q_t(\omega)(x) = u(t, x; \omega) \quad \text{for } \omega \in \mathcal{C}_p, t \geq 0, \text{ and } x \in \mathbb{R}, \quad (6.1)$$

where  $(t, x) \mapsto u(t, x; \omega)$  denotes the classical solution to the Cauchy problem (2.9)–(2.10) with initial condition  $u(0, \cdot; \omega) = \omega \in \mathcal{C}_p$ , given by Theorem 2.2. In particular,  $Q_0(\omega) = \omega$  for every  $\omega \in \mathcal{C}_p$ , and  $Q_t(0) = 0$  for every  $t \geq 0$ . Furthermore, since the continuous positive function  $p$  solves (2.14) and since  $u(t, \cdot; p)$  is periodic for every  $t \geq 0$  by Theorem 2.2, it follows with similar arguments as in Proposition A.2 (but this time in the class of periodic solutions) that  $u(t, \cdot; p) = p$  for each  $t \geq 0$ , that is,  $Q_t(p) = p$ . Theorem 2.2 then implies that  $u(t, \cdot; \omega) \in \mathcal{C}_p$  for every  $\omega \in \mathcal{C}_p$  and  $t \geq 0$ . In other words, for every  $t \geq 0$ ,  $Q_t$  maps  $\mathcal{C}_p$  into itself.

We recall that a family of maps  $\{Q_t\}_{t \geq 0}$  from  $\mathcal{C}_p$  into itself is said to be a semiflow in  $\mathcal{C}_p$  if it satisfies the following properties:

- (1)  $Q_0(\omega) = \omega$  for all  $\omega \in \mathcal{C}_p$ ;
- (2)  $Q_{t_1}(Q_{t_2}(\omega)) = Q_{t_1+t_2}(\omega)$  for all  $t_1, t_2 \geq 0$  and for all  $\omega \in \mathcal{C}_p$ ;
- (3) the map  $(t, \omega) \mapsto Q_t(\omega)$  is continuous from  $[0, +\infty) \times \mathcal{C}_p$  into  $\mathcal{C}_p$ , with  $\mathcal{C}_p$  equipped with the compact open topology, that is,  $Q_{t_m}(\omega_m) \rightarrow Q_t(\omega)$  as  $m \rightarrow +\infty$  locally uniformly in  $\mathbb{R}$  if  $t_m \rightarrow t$  and  $\omega_m \rightarrow \omega$  locally uniformly in  $\mathbb{R}$  as  $m \rightarrow +\infty$ , with  $(t_m, \omega_m) \in [0, +\infty) \times \mathcal{C}_p$ .

We also say that  $\{Q_t\}_{t \geq 0}$  is monotone in  $\mathcal{C}_p$  if, for every  $t \geq 0$ ,  $Q_t(\omega) \geq Q_t(\omega')$  in  $\mathbb{R}$  provided  $\omega \geq \omega'$  in  $\mathbb{R}$ , with  $\omega, \omega' \in \mathcal{C}_p$ . Lastly,  $\{Q_t\}_{t \geq 0}$  is called subhomogeneous if  $\gamma Q_t(\omega) \leq Q_t(\gamma \omega)$  in  $\mathbb{R}$  for every  $t \geq 0$ ,  $\gamma \in [0, 1]$ , and  $\omega \in \mathcal{C}_p$ .

The following proposition summarizes the properties of the family  $\{Q_t\}_{t \geq 0}$  defined in (6.1).

**Proposition 6.1.** *The family  $\{Q_t\}_{t \geq 0}$  defined in (6.1) is a monotone and subhomogeneous semiflow in  $\mathcal{C}_p$ . Furthermore, for every  $\omega \in \mathcal{C}_p$ ,  $a \in \mathbb{Z}$ ,  $t \geq 0$  and  $x \in \mathbb{R}$ , there holds  $Q_t(\omega(\cdot + a))(x) = Q_t(\omega)(x + a)$ .*

*Proof.* First of all, the property  $Q_0(\omega) = \omega$  is already known by definition, and the monotonicity of  $\{Q_t\}_{t \geq 0}$  follows from Theorem 2.2.

For the proof of the other properties, we consider a sequence  $(\delta^n)_{n \in \mathbb{N}}$  of continuous cut-off functions satisfying (3.20). Throughout the proof, for any  $\omega \in \mathcal{C}_p$  and  $n \in \mathbb{N}$ , we denote by  $(t, x) \mapsto u^n(t, x; \omega)$  the unique continuous solution of the truncated problem (3.1)–(3.4) in  $[0, +\infty) \times [-nl, nl]$  with initial condition  $u^n(0, \cdot; \omega) = \delta^n \omega|_{[-nl, nl]}$ . We know from the proof of Theorem 2.2 that, for each  $(t, x) \in [0, +\infty) \times \mathbb{R}$ , the sequence  $(u^n(t, x; \omega))_{n \geq |x|/l}$  is nonnegative, nondecreasing and converges to  $u(t, x; \omega)$  (and the convergence holds locally uniformly with respect to  $(t, x) \in [0, +\infty) \times \mathbb{R}$ ).

Let us first show that  $Q_{t_1}(Q_{t_2}(\omega)) = Q_{t_1+t_2}(\omega)$  for all  $t_1, t_2 \geq 0$  and for all  $\omega \in \mathcal{C}_p$ . The conclusion holds trivially when  $t_1$  or  $t_2$  is equal to 0, or when  $\omega \equiv 0$  in  $\mathbb{R}$ . So, let us assume that  $t_1 > 0$ ,  $t_2 > 0$  and  $\omega \in \mathcal{C}_p$  with  $\omega \not\equiv 0$  in  $\mathbb{R}$ . Call  $\omega_2 = u(t_2, \cdot; \omega)$ . For each  $n \in \mathbb{N}$ , one has

$$u^n(t_2, x; \omega) \leq u(t_2, x; \omega) = \omega_2(x) = \delta^{n+1}(x) \omega_2(x) = u^{n+1}(0, x; \omega_2) \text{ for all } x \in [-nl, nl],$$

and  $u^n(t_2 + t, \pm nl; \omega) = 0 \leq u^{n+1}(t, \pm nl; \omega_2)$  for all  $t \geq 0$ . Hence, Proposition A.2 implies that  $u^n(t_2 + t, x; \omega) \leq u^{n+1}(t, x; \omega_2)$  for all  $t \geq 0$  and  $x \in [-nl, nl]$ , hence  $u(t_2 + t, x; \omega) \leq u(t, x; \omega_2)$  for all  $(t, x) \in [0, +\infty) \times \mathbb{R}$ . In particular,

$$Q_{t_2+t_1}(\omega)(x) \leq Q_{t_1}(\omega_2)(x) = Q_{t_1}(Q_{t_2}(\omega))(x) \text{ for all } x \in \mathbb{R}. \quad (6.2)$$

To show the reverse inequality, consider any  $\eta \in (0, 1)$  and  $m \in \mathbb{N}$ . Since  $u^n(t_2, \cdot; \omega) \rightarrow u(t_2, \cdot; \omega) = \omega_2$  as  $n \rightarrow +\infty$  locally uniformly in  $\mathbb{R}$  and since the continuous function  $\omega_2$  is positive in  $\mathbb{R}$  by Theorem 2.2 (we here use  $\omega \geq \not\equiv 0$  in  $\mathbb{R}$  and the positivity of  $t_2$ ), there is an integer  $n_{\eta, m} \geq m$  such that

$$u^n(t_2, x; \omega) \geq \eta u(t_2, x; \omega) = \eta \omega_2(x) \geq u^m(0, x; \eta \omega_2) \text{ for all } x \in [-ml, ml] \text{ and } n \geq n_{\eta, m},$$

while  $u^n(t_2 + t, \pm ml; \omega) \geq 0 = u^m(t, \pm ml; \eta\omega_2)$  for all  $t \geq 0$ . Therefore, Proposition A.2 implies that

$$u^n(t_2 + t, x; \omega) \geq u^m(t, x; \eta\omega_2) \text{ for all } t \geq 0, x \in [-ml, ml], \text{ and } n \geq n_{\eta, m},$$

hence  $u(t_2 + t, x; \omega) \geq u^m(t, x; \eta\omega_2)$  for all  $t \geq 0$  and  $x \in [-ml, ml]$ . Since  $m \in \mathbb{N}$  was arbitrary, one gets that  $u(t_2 + t, x; \omega) \geq u(t, x; \eta\omega_2)$  for all  $(t, x) \in [0, +\infty) \times \mathbb{R}$ . Lastly, since from Theorem 2.2 the map  $\eta \mapsto u(t_1, \cdot; \eta\omega_2)$  is continuous from  $[0, 1]$  to the set of continuous bounded functions with the sup norm, one concludes that  $u(t_2 + t_1, x; \omega) \geq u(t_1, x; \omega_2)$  for all  $x \in \mathbb{R}$ , that is,  $Q_{t_2+t_1}(\omega)(x) \geq Q_{t_1}(\omega_2)(x) = Q_{t_1}(Q_{t_2}(\omega))(x)$  for all  $x \in \mathbb{R}$ . Together with (6.2), this yields the desired property  $Q_{t_2+t_1}(\omega) = Q_{t_1}(Q_{t_2}(\omega))$  in  $\mathbb{R}$ .

To show the continuity property, consider any  $(t, \omega) \in [0, +\infty) \times \mathcal{C}_p$  and any sequence  $(t_m, \omega_m)_{m \in \mathbb{N}}$  in  $[0, +\infty) \times \mathcal{C}_p$  such that  $t_m \rightarrow t$  and  $\omega_m \rightarrow \omega$  locally uniformly in  $\mathbb{R}$  as  $m \rightarrow +\infty$ . One has to show that  $u(t_m, \cdot; \omega_m) \rightarrow u(t, \cdot; \omega)$  locally uniformly in  $\mathbb{R}$  as  $m \rightarrow +\infty$ . Let  $T \in (0, +\infty)$  be such that  $0 \leq t \leq T$  and  $0 \leq t_m \leq T$  for all  $m \in \mathbb{N}$ . Take any  $A > 0$ , and let  $\varepsilon > 0$  be arbitrary. There is a function  $\bar{\omega} \in \mathcal{C}_p \cap C^3(\mathbb{R})$  such that  $\|\bar{\omega}\|_{C^3(\mathbb{R})} < +\infty$ ,  $\bar{\omega}' = 0$  at all points of  $S$ , and  $\|\bar{\omega} - \omega\|_{L^\infty(\mathbb{R})} \leq \varepsilon/2$ . For  $m \in \mathbb{N}$ , define  $\bar{\omega}_m : \mathbb{R} \rightarrow \mathbb{R}$  by

$$\bar{\omega}_m = \min(\max(\bar{\omega}, \omega_m - \varepsilon), \omega_m + \varepsilon),$$

that is,  $\bar{\omega}_m(x) = \bar{\omega}(x)$  if  $\omega_m(x) - \varepsilon \leq \bar{\omega}(x) \leq \omega_m(x) + \varepsilon$ ,  $\bar{\omega}_m(x) = \omega_m(x) + \varepsilon$  if  $\bar{\omega}(x) > \omega_m(x) + \varepsilon$ , and  $\bar{\omega}_m(x) = \omega_m(x) - \varepsilon$  if  $\bar{\omega}(x) < \omega_m(x) - \varepsilon$ . Each function  $\bar{\omega}_m$  belongs to  $\mathcal{C}_p$ , and  $\|\bar{\omega}_m - \omega_m\|_{L^\infty(\mathbb{R})} \leq \varepsilon$ . Furthermore, since  $\|\bar{\omega} - \omega\|_{L^\infty(\mathbb{R})} \leq \varepsilon/2$  and  $\omega_m \rightarrow \omega$  as  $m \rightarrow +\infty$  locally uniformly in  $\mathbb{R}$ , one infers that, for each compact set  $\mathcal{K} \subset \mathbb{R}$ , one has

$$\bar{\omega}_m|_{\mathcal{K}} = \bar{\omega}|_{\mathcal{K}} \text{ for all } m \text{ large enough.} \quad (6.3)$$

As  $\|\bar{\omega}\|_{C^3(\mathbb{R})}$  is finite and  $0 \leq u(t, x; \omega_m) \leq p(x)$  for all  $(t, x) \in [0, +\infty) \times \mathbb{R}$  and  $m \in \mathbb{N}$ , standard parabolic estimates yield the existence of a positive constant  $C_1$  such that, for each patch  $I = (a, b) \subset \mathbb{R}$ , the functions  $t \mapsto u(t, (a+b)/2; \bar{\omega}_m)$  belong to  $C^{1,1/4}([0, +\infty))$  for all  $m$  large enough, and  $\|u(\cdot, (a+b)/2; \bar{\omega}_m)\|_{C^{1,1/4}([0, +\infty))} \leq C_1$ . As in the proof of Theorem 2.2, and using here the fact that  $\bar{\omega}$  is bounded in  $C^3(\mathbb{R})$  and  $\bar{\omega}'$  vanishes on  $S$ , it then follows that there are  $\theta \in (0, 1)$  and a positive constant  $C_2$  such that, for each patch  $I \subset \mathbb{R}$  and from the above estimates applied at the middle points of the leftward and rightward adjacent patches, the functions  $u(\cdot, \cdot; \bar{\omega}_m)|_{[0, +\infty) \times \bar{I}}$  belong to  $C_{t;x}^{1,\theta;2,\theta}([0, +\infty) \times \bar{I})$  and

$$\|u(\cdot, \cdot; \bar{\omega}_m)|_{[0, +\infty) \times \bar{I}}\|_{C_{t;x}^{1,\theta;2,\theta}([0, +\infty) \times \bar{I})} \leq C_2$$

for all  $m$  large enough. Up to extraction of a subsequence, the continuous functions  $u(\cdot, \cdot; \bar{\omega}_m)$  converge as  $m \rightarrow +\infty$  locally uniformly in  $[0, +\infty) \times \mathbb{R}$  to a continuous function  $\bar{U}$  such that, for each patch  $I \subset \mathbb{R}$ ,  $\bar{U}|_{[0, +\infty) \times \bar{I}} \in C_{t;x}^{1,\theta;2,\theta}([0, +\infty) \times \bar{I})$ ,  $\|\bar{U}|_{[0, +\infty) \times \bar{I}}\|_{C_{t;x}^{1,\theta;2,\theta}([0, +\infty) \times \bar{I})} \leq C_2$ , and, for every  $\tau > 0$ ,  $u(\cdot, \cdot; \bar{\omega}_m)|_{[0, \tau] \times \bar{I}} \rightarrow \bar{U}|_{[0, \tau] \times \bar{I}}$  as  $m \rightarrow +\infty$  in  $C_{t;x}^{1,2}([0, \tau] \times \bar{I})$ . Notice that  $\bar{U}(0, \cdot) = \bar{\omega}$  from the above limits and (6.3). On the other hand, similarly, there is a positive constant  $C'_1$  such that, for each patch  $I = (a, b) \subset \mathbb{R}$ , the function  $t \mapsto u^n(t, (a+b)/2; \bar{\omega})$  belongs to  $C^{1,1/4}([0, +\infty))$  and  $\|u^n(\cdot, (a+b)/2; \bar{\omega})\|_{C^{1,1/4}([0, +\infty))} \leq C'_1$  for all  $n$  large enough. As above, there is then a positive constant  $C'_2$  such that, for each patch  $I \subset \mathbb{R}$ ,  $u^n(\cdot, \cdot; \bar{\omega})|_{[0, +\infty) \times \bar{I}}$  belongs to  $C_{t;x}^{1,\theta;2,\theta}([0, +\infty) \times \bar{I})$  and  $\|u^n(\cdot, \cdot; \bar{\omega})|_{[0, +\infty) \times \bar{I}}\|_{C_{t;x}^{1,\theta;2,\theta}([0, +\infty) \times \bar{I})} \leq C'_2$  for all  $n$  large enough. Therefore, for each patch  $I \subset \mathbb{R}$ , the restriction to  $[0, +\infty) \times \bar{I}$  of the limit function  $u(\cdot, \cdot; \bar{\omega})$  belongs to  $C_{t;x}^{1,\theta;2,\theta}([0, +\infty) \times \bar{I})$  and

$$\|u(\cdot, \cdot; \bar{\omega})|_{[0, +\infty) \times \bar{I}}\|_{C_{t;x}^{1,\theta;2,\theta}([0, +\infty) \times \bar{I})} \leq C'_2.$$

Since  $u(0, \cdot; \bar{\omega}) = \bar{\omega} = \bar{U}(0, \cdot)$ , Proposition A.4 then implies that  $u(\cdot, \cdot; \bar{\omega}) \equiv \bar{U}$  in  $[0, +\infty) \times \mathbb{R}$ . Therefore, since the limit of any subsequence of  $(u(\cdot, \cdot; \bar{\omega}_m))_{m \in \mathbb{N}}$  is unique, one gets that the whole sequence  $(u(\cdot, \cdot; \bar{\omega}_m))_{m \in \mathbb{N}}$  converges locally uniformly in  $[0, +\infty) \times \mathbb{R}$  to  $u(\cdot, \cdot; \bar{\omega})$ . In particular, there is  $m_0$  such that

$$\max_{[0, T] \times [-A, A]} |u(\cdot, \cdot; \bar{\omega}_m) - u(\cdot, \cdot; \bar{\omega})| \leq \varepsilon \quad \text{for all } m \geq m_0.$$

Finally, since  $\|\bar{\omega} - \omega\|_{L^\infty(\mathbb{R})} \leq \varepsilon/2$  and  $\|\bar{\omega}_m - \omega_m\|_{L^\infty(\mathbb{R})} \leq \varepsilon$  for each  $m$ , formula (3.27) of the proof of Theorem 2.2 yields  $\|u(\cdot, \cdot; \bar{\omega}) - u(\cdot, \cdot; \omega)\|_{L^\infty([0, T] \times \mathbb{R})} \leq \varepsilon e^{LT}$  and  $\|u(\cdot, \cdot; \bar{\omega}_m) - u(\cdot, \cdot; \omega_m)\|_{L^\infty([0, T] \times \mathbb{R})} \leq 2\varepsilon e^{LT}$ , with  $L := \max(\|f'_1\|_{L^\infty([0, \bar{K}] )}, \|f'_2\|_{L^\infty([0, \bar{K}] )})$  and  $\bar{K} := \max(K_1, K_2, \|p\|_{L^\infty(\mathbb{R})})$ . One infers that

$$\max_{[0, T] \times [-A, A]} |u(\cdot, \cdot; \omega_m) - u(\cdot, \cdot; \omega)| \leq \varepsilon (1 + 3e^{LT})$$

for all  $m \geq m_0$ . Since  $\varepsilon > 0$  was arbitrary and  $u(\cdot, \cdot; \omega)$  is continuous in  $[0, +\infty) \times \mathbb{R}$ , this shows that  $u(t_m, \cdot; \omega_m) \rightarrow u(t, \cdot; \omega)$  uniformly in  $[-A, A]$  as  $m \rightarrow +\infty$ , leading to the desired result.

Let us now show that the family  $\{Q_t\}_{t \geq 0}$  is subhomogeneous. So, let us consider any  $\gamma \in [0, 1]$  and  $\omega \in \mathcal{C}_p$ , and let us show that  $\gamma Q_t(\omega) \leq Q_t(\gamma\omega)$  in  $\mathbb{R}$  for all  $t \geq 0$ . From (2.12), it follows that, for every  $n \in \mathbb{N}$ ,  $\gamma f(x, u^n(t, x; \omega)) \leq f(x, \gamma u^n(t, x; \omega))$  for all  $(t, x) \in [0, +\infty) \times ((-nl, nl) \setminus S)$ , hence  $\gamma u^n(\cdot, \cdot; \omega)$  is a subsolution of the problem satisfied by  $u^n(\cdot, \cdot; \gamma\omega)$  in  $[0, +\infty) \times [-nl, nl]$  (with the same initial condition in  $[-nl, nl]$ ). Proposition A.2 implies that  $\gamma u^n(t, x; \omega) \leq u^n(t, x; \gamma\omega)$  for all  $n \in \mathbb{N}$  and  $(t, x) \in [0, +\infty) \times [-nl, nl]$ , and thus  $\gamma u(t, x; \omega) \leq u(t, x; \gamma\omega)$  for all  $(t, x) \in [0, +\infty) \times \mathbb{R}$ , that is,  $\gamma Q_t(\omega) \leq Q_t(\gamma\omega)$  in  $\mathbb{R}$  for all  $t \geq 0$ .

Finally, consider any  $\omega \in \mathcal{C}_p$  and  $a \in l\mathbb{Z}$ . Set  $\omega_a = \omega(\cdot + a)$ . For every  $n \in \mathbb{N}$  and  $x \in [-nl, nl]$ , one has  $\delta^{n+|a|/l+1}(x+a) = 1$ , hence

$$u^n(0, x; \omega_a) = \delta^n(x)\omega(x+a) \leq \omega(x+a) = \delta^{n+|a|/l+1}(x+a)\omega(x+a) = u^{n+|a|/l+1}(0, x+a; \omega).$$

Furthermore,  $u^n(t, \pm nl; \omega_a) = 0 \leq u^{n+|a|/l+1}(t, \pm nl + a; \omega)$  for all  $t \geq 0$ , and the function  $[0, +\infty) \times [-nl, nl] \ni (t, x) \mapsto u^{n+|a|/l+1}(t, x+a; \omega)$  satisfies the same partial differential equations as  $u^n(\cdot, \cdot; \omega_a)$ , as well as the same interface conditions, due to the periodicity of  $d(x)$  and  $f(x, s)$  with respect to  $x$ . Proposition A.2 then implies that  $u^n(t, x; \omega_a) \leq u^{n+|a|/l+1}(t, x+a; \omega)$  for all  $n \in \mathbb{N}$  and  $(t, x) \in [0, +\infty) \times [-nl, nl]$ , hence

$$u(t, x; \omega_a) \leq u(t, x+a; \omega) \quad \text{for all } (t, x) \in [0, +\infty) \times \mathbb{R}.$$

Since this property is valid for every  $a \in l\mathbb{Z}$  and  $\omega \in \mathcal{C}_p$ , one also gets that

$$u(t, x+a; \omega) = u(t, x+a; (\omega_a)_{-a}) \leq u(t, (x+a) - a; \omega_a) = u(t, x; \omega_a)$$

for every  $a \in l\mathbb{Z}$ ,  $\omega \in \mathcal{C}_p$  and  $(t, x) \in [0, +\infty) \times \mathbb{R}$ , and finally  $u(t, x; \omega_a) = u(t, x+a; \omega)$  for all  $(t, x) \in [0, +\infty) \times \mathbb{R}$ . This completes the proof of Proposition 6.1.  $\square$

As a consequence of Theorem 2.2 and Proposition 6.1, we conclude that the solution maps  $Q_t : \mathcal{C}_p \rightarrow \mathcal{C}_p$  satisfy the following properties:

- (E1) for each  $t \geq 0$ ,  $Q_t$  is periodic, that is,  $Q_t(T_a(\omega)) = T_a(Q_t(\omega))$  for all  $a \in l\mathbb{Z}$  and  $\omega \in \mathcal{C}_p$ , where  $T_y$  is the translation operator defined by  $T_y(\tilde{\omega}) = \tilde{\omega}(\cdot + y)$  for  $\tilde{\omega} \in \mathcal{C}_p$  and  $y \in l\mathbb{Z}$ ;
- (E2) the set  $\{Q_t(\mathcal{C}_p) : t \geq 0\} \subset \mathcal{C}_p$  is uniformly bounded and, for each  $t \geq 0$ ,  $Q_t : \mathcal{C}_p \rightarrow \mathcal{C}_p$  is continuous;
- (E3) for each  $t > 0$ , the map  $Q_t : \mathcal{C}_p \rightarrow \mathcal{C}$  is compact with respect to the compact open topology (as a consequence of the regularity estimates of Theorem 2.2);

(E4) for each  $t \geq 0$ ,  $Q_t$  is order-preserving (i.e., monotone);

(E5) for each  $t > 0$ ,  $Q_t$  admits exactly the two periodic fixed points 0 and  $p$  in  $\mathcal{C}_p$ : indeed, on the one hand, one knows that 0 and  $p$  are two fixed points; on the other hand, for each  $\omega \in \mathcal{C}_p \setminus \{0, p\}$ , one has  $Q_{mt}(\omega) \rightarrow p$  locally uniformly in  $\mathbb{R}$  as  $m \rightarrow +\infty$  by Theorem 2.5 (and even uniformly in  $\mathbb{R}$  if  $\omega$  is periodic, by Corollary 5.1), hence  $\omega$  can not be a fixed point of  $Q_t$ .

It then follows from [31, Theorem 5.1] that the time-1 map  $Q_1 : \mathcal{C}_p \rightarrow \mathcal{C}_p$  admits rightward and leftward asymptotic spreading speeds  $c_+^*$  and  $c_-^*$ , in the sense that, 1) if  $\omega \in \mathcal{C}_p$  is compactly supported then

$$\begin{cases} \|Q_n(\omega)\|_{L^\infty([nc, +\infty))} \xrightarrow{n \rightarrow +\infty} 0 & \text{for each } c > c_+^*, \\ \|Q_n(\omega)\|_{L^\infty((-\infty, -nc'])} \xrightarrow{n \rightarrow +\infty} 0 & \text{for each } c' > c_-^*, \end{cases} \quad (6.4)$$

and, 2) if  $c_+^* + c_-^* > 0$ , then, for any  $\delta > 0$ , there is  $r_\delta > 0$  such that

$$\|Q_n(\omega) - p\|_{L^\infty([-nc', nc])} \xrightarrow{n \rightarrow +\infty} 0 \text{ for each } c < c_+^* \text{ and } c' < c_-^* \text{ with } c + c' \geq 0, \quad (6.5)$$

and for each  $\omega \in \mathcal{C}_p$  with  $\omega \geq \delta$  on an interval of length  $2r_\delta$ . In the following, our goal is to give computational formulas for  $c_\pm^*$  via the linear operators approach of [30, 49], from which we eventually deduce that  $c_+^* = c_-^*$ .

Thus, in order to compute  $c_\pm^*$ , we consider the linearized problem of (2.9)–(2.10) at its zero solution:

$$\begin{cases} U_t = d(x)U_{xx} + f_s(x, 0)U, & t > 0, x \in \mathbb{R} \setminus S, \\ U(t, x^-) = U(t, x^+), U_x(t, x^-) = \sigma U_x(t, x^+), & t > 0, x = nl, \\ U(t, x^-) = U(t, x^+), \sigma U_x(t, x^-) = U_x(t, x^+), & t > 0, x = nl + l_2, \\ U(0, \cdot) = \omega \geq 0, \omega \in \mathcal{C}. \end{cases} \quad (6.6)$$

Let  $\{\mathbb{L}_t\}_{t \geq 0}$  be the linear solution maps generated by (6.6), namely,  $\mathbb{L}_t(\omega) = U(t, \cdot; \omega)$  where the function  $(t, x) \mapsto U(t, x; \omega)$  is the solution of (6.6) given by the same truncation and limit process as in the proof of Theorem 2.2 (this solution satisfies the same properties as the solution of the nonlinear problem (2.9)–(2.10) given in Theorem 2.2, with the exception of the global boundedness: the solutions of (6.6) are now bounded only locally with respect to  $t \in [0, +\infty)$  in general). For any given  $\mu \in \mathbb{R}$ , substituting  $U(t, x; \omega) = e^{-\mu x}v(t, x)$  in (6.6) yields

$$\begin{cases} v_t = d(x)v_{xx} - 2d(x)\mu v_x + (d(x)\mu^2 + f_s(x, 0))v, & t > 0, x \in \mathbb{R} \setminus S, \\ v(t, x^-) = v(t, x^+), [-\mu v + v_x](t, x^-) = \sigma[-\mu v + v_x](t, x^+), & t > 0, x = nl, \\ v(t, x^-) = v(t, x^+), \sigma[-\mu v + v_x](t, x^-) = [-\mu v + v_x](t, x^+), & t > 0, x = nl + l_2, \\ v(0, x) = \omega(x)e^{\mu x}, & x \in \mathbb{R}. \end{cases} \quad (6.7)$$

Let  $\{L_{\mu, t}\}_{t \geq 0}$  be the linear solution maps generated by (6.7) and obtained from the substitution  $v(t, x) = e^{\mu x}U(t, x; \omega)$ , that is, for any  $\omega \in \mathcal{C}$  with  $\omega \geq 0$  in  $\mathbb{R}$ ,

$$\mathbb{L}_t(y \mapsto e^{-\mu y}\omega(y))(x) = e^{-\mu x}L_{\mu, t}(\omega)(x), \text{ for } t \geq 0 \text{ and } x \in \mathbb{R}. \quad (6.8)$$

Substituting  $v(t, x) = e^{-\lambda t}\psi(x)$  into (6.7), with  $\psi$  periodic and positive, leads to the following periodic eigenvalue problem:

$$\begin{cases} \mathcal{L}_\mu \psi(x) := -d(x)\psi''(x) + 2d(x)\mu\psi'(x) - (d(x)\mu^2 + f_s(x, 0))\psi(x) = \lambda\psi(x), & x \in \mathbb{R} \setminus S, \\ \psi(x^-) = \psi(x^+), [-\mu\psi + \psi'](x^-) = \sigma[-\mu\psi + \psi'](x^+), & x = nl, \\ \psi(x^-) = \psi(x^+), \sigma[-\mu\psi + \psi'](x^-) = [-\mu\psi + \psi'](x^+), & x = nl + l_2, \\ \psi \text{ is periodic in } \mathbb{R}, \psi > 0, \|\psi\|_{L^\infty(\mathbb{R})} = 1. \end{cases} \quad (6.9)$$

**Lemma 6.2.** *For each  $\mu \in \mathbb{R}$ , the eigenvalue problem (6.9) has a simple principal eigenvalue  $\lambda = \lambda(\mu)$  corresponding to a unique positive continuous periodic principal eigenfunction  $\psi$ , which is such that  $\psi|_{\bar{I}} \in C^\infty(\bar{I})$  for each patch  $I$  in  $\mathbb{R}$ . Moreover, there is a max-inf characterization of  $\lambda(\mu)$ :*

$$\lambda(\mu) = \max_{\psi \in E_\mu} \inf_{x \in \mathbb{R} \setminus S} \frac{\mathcal{L}_\mu \psi(x)}{\psi(x)}, \quad (6.10)$$

where

$$E_\mu = \left\{ \psi \in \mathbb{P} : \psi|_{\bar{I}} \in C^2(\bar{I}) \text{ for each patch } I \subset \mathbb{R}, \psi > 0 \text{ in } \mathbb{R}, \right. \\ \left. \psi \text{ satisfies the interface conditions in (6.9)} \right\}$$

(we recall that  $\mathbb{P}$  is the set of all continuous and periodic functions from  $\mathbb{R}$  to  $\mathbb{R}$ ).<sup>5</sup> Lastly, the function  $\mu \mapsto \lambda(\mu)$  is concave in  $\mathbb{R}$ , and  $\lambda(0) = \lambda_1$ , where  $\lambda_1 < 0$  is the principal eigenvalue of the problem (2.13).

*Proof.* We first fix  $\mu \in \mathbb{R}$ . The existence of a unique principal eigenvalue for problem (6.9) can be shown similarly as for (4.5). This time, we introduce the space  $H$  of periodic functions belonging to  $H_{loc}^1(\mathbb{R})$ , with  $\|u\|_H^2 = \|u\|_{H^1(-l_1, 0)}^2 + (1/k)\|u\|_{H^1(0, l_2)}^2$ , and  $G$  the set of continuous periodic functions  $u$  such that  $u|_{[-l_1, 0]}$  and  $u|_{[0, l_2]}$  are of class  $C^1([-l_1, 0])$  and  $C^1([0, l_2])$  respectively, with  $\|u\|_G = \|u|_{[-l_1, 0]}\|_{C^1([-l_1, 0])} + \|u|_{[0, l_2]}\|_{C^1([0, l_2])}$ . We also set  $\Lambda := \max(f'_1(0) + d_1\mu^2, f'_2(0) + d_2\mu^2) + 1$ . For  $g \in G$ , we consider the following problem

$$\begin{cases} -d(x)u'' + 2d(x)\mu u' + (\Lambda - d(x)\mu^2 - f_s(x, 0))u = g, & \text{in } \mathbb{R} \setminus S, \\ u(x^-) = u(x^+), [-\mu u + u'](x^-) = \sigma[-\mu u + u'](x^+), & x = nl, \\ u(x^-) = u(x^+), \sigma[-\mu u + u'](x^-) = [-\mu u + u'](x^+), & x = nl + l_2, \\ u \text{ is periodic.} \end{cases} \quad (6.11)$$

We can solve this problem first in a weak sense, that is, we look for  $u \in H$  such that  $B(u, z) = \langle g, z \rangle$  for all  $z \in H$ , where the bilinear form  $B$  is defined by

$$B(u, z) = \int_{-l_1}^0 d_1 u' z' + d_1 \mu u' z - d_1 \mu u z' + (\Lambda - d_1 \mu^2 - f'_1(0)) u z \\ + \frac{1}{k} \int_0^{l_2} d_2 u' z' + d_2 \mu u' z - d_2 \mu u z' + (\Lambda - d_2 \mu^2 - f'_2(0)) u z,$$

and the scalar product  $\langle \cdot, \cdot \rangle$  is defined by

$$\langle g, z \rangle = \int_{-l_1}^0 g z + \frac{1}{k} \int_0^{l_2} g z.$$

Clearly, the map  $z \mapsto \langle g, z \rangle$  is continuous in  $H$ , and  $B$  is continuous in  $H \times H$ . Moreover, it is easily seen that, for any  $u \in H$ ,  $B(u, u) \geq \min(d_1, d_2, 1)\|u\|_H^2$ , whence  $B$  is coercive. The Lax-Milgram theorem implies the existence of a unique  $u \in H$  (hence,  $u$  can be identified with its unique continuous representative in  $\mathbb{R}$ ) satisfying  $B(u, z) = \langle g, z \rangle$  for all  $z \in H$ , and  $\|u\|_H \leq C_1 \|g\|_{L^2(-l_1, l_2)}$  for a positive constant  $C_1$  only depending on  $d_{1,2}$  and  $k$ . As for (4.5), one gets that  $u|_{[-l_1, 0]}$  and  $u|_{[0, l_2]}$  are in  $H^2([-l_1, 0])$  and  $H^2([0, l_2])$  and then in  $C^3([-l_1, 0])$

<sup>5</sup>In (6.10), even if the test functions  $\psi$  are positive, continuous in  $\mathbb{R}$ , and have restrictions to  $\bar{I}$  of class  $C^2(\bar{I})$  for each patch  $I \subset \mathbb{R}$ , the infimum of  $\mathcal{L}_\mu \psi(x)/\psi(x)$  is taken over the open set  $\mathbb{R} \setminus S$  and therefore is not a minimum in general. Notice that the quantity  $\mathcal{L}_\mu \psi(x)/\psi(x)$  is in general not defined when  $x \in S$ , even if the limits at  $x^\pm$  exist (but are different in general).



and  $C^3([0, l_2])$  respectively, that  $\|u|_{[-l_1, 0]}\|_{C^3([-l_1, 0])} + \|u|_{[0, l_2]}\|_{C^3([0, l_2])} \leq C_2 \|g\|_G$  with a positive constant  $C_2$  depending only on  $d_{1,2}$ ,  $k$ ,  $l_{1,2}$ ,  $f'_{1,2}(0)$  and  $\mu$ , and that the equations in (6.11) are satisfied pointwise. Therefore, the linear mapping  $T : g \in G \mapsto Tg := u \in G$  is compact. Let now  $\mathcal{K}$  be the cone  $\mathcal{K} = \{u \in G : u \geq 0 \text{ in } \mathbb{R}\}$ . Its interior  $\mathring{\mathcal{K}}$  is not empty, and  $\mathcal{K} \cap (-\mathcal{K}) = \{0\}$ . We claim that, if  $g \in \mathcal{K} \setminus \{0\}$ , then  $u \in \mathring{\mathcal{K}}$ . Indeed, by using the equality  $B(u, z) = \langle g, z \rangle$  with  $z := u^- = \max(-u, 0) \in H$ , one has

$$-\int_{-l_1}^0 (d_1((u^-)')^2 + (\Lambda - d_1\mu^2 - f'_1(0))(u^-)^2) - \int_0^{l_2} (d_2((u^-)')^2 + (\Lambda - d_2\mu^2 - f'_2(0))(u^-)^2) = \langle g, u^- \rangle \geq 0,$$

hence  $u^- \equiv 0$ , that is,  $u \geq 0$  in  $\mathbb{R}$ . From the strong elliptic maximum principle and the Hopf lemma, together with the fact that  $g \not\equiv 0$ , one concludes that  $u > 0$  in  $[-l_1, l_2]$  and then in  $\mathbb{R}$  by periodicity. Therefore,  $T(\mathcal{K} \setminus \{0\}) \subset \mathring{\mathcal{K}}$ . As for (4.5), one then infers from the Krein-Rutman theory the existence and uniqueness of a principal eigenpair  $(\lambda, \psi)$  solving (6.9). We then call  $\lambda(\mu)$  this principal eigenvalue  $\lambda$ . Notice that, for each patch  $I \subset \mathbb{R}$ , the function  $\psi|_{\bar{I}}$  is then of class  $C^\infty(\bar{I})$  since  $d$  and  $f_s(\cdot, 0)$  are constant in  $I$ .

Let us now prove the max-min representation (6.10) of  $\lambda(\mu)$ . Since  $\psi \in E_\mu$ , one has

$$\lambda(\mu) \leq \sup_{\psi \in E_\mu} \inf_{x \in \mathbb{R} \setminus S} \frac{\mathcal{L}_\mu \psi(x)}{\psi(x)}.$$

To show the reverse inequality, assume by way of contradiction that there is  $\varphi \in E_\mu$  such that

$$\lambda(\mu) < \inf_{x \in \mathbb{R} \setminus S} \frac{\mathcal{L}_\mu \varphi(x)}{\varphi(x)}.$$

Then there exists  $\eta > 0$  such that

$$-d(x)\varphi''(x) + 2d(x)\mu\varphi'(x) - (d(x)\mu^2 + f_s(x, 0))\varphi(x) - \lambda(\mu)\varphi(x) \geq \eta\varphi(x) > 0 \quad \text{for all } x \in \mathbb{R} \setminus S.$$

Since  $\psi, \varphi \in E_\mu$ , there exists  $\vartheta > 0$  such that  $\varphi \geq \vartheta\psi$  in  $\mathbb{R}$  with equality somewhere. Set  $w := \varphi - \vartheta\psi$ . Then  $w \in \mathbb{P}$ ,  $w \geq 0$  in  $\mathbb{R}$ ,  $w|_{\bar{I}} \in C^2(\bar{I})$  for each patch  $I \subset \mathbb{R}$ , and  $w$  satisfies

$$\begin{cases} -d(x)w''(x) + 2d(x)\mu w'(x) - (d(x)\mu^2 + f_s(x, 0))w(x) - \lambda(\mu)w(x) > 0, & x \in \mathbb{R} \setminus S, \\ w(x^-) = w(x^+), \quad [-\mu w + w'](x^-) = \sigma[-\mu w + w'](x^+), & x = nl, \\ w(x^-) = w(x^+), \quad \sigma[-\mu w + w'](x^-) = [-\mu w + w'](x^+), & x = nl + l_2, \\ w \text{ is periodic,} \end{cases} \quad (6.12)$$

and there exists  $x_0 \in \mathbb{R}$  such that  $w(x_0) = 0$ . The point  $x_0$  can not belong to  $\mathbb{R} \setminus S$  because of the strict inequality in the first line of (6.12). Therefore,  $w > 0$  in  $\mathbb{R} \setminus S$  and  $x_0 \in S$ . The Hopf lemma then implies that  $w'(x_0^+) > 0$  and  $w'(x_0^-) < 0$ , with  $w(x_0) = 0$ , contradicting the interface conditions in (6.12). One has then reached a contradiction. Hence,

$$\lambda(\mu) \geq \sup_{\psi \in E_\mu} \inf_{x \in \mathbb{R} \setminus S} \frac{\mathcal{L}_\mu \psi(x)}{\psi(x)}.$$

The max-min characterization (6.10) of  $\lambda(\mu)$  follows, and the supremum is a maximum since  $\psi \in E_\mu$ .

Next, we prove the concavity of the function  $\mu \mapsto \lambda(\mu)$ . With the change of functions  $\psi(x) = e^{\mu x} \tilde{\psi}(x)$  in (6.10), one has

$$\frac{\mathcal{L}_\mu \psi(x)}{\psi(x)} = \frac{-d(x)\tilde{\psi}''(x)}{\tilde{\psi}(x)} - f_s(x, 0) \quad \text{for all } x \in \mathbb{R} \setminus S,$$



hence

$$\lambda(\mu) = \max_{\tilde{\psi} \in \tilde{E}_\mu} \inf_{x \in \mathbb{R} \setminus S} \left( \frac{-d(x)\tilde{\psi}''(x)}{\tilde{\psi}(x)} - f_s(x, 0) \right), \quad (6.13)$$

where

$$\begin{aligned} \tilde{E}_\mu = \{ & \tilde{\psi} \in \mathcal{C} : x \mapsto e^{\mu x} \tilde{\psi}(x) \in \mathbb{P}, \tilde{\psi}|_{\bar{I}} \in C^2(\bar{I}) \text{ for each patch } I \subset \mathbb{R}, \tilde{\psi} > 0 \text{ in } \mathbb{R}, \\ & \tilde{\psi} \text{ satisfies the interface conditions in (2.13)} \}. \end{aligned}$$

Consider any real numbers  $\mu_1$  and  $\mu_2$ , and any  $t \in [0, 1]$ , and set  $\mu = t\mu_1 + (1-t)\mu_2$ . One has to verify that  $\lambda(\mu) \geq t\lambda(\mu_1) + (1-t)\lambda(\mu_2)$ . Let  $\tilde{\psi}_1$  and  $\tilde{\psi}_2$  be arbitrarily chosen in  $\tilde{E}_{\mu_1}$  and  $\tilde{E}_{\mu_2}$ , respectively. Define  $z_1 = \ln \tilde{\psi}_1$ ,  $z_2 = \ln \tilde{\psi}_2$ ,  $z = tz_1 + (1-t)z_2$  and  $\tilde{\psi} = e^z$ . We claim that  $\tilde{\psi} \in \tilde{E}_\mu$ . In fact, since  $\tilde{\psi} = \tilde{\psi}_1^t \tilde{\psi}_2^{1-t}$ , then  $\tilde{\psi} \in \mathcal{C}$  and  $\tilde{\psi}|_{\bar{I}} \in C^2(\bar{I})$  for each patch  $I \subset \mathbb{R}$ . Furthermore, the function  $x \mapsto e^{\mu x} \tilde{\psi}(x) = (e^{\mu_1 x} \tilde{\psi}_1(x))^t \times (e^{\mu_2 x} \tilde{\psi}_2(x))^{1-t}$  is periodic, and the flux conditions in (2.13) can be easily derived from  $\tilde{\psi}' = (\tilde{\psi}_1^t \tilde{\psi}_2^{1-t})' = t\tilde{\psi}_1^{t-1} \tilde{\psi}_1' \tilde{\psi}_2^{1-t} + (1-t)\tilde{\psi}_1^t \tilde{\psi}_2^{t-1} \tilde{\psi}_2'$  in  $\mathbb{R} \setminus S$  and from the fact that both  $\tilde{\psi}_1 \in \tilde{E}_{\mu_1}$  and  $\tilde{\psi}_2 \in \tilde{E}_{\mu_2}$  satisfy the interface conditions in (2.13). Therefore, by (6.13) we have

$$\lambda(\mu) \geq \inf_{x \in \mathbb{R} \setminus S} \left( \frac{-d(x)\tilde{\psi}''(x)}{\tilde{\psi}(x)} - f_s(x, 0) \right).$$

Notice that, for each  $x \in \mathbb{R} \setminus S$ , one has  $-d(x)\tilde{\psi}''(x)/\tilde{\psi}(x) = -d(x)z''(x) - d(x)(z'(x))^2$ , and

$$\begin{aligned} (z'(x))^2 &= (tz_1'(x) + (1-t)z_2'(x))^2 = t(z_1'(x))^2 + (1-t)(z_2'(x))^2 - t(1-t)(z_1'(x) - z_2'(x))^2 \\ &\leq t(z_1'(x))^2 + (1-t)(z_2'(x))^2, \end{aligned}$$

hence

$$\begin{aligned} \frac{-d(x)\tilde{\psi}''(x)}{\tilde{\psi}(x)} - f_s(x, 0) &\geq t(-d(x)z_1''(x) - d(x)(z_1'(x))^2 - f_s(x, 0)) \\ &\quad + (1-t)(-d(x)z_2''(x) - d(x)(z_2'(x))^2 - f_s(x, 0)). \end{aligned}$$

Eventually, we find that

$$\begin{aligned} \lambda(\mu) &\geq \inf_{x \in \mathbb{R} \setminus S} \left( \frac{-d(x)\tilde{\psi}''(x)}{\tilde{\psi}(x)} - f_s(x, 0) \right) \\ &\geq t \inf_{x \in \mathbb{R} \setminus S} \left( \frac{-d(x)\tilde{\psi}_1''(x)}{\tilde{\psi}_1(x)} - f_s(x, 0) \right) + (1-t) \inf_{x \in \mathbb{R} \setminus S} \left( \frac{-d(x)\tilde{\psi}_2''(x)}{\tilde{\psi}_2(x)} - f_s(x, 0) \right). \end{aligned}$$

Since  $\tilde{\psi}_1$  and  $\tilde{\psi}_2$  were arbitrarily chosen in  $\tilde{E}_{\mu_1}$  and  $\tilde{E}_{\mu_2}$  respectively, one infers from (6.13) that  $\lambda(\mu) \geq t\lambda(\mu_1) + (1-t)\lambda(\mu_2)$ . That is,  $\mu \mapsto \lambda(\mu)$  is concave in  $\mathbb{R}$ , which also yields the continuity of this function. Lastly, we also observe that the problem (6.9) coincides with (2.13) when  $\mu = 0$ , that is,  $\lambda(0) = \lambda_1$ , which is here negative by assumption. This completes the proof of Lemma 6.2.  $\square$

Since for each  $t > 0$  the linear operator  $L_{\mu,t}$  defined by (6.7)–(6.8) is strongly positive and compact, the Krein-Rutman theorem again implies that its spectral radius  $r(L_{\mu,t})$  is positive and is the principal eigenvalue of  $L_{\mu,t}$ , that is,  $r(L_{\mu,t}) = e^{-\lambda(\mu)t}$ .

We are now in a position to give variational formulas for the rightward and leftward asymptotic spreading speeds  $c_\pm^*$  given by (6.4)–(6.5) via the linear operators approach.

**Theorem 6.3.** Let  $c_+^*$  and  $c_-^*$  be the rightward and leftward asymptotic spreading speeds of  $Q_1$ , given by (6.4)–(6.5). Then,

$$c_+^* = \inf_{\mu > 0} \frac{-\lambda(\mu)}{\mu}, \quad c_-^* = \inf_{\mu > 0} \frac{-\lambda(-\mu)}{\mu}. \quad (6.14)$$

Furthermore, we have  $c_+^* = c_-^* > 0$ .

*Proof.* Due to assumption (2.12), and we have  $f(x, s) \leq f_s(x, 0)s$  for all  $x \in \mathbb{R} \setminus S$  and  $s \geq 0$ . Then, for every  $\omega \in \mathcal{C}_p$  and  $n \in \mathbb{N}$ , the solution  $u^n := u^n(\cdot, \cdot; \omega)$  of (3.1)–(3.4) satisfies

$$\begin{cases} u_t^n \leq d(x)u_{xx}^n + f_s(x, 0)u^n, & t > 0, x \in (-nl, nl) \setminus S, \\ u^n(t, x^-) = u^n(t, x^+), \quad u_x^n(t, x^-) = \sigma u_x^n(t, x^+), & t > 0, x \in S_1 \cap (-nl, nl), \\ u^n(t, x^-) = u^n(t, x^+), \quad \sigma u_x^n(t, x^-) = u_x^n(t, x^+), & t > 0, x \in S_2 \cap (-nl, nl). \end{cases}$$

Proposition A.2 and the construction of the solutions  $U(\cdot, \cdot; \omega)$  of (6.6) by using the same truncation and limit process as in the proof of Theorem 2.2 imply that  $u^n(t, x; \omega) \leq U^n(t, x; \omega)$  for all  $(t, x) \in [0, +\infty) \times [-nl, nl]$  and  $n \in \mathbb{N}$ , hence  $u(t, x; \omega) \leq U(t, x, \omega)$  for all  $(t, x) \in [0, +\infty) \times \mathbb{R}$ . In other words,  $Q_t(\omega) \leq \mathbb{L}_t(\omega)$  in  $\mathbb{R}$  for all  $t \geq 0$  and  $\omega \in \mathcal{C}_p$ . Particularly, by taking  $t = 1$ , one has  $Q_1(\omega) \leq \mathbb{L}_1(\omega)$  in  $\mathbb{R}$  for all  $\omega \in \mathcal{C}_p$ . For any  $\mu \in \mathbb{R}$ , one has

$$\mathbb{L}^\mu(\psi)(x) := e^{\mu x} \mathbb{L}_1(y \mapsto e^{-\mu y} \psi(y))(x) = L_{\mu, 1}(\psi)(x) \quad \text{for every } \psi \in \mathbb{P} \text{ and } x \in \mathbb{R},$$

thanks to (6.8). It then follows that  $e^{-\lambda(\mu)}$  is the principal eigenvalue of  $\mathbb{L}^\mu$ . On the other hand, by Lemma 6.2, the function  $\mu \mapsto \ln(e^{-\lambda(\mu)}) = -\lambda(\mu)$  is convex. With similar arguments as in [49, Theorem 2.5] and in [30, Theorem 3.10 (i)], one then obtains that

$$c_+^* \leq \inf_{\mu > 0} \frac{\ln(e^{-\lambda(\mu)})}{\mu} = \inf_{\mu > 0} \frac{-\lambda(\mu)}{\mu}. \quad (6.15)$$

On the other hand, for any given  $\varepsilon > 0$ , there is a  $\delta > 0$  such that

$$f(x, s) \geq (f_s(x, 0) - \varepsilon)s, \quad \text{for all } x \in \mathbb{R} \setminus S \text{ and } s \in [0, \delta].$$

From the continuity of the solutions of (2.9)–(2.10) with respect to the initial conditions, as stated in Theorem 2.2, there is a positive real number  $\eta$  such that  $\eta \leq p$  in  $\mathbb{R}$  and  $u(t, x; \eta) \leq \delta$  for all  $t \in [0, 1]$  and  $x \in \mathbb{R}$ . Define

$$\mathcal{C}_\eta = \{\omega \in \mathcal{C} : 0 \leq \omega \leq \eta \text{ in } \mathbb{R}\}.$$

It then follows from Theorem 2.2 that

$$0 \leq u(t, x; \omega) \leq u(t, x; \eta) \leq \delta, \quad \text{for all } \omega \in \mathcal{C}_\eta, t \in [0, 1] \text{ and } x \in \mathbb{R}.$$

Thus, for any  $\omega \in \mathcal{C}_\eta$ , the solution  $u(\cdot, \cdot; \omega)$  to (2.9)–(2.10) satisfies

$$\begin{cases} u_t \geq d(x)u_{xx} + (f_s(x, 0) - \varepsilon)u, & t \in (0, 1], x \in \mathbb{R} \setminus S, \\ u(t, x^-) = u(t, x^+), \quad u_x(t, x^-) = \sigma u_x(t, x^+), & t \in (0, 1], x = nl, \\ u(t, x^-) = u(t, x^+), \quad \sigma u_x(t, x^-) = u_x(t, x^+), & t \in (0, 1], x = nl + l_2. \end{cases}$$

Consider now the linear problem

$$\begin{cases} V_t = d(x)V_{xx} + (f_s(x, 0) - \varepsilon)V, & t > 0, x \in \mathbb{R} \setminus S, \\ V(t, x^-) = V(t, x^+), \quad V_x(t, x^-) = \sigma V_x(t, x^+), & t > 0, x = nl, \\ V(t, x^-) = V(t, x^+), \quad \sigma V_x(t, x^-) = V_x(t, x^+), & t > 0, x = nl + l_2. \end{cases} \quad (6.16)$$

Let  $\{\mathbb{L}_t^\varepsilon\}_{t \geq 0}$  be the solution maps generated by the above linear system, as for (6.6) above ( $\mathbb{L}_t^0 = \mathbb{L}_t$  for all  $t \geq 0$ ). Then, Proposition A.2 and the construction of the solutions of (6.16) as in the proof of Theorem 2.2 imply that  $\mathbb{L}_t^\varepsilon(\omega) \leq Q_t(\omega)$  in  $\mathbb{R}$  for all  $t \in [0, 1]$  and  $\omega \in \mathcal{C}_\eta$ . In particular,  $\mathbb{L}_1^\varepsilon(\omega) \leq Q_1(\omega)$  in  $\mathbb{R}$  for all  $\omega \in \mathcal{C}_\eta$ .

Denote by  $\lambda^\varepsilon(\mu)$  the first eigenvalue of the following eigenvalue problem:

$$\begin{cases} -d(x)\psi''(x) + 2\mu d(x)\psi'(x) - (d(x)\mu^2 + (f_s(x, 0) - \varepsilon))\psi(x) = \lambda^\varepsilon(\mu)\psi(x), & x \in \mathbb{R} \setminus S, \\ \psi(x^-) = \psi(x^+), \quad [-\mu\psi + \psi'](x^-) = \sigma[-\mu\psi + \psi'](x^+), & x = nl, \\ \psi(x^-) = \psi(x^+), \quad \sigma[-\mu\psi + \psi'](x^-) = [-\mu\psi + \psi'](x^+), & x = nl + l_2, \\ \psi \text{ is periodic in } \mathbb{R}, \quad \psi > 0, \quad \|\psi\|_{L^\infty(\mathbb{R})} = 1. \end{cases}$$

By uniqueness of the principal eigenvalue of (6.9), there holds  $\lambda^\varepsilon(\mu) = \lambda(\mu) + \varepsilon$ . From the convexity of the function  $\mu \mapsto -\lambda^\varepsilon(\mu) = -\lambda(\mu) - \varepsilon$  and the arguments in [49, Theorem 2.4] and in [30, Theorem 3.10 (ii)], one infers that

$$c_+^* \geq \inf_{\mu > 0} \frac{\ln(e^{-\lambda^\varepsilon(\mu)})}{\mu} = \inf_{\mu > 0} \frac{-\lambda(\mu) - \varepsilon}{\mu}, \quad (6.17)$$

and this property is valid for all  $\varepsilon > 0$ . Together with (6.15), (6.17), and the positivity of  $-\lambda(0) = -\lambda_1$ , it follows that

$$c_+^* = \inf_{\mu > 0} \frac{-\lambda(\mu)}{\mu}.$$

By the change of variable  $v(t, x; \omega) = u(t, -l_1 - x; \omega(-l_1 - \cdot))$ ,<sup>6</sup> one gets that  $c_-^*$  is the rightward asymptotic spreading speed of the resulting problem for the solutions  $v(\cdot, \cdot; \omega)$ . Therefore,

$$c_-^* = \inf_{\mu > 0} \frac{-\lambda(-\mu)}{\mu}.$$

Consequently, (6.14) is proved.

Next, for any  $\mu \in \mathbb{R}$ , if  $\psi_\mu$  is the principal eigenfunction of the problem (6.9), with principal eigenvalue  $\lambda(\mu)$ , then the function  $x \mapsto \hat{\psi}(x) := \psi_\mu(-l_1 - x)$  satisfies (6.9) with  $-\mu$  instead of  $\mu$  in the equations and the interface conditions, but with the same eigenvalue  $\lambda(\mu)$ . By uniqueness of the principal eigenvalue, one deduces that  $\lambda(-\mu) = \lambda(\mu)$ . Therefore, (6.14) yields  $c_+^* = c_-^*$ .

Lastly, consider any compactly supported  $\omega \in \mathcal{C}_p$  such that  $\omega \not\equiv 0$  in  $\mathbb{R}$  and  $\omega(x) < p(x)$  for all  $x \in \mathbb{R}$  (and remember that  $\omega$  and  $p$  are continuous, and that  $p$  is periodic and positive in  $\mathbb{R}$ ). From Theorem 2.5 (i), one knows that  $u(t, \cdot; \omega) \rightarrow p$  as  $t \rightarrow +\infty$  locally uniformly in  $\mathbb{R}$ . Hence, there is  $T \in \mathbb{N}$  such that  $u(T, \cdot; \omega) \geq \omega(\cdot \pm l)$  in  $\mathbb{R}$ . Theorem 2.2 and Proposition 6.1 then imply in particular that  $u(2T, \cdot; \omega) \geq \omega(\cdot \pm 2l)$  in  $\mathbb{R}$ , hence  $u(mT, \cdot; \omega) \geq \omega(\cdot \pm ml)$  in  $\mathbb{R}$  for all  $m \in \mathbb{N}$  by an immediate induction. In other words,  $Q_{mT}(\omega) \geq \omega(\cdot \pm ml)$  in  $\mathbb{R}$  for all  $m \in \mathbb{N}$ , and it follows from property (6.4) that  $c_\pm^* > 0$ . This completes the proof of Theorem 6.3.  $\square$

*Proofs of Theorems 2.6 and 2.8.* By [31, Theorems 5.2 and 5.3], together with Theorems 2.5 (i) and 6.3, one directly obtains Theorem 2.6, with spreading speed  $c^* := c_\pm^*$ , as well as the existence of time-nondecreasing periodic rightward and leftward traveling waves for problem (2.9)–(2.10) with all and only all speeds  $c \geq c^*$ . To complete the proof of Theorem 2.8, it is left to show that these periodic traveling waves are strictly monotone in time. For  $c \geq c^* > 0$ , consider a periodic rightward (the case of leftward waves can be handled similarly) traveling wave solving (2.9)–(2.10), written as  $u(t, x) = W(x - ct, x)$  (with  $u(t, x) = Q_{t-t'}(u(t', \cdot))(x)$  for all  $t' \leq t \in \mathbb{R}$  and

<sup>6</sup>Notice that  $p(x) = p(-l_1 - x)$  for all  $x \in \mathbb{R}$  by invariance of (2.14) with respect to this change of variable and by the uniqueness result of Theorem 2.4, hence  $x \mapsto \omega(-l_1 - x) \in \mathcal{C}_p$  for every  $\omega \in \mathcal{C}_p$ .

$x \in \mathbb{R}$ ), where  $W(s, x)$  is periodic in  $x$ , nonincreasing in  $s$ , and  $W(-\infty, x) = p(x)$ ,  $W(+\infty, x) = 0$  for all  $x \in \mathbb{R}$ . Notice in particular that  $0 \leq u(t, x) \leq p(x)$  for all  $(t, x) \in \mathbb{R} \times \mathbb{R}$ , that  $u(t, x) \rightarrow 0$  as  $t \rightarrow -\infty$  and  $u(t, x) \rightarrow p(x)$  as  $t \rightarrow +\infty$  for every  $x \in \mathbb{R}$ , and that  $u(t + h, x) \geq u(t, x)$  for every  $h > 0$  and  $(t, x) \in \mathbb{R}^2$ . From the Schauder estimates of Theorem 2.2 and Proposition A.4, it follows that, for every  $h > 0$  and  $t_0 \in \mathbb{R}$ , either  $u(\cdot + h, \cdot) \equiv u$  in  $(t_0, +\infty) \times \mathbb{R}$ , or  $u(\cdot + h, \cdot) > u$  in  $(t_0, +\infty) \times \mathbb{R}$ . Since  $u(-\infty, x) = 0 < p(x) = u(+\infty, x)$  for every  $x \in \mathbb{R}$ , one easily infers that, for every  $h > 0$ ,  $u(\cdot + h, \cdot) > u$  in  $\mathbb{R} \times \mathbb{R}$ . Therefore,  $u$  is increasing in  $t$  and the periodic rightward traveling wave  $W(x - ct, x)$  is decreasing in its first argument, and all properties of Definition 2.7 are therefore satisfied.  $\square$

## A Comparison principles

In this appendix, we prove comparison results for the problem (2.9)–(2.10), as well as for the patchy model in an interval  $(a, b) \subset \mathbb{R}$  composed of finitely many patches, say  $I_i$  for  $i = 1, \dots, n$ . For the latter, which we first deal with, the landscape  $(a, b)$  can be either bounded or unbounded. Set  $-\infty \leq a = x_0 < x_1 < \dots < x_n = b \leq +\infty$  and  $I_i = (x_{i-1}, x_i)$  for  $i = 1, \dots, n$ . Since the results will be used in the present paper and in the future work [21], we state them in more generality to cover different applications. We consider a one-dimensional parabolic operator

$$\mathcal{L}u := u_t - d(x)u_{xx} - c(t, x)u_x - F(x, u), \quad \text{for } t > 0 \text{ and } x \in (a, b) \setminus \{x_1, \dots, x_{n-1}\} = \bigcup_{i=1}^n I_i,$$

with interface conditions

$$u(t, x_i^-) = u(t, x_i^+) \text{ and } u_x(t, x_i^-) = \sigma_i u_x(t, x_i^+), \quad \text{for } t > 0 \text{ and } i = 1, \dots, n-1. \quad (\text{A.1})$$

If  $a$  or  $b$  is finite, we impose Dirichlet-type boundary conditions:

$$u(t, a) = \varphi^-(t) \text{ or } u(t, b) = \varphi^+(t), \quad \text{for } t \geq 0, \quad (\text{A.2})$$

where  $\varphi^\pm : [0, +\infty) \rightarrow \mathbb{R}$  are given continuous functions. Here, the function  $x \mapsto d(x)$  is assumed to be constant and positive in each patch, i.e.,  $d|_{I_i} = d_i > 0$  for some constant  $d_i$ ,<sup>7</sup> the function  $c$  is assumed to be continuous and bounded in  $(0, T_0) \times \bigcup_{i=1}^n I_i$  for every  $T_0 \in (0, +\infty)$ , the  $\sigma_i$ 's are given positive real numbers, and, for each  $1 \leq i \leq n$ ,  $F(x, s) = f_i(s)$  for  $(x, s) \in I_i \times \mathbb{R}$ , with  $f_i \in C^1(\mathbb{R})$ .

We first give the definition of super- and subsolutions of  $\mathcal{L}u = 0$  associated with the interface and boundary conditions (A.1)–(A.2).

**Definition A.1.** For  $T \in (0, +\infty]$ , we say that a continuous function  $\bar{u} : [0, T) \times \overline{(a, b)} \rightarrow \mathbb{R}$ ,<sup>8</sup> which is assumed to be bounded in  $[0, T_0] \times (a, b)$  for every  $T_0 \in (0, T)$ , is a supersolution for the problem  $\mathcal{L}u = 0$  with interface and boundary conditions (A.1)–(A.2), if  $\bar{u}|_{(0, T) \times \overline{I_i}} \in C_{t,x}^{1;2}((0, T) \times \overline{I_i})$  satisfies  $\mathcal{L}\bar{u}|_{(0, T) \times I_i} \geq 0$  in the classical sense for each  $1 \leq i \leq n$ , and if

$$\bar{u}_x(t, x_i^-) \geq \sigma_i \bar{u}_x(t, x_i^+), \quad \text{for } t \in (0, T) \text{ and } i = 1, \dots, n-1,$$

and

$$\bar{u}(t, a) \geq \varphi^-(t) \text{ or } \bar{u}(t, b) \geq \varphi^+(t), \quad \text{for } t \in [0, T),$$

provided that  $a$  or  $b$  is finite. A subsolution can be defined in a similar way with all the inequality signs above reversed.

<sup>7</sup>From the proofs below, it is easily seen that we can consider more general diffusion coefficients  $d(t, x)$  such that  $d|_{(0, +\infty) \times I_i}$  can be extended to a continuous and positive function in  $[0, +\infty) \times \overline{I_i}$ , for each  $1 \leq i \leq n$ .

<sup>8</sup>The notation  $\overline{(a, b)}$  covers all possible four cases when  $a$  or  $b$  is finite or not. If  $a$  and  $b$  are finite, then  $\overline{(a, b)} = [a, b]$ .

The first result of the appendix is a comparison principle between super- and subsolutions when the interval  $(a, b)$  is bounded.

**Proposition A.2** (Comparison principle in bounded intervals). *Assume that  $-\infty < a < b < +\infty$ . For  $T \in (0, +\infty]$ , let  $\bar{u}$  and  $\underline{u}$  be, respectively, a super- and a subsolution in  $[0, T] \times [a, b]$  of  $\mathcal{L}u = 0$  with (A.1)–(A.2), and assume that  $\bar{u}(0, \cdot) \geq \underline{u}(0, \cdot)$  in  $[a, b]$ . Then,  $\bar{u} \geq \underline{u}$  in  $[0, T] \times [a, b]$  and, if  $\bar{u}(0, \cdot) \not\equiv \underline{u}(0, \cdot)$ , then  $\bar{u} > \underline{u}$  in  $(0, T) \times (a, b)$ .*

*Proof.* Fix any  $T_0 \in (0, T)$  and set

$$M := \max(\|\bar{u}\|_{L^\infty([0, T_0] \times [a, b])}, \|\underline{u}\|_{L^\infty([0, T_0] \times [a, b])}) \quad \text{and} \quad \mu := \max_{1 \leq i \leq n} \|f'_i\|_{L^\infty([-M, M])} \quad (\text{A.3})$$

(notice that  $M$  and  $\mu$  are nonnegative real numbers owing to the assumptions on  $\bar{u}$ ,  $\underline{u}$  and  $f_i$ ). Define

$$w(t, x) := (\bar{u}(t, x) - \underline{u}(t, x)) e^{-\mu t} \quad \text{for } (t, x) \in [0, T_0] \times [a, b].$$

The function  $w$  is continuous in  $[0, T_0] \times [a, b]$ , with restriction in  $(0, T_0] \times \bar{I}_i$  of class  $C_{t;x}^{1;2}((0, T_0] \times \bar{I}_i)$  for each  $1 \leq i \leq n$ , and we see from the mean value theorem that  $w$  satisfies

$$\mathcal{N}w := w_t - d(x)w_{xx} - c(t, x)w_x + (\mu - F_s(x, \eta(t, x)))w \geq 0, \quad \text{for } (t, x) \in (0, T_0] \times \bigcup_{i=1}^n I_i, \quad (\text{A.4})$$

where  $\eta(t, x)$  is an intermediate value between  $\bar{u}(t, x)$  and  $\underline{u}(t, x)$  (hence,  $|\eta(t, x)| \leq M$  and  $\mu - F_s(x, \eta(t, x)) \geq 0$ ). Moreover, there holds

$$w_x(t, x_i^-) \geq \sigma_i w_x(t, x_i^+), \quad \text{for } t \in (0, T_0] \text{ and } i = 1, \dots, n-1, \quad (\text{A.5})$$

together with  $w(0, x) = \bar{u}(0, x) - \underline{u}(0, x) \geq 0$  for all  $x \in [a, b]$ ,  $w(t, a) \geq 0$  and  $w(t, b) \geq 0$  for all  $t \in [0, T_0]$ .

Consider now an arbitrary  $\varepsilon > 0$  and let us introduce the auxiliary function  $z$  defined by

$$z(t, x) := w(t, x) + \varepsilon(t+1) \quad \text{for } (t, x) \in [0, T_0] \times [a, b].$$

The function  $z$  has at least the same regularity as  $w$ , and  $z > 0$  in  $\{0\} \times [a, b]$  and in  $[0, T_0] \times \{a, b\}$ . Moreover,

$$\mathcal{N}z = \mathcal{N}w + \varepsilon + (\mu - F_s(x, \eta(t, x)))\varepsilon(t+1) \geq \varepsilon > 0, \quad \text{for } (t, x) \in (0, T_0] \times \bigcup_{i=1}^n I_i, \quad (\text{A.6})$$

with

$$z_x(t, x_i^-) \geq \sigma_i z_x(t, x_i^+), \quad \text{for } t \in (0, T_0] \text{ and } i = 1, \dots, n-1. \quad (\text{A.7})$$

We claim that  $z(t, x) > 0$  for all  $(t, x) \in [0, T_0] \times [a, b]$ . Assume not. Then, by continuity, there is a point  $(t_0, y_0) \in (0, T_0] \times (a, b)$  such that  $z(t_0, y_0) = \min_{[0, t_0] \times [a, b]} z = 0$ . We first assume that  $y_0 \in I_i$  for some  $1 \leq i \leq n$ . Since  $z_t(t_0, y_0) \leq 0$ ,  $z_x(t_0, y_0) = 0$  and  $z_{xx}(t_0, y_0) \geq 0$ , we see that

$$\mathcal{N}z(t_0, y_0) = z_t(t_0, y_0) - d_i z_{xx}(t_0, y_0) + c(t_0, y_0) z_x(t_0, y_0) + (\mu - f'_i(\eta(t_0, y_0))) z(t_0, y_0) \leq 0, \quad (\text{A.8})$$

which is impossible by (A.6). Thus, necessarily, we can assume without loss of generality that  $y_0 = x_i$  for some  $1 \leq i \leq n-1$  and that  $z > 0$  in  $[0, t_0] \times \bigcup_{i=1}^n I_i$ . Then, the Hopf lemma yields

$$z_x(t_0, x_i^-) < 0 \quad \text{and} \quad z_x(t_1, x_i^+) > 0,$$

which contradicts (A.7). Consequently,  $z > 0$  in  $[0, T_0] \times [a, b]$ . Since  $\varepsilon > 0$  was arbitrarily chosen, we obtain that  $w \geq 0$  in  $[0, T_0] \times [a, b]$ , which immediately implies  $\bar{u} \geq \underline{u}$  in  $[0, T_0] \times [a, b]$ , and then in  $[0, T) \times [a, b]$  since  $T_0 \in (0, T)$  was arbitrary.

Let us now further assume that  $\bar{u}(0, \cdot) \not\equiv \underline{u}(0, \cdot)$  in  $[a, b]$ , hence by continuity  $\bar{u}(0, \cdot) > \underline{u}(0, \cdot)$  in some non-empty open subinterval of  $(a, b)$  which has a non-empty intersection with  $I_i$ , for some  $1 \leq i \leq n$ . Since we already know from the previous paragraph that  $\bar{u} \geq \underline{u}$  in  $[0, T) \times [a, b]$ , it follows from the interior strong parabolic maximum principle that  $\bar{u} > \underline{u}$  in  $(0, T) \times I_i$ . If the interval  $(a, b)$  reduces to a single patch (that is,  $n = 1$ ), then we are done. Otherwise, either  $x_{i-1}$  or  $x_i$  belongs to the open interval  $(a, b)$ . Let us consider the case when  $x_i \in (a, b)$  (hence,  $i \leq n - 1$ ). We now claim that  $\bar{u}(t, x_i) > \underline{u}(t, x_i)$  for all  $t \in (0, T)$ . Indeed, otherwise, there is a time  $t_0 \in (0, T)$  such that  $\bar{u}(t_0, x_i) = \underline{u}(t_0, x_i)$ , and the Hopf lemma then implies that

$$\bar{u}_x(t_0, x_i^-) < \underline{u}_x(t_0, x_i^-).$$

But  $\bar{u}_x(t_0, x_i^+) \geq \underline{u}_x(t_0, x_i^+)$  since  $\bar{u} \geq \underline{u}$  in  $[0, T) \times \overline{I_{i+1}}$  and  $\bar{u}(t_0, x_i) = \underline{u}(t_0, x_i)$ . One finally gets a contradiction with the assumptions on the spatial derivatives of the super- and subsolutions  $\bar{u}$  and  $\underline{u}$  at  $x_i^\pm$ . Therefore,  $\bar{u}(t, x_i) > \underline{u}(t, x_i)$  for all  $t \in (0, T)$ . By continuity and by applying the strong interior parabolic maximum principle in  $(0, T) \times I_{i+1}$ , we infer that  $\bar{u} > \underline{u}$  in  $(0, T) \times I_{i+1}$ . By an immediate induction, going from one patch to the adjacent one in the left or right directions, we get that  $\bar{u} > \underline{u}$  in  $(0, T) \times (a, b)$ . The proof of Proposition A.2 is thereby complete.  $\square$

Then we prove in Proposition A.3 the comparison principle when  $(a, b) = \mathbb{R}$ , still in the case of a finite number of interfaces (the case when the domain is of the form  $(a, +\infty)$  with  $a \in \mathbb{R}$ , or  $(-\infty, b)$  with  $b \in \mathbb{R}$ , can be handled by a combination and a slight modification of the proofs of Propositions A.2 and A.3).

**Proposition A.3** (Comparison principle in  $\mathbb{R}$ ). *For  $T \in (0, +\infty]$ , let  $\bar{u}$  and  $\underline{u}$  be, respectively, a super- and a subsolution in  $[0, T) \times \mathbb{R}$  of  $\mathcal{L}u = 0$  with (A.1), and assume that  $\bar{u}(0, \cdot) \geq \underline{u}(0, \cdot)$  in  $\mathbb{R}$ . Then,  $\bar{u} \geq \underline{u}$  in  $[0, T) \times \mathbb{R}$  and, if  $\bar{u}(0, \cdot) \not\equiv \underline{u}(0, \cdot)$ , then  $\bar{u} > \underline{u}$  in  $(0, T) \times \mathbb{R}$ .*

*Proof.* Fix any  $T_0 \in (0, T)$  and define the nonnegative real numbers  $M$  and  $\mu$  as in (A.3) with this time  $\mathbb{R}$  instead of  $[a, b]$  in the definition of  $M$ . Denote  $w(t, x) := (\bar{u}(t, x) - \underline{u}(t, x))e^{-\mu t}$  for  $(t, x) \in [0, T_0] \times \mathbb{R}$ . The function  $w$  is continuous and bounded in  $[0, T_0] \times \mathbb{R}$ , with restriction in  $(0, T_0] \times \overline{I_i}$  of class  $C_{t;x}^{1;2}((0, T_0] \times \overline{I_i})$  for each  $1 \leq i \leq n$  (notice that, here,  $I_1 = (-\infty, x_1)$  and  $I_n = (x_{n-1}, +\infty)$  are unbounded), and  $w$  still satisfies (A.4)–(A.5), together with  $w(0, \cdot) = \bar{u}(0, \cdot) - \underline{u}(0, \cdot) \geq 0$  in  $\mathbb{R}$ . Set now  $R = \max_{1 \leq i \leq n-1} |x_i| + 1 > 0$ , and let  $\varrho : \mathbb{R} \rightarrow \mathbb{R}$  be a nonnegative  $C^2$  function with bounded first and second order derivatives, and satisfying

$$\begin{cases} \varrho = 0 \text{ in } [-R, R], & \lim_{x \rightarrow +\infty} \varrho(x) = +\infty, \\ \left( \max_{1 \leq i \leq n} d_i \right) \times \|\varrho''\|_{L^\infty(\mathbb{R})} + \|c\|_{L^\infty((0, T_0) \times \cup_{i=1}^n I_i)} \times \|\varrho'\|_{L^\infty(\mathbb{R})} \leq \frac{1}{2}. \end{cases}$$

Let us consider an arbitrary  $\varepsilon > 0$ , and introduce an auxiliary function  $z$  defined by

$$z(t, x) := w(t, x) + \varepsilon(\varrho(|x|) + t + 1) \quad \text{for } (t, x) \in [0, T_0] \times \mathbb{R}.$$

The function  $z$  has at least the same regularity as  $w$ , while  $z(0, x) \geq \varepsilon > 0$  for all  $x \in \mathbb{R}$  and  $z(t, x) \rightarrow +\infty$  as  $|x| \rightarrow +\infty$  uniformly in  $t \in [0, T_0]$ . Moreover,

$$\mathcal{N}z \geq \mathcal{N}w + \varepsilon - \varepsilon d(x)\varrho''(|x|) - \varepsilon |c(t, x)\varrho'(|x|)| + (\mu - F_s(x, \eta(t, x)))\varepsilon(\varrho(|x|) + t + 1) \geq \frac{\varepsilon}{2} > 0$$

for  $(t, x) \in (0, T_0] \times \bigcup_{i=1}^n I_i$ , and (A.7) still holds from (A.5), the definition of  $R$  and the choice of  $\varrho$ .

We claim that  $z(t, x) > 0$  for all  $(t, x) \in [0, T_0] \times \mathbb{R}$ . Assume not. Then, by continuity and the above properties of  $z$ , there is a point  $(t_0, y_0) \in (0, T_0] \times \mathbb{R}$  such that  $z(t_0, y_0) = \min_{[0, t_0] \times \mathbb{R}} z = 0$ . If  $y_0 \in I_i$  for some  $1 \leq i \leq n$ , then we see as in (A.8) that  $\mathcal{N}z(t_0, y_0) \leq 0$ , which is impossible. Thus, we can assume without loss of generality that  $y_0 = x_i$  for some  $1 \leq i \leq n-1$  and that  $z > 0$  in  $[0, t_0] \times \bigcup_{i=1}^n I_i$ . Then, the Hopf lemma yields  $z_x(t_0, x_i^-) < 0$  and  $z_x(t_1, x_i^+) > 0$ , contradicting (A.7). Consequently,  $z > 0$  in  $[0, T_0] \times \mathbb{R}$ . Hence, by passing to the limit as  $\varepsilon \rightarrow 0^+$ , we infer that  $w \geq 0$  in  $[0, T_0] \times \mathbb{R}$ , that is,  $\bar{u} \geq \underline{u}$  in  $[0, T_0] \times \mathbb{R}$ , and then  $\bar{u} \geq \underline{u}$  in  $[0, T) \times \mathbb{R}$  owing to the arbitrariness of  $T_0 \in (0, T)$ .

Lastly, if one further assumes that  $\bar{u}(0, \cdot) \not\equiv \underline{u}(0, \cdot)$ , then the proof of the strict inequality  $\bar{u} > \underline{u}$  in  $(0, T) \times \mathbb{R}$  follows similar lines as in the proof of the preceding proposition.  $\square$

The last statement is a comparison principle for the problem (2.9)–(2.10) involving the countably many interfaces  $S = l\mathbb{Z} \cup (l\mathbb{Z} + l_2)$ . For this problem with a given initial condition, Proposition A.4 also provides the uniqueness of the solutions satisfying the conditions of the statement.

**Proposition A.4** (Comparison principle for (2.9)–(2.10)). *For  $T \in (0, +\infty]$ , let  $\bar{u}$  and  $\underline{u}$  be, respectively, a super- and a subsolution of (2.9)–(2.10) in  $[0, T) \times \mathbb{R}$  (with a natural extension of Definition A.1) with  $\bar{u}(0, \cdot) \geq \underline{u}(0, \cdot)$  in  $\mathbb{R}$ , and assume that, for every  $T_0 \in (0, T)$ , there are  $\theta \in (0, 1)$  and  $C \geq 0$  such that  $\bar{u}|_{[0, T_0] \times \bar{I}}$  and  $\underline{u}|_{[0, T_0] \times \bar{I}}$  are of class  $C_{t;x}^{1,\theta;2,\theta}([0, T_0] \times \bar{I})$  with  $\|\bar{u}|_{[0, T_0] \times \bar{I}}\|_{C_{t;x}^{1,\theta;2,\theta}([0, T_0] \times \bar{I})} + \|\underline{u}|_{[0, T_0] \times \bar{I}}\|_{C_{t;x}^{1,\theta;2,\theta}([0, T_0] \times \bar{I})} \leq C$  for every patch  $I \subset \mathbb{R}$ . Then,  $\bar{u} \geq \underline{u}$  in  $[0, T) \times \mathbb{R}$ , and, if  $\bar{u}(0, \cdot) \not\equiv \underline{u}(0, \cdot)$ , then  $\bar{u} > \underline{u}$  in  $(0, T) \times \mathbb{R}$ .*

*Proof.* Fix any  $T_0 \in (0, T)$  and define the nonnegative real numbers  $M$  and  $\mu$  as in (A.3) with this time  $\mathbb{R}$  instead of  $[a, b]$  in the definition of  $M$ , and the only functions  $f_1$  and  $f_2$  in the definition of  $\mu$ . Denote  $w(t, x) := (\bar{u}(t, x) - \underline{u}(t, x))e^{-\mu t}$  for  $(t, x) \in [0, T_0] \times \mathbb{R}$ . The function  $w$  is continuous and bounded in  $[0, T_0] \times \mathbb{R}$ , and there are  $\theta \in (0, 1)$  and  $C' \geq 0$  such that  $w|_{[0, T_0] \times \bar{I}}$  is of class  $C_{t;x}^{1,\theta;2,\theta}([0, T_0] \times \bar{I})$  with

$$\|w|_{[0, T_0] \times \bar{I}}\|_{C_{t;x}^{1,\theta;2,\theta}([0, T_0] \times \bar{I})} \leq C' \quad (\text{A.9})$$

for every patch  $I \subset \mathbb{R}$ . The function  $w$  still satisfies inequalities similar to (A.4) (with here  $c \equiv 0$  and  $F = f$ ) and (A.5) (with here the countably many interface points  $S$ , and  $\sigma_i = \sigma$  for  $x_i \in S_1$  and  $\sigma_i = 1/\sigma$  for  $x_i \in S_2$ ), together with  $w(0, \cdot) = \bar{u}(0, \cdot) - \underline{u}(0, \cdot) \geq 0$  in  $\mathbb{R}$ . Let us consider an arbitrary  $\varepsilon > 0$ , and introduce an auxiliary function  $z$  defined by

$$z(t, x) := w(t, x) + \varepsilon(t + 1) \quad \text{for } (t, x) \in [0, T_0] \times \mathbb{R}.$$

The function  $z$  satisfies similar regularity estimates as  $w$  (even if it means replacing  $C'$  in (A.9) by another constant depending also on  $\varepsilon$ ), while  $z(0, x) \geq \varepsilon > 0$  for all  $x \in \mathbb{R}$ . Moreover, with the same notations as in (A.4) (with  $c \equiv 0$  and  $F = f$ ), one has

$$\mathcal{N}z(t, x) = \mathcal{N}w(t, x) + \varepsilon + (\mu - f_s(x, \eta(t, x)))\varepsilon(t + 1) \geq \varepsilon > 0 \quad \text{for } (t, x) \in [0, T_0] \times (\mathbb{R} \setminus S),$$

and (A.7) still holds from (A.5), at the interface points  $S$ . We claim that  $z > 0$  in  $[0, T_0] \times \mathbb{R}$ . Assume not. Then, by continuity and the regularity estimates of  $z$ , there are  $t_0 \in (0, T_0]$  and a sequence  $(y_m)_{m \in \mathbb{N}}$  in  $\mathbb{R}$  such that  $z(t_0, y_m) \rightarrow 0$  as  $m \rightarrow +\infty$ , and  $z > 0$  in  $[0, t_0] \times \mathbb{R}$ . Let  $\bar{y}_m \in (-l_1, l_2]$  be such that  $y_m - \bar{y}_m \in l\mathbb{Z}$  and  $\bar{y}_m \rightarrow \bar{y}_\infty \in [-l_1, l_2]$  as  $m \rightarrow +\infty$ , up to extraction of a subsequence. From the regularity estimates on  $z$ , there is a bounded continuous



function  $z_\infty : [0, T_0] \times \mathbb{R} \rightarrow \mathbb{R}$  such that  $z_\infty|_{[0, T_0] \times \bar{I}} \in C_{t;x}^{1;2}([0, T_0] \times \bar{I})$  for each patch  $I \subset \mathbb{R}$ , and, up to extraction of a subsequence,

$$z(\cdot, \cdot + y_m - \bar{y}_m)|_{[0, T_0] \times \bar{I}} \rightarrow z_\infty|_{[0, T_0] \times \bar{I}} \text{ as } m \rightarrow +\infty \text{ in } C_{t;x}^{1;2}([0, T_0] \times \bar{I}),$$

for every patch  $I \subset \mathbb{R}$ . Moreover,  $z_\infty \geq 0$  in  $[0, t_0] \times \mathbb{R}$ , and  $z_\infty(t_0, \bar{y}_\infty) = 0$ . Since the functions  $d(x)$  and  $f(x, s)$  are periodic with respect to  $x$ , the function  $z_\infty$  satisfies  $(z_\infty)_t - d(x)(z_\infty)_{xx} + 2\mu z_\infty \geq \varepsilon > 0$  in  $[0, t_0] \times (\mathbb{R} \setminus S)$ , as well as the interface conditions (A.7) at the interface points  $S$ . The previous partial differential inequality satisfied by  $z_\infty$  implies that  $\bar{y}_\infty \in S$  and that  $z_\infty(t_0, \cdot) > 0$  in  $\mathbb{R} \setminus S$ . The Hopf lemma then yields  $(z_\infty)_x(t_0, \bar{y}_\infty^+) > 0$  and  $(z_\infty)_x(t_0, \bar{y}_\infty^-) < 0$ , contradicting (A.7). As a consequence,  $z > 0$  in  $[0, T_0] \times \mathbb{R}$ , hence  $w \geq 0$  in  $[0, T_0] \times \mathbb{R}$  due to the arbitrariness of  $\varepsilon > 0$ , and finally  $\bar{u} \geq \underline{u}$  in  $[0, T] \times \mathbb{R}$  due to the arbitrariness of  $T_0 \in (0, T)$ .

Lastly, if one further assumes that  $\bar{u}(0, \cdot) \not\equiv \underline{u}(0, \cdot)$  in  $\mathbb{R}$ , then one concludes as in the proof of Proposition A.2 that  $\bar{u} > \underline{u}$  in  $(0, T) \times \mathbb{R}$ .  $\square$

## Acknowledgements

This work has been carried out in the framework of the A\*MIDEX project (ANR-11-IDEX-0001-02), funded by the "Investissements d'Avenir" French Government program managed by the French National Research Agency (ANR). The research leading to these results has also received funding from ANR project RESISTE (ANR-18-CE45-0019). M. Zhang acknowledges the China Scholarship Council for the two-year financial support during her study at Aix-Marseille Université. This work was initiated while F. Lutscher held the position of Professeur Invité at Aix-Marseille Université. The authors are grateful for this opportunity and the financial support from Aix-Marseille Université. The authors would also like to thank Prof. Xing Liang and Prof. Xiao-Qiang Zhao for many helpful discussions.

## References

- [1] Y. Alqawasmeh, F. Lutscher, *Movement behaviour of fish, harvesting-induced habitat degradation and the optimal size of marine reserves*, Theor. Ecol. 12, 453–466, 2019.
- [2] Y. Alqawasmeh, F. Lutscher, *Persistence and spread of stage-structured populations in heterogeneous landscapes*, J. Math. Biol. 78, 1485–1527, 2019.
- [3] D. G. Aronson, H. F. Weinberger, *Nonlinear diffusion in population genetics, combustion and nerve propagation*, In: Partial Differential Equations and Related Topics, Lecture Notes in Math 446, Springer, New York, 5–49, 1975.
- [4] D. G. Aronson, H. F. Weinberger, *Multidimensional nonlinear diffusions arising in population genetics*, Adv. Math. 30, 33–76, 1978.
- [5] H. Berestycki, F. Hamel, *Front propagation in periodic excitable media*, Comm. Pure Appl. Math. 55, 949–1032, 2002.
- [6] H. Berestycki, F. Hamel, N. Nadirashvili, *The speed of propagation for KPP type problems. I - periodic framework*, J. Europ. Math. Soc. 7, 173–213, 2005.
- [7] H. Berestycki, F. Hamel, L. Roques, *Analysis of the periodically fragmented environment model: I-Species persistence*, J. Math. Biol. 51, 75–113, 2005.
- [8] H. Berestycki, F. Hamel, L. Roques, *Analysis of the periodically fragmented environment model: II-Biological invasions and pulsating travelling fronts*, J. Math. Pures Appl. 84, 1101–1146, 2005.
- [9] H. Berestycki, B. Nicolaenko, B. Scheurer, *Traveling wave solutions to combustion models and their singular limits*, SIAM J. Math. Anal. 16, 1207–1242, 1985.

- [10] M. Bramson, *Convergence of solutions of the Kolmogorov equation to travelling waves*, Memoirs Amers. Math. Soc. 44, 1983.
- [11] P. Brunovský, P. Poláčik, B. Sandstede, *Convergence in general periodic parabolic equations in one space dimension*, Nonlinear Anal. 18, 209–215, 1992.
- [12] C. Cobbold, F. Lutscher, J. Sherratt, *Complex spatial patterns result from the Turing mechanism in a patchy landscape*, Ecol. Complexity 24, 69–81, 2015.
- [13] A. Ducrot, T. Giletti, H. Matano, *Existence and convergence to a propagating terrace in one-dimensional reaction-diffusion equations*, Trans. Amer. Math. Soc. 366, 5541–5566, 2014.
- [14] M. El Smaily, *Min-max formulas for the speeds of pulsating travelling fronts in periodic excitable media*, Ann. Mat. Pura Appl. 189, 47–66, 2010.
- [15] J. Fang, X.-Q. Zhao, *Bistable waves for monotone semiflows with applications*, J. Europ. Math. Soc. 17, 2243–2288, 2015.
- [16] P. C. Fife, *Mathematical aspects of reacting and diffusing systems*, Lecture Notes in Biomath. 28, Springer, 1979.
- [17] P. C. Fife, B. McLeod, *The approach of solutions of nonlinear diffusion equations to traveling front solutions*, Arch. Ration. Mech. Anal. 65, 335–361, 1977.
- [18] R. A. Fisher, *The advance of advantageous genes*, Ann. Eugenics 7, 335–369, 1937.
- [19] J.-S. Guo, F. Hamel, *Propagation and blocking in periodically hostile environments*, Arch. Ration. Mech. Anal. 204, 945–975, 2012.
- [20] F. Hamel, *Qualitative properties of KPP and monostable fronts: monotonicity and exponential decay*, J. Math. Pures Appl. 89, 355–399, 2008.
- [21] F. Hamel, F. Lutscher, M. Zhang, *Propagation phenomena in an environment with two patches*, in preparation
- [22] F. Hamel, J. Nolen, J.-M. Roquejoffre, L. Ryzhik, *A short proof of the logarithmic Bramson correction in Fisher-KPP equations*, Netw. Heterog. Media 8, 275–289, 2013.
- [23] F. Hamel, L. Roques, *Uniqueness and stability properties of monostable pulsating fronts*, J. Europ. Math. Soc. 13, 345–390, 2011.
- [24] D. Henry, *Geometric theory of semilinear parabolic equations*, Lectures Notes in Math., Springer Verlag, New York, 1981.
- [25] J. Huang, W. Shen, *Speeds of spread and propagation for KPP models in time almost and space periodic media*, SIAM J. Appl. Dyn. Syst. 8, 790–821, 2009.
- [26] W. Hudson, B. Zinner, *Existence of traveling waves for reaction-diffusion equations of Fisher type in periodic media*, In: Boundary Value Problems for Functional-Differential Equations. J. Henderson (ed.), World Scientific, 187–199, 1995.
- [27] Ya. I. Kanel', *Certain problems of burning-theory equations*, Soviet Math. Dokl. 2, 48–51, 1961.
- [28] A. N. Kolmogorov, I. G. Petrovsky, N. S. Piskunov, *Étude de l'équation de la diffusion avec croissance de la quantité de matière et son application à un problème biologique*, Bjul. Moskowskogo Gos. Univ. A 1, 1–26, 1937.
- [29] K.-S. Lau, *On the nonlinear diffusion equation of Kolmogorov, Petrovsky, and Piskounov*, J. Diff. Equations 59, 44–70, 1985.
- [30] X. Liang and X. Zhao, *Asymptotic speeds of spread and traveling waves for monotone semiflows with applications*, Comm. Pure Appl. Math. 60, 1–40, 2007.
- [31] X. Liang and X. Zhao, *Spreading speeds and traveling waves for abstract monostable evolution systems*, J. Funct. Anal. 259, 857–903, 2010.
- [32] A. Lunardi, *Analytic semigroups and optimal regularity in parabolic problems*, Birkhäuser, Basel, 1996.

- [33] G. Maciel, C. Cosner, R.S. Cantrell, F. Lutscher, *Evolutionarily stable movement strategies in reaction-diffusion models with edge behavior*, J. Math. Biol. 80, 61–92, 2020.
- [34] G. A. Maciel, F. Lutscher, *How individual movement response to habitat edges affects population persistence and spatial spread*, Am. Nat. 182, 42–52, 2013.
- [35] G. A. Maciel, F. Lutscher, *Allee effects and population spread in patchy landscapes*, J. Biol. Dyn. 9, 109–123, 2015.
- [36] G. A. Maciel, F. Lutscher, *Movement behavior determines competitive outcome and spread rates in strongly heterogeneous landscapes*, Theor. Ecol. 11, 351–365, 2018.
- [37] G. Nadin, *Travelling fronts in space-time periodic media*, J. Math. Pures Appl. 92, 232–262, 2009.
- [38] J. Nolen, M. Rudd, J. Xin, *Existence of KPP fronts in spatially-temporally periodic advection and variational principle for propagation speeds*, Dyn. Part. Diff. Equations 2, 1–24, 2005.
- [39] J. Nolen, J. Xin, *Existence of KPP type fronts in space-time periodic shear flows and a study of minimal speeds based on variational principle*, Disc. Cont. Dyn. Syst. 13, 1217–1234, 2005.
- [40] O. Ovaskainen, S. J. Cornell, *Biased movement at a boundary and conditional occupancy times for diffusion processes*, J. Appl. Probab. 40, 557–580, 2003.
- [41] A. Pazy, *Semigroups of linear operators and applications to partial differential equations*, In: Applied mathematical sciences, vol 44. Springer, Berlin, 1983.
- [42] D. H. Sattinger, *On the stability of waves of nonlinear parabolic systems*, Adv. Math. 22, 312–355, 1976.
- [43] G. Sell, Y. You, *Dynamics of evolutionary equations*, In: Applied mathematical sciences, vol 143. Springer, Berlin, 2002.
- [44] W. Shen, *Variational principle for spreading speeds and generalized propagating speeds in time almost periodic and space periodic KPP models*, Trans. Amer. Math. Soc. 362, 5125–5168, 2010.
- [45] N. Shigesada, K. Kawasaki, *Biological invasions: theory and practice*, Oxford Series in Ecology and Evolution, Oxford: Oxford University Press, 1997.
- [46] N. Shigesada, K. Kawasaki, E. Teramoto, *Traveling periodic waves in heterogeneous environments*, Theor. Population Biol. 30, 143–160, 1986.
- [47] J. G. Skellam, *Random dispersal in theoretical populations*, Biometrika 38, 196–218, 1951.
- [48] K. Uchiyama, *The behavior of solutions of some semilinear diffusion equation for large time*, J. Math. Kyoto Univ. 18, 453–508, 1978.
- [49] H. F. Weinberger, *On spreading speeds and traveling waves for growth and migration models in a periodic habitat*, J. Math. Biol. 45, 511–548, 2002.
- [50] X. Xin, *Existence and stability of travelling waves in periodic media governed by a bistable nonlinearity*, J. Dyn. Diff. Equations 3, 541–573, 1991.
- [51] X. Xin, *Existence of planar flame fronts in convective-diffusive periodic media*, Arch. Ration. Mech. Anal. 121, 205–233, 1992.
- [52] J. X. Xin, *Analysis and modeling of front propagation in heterogeneous media*, SIAM Review 42, 161–230, 2000.
- [53] S.-M. Zheng, *Nonlinear evolution equations*, Pitman Monographs and Surveys in Pure and Applied Mathematics, 133, Chapman & Hall/CRC, Boca Raton, Florida, 2004.