

REGULARITY FOR ORLICZ PHASE PROBLEMS

SUMIYA BAASANDORJ AND SUN-SIG BYUN

ABSTRACT. We provide comprehensive regularity results and optimal conditions for a general class of functionals involving Orlicz multi-phase of the type

$$v \mapsto \int_{\Omega} F(x, v, Dv) dx, \quad (0.1)$$

exhibiting non-standard growth conditions and non-uniformly elliptic properties.

The model functional under consideration is given by the Orlicz multi-phase integral

$$v \mapsto \int_{\Omega} f(x, v) \left[G(|Dv|) + \sum_{k=1}^N a_k(x) H_k(|Dv|) \right] dx, \quad N \geq 1, \quad (0.2)$$

where G, H_k are N -functions and $0 \leq a_k(\cdot) \in L^\infty(\Omega)$ with $0 < \nu \leq f(\cdot) \leq L$. Its ellipticity ratio varies according to the geometry of the level sets $\{a_k(x) = 0\}$ of the modulating coefficient functions $a_k(\cdot)$ for every $k \in \{1, \dots, N\}$.

We give a unified treatment to show various regularity results for such multi-phase problems with the coefficient functions $\{a_k(\cdot)\}_{k=1}^N$ not necessarily Hölder continuous even for a lower level of the regularity. Moreover, assuming that minima of the functional in (0.2) belong to better spaces such as $C^{0,\gamma}(\Omega)$ or $L^\kappa(\Omega)$ for some $\gamma \in (0, 1)$ and $\kappa \in (1, \infty]$, we address optimal conditions on nonlinearity for each variant under which we build comprehensive regularity results.

On the other hand, since there is a lack of homogeneity properties in the nonlinearity, we consider an appropriate scaling with keeping the structures of the problems under which we apply Harmonic type approximation in the setting varying on the a priori assumption on minima. We believe that the methods and proofs developed in this paper are suitable to build regularity theorems for a larger class of non-autonomous functionals.

CONTENTS

1. Introduction	2
2. Notations and preliminaries	7
2.1. Notations	7
2.2. Musielak-Orlicz and Musielak-Orlicz-Sobolev spaces	12
3. Absence of Lavrentiev phenomenon	13
4. Sobolev-Poincaré type inequalities	15
5. Basic regularity results	19
5.1. Local boundedness	20
5.2. Almost standard Caccioppoli inequality	23
5.3. Hölder continuity	27
5.4. The Harnack inequality	30
5.5. Higher integrability results	36
6. Harmonic type approximation	36
7. Comparison estimates	42
8. Proof of Theorem 1.2	57
9. Proof of Theorem 1.1.	64
10. Orlicz double phase problems	67

Date: March 19, 2022.

2010 *Mathematics Subject Classification*. Primary 49N60; Secondary 35B27, 35B65, 35J20.

Key words and phrases. Gradient Hölder regularity; Orlicz multi-phases problems; Lavrentiev phenomenon; non-standard growth.

This work was supported by NRF-2017R1A2B2003877.

11. Regularity results under additional integrability	82
References	88

1. INTRODUCTION

We aim to provide optimal and comprehensive regularity results for minimizers of functionals featuring a non-standard growth and a non-uniform ellipticity. The primary model case in mind under investigation is given by an Orlicz multi-phase functional

$$W^{1,1}(\Omega) \ni v \mapsto \mathcal{P}(v, \Omega) := \int_{\Omega} \Psi(x, |Dv|) dx \quad (1.1)$$

for a bounded open domain $\Omega \subset \mathbb{R}^n$ with $n \geq 2$, where throughout the paper we shall denote by

$$\Psi(x, t) := G(t) + a(x)H_a(t) + b(x)H_b(t) \quad (x \in \Omega, t \geq 0) \quad (1.2)$$

for functions $G, H_a, H_b \in \mathcal{N}$ in the sense of Definition 2.1 and $0 \leq a(\cdot), b(\cdot) \in L^\infty(\Omega)$. The Orlicz multi-phase functional \mathcal{P} in (1.1) is naturally defined for functions $v \in W^{1,1}(\Omega)$, which is a considerable one including the following examples of functionals for the regularity theory:

1. p -growth: $\Psi(x, t) := t^p$, $p > 1$, see for instance [49, 52, 63, 64, 66, 67, 84, 85].
2. Orlicz growth: $\Psi(x, t) := G(t)$, see for instance [8, 24, 42, 43, 65].
3. (p, q) -double phase: $\Psi(x, t) := t^p + a(x)t^q$ for $1 < p \leq q$, see for instance [11, 13, 30, 31, 32, 35].
4. Borderline case of double phase: $\Psi(x, t) := t^p + a(x)t^p \log(1 + t)$ for $1 < p$, see for instance [12, 21].
5. Multi-phase: $\Psi(x, t) := t^p + a(x)t^q + b(x)t^s$ for $1 < p \leq q, s$, see for instance [7, 38].
6. Orlicz double phase: $\Psi(x, t) := G(t) + a(x)H_a(t)$, see for instance [6, 22].
7. Orlicz multi-phase: $\Psi(x, t) := G(t) + a(x)H_a(t) + b(x)H_b(t)$, see for instance [7].

Over last several years a systematic analysis of the functionals aforementioned has been an object of intensive studies for the regularity theory. Among them the (p, q) -double phase functional is a significant example given by

$$W^{1,1}(\Omega) \ni v \mapsto \mathcal{P}_{p,q}(v, \Omega) := \int_{\Omega} [|Dv|^p + a(x)|Dv|^q] dx, \quad 1 < p \leq q. \quad (1.3)$$

Another example is the so-called borderline case of double phase defined by

$$W^{1,1}(\Omega) \ni v \mapsto \mathcal{P}_{\log}(v, \Omega) := \int_{\Omega} [|Dv|^p + a(x)|Dv|^p \log(1 + |Dv|)] dx, \quad 1 < p. \quad (1.4)$$

The last functional we would like to single out is the so-called multi-phase functional introduced in [38] is of type

$$W^{1,1}(\Omega) \ni v \mapsto \mathcal{P}_{p,q,s}(v, \Omega) := \int_{\Omega} [|Dv|^p + a(x)|Dv|^q + b(x)|Dv|^s] dx, \quad 1 < p \leq q, s. \quad (1.5)$$

The (p, q) -double phase functional was initially introduced by Zhikov [86, 87, 90] in order to study the feature of strongly anisotropic materials in the context of homogenization and nonlinear elasticity. A main feature of the functionals $\mathcal{P}_{p,q}$, $\mathcal{P}_{p,q,s}$ and \mathcal{P}_{\log} in (1.3)-(1.5) is that their integrand changes their growth and ellipticity ration depending on the geometric behavior of the coefficient functions $a(\cdot)$ and $b(\cdot)$, which determines the geometry of the mixture of different materials. As shown in [47, 50, 88, 89], such functionals exhibit Lavrentiev phenomenon which means that minimizers are discontinuous.

Each functional mentioned above belongs to a family of functionals satisfying nonstandard growth conditions of (p, q) -type. These are functionals of type

$$W^{1,1}(\Omega) \ni v \mapsto \int_{\Omega} F(x, Dv) dx,$$

whose energy density $F(x, z)$ satisfies

$$|z|^p \lesssim F(x, z) \lesssim |z|^q + 1, \quad 1 < p < q,$$

according to Marcellini's terminology [68, 69, 70]. Over the several decades, functionals with nonstandard growth have been extensively investigated, see for instance [18, 45, 46, 47, 50, 51, 80, 81] and the references therein. Those functionals aforementioned give a relevant example of the energy overlying in the so-called Musielak-Orlicz space which will be described in Section 2.2 below.

For the regularity theory, the optimal conditions for the gradient of a local minimizer v of the functional $\mathcal{P}_{p,q}$ to be Hölder continuous have been discovered in [13, 30, 31]. They are

$$\left\{ \begin{array}{ll} \frac{q}{p} \leq 1 + \frac{\alpha}{n} \quad \text{and} \quad 0 \leq a(\cdot) \in C^{0,\alpha}(\Omega) & \text{if } v \in W^{1,p}(\Omega), \end{array} \right. \quad (1.6a)$$

$$\left\{ \begin{array}{ll} q \leq p + \alpha \quad \text{and} \quad 0 \leq a(\cdot) \in C^{0,\alpha}(\Omega) & \text{if } v \in W^{1,p}(\Omega) \cap L^\infty(\Omega), \end{array} \right. \quad (1.6b)$$

$$\left\{ \begin{array}{ll} q < p + \frac{\alpha}{1-\gamma} \quad \text{and} \quad 0 \leq a(\cdot) \in C^{0,\alpha}(\Omega) & \text{if } v \in W^{1,p}(\Omega) \cap C^{0,\gamma}(\Omega) \text{ for some } \gamma \in (0, 1). \end{array} \right. \quad (1.6c)$$

Those conditions in (1.6a)-(1.6c) are essentially sharp in the sense of Lavrentiev gap for the functional $\mathcal{P}_{p,q}$. Here we refer to the very recent interesting paper [9] and see also [47, 50]. On the other hand, letting $a(\cdot) \in C^{\omega_a}(\Omega)$ with a concave function $\omega_a : [0, \infty) \rightarrow [0, \infty)$ vanishing at the origin, the conditions for a local minimizer v of the functional \mathcal{P}_{\log} to be regular have been discovered in [12], which are

$$\left\{ \begin{array}{ll} v \text{ is Hölder continuous} & \text{if } \limsup_{t \rightarrow 0^+} \omega_a(t) \log \left(\frac{1}{t} \right) < \infty, \end{array} \right. \quad (1.7a)$$

$$\left\{ \begin{array}{ll} v \text{ is Hölder continuous with an arbitrary exponent} & \text{if } \limsup_{t \rightarrow 0^+} \omega_a(t) \log \left(\frac{1}{t} \right) = 0, \end{array} \right. \quad (1.7b)$$

$$\left\{ \begin{array}{ll} Dv \text{ is Hölder continuous} & \text{if } \omega_a(t) \lesssim t^\alpha \text{ with some } \alpha \in (0, 1). \end{array} \right. \quad (1.7c)$$

Furthermore, the optimal condition for a local minimizer of the multi-phase functional $\mathcal{P}_{p,q,s}$ in (1.5) to be gradient Hölder continuous has been obtained in [38], that is

$$\frac{q}{p} \leq 1 + \frac{\alpha}{n}, \quad \frac{s}{p} \leq 1 + \frac{\beta}{n}, \quad 0 \leq a(\cdot) \in C^{0,\alpha}(\Omega) \quad \text{and} \quad 0 \leq b(\cdot) \in C^{0,\beta}(\Omega) \quad (1.8)$$

for some $\alpha, \beta \in (0, 1]$. This condition essentially is a natural outcome of the condition (1.6a) and sharp. So looking at the conditions presented in (1.6a)-(1.6c), (1.8) and the ones (1.7a)-(1.7c), there is a natural question as to whether the coefficient functions $a(\cdot), b(\cdot)$ in (1.3) and (1.5) are necessarily Hölder continuous even for a lower level of the regularity, depending on the a priori assumptions on a local minimizer under consideration. Here in this paper we intend to answer such questions by treating much more general class of functionals with Orlicz multi-phase growth below. Apart from the papers mentioned above, the regularity theory for the double phase problems has been the object of an intensive investigation over the last years, see for instance [14, 20, 25, 36, 37, 58, 73, 74] and the references therein.

In this paper we shall deal with a class of general functionals of type

$$W^{1,1}(\Omega) \ni v \mapsto \mathcal{F}(v, \Omega) := \int_{\Omega} F(x, v, Dv) dx, \quad (1.9)$$

where $F : \Omega \times \mathbb{R} \times \mathbb{R}^n \rightarrow \mathbb{R}$ is a Caratheodory function satisfying the double-sided bound

$$\nu \Psi(x, |z|) \leq F(x, y, z) \leq L \Psi(x, |z|) \quad (x \in \Omega, y \in \mathbb{R}, z \in \mathbb{R}^n), \quad (1.10)$$

where Ψ is the same as in (1.2). Under the growth conditions (1.10), local minimizers (Q -minimizers) of the functional \mathcal{F} in (1.9) for some number $Q \geq 1$ can be defined as follows:

Definition 1.1. A function $u \in W_{\text{loc}}^{1,1}(\Omega)$ is a local minimizer (Q -minimizer) of the functional \mathcal{F} defined in (1.9) if $\Psi(x, |Du|) \in L^1(\Omega)$ and the minimality condition $\mathcal{F}(u, \text{supp}(u - v)) \leq \mathcal{F}(v, \text{supp}(u - v))$ ($\mathcal{F}(u, \text{supp}(u - v)) \leq Q \mathcal{F}(v, \text{supp}(u - v))$) is satisfied, whenever $v \in W_{\text{loc}}^{1,1}(\Omega)$ with $\text{supp}(u - v) \Subset \Omega$.

In the rest of the paper we always assume $a(\cdot) \in C^{\omega_a}(\Omega)$ and $b(\cdot) \in C^{\omega_b}(\Omega)$, where $\omega_a, \omega_b : [0, \infty) \rightarrow [0, \infty)$ are concave functions such that $\omega_a(0) = 0$ and $\omega_b(0) = 0$ unless they are specified. Throughout the paper we define the function $\Lambda : (0, \infty) \times (0, \infty) \rightarrow (0, \infty)$ given by

$$\Lambda(s, t) := \frac{\omega_a(s)}{1 + \omega_a(s)} \frac{H_a(t)}{G(t)} + \frac{\omega_b(s)}{1 + \omega_b(s)} \frac{H_b(t)}{G(t)} \quad \text{for any } s, t > 0. \quad (1.11)$$

We shall consider a local Q -minimizer u of the functional \mathcal{P} in (1.1) or a local minimizer u of the functional \mathcal{F} in (1.9) under one of the following main assumptions:

$$\begin{cases} u \in W^{1, \Psi}(\Omega), \\ \lambda_1 := \sup_{t>0} \Lambda(t, G^{-1}(t^{-n})) < \infty, \end{cases} \quad (1.12)$$

$$\begin{cases} u \in W^{1, \Psi}(\Omega) \cap L^\infty(\Omega), \\ \lambda_2 := \sup_{t>0} \Lambda\left(t, \frac{1}{t}\right) < \infty, \end{cases} \quad (1.13)$$

$$\begin{cases} u \in W^{1, \Psi}(\Omega) \cap C^{0, \gamma}(\Omega) \quad \text{for some } \gamma \in (0, 1), \\ \lambda_3 := \sup_{t>0} \Lambda\left(t^{\frac{1}{1-\gamma}}, \frac{1}{t}\right) < \infty. \end{cases} \quad (1.14)$$

Here G^{-1} is the inverse function of G . We straightforwardly check that the conditions (1.12)-(1.14) read as (1.6a)-(1.6c), respectively, when $G(t) = t^p$, $H_a(t) = t^q$, $w_a(t) = t^\alpha$ and $b(\cdot) \equiv 0$ for some $1 < p \leq q$ and $\alpha \in (0, 1]$. Also the condition (1.8) is the same as (1.12) for relevant choices of the functions. Moreover, (1.12)₂ and (1.13)₂ are the same as the one in (1.7a) when $G(t) = t^p$ and $H_a(t) = t^p \log(1+t)$ for some $p > 1$ and $b(\cdot) \equiv 0$. It has been shown that the assumption (1.10) is not enough already in the special case of $G(t) = t^p$ for $p > 1$ together with $a(\cdot) \equiv 0$ and $b(\cdot) \equiv 0$ for obtaining higher regularity of minimizers of the functional \mathcal{F} in (1.9). In this regard we consider the energy density F in (1.9) of type

$$F(x, y, z) := F_G(x, y, z) + a(x)F_{H_a}(x, y, z) + b(x)F_{H_b}(x, y, z) \quad (1.15)$$

for every $x \in \Omega$, $y \in \mathbb{R}$ and $z \in \mathbb{R}^n$, where $F_G(\cdot)$, $F_{H_a}(\cdot)$ and $F_{H_b}(\cdot)$ are continuous functions belonging to $C^2(\mathbb{R}^n \setminus \{0\})$ with respect to z -variable and satisfying the following structure assumptions with fixed constants $0 < \nu \leq L$:

$$\begin{cases} |D_z F_\Phi(x, y, z)| |z| + |D_{zz}^2 F_\Phi(x, y, z)| |z|^2 \leq L \Phi(|z|), \\ \nu \frac{\Phi(|z|)}{|z|^2} |\xi|^2 \leq \langle D_{zz}^2 F_\Phi(x, y, z) \xi, \xi \rangle, \\ |D_z F_\Phi(x_1, y, z) - D_z F_\Phi(x_2, y, z)| |z| \leq L \omega(|x_1 - x_2|) \Phi(|z|), \\ |F_\Phi(x, y_1, z) - F_\Phi(x, y_2, z)| \leq L \omega(|y_1 - y_2|) \Phi(|z|) \end{cases} \quad (1.16)$$

for every $\Phi \in \{G, H_a, H_b\}$, whenever $x, x_1, x_2 \in \Omega$, $y, y_1, y_2 \in \mathbb{R}$, $z \in \mathbb{R}^n \setminus \{0\}$, $\xi \in \mathbb{R}^n$, here either

$$\omega(t) := \min\{t^\mu, 1\} \quad \text{with some } \mu \in (0, 1) \text{ for all } t \geq 0 \quad (1.17)$$

or

$$\omega : [0, +\infty) \rightarrow [0, +\infty) \text{ is concave such that } \omega(0) = 0 \text{ and } \omega(\cdot) \leq 1. \quad (1.18)$$

The structure conditions in (1.16) are satisfied for instance by the model functional

$$W^{1,1}(\Omega) \ni v \mapsto \int_{\Omega} f(x, v) \Psi(x, |Dv|) dx,$$

where $0 < \nu_1 \leq f(x, y) \leq L_1$ for some constants ν_1, L_1 and for some suitable continuous function $f(\cdot)$ satisfying the following inequality

$$|f(x_1, y_1) - f(x_2, y_2)| \leq L \omega(|x_1 - x_2| + |y_1 - y_2|)$$

whenever $x_1, x_2 \in \mathbb{R}^n$ and $y_1, y_2 \in \mathbb{R}$, where ω is the same as defined in (1.17) or (1.18). We also remark that those general functionals mentioned above have not been considered in the present literature for the regularity theory as far as we are concerned, moreover the functionals in (1.9) with structure assumptions (1.10) and (1.16) is not differentiable with respect to the second variable and so it can not be treated by its

Euler-Lagrange equation. In order to shorten the notations in this paper, for a given local minimizer u of the functional \mathcal{F} , we shall use a set of various basic parameters which is “data of the problem” depending on which assumption of (1.12)-(1.14) is considered as follows:

$$\mathbf{data} \equiv \begin{cases} n, \lambda_1, s(G), s(H_a), s(H_b), \nu, L, \|a\|_{C^{\omega_a}(\Omega)}, \|b\|_{C^{\omega_b}(\Omega)}, \omega(\cdot), \|\Psi(x, |Du|)\|_{L^1(\Omega)}, \\ \|u\|_{L^1(\Omega)}, \omega_a(1), \omega_b(1) & \text{if (1.12) is considered,} \\ n, \lambda_2, s(G), s(H_a), s(H_b), \nu, L, \|a\|_{C^{\omega_a}(\Omega)}, \|b\|_{C^{\omega_b}(\Omega)}, \omega(\cdot), \|u\|_{L^\infty(\Omega)}, \omega_a(1), \omega_b(1) \\ \text{if (1.13) is considered,} \\ n, \lambda_3, s(G), s(H_a), s(H_b), \nu, L, \|a\|_{C^{\omega_a}(\Omega)}, \|b\|_{C^{\omega_b}(\Omega)}, \omega(\cdot), [u]_{0,\gamma}, \omega_a(1), \omega_b(1) \\ \text{if (1.14) is considered,} \end{cases} \quad (1.19)$$

where $\lambda_1, \lambda_2, \lambda_3$ are the same numbers as defined in (1.12)-(1.14) and $s(G), s(H_a), s(H_b)$ are indices of the functions G, H_a, H_b in the sense of Definition 2.1, respectively. For a given local Q -minimizer u of the functional \mathcal{P} , \mathbf{data} is understood by the above set of parameters with the constants L, ν having been replaced by Q in any case of (1.12)-(1.14) into the consideration. With $\Omega_0 \Subset \Omega$ being a fixed open subset, we also denote by $\mathbf{data}(\Omega_0)$ the set of parameters in (1.19) together with $\text{dist}(\Omega_0, \partial\Omega)$ under one of the assumptions (1.12)-(1.14):

$$\mathbf{data}(\Omega_0) \equiv \mathbf{data}, \text{dist}(\Omega_0, \partial\Omega). \quad (1.20)$$

Now we are ready to state our main results in this paper.

Theorem 1.1 (Maximal regularity). *Let $u \in W^{1,\Psi}(\Omega)$ be a local minimizer of the functional \mathcal{F} defined in (1.9), under the assumptions (1.10), (1.16) and (1.17). Suppose that $\omega_a(t) = t^\alpha$ and $\omega_b(t) = t^\beta$ for some $\alpha, \beta \in (0, 1]$. If one of the following assumptions*

$$\begin{cases} (1.12), & (1.21a) \\ (1.13), & (1.21b) \\ (1.14) \text{ with } \limsup_{t \rightarrow 0^+} \Lambda\left(t^{\frac{1}{1-\gamma}}, \frac{1}{t}\right) = 0 & (1.21c) \end{cases}$$

is satisfied, then there exists $\theta \in (0, 1)$ depending only on $n, s(G), s(H_a), s(H_b), \nu, L, \alpha, \beta$ and μ such that $Du \in C_{\text{loc}}^{0,\theta}(\Omega)$.

Theorem 1.2 (Morrey decay). *Let $u \in W^{1,\Psi}(\Omega)$ be a local minimizer of the functional \mathcal{F} defined in (1.9), under the assumptions (1.10), (1.16) and (1.18). If one of the following assumptions*

$$\begin{cases} (1.12) \text{ with } \limsup_{t \rightarrow 0^+} \Lambda(t, G^{-1}(t^{-n})) = 0, & (1.22a) \\ (1.13) \text{ with } \limsup_{t \rightarrow 0^+} \Lambda\left(t, \frac{1}{t}\right) = 0, & (1.22b) \\ (1.14) \text{ with } \limsup_{t \rightarrow 0^+} \Lambda\left(t^{\frac{1}{1-\gamma}}, \frac{1}{t}\right) = 0, & (1.22c) \\ (1.12) \text{ with } \omega_a(t) = t^\alpha \text{ and } \omega_b(t) = t^\beta \text{ for some } \alpha, \beta \in (0, 1], & (1.22d) \\ (1.13) \text{ with } \omega_a(t) = t^\alpha \text{ and } \omega_b(t) = t^\beta \text{ for some } \alpha, \beta \in (0, 1] & (1.22e) \end{cases}$$

is satisfied, then

$$u \in C_{\text{loc}}^{0,\theta}(\Omega) \text{ for every } \theta \in (0, 1). \quad (1.23)$$

Moreover, for every $\sigma \in (0, n)$, there exists a positive constant $c \equiv c(\mathbf{data}(\Omega_0), \sigma)$ such that the decay estimate

$$\int_{B_\rho} \Psi(x, |Du|) dx \leq c \left(\frac{\rho}{R}\right)^{n-\sigma} \int_{B_R} \Psi(x, |Du|) dx \quad (1.24)$$

holds for every concentric balls $B_\rho \subset B_R \subset \Omega_0 \Subset \Omega$ with $R \leq 1$.

First of all we note that the gradient Hölder regularity of a local minimizer in Theorem 1.1 is already optimal in the classical p -Laplace case that $G(t) = t^p$ and $a(\cdot) \equiv b(\cdot) \equiv 0$ [84, 85]. The assumptions in (1.21a)-(1.21c) are optimal by the counterexamples given in [46, 50], see also [9]. The regularity results reported here complement in a unified way the main results of [12, 13, 30, 31, 38], where the functions in (1.3)-(1.5) are considered under the corresponding conditions we have discussed in (1.6a)-(1.6c), (1.7a)-(1.7c) and (1.8), respectively, and the arguments used in those papers are strongly dependent of the number of phases along with the Hölder continuity of the coefficient function in the non-linearity. Our approaches for proving the above theorems are in fact independent of this weakness. The assumptions of the above theorems lead to showing new instances of Lavrentiev phenomenon [86, 87, 88, 89, 90]. According to the classical definition, the Lavrentiev gap for the functional \mathcal{F} defined in (1.9) under the growth assumption (1.10) may appear if

$$\inf_{v \in v_0 + W_0^{1,G}(B)} \mathcal{F}(v, B) < \inf_{v \in v_0 + W_0^{1,G}(B) \cap W_{\text{loc}}^{1,\Psi_\Omega^+}(B)} \mathcal{F}(v, B) \quad (1.25)$$

holds for a ball $B \Subset \Omega$ and a function $v_0 \in W^{1,\infty}(B)$, where Ψ_Ω^+ is defined in (2.3) below. That is, local minimizers of \mathcal{F} do not belong to $W_{\text{loc}}^{1,\Psi_\Omega^+}(B)$ in general. To see this, let us look at the classical case that $G(t) = t^p$, $H_a(t) = t^q$, $\omega_a(t) = t^\alpha$ and $\omega_b(\cdot) \equiv 0$ for $1 < p < q$ and $\alpha \in (0, 1]$ such that

$$1 < p < n < n + \alpha < q. \quad (1.26)$$

Under classical double phase setting together with (1.26), the results of [30, Theorem 4.1] and [47, Section 3] provide us the existence of a coefficient function $a(\cdot) \in C^{0,\alpha}(\Omega)$ and a boundary datum $u_0 \in W^{1,p}(B) \cap L^\infty(B)$ such that the Lavrentiev phenomenon (1.25) is occurred. In this regard, we show that there is no Lavrentiev gap for the functional \mathcal{F} in (1.9) satisfying the basic structure assumption (1.10) under the one of assumptions (1.12)₂, (1.13)₂ and (1.14)₂, see Theorem 3.1 below. The approaches we present in this paper lead to avoiding the use of difference quotient methods employed in [30, 31] for obtaining various regularity properties of minimizers of the functional \mathcal{P} representing the (p, q) -double phase growth. In fact, the difference quotient techniques can deal with the case that the coefficient functions in the nonlinearity are Hölder continuous. On the other hand, we are treating the case of not necessarily having Hölder continuous coefficient functions in the nonlinearity by applying a Harmonic type approximation (see Lemma 6.1 below) for comparing a homogeneous equation with a limiting equation having the lipschitz regularity property (see Lemma 7.3 and 7.4).

Remark 1.1. We would like to point out, in the same spirit as this paper, the results of Theorem 1.1 and Theorem 1.2 can be restated and proved for the functional having a finite number of phases with replacing the function in (1.2) by

$$\Psi(x, t) := G(t) + \sum_{i=1}^N a_i(x) H_i(t), \quad m \geq 1, \quad (1.27)$$

where $G, H_i \in \mathcal{N}$ in the sense of the Definition 2.1 and $a_i(\cdot) \in C^{\omega_i}(\Omega)$ with $\omega_i : [0, \infty) \rightarrow [0, \infty)$ being a concave function vanishing at the origin for every $i \in \{1, \dots, N\}$. Under this setting we replace the function in (1.11) by

$$\Lambda(s, t) := \sum_{i=1}^N \frac{\omega_i(s)}{1 + \omega_i(s)} \frac{H_i(t)}{G(t)} \quad \text{for every } s, t > 0. \quad (1.28)$$

The coefficient functions in Theorem 1.1 along with (1.22d) and (1.22e) in Theorem 1.2 are understood by letting $\omega_i(t) = t^{\alpha_i}$ with some $\alpha_i \in (0, 1]$ for every $i \in \{1, \dots, N\}$.

The contents in this paper could provide a guideline to deal with a very general class of non-autonomous functionals whose energy density behaves like

$$F(x, y, z) \approx \Phi(x, |z|) \quad (1.29)$$

for Φ being a certain Young function as we shall introduce in Definition 2.1 below. The investigation of such problems has been a field of interest for research activities over the decades. In fact, a main difficulty lies in discovering the optimal conditions to be placed on $\Phi(x, t)$ with respect to (x, t) -variables. Here we mention a very recent and interesting paper [60] in which the authors give a reasonable answer to such a question

by considering a class of functionals of Uhlenbeck type without any a priori assumption on the minimizers involved. Essentially, the assumption [60, (VA1)] is not comparable with the assumption (1.12)₂. Moreover, the method used in [60] can not be applicable to treat the regularity of minimizers of the functional \mathcal{F} in (1.9) having the solution dependence. Besides the papers mentioned before, there is a rich literature, see for instance [3, 4, 5, 16, 17, 23, 25, 33, 34, 48, 61, 62, 76, 77, 83] and reference therein. We also refer to a survey paper [71].

Finally, we close the introduction part with outlining the organization of the paper. In the next section, we introduce our basic settings and some backgrounds. In Section 3 we shall deal with the absence of Lavrentiev phenomenon for the functional \mathcal{F} under our main assumptions (1.12)₂, (1.13)₂ and (1.14)₂. Section 4 is devoted to obtaining Sobolev-Poincaré type inequality. In Section 5 we start with showing basic regularity results of quasi-minimizers of the functional \mathcal{P} introduced in (1.1), while in Section 6 we provide a certain harmonic type approximation fitting in our settings. From Section 7 we start with making the comparison estimates in order to reach a limiting functional and in Section 8 and Section 9 we prove Theorem 1.1 and Theorem 1.2. In Section 10 we shall consider a more general class of functionals having the Orlicz double phase growth. Finally, we shall deal with our multi-phase problems under some additional integrability condition of Sobolev type and some relevant optimal condition in Section 11.

2. NOTATIONS AND PRELIMINARIES

2.1. Notations. In what follows we shall always denote by c to mean a generic positive constant, possibly varying from line to line, while special constants will be denoted by $c_1, \bar{c}, c_*, c_\varepsilon$, and so on. All such constants will be always not smaller than one; moreover relevant dependencies on parameters will be emphasized using parentheses, that is, for example $c \equiv c(n, s(G), \nu, L)$ means that c depends only on $n, s(G), \nu, L$. We denote by $B_R(x_0) = \{x \in \mathbb{R}^n : |x - x_0| < R\}$ the open ball in \mathbb{R}^n centered at $x_0 \in \mathbb{R}^n$ with a radius $R > 0$. If the center is clear in the context, we shall omit the center point by writing $B_R \equiv B_R(x_0)$. We shall also denote $B_1 \equiv B_1(0) \subset \mathbb{R}^n$ unless the center is specified. With $f : \mathcal{B} \rightarrow \mathbb{R}^N$ ($N \geq 1$) being a measurable map for a measurable subset $\mathcal{B} \subset \mathbb{R}^n$ having finite and positive measure, we denote by

$$(f)_{\mathcal{B}} \equiv \int_{\mathcal{B}} f(x) dx = \frac{1}{|\mathcal{B}|} \int_{\mathcal{B}} f(x) dx$$

its integral average over \mathcal{B} . For a measurable map $f : \Omega \rightarrow \mathbb{R}$ and an open subset $\mathcal{B} \subset \Omega$ with $\sigma : [0, \infty) \rightarrow [0, \infty)$ being a concave function such that $\sigma(0) = 0$, we shall use the notation as

$$[f]_{\sigma; \mathcal{B}} := \sup_{x, y \in \mathcal{B}, x \neq y} \frac{|f(x) - f(y)|}{\sigma(|x - y|)} \quad \text{and} \quad [f]_{\sigma} \equiv [f]_{\sigma; \Omega}.$$

We denote by $C^\sigma(\Omega)$ the space of uniformly continuous functions on Ω whose modulus of continuity does not exceed σ . The space $C^\sigma(\Omega)$ is endowed with the norm defined for a function f by

$$\|f\|_{C^\sigma(\Omega)} = \|f\|_{L^\infty(\Omega)} + [f]_{\sigma; \Omega}.$$

In particular, if $\sigma(t) = t^\alpha$ for some $\alpha \in (0, 1]$, then we denote

$$[f]_{0, \alpha; \mathcal{B}} := \sup_{x, y \in \mathcal{B}, x \neq y} \frac{|f(x) - f(y)|}{|x - y|^\alpha} \quad \text{and} \quad [f]_{0, \alpha} \equiv [f]_{0, \alpha; \Omega}.$$

For a given concave function $\sigma : [0, \infty) \rightarrow [0, \infty)$ vanishing at the origin, we shall use some elementary properties in the future as

$$\sigma(\lambda t) \leq \lambda \sigma(t) \quad \text{for every } \lambda \geq 1 \quad \text{and} \quad t \geq 0 \quad (2.1)$$

and

$$\frac{1}{\sigma(\lambda t)} \leq \frac{1}{\sigma(t)} + \frac{1}{\lambda \sigma(t)} \quad \text{for every } \lambda, t > 0 \quad \text{unless } \sigma \text{ is constant.} \quad (2.2)$$

Throughout the paper, for any given open subset $\mathcal{B} \subset \Omega$, we shall also use the notations by

$$\begin{aligned} a^-(\mathcal{B}) &:= \inf_{x \in \mathcal{B}} a(x), & a^+(\mathcal{B}) &:= \sup_{x \in \mathcal{B}} a(x), \\ b^-(\mathcal{B}) &:= \inf_{x \in \mathcal{B}} b(x), & b^+(\mathcal{B}) &:= \sup_{x \in \mathcal{B}} b(x), \\ \Psi_{\mathcal{B}}^-(t) &:= G(t) + \inf_{x \in \mathcal{B}} a(x)H_a(t) + \inf_{x \in \mathcal{B}} b(x)H_b(t), \\ \Psi_{\mathcal{B}}^+(t) &:= G(t) + \sup_{x \in \mathcal{B}} a(x)H_a(t) + \sup_{x \in \mathcal{B}} b(x)H_b(t) \end{aligned} \quad (2.3)$$

for every $t \geq 0$.

Definition 2.1. A measurable function $\Phi : \Omega \times [0, \infty) \rightarrow [0, \infty)$ is called a Young function if, for any fixed $x \in \Omega$, the function $\Phi(x, \cdot)$ increasing and convex such that

$$\Phi(x, 0) = 0, \quad \lim_{t \rightarrow \infty} \Phi(x, t) = +\infty, \quad \lim_{t \rightarrow 0^+} \frac{\Phi(x, t)}{t} = 0 \quad \text{and} \quad \lim_{t \rightarrow \infty} \frac{\Phi(x, t)}{t} = +\infty.$$

We denote by $\mathcal{N}(\Omega)$ the set of Young functions $\Phi : \Omega \times [0, \infty) \rightarrow [0, \infty)$ such that, for any fixed $x \in \Omega$, $\Phi(x, \cdot) \in C^1([0, \infty)) \cap C^2((0, \infty))$ and there exists a constant $s(\Phi) \geq 1$ with

$$\frac{1}{s(\Phi)} \leq \frac{\partial_{tt}^2 \Phi(x, t)t}{\partial_t \Phi(x, t)} \leq s(\Phi) \quad (2.4)$$

uniformly for all $x \in \Omega$ and $t > 0$, where in the future we shall call this number $s(\Phi)$ by an index of Φ . Furthermore, we denote also \mathcal{N} to mean the set of Young functions $\Phi \in \mathcal{N}(\Omega)$ such that Φ does not depend on the first variable x .

As a direct consequence of the above definition, for any $\Phi \in \mathcal{N}(\Omega)$ with an index $s(\Phi) \geq 1$ and any fixed point $x \in \Omega$, we can observe

$$t^2 \partial_{tt}^2 \Phi(x, t) \approx t \partial_t \Phi(x, t) \approx \Phi(x, t) \quad (2.5)$$

for uniformly all $t > 0$, where note that all implied constants only depend only on $s(\Phi)$. Now we state some important properties of functions of $\mathcal{N}(\Omega)$, see [6, 7, 22] for their proofs.

Lemma 2.1. Let $\Phi \in \mathcal{N}(\Omega)$ with an index $s(\Phi) \geq 1$. Then, for any fixed $x \in \Omega$, we have

1. $\Lambda_0^{1+\frac{1}{s(\Phi)}} \Phi(x, t) \leq \Phi(x, \Lambda_0 t) \leq \Lambda_0^{s(\Phi)+1} \Phi(x, t)$ for any $\Lambda_0 \geq 1$ and $t \geq 0$.
2. $\lambda_0^{1+s(\Phi)} \Phi(x, t) \leq \Phi(x, \lambda_0 t) \leq \lambda_0^{\frac{1}{s(\Phi)}+1} \Phi(x, t)$ for any $0 < \lambda_0 \leq 1$ and $t \geq 0$.
3. $\Lambda_0^{\frac{1}{1+s(\Phi)}} \Phi_t^{-1}(x, t) \leq \Phi_t^{-1}(x, \Lambda_0 t) \leq \Lambda_0^{\frac{s(\Phi)}{1+s(\Phi)}} \Phi_t^{-1}(x, t)$ for any $\Lambda_0 \geq 1$ and $t \geq 0$.
4. $\lambda_0^{\frac{s(\Phi)}{1+s(\Phi)}} \Phi_t^{-1}(x, t) \leq \Phi_t^{-1}(x, \lambda_0 t) \leq \lambda_0^{\frac{1}{1+s(\Phi)}} \Phi_t^{-1}(x, t)$ for any $0 < \lambda_0 \leq 1$ and $t \geq 0$.

In the above lemma, for a fixed point $x \in \Omega$, $\Phi_t^{-1}(x, t)$ is understood by the inverse function of $\Phi(x, t)$ with respect to t -variable.

Remark 2.1. For a given $\Phi \in \mathcal{N}(\Omega)$ with an index $s(\Phi) \geq 1$, we notice useful but direct consequences of Lemma 2.1 as

$$\Phi(x, t+s) \leq \Phi(x, 2t) + \Phi(x, 2s) \leq 2^{1+s(\Phi)} (\Phi(x, t) + \Phi(x, s)) \quad (2.6)$$

for every $x \in \Omega$ and $t, s \geq 0$. Furthermore, for any fixed $x \in \Omega$, we have

$$\partial_t \Phi(x, t) \leq (1+s(\Phi)) \frac{\Phi(x, t)}{t} \leq (1+s(\Phi)) [\Phi(x, 1)]^{\frac{s(\Phi)}{1+s(\Phi)}} [\Phi(x, t)]^{\frac{1}{1+s(\Phi)}} \quad \text{for every } 0 \leq t \leq 1$$

and

$$\partial_t \Phi(x, t) \leq (1+s(\Phi)) \frac{\Phi(x, t)}{t} \leq (1+s(\Phi)) [\Phi(x, 1)]^{\frac{1}{1+s(\Phi)}} [\Phi(x, t)]^{\frac{s(\Phi)}{1+s(\Phi)}} \quad \text{for every } t \geq 1.$$

Putting together the last two inequalities, we have the following very useful inequality which will be applied in the future

$$\frac{\Phi(x, t)}{t} \approx \partial_t \Phi(x, t) \leq (1+s(\Phi)) \left([\Phi(x, 1)]^{\frac{s(\Phi)}{1+s(\Phi)}} [\Phi(x, t)]^{\frac{1}{1+s(\Phi)}} + [\Phi(x, 1)]^{\frac{1}{1+s(\Phi)}} [\Phi(x, t)]^{\frac{s(\Phi)}{1+s(\Phi)}} \right) \quad (2.7)$$

for every $x \in \Omega$ and $t \geq 0$.

Lemma 2.2. *Let $\Phi, \tilde{\Phi} \in \mathcal{N}(\Omega)$ with indices $s(\Phi), s(\tilde{\Phi}) \geq 1$. Then,*

1. *For any non-negative real numbers a, b satisfying $a + b > 0$, $a\Phi + b\tilde{\Phi} \in \mathcal{N}(\Omega)$ with $s(a\Phi + b\tilde{\Phi}) = s(\Phi) + s(\tilde{\Phi})$ and $\Phi\tilde{\Phi} \in \mathcal{N}(\Omega)$ with $s(\Phi\tilde{\Phi}) = 4s(\Phi)s(\tilde{\Phi})(s(\Phi) + s(\tilde{\Phi}))$.*
2. *For any number $m \geq 1$, $\Phi^m \in \mathcal{N}(\Omega)$ with $s(\Phi^m) = s(\Phi) + (m-1)(s(\Phi) + 1)$.*
3. *For any number $\mu \geq 0$, $\Phi_\mu(x, t) := t^\mu \Phi(x, t) \in \mathcal{N}(\Omega)$ with $s(\Phi_\mu) = \mu + 3[s(\Phi)]^2$.*
4. *There exists $\theta_0 \in (0, 1)$ depending only on $s(\Phi)$ such that $\Phi^\theta \in \mathcal{N}(\Omega)$ for every $\theta \in (\theta_0, 1]$ with $s(\Phi^\theta)$ depending only on $s(\Phi)$ and θ .*

Lemma 2.3. *Let $\Phi \in \mathcal{N}$ with an index $s(\Phi) \geq 1$. Then $t \mapsto \Phi\left(t^{\frac{1}{s(\Phi)+1}}\right)$ is a concave function.*

Lemma 2.4. *Let $\Phi \in \mathcal{N}(\Omega)$ with an index $s(\Phi) \geq 1$. Then there exists a positive constant $c \equiv c(s(\Phi))$ such that*

$$s_1 \partial_t \Phi(x, s_2) + s_2 \partial_t \Phi(x, s_1) \leq \varepsilon \Phi(x, s_1) + \frac{c}{\varepsilon^{s(\Phi)}} \Phi(x, s_2)$$

holds for all $s_1, s_2 \geq 0$ and $0 < \varepsilon \leq 1$.

Remark 2.2. We note that $\Psi \in \mathcal{N}(\Omega)$ with an index $s(\Psi) = s(G) + s(H_a) + s(H_b)$ by Lemma 2.2. In particular, for every open subset $\mathcal{B} \subset \Omega$, it holds that $\Psi_{\mathcal{B}}^+, \Psi_{\mathcal{B}}^- \in \mathcal{N}$ with indices $s(\Psi_{\mathcal{B}}^+) = s(G) + s(H_a) + s(H_b)$ and $s(\Psi_{\mathcal{B}}^-) = s(G) + s(H_a) + s(H_b)$.

For a given Young function $\Phi \in \mathcal{N}(\Omega)$ with an index $s(\Phi) \geq 1$, we define the vector field $V_\Phi : \Omega \times \mathbb{R}^n \setminus \{0\} \rightarrow \mathbb{R}^n$ as follows

$$V_\Phi(x, z) := \left[\frac{\partial_t \Phi(x, |z|)}{|z|} \right]^{\frac{1}{2}} z. \quad (2.8)$$

Furthermore, we shall often use the following inequalities that

$$\int_0^1 \frac{\Phi(x, |\theta z_1 + (1-\theta)z_2|)}{|\theta z_1 + (1-\theta)z_2|^2} d\theta \approx \frac{\Phi(x, |z_1| + |z_2|)}{(|z_1| + |z_2|)^2}, \quad (2.9)$$

$$|V_\Phi(x, z_1) - V_\Phi(x, z_2)|^2 \approx \partial_{tt}^2 \Phi(x, |z_1| + |z_2|) |z_1 - z_2|^2 \approx \frac{\partial_t \Phi(x, |z_1| + |z_2|)}{|z_1| + |z_2|} |z_1 - z_2|^2, \quad (2.10)$$

$$\Phi(x, |z_1 - z_2|) \lesssim \Phi(x, |z_1| + |z_2|) \frac{|z_1 - z_2|}{|z_1| + |z_2|} \quad (2.11)$$

and

$$\left\langle \partial_t \Phi(x, |z_1|) \frac{z_1}{|z_1|} - \partial_t \Phi(x, |z_2|) \frac{z_2}{|z_2|}, z_1 - z_2 \right\rangle \approx |V_\Phi(x, z_1) - V_\Phi(x, z_2)|^2 \quad (2.12)$$

hold true, whenever $x \in \Omega$ and $z_1, z_2 \in \mathbb{R}^n \setminus \{0\}$, where all implied constants in (2.9)-(2.12) depend on n and $s(\Phi)$ (see [41] for further discussions). Moreover, we have the following useful inequality

$$|V_\Phi(x, z_2) - V_\Phi(x, z_1)|^2 \lesssim \int_0^1 |V_\Phi(x, \theta z_2 + (1-\theta)z_1) - V_\Phi(x, z_1)|^2 \frac{d\theta}{\theta} \quad (\forall x \in \Omega), \quad (2.13)$$

which follows from the following estimates that

$$\begin{aligned}
& \int_0^1 |V_\Phi(x, \theta z_2 + (1-\theta)z_1) - V_\Phi(x, z_1)|^2 \frac{d\theta}{\theta} \\
& \stackrel{(2.10)}{\gtrsim} \int_0^1 \frac{\Phi(x, |\theta z_2 + (1-\theta)z_1| + |z_1|)}{(|\theta z_2 + (1-\theta)z_1| + |z_1|)^2} \theta |z_2 - z_1|^2 d\theta \\
& \gtrsim \frac{|z_2 - z_1|^2}{(|z_2| + |z_1|)^2} \int_0^1 \Phi(x, |\theta z_2 + (1-\theta)z_1| + |z_1|) \theta d\theta \\
& \gtrsim \frac{|z_2 - z_1|^2}{(|z_2| + |z_1|)^2} \Phi \left(x, \int_0^1 (|\theta z_2 + (1-\theta)z_1| + |z_1|) \theta d\theta \right) \\
& \gtrsim \frac{|z_2 - z_1|^2}{(|z_2| + |z_1|)^2} \Phi(x, |z_2| + |z_1|) \stackrel{(2.10)}{\approx} |V_\Phi(x, z_2) - V_\Phi(x, z_1)|^2
\end{aligned}$$

hold with having all implied constants in the above display depending on n and $s(\Phi)$, whenever $x \in \Omega$ and $z_1, z_2 \in \mathbb{R}^n \setminus \{0\}$, where in the third inequality of the last display we have applied Jensen's inequality to the convex function $\Phi(x, \cdot)$ with respect to measure $\theta d\theta$.

Moreover, the maps introduced in (2.8) are very convenient to formulate the monotonicity properties of the vector field $D_z F(x, y, z)$ with respect to the gradient variable z and some growth properties of the integrand F defined in (1.9).

Lemma 2.5. *Let $F : \Omega \times \mathbb{R} \times \mathbb{R}^n \rightarrow \mathbb{R}$ be a function defined in (1.15) satisfying (1.10) and (1.16). Then there exist positive constants $c_1, c_2 \equiv c_1, c_2(n, s(G), s(H_a), s(H_b), \nu)$ and $c_3 \equiv c_3(n, s(G), s(H_a), s(H_b), L)$ such that the following inequalities*

$$\begin{aligned}
& |V_G(z_1) - V_G(z_2)|^2 + a(x)|V_{H_a}(z_1) - V_{H_a}(z_2)|^2 + b(x)|V_{H_b}(z_1) - V_{H_b}(z_2)|^2 \\
& \leq c_1 \langle D_z F(x, y, z_1) - D_z F(x, y, z_2), z_1 - z_2 \rangle,
\end{aligned} \tag{2.14}$$

$$\begin{aligned}
& |V_G(z_1) - V_G(z_2)|^2 + a(x)|V_{H_a}(z_1) - V_{H_a}(z_2)|^2 + b(x)|V_{H_b}(z_1) - V_{H_b}(z_2)|^2 \\
& + c_2 \langle D_z F(x, y, z_1), z_2 - z_1 \rangle \leq c_2 [F(x, y, z_2) - F(x, y, z_1)]
\end{aligned} \tag{2.15}$$

and

$$\begin{aligned}
& |F(x_1, y, z) - F(x_2, y, z)| \\
& \leq c_3 \omega(|x_1 - x_2|) [G(|z|) + \min\{a(x_1), a(x_2)\} H_a(|z|) + \min\{b(x_1), b(x_2)\} H_b(|z|)] \\
& + c_3 |a(x_1) - a(x_2)| H_a(|z|) + c_3 |b(x_1) - b(x_2)| H_b(|z|)
\end{aligned} \tag{2.16}$$

hold true, whenever $z, z_1, z_2 \in \mathbb{R}^n \setminus \{0\}$, $x, x_1, x_2 \in \Omega$ and $y \in \mathbb{R}$.

Proof. It follows from (1.16)₂ that

$$\begin{aligned}
& \langle D_z F(x, y, z_1) - D_z F(x, y, z_2), z_1 - z_2 \rangle = \int_0^1 \langle D_{zz}^2 F(x, y, \theta z_1 + (1-\theta)z_2) [z_1 - z_2], z_1 - z_2 \rangle d\theta \\
& \geq \nu \int_0^1 \frac{\Psi(x, \theta z_1 + (1-\theta)z_2)}{|\theta z_1 + (1-\theta)z_2|^2} |z_1 - z_2|^2 d\theta \\
& \geq c (|V_G(z_1) - V_G(z_2)|^2 + a(x)|V_{H_a}(z_1) - V_{H_a}(z_2)|^2 + b(x)|V_{H_b}(z_1) - V_{H_b}(z_2)|^2),
\end{aligned}$$

where in the last inequality of the last display we have used (2.9) and (2.10). Then (2.14) follows. The inequality (2.15) follows from the following observation that

$$\begin{aligned}
& [F(x, y, z_2) - F(x, y, z_1)] - \langle D_z F(x, y, z_1), z_2 - z_1 \rangle \\
&= \int_0^1 \langle D_z F(x, y, \theta z_2 + (1 - \theta)z_1) - D_z F(x, y, z_1), z_2 - z_1 \rangle d\theta \\
&\stackrel{(2.14)}{\geq} c \int_0^1 \frac{1}{\theta} (|V_G(\theta z_2 + (1 - \theta)z_1) - V_G(z_1)|^2 + a(x)|V_{H_a}(\theta z_2 + (1 - \theta)z_1) - V_{H_a}(z_1)|^2) d\theta \\
&\quad + \int_0^1 \frac{b(x)}{\theta} |V_{H_b}(\theta z_2 + (1 - \theta)z_1) - V_{H_b}(z_1)|^2 d\theta \\
&\stackrel{(2.13)}{\geq} c|V_G(z_1) - V_G(z_2)|^2 + a(x)|V_{H_a}(z_1) - V_{H_a}(z_2)|^2 + b(x)|V_{H_b}(z_1) - V_{H_b}(z_2)|^2.
\end{aligned}$$

Since $F(x, y, 0) = 0$ for every $x \in \Omega$ and $y \in \mathbb{R}$, using (1.15), we have

$$\begin{aligned}
& |F(x_1, y, z) - F(x_2, y, z)| = |(F(x_1, y, z) - F(x_1, y, 0)) - (F(x_2, y, z) - F(x_2, y, 0))| \\
&= \left| \int_0^1 \langle D_z F(x_1, y, \theta z), z \rangle d\theta - \int_0^1 \langle D_z F(x_2, y, \theta z), z \rangle d\theta \right| \\
&\leq \int_0^1 |D_z F(x_1, y, \theta z) - D_z F(x_2, y, \theta z)| |z| d\theta \\
&\leq \int_0^1 |D_z F_G(x_1, y, \theta z) - D_z F_G(x_2, y, \theta z)| |z| d\theta \\
&\quad + \int_0^1 |a(x_1)D_z F_{H_a}(x_1, y, \theta z) - a(x_2)D_z F_{H_a}(x_2, y, \theta z)| |z| d\theta \\
&\quad + \int_0^1 |b(x_1)D_z F_{H_b}(x_1, y, \theta z) - b(x_2)D_z F_{H_b}(x_2, y, \theta z)| |z| d\theta.
\end{aligned}$$

Without loss of generality, we can assume $a(x_2) \leq a(x_1)$ and $b(x_2) \leq b(x_1)$. Then using the structure assumption (1.16), we find

$$\begin{aligned}
& \int_0^1 |a(x_1)D_z F_{H_a}(x_1, y, \theta z) - a(x_2)D_z F_{H_a}(x_2, y, \theta z)| |z| d\theta \\
&\leq L|a(x_1) - a(x_2)| \int_0^1 \frac{H_a(\theta|z|)}{\theta} d\theta + a(x_2)\omega(|x_1 - x_2|) \int_0^1 \frac{H_a(\theta|z|)}{\theta} d\theta \\
&\leq ca(x_2)H_a(|z|) + c\omega(|x_1 - x_2|)H_a(|z|)
\end{aligned}$$

for some constant $c \equiv c(s(H_a), L)$. Similarly, we get

$$\begin{aligned}
& \int_0^1 |b(x_1)D_z F_{H_b}(x_1, y, \theta z) - b(x_2)D_z F_{H_b}(x_2, y, \theta z)| |z| d\theta \\
&\leq cb(x_2)H_b(|z|) + c\omega(|x_1 - x_2|)H_b(|z|),
\end{aligned}$$

where the validity of the last display is ensured by (1.16)₃. Combining the last three displays, (2.16) follows. \square

2.2. Musielak-Orlicz and Musielak-Orlicz-Sobolev spaces. We now introduce the Musielak-Orlicz spaces (generalized Orlicz spaces), which generalize the Orlicz spaces. Let $\Phi : \Omega \times [0, \infty) \rightarrow [0, \infty)$ be an Young function. Here we present some definitions and properties associated to Young functions.

Definition 2.2. Let Φ be a Young function.

- (1) Φ is said to satisfy the Δ_2 -condition, denoted by $\Phi \in \Delta_2$, if there exists a positive number $\Delta_2(\Phi)$ such that $\Phi(x, 2t) \leq \Delta_2(\Phi) \Phi(x, t)$ for all $x \in \Omega$ and $t \geq 0$.
- (2) Φ is said to satisfy the ∇_2 -condition, denoted by $\Phi \in \nabla_2$, if there exists a positive number $\nabla_2(\Phi) > 1$ such that $\Phi(x, \nabla_2(\Phi)t) \geq 2\nabla_2(\Phi) \Phi(x, t)$ for all $x \in \Omega$ and $t \geq 0$.
- (3) We write $\Phi \in \Delta_2 \cap \nabla_2$ if $\Phi \in \Delta_2$ and $\Phi \in \nabla_2$.

For a given Young function Φ , we define the complementary function Φ^* of Φ by, for each $x \in \Omega$ and $t \geq 0$,

$$\Phi^*(x, t) = \sup\{st - \Phi(x, s) : s \geq 0\}.$$

Then Φ^* satisfies all the conditions to be a Young function. One can see that $(\Phi^*)^* = \Phi$ and that $\Phi \in \nabla_2$ if and only if $\Phi^* \in \Delta_2$ with $2\nabla_2(\Phi) = \Delta_2(\Phi^*)$.

For an Young function Φ , the Musielak-Orlicz class $K^\Phi(\Omega; \mathbb{R}^N)$, $N \geq 1$, consists of all measurable functions $v : \Omega \rightarrow \mathbb{R}^N$ satisfying

$$\int_{\Omega} \Phi(x, |v(x)|) dx < +\infty.$$

The Musielak-Orlicz space $L^\Phi(\Omega; \mathbb{R}^N)$ is the vector space generated by $K^\Phi(\Omega; \mathbb{R}^N)$. If $\Phi \in \Delta_2$, then $K^\Phi(\Omega; \mathbb{R}^N) = L^\Phi(\Omega; \mathbb{R}^N)$ and this space is a Banach space under the Luxemburg norm

$$\|v\|_{L^\Phi(\Omega; \mathbb{R}^N)} = \inf \left\{ \sigma > 0 : \int_{\Omega} \Phi \left(x, \frac{|v(x)|}{\sigma} \right) dx \leq 1 \right\}.$$

The Musielak-Orlicz-Sobolev space $W^{1,\Phi}(\Omega; \mathbb{R}^N)$ is the function space of all measurable functions $v \in L^\Phi(\Omega; \mathbb{R}^N)$ such that its distributional gradient vector Dv belongs to $L^\Phi(\Omega; \mathbb{R}^{Nn})$. For $v \in W^{1,\Phi}(\Omega; \mathbb{R}^N)$, we define its norm to be

$$\|v\|_{W^{1,\Phi}(\Omega; \mathbb{R}^N)} = \|v\|_{L^\Phi(\Omega; \mathbb{R}^N)} + \|Dv\|_{L^\Phi(\Omega; \mathbb{R}^{Nn})}.$$

The space $W_0^{1,\Phi}(\Omega; \mathbb{R}^N)$ is defined as the closure of $C_0^\infty(\Omega; \mathbb{R}^N)$ in $W^{1,\Phi}(\Omega; \mathbb{R}^N)$. For $N = 1$, we simply write $L^\Phi(\Omega) := L^\Phi(\Omega; \mathbb{R})$ and $W^{1,\Phi}(\Omega) := W^{1,\Phi}(\Omega; \mathbb{R})$. For a detailed discussion of the Musielak-Orlicz spaces and the associated Sobolev spaces, we refer the reader to [2, 15, 29, 40, 55, 56, 72, 79, 82] and references therein.

We end up this preliminary section with presenting some standard technical lemmas which will be applied later, see for instance [53, 54, 63].

Lemma 2.6. Let $h : [\rho_1, \rho_2] \rightarrow \mathbb{R}$ be a non-negative and bounded function, and $\theta \in (0, 1)$, $A_0 \geq 0$, $\gamma_0 \geq 0$. Assume that

$$h(t) \leq \theta h(s) + \frac{A_0}{(s-t)^{\gamma_0}}$$

holds for $\rho_1 \leq t < s \leq \rho_2$. Then there is a constant $c \equiv c(\theta, \gamma_0)$ satisfying the following inequality:

$$h(\rho_0) \leq \frac{cA_0}{(\rho_1 - \rho_0)^{\gamma_0}}.$$

Lemma 2.7. Let $\{Y_i\}_{i=0}^\infty$ be a sequence of nonnegative numbers satisfying the following recursive inequalities

$$Y_{i+1} \leq Cb^i Y_i^{1+\tau_0}$$

with some fixed positive constant C , $b > 1$ and $\tau_0 > 0$ for every $i = 0, 1, 2, \dots$. If

$$Y_0 \leq C^{-\frac{1}{\tau_0}} b^{-\frac{1}{\tau_0}},$$

then $Y_i \rightarrow 0$ as $i \rightarrow \infty$.

Lemma 2.8. *Let $v \in W^{1,1}(B_\rho)$ for some ball $B_\rho \subset \mathbb{R}^n$. Then there exists $c \equiv c(n)$ such that*

$$(l-k)|B_\rho \cap \{v > l\}|^{1-\frac{1}{n}} \leq \frac{c|B_\rho|}{|B_\rho \setminus \{v > k\}|} \int_{B_\rho \cap \{k < v \leq l\}} |Dv| dx$$

holds, whenever l and k are real numbers with $l > k$.

3. ABSENCE OF LAVRENTIEV PHENOMENON

Here we deal with the absence of Lavrantiev phenomenon under the assumptions introduced in (1.12)₂, (1.13)₂ and (1.14)₂. The following theorem widely covers the results of [30, Theorem 4.1], [31, Proposition 3.6], [22, Theorem 3.1], [13, Theorem 4], [6, Theorem 4.1] and [7, Theorem 3.1].

Theorem 3.1. *Let \mathcal{P} be the functional defined in (1.1) with the coefficient functions $a(\cdot) \in C^{\omega_a}(\Omega)$ and $b(\cdot) \in C^{\omega_b}(\Omega)$ for the functions ω_a, ω_b being concave and vanishing at 0.*

1. *If the condition (1.12)₂ is satisfied, then for any function $v \in W_{\text{loc}}^{1,\Psi}(\Omega)$ and ball $B_R \equiv B_R(x_0) \Subset \tilde{B} \Subset \Omega$ with $\mathcal{P}(v, \tilde{B}) < \infty$, there exists a sequence of functions $\{v_k\}_{k=1}^\infty \subset W^{1,\infty}(B_R)$ such that*

$$v_k \rightarrow v \quad \text{in } W^{1,G}(B_R) \quad \text{and} \quad \mathcal{P}(v_k, B_R) \rightarrow \mathcal{P}(v, B_R). \quad (3.1)$$

2. *If the condition (1.13)₂ is satisfied, then for any function $v \in W_{\text{loc}}^{1,\Psi}(\Omega) \cap L_{\text{loc}}^\infty(\Omega)$ and ball $B_R \equiv B_R(x_0) \Subset \tilde{B} \Subset \Omega$ with $\mathcal{P}(v, \tilde{B}) < \infty$, there exists a sequence of functions $\{v_k\}_{k=1}^\infty \subset W^{1,\infty}(B_R)$ such that*

$$v_k \rightarrow v \quad \text{in } W^{1,G}(B_R), \quad \mathcal{P}(v_k, B_R) \rightarrow \mathcal{P}(v, B_R) \quad \text{and} \quad \limsup_{k \rightarrow \infty} \|v_k\|_{L^\infty(B_R)} \leq \|v\|_{L^\infty(B_R)}. \quad (3.2)$$

3. *Let $v \in W^{1,\Psi}(\Omega) \in C^{0,\gamma}(\Omega)$ with some $\gamma \in (0,1)$ be a local Q -minimizer of the functional \mathcal{P} under the assumption (1.14)₂. Then, for every ball $B_R \Subset \Omega$, there exists a sequence of functions $\{v_k\}_{k=1}^\infty \subset W^{1,\infty}(B_R)$ such that*

$$v_k \rightarrow v \quad \text{in } W^{1,G}(B_R) \quad \text{and} \quad \mathcal{P}(v_k, B_R) \rightarrow \mathcal{P}(v, B_R). \quad (3.3)$$

Proof. Essentially, the proof for the first two parts is similar to the one of [22, Theorem 3.1]. Since our assumptions are weaker than the assumptions considered there, we provide the detailed proof in any case. First we fix $\varepsilon_0 \in (0,1)$ such that $B_R \Subset B_{R+\varepsilon_0} \Subset \tilde{B} \Subset \Omega$. Let $\rho \in C_0^\infty(B_1)$ be a non-negative standard mollifier with $\int_{\mathbb{R}^n} \rho dx = 1$. Then we set $\rho_\varepsilon(x) := \frac{1}{\varepsilon^n} \rho\left(\frac{x}{\varepsilon}\right)$ for $x \in B_\varepsilon$ with $0 < \varepsilon < \varepsilon_0$. Clearly $\rho_\varepsilon \in C_0^\infty(B_\varepsilon)$, $\int_{\mathbb{R}^n} \rho_\varepsilon dx = 1$, $0 \leq \rho_\varepsilon \leq c(n)\varepsilon^{-n}$ and $|D\rho_\varepsilon| \leq c(n)\varepsilon^{-(n+1)}$. For every $0 < \varepsilon < \varepsilon_0/2$, we consider the following functions:

$$v_\varepsilon(x) := (v * \rho_\varepsilon)(x), \quad a_\varepsilon(x) := \inf_{y \in B_{2\varepsilon}(x)} a(y), \quad b_\varepsilon(x) := \inf_{y \in B_{2\varepsilon}(x)} b(y) \quad (3.4)$$

and

$$\Psi_\varepsilon(x, t) := G(t) + a_\varepsilon(x)H_a(t) + b_\varepsilon(x)H_b(t) \quad (3.5)$$

for every $x \in B_R$ and $t \geq 0$.

1. By Jensen's inequality, for a fixed $x \in B_R$, we have

$$G(|Dv_\varepsilon(x)|) = G(|(Dv * \rho_\varepsilon)(x)|) \leq \int_{\mathbb{R}^n} G(|Dv(x-y)|) \rho_\varepsilon(y) dy \leq c\varepsilon^{-n}.$$

It follows from (1.12)₂ and the last display that

$$\begin{aligned} H_a(|Dv_\varepsilon(x)|) &= \frac{(H_a \circ G^{-1})(G(|Dv_\varepsilon(x)|))}{G(|Dv_\varepsilon(x)|)} G(|Dv_\varepsilon(x)|) \\ &\leq \lambda_1 \left(1 + \left[\omega_a \left([G(|Dv_\varepsilon(x)|)]^{-\frac{1}{n}} \right) \right]^{-1} \right) G(|Dv_\varepsilon(x)|) \\ &\leq c \left(1 + [\omega_a(\varepsilon)]^{-1} \right) G(|Dv_\varepsilon(x)|) \leq c \left(1 + [\omega_a(\varepsilon)]^{-1} \right) \Psi_\varepsilon(x, |Dv_\varepsilon(x)|). \end{aligned} \quad (3.6)$$

Similarly as above, we have

$$H_b(|Dv_\varepsilon(x)|) \leq c(1 + [\omega_b(\varepsilon)]^{-1}) \Psi_\varepsilon(x, |Dv_\varepsilon(x)|). \quad (3.7)$$

2. Since v is locally bounded in Ω , we have

$$|Dv_\varepsilon(x)| = |(v * D\rho_\varepsilon)(x)| \leq \int_{\mathbb{R}^n} |v(x-y)| |D\rho_\varepsilon(y)| dy \leq c(n) \|v\|_{L^\infty(\tilde{B})} \varepsilon^{-1}.$$

Then the assumption (1.13)₂ and the last display imply

$$\begin{aligned} H_a(|Dv_\varepsilon(x)|) &= \frac{H_a(|Dv_\varepsilon(x)|)}{G(|Dv_\varepsilon(x)|)} G(|Dv_\varepsilon(x)|) \leq \lambda_2 \left(1 + [\omega_a(|Dv_\varepsilon(x)|^{-1})]^{-1}\right) G(|Dv_\varepsilon(x)|) \\ &\leq c(1 + [\omega_a(\varepsilon)]^{-1}) G(|Dv_\varepsilon(x)|) \leq c(1 + [\omega_a(\varepsilon)]^{-1}) \Psi_\varepsilon(x, |Dv_\varepsilon(x)|) \end{aligned} \quad (3.8)$$

with some constant $c \equiv c(n, \lambda_2, \|v\|_{L^\infty(\tilde{B})})$ for every $x \in B_R$. Arguing in the same way, for every $x \in B_R$, we have

$$H_b(|Dv_\varepsilon(x)|) \leq c(1 + [\omega_b(\varepsilon)]^{-1}) \Psi_\varepsilon(x, |Dv_\varepsilon(x)|). \quad (3.9)$$

Using the continuity of the coefficient functions $a(\cdot)$ and $b(\cdot)$ and recalling the very definition of Ψ_ε in (3.6), for every $x \in B_R$, we have

$$\begin{aligned} \Psi(x, |Dv_\varepsilon(x)|) &\leq \Psi_\varepsilon(x, |Dv_\varepsilon(x)|) + |a(x) - a_\varepsilon(x)| H_a(|Df_\varepsilon(x)|) + |b(x) - b_\varepsilon(x)| H_b(|Df_\varepsilon(x)|) \\ &\leq \Psi_\varepsilon(x, |Dv_\varepsilon(x)|) + 4[a]_{\omega_a} \omega_a(\varepsilon) H_a(|Dv_\varepsilon(x)|) + 4[b]_{\omega_b} \omega_b(\varepsilon) H_b(|Dv_\varepsilon(x)|). \end{aligned} \quad (3.10)$$

Therefore, taking into account (3.6)-(3.7) when the first case comes into play, and (3.8)-(3.9) when the second case is considered, in any case, it follows from (3.10) that

$$\begin{aligned} \Psi(x, |Dv_\varepsilon(x)|) &\leq c\Psi_\varepsilon(x, |Dv_\varepsilon(x)|) + c\omega_a(\varepsilon)(1 + [\omega_a(\varepsilon)]^{-1})\Psi_\varepsilon(x, |Dv_\varepsilon(x)|) \\ &\quad c\omega_b(\varepsilon)(1 + [\omega_b(\varepsilon)]^{-1})\Psi_\varepsilon(x, |Dv_\varepsilon(x)|) \leq c\Psi_\varepsilon(x, |Dv_\varepsilon(x)|) \end{aligned} \quad (3.11)$$

for some constant c being independent of ε . Therefore, by Jensen's inequality, we get

$$\begin{aligned} \Psi_\varepsilon(x, |Dv_\varepsilon(x)|) &\leq \int_{B_\varepsilon(x)} \Psi_\varepsilon(x, |Dv(y)|) \rho_\varepsilon(x-y) dy \leq \int_{B_\varepsilon(x)} \Psi(y, |Dv(y)|) \rho_\varepsilon(x-y) dy \\ &= [\Psi(\cdot, |Dv(\cdot)|) * \rho_\varepsilon](x) =: [\Psi(\cdot, |Dv(\cdot)|)]_\varepsilon(x). \end{aligned} \quad (3.12)$$

Hence, in any case, using (3.11)-(3.12), we conclude that

$$\Psi(x, |Dv_\varepsilon(x)|) \leq c[\Psi(\cdot, |Dv(\cdot)|)]_\varepsilon(x) \quad (3.13)$$

holds every $x \in B_R$ with a constant c independent of ε . Since $[\Psi(\cdot, |Dv(\cdot)|)]_\varepsilon \rightarrow \Psi(\cdot, |Dv(\cdot)|)$ strongly in $L^1(B_R)$, we are able to apply the general Lebesgue's dominated convergence theorem of [78, Theorem 19] to obtain a sequence of functions $\{v_k\} := \{v_{\varepsilon_k}\} \subset C_0^\infty(\tilde{B})$ satisfying (3.1) for the first case and (3.2)_{1,2} for the second case with some suitable choice of $\varepsilon_k \rightarrow 0$. Clearly, the assertion (3.2)₃ comes from the very definition of mollification of v defined in (3.4).

3. Now we turn our attention to proving the last part of the theorem. Applying a Caccioppoli type inequality of Lemma 5.2 under the assumption (1.14)₂ below, we see that

$$\int_{B_\varepsilon(x)} \Psi_\varepsilon(x, |Dv(z)|) dz \leq c \int_{B_{2\varepsilon}(x)} \Psi_\varepsilon\left(x, \left|\frac{v(z) - (v)_{B_{2\varepsilon}(x)}}{\varepsilon}\right|\right) dz \quad (3.14)$$

for a constant c independent of ε . Therefore, by the very definition of the convolution, the fact that $\Psi_\varepsilon(x, \cdot)$ is convex for any fixed $x \in B_R$ and (3.14), we have

$$\begin{aligned} |Dv_\varepsilon(x)| &\leq c(\Psi_\varepsilon(x, \cdot))_t^{-1} \circ \Psi_\varepsilon \left(x, \int_{B_\varepsilon(x)} |Dv(z)| dz \right) \\ &\leq c(\Psi_\varepsilon(x, \cdot))_t^{-1} \left(\int_{B_\varepsilon(x)} \Psi_\varepsilon(x, |Dv(z)|) dz \right) \\ &\leq c(\Psi_\varepsilon(x, \cdot))_t^{-1} \left(\int_{B_{2\varepsilon}(x)} \Psi_\varepsilon \left(x, \left| \frac{v(z) - (v)_{B_{2\varepsilon}(x)}}{\varepsilon} \right| \right) dz \right) \leq c\varepsilon^{\gamma-1} \end{aligned} \quad (3.15)$$

with some constant c independent of ε , whenever $x \in B_R$ and $\varepsilon \in (0, \varepsilon_0/4)$, where we have also used the assumption $v \in C^{0,\gamma}(\Omega)$ for some $\gamma \in (0, 1)$ and Lemma 2.1 together with Remark 2.2. Recalling the definition of Ψ_ε in (3.5), using the modulus of continuity of functions $a(\cdot)$, $b(\cdot)$ and the assumption (1.14)₂, for every $x \in B_R$, we estimate

$$\begin{aligned} \Psi(x, |Dv_\varepsilon(x)|) &\leq \Psi_\varepsilon(x, |Dv_\varepsilon(x)|) + |a_\varepsilon(x) - a(x)|H_a(|Dv_\varepsilon(x)|) + |b_\varepsilon(x) - b(x)|H_b(|Dv_\varepsilon(x)|) \\ &\leq \Psi_\varepsilon(x, |Dv_\varepsilon(x)|) + c\omega_a(\varepsilon) \left(1 + \left[\omega_a(|Dv_\varepsilon(x)|^{-\frac{1}{1-\gamma}}) \right]^{-1} \right) G(|Dv_\varepsilon(x)|) \\ &\quad + c\omega_b(\varepsilon) \left(1 + \left[\omega_b(|Dv_\varepsilon(x)|^{-\frac{1}{1-\gamma}}) \right]^{-1} \right) G(|Dv_\varepsilon(x)|) \\ &\leq c\Psi_\varepsilon(x, |Dv_\varepsilon(x)|) \end{aligned}$$

for a constant c independent of ε , where we have also used (3.15). Then arguing in the same way as in (3.12)-(3.13), we find a sequence of functions $\{v_k\}_{k=1}^\infty \subset W^{1,\infty}(B_R)$ satisfying (3.3). The proof is now finished. \square

4. SOBOLEV-POINCARÉ TYPE INEQUALITIES

In the present section we provide a Sobolev-Poincaré type inequality for functions $v \in W^{1,\Psi}(B_R)$ with some ball $B_R \subset \Omega$, which is one of key points for further investigations. For this, first we give a Sobolev-Poincaré type inequality for functions of $W^{1,\Phi}(B_R)$ with $\Phi \in \mathcal{N}$ and a ball $B_R \subset \mathbb{R}^n$.

Lemma 4.1. *Let $\Phi \in \mathcal{N}$ with an index $s(\Phi) \geq 1$. For any $d_0 \in [1, \frac{n}{n-1})$, there exists $\theta \equiv \theta(n, s(\Phi), d_0) \in (0, 1)$ such that*

$$\left(\int_{B_R} \left[\Phi \left(\left| \frac{v - (v)_{B_R}}{R} \right| \right) \right]^{d_0} dx \right)^{\frac{1}{d_0}} \leq c \left(\int_{B_R} [\Phi(|Dv|)]^\theta dx \right)^{\frac{1}{\theta}} \quad (4.1)$$

holds for some constant $c \equiv c(n, s(\Phi), d_0)$, whenever $v \in W^{1,\Phi}(B_R)$ and $B_R \subset \mathbb{R}^n$ is a ball. Moreover, the above estimate still holds with $v - (v)_{B_R}$ replaced by v if $v \in W_0^{1,\Phi}(B_R)$.

Proof. First note by Lemma 2.2₄ that there exists $\theta \equiv \theta(n, s(\Phi), d_0) \in \left(\frac{(n-1)d_0}{n}, 1 \right)$ such that $\Phi^\theta \in \mathcal{N}$ with an index $s(\Phi^\theta)$ depending on $n, s(\Phi), d_0$. Therefore, the following classical formula

$$|v(x) - (v)_{B_R}| \leq c(n) \int_{B_R} \frac{|Dv(y)|}{|x - y|^{n-1}} dy \quad (4.2)$$

holds for a.e $x \in B_R$, see for instance [54, Lemma 7.14]. Letting $E := \int_{B_R} \Phi^\theta(|Dv|) dx$, one can assume that $E > 0$, otherwise v is constant on B_R and the inequality (4.1) is trivial. Using (4.2), the fact that Φ is

increasing and Lemma 2.1, we have

$$I := \int_{B_R} \left[\Phi \left(\left| \frac{v - (v)_{B_R}}{R} \right| \right) \right]^{d_0} dx \leq c \int_{B_R} \left[\Phi \left(\int_{B_R} \frac{|Dv(y)|}{R|x-y|^{n-1}} dy \right) \right]^{d_0} dx$$

with $c \equiv c(n, s(\Phi), d_0)$. Since $\int_{B_R} \frac{1}{R|x-y|^{n-1}} dy < c(n)$, where this constant is independent of $x \in B_R$ and a ball B_R , we apply Jensen's inequality to the convex function Φ^θ with respect to the measure $R^{-1}|x-y|^{-(n-1)} dy$ to obtain

$$\begin{aligned} I &\leq c \int_{B_R} \left(\int_{B_R} \frac{[\Phi(|Dv(y)|)]^\theta}{R|x-y|^{n-1}} dy \right)^{\frac{d_0}{\theta}} dx = cR^{\frac{(n-1)d_0}{\theta}} E^{\frac{d_0}{\theta}} \int_{B_R} \left(\int_{B_R} \frac{[\Phi(|Dv(y)|)]^\theta}{|x-y|^{n-1}} E^{-1} dy \right)^{\frac{d_0}{\theta}} dx \\ &\leq cR^{\frac{(n-1)d_0}{\theta}} E^{\frac{d_0}{\theta}} \int_{B_R} \int_{B_R} \frac{[\Phi(|Dv(y)|)]^\theta}{|x-y|^{\frac{(n-1)d_0}{\theta}}} E^{-1} dy dx, \end{aligned} \quad (4.3)$$

where in the last estimate we have applied again Jensen's inequality to the convex function $t \mapsto t^{\frac{d_0}{\theta}}$ with respect to the probability measure $E^{-1}\Phi^\theta(|Dv(y)|) dy$. We observe that

$$\int_{B_R} \frac{1}{|x-y|^{\frac{(n-1)d_0}{\theta}}} dx \leq \frac{1}{|B_R|} \int_{B_{2R}(y)} \frac{1}{|x-y|^{\frac{(n-1)d_0}{\theta}}} dx \leq c(n, s(\Phi), d) R^{-\frac{(n-1)d_0}{\theta}},$$

which is possible since $\frac{(n-1)d_0}{\theta} < n$. Inserting the last estimate into (4.3), the inequality (4.1) follows. Finally, if we replace $v - (v)_{B_R}$ by v if $v \in W_0^{1,\Phi}(B_R)$, then the estimate (4.1) still holds true since the following classical formula

$$|v(x)| \leq c(n) \int_{B_R} \frac{|Dv(y)|}{|x-y|^{n-1}} dy$$

is valid for a.e $x \in B_R$, whenever $v \in W_0^{1,1}(B_R)$, see for instance [54, Lemma 7.14]. \square

Theorem 4.1. *Let $v \in W^{1,\Psi}(B_R)$ for a ball $B_R \subset \Omega$ with $R \leq 1$ under $a(\cdot) \in C^{\omega_a}(\Omega)$ and $b(\cdot) \in C^{\omega_b}(\Omega)$. Then, for any $d \in \left[1, \frac{n^2}{n^2-1}\right)$, there exist constants $\theta \equiv \theta(n, s(G), s(H_a), s(H_b), d) \in (0, 1)$ and $c \equiv c(n, s(G), s(H_a), s(H_b), \omega_a(1), \omega_b(1), d)$ such that the following Sobolev-Poincaré-type inequality holds:*

$$\left[\int_{B_R} \left[\Psi \left(x, \left| \frac{v - (v)_{B_R}}{R} \right| \right) \right]^d dx \right]^{\frac{1}{d}} \leq c\lambda_{sp} \left[\int_{B_R} [\Psi(x, |Dv|)]^\theta dx \right]^{\frac{1}{\theta}}, \quad (4.4)$$

where

$$\lambda_{sp} = \begin{cases} 1 + ([a]_{\omega_a} + [b]_{\omega_b}) \left(\lambda_1 + \lambda_1 \left(\int_{B_R} G(|Dv|) dx \right)^{\frac{1}{n}} \right) & \text{if } v \in W^{1,\Psi}(B_R) \text{ with (1.12)}_2. \quad (4.5a) \\ 1 + ([a]_{\omega_a} + [b]_{\omega_b}) \left(\lambda_2 + \lambda_2 \|v\|_{L^\infty(B_R)} \right) & \text{if } v \in L^\infty(B_R) \text{ with (1.13)}_2. \quad (4.5b) \\ 1 + ([a]_{\omega_a} + [b]_{\omega_b}) \left(\lambda_3 + \lambda_3 \left[R^{-\gamma} \text{osc}_{B_R} v \right]^{\frac{1}{1-\gamma}} \right) & \text{if } v \in C^{0,\gamma}(B_R) \text{ with (1.14)}_2. \quad (4.5c) \end{cases}$$

Moreover, the above estimate (4.4) is still valid with $v - (v)_{B_R}$ replaced by v depending on which one of (4.5a)-(4.5c) comes into play if $v \in W_0^{1,\Psi}(B_R)$.

Proof. The above theorem widely covers the results of [6, Theorem 4.2], [7, Theorem 32.] and also the results of [30, Theorem 1.6], which is a special case when $G(t) = t^p$, $H(t) = t^q$, $\omega_a(t) = t^\alpha$ and $\omega_b(\cdot) \equiv 0$ for some

constants $1 < p \leq q$ and $\alpha \in (0, 1]$. Then using the continuity of the coefficient functions $a(\cdot)$ and $b(\cdot)$, we find

$$\begin{aligned}
I &:= \left(\int_{B_R} \left[\Psi \left(x, \left| \frac{v - (v)_{B_R}}{R} \right| \right) \right]^d dx \right)^{\frac{1}{d}} \leq 18[a]_{\omega_a} \omega_a(R) \left(\int_{B_R} \left[H_a \left(\left| \frac{v - (v)_{B_R}}{R} \right| \right) \right]^d dx \right)^{\frac{1}{d}} \\
&\quad + 18[b]_{\omega_b} \omega_b(R) \left(\int_{B_R} \left[H_b \left(\left| \frac{v - (v)_{B_R}}{R} \right| \right) \right]^d dx \right)^{\frac{1}{d}} \\
&\quad + 9 \left(\int_{B_R} \left[\Psi_{B_R}^- \left(\left| \frac{v - (v)_{B_R}}{R} \right| \right) \right]^d dx \right)^{\frac{1}{d}} \\
&=: 18[a]_{\omega_a} I_1 + 18[b]_{\omega_b} I_2 + 9I_3,
\end{aligned} \tag{4.6}$$

where we have used the following elementary inequality

$$(t_1 + t_2 + t_3)^d \leq 3^d (t_1^d + t_2^d + t_3^d) \quad (\forall t_1, t_2, t_3 \geq 0).$$

We now estimate the terms I_i with $i \in \{1, 2, 3\}$ in (4.6) depending on which one of (1.12)₂, (1.13)₂ and (1.14)₂ is under consideration. In turn, using (1.12)₂ and (2.2), we see

$$\begin{aligned}
I_1 &= \omega_a(R) \left(\int_{B_R} \left[\frac{(H_a \circ G^{-1}) \left(G \left(\left| \frac{v - (v)_{B_R}}{R} \right| \right) \right)}{G \left(\left| \frac{v - (v)_{B_R}}{R} \right| \right)} G \left(\left| \frac{v - (v)_{B_R}}{R} \right| \right) \right]^d dx \right)^{\frac{1}{d}} \\
&\leq \lambda_1 \omega_a(R) \left(\int_{B_R} \left[\left(1 + \left[\omega_a \left(\left[G \left(\left| \frac{v - (v)_{B_R}}{R} \right| \right) \right]^{-\frac{1}{n}} \right) \right)^{-1} G \left(\left| \frac{v - (v)_{B_R}}{R} \right| \right) \right]^d dx \right)^{\frac{1}{d}} \\
&\leq \lambda_1 \omega_a(R) \left(\int_{B_R} \left[\left(1 + \left[\frac{1}{\omega_a(R)} + \frac{R}{\omega_a(R)} \left(G \left(\left| \frac{v - (v)_{B_R}}{R} \right| \right) \right)^{\frac{1}{n}} \right] G \left(\left| \frac{v - (v)_{B_R}}{R} \right| \right) \right]^d dx \right)^{\frac{1}{d}} \\
&\leq 9\lambda_1(1 + \omega_a(1)) \left(\int_{B_R} \left[G \left(\left| \frac{v - (v)_{B_R}}{R} \right| \right) \right]^d dx \right)^{\frac{1}{d}} \\
&\quad + 9\lambda_1 R \left(\int_{B_R} \left[G \left(\left| \frac{v - (v)_{B_R}}{R} \right| \right) \right]^{(1 + \frac{1}{n})d} dx \right)^{\frac{1}{d}},
\end{aligned} \tag{4.7}$$

where we have used also that $\omega_a(\cdot)$ is non decreasing and $R \leq 1$. In the same way, we have

$$\begin{aligned}
I_2 &\leq 9\lambda_1(1 + \omega_b(1)) \left(\int_{B_R} \left[G \left(\left| \frac{v - (v)_{B_R}}{R} \right| \right) \right]^d dx \right)^{\frac{1}{d}} \\
&\quad + 9\lambda_1 R \left(\int_{B_R} \left[G \left(\left| \frac{v - (v)_{B_R}}{R} \right| \right) \right]^{(1 + \frac{1}{n})d} dx \right)^{\frac{1}{d}}.
\end{aligned} \tag{4.8}$$

Adding the estimates coming from the last two displays and applying Lemma 4.1 with $\Phi \equiv G$ for $d_0 \equiv d$ and $d_0 \equiv (1 + \frac{1}{n})d < \frac{n}{n-1}$, there exists $\theta_1 \equiv \theta_1(n, s(G), d) \in (0, 1)$ such that

$$\begin{aligned} I_1 + I_2 &\leq c\lambda_1 \left(\int_{B_R} \left[G \left(\left| \frac{v - (v)_{B_R}}{R} \right| \right) \right]^d dx \right)^{\frac{1}{d}} + c\lambda_1 R \left(\int_{B_R} \left[G \left(\left| \frac{v - (v)_{B_R}}{R} \right| \right) \right]^{(1+\frac{1}{n})d} dx \right)^{\frac{1}{d}} \\ &\leq c\lambda_1 \left[\int_{B_R} [G(|Dv|)]^{\theta_1} dx \right]^{\frac{1}{\theta_1}} + c\lambda_1 R \left[\int_{B_R} [G(|Dv|)]^{\theta_1} dx \right]^{(1+\frac{1}{n})\frac{1}{\theta_1}} \\ &\leq c\lambda_1 \left[1 + \left(\int_{B_R} G(|Dv|) dx \right)^{\frac{1}{n}} \right] \left[\int_{B_R} G^{\theta_1}(|Dv|) dx \right]^{\frac{1}{\theta_1}} \end{aligned} \quad (4.9)$$

for some constant $c \equiv c(n, s(G), \omega_a(1), \omega_b(1), d)$, where in the last inequality of the above display we have used Hölder's inequality. Since $\Psi_{B_R}^- \in \mathcal{N}$ with an index $s(\Psi) = s(G) + s(H_a) + s(H_b)$ by Remark 2.2, we are able to apply Lemma 4.1 with $\Phi \equiv \Psi_{B_R}^-$ for $d_0 \equiv d$. In turn, there exists $\theta_2 \equiv \theta_2(n, s(\Psi), d)$ such that

$$I_3 \leq c \left[\int_{B_R} [\Psi_{B_R}^- (|Dv|)]^{\theta_2} dx \right]^{\frac{1}{\theta_2}} \quad (4.10)$$

with some constant $c \equiv c(n, s(\Psi), d)$. Inserting the estimates obtained in (4.9)-(4.10) into (4.6), recalling the very definition of $\Psi_{B_R}^-$ in (2.3) and setting $\theta := \max\{\theta_1, \theta_2\}$, we arrive at (4.5a). Now we turn our attention to proving (4.5b). For this, we estimate the terms I_i for $i \in \{1, 2, 3\}$ for $v \in L^\infty(B_R)$ under the assumption (1.13)₂. In turn, using (2.2) and the assumption (1.13)₂, we see

$$\begin{aligned} I_1 &= \omega_a(R) \left(\int_{B_R} \left[\frac{H_a \left(\left| \frac{v - (v)_{B_R}}{R} \right| \right)}{G \left(\left| \frac{v - (v)_{B_R}}{R} \right| \right)} G \left(\left| \frac{v - (v)_{B_R}}{R} \right| \right) \right]^d dx \right)^{\frac{1}{d}} \\ &\leq \lambda_2 \omega_a(R) \left(\int_{B_R} \left[\left(1 + \left[\omega_a \left(\left(\left| \frac{v - (v)_{B_R}}{R} \right| \right)^{-1} \right] \right)^{-1} \right] G \left(\left| \frac{v - (v)_{B_R}}{R} \right| \right) \right]^d dx \right)^{\frac{1}{d}} \\ &\leq \lambda_2 \omega_a(R) \left(\int_{B_R} \left[\left(1 + \left[\frac{1}{\omega_a(R)} + \frac{|v - (v)_{B_R}|}{\omega_a(R)} \right] \right) G \left(\left| \frac{v - (v)_{B_R}}{R} \right| \right) \right]^d dx \right)^{\frac{1}{d}} \\ &\leq 2\lambda_2 \left(1 + \omega_a(1) + \|v\|_{L^\infty(B_R)} \right) \left(\int_{B_R} \left[G \left(\left| \frac{v - (v)_{B_R}}{R} \right| \right) \right]^d dx \right)^{\frac{1}{d}}. \end{aligned} \quad (4.11)$$

In a similar way, one can see

$$I_2 \leq 2\lambda_2 \left(1 + \omega_b(1) + \|v\|_{L^\infty(B_R)} \right) \left(\int_{B_R} \left[G \left(\left| \frac{v - (v)_{B_R}}{R} \right| \right) \right]^d dx \right)^{\frac{1}{d}}. \quad (4.12)$$

Adding the estimates in (4.11)-(4.12) and applying Lemma 4.1 with $\Phi \equiv G$ for $d_0 \equiv d$, there exists an exponent $\theta_1 \equiv \theta_1(n, s(G), d) \in (0, 1)$ such that

$$I_1 + I_2 \leq c\lambda_2 \left(1 + \|v\|_{L^\infty(B_R)} \right) \left[\int_{B_R} [G(|Dv|)]^{\theta_1} dx \right]^{\frac{1}{\theta_1}} \quad (4.13)$$

for some constant $c \equiv c(n, s(G), \omega_a(1), \omega_b(1), d)$. This estimate together with (4.10) and the very definition of $\Psi_{B_R}^-$ in (2.3), we find (4.5b). It remains to prove (4.5c). Essentially, it can be proved in a similar manner we have shown in (4.11)-(4.12). So using the assumption (1.14)₂ and again (2.2), we see

$$\begin{aligned}
I_1 &= \omega_a(R) \left(\int_{B_R} \left[\frac{H_a\left(\left|\frac{v-(v)_{B_R}}{R}\right|\right)}{G\left(\left|\frac{v-(v)_{B_R}}{R}\right|\right)} G\left(\left|\frac{v-(v)_{B_R}}{R}\right|\right) \right]^d dx \right)^{\frac{1}{d}} \\
&\leq \lambda_3 \omega_a(R) \left(\int_{B_R} \left[\left(1 + \left[\omega_a \left(\left(\left| \frac{v-(v)_{B_R}}{R} \right| \right)^{-\frac{1}{1-\gamma}} \right] \right)^{-1} G\left(\left|\frac{v-(v)_{B_R}}{R}\right|\right) \right]^d dx \right)^{\frac{1}{d}} \\
&\leq \lambda_3 \omega_a(R) \left(\int_{B_R} \left[\left(1 + \left[\frac{1}{\omega_a(R)} + \frac{R^{-\frac{\gamma}{1-\gamma}}}{\omega_a(R)} |v-(v)_{B_R}|^{\frac{1}{1-\gamma}} \right] G\left(\left|\frac{v-(v)_{B_R}}{R}\right|\right) \right]^d dx \right)^{\frac{1}{d}} \\
&\leq 2\lambda_3 \left(1 + \omega_a(1) + \left[R^{-\gamma} \operatorname{osc}_{B_R} v \right]^{\frac{1}{1-\gamma}} \right) \left(\int_{B_R} \left[G\left(\left|\frac{v-(v)_{B_R}}{R}\right|\right) \right]^d dx \right)^{\frac{1}{d}}.
\end{aligned} \tag{4.14}$$

By arguing in the same way, we see

$$I_1 + I_2 \leq c\lambda_3 \left(1 + \left[R^{-\gamma} \operatorname{osc}_{B_R} v \right]^{\frac{1}{1-\gamma}} \right) \left(\int_{B_R} \left[G\left(\left|\frac{v-(v)_{B_R}}{R}\right|\right) \right]^d dx \right)^{\frac{1}{d}}. \tag{4.15}$$

for some constant $c \equiv c(s(G), \omega_a(1), \omega_b(1))$. Finally, this estimate together with (4.10) leads to (4.5c). The proof is complete. \square

Remark 4.1. We here remark that choosing $d \equiv 1$ in a Sobolev-Poincaré type inequality of Theorem 4.1, there exist an exponent $\theta \equiv \theta(n, s(G), s(H_a), s(H_b))$ such that

$$\int_{B_R} \Psi \left(x, \left| \frac{v-(v)_{B_R}}{R} \right| \right) dx \leq c\lambda_{sp} \left[\int_{B_R} [\Psi(x, |Dv|)]^\theta dx \right]^{\frac{1}{\theta}}, \tag{4.16}$$

holds for some constant $c \equiv c(n, s(G), s(H_a), s(H_b), \omega_a(1), \omega_b(1))$ where

$$\lambda_{sp} = \begin{cases} 1 + ([a]_{\omega_a} + [b]_{\omega_b}) \left(\lambda_1 + \lambda_1 \left(\int_{B_R} G(|Dv|) dx \right)^{\frac{1}{n}} \right) & \text{if } v \in W^{1,\Psi}(B_R) \text{ with (1.12)}_2. \quad (4.17a) \\ 1 + ([a]_{\omega_a} + [b]_{\omega_b}) \left(\lambda_2 + \lambda_2 \|v\|_{L^\infty(B_R)} \right) & \text{if } v \in L^\infty(B_R) \text{ with (1.13)}_2. \quad (4.17b) \\ 1 + ([a]_{\omega_a} + [b]_{\omega_b}) \left(\lambda_3 + \lambda_3 \left[R^{-\gamma} \operatorname{osc}_{B_R} v \right]^{\frac{1}{1-\gamma}} \right) & \text{if } v \in C^{0,\gamma}(B_R) \text{ with (1.14)}_2. \quad (4.17c) \end{cases}$$

5. BASIC REGULARITY RESULTS

We start this section by stating the following Caccioppoli inequality which is a fundamental result for the further investigations. In what follows let $Q = L/\nu$ for the convenience in the future but in general it could be any number larger than one.

Lemma 5.1 (Caccioppoli Inequality). *Let $u \in W^{1,\Psi}(\Omega)$ be a local Q -minimizer of the functional \mathcal{P} defined in (1.1) with $0 \leq a(\cdot), b(\cdot) \in L^\infty(\Omega)$. Then there exists a positive constant $c \equiv c(s(G), s(H_a), s(H_b), Q)$ such*

that the following Caccioppoli inequality

$$\int_{B_\rho} \Psi(x, |D(u-k)_\pm|) dx \leq c \int_{\tilde{B}_R} \Psi\left(x, \frac{(u-k)_\pm}{R-\rho}\right) dx \quad (5.1)$$

holds, whenever $B_\rho \Subset B_R \subset \Omega$ are concentric balls and $k \in \mathbb{R}$.

Proof. The proof is elementary as done for [22, Lemma 4.6]. The only difference lies in that we have an additional one phase. But the inequality (5.1) is still valid since $H_b \in \mathcal{N}$ with an index $s(H_b) \geq 1$. \square

Remark 5.1. As a direct consequence of Lemma 5.1, with $u \in W^{1,\Psi}(B_R)$ being a local Q -minimizer of the functional \mathcal{P} defined in (1.1) under the assumptions of Lemma 5.1, there exists a positive constant $c \equiv c(n, s(G), s(H_a), s(H_b), Q)$ such that

$$\int_{B_{R/2}} \Psi(x, |Du|) dx \leq c \int_{\tilde{B}_R} \Psi\left(x, \left| \frac{u - (u)_{B_R}}{R} \right| \right) dx$$

holds, whenever $B_R \subset \Omega$ is a ball.

5.1. Local boundedness. Now we focus on local boundedness of a local Q -minimizer u of the functional \mathcal{P} defined in (1.1) with obtaining precise estimates under the assumption (1.12)₂.

Theorem 5.1. *Let $u \in W^{1,\Psi}(\Omega)$ be a local Q -minimizer of the functional \mathcal{P} defined in (1.1) under the assumption (1.12). Then there exists a constant $c \equiv c(\text{data})$ such that*

$$\left\| \Psi_{B_R}^- \left(\left| \frac{(u - (u)_{B_R})_\pm}{R} \right| \right) \right\|_{L^\infty(B_{R/2})} \leq c \int_{\tilde{B}_R} \Psi\left(x, \left| \frac{(u - (u)_{B_R})_\pm}{R} \right| \right) dx \quad (5.2)$$

and

$$\Psi_{B_R}^- \left(\left| \frac{u(x_1) - u(x_2)}{R} \right| \right) \leq c \int_{\tilde{B}_R} \Psi(x, |Du|) dx \quad \text{for a.e. } x_1, x_2 \in B_{R/2}, \quad (5.3)$$

whenever $B_R \equiv B_R(x_0) \subset \Omega$ is a ball with $R \leq 1$. In particular, $u \in L^\infty_{\text{loc}}(\Omega)$.

Proof. Let us consider the following scaling:

$$\begin{aligned} \bar{u}(x) &:= \frac{u(x_0 + Rx) - (u)_{B_R}}{R}, \quad \bar{a}(x) := a(x_0 + Rx), \quad \bar{b}(x) := b(x_0 + Rx), \\ \bar{\Psi}(x, t) &:= G(t) + \bar{a}(x)H(t) + \bar{b}(x)H(t), \\ \bar{A}(k, s) &:= B_s(0) \cap \{\bar{u} > k\} \quad \text{and} \quad \bar{B}(k, s) := B_s(0) \cap \{\bar{u} < k\} \end{aligned} \quad (5.4)$$

for every $x \in B_1(0)$, $t \geq 0$, $s \in (0, 1)$ and $k \in \mathbb{R}$. The rest of the proof falls in 3 steps.

Step 1: Sobolev-Poincaré inequality under the scaling. Before going on further, let us consider a Sobolev-Poincaré type inequality under the new scaling introduced in (5.4). So we prove that there exists a positive exponent $\theta \equiv \theta(n, s(G), s(H_a), s(H_b)) \in (0, 1)$ such that

$$\int_{B_1} \bar{\Psi}(x, |f|) dx \leq c \bar{\kappa}_{sp} \left(\int_{\tilde{B}_1} [\bar{\Psi}(x, |Df|)]^\theta dx \right)^{\frac{1}{\theta}} \quad (5.5)$$

for some constant $c \equiv c(n, s(G), s(H_a), s(H_b), \omega_a(1), \omega_b(1))$, whenever $f \in W_0^{1,\bar{\Psi}}(B_1)$, where

$$\bar{\kappa}_{sp} = 1 + ([a]_{\omega_a} + [b]_{\omega_b}) \left(\lambda_1 + \lambda_1 R \left(\int_{\tilde{B}_1} G(|Df|) dx \right)^{\frac{1}{n}} \right). \quad (5.6)$$

Essentially, the proof of the inequality (5.5) comes from a careful revealing of the arguments used in (4.7)-(4.9). So using continuity properties of $\bar{a}(\cdot)$ and $\bar{b}(\cdot)$, we see

$$\begin{aligned} I &:= \int_{B_1} \bar{\Psi}(x, |f|) dx \leq 2[a]_{\omega_a} \omega_a(R) \int_{B_1} H_a(|f|) dx + 2[b]_{\omega_b} \omega_b(R) \int_{B_1} H_b(|f|) dx + \int_{B_1} \bar{\Psi}_{B_1}^-(|f|) dx \\ &: 2[a]_{\omega_a} I_1 + 2[b]_{\omega_b} I_2 + I_3, \end{aligned} \quad (5.7)$$

where

$$\bar{\Psi}_{B_1}^-(t) := G(t) + \inf_{x \in B_1} \bar{a}(x) H_a(t) + \inf_{x \in B_1} \bar{b}(x) H_b(t) \quad \text{for every } t \geq 0. \quad (5.8)$$

Now we estimate the terms I_i for $i \in \{1, 2, 3\}$ similarly as in the proof of Theorem 4.1. In turn, using the assumption (1.12)₂ and (2.2), we have

$$\begin{aligned} I_1 &= \omega_a(R) \int_{B_1} \frac{H_a(|f|)}{G(|f|)} G(|f|) dx \\ &\leq \lambda_1 \omega_a(R) \int_{B_1} \left(1 + \left[\omega_a \left([G(|f|)]^{-\frac{1}{n}} \right) \right]^{-1} \right) G(|f|) dx \\ &\leq \lambda_1 \omega_a(R) \int_{B_1} \left(1 + \left[\frac{1}{\omega_a(R)} + \frac{R}{\omega_a(R)} [G(|f|)]^{\frac{1}{n}} \right] \right) G(|f|) dx \\ &\leq \lambda_1 (1 + \omega_a(1)) \int_{B_1} G(|f|) dx + 2\lambda_1 R \int_{B_1} [G(|f|)]^{1+\frac{1}{n}} dx. \end{aligned} \quad (5.9)$$

In a similar manner, we find

$$I_2 \leq \lambda_1 (1 + \omega_b(1)) \int_{B_1} G(|f|) dx + \lambda_1 R \int_{B_1} [G(|f|)]^{1+\frac{1}{n}} dx. \quad (5.10)$$

Inserting the estimates (5.9)-(5.10) into (5.8), the inequality (5.5) follows from the similar arguments used in (4.9)-(4.10) and Lemma 4.1.

Step 2. Proof of (5.2). Since $u - (u)_{B_R}$ is a local Q -minimizer of the functional \mathcal{P} , we use a Caccioppoli inequality of Lemma 5.1 to see that

$$\int_{B_t} \bar{\Psi}(x, |D(\bar{u} - k)_{\pm}|) dx \leq c \int_{B_s} \bar{\Psi} \left(x, \frac{(\bar{u} - k)_{\pm}}{s - t} \right) dx \quad (5.11)$$

holds for some constant $c \equiv c(s(G), s(H_a), s(H_b), Q)$, whenever $0 < t < s \leq 1$ and $k \in \mathbb{R}$. Let us now consider the concentric balls $B_\rho \Subset B_t \Subset B_s$ with $1/2 \leq \rho < s \leq 1$ and $t := (\rho + s)/2$. Let $\eta \in C_0^\infty(B_t)$ be a standard cut-off function such that $\chi_{B_\rho} \leq \eta \leq \chi_{B_t}$ and $|D\eta| \leq \frac{2}{t-\rho} = \frac{4}{s-\rho}$. Now we apply inequality (5.5) from Step 1 above in order to have a positive exponent $\theta \equiv \theta(n, s(G), s(H_a), s(H_b))$ such that

$$\int_{\bar{A}(k, \rho)} \bar{\Psi}(x, \bar{u} - k) dx \leq \int_{B_1} \bar{\Psi}(x, \eta(\bar{u} - k)_+) dx \leq c \bar{k}_{sp} \left(\int_{B_1} [\bar{\Psi}(x, |D(\eta(\bar{u} - k)_+)|)]^\theta dx \right)^{\frac{1}{\theta}} \quad (5.12)$$

for some constant $c \equiv c(n, s(G), s(H_a), s(H_b), \omega_a(1), \omega_b(1))$, where

$$\bar{k}_{sp} = 1 + ([a]_{\omega_a} + [b]_{\omega_b}) \left(\lambda_1 + \lambda_1 R \left(\int_{B_1} G(|D(\eta(\bar{u} - k)_+)|) dx \right)^{\frac{1}{n}} \right). \quad (5.13)$$

By scaling back and using Lemma 2.1, for any $k \geq 0$, we have

$$\begin{aligned} \bar{\kappa}_{sp} &\leq c \left[1 + R \left(\oint_{\bar{B}_R} G(|Du|) dx \right)^{\frac{1}{n}} + \frac{R}{(s-\rho)^{\frac{s(G)+1}{n}}} \left(\oint_{\bar{B}_R} G \left(\left| \frac{u - (u)_{B_R}}{R} \right| \right) dx \right)^{\frac{1}{n}} \right] \\ &\leq \frac{c}{(s-\rho)^{s(G)+1}} \left[1 + \left(\int_{\bar{B}_R} G(|Du|) dx \right)^{\frac{1}{n}} \right] \end{aligned} \quad (5.14)$$

with a constant $c \equiv c(n, \lambda_1, [a]_{\omega_a} + [b]_{\omega_b})$, where we have also used Lemma 4.1 to $\Phi \equiv G$ for $d_0 \equiv 1$.

Then, inserting the last estimate into (5.12) and applying Hölder inequality together with (5.11) yield that

$$\begin{aligned} \int_{\bar{A}(k, \rho)} \bar{\Psi}(x, \bar{u} - k) dx &\leq c \frac{1}{(s-\rho)^{1+s(G)}} |\bar{A}(k, t)|^{\frac{1-\theta}{\theta}} \int_{\bar{A}(k, t)} \left(\bar{\Psi}(x, |D\bar{u}|) + \bar{\Psi} \left(x, \frac{\bar{u} - k}{s-\rho} \right) \right) dx \\ &\leq c \frac{1}{(s-\rho)^{1+s(G)}} |\bar{A}(k, s)|^{\frac{1-\theta}{\theta}} \int_{\bar{A}(k, s)} \bar{\Psi} \left(x, \frac{\bar{u} - k}{s-\rho} \right) dx \end{aligned} \quad (5.15)$$

holds with some constant $c \equiv c(\mathbf{data})$, where in the last display we have also used (2.6). By the very definition of \bar{A} in (5.4), we observe

$$|\bar{A}(k, s)| \leq \int_{\bar{A}(h, s)} \frac{\bar{\Psi}(x, \bar{u} - h)}{\bar{\Psi}(x, k - h)} dx \leq \frac{1}{\bar{\Psi}_{B_1}^-(k - h)} \int_{\bar{A}(h, s)} \bar{\Psi}(x, \bar{u} - h) dx$$

and

$$\int_{\bar{A}(k, s)} \bar{\Psi}(x, \bar{u} - k) dx \leq \int_{\bar{A}(h, s)} \bar{\Psi}(x, \bar{u} - h) dx$$

for any $h < k$. Putting the last two inequalities into (5.15) and applying Lemma 2.1, we have the following inequality:

$$\int_{\bar{A}(k, \rho)} \bar{\Psi}(x, \bar{u} - k) dx \leq \frac{c}{[\bar{\Psi}_{B_1}^-(k - h)]^{\frac{1-\theta}{\theta}} (s-\rho)^{2(\max\{s(G), s(H_a), s(H_b)\}+1)}} \left(\int_{\bar{A}(h, s)} \bar{\Psi}(x, \bar{u} - h) dx \right)^{\frac{1}{\theta}}. \quad (5.16)$$

Now we set sequences of numbers as follows:

$$\rho_i := \frac{1}{2} \left(1 + \frac{1}{2^i} \right), \quad k_i := 2l_0 \left(1 - \frac{1}{2^{i+1}} \right) \text{ and } M_i := \frac{1}{\bar{\Psi}_{B_1}^-(l_0)} \int_{\bar{A}(k_i, \rho_i)} \bar{\Psi}(x, \bar{u} - k_i) dx$$

for any integer $i \geq 0$ and some number $l_0 > 0$ to be chosen in a few lines. Then applying (5.16) with the choices $k \equiv k_{i+1}$, $h \equiv k_i$, $\rho \equiv \rho_{i+1}$ and $s \equiv \rho_i$, we have, for every $i \geq 0$,

$$\begin{aligned} M_{i+1} &\leq \frac{c}{[\bar{\Psi}_{B_1}^-(\frac{l_0}{2^{i+1}})]^{\frac{1-\theta}{\theta}} (\frac{1}{4^{i+2}})^{\max\{s(G), s(H_a), s(H_b)\}+1}} [\bar{\Psi}_{B_1}^-(l_0)]^{\frac{1-\theta}{\theta}} M_i^{\frac{1}{\theta}} \\ &\leq c_0 \left[4^{(\max\{s(G), s(H_a), s(H_b)\}+1)\frac{1}{\theta}} \right]^i M_i^{1+\frac{1-\theta}{\theta}} \end{aligned}$$

with $c_0 \equiv c_0(\mathbf{data})$, where in the last inequality of the last display we have used again Lemma 2.1. Now it's turn to apply a standard iteration of Lemma 2.7, which means that if

$$\frac{1}{\bar{\Psi}_{B_1}^-(l_0)} \int_{\bar{A}(l_0, 1)} \bar{\Psi}(x, \bar{u} - l_0) dx = M_0 \leq c_0^{-\frac{\theta}{1-\theta}} 4^{-(\max\{s(G), s(H_a), s(H_b)\}+1)\frac{\theta}{(1-\theta)^2}},$$

then we obtain

$$\|\bar{u}_+\|_{L^\infty(B_{1/2})} \leq 2l_0.$$

Consequently, choosing $l_0 > 0$ in such a way that

$$\bar{\Psi}_{B_1}^-(l_0) = c_0^{\frac{\theta}{1-\theta}} 4^{(\max\{s(G), s(H_a), s(H_b)\}+1)\frac{\theta}{(1-\theta)^2}} \int_{B_1} \bar{\Psi}(x, \bar{u}_+) dx,$$

we have

$$\|\bar{\Psi}_{B_1}^-(\bar{u}_+)\|_{L^\infty(B_{1/2})} \leq c \int_{B_1} \bar{\Psi}(x, \bar{u}_+) dx,$$

which implies that

$$\left\| \Psi_{B_R}^- \left(\frac{(u - (u)_{B_R})_+}{R} \right) \right\|_{L^\infty(B_{R/2})} \leq c \int_{B_R} \Psi \left(x, \frac{(u - (u)_{B_R})_+}{R} \right) dx$$

holds with $c \equiv c(\mathbf{data})$. Repeating the same argument for $-u$, which is also a local Q -minimizer of the functional \mathcal{P} defined in (1.9), the last inequality holds with $(u - (u)_{B_R})_+$ replaced by $(u - (u)_{B_R})_-$.

Step 3. Proof of (5.3). Using (5.2) and (2.6), for a.e. $x_1, x_2 \in B_{R/2}$, we have

$$\begin{aligned} \Psi_{B_R}^- \left(\left| \frac{u(x_1) - u(x_2)}{R} \right| \right) &\leq c \Psi_{B_R}^- \left(\left| \frac{u(x_1) - (u)_{B_R}}{R} \right| \right) + c \Psi_{B_R}^- \left(\left| \frac{u(x_2) - (u)_{B_R}}{R} \right| \right) \\ &\leq c \int_{B_R} \Psi \left(x, \left| \frac{u - (u)_{B_R}}{R} \right| \right) dx \leq c \int_{B_R} \Psi(x, |Du|) dx \end{aligned}$$

for some constant $c \equiv c(\mathbf{data})$, where in the last inequality of the above display we have used a Sobolev-Poincaré type inequality of Theorem 4.1. Clearly, the last display implies $u \in L_{\text{loc}}^\infty(\Omega)$. The proof is complete. \square

5.2. Almost standard Caccioppoli inequality. Now we present the primary results, the so-called almost standard Caccioppoli type inequality, for proving Hölder continuity of a local Q -minimizer of the functional \mathcal{P} .

Lemma 5.2 (Almost standard Caccioppoli inequality). *Let $u \in W^{1,\Psi}(\Omega)$ be a local Q -minimizer of the functional \mathcal{P} defined in (1.1) under one of the assumptions (1.12), (1.13) and (1.14). Let $B_{2R} \equiv B_{2R}(x_0) \subset \Omega$ be a ball with $R \leq 1$. Then there exists a constant $c \equiv c(\mathbf{data})$ such that*

$$\begin{aligned} \int_{B_{R_1}} \Psi_{B_R}^- (|D(u - k)_\pm|) dx &\leq \int_{B_{R_1}} \Psi(x, |D(u - k)_\pm|) dx \\ &\leq c \left(\frac{R}{R_2 - R_1} \right)^{s(\Psi)+1} \int_{B_{R_2}} \Psi_{B_R}^- \left(\frac{(u - k)_\pm}{R} \right) dx \end{aligned} \quad (5.17)$$

holds, whenever $B_{R_1} \Subset B_{R_2} \subset B_R(x_0)$ are concentric balls and $k \in \mathbb{R}$.

Proof. First we prove the inequality (5.17) for the values of $k \in \mathbb{R}$ with $\inf_{B_R} u \leq k \leq \sup_{B_R} u$, depending on which one of the assumptions (1.12)-(1.14) is in force. Firstly by the very definition of $\Psi_{B_R}^-$ in (2.3) and Lemma 5.1, we see

$$\begin{aligned} I &:= \int_{B_{R_1}} \Psi_{B_R}^- (|D(u - k)_\pm|) dx \leq \int_{B_{R_1}} \Psi(x, |D(u - k)_\pm|) dx \leq c_* \int_{B_{R_2}} \Psi \left(x, \frac{(u - k)_\pm}{R_2 - R_1} \right) dx \\ &\leq c_* \omega_a(R) \int_{B_{R_2}} H_a \left(\frac{(u - k)_\pm}{R_2 - R_1} \right) dx + c_* \omega_b(R) \int_{B_{R_2}} H_b \left(\frac{(u - k)_\pm}{R_2 - R_1} \right) dx \\ &\quad + c_* \int_{B_{R_2}} \Psi_{B_R}^- \left(\frac{(u - k)_\pm}{R_2 - R_1} \right) dx =: c_* (I_1 + I_2 + I_3) \end{aligned} \quad (5.18)$$

for some constant $c_* \equiv c_*(n, s(G), s(H_a), s(H_b), Q, [a]_{\omega_a}, [b]_{\omega_b})$. Now we shall estimate each term I_i for $i \in \{1, 2, 3\}$ in the above display. Then using Lemma 2.1, the assumption (1.12)₂, (2.2) and (5.3) of Lemma 5.1, we see

$$\begin{aligned}
I_1 &= \omega_a(R) \int_{B_{R_2}} \frac{H_a\left(\frac{(u-k)_\pm}{R_2-R_1}\right)}{G\left(\frac{(u-k)_\pm}{R_2-R_1}\right)} G\left(\frac{(u-k)_\pm}{R_2-R_1}\right) dx \\
&\leq \omega_a(R) \left(\frac{R}{R_2-R_1}\right)^{s(H_a)+1} \int_{B_{R_2}} \frac{(H_a \circ G^{-1})\left(G\left(\frac{(u-k)_\pm}{R}\right)\right)}{G\left(\frac{(u-k)_\pm}{R}\right)} G\left(\frac{(u-k)_\pm}{R_2-R_1}\right) dx \\
&\leq \lambda_1 \omega_a(R) \left(\frac{R}{R_2-R_1}\right)^{s(H_a)+1} \int_{B_{R_2}} \left(1 + \left[\omega_a\left(\left[G\left(\frac{(u-k)_\pm}{R}\right)\right]^{-\frac{1}{n}}\right)\right]^{-1}\right) G\left(\frac{(u-k)_\pm}{R_2-R_1}\right) dx \\
&\leq c \omega_a(R) \left(\frac{R}{R_2-R_1}\right)^{s(H_a)+1} \int_{B_{R_2}} \left[1 + \frac{1}{\omega_a(R)} + \frac{R}{\omega_a(R)} \left(G\left(\frac{\text{osc}_{B_R} u}{R}\right)\right)^{\frac{1}{n}}\right] G\left(\frac{(u-k)_\pm}{R_2-R_1}\right) dx \\
&\leq c \left[1 + \left(\int_{B_{2R}} \Psi(x, |Du|) dx\right)^{\frac{1}{n}}\right] \left(\frac{R}{R_2-R_1}\right)^{s(H_a)+1} \int_{B_{R_2}} G\left(\frac{(u-k)_\pm}{R_2-R_1}\right) dx \\
&\leq c \left(\frac{R}{R_2-R_1}\right)^{s(\Psi)+1} \int_{B_{R_2}} G\left(\frac{(u-k)_\pm}{R}\right) dx
\end{aligned} \tag{5.19}$$

for some constant $c \equiv c(\mathbf{data})$. In a totally similar way, it can be shown that

$$I_2 \leq c \left(\frac{R}{R_2-R_1}\right)^{s(\Psi)+1} \int_{B_{R_2}} G\left(\frac{(u-k)_\pm}{R}\right) dx \tag{5.20}$$

with a constant $c \equiv c(\mathbf{data})$. Clearly, recalling Remark 2.2 and using Lemma 2.1, we have

$$I_3 \leq \left(\frac{R}{R_2-R_1}\right)^{s(\Psi)+1} \int_{B_{R_2}} \Psi_{B_R}^- \left(\frac{(u-k)_\pm}{R}\right) dx \tag{5.21}$$

Inserting the estimates obtained in (5.19)-(5.21) into (5.18) and recalling the very definition of $\Psi_{B_R}^-$ in (2.3), we arrive at (5.17) under the assumption (1.12). The second part of the proof is to show (5.17) under the assumption (1.13). For this, we again estimate the terms I_i with $i \in \{1, 2, 3\}$ in (5.18). Applying Lemma

2.1, the assumption (1.13) and (2.2), we see

$$\begin{aligned}
I_1 &= \omega_a(R) \int_{B_{R_2}} \left(\frac{H_a}{G} \right) \left(\frac{(u-k)_\pm}{R_2 - R_1} \right) G \left(\frac{(u-k)_\pm}{R_2 - R_1} \right) dx \\
&\leq \omega_a(R) \left(\frac{R}{R_2 - R_1} \right)^{s(H_a)+1} \int_{B_{R_2}} \left(\frac{H_a}{G} \right) \left(\frac{(u-k)_\pm}{R} \right) G \left(\frac{(u-k)_\pm}{R_2 - R_1} \right) dx \\
&\leq 2\lambda_2 \omega_a(R) \left(\frac{R}{R_2 - R_1} \right)^{s(H_a)+1} \int_{B_{R_2}} \left(1 + \left[\omega_a \left(\frac{R}{(u-k)_\pm} \right) \right]^{-1} \right) G \left(\frac{(u-k)_\pm}{R_2 - R_1} \right) dx \\
&\leq c\lambda_2 \omega_a(R) \left(\frac{R}{R_2 - R_1} \right)^{s(H_a)+1} \int_{B_{R_2}} \left(1 + \left[\frac{1}{\omega_a(R)} + \frac{\|u\|_{L^\infty(B_R)}}{\omega_a(R)} \right] \right) G \left(\frac{(u-k)_\pm}{R_2 - R_1} \right) dx \\
&\leq c \left(\frac{R}{R_2 - R_1} \right)^{s(\Psi)+1} \int_{B_{R_2}} G \left(\frac{(u-k)_\pm}{R} \right) dx
\end{aligned} \tag{5.22}$$

for some constant $c \equiv c(\mathbf{data})$. Arguing similarly, we have

$$I_2 \leq c \left(\frac{R}{R_2 - R_1} \right)^{s(\Psi)+1} \int_{B_{R_2}} G \left(\frac{(u-k)_\pm}{R} \right) dx \tag{5.23}$$

with a constant $c \equiv c(\mathbf{data})$. Plugging the estimates (5.21)-(5.23) into (5.18), we conclude with (5.17) under the assumption (1.13). Finally, the remaining part of the proof is to obtain the inequality (5.17) under the assumption (1.14). In fact, we continue to estimate the terms I_i with $i \in \{1, 2, 3\}$ in (5.18). Therefore, using the assumption (1.14) and (2.2), we find

$$\begin{aligned}
I_1 &= \omega_a(R) \int_{B_{R_2}} \left(\frac{H_a}{G} \right) \left(\frac{(u-k)_\pm}{R_2 - R_1} \right) G \left(\frac{(u-k)_\pm}{R_2 - R_1} \right) dx \\
&\leq \omega_a(R) \left(\frac{R}{R_2 - R_1} \right)^{s(H_a)+1} \int_{B_{R_2}} \left(\frac{H_a}{G} \right) \left(\frac{(u-k)_\pm}{R} \right) G \left(\frac{(u-k)_\pm}{R_2 - R_1} \right) dx \\
&\leq 2\lambda_3 \omega_a(R) \left(\frac{R}{R_2 - R_1} \right)^{s(H_a)+1} \int_{B_{R_2}} \left(1 + \left[\omega_a \left(\left[\frac{R}{(u-k)_\pm} \right]^{\frac{1}{1-\gamma}} \right) \right]^{-1} \right) G \left(\frac{(u-k)_\pm}{R_2 - R_1} \right) dx \\
&\leq c\lambda_3 \omega_a(R) \left(\frac{R}{R_2 - R_1} \right)^{s(H_a)+1} \int_{B_{R_2}} \left(1 + \frac{1}{\omega_a(R)} \right) G \left(\frac{(u-k)_\pm}{R_2 - R_1} \right) dx \\
&\leq c \left(\frac{R}{R_2 - R_1} \right)^{s(\Psi)+1} \int_{B_{R_2}} G \left(\frac{(u-k)_\pm}{R} \right) dx
\end{aligned} \tag{5.24}$$

for some constant $c \equiv c(\mathbf{data})$, where we have used Lemma 2.1 several times. Using the same argument as above, we have

$$I_2 \leq c \left(\frac{R}{R_2 - R_1} \right)^{s(\Psi)+1} \int_{B_{R_2}} G \left(\frac{(u-k)_\pm}{R} \right) dx \tag{5.25}$$

with a constant $c \equiv c(\mathbf{data})$. Inserting the estimates (5.21), (5.24)-(5.25) into (5.18), we arrive at (5.17) under the assumption (1.14). So we have proved the inequality (5.17) for the values of $k \in \mathbb{R}$ such that $\inf_{B_R} u \leq k \leq \sup_{B_R} u$. Now we consider the remaining cases. Suppose $k < \inf_{B_R} u$. In this case, using (5.17) with $k \equiv \inf_{B_R} u$, we have

$$\begin{aligned}
\int_{B_{R_1}} \Psi_{B_R}^- (|D(u-k)_+|) dx &= \int_{B_{R_1}} \Psi_{B_R}^- \left(\left| D(u - \inf_{B_R} u)_+ \right| \right) dx \leq \int_{B_{R_1}} \Psi \left(x, \left| D(u - \inf_{B_R} u)_+ \right| \right) dx \\
&\leq c \left(\frac{R}{R_2 - R_1} \right)^{s(\Psi)+1} \int_{B_{R_2}} \Psi_{B_R}^- \left(\frac{\left(u - \inf_{B_R} u \right)_+}{R} \right) dx \\
&\leq c \left(\frac{R}{R_2 - R_1} \right)^{s(\Psi)+1} \int_{B_{R_2}} \Psi_{B_R}^- \left(\frac{(u-k)_+}{R} \right) dx
\end{aligned} \tag{5.26}$$

for some constant $c \equiv c(\mathbf{data})$. Similarly, it can be seen that (5.26) is valid for the values of $k > \sup_{B_R} u$. Since $-u$ is also the local Q -minimizer of the functional \mathcal{P} in (1.1), the inequality (5.17) is valid for all $k \in \mathbb{R}$. The proof is complete. \square

From now on also in the rest of paper, for a fixed ball $B_R \subset \Omega$, we say that

$$\left\{ \begin{array}{ll} G - \text{phase occurs in } B_R & \text{if } a^-(B_R) \leq 4[a]_{\omega_a} \omega_a(R) \text{ and } b^-(B_R) \leq 4[b]_{\omega_b} \omega_b(R). \end{array} \right. \tag{5.27a}$$

$$\left\{ \begin{array}{ll} (G, H_a) - \text{phase occurs in } B_R & \text{if } a^-(B_R) > 4[a]_{\omega_a} \omega_a(R) \text{ and } b^-(B_R) \leq 4[b]_{\omega_b} \omega_b(R). \end{array} \right. \tag{5.27b}$$

$$\left\{ \begin{array}{ll} (G, H_b) - \text{phase occurs in } B_R & \text{if } a^-(B_R) \leq 4[a]_{\omega_a} \omega_a(R) \text{ and } b^-(B_R) > 4[b]_{\omega_b} \omega_b(R). \end{array} \right. \tag{5.27c}$$

$$\left\{ \begin{array}{ll} (G, H_a, H_b) - \text{phase occurs in } B_R & \text{if } a^-(B_R) > 4[a]_{\omega_a} \omega_a(R) \text{ and } b^-(B_R) > 4[b]_{\omega_b} \omega_b(R). \end{array} \right. \tag{5.27d}$$

Then we have the following lemma which will be applied later, see Section 8.

Lemma 5.3. *Let $u \in W^{1,\Psi}(\Omega)$ be a local Q -minimizer of the functional \mathcal{P} defined in (1.1) under one of the assumptions (1.12), (1.13) and (1.14). Let $B_{2R} \equiv B_{2R}(x_0) \subset \Omega$ be a ball with $R \leq 1$. Then there exists a constant $c \equiv c(\mathbf{data})$ such that*

$$\begin{aligned}
\int_{B_{R_1}} \Psi_{B_R}^- (|D(u-k)_\pm|) dx &\leq \int_{B_{R_1}} \Psi(x, |D(u-k)_\pm|) dx \\
&\leq c \left(\frac{R}{R_2 - R_1} \right)^{s(\Psi)+1} \int_{B_{R_2}} \Phi \left(\frac{(u-k)_\pm}{R} \right) dx
\end{aligned} \tag{5.28}$$

holds, whenever $B_{R_1} \Subset B_{R_2} \subset B_R(x_0)$ are concentric balls and $k \in \mathbb{R}$, where

$$\Phi(t) = \begin{cases} G(t) & \text{if (5.27a) is satisfied in } B_R, \\ G(t) + a^-(B_R)H_a(t) & \text{if (5.27b) is satisfied in } B_R, \\ G(t) + b^-(B_R)H_b(t) & \text{if (5.27c) is satisfied in } B_R, \\ \Psi_{B_R}^-(t) & \text{if (5.27d) is satisfied in } B_R, \end{cases}$$

for every $t \geq 0$.

Proof. First we observe that if $a^-(B_R) > 4[a]_{\omega_a} \omega_a(R)$, then using the continuity of the function $a(\cdot)$, we have

$$a^-(B_R) \leq a(x) = a(x) - a^-(B_R) + a^-(B_R) \leq 2[a]_{\omega_a} \omega_a(R) + a^-(B_R) \leq 2a^-(B_R) \tag{5.30}$$

for every $x \in B_R$. On the other hand, if $a^-(B_R) \leq 4[a]_{\omega_a} \omega_a(R)$, then using again the continuity of $a(\cdot)$, we see

$$a(x) = a(x) - a^-(B_R) + a^-(B_R) \leq 6[a]_{\omega_a} \omega_a(R)$$

for every $x \in B_R$. Clearly, analogous estimates to the last two displays are valid for the function $b(\cdot)$ in B_R . After those observations, we argue similarly as in the proof of Lemma 5.2 depending on which case of (5.27b)-(5.27d) occurs in the ball B_R . \square

5.3. Hölder continuity. In this subsection we prove some local boundedness and Hölder continuity assertions of a local Q -minimizer of the functional \mathcal{P} in (1.1) with various constants having the precise dependencies.

Theorem 5.2. *Let $u \in W^{1,\Psi}(\Omega)$ be a local Q -minimizer of the functional \mathcal{P} defined in (1.1) under the coefficient functions $a(\cdot) \in C^{\omega_a}(\Omega)$ and $b(\cdot) \in C^{\omega_b}(\Omega)$ for ω_a, ω_b being non-negative concave functions vanishing at the origin.*

1. *If the assumption (1.12) is satisfied, then for every open subset $\Omega_0 \Subset \Omega$, there exists a Hölder continuity exponent $\gamma \equiv \gamma(\mathbf{data}(\Omega_0)) \in (0, 1)$ such that*

$$\|u\|_{L^\infty(\Omega_0)} + [u]_{0,\gamma;\Omega_0} \leq c(\mathbf{data}(\Omega_0)) \quad (5.31)$$

and the oscillation estimate

$$\text{osc}_{B_\rho} u \leq c \left(\frac{\rho}{R} \right)^\gamma \text{osc}_{B_R} u \quad (5.32)$$

holds for some $c \equiv c(\mathbf{data}(\Omega_0))$ and all concentric balls $B_\rho \Subset B_R \Subset \Omega_0 \Subset \Omega$ with $R \leq 1$.

2. *If the assumption (1.13) is satisfied, then there exists a Hölder continuity exponent $\gamma \equiv \gamma(\mathbf{data}) \in (0, 1)$ such that*

$$[u]_{0,\gamma;\Omega_0} \leq c(\mathbf{data}(\Omega_0)) \quad (5.33)$$

and the oscillation estimate

$$\text{osc}_{B_\rho} u \leq c \left(\frac{\rho}{R} \right)^\gamma \text{osc}_{B_R} u \quad (5.34)$$

holds for some $c \equiv c(\mathbf{data})$ and all concentric balls $B_\rho \Subset B_R \subset \Omega$ with $R \leq 1$.

Proof. Basically, we shall use De Giorgi's methods to prove the local Hölder continuity of u based on arguments employed in [22, 30]. For the convenience of the reader, we give a detailed proof. Note that, for any given ball $B_R \Subset \Omega$, either

$$\left| \left\{ x \in B_{R/2} : u(x) > \sup_{B_R} u - \frac{1}{2} \text{osc}_{B_R} u \right\} \right| \leq \frac{1}{2} |B_{R/2}| \quad (5.35)$$

or

$$\left| \left\{ x \in B_{R/2} : (-u(x)) > \sup_{B_R} (-u) - \frac{1}{2} \text{osc}_{B_R} u \right\} \right| \leq \frac{1}{2} |B_{R/2}| \quad (5.36)$$

holds true. It is enough to deal with only the case of (5.35) is valid since $-u$ is a local Q -minimizer of the functional \mathcal{P} . The proof falls in three steps. In what follows, let $B_{2R} \equiv B_{2R}(x_0) \subset \Omega_0 \Subset \Omega$ be a fixed ball such that $R \leq 1$. Let us also denote by

$$A(k, \rho) := \{x \in B_\rho : u(x) > k\} \quad \text{and} \quad B(k, \rho) := \{x \in B_\rho : u(x) < k\} \quad (5.37)$$

for every concentric ball $B_\rho \subset B_{2R}$ and $k \in \mathbb{R}$.

Step 1. We suppose that (5.35) is satisfied. Then in this step we prove that, for any $\varepsilon \in (0, 1)$, there exists a natural number $m \equiv m(\mathbf{data}(\Omega_0), \varepsilon) \geq 3$ if (1.12) is assumed, and $m \equiv m(\mathbf{data}, \varepsilon) \geq 3$ if (1.13) is assumed, such that

$$\left| \left\{ x \in B_{R/2} : u(x) > \sup_{B_R} u - \frac{1}{2^m} \text{osc}_{B_R} u \right\} \right| \leq \varepsilon |B_{R/2}|. \quad (5.38)$$

Let $m \geq 3$ be a natural number to be determined in a few lines. For every $i \in \{1, 2, \dots, m\}$, we set

$$k_i := \sup_{B_R} u - \frac{1}{2^i} \text{osc}_{B_R} u, \quad \mathcal{D}_i := A(k_i, R/2) \setminus A(k_{i+1}, R/2)$$

and

$$w_i(x) := \begin{cases} k_{i+1} - k_i & \text{if } u(x) > k_{i+1}, \\ u(x) - k_i & \text{if } k_i < u(x) \leq k_{i+1}, \\ 0 & \text{if } u(x) \leq k_i. \end{cases}$$

Clearly $w_i \in W^{1,\Psi}(B_{R/2})$ with $w_i \equiv 0$ in $B_{R/2} \setminus A(k_1, R/2)$ for all $i \in \{1, \dots, m\}$, and also $|B_{R/2} \setminus A(k_1, R/2)| \geq 1/2|B_{R/2}|$. Then applying Hölder's inequality, Sobolev's inequality and Lemma 2.4, for every $\tau \in (0, 1)$, we have

$$\begin{aligned}
|A(k_{i+1}, R/2)| \Psi_{B_R}^- \left(\frac{k_{i+1} - k_i}{R} \right) &\leq c \int_{A(k_i, R/2)} \Psi_{B_R}^- \left(\frac{w_i}{R} \right) dx \\
&\leq |A(k_i, R/2)|^{\frac{1}{n}} \left(\int_{A(k_i, R/2)} \left[\Psi_{B_R}^- \left(\frac{w_i}{R} \right) \right]^{\frac{n}{n-1}} dx \right)^{\frac{n-1}{n}} \\
&\leq cR \left(\int_{A(k_i, R/2)} \left[\Psi_{B_R}^- \left(\frac{w_i}{R} \right) \right]^{\frac{n}{n-1}} dx \right)^{\frac{n-1}{n}} \\
&\leq c \int_{\mathcal{D}_i} \partial_t \Psi_{B_R}^- \left(\frac{u - k_i}{R} \right) |Du| dx \\
&\leq \tau \int_{\mathcal{D}_i} \Psi_{B_R}^- (|Du|) dx + \frac{c}{\tau^{s(\Psi)}} \int_{\mathcal{D}_i} \Psi_{B_R}^- \left(\frac{u - k_i}{R} \right) dx.
\end{aligned} \tag{5.39}$$

Now we use a Caccioppoli type inequality of Lemma 5.2 in order to have

$$\begin{aligned}
\int_{\mathcal{D}_i} \Psi_{B_R}^- (|Du|) dx &\leq c \int_{A(k_i, R)} \Psi_{B_R}^- \left(\left| \frac{u - k_i}{R} \right| \right) dx \leq c \int_{A(k_i, R)} \Psi_{B_R}^- \left(\frac{\text{osc}_{B_R} u}{2^i R} \right) dx \\
&\leq c \Psi_{B_R}^- \left(\frac{k_{i+1} - k_i}{R} \right) |A(k_i, R)| \leq c \Psi_{B_R}^- \left(\frac{k_{i+1} - k_i}{R} \right) R^n.
\end{aligned}$$

One can see that

$$\int_{\mathcal{D}_i} \Psi_{B_R}^- \left(\frac{u - k_i}{R} \right) dx \leq \int_{\mathcal{D}_i} \Psi_{B_R}^- \left(\frac{k_{i+1} - k_i}{R} \right) dx = \Psi_{B_R}^- \left(\frac{k_{i+1} - k_i}{R} \right) |\mathcal{D}_i|.$$

Using the estimates coming from the last two displays in (5.39), for every $i \in \{1, \dots, m-1\}$, we see

$$|A(k_{m-1}, R/2)| \leq |A(k_{i+1}, R/2)| \leq c\tau R^n + \frac{c}{\tau^{s(\Psi)}} |\mathcal{D}_i|.$$

Summing for $i \in \{1, \dots, m-1\}$, it yields that

$$|A(k_{m-1}, R/2)| \leq \left(c\tau + \frac{c}{(m-1)\tau^{s(\Psi)}} \right) R^n.$$

Now taking small enough $\tau \equiv \tau(\mathbf{data}(\Omega_0), \varepsilon)$ and large enough $m \equiv m(\mathbf{data}(\Omega_0), \varepsilon)$, we arrive at (5.38) when (1.12) is assumed. But in the case that (1.13) is assumed, we choose small enough $\tau \equiv \tau(\mathbf{data}, \varepsilon)$ and large enough $m \equiv m(\mathbf{data}, \varepsilon)$ to conclude (5.38).

Step 2. In this step, we prove that there exists a small positive $\varepsilon_0 \equiv \varepsilon_0(\mathbf{data}(\Omega_0)) \in (0, 1/2^{n+1})$ such that if

$$0 < \nu_0 < \frac{1}{2} \text{osc}_{B_R} u \quad \text{and} \quad \left| \{x \in B_{R/2} : u(x) > \sup_{B_R} u - \nu_0\} \right| \leq \varepsilon_0 |B_{R/2}|, \tag{5.40}$$

then we have

$$\sup_{B_{R/4}} u \leq \sup_{B_R} u - \nu_0/2. \tag{5.41}$$

Now we set the sequences by

$$\rho_i := \frac{R}{4} \left(1 + \frac{1}{2^i} \right) \quad \text{and} \quad k_i := \sup_{B_R} u - \left(\frac{1}{2} + \frac{1}{2^{i+1}} \right) \nu_0 \quad \text{for every } i = 0, 1, 2, \dots,$$

and we define

$$\mathcal{D}_{i+1} := A(k_i, \rho_{i+1}) \setminus A(k_{i+1}, \rho_{i+1}) \quad \text{and} \quad Y_i := \frac{|A(k_i, \rho_i)|}{|B_{R/2}|}.$$

Applying Lemma 5.2 together with (5.40), we discover

$$\begin{aligned} \int_{A(k_i, \rho_{i+1})} \Psi_{B_R}^- (|Du|) dx &\leq c 2^{(i+3)(s(\Psi)+1)} \int_{A(k_i, \rho_i)} \Psi_{B_R}^- \left(\frac{(u - k_i)_+}{R} \right) dx \\ &\leq c 2^{i(s(\Psi)+1)} \Psi_{B_R}^- \left(\frac{\nu_0}{R} \right) |A(k_i, \rho_i)|, \end{aligned}$$

where we have also used the very definition of k_i and that $(u - k_i)_+ \leq \nu_0 \leq \|u\|_{L^\infty(B_R)}$. The last display and the convexity of $\Psi_{B_R}^-$ imply that

$$\begin{aligned} \Psi_{B_R}^- \left(\int_{\mathcal{D}_{i+1}} |Du| dx \right) &\leq \int_{\mathcal{D}_{i+1}} \Psi_{B_R}^- (|Du|) dx \leq c 2^{i(s(\Psi)+1)} \frac{|A(k_i, \rho_i)|}{|D_{i+1}|} \Psi_{B_R}^- \left(\frac{\nu_0}{R} \right) \\ &\leq \Psi_{B_R}^- \left(c 2^{i(s(\Psi)+1)} \frac{|A(k_i, \rho_i)|}{|D_{i+1}|} \frac{\nu_0}{R} \right). \end{aligned}$$

Therefore, we have

$$\int_{\mathcal{D}_{i+1}} |Du| dx \leq c 2^{i(s(\Psi)+1)} \frac{|A(k_i, \rho_i)|}{|D_{i+1}|} \frac{\nu_0}{R}.$$

On the other hand, applying Lemma 2.8 together with $\varepsilon_0 \in (0, 1/2^{n+1})$, we discover

$$\begin{aligned} \int_{\mathcal{D}_{i+1}} |Du| dx &\geq c(k_{i+1} - k_i) |A(k_{i+1}, \rho_{i+1})|^{1-\frac{1}{n}} |B_{\rho_{i+1}} \setminus A(k_i, \rho_{i+1})| \rho_{i+1}^{-n} \\ &\geq c 2^{-i} \nu_0 |A(k_{i+1}, \rho_{i+1})|^{1-\frac{1}{n}} (|B_{R/4}| - \varepsilon_0 |B_{R/2}|) R^{-n} \\ &\geq c 2^{-i} \nu_0 |A(k_{i+1}, \rho_{i+1})|^{1-\frac{1}{n}} \\ &\geq c 2^{-i} \nu_0 R^{n-1} Y_{i+1}^{1-\frac{1}{n}} \end{aligned}$$

for some constant $c \equiv c(\mathbf{data}(\Omega_0))$. Combining last two displays, we conclude

$$Y_{i+1} \leq c_* \left(2^{\frac{n(s(\Psi)+2)}{n-1}} \right)^i Y_i^{1+\frac{1}{n-1}}$$

for some constant $c_* \equiv c_*(\mathbf{data}(\Omega_0))$. Now we apply Lemma 2.7 in order to have $Y_i \rightarrow 0$ as $i \rightarrow \infty$, provided

$$Y_0 = \frac{|A(k_0, R/2)|}{|B_{R/2}|} \leq \varepsilon_0 \leq c_*^{-(n-1)} 2^{-n(n-1)(s(\Psi)+2)}.$$

Therefore, (5.41) is satisfied since

$$\left| A \left(\sup_{B_R} u - \frac{\nu_0}{2}, R/4 \right) \right| = 0.$$

Step 3: Proof of Hölder continuity. Finally, we are now ready to prove a local Hölder continuity of u . For this, let $m \geq 3$ be the natural number satisfying (5.38) for the choice $\varepsilon \equiv \varepsilon_0 \in (0, 1/2^{n+1})$, where ε_0 is determined via (5.40). Then we have

$$\text{osc}_{B_{R/4}} u \leq \left(1 - \frac{1}{2^{m+1}} \right) \text{osc}_{B_R} u$$

with $m \equiv m(\mathbf{data}(\Omega_0))$, whenever $B_{2R} \subset \Omega_0$ is a ball with $R \leq 1$. Clearly, the above display implies that there exists a positive exponent $\gamma \equiv \gamma(\mathbf{data}(\Omega_0)) \in (0, 1)$ such that, for any fixed ball $B_{8R_0} \subset \Omega_0$ with $8R_0 \leq 1$, the following oscillation

$$\text{osc}_{B_R} u \leq c \left(\frac{R}{R_0} \right)^\gamma \text{osc}_{B_{R_0}} u$$

holds with some constant $c \equiv c(\mathbf{data}(\Omega_0))$ for every $R \in (0, R_0]$. Here we note that in the case that the assumption (1.13) is in force, the constants appearing in the above lemma depend only on \mathbf{data} , but otherwise are independent of the subset Ω_0 . Finally, we have shown that

$$u \in C_{\text{loc}}^{0,\gamma}(\Omega_0)$$

if either the assumption (1.12) or (1.13) is satisfied. Therefore by a standard covering argument, the estimates (5.31) and (5.32) are satisfied. Clearly, if (1.13) is assumed instead of (1.12), γ in (5.33) depends only on \mathbf{data} since $\|u\|_{L^\infty(\Omega_0)} \leq \|u\|_{L^\infty(\Omega)}$. The proof is complete. \square

5.4. The Harnack inequality. In this subsection we prove the Harnack inequality for a local Q -minimizer u of the functional \mathcal{P} in (1.1) under one of the assumptions (1.12), (1.13) and (1.14). The analysis similar to the one in the Step 1 of the proof of Theorem 5.2 gives the following lemma.

Lemma 5.4. *Let $u \in W^{1,\Psi}(\Omega)$ be a non-negative local Q -minimizer of the functional \mathcal{P} in (1.1) under the coefficient functions $a(\cdot) \in C^{\omega_a}(\Omega)$ and $b(\cdot) \in C^{\omega_b}(\Omega)$ for ω_a, ω_b being non-negative concave functions vanishing at the origin. Suppose that one of the assumptions (1.12), (1.13) and (1.14) is satisfied. Let $B_{6R} \subset \Omega_0 \Subset \Omega$ be a ball with $6R \leq 1$. Then for any $\tau_1, \tau_2 \in (0, 1)$, there exists a large number m depending on \mathbf{data} and τ_1, τ_2 such that for any $0 < k \leq \|u\|_{L^\infty(B_{3R})}$, if*

$$|\{x \in B_R : u(x) \geq k\}| \geq \tau_1 |B_R| \quad (5.42)$$

holds, then

$$|\{x \in B_{2R} : u(x) \leq 2^{-m}k\}| \leq \tau_2 |B_{2R}|. \quad (5.43)$$

Proof. Let $m \geq 3$ be a large number to be determined later. We set, for $i = 0, 1, \dots, m$,

$$k_i := \frac{k}{2^i}, \quad \mathcal{D}_i := B(k_i, 2R) \setminus B(k_{i+1}, 2R)$$

and

$$w_i(x) := \begin{cases} k_i - k_{i+1} & \text{if } u(x) < k_{i+1}, \\ u(x) - k_{i+1} & \text{if } k_{i+1} \leq u(x) < k_i, \\ 0 & \text{if } u(x) \geq k_i. \end{cases}$$

We observe that $\Psi_{3R}^-(w_i) \in W^{1,1}(B_{2R})$ and $\Psi_{3R}^-(w_i) \equiv 0$ on $B_{2R} \setminus B(k_0, 2R)$ for every $i \in \{0, 1, \dots, m\}$ and $|B_{2R} \setminus B(k_0, 2R)| \geq \tau_1 |B_R|$. Then using Hölder's inequality, Sobolev's inequality and Lemma 2.4, we have that

$$\begin{aligned} B(k_{i+1}, 2R) \Psi_{3R}^- \left(\frac{k_i - k_{i+1}}{3R} \right) &\leq \int_{B(k_i, 2R)} \Psi_{3R}^- \left(\frac{w_i}{3R} \right) dx \\ &\leq |B(k_i, 2R)|^{\frac{1}{n}} \left(\int_{B(k_i, 2R)} \left[\Psi_{3R}^- \left(\frac{w_i}{3R} \right) \right]^{\frac{n}{n-1}} dx \right)^{\frac{n-1}{n}} \\ &\leq cR \left(\int_{B(k_i, 2R)} \left[\Psi_{3R}^- \left(\frac{w_i}{3R} \right) \right]^{\frac{n}{n-1}} dx \right)^{\frac{n-1}{n}} \\ &\leq c \int_{\mathcal{D}_i} (\Psi_{3R}^-)' \left(\frac{u - k_{i+1}}{3R} \right) |Du| dx \\ &\leq \varepsilon \int_{\mathcal{D}_i} \Psi_{3R}^- (|Du|) dx + \frac{c}{\varepsilon^{s(\Psi)}} \int_{\mathcal{D}_i} \Psi_{3R}^- \left(\frac{u - k_{i+1}}{3R} \right) dx \end{aligned} \quad (5.44)$$

for every $\varepsilon \in (0, 1)$ and some constant $c \equiv c(\mathbf{data}, \tau_1)$, where we have used Remark 2.2 that $\Psi_{3R}^- \in \mathcal{N}$ with an index $s(\Psi) = s(G) + s(H_a) + s(H_b)$. It follows from the almost standard Caccioppoli inequality of Lemma

5.2 that

$$\begin{aligned} \int_{\mathcal{D}_i} \Psi_{B_{3R}}^- (|Du|) dx &\leq c \int_{B(k_i, 2R)} \Psi_{3R}^- \left(\frac{k_i - u}{3R} \right) dx \leq c \int_{B(k_i, 2R)} \Psi_{3R}^- \left(\left| \frac{2(k_i - k_{i+1})}{3R} \right| \right) dx \\ &\leq c |B(k_i, 2R)| \Psi_{3R}^- \left(\left| \frac{2(k_i - k_{i+1})}{3R} \right| \right) \leq c R^n \Psi_{3R}^- \left(\left| \frac{k_i - k_{i+1}}{3R} \right| \right), \end{aligned} \quad (5.45)$$

where we have also used the assumption that u is non-negative. Clearly, by the very definition of \mathcal{D}_i , one can see that

$$\int_{\mathcal{D}_i} \Psi_{3R}^- \left(\frac{u - k_{i+1}}{3R} \right) dx \leq \int_{\mathcal{D}_i} \Psi_{3R}^- \left(\frac{k_i - k_{i+1}}{3R} \right) dx \leq c |\mathcal{D}_i| \int_{\mathcal{D}_i} \Psi_{3R}^- \left(\frac{k_i - k_{i+1}}{3R} \right) dx. \quad (5.46)$$

Combining the estimates obtained in (5.44)-(5.46), we find that

$$|B(k_m, 2R)| \leq |B(k_{i+1}, 2R)| \leq c \varepsilon R^n + \frac{c}{\varepsilon^{s(\Psi)}} |\mathcal{D}_i|$$

holds for some constant $c \equiv c(\mathbf{data}, \tau_1)$, whenever $\varepsilon \in (0, 1)$ and $i \in \{0, 1, \dots, m-1\}$. Summing the last inequality above over the index i from 0 to $m-1$ implies

$$|B(k_m, 2R)| \leq c \varepsilon R^n + \frac{c}{\varepsilon^{s(\Psi)} m} |B(k_0, 2R)| \leq \left(c_* \varepsilon + \frac{c_*}{\varepsilon^{s(\Psi)} m} \right) |B_{2R}|$$

for some constant $c_* \equiv c_*(\mathbf{data}, \tau_1)$. Now choosing small enough $\varepsilon \equiv (\mathbf{data}, \tau_1, \tau_2)$ and sufficiently large $m \equiv m(\mathbf{data}, \tau_1, \tau_2)$ such that

$$c_* \varepsilon + \frac{c_*}{\varepsilon^{s(\Psi)} m} \leq \tau_2,$$

we arrive at the desired estimate (5.43). □

Lemma 5.5. *Under the assumptions of Lemma 5.4, let $u \in W^{1, \Psi}(\Omega)$ be a non-negative Q -minimizer of the functional \mathcal{P} in (1.1). Suppose that one of the assumptions (1.12), (1.13) and (1.14) is satisfied. Then for any $\tau \in (0, 1)$, there exists a small $\delta_1 \equiv \delta_1(\mathbf{data}(\Omega_0))$ such that for any $0 < k \leq \|u\|_{L^\infty(B_{3R})}$, if*

$$|\{x \in B_R : u(x) \geq k\}| \geq \tau |B_R| \quad (5.47)$$

holds, then

$$\inf_{B_R} u \geq \delta_1 k. \quad (5.48)$$

Proof. It's enough to prove the lemma for $\tau \in (0, 2^{-(n+1)})$. Let us fix $m_0 \in \mathbb{N}$, and consider the sequences defined by

$$\rho_i := R \left(1 + \frac{1}{2^i} \right) \quad \text{and} \quad k_i := \left(\frac{1}{2} + \frac{1}{2^i} \right) 2^{-m_0} k \quad (i = 0, 1, 2, \dots). \quad (5.49)$$

Next we also define

$$\mathcal{D}_{i+1}^- := B(k_i, \rho_{i+1}) \setminus B(k_{i+1}, \rho_{i+1}) \quad \text{and} \quad Y_i := \frac{|B(k_i, \rho_i)|}{|B_{\rho_i}|}, \quad (5.50)$$

where the definition of $B(k_i, \rho_i)$ has been introduced in (5.37). By using the assumption that u is non-negative, we observe $(u - k_i)_- \leq 2^{-m_0} k$. Then by applying Lemma 5.2, we see

$$\begin{aligned} \int_{B(k_i, \rho_{i+1})} \Psi_{B_{2R}}^- (|Du|) dx &\leq c 2^{(i+3)(s(\Psi)+1)} \int_{B(k_i, \rho_i)} \Psi_{B_{2R}}^- \left(\frac{(u - k_i)_-}{2R} \right) dx \\ &\leq c 2^{(i+3)(s(\Psi)+1)} \Psi_{B_{2R}}^- \left(\frac{2^{-m_0} k}{R} \right) |B(k_i, \rho_i)| \end{aligned}$$

for some constant $c \equiv c(\mathbf{data})$. This estimate together with the convexity of $\Psi_{B_{2R}}^-$ implies

$$\begin{aligned} \Psi_{B_{2R}}^- \left(\int_{\mathcal{D}_{i+1}^-} |Du| dx \right) &\leq \int_{\mathcal{D}_{i+1}^-} \Psi_{B_{2R}}^- (|Du|) dx \leq c 2^{i(s(\Psi)+1)} \frac{|B(k_i, \rho_i)|}{|\mathcal{D}_{i+1}^-|} \Psi_{B_{2R}}^- \left(\frac{2^{-m_0} k}{R} \right) \\ &\leq c \Psi_{B_{2R}}^- \left(2^{i(s(\Psi)+1)} \frac{|B(k_i, \rho_i)|}{|\mathcal{D}_{i+1}^-|} \frac{2^{-m_0} k}{R} \right) \end{aligned}$$

for some constant $c \equiv c(\mathbf{data}(\Omega_0))$. Therefore, using the fact that the function $\Psi_{B_{2R}}^-$ is increasing and Lemma 2.1, we have

$$\int_{\mathcal{D}_{i+1}^-} |Du| dx \leq c 2^{i(s(\Psi)+1)} \frac{|B(k_i, \rho_i)|}{|\mathcal{D}_{i+1}^-|} \frac{2^{-m_0} k}{R}.$$

Now applying Lemma 2.8 together with the fact that $\tau \in (0, 2^{-(n+1)})$, we see

$$\begin{aligned} \int_{\mathcal{D}_{i+1}^-} |Du| dx &\geq c(k_i - k_{i+1}) |B(k_{i+1}, \rho_{i+1})|^{1-\frac{1}{n}} |B_{\rho_{i+1}} \setminus B(k_i, \rho_{i+1})| \rho_{i+1}^n \\ &\geq c 2^{-i} 2^{-m_0} k |B(k_{i+1}, \rho_{i+1})|^{1-\frac{1}{n}} (|B_{2R}| - \tau |B_R|) R^{-n} \\ &\geq c 2^{-i} 2^{-m_0} k |B(k_{i+1}, \rho_{i+1})|^{1-\frac{1}{n}} \\ &\leq c 2^{-i} 2^{-m_0} k R^{n-1} Y_{i+1}^{1-\frac{1}{n}}. \end{aligned}$$

The combination of the last two displays yields

$$Y_{i+1}^{1-\frac{1}{n}} \leq c 2^{i(s(\Psi)+1)} R^{-n} |B(k_i, \rho_i)| \leq c 2^{i(s(\Psi)+1)} Y_i$$

and then we conclude

$$Y_{i+1} \leq c_* 2^{i \frac{(s(\Psi)+1)n}{n-1}} Y_i^{1+\frac{1}{n-1}}$$

for some constant $c \equiv c(\mathbf{data}(\Omega_0))$. Now applying Lemma 5.4, we find a large natural number $m_0 \equiv m_0(\mathbf{data}(\Omega_0))$ such that

$$|\{x \in B_{2R} : u(x) \leq 2^{-m_0} k\}| \leq c_*^{-(n-1)} 2^{-n(n-1)(s(\Psi)+1)}.$$

With keeping the above choice of m_0 , we observe that

$$Y_0 = \frac{|B(k_0, 2R)|}{|B_{2R}|} = \frac{|x \in B_{2R} : u(x) \leq 2^{-m_0} k|}{|B_{2R}|} \leq c_*^{-(n-1)} 2^{-n(n-1)(s(\Psi)+1)}.$$

Now we are at stage in applying Lemma 2.7 to obtain that $Y_i \rightarrow 0$ as $i \rightarrow \infty$, which is equivalent to

$$|B(2^{-(m_0+1)} k, R)| = 0.$$

The last display implies the validity of (5.48) with the choice of $\delta_1 \equiv 2^{-(m_0+1)}$. \square

From Lemma 5.5 and the covering arguments in [62, Section 7], we obtain the following weak Harnack inequality for a local Q -minimizers of the functional \mathcal{P} defined in (1.1). We also refer to [11, 22, 59] for the proof.

Theorem 5.3 (The weak Harnack inequality). *Let $W^{1,\Psi}(\Omega)$ be a local non-negative Q -minimizer of the functional \mathcal{P} defined in (1.1) with the coefficient functions $a(\cdot) \in C^{\omega_a}(\Omega)$ and $b(\cdot) \in C^{\omega_b}(\Omega)$ satisfying (1.12)₂ for functions ω_a, ω_b being concave which vanish at 0. Suppose one of the assumptions (1.12), (1.13) and (1.14) is satisfied. Let $B_{9R} \equiv B_{9R}(x_0) \subset \Omega_0 \Subset \Omega$ be a ball with $9R \leq 1$. Then there exist $q_- > 0$ and a constant c depending on $\mathbf{data}(\Omega_0)$ such that*

$$\inf_{x \in B_R} u(x) \geq \frac{1}{c} \left(\int_{B_{2R}} u^{q_-} dx \right)^{\frac{1}{q_-}}. \quad (5.51)$$

To conclude the result of Theorem 5.4 below, we need to obtain a local sup-estimates for local quasiminimizers of \mathcal{P} .

Lemma 5.6. *Under the assumptions of Lemma 5.4, let $u \in W^{1,\Psi}(\Omega)$ be a non-negative local Q -minimizer of the functional \mathcal{P} in (1.1) with the coefficient functions $a(\cdot) \in C^{\omega_a}(\Omega)$ and $b(\cdot) \in C^{\omega_b}(\Omega)$ satisfying (1.12)₂ for functions ω_a, ω_b being concave which vanish at 0. Suppose that one of the assumptions (1.12), (1.13) and (1.14) is satisfied. Let $B_{9R} \equiv B_{9R}(x_0) \subset \Omega_0 \Subset \Omega$ be a ball with $9R \leq 1$. Then for any $q_+ > 0$, the local estimate holds*

$$\sup_{B_R} u \leq c \left(\int_{B_{2R}} |u|^{q_+} dx \right)^{\frac{1}{q_+}} \quad (5.52)$$

for some constant $c \equiv c(\mathbf{data}(\Omega_0))$.

Proof. The proof consists of two steps. For the convenience, let us consider the scaled functions

$$\bar{u}(x) := \frac{u(x_0 + Rx)}{R} \quad \text{for every } x \in B_4. \quad (5.53)$$

Then the almost standard Caccioppoli inequality (5.17) of Lemma 5.2 can be written in the view of \bar{u} as follows:

$$\int_{B_{r_1}} \Psi_{B_{2R}}^- (|D(\bar{u} - k)_\pm|) dx \leq \frac{c}{(r_2 - r_1)^{s(\Psi)+1}} \int_{B_{r_2}} \Psi_{B_{2R}}^- ((\bar{u} - k)_\pm) dx \quad (5.54)$$

with some constant $c \equiv c(\mathbf{data})$, whenever $B_{r_1} \Subset B_{r_2} \subset B_2(0)$ are concentric balls and $k \in \mathbb{R}$. Next for $1 \leq t \leq s \leq 2$, we set sequences by

$$\rho_i := \left(t + \frac{s-t}{2^i} \right) \quad \text{and} \quad k_i := 2l_0 \left(1 - \frac{1}{2^{i+1}} \right) \quad (5.55)$$

for some constant $d_0 > 0$ to be determined later. We also define

$$\bar{\rho}_i := \frac{\rho_i + \rho_{i+1}}{2} \quad \text{and} \quad Y_i := \frac{1}{\Psi_{B_{2R}}^-(l_0)} \int_{\bar{A}(k_i, \rho_i)} \Psi_{B_{2R}}^- ((u - k_i)_+) dx, \quad (5.56)$$

where

$$\bar{A}(k, \rho) := \{x \in B_\rho : \bar{u} > k\}. \quad (5.57)$$

Let $\eta_i \in C_0^\infty(B_{\bar{\rho}_i})$ be a cut-off function such that $0 \leq \eta_i \leq 1$, $\eta_i \equiv 1$ on $B_{\rho_{i+1}}$ and $|D\eta_i| \leq \frac{c(n)2^i}{(s-t)}$. Then using Hölder's inequality, Sobolev's inequality and Lemma 2.4, we have

$$\begin{aligned}
\Psi_{B_{2R}}^-(l_0)Y_{i+1} &\leq \int_{B_{\bar{\rho}_i}} \Psi_{B_{2R}}^-((\bar{u} - k_{i+1})\eta_i) dx \\
&\leq |\bar{A}(k_{i+1}, \rho_i)|^{\frac{1}{n}} \left(\int_{B_{\bar{\rho}_i}} [\Psi_{B_{2R}}^-((\bar{u} - k_{i+1})_+\eta_i)]^{\frac{n}{n-1}} dx \right)^{\frac{n-1}{n}} \\
&\leq c|\bar{A}(k_{i+1}, \rho_i)|^{\frac{1}{n}} \int_{B_{\bar{\rho}_i}} (\Psi_{B_{2R}}^-)'((\bar{u} - k_{i+1})_+\eta_i) [|D(\bar{u} - k_{i+1})_+|\eta_i + (\bar{u} - k_{i+1})_+|D\eta_i|] dx \\
&\leq c|\bar{A}(k_{i+1}, \rho_i)|^{\frac{1}{n}} \int_{B_{\bar{\rho}_i}} (\Psi_{B_{2R}}^-)'((\bar{u} - k_{i+1})_+) |D(\bar{u} - k_{i+1})_+| dx \\
&\quad + c|\bar{A}(k_{i+1}, \rho_i)|^{\frac{1}{n}} \frac{2^i}{s-t} \int_{B_{\bar{\rho}_i}} (\Psi_{B_{2R}}^-)'((\bar{u} - k_{i+1})_+) (\bar{u} - k_{i+1})_+ dx \\
&\leq c|\bar{A}(k_{i+1}, \rho_i)|^{\frac{1}{n}} \left(\int_{B_{\bar{\rho}_i}} \Psi_{B_{2R}}^- (|D(\bar{u} - k_{i+1})_+|) dx + \frac{2^i}{s-t} \int_{B_{\bar{\rho}_i}} \Psi_{B_{2R}}^-((\bar{u} - k_{i+1})_+) dx \right) \\
&\leq c|\bar{A}(k_{i+1}, \rho_i)|^{\frac{1}{n}} \left(\frac{2^i}{s-t} \right)^{s(\Psi)+1} \int_{B_{\rho_i}} \Psi_{B_{2R}}^-((\bar{u} - k_{i+1})_+) dx
\end{aligned}$$

for some constant $c \equiv c(\mathbf{data})$, where in the last inequality of the above display we also have used (5.54) and (5.57). Now applying Lemma 2.1, we see that

$$\begin{aligned}
|\bar{A}(k_{i+1}, \rho_i)| &\leq \frac{1}{\Psi_{B_{2R}}^-(k_{i+1} - k_i)} \int_{\bar{A}(k_{i+1}, \rho_i)} \Psi_{B_{2R}}^- (\bar{u} - k_i) dx \\
&\leq \frac{1}{\Psi_{B_{2R}}^-(l_0/2^{i+1})} \int_{\bar{A}(k_{i+1}, \rho_i)} \Psi_{B_{2R}}^- (\bar{u} - k_i) dx \\
&\leq \frac{\Psi_{B_{2R}}^-(l_0)}{\Psi_{B_{2R}}^-(l_0/2^{i+1})} Y_i \leq 2^{(i+1)(s(\Psi)+1)} Y_i \leq c \left(\frac{2^i}{s-t} \right)^{s(\Psi)+1} Y_i.
\end{aligned}$$

and

$$\begin{aligned}
\int_{B_{\rho_i}} \Psi_{B_{2R}}^-((\bar{u} - k_{i+1})_+) dx &= \int_{\bar{A}(k_{i+1}, \rho_i)} \Psi_{B_{2R}}^- (\bar{u} - k_{i+1}) dx \\
&\leq \int_{\bar{A}(k_i, \rho_i)} \Psi_{B_{2R}}^- (\bar{u} - k_i) dx = \Psi_{B_{2R}}^-(l_0) Y_i.
\end{aligned}$$

Combining the last three displays, we conclude with the following recursive inequality:

$$Y_{i+1} \leq c_0 \frac{2^{i(1+\frac{1}{n})(s(\Psi)+1)}}{(s-t)^{(1+\frac{1}{n})(s(\Psi)+1)}} Y_i^{1+\frac{1}{n}}$$

for some constant $c_0 \equiv c_0(\mathbf{data})$. Now we are at the stage to apply Lemma 2.7. In turn, we have $Y_i \rightarrow 0$ as $i \rightarrow \infty$, provided

$$Y_0 = \frac{1}{\Psi_{B_{2R}}^-(l_0)} \int_{\bar{A}(l_0, s)} \Psi_{B_{2R}}^- (\bar{u} - l_0) dx \leq \left[\frac{c_0}{(s-t)^{(1+\frac{1}{n})(s(\Psi)+1)}} \right]^{-n} 2^{-n(1+n)(s(\Psi)+1)}.$$

The inequality in the last display is satisfied if we choose $l_0 > 0$ in the following way

$$\Psi_{B_{2R}}^-(l_0) = \frac{c_0^n 2^{n(1+n)(s(\Psi)+1)}}{(s-t)^{(1+n)(s(\Psi)+1)}} \int_{B_s} \Psi_{B_{2R}}^-((\bar{u})_+) dx.$$

Therefore, we obtain $\bar{u} \leq 2l_0$ in B_t . This estimate together with the last display yields

$$\Psi_{B_{2R}}^-\left(\sup_{B_t}(\bar{u})_+\right) \leq \frac{c}{(s-t)^{(1+n)(s(\Psi)+1)}} \int_{B_s} \Psi_{B_{2R}}^-((\bar{u})_+) dx. \quad (5.58)$$

Recalling $\Psi_{B_{2R}}^- \in \mathcal{N}$ with an index $s(\Psi)$ and applying Lemma 2.3 for $\Psi_{B_{2R}}^-$, one can see that $t \mapsto \Psi_{B_{2R}}^-\left(t^{\frac{1}{s(\Psi)+1}}\right)$ is a concave function. Using this one together with Jensen's inequality in (5.58), we see

$$\begin{aligned} \Psi_{B_{2R}}^-\left(\sup_{B_t}(\bar{u})_+\right) &\leq \frac{c}{(s-t)^{(1+n)(s(\Psi)+1)}} \int_{B_s} \Psi_{B_{2R}}^-((\bar{u})_+) dx \\ &= \frac{c}{(s-t)^{(1+n)(s(\Psi)+1)}} \int_{B_s} \Psi_{B_{2R}}^-\left(\left[(\bar{u})_+^{s(\Psi)+1}\right]^{\frac{1}{s(\Psi)+1}}\right) dx \\ &\leq \frac{c}{(s-t)^{(1+n)(s(\Psi)+1)}} \Psi_{B_{2R}}^-\left(\left[\int_{B_s} (\bar{u})_+^{s(\Psi)+1} dx\right]^{\frac{1}{s(\Psi)+1}}\right) \\ &\leq \Psi_{B_{2R}}^-\left(\frac{c}{(s-t)^{(1+n)(s(\Psi)+1)}} \left[\int_{B_s} (\bar{u})_+^{s(\Psi)+1} dx\right]^{\frac{1}{s(\Psi)+1}}\right). \end{aligned}$$

Since $\Psi_{B_{2R}}^-$ is the increasing function, the last display implies

$$\sup_{B_t}(\bar{u})_+ \leq \frac{c}{(s-t)^{(1+n)(s(\Psi)+1)}} \left[\int_{B_s} (\bar{u})_+^{s(\Psi)+1} dx\right]^{\frac{1}{s(\Psi)+1}}.$$

Since $-u$ is a local Q -minimizer of the functional \mathcal{P} , we find

$$\sup_{B_t} |\bar{u}| \leq \frac{c}{(s-t)^{(1+n)(s(\Psi)+1)}} \left[\int_{B_s} |\bar{u}|^{s(\Psi)+1} dx\right]^{\frac{1}{s(\Psi)+1}}.$$

Therefore, for $0 < q_+ < s(\Psi) + 1$, we discover from Young's inequality that

$$\begin{aligned} \sup_{B_t} |\bar{u}| &\leq \frac{c}{(s-t)^{(1+n)(s(\Psi)+1)}} \left(\sup_{B_s} |\bar{u}|\right)^{1-\frac{q_+}{s(\Psi)+1}} \left[\int_{B_s} |\bar{u}|^{q_+} dx\right]^{\frac{1}{s(\Psi)+1}} \\ &\leq \frac{1}{2} \sup_{B_s} |\bar{u}| + \frac{c}{(s-t)^{\frac{(1+n)(s(\Psi)+1)^2}{q_+}}} \left[\int_{B_2} |\bar{u}|^{q_+} dx\right]^{\frac{1}{q_+}} \end{aligned}$$

holds for every $1 \leq t < s \leq 2$. Then we apply Lemma 2.6 for $h(t) = \sup_{B_t} |\bar{u}|$ in order to have

$$\sup_{B_1} |\bar{u}| \leq c \left[\int_{B_2} |\bar{u}|^{q_+} dx\right]^{\frac{1}{q_+}} \quad (5.59)$$

for $c \equiv c(\mathbf{data}, q_+)$. On the other hand, for $q_+ \geq s(\Psi) + 1$, the inequality (5.59) is still valid by using Hölder's inequality. Scaling back as we introduced in (5.53), we arrive at the desired estimate (5.52). \square

Finally, the main result of the this section is the following:

Theorem 5.4. *Let $u \in W^{1,\Psi}(\Omega)$ be a non-negative local Q -minimizer u of the functional \mathcal{P} defined in (1.1) under the coefficient functions $a(\cdot) \in C^{\omega_a}(\Omega)$ and $b(\cdot) \in C^{\omega_b}(\Omega)$ for ω_a, ω_b being non-negative concave functions vanishing at the origin. Suppose that one of the assumptions (1.12), (1.13) and (1.14) is satisfied. For every ball B_R with $B_{9R} \subset \Omega_0$ with $\Omega_0 \Subset \Omega$ being an open subset, there exists a positive constant $c \equiv c(\mathbf{data}(\Omega_0))$ such that*

$$\sup_{B_R} u \leq c \inf_{B_R} u \quad (5.60)$$

holds.

Proof. The proof is essentially based on the results we have obtained so far. In fact, applying Theorem 5.3 and Lemma 5.6 with $q_- = q_+$, we obtain (5.60). \square

Remark 5.2. The results of the above theorem refine the results of [58, Theorem 1.3] without any extra term in (5.60) under our multi-phase settings when the assumptions (1.12) and (1.13) come into play, and see also [57, 59].

5.5. Higher integrability results. Next, we provide a higher integrability result for a local minimizer of the functional \mathcal{F} defined in (1.9).

Theorem 5.5 (Higher Integrability). *Let $u \in W^{1,\Psi}(\Omega)$ be a local minimizer of the functional \mathcal{F} defined in (1.9) under the assumption (1.10). Assume that one of the assumptions (1.12), (1.13) and (1.14) is satisfied. Then there exists a higher integrability exponent $\delta \equiv \delta(\mathbf{data}) \in (0, 1)$ such that the following reverse type Hölder inequality*

$$\left(\int_{B_{R/2}} [\Psi(x, |Du|)]^{1+\delta} dx \right)^{\frac{1}{1+\delta}} \leq c \int_{B_R} \Psi(x, |Du|) dx \quad (5.61)$$

holds for a constant $c \equiv c(\mathbf{data})$, where \mathbf{data} is clarified in (1.19), whenever $B_R \Subset \Omega$ is a ball with $R \leq 1$. In particular, for any open subset $\Omega_0 \Subset \Omega$, it holds that

$$\|\Psi(x, |Du|)\|_{L^{1+\delta}(\Omega_0)} \leq c(\mathbf{data}(\Omega_0)). \quad (5.62)$$

Proof. Let $B_R \Subset \Omega$ be a ball with $R \leq 1$ as in the statement. Since u is a local $Q := L/\nu$ -minimizer of the functional \mathcal{P} in (1.1), we are able to apply Lemma 5.1 with the choices $\rho \equiv R/2$, $r \equiv R$ and $k \equiv (u)_{B_R}$ in order to get

$$\int_{B_{R/2}} \Psi(x, |Du|) dx \leq c \int_{B_R} \Psi\left(x, \left| \frac{u - (u)_{B_R}}{R} \right| \right) dx \quad (5.63)$$

with some constant $c \equiv c(n, s(G), s(H_a), s(H_b), \nu, L)$. Then, applying Remark 4.1 depending on which one of the assumptions (1.12), (1.13) and (1.14) is assumed, we obtain the following reverse Hölder inequality:

$$\int_{B_{R/2}} \Psi(x, |Du|) dx \leq c \left(\int_{B_R} [\Psi(x, |Du|)]^\theta dx \right)^{\frac{1}{\theta}}, \quad (5.64)$$

where $c \equiv c(\mathbf{data})$, and $\theta \in (0, 1)$ is the same appearing in Remark 4.1. At this point (5.61) follows using a variant of Gehring's lemma on reverse Hölder inequalities, see for instance [53, Theorem 6.6]. \square

6. HARMONIC TYPE APPROXIMATION

Here we discuss some important regularity results for the solution to the following Dirichlet boundary value problem:

$$\begin{cases} -\operatorname{div} A_0(Dh) = 0 & \text{in } B_R \\ h \in v + W_0^{1,\Psi_0}(B_R), \end{cases} \quad (6.1)$$

where $B_R \subset \mathbb{R}^n$ is a given ball with $n \geq 2$, $v \in W^{1,\Psi_0}(B_R)$ is a given function, and $A_0 : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is a vector field belonging to $C^0(\mathbb{R}^n) \cap C^1(\mathbb{R}^n \setminus \{0\})$ and satisfies the following ellipticity and coercivity assumptions:

$$\begin{cases} |A_0(z)||z| + |D_z A_0(z)||z|^2 \leq L\Psi_0(|z|) \\ \nu \frac{\Psi_0(|z|)}{|z|^2} |\xi|^2 \leq \langle D_z A_0(z)\xi, \xi \rangle \end{cases} \quad (6.2)$$

for fixed constants $0 < \nu \leq L$, whenever $z \in \mathbb{R}^n \setminus \{0\}$ and $\xi \in \mathbb{R}^n$, in which the function Ψ_0 is given by

$$\Psi_0(t) := G(t) + a_0 H_a(t) + b_0 H_b(t) \quad (6.3)$$

with fixed constants $a_0, b_0 \geq 0$ for every $t \geq 0$. By Lemma 2.2₁, we get the following

$$\frac{1}{s(G) + s(H_a) + s(H_b)} \leq \frac{\Psi_0''(t)t}{\Psi_0'(t)} \leq s(G) + s(H_a) + s(H_b) \quad (6.4)$$

for every $t > 0$, which means that $\Psi_0 \in \mathcal{N}$ with an index $s(\Psi_0) = s(G) + s(H_a) + s(H_b)$. Therefore, we note that the following monotonicity property that

$$\begin{aligned} |V_G(z_1) - V_G(z_2)|^2 + a_0 |V_{H_a}(z_1) - V_{H_a}(z_2)|^2 + b_0 |V_{H_b}(z_1) - V_{H_b}(z_2)|^2 &\approx |V_{\Psi_0}(z_1) - V_{\Psi_0}(z_2)|^2 \\ &\leq c \langle A_0(z_1) - A_0(z_2), z_1 - z_2 \rangle \end{aligned} \quad (6.5)$$

holds with some constant $c \equiv c(n, s(\Psi_0), \nu)$, whenever $z_1, z_2 \in \mathbb{R}^n \setminus \{0\}$, where the map V_Φ for a function $\Phi \in \mathcal{N}$ has been defined in (2.8).

Theorem 6.1. *Let $h \in W^{1,\Psi_0}(B_R)$ be the weak solution to (6.1) under the assumption (6.2). Suppose that there exists a higher integrability exponent $\delta_1 > 0$ such that*

$$\Psi_0(|Dv|) \in L^{1+\delta_1}(B_R) \quad \text{and} \quad \|\Psi_0(|Dv|)\|_{L^1(B_R)} \leq L_0 \quad (6.6)$$

for some constant $L_0 \geq 0$. Then there exists a positive exponent $\delta_0 \leq \delta_1$ depending on $n, s(\Psi_0), \nu, L$ and δ_1 such that the following inequality

$$\left(\int_{B_R} [\Psi_0(|Dh|)]^{1+\delta_0} dx \right)^{\frac{1}{1+\delta_0}} \leq c \left(\int_{B_R} [\Psi_0(|Dv|)]^{1+\delta_0} dx \right)^{\frac{1}{1+\delta_0}} \quad (6.7)$$

holds for some constant $c \equiv c(n, s(\Psi_0), \nu, L, L_0, \delta_1)$.

Proof. First the standard energy estimate implies that

$$\int_{B_R} \Psi_0(|Dh|) dx \leq c \int_{B_R} \Psi_0(|Dv|) dx \leq cL_0 \quad (6.8)$$

holds with some constant $c \equiv c(n, s(\Psi_0), \nu, L)$. For a fixed ball $B_{2\rho} \subset B_R$, let $\eta \in C_0^1(B_{2\rho})$ be a standard cut-off function satisfying $\chi_{B_\rho} \leq \eta \leq \chi_{B_{2\rho}}$ and $|D\eta| \leq 4/\rho$. Let us take the function $\varphi = \eta^{s(\Psi_0)+1} (h - (h)_{B_{2\rho}})$ as a test function in the equation (6.1). Then using the monotonicity property of $A_0(\cdot)$ and Lemma 2.4 with Ψ_0 , we have

$$\begin{aligned} \int_{B_{2\rho}} \eta^{s(\Psi_0)+1} \Psi_0(|Dh|) dx &\leq c \int_{B_{2\rho}} \eta^{s(\Psi_0)} \Psi_0'(|Dh|) \left| \frac{h - (h)_{B_{2\rho}}}{\rho} \right| dx \\ &\leq c \int_{B_{2\rho}} \eta^{s(\Psi_0)} \left((\varepsilon\eta) \Psi_0(|Dh|) + \frac{1}{(\varepsilon\eta)^{s(\Psi_0)}} \Psi_0 \left(\left| \frac{h - (h)_{B_{2\rho}}}{\rho} \right| \right) \right) dx. \end{aligned} \quad (6.9)$$

Choosing ε sufficiently small in the last display, we conclude that

$$\int_{B_\rho} \Psi_0(|Dh|) dx \leq c \int_{B_{2\rho}} \Psi_0 \left(\left| \frac{h - (h)_{B_{2\rho}}}{\rho} \right| \right) dx \quad (6.10)$$

for a constant $c \equiv c(n, s(\Psi_0), \nu, L)$. By applying Lemma 4.1 to $\Phi \equiv \Psi_0$ with $d_0 \equiv 1$, there exists $\theta_0 \equiv \theta_0(n, s(\Psi_0)) \in (0, 1)$ such that

$$\int_{B_\rho} \Psi_0(|Dh|) dx \leq c \int_{B_{2\rho}} \Psi_0 \left(\left| \frac{h - (h)_{B_{2\rho}}}{\rho} \right| \right) dx \leq c \left(\int_{B_{2\rho}} [\Psi_0(|Dh|)]^{\theta_0} dx \right)^{\frac{1}{\theta_0}} \quad (6.11)$$

holds for some constant $c \equiv c(n, s(\Psi_0), \nu, L)$, whenever $B_{2\rho} \subset B_R$ is a ball. Now we prove a version of the last inequality near the boundary of B_R . For this, let $B_{2\rho}(y) \subset \mathbb{R}^n$ be a ball such that $y \in B_R$ and $\frac{1}{10} < \frac{|B_{2\rho}(y) \setminus B_R|}{|B_{2\rho}(y)|}$. We take a test function by $\varphi \equiv \eta^{s(\Psi_0)+1}(h - v)$, where $\eta \in C_0^1(B_{2\rho})$ is a standard cut-off function as before so that $\chi_{B_\rho} \leq \eta \leq \chi_{B_{2\rho}}$ and $|D\eta| \leq 4/\rho$. This choice of φ is admissible since $\text{supp } \varphi \subseteq B_R \cap B_{2\rho}(y)$. Arguing similarly as we have done above, we have

$$\begin{aligned} \int_{B_R \cap B_{2\rho}(y)} \eta^{s(\Psi_0)+1} \Psi_0(|Dh|) dx &\leq c \int_{B_R \cap B_{2\rho}(y)} \eta^{s(\Psi_0)} \Psi_0'(|Dh|) \left| \frac{h - v}{\rho} \right| dx \\ &\quad + c \int_{B_R \cap B_{2\rho}(y)} \eta^{s(\Psi_0)} \Psi_0'(|Dh|) |Dv| dx \\ &\leq c \int_{B_R \cap B_{2\rho}} \eta^{s(\Psi_0)} \left((\varepsilon \eta) \Psi_0(|Dh|) + \frac{1}{(\varepsilon \eta)^{s(\Psi_0)}} \Psi_0 \left(\left| \frac{h - v}{\rho} \right| \right) \right) dx \\ &\quad + c \int_{B_R \cap B_{2\rho}} \eta^{s(\Psi_0)} \left((\varepsilon \eta) \Psi_0(|Dh|) + \frac{1}{(\varepsilon \eta)^{s(\Psi_0)}} \Psi_0(|Dv|) \right) dx. \end{aligned} \quad (6.12)$$

Again choosing ε small enough and reabsorbing the terms, we find that

$$\int_{B_R \cap B_{2\rho}(y)} \eta^{s(\Psi_0)+1} \Psi_0(|Dh|) dx \leq c \int_{B_R \cap B_{2\rho}(y)} \Psi_0 \left(\left| \frac{h - v}{\rho} \right| \right) dx + c \int_{B_R \cap B_{2\rho}(y)} \Psi_0(|Dv|) dx$$

for some constant $c \equiv c(n, s(\Psi_0), \nu, L)$. Redefining $h - v \equiv 0$ on $B_{2\rho}(y) \setminus B_R$, we are able to apply Lemma 4.1 to $\Phi \equiv \Psi_0$ with $d_0 \equiv 1$. In turn, there exists $\theta_0 \equiv \theta_0(n, s(\Psi_0)) \in (0, 1)$ as appearing in (6.11) such that

$$\begin{aligned} \int_{B_R \cap B_{2\rho}(y)} \Psi_0 \left(\left| \frac{h - v}{\rho} \right| \right) dx &\leq \left(\int_{B_R \cap B_{2\rho}(y)} [\Psi_0(|Dh - Dv|)]^{\theta_0} dx \right)^{\frac{1}{\theta_0}} \\ &\leq \left(\int_{B_R \cap B_{2\rho}(y)} [\Psi_0(|Dh|)]^{\theta_0} dx \right)^{\frac{1}{\theta_0}} + c \int_{B_R \cap B_{2\rho}(y)} [\Psi_0(|Dv|)] dx \end{aligned}$$

for some constant $c \equiv c(n, s(\Psi_0))$, where for the last inequality we have used (2.6) and Hölder's inequality. Combining the last two displays and (6.11), we have

$$\int_{B_\rho(y)} [V(x)]^{\frac{1}{\theta_0}} dx \leq c \left(\int_{B_{2\rho}(y)} V(x) dx \right)^{\frac{1}{\theta_0}} + c \int_{B_{2\rho}(y)} U(x) dx \quad (6.13)$$

for some $c \equiv c(n, s(\Psi_0), \nu, L)$, where

$$V(x) := [\Psi_0(|Dh|)]^{\theta_0} \chi_{B_{2\rho}(y)}(x) \quad \text{and} \quad U(x) := \Psi_0(|Dv|) \chi_{B_{2\rho}(y)}(x)$$

for every ball $B_{2\rho}(y) \subset \mathbb{R}^n$ satisfying either $B_{2\rho}(y) \subset B_R$ or $\frac{1}{10} < \frac{|B_{2\rho}(y) \setminus B_R|}{|B_{2\rho}(y)|}$ with $y \in B_R$. Applying a variant of Gehring's lemma and a standard covering argument, we arrive at the desired estimate (6.7). \square

Before going on further, we recall a classical truncation lemma due to [1]. The statement involves the Hardy-Littlewood maximal operator, defined as

$$M(f)(x) := \sup_{B_r(x) \subset \mathbb{R}^n} \int_{B_r(x)} |f(y)| dy, \quad x \in \mathbb{R}^n, \quad (6.14)$$

whenever $f \in L^1_{\text{loc}}(\mathbb{R}^n)$.

Theorem 6.2 ([1]). *Let $B_R \subset \mathbb{R}^n$ be a ball and $f \in W_0^{1,1}(B_R)$. Then, for every $\lambda > 0$, there exists $f_\lambda \in W_0^{1,\infty}(B_R)$ such that*

$$\|Df_\lambda\|_{L^\infty(B_R)} \leq c\lambda \quad (6.15)$$

for some constant c depending only on n . Moreover, it holds that

$$\{x \in B_R : f_\lambda(x) \neq f(x)\} \subset \{x \in B_R : M(|Df(x)|) > \lambda\} \cup \text{negligible set}. \quad (6.16)$$

We notice that in this theorem we may assume that f is defined on \mathbb{R}^n by redefining $f \equiv 0$ on $\mathbb{R}^n \setminus B_R$. We are now ready to state the main result of this section.

Lemma 6.1 (Harmonic type approximation). *Let $B_R \subset \mathbb{R}^n$ be a ball with $R \leq 1$, $\sigma \in (0, 1)$ and $v \in W^{1,\Psi_0}(B_{2R})$ be a function satisfying*

$$\int_{B_{2R}} \Psi_0(|Dv|) dx \leq c_0 \quad (6.17)$$

and

$$\int_{B_R} [\Psi_0(|Dv|)]^{1+\delta_1} dx \leq c_1 \quad (6.18)$$

for some constants $c_0, c_1 \geq 1$ and $\delta_1 > 0$. Suppose that $\Psi_0(1) \geq 1$. We further assume that

$$\left| \int_{B_R} \langle A_0(Dv), D\varphi \rangle dx \right| \leq \sigma \|D\varphi\|_{L^\infty(B_R)} \quad \text{holds for } \varphi \in C_0^\infty(B_R). \quad (6.19)$$

Then there exists $h \in v + W_0^{1,\Psi_0}(B_R)$ such that

$$\int_{B_R} \langle A_0(Dh), D\varphi \rangle dx = 0 \quad \text{for all } \varphi \in C_0^\infty(B_R), \quad (6.20)$$

$$\int_{B_R} [\Psi_0(|Dh|)]^{1+\delta_0} dx \leq c(n, s(\Psi_0), \nu, L, \delta_1, c_0, c_1) \quad (6.21)$$

for some positive $\delta_0 \equiv \delta_0(n, s(\Psi_0), \nu, L, \delta_1)$,

$$\int_{B_R} (|V_G(Dv) - V_G(Dh)|^2 + a_0|V_{H_a}(Dv) - V_{H_a}(Dh)|^2 + b_0|V_{H_b}(Dv) - V_{H_b}(Dh)|^2) dx \leq \bar{c}\sigma^{s_1}, \quad (6.22)$$

and

$$\int_{B_R} \Psi_0\left(\left|\frac{v-h}{R}\right|\right) dx \leq \bar{c}\sigma^{s_0} \quad (6.23)$$

for some constants with dependence as $s_1 \equiv s_1(n, s(\Psi_0), \delta_1, c_0) > 0$, $s_0 \equiv s_0(n, s(\Psi_0), \delta_1, c_0) > 0$ and $\bar{c} \equiv \bar{c}(n, s(\Psi_0), \nu, L, \delta_1, c_0, c_1) \geq 1$.

Proof. By the standard approximation argument, if (6.19) holds for all functions $\varphi \in C_0^\infty(B_R)$, then it also holds for all functions $\varphi \in W_0^{1,\infty}(B_R)$. The proof falls in three steps.

Step 1: Truncation. The standard energy estimate and (6.17) give us

$$\int_{B_R} \Psi_0(|Dh|) dx \leq \int_{B_R} \Psi_0(|Dv|) dx \leq c(n, s(\Psi_0), \nu, L) c_0. \quad (6.24)$$

By applying Theorem 6.1, there exists a positive exponent $\delta_0 \equiv \delta_0(n, s(\Psi_0), \nu, L, \delta_1)$ satisfying

$$\int_{B_R} [\Psi_0(|Dh|)]^{1+\delta_0} dx \leq c \int_{B_R} [\Psi_0(|Dv|)]^{1+\delta_0} dx \leq c(n, s(\Psi_0), \nu, L, \delta_1, c_0, c_1), \quad (6.25)$$

which is (6.21). We now set $f := v - h \in W_0^{1,\Psi_0}(B_R)$ and let $\lambda \geq 1$ to be chosen later. We consider $f_\lambda \in W_0^{1,\infty}(B_R)$ provided by Theorem 6.2, which satisfies (6.15) and (6.16). By these properties, Chebyshev's inequality and then the maximal function theorem for Orlicz spaces (see for instance [51, Proposition 1.2]), we have

$$\begin{aligned} \frac{|\{f \neq f_\lambda\}|}{|B_R|} &\leq \frac{|B_R \cap \{M(|Df|) > \lambda\}|}{|B_R|} \leq \frac{1}{[\Psi_0(\lambda)]^{1+\delta_0}} \int_{B_R} [\Psi_0(M(|Df|))]^{1+\delta_0} dx \\ &\leq \frac{c}{[\Psi_0(\lambda)]^{1+\delta_0}} \int_{B_R} [\Psi_0(|Df|)]^{1+\delta_0} dx \\ &\leq \frac{c}{[\Psi_0(\lambda)]^{1+\delta_0}} \left[\int_{B_R} [\Psi_0(|Dv|)]^{1+\delta_0} dx + \int_{B_R} [\Psi_0(|Dh|)]^{1+\delta_0} dx \right] \leq \frac{c}{[\Psi_0(\lambda)]^{1+\delta_0}} \end{aligned} \quad (6.26)$$

with $c \equiv c(n, s(\Psi_0), \nu, L, \delta_1, c_0, c_1)$, where we have used (6.21) and (6.25). Now we test the equation (6.1) against f_λ to obtain

$$\begin{aligned} \Gamma_1 &:= \int_{B_R} \langle A_0(Dv) - A_0(Dh), Df_\lambda \rangle \chi_{\{f=f_\lambda\}} dx \\ &= \int_{B_R} \langle A_0(Dv), Df_\lambda \rangle dx - \int_{B_R} \langle A_0(Dv) - A_0(Dh), Df_\lambda \rangle \chi_{\{f \neq f_\lambda\}} dx =: \Gamma_2 + \Gamma_3. \end{aligned} \quad (6.27)$$

Next we estimate each term appearing in the last equality. By using (6.5), we have

$$\Gamma_1 \geq c \int_{B_R} [|V_G(Dv) - V_G(Dh)|^2 + a_0 |V_{H_a}(Dv) - V_{H_a}(Dh)|^2 + b_0 |V_{H_b}(Dv) - V_{H_b}(Dh)|^2] \chi_{\{f=f_\lambda\}} dx$$

with $c \equiv c(n, s(\Psi_0))$. Using (6.19), and then (6.15), we get

$$|\Gamma_2| \leq \sigma \|Df_\lambda\|_{L^\infty(B_R)} \leq c(n) \sigma \lambda.$$

For Γ_3 , we fix $\varepsilon \in (0, 1)$ to be chosen later and we estimate

$$\begin{aligned} |\Gamma_3| &\leq \int_{B_R} (|A_0(Dh)| + |A_0(Dv)|) |Df_\lambda| \chi_{\{f \neq f_\lambda\}} dx \\ &\stackrel{(6.2)}{\leq} L \|Df_\lambda\|_{L^\infty(B_R)} \int_{B_R} \left[\frac{\Psi_0(|Dv|)}{|Dv|} + \frac{\Psi_0(|Dh|)}{|Dh|} \right] \chi_{\{f \neq f_\lambda\}} dx \\ &\leq \varepsilon \int_{B_R} [\Psi_0(|Dv|) + \Psi_0(|Dh|)] dx + \frac{c}{\varepsilon^{s(\Psi_0)}} \Psi_0(\|Df_\lambda\|_{L^\infty(B_R)}) \frac{|\{f \neq f_\lambda\}|}{|B_R|} \\ &\leq c \left(\varepsilon + \frac{1}{[\Psi_0(\lambda)]^{\delta_0} \varepsilon^{s(\Psi_0)}} \right) \end{aligned}$$

with some $c \equiv c(n, s(\Psi_0), \nu, L, \delta_1, c_0, c_1)$, where in the last two inequalities we have used Lemma 2.1 together with (6.15) and (6.24). Merging the estimates for Γ_1, Γ_2 and Γ_3 with (6.27), we deduce that

$$\begin{aligned} & \int_{B_R} (|V_G(Dv) - V_G(Dh)|^2 + a_0|V_{H_a}(Dv) - V_{H_a}(Dh)|^2 + b_0|V_{H_b}(Dv) - V_{H_b}(Dh)|^2) \chi_{\{f=f_\lambda\}} dx \\ & \leq c_* \left(\sigma\lambda + \varepsilon + \frac{1}{[\Psi_0(\lambda)]^{\delta_0} \varepsilon^{s(\Psi_0)}} \right) =: S(\sigma, \lambda, \varepsilon) \end{aligned} \quad (6.28)$$

for some constant $c_* \equiv c_*(n, s(\Psi_0), \nu, L, \delta_1, c_0, c_1)$, where $\varepsilon \in (0, 1)$ is still to be chosen later. Now let us use a short notation for the simplicity

$$Z^2 := |V_G(Dv) - V_G(Dh)|^2 + a_0|V_{H_a}(Dv) - V_{H_a}(Dh)|^2 + b_0|V_{H_b}(Dv) - V_{H_b}(Dh)|^2 \quad (6.29)$$

and fix $\theta \in (0, 1)$, again to be chosen later. Hölder's inequality and (6.28) imply

$$\left(\int_{B_R} Z^{2\theta} \chi_{\{f=f_\lambda\}} dx \right)^{\frac{1}{\theta}} \leq S(\sigma, \lambda, \varepsilon). \quad (6.30)$$

Again using Hölder's inequality, we get

$$\begin{aligned} & \left(\int_{B_R} Z^{2\theta} \chi_{\{f \neq f_\lambda\}} dx \right)^{\frac{1}{\theta}} \leq \left(\frac{|\{f \neq f_\lambda\}|}{|B_R|} \right)^{\frac{1-\theta}{\theta}} \int_{B_R} Z^2 dx \\ & \stackrel{(6.26)}{\leq} c[\Psi_0(\lambda)]^{-\frac{(1-\theta)(1+\delta_0)}{\theta}} \int_{B_R} [\Psi_0(|Dv|) + \Psi_0(|Dh|)] dx \stackrel{(6.24)}{\leq} c[\Psi_0(\lambda)]^{-\frac{(1-\theta)(1+\delta_0)}{\theta}} \end{aligned} \quad (6.31)$$

for some constant $c \equiv c(n, s(\Psi_0), \nu, L, \delta_1, c_0, c_1, \theta)$. Consequently, (6.30) and (6.31) yield that

$$\left(\int_{B_R} Z^{2\theta} dx \right)^{\frac{1}{\theta}} \leq c \left(S(\sigma, \lambda, \varepsilon) + [\Psi_0(\lambda)]^{-\frac{(1-\theta)(1+\delta_0)}{\theta}} \right)$$

holds with again $c \equiv c(n, s(\Psi_0), \nu, L, \delta_1, c_0, c_1, \theta)$. Recalling $S(\sigma, \lambda, \varepsilon)$ in (6.28) and using Lemma 2.1, we find

$$\left(\int_{B_R} Z^{2\theta} dx \right)^{\frac{1}{\theta}} dx \leq c \left(\sigma\lambda + \varepsilon + \lambda^{-\delta_0 \left(\frac{1}{s(\Psi_0)} + 1 \right)} \varepsilon^{-s(\Psi_0)} + \lambda^{-\left(\frac{1}{s(\Psi_0)} + 1 \right) \frac{(1-\theta)(1+\delta_0)}{\theta}} \right),$$

where at this moment we have used the assumption that $\Psi_0(1) \geq 1$. Choosing $\lambda = \sigma^{-\frac{1}{2}}$ and $\varepsilon = \sigma^s$ with $s = \frac{\delta_0}{4s(\Psi_0)} \left(\frac{1}{s(\Psi_0)} + 1 \right)$, we obtain

$$\left(\int_{B_R} (|V_G(Dv) - V_G(Dh)|^2 + a_0|V_{H_a}(Dv) - V_{H_a}(Dh)|^2 + b_0|V_{H_b}(Dv) - V_{H_b}(Dh)|^2)^\theta dx \right)^{\frac{1}{\theta}} \leq c\sigma^{m_0} \quad (6.32)$$

with constants $m_0 = \min\left\{\frac{1}{2}, \frac{\delta_0}{4s(\Psi_0)} \left(\frac{1}{s(\Psi_0)} + 1 \right), \left(\frac{1}{s(\Psi_0)} + 1 \right) \frac{(1-\theta)(1+\delta_0)}{2\theta}\right\}$ and $c \equiv c(n, s(\Psi_0), \nu, L, \delta_1, c_0, c_1, \theta)$. Recall that θ is yet to be chosen.

Step 2: Proof of (6.22). By taking θ properly, we can deduce (6.22) from (6.32). Hölder's inequality with conjugate exponents $\left(\frac{2(1+\delta_0)}{1+2\delta_0}, 2(1+\delta_0) \right)$ yields

$$\int_{B_R} Z^2 dx = \int_{B_R} Z \cdot Z dx \leq \left(\int_{B_R} Z^{\frac{2(1+\delta_0)}{1+2\delta_0}} dx \right)^{\frac{1+2\delta_0}{2(1+\delta_0)}} \left(\int_{B_R} Z^{2(1+\delta_0)} dx \right)^{\frac{1}{2(1+\delta_0)}}. \quad (6.33)$$

We now choose $\theta := \frac{1+\delta_0}{1+2\delta_0} \in (0, 1)$ in (6.32) in order to find that

$$\left(\int_{B_R} Z^{\frac{2(1+\delta_0)}{1+2\delta_0}} dx \right)^{\frac{1+2\delta_0}{2(1+\delta_0)}} \leq c\sigma^{\frac{m_0}{2}}. \quad (6.34)$$

On the other hand, recalling (6.29) and (6.25), we have

$$\begin{aligned} & \int_{B_R} Z^{2(1+\delta_0)} dx \\ &= \int_{B_R} (|V_G(Dv) - V_G(Dh)|^2 + a_0|V_{H_a}(Dv) - V_{H_a}(Dh)|^2 + b_0|V_{H_b}(Dv) - V_{H_b}(Dh)|^2)^{1+\delta_0} dx \\ &\leq c \int_{B_R} [\Psi_0(|Dv|)]^{1+\delta_0} dx + c \int_{B_R} [\Psi_0(|Dh|)]^{1+\delta_0} dx \leq c(n, s(\Psi_0), \nu, L, \delta_1, c_0, c_1). \end{aligned} \quad (6.35)$$

We combine the estimates (6.33)-(6.35) to discover

$$\int_{B_R} (|V_G(Dv) - V_G(Dh)|^2 + a_0|V_{H_a}(Dv) - V_{H_a}(Dh)|^2 + b_0|V_{H_b}(Dv) - V_{H_b}(Dh)|^2) dx \leq c\sigma^{s_1}, \quad (6.36)$$

where

$$s_1 = \frac{1}{2} \min \left\{ \frac{1}{2}, \frac{\delta_0}{4s(\Psi_0)} \left(\frac{1}{s(\Psi_0)} + 1 \right), \left(\frac{1}{s(\Psi_0)} + 1 \right) \frac{\delta_0}{2} \right\} \text{ and } c \equiv c(n, s(\Psi_0), \nu, L, \delta_0, c_0, c_1).$$

Step 3: Proof of (6.23). By applying Lemma 4.1 to $\Phi \equiv \Psi_0$ with $d_0 \equiv 1$, we see that there exists $\theta_0 \equiv \theta_0(n, s(\Psi_0)) \in (0, 1)$ such that

$$\begin{aligned} & \int_{B_R} \Psi_0 \left(\left| \frac{v-h}{R} \right| \right) dx \leq c \left(\int_{B_R} [\Psi_0(|Dv - Dh|)]^{\theta_0} dx \right)^{\frac{1}{\theta_0}} \\ &\leq c \left(\int_{B_R} \left([\Psi_0(|Dv| + |Dh|)]^{\frac{1}{2}} \frac{|Dv - Dh|}{(|Dv| + |Dh|)} \right)^{\theta_0} [\Psi_0(|Dv| + |Dh|)]^{\frac{\theta_0}{2}} dx \right)^{\frac{1}{\theta_0}} \\ &\leq c \left(\int_{B_R} \Psi_0(|Dv| + |Dh|) \frac{|Dv - Dh|^2}{(|Dv| + |Dh|)^2} dx \right)^{\frac{1}{2}} \left(\int_{B_R} [\Psi_0(|Dv| + |Dh|)]^{\frac{\theta_0}{2-\theta_0}} dx \right)^{\frac{2-\theta_0}{2\theta_0}} \\ &\leq c \left(\int_{B_R} Z^2 dx \right)^{\frac{1}{2}} \left(\int_{B_R} \Psi_0(|Dv| + |Dh|) dx \right)^{\frac{1}{2}} \\ &\leq c\sigma^{\frac{s_1}{2}} = c\sigma^{s_0} \end{aligned}$$

for some $c \equiv c(n, s(\Psi_0), \nu, L, \delta_1, c_0, c_1)$, where in the last display we have applied Hölder's inequality with conjugate exponents $\left(\frac{2}{\theta_0}, \frac{2}{2-\theta_0} \right)$, and finally used (2.10) with (6.36). This proves (6.23). The proof is complete. \square

7. COMPARISON ESTIMATES

Throughout this section we fix a ball $B_{2R} \equiv B_{2R}(x_0) \subset \Omega_0 \Subset \Omega$ with $R \leq 1$ and some open subset $\Omega_0 \Subset \Omega$. We consider the functional defined by

$$W^{1,1}(B_{2R}) \ni v \mapsto \mathcal{F}_{B_{2R}}(v) := \int_{B_{2R}} F(x, (u)_{B_{2R}}, Dv) dx, \quad (7.1)$$

where u is a local minimizer of the functional \mathcal{F} in (1.9). Now we consider a function $w \in u + W_0^{1,\Psi}(B_R)$ being the solution to the following variational Dirichlet problem:

$$\begin{cases} w \mapsto \min_v \mathcal{F}_{B_{2R}}(v) \\ v \in u + W_0^{1,\Psi}(B_{2R}). \end{cases} \quad (7.2)$$

In the following we shall deal with first comparison estimates in order to remove u -dependence in the original functional \mathcal{F} in (1.9).

Lemma 7.1. *Let $w \in W^{1,\Psi}(B_{2R})$ be the solution to the variational problem (7.2) under the assumptions (1.10), (1.16) and (1.18). Let the coefficient functions $a(\cdot) \in C^{\omega_a}(\Omega)$ and $b(\cdot) \in C^{\omega_b}(\Omega)$ for ω_a, ω_b being non-negative concave functions vanishing at the origin. Assume that one of the assumptions (1.12), (1.13) and (1.14) is satisfied. Then there exists a constant $c \equiv c(\mathbf{data}(\Omega_0))$ such that*

$$\begin{aligned} & \int_{B_{2R}} (|V_G(Du) - V_G(Dw)|^2 + a(x)|V_{H_a}(Du) - V_{H_a}(Dw)|^2 + b(x)|V_{H_b}(Du) - V_{H_b}(Dw)|^2) dx \\ & \leq c\omega(R^\gamma) \int_{B_{2R}} \Psi(x, |Du|) dx \end{aligned} \quad (7.3)$$

holds, where $\gamma \equiv \gamma(\mathbf{data}(\Omega_0))$ is the Hölder exponent determined via Theorem 5.2. Moreover, the following estimates hold true:

$$\int_{B_{2R}} \Psi(x, |Dw|) dx \leq \frac{L}{\nu} \int_{B_{2R}} \Psi(x, |Du|) dx, \quad (7.4)$$

$$\|w\|_{L^\infty(B_{2R})} \leq \|u\|_{L^\infty(B_{2R})}, \quad (7.5)$$

$$\text{osc}_{B_{2R}} w \leq \text{osc}_{B_{2R}} u \quad (7.6)$$

and

$$\int_{B_{2R}} \Psi\left(x, \left|\frac{u-w}{R}\right|\right) dx \leq c[\omega(R^\gamma)]^{\frac{1}{2}} \int_{B_{2R}} \Psi(x, |Du|) dx \quad (7.7)$$

for some constant $c \equiv c(\mathbf{data}(\Omega_0))$, where in the case that (1.14) is considered, γ appearing in (7.3) and (7.7) is the same as in the assumption (1.14).

Proof. The proof is very standard and we shall follow the structure of the proof of [13, Lemma 4]. The Euler-Lagrange equation of the functional $\mathcal{F}_{B_{2R}}$, which is

$$\int_{B_{2R}} \langle D_z F(x, (u)_{B_{2R}}, Dw), D\varphi \rangle dx = 0, \quad (7.8)$$

holds for any function $\varphi \in W_0^{1,\Psi}(B_{2R})$ (see for instance [6, Lemma 5.2]). The minimality and growth condition (1.10) imply that

$$\begin{aligned} \int_{B_{2R}} \Psi(x, |Dw|) dx & \leq \frac{1}{\nu} \int_{B_{2R}} F(x, (u)_{B_{2R}}, Dw) dx \\ & \leq \frac{1}{\nu} \int_{B_{2R}} F(x, (u)_{B_{2R}}, Du) dx \leq \frac{L}{\nu} \int_{B_{2R}} \Psi(x, |Du|) dx, \end{aligned} \quad (7.9)$$

which proves (7.4). Therefore, we conclude with

$$\int_{B_{2R}} \langle D_z F(x, (u)_{B_{2R}}, Dw), Du - Dw \rangle dx = 0. \quad (7.10)$$

Letting $u_{B_{2R}}^+ := \sup_{x \in B_{2R}} u(x)$ and $u_{B_{2R}}^- := \inf_{x \in B_{2R}} u(x)$, the minimality of w yields

$$\mathcal{F}_{B_{2R}}(w) \leq \mathcal{F}_{B_{2R}}(\min\{w, u_{B_{2R}}^+\}) \quad \text{and} \quad \mathcal{F}_{B_{2R}}(w) \leq \mathcal{F}_{B_{2R}}(\max\{w, u_{B_{2R}}^-\}).$$

Consequently, the last display together with (1.10) gives us

$$\int_{B_{2R} \cap \{w \geq u_{B_{2R}}^+\}} \Psi(x, |Dw|) dx = 0 \quad \text{and} \quad \int_{B_{2R} \cap \{w \leq u_{B_{2R}}^-\}} \Psi(x, |Dw|) dx = 0.$$

By coarea formula, we get that

$$\inf_{x \in B_{2R}} u \equiv u_{B_{2R}}^- \leq w(x) \leq u_{B_{2R}}^+ \equiv \sup_{x \in B_{2R}} u(x) \quad \text{a.e. } x \in B_{2R}. \quad (7.11)$$

This proves (7.5) and (7.6). Using (2.15) and (7.10) together with the minimality of u and w , we have that

$$\begin{aligned} & \int_{B_{2R}} (|V_G(Du) - V_G(Dw)|^2 + a(x)|V_{H_a}(Du) - V_{H_a}(Dw)|^2 + b(x)|V_{H_b}(Du) - V_{H_b}(Dw)|^2) dx \\ & \stackrel{(7.10)}{=} \int_{B_{2R}} (|V_G(Du) - V_G(Dw)|^2 + a(x)|V_H(Du) - V_H(Dw)|^2 + b(x)|V_{H_b}(Du) - V_{H_b}(Dw)|^2) dx \\ & \quad + c_* \int_{B_{2R}} \langle D_z F(x, (u)_{B_{2R}}, Dw), Du - Dw \rangle dx \\ & \leq c_* \int_{B_{2R}} [F(x, (u)_{B_{2R}}, Du) - F(x, (u)_{B_{2R}}, Dw)] dx \\ & = c_* \int_{B_{2R}} [F(x, (u)_{B_{2R}}, Du) - F(x, u, Du)] dx + c_* \int_{B_{2R}} [F(x, u, Du) - F(x, w, Dw)] dx \\ & \quad + c_* \int_{B_{2R}} [F(x, w, Dw) - F(x, (w)_{B_{2R}}, Dw)] dx + c_* \int_{B_{2R}} [F(x, (w)_{B_{2R}}, Dw) - F(x, (u)_{B_{2R}}, Dw)] dx \\ & =: c_* \sum_{i=1}^4 I_i \end{aligned} \quad (7.12)$$

with $c_* \equiv c_*(n, s(G), s(H_a), s(H_b), \nu)$. Now we estimate each term I_i for $i \in \{1, 2, 3, 4\}$ in the last display. We have

$$\begin{aligned} I_1 & \stackrel{(1.16)}{\leq} c \int_{B_{2R}} \omega(|u - (u)_{B_{2R}}|) \Psi(x, |Du|) dx \\ & \stackrel{(5.31), (5.33)}{\leq} c\omega(2[u]_{0, \gamma; \Omega_0} R^\gamma) \int_{B_{2R}} \Psi(x, |Du|) dx \\ & \stackrel{(2.1)}{\leq} c(\mathbf{data}(\Omega_0))\omega(R^\gamma) \int_{B_{2R}} \Psi(x, |Du|) dx, \end{aligned} \quad (7.13)$$

where in the last display we have also used the fact that $\omega(\cdot)$ is concave. The minimality of u implies

$$I_2 \leq 0. \quad (7.14)$$

We have therefore

$$\begin{aligned}
I_3 &\stackrel{(1.16)}{\leq} c \int_{B_{2R}} \omega(|w - (w)_{B_{2R}}|) \Psi(x, |Dw|) dx \\
&\leq c \int_{B_R} \omega \left(\operatorname{osc}_{B_{2R}} w \right) \Psi(x, |Dw|) dx \\
&\stackrel{(7.6)}{\leq} c \omega \left(\operatorname{osc}_{B_{2R}} u \right) \int_{B_{2R}} \Psi(x, |Dw|) dx \\
&\stackrel{(5.31), (5.33)}{\leq} c \omega(2[u]_{0,\gamma;\Omega_0} R^\gamma) \int_{B_{2R}} \Psi(x, |Du|) dx \\
&\stackrel{(2.1)}{\leq} c(\mathbf{data}(\Omega_0)) \omega(R^\gamma) \int_{B_{2R}} \Psi(x, |Du|) dx.
\end{aligned} \tag{7.15}$$

Observing that

$$|(w)_{B_{2R}} - (u)_{B_{2R}}| \stackrel{(7.11)}{\leq} \operatorname{osc}_{B_{2R}} u, \tag{7.16}$$

as in the estimate for I_1 , we still have

$$I_4 \leq c \omega(R^\gamma) \int_{B_{2R}} \Psi(x, |Du|) dx. \tag{7.17}$$

Inserting all the estimates obtained for I_i with $i \in \{1, 2, 3, 4\}$ into (7.12) completes the proof of (7.3).

Let us now prove (7.7). By Theorem 4.1 with $d \equiv 1$, there exists $\theta_1 \equiv \theta_1(n, s(G), s(H_a), s(H_b)) \in (0, 1)$ such that

$$\begin{aligned}
J &:= \int_{B_{2R}} \Psi \left(x, \left| \frac{u - w}{R} \right| \right) dx \leq c \left(\int_{B_{2R}} [\Psi(x, |Du - Dw|)]^{\theta_1} dx \right)^{\frac{1}{\theta_1}} \\
&\leq c \left(\int_{B_{2R}} \left([\Psi(x, |Du| + |Dw|)]^{\frac{1}{2}} \frac{|Du - Dw|}{|Du| + |Dw|} \right)^{\theta_1} [\Psi(x, |Du| + |Dw|)]^{\frac{\theta_1}{2}} dx \right)^{\frac{1}{\theta_1}},
\end{aligned} \tag{7.18}$$

where in the last inequality of the last display we have used (2.11) for Ψ . Applying Hölder's inequality with conjugate exponents $\left(\frac{2}{\theta_1}, \frac{2}{2-\theta_1}\right)$ to the right hand side of the last display and (2.10), we get

$$\begin{aligned}
J &\leq c \left(\int_{B_{2R}} \Psi(x, |Du| + |Dw|) \frac{|Du - Dw|^2}{(|Du| + |Dw|)^2} dx \right)^{\frac{1}{2}} \left(\int_{B_{2R}} [\Psi(x, |Du| + |Dw|)]^{\frac{\theta_1}{2-\theta_1}} dx \right)^{\frac{2-\theta_1}{2\theta_1}} \\
&\leq c \left(\int_{B_{2R}} |V_\Psi(x, Du) - V_\Psi(x, Dw)|^2 dx \right)^{\frac{1}{2}} \left(\int_{B_{2R}} \Psi(x, |Du| + |Dw|) dx \right)^{\frac{1}{2}} \\
&\leq c(\mathbf{data}(\Omega_0)) [\omega(R^\gamma)]^{\frac{1}{2}} \int_{B_{2R}} \Psi(x, |Du|) dx,
\end{aligned} \tag{7.19}$$

where in the last inequality of the above display we have used (7.3), and then (7.4). Combining the last two displays we arrive at (7.7). \square

Next we consider the functional defined by

$$W^{1,1}(B_R) \ni v \mapsto \mathcal{F}_c(v) := \int_{B_R} F_c(x, Dv) dx, \quad (7.20)$$

where the density function is given by

$$F_c(x, z) := F_G(x_c, (u)_{B_{2R}}, z) + a(x)F_{H_a}(x_c, (u)_{B_{2R}}, z) + b(x)F_{H_b}(x_c, (u)_{B_{2R}}, z) \quad (7.21)$$

for some fixed point $x_c \in B_R$ and for every $x \in \Omega$ and $z \in \mathbb{R}^n$. Now we consider a function $w_c \in w + W_0^{1,\Psi}(B_R)$ being the solution to the following variational Dirichlet problem:

$$\begin{cases} w_c \mapsto \min_v \mathcal{F}_c(v) \\ v \in w + W_0^{1,\Psi}(B_R), \end{cases} \quad (7.22)$$

where $w \in W^{1,\Psi}(B_{2R})$ is the solution to the variational problem (7.2).

Lemma 7.2. *Let $w_c \in W^{1,\Psi}(B_R)$ be the solution to the variational problem (7.22) under the assumptions (1.10), (1.16) and (1.18). Let the coefficient functions $a(\cdot) \in C^{\omega_a}(\Omega)$ and $b(\cdot) \in C^{\omega_b}(\Omega)$ for ω_a, ω_b being non-negative concave functions vanishing at the origin. Assume that one of the assumptions (1.12), (1.13) and (1.14) is satisfied. Then there exists a constant $c \equiv c(\mathbf{data}(\Omega_0))$ such that*

$$\begin{aligned} & \int_{B_R} (|V_G(Dw) - V_G(Dw_c)|^2 + a(x)|V_{H_a}(Dw) - V_{H_a}(Dw_c)|^2 + b(x)|V_{H_b}(Dw) - V_{H_b}(Dw_c)|^2) dx \\ & \leq c\omega(R) \int_{B_R} \Psi(x, |Du|) dx. \end{aligned} \quad (7.23)$$

Moreover, the following estimates hold true:

$$\int_{B_R} \Psi(x, |Dw_c|) dx \leq \frac{L}{\nu} \int_{B_R} \Psi(x, |Dw|) dx, \quad (7.24)$$

$$\|w_c\|_{L^\infty(B_R)} \leq \|w\|_{L^\infty(B_R)}, \quad (7.25)$$

$$\text{osc}_{B_R} w_c \leq \text{osc}_{B_R} w \quad (7.26)$$

and

$$\int_{B_R} \Psi\left(x, \left|\frac{w - w_c}{R}\right|\right) dx \leq c[\omega(R)]^{\frac{1}{2}} \int_{B_R} \Psi(x, |Dw|) dx \quad (7.27)$$

for some constant $c \equiv c(\mathbf{data})$. Finally, there exists a higher integrability exponent $\delta_0 \equiv \delta_0(\mathbf{data})$ with $\delta_0 \leq \delta$ with δ having been determined via Theorem 5.5, and a constant $c \equiv c(\mathbf{data})$ such that

$$\left(\int_{B_{R/2}} [\Psi(x, |Dw_c|)]^{1+\delta_0} dx \right)^{\frac{1}{1+\delta_0}} \leq c \int_{B_R} \Psi(x, |Dw_c|) dx. \quad (7.28)$$

Proof. Essentially, the proof is similar to the proof of Lemma 7.1. The estimates (7.24)-(7.26) can be obtained as for (7.4)-(7.6). We now focus on proving (7.23). The Euler-Lagrange equation arising from the functional \mathcal{F}_c defined in (7.20)

$$\int_{B_R} \langle D_z F_c(x, Dw_c), D\varphi \rangle dx = 0 \quad (7.29)$$

is valid, whenever $\varphi \in W_0^{1,\Psi}(B_R)$. Then using (1.16), we have

$$\begin{aligned}
& \int_{B_R} (|V_G(Dw) - V_G(Dw_c)|^2 + a(x)|V_{H_a}(Dw) - V_{H_a}(Dw_c)|^2 + b(x)|V_{H_b}(Dw) - V_{H_b}(Dw_c)|^2) dx \\
& \leq c \int_{B_R} \langle D_z F_c(x, Dw) - D_z F_c(x, Dw_c), Dw - Dw_c \rangle dx \\
& \leq c \int_{B_R} |D_z F_G(x_c, (u)_{B_{2R}}, Dw) - D_z F_G(x, (u)_{B_{2R}}, Dw)| |Dw - Dw_c| dx \\
& \quad + c \int_{B_R} a(x) |D_z F_{H_a}(x_c, (u)_{B_{2R}}, Dw) - D_z F_{H_a}(x, (u)_{B_{2R}}, Dw)| |Dw - Dw_c| dx \\
& \quad + c \int_{B_R} b(x) |D_z F_{H_b}(x_c, (u)_{B_{2R}}, Dw) - D_z F_{H_b}(x, (u)_{B_{2R}}, Dw)| |Dw - Dw_c| dx \\
& \leq c\omega(R) \int_{B_R} \Psi(x, |Dw|) dx
\end{aligned} \tag{7.30}$$

for some constant $c \equiv c(n, s(G), s(H_a), s(H_b), \nu, L)$. This proves (7.23), and (7.27) follows from this estimate together with applying the arguments used in (7.18)-(7.19). Since w_c is a L/ν -minimizer of the functional \mathcal{F}_c defined in (7.22), we are able to apply Lemma 5.1 with the choices of $v \equiv w_c$, $\rho \equiv R/2$, $r \equiv R$ and $k \equiv (w_c)_{B_R}$. In turn, it gives us that

$$\int_{B_{R/2}} \Psi(x, |Dw_c|) dx \leq c \int_{B_R} \Psi\left(x, \left| \frac{w_c - (w_c)_{B_R}}{R} \right| \right) dx \tag{7.31}$$

holds with $c \equiv c(n, s(G), s(H_a), s(H_b), L, \nu)$. Then applying Remark 4.1, there exists a positive exponent $\theta \equiv \theta(n, s(G), s(H_a), s(H_b)) \in (0, 1)$ such that

$$\int_{B_{R/2}} \Psi(x, |Dw_c|) dx \leq c\bar{\kappa}_{sp} \left[\int_{B_R} [\Psi(x, |Dw_c|)]^\theta dx \right]^{\frac{1}{\theta}} \tag{7.32}$$

holds with some constant $c \equiv c(n, s(G), s(H_a), s(H_b), L, \nu, \omega_a(1), \omega_b(1))$, where

$$\bar{\kappa}_{sp} = \begin{cases} 1 + ([a]_{\omega_a} + [b]_{\omega_b}) \left(\lambda_1 + \lambda_1 \left(\int_{B_R} G(|Dw_c|) dx \right)^{\frac{1}{n}} \right) & \text{if (1.12) is considered,} \\ 1 + ([a]_{\omega_a} + [b]_{\omega_b}) \left(\lambda_2 + \lambda_2 \|w_c\|_{L^\infty(B_R)} \right) & \text{if (1.13) is considered,} \\ 1 + ([a]_{\omega_a} + [b]_{\omega_b}) \left(\lambda_3 + \lambda_3 \left[R^{-\gamma} \text{osc}_{B_R} w_c \right]^{\frac{1}{1-\gamma}} \right) & \text{if (1.14) is considered.} \end{cases} \tag{7.33a}$$

$$\tag{7.33b}$$

$$\tag{7.33c}$$

Furthermore, taking into account (7.4)-(7.6) and (7.24)-(7.26) in the last display, we conclude that

$$\int_{B_{R/2}} \Psi(x, |Dw_c|) dx \leq c \left[\int_{B_R} [\Psi(x, |Dw_c|)]^\theta dx \right]^{\frac{1}{\theta}} \tag{7.34}$$

holds for some constant $\theta \equiv \theta(n, s(G), s(H_a), s(H_b)) \in (0, 1)$ and $c \equiv c(\mathbf{data})$. The estimate (7.28) follows from applying a variant of Gehring's lemma. \square

To go further let us introduce the excess functional defined by

$$E(v, B_r) := (\Psi_{B_{2r}}^-)^{-1} \left(\int_{B_r} \Psi_{B_{2r}}^- \left(\left| \frac{v - (v)_{B_r}}{2r} \right| \right) dx \right) \quad (7.35)$$

for any function $v \in L^1(B_{2r})$ and a ball $B_{2r} \subset \Omega$, where we note that $(\Psi_{B_{2r}}^-)^{-1}$ is the inverse function of $\Psi_{B_{2r}}^-$. By the convexity of $\Psi_{B_{2r}}^-$ together with Lemma 2.1 and Remark 2.2, one can see that

$$E(v, B_r) \leq c (\Psi_{B_{2r}}^-)^{-1} \left(\int_{B_r} \Psi_{B_{2r}}^- \left(\left| \frac{v - v_0}{2r} \right| \right) dx \right) \quad (7.36)$$

for some constant $c \equiv c(s(G) + s(H_a) + s(H_b))$, whenever $v_0 \in \mathbb{R}$ is an arbitrary number.

Lemma 7.3. *Let $u \in W^{1,\Psi}(\Omega)$ be a local minimizer of the functional \mathcal{F} defined in (1.9) under the assumptions (1.10), (1.16) and (1.18). Let $w_c \in W^{1,\Psi}(B_R)$ be the solution to the variational problem (7.22). If one of the assumptions (1.22a)-(1.22e) is satisfied, then for every $\varepsilon^* \in (0, 1)$, there exists a positive radius*

$$R^* \equiv R^*(\mathbf{data}(\Omega_0), \varepsilon^*) \quad (7.37)$$

such that

$$\int_{B_{\tau R}} \Psi_{B_R}^- \left(\left| \frac{w_c - (w_c)_{B_{\tau R}}}{\tau R} \right| \right) dx \leq c \left(1 + \tau^{-(n+s(\Psi)+1)} \varepsilon^* \right) \int_{B_{R/2}} \Psi_{B_R}^- \left(\left| \frac{w_c - (w_c)_{B_{R/2}}}{R} \right| \right) dx \quad (7.38)$$

for some constant $c \equiv c(\mathbf{data}(\Omega_0))$, whenever $\tau \in (0, 1/16)$ and $R \leq R^*$.

Proof. We assume $E(w_c, B_{R/2}) > 0$, otherwise (7.38) is trivial. For the sake of simplicity during the proof, we write

$$E(R) := E(w_c, B_{R/2}) = (\Psi_{B_R}^-)^{-1} \left(\int_{B_{R/2}} \Psi_{B_R}^- \left(\left| \frac{w_c - (w_c)_{B_{R/2}}}{R} \right| \right) dx \right). \quad (7.39)$$

The proof falls in several delicate steps.

Step 1: Initial settings on w_c . Applying Lemma 5.2 to $B_{R/2}$ with $k \equiv (w_c)_{B_{R/2}}$, we have

$$\int_{B_{R/4}} \Psi(x, |Dw_c|) dx \leq c \int_{B_{R/2}} \Psi_{B_R}^- \left(\left| \frac{w_c - (w_c)_{B_{R/2}}}{R} \right| \right) dx \quad (7.40)$$

for some constant $c \equiv c(\mathbf{data})$. Moreover, it follows from Lemma 7.2 that there exists a higher integrability exponent $\delta_0 \equiv \delta_0(\mathbf{data})$ such that

$$\left(\int_{B_{R/8}} [\Psi(x, |Dw_c|)]^{1+\delta_0} dx \right)^{\frac{1}{1+\delta_0}} \leq c \int_{B_{R/4}} \Psi(x, |Dw_c|) dx \quad (7.41)$$

for a constant $c \equiv c(\mathbf{data})$.

Step 2: Scaling. We set the scaled functions of $w_c(\cdot)$, $a(\cdot)$ and $b(\cdot)$ in the ball B_1 by

$$\begin{cases} \bar{w}_c(x) := \frac{w_c(x_0 + Rx) - (w_c)_{B_{R/2}}}{E(R)R}, \\ \bar{a}(x) := a(x_0 + Rx) \frac{H_a(E(R))}{\Psi_{B_R}^-(E(R))} \quad \text{and} \quad \bar{b}(x) := b(x_0 + Rx) \frac{H_b(E(R))}{\Psi_{B_R}^-(E(R))}. \end{cases} \quad (7.42a)$$

$$\quad (7.42b)$$

for every $x \in B_1$. Now we define the control function and energy integrand associated to our scaling in (7.42a)-(7.42b) as

$$\begin{cases} \bar{\Psi}(x, |z|) := \bar{G}(|z|) + \bar{a}(x)\bar{H}_a(|z|) + \bar{b}(x)\bar{H}_b(|z|), \end{cases} \quad (7.43a)$$

$$\bar{F}(x, z) := \bar{F}_G(z) + \bar{a}(x)\bar{F}_{H_a}(z) + \bar{b}(x)\bar{F}_{H_b}(z), \quad (7.43b)$$

$$\begin{cases} \bar{F}_G(z) := \frac{F_G(x_c, (u)_{B_{2R}}, E(R)z)}{\Psi_{B_R}^-(E(R))}, \quad \bar{F}_{H_a}(z) := \frac{F_{H_a}(x_c, (u)_{B_{2R}}, E(R)z)}{H_a(E(R))}, \end{cases} \quad (7.43c)$$

$$\begin{cases} \bar{F}_{H_b}(x, z) := \frac{F_{H_b}(x_c, (u)_{B_{2R}}, E(R)z)}{H_b(E(R))} \quad \text{and} \quad \bar{A}(x, z) := D_z \bar{F}(x, z) \end{cases} \quad (7.43d)$$

for every $x \in B_1$ and $z \in \mathbb{R}^n$, where the point $x_c \in B_R$ has been fixed in (7.22) and

$$\bar{G}(t) := \frac{G(E(R)t)}{\Psi_{B_R}^-(E(R))}, \quad \bar{H}_a(t) := \frac{H_a(E(R)t)}{H_a(E(R))}, \quad \bar{H}_b(t) := \frac{H_b(E(R)t)}{H_b(E(R))} \quad (7.44)$$

for every $t \geq 0$. Clearly, one can see that $\bar{G}, \bar{H}_a, \bar{H}_b \in \mathcal{N}$ with indices $s(G), s(H_a), s(H_b)$, respectively, and also that

$$\bar{G}(1) \leq 1, \quad \bar{H}_a(1) = 1 \quad \text{and} \quad \bar{H}_b(1) = 1. \quad (7.45)$$

Then one can check that the function \bar{w}_c minimizes the following functional

$$W^{1, \bar{\Psi}}(B_1) \ni v \mapsto \int_{B_1} \bar{F}(x, Dv) dx, \quad (7.46)$$

where the functions $\bar{\Psi}(\cdot)$ and $\bar{F}(\cdot)$ have been defined in (7.43a) and (7.43b), respectively. The Euler-Lagrange equation associated to the functional in (7.46) becomes

$$\oint_{B_1} \langle \bar{A}(x, D\bar{w}_c), D\varphi \rangle dx = \oint_{B_1} \langle D_z \bar{F}(x, D\bar{w}_c), D\varphi \rangle dx = 0 \quad (7.47)$$

for every $\varphi \in W_0^{1, \bar{\Psi}}(B_1)$. By the assumptions (1.10) and (1.16) via elementary computations, we have the following structure condition in the scaled settings:

$$\begin{cases} \nu \bar{\Psi}(x, |z|) \leq \bar{F}(x, z) \leq L \bar{\Psi}(x, |z|), \end{cases} \quad (7.48a)$$

$$\begin{cases} |\bar{A}(x, z)||z| + |D_z \bar{A}(x, z)||z|^2 \leq L \bar{\Psi}(x, |z|), \end{cases} \quad (7.48b)$$

$$\begin{cases} \nu \frac{\bar{\Psi}(x, |z|)}{|z|^2} |\xi|^2 \leq \langle D_z \bar{A}(x, z) \xi, \xi \rangle \end{cases} \quad (7.48c)$$

hold true for every $x \in B_1$ and $z \in \mathbb{R}^n \setminus \{0\}$.

Step 3: Freezing. Now we consider frozen functional and vector field associated to $\bar{F}(\cdot)$ and $\bar{A}(\cdot)$ defined in (7.43b)-(7.43d). Let $\bar{x}_a, \bar{x}_b \in \bar{B}_1$ be points such that $\bar{a}(\bar{x}_a) = \inf_{x \in B_1} \bar{a}(x)$ and $\bar{b}(\bar{x}_b) = \inf_{x \in B_1} \bar{b}(x)$. Then we denote by

$$\bar{F}_0(z) := \bar{F}_G(z) + \bar{a}(\bar{x}_a)\bar{F}_{H_a}(z) + \bar{b}(\bar{x}_b)\bar{F}_{H_b}(z), \quad (7.49)$$

$$\bar{A}_0(z) := D_z \bar{F}_0(z) \quad (7.50)$$

and

$$\bar{\Psi}_0(t) := \bar{G}(t) + \bar{a}(\bar{x}_a)\bar{H}_a(t) + \bar{b}(\bar{x}_b)\bar{H}_b(t) \quad (7.51)$$

for every $x \in B_1$, $z \in \mathbb{R}^n$ and $t \geq 0$. By the very definition in (7.43a)-(7.43d), straightforwardly one can see

$$\bar{\Psi}_0(1) = 1. \quad (7.52)$$

In our new scaled settings, we now consider the functional

$$W^{1, \bar{\Psi}_0}(B_{1/8}) \ni v \mapsto \int_{B_{1/8}} \bar{F}_0(Dv) dx. \quad (7.53)$$

We observe that the newly defined integrand $\bar{F}_0(\cdot)$ and vector field $\bar{A}_0(\cdot)$ satisfy the growth and ellipticity conditions as

$$\begin{cases} \nu \bar{\Psi}_0(|z|) \leq \bar{F}_0(z) \leq L \bar{\Psi}_0(|z|), \end{cases} \quad (7.54a)$$

$$\begin{cases} |\bar{A}_0(z)||z| + |D_z \bar{A}_0(z)||z|^2 \leq L \bar{\Psi}_0(|z|), \end{cases} \quad (7.54b)$$

$$\begin{cases} \nu \frac{\bar{\Psi}_0(|z|)}{|z|^2} |\xi|^2 \leq \langle D_z \bar{A}_0(z) \xi, \xi \rangle \end{cases} \quad (7.54c)$$

for every $z \in \mathbb{R}^n \setminus \{0\}$ and $\xi \in \mathbb{R}^n$. Therefore, the estimates (7.40) and (7.41) are written in the view of \bar{w}_c as

$$\int_{B_{1/4}} \bar{\Psi}(x, |D\bar{w}_c|) dx + \left(\int_{B_{1/8}} [\bar{\Psi}(x, |D\bar{w}_c|)]^{1+\delta_0} dx \right)^{\frac{1}{1+\delta_0}} \leq c(\mathbf{data}). \quad (7.55)$$

Step 4: Harmonic type approximation. Let $\varphi \in W_0^{1,\infty}(B_{1/8})$. Using (7.47), we see

$$\begin{aligned} I_0 &:= \left| \int_{B_{1/8}} \langle \bar{A}_0(D\bar{w}_c), D\varphi \rangle dx \right| = \left| \int_{B_{1/8}} \langle \bar{A}_0(D\bar{w}_c) - \bar{A}(x, D\bar{w}_c), D\varphi \rangle dx \right| \\ &\leq \int_{B_{1/8}} |\bar{A}_0(D\bar{w}_c) - \bar{A}(x, D\bar{w}_c)| dx \|D\varphi\|_{L^\infty(B_{1/8})} =: I_1 \|D\varphi\|_{L^\infty(B_{1/8})}. \end{aligned} \quad (7.56)$$

Now using (1.16), we see

$$I_1 \leq L \int_{B_{1/8}} |\bar{a}(x) - \bar{a}(\bar{x}_a)| \frac{\bar{H}_a(|D\bar{w}_c|)}{|D\bar{w}_c|} dx + L \int_{B_{1/8}} |\bar{b}(x) - \bar{b}(\bar{x}_b)| \frac{\bar{H}_b(|D\bar{w}_c|)}{|D\bar{w}_c|} dx =: L(I_{11} + I_{12}). \quad (7.57)$$

Now we estimate the terms appearing in the last display. In turn, using (2.7), (7.45) and (7.55), we have

$$\begin{aligned} I_{11} &\leq c \int_{B_{1/8}} |\bar{a}(x) - \bar{a}(\bar{x}_a)| \left([\bar{H}_a(|D\bar{w}_c|)]^{\frac{1}{s(H_a)+1}} + [\bar{H}_a(|D\bar{w}_c|)]^{\frac{s(H_a)}{s(H_a)+1}} \right) dx \\ &\leq c \|\bar{a} - \bar{a}(\bar{x}_a)\|_{L^\infty(B_{1/8})}^{\frac{s(H_a)}{s(H_a)+1}} \left(\int_{B_{1/8}} \bar{a}(x) \bar{H}_a(|D\bar{w}_c|) dx \right)^{\frac{1}{s(H_a)+1}} \\ &\quad + c \|\bar{a} - \bar{a}(\bar{x}_a)\|_{L^\infty(B_{1/8})}^{\frac{1}{s(H_a)+1}} \left(\int_{B_{1/8}} \bar{a}(x) \bar{H}_a(|D\bar{w}_c|) dx \right)^{\frac{s(H_a)}{s(H_a)+1}} \\ &\leq c(\mathbf{data}) \left(\|\bar{a} - \bar{a}(\bar{x}_a)\|_{L^\infty(B_{1/8})}^{\frac{1}{s(H_a)+1}} + \|\bar{a} - \bar{a}(\bar{x}_a)\|_{L^\infty(B_{1/8})}^{\frac{s(H_a)}{s(H_a)+1}} \right), \end{aligned} \quad (7.58)$$

where we have used also Hölder's inequality and the fact that $\bar{a}(\bar{x}_a) \leq \bar{a}(x)$ for every $x \in B_1$. In a similar way, we have

$$I_{12} \leq c(\mathbf{data}) \left(\|\bar{b} - \bar{b}(\bar{x}_b)\|_{L^\infty(B_{1/8})}^{\frac{1}{s(H_b)+1}} + \|\bar{b} - \bar{b}(\bar{x}_b)\|_{L^\infty(B_{1/8})}^{\frac{s(H_b)}{s(H_b)+1}} \right). \quad (7.59)$$

Inserting those estimates into (7.57) and then (7.56), we find

$$\begin{aligned} I_0 &\leq c(\mathbf{data}) \left(\|\bar{a} - \bar{a}(\bar{x}_a)\|_{L^\infty(B_{1/8})}^{\frac{1}{s(H_a)+1}} + \|\bar{a} - \bar{a}(\bar{x}_a)\|_{L^\infty(B_{1/8})}^{\frac{s(H_a)}{s(H_a)+1}} \right) \\ &\quad + c(\mathbf{data}) \left(\|\bar{b} - \bar{b}(\bar{x}_b)\|_{L^\infty(B_{1/8})}^{\frac{1}{s(H_b)+1}} + \|\bar{b} - \bar{b}(\bar{x}_b)\|_{L^\infty(B_{1/8})}^{\frac{s(H_b)}{s(H_b)+1}} \right). \end{aligned} \quad (7.60)$$

Now we estimate the terms $\|\bar{a} - \bar{a}(\bar{x}_a)\|_{L^\infty(B_{1/8})}$ and $\|\bar{b} - \bar{b}(\bar{x}_b)\|_{L^\infty(B_{1/8})}$ depending on which assumption of (1.22a)-(1.22e) comes into play. Recalling the definition of $\bar{a}(\cdot)$, $\bar{b}(\cdot)$ in (7.42b) and the excess functional in (7.39), we have

$$I_a := \|\bar{a} - \bar{a}(\bar{x}_a)\|_{L^\infty(B_{1/8})} \leq c\omega_a(R) \frac{H_a(E(R))}{\Psi_{B_R}^-(E(R))} \quad (7.61)$$

and

$$I_b := \|\bar{b} - \bar{b}(\bar{x}_b)\|_{L^\infty(B_{1/8})} \leq c\omega_b(R) \frac{H_b(E(R))}{\Psi_{B_R}^-(E(R))}. \quad (7.62)$$

Case 1: Assumption (1.22a) is in force. The assumption (1.22a)₂ implies that for any $\varepsilon \in (0, 1)$ there exists $\mu_1 > 0$ depending on ε such that

$$\Lambda(t, G^{-1}(t^{-n})) \leq \varepsilon \quad \text{for every } t \in (0, \mu_1). \quad (7.63)$$

Then using this one and (1.12), we continue to estimate I_a in (7.61) as

$$\begin{aligned} I_a &\leq c\omega_a(R) \frac{(H_a \circ G^{-1})(\Psi_{B_R}^-(E(R)))}{\Psi_{B_R}^-(E(R))} \\ &\leq c\omega_a(R)\varepsilon \left(1 + \frac{1}{\omega_a([\Psi_{B_R}^-(E(R))]^{-\frac{1}{n}})}\right) + c\omega_a(R) \left(1 + \frac{1}{\omega_a(\mu_1)}\right) \end{aligned} \quad (7.64)$$

with $c \equiv c([a]_{\omega_a}, \lambda_1)$, where we have used the fact that $(\Psi_{B_R}^-)^{-1}(t) \leq G^{-1}(t)$ for every $t \geq 0$. Using (2.2) together with the energy estimates (7.24) and (7.4), we observe that

$$\begin{aligned} \frac{1}{\omega_a([\Psi_{B_R}^-(E(R))]^{-\frac{1}{n}})} &\leq \frac{c}{\omega_a(R)} + \frac{c}{\omega_a(R)} \int_{B_{R/2}} \Psi_{B_R}^- \left(\left| \frac{w_c - (w_c)_{B_{R/2}}}{R} \right| \right) dx \\ &\leq \frac{c}{\omega_a(R)} + \frac{c}{\omega_a(R)} \int_{B_{2R}} \Psi(x, |Du|) dx \leq \frac{c(\mathbf{data})}{\omega_a(R)}. \end{aligned} \quad (7.65)$$

Combining the last two displays, we conclude

$$I_a \leq c \left(\varepsilon + \omega_a(R) \left(1 + \frac{1}{\omega_a(\mu_1)}\right) \right) \quad (7.66)$$

with some constant $c \equiv c(\mathbf{data})$. In the same manner, we see

$$I_b \leq c \left(\varepsilon + \omega_b(R) \left(1 + \frac{1}{\omega_b(\mu_1)}\right) \right) \quad (7.67)$$

for some constant $c \equiv c(\mathbf{data})$. Therefore, inserting the estimates in the last two displays into (7.60) and recalling (7.56), we have

$$\left| \int_{B_{1/8}} \langle \bar{A}_0(D\bar{w}_c), D\varphi \rangle dx \right| \leq c(\mathbf{data}) p_1(\varepsilon, R) \|D\varphi\|_{L^\infty(B_{1/8})}, \quad (7.68)$$

where

$$\begin{aligned} p_1(\varepsilon, R) &:= \left[\varepsilon + \omega_a(R) \left(1 + \frac{1}{\omega_a(\mu_1)}\right) \right]^{\frac{1}{s(H_a)+1}} + \left[\varepsilon + \omega_a(R) \left(1 + \frac{1}{\omega_a(\mu_1)}\right) \right]^{\frac{s(H_a)}{s(H_a)+1}} \\ &\quad + \left[\varepsilon + \omega_b(R) \left(1 + \frac{1}{\omega_b(\mu_1)}\right) \right]^{\frac{1}{s(H_b)+1}} + \left[\varepsilon + \omega_b(R) \left(1 + \frac{1}{\omega_b(\mu_1)}\right) \right]^{\frac{s(H_b)}{s(H_b)+1}}. \end{aligned} \quad (7.69)$$

Case 2: Assumption (1.22b) is in force. From the assumption (1.22b)₂ it holds that for every $\varepsilon \in (0, 1)$ there exists $\mu_2 > 0$ depending on ε such that

$$\Lambda\left(t, \frac{1}{t}\right) \leq \varepsilon \quad \text{for every } t \in (0, \mu_2). \quad (7.70)$$

Then by the very definition of $\Psi_{B_R}^-$ in (2.3) together with (7.70) and (1.13), we have

$$\begin{aligned} I_a &\leq c\omega_a(R) \frac{H_a(E(R))}{G(E(R))} \\ &\leq c\omega_a(R)\varepsilon \left(1 + \frac{1}{\omega_a([E(R)]^{-1})}\right) + c\omega_a(R) \left(1 + \frac{1}{\omega_a(\mu_2)}\right). \end{aligned} \quad (7.71)$$

Again using (2.1) together with taking into account (7.25) and (7.5), we see

$$\frac{1}{\omega_a([E(R)]^{-1})} \leq \frac{1}{\omega_a\left(\frac{R}{2\|w_c\|_{L^\infty(B_R)}}\right)} \leq \frac{c(\mathbf{data})}{\omega_a(R)}. \quad (7.72)$$

Combining the last two displays, we find

$$I_a \leq c \left(\varepsilon + \omega_a(R) \left(1 + \frac{1}{\omega_a(\mu_2)}\right) \right) \quad (7.73)$$

with some constant $c \equiv c(\mathbf{data})$. Similarly, it holds that

$$I_b \leq c \left(\varepsilon + \omega_b(R) \left(1 + \frac{1}{\omega_b(\mu_2)}\right) \right). \quad (7.74)$$

Then, plugging the estimates in the last two displays into (7.60) and recalling (7.56), we have

$$\left| \int_{B_{1/8}} \langle \bar{A}_0(D\bar{w}_c), D\varphi \rangle dx \right| \leq c(\mathbf{data}) p_2(\varepsilon, R) \|D\varphi\|_{L^\infty(B_{1/8})}, \quad (7.75)$$

where

$$\begin{aligned} p_2(\varepsilon, R) &:= \left[\varepsilon + \omega_a(R) \left(1 + \frac{1}{\omega_a(\mu_2)}\right) \right]^{\frac{1}{s(H_a)+1}} + \left[\varepsilon + \omega_a(R) \left(1 + \frac{1}{\omega_a(\mu_2)}\right) \right]^{\frac{s(H_a)}{s(H_a)+1}} \\ &\quad + \left[\varepsilon + \omega_b(R) \left(1 + \frac{1}{\omega_b(\mu_2)}\right) \right]^{\frac{1}{s(H_b)+1}} + \left[\varepsilon + \omega_b(R) \left(1 + \frac{1}{\omega_b(\mu_2)}\right) \right]^{\frac{s(H_b)}{s(H_b)+1}}. \end{aligned} \quad (7.76)$$

Case 3: Assumption (1.22c) is in force. The assumption (1.22c)₂ implies that for any $\varepsilon \in (0, 1)$ there exists $\mu_3 > 0$ depending on ε such that

$$\Lambda\left(t^{\frac{1}{1-\gamma}}, \frac{1}{t}\right) \leq \varepsilon \quad \text{for every } t \in (0, \mu_3). \quad (7.77)$$

This one together with recalling (7.61) and (1.14), we see

$$\begin{aligned} I_a &\leq c\omega(R) \frac{H_a(E(R))}{G(E(R))} \\ &\leq c\omega_a(R)\varepsilon \left(1 + \frac{1}{\omega_a([E(R)]^{-\frac{1}{1-\gamma}})}\right) + c\omega_a(R) \left(1 + \frac{1}{\omega_a\left(\mu_3^{\frac{1}{1-\gamma}}\right)}\right). \end{aligned} \quad (7.78)$$

Now using (7.26), (7.6) and (1.14), we have

$$\frac{1}{\omega_a([E(R)]^{-\frac{1}{1-\gamma}})} \leq \frac{1}{\omega_a\left(\left[\frac{\text{osc}_{B_{2R}} u}{R}\right]^{-\frac{1}{1-\gamma}}\right)} \leq \frac{c(\mathbf{data})}{\omega_a(R)}. \quad (7.79)$$

Combining the last two displays, we find

$$I_a \leq c \left(\varepsilon + \omega_a(R) \left(1 + \frac{1}{\omega_a \left(\mu_3^{\frac{1}{1-\gamma}} \right)} \right) \right) \quad (7.80)$$

for some constant $c \equiv c(\mathbf{data})$. In the same way, we show

$$I_b \leq c \left(\varepsilon + \omega_b(R) \left(1 + \frac{1}{\omega_b \left(\mu_3^{\frac{1}{1-\gamma}} \right)} \right) \right) \quad (7.81)$$

for some constant $c \equiv c(\mathbf{data})$. Using the estimates (7.80)-(7.81) in (7.60), we conclude

$$\left| \int_{B_{1/8}} \langle \bar{A}_0(D\bar{w}_c), D\varphi \rangle dx \right| \leq c(\mathbf{data}) p_3(\varepsilon, R) \|D\varphi\|_{L^\infty(B_{1/8})}, \quad (7.82)$$

where

$$\begin{aligned} p_3(\varepsilon, R) := & \left[\varepsilon + \omega_a(R) \left(1 + \frac{1}{\omega_a \left(\mu_3^{\frac{1}{1-\gamma}} \right)} \right) \right]^{\frac{1}{s(H_a)+1}} + \left[\varepsilon + \omega_a(R) \left(1 + \frac{1}{\omega_a \left(\mu_3^{\frac{1}{1-\gamma}} \right)} \right) \right]^{\frac{s(H_a)}{s(H_a)+1}} \\ & + \left[\varepsilon + \omega_b(R) \left(1 + \frac{1}{\omega_b \left(\mu_3^{\frac{1}{1-\gamma}} \right)} \right) \right]^{\frac{1}{s(H_b)+1}} + \left[\varepsilon + \omega_b(R) \left(1 + \frac{1}{\omega_b \left(\mu_3^{\frac{1}{1-\gamma}} \right)} \right) \right]^{\frac{s(H_b)}{s(H_b)+1}}. \end{aligned} \quad (7.83)$$

Case 4. Assumption (1.22d) is in force. We treat this case in a different way rather than the estimate used in (7.63)-(7.69). In fact, we take an advantage that $w_a(\cdot)$ is a power function. Then recalling I_a introduced in (7.61), we see that

$$\begin{aligned} I_a & \leq cR^\alpha \frac{(H_a \circ G^{-1})(\Psi_{B_R}^-(E(R)))}{\Psi_{B_R}^-(E(R))} \leq cR^\alpha \left(1 + \left[\int_{B_{R/2}} \Psi_{B_R}^- \left(\left| \frac{w_c - (w_c)_{B_{R/2}}}{R} \right| \right) dx \right]^{\frac{\alpha}{n}} \right) \\ & \leq cR^\alpha + c \left(\int_{B_{R/2}} \Psi_{B_R}^- (|Dw_c|) dx \right)^{\frac{\alpha}{n}} \leq cR^\alpha + c \left(\int_{B_{2R}} \Psi(x, |Du|) dx \right)^{\frac{\alpha}{n}} \\ & \leq cR^\alpha + cR^{\frac{\alpha\delta}{1+\delta}} \left(\int_{B_{2R}} [\Psi(x, |Du|)]^{1+\delta} dx \right)^{\frac{\alpha}{n(1+\delta)}} \\ & \leq c(\mathbf{data}(\Omega_0)) R^{\frac{\alpha\delta}{1+\delta}} \end{aligned} \quad (7.84)$$

for a higher integrability exponent δ coming from Theorem 5.5, where we have used (7.24), (7.4) together with (5.62). By arguing similarly, we estimate I_b in (7.62) as

$$I_b \leq c(\mathbf{data}(\Omega_0)) R^{\frac{\beta\delta}{1+\delta}}. \quad (7.85)$$

Using estimates from the last two displays in (7.60) and recalling $R \leq 1$, we see

$$\left| \int_{B_{1/8}} \langle \bar{A}_0(D\bar{w}_c), D\varphi \rangle dx \right| \leq c(\mathbf{data}(\Omega_0)) q_1(R) \|D\varphi\|_{L^\infty(B_{1/8})}, \quad (7.86)$$

where

$$q_1(R) := R^{\frac{\alpha\delta}{(1+\delta)(1+s(H_a))}} + R^{\frac{\beta\delta}{(1+\delta)(1+s(H_b))}}. \quad (7.87)$$

Case 5: Assumption (1.22e) is in force. Again we estimate I_a and I_b introduced in (7.61)-(7.62). Using the assumption (1.13), (7.26) and (7.6), we have

$$\begin{aligned} I_a &\leq cR^\alpha \frac{H_a(E(R))}{G(E(R))} \\ &\leq cR^\alpha \left(1 + \left[(\Psi_{B_R}^-)^{-1} \left(\oint_{B_{R/2}} \Psi_{B_R}^- \left(\left| \frac{w_c - (w_c)_{B_{R/2}}}{R} \right| \right) dx \right) \right]^\alpha \right) \\ &\leq c \left(R^\alpha + \left[\text{osc}_{B_{2R}} u \right]^\alpha \right) \leq c(\mathbf{data}(\Omega_0)) R^{\gamma\alpha}, \end{aligned} \quad (7.88)$$

where we have also used (5.33) and the Hölder continuity exponent γ came from Theorem 5.2. Similarly, we see

$$I_b \leq c(\mathbf{data}(\Omega_0)) R^{\gamma\beta}. \quad (7.89)$$

Inserting the estimates from the last two displays into (7.60) and recalling $R \leq 1$, we see

$$\left| \oint_{B_{1/8}} \langle \bar{A}_0(D\bar{w}_c), D\varphi \rangle dx \right| \leq c(\mathbf{data}(\Omega_0)) q_2(R) \|D\varphi\|_{L^\infty(B_{1/8})}, \quad (7.90)$$

where

$$q_2(R) := R^{\frac{\alpha\gamma}{1+s(H_a)}} + R^{\frac{\beta\gamma}{1+s(H_b)}} \quad (7.91)$$

Collecting the estimates obtained in (7.68), (7.75), (7.82), (7.86) and (7.90), we conclude with

$$\left| \oint_{B_{1/8}} \langle \bar{A}_0(D\bar{w}_c), D\varphi \rangle dx \right| \leq c_h d(\varepsilon, R) \|D\varphi\|_{L^\infty(B_{1/8})} \quad (7.92)$$

for some constant $c_h \equiv c_h(\mathbf{data}(\Omega_0))$, whenever $\varphi \in W_0^{1,\infty}(B_{1/8})$, where

$$d(\varepsilon, R) := \begin{cases} p_1(\varepsilon, R) & \text{if (1.22a) is assumed,} \\ p_2(\varepsilon, R) & \text{if (1.22b) is assumed,} \\ p_3(\varepsilon, R) & \text{if (1.22c) is assumed,} \\ q_1(R) & \text{if (1.22d) is assumed,} \\ q_2(R) & \text{if (1.22e) is assumed,} \end{cases} \quad (7.93)$$

in which p_1, p_2, p_3, q_1 and q_2 have been defined in (7.69), (7.76), (7.83), (7.87) and (7.91), respectively. By (7.52), (7.54a)-(7.55) and (7.92), we are able to apply Lemma 6.1 with $A_0(z) \equiv \bar{A}_0(z)$, $\Psi_0(t) \equiv \bar{\Psi}_0(t)$ with $a_0 \equiv \bar{a}(\bar{x}_a)$ and $b_0 \equiv \bar{b}(\bar{x}_b)$, to discover that there exists $\bar{h} \in \bar{w}_c + W_0^{1,\bar{\Psi}_0}(B_{1/8})$ such that

$$\oint_{B_{1/8}} \langle \bar{A}_0(D\bar{h}), D\varphi \rangle dx = 0 \quad \text{for all} \quad \varphi \in W_0^{1,\infty}(B_{1/8}), \quad (7.94)$$

$$\oint_{B_{1/4}} \bar{\Psi}_0(|D\bar{h}|) dx + \oint_{B_{1/8}} [\bar{\Psi}_0(|D\bar{h}|)]^{1+\delta_1} dx \leq c \text{ for some } \delta_1 \leq \delta_0, \quad (7.95)$$

$$\begin{aligned} &\oint_{B_{1/8}} (|V_{\bar{G}}(D\bar{w}_c) - V_{\bar{G}}(D\bar{h})|^2 + \bar{a}(\bar{x}_a)|V_{\bar{H}_a}(D\bar{w}_c) - V_{\bar{H}_a}(D\bar{h})|^2 + \bar{b}(\bar{x}_b)|V_{\bar{H}_b}(D\bar{w}_c) - V_{\bar{H}_b}(D\bar{h})|^2) dx \\ &\leq c[d(\varepsilon, R)]^{s_1} \end{aligned} \quad (7.96)$$

and finally

$$\int_{B_{1/8}} (\bar{G}(|\bar{w}_c - \bar{h}|) + \bar{a}(\bar{x}_a)\bar{H}_a(|\bar{w}_c - \bar{h}|) + \bar{b}(\bar{x}_b)\bar{H}_b(|\bar{w}_c - \bar{h}|)) dx \leq c_d[d(\varepsilon, R)]^{s_0} \quad (7.97)$$

with some constants $c, c_d \equiv c, c_d(\mathbf{data}(\Omega_0)) \geq 1$ and $s_0, s_1 \equiv s_0, s_1(\mathbf{data}) \in (0, 1)$, but they are all independent of R . Therefore, for a given $\varepsilon^* \in (0, 1)$ as in the statement of our lemma, we choose small enough ε and R^* to satisfy

$$c_d[d(\varepsilon, R^*)]^{s_0} \leq \varepsilon^*. \quad (7.98)$$

Since the constants c_d and s_0 only depend on $\mathbf{data}(\Omega_0)$ and \mathbf{data} , respectively, the last display gives us the dependence of R^* as in (7.37). Furthermore, by (7.97), we conclude with

$$\int_{B_{1/8}} [\bar{G}(|\bar{w}_c - \bar{h}|) + \bar{a}(\bar{x}_a)\bar{H}_a(|\bar{w}_c - \bar{h}|) + \bar{b}(\bar{x}_b)\bar{H}_b(|\bar{w}_c - \bar{h}|)] dx \leq \varepsilon^*. \quad (7.99)$$

Proof of (7.38). We observe that by a standard density argument, the relation in (7.94) still holds for every $\varphi \in W_0^{1,1}(B_{1/8})$ with $\bar{\Psi}_0(|D\varphi|) \in L^1(B_{1/8})$. Recalling (7.49) and (7.50), we see that \bar{h} is a local minimizer of the functional

$$W^{1,\bar{\Psi}_0}(B_{1/8}) \ni v \mapsto \int_{B_{1/8}} \bar{F}_0(Dv) dx. \quad (7.100)$$

Since the conditions (7.54a)-(7.54c) are satisfied for the integrand $\bar{F}_0(\cdot)$, we are in a position to apply the results from [65] to obtain the following a priori Lipschitz estimate:

$$\sup_{B_{1/16}} \bar{\Psi}_0(|D\bar{h}|) \leq c \int_{B_{1/8}} \bar{\Psi}_0(|D\bar{h}|) dx \quad (7.101)$$

with some constant $c \equiv c(n, s(G), s(H_a), s(H_b), \nu, L)$. For any $\tau \in (0, 1/16)$, we have that

$$\begin{aligned} \int_{B_\tau} \bar{\Psi}_0\left(\left|\frac{\bar{w}_c - (\bar{w}_c)_{B_\tau}}{\tau}\right|\right) dx &\leq \int_{B_\tau} \bar{\Psi}_0\left(\left|\frac{\bar{w}_c - (\bar{h})_{B_\tau}}{\tau}\right|\right) dx \\ &\leq \int_{B_\tau} \bar{\Psi}_0\left(\left|\frac{\bar{h} - (\bar{h})_{B_\tau}}{\tau}\right|\right) dx + \int_{B_\tau} \bar{\Psi}_0\left(\left|\frac{\bar{w}_c - \bar{h}}{\tau}\right|\right) dx \\ &\stackrel{(7.99)}{\leq} c \sup_{B_\tau} \bar{\Psi}_0(|D\bar{h}|) + c\tau^{-(n+s(\Psi)+1)}\varepsilon^* \\ &\stackrel{(7.101)}{\leq} c \int_{B_{1/8}} \bar{\Psi}_0(|D\bar{h}|) dx + c\tau^{-(n+s(\Psi)+1)}\varepsilon^* \\ &\stackrel{(7.95)}{\leq} c + c\tau^{-(n+s(\Psi)+1)}\varepsilon^*. \end{aligned} \quad (7.102)$$

By scaling back to w_c as introduced in (7.42a)-(7.42b), we obtain the desired estimate (7.38). The proof is complete. \square

Lemma 7.4. *Under the assumptions and notations of Lemma 7.3, let $w_c \in W^{1,\Psi}(B_R)$ be the solution to the problem defined in (7.22). If one of the assumptions (1.22a)-(1.22e) is satisfied, then there exists $h \in w_c + W_0^{1,\Psi}(B_{R/8})$ being a local minimizer of the functional defined by*

$$W^{1,1}(B_{R/8}) \ni v \mapsto \mathcal{F}_0(v) := \int_{B_{R/8}} F_0(Dv) dx, \quad (7.103)$$

where the integrand function is given by

$$F_0(z) := F_G(x_c, (u)_{B_{2R}}, z) + a(x_a)F_{H_a}(x_c, (u)_{B_{2R}}, z) + b(x_b)F_{H_b}(x_c, (u)_{B_{2R}}, z) \quad (7.104)$$

for some fixed point $x_c \in B_R$ having been fixed in (7.22) and $x_a, x_b \in \overline{B_R}$ being points such that $a(x_a) := \inf_{x \in B_R} a(x)$ and $b(x_b) := \inf_{x \in B_R} b(x)$, whenever $z \in \mathbb{R}^n$, such that

$$\begin{aligned} & \int_{B_{R/8}} [|V_G(Du) - V_G(Dh)|^2 + a(x_a)|V_{H_a}(Du) - V_{H_a}(Dh)|^2 + b(x_b)|V_{H_b}(Du) - V_{H_b}(Dh)|^2] dx \\ & \leq c(\omega(R^\gamma) + [d(\varepsilon, R)]^{s_1}) \int_{B_{2R}} \Psi(x, |Du|) dx \end{aligned} \quad (7.105)$$

for some constant $c \equiv c(\mathbf{data}(\Omega_0))$, where s_1 and $d(\varepsilon, R)$ have been defined in (7.96) and (7.93), respectively. Moreover, we have the energy estimate

$$\int_{B_{R/8}} \Psi_{B_R}^-(|Dh|) dx \leq c \int_{B_{2R}} \Psi(x, |Du|) dx \quad (7.106)$$

for some constant $c \equiv c(n, \nu, L)$.

Proof. We need to revisit the proof of Lemma 7.3, specially Step 3 and Step 4. Under the settings of the proof of Lemma 7.3, we consider a function $\bar{h} \in \bar{w}_c + W_0^{1, \Psi_0}(B_{1/8})$ satisfying (7.94)-(7.97). Let h be the scaled back function of \bar{h} in $B_{R/8}$ as

$$h(x) := E(w_c, B_{R/2}) R \bar{h} \left(\frac{x - x_0}{R} \right) \quad \text{for every } x \in B_{R/8}(x_0). \quad (7.107)$$

Clearly, $h \in w_c + W_0^{1, \Psi_{B_R}^-}(B_{R/8})$ is a local minimizer of the functional \mathcal{F}_0 defined in (7.103) which means that

$$\mathcal{F}_0(h) = \int_{B_{R/8}} F_0(Dh) dx \leq \int_{B_{R/8}} F_0(Dh + D\varphi) dx = \mathcal{F}_0(h + \varphi) \quad (7.108)$$

holds for every $\varphi \in W_0^{1, \Psi_{B_R}^-}(B_{R/8})$. As we have shown in (7.9), recalling (7.24) and (7.4), we see

$$\begin{aligned} \int_{B_{R/8}} \Psi_{B_R}^- (|Dh|) dx & \leq \frac{L}{\nu} \int_{B_{R/8}} \Psi_{B_R}^- (|Dw_c|) dx \leq \frac{8^n L}{\nu} \int_{B_R} \Psi(x, |Dw_c|) dx \\ & \leq c(n, \nu, L) \int_{B_R} \Psi(x, |Dw|) dx \leq c(n, \nu, L) \int_{B_{2R}} \Psi(x, |Du|) dx, \end{aligned} \quad (7.109)$$

which proves (7.106). We write the inequality (7.96) in view of G, H_a, H_b, w_c and h in order to have

$$\begin{aligned} & \int_{B_{R/8}} [|V_G(Dw_c) - V_G(Dh)|^2 + a(x_a)|V_{H_a}(Dw_c) - V_{H_a}(Dh)|^2 + b(x_b)|V_{H_b}(Dw_c) - V_{H_b}(Dh)|^2] dx \\ & \leq c[d(\varepsilon, R)]^{s_1} \int_{B_{R/2}} \Psi_{B_R}^- \left(\left| \frac{w_c - (w_c)_{B_{R/2}}}{R} \right| \right) dx \\ & \leq c[d(\varepsilon, R)]^{s_1} \int_{B_{R/2}} \Psi_{B_R}^- (|Dw_c|) dx \leq c[d(\varepsilon, R)]^{s_1} \int_{B_{R/2}} \Psi(x, |Du|) dx \end{aligned} \quad (7.110)$$

for some constant $c \equiv c(\mathbf{data}(\Omega_0))$, where we have applied Sobolev-Poincaré inequality and (7.109). Combining this estimate together with (7.3) and (7.23) via some elementary computations and recalling $R \leq 1$, we directly arrive at (7.105). \square

We finally finish this section with a crucial decay estimate on a local minimizer u of the functional \mathcal{F} .

Lemma 7.5. *Under the assumptions and notations of Lemma 7.3, if one of the conditions (1.22a)-(1.22e) is satisfied, then for every $\varepsilon_* \in (0, 1)$, there exists a positive radius R_* with the dependence as*

$$R_* \equiv R_*(\mathbf{data}(\Omega_0), \varepsilon_*) \quad (7.111)$$

such that if $R \leq R_*$, then there exists a constant $c \equiv c(\mathbf{data}(\Omega_0))$ such that

$$\int_{B_{\tau R}} \Psi_{B_R}^- \left(\left| \frac{u - (u)_{B_{\tau R}}}{\tau R} \right| \right) dx \leq c \left(\tau^n + \tau^{-(s(\Psi)+1)} \varepsilon_* \right) \int_{B_{2R}} \Psi(x, |Du|) dx \quad (7.112)$$

holds for every $\tau \in (0, 1/16)$.

Proof. First we apply Lemma 7.3 with $\varepsilon^* \in (0, 1)$ to be determined in a few lines, and we can use (7.38) provided

$$R \leq R^* \equiv R^*(\mathbf{data}(\Omega_0), \varepsilon^*)$$

is found via (7.37). Therefore, using the convexity of $\Psi_{B_R}^-$, Lemma 7.3 and a Sobolev-Poincaré inequality of Lemma 4.1 via some elementary manipulations, for every $\tau \in (0, 1/32)$, we have that

$$\begin{aligned} \int_{B_{\tau R}} \Psi_{B_R}^- \left(\left| \frac{u - (u)_{B_{\tau R}}}{\tau R} \right| \right) dx &\leq c \int_{B_{\tau R}} \Psi_{B_R}^- \left(\left| \frac{u - (w_c)_{B_{\tau R}}}{\tau R} \right| \right) dx \\ &\leq c \int_{B_{\tau R}} \Psi_{B_R}^- \left(\left| \frac{w_c - (w_c)_{B_{\tau R}}}{\tau R} \right| \right) dx + c\tau^{-(n+s(\Psi)+1)} \int_{B_R} \Psi_{B_R}^- \left(\left| \frac{u - w_c}{R} \right| \right) dx \\ &\leq c \left(1 + \tau^{-(n+s(\Psi)+1)} \varepsilon^* \right) \int_{B_{R/2}} \Psi_{B_R}^- \left(\left| \frac{w_c - (w_c)_{B_{R/2}}}{R} \right| \right) dx \\ &\quad + c\tau^{-(n+s(\Psi)+1)} \int_{B_R} \Psi_{B_R}^- \left(\left| \frac{u - w_c}{R} \right| \right) dx \\ &\leq c \left(1 + \tau^{-(n+s(\Psi)+1)} \varepsilon^* \right) \int_{B_R} \Psi_{B_R}^- (|Dw_c|) dx + c\tau^{-(n+s(\Psi)+1)} \int_{B_R} \Psi_{B_R}^- \left(\left| \frac{u - w_c}{R} \right| \right) dx \end{aligned} \quad (7.113)$$

with some constant $c \equiv c(\mathbf{data}(\Omega_0))$, where throughout the last display we have repeatedly used (2.6) and (7.36). The last display, (7.7) and (7.27) with some elementary manipulations yield

$$\int_{B_{\tau R}} \Psi_{B_R}^- \left(\left| \frac{u - (u)_{B_{\tau R}}}{\tau R} \right| \right) dx \leq c \left(\tau^n + \tau^{-(s(\Psi)+1)} \varepsilon^* + \tau^{-(s(\Psi)+1)} [\omega(R^\gamma)]^{\frac{1}{2}} \right) \int_{B_{2R}} \Psi(x, |Du|) dx$$

for every $\tau \in (0, 1/16)$ and some $c \equiv c(\mathbf{data}(\Omega_0))$. Then we choose $\varepsilon_* \equiv \varepsilon^*/2$ and $R_* \leq R^*$ in such a way that $[\omega(R_*^\gamma)]^{\frac{1}{2}} \leq \varepsilon_*/2$. This choice gives us the dependence as described in (7.111) and yields (7.112). \square

8. PROOF OF THEOREM 1.2

Now we are ready to provide the proof of Theorem 1.2. In fact, it comes from the combination of Lemma 5.3 and Lemma 7.5.

Step 1: Different alternatives. Now we consider the different alternatives depending on which phase of (5.27a)-(5.27d) occurs in some fixed ball $B_R \equiv B_R(x_0) \subset \Omega_0 \Subset \Omega$ with $R \leq R_* \equiv R_*(\mathbf{data}(\Omega_0), \varepsilon_*)$, which will be determined via Lemma 7.5 depending on $\varepsilon_* \in (0, 1)$.

Alternative 1. Let $\tau_{ab} \in (0, 1/64)$ to be chosen in a few lines. Assume that G -phase occurs in the ball $B_{\tau_{ab}R}$, which means that (5.27a) happens in $B_{\tau_{ab}R}$. In this situation, we have

$$a^-(B_{2\tau_{ab}R}) \leq 8[a]_{\omega_a} \omega_a(\tau_{ab}R) \quad \text{and} \quad b^-(B_{2\tau_{ab}R}) \leq 8[b]_{\omega_b} \omega_b(\tau_{ab}R). \quad (8.1)$$

Then we are able to apply Lemma 5.3 in the ball $B_{2\tau_{ab}R}$. In turn, this one together with applying Lemma 7.5 implies that

$$\begin{aligned} \int_{B_{\tau_{ab}R}} \Psi(x, |Du|) dx &\leq c \int_{B_{2\tau_{ab}R}} G\left(\left|\frac{u - (u)_{B_{2\tau_{ab}R}}}{2\tau_{ab}R}\right|\right) dx \\ &\leq c \int_{B_{2\tau_{ab}R}} \Psi_{B_R}^-\left(\left|\frac{u - (u)_{B_{2\tau_{ab}R}}}{2\tau_{ab}R}\right|\right) dx \\ &\leq c \left(\tau_{ab}^n + \tau_{ab}^{-(s(\Psi)+1)\varepsilon_*}\right) \int_{B_R} \Psi(x, |Du|) dx \end{aligned} \quad (8.2)$$

for $c \equiv c(\mathbf{data}(\Omega_0))$, provided $R \leq R_*(\mathbf{data}(\Omega_0), \varepsilon_*)$. Then, for every $\sigma \in (0, n)$, we write down the last inequality in the following form

$$\int_{B_{\tau_{ab}R}} \Psi(x, |Du|) dx \leq \tau_{ab}^{n-\sigma} \left(c_{ab} \tau_{ab}^\sigma + c_{ab} \tau_{ab}^{\sigma-(n+s(\Psi)+1)\varepsilon_*} \right) \int_{B_R} \Psi(x, |Du|) dx$$

for some constant $c_{ab} \equiv c_{ab}(\mathbf{data}(\Omega_0))$. We select small enough τ_{ab} , ε_* depending on $\mathbf{data}(\Omega_0)$ and σ in such a way that $c_{ab} \tau_{ab}^\sigma \leq 1/2$ and $c_{ab} \tau_{ab}^{\sigma-(n+s(\Psi)+1)\varepsilon_*} \leq 1/2$. Then we have

$$\int_{B_{\tau_{ab}R}} \Psi(x, |Du|) dx \leq \tau_{ab}^{n-\sigma} \int_{B_R} \Psi(x, |Du|) dx \quad (8.3)$$

for every $R \leq R_{ab} \equiv R_{ab}(\mathbf{data}(\Omega_0), \sigma)$.

Alternative 2. Let $\tau_b \in (0, 1/64)$ also to be determined later. This time we assume that (G, H_a) -phase occurs in B_R ((5.27b) happens in B_R) and that $b^-(B_{\tau_b R}) \leq 4[b]_{\omega_b} \omega_b(\tau_b R)$. Then we have

$$b^-(B_{2\tau_b R}) \leq 8[b]_{\omega_b} \omega_b(\tau_b R). \quad (8.4)$$

Also we can observe that

$$a^-(B_{\tau_b R}) \geq a^-(B_R) > 4[a]_{\omega_a} \omega_a(R) \geq 4[a]_{\omega_a} \omega_a(\tau_b R) \quad (8.5)$$

and

$$a^-(B_R) \leq a(x) \leq 2[a]_{\omega_a} \omega_a(R) + a^-(B_R) \leq 2a^-(B_R) \quad (\forall x \in B_R). \quad (8.6)$$

Applying Lemma 5.3 and then Lemma 7.5 together with recalling (8.6), we have

$$\begin{aligned} \int_{B_{\tau_b R}} \Psi(x, |Du|) dx &\leq c \int_{B_{2\tau_b R}} \left[G\left(\left|\frac{u - (u)_{B_{2\tau_b R}}}{2\tau_b R}\right|\right) + a^-(B_{2\tau_b R}) H_a\left(\left|\frac{u - (u)_{B_{2\tau_b R}}}{2\tau_b R}\right|\right) \right] dx \\ &\leq c \int_{B_{2\tau_b R}} \Psi_{B_R}^-\left(\left|\frac{u - (u)_{B_{2\tau_b R}}}{2\tau_b R}\right|\right) dx \\ &\leq c \left(\tau_b^n + \tau_b^{-(s(\Psi)+1)\varepsilon_*} \right) \int_{B_R} \Psi(x, |Du|) dx \end{aligned} \quad (8.7)$$

for some constant $c \equiv c(\mathbf{data}(\Omega_0))$, provided $R \leq R_*(\mathbf{data}(\Omega_0), \varepsilon_*)$. Then, for every $\sigma \in (0, n)$, we write down the last display as

$$\int_{B_{\tau_b R}} \Psi(x, |Du|) dx \leq \tau_b^{n-\sigma} \left(c_b \tau_b^\sigma + c_b \tau_b^{\sigma-(n+s(\Psi)+1)\varepsilon_*} \right) \int_{B_R} \Psi(x, |Du|) dx$$

for some constant $c_b \equiv c_b(\mathbf{data}(\Omega_0))$. We select small enough τ_b , ε_* depending on $\mathbf{data}(\Omega_0)$ and σ in such a way that $c_b \tau_b^\sigma \leq 1/2$ and $c_b \tau_b^{\sigma-(n+s(\Psi)+1)\varepsilon_*} \leq 1/2$. Then we have

$$\int_{B_{\tau_b R}} \Psi(x, |Du|) dx \leq \tau_b^{n-\sigma} \int_{B_R} \Psi(x, |Du|) dx \quad (8.8)$$

for every $R \leq R_b \equiv R_b(\mathbf{data}(\Omega_0), \sigma)$.

Alternative 3. Let $\tau_a \in (0, 1/64)$ to be fixed later. Assume that (G, H_b) -phase occurs in B_R ((5.27c) happens in B_R) and $a^-(B_{\tau_a R}) \leq 4[a]_{\omega_a} \omega_a(\tau_a R)$. Then we have

$$a^-(B_{2\tau_a R}) \leq 8[a]_{\omega_a} \omega_a(\tau_a R). \quad (8.9)$$

Applying Lemma 5.3 and then Lemma 7.5 together with recalling that $b^-(B_R) \leq b(x) \leq 2b^-(B_R)$ holds for every $x \in B_R$ if $b^-(B_R) > 4[b]_{\omega_b} \omega_b(R)$ likewise in (8.6), we have

$$\begin{aligned} \int_{B_{\tau_a R}} \Psi(x, |Du|) dx &\leq c \int_{B_{2\tau_a R}} \left[G\left(\left|\frac{u - (u)_{B_{2\tau_a R}}}{2\tau_a R}\right|\right) + b^-(B_{2\tau_a R}) H_b\left(\left|\frac{u - (u)_{B_{2\tau_a R}}}{2\tau_a R}\right|\right) \right] dx \\ &\leq c \int_{B_{2\tau_a R}} \Psi_{B_R}^-\left(\left|\frac{u - (u)_{B_{2\tau_a R}}}{2\tau_a R}\right|\right) dx \\ &\leq c \left(\tau_a^n + \tau_a^{-(s(\Psi)+1)} \varepsilon_* \right) \int_{B_R} \Psi(x, |Du|) dx \end{aligned} \quad (8.10)$$

for some constant $c \equiv c(\mathbf{data}(\Omega_0))$, provided $R \leq R_*(\mathbf{data}(\Omega_0), \varepsilon_*)$. Then, for every $\sigma \in (0, n)$, we write down the last display as

$$\int_{B_{\tau_a R}} \Psi(x, |Du|) dx \leq \tau_a^{n-\sigma} \left(c_a \tau_a^\sigma + c_a \tau_a^{\sigma - (n+s(\Psi)+1)} \varepsilon_* \right) \int_{B_R} \Psi(x, |Du|) dx$$

for some constant $c_a \equiv c_a(\mathbf{data}(\Omega_0))$. We select small enough τ_a, ε_* depending on $\mathbf{data}(\Omega_0)$ and σ in such a way that $c_a \tau_a^\sigma \leq 1/2$ and $c_a \tau_a^{\sigma - (n+s(\Psi)+1)} \varepsilon_* \leq 1/2$. Then we have

$$\int_{B_{\tau_a R}} \Psi(x, |Du|) dx \leq \tau_a^{n-\sigma} \int_{B_R} \Psi(x, |Du|) dx \quad (8.11)$$

for every $R \leq R_a \equiv R_a(\mathbf{data}(\Omega_0), \sigma)$.

Alternative 4. Let $\tau_0 \in (0, 1/64)$ to be chosen later. We assume that (G, H_a, H_b) -phase occurs in B_R , which means that (5.27d) happens in B_R . In this situation, from the observation in (8.6) we see that $a^-(B_R) \leq a(x) \leq 2a^-(B_R)$ and $b^-(B_R) \leq b(x) \leq 2b^-(B_R)$ for every $x \in B_R$. Then again applying Lemma 5.3 and Lemma 7.5, we find

$$\begin{aligned} \int_{B_{\tau_0 R}} \Psi(x, |Du|) dx &\leq c \int_{B_{2\tau_0 R}} \Psi_{B_{2\tau_0 R}}^-\left(\left|\frac{u - (u)_{B_{2\tau_0 R}}}{2\tau_0 R}\right|\right) dx \\ &\leq c \int_{B_{2\tau_0 R}} \Psi_{B_R}^-\left(\left|\frac{u - (u)_{B_{2\tau_0 R}}}{2\tau_0 R}\right|\right) dx \\ &\leq c \left(\tau_0^n + \tau_0^{-(s(\Psi)+1)} \varepsilon_* \right) \int_{B_R} \Psi(x, |Du|) dx \end{aligned} \quad (8.12)$$

for some constant $c \equiv c(\mathbf{data}(\Omega_0))$, provided $R \leq R_*(\mathbf{data}(\Omega_0), \varepsilon_*)$. Then, for every $\sigma \in (0, n)$, we write down the last display as

$$\int_{B_{\tau_0 R}} \Psi(x, |Du|) dx \leq \tau_0^{n-\sigma} \left(c_0 \tau_0^\sigma + c_0 \tau_0^{\sigma - (n+s(\Psi)+1)} \varepsilon_* \right) \int_{B_R} \Psi(x, |Du|) dx$$

for some constant $c_0 \equiv c_0(\mathbf{data}(\Omega_0))$. Then we choose τ_0, ε_* depending on $\mathbf{data}(\Omega_0)$ and σ in such a way that $c_0 \tau_0^\sigma \leq 1/2$ and $c_0 \tau_0^{\sigma - (n+s(\Psi)+1)} \varepsilon_* \leq 1/2$. Then we have

$$\int_{B_{\tau_0 R}} \Psi(x, |Du|) dx \leq \tau_0^{n-\sigma} \int_{B_R} \Psi(x, |Du|) dx \quad (8.13)$$

for every $R \leq R_0 \equiv R_0(\mathbf{data}(\Omega_0), \sigma)$. Next we consider the double nested exit time argument based on the proof of [38, Theorem 2].

Step 2: Double nested exit time and iteration. Now we shall combine all the alternatives we have discussed with the estimates (8.3), (8.8), (8.11) and (8.13). Take a ball $B_R \subset \Omega_0 \Subset \Omega$ such that $R \leq R_m$, where $R_m = \min\{R_{ab}, R_a, R_b, R_0\}$ depends on $\mathbf{data}(\Omega_0)$ and σ . We consider G -phase in $B_{\tau_{ab}^{k+1}R}$ for every integer $k \geq 0$ and define the exit time index

$$t_{ab} = \min\{k \in \mathbb{N} : G\text{-phase in the ball } B_{\tau_{ab}^{k+1}R} \text{ does not occur}\}. \quad (8.14)$$

If there does not exist such t_{ab} , then for any $0 < \rho < \tau_{ab}^2 R < R \leq R_m$, there exists an integer $m \geq 1$ such that $\tau_{ab}^{m+2} R \leq \rho < \tau_{ab}^{m+1} R$. Using iterative (8.3), we have

$$\begin{aligned} \int_{B_\rho} \Psi(x, |Du|) dx &\leq \int_{B_{\tau_{ab}^{m+1}R}} \Psi(x, |Du|) dx \\ &\leq \tau_{ab}^{(m-1)(n-\sigma)} \int_{\tau_{ab}^2 R} \Psi(x, |Du|) dx = \tau_{ab}^{(m+2)(n-\sigma)} \tau_{ab}^{-3(n-\sigma)} \int_{\tau_{ab}^2 R} \Psi(x, |Du|) dx \\ &\leq c(\mathbf{data}(\Omega_0), \sigma) \left(\frac{\rho}{R}\right)^{n-\sigma} \int_R \Psi(x, |Du|) dx. \end{aligned} \quad (8.15)$$

Clearly, the above inequality holds true when $\tau_{ab}^2 R \leq \rho \leq R \leq R_m$. So we consider the case of $t_{ab} < \infty$. For every $k \in \{1, \dots, t_{ab}\}$, we apply (8.3) repeatedly in order to obtain

$$\int_{B_{\tau_{ab}^k R}} \Psi(x, |Du|) dx \leq \tau_{ab}^{k(n-\sigma)} \int_{B_R} \Psi(x, |Du|) dx. \quad (8.16)$$

By the very definition of τ_{ab} in (8.14), we have three different scenarios: either (G, H_a) -phase occurs in $B_{\tau_{ab}^{t_{ab}+1}R}$, (G, H_b) -phase occurs in $B_{\tau_{ab}^{t_{ab}+1}R}$ or (G, H_a, H_b) -phase occurs in $B_{\tau_{ab}^{t_{ab}+1}R}$. Clearly, the last condition is stable for shrinking balls. Since the first two conditions can be considered similarly, we shall focus on the occurrence of (G, H_a) -phase in the ball $B_{\tau_{ab}^{t_{ab}+1}R}$. Let us define a second exit time index

$$t_b := \min\{k \in \mathbb{N} : (G, H_a) - \text{phase in the ball } B_{\tau_b^{k+1}\tau_{ab}^{t_{ab}+1}R} \text{ does not occur}\}. \quad (8.17)$$

Arguing similarly as in (8.15) by using (8.8) if there is no such a finite number $t_b \in \mathbb{N}$, we are able to arrive at the inequality (8.25) below. So we only focus on the case of $t_b < \infty$. Iterating (8.8) with B_R replaced by $B_{\tau_{ab}^{t_{ab}+1}R}$, we have

$$\int_{B_{\tau_b^k \tau_{ab}^{t_{ab}+1}R}} \Psi(x, |Du|) dx \leq \tau_b^{k(n-\sigma)} \int_{B_{\tau_{ab}^{t_{ab}+1}R}} \Psi(x, |Du|) dx \quad (8.18)$$

for every $k \in \{1, \dots, t_b\}$. By again the very definition of t_b , there is only one chance that (G, H_a, H_b) -phase occurs in the ball $B_{\tau_b^{t_b+1}\tau_{ab}^{t_{ab}+1}R}$. But as this condition is stable, we can iterate (8.13) for every $k \in \mathbb{N}$ in order to have

$$\int_{B_{\tau_0^k \tau_b^{t_b+1} \tau_{ab}^{t_{ab}+1}R}} \Psi(x, |Du|) dx \leq \tau_0^{k(n-\sigma)} \int_{B_{\tau_b^{t_b+1} \tau_{ab}^{t_{ab}+1}R}} \Psi(x, |Du|) dx. \quad (8.19)$$

Now we have all the needed estimates (8.16), (8.18) and (8.19). For $0 < \rho < R \leq R_m$, we consider the following cases.

Case 1: $R > \rho \geq \tau_{ab}^{t_{ab}+1} R$. There exists $m \in \{0, 1, \dots, t_{ab}\}$ such that $\tau_{ab}^{m+1} R \leq \rho < \tau_{ab}^m R$. Then from (8.16), we have

$$\begin{aligned}
\int_{B_\rho} \Psi(x, |Du|) dx &\leq \int_{B_{\tau_{ab}^m R}} \Psi(x, |Du|) dx \\
&\leq \tau_{ab}^{m(n-\sigma)} \int_{B_R} \Psi(x, |Du|) dx \\
&\leq \tau_{ab}^{(m+1)(n-\sigma)} \tau_{ab}^{\sigma-n} \int_{B_R} \Psi(x, |Du|) dx \\
&\leq c(\mathbf{data}(\Omega_0), \sigma) \left(\frac{\rho}{R}\right)^{n-\sigma} \int_{B_R} \Psi(x, |Du|) dx,
\end{aligned} \tag{8.20}$$

where the last inequality is valid since τ_{ab} depends on $\mathbf{data}(\Omega_0)$ and σ .

Case 2: $\tau_{ab}^{t_{ab}+1} R > \rho \geq \tau_b \tau_{ab}^{t_{ab}+1} R$. In this case, using (8.20), we see

$$\begin{aligned}
\int_{B_\rho} \Psi(x, |Du|) dx &\leq \int_{B_{\tau_{ab}^{t_{ab}+1} R}} \Psi(x, |Du|) dx \\
&= \tau_{ab}^{(t_{ab}+1)(n-\sigma)} \int_{B_R} \Psi(x, |Du|) dx \\
&\leq (\tau_b \tau_{ab}^{t_{ab}+1})^{n-\sigma} \tau_b^{\sigma-n} \int_{B_R} \Psi(x, |Du|) dx \\
&\leq c(\mathbf{data}(\Omega_0), \sigma) \left(\frac{\rho}{R}\right)^{n-\sigma} \int_{B_R} \Psi(x, |Du|) dx,
\end{aligned} \tag{8.21}$$

where again the last inequality is possible by the dependencies of τ_b .

Case 3: $\tau_b \tau_{ab}^{t_{ab}+1} R > \rho \geq \tau_b^{t_b+1} \tau_{ab}^{t_{ab}+1} R$. Again there exists a natural number $m \in \{1, \dots, t_b\}$ so that $\tau_b^m \tau_{ab}^{t_{ab}+1} R > \rho \geq \tau_b^{m+1} \tau_{ab}^{t_{ab}+1} R$. Therefore, using (8.18) and (8.20), we have

$$\begin{aligned}
\int_{B_\rho} \Psi(x, |Du|) dx &\leq \int_{B_{\tau_b^m \tau_{ab}^{t_{ab}+1} R}} \Psi(x, |Du|) dx \\
&\leq \tau_b^{m(n-\sigma)} \int_{B_{\tau_{ab}^{t_{ab}+1} R}} \Psi(x, |Du|) dx \\
&\leq \tau_b^{(m+1)(n-\sigma)} \tau_b^{\sigma-n} \tau_{ab}^{(t_{ab}+1)(n-\sigma)} \int_{B_R} \Psi(x, |Du|) dx \\
&\leq c(\mathbf{data}(\Omega_0), \sigma) \left(\frac{\rho}{R}\right)^{n-\sigma} \int_{B_R} \Psi(x, |Du|) dx.
\end{aligned} \tag{8.22}$$

Case 4: $\tau_b^{t_b+1}\tau_{ab}^{t_{ab}+1}R > \rho \geq \tau_b^{t_b+1}\tau_{ab}^{t_{ab}+1}\tau_0 R$. Now by (8.22), we find

$$\begin{aligned}
\int_{B_\rho} \Psi(x, |Du|) dx &\leq \int_{B_{\tau_b^{t_b+1}\tau_{ab}^{t_{ab}+1}R}} \Psi(x, |Du|) dx \\
&\leq c \left(\tau_b^{t_b+1}\tau_{ab}^{t_{ab}+1} \right)^{n-\sigma} \int_{B_R} \Psi(x, |Du|) dx \\
&\leq c \tau_0^{\sigma-n} \left(\tau_0 \tau_b^{t_b+1}\tau_{ab}^{t_{ab}+1} \right)^{n-\sigma} \int_{B_R} \Psi(x, |Du|) dx \\
&\leq c(\mathbf{data}(\Omega_0), \sigma) \left(\frac{\rho}{R} \right)^{n-\sigma} \int_{B_R} \Psi(x, |Du|) dx.
\end{aligned} \tag{8.23}$$

Case 5: $\tau_b^{t_b+1}\tau_{ab}^{t_{ab}+1}\tau_0 R > \rho > 0$. This condition implies that there exists a natural number $m \in \mathbb{N}$ such that $\tau_0^{m+1}\tau_b^{t_b+1}\tau_{ab}^{t_{ab}+1}R \leq \rho < \tau_0^m\tau_b^{t_b+1}\tau_{ab}^{t_{ab}+1}R$. This time we apply (8.19) and (8.23) in order to have

$$\begin{aligned}
\int_{B_\rho} \Psi(x, |Du|) dx &\leq \int_{B_{\tau_0^m\tau_b^{t_b+1}\tau_{ab}^{t_{ab}+1}R}} \Psi(x, |Du|) dx \\
&\leq \tau_0^{m(n-\sigma)} \int_{B_{\tau_b^{t_b+1}\tau_{ab}^{t_{ab}+1}R}} \Psi(x, |Du|) dx \\
&\leq \tau_0^{m(n-\sigma)} \left(\tau_b^{t_b+1}\tau_{ab}^{t_{ab}+1} \right)^{n-\sigma} \int_{B_R} \Psi(x, |Du|) dx \\
&\leq c \tau_0^{\sigma-n} \left(\frac{\rho}{R} \right)^{n-\sigma} \int_{B_R} \Psi(x, |Du|) dx = c(\mathbf{data}(\Omega_0), \sigma) \left(\frac{\rho}{R} \right)^{n-\sigma} \int_{B_R} \Psi(x, |Du|) dx.
\end{aligned} \tag{8.24}$$

As we discussed earlier after (8.16), we can proceed the same for the occurrence of (G, H_a) -phase in the ball $B_{\tau_{ab}^{\tau_{ab}+1}R}$ instead of the occurrence of (G, H_b) -phase in the ball $B_{\tau_{ab}^{\tau_{ab}+1}R}$. Then we can directly jump to the case that (G, H_a, H_b) -phase occurs in the ball $B_{\tau_{ab}^{\tau_{ab}+1}R}$, which is trivial by (8.13). Moreover, if we start with the occurrence of (G, H_a, H_b) -phase in B_R , then the procedure will be much easier by (8.13). Taking into account all the possible cases that we considered above, we can conclude that, for every $\sigma \in (0, 1)$, there exists $c \equiv c(\mathbf{data}(\Omega_0), \sigma)$ such that

$$\int_{B_\rho} \Psi(x, |Du|) dx \leq c \left(\frac{\rho}{R} \right)^{n-\sigma} \int_{B_R} \Psi(x, |Du|) dx \tag{8.25}$$

holds true, whenever $0 < \rho < R \leq R_m$, where R_m is some positive radius depending only on $\mathbf{data}(\Omega_0)$ and σ in the beginning of the proof. In order to complete the proof, we need to consider the remaining cases. If $0 < R_m \leq \rho < R \leq 1$, then we have

$$\begin{aligned}
\int_{B_\rho} \Psi(x, |Du|) dx &\leq \left(\frac{\rho}{R} \right)^{n-\sigma} \left(\frac{R}{\rho} \right)^{n-\sigma} \int_{B_R} \Psi(x, |Du|) dx \\
&\leq \left(\frac{\rho}{R} \right)^{n-\sigma} \left(\frac{R}{R_m} \right)^{n-\sigma} \int_{B_R} \Psi(x, |Du|) dx \\
&\leq c(\mathbf{data}(\Omega_0), \sigma) \left(\frac{\rho}{R} \right)^{n-\sigma} \int_{B_R} \Psi(x, |Du|) dx,
\end{aligned} \tag{8.26}$$

where we have used the dependence of R_m . Finally, if $0 < \rho < R_m \leq R \leq 1$, then by (8.25) and (8.26), we see

$$\begin{aligned} \int_{B_\rho} \Psi(x, |Du|) dx &\leq c \left(\frac{\rho}{R_m} \right)^{n-\sigma} \int_{B_{R_m}} \Psi(x, |Du|) dx \\ &\leq c \left(\frac{\rho}{R_m} \right)^{n-\sigma} \left(\frac{R_m}{R} \right)^{n-\sigma} \int_{B_R} \Psi(x, |Du|) dx. \\ &= c(\mathbf{data}(\Omega_0), \sigma) \left(\frac{\rho}{R} \right)^{n-\sigma} \int_{B_R} \Psi(x, |Du|) dx, \end{aligned} \quad (8.27)$$

All in all, collecting the estimates obtained in (8.25)-(8.27), we arrive at the validity of the Morrey type inequality (1.24). The proof is complete.

Now we consider a crucial outcome of Theorem 1.2, which plays a crucial role for proving Theorem 1.1 afterwards.

Lemma 8.1. *Under the assumptions of Lemma 7.3, let $w_c \in W^{1,\Psi}(B_R)$ be the solution to the problem defined in (7.22). Suppose that (1.22c) is satisfied for $\omega_a(t) = t^\alpha$ and $\omega_b(t) = t^\beta$ with some $\alpha, \beta \in (0, 1]$. Then there exists $h \in w_c + W_0^{1,\Psi_{B_R}}(B_{R/8})$ being a local minimizer of the functional \mathcal{F}_0 defined in (7.103) such that*

$$\begin{aligned} &\int_{B_{R/8}} [|V_G(Du) - V_G(Dh)|^2 + a(x_a)|V_{H_a}(Du) - V_{H_a}(Dh)|^2 + b(x_b)|V_{H_b}(Du) - V_{H_b}(Dh)|^2] dx \\ &\leq c \left(\omega(R^\gamma) + \left[R^{\frac{\alpha}{2(1+s(H_a))}} + R^{\frac{\beta}{2(1+s(H_b))}} \right]^{s_1} \right) \int_{B_{2R}} \Psi(x, |Du|) dx \end{aligned} \quad (8.28)$$

for some constant $c \equiv c(\mathbf{data}(\Omega_0))$ and $s_1 \equiv s_1(\mathbf{data})$, respectively. Moreover, the energy estimate

$$\int_{B_{R/8}} \Psi_{B_R}^- (|Dh|) dx \leq c \int_{B_{2R}} \Psi(x, |Du|) dx \quad (8.29)$$

holds for some constant $c \equiv c(n, \nu, L)$.

Proof. Essentially, the above lemma is a special case of Lemma 7.4 since we consider a particular case that $\omega_a(t) = t^\alpha$ and $\omega_b(t) = t^\beta$ for some $\alpha, \beta \in (0, 1)$. But our purpose here is to obtain an estimate such as (8.28) with a different multiplier containing some power of R , which will be used for proving Theorem 1.1. Therefore, we are able to apply Theorem 1.2. In turn, for every $\theta \in (0, 1)$ and open subset $\Omega_0 \Subset \Omega$, there exists a constant $c \equiv c(\mathbf{data}(\Omega_0), \theta)$ such that

$$[u]_{0,\theta;\Omega_0} \leq c(\mathbf{data}(\Omega_0), \theta). \quad (8.30)$$

In particular, we choose $\theta := (\gamma + 1)/2$. Now we need to revisit the proof of Lemma 7.3. Under the settings of the proof of Lemma 7.3, we turn our attention to estimating the terms I_a and I_b introduced in (7.59)-(7.60). Using (1.14), (7.26) and (7.6), we have

$$\begin{aligned} I_a &\leq cR^\alpha \left(1 + \left[(\Psi_{B_R}^-)^{-1} \left(\int_{B_{R/2}} \Psi_{B_R}^- \left(\left| \frac{w_c - (w_c)_{B_{R/2}}}{R} \right| \right) dx \right) \right]^{\frac{\alpha}{1-\gamma}} \right) \\ &\leq c \left(R^\alpha + R^{-\frac{\alpha\gamma}{1-\gamma}} \left[\text{osc}_{B_{2R}} u \right]^{\frac{\alpha}{1-\gamma}} \right) \leq c(\mathbf{data}(\Omega_0)) R^{\alpha/2}, \end{aligned} \quad (8.31)$$

where we have used (8.30) with the choice of $\theta := (1 + \gamma)/2$ and $B_{2R} \subset \Omega_0$ with $R \leq 1$. In the same way, we show

$$I_b \leq c(\mathbf{data}(\Omega_0)) R^{\beta/2}. \quad (8.32)$$

Inserting those estimates into (7.60), we see that

$$\left| \int_{B_{1/8}} \langle \bar{A}_0(D\bar{w}_c), D\varphi \rangle dx \right| \leq c(\mathbf{data}(\Omega_0)) q_3(R) \|D\varphi\|_{L^\infty(B_{1/8})}, \quad (8.33)$$

whenever $\varphi \in W_0^{1,\infty}(B_{1/8})$, where

$$q_3(R) := R^{\frac{\alpha}{2(1+s(H_a))}} + R^{\frac{\beta}{2(1+s(H_b))}}. \quad (8.34)$$

Note that the vector field \bar{A}_0 has been defined in (7.50). We consider a function $\bar{h} \in \bar{w}_c + W_0^{1,\Psi_0}(B_{1/8})$ satisfying (7.94)-(7.97) with the term $d(\varepsilon, R)$ replaced by $q_3(R)$. Let h be the scaled back function of \bar{h} in $B_{R/8}$ as

$$h(x) := E(w_c, B_{R/2}) R \bar{h} \left(\frac{x - x_0}{R} \right) \quad \text{for every } x \in B_{R/8}(x_0). \quad (8.35)$$

Clearly, $h \in w_c + W_0^{1,\Psi_{B_R}^-}(B_{R/8})$ is a local minimizer of the functional \mathcal{F}_0 defined in (7.103), which means that

$$\mathcal{F}_0(h) = \int_{B_{R/8}} F_0(Dh) dx \leq \int_{B_{R/8}} F_0(Dh + D\varphi) dx = \mathcal{F}_0(h + \varphi) \quad (8.36)$$

holds for every $\varphi \in W_0^{1,\Psi_{B_R}^-}(B_{R/8})$. As we have shown in (7.9), we recall (7.24) and (7.4) to see that

$$\begin{aligned} \int_{B_{R/8}} \Psi_{B_R}^- (|Dh|) dx &\leq \frac{L}{\nu} \int_{B_{R/8}} \Psi_{B_R}^- (|Dw_c|) dx \leq \frac{8^n L}{\nu} \int_{B_R} \Psi(x, |Dw_c|) dx \\ &\leq c(n, \nu, L) \int_{B_R} \Psi(x, |Dw|) dx \leq c(n, \nu, L) \int_{B_{2R}} \Psi(x, |Du|) dx, \end{aligned} \quad (8.37)$$

which proves (7.106). We write the inequality (7.96) in view of G, H_a, H_b, w_c and h in order to have

$$\begin{aligned} &\int_{B_{R/8}} [|V_G(Dw_c) - V_G(Dh)|^2 + a(x_a) |V_{H_a}(Dw_c) - V_{H_a}(Dh)|^2 + b(x_b) |V_{H_b}(Dw_c) - V_{H_b}(Dh)|^2] dx \\ &\leq c[q_3(R)]^{s_1} \int_{B_{R/2}} \Psi_{B_R}^- \left(\left| \frac{w_c - (w_c)_{B_{R/2}}}{R} \right| \right) dx \leq c[q_3(R)]^{s_1} \int_{B_{R/2}} \Psi_{B_R}^- (|Dw_c|) dx \\ &\leq c[q_3(R)]^{s_1} \int_{B_{R/2}} \Psi(x, |Du|) dx \end{aligned} \quad (8.38)$$

for some constant $c \equiv c(\mathbf{data}(\Omega_0))$, where we have applied Sobolev-Poincaré inequality and (7.109). Combining this estimate together with (7.3) and (7.23) alongside some elementary computations, we arrive at the desired estimate (8.28). \square

9. PROOF OF THEOREM 1.1.

Finally, we are ready to prove Theorem 1.1. First applying Theorem 1.2 and a standard covering argument, we find that for every open subset $\Omega_0 \Subset \Omega$ and any number $k > 0$, there exists a constant $c \equiv c(\mathbf{data}(\Omega_0), k)$ such that

$$\int_{B_{2R}} \Psi(x, |Du|) dx \leq cR^{-k}, \quad (9.1)$$

whenever $B_{2R} \subset \Omega_0$ is a ball with $R \leq 1$. Now we fix an open subset $\Omega_0 \Subset \Omega$ and a ball $B_{2R} \equiv B_{2R}(x_0) \subset \Omega_0$ with $R \leq 1$. Then applying Lemma 7.4 and Lemma 8.1,

$$\begin{aligned}
& \int_{B_{R/8}} (|V_G(Du) - V_G(Dh)|^2 + a(x_a)|V_{H_a}(Du) - V_{H_a}(Dh)|^2 + b(x_b)|V_{H_b}(Du) - V_{H_b}(Dh)|^2) dx \\
& \leq c(R^{\mu\gamma} + [q(R)]^{s_1}) \int_{B_{2R}} \Psi(x, |Du|) dx
\end{aligned} \tag{9.2}$$

for some constant $c \equiv c(\mathbf{data}(\Omega_0))$ and $s_1 \equiv s_1(\mathbf{data})$, where

$$q(R) := \begin{cases} R^{\frac{\alpha\delta}{(1+\delta)(1+s(H_a))}} + R^{\frac{\beta\delta}{(1+\delta)(1+s(H_b))}} & \text{if (1.21a) is assumed,} \\ R^{\frac{\alpha\gamma}{1+s(H_a)}} + R^{\frac{\beta\gamma}{1+s(H_b)}} & \text{if (1.21b) is assumed,} \\ R^{\frac{\alpha}{2(1+s(H_a))}} + R^{\frac{\beta}{2(1+s(H_b))}} & \text{if (1.21c) is assumed,} \end{cases} \tag{9.3}$$

in which γ is the Hölder continuity exponent determined via Theorem 5.2 and δ is the higher integrability exponent coming from Theorem 5.5. We denote by

$$d \equiv d(\mathbf{data}(\Omega_0)) := \begin{cases} \min \left\{ \mu\gamma, \frac{\alpha\delta s_1}{(1+\delta)(1+s(H_a))}, \frac{\beta\delta s_1}{(1+\delta)(1+s(H_b))} \right\} & \text{if (1.21a) is assumed,} \\ \min \left\{ \mu\gamma, \frac{\alpha\gamma s_1}{1+s(H_a)}, \frac{\beta\gamma s_1}{1+s(H_b)} \right\} & \text{if (1.21b) is assumed,} \\ \min \left\{ \mu\gamma, \frac{\alpha s_1}{2(1+s(H_a))}, \frac{\beta s_1}{2(1+s(H_b))} \right\} & \text{if (1.21c) is assumed,} \end{cases} \tag{9.4}$$

and $x_a, x_b \in \overline{B_R}$ are points such that $a(x_a) = \inf_{x \in B_R} a(x)$ and $b(x_b) = \inf_{x \in B_R} b(x)$. Now choosing $k \equiv d/4$ in (9.1), the inequality (9.2) can be written as

$$\begin{aligned}
& \int_{B_{R/8}} (|V_G(Du) - V_G(Dh)|^2 + a(x_a)|V_{H_a}(Du) - V_{H_a}(Dh)|^2 + b(x_b)|V_{H_b}(Du) - V_{H_b}(Dh)|^2) dx \\
& \leq cR^{3d/4}
\end{aligned} \tag{9.5}$$

for some constant $c \equiv c(\mathbf{data}(\Omega_0))$, where we again recall that the function h has been defined via Lemma 7.4 and Lemma 8.1. Recalling, (7.106) and (8.29), we have the energy estimate

$$\int_{B_{R/8}} \Psi_{B_R}^-(|Dh|) dx \leq c \int_{B_{2R}} \Psi(x, |Du|) dx \tag{9.6}$$

with a constant $c \equiv c(n, \nu, L)$. Now using repeatedly (2.10), we have

$$\begin{aligned}
& \int_{B_{R/8}} \Psi_{B_R}^-(|Du - Dh|) dx \leq c \int_{B_{R/8}} \left([\Psi_{B_R}^-(|Du| + |Dh|)]^{\frac{1}{2}} \frac{|Du - Dh|}{|Du| + |Dh|} \right) [\Psi_{B_R}^-(|Du| + |Dh|)]^{\frac{1}{2}} dx \\
& \leq c \left(\int_{B_{R/8}} \Psi_{B_R}^-(|Du| + |Dh|) \frac{|Du - Dh|^2}{(|Du| + |Dh|)^2} dx \right)^{\frac{1}{2}} \left(\int_{B_{R/8}} \Psi_{B_R}^-(|Du| + |Dh|) dx \right)^{\frac{1}{2}} \\
& \stackrel{(2.10), (9.5)}{\leq} cR^{3d/8} \left(\int_{B_{R/8}} \Psi_{B_R}^-(|Du| + |Dh|) dx \right)^{\frac{1}{2}} \stackrel{(9.6), (9.1)}{\leq} cR^{d/4}
\end{aligned} \tag{9.7}$$

with $c \equiv c(\mathbf{data}(\Omega_0))$, where d has been introduced in (9.3). Since h is a minimizer of functional \mathcal{F}_0 defined in (7.103), and this functional satisfies the growth and ellipticity conditions (1.16)_{1,2} with $a(x) \equiv a(x_a)$ and $b(x) \equiv b(x_b)$, we are able to apply the theory in [65], which provides the gradient Hölder regularity with the estimates

$$\begin{aligned}
& \int_{B_\rho} \Psi_{B_R}^-(|Dh - (Dh)_{B_\rho}|) dx \leq c \left(\frac{\rho}{R} \right)^{\beta_1} \int_{B_{R/8}} \Psi_{B_R}^-(|Dh|) dx \\
& \stackrel{(9.6)}{\leq} c \left(\frac{\rho}{R} \right)^{\beta_1} \int_{B_{2R}} \Psi(x, |Du|) dx,
\end{aligned} \tag{9.8}$$

whenever $0 < \rho \leq R/8$, where the constants c, β_1 depend only on $n, s(G), s(H_a), s(H_b), \nu, L$, but are independent of the values $a(x_a)$ and $b(x_b)$. Therefore, for every $0 < \rho \leq R/8$, we have

$$\begin{aligned}
& \oint_{B_\rho} G(|Du - (Du)_{B_\rho}|) dx \\
& \leq c \oint_{B_\rho} G(|Dh - (Dh)_{B_\rho}|) dx + c \oint_{B_\rho} G(|Du - Dh|) dx \\
& \stackrel{(9.8)}{\leq} c \left(\frac{\rho}{R}\right)^{\beta_1} \oint_{B_{2R}} \Psi(x, |Du|) dx + c \left(\frac{R}{\rho}\right)^n \oint_{B_{R/8}} G(|Du - Dh|) dx \\
& \stackrel{(9.1), (9.7)}{\leq} c \left(\frac{\rho}{R}\right)^{\beta_1} R^{-k} + c \left(\frac{R}{\rho}\right)^n R^{d/4}
\end{aligned} \tag{9.9}$$

with $c \equiv c(\mathbf{data}(\Omega_0), k)$. Notice that $k \in (0, 1)$ is still arbitrary and d has been defined in (9.3) depending only on $\mathbf{data}(\Omega_0)$. Taking $k \equiv d\beta_1/(32n)$ and $\rho \equiv (R/8)^{1+d/(16n)}$ in the last display, after some elementary manipulations, we get

$$\oint_{B_\rho} G(|Du - (Du)_{B_\rho}|) dx \leq c \rho^{\frac{d\beta_1}{64n}} \tag{9.10}$$

for every $\rho \in (0, 1/8)$, provided $B_{8\rho} \Subset \Omega_0$. In particular, using Jensen's inequality and Lemma 2.1₁, we have

$$\oint_{B_\rho} |Du - (Du)_{B_\rho}| dx \leq c \rho^{\frac{d\beta_1}{64n} \left(1 + \frac{1}{s(G)}\right)} \tag{9.11}$$

for every $\rho \in (0, 1/8)$ with $B_{8\rho} \Subset \Omega_0$. By the integral characterization of Hölder continuity due to Campanato and Meyers and a standard covering argument alongside (9.11), $Du \in C_{\text{loc}}^{0, \theta}(\Omega)$ for $\theta \equiv \frac{d\beta_1}{64n} \left(1 + \frac{1}{s(G)}\right)$. This proves the local Hölder continuity of Du . But the proof is not finished yet, since θ should be independent of Ω_0 as in the statement of Theorem 1.1. In order to obtain the full completeness, we apply some standard perturbation methods. Indeed, once we have that Du is locally bounded, we shall revisit the proof of Lemma 7.4 and Lemma 8.1. We also observe that the functional defined in (7.103) satisfies the bounded slope condition (see for instance [17]). Then there exists a constant $c \equiv c(n, s(G), s(H_a), s(H_b), \nu, L, \|Du\|_{L^\infty(B_R)})$ such that

$$\|Dh\|_{L^\infty(B_R)} \leq c.$$

Since Du is locally bounded, following the proof of Lemma 7.1, Lemma 7.4 and Lemma 8.1, specially the estimate in (7.3) can be modified with $\gamma \equiv 1$. Moreover, the estimates in (8.28) and (7.105) can be upgraded by

$$\begin{aligned}
& \oint_{B_{R/8}} (|V_G(Du) - V_G(Dh)|^2 + a(x_a)|V_{H_a}(Du) - V_{H_a}(Dh)|^2 + b(x_b)|V_{H_b}(Du) - V_{H_b}(Dh)|^2) dx \\
& \leq c R^{\min\{\mu, \alpha, \beta\}}
\end{aligned} \tag{9.12}$$

with some constant $c \equiv c(n, s(G), s(H_a), s(H_b), \nu, L, \|a\|_{L^\infty(\Omega_0)}, \|b\|_{L^\infty(\Omega_0)}, \|Du\|_{L^\infty(B_{2R})})$. In particular, the last estimate via (9.7) implies that

$$\oint_{B_R} G(|Du - Dh|) dx \leq c R^{\min\{\mu, \alpha, \beta\}/4}. \tag{9.13}$$

Therefore, (9.8) implies that

$$\oint_{B_\rho} G(|Dh - (Dh)_{B_\rho}|) dx \leq c \left(\frac{\rho}{R}\right)^{\beta_1}, \tag{9.14}$$

where β_1 depends on $n, s(G), s(H_a), s(H_b), \nu, L$ while the constant c depends only on $n, s(G), s(H_a), s(H_b), \nu, L, \|Du\|_{L^\infty(\Omega_0)}, \|a\|_{L^\infty(\Omega_0)}$ and $\|b\|_{L^\infty(\Omega_0)}$. Combining the last two estimates similarly as shown in (9.9), we deduce the gradient Hölder continuity with the exponent depending only on $n, s(G), s(H_a), s(H_b), \nu, L, \alpha, \beta$ and μ , which is the desired dependence as described in the statement. The proof is finally complete.

10. ORLICZ DOUBLE PHASE PROBLEMS

Let us consider a general class of functionals with double phase growth, which is essentially the case when $b(\cdot) \equiv 0$ in (1.2). The functionals we shall deal with is of type

$$W^{1,1}(\Omega) \ni v \mapsto \mathcal{F}_d(v, \Omega) := \int_{\Omega} F_d(x, v, Dv) dx, \quad (10.1)$$

where $F_d : \Omega \times \mathbb{R} \times \mathbb{R}^n \rightarrow \mathbb{R}$ is a Caratheódory function fulfilling the following double-sided growth

$$\nu \Psi_d(x, |z|) \leq F_d(x, y, z) \leq L \Psi_d(x, |z|), \quad (10.2)$$

whenever $x \in \Omega, y \in \mathbb{R}$ and $z \in \mathbb{R}^n$, in which here and in the rest of the paper we denote by

$$\Psi_d(x, t) := G(t) + a(x)H_a(t) \quad (\forall x \in \Omega, t \geq 0). \quad (10.3)$$

As we introduced before we assume $G, H_a \in \mathcal{N}$ with indices $s(G), s(H_a) \geq 1$ and $a(\cdot) \in C^{\omega_a}(\Omega)$ with $\omega_a : [0, \infty) \rightarrow [0, \infty)$ being a concave function such that $\omega_a(0) = 0$. We shall consider a local minimizer u of the functional \mathcal{F}_d in (10.1) under one of the assumptions (1.12), (1.13) and (1.14) with $\omega_b(\cdot) \equiv 0$. Since the double sided growth assumption (10.3) is not enough for higher regularity properties of a local minimizer u of the functional \mathcal{F}_d , we shall assume that F_d is a continuous integrand belonging to the space $C^2(\mathbb{R}^n \setminus \{0\})$ with respect to z -variable and having the the following structure assumptions:

$$\begin{cases} |D_z F_d(x, y, z)| |z| + |D_{zz}^2 F_d(x, y, z)| |z|^2 \leq L \Psi_d(x, |z|), \\ \nu \frac{\Psi_d(x, |z|)}{|z|^2} |\xi|^2 \leq \langle D_{zz}^2 F_d(x, y, z) \xi, \xi \rangle, \\ |D_z F_d(x_1, y, z) - D_z F_d(x_2, y, z)| |z| \leq L \omega(|x_1 - x_2|) [\Psi_d(x_1, |z|) + \Psi_d(x_2, |z|)] \\ \quad + L |\Psi_d(x_1, |z|) - \Psi_d(x_2, |z|)|, \\ |F_d(x, y_1, z) - F_d(x, y_2, z)| \leq L \omega(|y_1 - y_2|) \Psi_d(x, |z|), \end{cases} \quad (10.4)$$

whenever $x, x_1, x_2 \in \Omega, y, y_1, y_2 \in \mathbb{R}, z \in \mathbb{R}^n \setminus \{0\}, \xi \in \mathbb{R}^n$, where $0 < \nu \leq L$ are fixed constants, and the function ω is the same as defined in (1.17) or (1.18). The structure conditions in (10.4) are satisfied for instance by the model functional

$$W^{1,1}(\Omega) \ni v \mapsto \int_{\Omega} f_d(x, v) \Psi_d(x, |Dv|) dx, \quad (10.5)$$

where the continuous function $f_d(\cdot)$ satisfies $0 < \nu_0 \leq f(\cdot, \cdot) \leq L_0$ for some constants ν_0, L_0 and fulfills the following inequality

$$|f_d(x_1, y_1) - f_d(x_2, y_2)| \leq L_0 \omega(|x_1 - x_2| + |y_1 - y_2|),$$

whenever $x_1, x_2 \in \mathbb{R}^n$ and $y_1, y_2 \in \mathbb{R}$, in which ω is the same as defined in (1.17) or (1.18). Another model case is given by

$$W^{1,1}(\Omega) \ni v \mapsto \int_{\Omega} [F_G(x, v, Dv) + a(x)F_{H_a}(x, v, Dv)] dx, \quad (10.6)$$

where $F_G(\cdot)$ and $F_{H_a}(\cdot)$ have G -growth and H_a -growth respectively, and satisfy the following suitable structure assumptions that

$$\begin{cases} |D_z F_{\Phi}(x, y, z)| |z| + |D_{zz}^2 F_{\Phi}(x, y, z)| |z|^2 \leq L_0 \Phi(|z|), \\ \nu_0 \frac{\Phi(|z|)}{|z|^2} |\xi|^2 \leq \langle D_{zz}^2 F_{\Phi}(x, y, z) \xi, \xi \rangle, \\ |D_z F_{\Phi}(x_1, y, z) - D_z F_{\Phi}(x_2, y, z)| |z| \leq L_0 \omega(|x_1 - x_2|) \Phi(|z|), \\ |F_{\Phi}(x, y_1, z) - F_{\Phi}(x, y_2, z)| \leq L_0 \omega(|y_1 - y_2|) \Phi(|z|) \end{cases}$$

hold with $\Phi \in \{G, H_a\}$ for some positive constants ν_0, L_0 , where ω is as in (1.17) or (1.18). The reason we consider the double phase case independently is that we have discussed the various regularity properties of the functional \mathcal{F} in (1.9) in the sense of multi-phase of the type defined in (1.15) together with the structure assumptions (1.16), but this one is a special case of (10.1) together with the structure assumptions (10.4) in the sense of the double phase structures. Now we restate and prove Lemma 2.5 in the double phase settings which will be applied later.

Lemma 10.1. *Let $F_d : \Omega \times \mathbb{R} \times \mathbb{R}^n \rightarrow \mathbb{R}$ be a function defined in (10.1) which satisfies (10.2) and (10.4). There exist positive constants $c_1, c_2 \equiv c_1, c_2(n, s(G), s(H_a), \nu)$ such that the following inequalities*

$$|V_G(z_1) - V_G(z_2)|^2 + a(x)|V_{H_a}(z_1) - V_{H_a}(z_2)|^2 \leq c_1 \langle D_z F_d(x, y, z_1) - D_z F_d(x, y, z_2), z_1 - z_2 \rangle, \quad (10.7)$$

$$\begin{aligned} |V_G(z_1) - V_G(z_2)|^2 + a(x)|V_{H_a}(z_1) - V_{H_a}(z_2)|^2 &+ c_2 \langle D_z F_d(x, y, z_1), z_2 - z_1 \rangle \\ &\leq c_2 [F_d(x, y, z_2) - F_d(x, y, z_1)] \end{aligned} \quad (10.8)$$

and

$$|F_d(x_1, y, z) - F_d(x_2, y, z)| \leq L\omega(|x_1 - x_2|) [\Psi_d(x_1, |z|) + \Psi_d(x_2, |z|)] + L|a(x_1) - a(x_2)|H_a(|z|) \quad (10.9)$$

hold true, whenever $z, z_1, z_2 \in \mathbb{R}^n \setminus \{0\}$, $x, x_1, x_2 \in \Omega$ and $y \in \mathbb{R}$.

Proof. The arguments of the proof for (10.7) and (10.8) are essentially the same as done for Lemma 2.5. Only difference lies in the one for (10.9). Since $F_d(x, y, 0) = 0$ for every $x \in \Omega$ and $y \in \mathbb{R}$, we have

$$\begin{aligned} |F_d(x_1, y, z) - F_d(x_2, y, z)| &= |(F_d(x_1, y, z) - F_d(x_1, y, 0)) - (F_d(x_2, y, z) - F_d(x_2, y, 0))| \\ &= \left| \int_0^1 \langle D_z F_d(x_1, y, \theta z), z \rangle d\theta - \int_0^1 \langle D_z F_d(x_2, y, \theta z), z \rangle d\theta \right| \\ &\leq \int_0^1 |D_z F_d(x_1, y, \theta z) - D_z F_d(x_2, y, \theta z)| |z| d\theta \\ &\leq L\omega(|x_1 - x_2|) [\Psi_d(x_1, |z|) + \Psi_d(x_2, |z|)] + L|a(x_1) - a(x_2)|H_a(|z|), \end{aligned}$$

where the last inequality of the last display is implied by (10.4)₃. This proves (10.9). \square

In order to simplify the notations in the present section, we use the set of parameters for a minimizer u of the functional \mathcal{F}_d depending on which one of the assumptions (1.12)-(1.14) under $\omega_b(\cdot) \equiv 0$ comes into play as the **data** in this section.

$$\mathbf{data}_d \equiv \begin{cases} n, \lambda_1, s(G), s(H_a), \nu, L, \|a\|_{C^{\omega_a}(\Omega)}, \omega(\cdot), \|\Psi(x, |Du|)\|_{L^1(\Omega)}, \|u\|_{L^1(\Omega)}, \omega_a(1) \\ \text{if (1.12) is considered under } \omega_b(\cdot) \equiv 0, \\ n, \lambda_2, s(G), s(H_a), \nu, L, \|a\|_{C^{\omega_a}(\Omega)}, \omega(\cdot), \|u\|_{L^\infty(\Omega)}, \omega_a(1) \\ \text{if (1.13) is considered under } \omega_b(\cdot) \equiv 0, \\ n, \lambda_3, s(G), s(H_a), \nu, L, \|a\|_{C^{\omega_a}(\Omega)}, \omega(\cdot), [u]_{0,\gamma}, \omega_a(1) \\ \text{if (1.14) is considered under } \omega_b(\cdot) \equiv 0, \end{cases} \quad (10.10)$$

where $\lambda_1, \lambda_2, \lambda_3$ are the same as defined in (1.12)-(1.14) and $s(G), s(H_a)$ are indices of the functions G, H_a in the sense of Definition 2.1, respectively. With $\Omega_0 \Subset \Omega$ being a fixed open subset, we also denote by $\mathbf{data}_d(\Omega_0)$ the above set of parameters together with $\text{dist}(\Omega_0, \partial\Omega)$:

$$\mathbf{data}_d(\Omega_0) \equiv \mathbf{data}_d, \text{dist}(\Omega_0, \partial\Omega). \quad (10.11)$$

Now we provide the main results in this section, which correspond to Theorem 1.1 and Theorem 1.2.

Theorem 10.1 (Maximal regularity). *Let $u \in W^{1,\Psi_d}(\Omega)$ be a local minimizer of the functional \mathcal{F}_d defined in (10.1) under the assumptions (10.2), (10.4) and (1.17) with $\omega_b(\cdot) \equiv 0$. Suppose that $\omega_a(t) = t^\alpha$ for some*

$\alpha \in (0, 1]$. If one of the following assumptions

$$\begin{cases} (1.12), & (10.12a) \\ (1.13), & (10.12b) \\ (1.14) \text{ with } \limsup_{t \rightarrow 0^+} \Lambda \left(t^{\frac{1}{1-\gamma}}, \frac{1}{t} \right) = 0 & (10.12c) \end{cases}$$

is satisfied, then there exists $\theta \in (0, 1)$ depending only on $n, s(G), s(H_a), \nu, L, \alpha$ and μ such that $Du \in C_{\text{loc}}^{0, \theta}(\Omega)$.

Theorem 10.2 (Morrey decay). *Let $u \in W^{1, \Psi_d}(\Omega)$ be a local minimizer of the functional \mathcal{F}_d defined in (10.1), under the assumptions (10.2), (10.4) and (1.18). Assume that $\omega_b(\cdot) \equiv 0$ in what follows. If one of the following assumptions*

$$\begin{cases} (1.12) \text{ with } \limsup_{t \rightarrow 0^+} \Lambda(t, G^{-1}(t^{-n})) = 0, & (10.13a) \\ (1.13) \text{ with } \limsup_{t \rightarrow 0^+} \Lambda \left(t, \frac{1}{t} \right) = 0, & (10.13b) \\ (1.14) \text{ with } \limsup_{t \rightarrow 0^+} \Lambda \left(t^{\frac{1}{1-\gamma}}, \frac{1}{t} \right) = 0, & (10.13c) \\ (1.12) \text{ with } \omega_a(t) = t^\alpha \text{ for some } \alpha \in (0, 1], & (10.13d) \\ (1.13) \text{ with } \omega_a(t) = t^\alpha \text{ for some } \alpha \in (0, 1] & (10.13e) \end{cases}$$

is satisfied, then

$$u \in C_{\text{loc}}^{0, \theta}(\Omega) \text{ for every } \theta \in (0, 1). \quad (10.14)$$

Moreover, for every $\sigma \in (0, n)$, there exists a positive constant $c \equiv c(\text{data}_d(\Omega_0), \sigma)$ such that the decay estimate

$$\int_{B_\rho} \Psi_d(x, |Du|) dx \leq c \left(\frac{\rho}{R} \right)^{n-\sigma} \int_{B_R} \Psi_d(x, |Du|) dx \quad (10.15)$$

holds for every concentric balls $B_\rho \subset B_R \subset \Omega_0 \Subset \Omega$ with $R \leq 1$.

The above theorems completely cover the main results of [13], where the special case that $G(t) = t^p$, $H_a(t) = t^q$ and $\omega_a(t) = t^\alpha$ with some constants $q \geq p > 1$ and $\alpha \in (0, 1]$ is considered. Also the results of [12] can be considered for a general class of functionals not only for the model functional in (1.4). Let us now briefly overview our arguments employed in proving the above theorems comparing with the ones used in [12, 13]. We do not distinguish between the G -phase, where an inequality of the type $a(\cdot) \leq M\omega_a(R)$ is satisfied, and (G, H_a) -phase, where a complementary inequality $a(\cdot) \geq M\omega_a(R)$ holds in a certain ball B_R under consideration for some suitable large constant M , which has a drawback to deal with the multi-phase type problems and even double phase type problems that we consider. Instead we consider the function $[\Psi_d]_{B_R}^-(\cdot)$ defined in (10.29) for a ball $B_R \subset \Omega$ under the investigation to obtain various estimates, and the advantage of considering this function is that $[\Psi_d]_{B_R}^- \in \mathcal{N}$ with an index $s(\Psi_d) = s(G) + s(H_a)$ by Remark 2.2, which is independent of the considered ball B_R . Also the approach introduced in this paper may open a gate to study parabolic double phase equations of type

$$u_t - \operatorname{div} \left(G'(|Du|) \frac{Du}{|Du|} + a(x, t) H'_a(|Du|) \frac{Du}{|Du|} \right) = 0,$$

which would be one of attracting topics for the regularity theory in the future, we refer some recent results on this topic [19, 39]. Essentially, the idea of the proofs of Theorem 10.2 and Theorem 10.1 is based on the arguments previously used for proving Theorem 1.2 and Theorem 1.1, but the functional \mathcal{F}_d in (10.1) is much more general than the functional \mathcal{F} in (1.9) for the consideration under the double phase settings. In this regard, we need to take care of some points in more detail depending on the structure assumptions (10.4), specially Lemma 10.3 below. Since $u \in W^{1, \Psi_d}(\Omega)$ is a local L/ν -minimizer of the functional \mathcal{P} defined in (1.1) with $b(\cdot) \equiv 0$ if u is a minimizer of the functional \mathcal{F}_d in (10.1), we are able to rewrite the results

together with their proofs under the double phase settings up to the end of Section 6. Starting by Section 7, we shall investigate in a different way.

In what follows let $B_R \equiv B_R(x_0)$ be a ball such that $B_{2R} \subset \Omega_0 \Subset \Omega$, where Ω_0 is some fixed open subset of Ω . We define a functional given by

$$W^{1,1}(B_{2R}) \ni v \mapsto \mathcal{F}_{d,B_{2R}}(v) := \int_{B_{2R}} F_d(x, (u)_{B_{2R}}, Dv) dx \quad (10.16)$$

with u being a local minimizer of the functional \mathcal{F}_d defined in (10.1). Now we consider a function $w \in u + W_0^{1,\Psi_d}(B_{2R})$ being the solution to the following variational Dirichlet problem:

$$\begin{cases} w \mapsto \min_v \mathcal{F}_{d,B_{2R}}(v) \\ v \in u + W_0^{1,\Psi_d}(B_{2R}). \end{cases} \quad (10.17)$$

As in Lemma 7.1 we shall consider the first comparison estimates in order to remove u -dependence in the original functional \mathcal{F}_d defined in (10.1).

Lemma 10.2. *Let $w \in W^{1,\Psi}(B_{2R})$ be the solution to the variational problem (10.17) under the assumptions (10.2), (10.4) and (1.18). Let the coefficient function $a(\cdot) \in C^{\omega_a}(\Omega)$ for ω_a being non-negative concave function vanishing at the origin. Assume that one of the assumptions (1.12), (1.13) and (1.14) under $\omega_b(\cdot) \equiv 0$ is satisfied. Then there exists a constant $c \equiv c(\mathbf{data}_d(\Omega_0))$ such that*

$$\int_{B_{2R}} (|V_G(Du) - V_G(Dw)|^2 + a(x)|V_{H_a}(Du) - V_{H_a}(Dw)|^2) dx \leq c\omega(R^\gamma) \int_{B_{2R}} \Psi_d(x, |Du|) dx \quad (10.18)$$

holds, where $\gamma \equiv \gamma(\mathbf{data}_d(\Omega_0))$ is the Hölder exponent determined via Theorem 5.2 in the double phase settings. Moreover, the following estimates holds true:

$$\int_{B_{2R}} \Psi_d(x, |Dw|) dx \leq \frac{L}{\nu} \int_{B_{2R}} \Psi_d(x, |Du|) dx, \quad (10.19)$$

$$\|w\|_{L^\infty(B_{2R})} \leq \|u\|_{L^\infty(B_{2R})}, \quad (10.20)$$

$$\text{osc}_{B_{2R}} w \leq \text{osc}_{B_{2R}} u \quad (10.21)$$

and

$$\int_{B_{2R}} \Psi_d\left(x, \left|\frac{u-w}{R}\right|\right) dx \leq c[\omega(R^\gamma)]^{\frac{1}{2}} \int_{B_{2R}} \Psi_d(x, |Du|) dx \quad (10.22)$$

for some constant $c \equiv c(\mathbf{data}_d(\Omega_0))$. Moreover, there exist a positive higher integrability exponent $\delta_0 \equiv \delta_0(\mathbf{data}_d)$ with $\delta_0 \leq \delta$, where δ has been determined via Theorem 5.5 under the double phase settings, and a constant $c \equiv c(\mathbf{data}_d)$ satisfying the following reverse Hölder inequalities:

$$\left[\int_{B_{R/2}} [\Psi_d(x, |Dw|)]^{1+\delta_0} dx \right]^{\frac{1}{1+\delta_0}} \leq c \int_{B_R} \Psi_d(x, |Dw|) dx. \quad (10.23)$$

Here, in the case that (1.14) is considered, γ appearing in (10.18) and (10.22) is the same as in the assumption (1.14).

Proof. First of all the meaning of \mathbf{data}_d and $\mathbf{data}_d(\Omega_0)$ has been defined in (10.10) and (10.11), respectively. The proofs for (10.19)-(10.22) can be done by arguing similarly as in the proof Lemma 7.1 together with Lemma 10.1. Since w is a L/ν -minimizer of the functional $\mathcal{F}_{d,B_{2R}}$ defined in (10.16), we are able to apply Lemma 5.1 under the double phase settings. In turn, it gives us that

$$\int_{B_{R/2}} \Psi_d(x, |Dw|) dx \leq c \int_{B_R} \Psi_d\left(x, \left|\frac{w - (w)_{B_R}}{R}\right|\right) dx \quad (10.24)$$

holds with $c \equiv c(n, s(G), s(H_a), L, \nu)$. Then applying Theorem 4.1, there exists $\theta \equiv \theta(n, s(G), s(H_a)) \in (0, 1)$ such that

$$\int_{B_{R/2}} \Psi_d(x, |Dw|) dx \leq c \bar{\kappa}_{sp} \left[\int_{B_R} [\Psi_d(x, |Dw|)]^\theta dx \right]^{\frac{1}{\theta}} \quad (10.25)$$

holds with some constant $c \equiv c(n, s(G), s(H_a), L, \nu, \omega_a(1))$, where

$$\bar{\kappa}_{sp} = \begin{cases} 1 + \lambda_1[a]_{\omega_a} + \lambda_1[a]_{\omega_a} \left(\int_{B_R} G(|Dw|) dx \right)^{\frac{1}{n}} & \text{if (1.12) is in force with } \omega_b(\cdot) \equiv 0, \quad (10.26a) \\ 1 + \lambda_2[a]_{\omega_a} + \lambda_2[a]_{\omega_a} \|w\|_{L^\infty(B_R)} & \text{if (1.13) is in force with } \omega_b(\cdot) \equiv 0, \quad (10.26b) \\ 1 + \lambda_3[a]_{\omega_a} + \lambda_3[a]_{\omega_a} \left[R^{-\gamma} \text{osc}_{B_R} w \right]^{\frac{1}{1-\gamma}} & \text{if (1.14) is in force with } \omega_b(\cdot) \equiv 0. \quad (10.26c) \end{cases}$$

Furthermore, taking into account (10.19)-(10.22) in the last display, we conclude that

$$\int_{B_{R/2}} \Psi_d(x, |Dw|) dx \leq c \left[\int_{B_R} [\Psi_d(x, |Dw|)]^\theta dx \right]^{\frac{1}{\theta}} \quad (10.27)$$

holds for some constants $\theta \equiv \theta(n, s(G), s(H_a)) \in (0, 1)$ and $c \equiv c(\mathbf{data}_d)$. The last display follows (10.23) by applying a variant of Gehring's lemma. \square

At this stage, we do not need to consider Lemma 7.2 because we shall freeze x -variable in the non-linearity at once. For this, let us consider the excess functional given by

$$E_d(v, B_r) := \left([\Psi_d]_{B_{2r}}^- \right)^{-1} \left(\int_{B_r} [\Psi_d]_{B_{2r}}^- \left(\left| \frac{v - (v)_{B_r}}{2r} \right| \right) dx \right) \quad (10.28)$$

for any function $v \in L^1(B_{2r})$ and ball $B_{2r} \subset \Omega$, where now and in the rest of this section for every open subset $\mathcal{B} \subset \Omega$, we shall denote by

$$[\Psi_d]_{\mathcal{B}}^-(t) := G(t) + \inf_{x \in \mathcal{B}} a(x) H_a(t) \quad (\forall t \geq 0), \quad (10.29)$$

and $\left([\Psi_d]_{\mathcal{B}}^- \right)^{-1}$ is the inverse function of $[\Psi_d]_{\mathcal{B}}^-$. By convexity of the function $[\Psi_d]_{B_{2r}}^-$ and Lemma 2.1, there is a constant $c \equiv c(s(G) + s(H_a))$ such that

$$E_d(v, B_r) \leq c \left([\Psi_d]_{B_{2r}}^- \right)^{-1} \left(\int_{B_r} [\Psi_d]_{B_{2r}}^- \left(\left| \frac{v - v_0}{2r} \right| \right) dx \right) \quad (10.30)$$

holds for every $v_0 \in \mathbb{R}$. Now we consider the estimates corresponding to the outcome of Lemma 7.3 under our double phase settings.

Lemma 10.3. *Let $u \in W^{1, \Psi_d}(\Omega)$ be a local minimizer of the functional \mathcal{F}_d defined in (10.1) under the assumptions (10.2), (10.4) and (1.18). Let $w \in W^{1, \Psi}(B_{2R})$ be the solution to the variational problem (10.17). Suppose $\omega_b(\cdot) \equiv 0$ in what follows. If one of the assumptions (10.13a)-(10.13e) is satisfied, then for every $\varepsilon^* \in (0, 1)$, there exists a positive radius*

$$R^* \equiv R^*(\mathbf{data}_d(\Omega_0), \varepsilon^*) \quad (10.31)$$

such that

$$\int_{B_{\tau R}} [\Psi_d]_{B_R}^- \left(\left| \frac{w - (w)_{B_{\tau R}}}{\tau R} \right| \right) dx \leq c \left(1 + \tau^{-(n+s(\Psi_d)+1)} \varepsilon^* \right) \int_{B_{R/2}} [\Psi_d]_{B_R}^- \left(\left| \frac{w - (w)_{B_{R/2}}}{R} \right| \right) dx \quad (10.32)$$

for some constant $c \equiv c(\mathbf{data}_d(\Omega_0))$, whenever $\tau \in (0, 1/16)$ and $R \leq R^*$.

Proof. Again note that the meaning of \mathbf{data}_d and $\mathbf{data}_d(\Omega_0)$ already has been introduced in (10.10)-(10.11). We can always assume $E_d(w, B_{R/2}) > 0$, otherwise there is nothing to prove in (10.32). For the simplicity, we shall write

$$E_d(R) := E_d(w, B_{R/2}), \quad (10.33)$$

where the notion E_d has been defined in (10.28). The proof falls in several steps, similarly as we have done in the proof of Lemma 7.3. For the sake of completeness, we provide the proof in a full detail.

Step 1: Initial information on w . Applying Lemma 5.2 under the double phase settings to $B_{R/2}$ with $k \equiv (w)_{B_{R/2}}$, we have

$$\int_{B_{R/4}} \Psi_d(x, |Dw|) \, dx \leq c \int_{B_{R/2}} [\Psi_d]_{B_R}^- \left(\left| \frac{w - (w)_{B_{R/2}}}{R} \right| \right) \, dx \quad (10.34)$$

for some constant $c \equiv c(\mathbf{data}_d)$. Moreover, it follows from Lemma 10.2 that there exists a higher integrability exponent $\delta_0 \equiv \delta_0(\mathbf{data}_d)$ such that

$$\left(\int_{B_{R/8}} [\Psi_d(x, |Dw|)]^{1+\delta_0} \, dx \right)^{\frac{1}{1+\delta_0}} \leq c \int_{B_{R/4}} \Psi_d(x, |Dw|) \, dx \quad (10.35)$$

for a constant $c \equiv c(\mathbf{data}_d)$.

Step 2: Scaled functions. We consider scaled functions of $w(\cdot)$ and $a(\cdot)$ in the ball B_1 by setting

$$\bar{w}(x) := \frac{w(x_0 + Rx) - (w)_{B_{R/2}}}{E_d(R)R}, \quad (10.36a)$$

$$\bar{a}(x) := a(x_0 + Rx) \frac{H_a(E_d(R))}{[\Psi_d]_{B_R}^-(E_d(R))} \quad (10.36b)$$

for every $x \in B_1$. Now we introduce the control function and energy density associated to our scaling introduced above in (10.36a)-(10.36b) as

$$\begin{cases} \bar{\Psi}_d(x, |z|) := \bar{G}(|z|) + \bar{a}(x)\bar{H}_a(|z|), \\ \bar{F}_d(x, z) := \frac{F_d(x_0 + Rx, (u)_{B_{2R}}, E_d(R)z)}{[\Psi_d]_{B_R}^-(E_d(R))} \quad \text{and} \quad \bar{A}_d(x, z) := D_z \bar{F}_d(x, z) \end{cases} \quad (10.37a)$$

$$\bar{F}_d(x, z) := \frac{F_d(x_0 + Rx, (u)_{B_{2R}}, E_d(R)z)}{[\Psi_d]_{B_R}^-(E_d(R))} \quad \text{and} \quad \bar{A}_d(x, z) := D_z \bar{F}_d(x, z) \quad (10.37b)$$

for every $x \in B_1$ and $z \in \mathbb{R}^n$, where to the end of the proof of this lemma, we always shall understand by

$$\bar{G}(t) := \frac{G(E_d(R)t)}{[\Psi_d]_{B_R}^-(E_d(R))} \quad \text{and} \quad \bar{H}_a(t) := \frac{H_a(E_d(R)t)}{H_a(E_d(R))} \quad (10.38)$$

for every $t \geq 0$. By elementary computations, we can observe that $\bar{G}, \bar{H}_a \in \mathcal{N}$ with indices $s(\bar{G}), s(\bar{H}_a)$, respectively, and also that

$$\bar{G}(1) \leq 1 \quad \text{and} \quad \bar{H}_a(1) = 1. \quad (10.39)$$

Clearly, the function \bar{w} minimizes the following functional

$$W^{1, \bar{\Psi}_d}(B_1) \ni v \mapsto \int_{B_1} \bar{F}_d(x, Dv) \, dx, \quad (10.40)$$

where the functions $\bar{\Psi}_d(\cdot)$ and $\bar{F}_d(\cdot)$ have been defined in (10.37a) and (10.37b), respectively. The Euler-Lagrange equation arising from the functional in (10.40) can be written as

$$\int_{B_1} \langle \bar{A}_d(x, D\bar{w}), D\varphi \rangle \, dx = \int_{B_1} \langle D_z \bar{F}_d(x, D\bar{w}), D\varphi \rangle \, dx = 0 \quad (10.41)$$

for every $\varphi \in W_0^{1,\bar{\Psi}_d}(B_1)$. By the assumptions (10.2) and (10.4) via elementary computations, we have the following structure conditions in the scaled settings:

$$\begin{aligned} \left\{ \begin{aligned} \nu \bar{\Psi}_d(x, |z|) &\leq \bar{F}_d(x, z) \leq L \bar{\Psi}_d(x, |z|), \\ |\bar{A}_d(x, z)| |z| + |D_z \bar{A}_d(x, z)| |z|^2 &\leq L \bar{\Psi}_d(x, |z|), \\ \nu \frac{\bar{\Psi}_d(x, |z|)}{|z|^2} |\xi|^2 &\leq \langle D_z \bar{A}_d(x, z) \xi, \xi \rangle, \\ |\bar{A}_d(x_1, z) - \bar{A}_d(x_2, z)| |z| &\leq L \omega(R|x_1 - x_2|) [\bar{\Psi}_d(x_1, |z|) + \bar{\Psi}_d(x_2, |z|)] \\ &\quad + L |\bar{a}(x_1) - \bar{a}(x_2)| \bar{H}_a(|z|) \end{aligned} \right. \end{aligned} \quad \begin{aligned} (10.42a) \\ (10.42b) \\ (10.42c) \\ (10.42d) \\ (10.42e) \end{aligned}$$

for every $x, x_1, x_2 \in B_1$ and $z \in \mathbb{R}^n \setminus \{0\}$.

Step 3: Freezing. Now we shall consider frozen functional and vector field associated to $\bar{F}_d(\cdot)$ and $\bar{A}_d(\cdot)$ defined in (10.37b). Let $\bar{x}_a \in \bar{B}_1$ such that $\bar{a}(\bar{x}_a) = \inf_{x \in B_1} \bar{a}(x)$. Then we denote by

$$\bar{F}_0(z) := \bar{F}_d(\bar{x}_a, z), \quad \bar{A}_0(z) := D_z \bar{F}_d(\bar{x}_a, z), \quad (10.43)$$

and

$$\bar{\Psi}_0(t) := \bar{G}(t) + \bar{a}(\bar{x}_a) \bar{H}_a(t) \quad (10.44)$$

for every $x \in B_1$, $z \in \mathbb{R}^n$ and $t \geq 0$. Here we single out that here is a difference between Step 3 of the proof for Lemma 7.3 and our present situation. By the very definition in (10.37a) and (10.38), one can check

$$\bar{\Psi}_0(1) = 1. \quad (10.45)$$

In our newly scaled environment, let us now consider the functional

$$W^{1,\bar{\Psi}_0}(B_{1/8}) \ni v \mapsto \int_{B_{1/8}} \bar{F}_0(Dv) dx. \quad (10.46)$$

We observe that the newly defined integrand $\bar{F}_0(\cdot)$ and vector field $\bar{A}_0(\cdot)$ satisfy the growth and ellipticity conditions as

$$\begin{aligned} \left\{ \begin{aligned} \nu \bar{\Psi}_0(|z|) &\leq \bar{F}_0(z) \leq L \bar{\Psi}_0(|z|), \\ |\bar{A}_0(z)| |z| + |D_z \bar{A}_0(z)| |z|^2 &\leq L \bar{\Psi}_0(|z|), \\ \nu \frac{\bar{\Psi}_0(|z|)}{|z|^2} |\xi|^2 &\leq \langle D_z \bar{A}_0(z) \xi, \xi \rangle \end{aligned} \right. \end{aligned} \quad \begin{aligned} (10.47a) \\ (10.47b) \\ (10.47c) \end{aligned}$$

for every $z \in \mathbb{R}^n \setminus \{0\}$ and $\xi \in \mathbb{R}^n$. Therefore, the energy and higher integrability estimates in (10.34) and (10.35) can be seen in the view of \bar{w} as

$$\int_{B_{1/4}} \bar{\Psi}_d(x, |D\bar{w}|) dx + \left(\int_{B_{1/8}} [\bar{\Psi}_d(x, |D\bar{w}|)]^{1+\delta_0} dx \right)^{\frac{1}{1+\delta_0}} \leq c(\mathbf{data}_d). \quad (10.48)$$

Step 4: Harmonic type approximation. Let $\varphi \in W_0^{1,\infty}(B_{1/8})$ be any fixed function. Using (10.41), we see

$$\begin{aligned} I_0 &:= \left| \int_{B_{1/8}} \langle \bar{A}_0(D\bar{w}), D\varphi \rangle dx \right| = \left| \int_{B_{1/8}} \langle \bar{A}_0(D\bar{w}) - \bar{A}_d(x, D\bar{w}), D\varphi \rangle dx \right| \\ &\leq \int_{B_{1/8}} |\bar{A}_0(D\bar{w}) - \bar{A}_d(x, D\bar{w})| dx \|D\varphi\|_{L^\infty(B_{1/8})} =: I_1 \|D\varphi\|_{L^\infty(B_{1/8})}. \end{aligned} \quad (10.49)$$

Now we estimate I_1 in the last display using (10.42d)-(10.42e). In turn, we have

$$\begin{aligned}
I_1 &\leq L\omega(R) \int_{B_{1/8}} \left(\frac{\bar{\Psi}_d(\bar{x}_a, |D\bar{w}|)}{|D\bar{w}|} + \frac{\bar{\Psi}_d(x, |D\bar{w}|)}{|D\bar{w}|} \right) dx \\
&\quad + L \int_{B_{1/8}} |\bar{a}(x) - \bar{a}(\bar{x}_a)| \frac{\bar{H}_a(|D\bar{w}|)}{|D\bar{w}|} dx \\
&\leq 2L\omega(R) \int_{B_{1/8}} \frac{\bar{\Psi}_d(\bar{x}_a, |D\bar{w}|)}{|D\bar{w}|} dx + 2L(1 + \omega(R)) \int_{B_{1/8}} |\bar{a}(x) - \bar{a}(\bar{x}_a)| \frac{\bar{H}_a(|D\bar{w}|)}{|D\bar{w}|} dx \\
&=: 2L\omega(R)I_{11} + 2L(1 + \omega(R))I_{12}.
\end{aligned} \tag{10.50}$$

Now we estimate the terms appearing in the last display. Recalling (10.44) and (10.45) together with (2.7), we find

$$\begin{aligned}
I_{11} &\leq c \int_{B_{1/8}} \bar{\Psi}'_0(|D\bar{w}|) dx \leq c [\bar{\Psi}_0(1)]^{\frac{s(\Psi_d)}{1+s(\Psi_d)}} \int_{B_{1/8}} [\bar{\Psi}_0(|D\bar{w}|)]^{\frac{1}{1+s(\Psi_d)}} dx \\
&\quad + c [\bar{\Psi}_0(1)]^{\frac{1}{1+s(\Psi_d)}} \int_{B_{1/8}} [\bar{\Psi}_0(|D\bar{w}|)]^{\frac{s(\Psi_d)}{1+s(\Psi_d)}} dx \\
&\leq c \left(\int_{B_{1/8}} \bar{\Psi}_0(|D\bar{w}|) dx \right)^{\frac{1}{1+s(\Psi_d)}} + c \left(\int_{B_{1/8}} \bar{\Psi}_0(|D\bar{w}|) dx \right)^{\frac{s(\Psi_d)}{1+s(\Psi_d)}} \\
&\leq c(\mathbf{data}_d),
\end{aligned} \tag{10.51}$$

where we have applied the Hölder's inequality together with (10.48) and the fact that $\bar{\Psi}_0 \in \mathcal{N}$ with an index $s(\Psi_d) = s(G) + s(H_a)$. Next we shall deal with estimating the second term I_{12} in (10.50). In turn, using (2.7) and (10.39), we have

$$\begin{aligned}
I_{12} &\leq c \int_{B_{1/8}} |\bar{a}(x) - \bar{a}(\bar{x}_a)| \left([\bar{H}_a(|D\bar{w}|)]^{\frac{1}{s(H_a)+1}} + [\bar{H}_a(|D\bar{w}|)]^{\frac{s(H_a)}{s(H_a)+1}} \right) dx \\
&\leq c \|\bar{a} - \bar{a}(\bar{x}_a)\|_{L^\infty(B_{1/8})}^{\frac{s(H_a)}{s(H_a)+1}} \left(\int_{B_{1/8}} \bar{a}(x) \bar{H}_a(|D\bar{w}|) dx \right)^{\frac{1}{s(H_a)+1}} \\
&\quad + c \|\bar{a} - \bar{a}(\bar{x}_a)\|_{L^\infty(B_{1/8})}^{\frac{1}{s(H_a)+1}} \left(\int_{B_{1/8}} \bar{a}(x) \bar{H}_a(|D\bar{w}|) dx \right)^{\frac{s(H_a)}{s(H_a)+1}} \\
&\leq c(\mathbf{data}_d) \left(\|\bar{a} - \bar{a}(\bar{x}_a)\|_{L^\infty(B_{1/8})}^{\frac{1}{s(H_a)+1}} + \|\bar{a} - \bar{a}(\bar{x}_a)\|_{L^\infty(B_{1/8})}^{\frac{s(H_a)}{s(H_a)+1}} \right),
\end{aligned} \tag{10.52}$$

where we have used also Hölder's inequality and the fact that $\bar{a}(\bar{x}_a) \leq \bar{a}(x)$ for every $x \in B_1$. Inserting those estimates coming from the last two displays into (10.50) and then (10.49), we find

$$I_0 \leq c(\mathbf{data}_d(\Omega_0)) \left[\omega(R) + (1 + \omega(R)) \left(\|\bar{a} - \bar{a}(\bar{x}_a)\|_{L^\infty(B_{1/8})}^{\frac{1}{s(H_a)+1}} + \|\bar{a} - \bar{a}(\bar{x}_a)\|_{L^\infty(B_{1/8})}^{\frac{s(H_a)}{s(H_a)+1}} \right) \right]. \tag{10.53}$$

Now we shall estimate the term $\|\bar{a} - \bar{a}(\bar{x}_a)\|_{L^\infty(B_{1/8})}$ depending on which one of the assumptions (10.13a)-(10.13e) comes into play. Recalling the definition of $\bar{a}(\cdot)$ in (10.36b) and the excess functional in (10.33), we have

$$I_a := \|\bar{a} - \bar{a}(\bar{x}_a)\|_{L^\infty(B_{1/8})} \leq c\omega_a(R) \frac{H_a(E_d(R))}{[\Psi_d]_{B_R}^-(E_d(R))}. \tag{10.54}$$

Case 1: Assumption (10.13a) is in force. It follows from the assumption (10.13a)₂ that for any $\varepsilon \in (0, 1)$ there exists $\mu_1 > 0$ depending on ε such that

$$\Lambda(t, G^{-1}(t^{-n})) \leq \varepsilon \quad \text{for every } t \in (0, \mu_1). \quad (10.55)$$

Then using the last display, (1.12) and the fact that $([\Psi_d]_{B_R}^-)^{-1}(t) \leq G^{-1}(t)$ for every $t \geq 0$, I_a in (10.54) can be estimated as

$$\begin{aligned} I_a &\leq c\omega_a(R) \frac{(H_a \circ G^{-1})([\Psi_d]_{B_R}^-(E_d(R)))}{[\Psi_d]_{B_R}^-(E_d(R))} \\ &\leq c\omega_a(R)\varepsilon \left(1 + \frac{1}{\omega_a([\Psi_d]_{B_R}^-(E_d(R)))^{-\frac{1}{n}}}\right) + c\omega_a(R) \left(1 + \frac{1}{\omega_a(\mu_1)}\right) \end{aligned} \quad (10.56)$$

with $c \equiv c([a]_{\omega_a}, \lambda_1)$. Using (2.2) and the energy estimate (10.19), we see

$$\begin{aligned} \frac{1}{\omega_a([\Psi_d]_{B_R}^-(E_d(R)))^{-\frac{1}{n}}} &\leq \frac{c}{\omega_a(R)} + \frac{c}{\omega_a(R)} \int_{B_{R/2}} [\Psi_d]_{B_R}^- \left(\left| \frac{w - (w)_{B_{R/2}}}{R} \right| \right) dx \\ &\leq \frac{c}{\omega_a(R)} + \frac{c}{\omega_a(R)} \int_{B_{2R}} \Psi_d(x, |Du|) dx \leq \frac{c(\mathbf{data}_d)}{\omega_a(R)}. \end{aligned} \quad (10.57)$$

Combining the last two displays, we conclude

$$I_a \leq c \left(\varepsilon + \omega_a(R) \left(1 + \frac{1}{\omega_a(\mu_1)}\right) \right) \quad (10.58)$$

with some constant $c \equiv c(\mathbf{data}_d)$. Therefore, inserting the estimates in the last two displays into (10.53) and recalling (10.49), we have

$$\left| \int_{B_{1/8}} \langle \bar{A}_0(D\bar{w}), D\varphi \rangle dx \right| \leq c(\mathbf{data}_d) P_1(\varepsilon, R) \|D\varphi\|_{L^\infty(B_{1/8})}, \quad (10.59)$$

where

$$\begin{aligned} P_1(\varepsilon, R) &:= \omega(R) + (1 + \omega(R)) \left[\varepsilon + \omega_a(R) \left(1 + \frac{1}{\omega_a(\mu_1)}\right) \right]^{\frac{1}{s(H_a)+1}} \\ &\quad + (1 + \omega(R)) \left[\varepsilon + \omega_a(R) \left(1 + \frac{1}{\omega_a(\mu_1)}\right) \right]^{\frac{s(H_a)}{s(H_a)+1}} \end{aligned} \quad (10.60)$$

Case 2: Assumption (10.13b) is in force. From the assumption (10.13b)₂ it holds that for every $\varepsilon \in (0, 1)$ there exists $\mu_2 > 0$ depending on ε such that

$$\Lambda\left(t, \frac{1}{t}\right) \leq \varepsilon \quad \text{for every } t \in (0, \mu_2). \quad (10.61)$$

Then by the very definition of $[\Psi_d]_{B_R}^-$ in (10.29) together with (10.61) and (1.13) under $\omega_b \equiv 0$, we have

$$\begin{aligned} I_a &\leq c\omega_a(R) \frac{H_a(E_d(R))}{G(E_d(R))} \\ &\leq c\omega_a(R)\varepsilon \left(1 + \frac{1}{\omega_a([E_d(R)]^{-1})}\right) + c\omega_a(R) \left(1 + \frac{1}{\omega_a(\mu_2)}\right). \end{aligned} \quad (10.62)$$

Again using (2.1) together with taking into account (10.20), we see

$$\frac{1}{\omega_a([E_d(R)]^{-1})} \leq \frac{1}{\omega_a\left(\frac{R}{2\|w\|_{L^\infty(B_R)}}\right)} \leq \frac{c(\mathbf{data}_d)}{\omega_a(R)}. \quad (10.63)$$

Combining the last two displays, we find

$$I_a \leq c \left(\varepsilon + \omega_a(R) \left(1 + \frac{1}{\omega_a(\mu_2)} \right) \right) \quad (10.64)$$

with some constant $c \equiv c(\mathbf{data}_d)$. Then, plugging the estimates in the last two displays into (10.53) and recalling (10.49), we have

$$\left| \int_{B_{1/8}} \langle \bar{A}_0(D\bar{w}), D\varphi \rangle dx \right| \leq c(\mathbf{data}_d) P_2(\varepsilon, R) \|D\varphi\|_{L^\infty(B_{1/8})}, \quad (10.65)$$

where

$$\begin{aligned} P_2(\varepsilon, R) &:= \omega(R) + (1 + \omega(R)) \left[\varepsilon + \omega_a(R) \left(1 + \frac{1}{\omega_a(\mu_2)} \right) \right]^{\frac{1}{s(H_a)+1}} \\ &\quad + (1 + \omega(R)) \left[\varepsilon + \omega_a(R) \left(1 + \frac{1}{\omega_a(\mu_2)} \right) \right]^{\frac{s(H_a)}{s(H_a)+1}}. \end{aligned} \quad (10.66)$$

Case 3: Assumption (10.13c) is in force. The assumption (10.13c)₂ implies that for any $\varepsilon \in (0, 1)$, there exists $\mu_3 > 0$ depending on ε such that

$$\Lambda \left(t^{\frac{1}{1-\gamma}}, \frac{1}{t} \right) \leq \varepsilon \quad \text{for every } t \in (0, \mu_3). \quad (10.67)$$

This one together with using (10.54) and (1.14) under $\omega_b(\cdot) \equiv 0$ implies

$$\begin{aligned} I_a &\leq c\omega_a(R) \frac{H_a(E_d(R))}{G(E_d(R))} \\ &\leq c\omega_a(R)\varepsilon \left(1 + \frac{1}{\omega_a([E_d(R)]^{-\frac{1}{1-\gamma}})} \right) + c\omega_a(R) \left(1 + \frac{1}{\omega_a\left(\mu_3^{\frac{1}{1-\gamma}}\right)} \right). \end{aligned} \quad (10.68)$$

Now using (10.21) and (1.14), we have

$$\frac{1}{\omega_a([E_d(R)]^{-\frac{1}{1-\gamma}})} \leq \frac{1}{\omega_a\left(\left[\frac{\text{osc}_{B_{2R}} u}{R}\right]^{-\frac{1}{1-\gamma}}\right)} \leq \frac{c(\mathbf{data}_d)}{\omega_a(R)}. \quad (10.69)$$

Combining the last two displays, we find

$$I_a \leq c \left(\varepsilon + \omega_a(R) \left(1 + \frac{1}{\omega_a\left(\mu_3^{\frac{1}{1-\gamma}}\right)} \right) \right) \quad (10.70)$$

for some constant $c \equiv c(\mathbf{data}_d)$. Using the estimate (10.70) in (10.53), we conclude

$$\left| \int_{B_{1/8}} \langle \bar{A}_0(D\bar{w}), D\varphi \rangle dx \right| \leq c(\mathbf{data}_d) P_3(\varepsilon, R) \|D\varphi\|_{L^\infty(B_{1/8})}, \quad (10.71)$$

where

$$\begin{aligned}
 P_3(\varepsilon, R) &:= \omega(R) + (1 + \omega(R)) \left[\varepsilon + \omega_a(R) \left(1 + \frac{1}{\omega_a \left(\mu_3^{\frac{1}{1-\gamma}} \right)} \right) \right]^{\frac{1}{s(H_a)+1}} \\
 &+ (1 + \omega(R)) \left[\varepsilon + \omega_a(R) \left(1 + \frac{1}{\omega_a \left(\mu_3^{\frac{1}{1-\gamma}} \right)} \right) \right]^{\frac{s(H_a)}{s(H_a)+1}}.
 \end{aligned} \tag{10.72}$$

Case 4. Assumption (10.13d) is in force. Now we take the advantage that $w_a(\cdot)$ is the power function. Recalling I_a denoted in (10.54), we see that

$$\begin{aligned}
 I_a &\leq cR^\alpha \frac{(H_a \circ G^{-1}) \left([\Psi_d]_{B_R}^-(E_d(R)) \right)}{[\Psi_d]_{B_R}^-(E_d(R))} \leq cR^\alpha \left(1 + \left[\int_{B_{R/2}} [\Psi_d]_{B_R}^- \left(\left| \frac{w - (w)_{B_{R/2}}}{R} \right| \right) dx \right]^{\frac{\alpha}{n}} \right) \\
 &\leq cR^\alpha + c \left(\int_{B_{R/2}} [\Psi_d]_{B_R}^- (|Dw|) dx \right)^{\frac{\alpha}{n}} \leq cR^\alpha + cR^{\frac{\alpha\delta_0}{1+\delta_0}} \left(\int_{B_{R/2}} [\Psi_d]_{B_R}^- (|Dw|)^{1+\delta_0} dx \right)^{\frac{\alpha}{n(1+\delta_0)}} \\
 &\leq c(\mathbf{data}_d(\Omega_0)) R^{\frac{\alpha\delta_0}{1+\delta_0}},
 \end{aligned} \tag{10.73}$$

where we have used the higher integrability estimates (5.62) of Theorem 5.5 under the double phase settings. Using estimates from the last display in (10.53) and recalling $R \leq 1$, we see

$$\left| \int_{B_{1/8}} \langle \bar{A}_0(D\bar{w}), D\varphi \rangle dx \right| \leq c(\mathbf{data}_d(\Omega_0)) Q_1(R) \|D\varphi\|_{L^\infty(B_{1/8})}, \tag{10.74}$$

where

$$Q_1(R) := \omega(R) + (1 + \omega(R)) R^{\frac{\alpha\delta_0}{(1+\delta_0)(1+s(H_a))}}. \tag{10.75}$$

Case 5: Assumption (10.13e) is in force. Using the assumption (1.13) and (10.21), I_a in (10.54) can be estimated as

$$\begin{aligned}
 I_a &\leq cR^\alpha \frac{H_a(E_d(R))}{G(E_d(R))} \\
 &\leq cR^\alpha \left(1 + \left[\left([\Psi_d]_{B_R}^- \right)^{-1} \left(\int_{B_{R/2}} [\Psi_d]_{B_R}^- \left(\left| \frac{w - (w)_{B_{R/2}}}{R} \right| \right) dx \right) \right]^\alpha \right) \\
 &\leq c \left(R^\alpha + \left[\text{osc}_{B_{2R}} u \right]^\alpha \right) \leq c(\mathbf{data}_d(\Omega_0)) R^{\gamma\alpha},
 \end{aligned} \tag{10.76}$$

where we have also used (5.33) and γ is the Hölder continuity exponent coming from Theorem 5.2 under the double phase settings. Inserting the estimate from the last display into (10.53) and recalling $R \leq 1$, we see

$$\left| \int_{B_{1/8}} \langle \bar{A}_0(D\bar{w}), D\varphi \rangle dx \right| \leq c(\mathbf{data}_d(\Omega_0)) Q_2(R) \|D\varphi\|_{L^\infty(B_{1/8})}, \tag{10.77}$$

where

$$Q_2(R) := \omega(R) + (1 + \omega(R)) R^{\frac{\alpha\gamma}{1+s(H_a)}}. \tag{10.78}$$

Collecting the estimates obtained in (10.59), (10.65), (10.71), (10.74) and (10.77), we conclude with

$$\left| \oint_{B_{1/8}} \langle \bar{A}_0(D\bar{w}), D\varphi \rangle dx \right| \leq c_h D(\varepsilon, R) \|D\varphi\|_{L^\infty(B_{1/8})} \quad (10.79)$$

for some constant $c_h \equiv c_h(\mathbf{data}_d(\Omega_0))$ for every $\varphi \in W_0^{1,\infty}(B_{1/8})$, where

$$D(\varepsilon, R) := \begin{cases} P_1(\varepsilon, R) & \text{if (10.13a) is assumed,} \\ P_2(\varepsilon, R) & \text{if (10.13b) is assumed,} \\ P_3(\varepsilon, R) & \text{if (10.13c) is assumed,} \\ Q_1(R) & \text{if (10.13d) is assumed,} \\ Q_2(R) & \text{if (10.13e) is assumed,} \end{cases} \quad (10.80)$$

in which P_1, P_2, P_3, Q_1 and Q_2 have been defined in (10.60), (10.66), (10.72), (10.75) and (10.78), respectively. By (10.45), (10.47a)-(10.47c) and (10.79), we are able to apply Lemma 6.1 with $A_0(z) \equiv \bar{A}_0(z)$, $\Psi_0(t) \equiv \bar{\Psi}_0(t)$ with $a_0 \equiv \bar{a}(\bar{x}_a)$ and $b_0 \equiv 0$. By Lemma 6.1, there exists $\bar{h} \in \bar{w} + W_0^{1,\bar{\Psi}_0}(B_{1/8})$ such that

$$\oint_{B_{1/8}} \langle \bar{A}_0(D\bar{h}), D\varphi \rangle dx = 0 \quad \text{for all} \quad \varphi \in W_0^{1,\infty}(B_{1/8}), \quad (10.81)$$

$$\oint_{B_{1/4}} \bar{\Psi}_0(|D\bar{h}|) dx + \oint_{B_{1/8}} [\bar{\Psi}_0(|D\bar{h}|)]^{1+\delta_1} dx \leq c \text{ for some } \delta_1 \leq \delta_0, \quad (10.82)$$

$$\oint_{B_{1/8}} (|V_{\bar{G}}(D\bar{w}) - V_{\bar{G}}(D\bar{h})|^2 + \bar{a}(\bar{x}_a)|V_{\bar{H}_a}(D\bar{w}) - V_{\bar{H}_a}(D\bar{h})|^2) dx \leq c[D(\varepsilon, R)]^{s_1} \quad (10.83)$$

and finally

$$\oint_{B_{1/8}} (\bar{G}(|\bar{w} - \bar{h}|) + \bar{a}(\bar{x}_a)\bar{H}_a(|\bar{w} - \bar{h}|)) dx \leq c_d[D(\varepsilon, R)]^{s_0} \quad (10.84)$$

with some constants $c, c_d \equiv c, c_d(\mathbf{data}_d(\Omega_0)) \geq 1$ and $s_0, s_1 \equiv s_0, s_1(\mathbf{data}_d) \in (0, 1)$, but they are all independent of R . The rest of the proof is similar as the argument after (7.98) of Lemma 7.3. \square

Lemma 10.4. *Under the assumptions of Lemma 10.3, let $w \in W^{1,\Psi}(B_{2R})$ be the solution to the problem defined in (10.17). If one of the assumptions (10.13a)-(10.13e) is satisfied, then there exists $h \in w + W_0^{1, [\Psi_d]_{\bar{B}_R}}(B_{R/8})$ being a local minimizer of the functional defined by*

$$W^{1,1}(B_{R/8}) \ni v \mapsto \mathcal{F}_0(v) := \int_{B_{R/8}} F_0(Dv) dx, \quad (10.85)$$

where the integrand function is given by

$$F_0(z) := F(x_a, (u)_{B_{2R}}, z) \quad (10.86)$$

for $x_a \in \bar{B}_R$ being a point such that $a(x_a) := a^-(B_R)$, whenever $z \in \mathbb{R}^n$, such that

$$\begin{aligned} & \oint_{B_{R/8}} [|V_G(Du) - V_G(Dh)|^2 + a(x_a)|V_{H_a}(Du) - V_{H_a}(Dh)|^2] dx \\ & \leq c(\omega(R^\gamma) + [D(\varepsilon, R)]^{s_1}) \oint_{B_{2R}} \Psi_d(x, |Du|) dx \end{aligned} \quad (10.87)$$

for some constant $c \equiv c(\mathbf{data}_d(\Omega_0))$, where s_1 and $D(\varepsilon, R)$ have been defined in (10.83) and (10.80), respectively. Moreover, we have the energy estimate

$$\int_{B_{R/8}} [\Psi_d]_{B_R}^- (|Dh|) dx \leq c \int_{B_{2R}} \Psi_d(x, |Du|) dx \quad (10.88)$$

for some constant $c \equiv c(n, \nu, L)$.

Proof. We need to revisit the proof of Lemma 10.3, specially Step 3 and Step 4. We consider a function $\bar{h} \in \bar{w} + W_0^{1, \bar{\Psi}_0}(B_{1/8})$ satisfying (10.81)-(10.84). Let h be the scaled back function of \bar{h} in $B_{R/8}$ as

$$h(x) := E_d(w, B_{R/2}) R \bar{h} \left(\frac{x - x_0}{R} \right) \quad \text{for every } x \in B_{R/8}(x_0). \quad (10.89)$$

Clearly, $h \in w + W_0^{1, [\Psi_d]_{B_R}^-}(B_{R/8})$ is a local minimizer of the functional \mathcal{F}_0 defined in (10.85) which means that

$$\mathcal{F}_0(h) = \int_{B_{R/8}} F_0(Dh) dx \leq \int_{B_{R/8}} F_0(Dh + D\varphi) dx \leq \mathcal{F}_0(h + \varphi) \quad (10.90)$$

holds for every $\varphi \in W_0^{1, [\Psi_d]_{B_R}^-}(B_{R/8})$. As shown in (7.9), we recall (10.19) to discover that

$$\begin{aligned} \int_{B_{R/8}} [\Psi_d]_{B_R}^- (|Dh|) dx &\leq \frac{L}{\nu} \int_{B_{R/8}} [\Psi_d]_{B_R}^- (|Dw|) dx \\ &\leq \frac{8^n L}{\nu} \int_{B_R} \Psi_d(x, |Dw|) dx \leq c(n, \nu, L) \int_{B_{2R}} \Psi_d(x, |Du|) dx, \end{aligned} \quad (10.91)$$

which proves (10.88). We write the inequality (10.83) in view of G, H_a, w and h in order to have

$$\begin{aligned} &\int_{B_{R/8}} [|V_G(Du) - V_G(Dh)|^2 + a(x_a) |V_{H_a}(Du) - V_{H_a}(Dh)|^2] dx \\ &\leq c[D(\varepsilon, R)]^{s_1} \int_{B_{R/2}} [\Psi_d]_{B_R}^- \left(\left| \frac{w - (w)_{B_{R/2}}}{R} \right| \right) dx \leq c[D(\varepsilon, R)]^{s_1} \int_{B_{R/2}} [\Psi_d]_{B_R}^- (|Dw|) dx \\ &\leq c[D(\varepsilon, R)]^{s_1} \int_{B_{R/2}} \Psi_d(x, |Du|) dx \end{aligned} \quad (10.92)$$

for some constant $c \equiv c(\mathbf{data}_d(\Omega_0))$, where we have applied the Sobolev-Poincaré inequality and (10.91). Combining this estimate together with (10.18) via some elementary computations, we directly reach (10.87). \square

We finally finish the present subsection with a crucial decay estimate on u .

Lemma 10.5. *Under the assumptions of Lemma 10.3, if one of the conditions (10.13a)-(10.13e) is satisfied, then for every $\varepsilon_* \in (0, 1)$, there exists a positive radius R_* with the dependence as*

$$R_* \equiv R_*(\mathbf{data}_d(\Omega_0), \varepsilon_*) \quad (10.93)$$

such that if $R \leq R_*$, then there exists a constant $c_G \equiv c_G(\mathbf{data}_d(\Omega_0))$ such that

$$\int_{B_{\tau R}} [\Psi_d]_{B_R}^- \left(\left| \frac{u - (u)_{B_{\tau R}}}{\tau R} \right| \right) dx \leq c_G \left(\tau^n + \tau^{-(s(\Psi_d)+1)\varepsilon_*} \right) \int_{B_{2R}} \Psi_d(x, |Du|) dx \quad (10.94)$$

holds for every $\tau \in (0, 1/32)$.

Proof. For the proof, we apply Lemma 10.3 with $\varepsilon^* \in (0, 1)$ to be determined in a few lines, and we can use (10.32) provided

$$R \leq R^* \equiv R^*(\mathbf{data}_d(\Omega_0), \varepsilon^*)$$

is found via (10.31). For every $\tau \in (0, 1/32)$ with some elementary manipulations, we see that

$$\begin{aligned} \int_{B_{\tau R}} [\Psi_d]_{B_R}^- \left(\left| \frac{u - (u)_{B_{\tau R}}}{\tau R} \right| \right) dx &\leq c \int_{B_{\tau R}} [\Psi_d]_{B_R}^- \left(\left| \frac{u - (w)_{B_{\tau R}}}{\tau R} \right| \right) dx \\ &\leq c \int_{B_{\tau R}} [\Psi_d]_{B_R}^- \left(\left| \frac{w - (w)_{B_{\tau R}}}{\tau R} \right| \right) dx + c\tau^{-(n+s(\Psi_d)+1)} \int_{B_R} [\Psi_d]_{B_R}^- \left(\left| \frac{u - w}{R} \right| \right) dx \\ &\leq c \left(1 + \tau^{-(n+s(\Psi_d)+1)} \varepsilon^* \right) \int_{B_{R/2}} [\Psi_d]_{B_R}^- \left(\left| \frac{w - (w)_{B_{R/2}}}{R} \right| \right) dx \\ &\quad + c\tau^{-(n+s(\Psi_d)+1)} \int_{B_R} [\Psi_d]_{B_R}^- \left(\left| \frac{u - w}{R} \right| \right) dx \\ &\leq c \left(1 + \tau^{-(n+s(\Psi_d)+1)} \varepsilon^* \right) \int_{B_R} [\Psi_d]_{B_R}^- (|Dw|) dx + c\tau^{-(n+s(\Psi_d)+1)} \int_{B_R} [\Psi_d]_{B_R}^- \left(\left| \frac{u - w}{R} \right| \right) dx \end{aligned} \quad (10.95)$$

with some constant $c \equiv c(\mathbf{data}_d(\Omega_0))$, where throughout the last display we repeatedly used (2.6) and (7.36). The last display and (10.22) along with some elementary manipulations yield

$$\int_{B_{\tau R}} [\Psi_d]_{B_R}^- \left(\left| \frac{u - (u)_{B_{\tau R}}}{\tau R} \right| \right) dx \leq c \left(\tau^n + \tau^{-(s(\Psi_d)+1)} \varepsilon^* + \tau^{-(s(\Psi_d)+1)} [\omega(R^\gamma)]^{\frac{1}{2}} \right) \int_{B_{2R}} \Psi_d(x, |Du|) dx$$

for every $\tau \in (0, 1/16)$ and some $c \equiv c(\mathbf{data}_d(\Omega_0))$. Then we choose $\varepsilon^* \equiv \varepsilon_*/2$ and $R_* \leq R^*$ in such a way that $[\omega(R_*^\gamma)]^{\frac{1}{2}} \leq \varepsilon_*/2$. This choice gives us the dependence as described in (10.93) and yields (10.94). \square

We have now discovered all the necessary tools. They are Lemma 10.2, Lemma 10.3 and Lemma 10.5 in the double phase settings for proving Theorem 10.1 and Theorem 10.2. Applying those lemmas with arguing in a similar manner as in the proofs of Theorem 1.2 and Theorem 1.1, we are able to prove Theorem 10.1 and Theorem 10.2. For the sake of the completeness, we provide a sketch of the proofs.

Proof of Theorem 10.2. The proof of Theorem 10.2 can be done similarly as for the proof of Theorem 1.2. We just combine Lemma 5.3 under the double phase settings and Lemma 10.5, as we already have done in (8.1)-(8.27).

Lemma 10.6. *Under the assumptions and notations of Lemma 10.3 and Lemma 10.4, let $w \in W^{1, \Psi_d}(B_R)$ be the function defined in (10.17). Suppose that (10.13c) is satisfied for $\omega_a(t) = t^\alpha$ with some $\alpha \in (0, 1]$.*

Then there exists a function $h \in w + W_0^{1, [\Psi_d]_{B_R}^-}(B_{R/8})$ being a local minimizer of the functional \mathcal{F}_0 defined in (10.85) such that

$$\begin{aligned} \int_{B_{R/8}} [|V_G(Du) - V_G(Dh)|^2 + a(x_a) |V_{H_a}(Du) - V_{H_a}(Dh)|^2] dx \\ \leq c \left(\omega(R^\gamma) + \left[\omega(R) + (1 + \omega(R)) R^{\frac{\alpha}{2(1+s(H_a))}} \right]^{s_1} \right) \int_{B_{2R}} \Psi_d(x, |Du|) dx \end{aligned} \quad (10.96)$$

for some constant $c \equiv c(\mathbf{data}_d(\Omega_0))$ and $s_1 \equiv s_1(\mathbf{data}_d)$, respectively. Moreover, the energy estimate

$$\int_{B_{R/8}} [\Psi_d]_{B_R}^- (|Dh|) dx \leq c \int_{B_{2R}} \Psi_d(x, |Du|) dx \quad (10.97)$$

holds for some constant $c \equiv c(n, \nu, L)$.

Proof. First we apply Theorem 10.2 in order to obtain that, for every $\theta \in (0, 1)$ and every open subset $\Omega_0 \Subset \Omega$, there exists a constant $c \equiv c(\mathbf{data}_d(\Omega_0), \theta)$ such that

$$[u]_{0, \theta; \Omega_0} \leq c(\mathbf{data}_d(\Omega_0), \theta). \quad (10.98)$$

In particular, we choose $\theta \equiv (\gamma + 1)/2$. By revisiting the proof of Lemma 10.3, we shall estimate the term I_a introduced in (10.54). Using (1.14) and (10.21), we have

$$\begin{aligned} I_a &\leq cR^\alpha \left(1 + \left[\left([\Psi_d]_{B_R}^- \right)^{-1} \left(\int_{B_{R/2}} [\Psi_d]_{B_R}^- \left(\left| \frac{w - (w)_{B_{R/2}}}{R} \right| \right) dx \right) \right]^{\frac{\alpha}{1-\gamma}} \right) \\ &\leq c \left(R^\alpha + R^{-\frac{\alpha\gamma}{1-\gamma}} \left[\text{osc}_{B_{2R}} u \right]^{\frac{\alpha}{1-\gamma}} \right) \leq c(\mathbf{data}_d(\Omega_0)) R^{\alpha/2}, \end{aligned} \quad (10.99)$$

where we have used (10.98) with the choice of $\theta \equiv (1 + \gamma)/2$ and $B_{2R} \subset \Omega_0$ with $R \leq 1$. Plugging this estimate in (10.53), we find

$$\left| \int_{B_{1/8}} \langle \bar{A}_0(D\bar{w}), D\varphi \rangle dx \right| \leq c(\mathbf{data}_d(\Omega_0)) Q_3(R) \|D\varphi\|_{L^\infty(B_{1/8})}, \quad (10.100)$$

where

$$Q_3(R) := \omega(R) + (1 + \omega(R)) R^{\frac{\alpha}{2(1+s(H_a))}}, \quad (10.101)$$

where the vector field \bar{A}_0 has been defined in (10.43). We consider a function $\bar{h} \in \bar{w} + W_0^{1, \bar{\Psi}_0}(B_{1/8})$ satisfying (10.81)-(10.84) with the term $D(\varepsilon, R)$ replaced by $Q_3(R)$ defined above. Let h be the scaled back function of \bar{h} in $B_{R/8}$ as

$$h(x) := E_d(w, B_{R/2}) R \bar{h} \left(\frac{x - x_0}{R} \right) \quad \text{for every } x \in B_{R/8}(x_0). \quad (10.102)$$

Clearly, $h \in w + W_0^{1, [\Psi_d]_{B_R}^-}(B_{R/8})$ is a local minimizer of the functional \mathcal{F}_0 defined in (10.85) which means that

$$\mathcal{F}_0(h) = \int_{B_{R/8}} F_0(Dh) dx \leq \int_{B_{R/8}} F_0(Dh + D\varphi) dx \leq \mathcal{F}_0(h + \varphi) \quad (10.103)$$

holds for every $\varphi \in W_0^{1, [\Psi_d]_{B_R}^-}(B_{R/8})$. Arguing similarly as in the proof of Lemma 10.2 together with recalling (10.19), we see

$$\begin{aligned} \int_{B_{R/8}} [\Psi_d]_{B_R}^- (|Dh|) dx &\leq \frac{L}{\nu} \int_{B_{R/8}} [\Psi_d]_{B_R}^- (|Dw|) dx \leq \frac{8^n L}{\nu} \int_{B_R} \Psi_d(x, |Dw|) dx \\ &\leq c(n, \nu, L) \int_{B_{2R}} \Psi_d(x, |Du|) dx, \end{aligned} \quad (10.104)$$

which proves (10.97). We write the inequality (10.83) in view of G, H_a, w and h in order to have

$$\begin{aligned} &\int_{B_{R/8}} [|V_G(Dw) - V_G(Dh)|^2 + a(x_a) |V_{H_a}(Dw) - V_{H_a}(Dh)|^2] dx \\ &\leq c[Q_3(R)]^{s_1} \int_{B_{R/2}} [\Psi_d]_{B_R}^- \left(\left| \frac{w - (w)_{B_{R/2}}}{R} \right| \right) dx \leq c[Q_3(R)]^{s_1} \int_{B_{R/2}} [\Psi_d]_{B_R}^- (|Dw|) dx \\ &\leq c[Q_3(R)]^{s_1} \int_{B_{2R}} \Psi_d(x, |Du|) dx \end{aligned} \quad (10.105)$$

for some constant $c \equiv c(\mathbf{data}_d(\Omega_0))$, where we have applied the Sobolev-Poincaré inequality and (10.91). Combining this estimate together with (10.18) via some elementary computations implies (10.96). \square

Proof of Theorem 10.1. It follows from Theorem 10.2 and a standard covering argument that, for every open subset $\Omega_0 \Subset \Omega$ and any number $k > 0$, there exists a constant $c \equiv c(\mathbf{data}_d(\Omega_0), k)$ such that

$$\oint_{B_{2R}} \Psi_d(x, |Du|) dx \leq cR^{-k} \quad (10.106)$$

for every $B_{2R} \subset \Omega_0$ with $R \leq 1$. Now we fix an open subset $\Omega_0 \Subset \Omega$ and a ball $B_{2R} \equiv B_{2R}(x_0) \subset \Omega_0$ with $R \leq 1$. Then applying Lemma 10.4 and Lemma 10.6,

$$\begin{aligned} & \oint_{B_{R/8}} (|V_G(Du) - V_G(Dh)|^2 + a(x_a)|V_{H_a}(Du) - V_{H_a}(Dh)|^2) dx \\ & \leq c(R^{\mu\gamma} + [Q(R)]^{s_1}) \oint_{B_{2R}} \Psi_d(x, |Du|) dx \end{aligned} \quad (10.107)$$

for some constant $c \equiv c(\mathbf{data}_d(\Omega_0))$ and $s_1 \equiv s_1(\mathbf{data}_d)$, where

$$Q(R) := \begin{cases} R^\mu + (1 + R^\mu)R^{\frac{\alpha\delta_0}{(1+\delta_0)(1+s(H_a))}} & \text{if (10.12a) is assumed,} \\ R^\mu + (1 + R^\mu)R^{\frac{\alpha\gamma}{1+s(H_a)}} & \text{if (10.12b) is assumed,} \\ R^\mu + (1 + R^\mu)R^{\frac{\alpha}{2(1+s(H_a))}} & \text{if (10.12c) is assumed,} \end{cases} \quad (10.108)$$

in which γ is the Hölder continuity exponent determined via Theorem 5.2 under the double phase settings and δ_0 is the higher integrability exponent coming from Lemma 10.2. Denoting by

$$d \equiv d(\mathbf{data}_d(\Omega_0)) := \begin{cases} \min \left\{ \mu\gamma, s_1\mu, \frac{\alpha\delta_0 s_1}{(1+\delta_0)(1+s(H_a))} \right\} & \text{if (10.12a) is assumed,} \\ \min \left\{ \mu\gamma, s_1\mu, \frac{\alpha\gamma s_1}{1+s(H_a)} \right\} & \text{if (10.12b) is assumed,} \\ \min \left\{ \mu\gamma, s_1\mu, \frac{\alpha s_1}{2(1+s(H_a))} \right\} & \text{if (10.12c) is assumed,} \end{cases} \quad (10.109)$$

and choosing $k \equiv d/4$ in (10.106), the inequality (10.107) can be written as

$$\oint_{B_{R/8}} (|V_G(Du) - V_G(Dh)|^2 + a(x_a)|V_{H_a}(Du) - V_{H_a}(Dh)|^2) dx \leq cR^{3d/4} \quad (10.110)$$

for some constant $c \equiv c(\mathbf{data}_d(\Omega_0))$, where we again recall that the function h has been defined via Lemma 10.4 and Lemma 10.6. Recalling (10.88) and (10.97), we have the energy estimate

$$\oint_{B_{R/8}} [\Psi_d]_{B_R}^- (|Dh|) dx \leq c \oint_{B_{2R}} \Psi_d(x, |Du|) dx \quad (10.111)$$

with a constant $c \equiv c(n, \nu, L)$. Once we arrive at this stage, the rest of the proof is in the same way as argued in the proof of Theorem 1.1. The proof is complete.

11. REGULARITY RESULTS UNDER ADDITIONAL INTEGRABILITY

We turn our attention to studying properties of a local Q -minimizer of the functional \mathcal{P} defined in (1.1) under some additional Lebesgue integrability assumption. We shall consider a local Q -minimizer u of the functional \mathcal{P} under the following assumptions:

$$\begin{cases} u \in W^{1,\Psi}(\Omega) \cap L^\kappa(\Omega) \quad (\kappa \geq 1) \\ \lambda_4(\kappa) := \sup_{t>0} \Lambda \left(t^{\frac{\kappa}{n+\kappa}}, \frac{1}{t} \right) < \infty, \end{cases} \quad (11.1)$$

where the function $\Lambda : (0, \infty) \times (0, \infty) \rightarrow (0, \infty)$ has been defined in (1.11) together with $\omega_a, \omega_b : [0, \infty) \rightarrow [0, \infty)$ being concave functions vanishing at the origin such that $a(\cdot) \in C^{0,\omega_a}(\Omega)$ and $b(\cdot) \in C^{0,\omega_b}(\Omega)$. To see the meaning of the assumption (11.1)₂, let us consider the standard double phase that $G(t) = t^p$, $H_a(t) = t^q$

and $\omega_a(t) = t^\alpha$, $\omega_b \equiv 0$ for $1 < p \leq q$ and $\alpha \in (0, 1]$. Under these standard double phase settings, the assumption (11.1)₂ is equivalent to the following one:

$$q \leq p + \frac{\alpha\kappa}{n + \kappa}. \quad (11.2)$$

A local Q -minimizer $u \in W^{1,\Psi}(\Omega)$ implies that $u \in W^{1,p}(\Omega)$. It is clearly interesting point that $p < n$, otherwise we can prove $u \in L_{\text{loc}}^\infty(\Omega)$ by using Morrey-Embedding properties for $p > n$ and using a higher integrability for $p = n$. Then, for $1 < p < n$, applying Sobolev embedding properties, one can see that $u \in L_{\text{loc}}^{\frac{np}{n-p}}(\Omega)$. Choosing $\kappa \equiv \frac{np}{n-p}$, the condition (11.2) is equivalent to the following one

$$q \leq p + \frac{\alpha p}{n},$$

which generates the same condition as (1.6a), as we have discussed in the introduction part. Now if $\kappa > \frac{np}{n-p}$, then we would have

$$q \leq p + \frac{\alpha p}{n} < p + \frac{\alpha\kappa}{n + \kappa},$$

which tells us the possible range of q is larger than the one in (1.6a). Considering a local Q -minimizers of the functional \mathcal{P} under the assumption (11.1), we shall show that $u \in L_{\text{loc}}^\infty(\Omega)$. To do this, we start by proving a Sobolev-Poincaré inequality under the assumption (11.1)₂.

Theorem 11.1. *Let $v \in W^{1,\Psi}(B_R) \cap L^\kappa(B_R)$ for a ball $B_R \subset \Omega$ with $R \leq 1$ under the assumption (11.1)₂. Then, for any $d \in \left[1, \frac{n(n+\kappa)}{n(n+\kappa)-\kappa}\right)$, there exist constants $\theta \equiv \theta(n, s(G), s(H_a), s(H_b), \kappa, d) \in (0, 1)$ and $c \equiv c(n, s(G), s(H_a), s(H_b), \omega_a(1), \omega_b(1), \kappa, d)$ such that the following Sobolev-Poincaré-type inequality holds:*

$$\left[\int_{B_R} \left[\Psi \left(x, \left| \frac{v - (v)_{B_R}}{R} \right| \right) \right]^d dx \right]^{\frac{1}{d}} \leq c \lambda_{sp} \left[\int_{B_R} [\Psi(x, |Dv|)]^\theta dx \right]^{\frac{1}{\theta}}, \quad (11.3)$$

where

$$\lambda_{sp} = 1 + ([a]_{\omega_a} + [b]_{\omega_b}) \left(\lambda_4(\kappa) + \lambda_4(\kappa) \left(\int_{B_R} |v|^\kappa dx \right)^{\frac{1}{n+\kappa}} \right) \quad (11.4)$$

Moreover, the above estimate (11.3) is still valid with $v - (v)_{B_R}$ replaced by v if $v \in W_0^{1,\Psi}(B_R) \cap L^\kappa(B_R)$.

Proof. Note that the above theorem covers [75, Theorem 3.1], which is a special case when $G(t) = t^p$, $H(t) = t^q$, $\omega_a(t) = t^\alpha$ and $\omega_b(\cdot) \equiv 0$ for some constants $1 < p \leq q$ and $\alpha \in (0, 1]$. Also our proof is much more elementary comparing with the approach used there. Using the continuity of the coefficient functions $a(\cdot)$ and $b(\cdot)$ and arguing in the same way as in (4.6), we find

$$\begin{aligned} I &:= \left(\int_{B_R} \left[\Psi \left(x, \left| \frac{v - (v)_{B_R}}{R} \right| \right) \right]^d dx \right)^{\frac{1}{d}} \leq 6[a]_{\omega_a} \omega_a(R) \left(\int_{B_R} \left[H_a \left(\left| \frac{v - (v)_{B_R}}{R} \right| \right) \right]^d dx \right)^{\frac{1}{d}} \\ &\quad + 6[b]_{\omega_b} \omega_b(R) \left(\int_{B_R} \left[H_b \left(\left| \frac{v - (v)_{B_R}}{R} \right| \right) \right]^d dx \right)^{\frac{1}{d}} \\ &\quad + 3 \left(\int_{B_R} \left[\Psi_{B_R}^- \left(\left| \frac{v - (v)_{B_R}}{R} \right| \right) \right]^d dx \right)^{\frac{1}{d}} \\ &=: 6[a]_{\omega_a} I_1 + 6[b]_{\omega_b} I_2 + 3I_3. \end{aligned} \quad (11.5)$$

We now shall deal with estimating the terms I_i with $i \in 1, 2, 3$ in (11.5) using the additional a priori assumption $u \in L^\kappa(B_R)$ under (11.1)₂. In turn, using (2.2) and the assumption (11.1)₂, we see

$$\begin{aligned}
I_1 &= \omega_a(R) \left(\int_{B_R} \left[\frac{H_a \left(\left| \frac{v-(v)_{B_R}}{R} \right| \right)}{G \left(\left| \frac{v-(v)_{B_R}}{R} \right| \right)} G \left(\left| \frac{v-(v)_{B_R}}{R} \right| \right) \right]^d dx \right)^{\frac{1}{d}} \\
&\leq \lambda_4(\kappa) \omega_a(R) \left(\int_{B_R} \left[\left(1 + \left[\omega_a \left(\left(\left| \frac{v-(v)_{B_R}}{R} \right| \right)^{-\frac{\kappa}{n+\kappa}} \right] \right)^{-1} G \left(\left| \frac{v-(v)_{B_R}}{R} \right| \right) \right]^d dx \right)^{\frac{1}{d}} \\
&\leq \lambda_4(\kappa) \omega_a(R) \left(\int_{B_R} \left[\left(1 + \left[\frac{1}{\omega_a(R)} + \frac{R}{\omega_a(R)} \left| \frac{v-(v)_{B_R}}{R} \right|^{\frac{\kappa}{n+\kappa}} \right] G \left(\left| \frac{v-(v)_{B_R}}{R} \right| \right) \right]^d dx \right)^{\frac{1}{d}} \\
&\leq c_* \lambda_4(\kappa) \left(\int_{B_R} \left[G \left(\left| \frac{v-(v)_{B_R}}{R} \right| \right) \right]^d dx \right)^{\frac{1}{d}} \\
&\quad + c_* \lambda_4(\kappa) R^{\frac{n}{n+\kappa}} \left(\int_{B_R} |v-(v)_{B_R}|^{\frac{d\kappa}{n+\kappa}} \left[G \left(\left| \frac{v-(v)_{B_R}}{R} \right| \right) \right]^d dx \right)^{\frac{1}{d}}
\end{aligned}$$

for the constant $c_* = 2(1 + \omega_a(1))$. Using Hölder's inequality with conjugate exponents $\left(\frac{n+\kappa}{d}, \frac{n+\kappa}{n+\kappa-d} \right)$, we have

$$\begin{aligned}
&R^{\frac{n}{n+\kappa}} \left(\int_{B_R} |v-(v)_{B_R}|^{\frac{d\kappa}{n+\kappa}} \left[G \left(\left| \frac{v-(v)_{B_R}}{R} \right| \right) \right]^d dx \right)^{\frac{1}{d}} \\
&\leq R^{\frac{n}{n+\kappa}} \left(\int_{B_R} |v-(v)_{B_R}|^\kappa dx \right)^{\frac{1}{n+\kappa}} \left(\int_{B_R} \left[G \left(\left| \frac{v-(v)_{B_R}}{R} \right| \right) \right]^{\frac{(n+\kappa)d}{n+\kappa-d}} dx \right)^{\frac{n+\kappa-d}{(n+\kappa)d}} \\
&\leq c \left(\int_{B_R} |v|^\kappa dx \right)^{\frac{1}{n+\kappa}} \left(\int_{B_R} \left[G \left(\left| \frac{v-(v)_{B_R}}{R} \right| \right) \right]^{\frac{(n+\kappa)d}{n+\kappa-d}} dx \right)^{\frac{n+\kappa-d}{(n+\kappa)d}}
\end{aligned}$$

for some constant $c \equiv c(n)$. Combining the last two displays and arguing similarly for I_2 , we discover

$$\begin{aligned}
I_1 + I_2 &\leq c \lambda_4(\kappa) \left(\int_{B_R} \left[G \left(\left| \frac{v-(v)_{B_R}}{R} \right| \right) \right]^d dx \right)^{\frac{1}{d}} \\
&\quad + c \lambda_4(\kappa) \left(\int_{B_R} |v|^\kappa dx \right)^{\frac{1}{n+\kappa}} \left(\int_{B_R} \left[G \left(\left| \frac{v-(v)_{B_R}}{R} \right| \right) \right]^{\frac{(n+\kappa)d}{n+\kappa-d}} dx \right)^{\frac{n+\kappa-d}{(n+\kappa)d}}
\end{aligned}$$

for some constant $c \equiv c(n, \omega_a(1), \omega_b(1))$. Now we apply Lemma 4.1 to $\Phi \equiv G$ with $d_0 \equiv d$ and $d_0 \equiv \frac{n+\kappa-d}{(n+\kappa)d}$ in order to have an exponent $\theta_1 \equiv \theta_1(n, s(G), \kappa, d) \in (0, 1)$ such that

$$I_1 + I_2 \leq c \left(\lambda_4(\kappa) + \lambda_4(\kappa) \left(\int_{B_R} |v|^\kappa dx \right)^{\frac{1}{n+\kappa}} \right) \left(\int_{B_R} [G(|Dv|)]^{\theta_1} dx \right)^{\frac{1}{\theta_1}} \quad (11.6)$$

holds for some constant $c \equiv c(n, s(G), \omega_a(1), \omega_b(1), \kappa, d)$. On the other hand, since $\Psi_{B_R}^- \in \mathcal{N}$ with an index $s(\Psi) = s(G) + s(H_a) + s(H_b)$ by Remark 2.2, we are able to apply Lemma 4.1 with $\Phi \equiv \Psi_{B_R}^-$ for $d_0 \equiv d$. In

turn, there exists $\theta_2 \equiv \theta_2(n, s(\Psi), d)$ such that

$$I_3 \leq c \left[\int_{B_R} [\Psi_{B_R}^- (|Dv|)]^{\theta_2} dx \right]^{\frac{1}{\theta_2}} \quad (11.7)$$

with some constant $c \equiv c(n, s(\Psi), d)$. Taking into account the estimates obtained in (11.6)-(11.7) into (11.5), recalling the very definition of $\Psi_{B_R}^-$ in (2.3) and setting $\theta := \max\{\theta_1, \theta_2\}$, we arrive at (11.3). The proof is finished. \square

Remark 11.1. We here remark that choosing $d \equiv 1$ in a Sobolev-Poincaré type inequality of Theorem 11.1, we see that there exists an exponent $\theta \equiv \theta(n, s(G), s(H_a), s(H_b), \kappa)$ such that

$$\int_{B_R} \Psi \left(x, \left| \frac{v - (v)_{B_R}}{R} \right| \right) dx \leq c \lambda_{sp} \left[\int_{B_R} [\Psi(x, |Dv|)]^\theta dx \right]^{\frac{1}{\theta}} \quad (11.8)$$

holds for some constant $c \equiv c(n, s(G), s(H_a), s(H_b), \omega_a(1), \omega_b(1), \kappa)$, where λ_{sp} is the one same as in (11.4).

Remark 11.2. With $u \in W^{1,\Psi}(\Omega)$ being a local Q -minimizer of the functional \mathcal{P} , we here point out that it is also possible to suppose a priori $u \in W^\Phi(\Omega)$ for some Young function Φ . In this case, discovering a relevant assumption like (11.1)₂ would be an interesting point to find how it is connected to Embedding properties in Orlicz-Sobolev spaces [26, 27, 28] likewise we have discussed above in Lebesgue settings. Moreover, proving various regularity results under a new relevant condition may generate a different phenomenon even for a Lavrentiev gap. We can also a priori assume that local Q -minimizers belong to certain Campanato, BMO, VMO, or some other spaces. Under all those a priori assumptions, it should be necessary to discover out the relevant optimal conditions under which various regularity results are obtainable.

For a local Q -minimizer u of the functional \mathcal{P} under the assumption (11.1), the data of the problem is understood by the following set of parameters:

$$\mathbf{data}_i \equiv \{n, \lambda_4(\kappa), \kappa, s(G), s(H_a), s(H_b), \omega_a(1), \omega_b(1), \|u\|_{L^\kappa(\Omega)}, Q\}. \quad (11.9)$$

As usual, for any open subset $\Omega_0 \Subset \Omega$, we denote by $\mathbf{data}_i(\Omega_0)$ the set of parameters defined above together with $\text{dist}(\Omega_0, \partial\Omega)$. Now we focus on showing local boundedness estimates of a local Q -minimizer u of the functional \mathcal{P} in (1.1) under the assumption (11.1).

Theorem 11.2. *Let $u \in W^{1,\Psi}(\Omega)$ be a local Q -minimizer of the functional \mathcal{P} in (1.1) under the assumption (11.1). Then there exists a constant $c \equiv c(\mathbf{data}_i)$ such that*

$$\left\| \Psi_{B_R}^- \left(\left| \frac{(u - (u)_{B_R})_\pm}{R} \right| \right) \right\|_{L^\infty(B_{R/2})} \leq c \int_{B_R} \Psi \left(x, \left| \frac{(u - (u)_{B_R})_\pm}{R} \right| \right) dx \quad (11.10)$$

and

$$\Psi_{B_R}^- \left(\left| \frac{u(x_1) - u(x_2)}{R} \right| \right) \leq c \int_{B_R} \Psi(x, |Du|) dx \quad \text{for a.e. } x_1, x_2 \in B_{R/2}, \quad (11.11)$$

whenever $B_R \equiv B_R(x_0) \subset \Omega$ is a ball with $R \leq 1$. In particular, $u \in L_{\text{loc}}^\infty(\Omega)$.

Proof. The meaning of \mathbf{data}_i under the assumption (11.1), already has been introduced in (11.9). As in the proof of Theorem 5.1, we consider the following scaled functions as:

$$\begin{aligned} \bar{u}(x) &:= \frac{u(x_0 + Rx) - (u)_{B_R}}{R}, \quad \bar{a}(x) := a(x_0 + Rx), \quad \bar{b}(x) := b(x_0 + Rx), \\ \bar{\Psi}(x, t) &:= G(t) + \bar{a}(x)H(t) + \bar{b}(x)H(t), \\ \bar{A}(k, s) &:= B_s(0) \cap \{\bar{u} > k\} \quad \text{and} \quad \bar{B}(k, s) := B_s(0) \cap \{\bar{u} < k\} \end{aligned} \quad (11.12)$$

for every $x \in B_1(0)$, $t \geq 0$, $s \in (0, 1)$ and $k \in \mathbb{R}$. The remaining part of the proof consists of 3 steps as in the proof of Theorem 5.1.

Step 1: Sobolev-Poincaré under the scaling in (11.12). In this step, we prove that there exists a positive exponent $\theta \equiv \theta(n, s(G), s(H_a), s(H_b), \kappa) \in (0, 1)$ such that

$$\int_{B_1} \bar{\Psi}(x, |f|) dx \leq c \bar{\kappa}_{sp} \left(\int_{B_1} [\bar{\Psi}(x, |Df|)]^\theta dx \right)^{\frac{1}{\theta}} \quad (11.13)$$

for some constant $c \equiv c(n, s(G), s(H_a), s(H_b), \omega_a(1), \omega_b(1), \kappa)$, whenever $f \in W_0^{1, \bar{\Psi}}(B_1) \cap L^\kappa(B_1)$, where

$$\bar{\kappa}_{sp} = 1 + ([a]_{\omega_a} + [b]_{\omega_b}) \left(\lambda_1 + \lambda_1 R \left(\int_{B_1} |f|^\kappa dx \right)^{\frac{1}{n+\kappa}} \right).$$

Using the continuity properties of $\bar{a}(\cdot)$ and $\bar{b}(\cdot)$, we see

$$\begin{aligned} I &:= \int_{B_1} \bar{\Psi}(x, |f|) dx \\ &\leq 2[a]_{\omega_a} \omega_a(R) \int_{B_1} H_a(|f|) dx + 2[b]_{\omega_b} \omega_b(R) \int_{B_1} H_b(|f|) dx + \int_{B_1} \bar{\Psi}_{B_1}^-(|f|) dx \\ &=: 2[a]_{\omega_a} I_1 + 2[b]_{\omega_b} I_2 + I_3, \end{aligned}$$

where

$$\bar{\Psi}_{B_1}^-(t) := G(t) + \inf_{x \in B_1} \bar{a}(x) H_a(t) + \inf_{x \in B_1} \bar{b}(x) H_b(t) \quad \text{for every } t \geq 0.$$

Now we estimate the terms I_i for $i \in \{1, 2, 3\}$ similarly as in the proof of Theorem 11.1. In turn, using the assumption (11.1)₂ and (2.2), we have

$$\begin{aligned} I_1 &= \omega_a(R) \int_{B_1} \frac{H_a(|f|)}{G(|f|)} G(|f|) dx \\ &\leq \lambda_4(\kappa) \omega_a(R) \int_{B_1} \left(1 + \left[\omega_a \left(|f|^{-\frac{\kappa}{n+\kappa}} \right) \right]^{-1} \right) G(|f|) dx \\ &\leq \lambda_4(\kappa) \omega_a(R) \int_{B_1} \left(1 + \left[\frac{1}{\omega_a(R)} + \frac{R}{\omega_a(R)} |f|^{\frac{\kappa}{n+\kappa}} \right] \right) G(|f|) dx \\ &\leq \lambda_4(\kappa) (1 + \omega_a(1)) \int_{B_1} G(|f|) dx + 2\lambda_4(\kappa) R \int_{B_1} |f|^{\frac{\kappa}{n+\kappa}} G(|f|) dx. \end{aligned}$$

Arguing in the same way, we have

$$I_2 \leq \lambda_4(\kappa) (1 + \omega_b(1)) \int_{B_1} G(|f|) dx + 2\lambda_4(\kappa) R \int_{B_1} |f|^{\frac{\kappa}{n+\kappa}} G(|f|) dx.$$

Then the inequality (11.13) follows from the arguments used in the proof of Theorem 11.1 and Lemma 4.1.

Step 2. Proof of (11.10). Since $u - (u)_{B_R}$ is a local Q -minimizer of the functional \mathcal{P} in (1.1), using a Caccioppoli inequality of Lemma 5.1, one can see that

$$\int_{B_t} \bar{\Psi}(x, |D(\bar{u} - k)_\pm|) dx \leq c \int_{B_s} \bar{\Psi} \left(x, \frac{(\bar{u} - k)_\pm}{s - t} \right) dx \quad (11.14)$$

holds for some constant $c \equiv c(s(G), s(H_a), s(H_b), Q)$, whenever $0 < t < s \leq 1$ and $k \in \mathbb{R}$. Let us now consider the concentric balls $B_\rho \Subset B_t \Subset B_s$ with $1/2 \leq \rho < s \leq 1$ and $t := (\rho + s)/2$. Let $\eta \in C_0^\infty(B_t)$ be a

standard cut-off function such that $\chi_{B_\rho} \leq \eta \leq \chi_{B_t}$ and $|D\eta| \leq \frac{2}{t-\rho} = \frac{4}{s-\rho}$. Now we apply inequality (11.13) from Step 1 above in order to have a positive exponent $\theta \equiv \theta(n, s(G), s(H_a), s(H_b), \kappa)$ such that

$$\int_{\bar{A}(k, \rho)} \bar{\Psi}(x, \bar{u} - k) dx \leq \int_{\bar{B}_1} \bar{\Psi}(x, \eta(\bar{u} - k)_+) dx \leq c \bar{\kappa}_{sp} \left(\int_{\bar{B}_1} [\bar{\Psi}(x, |D(\eta(\bar{u} - k)_+)|)]^\theta dx \right)^{\frac{1}{\theta}}$$

for some constant $c \equiv c(n, s(G), s(H_a), s(H_b), \omega_a(1), \omega_b(1), \kappa)$, where

$$\bar{\kappa}_{sp} = 1 + ([a]_{\omega_a} + [b]_{\omega_b}) \left(\lambda_4(\kappa) + \lambda_4(\kappa) R \left(\int_{\bar{B}_1} [\eta(\bar{u} - k)_+]^\kappa dx \right)^{\frac{1}{n+\kappa}} \right).$$

By recalling the definition of \bar{u} in (11.12), we have

$$\bar{\kappa}_{sp} \leq c \left[1 + R \left(\int_{\bar{B}_R} \left| \frac{u - (u)_{B_R}}{R} \right|^\kappa dx \right)^{\frac{1}{n+\kappa}} \right] \leq c \left[1 + \left(\int_{\bar{B}_R} |u|^\kappa dx \right)^{\frac{1}{n+\kappa}} \right]$$

with a constant $c \equiv c(n, \lambda_4(\kappa), [a]_{\omega_a} + [b]_{\omega_b})$. Once we arrive at this stage the rest of the proof can be proceed in the same way as in the proof of Theorem 5.1. \square

Theorem 11.3. *Let $u \in W^{1, \Psi}(\Omega)$ be a local Q -minimizer of the functional \mathcal{P} defined in (1.1) under the coefficient functions $a(\cdot) \in C^{\omega_a}(\Omega)$ and $b(\cdot) \in C^{\omega_b}(\Omega)$ for ω_a, ω_b being non-negative concave functions vanishing at the origin. If the assumption (11.1) is satisfied, then for every open subset $\Omega_0 \Subset \Omega$, there exists a Hölder continuity exponent $\gamma \equiv \gamma(\mathbf{data}_i(\Omega_0)) \in (0, 1)$ such that*

$$\|u\|_{L^\infty(\Omega_0)} + [u]_{0, \gamma; \Omega_0} \leq c(\mathbf{data}_i(\Omega_0)) \quad (11.15)$$

and the oscillation estimate

$$\text{osc}_{B_\rho} u \leq c \left(\frac{\rho}{R} \right)^\gamma \text{osc}_{B_R} u \quad (11.16)$$

holds for some $c \equiv c(\mathbf{data}_i(\Omega_0))$ and all concentric balls $B_\rho \Subset B_R \Subset \Omega_0 \Subset \Omega$ with $R \leq 1$.

Proof. First let us observe that, for every $t \geq 1$, we have

$$\begin{aligned} \frac{\omega_a(t)}{1 + \omega_a(t)} \frac{1 + \omega_a(t^{\frac{\kappa}{n+\kappa}})}{\omega_a(t^{\frac{\kappa}{n+\kappa}})} &\leq 1 + \frac{\omega_a(t)}{\omega_a(t^{\frac{\kappa}{n+\kappa}}) + \omega_a(t)\omega_a(t^{\frac{\kappa}{n+\kappa}})} \\ &\leq 1 + \frac{1}{\omega_a(t^{\frac{\kappa}{n+\kappa}})} \leq 1 + \frac{1}{\omega_a(1)}. \end{aligned}$$

This same inequality holds true also for ω_b . Therefore, for every $t \geq 1$, we see that

$$\begin{aligned} \Lambda \left(t, \frac{1}{t} \right) &\leq \lambda_4(\kappa) \left(\frac{\omega_a(t)}{1 + \omega_a(t)} \frac{1 + \omega_a(t^{\frac{\kappa}{n+\kappa}})}{\omega_a(t^{\frac{\kappa}{n+\kappa}})} + \frac{\omega_b(t)}{1 + \omega_b(t)} \frac{1 + \omega_b(t^{\frac{\kappa}{n+\kappa}})}{\omega_b(t^{\frac{\kappa}{n+\kappa}})} \right) \\ &\leq \lambda_4(\kappa) \left(1 + \frac{1}{\omega_a(1)} + \frac{1}{\omega_b(1)} \right) =: \lambda_2, \end{aligned}$$

where we have used the assumption (11.1)₂. On the other hand, recalling that the functions ω_a and ω_b are increasing, we have

$$\Lambda \left(t, \frac{1}{t} \right) \leq \Lambda \left(t^{\frac{\kappa}{n+\kappa}}, \frac{1}{t} \right) \leq \lambda_4(\kappa) \leq \lambda_2$$

for every $t \in (0, 1]$. Recalling that $u \in L^\infty_{\text{loc}}(\Omega)$ by Theorem 11.2 and taking into account the last two displays, we are able to apply Theorem 5.2 in order to have (11.15) and (11.16). \square

Remark 11.3. As a consequence of the last two theorems like we have that if $u \in W^{1,\Psi}(\Omega)$ is a local Q -minimizer of the functional \mathcal{P} under the assumption (11.1), then $u \in L^\infty_{\text{loc}}(\Omega)$ and (1.13)₂ is satisfied. Therefore, the results of Theorem 1.1, Theorem 1.2, Theorem 10.1 and Theorem 10.2 are still available under the assumption (11.1). Furthermore, the results of the present section can be considered under multi-phase settings, as we have pointed out in Remark 1.1.

REFERENCES

1. E. Acerbi and N. Fusco, *An harmonic approximation lemma for $W^{1,p}$ functions*, Material Instabilities in Continuum Mechanics (Edinburg, 1985-1986), pp.1-5. Oxford Science Publications, Oxford University Press, New York (1988).
2. R.A. Adams and J.J.F. Fournier, *Sobolev spaces*, 2nd ed., Pure and Applied Mathematics (Amsterdam) **140**, Elsevier/Academic Press, Amsterdam, 2003.
3. E. Acerbi and G. Mingione, *Regularity results for a class of functionals with non-standard growth*, Arch. Ration. Mech. Anal., **156**, (2001) 121-140.
4. E. Acerbi and G. Mingione, *Regularity results for electrorheological fluids: the stationary case*, C. R. Math. Acad. Sci. Paris, **334**, (2002) 817-822.
5. E. Acerbi and G. Mingione, *Gradient estimates for a class of parabolic systems*, Duke Math. J., **136** (2) (2007), 285-320.
6. S. Baasandorj, S. Byun and J. Oh, *Calderón-Zygmund estimates for generalized double phase problems*, J. Funct. Anal. **279** (7) (2020), 108670.
7. S. Baasandorj, S. Byun and J. Oh, *Gradient estimates for multi-phase problems*, Calc. Var. Partial Differential Equations, to appear.
8. S. Baasandorj, S. Byun and H. Lee, *Global gradient estimates for a general class of quasilinear elliptic equations with Orlicz growth*, Proc. Amer. Math. Soc., to appear.
9. Anna Kh. Balci, L. Diening and M. Surnachev, *New Examples on Lavrentiev Gap Using Fractals*, Calc. Var. Partial Differential Equations **59:180** (2020).
10. P. Baroni, *Riesz potential estimates for a general class of quasilinear equations*, Calc. Var. Partial Differential Equations **53** (2015), 803-846.
11. P. Baroni, M. Colombo and G. Mingione, *Harnack inequalities for double phase functionals*, Nonlinear Anal. **121** (2015), 206-222.
12. P. Baroni, M. Colombo and G. Mingione, *Non-autonomous functionals, borderline cases and related function classes*, St. Petersburg Math. J. **27** (2016), 347-379.
13. P. Baroni, M. Colombo and G. Mingione, *Regularity for general functionals with double phase*, Calc. Var. Partial Differential Equations **57** (2) (2018), Art. 62, 48 pp.
14. L. Beck and G. Mingione, *Optimal Lipschitz criteria and local estimates for non-uniformly elliptic problems*, Atti Accad. Naz. Lincei Rend. Lincei Mat. Appl. **30** (2) (2019), 223-236.
15. A. Benkirane and M. Sidi El Vally, *Variational inequalities in Musielak-Orlicz-Sobolev spaces*, Bull. Belg. Math. Soc. Simon Stevin **21** (5) (2014), 787-811.
16. P. Bildhauer and M. Fuchs, *$C^{1,\alpha}$ -solutions to non-autonomous anisotropic variational problems*, Calc. Var. Partial Differential Equations **24**, (2005), 309-340.
17. P. Bousquet and L. Brasco, *Global Lipschitz continuity for minima of degenerate problems*, Math. Ann. **366**, (2016), 1403-1450.
18. D. Breit, *New regularity theorems for non-autonomous variational integrals with (p,q) -growth*, Calc. Var. Partial Differential Equations **44** (1-2) (2012), 101-129.
19. Kateryna O. Buryachenko and Igor I. Skrypnik, *Local continuity and Harnack's inequality for double-phase parabolic equations*, Potential Analysis **23** (2020).
20. S. Byun and J. Oh, *Global gradient estimates for non-uniformly elliptic equations*, Calc. Var. Partial Differential Equations **56** (2) (2017), Art. 46, 36 pp.
21. S. Byun and J. Oh, *Global gradient estimates for the borderline case of double phase problems with BMO coefficients in nonsmooth domains*, J. Differential Equations **263** (2) (2017), 1643-1693.
22. S. Byun and J. Oh, *Regularity results for generalized double phase functionals*, Anal. PDE **13** (5) (2020), 1269-1300.
23. S. Byun and S. Ryu, *Global weighted estimates for the gradient of solutions to nonlinear elliptic equations*, Ann. Inst. H. Poincaré Anal. Non Linéaire, **30** (2) (2013), 291-313.
24. S. Byun and Y. Cho, *Nonlinear gradient estimates for generalized elliptic equations with nonstandard growth in nonsmooth domains*, Nonlinear Analysis, **140** (2016), 145-165.
25. S. Byun and Y. Youn, *Riesz potential estimates for a class of double phase problems*, J. Differential Equations **264** (2) (2018), 1263-1316.
26. A. Cianchi and L. Pick, *Sobolev embeddings into spaces of Campanato, Morrey, and Hölder type*, J. Math. Anal. Appl. **282** (1) (2003), 128-150.
27. A. Cianchi, *Optimal Orlicz-Sobolev embeddings*, Rev. Mat. Iberoamericana **20** (2004), 427-474.
28. A. Cianchi, *Orlicz-Sobolev boundary trace embeddings*, Mathematische Zeitschrift **266** (2010), 431-449.
29. I. Chlebicka, *A pocket guide to nonlinear differential equations in Musielak-Orlicz spaces*, Nonlinear Analysis **175**(2018), 1-27.
30. M. Colombo and G. Mingione, *Regularity for double phase variational problems*, Arch. Ration. Mech. Anal. **215** (2) (2015), 443-496.

31. M. Colombo and G. Mingione, *Bounded minimisers of double phase variational integrals*, Arch. Ration. Mech. Anal. **218** (1) (2015), 219-273.
32. M. Colombo and G. Mingione, *Calderón-Zygmund estimates and non-uniformly elliptic operators*, J. Funct. Anal. **270** (4) (2016), 1416-1478.
33. G. Cupini, *Local boundedness of minimizers with limit growth conditions*, J. Optim. Theory Appl. **166** (1) (2015), 1-22.
34. G. Cupini, P. Marcellini, E. Mascolo, *Regularity of minimizers under limit growth conditions*, Nonlinear Anal. **153** (2017), 294-310.
35. C. De Filippis and G. Mingione, *A borderline case of Calderón-Zygmund estimates for nonuniformly elliptic problems*, St. Petersburg Math. J. **31** (3) (2019), 82-115.
36. C. De Filippis and G. Mingione, *Manifold constrained non-uniformly elliptic problems*, J. Geom. Anal., **30** (2020), 1661-1723.
37. C. De Filippis and G. Mingione, *On the regularity of minima of non-autonomous functionals*, J. Geom. Anal., **30** (2020), 1584-1626.
38. C. De Filippis and J. Oh, *Regularity for multi-phase variational problems*, J. Differential Equations **267** (3) (2019), 1631-1670.
39. C. De Filippis, *Gradient bounds for solutions to irregular parabolic equations with (p, q) -growth*, **59** (5) (2020), 1-32.
40. L. Diening, *Maximal function on Musielak-Orlicz spaces and generalized Lebesgue spaces*, Bull. Sci. Math. **129** (8) (2005), 657-700.
41. L. Diening and F. Ettwein, *Fractional estimates for non-differentiable elliptic systems with general growth*, Forum Math. **20** (3) (2008), 523-556.
42. L. Diening, B. Stroffolini and A. Verde, *Everywhere regularity of functionals with φ -growth*, Manuscripta Math. **129** (4) (2009), 449-481.
43. L. Diening, B. Stroffolini and A. Verde, *The φ -harmonic approximation and the regularity of φ -harmonic maps*, J. Differ. Equ. **253** (2012), 1943-1958.
44. F. Duzaar and G. Mingione, *Harmonic type approximation lemmas*, J. Math. Anal. Appl. **352** (2009), 301-335.
45. L. Esposito, F. Leonetti and G. Mingione, *Regularity for minimizers of functionals with $p-q$ growth*, Nonlinear Differential Equations Appl. **6** (2) (1999), 133-148.
46. L. Esposito, F. Leonetti and G. Mingione, *Regularity results for minimizers of irregular integrals with (p, q) growth*, Forum Math. **14** (2) (2002), 245-272.
47. L. Esposito, F. Leonetti and G. Mingione, *Sharp regularity for functionals with (p, q) growth*, J. Differential Equations **204** (1) (2004), 5-55.
48. L. Esposito, G. Mingione and C. Trombetti, *On the Lipschitz regularity for certain elliptic problems*, Forum Math. **18** (2) (2006), 263-292.
49. L. C. Evans, *A new proof of local $C^{1,\alpha}$ regularity for solutions of certain degenerate elliptic equations*, J. Differential Equations **45** (3) (1982), 356-373.
50. I. Fonseca, J. Malý and G. Mingione, *Scalar minimizers with fractal singular sets*, Arch. Ration. Mech. Anal. **172** (2) (2004), 295-307.
51. N. Fusco and C. Sbordone, *Higher integrability of the gradient of minimizers of functionals with nonstandard growth conditions*, Comm. Pure Appl. Math. **43** (5) (1990), 673-683.
52. M. Giaquinta and E. Giusti, *On the regularity of the minima of variational integrals*, Acta Math. **148** (1982), 31-46.
53. E. Giusti, *Direct methods in the calculus of variations*, World Scientific Publishing Co., Inc., River Edge, NJ, 2003.
54. D. Gilbarg and N.S. Trudinger, *Elliptic partial differential equations of second order*, Grundlehren der Mathematischen Wissenschaften, vol. **224**, Springer, Berlin, 1977.
55. P. Harjulehto and P. Hästö, *Orlicz Spaces and Generalized Orlicz Spaces*, Lecture notes in mathematics, **2236**, Springer, 2019.
56. P. Harjulehto, P. Hästö and R. Klén, *Generalized Orlicz spaces and related PDE*, Nonlinear Anal. **143** (2016), 155-173.
57. P. Harjulehto, P. Hästö and M. Lee *Hölder continuity of quasiminimizers and ω -minimizers of functionals with generalized Orlicz growth*, Ann. Sc. Norm. Super. Pisa Cl. Sci., to appear.
58. P. Harjulehto, P. Hästö and O. Toivanen, *Hölder regularity of quasiminimizers under generalized growth conditions*, Calc. Var. Partial Differential Equations **56** (2) (2017), Art. 22, 26 pp.
59. P. Harjulehto, T. Kuusi, T. Lukkari, N. Marola and M. Parviainen, *Harnack's inequality for quasiminimizers with non-standard growth conditions*, J. Math. Anal. Appl. **344**:1 (2008), 504-520.
60. P. Hästö and J. Ok, *Maximal regularity for local minimizers of non-autonomous functionals*, J. Eur. Math. Soc., to appear.
61. A. Karppinen *Global continuity and higher integrability of a minimizer of an obstacle under generalized Orlicz growth conditions*, Manuscripta Math. **164** (2021), 67-94.
62. J. Kinnunen and N. Shanmugalingam, *Regularity of quasi-minimizers of metric spaces*, Manuscripta Math. **105**:3 (2001), 401-423.
63. O. A. Ladyzhenskaya and N. N. Ural'tseva, *Linear and quasilinear elliptic equations*, Academic Press, New York, 1968.
64. J. L. Lewis, *Regularity of the derivatives of solutions to certain degenerate elliptic equations*, Indiana Univ. Math. J. **32** (6) (1983), 849-858.
65. G. M. Lieberman, *The natural generalization of the natural conditions of Ladyzhenskaya and Ural'tseva for elliptic equations*, Comm. Partial Differential Equations. **16** (1991), 311-361.
66. P. Lindqvist, *Notes on the p -Laplace equation*, Report Univ. Jyväskylä, vol. 102, Univ. Jyväskylä, Jyväskylä, 2006.
67. J. J. Manfredi, *Regularity for minima of functionals with p -growth*, J. Differential Equations **76** (2) (1988), 203-212.

68. P. Marcellini, *On the definition and the lower semicontinuity of certain quasiconvex integrals*, Ann. Inst. H. Poincaré Anal. Non Linéaire **3** (5) (1986), 391-409.
69. P. Marcellini, *Regularity of minimizers of integrals of the calculus of variations with nonstandard growth conditions*, Arch. Ration. Mech. Anal. **105** (3) (1989), 267-284.
70. P. Marcellini, *Regularity and existence of solutions of elliptic equations with p, q -growth conditions*, J. Differential Equations **90** (1) (1991), 1-30.
71. G. Mingione, *Regularity of minima: an invitation to the dark side of the calculus of variations*, Appl. Math. **51** (2006), 355-425.
72. J. Musielak, *Orlicz spaces and modular spaces*, Lecture Notes in Mathematics, 1034, Springer-Verlag, Berlin, 1983.
73. J. Ok, *Regularity of ω -minimizers for a class of functionals with non-standard growth*, Calc. Var. Partial Differential Equations **56** (2) (2017), Art. 48, 31 pp.
74. J. Ok, *Partial Hölder regularity for elliptic systems with non-standard growth*, J. Funct. Anal. **274** (3) (2018), 723-768.
75. J. Ok, *Regularity for double phase problems under additional integrability assumptions*, Nonlinear Analysis **194** (2020), 111408.
76. M.A. Ragusa and A. Tachikawa, *Partial regularity of $p(x)$ -harmonic maps*, Trans. Am. Math. Soc. **365** (2013), 3329-3353.
77. M.A. Ragusa and A. Tachikawa, *Regularity regularity of minimizers of $p(x)$ -energy functionals*, Ann. IHP Anal. non Linéaire **33** (2017), 451-476.
78. H.L. Royden and P.M. Fitzpatrick, *Real Analysis*, Upper Saddle River, NJ : Pearson Prentice Hall, 3rd ed.,(2010).
79. M. M. Rao and Z. D. Ren *Theory of Orlicz spaces*, Monogr. Textbooks Pure Appl. Math. **146**, Dekker, Newyork, 1991.
80. T. Schmidt, *Regularity of minimizers of $W^{1,p}$ -quasiconvex variational integrals with (p, q) -growth*, Calc. Var. Partial Differential Equations **32** (1) (2008), 1-24.
81. T. Schmidt, *Regularity of relaxed minimizers of quasiconvex variational integrals with (p, q) -growth*, Arch. Ration. Mech. Anal. **193** (2) (2009), 311-337.
82. M. Sidi El Vally, *Strongly nonlinear elliptic problems in Musielak-Orlicz-Sobolev spaces*, Adv. Dyn. Syst. Appl. **8** (1) (2013), 115-124.
83. A. Tachikawa and K. Usuba, *Regularity results up to the boundary for minimizers of $p(x)$ -energy with $p(x) > 1$* , Manus.Math. **152** (2017), 127-151.
84. K. Uhlenbeck, *Regularity for a class of non-linear elliptic systems*, Acta Math. **138**, (1977), 219-240.
85. N.N. Ural'tseva, *Degenerate quasilinear elliptic systems*, Zap. Na. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI) **7**, (1968), 184-222.
86. V.V. Zhikov, *Averaging of functionals of the calculus of variations and elasticity theory*, Izv. Akad. Nauk SSSR Ser. Mat. **50** (4) (1986), 675-710.
87. V.V. Zhikov, *Laurentiev phenomenon and homogenization for some variational problems*, C. R. Acad. Sci. Paris Sér. I Math. **316** (5) (1993), 435-439.
88. V.V. Zhikov, *On Lavrentiev's phenomenon*, Russian J. Math. Phys. **3** (2) (1995), 249-269.
89. V.V. Zhikov, *On some variational problems*, Russian J. Math. Phys. **5** (1) (1997), 105-116.
90. V.V. Zhikov, S.M. Kozlov and O.A. Oleinik, *Homogenization of differential operators and integral functionals*, Springer-Verlag, Berlin, 1994.

DEPARTMENT OF MATHEMATICAL SCIENCES, SEOUL NATIONAL UNIVERSITY, SEOUL 08826, KOREA

Email address: summa2017@snu.ac.kr

DEPARTMENT OF MATHEMATICAL SCIENCES AND RESEARCH INSTITUTE OF MATHEMATICS, SEOUL NATIONAL UNIVERSITY, SEOUL 08826, KOREA

Email address: byun@snu.ac.kr