

ESTIMATING GAUSSIAN MIXTURES USING SPARSE POLYNOMIAL MOMENT SYSTEMS

JULIA LINDBERG, CARLOS AMÉNDOLA, AND JOSE ISRAEL RODRIGUEZ

ABSTRACT. The method of moments is a statistical technique for density estimation that solves a system of moment equations to estimate the parameters of an unknown distribution. A fundamental question critical to understanding identifiability asks how many moment equations are needed to get finitely many solutions and how many solutions there are. We answer this question for classes of Gaussian mixture models using the tools of polyhedral geometry. Using these results, we present an algorithm that performs parameter recovery, and therefore density estimation, for high dimensional Gaussian mixture models that scales linearly in the dimension.

1. INTRODUCTION

A fundamental problem in statistics is to estimate the parameters of a density from samples. This problem is called *density estimation* and formally it asks: given n samples from an unknown distribution p , can we estimate p ? To have any hope of solving this problem we need to assume our density lives in a family of distributions. One family of densities known as Gaussian mixture models are a popular choice due to their broad expressive power.

Theorem 1.1. [21, Ch. 3] A Gaussian mixture model is a universal approximator of densities, in the sense that any smooth density can be approximated with any specific nonzero amount of error by a Gaussian mixture model with enough components.

Theorem 1.1 motivates our study of Gaussian mixture models. These are ubiquitous in the literature with applications in modeling geographic events [53], the spread of COVID-19 [44], the design of planar steel frame structures [18], speech recognition [40, 52, 42], image segmentation [9] and biometrics [25].

A *Gaussian random variable*, X , has a probability density function given by

$$f(x|\mu, \sigma^2) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(x-\mu)^2}{2\sigma^2}\right),$$

where $\mu \in \mathbb{R}$ is the mean and $\sigma \in \mathbb{R}_{>0}$ is the standard deviation. In this case we write $X \sim \mathcal{N}(\mu, \sigma^2)$. A random variable X is the *mixture of k Gaussians* if its probability density function is the convex combination of k Gaussian densities. Here we write $X \sim \sum_{\ell=1}^k \lambda_{\ell} \mathcal{N}(\mu_{\ell}, \sigma_{\ell}^2)$ where $\mu_{\ell} \in \mathbb{R}$, $\sigma_{\ell} \in \mathbb{R}_{>0}$ for all $\ell \in [k] = \{1, \dots, k\}$ and $(\lambda_1, \dots, \lambda_k) \in \Delta_{k-1} = \{\lambda \in \mathbb{R}_{>0}^k : \sum_{\ell=1}^k \lambda_{\ell} = 1\}$. Each λ_{ℓ} , $\ell \in [k]$, is the mixture weight of the ℓ th component.

A *Gaussian k -mixture model* is a collection of mixtures of k Gaussian densities. Often one imposes constraints on the means, variances or weights to define such

models. For example, one might assume that all variances are equal or that the mixture weights are all equal. The former is known as a *homoscedastic* mixture model, and the latter is called a *uniform* mixture model. In this paper we consider three classes of Gaussian mixture models.

- (1) The $\bar{\lambda}$ -weighted model, where the mixture weights are fixed for $\bar{\lambda} \in \Delta_{k-1}$.
- (2) The $\bar{\lambda}$ -weighted homoscedastic model, which is the $\bar{\lambda}$ -weighted model under the additional assumption that the variances are equal.
- (3) The $\bar{\lambda}$ -weighted known variance model, where the weights and variances are assumed known.

We wish to do parameter recovery for these classes of Gaussian mixture models, that is, we would like to solve the following problem.

Problem 1.2. Given y_1, \dots, y_N sampled from a mixture of k Gaussian densities, recover the parameters $\mu_\ell, \sigma_\ell^2, \lambda_\ell$ for $\ell \in [k]$.

These samples can be summarized by sample moments. Motivated by Problem 1.2 and the *method of moments*, in this paper we solve the following problem.

Problem 1.3. Given moments from a Gaussian k mixture density, recover the parameters of each Gaussian component.

The fact one can recover the mean, variance and weight of each component, up to relabeling, if given the mixture density function is known as *mixture identifiability* and was proven for univariate Gaussians in [46]. Multivariate Gaussians are still identifiable [51] and there exist finitely many moments that identify them [7]. In this paper we are interested in identifying the means and variances of each mixture from its moments up to a certain order.

It is important to distinguish parameter recovery from density estimation. For density estimation, one aims to estimate a density that is close to the true mixture density, with no restriction on how close each individual component is. In this paper we wish to do parameter recovery. Namely, we wish to recover accurate estimates of the mean, variance and mixing weight of each component. It is clear that density estimation follows once all of the parameters are known. The distinction between density estimation and parameter recovery is illustrated next.

Example 1.4. Consider the vector of eight moments (m_1, \dots, m_8) :

(0.1661, 2.133, 1.3785, 12.8629, 16.0203, 125.6864, 239.2856, 1695.5639).

There are two Gaussian 3-mixture densities with these eight moments. The weights and individual components for each of the mixture models are shown in Figure 2. These densities are shown in Figure 1 where it is seen that they are almost indistinguishable. In contrast, the individual components and weights are noticeably different. The weights and individual components for each of the mixture models are shown in Figure 2.

One idea to solve Problem 1.2 is to use *maximum likelihood estimation*. Maximum likelihood estimation aims to maximize the likelihood function by solving the following




FIGURE 1. Two distinct Gaussian mixture densities with $k = 3$ components and the same first eight moments.




FIGURE 2. Individual components of two Gaussian mixture models with similar mixture densities.

optimization problem:

$$(1.1) \quad \operatorname{argmax}_{\mu, \sigma^2, \lambda} \prod_{i=1}^N \sum_{i=1}^k \lambda_i \frac{1}{\sqrt{2\pi\sigma_i^2}} \exp\left(-\frac{(y_i - \mu_i)^2}{2\sigma_i^2}\right).$$

Unless $k = 1$, (1.1) is a nonconvex optimization problem and obtaining a global optimum is difficult or impossible. In general (1.1) is unbounded, so no global maximum exists. Iterative algorithms such as expectation maximization (EM) try to find the largest local maximum [17]. On top of being sensitive to the starting point, another downside of the EM algorithm is that it needs to access all data in each iteration, which is prohibitive for applications with large data sets. Further, there is no known bound on the number of critical points of the likelihood function, and it has been shown that no such bound can exist that is independent of the sample size [1].

Recent work has analyzed the local behavior of (1.1) by considering maximum likelihood estimation for two Gaussians in \mathbb{R}^n when $\lambda_1 = \lambda_2 = \frac{1}{2}$ and all of the covariance matrices are known and equal — this is a special case of the models we consider. It has been shown that in this regime the EM algorithm converges geometrically to the global optimum [16, 50]. Other work has studied the global landscape of the EM algorithm and the structure of local optima in this setting [13, 50]. Further work has considered inference for Gaussian mixture models with known mixing coefficients and identity covariance matrices [36] and clustering analysis of the mixture of two Gaussians where the covariance matrices are equal and unknown [12]. When these covariance matrices are further assumed to be spherical, [41] gives polynomial time approximation schemes for (1.1) so long as the mixtures are sufficiently well-separated.

Recently, techniques from numerical algebraic geometry have been used to identify the number of components in a Gaussian mixture model [43]. Further progress has been made on giving optimal sampling bounds needed to learn a Gaussian mixture model [5].

Another idea for density estimation in this set up is to use the *generalized method of moments*. The generalized method of moments was proposed in [23] and aims to minimize the difference between the fitted moments and the sample moments. For Gaussian mixture models, this again cannot be solved in a way guaranteeing global optimality due to the nonconvexity of the moment equations. Recently this method has been remedied for Gaussian mixture models in one dimension with the same variance parameter, where the authors provably and efficiently find the global optimum of the generalized method of moments [49]. It is important to note that in many of the cases above, assumptions are made on the values that each Gaussian component can take.

In this paper we propose using the method of moments to estimate the density arising from the mixture of k multivariate Gaussians. This methodology was first proposed and resolved for the mixture of two univariate Gaussians by Karl Pearson [39]. Pearson reduced solving this system of 6 polynomial equations in the 6 unknown density parameters, $\mu_\ell, \sigma_\ell^2, \lambda_\ell$, $\ell = 1, 2$, to understanding the roots of a degree nine polynomial with coefficients in the sample moments \bar{m}_i , $i = 1, \dots, 5$.

The method of moments for Gaussian mixture models was revisited in 2010 in a series of papers [28, 37]. The case of a $k = 2$ mixture model in n dimensions was handled in [28] where a polynomial time algorithm was presented. This approach was generalized in [37] where an algorithm for a general k -mixture model in n dimensional space was presented that scales polynomially in n and the number of samples required scales polynomially in the desired accuracy. When $k = 2, 3, 4$ the number of solutions to the corresponding moment system are given in [39, 2, 4] respectively.

1.1. Contributions. The main result of this paper is to present an algorithm (Algorithm 1) for parameter recovery for a mixture of k Gaussians in \mathbb{R}^n that scales linearly in n . Our results are organized as follows. We give a fresh presentation of the statistics and algebraic background needed for the study of method of moments. The main take away in this section is that the method of moments can be thought of as solving a system of sparse polynomial equations. By studying this system we can make conclusions about parameter recovery and density estimation. For example, we show that for a wide range of models the corresponding moment system has finitely many solutions, which leads to identifiability results. Moreover, the proofs of our results give rise to effective algorithms that implement the method of moments for mixtures of univariate Gaussians. As a surprising application of our theorems in the univariate case, we show that density estimation for Gaussian mixture models in \mathbb{R}^n can be done using homotopy continuation by tracking $\mathcal{O}(n)$ paths. In an Appendix, we also prove a new result concerning how many moments are needed to identify mixtures of Gaussians; namely, we show that Gaussian k -mixtures are rationally identifiable from $3k + 2$ moments.

2. PRELIMINARIES

We first introduce some necessary concepts from statistics and algebraic geometry.

2.1. Method of moments. This paper focuses on an approach for parameter recovery known as the *method of moments*. The method of moments for parameter estimation is based on the law of large numbers. This approach expresses the moments of a density as functions of its parameters.

For $r \geq 0$, we denote the r -th moment of a random variable X as $m_r = \mathbb{E}[X^r]$. We consider a statistical model with n unknown parameters, $\theta = (\theta_1, \dots, \theta_n)$, and consider the moments up to order n as functions of θ , $g_1(\theta), \dots, g_n(\theta)$. For random samples y_1, \dots, y_N we denote the sample moment by $\bar{m}_r = \frac{1}{N} \sum_{j=1}^N y_j^r$.

The method of moments works by using samples from the statistical model to calculate sample moments $\bar{m}_1, \dots, \bar{m}_n$, and then solve the corresponding system $\bar{m}_i = g_i(\theta)$, $i = 1, \dots, n$, for the parameters $\theta_1, \dots, \theta_n$.

The moments of Gaussian distributions are polynomials in the two indeterminants μ, σ^2 and can be calculated recursively as $M_0(\mu, \sigma^2) = 1$, $M_1(\mu, \sigma^2) = \mu$ and

$$(2.1) \quad M_i(\mu, \sigma^2) = \mu \cdot M_{i-1}(\mu, \sigma^2) + (i-1) \cdot \sigma^2 \cdot M_{i-2}(\mu, \sigma^2), \quad i \geq 2.$$

We calculate the i th moment of a mixture of k Gaussian densities as the convex combinations of $M_i(\mu_1, \sigma_1^2), \dots, M_i(\mu_k, \sigma_k^2)$.

For each nonnegative integer i , we define the polynomial f_i^k in the $3k$ indeterminants unknowns $\mu_\ell, \sigma_\ell, \lambda_\ell$, $\ell \in [k]$ and sample moment \bar{m}_i to be

$$(2.2) \quad f_i^k(\mu, \sigma^2, \lambda) := \lambda_1 M_i(\mu_1, \sigma_1^2) + \dots + \lambda_k M_i(\mu_k, \sigma_k^2) - \bar{m}_i.$$

We are then interested in solving the system

$$(2.3) \quad f_i^k(\mu, \sigma^2, \lambda) = 0$$

under assumptions on the parameters, μ, σ^2, λ where i varies over an appropriate index set.

We use an overline, e.g., \bar{x} to indicate a quantity is fixed and known. For the three classes of Gaussian mixture models defined in the introduction, we assume $\bar{\lambda} \in \Delta_{k-1}$ and we solve these respective systems:

$$(2.4) \quad f_i^k(\mu, \sigma^2, \bar{\lambda}) = 0, \quad i \in [2k], \quad (\text{Section 3.1})$$

$$(2.5) \quad f_i^k(\mu, \sigma^2, \bar{\lambda}) = 0, \quad i \in [k+1], \quad \sigma_1^2 = \sigma_2^2 = \dots = \sigma_k^2, \quad (\text{Section 3.2})$$

$$(2.6) \quad f_i^k(\mu, \bar{\sigma}^2, \bar{\lambda}) = 0, \quad i \in [k], \quad (\text{Section 3.4}).$$

2.2. Genericity. The concept of *generic* behavior is prevalent in applied algebraic geometry. It allows one to make mathematically precise statements without characterizing complicated algebraic sets. Informally, one says a property is *generic* if it can be described in terms of the complement of a proper algebraic subset. In particular, this algebraic subset is lower dimensional, so that non-generic behavior is restricted to a Lebesgue measure-zero set. We now define what genericity means in our context.

Consider a statistical model \mathcal{M} parametrized by Θ , where $\theta \mapsto p_\theta \in \mathcal{M}$ and $\dim(\Theta) = d$. Let

$$(2.7) \quad \phi : \Theta \rightarrow \mathbb{R}^d$$

be a *truncated moment map* that takes θ to a list of d moments of the distribution p_θ of prescribed order. We assume throughout the article that ϕ is a polynomial map.

We define the set of *generic parameters* in Θ (with respect to ϕ) to be

$$(2.8) \quad \Theta_{\phi, \text{gen}} := \{\theta \in \Theta : \det \text{Jac}(\phi)|_\theta \neq 0\}.$$

The set $\Theta_{\phi, \text{gen}}$ is the complement of an algebraic set restricted to Θ . Therefore, when $\Theta_{\phi, \text{gen}}$ is nonempty, we have $\Theta \setminus \Theta_{\phi, \text{gen}}$ is a Lebesgue measure-zero set.

Our next goal is to define generic moments. To do so, first we define the set \mathcal{B}_ϕ in the ambient space of moments to be the smallest algebraic set containing

$$(2.9) \quad \{\phi(\theta) \in \mathbb{R}^d : \det \text{Jac}(\phi)|_\theta = 0\} \subset \mathbb{C}^d.$$

The set of *generic moments* (with respect to ϕ) is taken to be the set complement

$$(2.10) \quad \mathbb{C}^d \setminus \mathcal{B}_\phi,$$

while the set of *realizable generic moments* is

$$(2.11) \quad V_\phi := \{(m_1, \dots, m_d) \in \phi(\Theta) : (m_1, \dots, m_d) \notin \mathcal{B}_\phi\}.$$

The set $\mathbb{C}^d \setminus \mathcal{B}_\phi$ is the complement of an algebraic set, which allows us to leverage tools from algebraic geometry. Proving statements about $\mathbb{C}^d \setminus \mathcal{B}_\phi$ allows us to make statements relevant for parameter recovery by restricting to realizable generic moments V_ϕ .

Example 2.1. Fix $\bar{\lambda} = (\frac{1}{2}, \frac{1}{2})$ in Δ_1 . As our model, we consider the collection of $\bar{\lambda}$ -weighted Gaussian 2-mixtures with mean 0 and second moment fixed to be 1. This model is parameterized by

$$\Theta = \left\{ (\mu, \sigma) : \frac{1}{2}\mu_1 + \frac{1}{2}\mu_2 = 0, \frac{1}{2}\sigma_1^2 + \frac{1}{2}\sigma_2^2 = 1 \right\}.$$

This model is two dimensional (since, e.g., μ_2, σ_2 can be expressed in terms of μ_1, σ_1). Choose ϕ to map $\theta \in \Theta$ to the third and fourth order moments: $\phi(\mu, \sigma) = (m_3, m_4)$ where

$$\begin{aligned} m_3 &= \frac{1}{2}\mu_1(\mu_1^2 + 3\sigma_1^2) + \frac{1}{2}\mu_2(\mu_2^2 + 3\sigma_2^2) \\ m_4 &= \frac{1}{2}(\mu_1^2(\mu_1^2 + 6\sigma_1^2) + 3\sigma_1^4) + \frac{1}{2}(\mu_2^2(\mu_2^2 + 6\sigma_2^2) + 3\sigma_2^4). \end{aligned}$$

The set of generic parameters are those where this 2×2 Jacobian matrix is invertible

$$\begin{bmatrix} 3\sigma_1^2 - 3 & 3\mu_1 \\ 4\mu_1^3 + 12\mu_1 & 6\sigma_1^2 - 6 \end{bmatrix}.$$

In other words, non-generic parameters are those such that the determinant of the matrix vanishes, i.e., this equation is satisfied

$$2\mu_1^4 + 6\mu_1^2 - 3\sigma_1^4 + 6\sigma_1^2 - 3 = 0.$$

For this model, the set of generic moments is the complement of the algebraic set

$$\mathcal{B}_\phi = \{(m_3, m_4) \in \mathbb{C}^2 : 3m_3^4 + 2(m_4 - 3)^3 = 0\}.$$

For example, the moments $(m_3, m_4) = (0, 3) \in \mathcal{B}_\phi$ are not generic. This makes sense from a statistical viewpoint, since $(m_1, m_2, m_3, m_4) = (0, 1, 0, 3)$ are precisely moments of a single standard Gaussian distribution, and as such it is not an honest Gaussian 2-mixture.

The set of realizable generic moments is the open set

$$V_\phi = \{(m_3, m_4) \in \phi(\Theta) : 3m_3^4 + 2(m_4 - 3)^3 \neq 0\}.$$

For $(m_3, m_4) \in V_\phi$, the corresponding moment system has only six complex solutions.

Definition 2.2. A statistical model parameterized by $\Theta \subset \mathbb{R}^d$ is said to be *algebraically identifiable* from ϕ in (2.7), if $\Theta_{\phi, \text{gen}}$ is d -dimensional.

In other words, for realizable generic moments $\phi(\theta^*)$, there are at most finitely many other parameters θ in Θ such that $\phi(\theta) = \phi(\theta^*)$. Similarly, we also consider a notion of identifiability where $\phi(\theta) = \phi(\theta^*)$ implies $\theta = \theta^*$.

Definition 2.3. A statistical model parameterized by $\Theta \subset \mathbb{R}^s$ is said to be *rationally identifiable* from moments m_1, \dots, m_d if there exists an algebraic set \mathcal{A} in \mathbb{C}^s such that $\Theta_{\text{gen}} := \Theta \setminus \mathcal{A}$ is s -dimensional and for each $\theta \in \Theta_{\text{gen}}$ the distribution p_θ is uniquely identifiable from the moments m_1, \dots, m_d .

The main result in [3] states that univariate Gaussian k -mixture models are algebraically identifiable from $\phi(\theta) = (m_1, \dots, m_{3k-1})$. This means that for generic moments m_1, \dots, m_{3k-1} , there are only finitely many Gaussian k -mixture distributions with exactly those moments. In particular, the polynomial system (2.3) for $i = 0, \dots, 3k - 1$ has finitely many solutions. Our results in Section 3 are for three classes of submodel of Gaussian k -mixtures. These are far from trivial as the moments needed to algebraically identify a submodel cannot be immediately determined. Moreover, the number of solutions to the moment systems for the submodel and model can be different as Example 2.1 illustrates.

2.3. Statistically meaningful solutions. We note that for any set of real-valued sample moments it is not guaranteed that the moment equations will give any statistically meaningful solutions. A *statistically meaningful solution* is a real valued solution with positive variances and mixing weights. In other words, it is a solution that corresponds to a true density. If the sample moments are inaccurate, it may happen that no solutions obtained from the method of moments is statistically meaningful. By the law of large numbers, as the number of samples goes to infinity the sample moments will converge to the true moments and the method of moments will return a consistent estimator [47, Theorem 9.6].

A property of parameterized polynomial systems is that the number of real solutions is constant in open Euclidean sets of the parameter space. Such open sets can be computed via cylindrical algebraic decomposition [14]. The constraints differing real solutions and statistically meaningful ones will further divide these cells. Therefore,

in any of these open cells the number of statistically meaningful solutions will be constant. So long as the sample moments lie in a cell that has at least one statistically meaningful solution, which will happen with increasing probability as the sample size increases, the method of moments will return a true density.

Example 2.4. Consider the setting from Example 2.1 with a mixture of two Gaussians with equal mixing weights, $\lambda_1 = \lambda_2 = \frac{1}{2}$ and with $m_1 = 0, m_2 = 1$. In this case there are four moment equations in the four unknowns $\mu_1, \sigma_1^2, \mu_2, \sigma_2^2$. There is one open cell that dictates whether or not there is a statistically meaningful solution. The region where there is precisely one statistically meaningful solution (up to symmetry) is shown in blue in Figure 3, where the horizontal and vertical axes correspond to m_3 and m_4 , respectively.




FIGURE 3. Blue region shows where there is one statistically meaningful solution for $k = 2$, $m_1 = 0, m_2 = 1$ and $\lambda_1 = \lambda_2 = \frac{1}{2}$.

2.4. Sparse polynomial systems. We now review concepts from algebraic geometry, but for an introduction to sparse polynomial systems see [45, Ch. 3]. A family of sparse polynomials is defined by its monomial support. For each $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}_{\geq 0}^n$, we have the monomial $x^\alpha := x_1^{\alpha_1} \cdots x_n^{\alpha_n}$ with exponent α . A *sparse polynomial* is a linear combination of monomials. Let $\mathcal{A}_\bullet = (\mathcal{A}_1, \dots, \mathcal{A}_k)$ denote a k -tuple of nonempty finite subsets of $\mathbb{N}_{\geq 0}^n$. A sparse polynomial system of equations with support \mathcal{A}_\bullet is given by

$$\sum_{\alpha \in \mathcal{A}_1} c_{1,\alpha} x^\alpha = \dots = \sum_{\alpha \in \mathcal{A}_k} c_{k,\alpha} x^\alpha = 0,$$

where $\{c_{i,\alpha}\}_{\alpha \in \mathcal{A}_i, i \in [k]}$ are the scalar coefficients. The terminology “sparse” is used because the monomial support \mathcal{A}_\bullet is typically much smaller compared to the number of monomials up to a specific degree.

2.5. Mixed volumes. Consider two polytopes $P, Q \in \mathbb{R}^n$. The *Minkowski sum* of P and Q is defined as

$$P + Q := \{p + q : p \in P, q \in Q\}.$$

A *scaling* by $t \in \mathbb{R}_{>0}$ of a polytope P is defined as $tP := \{tp : p \in P\}$. Given s convex polytopes in \mathbb{R}^n , K_1, \dots, K_s , we consider the standard n -dimensional Euclidean volume of a linear combination of these polytopes

$$\nu(t_1, \dots, t_s) = \text{Vol}_n\left(\sum_{i=1}^s t_i K_i\right)$$

where the sum here refers to the Minkowski sum and the t_i refers to a scaling. The polynomial $\nu(t_1, \dots, t_s)$ is homogeneous of degree n in t_1, \dots, t_s .

The *mixed volume* of s convex polytopes K_1, \dots, K_s in \mathbb{R}^n is the coefficient of the $t_1 t_2 \cdots t_s$ term in $\nu(t_1, \dots, t_s)$. It is denoted $\text{MVol}(K_1, \dots, K_s)$. More information on mixed volumes and polytopes can be found in [20].

Example 2.5. Consider $P = \text{Conv}\{(0, 0), (0, 1), (1, 0), (1, 1)\}$, the unit square. We compute $\text{MVol}(P, P)$ by considering

$$\nu(t_1, t_2) = \text{Vol}_2(t_1 P + t_2 P) = (t_1 + t_2)(t_1 + t_2) = t_1^2 + 2t_1 t_2 + t_2^2.$$

The coefficient in front of $t_1 t_2$ is 2, giving $\text{MVol}(P, P) = 2$.

Example 2.6. The mixed volume of K_1, \dots, K_n is easy to describe when K_i is a line segment from the origin to the vertex $v_i \in \mathbb{Z}^n$. The Minkowski sum $t_1 K_1 + \cdots + t_n K_n$ is a parallelepiped. Hence, its volume is given by a determinant:

$$\nu(t_1, t_2, \dots, t_n) = \left| \det \begin{bmatrix} t_1 v_1 & t_2 v_2 & \cdots & t_n v_n \end{bmatrix} \right|.$$

So $\text{MVol}(K_1, \dots, K_n)$ equals the absolute value of the determinant of the matrix with the vertices v_1, \dots, v_n as its columns.

2.6. BKK bound. A celebrated series of papers [8, 30, 31] gives the connection between sparse polynomial systems described in Section 2.4 and polyhedral geometry described in Section 2.5.

Consider a polynomial $f = \sum_{\alpha \in \mathcal{A}} c_\alpha x^\alpha \in \mathbb{R}[x_1, \dots, x_n]$ with monomial support $\mathcal{A} \subset \mathbb{N}^n$ where x^α denotes $x_1^{\alpha_1} \cdots x_n^{\alpha_n}$. The *Newton polytope* of f is the convex hull of its exponent vectors $\alpha \in \mathcal{A}$, denoted $\text{Newt}(f)$.

Theorem 2.7 (Bernstein-Khovanskii-Kouchnirenko Bound). Let $\mathcal{A} := (\mathcal{A}_1, \dots, \mathcal{A}_n)$ where $\mathcal{A}_i \subset \mathbb{N}^n$ and $0_n \in \mathcal{A}_i$ for all $i \in [n]$. Let $\mathcal{L}(\mathcal{A})$ be the collection of polynomials (f_1, \dots, f_n) where \mathcal{A}_i is the support of f_i and $P_i = \text{Conv}(\mathcal{A}_i)$ for all $i \in [n]$. Consider $F = (f_1, \dots, f_n)$ where $F \in \mathcal{L}(\mathcal{A})$ and the number of complex solutions to $F = 0$ is finite. Then the number of complex solutions to $F = 0$ is less than or equal to $\text{MVol}(P_1, \dots, P_n)$. Moreover, for generic choices of $F \in \mathcal{L}(\mathcal{A})$, equality holds.

We illustrate Theorem 2.7 on a basic example from statistics.

Example 2.8. The method of moments can be used to determine the parameters of a Gaussian distribution $\mathcal{N}(\mu, \sigma^2)$. For sample moments $\overline{m}_1, \overline{m}_2$, these parameters are determined by solving the sparse polynomial system

$$\begin{aligned} f_1^1(\mu, \sigma^2) &= \overline{m}_1 - \mu = 0, \\ f_2^1(\mu, \sigma^2) &= \overline{m}_2 - \mu^2 - \sigma^2 = 0 \end{aligned}$$

in the unknowns μ, σ^2 . The monomial supports corresponding to f_1^1, f_2^1 are

$$\mathcal{A}_1 = \{(0, 0), (1, 0)\}, \quad \mathcal{A}_2 = \{(0, 0), (2, 0), (0, 1)\}.$$

The mixed volume $\text{MVol}(\text{Conv}(\mathcal{A}_1), \text{Conv}(\mathcal{A}_2))$ and the number of solutions to the system are both one. This is no coincidence by the BKK bound in Theorem 2.7.

Remark 2.9. The BKK theorem is usually stated without the hypothesis $0_n \in \mathcal{A}_i$. In this case, the BKK bound counts the number of complex solutions with nonzero coordinates. To get a bound on the number of complex solutions, the assumption $0_n \in \mathcal{A}_i$ was added in [34].

Remark 2.10. A special case of the BKK bound is the Bézout bound [24, p. 47]. The Bézout bound says that if a polynomial system $F = (f_1, \dots, f_n)$ has finitely many solutions, then the number of complex solutions with nonzero coordinates is bounded above by $d_1 \cdots d_n$ where $\deg(f_i) = d_i$.

Theorem 2.7 states that for a generic sparse polynomial system in $\mathcal{L}(\mathcal{A})$, the number of complex solutions equals the mixed volume of the system. An obstacle to applying this theorem is that the mixed volume is #P-hard to compute [19].

An important property of mixed volumes is that they are *monotonic*. Namely, if $\hat{P}_1 \subseteq P_1$ then

$$\text{MVol}(\hat{P}_1, P_2, \dots, P_n) \leq \text{MVol}(P_1, P_2, \dots, P_n).$$

Therefore by Example 2.6, taking line segments $\text{Conv}(\{0, v_i\}) = Q_i \subseteq P_i$ for $i \in [n]$, is an easy way to get a lower bound on $\text{MVol}(P_1, \dots, P_n)$. We would like conditions under which such a lower bound is tight.

Definition 2.11. [38, Definition VII.32] Let P_1, \dots, P_m be convex polytopes in \mathbb{R}^n . We say P_1, \dots, P_m are *dependent* if there is a non-empty subset $\mathcal{I} \subseteq [m]$ such that $\dim(\sum_{i \in \mathcal{I}} P_i) < |\mathcal{I}|$. Otherwise we say P_1, \dots, P_m are *independent*.

This definition may be difficult to parse on a first read. But it is related to the usual definition of *linear independence*: if each P_i is a line through the origin, then the two ideas of dependent agree. Moreover, the collection of empty sets is independent.

Given a nonzero vector $w \in \mathbb{R}^n$ and a convex polytope $P \subseteq \mathbb{R}^n$, we consider the function $P \rightarrow \mathbb{R}$, $x \mapsto \langle w, x \rangle$ with $\langle (w_1, \dots, w_n), (x_1, \dots, x_n) \rangle := w_1 x_1 + \dots + w_n x_n$ denoting the standard inner product. We denote the minimum value w takes on P by $\text{val}_w(P)$, and the points of P where w is minimized by $\text{Init}_w(P)$. This set of points is a face of P , which we call the *face exposed* by w . Specifically, we have

$$\text{val}_w(P) = \min_{x \in P} \langle w, x \rangle \quad \text{and} \quad \text{Init}_w(P) = \{x \in P : \langle w, x \rangle \leq \langle w, y \rangle \text{ for all } y \in P\}.$$

If $f = \sum_{\alpha \in \text{Newt}(f)} c_\alpha x^\alpha$, we call

$$\text{Init}_w(f) := \sum_{\alpha \in \text{Init}_w(\text{Newt}(f))} c_\alpha x^\alpha$$

the *initial polynomial* of f . For more background on initial polynomials, see [15, Chapter 7].

Proposition 2.12. [38, Proposition VII.39] Let $P_i = \text{Conv}(\mathcal{A}_i)$ and $Q_i = \text{Conv}(\mathcal{B}_i) \subseteq P_i$ for $i \in [n]$. The following are equivalent:

- (1) $\text{MVol}(P_1, \dots, P_n) = \text{MVol}(Q_1, \dots, Q_n)$
- (2) One of the following holds:
 - (a) P_1, \dots, P_n are dependent i.e. $\text{MVol}(P_1, \dots, P_n) = 0$
 - (b) For each $w \in \mathbb{R}^n \setminus \{0\}$, the collection of polytopes

$$\{\text{Init}_w(Q_i) : Q_i \cap \text{Init}_w(P_i) \neq \emptyset\}$$

is dependent.

Proposition 2.12 gives conditions under which it suffices to consider the (potentially much simpler) polytopes $Q_i \subseteq P_i$ to compute the mixed volume of P_1, \dots, P_n .

Example 2.13. Consider the triangles $P_1 = P_2 = \text{Conv}(\{(0, 0), (1, 0), (0, 1)\})$, and the line segments

$$Q_1 = \text{Conv}(\{(0, 0), (1, 0)\}) \subset P_1, \quad Q_2 = \text{Conv}(\{(0, 0), (0, 1)\}) \subset P_2.$$

Direct computation shows

$$\text{MVol}(P_1, P_2) = 1 = \text{MVol}(Q_1, Q_2).$$

We can also use Example 2.6 to prove $\text{MVol}(Q_1, Q_2) = 1$ and Proposition 2.12 to prove $\text{MVol}(P_1, P_2) = \text{MVol}(Q_1, Q_2)$ since for any nonzero $w \in \mathbb{Z}^2$, the collection of polytopes $\{\text{Init}_w(Q_i) : Q_i \cap \text{Init}_w(P_i) \neq \emptyset\}$, $i = 1, 2$, contains a single point so the collection is dependent. This type of argument, where we use the dependence of polytopes and Proposition 2.12, is also used in the proofs of Theorem 3.3 and Theorem 3.7.

The following lemma will be of use later to apply Proposition 2.12 in the proof of our main results.

Lemma 2.14. Let $P_i \subseteq \mathbb{R}^n$ and $Q_i = \text{Conv}(\{0_n, v_i\}) \subseteq P_i$ for $i \in [n]$ be convex polytopes. Consider the set, W , of nonzero $w \in \mathbb{R}^n$ such that $Q_i \cap \text{Init}_w(P_i) \neq \emptyset$ for all $i \in [n]$. I.e.,

$$W = \{w \in \mathbb{R}^n \setminus \{0_n\} : Q_i \cap \text{Init}_w(P_i) \neq \emptyset, \forall i \in [n]\}.$$

If $\{v_1, \dots, v_n\}$ are linearly independent, then the polytopes

$$\{\text{Init}_w(Q_1), \dots, \text{Init}_w(Q_n)\}$$

are dependent for all $w \in W$.

Proof. Fix $w \in W$. By the definition of dependent, we need to show that

$$\dim \left(\sum_{i=1}^n \text{Init}_w(Q_i) \right) < n.$$

Since $\text{Init}_w(Q_i) \subseteq \mathbb{R}^n$, for all $i \in [n]$, $\sum_{i=1}^n \dim(\text{Init}_w(Q_i)) \leq n$. Furthermore, since each Q_i is one dimensional, $\sum_{i=1}^n \dim(\text{Init}_w(Q_i)) = n$ if and only if w minimizes all of Q_i for all $i \in [n]$. I.e., $\sum_{i=1}^n \dim(\text{Init}_w(Q_i)) = n$ if and only if $\text{Init}_w(Q_i) = Q_i$ for all $i \in [n]$.

Recalling $Q_i = \text{Conv}(\{0_n, v_i\})$, one sees $\text{Init}_w(Q_i) = Q_i$ if and only if

$$0 = \langle w, 0_n \rangle = \langle w, v_i \rangle,$$

for all $i \in [n]$. Since $\{v_1, \dots, v_n\}$ are linearly independent, the only w that satisfies this is $w = 0_n \notin W$. \square

3. PARAMETER RECOVERY IN ONE DIMENSION

We are now able to present our first set of results: efficiently finding all complex solutions stemming from the moment equations. As a consequence of each theorem, we show a model is algebraically identifiable from an appropriately chosen truncated moment map ϕ .

In each subsection, we use Θ to parameterizes the model covered in that section. In each case though ϕ (introduced in (2.7)) is the truncated moment map using the first $\dim(\Theta)$ moments.

3.1. Mixed volume of $\bar{\lambda}$ -weighted models. First consider a $\bar{\lambda}$ -weighted model with k mixture components. We study the moment system

$$(3.1) \quad f_1^k(\mu, \sigma^2, \bar{\lambda}) = 0, \dots, f_{2k}^k(\mu, \sigma^2, \bar{\lambda}) = 0,$$

where $\bar{\lambda}$ are known mixing coefficients, and f_i^k , $i \in [2k]$ is as defined in (2.3). In this set up the unknowns are μ_ℓ, σ_ℓ , $\ell \in [k]$.

First, we record the following fact about the moment functions $M_k(\mu, \sigma^2)$ of a Gaussian.

Lemma 3.1. The partial derivatives of M_k satisfy

$$\frac{\partial}{\partial \mu} M_k(\mu, \sigma^2) = k \cdot M_{k-1}(\mu, \sigma^2) \quad \text{and} \quad \frac{\partial}{\partial \sigma^2} M_k(\mu, \sigma^2) = \binom{k}{2} \cdot M_{k-2}(\mu, \sigma^2).$$

Proof. We verify both by induction. For both identities, the base case $k = 1$ is immediate. Suppose $\frac{\partial}{\partial \mu} M_{k-1}(\mu, \sigma^2) = (k-1) \cdot M_{k-2}(\mu, \sigma^2)$. By the recursive relationship (2.1) we have

$$\begin{aligned} \frac{\partial}{\partial \mu} M_k &= M_{k-1} + \mu \frac{\partial}{\partial \mu} M_{k-1} + \sigma^2 (k-1) \frac{\partial}{\partial \mu} M_{k-2} \\ &= M_{k-1} + \mu (k-1) M_{k-2} + \sigma^2 (k-1) (k-2) M_{k-3} \\ &= M_{k-1} + (k-1) M_{k-1} = k M_{k-1}. \end{aligned}$$

Similarly, suppose that $\frac{\partial}{\partial \sigma^2} M_{k-1}(\mu, \sigma^2) = \binom{k-1}{2} M_{k-3}(\mu, \sigma^2)$. Using (2.1) again,

$$\begin{aligned} \frac{\partial}{\partial \sigma^2} M_k &= \mu \frac{\partial}{\partial \sigma^2} M_{k-1} + (k-1) M_{k-2} + \sigma^2 (k-1) \frac{\partial}{\partial \sigma^2} M_{k-2} \\ &= \mu \binom{k-1}{2} M_{k-3} + (k-1) M_{k-2} + \sigma^2 (k-1) \binom{k-2}{2} M_{k-4} \\ &= (k-1) M_{k-2} + \binom{k-1}{2} (\mu M_{k-3} + \sigma^2 (k-3) M_{k-4}) \\ &= (k-1) M_{k-2} + \binom{k-1}{2} M_{k-2} = \binom{k}{2} M_{k-2}. \end{aligned}$$

□

Now we prove our first algebraic identifiability result.

Proposition 3.2. Fix $\bar{\lambda} \in \Delta_{k-1}$ and let Θ parameterize the $\bar{\lambda}$ -weighted Gaussian k -mixture model. For $(\bar{m}_1, \dots, \bar{m}_{2k}) \in V_\phi$, the set of complex solutions to (3.1) is finite and nonempty.

Proof. By [24, Ch. 1, Section 5] it suffices to show that the Jacobian of (3.1) has a nonzero determinant. This Jacobian, denoted by J_k , is a $2k \times 2k$ matrix with rows indexed by equations and columns indexed by the unknowns $\mu_1, \sigma_1, \dots, \mu_k, \sigma_k$:

$$(3.2) \quad J_k = \tilde{J}_k \cdot \tilde{D}_k$$

$$= \begin{bmatrix} \frac{\partial M_1}{\partial \mu_1}(\sigma_1, \mu_1) & \frac{\partial M_1}{\partial \sigma_1}(\sigma_1, \mu_1) & \dots & \frac{\partial M_1}{\partial \mu_k}(\sigma_k, \mu_k) & \frac{\partial M_1}{\partial \sigma_k}(\sigma_k, \mu_k) \\ \frac{\partial M_2}{\partial \mu_1}(\sigma_1, \mu_1) & \frac{\partial M_2}{\partial \sigma_1}(\sigma_1, \mu_1) & \dots & \frac{\partial M_2}{\partial \mu_k}(\sigma_k, \mu_k) & \frac{\partial M_2}{\partial \sigma_k}(\sigma_k, \mu_k) \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \frac{\partial M_{2k-1}}{\partial \mu_1}(\sigma_1, \mu_1) & \frac{\partial M_{2k-1}}{\partial \sigma_1}(\sigma_1, \mu_1) & \dots & \frac{\partial M_{2k-1}}{\partial \mu_k}(\sigma_k, \mu_k) & \frac{\partial M_{2k-1}}{\partial \sigma_k}(\sigma_k, \mu_k) \\ \frac{\partial M_{2k}}{\partial \mu_1}(\sigma_1, \mu_1) & \frac{\partial M_{2k}}{\partial \sigma_1}(\sigma_1, \mu_1) & \dots & \frac{\partial M_{2k}}{\partial \mu_k}(\sigma_k, \mu_k) & \frac{\partial M_{2k}}{\partial \sigma_k}(\sigma_k, \mu_k) \end{bmatrix} \cdot \tilde{D}_k,$$

where \tilde{D}_k is the diagonal matrix given by $(\bar{\lambda}_1, \bar{\lambda}_1, \bar{\lambda}_2, \bar{\lambda}_2, \dots, \bar{\lambda}_k, \bar{\lambda}_k)$. Note that for nonzero λ_ℓ , $\ell \in [k]$, J_k is full rank if and only if \tilde{J}_k is full rank.

We now show that \tilde{J}_k has a determinant that is not identically zero by induction on k . When $k = 1$, the determinant of $\tilde{J}_1 = \begin{bmatrix} 1 & 0 \\ 2\mu_1 & 1 \end{bmatrix}$ is 1.

Next note that μ_k, σ_k^2 only appear in the last two columns of \tilde{J}_k . Further, by Lemma 3.1, the nonzero entries of each row of \tilde{J}_k have higher degree than the previous row. Doing Laplace's cofactor expansion along the last two columns of \tilde{J}_k , we get

$$\det(\tilde{J}_k) = \det(\tilde{J}_{k-1}) \cdot \mu_k^{2k-2} \sigma_k^{2k-2} + \text{lower order terms}.$$

By induction, $\det(\tilde{J}_{k-1})$ is not identically zero and neither is $\det(\tilde{J}_k)$. □

We now use Theorem 2.7 and Proposition 2.12 to give an upper bound on the number of complex solutions to (3.1). Recall that if N is odd, the double factorial is defined as

$$N!! = 1 \cdot 3 \cdot 5 \cdots N.$$

Theorem 3.3 (Mixing coefficients known). Fix $\bar{\lambda} \in \Delta_{k-1}$, and let Θ parameterize the $\bar{\lambda}$ -weighted Gaussian k -mixture model. For $(\bar{m}_1, \dots, \bar{m}_{2k}) \in V_\phi$, the number of complex solutions to (3.1) is at most $(2k-1)!! \cdot k!$.

Proof. Consider the moment equations $f_1^k = 0, \dots, f_{2k}^k = 0$ as defined in (3.1), but now ordering the unknowns as $(\mu_1, \sigma_1^2, \dots, \mu_k, \sigma_k^2)$. Denote $P_i = \text{Newt}(f_i^k)$.

Let $Q_\ell \subset P_\ell$ be the line segment defined as:

$$\begin{aligned} Q_{2\ell-1} &= \text{Conv}(\{0_{2k}, (2\ell-1) \cdot e_{2\ell-1}\}), \quad \ell \in [k] \\ Q_{2\ell} &= \text{Conv}(\{0_{2k}, \ell \cdot e_{2\ell}\}), \quad \ell \in [k], \end{aligned}$$

where $0_{2k} \in \mathbb{R}^{2k}$ is the vector of all zeros and $e_\ell \in \mathbb{R}^{2k}$ is the ℓ th standard basis vector. By Example 2.6 we have

$$\text{MVol}(Q_1, \dots, Q_{2k}) = (1 \cdot 3 \cdot 5 \cdots (2k-1)) \cdot (1 \cdot 2 \cdot 3 \cdots k) = (2k-1)!!k!.$$

We want to use the equivalence of (1) and (2) in Proposition 2.12 to show

$$\text{MVol}(P_1, \dots, P_{2k}) = (2k-1)!!k!.$$

Theorem 2.7 then gives that the number of complex solutions to (3.1) is bounded above by $(2k-1)!!k!$.

For a nonzero vector $w \in \mathbb{R}^{2k}$, let

$$\mathcal{I}_w = \{i \in [2k] : Q_i \cap \text{Init}_w(P_i) \neq \emptyset\}.$$

We will show for each $w \in \mathbb{R}^n \setminus \{0\}$, the collection of polytopes

$$\{\text{Init}_w(Q_i) : i \in \mathcal{I}_w\}$$

is dependent.

By Lemma 3.4, (which we postpone to after this proof) \mathcal{I}_w is nonempty. Since each Q_i is a one dimensional line segment, it suffices to show that for any nonzero $w \in \mathbb{R}^{2k}$, w minimizes some Q_i at a unique point for $i \in \mathcal{I}_w$. This follows from the definition of dependent since each Q_i is one dimensional, so if $\text{Init}_w(Q_i)$ is a single point for some $i \in \mathcal{I}_w$, then $\sum_{i \in \mathcal{I}_w} \text{Init}_w(Q_i) < |\mathcal{I}_w|$.

We look at two cases.

- First, consider when $2\ell \notin \mathcal{I}_w$ for all $\ell \in [k]$. Since the origin is in Q_i , we have $0_{2k} \notin \text{Init}_w(P_{2\ell})$ for all $\ell \in [k]$. Since $P_{2\ell} \subset \mathbb{R}_{\geq 0}^{2k}$, this means $\text{val}_w(P_{2\ell}) < 0$ for all $\ell \in [k]$. Hence, $w_i < 0$ for some $i \in [2k]$.

Let i be the index of the smallest element of w . If i is odd, then

$$P_i = \text{Conv}\left(\left\{0_{2k}, \frac{i-1}{2}e_2, \dots, \frac{i-1}{2}e_{2k}, ie_1, \dots, ie_{2k-1}\right\}\right)$$

is in the nonnegative orthant. So

$$\text{val}_w(P_i) = \min\left\{0, \frac{i-1}{2}w_2, \dots, \frac{i-1}{2}w_k, iw_1, \dots, iw_{2k-1}\right\} = iw_i.$$

So $\text{Init}_w(Q_i) = \{ie_i\}$ for $i \in \mathcal{I}_w$ and we are done.

Now consider when i is even. Recall,

$$P_{2\ell} = \text{Conv}\left(\{0_{2k}, \ell e_2, \ell e_4, \dots, \ell e_{2k}, 2\ell e_1, 2\ell e_3, \dots, 2\ell e_{2k-1}\}\right) = \ell \cdot P_2,$$

and so $\text{Init}_w(P_{2\ell}) = \ell \cdot \text{Init}_w(P_2)$ for all $\ell \in [k]$. Therefore, $2\ell \notin \mathcal{I}_w$ for all $\ell \in [k]$ implies that P_2 cannot be minimized at e_j for any even j . This implies that if i is even, $0 > w_i > 2w_j$ for some odd j (otherwise P_i would be minimized at $\frac{i}{2}e_i$). Let j be the index of the smallest odd element of w . In this case, P_j would be minimized at je_j so $j \in \mathcal{I}_w$. Hence $\text{Init}_w(Q_j)$ is $\{je_j\}$.

- Second, suppose $2\ell \in \mathcal{I}_w$ for some $\ell \in [k]$. If $\text{Init}_w(P_{2\ell}) \cap Q_{2\ell}$ is a point, then we are done. Otherwise, we may assume $P_{2\ell}$ is minimized by w at a face containing the line segment $Q_{2\ell} = \text{Conv}(\{0_{2k}, \ell e_{2\ell}\})$.

This means

$$0 = w^T 0_{2k} \leq w^T a \quad \forall a \in P_{2\ell}.$$

So $w_i \geq 0$ for all $i \in [2k]$ because the vertices of $P_{2\ell}$ are scaled standard basis vectors. With the fact each P_i is in the nonnegative orthant, this implies

$$0 = \text{val}_w(P_i) \quad \text{for all } i \in [2k],$$

so $0_{2k} \in \text{Init}_w(P_i)$ for all $i \in [2k]$.

This shows that $\mathcal{I}_w = [2k]$ so by Lemma 2.14, we conclude that the collection of polytopes $\{\text{Init}_w(Q_1), \dots, \text{Init}_w(Q_{2k})\}$ is dependent. □

Lemma 3.4. The index set \mathcal{I}_w as defined in the proof of Theorem 3.3 is nonempty for any nonzero $w \in \mathbb{R}^{2k}$.

Proof. Recall that

$$P_2 = \text{Conv}(\{0_{2k}, e_2, \dots, e_{2k}, 2e_1, \dots, 2e_{2k-1}\}).$$

We consider three cases, depending on $\text{Init}_w(P_2)$.

- If either 0_{2k} or e_2 is in $\text{Init}_w(P_2)$, then $Q_2 \cap \text{Init}_w(P_2) \neq \emptyset$ and hence $2 \in \mathcal{I}_w$.
- Now suppose $e_j \in \text{Init}_w(P_2)$ for some even $j > 2$. This means $2w_j \leq w_i$ for all odd i and $w_j \leq w_m$ for all even m . Consider

$$P_j = \text{Conv}(\{0_{2k}, je_1, \frac{j}{2}e_2, \dots, je_{2k-1}, \frac{j}{2}e_{2k}\}).$$

Then $\frac{j}{2}e_j \in \text{Init}_w(P_j)$. Since $Q_j = \text{Conv}(\{0_{2k}, \frac{j}{2}e_j\})$, we have $j \in \mathcal{I}_w$.

- Now suppose $e_i \in \text{Init}_w(P_2)$ for some odd $i \geq 1$. This means $w_i \leq w_j$ for all odd j and $2w_i \leq w_m$ for all even m . Consider

$$P_i = \text{Conv}(\{0_{2k}, \frac{i-1}{2}e_2, \dots, \frac{i-1}{2}e_{2k}, ie_1, \dots, ie_{2k-1}\}).$$

Then $ie_i \in \text{Init}_w(P_i)$. Since $Q_i = \text{Conv}(\{0_{2k}, ie_i\})$, we have $i \in \mathcal{I}_w$. □

Remark 3.5. An instance of the previous theorem is when the mixing weights are all equal. In this case, there are $(2k-1)!!$ solutions up to the standard label-swapping symmetry. There exist special techniques in numerical nonlinear algebra that exploit this symmetry for computational speed up. One such technique known as *monodromy* was recently used for this problem with success [4, Section 4.1].

3.2. Mixed volume of $\bar{\lambda}$ -weighted homoscedastic models. We consider the $\bar{\lambda}$ -weighted homoscedastic model. In this setting the means are unknown and the variances are unknown but all equal. In this case, a k -mixture model has $k + 1$ unknowns. We address the high dimensional version of this problem in Section 4 which is also considered in recent work, for example [12].

We consider the moment system

$$(3.3) \quad f_1^k(\mu, \sigma^2, \bar{\lambda}) = 0, \dots, f_{k+1}^k(\mu, \sigma^2, \bar{\lambda}) = 0, \quad \sigma_1^2 = \sigma_2^2 = \dots = \sigma_k^2$$

where $\bar{\lambda}$ are known mixing coefficients, and f_i^k , $i \in [k + 1]$ is as defined in (2.3). In this set up, the $k + 1$ unknowns are μ_ℓ , $\ell \in [k]$ and σ^2 .

Again, the first step is to prove that this model is algebraically identifiable.

Proposition 3.6. Fix $\bar{\lambda} \in \Delta_{k-1}$, and let Θ parameterize the $\bar{\lambda}$ -weighted homoscedastic model. For $(\bar{m}_1, \dots, \bar{m}_{k+1}) \in V_\phi$, the number of solutions to (3.3) is finite.

Proof. This argument is similar to the one given in Proposition 3.2. Again, we consider the Jacobian of (3.3) with rows indexed by equations and columns indexed by the variables $\sigma^2, \mu_1, \dots, \mu_k$. It suffices to show that for the Jacobian, J_k is not identically zero. We proceed by induction on k . When $k = 1$,

$$J_1 = \begin{bmatrix} 0 & \bar{\lambda}_1 \\ \bar{\lambda}_1 & 2\bar{\lambda}_1\mu_1 \end{bmatrix}$$

has determinant $-\bar{\lambda}_1^2$ and is not identically zero because we assume $\bar{\lambda}_i \neq 0$. Now consider the $(k + 1) \times (k + 1)$ matrix J_k for any k . By cofactor expansion along the last column,

$$\det(J_k) = \mu_k^k \cdot \det(J_{k-1}) + \text{lower order terms in } \mu_k.$$

By induction, $\det(J_{k-1})$ is not identically zero. This shows that $\det(J_k)$ is a nonzero univariate polynomial in μ_k with nonzero leading coefficient $\det(J_{k-1})$. \square

We bound the number of solutions to (3.3) for a generic $\bar{\lambda}$ -weighted homoscedastic mixture model using mixed volumes.

Theorem 3.7 (Mixing coefficients known, variances equal). Fix $\bar{\lambda} \in \Delta_{k-1}$, and let Θ parameterize the $\bar{\lambda}$ -weighted homoscedastic Gaussian k -mixture model. For $(\bar{m}_1, \dots, \bar{m}_{k+1}) \in V_\phi$, the number of complex solutions to (3.3) is at most $\frac{(k+1)!}{2}$.

Proof. Let $P_i = \text{Newt}(f_i^k)$ where f_i^k is as defined in (3.3) with variable ordering $(\mu_1, \dots, \mu_k, \sigma^2)$ for $i \in [k + 1]$.

Define $Q_i \subset P_i$ as follows:

$$(3.4) \quad \begin{aligned} Q_1 &= \text{Conv}(\{0_{k+1}, e_1\}) \\ Q_2 &= \text{Conv}(\{0_{k+1}, e_{k+1}\}) \\ Q_i &= \text{Conv}(\{0_{k+1}, i \cdot e_{i-1}\}), \quad 3 \leq i \leq k + 1. \end{aligned}$$

where $0_{k+1} \in \mathbb{R}^{k+1}$ is the vector of all zeros and $e_i \in \mathbb{R}^{k+1}$ is the i th standard basis vector. By Example 2.6,

$$\text{MVol}(Q_1, \dots, Q_{k+1}) = 1 \cdot 3 \cdot 4 \cdots (k+1) = \frac{(k+1)!}{2}.$$

As in the proof of Theorem 3.3, we want to show the equality of mixed volumes $\text{MVol}(P_1, \dots, P_{k+1}) = \text{MVol}(Q_1, \dots, Q_{k+1})$ by using the equivalence of (1) and (2) in Proposition 2.12.

Let \mathcal{I}_w be the set of indices such that Q_i has a vertex in $\text{Init}_w(P_i)$. Specifically,

$$\mathcal{I}_w = \{i \in [k+1] : Q_i \cap \text{Init}_w(P_i) \neq \emptyset\}.$$

By Lemma 3.8, \mathcal{I}_w is nonempty. Now we want to show that for any $w \in \mathbb{Z}^{k+1} \setminus \{0\}$ the nonempty set of polytopes

$$\{\text{Init}_w(Q_i) : i \in \mathcal{I}_w\}$$

is dependent. Since Q_i is a one dimensional line segment, it suffices to show that for any w , there exists an $i \in \mathcal{I}_w$ such that Q_i is minimized at a single vertex. We consider 2 cases.

- Suppose $2 \in \mathcal{I}_w$. If $\text{Init}_w(P_2)$ is a single point, then we are done. Otherwise, assume P_2 is minimized at $Q_2 = \text{Conv}(\{0_{k+1}, e_{k+1}\})$. This means $w_{k+1} = 0$ and $w_j \geq 0$ for all $j \in [k]$, giving $\text{val}_w(P_i) = 0$ so $0 \in \text{Init}_w(P_i)$ for all $i \in [k]$. This shows that $\mathcal{I}_w = [k+1]$. By Lemma 2.14, the collection of polytopes $\{\text{Init}_w(Q_1), \dots, \text{Init}_w(Q_{k+1})\}$ is dependent.
- Now suppose $2 \notin \mathcal{I}_w$. If $1 \in \mathcal{I}_w$ and Q_1 is minimized at a single vertex then $\{\text{Init}_w(Q_i) : i \in \mathcal{I}_w\}$ is dependent, so we are done. If Q_1 is not minimized at a single vertex, then $w_1 = 0$ and $w_j \geq 0$ for all $2 \leq j \leq k$. Since $2 \notin \mathcal{I}_w$, this gives that $w_{k+1} > 2w_j$ for all $j \in [k]$. Therefore, $\text{val}_w(P_j) = 0$ for all $j \in [k+1]$ which shows that $0 \in \text{Init}_w(P_2)$. This contradicts $2 \notin \mathcal{I}_w$. On the other hand, if $i \in \mathcal{I}_w$ where $i \geq 3$, either Q_i is minimized at a single vertex or $w_{i-1} = 0$ and $w_j \geq 0$ for all $j \in [k+1] \setminus (i-1)$. In the latter case, this shows $\text{val}_w(P_i) = 0$ for all $i \in [k+1]$, contradicting $2 \notin \mathcal{I}_w$.

□

Lemma 3.8. The index set \mathcal{I}_w as defined in the proof of Theorem 3.7 is nonempty for any nonzero $w \in \mathbb{R}^{k+1}$.

Proof. If w is in the nonnegative orthant, then w minimizes P_i at the origin for all $i \in [k+1]$. In this case $\mathcal{I}_w = [k+1] \neq \emptyset$. Now let i be the index of the smallest element of w . If $i = k+1$ then w minimizes P_2 and Q_2 at $\{e_{k+1}\}$. If $i < k+1$ then w minimizes P_{i+1} and Q_{i+1} at $\{(i+1)e_i\}$, which shows \mathcal{I}_w is nonempty. □

3.3. Finding all solutions using homotopy continuation. To do parameter recovery for any of the set ups described in Section 3.1 and Section 3.2, it is not enough to know the number of complex solutions to the moment equations, we need a way to find all of them. Finding all complex solutions to a zero dimensional variety is

a well-studied problem in numerical algebraic geometry. We outline the basics of homotopy continuation below but give [6] for a detailed reference.

Consider the system of polynomial equations

$$F(x) = (f_1(x), f_2(x), \dots, f_n(x)) = 0.$$

where the number of complex solutions to $F(x) = 0$ is finite and $x = (x_1, \dots, x_n)$. The main idea is to construct a *homotopy* [33]

$$(3.5) \quad H(x; t) = \gamma(1 - t)G(x) + tF(x)$$

such that these three conditions hold:

- (1) the solutions to $H(x, 0) = G(x) = 0$ are trivial to find,
- (2) there are no singularities along the path $t \in [0, 1)$, and
- (3) all isolated solutions of $H(x, 1) = F(x) = 0$ can be reached.

Using a random $\gamma \in \mathbb{C}$ ensures that there are no singularities along the path $t \in [0, 1)$ so condition two is met. Continuation methods known as *predictor-corrector methods* are used to track the solutions from $G(x) = 0$ to $F(x) = 0$ as t varies from 0 to 1 [11].

It is standard terminology to call $G(x) = 0$ the *start system*. There are many choices for constructing a start system. A *total degree* start system is of the form

$$G(x) = \{x_1^{d_1} - 1, \dots, x_n^{d_n} - 1\} = 0,$$

where d_i is the degree of f_i . The number of solutions to $G(x) = 0$ is $d_1 \cdots d_n$, which is the Bézout bound of $G(x)$. Using this start system, one must track $d_1 \cdots d_n$ paths in order to find all solutions to $F(x) = 0$. If $F(x) = 0$ has close to $d_1 \cdots d_n$ solutions this is a reasonable choice.

If $F(x) = 0$ is a polynomial system with fewer solutions than the Bézout bound, then tracking $d_1 \cdots d_n$ paths is unnecessary. Instead, one can leverage the system's Newton polytopes to construct a start system that has as many solutions as the mixed volume. One example of this is the polyhedral homotopy method [26] which tracks N paths where N is the BKK bound described in Theorem 2.7. This method constructs several binomial start systems, that is, systems formed by polynomials with precisely two terms. The main drawback to polyhedral methods is that there is often a computational bottleneck associated with computing the start system. Our related approach circumvents this bottleneck and relies on the following lemma.

The collection of Newton polytopes of a polynomial system $F = (f_1, \dots, f_n)$ is denoted by $\text{Newt}(F) = (\text{Newt}(f_1), \dots, \text{Newt}(f_n))$.

Lemma 3.9. Suppose $G(x) = 0$ is a general sparse binomial system such that we have $\text{MVol}(\text{Newt}(G)) = \text{MVol}(\text{Newt}(F))$ and $\text{Newt}(G) \subseteq \text{Newt}(F)$ element-wise. If the origin is in each Newton polytope of G , then the three items above hold for the homotopy (3.5).

Proof. By Theorem 2.7 the number of solutions for $G(x) = 0$ equals the generic number of solutions for a polynomial system with Newton polytopes $\text{Newt}(F)$. Since $\text{Newt}(G) \subseteq \text{Newt}(F)$ and γ is generic, we have $\text{Newt}(F) = \text{Newt}(\gamma(1 - t)G + tF)$ for $t \in (0, 1]$. So the mixed volume, and therefore the number of solutions, of $(1 - t)G + tF$ agrees with the mixed volume of $\text{Newt}(F)$. \square

The fact that the total degree homotopy works is a special case of the previous lemma applied to polynomials with full monomial support. Combining Lemma 3.9 with Theorem 3.3 and Theorem 3.7 we get the following corollary.

Corollary 3.10. The binomial system induced by the polytopes Q_i in the proofs of Theorem 3.3 and Theorem 3.7 constructs an optimal homotopy continuation start system for the corresponding moment system.

Proof. In this proof we construct the homotopy; give its start points; and show that it is optimal. We only show the details for the case of Theorem 3.3 because the other statement's proof is analogous.

Consider the binomial system

$$\begin{aligned} g_{2\ell-1} &= a_\ell \mu_\ell^{2\ell-1} + b_\ell, & \ell \in [k] \\ g_{2\ell} &= c_\ell (\sigma_\ell^2)^\ell + d_\ell, & \ell \in [k] \end{aligned}$$

where $a_\ell, b_\ell, c_\ell, d_\ell \in \mathbb{C}^*$ are generic for $\ell \in [k]$.

Since each $g_{2\ell-1}$ and $g_{2\ell}$ is a univariate polynomial in distinct variables, multiplying the degrees we know that there are $(2k-1)!!k!$ solutions. This number agrees with the mixed volume of the respective moment system by Theorem 3.3. Moreover, the solutions are the start points of the homotopy

$$(3.6) \quad H(\mu, \sigma^2; t) := \begin{cases} (1-t)\gamma g_1 + t f_1^k = 0 \\ \vdots \\ (1-t)\gamma g_{2k} + t f_{2k}^k = 0. \end{cases}$$

where f_i^k are defined as in (3.1). By Lemma 3.9 the result follows. \square

Corollary 3.10 bypasses the computational bottleneck associated with polyhedral homotopy methods. Therefore, the proofs of each theorem give the number of complex solutions to the corresponding variety *and* provide an efficient way to find all of them.

Example 3.11. Consider (3.1) when $k = 2$ and $\bar{\lambda} = (1/2, 1/2)$. Here we have

$$\begin{aligned} f_1^2(\mu, \sigma^2, \bar{\lambda}) &= \frac{1}{2}\mu_1 + \frac{1}{2}\mu_2 - \bar{m}_1 \\ f_2^2(\mu, \sigma^2, \bar{\lambda}) &= \frac{1}{2}(\mu_1^2 + \sigma_1^2) + \frac{1}{2}(\mu_2^2 + \sigma_2^2) - \bar{m}_2 \\ f_3^2(\mu, \sigma^2, \bar{\lambda}) &= \frac{1}{2}(\mu_1^3 + 3\mu_1\sigma_1^2) + \frac{1}{2}(\mu_2^3 + 3\mu_2\sigma_2^2) - \bar{m}_3 \\ f_4^2(\mu, \sigma^2, \bar{\lambda}) &= \frac{1}{2}(\mu_1^4 + 6\mu_1^2\sigma_1^2 + 3\sigma_1^4) + \frac{1}{2}(\mu_2^4 + 6\mu_2^2\sigma_2^2 + 3\sigma_2^4) - \bar{m}_4 \end{aligned}$$

In this case we consider the start system:

$$\begin{aligned} g_1 &= \mu_1 - 10, & g_2 &= \sigma_1^2 - 12, \\ g_3 &= \mu_2^3 - 27, & g_4 &= \sigma_2^4 - 4. \end{aligned}$$

This gives six start solutions of the form $(\mu_1, \sigma_1^2, \mu_2, \sigma_2^2)$:

$$\begin{array}{lll}
(10, 12, \eta \cdot 3, 2), & (10, 12, \eta^2 \cdot 3, 2), & (10, 12, 3, 2) \\
(10, 12, \eta \cdot 3, -2), & (10, 12, \eta^2 \cdot 3, -2), & (10, 12, 3, -2)
\end{array}$$

where η is a primitive third root of unity. We chose integers as the coefficients for ease of exposition. In practice, random complex numbers with norm close to one are used as the coefficients.

For the λ -weighted model discussed in Section 3.1, the Bézout bound of (3.1) is $(2k)!$ but Theorem 3.3 showed that the mixed volume is $(2k-1)!!k!$. The limit of

$$\frac{(2k)!}{(2k-1)!!k!}$$

tends to infinity as $k \rightarrow \infty$, showing that for large k , our start system is significantly better than a total degree one.

Remark 3.12. For the λ -weighted homoscedastic model discussed in Section 3.2, Theorem 3.7 shows that there are at most $\frac{(k+1)!}{2}$ solutions even though the Bézout bound is $(k+1)!$. In this case using our start system, one would track half as many paths as a total degree start system would.

3.4. Means unknown. A final case of interest is the λ -weighted, known variance model. This is where only the means are unknown. This set up was considered in high dimensions in [13, 16, 50].

We consider the moment system

$$(3.7) \quad f_1^k(\mu, \bar{\sigma}_1^2, \bar{\lambda}) = \cdots = f_k^k(\mu, \bar{\sigma}_k^2, \bar{\lambda}) = 0$$

where $\bar{\lambda}$ are known mixing coefficients, $\bar{\sigma}_\ell^2$ is a known variance, and f_ℓ^k , $\ell \in [k]$ is as defined in (2.3). In this set up, the unknowns are μ_ℓ , $\ell \in [k]$.

Theorem 3.13 (Means unknown). Fix $\bar{\lambda} \in \Delta_{k-1}$, and let Θ parameterize the $\bar{\lambda}$ -weighted known variance Gaussian k -mixture model. For $(\bar{m}_1, \dots, \bar{m}_k) \in V_\phi$, the number of complex solutions to (3.7) is at most $k!$ where the inequality is tight.

Proof. First we observe that (3.7) for generic moments (as defined by Equation (2.10)) has finitely many solutions by the same arguments in the proof of Proposition 3.2 and Proposition 3.6. This proves the first part of the theorem.

It follows from the Bézout bound that there are at most $k!$ complex solutions to (3.7). We now show that this bound is achieved with equality for generic moments by giving parameter values where there are precisely $k!$ solutions.

Consider $\bar{\lambda}_\ell = \frac{1}{k}$, $\bar{\sigma}_\ell^2 = 1$ and $\bar{m}_\ell = \sum_{\ell=1}^k \frac{1}{k} M_\ell(1, \ell)$ for $\ell \in [k]$. It is clear that this has a solution of the form $\mu = (1, 2, \dots, k)$. Further, by the same induction argument involving the Jacobian of (3.7) referenced above, there are finitely many solutions for this set of parameters. We observe that in this case our solution set has the typical label-swapping symmetry. This shows that any action by the symmetric group on

k letters, S_k , on any solution is also a solution. Therefore, there are $k!$ solutions to (3.7) in this case, namely $\{\sigma \cdot (1, 2, \dots, k) : \sigma \in S_k\}$. \square

Corollary 3.14 (Equal mixture weights and variances). A Gaussian k -mixture model with uniform mixing coefficients and known and equal variances is identifiable up to label-swapping symmetry using moments $(m_1, \dots, m_k) \in V_\phi$.

As a consequence of Theorem 3.13, when only the means are unknown the Bézout bound is equal to the BKK bound. In this case using polyhedral homotopy gives no advantage to total degree.

As discussed in Corollary 3.14, when the mixture weights and variances are known and equal the standard label-swapping symmetry observed with mixture models gives only one solution up to symmetry. Tracking a single path from a total degree start system, this one solution is easy to find.

Example 3.15. When $k = 2$, $\lambda_1 = \lambda_2 = \frac{1}{2}$ and $\sigma_1^2 = \sigma_2^2 = \sigma^2$ is a known parameter, we can symbolically solve the corresponding moment system and see that up to symmetry

$$\mu_1 = \bar{m}_1 - \sqrt{-\bar{m}_1^2 + \bar{m}_2 - \sigma^2}, \quad \mu_2 = \bar{m}_1 + \sqrt{-\bar{m}_1^2 + \bar{m}_2 - \sigma^2}.$$

This shows that so long as $-\bar{m}_1^2 + \bar{m}_2 - \sigma^2 > 0$ we are guaranteed to get something statistically meaningful. A picture of that region in \mathbb{R}^3 is shown in Figure 4.




FIGURE 4. Region in the space of parameters $\bar{m}_1, \bar{m}_2, \sigma^2$ where there are statistically meaningful solutions for $k = 2$ mixture model with unknown means and $\lambda_1 = \lambda_2 = \frac{1}{2}$.

3.5. A bridge from infinite to finite samples. While the theoretical setup assumes we are working with population moments m_1, m_2, \dots, m_r , in practice one only has access to sample moments \bar{m}_r from data. It is thus natural to wonder how much

accuracy one gains or loses depending on the sample size N , and we make a couple of notes in this regard.

Since $\mathbb{E}[\overline{m}_r] = m_r$, by Chebyshev's inequality we have that for all $\epsilon > 0$,

$$(3.8) \quad \mathbb{P}(|\overline{m}_r - m_r| \geq \epsilon) \leq \frac{\text{Var}(\overline{m}_r)}{\epsilon^2} = \frac{m_{2r} - m_r^2}{N\epsilon^2},$$

Thus, as a rule, if we would like to have $\mathbb{P}(|\overline{m}_r - m_r| \geq \epsilon) \leq \delta$, for some (small) $\delta > 0$, we need the sample size to be $N \geq \frac{m_{2r} - m_r^2}{\epsilon^2 \delta}$.

In the Gaussian mixture setting, the moment m_r is a polynomial of degree r in the parameters μ_ℓ, σ_ℓ , $\ell \in [k]$, so that the value of m_r varies roughly on the order of $\max_{\ell \in [k]} \{|\mu_\ell|^r, \sigma_\ell^{2r}\}$. Bounding all μ_ℓ, σ_ℓ^2 parameters by a constant C , the numerator grows roughly like C^{2r} , giving a sample complexity of $O(\frac{C^{2r}}{\epsilon^2 \delta})$ to estimate the Gaussian mixture moments within $\pm\epsilon$ with probability $1 - \delta$.

Now we make a note on how the approximation of the moments affects the estimated parameters. Let $\overline{m} \in \mathbb{R}^{2k}$ be the vector of first $2k$ moments of a $\overline{\lambda}$ -weighted Gaussian mixture density with parameters $\theta^* = (\mu^*, (\sigma^*)^2)$. Let $F = (f_1^k, \dots, f_{2k}^k)$ be polynomials in the moment system defined in (2.3). Then $F(\theta^*) = 0$ and for population moments $m^* \in \mathbb{R}^{2k}$ a solution to $F(\theta) = m^* - \overline{m}$ yields an estimate for a $\overline{\lambda}$ -weighted Gaussian mixture density with first $2k$ moments m^* .

It follows from multivariate Taylor expansion that a first order approximation yields

$$(\theta^* - \hat{\theta}) \approx -J^{-1}(\hat{\theta}) \cdot (m^* - \overline{m})$$

where $J^{-1}(\hat{\theta})$ denotes the inverse of the Jacobian matrix of F evaluated at $\hat{\theta}$. Thus, the distance between $\hat{\theta}$ and θ^* is controlled by the Jacobian $J(\hat{\theta})$ of F in the following way:

$$\|\theta^* - \hat{\theta}\|_2 \approx \|J(\hat{\theta})^{-1}(m^* - \overline{m})\|_2.$$

Here it is crucial to note that the Jacobian of F , which appears in (3.2), does not depend on the choice of moments m or \overline{m} since they do not interact with the parameters $\theta = (\mu, \sigma^2)$. For generic moments \overline{m} , the associated $\hat{\theta}$ will yield an invertible Jacobian, and the farther away from \mathcal{B} we can expect the parameters estimated using the sample moments will be close to the population parameters.

4. PARAMETER RECOVERY FOR HIGH DIMENSIONAL GAUSSIAN MIXTURE MODELS

Section 3 gives upper bounds on the number of solutions to the moment equations for Gaussian mixture models where some parameters of the model are assumed known. Using homotopy continuation algorithms we can efficiently perform density estimation in these cases. We now use our results to do density estimation on Gaussian mixture models in high dimensions.

4.1. High-dimensional Gaussian mixture models. To set notation, recall a random variable $X \in \mathbb{R}^n$ is distributed as a *multivariate Gaussian* with mean $\mu \in \mathbb{R}^n$

and symmetric positive definite covariance matrix $\Sigma \in \mathbb{R}^{n \times n}$, if it has density

$$f_X(x_1, \dots, x_n | \mu, \Sigma) = ((2\pi)^n \det(\Sigma))^{-1/2} \exp\left(-\frac{1}{2}(x - \mu)^T \Sigma^{-1}(x - \mu)\right).$$

We denote $X \sim \mathcal{N}(\mu, \Sigma)$.

A random variable is distributed as the *mixture of k multivariate Gaussians* if it is the convex combination of k multivariate Gaussian densities. It has probability density function

$$f_X(x_1, \dots, x_n | \lambda_\ell, \mu_\ell, \Sigma_\ell)_{\ell=1, \dots, k} = \sum_{\ell=1}^k \lambda_\ell f_{X_\ell}(x_1, \dots, x_n | \mu_\ell, \Sigma_\ell)$$

where $(\lambda_1, \dots, \lambda_k) \in \Delta_{k-1}$, $\mu_\ell \in \mathbb{R}^n$, and $\Sigma_\ell \in \mathbb{R}^{n \times n}$ is symmetric and positive definite for $\ell \in [k]$. Here we write $X \sim \sum_{\ell=1}^k \lambda_\ell \mathcal{N}(\mu_\ell, \Sigma_\ell)$.

Let $f_X : \mathbb{R}^n \rightarrow \mathbb{R}$ be the probability density function of a random vector $X = (X_1, \dots, X_n)$. We denote, the *(i_1, \dots, i_n) -th moment* of X by

$$m_{i_1, \dots, i_n} = \mathbb{E}[X_1^{i_1} \cdots X_n^{i_n}] = \int_{\mathbb{R}^n} x_1^{i_1} \cdots x_n^{i_n} f_X(x_1, \dots, x_n) dx_1 \cdots dx_n$$

where $i_s \geq 0$ for all $s \in [n]$ and the non-negative integer $i_1 + \dots + i_n = d$ is the *order* of m_{i_1, \dots, i_n} .

We express m_{i_1, \dots, i_n} as a polynomial in the parameters of a Gaussian mixture using the moment generating function. These are given by the identity:

$$(4.1) \quad \sum_{i_1, \dots, i_n \geq 0} \frac{m_{i_1, \dots, i_n}}{i_1! \cdots i_n!} t_1^{i_1} \cdots t_n^{i_n} = \sum_{\ell=1}^k \lambda_\ell \exp(t_1 \mu_{\ell 1} + \dots + t_n \mu_{\ell n}) \cdot \exp\left(\frac{1}{2} \sum_{i,j=1}^n \sigma_{\ell ij} t_i t_j\right).$$

Using Taylor's formula, we can expand the left side of (4.1) and equate coefficients of each side to get m_{i_1, \dots, i_n} . As in the univariate case,

$$(4.2) \quad m_{i_1, \dots, i_n} = \lambda_1 M_{i_1, \dots, i_n}(\mu_1, \Sigma_1) + \dots + \lambda_k M_{i_1, \dots, i_n}(\mu_k, \Sigma_k)$$

where $M_{i_1, \dots, i_n}(\mu, \Sigma)$ denotes the (i_1, \dots, i_n) -th moment of a multivariate Gaussian random variable $X \sim \mathcal{N}(\mu, \Sigma)$.

Example 4.1. Consider a Gaussian mixture model with $k = 2$ components in \mathbb{R}^2 . Here we have $X \sim \lambda_1 \mathcal{N}(\mu_1, \Sigma_1) + \lambda_2 \mathcal{N}(\mu_2, \Sigma_2)$ where

$$\begin{aligned} \mu_1 &= \begin{pmatrix} \mu_{11} \\ \mu_{12} \end{pmatrix}, & \Sigma_1 &= \begin{pmatrix} \sigma_{111} & \sigma_{112} \\ \sigma_{112} & \sigma_{122} \end{pmatrix}, \\ \mu_2 &= \begin{pmatrix} \mu_{21} \\ \mu_{22} \end{pmatrix}, & \Sigma_2 &= \begin{pmatrix} \sigma_{211} & \sigma_{212} \\ \sigma_{212} & \sigma_{222} \end{pmatrix}. \end{aligned}$$

Observe that our convention is to use the first index of μ and σ to identify the component of the mixture. The moments up to order three are

$$\begin{aligned}
(4.3) \quad & m_{00} = \lambda_1 + \lambda_2 \\
& m_{10} = \lambda_1 \mu_{11} + \lambda_2 \mu_{21} \\
& m_{01} = \lambda_1 \mu_{12} + \lambda_2 \mu_{22} \\
& m_{20} = \lambda_1 (\mu_{11}^2 + \sigma_{111}) + \lambda_2 (\mu_{21}^2 + \sigma_{211}) \\
& m_{11} = \lambda_1 (\mu_{11} \mu_{12} + \sigma_{112}) + \lambda_2 (\mu_{21} \mu_{22} + \sigma_{212}) \\
& m_{02} = \lambda_1 (\mu_{12}^2 + \sigma_{122}) + \lambda_2 (\mu_{22}^2 + \sigma_{222}) \\
& m_{30} = \lambda_1 (\mu_{11}^3 + 3\mu_{11}\sigma_{111}) + \lambda_2 (\mu_{21}^3 + 3\mu_{21}\sigma_{211}) \\
& m_{21} = \lambda_1 (\mu_{11}^2 \mu_{12} + 2\mu_{11}\sigma_{112} + \mu_{12}\sigma_{111}) + \lambda_2 (\mu_{21}^2 \mu_{22} + 2\mu_{21}\sigma_{212} + \mu_{22}\sigma_{211}) \\
& m_{12} = \lambda_1 (\mu_{11} \mu_{12}^2 + \mu_{11}\sigma_{122} + 2\mu_{12}\sigma_{112}) + \lambda_2 (\mu_{21} \mu_{22}^2 + \mu_{21}\sigma_{222} + 2\mu_{22}\sigma_{212}) \\
& m_{03} = \lambda_1 (\mu_{12}^3 + 3\mu_{12}\sigma_{122}) + \lambda_2 (\mu_{22}^3 + 3\mu_{22}\sigma_{222}).
\end{aligned}$$

The $m_{0,0,\dots,0,i_s,0,\dots,0}$ -th moment is the same as the i_s -th order moment for the univariate Gaussian mixture model $\sum_{\ell=1}^k \lambda_\ell \mathcal{N}(\mu_{\ell s}, \sigma_{\ell ss})$. This observation is key to our proposed density estimation algorithm in Section 4.2 and it follows from the property that marginal distributions of a Gaussian are Gaussian themselves.

4.2. Dimension reduction and parameter recovery algorithm. We propose an algorithm for parameter estimation of multivariate Gaussian densities using the method of moments. The main idea is that if we use higher order moment equations, parameter estimation for multivariate Gaussian mixture models reduces to multiple instances of density estimation for univariate Gaussian mixture models. In this subsection we present Algorithm 1 to describe how to recover the parameters of a $\bar{\lambda}$ -weighted multivariate Gaussian.

Remark 4.2. There are many possible choices for the input $\bar{\mathbf{m}}_2$. For instance, if $k \leq \binom{n}{2}$, then one can use second order moments of the form $m_{e_i+e_j}$ for $i, j \in [n]$. If $k \gg n$, then one will have to use higher order moments to recover all off diagonal entries. In either case, one could use moments of the form $m_{te_i+e_j}$ for $t \in [k]$, $i, j \in [n]$.

Example 4.3. Suppose $X \sim 0.25 \cdot \mathcal{N}(\mu_1, \Sigma_1) + 0.75 \cdot \mathcal{N}(\mu_2, \Sigma_2)$ as in Example 4.1.

The first steps of Algorithm 1 use the sample moments $\bar{\mathbf{m}}_1$. Here we use the input

$$\begin{aligned}
\bar{\mathbf{m}}_{1,1} &= [\bar{m}_{10}, \bar{m}_{20}, \bar{m}_{30}, \bar{m}_{40}, \bar{m}_{50}] = [-0.25, 2.75, -1.0, 22.75, -6.5] \\
\bar{\mathbf{m}}_{1,2} &= [\bar{m}_{01}, \bar{m}_{02}, \bar{m}_{03}, \bar{m}_{04}, \bar{m}_{05}] = [2.5, 16.125, 74.5, 490.5625, 2921.25].
\end{aligned}$$

In Step 2, for the unknowns $\mu_{11}, \mu_{21}, \sigma_{111}, \sigma_{211}$ we solve (3.1) with $\bar{\mathbf{m}}_{1,1}$,

$$\begin{aligned}
-0.25 &= 0.25 \cdot \mu_{11} + 0.75 \cdot \mu_{21} \\
2.75 &= 0.25 \cdot (\mu_{11}^2 + \sigma_{111}) + 0.75 \cdot (\mu_{21}^2 + \sigma_{211}) \\
-1 &= 0.25 \cdot (\mu_{11}^3 + 3\mu_{11}\sigma_{111}) + 0.75 \cdot (\mu_{21}^3 + 3\mu_{21}\sigma_{211}) \\
22.75 &= 0.25 \cdot (\mu_{11}^4 + 6\mu_{11}^2\sigma_{111} + 3\sigma_{111}^2) + 0.75 \cdot (\mu_{21}^4 + 6\mu_{21}^2\sigma_{211} + 3\sigma_{211}^2),
\end{aligned}$$

Algorithm 1: Parameter recovery for Mixtures of $\bar{\lambda}$ -Weighted Multivariate Gaussians

Input: Mixing coefficients $\bar{\lambda}_\ell$, $\ell \in [k]$ and moments:

$$\bar{\mathbf{m}}_{1,i} := \{\bar{m}_{e_i}, \dots, \bar{m}_{De_i}\} \quad i \in [n] \quad \text{and} \quad D = 3k + 2.$$

$$\bar{\mathbf{m}}_2 := \{\bar{m}_{a_1}, \dots, \bar{m}_{a_N} : a_j \in \mathbb{N}^n\}$$

from a multivariate Gaussian mixture model:

$$\bar{\lambda}_1 \mathcal{N}(\mu_1, \Sigma_1) + \dots + \bar{\lambda}_k \mathcal{N}(\mu_k, \Sigma_k)$$

where $N = \frac{k}{2}(n^2 - n)$ and $\bar{\mathbf{m}}_2$ is a set of moments such that the polynomials m_{a_j} as in (4.1) are linear in the off diagonal entries of Σ_ℓ for $\ell \in [k]$.

Output: Parameters $\mu_\ell \in \mathbb{R}^n$, $\Sigma_\ell \succ 0$ for $\ell \in [k]$ such that $\bar{\mathbf{m}}_{1,i}, \bar{\mathbf{m}}_2$ are the moments of $\sum_{\ell=1}^k \bar{\lambda}_\ell \mathcal{N}(\mu_\ell, \Sigma_\ell)$.

- 1 **for** $i = 1, \dots, n$ **do**
 - 2 With mixing coefficients $\bar{\lambda}_\ell$ and moments $\bar{\mathbf{m}}_{1,i}$, find all solutions to (3.1) with $\mu_{\ell i} \in \mathbb{R}$ and $\sigma_{\ell ii} > 0$ for all $\ell \in [k]$.
 - 3 Select the solution consistent with the moment \bar{m}_{De_i} .
 - 4 Form a linear system of N equations in N unknowns by substituting $\mu_{\ell i}$ and $\sigma_{\ell ii}$ into $m_{a_j} = \bar{m}_{a_j}$.
 - 5 Solve the linear system for $\sigma_{\ell ij}$, $i \neq j$.
 - 6 Return μ_ℓ, Σ_ℓ , $\ell \in [k]$.
-

We find that there are two statistically meaningful solutions:

$$(\mu_{11}, \mu_{21}, \sigma_{111}, \sigma_{211}) = (-1, 0, 1, 3)$$

$$(\mu_{11}, \mu_{21}, \sigma_{111}, \sigma_{211}) = (0.978, -0.659, 2.690, 2.017).$$

The first solution has $m_{50} = -6.5$ and the second has $m_{50} = -1.55$ so in Step 3 we select the first solution. Proceeding with the second iteration of the for-loop in Step 1, we solve for the unknowns $\mu_{12}, \mu_{22}, \sigma_{122}, \sigma_{222}$ using the sample moments $\bar{\mathbf{m}}_{1,2}$ to obtain

$$(\mu_{12}, \mu_{22}, \sigma_{122}, \sigma_{222}) = (-2, 4, 2, 3.5).$$

Finally, to recover the off diagonal entries of Σ_1, Σ_2 , with sample moments

$$\bar{\mathbf{m}}_2 = [\bar{m}_{11}, \bar{m}_{21}] = [0.8125, 7.75],$$

we solve the linear system

$$0.8125 = 0.25 \cdot (2 + \sigma_{112}) + 0.75 \cdot \sigma_{212} \quad 7.75 = 0.25 \cdot (-4 - 2 \cdot \sigma_{112}) + 9$$

to find $(\sigma_{112}, \sigma_{212}) = (0.5, 0.25)$. We estimate that our samples came from density

$$0.25 \cdot \mathcal{N}\left(\begin{bmatrix} -1 \\ -2 \end{bmatrix}, \begin{bmatrix} 1 & 0.5 \\ 0.5 & 2 \end{bmatrix}\right) + 0.75 \cdot \mathcal{N}\left(\begin{bmatrix} 0 \\ 4 \end{bmatrix}, \begin{bmatrix} 3 & 0.25 \\ 0.25 & 3.5 \end{bmatrix}\right).$$

Using the homotopy continuation methods discussed in Section 3.3, we can efficiently solve Step 2 of Algorithm 1. Using these continuation methods for fixed k , the number of homotopy paths we need to track is $\mathcal{O}(n)$. One observation for implementation is that Steps 2 can be performed in parallel.

In order to prove Theorem 4.5 we need an additional result regarding rational identifiability of Gaussian k mixture models. The proof of this result requires additional results from algebraic geometry. We state the theorem here but defer the proof and relevant background to the appendix.

Theorem 4.4. Mixtures of k univariate Gaussians are rationally identifiable from moments m_1, \dots, m_{3k+2} .

Theorem 4.5. For a generic $\bar{\lambda}$ -Gaussian k mixture model in \mathbb{R}^n ,

$$\bar{\lambda}_1 \mathcal{N}(\mu_1, \Sigma_1) + \dots + \bar{\lambda}_k \mathcal{N}(\mu_k, \Sigma_k),$$

Algorithm 1 is guaranteed to recover the parameters μ_ℓ and diagonal entries of Σ_ℓ , $\ell \in [k]$. Furthermore, if $\bar{\mathbf{m}}_2$ is chosen as in Remark 4.2, then the off-diagonal entries of Σ_ℓ , $\ell \in [k]$ are guaranteed to be recovered as well.

Proof. From Theorem 3.3, in each iteration of the for-loop in Algorithm 1, we find finitely many prospective choices for each entry of the means μ_ℓ and diagonal entries of Σ_ℓ , $\ell \in [k]$. There is a unique one that agrees with all moments, due to the identifiability result in Theorem 4.4 in the appendix. In other words, after the for-loop we uniquely recover all mean vectors μ_ℓ and the diagonal entries of Σ_ℓ , $\ell \in [k]$. It suffices to show that the linear system in Step 5 of Algorithm 1 has a unique solution.

Fix $i, j \in [n]$, $i \neq j$, we show that using moment equations $\bar{\mathbf{m}}_2 = \{m_{te_i+e_j}\}$ for $t \in [k]$, the corresponding linear system in the variables $\sigma_{\ell ij}$ for $\ell \in [k]$ generically has a unique solution.

By [48, Equation 13] and the linearity of expectation, using the notation in (4.2) we have

$$m_{te_i+e_j} = \sum_{\ell=1}^k \mu_{\ell j} \cdot M_{te_i}(\mu_\ell, \Sigma_\ell) + t \sigma_{\ell ij} \cdot M_{(t-1)e_i}(\mu_\ell, \Sigma_\ell)$$

After substituting the values we have already obtained into the polynomials $m_{te_i+e_j}$, we get a linear system with a $k \times k$ coefficient matrix that we denote by A . Indexing the columns of A by $\sigma_{1ij}, \dots, \sigma_{kij}$, the pq -th entry of A is $p \cdot M_{(p-1)e_i}(\mu_q, \Sigma_q)$. Observe that the entries of A are polynomials in the parameters $\lambda_\ell, \mu_{\ell i}, \mu_{\ell j}$ for $\ell \in [k]$ and $i, j \in [n]$. Moreover, the entries in row p of A are polynomials of degree $p-1$ and the entries in column q of A contains the subset of parameters corresponding to the q th mixture component, $\lambda_q, \mu_{qi}, \mu_{qj}$, $i \in [n]$. Therefore, for generic values of $\mu_\ell, \lambda_\ell, \Sigma_\ell$, these rows will not be linearly dependent so the corresponding linear system is full rank, meaning there is a unique solution which recovers the off diagonal entries of $\sigma_{\ell ij}$, $\ell \in [k]$. □

Remark 4.6. Observe in the proof of Theorem 4.5 the linear system solved in Step 5 can be decomposed into $\frac{n(n-1)}{2}$ linear systems of size $k \times k$. Therefore, not only can

the for-loop in Algorithm 1 be parallelized, but solving for the off diagonal entries of $\Sigma_1, \dots, \Sigma_k$ can be as well.

Remark 4.7. Although Algorithm 1 uses higher order moments, this order does not exceed what we would need for the univariate case. In addition, it has been shown that there exist algebraic dependencies among lower order moment equations complicating the choice of which moment system to consider [3].

On the other hand, observe that Algorithm 1 can be specialized to both the λ -weighted homoscedastic and λ -weighted known variance models by applying the results of Theorem 3.7 and Theorem 3.13 to the framework outlined in Section 4.2. In each case, the number of homotopy paths tracked will be $\frac{(k+1)!}{2} + (n-1)k!$ and $nk!$ respectively. We note again that both situations give algorithms where the number of homotopy paths scales linearly in n .

4.3. Uniform mixtures with equal variances. In this section, we consider the special case of estimating the mixture of k Gaussians in \mathbb{R}^n where all of the means $\mu_i \in \mathbb{R}^n$ are unknown, each $\lambda_i = \frac{1}{k}$ and each covariance matrix $\Sigma_i \in \mathbb{R}^{n \times n}$ is equal and known. This is the set up in [13, 16, 36, 50].

Recall from Corollary 3.14 that in one dimension, there is generically a unique solution up to symmetry to (3.7) that can be found efficiently. We use this fact in Step 1 of Algorithm 2. In Step 2 there are other choices for sample moments that will give a square linear system. Some of these choices involve sample moments of lower order.

Algorithm 2: Density Estimation for Uniform Mixtures of Multivariate Gaussians with Equal Covariances

Input: The set of sample moments:

$$\begin{aligned}\bar{\mathbf{m}}_1 &:= \{\bar{m}_{e_1}, \dots, \bar{m}_{ke_1}\} \\ \bar{\mathbf{m}}_i &:= \{\bar{m}_{e_i}, \bar{m}_{e_1+e_i}, \bar{m}_{2e_1+e_i}, \dots, \bar{m}_{(k-1)e_1+e_i}\}, \quad 2 \leq i \leq n\end{aligned}$$

that are the moments to multivariate Gaussian mixture model:

$$\frac{1}{k}\mathcal{N}(\mu_1, \bar{\Sigma}) + \dots + \frac{1}{k}\mathcal{N}(\mu_k, \bar{\Sigma}).$$

Output: Parameters $\mu_\ell \in \mathbb{R}^n$, such that $\bar{\mathbf{m}}_i$, $i \in [n]$, are the moments of distribution $\sum_{\ell=1}^k \frac{1}{k}\mathcal{N}(\mu_\ell, \bar{\Sigma})$

- 1 Using mixing coefficients $\lambda_\ell = \frac{1}{k}$ for $\ell \in [k]$ and sample moments $\bar{\mathbf{m}}_1$ solve (3.7) to obtain $\mu_{\ell 1} \in \mathbb{R}$.
 - 2 Using sample moments $\bar{\mathbf{m}}_i$ solve the $k \times k$ linear system in $\mu_{i1}, \dots, \mu_{ik}$ for $2 \leq i \leq n$.
-

Observe that Step 1 of Algorithm 2 requires *tracking a single homotopy path*. This is in contrast to Step 2 of Algorithm 1 in which one needs to track $(2k-1)!!k!$ homotopy paths to obtain all complex solutions. Further, Algorithm 2 requires solving a $k \times k$

linear system $n - 1$ times (Step 2). This is again in contrast to Algorithm 1 where one needs to solve a nonlinear polynomial system that tracks $(2k - 1)!!k!$ paths $n - 1$ times (Step 2). In both cases, we see that we need to solve n polynomial systems, where n is the dimension of the Gaussian mixture model.

If we consider the tracking of a single homotopy path as unit cost, we consider the number of paths tracked as the complexity of the algorithm (as is customary in numerical algebraic geometry). With this choice, Algorithm 2 tracks n homotopy paths, while Algorithm 1 tracks $(2k - 1)!!k!n$ paths and solves a square linear system of size $\frac{kn(n-1)}{2}$. This highlights how Algorithm 2 is highly efficient and in this case the method of moments up can be effective for large n and k .

4.4. Beyond known weights. Algorithm 1 can be adapted to the case when the mixing coefficients λ_ℓ , $\ell \in [k]$ are not known a priori. One can use the method of moments for general Gaussian mixture models to recover them. When $k = 2, 3, 4$ the number of solutions to the corresponding moment system are given in [39, 2, 4] respectively. Since the number of complex solutions is known in each of these cases, state of the art polynomial system solvers can exploit the label swapping symmetry to find all solutions quickly, and select the best statistically meaningful one.

n	10	100	1,000	10,000	100,000
Time (s)	0.17	0.71	6.17	62.05	650.96
Error	7.8×10^{-15}	4.1×10^{-13}	5.7×10^{-13}	2.9×10^{-11}	1.8×10^{-9}
Normalized Error	1.9×10^{-16}	1.0×10^{-15}	1.4×10^{-16}	7.3×10^{-16}	4.5×10^{-15}

TABLE 1. Average running time and numerical error running Algorithm 1 on a mixture of 2 Gaussians in \mathbb{R}^n . The error is $\epsilon = \|v - \hat{v}\|_2$ where $v \in \mathbb{R}^{4n+2}$ is a vector of the true parameters and \hat{v} is a vector of the estimates. The normalized error is $\epsilon/(4n + 2)$.

n	10	100	1,000	10,000	100,000
Time (s)	4.71	10.87	73.74	845.55	8291.84
Error	3.6×10^{-13}	4.6×10^{-12}	1.3×10^{-10}	4.6×10^{-10}	9.6×10^{-9}
Normalized Error	1.1×10^{-14}	1.5×10^{-14}	4.2×10^{-14}	1.5×10^{-14}	3.2×10^{-14}

TABLE 2. Average running time and numerical error running Algorithm 1 on a mixture of 3 Gaussians in \mathbb{R}^n . The error is $\epsilon = \|v - \hat{v}\|_2$ where $v \in \mathbb{R}^{6n+3}$ is a vector of the true parameters and \hat{v} is a vector of the estimates. The normalized error is $\epsilon/(6n + 3)$.

We perform numerical experiments by running Algorithm 1 on randomly generated Gaussian mixture models with diagonal covariance matrices. We do not assume

the mixing coefficients are known a priori, so we solve for them first and then run Algorithm 1. We use `HomotopyContinuation.jl` to do all of the polynomial system solving [10]. The average running time and error for $k = 2$ are given in Table 1 and for $k = 3$ in Table 2. Overall, we see that the error incurred from doing homotopy continuation is negligible.

Acknowledgments. The authors would like to thank Fulvio Gesmundo and Nick Vannieuwenhoven for helpful discussions regarding Theorem 4.4, which took place within the Thematic Research Programme "Tensors in statistics, optimization and machine learning", University of Warsaw, Excellence Initiative – Research University and the Simons Foundation Award No. 663281 granted to the Institute of Mathematics of the Polish Academy of Sciences for the years 2021-2023. The authors would like to thank the anonymous referees, an Associate Editor and the Editor for their constructive comments that improved the quality of this paper.

5. APPENDIX

In this section we make significant progress on an open problem regarding moment identifiability of Gaussian mixtures. This improvement is the one that justifies the number of moments needed in the input of Algorithm 1.

Any univariate Gaussian k -mixture model is uniquely identifiable from at most its first $4k - 2$ moments m_1, \dots, m_{4k-2} [29] and algebraically identifiable from the moments m_1, \dots, m_{3k-1} [3, Theorem 1]. The number of moments needed for Gaussian k -mixtures to be rationally identifiable is therefore somewhere between $3k - 1$ and $4k - 2$.

We now show that indeed the correct asymptotic order of moments needed for rational identifiability grows like $3k$ and not like $4k$. The proof relies on concepts and recent results from algebraic geometry. We give a concise argument with detailed references.

Remark 5.1. For some historical background, note that when $k = 2$, the two expressions $3k$ and $4k - 2$ match at six moments. The unique identifiability of a 2-mixture from m_1, \dots, m_6 was implicitly assumed by Pearson in [39] and first shown to be true by Lazard in [32]. Regarding rationally identifiable, the number of moments needed for Gaussian k -mixtures to be conjectured [2, Conjecture 3] to be precisely $3k$.

Let \mathbb{P}^d be the projective space of dimension d whose coordinates are all $d + 1$ moments m_i with $i = 0, 1, \dots, d$. The main geometric object, first introduced in [2], is the *Gaussian moment surface* $\mathcal{G}_{1,d}$ in \mathbb{P}^d . It is the projective closure of the image of the parameterization $\mathbb{C}^2 \rightarrow \mathbb{C}^d$, $(\mu, \sigma^2) \mapsto (m_1, \dots, m_d)$ where $m_i = M_i(\mu, \sigma^2)$ with the right-hand side coming from (2.1). If we restrict the parameterization to $\mathbb{R} \times \mathbb{R}_{>0}$, then we have the image consisting of all univariate Gaussian moments up to order d .

The Gaussian moment surface $\mathcal{G}_{1,d}$ is an algebraic set in \mathbb{P}^d , also called a projective variety. This means it can be defined as the solution set to a system of homogeneous polynomials in m_0, \dots, m_d . As an example, for $d = 3$,

$$\mathcal{G}_{1,3} = \{(m_0, m_1, m_2, m_3) \in \mathbb{P}^3 : 2m_1^3 - 3m_0m_1m_2 + m_0^2m_3 = 0\}$$

is a surface in \mathbb{P}^3 , which is smooth everywhere except along the line defined by $m_0 = m_1 = 0$. Moreover, by [3, Lemma 4], for any $d \geq 3$, $\mathcal{G}_{1,d}$ is singular precisely on the line defined by

$$(5.1) \quad m_0 = m_1 = \dots = m_{d-2} = 0.$$

In addition, since $\mathcal{G}_{1,d}$ is the closure of the image of a parameterization by \mathbb{C}^2 , this projective variety is also irreducible, i.e., if any other closed algebraic sets A, B in \mathbb{P}^d such that $A \cup B = \mathcal{G}_{1,d}$ then either A or B equals $\mathcal{G}_{1,d}$. Now we are ready to prove a fact about the Gaussian moment surface $\mathcal{G}_{1,d}$.

Lemma 5.2. The Gaussian moment surface $\mathcal{G}_{1,d}$ has a non-degenerate Gauss map.

Proof. According to [22, Section 2] and [27, Theorem 3.4.6], if the Gauss map of a projective surface Z is degenerate, then Z must be one of the following:

- (a) a linearly embedded \mathbb{P}^2 ;

- (b) a cone over a curve;
- (c) the tangential variety to a curve.

We are not in case (a) because the degree of $\mathcal{G}_{1,d}$ is $\binom{d}{2} > 1$ for $d > 2$ by [2, Corollary 2]. In other words $\mathcal{G}_{1,d}$ is a nonlinear surface that cannot be linearly embedded into \mathbb{P}^2 . Case (b) is not possible for $\mathcal{G}_{1,d}$ either, by the proof of [3, Theorem 1] where great detail is given to exclude the cone over a curve scenario.

It remains to show we are not in case (c). If the surface $\mathcal{G}_{1,d}$ was a tangential variety to a curve, then it will contain said curve in its singular locus. Since the singular locus of $\mathcal{G}_{1,d}$ is a line (5.1), we must have $\mathcal{G}_{1,d}$ is the tangential variety to a line. This is a contradiction because the tangential variety of a line is the line itself and certainly not a surface. □

Secant varieties have a long history of appearing in applications and more recently in statistics. For a projective variety Z , by definition, the *kth secant variety* $\sigma_k(Z)$ is the Zariski closure of the union of the $(k-1)$ -planes spanned by collections of k points in Z .

Example 5.3. Let Z be the variety of $m \times n$ rank one matrices in the projective space \mathbb{P}^{mn-1} with coordinates z_{ij} being the (i, j) entry of the matrix. For $k \leq \min\{m, n\}$, the secant variety $\sigma_k(Z)$ consists of rank at most k matrices.

For our purposes, we are interested in the k th secant variety of the surface $\mathcal{G}_{1,d}$. Since the moment of a Gaussian k -mixture is a linear combination of the moments of each of the k components, points in $\sigma_k(\mathcal{G}_{1,d})$ represent the moments up to order d of a Gaussian k -mixture.

For $Z \subset \mathbb{P}^N$ the dimension of a k th secant variety of Z satisfies the inequality

$$(5.2) \quad \dim(\sigma_k(Z)) \leq \min\{k \cdot \dim(Z) + k - 1, N\}.$$

When the inequality is strict, we say that Z is *k-defective*. In geometric terms, a variety Z being k -defective means that there are *infinitely* many $(k-1)$ -planes passing through any k points in Z . On the other hand, we say Z is *k-identifiable* if there is a *unique* $(k-1)$ -plane that passes through k generic points in Z .

In the instance of (5.2) when $Z = \mathcal{G}_{1,d}$. By [3, Theorem 1], $\mathcal{G}_{1,d}$ is not k -defective, i.e.,

$$(5.3) \quad \dim(\sigma_k(\mathcal{G}_{1,d})) = \min\{3k - 1, d\}.$$

In the context of moment identifiability, from (5.3) it follows that Gaussian k -mixtures are algebraically identifiable from moments up to order $3k - 1$. Furthermore, if one has that $\mathcal{G}_{1,d}$ is k -identifiable, then we have rational identifiability from moments up to order d . We prove this for $d = 3k + 2$.

Theorem 5.4. Mixtures of k univariate Gaussians are rationally identifiable from moments m_1, \dots, m_{3k+2} .

Proof. We wish to apply [35, Theorem 1.5], which states that k -identifiability follows for irreducible and non-degenerate varieties $Z \subset \mathbb{P}^N$ of dimension n if the following conditions hold:

- (i) $(k+1)n + k \leq N$;
- (ii) Z has non-degenerate Gauss map;
- (iii) Z is not $(k+1)$ -defective.

We consider $Z = \mathcal{G}_{1,3k+2}$ which is $n = 2$ -dimensional in \mathbb{P}^{3k+2} so that $N = 3k+2$. The first item holds with equality as $(k+1)2 + k = N$. Item (ii) is the content of Lemma 5.2. Thus, it remains to prove item (iii). This follows from (5.2) and (5.3) by substituting $k+1$ for k to obtain

$$\dim(\sigma_{k+1}(\mathcal{G}_{1,3k+2})) = \min\{3(k+1) - 1, 3k+2\} = 3k+2,$$

thereby showing $\mathcal{G}_{1,3k+2}$ is not $(k+1)$ -defective. Then, by [35, Theorem 1.5], we have $\mathcal{G}_{1,3k+2}$ is k -identifiable. \square

REFERENCES

- [1] C. Améndola, M. Drton, and B. Sturmfels. Maximum likelihood estimates for Gaussian mixtures are transcendental. In *International Conference on Mathematical Aspects of Computer and Information Sciences*, pages 579–590. Springer, 2015.
- [2] C. Améndola, J.-C. Faugère, and B. Sturmfels. Moment varieties of Gaussian mixtures. *J. Algebr. Stat.*, 7(1):14–28, 2016.
- [3] C. Améndola, K. Ranestad, and B. Sturmfels. Algebraic identifiability of Gaussian mixtures. *Int. Math. Res. Not. IMRN*, (21):6556–6580, 2018.
- [4] C. Améndola, J. Lindberg, and J. I. Rodriguez. Solving parameterized polynomial systems with decomposable projections. *arXiv:1612.08807*, 2021.
- [5] H. Ashtiani, S. Ben-David, N. J. A. Harvey, C. Liaw, A. Mehrabian, and Y. Plan. Near-optimal sample complexity bounds for robust learning of Gaussian mixtures via compression schemes. *J. ACM*, 67(6):Art. 32, 42, 2020.
- [6] D. J. Bates, J. D. Hauenstein, A. J. Sommese, and C. W. Wampler. *Numerically solving polynomial systems with Bertini*, volume 25 of *Software, Environments, and Tools*. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2013.
- [7] M. Belkin and K. Sinha. Polynomial learning of distribution families. *SIAM J. Comput.*, 44(4):889–911, 2015.
- [8] D. N. Bernstein. The number of roots of a system of equations. *Funkcional. Anal. i Priložen.*, 9(3):1–4, 1975.
- [9] K. Blekas, A. Likas, N. Galatsanos, and I. Lagaris. A spatially constrained mixture model for image segmentation. *IEEE Transactions on Neural Networks*, 16(2):494–498, 2005.
- [10] P. Breiding and S. Timme. Homotopycontinuation.jl: A package for homotopy continuation in Julia. In *Mathematical Software – ICMS 2018*, pages 458–465. Springer International Publishing, 2018.

- [11] J. C. Butcher. *Numerical methods for ordinary differential equations*. John Wiley & Sons, Ltd., Chichester, 2003.
- [12] T. T. Cai, J. Ma, and L. Zhang. CHIME: clustering of high-dimensional Gaussian mixtures with EM algorithm and its optimality. *Ann. Statist.*, 47(3):1234–1267, 2019.
- [13] Y. Chen and X. Xi. Likelihood landscape and local minima structures of Gaussian mixture models. *arXiv:2009.13040*, 2020.
- [14] G. E. Collins. Quantifier elimination for real closed fields by cylindrical algebraic decomposition. In *Automata theory and formal languages (Second GI Conf., Kaiserslautern, 1975)*, pages 134–183. Lecture Notes in Comput. Sci., Vol. 33. 1975.
- [15] D. A. Cox, J. Little, and D. O’Shea. *Using algebraic geometry*, volume 185 of *Graduate Texts in Mathematics*. Springer, New York, second edition, 2005.
- [16] C. Daskalakis, C. Tzamos, and M. Zampetakis. Ten steps of EM suffice for mixtures of two Gaussians. In *Conference on Learning Theory*, pages 704–710, 2017.
- [17] A. Dempster, N. Laird, and D. Rubin. Maximum likelihood from incomplete data via the EM algorithm. *J. Roy. Statist. Soc. Ser. B*, 39(1):1–38, 1977.
- [18] B. Do and M. Ohsaki. Gaussian mixture model for robust design optimization of planar steel frames. *Struct. Multidiscip. Optim.*, 63(1):137–160, 2021.
- [19] M. Dyer, P. Gritzmann, and A. Hufnagel. On the complexity of computing mixed volumes. *SIAM J. Comput.*, 27(2):356–400, 1998.
- [20] G. Ewald. *Combinatorial convexity and algebraic geometry*, volume 168 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1996.
- [21] I. Goodfellow, Y. Bengio, and A. Courville. *Deep Learning*. MIT Press, 2016. <http://www.deeplearningbook.org>.
- [22] P. Griffiths and J. Harris. Algebraic geometry and local differential geometry. *Ann. Sci. École Norm. Sup. (4)*, 12(3):355–452, 1979.
- [23] L. P. Hansen. Large sample properties of generalized method of moments estimators. *Econometrica*, 50(4):1029–1054, 1982.
- [24] R. Hartshorne. *Algebraic geometry*. Springer-Verlag, New York-Heidelberg, 1977. Graduate Texts in Mathematics, No. 52.
- [25] D. Hosseinzadeh and S. Krishnan. Gaussian mixture modeling of keystroke patterns for biometric applications. *systems man and cybernetics*, 38(6):816–826, 2008.
- [26] B. Huber and B. Sturmfels. A polyhedral method for solving sparse polynomial systems. *Math. Comp.*, 64(212):1541–1555, 1995.
- [27] T. A. Ivey and J. M. Landsberg. *Cartan for beginners: differential geometry via moving frames and exterior differential systems*, volume 61 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2003.
- [28] A. T. Kalai, A. Moitra, and G. Valiant. Efficiently learning mixtures of two Gaussians. In *STOC’10—Proceedings of the 2010 ACM International Symposium on Theory of Computing*, pages 553–562. ACM, New York, 2010.

- [29] A. T. Kalai, A. Moitra, and G. Valiant. Disentangling Gaussians. *Communications of The ACM*, 55(2):113–120, 2012.
- [30] A. G. Khovanskii. Newton polyhedra, and the genus of complete intersections. *Funktsional. Anal. i Prilozhen.*, 12(1):51–61, 1978.
- [31] A. G. Kouchnirenko. Polyèdres de Newton et nombres de Milnor. *Invent. Math.*, 32(1):1–31, 1976.
- [32] D. Lazard. Injectivity of real rational mappings: the case of a mixture of two Gaussian laws. *Math. Comput. Simulation*, 67(1-2):67–84, 2004.
- [33] T. Y. Li. Numerical solution of multivariate polynomial systems by homotopy continuation methods. In *Acta numerica, 1997*, volume 6 of *Acta Numer.*, pages 399–436. Cambridge Univ. Press, Cambridge, 1997.
- [34] T. Y. Li and X. Wang. The BKK root count in \mathbf{C}^n . *Math. Comp.*, 65(216):1477–1484, 1996.
- [35] A. Massarenti and M. Mella. Bronowski’s conjecture and the identifiability of projective varieties, 2022.
- [36] S. Mei, Y. Bai, and A. Montanari. The landscape of empirical risk for nonconvex losses. *Ann. Statist.*, 46(6A):2747–2774, 2018.
- [37] A. Moitra and G. Valiant. Settling the polynomial learnability of mixtures of Gaussians. In *2010 IEEE 51st Annual Symposium on Foundations of Computer Science—FOCS 2010*, pages 93–102. IEEE Computer Soc., Los Alamitos, CA, 2010.
- [38] P. Mondal. *How Many Zeroes?: Counting Solutions of Systems of Polynomials Via Toric Geometry at Infinity*, volume 2. Springer Nature, 2021.
- [39] K. Pearson. Contributions to the mathematical theory of evolution. *Philosophical Transactions of the Royal Society A*, 185:71–110, 1894.
- [40] D. A. Reynolds. *A Gaussian mixture modeling approach to text-independent speaker identification*. PhD thesis, Georgia Institute of Technology, 1992.
- [41] A. Sanjeev and R. Kannan. Learning mixtures of arbitrary Gaussians. In *Proceedings of the thirty-third annual ACM symposium on Theory of computing*, pages 247–257, 2001.
- [42] F. Sha and L. Saul. Large margin Gaussian mixture modeling for phonetic classification and recognition. In *2006 IEEE International Conference on Acoustics Speech and Signal Processing Proceedings*, volume 1, pages 265–268, 2006.
- [43] S. Shirinkam, A. Alaeddini, and E. Gross. Identifying the number of components in Gaussian mixture models using numerical algebraic geometry. *J. Algebra Appl.*, 19(11):2050204, 21, 2020.
- [44] A. Singhal, P. Singh, B. Lall, and S. D. Joshi. Modeling and prediction of COVID-19 pandemic using Gaussian mixture model. *Chaos Solitons Fractals*, 138:110023, 8, 2020.
- [45] B. Sturmfels. *Solving systems of polynomial equations*, volume 97 of *CBMS Regional Conference Series in Mathematics*. Published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 2002.

- [46] H. Teicher. Identifiability of finite mixtures. *The Annals of Mathematical statistics*, pages 1265–1269, 1963.
- [47] L. Wasserman. *All of statistics*. Springer Texts in Statistics. Springer-Verlag, New York, 2004. A concise course in statistical inference.
- [48] R. Willink. Normal moments and hermite polynomials. *Statistics & Probability Letters*, 73(3):271–275, 2005.
- [49] Y. Wu and P. Yang. Optimal estimation of Gaussian mixtures via denoised method of moments. *Ann. Statist.*, 48(4):1981–2007, 2020.
- [50] J. Xu, D. J. Hsu, and A. Maleki. Global analysis of Expectation Maximization for mixtures of two Gaussians. In *Advances in Neural Information Processing Systems*, volume 29, pages 2676–2684, 2016.
- [51] S. J. Yakowitz and J. D. Spragins. On the identifiability of finite mixtures. *The Annals of Mathematical Statistics*, pages 209–214, 1968.
- [52] Y. Zhang, M. Alder, and R. Togneri. Using Gaussian mixture modeling in speech recognition. In *Proceedings of ICASSP '94. IEEE International Conference on Acoustics, Speech and Signal Processing*, pages 613–616, 1994.
- [53] S. Zhou, H. Bondell, A. Tordesillas, B. I. P. Rubinstein, and J. Bailey. Early identification of an impending rockslide location via a spatially-aided Gaussian mixture model. *Ann. Appl. Stat.*, 14(2):977–992, 2020.

JULIA LINDBERG, UNIVERSITY OF TEXAS AT AUSTIN, USA

Email address: julia.lindberg@math.utexas.edu

URL: <https://sites.google.com/view/julialindberg/home>

CARLOS AMÉNDOLA, TECHNISCHE UNIVERSITÄT BERLIN,, GERMANY

Email address: amendola@math.tu-berlin.de

URL: <http://www.carlos-amendola.com/>

JOSE ISRAEL RODRIGUEZ, UNIVERSITY OF WISCONSIN-MADISON, USA

Email address: jose@math.wisc.edu

URL: <https://sites.google.com/wisc.edu/jose>