# A proof of the continuous martingale convergence theorem

#### Joe Ghafari

#### Abstract

A proof of the continuous martingale convergence theorem is provided. It relies on a classical martingale inequality and the almost sure convergence of a uniformly bounded non-negative super-martingale, after a truncation argument.

Keywords: Martingale convergence, super-martingale. 2020 Mathematics Subject Classification: 60G44.

#### 1 Introduction

Unless otherwise stated, all processes in this paper are defined on a filtered probability space  $(\Omega, \mathcal{F}, (\mathcal{F}_r)_{r \in \mathbb{R}}, \mathbb{P})$ .

The continuous martingale convergence theorem states the following:

**Theorem 1.1.** Let  $(X_r)_{r \in \mathbb{R}}$  be a right-continuous integrable sub-martingale.

- 1. If  $\sup_{r \in \mathbb{R}_+} \mathbb{E}[|X_r|] < +\infty$ , then there exists an integrable random variable X such that  $\lim_{r \to +\infty} X_r = X$  a.s.
- 2. The limit  $\lim_{r\to\infty} X_r$  exists a.s. in  $[-\infty, +\infty]$ .

The discrete version of this result has various proofs (see [2], [4], [5], [7], and [8]). However, there doesn't seem to exist in the literature other proofs than Doob's original proof of Theorem 1.1. The latter uses the up-crossing inequality (for more details, see [9] and [11]).

In this paper we present a new elementary proof of theorem 1.1, avoiding the usual up-crossing lemma and using the classical inequality:

$$\delta \mathbb{P}\left(\sup_{r\in\mathbb{Q}\cap[u,u+v]}|Y_r|>\delta\right)\leq \mathbb{E}\left[|Y_u|\right]+2\mathbb{E}\left[|Y_{u+v}|\right],\tag{1}$$

where  $(u, v, \delta) \in \mathbb{R} \times (\mathbb{R}_+)^2$  and  $(Y_r)_{r \in \mathbb{R}}$  is an integrable super-martingale. The discrete martingale convergence theorem is also needed along with the following theorem due to Krickeberg. **Theorem 1.2** (Krickerberg decomposition). Let  $(Y_r)_{r \in \mathbb{R}_+}$  be an integrable submartingale for which  $\sup_{r \in \mathbb{R}_+} \mathbb{E}[|Y_r|] < +\infty$ . Then there exist a non-negative martingale  $(U_r)_{r \in \mathbb{R}_+}$  and a non-negative super-martingale  $(W_r)_{r \in \mathbb{R}_+}$  such that  $\sup_{r \in \mathbb{R}_+} \mathbb{E}[|U_r|] < +\infty$ ,  $\sup_{r \in \mathbb{R}_+} \mathbb{E}[|W_r|] < +\infty$ , and for all  $r \in \mathbb{R}_+$ ,  $Y_r = U_r - W_r$ .

The reader is referred to [9], [11], and [10] for a proof of inequality (1) and Theorem 1.2.

Lastly, we note that our approach gives a new proof of the theorem on the convergence of an integrable reversed discrete sub-martingale (see [1] and [6] for a general statement and a proof using the up-crossing lemma).

### 2 Preliminary tools

In this section we present the necessary tools for our proof of Theorem 1.1.

The following lemma is crucial for the proof of Theorem 2.2 below.

**Lemma 2.1.** If Y is an integrable random variable, then  $(\mathbb{E}[Y|\mathcal{F}_{-k}])_{k\in\mathbb{N}}$  converges to  $\mathbb{E}[Y|\bigcap_{k\in\mathbb{N}}\mathcal{F}_{-k}]$  in  $L^1$ .

*Proof.* For every  $k \in \mathbb{N}$ , let  $H_k := \mathbb{E}[Y|\mathcal{F}_{-k}]$ . We fix  $\mathcal{F}_{\infty} := \bigcap_{q \in \mathbb{N}} \mathcal{F}_{-q}$ . We consider first the case where  $Y \in L^2$ . We have

$$\forall (q,k) \in \mathbb{N}^2, \mathbb{E}\left[\left(H_{k+q} - H_q\right)^2\right] = \mathbb{E}\left[H_q^2\right] - \mathbb{E}\left[H_{k+q}^2\right].$$

We note that the sequence  $\left(\mathbb{E}\left[H_{k}^{2}\right]\right)_{k\in\mathbb{N}}$  is non-increasing and bounded below, hence it converges in  $\mathbb{R}$ .

Consequently, there exists  $H \in L^2$  such that  $\lim_{k \to +\infty} \mathbb{E}\left[ (H_k - H)^2 \right] = 0$  and H is  $\mathcal{F}_{\infty}$ -measurable.

Since

$$\forall G \in \mathcal{F}_{\infty}, \int_{G} H \, \mathrm{d}\mathbb{P} = \lim_{q \to +\infty} \int_{G} H_{q} \, \mathrm{d}\mathbb{P} = \int_{G} Y \, \mathrm{d}\mathbb{P},$$

it follows that  $H = \mathbb{E}[Y|\mathcal{F}_{\infty}]$  a.s.

Turning to the general case, we have for all  $(k,q) \in \mathbb{N}^2$ ,

$$\mathbb{E}\left[|H_{k} - \mathbb{E}\left[Y|\mathcal{F}_{\infty}\right]|\right] \leq 2\mathbb{E}\left[\left|Y - Y\mathbb{1}_{\{|Y| \leq q\}}\right|\right] + \mathbb{E}\left[\left|\mathbb{E}\left[Y\mathbb{1}_{\{|Y| \leq q\}}|\mathcal{F}_{-k}\right] - \mathbb{E}\left[Y\mathbb{1}_{\{|Y| \leq q\}}|\mathcal{F}_{\infty}\right]\right|\right]$$

Considering  $k \to +\infty$  and from the special case which we already proved, we deduce that for every  $q \in \mathbb{N}$ ,

$$\limsup_{k \to +\infty} \mathbb{E}\left[ |H_k - \mathbb{E}\left[ Y | \mathcal{F}_{\infty} \right] | \right] \le 2\mathbb{E}\left[ |Y - Y \mathbb{1}_{\{|Y| \le q\}} | \right].$$

Applying the dominated convergence theorem, we conclude the proof.

We provide next a theorem on the almost sure convergence of a uniformly bounded non-negative super-martingale. **Theorem 2.2.** If  $(Y_r)_{r \in \mathbb{R}}$  is an integrable super-martingale such that

$$\exists c \in \mathbb{R}_+, \forall r \in \mathbb{R}, 0 \le Y_r \le c,$$

then there exist integrable random variables Y and Y' such that  $\lim_{r\to+\infty,r\in\mathbb{Q}} Y_r = Y$  a.s. and  $\lim_{r\to-\infty,r\in\mathbb{Q}} Y_r = Y'$  a.s.

*Proof.* • First, we will prove the almost sure convergence at  $+\infty$ .

Since  $\sup_{k\in\mathbb{N}} \mathbb{E}[|Y_k|] \leq c < +\infty$ , it follows from the discrete martingale convergence theorem that  $(Y_k)_{k\in\mathbb{N}}$  converges a.s. to a random variable Y such that  $0 \leq Y \leq c$  a.s.

To prove the result in its generality, we begin by fixing  $k \in \mathbb{N}^*$  and  $\delta \in \mathbb{R}^*_+$ . Noticing that  $(Y_{r+k} - Y_k)_{r \in \mathbb{R}_+}$  is a super-martingale relative to  $(\mathcal{F}_{r+k})_{r \in \mathbb{R}_+}$ and applying inequality (1), we obtain that for all  $q \in \mathbb{N}^*$ ,

$$\begin{split} \delta \mathbb{P}\left(\sup_{r\in\mathbb{Q}\cap[0,q]}|Y_{r+k}-Y|>\delta\right) &\leq \delta \mathbb{P}\left(\sup_{r\in\mathbb{Q}\cap[0,q]}|Y_{r+k}-Y_k|>\frac{\delta}{2}\right) + \delta \mathbb{P}\left(|Y_k-Y|>\frac{\delta}{2}\right) \\ &\leq 4\mathbb{E}\left[|Y_{q+k}-Y_k|\right] + 2\mathbb{E}\left[|Y_k-Y|\right] \\ &\leq 4\mathbb{E}\left[|Y_{q+k}-Y|\right] + 6\mathbb{E}\left[|Y_k-Y|\right]. \end{split}$$

We note that  $\lim_{q\to+\infty} \mathbb{E}\left[|Y_{q+k} - Y|\right] = 0$  from the dominated convergence theorem.

Hence for any  $k \in \mathbb{N}^*$  and all  $\delta \in \mathbb{R}^*_+$ ,

$$\delta \mathbb{P}\left(\sup_{r\in\mathbb{Q}\cap[k,+\infty[}|Y_r-Y|>\delta\right)=\mathbb{P}\left(\bigcup_{q\in\mathbb{N}^*}\left\{\sup_{r\in\mathbb{Q}\cap[0,q]}|Y_{r+k}-Y|>\delta\right\}\right)\leq 6\mathbb{E}\left[|Y_k-Y|\right].$$

Taking  $k \to +\infty$  and applying the dominated convergence theorem, we conclude that for all  $\delta \in \mathbb{R}^*_+$ ,

$$\lim_{k \to +\infty} \mathbb{P}\left(\sup_{r \in \mathbb{Q} \cap [k, +\infty[} |Y_r - Y| > \delta\right) = 0,$$

in other words  $\lim_{r \to +\infty, r \in \mathbb{Q}} Y_r = Y$  a.s.

• The proof of the almost sure convergence at  $-\infty$  is essentially the same as before, therefore it's sufficient to show that the sequence  $(Y_{-k})_{k \in \mathbb{N}}$  converges a.s.

We will provide an elegant proof that doesn't use the up-crossing inequality.

For every  $k \in \mathbb{N}$ , let  $\Delta_k := Y_{-k-1} - \mathbb{E}[Y_{-k} | \mathcal{F}_{-k-1}]$  and  $V_k := \sup_{q \in \mathbb{N}} \left( \sum_{n=0}^q \Delta_{k+n} \right)$ . We note that the sequence  $(\mathbb{E}[Y_{-k}])_{k \in \mathbb{N}}$  is non-decreasing and bounded above, hence it has a finite limit denoted by l.

Also for every  $k \in \mathbb{N}, 0 \le V_k \le V_{k+1}$  a.s., therefore by monotone convergence theorem we have

$$\forall k \in \mathbb{N}, \mathbb{E}\left[V_k\right] = \sum_{q \in \mathbb{N}} \mathbb{E}\left[\Delta_{q+k}\right] = l - \mathbb{E}\left[Y_{-k}\right] < +\infty.$$

So  $(V_k)_{k \in \mathbb{N}}$  is a sequence of integrable random variable such that  $\lim_{k \to +\infty} \mathbb{E}[V_k] = 0$ .

Next, we check that  $(Y_k + V_{-k})_{k \in \mathbb{Z}_-}$  is an integrable martingale relative to  $(\mathcal{F}_k)_{k \in \mathbb{Z}_-}$ .

We fix 
$$k \in \mathbb{Z}_{-}$$
.

 $Y_k + V_{-k}$  is  $\mathcal{F}_k$ -measurable and integrable. We also have

$$\forall G \in \mathcal{F}_{k-1}, \int_{G} Y_{k} \, \mathrm{d}\mathbb{P} + \int_{G} V_{-k} \, \mathrm{d}\mathbb{P} = \int_{G} \mathbb{E}\left[Y_{k}|\mathcal{F}_{k-1}\right] \, \mathrm{d}\mathbb{P} + \sum_{q \in \mathbb{N}} \int_{G} \Delta_{q-k} \, \mathrm{d}\mathbb{P}$$
$$= \int_{G} Y_{k-1} \, \mathrm{d}\mathbb{P} - \int_{G} \Delta_{-k} \, \mathrm{d}\mathbb{P} + \sum_{q \in \mathbb{N}} \int_{G} \Delta_{q-k} \, \mathrm{d}\mathbb{P}$$
$$= \int_{G} Y_{k-1} \, \mathrm{d}\mathbb{P} + \sum_{q \in \mathbb{N}} \int_{G} \Delta_{q+1-k} \, \mathrm{d}\mathbb{P}$$
$$= \int_{G} Y_{k-1} \, \mathrm{d}\mathbb{P} + \int_{G} V_{1-k} \, \mathrm{d}\mathbb{P}.$$

Consequently,  $(Y_k + V_{-k})_{k \in \mathbb{Z}_-}$  is a martingale and hence for all  $k \in \mathbb{N}$ ,  $Y_{-k} = \mathbb{E}[Y_0 + V_0 | \mathcal{F}_{-k}] - V_k$  a.s.

It follows from Lemma 2.1 that  $(Y_{-k})_{k \in \mathbb{N}}$  converges to  $Y' := \mathbb{E} \left[ Y_0 + V_0 | \bigcap_{k \in \mathbb{N}} \mathcal{F}_{-k} \right]$ in  $L^1$ .

Lastly, noticing that  $(Y_k - Y')_{k \in \mathbb{Z}_-}$  is a super-martingale relative to  $(\mathcal{F}_k)_{k \in \mathbb{Z}_-}$ and using the discrete version of inequality (1), we have for every  $\delta \in \mathbb{R}^*_+$ and every  $(q, n) \in \mathbb{N}^2$ ,

$$\delta \mathbb{P}\left(\max_{0 \le k \le q} |Y_{-k-n} - Y'| > \delta\right) \le \mathbb{E}\left[|Y_{-q-n} - Y'|\right] + 2\mathbb{E}\left[|Y_{-n} - Y'|\right].$$

Letting  $q \to +\infty$  we obtain that for any  $\delta \in \mathbb{R}^*_+$  and all  $n \in \mathbb{N}$ ,

$$\mathbb{P}\left(\sup_{k\in\mathbb{N}}|Y_{-k-n}-Y'|>\delta\right)\leq\frac{2}{\delta}\mathbb{E}\left[|Y_{-n}-Y'|\right].$$

So for every  $\delta \in \mathbb{R}^*_+$ ,  $\lim_{n \to +\infty} \mathbb{P}(\sup_{k \in \mathbb{N}} |Y_{-k-n} - Y'| > \delta) = 0$ , concluding the proof.

We end this section by stating and proving a general version of Theorem 2.2.

**Theorem 2.3.** If  $(Y_r)_{r \in \mathbb{R}}$  is a non-negative super-martingale, then the limits  $\lim_{r \to +\infty, r \in \mathbb{Q}} Y_r$  and  $\lim_{r \to -\infty, r \in \mathbb{Q}} Y_r$  exist a.s. in  $\overline{\mathbb{R}}_+$ .

In particular, if the sample paths of  $(Y_r)_{r\in\mathbb{R}}$  are right-continuous, then  $\lim_{r\to+\infty} Y_r$ and  $\lim_{r\to-\infty} Y_r$  exist a.s. in  $\overline{\mathbb{R}}_+$ .

*Proof.* We will only prove the almost sure existence of the limit at  $+\infty$ , the proof is analogous at  $-\infty$ .

The idea is to truncate properly so that the super-martingale property is preserved.

We fix  $q \in \mathbb{N}$ .

 $(\min(q, Y_r))_{r \in \mathbb{R}}$  is a non-negative super-martingale uniformly bounded by q. We deduce from Theorem 2.2 that  $\limsup_{r \to +\infty, r \in \mathbb{Q}} \min(q, Y_r) = \liminf_{r \to +\infty, r \in \mathbb{Q}} \min(q, Y_r)$  a.s.

We also have the following relations:

$$\min\left(q, \limsup_{r \to +\infty, r \in \mathbb{Q}} Y_r\right) = \limsup_{r \to +\infty, r \in \mathbb{Q}} \min\left(q, Y_r\right),$$
$$\min\left(q, \liminf_{r \to +\infty, r \in \mathbb{Q}} Y_r\right) = \liminf_{r \to +\infty, r \in \mathbb{Q}} \min\left(q, Y_r\right).$$

So for every  $q \in \mathbb{N}$ , min  $(q, \limsup_{r \to +\infty, r \in \mathbb{Q}} Y_r) = \min(q, \liminf_{r \to +\infty, r \in \mathbb{Q}} Y_r)$ a.s. Hence  $\limsup_{r \to +\infty, r \in \mathbb{Q}} Y_r = \liminf_{r \to +\infty, r \in \mathbb{Q}} Y_r$  a.s., yielding that  $\lim_{r \to +\infty, r \in \mathbb{Q}} Y_r$  exists a.s. in  $\overline{\mathbb{R}}_+$ .

If  $(Y_r)_{r \in \mathbb{R}}$  is right-continuous, then  $\lim_{r \to +\infty} Y_r = \lim_{r \to +\infty, r \in \mathbb{Q}} Y_r$  a.s.  $\Box$ 

**Remark 2.4.** Theorem 2.3 holds true if  $\mathbb{Q}$  is replaced by a countable dense subset D of  $\mathbb{R}$ .

## 3 Proof of the continuous martingale convergence theorem

Finally, we are ready to prove our theorem.

Proof of Theorem 1.1. Since the sample paths of  $(X_r)_{r\in\mathbb{R}}$  are right-continuous, it's sufficient to prove that the limits of  $(X_r)_{r\in\mathbb{Q}}$  at  $+\infty$  and  $-\infty$  exist almost surely.

- 1. Applying Theorem 1.2, there exist a non-negative martingale  $(U_r)_{r \in \mathbb{R}_+}$ and a non-negative super-martingale  $(W_r)_{r \in \mathbb{R}_+}$  such that  $\sup_{r \in \mathbb{R}_+} \mathbb{E}\left[|U_r|\right] < +\infty$ ,  $\sup_{r \in \mathbb{R}_+} \mathbb{E}\left[|W_r|\right] < +\infty$ , and for every  $r \in \mathbb{R}_+, X_r = U_r - W_r$ . Theorem 2.3 yields that  $\lim_{r \to +\infty, r \in \mathbb{Q}} U_r$  and  $\lim_{r \to +\infty, r \in \mathbb{Q}} W_r$  exist a.s. in  $\mathbb{R}_+$ , we denote these almost sure limits by U and W, respectively. It follows by Fatou's lemma that U and W are integrable, in particular they are finite a.s. and hence  $\lim_{r \to +\infty, r \in \mathbb{Q}} X_r = U - W \in L^1$  a.s.
- 2. To verify the result, we need to write  $X_r$  suitably. We note that  $\left(\mathbb{E}\left[X_0^+|\mathcal{F}_r\right]\right)_{r\in\mathbb{R}_-}$  is an integrable martingale such that for all  $r\in\mathbb{R}_-,\mathbb{E}\left[X_0^+|\mathcal{F}_r\right]\geq\mathbb{E}\left[X_0|\mathcal{F}_r\right]\geq X_r$  a.s., so  $\left(\mathbb{E}\left[X_0^+|\mathcal{F}_r\right]-X_r\right)_{r\in\mathbb{R}_-}$ is a non-negative super-martingale. Applying again Theorem 2.3 and since  $X_0^+\in L^1$ , the limits  $\lim_{r\to-\infty,r\in\mathbb{Q}}\mathbb{E}\left[X_0^+|\mathcal{F}_r\right]$ and  $\lim_{r\to-\infty,r\in\mathbb{Q}}\left(\mathbb{E}\left[X_0^+|\mathcal{F}_r\right]-X_r\right)$  exist a.s. in  $\mathbb{R}$  and  $\overline{\mathbb{R}}_+$ , respectively. By writing for every  $r\in\mathbb{R}_-, X_r=\mathbb{E}\left[X_0^+|\mathcal{F}_r\right]-\left(\mathbb{E}\left[X_0^+|\mathcal{F}_r\right]-X_r\right)$ , we conclude that  $\lim_{r\to-\infty,r\in\mathbb{Q}}X_r$  exists a.s. in  $[-\infty,+\infty[$ .

**Remark 3.1.** Following the procedure used in proving Theorem 1.1 we can show that if  $u \in \mathbb{R}$ , D is a countable dense subset of  $\mathbb{R}$ , and  $(Y_r)_{r\in\mathbb{R}}$  is an integrable sub-martingale, then there exist integrable random variables V and V'such that  $\lim_{r\downarrow u, r\in D} Y_r = V$  a.s.,  $\lim_{r\uparrow u, r\in D} Y_r = V'$  a.s.,  $\mathbb{E}[V|\mathcal{F}_u] \geq Y_u$  a.s., and  $\mathbb{E}[Y_u|\mathcal{F}_{u-1}] \geq V'$  a.s. (see [9] for another proof applying the up-crossing inequality).

#### References

- Y. S. Chow and H. Teicher, Probability Theory: Independence, Interchangeability, Martingales, Third Edition, Springer-Verlag, New York, 1997.
- [2] J. L. Doob, Stochastic Processes, Wiley, New York, 1953.
- [3] K. Krickeberg, Probability Theory, Addison-Wesley, Reading, Mass., 1965.
- [4] R. Isaac, A Proof of the Martingale Convergence Theorem, Proc. Amer. Math. Soc. 16, 842-844, 1965.
- [5] C. W. Lamb, A Short Proof of the Martingale Convergence Theorem, Proc. Amer. Math. Soc. 38, 215-217, 1973.
- [6] J. Neveu, Discrete-Parameter Martingales, North-Holland, Amsterdam, 1975.
- [7] P. A. Meyer, *Martingales and Stochastic Integrals I*, Lecture Notes in Math. 284, Springer-Verlag, Berlin, 1972.
- [8] V. M. Kruglov, On the Kolmogorov theorem, J. Math. Sci. (N.Y.) 106 (2), 2830–2834, 2001.
- [9] C. Dellacherie and P. A. Meyer, Probabilities and Potential B: Theory of Martingales, North-Holland, Amsterdam, 1982.
- [10] D. Revuz and M. Yor, Continuous Martingales and Brownian Motion, Third Edition, Springer-Verlag, Berlin, 1999.
- [11] I. Karatzas and S. E. Shreve, Brownian Motion and Stochastic Calculus, Second Edition, Springer, New York, 1998.

Department of Mathematics and Statistics, University of Ottawa, Ottawa, K1N 6N5, Canada

Email: jghaf099@uottawa.ca