## Eavesdropping on the Decohering Environment: Quantum Darwinism, Amplification, and the Origin of Objective Classical Reality

Akram Touil, 1,2,\* Bin Yan,2,3 Davide Girolami,4 Sebastian Deffner,1,5 and Wojciech Hubert Zurek3

<sup>1</sup>Department of Physics, University of Maryland, Baltimore County, Baltimore, MD 21250, USA

<sup>2</sup>Center for Nonlinear Studies, Los Alamos National Laboratory, Los Alamos, New Mexico 87545

<sup>3</sup>Theoretical Division, Los Alamos National Laboratory, Los Alamos, New Mexico 87545

<sup>4</sup>Politecnico di Torino, Corso Duca degli Abruzzi 24, Torino, 10129, Italy

<sup>5</sup>Instituto de Física 'Gleb Wataghin', Universidade Estadual de Campinas, 13083-859, Campinas, São Paulo, Brazil

"How much information about a system S can one extract from a fragment F of the environment E that decohered it?" is the central question of Quantum Darwinism. To date, most answers relied on the quantum mutual information of SF, or on the Holevo bound on the channel capacity of F to communicate the classical information encoded in S. These are reasonable upper bounds on what is really needed but much harder to calculate – the accessible information in the fragment F about S. We consider a model based on imperfect c-not gates where all the above can be computed, and discuss its implications for the emergence of objective classical reality. We find that all relevant quantities, such as the quantum mutual information as well as various bounds on the accessible information exhibit similar behavior. In the regime relevant for the emergence of objective classical reality this includes scaling independent of the quality of the imperfect c-not gates or the size of E, and even nearly independent of the initial state of S.

Quantum Darwinism [1–5] explains the emergence of objective classical reality in our quantum Universe: The decohering environment  $\mathcal{E}$  is a "witness" who monitors and can reveal the state of the system  $\mathcal{S}$ . Agents like us never measure systems of interest directly. Rather, we accesses fragments  $\mathcal{F}$  of  $\mathcal{E}$  that carry information about them. Since its inception [1], Quantum Darwinism has advanced on both theory [6–25] and experimental fronts [26–29].

Quantum mutual information  $I(\mathcal{S}:\mathcal{F})$  between an environment fragment and the system yields an upper bound on what  $\mathcal{F}$  can reveal about  $\mathcal{S}$ . It has been used to estimate the capacity of the environment as a communication channel. We analyze a solvable model based on imperfect tunable c-not (or c-maybe) gates that couple  $\mathcal{S}$  to the subsystems of  $\mathcal{E}$ . We compute the mutual information  $I(\mathcal{S}:\mathcal{E})$  as well as the Holevo  $\chi(\mathcal{S}:\mathcal{F})$  [30, 31] – that characterize the accessible information in our c-maybe - based model. We also compute the quantum discord [1, 32–37] – the difference of  $I(\mathcal{S}:\mathcal{F})$  and  $\chi(\mathcal{S}:\mathcal{F})$  that quantifies the genuinely quantum correlations between  $\mathcal{S}$  and  $\mathcal{F}$  [38–41].

We find that  $I(\mathcal{S}:\mathcal{F})$  and  $\chi(\mathcal{S},\mathcal{F})$  exhibit strikingly similar dependence on the size of  $\mathcal{F}$ , with the initial steep rise followed by the classical plateau where, at the level set by the entropy  $H_{\mathcal{S}}$  of the system, the information  $\mathcal{F}$  has about  $\mathcal{S}$  saturates: Enlarging  $\mathcal{F}$  only confirms what is already known. This behavior is universal and nearly independent of the initial state of  $\mathcal{S}$  and the size of  $\mathcal{E}$ .

The model. The system  $\mathcal S$  is a qubit coupled to N independent non-interacting qubits of the environment  $\mathcal E$  via a c-maybe gate,

$$U_{\emptyset} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & s & c \\ 0 & 0 & c & -s \end{pmatrix}. \tag{1}$$

The parameters  $c = \cos(a)$  and  $s = \sin(a)$  (where a is the rotation angle of the target qubit) quantify the imperfection.

Our Quantum Universe SE starts in a pure state:

$$|\Psi^{0}_{S\mathcal{E}}\rangle = (\sqrt{p}|0_{S}\rangle + \sqrt{q}|1_{S}\rangle) \bigotimes_{i=1}^{N} |0^{i}\rangle,$$
 (2)

where p+q=1. The unitary  $U_{\odot}$  correlates each qubit in  $\mathcal E$  with  $\mathcal S$ , and we obtain a branching state [42],

$$|\Psi_{\mathcal{S}\mathcal{E}}^{\oslash}\rangle = \sqrt{p} |0_{\mathcal{S}}\rangle \bigotimes_{i=1}^{N} |0_{\mathcal{E}_{i}}\rangle + \sqrt{q} |1_{\mathcal{S}}\rangle \bigotimes_{i=1}^{N} |1_{\mathcal{E}_{i}}\rangle. \quad (3)$$

By construction  $|0_S\rangle$  and  $|1_S\rangle$  are the pointer states [43, 44]. They are orthogonal and immune to decoherence. The corresponding record states of  $\mathcal{E}$  are

$$|0_{\mathcal{E}_i}\rangle \equiv |0^i\rangle \quad \text{and} \quad |1_{\mathcal{E}_i}\rangle \equiv s|0^i\rangle + c|1^i\rangle \,, \qquad (4)$$

in terms of the orthogonal basis  $|0^i\rangle$  and  $|1^i\rangle$  of the *i*th qubit that defines  $U_{\varnothing}$ , so that  $\langle 0_{\varepsilon_i}|1_{\varepsilon_i}\rangle=s$ .

We will be interested in the correlations between the fragment  $\mathcal{F}$  and  $\mathcal{S}$ . The marginal states of  $\mathcal{S}$ , an m-qubit fragment  $\mathcal{F}_m$ , and a bipartition  $\mathcal{SF}_m$  are rank-two density matrices [45]:

$$\rho_{\mathcal{S}} \equiv \operatorname{tr}_{\mathcal{E}} \left\{ |\Psi_{\mathcal{S}\mathcal{E}}^{\oslash}\rangle \langle \Psi_{\mathcal{S}\mathcal{E}}^{\oslash}| \right\} = \begin{pmatrix} p & s^{N} \sqrt{pq} \\ s^{N} \sqrt{pq} & q \end{pmatrix}, \quad (5)$$

$$\rho_{\mathcal{F}_m} = \begin{pmatrix} p & s^m \sqrt{pq} \\ s^m \sqrt{pq} & q \end{pmatrix}, \tag{6}$$

$$\rho_{\mathcal{SF}_m} = \begin{pmatrix} p & s^{N-m} \sqrt{pq} \\ s^{N-m} \sqrt{pq} & q \end{pmatrix} . \tag{7}$$

Symmetric quantum mutual information is often used to estimate the accessible information in  $\mathcal{F}$  in Quantum Darwinism [2–4, 11–13, 46–48]. It is defined using the von Neumann entropy,  $H(\rho) = -\text{tr} \{\rho \log_2(\rho)\}$  as;

$$I(\mathcal{S}:\mathcal{F}_m) = H_{\mathcal{S}} + H_{\mathcal{F}_m} - H_{\mathcal{S},\mathcal{F}_m}.$$
 (8)

Joint entropy  $H_{\mathcal{S},\mathcal{F}_m}$  quantifies the ignorance about the state of  $\mathcal{S}\mathcal{F}_m$  in the tensor product of the Hilbert spaces of  $\mathcal{S}$  and  $\mathcal{F}$ .

In our model  $I(S : \mathcal{F}_m)$  can be computed exactly [49];

$$I(S: \mathcal{F}_m) = h(\lambda_{N,p}^+) + h(\lambda_{m,p}^+) - h(\lambda_{N-m,p}^+), \quad (9)$$

where  $h(x) = -x \log_2(x) - (1-x) \log_2(1-x)$  and  $\lambda_{k,p}^{\pm}$  are the eigenvalues of the density matrices (5) – (7),

$$\lambda_{k,p}^{\pm} = \frac{1}{2} \left( 1 \pm \sqrt{(q-p)^2 + 4s^{2k}pq} \right) .$$
 (10)

We thus have a closed expression for the mutual information  $I(S : \mathcal{F}_m)$ .

As seen in Fig. 1, symmetric mutual information  $I(S : \mathcal{F}_m)$  exhibits a steep initial rise with increasing fragment size m, as a larger  $\mathcal{F}_m$  provides more data about S. This initial rise is followed by a long *classical plateau*, where the additional information imprinted on the environment is redundant.

Note that, when  $\mathcal{SE}$  is in a pure state, the entropy of a fragment  $\mathcal{F}$  is equal to  $H_{\mathcal{SdF}}$ , that is the entropy  $\mathcal{S}$  would have if it was decohered only by the fragment  $\mathcal{F}$ . When we further assume  $good\ decoherence\ [42,49]\ -$  i.e., that the off-diagonal terms of  $\rho_{\mathcal{S}}$  and  $\rho_{\mathcal{SF}_m}$  are negligible (which in our model corresponds to  $s^{N-m}\ll s^m$ ) – we obtain an approximate equality;

$$I(\mathcal{S}:\mathcal{F}_m) = H_{\mathcal{F}_m} = H_{\mathcal{S}d\mathcal{F}_m},\tag{11}$$

since  $H_{\mathcal{S}}=H_{\mathcal{S}\mathcal{F}_m}$  cancel one another in Eq. (8). Furthermore, when the environment fragments are typical [50] (and in our model all fragments of the same size are identical – hence, each is typical) the plot of  $I(\mathcal{S}:\mathcal{F}_m)$  is antisymmetric around  $I(\mathcal{S}:\mathcal{F}_m)=H_{\mathcal{S}}$  and m=N/2 [42].

We will see that the behavior of  $I(\mathcal{S}:\mathcal{F}_m)$  is approximately universal. This means that after suitable re-scaling its functional form is nearly independent of the size of the environment N, of the quality of the c-maybe gate  $U_{\oslash}$ , and almost independent of the initial state of  $\mathcal{S}$ .

Agents generally do not insist on knowing the state of  $\mathcal{S}$  completely, but tolerate a finite information deficit  $\delta$ . When  $I(\mathcal{S}:\mathcal{F}_{m_{\delta}}) \geq (1-\delta)H_{\mathcal{S}}$  is attained already for a fragment with  $m_{\delta} \ll N$  subsystems, a fraction  $f_{\delta} = m_{\delta}/N$  of the environment, then there are many  $(1/f_{\delta})$  such fragments. We define redundancy of the information about  $\mathcal{S}$  in  $\mathcal{E}$  via:

$$\mathcal{R}_{\delta} \equiv N/m_{\delta}$$
 with  $I(\mathcal{S}:\mathcal{F}_{m_{\delta}}) = (1-\delta)H_{\mathcal{S}}$ . (12)

Redundancy  $\mathcal{R}_{\delta}$  is the length of the classical plateau in the units set by  $m_{\delta}$ , see Fig. (1). The beginning of the plateau is determined by the smallest  $m_{\delta}$  such that  $I(\mathcal{S}:\mathcal{F}_{m_{\delta}}) \geq (1-\delta)H_{\mathcal{S}}$ .

In realistic models  $I(S : \mathcal{F}) = H_S$  only when f = 1/2 (see [42]). Thus, significant redundancy appears only when the requirement of completeness of the information about S that can be extracted from F is relaxed. Moreover, Eq. (12) is an overestimate since  $I(S : \mathcal{F}_{m_\delta})$  is only an




FIG. 1. Approximate universality of mutual information: Symmetric  $I(\mathcal{S}:\mathcal{F}_m)$  and Holevo bound  $\chi(\check{\mathcal{S}}:\mathcal{F}_m)$  coincide until the fragment  $\mathcal{F}_m$  becomes almost as large as  $\mathcal{E}$ . Renormalized  $I(\mathcal{S}:\mathcal{F})/H_{\mathcal{S}}$  and  $\chi(\check{\mathcal{S}}:\mathcal{F})/H_{\mathcal{S}}$  depend only weakly on the probabilities of the outcomes (see inset). Their difference – quantum discord  $D(\check{\mathcal{S}}:\mathcal{F})$  – vanishes until  $\mathcal{F}_m$  begins to encompass almost all of  $\mathcal{E}$ ,  $m \sim N/R_{\delta}$ . The inset also compares the ratios  $\chi(\check{\mathcal{S}}:\mathcal{F})/H_{\mathcal{S}}$  and  $\chi(\mathcal{S}:\check{\mathcal{F}})/H_{\mathcal{S}}$  computed for several probabilities p of the pointer state  $|0_{\mathcal{S}}\rangle$  in Eq. (3). Note that the fragment sizes  $m_{\delta}$  that supply  $\sim$ 80% of information about  $\mathcal{S}$  are only modestly affected by p and quite similar for these two different information measures.

upper bound of what can be found out about S from F [51].

We will now consider better estimates: Inset in Fig. 1 compares  $I(S : \mathcal{F}_{m_{\delta}})$  with the two Holevo - like  $\chi$ 's we are about to discuss and illustrates resulting fragment sizes (hence, redundancies) they imply.

Asymmetric mutual information is defined using conditional entropy. We mark the system whose states are used for such conditioning by an inverted "hat", so when it is  $\check{\mathcal{S}}$  we consider the asymmetric mutual information:

$$J(\check{\mathcal{S}}:\mathcal{F}_m)_{\{|s_k\rangle\}} = H_{\mathcal{F}} - H_{\mathcal{F}|\check{\mathcal{S}}_{\{|s_k\rangle\}}}.$$
 (13)

Above,  $H_{\mathcal{F}_m|\check{\mathcal{S}}_{\{|s_k\rangle\}}}$  is the conditional entropy [31] that quantifies the missing information about  $\mathcal{F}$  remaining after the observable with the eigenstates  $\{|s_k\rangle\}$  was measured. Accordingly, the joint entropy in Eq. (8) is replaced by;

$$H_{\mathcal{F}_m, \check{\mathcal{S}}_{\{|s_k\rangle\}}} = H_{\mathcal{F}_m|\check{\mathcal{S}}_{\{|s_k\rangle\}}} + H_{\check{\mathcal{S}}_{\{|s_k\rangle\}}}. \tag{14}$$

The asymmetric joint entropy depends on whether  $\mathcal{S}$  or  $\mathcal{F}$  are measured and on the measurements that are used. The entropy increase associated with the wavepacket reduction means that the asymmetric entropy (14) is typically larger than the symmetric version  $H_{\mathcal{S},\mathcal{F}_m}$  in Eq. (8): Local measurements cannot extract information encoded in the quantum correlations between  $\mathcal{S}$  and  $\mathcal{F}_m$ , which is why the asymmetric  $J(\check{\mathcal{S}}:\mathcal{F}_m)$  is needed, [31]; see also [52].

For optimal measurements the asymmetric  $J(\mathcal{S} : \mathcal{F}_m)$  defines the Holevo bound [30],

$$J(\check{\mathcal{S}}:\mathcal{F}_m) = \max_{\{|s_k\rangle\}} J(\check{\mathcal{S}}:\mathcal{F}_m)_{\{|s_k\rangle\}} \equiv \chi(\check{\mathcal{S}}:\mathcal{F}_m).$$
(15)

In our model, measurement of the pointer observable of  $\mathcal S$  is optimal [49]. Indeed, Eq. (3) shows that in the pointer basis  $\{|0_{\mathcal S}\rangle, |1_{\mathcal S}\rangle\}$  the conditional entropy disappears,  $H_{\mathcal F_m|\check{\mathcal S}}=0$ , as states of  $\mathcal F_m$  correlated with pointer states of  $\mathcal S$  are pure.

The limit of large  $\mathcal{E}$   $(N \geq N - m \gg m)$  reflects typical situation of agents (who do not even know the size of  $\mathcal{E}$ , and only access "their  $\mathcal{F}_m$ ", with  $m \ll N$ ). This is good

decoherence,  $s^N \leq s^{N-m} \ll s^m$ , and equations simplify: Using  $H_{\mathcal{S},\mathcal{F}_m} = H_{\mathcal{S}}$  and Eq. (11) we can thus write,

$$I(\mathcal{S}:\mathcal{F}_m) \approx H_{\mathcal{F}_m} = h(\lambda_{m,n}^+) = \chi(\check{\mathcal{S}}:\mathcal{F}_m).$$
 (16)

An immediate important consequence is that  $H_{\mathcal{F}_m}$  determines both the symmetric  $I(\mathcal{S}:\mathcal{F}_m)$  (except for the final rise) as well as the asymmetric (optimal)  $J(\check{\mathcal{S}}:\mathcal{F}_m)=\chi(\check{\mathcal{S}}:\mathcal{F}_m)$ . We have:

$$\chi(\check{\mathcal{S}}:\mathcal{F}_m) = -\frac{1}{2}\log_2\left(pq\left(1-s^{2m}\right)\right) - \sqrt{1-4pq\left(1-s^{2m}\right)} \operatorname{Arctanh}_2\left(\sqrt{1-4pq\left(1-s^{2m}\right)}\right), \quad (17)$$

where "Arctanh<sub>2</sub>" denotes Arctanh  $/ \ln(2)$ .

Fig. 1 compares  $\chi(\check{\mathcal{S}}:\mathcal{F}_m)$  with  $I(\mathcal{S}:\mathcal{F}_m)$  for finite and infinite N and for different values of s and p. As it shows, Eq. (17) matches  $I(\mathcal{S}:\mathcal{F}_m)$  until the far end  $(N-m\ll m)$  of the classical plateau. This is a consequence of two scalings: (i) "vertically", the plateau appears at  $H_{\mathcal{S}}=-p\log_2(p)-q\log_2(q)$ , and it is easy to see that for  $s^N\ll s^m\ll 1$  we have  $\chi(\check{\mathcal{S}}:\mathcal{F}_m)=H_{\mathcal{S}}$  in Eq. (17); (ii) "horizontally",  $H_{\mathcal{F}_m}$  depends on  $s^m$ , so weakly entangling gates can be compensated by using more of them – larger m. What is surprising is how insensitive are these plots to p, the probability of the outcome.

This remarkably universal behavior is a consequence of good decoherence [49]. Both,  $\rho_{\mathcal{S}}$  and  $\rho_{\mathcal{S}\mathcal{F}_m}$ , Eqs. (5) and (7), become diagonal in the pointer basis. Moreover, the quality of  $U_{\oslash}$  (set by c and s) determines the "information flow rate" from  $\mathcal{S}$  to  $\mathcal{F}$ . Thus, when (at a fixed p) one demands the same  $H_{\mathcal{F}_m}$ , this translates into identical  $\rho_{\mathcal{F}_m}$  when  $s_1^{m_1} \simeq s_2^{m_2}$ . Therefore, less efficiently entangling gates can be compensated by relying on more of them – on a suitably enlarged  $\mathcal{F}$ , with  $m_2 = m_1 \log(s_1)/\log(s_2)$ .

Environment as a communication channel. While the mutual information  $I(S : \mathcal{F}_m)$  is easier to compute and a

safe upper bound on the accessible information in  $\mathcal{F}_m$ , it is important to verify it is also a reasonable estimate of that accessible information (as generally assumed in much of the Quantum Darwinism literature). The asymmetric mutual information extracted by optimal measurements on the environment fragment  $\mathcal{F}_m$  is:

$$J(\mathcal{S}: \check{\mathcal{F}}_m) = H_{\mathcal{S}} - H_{\mathcal{S}|\check{\mathcal{F}}_m} = \chi(\mathcal{S}: \check{\mathcal{F}}_m). \tag{18}$$

The joint entropy given in terms of the conditional entropy  $H_{S|\tilde{\mathcal{F}}_m}$  now becomes,

$$H_{\mathcal{S}, \check{\mathcal{F}}_m} = H_{\mathcal{S}|\check{\mathcal{F}}_m} + H_{\check{\mathcal{F}}_m}. \tag{19}$$

As in Eq. (14) above, all terms in Eq. (19) depend on how  $\mathcal F$  is measured. However, while measuring  $\mathcal S$  in the pointer basis simplified the analysis (since e.g.  $H_{\mathcal F|\mathcal S_{\{|s_k\rangle\}}}=0$ ) this is no longer the case when  $\mathcal F_m$  is measured.

To compute  $\chi(S : \mathcal{F}_m)$  we rely on the Koashi-Winter monogamy relation [53]. Details of that calculation are relegated to the supplementary material [54].

We focus again on the limit of large  $\mathcal{E}$  ( $N \geq N - m \gg m$ ): Agents only access "their  $\mathcal{F}_m$ ", a small fraction of  $\mathcal{E}$  with  $m \ll N$ . Assuming good decoherence we obtain

$$\chi(\mathcal{S}: \check{\mathcal{F}}_m) = H_{\mathcal{S}} + \frac{1}{2} \log_2 \left( pqs^{2m} \right) + \sqrt{1 - 4pqs^{2m}} \operatorname{Arctanh}_2 \left( \sqrt{1 - 4pqs^{2m}} \right). \tag{20}$$

Equation (20) constitutes our main result. We have decomposed the Holevo-like quantity  $\chi(\mathcal{S}:\check{\mathcal{F}}_m)$  into the plateau entropy  $H_{\mathcal{S}}$  and  $H_{\mathcal{S}|\check{\mathcal{F}}_m}$  – the ignorance about  $\mathcal{S}$  remaining in spite of the optimal measurement on  $\mathcal{F}_m$  [55]. Rather remarkably,  $H_{\mathcal{S}|\check{\mathcal{F}}_m} = H_{\mathcal{S}} - \chi(\mathcal{S}:\check{\mathcal{F}}_m)$  – the conditional entropy – scales *exactly* with  $pqs^{2m}$ . What remains to do is to quantify the differences of  $I(\mathcal{S}:\mathcal{F}_m)$  and  $\chi(\check{\mathcal{S}}:\mathcal{F}_m)$  with  $\chi(\mathcal{S}:\check{\mathcal{F}}_m)$ . In Fig. 2 we compare it with these other, easier to compute, quantities.

Redundancy of the information about S in the channel  $\mathcal{F}_m$  can be now computed using  $\chi(S:\check{\mathcal{F}}_m)$ , Eq. (20),

and compared with the estimates based on  $I(\mathcal{S}:\mathcal{F}_m)$ . The fragment  $\mathcal{F}_{m_\delta}$  can carry all but the deficit  $\delta$  of the classical information about the pointer state of  $\mathcal{S}$  when  $\chi(\mathcal{S}:\check{\mathcal{F}}_{m_\delta}) \geq (1-\delta)H_{\mathcal{S}}$ . This leads to a transcendental equation for  $m_\delta$  that we solve numerically:  $R_\delta = N/m_\delta$ , where N is the number of subsystems in  $\mathcal{E}$ .

The inset in Fig. 1 shows that – while  $m_\delta$  deduced using  $I(\mathcal{S}:\mathcal{F}_m)\approx\chi(\check{\mathcal{S}}:\mathcal{F}_m)$  do not coincide with those obtained using  $\chi(\mathcal{S}:\check{\mathcal{F}}_m)$  – the difference is modest, unlikely to materially affect conclusions about the emergence of objective classical reality. Indeed, in the supplementary

materials we estimate that the redundancy estimates based on  $I(S: \mathcal{F}_m)$  and  $\chi(S: \check{\mathcal{F}}_m)$  differ at most by  $\sim 37\%$  for  $\delta \leq 0.2$ , and by much less in the regime where  $\delta \to 0$ .

In situations relevant for observers who rely on photons,  $R_{\delta=0.1}\simeq 10^8$  is amassed when sunlight illuminates a  $1\mu m$  dust grain in a superposition with a  $1\mu m$  spatial separation for  $1\mu s$  [47, 48]. It may seem like we are stretching the applicability of our c-maybe model too far, but the equations for  $I(\mathcal{S}:\mathcal{F}_m)$  and  $\chi(\check{\mathcal{S}}:\mathcal{F}_m)$  derived for photon scattering coincide with our Eq. (17), see supplement [54]. Thus, it appears that the information transfer from  $\mathcal{S}$  to  $\mathcal{E}$  leading to the buildup of redundancy has universal features captured by our model.

Quantum discord is the difference between symmetric (9) and asymmetric quantum mutual information [32–41]. The systemic discord is defined as;

$$D(\check{\mathcal{S}}:\mathcal{F}_m) = I(\mathcal{S}:\mathcal{F}_m) - \chi(\check{\mathcal{S}}:\mathcal{F}_m). \tag{21}$$

The measurements on pointer observables of S are optimal. Mutual information for pure decoherence induced by non-interacting subsystems of E can be written as [49, 56]:

$$I(\mathcal{S}:\mathcal{F}) = \begin{pmatrix} local/classical & global/quantum \\ H_{\mathcal{F}} - H_{\mathcal{F}}(0) \end{pmatrix} + \begin{pmatrix} H_{\mathcal{S}d\mathcal{E}} - H_{\mathcal{S}d\mathcal{E}_{\backslash \mathcal{F}}} \end{pmatrix} . \tag{22}$$

As  $\mathcal{SE}$  is a pure product state, the initial entropy of  $\mathcal{F}$  is zero,  $H_{\mathcal{F}}(0)=0$ . Assuming good decoherence and conditioning on the pointer basis (hence,  $\chi(\check{\mathcal{S}}:\mathcal{F}_m)=H_{\mathcal{F}_m}$ ), Eq. (16) we have

$$I(\mathcal{S}:\mathcal{F}_m) - J(\check{\mathcal{S}}:\mathcal{F}_m) = H_{\mathcal{S}d\mathcal{E}} - H_{\mathcal{S}d\mathcal{E}_{\backslash \mathcal{F}_m}}, \quad (23)$$

where  $H_{Sd\mathcal{E}}$   $(H_{Sd\mathcal{E}_{\setminus \mathcal{F}}})$  is the entropy of the system *decohered* by  $\mathcal{E}$  (or just by  $\mathcal{E}_{\setminus \mathcal{F}}$  – i.e.,  $\mathcal{E}$  less the fragment  $\mathcal{F}$ ).

The global/quantum term represents quantum discord in the pointer basis of  $\mathcal{S}$  [49]. Good decoherence implies  $H_{\mathcal{S}d\mathcal{E}} \approx H_{\mathcal{S}d\mathcal{E}_{\backslash\mathcal{F}}}$ , so  $D(\check{\mathcal{S}}:\mathcal{F}_m) \approx 0$ . As long as  $\mathcal{E}_{\backslash\mathcal{F}}$  is large enough to induce good decoherence, Eq. (16) holds, and, hence, the systemic discord (21) vanishes [57].

Systemic quantum discord can become large again when  $\mathcal{F}_m$  encompasses almost all  $\mathcal{E}$ , as in this case  $H_{\mathcal{S}\mathcal{F}_m}$  approaches  $H_{\mathcal{S}\mathcal{E}}=0$  (given our assumption of a pure  $\mathcal{S}\mathcal{E}$ ). In this (unphysical) limit  $I(\mathcal{S}:\mathcal{F}_m)$  climbs to  $H_{\mathcal{F}_m}+H_{\mathcal{S}}=2H_{\mathcal{S}}$ , while  $\chi(\check{\mathcal{S}}:\mathcal{F}_m)\leq H_{\mathcal{F}_m}$ . As good decoherence implies  $\chi(\check{\mathcal{S}}:\mathcal{F}_m)\approx H_{\mathcal{F}_m}$ ,  $D(\check{\mathcal{S}}:\mathcal{F}_m)$  can reach  $H_{\mathcal{F}_m}$ . Indeed, when  $\mathcal{S}\mathcal{E}$  is pure,  $\chi(\check{\mathcal{S}}:\mathcal{F}_m)$  and  $D(\check{\mathcal{S}}:\mathcal{F}_m)$  – classical and quantum content of the correlation – are complementary [36], see Fig. 1.

The fragmentary discord is the difference between the mutual information I(S : F) and what can be extracted from SF by measuring only the fragment F:

$$I(S:F) - \chi(S:\check{F}) \approx \chi(\check{S}:F) - \chi(S:\check{F}).$$
 (24)

It can be evaluated:

$$D(\mathcal{S}: \check{\mathcal{F}}_m) \approx H_{\check{\mathcal{F}}_m} - \left(H_{\mathcal{S}} - H_{\mathcal{S}|\check{\mathcal{F}}_m}\right),$$
  
=  $\left(H_{\mathcal{S}|\check{\mathcal{F}}_m} + H_{\check{\mathcal{F}}_m}\right) - H_{\mathcal{S}}.$  (25)




FIG. 2. Accessible information in the environment fragment. Fragment  $\mathcal{F}_m$  carries at most  $\chi(\mathcal{S}:\check{\mathcal{F}})$  of classical information about the system it helped decohere. As seen above, this Holevo - like quantity is less than the symmetric mutual information  $I(\mathcal{S}:\mathcal{F})$  or the Holevo bound  $\chi(\check{\mathcal{S}}:\mathcal{F}_m)$ . Their difference (quantum discord  $D(\mathcal{S}:\check{\mathcal{F}})$ ) is significant already early on (in contrast to  $D(\check{\mathcal{S}}:\mathcal{F})$ ), but disappears as the plateau is reached. It reappears again (as did  $D(\check{\mathcal{S}}:\mathcal{F})$ ) when  $\mathcal{F}_m$  begins to encompass almost all of  $\mathcal{E}$ .

The bracketed terms in the last two expressions represent different quantities. The difference between the symmetric and asymmetric mutual information  $H_{\tilde{\mathcal{F}}_m} - (H_{\mathcal{S}} - H_{\mathcal{S}|\tilde{\mathcal{F}}_m})$  is the original definition of discord.

Note that initially decoherence does not suppress fragmentary discord  $D(\mathcal{S}:\check{\mathcal{F}}_m)$ . This is because the states of  $\mathcal{F}_m$  that are correlated with the pointer states of  $\mathcal{S}$  are not orthogonal: The scalar product of the branch fragments  $\mathcal{F}_m$  corresponding to  $|0_{\mathcal{S}}\rangle$  and  $|1_{\mathcal{S}}\rangle$  is  $s^m$ . Thus, while the symmetric mutual information increases with m, orthogonality is approached gradually, also as m increases. Perfect distinguishability, i.e., orthogonality of record states of  $\mathcal{F}$  is needed to pass on all the information about  $\mathcal{S}$  [58–60]. See again Fig. 2 for an illustration of these findings.

Concluding remarks. We found that in the pre-plateau regime relevant for emergence of objective reality (where  $I(\mathcal{S}:\mathcal{F})$  increases with the size of  $\mathcal{F}$ ) the mutual information as well as the Holevo bound  $\chi(\check{\mathcal{S}}:\mathcal{F})$  coincide and exhibit universal scaling behaviors independent of the size of  $\mathcal{E}$ , of how imperfect are the c-maybe's, and only weakly dependent on the probabilities of pointer states. The corresponding Holevo  $\chi(\check{\mathcal{S}}:\mathcal{F})$  and  $I(\mathcal{S}:\mathcal{F})$  coincide until  $\mathcal{F}$  encompasses almost all of  $\mathcal{E}$ .

However, the accessible information  $\chi(\mathcal{S}:\check{\mathcal{F}})$  in the environment fragments  $\mathcal{F}$  differs somewhat from  $I(\mathcal{S}:\mathcal{F})$  in the pre-plateau region. This difference tends to be small compared to, e.g., the level of the plateau, and disappears as the plateau is reached for larger fragments. This behavior – generic when many copies of the information about  $\mathcal{S}$  are deposited in the environment – facilitates estimates of the redundancy of the information about the system in the environment, as the differences between  $I(\mathcal{S}:\mathcal{F}) \approx \chi(\check{\mathcal{S}}:\mathcal{F})$ 

or  $\chi(\mathcal{S}:\check{\mathcal{F}})$  are noticeable but inconsequential.

To sum up, sensible measures of information flow lead to compatible conclusions about  $R_{\delta}$ . The differences in the estimates of redundancy based on these quantities are insignificant for the emergence of objective classical reality – the overarching goal of Quantum Darwinism. The functional dependence of the symmetric mutual information in the photon scattering model [47, 48] is the same as in our model. Thus, the universality we noted in scaling with s and p (approximate for  $I(S:\mathcal{F})=\chi(\check{S}:\mathcal{F}_m)$ ), exact for  $\chi(S:\check{\mathcal{F}}_m)$ ) may be a common attribute of the information that reaches us, human observers.

We acknowledge several discussions with Michael Zwolak, who has provided us with extensive and perceptive comments that have greatly improved presentation of our results. This research was supported by grants FQXiRFP-1808 and FQXiRFP-2020-224322 from the Foundational Questions Institute and Fetzer Franklin Fund, a donor advised fund of Silicon Valley Community Foundation (SD and WHZ, respectively), as well as by the Department of Energy under the LDRD program in Los Alamos. A.T., B.Y. and W.H.Z. also acknowledge support from U.S. Department of Energy, Office of Science, Basic Energy Sciences, Materials Sciences and Engineering Division, Condensed Matter Theory Program, and the Center for Nonlinear Studies. D. G. acknowledges financial support from the Italian Ministry of Research and Education (MIUR), grant number 54\_AI20GD01.

- \* akramt1@umbc.edu
- [1] W. H. Zurek, Einselection and decoherence from an information theory perspective, Ann. Phys. **9**, 855 (2000).
- [2] W. H. Zurek, Decoherence, einselection, and the quantum origins of the classical, Rev. Mod. Phys. **75**, 715 (2003).
- [3] H. Ollivier, D. Poulin, and W. H. Zurek, Objective properties from subjective quantum states: Environment as a witness, Phys. Rev. Lett. **93**, 220401 (2004).
- [4] H. Ollivier, D. Poulin, and W. H. Zurek, Environment as a witness: Selective proliferation of information and emergence of objectivity in a quantum universe, Phys. Rev. A 72, 042113 (2005).
- [5] W. H. Zurek, Quantum Darwinism, Nat. Phys. 5, 181 (2009).
- [6] G. L. Giorgi, F. Galve, and R. Zambrini, Quantum Darwinism and non-markovian dissipative dynamics from quantum phases of the spin-1/2 xx model, Phys. Rev. A **92**, 022105 (2015).
- [7] N. Balanesković, Random unitary evolution model of Quantum Darwinism with pure decoherence, Eur. Phys. J. D 69, 232 (2015).
- [8] N. Balaneskovic and M. Mendler, Dissipation, dephasing and Quantum Darwinism in qubit systems with random unitary interactions, Eur. Phys. J. D **70**, 177 (2016).
- [9] P. A. Knott, T. Tufarelli, M. Piani, and G. Adesso, Generic emergence of objectivity of observables in infinite dimen-

- sions, Phys. Rev. Lett. 121, 160401 (2018).
- [10] N. Milazzo, S. Lorenzo, M. Paternostro, and G. M. Palma, Role of information backflow in the emergence of Quantum Darwinism, Phys. Rev. A 100, 012101 (2019).
- [11] S. Campbell, B. Çakmak, O. E. Müstecaplıoğlu, M. Paternostro, and B. Vacchini, Collisional unfolding of Quantum Darwinism, Phys. Rev. A 99, 042103 (2019).
- [12] E. Ryan, M. Paternostro, and S. Campbell, Quantum Darwinism in a structured spin environment, arXiv preprint arrix:2011.13385 (2020).
- [13] G. García-Pérez, D. A. Chisholm, M. A. C. Rossi, G. M. Palma, and S. Maniscalco, Decoherence without entanglement and Quantum Darwinism, Phys. Rev. Research 2, 012061(R) (2020).
- [14] S. Lorenzo, M. Paternostro, and G. M. Palma, Anti-zeno-based dynamical control of the unfolding of Quantum Darwinism, Phys. Rev. Research 2, 013164 (2020).
- [15] M. Kiciński and J. K. Korbicz, Decoherence and objectivity in higher spin environments, arXiv preprint arXiv:2105.09093 (2021).
- [16] J. Korbicz, Roads to objectivity: Quantum Darwinism, spectrum broadcast structures, and strong Quantum Darwinism, arXiv preprint arXiv:2007.04276 (2020).
- [17] R. Blume-Kohout and W. H. Zurek, Quantum Darwinism in quantum brownian motion, Phys. Rev. Lett. 101, 240405 (2008).
- [18] M. Zwolak, C. J. Riedel, and W. H. Zurek, Amplification, redundancy, and quantum chernoff information, Phys. Rev. Lett. 112, 140406 (2014).
- [19] M. Zwolak, H. T. Quan, and W. H. Zurek, Quantum Darwinism in a mixed environment, Phys. Rev. Lett. 103, 110402 (2009).
- [20] J. P. Paz and A. J. Roncaglia, Redundancy of classical and quantum correlations during decoherence, Phys. Rev. A **80**, 042111 (2009).
- [21] C. J. Riedel, W. H. Zurek, and M. Zwolak, Objective past of a quantum universe: Redundant records of consistent histories, Phys. Rev. A 93, 032126 (2016).
- [22] C. J. Riedel, Classical branch structure from spatial redundancy in a many-body wave function, Phys. Rev. Lett. **118**, 120402 (2017).
- [23] F. G. Brandão, M. Piani, and P. Horodecki, Generic emergence of classical features in Quantum Darwinism, Nature communications 6, 7908 (2015).
- [24] X.-L. Qi and D. Ranard, Emergent classicality in general multipartite states and channels, arXiv preprint arXiv:2001.01507 (2020).
- [25] H.-F. Fu, Uniqueness of the observable leaving redundant imprints in the environment in the context of Quantum Darwinism, Phys. Rev. A 103, 042210 (2021).
- [26] M. A. Ciampini, G. Pinna, P. Mataloni, and M. Paternostro, Experimental signature of Quantum Darwinism in photonic cluster states, Phys. Rev. A 98, 020101(R) (2018).
- [27] M.-C. Chen, H.-S. Zhong, Y. Li, D. Wu, X.-L. Wang, L. Li, N.-L. Liu, C.-Y. Lu, and J.-W. Pan, Emergence of classical objectivity of Quantum Darwinism in a photonic quantum simulator, Science Bulletin **64**, 580 (2019).
- [28] T. K. Unden, D. Louzon, M. Zwolak, W. H. Zurek, and F. Jelezko, Revealing the emergence of classicality using nitrogen-vacancy centers, Phys. Rev. Lett. 123, 140402 (2019).

- [29] G. García-Pérez, M. A. C. Rossi, and S. Maniscalco, Ibm q experience as a versatile experimental testbed for simulating open quantum systems, npj Quantum Inf. 6, 1 (2020).
- [30] A. S. Holevo, Bounds for the quantity of information transmitted by a quantum communication channel, Problemy Peredachi Informatsii 9, 3 (1973).
- [31] M. A. Nielsen and I. Chuang, Quantum computation and quantum information (American Association of Physics Teachers, 2002).
- [32] H. Ollivier and W. H. Zurek, Quantum discord: A measure of the quantumness of correlations, Phys. Rev. Lett. 88, 017901 (2001).
- [33] L. Henderson and V. Vedral, Classical, quantum and total correlations, J. Phys. A: Math. Gen. 34, 6899 (2001).
- [34] P. Giorda and M. G. A. Paris, Gaussian quantum discord, Phys. Rev. Lett. 105, 020503 (2010).
- [35] M. Shi, W. Yang, F. Jiang, and J. Du, Quantum discord of two-qubit rank-2 states, J. Phys. A: Math. Theor. 44, 415304 (2011).
- [36] M. Zwolak and W. H. Zurek, Complementarity of quantum discord and classically accessible information, Sci. Rep. 3, 1729 (2013).
- [37] M. Zwolak, C. J. Riedel, and W. H. Zurek, Amplification, decoherence and the acquisition of information by spin environments, Sci. Rep. 6, 1 (2016).
- [38] A. Brodutch, A. Gilchrist, D. R. Terno, and C. J. Wood, Quantum discord in quantum computation, J. Phys.: Conf. Ser. 306, 012030 (2011).
- [39] G. Adesso, M. Cianciaruso, and T. R. Bromley, An introduction to quantum discord and non-classical correlations beyond entanglement, arXiv preprint arXiv:1611.01959 (2016).
- [40] A. Bera, T. Das, D. Sadhukhan, S. S. Roy, A. Sen(De), and U. Sen, Quantum discord and its allies: a review of recent progress, Rep. Prog. Phys. **81**, 024001 (2017).
- [41] K. Modi, A. Brodutch, H. Cable, T. Paterek, and V. Vedral, The classical-quantum boundary for correlations: Discord and related measures, Rev. Mod. Phys. 84, 1655 (2012).
- [42] R. Blume-Kohout and W. H. Zurek, A simple example of "Quantum Darwinism": Redundant information storage in many-spin environments, Found. Phys. **35**, 1857 (2005).
- [43] W. H. Zurek, Pointer basis of quantum apparatus: Into what mixture does the wave packet collapse?, Phys. Rev. D 24, 1516 (1981).
- [44] W. H. Zurek, Environment-induced superselection rules, Phys. Rev. D 26, 1862 (1982).
- [45] Deducing the reduced quantum states  $\rho_S$  and  $\rho_{SF_m}$  in Eqs. (5) and (7) is straightforward, whereas  $\rho_{F_m}$  is slightly more involved, see e.g. [49].
- [46] R. Blume-Kohout and W. H. Zurek, Quantum Darwinism: Entanglement, branches, and the emergent classicality of redundantly stored quantum information, Phys. Rev. A 73, 062310 (2006).
- [47] C. J. Riedel and W. H. Zurek, Quantum Darwinism in an everyday environment: Huge redundancy in scattered photons, Phys. Rev. Lett. **105**, 020404 (2010).
- [48] C. J. Riedel and W. H. Zurek, Redundant information from thermal illumination: Quantum Darwinism in scattered pho-

- tons, New J. Phys. 13, 073038 (2011).
- [49] M. Zwolak, H. T. Quan, and W. H. Zurek, Redundant imprinting of information in nonideal environments: Objective reality via a noisy channel, Phys. Rev. A **81**, 062110 (2010).
- [50] T. M. Cover and J. A. Thomas, *Elements of information the-ory* (John Wiley & Sons, 1999).
- [51] Note that  $H_S$  in the circumstances of interest to us is not the thermodynamic entropy of S. Rather, it is the missing information about the few *relevant* degrees of freedom of S. The thermodynamic entropy of a cat, for instance, will vastly exceed the information an observer is most interested in e.g., the one crucial bit in the 'diabolical contraption' envisaged by Schrödinger.
- [52] A. Touil, K. Weber, and S. Deffner, Quantum euler relation for local measurements, arXiv preprint arXiv:2106.04459 (2021).
- [53] M. Koashi and A. Winter, Monogamy of quantum entanglement and other correlations, Phys. Rev. A 69, 022309 (2004).
- [54] See Supplementary Material, which includes Refs.[61–64].
- [55] Strictly speaking, Holevo bound  $\chi$  on the channel capacity presumes conditioning on a predetermined classical basis (such as the pointer basis of  $\mathcal{S}$  in  $\chi(\check{\mathcal{S}}:\mathcal{F}_m)$ ). In general,  $\mathcal{F}_m$  in  $\chi(\mathcal{S}:\check{\mathcal{F}}_m)$  does not possess such a predetermined classical basis, hence the "Holevo-like" designation..
- [56] W. H. Zurek, Relative states and the environment: einselection, envariance, Quantum Darwinism, and the existential interpretation, arXiv preprint arXiv:0707.2832 (2007).
- [57] Some have advocated "strong Quantum Darwinism" [65] where the Holevo information  $\chi(\check{\mathcal{S}}:\mathcal{F}_m)$  based on the measurement of  $\mathcal{S}$  (rather than  $I(\mathcal{S}:\mathcal{F}_m)$ ) would play a key role. At least in case of good decoherence, and in view of Eq. (16) which implies  $\chi(\check{\mathcal{S}}:\mathcal{F}_m)\approx I(\mathcal{S}:\mathcal{F}_m)$  when  $m\ll N-m < N$ , such distinctions appear unnecessary.
- [58] W. H. Zurek, Quantum origin of quantum jumps: Breaking of unitary symmetry induced by information transfer in the transition from quantum to classical, Phys. Rev. A **76**, 052110 (2007).
- [59] W. H. Zurek, Wave-packet collapse and the core quantum postulates: Discreteness of quantum jumps from unitarity, repeatability, and actionable information, Phys. Rev. A 87, 052111 (2013).
- [60] B. Gardas, S. Deffner, and A. Saxena, Repeatability of measurements: Non-hermitian observables and quantum Coriolis force, Phys. Rev. A 94, 022121 (2016).
- [61] W. K. Wootters, Entanglement of formation and concurrence, Quantum Inf. Comput. 1, 27 (2001).
- [62] S. Hill and W. K. Wootters, Entanglement of a pair of quantum bits, Phys. Rev. Lett. 78, 5022 (1997).
- [63] W. K. Wootters, Entanglement of formation of an arbitrary state of two qubits, Phys. Rev. Lett. **80**, 2245 (1998).
- [64] V. Vedral, M. B. Plenio, M. A. Rippin, and P. L. Knight, Quantifying entanglement, Phys. Rev. Lett. 78, 2275 (1997).
- [65] T. P. Le and A. Olaya-Castro, Strong Quantum Darwinism and strong independence are equivalent to spectrum broadcast structure, Phys. Rev. Lett. 122, 010403 (2019).

# Supplemental Material Eavesdropping on the Decohering Environment: Quantum Darwinism, Amplification, and the Origin of Objective Classical Reality

Akram Touil, 1,2,\* Bin Yan, 2,3 Davide Girolami, Sebastian Deffner, 1,5 and Wojciech Hubert Zurek 1 Department of Physics, University of Maryland, Baltimore County, Baltimore, MD 21250, USA 2 Center for Nonlinear Studies, Los Alamos National Laboratory, Los Alamos, New Mexico 87545 3 Theoretical Division, Los Alamos National Laboratory, Los Alamos, New Mexico 87545 4 Politecnico di Torino, Corso Duca degli Abruzzi 24, Torino, 10129, Italy 5 Instituto de Física 'Gleb Wataghin', Universidade Estadual de Campinas, 13083-859, Campinas, São Paulo, Brazil

Within the framework of Quantum Darwinism, the emergence of classicality from the quantum substrate is a direct consequence of the redundant imprinting of classical information, about the system of interest  $\mathcal{S}$ , in different fragments  $\mathcal{F}$  of the environment. This classical information, obtained by eavesdropping on environmental fragments, is upper bounded by the Holevo quantity  $\chi(\mathcal{S}: \check{\mathcal{F}})$  (aka the Holevo bound), which can be expressed as

$$\chi(\mathcal{S}: \check{\mathcal{F}}) = H_{\mathcal{S}} - H_{\mathcal{S}|\check{\mathcal{F}}},\tag{S1}$$

such that  $H_i = -\mathrm{tr} \{ \rho_i \log_2(\rho_i) \}$  stands for the von Neumann entropy, and  $H_{\mathcal{S}|\tilde{\mathcal{F}}}$  is the conditional von Neumann entropy [R1] that accounts for the missing information about the system  $\mathcal{S}$  after optimal measurements are applied on the fragment  $\mathcal{F}$ .

In the present supplemental material, we provide the technical details leading to analytic expressions of  $\chi(\mathcal{S}:\tilde{\mathcal{F}})$ , in the physical model we described in the manuscript [R2]. We separate the supplemental material into five sections. First, we present a general overview of the Koashi-Winter relation as the main tool of our derivation. We then show how it applies to our specific physical model of a central qubit undergoing decoherence in a many-qubit environment. The second part of the supplemental material details the necessary steps leading to analytic expressions of the Holevo bound (in the general case and in the limit of good decoherence). Using this result, we illustrate the statements made in the manuscript regarding the redundancy of information and the breaking of the universal behavior of both the mutual information and the Holevo bound. Finally, in the final section we explicitly explore the connection, through the quantum mutual information, between our many-qubit model and the photon scattering model. The latter connection shows that our analysis captures universal attributes in cases when decoherence is caused by environments composed of noninteracting subsystems such as photons.

## THE KOASHI-WINTER RELATION

For any tripartite system ABC, such that  $\rho_{ABC}$  is pure, the Koashi-Winter relation can be written as [R3]

$$\chi(A:\check{B}) + E(A:C) = H_A,\tag{S2}$$

where "E(A:C)" refers to the entanglement of formation [R4–R6] of  $\rho_{AC}$ , which is an entanglement measure for bipartite mixed quantum states (defined as an extension of the entanglement entropy to mixed states). To elaborate, the entanglement of formation "E(A:C)" quantifies the amount of entanglement in the mixed state  $\rho_{AC}$ , since for such cases (i.e. for mixed states) the entanglement entropy is no longer a viable measure of entanglement. The latter is due to the fact that the von Neumann entropy is not symmetric for general mixed states [R1].

The Koashi-Winter relation (S2) represents a monogamy relation between the entanglement of formation and the classical correlations in an extended Hilbert space. In other words, this relation captures the trade-off between entanglement and classical correlations. For a given value of the von Neumann entropy  $H_A$ , the higher the entanglement subsystem "A" shares with "C", the less classical information about "A" is accessible through optimal measurements on "B".

For two-qubit states  $\rho_{AC}$ , the entanglement of formation can be expressed as a function of the concurrence [R6]

$$E(A:C) = h\left(\frac{1 + \sqrt{1 - (\text{Con}(A:C))^2}}{2}\right),$$
 (S3)

where "Con (A:C)" stands for the concurrence of the state  $\rho_{AC}$ , and  $h(x) = -x \log_2(x) - (1-x) \log_2(1-x)$ . Furthermore, the entanglement measure [R6, R7] "Con (A:C)" can be analytically determined by computing the eigenvalues

 $\mu_i$  of the 4x4 density matrix  $\rho_{AC}\tilde{\rho}_{AC}$ , where  $\tilde{\rho}_{AC} = (\sigma_y \otimes \sigma_y) \, \rho_{AC}^* \, (\sigma_y \otimes \sigma_y)$ . For two-qubit states where  $\mu_i \geq \mu_{i+1}$ , the analytic formula reads [R5]

$$Con(A:C) = \max\{0, \sqrt{\mu_1} - \sqrt{\mu_2} - \sqrt{\mu_3} - \sqrt{\mu_4}\}.$$
 (S4)

In the physical model of a central qubit S undergoing decoherence in a many-qubit environment E, assuming that the state of the quantum universe SE is pure implies that, by decomposing the environment into typical fragments, the state  $\rho_{SF_mF_{N-m}}$  is pure. Therefore, in this model, the Koashi-Winter relation is written as

$$\chi(\mathcal{S}: \check{\mathcal{F}}_m) + E(\mathcal{S}: \mathcal{F}_{N-m}) = H_{\mathcal{S}}.$$
 (S5)

Based on the above expression, in order to determine the Holevo bound we need to evaluate the entanglement of formation. In fact, given the structure of the branching state of the dynamics [R2]

$$|\Psi_{\mathcal{S}\mathcal{E}}\rangle = \sqrt{p}|0_{\mathcal{S}}\rangle \bigotimes_{i=1}^{N} |0_{\mathcal{E}_{i}}\rangle + \sqrt{q}|1_{\mathcal{S}}\rangle \bigotimes_{i=1}^{N} |1_{\mathcal{E}_{i}}\rangle , \tag{S6}$$

tracing out the degrees of freedom of a given partition of the environment, or the system of interest, results in a density matrix of rank-two at most, i.e. the resulting density matrices are regarded as virtual qubits [R8]. This implies that the bipartite system " $\mathcal{SF}_{N-m}$ " is a qubit-virtual qubit pair. Therefore, we have

$$E\left(S:\mathcal{F}_{N-m}\right) = h\left(\frac{1+\sqrt{1-\left(\operatorname{Con}\left(S:\mathcal{F}_{N-m}\right)\right)^{2}}}{2}\right). \tag{S7}$$

Additionally, since we are dealing with rank-two (at most) density matrices, when computing the concurrence there are only two nonzero eigenvalues ( $\mu_1$  and  $\mu_2$ ), which implies that Eq. (S4) simplifies to

$$\operatorname{Con}\left(\mathcal{S}:\mathcal{F}_{N-m}\right) = \left|\sqrt{\mu_1} - \sqrt{\mu_2}\right|. \tag{S8}$$

In summary, to determine the Holevo bound  $\chi(S: \hat{\mathcal{F}}_m)$  we first need to evaluate  $\mu_1$  and  $\mu_2$  from the expression of the 4x4 density matrix  $\rho_{S\mathcal{F}_{N-m}}$ . From these eigenvalues we can determine the concurrence (cf. Eq. (S8)), which directly results in an analytic expression for the entanglement of formation (cf. Eq. (S7)). Thus, we can infer the general form of the Holevo bound  $\chi(S: \hat{\mathcal{F}}_m)$ , and the corresponding expression in the good decoherence limit (presented in Equation (20) of the manuscript [R2]).

In what follows, we present the technical details that led to the expressions of  $\mu_1$  and  $\mu_2$ . We start with instructive case of m=1, and then generalize to arbitrary values of m such that  $1 \le m < N$ .

## THE HOLEVO BOUND

Single qubit of the environment  $\mathcal{F}_1$ : We split the branching state  $|\Psi_{\mathcal{S}\mathcal{E}}\rangle$  into two partitions. The first is composed of the central qubit  $\mathcal{S}$  and a single qubit of  $\mathcal{E}$ , while the second partition is the rest of the environment. We adopt the following notations,

$$|\mathbf{0}\rangle \equiv \bigotimes_{i=1}^{N-1} |0_{\mathcal{E}_i}\rangle, \ |\mathbf{1}\rangle \equiv \frac{1}{\mathcal{N}} \left( \bigotimes_{i=1}^{N-1} |1_{\mathcal{E}_i}\rangle - s^{N-1} \bigotimes_{i=1}^{N-1} |0_{\mathcal{E}_i}\rangle \right),$$

$$|\bar{0}\rangle \equiv |00\rangle, \ |\bar{1}\rangle \equiv s|10\rangle + c|11\rangle,$$
(S9)

with  $\mathcal{N}=\sqrt{1-s^{2(N-1)}}$ , and

$$(\forall i \in [1, N]); \ |0_{\mathcal{E}_i}\rangle \equiv |0\rangle, \ |1_{\mathcal{E}_i}\rangle \equiv s|0\rangle + c|1\rangle. \tag{S10}$$

The real parameters c and s, quantify the degree with which the environmental qubits monitor the central qubit  $\mathcal{S}$ . From the above definitions, we have  $\langle i|j\rangle=\delta_{i,j}$  and  $\langle \overline{i}|\overline{j}\rangle=\delta_{i,j}$  for  $(i,j)\in\{0,1\}^2$ , and the state of the universe  $\mathcal{S}\mathcal{E}$  can be written as

$$|\Psi_{\mathcal{S}\mathcal{E}}\rangle = \sqrt{p}|\overline{\mathbf{0}}\mathbf{0}\rangle + \sqrt{q}s^{N-1}|\overline{\mathbf{1}}\mathbf{0}\rangle + \sqrt{q}\sqrt{1 - s^{2(N-1)}}|\overline{\mathbf{1}}\mathbf{1}\rangle.$$
 (S11)

Therefore, the rank-two density matrix representing the composite state of the central qubit S and a single qubit of the environment  $F_1$  is

$$\rho_{\mathcal{SF}_1} = \begin{pmatrix} p & 0 & s^N \sqrt{pq} & s^{N-1} c \sqrt{pq} \\ 0 & 0 & 0 & 0 \\ s^N \sqrt{pq} & 0 & s^2 q & scq \\ s^{N-1} c \sqrt{pq} & 0 & scq & c^2 q \end{pmatrix}.$$
 (S12)

Now we generalize to the case of an arbitrary environmental fragment  $\mathcal{F}_m$  (with m qubits) in order to determine the density matrix  $\rho_{\mathcal{SF}_m}$ , for any  $1 \leq m < N$ , which directly implies the expression of  $\rho_{\mathcal{SF}_{N-m}}$  and the corresponding eigenvalues  $\mu_1$  and  $\mu_2$ .

Environmental fragment  $\mathcal{F}_m$ : Considering a partition of the environment (with m qubits), and similar to the previous single qubit analysis (cf. Eq. (S9)), we group the central qubit with "m" qubits from the environment to get

$$|\bar{0}\rangle \equiv |0\rangle \bigotimes_{i=1}^{m} |0_{\mathcal{E}_i}\rangle, \ |\bar{1}\rangle \equiv |1\rangle \bigotimes_{i=1}^{m} |1_{\mathcal{E}_i}\rangle,$$
 (S13)

which in turn can be written as a basis for "the qubit-virtual qubit" pair

$$|\overline{0}\rangle \equiv |00\rangle, |\overline{1}\rangle \equiv |1\rangle \left(s^m |0\rangle + \sqrt{1 - s^{2m}} |1\rangle\right),$$
 (S14)

with

$$|\mathfrak{o}\rangle \equiv \bigotimes_{i=1}^{m} |0_{\mathcal{E}_i}\rangle, \ |\mathfrak{1}\rangle \equiv \frac{1}{\mathcal{M}} \left( \bigotimes_{i=1}^{m} |1_{\mathcal{E}_i}\rangle - s^m \bigotimes_{i=1}^{m} |0_{\mathcal{E}_i}\rangle \right), \tag{S15}$$

such that  $\mathcal{M} = \sqrt{1 - s^{2m}}$  and  $\langle \mathfrak{i} | \mathfrak{j} \rangle = \delta_{\mathfrak{i},\mathfrak{j}}$ . Following this "qubit-virtual qubit" decomposition, we can infer the expression of the rank-two density matrix  $\rho_{\mathcal{SF}_{N-m}}$ ,

$$\rho_{\mathcal{S}\mathcal{F}_{N-m}} = \begin{pmatrix} p & 0 & s^N \sqrt{pq} & s^m \sqrt{1 - s^{2(N-m)}} \sqrt{pq} \\ 0 & 0 & 0 & 0 \\ s^N \sqrt{pq} & 0 & qs^{2(N-m)} & qs^{N-m} \sqrt{1 - s^{2(N-m)}} \\ s^m \sqrt{1 - s^{2(N-m)}} \sqrt{pq} & 0 & qs^{N-m} \sqrt{1 - s^{2(N-m)}} & q(1 - s^{2(N-m)}) \end{pmatrix}.$$
 (S16)

The eigenvalues,  $\mu_1$  and  $\mu_2$ , of the matrix  $\rho_{\mathcal{SF}_{N-m}}\tilde{\rho}_{\mathcal{SF}_{N-m}}$ , are

$$\mu_1 = pq (1 - s^m)^2 (1 - s^{2(N-m)}), \quad \mu_2 = pq (1 + s^m)^2 (1 - s^{2(N-m)}).$$
 (S17)

From the above eigenvalues, and based on Eqs. (S5), (S7) and (S8), we get the general analytic expression for the Holevo bound,

$$\chi(\mathcal{S}: \check{\mathcal{F}}_m) = h(r_p) - h(t_{p,m}), \tag{S18}$$

where

$$r_{p} = \frac{1}{2} \left( 1 + \sqrt{1 - 4pq \left( 1 - s^{2N} \right)} \right) \text{ and } t_{p,m} = \frac{1}{2} \left( 1 + \sqrt{1 - 4pq \left( s^{2m} - s^{2N} \right)} \right). \tag{S19}$$

From this expression, it is a simple exercise to show that for good decoherence (i.e. in the limit  $s^{N-m} \ll s^m$ ) we have the following simplification

$$\chi(\mathcal{S}: \check{\mathcal{F}}_m) = H_{\mathcal{S}} + \frac{1}{2}\log_2\left(pqs^{2m}\right) + \sqrt{1 - 4pqs^{2m}}\operatorname{Arctanh}_2\left(\sqrt{1 - 4pqs^{2m}}\right),\tag{S20}$$

such that  $H_{\mathcal{S}} = -p \log_2(p) - q \log_2(q)$  is the von Neumann entropy of the decohered state of the central qubit  $\mathcal{S}$ .

In the subsequent two sections, based on the above result, we will illustrate the following statements made in the manuscript. The first statement is that the redundancy evaluated through the quantum mutual information  $\mathcal{I}$  overestimates the redundancy from the viewpoint of observers eavesdropping on fragments  $\mathcal{F}$  of the environment via optimal measurements. The difference between the two quantities can be computed numerically. The second statement concerns the weak dependence of the re-scaled quantum mutual information and Holevo bounds on the state preparation of the central qubit  $\mathcal{S}$ , i.e. on the probability of the pointer states "p".

## REDUNDANCY

In Fig. S1, we illustrate the behavior of the redundancy  $R_{\delta}=N/m_{\delta}$  (with information deficit  $\delta=0.1$ ), evaluated using I (red curve) and  $\mathcal X$  (black curve), as a function of the parameter c. We observe that the redundancy is monotonically increasing as a function of c, such that the maximum redundancy is achieved when  $c\to 1$ . The latter is due to the fact that in the limit where  $c\to 1$  the coupling between the central qubit and each qubit from the environment is now modeled by the c-not gate, and the branching state is a GHZ state. We also observe that for almost all values of  $c\in [0,1]$  the redundancy computed using the quantum mutual information is an overestimate to the redundancy observed by eavesdropping on fragments of the environment through optimal measurements.




FIG. S1. Plots of the redundancy  $R_{0.1}$  computed through the mutual information I (red curve) and the Holevo bound  $\chi$  (black curve), as a function of c, for p = 1/2 and N = 100.

Using the analytic expressions of the mutual information and the Holevo bound, we can numerically determine the degree with which the redundancy obtained from the mutual information overestimates the actual redundancy. We can define  $\Delta m = m^{\chi} - m^{I}$ , such that  $m^{\chi}$  ( $m^{I}$ ) represents the minimum number of environmental qubits needed in order for the Holevo bound (mutual information) to reflect the missing information about the central qubit S, up to an information deficit  $\delta$ , i.e.,  $(1 - \delta)H_{S}$ . Therefore, the quantity  $\Delta m/m^{\chi}$  reflects the difference between the redundancies obtained from the two different information theoretic measures (I and  $\chi$ ). In Fig. S2, we illustrate, in the limit of large N ( $N \to \infty$ ),  $\Delta m/m^{\chi}$  as a function of  $\delta$  and p. In panel (a), we observe that the maximum difference, for a given value of  $\delta$ , is attained for p = 1/2. From panel (b), in the limit where  $\delta \to 0$  we conclude that the redundancies differ by at most  $\sim 13\%$ , and for  $\delta = 0.2$  (similar to the case presented in the inset of Figure 1 of the manuscript [R2]) the difference is at most  $\sim 37\%$ .




FIG. S2. Plots of  $\Delta m/m^{\chi}$  as a function of  $\delta \in [0, 0.2]$  and p. In panel (a), we plot  $\Delta m/m^{\chi}$  for all values of  $p \in [0.01, 0.99]$ , and in panel (b) we focus on the case of p = 1/2.

### BREAKING OF UNIVERSALITY

In the manuscript, we show that the quantum mutual information and the Holevo bound display a universal scaling behavior which is weakly dependent on the state preparation of the central qubit S. Namely, through appropriate rescaling, the two information theoretic quantities exhibit a universal rising behavior insensitive to the value of the parameter p, up to some limitations. Here, our goal is to quantify and illustrate these limitations, which shows the breaking of the aforementioned universality when the parameter p approaches a value of zero or one. To this end, we choose the case of the initial equal superposition (p=1/2) as a reference, and we compare its corresponding expressions of the mutual

information and the Holevo bound with those for arbitrary p', which leads to defining  $\Delta_I$  and  $\Delta_{\chi}$  as follows.

$$\Delta_I = I_{p=1/2}(\mathcal{S}: \mathcal{F}_m) - \frac{I_{p'}(\mathcal{S}: \mathcal{F}_m)}{H_{p'}(\mathcal{S})}, \tag{S21}$$

such that  $I_{p=1/2}(S:\mathcal{F}_m)$  and  $I_{p'}(S:\mathcal{F}_m)$  are the values of the mutual information for p=1/2 and arbitrary  $p'\in[0,1]$ , respectively. The quantity  $H_{p'}(S)$  is the maximum von Neumann entropy of the system S corresponding to p' (i.e. the plateau of the curve of  $I_{p'}(S:\mathcal{F}_m)$ ). Similarly, for the Holevo bound we define the quantity  $\Delta_{\chi}$  such that

$$\Delta_{\chi} = -\chi_{p=1/2}(\mathcal{S} : \check{\mathcal{F}}_m) + \frac{\chi_{p'}(\mathcal{S} : \check{\mathcal{F}}_m)}{H_{p'}(\mathcal{S})}.$$
(S22)

In Fig. S3, we plot  $\Delta_I$  and  $\Delta_\chi$  as functions of both m and p'. For small m, the breaking of the universal rise of both the mutual information and the Holevo bound is observed when p' is either close to zero or one, otherwise  $\Delta_I \approx 0$  and  $\Delta_\chi \approx 0$ . For completeness, we also plot our quantities for large values of m (i.e. when the environment fraction approaches one). In this limit,  $\Delta_\chi = 0$  for all p' while  $\Delta_I \neq 0$  for p' close to zero or one. This is due to the fact that the quantum mutual information rises beyond the plateau when we capture almost all of the environment, which is a direct consequence of the purity of the state of the universe  $\mathcal{SE}$ .




FIG. S3. Plots of  $\Delta_I$  (panel (a)) and  $\Delta_X$  (panel (b)), as a function of m and p', for N=100 and  $c=\sqrt{0.4}$ . Note that changing the value of the parameter c for a fixed total number of qubits N is equivalent to a straightforward re-scaling of the x-axis of our plots (displaying the number of qubits "m" in a typical environmental fragment), with  $\log(s)$  as the re-scaling factor [R2]. It is also noteworthy that the initial rise of the mutual information or the Holevo bound is independent of the total number of qubits N in our environment [R2].

#### CONNECTION TO THE PHOTON MODEL

As mentioned towards the end of the manuscript [R2], our results naturally extend to the realistic model of a photon environment, as studied in Refs. [R9, R10]. In fact, the quantum mutual information expression is the same as the one derived in the photon scattering model. More specifically, if we consider a dielectric sphere, as our system of interest  $\mathcal{S}$ , initially in a spatial superposition such that  $|\psi_{\mathcal{S}}\rangle = \sqrt{p}\delta\left(\vec{x}-\vec{x}_1\right) + \sqrt{q}\delta\left(\vec{x}-\vec{x}_2\right)$ . The environment is composed of N photons, originally emitted from a point source (for simplicity we assume point source illumination). In this case, it was shown [R9, R10] that the quantum mutual information has the following form,

$$I(S: \mathcal{F}_m) = 1 + \frac{1}{\log(2)} \sum_{i=1}^{\infty} \frac{\Gamma^{(1-f)i} - \Gamma^{fi} - \Gamma^i}{2i(2i-1)},$$
(S23)

where  $\Gamma^x=(p-q)^2+4pq\exp(-tx/\tau_D)$  such that  $x\in\{1,f,1-f\}$ ,  $\tau_D$  is the decoherence time, and f=m/N is the fraction of photons we access. The above expression is in direct agreement with our the results [R2]. In particular, from the specific form of equations (5-7) of the manuscript we can compute the analytic expression of the von Neumann entropies of the states  $\rho_{\mathcal{S}}$ ,  $\rho_{\mathcal{F}_m}$ , and  $\rho_{\mathcal{S}\mathcal{F}_m}$ ,

$$H_{\mathcal{S}} = -\lambda_{-}(N, p) \log_{2}(\lambda_{-}(N, p)) - \lambda_{+}(N, p) \log_{2}(\lambda_{+}(N, p)),$$
(S24)

$$H_{\mathcal{F}_m} = -\lambda_{-}(m, p) \log_2 (\lambda_{-}(m, p)) - \lambda_{+}(m, p) \log_2 (\lambda_{+}(m, p)),$$
 (S25)

$$H_{\mathcal{SF}_{m}} = -\lambda_{-}(N-m, p) \log_{2} (\lambda_{-}(N-m, p)) - \lambda_{+}(N-m, p) \log_{2} (\lambda_{+}(N-m, p)),$$
 (S26)

with  $\lambda_{\pm}(k,p)=\frac{1}{2}\left(1\pm\sqrt{\left(q-p\right)^{2}+4s^{2k}pq}\right)$ . Generally, we have, for |x|<1,

$$\log_2(1+x) = \frac{1}{\log(2)} \sum_{i=1}^{\infty} (-1)^{i+1} \frac{x^i}{i}, \text{ and } \log_2(1-x) = -\frac{1}{\log(2)} \sum_{i=1}^{\infty} \frac{x^i}{i}. \tag{S27}$$

Therefore,

$$-(1-x)\log_2(1-x) - (1+x)\log_2(1+x) = -\frac{2}{\log(2)} \sum_{i=1}^{\infty} \frac{x^{2i}}{2i(2i-1)}.$$
 (S28)

It is, then, a straightforward exercise to get

$$I(S: \mathcal{F}_m) = 1 + \frac{1}{\log(2)} \left( \sum_{i=1}^{\infty} \frac{X_{N-m}^{2i} - X_m^{2i} - X_N^{2i}}{2i(2i-1)} \right), \tag{S29}$$

such that

$$X_i = \sqrt{(p-q)^2 + 4s^{2i}pq} \le 1. {(S30)}$$

Comparing this expression of the quantum mutual information with that of the photon model (cf. Eq. (S23)) we recognize the exact similarity such that  $s^N \equiv \exp(-t/2\tau_D)$ . Therefore, the universal rise (one of our main results) of the quantum mutual information also applies to the realistic photon scattering model.

## \* akramt1@umbc.edu

- [R1] M. A. Nielsen and I. Chuang, Quantum computation and quantum information (2002).
- [R2] A. Touil, B. Yan, D. Girolami, S. Deffner, and W. H. Zurek, Eavesdropping on the decohering environment: Quantum darwinism, amplification, and the origin of objective classical reality, arXiv preprint (2021).
- [R3] M. Koashi and A. Winter, Monogamy of quantum entanglement and other correlations, Phys. Rev. A 69, 022309 (2004).
- [R4] S. Hill and W. K. Wootters, Entanglement of a pair of quantum bits, Phys. Rev. Lett. 78, 5022 (1997).
- [R5] W. K. Wootters, Entanglement of formation of an arbitrary state of two qubits, Phys. Rev. Lett. 80, 2245 (1998).
- [R6] W. K. Wootters, Entanglement of formation and concurrence, Quantum Inf. Comput. 1, 27 (2001).
- [R7] V. Vedral, M. B. Plenio, M. A. Rippin, and P. L. Knight, Quantifying entanglement, Phys. Rev. Lett. 78, 2275 (1997).
- [R8] Here, the notion of virtual qubit is a mathematical construct where the corresponding states can be mapped to a Hilbert space of dimension equal to two.
- [R9] C. J. Riedel and W. H. Zurek, Quantum darwinism in an everyday environment: Huge redundancy in scattered photons, Phys. Rev. Lett. **105**, 020404 (2010).
- [R10] C. J. Riedel and W. H. Zurek, Redundant information from thermal illumination: quantum darwinism in scattered photons, New J. Phys. 13, 073038 (2011).