

MULTIPLE SOLUTIONS FOR A CLASS OF QUASILINEAR PROBLEMS WITH DOUBLE CRITICALITY

KARIMA AIT-MAHIOUT, CLAUDIANOR O. ALVES AND PRASHANTA GARAIN

ABSTRACT. We establish multiplicity results for the following class of quasilinear problems

$$(P) \quad \begin{cases} -\Delta_{\Phi} u = f(x, u) & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

where $\Delta_{\Phi} u = \operatorname{div}(\varphi(x, |\nabla u|) \nabla u)$ for a generalized N-function $\Phi(x, t) = \int_0^{|t|} \varphi(x, s) s \, ds$. We consider $\Omega \subset \mathbb{R}^N$ to be a smooth bounded domain that contains two disjoint open regions Ω_N and Ω_p such that $\overline{\Omega_N} \cap \overline{\Omega_p} = \emptyset$. The main feature of the problem (P) is that the operator $-\Delta_{\Phi}$ behaves like $-\Delta_N$ on Ω_N and $-\Delta_p$ on Ω_p . We assume the nonlinearity $f : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ of two different types, but both behaves like $e^{\alpha|t|^{\frac{N}{N-1}}}$ on Ω_N and $|t|^{p^*-2}t$ on Ω_p as $|t|$ is large enough, for some $\alpha > 0$ and $p^* = \frac{Np}{N-p}$ being the critical Sobolev exponent for $1 < p < N$. In this context, for one type of nonlinearity f , we provide multiplicity of solutions in a general smooth bounded domain and for another type of nonlinearity f , in an annular domain Ω , we establish existence of multiple solutions for the problem (P) that are nonradial and rotationally nonequivalent.

1. INTRODUCTION

In this paper, we establish the existence of multiple solutions for the following class of quasilinear problems

$$(P) \quad \begin{cases} -\Delta_{\Phi} u = f(x, u) & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

where $\Omega \subset \mathbb{R}^N$, with $N \geq 2$, is a smooth bounded domain, $\Delta_{\Phi} u = \operatorname{div}(\varphi(x, |\nabla u|) \nabla u)$ is the Φ Laplace operator, where $\Phi(x, t) = \int_0^{|t|} \varphi(x, s) s \, ds$, $\varphi : \Omega \times [0, +\infty) \rightarrow [0, +\infty)$ and $f : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ are continuous functions that satisfy some hypothesis that will be mentioned later on.

Before proceeding further, let us go through some known results associated with the Φ Laplace equations. In the recent past, the study of such equations concerning the existence theory has been a research topic of considerable attention. Such operator extends the p -Laplace operator, the variable exponent p -Laplace operator, weighted p -Laplace operator, p, q -Laplace operator and indeed occur in many physical phenomena.

When Φ is independent of x , solutions of (P) are investigated in the Orlicz-Sobolev space [44] and we refer the reader to Alves, Figueiredo and Santos [4], Fukagai, Ito and Narukawa [29], Carvalho, Silva, Gonçalves and Goulart [15], Fukagai and Narukawa [30], Harjulehto and Hästö [33], and their references for the study of such PDEs. When Φ also depends on x , we are led to study the problems in variable exponent Sobolev [24, 38] or Musielak-Sobolev spaces [20, 35, 43, 44]. Differential equations in variable exponent Sobolev spaces have been studied extensively in the last years, most part of them involves the $p(x)$ -Laplacian operator, see for example, Alves and Barreiro [2], Alves and Ferreira [3], Alves and Souto [6], Alves

2010 *Mathematics Subject Classification.* 35A15; 35J62, 46E30.

Key words and phrases. Variational methods, Quasilinear problems, Musielak-Sobolev space.

C. O. Alves was partially supported by CNPq/Brazil 304804/2017-7 .

and Rădulescu [8], Chabrowski and Fu [19], Fan and Zhang [27], Fan [28], Rădulescu and Repovš [46] and their references. However, Differential equations in general Musielak-Sobolev spaces have been studied very little, see for instance, Azroul, Benkirane, Shimi and Srati [10], Benkirane and Sidi El Vally [11], Fan [26], Liu and Zhao [41], Wang and Liu [51] and the references therein.

In the present paper, we will apply some recent results involving the Musielak-Sobolev spaces to study the existence of nontrivial solution for the problem (P). Next, we will state our main hypothesis on the functions Φ and φ :

- (φ_1) For each $x \in \Omega$, $\varphi(x, \cdot)$ is a C^1 function in the interval $(0, +\infty)$.
- (φ_2) $\varphi(x, t)$, $\partial_t(\varphi(x, t)t) > 0$, for $x \in \Omega$ and $t > 0$.
- (φ_3) There exists $1 < p < N < q < p^*$ such that

$$p \leq \frac{\varphi(x, |t|)|t|^2}{\Phi(x, |t|)} \leq q, \text{ for } x \in \Omega \text{ and } t \neq 0.$$

Arguing as in Fukagai, Ito and Narukawa [29], it is possible to prove with few modifications that if φ satisfies the conditions (φ_1) – (φ_3), then the function Φ is a generalized N-function.

The complementary function $\tilde{\Phi}$ associated with Φ is given by the Legendre's transformation, that is,

$$(1.1) \quad \tilde{\Phi}(x, s) = \max_{t \geq 0} \{st - \Phi(x, t)\}, \quad x \in \Omega \text{ and } s \in \mathbb{R}.$$

The functions Φ and $\tilde{\Phi}$ are complement of each other and $\tilde{\Phi}$ is also a generalized N-function. Hereafter, we also assume that for some constant d_1 ,

- (φ_4) $\inf_{x \in \Omega} \Phi(x, 1) = d_1 > 0$.
- (φ_5) For each $t_0 \neq 0$, there is a constant $c_0 > 0$ such that

$$\frac{\Phi(x, t)}{t} \geq c_0 \text{ and } \frac{\tilde{\Phi}(x, t)}{t} \geq c_0 \text{ for } t \geq t_0 \text{ and } x \in \Omega.$$

The conditions (φ_1) – (φ_5) are very important in our approach, because they permit us to conclude that the Musielak-Orlicz space $L^\Phi(\Omega)$ and Musielak-Sobolev space $W^{1, \Phi}(\Omega)$ are reflexive and separable Banach spaces, for more details see Section 2.

In a recent paper, Alves, Garain and Rădulescu [7] proved the existence of at least one nontrivial solution for the following prototype problem

$$(Q) \quad \begin{cases} -\Delta_\Phi u = f(x, u) & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

provided Ω is a smooth bounded domain in \mathbb{R}^N with $N \geq 2$, f is a continuous function, $\varphi : \Omega \times [0, +\infty) \rightarrow [0, +\infty)$ satisfies the hypothesis (φ_1) – (φ_5) above (see [7] for precise assumptions).

In the present paper, as in [7], Ω satisfies the following conditions: There are three smooth domains $\Omega_N, \Omega_q, \Omega_p \subset \Omega$ with nonempty interior such that

$$\Omega = \Omega_N \cup \Omega_q \cup \Omega_p$$

and there is $\delta > 0$ such that

$$(\overline{\Omega_N})_\delta \cap (\overline{\Omega_p})_\delta = \emptyset.$$

Hereafter, if $A \subset \Omega$, we denote by A_δ to be the δ -neighborhood of A restricted to Ω , that is,

$$A_\delta = \{x \in \Omega : \text{dist}(x, A) < \delta\}.$$

Associated with the sets Ω_N, Ω_q and Ω_p , there are three continuous functions $\eta_N, \eta_q, \eta_p : \overline{\Omega} \rightarrow [0, 1]$ satisfying:

$$\begin{aligned}\eta_N(x) &= 1, & \forall x \in \overline{\Omega_N}, \\ \eta_p(x) &= 1, & \forall x \in \overline{\Omega_p},\end{aligned}$$

and

$$\eta_q(x) = 1, \quad \forall x \in \Omega_q = \Omega \setminus \overline{(\Omega_N \cup \Omega_p)},$$

$$\eta_N(x) = 0, \quad \forall x \in (\overline{\Omega_N})_\delta^c, \quad \eta_p(x) = 0, \quad \forall x \in (\overline{\Omega_p})_\delta^c, \quad \eta_q(x) > 0, \quad \forall x \in (\overline{\Omega_q})_\delta, \quad \eta_q(x) = 0, \quad \forall x \in (\overline{\Omega_q})_\delta^c$$

and for some positive constant c_4 ,

$$\eta_q(x) \leq c_4 \text{dist}(x, \partial(\Omega_q)_\delta \cap \Omega_p)^l, \quad \forall x \in \overline{\Omega_p} \cap (\Omega_q)_\delta,$$

where $l > q$ and $\text{dist}(x, \partial(\Omega_q)_\delta \cap \Omega_p) = \inf\{|x - y| : y \in \partial(\Omega_q)_\delta \cap \Omega_p\}$.

Related to the function $f : \overline{\Omega} \times \mathbb{R} \rightarrow \mathbb{R}$ we assume that it is a continuous function of one the the following forms:

(f₁)

$$f(x, t) = \lambda \eta_N(x) |t|^{\beta-2} t e^{\alpha|t|^{\frac{N}{N-1}}} + \mu \tilde{\eta}_q(x) g(x, t) + \eta_p(x) (\tau |t|^{\zeta-2} t + |t|^{p^*-2} t), \quad \forall (x, t) \in \Omega \times \mathbb{R},$$

or

$$(f_2) \quad f(x, t) = \lambda \eta_N(x) |t|^{\beta-2} t e^{\alpha|t|^{\frac{N}{N-1}}} + \tilde{\eta}_q(x) g(x, t) + \eta_p(x) |t|^{p^*-2} t, \quad \forall (x, t) \in \Omega \times \mathbb{R},$$

where λ, μ, τ are positive parameters, $\alpha > 0$, $p^* > \zeta > q > N > p > \frac{N}{2}$, $\beta > q$, where $p^* = \frac{Np}{N-p}$, $g : \overline{\Omega} \times \mathbb{R} \rightarrow \mathbb{R}$ and $\tilde{\eta}_q : \overline{\Omega} \rightarrow [0, 1]$ are continuous functions such that

$$\tilde{\eta}_q(x) = 1, \quad \forall x \in \Omega_q = \Omega \setminus \overline{(\Omega_N \cup \Omega_p)}$$

and

$$\tilde{\eta}_q(x) = 0, \quad \forall x \in (\overline{\Omega_q})_{\delta/2}^c.$$

Related to the function g , we assume the following conditions:

(g₀) g is odd with respect to the second variable t ,

(g₁) $g(x, t) = o(|t|^{q_1-1})$, as $t \rightarrow 0$, uniformly in $x \in (\overline{\Omega_q})_{\delta/2}$

for some $q_1 > q$ and there is $\theta > q$ such that

(g₂) $0 < \theta G(x, t) \leq g(x, t)t$, $\forall x \in (\overline{\Omega_q})_{\delta/2}$

where $G(x, t) = \int_0^t g(x, s) ds$, for $t \in \mathbb{R}$.

There exists a constant $c > 0$, such that

(g₃) $g(x, t) \geq ct^{q_2-1}$, $\forall t \geq 0, \forall x \in \Omega_q$,

for some $q_2 > q_1$,

With these notations, we are ready to mention the last conditions on φ . If f is the form (f₁), we assume for each $t > 0$ the following:

(φ_6) $\varphi(x, t) \geq t^{N-2}$, for $x \in \Omega_N$ and $c_1 t^{N-2} \geq \varphi(x, t)$, $x \in \Omega_N \setminus \overline{(\Omega_q)_\delta}$.

(φ_7) $\varphi(x, t) \geq \tau_1(x) t^{q-2}$, for $x \in (\Omega_q)_\delta$ where $\tau_1 : \overline{\Omega} \rightarrow \mathbb{R}$ is a continuous function satisfying:

$$\tau_1(x) > 0, \quad \forall x \in (\Omega_q)_\delta \quad \text{and} \quad \tau_1(x) = 0, \quad \forall x \in ((\Omega_q)_\delta)^c.$$

(φ_8) $\tau_2(x)t^{q-2} + c_2t^{p-2} \geq \varphi(x,t) \geq t^{p-2}$, $x \in \Omega_p$ where $\tau_2 : \overline{\Omega_p} \rightarrow \mathbb{R}$ is a nonnegative continuous function satisfying:

$$\tau_2(x) \leq c_3 \text{dist}(x, \partial(\Omega_q)_\delta \cap \Omega_p)^s, \quad \forall x \in \overline{\Omega_p} \cap (\Omega_q)_\delta$$

for some $s > q$ and

$$\tau_2(x) = 0, \quad \forall x \in \overline{\Omega_p} \setminus \overline{(\Omega_q)_\delta},$$

for some constants $c_i > 0$ with $i = 1, 2, 3$.

Now, if f is the form (f_2), the condition (φ_6) is assumed of the following way:

$$(\varphi_6) \varphi(x,t) \geq t^{N-2}, \quad \text{for } x \in \Omega_N.$$

As a model of a function that satisfies the conditions (φ_1) – (φ_8) is the function $\varphi : \Omega \times [0, +\infty) \rightarrow [0, +\infty)$ defined by

$$(1.2) \quad \varphi(x,t) = \eta_N(x)t^{N-2} + \eta_q(x)t^{q-2} + \eta_p(x)t^{p-2}, \quad \forall (x,t) \in \Omega \times [0, +\infty)$$

and so,

$$(1.3) \quad \Phi(x,t) = \frac{\eta_N(x)}{N}|t|^N + \frac{\eta_q(x)}{q}|t|^q + \frac{\eta_p(x)}{p}|t|^p, \quad \forall (x,t) \in \Omega \times \mathbb{R}.$$

Motivated by the study made in [7], we intend to prove existence of multiple solutions for the problem (Q) working with the same operator under the nonlinearities (f_1) and (f_2). Here we have two main results involving multiple solutions and their motivation are mentioned below.

Our first main result is motivated by the study made in Wei and Wu [52], where the authors showed the existence of multiple solutions for the following class of problems involving the p -Laplacian operator

$$(1.4) \quad \begin{cases} -\text{div}(|\nabla u|^{p-2}\nabla u) = f(x,u) + \lambda|u|^{p^*-2}u & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

where Ω is a bounded domain, λ is a positive parameter and f is a continuous function with subcritical growth and $p^* = \frac{Np}{N-p}$ for $N > p$. Using a version of an abstract theorem due to Ambrosetti and Rabinowitz [9] that involves the genus theory for C^1 even functional, it was proved that given $n \in \mathbb{N}$, there is $\lambda_* = \lambda_*(n) > 0$ such that problem (1.4) has at least n nontrivial solutions for $\lambda \in (0, \lambda_*)$. In [47], Silva and Xavier improved the main results proved in [52].

Here, we proved a version of the above mentioned result for the problem (P) and the statement of our result is the following.

Theorem 1.1. *Assume (g_0) – (g_3), (f_1) and (φ_1) – (φ_8). Then, for each $k \in \mathbb{N}$, there exists positive real numbers λ_k, μ_k and τ_k such that for $\lambda \geq \lambda_k, \mu \geq \mu_k$ and $\tau \geq \tau_k$, the problem (P) has at least k pairs of nontrivial solutions.*

Our second result involves the existence of many rotationally nonequivalent and nonradial solutions. We would like to point out that the existence of many rotationally nonequivalent and nonradial solutions was considered in some problems involving the Laplacian operator. Brézis and Nirenberg [14] proved the existence of nonradial positive solution for the following problem

$$(1.5) \quad \begin{cases} -\Delta u + u - u^p = 0 & \text{in } D, \\ u = 0 & \text{on } \partial D, \end{cases}$$

where

$$D = \{x \in \mathbb{R}^N : r < |x| < r + d\}$$

for some $d > 0$. This type of phenomenon is known as symmetry breaking. In [22], Coffman proved that the number of nonradial and rotationally nonequivalent positive solutions of (1.5) in D tends to $+\infty$ as r tends to $+\infty$, if $p > 1$ and $N = 2$ or $1 < p < N/(N - 2)$ and $N \geq 3$.

Motivated by the above papers, some authors have studied this class of problems. For the subcritical case, we cite the papers of Li [39], Lin [40], Suzuki [50] and references therein.

Related to the critical case, Wang and Willem [53] have showed the existence of multiple solutions for the following problem

$$(1.6) \quad \begin{cases} -\Delta u = \lambda u + u^{2^*-1} & \text{in } \Omega_r, \\ u = 0 & \text{on } \partial\Omega_r, \end{cases}$$

where

$$(1.7) \quad \Omega_r = \{x \in \mathbb{R}^N : r < |x| < r + 1\}.$$

The authors proved that for $0 < \lambda < \pi^2$ and $n \in \mathbb{N}$, there exists $R(\lambda, n)$ such that for $r > R(\lambda, n)$, the equation (1.6) has at least n nonradial and rotationally nonequivalent solutions. Motivated by [53], de Figueiredo and Miyagaki [25] have considered the following problem

$$(1.8) \quad \begin{cases} -\Delta u = f(|x|, u) + u^{2^*-1} & \text{in } \Omega_r, \\ u = 0 & \text{on } \partial\Omega_r, \end{cases}$$

where f is a C^1 function with subcritical growth.

In [5], Alves and de Freitas showed the existence of many rotationally nonequivalent and nonradial solutions for a large class of quasilinear problems that have in particular case the problem below

$$(R_1) \quad \begin{cases} -\Delta_N u = \lambda |u|^{\beta-2} \beta e^{\alpha|u|^{\frac{N}{N-1}}} & \text{in } \Omega_r, \\ u = 0 & \text{on } \partial\Omega_r. \end{cases}$$

Still related to this class of problem, we would like to cite the papers of Byeon [13], Castro and Finan [16], Catrina and Wang [17], Mizoguchi and Suzuki [42], Hirano and Mizoguchi [34] and references therein.

Motivated by bibliography cited above and more precisely, by results found in [5], [25] and [53], we are ready to state our second main result, however we need to fix some more conditions:

$$(\varphi_9) \quad \frac{\varphi(x, t)}{|t|^{q-3}t} \text{ is nonincreasing for } t \neq 0.$$

$$(\varphi_{10}) \quad \Phi \text{ is radial in relation with } x \text{ that is } \Phi(|x|, t) = \Phi(x, t) \text{ for all } t > 0.$$

$$(\varphi_{11}) \quad \text{There exists } \kappa \in (0, \frac{1}{2^{N+1}}) \text{ such that}$$

$$|\partial_s \Phi(s, t)| \leq \kappa \Phi(s, t), \forall (s, t) \in \mathbb{R}^2.$$

(η) The functions $\eta_N, \tilde{\eta}_q, \eta_p$ and g are radial in x , that is

$$\eta_N(x) = \eta_N(|x|), \tilde{\eta}_q(x) = \tilde{\eta}_q(|x|), \eta_p(x) = \eta_p(|x|)$$

and

$$g(x, t) = g(|x|, t),$$

for all $x \in \Omega$ and $t > 0$.

(Ω_N) There is $\delta_1 > 0$ such that

$$\mathcal{A} = \left\{ x \in \mathbb{R}^N : \frac{2r+1}{2} - \delta_1 \leq |x| \leq \frac{2r+1}{2} + \delta_1 \right\} \subset \Omega_N \setminus \overline{(\Omega_q)_\delta}.$$

(g_4) $g(|x|, \cdot)$ is a C^1 function in the interval $(0, +\infty)$ and $\frac{g(|x|, t)}{|t|^{q-1}}$ is increasing for $t \neq 0$ uniformly in $x \in (\overline{\Omega_q})_{\frac{\delta}{2}}$.

The reader is invited to see that Φ given in (1.3) also satisfies $(\varphi_9) - (\varphi_{11})$, provided (η) holds.

Our second main theorem has the following statement.

Theorem 1.2. *Assume $\Omega = \Omega_r$ with $N \geq 2$ and $N \neq 3$. Let $(f_2), (g_1), (g_2), (g_4), (\eta), (\Omega_N)$ and $(\varphi_1) - (\varphi_{11})$ holds. Then, for each $n \in \mathbb{N}$, there exists $r_0 = r_0(n) > 0$ and $\lambda_0 = \lambda_0(n) > 0$ such that for $\lambda \geq \lambda_0$ and $r \geq r_0$, the problem (P) has at least n nonradial, rotationally nonequivalent and nontrivial solutions.*

1.1. Our approach: To prove our main results (Theorem 1.1-1.2), we use variational methods. More precisely, for the proof of Theorem 1.1, we follow the approach from Wei and Wu [52] and Silva and Xavier [47]. To this end, we use a result from Ambrosetti and Rabinowitz (see Lemma 4.1). To obtain Theorem 1.2, we adapt for our problem some ideas found in de Figueiredo and Miyagaki [25] and Alves and de Freitas [5]. Here, we establish a Poincaré type inequality (Lemma 2.3) and a Strauss type result (Lemma 5.4) in the setting of Musielak-Sobolev spaces.

It is worth mentioning that, due to the double critical behavior, the energy functionals associated with the problem (P) do not satisfy the (PS) -condition at some levels, which brings some difficulties to apply variational methods. To overcome such difficulties, we closely follow the approach introduced in [7], where one needs to simultaneously employ the concentration compactness Lemma due to Lions in $W^{1,p}(\Omega_p)$ found in Medeiros [23], see Lemma 3.1, to obtain a useful estimate related to the critical exponent problem and a version of the Trundiger-Moser inequality in $W^{1,N}(\Omega_N)$ by Cianchi [21], see Lemma 3.3, to deal with the exponential growth. Another difficulty appears, since the trace of the functions on $\partial\Omega_p$ and $\partial\Omega_N$ may not vanish. We tackle this difficulty by applying the type of results that are used in the study of Neumann boundary value problems (Lemma 3.4-3.5).

1.2. Organization of the article. This article is organized as follows: In Section 2, we discuss some preliminary results for the Musielak-Sobolev spaces, while in Section 3, we show some technical results that will be used in our approach. In sections 4 and 5, we discuss some preliminaries required to prove our main results and finally, in Section 6, we prove our main results (Theorems 1.1 and 1.2).

1.3. Notations. Throughout the paper, for $t > 1$, we denote by $t' = \frac{t}{t-1}$. By C , we mean a constant which may vary from line to line or even over the same line. If C depends on the parameters r_1, r_2, \dots, r_k , we write $C = C(r_1, r_2, \dots, r_k)$.

2. A BRIEF REVIEW ABOUT THE MUSIELAK-SOBOLEV SPACES

In this section, we recall some results on Musielak-Orlicz and Musielak-Sobolev spaces. For more details, we refer to [20, 26, 33, 43] and their references.

Let $\Omega \subset \mathbb{R}^N$ be a smooth bounded domain and $\Phi(x, t) = \int_0^{|t|} \varphi(x, s) s ds$ be a generalized N-function, that is, for each $t \in \mathbb{R}$, the function $\Phi(\cdot, t)$ is measurable and for a.e. $x \in \Omega$, the function $\Phi(x, \cdot)$ is an N-function. For the reader's convenience, we recall that a continuous function $A : \mathbb{R} \rightarrow [0, +\infty)$ is an N-function if

- (i) A is convex.
- (ii) $A = 0 \Leftrightarrow t = 0$.

$$(iii) \lim_{t \rightarrow 0} \frac{A(t)}{t} = 0 \text{ and } \lim_{t \rightarrow +\infty} \frac{A(t)}{t} = +\infty .$$

(iv) A is even.

The Musielak-Orlicz space $L^\Phi(\Omega)$ is defined by

$$L^\Phi(\Omega) = \left\{ u : \Omega \rightarrow \mathbb{R} \mid u \text{ is measurable and } \exists \tau > 0 \text{ such that } \int_{\Omega} \Phi \left(x, \frac{|u|}{\tau} \right) dx < +\infty \right\}$$

endowed with the Luxemburg norm

$$|u|_{\Phi} = \inf \left\{ \lambda > 0 \mid \int_{\Omega} \Phi \left(x, \frac{|u|}{\lambda} \right) dx \leq 1 \right\} .$$

We say that an N-function Φ satisfies the Δ_2 -condition, denote by $\Phi \in \Delta_2$, if there exists a constant $K > 0$ such that

$$(2.1) \quad \Phi(x, 2t) \leq K\Phi(x, t) \quad \text{for } x \in \Omega \quad \text{and } t \in \mathbb{R} .$$

Arguing as in [44, Theorem 4.4.4], it follows that Φ satisfies the Δ_2 -condition if and only if,

$$\sup_{(x,t) \in \Omega \times (0, +\infty)} \frac{\varphi(x, |t|)|t|^2}{\tilde{\Phi}(x, |t|)} < +\infty .$$

Moreover, an important inequality involving Φ and its complementary function $\tilde{\Phi}$ (see (1.1)) is a Young's type inequality given by

$$(2.2) \quad st \leq \Phi(x, s) + \tilde{\Phi}(x, t), \quad x \in \Omega \quad \text{and } \forall s, t \geq 0 .$$

Using the above inequality, it is possible to prove a Hölder type inequality, that is,

$$\left| \int_{\Omega} uv dx \right| \leq 2\|u\|_{\Phi} \|v\|_{\tilde{\Phi}} \quad \forall u \in L^{\Phi}(\Omega) \quad \text{and } \forall v \in L^{\tilde{\Phi}}(\Omega) .$$

Arguing as in [29], if (φ_3) holds, we derive that

$$\frac{q}{q-1} \leq \frac{\tilde{\varphi}(x, |t|)|t|^2}{\tilde{\Phi}(x, |t|)} \leq \frac{p}{p-1}, \quad x \in \Omega \quad \text{and } t \neq 0 ,$$

where

$$\tilde{\Phi}(x, t) = \int_0^{|t|} \tilde{\varphi}(x, s) s ds ,$$

and

$$\tilde{\varphi}(x, s) = \sup\{t : \varphi(x, t)t \leq s\}, \quad x \in \bar{\Omega} \quad \text{and } s \geq 0 .$$

Hence, if (φ_3) holds, we have $\tilde{\Phi}$ also satisfies the Δ_2 -condition.

Arguing as in [29, Lemma A2], it is possible to prove that Φ and $\tilde{\Phi}$ satisfy the following inequality

$$(2.3) \quad \tilde{\Phi}(x, \varphi(x, t)t) \leq \Phi(x, 2t), \quad x \in \Omega \quad \text{and } t \geq 0 .$$

The condition (φ_3) is very important, because following the ideas of [29, Lemmas 2.1 and 2.5], it is possible to prove the following: Setting the functions

$$\xi_0(t) = \min\{t^p, t^q\}, \quad \xi_1(t) = \max\{t^p, t^q\}, \quad \xi_3(t) = \min\{t^{\frac{p}{p-1}}, t^{\frac{q}{q-1}}\} \quad \text{and} \quad \xi_4(t) = \max\{t^{\frac{p}{p-1}}, t^{\frac{q}{q-1}}\},$$

we have

$$(2.4) \quad \xi_0(s)\Phi(x, t) \leq \Phi(x, st) \leq \xi_1(s)\Phi(x, t) \quad \text{for } s, t \geq 0 ,$$

$$(2.5) \quad \xi_0(|u|_{\Phi}) \leq \int_{\Omega} \Phi(x, |u|) dx \leq \xi_1(|u|_{\Phi}) \quad \text{for } u \in L^{\Phi}(\Omega) ,$$

$$(2.6) \quad \xi_3(s)\tilde{\Phi}(x, t) \leq \tilde{\Phi}(x, st) \leq \xi_4(s)\tilde{\Phi}(x, t) \quad \text{for } s, t \geq 0,$$

and

$$(2.7) \quad \xi_3(|u|_{\tilde{\Phi}}) \leq \int_{\Omega} \tilde{\Phi}(x, |u|) dx \leq \xi_4(|u|_{\tilde{\Phi}}) \quad \text{for } u \in L^{\tilde{\Phi}}(\Omega).$$

The Musielak-Sobolev space $W^{1, \Phi}(\Omega)$ can be defined by

$$W^{1, \Phi}(\Omega) = \{u \in L^{\Phi}(\Omega) \mid |\nabla u| \in L^{\Phi}(\Omega)\}$$

with the norm

$$\|u\|_{1, \Phi} = |u|_{\Phi} + |\nabla u|_{\Phi}.$$

The conditions $(\varphi_1) - (\varphi_5)$ ensure that the spaces $L^{\Phi}(\Omega)$ and $W^{1, \Phi}(\Omega)$ are reflexive and separable Banach spaces, for more details see [26, Propositions 1.6 and 1.8]. In what follows, $W_0^{1, \Phi}(\Omega)$ is defined as the closure of $C_0^{\infty}(\Omega)$ in $W_0^{1, \Phi}(\Omega)$ with respect to the above norm. Moreover, $\|u\| = |\nabla u|_{\Phi}$ is a norm in $W_0^{1, \Phi}(\Omega)$ and if $(\varphi_1) - (\varphi_5)$ holds, by [32, Lemma 5.7], $\|\cdot\|$ is equivalent to the norm $\|u\|_{1, \Phi}$ in $W_0^{1, \Phi}(\Omega)$.

As a consequence of (2.5) we have the lemma below that will be used later on.

Proposition 2.1. *The functional $\rho : W_0^{1, \Phi}(\Omega) \rightarrow \mathbb{R}$ defined by*

$$(2.8) \quad \rho(u) = \int_{\Omega} \Phi(x, |\nabla u|) dx,$$

has the following properties:

- (i) If $\|u\| \geq 1$, then $\|u\|^p \leq \rho(u) \leq \|u\|^q$.
- (ii) If $\|u\| \leq 1$, then $\|u\|^q \leq \rho(u) \leq \|u\|^p$.

In particular, $\rho(u) = 1$ if and only if $\|u\| = 1$ and if $(u_n) \subset W_0^{1, \Phi}(\Omega)$, then $\|u_n\| \rightarrow 0$ if and only if $\rho(u_n) \rightarrow 0$.

Remark 1. *For the functional $\xi : L^{\Phi}(\Omega) \rightarrow \mathbb{R}$ given by*

$$\xi(u) = \int_{\Omega} \Phi(x, |u|) dx,$$

the conclusion of Proposition 2.1 also holds, for example, if $(u_n) \subset L^{\Phi}(\Omega)$, then $|u_n|_{\Phi} \rightarrow 0$ if and only if $\xi(u_n) \rightarrow 0$.

From the definition of $W^{1, \Phi}(\Omega)$ and properties of Φ , we have the continuous embedding

$$W^{1, \Phi}(\Omega) \hookrightarrow W^{1, q}((\Omega_q)_{\omega})$$

for all $\omega \in (0, \delta)$ and the compact embedding

$$W^{1, q}((\Omega_q)_{\delta}) \hookrightarrow C(\overline{(\Omega_q)_{\omega}}),$$

because $q > N$, from where it follows that

$$(2.9) \quad W^{1, \Phi}(\Omega) \hookrightarrow C(\overline{(\Omega_q)_{\omega}}),$$

is compact, which is crucial in our approach.

Next we would like to state our last result found in [26, Theorem 2.2], which says the operator $-\Delta_{\Phi} : W_0^{1, \Phi}(\Omega) \rightarrow (W_0^{1, \Phi}(\Omega))^*$ belongs to the Class (S_+) .

Lemma 2.2. *Assume the conditions $(\varphi_1) - (\varphi_8)$. If $u_n \rightharpoonup u$ in $W_0^{1, \Phi}(\Omega)$ and*

$$\lim_{n \rightarrow +\infty} \int_{\Omega} \langle \varphi(x, |\nabla u_n|) \nabla u_n, \nabla u_n - \nabla u \rangle dx = 0,$$

then $u_n \rightarrow u$ in $W_0^{1, \Phi}(\Omega)$.

Before concluding this section, we will show a version of Poincaré's inequality, which is a key point in the proof of Theorem 1.2.

Lemma 2.3. *Assume $(\varphi_1) - (\varphi_5)$ and (φ_{11}) . Then, there is $\Upsilon > 0$ independent of $r \geq 1$ such that*

$$\int_{\Omega_r} \Phi(x, |u|) dx \leq \Upsilon \int_{\Omega_r} \Phi(x, |\nabla u|) dx, \quad \forall u \in W_0^{1,\Phi}(\Omega_r).$$

Proof. Fix $p > 1$ and $v \in C_0^\infty(\Omega_r)$. Arguing as in [5, Lemma 3.1], we get

$$\int_{\Omega_r} |v|^p dx \leq \left(\frac{r+1}{r} \right)^{N-1} \int_{\Omega_r} |\nabla v|^p dx.$$

Now, taking the limit when $p \rightarrow 1$ and using the fact that $r \geq 1$, we derive that

$$\int_{\Omega_r} |v| dx \leq 2^{N-1} \int_{\Omega_r} |\nabla v| dx, \quad \forall v \in C_0^\infty(\Omega_r).$$

Since $C_0^\infty(\mathbb{R}^N)$ is dense in $W_0^{1,1}(\Omega_r)$, it follows that

$$\int_{\Omega_r} |w| dx \leq 2^{N-1} \int_{\Omega_r} |\nabla w| dx, \quad \forall w \in W_0^{1,1}(\Omega_r).$$

Now, for each $u \in W_0^{1,\Phi}(\Omega_r)$, we know that $w = \Phi(x, u) \in W_0^{1,1}(\Omega_r)$ and so,

$$\int_{\Omega_r} \Phi(x, |u|) dx \leq 2^{N-1} \int_{\Omega_r} |\nabla \Phi(x, u)| dx.$$

Since $|\nabla \Phi(x, u)| \leq |\partial_s \Phi(x, |u|)| + \varphi(|u|)|u||\nabla u|$, we obtain

$$\int_{\Omega_r} \Phi(x, |u|) dx \leq 2^{N-1} \int_{\Omega_r} (|\partial_s \Phi(x, |u|)| + \varphi(x, |u|)|u||\nabla u|) dx, \quad \forall u \in W_0^{1,\Phi}(\Omega_r).$$

Given $\epsilon > 0$, by Δ_2 condition, (φ_{11}) , (2.2) and (2.3), there is $C_\epsilon > 0$ such that

$$\int_{\Omega_r} \Phi(x, |u|) dx \leq 2^{N-1} \left[\epsilon \int_{\Omega_r} \Phi(x, |u|) dx + \kappa \int_{\Omega_r} \Phi(x, |u|) dx + C_\epsilon \int_{\Omega_r} \Phi(x, |\nabla u|) dx \right],$$

for all $u \in W_0^{1,\Phi}(\Omega_r)$. Thus, for $\epsilon = \frac{1}{2^{N+1}}$ and recalling that $\kappa < \frac{1}{2^{N+1}}$, there is $\Upsilon > 0$ independent of $r \geq 1$ such that

$$\int_{\Omega_r} \Phi(x, |u|) dx \leq \Upsilon \int_{\Omega_r} \Phi(x, |\nabla u|) dx, \quad \forall u \in W_0^{1,\Phi}(\Omega_r).$$

□

3. SOME TECHNICAL RESULTS

The main goal of this section is to recall and prove some technical results that are crucial in the proof of our main result. Since we are going to work with double criticality, which involves the exponential critical growth and the critical growth p^* , the next two results are crucial in our approach. The first one is a Concentration Compactness Lemma due to Lions for $W^{1,p}(\Theta)$ explored in Medeiros [23], where $\Theta \subset \mathbb{R}^N$ is a smooth bounded domain.

Lemma 3.1. *Let (u_n) be a sequence in $W^{1,p}(\Theta)$ with $1 < p < N$ and $u_n \rightharpoonup u$ in $W^{1,p}(\Theta)$. If*

(i) $|\nabla u_n|^p \rightarrow \mu$ weakly- $*$ in the sense of measure,

and

(ii) $|u_n|^{p^*} \rightarrow \nu$ weakly-* in the sense of measure, then for at most a countable index set J , we have

$$\left\{ \begin{array}{l} (a) \quad \nu = |u|^{p^*} + \sum_{j \in J} \nu_j \delta_{x_j}, \quad \nu_j \geq 0. \\ (b) \quad \mu \geq |\nabla u|^p + \sum_{j \in J} \mu_j \delta_{x_j}, \quad \mu_j \geq 0. \\ (c) \quad \text{If } x_j \in \Theta, \text{ then } S_p \nu_j^{\frac{p}{p^*}} \leq \mu_j. \\ (d) \quad \text{If } x_j \in \partial\Theta, \text{ then } \frac{S_p}{2^{p/N}} \nu_j^{\frac{p}{p^*}} \leq \mu_j, \end{array} \right.$$

where $p^* = \frac{Np}{N-p}$ and S_p denotes the best constant of the embedding $D^{1,p}(\mathbb{R}^N) \hookrightarrow L^{p^*}(\mathbb{R}^N)$ given by

$$(3.1) \quad S_p = \inf_{\substack{u \in D^{1,p}(\mathbb{R}^N) \\ u \neq 0}} \frac{\int_{\mathbb{R}^N} |\nabla u|^p dx}{\left(\int_{\mathbb{R}^N} |u|^{p^*} dx \right)^{\frac{p}{p^*}}}.$$

The proof of the above lemma follows by combining the arguments explored in Struwe [49, Chapter I, Section 4] and the following Cherrier's inequality [18] below.

Lemma 3.2. *Let $\Theta \subset \mathbb{R}^N$ be a smooth bounded domain and $p \in (1, N)$. Then for each $\tau > 0$, there is $M_\tau > 0$ such that*

$$\left[\frac{S_p}{2^{\frac{p}{N}}} - \tau \right] \|u\|_{L^{p^*}(\Theta)}^p \leq \|\nabla u\|_{L^p(\Theta)}^p + M_\tau \|u\|_{L^p(\Theta)}^p, \quad \forall u \in W^{1,p}(\Theta).$$

The second result that we would like to point out is a version of Trudinger-Moser inequality in $W^{1,N}(\Theta)$ due to Cianchi [21, Theorem 1.1].

Lemma 3.3. *Let $\Theta \subset \mathbb{R}^N$ be a smooth bounded domain for $N \geq 2$ and $u \in W^{1,N}(\Theta)$. Then, there is a constant $C(\Theta) > 0$ such that*

$$(3.2) \quad \int_{\Theta} e^{\alpha_N \left(\frac{|u - u_\Theta|}{\|\nabla u\|_{L^N(\Theta)}} \right)^{N'}} dx \leq C(\Theta),$$

where $u_\Theta = \frac{1}{|\Theta|} \int_{\Theta} u dx$ is the mean value of u in Θ , $\alpha_N = N \left(\frac{w_N}{2} \right)^{\frac{1}{N}}$ and w_N is the volume of sphere S^{N-1} . The integral on the left-hand of (3.2) is finite for each $u \in W^{1,N}(\Theta)$ even if α_N is replaced by any other small positive number, but no inequality of type (3.2) can hold with a large constant in the place of α_N .

From Lemma 3.3, for each $u \in W^{1,N}(\Theta)$, we have

$$(3.3) \quad e^{t|u|^{N'}} \in L^1(\Theta), \quad \forall t \geq 0.$$

For the reader interested in Trudinger-Moser inequality for functions in $W^{1,N}(\Theta)$, we would like to cite the papers due to Adimurthi and Yadava [1], Kaur and Sreenadh [37] and their references.

As a consequence of Lemma 3.3, we have the following two results whose proof can be found in [7].

Lemma 3.4. *Given $t > 1$ and $\alpha > 0$, there is $r \in (0, 1)$ and $C = C(t, r, N) > 0$ such that*

$$(3.4) \quad \sup \left\{ \int_{\Theta} e^{t\alpha|u|^{N'}} dx : u \in W^{1,N}(\Theta), \|\nabla u\|_{L^N(\Theta)} \leq r \quad \text{and} \quad \|u\|_{L^1(\Theta)} \leq r \right\} \leq C.$$

Lemma 3.5. *Let $\alpha > 0$ and $(u_n) \subset W^{1,N}(\Theta)$ be a sequence satisfying $\|\nabla u_n\|_{L^N(\Theta)}^{N'} \leq \frac{\tau}{2^{N'}} \frac{\alpha N}{\alpha}$ and $\|u_n\|_{L^1(\Theta)} \leq M$ for some $\tau \in (0, 1)$ and $M > 0$. Then, there is $t > 1$ with $t \approx 1$ such that*

$$(3.5) \quad \sup_{n \in \mathbb{N}} \int_{\Theta} e^{t\alpha |u_n|^{N'}} dx < +\infty.$$

Hence, the sequence $f_n(x) = e^{\alpha |u_n(x)|^{N'}}$ is bounded in $L^t(\Theta)$.

As a consequence of Lemma 3.5, we have the corollary below.

Corollary 3.6. *Let $(u_n) \subset W^{1,N}(\Theta)$ be a sequence as in Lemma 3.5. If $u_n(x) \rightarrow u(x)$ a.e. in Θ , then $f_n \rightarrow f$ in $L^t(\Theta)$ where $f(x) = e^{\alpha |u(x)|^{N'}}$, that is,*

$$\int_{\Theta} f_n \varphi dx \rightarrow \int_{\Theta} f \varphi dx, \quad \forall \varphi \in L^{t'}(\Theta),$$

where $\frac{1}{t} + \frac{1}{t'} = 1$.

Our next result will help us to conclude that the energy functional associated with the problem (P) is $C^1(W_0^{1,\Phi}(\Omega), \mathbb{R})$. Since it follows as in Bezerra do Ó, Medeiros and Severo [12, Proposition 1], we will omit its proof.

Lemma 3.7. *Let $(u_n) \subset W^{1,N}(\Theta)$ be a sequence such that $u_n \rightarrow u$ in $W^{1,N}(\Theta)$ for some $u \in W^{1,N}(\Theta)$. Then, for some subsequence, still denoted by itself, there is $v \in W^{1,N}(\Theta)$ such that:*

- (i) $u_n(x) \rightarrow u(x)$ a.e. in Θ .
- (ii) $|u_n(x)| \leq v(x)$ a.e. in Θ for all $n \in \mathbb{N}$.

4. PRELIMINARIES FOR THE PROOF OF THEOREM 1.1

To prove Theorem 1.1, we use the following result, whose proof follows similar arguments as in Ambrosetti and Rabinowitz [45]. Let X be a Banach space, $K \subset X$ be compact and $\gamma(Y)$ be the genus of $Y \subset \Sigma$, where

$$\Sigma := \{Y \in X \setminus \{0\} : Y \text{ is close in } X \text{ and symmetric with respect to the origin}\}.$$

Theorem 4.1. *Suppose $I \in C^1(X, \mathbb{R})$ satisfies:*

- (a) $I(0) = 0$, $I(u) = I(-u)$ for all $u \in X$.
- (b) there exists $\alpha, \rho > 0$ such that

$$I(u) \geq \alpha \quad \forall \|u\| = \rho.$$

- (c) for every $\hat{X} \subset X$ such that $\dim \hat{X} < \infty$, there exists $R = R(\hat{X}) > 0$, such that

$$I(u) \leq 0, \quad \text{for every } u \in \hat{X} \setminus B_R(0);$$

- (d) there exists $M > 0$, such that I satisfies $(PS)_c$ condition, for any $0 < c < M$.

For each $m \in \mathbb{N}$, fix a finite dimensional subspace X_m of X and consider $R_m = R(X_m) > 0$ given by condition (c). Now, define

$$(4.1) \quad D_m = B_{R_m} \cap X_m,$$

$$G_m := \{h \in C(D_m, X) : h \text{ is odd and } h(u) = u, \quad \forall u \in \partial B_{R_m} \cap X_m\},$$

$$(4.2) \quad \Gamma_j = \{h(\overline{D_m \setminus Y}); h \in G_m, m \geq j, Y \in \Sigma, \gamma(Y) \leq m - j\},$$

and

$$c_m := \inf_{K \in \Gamma_m} \max_{u \in K} I(u).$$

Then $0 < \alpha \leq c_m \leq c_{m+1}$, and if $c_m < M$, the levels c_j for $j \in \{1, 2, \dots, m\}$ are critical values of I . Moreover, if $c_1 = c_2 = \dots = c_r = c < M$, then $\gamma(K_0) > r$.

4.1. Functional setting. In what follows, we consider the associated energy functional $I : W_0^{1,\Phi}(\Omega) \rightarrow \mathbb{R}$ given by

$$I(u) = \int_{\Omega} \Phi(x, |\nabla u|) dx - \int_{\Omega} F(x, u) dx,$$

where $F(x, t) = \int_0^t f(x, s) ds$, $t \in \mathbb{R}$ and f is either of the form (f_1) or (f_2) . Here, we would like to mention that, for the rest of the article, whenever we deal with (f_1) , we assume Ω to be a smooth bounded domain in \mathbb{R}^N , $N \geq 2$ along with the hypothesis $(g_0) - (g_3)$ and $(\varphi_1) - (\varphi_8)$ as in Theorem 1.1. For (f_2) , we consider $\Omega = \Omega_r$, $N \geq 2$, $N \neq 3$ along with the hypothesis $(g_1), (g_2), (g_4), (\eta)$ and (Ω_N) and $(\varphi_1) - (\varphi_{11})$ as in Theorem 1.2.

Lemma 4.2. *Assume that f is of form (f_1) or (f_2) . Then, the functional I belongs to $C^1(W_0^{1,\Phi}(\Omega), \mathbb{R})$ and*

$$I'(u)v = \int_{\Omega} \varphi(x, |\nabla u|) \nabla u \nabla v dx - \int_{\Omega} f(x, u)v dx, \quad \forall u, v \in W_0^{1,\Phi}(\Omega).$$

Proof. See proof in [7, Lemma 3.8] □

Next, our goal is to prove that I satisfies the geometric conditions of Theorem 4.1 and the well known (PS) condition.

Lemma 4.3. *Assume that f is of the form (f_1) . Then,*

i) There are $r, \rho > 0$ such that

$$I(u) \geq \rho, \quad \text{for } \|u\| = r.$$

ii) For every $\hat{X} \subset W_0^{1,\Phi}(\Omega)$ with $\dim \hat{X} < \infty$, there exists $R = R(\hat{X}) > 0$, such that

$$I(u) \leq 0, \quad \text{for all } u \in \hat{X} \setminus B_R(0).$$

Proof. The proof of *i)* can be done as in [7, Lemma 3.9].

ii) Suppose for each $n \in \mathbb{N}$, there exists $u_n \in \hat{X} \setminus B_n(0)$ such that

$$(4.3) \quad I(u_n) > 0.$$

From (g_2) , it follows that

$$(4.4) \quad 0 < \chi F(x, t) \leq f(x, t)t, \quad \forall (x, t) \in \Omega \times (\mathbb{R} \setminus \{0\}),$$

where $\chi = \min\{\theta, \beta, \zeta\} > q$. Therefore, f satisfies the Ambrosetti-Rabinowitz condition. This gives the existence of positive constants C, D such that

$$(4.5) \quad F(x, t) \geq C|t|^\chi - D, \quad \forall t \in \mathbb{R} \quad \text{and} \quad \forall x \in \Omega.$$

Using Proposition 2.1-(ii) along with (4.5), we obtain

$$(4.6) \quad I(u_n) \leq \|u_n\|^q - C \int_{\Omega} |u_n|^\chi dx + D|\Omega|.$$

Since $\dim \hat{X} < \infty$ and $\chi > q$ letting $n \rightarrow \infty$ in (4.6), we arrive at a contradiction to our assumption (4.3). Hence *ii)* follows. □

Lemma 4.4. *Assume that f is of the type (f_1) or (f_2) . Then, every (PS) sequence (u_n) of the functional I is bounded in $W_0^{1,\Phi}(\Omega)$.*

Proof. Let $d > 0$ and (u_n) be a $(PS)_d$ sequence for I . Then, there are constants $C_1, C_2 > 0$ such that

$$(4.7) \quad I(u_n) - \frac{1}{\chi} I'(u_n)u_n \leq C_1 + C_2 \|u_n\|, \quad \forall n \in \mathbb{N}.$$

If f is of the type (f_1) or (f_2) it is easy to check that (4.4) holds. Hence, by (φ_3) and the definition of I ,

$$\begin{aligned} I(u_n) - \frac{1}{\chi} I'(u_n)u_n &\geq \int_{\Omega} \Phi(x, |\nabla u_n|) dx - \frac{1}{\chi} \int_{\Omega} \varphi(x, |\nabla u_n|) |\nabla u_n|^2 dx \\ &\geq \left(1 - \frac{q}{\chi}\right) \int_{\Omega} \Phi(x, |\nabla u_n|) dx. \end{aligned}$$

Therefore,

$$\left(1 - \frac{q}{\chi}\right) \int_{\Omega} \Phi(x, |\nabla u_n|) dx \leq C_1 + C_2 \|u_n\|, \quad \forall n \in \mathbb{N}.$$

If there is $(u_{n_j}) \subset (u_n)$ such that $\|u_{n_j}\| \geq 1$, then Proposition 2.1-(i) leads to

$$\left(1 - \frac{q}{\chi}\right) \|u_{n_j}\|^p \leq C_1 + C_2 \|u_{n_j}\|, \quad \forall j \in \mathbb{N},$$

from where it follows the boundedness of (u_{n_j}) . This implies the boundedness of (u_n) . \square

Corollary 4.5. *Assume that f is of the type (f_1) or (f_2) and let (u_n) be a $(PS)_d$ sequence of I with $d \in (0, M)$,*

$$M = \left(1 - \frac{q}{\chi}\right) \min \left\{ \frac{1}{N} \left(\frac{\alpha_N}{2^{N'}\alpha}\right)^{N-1}, \frac{1}{p} S_p^{\frac{N}{p}} \right\},$$

where $\chi = \min\{\theta, \beta, \zeta\}$. Then,

$$\limsup_{n \rightarrow +\infty} \|\nabla u_n\|_{L^N(\Omega_N)}^{N'} < \frac{\alpha_N}{2^{N'}\alpha}.$$

Hence, without loss of generality, we can assume that there is $\tau \in (0, 1)$ such that

$$\|\nabla u_n\|_{L^N(\Omega_N)}^{N'} \leq \frac{\tau \alpha_N}{2^{N'}\alpha}, \quad \forall n \in \mathbb{N}.$$

Proof. First of all, we must recall that

$$I(u_n) - \frac{1}{\chi} I'(u_n)u_n = d + o_n(1)\|u_n\| + o_n(1).$$

Therefore, by (φ_6) ,

$$\begin{aligned} d + o_n(1)\|u_n\| + o_n(1) &\geq \int_{\Omega} \left(\Phi(x, |\nabla u_n|) - \frac{1}{\chi} \varphi(x, |\nabla u_n|) |\nabla u_n|^2 \right) dx \\ &\geq \frac{1}{N} \left(1 - \frac{q}{\chi}\right) \int_{\Omega_N} |\nabla u_n|^N dx. \end{aligned}$$

Hence,

$$\limsup_{n \rightarrow +\infty} \frac{1}{N} \left(1 - \frac{q}{\chi}\right) \int_{\Omega_N} |\nabla u_n|^N dx \leq d < \min \left(1 - \frac{q}{\chi}\right) \left\{ \frac{1}{N} \left(\frac{\alpha_N}{2^{N'}\alpha}\right)^{N-1}, \frac{1}{p} S_p^{\frac{N}{p}} \right\}$$

leading to

$$\limsup_{n \rightarrow +\infty} \int_{\Omega_N} |\nabla u_n|^N dx < \left(\frac{\alpha_N}{2^{N'}\alpha}\right)^{N-1},$$

which proves the lemma. \square

Lemma 4.6. *Assume that f is of the type (f_1) or (f_2) . Then, the functional I verifies the $(PS)_d$ condition for $d \in (0, M)$, where M was given in Corollary 4.5.*

Proof. The proof this lemma follows as in [7, Lemma 3.13], however for the reader's convenience, we will write the proof for (f_1) , since for (f_2) , it follows with similar arguments. Let (u_n) be a $(PS)_d$ sequence for I . Then, by Lemma 4.4, (u_n) is bounded in $W_0^{1,\Phi}(\Omega)$. Since $W_0^{1,\Phi}(\Omega)$ is reflexive, we assume that for some subsequence, still denoted by itself, there is $u \in W_0^{1,\Phi}(\Omega)$ such that

$$u_n \rightharpoonup u \quad \text{in } W_0^{1,\Phi}(\Omega),$$

and

$$u_n(x) \rightarrow u(x) \quad \text{a.e. in } \Omega.$$

Let us set

$$P_n = \int_{\Omega} \langle \varphi(x, |\nabla u_n|) \nabla u_n, \nabla u_n - \nabla u \rangle dx,$$

that is,

$$P_n = I'(u_n)u_n + \int_{\Omega} f(x, u_n)u_n dx - I'(u_n)u - \int_{\Omega} f(x, u_n)u dx.$$

Consequently

$$P_n = \int_{\Omega} f(x, u_n)u_n dx - \int_{\Omega} f(x, u_n)u dx + o_n(1).$$

From the definition of f together with embedding (2.9),

$$\lim_{n \rightarrow +\infty} \int_{\Omega} \tilde{\eta}_q(x)g(x, u_n)u_n dx = \lim_{n \rightarrow +\infty} \int_{\Omega} \tilde{\eta}_q(x)g(x, u_n)u dx = \int_{\Omega} \tilde{\eta}_q(x)g(x, u)u dx,$$

$$\lim_{n \rightarrow +\infty} \int_{\Omega} \eta_p(x)|u_n|^{\zeta} dx = \lim_{n \rightarrow +\infty} \int_{\Omega} \eta_p(x)|u_n|^{\zeta-2}u_n u dx = \int_{\Omega} \eta_p(x)|u|^{\zeta} dx,$$

$$\lim_{n \rightarrow +\infty} \int_{\Omega \setminus \Omega_N} \eta_N(x)|u_n|^{\beta} e^{\alpha|u_n|^{N'}} dx = \int_{\Omega \setminus \Omega_N} \eta_N(x)|u|^{\beta} e^{\alpha|u|^{N'}} dx,$$

$$\lim_{n \rightarrow +\infty} \int_{\Omega \setminus \Omega_N} \eta_N(x)|u_n|^{\beta-2}u_n u e^{\alpha|u_n|^{N'}} dx = \int_{\Omega \setminus \Omega_N} \eta_N(x)|u|^{\beta} e^{\alpha|u|^{N'}} dx,$$

$$\lim_{n \rightarrow +\infty} \int_{\Omega \setminus \Omega_p} \eta_p(x)|u_n|^{p^*} dx = \int_{\Omega \setminus \Omega_p} \eta_p(x)|u|^{p^*} dx,$$

and

$$\lim_{n \rightarrow +\infty} \int_{\Omega \setminus \Omega_p} \eta_p(x)|u_n|^{p^*-2}u_n u dx = \int_{\Omega \setminus \Omega_p} \eta_p(x)|u|^{p^*} dx.$$

Consequently

$$\begin{aligned} P_n &= \lambda \int_{\Omega_N} |u_n|^{\beta} e^{\alpha|u_n|^{N'}} dx - \lambda \int_{\Omega_N} |u_n|^{\beta-2}u_n u e^{\alpha|u_n|^{N'}} dx + \int_{\Omega_p} |u_n|^{p^*} dx \\ &\quad - \int_{\Omega_p} |u_n|^{p^*-2}u_n u dx + o_n(1). \end{aligned}$$

By Corollary 4.5, the sequence (u_n) satisfies

$$\|\nabla u_n\|_{L^N(\Omega_N)}^{N'} \leq \frac{\tau \alpha_N}{2N' \alpha}, \quad \forall n \in \mathbb{N},$$

for some $\tau \in (0, 1)$. Employing Corollary 3.6, there is $t > 1$ and $t \approx 1$ such that the sequence $h_n(x) = e^{\alpha|u_n(x)|^{N'}}$ is weakly convergent to $h(x) = e^{\alpha|u(x)|^{N'}}$ in $L^t(\Omega_N)$, that is,

$$(4.8) \quad \int_{\Omega_N} h_n \varphi \, dx \rightarrow \int_{\Omega_N} h \varphi \, dx, \quad \forall \varphi \in L^t(\Omega_N).$$

As

$$|u_n|^\beta \rightarrow |u|^\beta \quad \text{in } L^t(\Omega_N),$$

it follows that

$$\int_{\Omega_N} h_n |u_n|^\beta \, dx \rightarrow \int_{\Omega_N} h |u|^\beta \, dx,$$

that is,

$$\int_{\Omega_N} |u_n|^\beta e^{\alpha|u_n|^{N'}} \, dx \rightarrow \int_{\Omega_N} |u|^\beta e^{\alpha|u|^{N'}} \, dx.$$

Now, using the fact that

$$|u_n|^{\beta-2} u_n u \rightarrow |u|^\beta \quad \text{in } L^t(\Omega_N),$$

we also derive that

$$\int_{\Omega_N} |u_n|^{\beta-2} u_n u e^{\alpha|u_n(x)|^{N'}} \, dx \rightarrow \int_{\Omega_N} |u|^{\beta-2} u u e^{\alpha|u(x)|^{N'}} \, dx.$$

The above analysis ensures that

$$\lim_{n \rightarrow +\infty} \int_{\Omega_N} |u_n|^\beta e^{\alpha|u_n(x)|^{N'}} \, dx = \lim_{n \rightarrow +\infty} \int_{\Omega_N} |u_n|^{\beta-2} u_n u e^{\alpha|u_n(x)|^{N'}} \, dx = \int_{\Omega_N} |u|^\beta e^{\alpha|u|^{N'}} \, dx,$$

and then,

$$P_n = \int_{\Omega_p} |u_n|^{p^*} \, dx - \int_{\Omega_p} |u_n|^{p^*-2} u_n u \, dx + o_n(1).$$

By [36, Lemma 4.8],

$$\lim_{n \rightarrow +\infty} \int_{\Omega_p} |u_n|^{p^*-2} u_n u \, dx = \int_{\Omega_p} |u|^{p^*} \, dx,$$

then

$$P_n = \int_{\Omega_p} |u_n|^{p^*} \, dx - \int_{\Omega_p} |u|^{p^*} \, dx + o_n(1).$$

Now, we are going to use the Concentration Compactness Lemma 3.1 to the sequence $(u_n) \subset W^{1,p}(\Omega_p)$. From (φ_7) , for each open ball $B \subset (\Omega_q)_\delta$ we have that the embedding $W^{1,\Phi}(\Omega) \hookrightarrow C(\overline{B})$ is compact, then as (u_n) is a bounded (PS) for I , it is possible to prove that for some subsequence there holds

$$\int_B \langle \varphi(x, |\nabla u_n|) \nabla u_n, \nabla u_n - \nabla u \rangle \, dx \rightarrow 0.$$

Since from $(\varphi_6) - (\varphi_8)$, the embedding $W^{1,\Phi}(B) \hookrightarrow L^\Phi(B)$ is compact, the last limit together with the Δ_2 -condition (2.1) implies that

$$u_n \rightarrow u \quad \text{in } W^{1,\Phi}(B).$$

Now, recalling that the embedding $W^{1,\Phi}(B) \hookrightarrow W^{1,p}(B)$ is continuous, we derive that

$$u_n \rightarrow u \quad \text{in } W^{1,p}(B),$$

from where it follows that $x_i \in \overline{\Omega_p} \setminus (\Omega_q)_\delta$ for all $i \in J$. Now, our goal is proving that J must be a finite set. Have this in mind, we will consider $J = J_1 \cup J_2$ where

$$J_1 = \{i \in J : x_i \in \overline{\Omega_p} \setminus \overline{(\Omega_q)_\delta}\}$$

and

$$J_2 = \{i \in J : x_i \in \partial(\Omega_q)_\delta \cap \Omega_p\}.$$

If $i \in J_1$, the condition (φ_7) says that $c_3 t^{p-2} \geq \varphi(x, t) \geq t^{p-2}$ for $x \in \overline{\Omega_p} \setminus \overline{(\Omega_q)_\delta}$. This fact permits us to repeat the same arguments explored in [31, Lemma 2.3] to conclude that J_1 is finite. Now, if $i \in J_2$, the situation is more subtle and we must be careful. In what follows let us consider $\tilde{\psi} \in C_0^\infty(\mathbb{R}^N)$ such that

$$\tilde{\psi} \equiv 1 \text{ on } B(0, 1) \text{ and } \tilde{\psi} \equiv 0 \text{ on } B(0, 2)^c.$$

For each $\epsilon > 0$, we set

$$\psi(x) = \tilde{\psi}((x - x_i)/\epsilon), \quad \forall x \in \mathbb{R}^N.$$

Since (u_n) is a bounded sequence in $W^{1,\Phi}(\Omega)$, the sequence (ψu_n) is also bounded in $W^{1,\Phi}(\Omega)$ and so, $I'(u_n)\psi u_n = o_n(1)$. Hence,

$$\int_{\Omega} \varphi(x, |\nabla u_n|) \nabla u_n \nabla(\psi u_n) dx = \int_{\Omega} \tilde{\eta}_q(x) g(x, u_n) \psi u_n dx + \int_{\Omega} \eta_p(x) |u_n|^{p^*} \psi dx + o_n(1).$$

Now, given $\xi > 0$, the Young's inequality (2.2) combined with (2.3) and Δ_2 -condition (2.1) gives

$$\int_{\Omega} |\varphi(x, |\nabla u_n|)|\nabla u_n||u_n||\nabla \psi| dx \leq \xi \int_{\Omega} \Phi(x, |\nabla u_n|) dx + C_\xi \int_{\Omega} \Phi(x, |\nabla \psi||u_n|) dx.$$

for some $C_\xi > 0$. Note that by (φ_8) ,

$$\int_{\Omega} \Phi(x, |\nabla \psi||u_n|) dx \leq C_1 \left(\int_{B(x_i, 2\epsilon)} |\nabla \psi|^p |u_n|^p dx + \int_{B(x_i, 2\epsilon)} \tau_2(x) |\nabla \psi|^q |u_n|^q dx \right).$$

By Hölder's inequality

$$\limsup_{n \rightarrow +\infty} \int_{B(x_i, 2\epsilon)} |u_n|^p |\nabla \psi|^p dx \leq C_2 \left(\int_{B(x_i, 2\epsilon)} |u|^{p^*} dx \right)^{\frac{N-p}{N}}$$

from where it follows that

$$(4.9) \quad \lim_{\epsilon \rightarrow 0} \left[\limsup_{n \rightarrow +\infty} \int_{B(x_i, 2\epsilon)} |u_n|^p |\nabla \psi|^p dx \right] \leq \lim_{\epsilon \rightarrow 0} C_2 \left(\int_{B(x_i, 2\epsilon)} |u|^{p^*} dx \right)^{\frac{N-p}{N}} = 0.$$

Arguing as above, we also have

$$\limsup_{n \rightarrow +\infty} \int_{\Omega} \tau_2(x) |u_n|^q |\nabla \psi|^q dx \leq \left(\int_{B(x_i, 2\epsilon)} \left| \tau_2^{\frac{1}{q}}(x) \nabla \psi \right|^{\frac{qp^*}{p^*-q}} dx \right)^{\frac{p^*-q}{p^*}} \left(\int_{B(x_i, 2\epsilon)} |u|^{p^*} dx \right)^{\frac{q}{p^*}}.$$

By change of variable,

$$\begin{aligned} \int_{B(x_i, 2\epsilon)} \left| \tau_2^{\frac{1}{q}}(x) \nabla \psi \right|^{\frac{qp^*}{p^*-q}} dx &= \left(\frac{1}{\epsilon} \right)^{\frac{qp^*}{p^*-q}} \int_{B(0, 2)} \left| \tau_2^{\frac{1}{q}}(\epsilon x + x_i) \nabla \tilde{\psi} \right|^{\frac{qp^*}{p^*-q}} dx \\ &\leq C_5 \left(\frac{1}{\epsilon} \right)^{\frac{qp^*}{p^*-q}} \int_{B(0, 2)} \left| \tau_2^{\frac{1}{q}}(\epsilon x + x_i) \right|^{\frac{qp^*}{p^*-q}} dx. \end{aligned}$$

Since $x_i \in \partial(\Omega_q)_\delta \cap \Omega_p$, it follows that

$$\tau_2(\epsilon x + x_i) \leq c_4 \epsilon^s |x|^s$$

and

$$\int_{B(x_i, 2\epsilon)} \left| \tau_2^{\frac{1}{q}}(x) \nabla \psi \right|^{\frac{qp^*}{p^*-q}} dx \leq C_6 \epsilon^{\frac{(s-q)p^*}{p^*-q}}.$$

As $s > q$, it follows that

$$(4.10) \quad \lim_{\epsilon \rightarrow 0} \left[\limsup_{n \rightarrow +\infty} \int_{\Omega} \tau_2(x) |u_n|^q |\nabla \psi|^q dx \right] = 0.$$

Now, the boundedness of (u_n) in $W^{1,\Phi}(\Omega)$ together with Proposition (2.1), (4.9) and (4.10) ensures that

$$\lim_{\epsilon \rightarrow 0} \left[\limsup_{n \rightarrow +\infty} \int_{\Omega} |\varphi(x, |\nabla u_n|)| \nabla u_n |u_n| |\nabla \psi| dx \right] \leq \xi C,$$

for some $C > 0$. Since $\xi > 0$ is arbitrary, we can deduce that

$$\lim_{\epsilon \rightarrow 0} \left[\limsup_{n \rightarrow +\infty} \int_{\Omega} |\varphi(x, |\nabla u_n|)| \nabla u_n |u_n| |\nabla \psi| dx \right] = 0.$$

The last limit together with the fact that $\varphi(x, t) \geq t^{p-2}$ for $x \in \Omega_p$ permit us to conclude as in [31, Lemma 2.3], that J_2 is also finite. Consequently, J is a finite set. However, in order to conclude the proof of the lemma, we need to show that J is in fact an empty set. Seeking by a contradiction, assume that there is $i \in J$. In this case, the argument explored in [31] also says for us that

$$\nu_i \geq S_p^{\frac{N}{p}}.$$

Hence, by Lemma 3.1-(d),

$$\mu_i \geq S_p^{\frac{N}{p}}.$$

As $|\nabla u_n|^p \rightarrow \mu$ weakly-* in the sense of measure, we have

$$\liminf_{n \rightarrow +\infty} \int_{\Omega_p} |\nabla u_n|^p dx \geq \mu_i$$

and so,

$$\liminf_{n \rightarrow +\infty} \int_{\Omega_p} |\nabla u_n|^p dx \geq S_p^{\frac{N}{p}}.$$

Now, using once more the equality

$$I(u_n) - \frac{1}{\chi} I'(u_n) u_n = d + o_n(1) \|u_n\| + o_n(1),$$

we get

$$d + o_n(1) \|u_n\| + o_n(1) \geq \frac{1}{p} \left(1 - \frac{q}{\chi} \right) \int_{\Omega_p} |\nabla u_n|^p dx.$$

Taking the limit of $n \rightarrow +\infty$, we find the inequality below

$$d \geq \frac{1}{p} \left(1 - \frac{q}{\chi} \right) S_p^{\frac{N}{p}},$$

which is a contradiction, showing that $J = \emptyset$. Thereby, by Lemma 3.1-(a), $\nu = |u|^{p^*}$ and

$$\int_{\Omega_p} |u_n|^{p^*} dx \rightarrow \int_{\Omega_p} |u|^{p^*} dx,$$

implying that $P_n = o_n(1)$, that is,

$$\lim_{n \rightarrow +\infty} \int_{\Omega} \langle \varphi(x, |\nabla u_n|) \nabla u_n, \nabla u_n - \nabla u \rangle dx = 0.$$

Now, it is enough to apply Lemma 2.2 to finish the proof. \square

Our last lemma in this section is as follows:

Lemma 4.7. *Assume that (f_1) holds. Then, for each $m \in \mathbb{N}$, there exists positive constants λ_m, μ_m and τ_m such that*

$$c_m^{\lambda, \mu, \tau} := \inf_{K \in \Gamma_m} \sup_{u \in K} I(u) < M,$$

for all $\lambda \geq \lambda_m, \mu \geq \mu_m$ and $\tau \geq \tau_m$, where M is given by Corollary 4.5.

Proof. First, we claim that for some positive constant $C > 0$, we have

$$(4.11) \quad \min_{u \in K, \|u\|=1} \left\{ \int_{\Omega_N} |u|^\beta dx + \int_{\Omega_q} |u|^{q_2} dx + \int_{\Omega_p} |u|^\zeta dx \right\} \geq C,$$

where $K \subset X_m$ is compact such that $\dim X_m < \infty$.

Indeed, if (4.11) does not hold, then there exists a sequence $\{u_n\} \subset K$ with $\|u_n\| = 1$ such that

$$(4.12) \quad \left\{ \int_{\Omega_N} |u_n|^\beta dx + \int_{\Omega_q} |u_n|^{q_2} dx + \int_{\Omega_p} |u_n|^\zeta dx \right\} \leq \frac{1}{n}, \quad \forall n \in \mathbb{N}.$$

Since $\dim X_m < \infty$, there exists a subsequence of $\{u_n\}$ still denoted by $\{u_n\}$ and $u \in K$ with $\|u\| = 1$ such that $u_{n_j} \rightarrow u$ in X_m . Then, letting $n \rightarrow \infty$ in (4.12), we obtain

$$\int_{\Omega_N} |u|^\beta dx + \int_{\Omega_q} |u|^{q_2} dx + \int_{\Omega_p} |u|^\zeta dx = 0.$$

Hence, we have $u = 0$ a.e. in each of the sets Ω_N, Ω_q and Ω_p . Now, since $\Omega = \Omega_N \cup \Omega_q \cup \Omega_p$, we have $u = 0$ a.e. in Ω . This contradicts the fact that $\|u\| = 1$. Hence the Claim (4.11) follows.

Now we choose $K = \overline{D}_m$ where D_m is given by (4.1). Since $h = I_d \in G_m$, the definition of Φ combined with (2.4) and (f_1) gives

$$c_m^{\lambda, \mu, \tau} \leq \sup_{u \in K} \left\{ \|u\|^p + \|u\|^q - \lambda \int_{\Omega_N} |u|^\beta dx - \mu \int_{\Omega_q} |u|^{q_2} dx - \tau \int_{\Omega_p} |u|^\zeta dx \right\},$$

or equivalently

$$c_m^{\lambda, \mu, \tau} \leq \sup_{u \in K} \left\{ \|u\|^p + \|u\|^q - \lambda \|u\|^\beta \int_{\Omega_N} \left| \frac{u}{\|u\|} \right|^\beta dx - \mu \|u\|^{q_2} \int_{\Omega_q} \left| \frac{u}{\|u\|} \right|^{q_2} dx - \tau \|u\|^\zeta \int_{\Omega_p} \left| \frac{u}{\|u\|} \right|^\zeta dx \right\}.$$

Now, when $\|u\| > 1$, we observe that

$$\begin{aligned} & \|u\|^p + \|u\|^q - \lambda \|u\|^\beta \int_{\Omega_N} \left| \frac{u}{\|u\|} \right|^\beta dx - \mu \|u\|^{q_2} \int_{\Omega_q} \left| \frac{u}{\|u\|} \right|^{q_2} dx - \tau \|u\|^\zeta \int_{\Omega_p} \left| \frac{u}{\|u\|} \right|^\zeta dx \\ & \leq \|u\|^p + \|u\|^q - \|u\|^{l_1} \chi_1 \left(\int_{\Omega_N} \left| \frac{u}{\|u\|} \right|^\beta dx + \int_{\Omega_q} \left| \frac{u}{\|u\|} \right|^{q_2} dx + \int_{\Omega_p} \left| \frac{u}{\|u\|} \right|^\zeta dx \right) \\ & \leq \|u\|^p + \|u\|^q - C \chi_1 \|u\|^{l_1}, \end{aligned}$$

where $l_1 = \min\{\beta, q_2, \zeta\}$, $\chi_1 = \min\{\lambda, \mu, \tau\}$ and the constant C is given by (4.11).

Moreover, when $\|u\| \leq 1$, we get

$$\begin{aligned} & \|u\|^p + \|u\|^q - \lambda \|u\|^\beta \int_{\Omega_N} \left| \frac{u}{\|u\|} \right|^\beta dx - \mu \|u\|^{q_2} \int_{\Omega_q} \left| \frac{u}{\|u\|} \right|^{q_2} dx - \tau \|u\|^r \int_{\Omega_p} \left| \frac{u}{\|u\|} \right|^\zeta dx \\ & \leq \|u\|^p + \|u\|^q - \|u\|^{l_2} \chi_1 \left(\int_{\Omega_N} \left| \frac{u}{\|u\|} \right|^\beta dx + \int_{\Omega_q} \left| \frac{u}{\|u\|} \right|^{q_2} dx + \int_{\Omega_p} \left| \frac{u}{\|u\|} \right|^\zeta dx \right) \\ & \leq \|u\|^p + \|u\|^q - C \chi_1 \|u\|^{l_2}, \end{aligned}$$

where $l_2 = \max\{\beta, q_2, \zeta\}$, $\chi_1 = \min\{\lambda, \mu, \tau\}$ and the constant C is given by (4.11). Hence,

$$c_m^{\lambda, \mu, \tau} \leq \|u\|^p + \|u\|^q - C \chi \|u\|^l,$$

where $l = l_1$ or l_2 .

Let, $w(t) = t^p + t^q - C \chi t^l$. Then using the fact that $l > q > p$, it can be easily seen that w achieves its maximum at $\hat{t} = \hat{t}(\lambda, \mu, \tau) > 0$ which goes to 0 as the parameters λ, μ, τ goes to infinity. Hence there exists λ_m, μ_m, τ_m such that for all $\lambda \geq \lambda_m, \mu \geq \mu_m$ and $\tau \geq \tau_m$, we have

$$c_m^{\lambda, \mu, \tau} < M,$$

where M is given by Corollary 4.5. Hence the Lemma follows. \square

5. PRELIMINARIES FOR THE PROOF OF THEOREM 1.2

5.1. Functional setting. In what follows f is of the type (f_2) , $\Omega = \Omega_r$, see (1.7), $N \geq 2$, $N \neq 3$ and the hypothesis $(g_1), (g_2), (g_4), (\varphi_1) - (\varphi_{11})$ will be assumed, unless otherwise mentioned. Let us denote by $O(N)$ the group of $N \times N$ orthogonal matrices. For any integer $1 \leq k < \infty$, let us consider the finite rotational subgroup O_k of $O(2)$ given by

$$O_k := \left\{ h \in O(2) : h(x) = \left(x_1 \cos \frac{2\pi l}{k} + x_2 \sin \frac{2\pi l}{k}, -x_1 \sin \frac{2\pi l}{k} + x_2 \cos \frac{2\pi l}{k} \right) \right\}$$

where $x = (x_1, x_2) \in \mathbb{R}^2$ and $l \in \{0, \dots, k-1\}$. We define the subgroups of $O(N)$

$$H_k := O_k \times O(N-2), \quad 1 \leq k < \infty \quad \text{and} \quad H_\infty := O(2) \times O(N-2).$$

Associated with the above subgroups, we set the subspaces

$$W_{0, H_k}^{1, \Phi}(\Omega_r) := \left\{ u \in W_0^{1, \Phi}(\Omega_r) : u(x) = u(h^{-1}x), \text{ for all } h \in H_k \right\}, \quad 1 \leq k \leq \infty,$$

endowed with the usual norm of $W_0^{1, \Phi}(\Omega_r)$, that is,

$$\|u\| = \|\nabla u\|_\Phi + \|u\|_\Phi.$$

Hereafter, we denote by $I : W_{0, H_k}^{1, \Phi}(\Omega_r) \rightarrow \mathbb{R}$ the functional given by

$$I(u) = \int_{\Omega_r} \Phi(x, |\nabla u|) dx - \int_{\Omega_r} F(x, u) dx.$$

Throughout this section, $J_{k,r}$ denotes the following real number

$$J_{k,r} = \inf_{u \in \mathcal{M}_{k,r}} I(u),$$

where

$$\mathcal{M}_{k,r} = \{u \in W_{0, H_k}^{1, \Phi}(\Omega_r) \setminus \{0\}, I'(u)u = 0\}.$$

5.2. Properties of the levels $J_{k,r}$. Our first result concerns the positivity of $J_{k,r}$.

Lemma 5.1. *For any $1 \leq k \leq \infty$ and $r > 0$, we have $J_{k,r} > 0$.*

Proof. We prove the result in two steps.

Step 1. We claim that for every fixed $1 \leq k \leq \infty$ and $r > 0$, there exists a constant $\eta > 0$ such that

$$(5.1) \quad \|u\| > \eta, \text{ for all } u \in \mathcal{M}_{k,r}.$$

Indeed, if (5.1) does not hold, there exists a sequence $(u_n) \in \mathcal{M}_{k,r}$ such that $\|u_n\| \rightarrow 0$ as $n \rightarrow \infty$. From $u_n \in \mathcal{M}_{k,r}$ we have $I'(u_n)u_n = 0$. Hence

$$(5.2) \quad \int_{\Omega_r} \varphi(x, \nabla u_n) |\nabla u_n|^2 dx = \int_{\Omega_r} f(|x|, u_n) u_n dx.$$

Due to the fact $\|u_n\| \rightarrow 0$ as $n \rightarrow \infty$, without loss of generality we may assume that $\|u_n\| < 1$ for all $n \in \mathbb{N}$. Hence from (φ_3) and Proposition 2.1, we can estimate the left hand side of (5.2) as follows:

$$(5.3) \quad p \|u_n\|^q \leq p \int_{\Omega_r} \Phi(x, u_n) dx \leq \int_{\Omega_r} \varphi(x, \nabla u_n) |\nabla u_n|^2 dx.$$

Now, we estimate the right hand side of (5.2). Indeed,

$$(5.4) \quad \begin{aligned} \int_{\Omega_r} f(|x|, u_n) u_n dx &= \lambda \int_{\Omega_N} |u_n|^\beta e^{\alpha|u_n|^{N'}} dx + \int_{\Omega_r \setminus \Omega_N} \lambda \eta_N(|x|) |u_n|^\beta e^{\alpha|u_n|^{N'}} dx \\ &\quad + \int_{\Omega_q} g(|x|, u_n) u_n dx + \int_{\Omega_r \setminus \Omega_q} \tilde{\eta}(|x|) g(|x|, u_n) u_n dx \\ &\quad + \int_{\Omega_p} |u_n|^{p^*} dx + \int_{\Omega_r \setminus \Omega_p} \eta_p(|x|) |u_n|^{p^*} dx \\ &\leq \lambda \int_{\Omega_N} |u_n|^\beta e^{\alpha|u_n|^{N'}} dx + \int_{\Omega_q} g(|x|, u_n) u_n dx + \int_{\Omega_p} |u_n|^{p^*} dx \\ &\quad + \int_{(\overline{\Omega_q})_{\delta/2}} f(|x|, u_n) u_n dx \\ &= I_1 + I_2 + I_3 + I_4. \end{aligned}$$

Estimate of I_1 : As $\|u_n\| \rightarrow 0$, by Hölder's inequality and Lemma 3.5 for some constant $C_1 > 0$ (independent of n), we have

$$(5.5) \quad \begin{aligned} I_1 &= \lambda \int_{\Omega_N} |u_n|^\beta e^{\alpha|u_n|^{N'}} dx \\ &\leq \lambda \left(\int_{\Omega_N} |u_n|^{2\beta} dx \right)^{\frac{1}{2}} \left(\int_{\Omega_N} e^{2\alpha|u_n|^{N'}} dx \right)^{\frac{1}{2}} \\ &\leq C_1 \|u_n\|^\beta. \end{aligned}$$

Estimate of I_2 : From the condition (g_1) and the embedding (2.9) for some constant $C_2 > 0$ (independent of n) we deduce that

$$(5.6) \quad I_2 = \int_{\Omega_q} g(|x|, u_n) u_n dx \leq C_2 \|u_n\|^{q_1}.$$

Estimate of I_3 : It is clear that

$$(5.7) \quad I_3 = \int_{\Omega_p} |u_n|^{p^*} dx \leq C_3 \|u_n\|^{p^*},$$

for some constant $C_3 > 0$ (independent of n).

Estimate of I_4 : By the embedding (2.9), the definition of f_2 and the condition (g_1) we have that

$$(5.8) \quad I_4 = \int_{(\overline{\Omega_q})_{\delta/2}} f(|x|, u_n) u_n dx \leq C_4 (\|u_n\|^{q_1} + \|u_n\|^\beta + \|u_n\|^{p^*}),$$

for some constant $C_4 > 0$ (independent of n).

Therefore, using the estimates (5.5), (5.6), (5.7) and (5.8) in (5.4), we obtain

$$(5.9) \quad \int_{\Omega_r} f(|x|, u_n) u_n dx \leq C (\|u_n\|^{q_1} + \|u_n\|^\beta + \|u_n\|^{p^*}),$$

for some constant $C > 0$ (independent of n). Using (5.9) in (5.3), we have

$$(5.10) \quad p \|u_n\|^q \leq p \int_{\Omega_r} \Phi(x, |\nabla u_n|) dx \leq C (\|u_n\|^{q_1} + \|u_n\|^\beta + \|u_n\|^{p^*}).$$

Since all the parameters β, q_1 and p^* are larger than q , from (5.10) for some constant $\widehat{C} > 0$ (independent of n), we have

$$\|u_n\| \geq \widehat{C}, \text{ for all } u_n \in \mathcal{M}_{k,r},$$

which is a contradiction to the fact $\|u_n\| \rightarrow 0$. Hence (5.1) holds.

Step 2. From the definition of I , (φ_3) and Proposition 2.1 for any $u \in \mathcal{M}_{k,r}$, we have

$$\begin{aligned} I(u) &= I(u) - \frac{1}{\chi} I'(u)u \\ &= \int_{\Omega_r} \Phi(x, |\nabla u|) dx - \frac{1}{\chi} \int_{\Omega_r} \phi(x, |\nabla u_n|) |\nabla u_n|^2 dx \\ &\geq \left(1 - \frac{q}{\chi}\right) \int_{\Omega_r} \Phi(x, |\nabla u|) dx \\ &\geq \left(1 - \frac{q}{\chi}\right) \max\{\|u\|^q, \|u\|^p\} \\ &\geq \left(1 - \frac{q}{\chi}\right) \max\{\eta^q, \eta^p\}, \end{aligned}$$

where in the last line, we have used the estimate (5.1) from Step 1 and $\chi = \min\{\theta, \beta, p^*\}$. This means that,

$$J_{k,r} \geq \left(1 - \frac{q}{r}\right) \max\{\eta^q, \eta^p\} > 0,$$

for every $1 \leq k \leq \infty$ and all $r > 0$. Hence the result follows. \square

Lemma 5.2. *For any integer $1 \leq k < \infty$, there exists $\lambda_0 = \lambda_0(k) > 0$, such that*

$$J_{k,r} < M = \left(1 - \frac{q}{\chi}\right) \min \left\{ \frac{1}{N} \left(\frac{\alpha_N}{2^{N'} \alpha} \right)^{N-1}, \frac{1}{p} S_p^{N/p} \right\}, \text{ for all } \lambda \geq \lambda_0, \text{ and } \chi = \min\{\theta, \beta, p^*\}.$$

Proof. Fix $1 \leq k < \infty$. Due to (η) , there exists $\gamma = \gamma(k) < \min\{\frac{1}{2}, \delta_1\}$ such that the ball $B_{\gamma,r} := B_\gamma\left(\frac{2r+1}{2}, 0, \dots, 0\right) \subset \Omega_N \setminus \overline{(\Omega_q)_\delta}$ satisfies

$$h^i B_{\gamma,r} \cap h^j B_{\gamma,r} = \emptyset, \text{ for all } h^i \in H_k, i \neq j \in \{0, 1, \dots, k-1\}.$$

Consider $v_r \in W_0^{1,\Phi}(B_{\delta,r}) \setminus \{0\}$ and define

$$v := \sum_{h \in H_k} h v_r \in W_{0,H_k}^{1,\Phi}(\Omega_N \setminus \overline{(\Omega_q)_\delta}) \setminus \{0\}.$$

By definition of I , we observe that

$$I'(tv)tv = \int_{\Omega_N \setminus \overline{(\Omega_q)_\delta}} \varphi(x, |t\nabla v|) t^2 |\nabla v|^2 dx - \int_{\Omega_N \setminus \overline{(\Omega_q)_\delta}} f(x, tv) tv dx.$$

Using

$$f(x, tv)tv \geq \lambda t^\beta |v|^\beta, \forall x \in \Omega_N \setminus \overline{(\Omega_q)_\delta} \text{ and } t \geq 0.$$

Therefore, using the hypothesis (φ_3) and (2.4), for every $t \geq 0$,

$$I'(tv)tv \leq q \xi_1(t) \int_{\Omega_N \setminus \overline{(\Omega_q)_\delta}} \Phi(|\nabla v|) dx - \lambda t^\beta \int_{\Omega_N \setminus \overline{(\Omega_q)_\delta}} |v|^\beta dx.$$

As $\beta > q > p$, we get that $I'(tv)tv \rightarrow -\infty$ as $t \rightarrow +\infty$ and $I'(tv)tv > 0$ for $t \approx 0$.

So, there exists $t_v > 0$ such that $t_v v \in W_{0,H_k}^{1,\Phi}(\Omega_N \setminus \overline{(\Omega_q)_\delta}) \setminus \{0\}$ with $I'(t_v v)t_v v = 0$. If we denote by $w = t_v v$, then

$$(5.11) \quad J_{k,r} \leq I(w) = kI(t_v v_r) = k \max_{t \geq 0} I(tv_r).$$

Following similar arguments as in the proof of [7, Lemma 3.11], we have

$$\max_{t \geq 0} I(t_v v_r) \leq \frac{1}{\lambda^{\frac{N}{\beta-N}}} \left(\frac{1}{N} - \frac{1}{\beta} \right) \frac{\left(c_1 \|\nabla v_r\|_{L^N(\Omega_N)}^N \right)^{\frac{\beta}{\beta-N}}}{\left(\|v_r\|_{L^\beta(\Omega_N)}^\beta \right)^{\frac{N}{\beta-N}}}.$$

Now, we fix $\lambda_0 = \lambda_0(k) > 0$ such that for all $\lambda \geq \lambda_0$, we have

$$(5.12) \quad \frac{k}{\lambda^{\frac{N}{\beta-N}}} \left(\frac{1}{N} - \frac{1}{\beta} \right) \frac{\left(c_1 \|\nabla v_r\|_{L^N(\Omega_N)}^N \right)^{\frac{\beta}{\beta-N}}}{\left(\|v_r\|_{L^\beta(\Omega_N)}^\beta \right)^{\frac{N}{\beta-N}}} < \left(1 - \frac{q}{\chi} \right) \min \left\{ \frac{1}{N} \left(\frac{\alpha_N}{2^{N'} \alpha} \right)^{N-1}, \frac{1}{p} S_p^{N/p} \right\}.$$

Therefore, from (5.11) and (5.12), the result follows. \square

Lemma 5.3. *If $1 \leq k < \infty$ and $\lambda \geq \lambda_0$, then $J_{k,r}$ is achieved.*

Proof. Let $(v_n) \subset \mathcal{M}_{k,r}$ a minimizing sequence for $J_{k,r}$, i.e., $(v_n) \subset W_{0,H_k}^{1,\Phi}(\Omega_r) \setminus \{0\}$ such that

$$I'(v_n)v_n = 0 \text{ and } I(v_n) \rightarrow J_{k,r}.$$

We claim that (v_n) is bounded. Assume there is some n such that $\|v_n\| \geq 1$, since otherwise (v_n) is bounded. Due to the fact $I(v_n) \rightarrow J_{k,r}$ and $J_{k,r} \leq M$, where M was given in Lemma 5.2, it follows that

$$\begin{aligned} M &\geq I(v_n) = I(v_n) - \frac{1}{\chi} I'(v_n)v_n \\ &\geq \left(1 - \frac{q}{\chi} \right) \|v_n\|^p. \end{aligned}$$

Therefore, $\|v_n\| \leq c$ if $\|v_n\| > 1$, for some constant $c > 0$ independent of n . This shows that (v_n) is bounded.

Claim:

$$I'(v_n) \rightarrow 0 \text{ in } (W_{0,H_k}^{1,\Phi}(\Omega_r))'.$$

Indeed, using the Ekeland variational Principle (see Willem [54]), there exists a sequence $(w_n) \subset \mathcal{M}_{k,r}$ such that

$$w_n = v_n + o_n(1), \quad I(w_n) \rightarrow J_{k,r}$$

and

$$(5.13) \quad I'(w_n) - \ell_n E'(w_n) = o_n(1),$$

where $(\ell_n) \subset \mathbb{R}$ and $E(w) = I'(w)w$ for $w \in W_{0,H_k}^{1,\Phi}(\Omega_r)$. Since (v_n) is bounded, we also have that (w_n) is bounded. Now, we prove that there exists $C > 0$ such that

$$(5.14) \quad |E'(w_n)w_n| > C \text{ for all } n \in \mathbb{N}.$$

Indeed, we observe that

$$(5.15) \quad \begin{aligned} -E'(w_n)w_n &= - \int_{\Omega_r} \left[\frac{\partial}{\partial t} \varphi(x, |\nabla w_n|) |\nabla w_n| + 2\varphi(x, |\nabla w_n|) \right] |\nabla w_n|^2 dx \\ &\quad + \int_{\Omega_r} \left[\frac{\partial}{\partial t} f(|x|, w_n) w_n^2 + f(|x|, w_n) w_n \right] dx \\ &\geq -q \int_{\Omega_r} \varphi(x, |\nabla w_n|) |\nabla w_n|^2 dx \\ &\quad + \int_{\Omega_r} \left[\frac{\partial}{\partial t} f(|x|, w_n) w_n^2 + f(|x|, w_n) w_n \right] dx \\ &= \int_{\Omega_r} \left[\frac{\partial}{\partial t} f(|x|, w_n) w_n^2 - (q-1) f(|x|, w_n) w_n \right] dx, \end{aligned}$$

where in the second step, we have used the hypothesis (φ_9) and in the third step, the property $I'(w_n)w_n = 0$, i.e.

$$\int_{\Omega_r} \varphi(x, |\nabla w_n|) |\nabla w_n|^2 dx = \int_{\Omega_r} f(|x|, w_n) w_n dx,$$

respectively. Since (w_n) is bounded in $W_0^{1,\Phi}(\Omega_r)$ and $J_{k,r} > 0$, there exists $w \in W^{1,\Phi}(\Omega_r) \setminus \{0\}$ such that $w_n \rightarrow w$ strongly in $L^\Phi(\Omega_r)$ and $w_n(x) \rightarrow w(x)$ a.e. in Ω_r for some subsequence. Then, by Fatou's lemma,

$$(5.16) \quad \begin{aligned} &\liminf_{n \rightarrow \infty} \int_{\Omega_r} \left[\frac{\partial}{\partial t} f(|x|, w_n) w_n^2 - (q-1) f(|x|, w_n) w_n \right] dx \\ &\geq \int_{\Omega_r} \left[\frac{\partial}{\partial t} f(|x|, w) w^2 - (q-1) f(|x|, w) w \right] dx. \end{aligned}$$

From the hypothesis (g_4) and the definition of f_2 , we obtain

$$(5.17) \quad \int_{\Omega_r} \left[\frac{\partial}{\partial t} f(|x|, w) w^2 - (q-1) f(|x|, w) w \right] dx > 0.$$

By contradiction, suppose

$$\lim_{n \rightarrow \infty} E'(w_n)w_n = 0.$$

Then letting $n \rightarrow \infty$ in (5.15) and using (5.16) along with (5.17), we obtain

$$0 = \lim_{n \rightarrow \infty} E'(w_n)w_n \geq \int_{\Omega_r} \left[\frac{\partial}{\partial t} f(|x|, w) w^2 - (q-1) f(|x|, w) w \right] dx > 0,$$

which is absurd. Therefore (5.14) holds.

From, (5.13)

$$\ell_n E'(w_n)w_n = o_n(1),$$

and so, $\ell_n = o_n(1)$. Since (w_n) is bounded we get $(E'(w_n))$ is bounded. Hence from (5.13)

$$I'(w_n) \rightarrow 0 \text{ in } \left(W_{0,H_k}^{1,\Phi}(\Omega_r)\right)'.$$

Thus, without loss generality, we may assume

$$I(v_n) \rightarrow J_{k,r} \text{ and } I'(v_n) \rightarrow 0.$$

Since (v_n) is bounded, there exists $v \in W_{0,H_k}^{1,\Phi}(\Omega_r)$ such that, for a subsequence we have

$$\begin{cases} v_n \rightharpoonup v & \text{in } W_{0,H_k}^{1,\Phi}(\Omega_r), \\ v_n(x) \rightarrow v(x), & \text{a.e. in } \Omega_r. \end{cases}$$

Now following exactly the proof of Lemma 4.6, we get $I(v_n) \rightarrow I(v) = J_{k,r}$. Hence, $J_{k,r}$ is achieved. \square

Now we establish the following Strauss-type result in Musielak-Sobolev space, which would be very useful to find a lower bound of $J_{\infty,r}$.

Lemma 5.4. *(A Strauss-type result in Musielak-Sobolev space) Assume that (φ_1) , (φ_2) , (φ_3) , $(\varphi_{10}) - (\varphi_{11})$ holds and let $v \in W^{1,\Phi}(\mathbb{R}^N)$ be a radial function. Then*

$$|v(x)| \leq \Phi^{-1} \left(x, \frac{C}{|x|^{N-1}} \int_{\mathbb{R}^N} [\Phi(x, |v|) + \Phi(x, |\nabla v|)] dx \right) \text{ a.e. in } \mathbb{R}^N,$$

where $\Phi^{-1}(x, \cdot)$ denotes the inverse function of $\Phi(x, \cdot)$ restricted to $[0, +\infty)$ and C is a positive constant independent of v .

Proof. We will establish the result for radial functions in $C_0^\infty(\mathbb{R}^N)$.

Let $v \in C_0^\infty(\mathbb{R}^N)$ be radial and let $|x| = r$, $w(r) = v(x)$. Then, from (φ_{10})

$$\Phi(b, w(b)) - \Phi(r, w(r)) = \int_r^b \left(\frac{d}{ds} \Phi(s, w(s)) \right) ds, \text{ for all } b > r > 0.$$

Since $w \in C_0^\infty([0, \infty))$, for b large enough,

$$\begin{aligned} \Phi(r, w(r)) &= - \int_r^\infty \frac{\partial}{\partial s} \Phi(s, w(s)) ds - \int_r^\infty \varphi(s, w(s)) w(s) w'(s) ds \\ (5.18) \quad &\leq \int_r^\infty \left| \frac{\partial}{\partial s} \Phi(s, w(s)) \right| ds + \int_r^\infty \varphi(s, |w(s)|) |w(s)| |w'(s)| ds. \end{aligned}$$

From (2.1), (2.2) and the Δ_2 condition (2.1), for all $s \geq 0$,

$$\begin{aligned} \varphi(s, |w(s)|) |w(s)| |w'(s)| &\leq \tilde{\Phi} \left(s, \varphi(s, |w(s)|) |w(s)| \right) + \Phi(s, |w'(s)|) \\ (5.19) \quad &\leq \Phi(s, 2|w(s)|) + \Phi(s, |w'(s)|) \\ &\leq K \Phi(s, |w(s)|) + \Phi(s, |w'(s)|). \end{aligned}$$

From (φ_{11}) , for all $s \geq 0$,

$$(5.20) \quad \left| \frac{\partial}{\partial s} \Phi(s, w(s)) \right| \leq M \Phi(s, w(s)).$$

Now using (5.19) and (5.20) in (5.18), we obtain

$$\Phi(r, w(r)) \leq (M + K + 1) \int_r^\infty [\Phi(s, |w(s)|) + \Phi(s, |w'(s)|)] ds.$$

Hence, we can conclude that

$$\Phi(r, w(r)) \leq \frac{(M + K + 1)}{r^{N-1}} \int_r^{+\infty} [\Phi(s, |w(s)|) + \Phi(s, |w'(s)|)] s^{N-1} ds.$$

From this, there is $C > 0$ such that

$$\Phi(x, v(x)) \leq \frac{C}{|x|^{N-1}} \int_{\mathbb{R}^N} [\Phi(x, |v|) + \Phi(x, |\nabla v|)] dx.$$

Since Φ is an even function, $\Phi(x, v(x)) = \Phi(x, |v(x)|)$ for all $x \in \mathbb{R}^N$ and so,

$$\Phi(x, |v(x)|) \leq \frac{C}{|x|^{N-1}} \int_{\mathbb{R}^N} [\Phi(x, |v|) + \Phi(x, |\nabla v|)] dx.$$

From this,

$$|v(x)| \leq \Phi^{-1} \left(x, \left(\frac{C}{|x|^{N-1}} \int_{\mathbb{R}^N} [\Phi(x, |v|) + \Phi(x, |\nabla v|)] dx \right) \right), \text{ for all } x \in \mathbb{R}^N \setminus \{0\},$$

where $\Phi^{-1}(x, \cdot)$ denotes the inverse function of $\Phi(x, \cdot)$ restricted to $[0, +\infty)$. Now the result follows from the density of $C_0^\infty(\mathbb{R}^N)$ in $W^{1, \Phi}(\mathbb{R}^N)$, because Φ satisfies the Δ_2 condition. \square

Lemma 5.5. *There exists $r_0 = r_0(\lambda) > 0$ such that for $\chi = \min\{\theta, \beta, p^*\}$,*

$$J_{\infty, r} \geq \left(1 - \frac{q}{\chi}\right) \min \left\{ \frac{1}{N} \left(\frac{\alpha_N}{2^{N'} \alpha} \right)^{N-1}, \frac{1}{p} S_p^{N/p} \right\}, \text{ for all } r > r_0.$$

Proof. By contradiction, suppose there exists a sequence (r_n) such that $r_n \rightarrow \infty$, satisfying

$$(5.21) \quad J_{\infty, r_n} < \left(1 - \frac{q}{\chi}\right) \min \left\{ \frac{1}{N} \left(\frac{\alpha_N}{2^{N'} \alpha} \right)^{N-1}, \frac{1}{p} S_p^{N/p} \right\}, \text{ for all } n \in \mathbb{N}.$$

First, we claim that J_{∞, r_n} is attained, for all $n \in \mathbb{N}$. In fact, for a fixed n , let $(v_k) \subset \mathcal{M}_{\infty, r_n}$ be a minimizing sequence for J_{∞, r_n} , i.e., $(v_k) \subset W_{0, H_\infty}^{1, \Phi}(\Omega_{r_n}) \setminus \{0\}$ and satisfies

$$I'(v_k)v_k = 0, \text{ and } I(v_k) \rightarrow J_{\infty, r_n}, \text{ as } k \rightarrow \infty.$$

Note that

$$(5.22) \quad \begin{aligned} o_k(1) + J_{\infty, r_n} &= I(v_k) - \frac{1}{\chi} I'(v_k)v_k \\ &\geq \left(1 - \frac{q}{\chi}\right) \int_{\Omega_{r_n}} \Phi(x, |\nabla v_k|) dx \\ &\geq \frac{1}{N} \left(1 - \frac{q}{\chi}\right) \int_{\Omega_N} |\nabla v_k|^N dx. \end{aligned}$$

Using (5.21) in (5.22),

$$\limsup_{k \rightarrow +\infty} \|\nabla v_k\|_{W^{1, N}(\Omega_N)}^N < \left(\frac{\alpha_N}{2^{N'} \alpha} \right)^{N-1}.$$

Now, we can repeat the same arguments employed in the proof of Lemma 5.3 to conclude that

$$I'(v_k) \rightarrow 0 \text{ in } (W_{0, H_\infty}^{1, \Phi}(\Omega_{r_n}))' \text{ and } v_k \rightarrow v \text{ in } W_{0, H_\infty}^{1, \Phi}(\Omega_{r_n})$$

where $v \in W_{0, H_\infty}^{1, \Phi}(\Omega_{r_n})$ is the limit of (v_k) in $W_{0, H_\infty}^{1, \Phi}(\Omega_{r_n})$. Then,

$$I(v_k) \rightarrow I(v) = J_{\infty, r_n} \text{ and } I'(v_k) \rightarrow I'(v) = 0.$$

Hence $I(v) = J_{\infty, r_n}$. Note that $v \neq 0$, since by Lemma 5.1 $J_{\infty, r_n} > 0$. Therefore $v \in \mathcal{M}_{\infty, r_n}$ and J_{∞, r_n} is attained at v .

Therefore, for each $n \in \mathbb{N}$ we can choose a sequence $\{u_n\} \subset W_{0,H^\infty}^{1,\Phi}(\Omega_{r_n}) \setminus \{0\}$ satisfying

$$I'(u_n)u_n = 0 \quad \text{and} \quad I(u_n) = J_{\infty,r_n}.$$

Proceeding as in (5.22)

$$\frac{1}{N} \left(1 - \frac{q}{\chi}\right) \left(\frac{\alpha_N}{2^{N'}\alpha}\right)^{N-1} > J_{\infty,r_n} = I(u_n) - \frac{1}{\chi} I'(u_n)u_n \geq \frac{1}{N} \left(1 - \frac{q}{\chi}\right) \int_{\Omega_N} |\nabla u_n|^N dx$$

which implies

$$(5.23) \quad \limsup_{k \rightarrow +\infty} \|\nabla u_n\|_{W^{1,N}(\Omega_N)}^N < \left(\frac{\alpha_N}{2^{N'}\alpha}\right)^{N-1}.$$

Let $\{\tilde{u}_n\}$ be a sequence given by

$$\tilde{u}_n(x) = \begin{cases} u_n(x), & \text{if } x \in \Omega_{r_n}, \\ 0, & \text{if } x \notin \Omega_{r_n}. \end{cases}$$

Observe that the following properties hold:

- (1) $\{\tilde{u}_n\} \subset W_{H^\infty}^{1,\Phi}(\mathbb{R}^N)$;
- (2) $\|\tilde{u}_n\|_{W_{H^\infty}^{1,\Phi}(\mathbb{R}^N)} = \|u_n\|_{W_{0,H^\infty}^{1,N}(\Omega_{r_n})}$;
- (3) $\tilde{u}_n \rightarrow 0$ in $W_{H^\infty}^{1,\Phi}(\mathbb{R}^N)$ because $\tilde{u}_n(x) \rightarrow 0$ a.e. in \mathbb{R}^N .

Therefore, we have

$$(5.24) \quad \int_{\mathbb{R}^N} \varphi(x, |\nabla \tilde{u}_n|) |\nabla \tilde{u}_n|^2 dx = \int_{\mathbb{R}^N} f(x, \tilde{u}_n) \tilde{u}_n dx.$$

From Lemma 5.4 we deduce that the sequence $\{\tilde{u}_n\}$ satisfies

$$|\tilde{u}_n|_\infty \rightarrow 0.$$

Using the fact that

$$\lim_{t \rightarrow 0} \frac{f(x, t)t}{\varphi(x, t)t^2} = 0, \quad \forall x \in \mathbb{R}^N,$$

and (φ_3) , it follows that given $\epsilon < \frac{p}{q\Upsilon}$, where Υ was given in Lemma 2.3, there exists $n_0 \in \mathbb{N}$ such that

$$\int_{\mathbb{R}^N} f(x, \tilde{u}_n) \tilde{u}_n dx \leq \epsilon \int_{\mathbb{R}^N} \varphi(x, |\tilde{u}_n|) |\tilde{u}_n|^2 dx \leq \epsilon q \int_{\mathbb{R}^N} \Phi(|\tilde{u}_n|) dx = \epsilon q \int_{\Omega_{r_n}} \Phi(|\tilde{u}_n|) dx, \quad n \geq n_0.$$

Since $r_n \rightarrow +\infty$, without loss of generality we can assume that $r_n \geq 1$ for all $n \in \mathbb{N}$. Therefore, by the Poincaré inequality from Lemma 2.3,

$$(5.25) \quad \int_{\mathbb{R}^N} f(x, \tilde{u}_n) \tilde{u}_n dx \leq \epsilon q \Upsilon \int_{\Omega_{r_n}} \Phi(|\nabla \tilde{u}_n|) dx, \quad \forall n \geq n_0.$$

From (5.24), (5.25) and (φ_3)

$$\begin{aligned} p \int_{\Omega_{r_n}} \Phi(|\nabla \tilde{u}_n|) dx &= p \int_{\mathbb{R}^N} \Phi(|\nabla \tilde{u}_n|) dx \\ &\leq \int_{\mathbb{R}^N} \varphi(x, |\nabla \tilde{u}_n|) |\nabla \tilde{u}_n|^2 dx \\ &\leq \epsilon q \Upsilon \int_{\Omega_{r_n}} \Phi(|\nabla \tilde{u}_n|) dx, \quad \forall n \geq n_0. \end{aligned}$$

As $\tilde{u}_n \neq 0$ for all $n \in \mathbb{N}$, we get $p \leq \epsilon q \Upsilon$, which is absurd. Hence, the result follows. \square

Lemma 5.6. *Suppose that $J_{km,r}$ is attained for some $1 \leq k < \infty$ and some $2 \leq m < \infty$. Suppose also that $J_{km,r} < J_{\infty,r}$. Then, $J_{k,r} < J_{km,r}$.*

Proof. Consider $u \in \mathcal{M}_{km,r}$ such that $I(u) = J_{km,r}$. Let $x = (\theta, \rho)$ be the polar coordinates of $x \in \mathbb{R}^2$. Then, $u = u(\theta, \rho, |y|)$, $y \in \mathbb{R}^{N-2}$. Note that

$$\Phi(\sqrt{\rho^2 + |y|^2}, |\nabla u|) = \Phi\left(\sqrt{\rho^2 + |y|^2}, \left(\frac{1}{\rho^2} u_\theta^2 + u_\rho^2 + |\nabla_y u|^2\right)^{1/2}\right).$$

Define

$$v(\theta, \rho, |y|) := u\left(\frac{\theta}{m}, \rho, |y|\right),$$

We observe that,

- (i) $v \in W_{0,H_k}^{1,\Phi}(\Omega_r)$;
- (ii) $\Phi(\sqrt{\rho^2 + |y|^2}, |\nabla v|) = \Phi\left(\sqrt{\rho^2 + |y|^2}, \left(\frac{1}{\rho^2 m^2} u_\theta^2 + u_\rho^2 + |\nabla_y u|^2\right)^{1/2}\right)$;
- (iii) $\int_{\Omega_r} F(|z|, v) dx dy = \int_{\Omega_r} F(|z|, u) dx dy$, where $z = (x, y) \in \Omega_r$.

Proceeding similarly as in the proof of Lemma 5.2, there exists $t_0 > 0$ such that $t_0 v \in \mathcal{M}_{k,r}$. For simplicity, let $v := t_0 v$. Now, since $v \in \mathcal{M}_{k,r}$,

$$J_{k,r} \leq I(v) = \int_{\Omega_r} \Phi(|z|, |\nabla v|) dx dy - \int_{\Omega_r} F(|z|, v) dx dy.$$

Using (ii) – (iii),

$$(5.26) \quad J_{k,r} \leq \int_{\Omega_r} \Phi\left(\sqrt{\rho^2 + |y|^2}, \left(\frac{1}{\rho^2 m^2} u_\theta^2 + u_\rho^2 + |\nabla_y u|^2\right)^{1/2}\right) dx dy - \int_{\Omega_r} F(|x|, u) dx dy.$$

If $I(u) = J_{km,r} < J_{\infty,r}$, then $u \notin W_{0,H_\infty}^{1,\Phi}(\Omega_r)$, then u_θ^2 is not identically zero. Therefore, using that $m > 1$,

$$\begin{aligned} & \int_{\Omega_r} \Phi\left(\sqrt{\rho^2 + |y|^2}, \left(\frac{1}{\rho^2 m^2} u_\theta^2 + u_\rho^2 + |\nabla_y u|^2\right)^{1/2}\right) dx dy \\ & < \int_{\Omega_r} \Phi\left(\sqrt{\rho^2 + |y|^2}, \left(\frac{1}{\rho^2} u_\theta^2 + u_\rho^2 + |\nabla_y u|^2\right)^{1/2}\right) dx dy, \end{aligned}$$

which together with (5.26) implies $J_{k,r} < I(u) = J_{km,r}$ and the proof is complete. \square

6. PROOF OF THE MAIN RESULTS

6.1. Proof of Theorem 1.1. By our assumption on f_1 , it is easy to verify that I is even and $I(0) = 0$. Moreover, from Lemma 4.2, Lemma 4.3 and Lemma 4.6, I satisfies all the properties in Lemma 4.1. Now from Lemma 4.7 we have there exists positive real numbers λ_m, μ_m, τ_m such that for all $\lambda \geq \lambda_m, \mu \geq \mu_m$ and $\tau \geq \tau_m$, we have

$$0 < c_1^{\lambda, \mu, \tau} \leq c_2^{\lambda, \mu, \tau} \leq \dots \leq c_m^{\lambda, \mu, \tau} < M.$$

Hence by Lemma 4.1, for every $\lambda \geq \lambda_m, \mu \geq \mu_m$ and $\tau \geq \tau_m$, the problem (P) has at least m pairs of nontrivial solutions. \square

6.2. Proof of Theorem 1.2. By Lemma 5.2, for each $n \in \mathbb{N}$, there exists $\lambda_0 = \lambda_0(n) > 0$ satisfying

$$J_{2^n, r} < \left(1 - \frac{q}{\chi}\right) \min \left\{ \frac{1}{N} \left(\frac{\alpha_N}{2^{N'\alpha}}\right)^{N-1}, \frac{1}{p} S_p^{N/p} \right\}, \quad \forall \lambda \geq \lambda_0.$$

On the other hand, by Lemma 5.5, there exists $r_0 = r_0(\lambda_0, n) > 0$ such that

$$J_{\infty, r} \geq \frac{1}{N} \left(1 - \frac{q}{\chi}\right) \left(\frac{\alpha_N}{2^{N'\alpha}}\right)^{N-1}, \quad \forall r > r_0.$$

Thus,

$$0 < J_{2^n, r} = J_{2, 2^{n-1}, r} < \left(1 - \frac{q}{\chi}\right) \min \left\{ \frac{1}{N} \left(\frac{\alpha_N}{2^{N'\alpha}}\right)^{N-1}, \frac{1}{p} S_p^{N/p} \right\} \leq J_{\infty, r},$$

for all $\lambda > \lambda_0$ and for all $r > r_0$. By Lemma 5.3, we have $J_{2^n, r}$ is attained. So, we can apply Lemma 5.6 to get

$$J_{2^{n-1}, r} < J_{2^n, r}, \quad \text{for all } \lambda > \lambda_0 \text{ and for all } r > r_0.$$

Now $J_{2^{n-2}, r}$ is also attained and satisfies

$$J_{2^{n-2}, r} = J_{2^{n-1}, r} < J_{2^n, r} < J_{\infty, r}.$$

Again, by Lemma 5.6, we get $J_{2^{n-2}, r} < J_{2^{n-1}, r}$. Inductively,

$$0 < J_{2, r} < J_{2^2, r} < \dots < J_{2^n, r} < J_{\infty, r},$$

for all $\lambda > \lambda_0$ and all $r > r_0$.

Noting Lemma 4.2, from Lemma 5.3, minimizers of $J_{k, m}$ are critical points of I in $W_{0, H_k}^{1, \Phi}(\Omega_r)$. Now, applying the Principle of symmetric criticality from [48], it follows that they are critical points of I in $W_0^{1, \Phi}(\Omega_r)$ and therefore are solutions of (P). Note that, due to Lemma 5.1, such solutions are nontrivial. Therefore, all minimizers of $J_{2^m, r}$, $m = 1, \dots, n$ are nonradial, rotationally nonequivalent and nontrivial solutions of (P). \square

REFERENCES

- [1] Adimurthi and S.L. Yadava, *Critical exponent problem in \mathbb{R}^2 with Neumann boundary condition*, Comm. Partial Differential Equations 15 (4) (1990) 461-501. [10](#)
- [2] C.O. Alves and J.L.P. Barreiro, *Existence and multiplicity of solutions for a $p(x)$ -Laplacian equation with critical growth*, J. Math. Anal. Appl. 403(2013), 143-154 [1](#)
- [3] C.O. Alves and M.C. Ferreira, *Multi-bump solutions for a class of quasilinear problems involving variable exponents*, Annali di Matematica 194(2015), 1563-1593. [1](#)
- [4] C.O. Alves, G.M. Figueiredo and J.A. Santos, *Strauss and Lions type results for a class of Orlicz-Sobolev spaces and applications*, Topol. Methods Nonlinear Anal. 44, no. 2, (2014), 435-456. [1](#)
- [5] C.O. Alves and L. R. de Freitas, *Multiplicity of nonradial solutions for a class of quasilinear equation on annulus with exponential critical growth*, Top. Meth. in Nonlinear Analysis 39(2012), 243-262. [5, 6, 9](#)
- [6] C.O. Alves and M.A.S. Souto, *Existence of solutions for a class of problems in \mathbb{R}^N involving the $p(x)$ -Laplacian*, Progr. Nonlinear Differential Equations Appl. 66 (2005) 17-32. [1](#)
- [7] C. O. Alves, P. Garain and V. D. Rădulescu, *High perturbations of quasilinear problems with double criticality*. To appear in Math. Zeitschrift. [2, 4, 6, 10, 12, 14, 22](#)
- [8] C.O. Alves and V. Rădulescu, *The Lane-Emden equation with variable double-phase and multiple regime*, Proc. Amer. Math. Soc. 148 (2020), 2937-2952. [2](#)
- [9] A. Ambrosetti and P.H. Rabinowitz, *Dual methods in critical point theory and applications*, J. Funct. Analysis 14, 347-381 (1973). [4](#)
- [10] E. Azroul, A Benkirane, M. Shimi and M. Srati, *Embedding and extension results in fractional Musielak-Sobolev spaces*, arXiv:2007.11043v1[Math AP] [2](#)
- [11] A. Benkirane and M. Sidi El Vally, *An existence result for nonlinear elliptic equations in Musielak-Orlicz-Sobolev spaces*, Bull. Belg. Math. Soc. 20 (1) (2013) 1-187. [2](#)
- [12] J.M. Bezerra do Ó, E. S. Medeiros and U. Severo, *On a quasilinear nonhomogeneous elliptic equation with critical growth in \mathbb{R}^N* , J. Diff. Equations 246 (2009), 1363-1386. [11](#)

- [13] J. Byeon, *Existence of many nonequivalent nonradial positive solutions of semilinear elliptic equations on three-dimensional annuli*, J. Differential Equations 136 (1997), 136–165. [5](#)
- [14] H. Brezis, L. Nirenberg, *Positive solutions of nonlinear elliptic equations involving critical Sobolev exponents*. Comm. Pure Appl. Math. 36, 437–477 (1983) [4](#)
- [15] M.L.M. Carvalho, E.D. Silva, J. V.A. Gonçalves and C. Goulart, *Critical elliptic problems using Concave-concave nonlinearities*, Ann. Mat. Pura Appl. (2019), 693-726. [1](#)
- [16] A. Castro and B.M. Finan, *Existence of many positive nonradial solutions for a superlinear Dirichlet problem on thin annuli*, Nonlinear Differential Equations 5 (2000), 21–31. [5](#)
- [17] F. Catrina and Z.-Q. Wang, *Nonlinear elliptic equations on expanding symmetric domains*, J. Differential Equations 156 (1999), 153–181. [5](#)
- [18] P. Cherrier, *Meilleures constantes dans les inegalites relatives aux espaces de Sobolev*, Bull. Sci. Math. 108 (1984) 225-262. [10](#)
- [19] J. Chabrowski and Y. Fu, *Existence of solutions for $p(x)$ -laplacian problems on a bounded domain*. J. Math. Anal. Appl 306 (2005), 604–618. [2](#)
- [20] I. Chlebicka, *A pocket guide to nonlinear differential equations in the Musielak-Orlicz spaces*, Nonlinear Anal. 175 (2018), 1-27. [1](#), [6](#)
- [21] A. Cianchi, *Moser-Trudinger inequalities without boundary conditions and isoperimetric problems*, Indiana Univ. Math. J. 54 (2004), 669-706 [6](#), [10](#)
- [22] C. COFFMAN, *A non-linear boundary value problem with many positive solutions*, J. Differential Equations 54 (1984), 429–437. [5](#)
- [23] E.S. de Medeiros, *Existência e concentração de solução para o p -Laplaciano com condição de Neumann*, Doctoral Dissertation, UNICAMP, 2001. [6](#), [9](#)
- [24] L. Diening, P. Hästö, P. Harjulehto and M. Ruzicka, *Lebesgue and Sobolev Spaces with Variable Exponents*, Springer Lecture Notes, vol. 2017, Springer-Verlag, Berlin, 2011. [1](#)
- [25] D.G. de Figueiredo and O.H. Miyagaki, *Multiplicity of non-radial solutions of critical elliptic problems in an annulus*, Proc. Roy. Soc. Edinburgh Sect. A 135 (2005), 25–37. [5](#), [6](#)
- [26] X.L. Fan, *Differential equations of divergence form in Musielak-Sobolev spaces and a sub-supersolution method*, J. Math. Anal. Appl. 386 (2012) 593-604 [2](#), [6](#), [8](#)
- [27] X.L. Fan and Q.H. Zhang, *Existence of solutions for $p(x)$ -Laplacian Dirichlet problem*, Nonlinear Anal. 52 (2003) 1843-1852. [2](#)
- [28] X.L. Fan, *On the sub-supersolution method for $p(x)$ -Laplacian equations*, J. Math. Anal. Appl. 330 (2007) 665-682. [2](#)
- [29] N. Fukagai, M. Ito and K. Narukawa, *Positive solutions of quasilinear elliptic equations with critical Orlicz-Sobolev nonlinearity on \mathbb{R}^N* , Funkcial. Ekvac. 49, no. 2, (2006) 235-267. [1](#), [2](#), [7](#)
- [30] N. Fukagai and K. Narukawa, *On the existence of multiple positive solutions of quasilinear elliptic eigenvalue problems*, Ann. Mat. Pura Appl. 186, no. 3, (2007) 539-564. [1](#)
- [31] J. Garcia Azorero and I. Peral Alonso, *Multiplicity of solutions for elliptic problems with critical exponent or with a nonsymmetric term*, Trans. Amer. Math. Soc. 2 (1991) 877-895. [16](#), [17](#)
- [32] J. Gossez, *Nonlinear elliptic boundary value problems for equations with rapidly (or slowly) increasing coefficients*, Trans. Amer. Math. Soc. 190 (1974), 163-205. [8](#)
- [33] P. Harjulehto and P. Hästö, *Orlicz Spaces and Generalized Orlicz Spaces*, Springer, 2019. [1](#), [6](#)
- [34] N. Hirano and N. Mizoguchi, *Nonradial solutions of semilinear elliptic equations on annuli*, J. Math. Soc. Japan 46 (1994), 111–117. [5](#)
- [35] H. Hudzik, *On generalized Orlicz-Sobolev space*, Funct. Approx. Comment. Math. 4 (1976) 37-51. [1](#)
- [36] O. Kavian, *Introduction a la Theorie Des Points Critiques: Et Applications Aux Problemes Elliptiques*, Springer, Heidelberg 1993. [15](#)
- [37] B. S. Kaur and K. Sreenadh, *Multiple positive solutions for a quasilinear elliptic equation with critical exponential nonlinearity*, Nonlinear Anal. 73 (2010) 2368-2382 [10](#)
- [38] O. Kováčik and J. Rákosník, *On spaces $L^{p(x)}$ and $W^{k,p(x)}$* , Czechoslovak Math. J., 41 (1991) 592–618 [1](#)
- [39] Y.Y. Li, *Existence of many positive solutions of semilinear elliptic equations on annulus*, J. Differential Equations 83 (1990), 348–367. [5](#)
- [40] S.S. Lin, *Existence of many positive nonradial solutions for nonlinear elliptic equations on an annulus*, J. Differential Equations 103 (1993), 338–349. [5](#)
- [41] D. Liu and P. Zhao, *Solutions for a quasilinear elliptic equation in Musielak-Sobolev spaces*, Nonlinear Analysis: Real World Applications, 26 (2015), 315-329. [2](#)
- [42] N. Mizoguchi and T. Suzuki, *Semilinear elliptic equations on annuli in three and higher dimensions*, Houston J. Math. 22 (1996), 199–215. [5](#)

- [43] J. Musielak, *Orlicz Spaces and Modular Spaces*, Lecture Notes in Mathematics, vol. 1034, Springer-Verlag, Berlin, 1983. [1](#), [6](#)
- [44] L. Pick, A. Kufner, O. John and S. Fučík, *Function spaces*, Volume 1, 2nd Revised and Extended Edition, De Gruyter, 2013. [1](#), [7](#)
- [45] Paul H. Rabinowitz, *Minimax methods in critical point theory with applications to differential equations*, volume 65 of *CBMS Regional Conference Series in Mathematics*. Published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 1986. [11](#)
- [46] V.D. Rădulescu and D.D. Repovš, *Partial Differential Equations with Variable Exponents: Variational Methods and Qualitative Analysis*, CRC Press, Taylor & Francis Group, Boca Raton FL, 2015. [2](#)
- [47] Elves A. B. Silva and Magda S. Xavier, Multiplicity of solutions for quasilinear elliptic problems involving critical Sobolev exponents. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 20(2):341–358, 2003. [4](#), [6](#)
- [48] M. Squassina, *On Palais' principle for non-smooth functionals*, *Nonlinear Analysis* 74 (2011) 3786–3804. [28](#)
- [49] M. Struwe, *Variational Methods*, Springer-Verlag, Berlin, 1990. [10](#)
- [50] T. Suzuki, *Positive solutions for semilinear elliptic equations on expanding annulus: mountain pass approach*, *Funkcial. Ekvac.* 39 (1996), 143–164. [5](#)
- [51] L. Wang and D. Liu, *On a compact trace embedding theorem in Musielak-Sobolev spaces*, arXiv:1911.10331v1[Math.FA] 23 Nov 2019. [2](#)
- [52] Zhi Hui Wei and Xin Min Wu, A multiplicity result for quasilinear elliptic equations involving critical Sobolev exponents. *Nonlinear Anal.*, 18(6):559–567, 1992. [4](#), [6](#)
- [53] Z. Wang and M. Willem, *Existence of many positive solutions of semilinear elliptic equations on an annulus*, *Proc. Amer. Math. Soc.* 127 (1999), 1711–1714. [5](#)
- [54] M. Willem, *Minimax Theorems*, Birkhäuser Boston, Inc., Boston, MA, 1996. [23](#)

(Karima Ait-Mahiout)
 LABORATOIRE "THÉORIE DU POINT FIXE ET APPLICATIONS",
 ÉCOLE NORMALE SUPÉRIEURE, BP 92,
 KOUBA, 16006, ALGIERS, ALGERIA
 EMAIL: karima_ait@hotmail.fr

(Claudianor O. Alves)
 UNIDADE ACADÊMICA DE MATEMÁTICA
 UNIVERSIDADE FEDERAL DE CAMPINA GRANDE
 58429-970, CAMPINA GRANDE - PB, BRAZIL
 EMAIL: coalves@mat.ufcg.edu.br

(Prashanta Garain)
 DEPARTMENT OF MATHEMATICS
 BEN-GURION UNIVERSITY OF THE NEGEV
 P.O.B. 653
 BEER SHEVA 8410501, ISRAEL
 EMAIL: pgarain92@gmail.com