A note on 1-2-3 and 1-2 Conjectures for 3-regular graphs

Jing-zhi Chang¹, Chao Yang^{1,*}, Zhi-xiang Yin¹, Bing Yao²

- School of Mathematics, Physics and Statistics; Center of Intelligent Computing and Applied Statistics,
 Shanghai University of Engineering Science, Shanghai, 201620, China
 - 2. College of Mathematics and Statistics, Northwest Normal University, Lanzhou, 730070, China

Abstract: The 1-2-3 Conjecture, posed by Karoński, Łuczak and Thomason, asked whether every connected graph G different from K_2 can be 3-edge-weighted so that every two adjacent vertices of G get distinct sums of incident weights. The 1-2 Conjecture states that if vertices also receive colors and the vertex color is added to the sum of its incident edges, then adjacent vertices can be distinguished using only $\{1,2\}$. In this paper we confirm 1-2 Conjecture for 3-regular graphs. Meanwhile, we show that every 3-regular graph can achieve a neighbor sum distinguishing edge coloring by using 4 colors, which answers 1-2-3 Conjecture positively.

Keywords: 1-2-3 Conjecture, 1-2 Conjecture, 3-regular graphs

MSC(2010): 05C15

1 Introduction

All considered graphs are finite, undirected, simple and connected. Let N(u) denote the set of neighbors of vertex u and $0 \le s < t$. Let $d_G(v)$ be the degree of vertex v of G. For general theoretic notations, we follow [3].

Karónski et al.[7] introduced and investigated a coloring of the edges of a graph with positive integers so that adjacent vertices have different sums of incident edge colors. More precisely, let $f: E \to [k] = \{1, 2, ..., k\}$ be an edge coloring of G (such a coloring is also called a k-coloring of G). For $x \in V$, set

$$\sigma(x) = \sum_{e \ni x} f(e).$$

A k-coloring f of G is called neighbor sum distinguishing edge coloring if $\sigma(x) \neq \sigma(y)$ for every edge $xy \in E(G)$. In other words, the vertex coloring σ induced by f in the above described way must be proper. The minimum integer k for which there is a neighbor sum distinguishing edge coloring of a graph G will be denoted by $\operatorname{gndi}_{\Sigma}(G)$.

In [7] Karónski, Łuczak and Thomason posed the following elegant problem, known as the 1-2-3 Conjecture.

Conjecture 1. Let G be a connected graph and $G \neq K_2$. Then $gndi_{\Sigma}(G) \leq 3$.

Supported by National Natural Science Foundation of China under Grant Nos. 61672001, 61662066, 62072296.

^{*}Corresponding author. E-mail address: yangchaomath0524@163.com

Toward the Conjecture 1, Kalkowski, Karoński and Pfender [6] showed that if G is a k-colorable graph with k odd then G admits a vertex-coloring k-edge-weighting. So, for the class of 3-colorable graphs, including bipartite graphs, the answer is affirmative. However, in general, this question is still open. Addario-Berry et al.[1] showed that every graph without isolated edges has a proper k-weighting when k = 30. After improvements to k = 15 in [2] and k = 13 in [10], Kalkowski, Karóński, and Pfender [6] showed that every graph without isolated edges has a proper 5-weighting. Przybylo [9] showed that every d-regular graph with $d \ge 2$ admits a vertex-coloring edge 4-weighting and every d-regular graph with $d \ge 10^8$ admits a vertex-coloring edge 3-weighting.

In [8] the following problem related to 1-2-3 Conjecture was introduced. Let $f: E \cup V \to [k] = \{1, 2, ..., k\}$ be a total k-coloring of a graph G = (V, E). For every vertex v, let

$$t(v) := f(v) + \sum_{u \in N(v)} f(uv) = f(v) + \sigma(v),$$

Thus, t(v) is the sum of incident colors of v and the color of v. We say that f is a neighbor total sum distinguishing total coloring of G if $t(u) \neq t(v)$ for all adjacent vertices u, v in G. Similarly as above, the minimum value of k for which there exists a total neighbor sum distinguishing coloring of a graph G will be denoted by $\operatorname{tgndi}_{\Sigma}(G)$.

In [8] Przybylo and Wózniak posed the following problem, known as the 1-2 Conjecture.

Conjecture 2. Let G be a connected graph. Then $\operatorname{tgndi}_{\Sigma}(G) \leq 2$.

Up to now, it is known that for every graph G, $\operatorname{tgndi}_{\Sigma}(G) \leq 3$ (see [5]).

This paper is organized as follows. In Section 2, we offer an important structural lemma that every connected graph G contains a m-partite spanning subgraph H such that $(1-\frac{1}{m})d_G(v) \leq d_H(v)$. Naturally, every 3-regular graph G has a maximal bipartite spanning subgraph H such that G - E(H) is either isolated vertices or isolated edges. Via the structural property between H and G - E(H), we design a coloring algorithm to compute $\operatorname{gndi}_{\Sigma}(G)$ and $\operatorname{tgndi}_{\Sigma}(G)$ for 3-regular graphs G in Section 3 and Section 4, respectively. We show that $\operatorname{gndi}_{\Sigma}(G) \leq 4$ and $\operatorname{tgndi}_{\Sigma}(G) \leq 2$ for any 3-regular graph G. Therfore, the first result answers 1-2-3 Conjecture positively and the second one confirms 1-2 Conjecture.

2 Structural Lemma

Lemma 3. Let G be a graph on n vertices. Then it exists a m-partite spanning subgraph H such that $(1-\frac{1}{m})d_G(v) \leq d_H(v)$ for all $v \in V(G)$, where m is a positive integer and $m \leq n$.

Proof Let H be a maximal m-partite spanning subgraph of G with the greatest possible number of edges. Let $\{V_1, V_2, \ldots, V_m\}$ be the m-partition of V(H) and let $v \in V_1$, $d_{V_i}(v) = |N_{V_i}(v)|$, $N_{V_i}(v) = \{u : u \in V_i, uv \in E(G)\}$, $i = 1, 2, \ldots, m$. Then $d_{V_1}(v) \leq d_{V_i}(v)$, $i = 1, 2, \ldots, m$. Otherwise, it exists an i_0 such that $d_{V_1}(v) > d_{V_{i_0}}(v)$, and we use $V_1 \setminus \{v\}$, $V_{i_0} \cup \{v\}$ instead of V_1 , V_{i_0} , respectively, and then it generates a new maximal m-partite spanning subgraph H' of G. Obviously, $\varepsilon(H') > \varepsilon(H)$, a contradiction. Therefore,

$$(m-1)d_{V_1}(v) \le \sum_{i=2}^m d_{V_i}(v) = d_H(v),$$

where $d_H(v) = |N_H(v)|$.

Let $d_G(v) = |N_G(v)|$. Then

$$d_G(v) = d_{V_1}(v) + d_H(v) \le \frac{1}{m-1}d_H(v) + d_H(v) = \frac{m}{m-1}d_H(v)$$

Hence

$$(1 - \frac{1}{m})d_G(v) \le d_H(v)$$

for all $v \in V(G)$.

By Lemma 3, every 3-regular graph G contains a maximal bipartite spanning subgraph H such that G - E(H) is either isolated vertices or isolated edges. Via the structural between H and G - E(H), we further study $gndi_{\Sigma}(G)$ and $tgndi_{\Sigma}(G)$ of 3-regular graphs G.

For the sake of narrative, we fix some natation. Let $G=(V_X,V_Y,E_X,E_Y,E_H)$ be a 3-regular graph with vertex partition (V_X,V_Y) and edge partition (E_X,E_Y,E_H) , where E_X,E_Y and E_H represent the edge sets in V_X,Y_Y and H, respectively. The maximal spanning bipartite graph H is the graph with vertex set $\{x_i:1\leq i\leq m\}\cup\{y_i:1\leq i\leq n\}$ and edge set $\{x_iy_i:1\leq i\leq m,1\leq i\leq n\}$. We use a_1 and b_1 to denote the number of vertices with degree 2 and 3 in X, respectively. Let a_2 and b_2 denote the number of vertices with degree 2 and 3 in Y, respectively. We use e_{x_i} for $i=1,2,\ldots,\frac{a_1}{2}$ to represent the edge with two endpoints v_{x_i}, v'_{x_i} in X. Let e_{y_j} for $j=1,2,\ldots,\frac{a_2}{2}$ to represent the edge with two endpoints v_{y_j}, v'_{y_j} in Y. In the bipartite graph H, let $v_2(x)$ and $v_3(x)$ be the vertex with degree 2 and 3 in X, respectively. Similarly, $v_2(y)$ and $v_3(y)$ denote the vertex with degree 2 and 3 in Y, respectively. The edge between $v_2(x)$ and $v_2(y)$ is denoted by $\widetilde{e_2}$, and the edge connects $v_2(x)$ (or $v_3(x)$) and $v_3(y)$ (or $v_2(y)$) is denoted by $\widetilde{e_{2-3}}$.

3 $\operatorname{gndi}_{\Sigma}(G)$ for 3-regular graphs

Theorem 4. For any 3-regular graph G, $gndi_{\Sigma}(G) \leq 4$.

Proof Case 1 $a_1 = a_2 = 0$.

This case implies that G is a 3-regular complete bipartite graph. This result has been proved by Chang et al. [4].

Case 2 $a_1 = b_2 = 0$ or $b_1 = a_2 = 0$.

Suppose that $a_1 = b_2 = 0$. We color all edges in E_H and E_Y with 1 and 2, respectively. Then $\sigma(v_x) = 3$, $\sigma(v_y) = 4$. To assure that $\sigma(v_{y_j}) \neq \sigma(v'_{y_j})$, recolor an incident edge of v_{y_j} (or v'_{y_j}) with 3. Then $\sigma(v_x)$ belongs to $\{3, 5, 7, 9\}$ and $\sigma(v_y)$ belongs to $\{4, 6\}$.

Case 3 $b_1 = b_2 = 0$.

Form Lemma 3, G contains a maximal bipartite spanning subgraph H such that G-E(H) is either isolated vertices or isolated edges. We color all edges in E_H and E_X with 1, and color all edges in E_Y with 2. For any edge $v_{x_i}v'_{x_i}$ in X, select one edge e_z from E_H such that v_{x_i} (or v'_{x_i}) is an endpoints of e_z and recolor edge e_z with 3. Assume that v_{x_i} and v_{y_j} are connected by e_z and all incident edges (except for $e_z, v_{x_i}v'_{x_i}, v_{y_j}v'_{y_j}$) of $v_{x_i}, v'_{x_i}, v_{y_j}, v'_{y_j}$ keep the color 1 as before, and we call these edges being dominated, see Fig.1. Suppose that v'_{x_i} and v_{y_j} are connected by e_z . Then $\sigma(v_{x_i}) = 5$, $\sigma(v'_{x_i}) = 3$, $\sigma(v_{y_j}) = 6$, $\sigma(v'_{y_j}) = 4$. Continue this procedure $\frac{a_1}{2}$ times until the weights of all adjacent vertices in G are distinct. Now we prove it feasibility, namely, it verifies that there exists at least one edge in E_H which can not be dominated after $\frac{a_1}{2} - 1$ operations. Suppose that all edges in H are dominated after $\frac{a_1}{2} - 1$ operations, and if there still exists a pair of adjacent vertices v_{x_k} and v'_{x_k} (or v_{y_k} and v'_{y_k}) having the same weights, then the four incident edges of v_{x_k} and v'_{x_k} in H must receive the same color 1. By our coloring rule, it is impossible.

Case 4 $a_1 \neq a_2, b_1 \neq b_2$ and they are positive integers.

Fig.1: Edges labelled by solid line are dominated.

We color all edges in E_H with 1, color all edges in E_X with 2 and color all edges in E_Y with 3. By the technique in Case 3, we can distinguish all adjacent vertices whose joined by $\widetilde{e_2}$. But we still need to distinguish adjacent vertices whose joined by $\widetilde{e_{2-3}}$. Select an incident edge from $\widetilde{e_{2-3}}$ and color it with 3, it deduces that $\sigma(v_{x_i}) \neq \sigma(v'_{x_i})$ and $\sigma(v_{y_j}) \neq \sigma(v'_{y_j})$. Moreover, $\sigma(v_2(x)) = 4$ or 6, $\sigma(v_2(y)) = 5$ or 7, $\sigma(v_3(x))$ belongs to set $\{3, 5, 7, 9\}$ and $\sigma(v_3(y))$ belongs to set $\{3, 5, 7, 9\}$. Possibly, there may exist some cases that the weight of adjacent vertices can not distinguish. We deal with it as follows.

Case 4.1 $\sigma(v_3(x)) = 3$.

- Case 4.1.1 Vertex $v_3(x)$ connects two vertices of degree 2 and one vertex of degree 3 in Y. Let $N(v_3(x)) = \{v_2(y_1), v_2(y_2), v_3(y_3)\}$. Then $\sigma(v_3(x)) = \sigma(v_3(y_3)) = 3$.
- 1. Vertex $v_2(y_1)$ or $v_2(y_2)$ has weight 7. Assume that $\sigma(v_2(y_1)) = 7$. Recolor $v_3(x)v_2(y_1)$ with 2, then $\sigma(v_3(x)) = 4$, $\sigma(v_2(y_1)) = 8$.
- 2. $\sigma(v_2(y_1)) = \sigma(v_2(y_2)) = 5$. We may recolor any incident edge of $v_2(y_1)$ or $v_2(y_2)$. Assume that recolor $v_3(x)v_2(y_1)$ with 4, then $\sigma(v_3(x)) = 4$ and $\sigma(v_2(x_1)) = 8$.
- Case 4.1.2 Vertex $v_3(x)$ connects two vertices of degree 3 and a vertex of degree 2 in Y. Let $N(v_3(x)) = \{v_3(y_1), v_3(y_2), v_2(y_3)\}$. Then $v_3(x)$ and its two neighbors all have weight 3 and $\sigma(v_2(y_3)) = 5$ (or 7). Recolor $v_3(x)v_2(y_3)$ with 2 (or 4). Then $\sigma(v_3(x)) = 4$ or 6, $\sigma(v_2(y_3)) = 8$.
- Case 4.1.3 Vertex $v_3(x)$ connects three vertices of degree 3 in Y. Denoted by $v_3(y_1)$, $v_3(y_2)$ and $v_3(y_3)$, respectively. If three neighbors of $v_3(x)$ have the same weight 3, then recolor all incident edges of $v_3(x)$ with 3, it follows that $\sigma(v_3(x)) = 9$ and $\sigma(v_3(y_1)) = \sigma(v_3(y_2)) = \sigma(v_3(y_3)) = 5$.

Case 4.2 $\sigma(v_3(y)) = 5$.

Vertex $v_3(x)$ connects a vertex of degree 2 with weight 7 in Y. Denote the vertex of degree 2 by $v_2(y_0)$. Recolor $v_3(x)v_2(y_0)$ with 4. Then $\sigma(v_3(x)) = 6$ and $\sigma(v_2(y_0)) = 8$.

Case 4.3 $\sigma(v_3(x)) = 7$.

Vertex $v_3(x)$ must connect two vertices of degree 2 in Y. Let $N(v_3(x)) = \{v_2(y_1), v_2(y_2), v_k\}$. Here we need not consider the weight of v_k . Recolor $v_3(x)v_2(y_1)$ and $v_3(x)v_2(y_2)$ with 4. Then $\sigma(v_3(x)) = 9$ and $\sigma(v_2(y_1)) = \sigma(v_2(y_2)) = 8$.

Przybylo [9] showed that every d-regular graph with $d \ge 2$ admits a vertex-coloring edge 4-weighting and every d-regular graph with $d \ge 10^8$ admits a vertex-coloring edge 3-weighting. From the proof of Theorem 4, we characterize the structural that some edges of 3-regular graphs receive color 4 when $a_1 \ne a_2$ and $b_1 \ne b_2$. In other words, if Conjecture 1 is true for every 3-regular graph, then it requires a method to deal with these edges which are colored by 4.

4 $\operatorname{tgndi}_{\Sigma}(G)$ for 3-regular graphs

Theorem 5. For any 3-regular graph G, $\operatorname{tgndi}_{\Sigma}(G) \leq 2$.

Proof Case 1 $a_1 = a_2 = 0$.

This case implies that G is a 3-regular complete bipartite graph. We color all vertices in X with 1, color all vertices in Y with 2 and color all edges in E_H with 1. Then $t(v_x) = 4$ and $t(v_y) = 5$.

Case 2 $a_1 = b_2 = 0$ or $b_1 = a_2 = 0$.

Without loss of generality, assume that $a_1 = b_2 = 0$. We color all vertices in H with 1, color all edges in E(H) with 1, and color all edges in E_Y with 2. To assure that $t(v_{y_j}) \neq t(v'_{y_j})$, recolor v_{y_j} (or v'_{y_j}) with 2. Then $t(v_x) = 4$ and $t(v_y)$ is 5 or 6.

Case 3 $b_1 = b_2 = 0$.

We color all vertices in X with 1, color all vertices in Y with 2, color all edges in E_H and E_X with 1, and color all edges in E_Y with 2. For any edge $v_{x_i}v'_{x_i}$ in G, select one edge e_z from E_H such that v_{x_i} (or v'_{x_i}) is an endpoints of e_z and recolor edge e_z with 2, meanwhile, all incident edges (except for $e_z, v_{x_i}v'_{x_i}, v_{y_j}v'_{y_j}$) of $v_{x_i}, v'_{x_i}, v_{y_j}, v'_{y_j}$ keep the color 1 as before, namely they are dominated, see Fig.1. Without loss of generality, assume that v_{x_i} and v_{y_j} are connected by e_z . Then $t(v_{x_i}) = 5$, $t(v'_{x_i}) = 4$, $t(v_{y_j}) = 6$, $t(v'_{y_j}) = 7$. Continue this procedure $\frac{a_1}{2}$ times until the weights of all adjacent vertices in G are distinct. The feasibility of this method is similar to Case 3 in Theorem 4.

Case 4 $a_1 \neq a_2, b_1 \neq b_2$ and they are positive integers.

We color all vertices in X with 1, color all vertices in Y with 2, color all edges in E_H with 2, color all edges in E_X with 1 and color all edges in E_Y with 2. Using the technique in Case 3 again, we change the color of e_z from 2 to 1, and change the weight of vertices which are connected only by $\widetilde{e_2}$. By this way, we can distinguish all adjacent vertices whose are connected by $\widetilde{e_2}$. But we still need to distinguish adjacent vertices whose joined by $\widetilde{e_{2-3}}$. Select an incident edge from $\widetilde{e_{2-3}}$ and color it with 1, it deduces that $t(v_{x_i}) \neq t(v_{x_i}')$ and $t(v_{y_i}) \neq t(v_{y_i}')$. Meanwhile, $t(v_2(x)) = 5$ or 6, $t(v_2(y)) = 7$ or 8, $t(v_3(x))$ belongs to set $\{7,6,5,4\}$ and $t(v_3(y))$ belongs to set $\{8,7,6,5\}$. Possibly, it still exist some cases that vertex can not distinguish. We deal it as follows.

Case 4.1 $t(v_3(y)) = 5$. Vertex $v_3(y)$ must join three vertices of degree 2 in X. Recolor $v_3(y)$ with 1, then $t(v_3(y)) = 4$ and $t(v_2(x)) = 5$.

Case 4.2 $t(v_3(y)) = 6$. Vertex $v_3(y)$ must connect two vertices of degree 2 in X, denoted by $v_2(x_1)$ and $v_2(x_2)$, respectively. Set $v_0 \in N(v_3(x)) - \{v_2(x_1), v_2(x_2)\}$. Let v_0' be the neighbor of v_0 .

Case 4.2.1 $t(v_0) = t(v_3(y)) = 6$ and $t(v_0') \neq 6$. Recolor $v_3(y)$, $v_0v_3(y)$ with 1 and v_0 with 2, then we have $t(v_3(y)) = 4$ and $t(v_0) = 6$.

Case 4.2.2 $t(v_0) = t(v_3(y)) = t(v'_0) = 6$. If $d_H(v_0) = 2$, then v_0 connects two vertices with degree 3 in Y and denoted by $v_3(y_1)$ and $v_3(y_2)$, respectively. Recolor v_0 with 2, it gets that $t(v_0) = 7$. If $d_H(v_0) = 3$, then v_0 connects two vertices of degree 3 in Y and denoted by $v_3(y_3)$ and $v_3(y_4)$, respectively. Denote $v_1 \in N(v_0) - \{v_3(y_3), v_3(y_4)\}$. Recolor v_0 and v_0v_1 with 2, and recolor v_1 with 1. Then $t(v_0) = 8$ and $t(v_1) = 7$.

Case 4.3 $t(v_3(x)) = 7$. Suppose that $N(v_3(x)) = \{v_{y_1}, v_{y_2}, v_{y_3}\}$. Let v_{x_1}, v'_{x_1} be two neighbors of v_{y_1} and v_{x_2}, v'_{x_2} be two neighbors of v_{y_2} . Then the following cases may appear adjacent vertices having the same weight.

Case 4.3.1 $t(v_{y_1})$, $t(v_{y_2})$ and $t(v_{y_3})$ are not equal to 8. Then it exists a vertex of $N(v_3(x))$ having weight 7 (otherwise, $v_3(x)$ and its neighbors can be distinguished immediately). Recolor $v_3(x)$ with 2, it follows that $t(v_3(x)) = 8$.

Case 4.3.2. One of $\{v_{y_1}, v_{y_2}, v_{y_3}\}$ has weight 8 and another two vertices have weight 7. Without loss of generality, suppose that $t(v_{y_3}) = 8$ and $t(v_{y_1}) = t(v_{y_2}) = 7$.

Case 4.3.2.1 $d_H(v_{y_1}) = d_H(v_{y_2}) = 2$.

1. If $t(v_{x_1}) = t(v_{x_2}) = 5$, recolor $v_{y_1}v_3(x)$ and $v_{y_2}v_3(x)$ with 1, then $t(v_{y_1}) = t(v_{y_2}) = 6$, $t(v_{y_3}) = 8$ and $t(v_{x_1}) = t(v_{x_2}) = t(v_3(x)) = 5$.

- 2. Without loss of generality, assume that $t(v_{x_1}) = 6$. Recolor v_{y_1} and $v_{y_1}v_3(x)$ with 1. Then $t(v_{y_1}) = 5$ and $t(v_3(x)) = 6$.
- Case 4.3.2.2 $d_H(v_{y_1}) = 2$ and $d_H(v_{y_2}) = 2$ hold not at the same time. Assume that $d_H(v_{y_1}) = 3$ and $t(v'_{x_1}) = 5$. Recolor $v_{y_1}v_{x_1}$, $v_{y_1}v_3(x)$ and v_{y_1} with 1, and recolor v_{x_1} with 2, then $t(v_{y_1}) = 4$, $t(v_3(x)) = 6$ and $t(v_{x_1})$ keep the same as before.
- Case 4.3.3 $\{t(v_{y_1}), t(v_{y_2}), t(v_{y_3})\} = \{6, 7, 8\}$. Assume that $t(v_{y_1}) = 8$, $t(v_{y_2}) = 6$ and $t(v_{y_3}) = 7$. Recolor $v_{y_2}v_3(x)$ and v_{y_2} with 1. Then $t(v_{y_2}) = 4$ and $t(v_3(x)) = 6$.
- Case 4.3.4 Two of $\{v_{y_1}, v_{y_2}, v_{y_3}\}$ have weights 8 and the remaining one has weight 7. Assume that $t(v_{y_1}) = 8$, $t(v_{y_2}) = 7$ and $t(v_{y_3}) = 8$.
- 1. If $d_H(v_{y_2}) = 2$ and $t(v_{x_2}) = 5$, recolor $v_{y_2}v_3(x)$ with 1 and recolor $v_3(x)$ with 2, then $t(v_3(x)) = 7$ and $t(v_{y_2}) = 6$.
 - 2. If $d_H(v_{y_2}) = 2$ and $t(v_{x_2}) = 6$, recolor $v_{y_2}v_3(x)$ and v_{y_2} with 1, then $t(v_3(x)) = 6$ and $t(v_{y_2}) = 5$.
- 3. If $d_H(v_{y_2}) = 3$, recolor $v_{y_2}v_{x_2}$, $v_{y_2}v_{3}(x)$ and v_{y_2} with 1 and recolor v_{x_2} with 2, then $t(v_{y_2}) = 4$, $t(v_3(x)) = 6$ and the weight of v_{x_2} does not change.

Note that Theorem 5 implies that 1-2 Conjecture is valid for 3-regular graphs. By Lemma 3, if G is 4-regular, then there exists a bipartite graph H such that $2 \le d_H(u) \le 4$ for all $u \in V(G)$. Let $v \in V(G - E(H))$. Then $0 \le d_{G-E(H)}(v) \le 2$. Suppose that X and Y are the two partitions of V(H), then the vertex-induced subgraph of X (or Y) consists of cycles or isolated vertices or isolated edges or paths. Therefore, it may offer an idea to solve 1-2-3 Conjecture and 1-2 Conjecture for 4-regular graphs.

5 References

- [1] L. Addario-Berry, K. Dalal, C. McDiarmid, B.A. Reed, and A. Thomason, Vertex-colouring edge-weightings[J]. Combinatorica, 2007, 27: 1-12.
- [2] L. Addario-Berry, K. Dalal, and B.A. Reed, Degree constrained subgraphs[J]. Discrete Appl. Math. 2008, 156: 1168-1174.
- [3] J.A. Bondy and U.S.R. Murty, Graph theory with applications. The MaCmillan Press ltd, London and Basingstoke, New York, 1976.
- [4] G.J. Chang, C.H. Lu, J.J Wu, Q.L. Yu. Vertex-coloring Edge-weightings of Graphs[J]. Taiwan. J. Math., 2011, 15(4): 1807-1813.
- [5] M. Kalkowski, A note on 1,2-Conjecture, to appear in Elect. Journ. of Comb.
- [6] M. Kalkowski, M. Karoński, F. Pfender, Vertex-coloring edge weightings: Towards the 1-2-3-conjecture[J]. J. Combin. Theory Ser. B, 2010, 100: 347-349.
- [7] M. Karoński, T. Łuczak, A. Thomason, Edge weights and vertex colours[J]. J. Combin. Theory ser. B, 2004, 91: 151-157
- [8] J. Przybylo, M. Woźniak, On a 1,2 Conjecture[J]. Discrete Math. Theor. Comput. Sci., 2010, 12(1): 101-108.
- [9] J. Przybylo, The 1-2-3 Conjecture almost holds for regular graphs[J]. J. Combin. Theory ser. B, 2020 https://doi.org/10.1016/j.jctb.2020.03.005.
- [10] T. Wang and Q. Yu, On vertex-coloring 13-edge-weighting[J]. Front. Math. China, 2008, 3: 581-587.