Dressing for generalised linear Hamiltonian systems depending rationally on the spectral parameter and some applications

Alexander Sakhnovich

Faculty of Mathematics, University of Vienna, Oskar-Morgenstern-Platz 1, A-1090 Vienna, Austria, e-mail: oleksandr.sakhnovych@univie.ac.at

Abstract

We construct so called Darboux matrices and fundamental solutions in the important case of the generalised Hamiltonian (or canonical) systems depending rationally on the spectral parameter. A wide class of explicit solutions is obtained in this way. Interesting results for dynamical systems depending on several variables and their explicit solutions follow. For these purposes we use our version of Bäcklund-Darboux transformation and square roots of the corresponding generalised matrix eigenvalues. Some new auxiliary results on the roots of matrices are included as well. An appendix is added to make the paper self-sufficient.

MSC(2020): 34B30, 34C14, 37C79, 37J06, 15A15

Keywords: Generalised Hamiltonian system, dynamical system, fundamental solution, conservation law, Darboux matrix, root of the matrix, discrete Dirac system.

1 Introduction

1. In this paper, we construct fundamental solutions and so called Darboux

matrices for generalised Hamiltonian (canonical) systems depending rationally on the spectral parameter:

$$y'(x,z) = i \sum_{k=1}^{r} (z - c_k)^{-1} j \beta_k(x)^* \beta_k(x) y(x,z) \quad (y(x,z) \in \mathbb{C}^m);$$
 (1.1)

$$j = \begin{bmatrix} I_{m_1} & 0 \\ 0 & -I_{m_2} \end{bmatrix}, \ \beta_k(x) \in \mathbb{C}^{p \times m} \quad (p \le m), \ c_k \in \mathbb{R}; \ m_1, m_2 \in \mathbb{N}_0, \quad (1.2)$$

$$\mathbb{N}_0 = \mathbb{N} \cup \{0\}, \quad m_1 + m_2 = m > 0. \tag{1.3}$$

Here, i is the imaginary unit (i² = -1), the $m \times m$ Hamiltonians $H_k(x) = \beta_k(x)^*\beta_k(x)$ are locally summable, $c_i \neq c_k$ for $i \neq k$, \mathbb{R} stands for the real axis, \mathbb{C} stands for the complex plane, \mathbb{N} stands for the set of positive integers, I_n denotes the $n \times n$ identity matrix, the class of $p \times m$ matrices with complex-valued entries is denoted by $\mathbb{C}^{p \times m}$, and $\mathbb{C}^p = \mathbb{C}^{p \times 1}$. We note that $j = I_m$ if $m_2 = 0$ and $j = -I_m$ if $m_1 = 0$.

The obtained results on systems (1.1) are used in order to construct solutions of the dynamical systems

$$\frac{\partial \psi}{\partial x} = i \sum_{k=1}^{r} j H_k(x) \frac{\partial \psi}{\partial \zeta_k}.$$
 (1.4)

When r=1 and $H_1(x)$ is invertible, system (1.4) is equivalent to the important dynamical canonical system studied, for instance, in [32,43]. Thus, this paper is a substantial development and generalisation of the work [43]. The so called transformed Hamiltonians \widetilde{H}_k treated here preserve the property which the Hamiltonian in [43] possessed. Namely, the matrices $j\widetilde{H}_k(x)$ are linear similar to $jH_k(x)$. (This property is essential for the research on the C_0 -semigroups related to dynamical canonical systems [32,43].)

Symmetries and conservation laws are important in the study of the dynamical systems (see, e.g., [5, 6, 10, 31] and references therein). The expression for energy from [43, Proposition 3.1] is not valid for system (1.4) but an interesting conservation law is presented in Proposition 4.4.

2. In order to construct fundamental solutions and Darboux matrices for systems (1.1), we use our GBDT (generalised Bäcklund-Darboux transformation) approach and square roots of generalised matrix eigenvalues. The

case r = 2, p = 1 and $j = I_2$ was studied earlier in [37]. Here, we greatly generalise the corresponding results from [37].

Dressing procedure (also called Bäcklund-Darboux transformation or commutation method) is a fruitful tool in the study of linear and nonlinear differential equations (see [3, 13, 23, 24, 26, 29, 35, 36, 39, 54] and numerous references therein). For the recent applications and references see, for instance, [7, 9, 21, 28, 43, 46]. Dressing procedure may be considered as an important part of the general symmetry theory [40, 41]. GBDT version of Bäcklund-Darboux transformation (with generalised matrix eigenvalues) was actively developed starting from our work [42] (see further references in [3, 17, 33, 34, 44–46, 48]).

The notion of Darboux matrix (matrix function) may be explained in the following way. Given some initial linear system y'(x,z) = G(x,z)y(x,z) ($y' := \frac{d}{dx}y$) and a transformed (via Bäcklund-Darboux transformation) system $\widetilde{y}'(x,z) = \widetilde{G}(x,z)\widetilde{y}(x,z)$, the corresponding Darboux matrix is a nondegenerate matrix function W(x,z) satisfying equation

$$W'(x,z) = \widetilde{G}(x,z)W(x,z) - W(x,z)G(x,z). \tag{1.5}$$

It is easy to see that the product $\widetilde{w}(x,z) = W(x,z)w(x,z)$, where w(x,z) is a fundamental solution of the initial system y'(x,z) = G(x,z)y(x,z), is a fundamental solution of the transformed system $\widetilde{y}'(x,z) = \widetilde{G}(x,z)\widetilde{y}(x,z)$. If G(x,z) has some simple form so that w(x,z) and Darboux matrix W(x,z) are known explicitly, then the fundamental solution $\widetilde{w}(x,z)$ of the transformed system is also known explicitly.

3. In Section 3, we construct GBDT for system (1.1), (1.2). If r=1 and $m_1=m_2$, system (1.1) is equivalent to the classical canonical system (see various references and results on canonical systems in [11, 27, 48, 50]). L. de Branges and M.G. Krein obtained some fundamental results for the case of canonical systems where m=2 (see, e.g., [11, 27]). The spectral theory of canonical systems, where m>2, is no less important but essentially more complicated (see, e.g., [1, 12, 45, 48, 50, 52] and references therein). In Section 3 we deal with the case of an arbitrary $m \in \mathbb{N}$.

In the classical works and classical problems, there is a linear dependence on the spectral parameter (see also some recent papers, for instance,

[1, 25, 55]). The case of the more complicated (including nonlinear) dependence on the spectral parameter z is of essential interest and was studied, for instance, in [16, 19, 30, 37, 38, 56]. Various other examples include auxiliary linear systems for integrable nonlinear equations. (See, e.g., [37] on the connection of the system (1.1), where r = m = 2 and p = 1, with the sine-Gordon equation.)

4. Further in Section 3, we study important cases, where Darboux matrices and fundamental solutions may be constructed explicitly. The recovery of the generalised eigenfunction $\Pi(x)$, that is, solving of the matrix differential equation (3.5) is the main part of the explicit construction of the Darboux matrices and fundamental solutions. We express this generalised eigenfunction in terms of the square root of the matrix eigenvalue (see Proposition 3.5). In this way, a much wider class of generalised eigenfunctions and explicit solutions (than in [43]) is constructed even for r = 1.

Matrix roots (especially, square roots of matrices) are actively studied and find various applications during more than one and half centuries (see the paper [2] from the year 1858). In particular, structured roots of structured matrices and constructive procedures to obtain these roots are of interest. Among important works on this topic one could mention [8,20,53]. See also interesting recent works [4,22] and references therein. In the next section "Preliminaries", we recall the construction of the matrix roots. Moreover, we modify this construction so that the roots possess certain commutation properties which are required in several GBDT cases [44–46].

5. In Section 4, we apply the results of Section 3 in order to construct solutions of the dynamical systems (1.4) and study some of their properties.

Appendix A is dedicated to the general GBDT results from [48] (see Theorem A.1) and to Corollary A.2 (for symmetric S-nodes), which we use in Section 3.

Appendix B deals with another important application of matrix roots, that is, an application to discrete Dirac systems and Verblunsky-type coefficients (see [47] and references therein).

Notations. Many notations were explained above. The notation $\sigma(A)$ stands for the spectrum of the matrix A, A^* stands for the conjugate transpose of A, and the matrix inequality S > 0 ($S \ge 0$) means that the eigenval-

ues of the matrix $S = S^*$ are positive (nonnegative). In a similar way, one interprets the inequalities S < 0 and $S \le 0$. The notation diag $\{A_1, A_2, \ldots, A_s\}$ means diagonal matrix with the entries (or blocks) A_1, A_2, \ldots on the main diagonal.

2 Preliminaries

1. An ℓ -th root Q ($\ell > 1$, $\ell \in \mathbb{N}$) of some invertible $n \times n$ matrix A is of interest in many applications. (Such roots are constructed, e.g., in §6 of [20, Chapter VIII]). If one takes a root of f(A) instead of A, it is natural to assume that A and Q still commute (see, e.g., [46, Proposition 3.1] where such commutation is required). In order to achieve the equality AQ = QA in this case, the construction in [20, Chapter VIII, §6] should be slightly modified. Namely, one should consider the matrices of the form

$$F = \mu I_p + T \quad (\mu \neq 0, \quad T \in \mathcal{NT}), \tag{2.1}$$

where \mathcal{NT} is the class of upper triangular nilpotent Toeplitz matrices, while Jordan cells are dealt with in [20]. Clearly, $p \times p$ matrix $T \in \mathcal{NT}$ admits representation

$$T = \sum_{i=1}^{p-1} t_i \mathcal{S}_i, \tag{2.2}$$

where S_i are shift matrices:

$$S_i := \{\delta_{k-r+i}\}_{k,r=1}^p \quad (i > 0), \quad S_i S_k = S_{i+k}, \tag{2.3}$$

 δ_s is Kronecker delta, and $S_i = 0$ for $i \geq p$. It is known (and follows immediately from $S_i S_k = S_{i+k}$) that the matrices from \mathcal{NT} commute.

Considering the branch of $h(\lambda) = (\lambda + \mu)^{1/\ell}$ (for small λ) near one of the roots $\mu^{1/\ell}$ and taking the partial sum of the first p terms of the Taylor expansion of $(\lambda + \mu)^{1/\ell}$ at $\lambda = 0$, we set

$$g(\lambda) = \mu^{1/\ell} + \frac{1}{\ell} \mu^{\left(\frac{1}{\ell} - 1\right)} \lambda + \frac{1}{\ell} \left(\frac{1}{\ell} - 1\right) \mu^{\left(\frac{1}{\ell} - 2\right)} \frac{\lambda^2}{2} + \dots + \frac{1}{\ell} \left(\frac{1}{\ell} - 1\right) \dots \left(\frac{1}{\ell} - p + 2\right) \mu^{\left(\frac{1}{\ell} - p + 1\right)} \frac{\lambda^{p-1}}{(p-1)!}.$$
 (2.4)

Then, we have $g(\lambda)^{\ell} = \mu + \lambda + O(\lambda^p)$ for $\lambda \to 0$. Thus, $g(\lambda)^{\ell}$ and $\mu + \lambda$ coincide on the spectrum of T (in the terminology of [20, Chapter V, §1]), and so

$$g(T)^{\ell} = \mu I_p + T = F. \tag{2.5}$$

Let the minimal polynomial of a matrix A have the form $\prod_{i=1}^{s} (\lambda - \lambda_i)^{p_i}$. Recall that $g(\lambda)$ is determined on the spectrum of this matrix A (i.e., g(A) exists) if $g(\lambda_i)$ and its derivatives up to $g^{(p_i)}(\lambda_i)$ exist in the points λ_i $(1 \le i \le s)$ of the spectrum of A.

2. Given some $n \times n$ matrix A (det $A \neq 0$), we consider its representation

$$A = uA_{\mathcal{J}}u^{-1}, \quad A_{\mathcal{J}} = \operatorname{diag}\{\mathcal{A}_1, \mathcal{A}_2, \dots, \mathcal{A}_s\}, \quad \mathcal{A}_i = \mu_i I + \mathcal{S}_1,$$
 (2.6)

where $A_{\mathcal{J}}$ is a Jordan normal form of A, \mathcal{A}_i are Jordan cells (of order p_i), I and \mathcal{S}_1 have the same order p_i as \mathcal{A}_i , and I is the identity matrix. (We omit sometimes the orders of I and \mathcal{S}_k in the notations.) We note that $g(\lambda)$ depends, in fact, on two additional parameters: $g(\lambda) = g(\lambda, \mu, k)$ ($0 \le k < \ell$). Here, k determines the chosen root $\mu^{1/\ell} = e^{2\pi i k/\ell} \xi_0$ and ξ_0 is some arbitrarily fixed root.

The calculation of square roots Q such that AQ = QA and either $Q^2 = (A - cI_n)^2 - |a|^2 I_n$ or $Q^2 = (A - cI_n)^2 + |a|^2 I_n$ was required for the construction of Darboux matrices and explicit solutions of Dirac systems in [46, Proposition 3.1]. Similar problems appear in [45, Proposition 4.1].

Proposition 2.1 Let Jordan normal form of the $n \times n$ matrix A (det $A \neq 0$) be given by (2.6) and let the function f be defined on the spectrum of A.

Then, the matrices Q of the form

$$Q = u \operatorname{diag}\{\mathcal{B}_1, \mathcal{B}_2, \dots, \mathcal{B}_s\} u^{-1}, \quad \mathcal{B}_i = g(f(\mathcal{A}_i) - f(\mu_i)I, f(\mu_i), k_i), \quad (2.7)$$

where k_i may take fixed arbitrary values between 0 and $\ell-1$, satisfy relations

$$Q^{\ell} = f(A), \quad AQ = QA. \tag{2.8}$$

Proof. It is easy to see that

$$f(A) = u \operatorname{diag}\{f(A_1), f(A_2), \dots, f(A_s)\} u^{-1}.$$
 (2.9)

Moreover, in view of the last equalities in (2.3) and (2.6) we have

$$f(\mathcal{A}_i) - f(\mu_i)I \in \mathcal{NT}.$$

Hence, formulas (2.5), (2.7) and (2.9) yield the first equality in (2.8).

Finally, since $\mathcal{B}_i - (f(\mu_i))^{1/\ell} I \in \mathcal{NT}$, the last relation in (2.6) implies that $\mathcal{A}_i \mathcal{B}_i = \mathcal{B}_i \mathcal{A}_i$ and the second equality in (2.8) follows.

Remark 2.2 Clearly, if u and $A_{\mathcal{J}}$ are given, the roots Q of f(A) are constructed in Proposition 2.1 explicitly.

Remark 2.3 It follows from $\mathcal{B}_i - (f(\mu_i))^{1/\ell}I \in \mathcal{NT}$ that different roots of f(A) constructed in Proposition 2.1 commute.

Remark 2.4 It is easy to see that the equality AQ = QA does not necessarily hold for all the roots of $Q^{\ell} = f(A)$ (and so not necessarily all the roots are constructed in in Proposition 2.1). For instance, setting n = 3, $A = I_3 + S_1$, $f(\lambda) = (\lambda - 1)^2 + 4$, and $\ell = 2$, we can define u and one of the roots Q as follows:

$$u = u^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}, \quad Q = \begin{bmatrix} 2 & 0 & 1/4 \\ 0 & -2 & 0 \\ 0 & 0 & 2 \end{bmatrix} \quad (Q^2 = (A - I_3)^2 + 4I_3).$$
(2.10)

One can easily check that $S_1Q \neq QS_1$ and so $AQ \neq QA$.

3. Another (related to AQ = QA) commutation property is required for the explicit construction of solutions of sigma models and gravitational equations in [44]. Namely (after a change of notations for the ones used in this paper), it is required in [44, (B.1) and (B.2)] that

$$Q(z)^{\ell} = A - zI_n, \quad Q(z_1)Q(z_2) = Q(z_2)Q(z_1) \quad (z_1, z_2 \in \mathbb{R}),$$
 (2.11)

where $z, z_1, z_2 \notin \sigma(A)$ and $\ell = 2$.

In order to solve (2.11), we fix u (independently of z) in the procedures to find the roots Q(z) of $A-zI_n$ in Proposition 2.1. Then, we find the blocks

 $\mathcal{B}_i(z_1)$ setting $f(\lambda) = \lambda - z_1$ and find the blocks $\mathcal{B}_i(z_2)$ setting $f(\lambda) = \lambda - z_2$. The obtained blocks are upper triangular Toeplitz matrices. Thus, the second (commutation) equality in (2.11) holds. In this way, the following corollary of Proposition 2.1 is obtained.

Corollary 2.5 Let $\ell > 1$, $\ell \in \mathbb{N}$ and assume that $z, z_1, z_2 \notin \sigma(A)$. Then, the roots Q(z) satisfying (2.11) are constructed using procedures from Proposition 2.1.

3 GBDT for generalised Hamiltonian systems

1. Recall that we consider generalised Hamiltonian systems of the form

$$y'(x,z) = G(x,z)y(x,z), \quad G(x,z) = i\sum_{k=1}^{r} (z-c_k)^{-1}j\beta_k(x)^*\beta_k(x), \quad (3.1)$$

where (1.2) holds. Systems (3.1) are considered on some interval \mathcal{I} (0 $\in \mathcal{I}$).

Given initial system of the form (3.1), we construct GBDT-transformed system, its fundamental solution and the corresponding Darboux matrix using general results on GBDT from Appendix A. For this purpose, we set J = j in (A.14) (and further in Appendix A) and fix some triple of matrices $\{A, S(0), \Pi(0)\}$ such that

$$A, S(0) \in \mathbb{C}^{n \times n}, \quad \Pi(0) \in \mathbb{C}^{n \times m}, \quad n \in \mathbb{N}, \quad c_k \notin \sigma(A) \ (1 \le k \le r), \quad (3.2)$$

$$AS(0) - S(0)A^* = i\Pi(0)j\Pi(0)^*, \quad S(0) = S(0)^*.$$
(3.3)

We note that some notations here are different from Appendix A. In particular, the summation in (3.1) (from k = 1 till r) is substituted by the summation from s = 1 till ℓ in (A.1). Lower indices below are enumerated sometimes with commas in order to separate one index from another. Clearly, the equalities on $q_{1,k}$ in (A.16) hold for

$$q_{1,k} = -ijH_k(x), \quad H_k(x) := \beta_k(x)^*\beta_k(x).$$
 (3.4)

According to (3.1)–(3.4), the conditions of Theorem A.1 and Corollary A.2 for GBDT determined by the triple $\{A, S(0), \Pi(0)\}$ are fulfilled.

Taking into account the form of G(x, z) in (3.1), equalities (A.15) and the first equality in (A.17), we rewrite formulas (A.3), (A.19), (A.6), and (A.12) (in this order) in the form

$$\Pi'(x) = -i \sum_{k=1}^{r} (A - c_k I_n)^{-1} \Pi(x) j H_k(x),$$
(3.5)

$$S'(x) = -\sum_{k=1}^{r} (A - c_k I_n)^{-1} \Pi(x) j H_k(x) j \Pi(x)^* (A^* - c_k I_n)^{-1}, \qquad (3.6)$$

$$AS(x) - S(x)A^* = i\Pi(x)j\Pi(x)^*, \tag{3.7}$$

$$w_A(x,z) = I_m - ij\Pi(x)^* S(x)^{-1} (A - zI_n)^{-1} \Pi(x).$$
(3.8)

The matrix functions $\Pi(x)$ and S(x) are uniquely determined by the triple $\{A, S(0), \Pi(0)\}$ and systems (3.5), (3.6).

Remark 3.1 We will express the GBDT-transformed system via the transfer function w_A , which, in turn, is expressed via $\Pi(x)$ and S(x). Moreover, when $\sigma(A) \cap \sigma(A^*) = \emptyset$, S(x) is uniquely recovered from (3.7). The Darboux matrix and fundamental solution of the GBDT-transformed system are expressed via w_A as well (see Appendix A and the text below in this section). Summing up, we see that explicit construction of $\Pi(x)$ is one of the main components of this theory (and construction of S(x) and $w_A(x,z)$ easily follows). Matrix roots are essential for the construction of $\Pi(x)$ as demonstrated in Proposition 3.5.

Using (3.4) and (A.9), we see that the transformed system (for the initial system (3.1)) has the form

$$\widetilde{y}'(x,z) = i \sum_{k=1}^{r} (z - c_k)^{-1} j \widetilde{H}_k(x) \widetilde{y}(x,z), \tag{3.9}$$

$$\widetilde{H}_{k} = H_{k} + H_{k}Y_{k,-1} - jX_{k,-1}jH_{k} - jX_{k,-1}jH_{k}Y_{k,-1}
= (I_{m} - jX_{k,-1}j)H_{k}(I_{m} + Y_{k,-1}).$$
(3.10)

Relations (A.11) and (A.12) (together with the formula [48, (1.76)] show that

$$I_m - X_{k,-1}(x) = w_A(x, c_k), \quad (I_m - X_{k,-1}(x))(I_m + Y_{k,-1}(x)) = I_m.$$
 (3.11)

On the other hand, formula [48, (1.87)] for symmetric S-nodes yields

$$w_A(x, c_k)^{-1} = jw_A(x, c_k)^*j.$$
 (3.12)

Equalities (3.4) and (3.10)–(3.12) imply that

$$\widetilde{H}_k(x) = \widetilde{\beta}_k(x)^* \widetilde{\beta}_k(x), \quad \widetilde{\beta}_k(x) := \beta_k(x) j w_A(x, c_k)^* j.$$
 (3.13)

The matrices $\widetilde{H}_k(x)$ are nonnegative and may be considered as the transformed generalised Hamiltonians. Our next statement follows from Theorem A.1.

Theorem 3.2 Let the initial generalised Hamiltonian system (3.1) and the triple $\{A, S(0), \Pi(0)\}$ be given, where (1.2), (3.2) and (3.3) hold, and $c_k \neq c_l$ for $k \neq l$. Then, the GBDT-transformed generalised Hamiltonian system is given by the formulas (3.9) and (3.13), and $w_A(x, z)$ determined by (3.5), (3.6) and (3.8) is the corresponding Darboux matrix. If S(x) is invertible on the interval \mathcal{I} where our systems are considered, and w(x, z) is a fundamental solution of the initial system, then the fundamental solution $\widetilde{w}(x, z)$ of the transformed system has the form:

$$\widetilde{w}(x,z) = w_A(x,z)w(x,z). \tag{3.14}$$

Remark 3.3 According to (3.6), $S'(x) \leq 0$. Thus, S(x) < 0 in the case

$$S(0) < 0, \quad \mathcal{I} = [0, a] \quad (a > 0),$$

and S(x) > 0 in the case

$$S(0) > 0$$
, $\mathcal{I} = [-a, 0]$ $(a > 0)$.

In both cases, S(x) is invertible.

2. Let us consider some examples, where the conditions of Theorem 3.2 hold.

Example 3.4 Consider a simple example of the trivial initial Hamiltonians, that is, the case of $H_k(x) \equiv I_m$ ($1 \le k \le r$). Assuming for simplicity that $m_1 > 0$ and $m_2 > 0$, the case may be treated similar (in many details) to the example with r = 1 in [43].

We partition $\Pi(0)$ into the $n \times m_1$ and $n \times m_2$ blocks ϑ_1 and ϑ_2 , respectively: $\Pi(0) = \begin{bmatrix} \vartheta_1 & \vartheta_2 \end{bmatrix}$. Hence, it follows from (3.5) that

$$\Pi(x) = \left[\exp\left\{ -ix \sum_{k=1}^{r} (A - c_k I_n)^{-1} \right\} \vartheta_1 \quad \exp\left\{ ix \sum_{k=1}^{r} (A - c_k I_n)^{-1} \right\} \vartheta_2 \right]. \tag{3.15}$$

If $\sigma(A) \cap \sigma(A^*) = \emptyset$, there are unique matrices C_i (i = 1, 2) such that

$$AC_i - C_i A^* = i\vartheta_i \vartheta_i^*. \tag{3.16}$$

Moreover, in view of (3.15), (3.16) and equality $\sigma(A) \cap \sigma(A^*) = \emptyset$, there is a unique solution S(x) of (3.7) which is given by

$$S(x) = \exp\left\{-ix\sum_{k=1}^{r} (A - c_k I_n)^{-1}\right\} C_1 \exp\left\{ix\sum_{k=1}^{r} (A^* - c_k I_n)^{-1}\right\}$$

$$-\exp\left\{ix\sum_{k=1}^{r} (A - c_k I_n)^{-1}\right\} C_2 \exp\left\{-ix\sum_{k=1}^{r} (A^* - c_k I_n)^{-1}\right\}.$$
(3.17)

Here, equality (3.7) for S(x) of the form (3.17) is checked by direct substitution. In particular, we have $S(0) = C_1 - C_2$.

3. Next, we study a nontrivial example, where the matrix roots are involved. The following proposition is valid.

Proposition 3.5 Let the conditions of Theorem 3.2 hold. Assume that r = 2, $m_1 = m_2 = p > 0$, $\beta_k(x)$ are constant matrices β_k (k = 1, 2) and

$$\beta_k j \beta_k^* = 0 \quad (k = 1, 2), \quad \beta_1 j \beta_2^* = I_p.$$
 (3.18)

Then, the solution $\Pi(x)$ of (3.5) is given explicitly by the formulas

$$\Pi(x)j \begin{bmatrix} \beta_1^* & \beta_2^* \end{bmatrix} = \begin{bmatrix} \Phi_1(x) & \Phi_2(x) \end{bmatrix}; \tag{3.19}$$

$$\Phi_1(x) = e^{ixQ} h_1 + e^{-ixQ} h_2, \quad h_1, h_2, \Phi_1(x) \in \mathbb{C}^{n \times p},$$
(3.20)

$$\Phi_2(x) = -(A - c_2 I_n) Q(e^{ixQ} h_1 - e^{-ixQ} h_2), \quad \Phi_2(x) \in \mathbb{C}^{n \times p};$$
(3.21)

$$h_1 + h_2 = \Pi(0)j\beta_1^*, \quad (A - c_2 I_n)Q(h_2 - h_1) = \Pi(0)j\beta_2^*;$$
 (3.22)

$$Q^{2} = (A - c_{1}I_{n})^{-1}(A - c_{2}I_{n})^{-1}.$$
(3.23)

Proof. First note that

$$\begin{bmatrix} \beta_1 \\ \beta_2 \end{bmatrix} j \begin{bmatrix} \beta_1^* & \beta_2^* \end{bmatrix} = \begin{bmatrix} 0 & I_p \\ I_p & 0 \end{bmatrix}.$$

Thus, $\begin{bmatrix} \beta_1^* & \beta_2^* \end{bmatrix}$ is invertible (and $\begin{bmatrix} \beta_1^* & \beta_2^* \end{bmatrix}^{-1} = \begin{bmatrix} \beta_2 \\ \beta_1 \end{bmatrix} j$). Hence, in view of (3.18) and (3.19), formula (3.5) is equivalent to the equalities

$$\Phi_1'(x) = -i(A - c_2 I_n)^{-1} \Phi_2(x), \quad \Phi_2'(x) = -i(A - c_1 I_n)^{-1} \Phi_1(x).$$
 (3.24)

The first equality in (3.24) is immediate from (3.20) and (3.21). The second equality in (3.24) follows from (3.20), (3.21), and (3.23). Now, we see that

$$\Pi(x) = \begin{bmatrix} \Phi_1(x) & \Phi_2(x) \end{bmatrix} \begin{bmatrix} \beta_2 \\ \beta_1 \end{bmatrix}$$

satisfies (3.5).

Relations (3.22) follow from (3.19) and from the equalities (3.20) and (3.21) at x = 0. If $\Pi(0)$ is given, h_1 and h_2 are uniquely recovered from (3.22). (Vice versa, starting from h_1 and h_2 we easily obtain $\Pi(0)$.)

Remark 3.6 We note that commutation properties are not required from the square root Q in (3.23). The requirement AQ = QA is convenient but not necessary in [45, 46]. However, in using GBDT for the construction of explicit solutions of integrable nonlinear PDEs the commutation requirements are often unavoidable (see, e.g., paragraph 3 of Section 2).

4 The case of several variables

1. Let the conditions of Theorem 3.2 be fulfilled. That is, let the corresponding set of matrix functions $\beta_k(x)$ and the triple $\{A, S(0), \Pi(0)\}$ be given. It follows from Corollary A.2 that, for our case of symmetric S-nodes and system (3.1), formula [48, (7.61)] takes the form

$$\left(j\Pi(x)^*S(x)^{-1}\right)' = i\sum_{k=1}^r j\widetilde{H}_k(x)j\Pi(x)^*S(x)^{-1}(A - c_kI_n)^{-1}.$$
 (4.1)

Relation (4.1) yields our next theorem.

Theorem 4.1 Let a set of numbers c_k and matrices $\beta_k(x)$ $(1 \le k \le r)$ as well as a triple $\{A, S(0), \Pi(0)\}$ be given. Assume that relations (1.2), (3.2) and (3.3) hold and that $\Pi(x)$ and S(x) satisfy (3.5) and (3.6), respectively.

Then, in the points of invertibility of S(x), the matrix function

$$\widetilde{\psi}(x,\zeta_1,\ldots,\zeta_r) = j\Pi(x)^*S(x)^{-1}\exp\left\{\sum_{k=1}^r \zeta_k(A-c_kI_n)^{-1}\right\}$$
 (4.2)

satisfies the system

$$\frac{\partial \widetilde{\psi}}{\partial x} = i \sum_{k=1}^{r} j \widetilde{H}_{k}(x) \frac{\partial \widetilde{\psi}}{\partial \zeta_{k}}, \tag{4.3}$$

where \widetilde{H}_k is given by (3.13).

Remark 4.2 Note that $\widetilde{\psi}$ is an $m \times n$ matrix function. Thus, (4.3) means that each column of $\widetilde{\psi}$ satisfies our transformed dynamical system (of several variables).

Remark 4.3 Under conditions of Theorem 4.1, equalities (3.12) and (3.13) (where w_A is expressed via Π and S in (3.8)) are fulfilled. Hence, one may rewrite (3.13) in the form:

$$j\widetilde{H}_k(x) = w_A(x, c_k)jH_k(x)w_A(x, c_k)^{-1},$$
 (4.4)

that is, $j\widetilde{H}_k(x)$ is linear similar to $jH_k(x)$.

2. The following conservation law is valid for the GBDT-transformations in this paper.

Proposition 4.4 Let the conditions of Theorem 4.1 hold. Then,

$$(\Pi(x)^* S(x)^{-1} \Pi(x))' = \sum_{k=1}^r (\widetilde{H}_k(x) - H_k(x)).$$
(4.5)

Proof. Relations (3.8) and (4.1) yield

$$\left(\Pi(x)^* S(x)^{-1}\right)' \Pi(x) = \sum_{k=1}^r \widetilde{H}_k(x) \left(I_m - w_A(x, c_k)\right). \tag{4.6}$$

From (3.5) and (4.6), it follows that

$$(\Pi(x)^* S(x)^{-1} \Pi(x))' = \sum_{k=1}^r \widetilde{H}_k(x) (I_m - w_A(x, c_k))$$
$$+ j \sum_{k=1}^r (w_A(x, c_k) - I_m) j H_k(x). \tag{4.7}$$

Finally, the equalities (4.4) and (4.7) imply (4.5).

.

Remark 4.5 If $\Pi(x)$ is known explicitly (as in Example 3.4 or in Proposition 3.5), we obtain explicit expressions for $\widetilde{\psi}$ and system (4.3).

A GBDT - general case

1. Following [48, pp. 219–221] (see also the references therein), let us consider a general case of the first order system depending rationally on the spectral parameter z, that is, the case of multiple poles with respect to z:

$$y' = Gy$$
, $G(x, z) = -\left(\sum_{k=0}^{r} z^k q_k(x) + \sum_{s=1}^{\ell} \sum_{k=1}^{r_s} (z - c_s)^{-k} q_{sk}(x)\right)$. (A.1)

Similar to [48], we assume that $x \in \mathcal{I}$, where \mathcal{I} is an interval such that $0 \in \mathcal{I}$, and that the coefficients $q_k(x)$ and $q_{sk}(x)$ are $m \times m$ locally summable matrix functions. The GBDT of the initial system (A.1) is determined by a number $n \in \mathbb{N}$, by $n \times n$ matrices A_k (k = 1, 2) and S(0), and by $n \times m$ matrices $\Pi_k(0)$ (k = 1, 2). It is required that these matrices form an S-node, that is, the matrix identity

$$A_1S(0) - S(0)A_2 = \Pi_1(0)\Pi_2(0)^*$$
(A.2)

holds. The matrix functions $\Pi_k(x)$ are introduced via their values at x=0

and equations

$$(\Pi_1)' = \sum_{k=0}^r A_1^k \Pi_1 q_k + \sum_{s=1}^\ell \sum_{k=1}^{r_s} (A_1 - c_s I_n)^{-k} \Pi_1 q_{sk}, \tag{A.3}$$

$$\left(\Pi_2^*\right)' = -\left(\sum_{k=0}^r q_k \Pi_2^* A_2^k + \sum_{s=1}^\ell \sum_{k=1}^{r_s} q_{sk} \Pi_2^* (A_2 - c_s I_n)^{-k}\right). \tag{A.4}$$

Compare (A.1) with (A.4) to see that Π_2^* can be viewed as a generalized eigenfunction of the system u' = Gu.

Matrix function S(x) is introduced via S(0) and via S'(x) given by the equality

$$S' = \sum_{k=1}^{r} \sum_{j=1}^{k} A_1^{k-j} \Pi_1 q_k \Pi_2^* A_2^{j-1} - \sum_{s=1}^{\ell} \sum_{k=1}^{r_s} \sum_{j=1}^{k} (A_1 - c_s I_n)^{j-k-1} \times \Pi_1 q_{sk} \Pi_2^* (A_2 - c_s I_n)^{-j}.$$
(A.5)

Equations (A.2)–(A.5)) yield the identity

$$A_1S(x) - S(x)A_2 = \Pi_1(x)\Pi_2(x)^*, \quad x \in \mathcal{I}$$
 (A.6)

of which (A.2) is a particular case of x = 0. For initial system (A.1), the GBDT-transformed system is

$$\widetilde{y}' = \widetilde{G}\widetilde{y}, \quad \widetilde{G}(x,z) = -\Big(\sum_{k=0}^{r} z^{k} \widetilde{q}_{k}(x) + \sum_{s=1}^{\ell} \sum_{k=1}^{r_{s}} (z - c_{s})^{-k} \widetilde{q}_{sk}(x)\Big), \quad (A.7)$$

where the transformed coefficients \widetilde{q}_k and \widetilde{q}_{sk} are given by the formulas

$$\widetilde{q}_k = q_k - \sum_{j=k+1}^r \left(q_j Y_{j-k-1} - X_{j-k-1} q_j + \sum_{i=k+2}^j X_{j-i} q_j Y_{i-k-2} \right),$$
(A.8)

$$\widetilde{q}_{sk} = q_{sk} + \sum_{j=k}^{r_s} \left(q_{sj} Y_{s,k-j-1} - X_{s,k-j-1} q_{sj} - \sum_{j=k}^{j} X_{s,i-j-1} q_{sj} Y_{s,k-i-1} \right), \quad (A.9)$$

and the matrix functions $X_k(x)$, $Y_k(x)$ and $X_{sk}(x)$, $Y_{sk}(x)$ have the form

$$X_k = \Pi_2^* S^{-1} A_1^k \Pi_1, \quad Y_k = \Pi_2^* A_2^k S^{-1} \Pi_1, \tag{A.10}$$

$$X_{sk} = \Pi_2^* S^{-1} (A_1 - c_s I_n)^k \Pi_1, \quad Y_{sk} = \Pi_2^* (A_2 - c_s I_n)^k S^{-1} \Pi_1.$$
 (A.11)

Our theorem below shows that the transfer function

$$w_A(z) = I_m - \Pi_2^* S^{-1} (A_1 - z I_n)^{-1} \Pi_1, \tag{A.12}$$

(that is, the transfer matrix function in Lev Sakhnovich form [49,50]) is the Darboux matrix for the systems (A.1) and (A.7).

Theorem A.1 [48] Let the first order system (A.1) and five matrices S(0), A_1 , A_2 and $\Pi_1(0)$, $\Pi_2(0)$ be given. Assume that the identity (A.2) holds and that $\{c_s\} \cap \sigma(A_k) = \emptyset$ (k = 1, 2). Then, in the points of invertibility of S, the transfer matrix function $w_A(x, z)$ given by (A.12), where S(x) and $\Pi_k(x)$ are determined by (A.3)–(A.5), satisfies the equation

$$w'_A(x,z) = \tilde{G}(x,z)w_A(x,z) - w_A(x,z)G(x,z),$$
 (A.13)

where \widetilde{G} is determined by the formulas (A.7)-(A.11).

Compare system (A.13) with the system (1.5) in introduction.

We note that the transfer matrix function $w_A(z)$ given by (A.12) takes roots in M. Livšic's characteristic matrix function (see the discussion in [51]).

2. In this paper, we are interested in the case of the *symmetric S-nodes*, that is, in the case

$$A_2 = A_1^*, \quad S(0) = S(0)^*, \quad \Pi_2(0) = iJ\Pi_1(0)^*, \quad J = J^* = J^{-1}.$$
 (A.14)

In this case, we also change notations and set

$$A_1 = A_2^* = A, \quad \Pi_1(x) = \Pi(x).$$
 (A.15)

Moreover, we assume that

$$Jq_k^*J = -q_k, \quad Jq_{sk}^*J = -q_{sk}, \quad c_s \in \mathbb{R} \quad (1 \le s \le \ell, \quad 1 \le k \le r_s).$$
 (A.16)

The following corollary is valid under conditions above.

Corollary A.2 Let the conditions of Theorem A.1 and relations (A.14)–(A.16) hold. Then, we have

$$\Pi_2(x)^* = iJ\Pi(x)^*, \quad S(x) = S(x)^*.$$
 (A.17)

Proof. It is immediate from (A.3), from (A.14), and from $c_s \in \mathbb{R}$ that

$$(iJ\Pi_1^*)' = \sum_{k=0}^r Jq_k^* J(iJ\Pi_1^*) A_2^k + \sum_{s=1}^\ell \sum_{k=1}^{r_s} Jq_{sk}^* J(iJ\Pi_1^*) (A_2 - c_s I_n)^{-k}.$$
 (A.18)

Hence, in view of (A.4), (A.16) and the third equality in (A.14), $ij\Pi_1(x)^*$ and $\Pi_2(x)^*$ satisfy the same first order linear differential system and coincide at x = 0. Thus, we obtain the first equality in (A.17). Therefore, taking into account (A.15), we rewrite (A.5) in the form

$$S' = \sum_{k=1}^{r} \sum_{j=1}^{k} A^{k-j} \Pi(iq_k J) \Pi^*(A^*)^{j-1} - \sum_{s=1}^{\ell} \sum_{k=1}^{r_s} \sum_{j=1}^{k} (A - c_s I_n)^{j-k-1} \times \Pi(iq_{sk} J) \Pi^*(A^* - c_s I_n)^{-j}.$$
(A.19)

Changing the order of summation, that is, setting

$$k-j=p-1$$
, $Q_k = \Pi(iq_k J)\Pi^*$, $Q_{sk} = \Pi(iq_{sk} J)\Pi^*$,

we have

$$\sum_{j=1}^{k} A^{k-j} Q_k (A^*)^{j-1} = \sum_{p=1}^{k} A^{p-1} Q_k (A^*)^{k-p}$$

$$= \frac{1}{2} \sum_{j=1}^{k} \left(A^{k-j} Q_k (A^*)^{j-1} + A^{j-1} Q_k (A^*)^{k-j} \right), \quad (A.20)$$

and in, in a similar way,

$$\sum_{j=1}^{k} (A - c_s I_n)^{j-k-1} Q_{sk} (A^* - c_s I_n)^{-j}$$

$$= \frac{1}{2} \sum_{j=1}^{k} ((A - c_s I_n)^{j-k-1} Q_{sk} (A^* - c_s I_n)^{-j} + (A - c_s I_n)^{-j} Q_{sk} (A^* - c_s I_n)^{j-k-1}). \tag{A.21}$$

Using (A.16), it is easy to see that $Q_k = Q_k^*$, $Q_{sk} = Q_{sk}^*$, and so the right-hand sides of (A.20) and (A.21) are Hermitian. It follows that the right-hand side

of (A.19) is Hermitian as well, that is, S' is Hermitian. Since $S(0) = S(0)^*$ and S' is Hermitian, the second equality in (A.17) is also valid.

In the case of symmetric S-nodes, formula (A.14) implies that GBDT is determined by the triple $\{A, S(0), \Pi(0)\}$ of matrices.

B Matrix roots and discrete Dirac systems

Discrete Dirac systems

$$y_{k+1}(z) = (I_m - \frac{i}{z}jC_k)y_k(z), \quad C_k > 0, \quad C_k jC_k = j,$$
 (B.1)

present (in the case of $m_1 = m_2$) a fruitful approach to Toeplitz matrices and measure theory (see [47] and references therein), which completes in an interesting way the classical approach via Szegő recurrences. (Recall that j is introduced in (1.2).)

The positive roots of the matrices C_k play an essential role in the study of discrete Dirac systems. Using [18, Propositions 2.1, 2.4] (and completing the proof of [18, Proposition 2.1] with the fact that there is a unique positive root $D^{1/\ell} > 0$ of the matrix D > 0, see [20, (62), p. 232]), we have the following proposition.

Proposition B.1 Let some $m \times m$ matrix C > 0 satisfy condition CjC = j. Then, there is a unique positive root $C^{1/\ell}$ ($\ell \in \mathbb{N}$) and this root has the property $C^{1/\ell}jC^{1/\ell} = j$. The matrix C (and $C^{1/\ell}$ as well) admits so called Halmos extention representation:

$$C = \operatorname{diag} \left\{ \left(I_{m_1} - \rho \rho^* \right)^{-\frac{1}{2}}, \ \left(I_{m_2} - \rho^* \rho \right)^{-\frac{1}{2}} \right\} \left[\begin{array}{cc} I_{m_1} & \rho \\ \rho^* & I_{m_2} \end{array} \right], \quad \rho^* \rho < I_{m_2},$$

where ρ is an $m_1 \times m_2$ matrix.

The properties of Halmos extensions are well-known (see, e.g., [14,15]).

Acknowledgments This research was supported by the Austrian Science Fund (FWF) under Grant No. Y-963.

References

- K.R. Acharya, A. Ludu, Some theory and applications of 2N-dimensional canonical systems: periodic and nonperiodic, J. Phys. A 54 (2021) Paper No. 085202.
- [2] A. Cayley, A memoir on the theory of matrices, Philos. Trans. R. Soc. 148 (1858) 17–37.
- [3] J.L. Cieslinski, Algebraic construction of the Darboux matrix revisited, J. Phys. A 42 (2009) Paper No. 404003.
- [4] D. Choi, A. Greenbaum, Roots of matrices in the study of GMRES convergence and Crouzeix's conjecture, SIAM J. Matrix Anal. Appl. 36 (2015) 289–301.
- [5] R.M. Colombo, Well posedness and control in models based on conservation laws. In: Nonlinear conservation laws and applications (eds. A. Bressan et al.), pp. 267–277, Springer, 2011.
- [6] A. Constantin, A dynamical systems approach towards isolated vorticity regions for Tsunami background states, Arch. Ration. Mech. Anal. 200 (2011) 239–253.
- [7] A. Constantin, R. Ivanov, Dressing method for the Degasperis-Procesi equation, Stud. Appl. Math. 138 (2017) 205–226.
- [8] G.W. Cross, P. Lancaster, Square roots of complex matrices, Linear Multilinear Algebra 1 (1974) 289–293.
- [9] S. Cuenda, N.R. Quintero, A. Sanchez, Sine-Gordon wobbles through Bäcklund transformations, Discrete Contin. Dyn. Syst. Ser. S 4 (2011) 1047–1056.
- [10] C.M. Dafermos, Hyperbolic conservation laws in continuum physics. Fourth edition. Springer, Berlin, 2016.
- [11] L. de Branges, Hilbert spaces of entire functions, Prentice-Hall, Englewood Cliffs, N.J., 1968.

- [12] L. de Branges, The expansion theorem for Hilbert spaces of entire functions. In: Entire Functions and Related Parts of Analysis (Proc. Sympos. Pure Math., La Jolla, Calif., 1966), pp. 79–148, Amer. Math. Soc., Providence, RI, 1968.
- [13] P.A. Deift, Applications of a commutation formula, Duke Math. J. 45 (1978) 267–310.
- [14] V.K. Dubovoj, B. Fritzsche, B. Kirstein, Matricial version of the classical Schur problem, Teubner-Texte zur Mathematik 129, B.G. Teubner Verlagsgesellschaft, Stuttgart, 1992.
- [15] H. Dym, J contractive matrix functions, reproducing kernel Hilbert spaces and interpolation, [J] CBMS Reg. Conf. Ser. in Math. 71, American Mathematical Society, Providence, RI, 1989.
- [16] J. Elyseeva, R. Šimon Hilscher, Discrete oscillation theorems for symplectic eigenvalue problems with general boundary conditions depending nonlinearly on spectral parameter, Linear Algebra Appl. 558 (2018) 108–114.
- [17] B. Fritzsche, B. Kirstein, A.L. Sakhnovich, Completion problems and scattering problems for Dirac type differential equations with singularities, J. Math. Anal. Appl. 317 (2006) 510–525.
- [18] B. Fritzsche, B. Kirstein, I. Roitberg, A.L. Sakhnovich, Discrete Dirac system: rectangular Weyl, functions, direct and inverse problems, Oper. Matrices 8 (2014) 799–819.
- [19] C.T. Fulton, Singular eigenvalue problems with eigenvalue parameter contained in the boundary conditions, Proc. R. Soc. Edinb. A 87 (1980) 1–34.
- [20] F.R. Gantmacher, The Theory of Matrices, Chelsea, New York, 1959.
- [21] J.C. García-Ardila, F. Marcellán, P.H. Villamil-Hernández, Associated orthogonal polynomials of the first kind and Darboux transformations, J. Math. Anal. Appl. 508 (2022) Paper No. 125883.

- [22] E.S. Gawlik, Rational minimax iterations for computing the matrix pth root, Constr. Approx. 54 (2021) 1–34.
- [23] F. Gesztesy, A complete spectral characterization of the double commutation method, J. Funct. Anal. 117 (1993) 401–446.
- [24] F. Gesztesy, H. Holden, Soliton equations and their algebrogeometric solutions, Vol. I, Cambridge University Press, Cambridge, 2003
- [25] F. Gesztesy, A.L. Sakhnovich, The inverse approach to Dirac-type systems based on the A-function concept, J. Funct. Anal. 279 (2020) Paper No. 108609.
- [26] F. Gesztesy, G. Teschl, On the double commutation method, Proc. Amer. Math. Soc. 124 (1996) 1831–1840.
- [27] I. Gohberg, M.G. Krein, Theory and applications of Volterra operators in Hilbert space, Transl. of math. monographs 24, American Mathematical Society, Providence, RI, 1970.
- [28] F.A. Grünbaum, M.D. de la Iglesia, Stochastic Darboux transformations for quasi-birth-and-death processes and urn models, J. Math. Anal. Appl. 478 (2019) 634–654.
- [29] C.H. Gu, H. Hu, Z. Zhou, Darboux transformations in integrable systems. Theory and their applications to geometry, Mathematical Physics Studies 26, Springer, Dordrecht, 2005.
- [30] S. Hassi, M. Möller, H. de Snoo, Singular Sturm-Liouville problems whose coefficients depend rationally on the eigenvalue parameter, J. Math. Anal. Appl. 295 (2004) 258–275.
- [31] H. Holden, N.K. Risebro, Front tracking for hyperbolic conservation laws. Second edition. Springer, Heidelberg, 2015.
- [32] B. Jacob, K. Morris, H. Zwart, C_0 -semigroups for hyperbolic partial differential equations on a one-dimensional spatial domain, J. Evol. Equ., 15 (2015) 493–502.

- [33] M.A. Kaashoek, A.L. Sakhnovich, Discrete skew self-adjoint canonical system and the isotropic Heisenberg magnet model, J. Funct. Anal. 228 (2005) 207–233.
- [34] A. Kostenko, A. Sakhnovich, G. Teschl, Commutation methods for Schrödinger operators with strongly singular potentials, Math. Nachr. 285 (2012) 392–410.
- [35] V.A. Marchenko, Nonlinear equations and operator algebras, Mathematics and Its Applications (Soviet Series) 17, D. Reidel, Dordrecht, 1988.
- [36] V.B. Matveev, M.A. Salle, Darboux transformations and solitons, Springer, Berlin, 1991.
- [37] R. Mennicken, A.L. Sakhnovich, C. Tretter, Direct and inverse spectral problem for a system of differential equations depending rationally on the spectral parameter, Duke Math. J. 109 (2001) 413–449.
- [38] R. Mennicken, H. Schmid, A.A. Shkalikov, On the eigenvalue accumulation of Sturm–Liouville problems depending nonlinearly on the spectral parameter, Math. Nachr. 189 (1998) 157–170.
- [39] R. Miura (ed.), Bäcklund Transformations, Lecture Notes in Math. 515, Springer, Berlin, 1976.
- [40] P.J. Olver, Equivalence, invariants, and symmetry, Cambridge University Press, Cambridge, 1995.
- [41] R.O. Popovych, Point and contact equivalence groupoids of twodimensional quasilinear hyperbolic equations, Appl. Math. Lett. 116 (2021) Paper No. 107068.
- [42] A.L. Sakhnovich, Dressing procedure for solutions of nonlinear equations and the method of operator identities, Inverse problems 10 (1994) 699–710.
- [43] A.L. Sakhnovich, Dynamical canonical systems and their explicit solutions, Discrete Contin. Dyn. Syst. 37 (2017) 1679–1689.

- [44] A.L. Sakhnovich, Einstein, σ -model and Ernst-type equations and non-isospectral GBDT version of Darboux transformation, arXiv:2003.13024 (Adv. Theor. Math. Phys. to appear).
- [45] A.L. Sakhnovich, On a class of canonical systems corresponding to matrix string equations: general-type and explicit fundamental solutions and Weyl–Titchmarsh theory, Doc. Math. 26 (2021) 583–615.
- [46] A.L. Sakhnovich, On the classes of explicit solutions of Dirac, dynamical Dirac and Dirac-Weyl systems with non-vanishing at infinity potentials, their properties and applications, J. Differential Equations 275 (2021) 250–269.
- [47] A.L. Sakhnovich, Discrete self-adjoint Dirac systems: asymptotic relations, Weyl functions and Toeplitz matrices, Constr. Approx. 55 (2022) 641–659.
- [48] A.L. Sakhnovich, L.A. Sakhnovich, I.Ya. Roitberg, Inverse Problems and Nonlinear Evolution Equations. Solutions, Darboux Matrices and Weyl-Titchmarsh Functions, De Gruyter Studies in Mathematics 47, De Gruyter, Berlin, 2013.
- [49] L.A. Sakhnovich, On the factorization of the transfer matrix function, Sov. Math. Dokl. 17 (1976) 203–207.
- [50] L.A. Sakhnovich, Spectral theory of canonical differential systems, method of operator identities, Birkhäuser, Basel, 1999.
- [51] L.A. Sakhnovich, Characteristic function of M. Livšic and some developments, arXiv:2104.12694.
- [52] P. Šepitka, Riccati equations for linear Hamiltonian systems without controllability condition, Discrete Contin. Dyn. Syst. 39 (2019) 1685–1730.
- [53] J. H. M. Wedderburn, Lectures on Matrices, American Mathematical Society, New York, 1934.

- [54] V.E. Zakharov, A.V. Mikhailov, On the integrability of classical spinor models in two-dimensional space-time, Comm. Math. Phys. 74 (1980) 21–40.
- [55] P. Zemánek, Eigenfunctions expansion for discrete symplectic systems with general linear dependence on spectral parameter, J. Math. Anal. Appl. 499 (2021) Paper No. 125054.
- [56] Xin Zhou, Inverse scattering transform for systems with rational spectral dependence, J. Differential Equations 115 (1995) 277–303.