LOCAL MINIMIZERS FOR A CLASS OF FUNCTIONALS OVER THE NEHARI SET

HUMBERTO RAMOS QUOIRIN AND KAYE SILVA

ABSTRACT. We analyze the topological structure of the Nehari set for a class of functionals depending on a real parameter λ , and having two degrees of homogeneity. A special attention is paid to the extremal parameter λ^* , which is the threshold value for the Nehari set to be given by a *natural constraint*. The main difficulty arises when $\lambda > \lambda^*$, as the energy functional may be unbounded from below over the Nehari set. In such situation we prove the existence of local minimizers of the functional constrained to this set. We unify and extend previous existence and multiplicity results for critical points of indefinite, (p, q)-Laplacian, and Kirchhoff type problems.

CONTENTS

1. Introduction	1
2. Proofs	4
2.1. Basic properties	5
2.2. Minimization up to λ^*	6
2.3. Minimization at λ^*	10
2.4. Local Minimization Beyond λ^*	12
2.5. A mountain-pass critical point for $\lambda > \lambda^*$	15
3. Applications	17
3.1. Indefinite <i>p</i> -Laplace equations	17
3.2. (p,q) -Laplacian problems	20
3.3. Kirchhoff type problems	21
Acknowledgements	22
Data availability	22
References	22

1. INTRODUCTION

This article is devoted to the analysis of the Nehari set (usually known as Nehari manifold) associated to a functional depending on a real parameter. We shall proceed with the investigation on the extremal parameter carried out in [3, 20], where a class of indefinite and superlinear type problems has been investigated. We aim at unifying and extending the results of [3, 20] by dealing with a general class of functionals having two degrees of homogeneity. More precisely, we consider the family

$$\Phi_{\lambda} = \frac{1}{p} \left(P_1 - \lambda P_2 \right) - \frac{1}{\gamma} F, \tag{1.1}$$

²⁰¹⁰ Mathematics Subject Classification. Primary 35J50, 35J91, 35Q60,

Key words and phrases. Quasilinear pde, variational methods, Nehari manifold, indefinite problems.

Kaye Silva was partially supported by CNPq/Brazil under Grant 408604/2018-2.

where P_1, P_2, F are C^1 functionals acting on a uniformly convex Banach space X, λ is a real parameter, and $p, \gamma > 1$ with $p \neq \gamma$. Furthermore the following basic conditions shall be assumed:

- P_1, P_2 are *p*-homogeneous and *F* is γ -homogeneous, i.e. $P_1(tu) = t^p P_1(u), P_2(tu) = t^p P_2(u)$ and $F(tu) = t^{\gamma} F(u)$ for all t > 0 and $u \in X$.
- $P_2(u) \ge 0$ for all $u \in X \setminus \{0\}$ and F(0) = 0.
- There exists $C_1, C_2, C_3 > 0$ such that $P_1(u) \ge C_1 ||u||^p$, $P_2(u) \le C_2 ||u||^p$ and $F(u) \le C_3 ||u||^{\gamma}$ for all $u \in X$.
- There exists $u_1, u_2, u_3 \in X \setminus \{0\}$ such that $F(u_1) > 0 > F(u_2)$ and $F(u_3) = 0$.
- P_1 is weakly lower semi-continuous, P_2 is weakly continuous, and F is weakly upper semi-continuous.

This class of functionals appears as the energy functional in several elliptic problems, among which the main prototype is

$$\Phi_{\lambda}(u) = \frac{1}{p} \int_{\Omega} \left(|\nabla u|^p - \lambda |u|^p \right) - \frac{1}{\gamma} \int_{\Omega} f(x) |u|^{\gamma}, \qquad (1.2)$$

defined for $u \in X = W_0^{1,p}(\Omega)$. Here Ω is a bounded domain of \mathbb{R}^N , $N \ge 1$, p > 1 and $\gamma \neq p$ with $1 < \gamma < p^*$ (the critical Sobolev exponent), and $f \in L^{\infty}(\Omega)$. The Euler-Lagrange equation for this functional is

$$-\Delta_p u = \lambda |u|^{p-2} u + f(x) |u|^{\gamma-2} u, \quad u \in W_0^{1,p}(\Omega),$$
(1.3)

which has been studied by several authors [2, 6, 10, 11, 17, 19, 20, 24]. Inspired by the approach used in [20], where the functional (1.2) is considered in the superhomogeneous (or superlinear, if p = 2) case $\gamma > p$, we shall investigate the general class (1.1). In particular, we shall complement [20] by including the subhomogeneous case $\gamma < p$. We shall also consider some variations of (1.3), namely, the Neumann problem

$$\begin{cases} -\Delta_p u = \lambda |u|^{p-2} u + f(x)|u|^{\gamma-2} u & \text{in} \quad \Omega, \\ \partial_n u = 0 & \text{on} \quad \partial\Omega, \end{cases}$$

where $\partial_n u$ denotes the outer normal derivative of u, as well as problems with nonlinear boundary conditions.

Furthermore, our results also apply to the (p, q)-Laplacian problem

$$-\Delta_p u - \Delta_q u = \alpha |u|^{p-2} u + \beta |u|^{q-2} u, \quad u \in W_0^{1,p}(\Omega),$$

where 1 < q < p and $\alpha, \beta \in \mathbb{R}$, and to the Kirchhoff type equation

$$-\left(a+b\int_{\Omega}|\nabla u|^{2}dx\right)\Delta u=\lambda u+\mu|u|^{2}u,\quad u\in H_{0}^{1}(\Omega),$$

where a, b > 0, and $\lambda, \mu \in \mathbb{R}$.

Our main purpose is to analyze the topological structure (with respect to λ) of the Nehari set associated to Φ_{λ} , which is defined by

$$\mathcal{N}_{\lambda} := \{ u \in X \setminus \{0\} : \Phi_{\lambda}'(u)u = 0 \}.$$

It can also be written as

$$\mathcal{N}_{\lambda} = \{ u \in X \setminus \{0\} : \varphi_{\lambda,u}'(1) = 0 \} = \{ tu \in X \setminus \{0\} : \varphi_{\lambda,u}'(t) = 0 \},\$$

where $\varphi_{\lambda,u}: [0,\infty) \to \mathbb{R}$ is the *fibering map* given by $\varphi_{\lambda,u}(t) = \Phi_{\lambda}(tu)$ for $t \ge 0$ and $u \in X$. A basic issue related to the Nehari set is to know whether it is given by a natural constraint, i.e. if any critical point of the restriction of Φ_{λ} to \mathcal{N}_{λ} is a critical point of Φ_{λ} . To discuss this issue, let us recall the splitting

$$\mathcal{N}_{\lambda} = \mathcal{N}_{\lambda}^{+} \cup \mathcal{N}_{\lambda}^{0} \cup \mathcal{N}_{\lambda}^{-},$$

where

$$\mathcal{N}_{\lambda}^{+} = \{ u \in \mathcal{N}_{\lambda} : \varphi_{\lambda,u}''(1) > 0 \}, \quad \mathcal{N}_{\lambda}^{-} = \{ u \in \mathcal{N}_{\lambda} : \varphi_{\lambda,u}''(1) < 0 \},$$

and

$$\mathcal{N}_{\lambda}^{0} = \{ u \in \mathcal{N}_{\lambda} : \varphi_{\lambda,u}''(1) = 0 \}.$$

are mutually disjoint sets. By the implicit function theorem, it is promptly seen that (whenever non-empty) \mathcal{N}^+_{λ} and \mathcal{N}^-_{λ} are C^1 manifolds in X, and critical points of Φ_{λ} restricted to $\mathcal{N}^+_{\lambda} \cup \mathcal{N}^-_{\lambda}$ are critical points of Φ_{λ} .

In view of these facts we may refer to $\mathcal{N}_{\lambda}^{+}$ and $\mathcal{N}_{\lambda}^{-}$ as Nehari manifolds, and we see that \mathcal{N}_{λ} is given by a natural constraint if and only if $\mathcal{N}_{\lambda}^{0} = \emptyset$. Let us note that most of the applications of the Nehari manifold method in the litterature (in particular the abstract results in [4,26]) occur when $\mathcal{N}_{\lambda}^{0} = \emptyset$, which prevents the difficulty previously described. This is the case in [5,10,11], which deal with the functional (1.2). The situation where $\mathcal{N}_{\lambda}^{0} \neq \emptyset$ also brings other difficulties, which are related with the behavior of Φ_{λ} on $\mathcal{N}_{\lambda}^{+}$ and $\mathcal{N}_{\lambda}^{-}$. As a matter of fact, in some situations we shall see that Φ_{λ} is unbounded from below on $\mathcal{N}_{\lambda}^{+}$ and its infimum over $\mathcal{N}_{\lambda}^{-}$ is not achieved (see Lemma 2.9 and Remark 2.10), which obviously makes impossible to use a standard minimization technique in these sets. Instead, we shall see that a local minimization procedure can be carried out in $\mathcal{N}_{\lambda}^{+}$.

Let us decribe in the sequel our main results. First we observe that \mathcal{N}^0_{λ} becomes nonempty as soon as λ crosses the threshold value

$$\lambda^* := \inf \left\{ \frac{P_1(u)}{P_2(u)} : u \in X \setminus \{0\}, F(u) = 0 \right\}.$$

Since we intend to minimize Φ_{λ} over \mathcal{N}_{λ}^+ and \mathcal{N}_{λ}^- , let us fix the following notation:

$$c_{\lambda}^{\pm} := \inf_{\mathcal{N}_{\lambda}^{\pm}} \Phi_{\lambda}$$

Under the condition

(H1)
$$\lambda^* = \inf\left\{\frac{P_1(u)}{P_2(u)} : F(u) \ge 0\right\}$$

it turns out that c_{λ}^+ or c_{λ}^- provide a critical point of Φ_{λ} for $\lambda < \lambda^*$. Furthermore, in case λ^* is larger than

$$\mu_* := \inf \left\{ \frac{P_1(u)}{P_2(u)} : u \in X, F(u) < 0 \right\}.$$

both c_{λ}^{+} and c_{λ}^{-} are achieved for $\mu_{*} < \lambda < \lambda^{*}$:

Theorem 1.1 (Minimization up to λ^*). Assume that $\lambda < \lambda^*$. Then $\mathcal{N}^0_{\lambda} = \emptyset$. Moreover, under (H1) the following holds:

- (1) If $\gamma > p$ then c_{λ}^{-} is achieved, i.e. there exists $u_{\lambda} \in \mathcal{N}_{\lambda}^{-}$ such that $\Phi_{\lambda}(u_{\lambda}) = c_{\lambda}^{-} > 0$. If, in addition, $\mu_{*} < \lambda$ then c_{λ}^{+} is achieved, i.e. there exists $w_{\lambda} \in \mathcal{N}_{\lambda}^{+}$ such that $\Phi_{\lambda}(w_{\lambda}) = c_{\lambda}^{+} < 0$.
- (2) If $\gamma < p$ then c_{λ}^{+} is achieved, i.e. there exists $u_{\lambda} \in \mathcal{N}_{\lambda}^{+}$ such that $\Phi_{\lambda}(u_{\lambda}) = c_{\lambda}^{+} < 0$. If, in addition, $\mu_{*} < \lambda$ then c_{λ}^{-} is achieved, i.e. there exists $w_{\lambda} \in \mathcal{N}_{\lambda}^{-}$ such that $\Phi_{\lambda}(w_{\lambda}) = c_{\lambda}^{-} > 0$.

For $\lambda \geq \lambda^*$ we shall be concerned only with minimization over \mathcal{N}_{λ}^+ . Indeed, it turns out that $c_{\lambda}^{-} = 0$ for $\lambda > \lambda^{*}$, see Lemma 2.9 below. The following conditions play an important role in this case:

- (C1) If λ^* is achieved by u then $F'(u) \neq 0$.
- (C2) If λ^* is achieved by u then $H'_{\lambda^*}(u) \neq 0$.
- (S) If $w_n \rightharpoonup w$ in X and $P_1(w_n) \rightarrow P_1(w)$, then $w_n \rightarrow w$ in X.

We point out that (S) is a structural condition needed in our minimization arguments, which is satisfied for instance if $P_1(u) = ||u||^p$, in view of the uniform convexity of X. On the other hand, (C1) and (C2) guarantee, in combination with (H1), that λ^* is achieved by some $u \in \mathcal{N}^0_{\lambda}$, which up to some multiplicative constant yields a critical point of Φ_{λ^*} . A second critical point may still be found in $\mathcal{N}_{\lambda^*}^+$:

Theorem 1.2 (Minimization at λ^*). Suppose (H1), (S), (C1), (C2), and $\mu_* < \lambda^*$. Then:

- (1) λ^* is achieved and its minimizers satisfy F(u) = 0. Moreover there exists t > 0satisfying $tu \in \mathcal{N}_{\lambda^*}^0$ and $\Phi'_{\lambda^*}(tu) = \Phi_{\lambda^*}(tu) = 0$.
- (2) $c_{\lambda^*}^+$ is a critical value of Φ'_{λ^*} , i.e. there exists $u \in \mathcal{N}_{\lambda^*}^+$ such that $\Phi_{\lambda^*}(u) = c_{\lambda^*}^+ < 0$ and $\Phi'_{\lambda^*}(u) = 0$.

Finally, for λ larger than λ^* , the functional Φ_{λ} is no longer bounded from below on \mathcal{N}_{λ}^+ , at least for $\gamma > p$ (see Lemma 2.9 and Remark 2.10). Yet it has a local minimizer therein for λ close to λ^* , which generates then a mountain-pass critical point:

Theorem 1.3 (Minimization beyond λ^*). Suppose (H1), (S), (C1), (C2), and $\mu_* < \lambda^*$. Then there exists $\varepsilon > 0$ having the following properties:

- (1) $c_{\lambda}^{+} = -\infty$ and Φ_{λ} has a local minimizer $u_{\lambda} \in \mathcal{N}_{\lambda}^{+}$ for $\lambda \in (\lambda^{*}, \lambda^{*} + \varepsilon)$. (2) Assume, in addition, the Palais-Smale condition:
- - (PS) If $(u_n) \subset X$ is a sequence such that $(\Phi_{\lambda}(u_n))$ is bounded and $\Phi'_{\lambda}(u_n) \to 0$, then (u_n) has a convergent subsequence.

Then Φ_{λ} has a second critical point (of mountain-pass type) for $\lambda \in (\lambda^*, \lambda^* + \varepsilon)$.

Remark 1.4. Let us make some comments on the assumptions of Theorems 1.2 and 1.3 in the context of our applications. In the case of the problem (1.3), many of these conditions are satisfied if $\int_{\Omega} f(x) |\phi_1|^{\gamma} < 0$, where ϕ_1 is a positive eigenfunction associated to $\lambda_1(p)$, the first eigenvalue of the Dirichlet p-Laplacian. Indeed, in this case we clearly have $\mu_* = \lambda_1(p) < \lambda^*$. Moreover, the fact that $\lambda_1(p)$ is the only eigenvalue associated to a positive eigenfunction of the Dirichlet p-Laplacian implies that (H1) and (C2) satisfied. We refer to Section 3 for more details on this issue as well as the verification of these conditions for our further applications.

The work is organized as follows: in section 2 we prove Theorems 1.1, 1.2 and 1.3. In section 3 we apply these theorems to unify and extend previous existence and multiplicity results for critical points of indefinite, (p, q)-Laplacian, and Kirchhoff type problems.

2. Proofs

To simplify the notation we set

$$H_{\lambda} := P_1 - \lambda P_2$$
, i.e. $\Phi_{\lambda} = \frac{1}{p} H_{\lambda} - \frac{1}{\gamma} F$.

2.1. **Basic properties.** The following result shall be used repeatedly:

Lemma 2.1. If $(u_n) \subset S$, $\lambda_n \to \lambda \ge 0$, and $\limsup H_{\lambda_n}(u_n) \le 0$ then, up to a subsequence, $u_n \rightharpoonup u$ in X and $P_2(u) > 0$. In particular $u \ne 0$.

Proof. Since (u_n) is bounded and X is reflexive, up to a subsequence, $u_n \to u$. If $P_2(u) \leq 0$ then $H_{\lambda}(u) \leq \liminf H_{\lambda_n}(u_n) \leq \limsup H_{\lambda_n}(u_n) \leq 0 \leq P_1(u) \leq H_{\lambda}(u)$, i.e. $H_{\lambda_n}(u_n) \to H_{\lambda}(u) = 0$. Since $P_1 = H_{\lambda} + \lambda P_2$ and $P_2(u_n) \to P_2(u)$, it follows that $P_1(u_n) \to \lambda P_2(u) \leq 0$. But $P_1(u_n) \geq C_1 > 0$, and we reach a contradiction.

Since Φ_{λ} is composed by two homogeneous terms, one may easily formulate a necessary and sufficient condition to have $\mathcal{N}_{\lambda} \setminus \mathcal{N}_{\lambda}^{0} \neq \emptyset$:

Lemma 2.2. If $u \in \mathcal{N}_{\lambda} \setminus \mathcal{N}_{\lambda}^{0}$ then $H_{\lambda}(u)F(u) > 0$. Conversely, if $H_{\lambda}(u)F(u) > 0$ then there exists a unique $t = t_{\lambda}(u) > 0$ such that $tu \in \mathcal{N}_{\lambda} \setminus \mathcal{N}_{\lambda}^{0}$, which is given by $t_{\lambda}(u) = (H_{\lambda}(u)/F(u))^{\frac{1}{\gamma-p}}$.

Proof. If $u \in \mathcal{N}_{\lambda}$, then $J_{\lambda}(u) = 0$ and thus $H_{\lambda}(u) = F(u)$. Conversely, if $H_{\lambda}(u)$ and F(u) are nonzero and have the same sign, then the fibering map $\varphi_{\lambda,u}$ has a unique positive critical point t, so that $tu \in \mathcal{N}_{\lambda}$. The equation $\varphi'_{\lambda,u}(t) = 0$ yields the desired expression of $t_{\lambda}(u)$. \Box

Let us set

$$D_{\lambda}^{+} := \{ u \in X \setminus \{0\} : H_{\lambda}(u), F(u) > 0 \},\$$

$$D_{\lambda}^{-} := \{ u \in X \setminus \{0\} : H_{\lambda}(u), F(u) < 0 \},\$$

and

$$D_{\lambda}^{0} := \{ u \in X \setminus \{0\} : H_{\lambda}(u) = F(u) = 0 \}.$$

Since H_{λ} and F are homogeneous, we see that D_{λ}^+ , D_{λ}^- and D_{λ}^0 are cones, i.e. $u \in D_{\lambda}^+$ if and only if $tu \in D_{\lambda}^+$ for any t > 0. The following properties are straightforward, so we omit their proofs:

Proposition 2.3. There holds $\mathcal{N}^0_{\lambda} = D^0_{\lambda}$. Furthermore:

- (1) If $\gamma > p$, then: (a) For each $u \in D_{\lambda}^{+}$, the point $t_{\lambda}(u)$ is a non-degenerate global maximum point of $\varphi_{\lambda,u}$. Moreover $\mathcal{N}_{\lambda}^{-} = \{t_{\lambda}(u)u : u \in D_{\lambda}^{+}\}$. In particular $\mathcal{N}_{\lambda}^{-} \subset D_{\lambda}^{+}$.
 - (b) For each $u \in D_{\lambda}^{-}$, the point $t_{\lambda}(u)$ is a non-degenerate global minimum point of $\varphi_{\lambda,u}$. Moreover $\mathcal{N}_{\lambda}^{+} = \{t_{\lambda}(u)u : u \in D_{\lambda}^{-}\}$. In particular $\mathcal{N}_{\lambda}^{+} \subset D_{\lambda}^{-}$.
- (2) If $\gamma < p$, then:
 - (a) For each $u \in D_{\lambda}^+$, the point $t_{\lambda}(u)$ is a non-degenerate global minimum point of $\varphi_{\lambda,u}$. Moreover $\mathcal{N}_{\lambda}^+ = \{t_{\lambda}(u)u : u \in D_{\lambda}^+\}$. In particular $\mathcal{N}_{\lambda}^+ \subset D_{\lambda}^+$.
 - (b) For each $u \in D_{\lambda}^{-}$, the point $t_{\lambda}(u)$ is a non-degenerate global maximum point of $\varphi_{\lambda,u}$. Moreover $\mathcal{N}_{\lambda}^{-} = \{t_{\lambda}(u)u : u \in D_{\lambda}^{-}\}$. In particular $\mathcal{N}_{\lambda}^{-} \subset D_{\lambda}^{-}$.

Recall that

$$\lambda^* := \inf \left\{ \frac{P_1(u)}{P_2(u)} : u \in X \setminus \{0\}, F(u) = 0 \right\},$$

and

$$\mu_* := \inf \left\{ \frac{P_1(u)}{P_2(u)} : u \in X, F(u) < 0 \right\}.$$

We also introduce

$$\mu^* := \sup\left\{\frac{P_1(u)}{P_2(u)} : u \in X, F(u) > 0\right\}.$$

Let us prove now that λ^* is the threshold value for \mathcal{N}_{λ} to be a manifold and that μ_*, μ^* determine whether \mathcal{N}_{λ}^+ and \mathcal{N}_{λ}^- are empty or not:

Proposition 2.4. We have $\mathcal{N}^0_{\lambda} = \emptyset$ for any $\lambda < \lambda^*$. In addition:

Proof. If $\lambda < \lambda^*$, then for any $u \in X \setminus \{0\}$ satisfying F(u) = 0 we have $P_1(u)/P_2(u) > \lambda$ or, equivalently, $H_{\lambda}(u) > 0$, so that $\mathcal{N}_{\lambda}^0 = \mathcal{D}_{\lambda}^0 = \emptyset$. Let us assume that $\gamma > p$ and prove (1).

- (a) Indeed, if $\lambda < \mu^*$ then there exists $u \in X \setminus \{0\}$ such that F(u) > 0 and $P_1(u)/P_2(u) > \lambda$ or, equivalently, $H_{\lambda}(u) > 0$ and thus, by Proposition 2.3 it follows that $\mathcal{N}_{\lambda}^- \neq \emptyset$. Now, if $\lambda \ge \mu^*$, then it is clear that for every u satisfying F(u) > 0 we must have $H_{\lambda}(u) \le 0$, i.e. $D_{\lambda}^+ = \emptyset$, so that by Proposition 2.3, we conclude that $\mathcal{N}_{\lambda}^- = \emptyset$.
- (b) If $\lambda > \mu_*$ then there exists $u \in X \setminus \{0\}$ such that F(u) < 0 and $P_1(u)/P_2(u) < \lambda$ or, equivalently, $H_{\lambda}(u) < 0$ and thus, by Lemma 2.2 it follows that $\mathcal{N}_{\lambda}^+ \neq \emptyset$. Now, if $\lambda < \mu_*$, then it is clear that for all u satisfying F(u) < 0 we must have $H_{\lambda}(u) \ge 0$, i.e. $D_{\lambda}^- = \emptyset$, so that by Proposition 2.3, we have $\mathcal{N}_{\lambda}^+ = \emptyset$.

The proof of (2) is completely similar, so we omit it.

2.2. Minimization up to λ^* . In the sequel we shall use the assumption (H1), which we recall below:

(H1)
$$\lambda^* = \inf \left\{ \frac{P_1(u)}{P_2(u)} : u \in X, F(u) \ge 0 \right\}.$$

It is straightforward that this condition implies that $\lambda^* < \mu^*$ and the following properties:

$$H_{\lambda}(u) > 0$$
 for any $u \in X \setminus \{0\}$ such that $F(u) \ge 0$, and any $\lambda < \lambda^*$, (2.1)

and

$$H_{\lambda^*}(u) \ge 0$$
 for any $u \in X$ such that $F(u) \ge 0$. (2.2)

The next result shows in particular that (2.1) actually holds in a stronger form under (H1), and that N_{λ}^{-} , N_{λ}^{+} are away from zero and infinity, respectively, if $\lambda < \lambda^{*}$:

Proposition 2.5. Suppose (H1) and $\lambda < \lambda^*$. Then:

- (1) There exist C, D > 0 such that (a) $H_{\lambda}(u) \ge C ||u||^{p}$, for every $u \in X$ such that $F(u) \ge 0$. (b) $F(u) \le -D ||u||^{\gamma}$ for every $u \in X$ such that $H_{\lambda}(u) \le 0$.
- (2) $\mathcal{N}_{\lambda}^{-}$ is away from zero, i.e. there exists c > 0 such that $c \leq ||u||$ for all $u \in \mathcal{N}_{\lambda}^{-}$. Moreover Φ_{λ} is coercive over $\mathcal{N}_{\lambda}^{-}$.

 $\mathbf{6}$

(3) \mathcal{N}_{λ}^+ is bounded, i.e. there exists C > 0 such that $||u|| \leq C$ for all $u \in \mathcal{N}_{\lambda}^+$.

Proof.

(1) We prove only (a), since (b) is similar. Suppose, on the contrary, that there exists a sequence $(u_n) \subset S$ such that $H_{\lambda}(u_n) < 1/n$ and $F(u_n) \ge 0$ for all n. By Lemma 2.1 we can assume that $u_n \rightharpoonup u \ne 0$ in X. Therefore

$$H_{\lambda}(u) \le \liminf H_{\lambda}(u_n) \le 0 \le \limsup F(u_n) \le F(u),$$

which contradicts (2.1).

(2) Indeed, note that

$$H_{\lambda}(u) = F(u), \quad \forall u \in \mathcal{N}_{\lambda},$$

$$(2.3)$$

and thus

$$\Phi_{\lambda}(u) = \left(\frac{\gamma - p}{p\gamma}\right) H_{\lambda}(u) = \left(\frac{\gamma - p}{p\gamma}\right) F(u), \quad \forall u \in \mathcal{N}_{\lambda}.$$
(2.4)

Case 1: $\gamma > p$. Recall that F(u) > 0 and $H_{\lambda}(u) > 0$ for all $u \in \mathcal{N}_{\lambda}^{-}$. By (1) we conclude from (2.3) that

$$C||u||^{p} \le H_{\lambda}(u) = F(u) \le C_{3}||u||^{\gamma}, \quad \forall u \in \mathcal{N}_{\lambda}^{-}.$$

and since $\gamma > p$, it follows that there exists c > 0 such that $c \leq ||u||$ for all $u \in \mathcal{N}_{\lambda}^{-}$. Clearly by (2.4) and (1) we also have that Φ_{λ} is coercive.

Case 2: $\gamma < p$. Recall that F(u) < 0 and $H_{\lambda}(u) < 0$ for all $u \in \mathcal{N}_{\lambda}^{-}$. By (1) and (2.3) we infer that

$$-d\|u\|^{p} \leq H_{\lambda}(u) = F(u) \leq -C\|u\|^{\gamma}, \quad \forall u \in \mathcal{N}_{\lambda}^{-},$$

where d > 0. Since $\gamma < p$, we obtain the desired conclusion.

(3) Case 1: $\gamma > p$. Recall that F(u) < 0 and $H_{\lambda}(u) < 0$ for all $u \in \mathcal{N}_{\lambda}^+$. By (1) we conclude from (2.3) that

$$-c\|u\|^{p} \le H_{\lambda}(u) = F(u) \le -C\|u\|^{\gamma}, \quad \forall u \in \mathcal{N}_{\lambda}^{+},$$

where c > 0 and since $\gamma > p$, we obtain the desired inequality.

Case 2: $\gamma < p$. Recall that F(u) > 0 and $H_{\lambda}(u) > 0$ for all $u \in \mathcal{N}_{\lambda}^+$. By (1) we conclude from (2.3) that

$$C||u||^{p} \le H_{\lambda}(u) = F(u) \le C_{3}||u||^{\gamma}, \quad \forall u \in \mathcal{N}_{\lambda}^{-}.$$

and since $\gamma < p$, it follows that there exists C > 0 such that $||u|| \leq C$ for all $u \in \mathcal{N}_{\lambda}^+$.

Proof of Theorem 1.1.

Let $\lambda < \lambda^*$. By Proposition 2.4 we have $\mathcal{N}_{\lambda}^0 = \emptyset$ so that \mathcal{N}_{λ} is a C^1 manifold. Note also that $\lambda^* \leq \mu^*$, so according to Proposition 2.4 we have $\mathcal{N}_{\lambda}^- \neq \emptyset$ if $\gamma > p$ and $\mathcal{N}_{\lambda}^+ \neq \emptyset$ if $\gamma < p$. If, in addition, $\lambda > \mu^*$ then $\mathcal{N}_{\lambda}^+ \neq \emptyset$ if $\gamma > p$ and $\mathcal{N}_{\lambda}^- \neq \emptyset$ if $\gamma < p$. It remains to show that c_{λ}^+ and c_{λ}^- are achieved whenever \mathcal{N}_{λ}^+ and \mathcal{N}_{λ}^- are nonempty, respectively:

(1) First we deal with c_{λ}^- . Proposition 2.3 yields that $c_{\lambda}^- \ge 0$. Let $(u_n) \subset \mathcal{N}_{\lambda}^-$ be a minimizing sequence for c_{λ}^- . By Proposition 2.5(2) we can assume that $u_n \rightharpoonup u$ in X. We claim that $u \neq 0$. Indeed, note by Proposition 2.5(1) that

$$C||u||^{p} \le \liminf C||u_{n}||^{p} \le \liminf H_{\lambda}(u_{n}) = \liminf F(u_{n}) \le F(u).$$

$$(2.5)$$

Thus if u = 0, then $u_n \to 0$, which contradicts Proposition 2.5(2). Hence $u \neq 0$. Now we consider two cases:

Case 1: $\gamma > p$. We claim that $u \in D_{\lambda}^+$. In fact, note that $F(u) \ge \limsup F(u_n) \ge 0$, which implies that $H_{\lambda}(u) > 0$ by (2.1). From

$$H_{\lambda}(u) \le \liminf H_{\lambda}(u_n) = \liminf F(u_n) \le F(u)$$
(2.6)

it follows that F(u) > 0. Thus Proposition 2.3 provides us with $t_{\lambda}(u) > 0$ such that $t_{\lambda}(u)u \in \mathcal{N}_{\lambda}^{-}$. Therefore

$$\Phi_{\lambda}(t_{\lambda}(u)u) \leq \liminf \Phi_{\lambda}(t_{\lambda}(u)u_n) \leq \liminf \Phi_{\lambda}(u_n) = c_{\lambda}^{-},$$

and so $\Phi_{\lambda}(t_{\lambda}(u)u) = c_{\lambda}^{-}$.

Case 2: $\gamma < p$. We claim that $u \in D_{\lambda}^-$. Indeed, note that $H_{\lambda}(u) \leq \liminf H_{\lambda}(u_n) \leq 0$, which implies that F(u) < 0, by (2.1). It follows from (2.6) that $H_{\lambda}(u) < 0$. Thus by Proposition 2.3, there exists $t_{\lambda}(u) > 0$ such that $t_{\lambda}(u)u \in \mathcal{N}_{\lambda}^-$. Therefore

 $\Phi_{\lambda}(t_{\lambda}(u)u) \leq \liminf \Phi_{\lambda}(t_{\lambda}(u)u_n) \leq \liminf \Phi_{\lambda}(u_n) = c_{\lambda}^{-},$

and hence $\Phi_{\lambda}(t_{\lambda}(u)u) = c_{\lambda}^{-}$.

(2) Let us consider now c_{λ}^+ . Propositions 2.3 and 2.5(3) yield that $-\infty < c_{\lambda}^+ < 0$. Let $(u_n) \subset \mathcal{N}_{\lambda}^+$ be a minimizing sequence for c_{λ}^+ . By Proposition 2.5(3) we can assume that $u_n \rightharpoonup u$ in X and it is clear that $u \neq 0$. Once again we consider two cases:

Case 1: $\gamma > p$. We claim that $u \in D_{\lambda}^-$. Indeed, note that $H_{\lambda}(u) \leq \liminf H_{\lambda}(u_n) \leq 0$, and (2.1) implies that F(u) < 0. From (2.6) it follows that $H_{\lambda}(u) < 0$. Thus from Proposition 2.3, there exists $t_{\lambda}(u) > 0$ such that $t_{\lambda}(u)u \in \mathcal{N}_{\lambda}^+$. Therefore

$$\Phi_{\lambda}(t_{\lambda}(u)u) \le \Phi_{\lambda}(u) \le \liminf \Phi_{\lambda}(u_n) = c_{\lambda}^+,$$

and hence $\Phi_{\lambda}(t_{\lambda}(u)u) = c_{\lambda}^{+}$.

Case 2: $\gamma < p$. We claim that $u \in D_{\lambda}^+$. Indeed, note that $F(u) \ge \limsup F(u_n) \ge 0$, which implies that $H_{\lambda}(u) > 0$. By (2.6) it follows that F(u) > 0. Thus from Proposition 2.3, there exists $t_{\lambda}(u) > 0$ such that $t_{\lambda}(u)u \in \mathcal{N}_{\lambda}^+$. Therefore

$$\Phi_{\lambda}(t_{\lambda}(u)u) \leq \Phi_{\lambda}(u) \leq \liminf \Phi_{\lambda}(u_{n}) = c_{\lambda}^{+},$$

and therefore $\Phi_{\lambda}(t_{\lambda}(u)u) = c_{\lambda}^{+}.$

We conclude this subsection showing that for $\mu_* < \lambda < \lambda^*$ the minimization procedure over \mathcal{N}_{λ}^+ and \mathcal{N}_{λ}^- is equivalent to minimizing H_{λ} under the constraints $F(u) = \pm 1$. The latter method has been used to deal with the problem (1.3) in [24] for $\gamma > p = 2$, and in [23] for $\gamma < p$.

Lemma 2.6. Let $\mu_* < \lambda < \lambda^*$ and $m_{\lambda}^{\pm} := \inf \{H_{\lambda}(u) : u \in X, F(u) = \pm 1\}$. Then: (1) m_{λ}^{\pm} are achieved, and $m_{\lambda}^{\pm} > 0 > m_{\lambda}^{-}$.

(2) There holds

$$c_{\lambda}^{+} = \begin{cases} \frac{p - \gamma}{p \gamma} (-m_{\lambda}^{-})^{\frac{\gamma}{\gamma - p}} & \text{if } \gamma > p, \\ \frac{\gamma - p}{p \gamma} (m_{\lambda}^{+})^{\frac{\gamma}{\gamma - p}} & \text{if } \gamma < p. \end{cases}$$
(2.7)

and

$$c_{\lambda}^{-} = \begin{cases} \frac{\gamma - p}{p\gamma} (m_{\lambda}^{+})^{\frac{\gamma}{\gamma - p}} & \text{if } \gamma > p, \\ \frac{p - \gamma}{p\gamma} (-m_{\lambda}^{-})^{\frac{\gamma}{\gamma - p}} & \text{if } \gamma < p. \end{cases}$$
(2.8)

(3) The maps $\lambda \mapsto m_{\lambda}^{\pm}$ are concave (therefore continuous) and decreasing in (μ_*, λ^*) . In particular, the maps $\lambda \mapsto c_{\lambda}^{\pm}$ are decreasing and continuous in (μ_*, λ^*) .

Proof.

- (1) Let us first show that m_{λ}^{\pm} are finite. Indeed, if $(u_n) \subset X$ is such that $H_{\lambda}(u_n) \to -\infty$ and $F(u_n) = \pm 1$ then (u_n) is unbounded, so we can assume that $||u_n|| \to \infty$ and $v_n := \frac{u_n}{||u_n||} \rightharpoonup v$. From $H_{\lambda}(v_n) \leq 0$ we have $v \neq 0$ and $H_{\lambda}(v) \leq 0 \leq F(v)$, which contradicts (2.1). Thus m_{λ}^{\pm} are both finite. The previous discussion also shows that any minimizing sequence (u_n) for m_{λ}^{\pm} is bounded, so we can assume that $u_n \to u$ in X. If $H_{\lambda}(u_n) \to m_{\lambda}^+$ and $F(u_n) = 1$ then $H_{\lambda}(u) \leq m_{\lambda}^+$ and $F(u) \geq 1$. Hence $\tilde{u} := \frac{u}{F(u)^{\frac{1}{\gamma}}}$ satisfies $F(\tilde{u}) = 1$ and $H_{\lambda}(\tilde{u}) \leq H_{\lambda}(u) \leq m_{\lambda}^+$, i.e. it achieves m_{λ}^+ . A similar argument shows that m_{λ}^- is also achieved. Moreover, from (2.1) it is clear that $m_{\lambda}^+ > 0$. Note also that $m_{\lambda}^- < 0$ if and only if there exists $u \in X$ such that F(u) = -1 and $H_{\lambda}(u) < 0$, which is equivalent to have $\lambda > \frac{P_1(u)}{P_2(u)}$ for some $u \in X$ such that F(u) < 0, and this holds since $\lambda > \mu_*$.
- (2) We prove (2.7) for $\gamma > p$ (the case $\gamma < p$ and (2.8) are similar). Let v_{λ} achieve m_{λ}^{-} , i.e. $H_{\lambda}(v_{\lambda}) = m_{\lambda}^{-} < 0$ and $F(v_{\lambda}) = -1$. Thus $v_{\lambda} \in D_{\lambda}^{-}$ and $t_{\lambda}(v_{\lambda}) = (-m_{\lambda}^{-})^{\frac{1}{\gamma-p}}$, so

$$c_{\lambda}^{+} \leq \Phi_{\lambda}(t_{\lambda}(v_{\lambda})v_{\lambda}) = \frac{1}{p}t_{\lambda}(v_{\lambda})^{p}m_{\lambda}^{-} + \frac{1}{\gamma}t_{\lambda}(v_{\lambda})^{\gamma} = \frac{p-\gamma}{p\gamma}(-m_{\lambda}^{-})^{\frac{\gamma}{\gamma-p}}.$$

On the other hand, since $H(u_{\lambda}) = F(u_{\lambda}) < 0$ we have

$$m_{\lambda}^{-} \leq H_{\lambda}\left(\frac{u_{\lambda}}{(-F(u_{\lambda}))^{\frac{1}{\gamma}}}\right) = -(-H(u_{\lambda}))^{\frac{\gamma-p}{\gamma}}$$

It follows that $(-m_{\lambda}^{-})^{\frac{\gamma}{\gamma-p}} \ge -H(u_{\lambda})$, so that $c_{\lambda}^{+} = -\frac{p-\gamma}{p\gamma}H(u_{\lambda}) \ge \frac{p-\gamma}{p\gamma}(-m_{\lambda}^{-})^{\frac{\gamma}{\gamma-p}}$.

(3) The concavity of $\lambda \mapsto m_{\lambda}^{\pm}$ follows from the fact that m_{λ}^{\pm} are pointwise infima of $H_{\lambda}(u)$, which is affine (therefore concave) with respect to λ . Let $\mu_* < \lambda < \lambda' < \lambda^*$ and v_{λ} achieve m_{λ}^{\pm} . Then

$$m_{\lambda'}^{\pm} \le H_{\lambda'}(v_{\lambda}) = H_{\lambda}(v_{\lambda}) + (\lambda - \lambda')P_2(v_{\lambda}) < H_{\lambda}(v_{\lambda}) = m_{\lambda}^{\pm}.$$

The assertions on $\lambda \mapsto c_{\lambda}^{+}$ follow from (2.7).

2.3. Minimization at λ^* . We start proving that λ^* is achieved and provides us with a critical point of Φ_{λ^*} with zero energy if the following conditions are satisfied:

(C1): If λ^* is achieved by u, then $F'(u) \neq 0$. (C2): If λ^* is achieved by u, then $H'_{\lambda^*}(u) \neq 0$.

Proof of Theorem 1.2 (1).

First we show that λ^* is achieved under (H1). Since P_1/P_2 is 0-homogeneous, we can find a minimizing sequence for λ^* in S. By the weak (semi)continuity properties of P_1 , P_2 and F, it follows that λ^* is achieved by some u. If F(u) > 0 then u is a local minimizer of P_1/P_2 over X, and consequently a critical point of this quotient, so that $P'_1(u) - \frac{P_1(u)}{P_2(u)}P'_2(u) = 0$, i.e. $H'_{\lambda^*}(u) = 0$, which contradicts (C2). Thus F(u) = 0. By (C1) we may apply Lagrange's multiplier rule to find some $\alpha \in \mathbb{R}$ such that

$$\left(\frac{P_1(u)}{P_2(u)}\right)' = \alpha F'(u)$$

which implies that

$$H'_{\lambda^*}(u) = P_2(u)\alpha F'(u).$$
 (2.9)

(C2) yields that $\alpha \neq 0$. We claim that $\alpha > 0$. Otherwise there exists $v \in X$ such that $H'_{\lambda^*}(u)v < 0 < F'(u)v$. Thus

$$\lim_{h \to 0^+} \frac{H_{\lambda^*}(u+hv)}{h} = \lim_{h \to 0^+} \frac{H_{\lambda^*}(u+hv) - H_{\lambda^*}(u)}{h} < 0$$

and

$$\lim_{h \to 0^+} \frac{F(u+hv)}{h} = \lim_{h \to 0^+} \frac{F(u+hv) - F(u)}{h} > 0,$$

which implies that there exists $\delta > 0$ such that $H_{\lambda^*}(u + \delta v) < 0 < F(u + \delta v)$ and hence, for $\lambda < \lambda^*$ and close to λ^* we conclude that $H_{\lambda}(u + \delta v) < 0$ and $F(u + \delta v) > 0$, a contradiction with (2.1). Therefore $\alpha > 0$ and setting $t = (\alpha P_2(u))^{-1/\gamma}$ we conclude from (2.9) that tu is a critical point of Φ_{λ^*} . Moreover it is clear that $tu \in \mathcal{N}_{\lambda^*}^0$ and $\Phi_{\lambda^*}(tu) = 0$.

Let us set

$$\Theta_{\lambda} = \{ u / \| u \| : u \in \mathcal{N}_{\lambda}^+ \}.$$

From Proposition 2.4 we have $\Theta_{\lambda} \neq \emptyset$ for $\lambda > \mu_*$ if $\gamma > p$, and for $\lambda < \mu^*$ if $\gamma < p$.

Lemma 2.7.

(1) If $\gamma > p$ then Θ_{λ} is increasing with respect to λ , i.e. $\Theta_{\lambda} \subset \Theta_{\lambda'}$ for $\mu_* < \lambda < \lambda'$.

(2) If $\gamma < p$ then Θ_{λ} is decreasing with respect to λ , i.e. $\Theta_{\lambda'} \subset \Theta_{\lambda}$ for $\lambda < \lambda' < \mu^*$.

Proof. Let $\gamma > p$ and $\mu_* < \lambda < \lambda'$. If $v \in \Theta_{\lambda}$, then $H_{\lambda}(v) < 0$ and F(v) < 0. Thus $H_{\lambda'}(v) < 0$ and consequently $v \in \Theta_{\lambda'}$. The proof of (2) is similar (recall that $H_{\lambda'}(v) > 0$ in this case).

Lemma 2.8. Let $\lambda_n \nearrow \lambda^*$ and $u_n \in \mathcal{N}_{\lambda_n}^+$ be such that $\Phi_{\lambda_n}(u_n) = c_{\lambda_n}^+$. If $\gamma > p$ and $u_n \rightharpoonup u \in \mathcal{D}_{\lambda^*}^-$ (respect. $\gamma < p$ and $u_n \rightharpoonup u \in \mathcal{D}_{\lambda^*}^+$) then $u \in \mathcal{N}_{\lambda^*}^+$, $\Phi_{\lambda^*}(u) = c_{\lambda^*}^+$ and $\Phi'_{\lambda^*}(u) = 0$.

Proof. It suffices to show that $\Phi_{\lambda^*}(u) = c_{\lambda^*}^+$. Since $t_{\lambda^*}(u)$ is the global minimum point of $\varphi_{\lambda^*,u}$ we have $\Phi_{\lambda^*}(u) \ge \Phi_{\lambda^*}(t_{\lambda^*}(u)u) \ge c_{\lambda^*}^+$. Suppose, by contradiction, that $\Phi_{\lambda^*}(u) > c_{\lambda^*}^+$.

Case 1: $\gamma > p$

Given $\alpha \in (c_{\lambda^*}^+, \Phi_{\lambda^*}(u))$, we choose $v \in \Theta_{\lambda^*}$ such that $c_{\lambda^*}^+ \leq \Phi_{\lambda^*}(t_{\lambda^*}(v)v) < \alpha$. By continuity there exists $\delta > 0$ such that $\Phi_{\lambda}(t_{\lambda}(v)v) < \alpha$ for all $\lambda \in (\lambda^* - \delta, \lambda^*)$. Therefore, for *n* sufficiently large we have

$$\Phi_{\lambda_n}(u_n) = c_{\lambda_n}^+ \le \Phi_{\lambda_n}(t_{\lambda_n}(v)v) < \alpha < \Phi_{\lambda^*}(u),$$

which contradicts the fact that $\Phi_{\lambda^*}(u) \leq \liminf \Phi_{\lambda_n}(u_n)$.

Case 2: $\gamma < p$

Given $\alpha \in (c_{\lambda^*}^+, \Phi_{\lambda^*}(u))$, choose $v \in \Theta_{\lambda^*}$ such that $c_{\lambda^*}^+ \leq \Phi_{\lambda^*}(t_{\lambda^*}(v)v) < \alpha$. Take $\delta > 0$ such that $t_{\lambda}(v)$ is well defined and $\Phi_{\lambda}(t_{\lambda}(v)v) < \alpha$ for $\lambda \in (\lambda^* - \delta, \lambda^*)$. One may argue as in the previous case to reach a contradiction.

We are now in position to provide the

Proof of Theorem 1.2 (2).

Choose a sequence $\lambda_n \nearrow \lambda^*$ and $u_n \in \mathcal{N}_{\lambda_n}^+$ such that $\Phi_{\lambda_n}(u_n) = c_{\lambda_n}^+$ (which exists by Theorem 1.1). We write $u_n = t_n w_n$, where $w_n \in \Theta_{\lambda_n}$ and $t_n := t_{\lambda_n}(w_n)$ is given by

$$t_n = \left(\frac{H_{\lambda_n}(w_n)}{F(w_n)}\right)^{\frac{1}{\gamma-p}}.$$
(2.10)

Note also that by Lemma 2.6(3) there exists c < 0 such that

$$c > c_{\lambda_n}^+ = \Phi_{\lambda_n}(t_n w_n) = \pm \frac{\gamma - p}{p\gamma} t_n^p |H_{\lambda_n}(w_n)|, \qquad (2.11)$$

where the sign – is for the case $\gamma > p$ and the sign + corresponds to $\gamma < p$. Let us show that $u_n \rightharpoonup u \in \mathcal{N}_{\lambda^*}^+$, so that Lemma 2.8 yields the desired conclusion:

Case 1: $\gamma > p$

Since $H_{\lambda_n}(w_n) < 0$, by Lemma 2.1 we can assume that $w_n \rightharpoonup w \neq 0$ in X. We claim that (t_n) is bounded and away from zero. Indeed, suppose by contradiction that, up to a subsequence $t_n \rightarrow \infty$. Then, by (2.10), we conclude that $F(w_n) \rightarrow 0$ and hence $F(w) \geq \limsup F(w_n) = 0$. Since $H_{\lambda^*}(w) \leq \liminf H_{\lambda_n}(w_n) \leq 0$, we deduce from (2.2) that $H_{\lambda^*}(w) = 0$. It follows from Theorem 1.2(1) that $H_{\lambda^*}(w) = F(w) = 0$ and w achieves λ^* . Moreover since

$$0 = H_{\lambda^*}(w) \le \liminf H_{\lambda_n}(w_n) \le 0,$$

we have $H_{\lambda_n}(w_n) \to 0$, so $P_1(w_n) \to P_1(w)$. Hence, by (S2) we obtain $w_n \to w$ in X, and from $t_n^{p-1} H'_{\lambda_n}(w_n) = t_n^{\gamma-1} F'(w_n), \quad \forall n,$

we infer that F'(w) = 0, which contradicts (C1). So (t_n) is bounded. Now it is clear from (2.11) that $t_n \neq 0$, so the claim is proved and we can suppose that $t_n \to t \in (0, \infty)$. Thus $u_n = t_n w_n \rightarrow u := tw$. Note again by (2.11) that $H_{\lambda^*}(u) = t^p H_{\lambda^*}(w) \leq t^p \liminf H_{\lambda_n}(w_n) < 0$ and consequently F(u) < 0, i.e. $u \in \mathcal{D}_{\lambda^*}^-$. The conclusion follows from Lemma 2.8.

Case 2: $\gamma < p$

Let us show again that (t_n) is bounded and away from zero. Indeed, suppose that up to a subsequence $t_n \to \infty$, so that by (2.10) we conclude that $H_{\lambda_n}(w_n) \to 0$ and thus by Lemma 2.1 we can assume that $w_n \rightharpoonup w \neq 0$ in X. Since $F(w) \geq \limsup F(w_n) = 0$ and $H_{\lambda^*}(w) \leq \liminf_{n \to \infty} H_{\lambda_n}(w_n) \leq 0$, we deduce from (2.2) and Theorem 1.2(1) that $H_{\lambda^*}(w) = F(w) = 0$ and therefore w achieves λ^* . Moreover $w_n \to w$ in X and since

$$t_n^{p-1}H'_{\lambda_n}(w_n) = t_n^{\gamma-1}F'(w_n), \quad \forall n,$$

we conclude that $H'_{\lambda^*}(w) = 0$, which contradicts (C2). Thus (t_n) is bounded, and the proof can then be concluded as in the previous case.

2.4. Local Minimization Beyond λ^* . In this Section we look for critical points of Φ_{λ} for $\lambda > \lambda^*$. We start with the following observation, which shows that c_{λ}^+ (for $\gamma > p$) and c_{λ}^- (for $\gamma < p$) are no longer achieved.

Lemma 2.9. Suppose (H1), (C1), and (C2).

- (1) If $\gamma > p$, then $c_{\lambda}^{+} = -\infty$ for all $\lambda > \lambda^{*}$. (2) If $\gamma < p$, then $c_{\overline{\lambda}}^{-} = 0$ for all $\lambda > \lambda^{*}$.

Proof.

(1) By Theorem 1.2(1) there exists $u \in X$ such that $H_{\lambda^*}(u) = F(u) = 0$ and

$$\frac{1}{p}H'_{\lambda^*}(u) = \frac{1}{\gamma}F'(u) \neq 0.$$
(2.12)

We choose $v \in X$ such that $H'_{\lambda^*}(u)v < 0$ and F'(u)v < 0. As in the proof of Theorem 1.2(1), we have

 $H_{\lambda^*}(u+sv) < 0$ and F(u+sv) < 0,

for s > 0 small enough. Now fix $\lambda > \lambda^*$ and note that $H_{\lambda}(u + sv) < 0$, which implies, in particular, that $u + sv \in D_{\lambda}^{-}$. Moreover, by continuity $H_{\lambda}(u + sv) \to H_{\lambda}(u) < 0$ $H_{\lambda^*}(u) = 0$ as $s \to 0^+$. Therefore $t_{\lambda}(u + sv)(u + sv) \in \mathcal{N}_{\lambda}^+$ and

$$\lim_{s \to 0^+} \Phi_{\lambda}(t_{\lambda}(u+sv)(u+sv)) = \lim_{s \to 0^+} -\frac{\gamma-p}{\gamma p} \frac{|H_{\lambda}(u+sv)|^{\frac{1}{\gamma-p}}}{|F(u+sv)|^{\frac{p}{\gamma-p}}} = -\infty$$

(2) The argument is similar to the previous one. Note that now if $u + sv \in D^{\pm}_{\lambda}$ then $t_{\lambda}(u+sv)(u+sv) \in \mathcal{N}_{\lambda}^{\pm}$ and $\gamma - p < 0$.

Remark 2.10.

- (1) Proving that $c_{\lambda}^{+} = -\infty$ for $\lambda > \lambda^{*}$ when $\gamma < p$ (as well as $c_{\lambda}^{-} = 0$ for $\lambda > \lambda^{*}$ when $\gamma > p$) is more delicate and we are not able to do it in this general setting. However, this is indeed the case in our applications (see Remark 3.8). Finally, let us observe that this result can be proved if we assume that the set $\{u \in X : F(u) > 0\}$ is pathwise connected.
- (2) An argument similar to the one in the previous proof shows that $c_{\lambda^*} = 0$ for $\gamma < p$.

To overcome the problem posed by Lemma 2.9, we show that Φ_{λ} has a local minimizer over \mathcal{N}^+_{λ} . Let us first extend Proposition 2.5:

Proposition 2.11. Suppose (H1), (C1), (C2), and $\mu_* < \lambda^*$.

(1) Given $\mu > 0$ there exist $C, \varepsilon > 0$ such that $H_{\lambda}(u) \ge C \|u\|^p$ for any $u \in X$ satisfying $F(u) \ge \mu ||u||^{\gamma}$, and any $\lambda \in [\lambda^*, \lambda^* + \varepsilon]$.

(2) Given $\mu \in (\mu_*, \lambda^*)$ there exists D > 0 such that $F(u) \leq -D \|u\|^{\gamma}$ for any $u \in X$ such that $H_{\mu}(u) \leq 0$.

Proof.

- (1) Suppose, on the contrary, that there exists a sequence $\lambda_n \to \lambda^*$ with $\lambda_n > \lambda^*$ and $(u_n) \subset X$ such that $F(v_n) \ge \mu$ and $H_{\lambda_n}(v_n) \to 0$, where $v_n = u_n/||u_n||$, for every n. Thus $v_n \rightharpoonup v \ne 0$, by Lemma 2.1. Therefore $H_{\lambda^*}(v) \le 0 < \mu \le F(v)$ and by definition of λ^* we have $H_{\lambda^*}(v) = 0$, i.e. λ^* is achieved by v. Theorem 1.2 (1) implies that F(v) = 0, which yields a contradiction.
- (2) Arguing by contradiction we find a sequence $(u_n) \subset S$ such that $H_{\mu}(u_n) \leq 0$ and $F(u_n) \geq -1/n$ for all n. By Lemma 2.1 we can assume that $u_n \rightharpoonup u \neq 0$ in X. Therefore $H_{\mu}(u) \leq 0 \leq F(u)$, which contradicts (2.1).

Next we fix $(\lambda, \mu) \in (\lambda^*, \infty) \times (\mu_*, \lambda^*)$ and set

$$\mathcal{N}_{\lambda,\mu}^{+} := \begin{cases} \{u \in \mathcal{N}_{\lambda}^{+} : H_{\mu}(u) < 0\} & \text{if } \gamma > p, \\ \{u \in \mathcal{N}_{\lambda}^{+} : F(u/||u||) > \mu\} & \text{if } \gamma < p. \end{cases}$$

In the next result ε is given by Proposition 2.11.

Proposition 2.12. Suppose (H1), (C1), (C2), and $\mu_* < \lambda^*$. Then $\mathcal{N}^+_{\lambda,\mu}$ is bounded for $\lambda \ge \lambda^*$ if $\gamma > p$ (respect. for $\lambda \in [\lambda^*, \lambda^* + \varepsilon]$ if $\gamma < p$).

Proof. Let $\gamma > p$. By using Proposition 2.11 (2), we can argue as in the proof of Proposition 2.5(3). Now, if $\gamma < p$ then we argue by contradiction. Assume that there exist $\lambda_n \to \lambda^*$ with $\lambda_n > \lambda^*$ and $\mathcal{N}_{\lambda_n}^+$ unbounded. So we can find a sequence (u_n) such that $u_n \in \mathcal{N}_{\lambda_n,\mu}^+$ for every n, and $||u_n|| \to \infty$. Setting $v_n = \frac{u_n}{||u_n||}$, we may assume that $v_n \rightharpoonup v$ in X. Since $t_{\lambda_n}(v_n) = ||u_n||$ we have that

$$t_{\lambda_n}(v_n) = \left(\frac{F(v_n)}{H_{\lambda_n}(v_n)}\right)^{\frac{1}{p-\gamma}} \to \infty.$$

It follows that $H_{\lambda_n}(v_n) \to 0$. On the other hand, since $F(v_n) > \mu$ we reach a contradiction as in the proof of Proposition 2.11 (1).

Let us now introduce

$$c_{\lambda,\mu}^+ := \inf_{\mathcal{N}_{\lambda,\mu}^+} \Phi_{\lambda}$$

Theorem 2.13. Suppose (H1), (C1), (C2), and $\mu_* < \lambda^*$.

(1) If $\gamma > p$ and $\lambda \ge \lambda^* > \mu > \mu_*$, then there exists $u_\lambda \in \overline{\mathcal{N}_{\lambda,\mu}^+}$ such that $\Phi_\lambda(u_\lambda) = c_{\lambda,\mu}^+ < 0$. (2) If $\gamma < p$ and $\lambda \in [\lambda^*, \lambda^* + \varepsilon]$, then there exists $u_\lambda \in \overline{\mathcal{N}_{\lambda,\mu}^+}$ such that $\Phi_\lambda(u_\lambda) = c_{\lambda,\mu}^+ < 0$.

Proof. Propositions 2.3 and 2.12 imply that $-\infty < c_{\lambda,\mu}^+ < 0$. Let $(u_n) \subset \mathcal{N}_{\lambda,\mu}^+$ be a minimizing sequence for $c_{\lambda,\mu}^+$. By Proposition 2.12 we can assume that $u_n \rightharpoonup u$ in X and it is clear that $u \neq 0$.

(1) Note that $H_{\mu}(u) \leq \liminf H_{\mu}(u_n) \leq 0$, which implies that F(u) < 0 by (2.1). Since $H_{\lambda}(u) < H_{\mu}(u) \leq 0$, by Proposition 2.3 there exists $t_{\lambda}(u) > 0$ such that $t_{\lambda}(u)u \in \mathcal{N}_{\lambda}^+$, and then $t_{\lambda}(u)u \in \overline{\mathcal{N}_{\lambda,\mu}^+}$ since $H_{\mu}(t_{\lambda}(u)u) = t_{\lambda}(u)^p H_{\mu}(u) \leq 0$. Therefore $\Phi_{\lambda}(t_{\lambda}(u)u) \leq \Phi_{\lambda}(u) \leq \liminf \Phi_{\lambda}(u_n) = c_{\lambda,\mu}^+$,

i.e. $\Phi_{\lambda}(t_{\lambda}(u)u) = c_{\lambda,\mu}^+$.

so that $\Phi_{\lambda}($

(2) Note that $F(u/||u||) \ge \limsup F(u_n/||u_n||) \ge \mu$, which implies by Proposition 2.11 that $H_{\lambda}(u/||u||) > 0$. Thus by Proposition 2.3, there exists $t_{\lambda}(u) > 0$ such that $t_{\lambda}(u)u \in \overline{\mathcal{N}^+_{\lambda,\mu}}$. Therefore

$$\Phi_{\lambda}(t_{\lambda}(u)u) \le \Phi_{\lambda}(u) \le \liminf \Phi_{\lambda}(u_n) = c_{\lambda,\mu}^+,$$

$$t_{\lambda}(u)u) = c_{\lambda,\mu}^+.$$

We introduce now the set of minimizers associated to c_{λ}^{+} , i.e.

$$\mathcal{S}_{\lambda} = \{ u \in \mathcal{N}_{\lambda}^+ : \Phi_{\lambda}(u) = c_{\lambda}^+ \}.$$

Lemma 2.14. Suppose (H1), $\mu_* < \lambda^*$, and (C1) if $\gamma > p$ (respect. (C2) if $\gamma < p$). Then S_{λ^*} is compact.

Proof. Indeed, take $(u_n) \subset S_{\lambda^*}$ and note that $\Phi'_{\lambda^*}(u_n) = 0$. Writing $u_n = t_n w_n$ with $w_n \in \Theta_{\lambda^*}$, and arguing as in the proof of Theorem 1.2, we conclude that up to a subsequence $u_n \to u \in \mathcal{N}_{\lambda^*}^+$ and the proof is complete.

Corollary 2.15. Suppose (H1), $\mu_* < \lambda^*$, and $S_{\lambda^*} \neq \emptyset$.

- (1) Assume $\gamma > p$ and (C1). Then there exists $\mu \in (\mu_*, \lambda^*)$ such that $H_{\mu}(u) < 0$ for all $u \in S_{\lambda^*}$.
- (2) Assume $\gamma < p$ and (C2). Then there exists $\mu > 0$ such that $F(u) > \mu$ for all $u \in S_{\lambda^*}$.

Proof.

- (1) By Lemma 2.14 the set S_{λ^*} is compact and $H_{\lambda^*}(u) = \frac{p\gamma}{\gamma p} c_{\lambda^*} < 0$ for all $u \in S_{\lambda^*}$. Now choose $\mu \in (\mu_*, \lambda^*)$ such that $H_{\mu}(u) < 0$ for all $u \in S_{\lambda^*}$ and the proof is complete.
- (2) Indeed, we have $F(u) = \frac{p\gamma}{\gamma p} c_{\lambda^*} > 0$ for all $u \in \mathcal{S}_{\lambda^*}$.

Next we deal with μ given by Corollary 2.15 and u_{λ} given by Theorem 2.13:

Lemma 2.16. Let $\lambda_n \searrow \lambda^*$ and $u_n \in \overline{\mathcal{N}_{\lambda_n,\mu}^+}$ be such that $\Phi_{\lambda_n}(u_n) = c_{\lambda_n,\mu}^+$. If $\gamma > p$ and $u_n \rightharpoonup u \in \mathcal{D}_{\lambda^*}^-$ (respect. $\gamma < p$ and $u_n \rightharpoonup u \in \mathcal{D}_{\lambda^*}^+$) then $u_n \rightarrow u \in \mathcal{S}_{\lambda^*}$.

Proof. We argue as in the proof of Lemma 2.8. First note that $\Phi_{\lambda^*}(u) \geq c_{\lambda^*}^+$. Suppose, by contradiction, that $\Phi_{\lambda^*}(u) > c_{\lambda^*}^+$. Recall from Theorem 1.2 that there exists $v \in S_{\lambda^*}$, i.e. $v \in \mathcal{N}_{\lambda^*}^+$ with $c_{\lambda^*}^+ = \Phi_{\lambda^*}(v)$.

Case 1: $\gamma > p$

By Corollary 2.15 we have $H_{\mu}(v) < 0$. Then $w := v/||v|| \in \Theta_{\lambda^*}$ and $H_{\mu}(w) < 0$. From Lemma 2.6(3) we know that $\Phi_{\lambda}(t_{\lambda}(w)w) < c_{\lambda^*}^+$ for all $\lambda > \lambda^*$. Since $t_{\lambda_n}(w)w \in \mathcal{N}_{\lambda_n,\mu}^+$ we find that

$$\Phi_{\lambda_n}(u_n) = c^+_{\lambda_n,\mu} \le \Phi_{\lambda_n}(t_{\lambda_n}(w)w) < c^+_{\lambda^*} < \Phi_{\lambda^*}(u),$$

which contradicts $\Phi_{\lambda^*}(u) \leq \liminf \Phi_{\lambda_n}(u_n)$. Therefore $u \in S_{\lambda^*}$, and repeating the argument above with u instead of v, we find that $\Phi_{\lambda_n}(u_n) \to \Phi_{\lambda^*}(u)$, so that $P_1(u_n) \to P_1(u)$. Condition (S2) implies that $u_n \to u$ in X. Case 2: $\gamma < p$

We have now $\limsup F(t_{\lambda_n}(w)w) = \limsup t_{\lambda_n}(w)^{\gamma}F(w) \ge F(v) > \mu$, i.e. $t_{\lambda_n}(w)w \in \mathcal{N}^+_{\lambda_n,\mu}$. From Lemma 2.6(3) we know that there exists $\varepsilon > 0$ such that $\Phi_{\lambda}(t_{\lambda}(w)w) < c^+_{\lambda^*}$ for all $\lambda \in (\lambda^*, \lambda^* + \varepsilon)$. One can argue as in the previous case to reach a contradiction.

Proof of Theorem 1.3 (1).

Let u_{λ} be given by Theorem 2.13, i.e. $u_{\lambda} \in \overline{\mathcal{N}_{\lambda,\mu}^+}$ satisfies $\Phi_{\lambda}(u_{\lambda}) = c_{\lambda,\mu}^+ < 0$. We claim that there exists $\varepsilon > 0$ such that $u_{\lambda} \in \mathcal{N}_{\lambda,\mu}^+$ for all $\lambda \in (\lambda^*, \lambda^* + \varepsilon)$, i.e. $c_{\lambda,\mu}^+$ is achieved for these values of λ .

Case 1: $\gamma > p$

Let us prove that $H_{\mu}(u_{\lambda}) < 0$ if $\lambda \in (\lambda^*, \lambda^* + \varepsilon)$, for some $\varepsilon > 0$. Indeed, suppose on the contrary, that there exists a sequence $\lambda_n \searrow \lambda^*$ such that $u_n := u_{\lambda_n}$ satisfies $H_{\mu}(u_n) = 0$ and $\Phi_{\lambda_n}(u_n) = c_{\lambda_n,\mu}^+$. As in the proof of Theorem 1.2(2), we can show that writing $u_n = t_n w_n$, where $w_n \in \Theta_{\lambda_n}$ and $t_n := t_{\lambda_n}(w_n)$, up to a subsequence $u_n \rightharpoonup u \in \mathcal{D}_{\lambda^*}^-$. Lemma 2.16 yields that $u_n \rightarrow u \in S_{\lambda^*}$. It follows from Corollary 2.15 that $H_{\mu}(u) < 0$. However, this is a contradiction, since $H_{\mu}(u) = \lim H_{\mu}(u_n) = 0$.

Case 2: $\gamma < p$

Arguing by contradiction as in the previous case, we find that $u_n \to u \in S_{\lambda^*}$ and by Corollary 2.15 it follows that $F(u) > \mu$. However, this is a contradiction, since $F(u) = \lim F(u_n) = \mu$. Thus the existence of ε is guaranteed and the proof is complete.

2.5. A mountain-pass critical point for $\lambda > \lambda^*$. All over this section we assume (H1), (S), (C1), (C2), and $\mu_* < \lambda^*$. Recall that

$$\Phi_{\lambda}(u_{\lambda}) = c_{\lambda,\mu}^{+} := \inf_{\mathcal{N}_{\lambda,\mu}^{+}} \Phi_{\lambda},$$

and

$$\mathcal{N}_{\lambda,\mu}^{+} = \begin{cases} \{u \in \mathcal{N}_{\lambda}^{+} : H_{\mu}(u) < 0\} & \text{if } \gamma > p, \\ \{u \in \mathcal{N}_{\lambda}^{+} : F(u) > \mu\} & \text{if } \gamma < p. \end{cases}$$

for $\lambda \in (\lambda^*, \lambda^* + \varepsilon)$. The next result follows from the proof of Theorem 1.3(1):

Corollary 2.17. For each $\lambda \in (\lambda^*, \lambda^* + \varepsilon)$ there holds $\inf_{\partial \mathcal{N}^+_{\lambda,\mu}} \Phi_{\lambda} > c^+_{\lambda,\mu}$.

Next we deal with the set

$$B_{\delta} := \{ tu : u \in \overline{\mathcal{N}_{\lambda,\mu}^+}, \ t \in (1 - \delta, 1 + \delta) \},\$$

defined for $\delta > 0$.

Proposition 2.18. For each $\lambda \in (\lambda^*, \lambda^* + \varepsilon)$ there exists $\delta > 0$ such that $\inf_{\partial B_{\delta}} \Phi_{\lambda} > c_{\lambda,\mu}^+$.

Proof. Indeed, first we claim that there exists c > 0 such that $\varphi_{\lambda,u}''(1) > c$ for all $u \in \mathcal{N}_{\lambda,\mu}^+$. Since $\overline{\mathcal{N}_{\lambda,\mu}^+} \subset \mathcal{N}_{\lambda}^+$ it is clear that $\varphi_{\lambda,u}''(1) > 0$ for $u \in \overline{\mathcal{N}_{\lambda,\mu}^+}$.

Case 1: $\gamma > p$

Suppose, on the contrary, that there exists $(u_n) \subset \overline{\mathcal{N}_{\lambda,\mu}^+}$ such that $\varphi_{\lambda,u_n}'(1) \to 0$. Since $\varphi_{\lambda,u_n}'(1) = 0$ for all n, it follows that $H_{\lambda}(u_n) \to 0$. We write $u_n = t_n w_n$ where $w_n \in \Theta_{\lambda}$ and

$$t_n := t_\lambda(w_n) = \left(\frac{H_\lambda(w_n)}{F(w_n)}\right)^{\frac{1}{\gamma-p}}.$$
(2.13)

Thus $t_n^p H_{\lambda}(w_n) = H_{\lambda}(u_n) \to 0$ which, combined with (2.13), implies that $H_{\lambda}(w_n) \to 0$. Therefore, we can assume that $w_n \rightharpoonup w \neq 0$ and moreover

$$H_{\mu}(w_n) = H_{\lambda}(w_n) + (\lambda - \mu)P_2(w_n)$$

which implies that $H_{\mu}(w_n) > 0$ for *n* sufficiently large. This yields a contradiction, since $H_{\mu}(w_n) = t_n^{-p} H_{\mu}(u_n) \leq 0$ for all *n*.

Case 2: $\gamma < p$

It is enough to note that for $u \in \overline{\mathcal{N}^+_{\lambda_n,\mu}}$ we have $\varphi''_{\lambda,u}(1) = (p-\gamma)F(u) \ge (p-\gamma)\mu$.

Thus the claim is proved and there exists c > 0 such that $\varphi_{\lambda,u}''(1) > c$ for all $u \in \overline{\mathcal{N}_{\lambda,\mu}^+}$. This implies in particular that there exists c > 0 such that $||u|| \ge c$ for all $u \in \overline{\mathcal{N}_{\lambda,\mu}^+}$. By making c smaller if necessary, we can choose $\delta > 0$ such that $\varphi_{\lambda,w}''(1) > c$ for all $w \in \overline{B_{\delta}}$. Therefore, for any $tu \in \overline{B_{\delta}}$ there holds

$$\begin{split} \Phi_{\lambda}(tu) - \Phi_{\lambda}(u) &= \varphi_{\lambda,u}(t) - \varphi_{\lambda,u}(1) = \varphi_{\lambda,u}'(1)(t-1) + \frac{1}{2}\varphi_{\lambda,u}''(\theta)(t-1)^2 \\ &= \frac{1}{2}\varphi_{\lambda,u}''(\theta)(t-1)^2 > \frac{c}{2}(t-1)^2, \end{split}$$

where $\theta \in (\min\{1, t\}, \max\{1, t\})$. Now observe that

$$\partial B_{\delta} = \{ tu : u \in \partial \mathcal{N}_{\lambda,\mu}^{+}, \ t \in [-\delta, \delta] \} \cup \{ su : u \in \mathcal{N}_{\lambda,\mu}^{+}, \ s \in \{-\delta, \delta\} \}.$$

From Corollary 2.17 we have, for $t \in [-\delta, \delta]$ and $u \in \partial \mathcal{N}^+_{\lambda, u}$, that

$$\Phi_{\lambda}(tu) > \Phi_{\lambda}(u) + \frac{c}{2}(t-1)^2 > c^+_{\lambda,\mu}$$

On the other hand, if $u \in \mathcal{N}_{\lambda,\mu}^+$ and $s \in \{-\delta, \delta\}$, then

$$\Phi_{\lambda}(su) > \Phi_{\lambda}(u) + \frac{c}{2}(\pm\delta - 1)^2 \ge c_{\lambda,\mu}^+ + \frac{c}{2}(\pm\delta - 1)^2,$$

and the proof is complete.

Proof of Theorem 1.3 (2).

Let $\delta > 0$ be given by Proposition 2.18. Since B_{δ} is bounded and Φ_{λ} is unbounded from below, we can find some $v_{\lambda} \in X \setminus B_{\delta}$ such that $\Phi_{\lambda}(v_{\lambda}) < c_{\lambda}$.

Therefore, setting

$$\Gamma_{\lambda} = \{\eta \in C([0,1], X) : \eta(0) = u_{\lambda}, \ \eta(1) = v_{\lambda}\}$$

we infer that

$$d_{\lambda} = \inf_{\eta \in \Gamma_{\lambda}} \max_{t \in [0,1]} \Phi_{\lambda}(\eta(t))$$

is a critical value of Φ_{λ} for $\lambda \in (\lambda^*, \lambda^* + \varepsilon)$.

3. Applications

Let us provide some applications of Theorems 1.1, 1.2 and 1.3. Throughout this section Ω is a bounded domain of \mathbb{R}^N , with $N \geq 1$.

3.1. Indefinite *p*-Laplace equations. We consider the functional associated to (1.3), i.e.

$$\Phi_{\lambda}(u) = \frac{1}{p} \int_{\Omega} \left(|\nabla u|^p - \lambda(u^+)^p \right) - \frac{1}{\gamma} \int_{\Omega} f(x)(u^+)^{\gamma}, \quad u \in X = W_0^{1,p}(\Omega),$$

and some variations of it. Here p > 1, $1 < \gamma < p^*$ with $\gamma \neq p$, and $f \in L^{\infty}(\Omega)$. Recall that $p^* = \frac{Np}{N-p}$ if p < N and $p^* = \infty$ if $p \ge N$.

Let $H_{\lambda}(u) = \int_{\Omega} (|\nabla u|^p - \lambda(u^+)^p)$, i.e. $P_1(u) = \int_{\Omega} |\nabla u|^p = ||u||^p$, $P_2(u) = \int_{\Omega} (u^+)^p$, and $F(u) = \int_{\Omega} f(x)(u^+)^\gamma$ for $u \in X$. It is standard to check that P_1 , P_2 and F satisfy our basic assumptions. Note also that critical points of Φ_{λ} are nonnegative weak solutions of (1.3). For this functional we have

$$\lambda^* := \inf \left\{ \frac{\int_{\Omega} |\nabla u|^p}{\int_{\Omega} (u^+)^p} : u \in W_0^{1,p}(\Omega) \setminus \{0\}, \int_{\Omega} f(x) (u^+)^{\gamma} = 0 \right\}.$$

It is clear that

$$\lambda^* \ge \inf\left\{\frac{\int_{\Omega} |\nabla u|^p}{\int_{\Omega} (u^+)^p} : u \in W_0^{1,p}(\Omega)\right\} = \inf\left\{\frac{\int_{\Omega} |\nabla u|^p}{\int_{\Omega} |u|^p} : u \in W_0^{1,p}(\Omega)\right\} = \lambda_1(p)$$

the first eigenvalue of $-\Delta_p$ in $W_0^{1,p}(\Omega)$. Indeed, the equality of the infima above follows from the fact that $\lambda_1(p)$ is achieved by a positive eigenfunction $\varphi_1 = \varphi_1(p)$, and the inequality $\int_{\Omega} |\nabla u^+|^p \leq \int_{\Omega} |\nabla u|^p$ for any $u \in X$.

We shall assume that $\int_{\Omega} f(x)\varphi_1^{\gamma} < 0$, which clearly yields $\mu_* = \lambda_1(p) < \lambda^*$. Let us show that (H1) also holds. Indeed, otherwise we would have

$$\lambda^* > \Lambda := \inf \left\{ \frac{\int_{\Omega} |\nabla u|^p}{\int_{\Omega} (u^+)^p} : u \in W_0^{1,p}(\Omega) \setminus \{0\}, \int_{\Omega} f(x)(u^+)^{\gamma} \ge 0 \right\},$$

so Λ would be achieved by some u_0 such that $\int_{\Omega} f(x)(u_0^+)^{\gamma} > 0$. In particular, u_0 minimizes $\frac{\int_{\Omega} |\nabla u|^p}{\int_{\Omega} (u^+)^p}$ over the open set $\{u \in W_0^{1,p}(\Omega) : \int_{\Omega} f(x)(u^+)^{\gamma} > 0\}$. Thus u_0 would be a critical point of $\frac{\int_{\Omega} |\nabla u|^p}{\int_{\Omega} (u^+)^p}$, and consequently a nonnegative eigenfunction of $-\Delta_p$ associated to the eigenvalue Λ . However, the assumption $\int_{\Omega} f(x)\varphi_1^{\gamma} < 0$ entails that $\Lambda > \lambda_1$. Finally, it is known that λ_1 is the only principal eigenvalue of $-\Delta_p$, i.e. it is the only eigenvalue associated to a nonnegative eigenfunction. So we reach a contradiction, and (H1) is proved.

The latter argument also shows that the condition $\int_{\Omega} f(x)\varphi_1^{\gamma} < 0$ implies (C2), since $H'_{\lambda^*}(u) = 0$ and $u^+ \neq 0$ means that λ^* is a principal eigenvalue of $-\Delta_p$, i.e. $\lambda^* = \lambda_1(p)$, which is impossible.

It is also clear that (S) holds since $P_1(u) = ||u||^p$ and $W_0^{1,p}(\Omega)$ is a uniformly convex space.

As for the (PS) condition, it suffices to show that any (PS) sequence is bounded (the (S+) property of the *p*-Laplacian gives then the desired conclusion). To this end, let us introduce the notation

 $\Omega^{0} = \{ x \in \Omega : f(x) = 0 \} \text{ and } \Omega^{+} = \{ x \in \Omega : f(x) > 0 \}.$

More precisely, Ω_+ is the largest open set where f > 0 a.e. We denote by $int(\Omega^0)$ the interior of Ω^0 and similarly we define $int(\Omega^0 \cup \Omega^+)$. Given an open, bounded and smooth set U, we denote by $(\lambda_1(p, U), \phi_1(U))$ the first eigenpair of $(-\Delta_p, U)$. We shall assume that $int(\Omega^0)$ is smooth, so that the following property holds: (h₀) If $v \in W_0^{1,p}(\Omega)$ and $fv \equiv 0$ then $v \in W_0^{1,p}(\operatorname{int}(\Omega^0))$.

This property holds, for instance, if $int(\Omega^0)$ is a *p*-stable set, in the capacity sense (see e.g. [15, Proposition 11]).

For $\gamma > p$ we shall prove that (PS) holds for $\lambda < \lambda_1(p, \operatorname{int}(\Omega^0))$. Indeed, let $(u_n) \subset X$ be such that $(\Phi_\lambda(u_n))$ is bounded and $\Phi'_\lambda(u_n) \to 0$. Assume by contradiction that $||u_n|| \to \infty$ and $v_n := \frac{u_n}{||u_n||} \rightharpoonup v$ in X. Since $\Phi'_\lambda(u_n)\phi \to 0$ we find that $\int_\Omega f(v^+)^{\gamma-1}\phi = \lim \int_\Omega f(v_n^+)^{\gamma-1}\phi = 0$ for every $\phi \in X$. It follows that $fv^+ \equiv 0$ so $v^+ \in W_0^{1,p}(\operatorname{int}(\Omega^0))$. Moreover, combining the fact that $(\Phi_\lambda(u_n))$ is bounded and $|\Phi'_\lambda(u_n)u_n| \leq \varepsilon_n ||u_n||$ with $\varepsilon_n \to 0$, we derive that $H_\lambda(v_n) \to 0$, so $v^+ \neq 0$ and $H_\lambda(v) \leq 0$. It follows that $\lambda \geq \lambda_1(p, \operatorname{int}(\Omega^0))$, a contradiction.

Now, if $\gamma < p$ then we have now $H'_{\lambda}(v)\phi = \lim H'_{\lambda}(v_n)\phi = 0$ for every $\phi \in X$. From the boundedness of $(\Phi^+_{\lambda}(u_n))$ it follows that $H_{\lambda}(v_n) \to 0$, which yields $v^+ \neq 0$. Thus $-\Delta_p v = \lambda(v^+)^{p-1}$ so that $v \ge 0$, and therefore $\lambda = \lambda_1(p)$ and v is an eigenfunction associated to $\lambda_1(p)$. Now, the fact that $(\Phi_{\lambda}(u_n))$ is bounded and $|\Phi'_{\lambda}(u_n)u_n| \le \varepsilon_n ||u_n||$ with $\varepsilon_n \to 0$, yields that $\int_{\Omega} f(v^+)^{\gamma} = \lim \int_{\Omega} f(v_n^+)^{\gamma} = 0$, which contradicts the assumption $\int_{\Omega} f(x)\varphi_1^{\gamma} < 0$. Therefore in this case (PS) holds for $\lambda \neq \lambda_1(p)$.

Finally we shall prove that (C1) holds under the following additional condition:

 (f_0) If $|\Omega^0| > 0$ then $int(\Omega^0)$ is an open, bounded and smooth set, satisfying

$$\lambda^* < \lambda_1(p, \operatorname{int}(\Omega^0)).$$

Proposition 3.1. Assume (h_0) and (f_0) . Then (C1) holds true.

Proof. Indeed, suppose on the contrary that F'(u) = 0 and u achieves λ^* . It follows that $f(u^+)^{\gamma-1} \equiv 0$ and hence $u^+ \in W_0^{1,p}(\operatorname{int}(\Omega^0))$ by (h_0) . It follows that

$$\lambda_1(p, \operatorname{int}(\Omega^0)) \le \frac{\int_{\operatorname{int}(\Omega^0)} |\nabla u^+|^p}{\int_{\operatorname{int}(\Omega^0)} (u^+)^p} \le \frac{\int_{\operatorname{int}(\Omega^0)} |\nabla u|^p}{\int_{\operatorname{int}(\Omega^0)} (u^+)^p} = \lambda^*,$$
(3.1)

which contradicts (f_0) . Therefore $F'(u) \neq 0$.

Let us discuss on the condition (f_0) . It is clear that

 $\lambda^* \leq \lambda_1(p, \operatorname{int}(\Omega^0)).$

We shall provide some conditions that ensure the inequality.

Corollary 3.2. Suppose that $int(\Omega^0)$ and $int(\overline{\Omega^0} \cup \overline{\Omega^+})$ are bounded and smooth sets. If (h_0) holds, and

$$\lambda_1(p, \operatorname{int}(\overline{\Omega^0} \cup \overline{\Omega^+})) < \lambda_1(p, \operatorname{int}(\Omega^0)),$$

then (C1) holds true.

Proof. Indeed, it is clear that

$$\lambda^* \le \lambda_1(p, \operatorname{int}(\overline{\Omega^0} \cup \overline{\Omega^+})),$$

and hence (f_0) is satisfied, which implies, by Proposition 3.1, the condition (C1).

Corollary 3.3. Suppose that $\operatorname{int}(\Omega^0)$ and $\operatorname{int}(\overline{\Omega^0} \cup \overline{\Omega^+})$ are bounded smooth domains. If (h_0) holds and $\operatorname{int}(\Omega^0)$ is a proper subset of $\operatorname{int}(\overline{\Omega^0} \cup \overline{\Omega^+})$, then (C1) holds true. In particular, if there exists a ball $B \subset \operatorname{int}(\overline{\Omega^0} \cup \overline{\Omega^+})$ such that $B \cap \Omega^0 \neq \emptyset$ and $B \cap \Omega^+ \neq \emptyset$, then (C1) holds true.

Proof. Indeed, if $\operatorname{int}(\Omega^0)$ is a proper subset of $\operatorname{int}(\overline{\Omega^0} \cup \overline{\Omega^+})$, then (see e.g. [14, Proposition 4.4]) $\lambda_1(p, \operatorname{int}(\overline{\Omega^0} \cup \overline{\Omega^+})) < \lambda_1(p, \operatorname{int}(\Omega^0)),$

and from Corollary 3.2 we obtain (C1). To conclude, it is clear that if such a ball exists, then $int(\Omega^0)$ is a proper subset of $int(\overline{\Omega^0} \cup \overline{\Omega^+})$.

Remark 3.4. It is worth pointing out that if $\gamma > p$, $f \leq 0$ and $|\Omega_0| > 0$ then one may still consider $c_{\lambda^*}^+$. However, condition (C1) fails in this case, since F(u) = 0 clearly implies F'(u) = 0. This fact is not a technical issue, since one can show that Φ_{λ} has no nontrivial critical point for $\lambda \geq \lambda_1(p, \operatorname{int}(\Omega^0))$.

Thus we infer the following result:

Corollary 3.5. Let p > 1 and $\gamma \in (1, p^*)$ with $\gamma \neq p$. Assume that $f \in L^{\infty}(\Omega)$ satisfies $(h_0), (f_0), and \int_{\Omega} f(x)\varphi_1^{\gamma} < 0$. Then the conclusions of Theorems 1.1, 1.2 and 1.3 hold true.

The previous result has been established for $\gamma > p$ in [20], whereas for $\gamma < p$ it extends the analysis carried out in [10] for p = 2 and in [23] for p > 1, both dealing with $\lambda < \lambda^*$.

Let us consider now the Neumann problem

$$-\Delta_p u = \lambda |u|^{p-2} u + f(x)|u|^{\gamma-2} u, \quad u \in W^{1,p}(\Omega),$$
(3.2)

In comparison with the functional of the Dirichlet problem, a slight modification is needed. Namely, we set

$$\Phi_{\lambda}(u) = \frac{1}{p} \int_{\Omega} \left(|\nabla u|^{p} + (u^{-})^{p} - \lambda (u^{+})^{p} \right) - \frac{1}{\gamma} \int_{\Omega} f(x) (u^{+})^{\gamma},$$
(3.3)

for $u \in W^{1,p}(\Omega)$. It is clear that critical poins of this functional are nonnegative and thus solutions of (3.2). We have then

$$\lambda^* := \inf\left\{\frac{\int_{\Omega} |\nabla u|^p + (u^-)^p}{\int_{\Omega} (u^+)^p} : u \in W^{1,p}(\Omega) \setminus \{0\}, \int_{\Omega} f(x)(u^+)^{\gamma} = 0\right\}.$$

As in the previous problem, one can show that

$$\lambda^* \ge \inf\left\{\frac{\int_{\Omega} |\nabla u|^p + (u^-)^p}{\int_{\Omega} (u^+)^p} : u \in W^{1,p}(\Omega)\right\} = \inf\left\{\frac{\int_{\Omega} |\nabla u|^p}{\int_{\Omega} |u|^p} : u \in W^{1,p}(\Omega)\right\} = 0,$$

and the inequality holds if $\int_{\Omega} f < 0$. This condition also yields $\mu_* = 0 < \lambda^*$, (H1), and (C2). It is also clear that (S) is satisfied and proceeding as in the Dirichlet case one can show that (PS) holds for $\lambda \neq 0$. Finally, we note that (f_0) is weaker than in the Dirichlet case, since the infimum in the definition of λ^* is taken over $W^{1,p}(\Omega)$ whereas $\lambda_1(p, \operatorname{int}(\Omega^0))$ remains unchanged.

A similar analysis applies to the functionals

$$\Phi^{1}_{\lambda}(u) = \frac{1}{p} \int_{\Omega} \left(|\nabla u|^{p} + (u^{-})^{p} - \lambda(u^{+})^{p} \right) - \frac{1}{\gamma} \int_{\partial\Omega} f(x)(u^{+})^{\gamma}, \tag{3.4}$$

and

$$\Phi_{\lambda}^{2}(u) = \frac{1}{p} \left(\int_{\Omega} |\nabla u|^{p} + (u^{-})^{p} - \lambda \int_{\partial \Omega} (u^{+})^{p} \right) - \frac{1}{\gamma} \int_{\Omega} f(x)(u^{+})^{\gamma},$$
(3.5)

defined in $X = W^{1,p}(\Omega)$. These functionals are respectively associated to the problems

$$\begin{cases} -\Delta_p u = \lambda |u|^{p-2} u & \text{in } \Omega, \\ |\nabla u|^{p-2} \partial_n u = f(x)|u|^{\gamma-2} u & \text{on } \partial\Omega. \end{cases} \text{ and } \begin{cases} -\Delta_p u = f(x)|u|^{\gamma-2} u & \text{in } \Omega, \\ |\nabla u|^{p-2} \partial_n u = \lambda |u|^{p-2} u & \text{on } \partial\Omega. \end{cases}$$

We refer to [1, 12, 21, 22] for previous results on (3.2), (3.4), and (3.5) with $1 < \gamma < p = 2$.

3.2. (p,q)-Laplacian problems. We consider now the functional

$$\Phi_{\lambda}(u) = \frac{1}{p} \int_{\Omega} \left(|\nabla u|^p - \lambda (u^+)^p \right) + \frac{1}{q} \int_{\Omega} \left(|\nabla u|^q - \beta(x)(u^+)^q \right) \quad u \in W_0^{1,p}(\Omega),$$
(3.6)

whose critical points are nonnegative solutions of the (p, q)-Laplacian problem

$$-\Delta_p u - \Delta_q u = \lambda |u|^{p-2} u + \beta(x) |u|^{q-2} u, \quad u \in W_0^{1,p}(\Omega),$$

$$(3.7)$$

where 1 < q < p, $\lambda \in \mathbb{R}$ and $\beta \in L^{\infty}(\Omega)$ is nonnegative and nontrivial. This problem, with β constant, has been recently studied in [7–9].

Here $H_{\lambda}(u) = \int_{\Omega} (|\nabla u|^p - \lambda(u^+)^p)$, whereas F is given now by

$$F(u) = -\int_{\Omega} \left(|\nabla u|^q - \beta(x)(u^+)^q \right),$$

so that

$$\lambda^* = \lambda^*(\beta) := \inf \left\{ \frac{\int_{\Omega} |\nabla u|^p}{\int_{\Omega} (u^+)^p} : u \in W_0^{1,p}(\Omega) \setminus \{0\}, \int_{\Omega} \left(|\nabla u|^q - \beta(x)(u^+)^q \right) = 0 \right\}.$$

We still have $\lambda^* \geq \lambda_1(p)$, and the inequality holds if, and only if, $F(\varphi_p) < 0$, i.e. $\int_{\Omega} (|\nabla \varphi_p|^q - \beta(x)\varphi_p^q) > 0$, where $\varphi_p := \varphi_1(p)$. Arguing as in the previous subsection, we can show that this condition implies (H1) and (C2).

To check (C1), assume by contradiction that λ^* is achieved by some u such that F'(u) = 0. It follows that $\int_{\Omega} |\nabla u|^{q-2} \nabla u \nabla \phi = \int_{\Omega} \beta(x) (u^+)^{q-1} \phi$ for every $\phi \in X$. which yields $u \ge 0$. Moreover, even though this equation holds in $(W_0^{1,p}(\Omega))^*$, one can show that $\lambda_1(\beta, q) = 1$, where

$$\lambda_1(\beta,q) := \inf \left\{ \frac{\int_{\Omega} |\nabla u|^q}{\int_{\Omega} \beta(x) |u|^q} : u \in W_0^{1,q}(\Omega) \right\}.$$

Thus (C1) holds if $\lambda_1(\beta, q) \neq 1$. Note also that F takes positive values if, and only if, $\lambda_1(\beta, q) < 1$. Finally, proceeding as in the previous subsection (in the case $\gamma < p$), one can show that (PS) holds for any $\lambda \neq \lambda_1(p)$ if $F(\varphi_p) < 0$. Summing up, we derive the following result:

Corollary 3.6. Let 1 < q < p, $\lambda \in \mathbb{R}$ and $\beta \in L^{\infty}(\Omega)$ be nonnegative with $\lambda_1(\beta, q) < 1$.

- (1) If $\lambda < \lambda_1(p)$ then c_{λ}^+ is achieved, i.e. (3.7) has a nonnegative solution $u_+ \in \mathcal{N}_{\lambda}^+$.
- (2) If $\int_{\Omega} (|\nabla \varphi_p|^q \beta(x) \varphi_p^q) > 0$ then $\lambda^* > \lambda_1(p)$, and (3.7) has:
 - (a) one nonnnegative solution $u_+ \in \mathcal{N}^+_{\lambda}$ for $\lambda = \lambda_1(p)$.
 - (b) two nonnegative solutions $u_+ \in \mathcal{N}^+_{\lambda}$ and $u_- \in \mathcal{N}^-_{\lambda}$ for $\lambda_1(p) < \lambda < \lambda^*$. Moreover, there exists $\varepsilon > 0$ such that (3.7) has two nonnegative solutions for $\lambda^* \leq \lambda < \lambda^* + \varepsilon$.

The previous result extends [8, Theorem 2.7], which deals with β constant. In this case, the condition $\lambda_1(\beta, q) < 1$ reads as $\beta > \lambda_1(q)$, whereas $\int_{\Omega} (|\nabla \varphi_p|^q - \beta(x)\varphi_p^q) > 0$ becomes now $\beta < \beta_* := \frac{\int_{\Omega} |\nabla \varphi_p|^q}{\int_{\Omega} \varphi_p^q}$. Let us note that in [8, Theorem 2.7] the roles of λ and β are interchanged (see [18, Section 3.2] for more details).

Similarly to the problem of the previous section, we may also consider the functional

$$\Phi_{\lambda}(u) = \frac{1}{p} \int_{\Omega} \left(|\nabla u|^p + (u^-)^p - \lambda(u^+)^p \right) + \frac{1}{q} \int_{\Omega} \left(|\nabla u|^q - \beta(x)(u^+)^q \right) \quad u \in W_0^{1,p}(\Omega), \quad (3.8)$$

on $X = W^{1,p}(\Omega)$. In this case, we have $\lambda_1(p) = 0$ and φ_p is a positive constant. Hence the condition $\int_{\Omega} (|\nabla \varphi_p|^q - \beta(x)\varphi_p^q) > 0$ reads $\int_{\Omega} \beta < 0$, so that we need $\beta^- \neq 0$. On the other

hand, F take positive values only if $\beta^+ \neq 0$. Thus β has to change sign. We derive then the following result on the problem

$$-\Delta_p u - \Delta_q u = \lambda |u|^{p-2} u + \beta(x) |u|^{q-2} u, \quad u \in W^{1,p}(\Omega).$$

$$(3.9)$$

Corollary 3.7. Let 1 < q < p, $\lambda \in \mathbb{R}$ and $\beta \in L^{\infty}(\Omega)$ be sign-changing and such that

$$\inf\left\{\frac{\int_{\Omega}|\nabla u|^{q}}{\int_{\Omega}\beta(x)|u|^{q}}: u \in W^{1,q}(\Omega), \int_{\Omega}\beta(x)|u|^{q} > 0\right\} < 1$$

- (1) If $\lambda < 0$ then c_{λ}^+ is achieved, i.e. (3.9) has a nonnegative solution $u_+ \in \mathcal{N}_{\lambda}^+$.
- (2) If $\int_{\Omega} \beta < 0$ then $\lambda^* > 0$, and (3.9) has:
 - (a) one nonnnegative solution $u_+ \in \mathcal{N}^+_{\lambda}$ for $\lambda = 0$.
 - (b) two nonnegative solutions $u_+ \in \mathcal{N}^+_{\lambda}$ and $u_- \in \mathcal{N}^-_{\lambda}$ for $0 < \lambda < \lambda^*$. Moreover, there exists $\varepsilon > 0$ such that (3.9) has two nonnegative solutions for $\lambda^* \le \lambda < \lambda^* + \varepsilon$.

Remark 3.8. As mentioned in Remark 2.10, we have $c_{\lambda}^{+} = -\infty$ if $\gamma < p$ and $c_{\lambda}^{-} = 0$ if $\gamma > p$ and $\lambda > \lambda^{*}$, for the functional of Section 3.1, and $c_{\lambda}^{+} = -\infty$ if $\lambda > \lambda^{*}$, for the functional of Section 3.2. Indeed, in the first case one may argue as in the proof of [10, Lemma 4] to show that $c_{\lambda}^{+} = -\infty$ for $\lambda > \lambda^{*}$ if $\gamma < p$. To this end, it suffices to prove that if $\lambda > \lambda^{*}$, then there exists u such that $H_{\lambda}(w) < 0$ and F(w) > 0. We choose $u \neq 0$ such that $H_{\lambda^{*}}(u) = F(u) = 0$. It is clear that $H_{\lambda}(u) < 0$. Choose r > 0 such that $H_{\lambda}(w) < 0$ for all $w \in B$, where B is the ball centered at u, with radius r. If $F(w) \leq 0$ for all $w \in B$, then u is a local maximizer of F, so that F'(u) = 0, which contradicts (C1). Therefore there exists w such that $H_{\lambda}(w) < 0$ and F(w) > 0 which completes the proof. For the functional of Section 3.2 we refer to [8, Theorem 2.5(ii)].

3.3. Kirchhoff type problems. We consider now the functional

$$\Phi_{\lambda}(u) = \frac{1}{2} \int_{\Omega} \left(a |\nabla u|^2 - \lambda (u^+)^2 \right) + \frac{b}{4} \left(\int_{\Omega} |\nabla u|^2 \right)^2 - \frac{1}{4} \int_{\Omega} \beta(x) (u^+)^4 \quad u \in W_0^{1,2}(\Omega), \quad (3.10)$$

whose critical points are nonnegative solutions of the Kirchhoff type problem

$$-\left(a+b\int_{\Omega}|\nabla u|^{2}\right)\Delta u = \lambda|u|^{p-2}u+\beta(x)|u|^{q-2}u, \quad u \in H_{0}^{1}(\Omega).$$

$$(3.11)$$

Here $a, b > 0, \lambda \in \mathbb{R}$ and $\beta \in L^{\infty}(\Omega)$ is nonnegative and nontrivial, and $N \leq 3$. This problem, with β constant, has been recently investigated in [13, 25].

Here $H_{\lambda}(u) = \int_{\Omega} \left(a |\nabla u|^2 - \lambda (u^+)^2 \right)$, whereas $F(u) = -b \left(\int_{\Omega} |\nabla u|^2 \right)^2 + \int_{\Omega} \beta(x) (u^+)^4$, so that

$$\lambda^* = \lambda^*(\beta) := \inf\left\{\frac{\int_{\Omega} |\nabla u|^2}{\int_{\Omega} (u^+)^2} : u \in H_0^1(\Omega) \setminus \{0\}, b\left(\int_{\Omega} |\nabla u|^2\right)^2 - \int_{\Omega} \beta(x)(u^+)^4 = 0\right\}.$$

We still have $\lambda^* \geq \lambda_1 := \lambda_1(2)$, and the inequality holds if, and only if, $F(\varphi_1) < 0$, i.e. $b \left(\int_{\Omega} |\nabla \varphi_1|^2 \right)^2 - \int_{\Omega} \beta(x) (\varphi_1^+)^4 > 0$, where $\varphi_1 := \varphi_1(2)$. Arguing as in the previous subsection, we can show that this condition implies (H1) and (C2).

To check (C1), assume by contradiction that λ^* is achieved by some u such that F'(u) = 0. It follows that $b \int_{\Omega} |\nabla u|^2 \int_{\Omega} \nabla u \nabla \phi = \int_{\Omega} \beta(x) (u^+)^3 \phi$ for every $\phi \in X$, which yields $u \ge 0$. Therefore $\lambda_1(\beta) = 1$, where

$$\lambda_1(\beta) := \inf \left\{ \frac{b \left(\int_{\Omega} |\nabla u|^2 \right)^2}{\int_{\Omega} \beta(x) |u|^4} : u \in H_0^1(\Omega) \right\}.$$

Thus (C1) holds if $\lambda_1(\beta) \neq 1$. Note also that F takes positive values if, and only if, $\lambda_1(\beta) < 1$. Finally, proceeding as in the previous subsection (in the case $\gamma < p$), one can show that (PS) holds for any $\lambda \neq \lambda_1$ if $F(\varphi_1) < 0$. Summing up, we derive the following result:

Corollary 3.9. Let $\lambda \in \mathbb{R}$ and $\beta \in L^{\infty}(\Omega)$ be nonnegative with $\lambda_1(\beta) < 1$.

- (1) If $\lambda < \lambda_1$ then c_{λ}^+ is achieved, i.e. (3.12) has a nonnegative solution $u_+ \in \mathcal{N}_{\lambda}^+$. (2) If $b \left(\int_{\Omega} |\nabla \varphi_1|^2 \right)^2 \int_{\Omega} \beta(x) (\varphi_1^+)^4 > 0$ then $\lambda^* > \lambda_1$, and (3.12) has: (a) one nonnnegative solution $u_+ \in \mathcal{N}_{\lambda}^+$ for $\lambda = \lambda_1$.

 - (b) two nonnegative solutions $u_+ \in \mathcal{N}_{\lambda}^+$ and $u_- \in \mathcal{N}_{\lambda}^-$ for $\lambda_1 < \lambda < \lambda^*$. Moreover, there exists $\varepsilon > 0$ such that (3.12) has two nonnegative solutions for $\lambda^* \leq \lambda < \varepsilon$ $\lambda^* + \varepsilon$.

The previous result extends [25, Theorem 2], which deals with β constant. In this case, the condition $\lambda_1(\beta) < 1$ reads as $\beta > b\mu_1$, where μ_1 is defined by

$$\mu_1 = \inf \left\{ \frac{b \left(\int_{\Omega} |\nabla u|^2 \right)^2}{\int_{\Omega} |u|^4} : u \in H_0^1(\Omega) \right\}.$$

Let us observe that μ_1 is the first eigenvalue of the problem

$$-b\left(\int_{\Omega} |\nabla u|^2\right) \Delta u = \lambda |u|^2 u, \quad u \in H_0^1(\Omega), \tag{3.12}$$

see [16]. Moreover $b\left(\int_{\Omega} |\nabla \varphi_1|^2\right)^2 - \int_{\Omega} \beta(x)(\varphi_1^+)^4 > 0$ becomes now $\beta < \beta_* := \frac{b\left(\int_{\Omega} |\nabla \varphi_1|^2\right)^2}{\int_{\Omega} \varphi_1^4}$.

Acknowledgements. The authors are thankful to V. Bobkov for pointing out an error in a former version of Proposition 2.11(1).

Data availability. Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

References

- [1] S. Alama, Semilinear elliptic equations with sublinear indefinite nonlinearities, Adv. Differential Equations 4 (1999), 813-842. 19
- [2] S. Alama, G. Tarantello, On semilinear elliptic equations with indefinite nonlinearities, Calc. Var. Partial Differential Equations 1 (1993), 439–475. 2
- [3] J. C. de Albuquerque, K. Silva, On the extreme value of the Nehari manifold method for a class of Schrödinger equations with indefinite weight functions. J. Differential Equations 269 (2020), no. 7, 5680-5700. 1
- [4] A. Ambrosetti, Critical points and nonlinear variational problems. Mémoires de la S. M. F. 2e série, tome 49 (1992). 3
- [5] P. A. Binding, P. Drábek, Y. X. Huang, On Neumann boundary value problems for some quasilinear elliptic equations. Electron. J. Differential Equations (1997), No. 05, approx. 11 pp. 3
- [6] I. Birindelli, F. Demengel, Existence of solutions for semi-linear equations involving the p-Laplacian: the non coercive case. Calc. Var. Partial Differential Equations 20 (2004), 343–366. 2
- [7] V. Bobkov, M. Tanaka, On positive solutions for (p, q)-Laplace equations with two parameters. Calculus of Variations and Partial Differential Equations, 54(3), 3277-3301. 20
- [8] V. Bobkov, M. Tanaka, Remarks on minimizers for (p,q)-Laplace equations with two parameters. Communications on Pure and Applied Analysis, 17(3), (2018) 1219-1253. 20, 21
- [9] V. Bobkov, M. Tanaka, Multiplicity of positive solutions for (p,q)-Laplace equations with two parameters. (2020), to appear in Comm. Contemp. Math. 20
- [10] K.J. Brown, The Nehari manifold for a semilinear elliptic equation involving a sublinear term. Calc. Var. Partial Differential Equations 22 (2005), 483–494. 2, 3, 19, 21

- [11] K.J. Brown, Y. Zhang, The Nehari manifold for a semilinear elliptic equation with a sign-changing weight function. J. Differential Equations 193 (2003), no. 2, 481–499. 2, 3
- [12] J. Chabrowski, C. Tintarev, An elliptic problem with an indefinite nonlinearity and a parameter in the boundary condition, NoDEA Nonlinear Differ. Equ. Appl. 21 (2014), 519–540. 19
- [13] B. Chen, Z. Q. Ou, Existence and bifurcation behavior of positive solutions for a class of Kirchhoff-type problems. Comput. Math. Appl. 77(10), 2859–2866 (2019). 21
- [14] M. Cuesta, Eigenvalue problems for the p-Laplacian with indefinite weights. Electron. J. Differential Equations 2001, No. 33, 9 pp. 19
- [15] M. Cuesta, H. Ramos Quoirin, A weighted eigenvalue problem for the p-Laplacian plus a potential. NoDEA Nonlinear Differential Equations Appl. 16 (2009), no. 4, 469–491. 18
- [16] G. Dai, Eigenvalues, global bifurcation and positive solutions for a class of nonlocal elliptic equations. Topol. Methods Nonlinear Anal. 48(1), 213–233 (2016) 22
- [17] P. Drábek, S. I. Pohozaev, Positive solutions for the p-Laplacian: application of the fibering method. Proc. Roy. Soc. Edinburgh Sect. A 127 (1997), no. 4, 703–726. 2
- [18] G. M. Figueiredo, H. Ramos Quoirin, K. Silva, Ground states of elliptic problems over cones. Calc. Var. Partial Differential Equations 60 (2021), no. 5, Paper No. 189, 29 pp. 20
- [19] Y. Il'yasov, On positive solutions of indefinite elliptic equations, Comptes Rendus de l'Académie des Sciences-Series I-Mathematics 333 (2001), 533–538. 2
- [20] Y. Il'yasov, K. Silva, On branches of positive solutions for p-Laplacian problems at the extreme value of the Nehari manifold method, Proc. Amer. Math. Soc. 146 (2018), no. 7, 2925–2935. 1, 2, 19
- [21] U. Kaufmann, H. Ramos Quoirin, K. Umezu, Nonnegative solutions of an indefinite sublinear Robin problem I: positivity, exact multiplicity, and existence of a subcontinuum, Annali di Matematica 199, 2015–2038 (2020). 19
- [22] U. Kaufmann, H. Ramos Quoirin, K. Umezu, Nonnegative solutions of an indefinite sublinear Robin problem II: local and global exactness results, to appear in Israel J. Math. 19
- [23] U. Kaufmann, H. Ramos Quoirin, K. Umezu, Uniqueness and positivity issues in a quasilinear indefinite problem. Calc. Var. Partial Differential Equations 60 (2021), no. 5, Paper No. 187, 21 pp. 8, 19
- [24] T. Ouyang, On the positive solutions of semilinear equations $\Delta u + \lambda u + hu^p = 0$ on compact manifolds. II. Indiana Univ. Math. J. 40 (1991), no. 3, 1083–1141. 2, 8
- [25] K. Silva, S. M. Sousa, Finer analysis of the Nehari set associated to a class of Kirchhoff-type equations. SN Partial Differ. Equ. Appl. 1, 43 (2020). 21, 22
- [26] A. Szulkin, T. Weth, The method of Nehari manifold. Handbook of nonconvex analysis and applications, 597–632, Int. Press, Somerville, MA, (2010). 3

H. RAMOS QUOIRIN

CIEM-FAMÁF UNIVERSIDAD NACIONAL DE CÓRDOBA, (5000) CÓRDOBA, ARGENTINA Email address: humbertorq@gmail.com

K. SILVA INSTITUTO DE MATEMÁTICA E ESTATÍSTICA. UNIVERSIDADE FEDERAL DE GOIÁS, RUA SAMAMBAIA, 74001-970, GOIÂNIA, GO, BRAZIL Email address: kayesilva@ufg.br