Moments of Subsets of General Equiangular Tight Frames

Marina Haikin
Amazon, Tel Aviv, Israel

mkokotov@gmail.com
Ram Zamir
Tel Aviv University, Israel
zamir@eng.tau.ac.il

Abstract

This note outlines the steps for proving that the moments of a randomly-selected subset of a general ETF (complex, with aspect ratio $0 < \gamma < 1$) converge to the corresponding MANOVA moments. We bring here an extension for the proof of the 'Kesten-Mckay' moments (real ETF, $\gamma = 1/2$) [1]. In particular, we establish a recursive computation of the rth moment, for $r = 1, 2, \ldots$, and verify, using a symbolic program, that the recursion output coincides with the MANOVA moments.

I. RECURSIVE COMPUTATION OF MOMENTS

Let F be a unit-norm equiangular tight frame comprised of n vectors in C^m , $\gamma = \frac{m}{n} < 1$ is the aspect ratio of the frame. We define $s \stackrel{\Delta}{=} (1-x)/2\sqrt{x}$, where $x = \gamma^{-1} - 1$ and denote a generalized conference matrix as

$$S = \sqrt{\frac{n-1}{x}} \cdot \left(F'F - \frac{1}{2\gamma} \cdot I_n \right) \tag{1}$$

which satisfies:

- (i) $S_{i,i} = \text{constant} = \sqrt{n-1}(1-x)/2\sqrt{x} \stackrel{\Delta}{=} \sqrt{n-1} \cdot s$,
- (ii) S is conjugate-symmetric and $|S_{i,j}| = 1$ for every $i, j \in [n]$ with $i \neq j$, and
- (iii) $S'S \propto I_n$, i.e., the off-diagonal elements of S^2 are zero, while the diagonal elements are equal to $(n-1)(x+1)^2/4x$.

Note that x/(n-1) is the Welch bound, i.e., the squared absolute cross correlation between the frame vectors. For $\gamma = 0.5$, we have x = 1, s = 0, so $S = \sqrt{n-1} \cdot (F'F - I_n)$, and S'S = (n-1)I, in agreement with the properties of the conference matrix in [1].

§Now at Amazon. Previously at Tel Aviv University where this work was performed.

We extend the proof in [1] to derive the expression for moments of the random submatrix $X_S = PSP$ defined as

$$m_k^S = \frac{1}{n^{k/2+1}} \cdot E\left\{\operatorname{trace}(X_S^k)\right\} = \sum_{t=1}^k \left(\sum_{\pi \in \prod(k,t)} V_n(\pi)\right) \cdot p^t \tag{2}$$

where P is a diagonal matrix such that the diagonal elements are independent random variables Bernoulli(p). For every partition π of [k] into t blocks, $V_n(\pi)$ is sum of products of k elements from the generalized conference matrix S, as a function of the partition. The sum is over all possible t distinct values from [n] according to the blocks of π (value per block),

$$V_n(\pi) \stackrel{\Delta}{=} \frac{1}{n^{k/2+1}} \sum_{a \in L_n(\pi)} S_{a(1),a(2)} S_{a(2),a(3)} \cdots S_{a(k),a(1)}, \tag{3}$$

$$L_n(\pi) \stackrel{\Delta}{=} \{a : [k] \to [n] : \{a^{-1}(a(i)) : i \in [k]\} = \pi\}. \tag{4}$$

The main difference from the KM case will be the computation of $V_n(\pi)$. Lemma 16 in [1] will be replaced by

Lemma 1. For every non-crossing partition $\pi \in \prod(k,t)$, the following holds: if the edges of G_{π} partition into m simple cycles of sizes l_1, \ldots, l_m , then as $n \to \infty$,

$$V_n(\pi) \to A_{l_1} \cdots A_{l_m} \tag{5}$$

where

$$A_{l+1} = -\sum_{i=1}^{l} A_i A_{l+1-i}, \quad A_1 = s, A_2 = 1.$$
(6)

Note that for $\gamma=0.5$, s=0 and we get that for odd cycle, i.e., odd l_i , $A_{l_i}=0$, and for even l_i , $A_{l_i}=(-1)^{l_i/2-1}\cdot C_{l_i/2-1}$. Thus if G_{π} contains any odd cycle, then $V_n(\pi)\to 0$ otherwise $V_n(\pi)$ converges to a product of Catalan numbers, in agreement with Lemma 16 in [1].

For general γ ($s \neq 0$), we don't have yet an a analogy to Lemma 8, but $\sum_{\pi \in \prod(k,t)} V_n(\pi)$ can be computed by counting all N(k,t) (Narayana number) non-crossing partitions of [k] into t blocks and for each induced cactus to compute $V_n(\pi)$ according to Lemma 1.

The desired moments of a (Bernoulli-p selected) subset of ETF F are then derived by using (1) and the binom formula:

$$m_k \stackrel{\Delta}{=} E\left\{\frac{1}{n} \cdot \operatorname{trace}((PF'FP)^k)\right\}$$

$$= \sum_{j=0}^k \binom{k}{j} \cdot x^{j/2} \cdot \left(\frac{x+1}{2}\right)^{k-j} p^{j=0} \cdot \frac{1}{n^{j/2+1}} \cdot E\left\{\operatorname{trace}(X_S^j)\right\}$$

$$= \left(\frac{x+1}{2}\right)^k \cdot p + \sum_{j=1}^k \binom{k}{j} \cdot x^{j/2} \cdot \left(\frac{x+1}{2}\right)^{k-j} \cdot m_j^S$$
(7)

where m_j^S is defined in (2), and j==0 denotes 1 if j=0 or 0 otherwise.

Before we turn to the proof, we apply the recursion for several first moments:

$$m_0^S = 1 \qquad m_0 = 1$$

$$A_1 = s \qquad m_1^S = sp \qquad m_1 = p$$

$$A_2 = 1 \qquad m_2^S = s^2p + p^2 \qquad m_2 = p + p^2x$$

$$A_3 = -2s \qquad m_3^S = s^3p + 3sp^2 + (-2s)p^3 \qquad \vdots$$

$$A_4 = 4s^2 - 1 \qquad m_4^S = s^4p + 6s^2p^2 + (-8s^2 + 2)p^3 + (4s^2 - 1)p^4$$

$$A_5 = -8s^3 + 6s \qquad \vdots$$

$$A_6 = 16s^4 - 24s^2 + 2$$

Proof. We follow the notations in [1] $\Delta(a(1), a(2), \dots, a(k)) = S_{a(1), a(2)} S_{a(2), a(3)} \cdots S_{a(k), a(1)}$. If $a_j = a_{j+1}$ for any $j \in [k-1]$ or $a_k = a_1$,

$$V_n(\pi) = \frac{1}{n^{k/2+1}} \sum_{a \in L_n(\pi)} \Delta(a(1), \dots a(j-1), a(j+1), a(j+1), \dots, a(k))$$
(8)

$$= \frac{\sqrt{n-1}s}{n^{k/2+1}} \sum_{a \in L_n(\pi)} \Delta(a(1), \dots, a(j-1), a(j+1), \dots, a(k))$$
(9)

$$= \frac{s}{n^{(k-1)/2+1}} \sum_{a \in L_n(\pi \setminus \{j\})} \Delta(a(1), \dots, a(j-1), a(j+1), \dots, a(k)) + o(1)$$
(10)

were the first equality follows from the fact that the diagonal entries of S are $s\sqrt{n-1}$, thus $S_{a(j+1)a(j+1)} = s\sqrt{n-1}$. The restriction of $\pi \setminus \{j\}$ to $[k] \setminus j$ results in a partition π' of $[k] \setminus j$ into same t blocks besides removing j from the block $\pi(j)$. Thus the above expression of $V_n(\pi)$ implies

$$V_n(\pi) \to s \cdot V_n(\pi'),$$
 (11)

where throughout the proof, \to means as $n \to \infty$. This means that two consecutive indices which belong to same block and contribute a loop to the graph G_{π} , contribute a constant factor s to $V_n(\pi)$. This claim, as well as the derivation in (8)-(10), extends for several loops. Given partition π with r loops in total (in one or more blocks), we can consider a squeezed partition π' with same number of blocks and k' = k - r. Each block B'_i for $i = 1, \ldots, t$, is a squeezed block B_i , with only one representative index of sequence of several consecutive equal indices.

$$V_n(\pi) \to s^r \cdot V_n(\pi') \tag{12}$$

For example a partition 122234422333341 translated to 1234234 with a factor of s^8 .

Now we focus on estimation of $V_n(\pi')$. $G_{\pi'}$ is a loop-free graph and thus Lemma 6 in [1] holds for the same reasons. Lemma 12(iv),12(v) also holds due to similar the properties of S, and thus Lemma 14 - which relies on all of the above - holds. This proves the claim of vanishing crossing partitions, i.e., let $\pi' \in \prod(k',t)$ be a crossing partition, then $V_n(\pi') \to 0$.

We follow the proof by induction of Lemma 16 in [1]. Lemma 12(i) holds again by S properties and thus for t=2, $V_n(\pi')=V_n(\{1\},\{2\})=1$. As for G_π , with assumed r cycles of sizes 1, proving the lemma for $V_n(\pi')$, will immediately imply the proof for $V_n(\pi)$, as $A_1^r=s^r$. Assuming that the (loops-removed) partition π' is non-crossing,

it follows that the graph $G_{\pi'}$ is a cactus, and by Lemma 15 it is guaranteed to contain a singleton leaf (block in the partition). As we deal with π' , with m' = m - r cycles, we aim to prove that

$$V_n(\pi') \to A_{l_1} \cdots A_{l_{m'}}. \tag{13}$$

Case I of the proof refers to the case when the singleton block resides in a cycle of length 2, i.e., if the singleton is $\{k\} \in \pi$, $\pi(1) = \pi(k)$. A similar derivation as in [1] implies

$$V_n(\pi') = V_n(\pi'') + o(1). \tag{14}$$

Our induction hypothesis and (14), imply

$$V_n(\pi') \to A_{l_1} \cdots A_{l_{m'-1}}.$$
 (15)

Since $l_m = 2$ and $A_2 = 1$, this establishes (13).

Case II of the proof refers to the case when $\{k\} \in \pi$ (the singleton block) resides in a cycle of length $l \geq 3$. With similar analysis to [1] we get:

$$V_n(\pi') = -\sum_{i=2}^{l} V_n(\pi'^i) + o(1).$$
(16)

Assuming $l_m = l$ and applying the induction hypothesis, we have

$$V_n(\pi') \to -A_{l_1} \cdots A_{l_{m'-1}} \cdot \sum_{i=2}^l A_{l-1} A_{l-i+1} = -A_{l_l} \cdots A_{l_{m'-1}} \cdot \sum_{i=1}^{l-1} A_l A_{l-i}. \tag{17}$$

Applying the identity (6), we get $V_n(\pi') \to A_{l_1} \cdots A_{l_{m'}}$, thereby establishing (13).

II. IMPLEMENTATION OF THE DERIVED ALGORITHM

```
import numpy as np
import sympy
from sympy.utilities.iterables import multiset_partitions
import scipy.special
# recursive calculation of A(l) for l=1..n, arr_A[i]=A(i+1)
def calc_A(n,s):
   arr_A = np.zeros(n)
    arr_A[0] = s
   arr_A[1] = 1
    for i in np.arange(n-2)+2:
        arr_A[i] = -sum([arr_A[j]*arr_A[i-j-1] for j in np.arange(i)])
    return arr A
# generation of all non crossing partitions of k to t blocks
def calc_partitions_k_t(k,t):
    partitions = list(multiset_partitions(np.arange(k),t))
    non_crossing = []
    non_crossing_len = []
   for part in partitions:
        cross = 0
        n = len(part)
        if n == 1:
            non_crossing.append(part)
            non_crossing_len.append([len(b) for b in part])
        else:
```

```
for b1 in part:
                for b2 in part:
                    if (b2[0]>b1[0]) & (len(np.array(np.where((b1>b2[0]) & (b1<b2[-1]))) [0])>0):
                        cross = 1
                        break
                if cross==1:
                    break
            if cross==0:
                non_crossing.append(part)
                non_crossing_len.append([len(b) for b in part])
    return non_crossing
Find length of cactus leafs which correspond to a non-crossing partition
# genrate lists off all intervals within blocks of partition
# list_pairs: a list of consecutive pairs in blocks [block[i],block[i+1]] for every block in the partition
# list_pairs_intervals: a list of intervals between consecutive pairs in blocks block[i+1]-block[i] for every
# block in the partition
def create_pairs_lists(part):
   list_pairs = []
    list_pairs_intervals = []
    for block in part:
        b_len = len(block)
        if b_len > 1:
            for i in range(b_len-1):
                list_pairs.append([block[i],block[i+1]])
                list_pairs_intervals.append(block[i+1]-block[i])
    return [list_pairs, list_pairs_intervals]
# sort the lists by the length of the intervals.
# start computing the the length of cycle which corresponds to an interval, from shorter to longer -
# for each interval substruct the cycles which correspond to smaller intervals contained in the interval
def update_lists(list_pairs, list_intervals):
    zipped_lists = zip(list_intervals, list_pairs)
    sorted_pairs = sorted(zipped_lists)
   tuples = zip(*sorted_pairs)
   list_intervals, list_pairs = [list(tuple) for tuple in tuples]
    for i in range(len(list_intervals)):
        intersections\_length = 0
        for j in range(i):
            if (list_pairs[j][0] > list_pairs[i][0]) & (list_pairs[j][1] < list_pairs[i][1]):</pre>
                intersections_length = intersections_length + list_intervals[j]
        list_intervals[i] = list_intervals[i] - intersections_length
    return list_intervals
# generated the updated lists and add the cycle which correspond to an outer interval [k, 0],
\# the length of which is k - sum of all cycles
def partition_to_cactus_sizes(part,k):
    [list_pairs, list_intervals] = create_pairs_lists(part)
    if len(list_pairs)>0:
        list_intervals = update_lists(list_pairs, list_intervals)
        cactus_sizes = list_intervals
        cactus_sizes.append(k - sum(cactus_sizes))
        cactus\_sizes = [k]
    return cactus sizes
calculation of the 'centralized' moments
# calculation of asymptotic V_n(pi) for a given partition by eq. 5 and sum over all possible (k,t) non-crossing
# partitions
```

```
V_pi_sum = V_pi_sum + np.prod(arr_A[np.array(cactus)-1])
   return V_pi_sum
\# calculation of m_s_k by eq. 2
def calc_m_s(k,p):
   m_s = 0
   for t in np.arange(k)+1:
      m_s = m_s + calc_V_{pi_sum(k,t)*p**t}
   return m s
binom formula for 'de-centralization' of the moments, eq.7
def calc_m(k, x, p):
   m = ((x+1)/2) * *k*p
   for j in np.arange(k)+1:
       m = m + scipy.special.binom(k,j) *x**(j/2)*((x+1)/2)**(k-j)*(m_s_vec[j-1])
   return m
# run calculation of 10 first moments
gamma = np.sqrt(0.5)
p = 0.6
x = 1/qamma - 1
s = (1-x)/(2*np.sqrt(x))
d = 10
arr_A = calc_A(d,s)
m_s_vec = [calc_m_s(k,p) \text{ for } k \text{ in } np.arange(d)+1]
m_{vec} = [calc_m(k, x, p) \text{ for } k \text{ in } np.arange(d)+1]
And the output is:
gamma = 0.7071067811865476
p = 0.6
x = 0.4142135623730949
s = 0.45508986056222756
_____
calc ETF moments until d = 10
A[1] for 1 = 1..10
-2.1862701245144653, 10.126983722080917, -8.324794719173925, -21.347181155833745]
A[1] for 1 = 1...6 by manual application of eq.6
[0.45508986056222756, \ 1.0, \ -0.9101797211244551, \ -0.17157287525380915, \ 1.9765215939999432, \ -2.2842712474619047]
example for all partitions (5,3) and the corresponding cactus lengths
[[0, 1, 2], [3], [4]] [1, 1, 3]
[[0, 1, 3], [2], [4]]
                     [1, 2, 2]
[[0, 1], [2, 3], [4]]
                     [1, 1, 3]
[[0, 1, 4], [2], [3]]
                      [1, 3, 1]
[[0, 1], [2, 4], [3]]
                      [1, 2, 2]
[[0, 1], [2], [3, 4]]
                      [1, 1, 3]
[[0, 2, 3], [1], [4]]
                      [1, 2, 2]
[[0, 2, 4], [1], [3]]
                     [2, 2, 1]
[[0, 2], [1], [3, 4]]
                     [1, 2, 2]
[[0, 3], [1, 2], [4]] [1, 2, 2]
```

cactuses = [partition_to_cactus_sizes(part,k) for part in calc_partitions_k_t(k,t)]

def calc_V_pi_sum(k,t):
 V_pi_sum = 0

for cactus in cactuses:

```
[1, 1, 3]
[[0], [1, 2, 3], [4]]
                  [1, 3, 1]
[[0, 4], [1, 2], [3]]
[[0], [1, 2, 4], [3]]
                   [1, 2, 2]
[[0], [1, 2], [3, 4]]
                   [1, 1, 3]
                   [1, 3, 1]
[[0, 3, 4], [1], [2]]
[[0, 4], [1, 3], [2]]
                   [2, 2, 1]
[[0], [1, 3, 4], [2]]
                  [1, 2, 2]
[[0, 4], [1], [2, 3]]
                  [1, 3, 1]
[[0], [1, 4], [2, 3]]
                  [1, 2, 2]
[[0], [1], [2, 3, 4]]
                   [1, 1, 3]
example of cactus length for partition [[0,2],[1],[3],[4,5,7,10],[6],[8,9],[11]]
[1, 1, 2, 2, 2, 4]
m_s[k] for k = 1..10
0.6031675926696468, 0.5443213610399722, 0.7080038062958746, 0.6683603744969959, 0.8395068505692707]
m_s[k] for l = 1..4 by manual application of eq.2
[0.2730539163373365, 0.4842640687119286, 0.35144954734733014, 0.5249702161277665]
_____
m[k] for k = 1..10
3.633212856926222, \ 5.091635541233709, \ 7.151079693455817, \ 10.05923853882072]
m[k] for l = 1..10 by Dubbs&Edelman (Manova moments triangles in Fig.1)
[0.6,\ 0.7491168824543142,\ 0.994940258945177,\ 1.355364155367106,\ 1.8704311672940082,\ 2.6006998421840386,
3.633212856926222, 5.091635541233708, 7.1510796934558165, 10.059238538820706]
```

The output of a symbolic calculation is:

```
gamma = gamma
p = p
                                      _____
x = x
s = s
calc ETF moments until d = 10
 _____
A[1] for 1 = 1..10
[s, 1, -2*s, 4*s**2 - 1, -8*s**3 + 6*s, 16*s**4 - 24*s**2 + 2,
    -32 \times s \times *5 + 80 \times s \times *3 - 20 \times s, \ 64 \times s \times *6 - 240 \times s \times *4 + 120 \times s \times *2 - 5, \ -128 \times s \times *7 + 672 \times s \times *5 - 560 \times s \times *3 + 70 \times s, \ -128 \times s \times *7 + 672 \times s \times *5 + 80 \times s \times 5 + 80 \times s \times 5
   256*s**8 - 1792*s**6 + 2240*s**4 - 560*s**2 + 141
A[l] for l = 1..6 by manual application of eq.6
[s, 1, -2*s, 4*s**2 - 1, -8*s**3 + 6*s, 16*s**4 - 24*s**2 + 2]
  _____
m_s[k] for k = 1..10 [to save space we show only first 6]
[p*s, p**2 + p*s**2, -2*p**3*s + 3*p**2*s + p*s**3,
 p**4*(4*s**2 - 1) + p**3*(-8*s**2 + 2) + 6*p**2*s**2 + p*s**4,
  p**5*(-8*s**3 + 6*s) + p**4*(5*s*(4*s**2 - 1) - 10*s) + p**3*(-20*s**3 + 10*s) + 10*p**2*s**3 + p*s**5,
   p**6*(16*s**4 - 24*s**2 + 2) + p**5*(36*s**2 + 6*s*(-8*s**3 + 6*s) - 6) +
   p**4*(15*s**2*(4*s**2-1)-60*s**2+5)+p**3*(-40*s**4+30*s**2)+15*p**2*s**4+p*s**6, ...]
m_s[k] for l = 1..4 by manual application of eq.2
 [p*s, p**2 + p*s**2, -2*p**3*s + 3*p**2*s + p*s**3,
  p**4*(4*s**2 - 1) + p**3*(-8*s**2 + 2) + 6*p**2*s**2 + p*s**4]
_____
gamma = gamma
q = q
s = (-x + 1) / (2*sqrt(x))
calc ETF moments until d = 10
```

```
 \begin{aligned} &\text{m[k] for k = 1..10} \\ &[0.5*p*(-x+1)+p*(x/2+1/2),\\ &1.0*p*(-x+1)*(x/2+1/2)+p*(x/2+1/2)**2+1.0*x**1.0*(p**2+p*(-x+1)**2/(4*x)),\\ &1.5*p*(-x+1)*(x/2+1/2)**2+p*(x/2+1/2)**3+3.0*x**1.0*(p**2+p*(-x+1)**2/(4*x))*(x/2+1/2)+\\ &1.0*x**1.5*(p**3*(x-1)/sqrt(x)+3*p**2*(-x+1)/(2*sqrt(x))+p*(-x+1)**3/(8*x**(3/2))),\\ &2.0*p*(-x+1)*(x/2+1/2)**3+p*(x/2+1/2)**4+6.0*x**1.0*(p**2+p*(-x+1)**2/(4*x))*(x/2+1/2)**2+\\ &4.0*x**1.5*(x/2+1/2)*(p**3*(x-1)/sqrt(x)+3*p**2*(-x+1)/(2*sqrt(x))+p*(-x+1)**3/(8*x**(3/2)))+\\ &1.0*x**2.0*(p**4*(-x+(x-1)**2)/x+p**3*(2+2*(-x+1)*(x-1)/x)+3*p**2*(-x+1)**2/(2*x)+\\ &p*(-x+1)**4/(16*x**2)), \ldots] \end{aligned} \\ &\text{m[k] for 1 = 1..10 by Dubbs&Edelman (Manova moments triangles in Fig.1)} \\ &[p,\\ &p**2*x+p,\\ &p**4*(x**3-3*x**2+x)+p**3*(6*x**2-4*x)+6*p**2*x+p, \ldots] \end{aligned}
```

REFERENCES

[1] M. Magsino, D. G. Mixon, and H. Parshall, "Kesten-Mckay law for random subensembles of paley equiangular tight frames," *Constructive Approximation*, pp. 1–22, 2020.