

The fractional chromatic number of K_Δ -free graphs

Xiaolan Hu*

Xing Peng†

Abstract

For a simple graph G , let $\chi_f(G)$ be the fractional chromatic number of G . In this paper, we aim to establish upper bounds on $\chi_f(G)$ for those graphs G with restrictions on the clique number. Namely, we prove that for $\Delta \geq 4$, if G has maximum degree at most Δ and is K_Δ -free, then $\chi_f(G) \leq \Delta - \frac{1}{8}$ unless $G = C_8^2$ or $G = C_5 \boxtimes K_2$. This improves the result in [King, Lu, and Peng, *SIAM J. Discrete Math.*, 26(2) (2012), pp. 452-471] for $\Delta \geq 4$ and the result in [Katherine and King, *SIAM J. Discrete Math.*, 27(2) (2013), pp. 1184-1208] for $\Delta \in \{6, 7, 8\}$.

1 Introduction

In this paper, we use $\chi(G)$ to denote the chromatic number of a simple graph G . It is well-known that $\Delta + 1$ is a trivial upper bound for $\chi(G)$, where Δ is the maximum degree of G . The famous Brooks' Theorem states that if the maximum degree of a connected graph G is Δ , then $\chi(G) \leq \Delta$ unless G is $K_{\Delta+1}$ or an odd cycle. This seminal result leads to the idea bounding the chromatic number $\chi(G)$ in terms of the maximum degree $\Delta(G)$ and the clique number $\omega(G)$. For $\Delta \geq 3$, Brooks' Theorem asserts that if G is $K_{\Delta+1}$ -free (i.e., $\omega(G) \leq \Delta(G)$), then $\chi(G) \leq \Delta(G)$. One may ask to show an upper bound for $\chi(G)$ if G is K_Δ -free. This is exactly the following conjecture by Borodin and Kostochka [1].

Conjecture 1 *For a connected graph G , if $\omega(G) \leq \Delta(G) - 1$ and $\Delta(G) \geq 9$, then we have $\chi(G) \leq \Delta(G) - 1$.*

If the conjecture is true, then it is best possible since there is a K_8 -free graph $G = C_5 \boxtimes K_3$ (see Figure 1) with $\Delta(G) = 8$, $\omega(G) = 6$, and $\chi(G) = 8$.




Figure 1: The graphs $C_5 \boxtimes K_3$, C_8^2 , and $C_5 \boxtimes K_2$.

*School of Mathematics and Statistics & Hubei Key Laboratory of Mathematical Sciences, Central China Normal University, Wuhan 430079, P. R. China. E-mail: xlu@mail.ccnu.edu.cn, Research is partially supported by National Natural Science Foundation of China (No. 11971196).

†School of Mathematical Sciences, Anhui University, Hefei 230601, P. R. China. E-mail: x2peng@ahu.edu.cn, Research is supported in part by National Natural Science Foundation of China (No. 12071002) and a Start-up Fund from Anhui University.

Given two graphs G and H , the *strong product* $G \boxtimes H$ is the graph with vertex set $V(G) \times V(H)$, and (u, x) is connected to (v, y) if one of the following holds

- 1: $u = v$ and $xy \in E(H)$,
- 2: $uv \in E(G)$ and $x = y$,
- 3: $uv \in E(G)$ and $xy \in E(H)$.

Using the probabilistic methods, Reed [17] confirmed Conjecture 1 for $\Delta \geq 10^{14}$. In the same paper, Reed claimed that a careful calculation could reduce 10^{14} to 10^3 . Let us assume that Conjecture 1 is true for $\Delta \geq 10^3$. This is the state of the art of Conjecture 1. King, Lu, and Peng first considered to bound the fractional chromatic number of K_Δ -free graphs, which can be viewed as the fractional analog of Conjecture 1.

The fractional chromatic number $\chi_f(G)$ of a graph G can be defined as follows. For positive integers a and b with $b \leq a$, we use $[a]$ to denote the set of first a positive integers and $\binom{[a]}{b}$ to denote the set of b -subsets of $[a]$. An $(a:b)$ -coloring of a graph G is a mapping $\psi : V(G) \rightarrow \binom{[a]}{b}$ such that $\psi(u) \cap \psi(v) = \emptyset$ whenever $uv \in E(G)$. The b -fold coloring number of G , denoted by $\chi_b(G)$, is the smallest integer a such that G has an $(a:b)$ -coloring. Note that $\chi_1(G) = \chi(G)$. The *fractional chromatic number* $\chi_f(G)$ of G is $\lim_{b \rightarrow \infty} \frac{\chi_b(G)}{b}$. If G is vertex transitive, then $\chi_f(G) = \frac{|V(G)|}{\alpha(G)}$, where $\alpha(G)$ is the independence number of G .

For $\Delta \geq 3$, we define

$$f(\Delta) = \min_G \{ \Delta - \chi_f(G) : \Delta(G) \leq \Delta, \omega(G) \leq \Delta - 1, G \notin \{C_8^2, C_5 \boxtimes K_2\} \}.$$

The value of $f(3)$ received a great deal of attention. Actually, based on the study of independent sets in triangle-free subcubic graphs [4, 7, 9, 19], Heckman and Thomas [9] conjectured $f(3) = \frac{1}{5}$. Before Dvořák, Sereni, and Volec [2] proved Heckman and Thomas' Conjecture, progress towards this conjecture can be found in [5, 8, 11, 13, 14]. King, Lu, and Peng [12] first proved that $f(\Delta) \geq \frac{2}{67}$ for each $\Delta \geq 4$. Edwards and King [3] established an improved lower bound on $f(\Delta)$ for all $\Delta \geq 6$. In particular, they showed $f(6) \geq \frac{1}{22.5}$, $f(7) \geq \frac{1}{11.2}$, $f(8) \geq \frac{1}{8.9}$, and $f(\Delta) > \frac{1}{7.7}$ for each $\Delta \geq 9$.

We prove the following theorem in this paper.

Theorem 1 *For each $\Delta \geq 4$, if G has maximum degree at most Δ and is K_Δ -free, then $\chi_f(G) \leq \Delta - \frac{1}{8}$ unless $G = C_8^2$ or $G = C_5 \boxtimes K_2$ (see Figure 1).*

By the theorem above, we get $f(\Delta) \geq \frac{1}{8}$ for each $\Delta \geq 4$. Therefore, Theorem 1 improves the result of King, Lu, and Peng [12] for each $\Delta \geq 4$ and the result of Edwards and King [3] for $\Delta \in \{6, 7, 8\}$.

The rest of this paper is organized as follows. In Section 2, we will prove Theorem 1. Section 3 is devoted to prove technical lemmas.

2 Proof of Theorem 1

Recall the definition of $f(\Delta)$. King, Lu, and Peng [12] proved the following lemma on $f(\Delta)$.

Lemma 1 *For each $\Delta \geq 6$, we have $f(\Delta) \geq \min \{f(\Delta - 1), \frac{1}{2}\}$. We also have $f(5) \geq \min \{f(4), \frac{1}{3}\}$.*

In view of Lemma 1, in order to prove Theorem 1, we need only to show $f(4) \geq \frac{1}{8}$. We define

$$\mathcal{G} = \{G : \Delta(G) \leq 4, \omega(G) \leq 3, \text{ and } G \neq C_8^2\}.$$

Equivalently, we have to show $\chi_f(G) \leq 4 - \frac{1}{8}$ for each $G \in \mathcal{G}$. Actually, we will prove a stronger result. To state our result, we need to introduce some definitions.

Let G be a graph. For $v \in V(G)$, the *degree* $d_G(v)$ of v is the number of edges incident with v in G . For two vertices u and v of G , let $\text{dist}_G(u, v)$ be the length of a shortest path between u and v . For $v \in V(G)$ and a positive integer j , let $N_G^j(v) = \{u \in V(G) : \text{dist}_G(u, v) = j\}$. If S and T are two disjoint subsets of $V(G)$, then the number of edges between S and T is denoted by $e(S, T)$. A *demand function* h_G is a function from $V(G)$ to $[0, 1]$ taking rational values. For positive integers a and b with $a < b$, we use $[a, b]$ to denote the set $\{a, a+1, \dots, b\}$. If $a = 1$, then we write $[b]$ instead of $[1, b]$ for short. Suppose h_G is a demand function and N is a positive integer such that $N \cdot h_G(v)$ is a positive integer for each $v \in V(G)$. Let ψ be a function from $V(G)$ to the set of all subsets of $[N]$. If $|\psi(v)| \geq N h_G(v)$ holds for each $v \in V(G)$, then we say ψ is an (h_G, N) -coloring of G . We mention a few remarks on these definitions.

- (i) If G has an (h_G, N) -coloring, then G also has an (h_G, M) -coloring provided M is divisible by N . Actually, we can duplicate each color from an (h_G, N) -coloring M/N times to get an (h_G, M) -coloring.
- (ii) If ψ is an (h_G, N) -coloring of G , then we can assume that $|\psi(v)| = N h_G(v)$ for each $v \in V(G)$ by deleting colors from $\psi(v)$. We will keep this assumption in the rest of this paper.
- (iii) To show $\chi_f(G) \leq r$ for a rational number r , it is enough to show G has an (h_G, N) -coloring such that $h_G(v) \geq 1/r$ for each $v \in V(G)$.

In order to prove $\chi_f(G) \leq 4 - \frac{1}{8}$ for each $G \in \mathcal{G}$, we define a demand function h_G as following:

- (i) If $1 \leq d_G(v) \leq 4$, then $h_G(v) = \frac{12-d_G(v)}{31}$;
- (ii) If $d_G(v) = 0$, then $h_G(v) = 1$.

We will prove the following stronger result.

Theorem 2 *For each $G \in \mathcal{G}$, there is a positive integer N_G such that G has an (h_G, N_G) -coloring.*

It is easy to see Theorem 2 implies that $\chi_f(G) \leq 4 - \frac{1}{8}$ for each $G \in \mathcal{G}$. We next prove Theorem 2 by contradiction. Let $G \in \mathcal{G}$ be a counterexample with the smallest number of vertices. The following simple remark follows from the remark (i) on the definition of (h_G, N) -coloring.

Remark 1 *There exists a positive integer M such that each graph G' with $|V(G')| < |V(G)|$ has an $(h_{G'}, M)$ -coloring.*

As G is a minimum counterexample, G' has an $(h_{G'}, N_{G'})$ -coloring for each G' with $|V(G')| < |V(G)|$. Thus we can take M as the least common multiplier of $N_{G'}$, where G' ranges over all graphs satisfying $|V(G')| < |V(G)|$.

We will show that G satisfies the following properties.

Lemma 2 *The graph G is 2-connected and satisfies $\delta(G) \geq 3$.*

Proof: Clearly, G is connected. If G contains a cut-vertex v , then there are two subgraphs G_1 and G_2 such that $V(G_1) \cup V(G_2) = V(G)$ and $V(G_1) \cap V(G_2) = \{v\}$. As G is a minimum counterexample, the graph G_i has an (h_{G_i}, M) -coloring ψ_i for $1 \leq i \leq 2$. By deleting colors if it is necessary, we can assume that $|\psi_1(v)| = |\psi_2(v)| = M \cdot h_G(v)$, here $h_G(v) = \frac{12-d_G(v)}{31}$.

By permuting colors in ψ_2 , we can further assume that $\psi_1(v) = \psi_2(v)$. Thus the union of ψ_1 and ψ_2 is an (h_G, M) -coloring of G . A contradiction.

Since G is 2-connected, we get $\delta(G) \geq 2$. Suppose that there exists $v \in V(G)$ with $d_G(v) = 2$. Let u and w be the neighbors of v . By the minimality of G , the graph $G' = G - v$ has an $(h_{G'}, M)$ -coloring, say ψ . Deleting colors from $\psi(u)$ and $\psi(w)$, we can assume that $|\psi(u)| = M \cdot \frac{12-d_G(u)}{31}$ and $|\psi(w)| = M \cdot \frac{12-d_G(w)}{31}$. Since $d_G(u) \geq 2$ and $d_G(w) \geq 2$, it follows that $|\psi(u)| \leq \frac{10}{31} \cdot M$ and $|\psi(w)| \leq \frac{10}{31} \cdot M$. As $|[M] \setminus (\psi(u) \cup \psi(w))| \geq \frac{11}{31} \cdot M \geq h_G(v) \cdot M$, we can extend the coloring ψ to v and obtain an (h_G, M) -coloring of G . A contradiction. \square

We also need the following lemma on the local structures of G .

Lemma 3 *For each $v \in V(G)$, if $d_G(v) = 3$, then we have $e(N_G(v), N_G^2(v)) \geq 5$; if $d_G(v) = 4$, then we have $e(N_G(v), N_G^2(v)) \geq 9$.*

The proof of Lemma 3 is quite involved, we postpone it until Section 3 and proceed to prove Theorem 2.

Proof of Theorem 2: We will prove that G has an (h_G, M') -coloring for some positive integer M' . This will be a contradiction to the assumption of G . For each $v \in V(G)$, let H_v be the proper subgraph of G obtained from G by deleting vertices from $N_G(v)$. By Remark 1 and the choice of G , the graph H_v has an (h_{H_v}, M) -coloring, say ψ_v . Let ψ be the union of ψ_v , where v ranges over all vertices of G and colors used for different ψ_v are distinct. Let $n = |V(G)|$. We next show ψ is an (h_G, M') -coloring, where $M' = nM$. To show this, we have to verify $|\psi(u)| \geq M'h_G(u)$ for each vertex u . By Lemma 2, there are two cases depending on the degree of u .

If $d_G(u) = 3$, then we write $N_G^2(u) = \cup_{i=1}^3 A_i$, where $w \in A_i$ if w has i neighbors in $N_G(u)$. We can compute the number of colors u received as follows. In the graph H_u , the vertex u is an isolated one and it receives M colors from ψ_u . For each $v \in N_G(u)$, the vertex u is not in the graph H_v and does not receive any color from ψ_v . If $v \in N_G^2(u) \cap A_1$, then u has degree two in H_v and u receives $10M/31$ colors from ψ_v . Similarly, for $v \in N_G^2(u) \cap A_2$, the coloring ψ_v gives u exactly $11M/31$ color as u has degree one in H_v . For $v \in N_G^2(u) \cap A_3$, the vertex u receives M colors from ψ_v as u is an isolated vertex in H_v . If $v \in V(G) \setminus (\{u\} \cup N_G(u) \cup N_G^2(u))$, then u has degree three in H_v and so it receives $9M/31$ colors from ψ_v . For $1 \leq i \leq 3$, let $a_i = |A_i|$. The number of colors for u can be calculated as follows:

$$\begin{aligned} |\psi_u| &= M + a_1 \cdot (10M)/31 + a_2 \cdot (11M)/31 + a_3 \cdot M \\ &\quad + (n - 4 - a_1 - a_2 - a_3) \cdot (9M)/31 \\ &= (9n + a_1 + 2a_2 + 22a_3 - 5) \cdot M/31 \\ &\geq (9n + a_1 + 2a_2 + 3a_3 - 5) \cdot M/31 \\ &\geq (9nM)/31, \end{aligned}$$

in the last step, we notice $e(N_G(u), N_G^2(u)) = a_1 + 2a_2 + 3a_3 \geq 5$ as the first part of Lemma 3.

If $d_G(u) = 4$, then we write $N_G^2(u) = \cup_{i=1}^4 A_i$, where $w \in A_i$ if w has i neighbors in $N_G(u)$. Let $a_i = |A_i|$ for each $1 \leq i \leq 4$. As we did in the previous case, we can estimate the size of ψ_u as follows:

$$\begin{aligned} |\psi_u| &= M + a_1 \cdot (9M)/31 + a_2 \cdot (10M)/31 + a_3 \cdot (11M)/31 + a_4 \cdot M \\ &\quad + (n - 5 - a_1 - a_2 - a_3 - a_4) \cdot (8M)/31 \\ &= (8n + a_1 + 2a_2 + 3a_3 + 23a_4 - 9) \cdot M/31 \\ &\geq (8n + a_1 + 2a_2 + 3a_3 + 4a_4 - 9) \cdot M/31 \\ &\geq (8nM)/31, \end{aligned}$$

in the last step, we note that $e(N_G(u), N_G^2(u)) = a_1 + 2a_2 + 3a_3 + 4a_4 \geq 9$ by the second part of Lemma 3.

Therefore, ψ is an (h_G, M') -coloring for G . This is a contradiction and the theorem is proved. \square

3 Proof of Lemma 3

We start with some definitions and notations. Let G be a graph. A subset $S \subseteq V(G)$ is called an *independent set* if there are no edges with both endpoints in S . The *contraction* of S is to replace vertices in S by a single fat vertex, denoted by \underline{S} , whose incident edges are all edges that are incident with at least one vertex in S , except edges with both endpoints in S . The new graph obtained by contracting S is denoted by G/S . This operation is also known as identifying vertices of S in the literature. For convenience, if $S = \{u, v\}$, then the fat vertex will be denoted by \underline{uv} . We use $G - S$ to denote the subgraph of G induced by $V(G) - S$. For two nonadjacent vertices u and v , let $G + uv$ be the graph obtained from G by adding an edge uv . The square G^2 of a graph G is the graph with vertex set $V(G)$ and two vertices are adjacent in G^2 if and only if their distance in G is at most 2.

Recall that G is a counterexample with the smallest number of vertices. If H is a subgraph of G with $|V(H)| < |V(G)|$, then H has an (h_H, M) -coloring ψ by the definition of G . For each $v \in V(G) - V(H)$, we define $L(v) = M \setminus \bigcup_{u \in N_H(v)} \psi(u)$. Since $d_H(v) \leq d_G(v)$ for any $v \in V(H)$, we get $|\psi(v)| = (12 - d_H(v))/31 \cdot M \geq (12 - d_G(v))/31 \cdot M$. If ψ' is an (h_G, M) coloring of H such that $\psi'(v) \subseteq \psi(v)$ for each $v \in V(H)$, then we say ψ' is a *limitation* of ψ on H . Given a limitation ψ' and a vertex $v \in V(G) - V(H)$, let $L'(v) = M \setminus \bigcup_{u \in N_H(v)} \psi'(u)$. Roughly speaking, if we have the partial coloring ψ' of G and we wish to color u , then $L'(u)$ is the set of colors which could be assigned to the vertex u . We also have the following simple fact which will be frequently used in our proof. Namely, we are able to prescribe a set of colors for $L'(u)$ by defining the limitation carefully.

Fact 1 *Let $S \subset V(G)$ and H be the subgraph of G induced by the vertex set $V(G) - S$. Suppose that ψ is an (h_H, M) -coloring of H . For $v_1, v_2 \in S$, if $A_1, A_2 \subset [1, M]$ with $|A_1| \leq M/31$ and $|A_2| \leq M/31$, then there is a limitation ψ' of ψ on H such that $A_1 \subset L'(v_1)$ and $A_2 \subset L'(v_2)$. Here, A_1 and A_2 are disjoint if v_1 and v_2 are adjacent.*

Proof: For each $u \in H$, if $N_G(u) \cap \{v_1, v_2\} = \emptyset$, then to define $\psi'(u)$, we delete colors from $\psi(u)$ arbitrarily if it is necessary. If $|N_G(u) \cap \{v_1, v_2\}| = 1$, say u is adjacent to v_1 , then we note that $h_H(u) \geq h_G(v) + 1/31$. In order to define $\psi'(u)$, we need to delete at least $M/31$ colors from $\psi(u)$. Therefore, we can first remove colors of A_1 from $\psi(u)$ and then delete extra colors if it is necessary. If $|N_G(u) \cap \{v_1, v_2\}| = 2$, then we have $h_H(u) \geq h_G(u) + 2/31$. To define $\psi'(u)$, we remove colors of $A_1 \cup A_2$ from $\psi(u)$ first and delete extra colors if it is necessary. It is easy to see that the limitation defined above is a desired one. \blacksquare

We remark that one of A_1 and A_2 could be the empty set when we apply Fact 1. Next, we show properties that satisfied by the minimal counterexample G .

Fact 2 *Suppose that G contains a subgraph consisting of a triangle uvw and two edges uu', vv' . Let ψ be an $(h_{G'}, M)$ -coloring of $G' = G - \{u, v, w\}$. If $d_G(u) = d_G(v) = 3$ and $|\psi(u') \setminus \psi(v')| \geq (8 - d_G(w))M/31$, then any limitation of ψ on G' can be extended as an (h_G, M) -coloring of G .*

Proof: We choose $A \subset \psi(u') \setminus \psi(v')$ with $|A| = (8 - d_G(w))M/31$. Let ψ' be any limitation of ψ on G' . Then $A \subseteq L'(v)$. Note that $|L'(w)| \geq M - (d_G(w) - 2) \cdot 9M/31$ and $|L'(w) \setminus A| \geq (41 - 8d_G(w))/31 \geq (12 - d_G(w))/31$. We can first extend ψ' to w such that $\psi'(w) \cap A = \emptyset$, then extend ψ' to v such that $A \subseteq \psi'(v)$. Notice that $|L'(u)| \geq M - (12 - d_G(w))M/31 -$

$9M/31 \times 2 + (8 - d_G(w))M/31 = 9M/31$. Thus we can color u properly and get an (h_G, M) -coloring of G , a contradiction. ■

Claim 1 $G + uv$ contains no C_8^2 for any $uv \notin E(G)$.

Proof: Suppose that there exists $uv \notin E(G)$ such that $G + uv$ contains a C_8^2 . Then uv must be contained in this C_8^2 . Assume that $V(C_8^2) = \{v_1, v_2, \dots, v_8\}$, where $v_i v_{i+1}$ and $v_i v_{i+2}$ are edges for each $1 \leq i \leq 8$ while the addition is under module 8. Let G' be the graph obtained from $G - \{v_1, v_2, \dots, v_8\} \setminus \{u, v\}$ by identifying u and v . Note that \underline{uv} has degree at most two in G' . Therefore, G' is K_4 -free and C_8^2 -free. Thus G' has an $(h_{G'}, M)$ -coloring ψ by the definition of G and the fact $|V(G')| < |V(G)|$. Let ψ' be a limitation of ψ on $G - \{v_1, v_2, \dots, v_8\} \setminus \{u, v\}$. Notice that $\min\{d_G(u), d_G(v)\} \geq 3$ and $d_{G'}(\underline{uv}) \leq 2$. We can suppose that $\psi'(u), \psi'(v) \subseteq [1, 9M/31]$. Without loss of generality, we assume that $u = v_1$. Then either $v = v_2$ or $v = v_3$ by the symmetry.

If $v = v_2$, then we extend ψ' as follows. Let $\psi'(v_3) = [1 + 9M/31, 17M/31]$, $\psi'(v_4) = [1 + 17M/31, 25M/31]$, $\psi'(v_5) = [1, 2M/31] \cup [1 + 25M/31, M]$, $\psi'(v_6) = [1 + 3M/31, 11M/31]$, $\psi'(v_7) = [1 + 11M/31, 19M/31]$, and $\psi'(v_8) = [1 + 19M/31, 27M/31]$. We obtained an (h_G, M) -coloring of G , which is a contradiction.

If $v = v_3$, then we extend ψ' as follows. Let $\psi'(v_2) = [1 + 23M/31, M]$, $\psi'(v_4) = [1 + 9M/31, 17M/31]$, $\psi'(v_5) = [1 + 17M/31, 25M/31]$, $\psi'(v_6) = [1, 8M/31]$, $\psi'(v_7) = [1 + 26M/31, M] \cup [1 + 9M/31, 12M/31]$, and $\psi'(v_8) = [1 + 14M/31, 22M/31]$. Thus G has an (h_G, M) -coloring of G , which is a contradiction. ■

Claim 2 $G/\{x, y\}$ contains no C_8^2 for any $xy \notin E(G)$.

Proof: Suppose that $G/\{x, y\}$ contains a C_8^2 for some $xy \notin E(G)$. Then \underline{xy} must be contained in this C_8^2 . Assume that $V(C_8^2) = \{v_1, v_2, \dots, v_8\}$, where $v_i v_{i+1}$ and $v_i v_{i+2}$ are edges for each $1 \leq i \leq 8$ while the addition is under module 8. Without loss of generality, we suppose that $\underline{xy} = v_8$ and $|N_G(x) \cap \{v_1, v_2, v_6, v_7\}| \geq |N_G(y) \cap \{v_1, v_2, v_6, v_7\}|$. Let $G' = G - \{v_1, v_2, \dots, v_7, x\}$. Clearly, G' has an $(h_{G'}, M)$ -coloring ψ by the definition of G . Let ψ' be a limitation of ψ on G' . As x has at most two neighbors in G' , it follows that $|L'(x)| \geq M - 18M/31 = 13M/31$. Notice that $|\psi'(y)| \leq 9M/31$. Thus ψ' can be extended to x satisfying $|\psi'(x) \setminus \psi'(y)| \geq 4M/31$.

For $u \in \{v_1, v_2, v_6, v_7\}$, we get $|L'(u)| \geq M - 9M/31 = 22M/31$ as u has exactly one neighbor in $\{x, y\}$. Moreover, we may assume that either $|L'(v_2) \setminus L'(v_6)| \geq 4M/31$ or $|L'(v_2) \setminus L'(v_7)| \geq 4M/31$ because of $|\psi'(x) \setminus \psi'(y)| \geq 4M/31$.

If $|L'(v_2) \setminus L'(v_6)| \geq 4M/31$, then we choose $A \subseteq L'(v_2) \setminus L'(v_6)$ with $|A| = 4M/31$ and $B \subseteq L'(v_1) \setminus A$ with $|B| = 5M/31$. To define color sets for remaining vertices, we first set $A \subset \psi'(v_2)$, $A \subset \psi'(v_5)$, $B \subset \psi'(v_1)$, and $B \subset \psi'(v_4)$. Then we can complete the extension of ψ' by coloring $v_1, v_2, v_7, v_6, v_5, v_4, v_3$ one by one greedily. When we color v_6 , note that $|L'(v_6) \setminus (\psi'(v_7) \cup A \cup B)| \geq M - 9M/31 - 8M/31 - 5M/31 = 9M/31$ as $A \cap L'(v_6) = \emptyset$. When we color v_3 , notice that $|\cup_{u \in N_G(v_3)} \psi'(u)| \leq 4 \cdot (8M/31) - 9M/31 = 23M/31$. Therefore, we can extend ψ' as an (h_G, M) -coloring of G . This is a contradiction.

If $|L'(v_2) \setminus L'(v_7)| \geq 4M/31$, then let $A \subseteq L'(v_2) \setminus L'(v_7)$ with $|A| = 4M/31$ and $B \subseteq L'(v_6) \setminus A$ with $|B| = 5M/31$. As we did in the previous case, we first set $A \subset \psi'(v_2)$, $A \subset \psi'(v_5)$, $B \subset \psi'(v_3)$, and $B \subset \psi'(v_6)$. We are also able to color vertices $v_1, v_2, v_3, v_7, v_6, v_5, v_4$ one by one greedily. When we color v_7 , note that $|L'(v_7) \setminus (\psi'(v_1) \cup A \cup B)| \geq M - 9M/31 - 8M/31 - 5M/31 = 9M/31$ as $A \cap L'(v_7) = \emptyset$. Thus ψ' can be extended as an (h_G, M) -coloring of G , which is a contradiction. ■

Claim 3 G is P_5^2 -free.




Figure 2: The related graphs in the proof of Claim 3

Proof: Suppose that G contains a P_5^2 . Let v_1, v_2, \dots, v_5 be the vertices of this P_5^2 . Since G contains no K_4 , we have $v_1v_4, v_2v_5 \notin E(G)$.

Case 1: $v_1v_5 \in E(G)$. Let $G' = G - \{v_1, v_2, v_3, v_4, v_5\}$. Clearly, G' has an $(h_{G'}, M)$ -coloring ψ by the definition of G . Let ψ' be a limitation of ψ on G' . Note that $|L'(v_i)| \geq M - 9M/31 = 22M/31$ for $i \in \{1, 2, 4, 5\}$. It follows that $|L'(v_1) \cap L'(v_4)| \geq 13M/31$ and $|L'(v_2) \cap L'(v_5)| \geq 13M/31$. Let $A \subseteq L'(v_1) \cap L'(v_4)$ with $|A| = 6M/31$ and $B \subseteq (L'(v_2) \cap L'(v_5)) \setminus A$ with $|B| = 7M/31$. We set $A \subset \psi'(v_1)$, $A \subset \psi'(v_4)$, $B \subset \psi'(v_2)$ and $B \subset \psi'(v_5)$, which enable us to color vertices v_1, v_5, v_2, v_4 one by one greedily. When we color v_3 , we note that $|\cup_{u \in N_G(v_3)} \psi(u)| \leq 4 \cdot (9M/31) - 13M/31 = 23M/31$. Therefore, we can define $\psi'(v_3)$ properly and extend ψ' as an (h_G, M) -coloring of G . This is a contradiction.

Case 2: $v_1v_5 \notin E(G)$. In this case, we first prove that $d_G(v_i) = 4$ for each $i \in \{1, 2, 4, 5\}$ step by step. We begin to show $d_G(v_2) = d_G(v_4) = 4$. If $d_G(v_2) = 3$, then let $G' = G - \{v_1, v_2, v_3, v_4, v_5\}$. As the definition of G , the graph G' has an $(h_{G'}, M)$ -coloring ψ . Let ψ' be a limitation of ψ on G' . Note that $|L'(v_1)| \geq M - 9M/31 \times 2 = 13M/31$, $|L'(v_5)| \geq M - 9M/31 \times 2 = 13M/31$, and $|L'(v_4)| \geq M - 9M/31 = 22M/31$. It follows that $|L'(v_1) \cap L'(v_4)| \geq 4M/31$. Let $A \subseteq L'(v_1) \cap L'(v_4)$ with $|A| = 4M/31$. We first set $A \subset \psi'(v_1)$ and $A \subset \psi'(v_4)$. Now, we are able to color vertices v_5, v_1, v_4, v_3 one by one greedily. When we color v_2 , we note that $|\cup_{u \in N_G(v_2)} \psi'(u)| \leq 2 \cdot (9M/31) + 8M/31 - 4M/31 = 22M/31$. Therefore, we can define $\psi'(v_2)$ properly and extend ψ' as an (h_G, M) -coloring of G . This is a contradiction. Therefore $d_G(v_2) = 4$. By the symmetry, we can show $d_G(v_4) = 4$.

We next show $d_G(v_1) = d_G(v_5) = 4$. If $d_G(v_1) = 3$, then let u_1 be another neighbor of v_1 and u_2 be another neighbor of v_2 . Let $G' = G - \{v_1, v_2, v_3, v_4, v_5\}$. Then G' has an $(h_{G'}, M)$ -coloring ψ by the definition of G . Let ψ' be a limitation of ψ on G' . If $|\psi'(u_1) \cap \psi'(u_2)| \geq 3M/31$, then we define $A \subseteq \psi'(u_1) \cap \psi'(u_2)$ with $|A| = 3M/31$. We set $A \subset \psi'(v_3)$, which allows us to color vertices v_5, v_4, v_3, v_2, v_1 one by one greedily. This is a contradiction. If $|\psi'(u_1) \cap \psi'(u_2)| < 3M/31$, then there is a set $A \subseteq \psi'(u_1) \setminus \psi'(u_2)$ with $|A| = 3M/31$ and a set $B \subseteq \psi'(u_2) \setminus A$ with $|B| = 2M/31$. We first set $A \subset \psi'(v_2)$ and $B \subset \psi'(v_3)$, then we are able to color vertices v_5, v_4, v_3, v_2, v_1 one by one greedily. This is a contradiction. Therefore, it follows that $d_G(v_1) = 4$. By the symmetry of v_1 and v_5 , we have $d_G(v_5) = 4$. Let us proceed with the assumption $d_G(v_i) = 4$ for each $i \in \{1, 2, 4, 5\}$.

Let G' be the graph obtained from G by deleting v_2, v_3 , and v_5 followed by identifying v_1 and v_4 . Similarly, let G'' be the graph obtained from G by deleting $\{v_2, v_3, v_4\}$ followed by identifying $\{v_1, v_5\}$. Claim 2 implies that both G' and G'' are C_8^2 -free. We further assert that one of G' and G'' is K_4 -free. If it is not the case, then the existence of a K_4 in G'' implies that there is a triangle xyz outside P_5^2 such that $v_1x, v_5y, v_5z \in E(G)$, see the first picture in Figure 2. Similarly, a K_4 in G' indicates that there exists a triangle $x'y'z'$ outside P_5^2 such that $v_1x', v_1y', v_4z' \in E(G)$. Since $\Delta(G) \leq 4$, we may assume that $x' = x$. Note that $\{y, z, v_1\} \subseteq N_G(x)$ and $\{y', z', v_1\} \subseteq N_G(x') = N_G(x)$. Thus we have either $y' = y$ or $y' = z$, say $y' = y$. In this case, $z' = z$ and $v_4z \in E(G)$ (see the second picture in Figure 2), but now $G + v_2x$ contains a C_8^2 (see the last picture in Figure 2). This is a contradiction to

Claim 1 and the assertion follows.

If G' is K_4 -free, then let ψ be an $(h_{G'}, M)$ -coloring of G' . We may assume that $\psi(v_1) = \psi(v_4) = [1, 9M/31]$. We will color neighbors of v_3 carefully such that each color from $[1, 9M/31]$ appears twice in the neighbors of v_3 . This will ensure that we can color v_3 properly. By Fact 1, there is a limitation ψ' of ψ on G' such that $[1, M/31] \subseteq L'(v_2)$ and $[1, M/31] \subseteq L'(v_5)$. If we define $A = L'(v_5) \setminus [1, 9M/31]$, then it is true that $|A| \geq 4M/31$ as $|L'(v_5)| \geq 13M/31$. To define $\psi'(v_5)$, we first put $A \cup [1, M/31] \subset \psi'(v_5)$. In order to make $|\psi'(v_5)| = 8M/31$, we choose at most $3M/31$ colors from $[1 + M/31, 9M/31]$ and add them to $\psi'(v_5)$. To color v_4 , we set $([1 + M/31, 9M/31] \setminus \psi'(v_5)) \subseteq \psi'(v_4)$ and add extra colors greedily such that $[1, M/31] \cap \psi'(v_4) = \emptyset$. Define $\psi'(v_1) = [1 + M/31, 9M/31]$. Note that $|\psi'(v_1) \cap \psi'(v_4)| \geq 3M/31$. It follows that ψ' can be extended to v_2 such that $[1, M/31] \subseteq \psi'(v_2)$. Since each color in $[1, 9M/31]$ appears twice in neighbors of v_3 , then $|L'(v_3)| \geq M - (4 \times 8M/31 - 9M/31) = 8M/31$. Therefore, we can define $\psi'(v_3)$ properly to get an (h_G, M) -coloring of G , a contradiction.

If G'' is K_4 -free, then we next show that G has an (h_G, M) -coloring. Let ψ be an $(h_{G''}, M)$ -coloring of G'' . We may assume that $[1, 8M/31] \subseteq \psi(v_1) = \psi(v_5)$. Consequently, $[1, 8M/31] \subseteq L(v_1)$ and $[1, 8M/31] \subseteq L(v_5)$. Note that $|L(v_1)| \geq 11M/31$, which enables us to choose $A \subseteq L(v_1)$ such that $|A| = M/31$ and $A \cap [1, 8M/31] = \emptyset$. By Fact 1, we can define ψ' as a limitation of $G - \{v_1, v_2, v_3, v_4, v_5\}$ on ψ such that $A \subseteq L'(v_4)$. Observe that $|L'(v_2)| \geq M - 9M/31 = 22M/31$ and $|L'(v_5)| \geq M - 2 \times 9M/31 = 13M/31$. Thus we get $|L'(v_2) \cap L'(v_5)| \geq 4M/31$. Let $B \subseteq L'(v_2) \cap L'(v_5)$ such that $|B| = 2M/31$ and $A \cap B = \emptyset$. To color v_5 , we first put B in $\psi'(v_5)$ and add extra colors from $[1, 8M/31] \setminus B$ to $\psi'(v_5)$ so that $|\psi'(v_5)| = 8M/31$. To color v_1 , we put $(\psi'(v_5) \setminus B) \cup A$ in $\psi'(v_1)$ followed by adding extra colors to $\psi'(v_1)$ so that $|\psi'(v_1)| = 8M/31$ and $\psi'(v_1) \cap B = \emptyset$. Extend ψ' to v_2 such that $B \subseteq \psi'(v_2)$ and $\psi'(v_2) \cap A = \emptyset$. Since $B \subseteq \psi'(v_2) \cap \psi'(v_5)$, we can define $\psi'(v_4)$ properly with $A \subseteq \psi'(v_4)$. Since each color in $A \cup \psi'(v_5)$ appears twice in neighbors of v_3 , then $|L'(v_3)| \geq M - (4 \times 8M/31 - 9M/31) = 8M/31$. We can extend ψ' to v_3 to get an (h_G, M) -coloring of G , a contradiction. ■

Let F_4 be a graph consisting of a triangle uvw and a vertex x adjacent to u and v . We will keep the labelling of vertices of this graph in the rest of this paper.

Claim 4 G contains no F_4 such that $d_G(u) = d_G(v) = d_G(w) = 3$.

Proof: Let $G' = G - \{u, v, w, x\}$. Then G' has an $(h_{G'}, M)$ -coloring ψ . Let ψ' be a limitation of ψ on G' . As $\delta(G) \geq 3$, it follows that either $|\psi'(y)| = 8M/31$ or $|\psi'(y)| = 9M/31$ for each $y \in V(G')$. Note that $|L'(w)| \geq 22M/31$ and $|L'(x)| \geq M - (d_G(x) - 2) \times 9M/31 = (49 - 9d_G(x))M/31$. Then $|L'(w) \cap L'(x)| \geq 22M/31 + (49 - 9d_G(x))M/31 - M = (40 - 9d_G(x))M/31 \geq (8 - d_G(x))M/31$. We can first color w and x such that $|\psi'(x) \cap \psi'(w)| = (8 - d_G(x))M/31$. As $|\psi'(x) \cup \psi'(w)| = 9M/31 + (12 - d_G(x))M/31 - (8 - d_G(x))M/31 = 13M/31$, we can define the coloring of u and v greedily. Thus ψ' can be extended as an (h_G, M) -coloring of G , which is a contradiction and the claim follows. ■

Claim 5 G does not contain F_7 with $d_G(u) = 3$ as a subgraph, where F_7 is shown in Figure 3.

Proof: Suppose that G contains F_7 as a subgraph. By Claim 3, F_7 is an induced subgraph of G , and by Claim 4, $d_G(a) + d_G(b) + d_G(c) \geq 3 + 3 + 4 = 10$. We claim that there is a set coloring ψ' of $G - \{u, v\}$ such that $|\psi'(y)| = h_G(y)M$ for each $y \in V(G) - \{u, v\}$ and $|\psi'(x) \cap \psi'(w)| \geq 3M/31$.

Let $G' = G - \{u, v, w, x\}$. As G is a minimum counterexample, G' has an $(h_{G'}, M)$ -coloring ψ . For each $y \in \{a, b, c\}$, the fact $d_{G'}(y) \leq d_G(y) - 1$ implies that $h_{G'}(y) \geq h_G(y) + 1/31$. There are two cases depending on the degree of w .




Figure 3: The graph F_7

Case 1: $d_G(w) = 4$. As $\{u, v, c\} \subset N_G(w)$, we assume that w' is another neighbor of w . The assumption $d_G(a) + d_G(b) + d_G(c) \geq 10$ implies that $d_{G'}(a) + d_{G'}(b) + d_{G'}(c) \geq 7$ and $|\psi(a) \cup \psi(b) \cup \psi(c)| \leq 29M/31$. Let $A \subseteq M \setminus (\psi(a) \cup \psi(b) \cup \psi(c))$ with $|A| = 2M/31$. Denote $B = A \cap \psi(w')$ and $C = A \setminus B$. Note that $|(\psi(a) \cup \psi(b)) \setminus \psi(w')| \geq 8M/31$.

If $|B| \leq M/31$, then let $D \subseteq (\psi(a) \cup \psi(b)) \setminus \psi(w')$ such that $|D| = M/31$. By Fact 1, we can define a limitation ψ' of ψ on G' such that $B \subseteq L'(w)$ and $D \subseteq L'(x)$. Set $(A \cup D) \subset \psi'(w)$ and $(A \cup D) \subset \psi'(x)$. Then we pick extra colors greedily for $\psi'(w)$ and $\psi'(x)$ to satisfy $|\psi'(w)| = Mh_G(w)$ and $|\psi'(x)| = Mh_G(x)$. We note that $|\psi'(x) \cap \psi'(w)| \geq |A| + |D| = 3M/31$.

If $|B| > M/31$, then let $B' \subseteq B$ such that $|B'| = M/31$. If there is a vertex $y \in \{a, b\}$ such that $|\psi(y) \setminus \psi(w')| < M/31$, say $y = a$ by the symmetry, then there are sets $C \subset \psi(b) \setminus \psi(w')$ and $D \subset \psi(c) \setminus \psi(w')$ with $|C| = |D| = M/31$ as $\psi(a)$, $\psi(b)$, and $\psi(c)$ are pairwise disjoint. To define a limitation ψ' of ψ on G' , we remove colors of C from $\psi(b)$, colors of D from $\psi(c)$, and colors of B' from $\psi(w')$. Thus we have $B' \cup C \cup D \subset L'(w)$ and $B' \cup C \cup D \subset L'(x)$. Set $B' \cup C \cup D \subset \psi'(w)$ and $B' \cup C \cup D \subset \psi'(x)$ in this case. If $|\psi(a) \setminus \psi(w')| \geq M/31$ and $|\psi(b) \setminus \psi(w')| \geq M/31$, then let $C \subset \psi(a) \setminus \psi(w')$ and $D \subset \psi(b) \setminus \psi(w')$ with $|C| = |D| = M/31$. We can define a limitation ψ' of ψ on G' such that $B' \subset L'(w)$ by Fact 1. Moreover, we can assume that $C \cup D \subset L'(x)$ as we can remove colors of C from $\psi(a)$ and colors of D from $\psi(b)$ when we define the limitation. Similarly, set $B' \cup C \cup D \subset \psi'(w)$ and $B' \cup C \cup D \subset \psi'(x)$. Then we pick extra colors greedily for $\psi'(w)$ and $\psi'(x)$ to satisfy $|\psi'(w)| = Mh_G(w)$ and $|\psi'(x)| = Mh_G(x)$. We note that $|\psi'(x) \cap \psi'(w)| \geq |B'| + |C| + |D| = 3M/31$.

Case 2: $d_G(w) = 3$. Let ψ' be a limitation of ψ on $G - \{u, v, w, x\}$. As $|\psi'(a) \cup \psi'(b) \cup \psi'(c)| \leq 27M/31$, we are able to choose $A \subseteq M \setminus (\psi'(a) \cup \psi'(b) \cup \psi'(c))$ with $|A| = 3M/31$. Set $A \subseteq \psi'(w)$ and $A \subseteq \psi'(x)$. Then we pick extra colors greedily for $\psi'(w)$ and $\psi'(x)$ to satisfy $|\psi'(w)| = Mh_G(w)$ and $|\psi'(x)| = Mh_G(x)$. We note that $|\psi'(x) \cap \psi'(w)| \geq 3M/31$.

To finish the coloring, Claim 4 implies that $d_G(w) + d_G(v) \geq 7$. We first color v greedily. Note that $|\psi'(x) \cup \psi'(v) \cup \psi'(w)| \leq 2 \cdot 8M/31 + 9M/31 - 3M/31 = 22M/31$ in each case. Thus we can color u properly and get an (h_G, M) -coloring of G , which is a contradiction. ■

Claim 6 G contains no F_4 such that $d_G(u) = 3$.

Proof: Suppose that G contains an F_4 such that $d_G(u) = 3$. Let G' be the graph obtained from $G - \{u, v\}$ by identifying w and x . Claim 2 tells us that G' contains no C_8^2 . If G' contains a K_4 , then G contains an F_7 , which is a contradiction to Claim 5. Therefore, G' is K_4 -free and G' has an $(h_{G'}, M)$ -coloring ψ . Note that $\psi(w) = \psi(x)$. We can extend ψ as an (h_G, M) -coloring of G by coloring v and u greedily, which is a contradiction. ■

Let F_5 be a graph consisting of an edge uv and three common neighbors a , b , and c .

Claim 7 G contains no F_5 .

Proof: Suppose that G contains an F_5 . As G is K_4 -free, we get that $\{a, b, c\}$ is an independent set. There are three cases depending on the number of three vertices from $\{a, b, c\}$.

Case 1: There are two vertices from $\{a, b, c\}$ having degree three. We can assume that $d_G(a) = d_G(b) = 3$ and $3 \leq d_G(c) \leq 4$. Let $G' = G - \{u, v, a, b, c\}$. Then G' has an $(h_{G'}, M)$ -coloring ψ . Let ψ' be a limitation of ψ on $G - \{u, v, a, b, c\}$. Note that $|L'(a)| \geq 22M/31$, $|L'(b)| \geq 22M/31$, and $|L'(c)| \geq 13M/31$. Thus $|L'(a) \cap L'(b)| \geq 13M/31$, $|L'(a) \cap L'(c)| \geq 4M/31$, and $|L'(b) \cap L'(c)| \geq 4M/31$. Let $A \subseteq L'(a) \cap L'(c)$ with $|A| = 4M/31$ and $B \subseteq L'(b) \cap L'(c)$ such that $|B| = 4M/31$. Moreover, we choose $C \subseteq L(a) \cap L(b)$ such that $|C| = 5M/31$ and $C \cap (A \cup B) = \emptyset$. Set $A \cup C \subset \psi'(a)$, $B \cup C \subset \psi'(b)$, and $A \cup B \subset \psi'(c)$. Then ψ' can be extended as an (h_G, M) -coloring of G by choosing colors for a, b, c, u, v one by one greedily, which is a contradiction.

Case 2: There is exactly one vertex of $\{a, b, c\}$ with degree three. Then we assume that $d_G(a) = 3$ and $d_G(b) = d_G(c) = 4$. Let G' (and G'') be the graph obtained from $G - \{u, v, c\}$ (and $\{u, v, b\}$) by identifying a and b (by identifying a and c) respectively. Note that both G' and G'' are C_8^2 -free by Claim 2. There are two subcases.

Subcase 2.1: One of G' and G'' is K_4 -free. By the symmetry, we can assume that G' is K_4 -free. Thus G' has an $(h_{G'}, M)$ -coloring ψ . Note that \underline{ab} is a 3-vertex of G' . Then we may assume that $\psi(\underline{ab}) = [1, 9M/31]$. Let $A = [1, M/31]$. By Fact 1, we can let ψ' be a limitation of ψ on $G - \{u, v, a, b, c\}$ such that $A \subseteq L'(c)$. Note that $[1, 9M/31] \subseteq L'(a) \cap L'(b)$, $|L'(a)| \geq 22M/31$, and $|L'(c)| \geq 13M/31$. It follows that $|L'(a) \cap L'(c)| \geq 4M/31$. Choose $B \subset L'(a) \cap L'(c)$ such that $|B| = M/31$ and $A \cap B = \emptyset$. Define $\psi'(b) = [1, 8M/31]$. We put $[1, 8M/31] \cup B \subseteq \psi'(a)$ and $A \cup B \subseteq \psi'(c)$ followed by adding extra colors to $\psi'(a)$ and $\psi'(c)$ greedily. We can construct an (h_G, M) -coloring of G by choosing colors for u and v properly. A contradiction.

Subcase 2.2: Both G' and G'' contain a K_4 . In this case, by the symmetry, there are vertices x, y , and z outside F_5 such that $\{xy, xz, yz, ax, yb, yc, zb, zc\} \subset E(G)$. Since $\Delta(G) \leq 4$ and G is 2-connected, we have G is the graph F_8 as shown in Figure 4. It is easy to see that G has an $(h_G, 31)$ -coloring. A contradiction.




Figure 4: The graph F_8 and an $(h_{F_8}, 31)$ -coloring of F_8

Case 3: $d_G(a) = d_G(b) = d_G(c) = 4$. Let G' be the graph obtained from G by deleting vertices from $\{u, v, c\}$ followed by identifying vertices from $\{a, b\}$. Similarly, let G'' (and G''') be the graph obtained from G by deleting vertices from $\{u, v, b\}$ (and $\{u, v, a\}$) followed by identifying vertices from $\{a, c\}$ (and vertices from $\{b, c\}$) respectively. Note that Claim 2 implies that all of them are C_8^2 -free. We proceed to complete the proof with the assumption that one of G' , G'' , and G''' is K_4 -free. By the symmetry, we suppose that G' is K_4 -free. Let ψ be an $(h_{G'}, M)$ -coloring of G' . We assume that $\psi'(a) = \psi'(b) = [1, 8M/31]$ and fix $C \subset [1, 8M/31]$ with $|C| = M/31$. By Fact 1, we can let ψ' be a limitation of ψ on $G - \{u, v, a, b, c\}$ such that $C \subseteq L'(c)$. Set $C \subset \psi'(c)$ and add extra colors greedily. Now,

$|\psi'(a) \cup \psi'(b) \cup \psi'(c)| \leq 8M/31 + 7M/31 = 15M/31$ and we can choose colors for u and v greedily to get an (h_G, M) -coloring of G , a contradiction.

It remains to show the assertion that one of G' , G'' , and G''' is K_4 -free. If G' contains a K_4 , then there is a triangle with vertex set $\{x, y, z\}$ outside F_5 . By the symmetry, we can assume that $\{ax, by, bz\} \subset E(G)$. Since $G[\{a, b, x, y, z\}]$ is not a P_5^2 by Claim 3, we have $ay, az \notin E(G)$. If $cx \in E(G)$, then G'' is K_4 -free as G is P_5^2 -free. Now we assume that $cx \notin E(G)$. The existence of a K_4 in G''' implies that there exists x' such that $cx', x'y, x'z \in E(G)$. Now we can observe that G'' is K_4 -free. The assertion is proved and the claim follows. ■

Claim 8 G contains no triangle uvw such that $d_G(u) = d_G(v) = d_G(w) = 3$.

Proof: Suppose that G contains a triangle uvw with $d_G(u) = d_G(v) = d_G(w) = 3$. Let u', v' , and w' be the neighbors of u, v, w which is not on the triangle, respectively. Claim 6 implies that u', v' , and w' are distinct. Let $G' = G - \{u, v, w\}$. By Claim 1, $G' + u'v'$, $G' + u'w'$, and $G' + v'w'$ are C_8^2 -free. We claim that either $G' + u'v'$, or $G' + u'w'$, or $G' + v'w'$ is K_4 -free. Otherwise, either G contains a K_4 , or there exists $xy \in E(G')$ such that $\{x, y\}$ together with each of u', v' , and w' form a triangle, which is a contradiction to Claim 7.

Assume by the symmetry that $G' + u'v'$ is K_4 -free. As the definition of G , the graph $G' + u'v'$ has an $(h_{G'}, M)$ -coloring ψ . Since $\psi(u') \cap \psi(v') = \emptyset$, any limitation ψ' of ψ on G' can be extended as an (h_G, M) -coloring of G by Fact 2, a contradiction. ■

Claim 9 G contains no path $u_1u_2u_3$ such that $d_G(u_1) = d_G(u_2) = d_G(u_3) = 3$.

Proof: Suppose that G contains a path $u_1u_2u_3$ such that $d_G(u_1) = d_G(u_2) = d_G(u_3) = 3$. By Claim 8, we have $u_1u_3 \notin E(G)$. For $i \in \{1, 3\}$, we assume that u'_i and u''_i are the other two neighbors of u_i .

If $d_G(u'_1) + d_G(u''_1) = d_G(u'_3) + d_G(u''_3) = 6$, then let G' be the graph obtained from G by deleting u_2 followed by identifying u_1 and u_3 . As $d_G(u'_1) + d_G(u''_1) = d_G(u'_3) + d_G(u''_3) = 6$ and Claim 8, the graph G' is K_4 -free. Claim 2 implies that G' is also C_8^2 -free. Thus G' has an $(h_{G'}, M)$ -coloring ψ . We can easily extend ψ as an (h_G, M) -coloring of G , a contradiction.

If either $d_G(u'_1) + d_G(u''_1) \geq 7$ or $d_G(u'_3) + d_G(u''_3) \geq 7$, then by the symmetry, we assume that $d_G(u'_1) + d_G(u''_1) \geq 7$. Let $G' = G - \{u_1, u_2, u_3\}$. Then G' has an $(h_{G'}, M)$ -coloring ψ and let ψ' be a limitation of ψ on $G - \{u_1, u_2, u_3\}$. Note that $|L'(u_1)| \geq 14M/31$, $|L'(u_2)| \geq 22M/31$, and $|L'(u_3)| \geq 13M/31$, here we applied the assumption $d_G(u'_1) + d_G(u''_1) \geq 7$. Let $A \subseteq L'(u_2) \setminus L'(u_3)$ such that $|A| = 5M/31$. We first extend ψ' to u_1 such that $\psi'(u_1) \cap A = \emptyset$ followed by extending ψ' to u_2 such that $A \subseteq \psi'(u_2)$. Since $|L'(u_3) \setminus \psi'(u_2)| \geq 9M/31$, we can extend ψ' to u_3 and get an (h_G, M) -coloring of G , a contradiction. ■

Claim 10 G contains no triangle uvw such that $d_G(u) = d_G(v) = 3$.

Proof: Suppose that G contains a triangle uvw such that $d_G(u) = d_G(v) = 3$. Then $d_G(w) = 4$ by Claim 8. Let u' and v' be the neighbors of u and v not on the triangle, respectively. Claim 6 gives us that $u' \neq v'$, and $wu', wv' \notin E(G)$. Claim 9 implies that $d_G(u') = d_G(v') = 4$.

If $u'v' \in E(G)$, then let $G' = G - \{u, v\}$. Clearly, G' has an $(h_{G'}, M)$ -coloring ψ , where $\psi(u') \cap \psi(v') = \emptyset$. Thus a limitation of ψ on G' can be extended as an (h_G, M) -coloring of G by Fact 2, a contradiction. We need only to consider the case where $u'v' \notin E(G)$. Let $G' = G - \{u, v\} + u'v'$. By Claim 1, G' is C_8^2 -free. If G' is also K_4 -free, then G' has an $(h_{G'}, M)$ -coloring ψ . Since $\psi(u') \cap \psi(v') = \emptyset$, a limitation of ψ on G' can be extended as an (h_G, M) -coloring of G by Fact 2, a contradiction. Therefore, G' contains a K_4 , which implies that there exist $xy \in E(G)$ such that $u'x, u'y, v'x, v'y \in E(G)$. If $u'w \in E(G')$, then

G contains F_7 as a subgraph, which is a contradiction to Claim 5. Thus $u'w \notin E(G')$. By the symmetry, we also have $v'w \notin E(G')$.

We first assume that $wx \in E(G)$. Claim 7 gives that $wy \notin E(G)$. Let w' be the other neighbor of w . Without raising any confusion, we define $G' = G - \{u, v, w\} + v'w'$. By Claim 1, G' is C_8^2 -free. Note that G' is also K_4 -free by checking degrees of neighbors of v' . Thus G' has an $(h_{G'}, M)$ -coloring ψ . Let ψ' be a limitation of ψ on $G - \{u, v, w\}$. If $|\psi'(u') \setminus \psi'(v')| \geq 4M/31$, then ψ' can be extended as an (h_G, M) -coloring of G by Fact 2, a contradiction. Therefore $|\psi'(u') \setminus \psi'(v')| < 4M/31$, which implies $|\psi'(u') \cap \psi'(v')| > 4M/31$. Let $B = \psi'(u') \cap \psi'(v')$ with $|B| = 4M/31$. We first put $B \subset \psi'(w)$ and add extra colors greedily to ensure $|\psi'(w)| = 8M/31$. This is possible because $u'x \in E(G)$ and $w'v' \in E(G')$. Then we can define $\psi'(u)$ and $\psi'(v)$ one by one greedily. In this case, we can construct an (h_G, M) -coloring of G , which is a contradiction.

Now we can assume that $wx \notin E(G)$. By the symmetry, we have $wy \notin E(G)$. Let w_1 and w_2 be the neighbors of w not on the triangle. Define $G' = G - \{u, v, w\} + w_1u' + w_2v'$. We first claim that G' is K_4 -free. Otherwise, this K_4 must contain exactly one of the edges $\{w_1u', w_2v'\}$ as $u'v' \notin E(G)$. We can assume that K_4 contains the edge w_1u' . As $\Delta(G) \leq 4$, the only possibility for the vertex set of this K_4 is $\{w_1, u', x, y\}$. Now x and y have three common neighbors u', v' and w_1 , which is a contradiction to Claim 7.

If G' contains no C_8^2 , then G' has an $(h_{G'}, M)$ -coloring ψ . Let ψ' be a limitation of ψ on $G - \{u, v, w\}$. If $|\psi'(v') \cap \psi'(u')| \geq 3M/31$, then let $A \subset \psi'(v') \cap \psi'(u')$ with $|A| = 3M/31$. This is possible as $w_1u', w_2v' \in E(G')$. We put A in $\psi'(w)$ and add extra colors to $\psi'(w)$ such that $|\psi'(w)| = 8M/31$. Then ψ' can be extended as an (h_G, M) -coloring of G by defining $\psi'(u)$ and $\psi'(v)$ one by one greedily. If $|\psi'(v') \cap \psi'(u')| < 3M/31$, then $|\psi'(v') \setminus \psi'(u')| > 5M/31$. By Fact 2, ψ' can be extended to get an (h_G, M) -coloring of G , a contradiction.

Now we assume that G' contains a C_8^2 . By Claim 1, this C_8^2 must contain edges w_1u' and w_2v' . We assume that $N_G(u') = \{a, w_1, x, y\}$ and $N_G(v') = \{b, w_2, x, y\}$. The fact that all vertices in G' have degree at most four implies that G' is the whole graph C_8^2 . Since $G[\{a, u', v', x, y\}]$ contains no P_5^2 as a subgraph by Claim 3, it follows that $ax, ay \notin E(G)$. Then we get $aw_1, aw_2, ab \in E(G)$. By the symmetry of a and b , we have $bw_1, bw_2 \in E(G)$. Therefore, G must be the graph F_{11} as shown in Figure 5. However, F_{11} has an $(h_{F_{11}}, 31)$ -coloring as shown by Figure 5, a contradiction.

The proof of the claim is complete. ■




Figure 5: The graph F_{11} and an $(h_{F_{11}}, 31)$ -coloring of F_{11}

Claim 11 G contains no path $u_1u_2u_3u_4$ such that $d_G(u_1) = d_G(u_2) = d_G(u_4) = 3$.

Proof: Suppose that G contains a path $u_1u_2u_3u_4$ such that $d_G(u_1) = d_G(u_2) = d_G(u_4) = 3$. Claim 10 implies that $u_1u_2u_3u_4$ is an induced path. By Claim 9, $d_G(u_3) = 4$ while other neighbors of u_1 and u_2 not in this path are 4-vertices.

Let $G' = G - \{u_1, u_2\}$. Clearly, G' has an $(h_{G'}, M)$ -coloring ψ . Suppose that ψ' is a limitation of ψ on $G - \{u_1, u_2, u_3, u_4\}$. Note that $|L'(u_1)| = 15M/31$, $|L'(u_2)| = 23M/31$, $|L'(u_3)| \geq 13M/31$, and $|L'(u_4)| \geq 13M/31$. Moreover, $|L'(u_3) \cup L'(u_4)| \geq 17M/31$ as it contains $\psi(u_3) \cup \psi(u_4)$ as a subset. By deleting colors not in $\psi(u_3) \cup \psi(u_4)$ if it necessary, we may assume that $|L'(u_3)| = |L'(u_4)| = 13M/31$. Then $|L'(u_3) \setminus L'(u_4)| \geq 4M/31$ because of $|L'(u_3) \cup L'(u_4)| \geq 17M/31$. Let $A \subseteq L'(u_3) \setminus L'(u_4)$ and $B \subseteq L'(u_2) \setminus L'(u_1)$ such that $|A| = 4M/31$, $|B| = 3M/31$, and $A \cap B = \emptyset$. Put $A \subset \psi'(u_3)$ and $B \subset \psi'(u_2)$. Choosing extra colors for u_3, u_2, u_1 and u_4 one by one greedily, we obtain an (h_G, M) -coloring of G , a contradiction. ■

Claim 12 G contains no F_4 such that $d_G(w) = d_G(x) = 3$.

Proof: Suppose that G contains an F_4 such that $d_G(w) = d_G(x) = 3$. Claim 10 indicates that $d_G(u) = d_G(v) = 4$. For each $z \in \{u, v, w, x\}$, let z' be the neighbor of z which is not in F_4 . Claim 11 implies that $d_G(w') = d_G(x') = 4$. Moreover, u', v' , and w' are all distinct by Claim 3 and Claim 7. Similarly, u', v' , and x' are distinct. Let $G' = G - \{u, v, w, x\}$. Then G' is C_8^2 -free and K_4 -free. Thus G' has an $(h_{G'}, M)$ -coloring ψ . There are two cases depending on the size of $\psi(w') \cap \psi(x')$.

Case 1: $|\psi(w') \cap \psi(x')| \geq 2M/31$. Let $A \subseteq (\psi(w') \cap \psi(x'))$ and $B \subseteq (\psi(w') \cap \psi(x'))$ such that $A \cap B = \emptyset$ and $|A| = |B| = M/31$. By Fact 1, there exists a limitation ψ' of ψ on $G - \{u, v, w, x\}$ such that $A \subseteq L'(u)$ and $B \subseteq L'(v)$. Therefore, it is proper to put A in $\psi'(u)$ and B in $\psi'(v)$. We add extra colors to $\psi'(u)$ and $\psi'(v)$ such that $|\psi'(u)| = |\psi'(v)| = 8M/31$. Then ψ' can be extended as an (h_G, M) -coloring of G by picking colors for $\psi'(w)$ and $\psi'(x)$ one by one greedily. Thus we get an (h_G, M) -coloring of G , which is a contradiction.

Case 2: $|\psi(w') \cap \psi(x')| < 2M/31$. We have $w' \neq x'$ in this case. Note that $|\psi(w') \cup \psi(x')| > 16M/31$ and $|L(u)| \geq 21M/31$. It follows that $|L(u) \cap (\psi(w') \cup \psi(x'))| > 6M/31$. By the symmetry, we may assume that $|L(u) \cap \psi(w')| > 3M/31$. Let $A \subset \psi(x')$ and $B \subset \psi(x')$ such that $A \cap B = \emptyset$ and $|A| = |B| = M/31$. Additionally, we choose $C \subset L(u) \cap \psi(w')$ with $|C| = 3M/31$. Fact 1 gives a limitation ψ' of ψ on $G - \{u, v, w, x\}$ with the property $A \subset L'(u)$ and $B \subset L'(v)$. Furthermore, we have $C \subset L'(u) \cap \psi'(w')$ by deleting colors from $\psi(w')$ properly. To color u and v , we put $A \cup C$ in $\psi'(u)$ and B in $\psi'(v)$ followed by adding extra colors greedily. As the selection of sets A, B , and C , we are able to color x and w one by one, and we obtained an (h_G, M) -coloring of G . A contradiction. ■

Claim 13 G contains no F_4 such that $d_G(x) = 3$.

Proof: Suppose that G contains an F_4 such that $d_G(x) = 3$. Claim 10 and Claim 12 yield that $d_G(u) = d_G(v) = d_G(w) = 4$. For each $y \in \{u, v, x\}$, let y' be another neighbor of y not in F_4 . By Claim 3 and Claim 7, u', v' , and x' are distinct. Similarly, if w' and w'' are neighbors of w not in F_4 , then u', v', w', w'' are distinct.

It is easy to see that either $G - \{u, v, w, x\} + u'x'$ or $G - \{u, v, w, x\} + v'x'$ is K_4 -free. Assume that $G' = G - \{u, v, w, x\} + u'x'$ is K_4 -free. Claim 1 implies that G' is also C_8^2 -free. Thus G' has an $(h_{G'}, M)$ -coloring ψ . Let $A \subseteq \psi(u')$ and $B \subseteq \psi(u')$ such that $A \cap B = \emptyset$ and $|A| = |B| = M/31$, and let $C \subseteq \psi(x')$ with $|C| = 3M/31$. Note that A, B , and C are pairwise disjoint. By Fact 1, we may assume that ψ' is a limitation of ψ on $G - \{u, v, w, x\}$ such that $A \subseteq L'(w)$ and $B \subseteq L'(v)$. Put A in $\psi'(w)$, B in $\psi'(v)$, and C in $\psi'(u)$. Choosing extra colors for w, v, u and x one by one greedily, we obtain an (h_G, M) -coloring of G , a contradiction. ■

Claim 14 G contains no triangle uvw such that $d_G(u) = 3$.

Proof: Suppose that G contains a triangle uvw such that $d_G(u) = 3$. By Claim 10, we have $d_G(v) = d_G(w) = 4$. Let x be the neighbor of u not on the triangle. Claim 6 implies that

$vx, wx \notin E(G)$. Suppose that $N_G(w) = \{u, v, w_1, w_2\}$ and $N_G(v) = \{u, w, v_1, v_2\}$. By Claim 13, vertices v_1, v_2, w_1 and w_2 are all distinct.

Case 1: $d_G(x) = 3$. Claim 11 implies that $d_G(v_1) = d_G(v_2) = d_G(w_1) = d_G(w_2) = 4$. Denote $N_G(x) = \{u, y, z\}$ and $G' = G - \{v, w, u, x\}$. Then G' has an $(h_{G'}, M)$ -coloring ψ . If $|L(w) \cap L(x)| \geq 2M/31$, then we choose $A \subseteq L(w) \cap L(x)$ such that $|A| = 2M/31$. Let $B \subseteq L(x) \setminus A$ such that $|B| = M/31$ as well as $C \subset \psi(w_1) \cup \psi(w_2)$ such that $|C| = M/31$ and $C \cap A = C \cap B = \emptyset$. By Fact 1, there is a limitation ψ' of ψ on G' such that $B \subset L'(w)$ and $C \subset L'(v)$. We set $C \subset \psi'(v)$, $A \cup B \subset \psi'(w)$, and $A \cup B \subset \psi'(x)$. Then we add colors to $\psi'(v)$, $\psi'(w)$, and $\psi'(x)$ one by one greedily. Now we can define $\psi'(u)$ properly because each color in $A \cup B$ appears at least twice in neighbors of u . A contradiction.

Let us assume that $|L(w) \cap L(x)| < 2M/31$. Note that $|L(w)| \geq 13M/31$ and $|L(x)| \geq 13M/31$. We assert that y, z, w_1 , and w_2 are distinct. Otherwise, suppose that $y = w_1$. Then $L(w) \cup L(x) \subset M \setminus \psi(w_1)$ and $|L(w) \cup L(x)| \leq 22M/31$. This implies that $|L(w) \cap L(x)| \geq 2M/31$, which is a contradiction and the assertion follows. Since $\psi(y) \cup \psi(z) \cup L(x) = M$, we get $|L(w) \cap (\psi(y) \cup \psi(z))| > 11M/31$. This implies that there are $A_1 \subseteq L(w) \cap \psi(y)$ and $A_2 \subseteq L(w) \cap \psi(z)$ such that $|A_1| = |A_2| = M/31$ and $A_1 \cap A_2 = \emptyset$. We fix $B \subseteq L(x)$ such that $|B| = M/31$ and $B \cap (A_1 \cup A_2) = \emptyset$. In addition, we choose $C \subset \psi(w_1) \cup \psi(w_2)$ such that $|C| = M/31$ and $C \cap (A_1 \cup A_2 \cup B) = \emptyset$. By Fact 1, there is a limitation ψ' of ψ on G' such that $B \subset L'(w)$ and $C \subset L'(v)$. Deleting colors of A_1 first for $\psi'(y)$ and colors of A_2 first for $\psi'(z)$, we can assume further $A_1 \cup A_2 \subset L'(x)$. We put $A_1 \cup A_2 \cup B \subset \psi'(x)$, $A_1 \cup A_2 \cup B \subset \psi'(w)$, and $C \subset \psi'(v)$ followed by adding colors to $\psi'(v)$, $\psi'(w)$, and $\psi'(x)$ one by one greedily. As the choices of A_1, A_2 , and B , we can color u properly to get an (h_G, M) -coloring of G , a contradiction.

Case 2: $d_G(x) = 4$. If each vertex from $\{v_1, v_2, w_1, w_2\}$ have degree four, then let $G' = G - \{u, v, w\}$. Suppose that ψ is an $(h_{G'}, M)$ -coloring of G' . If $|\psi(x) \cap (\psi(v_1) \cup \psi(v_2))| \geq M/31$, then let $A \subset \psi(x) \cap (\psi(v_1) \cup \psi(v_2))$ with $|A| = M/31$. We choose $B \subset \psi(x)$ such that $A \cap B = \emptyset$ and $|B| = M/31$. By Fact 1, there is a limitation ψ' of ψ on G' such that $A \subset L'(w)$ and $B \subset L'(v)$. Clearly, $A \cup B \subset \psi'(x)$. We set $A \subset \psi'(w)$ and $B \subset \psi'(v)$ followed by adding colors to $\psi'(w)$, $\psi'(v)$, and $\psi'(u)$ one by one greedily. Then we can define $\psi'(u)$ properly because of sets A and B . We obtain an (h_G, M) -coloring of G , a contradiction. Thus we need only to consider the case where $|\psi(x) \cap (\psi(v_1) \cup \psi(v_2))| < M/31$. In this case, $|L(v) \cap \psi(x)| > 8M/31$ as $L(v) \cup (\psi(v_1) \cup \psi(v_2)) = M$. We choose $C \subset (\psi(w_1) \cup \psi(w_2))$ such that $|C| = M/31$. By Fact 1, there is a limitation ψ' of ψ on G' such that $C \subseteq L'(v)$. We define $\psi'(v)$ and $\psi'(x)$ such that $C \subset \psi'(v)$ and $|\psi'(v) \cap \psi'(x)| \geq 2M/31$. To complete the extension of ψ' , we pick colors for w and u one by one. Therefore, we can construct an (h_G, M) -coloring for G , which is a contradiction.

For the case where one of v_1, v_2, w_1 , and w_2 has degree three, we suppose that $d_G(v_1) = 3$. Let v'_1 and v''_1 be the other two neighbors of v_1 . Then $d_G(v'_1) = d_G(v''_1) = 4$ by Claim 11. Let $G' = G - \{u, v, w, v_1\}$ and ψ be an $(h_{G'}, M)$ -coloring of G' . If $|\psi(v'_1) \cap \psi(v''_1)| \geq 2M/31$, then we fix $A \subset \psi(v'_1) \cap \psi(v''_1)$ with $|A| = 2M/31$. Let $B \subset \psi(x)$ and $C \subset \psi(x)$ such that $|B| = |C| = M/31$ as well as A, B, C are pairwise disjoint. By Fact 1, there is a limitation ψ' of ψ on $G - \{u, v, w, v_1\}$ such that $B \subset L'(w)$ and $C \subset L'(v)$. Furthermore, it satisfies that $B \cup C \subset \psi'(x)$ and $A \subset \psi'(v'_1) \cap \psi'(v''_1)$. We put $B \subset \psi'(w)$ and $C \subset \psi'(v)$ followed by adding colors to $\psi'(w)$ and $\psi'(v)$ one by one greedily. To complete the extension, we color v_1 and u one by one greedily. We obtain an (h_G, M) -coloring of G , a contradiction.

If $|\psi(v'_1) \cap \psi(v''_1)| < 2M/31$, then we choose $B, C \subset \psi(x)$ as above. Additionally, we fix $D \subset (\psi(v'_1) \cup \psi(v''_1)) \setminus \psi(v_2)$ such that $|D| = 2M/31$ as well as B, C , and D are pairwise disjoint. As we did above, we can define a limitation ψ' of ψ on $G - \{u, v, w, v_1\}$ such that $B \cup C \subset \psi'(x)$, $B \subset L(w)$, and $C \cup D \subset L'(v)$. We add extra colors to $\psi'(w)$ and $\psi'(v)$ one by one greedily. Then we can get an (h_G, M) -coloring for G by picking colors for v_1 and u greedily, which is a contradiction. ■

Claim 15 G contains no F_4 such that u has a 3-neighbor.

Proof: Claim 14 implies that each vertex from $\{u, v, w, x\}$ is a 4-vertex. Suppose that $N_G(u) = \{w, v, x, u'\}$ and $N_G(v) = \{w, v, x, v'\}$, where $d_G(u') = 3$. Claim 7 gives us that $u' \neq v'$. Moreover, Claim 3 implies that $wu', wv', xu', xv' \notin E(G)$. Let G' be the graph obtained from $G - \{u, x, w\}$ by identifying u' and v . Claim 2 together with Claim 13 yield that G' is C_8^2 -free and K_4 -free. Therefore, G' has an $(h_{G'}, M)$ -coloring ψ , where we assume $\psi(\underline{u'v}) = [1, 9M/31]$. We define $A = [1 + 8M/31, 9M/31]$.

If either $d_G(v') = 4$ or $|L(w) \cap L(x)| \geq 2M/31$, then in the former case, there is a limitation ψ' of ψ on $G - \{u, v, w, x\}$ with $A \subseteq L'(w) \cap L'(x)$ by Fact 1. Note that $[1, 8M/31] \subset L'(v)$. We put A in $\psi'(x)$ and $\psi'(w)$ followed by adding extra colors to $\psi'(x)$ and $\psi'(w)$ greedily. Let $B = [1, 8M/31] \setminus (\psi'(x) \cup \psi'(w))$. We put $B \subset \psi'(v)$ and choose extra colors greedily. This is possible because of $d_G(v') = 4$ and the definition of A . Now, we are able to color u properly because each color from $[1, 9M/31]$ appears twice in the neighbors of u . In the latter case, let ψ' be a limitation of ψ on $G - \{u, v, w, x\}$ such that $A \subseteq L'(w) \cap L'(x)$ and $|L'(w) \cap L'(x)| \geq 2M/31$. To color x and w , we set $A \subset \psi'(x)$ and $A \subset \psi'(w)$ followed by choosing extra colors satisfying $|\psi'(x) \cap \psi'(w)| \geq 2M/31$. Similarly, let $B = [1, 8M/31] \setminus (\psi'(x) \cup \psi'(w))$. We put $B \subset \psi'(v)$ and pick additional colors for $\psi'(v)$. We are able to construct an (h_G, M) -coloring for G by picking colors for u greedily, which is a contradiction.

If $d_G(v') = 3$ and $|L(w) \cap L(x)| < 2M/31$, then let ψ' be a limitation of ψ on $G - \{u, v, w, x\}$ such that $A \subseteq L'(w) \cap L'(x)$. Note that $|(L'(w) \cup L'(x)) \cap \psi'(v')| \geq M/31$. We choose $B \subseteq (L'(w) \cup L'(x)) \cap \psi'(v')$ with $|B| = M/31$ and denote $B_1 = B \cap L'(x)$ as well as $B_2 = B \cap L'(w)$. We put $A \cup B_1 \subset \psi'(x)$ and $A \cup B_2 \subset \psi'(w)$ followed by adding extra colors to $\psi'(x)$ and $\psi'(w)$ greedily. Similarly, to color v , we first put $[1, 9M/31] \setminus (\psi'(w) \cup \psi'(x))$ in $\psi'(v)$ and choose extra colors greedily. This is possible because of sets A , B_1 , and B_2 . Then we can get an (h_G, M) -coloring for G by coloring u greedily, which is a contradiction. ■

Claim 16 The graph G contains no F_7 , see Figure 3.

Proof: Suppose that G contains an F_7 . Claim 3 tells us that F_7 is an induced subgraph of G . Furthermore, each vertex in F_7 is a 4-vertex by Claim 14. For $z \in \{u, v, a, b\}$, let z' be the neighbor of z not in F_7 . Claim 15 gives us that u', v', a' , and b' are 4-vertices. Moreover, Claim 7 implies that $u' \neq v'$ and $a' \neq b'$.

Let G' be the graph obtained from $G - \{x, v, b\}$ by identifying u and a . Clearly, Claim 2 and Claim 3 tell us that G' is K_4 -free and C_8^2 -free. Thus G' has an $(h_{G'}, M)$ -coloring ψ . Suppose that $\psi(\underline{ua}) = [1, 8M/31]$. We define the limitation ψ' of ψ on $G - \{x, u, v, a, b\}$ with some restrictions. Through removing extra colors properly, we can assume that $|L'(u)| = |L'(a)| = 15M/31$, $[1, 8M/31] \subset L'(u)$, and $[1, 8M/31] \subset L'(a)$. Let A be a set such that $|A| = M/31$ and $A \cap (L'(u) \cup L'(a)) = \emptyset$. We can ensure $A \subseteq L'(v) \cap L'(b)$ by deleting colors of A from $\psi(v')$ and $\psi(b')$ first. Similarly, we can assume that $|L'(v)| = |L'(b)| = 15M/31$ by removing colors. We first set $A \subset \psi'(v)$ and then define $\psi'(u)$ and $\psi'(v)$ such that $[1, 8M/31] \subset \psi'(u) \cup \psi'(v)$. If $|[1, 8M/31] \cap L'(v)| \leq 7M/31$, then we set $A \cup ([1, 8M/31] \cap L'(v)) \subset \psi'(v)$ and choose extra colors for $\psi'(v)$ greedily. Note that $|L'(u) \setminus \psi'(v)| \geq 8M/31$ and we can define $\psi'(u)$ properly by including colors from $[1, 8M/31] \setminus \psi'(v)$ first. If $|[1, 8M/31] \cap L'(v)| > 7M/31$, then let $B \subseteq [1, 8M/31] \cap L'(v)$ such that $|B| = 7M/31$. Set $\psi'(v) = A \cup B$. To define $\psi'(u)$, we first put $[1, 8M/31] \setminus \psi'(v)$ in $\psi'(u)$ and add extra colors to $\psi'(u)$ such that $|\psi'(u)| = 8M/31$. This is possible because of $|L'(u) \setminus B| = 8M/31$. We define $\psi'(a)$ and $\psi'(b)$ similarly. As each color from $[1, 8M/31] \cup A$ appears twice in the neighbors of x , we can color x properly. Therefore, we get an (h_G, M) -coloring for G , a contradiction. ■

Claim 17 There are no two triangles sharing an edge.

Proof: Suppose that G contains a subgraph consisting of a triangle uvw and a vertex x adjacent to u and v . By Claim 14, each vertex from $\{u, v, w, x\}$ is a 4-vertex. Let u' and v' be the other neighbor of u and v respectively. Claim 7 tells us that $u' \neq v'$ and Claim 15 implies that $d_G(u') = d_G(v') = 4$. Let G' be a graph obtained from G by deleting u and v followed by identifying w and x . Claim 16 together with Claim 2 yield that G' is both K_4 -free and C_8^2 -free. Thus G' has $(h_{G'}, M)$ -coloring ψ . Note that $|\psi(\underline{wx})| = 8M/31$ and $|\psi(u')| = 9M/31$. Let $A \subset \psi(u') \setminus \psi(\underline{wx})$ such that $|A| = M/31$. By Fact 1, we may let ψ' be a limitation of ψ on $G' - \{u, v, w, x\}$ such that $A \subseteq L'(v)$. Set $\psi'(w) = \psi(\underline{wx})$ and $\psi'(x) = \psi(\underline{wx})$. In order to define $\psi'(v)$, we first put A in $\psi'(v)$ and followed by adding extra colors greedily. As $d_G(u) = 4$ and the definition of A , we can color u properly to get an (h_G, M) -coloring of G , a contradiction. ■

A 4-vertex v is *bad* if it is contained in a triangle and has two neighbors with degree three.

Claim 18 G contains no bad 4-vertex.

Proof: Suppose that G contains a bad 4-vertex v . Denote $N_G(v) = \{v_1, v_2, v_3, v_4\}$. Suppose that $d_G(v_1) = d_G(v_2) = 3$. By Claim 14, both $\{v_1, v_2, v_3\}$ and $\{v_1, v_2, v_4\}$ are independent sets. Then we have $v_3v_4 \in E(G)$ and $d_G(v_3) = d_G(v_4) = 4$. For each $1 \leq i \leq 4$, let v'_i and v''_i be the other neighbors of v_i . By Claim 11 and Claim 14, we get $d_G(v'_i) = d_G(v''_i) = 4$ for $1 \leq i \leq 2$. As Claim 17 shows that there are no two triangles sharing an edge, we get that vertices v'_3, v''_3, v'_4 , and v''_4 are distinct.

Let G' be the graph obtained from $G - v$ by identifying v_1 and v_2 . Then G' is C_8^2 -free and K_4 -free by Claim 2 and Claim 14. Thus G' has an $(h_{G'}, M)$ -coloring ψ . Note that $|L(v_1)| \geq 15M/31$ and $|L(v_2)| \geq 15M/31$. We have $|L(v_1) \cap (\psi(v_3) \cup \psi(v_4))| \geq 2M/31$ and $|L(v_2) \cap (\psi(v_3) \cup \psi(v_4))| \geq 2M/31$. Let $A \subseteq L(v_1) \cap (\psi(v_3) \cup \psi(v_4))$ such that $|A| = 2M/31$. Similarly, let $B \subseteq L(v_2) \cap (\psi(v_3) \cup \psi(v_4))$ such that $|B| = 2M/31$.

Let ψ' be a limitation of ψ on $G - \{v, v_1, v_2\}$ such that $A \cup B \subseteq \psi'(v_3) \cup \psi'(v_4)$. We extend ψ' to v_1 and v_2 such that $A \subseteq \psi'(v_1)$, $B \subseteq \psi'(v_2)$, and $|\psi'(v_1) \cap \psi'(v_2)|$ as large as possible. Note that $\psi(v_1v_2) \subseteq L'(v_1) \cap L'(v_2)$ and $|\psi'(v_1)| = |\psi'(v_2)| = 9M/31$. Then $|\psi'(v_1) \cap \psi'(v_2)| \geq 7M/31$. Now, $|\cup_{i=1}^4 \psi'(v_i)| \leq 23M/31$ and we can define $\psi'(v)$ properly to obtain an (h_G, M) -coloring of G , a contradiction. ■

Claim 19 There are no two triangles in G sharing a vertex.

Proof: Suppose that G contains two triangles vv_1v_2 and vv_3v_4 sharing a vertex v . Claim 14 implies that $d_G(v_i) = 4$ for $1 \leq i \leq 4$. We next show that G has an (h_G, M) -coloring. The rough idea is that we will start with a coloring of $G - \{v, v_2, v_3\}$, where v_1 and v_4 are colored with the same colors. We extend the coloring to v_2, v_3 , and v one by one. As each v_i has degree four, if there are $9M/31$ colors such that each of them appears twice in the neighbors of v , then we can color v properly. To achieve this, when we color v_3 , we will choose $M/31$ colors from the color set of v_2 and assign them to the color set of v_3 . In the process of coloring, we may need to redefine the color sets of v_1 and v_4 .

For each $1 \leq i \leq 4$, let v'_i and v''_i be the other two neighbors of v_i . By Claim 17, vertices v'_1, v''_1, v'_2 , and v''_2 are distinct. Similarly, v'_3, v''_3, v'_4 , and v''_4 are distinct. For each $1 \leq i \leq 4$, we have $d_G(v'_i) + d_G(v''_i) \geq 7$ by Claim 18. There are two cases.

Case 1: There is some $j \in \{1, 2\}$ and some $k \in \{3, 4\}$ such that $N_G(v_j) \cap N_G(v_k) = v$. By the symmetry, we can assume that $j = 1$ and $k = 3$. Let G' be the graph obtained from $G - \{v, v_2, v_3\}$ by identifying v_1 and v_4 . Claim 2 and Claim 17 indicate that G' is C_8^2 -free and K_4 -free. Thus G' has an $(h_{G'}, M)$ -coloring ψ . Let ψ' be a limitation of ψ on $G - \{v, v_2, v_3, v'_2, v''_2, v'_3, v''_3\}$. Without loss of generality, we assume that $\psi'(v_1) = \psi'(v_4) = [1, 8M/31]$. There are two subcases.

Subcase 1.1: $|N_G(v_2) \cap N_G(v_3)| = 1$. It implies that v is the only common neighbor of v_2 and v_3 . Notice that v'_2, v''_2, v'_3 , and v''_3 are distinct as well as $d_{G'}(u) = d_G(u) - 1$ for each $u \in \{v'_2, v''_2, v'_3, v''_3\}$. Let $A_2 = [1, 8M/31] \cap (\psi(v'_2) \cup \psi(v''_2))$ and $B_2 = \psi(v'_2) \cap \psi(v''_2)$. Similarly, let $A_3 = [1, 8M/31] \cap (\psi(v'_3) \cup \psi(v''_3))$ and $B_3 = \psi(v'_3) \cap \psi(v''_3)$. We next color v_2, v_3 , and v one by one.

Let us start with the case where either $|A_2| \geq (9 - d_G(v'_2) - d_G(v''_2))M/31$ or $|B_2| \geq (9 - d_G(v'_2) - d_G(v''_2))M/31$. In the former case, we delete $M/31$ colors from $\psi(v'_2)$ and $M/31$ colors from $\psi(v''_2)$ respectively, while keeping A_2 as large as possible. In the latter case, we delete colors from $\psi(v'_2)$ and $\psi(v''_2)$ while keeping B_2 as large as possible. In each case, we can choose colors for $\psi'(v_2)$ greedily. We next color v_3 . If either $|A_3| \geq (9 - d_G(v'_3) - d_G(v''_3))M/31$ or $|B_3| \geq (9 - d_G(v'_3) - d_G(v''_3))M/31$, then we pick a set $C \subset \psi'(v_2)$ such that $|C| = M/31$. Notice that $C \cap [1, 8M/31] = \emptyset$. In the latter case, we further require $|C \cap B_3|$ as small as possible. Let us put $C \subset \psi'(v_3)$. When we define $\psi'(v'_3)$ and $\psi'(v''_3)$, we remove colors from C first while keeping A_3 (and B_3 in the latter case) as large as possible. In each case, we can choose extra colors for $\psi'(v_3)$ greedily. Note that each color from $[1, 8M/31] \cup C$ appears twice in the neighbors of v , we can color v properly to get an (h_G, M) -coloring of G , a contradiction. Therefore, we assume that $|A_3| < (9 - d_G(v'_3) - d_G(v''_3))M/31$ and $|B_3| < (9 - d_G(v'_3) - d_G(v''_3))M/31$. Notice that $|\psi(v'_3) \cup \psi(v''_3)| = (26 - d_G(v'_3) - d_G(v''_3))M/31 - |B_3|$. Recall $d_G(v'_4) + d_G(v''_4) \geq 7$. Therefore, $|\psi'(v'_4) \cup \psi'(v''_4)| \leq 17M/31$, which yields that there exists a set $D \subset (\psi(v'_3) \cup \psi(v''_3)) \setminus (\psi'(v'_4) \cup \psi'(v''_4))$ with $|D| = (9 - d_G(v'_3) - d_G(v''_3))M/31 - |B_3|$. The fact $d_G(v'_3) + d_G(v''_3) \geq 7$ implies that $|B_3| + |D| \leq 2M/31$. Note that $\psi'(v_2) \cap [1, 8M/31] = \emptyset$. Now we are able to choose $C \subset \psi'(v_2)$ such that $|C| = M/31$ and $C \cap B_3 = C \cap D = \emptyset$. Let $E \subset [1, 8M/31]$ such that $|E| = |D|$ and $E \cap A_3 = E \cap C = \emptyset$. We redefine $\psi'(v_4)$ by removing colors of E and adding colors from D . To color v_3 , we put $C \cup E \subset \psi'(v_3)$. We next define $\psi'(v'_3)$ and $\psi'(v''_3)$ by removing colors from C first and extra colors if it is needed, here we keep colors from B_3 . As sets B_3 and D , we can color v_3 properly. Since each color from $[1, 8M/31] \cup C$ appears twice in the neighbors of v , we can color v properly to get an (h_G, M) -coloring of G , a contradiction.

Now, we consider the case where $|A_2| < (9 - d_G(v'_2) - d_G(v''_2))M/31$ and $|B_2| < (9 - d_G(v'_2) - d_G(v''_2))M/31$. In this case, to color v_2 , we can use the idea to color v_3 when $|A_3| < (9 - d_G(v'_3) - d_G(v''_3))M/31$ and $|B_3| < (9 - d_G(v'_3) - d_G(v''_3))M/31$. Namely, we move a set of at most $2M/31$ colors from $[1, 8M/31]$ to $\psi'(v_2)$ and replace them by a set of at most $2M/31$ from $(\psi'(v'_2) \cup \psi'(v''_2)) \setminus (\psi'(v'_1) \cup \psi'(v''_1))$. We repeat the argument above to color v_3 by case analysis according to the sizes of A_3 and B_3 . No matter whether $|A_3| < (9 - d_G(v'_3) - d_G(v''_3))M/31$ or $|B_3| < (9 - d_G(v'_3) - d_G(v''_3))M/31$, the existence of a desired subset $C \subset \psi'(v_2) \setminus [1, 8M/31]$ is ensured by facts $|D| + |B_3| \leq 2M/31$ and $|\psi'(v_2) \setminus [1, 8M/31]| \geq 6M/31$. As above, each color from $[1, 8M/31] \cup C$ appears twice in the neighbors of v . Thus we can color v greedily and obtain an (h_G, M) -coloring of G , a contradiction.

Subcase 1.2: $|N_G(v_2) \cap N_G(v_3)| \geq 2$. We define A_2, B_2, A_3 , and B_3 as we did in Subcase 1.1. We can reuse ideas in Subcase 1.1 to extend the coloring ψ' to G . For the case where $v'_2 = v'_3$ and $v''_2 \neq v''_3$, note that $|\psi(v'_2)| = (14 - d_G(v'_2))M/31$ as both v_2 and v_3 are adjacent to v'_2 . After we finish the coloring of v'_2 , if we still use $\psi(v'_2)$ to denote the color set of v'_2 , then we require $|\psi(v'_2)| = (13 - d_G(v'_2))M/31$. Otherwise, we have troubles to extend the coloring to v_3 . Therefore, to color v_2 , we need to consider cases depending on whether $|A_2| \geq (10 - d_G(v'_2) - d_G(v''_2))M/31$ or $|B_2| \geq (10 - d_G(v'_2) - d_G(v''_2))M/31$. Note that when $|A_2| < (10 - d_G(v'_2) - d_G(v''_2))M/31$ and $|B_2| < (10 - d_G(v'_2) - d_G(v''_2))M/31$, it holds $|(\psi(v'_2) \cup \psi(v''_2)) \setminus (\psi(v'_1) \cup \psi(v''_1))| \geq (10 - d_G(v'_2) - d_G(v''_2))M/31 - |B_2|$ as $|\psi(v'_2)| = (14 - d_G(v'_2))M/31$ and $|\psi(v''_2)| = (13 - d_G(v''_2))M/31$. For the case where $v'_2 = v'_3$ and $v''_2 = v''_3$, to define $\psi'(v_2)$, we have to consider cases depending on whether $|A_2| \geq (11 - d_G(v'_2) - d_G(v''_2))M/31$ or $|B_2| \geq (11 - d_G(v'_2) - d_G(v''_2))M/31$ similarly. In each case, we can construct an (h_G, M) -

coloring of G , a contradiction.

Case 2: For each $j \in \{1, 2\}$ and each $k \in \{3, 4\}$, we have $|N_G(v_j) \cap N_G(v_k)| \geq 2$. Recall Claim 19. In this case, there are distinct vertices x, y, w, z such that $x \in N_G(v_1) \cap N_G(v_3)$, $y \in N_G(v_1) \cap N_G(v_4)$, $w \in N_G(v_2) \cap N_G(v_3)$, and $z \in N_G(v_2) \cap N_G(v_4)$. By the symmetry, we assume that $d_G(x) + d_G(y) \geq d_G(w) + d_G(z)$. Let G' be the graph obtained from $G - \{v, v_2, v_3\}$ by identifying v_1 and v_4 . It is easy to check that G' is K_4 -free and C_8^2 -free. Let ψ be an $(h_{G'}, M)$ -coloring of G' and ψ' be a limitation of ψ on $G - \{v, v_2, v_3, x, w, z\}$. We also assume that $\psi'(v_1) = \psi'(v_4) = [1, 8M/31]$ and define $A_2 = [1, 8M/31] \cap (\psi(w) \cup \psi(z))$ as well as $B_2 = \psi(w) \cap \psi(z)$. To color v_2 , we reuse ideas in Subcase 1.1 and consider cases depending on whether $|A_2| \geq (10 - d_G(w) - d_G(z))M/31$ or $|B_2| \geq (10 - d_G(w) - d_G(z))M/31$. After the coloring of $\psi'(v_2)$, we still use $\psi(w)$ to denote the color set of w , here we similarly require $|\psi(w)| = (13 - d_G(w))M/31$. To define $\psi'(v_2')$, when $|A_2| < (10 - d_G(w) - d_G(z))M/31$ and $|B_2| < (10 - d_G(w) - d_G(z))M/31$, it holds $|(\psi(w) \cup \psi(z)) \setminus (\psi(x) \cup \psi(y))| \geq (10 - d_G(w) - d_G(z))M/31 - |B_2|$ as the assumption $d_G(x) + d_G(y) \geq d_G(w) + d_G(z)$. To color v_3 , we define $A_3 = [1, 8M/31] \cap (\psi(x) \cup \psi(w))$ and $B_3 = \psi(x) \cap \psi(w)$ followed by considering cases depending on $|A_3|$ and $|B_3|$. In each case, we can guarantee that each color of $[1, 8M/31] \cup C$ appears twice in the neighbors of v for some $C \subset \psi'(v_2) \setminus [1, 8M/31]$ with $|C| = M/31$. Thus we can color v greedily to get an (h_G, M) -coloring of G , a contradiction. ■

Claim 20 *Each 4-vertex has at least one 4-neighbor.*

Proof: Let v be a 4-vertex of G with $N_G(v) = \{v_1, v_2, v_3, v_4\}$. Suppose that $d_G(v_i) = 3$ for $1 \leq i \leq 4$. Then $\{v_1, v_2, v_3, v_4\}$ is an independent set and the neighbors of v_1, v_2, v_3, v_4 are 4-vertices by Claim 11 and Claim 14. Let G' be the graph obtained from $G - v$ by identifying v_1 and v_2 . Claim 14 and Claim 2 imply that G' is K_4 -free and contains no C_8^2 . Then G' has an $(h_{G'}, M)$ -coloring ψ . Note that $|\psi(v_1 v_2)| = 8M/31$. Let ψ' be a limitation of ψ on $G - \{v, v_1, v_2, v_3, v_4\}$. To define $\psi'(v_1)$ and $\psi'(v_2)$, we add $M/31$ extra colors to $\psi(v_1 v_2)$ greedily. We can assume that $|L'(v)| = 21M/31$, $|L'(v_3)| = 15M/31$ and $|L'(v_4)| = 15M/31$ by deleting colors if it is necessary. Let $A \subseteq L'(v) \setminus L'(v_3)$ with $|A| = 2M/31$ and $B \subseteq L'(v) \setminus L'(v_4)$ with $|B| = 2M/31$. To color v , we put $A \cup B$ in $\psi'(v)$ and add extra colors to $\psi'(v)$ such that $|\psi'(v)| = 8M/31$. Now v_3 and v_4 can be colored properly because of sets A and B . We obtained an (h_G, M) -coloring of G , a contradiction. ■

Proof of Lemma 3: Claim 2 implies that $d_G(v) \geq 3$ for each $v \in V(G)$. If $d_G(v) = 3$, then $N_G(v)$ is an independent set by Claim 14. Consequently, $e(N_G(v), N_G^2(v)) \geq 6$. If $d_G(v) = 4$, then v has at least a 4-neighbor by Claim 20. If v has exactly one 4-neighbor, then v is not in any triangles by Claim 14, which yields $e(N_G(v), N_G^2(v)) \geq 9$. If v has exactly two 4-neighbors, then v is not in any triangles by Claims 14 and 18. It follows that $e(N_G(v), N_G^2(v)) \geq 2 + 2 + 3 + 3 = 10$. If v has at least three 4-neighbors, then v is in at most one triangle by Claim 17 and Claim 19 and we have $e(N_G(v), N_G^2(v)) \geq 9$. The lemma is proved. ■

References

- [1] O. Borodin and A. Kostochka, On an upper bound on a graph's chromatic number, depending on the graph's degree and density, *J. Combin. Theory Ser. B*, **23** (1977), 247–250.
- [2] Z. Dvořák, J. S. Sereni, and J. Volec, Subcubic triangle-free graphs have fractional chromatic number at most $14/5$, *J. Lond. Math. Soc. (2)*, **89**(3) (2014), 641–662.
- [3] K. Edwards and A. D. King, Bounding the fractional chromatic number of K_Δ -free graphs, *SIAM J. Discrete Math.*, **27**(2) (2013), 1184–1208.

- [4] S. Fajtlowicz, On the size of independent sets in graphs, in *Proceedings of the Ninth Southeastern Conference on Combinatorics, Graph Theory and Computing* (Florida Atlantic Univ, Boca Raton, FL, 1978), Congress, Numer, XXI, Utilitas Math, Winnipeg, MB, 1978, 269–274.
- [5] D. Ferguson, T. Kaiser, and D. Král', The fractional chromatic number of triangle-free subcubic graphs, *European J. Combin.*, **35** (2014), 184–220.
- [6] T. Gallai, Kritische graphen I, *Magyar Tud. Akad. Mat. Kutat ó Int. Közl*, 8 (1963), 165–192.
- [7] J. Griggs and O. Murphy, Edge density and independence ratio in triangle-free graphs with maximum degree three, *Discrete Math.*, **152** (1996), 157–170.
- [8] H. Hatami and X. Zhu, The fractional chromatic number of graphs of maximum degree at most three, *SIAM J. Discrete Math.*, **24**(4) (2009/2010), 1762–1775.
- [9] C. C. Heckman and R. Thomas, A new proof of the independence ratio of triangle-free cubic graphs, *Discrete Math.*, **233** (2001), 233–237.
- [10] A. Johansson, Asymptotic choice number for triangle free graphs, *DIMACS Technical Report*, **91-5**.
- [11] F. Kardoš, D. Král', and J. Volec, Fractional colorings of cubic graphs with large girth, *SIAM J. Discrete Math.*, **25**(3) (2011), 1454–1476.
- [12] A. D. King, L. Lu, and X. Peng, A fractional analogue of Brooks' Theorem, *SIAM J. Discrete Math.*, **26-2** (2012), 452–471.
- [13] C. H. Liu, An upper bound on the fractional chromatic number of triangle-free subcubic graphs. *SIAM J. Discrete Math.*, **28**(3) (2014), 1102–1136.
- [14] L. Lu and X. Peng, The fractional chromatic number of triangle-free graphs with $\Delta \leq 3$, *Discrete Math.*, **312**(24) (2012), 3502–3516.
- [15] M. Molloy and B. Reed, *Graph colouring and the probabilistic method*, volume 23 of Algorithms and Combinatorics. Springer-Verlag, Berlin, 2002.
- [16] B. Reed, ω , Δ , and χ , *J. of Graph Theory*, vol **27-4** (1998), 177–227.
- [17] B. Reed, A strengthening of Brooks' theorem, *J. Combin. Theory Ser. B*, **76** (1999), 136–149.
- [18] E. R. Scheinerman and D. H. Ullman, *Fractional Graph Theory. A Rational Approach to the Theory of Graphs*, Wiley-Intersci. Ser. Discrete Math. Optim, John Wiley & Sons, Inc, New York, 1997.
- [19] W. Staton, Some Ramsey-type numbers and the independence ratio, *Trans. Amer. Math. Soc.*, **256** (1979), 353–370.