UNCERTAINTY PRINCIPLES FOR THE WINDOWED OPDAM-CHEREDNIK TRANSFORM

SHYAM SWARUP MONDAL AND ANIRUDHA PORIA

Shyam Swarup Mondal, Department of Mathematics, Indian Institute of Technology Delhi, New Delhi 110016, India

Anirudha Poria, Department of Mathematics, Bar-Ilan University, Ramat-Gan 5290002, Israel

Department of Mathematics, SRM Institute of Science and Technology, Kattankulathur 603203, Tamil Nadu, India

Corresponding Author: Anirudha Poria, Department of Mathematics, Bar-Ilan University, Ramat-Gan 5290002, Israel anirudhamath@gmail.com, anirudhp@srmist.edu.in

ABSTRACT. In this paper, we study a few versions of the uncertainty principle for the windowed Opdam–Cherednik transform. In particular, we establish the uncertainty principle for orthonormal sequences, Donoho–Stark's uncertainty principle, Benedicks-type uncertainty principle, Heisenberg-type uncertainty principle and local uncertainty inequality for this transform. We also obtain the Heisenberg-type uncertainty inequality using the k-entropy of the windowed Opdam–Cherednik transform.

1. INTRODUCTION

Uncertainty principles play a fundamental role in the field of mathematics, physics, in addition to some engineering areas such as signal processing, image processing, quantum theory, optics, and many other well known areas [6, 11, 13, 18, 41]. In quantum physics, it says that a particle's speed and position cannot both be measured with infinite precision. The classical Heisenberg uncertainty principle was established in the Schrödinger space (the square-integrable function space). It states that a non-zero function and its Fourier transform cannot be both sharply localized, i.e., it is impossible for a non-zero function and its Fourier transform to be simultaneously small. This phenomenon has been under intensive study for almost a century now and extensively investigated in different settings. One can formulate different forms of the uncertainty principle depending on various ways of measuring the localization of a function. Uncertainty principles can be divided into two categories: quantitative and qualitative uncertainty principles. Quantitative uncertainty principles are some special types of inequalities that tells us how a function and its Fourier transform relate. For example, Benedicks [5], Donoho and Stark [13], and Slepian and Pollak [43] gave quantitative uncertainty principles for the Fourier transforms. On the other hand, qualitative uncertainty principles imply the vanishing of a function under some strong conditions on the function. In particular, Hardy [21], Morgan [36], Cowling and Price [9], and Beurling [25] theorems are the examples of qualitative uncertainty principles. For a more detailed study on the history of the uncertainty principle, and for many other generalizations and variations of the uncertainty principle, we refer to the book of Havin and Jöricke [22],

Date: August 22, 2023.

²⁰¹⁰ Mathematics Subject Classification. Primary 44A15; Secondary 42C20, 33C45.

Key words and phrases. Opdam-Cherednik transform; windowed Opdam-Cherednik transform; Heisenbergtype uncertainty inequality; Donoho-Stark's uncertainty principle; Benedicks-type uncertainty principle.

Research supported by ERC Starting Grant No. 713927.

and the excellent survey of Folland and Sitaram [15].

Considerable attention has been devoted to discovering generalizations to new contexts for quantitative uncertainty principles. For example, quantitative uncertainty principles were studied in [31] for the generalized Fourier transform. Over the years, discovering new mathematical formulations of the uncertainty principle for the windowed Fourier transform have drawn significant attention among many researchers, see for example [7, 22, 44] and the references therein. Uncertainty principles are further extended and studied in different settings by many authors. For instance, one can look [32] for the Dunkl Gabor transform, [8] for two sided quaternion windowed Fourier transform, [4, 19] for the continuous Hankel transform, [16, 17] for the windowed Hankel transform, [20] for the Hankel–Gabor transform, and [28] for the Heckman–Opdam transform.

As a generalization of Euclidean uncertainty principles for the Fourier transform, Daher et al. [10] have obtained some qualitative uncertainty principles for the Cherednik transform. Also, qualitative uncertainty principles for the Opdam–Cherednik transform have been studied by Mejjaoli in [29] using classical uncertainty principles for the Fourier transform and composition properties of the Opdam–Cherednik transform. These results have been further extended to modulation spaces by the second author in [39]. Moreover, the Benedicks-type uncertainty principle for the Opdam–Cherednik transform has been investigated by Achak and Daher in [1]. Recently, the second author introduced the windowed Opdam–Cherednik transform and discussed the time-frequency analysis of localization operators associated with the windowed Opdam–Cherednik transform in [40]. However, up to our knowledge, quantitative uncertainty principles have not been studied for the windowed Opdam–Cherednik transform. In this paper, we extend some quantitative uncertainty principles for this transform.

The motivation to prove these types of quantitative uncertainty principles for the windowed Opdam–Cherednik transform in the framework of the Opdam–Cherednik transform arises from the classical uncertainty principles for the windowed Fourier transform and the remarkable contribution of the Opdam–Cherednik transform in harmonic analysis (see [2,37,38,42]). Another important motivation to study the Jacobi–Cherednik operators arises from their relevance in the algebraic description of exactly solvable quantum many-body systems of Calogero–Moser–Sutherland type (see [12,24]) and they provide a useful tool in the study of special functions with root systems (see [14,23]). These describe algebraically integrable systems in one dimension and have gained considerable interest in mathematical physics. Other motivation for the investigation of the Jacobi–Cherednik operator and the windowed Opdam–Cherednik transform is to generalize the previous subjects which are bound with the physics. For a more detailed discussion, one can see [33].

Uncertainty inequalities are some special class of uncertainty principles that give us information about how a function and its Fourier transform relate and have got considerable importance in signal analysis, physics, optics, and many other well-known areas [6,13,18,26,41]. A well-known example of uncertainty inequality is the Heisenberg inequality. Considerable attention has been devoted to discovering generalizations to new contexts for uncertainty inequalities for various generalized transforms by several researchers. For instance, these uncertainty inequalities were obtained in [28] for the Heckman–Opdam transform, and in [35] for the Opdam–Cherednik transform. Some recent work on the windowed Opdam–Cherednik transform [34,40] motivates us to study a few quantitative uncertainty principles for this transform.

The main aim of this paper is to study a few uncertainty principles related to the windowed Opdam–Cherednik transform. More preciously, we prove the uncertainty principle for orthonormal sequences, Donoho–Stark's uncertainty principle, Benedicks-type uncertainty principle, Heisenberg-type uncertainty principle and local uncertainty inequality for the windowed Opdam–Cherednik transform. We study the version of Donoho–Stark's uncertainty principle and show that the windowed Opdam–Cherednik transform cannot be concentrated in any small set. Also, we obtain an estimate for the size of the essential support of this transform under the condition that the windowed Opdam–Cherednik transform of a non-zero function is time-frequency concentrated on a measurable set. Then, we investigate the Benedicks-type uncertainty principle and show that the windowed Opdam–Cherednik transform cannot be concentrated inside a set of measures arbitrarily small. Further, we study the Heisenberg-type uncertainty inequality for a general magnitude and provide the result related to the $L^2(\mathbb{R}, A_{\alpha,\beta})$ -mass of the windowed Opdam–Cherednik transform outside sets of finite measure. Finally, we obtain the Heisenbergtype uncertainty inequality using the k-entropy and study the localization of the k-entropy of the windowed Opdam–Cherednik transform.

The presentation of this manuscript is divided into four sections apart from the introduction. In Section 2, we present some preliminaries related to the Opdam–Cherednik transform. In Section 3, we recall some essential properties related to the windowed Opdam–Cherednik transform. Some different types of uncertainty principles associated with the windowed Opdam–Cherednik transform are provided in Section 4. In particular, we prove the uncertainty principle for orthonormal sequences, Donoho–Stark's uncertainty principle, Benedicks-type uncertainty principle, Heisenberg-type uncertainty principle and local uncertainty inequality for the windowed Opdam–Cherednik transform. We conclude the paper with the Heisenberg-type uncertainty inequality using the k-entropy of this transform.

2. HARMONIC ANALYSIS AND THE OPDAM-CHEREDNIK TRANSFORM

In this section, we recall some necessary definitions and results from the harmonic analysis related to the Opdam–Cherednik transform, which will be used frequently. A complete account of harmonic analysis related to this transform can be found in [2,27,30,37–40,42]. However, we will use the notations given in [39].

Let $T_{\alpha,\beta}$ denote the Jacobi–Cherednik differential–difference operator (also called the Dunkl–Cherednik operator)

$$T_{\alpha,\beta}f(x) = \frac{d}{dx}f(x) + \left[(2\alpha + 1)\coth x + (2\beta + 1)\tanh x \right] \frac{f(x) - f(-x)}{2} - \rho f(-x)$$

where α, β are two parameters satisfying $\alpha \geq \beta \geq -\frac{1}{2}$ and $\alpha > -\frac{1}{2}$, and $\rho = \alpha + \beta + 1$. Let $\lambda \in \mathbb{C}$. The Opdam hypergeometric functions $G_{\lambda}^{\alpha,\beta}$ on \mathbb{R} are eigenfunctions $T_{\alpha,\beta}G_{\lambda}^{\alpha,\beta}(x) = i\lambda G_{\lambda}^{\alpha,\beta}(x)$ of $T_{\alpha,\beta}$ that are normalized such that $G_{\lambda}^{\alpha,\beta}(0) = 1$. The eigenfunction $G_{\lambda}^{\alpha,\beta}$ is given by

$$G_{\lambda}^{\alpha,\beta}(x) = \varphi_{\lambda}^{\alpha,\beta}(x) - \frac{1}{\rho - i\lambda} \frac{d}{dx} \varphi_{\lambda}^{\alpha,\beta}(x) = \varphi_{\lambda}^{\alpha,\beta}(x) + \frac{\rho + i\lambda}{4(\alpha + 1)} \sinh 2x \; \varphi_{\lambda}^{\alpha + 1,\beta + 1}(x),$$

where $\varphi_{\lambda}^{\alpha,\beta}(x) = {}_{2}F_{1}\left(\frac{\rho+i\lambda}{2}, \frac{\rho-i\lambda}{2}; \alpha+1; -\sinh^{2}x\right)$ is the classical Jacobi function.

For every $\lambda \in \mathbb{C}$ and $x \in \mathbb{R}$, the eigenfunction $G_{\lambda}^{\alpha,\beta}$ satisfy

$$|G_{\lambda}^{\alpha,\beta}(x)| \le C \ e^{-\rho|x|} e^{|\mathrm{Im}(\lambda)||x|}$$

where C is a positive constant. Since $\rho > 0$, we have

$$|G_{\lambda}^{\alpha,\beta}(x)| \le C \ e^{|\operatorname{Im}(\lambda)||x|}.$$
(2.1)

Let us denote by $C_c(\mathbb{R})$ the space of continuous functions on \mathbb{R} with compact support. The Opdam–Cherednik transform is the Fourier transform in the trigonometric Dunkl setting, and it is defined as follows.

Definition 2.1. Let $\alpha \geq \beta \geq -\frac{1}{2}$ with $\alpha > -\frac{1}{2}$. The Opdam-Cherednik transform $\mathcal{H}_{\alpha,\beta}(f)$ of a function $f \in C_c(\mathbb{R})$ is defined by

$$\mathcal{H}_{\alpha,\beta}(f)(\lambda) = \int_{\mathbb{R}} f(x) \ G_{\lambda}^{\alpha,\beta}(-x) \ A_{\alpha,\beta}(x) dx \quad \text{for all } \lambda \in \mathbb{C},$$

where $A_{\alpha,\beta}(x) = (\sinh |x|)^{2\alpha+1} (\cosh |x|)^{2\beta+1}$. The inverse Opdam-Cherednik transform for a suitable function g on \mathbb{R} is given by

$$\mathcal{H}_{\alpha,\beta}^{-1}(g)(x) = \int_{\mathbb{R}} g(\lambda) \ G_{\lambda}^{\alpha,\beta}(x) \ d\sigma_{\alpha,\beta}(\lambda) \quad \text{for all } x \in \mathbb{R},$$

where

$$d\sigma_{\alpha,\beta}(\lambda) = \left(1 - \frac{\rho}{i\lambda}\right) \frac{d\lambda}{8\pi |C_{\alpha,\beta}(\lambda)|^2}$$

and

$$C_{\alpha,\beta}(\lambda) = \frac{2^{\rho-i\lambda}\Gamma(\alpha+1)\Gamma(i\lambda)}{\Gamma\left(\frac{\rho+i\lambda}{2}\right)\Gamma\left(\frac{\alpha-\beta+1+i\lambda}{2}\right)}, \quad \lambda \in \mathbb{C} \setminus i\mathbb{N}.$$

The Plancherel formula is given by

$$\int_{\mathbb{R}} |f(x)|^2 A_{\alpha,\beta}(x) dx = \int_{\mathbb{R}} \mathcal{H}_{\alpha,\beta}(f)(\lambda) \overline{\mathcal{H}_{\alpha,\beta}(\check{f})(-\lambda)} \, d\sigma_{\alpha,\beta}(\lambda), \tag{2.2}$$

where $\check{f}(x) := f(-x)$.

Let $L^p(\mathbb{R}, A_{\alpha,\beta})$ (resp. $L^p(\mathbb{R}, \sigma_{\alpha,\beta})$), $p \in [1, \infty]$, denote the L^p -spaces corresponding to the measure $A_{\alpha,\beta}(x)dx$ (resp. $d|\sigma_{\alpha,\beta}|(x)$).

The generalized translation operator associated with the Opdam–Cherednik transform is defined by [3]

$$\tau_x^{(\alpha,\beta)}f(y) = \int_{\mathbb{R}} f(z) \, d\mu_{x,y}^{(\alpha,\beta)}(z), \qquad (2.3)$$

where $d\mu_{x,y}^{(\alpha,\beta)}$ is given by

$$d\mu_{x,y}^{(\alpha,\beta)}(z) = \begin{cases} \mathcal{K}_{\alpha,\beta}(x,y,z) \ A_{\alpha,\beta}(z) \ dz & \text{if } xy \neq 0\\ d\delta_x(z) & \text{if } y = 0\\ d\delta_y(z) & \text{if } x = 0 \end{cases}$$
(2.4)

and

$$\mathcal{K}_{\alpha,\beta}(x,y,z) = M_{\alpha,\beta} |\sinh x \cdot \sinh y \cdot \sinh z|^{-2\alpha} \int_0^\pi g(x,y,z,\chi)_+^{\alpha-\beta-1} \\ \times \left[1 - \sigma_{x,y,z}^{\chi} + \sigma_{x,z,y}^{\chi} + \sigma_{z,y,x}^{\chi} + \frac{\rho}{\beta + \frac{1}{2}} \coth x \cdot \coth y \cdot \coth z (\sin \chi)^2 \right] \times (\sin \chi)^{2\beta} d\chi$$

if $x, y, z \in \mathbb{R} \setminus \{0\}$ satisfy the triangular inequality ||x| - |y|| < |z| < |x| + |y|, and $\mathcal{K}_{\alpha,\beta}(x, y, z) = 0$ otherwise. Here

$$\sigma_{x,y,z}^{\chi} = \begin{cases} \frac{\cosh x \cdot \cosh y - \cosh z \cdot \cos \chi}{\sinh x \cdot \sinh y} & \text{if } xy \neq 0\\ 0 & \text{if } xy = 0 \end{cases} \quad \text{for } x, y, z \in \mathbb{R}, \ \chi \in [0,\pi], \end{cases}$$

 $g(x, y, z, \chi) = 1 - \cosh^2 x - \cosh^2 y - \cosh^2 z + 2 \cosh x \cdot \cosh y \cdot \cosh z \cdot \cos \chi, \text{ and } x + 2 \cosh x \cdot \cosh y \cdot \cosh z \cdot \cos \chi,$

$$g_{+} = \begin{cases} g & \text{if } g > 0\\ 0 & \text{if } g \le 0. \end{cases}$$

The kernel $\mathcal{K}_{\alpha,\beta}(x,y,z)$ satisfies the following symmetry properties:

 $\mathcal{K}_{\alpha,\beta}(x,y,z) = \mathcal{K}_{\alpha,\beta}(y,x,z), \ \mathcal{K}_{\alpha,\beta}(x,y,z) = \mathcal{K}_{\alpha,\beta}(-z,y,-x), \ \mathcal{K}_{\alpha,\beta}(x,y,z) = \mathcal{K}_{\alpha,\beta}(x,-z,-y).$ For every $x, y \in \mathbb{R}$, we have

$$\tau_x^{(\alpha,\beta)}f(y) = \tau_y^{(\alpha,\beta)}f(x), \tag{2.5}$$

and

$$\mathcal{H}_{\alpha,\beta}(\tau_x^{(\alpha,\beta)}f)(\lambda) = G_{\lambda}^{\alpha,\beta}(x) \mathcal{H}_{\alpha,\beta}(f)(\lambda), \qquad (2.6)$$

for $f \in C_c(\mathbb{R})$.

If $f \in L^1(\mathbb{R}, A_{\alpha,\beta})$, then

$$\int_{\mathbb{R}} \tau_x^{(\alpha,\beta)} f(y) \ A_{\alpha,\beta}(y) \ dy = \int_{\mathbb{R}} f(y) \ A_{\alpha,\beta}(y) \ dy.$$
(2.7)

For every $f \in L^p(\mathbb{R}, A_{\alpha,\beta})$ and every $x \in \mathbb{R}$, the function $\tau_x^{(\alpha,\beta)} f$ belongs to the space $L^p(\mathbb{R}, A_{\alpha,\beta})$ and

$$\left\|\tau_x^{(\alpha,\beta)}f\right\|_{L^p(\mathbb{R},A_{\alpha,\beta})} \le C_{\alpha,\beta} \|f\|_{L^p(\mathbb{R},A_{\alpha,\beta})},\tag{2.8}$$

where $C_{\alpha,\beta}$ is a positive constant.

The convolution product associated with the Opdam–Cherednik transform is defined for two suitable functions f and g by [3]

$$(f *_{\alpha,\beta} g)(x) = \int_{\mathbb{R}} \tau_x^{(\alpha,\beta)} f(-y) g(y) A_{\alpha,\beta}(y) dy$$

and

$$\mathcal{H}_{\alpha,\beta}(f*_{\alpha,\beta}g) = \mathcal{H}_{\alpha,\beta}(f) \ \mathcal{H}_{\alpha,\beta}(g).$$
(2.9)

3. The windowed Opdam-Cherednik transform

In this section, we collect the necessary definitions and results from the harmonic analysis related to the windowed Opdam–Cherednik transform. For a detailed discussion on this transform, we refer to [40].

Let $g \in L^2(\mathbb{R}, A_{\alpha,\beta})$ and $\xi \in \mathbb{R}$, the modulation operator of g associated with the Opdam– Cherednik transform is defined by

$$\mathcal{M}_{\xi}^{(\alpha,\beta)}g = \mathcal{H}_{\alpha,\beta}^{-1}\left(\sqrt{\tau_{\xi}^{(\alpha,\beta)}|\mathcal{H}_{\alpha,\beta}(g)|^2}\right).$$
(3.1)

Then, for every $g \in L^2(\mathbb{R}, A_{\alpha,\beta})$ and $\xi \in \mathbb{R}$, by using the Plancherel formula (2.2) and the translation invariance of the Plancherel measure $d\sigma_{\alpha,\beta}$, we obtain

$$\left\|\mathcal{M}_{\xi}^{(\alpha,\beta)}g\right\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})} = \|g\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})}.$$
(3.2)

Now, for a non-zero window function $g \in L^2(\mathbb{R}, A_{\alpha,\beta})$ and $(x,\xi) \in \mathbb{R}^2$, we consider the function $g_{x,\xi}^{(\alpha,\beta)}$ defined by

$$g_{x,\xi}^{(\alpha,\beta)} = \tau_x^{(\alpha,\beta)} \mathcal{M}_{\xi}^{(\alpha,\beta)} g.$$
(3.3)

For any function $f \in L^2(\mathbb{R}, A_{\alpha,\beta})$, we define the windowed Opdam–Cherednik transform by

$$\mathcal{W}_{g}^{(\alpha,\beta)}(f)(x,\xi) = \int_{\mathbb{R}} f(s) \,\overline{g_{x,\xi}^{(\alpha,\beta)}(-s)} \,A_{\alpha,\beta}(s) \,ds, \quad (x,\xi) \in \mathbb{R}^{2},\tag{3.4}$$

which can be also written in the form

$$\mathcal{W}_{g}^{(\alpha,\beta)}(f)(x,\xi) = \left(f *_{\alpha,\beta} \overline{\mathcal{M}_{\xi}^{(\alpha,\beta)}g}\right)(x).$$
(3.5)

We define the measure $A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta}$ on \mathbb{R}^2 by

$$d(A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta})(x,\xi) = A_{\alpha,\beta}(x)dx \ d\sigma_{\alpha,\beta}(\xi).$$
(3.6)

The windowed Opdam–Cherednik transform satisfies the following properties.

Proposition 3.1. [40] Let $g \in L^2(\mathbb{R}, A_{\alpha,\beta})$ be a non-zero window function. Then we have (1) (Plancherel's formula) For every $f \in L^2(\mathbb{R}, A_{\alpha,\beta})$,

$$\left\| \mathcal{W}_{g}^{(\alpha,\beta)}(f) \right\|_{L^{2}(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta})} = \|f\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})} \|g\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})}.$$
(3.7)

(2) (Orthogonality relation) For every $f, h \in L^2(\mathbb{R}, A_{\alpha,\beta})$, we have

$$\iint_{\mathbb{R}^2} \mathcal{W}_g^{(\alpha,\beta)}(f)(x,\xi) \ \overline{\mathcal{W}_g^{(\alpha,\beta)}(h)(x,\xi)} \ d(A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta})(x,\xi) = \|g\|_{L^2(\mathbb{R},A_{\alpha,\beta})}^2 \int_{\mathbb{R}} f(s)\overline{h(s)} \ A_{\alpha,\beta}(s) \ ds.$$
(3.8)

(3) (Reproducing kernel Hilbert space) The space $\mathcal{W}_{g}^{(\alpha,\beta)}(L^{2}(\mathbb{R}, A_{\alpha,\beta}))$ is a reproducing kernel Hilbert space in $L^{2}(\mathbb{R}^{2}, A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta})$ with kernel function K_{g} defined by

$$K_{g}((x',\xi');(x,\xi)) = \frac{1}{\|g\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})}^{2}} \left(g_{x,\xi}^{(\alpha,\beta)}(-\cdot) *_{\alpha,\beta} \overline{\mathcal{M}_{\xi'}^{(\alpha,\beta)}g}\right)(x')$$
$$= \frac{1}{\|g\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})}^{2}} \mathcal{W}_{g}^{(\alpha,\beta)} \left(g_{x,\xi}^{(\alpha,\beta)}(-\cdot)\right)(x',\xi').$$
(3.9)

Furthermore, the kernel is pointwise bounded

$$|K_g((x',\xi');(x,\xi))| \le 1, \quad \text{for all } (x,\xi); (x',\xi') \in \mathbb{R}^2.$$
 (3.10)

The windowed Opdam–Cherednik transform also satisfies the following boundedness properties.

Proposition 3.2. [40] Let $g \in L^2(\mathbb{R}, A_{\alpha,\beta})$ be a non-zero window function. Then for every $f \in L^2(\mathbb{R}, A_{\alpha,\beta})$, we have (1)

$$\left\| \mathcal{W}_{g}^{(\alpha,\beta)}(f) \right\|_{L^{\infty}(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta})} \leq \|f\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})} \|g\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})}.$$
(3.11)

(2) The function $\mathcal{W}_{g}^{(\alpha,\beta)}(f) \in L^{p}(\mathbb{R}^{2}, A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta}), \ p \in [2,\infty)$ and $\left\| \mathcal{W}_{g}^{(\alpha,\beta)}(f) \right\|_{L^{p}(\mathbb{R}^{2}, A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta})} \leq \|f\|_{L^{2}(\mathbb{R}, A_{\alpha,\beta})} \|g\|_{L^{2}(\mathbb{R}, A_{\alpha,\beta})}.$ (3.12)

4. Uncertainty principles for the windowed Opdam-Cherednik transform

In this section, we obtain various uncertainty principles for the windowed Opdam–Cherednik transform.

4.1. Uncertainty principle for orthonormal sequences. In this subsection, we establish the uncertainty principle for orthonormal sequences associated with the windowed Opdam–Cherednik transform. First, we consider the following orthogonal projections:

- (1) Let P_g be the orthogonal projection from $L^2(\mathbb{R}^2, A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta})$ onto $\mathcal{W}_g^{(\alpha,\beta)}(L^2(\mathbb{R}, A_{\alpha,\beta}))$ and Im P_g denotes the range of P_g .
- (2) Let P_{Σ} be the orthogonal projection on $L^2(\mathbb{R}^2, A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta})$ defined by

$$P_{\Sigma}F = \chi_{\Sigma}F, \quad F \in L^2(\mathbb{R}^2, A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta}), \tag{4.1}$$

where $\Sigma \subset \mathbb{R}^2$ and $\operatorname{Im} P_{\Sigma}$ is the range of P_{Σ} .

Also, we define

$$\|P_{\Sigma}P_{g}\| = \sup\left\{\|P_{\Sigma}P_{g}(F)\|_{L^{2}(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta})} : F \in L^{2}(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta}), \|F\|_{L^{2}(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta})} = 1\right\}.$$

We first need the following result.

Theorem 4.1. Let $g \in L^2(\mathbb{R}, A_{\alpha,\beta})$ be a non-zero window function. Then for any $\Sigma \subset \mathbb{R}^2$ of finite measure $A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta}(\Sigma) < \infty$, the operator $P_{\Sigma}P_g$ is a Hilbert–Schmidt operator. Moreover, we have the following estimation

$$\left\|P_{\Sigma}P_{g}\right\|^{2} \leq A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta}(\Sigma).$$

Proof. Since P_g is a projection onto a reproducing karnel Hilbert space, for any function $F \in L^2(\mathbb{R}^2, A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta})$, the orthogonal projection P_g can be expressed as

$$P_g(F)(x,\xi) = \iint_{\mathbb{R}^2} F(x',\xi') K_g((x',\xi');(x,\xi)) \ d(A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta})(x',\xi'),$$

where $K_g((x',\xi');(x,\xi))$ is given by (3.9). Using the relation (4.1), we obtain

$$P_{\Sigma}P_g(F)(x,\xi) = \iint_{\mathbb{R}^2} \chi_{\Sigma}(x,\xi)F(x',\xi')K_g((x',\xi');(x,\xi)) \ d(A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta})(x',\xi').$$

This shows that the operator $P_{\Sigma}P_g$ is an integral operator with kernel $K((x',\xi');(x,\xi)) = \chi_{\Sigma}(x,\xi)K_g((x',\xi');(x,\xi))$. Using the relation (3.9), Plancherel's formula (3.7), and Fubini's theorem, we have

$$\begin{split} \|P_{\Sigma}P_{g}\|_{HS}^{2} &= \iint_{\mathbb{R}^{2}} \iint_{\mathbb{R}^{2}} \left| K((x',\xi');(x,\xi)) \right|^{2} \, d(A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta})(x',\xi') \, d(A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta})(x,\xi) \\ &= \iint_{\mathbb{R}^{2}} \iint_{\mathbb{R}^{2}} \left| \chi_{\Sigma}(x,\xi) \right|^{2} \left| K_{g}((x',\xi');(x,\xi)) \right|^{2} \, d(A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta})(x',\xi') \, d(A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta})(x,\xi) \\ &= \frac{1}{\|g\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})}^{4}} \iint_{\Sigma} \left(\iint_{\mathbb{R}^{2}} \left| \mathcal{W}_{g}^{(\alpha,\beta)} \left(g_{x,\xi}^{(\alpha,\beta)}(-\cdot) \right) (x',\xi') \right|^{2} \, d(A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta})(x',\xi') \right) \\ &\times d(A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta})(x,\xi) \\ &\leq \frac{\|g\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})}^{4}}{\|g\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})}^{4}} A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta}(\Sigma) = A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta}(\Sigma). \end{split}$$

Thus, the operator $P_{\Sigma}P_g$ is a Hilbert–Schmidt operator. Now, the proof follows from the fact that $\|P_{\Sigma}P_g\| \leq \|P_{\Sigma}P_g\|_{HS}$.

In the following, we obtain the uncertainty principle for orthonormal sequences associated with the windowed Opdam–Cherednik transform.

Theorem 4.2. Let $g \in L^2(\mathbb{R}, A_{\alpha,\beta})$ be a non-zero window function and $\{\phi_n\}_{n\in\mathbb{N}}$ be an orthonormal sequence in $L^2(\mathbb{R}, A_{\alpha,\beta})$. Then for any $\Sigma \subset \mathbb{R}^2$ of finite measure $A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta}(\Sigma) < \infty$, we have

$$\sum_{n=1}^{N} \left(1 - \left\| \chi_{\Sigma^{c}} \mathcal{W}_{g}^{(\alpha,\beta)}(\phi_{n}) \right\|_{L^{2}(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta})} \right) \leq A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta}(\Sigma),$$

for every $N \in \mathbb{N}$.

Proof. Let $\{e_n\}_{n\in\mathbb{N}}$ be an orthonormal basis for $L^2(\mathbb{R}^2, A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta})$. Since $P_{\Sigma}P_g$ is a Hilbert–Schmidt operator, from Theorem 4.1, we get

$$tr\left(P_{g}P_{\Sigma}P_{g}\right) = \sum_{n \in \mathbb{N}} \left\langle P_{g}P_{\Sigma}P_{g}e_{n}, e_{n} \right\rangle_{L^{2}(\mathbb{R}^{2}, A_{\alpha, \beta} \otimes \sigma_{\alpha, \beta})} = \left\| P_{\Sigma}P_{g} \right\|_{HS}^{2} \leq A_{\alpha, \beta} \otimes \sigma_{\alpha, \beta}(\Sigma),$$

where $tr(P_g P_{\Sigma} P_g)$ denotes the trace of the operator $P_g P_{\Sigma} P_g$ and $\langle \cdot, \cdot \rangle_{L^2(\mathbb{R}^2, A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta})}$ denotes the inner product of $L^2(\mathbb{R}^2, A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta})$. Since $\{\phi_n\}_{n \in \mathbb{N}}$ be an orthonormal sequence in $L^2(\mathbb{R}, A_{\alpha,\beta})$, from the orthogonality relation (3.8), we obtain that $\{\mathcal{W}_g^{(\alpha,\beta)}(\phi_n)\}_{n \in \mathbb{N}}$ is also an orthonormal

sequence in $L^2(\mathbb{R}^2, A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta})$. Therefore

$$\sum_{n=1}^{N} \left\langle P_{\Sigma} \mathcal{W}_{g}^{(\alpha,\beta)}(\phi_{n}), \mathcal{W}_{g}^{(\alpha,\beta)}(\phi_{n}) \right\rangle_{L^{2}(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta})}$$
$$= \sum_{n=1}^{N} \left\langle P_{g} P_{\Sigma} P_{g} \mathcal{W}_{g}^{(\alpha,\beta)}(\phi_{n}), \mathcal{W}_{g}^{(\alpha,\beta)}(\phi_{n}) \right\rangle_{L^{2}(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta})}$$
$$\leq tr\left(P_{g} P_{\Sigma} P_{g}\right).$$

Thus, we have

$$\sum_{n=1}^{N} \left\langle P_{\Sigma} \mathcal{W}_{g}^{(\alpha,\beta)}(\phi_{n}), \mathcal{W}_{g}^{(\alpha,\beta)}(\phi_{n}) \right\rangle_{L^{2}(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta})} \leq A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta}(\Sigma).$$

Moreover, for any n with $1 \le n \le N$, using the Cauchy–Schwarz inequality, we get

$$\left\langle P_{\Sigma} \mathcal{W}_{g}^{(\alpha,\beta)}(\phi_{n}), \mathcal{W}_{g}^{(\alpha,\beta)}(\phi_{n}) \right\rangle_{L^{2}(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta})}$$

$$= 1 - \left\langle P_{\Sigma^{c}} \mathcal{W}_{g}^{(\alpha,\beta)}(\phi_{n}), \mathcal{W}_{g}^{(\alpha,\beta)}(\phi_{n}) \right\rangle_{L^{2}(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta})}$$

$$\geq 1 - \left\| \chi_{\Sigma^{c}} \mathcal{W}_{g}^{(\alpha,\beta)}(\phi_{n}) \right\|_{L^{2}(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta})}.$$

Therefore

$$\sum_{n=1}^{N} \left(1 - \left\| \chi_{\Sigma^{c}} \mathcal{W}_{g}^{(\alpha,\beta)}(\phi_{n}) \right\|_{L^{2}(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta})} \right) \leq \sum_{n=1}^{N} \left\langle P_{\Sigma} \mathcal{W}_{g}^{(\alpha,\beta)}(\phi_{n}), \mathcal{W}_{g}^{(\alpha,\beta)}(\phi_{n}) \right\rangle_{L^{2}(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta})} \leq A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta}(\Sigma).$$

This completes the proof of the theorem.

4.2. Donoho–Stark's uncertainty principle for the windowed Opdam–Cherednik transform. Here, we study the version of Donoho–Stark's uncertainty principle for the windowed Opdam–Cherednik transform. In particular, we investigate the case where f and $\mathcal{W}_{g}^{(\alpha,\beta)}(f)$ are close to zero outside measurable sets. We start with the following result.

Theorem 4.3. Let $g \in L^2(\mathbb{R}, A_{\alpha,\beta})$ be a non-zero window function and $f \in L^2(\mathbb{R}, A_{\alpha,\beta})$ such that $f \neq 0$. Then for any $\Sigma \subset \mathbb{R}^2$ and $\varepsilon \geq 0$ such that

$$\iint_{\Sigma} \left| \mathcal{W}_{g}^{(\alpha,\beta)}(f)(x,\xi) \right|^{2} d(A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta})(x,\xi) \geq (1-\varepsilon) \|f\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})}^{2} \|g\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})}^{2},$$

we have

$$A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta}(\Sigma) \ge 1 - \varepsilon.$$

Proof. Using the relation (3.11), we get

$$(1-\varepsilon)\|f\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})}^{2}\|g\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})}^{2} \leq \iint_{\Sigma} \left|\mathcal{W}_{g}^{(\alpha,\beta)}(f)(x,\xi)\right|^{2} d(A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta})(x,\xi)$$
$$\leq \left\|\mathcal{W}_{g}^{(\alpha,\beta)}(f)\right\|_{L^{\infty}(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta})}^{2} A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta}(\Sigma)$$
$$\leq A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta}(\Sigma) \|f\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})}^{2}\|g\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})}^{2}.$$

Therefore, $A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta}(\Sigma) \geq 1 - \varepsilon$.

The following proposition shows that the windowed Opdam–Cherednik transform cannot be concentrated in any small set.

Proposition 4.4. Let $g \in L^2(\mathbb{R}, A_{\alpha,\beta})$ be a non-zero window function. Then for any function $f \in L^2(\mathbb{R}, A_{\alpha,\beta})$ and for any $\Sigma \subset \mathbb{R}^2$ such that $A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta}(\Sigma) < 1$, we have

$$\left\|\chi_{\Sigma^{c}}\mathcal{W}_{g}^{(\alpha,\beta)}(f)\right\|_{L^{2}\left(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta}\right)} \geq \sqrt{1-A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta}(\Sigma)} \|f\|_{L^{2}\left(\mathbb{R},A_{\alpha,\beta}\right)} \|g\|_{L^{2}\left(\mathbb{R},A_{\alpha,\beta}\right)}$$

Proof. For any function $f \in L^2(\mathbb{R}, A_{\alpha,\beta})$, using the relation (3.11), we get

$$\begin{split} \left\| \mathcal{W}_{g}^{(\alpha,\beta)}(f) \right\|_{L^{2}\left(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta}\right)}^{2} &= \left\| \chi_{\Sigma}\mathcal{W}_{g}^{(\alpha,\beta)}(f) + \chi_{\Sigma^{c}}\mathcal{W}_{g}^{(\alpha,\beta)}(f) \right\|_{L^{2}\left(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta}\right)}^{2} \\ &= \left\| \chi_{\Sigma}\mathcal{W}_{g}^{(\alpha,\beta)}(f) \right\|_{L^{2}\left(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta}\right)}^{2} + \left\| \chi_{\Sigma^{c}}\mathcal{W}_{g}^{(\alpha,\beta)}(f) \right\|_{L^{2}\left(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta}\right)}^{2} \\ &\leq A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta}(\Sigma) \left\| \mathcal{W}_{g}^{(\alpha,\beta)}(f) \right\|_{L^{\infty}\left(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta}\right)}^{2} + \left\| \chi_{\Sigma^{c}}\mathcal{W}_{g}^{(\alpha,\beta)}(f) \right\|_{L^{2}\left(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta}\right)}^{2} \\ &\leq A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta}(\Sigma) \left\| f \right\|_{L^{2}\left(\mathbb{R},A_{\alpha,\beta}\right)}^{2} \left\| g \right\|_{L^{2}\left(\mathbb{R},A_{\alpha,\beta}\right)}^{2} + \left\| \chi_{\Sigma^{c}}\mathcal{W}_{g}^{(\alpha,\beta)}(f) \right\|_{L^{2}\left(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta}\right)}^{2} . \end{split}$$

Thus, using Plancherel's formula (3.7), we obtain

$$\left\|\chi_{\Sigma^{c}}\mathcal{W}_{g}^{(\alpha,\beta)}(f)\right\|_{L^{2}\left(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta}\right)} \geq \sqrt{1-A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta}(\Sigma)} \|f\|_{L^{2}\left(\mathbb{R},A_{\alpha,\beta}\right)}\|g\|_{L^{2}\left(\mathbb{R},A_{\alpha,\beta}\right)}.$$

Definition 4.5. Let *E* be a measurable subset of \mathbb{R} and $0 \leq \varepsilon_E < 1$. Then we say that a function $f \in L^p(\mathbb{R}, A_{\alpha,\beta}), 1 \leq p \leq 2$, is ε_E -concentrated on *E* in $L^p(\mathbb{R}, A_{\alpha,\beta})$ -norm, if

$$\|\chi_{E^c}f\|_{L^p(\mathbb{R},A_{\alpha,\beta})} \le \varepsilon_E \|f\|_{L^p(\mathbb{R},A_{\alpha,\beta})}.$$

If $\varepsilon_E = 0$, then E contains the support of f.

Definition 4.6. Let Σ be a measurable subset of \mathbb{R}^2 and $0 \leq \varepsilon_{\Sigma} < 1$. Let $f, g \in L^2(\mathbb{R}, A_{\alpha,\beta})$ be two non-zero functions. We say that $\mathcal{W}_q^{(\alpha,\beta)}(f)$ is ε_{Σ} -time-frequency concentrated on Σ , if

$$\left\|\chi_{\Sigma^{c}}\mathcal{W}_{g}^{(\alpha,\beta)}(f)\right\|_{L^{2}\left(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta}\right)}\leq\varepsilon_{\Sigma}\left\|\mathcal{W}_{g}^{(\alpha,\beta)}(f)\right\|_{L^{2}\left(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta}\right)}$$

If $\mathcal{W}_{g}^{(\alpha,\beta)}(f)$ is ε_{Σ} -time-frequency concentrated on Σ , then in the following, we obtain an estimate for the size of the essential support of the windowed Opdam–Cherednik transform.

Theorem 4.7. Let $g \in L^2(\mathbb{R}, A_{\alpha,\beta})$ be a non-zero window function and $f \in L^2(\mathbb{R}, A_{\alpha,\beta})$ such that $f \neq 0$. Let $\Sigma \subset \mathbb{R}^2$ such that $A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta}(\Sigma) < \infty$ and $\varepsilon_{\Sigma} \ge 0$. If $\mathcal{W}_g^{(\alpha,\beta)}(f)$ is ε_{Σ} -time-frequency concentrated on Σ , then

$$A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta}(\Sigma) \ge (1 - \varepsilon_{\Sigma}^2).$$

Proof. Since $\mathcal{W}_g^{(\alpha,\beta)}(f)$ is ε_{Σ} -time-frequency concentrated on Σ , using Plancherel's formula (3.7), we deduce that

$$\begin{split} \|f\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})}^{2} \|g\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})}^{2} &= \left\|\mathcal{W}_{g}^{(\alpha,\beta)}(f)\right\|_{L^{2}(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta})}^{2} \\ &= \left\|\chi_{\Sigma^{c}}\mathcal{W}_{g}^{(\alpha,\beta)}(f)\right\|_{L^{2}(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta})}^{2} + \left\|\chi_{\Sigma}\mathcal{W}_{g}^{(\alpha,\beta)}(f)\right\|_{L^{2}(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta})}^{2} \\ &\leq \varepsilon_{\Sigma}^{2} \left\|\mathcal{W}_{g}^{(\alpha,\beta)}(f)\right\|_{L^{2}(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta})}^{2} + \left\|\chi_{\Sigma}\mathcal{W}_{g}^{(\alpha,\beta)}(f)\right\|_{L^{2}(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta})}^{2}. \end{split}$$

Hence, using the relation (3.11), we obtain

$$(1 - \varepsilon_{\Sigma}^{2}) \|f\|_{L^{2}(\mathbb{R}, A_{\alpha, \beta})}^{2} \|g\|_{L^{2}(\mathbb{R}, A_{\alpha, \beta})}^{2} \leq \left\|\chi_{\Sigma} \mathcal{W}_{g}^{(\alpha, \beta)}(f)\right\|_{L^{2}(\mathbb{R}^{2}, A_{\alpha, \beta} \otimes \sigma_{\alpha, \beta})}^{2} \qquad (4.2)$$

$$\leq \left\|\mathcal{W}_{g}^{(\alpha, \beta)}(f)\right\|_{L^{\infty}(\mathbb{R}^{2}, A_{\alpha, \beta} \otimes \sigma_{\alpha, \beta})}^{2} A_{\alpha, \beta} \otimes \sigma_{\alpha, \beta}(\Sigma)$$

$$\leq \|f\|_{L^{2}(\mathbb{R}, A_{\alpha, \beta})}^{2} \|g\|_{L^{2}(\mathbb{R}, A_{\alpha, \beta})}^{2} A_{\alpha, \beta} \otimes \sigma_{\alpha, \beta}(\Sigma),$$

which completes the proof.

Theorem 4.8. Let $\Sigma \subset \mathbb{R}^2$ such that $A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta}(\Sigma) < \infty$, $\varepsilon_{\Sigma} \geq 0$, $g \in L^2(\mathbb{R}, A_{\alpha,\beta})$ be a nonzero window function, and $f \in L^2(\mathbb{R}, A_{\alpha,\beta})$ such that $f \neq 0$. If $\mathcal{W}_g^{(\alpha,\beta)}(f)$ is ε_{Σ} -time-frequency concentrated on Σ , then for every p > 2, we have

$$A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta}(\Sigma) \ge (1 - \varepsilon_{\Sigma}^2)^{\frac{p}{p-2}}$$

Proof. Since $\mathcal{W}_{g}^{(\alpha,\beta)}(f)$ is ε_{Σ} -time-frequency concentrated on Σ , from (4.2), we have

$$(1 - \varepsilon_{\Sigma}^2) \|f\|_{L^2(\mathbb{R}, A_{\alpha, \beta})}^2 \|g\|_{L^2(\mathbb{R}, A_{\alpha, \beta})}^2 \le \left\|\chi_{\Sigma} \mathcal{W}_g^{(\alpha, \beta)}(f)\right\|_{L^2(\mathbb{R}^2, A_{\alpha, \beta} \otimes \sigma_{\alpha, \beta})}^2$$

Again, aplying Hölder's inequality for the conjugate exponents $\frac{p}{2}$ and $\frac{p}{p-2}$, we get

$$\left\|\chi_{\Sigma}\mathcal{W}_{g}^{(\alpha,\beta)}(f)\right\|_{L^{2}(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta})}^{2} \leq \left\|\mathcal{W}_{g}^{(\alpha,\beta)}(f)\right\|_{L^{p}(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta})}^{2} \left(A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta}(\Sigma)\right)^{\frac{p-2}{p}}.$$

Now, using the relation (3.12), we obtain

$$\left\|\chi_{\Sigma}\mathcal{W}_{g}^{(\alpha,\beta)}(f)\right\|_{L^{2}(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta})}^{2} \leq \|f\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})}^{2}\|g\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})}^{2}(A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta}(\Sigma))^{\frac{p-2}{p}}.$$

Hence,

$$A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta}(\Sigma) \ge (1 - \varepsilon_{\Sigma}^2)^{\frac{p}{p-2}}.$$

Theorem 4.9. Let $\Sigma \subset \mathbb{R}^2$ such that $A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta}(\Sigma) < \infty$, $g \in L^2(\mathbb{R}, A_{\alpha,\beta})$ be a non-zero window function, and $f \in L^1(\mathbb{R}, A_{\alpha,\beta}) \cap L^2(\mathbb{R}, A_{\alpha,\beta})$ such that $\|\mathcal{W}_g^{(\alpha,\beta)}(f)\|_{L^2(\mathbb{R}^2, A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta})} = 1$. Let $E \subset \mathbb{R}$ such that $A_{\alpha,\beta}(E) < \infty$. If f is ε_E -concentrated on E in $L^1(\mathbb{R}, A_{\alpha,\beta})$ -norm and $\mathcal{W}_q^{(\alpha,\beta)}(f)$ is ε_{Σ} -time-frequency concentrated on Σ , then

$$A_{\alpha,\beta}(E) \ge (1-\varepsilon_E)^2 \|f\|_{L^1(\mathbb{R},A_{\alpha,\beta})}^2 \|g\|_{L^2(\mathbb{R},A_{\alpha,\beta})}^2,$$

and

$$\|f\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})}^{2} \|g\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})}^{2} A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta}(\Sigma) \geq (1-\varepsilon_{\Sigma}^{2}).$$

In particular,

$$A_{\alpha,\beta}(E) \ A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta}(\Sigma) \ \|f\|_{L^2(\mathbb{R},A_{\alpha,\beta})}^2 \ge (1-\varepsilon_E)^2 \ (1-\varepsilon_{\Sigma}^2) \ \|f\|_{L^1(\mathbb{R},A_{\alpha,\beta})}^2$$

Proof. Since $\mathcal{W}_{g}^{(\alpha,\beta)}(f)$ is ε_{Σ} -time-frequency concentrated on Σ , from (4.2), we have

$$(1 - \varepsilon_{\Sigma}^{2}) \|f\|_{L^{2}(\mathbb{R}, A_{\alpha, \beta})}^{2} \|g\|_{L^{2}(\mathbb{R}, A_{\alpha, \beta})}^{2} \leq \left\|\chi_{\Sigma} \mathcal{W}_{g}^{(\alpha, \beta)}(f)\right\|_{L^{2}(\mathbb{R}^{2}, A_{\alpha, \beta} \otimes \sigma_{\alpha, \beta})}^{2}$$

Since $\|\mathcal{W}_{g}^{(\alpha,\beta)}(f)\|_{L^{2}(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta})} = 1$, using the relation (3.11) and Plancherel's formula (3.7), we obtain

$$(1 - \varepsilon_{\Sigma}^{2}) \leq \left\| \chi_{\Sigma} \mathcal{W}_{g}^{(\alpha,\beta)}(f) \right\|_{L^{2}(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta})}^{2} \\ \leq \left\| \mathcal{W}_{g}^{(\alpha,\beta)}(f) \right\|_{L^{\infty}(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta})}^{2} A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta}(\Sigma) \\ \leq \|f\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})}^{2} \|g\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})}^{2} A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta}(\Sigma).$$

$$(4.3)$$

Similarly, since f is ε_E -concentrated on E in $L^1(\mathbb{R}, A_{\alpha,\beta})$ -norm, using the Cauchy–Schwarz inequality and the fact that $\|f\|_{L^2(\mathbb{R}, A_{\alpha,\beta})} \|g\|_{L^2(\mathbb{R}, A_{\alpha,\beta})} = 1$, we get

$$(1 - \varepsilon_E) \|f\|_{L^1(\mathbb{R}, A_{\alpha, \beta})} \le \|\chi_E f\|_{L^1(\mathbb{R}, A_{\alpha, \beta})} \le \|f\|_{L^2(\mathbb{R}, A_{\alpha, \beta})} A_{\alpha, \beta}(E)^{\frac{1}{2}} = \frac{A_{\alpha, \beta}(E)^{\frac{1}{2}}}{\|g\|_{L^2(\mathbb{R}, A_{\alpha, \beta})}}.$$
 (4.4)

Finally, from (4.3) and (4.4), we have

$$A_{\alpha,\beta}(E) \ A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta}(\Sigma) \ \|f\|_{L^2(\mathbb{R},A_{\alpha,\beta})}^2 \ge (1-\varepsilon_E)^2 \ (1-\varepsilon_{\Sigma}^2) \ \|f\|_{L^1(\mathbb{R},A_{\alpha,\beta})}^2.$$

This completes the proof of the theorem.

4.3. Benedicks-type uncertainty principle for the windowed Opdam–Cherednik transform. In this subsection, we study Benedicks-type uncertainty principle for the windowed Opdam–Cherednik transform. The following proposition shows that this transform cannot be concentrated inside a set of measure arbitrarily small.

Proposition 4.10. Let $g \in L^2(\mathbb{R}, A_{\alpha,\beta})$ be a non-zero window function and $\Sigma \subset \mathbb{R}^2$. If $\|P_{\Sigma}P_g\| < 1$, then there exists a constant $c(\Sigma, g) > 0$ such that for every $f \in L^2(\mathbb{R}, A_{\alpha,\beta})$, we have

$$\|f\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})}\|g\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})} \leq c(\Sigma,g) \left\|\chi_{\Sigma^{c}}\mathcal{W}_{g}^{(\alpha,\beta)}(f)\right\|_{L^{2}\left(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta}\right)}.$$

Proof. Since P_{Σ} is an orthogonal projection on $L^2(\mathbb{R}^2, A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta})$, for any $F \in L^2(\mathbb{R}^2, A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta})$, we get

$$\|P_g(F)\|_{L^2(\mathbb{R}^2, A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta})}^2 = \|P_{\Sigma}P_g(F)\|_{L^2(\mathbb{R}^2, A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta})}^2 + \|P_{\Sigma^c}P_g(F)\|_{L^2(\mathbb{R}^2, A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta})}^2$$

Again, using the identity $P_{\Sigma}P_g(F) = P_{\Sigma}P_g \cdot P_g(F)$, we get

$$\|P_{\Sigma}P_g(F)\|^2_{L^2(\mathbb{R}^2,A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta})} \leq \|P_{\Sigma}P_g\|^2 \|P_g(F)\|^2_{L^2(\mathbb{R}^2,A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta})}.$$

Hence,

$$\|P_g(F)\|^2_{L^2\left(\mathbb{R}^2, A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta}\right)} \le \frac{1}{1 - \|P_{\Sigma}P_g\|^2} \|P_{\Sigma^c}P_g(F)\|^2_{L^2\left(\mathbb{R}^2, A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta}\right)}.$$
(4.5)

Since P_g is an orthogonal projection from $L^2(\mathbb{R}^2, A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta})$ onto $\mathcal{W}_g^{(\alpha,\beta)}(L^2(\mathbb{R}, A_{\alpha,\beta}))$, for any $f \in L^2(\mathbb{R}, A_{\alpha,\beta})$, using the relation (4.5) and Plancherel's formula (3.7), we get

$$\|f\|_{L^2(\mathbb{R},A_{\alpha,\beta})}\|g\|_{L^2(\mathbb{R},A_{\alpha,\beta})} \leq \frac{1}{\sqrt{1-\|P_{\Sigma}P_g\|^2}} \left\|\chi_{\Sigma^c}\mathcal{W}_g^{(\alpha,\beta)}(f)\right\|_{L^2(\mathbb{R}^2,A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta})}.$$

The desired result follows by choosing the constant $c(\Sigma, g) = \frac{1}{\sqrt{1 - \|P_{\Sigma}P_g\|^2}}$.

Next, we obtain Benedicks-type uncertainty principle for the windowed Opdam–Cherednik transform.

11

Theorem 4.11. Let $g \in L^2(\mathbb{R}, A_{\alpha,\beta})$ be a non-zero window function such that

$$\sigma_{\alpha,\beta}\left(\{\mathcal{H}_{\alpha,\beta}(g)\neq 0\}\right)<\infty.$$

Then for any $\Sigma \subset \mathbb{R}^2$ such that for almost every $\xi \in \mathbb{R}$, $\int_{\mathbb{R}} \chi_{\Sigma}(x,\xi) A_{\alpha,\beta}(x) dx < \infty$, we have

$$\mathcal{W}_{q}^{(\alpha,\beta)}\left(L^{2}(\mathbb{R},A_{\alpha,\beta})\right)\cap\operatorname{Im}P_{\Sigma}=\{0\}.$$

Proof. Let $F \in \mathcal{W}_g^{(\alpha,\beta)}(L^2(\mathbb{R}, A_{\alpha,\beta})) \cap \operatorname{Im} P_{\Sigma}$ be a non-trivial function. Then there exists a function $f \in L^2(\mathbb{R}, A_{\alpha,\beta})$ such that $F = \mathcal{W}_g^{(\alpha,\beta)}(f)$ and $Supp F \subset \Sigma$. For any $\xi \in \mathbb{R}$, we consider the function

$$F_{\xi}(x) = \mathcal{W}_{g}^{(\alpha,\beta)}(f)(x,\xi), \qquad x \in \mathbb{R}.$$

Then, we get $SuppF_{\xi} \subset \{x \in \mathbb{R} : (x,\xi) \in \Sigma\}$. Since for almost every $\xi \in \mathbb{R}, \int_{\mathbb{R}} \chi_{\Sigma}(x,\xi) A_{\alpha,\beta}(x) dx < \infty$, we have $A_{\alpha,\beta}(SuppF_{\xi}) < \infty$. Using the relation (3.5), we get

$$\mathcal{H}_{\alpha,\beta}(F_{\xi}) = \mathcal{H}_{\alpha,\beta}(f) \ \mathcal{H}_{\alpha,\beta}(\mathcal{M}_{\xi}^{(\alpha,\beta)}g) \quad a.e.$$

Hence,

$$Supp\mathcal{H}_{\alpha,\beta}(F_{\xi}) \subset Supp\tau_{\xi}^{(\alpha,\beta)}|\mathcal{H}_{\alpha,\beta}(g)|^{2},$$

and from the hypothesis, we get $\sigma_{\alpha,\beta}(\{\mathcal{H}_{\alpha,\beta}(F_{\xi})\neq 0\})<\infty$. From Benedicks-type result for the Opdam–Cherednik transform [1], we conclude that, for every $\xi \in \mathbb{R}, F(\cdot,\xi) = 0$, which eventually implies that F = 0.

Remark 4.12. Let $g \in L^2(\mathbb{R}, A_{\alpha,\beta})$ be a non-zero window function such that

$$\sigma_{\alpha,\beta}\left(\{\mathcal{H}_{\alpha,\beta}(g)\neq 0\}\right)<\infty.$$

Then, for any non-zero function $f \in L^2(\mathbb{R}, A_{\alpha,\beta})$, we have $A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta}(Supp\mathcal{W}_g^{(\alpha,\beta)}(f)) = \infty$, i.e., the support of $\mathcal{W}_g^{(\alpha,\beta)}(f)$ cannot be of finite measure.

Proposition 4.13. Let $g \in L^2(\mathbb{R}, A_{\alpha,\beta})$ be a non-zero window function such that

$$\sigma_{\alpha,\beta}\left(\{\mathcal{H}_{\alpha,\beta}(g)\neq 0\}\right)<\infty.$$

Let $\Sigma \subset \mathbb{R}^2$ such that $A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta}(\Sigma) < \infty$, then there exists a constant $c(\Sigma,g) > 0$ such that

$$\|f\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})} \leq c(\Sigma,g) \left\|\chi_{\Sigma^{c}}\mathcal{W}_{g}^{(\alpha,\beta)}(f)\right\|_{L^{2}\left(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta}\right)}$$

Proof. Assume that

$$\|P_{\Sigma}P_g(F)\|_{L^2(\mathbb{R}^2,A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta})} = \|F\|_{L^2(\mathbb{R}^2,A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta})}, \ F \in L^2(\mathbb{R}^2,A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta}).$$

Since P_{Σ} and P_g are orthogonal projections, we obtain $P_{\Sigma}(F) = P_g(F) = F$. Again, since $A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta}(\Sigma) < \infty$, for almost every $\xi \in \mathbb{R}$, $\int_{\mathbb{R}} \chi_{\Sigma}(x,\xi) A_{\alpha,\beta}(x) dx < \infty$ and from Theorem 4.11, we get F = 0. Hence, for $F \neq 0$, we obtain

$$\|P_{\Sigma}P_g(F)\|_{L^2(\mathbb{R}^2,A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta})} < \|F\|_{L^2(\mathbb{R}^2,A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta})}.$$

Since $P_{\Sigma}P_g$ is a Hilbert–Schmidt operator, we obtain that the largest eigenvalue $|\lambda|$ of the operator $P_{\Sigma}P_g$ satisfy $|\lambda| < 1$ and $||P_{\Sigma}P_g|| = |\lambda| < 1$. Finally, using Proposition 4.10, we get the desired result.

4.4. Heisenberg-type uncertainty principle for the windowed Opdam–Cherednik transform. This subsection is devoted to study Heisenberg-type uncertainty inequality for the windowed Opdam–Cherednik transform for general magnitude s > 0. Indeed, we have the following result.

Theorem 4.14. Let s > 0. Then there exists a constant $c(s, \alpha, \beta) > 0$ such that for all $f, g \in L^2(\mathbb{R}, A_{\alpha,\beta})$, we have

$$\begin{aligned} \left\| x^{s} \mathcal{W}_{g}^{(\alpha,\beta)}(f) \right\|_{L^{2}(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta})}^{2} + \left\| \xi^{s} \mathcal{W}_{g}^{(\alpha,\beta)}(f) \right\|_{L^{2}(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta})}^{2} \\ \geq c(s,\alpha,\beta) \left\| f \right\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})}^{2} \left\| g \right\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})}^{2}. \end{aligned}$$

Proof. Let r > 0 and $B_r = \{(x,\xi) \in \mathbb{R}^2 : |(x,\xi)| < r\}$ be the ball with centre at origin and radius r in \mathbb{R}^2 . Fix $\varepsilon_0 \leq 1$ small enough such that $A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta}(B_{\varepsilon_0}) < 1$. From Proposition 4.4, we have

$$\begin{split} \|f\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})}^{2} \|g\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})}^{2} \\ &\leq \frac{1}{\left[1-A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta}(B_{\varepsilon_{0}})\right]} \iint_{|(x,\xi)|\geq\varepsilon_{0}} \left|\mathcal{W}_{g}^{(\alpha,\beta)}(f)(x,\xi)\right|^{2} \, d(A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta})(x,\xi) \\ &\leq \frac{1}{\varepsilon_{0}^{2s}\left[1-A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta}(B_{\varepsilon_{0}})\right]} \iint_{|(x,\xi)|\geq\varepsilon_{0}} |(x,\xi)|^{2s} \left|\mathcal{W}_{g}^{(\alpha,\beta)}(f)(x,\xi)\right|^{2} \, d(A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta})(x,\xi) \\ &\leq \frac{1}{\varepsilon_{0}^{2s}\left[1-A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta}(B_{\varepsilon_{0}})\right]} \iint_{\mathbb{R}^{2}} |(x,\xi)|^{2s} \left|\mathcal{W}_{g}^{(\alpha,\beta)}(f)(x,\xi)\right|^{2} \, d(A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta})(x,\xi). \end{split}$$

Consequentely,

$$\left\| |(x,\xi)|^{s} \mathcal{W}_{g}^{(\alpha,\beta)}(f) \right\|_{L^{2}(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta})}^{2} \geq \varepsilon_{0}^{2s} \left[1 - A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta}(B_{\varepsilon_{0}}) \right] \|f\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})}^{2} \|g\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})}^{2} \cdot \left(4.6 \right)$$

$$(4.6)$$

Finally, using the fact that $|a+b|^s \leq 2^s (|a|^s + |b|^s)$ and from (4.6), we get

$$\begin{split} & \varepsilon_{0}^{2s} \left[1 - A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta}(B_{\varepsilon_{0}}) \right] \left\| f \right\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})}^{2} \left\| g \right\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})}^{2} \\ & \leq \left\| |(x,\xi)|^{s} \mathcal{W}_{g}^{(\alpha,\beta)}(f) \right\|_{L^{2}(\mathbb{R}^{2},A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta})}^{2} \\ & \leq 2^{s} \left\| x^{s} \mathcal{W}_{g}^{(\alpha,\beta)}(f) \right\|_{L^{2}(\mathbb{R}^{2},A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta})}^{2} + 2^{s} \left\| \xi^{s} \mathcal{W}_{g}^{(\alpha,\beta)}(f) \right\|_{L^{2}(\mathbb{R}^{2},A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta})}^{2}. \end{split}$$

Hence, we get the desired result with $c(s, \alpha, \beta) = \varepsilon_0^{2s} 2^{-s} \left[1 - A_{\alpha, \beta} \otimes \sigma_{\alpha, \beta}(B_{\varepsilon_0})\right].$

4.5. Local uncertainty inequality for the windowed Opdam–Cherednik transform. Here, we discuss results related to the $L^2(\mathbb{R}, A_{\alpha,\beta})$ -mass of the windowed Opdam–Cherednik transform outside sets of finite measure. Indeed, we establish the following result.

Theorem 4.15. Let s > 0. Then there exists a constant $c(s, \alpha, \beta) > 0$ such that for any $f, g \in L^2(\mathbb{R}, A_{\alpha,\beta})$ and any $\Sigma \subset \mathbb{R}^2$ such that $A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta}(\Sigma) < \infty$, we have

$$\left\| \mathcal{W}_{g}^{(\alpha,\beta)}(f) \right\|_{L^{2}(\Sigma,A_{\alpha,\beta})} \leq c(s,\alpha,\beta) \left[A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta}(\Sigma) \right]^{\frac{1}{2}} \left\| |(x,\xi)|^{s} \mathcal{W}_{g}^{(\alpha,\beta)}(f) \right\|_{L^{2}(\mathbb{R}^{2},A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta})}.$$

Proof. Since $\|\mathcal{W}_{g}^{(\alpha,\beta)}(f)\|_{L^{2}(\Sigma,A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta})} \leq \|\mathcal{W}_{g}^{(\alpha,\beta)}(f)\|_{L^{\infty}(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta})} [A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta}(\Sigma)]^{\frac{1}{2}}$, using the relation (3.11), we get

$$\left| \mathcal{W}_{g}^{(\alpha,\beta)}(f) \right\|_{L^{2}\left(\Sigma,A_{\alpha,\beta}\right)} \leq \left[A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta}(\Sigma) \right]^{\frac{1}{2}} \|f\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})} \|g\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})}$$

Thus, using Heisenberg's inequality (4.6), we get

$$\left\| \mathcal{W}_{g}^{(\alpha,\beta)}(f) \right\|_{L^{2}\left(\Sigma,A_{\alpha,\beta}\right)} \leq \frac{\left[A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta}(\Sigma)\right]^{\frac{1}{2}}}{\varepsilon_{0}^{s} \left[1 - A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta}(B_{\varepsilon_{0}})\right]^{\frac{1}{2}}} \left\| |(x,\xi)|^{s} \mathcal{W}_{g}^{(\alpha,\beta)}(f) \right\|_{L^{2}\left(\mathbb{R}^{2},A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta}\right)}.$$

This completes the proof of the theorem with $c(s, \alpha, \beta) = \varepsilon_0^{-s} \left[1 - A_{\alpha, \beta} \otimes \sigma_{\alpha, \beta}(B_{\varepsilon_0})\right]^{-\frac{1}{2}}$.

The following result shows that, Theorem 4.15 gives a general form of Heisenberg-type inequality with a different constant.

Corollary 4.16. Let s > 0. Then there exists a constant $c_{s,\alpha,\beta} > 0$ such that, for all $f,g \in L^2(\mathbb{R}, A_{\alpha,\beta})$, we have

$$\left\| |(x,\xi)|^{s} \mathcal{W}_{g}^{(\alpha,\beta)}(f) \right\|_{L^{2}(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta})} \geq c_{s,\alpha,\beta} \|f\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})} \|g\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})}.$$
(4.7)

In particular,

$$\begin{split} & \left\| x^{s} \mathcal{W}_{g}^{(\alpha,\beta)}(f) \right\|_{L^{2}(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta})}^{2} + \left\| \xi^{s} \mathcal{W}_{g}^{(\alpha,\beta)}(f) \right\|_{L^{2}(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta})}^{2} \\ & \geq \frac{c_{s,\alpha,\beta}^{2}}{2^{s}} \left\| f \right\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})}^{2} \left\| g \right\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})}^{2}. \end{split}$$

Proof. Let r > 0 and $B_r = \{(x,\xi) \in \mathbb{R}^2 : |(x,\xi)| < r\}$ be the ball with centre at origin and radius r in \mathbb{R}^2 . Using Plancherel's formula (3.7) and Theorem 4.15, we obtain

$$\begin{split} \|f\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})}^{2} \|g\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})}^{2} &= \left\|\mathcal{W}_{g}^{(\alpha,\beta)}(f)\right\|_{L^{2}(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta})}^{2} \\ &= \left\|\chi_{B_{r}}\mathcal{W}_{g}^{(\alpha,\beta)}(f)\right\|_{L^{2}(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta})}^{2} + \left\|\chi_{B_{r}^{c}}\mathcal{W}_{g}^{(\alpha,\beta)}(f)\right\|_{L^{2}(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta})}^{2} \\ &\leq c(s,\alpha,\beta)^{2} \left[A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta}(B_{r})\right] \left\||(x,\xi)|^{s}\mathcal{W}_{g}^{(\alpha,\beta)}(f)\right\|_{L^{2}(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta})}^{2} \\ &+ r^{-2s} \left\||(x,\xi)|^{s}\mathcal{W}_{g}^{(\alpha,\beta)}(f)\right\|_{L^{2}(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta})}^{2}. \end{split}$$

We get the inequality (4.7) by minimizing the right-hand side of the above inequality over r > 0. Now, proceeding similarly as in the proof of Theorem 4.14, we obtain

$$2^{s} \left\| x^{s} \mathcal{W}_{g}^{(\alpha,\beta)}(f) \right\|_{L^{2}(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta})}^{2} + 2^{s} \left\| \xi^{s} \mathcal{W}_{g}^{(\alpha,\beta)}(f) \right\|_{L^{2}(\mathbb{R}^{2},A_{\alpha,\beta}\otimes\sigma_{\alpha,\beta})}^{2}$$
$$\geq c_{s,\alpha,\beta}^{2} \left\| f \right\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})}^{2} \left\| g \right\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})}^{2},$$

and this completes the proof.

4.6. Heisenberg-type uncertainty inequality via the k-entropy. In this subsection, we study the localization of the k-entropy of the windowed Opdam–Cherednik transform over the space \mathbb{R}^2 . Before going to prove the main result, we first need the following definition.

Definition 4.17.

(1) A probability density function ρ on \mathbb{R}^2 is a non-negative measurable function on \mathbb{R}^2 satisfying

$$\iint_{\mathbb{R}^2} \rho(x,\xi) \ d(A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta})(x,\xi) = 1.$$

(2) Let ρ be a probability density function on \mathbb{R}^2 . Then the k-entropy of ρ is defined by

$$E_k(\rho) := -\iint_{\mathbb{R}^2} \ln(\rho(x,\xi)) \ \rho(x,\xi) \ d(A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta})(x,\xi),$$

whenever the integral on the right-hand side is well defined.

In the following, we prove the main result of this subsection.

Theorem 4.18. Let $g \in L^2(\mathbb{R}, A_{\alpha,\beta})$ be a non-zero window function and $f \in L^2(\mathbb{R}, A_{\alpha,\beta})$ such that $f \neq 0$. Then, we have

$$E_{k}(|\mathcal{W}_{g}^{(\alpha,\beta)}(f)|^{2}) \geq -2\ln\left(\|f\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})} \|g\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})}\right) \|f\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})}^{2} \|g\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})}^{2}.$$
(4.8)

Proof. First, we assume that $||f||_{L^2(\mathbb{R},A_{\alpha,\beta})} = ||g||_{L^2(\mathbb{R},A_{\alpha,\beta})} = 1$. For any $(x,\xi) \in \mathbb{R}^2$, using the relation (3.11), we get

$$|\mathcal{W}_g^{(\alpha,\beta)}(f)(x,\xi)| \le \|f\|_{L^2(\mathbb{R},A_{\alpha,\beta})} \|g\|_{L^2(\mathbb{R},A_{\alpha,\beta})} = 1.$$

Consequently, $\ln(|\mathcal{W}_g^{(\alpha,\beta)}(f)|) \leq 0$ and therefore $E_k(|\mathcal{W}_g^{(\alpha,\beta)}(f)|) \geq 0$. The desired inequality (4.8) holds trivially if the entropy $E_k(|\mathcal{W}_g^{(\alpha,\beta)}(f)|)$ is infinite. Now, suppose that the entropy $E_k(|\mathcal{W}_g^{(\alpha,\beta)}(f)|)$ is finite. Let f and g be two non-zero functions in $L^2(\mathbb{R}, A_{\alpha,\beta})$. We define

$$\phi = \frac{f}{\|f\|_{L^2(\mathbb{R}, A_{\alpha, \beta})}} \quad \text{and} \quad \psi = \frac{g}{\|g\|_{L^2(\mathbb{R}, A_{\alpha, \beta})}}$$

Then ϕ , $\psi \in L^2(\mathbb{R}, A_{\alpha,\beta})$ with $\|\phi\|_{L^2(\mathbb{R}, A_{\alpha,\beta})} = \|\psi\|_{L^2(\mathbb{R}, A_{\alpha,\beta})} = 1$ and consequently

$$E_k(|\mathcal{W}_{\psi}^{(\alpha,\beta)}(\phi)|) \ge 0.$$
(4.9)

Since
$$\mathcal{W}_{\psi}^{(\alpha,\beta)}(\phi) = \frac{1}{\|f\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})} \|g\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})}} \mathcal{W}_{g}^{(\alpha,\beta)}(f)$$
, we have

$$E_{k}(|\mathcal{W}_{\psi}^{(\alpha,\beta)}(\phi)|^{2}) = -\iint_{\mathbb{R}^{2}} \ln(|\mathcal{W}_{\psi}^{(\alpha,\beta)}(\phi)(x,\xi)|^{2}) |\mathcal{W}_{\psi}^{(\alpha,\beta)}(\phi)(x,\xi)|^{2} d(A_{\alpha,\beta} \otimes \sigma_{\alpha,\beta})(x,\xi)$$

$$= \frac{1}{\|f\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})}^{2} \|g\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})}^{2}} E_{k}(|\mathcal{W}_{g}^{(\alpha,\beta)}(f)|^{2})$$

$$+ 2\ln(\|f\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})} \|g\|_{L^{2}(\mathbb{R},A_{\alpha,\beta})}^{2}).$$

Finally, from (4.9), we obtain

$$E_k(|\mathcal{W}_g^{(\alpha,\beta)}(f)|^2) \ge -2\ln\left(\|f\|_{L^2(\mathbb{R},A_{\alpha,\beta})} \|g\|_{L^2(\mathbb{R},A_{\alpha,\beta})}\right) \|f\|_{L^2(\mathbb{R},A_{\alpha,\beta})}^2 \|g\|_{L^2(\mathbb{R},A_{\alpha,\beta})}^2.$$

completes the proof of the theorem.

This completes the proof of the theorem.

4.7. Application in signal processing. Here, we present an application of these uncertainty principles in compressive sensing. Mainly, we show that uncertainty principles can be used for the separation of signals. The signal separation problem is an extremely ill-defined signal processing problem, which is also important in many engineering problems. It consists in splitting a signal f into a sum of components f_k of different nature: $f = f_1 + f_2 + \cdots + f_n$. Since this notion of different nature often makes sense in applied domains, it is generally extremely difficult to formalize mathematically. Sparsity offers a convenient framework for approaching such a notion. More precisely, signals of different natures can be sparsely represented in different waveform systems. Given a union of several frames $\mathcal{U}^{(1)}, \mathcal{U}^{(2)}, \cdots, \mathcal{U}^{(n)}$ in a Hilbert space \mathbb{H} , the separation problem can be given various formulations, among which the analysis and synthesis formulations (see [41]).

Let $U^{(k)}$ denotes the analysis operator of frame k. In the case of n frames, applying these uncertainty principles and using the similar method as in [41], it can be proven that if one frame gives a splitting $f = f_1 + f_2 + \dots + f_n$, obtained via any algorithm, if $||U^{(1)}f_1|| + ||U^{(2)}f_2|| +$ $\cdots + \|U^{(n)}f_n\|$ is small enough, then this splitting is necessarily optimal. More precisely, we have the following result. Let $\mathcal{U}^{(1)}, \mathcal{U}^{(2)}, \cdots, \mathcal{U}^{(n)}$ denote n frames in \mathbb{H} . For any $f \in \mathbb{H}$, let $f = f_1 + f_2 + \cdots + f_n$ denote a splitting such that

$$\left\| U^{(1)}f_1 \right\| + \left\| U^{(2)}f_2 \right\| + \dots + \left\| U^{(n)}f_n \right\| < \frac{1}{\mu_{\star}}$$

where μ_{\star} is the generalized coherence function (see [41]). Then, using these uncertainty principles for the windowed Opdam–Cherednik transform, we obtain that this splitting minimizes $\|U^{(1)}f_1\| + \|U^{(2)}f_2\| + \cdots + \|U^{(n)}f_n\|$. Similarly, using these uncertainty principles one can study sparsity-based algorithms for window optimization in time-frequency analysis. Many other applications can be given using uncertainty principles for the windowed Opdam–Cherednik transform.

Acknowledgments

The first author gratefully acknowledges the support provided by IIT Guwahati, Government of India. The second author is deeply indebted to Prof. Nir Lev for several fruitful discussions and generous comments. The authors wish to thank the anonymous referees for their helpful comments and suggestions that helped to improve the quality of the paper.

Conflict of interest

No potential conflict of interest was reported by the authors.

ORCID

Shyam Swarup Mondal https://orcid.org/0000-0002-9778-0757 Anirudha Poria https://orcid.org/0000-0002-0224-3642

References

- A. Achak and R. Daher, Benedicks-Amrein-Berthier type theorem related to Opdam-Cherednik transform, J. Pseudo-Differ. Oper. Appl. 9:431-441 (2018). [2, 12]
- N.B. Andersen, Real Paley-Wiener theorems and Roe's theorem associated with the Opdam-Cherednik transform, J. Math. Anal. Appl. 427(1):47–59 (2015). [2, 3]
- [3] J.-P. Anker, F. Ayadi and M. Sifi, Opdam's hypergeometric functions: product formula and convolution structure in dimension 1, Adv. Pure Appl. Math. 3(1):11-44 (2012). [4, 5]
- [4] C. Baccar, Uncertainty principles for the continuous Hankel Wavelet transform, Integral Trans. Spec. Funct. 27(6):413-429 (2016).
- [5] M. Benedicks, On Fourier transforms of functions supported on sets of finite Lebesgue measure, J. Math. Anal. Appl. 106:180-183 (1985). [1]
- [6] I. Bialynicki-Birula, Entropic Uncertainty Relations in Quantum Mechanics, Quantum Probability and Applications II, Springer, Berlin 90–103 (1985). [1, 2]
- [7] A. Bonami, B. Demange and P. Jaming, Hermite functions and uncertainty principles for the Fourier and the windowed Fourier transforms, Rev. Mat. Iberoam. 19:23–55 (2003). [2]
- [8] K. Brahim and E. Tefjeni, Uncertainty principle for the two sided quaternion windowed Fourier transform, J. Pseudo-Differ. Oper. Appl. 11:159–185 (2020). [2]
- M.G. Cowling and J.F. Price, Generalizations of Heisenberg's inequality, In: Harmonic analysis (G. Mauceri, F. Ricci and G. Weiss (eds.)), Springer, Berlin 443–449 (1983). [1]
- [10] R. Daher, S.L. Hamad, T. Kawazoe and N. Shimeno, Uncertainty principles for the Cherednik transform, Proc. Indian Acad. Sci. Math. Sci. 122(3):429–436 (2012). [2]
- [11] L. Debnath and F.A. Shah, Wavelet Transforms and Their Applications, Birkhäuser, New York (2015). [1]
- [12] J.F. van Diejen and L. Vinet, Calogero-Moser-Sutherland Models, CRM Series in Mathematical Physics, Springer, New York (2000). [2]
- [13] D.L. Donoho and P.B. Stark, Uncertainty principles and signal recovery, SIAM J. Appl. Math. 49(3):906–931 (1989).
 [1, 2]
- [14] C.F. Dunkl, Hankel transforms associated to finite reflection groups, Contemp. Math. 138:123–138 (1992). [2]
- [15] G.B. Folland and A. Sitaram, The uncertainty principle: A mathematical survey, J. Fourier Anal. Appl. 3:207–238 (1997). [2]
- [16] W.B. Gao and B.Z. Li, Uncertainty principles for the windowed Hankel transform, Integral Trans. Spec. Funct. 31(12):982–997 (2020). [2]
- [17] S. Ghobber and S. Omri, *Time-frequency concentration of the windowed Hankel transform*, Integral Trans. Spec. Funct. 25(6):481–496 (2014). [2]
- [18] K. Gröchenig, Foundations of Time-Frequency Analysis, Birkhäuser, Boston (2001). [1, 2]
- [19] N.B. Hamadi and S. Omri, Uncertainty principles for the continuous wavelet transform in the Hankel setting, Appl. Anal. 97(4):513–527 (2017). [2]
- [20] A. Hammami, Uncertainty principles for the Hankel-Gabor transform, Indian J. Pure Appl. Math. 51(1):251– 264 (2020). [2]

- [21] G.H. Hardy, A theorem concerning Fourier transforms, J. Lond. Math. Soc. 8:227–231 (1933). [1]
- [22] V. Havin and B. Jöricke, The uncertainty principle in harmonic analysis, Springer-Verlag, Berlin (1994). [1, 2]
- [23] G.J. Heckman, An elementary approach to the hypergeometric shift operators of Opdam, Invent. Math. 103:341–350 (1991). [2]
- [24] K. Hikami, Dunkl operators formalism for quantum many-body problems associated with classical root systems, Phys. Soc. Japan 65:394–401 (1996). [2]
- [25] L. Hörmander, A uniqueness theorem of Beurling for Fourier transform pairs, Ark. Mat. 29:237–240 (1991).
 [1]
- [26] L.-L. Huang, D. Baleanu, Z.-W. Mo and G.-C. Wu, Fractional discrete-time diffusion equation with uncertainty: Applications of fuzzy discrete fractional calculus, Phys. A: Stat. Mech. Appl. 508:166–175 (2018).
 [2]
- [27] T.R. Johansen, Remarks on the inverse Cherednik–Opdam transform on the real line, arXiv:1502.01293 (2015). [3]
- [28] T.R. Johansen, Uncertainty principles for the Heckman-Opdam transform, Bull. Sci. Math. 140:687–717 (2016). [2]
- [29] H. Mejjaoli, Qualitative uncertainty principles for the Opdam-Cherednik transform, Integral Trans. Spec. Funct. 25(7):528-546 (2014). [2]
- [30] H. Mejjaoli, Spectral theorems associated with the Jacobi-Cherednik operator, Bull. Sci. Math. 138(3):416–439 (2014).
- [31] H. Mejjaoli, Uncertainty principles for the generalized Fourier transform associated to a Dunkl-type operator, Appl. Anal. 95(9):1930–1956 (2016). [2]
- [32] H. Mejjaoli, N. Sraieb and K. Trimèche, Inversion theorem and quantitative uncertainty principles for the Dunkl Gabor transform on R^d, J. Pseudo-Differ. Oper. Appl. 10:883–913 (2019). [2]
- [33] H. Mejjaoli and K. Trimèche, Qualitative uncertainty principles for the generalized Fourier transform associated to a Dunkl type operator on the real line, Anal. Math. Phys. 6:141–162 (2016). [2]
- [34] S.S. Mondal and A. Poria, Qualitative uncertainty principles for the windowed Opdam-Cherednik transform on weighted modulation spaces, Math. Meth. Appl. Sci. 45(16):10424-10439 (2022). [2]
- [35] S.S. Mondal and A. Poria, Weighted norm inequalities for the Opdam-Cherednik transform, Internat. J. Math. 33(9): Paper No. 2250066, 21 pp. (2022). [2]
- [36] M.W. Morgan, A note on Fourier transforms, J. Lond. Math. Soc. 9:188–192 (1934). [1]
- [37] E.M. Opdam, Harmonic analysis for certain representations of graded Hecke algebras, Acta. Math. 175(1):75– 121 (1995). [2, 3]
- [38] E.M. Opdam, Lecture notes on Dunkl operators for real and complex reflection groups, In: Mem. Math. Soc. Japan 8, Tokyo (2000). [2, 3]
- [39] A. Poria, Uncertainty principles for the Opdam-Cherednik transform on modulation spaces, Integral Trans. Spec. Funct. 32(3):191–206 (2021). [2, 3]
- [40] A. Poria, Localization operators associated with the windowed Opdam-Cherednik transform on modulation spaces, Complex Var. Elliptic Equ. 1–24 (2022). [2, 3, 5, 6]
- [41] B. Ricaud and B. Torrsani, A survey of uncertainty principles and some signal processing applications, Adv. Comput. Math. 40:629–650 (2014). [1, 2, 15, 16]
- [42] B. Schapira, Contributions to the hypergeometric function theory of Heckman and Opdam: sharp estimates, Schwartz space, heat kernel, Geom. Funct. Anal. 18:222–250 (2008). [2, 3]
- [43] D. Slepian and H.O. Pollak, Prolate spheroidal wave functions, Fourier analysis and uncertainty I, Bell Syst. Tech. J. 40:43–63 (1961). [1]
- [44] E. Wilczok, New uncertainty principles for the continuous Gabor transform and the continuous Wavelet transform, Doc. Math. 5:201–226 (2000). [2]