

# $L(n)$ graphs are vertex-pancyclic and Hamilton-connected

S.Morteza Mirafzal\*, Sara Kouhi

Department of Mathematics

Lorestan University, Khorramabad, Iran

E-mail: mirafzal.m@lu.ac.ir

E-mail: smortezamirafzal@yahoo.com

Sara Kouhi

Department of Mathematics

Lorestan University, Khorramabad, Iran

E-mail:sarakouhi22@gmail.com

## Abstract

A graph  $G$  of order  $n > 2$  is pancyclic if  $G$  contains a cycle of length  $l$  for each integer  $l$  with  $3 \leq l \leq n$  and it is called vertex-pancyclic if every vertex is contained in a cycle of length  $l$  for every  $3 \leq l \leq n$ . A graph  $G$  of order  $n > 2$  is Hamilton-connected if for any pair of distinct vertices  $u$  and  $v$ , there is a Hamilton  $u$ - $v$  path, namely, there is a  $u$ - $v$  path of length  $n - 1$ . The graph  $B(n)$  is a graph with the vertex set  $V = \{v \mid v \subset [n], |v| \in \{1, 2\}\}$  and the edge set  $E = \{\{v, w\} \mid v, w \in V, v \subset w \text{ or } w \subset v\}$ , where  $[n] = \{1, 2, \dots, n\}$ . We denote by  $L(n)$  the line graph of  $B(n)$ , that is,  $L(n) = L(B(n))$ . In this paper, we show that the graph  $L(n)$  is vertex-pancyclic and Hamilton-connected whenever  $n \geq 6$ .

---

2010 *Mathematics Subject Classification*: 05C38, 90B10

*Keywords*:  $L(n)$  graph, path and cycle, vertex-pancyclic, Hamilton-connected

*\*Corresponding author.*

*Date*: 26 3 2022

# 1 Introduction and Preliminaries

In this paper, a graph  $G = (V, E)$  is considered as an undirected simple graph where  $V = V(G)$  is the vertex-set and  $E = E(G)$  is the edge-set. For all the terminology and notation not defined here, we follow [3,4].

Let  $n \geq 3$  be an integer and  $[n] = \{1, 2, \dots, n\}$ . The graph  $B(n)$  is a graph with the vertex set  $V = \{v \mid v \subset [n], |v| \in \{1, 2\}\}$  and the edge set  $E = \{\{v, w\} \mid v, w \in V, v \subset w \text{ or } w \subset v\}$ . We denote by  $L(n)$  the line graph of  $B(n)$ , that is,  $L(n) = L(B(n))$ . It is easy to check that  $L(n)$  is a connected regular graph of regularity  $n - 1$ . Hence when  $n = 3$ ,  $L(n)$  is the cycle  $C_6$ . It can easily be seen that the graph  $B(n)$  is an edge-transitive graph [8], thus  $L(n)$  is a vertex-transitive graph [3,4]. The graphs  $B(n)$  and  $L(n)$  have some interesting properties and have been studied in some aspects [6,7,8,9]. In [8] it has been shown that the graph  $L(n)$  is a Cayley graph if and only if  $n$  is a power of a prime. The graph  $G$  is called an *integral* graph whenever each of the eigenvalues of its adjacency matrix is an integer. In [6] it has been proved that the graph  $L(n)$  is an integral graph. In fact the set of eigenvalues of  $L(n)$  is  $\{-2, -1, 0, n-2, n-1\}$ . An interesting property of the graph  $B(n)$  has been appeared in [7], that is,  $L(n)$  is (isomorphic to) the square root of the Johnson graph  $J(n+1, 2)$ , namely,  $B(n)^2$  is isomorphic to  $J(n+1, 2)$ . If  $v$  is a vertex of the graph  $B(n)$ , then  $\deg(v) \in \{2, n-1\}$ . Therefore, if  $n$  is an odd integer then  $B(n)$  is an *eulerian* graph [4]. Hence  $L(n)$  is a *Hamilton* graph when  $n$  is an odd integer. A graph  $G$  of order  $n > 2$  is *Hamilton-connected* if for any pair of distinct vertices  $u$  and  $v$ , there is a Hamilton  $u$ - $v$  path, namely, there is a  $u$ - $v$  path of length  $n - 1$ . Note that if the graph  $G$  is Hamilton-connected then it is hamiltonian. Also note that a hamiltonian graph may not be Hamilton-connected. For instance the cycle  $C_6$  is hamiltonian, but it is easy to see that  $C_6$  is not Hamilton-connected. A graph  $G$  of order  $n > 2$  is *panconnected* if for every two vertices  $u$  and  $v$ , there is a  $u$ - $v$  path of length  $l$  for every integer  $l$  with  $d(u, v) \leq l \leq n - 1$ . It is trivial that if  $G$  is a panconnected graph, then it is Hamilton-connected. Alspach [1] showed that the Johnson graph  $J(n, m)$  is Hamilton-connected. This result has been generalized in [7] where it has been shown that the Johnson graphs are panconnected. It is easy to check that the graph  $L(n)$  is not panconnected. In fact, it is not hard to show that for adjacent vertices  $v = [1, 12]$  and  $w = [2, 12]$ , there is not a  $v$ - $w$  path of length 3 (or 4) in the graph  $L(n)$ . In this paper, we wish to show that the graph  $L(n)$  has another interesting properties, that is, if  $n \geq 6$ , then it is vertex-pancyclic and Hamilton-connected.

The group of all permutations of a set  $V$  is denoted by  $Sym(V)$  or just  $Sym(n)$  when  $|V| = n$ . A *permutation group*  $\Gamma$  on  $V$  is a subgroup of  $Sym(V)$ . In this case we say that  $\Gamma$  acts on  $V$ . If  $G$  is a graph with vertex-set  $V$ , then we can view each automorphism of  $G$  as a permutation of  $V$ , and so  $Aut(G)$  is a permutation group. Let the group  $\Gamma$  act on  $V$ . We say that  $\Gamma$  is *transitive* (or  $\Gamma$  acts *transitively* on  $V$ ) if there is just one orbit. This means that given any two element  $u$  and  $v$  of  $V$ , there is an element  $\beta$  of  $\Gamma$  such that  $\beta(u) = v$ .

The graph  $G$  is called *vertex-transitive* if  $Aut(G)$  acts transitively on

$V(G)$ . The action of  $Aut(G)$  on  $V(G)$  induces an action on  $E(G)$ , by the rule,  $\beta\{x, y\} = \{\beta(x), \beta(y)\}$ ,  $\beta \in Aut(G)$ , and  $G$  is called *edge-transitive* if this action is transitive. The graph  $G$  is called *symmetric*, if for all vertices  $u, v, x, y$ , of  $G$  such that  $u$  and  $v$  are adjacent, and  $x$  and  $y$  are adjacent, there is an automorphism  $\alpha$  such that  $\alpha(u) = x$ , and  $\alpha(v) = y$ . It is clear that a connected symmetric graph is vertex-transitive and edge-transitive. It is easy to show that the graph  $L(n)$  is not a symmetric graph.

Let  $\Gamma$  be any abstract finite group with identity 1, and suppose  $\Omega$  is a subset of  $\Gamma$  with the properties: (i)  $x \in \Omega \implies x^{-1} \in \Omega$ , (ii)  $1 \notin \Omega$ . The *Cayley graph*  $G = G(\Gamma; \Omega)$  is the (simple) graph whose vertex-set and edge-set are defined as follows:  $V(G) = \Gamma$ ,  $E(G) = \{\{g, h\} \mid g^{-1}h \in \Omega\}$ . As we have already stated, the graph  $L(n)$  is a Cayley graph if and only if  $n$  is a power of a prime integer [8].


## 2 Main results

**Definition 2.1.** Let  $n \geq 3$  be an integer and  $[n] = \{1, 2, \dots, n\}$ . The graph  $B(n)$  is a graph with the vertex set  $V = \{v \mid v \subset [n], |v| \in \{1, 2\}\}$  and the edge set  $E = \{\{v, w\} \mid v, w \in V, v \subset w \text{ or } w \subset v\}$ . We denote by  $L(n)$  the line graph of  $B(n)$ , that is,  $L(n) = L(B(n))$ .

From Definition 2.1, it follows that every vertex of the graph  $L(n)$  is of the form  $\{\{i\}, \{i, j\}\}$ , where  $i, j \in [n]$ , and  $i \neq j$ . In this paper, we denote  $\{\{i\}, \{i, j\}\}$  by  $[i, ij]$ . Hence  $L(n)$  is the graph with the vertex-set  $V(L(n)) = V = \{[i, ij], i, j \in [n], i \neq j\}$ , in which two vertices  $[i, ij]$  and  $[r, rs]$  are adjacent if and only if  $i = r$  or  $\{i, j\} = \{r, s\}$ . Thus, if  $[i, ij]$  is a vertex of  $L(n)$ , then

$$N([i, ij]) = \{[i, ik] \mid k \in [n], k \neq j, i\} \cup \{[j, ij]\},$$

hence,  $\deg([i, ij]) = n - 1$ . Therefore  $L(n)$  is a regular graph of valency  $n - 1$ . In fact,  $L(n)$  is a vertex-transitive graph [6,8]. By an easy argument, we can show that the graph  $L(n)$  is a connected graph with diameter 3. Also, its girth is 3 and hence it is not a bipartite graph. Figure 1, shows  $L(4)$  in the plane.

Figure 1: The graph  $L(4)$ 

Amongst the various properties of the graph  $L(n)$ , we interested in its cycle structure and Hamilton-connectivity. Since for  $n = 3$  the graph  $L(n)$  is isomorphic with  $C_6$ , and the structure of this graph is simple, in the sequel we let  $n \geq 4$ . As we can see in Figure 1, there are 4 cliques of order 3 in  $L(4)$  such that they construct a partition for the vertex-set of this graph. We can easily check that by these cliques we can construct a  $u-v$  paths of length  $m$  if  $5 \leq m \leq 11$  for any two vertices  $u$  and  $v$  in the graph  $L(4)$ . Let  $u = [1, 12]$  and  $v = [2, 12]$ . It is easy to check that there is no any  $u-v$  path of length 4 (or 3) in  $L(4)$ . Hence  $L(4)$  is not a panconnected graph. In the graph  $L(n)$  the subgraph induced by the set  $C_i = \{[i, ij] \mid j \in [n], j \neq i\}$ ,  $1 \leq i \leq n$ , is a clique of order  $n - 1$ . It is clear that  $P = \{C_i \mid 1 \leq i \leq n\}$  is a partition for the vertex-set of  $L(n)$ . Note that for any two cliques  $C_i$  and  $C_j$  there is a unique pair of adjacent vertices  $(u, v)$  such that  $u \in C_i$ ,  $v \in C_j$  (indeed  $u = [i, ij]$  and  $v = [j, ij]$ ). Moreover if  $v$  is a vertex in the clique  $C_i$ , then there is exactly one vertex  $w$  in  $V(L(n)) - C_i$  such that  $w$  is adjacent to  $v$ . In fact if  $v = [i, ij] \in C_i$ , then  $w = [j, ij]$  is the unique vertex which is not in  $C_i$  ( $w \in C_j$ ) and adjacent to  $v$ .

**Lemma 2.2.** *Let  $n \geq 4$  be an integer and  $P = \{C_i \mid 1 \leq i \leq n\}$  be the clique partition of the vertex-set of the graph  $L(n)$  where  $C_i = \{[i, ij] \mid j \in [n], j \neq i\}$ . Let  $C \in P$  and let  $v, w$  be two vertices in  $C$ . Let  $t$  be an integer such that,  $2 \leq t \leq n - 1$ . Then there is a  $t$ -subset  $P_1 = \{A_1, \dots, A_t\} \subseteq P - \{C\}$ , and a  $v-w$  path  $Q : v, v_1, w_1, v_2, w_2, \dots, v_t, w_t, w$  of length  $2t + 1$  such that  $v_i, w_i \in A_i$ .*

*Proof.* We prove the lemma by induction on  $t$ . Let  $t = 2$ . Since  $L(n)$  is a vertex-transitive graph, without loss of generality, we can assume that  $v = [1, 12]$  (and hence  $C = C_1$ ),  $w = [1, 1i]$ ,  $i \neq 1, 2$ . Consider the vertices  $v_1 = [2, 12] \in C_2$  and  $w_2 = [i, 1i] \in C_i$ . Thus there is a unique adjacent pair  $(w_1, v_2)$  such that  $w_1 \in C_2$  and  $v_2 \in C_i$  ( $w_1 = [2, 2i]$ ,  $v_2 = [i, 2i]$ ). Now,

it is clear that the path  $Q : v, v_1, w_1, v_2, w_2, w$  is a desired path of length  $5 = 2t + 1$ . Now let  $2 \leq m < t$  and the claim is true for  $m$ . By induction hypothesis there are cliques  $A_1, A_2, \dots, A_{t-1} \subset P - C_1$  and a  $v$ - $w$  path  $Q : v, v_1, w_1, v_2, w_2, \dots, v_{t-1}, w_{t-1}, w$  of length  $2t - 1$  such that  $v_i, w_i \in A_i$ . Let  $A_t \in P - \{C_1, A_1, A_2, \dots, A_{t-1}\}$  be an arbitrary clique. There is a unique adjacent pair  $(u_{t-2}, z_{t-1})$  such that  $u_{t-2} \in A_{t-2}$  and  $z_{t-1} \in A_t$ . Note that  $u_{t-2} \neq v_{t-2}, w_{t-2}$ . Moreover there is a unique adjacent pair  $(u_{t-1}, z_t)$  such that  $u_{t-1} \in A_t$  and  $z_t \in A_{t-1}$ . Again Note that  $z_t \neq v_{t-1}, w_{t-1}$ . Now it is clear that the  $v$ - $w$  path  $Q_0 : v, v_1, w_1, v_2, w_2, \dots, v_{t-2}, u_{t-2}, z_{t-1}, u_{t-1}, z_t, w_{t-1}, w$  of length  $2t + 1$  is a desired path if we rename  $A_{t-1}$  by  $A_t$  and  $A_t$  by  $A_{t-1}$ .  $\square$

A graph  $G$  of order  $n$  is  $k$ -pancyclic ( $k \leq n$ ) if it contains cycles of every length from  $k$  to  $n$  inclusive, and  $G$  is pancyclic if it is  $g$ -pancyclic, where  $g = g(G)$  is the girth of  $G$ . A graph is of pancyclicity if it is pancyclic. Note that if  $G$  is pancyclic, then  $G$  is hamiltonian. The pancyclicity is an important property to determine if a topology of a network is suitable for some applications where mapping cycles of any length into the topology of the network is required [10,11]. The concept of pancyclicity, proposed first by Bondy [5], has been extended to vertex-pancyclicity and edge-pancyclicity [1]. A graph  $G$  of order  $n$  is vertex-pancyclic (resp. edge-pancyclic) if any vertex (resp. edge) lies on cycles of every length from  $g(G)$  to  $n$  inclusive. Obviously, an edge pancyclic graph is certainly vertex-pancyclic.

It is easy to check that  $L(4)$  has no cycles of lengths 4 and 5. Also  $L(5)$  has no cycle of length 5. But by lemma 2.2, we can show that if  $n \geq 6$ , then  $L(n)$  is a vertex-pancyclic graph. Moreover, we can check that  $L(n)$  is not edge-pancyclic, since the edge  $e = \{[1, 12], [2, 12]\}$  does not lie on a 3-cycle (or 4-cycle). In fact we have the following result.

**Theorem 2.3.** *Let  $n \geq 6$  be an integer. Then  $L(n)$  is a vertex-pancyclic graph.*

*Proof.* Let  $m$  be an integer such that  $n(n-1) \geq m \geq 3$ . Let  $v$  be an arbitrary vertex in the graph  $L(n)$ . Let  $C$  be the  $(n-1)$ -clique containing  $v$  in  $L(n)$ . Hence if  $3 \leq m \leq 5$ , then there is a cycle  $C_m$  containing  $v$  in  $C$ . Now assume that  $6 \leq m \leq 3(n-1)$ . Let  $w \in C$  be such that  $w \neq v$ . By Lemma 2.2, there are two  $(n-1)$ -cliques  $A_1$  and  $A_2$  and a path  $Q : v, v_1, w_1, v_2, w_2, w$  of length 5 such that  $v_i, w_i \in A_i$  for each  $i \in \{1, 2\}$ . Let  $1 \leq t \leq n-3$ . It is clear that we can insert  $t$  vertices of the clique  $C$  between  $v$  and  $w$ . Also, we can insert  $t$  vertices of the clique  $A_i$ ,  $i \in \{1, 2\}$  between  $v_i$  and  $w_i$ . Now it is clear how we can construct an  $m$ -cycle containing the vertex  $v$  in the graph  $L(n)$ . Now let  $m > 3(n-1)$ . By Lemma 2.2, there are cliques  $A_1, \dots, A_{(n-1)}$  and the path  $Q : v, v_1, w_1, v_2, w_2, \dots, v_{(n-1)}, w_{(n-1)}, w$  in  $L(n)$  such that  $v_i, w_i \in A_i$ ,  $1 \leq i \leq n-1$ . Note that  $3(n-1) > 2n$ . By inserting adequate number of vertices of each  $A_i$  (and  $C$ ) between each pair of vertices  $v_i$  and  $w_i$  ( $v$  and  $w$ ), we can construct an  $m$ -cycle containing the vertex  $v$  in the graph  $L(n)$ .  $\square$

We now want to show that the graph  $L(n)$  is a Hamilton-connected graph.

**Theorem 2.4.** *Let  $n \geq 4$  be an integer. Then  $L(n)$  is a Hamilton-connected graph.*

*Proof.* It is easy to check that the assertion of the theorem is true for the case  $n = 4$ , hence in the sequel we assume that  $n \geq 5$ . Let  $v, w$  be two vertices in  $L(n)$ . We show that there is a hamiltotian  $v$ - $w$  path in  $L(n)$ . Let  $P = \{C_i \mid 1 \leq i \leq n\}$  be the clique partition of the vertex-set of the graph  $L(n)$  where  $C_i = \{[i, ij] \mid j \in [n], j \neq i\}$ . There are two cases, namely,

- (i)  $v$  and  $w$  are in the same  $(n-1)$ -clique in the graph  $L(n)$ , or
  - (ii)  $v$  and  $w$  are in distinct  $(n-1)$ -cliques.
- (i) Assume that  $A_1 \in P$  and  $v, w \in A_1$ . Let  $w_1 \in A_1$  be such that  $w_1 \neq v, w$ . Therefore by Lemma 2.2, for subset  $P_1 = \{A_2, \dots, A_n\} \subseteq P - \{A_1\}$ , there is a  $v$ - $w_1$  path  $Q_0 : v = v_1, v_2, w_2, \dots, v_n, w_n, w_1$  of length  $2n-1$  such that  $v_i, w_i \in A_i$ . Hence  $Q_1 : v = v_1, v_2, w_2, \dots, v_n, w_n, w_1, w$  is a  $v$ - $w$  path of length  $2n$  in  $L(n)$ . For each  $i, 2 \leq i \leq n$  we insert all other vertices in the clique  $A_i$  between  $v_i$  and  $w_i$  and again obtain a  $v$ - $w$  path of greater length. Now by inserting all the vertices in the set  $A_1 - \{v, w, w_1\}$  between the vertices  $w_1$  and  $w$ , we obtain a hamiltonian  $v$ - $w$  path in the graph  $L(n)$ .
- (ii) We now assume that  $v$  and  $w$  are in distinct  $(n-1)$ -cliques. Let  $A_1, A_n \in P$  are such that  $v \in A_1$  and  $w \in A_n$ . We know that for each vertex  $x$  in  $L(n)$  there is a unique  $(n-1)$ -clique  $C_x$  in the graph  $L(n)$  such that  $x \notin C_x$  but  $x$  is adjacent to a unique vertex in  $C_x$ . Let  $P - \{A_1, A_n\} = \{A_2, \dots, A_{n-1}\}$  be such that  $A_2 \neq C_v$  and  $A_{n-1} \neq C_w$ . Therefore  $v$  is adjacent to no vertex in  $A_2$  and  $w$  is adjacent to no vertex in  $A_{n-1}$ . On the other hand, for each  $i, 1 \leq i \leq n-1$  there is a unique adjacent pair  $w_i, v_{i+1}$  such that  $w_i \in A_i$  and  $v_{i+1} \in A_{i+1}$ . Note that  $w_1 \neq v$  and  $v_n \neq w$ . Hence the path  $Q_0 : v = v_1, w_1, v_2, w_2, \dots, v_n, w_n = w$  is a  $v$ - $w$  path in the graph  $L(n)$ . For each  $i, 1 \leq i \leq n$ , if we insert between  $v_i$  and  $w_i$  all other vertices in the clique  $A_i$ , we obtain a hamiltonian  $v$ - $w$  path in the graph  $L(n)$ . □

## References

- [1] Alspach B, Hare D, Edge-pancyclic block-intersection graphs. Discrete Math, 1991, 97(1-3): 17-24.
- [2] Alspach B, Johnson graphs are Hamilton-connected. Ars Math. Contemp 6 (2013), 21-23.
- [3] Biggs N.L, Algebraic Graph Theory 1993 (Second edition), Cambridge Mathematical Library (Cambridge University Press; Cambridge).
- [4] Bondy J.A, Murty U. S. R, Graph Theory, Springer, Berlin, (2008).
- [5] Bondy J A, Pancyclic graphs. I. J. Combin. Theory 11, 1971, 80-84.

- [6] Mirafzal S.M, A new class of integral graphs constructed from the hypercube, *Linear Algebra Appl.* 2018;558:186–194.
- [7] Mirafzal S.M, Heidari A, Johnson graphs are panconnected, *Proceedings-Mathematical Sciences*, (2019), <https://doi.org/10.1007/s12044-019-0527-3>.
- [8] Mirafzal S.M, Cayley properties of the line graphs induced by consecutive layers of the hypercube, *Bull. Malaysian Math. Sci.* (2020), <https://doi.org/10.1007/s40840-020-01009-3>.
- [9] Mirafzal S.M, On the automorphism groups of connected bipartite irreducible graphs. *Proc. Math. Sci.* (2020). <https://doi.org/10.1007/s12044-020-0589-1>.
- [10] Xu J.-M, *Topological Structure and Analysis of Interconnection Networks*, Kluwer Academic Publishers., Dordrecht/Boston/London, 2001.
- [11] Xu J.-M, Ma M, Survey on path and cycle embedding in some networks, *Front. Math. China* 4 (2) (2009) 217–252.