

Locally octahedral and locally almost square Köthe-Bochner spaces

Jan-David Hardtke

ABSTRACT. It has been proved in [19] that a Köthe-Bochner space $E(X)$ is locally octahedral/locally almost square if X has the respective property and the simple functions are dense in $E(X)$. Here we show that the result still holds true without the density assumption. The proof makes use of the Kuratowski-Ryll-Nardzewski Theorem on measurable selections.

1 Introduction

Let X be a real Banach space. We denote by X^* its topological dual, by B_X its closed unit ball and by S_X its unit sphere.

X is locally octahedral (LOH) if the following holds: for every $x \in S_X$ and every $\varepsilon > 0$ there exists $y \in S_X$ such that $\|x \pm y\| \geq 2 - \varepsilon$. This notion was introduced in [15] in connection with the so called diameter-two-properties.

X is locally almost square (LASQ) if the following holds: for every $x \in S_X$ and every $\varepsilon > 0$ there exists $y \in S_X$ such that $\|x \pm y\| \leq 1 + \varepsilon$. This notion was introduced in [2].

For more information on these and related properties the reader may consult [1–9, 11, 13–22, 24], [10, Theorem 2.5, p.106] and references therein.

Now consider a complete, σ -finite measure space (S, \mathcal{A}, μ) . For a set $A \subseteq S$ we denote by χ_A the characteristic function of A . Let $(E, \|\cdot\|_E)$ be a Banach space of real-valued measurable functions on S (modulo equality μ -almost everywhere) such that the following holds:

- (i) $\chi_A \in E$ for every set $A \in \mathcal{A}$ with $\mu(A) < \infty$.
- (ii) If $f \in E$ and $A \in \mathcal{A}$ with $\mu(A) < \infty$, then f is μ -integrable over A .
- (iii) If g is measurable and $f \in E$ such that $|g(t)| \leq |f(t)|$ μ -a. e. then $g \in E$ and $\|g\|_E \leq \|f\|_E$.

Then $(E, \|\cdot\|_E)$ is called a Köthe function space over (S, \mathcal{A}, μ) .

Keywords: Köthe-Bochner spaces; locally octahedral spaces; locally almost square spaces; measurable selections

AMS Subject Classification (2020): 46B20 46E30 46E40 28B20

Standard examples are the spaces $L^p(\mu)$ for $1 \leq p \leq \infty$. More generally, Orlicz spaces with the Luxemburg norm (see [25]) are examples of Köthe function spaces.

A function $f : S \rightarrow X$ is called simple if there are finitely many pairwise disjoint sets $A_1, \dots, A_n \in \mathcal{A}$ such that $\mu(A_i) < \infty$ for all $i = 1, \dots, n$, f is constant on each A_i and $f(t) = 0$ for every $t \in S \setminus \bigcup_{i=1}^n A_i$.

f is said to be Bochner-measurable if there exists a sequence $(f_n)_{n \in \mathbb{N}}$ of simple functions such that $\lim_{n \rightarrow \infty} \|f_n(t) - f(t)\| = 0$ μ -a. e.

According to the well-known Pettis Measurability Theorem, the following assertions are equivalent:

- (i) f is Bochner-measurable.
- (ii) $x^* \circ f$ is measurable for every $x^* \in X^*$ and there is a separable, closed subspace Y of X such that $f(s) \in Y$ for μ -a. e. $s \in S$.
- (iii) f is measurable (with respect to \mathcal{A} and the Borel- σ -algebra of X) and there is a separable, closed subspace Y of X such that $f(s) \in Y$ for μ -a. e. $s \in S$.

We denote by $E(X)$ the space of all Bochner-measurable functions $f : S \rightarrow X$ (modulo equality a. e.) such that $\|f(\cdot)\| \in E$.

We define $\|f\|_{E(X)} = \|\|f(\cdot)\|\|_E$ for $f \in E(X)$. Then $E(X)$ becomes a Banach space, the so called Köthe-Bochner space induced by E and X .

For $E = L^p(\mu)$ we obtain the usual Lebesgue-Bochner spaces $L^p(\mu, X)$ for $1 \leq p \leq \infty$. For more information on Köthe-Bochner spaces the reader is referred to the book [23].

In [19, Theorems 4.1 and 4.5] the author proved the following results:

- (a) If X is LOH/LASQ and the simple functions are dense in $E(X)$, then $E(X)$ is also LOH/LASQ.
- (b) If X is LOH/LASQ, then $L^\infty(\mu, X)$ is also LOH/LASQ.

The proofs in [19] are based on a general reduction theorem and corresponding results for absolute sums of LOH/LASQ spaces that were obtained in [2, Proposition 5.3].

The assumption that the simple functions are dense in $E(X)$ holds true whenever E is order continuous, in particular for $E = L^p(\mu)$ with $1 \leq p < \infty$.

Here we will show that the result is still true without any additional assumptions on E or X . The proof makes use of the Kuratowski-Ryll-Nardzewski Theorem on the existence of measurable selections, which we will recall in the next section.

2 Measurable selections

Let (S, \mathcal{A}) be a measurable space and (Y, d) a metric space. Denote by $\mathcal{P}(Y)$ the power-set of Y and by $\mathcal{B}(Y)$ the Borel- σ -Algebra of (Y, d) . For a subset $M \subseteq Y$ we denote by \overline{M} the closure of M in (Y, d) .

Let $F : S \rightarrow \mathcal{P}(Y)$ be a set-valued map. For $M \subseteq Y$ we put

$$F_-(M) := \{s \in S : F(s) \cap M \neq \emptyset\}.$$

F is said to be weakly \mathcal{A} -measurable if $F_-(U) \in \mathcal{A}$ for every open set $U \subseteq Y$. If $F_-(C) \in \mathcal{A}$ for every closed set $C \subseteq Y$, then F is called \mathcal{A} -measurable. \mathcal{A} -measurability of F implies weak \mathcal{A} -measurability (this follows from the fact that every open subset in a metric space is an F_σ -set).

The following standard lemma follows directly from the definition.

Lemma 2.1. *Suppose that $F, G : S \rightarrow \mathcal{P}(Y)$ are two set-valued maps such that $F(s) \subseteq G(s) \subseteq \overline{F(s)}$ for every $s \in S$. Then F is weakly \mathcal{A} -measurable if and only if G is weakly \mathcal{A} -measurable.*

The next lemma is also standard but we include a sketch of the proof here for the reader's convenience.

Lemma 2.2. *Suppose that (Y, d) is separable and $g : S \times Y \rightarrow \mathbb{R}$ is a Carathéodory function, i. e.*

- (i) $g(s, \cdot)$ is continuous for every $s \in S$,
- (ii) $g(\cdot, y)$ is \mathcal{A} -measurable for every $y \in Y$.

Let $\alpha \in \mathbb{R}$ and put

$$F(s) := \{y \in Y : g(s, y) > \alpha\} \quad \text{for all } s \in S.$$

Then $F : S \rightarrow \mathcal{P}(Y)$ is \mathcal{A} -measurable.

Proof. Let $C \subseteq Y$ be nonempty and closed. Choose a sequence $(y_n)_{n \in \mathbb{N}}$ such that $C = \overline{\{y_n : n \in \mathbb{N}\}}$ and put $A_n := \{s \in S : g(s, y_n) > \alpha\}$ for every $n \in \mathbb{N}$. Then we have $A_n \in \mathcal{A}$ for each n and it is easy to see that $F_-(C) = \bigcup_{n \in \mathbb{N}} A_n$. Thus $F_-(C) \in \mathcal{A}$. \square

A classical result on the existence of measurable selections is the Kuratowski-Ryll-Nardzewski Theorem (see for instance [12, Theorem 2.1]).

Theorem 2.3 (Kuratowski-Ryll-Nardzewski Selection Theorem). *Let (S, \mathcal{A}) be a measurable space and (Y, d) a complete, separable metric space. Let $F : S \rightarrow \mathcal{P}(Y)$ be a weakly \mathcal{A} -measurable set-valued map such that $F(s)$ is non-empty and closed in Y for every $s \in S$. Then there is an \mathcal{A} - $\mathcal{B}(Y)$ -measurable map $f : S \rightarrow Y$ such that $f(s) \in F(s)$ for every $s \in S$.*

3 LOH and LASQ Köthe-Bochner spaces

Now we are ready to prove the general stability result.

Theorem 3.1. *Let (S, \mathcal{A}, μ) be a complete, σ -finite measure space and let E be a Köthe function space over (S, \mathcal{A}, μ) . If X is a real Banach space which is LOH/LASQ, then the Köthe-Bochner space $E(X)$ is also LOH/LASQ.*

Proof. 1) Assume that X is LOH. Fix $\varepsilon > 0$ and $f \in E(X)$ with $\|f\|_{E(X)} = 1$. Since $f : S \rightarrow X$ is Bochner-measurable there is a separable, closed subspace Y of X such that $f(s) \in Y$ for μ -a. e. $s \in S$ (Pettis Measurability Theorem). Without loss of generality we may assume that this holds even for all $s \in S$.

Put $S' := \{s \in S : f(s) \neq 0\}$.

Since Y is separable the unit sphere S_Y is also separable. We fix a sequence $(y_n)_{n \in \mathbb{N}}$ which is dense in S_Y .

Because X is LOH we can find a sequence $(z_n)_{n \in \mathbb{N}}$ in S_X such that $\|y_n \pm z_n\| \geq 2 - \varepsilon/2$ for every $n \in \mathbb{N}$. We put $Z := \overline{\text{span}}\{z_n : n \in \mathbb{N}\}$. This is again a separable, closed subspace of X .

Next we define $F : S' \rightarrow \mathcal{P}(S_Z)$ by

$$F(s) := \overline{\left\{ z \in S_Z : \left\| \frac{f(s)}{\|f(s)\|} \pm z \right\| > 2 - \varepsilon \right\}} \quad \text{for all } s \in S'.$$

We have $F(s) \neq \emptyset$ for every $s \in S'$. To see this note that $f(s)/\|f(s)\| \in S_Y$ and hence there is some index $N \in \mathbb{N}$ such that $\|f(s)/\|f(s)\| - y_N\| < \varepsilon/2$. It follows that

$$\left\| \frac{f(s)}{\|f(s)\|} \pm z_N \right\| \geq \|y_N \pm z_N\| - \left\| \frac{f(s)}{\|f(s)\|} - y_N \right\| > 2 - \frac{\varepsilon}{2} - \frac{\varepsilon}{2} = 2 - \varepsilon.$$

If we define

$$g(s, z) := \min \left\{ \left\| \frac{f(s)}{\|f(s)\|} + z \right\|, \left\| \frac{f(s)}{\|f(s)\|} - z \right\| \right\} \quad \text{for } s \in S', z \in S_Z,$$

then $F(s) = \overline{\{z \in S_Z : g(s, z) > 2 - \varepsilon\}}$ and since g is a Carathéodory function, it follows from Lemma 2.2 and Lemma 2.1 that F is weakly measurable.

Thus by the Kuratowski-Ryll-Nardzewski Selection Theorem there exists a measurable function $\tilde{f} : S' \rightarrow S_Z$ such that $\tilde{f}(s) \in F(s)$ for every $s \in S'$. Note that by Pettis Measurability Theorem \tilde{f} is also Bochner-measurable. We put $h(s) := \|f(s)\|\tilde{f}(s)$ for $s \in S'$ and $h(s) := 0$ for $s \in S \setminus S'$. Then h is Bochner-measurable and $\|h(s)\| = \|f(s)\|$ for every $s \in S$ and thus $\|h\|_{E(X)} = \|f\|_{E(X)} = 1$.

We further have $\|f(s) \pm h(s)\| \geq (2 - \varepsilon)\|f(s)\|$ for every $s \in S$. This implies $\|f \pm h\|_{E(X)} \geq (2 - \varepsilon)\|f\|_{E(X)} = 2 - \varepsilon$.

This proves that $E(X)$ is LOH.

2) Now assume that X is LASQ. We take ε , f , Y , S' and $(y_n)_{n \in \mathbb{N}}$ as in part 1). Since X is LASQ there is a sequence $(z_n)_{n \in \mathbb{N}}$ in S_X such that $\|y_n \pm z_n\| \leq 1 + \varepsilon/2$ for every $n \in \mathbb{N}$ and we put $Z := \overline{\text{span}\{z_n : n \in \mathbb{N}\}}$ and

$$F(s) := \overline{\left\{ z \in S_Z : \left\| \frac{f(s)}{\|f(s)\|} \pm z \right\| < 1 + \varepsilon \right\}} \quad \text{for all } s \in S'.$$

Analogously to the proof in part 1) we can see that each set $F(s)$ is non-empty and also that F is weakly measurable. For the latter, put

$$g(s, z) := \max \left\{ \left\| \frac{f(s)}{\|f(s)\|} + z \right\|, \left\| \frac{f(s)}{\|f(s)\|} - z \right\| \right\} \quad \text{for } s \in S', z \in S_Z.$$

Then g is a Carathéodory function and $F(s) = \overline{\{z \in S_Z : g(s, z) < 1 + \varepsilon\}}$.

By the Kuratowski-Ryll-Nardzewski Selection Theorem we find a measurable function $\tilde{f} : S' \rightarrow S_Z$ such that $\tilde{f}(s) \in F(s)$ for every $s \in S'$. Then if we define h as in 1), we find that $h \in S_{E(X)}$ and $\|f(s) \pm h(s)\| \leq \|f(s)\|(1 + \varepsilon)$ for every $s \in S$. This implies $\|f \pm h\|_{E(X)} \leq 1 + \varepsilon$ and the proof is finished. \square

References

- [1] T. Abrahamsen, V. Lima, and O. Nygaard, *Remarks on diameter 2 properties*, J. Conv. Anal. **20** (2013), 439–452.
- [2] T. A. Abrahamsen, J. Langemets, and V. Lima, *Almost square Banach spaces*, J. Math. Anal. Appl. **434** (2016), no. 2, 1549–1565.
- [3] T. A. Abrahamsen, P. Hájek, and S. Troyanski, *Almost square dual Banach spaces*, J. Math. Anal. Appl. **487** (2020), no. 2, 11 p.
- [4] M. D. Acosta, J. Becerra Guerrero, and G. López-Pérez, *Stability results of diameter two properties*, J. Conv. Anal. **22** (2015), no. 1, 1–17.
- [5] J. Becerra Guerrero and G. López-Pérez, *Relatively weakly open subsets of the unit ball in function spaces*, J. Math. Anal. Appl. **315** (2006), 544–554.
- [6] J. Becerra Guerrero, G. López-Pérez, and A. Rueda Zoca, *Octahedral norms and convex combination of slices in Banach spaces*, J. Funct. Anal. **266** (2014), no. 4, 2424–2435.
- [7] ———, *Big slices versus big relatively weakly open subsets in Banach spaces*, J. Math. Anal. Appl. **428** (2015), 855–865.
- [8] ———, *Octahedral norms in spaces of operators*, J. Math. Anal. Appl. **427** (2015), 171–184.
- [9] ———, *Some results on almost square Banach spaces*, J. Math. Anal. Appl. **438** (2016), no. 2, 1030–1040.
- [10] R. Deville, G. Godefroy, and V. Zizler, *Smoothness and renormings in Banach spaces*, Pitman Monographs and Surveys in Pure and Applied Mathematics, vol. 64, Longman Scientific & Technical, Harlow, 1993.
- [11] G. Godefroy, *Metric characterization of first Baire class linear forms and octahedral norms*, Studia Math. **95** (1989), no. 1, 1–15.

- [12] S. Graf, *Selected results on measurable selections*, Proceedings of the 10th Conference on Abstract Analysis (Z. Frolík, ed.), Circolo Matematico di Palermo, Palermo, 1982, pp. 87–122.
- [13] R. Haller and J. Langemets, *Two remarks on diameter 2 properties*, Proc. Estonian Acad. Sci. **63** (2014), no. 1, 2–7.
- [14] ———, *Geometry of Banach spaces with an octahedral norm*, Acta Comment. Univ. Tartuensis Math. **18** (2014), no. 1, 125–133.
- [15] R. Haller, J. Langemets, and M. Põldvere, *On duality of diameter 2 properties*, J. Conv. Anal. **22** (2015), no. 2, 465–483.
- [16] ———, *Rough norms in spaces of operators*, Math. Nachr. (2017), 11p. doi:10.1002/mana.201600409.
- [17] R. Haller, J. Langemets, and R. Nadel, *Stability of average roughness, octahedrality, and strong diameter two properties of Banach spaces with respect to absolute sums*, Banach J. Math. Anal. **12** (2018), 222–239.
- [18] J.-D. Hardtke, *Summands in locally almost square and locally octahedral spaces*, Acta Comment. Univ. Tartu. Math. **22** (2018), 149–162.
- [19] ———, *On certain geometric properties in Banach spaces of vector-valued functions*, J. Math. Phys. Anal. Geom. **16** (2020), no. 2, 119–137.
- [20] D. Kubiak, *Some geometric properties of the Cesàro function spaces*, J. Convex Anal. **21** (2014), no. 1, 189–200.
- [21] J. Langemets, V. Lima, and A. Rueda Zoca, *Almost square and octahedral norms in tensor products of Banach spaces*, Rev. Real Acad. Cienc. Exact. Fisc. Nat. Serie A Mat. **111** (2017), 841–853.
- [22] ———, *Octahedral norms in tensor products of Banach spaces*, Quart. J. Math. **68** (2017), no. 4, 1247–1260.
- [23] P. K. Lin, *Köthe-Bochner function spaces*, Birkhäuser, Boston-Basel-Berlin, 2004.
- [24] G. López-Pérez, *The big slice phenomena in M -embedded and L -embedded spaces*, Proc. Amer. Math. Soc. **134** (2005), 273–282.
- [25] M. M. Rao and Z. D. Ren, *Theory of Orlicz spaces*, Monographs and Textbooks in Pure and Applied Mathematics, vol. 146, Marcel Dekker, Inc., New York, 1991.

DEPARTMENT OF MATHEMATICS
UNIVERSITÄT LEIPZIG
AUGUSTUSPLATZ 10, 04109 LEIPZIG
GERMANY

E-mail address: hardtke@math.uni-leipzig.de