

Equilibrium modeling and solution approaches inspired by nonconvex bilevel programming

Stuart Harwood^{*1}, Francisco Trespalacios², Dimitri Papageorgiou¹, and Kevin Furman²

¹ExxonMobil Corporate Strategic Research, Annandale, NJ 08801, USA

²ExxonMobil Upstream Research Company, Spring, TX 77389, USA

November 7, 2021

Abstract

This paper introduces the concept of *optimization equilibrium* as an equivalently versatile definition of a generalized Nash equilibrium for multi-agent non-cooperative games. Through this modified definition of equilibrium, we draw precise connections between generalized Nash equilibria, feasibility for bilevel programming, the Nikaido-Isoda function, and classic arguments involving Lagrangian duality and social welfare maximization. Significantly, this is all in a general setting without the assumption of convexity. Along the way, we introduce the idea of *minimum disequilibrium* as a solution concept that reduces to traditional equilibrium when equilibrium exists. The connections with bilevel programming and related semi-infinite programming permit us to adapt global optimization methods for those classes of problems, such as constraint generation or cutting plane methods, to the problem of finding a minimum disequilibrium solution. We show that this method works, both theoretically and with a numerical example, even when the agents are modeled by mixed-integer programs.

1 Introduction

The concept of equilibrium appears in many fields. In the context of game theory, Nash's celebrated results [26, 27] and further generalizations [7] regarding equilibrium between non-cooperative agents or players put equilibrium on firm mathematical foundations and have inspired further research in the ensuing decades. See [8] for a review of the history, applications, theory, and numerical methods for generalized Nash equilibrium. While that review lays out many different numerical approaches for finding equilibria, it does highlight two main classes of methods: methods involving KKT conditions, complementarity problems, or variational inequalities; and methods involving an optimization reformulation of the equilibrium problem, usually involving the Nikaido-Isoda function.

When the agents are modeled by convex optimization problems, the methods involving KKT conditions and complementarity problems are natural approaches. Similarly, as noted in [8], the use of an optimization reformulation involving the Nikaido-Isoda function has seen success, but once again assuming a certain amount of convexity; see [21, 23, 34, 35]. With the assumption of convexity, strong claims about the existence of equilibria can be made.

^{*}Corresponding author: stuart.m.harwood@exxonmobil.com

However, when the agents do not have convex structure, modeling with complementarity problems presents significant challenges, among them a certain inflexibility to define appropriate generalized solution concepts and numerical approaches. For instance, the approaches in [12, 13] combine integer decisions with complementarity models to arrive at a mixed-integer complementarity problem. The solution methods then relax certain integrality and complementarity conditions so that they obtain a mixed-integer linear program. While this is a clear approach to a numerical implementation, the solution that one arrives at is not guaranteed to be a Nash equilibrium, if one even exists. In contrast, [10] introduces and motivates “minimum total opportunity cost” as a solution concept for unit commitment problems in electricity markets with nonconvexities arising from integer decisions. However, the structure of the resulting problem led the authors to propose a heuristic solution method, and ultimately show that the “minimum complementarity” approach of [12] is an approximate solution for their minimum total opportunity cost idea.

Approaches that use the Nikaido-Isoda (NI) or some other “gap” function typically reformulate the equilibrium problem as an optimization or fixed-point problem [20, 21, 23, 34, 35, 36]. While the basic reformulations are often applicable even when there is a certain amount of nonconvexity [34, 21], the numerical methods are limited to the setting where the agent problems are convex, or at least have convex constraint sets. For the reformulations as optimization problems, this is in part because an equilibrium is a global minimizer of an implicitly defined and in general nonconvex function; finding such a point is a challenge.

One of the ultimate goals of this paper is to describe a solution method for equilibrium problems that applies even when the agents are nonconvex or even integer-constrained. The method finds an equilibrium when one exists, and if an equilibrium does not exist, the method will confirm this nonexistence while the solution obtained still has a well-defined meaning. For instance, in application to unit commitment problems, our method finds the minimum total opportunity cost solution from [10]. Central to the development of this method are theoretical and numerical approaches related to bilevel programs and semi-infinite programs [15, 32]. Through this perspective, we draw precise connections between generalized Nash equilibria, feasibility for bilevel programming, the NI function, and classic arguments involving Lagrangian duality and social welfare maximization [29]. Once again, this is all in a general setting without the assumption of convexity.

Critical to these developments is a modified definition of equilibrium which we call *optimization equilibrium*; see Definition 1. While we will show that it is equivalent to generalized Nash equilibrium (Section 2.3), optimization equilibrium has at least two benefits: one regards the simplicity of modeling certain situations; another regards the clarity of the connections to other math programs, which ultimately provide the insight into our reformulation and solution techniques.

First, optimization equilibrium introduces “global” constraints that no agent explicitly sees. This is a natural feature to allow when modeling certain equilibrium phenomena and makes the assumptions of the model clearer. For instance, Arrow and Debreu’s work in [1] defines a competitive equilibrium of an economy in terms of a set of conditions, some of which involve the optimality of certain quantities for various players. They introduce what they admit is a fictitious player, termed the “market participant” which is modeled as an optimization problem, deciding prices to maximize a payoff. This enables them to show that a Nash equilibrium exists for a related problem, which is equivalent to their definition of competitive equilibrium for an economy. While we will not try to repeat or generalize their results, we point out that the definition that they started with, competitive equilibrium, naturally includes conditions (related to supply and demand balance), that no player or agent considers, and yet are fundamental economic concepts that must be included in a sensible definition of “equilibrium.” Permitting such conditions to be included directly in a modeling framework motivates the definition of optimization equilibrium.

Second, the format of optimization equilibrium naturally leads to our consideration of bilevel

programming. Optimization equilibrium introduces variables that no agent explicitly controls. This suggests a leader-follower or outer-inner structure, akin to Stackelberg games or bilevel programs, and in fact we will discuss in Section 3 that an optimization equilibrium is equivalent to a feasible point of a certain bilevel program. This connection is not exactly new; see, e.g., the basic exposition from a complementarity viewpoint in Sections 2.7 and 2.8 of [11], which mentions mathematical programs with equilibrium constraints as a formalism for hierarchical optimization, and as a generalization of equilibrium problems. While Stackelberg/bilevel problems with multiple followers have been considered in [6, 18, 22, 37], the idea that the followers arrive at a Nash equilibrium is typically not considered, or at least, not by that terminology. In a more specific setting of controller design, multiple followers arriving at a Nash equilibrium is considered in [3]. Further, the specialization of bilevel programming and related semi-infinite programming algorithms to the (presumably simpler) class of equilibrium problems has not been explored in the literature very widely, especially in the case of nonconvex problems.

The present work has some connections with the work in [9, 10]. Specifically, our notion of *minimum disequilibrium* (Section 3) agrees with the minimum total opportunity cost idea in [10]. Once again, we arrive at this formulation through purely mathematical considerations for handling a bilevel feasibility problem. The specialization of the minimum disequilibrium problem to the economic situation of price-taking agents reveals connections to social welfare maximization. In addition, we show that optimal solutions of a certain dualized version of the max social welfare problem yield prices corresponding to a minimum disequilibrium solution, even in the nonconvex case (Section 4).

Finally, we leverage the clear connections with semi-infinite programming and propose our solution method for finding a minimum disequilibrium solution under very general conditions, for instance, even when an agent may be modeled by a mixed-integer quadratic program (Section 5). Since a minimum disequilibrium solution coincides with a Nash equilibrium when the latter exists, our method will find an equilibrium if one exists. We provide some examples to illustrate the application of this solution method, including an example from the modeling of electricity markets (Section 6).

2 Definitions and equivalence of equilibria

In this section we define optimization equilibrium and generalized Nash equilibrium, and show their equivalence. We also establish some notation that will be used throughout the work.

2.1 Optimization equilibrium definition

Consider the (nonempty) set $F \subset \mathbb{R}^{n_x} \times \mathbb{R}^{n_1} \times \dots \times \mathbb{R}^{n_m}$, and collection of m agent optimization problems parametric in $x \in \mathbb{R}^{n_x}$

$$\begin{aligned} S_i(x) &= \arg \min_{z_i} g_i(x, z_i) \\ \text{s.t. } z_i &\in \widehat{Y}_i(x), \end{aligned} \tag{A_i}$$

where $g_i : \mathbb{R}^{n_x} \times \mathbb{R}^{n_i} \rightarrow \mathbb{R}$ and \widehat{Y}_i is a set-valued mapping into \mathbb{R}^{n_i} . We will typically assume that \widehat{Y}_i takes the form

$$\widehat{Y}_i(x) = \{z_i \in Y_i : h_i(x, z_i) \leq 0\}$$

for nonempty $Y_i \subset \mathbb{R}^{n_i}$, $h_i : \mathbb{R}^{n_x} \times \mathbb{R}^{n_i} \rightarrow \mathbb{R}^{p_i}$, and inequalities between vectors holding componentwise. Also important is the optimal value function of problem (\mathcal{A}_i)

$$g_i^*(x) = \inf \{g_i(x, z_i) : z_i \in Y_i, h_i(x, z_i) \leq 0\}.$$

As usual, for any i , define $g_i^*(x) = +\infty$ if optimization problem (\mathcal{A}_i) is infeasible, and $g_i^*(x) = -\infty$ if the problem is unbounded. This formulation for the agent problems is quite general, and in particular it allows that Y_i might enforce certain decision variables to be integer-valued. We have the following definition.

Definition 1 (Optimization equilibrium). A point (x, y_1, \dots, y_m) is an optimization equilibrium of the collection of (\mathcal{A}_i) with respect to F if $(x, y_1, \dots, y_m) \in F$ and y_i solves (\mathcal{A}_i) for each $i \in \{1, \dots, m\}$ (that is, $y_i \in S_i(x)$).

We may simply say that (x, y_1, \dots, y_m) is an optimization equilibrium if the set F and parametric problems are clear from context. Further, as a brief note on notation, we will sometimes write

$$\begin{aligned} (y_1, \dots, y_m) &= y \in \mathbb{R}^{(\sum_i n_i)} \text{ and} \\ (x, y_1, \dots, y_m) &= (x, y) \in \mathbb{R}^{n_x} \times \mathbb{R}^{(\sum_i n_i)}. \end{aligned}$$

More generally, a symbol without a subscript refers to a tuple/block vector of the subscripted objects; for example we have $z = (z_1, z_2, \dots, z_J)$ for some vectors or scalars z_j .

Throughout this work we will not take for granted the existence of solutions to various mathematical optimization problems. Further, we will often deal with the optimal value of a math program. This means that, for consistency if nothing else, we will denote the general math program of “minimize the function f on the set G ” as “ $\inf_x \{f(x) : x \in G\}$.” As usual, “arg min” refers to the solution set, empty or not.

2.2 Generalized Nash equilibrium definition

Nash’s original definition of equilibrium in the game-theoretic context from [26, 27] is limited to players with constant decision sets (specifically, probability distributions over finite sets of actions). A subsequent generalization, due to Debreu [7], has proved popular, appearing largely unchanged as the central definition in a recent review of Nash equilibrium problems in [8]. We adopt and present this latter, generalized, definition.

Let $u = (u_1, \dots, u_{\widehat{m}}) \in \mathbb{R}^{\widehat{n}_1} \times \dots \times \mathbb{R}^{\widehat{n}_{\widehat{m}}}$ and for each i let u_{-i} denote the vector of variables excluding the vector u_i :

$$u_{-i} = (u_1, \dots, u_{i-1}, u_{i+1}, \dots, u_{\widehat{m}}).$$

Consider a set of \widehat{m} players; the i^{th} player is defined by an optimization problem parametric in u_{-i} :

$$\begin{aligned} R_i(u_{-i}) &= \arg \min_{v_i} f_i(v_i, u_{-i}) \\ &\text{s.t. } v_i \in U_i(u_{-i}), \end{aligned} \tag{\mathcal{B}_i}$$

where f_i is a real-valued function and U_i is a set-valued mapping into $\mathbb{R}^{\widehat{n}_i}$. A generalized Nash equilibrium (GNE) of (\mathcal{B}_i) is a point $(u_1, \dots, u_{\widehat{m}})$ such that for all $i \in \{1, \dots, \widehat{m}\}$,

$$u_i \in R_i(u_{-i}).$$

In words, a GNE is a point from which each player (\mathcal{B}_i) has no incentive to deviate, given the decisions of the other players. We may more colloquially refer to such a point as simply a Nash equilibrium.

2.3 Equivalence of Nash and optimization equilibria

We proceed to show that Nash equilibria and optimization equilibria are equally flexible modeling concepts; given the problem of finding a Nash equilibrium, one can easily define an equivalent problem in finding an optimization equilibrium and vice versa. More explicitly, Theorem 1 establishes that the set of optimization equilibrium problems is a subset of the set of Nash equilibrium problems, while Theorem 2 conversely establishes that the set of Nash equilibrium problems is a subset of the set of optimization equilibrium problems. Together, these results establish the equivalence of the equilibrium concepts.

First, we show that to go from an optimization equilibrium to a Nash equilibrium, we essentially add an extra player with trivial objective and constraints that enforce $(x, y) \in F$.

Theorem 1. Consider the optimization problems (\mathcal{A}_i) and set $F \subset \mathbb{R}^{n_x} \times \mathbb{R}^{n_1} \times \dots \times \mathbb{R}^{n_m}$. For $u = (u_1, \dots, u_m, u_{m+1}) \in \mathbb{R}^{n_1} \times \dots \times \mathbb{R}^{n_m} \times \mathbb{R}^{n_x}$ and $i \in \{1, \dots, m\}$, let

$$\begin{aligned} R'_i(u_{-i}) &= \arg \min_{v_i} g_i(u_{m+1}, v_i) \\ &\text{s.t. } v_i \in \widehat{Y}_i(u_{m+1}), \end{aligned} \tag{\mathcal{B}'_i}$$

and

$$\begin{aligned} R'_{m+1}(u_{-(m+1)}) &= \arg \min_{v_{m+1}} 0 \\ &\text{s.t. } v_{m+1} \in \mathbb{R}^{n_x}, \\ &\quad (v_{m+1}, u_1, \dots, u_m) \in F. \end{aligned} \tag{\mathcal{B}'_{m+1}}$$

Then $(x^*, y_1^*, \dots, y_m^*)$ is an optimization equilibrium of (\mathcal{A}_i) with respect to F if and only if $(u_1^*, \dots, u_m^*, u_{m+1}^*) \equiv (y_1^*, \dots, y_m^*, x^*)$ is a GNE of (\mathcal{B}'_i) ($i \in \{1, \dots, m+1\}$).

Proof. Let $(x^*, y_1^*, \dots, y_m^*)$ be an optimization equilibrium; by definition we have $(x^*, y_1^*, \dots, y_m^*) \in F$, and for each i , $y_i^* \in S_i(x^*)$. With the definitions given in the theorem, it is easy to see that for each $i \in \{1, \dots, m\}$ we have $R'_i(u_{-i}^*) = S_i(x^*)$ and so $u_i^* = y_i^*$ is optimal for (\mathcal{B}'_i) . Further, the player (\mathcal{B}'_{m+1}) is a feasibility problem, for which $u_{m+1}^* = x^*$ is feasible and thus optimal: $u_{m+1}^* \in R'_{m+1}(u_{-(m+1)}^*)$. We conclude that $(u_1^*, \dots, u_{m+1}^*)$ is a Nash equilibrium.

Conversely, let $u^* = (u_1^*, \dots, u_m^*, u_{m+1}^*)$ be a GNE of (\mathcal{B}'_i) : $u_i^* \in R'_i(u_{-i}^*)$. Then $x^* = u_{m+1}^* \in R'_{m+1}(u_{-(m+1)}^*)$; again this implies $(x^*, y^*) \in F$. As before, we have $S_i(x^*) = R'_i(u_{-i}^*)$, and so $y_i^* = u_i^*$ is optimal for (\mathcal{A}_i) for each i . Thus $(x^*, y_1^*, \dots, y_m^*)$ is an optimization equilibrium. \square

Next, to go from a Nash equilibrium to an optimization equilibrium, we must define F so that the solution components of each player can be “communicated” to one another appropriately, essentially by defining the x variables as a copy of the y variables.

Theorem 2. Consider the optimization problems (\mathcal{B}_i) . Define

$$\begin{aligned} F' &= \{(x_1, \dots, x_{\widehat{m}}, y_1, \dots, y_{\widehat{m}}) : x_i = y_i, \forall i \in \{1, \dots, \widehat{m}\}\} \\ &\subset \left(\mathbb{R}^{\widehat{n}_1} \times \dots \times \mathbb{R}^{\widehat{n}_{\widehat{m}}} \right)^2, \end{aligned}$$

and for $x = (x_1, \dots, x_{\widehat{m}})$ and $i \in \{1, \dots, \widehat{m}\}$, let

$$\begin{aligned} S'_i(x) &= \arg \min_{z_i} f_i(z_i, x_{-i}) \\ &\text{s.t. } z_i \in U_i(x_{-i}). \end{aligned} \tag{\mathcal{A}'_i}$$

Then $u^* = (u_1^*, \dots, u_{\widehat{m}}^*)$ is a GNE of (\mathcal{B}_i) if and only if $(x^*, y^*) \equiv (u^*, u^*)$ is an optimization equilibrium of (\mathcal{A}'_i) ($i \in \{1, \dots, \widehat{m}\}$) with respect to F' .

Proof. Let (u_1^*, \dots, u_m^*) be a GNE: $u_i^* \in R_i(u_{-i}^*)$ for each i . Since $x^* = u^*$, we see that $S'_i(x^*) = R_i(u_{-i}^*)$ and thus $y_i^* = u_i^*$ is optimal for (\mathcal{A}'_i) , for each i . Since $x^* = y^*$, we have $(x^*, y^*) \in F'$ by definition of F' . Thus $(x^*, y_1^*, \dots, y_m^*)$ is an optimization equilibrium.

Conversely, let (x^*, y^*) be an optimization equilibrium of (\mathcal{A}'_i) with respect to F' : $(x^*, y^*) \in F'$, and $y_i^* \in S'_i(x^*)$. Once again, since $u^* = x^*$, we have $R_i(u_{-i}^*) = S'_i(x^*)$ and so $u_i^* = y_i^*$ is optimal for (\mathcal{B}_i) for each i . Thus (u_1^*, \dots, u_m^*) is a Nash equilibrium. \square

2.4 Terminology

Given the equivalence between the concepts of Nash and optimization equilibrium established in the previous subsection, we will drop the modifier “optimization” and just call a point an *equilibrium* if it satisfies Definition 1 for appropriate data. Furthermore, all subsequent problems will be posed in the optimization equilibrium form of Section 2.1, and so there should be no risk of confusion.

3 Characterization of equilibrium and minimum disequilibrium

In this section we pose an abstract math program that can be used to characterize an equilibrium, and as a by-product, we introduce the concept of minimum disequilibrium. Connections to the Nikaido-Isoda function are discussed as well.

We can express the problem of finding an equilibrium as a feasibility problem for the following bilevel program:

$$\begin{aligned} & \inf_{x, y_1, \dots, y_m} && 0 \\ & \text{s.t.} && (x, y_1, \dots, y_m) \in F, \\ & && y_i \in \arg \min_{z_i} \{g_i(x, z_i) : z_i \in Y_i, h_i(x, z_i) \leq 0\}, \quad \forall i. \end{aligned}$$

We can reformulate this bilevel problem as

$$\begin{aligned} & \inf_{x, y_1, \dots, y_m} && 0 && (1a) \\ & \text{s.t.} && (x, y_1, \dots, y_m) \in F, && (1b) \\ & && h_i(x, y_i) \leq 0, \forall i, && (1c) \\ & && y_i \in Y_i, \forall i, && (1d) \\ & && g_i(x, y_i) \leq g_i(x, z_i), \quad \forall z_i \in Y_i : h_i(x, z_i) \leq 0, \forall i. && (1e) \end{aligned}$$

Such a reformulation has been considered in, for instance, [2, 32, 33], although it is easy to see that y_i is optimal for the i^{th} lower-level agent problem if and only if it is feasible (Constraints (1c) and (1d)) and its objective value is less than that of any other feasible point (Constraint (1e)). Depending on the cardinality of the Y_i sets, Problem (1) may have an infinite number of constraints (for each i , z_i is effectively an index for the last set of constraints (1e)). Thus, (1) may be categorized as a generalized semi-infinite program (GSIP) (see e.g. [15] for a review). Whether or not (1) is truly infinitely constrained depends on the nature of the Y_i sets; however, whether or not any of the Y_i are infinite does not complicate the following analysis and discussion. Thus, to allow for the most general context possible, we will refer to problem (1) as a GSIP.

Once again, (1) is a feasibility problem; evidently the most difficult constraint to handle is the “infinite” one (1e). Thus, we can consider replacing the trivial objective with one which aims to

minimize the violation of the infinite constraint. To this end, let

$$\begin{aligned} g &: (x, y) \mapsto (g_1(x, y_1), g_2(x, y_2), \dots, g_m(x, y_m)), \\ g^* &: (x) \mapsto (g_1^*(x), g_2^*(x), \dots, g_m^*(x)). \end{aligned}$$

When an agent problem is unbounded, g^* is extended real-valued. Define $\bar{\mathbb{R}} = \mathbb{R} \cup \{-\infty, +\infty\}$ (with the usual order), their Cartesian product $\bar{\mathbb{R}}^m = \bar{\mathbb{R}} \times \dots \times \bar{\mathbb{R}}$, and the non-negative orthant $\bar{\mathbb{R}}_+^m = \{(v_1, \dots, v_m) \in \bar{\mathbb{R}}^m : v_i \geq 0, \forall i\}$.

In the following result, which characterizes equilibrium as the solution of an optimization problem, the objective can be thought of as a measure of “disequilibrium,” or roughly the dissatisfaction of all the players in aggregate. This depends on an $\bar{\mathbb{R}}$ -valued function μ ; a convenient choice is

$$\mu : (v_1, \dots, v_m) \mapsto \sum_{i=1}^m v_i,$$

although we allow for different forms (for instance, any norm satisfies the required properties).

Proposition 1. *Let $\mu : \bar{\mathbb{R}}^m \rightarrow \bar{\mathbb{R}}$ be any function satisfying:*

1. *if $v \in \bar{\mathbb{R}}_+^m$ then $\mu(v) \geq 0$;*
2. *$v \in \bar{\mathbb{R}}_+^m$ and $\mu(v) = 0$ if and only if $v_i = 0$ for all i .*

Consider

$$\begin{aligned} \eta &= \inf_{x, y_1, \dots, y_m} \mu(g(x, y) - g^*(x)) \\ &\text{s.t. } (x, y_1, \dots, y_m) \in F^{\text{MD}} \end{aligned} \tag{MD}$$

where the feasible set F^{MD} is defined as

$$F^{\text{MD}} = \{(x, y_1, \dots, y_m) \in F : y_i \in Y_i, h_i(x, y_i) \leq 0, \forall i\}. \tag{2}$$

Any (x, y_1, \dots, y_m) is an equilibrium if and only if it is a solution of (MD) with $\eta = 0$.

Proof. First, note that for any (x, y_1, \dots, y_m) feasible in (MD), we must have $g_i(x, y_i) \geq g_i^*(x)$ (since y_i is feasible in (\mathcal{A}_i)). Thus, since $\mu(v) \geq 0$ for all $v \geq 0$, the objective value of (MD) is bounded below by zero; that is, $\eta \geq 0$.

Let (x, y_1, \dots, y_m) be an equilibrium. Then $(x, y_1, \dots, y_m) \in F$ and for each i , y_i is optimal for (\mathcal{A}_i) ; thus $g_i(x, y_i) = g_i^*(x)$ for each i , and so $\mu(g(x, y) - g^*(x)) = 0$. Furthermore, y_i is feasible in (\mathcal{A}_i) so $y_i \in Y_i$ and $h_i(x, y_i) \leq 0$ for each i ; therefore, (x, y_1, \dots, y_m) is feasible in (MD), and has an objective value equal to zero. It follows that $\eta \leq 0$, but using the reverse inequality established above, we have $\eta = 0$, and (x, y_1, \dots, y_m) is a solution of (MD).

Let (x, y_1, \dots, y_m) be a solution of (MD) with $\eta = 0$. As noted above, by feasibility of (x, y) , we have $g(x, y) - g^*(x) \geq 0$. Using the properties of μ , we must have for all i that $g_i(x, y_i) = g_i^*(x)$; but further, y_i is feasible for (\mathcal{A}_i) and thus optimal. Finally, since we also have $(x, y_1, \dots, y_m) \in F$, it follows that (x, y_1, \dots, y_m) is an equilibrium. \square

We can interpret a solution of Problem (MD) as the “closest” point to being an equilibrium. Specifically, its solution is a point $(x, y_1, \dots, y_m) \in F$ such that y_i is ϵ_i -optimal in (\mathcal{A}_i) for each i , that is,

$$g_i(x, y_i) = g_i^*(x) + \epsilon_i,$$

and $\mu(\epsilon_1, \dots, \epsilon_m)$ is at a minimum. This perspective suggests that finding a solution with $\eta > 0$ may still be useful and provide a meaningful or practically useful point. In general, we will refer to a solution of Problem (MD) as a “minimum disequilibrium” solution.

As mentioned, minimum disequilibrium agrees with the minimum total opportunity cost idea in [10]. Thus, that work provides excellent motivation for the value of a minimum disequilibrium solution for decision making in certain electricity markets; we provide a brief discussion here. In some electricity markets, a market operator has the problem of setting short-term electricity prices. The agents in this setting are electricity generators who decide their level of generation given prices, and the linking constraints F may model a required minimum level of electricity generation to meet demand, as well as limits on prices. Generators may have fixed start-up costs; this introduces nonconvexities, and a competitive equilibrium may not exist. To address this, different approaches can be taken. For instance, in the PJM electricity market “uplift” or “make whole” payments are introduced, to ensure that generators operate at the required levels to meet demand without losing money. Naturally, this leads to the idea of minimizing the total of the uplift payments. A related formulation, the minimum total opportunity cost, aims to find prices (x), and for each generator i , operational levels and other decisions (y_i), so that the total discrepancy between the generators’ realized profits ($-g_i(x, y_i)$) and what they *could* have made ($-g_i^*(x)$) is minimized. The result is a formulation that fits into the form of Problem (\mathcal{MD}). We will explore this further in Section 6.2.

Once again, characterization of a Nash equilibrium in terms of the solution of an optimization problem has been proposed before, typically utilizing the NI function. See Appendix A for further discussion on this connection as well an alternate proof of Proposition 1 from this perspective. However, we reiterate the practical use of an optimization problem such as (\mathcal{MD}) to characterize equilibrium has largely been confined to cases assuming a certain amount of convexity.

4 Application: Competitive economic equilibrium

After specifying a general problem form consistent with spatial price equilibrium problems (see also Section 6.1), we analyze an approach to solving this problem that will highlight connections to Lagrangian duality. In particular, we see that our characterization of equilibrium permits a generalization to the nonconvex case of the classic result that optimal dual variables can be interpreted as equilibrium prices for spatial price equilibrium problems; see Theorem 3.

4.1 General problem structure

First, assume that each agent may be written as

$$\inf_{z_i} \left\{ g_i^a(z_i) + \sum_{k \in K} g_{i,k}^b(z_i) g_k^c(x) : z_i \in Y_i \right\},$$

for appropriate \mathbb{R} -valued functions g_i^a , $g_{i,k}^b$, g_k^c and a finite set $K = \{1, \dots, n_K\}$. That is, their feasible sets are independent of x , and their objectives have related forms. Further, assume that the global constraint set F has the following form:

$$F = \mathbb{R}^{n_x} \times \left\{ (y_1, \dots, y_m) : \sum_i g_{i,k}^b(y_i) = 0, \forall k \in K \right\}.$$

Similarly, F is “independent” of x .

This particular structure is relevant for modeling price-taking behavior of agents in multiple market structures. For example, single commodity spatial price equilibrium considered in Section 6.1 fits into this form.

The significance of these assumptions becomes clear when writing down problem (\mathcal{MD}) under these assumptions. If we let $\mu(v) = \sum_i v_i$, what is important to note is that for all $(x, y_1, \dots, y_m) \in$

F , we have

$$\sum_i g_i(x, y_i) = \sum_i g_i^a(y_i),$$

since the other terms in the objectives cancel. Thus we obtain

$$\begin{aligned} \eta &= \inf_{x, y_1, \dots, y_m} \sum_i g_i^a(y_i) - \sum_i g_i^*(x) \\ &\text{s.t. } \sum_i g_{i,k}^b(y_i) = 0, \forall k \in K, \\ &\quad y_i \in Y_i, \forall i. \end{aligned} \tag{3}$$

4.2 Primal-dual optimality and minimum disequilibrium

Inspecting problem (3), we see that its feasible set is $\mathbb{R}^{n_x} \times \left\{ y \in \prod_i Y_i : \sum_i g_{i,k}^b(y_i) = 0, \forall k \in K \right\}$.

Thus we can write $\eta = \eta^P - \eta^{\tilde{D}}$ where

$$\begin{aligned} \eta^P &= \inf_{y_1, \dots, y_m} \sum_i g_i^a(y_i) \\ &\text{s.t. } \sum_i g_{i,k}^b(y_i) = 0, \forall k \in K, \\ &\quad y_i \in Y_i, \forall i, \end{aligned} \tag{P}$$

and

$$\begin{aligned} \eta^{\tilde{D}} &= -\inf_x -\sum_i g_i^*(x) \\ &= \sup_x \inf_z \sum_i g_i^a(z_i) + \sum_{k \in K} g_k^c(x) \sum_i g_{i,k}^b(z_i) \\ &\text{s.t. } z_i \in Y_i, \forall i \end{aligned} \tag{\tilde{D}}$$

where we have used the definition of g_i^* and the fact that $\sup_x f(x) = -\inf_x (-f(x))$ for any real function f . We note a striking similarity between (\tilde{D}) and the Lagrangian dual problem of (P) :

$$\eta^D = \sup_{\lambda} \inf_z \left\{ \sum_i g_i^a(z_i) + \sum_k \lambda_k \sum_i g_{i,k}^b(z_i) : z_i \in Y_i, \forall i \right\}. \tag{D}$$

In particular, we have that $\eta^{\tilde{D}} \leq \eta^D$, since (\tilde{D}) is a restriction of (D) ; the supremum in (D) is over all $\lambda \in \mathbb{R}^{n_K}$, including the range/image of g_k^c . Meanwhile, we always have $\eta^D \leq \eta^P$, which is known as weak duality (see for instance [4, Prop. 5.1.3]). Thus we have

$$\eta^{\tilde{D}} \leq \eta^D \leq \eta^P.$$

These considerations inspire us to characterize a minimum disequilibrium solution in terms of optimal primal-dual solutions. The next result formally states that this decomposition of problem (3) is correct; although not particularly profound on its own, the subsequent results are far more useful and basically fall out as corollaries.

Proposition 2. *If (x^*, y^*) is a minimum disequilibrium solution of (3), then $\eta = \eta^P - \eta^{\tilde{D}}$. Further, (x^*, y^*) is a minimum disequilibrium solution of (3) if and only if $y^* = (y_1^*, \dots, y_m^*)$ is optimal for (P) and x^* is optimal for (\tilde{D}) .*

Proof. First we show the equivalence of solutions of (3) and solutions of (\mathcal{P}) and $(\tilde{\mathcal{D}})$. Let y^* and x^* be optimal for (\mathcal{P}) and $(\tilde{\mathcal{D}})$, respectively. For a contradiction, assume that (x^*, y^*) is not optimal for (3). Then there exists (x, y) feasible in (3) with $\sum_i g_i^a(y_i) - \sum_i g_i^*(x) < \sum_i g_i^a(y_i^*) - \sum_i g_i^*(x^*)$. This implies that either $-\sum_i g_i^*(x) < -\sum_i g_i^*(x^*)$, or $\sum_i g_i^a(y_i) < \sum_i g_i^a(y_i^*)$, or both; in any case this contradicts the optimality of x^* , y^* , or both. Thus (x^*, y^*) is optimal for (3) (that is, (x^*, y^*) is a minimum disequilibrium solution).

Conversely, assume that (x^*, y^*) is optimal for (3). Once again, for a contradiction assume that y^* is not optimal in (\mathcal{P}) , or x^* is not optimal in $(\tilde{\mathcal{D}})$. Take the first case; then there exists y feasible in (\mathcal{P}) such that $\sum_i g_i^a(y_i) - \sum_i g_i^*(x^*) < \sum_i g_i^a(y_i^*) - \sum_i g_i^*(x^*)$, which contradicts the optimality of (x^*, y^*) . Similar reasoning holds in the second case that x^* is not optimal in $(\tilde{\mathcal{D}})$. Thus y^* and x^* are optimal for (\mathcal{P}) and $(\tilde{\mathcal{D}})$, respectively.

To show the first claim, let (x^*, y^*) be a minimum disequilibrium solution of (3). Then as already shown, y^* and x^* are optimal for (\mathcal{P}) and $(\tilde{\mathcal{D}})$. In this case $\eta^P = \sum_i g_i^a(y_i^*)$ and $\eta^{\tilde{D}} = \sum_i g_i^*(x^*)$ and so $\eta^P - \eta^{\tilde{D}} = \sum_i g_i^a(y_i^*) - \sum_i g_i^*(x^*) = \eta$. \square

The next result takes advantage of the fact that the dual problem (\mathcal{D}) is a relaxation of $(\tilde{\mathcal{D}})$.

Proposition 3. *If $y^* = (y_1^*, \dots, y_m^*)$ is optimal for (\mathcal{P}) , and $\lambda^* \equiv (g_1^c(x^*), \dots, g_{n_K}^c(x^*))$ is optimal for (\mathcal{D}) , then $\eta^{\tilde{D}} = \eta^D$ and (x^*, y^*) is a minimum disequilibrium solution of (3).*

Proof. This follows from Proposition 2 and the fact that (\mathcal{D}) is a relaxation of $(\tilde{\mathcal{D}})$; x^* is a point that achieves the upper bound η^D on the optimal value of $(\tilde{\mathcal{D}})$, and so must be optimal for $(\tilde{\mathcal{D}})$. \square

Proposition 3 suggests the following procedure to find a minimum disequilibrium: solve the primal problem (\mathcal{P}) ; solve its Lagrangian dual problem (\mathcal{D}) to get an optimal dual solution λ^* ; find x^* such that $g_k^c(x^*) = \lambda_k^*$ for each k . Optimization of the dual problem entails all the usual caveats, but may be tractable for certain problem structures. When strong duality holds, $\eta^P = \eta^D$, and we see that disequilibrium is zero: $\eta^P - \eta^{\tilde{D}} = 0$.

In Section 6.1, x is a vector of prices, and $(g_1^c, \dots, g_{n_K}^c)$ is the identity mapping. In this case, problems $(\tilde{\mathcal{D}})$ and (\mathcal{D}) coincide and Proposition 2 then implies that optimal dual variables λ^* are the prices corresponding to a minimum disequilibrium solution, and the duality gap equals the minimized disequilibrium. When an equilibrium exists, we regain the classic interpretation of optimal dual variables as equilibrium prices. This is summarized in the following result.

Theorem 3. *Suppose that $n_x = n_K$ and $(g_1^c, \dots, g_{n_K}^c)$ is the identity mapping (that is, $g^c(x) = x$). Then (λ^*, y^*) is a minimum disequilibrium solution of (3) if and only if $y^* = (y_1^*, \dots, y_m^*)$ is optimal for (\mathcal{P}) and λ^* is optimal for (\mathcal{D}) . If y^* is optimal for (\mathcal{P}) and λ^* is optimal for (\mathcal{D}) , then η equals the duality gap $\eta^P - \eta^D$.*

Proof. As noted, $(\tilde{\mathcal{D}})$ and (\mathcal{D}) coincide, and the result follows from Proposition 2. \square

The example in Section 6.1 elaborates on this result and we show that the primal problem (\mathcal{P}) is the “maximum social welfare” problem that appears in many applications of economic equilibrium.

Theorem 3 is a rather clean version of various results in the literature dealing with pricing in electricity markets. For instance, Theorem 8 of [9] addresses a slightly different problem form and asserts that a point is an equilibrium if and only if it is primal and dual optimal and there is zero duality gap. Certain results about the properties of convex hull pricing, introduced in [14] and studied further in [30], go through similar arguments. The application of the present work to that literature is a fertile area of future research.

5 Solution method for minimum disequilibrium

Problem (\mathcal{MD}) may be nontrivial to solve. Evidently, the challenge stems from the fact that the agents' optimal value functions, g_i^* , are in general implicitly defined. If the agent problems have some known structure that permits a more explicit description of the optimal value function, then that could be used to reformulate the problems. Notably, if the agent problems are convex, arguments from duality theory or KKT conditions can be used to obtain various mathematical programs; see for instance [11], or from the perspective of GSIP, [31].

However, our aim in this section is to develop methods that can solve Problem (\mathcal{MD}) (or an equivalent reformulation) under fairly broad assumptions, including the cases that the agent problems are nonconvex. In particular, since verification of equilibrium requires the global solution of (\mathcal{MD}) , our discussion is motivated by methods for global optimization.

For the sake of clarity, we will make two simplifying assumptions. First, for the function μ appearing in Proposition 1, we use the previously mentioned choice $\mu : v \mapsto \sum_i v_i$. Second, we focus on developing a solution algorithm for finding a minimum disequilibrium solution when all the agents have constant feasible sets (some authors refer to this setting as a Nash, as opposed to generalized Nash, problem). The solution method is in essence a method for solving a semi-infinite program (SIP). In moving to the general case, we obtain a GSIP (recall problem (1)). We refer to recent work in [25, 19] for adapting this basic solution method for SIP to GSIP, and leave their application to equilibrium problems for future work.

Under the assumption that the agents have constant feasible sets, the i^{th} agent problem (\mathcal{A}_i) is

$$g_i^*(x) = \inf_{z_i} \{g_i(x, z_i) : z_i \in Y_i\}.$$

Further, we assume that $g_i^*(x)$ is finite valued for all x and i . This holds, for instance, when g_i is continuous and Y_i is compact for each i (which is required to show that the main algorithm converges). Thus, for each i we introduce a new scalar variable w_i which essentially approximates g_i^* from below, and Problem (\mathcal{MD}) becomes

$$\begin{aligned} \eta &= \inf_{x, y, w} \sum_i (g_i(x, y_i) - w_i) & (\mathcal{MD}^N) \\ \text{s.t. } & (x, y_1, \dots, y_m) \in F, \\ & y_i \in Y_i, \quad \forall i, \\ & w_i \leq g_i^*(x), \quad \forall i. \end{aligned}$$

For some sets $Y_i^L \subset Y_i$, consider the following problem:

$$\begin{aligned} \eta^L &= \inf_{x, y, w} \sum_i (g_i(x, y_i) - w_i) & (4) \\ \text{s.t. } & (x, y_1, \dots, y_m) \in F, \\ & y_i \in Y_i, \quad \forall i, \\ & w_i \leq g_i(x, z_i), \quad \forall z_i \in Y_i^L, \forall i. \end{aligned}$$

Since Y_i^L is a subset of the feasible set of the i^{th} agent, we have that $g_i^*(x) \leq \inf \{g_i(x, z_i) : z_i \in Y_i^L\}$. Thus, Problem (4) is a relaxation of Problem (\mathcal{MD}^N) , and thus η^L is a lower bound on η .

The proposed solution method is given in Algorithm 1. This algorithm is in essence the general algorithm from [5] for the solution of SIP. That work and subsequent generalizations (see [24, 17]) explore this general class of methods further and other details of their implementation. For instance,

Algorithm 1 Solution method for Problem (\mathcal{MD}^N)

Require: $\varepsilon > 0$, $Y_i^L \subset Y_i$, $Y_i^L \neq \emptyset$, for each i

```

1:  $\eta^U = +\infty$ 
2: loop
3:   Solve Problem (4) to obtain optimal solution  $(x, y, w)$  and value  $\eta^L$ .
4:   For this value of  $x$ , solve agent problems  $(\mathcal{A}_i)$  to obtain optimal solution  $z_i$  and value  $g_i^*(x)$ ,
     for each  $i$ .
5:   if  $w_i \leq g_i^*(x)$ , for all  $i$  then
6:      $(x^*, y^*) \leftarrow (x, y)$ 
7:     return  $(x^*, y^*)$ 
8:   else
9:      $Y_i^L \leftarrow Y_i^L \cup \{z_i\}$  for each  $i$ 
10:  end if
11:  if  $\sum_i (g_i(x, y_i) - g_i^*(x)) < \eta^U$  then
12:     $\eta^U \leftarrow \sum_i (g_i(x, y_i) - g_i^*(x))$ 
13:     $(x^*, y^*) \leftarrow (x, y)$ 
14:  end if
15:  if  $\eta^U - \max\{\eta^L, 0\} < \varepsilon$  then
16:    return  $(x^*, y^*)$ 
17:  end if
18: end loop

```

convex structures permit some specializations of the algorithm (see [5]), although we stress that this method applies even without convexity assumptions.

Algorithm 1 can terminate in one of two ways. Either:

1. The solution to Problem (4) at a certain iteration is feasible in Problem (\mathcal{MD}^N) , indicated by $w_i \leq g_i^*(x)$ for each i (note that this relies on the global minimizer of (4) being found, otherwise some w_i might be “too small”). Then the solution to Problem (4) is optimal for (\mathcal{MD}^N) , since (4) is a relaxation. Thus $\eta^L = \eta$, and the value (x^*, y^*) returned is a solution.

If one only cares about the existence of an equilibrium solution, the algorithm can terminate early if it happens that $\eta^L > 0$; since we have that η^L is a lower bound on η , we have by Proposition 1 that an equilibrium solution does not exist.

2. An ε -optimal solution (x^*, y^*) of Problem (\mathcal{MD}^N) is found. Note that at Step 12, $\sum_i (g_i(x, y_i) - g_i^*(x))$ is an upper bound for η ; we have merely evaluated the objective at a feasible point of Problem (\mathcal{MD}^N) . More specifically, at any iteration for which the upper bound η^U is finite, we have a point (x^*, y^*) which came from a solution of Problem (4); thus, $(x^*, y^*) \in F$ and $y_i^* \in Y_i$ for each i and so $(x, y, w) = (x^*, y^*, g^*(x^*))$ is feasible in Problem (\mathcal{MD}^N) , and its objective value is an upper bound on η . The upper bound η^U tracks the best of these upper bounds.

The termination criterion at Step 15 is justified since 0 is an *a priori* lower bound on η , thus we can terminate if $\eta^U < \varepsilon$.

The following result establishes when the algorithm produces an ε -optimal solution in finite iterations. To be clear, we obtain a point $(x^*, y^*) \in F$ satisfying $y_i^* \in Y_i$ for each i , and if $g_i(x^*, y_i^*) = g_i^*(x^*) + \varepsilon_i$ for each i , then $\sum_i \varepsilon_i < \eta + \varepsilon$. Thus if $\sum_i \varepsilon_i$ is small, then as in the

discussion following Proposition 1, we have a point that is “close” to being an equilibrium. In general, we obtain an ε -approximate minimum disequilibrium solution.

Theorem 4. *Assume that F^{MD} (which in this setting reduces to $\{(x, y) : y_i \in Y_i, (x, y_1, \dots, y_m) \in F\}$) is compact and nonempty. Assume that for each i , g_i is continuous and Y_i is compact. Then for any $\varepsilon > 0$, Algorithm 1 produces an ε -optimal solution (x^*, y^*) of Problem (MD^N) in finite iterations.*

See Appendix B for its proof.

6 Examples

Here we present two examples in order to illustrate how to apply our theoretical and numerical developments.

6.1 Single commodity spatial price equilibrium

We apply the theoretical developments to an example from spatial price equilibrium (SPE), a classic problem with a long history going back to e.g. [29]. We show how social welfare maximization corresponds to the primal problem (\mathcal{P}) and how Algorithm 1 specializes to a method for solving the dual problem.

First we define the data of the SPE problem so that it fits in the form considered in Section 4.1. Consider a single commodity market with m agents. Each agent’s optimization problem is represented as follows:

$$S_i(x) = \arg \min_{z_i^P, z_i^C, z_i^I, z_i^E} c_i^P(z_i^P) + x_i z_i^I - u_i(z_i^C) - \sum_{j:j \neq i} (x_j - c_{ij}^E) z_{ij}^E \quad (\mathcal{A}_i^{\text{SPE}})$$

s.t. $(z_i^P, z_i^C, z_i^I, z_i^E) \in Y_i,$

where z_i^P is the amount produced, z_i^C is the amount consumed, z_i^I is the amount imported, and z_{ij}^E is the amount exported to agent j . The objective function models the individual agent’s social welfare; $c_i^P(z_i^P)$ is the production cost function, $u_i(z_i^C)$ is the consumer benefit function of the agent¹, x_j is the price that agent j accepts, and c_{ij}^E the exporting cost to agent j . The constraint set Y_i can enforce limits on any of the quantities, as well as intra-agent supply-demand balance conditions like

$$z_i^P + z_i^I = z_i^C + \sum_{j:j \neq i} z_{ij}^E.$$

A common modeling assumption is that each agent is exclusively a producer or consumer. For a producer, we can model this in the present setting by defining u_i as a constant function taking the value zero and defining Y_i so that import levels z_i^I are restricted to be zero. Similar considerations apply for a consumer.

The decisions z_i of agent i do not appear as parameters in the problem of any other agent j ; each agent only depends on the prices x . Thus, they are price-takers. However, the agents are

¹Specifically, we might have

$$u_i(z_i^C) = \int_0^{z_i^C} p_i(\zeta) d\zeta,$$

i.e. it is the integral under the inverse demand curve p_i .

importing and exporting the commodity, and the notion of equilibrium depends on these quantities balancing. Thus F is defined by

$$F = \left\{ (x, y^P, y^C, y^I, y^E) : y_j^I = \sum_{i:i \neq j} y_{ij}^E, \forall j \right\},$$

which enforces that the amount imported by agent j , y_j^I , does in fact equal the sum of the export amounts of the other agents.

This fits into the problem form from Section 4.1 by letting $K = \{1, \dots, m\}$, $n_x = m$, and for each i , $z_i = (z_i^P, z_i^C, z_i^I, z_{i1}^E, \dots, z_{im}^E)$, $g_i^a(z_i) = c_i^P(z_i^P) - u_i(z_i^C) + \sum_{j:j \neq i} c_{ij}^E z_{ij}^E$, $g_i^c(x) = x_i$, $g_{ii}^b(z_i) = z_i^I$, and $g_{ij}^b(z_i) = -z_{ij}^E$, for $j \neq i$. In that way, $g_i^a(z_i) + \sum_j g_{ij}^b(z_i) g_j^c(x)$ equals the objective of (\mathcal{A}_i^{SPE}) . Further, for each j , $\sum_i g_{ij}^b(z_i) = 0$ is equivalent to $g_{jj}^b(z_j) + \sum_{i:i \neq j} g_{ij}^b(z_i) = z_j^I - \sum_{i:i \neq j} z_{ij}^E = 0$, which are indeed the conditions enforced by the definition of F above.

The next question is what the analysis in Section 4.2 looks like for this specific problem. Inspired by Theorem 3, we may want to solve the primal problem (\mathcal{P}) . For the present SPE problem, problem (\mathcal{P}) takes the form

$$\eta^P = \inf_{y_1, \dots, y_m} \sum_i \left(c_i^P(y_i^P) - u_i(y_i^C) + \sum_{j:j \neq i} c_{ij}^E y_{ij}^E \right) \quad (5a)$$

$$\text{s.t. } y_i^I = \sum_{j:j \neq i} y_{ji}^E, \forall i, \quad (5b)$$

$$y_i = (y_i^P, y_i^C, y_i^I, y_i^E) \in Y_i, \forall i. \quad (5c)$$

We recognize this as the max social welfare problem that is often at the heart of SPE problems and other forms of equilibrium problems.

Then if Problem (5) is feasible and bounded, and for each i , c_i^P is convex, u_i is concave, and Y_i is polyhedral, classic Lagrangian duality results imply that strong duality holds (see for instance [4, Prop. 5.2.1]), and by Theorem 3, the optimal solution of (5) along with the optimal Lagrange multipliers of constraint (5b) yield an equilibrium solution.

Of course, Theorem 3 holds more generally. While it is difficult to establish conditions under which strong duality holds for general nonconvex problems (although see [16] for the notion of ‘‘copositive duality’’ for mixed-binary quadratic programs), Theorem 3 establishes that if we solve the max social welfare problem and its dual, we will arrive at a minimum disequilibrium solution that may have value in certain situations or as an approximate equilibrium.

How do we solve the dual problem? Algorithm 1 provides the answer. By similar reasoning as in Section 4.2, for the SPE setting the lower-bounding problem (4) decomposes into the max social welfare problem (5) above and

$$\begin{aligned} \eta^{D,U} &= \sup_{x,w} \sum_i w_i \\ \text{s.t. } w_i &\leq c_i^P(z_i^P) - u_i(z_i^C) + \sum_{j:j \neq i} c_{ij}^E z_{ij}^E + x_i(z_i^I - \sum_{j:j \neq i} z_{ij}^E), \quad \forall z_i \in Y_i^L, \forall i. \end{aligned} \quad (6)$$

The max social welfare problem is static and only needs to be solved once; Algorithm 1 reduces to iteratively solving the problem over (x, w) above and the agent problems. Solving the agent problems may be seen as equivalently minimizing the Lagrangian of (5):

$$\begin{aligned} \sum_i g_i^*(x) &= \inf_{z_1, \dots, z_m} \sum_i \left(c_i^P(z_i^P) - u_i(z_i^C) + \sum_{j:j \neq i} c_{ij}^E z_{ij}^E + x_i(z_i^I - \sum_{j:j \neq i} z_{ij}^E) \right) \\ \text{s.t. } z_i &\in Y_i, \forall i. \end{aligned} \quad (7)$$

Table 1: Data for producer agents

Symbol	$i = 1$	$i = 2$	$i = 3$
γ_i	10	45	35
δ_i	0.05	0.1	0.002
d_i	4000	100	2000
K_i^L	400	200	300
K_i^U	600	250	500

We recognize this overall procedure as a bundle subgradient method or cutting plane method for maximizing the concave dual function. See for instance Section 6.3.3 of [4].

6.2 Unit commitment in electricity markets

As discussed at the end of Section 3, models of electricity markets are a source of equilibrium problems with nonconvex agents. Here, we present a simple example from [10] to demonstrate how Algorithm 1 works in practice.

The example is the “single period unit commitment model” from [10]. We have three price-taking, profit-maximizing power producers modeled by mixed-integer quadratic programs (MIQP). An integer (specifically binary) decision is required to model a fixed cost for starting up generation. Thus, the producer agents are parametric in the price x^p ; for $i \in \{1, 2, 3\}$, we have

$$\begin{aligned}
\min_{z_i} \quad & \gamma_i z_i^c + (1/2)\delta_i(z_i^c)^2 + d_i z_i^b - x^p z_i^c \\
\text{s.t.} \quad & z_i^b K_i^L \leq z_i^c \leq z_i^b K_i^U, \\
& z_i^c \in \mathbb{R}, \\
& z_i^b \in \{0, 1\}.
\end{aligned}$$

Data are in Table 1.

In the terminology of [10], the consumption side of this problem is “non-dispatchable.” In other words, consumption should *not* be treated as an agent potentially contributing to disequilibrium². Consumption is modeled instead by an inverse demand curve that provides constraints between the level of consumption x^q and price x^p that are included in the global constraints F . Thus we have $x = (x^p, x^q)$ and

$$F = \left\{ (x, y_1, y_2, y_3) : x^p = \alpha - \beta x^q, x^q = \sum_{i=1}^3 y_i^c, x^q \in [0, K_1^U + K_2^U + K_3^U] \right\},$$

where $\alpha = 200$ and $\beta = 0.2$. Note that F also includes the basic requirement that production and consumption balance, as well as bounds on consumption implied by production bounds.

The ultimate goal is to determine price and consumption that minimize disequilibrium, which in this case equals total opportunity cost defined in [10]. Algorithm 1 is applicable. We first guess a consumption level of $x^q = 100$, set the price accordingly (i.e. $x^p = \alpha - \beta x^q$), solve the agent problems, and use the optimal solutions to initialize the Y_i^L sets. This initial guess of consumption is arbitrary and does not affect the answer provided by Algorithm 1. We can then solve the

²Our modeling framework handles the other case, i.e. dispatchable consumption, equally well. In this case we proceed by the analysis in Section 6.1.

lower-bounding problem (4) which takes the form

$$\begin{aligned}
\eta^L = \inf_{x,y,w} & \sum_i (\gamma_i y_i^c + (1/2)\delta_i (y_i^c)^2 + d_i y_i^b - x^p y_i^c - w_i) \\
\text{s.t. } & x^p = \alpha - \beta x^q, \\
& x^q = \sum_i y_i^c, \\
& x^q \in [0, K_1^U + K_2^U + K_3^U], \\
& y_i^b K_i^L \leq y_i^c \leq y_i^b K_i^U, \quad \forall i, \\
& y_i^c \in \mathbb{R}, \quad \forall i, \\
& y_i^b \in \{0, 1\}, \quad \forall i, \\
& w_i \leq \gamma_i z_i^c + (1/2)\delta_i (z_i^c)^2 + d_i z_i^b - x^p z_i^c, \quad \forall z_i \in Y_i^L, \forall i.
\end{aligned}$$

Note that we may use the constraint $x^p = \alpha - \beta x^q$ to eliminate the variable x^p . Further, we can transform the objective to

$$-(\alpha - \beta x^q)x^q + \sum_i (\gamma_i y_i^c + (1/2)\delta_i (y_i^c)^2 + d_i y_i^b - w_i)$$

where we have used the constraint $x^q = \sum_i y_i^c$. Thus, the lower-bounding problem is an MIQP.

The method converges after three iterations when we have $\eta^L = \eta^U = 931.41$. The minimum disequilibrium price and consumption are $x^{p,*} = 39.5$ and $x^{q,*} = 802.5$. These values agree with those found by a heuristic procedure in [10], but thanks to Theorem 4, we can guarantee that these are in fact the minimum disequilibrium values.

7 Conclusions

This work has analyzed and expanded upon the connections between Nash equilibrium and bilevel programming. Some of these connections seem to exist in the literature in vague terms; by posing them independently of the typical settings of complementarity problems, variational inequalities, or Nikaido-Isoda functions, we have been able to leverage ideas from the broader literature on nonconvex bilevel programs in order to propose solution algorithms for equilibrium problems with potentially nonconvex agents. The connections with bilevel programming and GSIP provide directions for future research in applying solution methods for those problems to general equilibrium problems.

The concept of minimum disequilibrium was introduced as an alternative solution when no equilibrium exists. It was shown with an example that this solution concept can handle the minimum total opportunity cost idea in unit commitment problems. Furthermore, special but still economically relevant situations (competitive spatial price equilibrium) can be analyzed to show that optimal dual variables have significance as part of a minimum disequilibrium solution. In general, the examples have demonstrated the flexibility of the optimization equilibrium modeling framework and how specialization of Algorithm 1 to various settings provides a principled way to find practical solution methods to nonconvex equilibrium problems.

Acknowledgments

The authors would like to thank their colleague Nicolas Sawaya for introducing some of the challenges associated with equilibrium modeling within a real-world setting, as well as the various fruitful discussions on this topic over the past several years. The authors would also like to thank their colleague Myun-Seok Cheon for similarly fruitful discussions while developing this work.

A Connections with the Nikaido-Isoda function

Optimization formulations of equilibrium problems have been presented in the literature before. Many rely on the Nikaido-Isoda (NI) function, first proposed by Nikaido and Isoda in [28], although see [8, 35] for those results expressed in terms of a generalized Nash problem as defined in Section 2.2. We show that we can characterize equilibrium with a result similar to Proposition 1 using an NI function.

Consider the generalized Nash equilibrium problem defined by the agents (\mathcal{B}_i) ($i \in \{1, \dots, \widehat{m}\}$), from Section 2.2. For $u = (u_1, \dots, u_{\widehat{m}})$, define

$$\begin{aligned} \phi(u) = \sup_{v_1, \dots, v_{\widehat{m}}} \sum_{i=1}^{\widehat{m}} (f_i(u_i, u_{-i}) - f_i(v_i, u_{-i})) \\ \text{s.t. } v_i \in U_i(u_{-i}), \forall i \in \{1, \dots, \widehat{m}\}. \end{aligned}$$

The objective function in this optimization problem defining ϕ is the NI function as defined by [8, 35]. Theorem 3.2 of [8] states that $u^* = (u_1^*, \dots, u_{\widehat{m}}^*)$ is a GNE of (\mathcal{B}_i) if and only if $\phi(u^*) = 0$ and

$$u^* \in \arg \min_u \{\phi(u) : u_i \in U_i(u_{-i}), i \in \{1, \dots, \widehat{m}\}\}. \quad (8)$$

This minimization problem is over the set $\prod_i U_i(u_{-i})$, which is parameterized by the decision variables u . This leads the authors of [8] to refer to such a problem in the general case as a “quasi-optimization” problem. Avoiding this non-standard concept and notation is another benefit of the optimization equilibrium formalism.

Now consider the problem of finding an optimization equilibrium defined by the agent problems (\mathcal{A}_i) and set F , and use Theorem 1 to obtain an equivalent Nash equilibrium problem (defined by (\mathcal{B}'_i) , for $i \in \{1, \dots, m+1\}$). Theorem 1 implies that $U_i(u_{-i}) = \widehat{Y}_i(u_{m+1})$, and $U_{m+1}(u_{-(m+1)}) = \{x : (x, u_1, \dots, u_m) \in F\}$, and so letting $\widehat{Y}_i(x) = \{z_i \in Y_i : h_i(x, z_i) \leq 0\}$, we can write the feasible set of (8) in terms of the data of the optimization equilibrium as

$$\begin{aligned} \Phi = \{ & (u_1, \dots, u_{m+1}) : \\ & u_i \in Y_i, \forall i \in \{1, \dots, m\}, \\ & h_i(u_{m+1}, u_i) \leq 0, \forall i \in \{1, \dots, m\}, \\ & (u_{m+1}, u_1, \dots, u_m) \in F\}, \end{aligned}$$

which coincides with the feasible set of Problem (\mathcal{MD}) (map the variables as $(x, y_1, \dots, y_m) = (u_{m+1}, u_1, \dots, u_m)$). Further, Theorem 1 implies that $f_i(v_i, u_{-i}) = g_i(u_{m+1}, v_i)$ and in particular f_{m+1} is identically zero; we transform ϕ into

$$\begin{aligned} \phi(u) = \sum_{i=1}^m g_i(u_{m+1}, u_i) - \inf_v \{ \sum_{i=1}^m g_i(u_{m+1}, v_i) : \\ v_i \in Y_i, \forall i \in \{1, \dots, m\}, \\ h_i(u_{m+1}, v_i) \leq 0, \forall i \in \{1, \dots, m\}, \\ (v_{m+1}, u_1, \dots, u_m) \in F\}. \end{aligned}$$

For $u \in \Phi$, the infimum above is over a nonempty set, and furthermore, does not depend on the v_{m+1} variable. Consequently, we can ignore the global constraints encoded in F and decompose the minimization. Thus, for $u \in \Phi$, the expression for $\phi(u)$ simplifies to

$$\phi(u) = \sum_{i=1}^m g_i(u_{m+1}, u_i) - \sum_{i=1}^m g_i^*(u_{m+1})$$

where we recall the optimal agent value function g_i^* defined in Section 2.1. Finally, note that this expression equals the objective function of Problem (\mathcal{MD}) when $\mu : v \mapsto \sum_i v_i$. Thus, Problems (\mathcal{MD}) and (8) coincide, and the statement that u^* is an equilibrium iff $\phi(u^*) = 0$ and $u^* \in \arg \min \{\phi(u) : u \in \Phi\}$, essentially provides an alternate proof of Proposition 1.

B Proof of Theorem 4

Theorem (4). *Assume that F^{MD} (which in this setting reduces to $\{(x, y) : y_i \in Y_i, (x, y_1, \dots, y_m) \in F\}$) is compact and nonempty. Assume that for each i , g_i is continuous and Y_i is compact. Then for any $\varepsilon > 0$, Algorithm 1 produces an ε -optimal solution (x^*, y^*) of Problem (\mathcal{MD}^N) in finite iterations.*

Theorem 4 is a consequence of the following results, which establish convergence of the upper and lower bounds. The analysis proceeds similarly to that in [5]. In particular, the convergence of the lower bound is nearly identical to the proof of Theorem 2.1 in [5]. We include it for completeness, and further, the constructions employed are useful for establishing a simple convergent upper bound, which distinguishes this analysis from [5, 24].

First we have convergence of the lower bounds.

Proposition 4. *Assume that the set*

$$F^{\text{MD}} = \{(x, y) : y_i \in Y_i, (x, y_1, \dots, y_m) \in F\}$$

is compact and nonempty. Assume that for each i , g_i is continuous and Y_i is compact. If the sequence of lower bounds produced by Algorithm 1 is $(\eta^{L,k})_{k \in \mathbb{N}}$, then $\eta^{L,k} \rightarrow \eta$.

Proof. Let $((x^k, y^k, w^k))_{k \in \mathbb{N}}$ be the sequence of solutions of Problem (4) produced by Algorithm 1. Since more elements are added to Y_i^L at each iteration, the part of the solution sequence $(w_i^k)_k$ is decreasing, but bounded below by the minimum of g_i on F^{MD} ; continuity and compactness ensure that this is finite (specifically, $\inf_{x,y,z_i} \{g_i(x, z_i) : z_i \in Y_i, (x, y) \in F^{\text{MD}}\} > -\infty$). Thus, for each i , $\{w_i^k : k \in \mathbb{N}\}$ is contained in a compact set, and so the entire solution sequence is in a compact set. For each i , let $(z_i^k)_{k \in \mathbb{N}}$ be the corresponding sequence of solutions to the agent problem (\mathcal{A}_i) . Again, the image of this sequence $\{z_i^k : k \in \mathbb{N}\}$ is in a compact set for each i . Consequently, we have that a subsequence of solutions converges to some point. Abusing notation, we have that $(x^k, y^k, w^k) \rightarrow (x^*, y^*, w^*)$ and $z^k \rightarrow z^*$. Note that we have $(x^*, y^*) \in F^{\text{MD}}$.

First, we establish that (x^*, y^*, w^*) is feasible in Problem (\mathcal{MD}^N) . Since z_i^k is added to Y_i^L at the end of each iteration, we have for each i

$$w_i^\ell \leq g_i(x^\ell, z_i^k), \quad \forall \ell, k : \ell > k.$$

By taking the limit over ℓ , and then the limit over k , we get for each i

$$w_i^* \leq g_i(x^*, z_i^*). \tag{9}$$

Now, for a contradiction, assume that for some i , $w_i^* > g_i^*(x^*)$, indicating that (x^*, y^*, w^*) is not feasible in Problem (\mathcal{MD}^N) . This means that there exists $z_i^\dagger \in Y_i$ (feasible in the agent problem) with

$$w_i^* > g_i(x^*, z_i^\dagger).$$

By continuity of g_i , for all k sufficiently large we have

$$w_i^k > g_i(x^k, z_i^\dagger).$$

By definition of z_i^k as the minimizer of (\mathcal{A}_i) for $x = x^k$, we have

$$g_i(x^k, z_i^\dagger) \geq g_i(x^k, z_i^k).$$

Taking limits we have

$$w_i^* > g_i(x^*, z_i^\dagger) \geq g_i(x^*, z_i^*),$$

which contradicts (9). Thus, (x^*, y^*, w^*) is feasible in Problem (\mathcal{MD}^N) .

As mentioned, $\eta^{L,k} \leq \eta$ for all k , and since constraints are added to Problem (4) at each iteration, $\eta^{L,k}$ is increasing (this is true even without going to a subsequence). Thus, $(\eta^{L,k})_k$ converges to $\eta^{L,*} \leq \eta$, but of course we can use the subsequence from before to get $\sum_i (g_i(x^k, y_i^k) - w_i^k) = \eta^{L,k}$, and so

$$\sum_i (g_i(x^*, y_i^*) - w_i^*) = \eta^{L,*} \leq \eta.$$

But by the previously established feasibility,

$$\sum_i (g_i(x^*, y_i^*) - w_i^*) \geq \eta$$

which establishes $\eta^{L,k} \rightarrow \eta$. □

Next we establish convergence of the upper bounds.

Proposition 5. *Assume that the set*

$$F^{\text{MD}} = \{(x, y) : y_i \in Y_i, (x, y_1, \dots, y_m) \in F\}$$

is compact and nonempty. Assume that for each i , g_i is continuous and Y_i is compact. If the sequence of upper bounds produced by Algorithm 1 is $(\eta^{U,k})_{k \in \mathbb{N}}$, then $\eta^{U,k} \rightarrow \eta$.

Proof. As in the proof of Proposition 4, we can assume that we have the convergent (sub)sequences $(x^k, y^k, w^k) \rightarrow (x^*, y^*, w^*)$ and $z^k \rightarrow z^*$. At a particular iteration, we have evaluated $g_i^*(x^k) = g_i(x^k, z_i^k)$. Then as noted earlier,

$$\bar{\eta}^k = \sum_i (g_i(x^k, y_i^k) - g_i^*(x^k))$$

is an upper bound for η . Then using the continuity of each g_i , we have

$$\bar{\eta}^k = \sum_i (g_i(x^k, y_i^k) - g_i(x^k, z_i^k)) \rightarrow \sum_i (g_i(x^*, y_i^*) - g_i(x^*, z_i^*)) \equiv \bar{\eta}^*.$$

By Inequality (9)

$$\sum_i (g_i(x^*, y_i^*) - w_i^*) \geq \sum_i (g_i(x^*, y_i^*) - g_i(x^*, z_i^*)),$$

but as in the proof of Proposition 4, $\sum_i (g_i(x^*, y_i^*) - w_i^*) = \eta$, and so $\eta \geq \bar{\eta}^*$. Since $\eta \leq \bar{\eta}^k$ for all k , we have $\eta \leq \bar{\eta}^*$, and combined with the opposite inequality we have $\bar{\eta}^k \rightarrow \eta$.

Finally, the sequence $\eta^{U,k}$ is constructed as $\eta^{U,k} = \min \{\eta^{U,k-1}, \bar{\eta}^k\}$. Thus $\eta^{U,k}$ is also an upper bound of η , and forms a decreasing sequence. Furthermore $\bar{\eta}^k \geq \eta^{U,k}$, and since a subsequence of $(\bar{\eta}^k)_k$ converges to η , we also have that $(\eta^{U,k})_k$ converges to η . □

References

- [1] Kenneth J. Arrow and Gerard Debreu. Existence of an equilibrium for a competitive economy. *Econometrica: Journal of the Econometric Society*, pages 265–290, 1954.
- [2] Jonathan F. Bard. An algorithm for solving the general bilevel programming problem. *Mathematics of Operations Research*, 8(2):260–272, 1983.
- [3] Tamer Basar and Hasan Selbuz. Closed-loop Stackelberg strategies with applications in the optimal control of multilevel systems. *IEEE Transactions on Automatic Control*, 24(2):166–179, 1979.
- [4] Dimitri P. Bertsekas. *Nonlinear Programming*. Athena Scientific, Belmont, Massachusetts, second edition, 1999.
- [5] Jerry W. Blankenship and James E. Falk. Infinitely constrained optimization problems. *Journal of Optimization Theory and Applications*, 19(2):261–281, 1976.
- [6] Herminia I. Calvete and Carmen Galé. Linear bilevel multi-follower programming with independent followers. *Journal of Global Optimization*, 39(3):409–417, 2007.
- [7] Gerard Debreu. A social equilibrium existence theorem. *Proceedings of the National Academy of Sciences*, 38(10):886–893, 1952.
- [8] Francisco Facchinei and Christian Kanzow. Generalized Nash equilibrium problems. *Annals of Operations Research*, 175(1):177–211, 2010.
- [9] J. David Fuller. Market equilibrium models with continuous and binary variables. Technical report, 2008.
- [10] J. David Fuller and Emre Çelebi. Alternative models for markets with nonconvexities. *European Journal of Operational Research*, 261(2):436–449, 2017.
- [11] Steven A. Gabriel, Antonio J. Conejo, J. David Fuller, Benjamin F. Hobbs, and Carlos Ruiz. *Complementarity Modeling in Energy Markets*. Springer, New York, 2013.
- [12] Steven A. Gabriel, Antonio J. Conejo, Carlos Ruiz, and Sauleh Siddiqui. Solving discretely constrained, mixed linear complementarity problems with applications in energy. *Computers & Operations Research*, 40(5):1339–1350, 2013.
- [13] Steven A. Gabriel, Sauleh Ahmad Siddiqui, Antonio J. Conejo, and Carlos Ruiz. Solving discretely-constrained Nash-Cournot games with an application to power markets. *Networks and Spatial Economics*, 13(3):307–326, 2013.
- [14] Paul R. Gribik, William W. Hogan, and Susan L. Pope. Market-clearing electricity prices and energy uplift. Technical report, Harvard Electricity Policy Group, 2007.
- [15] Francisco Guerra-Vázquez, J-J Rückmann, Oliver Stein, and Georg Still. Generalized semi-infinite programming: a tutorial. *Journal of Computational and Applied Mathematics*, 217(2):394–419, 2008.
- [16] Cheng Guo, Merve Bodur, and Joshua A. Taylor. Copositive duality for discrete markets and games. *arXiv preprint arXiv:2101.05379*, 2021.

- [17] Stuart M. Harwood and Paul I. Barton. Lower level duality and the global solution of generalized semi-infinite programs. *Optimization*, 65(6):1129–1149, 2016.
- [18] Yuan-wei Jing and Si-ying Zhang. The solution to a kind of Stackelberg game systems with multi-follower: Coordinative and incentive. In *Analysis and optimization of systems*, pages 593–602. Springer, 1988.
- [19] Peter Kirst and Oliver Stein. Global optimization of generalized semi-infinite programs using disjunctive programming. *Journal of Global Optimization*, 73(1):1–25, 2019.
- [20] Jacek B. Krawczyk. Coupled constraint Nash equilibria in environmental games. *Resource and Energy Economics*, 27(2):157–181, 2005.
- [21] Jacek B. Krawczyk and Stanislav Uryasev. Relaxation algorithms to find Nash equilibria with economic applications. *Environmental Modeling & Assessment*, 5(1):63–73, 2000.
- [22] Jie Lu, Chenggen Shi, and Guangquan Zhang. On bilevel multi-follower decision making: General framework and solutions. *Information Sciences*, 176(11):1607–1627, 2006.
- [23] Giandomenico Mastroeni. Gap functions for equilibrium problems. *Journal of Global Optimization*, 27(4):411–426, 2003.
- [24] Alexander Mitsos. Global optimization of semi-infinite programs via restriction of the right-hand side. *Optimization*, 60(10-11):1291–1308, 2011.
- [25] Alexander Mitsos and Angelos Tsoukalas. Global optimization of generalized semi-infinite programs via restriction of the right hand side. *Journal of Global Optimization*, 61(1):1–17, 2015.
- [26] John Nash. Equilibrium points in n-person games. *Proceedings of the National Academy of Sciences*, 36(1):48–49, 1950.
- [27] John Nash. Non-cooperative games. *Annals of Mathematics*, pages 286–295, 1951.
- [28] Hukukane Nikaidō and Kazuo Isoda. Note on non-cooperative convex games. *Pacific Journal of Mathematics*, 5(Suppl. 1):807–815, 1955.
- [29] Paul A. Samuelson. Spatial price equilibrium and linear programming. *The American Economic Review*, 42(3):283–303, 1952.
- [30] Dane A. Schiro, Tongxin Zheng, Feng Zhao, and Eugene Litvinov. Convex hull pricing in electricity markets: Formulation, analysis, and implementation challenges. *IEEE Transactions on Power Systems*, 31(5):4068–4075, 2015.
- [31] Oliver Stein and Georg Still. Solving semi-infinite optimization problems with interior point techniques. *SIAM Journal on Control and Optimization*, 42(3):769–788, 2003.
- [32] Angelos Tsoukalas, Berç Rustem, and Efstratios N. Pistikopoulos. A global optimization algorithm for generalized semi-infinite, continuous minimax with coupled constraints and bilevel problems. *Journal of Global Optimization*, 44(2):235–250, 2009.
- [33] Hoang Tuy, Athanasios Migdalas, and Peter Värbrand. A global optimization approach for the linear two-level program. *Journal of Global Optimization*, 3(1):1–23, 1993.

- [34] Stanislav Uryas'ev and Reuven Y. Rubinstein. On relaxation algorithms in computation of noncooperative equilibria. *IEEE Transactions on Automatic Control*, 39(6):1263–1267, 1994.
- [35] Anna von Heusinger and Christian Kanzow. Optimization reformulations of the generalized Nash equilibrium problem using Nikaido-Isoda-type functions. *Computational Optimization and Applications*, 43(3):353–377, 2009.
- [36] Anna von Heusinger, Christian Kanzow, and Masao Fukushima. Newton's method for computing a normalized equilibrium in the generalized Nash game through fixed point formulation. *Mathematical Programming*, 132(1):99–123, 2012.
- [37] Guangquan Zhang and Jie Lu. Fuzzy bilevel programming with multiple objectives and cooperative multiple followers. *Journal of Global Optimization*, 47(3):403–419, 2010.