

# MAXIMUM LIKELIHOOD FOR HIGH-NOISE GROUP ORBIT ESTIMATION AND SINGLE-PARTICLE CRYO-EM

ZHOU FAN, ROY R. LEDERMAN, YI SUN, TIANHAO WANG, AND SHENG XU

**ABSTRACT.** Motivated by applications to single-particle cryo-electron microscopy (cryo-EM), we study several problems of function estimation in a high noise regime, where samples are observed after random rotation and possible linear projection of the function domain. We describe a stratification of the Fisher information eigenvalues according to transcendence degrees of graded pieces of the algebra of group invariants, and we relate critical points of the log-likelihood landscape to a sequence of moment optimization problems, extending previous results for a discrete rotation group without projections.

We then compute the transcendence degrees and forms of these optimization problems for several examples of function estimation under  $\text{SO}(2)$  and  $\text{SO}(3)$  rotations, including a simplified model of cryo-EM as introduced by Bandeira, Blum-Smith, Kileel, Perry, Weed, and Wein. We affirmatively resolve conjectures that 3<sup>rd</sup>-order moments are sufficient to locally identify a generic signal up to its rotational orbit in these examples.

For low-dimensional approximations of the electric potential maps of two small protein molecules, we empirically verify that the noise-scalings of the Fisher information eigenvalues conform with our theoretical predictions over a range of SNR, in a model of  $\text{SO}(3)$  rotations without projections.

## CONTENTS

1. Introduction	1
2. The general orbit recovery model in high noise	5
3. Continuous multi-reference alignment	12
4. Spherical registration and cryo-EM	14
5. Numerical evaluations of the Fisher information	18
6. Conclusion	21
Appendix A. Proofs for general results on the orbit recovery model	21
Appendix B. Analysis of orthogonal Procrustes alignment	37
Appendix C. Analysis of continuous multi-reference alignment	39
Appendix D. Analyses of function estimation under an $\text{SO}(3)$ rotation	50
Appendix E. Details of the numerical simulations	83
Appendix F. Cryogenic Electron Microscopy (cryo-EM)	85
References	86

## 1. INTRODUCTION

We study several problems of function estimation in low dimensions, where the function is observed under random and unknown rotations of its domain. Let  $f : \mathcal{X} \rightarrow \mathbb{R}$  be a function on the unit circle  $\mathcal{X} = \mathcal{S}^1$ , the unit sphere  $\mathcal{X} = \mathcal{S}^2$ , or  $\mathcal{X} = \mathbb{R}^3$ . Let  $G$  be a rotation group acting on  $\mathcal{X}$ . We seek to estimate  $f$  from samples of the form

$$f_{\mathbf{g}}(x) + \text{white noise}$$

where each sample consists of the function  $f_{\mathbf{g}}(x) = f(\mathbf{g}^{-1} \cdot x)$  rotated by a uniformly random element  $\mathbf{g} \in G$  and observed with continuous Gaussian white noise on  $\mathcal{X}$ . Equivalently, choosing an orthonormal basis

---

ZF, RL, TW, SX: DEPARTMENT OF STATISTICS AND DATA SCIENCE, YALE UNIVERSITY

YS: DEPARTMENT OF STATISTICS, UNIVERSITY OF CHICAGO

*E-mail addresses:* zhou.fan@yale.edu, roy.lederman@yale.edu, yisun@statistics.uchicago.edu, tianhao.wang@yale.edu, sheng.xu@yale.edu.

*Date:* October 6, 2022.

for  $L_2(\mathcal{X})$ , the basis coefficients of  $f_g$  are observed with i.i.d. Gaussian noise, having some entrywise noise variance  $\sigma^2 > 0$ . We focus on a regime of this problem where each sample has high noise  $\sigma^2 \gtrsim \|f\|_{L_2}^2$ , and the information from many rotated samples must be combined to obtain an accurate estimate of  $f$ . We study also a variant of this model where samples are observed under an additional linear projection.

Our primary motivation is a formulation of this problem that models molecular reconstruction in single-particle cryo-electron microscopy (cryo-EM) [DAC<sup>+</sup>88, HBC<sup>+</sup>90, Fra06]. In this application,  $f : \mathbb{R}^3 \rightarrow \mathbb{R}$  is the electric potential of an unknown molecular structure. Tomographic projections of this potential are measured for many samples of the molecule, each in a different and unknown rotated orientation, typically with a high level of measurement noise. The molecular structure is determined by estimating this electric potential  $f$  from the rotated and projected samples, and then fitting an atomic model [BBS20, SS20]. A brief introduction to cryo-EM and a discussion of its relation to the problems studied in this work are presented in Appendix F.

Among computational procedures for solving this reconstruction problem, regularized versions of maximum likelihood estimation (MLE), as implemented via expectation-maximization or stochastic gradient descent, are commonly used [Sig98, SGV<sup>+</sup>07, Sch12, PRFB17]. However, many theoretical properties of the optimization landscape and reconstruction errors of these procedures are not fully understood in cryo-EM applications [BBS20].

In this work, we study the Fisher information matrix and log-likelihood function landscape associated with maximum likelihood estimation for a basic model of the cryo-EM reconstruction problem, as well as several simpler statistical models with qualitative similarities. These models may be of independent interest while building up to the complexity of cryo-EM:

- (Continuous multi-reference alignment, Section 3.) Estimating a function on the unit circle  $\mathcal{X} = \mathcal{S}^1$ , under  $\text{SO}(2)$  rotations of the circle [BNWR20, BJL<sup>+</sup>20].
- (Spherical registration, Section 4.1.) Estimating a function on the unit sphere  $\mathcal{X} = \mathcal{S}^2$ , under  $\text{SO}(3)$  rotations of the sphere [BBSK<sup>+</sup>17, Section 5.4].
- (Unprojected cryo-EM, Section 4.2.) Estimating a function on  $\mathbb{R}^3$  under  $\text{SO}(3)$  rotations about the origin, without tomographic projection [BBSK<sup>+</sup>17, Appendix B]. Such a problem arises in a related application of cryo-ET, discussed in Appendix F.1.
- (Cryo-EM, Section 4.3.) Estimating a function on  $\mathbb{R}^3$  under  $\text{SO}(3)$  rotations about the origin, with tomographic projection [BBSK<sup>+</sup>17, Section 5.5].

**1.1. Group orbit recovery and related literature.** Classical literature on function estimation has explored the rich interplay between the complexity of infinite-dimensional function classes, the statistical difficulty of estimation, and the role of regularization [IH81, Tsy08, Joh17]. We restrict attention instead to a finite-dimensional function space for each of the above models, with the goal of understanding connections between estimation in these models having latent rotations and the algebraic structure of the underlying rotational group.

Choosing a  $d$ -dimensional function basis, each of the above function estimation problems may be restated as an *orbit recovery problem* [APS18, BBSK<sup>+</sup>17] of estimating the coefficients  $\theta_* \in \mathbb{R}^d$  of  $f$  in this basis, from noisy observations of  $\theta_*$  that are rotated by elements of a subgroup  $G \subset \text{O}(d)$ . This subgroup  $G$  represents the transformation of basis coefficients under rotations of the function domain  $\mathcal{X}$ . A body of recent literature has studied both specific and general instances of this orbit recovery problem [PWB<sup>+</sup>19, ABL<sup>+</sup>19, BNWR20, BBSK<sup>+</sup>17, APS18, PSB19, Bru19, FSWW20, SKK<sup>+</sup>20, RBO21]. When  $G$  is the group of cyclic rotations of coordinates (a.k.a. discrete multi-reference alignment), [PWB<sup>+</sup>19] first proved that the optimal squared error for estimating generic signals  $\theta_* \in \mathbb{R}^d$  in high noise is significantly larger than that in the model without latent rotations, scaling as  $\sigma^6$  rather than as just the noise variance  $\sigma^2$ . This analysis was extended to non-generic signals for continuous multi-reference alignment in [BNWR20] and to general group actions in [APS18, BBSK<sup>+</sup>17].

Our current work is inspired, in particular, by results of Bandeira et al. [BBSK<sup>+</sup>17], which placed cryo-EM and other examples of function estimation in this context, and connected statistical properties of method-of-moments estimators in these problems to properties of the invariant algebra of the group action. Further connections between the algebraic structure of invariants and the geometry of the log-likelihood function landscape were developed in [FSWW20, KB20]. As a central technical ingredient, these papers derived a series expansion of the log-likelihood function in powers of  $\sigma^{-1}$ , in [FSWW20] for orbit recovery models

without linear projection, and in [KB20] for more general Gaussian mixture models that include the models we study in this work. We discuss below several relevant results of [BBSK<sup>+</sup>17, FSWW20, KB20] in further detail.

**1.2. Overview of results.** In Section 2, we introduce the general orbit recovery model both with and without a linear projection, and describe results that connect geometric properties of the log-likelihood function to properties of the invariant algebra of the group action. In Sections 3 and 4, we apply this connection to study the preceding problems of function estimation, including continuous multi-reference alignment (MRA) and cryo-EM. In Section 5, we report results of numerical simulations for estimating the electric potential functions of two small protein molecules in an unprojected cryo-EM model, which corroborate predictions of our theory for the spectra of the Fisher information matrices.

Here, we provide a brief overview of these results in the context of related literature.

### Fisher information and log-likelihood function landscape.

For general orbit recovery problems with  $\theta_* \in \mathbb{R}^d$ , in a high-noise regime  $\sigma^2 \gtrsim \|\theta_*\|^2$ , results of [FSWW20, KB20] demonstrated that it is informative to study properties of the (negative) population log-likelihood function  $R(\theta)$  via a series expansion in powers of  $\sigma^{-1}$  of the form

$$R(\theta) = \sum_{k=0}^{\infty} \frac{1}{\sigma^{2k}} R_k(\theta). \quad (1.1)$$

Each term  $R_k(\theta)$  is a  $G$ -invariant polynomial function of  $\theta$  which may depend on a number of “degrees-of-freedom” of  $\theta$  strictly smaller than the total dimension  $d$ . For a model with discrete group  $G$  and no linear projection, [FSWW20] showed this number exactly coincides with  $\text{trdeg } \mathcal{R}_{\leq k}^G$ , the transcendence degree of the invariant subalgebra generated by all  $G$ -invariant polynomials of degree  $\leq k$ . This implies a graded structure of the Fisher information matrix  $I(\theta_*)$  for generic signal vectors  $\theta_*$ , where eigenvalues corresponding to different degrees-of-freedom have different scalings with  $\sigma^{-1}$ . Furthermore, local minimizers of  $R(\theta)$  have a certain correspondence with successive local minimizers of each function  $R_k(\theta)$ .

In this work, we first extend these results to a model where  $G \subseteq O(d)$  may be continuous, and samples may be observed with an additional linear projection. This extension encompasses a basic formulation of the molecular reconstruction problem in cryo-EM. The main result of [KB20] implies that a series expansion analogous to (1.1) continues to hold for the population log-likelihood function  $R(\theta)$  in such a model. However, as anticipated from the structure of the expansion in [KB20], the algebraic properties of its terms differ from the unprojected setting studied in [FSWW20]. We show here that the number of degrees-of-freedom associated to each function  $R_k(\theta)$  coincides with the transcendence degree of a possibly reduced subalgebra generated by order- $k$  moments of the projected signal (Theorem 2.7). In addition, a version of the correspondence between local minimizers of  $R(\theta)$  and of successive local minimizers of  $R_k(\theta)$  remains true over a bounded domain of  $\mathbb{R}^d$  (Theorems 2.11 and 2.13). When the group  $G$  is continuous, we extend the arguments of [FSWW20] to address technical issues arising from the Fisher information matrix  $I(\theta_*)$  being singular, and the locus of minimizers of  $R(\theta)$  being a manifold of positive rather than zero dimension.

### Multi-reference alignment and cryo-EM.

These general results enable our study of maximum likelihood procedures in specific function estimation problems, the main focus of our work. In high-noise regimes of these problems, statistical properties of the MLE are related to the structures of the subalgebras  $\mathcal{R}_{\leq k}^G$  and to their transcendence degrees. In particular, the squared-error risk of the MLE is dictated by the smallest non-zero eigenvalue of the Fisher information matrix  $I(\theta_*)$ , and scales as  $\sigma^{2K}$  for generic signals  $\theta_*$  where  $K$  is the smallest integer for which  $\text{trdeg } \mathcal{R}_{\leq K}^G$  equals  $\text{trdeg } \mathcal{R}^G$ , the transcendence degree of the full  $G$ -invariant algebra. This connects with a central result of [BBSK<sup>+</sup>17], which showed that  $K$  is the lowest order moment needed to identify  $\theta_*$  up to a finite list of group orbits, and that  $\sigma^{2K}$  is also the scaling of the sample complexity required for estimating  $\theta_*$  up to such a finite list. We apply our general results to determine the explicit value of  $K$  in several function estimation examples.

For our model of continuous MRA on  $\mathcal{S}^1$ , we verify that  $K = 3$  (Theorem 3.1). This is expected from known results about estimation using 3<sup>rd</sup>-order moments in similar observation models for both discrete and continuous MRA in [PWB<sup>+</sup>19, BNWR20]. We also show that the optimization landscape of  $R(\theta)$  may

possess spurious local minimizers even for generic Fourier coefficient vectors  $\theta_* \in \mathbb{R}^d$  (Theorem 3.4) when the number of Fourier basis functions  $d$  exceeds a small constant. This statement is analogous to results shown for discrete MRA in [FSWW20], although our construction here in the continuous setting has a different structure.

For spherical registration and projected and unprojected cryo-EM under an  $\text{SO}(3)$ -action, a primary contribution of our work is proving also that  $\text{trdeg } \mathcal{R}_{\leq K}^G = \text{trdeg } \mathcal{R}^G$  for  $K = 3$ . An iterative algorithm for estimating  $\theta_*$  from 3<sup>rd</sup>-order moments in cryo-EM was first proposed by Kam in [Kam80], which implicitly assumed that these moments are sufficient to identify  $\theta_*$  (up to symmetries such as chirality). Formal conjectures that  $K = 3$  were stated in [BBSK<sup>+</sup>17] and verified numerically for small values of the basis dimension  $d$  in exact-precision arithmetic. We prove that  $K = 3$  for  $d$  exceeding small absolute constants (Theorems 4.1, 4.6, and 4.10), hence resolving several of these conjectures that 3<sup>rd</sup>-order moments are sufficient to locally identify the orbit of  $\theta_*$ .

Writing the terms  $R_k(\theta)$  of (1.1) as

$$R_k(\theta) = s_k(\theta) + q_k(\theta)$$

where  $s_k(\theta)$  depends on the additional degrees-of-freedom of  $\theta$  beyond those which define  $R_{k-1}(\theta)$ , our proofs of  $K = 3$  leverage a connection between  $\text{trdeg } \mathcal{R}_{\leq k}^G$  and the generic ranks of the Hessians  $\nabla^2 s_k(\theta)$  (Lemma 2.8). We show that  $\nabla^2 s_3(\theta)$  is generically of full rank by using an inductive “frequency marching” argument on the dimension  $d$  and explicitly analyzing  $\text{rank}(\nabla^2 s_3(\theta))$  for special choices of  $\theta \in \mathbb{R}^d$ . As a by-product of these analyses, we derive explicit forms for  $s_3(\theta)$ , which define optimization problems analogous to bispectrum inversion problems studied in MRA models [BBM<sup>+</sup>17].

In an unprojected spherical registration model over  $\mathcal{S}^2$ , recent independent work of [LM21] provides a more quantitative version of this inductive frequency marching argument. The result of [LM21, Lemma 5.6] implies that above some absolute constant dimension  $\underline{d}$ , the increase in  $\text{rank}(\nabla^2 s_3(\theta))$  from dimension  $\underline{d}$  to  $d$  must be exactly  $d - \underline{d}$  for generic  $\theta \in \mathbb{R}^d$ , and [LM21] obtained a quantitative lower bound on the smallest singular value in a smoothed analysis over  $\theta$ . Our proofs show versions of this statement that are less quantitative but more explicit about the value of  $\underline{d}$ , holding down to small enough  $\underline{d}$  where the full-rank condition for  $\nabla^2 s_3(\theta)$  may be explicitly checked. We carry this out for both spherical registration and cryo-EM, and in particular, our inductive argument in the projected cryo-EM model is more complex than in the unprojected models and uses different ideas.

### Simulations of the Fisher information for small proteins.

To empirically investigate the predictions of this body of theory in a cryo-EM example, we computed in simulation the observed Fisher information matrices for the electric potential functions of two small protein molecules—a rotavirus VP6 trimer and hemoglobin—in a model without tomographic projection.

We developed and employed a procedure of adaptively constructing a radial function basis in the Fourier domain (Appendix E) so as to reduce the dimension of the function space needed to approximate the true potential. Applying this construction, we obtained function bases of dimension  $d \approx 400$  that capture the coarse trimer structure of the rotavirus example, and of dimension  $d \approx 4000$  that capture the secondary structures of both proteins up to spatial resolutions of 7–8 Angstroms. At these dimensions and spatial resolutions, the theoretically predicted  $\sigma^{-2}$ ,  $\sigma^{-4}$ , and  $\sigma^{-6}$  scalings of the Fisher information eigenvalues were apparent in simulation for sufficiently high noise. We observe deviations from these predictions at lower levels of noise, and also in higher-dimensional function spaces that may be necessary to approximate the potentials to better spatial resolutions.

**Notation.** We use the conventions  $\langle u, v \rangle = \sum_i \bar{u}_i v_i$  for the complex inner product,  $\|u\|$  for the (real or complex)  $\ell_2$ -norm,  $\|M\|$  for the  $\ell_2 \rightarrow \ell_2$  operator norm for matrices, and  $\mathbf{i} = \sqrt{-1}$  for the imaginary unit.

For a measure space  $(X, \mu)$ ,  $L_2(X, \mathbb{C})$  is the  $L_2$ -space of functions  $f : X \rightarrow \mathbb{C}$  with inner-product  $\int_X \overline{f(x)} g(x) \mu(dx)$ . We write  $L_2(X) = L_2(X, \mathbb{R})$  for the analogous  $L_2$ -space of real-valued functions.  $\mathcal{S}^1$  and  $\mathcal{S}^2$  are the unit circle and unit sphere.

For differentiable  $f : \mathbb{R}^d \rightarrow \mathbb{R}^k$ ,  $df(x) \in \mathbb{R}^{k \times d}$  is its derivative or Jacobian at  $x$ . For twice-differentiable  $f : \mathbb{R}^d \rightarrow \mathbb{R}$ ,  $\nabla f(x) = df(x)^\top \in \mathbb{R}^d$  is its gradient, and  $\nabla^2 f(x) \in \mathbb{R}^{d \times d}$  is its Hessian. We will write  $d_x, \nabla_x, \nabla_x^2$  to clarify that the variable of differentiation is  $x$ . For a subset of coordinates  $y$ , we write  $\nabla_y f(x)$  and  $\nabla_y^2 f(x)$  as the components of this gradient and Hessian in  $y$ .

For a smooth manifold  $\mathcal{M}$  and twice-differentiable  $f : \mathcal{M} \rightarrow \mathbb{R}$ , we write  $\nabla f(x)|_{\mathcal{M}}$  and  $\nabla^2 f(x)|_{\mathcal{M}}$  for its gradient and Hessian evaluated in any choice of local chart at  $x \in \mathcal{M}$ . We will often not make the choice of chart explicit when referring to properties of  $\nabla f(x)|_{\mathcal{M}}$  and  $\nabla^2 f(x)|_{\mathcal{M}}$  that do not depend on the specific choice of chart.

**Acknowledgments.** We would like to thank Fred Sigworth for helpful discussions about cryo-EM, and for suggesting to us the hemoglobin example. ZF was supported in part by NSF DMS-1916198. RRL was supported in part by NIH/NIGMS 1R01GM136780-01. YS was supported in part by NSF DMS-1701654, DMS-2039183, and DMS-2054838.

## 2. THE GENERAL ORBIT RECOVERY MODEL IN HIGH NOISE

**2.1. Model and likelihood.** Let  $\theta_* \in \mathbb{R}^d$  be an unknown signal of interest. Let  $\mathbf{G} \subseteq \mathbf{O}(d)$  be a known compact subgroup of the orthogonal group of dimension  $d$ . We denote by  $\Lambda$  the unique Haar probability measure on  $\mathbf{G}$ , satisfying

$$\Lambda(\mathbf{G}) = 1, \quad \Lambda(g \cdot S) = \Lambda(S \cdot g) = \Lambda(S),$$

for any  $g \in \mathbf{G}$  and Borel measurable subset  $S \subseteq \mathbf{G}$ . In the *unprojected orbit recovery model*, we observe  $n$  noisy and rotated samples of  $\theta_*$ , given by

$$Y_i = g_i \cdot \theta_* + \sigma \varepsilon_i \in \mathbb{R}^d, \quad i = 1, \dots, n \quad (2.1)$$

where  $g_1, \dots, g_n \stackrel{iid}{\sim} \Lambda$  are Haar-uniform random elements of  $\mathbf{G}$ , and  $\varepsilon_1, \dots, \varepsilon_n \stackrel{iid}{\sim} \mathcal{N}(0, \text{Id}_{d \times d})$  are Gaussian noise vectors independent of  $g_1, \dots, g_n$ . The signal  $\theta_*$  is identifiable only up to an arbitrary rotation in  $\mathbf{G}$ , i.e. it is identifiable up to its orbit

$$\mathcal{O}_{\theta_*} = \{g \cdot \theta_* : g \in \mathbf{G}\}.$$

Our goal is to estimate  $\mathcal{O}_{\theta_*}$  from the observed rotated samples  $Y_1, \dots, Y_n$ .

In the *projected orbit recovery model*, we consider an additional known linear map  $\Pi : \mathbb{R}^d \rightarrow \mathbb{R}^{\tilde{d}}$ . (Note that  $\Pi$  may not necessarily be an orthogonal projection; our terminology is borrowed from the example of tomographic projection in cryo-EM.) We observe  $n$  samples

$$Y_i = \Pi(g_i \cdot \theta_*) + \sigma \varepsilon_i \in \mathbb{R}^{\tilde{d}}, \quad i = 1, \dots, n \quad (2.2)$$

where  $g_1, \dots, g_n \stackrel{iid}{\sim} \Lambda$  as before, and  $\varepsilon_1, \dots, \varepsilon_n \stackrel{iid}{\sim} \mathcal{N}(0, \text{Id}_{\tilde{d} \times \tilde{d}})$  are Gaussian noise vectors in the projected dimension  $\tilde{d}$ . Our goal is again to estimate  $\mathcal{O}_{\theta_*}$  from  $Y_1, \dots, Y_n$ .

The unprojected and projected orbit recovery models are both Gaussian mixture models, where the distribution of mixture centers is the law of  $g \cdot \theta_* \in \mathbb{R}^d$  or of  $\Pi(g \cdot \theta_*) \in \mathbb{R}^{\tilde{d}}$  induced by the uniform law  $g \sim \Lambda$  over  $\mathbf{G}$ . This mixture distribution may be continuous if  $\mathbf{G} \subseteq \mathbf{O}(d)$  is a continuous subgroup. In both models, we denote the negative sample log-likelihood as

$$R_n(\theta) = -\frac{1}{n} \sum_{i=1}^n \log p_\theta(Y_i),$$

where  $p_\theta(Y_i)$  is the Gaussian mixture density for  $Y_i$ , marginalizing over the unknown rotation  $g_i \sim \Lambda$ . This density is given in the projected setting by

$$p_\theta(y) = \int_{\mathbf{G}} \frac{1}{(2\pi\sigma^2)^{\tilde{d}/2}} \exp\left(-\frac{\|y - \Pi(g \cdot \theta)\|^2}{2\sigma^2}\right) d\Lambda(g), \quad (2.3)$$

and in the unprojected setting by the same expression with  $\Pi = \text{Id}$  and  $\tilde{d} = d$ . The maximum likelihood estimator (MLE) of  $\theta_*$  is  $\hat{\theta}_n = \arg \min_{\theta \in \mathbb{R}^d} R_n(\theta)$ . Since  $R_n$  satisfies the invariance  $R_n(\theta) = R_n(g \cdot \theta)$  for all  $g \in \mathbf{G}$ , the MLE is also only defined up to its orbit  $\mathcal{O}_{\hat{\theta}_n}$ .

**Remark 2.1** (Identifiability of the orbit). The parameter  $\theta_*$  is identifiable up to the distribution of the mixture centers  $g \cdot \theta_*$  or  $\Pi(g \cdot \theta_*)$ . In the unprojected model, the equality in law  $g \cdot \theta \stackrel{L}{=} g \cdot \theta'$  over  $g \sim \Lambda$  holds if and only if  $\mathcal{O}_\theta = \mathcal{O}_{\theta'}$ , so  $\theta_*$  is identifiable exactly up to its orbit.

In projected models, there may be further non-identifiability. For instance, under the tomographic projection arising in cryo-EM, we have  $\Pi(g \cdot \theta) \stackrel{L}{=} \Pi(g \cdot \theta')$  when  $\theta'$  represents the mirror reflection of  $\theta$  [BBS20].

Thus in this setting there may be two distinct orbits which cannot be further identified, and  $\theta_*$  is recovered only up to chirality.

In general, the number of distinct orbits with the same image under  $\Pi$  depends on the interaction between the structures of  $\mathbf{G}$  and  $\Pi$ , and can be infinite. For example, for the trivial group  $\mathbf{G} = \{\text{Id}\}$  and the projection  $\Pi : \mathbb{R}^d \rightarrow \mathbb{R}^{d-k}$  that removes the last  $k$  coordinates of  $\theta$ ,  $\mathcal{O}_\theta = \{\theta\}$  and  $\Pi(\mathcal{O}_\theta) = \Pi(\mathcal{O}_{\theta_*})$  for any  $\theta$  sharing the same first  $d - k$  coordinates as  $\theta_*$ .

We use the equivalence notation

$$\Pi(\mathcal{O}_\theta) \equiv \Pi(\mathcal{O}_{\theta_*})$$

to mean that  $\Pi(\mathcal{O}_\theta) = \Pi(\mathcal{O}_{\theta_*})$  as subsets of  $\mathbb{R}^{\tilde{d}}$ , and in addition,  $\Pi(g \cdot \theta) \stackrel{L}{=} \Pi(g \cdot \theta_*)$  under the Haar-uniform law  $g \sim \Lambda$ . Thus  $\theta_*$  is identifiable up to this equivalence. We will restrict attention to projected models where

$$\text{there are a finite number of orbits } \mathcal{O}_\theta \text{ such that } \Pi(\mathcal{O}_\theta) \equiv \Pi(\mathcal{O}_{\theta_*}), \text{ for generic } \theta_* \in \mathbb{R}^d. \quad (2.4)$$

An equivalent algebraic characterization is provided in Proposition 2.6(b) below.

We denote the negative *population* log-likelihood function by

$$R(\theta) = \mathbb{E}[R_n(\theta)] = -\mathbb{E}[\log p_\theta(Y)], \quad (2.5)$$

where the expectation is taken under the true model  $Y \sim p_{\theta_*}$ .  $R(\theta)$  depends implicitly on  $\theta_*$ , but we will omit this dependence in the notation. This population log-likelihood is minimized at  $\theta \in \mathcal{O}_{\theta_*}$  in the unprojected model, and at  $\{\theta : \Pi(\mathcal{O}_\theta) \equiv \Pi(\mathcal{O}_{\theta_*})\}$  in projected models.

**2.2. Invariant polynomials and the high-noise expansion.** For sufficiently high noise  $\sigma^2$ , it is informative to study  $R(\theta)$  via a series expansion of the Gaussian density of (2.3) in powers of  $\sigma^{-1}$ , as developed in [FSWW20, KB20]. We review this expansion in this section.

Let  $\mathcal{R}^{\mathbf{G}}$  be the (real) algebra of all  $\mathbf{G}$ -invariant polynomial functions  $p : \mathbb{R}^d \rightarrow \mathbb{R}$ . These are the polynomials  $p$  that satisfy

$$p(\theta) = p(g \cdot \theta) \text{ for all } \theta \in \mathbb{R}^d \text{ and } g \in \mathbf{G}.$$

For each integer  $k \geq 0$ , let  $\mathcal{R}_{\leq k}^{\mathbf{G}}$  be the subalgebra generated by the  $\mathbf{G}$ -invariant polynomials having total degree at most  $k$ . This subalgebra consists of the polynomials  $p \in \mathcal{R}^{\mathbf{G}}$  that may be expressed as  $p(\theta) = q(p_1(\theta), \dots, p_j(\theta))$  for some polynomial  $q$  and some  $p_1, \dots, p_j \in \mathcal{R}^{\mathbf{G}}$  each having degree  $\leq k$  (where  $p$  itself may have degree larger than  $k$ ).

Examples of polynomials in  $\mathcal{R}_{\leq k}^{\mathbf{G}}$  include the entries of the symmetric moment tensors

$$T_k(\theta) = \int_{\mathbf{G}} (g \cdot \theta)^{\otimes k} d\Lambda(g) \in \mathbb{R}^{d \times \dots \times d} \quad (2.6)$$

where  $T_k(\theta)$  is a tensor of order  $k$ . The entries of  $T_k(\theta)$  are the  $k^{\text{th}}$ -order mixed moments of the distribution of Gaussian mixture centers  $g \cdot \theta$ . Conversely, any  $\mathbf{G}$ -invariant polynomial  $p(\theta)$  of degree  $\leq k$  satisfies the identity

$$p(\theta) = \int_{\mathbf{G}} p(g \cdot \theta) d\Lambda(g),$$

and decomposing  $p$  on the right side into a sum of monomials shows that  $p(\theta)$  is an affine linear combination of entries of  $T_1, \dots, T_k$ . Hence  $\mathcal{R}_{\leq k}^{\mathbf{G}}$  is generated by  $T_1, \dots, T_k$ , and the subalgebra  $\mathcal{R}_{\leq k}^{\mathbf{G}}$  may be intuitively understood as containing all information in the moments of orders 1 to  $k$  for the Gaussian mixture defined by  $\theta$ .

For the projected model with projection  $\Pi$ , we define analogously the projected moment tensors

$$\tilde{T}_k(\theta) = \int_{\mathbf{G}} (\Pi \cdot g \cdot \theta)^{\otimes k} d\Lambda(g) \in \mathbb{R}^{\tilde{d} \times \dots \times \tilde{d}},$$

which are again the mixed moments of the Gaussian mixture centers  $\Pi \cdot g \cdot \theta$ . We then define

$$\tilde{\mathcal{R}}_{\leq k}^{\mathbf{G}} = \text{subalgebra of } \mathcal{R}^{\mathbf{G}} \text{ generated by the entries of } \tilde{T}_1, \dots, \tilde{T}_k.$$

Since each entry of  $\tilde{T}_k$  is a  $\mathbf{G}$ -invariant polynomial of degree  $k$ , we have  $\tilde{\mathcal{R}}_{\leq k}^{\mathbf{G}} \subseteq \mathcal{R}_{\leq k}^{\mathbf{G}}$ , but equality does not necessarily hold.

We denote by  $\langle \cdot, \cdot \rangle$  the Euclidean inner-product in the vectorization of these tensor spaces  $\mathbb{R}^{d \times \dots \times d}$  and  $\mathbb{R}^{\tilde{d} \times \dots \times \tilde{d}}$ , and by  $\|\cdot\|_{\text{HS}}^2$  the corresponding squared Euclidean norm. We will use the following general form

of the large- $\sigma$  series expansion of the population log-likelihood  $R(\theta)$ . We explain how the results of [KB20] yield this form in Appendix A.

**Theorem 2.2.** *Let  $G \subseteq O(d)$  be any compact subgroup. Fix any  $\theta_* \in \mathbb{R}^d$  and any integer  $K \geq 0$ .*

(a) *In the unprojected orbit recovery model,  $R(\theta)$  admits an expansion*

$$R(\theta) = C_0 + \sum_{k=1}^K \frac{1}{\sigma^{2k}} \left( s_k(\theta) + q_k(\theta) \right) + q(\theta). \quad (2.7)$$

Here  $C_0 \in \mathbb{R}$ ,  $q_k \in \mathcal{R}_{\leq k-1}^G$  is a polynomial of degree at most  $2k$ , and  $s_k \in \mathcal{R}_{\leq k}^G$  is the polynomial

$$s_k(\theta) = \frac{1}{2(k!)} \|T_k(\theta) - T_k(\theta_*)\|_{\text{HS}}^2. \quad (2.8)$$

The remainder  $q(\theta)$  is  $G$ -invariant and satisfies, for all  $\theta \in \mathbb{R}^d$  with  $\|\theta\| \leq \sigma$ ,

$$|q(\theta)| \leq \frac{C_K(1 \vee \|\theta\|)^{2K+2}}{\sigma^{2K+2}}, \quad \|\nabla q(\theta)\| \leq \frac{C_K(1 \vee \|\theta\|)^{2K+1}}{\sigma^{2K+2}}, \quad \|\nabla^2 q(\theta)\| \leq \frac{C_K(1 \vee \|\theta\|)^{2K}}{\sigma^{2K+2}}. \quad (2.9)$$

(b) *In the projected orbit recovery model,  $R(\theta)$  admits an expansion*

$$R(\theta) = C_0 + \sum_{k=1}^K \frac{1}{\sigma^{2k}} \left( \tilde{s}_k(\theta) + \langle \tilde{T}_k(\theta), P_k(\theta) \rangle + q_k(\theta) \right) + q(\theta). \quad (2.10)$$

Here  $C_0 \in \mathbb{R}$ ,  $q_k \in \tilde{\mathcal{R}}_{\leq k-1}^G$  is a polynomial of degree at most  $2k$ , all entries of  $P_k$  are polynomials of degree at most  $k$  belonging to  $\tilde{\mathcal{R}}_{\leq k-1}^G$ ,  $P_k$  satisfies  $P_k(\theta_*) = 0$ , and  $\tilde{s}_k \in \tilde{\mathcal{R}}_{\leq k}^G$  is the polynomial

$$\tilde{s}_k(\theta) = \frac{1}{2(k!)} \|\tilde{T}_k(\theta) - \tilde{T}_k(\theta_*)\|_{\text{HS}}^2. \quad (2.11)$$

The remainder  $q(\theta)$  is  $G$ -invariant and satisfies (2.9) for all  $\theta \in \mathbb{R}^d$  with  $\|\theta\| \leq \sigma$ .

The above constants  $C_0, C_K$ , the coefficients of the polynomials  $q_k(\theta)$  and  $P_k(\theta)$ , and the forms of the functions  $q(\theta)$  may all depend on  $\theta_*, G, d, \tilde{d}$ , and the projection  $\Pi$ .

The exact forms of  $q_k(\theta)$  and  $P_k(\theta)$  can be explicitly derived—see [FSWW20, Section 4.2] for these derivations in the unprojected setting—but we will not require them in what follows. Our arguments will only require the forms of the “leading” terms  $s_k(\theta)$  and  $\tilde{s}_k(\theta)$  defined in (2.8) and (2.11).

**2.3. Fisher information in high noise.** Consider the Fisher information matrix

$$I(\theta_*) = \nabla^2 R(\theta)|_{\theta=\theta_*}.$$

In this section, we characterize the eigenvalues and eigenvectors of  $I(\theta_*)$  for high noise and generic  $\theta_* \in \mathbb{R}^d$ . This generalizes [FSWW20, Theorem 4.14] for the unprojected model and a discrete group.

**Definition 2.3.** A subset  $S \subseteq \mathbb{R}^d$  is *generic* if  $\mathbb{R}^d \setminus S$  is contained in the zero set of some non-zero analytic function  $\psi : \mathbb{R}^d \rightarrow \mathbb{R}^k$ , for some  $k \geq 1$ .

If  $S \subseteq \mathbb{R}^d$  is generic, then  $\mathbb{R}^d \setminus S$  has zero Lebesgue measure [Mit20]. We say that a statement holds for generic  $\theta_* \in \mathbb{R}^d$  if it holds for all  $\theta_*$  in some generic subset of  $\mathbb{R}^d$ .

Our characterization of  $I(\theta_*)$  is in terms of the number of distinct “degrees-of-freedom” captured by the moments of the Gaussian mixture model up to each order  $k$ . This is formalized by the notion of the transcendence degrees of the subalgebras  $\mathcal{R}_{\leq k}^G$  and  $\tilde{\mathcal{R}}_{\leq k}^G$ .

**Definition 2.4.** Polynomials  $p_1, \dots, p_k : \mathbb{R}^d \rightarrow \mathbb{R}$  are *algebraically independent* (over  $\mathbb{R}$ ) if there is no non-zero polynomial  $q : \mathbb{R}^k \rightarrow \mathbb{R}$  such that  $q(p_1(\theta), \dots, p_k(\theta))$  is identically 0 for all  $\theta \in \mathbb{R}^d$ .

For any  $\mathcal{A} \subseteq \mathcal{R}^G$ , its *transcendence degree*  $\text{trdeg}(\mathcal{A})$  is the maximum cardinality of any algebraically independent subset  $A \subseteq \mathcal{A}$ . Any maximal such subset  $A \subseteq \mathcal{A}$  is a *transcendence basis* for  $\mathcal{A}$ .

Geometrically, by the Jacobian criterion for algebraic independence (c.f. Lemma A.2), the transcendence degree coincides with the maximum number of linearly independent gradient vectors of the polynomials in  $\mathcal{A}$ , evaluated at any generic point  $\theta \in \mathbb{R}^d$ .

As a simple example, if  $\mathbf{G}$  is the symmetric group of all permutations of  $d$  coordinates, then  $\mathcal{R}^{\mathbf{G}}$  is the algebra of all symmetric polynomials in  $d$  variables. Each subalgebra  $\mathcal{R}_{\leq k}^{\mathbf{G}}$  for  $k \leq d$  has transcendence degree exactly equal to  $k$ , and one choice of a transcendence basis for  $\mathcal{R}_{\leq k}^{\mathbf{G}}$  is the set of symmetric power sums  $\{\theta_1^j + \dots + \theta_d^j : j = 1, \dots, k\}$ .

For the full invariant algebra  $\mathcal{R}^{\mathbf{G}}$ , if  $\mathbf{G} \subset \mathbf{O}(d)$  is any discrete subgroup as studied in [FSWW20], then  $\text{trdeg}(\mathcal{R}^{\mathbf{G}}) = d$ . More generally, we have the following.

**Proposition 2.5.** *Let  $\mathbf{G}$  be a compact subgroup of  $\mathbf{O}(d)$ . Then*

$$\text{trdeg}(\mathcal{R}^{\mathbf{G}}) = d - \max_{\theta \in \mathbb{R}^d} \dim(\mathcal{O}_{\theta})$$

where  $\dim(\mathcal{O}_{\theta})$  is the dimension of the orbit  $\mathcal{O}_{\theta}$  as a submanifold of  $\mathbb{R}^d$ . Here, the maximum orbit dimension  $\max_{\theta \in \mathbb{R}^d} \dim(\mathcal{O}_{\theta})$  is also the orbit dimension for generic points  $\theta \in \mathbb{R}^d$ .

We will mostly consider group actions where this generic orbit dimension equals the group dimension  $\dim(\mathbf{G})$ , so that  $\text{trdeg}(\mathcal{R}^{\mathbf{G}}) = d - \dim(\mathbf{G})$ . In particular, for the function estimation examples to be discussed in Sections 3 and 4, we will have  $\text{trdeg}(\mathcal{R}^{\mathbf{G}}) = d - 1$  for an action of  $\mathbf{G}$  that is isomorphic to  $\mathbf{SO}(2)$ , and  $\text{trdeg}(\mathcal{R}^{\mathbf{G}}) = d - 3$  for an action of  $\mathbf{G}$  that is isomorphic to  $\mathbf{SO}(3)$ .

It was shown in [BBSK<sup>+</sup>17, Theorem 4.9], for generic signals  $\theta_* \in \mathbb{R}^d$ , that the values of the moment tensors  $T_1(\theta_*), \dots, T_k(\theta_*)$  are sufficient to identify  $\theta_*$  up to a finite list of possible orbits if and only if  $\text{trdeg}(\mathcal{R}_{\leq k}^{\mathbf{G}}) = \text{trdeg}(\mathcal{R}^{\mathbf{G}})$ . More informally, the order of moments needed to “locally” identify the orbit of  $\theta_*$  coincides with the order of moments needed to capture all  $\text{trdeg}(\mathcal{R}^{\mathbf{G}})$  degrees-of-freedom of the invariant algebra. Throughout this paper, we will denote this number as  $K$  in the unprojected model and as  $\tilde{K}$  in the projected model, which are well-defined by the following proposition. We defer proofs of Propositions 2.5 and 2.6 to Appendix A.

**Proposition 2.6.** *For any compact subgroup  $\mathbf{G} \subseteq \mathbf{O}(d)$ ,*

- (a) *There is a smallest integer  $K < \infty$  for which  $\text{trdeg}(\mathcal{R}_{\leq K}^{\mathbf{G}}) = \text{trdeg}(\mathcal{R}^{\mathbf{G}})$ .*
- (b)  *$\Pi$  satisfies (2.4) if and only if there is a smallest integer  $\tilde{K} < \infty$  for which  $\text{trdeg}(\tilde{\mathcal{R}}_{\leq \tilde{K}}^{\mathbf{G}}) = \text{trdeg}(\mathcal{R}^{\mathbf{G}})$ .*

In the unprojected model, let us now denote

$$d_0 = \max_{\theta \in \mathbb{R}^d} \dim(\mathcal{O}_{\theta}), \quad d_k = \text{trdeg} \mathcal{R}_{\leq k}^{\mathbf{G}} - \text{trdeg} \mathcal{R}_{\leq k-1}^{\mathbf{G}} \text{ for } k = 1, \dots, K \quad (2.12)$$

to decompose the total dimension of  $\theta_*$  as  $d = d_0 + d_1 + \dots + d_K$ . In the projected model, assuming the condition (2.4), let us similarly denote

$$\tilde{d}_0 = \max_{\theta \in \mathbb{R}^d} \dim(\mathcal{O}_{\theta}), \quad \tilde{d}_k = \text{trdeg} \tilde{\mathcal{R}}_{\leq k}^{\mathbf{G}} - \text{trdeg} \tilde{\mathcal{R}}_{\leq k-1}^{\mathbf{G}} \text{ for } k = 1, \dots, \tilde{K} \quad (2.13)$$

to decompose the total dimension as  $d = \tilde{d}_0 + \tilde{d}_1 + \dots + \tilde{d}_{\tilde{K}}$ . The following result expresses the spectral properties of the Fisher information matrix in terms of these decompositions.

**Theorem 2.7.** *For generic  $\theta_* \in \mathbb{R}^d$ , some  $(\theta_*, \mathbf{G}, \Pi)$ -dependent constants  $\sigma_0, C, c > 0$  and function  $\varepsilon(\sigma)$  satisfying  $\varepsilon(\sigma) \rightarrow 0$  as  $\sigma \rightarrow \infty$ , and all  $\sigma > \sigma_0$ :*

- (a) *In the unprojected orbit recovery model,*
  1. *The Fisher information matrix  $I(\theta_*)$  has rank exactly  $\text{trdeg}(\mathcal{R}^{\mathbf{G}}) = d - d_0$ . Defining  $K$  by Proposition 2.6(a), for each  $k = 1, \dots, K$ ,*

$$\text{exactly } d_k \text{ eigenvalues of } I(\theta_*) \text{ belong to } [c\sigma^{-2k}, C\sigma^{-2k}].$$

2. *For each  $k = 1, \dots, K$ , let  $V_k$  be the subspace spanned by the leading  $d_1 + \dots + d_k$  eigenvectors of  $I(\theta_*)$ , and let  $W_k$  be the subspace spanned by the gradient vectors  $\{\nabla p(\theta_*) : p \in \mathcal{R}_{\leq k}^{\mathbf{G}}\}$ . Then the sin-theta distance between  $V_k$  and  $W_k$  is bounded as*

$$\|\sin \Theta(V_k, W_k)\| < \varepsilon(\sigma).$$

3. For any  $k = 1, \dots, K$  and any polynomial  $p \in \mathcal{R}_{\leq k}^G$ , the gradient  $\nabla p(\theta_*) \in \mathbb{R}^d$  is orthogonal to the null space of  $I(\theta_*)$  and satisfies

$$\nabla p(\theta_*)^\top I(\theta_*)^\dagger \nabla p(\theta_*) \leq C\sigma^{2k}$$

where  $I(\theta_*)^\dagger$  is the Moore-Penrose pseudo-inverse.

- (b) In the projected orbit recovery model satisfying condition (2.4), the same statements hold with  $\mathcal{R}_{\leq k}^G$ ,  $K$ , and  $d_k$  replaced by  $\tilde{\mathcal{R}}_{\leq k}^G$ ,  $\tilde{K}$ , and  $\tilde{d}_k$ , where  $\tilde{K}$  is defined by Proposition 2.6(b).

**Remark.** Theorem 2.7(a1) states that  $I(\theta_*)$  has eigenvalues on differing scales of  $\sigma^{-2}$  in high noise, with  $d_k$  such eigenvalues scaling as  $\sigma^{-2k}$ , and  $d_0 = \dim(\mathcal{O}_{\theta_*})$  eigenvalues of 0 representing the non-identifiable degrees-of-freedom tangent to  $\mathcal{O}_{\theta_*}$ . Thus there are  $d_k$  degrees-of-freedom in  $\theta_*$  that are estimated with asymptotic variance  $O(\sigma^{2k}/n)$  by the MLE. The largest such variance is  $O(\sigma^{2K}/n)$ , which is in accordance with results about list-recovery of generic signals in [BBSK<sup>+</sup>17] and with the  $\sigma^6$  sample complexity established in [PWB<sup>+</sup>19] for multi-reference alignment, where  $K = 3$ .

**Remark.** Theorem 2.7(a2) describes also the associated spaces of eigenvectors of  $I(\theta_*)$ , where the eigenspaces corresponding to eigenvalues at scales  $\sigma^{-2}, \dots, \sigma^{-2k}$  coincide approximately with the span of the gradients of  $G$ -invariant polynomials up to degree  $k$ . Theorem 2.7(a3) then implies that the functional  $p(\theta_*)$  for any  $p \in \mathcal{R}_{\leq k}^G$  is estimated by the plug-in MLE  $p(\hat{\theta}_n)$  with asymptotic variance  $O(\sigma^{2k}/n)$ . Similar statements hold for projected models by Theorem 2.7(b).

The following result connects the above sequences of transcendence degrees and gradients  $\{\nabla p(\theta_*) : p \in \mathcal{R}_{\leq k}^G\}$  to the terms  $s_k(\theta)$  and  $\tilde{s}_k(\theta)$  in the series expansions of  $R(\theta)$  in Theorem 2.2. We will use this to deduce the values of these transcendence degrees for the function estimation examples of Sections 3 and 4.

**Lemma 2.8.** (a) In the unprojected orbit recovery model, let  $s_k(\theta)$  be defined by (2.8). Then each matrix  $\nabla^2 s_k(\theta)|_{\theta=\theta_*}$  is positive semidefinite. For any  $k \geq 1$ , at generic  $\theta_* \in \mathbb{R}^d$ ,

$$\text{trdeg}(\mathcal{R}_{\leq k}^G) = \text{rank} \left( \nabla^2 s_1(\theta) + \dots + \nabla^2 s_k(\theta) \Big|_{\theta=\theta_*} \right), \quad (2.14)$$

and the span of  $\{\nabla p(\theta_*) : p \in \mathcal{R}_{\leq k}^G\}$  is the column span of  $\nabla^2 s_1(\theta) + \dots + \nabla^2 s_k(\theta)|_{\theta=\theta_*}$ .

- (b) In the projected orbit recovery model, the same holds for  $\tilde{\mathcal{R}}_{\leq k}^G$  and  $\tilde{s}_k(\theta)$  as defined by (2.11).

**Remark 2.9.** We restrict attention to generic signals  $\theta_* \in \mathbb{R}^d$  in this work. The specific condition for  $\theta_*$  that we use in Theorem 2.7, and in Theorems 2.11 and 2.13 to follow, is that the gradient vectors  $\{\nabla p(\theta_*) : p \in \mathcal{R}_{\leq K}^G\}$  or  $\{\nabla p(\theta_*) : p \in \tilde{\mathcal{R}}_{\leq \tilde{K}}^G\}$  span a subspace of dimension  $\text{trdeg}(\mathcal{R}^G)$  or  $\text{trdeg}(\tilde{\mathcal{R}}^G)$ , respectively.

Different behavior may be observed for non-generic signals: For  $G = \{+ \text{Id}, - \text{Id}\}$ , which has been studied in [XHM16, WZ19], the Fisher information  $I(\theta_*)$  is singular at  $\theta_* = 0$  (even though  $d_0 = 0$ , as the group is discrete). This leads to a  $n^{-1/4}$  rate of estimation error near  $\theta_* = 0$ , instead of the  $n^{-1/2}$  parametric rate. This  $n^{-1/4}$  rate holds more generally for any discrete group  $G$  at signals  $\theta_*$  whose orbit points are not pairwise distinct, which are precisely those signals where the Fisher information  $I(\theta_*)$  is singular [Bru19].

A different distinction between generic and non-generic signals was highlighted in [PWB<sup>+</sup>19] when  $G$  is the group of cyclic rotations of coordinates in  $\mathbb{R}^d$ . There, orbits of generic signals are uniquely identified by moments up to the order  $K = 3$ , but identification of non-generic signals having zero power in certain Fourier frequencies may require moments up to the order  $d - 1$ . For such non-generic signals, we expect  $I(\theta_*)$  to be non-singular and the MLE to attain the parametric rate, but with asymptotic variance scaling as  $\sigma^{2(d-1)}/n$  rather than  $\sigma^{2K}/n = \sigma^6/n$ . In a related model of continuous MRA, this asymptotic scaling is implied by the results of [BNWR20].

**2.4. Global likelihood landscape.** In this section, we establish correspondences between global and local minimizers of the population negative log-likelihood  $R(\theta)$  with those of a sequence of moment optimization problems. These results are similar to results of [FSWW20, Sections 4.3 and 4.5] for discrete groups  $G$ , with a distinction that when  $G$  is continuous, these minimizers are not isolated points but rather manifolds of positive dimension.

We recall the following structural property for smooth non-convex optimization landscapes, under which convergence to the global optimum from a random initialization is guaranteed for various descent-based optimization algorithms [GHJY15, LSJR16, JGN<sup>+</sup>17].

**Definition 2.10.** The problem of minimizing a twice-continuously differentiable function  $f : \mathcal{V} \rightarrow \mathbb{R}$  over a smooth manifold  $\mathcal{V}$  is *globally benign* if each point  $x \in \mathcal{V}$  where  $\nabla f(x)|_{\mathcal{V}} = 0$  is either a global minimizer of  $f$  over  $\mathcal{V}$ , or has a direction of strictly negative curvature,  $\lambda_{\min}(\nabla^2 f(x)|_{\mathcal{V}}) < 0$ .

Here  $\nabla f(x)|_{\mathcal{V}}$  and  $\nabla^2 f(x)|_{\mathcal{V}}$  denote the gradient and Hessian of  $f$  on  $\mathcal{V}$ , which may be taken in any choice of a smooth local chart around  $x \in \mathcal{V}$ .

Minimizing  $R(\theta)$  in high noise may be viewed as successively solving a sequence of moment optimizations defined by the terms of its expansion in Theorem 2.2. To ease notation, let us collect the vectorized moment tensors up to order  $k$  as

$$M_k(\theta) = \text{vec} \left( T_1(\theta), \dots, T_k(\theta) \right) \in \mathbb{R}^{d+d^2+\dots+d^k}, \quad (2.15)$$

$$\widetilde{M}_k(\theta) = \text{vec} \left( \widetilde{T}_1(\theta), \dots, \widetilde{T}_k(\theta) \right) \in \mathbb{R}^{\widetilde{d}+\widetilde{d}^2+\dots+\widetilde{d}^k}. \quad (2.16)$$

Fixing the true signal  $\theta_* \in \mathbb{R}^d$ , we define the moment varieties

$$\mathcal{V}_k(\theta_*) = \left\{ \theta \in \mathbb{R}^d : M_k(\theta) = M_k(\theta_*) \right\}, \quad \mathcal{V}_0(\theta_*) = \mathbb{R}^d, \quad (2.17)$$

$$\widetilde{\mathcal{V}}_k(\theta_*) = \left\{ \theta \in \mathbb{R}^d : \widetilde{M}_k(\theta) = \widetilde{M}_k(\theta_*) \right\}, \quad \widetilde{\mathcal{V}}_0(\theta_*) = \mathbb{R}^d. \quad (2.18)$$

These are the points  $\theta \in \mathbb{R}^d$  for which the mixed moments of the Gaussian mixture model defined by  $\theta$  match those of the true signal  $\theta_*$  up to order  $k$ .

We state a general result on the optimization landscape, assuming that the Jacobian matrices  $dM_k$  and  $d\widetilde{M}_k$  have constant rank over  $\mathcal{V}_k(\theta_*)$  and  $\widetilde{\mathcal{V}}_k(\theta_*)$ , so that  $\mathcal{V}_k(\theta_*)$  and  $\widetilde{\mathcal{V}}_k(\theta_*)$  are smooth manifolds. Then, recalling  $s_k(\theta)$  and  $\tilde{s}_k(\theta)$  from (2.8) and (2.11), we consider the optimization problem

$$\text{minimize } s_k(\theta) \text{ over } \theta \in \mathcal{V}_{k-1}(\theta_*) \quad (2.19)$$

in the unprojected setting, and

$$\text{minimize } \tilde{s}_k(\theta) \text{ over } \theta \in \widetilde{\mathcal{V}}_{k-1}(\theta_*) \quad (2.20)$$

in the projected setting. These are polynomial optimization problems in  $\theta$  that are defined independently of the noise level  $\sigma^2$ . The following theorem guarantees that the landscape of  $R(\theta)$  is globally benign in high noise, as long as the landscape of each problem (2.19) or (2.20) is globally benign, and the final moment variety  $\mathcal{V}_K(\theta_*)$  or  $\widetilde{\mathcal{V}}_{\widetilde{K}}(\theta_*)$  contains only the points which globally minimize  $R(\theta)$ . We illustrate part (a) of this result using a simple example of orthogonal Procrustes alignment at the conclusion of this section.

**Theorem 2.11.** *For generic  $\theta_* \in \mathbb{R}^d$ :*

- (a) *In the unprojected model, define  $K$  by Proposition 2.6(a). Suppose that  $\mathcal{V}_K(\theta_*) = \mathcal{O}_{\theta_*}$ . Suppose also that for each  $k = 1, \dots, K$ , the derivative matrix  $dM_k(\theta)$  has constant rank over  $\mathcal{V}_k(\theta_*)$ , and the minimization of  $s_k(\theta)$  over  $\mathcal{V}_{k-1}(\theta_*)$  is globally benign. Then for some  $\sigma_0 \equiv \sigma_0(\theta_*, \mathbf{G})$  and any  $\sigma > \sigma_0$ , the minimization of  $R(\theta)$  is also globally benign.*
- (b) *In the projected model satisfying (2.4), define  $\widetilde{K}$  by Proposition 2.6(b). Suppose that  $\widetilde{\mathcal{V}}_{\widetilde{K}}(\theta_*) = \{\theta : \Pi(\mathcal{O}_{\theta}) \equiv \Pi(\mathcal{O}_{\theta_*})\}$ . Suppose also that for each  $k = 1, \dots, \widetilde{K}$ , the derivative matrix  $d\widetilde{M}_k(\theta)$  has constant rank over  $\widetilde{\mathcal{V}}_k(\theta_*)$ , and the minimization of  $\tilde{s}_k(\theta)$  over  $\widetilde{\mathcal{V}}_{k-1}(\theta_*)$  is globally benign. Then for any constant  $B > 0$ , some  $\sigma_0 \equiv \sigma_0(\theta_*, \mathbf{G}, \Pi, B)$ , and any  $\sigma > \sigma_0$ , the minimization of  $R(\theta)$  is globally benign over the domain  $\{\theta \in \mathbb{R}^d : \|\theta\| < B(\|\theta_*\| + \sigma)\}$ .*

In Theorem 2.11(b), we have restricted to a ball  $\{\theta \in \mathbb{R}^d : \|\theta\| < B(\|\theta_*\| + \sigma)\}$ , as the landscape of  $R(\theta)$  outside this ball may depend on the specific interaction between  $\mathbf{G}$  and  $\Pi$ . In practice, such a bound for  $\|\theta\|$  may be known a priori, so that optimization may indeed be restricted to this ball. (In unprojected models, we show that  $R(\theta)$  cannot have critical points outside this ball for any group  $\mathbf{G}$ , allowing us to remove such a restriction in part (a).)

Whether the conditions of Theorem 2.11 hold depends on the specific model, and both positive and negative examples for discrete groups were exhibited in [FSWW20]. In models where they do not hold,  $R(\theta)$  may in fact have spurious local minimizers in high noise, and Theorem 2.2 can be used to further establish a correspondence between the local minimizers of  $R(\theta)$  and those of the above moment optimizations. We formalize one such result—not fully general, but sufficient to study many examples of interest—as follows.

**Definition 2.12.** Suppose  $\mathcal{V}_{K-1}(\theta_*)$  is a smooth manifold. A critical point  $\theta$  of  $s_K(\theta)|_{\mathcal{V}_{K-1}(\theta_*)}$  is *non-degenerate up to orbit* if  $\mathcal{O}_\theta$  is a smooth manifold of dimension  $d_0$  in a local neighborhood of  $\theta$ , and

$$\text{rank}(\nabla^2 s_K(\theta)|_{\mathcal{V}_{K-1}(\theta_*)}) = \dim(\mathcal{V}_{K-1}(\theta_*)) - d_0.$$

Note that  $\nabla^2 s_K(\theta)|_{\mathcal{V}_{K-1}(\theta_*)}$  is a symmetric matrix of dimension  $\dim(\mathcal{V}_{K-1}(\theta_*))$ . For any critical point  $\theta$  of  $s_K|_{\mathcal{V}_{K-1}(\theta_*)}$ , the null space of this Hessian must contain the tangent space to  $\mathcal{O}_\theta$ , and Definition 2.12 ensures that this Hessian has no further rank degeneracy.

**Theorem 2.13.** For generic  $\theta_* \in \mathbb{R}^d$ :

- (a) In the unprojected model, suppose that  $dM_k(\theta)$  has constant rank over  $\mathcal{V}_k(\theta_*)$ , and the minimization of  $s_k(\theta)$  over  $\mathcal{V}_{k-1}(\theta_*)$  is globally benign for each  $k = 1, \dots, K-1$ . Then for some  $(\theta_*, \mathbf{G})$ -dependent constant  $\sigma_0 > 0$  and function  $\varepsilon(\sigma)$  satisfying  $\varepsilon(\sigma) \rightarrow 0$  as  $\sigma \rightarrow \infty$ , and for all  $\sigma > \sigma_0$ :
1. Let  $\theta_+$  be any local minimizer of  $s_K(\theta)$  over  $\mathcal{V}_{K-1}(\theta_*)$  that is non-degenerate up to orbit. Then there exists a local minimizer  $\theta'_+$  of  $R(\theta)$  where  $\|\theta_+ - \theta'_+\| < \varepsilon(\sigma)$ .
  2. Conversely, suppose that all critical points of  $s_K(\theta)$  over  $\mathcal{V}_{K-1}(\theta_*)$  are non-degenerate up to orbit. Let  $\theta_+$  be any local minimizer of  $R(\theta)$ . Then there exists a local minimizer  $\theta'_+$  of  $s_K(\theta)$  over  $\mathcal{V}_{K-1}(\theta_*)$  where  $\|\theta_+ - \theta'_+\| < \varepsilon(\sigma)$ .
- (b) In the projected model satisfying (2.4), statement (1.) holds with  $K$ ,  $M_k$ ,  $\mathcal{V}_k$ , and  $s_k$  replaced by  $\tilde{K}$ ,  $\tilde{M}_k$ ,  $\tilde{\mathcal{V}}_k$ , and  $\tilde{s}_k$ , where  $\sigma_0$  and  $\varepsilon(\sigma)$  may depend also on the projection  $\Pi$ . Statement (2.) holds for local minimizers  $\theta_+$  of  $R(\theta)$  satisfying  $\|\theta_+\| < B(\|\theta_*\| + \sigma)$  for any constant  $B > 0$ , where  $\sigma_0$  and  $\varepsilon(\sigma)$  may depend also on  $\Pi$  and  $B$ .

The guarantees of Theorems 2.11 and 2.13 may be translated to the sample log-likelihood  $R_n(\theta)$  by establishing concentration of  $\nabla R_n(\theta)$  and  $\nabla^2 R_n(\theta)$  around  $\nabla R(\theta)$  and  $\nabla^2 R(\theta)$  [MBM18]. For orbit recovery models in the high-noise regime, we believe that it may be possible to obtain sharp concentration bounds by deriving a series expansion also of the empirical log-likelihood function  $R_n(\theta)$  in powers of  $\sigma^{-1}$ , and analyzing the concentration term-by-term. Some results of this form were obtained for models without linear projection in [FSWW20, Lemma 4.11 and Corollary 4.18], and we leave the analysis of the empirical log-likelihood function and landscape for more general models as an open problem for future work.

**Example 2.14** (Landscape of orthogonal Procrustes alignment). We illustrate Theorems 2.11 and 2.13 using a simple example of *orthogonal Procrustes alignment* [Gow75, Goo91, PSB19].

In this problem, samples of an object consisting of  $m \geq 3$  atoms in  $\mathbb{R}^3$  are observed under random orthogonal rotations and reflections. We represent the object as  $\theta_* \in \mathbb{R}^{3 \times m} \cong \mathbb{R}^d$  where  $d = 3m$ . The rotational group is  $\mathbf{G} = \mathbf{O}(3) \otimes \text{Id}_m \subset \mathbf{O}(d)$ , where a common orthogonal matrix in 3-dimensions is applied to all  $m$  atoms. Assuming the generic condition that  $\text{rank}(\theta_*) = 3$ , i.e. these  $m$  atoms do not lie on a common 2-dimensional subspace, we study the likelihood landscape for estimating  $\theta_*$  from many independently rotated samples.

In this model, we check in Appendix B that  $K = 2$ ,  $(d_0, d_1, d_2) = (3, 0, d - 3)$ ,  $\mathcal{V}_1(\theta_*) = \mathbb{R}^d$ , and  $\mathcal{V}_2(\theta_*) = \{g \cdot \theta_* : g \in \mathbf{G}\} = \mathcal{O}_{\theta_*}$ . The first two moment tensors  $T_1(\theta)$  and  $T_2(\theta)$  are given by  $T_1(\theta) = 0$  and  $T_2(\theta) = \frac{1}{3} \text{Id}_{3 \times 3} \otimes (\theta^\top \theta) \in \mathbb{R}^{d \times d}$ , and the terms  $s_1(\theta)$  and  $s_2(\theta)$  in (2.7) are given by  $s_1(\theta) = 0$  and

$$s_2(\theta) = \frac{1}{12} \|\theta^\top \theta - \theta_*^\top \theta_*\|_{\text{HS}}^2,$$

where  $\theta^\top \theta, \theta_*^\top \theta_* \in \mathbb{R}^{m \times m}$ . The minimization of  $s_1(\theta)$  over  $\mathcal{V}_0(\theta_*) = \mathbb{R}^d$  is trivially globally benign. We show in Appendix B that  $dM_2(\theta)$  has constant rank over  $\mathcal{V}_2(\theta_*)$ , and that the minimization of  $s_2(\theta)$  over  $\mathcal{V}_1(\theta_*) = \mathbb{R}^d$  is also globally benign, with minimizers given exactly by  $\mathcal{V}_2(\theta_*) = \mathcal{O}_{\theta_*}$ . Thus, Theorem 2.11(a) implies that the landscape of  $R(\theta)$  is also globally benign for sufficiently high noise, and the only local minimizers of  $R(\theta)$  are rotations and reflections of the true object.

A variation of this problem is the rotation-only variant, where we observe 3-dimensional rotations (but not reflections) of the object. Then the rotational group is instead  $\mathbf{G} = \text{SO}(3) \otimes \text{Id}_m \subset \mathbf{O}(d)$ . We show in Appendix B that still  $K = 2$ ,  $(d_0, d_1, d_2) = (3, 0, d - 3)$ , and the forms of  $T_1(\theta), T_2(\theta), \mathcal{V}_1(\theta_*), \mathcal{V}_2(\theta_*), s_1(\theta), s_2(\theta)$  are identical to the above (even though the full log-likelihood  $R(\theta)$  is not). Thus the minimization of  $s_2(\theta)$  over  $\mathcal{V}_1(\theta_*) = \mathbb{R}^d$  is still globally benign, with minimizers  $\mathcal{V}_2(\theta_*)$ . However, this set of minimizers is now written as

$$\mathcal{V}_2(\theta_*) = \{g \cdot \theta_* : g \in \mathbf{G}\} \cup \{-g \cdot \theta_* : g \in \mathbf{G}\} = \mathcal{O}_{\theta_*} \cup \mathcal{O}_{-\theta_*}$$

constituting two distinct orbits under this more restrictive group action. The first orbit  $\mathcal{O}_{\theta_*}$  are the global minimizers of  $R(\theta)$ . The second orbit corresponds to the mirror reflection  $-\theta_*$ , which does not globally minimize  $R(\theta)$ , but the difference between  $R(\theta_*)$  and  $R(-\theta_*)$  lies in the remainder term of the expansion (2.7). Theorem 2.13(a) shows that for high noise,  $R(\theta)$  will have spurious local minimizers near (but not exactly equal to) this second orbit  $\mathcal{O}_{-\theta_*}$ .

### 3. CONTINUOUS MULTI-REFERENCE ALIGNMENT

We now specialize the preceding general results to the problem of estimating a periodic function on the circle, observed under  $\text{SO}(2)$  rotations of its domain. We will refer to this as the *continuous MRA* model. We study the unprojected model in this section, and a version with a two-fold projection in Appendix C.3. These provide simpler 1-dimensional analogues of the 2-dimensional and 3-dimensional problems that we will discuss in Section 4.

To describe the model, let  $f : \mathcal{S}^1 \rightarrow \mathbb{R}$  be a periodic function on the unit circle  $\mathcal{S}^1 \cong [0, 1)$ . We identify the rotational group  $\text{SO}(2)$  also with  $[0, 1)$ , and represent the rotation of  $f$  by an element  $\mathbf{g} \in \text{SO}(2) \cong [0, 1)$  as  $f_{\mathbf{g}}(t) = f(t + \mathbf{g} \bmod 1)$ . Each sample is an observation of the rotated function  $f_{\mathbf{g}}$  with additive white noise,

$$f_{\mathbf{g}}(t)dt + \sigma dW(t)$$

where  $\mathbf{g} \sim \text{Unif}([0, 1))$  and  $dW(t)$  denotes a standard Gaussian white noise process on  $\mathcal{S}^1$ . This may be understood as observing a realization of the Gaussian process  $\{F(h)\}_{h \in L_2(\mathcal{S}^1)} = \{\int h(t)[f_{\mathbf{g}}(t)dt + \sigma dW(t)]\}_{h \in L_2(\mathcal{S}^1)}$  with mean and covariance functions

$$\mathbb{E}[F(h)] = \int_0^1 h(t)f_{\mathbf{g}}(t)dt, \quad \text{Cov}[F(h_1), F(h_2)] = \sigma^2 \int_0^1 h_1(t)h_2(t)dt, \quad (3.1)$$

or equivalently as observing all coefficients of  $f_{\mathbf{g}}$  in a complete orthonormal basis  $\{h_j(t)\}_{j=1}^\infty$  of  $L_2(\mathcal{S}^1)$  with independent  $\mathcal{N}(0, \sigma^2)$  noise for each basis coefficient.

We consider the real Fourier basis on  $\mathcal{S}^1$ , given by

$$h_0(t) = 1, \quad h_{l1}(t) = \sqrt{2} \cos 2\pi lt, \quad h_{l2}(t) = \sqrt{2} \sin 2\pi lt \quad \text{for } l = 1, 2, 3, \dots \quad (3.2)$$

We then restrict our model to the finite-dimensional space of functions  $f : \mathcal{S}^1 \rightarrow \mathbb{R}$  that have finite bandlimit  $L \geq 1$  in this basis, i.e.  $f$  admits a representation

$$f(t) = \theta^{(0)}h_0(t) + \sum_{l=1}^L \theta_1^{(l)}h_{l1}(t) + \sum_{l=1}^L \theta_2^{(l)}h_{l2}(t). \quad (3.3)$$

Importantly, the space of such bandlimited functions is closed under rotations of  $\mathcal{S}^1$ . Writing

$$\theta = (\theta^{(0)}, \theta_1^{(1)}, \theta_2^{(1)}, \dots, \theta_1^{(L)}, \theta_2^{(L)}) \in \mathbb{R}^d, \quad d = 2L + 1$$

for the vector of Fourier coefficients, the rotation  $f \mapsto f_{\mathbf{g}}$  corresponds to  $\theta \mapsto g \cdot \theta$ , where  $g$  belongs to the block-diagonal representation

$$G = \left\{ \text{diag} \left( 1, \begin{pmatrix} \cos 2\pi \mathbf{g} & \sin 2\pi \mathbf{g} \\ -\sin 2\pi \mathbf{g} & \cos 2\pi \mathbf{g} \end{pmatrix}, \dots, \begin{pmatrix} \cos 2\pi L \mathbf{g} & \sin 2\pi L \mathbf{g} \\ -\sin 2\pi L \mathbf{g} & \cos 2\pi L \mathbf{g} \end{pmatrix} \right) : \mathbf{g} \in [0, 1) \right\} \quad (3.4)$$

of  $\text{SO}(2)$ . The observation model for the Fourier coefficients of  $f$  then takes the form of (2.1), where we observe coordinates of  $g \cdot \theta$  with entrywise i.i.d.  $\mathcal{N}(0, \sigma^2)$  noise.

Theorem 3.1 below first characterizes, for this model, the decomposition of total dimension described in Theorem 2.7. As a direct consequence of this result, we state Corollary 3.2 which summarizes the implications for identifying  $\theta_*$  based on its low-order moments, and for the spectral structure of the Fisher information matrix  $I(\theta_*)$ .

**Theorem 3.1.** *For any  $L \geq 1$ , we have*

$$\text{trdeg}(\mathcal{R}_{\leq 1}^G) = 1, \quad \text{trdeg}(\mathcal{R}_{\leq 2}^G) = L + 1, \quad \text{trdeg}(\mathcal{R}_{\leq 3}^G) = \text{trdeg}(\mathcal{R}^G) = 2L = d - 1.$$

**Corollary 3.2.** *A generic signal  $\theta_* \in \mathbb{R}^d$  in this continuous MRA model has the following properties:*

- (a)  $\theta_*$  is identified up to a finite list of orbits by the moments of  $g \cdot \theta_*$  up to order  $K = 3$  when  $L \geq 2$ , and order  $K = 2$  when  $L = 1$ .

- (b) For  $(\theta_*, \mathbf{G})$ -dependent constants  $C, c > 0$  independent of  $\sigma$ , the Fisher information  $I(\theta_*)$  has  $d_0 = 1$  eigenvalue of 0 and  $d_k$  eigenvalues in  $[c\sigma^{-2k}, C\sigma^{-2k}]$  for  $k = 1, 2, 3$  and  $(d_1, d_2, d_3) = (1, L, L - 1)$ .

Part (a) of this corollary follows immediately from [BBSK<sup>+</sup>17, Theorem 4.9] (which we review in Appendix A.2), and part (b) follows from Theorem 2.7.

In Appendix C.1, we provide a proof of Theorem 3.1 using our general result of Lemma 2.8, as a warm-up for our analyses of the  $\text{SO}(3)$ -rotational models to follow. We note that for a similar observation model of continuous MRA studied in [BNWR20], a stronger form of Corollary 3.2(a) is already known, namely that 3<sup>rd</sup>-order moments are sufficient to identify generic signals  $\theta_* \in \mathbb{R}^d$  up to a single unique orbit.

Next, we study the moment optimization problems of (2.19), and we describe more explicit forms for these optimization problems in this continuous MRA example. Denote the Fourier coefficients of the true function  $f$  by  $\theta_* \in \mathbb{R}^d$ . Define the complex Fourier coefficients

$$u^{(0)}(\theta) = \theta^{(0)} \in \mathbb{R}, \quad u^{(l)}(\theta) = \theta_1^{(l)} + \mathbf{i}\theta_2^{(l)} = r_l(\theta)e^{\mathbf{i}\lambda_l(\theta)} \in \mathbb{C},$$

where  $(r_l(\theta), \lambda_l(\theta))$  for  $l \geq 1$  are the magnitude and phase of  $u^{(l)}(\theta)$ . Write as shorthand

$$r_{l,l',l''}(\theta) = r_l(\theta)r_{l'}(\theta)r_{l''}(\theta), \quad \lambda_{l,l',l''}(\theta) = \lambda_l(\theta) - \lambda_{l'}(\theta) - \lambda_{l''}(\theta).$$

Here  $\lambda_{l,l',l''}(\theta)$  are the elements of the Fourier bispectrum of  $\theta$ .

**Theorem 3.3.** *For any  $L \geq 1$ ,*

$$\begin{aligned} s_1(\theta) &= \frac{1}{2} \left( \theta^{(0)} - \theta_*^{(0)} \right)^2 \\ s_2(\theta) &= \frac{1}{4} \left( (\theta^{(0)})^2 - (\theta_*^{(0)})^2 \right)^2 + \frac{1}{8} \sum_{l=1}^L \left( r_l(\theta)^2 - r_l(\theta_*)^2 \right)^2 \\ s_3(\theta) &= \frac{1}{48} \left( (u^{(0)}(\theta))^3 - (u^{(0)}(\theta_*))^3 \right)^2 + \frac{1}{16} \sum_{\substack{l,l',l''=0 \\ l=l'+l''}}^L \left| u^{(l)}(\theta) \overline{u^{(l')}(\theta) u^{(l'')}(\theta)} - u^{(l)}(\theta_*) \overline{u^{(l')}(\theta_*) u^{(l'')}(\theta_*)} \right|^2 \\ &= \frac{1}{12} \left( (\theta^{(0)})^3 - (\theta_*^{(0)})^3 \right)^2 + \frac{1}{8} \sum_{l=1}^L \left( \theta^{(0)} \cdot r_l(\theta)^2 - \theta_*^{(0)} \cdot r_l(\theta_*)^2 \right)^2 \\ &\quad + \frac{1}{16} \sum_{\substack{l,l',l''=1 \\ l=l'+l''}}^L \left( r_{l,l',l''}(\theta)^2 + r_{l,l',l''}(\theta_*)^2 - 2r_{l,l',l''}(\theta)r_{l,l',l''}(\theta_*) \cos(\lambda_{l,l',l''}(\theta) - \lambda_{l,l',l''}(\theta_*)) \right). \end{aligned}$$

Since each moment variety  $\mathcal{V}_k(\theta_*)$  in (2.17) is precisely the set of points  $\{\theta \in \mathbb{R}^d : s_1(\theta) = 0, \dots, s_k(\theta) = 0\}$ , this implies also that

$$\begin{aligned} \mathcal{V}_0(\theta_*) &= \mathbb{R}^d, \quad \mathcal{V}_1(\theta_*) = \{\theta : \theta^{(0)} = \theta_*^{(0)}\}, \\ \mathcal{V}_2(\theta_*) &= \{\theta : \theta^{(0)} = \theta_*^{(0)} \text{ and } r_l(\theta) = r_l(\theta_*) \text{ for each } l = 1, \dots, L\}. \end{aligned}$$

Thus the minimization of  $s_1(\theta)$  on  $\mathcal{V}_0(\theta_*)$  is over the global function mean  $\theta^{(0)}$ , the minimization of  $s_2(\theta)$  on  $\mathcal{V}_1(\theta_*)$  is over the Fourier power spectrum  $\{r_l(\theta) : l = 1, \dots, L\}$ , and the minimization of  $s_3(\theta)$  on  $\mathcal{V}_2(\theta_*)$  is over the Fourier bispectrum  $\{\lambda_{l,l',l''}(\theta) : l = l' + l''\}$ .

In high noise, minimizing the population log-likelihood function  $R(\theta)$  becomes similar to successively minimizing  $s_1(\theta)$ ,  $s_2(\theta)$ , and  $s_3(\theta)$ . The following result describes the nature of these three optimization landscapes.

**Theorem 3.4.** *For any  $L \geq 1$  and generic  $\theta_* \in \mathbb{R}^d$ , the minimizations of  $s_1(\theta)$  over  $\mathcal{V}_0(\theta_*)$  and of  $s_2(\theta)$  over  $\mathcal{V}_1(\theta_*)$  are globally benign. However, for any  $L \geq 30$ , there exists a non-empty open subset  $U \subset \mathbb{R}^d$  such that for any  $\theta_* \in U$ , the minimization of  $s_3(\theta)$  over  $\mathcal{V}_2(\theta_*)$  has a local minimizer outside  $\mathcal{O}_{\theta_*}$  that is non-degenerate up to orbit.*

The correspondence between optimization landscapes shown in Theorem 2.13(a) then implies that, for the class of signals  $\theta_* \in U$  described in Theorem 3.4 and in sufficiently high noise, the landscape of the population negative log-likelihood function  $R(\theta)$  must also have spurious local minimizers near those of  $s_3(\theta)$ . The particular local minimizers of  $s_3(\theta)$  that we exhibit in the proof of Theorem 3.4 correspond to certain Fourier phase shifts of the true signal. This example is somewhat analogous to the spurious local minimizers discovered in dimensions  $d \geq 53$  for the log-likelihood landscape of discrete MRA in [FSWW20, Section 4.6].

We conjecture, based on the algebraic similarities between these models, that spurious local minimizers of  $R(\theta)$  may also exist for generic  $\theta_* \in \mathbb{R}^d$  in the  $\text{SO}(3)$ -rotational models to be discussed in Section 4, and we leave this as an open question.

#### 4. SPHERICAL REGISTRATION AND CRYO-EM

We now describe examples of estimating a function in 2 or 3 dimensions, observed under  $\text{SO}(3)$  rotations of its domain. Section 4.1 studies estimation on the sphere, Section 4.2 studies estimation in  $\mathbb{R}^3$ , and Section 4.3 studies a simplified “cryo-EM model” of estimation in  $\mathbb{R}^3$  with a tomographic projection onto a 2-dimensional plane.

**4.1. Spherical registration.** Let  $\mathcal{S}^2 \subset \mathbb{R}^3$  be the unit sphere, and let  $f : \mathcal{S}^2 \rightarrow \mathbb{R}$  be a function on this sphere. We parametrize  $\mathcal{S}^2$  by the latitude  $\phi_1 \in [0, \pi]$  and longitude  $\phi_2 \in [0, 2\pi)$ . Writing  $f_{\mathbf{g}}(\phi_1, \phi_2) = f(\mathbf{g}^{-1} \cdot (\phi_1, \phi_2))$  for the rotation of the function  $f$ , we consider the observation model with samples

$$f_{\mathbf{g}}(\phi_1, \phi_2) d(\phi_1, \phi_2) + \sigma dW(\phi_1, \phi_2)$$

where  $\mathbf{g} \in \text{SO}(3)$  is a uniform random rotation for each sample,  $d(\phi_1, \phi_2) = \sin \phi_1 d\phi_1 d\phi_2$  denotes the surface area measure on  $\mathcal{S}^2$ , and  $dW(\phi_1, \phi_2)$  is a standard Gaussian white noise process on  $\mathcal{S}^2$ . This observation model may be understood as observing a realization of the Gaussian process  $\{\int h(\phi_1, \phi_2)[f_{\mathbf{g}}(\phi_1, \phi_2)d(\phi_1, \phi_2) + \sigma dW(\phi_1, \phi_2)]\}_{h \in L_2(\mathcal{S}^2)}$  defined analogously to (3.1), or equivalently, as observing each coefficient of  $f_{\mathbf{g}}$  in an orthonormal basis of  $L_2(\mathcal{S}^2)$  with i.i.d.  $\mathcal{N}(0, \sigma^2)$  noise.

We choose as our orthonormal basis the real spherical harmonics

$$h_{lm}(\phi_1, \phi_2) \quad \text{for } l = 0, 1, 2, \dots \text{ and } m = -l, -l+1, \dots, l-1, l.$$

We assume that  $f : \mathcal{S}^2 \rightarrow \mathbb{R}$  has a finite bandlimit  $L \geq 1$  in this basis, i.e. it takes the form

$$f(\phi_1, \phi_2) = \sum_{l=0}^L \sum_{m=-l}^l \theta_m^{(l)} h_{lm}(\phi_1, \phi_2). \quad (4.1)$$

We may then represent  $f$  by its vector of real spherical harmonic coefficients

$$\theta = (\theta_m^{(l)} : l = 0, \dots, L \text{ and } m = -l, \dots, l) \in \mathbb{R}^d, \quad d = (L+1)^2.$$

This subspace of bandlimited functions is closed under  $\text{SO}(3)$ -rotations of  $\mathcal{S}^2$ , and we review the forms of  $h_{lm}$  and of the rotational action on the basis coefficients in Appendix D.2.

The following result describes the decomposition of total dimension in Theorem 2.7(a) for bandlimits  $L \geq 10$ .

**Theorem 4.1.** *For any  $L \geq 10$ , we have*

$$\text{trdeg}(\mathcal{R}_{\leq 1}^G) = 1, \quad \text{trdeg}(\mathcal{R}_{\leq 2}^G) = L+1, \quad \text{trdeg}(\mathcal{R}_{\leq 3}^G) = \text{trdeg}(\mathcal{R}^G) = d-3.$$

**Corollary 4.2.** *A generic signal  $\theta_* \in \mathbb{R}^d$  in this spherical registration model for  $L \geq 10$  has the following properties:*

- (a)  $\theta_*$  may be identified up to a finite list of orbits by the moments of  $g \cdot \theta_*$  up to order  $K = 3$ .
- (b) For  $(\theta_*, \mathbf{G})$ -dependent constants  $C, c > 0$  independent of  $\sigma$ , the Fisher information  $I(\theta_*)$  has  $d_0 = 3$  eigenvalues of 0 and  $d_k$  eigenvalues in  $[c\sigma^{-2k}, C\sigma^{-2k}]$  for  $k = 1, 2, 3$  and  $(d_1, d_2, d_3) = (1, L, L(L+1)-3)$ .

**Remark 4.3.** The result of Theorem 4.1 was conjectured for all bandlimits  $L \geq 10$  in [BBSK<sup>+</sup>17, Conjecture 5.6], and it was verified numerically in exact-precision arithmetic for  $L \in \{10, \dots, 16\}$ . Our result resolves this conjecture for all  $L \geq 10$ . Conversely, for low bandlimits  $L \leq 9$ , it was shown in [BBSK<sup>+</sup>17, Section 5.4] that  $K > 3$  strictly, meaning that moments up to 3<sup>rd</sup> order are insufficient to locally identify  $\theta_*$  up to its orbit.

Turning to the forms of  $s_k(\theta)$  in (2.19), let us denote the real spherical harmonic coefficients of the true function by  $\theta_* \in \mathbb{R}^d$ . We write as shorthand

$$u^{(l)}(\theta) = (u_m^{(l)}(\theta) : m = -l, \dots, l) \in \mathbb{C}^{2l+1}$$

for the complex spherical harmonic coefficients at frequency  $l$ , which are defined from the real coefficients  $(\theta_m^{(l)} : m = -l, \dots, l)$  by a unitary transform described in (D.11). We denote

$$B_{l,l',l''}(\theta) = \sum_{m=-l}^l \sum_{\substack{m'=-l' \\ m''=-l'' \\ m'=m+m'}}^{l'} \sum_{m''=-l''}^{l''} \langle l, m; l', m' | l'', m'' \rangle \overline{u_m^{(l)}(\theta) u_{m'}^{(l')}(\theta)} u_{m''}^{(l'')}(\theta) \quad (4.2)$$

where  $\langle l, m; l', m' | l'', m'' \rangle \in \mathbb{R}$  is the Clebsch-Gordan coefficient. These quantities express the integrals of three-fold products of spherical harmonics over  $\mathcal{S}^2$  and arise naturally in the computation of 3<sup>rd</sup>-order moments of  $g \cdot \theta$ . We review their definition in Appendix D.1. The functions  $B_{l,l',l''}(\theta)$  are analogous to the scaled components  $r_{l,l',l''}(\theta) \lambda_{l,l',l''}(\theta)$  of the Fourier bispectrum that appeared in the 1-dimensional MRA example of Section 3. The minimizations of  $s_1(\theta)$ ,  $s_2(\theta)$ , and  $s_3(\theta)$  described in Theorem 2.13 may then be analogously understood as minimizing the global function mean, the power in each spherical harmonic frequency, and certain “bispectrum” variables for each frequency.

**Theorem 4.4.** *For any  $L \geq 1$ ,*

$$\begin{aligned} s_1(\theta) &= \frac{1}{2} \left( u^{(0)}(\theta) - u^{(0)}(\theta_*) \right)^2 \\ s_2(\theta) &= \frac{1}{4} \sum_{l=0}^L \frac{1}{2l+1} \left( \|u^{(l)}(\theta)\|^2 - \|u^{(l)}(\theta_*)\|^2 \right)^2 \\ s_3(\theta) &= \frac{1}{6} \sum_{\substack{l,l',l''=0 \\ |l-l'| \leq l'' \leq l+l'}}^L \frac{1}{2l''+1} \left| B_{l,l',l''}(\theta) - B_{l,l',l''}(\theta_*) \right|^2. \end{aligned}$$

We prove Theorems 4.1 and 4.4 in Appendix D.2. Here, let us describe the high-level proof idea for Theorem 4.1, which is used also in our analyses of the cryo-EM models to follow. By Lemma 2.8, it suffices to analyze the ranks of the Hessians  $\nabla^2 s_1(\theta_*)$ ,  $\nabla^2 s_2(\theta_*)$ , and  $\nabla^2 s_3(\theta_*)$  at a generic point  $\theta_* \in \mathbb{R}^d$ . This analysis is straightforward for  $s_1, s_2$ , and the core of the proof is to show that  $\nabla^2 s_3(\theta_*)$  has full rank  $d-3$  (which accounts for the 3-dimensional orbit of  $\theta_*$ ) when  $L \geq 10$ .

Importantly, for any matrix  $M(\theta)$  that is analytic in  $\theta$ , we have  $\text{rank}(M(\theta)) < k$  if and only if every  $k \times k$  submatrix of  $M(\theta)$  has determinant 0. Because the  $k \times k$  minors are themselves analytic in  $\theta$ , this holds either for all  $\theta \in \mathbb{R}^d$ , or only for  $\theta$  outside a generic subset of  $\mathbb{R}^d$ . This implies the following fact.

**Fact 4.5.** *For any  $k \geq 1$  and matrix  $M(\theta)$  whose entries are analytic in  $\theta$ , we have  $\text{rank } M(\theta_*) \geq k$  for generic points  $\theta_* \in \mathbb{R}^d$  if and only if there exists at least one point  $\theta_* \in \mathbb{R}^d$  for which this inequality holds.*

Thus, to show that  $\text{rank}(\nabla^2 s_3(\theta_*)) \geq d-3$  for generic  $\theta_* \in \mathbb{R}^d$ , it suffices to construct a single point  $\theta_* \in \mathbb{R}^d$  where this holds. We do this by analyzing the explicit form of  $\nabla^2 s_3(\theta_*)$  derived from Theorem 4.4. For  $L = 10$ , we exhibit such a point  $\theta_*$  numerically. We then use this as a base case to inductively construct  $\theta_*$  for all  $L \geq 10$ , by carefully choosing certain coordinates of  $\theta_*$  to be 0 so that  $\nabla^2 s_3(\theta_*)$  has a sparse structure and its rank may be explicitly deduced from the ranks of  $2 \times 2$  submatrices.

**4.2. Unprojected cryo-EM.** Consider now a function  $f : \mathbb{R}^3 \rightarrow \mathbb{R}$ , and the action of  $\text{SO}(3)$  on  $\mathbb{R}^3$  given by rotation about the origin. Write  $f_{\mathbf{g}}(x) = f(\mathbf{g}^{-1} \cdot x)$  for the rotated function. We consider the observation model with samples

$$f_{\mathbf{g}}(x) dx + \sigma dW(x)$$

where  $\mathbf{g} \in \text{SO}(3)$  is uniformly random for each sample, and  $dW(x)$  is a standard Gaussian white noise process on  $\mathbb{R}^3$ . This is an unprojected model of the single-particle reconstruction problem in cryo-EM, to which we will add a tomographic projection in the next section. This model may be of independent interest for applications to cryo-ET, described in Appendix F.1.

We model  $f$  using a basis representation for its Fourier transform  $\hat{f} : \mathbb{R}^3 \rightarrow \mathbb{C}$ , similar to the approach of [BBSK<sup>+</sup>17, Section 5.5]. We parametrize the Fourier domain  $\mathbb{R}^3$  by spherical coordinates  $(\rho, \phi_1, \phi_2)$  with radius  $\rho \geq 0$ , latitude  $\phi_1 \in [0, \pi]$  and longitude  $\phi_2 \in [0, 2\pi)$ , and decompose  $\hat{f}(\rho, \phi_1, \phi_2)$  in a complex basis

$\{\hat{j}_{lsm}\}$  given by the product of the complex spherical harmonics  $y_{lm}(\phi_1, \phi_2)$  (reviewed in Appendix D.1) with radial functions  $z_s(\rho)$ :

$$\hat{j}_{lsm}(\rho, \phi_1, \phi_2) = z_s(\rho) y_{lm}(\phi_1, \phi_2) \quad \text{for } s \geq 1, \quad l \geq 0, \quad m \in \{-l, \dots, l\}. \quad (4.3)$$

Here  $\{z_s : s \geq 1\}$  may be any system of radial basis functions  $z_s : [0, \infty) \rightarrow \mathbb{R}$  satisfying the orthogonality relation

$$\int_0^\infty \rho^2 z_s(\rho) z_{s'}(\rho) d\rho = \mathbf{1}\{s = s'\}, \quad (4.4)$$

so that  $\{\hat{j}_{lsm}\}$  are orthonormal over  $L_2(\mathbb{R}^3, \mathbb{C})$ . The inverse Fourier transforms  $\{j_{lsm}\}$  of  $\{\hat{j}_{lsm}\}$  then provide a complex orthonormal basis in the original signal domain of  $f$ .

Fixing integer bandlimits  $L \geq 1$  and  $S_0, \dots, S_L \geq 1$ , we define the index set

$$\mathcal{I} = \{(l, s, m) : 0 \leq l \leq L, 1 \leq s \leq S_l, -l \leq m \leq l\}, \quad d = |\mathcal{I}| = \sum_{l=0}^L (2l+1) S_l \quad (4.5)$$

and assume that  $f$  is  $(L, S_0, \dots, S_L)$ -bandlimited in the sense of admitting the finite basis representation

$$f = \sum_{(l,s,m) \in \mathcal{I}} u_m^{(ls)} \cdot j_{lsm}, \quad u = (u_m^{(ls)} : (l, s, m) \in \mathcal{I}) \in \mathbb{C}^d. \quad (4.6)$$

This corresponds to modeling the Fourier transform  $\hat{f}$  up to the spherical frequency  $L$ , and up to the radial frequency  $S_l$  for each spherical component  $l = 0, 1, \dots, L$ . For real-valued functions  $f$ , writing  $u = \hat{V}^* \theta$  for a unitary transform  $\hat{V} \in \mathbb{C}^{d \times d}$  defined explicitly in (D.31), we then obtain a real sequence representation

$$f = \sum_{(l,s,m) \in \mathcal{I}} \theta_m^{(ls)} \cdot h_{lsm}, \quad \theta = (\theta_m^{(ls)} : (l, s, m) \in \mathcal{I}) \in \mathbb{R}^d \quad (4.7)$$

for a real-valued orthonormal basis  $\{h_{lsm}\}$ . We describe the forms of  $h_{lsm}$  and the rotational action on the basis coefficients  $\theta \in \mathbb{R}^d$  in Appendix D.3.

The following result describes the decomposition of total dimension in Theorem 2.7(a), assuming  $L \geq 1$  and  $S_l \geq 2$  for each  $l = 0, \dots, L$ . (Note that the case of  $S_0 = \dots = S_L = 1$  would be similar to the spherical registration example of Section 4.1, and a lower bound of  $L \geq 10$  would be needed in this case to ensure  $K = 3$ .)

**Theorem 4.6.** *For any  $L \geq 1$  and  $S_0, \dots, S_L \geq 2$ , we have*

$$\begin{aligned} \text{trdeg}(\mathcal{R}_{\leq 1}^G) &= S_0 \\ \text{trdeg}(\mathcal{R}_{\leq 2}^G) &= \sum_{l=0}^L d(S_l), \quad d(S_l) \equiv \begin{cases} \frac{S_l(S_l+1)}{2} & \text{for } S_l < 2l+1 \\ (2l+1)(S_l-l) & \text{for } S_l \geq 2l+1 \end{cases} \\ \text{trdeg}(\mathcal{R}_{\leq 3}^G) &= \text{trdeg}(\mathcal{R}^G) = d - 3. \end{aligned}$$

**Corollary 4.7.** *In this unprojected cryo-EM model with  $S_0, \dots, S_L \geq 2$ , a generic signal  $\theta_* \in \mathbb{R}^d$  may be identified up to a finite list of orbits by the moments of  $g \cdot \theta_*$  up to order  $K = 3$  if  $L \geq 2$ , and up to order  $K = 2$  if  $L = 1$ .*

**Remark 4.8.** In [BBSK<sup>+</sup>17, Conjecture B.1], the authors conjectured that a generic signal  $\theta_*$  in this model may be identified by 3<sup>rd</sup>-order moments up to a *single unique* orbit when  $L \geq 1$  and  $S_0 = \dots = S_L \geq 3$ . As discussed in [BBSK<sup>+</sup>17], this would hold if the fraction field of all  $G$ -invariant rational functions coincides with that generated by  $\mathcal{R}_{\leq 3}^G$ . Theorem 4.6 shows the weaker statement that these fraction fields have the same transcendence degree. Thus Corollary 4.7(a) guarantees only that  $\theta_*$  may be identified up to a finite list of orbits, and we show this under a slightly weaker requirement that  $S_0, \dots, S_L \geq 2$ .

Turning to the forms of  $s_k(\theta)$  that define the moment optimization (2.19), write  $\theta_* \in \mathbb{R}^d$  for the true coefficients in the above real basis  $\{h_{lsm}\}$ . Let

$$u^{(ls)}(\theta) = (u_m^{(ls)}(\theta) : m = -l, \dots, l) \in \mathbb{C}^{2l+1} \quad (4.8)$$

be the components of the *complex* coefficients  $u = \hat{V}^* \theta$  for the frequency pair  $(l, s)$ , and define analogously to (4.2)

$$B_{(l,s),(l',s'),(l'',s'')}(\theta) = \sum_{m=-l}^l \sum_{\substack{m'=-l' \\ m''=m+m'}}^{l'} \sum_{m''=-l''}^{l''} \langle l, m; l', m' | l'', m'' \rangle \overline{u_m^{(ls)}(\theta)} u_{m'}^{(l's')}(\theta) u_{m''}^{(l''s'')}(\theta). \quad (4.9)$$

When the original function  $f : \mathbb{R}^3 \rightarrow \mathbb{R}$  is real-valued, we verify in the proof of Theorem 4.9 below that each  $B_{(l,s),(l',s'),(l'',s'')}(\theta)$  is also real-valued.

**Theorem 4.9.** *For any  $L \geq 1$  and  $S_0, \dots, S_L \geq 1$ ,*

$$\begin{aligned} s_1(\theta) &= \frac{1}{2} \sum_{s=1}^{S_0} \left( u^{(0s)}(\theta) - u^{(0s)}(\theta_*) \right)^2 \\ s_2(\theta) &= \frac{1}{4} \sum_{l=0}^L \frac{1}{2l+1} \sum_{s,s'=1}^{S_l} \left( \langle u^{(ls)}(\theta), u^{(ls')}(\theta) \rangle - \langle u^{(ls)}(\theta_*), u^{(ls')}(\theta_*) \rangle \right)^2 \\ s_3(\theta) &= \frac{1}{12} \sum_{\substack{l,l',l''=0 \\ |l-l'|\leq l''\leq l+l'}}^L \frac{1}{2l''+1} \sum_{s=1}^{S_l} \sum_{s'=1}^{S_{l'}} \sum_{s''=1}^{S_{l''}} \left( B_{(l,s),(l',s'),(l'',s'')}(\theta) - B_{(l,s),(l',s'),(l'',s'')}(\theta_*) \right)^2. \end{aligned}$$

In this model, the optimization of  $s_1(\theta)$  is over the mean component  $u^{(0s)}(\theta)$  corresponding to each radial frequency  $s$ . The optimization of  $s_2(\theta)$  is over not just the power  $\|u^{(ls)}(\theta)\|^2$  within each frequency pair  $(l, s)$ , but also the cross-correlations between  $u^{(ls)}$  and  $u^{(ls')}$  for different radial frequencies  $s$  and  $s'$ .

The proofs of Theorems 4.6 and 4.9 are deferred to Appendix D.3. The argument for Theorem 4.6 is similar to that of Theorem 4.1: When  $S_0, \dots, S_L \geq 2$ , the claim that  $\text{rank}(\nabla^2 s_3(\theta_*)) \geq d-3$  may be established by induction on  $L$  down to the base case of  $L=1$  rather than  $L=10$ , using a different construction of the point  $\theta_* \in \mathbb{R}^d$  that induces a sparse structure in  $\nabla^2 s_3(\theta_*)$ .

**4.3. Projected cryo-EM.** We now extend the model of the preceding section to include the tomographic projection arising in cryo-EM. In this projected model, we observe samples

$$(\Pi \cdot f_g)(x) dx + \sigma dW(x) \quad (4.10)$$

on  $\mathbb{R}^2$  where, for  $x = (x_1, x_2) \in \mathbb{R}^2$ , the tomographic projection  $\Pi$  is defined by

$$(\Pi \cdot f_g)(x_1, x_2) = \int_{-\infty}^{\infty} f_g(x_1, x_2, x_3) dx_3, \quad (4.11)$$

and  $dW(x)$  in (4.10) is a standard Gaussian white noise process on the projected domain  $\mathbb{R}^2$ .

Our model setup is similar to [BBSK<sup>+</sup>17, Section 5.5 and Appendix A.4]. We again model the Fourier transform of  $f$  in a basis  $\{\hat{j}_{lsm}\}$  given by the product of complex spherical harmonics with radial functions. We restrict  $f$  to a space of  $(L, S_0, \dots, S_L)$ -bandlimited functions with representation

$$f = \sum_{(l,s,m) \in \mathcal{I}} u_m^{(ls)} \cdot j_{lsm} = \sum_{(l,s,m) \in \mathcal{I}} \theta_m^{(ls)} \cdot h_{lsm} \quad (4.12)$$

for the index set  $\mathcal{I}$  defined in (4.5), where  $\{j_{lsm}\}$  are the inverse Fourier transforms of  $\{\hat{j}_{lsm}\}$ , and the second equality describes a parametrization by an equivalent real orthonormal basis  $\{h_{lsm}\}$  as before. In Appendix D.4, we apply the Fourier slice theorem to derive basis representations for the tomographic projection  $\Pi \cdot f$ . These take the forms

$$\Pi \cdot f = \sum_{(s,m) \in \tilde{\mathcal{I}}} \tilde{u}_m^{(s)} j_{sm} = \sum_{(s,m) \in \tilde{\mathcal{I}}} \tilde{\theta}_m^{(s)} h_{sm}, \quad (4.13)$$

where  $\{j_{sm}\}$  and  $\{h_{sm}\}$  are (complex and real, resp.) basis functions over  $\mathbb{R}^2$ , and  $\Pi \cdot f$  is bandlimited to an index set

$$\tilde{\mathcal{I}} = \left\{ (s, m) : 1 \leq s \leq S, -L \leq m \leq L \right\}, \quad \tilde{d} = |\tilde{\mathcal{I}}| = S(2L+1) \quad (4.14)$$

for  $S = \max(S_0, \dots, S_L)$ . This expresses  $\Pi$  as a linear map from  $\theta \in \mathbb{R}^d$  to  $\tilde{\theta} \in \mathbb{R}^{\tilde{d}}$ , and we give its explicit form in (D.57). We choose radial functions to ensure that the basis  $\{h_{sm}\}$  is orthonormal in  $L_2(\mathbb{R}^2)$ , so

that (4.10) is equivalent to observing the coefficients of  $\Pi \cdot f$  in this basis with i.i.d.  $\mathcal{N}(0, \sigma^2)$  noise. Further details of the setup are described in Appendix D.4.

The following result verifies that when the bandlimits satisfy  $L \geq 1$  and  $S_1, \dots, S_L \geq 4$ , we have also  $\text{trdeg } \tilde{\mathcal{R}}_{\leq \tilde{K}}^G = \text{trdeg } \mathcal{R}^G$  for  $\tilde{K} = 3$ .

**Theorem 4.10.** *For any  $L \geq 1$  and  $S_0, \dots, S_L \geq 4$ , we have*

$$\begin{aligned} \text{trdeg}(\tilde{\mathcal{R}}_{\leq 1}^G) &= S_0 \\ \text{trdeg}(\tilde{\mathcal{R}}_{\leq 2}^G) &= \sum_{l=0}^L d(S_l), \quad d(S_l) \equiv \begin{cases} \frac{S_l(S_l+1)}{2} & \text{for } S_l < 2l+1 \\ (2l+1)(S_l-l) & \text{for } S_l \geq 2l+1 \end{cases} \\ \text{trdeg}(\tilde{\mathcal{R}}_{\leq 3}^G) &= \text{trdeg}(\mathcal{R}^G) = d-3, \end{aligned}$$

which matches the values of  $\text{trdeg}(\mathcal{R}_{\leq 1}^G)$ ,  $\text{trdeg}(\mathcal{R}_{\leq 2}^G)$ , and  $\text{trdeg}(\mathcal{R}_{\leq 3}^G)$  in the unprojected setting of Theorem 4.6.

**Corollary 4.11.** *In this projected cryo-EM model with  $S_0, \dots, S_L \geq 4$ , a generic signal  $\theta_* \in \mathbb{R}^d$  may be identified up to a finite list of orbits by the moments of  $\Pi(g \cdot \theta_*)$  up to order  $\tilde{K} = 3$  if  $L \geq 2$ , and order  $\tilde{K} = 2$  if  $L = 1$ .*

We prove Theorem 4.10 in Appendix D.4, where we also state an analogue of Theorem 4.9 that describes the explicit forms of  $\tilde{s}_k(\theta)$  for  $k = 1, 2, 3$  in this projected model.

Our proof of Theorem 4.10 again constructs a point  $\theta_* \in \mathbb{R}^d$  where  $\text{rank}(\nabla^2 \tilde{s}_3(\theta_*)) \geq d-3$ . However, the form of  $\nabla^2 \tilde{s}_3(\theta_*)$  now involves the precise form of the projection  $\Pi$ , and our choice of  $\theta_*$  does not induce sparsity in this Hessian. Instead, we choose  $\theta_*$  to have many coordinates equal to 0, and track the dependence of minors of  $\nabla^2 \tilde{s}_3(\theta_*)$  on the non-zero coordinates of  $\theta_*$  to show they are generically non-vanishing. We give this argument in the proof of Lemma D.10 in Appendix D.4.

**Remark 4.12.** Taking  $S_0 = \dots = S_L = S$  yields a model equivalent to the projected cryo-EM model with  $S$  spherical shells in [BBSK<sup>+</sup>17, Section 5.5]. In [BBSK<sup>+</sup>17, Conjecture 5.11], the authors conjectured that a generic signal  $\theta_*$  may be identified up to a finite list of orbits by 3<sup>rd</sup>-order moments if  $S \geq 2$ . Corollary 4.11(a) thus resolves this conjecture positively when  $S \geq 4$ . The constraint  $S \geq 4$  is technical, and we believe that the conjecture holds as stated for  $S \in \{2, 3\}$  as well, but we do not pursue these cases in this work.

## 5. NUMERICAL EVALUATIONS OF THE FISHER INFORMATION

We conclude with an empirical investigation of the spectrum of the Fisher information matrix in two simulated examples of the unprojected cryo-EM model described in Section 4.2.

In each example, we begin with a near-atomic-resolution electric potential map estimated from a cryo-EM experiment. We obtain a finite-dimensional approximation to this map by applying a low-pass filter to its Fourier transform, followed by a basis approximation for the filtered map. We simulate noisy and rotated samples using this finite-dimensional approximation as the underlying true signal, for various inverse-SNR parameters

$$\alpha \equiv \sigma^2 / \|\theta_*\|^2.$$

We then study the dependence of eigenvalues of the observed information matrix  $\nabla^2 R_n(\theta_*)$  on  $\alpha$ .

**Rotavirus VP6 trimer.** We consider a map of the VP6 trimer in bovine rotavirus, reported in [ZSX<sup>+</sup>08] (EMDB:1461). A contour plot of this map is overlaid with the atomic structure previously obtained by [MPN<sup>+</sup>01] (PDB:1QHD), in Figure 5.1(a). We applied low-pass filters in the Fourier domain at two different cutoff frequencies, a “low-resolution” frequency of  $(24.6\text{\AA})^{-1}$  and a “medium-resolution” frequency of  $(8.2\text{\AA})^{-1}$ . The corresponding smoothed maps in the spatial domain are depicted in Figure E.1 of Appendix E.

We approximated each smoothed map using a finite basis of the form (4.3), with an adaptive construction of the radial functions  $\{z_s\}$  to maximize the power captured by each successive radial frequency. Details of our numerical procedures are described in Appendix E. Choosing bandlimits  $(S, L) = (5, 8)$  and total dimension  $d = L(S+1)^2 = 405$  gave an accurate approximation to the 24.6Å-resolution map that reveals the trimer composition of the VP6 complex, as depicted in Figure 5.1(b). Choosing bandlimits  $(S, L) = (10, 20)$




FIGURE 5.1. (a) 3.8Å-resolution cryo-EM map of the rotavirus VP6 trimer, overlaid with the atomic structure. (b) A finite-dimensional approximation using 405 basis functions at 24.6Å-resolution (displayed in a rotated orientation for clarity). (c) An approximation using 4410 basis functions at 8.2Å-resolution. (d–f) We stratify the eigenvalues of the 405-dimensional observed Fisher information corresponding to (b) into three “eigenvalue tiers” according to Theorem 4.6, and plot the scalings of the 10<sup>th</sup>, 30<sup>th</sup>, 50<sup>th</sup>, 70<sup>th</sup>, and 90<sup>th</sup> percentiles of eigenvalues in each tier against  $1/\alpha \propto \sigma^{-2}$ ,  $1/\alpha^2 \propto \sigma^{-4}$ , and  $1/\alpha^3 \propto \sigma^{-6}$ . (These quantiles nearly overlap for Tier 1.) Linear trends fitted using least squares are shown as dashed lines. (g–i) The same for the 4410-dimensional Fisher information matrix corresponding to (c).

and total dimension  $d = L(S + 1)^2 = 4410$  gave an accurate approximation of the 8.2Å-resolution map that captures interesting aspects of the tertiary and secondary structure, as shown in Figure 5.1(c). We denote the basis coefficients of these approximated maps as  $\theta_* \in \mathbb{R}^d$ .

We computed the Hessians  $\nabla^2 R_n(\theta_*)$  of the empirical negative log-likelihood functions from  $n = 500,000$  simulated samples, with inverse-SNR  $\alpha = \sigma^2 / \|\theta_*\|^2 \in [0.04, 0.10]$ . We then separated the largest  $d - 3$  eigenvalues of  $\nabla^2 R_n(\theta_*)$  into three “tiers” with cardinalities  $(d_1, d_2, d_3)$  as implied by Theorem 4.6. Figure 5.1(d–f) depicts representative eigenvalues in each tier, plotted against  $1/\alpha \propto \sigma^{-2}$ ,  $1/\alpha^2 \propto \sigma^{-4}$ , and  $1/\alpha^3 \propto \sigma^{-6}$ . A linear trend is observed in all settings, in agreement with the prediction of Theorem 2.7. This may be contrasted with Figure E.2 in Appendix E, which instead plots eigenvalues in all three tiers against  $1/\alpha \propto \sigma^{-2}$ , and where non-linearity of the scaling is visually apparent for Tiers 2 and 3.

**Hemoglobin.** We consider a map of hemoglobin, reported in [KRBD17] (EMDB:3650, PDB:5NI1). A contour plot overlaid with the atomic structure is presented in Figure 5.2(a). We applied a low-pass filter




FIGURE 5.2. (a) 3.4Å-resolution cryo-EM map of hemoglobin, overlaid with the atomic structure. (b) A finite-dimensional approximation using 3528 basis functions at 7.0Å-resolution. (c-e) The 10<sup>th</sup>, 30<sup>th</sup>, 50<sup>th</sup>, 70<sup>th</sup>, and 90<sup>th</sup> percentiles of eigenvalues within each “eigenvalue tier” of the 3528-dimensional observed Fisher information, plotted against  $1/\alpha \propto \sigma^{-2}$ ,  $1/\alpha^2 \propto \sigma^{-4}$ ,  $1/\alpha^3 \propto \sigma^{-6}$  as in Figure 5.1.

with cutoff frequency  $(7.0\text{\AA})^{-1}$  in the Fourier domain, depicted in Figure E.1. We then applied a basis approximation with bandlimits  $(S, L) = (8, 20)$  and total dimension  $d = 3528$ . The approximated map is shown in Figure 5.2(b), and captures important aspects of the secondary structure including the locations of the  $\alpha$ -helices and embedded prosthetic heme groups. We denote the basis coefficients of this approximation as  $\theta_*$ .

Figure 5.2(c-e) again depicts the leading  $d - 3$  eigenvalues of  $\nabla^2 R_n(\theta_*)$  computed from  $n = 500,000$  simulated samples, stratified into three tiers of sizes  $(d_1, d_2, d_3)$ . Linear trends with  $1/\alpha \propto \sigma^{-2}$ ,  $1/\alpha^2 \propto \sigma^{-4}$ , and  $1/\alpha^3 \propto \sigma^{-6}$  are again observed, and may be contrasted with the non-linear scalings of eigenvalues in Tiers 2 and 3 with  $1/\alpha$  as depicted in Figure E.2.

We note that although the eigenvalues of  $\nabla^2 R_n(\theta_*)$  do scale with powers of the SNR  $1/\alpha$  according to our theoretical predictions, at any fixed SNR and for basis dimensions exceeding  $d \approx 100$ , we do not observe a clear separation between the eigenvalues of Tier 2 and of Tier 3, due to the variation in magnitude of eigenvalues corresponding to differing radial frequencies within each tier.

In these examples, we also begin to observe some deviations from the predicted eigenvalue scalings at the higher and lower ends of tested SNR. Deviations in higher basis dimensions  $d$  and at lower SNR  $1/\alpha$  (seen in Figures 5.1(h-i) and 5.2(d)) are likely finite-sample effects due to differences between the observed information matrix  $\nabla^2 R_n(\theta_*)$  and the (population) Fisher information  $I(\theta_*) = \nabla^2 R(\theta_*)$ . We believe that deviations at higher SNR  $1/\alpha$  (seen in Figures 5.1(e-f) and 5.2(e)) reflect a departure of the behavior of the population Fisher information  $I(\theta_*)$  from the predictions of the large- $\sigma$  theoretical regime. Our largest tested SNR  $1/\alpha = 25$  yields a spectral SNR (average power of signal / average power of noise at a fixed Fourier radius) of 0.2-0.4 near the origin of the Fourier domain, which we believe reflects a level of noise that may be slightly higher than that of modern cryo-EM experiments.

## 6. CONCLUSION

In this work, we characterized properties of the Fisher information matrix and log-likelihood function landscape for continuous group orbit estimation problems in a high noise regime, showing that they are related to the structure of the invariant algebra of the rotational group. We applied these results to study several models of function estimation in finite-dimensional function spaces, in particular establishing that 3<sup>rd</sup>-order moment information is sufficient to locally identify generic signals in these models.

In many interesting applications including single-particle cryo-EM, the target function at full spatial resolution may not admit an accurate low-dimensional approximation. In such settings, our theoretical results may have relevance to estimating lower-dimensional smoothed approximations of the function. We demonstrated in simulation that this theory can accurately predict the noise scalings of the Fisher information eigenvalues for two small protein molecules over a range of sufficiently high noise, or low SNR. We highlight the theoretical understanding of likelihood-based estimation in high-dimensional and infinite-dimensional settings and over a broader range of SNR as a question for future work.

## APPENDIX A. PROOFS FOR GENERAL RESULTS ON THE ORBIT RECOVERY MODEL

**A.1. High-noise expansion.** We first provide a form for  $s_k(\theta)$  and  $\tilde{s}_k(\theta)$  that will be more convenient for later computations. We then prove Theorem 2.2.

**Lemma A.1.** *The expressions  $s_k(\theta)$  and  $\tilde{s}_k(\theta)$  from (2.8) and (2.11) have the equivalent forms*

$$s_k(\theta) = \frac{1}{2(k!)} \mathbb{E}_g [\langle \theta, g \cdot \theta \rangle^k - 2\langle \theta, g \cdot \theta_* \rangle^k + \langle \theta_*, g \cdot \theta_* \rangle^k] \quad (\text{A.1})$$

$$\tilde{s}_k(\theta) = \frac{1}{2(k!)} \mathbb{E}_{g,h} [\langle \Pi \cdot g \cdot \theta, \Pi \cdot h \cdot \theta \rangle^k - 2\langle \Pi \cdot g \cdot \theta, \Pi \cdot h \cdot \theta_* \rangle^k + \langle \Pi \cdot g \cdot \theta_*, \Pi \cdot h \cdot \theta_* \rangle^k] \quad (\text{A.2})$$

where the expectations are over independent Haar-uniform random elements  $g, h \in \mathbf{G}$ .

*Proof.* For the first statement, expanding the square in the definition of  $s_k$  from (2.8), we have

$$\begin{aligned} s_k(\theta) &= \frac{1}{2(k!)} \|T_k(\theta) - T_k(\theta_*)\|_{\text{HS}}^2 \\ &= \frac{1}{2(k!)} \|\mathbb{E}_g[(g \cdot \theta)^{\otimes k}] - \mathbb{E}_g[(g \cdot \theta_*)^{\otimes k}]\|_{\text{HS}}^2 \\ &= \frac{1}{2(k!)} \mathbb{E}_{g,h} [\langle (g \cdot \theta)^{\otimes k}, (h \cdot \theta)^{\otimes k} \rangle - 2\langle (g \cdot \theta)^{\otimes k}, (h \cdot \theta_*)^{\otimes k} \rangle + \langle (g \cdot \theta_*)^{\otimes k}, (h \cdot \theta_*)^{\otimes k} \rangle] \\ &= \frac{1}{2(k!)} \mathbb{E}_{g,h} [\langle g \cdot \theta, h \cdot \theta \rangle^k - 2\langle g \cdot \theta, h \cdot \theta_* \rangle^k + \langle g \cdot \theta_*, h \cdot \theta_* \rangle^k] \\ &= \frac{1}{2(k!)} \mathbb{E}_g [\langle \theta, g \cdot \theta \rangle^k - 2\langle \theta, g \cdot \theta_* \rangle^k + \langle \theta_*, g \cdot \theta_* \rangle^k]. \end{aligned}$$

The last step above applies  $\langle g \cdot u, h \cdot v \rangle = \langle u, (g^\top h) \cdot v \rangle$  and the equality in law  $g^\top h \stackrel{L}{=} g$ . The second statement follows similarly from (2.11), omitting this last step.  $\square$

*Proof of Theorem 2.2.* Part (a) follows from specializing part (b) to  $\tilde{d} = d$  and  $\Pi = \text{Id}$ , and observing that in this case, the term  $\langle \tilde{T}_k(\theta), P_k(\theta) \rangle$  in (2.10) also belongs to  $\mathcal{R}_{\leq k-1}^{\mathbf{G}}$  and hence may be absorbed into  $q_k(\theta)$ —see [KB20, Proposition 2.3] or [FSWW20, Lemma 4.8].

Most of the claims in part (b) follow directly from [KB20, Lemma 2.2]: Specializing to our setting (where  $\rho$  and  $\rho_*$  in [KB20] are the distributions of  $\Pi \cdot g \cdot \theta$  and  $\Pi \cdot h \cdot \theta_*$  for  $g, h \sim \Lambda$ , and where  $\delta$  in [KB20, Eqs. (2.12–2.13)] is bounded as  $\delta \leq C(1 \vee \|\theta\|)$  for a  $(\Pi, \mathbf{G}, \theta_*)$ -dependent constant  $C > 0$  and all  $\theta \in \mathbb{R}^d$ ), this result guarantees that the expansion (2.10) holds for  $\tilde{s}_k$ ,  $\tilde{T}_k$ ,  $P_k$ , and  $q_k$  having all of the stated properties, and for a remainder  $q(\theta)$  that satisfies

$$|q(\theta)| \leq \frac{C_K(1 \vee \|\theta\|)^{2K+2}}{\sigma^{2K+2}} \quad (\text{A.3})$$

when  $\|\theta\| \leq \sigma$ . This remainder  $q(\theta)$  must also be  $\mathbf{G}$ -invariant, as all of the other terms in (2.10) are  $\mathbf{G}$ -invariant.

It remains to verify the bounds for  $\|\nabla q(\theta)\|$  and  $\|\nabla^2 q(\theta)\|$  in (2.9). These types of bounds were shown in the unprojected setting of  $\Pi = \text{Id}$  in [FSW20, Lemma 4.7]. They were not stated explicitly in [KB20], but may be deduced from a small extension of the analysis: Denote by  $\mathbb{E}_g, \mathbb{E}_h$  the expectations over  $g, h \sim \Lambda$ , and by  $\mathbb{E}_\varepsilon, \mathbb{E}_{\varepsilon'}$  those over  $\varepsilon, \varepsilon' \sim \mathcal{N}(0, \text{Id})$ . Write

$$t = 1/\sigma, \quad Y = \Pi \cdot h \cdot \theta_* + t^{-1}\varepsilon, \quad w = \Pi \cdot g \cdot \theta - \Pi \cdot h \cdot \theta_* \in \mathbb{R}^{\tilde{d}},$$

and define

$$f(t) = -\log M(t), \quad M(t) = \mathbb{E}_g \left[ \exp \left( -\frac{t^2 \|w\|^2}{2} + tw^\top \varepsilon \right) \right]$$

Comparing with (2.3), this function  $f(t)$  is the negative log-likelihood for the single sample  $Y$ , up to a  $\theta$ -independent constant and viewed as a function of  $t = 1/\sigma$ . Applying a Taylor expansion of  $f(t)$  around  $t = 0$ , and then taking expectations over  $(h, \varepsilon)$  that define  $Y$ , we have

$$R(\theta) = \text{constant} + \sum_{p=1}^{2K+1} \frac{t^p}{p!} \mathbb{E}_{h,\varepsilon} [f^{(p)}(0)] + \frac{t^{2K+2}}{(2K+2)!} \mathbb{E}_{h,\varepsilon} [f^{(2K+2)}(\xi(h, \varepsilon))] \quad (\text{A.4})$$

for a random point  $\xi(h, \varepsilon)$  between 0 and  $t = 1/\sigma$ . This is a rewriting of the Taylor expansion in [KB20, Eq. (5.7)]. It is shown in [KB20] that the leading terms in (A.4) of orders  $t^1, \dots, t^{2K+1}$  give exactly the leading terms of (2.10), and the last term of (A.4) is the remainder  $q(\theta)$  in (2.10). The bound (A.3) for  $q(\theta)$  follows from [KB20, Eq. (5.22)].

To bound  $\partial_{\theta_a} q(\theta)$  for any index  $a \in \{1, \dots, d\}$ , we may apply a similar Taylor expansion for  $\partial_{\theta_a} R(\theta)$ , and write

$$\partial_{\theta_a} R(\theta) = \sum_{p=1}^{2K+1} \frac{t^p}{p!} \mathbb{E}_{h,\varepsilon} [\partial_t^p \partial_{\theta_a} f(0)] + \frac{t^{2K+2}}{(2K+2)!} \mathbb{E}_{h,\varepsilon} [\partial_t^{2K+2} \partial_{\theta_a} f(\xi(h, \varepsilon))]$$

for a possibly different point  $\xi(h, \varepsilon) \in (0, t)$  depending on the index  $a$ . Then

$$\partial_{\theta_a} q(\theta) = \frac{t^{2K+2}}{(2K+2)!} \mathbb{E}_{h,\varepsilon} [\partial_t^{2K+2} \partial_{\theta_a} f(\xi(h, \varepsilon))] \quad (\text{A.5})$$

and we wish to bound this term for each  $a \in \{1, \dots, d\}$ . The function  $\partial_{\theta_a} f(t)$  is the  $\theta_a$ -derivative of  $f$ , given by

$$\partial_{\theta_a} f(t) = -\frac{\partial_{\theta_a} M(t)}{M(t)}.$$

Then differentiating  $2K+2$  times in  $t$ , we see that  $\partial_t^{2K+2} \partial_{\theta_a} f(t)$  is a sum of at most  $C_K$  terms of the form

$$C_{\ell_0, \dots, \ell_j} \cdot \frac{\partial_t^{\ell_0} \partial_{\theta_a} M(t)}{M(t)} \cdot \frac{\partial_t^{\ell_1} M(t)}{M(t)} \cdot \dots \cdot \frac{\partial_t^{\ell_j} M(t)}{M(t)}$$

for some integers  $j \geq 0$  and  $\ell_0, \dots, \ell_j \geq 0$  such that  $\ell_0 + \dots + \ell_j = 2K+2$ , and for some universal constants  $C_{\ell_0, \dots, \ell_j}$  depending only on  $\ell_0, \dots, \ell_j$ .

From [KB20, Eq. (5.20)], we have

$$\left| \frac{\partial_t^\ell M(\xi)}{M(\xi)} \right| \leq \delta^\ell \mathbb{E}_{\varepsilon'} \left[ (\|W\| + |\xi| \delta)^\ell \right], \quad W = \varepsilon + \mathbf{i} \varepsilon', \quad \delta = \sup_{g, h \in \mathbf{G}} \|\Pi \cdot g \cdot \theta - \Pi \cdot h \cdot \theta_*\|$$

where  $\varepsilon' \sim \mathcal{N}(0, \text{Id}_{\tilde{d} \times \tilde{d}})$  is an independent copy of  $\varepsilon$ . Applying this to any  $\xi \in (0, t)$ , and applying  $\delta \leq C(1 \vee \|\theta\|) \leq C\sigma = Ct^{-1}$ , we obtain

$$\left| \frac{\partial_t^\ell M(\xi)}{M(\xi)} \right| \leq C_\ell \delta^\ell (1 + \|\varepsilon\|)^\ell. \quad (\text{A.6})$$

We may bound the  $t$ -derivatives of  $\partial_{\theta_a} M(t)$  using a similar argument: Introducing the  $a^{\text{th}}$  standard basis vector  $e_a \in \mathbb{R}^d$ , observe that

$$\begin{aligned} \partial_{\theta_a} M(t) &= \mathbb{E}_g \left[ e_a^\top g^\top \Pi^\top (t\varepsilon - t^2 w) \cdot \exp \left( -\frac{t^2 \|w\|^2}{2} + tw^\top \varepsilon \right) \right] \\ &= \mathbb{E}_g \left[ e_a^\top g^\top \Pi^\top (t\varepsilon - t^2 w) \cdot \mathbb{E}_{\varepsilon'} [e^{tw^\top W}] \right], \quad \text{where } W = \varepsilon + \mathbf{i} \varepsilon'. \end{aligned}$$

Then applying Leibniz' rule and the same argument as [KB20, Eqs. (5.18–5.19)] to differentiate in  $t$ , we obtain

$$\begin{aligned}\partial_t^\ell \partial_{\theta_a} M(t) &= \mathbb{E}_g \left[ e_a^\top g^\top \Pi^\top (t\varepsilon - t^2 w) \cdot \mathbb{E}_{\varepsilon'}[e^{tw^\top W}] \cdot \mathbb{E}_{\varepsilon'}[(w^\top (W - tw))^\ell] \right] \\ &\quad + \ell \cdot \mathbb{E}_g \left[ e_a^\top g^\top \Pi^\top (\varepsilon - 2tw) \cdot \mathbb{E}_{\varepsilon'}[e^{tw^\top W}] \cdot \mathbb{E}_{\varepsilon'}[(w^\top (W - tw))^{\ell-1}] \right] \\ &\quad + \binom{\ell}{2} \cdot \mathbb{E}_g \left[ e_a^\top g^\top \Pi^\top (-2w) \cdot \mathbb{E}_{\varepsilon'}[e^{tw^\top W}] \cdot \mathbb{E}_{\varepsilon'}[(w^\top (W - tw))^{\ell-2}] \right].\end{aligned}$$

Applying also  $M(t) = \mathbb{E}_g[\mathbb{E}_{\varepsilon'}[e^{tw^\top W}]]$ , so that  $\mathbb{E}_g[(\cdot)\mathbb{E}_{\varepsilon'}[e^{tw^\top W}]]/M(t)$  is a reweighted average over  $g \in \mathbf{G}$ , this yields analogously to [KB20, Eq. (5.20)] that

$$\begin{aligned}\left| \frac{\partial_t^\ell \partial_{\theta_a} M(\xi)}{M(\xi)} \right| &\leq C_\ell \left[ (|\xi| \|\varepsilon\| + \xi^2 \delta) \cdot \delta^\ell \mathbb{E}_{\varepsilon'}[(\|W\| + |\xi| \delta)^\ell] \right. \\ &\quad \left. + (\|\varepsilon\| + |\xi| \delta) \cdot \delta^{\ell-1} \mathbb{E}_{\varepsilon'}[(\|W\| + |\xi| \delta)^{\ell-1}] + \delta \cdot \delta^{\ell-2} \mathbb{E}_{\varepsilon'}[(\|W\| + |\xi| \delta)^{\ell-2}] \right]\end{aligned}$$

where we have absorbed  $\|e_a^\top g^\top \Pi\|$  into the constant  $C_\ell$ . Applying this with  $\xi \in (0, t)$  and  $\delta \leq C(1 \vee \|\theta\|) \leq Ct^{-1}$ , we get

$$\left| \frac{\partial_t^\ell \partial_{\theta_a} M(\xi)}{M(\xi)} \right| \leq C_\ell \delta^{\ell-1} (1 + \|\varepsilon\|)^{\ell+1}. \quad (\text{A.7})$$

Then combining with (A.6) and applying to the previously stated form of  $\partial_t^{2K+2} \partial_{\theta_a} f$ ,

$$|\partial_t^{2K+2} \partial_{\theta_a} f(\xi)| \leq C_K \sum_{\substack{\ell_0, \dots, \ell_j \geq 0 \\ \ell_0 + \dots + \ell_j = 2K+2}} \delta^{\ell_0 + \dots + \ell_j - 1} (1 + \|\varepsilon\|)^{\ell_0 + \dots + \ell_j + 1} \leq C'_K (1 \vee \|\theta\|)^{2K+1} (1 + \|\varepsilon\|)^{2K+3}.$$

Finally, taking the expectation over  $\varepsilon$  and applying this to (A.5) for each  $a \in \{1, \dots, d\}$ , we obtain the desired bound

$$\|\nabla q(\theta)\| \leq \frac{C_K (1 \vee \|\theta\|)^{2K+1}}{\sigma^{2K+2}}.$$

The argument to bound  $\|\nabla^2 q(\theta)\|$  is similar: For any  $a, b \in \{1, \dots, d\}$ , applying a Taylor expansion of  $\partial_{\theta_a, \theta_b}^2 R(\theta)$ , we wish to bound

$$\partial_{\theta_a, \theta_b}^2 q(\theta) = \frac{t^{2K+2}}{(2K+2)!} \mathbb{E}_{h, \varepsilon} [\partial_t^{2K+2} \partial_{\theta_a, \theta_b}^2 f(\xi(h, \varepsilon))]$$

for some  $\xi(h, \varepsilon) \in (0, t)$  depending on  $a, b$ . We may compute

$$\partial_{\theta_a, \theta_b}^2 f(t) = -\frac{\partial_{\theta_a, \theta_b}^2 M(t)}{M(t)} + \frac{\partial_{\theta_a} M(t)}{M(t)} \frac{\partial_{\theta_b} M(t)}{M(t)},$$

differentiate this  $2K+2$  times in  $t$ , and apply (A.6), (A.7), and the analogous bound

$$\left| \frac{\partial_t^\ell \partial_{\theta_a, \theta_b}^2 M(\xi)}{M(\xi)} \right| \leq C_\ell \delta^{\ell-2} (1 + \|\varepsilon\|)^{\ell+2}$$

which is derived similarly. This yields  $|\partial_t^{2K+2} \partial_{\theta_a} f(\xi)| \leq C_K (1 \vee \|\theta\|)^{2K} (1 + \|\varepsilon\|)^{2K+4}$ , and taking the expectation over  $\varepsilon$  gives the desired bound for  $\|\nabla^2 q(\theta)\|$ .  $\square$

**A.2. Identifiability and transcendence degree.** We prove Propositions 2.5 and 2.6. An analogue of Proposition 2.5 for algebraic groups over algebraically closed fields may be found in e.g. [PV94, Section 2.3]; we provide here an argument in our context of a compact subgroup  $\mathbf{G} \subseteq \mathbf{O}(d)$  acting on  $\mathbb{R}^d$ .

*Proof of Proposition 2.6(a).* The algebra of all polynomials  $\mathbb{R}[\theta_1, \dots, \theta_d]$  has transcendence degree  $d$  over  $\mathbb{R}$ , so also  $\text{trdeg } \mathcal{R}^G \leq d < \infty$  for the subalgebra  $\mathcal{R}^G$ . Taking any finite transcendence basis  $\varphi$  of  $\mathcal{R}^G$  and letting  $k$  be the maximum degree of polynomials constituting  $\varphi$ , we have  $\text{trdeg } \mathcal{R}_{\leq k}^G = |\varphi| = \text{trdeg } \mathcal{R}^G$ . Then there must be a smallest integer  $k$  for which this holds.  $\square$

For the remaining statements of Propositions 2.5 and 2.6, we will use the following Jacobian criterion for algebraic independence, and a relation between generic list recovery of  $\theta_*$  and transcendence degree.

**Lemma A.2** (Jacobian criterion, [BMS13] Theorem 8). *For any real polynomials  $p_1, \dots, p_m$  in  $\theta_1, \dots, \theta_d$ ,*

$$\text{trdeg}(\{p_1, \dots, p_m\}) = \text{rank}_{\mathbb{R}(\theta_1, \dots, \theta_d)}[\nabla p_1, \dots, \nabla p_m]^\top$$

*where the right side denotes the rank of the Jacobian matrix  $[\nabla p_1, \dots, \nabla p_m]^\top \in \mathbb{R}^{m \times d}$  over the field of rational functions  $\mathbb{R}(\theta_1, \dots, \theta_d)$ .*

Note that this implies  $\text{trdeg}(\{p_1, \dots, p_m\}) \geq \text{rank}[\nabla p_1(\theta), \dots, \nabla p_m(\theta)]^\top$  (the rank over  $\mathbb{R}$ ) evaluated at any  $\theta \in \mathbb{R}^d$ , with equality holding at generic points  $\theta \in \mathbb{R}^d$ .

**Lemma A.3** (Generic list recovery, [BBSK<sup>+</sup>17] Theorem 4.9). *Let  $G \subseteq O(d)$  be a compact subgroup, and let  $U$  be a finite-dimensional subspace of  $\mathcal{R}^G$ . If  $\text{trdeg } U = \text{trdeg } \mathcal{R}^G$ , then for generic  $\theta_* \in \mathbb{R}^d$ , the set of points  $\{\theta \in \mathbb{R}^d : P(\theta) = P(\theta_*) \text{ for all } P \in U\}$  is a union of a finite number of orbits.*

(Theorem 4.9 of [BBSK<sup>+</sup>17] shows also a converse of this statement, but we will not directly use this converse.)

Recall from (2.12–2.13) the values

$$d_k = \text{trdeg } \mathcal{R}_{\leq k}^G - \text{trdeg } \mathcal{R}_{\leq k-1}^G, \quad \tilde{d}_k = \text{trdeg } \tilde{\mathcal{R}}_{\leq k}^G - \text{trdeg } \tilde{\mathcal{R}}_{\leq k-1}^G$$

and from (2.15–2.16) the combined moment functions  $M_k(\theta)$  and  $\tilde{M}_k(\theta)$ . The following structural lemma is an important ingredient for our proofs, and provides explicit coordinate systems in local neighborhoods of points  $\theta \in \mathbb{R}^d$  using  $G$ -invariant polynomials. Versions of these statements were shown in [FSWW20, Section 4] when  $G$  is a discrete group and generic orbits have dimension 0, and the following lemma provides an extension to models where orbits have positive dimension.

**Lemma A.4.** *Let  $K$  be the smallest integer for which  $\text{trdeg } \mathcal{R}_{\leq K}^G = \text{trdeg } \mathcal{R}^G$ . In the unprojected model, for generic  $\theta \in \mathbb{R}^d$  and every  $k = 1, \dots, K$ , the rank of  $dM_k(\theta)$  equals  $d_1 + \dots + d_k$ .*

*Furthermore, for any  $k \in \{1, \dots, K\}$  and any  $\theta \in \mathbb{R}^d$  where  $\text{rank } dM_k(\theta) = d_1 + \dots + d_k$ , there exist functions  $\varphi^j : \mathbb{R}^d \rightarrow \mathbb{R}^{d_j}$  for each  $j = 1, \dots, k$  and  $\bar{\varphi} : \mathbb{R}^d \rightarrow \mathbb{R}^{d-d_1-\dots-d_k}$ , such that:*

- (a) *For each  $j = 1, \dots, k$ , the  $d_j$  coordinates of  $\varphi^j$  are entries of the moment tensor  $T_j$ .*
  - (b) *The combined map  $\varphi = (\varphi^1, \dots, \varphi^k, \bar{\varphi}) : \mathbb{R}^d \rightarrow \mathbb{R}^d$  has non-singular derivative  $d\varphi(\theta) \in \mathbb{R}^{d \times d}$  at this point  $\theta$ , and hence an analytic inverse function  $\theta(\varphi)$  over a neighborhood  $U$  of  $\theta$ .*
  - (c) *For any  $j = 1, \dots, k$ , any polynomial  $p \in \mathcal{R}_{\leq j}^G$ , and sufficiently small such neighborhood  $U$ ,  $\varphi \mapsto p(\theta(\varphi))$  is a function only of the  $d_1 + \dots + d_j$  coordinates  $\varphi^1, \dots, \varphi^j$ , over  $\varphi(U)$ .*
  - (d) *Suppose  $k = K$ . Then for any  $G$ -invariant continuous function  $f : \mathbb{R}^d \rightarrow \mathbb{R}$  and sufficiently small such neighborhood  $U$ ,  $\varphi \mapsto f(\theta(\varphi))$  is a function only of the  $d_1 + \dots + d_K$  coordinates  $\varphi^1, \dots, \varphi^K$ , over  $\varphi(U)$ .*
- Also,*

$$\left\{ \theta' \in U : (\varphi^1(\theta'), \dots, \varphi^K(\theta')) = (\varphi^1(\theta), \dots, \varphi^K(\theta)) \right\} = U \cap \mathcal{O}_\theta. \quad (\text{A.8})$$

*In the projected model, suppose there exists a smallest integer  $\tilde{K} < \infty$  for which  $\text{trdeg } \tilde{\mathcal{R}}_{\leq \tilde{K}}^G = \text{trdeg } \mathcal{R}^G$ .*

*Then the same statements hold with  $\tilde{K}$ ,  $\tilde{M}_j$ ,  $\tilde{T}_j$ ,  $\tilde{d}_j$ , and  $\tilde{\mathcal{R}}_{\leq j}^G$  in place of  $K$ ,  $M_j$ ,  $T_j$ ,  $d_j$ , and  $\mathcal{R}_{\leq j}^G$ .*

*Proof.* In the unprojected model,  $\mathcal{R}_{\leq k}^G$  is generated by the entries of  $T_1, \dots, T_k$ , and  $\text{trdeg } \mathcal{R}_{\leq k}^G = d_1 + \dots + d_k$ . Thus there are exactly  $d_1 + \dots + d_k$  algebraically independent entries of  $T_1, \dots, T_k$  (c.f. [Lan02, Chapter 8, Theorem 1.1]). Then by Lemma A.2,  $dM_k(\theta)$  has rank  $d_1 + \dots + d_k$  at generic  $\theta \in \mathbb{R}^d$ , and rank at most  $d_1 + \dots + d_k$  at every  $\theta \in \mathbb{R}^d$ .

If  $\theta \in \mathbb{R}^d$  has  $\text{rank } dM_k(\theta) = d_1 + \dots + d_k$ , then this implies that also  $\text{rank } dM_j(\theta) = d_1 + \dots + d_j$  for each  $j = 1, \dots, k$ . Then for each  $j = 1, \dots, k$ , we may pick  $d_j$  entries of  $T_j$  to be  $\varphi^j$ , such that  $(\varphi^1, \dots, \varphi^k)$  have linearly independent gradients at  $\theta$ . We may arbitrarily pick  $d - d_1 - \dots - d_k$  additional analytic functions to be  $\bar{\varphi}$ , so that  $\varphi = (\varphi^1, \dots, \varphi^k, \bar{\varphi})$  has non-singular derivative  $d\varphi(\theta)$ . This shows properties (a) and (b), where the existence of an analytic inverse  $\theta(\varphi)$  on  $U$  follows from the inverse function theorem.

For (c), denote  $q(\varphi) = p(\theta(\varphi))$ . Applying the chain rule, for any  $x \in U$ ,

$$\nabla p(x) = d_x \varphi(x)^\top \nabla_\varphi q(\varphi(x)).$$

Since  $p$  is a function of  $T_1, \dots, T_j$ , its gradient is a linear combination of the gradients of the entries of  $T_1, \dots, T_j$ , and hence linearly dependent with the gradients of  $\varphi^1, \dots, \varphi^j$ . Thus  $\nabla p(x)$  belongs to the span

of the columns of  $d_x \varphi(x)^\top$  corresponding to  $\varphi^1, \dots, \varphi^j$ , implying that  $\nabla_\varphi q(\varphi(x))$  is 0 in the remaining coordinates  $\varphi^{j+1}, \dots, \varphi^k, \bar{\varphi}$ . This holds at every  $x \in U$ , so  $q$  is a function only of  $\varphi^1, \dots, \varphi^j$ .

For (d), denote  $h(\varphi) = f(\theta(\varphi))$ . Suppose first that  $f \in \mathcal{R}^G$  is a  $G$ -invariant polynomial. Since  $d_1 + \dots + d_K = \text{trdeg } \mathcal{R}_{\leq K}^G = \text{trdeg } \mathcal{R}^G$ , we must have that  $(\varphi^1, \dots, \varphi^K, f)$  are algebraically dependent. Then their gradients are linearly dependent at every  $x \in U$ . Then the same argument as in (c) shows that  $h(\varphi)$  depends only on  $\varphi^1, \dots, \varphi^K$ . For a general  $G$ -invariant continuous function  $f$ , let  $r > 0$  be large enough such that  $U \subset \overline{B_r} = \{x \in \mathbb{R}^d : \|x\| \leq r\}$ . For any  $\varepsilon > 0$ , by the Stone-Weierstrass theorem, there is a polynomial  $p$  such that  $|p(x) - f(x)| < \varepsilon$  for all  $x \in \overline{B_r}$ . Applying the Reynolds operator  $\bar{p}(x) = \int p(g \cdot x) d\Lambda(g)$ , we then have  $\bar{p} \in \mathcal{R}^G$ , and also

$$|\bar{p}(x) - f(x)| = \left| \int (p(g \cdot x) - f(g \cdot x)) d\Lambda(g) \right| < \varepsilon \quad \text{for all } x \in \overline{B_r}$$

because  $g \cdot x \in \overline{B_r}$  for any orthogonal matrix  $g$ . Writing  $\bar{q}(\varphi) = \bar{p}(\theta(\varphi))$ , we have shown that  $\bar{q}$  depends only on  $\varphi^1, \dots, \varphi^K$ . So for any  $\varphi, \varphi' \in \varphi(U)$  differing in only the coordinates of  $\bar{\varphi}$ , we have  $\bar{q}(\varphi) = \bar{q}(\varphi')$  and  $\theta(\varphi), \theta(\varphi') \in U \subset \overline{B_r}$ , hence  $|h(\varphi) - h(\varphi')| \leq |h(\varphi) - \bar{q}(\varphi)| + |h(\varphi') - \bar{q}(\varphi')| < 2\varepsilon$ . Here  $\varepsilon > 0$  is arbitrary, so in fact  $h(\varphi) = h(\varphi')$ . Thus  $h$  depends only on  $\varphi^1, \dots, \varphi^K$ .

To show (A.8), clearly if  $\mathcal{O}_{\theta'} = \mathcal{O}_\theta$ , then  $(\varphi^1(\theta'), \dots, \varphi^K(\theta')) = (\varphi^1(\theta), \dots, \varphi^K(\theta))$ . For the converse direction, if  $\mathcal{O}_{\theta'}$  and  $\mathcal{O}_\theta$  are distinct, then they are disjoint compact subsets of  $\mathbb{R}^d$ . Then there is a continuous function  $f : \mathbb{R}^d \rightarrow \mathbb{R}$  taking value 1 on  $\mathcal{O}_{\theta'}$  and 0 on  $\mathcal{O}_\theta$ . Then  $\bar{f}(x) = \int f(g \cdot x) d\Lambda(g)$  is a  $G$ -invariant continuous function with the same property. Thus  $\bar{f}$  depends only on  $\varphi^1, \dots, \varphi^K$ , implying that  $(\varphi^1(\theta'), \dots, \varphi^K(\theta')) \neq (\varphi^1(\theta), \dots, \varphi^K(\theta))$ . This shows (A.8), and concludes the proof in the unprojected setting.

The proof in the projected setting is the same, where the given condition for  $\tilde{K}$  is used in part (d) to show  $\tilde{d}_1 + \dots + \tilde{d}_{\tilde{K}} = \text{trdeg } \mathcal{R}^G$ , and hence  $(\varphi^1, \dots, \varphi^{\tilde{K}}, f)$  are algebraically dependent for any  $f \in \mathcal{R}^G$ .  $\square$

*Proof of Proposition 2.5.* Since  $G$  is a compact Lie group acting smoothly on  $\mathbb{R}^d$ , its action is proper. Then each orbit  $\mathcal{O}_\theta$  is an embedded submanifold of  $\mathbb{R}^d$ , and the tangent space to  $\mathcal{O}_\theta$  at  $\theta$  is given by  $T_\theta \mathcal{O}_\theta = \{\mathfrak{g} \cdot \theta : \mathfrak{g} \in T_{\text{Id}} G\}$  where  $T_{\text{Id}} G$  is the Lie algebra, i.e. the tangent space to  $G$  at  $g = \text{Id}$  (c.f. [Aud04, Section I.1.b and Corollary I.1.2]). Parametrizing  $G$  around  $g = \text{Id}$  by any local chart  $x$  such that  $g(x) = \text{Id}$  at  $x = 0$ , we then have  $\dim(\mathcal{O}_\theta) = \dim(\{\mathfrak{g} \cdot \theta : \mathfrak{g} \in T_{\text{Id}} G\}) = \text{rank}(d_x[g(x) \cdot \theta]_{x=0})$ . Since  $d_x[g(x) \cdot \theta]_{x=0}$  is an analytic matrix in  $\theta$ , this implies that  $\max_{\theta \in \mathbb{R}^d} \dim(\mathcal{O}_\theta)$  is attained at generic points  $\theta \in \mathbb{R}^d$  (c.f. Fact 4.5). On the other hand, for generic  $\theta \in \mathbb{R}^d$ , the statement (A.8) of Lemma A.4(d) shows that  $\dim(\mathcal{O}_\theta) = d - (d_1 + \dots + d_K) = d - \text{trdeg } \mathcal{R}_{\leq K}^G = d - \text{trdeg } \mathcal{R}^G$ . Hence  $\text{trdeg } \mathcal{R}^G = d - \max_{\theta \in \mathbb{R}^d} \dim(\mathcal{O}_\theta)$ .  $\square$

*Proof of Proposition 2.6(b).*  $\Pi(\mathcal{O}_\theta)$  is a continuous image of the compact group  $G$ , and hence is also compact. Then the distribution of  $\Pi \cdot g \cdot \theta$  is uniquely determined by its sequence of mixed moments. Thus

$$\left\{ \theta : \Pi(\mathcal{O}_\theta) \equiv \Pi(\mathcal{O}_{\theta_*}) \right\} = \left\{ \theta : \tilde{T}_k(\theta) = \tilde{T}_k(\theta_*) \text{ for all } k \geq 1 \right\}. \quad (\text{A.9})$$

Suppose that  $\text{trdeg } \tilde{\mathcal{R}}_{\leq k}^G = \text{trdeg } \mathcal{R}^G$  for some integer  $k$ . Let  $U_k$  be the linear subspace of  $\mathcal{R}^G$  spanned by 1 and all entries of  $\tilde{T}_1, \dots, \tilde{T}_k$ . Since  $U_k$  generates  $\tilde{\mathcal{R}}_{\leq k}^G$ , we have  $\text{trdeg } U_k = \text{trdeg } \tilde{\mathcal{R}}_{\leq k}^G$  (c.f. [Lan02, Chapter 8, Theorem 1.1]). Then Lemma A.3 implies that for generic  $\theta_* \in \mathbb{R}^d$ , there are only finitely many orbits  $\mathcal{O}_\theta$  such that  $P(\theta) = P(\theta_*)$  for all  $P \in U_k$ . This condition must hold for all  $\theta$  belonging to (A.9), so (A.9) also consists of finitely many orbits.

Conversely, suppose  $\text{trdeg } \tilde{\mathcal{R}}_{\leq k}^G < \text{trdeg } \mathcal{R}^G$  for all  $k \geq 1$ . Consider  $\tilde{\mathcal{R}}^G = \bigcup_{k \geq 1} \tilde{\mathcal{R}}_{\leq k}^G$  (the subalgebra generated by entries of  $\tilde{T}_k$  for all  $k \geq 1$ ). By the same argument as in the proof of Proposition 2.6(a), we have  $\text{trdeg } \tilde{\mathcal{R}}^G \leq d < \infty$ , so  $\text{trdeg } \tilde{\mathcal{R}}^G = \text{trdeg } \tilde{\mathcal{R}}_{\leq k}^G$  for some integer  $k$ . Then also  $\text{trdeg } \tilde{\mathcal{R}}^G < \text{trdeg } \mathcal{R}^G$ . We now apply an argument similar to that of the proof of Lemma A.4(c): Fix any transcendence basis  $\varphi^0$  of  $\tilde{\mathcal{R}}^G$ . Then the gradient vectors of  $\varphi^0$  are linearly independent at generic  $\theta_* \in \mathbb{R}^d$  by Lemma A.2. Fix any such  $\theta_*$ . In a sufficiently small open neighborhood  $O$  of  $\theta_*$ , we claim that

$$\left\{ \theta \in O : \tilde{T}_k(\theta) = \tilde{T}_k(\theta_*) \text{ for all } k \geq 1 \right\} = \left\{ \theta \in O : \varphi^0(\theta) = \varphi^0(\theta_*) \right\}. \quad (\text{A.10})$$

To see this, choose any  $d - |\varphi^0|$  additional functions  $\bar{\varphi}$  for which  $\varphi = (\varphi^0, \bar{\varphi})$  has non-singular derivative at  $\theta_*$ , and hence forms an invertible local reparametrization over a sufficiently small such neighborhood  $O$ , by

$$\begin{pmatrix} \sigma^{-2} & \sigma^{-4} & \sigma^{-6} & \sigma^{-8} & \dots & \sigma^{-2K} \\ \sigma^{-4} & \sigma^{-4} & \sigma^{-6} & \sigma^{-8} & \dots & \sigma^{-2K} \\ \sigma^{-6} & \sigma^{-6} & \sigma^{-6} & \sigma^{-8} & \dots & \sigma^{-2K} \\ \sigma^{-8} & \sigma^{-8} & \sigma^{-8} & \sigma^{-8} & \dots & \sigma^{-2K} \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ \sigma^{-2K} & \sigma^{-2K} & \sigma^{-2K} & \sigma^{-2K} & \dots & \sigma^{-2K} \end{pmatrix}$$

FIGURE A.1. An illustration of the block scalings with  $\sigma^{-2}$  for matrices  $H$  with graded block structure, where each  $(k, \ell)$  entry represents the scaling for a single block  $H_{k\ell}$  of  $H$ .

the inverse function theorem. Let  $p(\theta)$  be any entry of  $\tilde{T}_k$  for any  $k \geq 1$ , and write  $q(\varphi) = p(\theta(\varphi))$  for its reparametrization by the local coordinates  $\varphi$  on  $O$ . By the chain rule,

$$\nabla p(\theta) = d_\theta \varphi(\theta)^\top \nabla_\varphi q(\varphi(\theta)).$$

Since  $p \in \tilde{\mathcal{R}}^\mathbb{G}$ , and  $\varphi^0$  is a transcendence basis for  $\tilde{\mathcal{R}}^\mathbb{G}$ , we have that  $(p, \varphi^0)$  is algebraically dependent. Then the gradients of  $p$  and  $\varphi^0$  are linearly dependent at every  $\theta \in O$  by Lemma A.2, so  $\nabla p(\theta)$  belongs to the span of columns of  $d_\theta \varphi(\theta)^\top$  corresponding to only the coordinates of  $\varphi^0$ . Then  $\nabla_\varphi q(\varphi(\theta))$  must be 0 in the remaining coordinates  $\bar{\varphi}$ . This holds for all  $\theta \in O$ , so  $q(\varphi)$  is a function only of  $\varphi^0$  in this local parametrization over  $O$ . This shows our claim (A.10). The set (A.10) forms a manifold of dimension  $d - |\varphi^0| = d - \text{trdeg } \tilde{\mathcal{R}}^\mathbb{G}$ . On the other hand, Proposition 2.5 shows that every orbit  $\mathcal{O}_\theta$  has dimension at most  $d - \text{trdeg } \mathcal{R}^\mathbb{G}$ , which is strictly smaller when  $\text{trdeg } \tilde{\mathcal{R}}^\mathbb{G} < \text{trdeg } \mathcal{R}^\mathbb{G}$ . Thus (A.9) must contain infinitely many orbits corresponding to  $\theta \in O$ .  $\square$

**A.3. Fisher information.** We prove Theorem 2.7 and Lemma 2.8. Throughout, we assume that (2.4) holds in the projected setting, and we denote by  $K$  and  $\tilde{K}$  the (smallest) integers satisfying Proposition 2.6. All constants  $C, C', c, c', \sigma_0 > 0$  in the proofs may depend implicitly on  $(\theta_*, \mathbb{G}, \Pi)$ .

The proofs are analogous to the arguments of [FSWW20, Section 4.4]: Locally around any generic point  $\theta_* \in \mathbb{R}^d$ , we reparametrize  $\theta$  by a transcendence basis  $\varphi^1, \dots, \varphi^K$  for  $\mathcal{R}^\mathbb{G}$  having full-rank Jacobian, and then compute the Hessian of  $R(\theta(\varphi))$  in  $\varphi$  by taking derivatives of the series expansion (2.7) or (2.10) term-by-term. When  $\mathbb{G}$  is a continuous group, we extend this transcendence basis using the additional analytic functions  $\bar{\varphi} \in \mathbb{R}^{d_0}$  provided in Lemma A.4 to obtain a complete system of coordinates. The properties stated in Theorem 2.2 will guarantee that the Hessian of each term of order  $\sigma^{-2k}$  depends only on  $\varphi^1, \dots, \varphi^k$  in this system of coordinates, and that at the true parameter  $\theta = \theta_*$ , the block of this Hessian corresponding to the coordinates of  $\varphi^k$  is strictly positive definite. In the projected setting, this latter property uses the condition given in Theorem 2.2(b) that  $P_k(\theta_*) = 0$ . Then Theorem 2.7 will follow from the chain rule and some linear algebra.

We recall here the following definition and elementary linear-algebraic result from [FSWW20].

**Definition A.5** ([FSWW20] Definition 4.14). Let  $(\varphi^1, \dots, \varphi^K)$  be a partition of coordinates for  $\mathbb{R}^{d'}$ . Let  $H \equiv H(\sigma) \in \mathbb{R}^{d' \times d'}$  be a symmetric matrix, and write its  $K \times K$  block decomposition with respect to this partition as

$$H = \begin{pmatrix} H_{11} & \dots & H_{1K} \\ \vdots & \ddots & \vdots \\ H_{K1} & \dots & H_{KK} \end{pmatrix}.$$

The matrix  $H(\sigma)$  has a **graded block structure** with respect to this partition if there are constants  $C, c, \sigma_0 > 0$  such that for all  $\sigma > \sigma_0$  and all  $k, \ell \in \{1, \dots, K\}$  where  $\varphi^k$  and  $\varphi^\ell$  have non-zero dimension,

$$C\sigma^{-2k} \geq \lambda_{\max}(H_{kk}) \geq \lambda_{\min}(H_{kk}) \geq c\sigma^{-2k} \quad \text{and} \quad \|H_{k\ell}\| \leq C\sigma^{-2 \max(k, \ell)}$$

where  $\lambda_{\max}, \lambda_{\min}$  denote the largest and smallest eigenvalues.

A visual illustration of this structure is depicted in Figure A.1.

**Lemma A.6** ([FSWW20] Lemma 4.17). *Suppose  $H \equiv H(\sigma) \in \mathbb{R}^{d' \times d'}$  has a graded block structure with respect to  $(\varphi^1, \dots, \varphi^K)$ . Let  $d_k \geq 0$  be the dimension of  $\varphi^k$ . Let  $H_{:,k,:k}$  and  $(H^{-1})_{:,k,:k}$  be the submatrices of upper-left  $k \times k$  blocks of  $H$  and  $H^{-1}$ . Then for some constants  $C, c, \sigma_0 > 0$  and all  $\sigma > \sigma_0$ :*

- (a)  *$H$  has  $d_k$  eigenvalues belonging to  $[c\sigma^{-2k}, C\sigma^{-2k}]$  for each  $k = 1, \dots, K$ .*
- (b) *For each  $k$  where  $d_1 + \dots + d_k > 0$ ,  $\lambda_{\min}(H_{:,k,:k}) \geq c\sigma^{-2k}$ .*
- (c) *For each  $k$  where  $d_1 + \dots + d_k > 0$ ,  $\lambda_{\max}((H^{-1})_{:,k,:k}) \leq C\sigma^{2k}$ .*

Let us now fix  $d_k$  and  $\tilde{d}_k$  as the constants defined by (2.12–2.13). Recall the combined moment functions  $M_k(\theta)$  and  $\tilde{M}_k(\theta)$  from (2.15–2.16), and the moment varieties  $\mathcal{V}_k(\theta_*)$  and  $\tilde{\mathcal{V}}_k(\theta_*)$  from (2.17–2.18).

**Lemma A.7.** *In the unprojected model, fix any  $\theta_* \in \mathbb{R}^d$  and any  $k \in \{1, \dots, K\}$ . Let  $\tilde{\theta} \in \mathcal{V}_k(\theta_*)$  be such that  $\text{rank } dM_k(\tilde{\theta}) = d_1 + \dots + d_k$ , and let  $\varphi = (\varphi^1, \dots, \varphi^k, \bar{\varphi})$  be the map defined by Lemma A.4 with inverse  $\theta(\varphi)$  in a neighborhood of  $\tilde{\theta}$ . Let  $s_k(\theta)$  be as defined in (2.8). Then in the parametrization by  $\varphi$ ,*

$$\nabla_{\varphi^k}^2 s_k(\theta(\varphi)) \Big|_{\varphi=\varphi(\tilde{\theta})} \text{ has full rank } d_k \text{ and is positive definite.}$$

*In the projected model, the same statements hold for  $\tilde{\mathcal{V}}_k$ ,  $\tilde{M}_k$ ,  $\tilde{d}_k$ , and  $\tilde{s}_k$  in place of  $\mathcal{V}_k$ ,  $M_k$ ,  $d_k$ , and  $s_k$ .*

*Proof.* We focus on the unprojected model; the proof in the projected model is the same.

Since  $s_k$  is globally minimized at all points of  $\mathcal{V}_k(\theta_*)$ , we must have  $\nabla_{\varphi^k}^2 s_k(\theta(\varphi)) \Big|_{\varphi=\varphi(\tilde{\theta})} \succeq 0$ . To show this has full rank  $d_k$ , observe that  $\varphi^k$  consists of a subset of entries of  $T_k$ . Thus the corresponding  $d_k \times d_k$  submatrix of  $d_{\varphi^k} T_k$  is the identity, so  $d_{\varphi^k} T_k$  has full column rank  $d_k$ . Applying the chain rule and the observation  $T_k(\tilde{\theta}) - T_k(\theta_*) = 0$  because  $\tilde{\theta} \in \mathcal{V}_k(\theta_*)$ , we may differentiate  $s_k(\theta(\varphi))$  twice in  $\varphi^k$  to obtain

$$\nabla_{\varphi^k}^2 s_k(\theta(\varphi)) \Big|_{\varphi=\varphi(\tilde{\theta})} = \frac{1}{k!} \cdot d_{\varphi^k} T_k(\theta(\varphi))^\top d_{\varphi^k} T_k(\theta(\varphi)) \Big|_{\varphi=\varphi(\tilde{\theta})}.$$

Thus this matrix has full rank  $d_k$ . □

*Proof of Theorem 2.7.* Consider the unprojected setting of part (a). For generic  $\theta_* \in \mathbb{R}^d$ , by Lemma A.4, we have  $\text{rank } dM_K(\theta_*) = d_1 + \dots + d_K$ . Let  $\varphi = (\varphi^1, \dots, \varphi^K, \bar{\varphi})$  be the map defined by Lemma A.4, with inverse  $\theta(\varphi)$  in a neighborhood  $U$  of  $\theta_*$ . We may assume without loss of generality that

$$A = d\varphi(\theta_*) \text{ is orthogonal} \tag{A.11}$$

upon replacing each function in  $\varphi$  by a  $(\theta_*$ -dependent) linear combination of itself and its preceding functions. Note that statements (c) and (d) of Lemma A.4 continue to hold after such a replacement. We denote  $\varphi_* = \varphi(\theta_*)$ . With slight abuse of notation, we write as shorthand  $f(\varphi)$  for  $f(\theta(\varphi))$ . In particular, recalling the expansion (2.7), we denote by  $q_k(\varphi), s_k(\varphi), q(\varphi)$  the terms of this expansion parametrized by  $\varphi \in U$ .

For (a1), observe that Theorem 2.2(a) guarantees  $s_k \in \mathcal{R}_{\leq k}^G$ , so Lemma A.4(c) shows that  $s_k(\varphi)$  depends only on  $\varphi^1, \dots, \varphi^k$  in the reparametrization by  $\varphi$ . Similarly,  $q_k \in \mathcal{R}_{\leq k-1}^G$ , so  $q_k(\varphi)$  depends only on  $\varphi^1, \dots, \varphi^{k-1}$ , and  $q$  is a continuous  $G$ -invariant function, so  $q(\varphi)$  depends only on  $\varphi^1, \dots, \varphi^K$  by Lemma A.4(d). Let us decompose  $\nabla_{\varphi}^2 R(\varphi_*)$  into  $(K+1) \times (K+1)$  blocks according to the partition  $(\varphi^1, \dots, \varphi^K, \bar{\varphi})$ , of sizes  $(d_1, \dots, d_K, d_0)$ . Differentiating the expansion (2.7) term-by-term, it then follows that the entries of  $\nabla_{\varphi}^2 R(\varphi_*)$  are non-zero only in the upper-left  $K \times K$  blocks, and that the  $(k, \ell)$  block corresponding to  $\nabla_{\varphi^k, \varphi^\ell}^2 R(\varphi_*)$  has operator norm bounded above by  $C\sigma^{-2 \max(k, \ell)}$  for a constant  $C > 0$ . Furthermore, as  $\nabla_{\varphi^k}^2 q_k(\varphi) = 0$  and  $\nabla_{\varphi^k}^2 s_k(\varphi_*)$  is strictly positive definite by Lemma A.7 (applied with  $\tilde{\theta} = \theta_*$ ), there are constants  $c, \sigma_0 > 0$  such that for all  $\sigma > \sigma_0$ ,

$$\lambda_{\min} \left( \nabla_{\varphi^k}^2 R(\varphi_*) \right) \geq c\sigma^{-2k} > 0 \text{ for all } k = 1, \dots, K. \tag{A.12}$$

So the upper-left  $K \times K$  blocks of  $\nabla_{\varphi}^2 R(\varphi_*)$  have the graded block structure of Definition A.5 with  $d' = d - d_0$ . Then by Lemma A.6(a),  $\nabla_{\varphi}^2 R(\varphi_*)$  has  $d_0$  eigenvalues equal to 0, and  $d_k$  eigenvalues in  $[c\sigma^{-2k}, C\sigma^{-2K}]$  for each  $k = 1, \dots, K$ . For the Hessian in  $\theta$  rather than in  $\varphi$ , since  $\nabla_{\theta} R(\theta_*) = 0$ , we have by the chain rule

$$I(\theta_*) = \nabla_{\theta}^2 R(\theta_*) = A^\top \cdot \nabla_{\varphi}^2 R(\varphi_*) \cdot A, \quad A = d\varphi(\theta_*). \tag{A.13}$$

By the orthogonality of  $A$  assumed in (A.11), the eigenvalues of  $I(\theta_*)$  are the same as those of  $\nabla_{\varphi}^2 R(\varphi_*)$ , and this shows (a1).

For (a2), observe from (A.13) that the subspace  $V_k$  spanned by the  $d_1 + \dots + d_k$  leading eigenvectors of  $I(\theta_*)$  is given by  $V_k = A^\top \cdot V_{k,\varphi}$ , where  $V_{k,\varphi}$  is the subspace spanned by the  $d_1 + \dots + d_k$  leading eigenvectors of  $\nabla_\varphi^2 R(\varphi_*)$ . By Lemma A.6(b), the submatrix of upper-left  $k \times k$  blocks of  $\nabla_\varphi^2 R(\varphi_*)$  has smallest eigenvalue at least  $c\sigma^{-2k}$ , while the remaining blocks have operator norm at most  $C\sigma^{-2(k+1)}$  by Definition A.5. Let  $W_{k,\varphi}$  be the subspace of vectors having only first  $d_1 + \dots + d_k$  coordinates non-zero. Then the Davis-Kahan theorem implies that, for a function  $\varepsilon(\sigma)$  satisfying  $\varepsilon(\sigma) \rightarrow 0$  as  $\sigma \rightarrow \infty$ ,

$$\|\sin \Theta(V_{k,\varphi}, W_{k,\varphi})\| < \varepsilon(\sigma).$$

For any polynomial  $p \in \mathcal{R}_{\leq k}^G$ , we have  $\nabla_{\theta} p(\theta_*) = A^\top \nabla_\varphi p(\varphi_*)$  by the chain rule. Lemma A.4(d) shows that  $p$  depends only on  $\varphi^1, \dots, \varphi^k$  in the parametrization by  $\varphi$ , so  $\nabla_\varphi p(\varphi_*) \in W_{k,\varphi}$ , and hence  $\nabla_{\theta} p(\theta_*) \in A^\top \cdot W_{k,\varphi}$ . For generic  $\theta_* \in \mathbb{R}^d$ , the linear span  $W_k$  of all such gradient vectors  $\nabla_{\theta} p(\theta_*)$  has dimension exactly  $d_1 + \dots + d_k$  by Lemma A.2, so this shows  $W_k = A^\top \cdot W_{k,\varphi}$ . Thus also

$$\|\sin \Theta(V_k, W_k)\| = \|\sin \Theta(V_{k,\varphi}, W_{k,\varphi})\| < \varepsilon(\sigma).$$

For (a3), observe that since  $A$  is orthogonal, we have from (A.13) that  $I(\theta_*)^\dagger = A^\top \cdot \nabla_\varphi^2 R(\varphi_*)^\dagger \cdot A$  for the Moore-Penrose pseudo-inverse. Combining this with  $\nabla_{\theta} p(\theta_*) = A^\top \nabla_\varphi p(\varphi_*)$ , we have

$$\nabla_{\theta} p(\theta_*)^\top I(\theta_*)^\dagger \nabla_{\theta} p(\theta_*) = \nabla_\varphi p(\varphi_*)^\top \cdot \nabla_\varphi^2 R(\varphi_*)^\dagger \cdot \nabla_\varphi p(\varphi_*).$$

Lemma A.6(c) shows that the maximum eigenvalue of the upper-left  $k \times k$  blocks of  $\nabla_\varphi^2 R(\varphi_*)^\dagger$  is at most  $C\sigma^{2k}$ . Since  $\nabla_\varphi p(\varphi_*)$  is non-zero only in its first  $k$  blocks, this implies

$$\nabla_{\theta} p(\theta_*)^\top I(\theta_*)^\dagger \nabla_{\theta} p(\theta_*) \leq C\sigma^{2k}.$$

Finally, if  $w$  is in the null space of  $I(\theta_*)$ , then  $Aw$  is in the null space of  $\nabla_\varphi^2 R(\varphi_*)$ , i.e.  $Aw$  is non-zero only in the last block corresponding to  $\bar{\varphi}$ . Then  $\nabla_{\theta} p(\theta_*)^\top w = \nabla_\varphi p(\varphi_*)^\top Aw = 0$ , so  $\nabla_{\theta} p(\theta_*)$  is orthogonal to the null space of  $I(\theta_*)$ . This shows (a3).

The proof of part (b) in the projected setting is similar: We let  $\varphi = (\varphi^1, \dots, \varphi^{\tilde{K}}, \bar{\varphi})$  be the map defined by Lemma A.4, and compute the Hessian of (2.10) in the parametrization by  $\varphi$  term-by-term. In this computation, there is an additional contribution from each term  $\sigma^{-2k} \langle \tilde{T}_k(\varphi), P_k(\varphi) \rangle$ . This term depends only on  $\varphi^1, \dots, \varphi^k$ , so its Hessian lies only in the upper-left  $k \times k$  blocks of the  $(\tilde{K} + 1) \times (\tilde{K} + 1)$  block decomposition of  $\nabla_\varphi^2 R(\varphi_*)$ . Hence each block  $\nabla_{\varphi^k, \varphi^\ell}^2 R(\varphi_*)$  still has operator norm bounded above by  $C\sigma^{-2 \max(k, \ell)}$ . Furthermore, the Hessian of  $\sigma^{-2k} \langle \tilde{T}_k(\varphi), P_k(\varphi) \rangle$  is the sum of three terms, corresponding to differentiating twice  $\tilde{T}_k(\varphi)$ , twice  $P_k(\varphi)$ , and once each  $\tilde{T}_k(\varphi)$  and  $P_k(\varphi)$ . The first term vanishes upon evaluating at  $\varphi = \varphi_*$ , because  $P_k(\varphi_*) = 0$  by its characterization in Theorem 2.2(b). The remaining two terms are 0 on the  $(k, k)$  block, because  $P_k(\varphi)$  depends only on  $\varphi^1, \dots, \varphi^{k-1}$ . Thus the Hessian of  $\sigma^{-2k} \langle \tilde{T}_k(\varphi), P_k(\varphi) \rangle$  at  $\varphi = \varphi_*$  is 0 in the  $(k, k)$  block, so we still have  $\nabla_{\varphi^k}^2 R(\varphi_*) \succeq c\sigma^{-2k}$ . Then the upper-left  $\tilde{K} \times \tilde{K}$  blocks of  $\nabla_\varphi^2 R(\varphi_*)$  still have the graded block structure of Definition A.5, and the remainder of the proof is the same as in the unprojected setting of part (a).  $\square$

*Proof of Lemma 2.8.* We focus on the unprojected setting; the proof in the projected setting is the same.

Note that  $\theta_*$  is a global minimizer of  $s_k(\theta)$ , so  $\nabla_{\theta} s_k(\theta_*) = 0$  and  $\nabla_{\theta}^2 s_k(\theta_*) \succeq 0$ . For generic  $\theta_*$ , we have  $\text{rank } dM_K(\theta_*) = d_1 + \dots + d_K$  by Lemma A.4. Let  $\varphi = (\varphi^1, \dots, \varphi^K, \bar{\varphi})$  be the map defined by Lemma A.4, with inverse  $\theta(\varphi)$  in a neighborhood  $U$  of  $\theta_*$ . Let  $\varphi_* = \varphi(\theta_*)$ . Then by the chain rule,

$$\nabla_{\varphi}^2 s_k(\theta(\varphi))|_{\varphi=\varphi_*} = d_{\varphi} \theta(\varphi_*)^\top \cdot \nabla_{\theta}^2 s_k(\theta_*) \cdot d_{\varphi} \theta(\varphi_*). \quad (\text{A.14})$$

Since  $d_{\varphi} \theta(\varphi_*)$  is non-singular, this yields

$$\text{rank} \left( \nabla_{\theta}^2 s_1(\theta_*) + \dots + \nabla_{\theta}^2 s_k(\theta_*) \right) = \text{rank} \left( \nabla_{\varphi}^2 s_1(\theta(\varphi)) + \dots + \nabla_{\varphi}^2 s_k(\theta(\varphi)) \Big|_{\varphi=\varphi_*} \right).$$

Lemma A.4(c) ensures that  $s_1(\theta(\varphi)), \dots, s_k(\theta(\varphi))$  depend only on  $\varphi^1, \dots, \varphi^k$ , so

$$\text{rank} \left( \nabla_{\varphi}^2 s_1(\theta(\varphi)) + \dots + \nabla_{\varphi}^2 s_k(\theta(\varphi)) \Big|_{\varphi=\varphi_*} \right) \leq d_1 + \dots + d_k = \text{trdeg}(\mathcal{R}_{\leq k}^G).$$

To show that this holds with equality, consider any non-zero vector  $v = (v_1, \dots, v_k, 0, \dots, 0) \in \mathbb{R}^d$ , where  $v_j$  is the subvector corresponding to the coordinates of  $\varphi^j$ . Let  $j \in \{1, \dots, k\}$  be the smallest index for which  $v_j \neq 0$ . Lemma A.7 applied with  $\tilde{\theta} = \theta_*$  shows that  $\nabla_{\varphi^j}^2 s_j(\theta(\varphi))|_{\varphi=\varphi_*} \succ 0$  strictly, so

$$v^\top \left[ \nabla_{\varphi}^2 s_j(\theta(\varphi)) \Big|_{\varphi=\varphi_*} \right] v = v_j^\top \left[ \nabla_{\varphi^j}^2 s_j(\theta(\varphi)) \Big|_{\varphi=\varphi_*} \right] v_j > 0$$

where the first equality holds because  $v_1 = \dots = v_{j-1} = 0$ , whereas  $s_j(\theta(\varphi))$  depends only on  $\varphi^1, \dots, \varphi^j$ . Furthermore,  $v^\top [\nabla_{\varphi}^2 s_i(\theta(\varphi))|_{\varphi=\varphi_*}] v \geq 0$  for all  $i \neq j$ , because (A.14) and the condition  $\nabla_{\theta}^2 s_i(\theta_*) \succeq 0$  imply that  $\nabla_{\varphi}^2 s_i(\theta(\varphi))|_{\varphi=\varphi_*}$  is positive semidefinite. Then  $v^\top [\nabla_{\varphi}^2 s_1(\theta(\varphi)) + \dots + \nabla_{\varphi}^2 s_k(\theta(\varphi))|_{\varphi=\varphi_*}] v > 0$  strictly. This holds for every non-zero vector  $v = (v_1, \dots, v_k, 0, \dots, 0) \in \mathbb{R}^d$ , so in fact

$$\text{rank} \left( \nabla_{\varphi}^2 s_1(\theta(\varphi)) + \dots + \nabla_{\varphi}^2 s_k(\theta(\varphi)) \Big|_{\varphi=\varphi_*} \right) = d_1 + \dots + d_k = \text{trdeg}(\mathcal{R}_{\leq k}^{\mathcal{E}}).$$

This shows also that the column span of  $\nabla_{\varphi}^2 s_1(\theta(\varphi)) + \dots + \nabla_{\varphi}^2 s_k(\theta(\varphi))|_{\varphi=\varphi_*}$  is exactly the space of vectors  $v$  with only its first  $d_1 + \dots + d_k$  coordinates non-zero. Then by (A.14), the column span of  $\nabla_{\theta}^2 s_1(\theta_*) + \dots + \nabla_{\theta}^2 s_k(\theta_*)$  is the span of the first  $d_1 + \dots + d_k$  rows of  $d_{\varphi} \theta(\varphi_*)^{-1} = d_{\theta} \varphi(\theta_*)$ , which are the gradients of  $\varphi^1(\theta_*), \dots, \varphi^k(\theta_*)$ . By Lemma A.4, the span of these gradients is exactly the span of  $\{\nabla p(\theta_*) : p \in \mathcal{R}_{\leq k}^{\mathcal{E}}\}$ , concluding the proof.  $\square$

**A.4. Global landscape.** We prove Theorems 2.11 and 2.13. The following lemma first shows that all critical points of  $R(\theta)$  described by Theorems 2.11 and 2.13 in fact belong to a ball of constant radius  $M > 0$ , independent of  $\sigma$ . In unprojected models, this result was proven in [FSWW20, Lemmas 2.10 and 4.19]. The argument is reviewed and extended in the proof below, to the domain  $\{\theta : \|\theta\| < B(\|\theta_*\| + \sigma)\}$  for projected models under the assumption (2.4) for the projection  $\Pi$ .

**Lemma A.8.** *In the unprojected model, for some constants  $M, c, \sigma_0 > 0$  depending on  $\theta_*, \mathbf{G}$  and for all  $\sigma > \sigma_0$ ,*

$$\|\nabla R(\theta)\| \geq c\sigma^{-4} \quad \text{for all } \theta \text{ satisfying } \|\theta\| > M.$$

*In the projected model with projection  $\Pi$ , for any  $B > 0$ , some constants  $M, c, \sigma_0 > 0$  depending on  $\theta_*, \mathbf{G}, \Pi, B$ , and all  $\sigma > \sigma_0$ ,*

$$\|\nabla R(\theta)\| \geq c\sigma^{-4} \quad \text{for all } \theta \text{ satisfying } B(\|\theta_*\| + \sigma) > \|\theta\| > M.$$

*Proof. Step 1: Forms of  $\nabla R(\theta)$ .* We consider the projected model, which will reduce to the unprojected model when  $\Pi = \text{Id}$ . Write  $\mathbb{E}_g$  and  $\mathbb{E}_{g, g'}$  for expectations over independent group elements  $g, g' \sim \Lambda$ , and  $\mathbb{E}_Y$  for that over the sample  $Y \sim p_{\theta_*}$ . Introduce the weight

$$p(g, Y) = \frac{\exp(-\frac{1}{2\sigma^2} \|Y - \Pi g \theta\|^2)}{\mathbb{E}_{g'} [\exp(-\frac{1}{2\sigma^2} \|Y - \Pi g' \theta\|^2)]} = \frac{\exp(\frac{1}{\sigma^2} Y^\top \Pi g \theta - \frac{1}{2\sigma^2} \|\Pi g \theta\|^2)}{\mathbb{E}_{g'} [\exp(\frac{1}{\sigma^2} Y^\top \Pi g' \theta - \frac{1}{2\sigma^2} \|\Pi g' \theta\|^2)]}.$$

Then

$$\begin{aligned} \sigma^2 \nabla R(\theta) &= \sigma^2 \nabla_{\theta} \mathbb{E}_Y [-\log p_{\theta}(Y)] = -\sigma^2 \nabla_{\theta} \mathbb{E}_Y \left[ \log \mathbb{E}_g \left[ \exp \left( -\frac{\|Y - \Pi g \theta\|^2}{2\sigma^2} \right) \right] \right] \\ &= -\mathbb{E}_Y [\mathbb{E}_g [p(g, Y) g^\top \Pi^\top (Y - \Pi g \theta)]] . \end{aligned} \quad (\text{A.15})$$

We derive a second alternative form for  $\sigma^2 \nabla R(\theta)$  using Gaussian integration by parts. Let us represent  $Y = \Pi h \theta_* + \sigma \varepsilon$ , where  $h \sim \Lambda$  and  $\varepsilon \sim \mathcal{N}(0, \text{Id})$ , and write  $\mathbb{E}_Y = \mathbb{E}_{h, \varepsilon}$ . It follows from (A.15) that

$$\begin{aligned} \sigma^2 \nabla R(\theta) &= -\mathbb{E}_{h, \varepsilon} [\mathbb{E}_g [p(g, Y) g^\top \Pi^\top \Pi h \theta_*]] + \mathbb{E}_{h, \varepsilon} [\mathbb{E}_g [p(g, Y) g^\top \Pi^\top \Pi g \theta]] \\ &\quad - \mathbb{E}_{h, \varepsilon} [\mathbb{E}_g [p(g, Y) g^\top \Pi^\top (\sigma \varepsilon)]] . \end{aligned} \quad (\text{A.16})$$

For the third term above, applying the integration by parts identity  $\mathbb{E}_{\xi \sim \mathcal{N}(0, 1)} [\xi f(\xi)] = \mathbb{E}_{\xi \sim \mathcal{N}(0, 1)} [f'(\xi)]$  to each coordinate of  $\varepsilon$ , we have for  $Y = \Pi h \theta_* + \sigma \varepsilon$  and any fixed  $h$ ,

$$\mathbb{E}_{\varepsilon} [\mathbb{E}_g [p(g, Y) g^\top \Pi^\top (\sigma \varepsilon)]] = \sigma \mathbb{E}_{\varepsilon} [\mathbb{E}_g [g^\top \Pi^\top \nabla_{\varepsilon} p(g, Y)]] .$$

Explicitly differentiating  $p(g, Y) = p(g, \Pi h \theta_* + \sigma \varepsilon)$  in  $\varepsilon$  gives

$$\nabla_{\varepsilon} p(g, Y) = \sigma \nabla_Y p(g, Y) = \frac{1}{\sigma} \left( p(g, Y) \Pi g \theta - p(g, Y) \mathbb{E}_{g'} [p(g', Y) \Pi g' \theta] \right).$$

Thus,

$$\mathbb{E}_\varepsilon [\mathbb{E}_g [p(g, Y)g^\top \Pi^\top (\sigma\varepsilon)]] = \mathbb{E}_\varepsilon [\mathbb{E}_g [p(g, Y)g^\top \Pi^\top \Pi g\theta] - \mathbb{E}_{g, g'} [p(g, Y)p(g', Y)g^\top \Pi^\top \Pi g'\theta]].$$

Then, taking the expectation also over  $h \sim \Lambda$  and substituting this for the third term in (A.16),

$$\sigma^2 \nabla R(\theta) = \mathbb{E}_{h, \varepsilon} [\mathbb{E}_{g, g'} [p(g, Y)p(g', Y)g^\top \Pi^\top \Pi g'\theta] - \mathbb{E}_g [p(g, Y)g^\top \Pi^\top \Pi h\theta_*]]. \quad (\text{A.17})$$

The expressions (A.15) and (A.17) hold also in the unprojected model upon substituting  $\Pi = \text{Id}$ , where they may be further reduced to [FSWW20, Eqs. (2.8–2.9)].

**Step 2: Gradient bound for  $\|\theta\| \geq B(\|\theta_*\| + \sigma)$ .** In the unprojected model, fixing a sufficiently large constant  $B > 0$ , let us first derive a bound  $\|\nabla R(\theta)\| \geq c\sigma^{-1}$  for  $\|\theta\| \geq B(\|\theta_*\| + \sigma)$ . (This is the same argument as in [FSWW20, Lemma 2.9], which for convenience we reproduce here.) Restricting to  $\Pi = \text{Id}$  and taking the inner-product of (A.15) with  $\theta$ ,

$$\sigma^2 \|\theta\| \cdot \|\nabla R(\theta)\| \geq \sigma^2 \theta^\top \nabla R(\theta) = \|\theta\|^2 - \mathbb{E}_Y [\mathbb{E}_g [p(g, Y)\theta^\top g^\top Y]] \geq \|\theta\|^2 - C(\|\theta_*\| + \sigma)\|\theta\|$$

for a constant  $C = C(\theta_*, \mathbf{G}) > 0$ . Then for sufficiently large  $B > 0$  and large  $\sigma$ , this shows  $\|\nabla R(\theta)\| \geq c\sigma^{-1}$  as claimed.

**Step 3: Gradient bound for  $\|\theta\| \geq C_0\sigma^{2/3}$ .** We now show the lower bound  $\|\nabla R(\theta)\| \geq c\sigma^{-2}$  when  $B(\|\theta_*\| + \sigma) \geq \|\theta\| \geq C_0\sigma^{2/3}$ , for a large enough constant  $C_0 > 0$  and  $\sigma > \sigma_0(\theta_*, \mathbf{G}, \Pi, B)$ . The bound in the unprojected model follows from specializing to  $\Pi = \text{Id}$ .

Define unit vectors  $\bar{\theta} = \theta/\|\theta\|$  and  $\bar{Y} = Y/\|Y\|$ . Now taking the inner-product of (A.17) with  $\bar{\theta}$ ,

$$\begin{aligned} \sigma^2 \|\nabla R(\theta)\| &\geq \sigma^2 \cdot \bar{\theta}^\top \nabla R(\theta) \\ &= \|\theta\| \cdot \mathbb{E}_{h, \varepsilon} [\|\mathbb{E}_g [p(g, Y)\Pi g\bar{\theta}]\|^2] - \bar{\theta}^\top \mathbb{E}_{h, \varepsilon} [\mathbb{E}_g [p(g, Y)g^\top \Pi^\top \Pi h]] \cdot \theta_* \\ &\geq \|\theta\| \cdot \mathbb{E}_{h, \varepsilon} [(\bar{Y}^\top \mathbb{E}_g [p(g, Y)\Pi g\bar{\theta}])^2] - \|\Pi\|^2 \cdot \|\theta_*\|. \end{aligned}$$

For fixed  $\theta$  and  $Y$ , define

$$K(t) = \log \mathbb{E}_g \left[ \exp \left( t \bar{Y}^\top \Pi g\bar{\theta} - \frac{1}{2\sigma^2} \|\Pi g\bar{\theta}\|^2 \right) \right], \quad t(Y, \theta) = \frac{\|Y\| \cdot \|\theta\|}{\sigma^2}.$$

Then  $\bar{Y}^\top \mathbb{E}_g [p(g, Y)\Pi g\bar{\theta}] = \mathbb{E}_g [p(g, Y)\bar{Y}^\top \Pi g\bar{\theta}] = K'(t(Y, \theta))$ , so

$$\sigma^2 \|\nabla R(\theta)\| \geq \|\theta\| \cdot \mathbb{E}_{h, \varepsilon} [K'(t(Y, \theta))^2] - \|\Pi\|^2 \cdot \|\theta_*\|. \quad (\text{A.18})$$

Define a tilted probability distribution  $\bar{\Lambda}$  for  $g \in \mathbf{G}$ , having density  $d\bar{\Lambda}(g) \propto \exp(-\frac{\|\Pi g\bar{\theta}\|^2}{2\sigma^2})d\Lambda(g)$  with respect to the Haar measure  $\Lambda$ . Observe that  $K(t) - K(0)$  is the cumulant generating function for the law of  $\bar{Y}^\top \Pi g\bar{\theta}$  (fixing  $\bar{Y}$ ) that is induced by  $g \sim \bar{\Lambda}$ . We proceed to analyze the cumulants of  $\bar{Y}^\top \Pi g\bar{\theta}$  under this law. This is simpler in the unprojected setting of  $\Pi = \text{Id}$ , where  $\bar{\Lambda} = \Lambda$  and  $\bar{Y}^\top \Pi g\bar{\theta} = \bar{Y}^\top g\bar{\theta}$ ; in this setting, upper and lower bounds for the cumulants were established in the proof of [FSWW20, Lemma 2.10]. Here, we extend these bounds to the setting of a general projection  $\Pi$  that satisfies (2.4).

Note that for  $\|\theta\| < B(\|\theta_*\| + \sigma)$  and  $\sigma > \sigma_0$ , we have

$$\frac{d\bar{\Lambda}}{d\Lambda}(g) \in [c, C] \quad (\text{A.19})$$

for some  $(\theta_*, \mathbf{G}, \Pi, B)$ -dependent constants  $C, c, \sigma_0 > 0$ . The random variable  $\bar{Y}^\top \Pi g\bar{\theta}$  is bounded as  $|\bar{Y}^\top \Pi g\bar{\theta}| \leq \|\Pi\|$ , so for  $|t| < 1/(\|\Pi\|e)$ ,  $K(t) - K(0)$  is defined equivalently by the convergent cumulant series

$$K(t) - K(0) = \sum_{\ell \geq 1} \kappa_\ell(\bar{Y}^\top \Pi g\bar{\theta}) \frac{t^\ell}{\ell!}.$$

Here,  $\kappa_\ell = \kappa_\ell(\bar{Y}^\top \Pi g\bar{\theta})$  is the  $\ell^{\text{th}}$  cumulant of  $\bar{Y}^\top \Pi g\bar{\theta}$  under its law induced by  $g \sim \bar{\Lambda}$ , satisfying  $|\kappa_\ell| \leq (\|\Pi\|e)^\ell$  (c.f. [FSWW20, Lemma A.1]). In particular,  $\kappa_1$  is the mean and  $\kappa_2$  is the variance. Then for any

$0 < t < 1/(\|\Pi\|e)$ , applying also  $\ell! \geq (\ell/e)^\ell$  and convexity of the cumulant generating function  $K(t)$ ,

$$K'(t) \geq \frac{K(t) - K(0)}{t} = \sum_{\ell \geq 1} \kappa_\ell \frac{t^{\ell-1}}{\ell!} \geq \kappa_1 + \frac{t}{2} \kappa_2 - \sum_{\ell \geq 3} (\|\Pi\|e)^\ell t^{\ell-1}. \quad (\text{A.20})$$

We now lower-bound the mean and variance  $\kappa_1, \kappa_2$  when  $\bar{Y}$  belongs to some “good” subset  $U$  of the unit sphere: First note that for any non-zero  $\theta \in \mathbb{R}^d$ ,  $\Pi g \theta$  cannot be identically 0 over all  $g \in \mathbf{G}$ . This is because otherwise,  $\Pi(\mathcal{O}_{\theta_*}) \equiv \Pi(\mathcal{O}_{\theta_* + c\theta})$  for any  $\theta_* \in \mathbb{R}^d$  and any  $c \in \mathbb{R}$ , where  $\{\mathcal{O}_{\theta_* + c\theta} : c \in \mathbb{R}\}$  is an infinite family of distinct orbits, violating (2.4). Thus, denoting by  $\mathcal{S}^{d-1}$  the unit sphere in  $\mathbb{R}^d$ ,

$$\sup_{\bar{y}: \|\bar{y}\|=1} \sup_{g \in \mathbf{G}} \bar{y}^\top \Pi g \bar{\theta} > 0 \text{ for all } \bar{\theta} \in \mathcal{S}^{d-1}$$

By continuity of the left side as a function of  $\bar{\theta}$  and by compactness of  $\mathcal{S}^{d-1}$ , there is then a constant  $c = c(\Pi, \mathbf{G}) > 0$  such that

$$\sup_{\bar{y}: \|\bar{y}\|=1} \sup_{g \in \mathbf{G}} \bar{y}^\top \Pi g \bar{\theta} > c \text{ for all } \bar{\theta} \in \mathcal{S}^{d-1}.$$

Let  $\Gamma$  denote the uniform probability measure on  $\mathcal{S}^{d-1}$ . Since  $\{(\bar{y}, g) : \bar{y}^\top \Pi g \bar{\theta} > c\}$  is an open subset of  $\mathcal{S}^{d-1} \times \mathbf{G}$ , the above implies

$$\Gamma \times \Lambda\left((\bar{y}, g) : \bar{y}^\top \Pi g \bar{\theta} > c\right) > 0 \text{ for all } \bar{\theta} \in \mathcal{S}^{d-1}.$$

By the bounded convergence theorem and lower-semicontinuity of  $x \mapsto \mathbf{1}\{x > c\}$ , if  $\bar{\theta}_k \in \mathcal{S}^{d-1}$  is a sequence converging to  $\bar{\theta} \in \mathcal{S}^{d-1}$ , then

$$\liminf_{k \rightarrow \infty} \Gamma \times \Lambda\left((\bar{y}, g) : \bar{y}^\top \Pi g \bar{\theta}_k > c\right) = \mathbb{E}_{\bar{y}, g \sim \Gamma \times \Lambda} \left[ \liminf_{k \rightarrow \infty} \mathbf{1}\{\bar{y}^\top \Pi g \bar{\theta}_k > c\} \right] \geq \Gamma \times \Lambda\left((\bar{y}, g) : \bar{y}^\top \Pi g \bar{\theta} > c\right).$$

Thus  $\bar{\theta} \mapsto \Gamma \times \Lambda((\bar{y}, g) : \bar{y}^\top \Pi g \bar{\theta} > c)$  is lower-semicontinuous on  $\mathcal{S}^{d-1}$ , so again by compactness of  $\mathcal{S}^{d-1}$ , there is a constant  $\delta = \delta(\Pi, \mathbf{G}) > 0$  such that

$$\Gamma \times \Lambda\left((\bar{y}, g) : \bar{y}^\top \Pi g \bar{\theta} > c\right) > \delta \text{ for all } \bar{\theta} \in \mathcal{S}^{d-1}.$$

Then by (A.19), for a constant  $\delta' = \delta'(\theta_*, \Pi, \mathbf{G}, B)$ , we get  $\Gamma \times \bar{\Lambda}((\bar{y}, g) : \bar{y}^\top \Pi g \bar{\theta} > c) > \delta'$ . Define the  $\bar{\theta}$ -dependent subset of the unit sphere

$$U' = \left\{ \bar{y} : \bar{\Lambda}(g \in \mathbf{G} : \bar{y}^\top \Pi g \bar{\theta} > c) > \delta'/2 \right\}.$$

Then the above implies  $\Gamma(U') + (\delta'/2)(1 - \Gamma(U')) > \delta'$ , so  $\Gamma(U') > \delta'/2$ . If any random variable  $X \in \mathbb{R}$  satisfies  $\mathbb{P}[X > c] > \delta'/2$  for constants  $c, \delta' > 0$ , then  $\max(\mathbb{E}[X], \text{Var}[X]) > c'$  for a constant  $c' = c'(c, \delta') > 0$ , and furthermore either  $\mathbb{E}[X] \geq 0$  or  $\mathbb{E}[-X] \geq 0$ . Thus, defining  $U$  from  $U'$  by multiplying each element  $\bar{y} \in U'$  by an appropriate choice of  $\pm$  sign, we obtain  $\Gamma(U) > \delta'/4$  and

$$\max\left(\kappa_1(\bar{Y}^\top \Pi g \bar{\theta}), \kappa_2(\bar{Y}^\top \Pi g \bar{\theta})\right) > c' \text{ and } \kappa_1(\bar{Y}^\top \Pi g \bar{\theta}), \kappa_2(\bar{Y}^\top \Pi g \bar{\theta}) \geq 0 \text{ whenever } \bar{Y} \in U. \quad (\text{A.21})$$

Recalling that  $Y = \Pi h \theta_* + \sigma \varepsilon$  and  $\bar{Y} = Y/\|Y\|$ , the law of  $\bar{Y}$  converges to the uniform measure  $\Gamma$  on the sphere as  $\sigma \rightarrow \infty$ . Thus, for  $\sigma > \sigma_0(\Pi, \mathbf{G}, \theta_*, B)$ , we have  $\mathbb{P}[\bar{Y} \in U] \geq \Gamma(U)/2 > \delta'/8$ . For a constant  $C_0 = C_0(\Pi, \theta_*, \delta', c') > 0$  large enough and to be determined, if  $\|\theta\| \geq C_0 \sigma^{2/3}$ , then

$$t(Y, \theta) = \frac{\|Y\| \cdot \|\theta\|}{\sigma^2} = \frac{\|\Pi h \theta_* + \sigma \varepsilon\| \cdot \|\theta\|}{\sigma^2} \geq \sigma^{-1/3}$$

with probability at least  $1 - \delta'/16$ . Then, on an event of probability at least  $\delta'/16$  where both  $\bar{Y} \in U$  and  $t(Y, \theta) \geq \sigma^{-1/3}$ , and for  $\sigma > \sigma_0(\Pi, \mathbf{G}, \theta_*, B)$ , we have

$$K'(t(Y, \theta)) \geq K'(\sigma^{-1/3}) \geq (c'/3)\sigma^{-1/3},$$

the first inequality applying convexity of  $K(t)$  and the second applying (A.20) with the bound (A.21). Then, applying this to (A.18),

$$\sigma^2 \|\nabla R(\theta)\|^2 \geq C_0 \sigma^{2/3} \cdot (\delta'/16) \cdot (c'/3)^2 \sigma^{-2/3} - \|\Pi\|^2 \cdot \|\theta_*\|.$$

Here, the constants  $c', \delta' > 0$  are as defined in the argument leading to (A.21), and do not depend on  $C_0$ . Then taking  $C_0 = C_0(\Pi, \theta_*, \delta', c')$  large enough ensures that  $\|\nabla R(\theta)\|^2 \geq c\sigma^{-2}$  as desired.

**Step 4: Gradient bound for  $\|\theta\| > M$ .** Finally, we show  $\|\nabla R(\theta)\| \geq c\sigma^{-4}$  for  $C_0\sigma^{2/3} \geq \|\theta\| > M$  and a sufficiently large constant  $M > 0$ . This is again simpler in the unprojected setting of  $\Pi = \text{Id}$ , and was shown in [FSWW20, Lemma 4.19]. Here, we extend the argument to the projected model using the series expansion of Theorem 2.2(b), and specializing to  $\Pi = \text{Id}$  again recovers the result in the unprojected setting.

Define  $v = (\theta - \theta_*)/\|\theta - \theta_*\|$ , and suppose first that  $\|\mathbb{E}_g[\Pi g v]\| \geq c_0$  for any  $(\Pi, \mathbf{G}, \theta_*)$ -dependent constant  $c_0 > 0$ . We apply the expansion (2.10) to the order  $K = 1$ . Noting that  $P_1(\theta) = 0$  (because it is a constant that is 0 at  $\theta = \theta_*$ ) and  $q_1(\theta)$  is a constant,

$$\nabla R(\theta) = \frac{1}{\sigma^2} \nabla \tilde{s}_1(\theta) + \nabla q(\theta), \quad \|\nabla q(\theta)\| \leq \frac{C\|\theta\|^3}{\sigma^4}.$$

Applying the form of  $\tilde{s}_1(\theta)$  in Lemma A.1,

$$\nabla R(\theta) = \frac{1}{\sigma^2} \mathbb{E}_{g,h}[g^\top \Pi^\top \Pi h](\theta - \theta_*) + \nabla q(\theta).$$

Then

$$\sigma^2 \|\nabla R(\theta)\| \geq \sigma^2 v^\top \nabla R(\theta) \geq \|\theta - \theta_*\| \cdot \|\mathbb{E}_g[\Pi g v]\|^2 - C\|\theta\|^3/\sigma^2.$$

When  $C_0\sigma^{2/3} \geq \|\theta\| > M$ ,  $\|\mathbb{E}_g[\Pi g v]\| \geq c_0$ , and  $M = M(C_0, c_0, \Pi, \theta_*)$  is large enough, this is lower-bounded by a positive constant, so  $\|\nabla R(\theta)\| \geq c\sigma^{-2}$ .

Now suppose  $\|\mathbb{E}_g[\Pi g v]\| < c_0$ , where we will choose this  $(\Pi, \mathbf{G}, \theta_*)$ -dependent constant  $c_0 \in (0, 1)$  sufficiently small and to be determined. Constants  $C, C', c, c' > 0$  below are independent of  $c_0$ , and we will track explicitly the dependence of the argument on  $c_0$ . We apply the expansion (2.10) to the order  $K = 2$ . Then similarly,

$$\sigma^4 \|\nabla R(\theta)\| \geq \sigma^4 v^\top \nabla R(\theta) \geq v^\top \nabla \tilde{s}_2(\theta) + v^\top \nabla [\langle \tilde{T}_2(\theta), P_2(\theta) \rangle] + v^\top \nabla q_2(\theta) - C\|\theta\|^5/\sigma^2, \quad (\text{A.22})$$

where we have applied  $v^\top \nabla \tilde{s}_1(\theta) \geq 0$  from the form of  $\nabla \tilde{s}_1(\theta)$  above to drop the contribution from the  $k = 1$  term. We bound each expression on the right side of (A.22): First, applying the form of  $\tilde{s}_2(\theta)$  in Lemma A.1,

$$\begin{aligned} v^\top \nabla \tilde{s}_2(\theta) &= v^\top \mathbb{E}_{g,h} \left[ g^\top \Pi^\top \Pi h \theta \cdot \theta^\top g^\top \Pi^\top \Pi h \theta - g^\top \Pi^\top \Pi h \theta_* \cdot \theta^\top g^\top \Pi^\top \Pi h \theta_* \right] \\ &\geq \|\theta - \theta_*\|^3 \cdot \mathbb{E}_{g,h}[(v^\top g^\top \Pi^\top \Pi h v)^2] - C\|\theta\|^2, \end{aligned}$$

where the second line is obtained by writing each  $\theta$  as  $(\theta - \theta_*) + \theta_*$  and absorbing all but the term with cubic dependence on  $(\theta - \theta_*)$  into the  $C\|\theta\|^2$  remainder. As noted in Step 2 above,  $\Pi g v$  is not identically 0 over  $g \in \mathbf{G}$ , for any unit vector  $v$ . Then  $(v^\top g^\top \Pi^\top \Pi h v)^2$  is the squared-inner product between two i.i.d. non-zero vectors, and hence is strictly positive with positive probability. So  $\mathbb{E}_{g,h}[(v^\top g^\top \Pi^\top \Pi h v)^2] > 0$ . Then by compactness of the unit sphere,  $\mathbb{E}_{g,h}[(v^\top g^\top \Pi^\top \Pi h v)^2] > c > 0$  for every unit vector  $v$  and some constant  $c = c(\Pi, \mathbf{G}) > 0$ . So for  $\|\theta\| > M$  and large enough  $M = M(\Pi, \mathbf{G}, \theta_*)$ , this shows

$$v^\top \nabla \tilde{s}_2(\theta) \geq c'\|\theta\|^3. \quad (\text{A.23})$$

Next, consider  $v^\top \nabla q_2(\theta)$ . Since  $q_2 \in \tilde{\mathcal{R}}_{\leq 1}^{\mathbf{G}}$  which is generated by  $\tilde{T}_1(\theta) = \mathbb{E}_g[\Pi g \theta]$ ,  $q_2(\theta)$  is a quartic polynomial of the entries of  $\mathbb{E}_g[\Pi g \theta]$ , whose specific form depends only on  $\Pi, \mathbf{G}, \theta_*$ . Then, applying the chain rule to differentiate  $q_2(\theta)$ , we have

$$\|\nabla q_2(\theta)\| \leq C(\|\mathbb{E}_g[\Pi g \theta]\|^3 + 1).$$

Now applying  $\theta = (\theta - \theta_*) + \theta_* = \|\theta - \theta_*\|v + \theta_*$  and  $\|\mathbb{E}_g[\Pi g v]\| < c_0$ , we have

$$\|\mathbb{E}_g[\Pi g \theta]\| \leq c_0\|\theta - \theta_*\| + \|\Pi\|\|\theta_*\| \leq C'(c_0\|\theta\| + 1).$$

Then for a  $(\theta_*, \mathbf{G}, \Pi)$ -dependent constant  $C > 0$  independent of  $c_0$ , we get

$$|v^\top \nabla q_2(\theta)| \leq \|\nabla q_2(\theta)\| \leq C(c_0^3\|\theta\|^3 + 1). \quad (\text{A.24})$$

Similarly, consider  $v^\top \nabla [\langle \tilde{T}_2(\theta), P_2(\theta) \rangle]$ . Each entry of  $P_2(\theta)$  is a quadratic polynomial of  $\mathbb{E}_g[\Pi g \theta]$ . Noting that  $\tilde{T}_2(\theta)$  is also quadratic in  $\theta$ , by a similar argument as above,

$$\begin{aligned} |v^\top \nabla [\langle \tilde{T}_2(\theta), P_2(\theta) \rangle]| &\leq C' \left( (\|\mathbb{E}_g[\Pi g \theta]\| + 1) \cdot \|\theta\|^2 + (\|\mathbb{E}_g[\Pi g \theta]\| + 1)^2 \cdot \|\theta\| \right) \\ &\leq C'(c_0\|\theta\|^3 + \|\theta\|^2). \end{aligned} \quad (\text{A.25})$$

Finally, we may apply the condition  $\|\theta\| \leq C_0 \sigma^{2/3}$  to bound

$$\|\theta\|^5 / \sigma^2 \leq \|\theta\|^3 \cdot C_0^2 \sigma^{-2/3}. \quad (\text{A.26})$$

Applying (A.23), (A.24), (A.25), and (A.26) to (A.22), for sufficiently small  $c_0 = c_0(\Pi, G, \theta_*) \in (0, 1)$ , sufficiently large  $M = M(\Pi, G, \theta_*) > 0$ , any  $\theta$  satisfying  $C_0 \sigma^{2/3} \geq \|\theta\| > M$ , and all sufficiently large  $\sigma > \sigma_0$ , we obtain that  $\sigma^4 \|\nabla R(\theta)\|$  is lower-bounded by a  $(\Pi, G, \theta_*)$ -dependent constant, so  $\|\nabla R(\theta)\| \geq c \sigma^{-4}$  as desired.  $\square$

To complete the proofs of Theorems 2.11 and 2.13, it remains to analyze the optimization landscape of  $R(\theta)$  over the ball  $\{\theta : \|\theta\| \leq M\}$ . Our arguments are similar to those of [FSWW20, Sections 4.3 and 4.5]: Fixing any  $\tilde{\theta}$  in this ball, we may apply Lemma A.4 to reparametrize  $\theta$  by  $G$ -invariant polynomials in a sufficiently small neighborhood  $U_{\tilde{\theta}}$  of  $\tilde{\theta}$ , and then deduce statements about the landscape of  $R(\theta)$  within  $U_{\tilde{\theta}}$  by sequentially analyzing the landscapes of the terms  $s_1, s_2, s_3, \dots$  in this new system of coordinates. The conclusions about the landscape of  $R(\theta)$  over the full ball  $\{\theta : \|\theta\| \leq M\}$  then follow from patching together these analyses for local neighborhoods  $U_{\tilde{\theta}}$  that form a finite cover of this ball. Importantly, this argument requires both the radius  $M$  and the local neighborhoods  $U_{\tilde{\theta}}$  to be independent of  $\sigma$ .

To describe the full landscape of  $R(\theta)$  in this ball  $\{\theta : \|\theta\| \leq M\}$ , we must consider non-generic points  $\theta$  (even if  $\theta_*$  is generic). Thus it may be necessary to use non-generic points  $\tilde{\theta}$  in constructing this finite cover, where the system of coordinates given by Lemma A.4 locally around  $\tilde{\theta}$  may not contain a complete transcendence basis for  $\mathcal{R}^G$ . Instead, we separate the cases for  $\tilde{\theta}$  by the largest index  $k$  for which  $\tilde{\theta} \in \mathcal{V}_{k-1}(\theta_*)$  but  $\tilde{\theta} \notin \mathcal{V}_k(\theta_*)$ , and we provide a separate analysis for each  $k$ , using the given condition that  $dM_k(\theta)$  has constant rank on  $\mathcal{V}_k(\theta_*)$ . Our argument in the final case  $\tilde{\theta} \in \mathcal{V}_K(\theta_*)$  extends the analyses of [FSWW20] to handle orbits of positive dimension arising in the setting of a continuous group, and we then explain how these arguments may be adapted to use the expansion of (2.10) in models with projection.

*Proof of Theorem 2.11.* Consider the unprojected setting of part (a). Fixing a generic point  $\theta_* \in \mathbb{R}^d$ , Lemma A.4 shows that for any  $k = 1, \dots, K$ , the rank of  $dM_k(\theta_*)$  is  $d_1 + \dots + d_k$ . Then by the assumption that  $dM_k(\theta)$  has constant rank over  $\mathcal{V}_k(\theta_*)$ , since  $\theta_* \in \mathcal{V}_k(\theta_*)$ , this constant rank must be  $d_1 + \dots + d_k$ .

We now consider two cases for a (possibly non-generic) point  $\tilde{\theta} \in \mathbb{R}^d$ :

**Case 1:**  $\tilde{\theta} \in \mathcal{V}_{k-1}(\theta_*) \subseteq \dots \subseteq \mathcal{V}_0(\theta_*) = \mathbb{R}^d$ , but  $\tilde{\theta} \notin \mathcal{V}_k(\theta_*)$ , for some  $k \in \{1, \dots, K\}$ . The argument in this case is the same as that of [FSWW20, Theorem 4.27], and we reproduce it here for the reader's convenience. By the constant rank assumption,  $dM_{k-1}(\tilde{\theta})$  has rank  $d_1 + \dots + d_{k-1}$ . Let  $\varphi = (\varphi^1, \dots, \varphi^{k-1}, \bar{\varphi})$  be the map of Lemma A.4, with inverse  $\theta(\varphi)$  in a neighborhood  $U_{\tilde{\theta}}$  of  $\tilde{\theta}$ . (If  $k = 1$ , we take  $\varphi = \bar{\varphi} : \mathbb{R}^d \rightarrow \mathbb{R}^d$  to be an arbitrary invertible map, say the identity map.) We write  $f(\varphi)$  as shorthand for  $f(\theta(\varphi))$ .

In the parametrization by  $\varphi$ , each entry of  $T_1, \dots, T_{k-1}$  depends only on the coordinates  $\varphi^1, \dots, \varphi^{k-1}$ , by Lemma A.4(c). Thus

$$\mathcal{V}_{k-1}(\theta_*) \cap U_{\tilde{\theta}} = \left\{ \theta \in U_{\tilde{\theta}} : \varphi^1(\theta) = \varphi^1(\theta_*), \dots, \varphi^{k-1}(\theta) = \varphi^{k-1}(\theta_*) \right\},$$

and the remaining coordinates  $\bar{\varphi}$  form a local chart for the manifold  $\mathcal{V}_{k-1}(\theta_*)$  over  $U_{\tilde{\theta}}$ . This holds trivially also for  $k = 1$ .

Consider now the minimization of  $s_k$  over  $\mathcal{V}_{k-1}(\theta_*)$ . By the form of  $s_k$  in (2.8), its global minimizers over  $\mathcal{V}_{k-1}(\theta_*)$  are exactly the points of  $\mathcal{V}_k(\theta_*)$ . Since  $\tilde{\theta} \notin \mathcal{V}_k(\theta_*)$ , and the minimization of  $s_k$  over  $\mathcal{V}_{k-1}(\theta_*)$  is globally benign by assumption, this implies that

$$\text{either} \quad \nabla_{\bar{\varphi}} s_k(\varphi(\tilde{\theta})) \neq 0 \quad \text{or} \quad \lambda_{\min} \left( \nabla_{\bar{\varphi}}^2 s_k(\varphi(\tilde{\theta})) \right) < 0.$$

Applying continuity of  $s_k$  and its derivatives, and reducing the size of  $U_{\tilde{\theta}}$  as necessary, we may then ensure

$$\text{either} \quad \|\nabla_{\bar{\varphi}} s_k(\varphi)\| > c \quad \text{or} \quad \lambda_{\min} \left( \nabla_{\bar{\varphi}}^2 s_k(\varphi) \right) < -c \quad \text{for all } \varphi \in \varphi(U_{\tilde{\theta}}).$$

Here, the size of the neighborhood  $U_{\tilde{\theta}}$  and the constant  $c > 0$  are independent of  $\sigma$ , as  $s_k$  does not depend on  $\sigma$ . Now applying the expansion (2.7) to the order  $k$  and differentiating term-by-term in

$\varphi = (\varphi^1, \dots, \varphi^{k-1}, \bar{\varphi})$ , observe that  $\{s_j : j \leq k-1\}$  and  $\{q_j : j \leq k\}$  depend only on  $(\varphi^1, \dots, \varphi^{k-1})$  and not on  $\bar{\varphi}$ . Then for some constant  $\sigma_0 = \sigma_0(\tilde{\theta}) > 0$ , all  $\sigma > \sigma_0$ , and all  $\varphi \in \varphi(U_{\tilde{\theta}})$ ,

$$\text{either } \|\nabla_{\varphi} R(\varphi)\| > (c/2)\sigma^{-2k} \quad \text{or} \quad \lambda_{\min}(\nabla_{\varphi}^2 R(\varphi)) < -(c/2)\sigma^{-2k}. \quad (\text{A.27})$$

Finally, changing variables back to  $\theta$  by the chain rule, this implies

$$\text{either } \nabla_{\theta} R(\theta) \neq 0 \quad \text{or} \quad \lambda_{\min}(\nabla_{\theta}^2 R(\theta)) < 0 \quad \text{for all } \theta \in U_{\tilde{\theta}}. \quad (\text{A.28})$$

**Case 2:**  $\tilde{\theta} \in \mathcal{V}_K(\theta_*)$ . By the constant rank assumption,  $dM_K(\tilde{\theta}) = d_1 + \dots + d_K$ . Let  $\varphi = (\varphi^1, \dots, \varphi^K, \bar{\varphi})$  be the map of Lemma A.4 in a neighborhood  $U_{\tilde{\theta}}$  of  $\tilde{\theta}$ , with inverse  $\theta(\varphi)$ . We again write as shorthand  $f(\varphi) = f(\theta(\varphi))$ .

Let us write  $\nabla_{\varphi}^2 R(\varphi)$  in the  $(K+1) \times (K+1)$  block decomposition corresponding to  $(\varphi^1, \dots, \varphi^K, \bar{\varphi})$ . Applying the expansion (2.7) now to order  $K$ , each  $s_k(\varphi)$  depends only on  $\varphi^1, \dots, \varphi^k$ , each  $q_k(\varphi)$  depends only on  $\varphi^1, \dots, \varphi^{k-1}$ , and  $q(\varphi)$  and  $R(\varphi)$  depend only on  $\varphi^1, \dots, \varphi^K$ . Furthermore, Lemma A.7 shows  $\nabla_{\varphi^k}^2 s_k(\varphi(\tilde{\theta})) \succ 0$  strictly for each  $k = 1, \dots, K$ , so  $\nabla_{\varphi^k}^2 s_k(\varphi) \succ c \text{Id}$  for all  $\varphi \in \varphi(U_{\tilde{\theta}})$  by continuity, for a sufficiently small neighborhood  $U_{\tilde{\theta}}$  and constant  $c > 0$ . Then differentiating (2.7) term-by-term,  $\nabla_{\varphi}^2 R(\varphi)$  is zero outside the upper-left  $K \times K$  blocks, and these  $K \times K$  blocks have a graded block structure in the sense of Definition A.5 with  $d' = d - d_0$ , for any  $\sigma > \sigma_0 = \sigma_0(\tilde{\theta})$  and all points  $\varphi \in \varphi(U_{\tilde{\theta}})$ . Then, applying Lemma A.6(b), the upper-left  $K \times K$  blocks of  $\nabla_{\varphi}^2 R(\varphi)$  form a strictly positive-definite matrix. Recalling that  $R(\varphi)$  depends only on  $\varphi^1, \dots, \varphi^K$  and not on  $\bar{\varphi}$ , let us write  $\bar{R}(\varphi^1, \dots, \varphi^K) = R(\varphi)$ , and also reduce to a smaller neighborhood  $U_{\tilde{\theta}}$  such that  $\varphi(U_{\tilde{\theta}})$  has a product form  $V \times W$ , where  $V \subset \mathbb{R}^{d_1 + \dots + d_K}$  and  $W \subset \mathbb{R}^{d_0}$ . Then this shows that

$$\bar{R}(\varphi^1, \dots, \varphi^K) \text{ is strictly convex on } V. \quad (\text{A.29})$$

Now applying the assumption that  $\mathcal{V}_K(\theta_*) = \mathcal{O}_{\theta_*}$ , we have that  $\tilde{\theta} \in \mathcal{O}_{\theta_*}$  is a global minimizer of  $R(\theta)$ . Thus  $(\varphi^1(\tilde{\theta}), \dots, \varphi^K(\tilde{\theta}))$  is the global minimizer and unique critical point of  $\bar{R}$  on  $V$ . Changing coordinates back to  $\theta$  by the chain rule, the critical points of  $R(\theta)$  on  $U_{\tilde{\theta}}$  are then given exactly by

$$\left\{ \theta \in U_{\tilde{\theta}} : (\varphi^1(\theta), \dots, \varphi^K(\theta)) = (\varphi^1(\tilde{\theta}), \dots, \varphi^K(\tilde{\theta})) \right\}.$$

Applying (A.8), this shows

$$\left\{ \theta \in U_{\tilde{\theta}} : \nabla R(\theta) = 0 \right\} = U_{\tilde{\theta}} \cap \mathcal{O}_{\tilde{\theta}} = U_{\tilde{\theta}} \cap \mathcal{O}_{\theta_*} \quad (\text{A.30})$$

Finally, we combine these two cases using a compactness argument: By Lemma A.8, there are no critical points of  $R(\theta)$  outside a sufficiently large ball  $\overline{B}_M = \{\theta : \|\theta\| \leq M\}$ . For each point  $\tilde{\theta} \in \overline{B}_M$ , construct the neighborhood  $U_{\tilde{\theta}}$  as above, and take a finite set  $S$  of such points  $\tilde{\theta}$  for which  $\bigcup_{\tilde{\theta} \in S} U_{\tilde{\theta}}$  covers  $\overline{B}_M$ . Set  $\sigma_0 = \max_{\tilde{\theta} \in S} \sigma_0(\tilde{\theta})$ , where  $\sigma_0(\tilde{\theta})$  is as defined in the two cases above. Then for any  $\sigma > \sigma_0$ , the conditions (A.28) and (A.30) combine to show that any critical point of  $R(\theta)$  inside  $\overline{B}_M$  either belongs to the locus  $\mathcal{O}_{\theta_*}$  of global minimizers, or has  $\lambda_{\min}(\nabla^2 R(\theta)) < 0$ . Thus the minimization of  $R(\theta)$  is globally benign, concluding the proof of part (a).

The proof in the projected setting of part (b) is similar, with the following modifications: For the first case where  $\tilde{\theta} \in \tilde{\mathcal{V}}_{k-1}(\theta_*) \subseteq \dots \subseteq \tilde{\mathcal{V}}_0(\theta_*) = \mathbb{R}^d$  but  $\tilde{\theta} \notin \tilde{\mathcal{V}}_k(\theta_*)$ , Lemma A.4 still yields a local parametrization  $\varphi = (\varphi^1, \dots, \varphi^{k-1}, \bar{\varphi})$  where  $\bar{\varphi}$  forms a local chart for  $\tilde{\mathcal{V}}_{k-1}(\theta_*)$ . Differentiating (2.10) applied to the order  $k$  term-by-term in  $\varphi = (\varphi^1, \dots, \varphi^{k-1}, \bar{\varphi})$ , the gradient and Hessian of  $R(\varphi)$  in  $\bar{\varphi}$  have an additional contribution from  $\sigma^{-2k} \langle \tilde{T}_k(\varphi), P_k(\varphi) \rangle$ . Since  $P_k$  depends only on  $\varphi^1, \dots, \varphi^{k-1}$  and not on  $\bar{\varphi}$ , the gradient and Hessian in  $\bar{\varphi}$  are obtained by differentiating only  $\tilde{T}_k$ . Then both  $\nabla_{\bar{\varphi}}[\langle \tilde{T}_k(\varphi), P_k(\varphi) \rangle]|_{\varphi=\varphi(\tilde{\theta})} = 0$  and  $\nabla_{\bar{\varphi}}^2[\langle \tilde{T}_k(\varphi), P_k(\varphi) \rangle]|_{\varphi=\varphi(\tilde{\theta})} = 0$ , because  $P_k(\tilde{\theta}) = P_k(\theta_*) = 0$  for any  $\tilde{\theta} \in \tilde{\mathcal{V}}_{k-1}(\theta_*)$ . Then for a sufficiently small neighborhood  $U_{\tilde{\theta}}$ , we still obtain (A.27) for all  $\varphi \in \varphi(U_{\tilde{\theta}})$ , and hence (A.28) still holds.

For the second case where  $\tilde{\theta} \in \tilde{\mathcal{V}}_{\tilde{K}}(\theta_*)$ , similarly when computing the Hessian of (2.10) applied to the order  $\tilde{K}$  term-by-term in  $\varphi = (\varphi^1, \dots, \varphi^{\tilde{K}}, \bar{\varphi})$ , we have additional contributions from the terms  $\sigma^{-2k} \langle \tilde{T}_k(\varphi), P_k(\varphi) \rangle$ . In each  $(k, k)$  block, we again have  $\nabla_{\varphi^k}^2[\langle \tilde{T}_k(\varphi), P_k(\varphi) \rangle]|_{\varphi=\varphi(\tilde{\theta})} = 0$ , because only  $\tilde{T}_k$  depends on  $\varphi^k$  whereas  $P_k(\tilde{\theta}) = P_k(\theta_*) = 0$ . Then  $\nabla_{\varphi}^2 R(\varphi)$  still has a graded block structure in its upper-left  $\tilde{K} \times \tilde{K}$  blocks,

implying (A.29) for  $\varphi(U_{\tilde{\theta}}) = V \times W$  and a sufficiently small neighborhood  $U_{\tilde{\theta}}$  of  $\tilde{\theta}$ . Now by the assumption given in part (b) of the theorem, we have  $\Pi(\mathcal{O}_{\tilde{\theta}}) \equiv \Pi(\mathcal{O}_{\theta_*})$  for  $\tilde{\theta} \in \tilde{\mathcal{V}}_{\tilde{K}}(\theta_*)$ . Then  $\tilde{\theta}$  is a global minimizer of  $R(\theta)$ . The convexity of (A.29) then implies, by the same argument as in the unprojected setting, that  $\{\theta \in U_{\tilde{\theta}} : \nabla R(\theta) = 0\} = U_{\tilde{\theta}} \cap \mathcal{O}_{\tilde{\theta}}$ . Over the given domain  $\{\theta : \|\theta\| < B(\|\theta_*\| + \sigma)\}$ , Lemma A.8 ensures that there are no critical points of  $R(\theta)$  outside the smaller ball  $\{\theta : \|\theta\| \leq M\}$ , which is independent of  $\sigma$ . We conclude the proof by applying the same compactness argument over  $\{\theta : \|\theta\| \leq M\}$  as in the unprojected setting.  $\square$

Finally, we show Theorem 2.13 on a correspondence between local minimizers of  $R(\theta)$  and  $s_K(\theta)|_{\mathcal{V}_{K-1}(\theta_*)}$ . We remark that in the preceding proof of Theorem 2.11, *global* minimizers of  $R(\theta)$  must also minimize each function  $s_k(\theta)$  over  $\mathcal{V}_{k-1}(\theta_*)$ , and in particular, they are also exactly the minimizers of  $s_K(\theta)|_{\mathcal{V}_{K-1}(\theta_*)}$ . Such a statement is not true for local minimizers, and we will instead show that local minimizers of  $R(\theta)$  and  $s_K(\theta)|_{\mathcal{V}_{K-1}(\theta_*)}$  are close for large  $\sigma$ . We will use the following elementary lemma from [FSWW20], which ensures that minimizers of convex functions are close if the functions are pointwise close to each other.

**Lemma A.9** ([FSWW20] Lemma 2.8). *Let  $B_\varepsilon(\theta_0)$  be the ball of radius  $\varepsilon > 0$  around  $\theta_0 \in \mathbb{R}^d$ . Let  $f_1, f_2 : B_\varepsilon(\theta_0) \rightarrow \mathbb{R}$  be two functions which are twice continuously differentiable. Suppose that  $\theta_0$  is a critical point of  $f_1$ , and  $\lambda_{\min}(\nabla^2 f_1(\theta_0)) \geq c_0$  for a constant  $c_0 > 0$  and all  $\theta \in B_\varepsilon(\theta_0)$ . If*

$$|f_1(\theta) - f_2(\theta)| \leq \delta \quad \text{and} \quad \|\nabla^2 f_1(\theta) - \nabla^2 f_2(\theta)\| \leq \delta$$

*for some  $\delta < \min(c_0, c_0\varepsilon^2/4)$  and all  $\theta \in B_\varepsilon(\theta_0)$ , then  $f_2$  has a unique critical point in  $B_\varepsilon(\theta_0)$ , which is a local minimizer of  $f_2$ .*

*Proof of Theorem 2.13.* Consider the unprojected setting of part (a). Fix a generic point  $\theta_* \in \mathbb{R}^d$ . Lemma A.4 shows that  $\text{rank } dM_k(\theta_*) = d_1 + \dots + d_k$  for each  $k = 1, \dots, K$ . Then the given constant rank assumption ensures that  $\text{rank } dM_k(\theta) = d_1 + \dots + d_k$  for all  $\theta \in \mathcal{V}_k(\theta_*)$  and  $k = 1, \dots, K-1$ .

Statement (a2) is established by a small extension of the argument in Theorem 2.11, using the given condition that critical points of  $s_K(\theta)|_{\mathcal{V}_{K-1}(\theta_*)}$  are non-degenerate up to orbit: Lemma A.8 ensures that all critical points of  $R(\theta)$  belong to the ball  $\overline{B_M} = \{\theta : \|\theta\| \leq M\}$ . Fix any constant  $\varepsilon > 0$ , and let  $N_{\varepsilon, M}$  be the points in  $\overline{B_M}$  at distance  $\geq \varepsilon$  from all critical points of  $s_K(\theta)|_{\mathcal{V}_{K-1}(\theta_*)}$ . We consider two cases for a point  $\tilde{\theta} \in N_{\varepsilon, M}$ :

**Case 1:**  $\tilde{\theta} \in \mathcal{V}_{k-1}(\theta_*) \subseteq \dots \subseteq \mathcal{V}_0(\theta_*)$ , but  $\tilde{\theta} \notin \mathcal{V}_k(\theta_*)$ , for some  $k \in \{1, \dots, K-1\}$ . Then we have

$$\text{either } \nabla_{\theta} R(\theta) \neq 0 \quad \text{or} \quad \lambda_{\min}(\nabla_{\theta}^2 R(\theta)) < 0 \quad \text{for all } \theta \in U_{\tilde{\theta}}$$

by the same argument as leading to (A.28) in Theorem 2.11.

**Case 2:**  $\tilde{\theta} \in \mathcal{V}_{K-1}(\theta_*)$ . Then Lemma A.4 provides a local reparametrization  $\varphi = (\varphi^1, \dots, \varphi^{K-1}, \bar{\varphi})$  on a neighborhood  $U_{\tilde{\theta}}$  of  $\tilde{\theta}$ , where  $\varphi^j : \mathbb{R}^d \rightarrow \mathbb{R}^{d_j}$  and  $T_j$  depends only on  $(\varphi^1, \dots, \varphi^j)$  for each  $j = 1, \dots, K-1$ . Then  $\bar{\varphi}$  forms a local chart for  $\mathcal{V}_{K-1}(\theta_*)$  at  $\tilde{\theta}$ . Let  $\tilde{\varphi} = \varphi(\tilde{\theta})$ .

If  $\nabla_{\tilde{\varphi}} s_K(\tilde{\varphi}) = 0$ , then  $\tilde{\theta}$  is a critical point of  $s_K|_{\mathcal{V}_{K-1}(\theta_*)}$ , which by the given assumption must be non-degenerate up to orbit. Hence  $\mathcal{O}_{\tilde{\theta}}$  is locally a manifold of dimension  $d_0$  at  $\tilde{\theta}$ , so we may choose the parametrization  $\bar{\varphi}$  above to have a decomposition  $\bar{\varphi} = (\varphi^K, \varphi^0)$ , where  $\varphi^0$  has  $d_0$  coordinates forming a local chart for  $\mathcal{O}_{\tilde{\theta}}$ , and  $\varphi^K$  has  $d_K$  remaining coordinates. Since  $s_K$  is constant over  $\mathcal{O}_{\tilde{\theta}}$ , we must have  $\nabla_{\varphi^0} s_K = 0$ , so  $s_K$  depends only on  $\varphi^K$  and not on  $\varphi^0$  in this chart  $\bar{\varphi} = (\varphi^K, \varphi^0)$  for  $\mathcal{V}_{K-1}(\theta_*)$ . Then non-degeneracy of  $\tilde{\theta}$  up to orbit further implies that  $\nabla_{\varphi^K}^2 s_K(\tilde{\varphi})$  is a  $d_K \times d_K$  matrix of full rank  $d_K$ . If this were positive definite, then  $\tilde{\theta}$  would be a local minimizer of  $s_K$  on  $\mathcal{V}_{K-1}(\theta_*)$ , but we have assumed  $\tilde{\theta} \in N_{\varepsilon, M}$  which does not include such local minimizers. Therefore  $\nabla_{\varphi^K}^2 s_K(\tilde{\varphi})$  must have a negative eigenvalue. This shows that

$$\text{either } \nabla_{\tilde{\varphi}} s_K(\tilde{\varphi}) \neq 0 \quad \text{or} \quad \lambda_{\min}(\nabla_{\tilde{\varphi}}^2 s_K(\tilde{\varphi})) < 0.$$

Thus, differentiating (2.7) applied to the order  $K-1$  term-by-term in  $\varphi = (\varphi^1, \dots, \varphi^{K-1}, \bar{\varphi})$ , also in this case

$$\text{either } \nabla_{\theta} R(\theta) \neq 0 \quad \text{or} \quad \lambda_{\min}(\nabla_{\theta}^2 R(\theta)) < 0 \quad \text{for all } \theta \in U_{\tilde{\theta}}.$$

Combining these two cases, taking a finite cover of  $N_{\varepsilon, M}$  by such neighborhoods  $U_{\tilde{\theta}}$ , this shows that for any  $\varepsilon > 0$  and all  $\sigma > \sigma_0(\varepsilon)$ , all local minimizers of  $R(\theta)$  must be  $\varepsilon$ -close to some local minimizer of  $s_K(\theta)$  on  $\mathcal{V}_{K-1}(\theta_*)$ . Then there exists a slowly decreasing sequence  $\varepsilon(\sigma) \rightarrow 0$  as  $\sigma \rightarrow \infty$ , for which each local minimizer of  $R(\theta)$  is  $\varepsilon(\sigma)$ -close to a local minimizer of  $s_K(\theta)$ . This establishes (a2). The proof of (b2) in the projected setting is the same, where the gradients and Hessians in  $\bar{\varphi}$  of the additional terms  $\sigma^{-2k} \langle \tilde{T}_k(\varphi), P_k(\varphi) \rangle$  from (2.10) are handled in the same way as in the proof of Theorem 2.11.

We now show the converse direction (a1). Let  $\theta_+ \in \mathcal{V}_{K-1}(\theta_*)$  be a local minimizer of  $s_K|_{\mathcal{V}_{K-1}(\theta_*)}$  that is non-degenerate up to orbit. By Lemma A.4 and the same argument as above, there is a local reparametrization  $\varphi = (\varphi^1, \dots, \varphi^{K-1}, \varphi^K, \varphi^0)$  on a neighborhood  $U_{\theta_+}$  of  $\theta_+$  such that  $T_k(\varphi)$  depends only on  $\varphi^1, \dots, \varphi^k$  for each  $k = 1, \dots, K-1$ , and  $s_K(\varphi)$  and  $R(\varphi)$  depend only on  $\varphi^1, \dots, \varphi^K$ . Let  $\varphi_+ = \varphi(\theta_+)$ . For  $k = 1, \dots, K-1$ , some constant  $c > 0$ , and all  $\varphi \in \varphi(U_{\theta_+})$ ,

$$\lambda_{\min}(\nabla_{\varphi^k}^2 s_k(\varphi)) > c \quad (\text{A.31})$$

by Lemma A.7 applied with  $\tilde{\theta} = \theta_+$  and by continuity of this Hessian. This holds also for  $k = K$ , by the non-degeneracy of  $\theta_+$  up to orbit. Then, writing the Hessian  $\nabla_{\varphi}^2 R(\varphi)$  in the  $(K+1) \times (K+1)$  block structure corresponding to  $(\varphi^1, \dots, \varphi^K, \varphi^0)$ , we obtain as in Theorem 2.11 that the upper-left  $K \times K$  blocks have a graded block structure, in a sufficiently small neighborhood  $U_{\theta_+}$  where  $\varphi(U_{\theta_+}) = V \times W$  has a product form. So, defining  $\bar{R}(\varphi^1, \dots, \varphi^K) = R(\varphi)$ ,  $\bar{R}$  is strictly convex over  $V$ .

However, in contrast to Theorem 2.11,  $\theta_+$  is not necessarily a global (or local) minimizer of  $R(\theta)$ , so the existence of a local minimizer of  $\bar{R}(\varphi^1, \dots, \varphi^K)$  in  $V$  is less immediate. By further reducing  $U_{\theta_+}$ , we may assume  $V$  takes a product form  $V = V_1 \times \dots \times V_K$  where  $V_k$  corresponds to the coordinates of  $\varphi^k$ . Let  $\bar{V}, \bar{V}_k$  be the closures of  $V, V_k$ , which are compact. Let  $\hat{\varphi} = (\hat{\varphi}^1, \dots, \hat{\varphi}^K)$  be a point which minimizes  $\bar{R}$  over  $\bar{V}$ . We aim to show that  $\hat{\varphi}$  in fact belongs to the interior of  $\bar{V}$ , and hence is a critical point and local minimizer of  $\bar{R}$  in  $V$ . To show this, we will inductively show that each subvector  $\hat{\varphi}^k$  belongs to the interior of  $\bar{V}_k$ , by using Lemma A.9 to argue that it is close to  $\varphi_+^k$  where  $(\varphi_+^1, \dots, \varphi_+^k)$  minimizes  $s_k(\varphi)$ . The argument is similar to that of [FSW20, Lemma 4.15] (fixing a minor error therein), and we reproduce this argument here.

Let  $C, C', c > 0$  denote  $(\theta_*, \mathbb{G})$ -dependent constants changing from instance to instance. Let us write  $s_k(\varphi^1, \dots, \varphi^k) = s_k(\varphi)$ , as this does not depend on the remaining coordinates of  $\varphi$ . Because  $\theta_+ \in \mathcal{V}_1(\theta_*)$ ,  $\theta_+$  is a global minimizer of  $s_1$  over  $\bar{V}_1$ . Then, applying (A.31) for  $k = 1$ , we get for some constant  $c > 0$  and sufficiently small neighborhood  $V_1$  that

$$s_1(\hat{\varphi}^1) - s_1(\varphi_+^1) \geq c \|\hat{\varphi}^1 - \varphi_+^1\|^2.$$

Applying the series expansion (2.7) to order  $K = 1$ , and noting that  $q_1 \in \mathcal{R}_{\leq 0}^{\mathbb{G}}$  must be a constant, this implies that

$$\bar{R}(\hat{\varphi}^1, \dots, \hat{\varphi}^K) - \bar{R}(\varphi_+^1, \dots, \varphi_+^K) \geq c\sigma^{-2} \|\hat{\varphi}^1 - \varphi_+^1\|^2 - C\sigma^{-4}, \quad (\text{A.32})$$

for all  $\sigma > \sigma_0$  and large enough  $\sigma_0$ . Since  $\hat{\varphi}$  minimizes  $\bar{R}(\varphi)$  over  $\bar{V}$ , the left side is non-positive, so we obtain  $c\sigma^{-2} \|\hat{\varphi}^1 - \varphi_+^1\|^2 - C\sigma^{-4} \leq 0$ . This shows

$$\|\hat{\varphi}^1 - \varphi_+^1\| \leq C'\sigma^{-1} < \sigma^{-\eta_1} \quad (\text{A.33})$$

for large enough  $\sigma$  and, say,  $\eta_1 = 0.1$ .

Suppose inductively that we have shown

$$\|\hat{\varphi}^1 - \varphi_+^1\|, \dots, \|\hat{\varphi}^{k-1} - \varphi_+^{k-1}\| < \sigma^{-\eta_{k-1}} \quad (\text{A.34})$$

for some constant  $\eta_{k-1} > 0$ . Consider the functions  $h_+(\varphi^k) = s_k(\varphi_+^1, \dots, \varphi_+^{k-1}, \varphi^k)$  and  $\hat{h}(\varphi^k) = s_k(\hat{\varphi}^1, \dots, \hat{\varphi}^{k-1}, \varphi^k)$  on  $\bar{V}_k$ . Since  $s_k$  is a polynomial function of its arguments, both  $s_k$  and its Hessian are Lipschitz over the bounded domain  $\bar{V}_1 \times \dots \times \bar{V}_k$ . Then applying (A.34), for a constant  $C > 0$  depending on  $\bar{V}_1 \times \dots \times \bar{V}_k$  and  $s_k$  but not on  $\sigma$ , we have

$$\sup_{\varphi^k \in \bar{V}_k} |h_+(\varphi^k) - \hat{h}(\varphi^k)| < C\sigma^{-\eta_{k-1}}, \quad \sup_{\varphi^k \in \bar{V}_k} \|\nabla^2 h_+(\varphi^k) - \nabla^2 \hat{h}(\varphi^k)\| < C\sigma^{-\eta_{k-1}}. \quad (\text{A.35})$$

Both  $h_+$  and  $\hat{h}$  are strongly convex over  $\bar{V}_k$ , with Hessians lower bounded as (A.31). For  $k \leq K-1$ , since  $\theta_+ \in \mathcal{V}_k(\theta_*)$ , we know that  $\varphi_+^k$  minimizes  $h_+$  on  $\bar{V}_k$ . For  $k = K$ , since  $\theta_+$  is a local minimizer of  $s_K|_{\mathcal{V}_{K-1}(\theta_*)}$  by assumption, we know also that  $\varphi_+^K$  minimizes  $h_+$  on a sufficiently small neighborhood  $\bar{V}_K$ . Then (A.35)

and Lemma A.9 applied with  $\varepsilon = C'\sqrt{\sigma^{-\eta_{k-1}}}$  for a large enough constant  $C' > 0$  guarantee that the global minimizer  $\bar{\varphi}^k$  of  $\hat{h}$  over  $\bar{V}_k$  satisfies

$$\|\bar{\varphi}^k - \varphi_+^k\| < \varepsilon = C'\sqrt{\sigma^{-\eta_{k-1}}}. \quad (\text{A.36})$$

In particular,  $\bar{\varphi}^k$  must be in the interior of  $\bar{V}_k$  and is a critical point of  $\hat{h}$ , for sufficiently large  $\sigma > \sigma_0$ . So (A.31) implies for a constant  $c > 0$  that

$$s_k(\hat{\varphi}^1, \dots, \hat{\varphi}^{k-1}, \hat{\varphi}^k) - s_k(\hat{\varphi}^1, \dots, \hat{\varphi}^{k-1}, \bar{\varphi}^k) \geq c\|\hat{\varphi}^k - \bar{\varphi}^k\|^2.$$

Applying the series expansion (2.7) to the order  $k$ , and recalling that  $s_1, \dots, s_{k-1}$  and  $q_1, \dots, q_k$  depend only on  $\varphi^1, \dots, \varphi^{k-1}$ , we get

$$\bar{R}(\hat{\varphi}^1, \dots, \hat{\varphi}^K) - \bar{R}(\hat{\varphi}^1, \dots, \hat{\varphi}^{k-1}, \bar{\varphi}^k, \varphi_+^{k+1}, \dots, \varphi_+^K) \geq c\sigma^{-2k}\|\hat{\varphi}^k - \bar{\varphi}^k\|^2 - C\sigma^{-2(k+1)}. \quad (\text{A.37})$$

This is again non-positive because  $\hat{\varphi}$  minimizes  $\bar{R}$  over  $\bar{V}$ . So  $\|\hat{\varphi}^k - \bar{\varphi}^k\| \leq C'\sigma^{-1}$ . Combining with (A.36) and (A.34), we obtain

$$\|\hat{\varphi}^1 - \varphi_+^1\|, \dots, \|\hat{\varphi}^k - \varphi_+^k\| < \sigma^{-\eta_k} \quad (\text{A.38})$$

for large enough  $\sigma$  and, say,  $\eta_k = \eta_{k-1}/3$ . This completes the induction, showing that (A.38) holds up to  $k = K$ , and hence that  $\hat{\varphi} = (\hat{\varphi}^1, \dots, \hat{\varphi}^K)$  is in the interior of  $\bar{V}$  for  $\sigma > \sigma_0$  sufficiently large. Hence  $\hat{\varphi}$  is a critical point and local minimizer of  $\bar{R}$ . Then the point  $\theta(\hat{\varphi}^1, \dots, \hat{\varphi}^K, \varphi_+^0)$  is a local minimizer of  $R(\theta)$  on  $\mathbb{R}^d$ , which is  $\varepsilon(\sigma)$ -close to  $\theta_+$  for  $\varepsilon(\sigma) = C\sigma^{-\eta_K}$ . This shows (a1).

The proof of (b1) in the projected setting is similar, applying (2.10) in place of (2.7). In the first step for  $k = 1$  we observe, in addition to  $q_1 \in \mathcal{R}_{\leq 0}^G$  being constant, that  $P_1 = 0$  because it is also constant and equals 0 at  $\theta_*$ . Thus we obtain (A.32) and (A.33) without modification. In the inductive step, in place of (A.37), we have

$$\begin{aligned} & \bar{R}(\hat{\varphi}^1, \dots, \hat{\varphi}^K) - \bar{R}(\hat{\varphi}^1, \dots, \hat{\varphi}^{k-1}, \bar{\varphi}^k, \varphi_+^{k+1}, \dots, \varphi_+^K) \\ & \geq \sigma^{-2k} \left( c\|\hat{\varphi}^k - \bar{\varphi}^k\|^2 + \left\langle \tilde{T}_k(\hat{\varphi}^1, \dots, \hat{\varphi}^k) - \tilde{T}_k(\hat{\varphi}^1, \dots, \hat{\varphi}^{k-1}, \bar{\varphi}^k), P_k(\hat{\varphi}^1, \dots, \hat{\varphi}^{k-1}) \right\rangle \right) - C\sigma^{-2(k+1)}. \end{aligned}$$

We may bound

$$\|\tilde{T}_k(\hat{\varphi}^1, \dots, \hat{\varphi}^k) - \tilde{T}_k(\hat{\varphi}^1, \dots, \hat{\varphi}^{k-1}, \bar{\varphi}^k)\|_{\text{HS}} \leq C\|\hat{\varphi}^k - \bar{\varphi}^k\|$$

and

$$\begin{aligned} \|P_k(\hat{\varphi}^1, \dots, \hat{\varphi}^{k-1})\|_{\text{HS}} &= \|P_k(\hat{\varphi}^1, \dots, \hat{\varphi}^{k-1}) - P_k(\varphi_+^1, \dots, \varphi_+^{k-1})\|_{\text{HS}} \\ &\leq C(\|\hat{\varphi}^1 - \varphi_+^1\| + \dots + \|\hat{\varphi}^{k-1} - \varphi_+^{k-1}\|) \leq C'\sigma^{-\eta_{k-1}}, \end{aligned}$$

where the first equality holds because  $\theta_+ \in \mathcal{V}_{k-1}(\theta_*)$  so  $P_k(\varphi_+^1, \dots, \varphi_+^{k-1}) = P_k(\varphi_*^1, \dots, \varphi_*^{k-1}) = 0$ . This yields

$$\begin{aligned} 0 &\geq \bar{R}(\hat{\varphi}^1, \dots, \hat{\varphi}^K) - \bar{R}(\hat{\varphi}^1, \dots, \hat{\varphi}^{k-1}, \bar{\varphi}^k, \varphi_+^{k+1}, \dots, \varphi_+^K) \\ &\geq \sigma^{-2k} \left( c\|\hat{\varphi}^k - \bar{\varphi}^k\|^2 - C\sigma^{-\eta_{k-1}}\|\hat{\varphi}^k - \bar{\varphi}^k\| \right) - C\sigma^{-2(k+1)}. \end{aligned}$$

Viewing the right side as a quadratic function in  $\|\hat{\varphi}^k - \bar{\varphi}^k\|$ , this implies that  $\|\hat{\varphi}^k - \bar{\varphi}^k\|$  is at most the larger of the two roots of this quadratic function, which still gives (A.38). This completes the induction, and the proof is concluded as in the unprojected setting of (a1).  $\square$

## APPENDIX B. ANALYSIS OF ORTHOGONAL PROCRUSTES ALIGNMENT

We provide the details for Example 2.14 on the Procrustes alignment model, either with or without reflections. We consider a generic point  $\theta_* \in \mathbb{R}^{3 \times m}$  satisfying  $\text{rank}(\theta_*) = 3$  (where this is a generic condition because  $\text{rank}(\theta_*) < 3$  is equivalent to  $\det \theta_* \theta_*^\top = 0$ ). Recall that the group is either  $G = O(3) \otimes \text{Id}_m$  or  $G = SO(3) \otimes \text{Id}_m$ , acting on  $\mathbb{R}^{3 \times m} \cong \mathbb{R}^d$ . We represent an element of this group as  $g \otimes \text{Id}_m$ , where  $g \in O(3)$  or  $g \in SO(3)$  is a  $3 \times 3$  matrix. For the Haar-uniform law on both  $O(3)$  and  $SO(3)$ , we have the moment identities

$$\mathbb{E}_g[g_{ij}] = 0, \quad \mathbb{E}_g[g_{ij}g_{i'j'}] = \frac{1}{3} \cdot \mathbf{1}\{(i, j) = (i', j')\}. \quad (\text{B.1})$$

The first identity and the second for  $(i, j) \neq (i', j')$  follow from the fact that  $g$  is invariant in law under negation of any two rows or two columns. The second identity for  $(i, j) = (i', j')$  follows from  $\text{Tr } g^\top g =$

$\sum_{i,j=1}^3 g_{ij}^2 = 3$ , and the equality in law of the entries  $g_{ij}$ . The action of this element  $g \otimes \text{Id}_m$  on  $\theta \in \mathbb{R}^{3 \times m}$  is given by the matrix product

$$\theta \mapsto g\theta,$$

and the Euclidean inner-product may be written as  $\langle \theta, \theta' \rangle = \text{Tr } \theta^\top \theta' = \text{Tr } \theta' \theta^\top$ .

Let us first compute  $d_0 = \max_\theta \dim(\mathcal{O}_\theta)$ . Note that  $\dim(\mathcal{O}_\theta) = \dim(\mathbf{G}) - \dim(\mathbf{G}_\theta)$ , where  $\mathbf{G}_\theta = \{g \in \mathbf{G} : g\theta = \theta\}$  is the stabilizer subgroup of  $\theta$  [Aud04, Section I.1.b]. For both  $\mathbf{G} = \text{O}(3) \otimes \text{Id}_m$  and  $\mathbf{G} = \text{SO}(3) \otimes \text{Id}_m$ , we have  $\dim(\mathbf{G}) = 3$ . For any  $\theta \in \mathbb{R}^{3 \times m}$  having full rank 3, there is a right inverse  $\theta^\dagger \in \mathbb{R}^{m \times 3}$  for which  $\theta\theta^\dagger = \text{Id}$ . Thus  $g\theta = \theta$  requires  $g = \text{Id}$ , so that  $\dim(\mathbf{G}_\theta) = 0$ . Thus, we obtain

$$d_0 = 3, \quad \text{trdeg } \mathcal{R}^{\mathbf{G}} = d - d_0 = d - 3.$$

We now verify the values of  $d_1, d_2$  and the forms of  $T_1(\theta)$ ,  $T_2(\theta)$ ,  $\mathcal{V}_1(\theta_*)$ ,  $\mathcal{V}_2(\theta_*)$ ,  $s_1(\theta)$ , and  $s_2(\theta)$  as stated in Example 2.14. For  $T_1(\theta)$ ,  $\mathcal{V}_1(\theta_*)$ , and  $s_1(\theta)$ , by the first identity of (B.1),  $T_1(\theta) = \mathbb{E}_g[g\theta] = 0$ . So by the definitions of  $s_1(\theta)$  and  $\mathcal{V}_1(\theta_*)$  and Lemma 2.8,

$$s_1(\theta) = 0, \quad d_1 = 0, \quad \mathcal{V}_1(\theta_*) = \mathbb{R}^d.$$

For  $T_2(\theta)$ ,  $\mathcal{V}_2(\theta_*)$ , and  $s_2(\theta)$ , let us write  $[\theta]_i, [g\theta]_i \in \mathbb{R}^m$  for the  $i^{\text{th}}$  rows of  $\theta, g\theta \in \mathbb{R}^{3 \times m}$ , and  $[T_2(\theta)]_{ii'} \in \mathbb{R}^{m \times m}$  for the  $(i, i')$  block of  $T_2(\theta)$  in the  $3 \times 3$  block decomposition of  $\mathbb{R}^{d \times d} \cong \mathbb{R}^{3m \times 3m}$ , where  $i, i' \in \{1, 2, 3\}$ . Then

$$[T_2(\theta)]_{ii'} = \mathbb{E}_g [[g\theta]_i \cdot [g\theta]_{i'}^\top].$$

When  $i \neq i'$ , the second identity of (B.1) yields  $[T_2(\theta)]_{ii'} = 0$ . When  $i = i'$ , it yields

$$[T_2(\theta)]_{ii} = \sum_{j,j'=1}^3 \mathbb{E}_g [g_{ij}[\theta]_j \cdot g_{ij'}[\theta]_{j'}^\top] = \frac{1}{3} ([\theta]_1[\theta]_1^\top + [\theta]_2[\theta]_2^\top + [\theta]_3[\theta]_3^\top) = \frac{1}{3} \theta^\top \theta.$$

Thus we get  $T_2(\theta) = \frac{1}{3} \text{Id}_{3 \times 3} \otimes (\theta^\top \theta)$ . Then also

$$\begin{aligned} s_2(\theta) &= \frac{1}{4} \|T_2(\theta) - T_2(\theta_*)\|_{\text{HS}}^2 = \frac{1}{4} \cdot \frac{1}{9} \|\text{Id}_{3 \times 3} \otimes (\theta^\top \theta) - \text{Id}_{3 \times 3} \otimes (\theta_*^\top \theta_*)\|_{\text{HS}}^2 \\ &= \frac{1}{12} \|\theta^\top \theta - \theta_*^\top \theta_*\|_{\text{HS}}^2. \end{aligned} \tag{B.2}$$

This shows that  $\mathcal{V}_2(\theta_*) = \{\theta : s_2(\theta) = 0\} = \{\theta : \theta^\top \theta = \theta_*^\top \theta_*\}$ . For such  $\theta$ , the row span of  $\theta$  coincides with that of  $\theta_*$ . Since  $\text{rank}(\theta_*) = 3$ , this implies  $\theta = A\theta_*$  for some invertible matrix  $A \in \mathbb{R}^{3 \times 3}$ . Then  $\theta^\top \theta = \theta_*^\top \theta_*$  requires  $0 = \theta_*^\top (A^\top A - \text{Id})\theta_*$ , so  $A^\top A - \text{Id} = 0$ . Then  $A$  is orthogonal, and we obtain that  $\theta \in \{g\theta_* : g \in \text{O}(3)\}$ . This verifies

$$\mathcal{V}_2(\theta_*) = \{g\theta_* : g \in \text{O}(3)\},$$

which is exactly  $\mathcal{O}_{\theta_*}$  if  $\mathbf{G} = \text{O}(3) \otimes \text{Id}_m$ , and  $\mathcal{O}_{\theta_*} \cup \mathcal{O}_{-\theta_*}$  if  $\mathbf{G} = \text{SO}(3) \otimes \text{Id}_m$ .

To compute  $d_2$ , we apply Lemma 2.8. Differentiating the expression (B.2) twice at  $\theta = \theta_*$  by the chain rule,

$$\nabla^2 s_2(\theta_*) = \frac{1}{6} d_\theta [\theta^\top \theta]^\top \cdot d_\theta [\theta^\top \theta] \Big|_{\theta=\theta_*}$$

where  $d_\theta [\theta^\top \theta]$  denotes the Jacobian of the vectorization of  $\theta^\top \theta$  as a function of  $\theta$ . For generic  $\theta_* \in \mathbb{R}^{3 \times m}$ , specializing Lemma D.6 to follow with  $l = 1$  and  $S = m \geq 3$ , we then get  $\text{rank}[\nabla^2 s_2(\theta_*)] = \text{rank}(d_\theta [\theta^\top \theta]|_{\theta=\theta_*}) = 3(m-1) = d-3$ . Thus, recalling  $d_1 = 0$ , Lemma 2.8 shows

$$d_2 = \text{trdeg } \mathcal{R}_{\leq 2}^{\mathbf{G}} = d - 3.$$

This coincides with  $\text{trdeg } \mathcal{R}^{\mathbf{G}}$ , so also  $K = 2$  is the smallest integer satisfying Proposition 2.6.

Finally, we analyze the optimization landscape of  $s_2(\theta)$  over  $\mathcal{V}_1(\theta_*) = \mathbb{R}^d$ , and show that its only critical points are strict saddles or the global minimizers  $\mathcal{V}_2(\theta_*)$ . Computing the gradient of (B.2), we have

$$\nabla s_2(\theta) = \frac{1}{3} \theta \left( \theta^\top \theta - \theta_*^\top \theta_* \right) \in \mathbb{R}^{3 \times m}. \tag{B.3}$$

The row span of  $\theta\theta^\top \theta$  is the same as that of  $\theta$  (regardless of the rank of  $\theta$ ), while the row span of  $\theta\theta_*^\top \theta_*$  is contained in the row span of  $\theta_*$ . Thus, at any critical point  $\theta$  satisfying  $\nabla s_2(\theta) = 0$ , the row span of  $\theta$  is contained in that of  $\theta_*$ , i.e. we have

$$\theta = A\theta_* \tag{B.4}$$

for some (possibly singular) matrix  $A \in \mathbb{R}^{3 \times 3}$ . Applying this form to (B.3), we see that  $\nabla s_2(\theta) = 0$  implies

$$0 = A\theta_*\theta_*^\top A^\top A\theta_* - A\theta_*\theta_*^\top \theta_* = A\theta_*\theta_*^\top (A^\top A - \text{Id})\theta_*.$$

When  $\text{rank}(\theta_*) = 3$ , the rows of  $\theta_*$  are linearly independent, so this implies

$$0 = A\theta_*\theta_*^\top (A^\top A - \text{Id}). \quad (\text{B.5})$$

We now consider two cases for a critical point  $\theta$  of  $s_2(\theta)$ :

**Case 1:**  $\theta$  has full rank 3. Then by (B.4),  $A$  must be nonsingular. Multiplying (B.5) by  $(\theta_*\theta_*^\top)^{-1}A^{-1}$ , we get  $0 = A^\top A - \text{Id}$ , so  $A^\top A = \text{Id}$ . Thus,  $A$  is an orthogonal matrix, so  $\theta \in \mathcal{V}_2(\theta_*)$ , and this is a global minimizer of  $s_2(\theta)$ .

**Case 2:**  $\theta$  has some rank  $k \leq 2$ . Then by (B.4), also  $\text{rank}(A) = \text{rank}(A^\top A) = k$ . Then  $P = \text{Id} - A^\top A$  has  $3-k$  eigenvalues equal to 1, and in particular,  $\text{rank } P \geq 3-k$ . On the other hand, since  $\text{rank}(A\theta_*\theta_*^\top) = k$  and  $0 = A\theta_*\theta_*^\top P$  by (B.5), the kernel of  $P$  has dimension at least  $k$ . Then also  $\text{rank } P \leq 3-k$ , so  $\text{rank } P = 3-k$  exactly. Then  $P$  has  $3-k$  eigenvalues 1 and  $k$  eigenvalues 0, so it is an orthogonal projection onto a subspace of dimension  $3-k$  in  $\mathbb{R}^3$ .

Using this observation, we now exhibit a direction of negative curvature in  $\nabla^2 s_2(\theta)$ : Let  $\Delta = uv^\top \in \mathbb{R}^{3 \times m}$  be any rank-one non-zero matrix where  $A^\top u = 0$  and  $P\theta_*v \neq 0$ . Such vectors  $u$  and  $v$  exist because  $\text{rank } A < 3$ ,  $\text{rank } P > 0$ , and  $\theta_*$  has full rank 3. Consider

$$\theta_t = \theta + t\Delta = A\theta_* + t\Delta,$$

and the Hessian  $\nabla^2 s_2(\theta)$  in the direction of  $\Delta$ , given by  $\partial_t^2 s_2(\theta_t)|_{t=0}$ . Applying  $A^\top \Delta = A^\top uv^\top = 0$  by the condition  $A^\top u = 0$ , observe that  $\theta_t^\top \theta_t = \theta_*^\top A^\top A\theta_* + t^2 \Delta^\top \Delta$ , so from (B.2),

$$s_2(\theta_t) = \frac{1}{12} \left\| \theta_t^\top \theta_t - \theta_*^\top \theta_* \right\|_{\text{HS}}^2 = \frac{1}{12} \left\| t^2 \Delta^\top \Delta - \theta_*^\top P\theta_* \right\|_{\text{HS}}^2.$$

This is a polynomial in  $t$ , whose quadratic term is

$$[t^2]s_2(\theta_t) = -\frac{1}{6} \text{Tr } \theta_*^\top P\theta_* \Delta^\top \Delta.$$

So

$$\partial_t^2 s_2(\theta_t) \Big|_{t=0} = -\frac{1}{3} \text{Tr } \theta_*^\top P\theta_* \Delta^\top \Delta = -\frac{1}{3} \|P\theta_* \Delta^\top\|_{\text{HS}}^2.$$

Finally, note that  $P\theta_* \Delta^\top = (P\theta_*v)u^\top \neq 0$  because  $P\theta_*v \neq 0$  and  $u \neq 0$ . Then this is strictly negative, so  $\lambda_{\min}(\nabla^2 s_2(\theta)) < 0$ .

Combining these two cases, this verifies the claim in Example 2.14 that for generic  $\theta_*$ , the minimization of  $s_2(\theta)$  over  $\mathcal{V}_1(\theta_*) = \mathbb{R}^d$  is globally benign. For the claims about the landscape of  $R(\theta)$ , observe that for any  $\theta = g\theta_*$  where  $g \in \text{O}(3)$ , we have  $s_2(\theta) = 0$ , so  $T_2(\theta) = T_2(\theta_*)$ . Then applying the chain rule to differentiate twice  $s_2(\theta) = \|T_2(\theta) - T_2(\theta_*)\|_{\text{HS}}^2/4$ , and applying also  $s_1(\theta) = 0$ , we obtain  $\text{rank } \nabla^2 s_2(\theta) = \text{rank } dT_2(\theta) = \text{rank } dM_2(\theta)$ . On the other hand, for any such  $\theta$ , the preceding computation shows also  $\text{rank } \nabla^2 s_2(\theta) = \text{rank } d_\theta[\theta^\top] = d-3$ , as this rank is the same at  $\theta = g\theta_*$  as at  $\theta_*$ . Therefore  $\text{rank } dM_2(\theta) = d-3$ . Thus  $dM_2(\theta)$  has constant rank on  $\mathcal{V}_2(\theta_*)$ , which is a manifold of dimension  $d_0 = 3$ , so the minimizers  $\mathcal{V}_2(\theta_*) = \{g\theta_* : g \in \text{O}(3)\}$  of  $s_2(\theta_*)$  are non-degenerate up to orbit. Then the claims about the landscape of  $R(\theta)$  for large  $\sigma$  follow from Theorems 2.11 and 2.13.

## APPENDIX C. ANALYSIS OF CONTINUOUS MULTI-REFERENCE ALIGNMENT

**C.1. Unprojected continuous MRA.** In this section we prove Theorems 3.1 and 3.3.

*Proof of Theorem 3.3.* Recall by Lemma A.1 that

$$s_k(\theta) = \frac{1}{2(k!)} \mathbb{E}_g \left[ \langle \theta, g \cdot \theta \rangle^k - 2 \langle \theta, g \cdot \theta_* \rangle^k + \langle \theta_*, g \cdot \theta_* \rangle^k \right]. \quad (\text{C.1})$$

Let  $\theta, \vartheta \in \mathbb{R}^d$ . Computing the form of  $\mathbb{E}_g[\langle \theta, g \cdot \vartheta \rangle^k]$  will give the form of  $s_k(\theta)$ .

**Case  $k = 1$ :** By (3.4),  $\mathbb{E}_g[g] = \text{diag}(1, 0, \dots, 0)$ . It follows from (C.1) that

$$s_1(\theta) = \frac{1}{2} \left( (\theta^{(0)})^2 - 2\theta^{(0)}\theta_*^{(0)} + (\theta_*^{(0)})^2 \right) = \frac{1}{2} \left( \theta^{(0)} - \theta_*^{(0)} \right)^2.$$

**Case  $k = 2$ :** Recall  $u^{(0)}(\theta) = \theta^{(0)}$  and  $u^{(l)}(\theta) = \theta_1^{(l)} + \mathbf{i}\theta_2^{(l)}$  for  $l = 1, \dots, L$ . From (3.4), we may check that

$$u^{(l)}(g \cdot \theta) = e^{-\mathbf{i}2\pi l \mathbf{g}} u^{(l)}(\theta). \quad (\text{C.2})$$

Then, applying the identity  $\operatorname{Re} a = (a + \bar{a})/2$ ,

$$\langle \theta, g \cdot \vartheta \rangle = \sum_{l=0}^L \operatorname{Re} \left[ u^{(l)}(\theta) \cdot \overline{u^{(l)}(g \cdot \vartheta)} \right] = \frac{1}{2} \sum_{l=0}^L \left[ u^{(l)}(\theta) \overline{u^{(l)}(\vartheta)} e^{\mathbf{i}2\pi l \mathbf{g}} + \overline{u^{(l)}(\theta)} u^{(l)}(\vartheta) e^{-\mathbf{i}2\pi l \mathbf{g}} \right]. \quad (\text{C.3})$$

Taking the expected square on both sides gives

$$\begin{aligned} \mathbb{E}_g[\langle \theta, g \cdot \vartheta \rangle^2] &= \frac{1}{4} \mathbb{E}_{\mathbf{g}} \left[ \sum_{l_1, l_2=0}^L u^{(l_1)}(\theta) u^{(l_2)}(\theta) \overline{u^{(l_1)}(\vartheta) u^{(l_2)}(\vartheta)} e^{\mathbf{i}2\pi(l_1+l_2)\mathbf{g}} \right] \\ &\quad + \frac{1}{4} \mathbb{E}_{\mathbf{g}} \left[ \sum_{l_1, l_2=0}^L \overline{u^{(l_1)}(\theta) u^{(l_2)}(\theta)} u^{(l_1)}(\vartheta) u^{(l_2)}(\vartheta) e^{-\mathbf{i}2\pi(l_1+l_2)\mathbf{g}} \right] \\ &\quad + \frac{1}{4} \mathbb{E}_{\mathbf{g}} \left[ \sum_{l_1, l_2=0}^L u^{(l_1)}(\theta) \overline{u^{(l_2)}(\theta) u^{(l_1)}(\vartheta) u^{(l_2)}(\vartheta)} e^{\mathbf{i}2\pi(l_1-l_2)\mathbf{g}} \right] \\ &\quad + \frac{1}{4} \mathbb{E}_{\mathbf{g}} \left[ \sum_{l_1, l_2=0}^L \overline{u^{(l_1)}(\theta) u^{(l_2)}(\theta)} u^{(l_1)}(\vartheta) \overline{u^{(l_2)}(\vartheta)} e^{-\mathbf{i}2\pi(l_1-l_2)\mathbf{g}} \right], \end{aligned}$$

where  $\mathbf{g}$  is uniformly distributed over  $[0, 1)$ . Applying the property

$$\mathbb{E}_{\mathbf{g}}[e^{\mathbf{i}2\pi l \mathbf{g}}] = \begin{cases} 1 & \text{for } l = 0, \\ 0 & \text{for } l \neq 0, \end{cases} \quad (\text{C.4})$$

gives

$$\begin{aligned} \mathbb{E}_g[\langle \theta, g \cdot \vartheta \rangle^2] &= (u^{(0)}(\theta))^2 (u^{(0)}(\vartheta))^2 + \frac{1}{4} \sum_{l=1}^L \left[ u^{(l)}(\theta) \overline{u^{(l)}(\theta) u^{(l)}(\vartheta) u^{(l)}(\vartheta)} + \overline{u^{(l)}(\theta) u^{(l)}(\theta) u^{(l)}(\vartheta) u^{(l)}(\vartheta)} \right] \\ &= (\theta^{(0)})^2 (\vartheta^{(0)})^2 + \frac{1}{2} \sum_{l=1}^L r_l(\theta)^2 r_l(\vartheta)^2. \end{aligned}$$

Then from (C.1),

$$s_2(\theta) = \frac{1}{4} \left( (\theta^{(0)})^2 - (\theta_*^{(0)})^2 \right)^2 + \frac{1}{8} \sum_{l=1}^L \left( r_l(\theta)^2 - r_l(\theta_*)^2 \right)^2.$$

**Case  $k = 3$ :** We now take the expected cube on both sides of (C.3). Applying (C.4),

$$\begin{aligned} \mathbb{E}_g[\langle \theta, g \cdot \vartheta \rangle^3] &= \frac{1}{4} (u^{(0)}(\theta))^3 (u^{(0)}(\vartheta))^3 + \frac{3}{8} \sum_{\substack{l, l', l''=0 \\ l=l'+l''}}^L u^{(l)}(\theta) \overline{u^{(l')}(\theta) u^{(l'')}(\theta) u^{(l)}(\vartheta) u^{(l')}(\vartheta) u^{(l'')}(\vartheta)} \\ &\quad + \frac{3}{8} \sum_{\substack{l, l', l''=0 \\ l=l'+l''}}^L \overline{u^{(l)}(\theta) u^{(l')}(\theta) u^{(l'')}(\theta) u^{(l)}(\vartheta) u^{(l')}(\vartheta) u^{(l'')}(\vartheta)}. \end{aligned}$$

Let us write as shorthand  $u^{(l)} = u^{(l)}(\theta)$ ,  $u_*^{(l)} = u^{(l)}(\theta_*)$ , and similarly for  $r_l, \lambda_l, r_{l, l', l''}, \lambda_{l, l', l''}$  and  $r_{*, l}, \lambda_{*, l}, r_{*, l, l', l''}, \lambda_{*, l, l', l''}$ . Then from (C.1),

$$\begin{aligned} s_3(\theta) &= \frac{1}{12} \mathbb{E}_g \left[ \langle \theta, g \cdot \theta \rangle^3 - 2 \langle \theta, g \cdot \theta_* \rangle^3 + \langle \theta_*, g \cdot \theta_* \rangle^3 \right] \\ &= \frac{1}{48} \left( (u^{(0)})^3 - (u_*^{(0)})^3 \right)^2 + \frac{1}{16} \sum_{\substack{l, l', l''=0 \\ l=l'+l''}}^L \left| u^{(l)} u^{(l')} u^{(l'')} \right|^2 + \left| u_*^{(l)} u_*^{(l')} u_*^{(l'')} \right|^2 \end{aligned}$$

$$\begin{aligned}
& -\frac{1}{16} \sum_{\substack{l,l',l''=0 \\ l=l'+l''}}^L u^{(l)} \overline{u^{(l')} u^{(l'')}} u_*^{(l)} u_*^{(l')} u_*^{(l'')} + \overline{u^{(l)} u^{(l')} u^{(l'')}} u_*^{(l)} u_*^{(l')} u_*^{(l'')} \\
& = \frac{1}{48} \left( (u^{(0)})^3 - (u_*^{(0)})^3 \right)^2 + \frac{1}{16} \sum_{\substack{l,l',l''=0 \\ l=l'+l''}}^L \left| u^{(l)} \overline{u^{(l')} u^{(l'')}} - u_*^{(l)} \overline{u_*^{(l')} u_*^{(l'')}} \right|^2.
\end{aligned} \tag{C.5}$$

This verifies the first expression for  $s_3(\theta)$ . For the second expression, we split the second sum of (C.5) into the cases  $l = l' = l'' = 0$ , only  $l' = 0$  and  $l = l'' \geq 1$ , only  $l'' = 0$  and  $l = l' \geq 1$ , and all  $l, l', l'' \geq 1$ . The first three cases are easily rewritten in terms of  $u^{(0)}, r_l, u_*^{(0)}, r_{*,l}$ . Each term of the last case  $l, l', l'' \geq 1$  may be written as

$$\left| u^{(l)} \overline{u^{(l')} u^{(l'')}} - u_*^{(l)} \overline{u_*^{(l')} u_*^{(l'')}} \right|^2 = r_{l,l',l''}^2 + r_{*,l,l',l''}^2 - 2r_{l,l',l''} r_{*,l,l',l''} \cos(\lambda_{*,l,l',l''} - \lambda_{l,l',l''}),$$

and this yields the second expression for  $s_3(\theta)$ .  $\square$

*Proof of Theorem 3.1.* Note that for generic  $\theta \in \mathbb{R}^d$ , for example having  $(\theta_1^{(1)}, \theta_2^{(1)}) \neq (0, 0)$ , its stabilizer  $G_\theta = \{g \in G : g \cdot \theta = \theta\} = \{\text{Id}\}$  is trivial. Thus  $\dim \mathcal{O}_\theta = \dim G = 1$ , so  $\text{trdeg } \mathcal{R}^G = d - 1 = 2L$ .

We compute  $\text{trdeg}(\mathcal{R}_{\leq k}^G)$  for  $k = 1, 2, 3$  by applying Lemma 2.8 at any generic point  $\theta_* \in \mathbb{R}^d$  where  $r_l(\theta_*) > 0$  for each  $l = 1, \dots, L$ . Write as shorthand  $u^{(l)}(\theta) = r_l e^{i\lambda_l}$ ,  $u^{(l)}(\theta_*) = r_{*,l} e^{i\lambda_{*,l}}$ , and define  $t_l = \lambda_l - \lambda_{*,l} \in [-\pi, \pi)$  for each  $l = 1, \dots, L$ . Setting

$$\zeta(\theta) = (\theta_0, r_1, \dots, r_L, t_1, \dots, t_L),$$

this map  $\zeta(\theta)$  has non-singular derivative at  $\theta_*$ . Then by the inverse function theorem, the coordinates  $\zeta \in \mathbb{R}^{2L+1}$  provide an invertible reparametrization of  $\theta$  in a local neighborhood of  $\theta_*$ , with some inverse function  $\theta(\zeta)$ . Let  $\zeta_* = \zeta(\theta_*)$ . Note that  $\theta_*$  is a global minimizer and hence critical point of  $s_k(\theta)$  for each  $k \geq 1$ . Then by the chain rule,

$$\nabla_\zeta^2 s_k(\theta(\zeta)) \Big|_{\zeta=\zeta_*} = d_\zeta \theta(\zeta_*)^\top \cdot \nabla_{\theta_*}^2 s_k(\theta_*) \cdot d_\zeta \theta(\zeta_*).$$

Applying Lemma 2.8 and the fact that  $d_\zeta \theta(\zeta_*)$  is non-singular, this gives

$$\text{trdeg}(\mathcal{R}_{\leq k}^G) = \text{rank} \left( \nabla_{\theta_*}^2 s_1(\theta_*) + \dots + \nabla_{\theta_*}^2 s_k(\theta_*) \right) = \text{rank} \left( \nabla_\zeta^2 s_1(\theta(\zeta)) + \dots + \nabla_\zeta^2 s_k(\theta(\zeta)) \Big|_{\zeta=\zeta_*} \right).$$

For  $k = 1$  and  $k = 2$ , by Theorem 3.3,

$$s_1(\theta(\zeta)) = \frac{1}{2}(\theta_0 - \theta_{*,0})^2, \quad s_2(\theta(\zeta)) = \frac{1}{4}(\theta_0^2 - \theta_{*,0}^2)^2 + \frac{1}{8} \sum_{l=1}^L (r_l^2 - r_{*,l}^2)^2.$$

Then taking the Hessians yields

$$\text{trdeg}(\mathcal{R}_{\leq 1}^G) = \text{rank} \left( \nabla_\zeta^2 s_1(\theta(\zeta)) \Big|_{\zeta=\zeta_*} \right) = \text{rank} \left( \text{diag}(1, 0, \dots, 0) \right) = 1,$$

and

$$\begin{aligned}
\text{trdeg}(\mathcal{R}_{\leq 2}^G) &= \text{rank} \left( \nabla_\zeta^2 s_1(\theta(\zeta)) + \nabla_\zeta^2 s_2(\theta(\zeta)) \Big|_{\zeta=\zeta_*} \right) \\
&= \text{rank} \left( \text{diag}(1, 0, \dots, 0) + \text{diag}(2\theta_{*,0}^2, r_{*,1}^2, \dots, r_{*,L}^2) \right) = L + 1.
\end{aligned}$$

For  $k = 3$ , noting that  $\text{trdeg}(\mathcal{R}_{\leq 3}^G) \leq \text{trdeg}(\mathcal{R}^G) = 2L$ , it remains to show  $\text{trdeg}(\mathcal{R}_{\leq 3}^G) \geq 2L$ . Denote

$$H(\zeta_*) = \nabla_\zeta^2 s_1(\theta(\zeta)) + \nabla_\zeta^2 s_2(\theta(\zeta)) + \nabla_\zeta^2 s_3(\theta(\zeta)) \Big|_{\zeta=\zeta_*},$$

so that  $\text{trdeg}(\mathcal{R}_{\leq 3}^G) = \text{rank}(H(\zeta_*))$ . Group the coordinates of  $\zeta$  as  $r = (\theta_0, r_1, \dots, r_L) \in \mathbb{R}^{L+1}$  and  $t = (t_1, \dots, t_L) \in \mathbb{R}^L$ , and define the corresponding block decomposition

$$H(\zeta_*) = \begin{pmatrix} H_{rr}(\zeta_*) & H_{rt}(\zeta_*) \\ H_{tr}(\zeta_*) & H_{tt}(\zeta_*) \end{pmatrix}.$$

Since  $s_1(\theta(\zeta))$  and  $s_2(\theta(\zeta))$  are functions only of  $r$  and not of  $t$ , we have

$$\begin{aligned} H_{rr}(\zeta_*) &= \nabla_r^2 s_1(\theta(\zeta)) + \nabla_r^2 s_2(\theta(\zeta)) + \nabla_r^2 s_3(\theta(\zeta)) \Big|_{\zeta=\zeta_*}, \\ H_{rt}(\zeta_*) &= \nabla_{rt}^2 s_3(\theta(\zeta)) \Big|_{\zeta=\zeta_*}, \\ H_{tr}(\zeta_*) &= \nabla_{tr}^2 s_3(\theta(\zeta)) \Big|_{\zeta=\zeta_*}, \\ H_{tt}(\zeta_*) &= \nabla_t^2 s_3(\theta(\zeta)) \Big|_{\zeta=\zeta_*}. \end{aligned}$$

For the upper-left block  $H_{rr}(\zeta_*)$  of size  $(L+1) \times (L+1)$ , Lemma 2.8 ensures that each matrix  $\nabla_{\zeta}^2 s_k(\theta(\zeta))$  is positive semidefinite, and hence so is each submatrix  $\nabla_r^2 s_k(\theta(\zeta))$ . Then from the analysis for  $\text{trdeg}(\mathcal{R}_{\leq 2}^G)$ ,

$$\text{rank}(H_{rr}(\zeta_*)) \geq \text{rank} \left( \nabla_r^2 s_1(\theta(\zeta)) + \nabla_r^2 s_2(\theta(\zeta)) \Big|_{\zeta=\zeta_*} \right) = L+1, \quad (\text{C.6})$$

and equality must hold because  $H_{rr}(\zeta_*)$  has dimension  $L+1$ . For the blocks  $H_{rt}(\zeta_*)$  and  $H_{tr}(\zeta_*)$ , recall from the form of  $s_3(\theta)$  in Theorem 3.3 that

$$s_3(\theta(\zeta)) = f(r) - \frac{1}{8} \sum_{\substack{l, l', l''=1 \\ l=l'+l''}}^L r_{l, l', l''} r_{*, l, l', l''} \cos(t_l - t_{l'} - t_{l''}),$$

where  $f(r)$  is a function depending only on  $r$  and not on  $t$ . Then, noting that  $t_l = 0$  for all  $l = 1, \dots, L$  at  $\zeta = \zeta_*$ , we get

$$\nabla_{r\lambda}^2 s_3(\theta(\zeta)) \Big|_{\zeta=\zeta_*} = 0 \text{ and } \nabla_{\lambda r}^2 s_3(\theta(\zeta)) \Big|_{\zeta=\zeta_*} = 0,$$

which further implies

$$H_{rt}(\zeta_*) = 0 \text{ and } H_{tr}(\zeta_*) = 0. \quad (\text{C.7})$$

Finally, the lower-right block  $H_{tt}(\zeta_*)$  of size  $L \times L$  is given explicitly by

$$H_{tt}(\zeta_*) = -\frac{1}{8} \sum_{\substack{l, l', l''=1 \\ l=l'+l''}}^L r_{*, l, l', l''}^2 \cdot \nabla_t^2 [\cos(t_l - t_{l'} - t_{l''})] \Big|_{t=0}.$$

To show that its rank is at least  $L-1$  for generic  $\theta_* \in \mathbb{R}^d$ , it suffices to exhibit a single such point  $\theta_*$ . For simplicity, we pick  $\theta_*$  such that  $\theta_*^{(0)} = 1$  and  $r_{*, l} = 1$  for all  $l = 1, \dots, L$ . Let  $e_l$  be the  $l^{\text{th}}$  standard basis vector, and define the vector

$$w_{l, l', l''} = \nabla_t [t_l - t_{l'} - t_{l''}] = e_l - e_{l'} - e_{l''} \in \mathbb{R}^L. \quad (\text{C.8})$$

By the chain rule,

$$\nabla_t^2 [\cos(t_l - t_{l'} - t_{l''})] \Big|_{t=0} = -\cos(t_l - t_{l'} - t_{l''}) \Big|_{t=0} \cdot w_{l, l', l''} w_{l, l', l''}^\top = -w_{l, l', l''} w_{l, l', l''}^\top.$$

Define the index set  $\mathcal{L} = \{l, l', l'' \in \{1, \dots, L\} : l = l' + l''\}$ , and let  $W \in \mathbb{R}^{L \times |\mathcal{L}|}$  be the matrix with the vectors  $w_{l, l', l''}$  as columns. Then

$$H_{tt}(\zeta_*) = \frac{1}{8} W W^\top.$$

Note that, in particular,  $W$  has a subset of  $L-1$  columns corresponding to  $l' = 1, l'' \in \{1, \dots, L-1\}$ , and  $l = 1 + l''$ . These columns (in  $\mathbb{R}^L$ ) are given by

$$\begin{pmatrix} 2 \\ -1 \\ 0 \\ 0 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad \begin{pmatrix} 1 \\ 1 \\ -1 \\ 0 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \quad \begin{pmatrix} 1 \\ 0 \\ 1 \\ -1 \\ \vdots \\ 0 \\ 0 \end{pmatrix}, \dots, \quad \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \\ \vdots \\ 1 \\ -1 \end{pmatrix}.$$

The bottom  $L - 1$  rows of these columns form an upper-triangular matrix with non-zero diagonal, and hence these columns are linearly independent. Thus  $\text{rank}(W) \geq L - 1$ , so also

$$\text{rank}(H_{tt}(\zeta_*)) \geq L - 1. \quad (\text{C.9})$$

Combining (C.6), (C.7), and (C.9), we obtain  $\text{trdeg } \mathcal{R}_{\leq 3}^G = \text{rank}(H(\zeta_*)) \geq 2L$ . Hence  $\text{trdeg } \mathcal{R}_{\leq 3}^G = 2L$ , as desired. This proves Theorem 3.1.

The statement  $(d_0, d_1, d_2, d_3) = (1, 1, L, L - 1)$  in Corollary 3.2 now follows from these transcendence degrees. The smallest  $K$  for which  $\text{trdeg } \mathcal{R}_{\leq K}^G = \text{trdeg } \mathcal{R}^G = 2L$  is  $K = 3$ , except in the case  $L = 1$  where it is instead  $K = 2$ . This proves Corollary 3.2, in light of Theorem 2.7 and [BBSK<sup>+</sup>17, Theorem 4.9] (as reviewed in Lemma A.3).  $\square$

**C.2. Spurious local minimizers for continuous MRA.** In this section, we prove Theorem 3.4 showing that in the continuous MRA model where  $L \geq 30$ , for some open subset of true signal vectors  $\theta_* \in \mathbb{R}^d$  and for sufficiently high noise  $\sigma^2 > 0$ , there must exist spurious local minimizers of the negative population log-likelihood function  $R(\theta)$ .

*Proof of Theorem 3.4.* Recall the forms of  $s_1(\theta)$ ,  $s_2(\theta)$ , and  $s_3(\theta)$  from Theorem 3.3. By definition of  $s_k(\theta)$  in (2.8), each moment variety  $\mathcal{V}_k(\theta_*)$  is the intersection of the global minimizers of  $s_1(\theta), \dots, s_k(\theta)$ . Then

$$\mathcal{V}_1(\theta_*) = \{\theta : \theta^{(0)} = \theta_*^{(0)}\}, \quad \mathcal{V}_2(\theta_*) = \{\theta : \theta^{(0)} = \theta_*^{(0)}, r_l(\theta) = r_l(\theta_*) \text{ for each } l = 1, \dots, L\}.$$

On  $\mathcal{V}_0(\theta_*) = \mathbb{R}^d$ , we have  $\nabla s_1(\theta)|_{\mathcal{V}_0(\theta_*)} = \theta^{(0)} - \theta_*^{(0)}$ , and this vanishes exactly when  $\theta \in \mathcal{V}_1(\theta_*)$ . Differentiating in the coordinates  $\{(\theta_1^{(l)}, \theta_2^{(l)}) : l = 1, \dots, L\}$  that parametrize  $\mathcal{V}_1(\theta_*)$ , and applying  $r_l(\theta)^2 = (\theta_1^{(l)})^2 + (\theta_2^{(l)})^2$ , we have

$$\nabla s_2(\theta)|_{\mathcal{V}_1(\theta_*)} = \frac{1}{2} \left( (r_l(\theta)^2 - r_l(\theta_*)^2) \cdot (\theta_1^{(l)}, \theta_2^{(l)}) : l = 1, \dots, L \right).$$

Suppose  $\theta_*$  satisfies the generic condition  $r_l(\theta_*) > 0$  for each  $l = 1, \dots, L$ . Then  $\nabla s_2(\theta)|_{\mathcal{V}_1(\theta_*)} = 0$  if and only if, for each  $l = 1, \dots, L$ , either  $r_l(\theta) = r_l(\theta_*)$  or  $(\theta_1^{(l)}, \theta_2^{(l)}) = (0, 0)$ . If the latter holds for any  $l = 1, \dots, L$ , then differentiating in  $(\theta_1^{(l)}, \theta_2^{(l)})$  a second time shows that the Hessian of  $s_2(\theta)$  in  $(\theta_1^{(l)}, \theta_2^{(l)})$  is negative-definite at  $(\theta_1^{(l)}, \theta_2^{(l)}) = (0, 0)$ , and hence  $\lambda_{\min}(\nabla^2 s_2(\theta)|_{\mathcal{V}_1(\theta_*)}) < 0$ . On the other hand, if  $r_l(\theta) = r_l(\theta_*)$  for every  $l = 1, \dots, L$ , then  $\theta \in \mathcal{V}_2(\theta_*)$  and  $\theta$  is a global minimizer of  $s_2(\theta)$ . Thus, the minimizations of  $s_1(\theta)$  and  $s_2(\theta)$  on  $\mathcal{V}_0(\theta_*)$  and  $\mathcal{V}_1(\theta_*)$  are globally benign.

We now take  $L \geq 30$ , and construct the example of  $\theta_*$  where  $s_3(\theta)$  has a spurious local minimizer in  $\mathcal{V}_2(\theta_*)$  that is nondegenerate up to orbit. Consider  $\theta_*$  such that  $r_{*,l} := r_l(\theta_*) > 0$  for each  $l = 1, \dots, L$ . Then  $\mathcal{V}_2(\theta_*)$  may be smoothly parametrized by the coordinates  $t = (t_1, \dots, t_L)$  where  $t_l = \lambda_l(\theta) - \lambda_l(\theta_*)$ . The function  $8s_3(\theta)$  restricted to  $\mathcal{V}_2(\theta_*)$  is given as a function of  $t$  by

$$s(t) = - \sum_{\substack{l, l', l''=1 \\ l=l'+l''}}^L r_{*,l}^2 r_{*,l'}^2 r_{*,l''}^2 \cos(t_l - t_{l'} - t_{l''}) + \text{constant}.$$

The orbit  $\mathcal{O}_{\theta_*} \cap \mathcal{V}_2(\theta_*)$  is defined by

$$\{(t_1, \dots, t_L) : t_l \equiv \tau \cdot l \pmod{2\pi} \text{ for all } l = 1, \dots, L \text{ and some } \tau \in \mathbb{R}\}, \quad (\text{C.10})$$

where  $\tau = 0$  corresponds to the point  $\theta_*$  itself. Thus, our goal is to exhibit a point  $\theta_*$  for which

$$\nabla s(\hat{t}) = 0, \quad \nabla^2 s(\hat{t}) \succeq 0, \quad \text{and} \quad \text{rank}(\nabla^2 s(\hat{t})) = L - 1 \quad (\text{C.11})$$

at some  $\hat{t}$  not belonging to this orbit (C.10). Then the corresponding point  $\theta \in \mathbb{R}^d$  where  $\theta^{(0)} = \theta_*^{(0)}$ ,  $r_l(\theta) = r_{*,l}$ , and  $t_l(\theta) = \hat{t}_l$  for each  $l = 1, \dots, L$  is our desired spurious local minimizer for  $s_3(\theta)$ . Note that the condition (C.11) depends on  $\theta_*$  only via  $r_* = (r_{*,1}, \dots, r_{*,L})$ , so equivalently, our goal is to construct an appropriate such vector  $r_*$ .

We split the construction into two steps: First, we construct  $\tilde{r}_*$  for which  $\hat{t} = (\pi, 0, \dots, 0)$  satisfies

$$\nabla s(\hat{t}) = 0, \quad \nabla^2 s(\hat{t}) \succeq 0, \quad \text{rank}(\nabla^2 s(\hat{t})) = L - 2.$$

This Hessian  $\nabla^2 s(\hat{t})$  will have a dimension-2 null space spanned by the vectors  $e_3 = (0, 0, 1, 0, \dots, 0)$  and  $e = (1, 2, \dots, L)$ . Second, we make a small perturbation of the third coordinate of  $\tilde{r}_*$ , to eliminate the null

vector  $e_3$  while preserving  $\nabla s(\hat{t}) = 0$  and  $\nabla^2 s(\hat{t}) \succeq 0$ . This yields  $r_*$  satisfying (C.11).

**Step I.** Clearly  $\hat{t} = (\pi, 0, \dots, 0)$  satisfies  $\nabla s(\hat{t}) = 0$  for any choice of  $r_*$ , because  $\sin(k\pi) = 0$  for any integer  $k$ . The Hessian of  $s(t)$  is given by

$$\nabla^2 s(t) = \sum_{\substack{l, l', l''=1 \\ l=l'+l''}}^L r_{*,l}^2 r_{*,l'}^2 r_{*,l''}^2 \cos(t_l - t_{l'} - t_{l''}) (e_l - e_{l'} - e_{l''})(e_l - e_{l'} - e_{l''})^\top,$$

where  $e_l$  is the  $l^{\text{th}}$  standard basis vector. Then for the above choice of  $\hat{t}$  and for any vectors  $u, v \in \mathbb{R}^L$ ,

$$u^\top \nabla^2 s(\hat{t}) v = \sum_{\substack{l, l', l''=1 \\ l=l'+l''}}^L r_{*,l}^2 r_{*,l'}^2 r_{*,l''}^2 (u_l - u_{l'} - u_{l''})(v_l - v_{l'} - v_{l''}) \times \begin{cases} -1 & \text{if exactly one of } l', l'' \text{ equals } 1, \\ 1 & \text{otherwise.} \end{cases} \quad (\text{C.12})$$

Consider  $\tilde{r}_* = (\tilde{r}_{*,1}, \tilde{r}_{*,2}, \tilde{r}_{*,3}, \tilde{r}_{*,4}, \dots, \tilde{r}_{*,L}) = (1, L^{\kappa/2}, 0, 1, \dots, 1)$  where  $\tilde{r}_{*,\ell} = 1$  for all  $\ell \geq 4$ , and for a constant  $\kappa > 0$  to be determined later. Then from (C.12) applied with  $v = e_3$  and  $v = e = (1, 2, \dots, L)$ , it is immediate that  $u^\top \nabla^2 s(\hat{t}) v = 0$  for any vector  $u \in \mathbb{R}^L$ , so these two vectors  $v = e_3$  and  $v = e$  belong to the null space of  $\nabla^2 s(\hat{t})$ .

We now check that for any other unit vector  $v \in \mathbb{R}^L$  orthogonal to both  $e_3$  and  $e$ , we have  $v^\top \nabla^2 s(\hat{t}) v > 0$  strictly. Observe from (C.12) that for  $r_* = \tilde{r}_*$ ,

$$\begin{aligned} v^\top \nabla^2 s(\hat{t}) v &= \sum_{\substack{l, l', l''=4 \\ l=l'+l''}}^L (v_l - v_{l'} - v_{l''})^2 + L^\kappa (v_2 - 2v_1)^2 + L^{2\kappa} (v_4 - 2v_2)^2 \\ &\quad + 2L^\kappa \sum_{l=4}^{L-2} (v_{l+2} - v_l - v_2)^2 - 2 \sum_{l=4}^{L-1} (v_{l+1} - v_l - v_1)^2. \end{aligned} \quad (\text{C.13})$$

Note that only the last term is negative. Denote  $\epsilon = \sqrt{6}L^{\frac{1-\kappa}{2}}$ . We consider two cases:

**Case 1:** Any one of  $|v_2 - 2v_1|$ ,  $|v_4 - 2v_2|$ ,  $\{|v_{l+2} - v_l - v_2| : l \geq 4\}$  is larger than  $\epsilon$ . Then let us upper bound the last term of (C.13) by

$$2 \sum_{l=4}^{L-1} (v_{l+1} - v_l - v_1)^2 \leq 6 \sum_{l=4}^{L-1} (v_{l+1}^2 + v_l^2 + v_1^2) \leq 6L,$$

where the last inequality follows from  $\|v\|_2^2 = 1$ . Then

$$v^\top \nabla^2 s(\hat{t}) v > L^\kappa \epsilon^2 - 2 \sum_{l=4}^{L-1} (v_{l+1} - v_l - v_1)^2 \geq 0.$$

**Case 2:** We have instead

$$|v_2 - 2v_1| \leq \epsilon, \quad |v_4 - 2v_2| \leq \epsilon, \quad |v_{l+2} - v_l - v_2| \leq \epsilon \text{ for } l \geq 4. \quad (\text{C.14})$$

In this case, we aim to show that the first term in (C.13) is large enough to compensate for the negative last term of (C.13).

For all  $m \geq 1$ , the second and third inequalities of (C.14) imply  $|v_{2m} - mv_2| \leq (m-1)\epsilon$ . Similarly, for all  $m \geq 0$ , the last inequality implies  $|v_{2m+5} - v_5 - mv_2| \leq m\epsilon$ . Combining with  $|mv_2 - 2mv_1| \leq m\epsilon$  by the first inequality of (C.14), we obtain

$$|v_{2m} - 2mv_1| \leq L\epsilon \text{ for } 1 \leq m \leq L/2, \quad |v_{2m+5} - v_5 - 2mv_1| \leq L\epsilon \text{ for } 0 \leq m \leq (L-5)/2. \quad (\text{C.15})$$

For the summands of the first term in (C.13), if  $l' = 2m' + 5$  and  $l'' = 2m'' + 5$  are both odd where  $m', m'' \geq 0$ , then  $l = 2m + 10$  for  $m = m' + m''$ , and we have

$$|v_l - v_{l'} - v_{l''}| \geq |10v_1 - 2v_5| - 3L\epsilon$$

by (C.15) and the triangle inequality

$$|v_l - v_{l'} - v_{l''}| + |-v_l + (2m + 10)v_1| + |v_{l'} - v_5 - 2m'v_1| + |v_{l''} - v_5 - 2m''v_1| \geq |10v_1 - 2v_5|.$$

For any even  $l = 2m + 10$  with  $m \geq 0$ , the number of odd pairs  $l', l'' \geq 5$  where  $l' + l'' = 2m + 10$  is  $m + 1$ . Then the total number of tuples  $(l, l', l'')$  in the first term of (C.13) where  $l', l''$  are odd is

$$\sum_{m=0}^{\lfloor L/2 \rfloor - 5} (m + 1) = \frac{1}{2}(\lfloor L/2 \rfloor - 3)(\lfloor L/2 \rfloor - 4) \geq \frac{1}{8}(L - 8)(L - 10).$$

Hence, we may lower bound the first term in (C.13) by

$$\sum_{\substack{l, l', l''=4 \\ l=l'+l''}}^L (v_l - v_{l'} - v_{l''})^2 \geq \frac{1}{8}(L - 8)(L - 10) \left( |10v_1 - 2v_5| - 3L\epsilon \right)_+^2. \quad (\text{C.16})$$

Similarly, the summands of the last term in (C.13) for  $l \geq 4$  may be upper bounded as

$$|v_{l+1} - v_l - v_1| \leq |5v_1 - v_5| + 2L\epsilon \quad (\text{C.17})$$

by applying (C.15) to approximate both  $v_{l+1}$  and  $v_l$  via linear combinations of  $v_1, v_5$ , and using the triangle inequality. (This may be checked separately in the cases where  $l$  is odd and even.) The number of summands in this last term is  $L - 4$ . Then, combining (C.16) and (C.17) gives

$$v^\top \nabla^2 s(\hat{t}) v \geq \frac{1}{8}(L - 8)(L - 10) \left( |10v_1 - 2v_5| - 3L\epsilon \right)_+^2 - 2(L - 4) \left( |5v_1 - v_5| + 2L\epsilon \right)^2 \quad (\text{C.18})$$

We now claim that for sufficiently large  $\kappa > 0$ , we must have

$$|5v_1 - v_5| > 5L\epsilon. \quad (\text{C.19})$$

To show this claim, since  $v \perp e_3$  and  $v \perp e$ ,

$$0 = -3v_3 = \sum_{\substack{l=1 \\ l \neq 3}}^L l v_l = v_1 + 5v_5 + \sum_{m=1}^{\lfloor L/2 \rfloor} 2m \cdot v_{2m} + \sum_{m=1}^{\lfloor (L-5)/2 \rfloor} (2m + 5) \cdot v_{2m+5}.$$

Denote  $a_{2m} = v_{2m} - 2m \cdot v_1$  and  $a_{2m+5} = v_{2m+5} - v_5 - 2m \cdot v_1$  for  $m \geq 1$ , where these satisfy  $|a_l| \leq L\epsilon$  by (C.15). Let us define

$$M_1 = 1 + \sum_{m=1}^{\lfloor L/2 \rfloor} 4m^2 + \sum_{m=1}^{\lfloor (L-5)/2 \rfloor} 2m(2m + 5), \quad M_5 = 5 + \sum_{m=1}^{\lfloor (L-5)/2 \rfloor} (2m + 5).$$

Then we may write the above as

$$\begin{aligned} 0 &= v_1 + 5v_5 + \sum_{m=1}^{\lfloor L/2 \rfloor} 2m \cdot (a_{2m} + 2m \cdot v_1) + \sum_{m=1}^{\lfloor (L-5)/2 \rfloor} (2m + 5) \cdot (a_{2m+5} + v_5 + 2m \cdot v_1) \\ &= M_1 v_1 + M_5 v_5 + \sum_{\substack{l=1 \\ l \neq 1, 3, 5}}^L l a_l. \end{aligned}$$

This may be rearranged as

$$v_1 = \frac{M_5(5v_1 - v_5) - \sum_{l: l \neq 1, 3, 5} l a_l}{M_1 + 5M_5}.$$

Now suppose by contradiction that  $|5v_1 - v_5| \leq 5L\epsilon$ . Then this implies

$$|v_1| \leq \frac{5M_5 L + L^2(L + 1)/2}{M_1 + 5M_5} \epsilon < C\epsilon,$$

where the second inequality holds for a universal constant  $C > 0$  and any  $L \geq 1$ . Then  $|v_5| \leq |5v_1| + 5L\epsilon < 5(L + C)\epsilon$ , and combining with (C.15) gives

$$|v_l| \leq C' L\epsilon \text{ for all } l \in \{1, \dots, L\} \setminus \{3\}$$

and a different universal constant  $C' > 0$ . Recalling  $v_3 = 0$  and  $\epsilon = \sqrt{6}L^{\frac{1-\kappa}{2}}$ , this contradicts that  $\|v\|_2 = 1$  for sufficiently large  $\kappa > 0$ . Thus, (C.19) holds.

Finally, this bound (C.19) implies  $|10v_1 - 2v_5| - 3L\epsilon > |5v_1 - v_5| + 2L\epsilon > 0$ . For  $L \geq 30$ , we have  $(L-8)(L-10)/8 \geq 2(L-4)$ . Applying these to (C.18) yields  $v^\top s(\hat{t})v > 0$  as desired.

**Step II.** We now show that making a small positive perturbation to  $\tilde{r}_{*,3}$  yields a point  $r_*$  which satisfies (C.11) at  $\hat{t} = (\pi, 0, \dots, 0)$ . Denote  $q_* = r_{*,3}^2$  and set  $\tilde{q}_* = \tilde{r}_{*,3}^2 = 0$ . Let  $\Pi \in \mathbb{R}^{L \times (L-1)}$  have orthonormal columns spanning the orthogonal complement of  $e = (1, 2, \dots, L)$ , and consider the projected Hessian

$$H(q_*) = \Pi^\top \nabla^2 s(\hat{t}) \Pi \in \mathbb{R}^{(L-1) \times (L-1)}$$

now as a function of  $q_*$ . By the result of Step I,  $H(\tilde{q}_*) \succeq 0$ , and  $H(\tilde{q}_*)$  has a simple eigenvalue  $\mu = 0$  with eigenvector  $v = \Pi^\top e_3 / \|\Pi^\top e_3\|$ . Then this eigenvalue  $\mu = \mu(q_*)$  is differentiable in  $q_*$ , with derivative given by  $\partial_{q_*} \mu(q_*) = v^\top \partial_{q_*} H(q_*) v$ . Applying  $\Pi \cdot \Pi^\top = \text{Id} - ee^\top / \|e\|^2$  and the fact that  $e = (1, 2, \dots, L)$  belongs to the null space of  $\nabla_t^2 s(\hat{t})$  for any  $q_*$ , this is

$$\partial_{q_*} \mu(q_*) = \frac{e_3^\top (\text{Id} - ee^\top / \|e\|^2) \cdot \partial_{q_*} \nabla_t^2 s(\hat{t}) \cdot (\text{Id} - ee^\top / \|e\|^2) e_3}{e_3^\top (\text{Id} - ee^\top / \|e\|^2) e_3} = \frac{e_3^\top \cdot \partial_{q_*} \nabla_t^2 s(\hat{t}) \cdot e_3}{e_3^\top (\text{Id} - ee^\top / \|e\|^2) e_3}.$$

By (C.12) applied with  $u = v = e_3$ , for general  $r_*$ , we have

$$e_3^\top \nabla_t^2 s(\hat{t}) e_3 = -2r_{*,1}^2 r_{*,2}^2 r_{*,3}^2 - 2r_{*,1}^2 r_{*,3}^2 r_{*,4}^2 + 2 \sum_{l=2}^{L-3} r_{*,3}^2 r_{*,l}^2 r_{*,l+3}^2 (1 + \mathbf{1}\{l=3\}).$$

Then differentiating in  $q_* = r_{*,3}^2$  and evaluating at  $\tilde{r}_* = (1, L^\kappa, 0, 1, \dots, 1)$ ,

$$e_3^\top \cdot \partial_{q_*} \nabla_t^2 s(\hat{t}) \cdot e_3 \Big|_{q_* = \tilde{q}_*} = -2L^{2\kappa} - 2 + 2L^{2\kappa} + 2(L-6) > 0.$$

Thus, for some sufficiently small  $\delta > 0$ , setting  $q_* = \delta$  yields  $\mu(\delta) > 0$ , and hence  $H(\delta) \succ 0$  strictly. Then at the point  $r_* = (1, L^\kappa, \delta, 1, \dots, 1)$ , we obtain that (C.11) holds.

Combining Steps I and II, we have shown that  $\theta_*$  given by  $(r_1(\theta_*), \dots, r_L(\theta_*)) = (1, L^\kappa, \delta, 1, \dots, 1)$  and (say)  $\theta_*^{(0)} = 0$  and  $(\lambda_1(\theta_*), \dots, \lambda_L(\theta_*)) = 0$  satisfies (C.11). Then (C.11) holds also in a sufficiently small open neighborhood  $U$  of  $\theta_*$ , by continuity, where  $e = (1, 2, 3, \dots, L)$  is the fixed vector in the null space of  $\nabla^2 s(\hat{t})$  for every  $\theta_* \in U$ . Then for any  $\theta_* \in U$ , the function  $s_3(\theta)$  has a spurious local minimizer  $\theta \in \mathcal{V}_2(\theta_*)$  that is non-degenerate up to orbit, where  $\theta^{(0)} = \theta_*^{(0)}$ ,  $r_l(\theta) = r_l(\theta_*)$  for all  $l \geq 1$ ,  $\lambda_1(\theta) = \lambda_1(\theta_*) + \pi$ , and  $\lambda_l(\theta) = \lambda_l(\theta_*)$  for all  $l \geq 2$ . This concludes the proof.  $\square$

**C.3. Projected continuous MRA.** In this section, we now describe a projected version of the continuous MRA problem with a two-fold projection onto an interval. We analyze this as a simpler example of a model with projection, before diving into the projected cryo-EM model to follow.

Again writing  $\mathcal{S}^1 \cong [0, 1)$  for the unit circle and  $f_g(t)$  for the periodic function  $f : \mathcal{S}^1 \rightarrow \mathbb{R}$  rotated by  $g \in \text{SO}(2) \cong [0, 1)$ , we consider the observations

$$(\Pi \cdot f_g)(t) dt + \sigma dW(t)$$

over  $t \in (0, 1/2)$  where

$$(\Pi \cdot f_g)(t) = f_g(t) + f_g(1-t)$$

and  $dW(t)$  is a standard Gaussian white noise process on the interval  $(0, 1/2)$ . The map  $\Pi$  represents a two-fold projection of the circle  $\mathcal{S}^1$  onto the interval  $(0, 1/2)$ .

To represent this projected model in a Gaussian sequence space, observe that for the Fourier basis (3.2), we have  $\Pi \cdot h_{l2} = 0$  for all  $l \geq 1$ , while  $(\Pi \cdot h_0)/\sqrt{2}$  and  $(\Pi \cdot h_{11})/\sqrt{2}$  form an orthonormal basis over  $(0, 1/2)$ . Thus, expressing  $\Pi \cdot f$  in this projected basis,  $\Pi$  may be represented as a linear map  $\Pi : \mathbb{R}^d \rightarrow \mathbb{R}^{\tilde{d}}$  for  $\tilde{d} = L + 1$ , where

$$\Pi(\theta) = \sqrt{2}(\theta^{(0)}, \theta_1^{(1)}, \dots, \theta_1^{(L)}). \quad (\text{C.20})$$

In this projected basis, the above observation model corresponds to the projected orbit recovery model (2.2) where  $g \in \mathbf{G}$  is a random rotation from the same group  $\mathbf{G}$  as in (3.4).

The following result shows that the decomposition of total dimension in Theorem 2.7 is the same as in the unprojected setting. In particular,  $\text{trdeg } \tilde{\mathcal{R}}_{\leq \tilde{K}}^{\mathcal{G}} = \text{trdeg } \mathcal{R}^{\mathcal{G}}$  for  $\tilde{K} = 3$ . This model is a continuous analogue of the projected discrete MRA model studied in [BBSK<sup>+</sup>17, Section 5.3.1], where an analogous conclusion was described as [BBSK<sup>+</sup>17, Conjecture 5.3].

**Theorem C.1.** *For any  $L \geq 1$ , we have*

$$\text{trdeg}(\tilde{\mathcal{R}}_{\leq 1}^{\mathcal{G}}) = 1, \quad \text{trdeg}(\tilde{\mathcal{R}}_{\leq 2}^{\mathcal{G}}) = L + 1, \quad \text{trdeg}(\tilde{\mathcal{R}}_{\leq 3}^{\mathcal{G}}) = \text{trdeg}(\mathcal{R}^{\mathcal{G}}) = 2L,$$

which match the values of  $\text{trdeg}(\mathcal{R}_{\leq 1}^{\mathcal{G}})$ ,  $\text{trdeg}(\mathcal{R}_{\leq 2}^{\mathcal{G}})$ , and  $\text{trdeg}(\mathcal{R}_{\leq 3}^{\mathcal{G}})$  in the unprojected setting of Theorem 3.1.

The following result describes the forms of  $\tilde{s}_k(\theta)$  for  $k = 1, 2, 3$ , which are similar to those in the unprojected setting. The minimizations of  $\tilde{s}_1(\theta)$ ,  $\tilde{s}_2(\theta)$ , and  $\tilde{s}_3(\theta)$  are also optimization problems over the signal mean, Fourier power spectrum, and Fourier bispectrum respectively, although the specific forms are different from the unprojected counterparts.

**Theorem C.2.** *For any  $L \geq 1$ ,*

$$\begin{aligned} \tilde{s}_1(\theta) &= \left( \theta^{(0)} - \theta_*^{(0)} \right)^2 \\ \tilde{s}_2(\theta) &= \left( (\theta^{(0)})^2 - (\theta_*^{(0)})^2 \right)^2 + \frac{1}{4} \sum_{l=1}^L \left( r_l(\theta)^2 - r_l(\theta_*)^2 \right)^2 \\ \tilde{s}_3(\theta) &= \frac{2}{3} \left( (\theta^{(0)})^3 - (\theta_*^{(0)})^3 \right)^2 + \frac{1}{2} \sum_{l=1}^L \left( \theta^{(0)} r_l(\theta)^2 - \theta_*^{(0)} r_l(\theta_*)^2 \right)^2 \\ &\quad + \frac{1}{16} \sum_{\substack{l, l', l''=1 \\ l=l'+l''}}^L \left( r_{l, l', l''}(\theta)^2 + r_{l, l', l''}(\theta_*)^2 - 2r_{l, l', l''}(\theta)r_{l, l', l''}(\theta_*) \cos(\lambda_{l, l', l''}(\theta) - \lambda_{l, l', l''}(\theta_*)) \right. \\ &\quad \left. + r_{l, l', l''}(\theta)^2 \cos(2\lambda_{l, l', l''}(\theta)) + r_{l, l', l''}(\theta_*)^2 \cos(2\lambda_{l, l', l''}(\theta_*)) \right. \\ &\quad \left. - 2r_{l, l', l''}(\theta)r_{l, l', l''}(\theta_*) \cos(\lambda_{l, l', l''}(\theta) + \lambda_{l, l', l''}(\theta_*)) \right). \end{aligned}$$

In the remainder of this section, we prove Theorems C.2 and C.1.

*Proof of Theorem C.2.* Recall by Lemma A.1 that

$$\tilde{s}_k(\theta) = \frac{1}{2(k!)} \mathbb{E}_{g, h} [\langle \Pi \cdot g \cdot \theta, \Pi \cdot h \cdot \theta \rangle^k - 2\langle \Pi \cdot g \cdot \theta, \Pi \cdot h \cdot \theta_* \rangle^k + \langle \Pi \cdot g \cdot \theta_*, \Pi \cdot h \cdot \theta_* \rangle^k]. \quad (\text{C.21})$$

We compute  $\mathbb{E}_{g, h} [\langle \Pi \cdot g \cdot \theta, \Pi \cdot h \cdot \vartheta \rangle^k]$  for  $\theta, \vartheta \in \mathbb{R}^d$ .

**Case  $k = 1$ :** Recall  $u^{(0)}(\theta) = \theta^{(0)}$  and  $u^{(l)}(\theta) = \theta_1^{(l)} + \mathbf{i}\theta_2^{(l)}$  for  $l = 1, \dots, L$ . Write  $g, h \in \mathcal{G}$  corresponding to the rotations  $\mathbf{g}, \mathbf{h} \in [0, 1)$ . Then, applying (C.20), (C.2), and the identity  $\text{Re } a \cdot \text{Re } b = (ab + \bar{a}\bar{b} + a\bar{b} + \bar{a}b)/4$ , we have

$$\langle \Pi \cdot g \cdot \theta, \Pi \cdot h \cdot \vartheta \rangle = 2 \sum_{l=0}^L \text{Re } u^{(l)}(g \cdot \theta) \cdot \text{Re } u^{(l)}(h \cdot \vartheta) = \frac{1}{2} M_1 + \frac{1}{2} M_2 \quad (\text{C.22})$$

where

$$\begin{aligned} M_1 &:= \sum_{l=0}^L u^{(l)}(\theta) u^{(l)}(\vartheta) e^{-2\mathbf{i}\pi l(\mathbf{g}+\mathbf{h})} + \overline{u^{(l)}(\theta) u^{(l)}(\vartheta)} e^{2\mathbf{i}\pi l(\mathbf{g}+\mathbf{h})}, \\ M_2 &:= \sum_{l=0}^L u^{(l)}(\theta) \overline{u^{(l)}(\vartheta)} e^{-2\mathbf{i}\pi l(\mathbf{g}-\mathbf{h})} + \overline{u^{(l)}(\theta)} u^{(l)}(\vartheta) e^{2\mathbf{i}\pi l(\mathbf{g}-\mathbf{h})}. \end{aligned}$$

For independent and uniformly random  $\mathbf{g}, \mathbf{h} \in [0, 1)$ , taking the expected value on both sides using (C.4) gives  $\mathbb{E}_{g, h} [\langle \Pi \cdot g \cdot \theta, \Pi \cdot h \cdot \vartheta \rangle] = 2u^{(0)}(\theta)u^{(0)}(\vartheta)$ . Then from (C.21), we obtain

$$\tilde{s}_1(\theta) = \left( u^{(0)}(\theta) - u^{(0)}(\theta_*) \right)^2 = \left( \theta^{(0)} - \theta_*^{(0)} \right)^2. \quad (\text{C.23})$$

**Case  $k = 2$ :** Taking the expected square on both sides of (C.22), we have

$$\mathbb{E}_{g,h}[\langle \Pi \cdot g \cdot \theta, \Pi \cdot h \cdot \vartheta \rangle^2] = \frac{1}{4} \left\{ \mathbb{E}_{g,h}[M_1^2] + 2\mathbb{E}_{g,h}[M_1 M_2] + \mathbb{E}_{g,h}[M_2^2] \right\}.$$

Applying (C.4) and an argument similar to the  $k = 2$  computation in the proof of Theorem 3.3,

$$\begin{aligned} \mathbb{E}_{g,h}[M_1^2] &= 2(u^{(0)}(\theta))^2(u^{(0)}(\vartheta))^2 + 2 \sum_{l=0}^L |u^{(l)}(\theta)|^2 |u^{(l)}(\vartheta)|^2, \\ \mathbb{E}_{g,h}[M_1 M_2] &= 4(u^{(0)}(\theta))^2(u^{(0)}(\vartheta))^2, \\ \mathbb{E}_{g,h}[M_2^2] &= 2(u^{(0)}(\theta))^2(u^{(0)}(\vartheta))^2 + 2 \sum_{l=0}^L |u^{(l)}(\theta)|^2 |u^{(l)}(\vartheta)|^2. \end{aligned}$$

Then, separating the  $l = 0$  terms from these sums,  $\mathbb{E}_{g,h}[\langle \Pi \cdot g \cdot \theta, \Pi \cdot h \cdot \vartheta \rangle^2] = 4(\theta^{(0)})^2(\vartheta^{(0)})^2 + \sum_{l=1}^L r_l(\theta)^2 r_l(\vartheta)^2$ , so by (C.21),

$$\tilde{s}_2(\theta) = \left( (\theta^{(0)})^2 - (\theta_*^{(0)})^2 \right)^2 + \frac{1}{4} \sum_{l=1}^L \left( r_l(\theta)^2 - r_l(\theta_*)^2 \right)^2. \quad (\text{C.24})$$

**Case  $k = 3$ :** Taking the expected cube on both sides of (C.22), we have

$$\mathbb{E}_{g,h}[\langle \Pi \cdot g \cdot \theta, \Pi \cdot h \cdot \vartheta \rangle^3] = \frac{1}{8} \left\{ \mathbb{E}_{g,h}[M_1^3] + 3\mathbb{E}_{g,h}[M_1^2 M_2] + 3\mathbb{E}_{g,h}[M_1 M_2^2] + \mathbb{E}_{g,h}[M_2^3] \right\}.$$

Applying (C.4) and an argument similar to the  $k = 3$  computation in the proof of Theorem 3.3,

$$\begin{aligned} \mathbb{E}_{g,h}[M_1^3] &= 2(u^{(0)}(\theta))^3(u^{(0)}(\vartheta))^3 + 3 \sum_{\substack{l,l',l''=0 \\ l=l'+l''}}^L u^{(l)}(\theta) \overline{u^{(l')}(\theta) u^{(l'')}(\theta)} u^{(l)}(\vartheta) \overline{u^{(l')}(\vartheta) u^{(l'')}(\vartheta)} \\ &\quad + 3 \sum_{\substack{l,l',l''=0 \\ l=l'+l''}}^L \overline{u^{(l)}(\theta) u^{(l')}(\theta) u^{(l'')}(\theta)} \overline{u^{(l)}(\vartheta) u^{(l')}(\vartheta) u^{(l'')}(\vartheta)}, \\ \mathbb{E}_{g,h}[M_1^2 M_2] &= \mathbb{E}_{g,h}[M_1 M_2^2] = 8(u^{(0)}(\theta))^3(u^{(0)}(\vartheta))^3 + 4u^{(0)}(\theta)u^{(0)}(\vartheta) \cdot \sum_{l=1}^L |u^{(l)}(\theta)|^2 |u^{(l)}(\vartheta)|^2, \\ \mathbb{E}_{g,h}[M_2^3] &= 2(u^{(0)}(\theta))^3(u^{(0)}(\vartheta))^3 + 3 \sum_{\substack{l,l',l''=0 \\ l=l'+l''}}^L u^{(l)}(\theta) \overline{u^{(l')}(\theta) u^{(l'')}(\theta) u^{(l)}(\vartheta) u^{(l')}(\vartheta) u^{(l'')}(\vartheta)} \\ &\quad + 3 \sum_{\substack{l,l',l''=0 \\ l=l'+l''}}^L \overline{u^{(l)}(\theta) u^{(l')}(\theta) u^{(l'')}(\theta) u^{(l)}(\vartheta) u^{(l')}(\vartheta) u^{(l'')}(\vartheta)}. \end{aligned}$$

In these expressions for  $\mathbb{E}_{g,h}[M_1^3]$  and  $\mathbb{E}_{g,h}[M_2^3]$ , separating out the three cases of  $l = l' = l'' = 0$ , only  $l' = 0$  and  $l = l'' \geq 1$ , and only  $l'' = 0$  and  $l = l' \geq 1$ , this gives

$$\begin{aligned} &\mathbb{E}_{g,h}[\langle \Pi \cdot g \cdot \theta, \Pi \cdot h \cdot \vartheta \rangle^3] \\ &= 8(u^{(0)}(\theta))^3(u^{(0)}(\vartheta))^3 + 6u^{(0)}(\theta)u^{(0)}(\vartheta) \cdot \sum_{l=1}^L |u^{(l)}(\theta)|^2 |u^{(l)}(\vartheta)|^2 \\ &\quad + \frac{3}{8} \sum_{\substack{l,l',l''=1 \\ l=l'+l''}}^L u^{(l)}(\theta) \overline{u^{(l')}(\theta) u^{(l'')}(\theta) u^{(l)}(\vartheta) u^{(l')}(\vartheta) u^{(l'')}(\vartheta)} + \overline{u^{(l)}(\theta) u^{(l')}(\theta) u^{(l'')}(\theta) u^{(l)}(\vartheta) u^{(l')}(\vartheta) u^{(l'')}(\vartheta)} \\ &\quad + \frac{3}{8} \sum_{\substack{l,l',l''=1 \\ l=l'+l''}}^L u^{(l)}(\theta) \overline{u^{(l')}(\theta) u^{(l'')}(\theta) u^{(l)}(\vartheta) u^{(l')}(\vartheta) u^{(l'')}(\vartheta)} + \overline{u^{(l)}(\theta) u^{(l')}(\theta) u^{(l'')}(\theta) u^{(l)}(\vartheta) u^{(l')}(\vartheta) u^{(l'')}(\vartheta)}. \end{aligned}$$

Applying this to (C.21) using  $(x^2 + \bar{x}^2) - 2(xx_* + \bar{x}\bar{x}_*) + (x_*^2 + \bar{x}_*^2) = (x - x_*)^2 + (\bar{x} - \bar{x}_*)^2 = 2 \operatorname{Re}[(x - x_*)^2]$ , and writing as shorthand  $u^{(l)} = u^{(l)}(\theta)$ ,  $u_*^{(l)} = u^{(l)}(\theta_*)$  and similarly for  $r_l, \lambda_l, r_{l,l'}, r_{l,l'}, \lambda_{l,l'}, r_{l,l''}$ ,

$$\begin{aligned} \tilde{s}_3(\theta) &= \frac{2}{3} \left( (u^{(0)})^3 - (u_*^{(0)})^3 \right)^2 + \frac{1}{2} \sum_{l=1}^L \left( u^{(0)} |u^{(l)}|^2 - u_*^{(0)} |u_*^{(l)}|^2 \right)^2 \\ &\quad + \frac{1}{16} \sum_{\substack{l,l',l''=1 \\ l=l'+l''}}^L \left| u^{(l)} \overline{u^{(l')} u^{(l'')}} - u_*^{(l)} \overline{u_*^{(l')} u_*^{(l'')}} \right|^2 + \operatorname{Re} \left[ \left( u^{(l)} \overline{u^{(l')} u^{(l'')}} - u_*^{(l)} \overline{u_*^{(l')} u_*^{(l'')}} \right)^2 \right] \\ &= \frac{2}{3} \left( (\theta^{(0)})^3 - (\theta_*^{(0)})^3 \right)^2 + \frac{1}{2} \sum_{l=1}^L \left( \theta^{(0)} r_l^2 - \theta_*^{(0)} r_{*,l}^2 \right)^2 \\ &\quad + \frac{1}{16} \sum_{\substack{l,l',l''=1 \\ l=l'+l''}}^L r_{l,l'}^2 + r_{*,l,l'}^2 - 2r_{l,l'} r_{*,l,l'} \cos(\lambda_{*,l,l'} - \lambda_{l,l'}) \\ &\quad + r_{l,l'}^2 \cos(2\lambda_{l,l'}) + r_{*,l,l'}^2 \cos(2\lambda_{*,l,l'}) - 2r_{l,l'} r_{*,l,l'} \cos(\lambda_{*,l,l'} + \lambda_{l,l'}). \end{aligned}$$

□

**Lemma C.3.** Let  $H \in \mathbb{R}^{n \times n}$  and  $n = n_1 + n_2$  with  $n_1, n_2 \geq 1$ . Suppose  $H$  can be decomposed as the sum of two positive semidefinite matrices  $A$  and  $B$  with

$$A = \begin{pmatrix} A_{11} & 0 \\ 0 & 0 \end{pmatrix} \text{ and } B = \begin{pmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{pmatrix},$$

where  $A_{11}, B_{11} \in \mathbb{R}^{n_1 \times n_1}$ ,  $B_{12} \in \mathbb{R}^{n_1 \times n_2}$ ,  $B_{21} \in \mathbb{R}^{n_2 \times n_1}$ , and  $B_{22} \in \mathbb{R}^{n_2 \times n_2}$ . Then

$$\operatorname{rank}(H) \geq \operatorname{rank}(A_{11}) + \operatorname{rank}(B_{22}).$$

*Proof.* Since  $A$  and  $B$  are positive semidefinite, so are  $A_{11}, B_{11}, B_{22}$ . There are  $\operatorname{rank}(A_{11})$  linearly independent vectors  $v \in \mathbb{R}^{n_1}$  where  $v^\top A_{11} v > 0$  strictly. Then  $(v, 0)^\top H (v, 0) > 0$  strictly for each such vector  $v$ . There are also  $\operatorname{rank}(B_{22})$  linearly independent vectors  $w \in \mathbb{R}^{n_2}$  where  $w^\top B_{22} w > 0$  strictly. Then  $(0, w)^\top H (0, w) > 0$  strictly for each such vector  $w$ . Thus  $u^\top H u > 0$  for  $\operatorname{rank}(A_{11}) + \operatorname{rank}(B_{22})$  linearly independent vectors  $u \in \mathbb{R}^n$ , so  $\operatorname{rank}(H) \geq \operatorname{rank}(A_{11}) + \operatorname{rank}(B_{22})$ . □

*Proof of Theorem C.1.* As in Theorem 3.1, we have  $\operatorname{trdeg} \mathcal{R}^G = 2L$ . We compute  $\operatorname{trdeg}(\tilde{\mathcal{R}}_{\leq k}^G)$  for  $k = 1, 2, 3$  by applying Lemma 2.8 at a generic point  $\theta_* \in \mathbb{R}^d$  with  $r_l(\theta_*) > 0$  for each  $l = 1, \dots, L$ . Recall from the proof of Theorem 3.1 the map

$$\zeta(\theta) = (\theta_0, r_1, \dots, r_L, t_1, \dots, t_L), \quad \zeta_* = \zeta(\theta_*),$$

with inverse function  $\theta(\zeta)$  in a local neighborhood of  $\theta_*$ . The forms of  $\tilde{s}_1(\theta(\zeta))$  and  $\tilde{s}_2(\theta(\zeta))$  are similar to those of  $s_1(\theta(\zeta))$  and  $s_2(\theta(\zeta))$  in Theorem 3.1, and the same arguments as in the proof of Theorem 3.1 show

$$\begin{aligned} \operatorname{trdeg}(\tilde{\mathcal{R}}_{\leq 1}^G) &= \operatorname{rank} \left( \nabla_\zeta^2 \tilde{s}_1(\theta(\zeta)) \Big|_{\zeta=\zeta_*} \right) = 1, \\ \operatorname{trdeg}(\tilde{\mathcal{R}}_{\leq 2}^G) &= \operatorname{rank} \left( \nabla_\zeta^2 \tilde{s}_1(\theta(\zeta)) + \nabla_\zeta^2 \tilde{s}_2(\theta(\zeta)) \Big|_{\zeta=\zeta_*} \right) = L + 1, \end{aligned}$$

and  $\operatorname{trdeg}(\tilde{\mathcal{R}}_{\leq 3}^G) = \operatorname{rank}(\tilde{H}(\zeta_*))$  for the Hessian

$$\tilde{H}(\zeta_*) = \nabla_\zeta^2 \tilde{s}_1(\theta(\zeta)) + \nabla_\zeta^2 \tilde{s}_2(\theta(\zeta)) + \nabla_\zeta^2 \tilde{s}_3(\theta(\zeta)) \Big|_{\zeta=\zeta_*}.$$

Writing this Hessian in the block decomposition according to  $r = (\theta_0, r_1, \dots, r_L)$  and  $t = (t_1, \dots, t_L)$ , and noting that  $\tilde{s}_1, \tilde{s}_2$  depend only on  $r$  and not on  $t$ , we have the decomposition

$$\tilde{H}(\zeta_*) = \begin{pmatrix} \nabla_r^2 \tilde{s}_1(\theta(\zeta)) + \nabla_r^2 \tilde{s}_2(\theta(\zeta)) \Big|_{\zeta=\zeta_*} & 0 \\ 0 & 0 \end{pmatrix} + \begin{pmatrix} \nabla_r^2 \tilde{s}_3(\theta(\zeta)) \Big|_{\zeta=\zeta_*} & \nabla_{rt}^2 \tilde{s}_3(\theta(\zeta)) \Big|_{\zeta=\zeta_*} \\ \nabla_{tr}^2 \tilde{s}_3(\theta(\zeta)) \Big|_{\zeta=\zeta_*} & \nabla_t^2 \tilde{s}_3(\theta(\zeta)) \Big|_{\zeta=\zeta_*} \end{pmatrix}$$

The second matrix is positive semidefinite by Lemma 2.8, and the first matrix has  $\nabla_r^2 \tilde{s}_1(\theta(\zeta)) + \nabla_r^2 \tilde{s}_2(\theta(\zeta))|_{\zeta=\zeta_*} \succ 0$  strictly by the analysis of  $\text{trdeg}(\tilde{\mathcal{R}}_{\leq 2}^G)$ , with rank exactly  $L + 1$ . Then by Lemma C.3,

$$\text{rank}(\tilde{H}(\zeta_*)) \geq L + 1 + \text{rank}(\nabla_t^2 \tilde{s}_3(\theta(\zeta))|_{\zeta=\zeta_*}). \quad (\text{C.25})$$

As in the proof of Theorem 3.1, let us show  $\text{rank}(\nabla_t^2 \tilde{s}_3(\theta(\zeta))|_{\zeta=\zeta_*}) \geq L - 1$  for generic  $\theta_* \in \mathbb{R}^d$  by exhibiting a single point  $\theta_*$  where this holds.

We may write the expression for  $\tilde{s}_3(\theta)$  in Theorem C.2 as

$$\begin{aligned} \tilde{s}_3(\theta(\zeta)) = f(r) + \frac{1}{16} \sum_{\substack{l, l', l''=1 \\ l=l'+l''}}^L & -2r_{l, l', l''} r_{*, l, l', l''} \cos(\lambda_{l, l', l''} - \lambda_{*, l, l', l''}) + r_{l, l', l''}^2 \cos(2\lambda_{l, l', l''}) \\ & - 2r_{l, l', l''} r_{*, l, l', l''} \cos(\lambda_{l, l', l''} + \lambda_{*, l, l', l''}), \end{aligned}$$

for a function  $f(r)$  depending only on  $r$  and not  $t$ . We pick  $\theta_*$  such that  $\theta_*^{(0)} = 1$  and  $r_{*, l} = 1$  for all  $l = 1, \dots, L$ . Then, recalling  $t_l = \lambda_l - \lambda_{*, l}$  and  $\lambda_{l, l', l''} = \lambda_l - \lambda_{l'} - \lambda_{l''}$ , and differentiating twice in  $t$  at  $(r, t) = (r_*, t_*) = (r_*, 0)$ ,

$$\begin{aligned} \nabla_t^2 \tilde{s}_3(\theta(\zeta))|_{\zeta=\zeta_*} &= \frac{1}{16} \nabla_t^2 \left( \sum_{\substack{l, l', l''=1 \\ l=l'+l''}}^L -2 \cos(t_l - t_{l'} - t_{l''}) + \cos(2t_l - 2t_{l'} - 2t_{l''} + 2\lambda_{*, l} - 2\lambda_{*, l'} - 2\lambda_{*, l''}) \right. \\ &\quad \left. - 2 \cos(t_l - t_{l'} - t_{l''} + 2\lambda_{*, l} - 2\lambda_{*, l'} - 2\lambda_{*, l''}) \right) \Big|_{t=0} \\ &= \frac{1}{16} \sum_{\substack{l, l', l''=1 \\ l=l'+l''}}^L \left( 2 - 4 \cos(2\lambda_{*, l, l', l''}) + 2 \cos(2\lambda_{*, l, l', l''}) \right) \cdot w_{l, l', l''} w_{l, l', l''}^\top \\ &= \frac{1}{8} \sum_{\substack{l, l', l''=1 \\ l=l'+l''}}^L \left( 1 - \cos(2\lambda_{*, l, l', l''}) \right) \cdot w_{l, l', l''} w_{l, l', l''}^\top, \end{aligned}$$

where  $w_{l, l', l''}$  is defined as (C.8). Stacking  $w_{l, l', l''}$  as the columns of  $W \in \mathbb{R}^{L \times |\mathcal{L}|}$  as in the proof of Theorem 3.1, and defining the diagonal matrix  $D = \text{diag}(1 - \cos(2\lambda_{*, l, l', l''})) \in \mathbb{R}^{|\mathcal{L}| \times |\mathcal{L}|}$ , this shows

$$\nabla_t^2 \tilde{s}_3(\theta(\zeta))|_{\zeta=\zeta_*} = \frac{1}{8} W D W^\top.$$

Note that, for generic  $\lambda_* = (\lambda_{*, 1}, \dots, \lambda_{*, L})$ , we have

$$2 - 2 \cos(2\lambda_{*, l} - 2\lambda_{*, l'} - 2\lambda_{*, l''}) > 0$$

for each fixed tuple  $(l, l', l'') \in \mathcal{L}$ . Hence we may pick  $\lambda_*$  so that this holds simultaneously for all tuples  $(l, l', l'') \in \mathcal{L}$ . Then  $\text{rank}(W D W^\top) = \text{rank}(W W^\top) \geq L - 1$  as shown in Theorem 3.1. Applying this back to (C.25), we have shown  $\text{trdeg} \tilde{\mathcal{R}}_{\leq 3}^G = \text{rank}(\tilde{H}(\zeta_*)) \geq 2L$ . Since also  $\text{trdeg} \tilde{\mathcal{R}}_{\leq 3}^G \leq \text{trdeg} \mathcal{R}^G = 2L$ , this shows  $\text{trdeg} \tilde{\mathcal{R}}_{\leq 3}^G = 2L$ .  $\square$

#### APPENDIX D. ANALYSES OF FUNCTION ESTIMATION UNDER AN $\text{SO}(3)$ ROTATION

This appendix contains further details on the setups of the models and the proofs of the main results in Section 4 on estimating a function in 2 or 3 dimensions under  $\text{SO}(3)$ -rotations.

Appendix D.1 first reviews the complex spherical harmonics basis and the associated calculus of Wigner D-matrices and Clebsch-Gordan coefficients. Appendix D.2 contains further details and proofs for Section 4.1 on spherical registration. Appendix D.3 contains further details and proofs for Section 4.2 on the unprojected cryo-EM model. Finally, Appendix D.3 contains further details and proofs for Section 4.3 on the projected cryo-EM model.

**D.1. Calculus of spherical harmonics.** We first fix notations for some special functions related to the action of  $\text{SO}(3)$  and present some identities between them which will appear in the proofs.

D.1.1. *Complex spherical harmonics.* Let  $P_{lm}(x)$  denote the associated Legendre polynomials (without Cordon-Shortley phase)

$$P_{lm}(x) = \frac{1}{2^l l!} (1-x^2)^{m/2} \frac{d^{l+m}}{dx^{l+m}} (x^2-1)^l \quad \text{for } m = -l, -l+1, \dots, l-1, l. \quad (\text{D.1})$$

Let  $\mathcal{S}^2 \subset \mathbb{R}^3$  be the unit sphere, parametrized by the latitude  $\phi_1 \in [0, \pi]$  and longitude  $\phi_2 \in [0, 2\pi)$ . The complex spherical harmonics basis on  $\mathcal{S}^2$  is given by (see [Ros95, Eq. (III.20)])

$$y_{lm}(\phi_1, \phi_2) = (-1)^m \sqrt{\frac{2l+1}{4\pi} \cdot \frac{(l-m)!}{(l+m)!}} \cdot P_{lm}(\cos \phi_1) e^{im\phi_2} \quad \text{for } l \geq 0 \text{ and } m = -l, \dots, l. \quad (\text{D.2})$$

(We will use interchangeably notations such as  $y_{lm}$  and  $y_{l,m}$  when the meaning is clear.) The index  $l$  is the frequency, and there are  $2l+1$  basis functions at each frequency  $l$ . These functions are orthonormal in  $L_2(\mathcal{S}^2, \mathbb{C})$  with respect to the surface area measure  $\sin \phi_1 d\phi_1 d\phi_2$ , and satisfy the conjugation symmetry (see [Ros95, Eq. (III.23)])

$$\overline{y_{l,m}(\phi_1, \phi_2)} = (-1)^m y_{l,-m}(\phi_1, \phi_2). \quad (\text{D.3})$$

**Lemma D.1.** *For all  $m$ , the associated Legendre polynomials in (D.1) satisfy*

$$P_{lm}(0) = \mathbf{1}\{l+m \text{ is even}\} \cdot \frac{(-1)^{(l-m)/2}}{2^l l!} \binom{l}{(l+m)/2} (l+m)!$$

*Proof.* This follows from applying a binomial expansion of  $(x^2-1)^l$ , and then differentiating in  $x$ —see also [BBSK<sup>+</sup>17, Eq. (14)].  $\square$

D.1.2. *Wigner D-matrices.* Let  $f \in L_2(\mathcal{S}^2, \mathbb{C})$ . Then  $f$  may be decomposed in the complex spherical harmonics basis (D.2) as

$$f = \sum_{l=0}^{\infty} \sum_{m=-l}^l u_m^{(l)} y_{lm}.$$

Writing  $u^{(l)} = \{u_m^{(l)} : -l \leq m \leq l\}$ , the rotation  $f \mapsto f_{\mathbf{g}}$  given by  $f_{\mathbf{g}}(\phi_1, \phi_2) = f(\mathbf{g}^{-1} \cdot (\phi_1, \phi_2))$  for  $\mathbf{g} \in \text{SO}(3)$  is described by the map of spherical harmonic coefficients (see [Ros95, Eq. (4.28a)])

$$u^{(l)} \mapsto D^{(l)}(\mathbf{g}) u^{(l)} \quad \text{for each } l = 0, 1, 2, \dots,$$

where  $D^{(l)}(\mathbf{g}) \in \mathbb{C}^{(2l+1) \times (2l+1)}$  is the *complex Wigner D-matrix* at frequency  $l$  corresponding to  $\mathbf{g}$ . We index the rows and columns of  $D^{(l)}(\mathbf{g})$  by  $-l, \dots, l$ .

Our computations will not require the explicit forms of  $D^{(l)}(\mathbf{g})$ , but only the following moment identities when  $\mathbf{g} \in \text{SO}(3)$  is a Haar-uniform random rotation (see [Ros95, Section 16] and [BBSK<sup>+</sup>17, Appendix A.3]):

(1) Mean identity:

$$D^{(0)}(\mathbf{g}) = 1, \quad \mathbb{E}_{\mathbf{g}}[D^{(l)}(\mathbf{g})] = 0 \text{ for all } l \geq 1. \quad (\text{D.4})$$

(2) Orthogonality: for any  $l, l' \geq 0$  and  $-l \leq q, m \leq l$  and  $-l' \leq q', m' \leq l'$ ,

$$\mathbb{E}_{\mathbf{g}} \left[ D_{qm}^{(l)}(\mathbf{g}) D_{q'm'}^{(l')}(\mathbf{g}) \right] = \frac{(-1)^{m+q}}{2l+1} \mathbf{1}\{l = l', q = -q', m = -m'\}. \quad (\text{D.5})$$

(3) Third order identity: for any  $l, l', l'' \geq 0$  and  $-l \leq q, m \leq l$  and  $-l' \leq q', m' \leq l'$  and  $-l'' \leq q'', m'' \leq l''$ ,

$$\begin{aligned} \mathbb{E}_{\mathbf{g}} \left[ D_{qm}^{(l)}(\mathbf{g}) D_{q'm'}^{(l')}(\mathbf{g}) D_{q''m''}^{(l'')}(\mathbf{g}) \right] &= \mathbf{1}\{q + q' = -q''\} \cdot \mathbf{1}\{m + m' = -m''\} \cdot \mathbf{1}\{|l - l'| \leq l'' \leq l + l'\} \\ &\quad \cdot \frac{(-1)^{m''+q''}}{2l''+1} \langle l, q; l', q' | l'', -q'' \rangle \langle l, m; l', m' | l'', -m'' \rangle, \end{aligned} \quad (\text{D.6})$$

where  $\langle l, m; l', m' | l'', m'' \rangle$  is a Clebsch-Gordan coefficient, defined in the following section.

D.1.3. *Clebsch-Gordan coefficients.* The Clebsch-Gordan coefficients  $\langle l, m; l', m' | l'', m'' \rangle$  are defined for integer arguments  $l, l', l'', m, m', m''$  where

$$|m| \leq l, \quad |m'| \leq l', \quad |m''| \leq l'' \quad \text{and} \quad |l - l'| \leq l'' \leq l + l'. \quad (\text{D.7})$$

The latter condition  $|l - l'| \leq l'' \leq l + l'$  is equivalent to the three symmetric triangle inequality conditions  $l + l' \geq l''$ ,  $l + l'' \geq l'$ , and  $l' + l'' \geq l$ . For such arguments,  $\langle l, m; l', m' | l'', m'' \rangle$  is given explicitly by (see [Böh13, Eq. (2.41)] and [BBSK<sup>+</sup>17, Appendix A.2])

$$\begin{aligned} & \langle l, m; l', m' | l'', m'' \rangle \\ &= \mathbf{1}\{m'' = m + m'\} \times \sqrt{\frac{(2l'' + 1)(l + l' - l'')!(l + l'' - l')!(l' + l'' - l)!}{(l + l' + l'' + 1)!}} \\ & \quad \times \sqrt{(l - m)!(l + m)!(l' - m')!(l' + m')!(l'' - m'')!(l'' + m'')!} \\ & \quad \times \sum_k \frac{(-1)^k}{k!(l + l' - l'' - k)!(l - m - k)!(l' + m' - k)!(l'' - l' + m + k)!(l'' - l - m' + k)!} \end{aligned} \quad (\text{D.8})$$

where the summation is over all integers  $k$  for which the argument of every factorial is nonnegative. We extend the definition to all integer arguments by

$$\langle l, m; l', m' | l'', m' \rangle = 0 \quad \text{if (D.7) does not hold.} \quad (\text{D.9})$$

We will use the notational shorthand

$$C_{m, m', m''}^{l, l', l''} = \langle l, m; l', m' | l'', m'' \rangle.$$

These coefficients satisfy the sign symmetry (see [Ros95, Eq. (3.16a)] and [Böh13, Eq. (2.47)])

$$C_{m, m', m''}^{l, l', l''} = (-1)^{l+l'+l''} C_{-m, -m', -m''}^{l, l', l''}. \quad (\text{D.10})$$

Note that we may have  $C_{m, m', m''}^{l, l', l''} = 0$  even if (D.7) holds and  $m'' = m + m'$ . For example,  $C_{2, -1, 1}^{3, 2, 2} = 0$ . In our later proofs, we will require that certain Clebsch-Gordan coefficients are *non-zero*, and the following lemma provides a sufficient condition for this to hold.

**Lemma D.2.** *Let  $l, l', l'', m, m', m''$  satisfy (D.7), where  $m'' = m + m'$ . In addition, suppose the following conditions all hold:*

- $l \geq l'$  and  $l \geq l'' + 1$ .
- $|m'| \in \{l' - 1, l'\}$  and  $|m''| \in \{l'' - 1, l''\}$ .
- We do not have simultaneously  $|m| = l - 1$ ,  $|m'| = l' - 1$ ,  $|m''| = l'' - 1$ , and  $l' = l''$ .

Then

$$C_{m, m', m''}^{l, l', l''} \neq 0.$$

*Proof.* Applying the sign symmetry (D.10), we may assume  $m' \in \{-l', -l' + 1\}$ . We consider separately these cases.

**Case I:**  $m' = -l'$ . Then  $m = m'' - m' = m'' + l'$ . The condition that  $k$  and  $l' + m' - k$  are both nonnegative in (D.8) requires  $k = 0$ , so the sum in (D.8) consists of just this single term. The remaining factorials in (D.8) are also nonnegative, because  $l + l' - l'' \geq 0$ ,  $l - m \geq 0$ ,  $l'' - l' + m = l'' + m'' \geq 0$ , and  $l'' - l - m' = l'' + l' - l \geq 0$ . Thus (D.8) is non-zero.

**Case II:**  $m' = -l' + 1$ . Then  $m = m'' + l' - 1$ . The condition that  $k$  and  $l' + m' - k$  are both nonnegative then requires  $k \in \{0, 1\}$ . Substituting  $m' = -l' + 1$  and  $m = m'' + l' - 1$ , these two terms in (D.8) for  $k \in \{0, 1\}$  are

$$\frac{1}{(l+l'-l'')!(l-m)!(l''+m''-1)!(l'+l''-l-1)!} - \frac{1}{(l+l'-l''-1)!(l-m-1)!(l''+m'')!(l'+l''-l)!}$$

where each term is understood as 0 if an argument to one of its factorials is negative. Here  $l + l' - l'' \geq l + l' - l'' - 1 \geq 0$  always, because  $l \geq l'' + 1$ .

- If  $m = l$ , then the second term is 0. Also  $l'' + m'' - 1 = l'' + (m + m') - 1 = l'' + l - l' \geq 0$  and  $l' + l'' - l - 1 = (m - m'' + 1) + l'' - l - 1 = l'' - m'' \geq 0$ , so the first term is non-zero.
- If  $m < l$  but  $m'' = -l''$  or  $l = l' + l''$ , then the first term is 0, but the second term is non-zero.

It remains to consider  $m < l$ ,  $m'' \in \{-l'' + 1, l'' - 1, l''\}$ , and  $l < l' + l''$ . Then both terms are non-zero, and their sum is

$$\frac{1}{(l+l'-l'')!(l-m)!(l''+m'')!(l'+l''-l)!} ((l'' + m'')(l' + l'' - l) - (l + l' - l'')(l - m)).$$

We now consider the three cases of  $m''$ :

- $m'' = l''$  is not possible, because this would imply  $m = l' + l'' - 1 \geq l$ , contradicting  $m < l$ .
- If  $m'' = l'' - 1$ , then  $m = l' + l'' - 2 \geq l - 1$ . Hence we must have the equalities  $m = l - 1$  and  $l' + l'' - 1 = l$ . Then

$$(l'' + m'')(l' + l'' - l) - (l + l' - l'')(l - m) = (2l'' - 1) - (2l' - 1),$$

which is non-zero because our third given condition implies  $l' \neq l''$  when  $|m| = l - 1$ ,  $|m'| = l' - 1$ , and  $|m''| = l'' - 1$ .

- If  $m'' = -l'' + 1$ , then

$$(l'' + m'')(l' + l'' - l) - (l + l' - l'')(l - m) = (l' + l'' - l) - (l + l' - l'')(l - m).$$

This is non-zero because  $l - m \geq 1$  and  $l + l' - l'' > l' + l'' - l$  strictly.

Thus we obtain that (D.8) is non-zero in all cases.  $\square$

## D.2. Spherical registration.

D.2.1. *Function basis.* We review the real spherical harmonics basis that we use for this example, and the action of  $\text{SO}(3)$  on the coefficients in this basis that is induced by rotation of the function domain  $\mathcal{S}^2$ . The setup is the same as that of the spherical registration model discussed in [BBSK<sup>+</sup>17, Sections 5.4 and A.1].

We define the real spherical harmonics basis  $\{h_{lm} : l \geq 0, m \in \{-l, \dots, l\}\}$  from the complex spherical harmonics basis (D.2) by

$$h_{l,m} = \begin{cases} \frac{1}{\sqrt{2}}(y_{l,-m} + (-1)^m y_{l,m}) & \text{if } m > 0 \\ y_{l,0} & \text{if } m = 0 \\ \frac{i}{\sqrt{2}}(y_{l,m} - (-1)^m y_{l,-m}) & \text{if } m < 0. \end{cases}$$

It may be checked from (D.3) that these functions  $\{h_{lm}\}$  are real-valued and form an orthonormal basis for  $L_2(\mathcal{S}^2, \mathbb{R})$ . For any function  $f \in L_2(\mathcal{S}^2, \mathbb{C})$ , writing its orthogonal decompositions in the bases  $\{y_{lm}\}$  and  $\{h_{lm}\}$  as

$$f = \sum_{l=0}^{\infty} \sum_{m=-l}^l u_m^{(l)} y_{lm} = \sum_{l=0}^{\infty} \sum_{m=-l}^l \theta_m^{(l)} h_{lm},$$

its real and complex spherical harmonic coefficients  $\{u_m^{(l)}\}$  and  $\{\theta_m^{(l)}\}$  are then related by

$$u_m^{(l)} = \begin{cases} \frac{(-1)^m}{\sqrt{2}}(\theta_{|m|}^{(l)} - i\theta_{-|m|}^{(l)}) & \text{if } m > 0 \\ \theta_0^{(l)} & \text{if } m = 0 \\ \frac{1}{\sqrt{2}}(\theta_{|m|}^{(l)} + i\theta_{-|m|}^{(l)}) & \text{if } m < 0. \end{cases} \quad (\text{D.11})$$

Up to the finite bandlimit  $L \geq 1$ , this relation (D.11) is a linear map  $u = V^* \theta$  between  $u \in \mathbb{C}^d$  and  $\theta \in \mathbb{C}^d$ , where  $V \in \mathbb{C}^{d \times d}$  is a unitary matrix. If  $f$  is real-valued, then  $\{\theta_m^{(l)}\}$  are real, and hence  $\{u_m^{(l)}\}$  satisfy the sign symmetry

$$u_m^{(l)} = (-1)^m \overline{u_{-m}^{(l)}}. \quad (\text{D.12})$$

The space of bandlimited functions (4.1) is closed under the action of  $\text{SO}(3)$ , and the rotation  $f \mapsto f_{\mathbf{g}}$  is represented by the following subgroup  $G \subset \text{O}(d)$  acting on  $\theta \in \mathbb{R}^d$ .

**Lemma D.3.** *The action of  $\text{SO}(3)$  on the real spherical harmonic coefficients  $\theta \in \mathbb{R}^d$  admits the representation*

$$G = \left\{ V \cdot D(\mathbf{g}) \cdot V^* : \mathbf{g} \in \text{SO}(3) \right\} \subset \text{O}(d),$$

where  $V \in \mathbb{C}^{d \times d}$  is the unitary transform describing the map  $u = V^* \theta$  in (D.11), and  $D(\mathfrak{g})$  is the block-diagonal matrix

$$D(\mathfrak{g}) = \bigoplus_{l=0}^L D^{(l)}(\mathfrak{g}) \in \mathbb{C}^{d \times d} \quad (\text{D.13})$$

with diagonal blocks  $D^{(l)}(\mathfrak{g}) \in \mathbb{C}^{(2l+1) \times (2l+1)}$  given by the complex Wigner D-matrices at frequencies  $l = 0, \dots, L$  (defined in Appendix D.1).

*Proof.* As described in Appendix D.1.2, the rotation by  $\mathfrak{g} \in \text{SO}(3)$  acts on the complex spherical harmonic coefficients  $u \in \mathbb{C}^d$  by

$$u \mapsto D(\mathfrak{g})u,$$

where  $D(\mathfrak{g})$  is the block-diagonal matrix defined in (D.13). Since  $u$  and  $\theta$  are related by the unitary transformations  $u = V^* \theta$  and  $\theta = V u$ , the action on  $\theta$  is then given by  $\theta \mapsto V \cdot D(\mathfrak{g}) \cdot V^* \theta$ .  $\square$

**D.2.2. Terms of the high-noise series expansion.** We prove Theorem 4.4 on the forms of  $s_1(\theta)$ ,  $s_2(\theta)$ , and  $s_3(\theta)$ .

*Proof of Theorem 4.4.* Recall by Lemma A.1 that

$$s_k(\theta) = \frac{1}{2(k!)} \mathbb{E}_g [\langle \theta, g \cdot \theta \rangle^k - 2\langle \theta, g \cdot \theta_* \rangle^k + \langle \theta_*, g \cdot \theta_* \rangle^k]. \quad (\text{D.14})$$

Consider two different real spherical harmonic coefficient vectors  $\theta, \vartheta \in \mathbb{R}^d$ , and the corresponding complex coefficients  $u = V^* \theta \in \mathbb{C}^d$  and  $v = V^* \vartheta \in \mathbb{C}^d$ . We compute  $\mathbb{E}_g[\langle \theta, g \cdot \vartheta \rangle^k]$  for  $k = 1, 2, 3$ .

**Case  $k = 1$ :** By Lemma D.3, for any  $g \in \mathbf{G}$ ,

$$\langle \theta, g \cdot \vartheta \rangle = \langle \theta, V D(\mathfrak{g}) V^* \vartheta \rangle = \langle u, D(\mathfrak{g}) v \rangle.$$

From the block-diagonal form for  $D(\mathfrak{g})$  in (D.13), we obtain

$$\langle \theta, g \cdot \vartheta \rangle = \sum_{l=0}^L \langle u^{(l)}, D^{(l)}(\mathfrak{g}) v^{(l)} \rangle = \sum_{l=0}^L \sum_{q, m=-l}^l \overline{u_q^{(l)}} D_{qm}^{(l)}(\mathfrak{g}) v_m^{(l)}. \quad (\text{D.15})$$

Applying the identities (D.4) yields

$$\mathbb{E}_g[\langle \theta, g \cdot \vartheta \rangle] = \overline{u_0^{(0)}} v_0^{(0)}.$$

Write the shorthands  $u^{(0)} = u_0^{(0)}$  and  $v^{(0)} = v_0^{(0)}$ , and recall from (D.12) that  $u^{(0)}, v^{(0)}$  are real. Then applying this to (D.14),

$$s_1(\theta) = \frac{1}{2} \mathbb{E}_g[\langle \theta, g \cdot \theta \rangle] - \mathbb{E}_g[\langle \theta_*, g \cdot \theta \rangle] + \frac{1}{2} \mathbb{E}_g[\langle \theta_*, g \cdot \theta_* \rangle] = \frac{1}{2} \left( u^{(0)}(\theta) - u^{(0)}(\theta_*) \right)^2.$$

**Case  $k = 2$ :** We take the expected square on both sides of (D.15), applying (D.12) and the relation (D.5). Then

$$\begin{aligned} \mathbb{E}_g[\langle \theta, g \cdot \vartheta \rangle^2] &= \sum_{l=0}^L \sum_{q, m=-l}^l \frac{(-1)^{m+q}}{2l+1} \overline{u_q^{(l)}} u_{-q}^{(l)} v_m^{(l)} v_{-m}^{(l)} \\ &= \sum_{l=0}^L \sum_{q, m=-l}^l \frac{1}{2l+1} \overline{u_q^{(l)}} u_q^{(l)} v_m^{(l)} \overline{v_m^{(l)}} = \sum_{l=0}^L \frac{1}{2l+1} \|u^{(l)}\|^2 \cdot \|v^{(l)}\|^2. \end{aligned}$$

Applying this to (D.14),

$$\begin{aligned} s_2(\theta) &= \frac{1}{4} \mathbb{E}_g[\langle \theta, g \cdot \theta \rangle^2] - \frac{1}{2} \mathbb{E}_g[\langle \theta_*, g \cdot \theta \rangle^2] + \frac{1}{4} \mathbb{E}_g[\langle \theta_*, g \cdot \theta_* \rangle^2] \\ &= \sum_{l=0}^L \frac{1}{4(2l+1)} \left( \|u^{(l)}(\theta)\|^2 - \|u^{(l)}(\theta_*)\|^2 \right)^2. \end{aligned}$$

**Case  $k = 3$ :** We now take the expected cube on both sides of (D.15), applying the relation (D.6). Recall the convention (D.9). Then

$$\mathbb{E}_g[\langle \theta, g \cdot \vartheta \rangle^3]$$

$$\begin{aligned}
&= \sum_{\substack{l, l', l''=0 \\ |l-l'|\leq l''\leq l+l'}}^L \sum_{q, m=-l}^l \sum_{q', m'=-l'}^{l'} \frac{(-1)^{m+m'+q+q'}}{2l''+1} \cdot C_{q, q', q+q'}^{l, l', l''} C_{m, m', m+m'}^{l, l', l''} \overline{u_q^{(l)} u_{q'}^{(l')} u_{-q-q'}^{(l'')}} v_m^{(l)} v_{m'}^{(l')} v_{-m-m'}^{(l'')} \\
&= \sum_{\substack{l, l', l''=0 \\ |l-l'|\leq l''\leq l+l'}}^L \sum_{q, m=-l}^l \sum_{q', m'=-l'}^{l'} \frac{1}{2l''+1} \cdot C_{q, q', q+q'}^{l, l', l''} C_{m, m', m+m'}^{l, l', l''} \overline{u_q^{(l)} u_{q'}^{(l')} u_{q+q'}^{(l'')}} v_m^{(l)} v_{m'}^{(l')} \overline{v_{m+m'}^{(l'')}}.
\end{aligned}$$

Recall that

$$B_{l, l', l''}(\theta) = \sum_{m=-l}^l \sum_{m'=-l'}^{l'} C_{m, m', m+m'}^{l, l', l''} \overline{u_m^{(l)} u_{m'}^{(l')} u_{m+m'}^{(l'')}}, \quad u = V^* \theta.$$

Then the above may be written as

$$\mathbb{E}_g[\langle \theta, g \cdot \vartheta \rangle^3] = \sum_{\substack{l, l', l''=0 \\ |l-l'|\leq l''\leq l+l'}}^L \frac{1}{2l''+1} B_{l, l', l''}(\theta) \overline{B_{l, l', l''}(\vartheta)}.$$

Changing indices  $(m, m') \mapsto (-m, -m')$  and applying the symmetries (D.10) and (D.12), we have

$$\begin{aligned}
B_{l, l', l''}(\theta) &= \sum_{m=-l}^l \sum_{m'=-l'}^{l'} C_{-m, -m', -m-m'}^{l, l', l''} \overline{u_{-m}^{(l)} u_{-m'}^{(l')} u_{-m-m'}^{(l'')}} \\
&= \sum_{m=-l}^l \sum_{m'=-l'}^{l'} (-1)^{l+l'+l''} C_{m, m', m+m'}^{l, l', l''} \cdot (-1)^{m+m'+(m+m')} u_m^{(l)} u_{m'}^{(l')} \overline{u_{m+m'}^{(l'')}} \\
&= (-1)^{l+l'+l''} \overline{B_{l, l', l''}(\theta)}.
\end{aligned}$$

Thus  $B_{l, l', l''}(\theta)$  is real-valued if  $l + l' + l''$  is even and pure imaginary if  $l + l' + l''$  is odd. Applying this to (D.14),

$$\begin{aligned}
s_3(\theta) &= \frac{1}{12} \mathbb{E}_g[\langle \theta, g \cdot \theta \rangle^3] - \frac{1}{6} \mathbb{E}_g[\langle \theta_*, g \cdot \theta \rangle^3] + \frac{1}{12} \mathbb{E}_g[\langle \theta_*, g \cdot \theta_* \rangle^3] \\
&= \frac{1}{12} \sum_{\substack{l, l', l''=0 \\ |l-l'|\leq l''\leq l+l'}}^L \frac{1}{2l''+1} (B_{l, l', l''}(\theta) - B_{l, l', l''}(\theta_*)) \left( \overline{B_{l, l', l''}(\theta)} - \overline{B_{l, l', l''}(\theta_*)} \right) \\
&= \frac{1}{12} \sum_{\substack{l, l', l''=0 \\ |l-l'|\leq l''\leq l+l'}}^L \frac{1}{2l''+1} \left| B_{l, l', l''}(\theta) - B_{l, l', l''}(\theta_*) \right|^2.
\end{aligned}$$

□

**D.2.3. Transcendence degrees.** We now prove Theorem 4.1 on the sequences of transcendence degrees.

*Proof of Theorem 4.1.* Recall the form of  $\mathbf{G}$  in Lemma D.3. Denote the diagonal blocks of  $g \in \mathbf{G}$  by  $g^{(l)}(\mathbf{g}) = V^{(l)} \cdot D^{(l)}(\mathbf{g}) \cdot V^{(l)*}$ , where each  $V^{(l)} \in \mathbb{C}^{(2l+1) \times (2l+1)}$  represents the unitary map  $u^{(l)} \mapsto \theta^{(l)}$ , and  $g^{(l)}$  is the irreducible representation of  $\mathrm{SO}(3)$  acting on the subvector  $\theta^{(l)} \in \mathbb{R}^{2l+1}$ . For  $l = 3$ , it is known that any generic point  $\theta^{(3)} \in \mathbb{R}^7$  has a trivial stabilizer subgroup  $\{\mathrm{Id}\}$  and a 3-dimensional orbit under this action [Bry16, Proposition 1]. Defining the pre-image  $\mathbf{H} := \{\mathbf{g} \in \mathrm{SO}(3) : g^{(3)}(\mathbf{g}) = \mathrm{Id}\}$ , for  $L \geq 3$  and any extension of  $\theta^{(3)}$  to  $\theta \in \mathbb{R}^d$ , the stabilizer of  $\theta$  in  $\mathbf{G}$  must then satisfy  $\mathbf{G}_\theta \subseteq \{V \cdot D(\mathbf{g}) \cdot V^* : \mathbf{g} \in \mathbf{H}\}$ . Here  $\mathbf{H}$  is a discrete subgroup of  $\mathrm{SO}(3)$ , so  $\mathbf{G}_\theta$  is a discrete subgroup of  $\mathbf{G}$ , and hence  $\dim(\mathbf{G}_\theta) = 0$  and  $d_0 = \max_\theta \dim(\mathcal{O}_\theta) = \dim(\mathbf{G}) - \min_\theta \dim(\mathbf{G}_\theta) = 3$ .

We compute  $\mathrm{trdeg}(\mathcal{R}_{\leq k}^{\mathbf{G}})$  for  $k = 1, 2, 3$  using Lemma 2.8. Recall the forms of  $s_1(\theta)$  and  $s_2(\theta)$  in Theorem 4.4, where  $u^{(0)}(\theta) = \theta_0^{(0)}$  and  $\|u^{(l)}(\theta)\|^2 = \|\theta^{(l)}\|^2$  for  $\theta^{(l)} = (\theta_m^{(l)} : m = -l, \dots, l)$ . Then we obtain directly that for generic  $\theta_* \in \mathbb{R}^d$ ,

$$\begin{aligned}
\mathrm{trdeg}(\mathcal{R}_{\leq 1}^{\mathbf{G}}) &= \mathrm{rank}(\nabla^2 s_1(\theta_*)) = 1 \\
\mathrm{trdeg}(\mathcal{R}_{\leq 2}^{\mathbf{G}}) &= \mathrm{rank}(\nabla^2 s_1(\theta_*) + \nabla^2 s_2(\theta_*)) = L + 1.
\end{aligned}$$

It remains to show  $\text{trdeg}(\mathcal{R}_{\leq 3}^G) = d - 3$ . Note that  $\text{trdeg}(\mathcal{R}_{\leq 3}^G) \leq \text{trdeg}(\mathcal{R}^G) = d - 3$ , so it suffices to show the lower bound  $\text{trdeg}(\mathcal{R}_{\leq 3}^G) \geq d - 3$ .

By Lemma 2.8 and the fact that each Hessian  $\nabla^2 s_k(\theta_*)$  is positive semidefinite,

$$\text{trdeg}(\mathcal{R}_{\leq 3}^G) \geq \text{rank}(\nabla^2 s_3(\theta_*)).$$

Writing the index set

$$\mathcal{J} = \{(l, l', l'') : 0 \leq l, l', l'' \leq L, |l - l'| \leq l'' \leq l + l'\}, \quad (\text{D.16})$$

we have by Theorem 4.4

$$s_3(\theta) = \frac{1}{12} \sum_{(l, l', l'') \in \mathcal{J}} \frac{1}{2l'' + 1} (B_{l, l', l''}(\theta) - B_{l, l', l''}(\theta_*)) (\overline{B_{l, l', l''}(\theta)} - \overline{B_{l, l', l''}(\theta_*)}).$$

Let us denote

$$B(\theta) = (B_{l, l', l''}(\theta) : (l, l', l'') \in \mathcal{J}), \quad B : \mathbb{R}^d \rightarrow \mathbb{C}^{|\mathcal{J}|},$$

and write  $\text{dB}(\theta) \in \mathbb{C}^{|\mathcal{J}| \times d}$  for its the derivative in  $\theta$ . Then, applying the chain rule to differentiate  $s_3(\theta)$  twice at  $\theta = \theta_*$ , we obtain

$$\nabla^2 s_3(\theta_*) = \text{dB}(\theta_*)^\top \cdot \text{diag} \left( \frac{1}{6(2l'' + 1)} : (l, l', l'') \in \mathcal{J} \right) \cdot \overline{\text{dB}(\theta_*)}.$$

The diagonal matrix in the middle has full rank, so

$$\text{rank}(\nabla^2 s_3(\theta_*)) = \text{rank}(\text{dB}(\theta_*)). \quad (\text{D.17})$$

To analyze this rank, recall the complex parametrization  $u = V^* \theta \in \mathbb{C}^d$  from (D.11), satisfying the symmetry (D.12). Let us write the real and imaginary parts of  $u$  as

$$u_m^{(l)} = v_m^{(l)} + \mathbf{i} w_m^{(l)}$$

so that this symmetry (D.12) is equivalent to

$$v_{-m}^{(l)} = (-1)^m v_m^{(l)}, \quad w_{-m}^{(l)} = (-1)^{m+1} w_m^{(l)}. \quad (\text{D.18})$$

Note that for  $m = 0$ , this implies  $w_0^{(l)} = 0$ . Then, setting

$$\eta^{(l)}(\theta) = (v_0^{(l)}, v_1^{(l)}, w_1^{(l)}, v_2^{(l)}, w_2^{(l)}, \dots, v_l^{(l)}, w_l^{(l)}),$$

these coordinates  $\eta^{(l)} \in \mathbb{R}^{2l+1}$  provide a (linear) invertible reparametrization of  $\theta^{(l)}$ . This defines a reparametrization

$$\eta(\theta) = (\eta^{(l)}(\theta) : 0 \leq l \leq L) \in \mathbb{R}^d$$

with inverse function  $\theta(\eta)$ . Writing as shorthand  $\eta_* = \eta(\theta_*)$  and  $B(\eta) \equiv B(\theta(\eta))$ , and denoting by  $\text{d}_\eta B(\eta)$  the derivative of  $B$  in the new variables  $\eta$ , (D.17) is equivalent to

$$\text{rank}(\nabla^2 s_3(\theta_*)) = \text{rank}(\text{d}_\eta B(\eta_*)).$$

Denote

$$\tilde{B}(\eta) = (B_{l, l', l''}(\eta) : (l, l', l'') \in \mathcal{J}, \max(l, l', l'') \leq 10).$$

Let us group the columns and rows of  $\text{d}_\eta B$  into blocks indexed by  $\{\sim, 11, 12, \dots, L\}$  as follows: The column block  $\sim$  corresponds to  $\text{d}_{\eta^{(0)}, \dots, \eta^{(10)}}$ . The row block  $\sim$  corresponds to  $\tilde{B}(\eta)$ . For  $l \geq 11$ , the column block  $l$  corresponds to  $\text{d}_{\eta^{(l)}}$ , and the row block  $l$  corresponds to  $B^{(l)}$  as defined below in Lemma D.4. (These blocks  $\tilde{B}$  and  $B^{(l)}$  for  $l \geq 11$  are disjoint by definition, and we may discard the remaining rows of  $\text{d}_\eta B$  not corresponding to any such block to produce a lower bound for its rank.) Ordering the blocks by  $\sim, 11, 12, \dots, L$ , the resulting matrix  $\text{d}_\eta B$  is block lower-triangular, because each  $B^{(l)}$  does not depend on the variables  $\eta^{(l+1)}, \dots, \eta^{(L)}$ . Thus  $\text{rank}(\text{d}_\eta B)$  is lower-bounded by the sum of ranks of all diagonal blocks, i.e.

$$\text{rank}(\text{d}_\eta B(\eta_*)) \geq \text{rank}(\text{d}_{\eta^{(0)}, \dots, \eta^{(10)}} \tilde{B}(\eta_*)) + \sum_{l=11}^L \text{rank}(\text{d}_{\eta^{(l)}} B^{(l)}(\eta_*)).$$

A direct numerical evaluation of the matrix  $d_{\eta^{(0)}, \dots, \eta^{(10)}} \tilde{B}(\eta_*)$  verifies that for  $\eta_* \in \mathbb{R}^d$  with all entries of  $\eta_*^{(0)}, \dots, \eta_*^{(10)}$  equal to 1, we have<sup>1</sup>

$$\text{rank} \left( d_{\eta^{(0)}, \dots, \eta^{(10)}} \tilde{B}(\eta_*) \right) = \sum_{l=0}^{10} (2l+1) - 3 = 118.$$

Then also for generic  $\eta_* \in \mathbb{R}^d$ , by Fact 4.5,

$$\text{rank} \left( d_{\eta^{(0)}, \dots, \eta^{(10)}} \tilde{B}(\eta_*) \right) \geq \sum_{l=0}^{10} (2l+1) - 3 = 118.$$

In particular, this establishes the desired result that  $\text{rank}(\nabla^2 s_3(\theta_*)) = \text{rank}(d_\eta B(\eta_*)) \geq d-3$  for  $L=10$ . If  $L \geq 11$ , then by Lemma D.4 below, we also have for generic  $\eta_* \in \mathbb{R}^d$ ,

$$\sum_{l=11}^L \text{rank}(d_{\eta^{(l)}} B^{(l)}(\eta_*)) = \sum_{l=11}^L (2l+1).$$

Combining the above,

$$\text{rank}(\nabla^2 s_3(\theta_*)) = \text{rank}(d_\eta B(\eta_*)) \geq \sum_{l=0}^{10} (2l+1) - 3 + \sum_{l=11}^L (2l+1) = d-3,$$

which completes the proof that  $\text{trdeg}(\mathcal{R}_{\leq 3}^G) = d-3$ .  $\square$

**Lemma D.4.** *Suppose  $L \geq 11$ . For each  $l \in \{11, \dots, L\}$ , let  $\mathcal{J}^{(l)}$  be the set of tuples  $(l, l', l'') \in \mathcal{J}$  where  $l$  takes this fixed value, and where  $l' \leq l$  and  $l'' \leq l$ . Denote*

$$B^{(l)}(\eta) = \left( B_{l, l', l''}(\eta) : (l, l', l'') \in \mathcal{J}^{(l)} \right) \in \mathbb{C}^{|\mathcal{J}^{(l)}|}$$

and let  $d_{\eta^{(l)}} B^{(l)} \in \mathbb{C}^{|\mathcal{J}^{(l)}| \times (2l+1)}$  be the submatrix of  $d_\eta B$  corresponding to the derivative of  $B^{(l)}$  in  $\eta^{(l)}$ . Then for all generic  $\eta_* \in \mathbb{R}^d$ ,

$$\text{rank} \left( d_{\eta^{(l)}} B^{(l)}(\eta_*) \right) = 2l+1.$$

*Proof.* By Fact 4.5, it suffices to show  $\text{rank}(d_{\eta^{(l)}} B^{(l)}(\eta_*)) = 2l+1$  for a single point  $\eta_* \in \mathbb{R}^d$ . Our strategy is to choose  $\eta_*$  with many coordinates equal to 0, such that  $d_{\eta^{(l)}} B^{(l)}(\eta_*)$  has a sparse structure and its rank may be explicitly analyzed. Specifically, we choose  $\eta_*$  so that

$$\begin{aligned} \text{For } l' \in \{l-1, l\} : v_{*, m'}^{(l')}, w_{*, m'}^{(l')} &= 0 \text{ unless } m' = l-1 \\ \text{For } l' \in \{0, 1, 4, 5, \dots, l-2\} : v_{*, m'}^{(l')}, w_{*, m'}^{(l')} &= 0 \text{ unless } m' = l' \\ \text{For } l' \in \{2, 3\} : v_{*, m'}^{(l')}, w_{*, m'}^{(l')} &= 0 \text{ unless } m' \in \{l', l'-1\}. \end{aligned} \quad (\text{D.19})$$

We choose the values of the non-zero coordinates of  $\eta_*$  to be generic. The rest of this proof checks that  $\text{rank}(d_{\eta^{(l)}} B^{(l)}(\eta_*)) = 2l+1$  holds under this choice.

Recall the form of  $B_{l, l', l''}$  from (4.2). We first compute  $d_{\eta^{(l)}} B_{l, l', l''}$  in the two cases: (i)  $l', l'' < l$  and (ii)  $l' = l$  and  $l'' < l$ .

**Case I:**  $l', l'' < l$ . For each  $k = 0, \dots, l$ , the derivatives  $\partial_{v_k^{(l)}}, \partial_{w_k^{(l)}}$  apply only to the terms  $\overline{u_m^{(l)}}$  in (4.2) for  $m \in \{+k, -k\}$ . We have  $u_m^{(l)} = v_m^{(l)} + \mathbf{i} w_m^{(l)} = (-1)^m v_{-m}^{(l)} - \mathbf{i} \cdot (-1)^m w_{-m}^{(l)}$  where the second equality applies the sign symmetry (D.37). Thus (relabeling  $k$  by  $m$ ), for  $m > 0$  strictly,

$$\partial_{v_m^{(l)}} B_{l, l', l''} = \sum_{m'=-l'}^{l'} C_{m, m', m+m'}^{l, l', l''} \overline{u_{m'}^{(l')}} u_{m+m'}^{(l'')} + (-1)^m C_{-m, m', -m+m'}^{l, l', l''} \overline{u_{m'}^{(l')}} u_{-m+m'}^{(l'')}$$

<sup>1</sup>An equivalent statement was verified in [BBSK<sup>+</sup>17, Theorem 5.5] corresponding to the case  $F=10$ , using exact-precision numerical arithmetic.

$$\partial_{w_m^{(l)}} B_{l,l',l''} = \sum_{m'=-l'}^{l'} -\mathbf{i} \cdot C_{m,m',m+m'}^{l,l',l''} \overline{u_{m'}^{(l')}} u_{m+m'}^{(l'')} + (-1)^m \cdot \mathbf{i} \cdot C_{-m,m',-m+m'}^{l,l',l''} \overline{u_{m'}^{(l')}} u_{-m+m'}^{(l'')}.$$

Re-indexing  $m' \mapsto -m'$  for the summations of the second terms, and applying the symmetries (D.10) and (D.12), we obtain

$$\partial_{v_m^{(l)}} B_{l,l',l''} = \sum_{m'=-l'}^{l'} 2C_{m,m',m+m'}^{l,l',l''} \times \begin{cases} \operatorname{Re} \overline{u_{m'}^{(l')}} u_{m+m'}^{(l'')} & \text{if } l+l'+l'' \text{ is even} \\ \mathbf{i} \cdot \operatorname{Im} \overline{u_{m'}^{(l')}} u_{m+m'}^{(l'')} & \text{if } l+l'+l'' \text{ is odd,} \end{cases} \quad (\text{D.20})$$

$$\partial_{w_m^{(l)}} B_{l,l',l''} = \sum_{m'=-l'}^{l'} 2C_{m,m',m+m'}^{l,l',l''} \times \begin{cases} \operatorname{Im} \overline{u_{m'}^{(l')}} u_{m+m'}^{(l'')} & \text{if } l+l'+l'' \text{ is even} \\ (-\mathbf{i}) \cdot \operatorname{Re} \overline{u_{m'}^{(l')}} u_{m+m'}^{(l'')} & \text{if } l+l'+l'' \text{ is odd.} \end{cases} \quad (\text{D.21})$$

For  $m = 0$ , we have similarly

$$\partial_{v_0^{(l)}} B_{l,l',l''} = \sum_{m'=-l'}^{l'} C_{0,m',m'}^{l,l',l''} \overline{u_{m'}^{(l')}} u_{m'}^{(l'')}. \quad (\text{D.22})$$

**Case II:**  $l' = l$  and  $l'' < l$ . An additional contribution to each derivative  $\partial_{v_k^{(l)}}, \partial_{w_k^{(l)}}$  arises from differentiating  $u_{m'}^{(l')}$  in (4.2). By symmetry of (4.2) with respect to interchanging  $l$  and  $l'$ , this has the effect of doubling each of the above expressions. So for  $m > 0$ , we have

$$\partial_{v_m^{(l)}} B_{l,l,l''} = \sum_{m'=-l}^l 4C_{m,m',m+m'}^{l,l,l''} \times \begin{cases} \operatorname{Re} \overline{u_{m'}^{(l)}} u_{m+m'}^{(l'')} & \text{if } l'' \text{ is even} \\ \mathbf{i} \cdot \operatorname{Im} \overline{u_{m'}^{(l)}} u_{m+m'}^{(l'')} & \text{if } l'' \text{ is odd,} \end{cases} \quad (\text{D.23})$$

$$\partial_{w_m^{(l)}} B_{l,l,l''} = \sum_{m'=-l}^l 4C_{m,m',m+m'}^{l,l,l''} \times \begin{cases} \operatorname{Im} \overline{u_{m'}^{(l)}} u_{m+m'}^{(l'')} & \text{if } l'' \text{ is even} \\ (-\mathbf{i}) \cdot \operatorname{Re} \overline{u_{m'}^{(l)}} u_{m+m'}^{(l'')} & \text{if } l'' \text{ is odd.} \end{cases} \quad (\text{D.24})$$

For  $m = 0$  we have

$$\partial_{v_0^{(l)}} B_{l,l,l''} = \sum_{m'=-l}^l 2C_{0,m',m'}^{l,l,l''} \overline{u_{m'}^{(l)}} u_{m'}^{(l'')}. \quad (\text{D.25})$$

Now specializing these derivatives to  $\eta_*$  of the form (D.19), we observe for example the following: If  $l' \in \{l-1, l\}$  and  $l'' \in \{4, \dots, l-2\}$ , then  $\partial_{v_m^{(l)}} B_{l,l',l''}, \partial_{w_m^{(l)}} B_{l,l',l''}$  are 0 unless  $|m+m'| = l''$  for either  $m' = l-1$  or  $m' = -(l-1)$ . Since  $0 \leq m \leq l$ , this occurs for only the single index  $m = (l-1) - l''$ . Thus, only two entries in the row  $d_{\eta^{(l)}} B_{l,l',l''}$  are non-zero, corresponding to  $\partial_{v_m^{(l)}}, \partial_{w_m^{(l)}}$  for this  $m$ . More generally, let us write the condition (D.19) succinctly as

$$\text{Type}(0), \text{Type}(1), \text{Type}(4), \dots, \text{Type}(l-2), \text{Type}(l-1) = 0, \quad \text{Type}(l) = 1, \quad \text{Type}(2, 3) = \{0, 1\}$$

where  $\text{Type}(l') = i$  means that  $v_{*,m'}^{(l')}, w_{*,m'}^{(l')} = 0$  except for  $m' \in l' - i$ . Then for  $\partial_{v_m^{(l)}} B_{l,l',l''}, \partial_{w_m^{(l)}} B_{l,l',l''}$  to be non-zero, we require  $|m+m'| \in l'' - \text{Type}(l'')$  for some  $m'$  satisfying  $|m'| \in l' - \text{Type}(l')$ . This occurs for the indices

$$m \in \left\{ |(l' - \text{Type}(l')) - (l'' - \text{Type}(l''))|, (l' - \text{Type}(l')) + (l'' - \text{Type}(l'')) \right\} \cap \{0, \dots, l\} \quad (\text{D.26})$$

where we use the set notations  $|A| = \{|a| : a \in A\}$ ,  $A - B = \{a - b : a \in A, b \in B\}$ , and  $A + B = \{a + b : a \in A, b \in B\}$ .

For the given value of  $l$ , note that

$$\mathcal{J}^{(l)} = \left\{ (l', l'') : 0 \leq l' \leq l, 0 \leq l'' \leq l, l' + l'' \geq l \right\}.$$

We label rows of  $d_{\eta^{(l)}} B^{(l)}(\eta_*)$  by the pairs  $(l', l'')$  where  $(l', l'') \in \mathcal{J}^{(l)}$ . We now choose  $2l+1$  such rows  $(l', l'')$  and check that the corresponding square  $(2l+1) \times (2l+1)$  submatrix is non-singular. These rows are indicated in the left column of the below table. The right column displays all values of  $m$  satisfying (D.26), i.e. for which  $\partial_{v_m^{(l)}}, \partial_{w_m^{(l)}}$  are non-zero in that row.

TABLE 1.

$(l', l'')$	Values of $m$
$(l-1, l-1)$ if $l$ is even or $(l, l-1)$ if $l$ is odd	0
$(l, l-2)$	1
$(l-1, l-2)$	1
$(l, l-3)$	2
$(l-1, l-3)$	2
$\vdots$	$\vdots$
$(l, 4)$	$l-5$
$(l-1, 4)$	$l-5$
$(l-5, 5)$	$l-10$ and $l$
$(l-4, 4)$	$l-8$ and $l$
$(l, 1)$	$l$ and $l-2$
$(l-1, 1)$	$l$ and $l-2$
$(l, 2)$	$l, l-2$ , and $l-3$
$(l-1, 2)$	$l, l-2$ , and $l-3$
$(l, 3)$	$l-3$ and $l-4$
$(l-1, 3)$	$l-3$ and $l-4$
$(l-3, 3)$	$l-6, l-5, l$ , and $l-1$
$(l-2, 2)$	$l-4, l-3, l$ , and $l-1$

To verify that this selected  $(2l+1) \times (2l+1)$  submatrix of  $d_{\eta^{(l)}} B^{(l)}(\eta_*)$  is non-singular, let us order its rows in the order of the above table, and its columns according to the ordering of variables

$$v_0^{(l)}, v_1^{(l)}, w_1^{(l)}, \dots, v_{l-5}^{(l)}, w_{l-5}^{(l)}, v_l^{(l)}, w_l^{(l)}, v_{l-2}^{(l)}, w_{l-2}^{(l)}, v_{l-3}^{(l)}, w_{l-3}^{(l)}, v_{l-4}^{(l)}, w_{l-4}^{(l)}, v_{l-1}^{(l)}, w_{l-1}^{(l)}$$

as they appear in the right column above. Then the table implies that this  $(2l+1) \times (2l+1)$  submatrix has a block lower-triangular structure with respect to  $2l+1 = 1 + 2 + \dots + 2$ . So it suffices to check that each  $1 \times 1$  and  $2 \times 2$  diagonal block is non-singular. It is tedious but straightforward to verify this explicitly, by computing their forms:

**Block corresponding to  $v_0^{(l)}$ :** For even  $l$ , we have that  $l + (l-1) + (l-1)$  is even. Then applying (D.22) and the symmetries (D.10) and (D.12), this  $1 \times 1$  matrix is

$$\partial_{v_0^{(l)}} B_{l,l-1,l-1}(\eta_*) = \left( C_{0,l-1,l-1}^{l,l-1,l-1} + C_{0,-(l-1),-(l-1)}^{l,l-1,l-1} \right) |u_{*,l-1}^{(l-1)}|^2 = 2C_{0,l-1,l-1}^{l,l-1,l-1} |u_{*,l-1}^{(l-1)}|^2.$$

For odd  $l$ , we have that  $l + l + (l-1)$  is even. Then applying instead (D.25), this  $1 \times 1$  matrix is

$$\partial_{v_0^{(l)}} B_{l,l,l-1}(\eta_*) = 2C_{0,l-1,l-1}^{l,l,l-1} \overline{u_{*,l-1}^{(l)}} u_{*,l-1}^{(l-1)} + 2C_{0,-(l-1),-(l-1)}^{l,l,l-1} u_{*,l-1}^{(l)} \overline{u_{*,l-1}^{(l-1)}} = 4C_{0,l-1,l-1}^{l,l,l-1} \operatorname{Re} \overline{u_{*,l-1}^{(l)}} u_{*,l-1}^{(l-1)}.$$

These two coefficients  $C_{0,l-1,l-1}^{l,l,l-1}$  and  $C_{0,-(l-1),-(l-1)}^{l,l,l-1}$  are non-zero by Lemma D.2, so this block is non-zero for generic values of the non-zero coordinates  $v_{*,l-1}^{(l-1)}, w_{*,l-1}^{(l-1)}, v_{*,l-1}^{(l)}, w_{*,l-1}^{(l)}$  of  $\eta_*$ .

**Blocks corresponding to  $(v_1^{(l)}, w_1^{(l)}), \dots, (v_{l-2}^{(l)}, w_{l-2}^{(l)})$ :** The calculations for all these blocks are similar. We demonstrate the case  $v_{l-3}^{(l)}, w_{l-3}^{(l)}$ : Applying (D.20–D.21) and (D.23–D.24), this  $2 \times 2$  block is

$$\begin{aligned} & \partial_{v_{l-3}^{(l)}, w_{l-3}^{(l)}} \left( B_{l,l,2}, B_{l,l-1,2} \right) (\eta_*) \\ &= \begin{pmatrix} 4C_{l-3,-(l-1),-2}^{l,l,2} \cdot \operatorname{Re} \overline{u_{*,-(l-1)}^{(l)}} u_{*,2}^{(2)} & 4C_{l-3,-(l-1),-2}^{l,l,2} \cdot \operatorname{Im} \overline{u_{*,-(l-1)}^{(l)}} u_{*,2}^{(2)} \\ 2iC_{l-3,-(l-1),-2}^{l,l-1,2} \cdot \operatorname{Im} \overline{u_{*,-(l-1)}^{(l-1)}} u_{*,2}^{(2)} & -2iC_{l-3,-(l-1),-2}^{l,l-1,2} \cdot \operatorname{Re} \overline{u_{*,-(l-1)}^{(l-1)}} u_{*,2}^{(2)} \end{pmatrix} \\ &= \begin{pmatrix} 4C_{l-3,-(l-1),-2}^{l,l,2} & 0 \\ 0 & 2iC_{l-3,-(l-1),-2}^{l,l-1,2} \end{pmatrix} \begin{pmatrix} \operatorname{Re} \overline{u_{*,-(l-1)}^{(l)}} u_{*,2}^{(2)} & \operatorname{Im} \overline{u_{*,-(l-1)}^{(l)}} u_{*,2}^{(2)} \\ \operatorname{Im} \overline{u_{*,-(l-1)}^{(l-1)}} u_{*,2}^{(2)} & -\operatorname{Re} \overline{u_{*,-(l-1)}^{(l-1)}} u_{*,2}^{(2)} \end{pmatrix} \end{aligned}$$

The coefficients  $C_{l-3,-(l-1),-2}^{l,l,2}$  and  $C_{l-3,-(l-1),-2}^{l,l-1,2}$  are both non-zero by Lemma D.2, so the first matrix of this product is non-singular. It is direct to check that the determinant of the second matrix is a non-zero polynomial of the six non-zero coordinates  $v_{*,l-1}^{(l)}, w_{*,l-1}^{(l)}, v_{*,l-1}^{(l-1)}, w_{*,l-1}^{(l-1)}, v_{*,2}^{(2)}, w_{*,2}^{(2)}$  of  $\eta_*$ . Then for generic values of these six coordinates, the determinant is non-zero, and this matrix is also non-singular.

**Blocks corresponding to  $(v_{l-1}^{(l)}, w_{l-1}^{(l)}), (v_l^{(l)}, w_l^{(l)})$ :** Applying (D.20–D.21), these  $2 \times 2$  matrices are

$$\begin{aligned} & \partial_{v_{l-1}^{(l)}, w_{l-1}^{(l)}} \left( B_{l,l-3,3}, B_{l,l-2,2} \right) (\eta_*) \\ &= \begin{pmatrix} 2C_{l-1,-(l-3),2}^{l,l-3,3} \cdot \operatorname{Re} \overline{u_{*,-(l-3)}^{(l-3)}} u_{*,2}^{(3)} & 2C_{l-1,-(l-3),2}^{l,l-3,3} \cdot \operatorname{Im} \overline{u_{*,-(l-3)}^{(l-3)}} u_{*,2}^{(3)} \\ 2C_{l-1,-(l-2),1}^{l,l-2,2} \cdot \operatorname{Re} \overline{u_{*,-(l-2)}^{(l-2)}} u_{*,1}^{(2)} & 2C_{l-1,-(l-2),1}^{l,l-2,2} \cdot \operatorname{Im} \overline{u_{*,-(l-2)}^{(l-2)}} u_{*,1}^{(2)} \end{pmatrix}, \\ & \partial_{v_l^{(l)}, w_l^{(l)}} \left( B_{l,l-5,5}, B_{l,l-4,4} \right) (\eta_*) \\ &= \begin{pmatrix} 2C_{l,-(l-5),5}^{l,l-5,5} \cdot \operatorname{Re} \overline{u_{*,-(l-5)}^{(l-5)}} u_{*,5}^{(5)} & 2C_{l,-(l-5),5}^{l,l-5,5} \cdot \operatorname{Im} \overline{u_{*,-(l-5)}^{(l-5)}} u_{*,5}^{(5)} \\ 2C_{l,-(l-4),4}^{l,l-4,4} \cdot \operatorname{Re} \overline{u_{*,-(l-4)}^{(l-4)}} u_{*,4}^{(4)} & 2C_{l,-(l-4),4}^{l,l-4,4} \cdot \operatorname{Im} \overline{u_{*,-(l-4)}^{(l-4)}} u_{*,4}^{(4)} \end{pmatrix}. \end{aligned}$$

The coefficients  $C_{l-1,-(l-3),2}^{l,l-3,3}, C_{l-1,-(l-2),1}^{l,l-2,2}, C_{l,-(l-5),5}^{l,l-5,5}, C_{l,-(l-4),4}^{l,l-4,4}$  are non-zero by Lemma D.2. Then the determinant of the first matrix is a non-zero polynomial of the eight coordinates

$$v_{*,l-3}^{(l-3)}, w_{*,l-3}^{(l-3)}, v_{*,l-2}^{(l-2)}, w_{*,l-2}^{(l-2)}, v_{*,1}^{(2)}, w_{*,1}^{(2)}, v_{*,2}^{(3)}, w_{*,2}^{(3)},$$

and that of the second matrix is a non-zero polynomial of the eight coordinates

$$v_{*,l-5}^{(l-5)}, w_{*,l-5}^{(l-5)}, v_{*,l-4}^{(l-4)}, w_{*,l-4}^{(l-4)}, v_{*,5}^{(5)}, w_{*,5}^{(5)}, v_{*,4}^{(4)}, w_{*,4}^{(4)}.$$

(Note that these eight coordinates are distinct when  $l \geq 11$ .) Thus for generic values of these coordinates, these matrices are non-singular.

Combining these cases, we have shown that each  $1 \times 1$  and  $2 \times 2$  diagonal block of this  $(2l+1) \times (2l+1)$  submatrix is nonsingular for generic choices of the non-zero coordinates of  $\eta_*$ . Then they are also simultaneously nonsingular for generic choices of these coordinates, so in particular there exists  $\eta_* \in \mathbb{R}^d$  where  $d_{\eta^{(l)}} B^{(l)}(\eta_*)$  has full column rank  $2l+1$ . Then  $d_{\eta^{(l)}} B^{(l)}(\eta_*)$  must also have full column rank  $2l+1$  for all generic  $\eta_* \in \mathbb{R}^d$ , concluding the proof.  $\square$

### D.3. Unprojected cryo-EM.

**D.3.1. Function basis.** We first describe in further detail the function basis and rotational action of  $\mathrm{SO}(3)$  on basis coefficients in the unprojected cryo-EM example of Section 4.2. This is similar to the setup of the model with  $S$  spherical shells in [BBSK<sup>+</sup>17, Section 5.5].

For  $f \in L_2(\mathbb{R}^3, \mathbb{C})$ , denote its Fourier transform

$$\hat{f}(k_1, k_2, k_3) = \int_{\mathbb{R}^3} e^{-2\pi i(k_1 x_1 + k_2 x_2 + k_3 x_3)} f(x_1, x_2, x_3) dx_1 dx_2 dx_3. \quad (\text{D.27})$$

We reparametrize  $k = (k_1, k_2, k_3) \in \mathbb{R}^3$  in the Fourier domain by spherical coordinates  $(\rho, \phi_1, \phi_2)$ , and write with a slight abuse of notation  $\hat{f}(\rho, \phi_1, \phi_2)$  for this parametrization.

Let  $\hat{j}_{lsm}$  be as defined in (4.3), where  $y_{lm}$  are the complex spherical harmonics in (D.2) and  $\{z_s : s \geq 1\}$  are any functions  $z_s : [0, \infty) \rightarrow \mathbb{R}$  satisfying the orthogonality relation (4.4). By the spherical change-of-coordinates  $dk_1 dk_2 dk_3 = \rho^2 \sin \phi_1 d\rho d\phi_1 d\phi_2$ , these functions  $\{\hat{j}_{lsm}\}$  are orthonormal in  $L_2(\mathbb{R}^3, \mathbb{C})$ . Then so are their inverse Fourier transforms  $\{j_{lsm}\}$ , by the Parseval relation.

Recall the space of  $(L, S_0, \dots, S_L)$ -bandlimited functions (4.6). By linearity of the Fourier transform, the basis representation (4.6) is equivalent to

$$\hat{f} = \sum_{(l,s,m) \in \mathcal{I}} u_m^{(ls)} \hat{j}_{lsm} \quad (\text{D.28})$$

in the Fourier domain. A function  $f \in L_2(\mathbb{R}^3, \mathbb{C})$  is real-valued if and only if

$$\hat{f}(\rho, \phi_1, \phi_2) = \overline{\hat{f}(\rho, \pi - \phi_1, \pi + \phi_2)},$$

where  $(\rho, \pi - \phi_1, \pi + \phi_2)$  are the coordinates for the reflection of  $(\rho, \phi_1, \phi_2)$  about the origin. Applying  $P_{lm}(x) = (-1)^{l+m} P_{lm}(-x)$  by (D.1) and hence  $y_{l,m}(\pi - \phi_1, \pi + \phi_2) = (-1)^l y_{l,m}(\phi_1, \phi_2) = (-1)^{l+m} \overline{y_{l,-m}(\phi_1, \phi_2)}$  by (D.2) and (D.3), it may be checked that this condition is equivalent to the sign symmetry

$$u_m^{(ls)} = (-1)^{l+m} \overline{u_{-m}^{(ls)}} \quad (\text{D.29})$$

in the basis representations (4.6) and (D.28). We may then define a real basis  $\{h_{lsm}\}$  by

$$h_{l,s,m} = \begin{cases} \frac{1}{\sqrt{2}} (j_{l,s,-m} + (-1)^{l+m} j_{l,s,m}) & \text{if } m > 0 \\ \mathbf{i}^l \cdot j_{l,s,0} & \text{if } m = 0 \\ \frac{1}{\sqrt{2}} (j_{l,s,m} - (-1)^{l+m} j_{l,s,-m}) & \text{if } m < 0. \end{cases} \quad (\text{D.30})$$

Note that by this definition,  $h_{lsm}$  satisfies (D.29) for its coefficients  $u_m^{(ls)}$  in the basis  $\{j_{lsm}\}$ , and hence is real-valued. Thus  $\{h_{lsm}\}$  forms an orthonormal basis for  $L_2(\mathbb{R}^3, \mathbb{R})$ . For any  $(L, S_0, \dots, S_L)$ -bandlimited function  $f \in L_2(\mathbb{R}^3, \mathbb{R})$ , writing its orthogonal decompositions

$$f = \sum_{(l,s,m) \in \mathcal{I}} u_m^{(ls)} \cdot j_{lsm} = \sum_{(l,s,m) \in \mathcal{I}} \theta_m^{(ls)} \cdot h_{lsm},$$

the coefficients  $\{u_m^{(ls)}\}$  and  $\{\theta_m^{(ls)}\}$  are then related by a unitary transform  $u = \hat{V}^* \theta$  defined as

$$u_m^{(ls)} = \begin{cases} \frac{(-1)^{l+m}}{\sqrt{2}} (\theta_{|m|}^{(ls)} - \mathbf{i} \theta_{-|m|}^{(ls)}) & \text{if } m > 0 \\ \mathbf{i}^l \cdot \theta_0^{(ls)} & \text{if } m = 0 \\ \frac{1}{\sqrt{2}} (\theta_{|m|}^{(ls)} + \mathbf{i} \theta_{-|m|}^{(ls)}) & \text{if } m < 0. \end{cases} \quad (\text{D.31})$$

Here, the sign symmetry (D.29) and transform  $\hat{V}$  are different from (D.12) and the transform  $V$  defined by (D.11) in the example of spherical registration, because we are modeling the Fourier transform  $\hat{f}$  rather than  $f$  in the spherical harmonics basis, but we assume that  $f$  rather than  $\hat{f}$  is real-valued.

The rotation of  $\mathbb{R}^3$  induces the following rotational action on the coefficient vector  $\theta \in \mathbb{R}^d$ .

**Lemma D.5.** *The action of  $\text{SO}(3)$  on the space of real-valued  $(L, S_0, \dots, S_L)$ -bandlimited functions is given by*

$$\mathbf{G} = \left\{ \hat{V} \cdot D(\mathbf{g}) \cdot \hat{V}^* : \mathbf{g} \in \text{SO}(3) \right\} \subset \text{O}(d) \quad (\text{D.32})$$

where  $\hat{V} \in \mathbb{C}^{d \times d}$  is the unitary transform defined in (D.31) for which  $u = \hat{V}^* \theta$ , and  $D(\mathbf{g})$  is the block-diagonal matrix

$$D(\mathbf{g}) = \bigoplus_{l=0}^L \bigoplus_{s=1}^{S_l} D^{(l)}(\mathbf{g}) \quad (\text{D.33})$$

with diagonal blocks  $D^{(l)}(\mathbf{g}) \in \mathbb{C}^{(2l+1) \times (2l+1)}$  given by the complex Wigner  $D$ -matrices.

*Proof.* Note that if  $f_{\mathbf{g}}(x) = f(\mathbf{g}^{-1} \cdot x)$ , then its Fourier transform undergoes the same rotation  $\hat{f}_{\mathbf{g}}(k) = \hat{f}(\mathbf{g}^{-1} \cdot k)$ , by (D.27). Writing

$$\hat{f}(\rho, \phi_1, \phi_2) = \sum_{l=0}^L \sum_{s=1}^{S_l} z_s(\rho) \cdot \hat{f}_{ls}(\phi_1, \phi_2), \quad \hat{f}_{ls}(\phi_1, \phi_2) = \sum_{m=-l}^l u_m^{(ls)} y_{lm}(\phi_1, \phi_2),$$

each function  $\hat{f}_{ls}$  is defined on the unit sphere, and the rotation by  $\mathbf{g}$  acts separately on each such function  $\hat{f}_{ls}$  via the map  $u^{(ls)} \mapsto D^{(l)}(\mathbf{g}) u^{(ls)}$  described in Appendix D.1.2. Thus rotation by  $\mathbf{g}$  induces the transformation  $u \mapsto D(\mathbf{g})u$  on the complex coefficient vector  $u$ , for  $D(\mathbf{g})$  as defined in (D.33). Applying the unitary relations  $u = \hat{V}^* \theta$  and  $\theta = \hat{V} u$ , this rotation then induces the transformation  $\theta \mapsto \hat{V} \cdot D(\mathbf{g}) \cdot \hat{V}^* \theta$  on the real coefficients  $\theta$ .  $\square$

**D.3.2. Terms of the high-noise series expansion.** We prove Theorem 4.9 on the forms of  $s_1(\theta)$ ,  $s_2(\theta)$ , and  $s_3(\theta)$ .

*Proof of Theorem 4.9.* Similar to the proof of Theorem 4.4, consider two different real coefficient vectors  $\theta, \vartheta \in \mathbb{R}^d$ , with corresponding complex coefficients  $u = \hat{V}^* \theta$  and  $v = \hat{V}^* \vartheta$ . We compute  $\mathbb{E}_g[\langle \theta, g \cdot \vartheta \rangle^k]$  for  $k = 1, 2, 3$ .

**Case  $k = 1$ :** By Lemma D.5, for any  $g \in \mathbf{G}$ ,

$$\langle \theta, g \cdot \vartheta \rangle = \langle \theta, \hat{V} D(\mathbf{g}) \hat{V}^* \vartheta \rangle = \langle u, D(\mathbf{g}) v \rangle.$$

From the block-diagonal form of  $D(\mathbf{g})$  in (D.33), we obtain

$$\langle \theta, g \cdot \vartheta \rangle = \sum_{l=0}^L \sum_{s=1}^{S_l} \langle u^{(ls)}, D^{(l)}(\mathbf{g}) v^{(ls)} \rangle = \sum_{l=0}^L \sum_{s=1}^{S_l} \sum_{q, m=-l}^l \overline{u_q^{(ls)}} D_{qm}^{(l)}(\mathbf{g}) v_m^{(ls)}. \quad (\text{D.34})$$

Applying the identities (D.4) yields

$$\mathbb{E}_g[\langle \theta, g \cdot \vartheta \rangle] = \sum_{s=1}^{S_0} \overline{u_0^{(0s)}} v^{(0s)}.$$

Write as shorthand  $u^{(0s)} = u_0^{(0s)}$ ,  $v^{(0s)} = v_0^{(0s)}$ , and observe from (D.29) that these are real-valued. Then applying this to (D.14), we have

$$s_1(\theta) = \frac{1}{2} \mathbb{E}_g[\langle \theta, g \cdot \theta \rangle] - \mathbb{E}_g[\langle \theta_*, g \cdot \theta \rangle] + \frac{1}{2} \mathbb{E}_g[\langle \theta_*, g \cdot \theta_* \rangle] = \frac{1}{2} \sum_{s=1}^{S_0} \left( u^{(0s)}(\theta) - u^{(0s)}(\theta_*) \right)^2.$$

**Case  $k = 2$ :** We take the expected square on both sides of (D.34), applying (D.29) and the relation (D.5). Then

$$\begin{aligned} \mathbb{E}_g[\langle \theta, g \cdot \vartheta \rangle^2] &= \sum_{l=0}^L \sum_{s, s'=1}^{S_l} \sum_{q, m=-l}^l \frac{(-1)^{m+q}}{2l+1} \overline{u_q^{(ls)} u_{-q}^{(ls')}} v_m^{(ls)} v_{-m}^{(ls')} \\ &= \sum_{l=0}^L \sum_{s, s'=1}^{S_l} \sum_{q, m=-l}^l \frac{1}{2l+1} \overline{u_q^{(ls)} u_q^{(ls')}} v_m^{(ls)} \overline{v_m^{(ls')}} \\ &= \sum_{l=0}^L \frac{1}{2l+1} \sum_{s, s'=1}^{S_l} \langle u^{(ls)}, u^{(ls')} \rangle \cdot \overline{\langle v^{(ls)}, v^{(ls')} \rangle}. \end{aligned}$$

Note that from the isometry  $\langle u^{(ls)}, u^{(ls')} \rangle = \langle \theta^{(ls)}, \theta^{(ls')} \rangle$ , the inner-products on the last line are real. Then applying this to (D.14),

$$\begin{aligned} s_2(\theta) &= \frac{1}{4} \mathbb{E}_g[\langle \theta, g \cdot \theta \rangle^2] - \frac{1}{2} \mathbb{E}_g[\langle \theta_*, g \cdot \theta \rangle^2] + \frac{1}{4} \mathbb{E}_g[\langle \theta_*, g \cdot \theta_* \rangle^2] \\ &= \sum_{l=0}^L \frac{1}{4(2l+1)} \sum_{s, s'=1}^{S_l} \left( \langle u^{(ls)}(\theta), u^{(ls')}(\theta) \rangle - \langle u^{(ls)}(\theta_*), u^{(ls')}(\theta_*) \rangle \right)^2. \end{aligned}$$

**Case  $k = 3$ :** We now take the expected cube on both sides of (D.34) and apply the relations (D.6) and (D.29). Then

$$\begin{aligned} \mathbb{E}_g[\langle \theta, g \cdot \vartheta \rangle^3] &= \sum_{\substack{l, l', l''=0 \\ |l-l'| \leq l'' \leq l+l'}}^L \sum_{s=1}^{S_l} \sum_{s'=1}^{S_{l'}} \sum_{s''=1}^{S_{l''}} \sum_{q, m=-l}^l \sum_{q', m'=-l'}^{l'} \frac{1}{2l''+1} \\ &\quad \cdot C_{q, q', q+q'}^{l, l', l''} C_{m, m', m+m'}^{l, l', l''} \overline{u_q^{(ls)} u_{q'}^{(l's')}} u_{q+q'}^{(l'' s'')} v_m^{(ls)} v_{m'}^{(l's')} \overline{v_{m+m'}^{(l'' s'')}}. \end{aligned}$$

Recall that

$$B_{(l, s), (l', s'), (l'', s'')}(\theta) = \sum_{m=-l}^l \sum_{m'=-l'}^{l'} C_{m, m', m+m'}^{l, l', l''} \overline{u_m^{(ls)} u_{m'}^{(l's')}} u_{m+m'}^{(l'' s'')}, \quad u = \hat{V}^* \theta$$

with the convention (D.9). Changing indices  $(m, m') \mapsto (-m, -m')$  and applying the symmetries (D.10) and (D.29), we have

$$\begin{aligned} B_{(l,s),(l',s'),(l'',s'')}(\theta) &= \sum_{m=-l}^l \sum_{m'=-l'}^{l'} C_{-m,-m',-m-m'}^{l,l',l''} \overline{u_{-m}^{(ls)} u_{-m'}^{(l's')}} u_{-m-m'}^{(l''s'')} \\ &= \sum_{m=-l}^l \sum_{m'=-l'}^{l'} (-1)^{l+l'+l''} C_{m,m',m+m'}^{l,l',l''} \cdot (-1)^{l+l'+l''+m+m'+(m+m')} u_m^{(ls)} u_{m'}^{(l's')} \overline{u_{m+m'}^{(l''s'')}} \\ &= \overline{B_{(l,s),(l',s'),(l'',s'')}(\theta)}. \end{aligned}$$

Thus (in contrast to spherical registration in Appendix D.2)  $B_{(l,s),(l',s'),(l'',s'')}(\theta)$  is always real-valued. Then the above may be written as

$$\mathbb{E}_g[\langle \theta, g \cdot \vartheta \rangle^3] = \sum_{\substack{l,l',l''=0 \\ |l-l'| \leq l'' \leq l+l'}}^L \frac{1}{2l''+1} \sum_{s=1}^{S_l} \sum_{s'=1}^{S_{l'}} \sum_{s''=1}^{S_{l''}} B_{(l,s),(l',s'),(l'',s'')}(\theta) B_{(l,s),(l',s'),(l'',s'')}(\vartheta).$$

Then applying this to (D.14),

$$\begin{aligned} s_3(\theta) &= \frac{1}{12} \mathbb{E}_g[\langle \theta, g \cdot \theta \rangle^3] - \frac{1}{6} \mathbb{E}_g[\langle \theta_*, g \cdot \theta \rangle^3] + \frac{1}{12} \mathbb{E}_g[\langle \theta_*, g \cdot \theta_* \rangle^3] \\ &= \frac{1}{12} \sum_{\substack{l,l',l''=0 \\ |l-l'| \leq l'' \leq l+l'}}^L \frac{1}{2l''+1} \sum_{s=1}^{S_l} \sum_{s'=1}^{S_{l'}} \sum_{s''=1}^{S_{l''}} (B_{(l,s),(l',s'),(l'',s'')}(\theta) - B_{(l,s),(l',s'),(l'',s'')}(\theta_*))^2. \end{aligned} \quad (\text{D.35})$$

□

**D.3.3. Transcendence degrees.** We now prove Theorem 4.6 on the sequences of transcendence degrees.

*Proof of Theorem 4.6.* Recall the form of  $\mathbf{G}$  in Lemma D.5, let  $\hat{V}^{(l)} \in \mathbb{C}^{(2l+1) \times (2l+1)}$  represent the unitary map  $u^{(ls)} \mapsto \theta^{(ls)}$  (which has the same form for each  $s = 1, \dots, S_l$ ), and denote by  $g^{(l)}(\mathbf{g}) = \hat{V}^{(l)} \cdot D^{(l)}(\mathbf{g}) \cdot (\hat{V}^{(l)})^*$  the irreducible representation of  $\text{SO}(3)$  acting on each subvector  $\theta^{(ls)} \in \mathbb{R}^{2l+1}$ . For  $l = 1$ , note that this representation  $g^{(1)}$  is isomorphic to  $\text{SO}(3)$  acting on  $\mathbb{R}^3$  by rotations, so its action on the pair of subvectors  $(\theta^{(1,1)}, \theta^{(1,2)}) \in \mathbb{R}^3 \oplus \mathbb{R}^3$  has the trivial stabilizer subgroup  $\{\text{Id}\}$  for generic  $(\theta^{(1,1)}, \theta^{(1,2)})$ . This implies for any  $L \geq 1$  and  $S_1 \geq 2$  that, as in the proof of Theorem 4.1, the stabilizer subgroup  $\mathbf{G}_\theta$  of any extension of  $(\theta^{(1,1)}, \theta^{(1,2)})$  to  $\theta \in \mathbb{R}^d$  satisfies  $\dim(\mathbf{G}_\theta) = 0$ , and hence  $d_0 = \dim(\mathbf{G}) - \min_\theta \dim(\mathbf{G}_\theta) = 3$ .

We now compute  $\text{trdeg}(\mathcal{R}_{\leq k}^{\mathbf{G}})$  for  $k = 1, 2, 3$  using Lemma 2.8. Recall the forms of  $s_1(\theta)$  and  $s_2(\theta)$  in Theorem 4.9. For  $k = 1$ , differentiating twice at  $\theta = \theta_*$  yields

$$\nabla^2 s_1(\theta_*) = \sum_{s=1}^{S_0} \nabla u^{(0s)}(\theta_*) \nabla u^{(0s)}(\theta_*)^\top.$$

Recalling  $u^{(0s)}(\theta) = \theta_0^{(0s)}$ , these vectors  $\{\nabla u^{(0s)}(\theta_*)\}_{s=1}^{S_0}$  are  $S_0$  different standard basis vectors, so

$$\text{trdeg}(\mathcal{R}_{\leq 1}^{\mathbf{G}}) = \text{rank}(\nabla^2 s_1(\theta_*)) = S_0.$$

For  $k = 2$ , differentiating twice at  $\theta = \theta_*$  yields

$$\begin{aligned} \nabla^2 s_1(\theta_*) + \nabla^2 s_2(\theta_*) &= \sum_{s=1}^{S_0} \nabla u^{(0s)}(\theta_*) \nabla u^{(0s)}(\theta_*)^\top \\ &\quad + \sum_{l=0}^L \frac{1}{2(2l+1)} \sum_{s,s'=1}^{S_l} \nabla[\langle u^{(ls)}(\theta), u^{(ls')}(\theta) \rangle] \nabla[\langle u^{(ls)}(\theta), u^{(ls')}(\theta) \rangle]^\top \Big|_{\theta=\theta_*}. \end{aligned}$$

Defining matrices  $G^0$  and  $G$  with the columns

$$G_s^0 := \nabla u^{(0s)}(\theta_*) \quad \text{for } 1 \leq s \leq S_0$$

$$G_{ls s'} := \frac{1}{\sqrt{2(2l+1)}} \nabla[\langle u^{(ls)}(\theta), u^{(ls')}(\theta) \rangle] \Big|_{\theta=\theta_*} \quad \text{for } 0 \leq l \leq L, 1 \leq s, s' \leq S_l,$$

this may be written as

$$\nabla^2 s_1(\theta_*) + \nabla^2 s_2(\theta_*) = G^0(G^0)^\top + GG^\top = [G \mid G^0][G \mid G^0]^\top.$$

For generic  $\theta_*$  where  $\theta_0^{(0s)} \neq 0$ , the column span of  $G^0$  coincides with the span of columns  $\{G_{0ss} : s = 1, \dots, S_0\}$  of  $G$ . Thus

$$\text{trdeg}(\mathcal{R}_{\leq 2}^G) = \text{rank}(\nabla^2 s_1(\theta_*) + \nabla^2 s_2(\theta_*)) = \text{rank}([G \mid G^0]) = \text{rank}(G).$$

Applying the isometry  $\langle u^{(ls)}(\theta), u^{(ls')}(\theta) \rangle = \langle \theta^{(ls)}, \theta^{(ls')} \rangle$ , Lemma D.6 below shows that

$$\text{rank}(G) = \sum_{l=0}^L \begin{cases} \frac{S_l(S_l+1)}{2} & \text{if } S_l < 2l+1, \\ (2l+1)(S_l-l) & \text{if } S_l \geq 2l+1, \end{cases}$$

establishing the desired form for  $k=2$ .

For  $k=3$ , we have  $\text{trdeg}(\mathcal{R}_{\leq 3}^G) \leq \text{trdeg}(\mathcal{R}^G) = d-d_0 = d-3$ , so it suffices to show  $\text{rank}(\nabla^2 s_3(\theta_*)) \geq d-3$ . Writing the index set

$$\mathcal{K} = \left\{ ((l, s), (l', s'), (l'', s'')) : 0 \leq l, l', l'' \leq L, |l-l'| \leq l'' \leq l+l', 1 \leq s \leq S_l, 1 \leq s' \leq S_{l'}, 1 \leq s'' \leq S_{l''} \right\},$$

by Theorem 4.9 we have

$$s_3(\theta) = \sum_{((l,s),(l',s'),(l'',s'')) \in \mathcal{K}} \frac{1}{12} \cdot \frac{1}{2l''+1} \cdot (B_{(l,s),(l',s'),(l'',s'')}(\theta) - B_{(l,s),(l',s'),(l'',s'')}(\theta_*))^2$$

for the function  $B_{(l,s),(l',s'),(l'',s'')}(\theta)$  defined in (4.9). Recall from the proof of Theorem 4.9 that  $B_{(l,s),(l',s'),(l'',s'')}(\theta)$  is real-valued. Let us denote

$$B(\theta) = \left( B_{(l,s),(l',s'),(l'',s'')}(\theta) : ((l,s),(l',s'),(l'',s'')) \in \mathcal{K} \right), \quad B : \mathbb{R}^d \rightarrow \mathbb{R}^{|\mathcal{K}|},$$

and write  $dB(\theta) \in \mathbb{R}^{|\mathcal{K}| \times d}$  for its the derivative in  $\theta$ . Then, applying the chain rule to differentiate  $s_3(\theta)$  twice at  $\theta = \theta_*$ , we obtain

$$\nabla^2 s_3(\theta_*) = dB(\theta_*)^\top \cdot \text{diag} \left( \frac{1}{6(2l''+1)} : ((l,s),(l',s'),(l'',s'')) \in \mathcal{K} \right) \cdot dB(\theta_*).$$

The diagonal matrix in the middle has full rank, so

$$\text{rank}(\nabla^2 s_3(\theta_*)) = \text{rank}(dB(\theta_*)). \quad (\text{D.36})$$

To analyze this rank, recall the complex parametrization  $u = \hat{V}^* \theta \in \mathbb{C}^d$  from (D.31), where  $u$  satisfies the symmetry (D.29). Let us write the real and imaginary parts of  $u$  as

$$u_m^{(ls)} = v_m^{(ls)} + \mathbf{i} w_m^{(ls)}$$

so that this symmetry (D.29) is equivalent to

$$v_{-m}^{(ls)} = (-1)^{l+m} v_m^{(ls)}, \quad w_{-m}^{(ls)} = (-1)^{l+m+1} w_m^{(ls)}. \quad (\text{D.37})$$

Note that for  $m=0$ , this implies  $v_0^{(ls)} = 0$  when  $l$  is odd and  $w_0^{(ls)} = 0$  when  $l$  is even. Then, setting

$$\eta^{(ls)}(\theta) = \begin{cases} (v_0^{(ls)}, v_1^{(ls)}, w_1^{(ls)}, \dots, v_l^{(ls)}, w_l^{(ls)}) & \text{if } l \text{ is even} \\ (w_0^{(ls)}, v_1^{(ls)}, w_1^{(ls)}, \dots, v_l^{(ls)}, w_l^{(ls)}) & \text{if } l \text{ is odd,} \end{cases} \quad (\text{D.38})$$

these coordinates  $\eta^{(ls)} \in \mathbb{R}^{2l+1}$  provide a (linear) invertible reparametrization of  $\theta^{(ls)}$ . This defines a reparametrization

$$\eta(\theta) = \left( \eta^{(ls)}(\theta) : 0 \leq l \leq L, 1 \leq s \leq S_l \right) \in \mathbb{R}^d \quad (\text{D.39})$$

with inverse function  $\theta(\eta)$ . Writing as shorthand  $\eta_* = \eta(\theta_*)$  and  $B(\eta) \equiv B(\theta(\eta))$ , and denoting by  $d_\eta B(\eta)$  the derivative of  $B$  in the new variables  $\eta$ , (D.36) is equivalent to

$$\text{rank}(\nabla^2 s_3(\theta_*)) = \text{rank}(d_\eta B(\eta_*)). \quad (\text{D.40})$$

Let us group the columns and rows of  $d_\eta B$  into blocks indexed by  $(l, s)$ , where the  $(l, s)$  column block corresponds to  $d_{\eta^{(ls)}}$  and the  $(l, s)$  row block corresponds to  $B^{(ls)}$  as defined below in Lemma D.7. (These blocks  $B^{(ls)}$  are disjoint by definition, and we may discard the remaining rows of  $d_\eta B$  not belonging to any such block to produce a lower bound for its rank.) Ordering the pairs  $(l, s)$  as in Lemma D.7, the resulting matrix  $d_\eta B$  is block lower-triangular. Thus its rank is lower-bounded by the total rank of all blocks along the diagonal, i.e.

$$\text{rank}(d_\eta B(\eta_*)) \geq \sum_{l=0}^L \sum_{s=1}^{S_l} \text{rank}(d_{\eta^{(ls)}} B^{(ls)}(\eta_*)).$$

Lemma D.7 shows that for generic  $\eta_* \in \mathbb{R}^d$ ,

$$\begin{aligned} \sum_{l=2}^L \sum_{s=1}^{S_l} \text{rank}(d_{\eta^{(ls)}} B^{(ls)}(\eta_*)) &= \sum_{l=2}^L \sum_{s=1}^{S_l} (2l+1) = \sum_{l=2}^L (2l+1)S_l, \\ \sum_{s=1}^{S_1} \text{rank}(d_{\eta^{(1s)}} B^{(1s)}(\eta_*)) &\geq 1 + 2 + \sum_{s=3}^{S_1} 3 = 3S_1 - 3 \\ \sum_{s=1}^{S_0} \text{rank}(d_{\eta^{(0s)}} B^{(0s)}(\eta_*)) &= S_0. \end{aligned}$$

Combining these,

$$\text{trdeg } \mathcal{R}_{\leq 3}^G \geq \text{rank}(\nabla^2 s_3(\theta_*)) = \text{rank}(d_\eta B(\eta_*)) \geq \left( \sum_{l=0}^L (2l+1)S_l \right) - 3 = d - 3.$$

Thus  $\text{trdeg } \mathcal{R}_{\leq 3}^G = d - 3$ . □

**Lemma D.6.** For any  $l \geq 0$  and  $S \geq 2$ , consider  $\theta^{(1)}, \dots, \theta^{(S)} \in \mathbb{R}^{2l+1}$  and the Jacobian matrix of all their pairwise inner-products with respect to  $\theta = (\theta^{(1)}, \dots, \theta^{(S)}) \in \mathbb{R}^{(2l+1)S}$ ,

$$d_\theta[\langle \theta^{(s)}, \theta^{(s')} \rangle : 1 \leq s \leq s' \leq S] \in \mathbb{R}^{\frac{S(S+1)}{2} \times (2l+1)S}.$$

At generic  $\theta_* \in \mathbb{R}^{(2l+1)S}$ , this matrix has rank

$$\text{rank} \left( d_\theta[\langle \theta^{(s)}, \theta^{(s')} \rangle : 1 \leq s \leq s' \leq S] \Big|_{\theta=\theta_*} \right) = \begin{cases} \frac{S(S+1)}{2} & S < 2l+1 \\ (2l+1)(S-l) & S \geq 2l+1. \end{cases}$$

*Proof.* Let  $q = \min(2l+1, S)$  and consider the rows of the Jacobian for pairs  $(s, s')$  given by

$$(1, 1), (1, 2), \dots, (1, S), (2, 2), (2, 3), \dots, (2, S), \dots, (q, q), \dots, (q, S).$$

It may be checked that the number of such rows is exactly the desired formula for the rank. Consider  $\theta_*$  where  $\theta_{*,1}^{(1)} = \theta_{*,2}^{(2)} = \dots = \theta_{*,q}^{(q)} = 1$ , and all other coordinates are 0. For this  $\theta_*$ , the entries of the Jacobian are given by

$$\partial_{\theta_m^{(p)}}[\langle \theta^{(s)}, \theta^{(s')} \rangle] \Big|_{\theta=\theta_*} = \mathbf{1}\{p=s\} \cdot \theta_{*,m}^{(s')} + \mathbf{1}\{p=s'\} \cdot \theta_{*,m}^{(s)} = \mathbf{1}\{p=s, m=s'\} + \mathbf{1}\{p=s', m=s\}.$$

Thus for each row  $(s, s')$  above where  $s \leq q \leq 2l+1$  and  $s' \leq S$ , there are either 1 or 2 non-zero entries, in the column  $\partial_{\theta_s^{(s)}}$  and also in the column  $\partial_{\theta_{s'}^{(s)}}$  if  $s' \leq 2l+1$ . These columns are distinct for different rows  $(s, s')$ , so the submatrix of these columns has full row rank. This shows

$$\text{rank} \left( d_\theta[\langle \theta^{(s)}, \theta^{(s')} \rangle : 1 \leq s \leq s' \leq S] \Big|_{\theta=\theta_*} \right) \geq \begin{cases} \frac{S(S+1)}{2} & S < 2l+1 \\ (2l+1)(S-l) & S \geq 2l+1 \end{cases}$$

for this choice of  $\theta_*$ , and hence also at any generic  $\theta_*$  by Fact 4.5.

For the corresponding upper bound, for  $S < 2l+1$  this follows because the Jacobian has only  $S(S+1)/2$  rows. For  $S \geq 2l+1$ , consider the action of  $\text{SO}(2l+1)$  on  $\theta$  by simultaneous rotation of the vectors  $\theta^{(1)}, \dots, \theta^{(S)}$ . For  $S \geq 2l+1$  and at generic  $\theta_*$  where  $\theta_*^{(1)}, \dots, \theta_*^{(S)}$  span all of  $\mathbb{R}^{2l+1}$ , this action has trivial

stabilizer, so  $\dim(\mathcal{O}_{\theta_*}) = \dim(\mathrm{SO}(2l+1)) = (2l+1)l$  where  $\mathcal{O}_{\theta_*}$  is the orbit of  $\theta_*$  under this action. Since  $\langle \theta^{(s)}, \theta^{(s')} \rangle$  is constant on  $\mathcal{O}_{\theta_*}$ , for each vector  $v$  in the dimension- $(2l+1)l$  tangent space to  $\mathcal{O}_{\theta_*}$ , we have

$$d_\theta[\langle \theta^{(s)}, \theta^{(s')} \rangle : 1 \leq s \leq s' \leq S] \Big|_{\theta=\theta_*} \cdot v = 0.$$

Thus the dimension of the row span of the Jacobian is at most  $(2l+1)S - (2l+1)l = (2l+1)(S-l)$ , as desired.  $\square$

**Lemma D.7.** *Suppose  $L \geq 1$  and  $S_0, \dots, S_L \geq 2$ . Order the pairs  $(l, s)$  by  $(l, s) < (l', s')$  if  $l < l'$  or if  $l = l'$  and  $s < s'$ . Fix any  $l \in \{1, \dots, L\}$  and  $s \in \{1, \dots, S_l\}$ , and let  $\mathcal{K}^{(ls)}$  be the set of tuples  $((l, s), (l', s'), (l'', s'')) \in \mathcal{K}$  where  $(l, s)$  take these fixed values, and  $(l', s') \leq (l, s)$  and  $(l'', s'') \leq (l, s)$ . Denote*

$$B^{(ls)}(\eta) = \left( B_{(l,s),(l',s'),(l'',s'')}(\eta) : ((l, s), (l', s'), (l'', s'')) \in \mathcal{K}^{(ls)} \right) \in \mathbb{R}^{|\mathcal{K}^{(ls)}|}$$

and let  $d_{\eta^{(ls)}} B^{(ls)} \in \mathbb{R}^{|\mathcal{K}^{(ls)}| \times (2l+1)}$  be its Jacobian in  $\eta^{(ls)}$ . Then for any generic  $\eta_* \in \mathbb{R}^d$ :

- (a) If  $l \geq 2$ , then  $\mathrm{rank}(d_{\eta^{(ls)}} B^{(ls)}(\eta_*)) = 2l + 1$ .
- (b) If  $l \geq 1$  and  $s \geq 3$ , then also  $\mathrm{rank}(d_{\eta^{(ls)}} B^{(ls)}(\eta_*)) = 2l + 1 = 3$ . Furthermore for  $s = 2$ , removing  $w_0^{(12)}$  from  $\eta^{(12)}$ , we have  $\mathrm{rank}(d_{v_1^{(12)}, w_1^{(12)}} B^{(12)}(\eta_*)) = 2$ . For  $s = 1$ , removing  $w_0^{(11)}, w_1^{(11)}$  from  $\eta^{(11)}$ , we have  $\mathrm{rank}(d_{v_1^{(11)}} B^{(11)}(\eta_*)) = 1$ .
- (c) If  $l = 0$ , then  $\mathrm{rank}(d_{\eta^{(0s)}} B^{(0s)}(\eta_*)) = 1$ .

*Proof.* The strategy is similar to the proof of Lemma D.4. For each statement, by Fact 4.5, it suffices to exhibit a single point  $\eta_* \in \mathbb{R}^d$  where the rank equality holds. We choose  $\eta_*$  having most coordinates 0, to allow an explicit computation of the rank.

Recall  $B_{(l,s),(l',s'),(l'',s'')}$  from (4.9). We first compute  $d_{\eta^{(ls)}} B_{(l,s),(l',s'),(l'',s'')}$ : If  $(l', s'), (l'', s'') < (l, s)$  strictly, then the derivative  $d_{\eta^{(ls)}}$  applies to only the term  $\overline{u_m^{(ls)}}$  in (4.9). A computation analogous to (D.20–D.21) using the sign symmetry (D.29) shows, for  $m > 0$  strictly,

$$\partial_{v_m^{(ls)}} B_{(l,s),(l',s'),(l'',s'')} = \sum_{m'=-l'}^{l'} 2C_{m,m',m+m'}^{l,l',l''} \cdot \mathrm{Re} \overline{u_{m'}^{(l's')}} u_{m+m'}^{(l''s'')} \quad (\text{D.41})$$

$$\partial_{w_m^{(ls)}} B_{(l,s),(l',s'),(l'',s'')} = \sum_{m'=-l'}^{l'} 2C_{m,m',m+m'}^{l,l',l''} \cdot \mathrm{Im} \overline{u_{m'}^{(l's')}} u_{m+m'}^{(l''s'')} \quad (\text{D.42})$$

For  $m = 0$ , recalling that  $\eta_0^{(ls)} = v_0^{(ls)}$  if  $l$  is even and  $\eta_0^{(ls)} = w_0^{(ls)}$  if  $l$  is odd, we also have

$$\partial_{\eta_0^{(ls)}} B_{(l,s),(l',s'),(l'',s'')} = \sum_{m'=-l'}^{l'} C_{0,m',m'}^{l,l',l''} \overline{u_{m'}^{(l's')}} u_{m'}^{(l''s'')} \times \begin{cases} 1 & \text{if } l \text{ is even} \\ -\mathbf{i} & \text{if } l \text{ is odd.} \end{cases} \quad (\text{D.43})$$

If  $(l', s') = (l, s)$  and  $(l'', s'') < (l, s)$  strictly, an additional contribution to the derivatives arise from differentiating  $u_{m'}^{(l's')}$ . This doubles the above expressions, and we have

$$\partial_{v_m^{(ls)}} B_{(l,s),(l,s),(l'',s'')} = \sum_{m'=-l}^l 4C_{m,m',m+m'}^{l,l,l''} \cdot \mathrm{Re} \overline{u_{m'}^{(ls)}} u_{m+m'}^{(l''s'')} \quad (\text{D.44})$$

$$\partial_{w_m^{(ls)}} B_{(l,s),(l,s),(l'',s'')} = \sum_{m'=-l}^l 4C_{m,m',m+m'}^{l,l,l''} \cdot \mathrm{Im} \overline{u_{m'}^{(ls)}} u_{m+m'}^{(l''s'')} \quad (\text{D.45})$$

$$\partial_{\eta_0^{(ls)}} B_{(l,s),(l,s),(l'',s'')} = \sum_{m'=-l}^l 2C_{0,m',m'}^{l,l,l''} \overline{u_{m'}^{(ls)}} u_{m'}^{(l''s'')} \times \begin{cases} 1 & \text{if } l \text{ is even} \\ -\mathbf{i} & \text{if } l \text{ is odd.} \end{cases} \quad (\text{D.46})$$

We now use a different construction of  $\eta_*$  for different values of  $(l, s)$ :

**Part (a),  $l \geq 4$ :** Let us fix two radial frequencies  $(A, B) = (1, 2)$ . (We use here the condition  $S_0, \dots, S_L \geq 2$ , so that these frequencies exist for each  $l = 0, \dots, L$ .) We choose  $\eta_*$  such that for all  $l' \in \{1, \dots, l-1\}$  and  $m' \in \{0, \dots, l'\}$ ,

$$v_{*,m'}^{(l'A)} = w_{*,m'}^{(l'A)} = 0 \text{ unless } m' = l', \quad v_{*,m'}^{(l'B)} = w_{*,m'}^{(l'B)} = 0 \text{ unless } m' = l' - 1.$$

We choose the non-zero coordinates of  $\eta_*$  to be generic. Using similar notation as in Lemma D.4, we write this as

$$\text{Type}(l', A) = 0, \quad \text{Type}(l', B) = 1 \quad \text{for all } l' = 1, \dots, l-1,$$

where  $\text{Type}(l', s') = i$  indicates that  $v_{*,m}^{(l's')}, w_{*,m}^{(l's')} = 0$  unless  $m = l' - i$ . (In contrast to Lemma D.4, here  $\text{Type}(\cdot)$  is a single integer rather than a set.) For  $\partial_{v_m^{(ls)}} B_{(l,s),(l',s')}, \partial_{w_m^{(ls)}} B_{(l,s),(l',s')}, \partial_{v_m^{(ls)}} B_{(l,s),(l'',s'')}, \partial_{w_m^{(ls)}} B_{(l,s),(l'',s'')}$  to be non-zero, we require  $|m + m'| = l'' - \text{Type}(l'', s'')$  and  $|m'| = l' - \text{Type}(l', s')$ . This requires analogously to (D.26)

$$m \in \left\{ |(l' - \text{Type}(l', s')) - (l'' - \text{Type}(l'', s''))|, (l' - \text{Type}(l', s')) + (l'' - \text{Type}(l'', s'')) \right\}. \quad (\text{D.47})$$

Rows of  $d_{\eta^{(ls)}} B^{(ls)}(\eta_*)$  may be indexed by  $(l', s'), (l'', s'')$  for which  $((l, s), (l', s'), (l'', s'')) \in \mathcal{K}^{(ls)}$ . We select  $2l + 1$  such rows, given by the left column of the following Table 3. Note that when  $l \geq 4$ , these rows satisfy the requirement  $l' + l'' \geq l$  in the definition of  $\mathcal{K}$ . For each row, the right column indicates the values of  $m$  satisfying (D.47), for which  $\partial_{v_m^{(ls)}}, \partial_{w_m^{(ls)}}$  are non-zero.

TABLE 3.

$(l', s') \text{ and } (l'', s'')$	Values of $m$
$(l-1, A) \text{ and } (l-1, A) \text{ if } l \text{ is even; } (l-2, A) \text{ and } (l-1, B) \text{ if } l \text{ is odd}$	0
$(l-1, B) \text{ and } (l-1, A)$	1
$(l-2, A) \text{ and } (l-1, A)$	1
$(l-2, B) \text{ and } (l-1, A)$	2
$(l-3, A) \text{ and } (l-1, A)$	2
$\vdots$	$\vdots$
$(3, B) \text{ and } (l-1, A)$	$l-3$
$(2, A) \text{ and } (l-1, A)$	$l-3$
$(3, B) \text{ and } (l-1, B)$	$l-4 \text{ and } l$
$(2, A) \text{ and } (l-1, B)$	$l-4 \text{ and } l$
$(2, B) \text{ and } (l-1, B)$	$l-3 \text{ and } l-1$
$(1, A) \text{ and } (l-1, B)$	$l-3 \text{ and } l-1$
$(2, B) \text{ and } (l-1, A)$	$l \text{ and } l-2$
$(1, A) \text{ and } (l-1, A)$	$l \text{ and } l-2$

To verify that this selected  $(2l+1) \times (2l+1)$  submatrix of  $d_{\eta^{(ls)}} B^{(ls)}(\eta_*)$  is non-singular, we order its rows as in Table 3, and its columns by the ordering of variables

$$\eta_0^{(ls)}, v_1^{(ls)}, w_1^{(ls)}, \dots, v_{l-3}^{(ls)}, w_{l-3}^{(ls)}, v_l^{(ls)}, w_l^{(ls)}, v_{l-1}^{(ls)}, w_{l-1}^{(ls)}, v_{l-2}^{(ls)}, w_{l-2}^{(ls)}$$

as they appear in the right column above. Then this  $(2l+1) \times (2l+1)$  submatrix is block lower-triangular in the decomposition  $2l+1 = 1 + 2 + 2 + \dots + 2$ . It suffices to check that each  $1 \times 1$  and  $2 \times 2$  diagonal block is non-singular.

**Block corresponding to  $\eta_0^{(l)}$ :** Applying (D.43) and the symmetries (D.10) and (D.29), the first  $1 \times 1$  diagonal block is

$$\begin{aligned} \partial_{\eta_0}^{(ls)} B_{(l,s),(l-1,A),(l-1,A)}(\eta_*) &= 2C_{0,l-1,l-1}^{l,l-1,l-1} \cdot |u_{*,l-1}^{(l-1,A)}|^2 \text{ for even } l \\ \partial_{\eta_0}^{(ls)} B_{(l,s),(l-2,A),(l-1,B)}(\eta_*) &= 2C_{0,l-2,l-2}^{l,l-2,l-1} \cdot \text{Im} \overline{u_{*,l-2}^{(l-2,A)}} u_{*,l-2}^{(l-1,B)} \text{ for odd } l. \end{aligned}$$

By Lemma D.2, these Clebsch-Gordan coefficients are non-zero, so this block is generically non-zero.

**Blocks corresponding to  $(v_1^{(ls)}, w_1^{(ls)}), \dots, (v_{l-3}^{(ls)}, w_{l-3}^{(ls)})$ :** The arguments for these blocks are similar, so we consider only  $v_{l-3}^{(ls)}, w_{l-3}^{(ls)}$ . Applying (D.41–D.42), this  $2 \times 2$  block is

$$\begin{aligned} & \partial_{v_{l-3}^{(ls)}, w_{l-3}^{(ls)}} \left( B_{(l,s),(3,B),(l-1,A)}, B_{(l,s),(2,A),(l-1,A)} \right) (\eta_*) \\ &= \begin{pmatrix} 2C_{l-3,2,l-1}^{l,3,l-1} \cdot \operatorname{Re} \overline{u_{*,2}^{(3,B)}} u_{*,l-1}^{(l-1,A)} & 2C_{l-3,2,l-1}^{l,3,l-1} \cdot \operatorname{Im} \overline{u_{*,2}^{(3,B)}} u_{*,l-1}^{(l-1,A)} \\ 2C_{l-3,2,l-1}^{l,2,l-1} \cdot \operatorname{Re} \overline{u_{*,2}^{(2,A)}} u_{*,l-1}^{(l-1,A)} & 2C_{l-3,2,l-1}^{l,2,l-1} \cdot \operatorname{Im} \overline{u_{*,2}^{(2,A)}} u_{*,l-1}^{(l-1,A)} \end{pmatrix} \end{aligned}$$

These Clebsch-Gordan coefficients are again non-zero by Lemma D.2, so the determinant of this matrix is a non-zero polynomial in the six coefficients of  $\eta_*$

$$v_{*,2}^{(3,B)}, w_{*,2}^{(3,B)}, v_{*,2}^{(2,A)}, w_{*,2}^{(2,A)}, v_{*,l-1}^{(l-1,A)}, w_{*,l-1}^{(l-1,A)}.$$

(These coefficients are distinct for  $l \geq 4$ .) Hence this determinant is generically non-zero.

**Blocks corresponding to  $(v_l^{(ls)}, w_l^{(ls)}), (v_{l-1}^{(ls)}, w_{l-1}^{(ls)}), (v_{l-2}^{(ls)}, w_{l-2}^{(ls)})$ :** Applying (D.41–D.42), these  $2 \times 2$  blocks are

$$\begin{aligned} & \partial_{v_l^{(ls)}, w_l^{(ls)}} \left( B_{(l,s),(3,B),(l-1,B)}, B_{(l,s),(2,A),(l-1,B)} \right) (\eta_*) \\ &= \begin{pmatrix} 2C_{l-2,l-2}^{l,3,l-1} \cdot \operatorname{Re} \overline{u_{*,2}^{(3,B)}} u_{*,l-2}^{(l-1,B)} & 2C_{l-2,l-2}^{l,3,l-1} \cdot \operatorname{Im} \overline{u_{*,2}^{(3,B)}} u_{*,l-2}^{(l-1,B)} \\ 2C_{l-2,l-2}^{l,2,l-1} \cdot \operatorname{Re} \overline{u_{*,2}^{(2,A)}} u_{*,l-2}^{(l-1,B)} & 2C_{l-2,l-2}^{l,2,l-1} \cdot \operatorname{Im} \overline{u_{*,2}^{(2,A)}} u_{*,l-2}^{(l-1,B)} \end{pmatrix} \\ & \partial_{v_{l-1}^{(ls)}, w_{l-1}^{(ls)}} \left( B_{(l,s),(2,B),(l-1,B)}, B_{(l,s),(1,A),(l-1,B)} \right) (\eta_*) \\ &= \begin{pmatrix} 2C_{l-1,-1,l-2}^{l,2,l-1} \cdot \operatorname{Re} \overline{u_{*,1}^{(2,B)}} u_{*,l-2}^{(l-1,B)} & 2C_{l-1,-1,l-2}^{l,2,l-1} \cdot \operatorname{Im} \overline{u_{*,1}^{(2,B)}} u_{*,l-2}^{(l-1,B)} \\ 2C_{l-1,-1,l-2}^{l,1,l-1} \cdot \operatorname{Re} \overline{u_{*,1}^{(1,A)}} u_{*,l-2}^{(l-1,B)} & 2C_{l-1,-1,l-2}^{l,1,l-1} \cdot \operatorname{Im} \overline{u_{*,1}^{(1,A)}} u_{*,l-2}^{(l-1,B)} \end{pmatrix} \\ & \partial_{v_{l-2}^{(ls)}, w_{l-2}^{(ls)}} \left( B_{(l,s),(2,B),(l-1,A)}, B_{(l,s),(1,A),(l-1,A)} \right) (\eta_*) \\ &= \begin{pmatrix} 2C_{l-2,1,l-1}^{l,2,l-1} \cdot \operatorname{Re} \overline{u_{*,1}^{(2,B)}} u_{*,l-1}^{(l-1,A)} & 2C_{l-2,1,l-1}^{l,2,l-1} \cdot \operatorname{Im} \overline{u_{*,1}^{(2,B)}} u_{*,l-1}^{(l-1,A)} \\ 2C_{l-2,1,l-1}^{l,1,l-1} \cdot \operatorname{Re} \overline{u_{*,1}^{(1,A)}} u_{*,l-1}^{(l-1,A)} & 2C_{l-2,1,l-1}^{l,1,l-1} \cdot \operatorname{Im} \overline{u_{*,1}^{(1,A)}} u_{*,l-1}^{(l-1,A)} \end{pmatrix} \end{aligned}$$

The Clebsch-Gordan coefficients here are non-zero by Lemma D.2 (for the term  $C_{l-1,-1,l-2}^{l,2,l-1}$  of the second matrix, this uses the condition  $2 \neq l-1$  when  $l \geq 4$ ). Then the determinants of all three matrices are non-zero polynomials of six distinct coordinates of  $\eta_*$ , except in the case of the first matrix for  $l = 4$ . In this case,  $u_2^{(3,B)}$  and  $u_{l-2}^{(l-1,B)}$  coincide, and the determinant may be checked to be a non-zero polynomial of the four distinct coordinates  $v_{*,2}^{(3,B)}, w_{*,2}^{(3,B)}, v_{*,2}^{(2,A)}, w_{*,2}^{(2,A)}$ .

Combining these cases shows that  $d_{\eta^{(ls)}} B^{(ls)}(\eta_*)$  has full column rank  $2l+1$  as desired.

**Part (a),  $l = 3$ :** We again fix  $(A, B) = (1, 2)$ , and specialize to a point  $\eta_*$  such that

$$\operatorname{Type}(3, A), \operatorname{Type}(2, A), \operatorname{Type}(1, A) = 0, \quad \operatorname{Type}(2, B), \operatorname{Type}(1, B) = 1.$$

We then pick 7 rows of  $d_{\eta^{(3s)}} B^{(3s)}(\eta_*)$ , indicated by the left column of the below table. Applying (D.47), the derivatives in  $(v_m^{(3s)}, w_m^{(3s)})$  are non-zero for only the values of  $m$  in the right column.

TABLE 4.

$(l', s')$ and $(l'', s'')$	Values of $m$
$(1, B)$ and $(2, B)$	1
$(3, A)$ and $(2, A)$	1
$(1, B)$ and $(2, A)$	2
$(3, A)$ and $(1, A)$	2
$(1, A)$ and $(2, A)$	1 and 3
$(2, B)$ and $(2, A)$	1 and 3
$(1, A)$ and $(2, B)$	2 and 0

Ordering the columns by  $v_1^{(3s)}, w_1^{(3s)}, v_2^{(3s)}, w_2^{(3s)}, v_3^{(3s)}, w_3^{(3s)}, w_0^{(3s)}$ , this  $7 \times 7$  submatrix has a block lower-triangular structure in the decomposition  $7 = 2 + 2 + 2 + 1$ . If  $s \neq A$ , then applying (D.41–D.43) and (D.44–D.45), its diagonal blocks are given explicitly by

$$\begin{aligned}
\partial_{v_1^{(3s)}, w_1^{(3s)}} \left( B_{(3,s),(1,B),(2,B)}, B_{(3,s),(3,A),(2,A)} \right) (\eta_*) &= \begin{pmatrix} 2C_{1,0,1}^{3,1,2} \cdot \text{Re } \overline{u_{*,0}^{(1B)}} u_{*,1}^{(2B)} & 2C_{1,0,1}^{3,1,2} \cdot \text{Im } \overline{u_{*,0}^{(1B)}} u_{*,1}^{(2B)} \\ 2C_{1,-3,-2}^{3,3,2} \cdot \text{Re } \overline{u_{*, -3}^{(3A)}} u_{*, -2}^{(2A)} & 2C_{1,-3,-2}^{3,3,2} \cdot \text{Im } \overline{u_{*, -3}^{(3A)}} u_{*, -2}^{(2A)} \end{pmatrix} \\
\partial_{v_2^{(3s)}, w_2^{(3s)}} \left( B_{(3,s),(1,B),(2,A)}, B_{(3,s),(3,A),(1,A)} \right) (\eta_*) &= \begin{pmatrix} 2C_{2,0,2}^{3,1,2} \cdot \text{Re } \overline{u_{*,0}^{(1B)}} u_{*,2}^{(2A)} & 2C_{2,0,2}^{3,1,2} \cdot \text{Im } \overline{u_{*,0}^{(1B)}} u_{*,2}^{(2A)} \\ 2C_{2,-3,-1}^{3,3,1} \cdot \text{Re } \overline{u_{*, -3}^{(3A)}} u_{*, -1}^{(1A)} & 2C_{2,-3,-1}^{3,3,1} \cdot \text{Im } \overline{u_{*, -3}^{(3A)}} u_{*, -1}^{(1A)} \end{pmatrix} \\
\partial_{v_3^{(3s)}, w_3^{(3s)}} \left( B_{(3,s),(1,A),(2,A)}, B_{(3,s),(2,B),(2,A)} \right) (\eta_*) &= \begin{pmatrix} 2C_{3,-1,2}^{3,1,2} \cdot \text{Re } \overline{u_{*, -1}^{(1A)}} u_{*,2}^{(2A)} & 2C_{3,-1,2}^{3,1,2} \cdot \text{Im } \overline{u_{*, -1}^{(1A)}} u_{*,2}^{(2A)} \\ 2C_{3,-1,2}^{3,2,2} \cdot \text{Re } \overline{u_{*, -1}^{(2B)}} u_{*,2}^{(2A)} & 2C_{3,-1,2}^{3,2,2} \cdot \text{Im } \overline{u_{*, -1}^{(2B)}} u_{*,2}^{(2A)} \end{pmatrix} \\
\partial_{w_0^{(3s)}} B_{(3,s),(1,A),(2,B)} (\eta_*) &= 2C_{0,1,1}^{3,1,2} \cdot \text{Im } \overline{u_{*,1}^{1A}} u_{*,1}^{2B}
\end{aligned}$$

Here  $u_{*,0}^{(1B)} = i w_{*,0}^{(1B)}$  depends on only one rather than two non-zero coordinate of  $\eta_*$ ; nonetheless, one may still check that the determinants of the above three matrices are generically non-zero. If  $s = A$ , then the second rows of the first two matrices above have coefficients 4 instead of 2, from applying (D.44–D.45) in place of (D.41–D.42), but this does not affect their ranks. Thus these blocks are generically non-singular, so  $d_{\eta^{(3s)}} B^{(3s)}(\eta_*)$  has full column rank 7.

**Part (a),  $l = 2$ :** We specialize to a point  $\eta_*$  such that

$$\text{Type}(2, A), \text{Type}(1, A), \text{Type}(0, A) = 0, \quad \text{Type}(2, B), \text{Type}(1, B) = 1.$$

(Here  $\text{Type}(0, A) = 0$  means simply that  $v_{*,0}^{(0A)}$  is non-zero.) We pick the following 5 rows of  $d_{\eta^{(2s)}} B^{(2s)}(\eta_*)$ , for which the derivatives in  $(v_m^{(2s)}, w_m^{(2s)})$  are non-zero for only the following corresponding values of  $m$ .

TABLE 5.

$(l', s')$ and $(l'', s'')$	Values of $m$
$(1, B)$ and $(1, B)$	0
$(1, B)$ and $(1, A)$	1
$(2, B)$ and $(0, A)$	1
$(2, A)$ and $(0, A)$	2
$(1, A)$ and $(1, A)$	0 and 2

Ordering the columns by  $v_0^{(2s)}, v_1^{(2s)}, w_1^{(2s)}, v_2^{(2s)}, w_2^{(2s)}$ , this  $5 \times 5$  submatrix has a block lower-triangular structure in the decomposition  $5 = 1 + 2 + 2$ . If  $s \notin \{A, B\}$ , these blocks are

$$\begin{aligned}
\partial_{v_0^{(2s)}} B_{(2,s),(1,B),(1,B)} (\eta_*) &= C_{0,0,0}^{2,1,1} \cdot \left| u_{*,0}^{(1B)} \right|^2 \\
\partial_{v_1^{(2s)}, w_1^{(2s)}} \left( B_{(2,s),(1,B),(1,A)}, B_{(2,s),(2,B),(0,A)} \right) (\eta_*) &= \begin{pmatrix} 2C_{1,0,1}^{2,1,1} \cdot \text{Re } \overline{u_{*,0}^{(1B)}} u_{*,1}^{(1A)} & 2C_{1,0,1}^{2,1,1} \cdot \text{Im } \overline{u_{*,0}^{(1B)}} u_{*,1}^{(1A)} \\ 2C_{1,-1,0}^{2,2,0} \cdot \text{Re } \overline{u_{*, -1}^{(2B)}} u_{*,0}^{(0A)} & 2C_{1,-1,0}^{2,2,0} \cdot \text{Im } \overline{u_{*, -1}^{(2B)}} u_{*,0}^{(0A)} \end{pmatrix} \\
\partial_{v_2^{(2s)}, w_2^{(2s)}} \left( B_{(2,s),(2,A),(0,A)}, B_{(2,s),(1,A),(1,A)} \right) (\eta_*) &= \begin{pmatrix} 2C_{2,-2,0}^{2,2,0} \cdot \text{Re } \overline{u_{*, -2}^{(2A)}} u_{*,0}^{(0A)} & 2C_{2,-2,0}^{2,2,0} \cdot \text{Im } \overline{u_{*, -2}^{(2A)}} u_{*,0}^{(0A)} \\ 2C_{2,-1,1}^{2,1,1} \cdot \text{Re } \overline{u_{*, -1}^{(1A)}} u_{*,1}^{(1A)} & 2C_{2,-1,1}^{2,1,1} \cdot \text{Im } \overline{u_{*, -1}^{(1A)}} u_{*,1}^{(1A)} \end{pmatrix}
\end{aligned}$$

If  $s = A$  or  $s = B$ , then the first row of the third matrix or second row of the second matrix should have coefficient 4 in place of 2, but this does not affect their ranks. These blocks are generically non-singular, so  $d_{\eta^{(2s)}} B^{(2s)}(\eta_*)$  has full column rank 5.

**Part (b),  $l = 1, s \geq 3$ :** Note that  $(A, B, s) = (1, 2, s)$  are distinct indices because  $s \geq 3$ . We specialize to a point  $\eta_*$  such that

$$\text{Type}(1, s), \text{Type}(1, A), \text{Type}(0, A), \text{Type}(0, B) = 0, \quad \text{Type}(1, B) = 1.$$

We pick the following 3 rows of  $d_{\eta^{(1s)}} B^{(1s)}(\eta_*)$ , for which the derivatives in  $(v_m^{(1s)}, w_m^{(1s)})$  are non-zero for only the following corresponding values of  $m$ .

TABLE 6.

$(l', s')$ and $(l'', s'')$	Values of $m$
$(1, s)$ and $(0, A)$	1
$(1, A)$ and $(0, B)$	1
$(1, B)$ and $(0, B)$	0

Ordering the columns by  $v_1^{(1s)}, w_1^{(1s)}, w_0^{(1s)}$ , this  $3 \times 3$  submatrix has a block lower-triangular structure in the decomposition  $3 = 2 + 1$ , with diagonal blocks

$$\begin{aligned} \partial_{v_1^{(1s)}, w_1^{(1s)}} \left( B_{(1,s),(1,s),(0,A)}, B_{(1,s),(1,A),(0,B)} \right) (\eta_*) &= \begin{pmatrix} 4C_{1,-1,0}^{1,1,0} \cdot \text{Re } \overline{u_{*, -1}^{(1s)}} u_{*, 0}^{(0A)} & 4C_{1,-1,0}^{1,1,0} \cdot \text{Im } \overline{u_{*, -1}^{(1s)}} u_{*, 0}^{(0A)} \\ 2C_{1,-1,0}^{1,1,0} \cdot \text{Re } \overline{u_{*, -1}^{(1A)}} u_{*, 0}^{(0B)} & 2C_{1,-1,0}^{1,1,0} \cdot \text{Im } \overline{u_{*, -1}^{(1A)}} u_{*, 0}^{(0B)} \end{pmatrix} \\ \partial_{w_0^{(1s)}} B_{(1,s),(1,B),(0,B)} (\eta_*) &= -i C_{0,0,0}^{1,1,0} \cdot \overline{u_{*, 0}^{(1B)}} u_{*, 0}^{(0A)} \end{aligned}$$

These blocks are generically non-singular (where we use that  $s$  and  $A$  are distinct for the first block), so  $d_{\eta^{(1s)}} B^{(1s)}(\eta_*)$  has full column rank 3.

**Part (b),  $l = 1, s = 2$ :** Consider the two columns of  $d_{\eta^{(12)}} B^{(12)}$  corresponding to  $\partial_{v_1^{(12)}, w_1^{(12)}}$  and the two rows corresponding to  $B_{(1,2),(1,1),(0,1)}, B_{(1,2),(1,2),(0,1)}$ . This  $2 \times 2$  submatrix is (for any  $\eta_*$ )

$$\partial_{v_1^{(12)}, w_1^{(12)}} \left( B_{(1,2),(1,1),(0,1)}, B_{(1,2),(1,2),(0,1)} \right) (\eta_*) = \begin{pmatrix} 2C_{1,-1,0}^{1,1,0} \cdot \text{Re } \overline{u_{*, -1}^{(11)}} u_{*, 0}^{(01)} & 2C_{1,-1,0}^{1,1,0} \cdot \text{Im } \overline{u_{*, -1}^{(11)}} u_{*, 0}^{(01)} \\ 4C_{1,-1,0}^{1,1,0} \cdot \text{Re } \overline{u_{*, -1}^{(12)}} u_{*, 0}^{(01)} & 4C_{1,-1,0}^{1,1,0} \cdot \text{Im } \overline{u_{*, -1}^{(12)}} u_{*, 0}^{(01)} \end{pmatrix}$$

This is generically non-singular, so  $d_{v_1^{(12)}, w_1^{(12)}} B^{(12)}(\eta_*)$  has full column rank 2.

**Part (b),  $l = 1, s = 1$ :** Consider the column and row of  $d_{\eta^{(11)}} B^{(11)}$  corresponding to

$$\partial_{v_1^{(11)}} B_{(1,1),(1,1),(0,1)} (\eta_*) = 4C_{1,-1,0}^{1,1,0} \text{Re } \overline{u_{*, -1}^{(11)}} u_{*, 0}^{(01)}.$$

This is generically non-zero, so  $d_{v_1^{(11)}} B^{(11)}(\eta_*)$  has full column rank 1.

**Part (c),  $l = 0$ :** For any  $s \in \{1, \dots, S_0\}$ ,  $\eta^{(0s)} = v_0^{(0s)}$  is a single real variable. Applying  $\langle 0, 0; 0, 0 | 0, 0 \rangle = 1$ , we have from (4.9) that

$$\partial_{\eta^{(0s)}} B_{(0,s),(0,s),(0,s)} (\eta_*) = \partial_{v_0^{(0s)}} (v_{*, 0}^{(0s)})^3 = 3(v_{*, 0}^{(0s)})^2,$$

which is generically non-zero. Thus  $d_{\eta^{(0s)}} B^{(0s)}(\eta_*)$  has full column rank 1.  $\square$

#### D.4. Projected cryo-EM.

D.4.1. *Function bases.* We describe in further detail the function bases and the forms of the projection map and rotational action on the basis coefficients in the projected cryo-EM model of Section 4.3, following a setup of the model that is similar to that of [BBSK<sup>+</sup>17, Section 5.5 and Appendix A.4].

Parametrizing  $\mathbb{R}^3$  by spherical coordinates  $(\rho, \phi_1, \phi_2)$ , define a product basis

$$\hat{j}_{ism}(\rho, \phi_1, \phi_2) = \tilde{z}_s(\rho) y_{lm}(\phi_1, \phi_2) \quad \text{for } s \geq 1, \quad l \geq 0, \quad m \in \{-l, \dots, l\} \quad (\text{D.48})$$

where  $\{y_{lm}\}$  are the complex spherical harmonics (D.2), and  $\{\tilde{z}_s : s \geq 1\}$  is a system of radial basis functions  $\tilde{z}_s : [0, \infty) \rightarrow \mathbb{R}$ . Similarly, parametrizing  $\mathbb{R}^2$  by polar coordinates  $(\rho, \phi_2)$  where  $\rho \geq 0$  is the radius and  $\phi_2 \in [0, 2\pi)$  is the angle, let

$$b_m(\phi_2) = (2\pi)^{-1/2} e^{im\phi_2}$$

and define a corresponding product basis

$$\hat{j}_{sm}(\rho, \phi_2) = \tilde{z}_s(\rho) b_m(\phi_2) \quad \text{for } s \geq 1, \quad m \in \mathbb{Z}. \quad (\text{D.49})$$

We choose the radial basis functions  $\{\tilde{z}_s : s \geq 1\}$  to satisfy a modified orthogonality relation

$$\int_0^\infty \rho \tilde{z}_s(\rho) \tilde{z}_{s'}(\rho) d\rho = \mathbf{1}\{s = s'\} \quad (\text{D.50})$$

with weight  $\rho$  instead of  $\rho^2$ , which ensures that  $\{\hat{j}_{sm}\}$  are orthonormal in  $L_2(\mathbb{R}^2, \mathbb{C})$ . We write  $\{j_{lsm}\}$  for the 3-D inverse Fourier transform of  $\{\hat{j}_{lsm}\}$ , and  $\{j_{sm}\}$  for the 2-D inverse Fourier transform of  $\{\hat{j}_{sm}\}$ .

For any function  $f \in L_2(\mathbb{R}^3)$ , let  $\hat{f}$  be its Fourier transform as defined in (D.27), and let

$$\widehat{\Pi \cdot f}(k_1, k_2) = \int_{\mathbb{R}^2} e^{-2\pi i(k_1 x_1 + k_2 x_2)} (\Pi \cdot f)(x_1, x_2) dx_1 dx_2$$

be the 2-D Fourier transform of its tomographic projection. By the Fourier-slice relation,

$$\widehat{\Pi \cdot f}(k_1, k_2) = \hat{f}(k_1, k_2, 0). \quad (\text{D.51})$$

In spherical coordinates  $(\rho, \phi_1, \phi_2)$  for  $\mathbb{R}^3$  and polar coordinates  $(\rho, \phi_2)$  for  $\mathbb{R}^2$ , this corresponds to

$$\widehat{\Pi \cdot f}(\rho, \phi_2) = \hat{f}(\rho, \pi/2, \phi_2)$$

where we restrict  $\phi_1 = \pi/2$ . This restriction of each complex spherical harmonic  $y_{lm}$  in (D.2) is given by

$$y_{lm}(\pi/2, \phi_2) = p_{lm} \cdot b_m(\phi_2)$$

where  $b_m$  is the function defined above, and

$$\begin{aligned} p_{lm} &= (-1)^m \sqrt{\frac{(2l+1)(l-m)!}{2(l+m)!}} \cdot P_{lm}(0) \\ &= \mathbf{1}\{l+m \text{ is even}\} \times \frac{(-1)^{(l+m)/2} \sqrt{(2l+1)/2}}{2^l l!} \binom{l}{(l+m)/2} \sqrt{(l-m)!(l+m)!} \end{aligned} \quad (\text{D.52})$$

the second equality applying Lemma D.1. Note that these coefficients satisfy a sign symmetry

$$p_{lm} = (-1)^m p_{l, -m}. \quad (\text{D.53})$$

Specializing (D.51) to  $f = j_{lsm}$ , we then have

$$\widehat{\Pi \cdot j_{lsm}}(\rho, \phi_2) = p_{lm} \cdot \tilde{z}_s(\rho) b_m(\phi_2) = p_{lm} \cdot \hat{j}_{sm}(\rho, \phi_2).$$

Then taking inverse Fourier transforms,

$$\Pi \cdot j_{lsm} = p_{lm} \cdot j_{sm}. \quad (\text{D.54})$$

We consider the space of functions  $f : \mathbb{R}^3 \rightarrow \mathbb{R}$  that are  $(L, S_0, \dots, S_L)$ -bandlimited in the basis  $\{j_{lsm}\}$ , admitting the first representation in (4.12),

$$f = \sum_{(l,s,m) \in \mathcal{I}} u_m^{(ls)} \cdot j_{lsm}$$

where the index set  $\mathcal{I}$  is defined in (4.5). Then (D.54) shows that  $\Pi \cdot f$  is also bandlimited in the basis  $\{j_{sm}\}$ , admitting the first representation in (4.13),

$$\Pi \cdot f = \sum_{(s,m) \in \tilde{\mathcal{I}}} \tilde{u}_m^{(s)} \cdot j_{sm}$$

where the index set  $\tilde{\mathcal{I}}$  is defined in (4.14). Define a real basis  $\{h_{lsm}\}$  from  $\{j_{lsm}\}$  by (D.30), so that the coefficients  $u \in \mathbb{C}^d$  for the former and  $\theta \in \mathbb{R}^d$  for the latter are related by the unitary transform  $u = \hat{V}^* \theta$  in (D.31). Similarly, define a real basis  $\{h_{sm}\}$  from  $\{j_{sm}\}$  by

$$h_{sm} = \begin{cases} \frac{1}{\sqrt{2}}(j_{s,-m} + (-1)^m j_{sm}) & \text{if } m > 0 \\ j_{s0} & \text{if } m = 0 \\ \frac{i}{\sqrt{2}}(j_{sm} - (-1)^m j_{s,-m}) & \text{if } m < 0, \end{cases}$$

where the coefficients  $u \in \mathbb{C}^{\tilde{d}}$  for the former and  $\theta \in \mathbb{R}^{\tilde{d}}$  for the latter are also related by a unitary transform  $\tilde{u} = \tilde{V}^* \theta$ , defined as

$$\tilde{u}_m^{(s)} = \begin{cases} \frac{(-1)^m}{\sqrt{2}}(\tilde{\theta}_{|m|}^{(s)} - i\tilde{\theta}_{-|m|}^{(s)}) & \text{if } m > 0 \\ \tilde{\theta}_0^{(s)} & \text{if } m = 0 \\ \frac{1}{\sqrt{2}}(\tilde{\theta}_{|m|}^{(s)} + i\tilde{\theta}_{|m|}^{(s)}) & \text{if } m < 0. \end{cases} \quad (\text{D.55})$$

It may be checked from the orthonormality of  $\{\hat{j}_{sm}\}$  and  $\{j_{sm}\}$  in  $L_2(\mathbb{R}^2, \mathbb{C})$  that these functions  $\{h_{sm}\}$  are also orthonormal in  $L_2(\mathbb{R}^2, \mathbb{R})$ . This yields the latter two real representations in (4.12) and (4.13).

Recall that since  $f$  is real-valued, the coefficients  $u_m^{(ls)}$  satisfy the sign symmetry (D.29). Similarly, since  $\Pi \cdot f$  is real-valued, its Fourier transform satisfies

$$\widehat{\Pi \cdot f}(\rho, \phi_2) = \overline{\widehat{\Pi \cdot f}(\rho, \pi + \phi_2)}$$

where  $(\rho, \pi + \phi_2)$  is the reflection of  $(\rho, \phi_2)$  about the origin. Then the coefficients  $\tilde{u}_m^{(s)}$  of  $\widehat{\Pi \cdot f}$  in the basis  $\{\hat{j}_{sm}\}$  must satisfy the analogous sign symmetry

$$\tilde{u}_m^{(s)} = (-1)^m \overline{\tilde{u}_{-m}^{(s)}}. \quad (\text{D.56})$$

The identity (D.54) and the relations  $u = \hat{V}^* \theta$  and  $\tilde{u} = \tilde{V}^* \tilde{\theta}$  show that the tomographic projection  $\Pi$  is a linear map from  $\theta \in \mathbb{R}^d$  to  $\tilde{\theta} \in \mathbb{R}^{\tilde{d}}$ , defined as

$$\Pi = \tilde{V} \cdot \Pi^{\mathbb{C}} \cdot \hat{V}^* \in \mathbb{R}^{\tilde{d} \times d}, \quad \Pi_{(s', m'), (l, s, m)}^{\mathbb{C}} = \mathbf{1}\{s = s'\} \cdot \mathbf{1}\{m = m'\} \cdot p_{lm}. \quad (\text{D.57})$$

Here,  $\Pi^{\mathbb{C}}$  is the corresponding linear map from  $u \in \mathbb{C}^d$  to  $\tilde{u} \in \mathbb{C}^{\tilde{d}}$ , and  $p_{lm}$  are the values defined in (D.52). The action of the rotation  $f \mapsto f_{\mathbf{g}}$  on  $\theta \in \mathbb{R}^d$  is as previously described in Lemma D.5, and this expresses the model in the general form of (2.2) for projected orbit recovery.

**Remark D.8.** Note that if  $\{\tilde{z}_s : s = 1, \dots, S\}$  has the same linear span as  $\{z_s : s = 1, \dots, S\}$  used in the unprojected cryo-EM model of Section 4.2, then the two spaces of bandlimited functions (4.12) and (4.7) coincide. However, we caution that here under the orthogonality relation (D.50), the *unprojected* basis  $\{h_{lsm}\}$  is not orthonormal for this function space, so our parametrization  $f \mapsto \theta$  here is not an isometric parametrization of  $f \in L_2(\mathbb{R}^3)$ .

**D.4.2. Terms of the high noise series expansion.** We describe the explicit forms of  $\tilde{s}_1(\theta)$ ,  $\tilde{s}_2(\theta)$ , and  $\tilde{s}_3(\theta)$ . Recalling the entries  $p_{lm}$  in (D.52), define

$$Q_{kl} = \frac{(-1)^{k+l}}{(2k+1)(2l+1)} \sum_{q=-(k \wedge l)}^{k \wedge l} p_{kq}^2 p_{lq}^2, \quad (\text{D.58})$$

$$M_{k, k', k'', l, l', l''} = \frac{(-1)^{k''+l''}}{(2k''+1)(2l''+1)} \sum_{q=-(k \wedge l)}^{k \wedge l} \sum_{\substack{q'=(k' \wedge l') \\ q''=q+q'}}^{k' \wedge l'} \sum_{q''=-(k'' \wedge l'')}^{k'' \wedge l''} \langle k, q; k', q' | k'', q'' \rangle \langle l, q; l', q' | l'', q'' \rangle p_{kq} p_{k'q'} p_{k''q''} p_{lq} p_{l'q'} p_{l''q''}. \quad (\text{D.59})$$

Recall also  $u^{(ls)}(\theta)$  and  $B_{(l, s), (l', s'), (l'', s'')}(\theta)$  from (4.8) and (4.9).

**Theorem D.9.** For any  $L \geq 1$  and  $S_0, \dots, S_L \geq 1$ ,

$$\tilde{s}_1(\theta) = \frac{p_{00}^2}{2} \sum_{s=1}^{S_0} \left( u^{(0s)}(\theta) - u^{(0s)}(\theta_*) \right)^2$$

$$\begin{aligned}
\tilde{s}_2(\theta) &= \frac{1}{4} \sum_{k,l=0}^L Q_{kl} \sum_{s,s'=1}^{S_k \wedge S_l} \left( \langle u^{(ks)}(\theta), u^{(ks')}(\theta) \rangle - \langle u^{(ks)}(\theta_*), u^{(ks')}(\theta_*) \rangle \right) \\
&\quad \times \left( \langle u^{(ls)}(\theta), u^{(ls')}(\theta) \rangle - \langle u^{(ls)}(\theta_*), u^{(ls')}(\theta_*) \rangle \right) \\
\tilde{s}_3(\theta) &= \frac{1}{12} \sum_{\substack{k,k',k'',l,l',l''=0 \\ |k-k'| \leq k'' \leq k+k' \text{ and } |l-l'| \leq l'' \leq l+l'}}^L M_{k,k',k'',l,l',l''} \sum_{s=1}^{S_k \wedge S_l} \sum_{s'=1}^{S_{k'} \wedge S_{l'}} \sum_{s''=1}^{S_{k''} \wedge S_{l''}} \\
&\quad \left( B_{(k,s),(k',s'),(k'',s'')}(\theta) - B_{(k,s),(k',s'),(k'',s'')}(\theta_*) \right) \left( B_{(l,s),(l',s'),(l'',s'')}(\theta) - B_{(l,s),(l',s'),(l'',s'')}(\theta_*) \right).
\end{aligned}$$

*Proof.* Recall from Lemma A.1 that

$$\tilde{s}_k(\theta) = \frac{1}{2(k!)} \mathbb{E}_{g,h} [\langle \Pi g \theta, \Pi h \theta \rangle^k - 2 \langle \Pi g \theta, \Pi h \theta_* \rangle^k + \langle \Pi g \theta_*, \Pi h \theta_* \rangle^k]. \quad (\text{D.60})$$

Consider two different real coefficient vectors  $\theta, \vartheta \in \mathbb{R}^d$ , with corresponding complex coefficients  $u = \hat{V}^* \theta$  and  $v = \hat{V}^* \vartheta$ .

**Case  $k = 1$ :** Notice that

$$\langle \Pi g \theta, \Pi h \vartheta \rangle = \left\langle (\tilde{V}^* \Pi \hat{V})(\hat{V}^* g \hat{V})u, (\tilde{V}^* \Pi \hat{V})(\hat{V}^* h \hat{V})v \right\rangle = \langle D(\mathfrak{g})u, (\Pi^{\mathbb{C}})^* \Pi^{\mathbb{C}} D(\mathfrak{h})v \rangle,$$

where  $D(\mathfrak{g}), D(\mathfrak{h})$  are the block-diagonal matrices in (D.33). The form of  $\Pi^{\mathbb{C}}$  from (D.57) yields

$$(\Pi^{\mathbb{C}*} \Pi^{\mathbb{C}})_{lsm, l' s' m'} = \mathbf{1}\{s = s'\} \cdot \mathbf{1}\{m = m'\} p_{lm} p_{l' m'}$$

so that

$$\langle \Pi g \theta, \Pi h \vartheta \rangle = \sum_{k,l=0}^L \sum_{s=1}^{S_k \wedge S_l} \sum_{m,q=-k}^k \sum_{n,r=-l}^l \overline{D_{qm}^{(k)}(\mathfrak{g}) u_m^{(ks)}} \cdot \mathbf{1}\{q = r\} p_{kq} p_{lr} \cdot D_{rn}^{(l)}(\mathfrak{h}) v_n^{(ls)}. \quad (\text{D.61})$$

Applying (D.4) to take the expectation, we preserve only the terms for  $k = l = m = q = n = r = 0$ , yielding

$$\mathbb{E}_{g,h} [\langle \Pi g \theta, \Pi h \vartheta \rangle] = \sum_{s=1}^{S_0} p_{00}^2 \overline{u_0^{(0s)}} v_0^{(0s)}.$$

Recalling that  $u^{(0s)} = u_0^{(0s)}$  and  $v^{(0s)} = v_0^{(0s)}$  are real-valued by (D.31), and substituting into (D.60), we obtain

$$\tilde{s}_1(\theta) = \frac{p_{00}^2}{2} \sum_{s=1}^{S_0} \left( u^{(0s)}(\theta) - u^{(0s)}(\theta_*) \right)^2.$$

**Case  $k = 2$ :** We square both sides of (D.61) and apply the relations (D.5), (D.29), and (D.53) to get

$$\begin{aligned}
&\mathbb{E}_{g,h} [\langle \Pi g \theta, \Pi h \vartheta \rangle^2] \\
&= \sum_{k,l=0}^L \sum_{s,s'=1}^{S_k \wedge S_l} \sum_{m,q=-k}^k \sum_{n,r=-l}^l \frac{(-1)^{m+q+n+r}}{(2k+1)(2l+1)} \overline{u_m^{(ks)} u_{-m}^{(ks')}} \cdot \mathbf{1}\{q = r\} p_{k,q} p_{k,-q} p_{l,r} p_{l,-r} \cdot v_n^{(ls)} v_{-n}^{(ls')} \\
&= \sum_{k,l=0}^L \sum_{s,s'=1}^{S_k \wedge S_l} \sum_{m,q=-k}^k \sum_{n,r=-l}^l \frac{(-1)^{k+l}}{(2k+1)(2l+1)} \overline{u_m^{(ks)} u_{-m}^{(ks')}} \cdot \mathbf{1}\{q = r\} p_{kq}^2 p_{lr}^2 \cdot v_n^{(ls)} \overline{v_{-n}^{(ls')}} \\
&= \sum_{k,l=0}^L \frac{(-1)^{k+l}}{(2k+1)(2l+1)} \sum_{s,s'=1}^{S_k \wedge S_l} \langle u^{(ks)}, u^{(ks')} \rangle \cdot \overline{\langle v^{(ls)}, v^{(ls')} \rangle} \sum_{q=-(k \wedge l)}^{k \wedge l} p_{kq}^2 p_{lq}^2 \\
&= \sum_{k,l=0}^L Q_{kl} \sum_{s,s'=1}^{S_k \wedge S_l} \langle u^{(ks)}, u^{(ks')} \rangle \cdot \overline{\langle v^{(ls)}, v^{(ls')} \rangle}
\end{aligned}$$

where we have substituted  $Q_{kl}$  from (D.58) in the last equality. By the isometry  $\langle u^{(ks)}, u^{(ks')} \rangle = \langle \theta^{(ks)}, \theta^{(ks')} \rangle$ , both inner products on the last line are real. Then applying this to (D.60),

$$\begin{aligned} \tilde{s}_2(\theta) = \frac{1}{4} \sum_{k,l=0}^L Q_{kl} \sum_{s,s'=1}^{S_k \wedge S_l} \left( \langle u^{(ks)}(\theta), u^{(ks')}(\theta) \rangle - \langle u^{(ks)}(\theta_*), u^{(ks')}(\theta_*) \rangle \right) \\ \times \left( \langle u^{(ls)}(\theta), u^{(ls')}(\theta) \rangle - \langle u^{(ls)}(\theta_*), u^{(ls')}(\theta_*) \rangle \right). \end{aligned}$$

**Case  $k = 3$ :** Let us introduce the abbreviations

$$\sum_{k,k',k'',l,l',l''} = \sum_{\substack{k,k',k'',l,l',l''=0 \\ |k-k'| \leq k'' \leq k+k', |l-l'| \leq l'' \leq l+l'}}^L \quad \text{and} \quad \sum_{s,s',s''} = \sum_{s=1}^{S_k \wedge S_l} \sum_{s'=1}^{S_{k'} \wedge S_{l'}} \sum_{s''=1}^{S_{k''} \wedge S_{l''}}.$$

For given indices  $m, m', q, q', n, n', r, r'$ , let us write as shorthand  $m'' = m + m'$ ,  $n'' = n + n'$ ,  $q'' = q + q'$ , and  $r'' = r + r'$ . We cube both sides of (D.61) and apply the relation (D.6) to obtain

$$\begin{aligned} \mathbb{E}_{g,h}[\langle \Pi g \theta, \Pi h \vartheta \rangle^3] = \sum_{k,k',k'',l,l',l''} \sum_{s,s',s''} \sum_{m,q=-k}^k \sum_{m',q'=-k'}^{k'} \sum_{n,r=-l}^l \sum_{n',r'=-l'}^{l'} \frac{(-1)^{m''+q''+n''+r''}}{(2k''+1)(2l''+1)} \\ \times C_{q,q'}^{k,k',k''} C_{m,m',m''}^{k,k',k''} C_{r,r',r''}^{l,l',l''} C_{n,n',n''}^{l,l',l''} \mathbf{1}\{q=r\} \mathbf{1}\{q'=r'\} \\ \times \overline{u_{-m}^{(ks)} u_{-m'}^{(k's')} u_{-m''}^{(k''s'')}} v_n^{(ls)} v_{n'}^{(l's')} v_{-n''}^{(l''s'')} p_{k,q} p_{k',q'} p_{k'',-q''} p_{l,r} p_{l',r'} p_{l'',-r''}. \end{aligned}$$

Let us apply, by (D.29) and (D.53),

$$\begin{aligned} \overline{u_{-m''}^{(k''s'')}} &= (-1)^{m''+k''} u_{m''}^{(k''s'')}, & \overline{v_{-n''}^{(l''s'')}} &= (-1)^{n''+l''} v_{n''}^{(l''s'')}, \\ p_{k'',-q''} &= (-1)^{q''} p_{k'',q''}, & p_{l'',-r''} &= (-1)^{r''} p_{l'',r''}. \end{aligned}$$

Recalling  $B_{(l,s),(l',s'),(l'',s'')}( \theta )$  from (4.9), which is real-valued, and substituting the form of  $M_{k,k',k'',l,l',l''}$  in (D.59), the above may be written succinctly as

$$\mathbb{E}_{g,h}[\langle \Pi g \theta, \Pi h \vartheta \rangle^3] = \sum_{k,k',k'',l,l',l''} M_{k,k',k'',l,l',l''} \sum_{s,s',s''} B_{(k,s),(k',s'),(k'',s'')}( \theta ) B_{(l,s),(l',s'),(l'',s'')}( \vartheta ).$$

Then by (D.60), we find

$$\begin{aligned} \tilde{s}_3(\theta) = \frac{1}{12} \sum_{k,k',k'',l,l',l''} M_{k,k',k'',l,l',l''} \sum_{s,s',s''} \\ \left( B_{(k,s),(k',s'),(k'',s'')}( \theta ) - B_{(k,s),(k',s'),(k'',s'')}( \theta_* ) \right) \left( B_{(l,s),(l',s'),(l'',s'')}( \theta ) - B_{(l,s),(l',s'),(l'',s'')}( \theta_* ) \right). \end{aligned}$$

□

**D.4.3. Transcendence degrees.** We now prove Theorem 4.10 on the sequences of transcendence degrees.

*Proof of Theorem 4.10.* Theorem 4.6 shows  $\text{trdeg}(\mathcal{R}^G) = d - 3$ . We compute  $\text{trdeg}(\tilde{\mathcal{R}}_{\leq m}^G)$  for  $m = 1, 2, 3$  using Lemma 2.8. For  $m = 1$ ,

$$\nabla^2 \tilde{s}_1(\theta_*) = p_{00}^2 \sum_{s=1}^{S_0} \nabla u^{(0s)}(\theta_*) \nabla u^{(0s)}(\theta_*)^\top.$$

Since  $p_{00} \neq 0$ , this shows  $\text{trdeg}(\tilde{\mathcal{R}}_{\leq 1}^G) = \text{rank}(\nabla^2 \tilde{s}_1(\theta_*)) = S_0$  as in Theorem 4.6.

For  $m = 2$ ,

$$\begin{aligned} \nabla^2 \tilde{s}_1(\theta_*) + \nabla^2 \tilde{s}_2(\theta_*) &= p_{00}^2 \sum_{s=1}^{S_0} \nabla u^{(0s)}(\theta_*) \nabla u^{(0s)}(\theta_*)^\top \\ &+ \frac{1}{2} \sum_{k,l=0}^L Q_{kl} \sum_{s,s'=1}^{S_k \wedge S_l} \nabla [\langle u^{(ks)}(\theta), u^{(ks')}(\theta) \rangle] \nabla [\langle u^{(ls)}(\theta), u^{(ls')}(\theta) \rangle]^\top \Big|_{\theta=\theta_*}. \end{aligned}$$

Recall the form of  $Q_{kl}$  from (D.58). Define the index sets

$$\mathcal{J} = \{(k, r, r') : 0 \leq k \leq L, 1 \leq r, r' \leq S_k\}, \quad \mathcal{L} = \{(q, t, t') : -L \leq q \leq L, 1 \leq t, t' \leq S\}$$

where  $S = \max_{l=0}^L S_l$ , and define a matrix  $D \in \mathbb{R}^{|\mathcal{J}| \times |\mathcal{L}|}$  with the entries

$$D_{krr', qtt'} := \frac{(-1)^k}{2k+1} \cdot \frac{p_{kq}^2}{\sqrt{2}} \cdot \mathbf{1}_{t=r} \mathbf{1}_{t'=r'} \mathbf{1}_{-k \leq q \leq k}.$$

This definition satisfies, for any  $(k, r, r'), (l, s, s') \in \mathcal{J}$ ,

$$\frac{1}{2} Q_{kl} \mathbf{1}_{r=s} \mathbf{1}_{r'=s'} = \sum_{(q, t, t') \in \mathcal{L}} D_{krr', qtt'} D_{lss', qtt'} = (DD^\top)_{krr', lss'}.$$

Then, defining the matrices  $G^0$  and  $G$  with columns

$$\begin{aligned} G_s^0 &:= p_{00} \nabla u^{(0s)}(\theta_*) \quad \text{for } 1 \leq s \leq S_0 \\ G_{krr'} &:= \nabla [\langle u^{(kr)}(\theta), u^{(kr')}(\theta) \rangle] \Big|_{\theta=\theta_*} \quad \text{for } (k, r, r') \in \mathcal{J}, \end{aligned}$$

we have

$$\begin{aligned} \nabla^2 \tilde{s}_1(\theta_*) + \nabla^2 \tilde{s}_2(\theta_*) &= G^0 (G^0)^\top + \sum_{(k, r, r') \in \mathcal{J}} \sum_{(l, s, s') \in \mathcal{J}} \frac{1}{2} Q_{kl} \mathbf{1}_{r=s} \mathbf{1}_{r'=s'} G_{krr'} G_{lss'}^\top \\ &= G^0 (G^0)^\top + G D D^\top G^\top = [GD \mid G^0] [GD \mid G^0]^\top \end{aligned}$$

Here, the square submatrix of  $D$  consisting of the columns  $(q, t, t') \in \mathcal{J} \subset \mathcal{L}$  is lower triangular with non-zero diagonal, because  $p_{kk} \neq 0$  for any  $k = 0, \dots, L$ . Then the column span of  $GD$  coincides with that of  $G$ . Since the column span of  $G^0$  is also contained in that of  $G$  for generic  $\theta_*$ , we conclude that

$$\text{trdeg}(\tilde{\mathcal{R}}_{\leq 2}^{\mathcal{G}}) = \text{rank}(\nabla^2 \tilde{s}_1(\theta_*) + \nabla^2 \tilde{s}_2(\theta_*)) = \text{rank}([GD \mid G^0]) = \text{rank}(G).$$

Then  $\text{trdeg}(\tilde{\mathcal{R}}_{\leq 2}^{\mathcal{G}}) = \text{trdeg}(\mathcal{R}_{\leq 2}^{\mathcal{G}})$  as in Theorem 4.6.

For  $m = 3$ , we have  $\text{trdeg}(\mathcal{R}_{\leq 3}^{\mathcal{G}}) \leq \text{trdeg}(\mathcal{R}^{\mathcal{G}}) = d - 3$ , so it suffices to show  $\text{rank}(\nabla^2 \tilde{s}_3(\theta_*)) \geq d - 3$ . For this, we first write a more convenient form for  $\tilde{s}_3(\theta)$  and its Hessian at  $\theta = \theta_*$ . Recall  $S = \max_{l=0}^L S_l$  and define the index sets

$$\begin{aligned} \mathcal{Q} &= \left\{ ((q, r), (q', r'), (q'', r'')) : -L \leq q, q', q'' \leq L, \quad q + q' + q'' = 0, \quad 1 \leq r, r', r'' \leq S \right\} \\ \mathcal{H} &= \left\{ ((l, s), (l', s'), (l'', s'')) : 0 \leq l, l', l'' \leq L, \quad |l - l'| \leq l'' \leq l + l', \quad 1 \leq s \leq S_l, \quad 1 \leq s' \leq S_{l'}, \quad 1 \leq s'' \leq S_{l''} \right\}. \end{aligned}$$

Define a matrix  $N \in \mathbb{R}^{|\mathcal{Q}| \times |\mathcal{H}|}$  entrywise by

$$\begin{aligned} N_{(q, r), (q', r'), (q'', r''), (l, s), (l', s'), (l'', s'')}^{(l, s), (l', s'), (l'', s'')} &= \mathbf{1}\{r = s, r' = s', r'' = s''\} \cdot \mathbf{1}\{|q| \leq l, |q'| \leq l', |q''| \leq l''\} \\ &\quad \cdot \frac{(-1)^{l''+q''}}{2l''+1} \cdot \langle l, q; l', q' | l'', -q'' \rangle p_{lq} p_{l'q'} p_{l''q''}, \end{aligned} \quad (\text{D.62})$$

where the subscript is the row index in  $\mathcal{Q}$  and the superscript is the column index in  $\mathcal{H}$ . In the expression (D.59) for  $M_{k, k', k'', l, l', l''}$ , let us flip the sign of  $q''$  and apply (D.53) to write this as

$$\begin{aligned} M_{k, k', k'', l, l', l''} &= \frac{(-1)^{k''+q''}}{2k''+1} \frac{(-1)^{l''+q''}}{2l''+1} \sum_{|q| \leq k \wedge l} \sum_{|q'| \leq k' \wedge l'} \sum_{\substack{|q''| \leq k'' \wedge l'' \\ q+q'+q''=0}} \sum_{q+q'+q''=0} \\ &\quad \langle k, q; k', q' | k'', -q'' \rangle \langle l, q; l', q' | l'', -q'' \rangle p_{kq} p_{k'q'} p_{k''q''} p_{lq} p_{l'q'} p_{l''q''}. \end{aligned}$$

Then

$$\tilde{s}_3(\theta) = \frac{1}{12} (B(\theta) - B(\theta_*))^\top N^\top N (B(\theta) - B(\theta_*))$$

Applying the chain rule to differentiate this twice at  $\theta = \theta_*$ , we obtain

$$\nabla^2 \tilde{s}_3(\theta_*) = \frac{1}{6} dB(\theta_*)^\top N^\top N dB(\theta_*),$$

so  $\text{rank}(\nabla^2 \tilde{s}_3(\theta_*)) = \text{rank}(N \cdot dB(\theta_*))$ . We remark that, here, we cannot reduce this directly to  $\text{rank}(dB(\theta_*))$  using the above argument that showed  $\text{rank}(GDD^\top G^\top) = \text{rank}(G)$  for  $m = 2$ . This is because for large  $L$ , the matrix  $N$  may have  $O(L^3)$  columns but only  $O(L^2)$  rows. Then  $N^\top N$  is of low-rank, in contrast to  $DD^\top$  above which is nonsingular.

To analyze  $\text{rank}(N \cdot dB(\theta_*))$ , recall the linear reparametrization by the coordinates  $\eta(\theta)$  in (D.38) and (D.39). Then equivalently

$$\text{rank}(\nabla^2 \tilde{s}_3(\theta_*)) = \text{rank}(N \cdot d_\eta B(\eta_*))$$

The proof of Theorem 4.6 verified that  $\text{rank}(d_\eta B(\eta_*)) = d - 3$  for generic  $\eta_* \in \mathbb{R}^d$ . In fact, let

$$D(\eta_*) = \text{submatrix of } d_\eta B(\eta_*) \text{ with columns } \partial_{w_0^{(12)}}, \partial_{w_1^{(11)}}, \partial_{w_0^{(11)}} \text{ removed.}$$

Then Lemma D.7 shows that  $D(\eta_*)$  has full column rank  $d - 3$  for generic  $\eta_* \in \mathbb{R}^d$ . Applying

$$\text{rank}(N \cdot d_\eta B(\eta_*)) \geq \text{rank}(N \cdot D(\eta_*)),$$

it then suffices to show that  $N \cdot D(\eta_*)$  also has full column rank  $d - 3$  for generic  $\eta_* \in \mathbb{R}^d$ .

For this, we define the following submatrices of  $N$  and  $D(\eta_*)$ . For each  $k \in \{0, 1, \dots, L\}$ , define the index sets

$$\begin{aligned} \mathcal{H}^{(k)} &= \left\{ ((l, s), (l', s'), (l'', s'')) \in \mathcal{H} : \max(l, l', l'') = k \right\}, \\ \mathcal{Q}^{(k)} &= \left\{ ((q, r), (q', r'), (q'', r'')) \in \mathcal{Q} : \max(|q|, |q'|, |q''|) = k \right\}, \\ \mathcal{V}^{(k)} &= \left\{ \text{coordinates } v_m^{(ls)}, w_m^{(ls)} \text{ of } \eta : l = k \right\} \quad \text{if } k \neq 1, \\ \mathcal{V}^{(1)} &= \left\{ \text{coordinates } v_m^{(1s)}, w_m^{(1s)} \text{ of } \eta \right\} \setminus \left\{ w_0^{(12)}, w_1^{(11)}, w_0^{(11)} \right\}. \end{aligned}$$

Let  $N_k \in \mathbb{R}^{|\mathcal{Q}^{(k)}| \times |\mathcal{H}^{(k)}|}$  be the submatrix of  $N$  containing the rows in  $\mathcal{Q}^{(k)}$  and columns in  $\mathcal{H}^{(k)}$ , and let  $D_k(\eta_*) \in \mathbb{R}^{|\mathcal{H}^{(k)}| \times |\mathcal{V}^{(k)}|}$  be the submatrix of  $D(\eta_*)$  containing the rows in  $\mathcal{H}^{(k)}$  and columns in  $\mathcal{V}^{(k)}$ . Similarly, define  $N_{\leq k}$  and  $D_{\leq k}(\eta_*)$  to contain rows and columns of  $\mathcal{Q}^{(l)}, \mathcal{H}^{(l)}, \mathcal{V}^{(l)}$  for  $l \leq k$ . Note that  $D_k(\eta_*)$  and  $D_{\leq k}(\eta_*)$  depend only on the coordinates of  $v_m^{(ls)}$  and  $w_m^{(ls)}$  where  $l \leq k$ , by the definition of  $\mathcal{H}^{(k)}$  and the form of each function  $B_{(l,s),(l',s'),(l'',s'')}$ .

We prove by induction on  $L$  the claim that  $N \cdot D(\eta_*)$  has full column rank for generic  $\eta_* \in \mathbb{R}^d$ . Lemma D.10(a) below shows that for  $L = 1$ , there exists some  $\eta_*$  where  $N \cdot D(\eta_*)$  has full column rank. Then  $N \cdot D(\eta_*)$  has full column rank also for generic  $\eta_*$  by Fact 4.5, establishing the base case  $L = 1$ .

For the inductive step, we establish a block structure on  $N$  and  $D(\eta_*)$ . Block the rows and columns of  $N$  by  $(\mathcal{Q} \setminus \mathcal{Q}^{(L)}, \mathcal{Q}^{(L)})$  and  $(\mathcal{H} \setminus \mathcal{H}^{(L)}, \mathcal{H}^{(L)})$ , and those of  $D(\eta_*)$  by  $(\mathcal{H} \setminus \mathcal{H}^{(L)}, \mathcal{H}^{(L)})$  and  $(\eta \setminus \mathcal{V}^{(L)}, \mathcal{V}^{(L)})$ . Note that  $N_{(q,r),(q',r'),(q'',r'')}^{(l,s),(l',s'),(l'',s'')} = 0$  unless  $\max(|q|, |q'|, |q''|) \leq \max(l, l', l'')$ , and also  $B_{(l,s),(l',s'),(l'',s'')}$  does not depend on any variable  $v_m^{(ks)}$  or  $w_m^{(ks)}$  where  $k > \max(l, l', l'')$ . Thus  $N$  and  $D(\eta_*)$  have the block structures

$$N = \begin{pmatrix} A & B \\ 0 & N_L \end{pmatrix}, \quad D(\eta_*) = \begin{pmatrix} X(\eta_*) & 0 \\ Y(\eta_*) & D_L(\eta_*) \end{pmatrix}$$

for some matrices  $A, B, X(\eta_*), Y(\eta_*)$ .

Let us now specialize to  $\eta_* \in \mathbb{R}^d$  where

$$v_{*,m}^{(Ls)} = w_{*,m}^{(Ls)} = 0 \text{ for all } s = 1, \dots, S_L \text{ and } m = -L, \dots, L. \quad (\text{D.63})$$

The above matrix  $Y(\eta_*)$  contains the derivatives in variables  $\{v_m^{(ks)}, w_m^{(ks)} : k < L\}$  of the functions  $B_{(l,s),(l',s'),(l'',s'')}$  where  $\max(l, l', l'') = L$ . By the form of  $B_{(l,s),(l',s'),(l'',s'')}$ , any such derivative vanishes for  $\eta_*$  satisfying (D.63), so  $Y(\eta_*) = 0$  and

$$N \cdot D(\eta_*) = \begin{pmatrix} A \cdot X(\eta_*) & B \cdot D_L(\eta_*) \\ 0 & N_L \cdot D_L(\eta_*) \end{pmatrix}.$$

The induction hypothesis for  $L - 1$  is exactly the statement that the upper-left block  $A \cdot X(\eta_*)$  has full column rank for all generic values of the coordinates  $\{v_{*,m}^{(ks)}, w_{*,m}^{(ks)} : k \leq L - 1\}$ . Applying Fact 4.5 and Lemma D.10(b) below with  $k = L$ , restricting to  $\eta_*$  satisfying (D.63) and for generic values of the remaining coordinates  $\{v_{*,m}^{(ks)}, w_{*,m}^{(ks)} : k \leq L - 1\}$ , the lower-right block  $N_L \cdot D_L(\eta_*)$  also has full column rank. Then

there exists a point  $\eta_* \in \mathbb{R}^d$  satisfying (D.63) where  $N \cdot D(\eta_*)$  has full column rank. Then  $N \cdot D(\eta_*)$  has full column rank also for generic  $\eta_* \in \mathbb{R}^d$ , completing the induction and the proof.  $\square$

**Lemma D.10.** *If  $S_l \geq 4$  for  $0 \leq l \leq L$ , then we have the following.*

- (a) *There exists a point  $\eta_* \in \mathbb{R}^d$  such that  $N_{\leq 1} \cdot D_{\leq 1}(\eta_*)$  has full column rank.*
- (b) *For each  $k \geq 2$ , there exists a point  $\eta_* \in \mathbb{R}^d$  such that  $v_{*,m}^{(ks)} = w_{*,m}^{(ks)} = 0$  for all  $s \in \{1, \dots, S_k\}$  and  $m \in \{-k, \dots, k\}$ , and  $N_k \cdot D_k(\eta_*)$  has full column rank.*

*Proof of Lemma D.10. Part (a):* Recall from the form of  $p_{lm}$  in (D.52) that  $p_{lm} = 0$  if  $l + m$  is odd and  $p_{lm} \neq 0$  if  $l + m$  is even. Then, for  $\max\{l, l', l''\} \leq 1$ , the non-vanishing of Clebsch-Gordon coefficients in Lemma D.2 and the definition of  $N$  in (D.62) imply that

$$N_{(q,r),(q',r'),(q'',r'')}^{(l,s),(l',s'),(l'',s'')} \neq 0 \text{ if and only if } l = |q|, l' = |q'|, l'' = |q''|, r = s, r' = s', r'' = s''. \quad (\text{D.64})$$

For each  $((l, s), (l', s'), (l'', s'')) \in \mathcal{H}^{(0)} \cup \mathcal{H}^{(1)}$  where  $l = l' + l''$ , take the row  $((-l, s), (l', s'), (l'', s'')) \in \mathcal{Q}^{(0)} \cup \mathcal{Q}^{(1)}$  of  $N_{\leq 1}$ . It suffices to exhibit  $\eta_*$  such that the submatrix of corresponding rows of  $N_{\leq 1} \cdot D_{\leq 1}(\eta_*)$  has full column rank. Observation (D.64) implies that each such row of  $N_{\leq 1}$  has exactly one non-zero entry, which is given by  $N_{(-l,s),(l',s'),(l'',s'')}^{(l,s),(l',s'),(l'',s'')}$ . Then it suffices to show that the submatrix of  $D_{\leq 1}(\eta_*)$  consisting of the rows  $((l, s), (l', s'), (l'', s'')) \in \mathcal{H}^{(0)} \cup \mathcal{H}^{(1)}$  where  $l = l' + l''$  has full column rank. But this has been exhibited already in Lemma D.7, because the proof of Lemma D.7(b–c) in fact only used rows of  $d_{\eta(0s)} B^{(0s)}$  and  $d_{\eta(1s)} B^{(1s)}$  for which  $(l, l', l'') = (0, 0, 0)$  or  $(1, 1, 0)$ , both satisfying  $l = l' + l''$ . This completes the proof of (a).

**Part (b),  $k = 2$  and  $k = 3$ :** The argument is similar to part (a). Observe first that when  $\eta_*$  satisfies  $v_{*,m}^{(ks)} = w_{*,m}^{(ks)} = 0$  for all  $s$  and  $m$ , the rows of  $D_k(\eta_*)$  indexed by  $((l, s), (l', s'), (l'', s'')) \in \mathcal{H}^{(k)}$  having more than one index  $l, l', l''$  equal to  $k$  are identically 0. Let  $D_k(\eta_*)'$  be the submatrix of  $D_k(\eta_*)$  with these rows removed, and let  $N'_k$  be the submatrix of  $N_k$  with the corresponding columns removed. Then  $N_k \cdot D_k(\eta_*) = N'_k \cdot D_k(\eta_*)'$ .

For each remaining tuple  $((l, s), (l', s'), (l'', s'')) \in \mathcal{H}^{(k)}$  where  $l = l' + l''$ , consider the row

$$((-l, s), (l', s'), (l'', s'')) \in \mathcal{Q}^{(k)}$$

of  $N'_k$ . Note that we must have  $(l, l', l'') = (2, 1, 1)$  if  $k = 2$ , and  $(l, l', l'') = (3, 1, 2)$  or  $(3, 2, 1)$  if  $k = 3$ . Each such row has the non-zero entry  $N_{(-l,s),(l',s'),(l'',s'')}^{(l,s),(l',s'),(l'',s'')}$  as above, and this is the only non-zero entry in the row: Indeed, if  $((j, r), (j', r'), (j'', r''))$  is a column of  $N'_k$  where  $N_{(-l,s),(l',s'),(l'',s'')}^{(j,r),(j',r'),(j'',r'')} \neq 0$ , then by definition of  $N$  in (D.62) we must have  $(s, s', s'') = (r, r', r'')$ ,  $j \geq l$ ,  $j' \geq l'$ ,  $j'' \geq l''$ , and each of  $j - l$ ,  $j' - l'$ ,  $j'' - l''$  is even. Columns of  $N'_k$  must satisfy  $(j, j', j'') \in \{(2, 1, 1), (3, 1, 2), (3, 2, 1)\}$ , and this forces  $(j, j', j'') = (l, l', l'')$ . So the non-zero entry in this row of  $N'_k$  is unique, as claimed.

Then it suffices to check that the submatrix of rows of  $D_k(\eta_*)$  indexed by  $((l, s), (l', s'), (l'', s'')) \in \mathcal{H}^{(k)}$  where  $l', l'' < k$  and  $k = l = l' + l''$  has full column rank. This was not exhibited in the proof of Lemma D.7 (which used rows where  $l' + l'' > l$  strictly) but we may show this here by a similar argument, assuming now the availability of 4 different spherical frequencies: Fix spherical frequencies  $(A, B, C, D) = (1, 2, 3, 4)$ , and consider  $\eta_*$  satisfying

$$\text{Type}(l', A), \text{Type}(l', B) = 0, \quad \text{Type}(l', C), \text{Type}(l', D) = 1 \quad \text{for all } l' \in \{1, \dots, k-1\}.$$

Recall that this means  $v_{*,m}^{(l'A)}, w_{*,m}^{(l'A)}, v_{*,m}^{(l'B)}, w_{*,m}^{(l'B)} = 0$  unless  $m = l'$ , and  $v_{*,m}^{(l'C)}, w_{*,m}^{(l'C)}, v_{*,m}^{(l'D)}, w_{*,m}^{(l'D)} = 0$  unless  $m = l' - 1$ . Then, for  $\partial_{v_m^{(ls)}} B_{(l,s),(l',s'),(l'',s'')}, \partial_{w_m^{(ls)}} B_{(l,s),(l',s'),(l'',s'')}$  to be non-zero, this requires as in (D.47)

$$m \in \left\{ |(l' - \text{Type}(l', s')) - (l'' - \text{Type}(l'', s''))|, (l' - \text{Type}(l', s')) + (l'' - \text{Type}(l'', s'')) \right\}. \quad (\text{D.65})$$

For  $k = 2$ , we choose the following 5 rows of  $D_k(\eta_*)$ , with the following corresponding values of  $m$  satisfying (D.65):

TABLE 7.

$(l', s')$ and $(l'', s'')$	Values of $m$
$(1, C)$ and $(1, C)$	0
$(1, A)$ and $(1, A)$	0, 2
$(1, B)$ and $(1, B)$	0, 2
$(1, A)$ and $(1, C)$	1
$(1, B)$ and $(1, D)$	1

Ordering the columns by  $v_0^{(2s)}, v_2^{(2s)}, w_2^{(2s)}, v_1^{(2s)}, w_1^{(2s)}$ , the resulting  $5 \times 5$  submatrix is block lower-triangular with diagonal blocks

$$\begin{aligned} \partial_{v_0^{(2s)}} B_{(2,s),(1,C),(1,C)}(\eta_*) &= C_{0,0,0}^{2,1,1} \cdot \left| u_{*,0}^{(1C)} \right|^2 \\ \partial_{v_2^{(2s)}, w_2^{(2s)}} \left( B_{(2,s),(1,A),(1,A)}, B_{(2,s),(1,B),(1,B)} \right)(\eta_*) &= \begin{pmatrix} 2C_{2,-1,1}^{2,1,1} \cdot \operatorname{Re} \overline{u_{*,1}^{(1A)}} u_{*,1}^{(1A)} & 2C_{2,-1,1}^{2,1,1} \cdot \operatorname{Im} \overline{u_{*,1}^{(1A)}} u_{*,1}^{(1A)} \\ 2C_{2,-1,1}^{2,1,1} \cdot \operatorname{Re} \overline{u_{*,1}^{(1B)}} u_{*,1}^{(1B)} & 2C_{2,-1,1}^{2,1,1} \cdot \operatorname{Im} \overline{u_{*,1}^{(1B)}} u_{*,1}^{(1B)} \end{pmatrix} \\ \partial_{v_1^{(2s)}, w_1^{(2s)}} \left( B_{(2,s),(1,A),(1,C)}, B_{(2,s),(1,B),(1,D)} \right)(\eta_*) &= \begin{pmatrix} 2C_{1,-1,0}^{2,1,1} \cdot \operatorname{Re} \overline{u_{*,0}^{(1A)}} u_{*,0}^{(1C)} & 2C_{1,-1,0}^{2,1,1} \cdot \operatorname{Im} \overline{u_{*,0}^{(1A)}} u_{*,0}^{(1C)} \\ 2C_{1,-1,0}^{2,1,1} \cdot \operatorname{Re} \overline{u_{*,0}^{(1B)}} u_{*,0}^{(1D)} & 2C_{1,-1,0}^{2,1,1} \cdot \operatorname{Im} \overline{u_{*,0}^{(1B)}} u_{*,0}^{(1D)} \end{pmatrix} \end{aligned}$$

These blocks are generically non-singular, so this submatrix of  $D_2(\eta_*)$  has full column rank.

For  $k = 3$ , we choose the following 7 rows of  $D_k(\eta_*)$ , with the following corresponding values of  $m$  satisfying (D.65):

TABLE 8.

$(l', s')$ and $(l'', s'')$	values of $m$
$(1, D)$ and $(2, B)$	2
$(1, C)$ and $(2, A)$	2
$(1, C)$ and $(2, C)$	1
$(1, D)$ and $(2, D)$	1
$(1, A)$ and $(2, A)$	1, 3
$(1, B)$ and $(2, B)$	1, 3
$(1, A)$ and $(2, C)$	2, 0

Ordering the columns by  $v_2^{(3s)}, w_2^{(3s)}, v_1^{(3s)}, w_1^{(3s)}, v_3^{(3s)}, w_3^{(3s)}, w_0^{(3s)}$ , the resulting  $7 \times 7$  submatrix is block lower-triangular with diagonal blocks

$$\begin{aligned} \partial_{v_2^{(3s)}, w_2^{(3s)}} \left( B_{(3,s),(1,D),(2,B)}, B_{(3,s),(1,C),(2,A)} \right)(\eta_*) &= \begin{pmatrix} 2C_{2,0,2}^{3,1,2} \cdot \operatorname{Re} \overline{u_{*,2}^{(1D)}} u_{*,2}^{(2B)} & 2C_{2,0,2}^{3,1,2} \cdot \operatorname{Im} \overline{u_{*,2}^{(1D)}} u_{*,2}^{(2B)} \\ 2C_{2,0,2}^{3,1,2} \cdot \operatorname{Re} \overline{u_{*,2}^{(1C)}} u_{*,2}^{(2A)} & 2C_{2,0,2}^{3,1,2} \cdot \operatorname{Im} \overline{u_{*,2}^{(1C)}} u_{*,2}^{(2A)} \end{pmatrix} \\ \partial_{v_1^{(3s)}, w_1^{(3s)}} \left( B_{(3,s),(1,C),(2,C)}, B_{(3,s),(1,D),(2,D)} \right)(\eta_*) &= \begin{pmatrix} 2C_{1,0,1}^{3,1,2} \cdot \operatorname{Re} \overline{u_{*,1}^{(1C)}} u_{*,1}^{(2C)} & 2C_{1,0,1}^{3,1,2} \cdot \operatorname{Im} \overline{u_{*,1}^{(1C)}} u_{*,1}^{(2C)} \\ 2C_{1,0,1}^{3,1,2} \cdot \operatorname{Re} \overline{u_{*,1}^{(1D)}} u_{*,1}^{(2D)} & 2C_{1,0,1}^{3,1,2} \cdot \operatorname{Im} \overline{u_{*,1}^{(1D)}} u_{*,1}^{(2D)} \end{pmatrix} \\ \partial_{v_3^{(3s)}, w_3^{(3s)}} \left( B_{(3,s),(1,A),(2,A)}, B_{(3,s),(1,B),(2,B)} \right)(\eta_*) &= \begin{pmatrix} 2C_{3,-1,2}^{3,1,2} \cdot \operatorname{Re} \overline{u_{*,2}^{(1A)}} u_{*,2}^{(2A)} & 2C_{3,-1,2}^{3,1,2} \cdot \operatorname{Im} \overline{u_{*,2}^{(1A)}} u_{*,2}^{(2A)} \\ 2C_{3,-1,2}^{3,1,2} \cdot \operatorname{Re} \overline{u_{*,2}^{(1B)}} u_{*,2}^{(2B)} & 2C_{3,-1,2}^{3,1,2} \cdot \operatorname{Im} \overline{u_{*,2}^{(1B)}} u_{*,2}^{(2B)} \end{pmatrix} \\ \partial_{w_0^{(3s)}} B_{(3,s),(1,A),(2,C)}(\eta_*) &= 2C_{0,1,1}^{3,1,2} \cdot \operatorname{Im} \overline{u_{*,1}^{(1A)}} u_{*,1}^{2C} \end{aligned}$$

These blocks are again generically non-singular, so this submatrix of  $D_3(\eta_*)$  has full column rank.

This verifies that  $N_k \cdot D_k(\eta_*)$  has full column rank for  $k = 2, 3$ .

**Part (b),  $k \geq 4$ :** As above, we fix  $(A, B, C, D) = (1, 2, 3, 4)$  and consider  $\eta_*$  satisfying both  $v_{*,m}^{(ks)}, w_{*,m}^{(ks)} = 0$  for all  $m, s$  and

$$\operatorname{Type}(l', A), \operatorname{Type}(l', B) = 0, \quad \operatorname{Type}(l', C), \operatorname{Type}(l', D) = 1 \quad \text{for all } l' \in \{1, \dots, k-1\}. \quad (\text{D.66})$$

Columns of  $D_k(\eta_*)$  correspond to derivatives in the coordinates  $\mathcal{V}^{(k)}$ . We partition these coordinates into blocks  $s = 1, \dots, S_k$  and write

$$D_k(\eta_*) = [D_k^{(s)}(\eta_*) : s = 1, \dots, S_k],$$

where columns of each  $D_k^{(s)}(\eta_*)$  are indexed by  $\eta_0^{(ks)}, v_1^{(ks)}, w_1^{(ks)}, \dots, v_k^{(ks)}, w_k^{(ks)}$ . It suffices to show that  $N_k \cdot D_k^{(s)}(\eta_*)$  has full column rank  $2k + 1$  for generic  $\eta_*$  satisfying (D.66), for each fixed  $s$ . We do this by choosing  $2k + 1$  rows of  $N_k$ —call this submatrix  $N'_k$ —and verifying that the corresponding  $(2k + 1) \times (2k + 1)$  submatrix  $N'_k \cdot D_k^{(s)}(\eta_*)$  is non-singular.

The argument for verifying non-singularity is different from our preceding approaches in Lemmas D.4 and D.7. Let us first explain the high-level idea: Rather than exhibiting a sparse structure for  $N'_k \cdot D_k^{(s)}(\eta_*)$  where the rank may be explicitly checked, we study the determinant

$$P(\eta_*) = \det[N'_k \cdot D_k^{(s)}(\eta_*)] \quad (\text{D.67})$$

and show that this is not identically 0 as a polynomial of the non-zero coordinates of  $\eta_*$ . We introduce a special degree- $(2k + 1)$  monomial

$$M = \left( w_0^{(1C)} w_0^{(1D)} v_1^{(1A)} v_1^{(1B)} \right)^2 \left( \prod_{j=2}^{\lfloor k/2 \rfloor - 1} v_j^{(jA)} v_j^{(jB)} v_{j-1}^{(jC)} v_{j-1}^{(jD)} \right) v_{\lfloor k/2 \rfloor - 1}^{(\lfloor k/2 \rfloor C)} v_{\lfloor k/2 \rfloor - 1}^{(\lfloor k/2 \rfloor A)} v_{\lfloor k/2 \rfloor - 1}^{(\lfloor k/2 \rfloor D)} \mathbf{1}_{\{k \text{ odd}\}}, \quad (\text{D.68})$$

where all variables appearing in  $M$  are coordinates of  $\eta_*$  which are not fixed to be zero. We then write

$$P = (P/M) \cdot M + Q \quad (\text{D.69})$$

where  $Q$  are the terms of  $P$  not divisible by  $M$ ,  $(P/M) \cdot M$  are the terms which are divisible by  $M$ , and  $P/M$  denotes their quotient by  $M$ . It suffices to show that  $P/M$  is not identically 0.

We now describe the choice of  $2k + 1$  rows of  $N'_k$  that allows us to verify this claim  $P/M \neq 0$ . We restrict to rows  $((-k, s), (q', s'), (q'', s'')) \in \mathcal{Q}^{(k)}$  of  $N_k$  where the first pair is fixed to be  $(-k, s)$ , and where  $q' \in \{1, \dots, \lfloor k/2 \rfloor\}$ . This requires  $-k + q' + q'' = 0$ , so  $q'' = k - q' \in \{k - 1, \dots, \lceil k/2 \rceil\}$ . We index such rows by  $(q', s'), (q'', s'')$ . For any such row  $(q', s'), (q'', s'')$ , we apply the following two observations:

- By definition of  $N$  in (D.62), each non-zero entry in this row of  $N_k$  belongs to a column  $((k, s), (l', s'), (l'', s'')) \in \mathcal{H}^{(k)}$  where  $l', l'' \in \{1, \dots, k - 1\}$  and

$$l' \geq q', \quad l'' \geq q'', \quad l' - q', l'' - q'' \text{ are even.} \quad (\text{D.70})$$

- As in (D.47), for this row  $((k, s), (l', s'), (l'', s'')) \in \mathcal{H}^{(k)}$  of  $D_k^{(s)}(\eta_*)$ , the entries in the columns  $\partial_{v_m^{(ks)}}, \partial_{w_m^{(ks)}}$  can be non-zero only when

$$m \in \left\{ |(l' - \text{Type}(l', s')) - (l'' - \text{Type}(l'', s''))|, (l' - \text{Type}(l', s')) + (l'' - \text{Type}(l'', s'')) \right\}. \quad (\text{D.71})$$

Combined, these yield the important observation that, fixing a row  $(q', s'), (q'', s'')$  of  $N_k \cdot D_k^{(s)}(\eta_*)$  and a pair of columns  $\partial_{v_m^{(ks)}}, \partial_{w_m^{(ks)}}$  (or a single column in the case  $m = 0$ ) for a specific index  $m \in \{0, \dots, k\}$ , these two entries (or one entry) of  $N_k \cdot D_k^{(s)}(\eta_*)$  are homogenous degree-2 polynomials, whose degree-2 monomials are each a product of some variable  $v^{l' \cdot}, w^{l' \cdot}$  and some variable  $v^{l'' \cdot}, w^{l'' \cdot}$  where  $l', l''$  satisfy both conditions (D.70) and (D.71).

Table 9 now specifies an explicit choice of  $2k + 1$  rows  $(q', s'), (q'', s'')$  of  $N_k$  to form  $N'_k$ , and indicates which columns  $\partial_{v_m^{(ks)}}, \partial_{w_m^{(ks)}}$  of each corresponding row of  $N_k \cdot D_k^{(s)}(\eta_*)$  can depend on some variable  $v^{q' \cdot}, w^{q' \cdot}$  with the same spherical frequency as the first row index  $q'$ . For example: If  $(s', s'', q', q'') = (A, A, 1, k - 1)$ , then (D.70) forces  $l'' = k - 1$ . In order for a term of this row to depend on  $v^{1 \cdot}, w^{1 \cdot}$ , we must then have  $l' = 1$ . Then  $\text{Type}(l', s') = 1$  and  $\text{Type}(l'', s'') = k - 1$ , so the condition (D.71) implies that only columns corresponding to  $m \in \{k, k - 2\}$  can depend on such variables  $v^{1 \cdot}, w^{1 \cdot}$ . This yields the first row of the table. If  $(s', s'', q', q'') = (C, C, 3, k - 3)$ , then (D.70) forces  $l'' \in \{k - 3, k - 1\}$ . For this row to depend on  $v^{3 \cdot}, w^{3 \cdot}$ , we must have  $l' = 3$ . Then  $\text{Type}(l', s') = 2$  and  $\text{Type}(l'', s'') \in \{k - 4, k - 2\}$ , so  $m \in \{k - 6, k - 4, k - 2, k\}$ , and this yields the 13<sup>th</sup> row of the table. The remaining rows are deduced by the same type of reasoning. (The sequences  $(2, 4, \dots)$ ,  $(3, 5, \dots)$ , etc. in Table 9 denote some sequences of consecutive even and odd integers, whose exact last elements will not be important for our later arguments.)

Order the rows of  $N'_k \cdot D_k^{(s)}(\eta_*)$  in the order listed in Table 9, and the columns in the ordering of *decreasing*  $m$ :

$$v_k^{(ks)}, w_k^{(ks)}, v_{k-1}^{(ks)}, w_{k-1}^{(ks)}, v_{k-2}^{(ks)}, w_{k-2}^{(ks)}, \dots, v_1^{(ks)}, w_1^{(ks)}, \eta_0^{(ks)}.$$

TABLE 9.

	$s'$	$s''$	$q'$	$q''$	$m$ s.t. $\partial_{v_m^{(ks)}}, \partial_{w_m^{(ks)}}$ can depend on $v^{q' \cdot}, w^{q' \cdot}$
	$A$	$A$	1	$k-1$	$k, k-2$
	$B$	$B$	1	$k-1$	$k, k-2$
	$C$	$C$	1	$k-1$	$k-2$
	$D$	$D$	1	$k-1$	$k-2$
	$A$	$C$	1	$k-1$	$k-1, k-3$
	$B$	$D$	1	$k-1$	$k-1, k-3$
	$C$	$A$	1	$k-1$	$k-1$
	$D$	$B$	1	$k-1$	$k-1$
	$C$	$C$	2	$k-2$	$k-4$ and $k-2$
	$D$	$D$	2	$k-2$	$k-4$ and $k-2$
	$A$	$C$	2	$k-2$	$k-5$ and $k-1$
	$B$	$D$	2	$k-2$	$k-5$ and $k-1$
	$C$	$C$	3	$k-3$	$k-6, k-4, k-2$ and $k$
	$D$	$D$	3	$k-3$	$k-6, k-4, k-2$ and $k$
	$A$	$C$	3	$k-3$	$k-7, k-5$ and $k-1$
	$B$	$D$	3	$k-3$	$k-7, k-5$ and $k-1$
			$\vdots$	$\vdots$	$\vdots$
	$C$	$C$	$\lfloor k/2 \rfloor - 1$	$\lceil k/2 \rceil + 1$	$2, 4, \dots$ and $k-2, k$ if $k$ even; $3, 5, \dots$ and $k-2, k$ if $k$ odd
	$D$	$D$	$\lfloor k/2 \rfloor - 1$	$\lceil k/2 \rceil + 1$	$2, 4, \dots$ and $k-2, k$ if $k$ even; $3, 5, \dots$ and $k-2, k$ if $k$ odd
	$A$	$C$	$\lfloor k/2 \rfloor - 1$	$\lceil k/2 \rceil + 1$	$1, 3, \dots$ and $k-1$ if $k$ even; $2, 4, \dots$ and $k-1$ if $k$ odd
	$B$	$D$	$\lfloor k/2 \rfloor - 1$	$\lceil k/2 \rceil + 1$	$1, 3, \dots$ and $k-1$ if $k$ even; $2, 4, \dots$ and $k-1$ if $k$ odd
(if $k$ odd)	$C$	$C$	$\lfloor k/2 \rfloor$	$\lceil k/2 \rceil$	$0, 2, \dots$ and $k-2, k$ if $k$ even; $1, 3, \dots$ and $k-2, k$ if $k$ odd
(if $k$ odd)	$D$	$D$	$\lfloor k/2 \rfloor$	$\lceil k/2 \rceil$	$1, 3, \dots$ and $k-2, k$
(if $k$ odd)	$A$	$C$	$\lfloor k/2 \rfloor$	$\lceil k/2 \rceil$	$0, 2, \dots$ and $k-1$

Consider the block decomposition for both rows and columns of  $N'_k \cdot D_k^{(s)}(\eta_*)$  with respect to

$$2k+1 = 8 + 4 + 4 + \dots + 4 + \begin{cases} 1 & \text{if } k \text{ even} \\ 3 & \text{if } k \text{ odd.} \end{cases}$$

Let  $P = \det(N'_k \cdot D_k^{(s)}(\eta_*))$  as defined in (D.67), and let  $P_1, P_2, \dots, P_{\lfloor k/2 \rfloor}$  be the determinants of the diagonal blocks in this decomposition. Recall that we wish to show  $P/M \neq 0$ . Let us factor  $M$  in (D.68) correspondingly as  $M = M_1 M_2 \dots M_{\lfloor k/2 \rfloor}$  where

$$\begin{aligned} M_1 &= (w_0^{(1C)} w_0^{(1D)} v_1^{(1A)} v_1^{(1B)})^2 \\ M_j &= v_j^{(jA)} v_j^{(jB)} v_{j-1}^{(jC)} v_{j-1}^{(jD)} \quad \text{for } j = 2, \dots, \lfloor k/2 \rfloor - 1 \\ M_{\lfloor k/2 \rfloor} &= v_{\lfloor k/2 \rfloor - 1}^{(\lfloor k/2 \rfloor C)} \left( v_{\lfloor k/2 \rfloor}^{(\lfloor k/2 \rfloor A)} v_{\lfloor k/2 \rfloor - 1}^{(\lfloor k/2 \rfloor D)} \right)^{\mathbf{1}_{\{k \text{ odd}\}}} \end{aligned}$$

The degrees of  $M_1, M_2, M_3, \dots$  coincide with the above block sizes  $8, 4, 4, \dots$ . Furthermore, each  $M_j$  depends on only variables  $v^{j \cdot}, w^{j \cdot}$  having spherical frequency  $j$ .

Observe now that:

- Only the  $8 \times 8$  upper-left diagonal block of  $N'_k \cdot D_k^{(s)}(\eta_*)$  has entries depending on variables  $v^{1 \cdot}, w^{1 \cdot}$ . This is because all monomials in rows 9 onwards are a product of some  $v^{l' \cdot}, w^{l' \cdot}$  with some  $v^{l'' \cdot}, w^{l'' \cdot}$  where  $l' \geq q' \geq 2$  and  $l'' \geq q'' \geq 2$ . In the first 8 rows, the table indicates that only the first 8 columns (corresponding to  $m = k, k-1, k-2, k-3$ ) can depend on  $v^{1 \cdot}, w^{1 \cdot}$ .
- Furthermore, any degree-2 monomial in this  $8 \times 8$  block that depends on  $v^{1 \cdot}, w^{1 \cdot}$  must have as its second variable  $v^{l'' \cdot}, w^{l'' \cdot}$  for some  $l'' \geq k-1 > \lfloor k/2 \rfloor$  strictly.
- Removing this first row block and column block of size 8, only the  $4 \times 4$  upper-left diagonal block of the remaining matrix has entries depending on  $v^{2 \cdot}, w^{2 \cdot}$ , by the same reasoning. Furthermore, any

degree-2 monomial in this  $4 \times 4$  block that depends on  $v^{2\cdot}, w^{2\cdot}$  must have as its second variable  $v^{l''\cdot}, w^{l''\cdot}$  for some  $l'' \geq k-2 > \lfloor k/2 \rfloor$  strictly.

- Removing also this second row block and column block of size 4, only the  $4 \times 4$  upper-left remaining diagonal block has entries depending on  $v^{3\cdot}, w^{3\cdot}$ , etc. This argument can be continued inductively until the last block.

These observations imply that the terms of  $P$  divisible by  $M$  must have the factorization

$$P/M = (P_1/M_1)(P_2/M_2) \dots (P_{\lfloor k/2 \rfloor}/M_{\lfloor k/2 \rfloor}), \quad (\text{D.72})$$

where, analogously to (D.69), each factor  $P_j/M_j$  is the polynomial that is the quotient by  $M_j$  of those terms of  $P_j$  which are exactly divisible by  $M_j$ .

To complete the proof, we check by direct computation that each polynomial  $P_j/M_j$  on the right side of (D.72) is non-zero.

**Verification that  $P_1/M_1 \neq 0$ :** Consider, as an example, the entry in the first row  $(A, A, 1, k-1)$  and first column  $\partial_{v_k^{(ks)}}$  of  $N'_k \cdot D_k^{(s)}(\eta_*)$ . This entry is the inner product

$$(N_{(-k,s),(1,A),(k-1,A)})^\top (\partial_{v_k^{(ks)}} B(\eta_*)) = \sum_{((k,s),(l',s'),(l'',s'')) \in \mathcal{H}^{(k)}} N_{(-k,s),(1,A),(k-1,A)}^{(k,s),(l',s'),(l'',s'')} \cdot \partial_{v_k^{(ks)}} B_{(k,s),(l',s'),(l'',s'')}(\eta_*).$$

Importantly, only the single term indexed by  $((k,s),(l',s'),(l'',s'')) = ((k,s),(1,A),(k-1,A))$  of this sum depends on any variable appearing in the monomial  $M_1$ . This is because, from the definition of  $N$  in (D.62), all other non-zero entries  $N_{(-k,s),(1,A),(k-1,A)}^{(k,s),(l',s'),(l'',s'')}$  of this row of  $N$  have  $l' \geq 3$  and  $l'' = k-1$ . We introduce the shorthand

$$Y_q = N_{(-k,s),(q,s'),(k-q,s'')}^{(k,s),(q,s'),(k-q,s'')},$$

where this notation fixes  $k$  and uses that this value does not actually depend on  $(s,s',s'')$ . Note that by (D.62) and the non-vanishing of Clebsch-Gordan coefficients in Lemma D.2,  $Y_q \neq 0$  for every  $q = 0, 1, \dots, k$ . Then, applying (D.41), (D.66), and the sign symmetry (D.29), the above single term is

$$\begin{aligned} N_{(-k,s),(1,A),(k-1,A)}^{(k,s),(1,A),(k-1,A)} \cdot \partial_{v_k^{(ks)}} B_{(k,s),(1,A),(k-1,A)}(\eta_*) &= Y_1 \cdot 2C_{k,-1,k-1}^{k,1,k-1} \overline{\text{Re } u_{-1}^{(1A)}} u_{k-1}^{(k-1,A)} \\ &= 2Y_1 C_{k,-1,k-1}^{k,1,k-1} \left( v_1^{(1A)} v_{k-1}^{(k-1,A)} - w_1^{(1A)} w_{k-1}^{(k-1,A)} \right). \end{aligned}$$

Only the first of these two summands depends on a variable in  $M_1$ , namely  $v_1^{(1A)}$ . We write its quotient by this variable  $v_1^{(1A)}$  in the upper-left entry of the first table below.

As a second example, consider the entry in row  $(C, C, 1, k-1)$  and column  $\partial_{v_{k-2}^{(ks)}}$  of  $N'_k \cdot D_k^{(s)}(\eta_*)$ . By the same reasoning as above, the only term of this entry which depends on a variable in  $M_1$  is

$$\begin{aligned} N_{(-k,s),(1,C),(k-1,C)}^{(k,s),(1,C),(k-1,C)} \cdot \partial_{v_{k-2}^{(ks)}} B_{(k,s),(1,C),(k-1,C)} &= Y_1 \cdot 2C_{k-2,0,k-2}^{k,1,k-1} \cdot \overline{\text{Re } u_0^{(1C)}} u_{k-2}^{(k-1,C)} \\ &= 2Y_1 C_{k-2,0,k-2}^{k,1,k-1} \cdot w_0^{(1C)} w_{k-2}^{(k-1,C)}. \end{aligned}$$

Its quotient by the variable  $w_0^{(1C)}$  appearing in  $M_1$  is the  $(3,3)$  entry of the first table below.

The entries of the  $8 \times 8$  block for  $P_1$  which depend on some variable in  $M_1$  are contained within two  $4 \times 4$  submatrices, corresponding to the below two tables. Similar to the above computations, each entry of each submatrix has at most 1 term depending on some variable in  $M_1$ . We indicate the quotient of this term by the corresponding variable of  $M_1$  in the two tables below. For entries that have no dependence on variables of  $M_1$ , we write this quotient as 0.

	$v_k^{k,s}$	$w_k^{k,s}$	$v_{k-2}^{k,s}$	$w_{k-2}^{k,s}$
$A, A, 1, k-1$	$2Y_1 C_{k-1,1,k-1}^{k-1,A} v_{k-1}^{k-1,A}$	$2Y_1 C_{k-1,1,k-1}^{k-1,A} w_{k-1}^{k-1,A}$	$2Y_1 C_{k-2,1,k-1}^{k-1,A} v_{k-1}^{k-1,A}$	$2Y_1 C_{k-2,1,k-1}^{k-1,A} w_{k-1}^{k-1,A}$
$B, B, 1, k-1$	$2Y_1 C_{k-1,1,k-1}^{k-1,B} v_{k-1}^{k-1,B}$	$2Y_1 C_{k-1,1,k-1}^{k-1,B} w_{k-1}^{k-1,B}$	$2Y_1 C_{k-2,1,k-1}^{k-1,B} v_{k-1}^{k-1,B}$	$2Y_1 C_{k-2,1,k-1}^{k-1,B} w_{k-1}^{k-1,B}$
$C, C, 1, k-1$	0	0	$2Y_1 C_{k-2,0,k-2}^{k-1,C} v_{k-2}^{k-1,C}$	$-2Y_1 C_{k-2,0,k-2}^{k-1,C} w_{k-2}^{k-1,C}$
$D, D, 1, k-1$	0	0	$2Y_1 C_{k-2,0,k-2}^{k-1,D} v_{k-2}^{k-1,D}$	$-2Y_1 C_{k-2,0,k-2}^{k-1,D} w_{k-2}^{k-1,D}$
	$v_{k-1}^{k,s}$	$w_{k-1}^{k,s}$	$v_{k-3}^{k,s}$	$w_{k-3}^{k,s}$
$A, C, 1, k-1$	$2Y_1 C_{k-1,1,k-2}^{k-1,C} v_{k-2}^{k-1,C}$	$2Y_1 C_{k-1,1,k-2}^{k-1,C} w_{k-2}^{k-1,C}$	$2Y_1 C_{k-3,1,k-2}^{k-1,C} v_{k-2}^{k-1,C}$	$2Y_1 C_{k-3,1,k-2}^{k-1,C} w_{k-2}^{k-1,C}$
$B, D, 1, k-1$	$2Y_1 C_{k-1,1,k-2}^{k-1,D} v_{k-2}^{k-1,D}$	$2Y_1 C_{k-1,1,k-2}^{k-1,D} w_{k-2}^{k-1,D}$	$2Y_1 C_{k-3,1,k-2}^{k-1,D} v_{k-2}^{k-1,D}$	$2Y_1 C_{k-3,1,k-2}^{k-1,D} w_{k-2}^{k-1,D}$
$C, A, 1, k-1$	$2Y_1 C_{k-1,0,k-1}^{k-1,A} w_{k-1}^{k-1,A}$	$-2Y_1 C_{k-1,0,k-1}^{k-1,A} v_{k-1}^{k-1,A}$	0	0
$D, B, 1, k-1$	$2Y_1 C_{k-1,0,k-1}^{k-1,B} w_{k-1}^{k-1,B}$	$-2Y_1 C_{k-1,0,k-1}^{k-1,B} v_{k-1}^{k-1,B}$	0	0

Now  $P_1/M_1$  is the product of determinants of the above two  $4 \times 4$  matrices. By Lemma D.2, each Clebsch-Gordan coefficient here is non-zero. Then the determinant of each  $4 \times 4$  matrix is the product of two  $2 \times 2$  determinants, each of which is a non-vanishing quadratic, so  $P_1/M_1 \neq 0$ .

**Verification that  $P_j/M_j \neq 0$  for  $j = 2, \dots, \lfloor k/2 \rfloor - 1$ :** Consider, as an example, the entry of row  $(C, C, 3, k-3)$  and column  $\partial_{w_{k-6}}^{(ks)}$ . This entry is the inner product

$$\left(N_{(-k,s),(3,C),(k-3,C)}\right)^\top \left(\partial_{w_{k-6}}^{(ks)} B(\eta_*)\right).$$

The non-zero entries  $N_{(-k,s),(3,C),(k-3,C)}^{(k,s),(l',C),(l'',C)}$  of this row of  $N$  must have  $l' \geq 3$  odd and  $l'' \in \{k-1, k-3\}$ . We must then have  $l' = 3$  in order for the corresponding entry  $\partial_{w_{k-6}}^{(ks)} B_{(k,s),(3,C),(l',C),(l'',C)}(\eta_*)$  to depend on some variable of  $M_3$ , which would be  $v_2^{(3C)}$ . Recalling the form of this derivative in (D.42), this forces  $|m'| = 2$  for the summation index of (D.42) corresponding to any term depends on  $v_2^{(3C)}$ . Then  $m + m' \in \{k-8, k-4\}$ , since  $m = k-6$ . Applying our specialization (D.71) to the derivative (D.42), this requires  $l'' \in \{k-7, k-3\}$  in order for this term to be non-zero. Combining with the above condition  $l'' \in \{k-1, k-3\}$ , we must have  $l'' = k-3$ . Thus, to summarize, again only a single term of the sum constituting the above inner-product depends on the variable  $v_2^{(3C)}$  of  $M_3$ . This term is

$$\begin{aligned} N_{(-k,s),(3,C),(k-3,C)}^{(k,s),(3,C),(k-3,C)} \cdot \partial_{w_{k-6}}^{(ks)} B_{(k,s),(3,C),(k-3,C)}(\eta_*) &= Y_3 \cdot 2C_{k-6,2,k-4}^{k,3,k-3} \operatorname{Im} \overline{u_2^{(3C)}} u_{k-4}^{(k-3,C)} \\ &= 2Y_3 C_{k-6,2,k-4}^{k,3,k-3} \left( v_2^{(3C)} w_{k-4}^{(k-3,C)} - w_2^{(3C)} v_{k-4}^{(k-3,C)} \right). \end{aligned}$$

Only the first summand depends on  $v_2^{(3C)}$ , and its quotient by  $v_2^{(3C)}$  is recorded in row 1 and column 2 of the table below, corresponding to  $j = 3$ .

By the same reasoning, a similar simplification occurs for every  $j = 2, \dots, \lfloor k/2 \rfloor - 1$  and every entry of  $P_j/M_j$ . For general  $j$ , we may compute each entry of this  $4 \times 4$  block that depends on a variable of  $M_j$ , and the table records the quotient of this entry by the corresponding variable of  $M_j$ .

	$v_{k-2j}^{k,s}$	$w_{k-2j}^{k,s}$	$v_{k-2j-1}^{k,s}$	$w_{k-2j-1}^{k,s}$
$C, C, j, k-j$	$2Y_j C_{k-2j,j-1,k-j-1}^{k-j,C} v_{k-j-1}^{k-j,C}$	$2Y_j C_{k-2j,j-1,k-j-1}^{k-j,C} w_{k-j-1}^{k-j,C}$	0	0
$D, D, j, k-j$	$2Y_j C_{k-2j,j-1,k-j-1}^{k-j,D} v_{k-j-1}^{k-j,D}$	$2Y_j C_{k-2j,j-1,k-j-1}^{k-j,D} w_{k-j-1}^{k-j,D}$	0	0
$A, C, j, k-j$	0	0	$2Y_j C_{k-2j-1,j,k-j-1}^{k-j,C} v_{k-j-1}^{k-j,C}$	$2Y_j C_{k-2j-1,j,k-j-1}^{k-j,C} w_{k-j-1}^{k-j,C}$
$B, D, j, k-j$	0	0	$2Y_j C_{k-2j-1,j,k-j-1}^{k-j,D} v_{k-j-1}^{k-j,D}$	$2Y_j C_{k-2j-1,j,k-j-1}^{k-j,D} w_{k-j-1}^{k-j,D}$

Then  $P_j/M_j$  is the determinant of this  $4 \times 4$  matrix, which is a product of two  $2 \times 2$  determinants. By Lemma D.2, these Clebsch-Gordan coefficients are non-zero, so each  $2 \times 2$  determinant is a non-vanishing quadratic, and  $P_j/M_j \neq 0$ .

**Verification that  $P_{\lfloor k/2 \rfloor}/M_{\lfloor k/2 \rfloor} \neq 0$ :** If  $k$  is even, we have  $\eta_0^{(ks)} = v_0^{(ks)}$ , and

$$P_{k/2}(\eta_*) = \left(N_{(-k,s),(k/2,C),(k/2,C)}\right)^\top \left(\partial_{v_0}^{(ks)} B(\eta_*)\right).$$

The non-zero elements  $N_{(-k,s),(k/2,C),(k/2,C)}^{(k,s),(l',C),(l'',C)}$  have  $l' \geq k/2$  and  $l'' \geq k/2$  with  $l' - k/2$  and  $l'' - k/2$  both even. From (D.43) and the specialization (D.71), the only term of this inner-product depending on  $M_{k/2} = v_{(k/2)-1}^{(k/2)C}$  arises from  $l' = l'' = k/2$ , and this term is

$$N_{(-k,s),(k/2,C),(k/2,C)}^{(k,s),(k/2,C),(k/2,C)} \cdot \partial_{v_0}^{(ks)} B_{(k,s),(k/2,C),(k/2,C)}(\eta_*)$$




FIGURE E.1. (a) 24.6Å-resolution and (b) 8.2Å-resolution low-pass filtered maps for the rotavirus VP6 trimer, prior to performing basis approximation as depicted in Figure 5.1. (c) 7.0Å-resolution low-pass filtered map for hemoglobin, prior to performing basis approximation as depicted in Figure 5.2.




FIGURE E.2. Leading  $d-3$  eigenvalues of the observed Fisher information matrices depicted in Figures 5.1 and 5.2, plotted against a common scaling  $1/\alpha \propto \sigma^{-2}$ , for (a) the 405-dimensional approximation for rotavirus VP6, (b) the 4410-dimensional approximation for rotavirus VP6, and (c) the 3528-dimensional approximation for hemoglobin. Lines depict the median within each of the three tiers, and bands depict the 10th to 90th percentiles.

$$= Y_{k/2} C_{0,k/2-1,k/2-1}^{k,k/2,k/2} \left| u_{k/2-1}^{(k/2)C} \right|^2 = Y_{k/2} C_{0,k/2-1,k/2-1}^{k,k/2,k/2} \left( (v_{k/2-1}^{(k/2)C})^2 + (w_{k/2-1}^{(k/2)C})^2 \right).$$

So  $P_{k/2}/M_{k/2} = Y_{k/2} C_{0,k/2-1,k/2-1}^{k,k/2,k/2} \cdot v_{k/2-1}^{(k/2)C}$ , which is non-zero.

If  $k$  is odd, then  $\eta_0^{(ks)} = w_0^{(ks)}$ . In the  $3 \times 3$  submatrix corresponding to  $P_{[k/2]}$ , again each entry has at most 1 term depending on some variable of  $M_{[k/2]}$ . The below table records the quotient of this term by the corresponding variable.

	$v_{1,[k/2]}^{k,s}$	$w_{1,[k/2]}^{k,s}$	$w_0^{k,s}$
$C, C, [k/2], [k/2]$	$2Y_{[k/2]} C_{1,[k/2]-1,[k/2]-1}^{k,[k/2],[k/2]} v_{[k/2]-1}^{[k/2],C}$	$2Y_{[k/2]} C_{1,[k/2]-1,[k/2]-1}^{k,[k/2],[k/2]} w_{[k/2]-1}^{[k/2],C}$	0
$D, D, [k/2], [k/2]$	$2Y_{[k/2]} C_{1,[k/2]-1,[k/2]-1}^{k,[k/2],[k/2]} v_{[k/2]-1}^{[k/2],D}$	$2Y_{[k/2]} C_{1,[k/2]-1,[k/2]-1}^{k,[k/2],[k/2]} w_{[k/2]-1}^{[k/2],D}$	0
$A, C, [k/2], [k/2]$	0	0	$2Y_{[k/2]} C_{0,[k/2],[k/2]}^{k,[k/2],[k/2]} w_{[k/2]-1}^{[k/2],C}$

Then  $P_{[k/2]}/M_{[k/2]}$  is the determinant of this  $3 \times 3$  matrix, which is non-zero.

Combining the above, we have shown  $P/M \neq 0$  as desired. This completes the proof of part (b) also for  $k \geq 4$ .  $\square$

#### APPENDIX E. DETAILS OF THE NUMERICAL SIMULATIONS

We used rotavirus VP6 and hemoglobin maps publicly available on EMDB (EMDB-1461 and EMDB-3650). We recentered the rotavirus map EMDB-1461 to have center-of-mass at the origin, and zero-padded it into a cubical volume of dimensions  $141 \times 141 \times 141$ . The hemoglobin volume EMDB-3650 is already cubical. We centered the values of both maps to have mean value 0.

**Fourier quadrature.** We computed the Fourier transform  $\hat{f}$  of both maps over a spherical grid in the Fourier domain, using the FINUFFT library developed in [BMaK19]. Parametrizing Fourier space by spherical coordinates  $(\rho, \phi_1, \phi_2)$ , we computed  $\hat{f}$  on a linearly spaced grid of  $150 \times 250 \times 250$  points  $(\rho, \phi_1, \phi_2) \in [0, 1/R] \times [0, \pi] \times [0, 2\pi)$ , where  $R$  is the spatial-domain distance from the origin to the boundary of the cubical volume. All integrals in the Fourier domain were computed using the weighted quadrature defined by this discrete grid with weight proportional to  $\rho^2 \sin \phi_1$ .

**Low-pass filter and basis approximation.** For each frequency threshold  $v$ , we performed low-pass filtering by simple truncation of the Fourier transform to radii  $\rho \in [0, v]$ . We then iteratively defined radial functions  $z_1, \dots, z_S : [0, v] \rightarrow \mathbb{R}$  satisfying the orthogonality (4.4), so that for each  $s = 1, \dots, S$ , the partial basis  $\{z_1, \dots, z_s\}$  maximizes the total power of the projection of the Fourier transform  $\hat{f}$  onto the function space

$$\left\{ z_1(\rho)h_1(\phi_1, \phi_2) + \dots + z_s(\rho)h_s(\phi_1, \phi_2) : h_1, \dots, h_s \in L_2(\mathcal{S}^2, \mathbb{C}) \right\}.$$

In detail, let us denote  $u = (\phi_1, \phi_2)$ , and  $du = \sin \phi_1 d\phi_1 d\phi_2$  as the surface area measure on  $\mathcal{S}^2$ . Then the projection of  $\hat{f}$  onto the above space is defined explicitly by

$$h_i(u) = \int_0^v \hat{f}(\rho, u) \overline{z_i(\rho)} \cdot \rho^2 d\rho \quad \text{for each } i = 1, \dots, s.$$

The projected power is then

$$\begin{aligned} \text{Power} &= \int_{\mathcal{S}^2} \int_0^v \left| \sum_{i=1}^s z_i(\rho) h_i(u) \right|^2 \cdot \rho^2 d\rho du = \int_{\mathcal{S}^2} \sum_{i=1}^s |h_i(u)|^2 du \\ &= \int_0^v \int_0^v \sum_{i=1}^s \overline{z_i(\rho)} C(\rho, \rho') z_i(\rho') \cdot \rho^2 d\rho \cdot \rho'^2 d\rho' \end{aligned}$$

where we have defined the cross-covariance of  $\hat{f}(\rho, \cdot)$  and  $\hat{f}(\rho', \cdot)$  as

$$C(\rho, \rho') = \int_{\mathcal{S}^2} \hat{f}(\rho, u) \overline{\hat{f}(\rho', u)} du.$$

From the orthogonality (4.4), the maximizing functions  $z_1, \dots, z_S$  are such that  $\{\rho z_1(\rho), \dots, \rho z_S(\rho)\}$  are the  $S$  leading eigenfunctions (orthogonal with respect to the standard unweighted  $L_2$ -inner-product on  $[0, v]$ ) of the weighted cross-covariance kernel

$$K(g, h) = \int_0^v \int_0^v \overline{g(\rho)} C(\rho, \rho') h(\rho') \cdot \rho d\rho \cdot \rho' d\rho'.$$

We approximated this kernel  $K$  by its  $M \times M$  matrix discretization  $K_{\text{mat}} = (C(\rho, \rho') \rho \rho')_{\rho, \rho'}$  where  $M$  is the number of radial quadrature points  $\rho \in [0, v]$ . We approximated its eigenfunctions by the eigenvectors of  $K_{\text{mat}}$ . As the eigenfunctions of  $K$  correspond to  $\rho z_i(\rho)$ , we divided the eigenvectors of  $K_{\text{mat}}$  by  $\rho$  to obtain the values of the radial basis functions  $z_1, \dots, z_S$  along the above radial quadrature.

The final function basis over  $\mathbb{R}^3$  was obtained as a product of  $\{z_s : s = 1, \dots, S\}$  with the spherical harmonics as described in Section 4.2. We computed the spherical harmonics on the above  $250 \times 250$  quadrature points  $(\phi_1, \phi_2)$  using the implementation of the spherical harmonics provided by the `sph_harm` function in `scipy`. Finally, basis coefficients  $\theta_*$  were computed by integration in the Fourier domain, as approximated by the above quadrature.

**SO(3) quadrature.** We computed the empirical Hessian  $\nabla^2 R_n(\theta_*)$  by approximating the integral over  $\text{SO}(3)$  in the definition of the log-likelihood using a weighted discrete quadrature on  $\text{SO}(3)$ . Parametrizing  $\text{SO}(3)$  by the Euler angles  $(\alpha, \beta, \gamma)$ , we used a discrete grid of  $40 \times 40 \times 40$  values  $(\alpha, \beta, \gamma) \in [0, 2\pi) \times [0, \pi] \times [0, 2\pi)$ , with linearly-spaced grid points and equal weights for  $(\alpha, \gamma)$ . For  $\beta$  we also used linearly-spaced points  $\beta_1, \dots, \beta_{40}$ , with weights  $w_1, \dots, w_{40}$  computed by numerically solving the equations

$$\sum_{i=1}^{40} w_i D_{0,0}^{(l)}((0, \beta_i, 0)) = \begin{cases} 1, & \text{if } l = 0 \\ 0, & \text{otherwise} \end{cases} \quad (\text{E.1})$$

for  $0 \leq l < 40$ . Here  $D_{0,0}^{(l)}((0, \beta_i, 0))$  is the middle element of the Wigner-D matrix  $D^{(l)}(\mathbf{g})$ , evaluated at  $\mathbf{g} \in \text{SO}(3)$  having Euler angles  $(0, \beta_i, 0)$ . With proper normalization, this yields an approximate quadrature for band-limited functions over  $\text{SO}(3)$ .

We note that computing  $\nabla^2 R_n(\theta_*)$  for many samples is quite computationally intensive, scaling as  $O(nd^2 \cdot |\text{quad}|)$  where  $|\text{quad}|$  is the number of  $\text{SO}(3)$  quadrature points, and we were consequently limited in the size of this quadrature. Since this quadrature may provide an imperfect approximation to integration over true Haar measure on  $\text{SO}(3)$ , to mitigate some of the discretization effects, we generated samples  $y_1, \dots, y_n$  using random rotations also drawn from the weighted discrete distribution over  $\text{SO}(3)$  defined by this quadrature, rather than from the Haar measure. This does not fully address the numerical inaccuracy, but at least ensures that the true rotation for each sample  $y_i$  belongs to our discretization of  $\text{SO}(3)$ . This becomes numerically important for smaller values of the noise variance  $\sigma^2$ , when the posterior distribution of the rotation  $\mathbf{g} \in \text{SO}(3)$  for each sample becomes more localized over  $\text{SO}(3)$ .

We translated each quadrature point of  $\text{SO}(3)$  to a rotational element in  $\text{O}(d)$  via the representation described in Lemma D.3. We used the implementation of Wigner-D matrices provided by the third-party `quaternion` and `spherical_functions` Python libraries. These elements of  $\text{O}(d)$  were applied to  $\theta_*$  (with additive Gaussian noise) to generate the samples  $y_1, \dots, y_n$ , and the corresponding quadrature on  $\text{O}(d)$  was used also to compute the integrals over  $\text{SO}(3)$  in the analytic expression for  $\nabla^2 R_n(\theta_*)$ .

**Visualizations.** The molecular graphics in Figures 5.1 and 5.2 were rendered using the UCSF ChimeraX software [PGH<sup>+</sup>21]. We used atomic structures publicly available on PDB (PDB:1QHD and PDB:5NI1). We aligned the rotavirus VP6 structure PDB:1QHD to the 8.2Å-resolution map depicted in Figure E.1(a) using the ChimeraX “Fit in Map” tool. The hemoglobin structure PDB:5NI1 is already pre-aligned to the analyzed map.

## APPENDIX F. CRYOGENIC ELECTRON MICROSCOPY (CRYO-EM)

Cryo-EM is a technology for determining the spatial structure of macromolecules. In recent years, cryo-EM has become increasingly popular in structural biology. Thanks to technological advancement in hardware and algorithms in the last decade, cryo-EM now allows scientists to routinely recover structures at near-atomic resolutions. Unlike popular X-ray crystallography techniques for structure determination, cryo-EM does not require the samples to be crystallized. This gives cryo-EM an advantage, in particular, for molecules that are difficult to crystallize, and in heterogeneous samples.

In a typical cryo-EM study, a solution with the molecule of interest is flash-frozen in a thin layer of ice. The particles are sufficiently sparse and the layer sufficiently thin so that when viewed from above, the molecules rarely overlap. Each particle is trapped in the ice at an unknown random orientation. The sample is then inserted into the microscope; an electron beam is transmitted through the sample and then recorded by a camera. A detailed description of the procedure can be found, inter alia, in [GNC21].

The procedure produces a 2D tomographic projection of each particle. For simplicity, in this work we have omitted some effects such as the filters applied in the process (i.e. the contrast transfer function) and the problem of centering the particles, which is less crucial at lower resolutions. In addition, we assume the experimental distribution of viewing directions is uniform. A more comprehensive description can be found in [GNC21], and a more detailed mathematical description can be found in [BBS20]. The simplified imaging model is summarized by the equation

$$I(r_1, r_2) = \int_{\mathbb{R}} f(\mathbf{g}^{-1} \mathbf{r}) d\mathbf{r}_3 = \int_{\mathbb{R}} (\mathbf{g} \cdot f)((r_1, r_2, r_3)) d\mathbf{r}_3, \quad (\text{F.1})$$

where  $\mathbf{r} = (r_1, r_2, r_3)$ , the function  $f : \mathbb{R}^3 \rightarrow \mathbb{R}$  is the electric potential, and  $\mathbf{g} \in \text{SO}(3)$  is a rotation. We have expressed this equivalently in the main text as (4.11). In other words, the image is obtained by integrating the  $z$  axis of the volume rotated to viewing direction  $\mathbf{g}$  (which is not known to us). The interactions of the electrons with the sample lead to rapid deterioration in the quality of the sample, and very few electrons can be used to record the images before the sample becomes unusable. Therefore, the measurements are characterized by low signal-to-noise ratios.

Taking the Fourier transform of Eq. (F.1) with appropriate normalization yields

$$\hat{I}(\omega_1, \omega_2) = (\mathbf{g} \cdot \hat{f})((\omega_1, \omega_2, 0)), \quad (\text{F.2})$$

where  $\hat{I}$  is the Fourier transform of the image and  $\hat{f}$  is the Fourier transform of the density map. In other words, in the Fourier domain, the tomographic projection can be conveniently described as taking a slice of the Fourier transform of the volume, at the plane that passes through the origin and is perpendicular to the viewing direction. We have expressed this relation in our analysis as (D.51), which is characterized in our specific function bases by the projection operator  $\Pi^C$  in (D.57).

**F.1. Cryo-Electron Tomography (Cryo-ET, “Unprojected Cryo-EM”).** Cryo-ET is based on the same technology as cryo-EM. However, in cryo-ET several images are taken of each particle, with the sample tilted in a different direction for every image. As in the classic cryo-EM problem above, the relative rotation angles of the different particles are unknown. However, the relative tilt angles of images of *the same particle* are known. By the Fourier-slice relation (F.2), each image is a slice of the Fourier domain, and thus a dense set of slices from different viewing directions of *the same particle* can be used to reconstruct an entire 3D volume. Unfortunately, due to physical limitations, the tilt angles cannot cover all viewing directions, and a series of cryo-ET tilt images is typically noisier than a cryo-EM image. Thus cryo-ET provides a method of obtaining noisy 3D maps of individual particles, whose relative rotations across different particles are unknown as in the cryo-EM problem. Cryo-ET is more commonly used to study larger samples (e.g. entire cells), but is also used in the study of smaller particles. For additional information, see [EDP19, TB20]. In the main text, we have referred to this problem of reconstructing a map from *unprojected* and rotated 3D volumes also as the “unprojected cryo-EM” model.

A simplified model of cryo-ET, after the tilt series has been reassembled to a 3D function, has the form:

$$f_{\mathbf{g}}(\omega) = (\mathbf{g} \cdot f)(\omega) = f(\mathbf{g}^{-1} \cdot \omega), \quad (\text{F.3})$$

or, in the Fourier domain:

$$\hat{f}_{\mathbf{g}}(\omega) = (\mathbf{g} \cdot \hat{f})(\omega) = \hat{f}(\mathbf{g}^{-1} \cdot \omega). \quad (\text{F.4})$$

The (projected) cryo-EM model, is related to the cryo-ET model (“unprojected cryo-EM”) through the tomographic projection operator  $\Pi$ ; in the Fourier domain,  $\Pi$  has the form

$$(\Pi \hat{f})(\omega_1, \omega_2) = \hat{f}(\omega_1, \omega_2, 0). \quad (\text{F.5})$$

## REFERENCES

- [ABL<sup>+</sup>19] Emmanuel Abbe, Tamir Bendory, William Leeb, João M. Pereira, Nir Sharon, and Amit Singer. Multireference alignment is easier with an aperiodic translation distribution. *IEEE Transactions on Information Theory*, 65(6):3565–3584, 2019.
- [APS18] Emmanuel Abbe, João M Pereira, and Amit Singer. Estimation in the group action channel. In *2018 IEEE International Symposium on Information Theory (ISIT)*, pages 561–565. IEEE, 2018.
- [Aud04] Michele Audin. *Torus actions on symplectic manifolds*, volume 93. Springer Science & Business Media, 2004.
- [BBM<sup>+</sup>17] Tamir Bendory, Nicolas Boumal, Chao Ma, Zhizhen Zhao, and Amit Singer. Bispectrum inversion with application to multireference alignment. *IEEE Transactions on signal processing*, 66(4):1037–1050, 2017.
- [BBS20] Tamir Bendory, Alberto Bartsaghi, and Amit Singer. Single-particle cryo-electron microscopy: Mathematical theory, computational challenges, and opportunities. *IEEE signal processing magazine*, 37(2):58–76, 2020.
- [BBSK<sup>+</sup>17] Afonso S Bandeira, Ben Blum-Smith, Joe Kileel, Amelia Perry, Jonathan Weed, and Alexander S Wein. Estimation under group actions: Recovering orbits from invariants. *arXiv preprint arXiv:1712.10163*, 2017.
- [BJL<sup>+</sup>20] Tamir Bendory, Ariel Jaffe, William Leeb, Nir Sharon, and Amit Singer. Super-resolution multi-reference alignment. *arXiv preprint arXiv:2006.15354*, 2020.
- [BMaK19] Alexander H Barnett, Jeremy Magland, and Ludvig af Klinteberg. A parallel nonuniform fast fourier transform library based on an “exponential of semicircle” kernel. *SIAM Journal on Scientific Computing*, 41(5):C479–C504, 2019.
- [BMS13] Malte Breecken, Johannes Mittmann, and Nitin Saxena. Algebraic independence and blackbox identity testing. *Information and Computation*, 222:2–19, 2013.
- [BNWR20] Afonso S Bandeira, Jonathan Niles-Weed, and Philippe Rigollet. Optimal rates of estimation for multi-reference alignment. *Mathematical Statistics and Learning*, 2(1):25–75, 2020.
- [Böh13] Arno Böhm. *Quantum mechanics: foundations and applications*. Springer Science & Business Media, 2013.
- [Bru19] Victor-Emmanuel Brunel. Learning rates for Gaussian mixtures under group action. In *Conference on Learning Theory*, pages 471–491, 2019.
- [Bry16] Robert L Bryant. Second order families of special Lagrangian 3-folds. In *Perspectives in Riemannian Geometry, CRM Proceedings and Lecture Notes*, edited by Vestislav Apostolov, Andrew Dancer, Nigel Hitchin, and McKenzie Wang, vol. 40 (2006), American Mathematical Society. AMER MATHEMATICAL SOC, 2016.

- [DAC<sup>+</sup>88] Jacques Dubochet, Marc Adrian, Jiin-Ju Chang, Jean-Claude Homo, Jean Lepault, Alasdair W McDowell, and Patrick Schultz. Cryo-electron microscopy of vitrified specimens. *Quarterly reviews of biophysics*, 21(2):129–228, 1988.
- [EDP19] Fabian Eisenstein, Radostin Danev, and Martin Pilhofer. Improved applicability and robustness of fast cryo-electron tomography data acquisition. *Journal of structural biology*, 208(2):107–114, 2019.
- [Fra06] Joachim Frank. *Three-dimensional electron microscopy of macromolecular assemblies: visualization of biological molecules in their native state*. Oxford University Press, 2006.
- [FSWW20] Zhou Fan, Yi Sun, Tianhao Wang, and Yihong Wu. Likelihood landscape and maximum likelihood estimation for the discrete orbit recovery model. *Communications on Pure and Applied Mathematics*, 2020.
- [GHJY15] Rong Ge, Furong Huang, Chi Jin, and Yang Yuan. Escaping from saddle points—online stochastic gradient for tensor decomposition. In *Conference on Learning Theory*, pages 797–842, 2015.
- [GNC21] Robert M Glaeser, Eva Nogales, and Wah Chiu, editors. *Single-particle Cryo-EM of Biological Macromolecules*. 2053-2563. IOP Publishing, 2021.
- [Goo91] Colin Goodall. Procrustes methods in the statistical analysis of shape. *Journal of the Royal Statistical Society: Series B (Methodological)*, 53(2):285–321, 1991.
- [Gow75] John C Gower. Generalized procrustes analysis. *Psychometrika*, 40(1):33–51, 1975.
- [HBC<sup>+</sup>90] Richard Henderson, J M Baldwin, T A Ceska, F Zemlin, E A Beckmann, and Kenneth H Downing. Model for the structure of bacteriorhodopsin based on high-resolution electron cryo-microscopy. *Journal of molecular biology*, 213(4):899–929, 1990.
- [IH81] IA Ibragimov and RZ Has'minskii. *Statistical estimation: Asymptotic theory*. Springer, 1981.
- [JGN<sup>+</sup>17] Chi Jin, Rong Ge, Praneeth Netrapalli, Sham M Kakade, and Michael I Jordan. How to escape saddle points efficiently. In *Proceedings of the 34th International Conference on Machine Learning-Volume 70*, pages 1724–1732. JMLR. org, 2017.
- [Joh17] Iain M Johnstone. *Gaussian estimation: Sequence and wavelet models*. unpublished draft, 2017.
- [Kam80] Zvi Kam. The reconstruction of structure from electron micrographs of randomly oriented particles. *Journal of Theoretical Biology*, 82(1):15–39, 1980.
- [KB20] Anya Katsevich and Afonso S Bandeira. Likelihood maximization and moment matching in low snr gaussian mixture models. *Communications on Pure and Applied Mathematics*, 2020.
- [KRBD17] Maryam Khoshouei, Mazdak Radjainia, Wolfgang Baumeister, and Radostin Danev. Cryo-EM structure of haemoglobin at 3.2 Å determined with the Volta phase plate. *Nature communications*, 8(1):1–6, 2017.
- [Lan02] Serge Lang. *Algebra*. Springer, 3rd edition, 2002.
- [LM21] Allen Liu and Ankur Moitra. Algorithms from invariants: Smoothed analysis of orbit recovery over  $so(3)$ . *arXiv e-prints*, pages arXiv–2106, 2021.
- [LSJR16] Jason D Lee, Max Simchowitz, Michael I Jordan, and Benjamin Recht. Gradient descent only converges to minimizers. In *Conference on learning theory*, pages 1246–1257, 2016.
- [MBM18] Song Mei, Yu Bai, and Andrea Montanari. The landscape of empirical risk for nonconvex losses. *Annals of Statistics*, 46(6A):2747–2774, 2018.
- [Mit20] Boris Samuilovich Mityagin. The zero set of a real analytic function. *Matematicheskie Zametki*, 107(3):473–475, 2020.
- [MPN<sup>+</sup>01] Magali Mathieu, Isabelle Petitpas, Jorge Navaza, Jean Lepault, Evelyne Kohli, Pierre Pothier, BV Venkataram Prasad, Jean Cohen, and Félix A Rey. Atomic structure of the major capsid protein of rotavirus: Implications for the architecture of the virion. *The EMBO Journal*, 20(7):1485–1497, 2001.
- [PGH<sup>+</sup>21] Eric F Pettersen, Thomas D Goddard, Conrad C Huang, Elaine C Meng, Gregory S Couch, Tristan I Croll, John H Morris, and Thomas E Ferrin. Ucsf ChimeraX: Structure visualization for researchers, educators, and developers. *Protein Science*, 30(1):70–82, 2021.
- [PRFB17] Ali Punjani, John L Rubinstein, David J Fleet, and Marcus A Brubaker. cryoSPARC: algorithms for rapid unsupervised cryo-EM structure determination. *Nature methods*, 14(3):290–296, 2017.
- [PSB19] Thomas Pumis, Amit Singer, and Nicolas Boumal. The generalized orthogonal Procrustes problem in the high noise regime. *arXiv preprint arXiv:1907.01145*, 2019.
- [PV94] Vladimir L Popov and Ernest B Vinberg. Invariant theory. In *Algebraic geometry IV*, pages 123–278. Springer, 1994.
- [PWB<sup>+</sup>19] Amelia Perry, Jonathan Weed, Afonso S Bandeira, Philippe Rigollet, and Amit Singer. The sample complexity of multireference alignment. *SIAM Journal on Mathematics of Data Science*, 1(3):497–517, 2019.
- [RBO21] Elad Romanov, Tamir Bendory, and Or Ordentlich. Multi-reference alignment in high dimensions: sample complexity and phase transition. *SIAM Journal on Mathematics of Data Science*, 3(2):494–523, 2021.
- [Ros95] Morris Edgar Rose. *Elementary theory of angular momentum*. Courier Corporation, 1995.
- [Sch12] Sjors HW Scheres. RELION: implementation of a Bayesian approach to cryo-EM structure determination. *Journal of structural biology*, 180(3):519–530, 2012.
- [SGV<sup>+</sup>07] Sjors HW Scheres, Haixiao Gao, Mikel Valle, Gabor T Herman, Paul PB Eggermont, Joachim Frank, and Jose-Maria Carazo. Disentangling conformational states of macromolecules in 3D-EM through likelihood optimization. *Nature methods*, 4(1):27–29, 2007.
- [Sig98] Fred J Sigworth. A maximum-likelihood approach to single-particle image refinement. *Journal of structural biology*, 122(3):328–339, 1998.
- [SKK<sup>+</sup>20] Nir Sharon, Joe Kileel, Yuehaw Khoo, Boris Landa, and Amit Singer. Method of moments for 3D single particle ab initio modeling with non-uniform distribution of viewing angles. *Inverse Problems*, 36(4):044003, 2020.

- [SS20] Amit Singer and Fred J Sigworth. Computational methods for single-particle electron cryomicroscopy. *Annual Review of Biomedical Data Science*, 3, 2020.
- [TB20] Martin Turk and Wolfgang Baumeister. The promise and the challenges of cryo-electron tomography. *FEBS letters*, 594(20):3243–3261, 2020.
- [Tsy08] Alexandre B Tsybakov. *Introduction to nonparametric estimation*. Springer Science & Business Media, 2008.
- [WZ19] Yihong Wu and Harrison H Zhou. Randomly initialized EM algorithm for two-component Gaussian mixture achieves near optimality in  $o(\sqrt{n})$  iterations. *arXiv preprint arXiv:1908.10935*, 2019.
- [XHM16] Ji Xu, Daniel Hsu, and Arian Maleki. Global analysis of expectation maximization for mixtures of two Gaussians. In *Proceedings of the 30th International Conference on Neural Information Processing Systems*, pages 2684–2692, 2016.
- [ZSX<sup>+</sup>08] Xing Zhang, Ethan Settembre, Chen Xu, Philip R Dormitzer, Richard Bellamy, Stephen C Harrison, and Nikolaus Grigorieff. Near-atomic resolution using electron cryomicroscopy and single-particle reconstruction. *Proceedings of the National Academy of Sciences*, 105(6):1867–1872, 2008.