

# A-NUMERICAL RADIUS : NEW INEQUALITIES AND CHARACTERIZATION OF EQUALITIES

PINTU BHUNIA AND KALLOL PAUL

ABSTRACT. We develop new lower bounds for the  $A$ -numerical radius of semi-Hilbertian space operators, and applying these bounds we obtain upper bounds for the  $A$ -numerical radius of the commutators of operators. The bounds obtained here improve on the existing ones. Further, we provide characterizations for the equality of the existing  $A$ -numerical radius inequalities of semi-Hilbertian space operators.

## 1. INTRODUCTION

Let  $\mathcal{B}(\mathcal{H})$  denote the  $\mathbb{C}^*$ -algebra of all bounded linear operators on a complex Hilbert space  $\mathcal{H}$  with inner product  $\langle \cdot, \cdot \rangle$  and the corresponding norm  $\|\cdot\|$ . For  $T \in \mathcal{B}(\mathcal{H})$ , the range and the kernel of  $T$  are denoted by  $R(T)$  and  $N(T)$ , respectively. By  $\overline{R(T)}$ , we denote the norm closure of  $R(T)$ . The Hilbert-adjoint of  $T$  is denoted by  $T^*$ . Let  $A \in \mathcal{B}(\mathcal{H})$  be a positive operator, henceforth we reserve the symbol  $A$  for positive operator on  $\mathcal{H}$ . Clearly,  $A$  induces a positive semidefinite sesquilinear form  $\langle \cdot, \cdot \rangle_A : \mathcal{H} \times \mathcal{H} \rightarrow \mathbb{C}$ , defined by  $\langle x, y \rangle_A = \langle Ax, y \rangle$  for all  $x, y \in \mathcal{H}$ . This sesquilinear form induces a seminorm  $\|\cdot\|_A : \mathcal{H} \rightarrow \mathbb{R}^+$ , defined by  $\|x\|_A = \sqrt{\langle x, x \rangle_A}$  for all  $x \in \mathcal{H}$ . Clearly,  $\|\cdot\|_A$  is a norm if and only if  $A$  is injective, and  $(\mathcal{H}, \|\cdot\|_A)$  is complete if and only if  $R(A)$  is closed in  $\mathcal{H}$ . An operator  $T \in \mathcal{B}(\mathcal{H})$  is said to be  $A$ -bounded if there exists  $c > 0$  such that  $\|Tx\|_A \leq c\|x\|_A$  for all  $x \in \overline{R(A)}$ , and in this case

$$\|T\|_A = \sup_{x \in \overline{R(A)}, x \neq 0} \frac{\|Tx\|_A}{\|x\|_A} = \sup_{x \in \overline{R(A)}, \|x\|_A=1} \|Tx\|_A < +\infty.$$

Let  $\mathcal{B}^A(\mathcal{H})$  denote the collection of all  $A$ -bounded operators, i.e.,  $\mathcal{B}^A(\mathcal{H}) = \{T \in \mathcal{B}(\mathcal{H}) : \|T\|_A < +\infty\}$ . It is well known that  $\mathcal{B}^A(\mathcal{H})$  is, in general, not a subalgebra of  $\mathcal{B}(\mathcal{H})$ . Note that  $\|T\|_A = 0$  if and only if  $ATA = 0$ . The  $A$ -adjoint of  $T$ , if it exists, is defined as follows:

**Definition 1.1.** ([1]) For  $T \in \mathcal{B}(\mathcal{H})$ , an operator  $S \in \mathcal{B}(\mathcal{H})$  is said to be an  $A$ -adjoint of  $T$  if  $\langle Tx, y \rangle_A = \langle x, Sy \rangle_A$  for all  $x, y \in \mathcal{H}$ , i.e,  $AS = T^*A$ .

Let  $\mathcal{B}_A(\mathcal{H})$  denote the collection of all operators in  $\mathcal{B}(\mathcal{H})$ , which admit  $A$ -adjoint. By Douglas theorem [10], it follows that

$$\mathcal{B}_A(\mathcal{H}) = \{T \in \mathcal{B}(\mathcal{H}) : R(T^*A) \subseteq R(A)\}.$$

---

2010 *Mathematics Subject Classification.* 47A12; 47A30; 47A63.

*Key words and phrases.*  $A$ -numerical radius;  $A$ -operator seminorm; Semi-Hilbertian space; Inequality.

First author would like to thank UGC, Govt. of India for the financial support in the form of SRF.

For  $T \in \mathcal{B}_A(\mathcal{H})$ , the operator equation  $AX = T^*A$  has a unique solution, denoted by  $T^{\sharp A}$ , satisfying  $R(T^{\sharp A}) \subseteq \overline{R(A)}$ . Note that  $T^{\sharp A} = A^\dagger T^*A$ , where  $A^\dagger$  is the Moore-Penrose inverse of  $A$ . We note that  $\mathcal{B}_A(\mathcal{H}) (\subseteq \mathcal{B}^A(\mathcal{H}))$  is a sub-algebra of  $\mathcal{B}(\mathcal{H})$ . For  $T \in \mathcal{B}_A(\mathcal{H})$ , we have,  $AT^{\sharp A} = T^*A$  and  $N(T^{\sharp A}) = N(T^*A)$ . If  $T \in \mathcal{B}_A(\mathcal{H})$ , then  $T^{\sharp A} \in \mathcal{B}_A(\mathcal{H})$  and  $(T^{\sharp A})^{\sharp A} = P_{\overline{R(A)}}TP_{\overline{R(A)}}$  where  $P_{\overline{R(A)}}$  is the orthogonal projection onto  $\overline{R(A)}$ . An operator  $T \in \mathcal{B}_A(\mathcal{H})$  is said to be  $A$ -self-adjoint if  $AT$  is self-adjoint, i.e.,  $AT = T^*A$ . For further study on the  $A$ -adjoint operator, we refer to [1]. Note that, for  $T, S \in \mathcal{B}_A(\mathcal{H})$ ,  $(TS)^{\sharp A} = S^{\sharp A}T^{\sharp A}$ ,  $\|TS\|_A \leq \|T\|_A\|S\|_A$  and  $\|Tx\|_A \leq \|T\|_A\|x\|_A$  for all  $x \in \mathcal{H}$ . Clearly, for  $T \in \mathcal{B}_A(\mathcal{H})$ ,  $\|TT^{\sharp A}\|_A = \|T^{\sharp A}T\|_A = \|T^{\sharp A}\|_A^2 = \|T\|_A^2$ .

For  $T \in \mathcal{B}_A(\mathcal{H})$ , the  $A$ -numerical range of  $T$ , denoted by  $W_A(T)$ , is defined as

$$W_A(T) = \{\langle Tx, x \rangle_A : x \in \mathcal{H}, \|x\|_A = 1\}$$

and the  $A$ -numerical radius of  $T$ , denoted by  $w_A(T)$ , is defined as

$$w_A(T) = \sup\{|\langle Tx, x \rangle_A| : x \in \mathcal{H}, \|x\|_A = 1\}.$$

It was shown in [15] that for  $T \in \mathcal{B}_A(\mathcal{H})$ ,  $w_A(T) = \sup_{\theta \in \mathbb{R}} \left\| \frac{e^{i\theta}T + (e^{i\theta}T)^{\sharp A}}{2} \right\|_A$ . For  $T \in \mathcal{B}_A(\mathcal{H})$ , we have

$$\|T\|_A = \sup\{\|Tx\|_A : \|x\|_A = 1\} = \sup\{|\langle Tx, y \rangle_A| : \|x\|_A = \|y\|_A = 1\}.$$

It is well-known that  $w_A(\cdot)$  and  $\|\cdot\|_A$  are equivalent seminorms on  $\mathcal{B}_A(\mathcal{H})$  satisfying the inequality (see [2, Prop. 2.5])

$$\frac{1}{2}\|T\|_A \leq w_A(T) \leq \|T\|_A. \quad (1.1)$$

The inequalities in (1.1) are sharp (see [11]),  $w_A(T) = \frac{1}{2}\|T\|_A$  if  $AT^2 = 0$  and  $w_A(T) = \|T\|_A$  if  $AT = T^*A$ . An improvement of (1.1) is given in [12, 15], which is

$$\frac{1}{4}\|T^{\sharp A}T + TT^{\sharp A}\|_A \leq w_A^2(T) \leq \frac{1}{2}\|T^{\sharp A}T + TT^{\sharp A}\|_A. \quad (1.2)$$

More refinements in this direction are also given in [3, 4, 5, 6, 7, 13, 14]. Inspired by the inequalities obtained in the article [8, 9], here we obtain new refinements of the first inequalities in (1.1) and (1.2). By applying the new refinements developed here, we obtain upper bounds for the  $A$ -numerical radius of the commutators of operators. Also, we obtain characterizations for the equality of first inequalities in (1.1) and (1.2).

## 2. LOWER BOUNDS FOR $A$ -NUMERICAL RADIUS

We begin with the observation that any  $T \in \mathcal{B}_A(\mathcal{H})$  can be expressed as  $T = \Re_A(T) + i\Im_A(T)$ , where  $\Re_A(T) = \frac{T+T^{\sharp A}}{2}$  and  $\Im_A(T) = \frac{T-T^{\sharp A}}{2i}$ . It is easy to verify that  $\Re_A(T)$  and  $\Im_A(T)$  both are  $A$ -self-adjoint, i.e.,  $A\Re_A(T) = (\Re_A(T))^*A$  and  $A\Im_A(T) = (\Im_A(T))^*A$ . Therefore,  $w_A(\Re_A(T)) = \|\Re_A(T)\|_A$  and  $w_A(\Im_A(T)) = \|\Im_A(T)\|_A$ . Now, we are in a position to prove our first improvement.

**Theorem 2.1.** *If  $T \in \mathcal{B}_A(\mathcal{H})$ , then*

$$w_A(T) \geq \frac{\|T\|_A}{2} + \frac{|\|\Re_A(T)\|_A - \|\Im_A(T)\|_A|}{2}.$$

*Proof.* Let  $x \in \mathcal{H}$  with  $\|x\|_A = 1$ . Then from  $T = \Re_A(T) + i\Im_A(T)$ , we have

$$|\langle Tx, x \rangle_A|^2 = |\langle \Re_A(T)x, x \rangle_A|^2 + |\langle \Im_A(T)x, x \rangle_A|^2.$$

This implies that

$$|\langle Tx, x \rangle_A| \geq |\langle \Re_A(T)x, x \rangle_A| \text{ and } |\langle Tx, x \rangle_A| \geq |\langle \Im_A(T)x, x \rangle_A|.$$

Considering supremum over  $\|x\|_A = 1$ , we get

$$w_A(T) \geq \|\Re_A(T)\|_A \text{ and } w_A(T) \geq \|\Im_A(T)\|_A.$$

Hence,

$$\begin{aligned} w_A(T) &\geq \max\{\|\Re_A(T)\|_A, \|\Im_A(T)\|_A\} \\ &= \frac{\|\Re_A(T)\|_A + \|\Im_A(T)\|_A}{2} + \frac{\|\Re_A(T)\|_A - \|\Im_A(T)\|_A}{2} \\ &\geq \frac{\|\Re_A(T) + i\Im_A(T)\|_A}{2} + \frac{\|\Re_A(T)\|_A - \|\Im_A(T)\|_A}{2} \\ &= \frac{\|T\|_A}{2} + \frac{\|\Re_A(T)\|_A - \|\Im_A(T)\|_A}{2}. \end{aligned}$$

Thus, we complete the proof.  $\square$

**Remark 2.2.** Clearly, the inequality in Theorem 2.1 is sharper than the first inequality in (1.1), i.e.,  $w_A(T) \geq \frac{\|T\|_A}{2}$ .

In the next theorem we provide a characterization for the equality of lower bound of  $A$ -numerical radius mentioned in (1.1).

**Theorem 2.3.** *Let  $T \in \mathcal{B}_A(\mathcal{H})$ .*

(i) *If  $w_A(T) = \frac{\|T\|_A}{2}$ , then*

$$\|\Re_A(T)\|_A = \|\Im_A(T)\|_A = \frac{\|T\|_A}{2},$$

*but the converse is not necessarily true.*

(ii)  *$w_A(T) = \frac{\|T\|_A}{2}$  if and only if  $\|\Re_A(e^{i\theta}T)\|_A = \|\Im_A(e^{i\theta}T)\|_A = \frac{\|T\|_A}{2}$  for all  $\theta \in \mathbb{R}$ .*

*Proof.* (i) It follows from Theorem 2.1 that if  $w_A(T) = \frac{\|T\|_A}{2}$ , then

$$\|\Re_A(T)\|_A = \|\Im_A(T)\|_A.$$

Also, we get

$$\begin{aligned} \|\Re_A(T)\|_A \leq w_A(T) &= \frac{\|T\|_A}{2} = \frac{\|\Re_A(T) + i\Im_A(T)\|_A}{2} \leq \frac{\|\Re_A(T)\|_A + \|\Im_A(T)\|_A}{2} \\ &= \|\Re_A(T)\|_A. \end{aligned}$$

This implies that  $\|\Re_A(T)\|_A = \frac{\|T\|_A}{2}$ , and so  $\|\Im_A(T)\|_A = \frac{\|T\|_A}{2}$ .

(ii) The ‘‘if’’ part follows from  $w_A(T) = \sup_{\theta \in \mathbb{R}} \|\Re_A(e^{i\theta}T)\|_A$ , and so we only need to prove the ‘‘only if’’ part. Let  $w_A(T) = \frac{\|T\|_A}{2}$ . Clearly  $e^{i\theta}T \in \mathcal{B}_A(\mathcal{H})$  for all  $\theta \in \mathbb{R}$ . Now,  $w_A(e^{i\theta}T) = w_A(T)$  and  $\|e^{i\theta}T\|_A = \|T\|_A$  for all  $\theta \in \mathbb{R}$ . Therefore, it follows from (i) that  $\|\Re_A(e^{i\theta}T)\|_A = \|\Im_A(e^{i\theta}T)\|_A = \frac{\|T\|_A}{2}$  for all  $\theta \in \mathbb{R}$ .  $\square$

Our next improvement of the first inequality in (1.2) reads as follows.

**Theorem 2.4.** *If  $T \in \mathcal{B}_A(\mathcal{H})$ , then*

$$w_A(T) \geq \sqrt{\frac{1}{4} \|T^{\sharp_A}T + TT^{\sharp_A}\|_A + \frac{1}{2} | \|\Re_A(T)\|_A^2 - \|\Im_A(T)\|_A^2 |}.$$

*Proof.* We have  $w_A(T) \geq \|\Re_A(T)\|_A$  and  $w_A(T) \geq \|\Im_A(T)\|_A$  and so

$$\begin{aligned} w_A^2(T) &\geq \max \{ \|\Re_A(T)\|_A^2, \|\Im_A(T)\|_A^2 \} \\ &= \frac{\|\Re_A(T)\|_A^2 + \|\Im_A(T)\|_A^2}{2} + \frac{|\|\Re_A(T)\|_A^2 - \|\Im_A(T)\|_A^2|}{2} \\ &\geq \frac{\|(\Re_A(T))^2\|_A + \|(\Im_A(T))^2\|_A}{2} + \frac{|\|\Re_A(T)\|_A^2 - \|\Im_A(T)\|_A^2|}{2} \\ &\geq \frac{\|(\Re_A(T))^2 + (\Im_A(T))^2\|_A}{2} + \frac{|\|\Re_A(T)\|_A^2 - \|\Im_A(T)\|_A^2|}{2} \\ &= \frac{1}{4} \|T^{\sharp_A}T + TT^{\sharp_A}\|_A + \frac{1}{2} | \|\Re_A(T)\|_A^2 - \|\Im_A(T)\|_A^2 |. \end{aligned}$$

This completes the proof.  $\square$

**Remark 2.5.** Clearly, the inequality in Theorem 2.4 is sharper than the first inequality in (1.2), i.e.,  $w_A^2(T) \geq \frac{1}{4} \|T^{\sharp_A}T + TT^{\sharp_A}\|_A$ .

Next, we prove an equivalent condition for  $w_A(T) = \sqrt{\frac{1}{4} \|T^{\sharp_A}T + TT^{\sharp_A}\|_A}$ .

**Theorem 2.6.** *Let  $T \in \mathcal{B}_A(\mathcal{H})$ . Then,  $w_A(T) = \sqrt{\frac{1}{4} \|T^{\sharp_A}T + TT^{\sharp_A}\|_A}$  if and only if  $\|\Re_A(e^{i\theta}T)\|_A^2 = \|\Im_A(e^{i\theta}T)\|_A^2 = \frac{1}{4} \|T^{\sharp_A}T + TT^{\sharp_A}\|_A$  for all  $\theta \in \mathbb{R}$ .*

*Proof.* The ‘‘if’’ part is trivial, we only prove the ‘‘only if’’ part. Let  $w_A^2(T) = \frac{1}{4} \|T^{\sharp_A}T + TT^{\sharp_A}\|_A$ . Now,  $(\Re_A(e^{i\theta}T))^2 + (\Im_A(e^{i\theta}T))^2 = \frac{T^{\sharp_A}T + TT^{\sharp_A}}{2}$  for all  $\theta \in \mathbb{R}$ . Therefore, we have

$$\begin{aligned} \frac{1}{4} \|T^{\sharp_A}T + TT^{\sharp_A}\|_A &= \frac{1}{2} \left\| (\Re_A(e^{i\theta}T))^2 + (\Im_A(e^{i\theta}T))^2 \right\|_A \\ &\leq \frac{1}{2} \left( \|\Re_A(e^{i\theta}T)\|_A^2 + \|\Im_A(e^{i\theta}T)\|_A^2 \right) \\ &\leq w_A^2(T) = \frac{1}{4} \|T^{\sharp_A}T + TT^{\sharp_A}\|_A. \end{aligned}$$

Hence,  $\|\Re_A(e^{i\theta}T)\|_A^2 + \|\Im_A(e^{i\theta}T)\|_A^2 = \frac{1}{2} \|T^{\sharp_A}T + TT^{\sharp_A}\|_A$ . Now,  $\sup_{\theta \in \mathbb{R}} \|\Re_A(e^{i\theta}T)\|_A^2 = \frac{1}{4} \|T^{\sharp_A}T + TT^{\sharp_A}\|_A = \sup_{\theta \in \mathbb{R}} \|\Im_A(e^{i\theta}T)\|_A^2$ . Therefore,  $\|\Re_A(e^{i\theta}T)\|_A^2 = \|\Im_A(e^{i\theta}T)\|_A^2 = \frac{1}{4} \|T^{\sharp_A}T + TT^{\sharp_A}\|_A$  for all  $\theta \in \mathbb{R}$ .  $\square$

Now we obtain another characterizations for the equalities  $w_A(T) = \frac{1}{2} \|T\|_A$  and  $w_A(T) = \sqrt{\frac{1}{4} \|T^{\sharp_A}T + TT^{\sharp_A}\|_A}$ . First we need to prove the following lemma.

**Lemma 2.7.** *Let  $T \in \mathcal{B}_A(\mathcal{H})$ . Then,  $\|\Re_A(e^{i\theta}T)\|_A = k$  (i.e., a constant) for all  $\theta \in \mathbb{R}$  if and only if  $W_A(T)$  is a circular disk with center at the origin and radius  $k$ .*

*Proof.* The “if” part is trivial, we only prove the “only if” part. Let  $\|\Re_A(e^{i\theta}T)\|_A = k$  for all  $\theta \in \mathbb{R}$ . Then,  $\sup_{\|x\|_A=1} |\langle \Re_A(e^{i\theta}T)x, x \rangle_A| = k$  for all  $\theta \in \mathbb{R}$ , i.e.,  $\sup_{\|x\|_A=1} |Re(e^{i\theta} \langle Tx, x \rangle_A)| = k$  for all  $\theta \in \mathbb{R}$ . Thus, for each  $\theta \in \mathbb{R}$ , there exists a sequence  $\{x_n^\theta\} \subseteq \mathcal{H}$  with  $\|x_n^\theta\|_A = 1$  such that  $|Re(e^{i\theta} \langle Tx_n^\theta, x_n^\theta \rangle_A)| \rightarrow k$ . This implies that the boundary of  $W_A(T)$  must be a circle with center at the origin and radius  $k$ . Since  $W_A(T)$  is a convex subset of  $\mathbb{C}$  (see in [2, Th. 2.1]), so  $W_A(T)$  is a circular disk with center at the origin and radius  $k$ . □

**Theorem 2.8.** *Let  $T \in \mathcal{B}_A(\mathcal{H})$ . Then, the following results hold.*

(i)  $w_A(T) = \frac{1}{2}\|T\|_A$  if and only if  $W_A(T)$  is a circular disk with center at the origin and radius  $\frac{1}{2}\|T\|_A$ .

(ii)  $w_A(T) = \sqrt{\frac{1}{4}\|T^{\sharp_A}T + TT^{\sharp_A}\|_A}$  if and only if  $W_A(T)$  is a circular disk with center at the origin and radius  $\sqrt{\frac{1}{4}\|T^{\sharp_A}T + TT^{\sharp_A}\|_A}$ .

*Proof.* The proof of (i) and (ii) follow from Theorems 2.3 and 2.6, respectively, by using Lemma 2.7. □

Another improvement of the first inequality in (1.1) reads as:

**Theorem 2.9.** *If  $T \in \mathcal{B}_A(\mathcal{H})$ , then*

$$w_A(T) \geq \frac{\|T\|_A}{2} + \frac{|\|\Re_A(T) + \Im_A(T)\|_A - \|\Re_A(T) - \Im_A(T)\|_A|}{2\sqrt{2}}.$$

*Proof.* Let  $x \in \mathcal{H}$  with  $\|x\|_A = 1$ . Then, we have

$$\begin{aligned} |\langle Tx, x \rangle_A| &= \sqrt{|\langle \Re_A(T)x, x \rangle_A|^2 + |\langle \Im_A(T)x, x \rangle_A|^2} \\ &\geq \frac{1}{\sqrt{2}}(|\langle \Re_A(T)x, x \rangle_A| + |\langle \Im_A(T)x, x \rangle_A|) \\ &\geq \frac{1}{\sqrt{2}}|\langle (\Re_A(T) \pm \Im_A(T))x, x \rangle_A|. \end{aligned}$$

Taking supremum over  $\|x\|_A = 1$ , we get

$$w_A(T) \geq \frac{1}{\sqrt{2}}\|\Re_A(T) \pm \Im_A(T)\|_A.$$

Therefore, we have

$$\begin{aligned}
w_A(T) &\geq \frac{1}{\sqrt{2}} \max\{\|\Re_A(T) + \Im_A(T)\|_A, \|\Re_A(T) - \Im_A(T)\|_A\} \\
&= \frac{1}{\sqrt{2}} \left\{ \frac{\|\Re_A(T) + \Im_A(T)\|_A + \|\Re_A(T) - \Im_A(T)\|_A}{2} \right. \\
&\quad \left. + \frac{|\|\Re_A(T) + \Im_A(T)\|_A - \|\Re_A(T) - \Im_A(T)\|_A|}{2} \right\} \\
&\geq \frac{1}{\sqrt{2}} \left\{ \frac{\|(\Re_A(T) + \Im_A(T)) - i(\Re_A(T) - \Im_A(T))\|_A}{2} \right. \\
&\quad \left. + \frac{|\|\Re_A(T) + \Im_A(T)\|_A - \|\Re_A(T) - \Im_A(T)\|_A|}{2} \right\} \\
&= \frac{1}{\sqrt{2}} \left\{ \frac{\|(1-i)T\|_A}{2} + \frac{|\|\Re_A(T) + \Im_A(T)\|_A - \|\Re_A(T) - \Im_A(T)\|_A|}{2} \right\} \\
&= \frac{\|T\|_A}{2} + \frac{|\|\Re_A(T) + \Im_A(T)\|_A - \|\Re_A(T) - \Im_A(T)\|_A|}{2\sqrt{2}},
\end{aligned}$$

as desired.  $\square$

**Remark 2.10.** (i) Clearly, the inequality in Theorem 2.9 is sharper than the first inequality in (1.1), i.e.,  $w_A(T) \geq \frac{\|T\|_A}{2}$ .

(ii) The inequalities obtained in Theorem 2.1 and Theorem 2.9 are not comparable, in general.

Another refinement of the first inequality in (1.2) reads as follows:

**Theorem 2.11.** *If  $T \in \mathcal{B}_A(\mathcal{H})$ , then*

$$w_A(T) \geq \sqrt{\frac{\|T^{\sharp_A}T + TT^{\sharp_A}\|_A}{4} + \frac{|\|\Re_A(T) + \Im_A(T)\|_A^2 - \|\Re_A(T) - \Im_A(T)\|_A^2|}{4}}.$$

*Proof.* Following the proof of Theorem 2.9, we have

$$\begin{aligned}
w_A^2(T) &\geq \frac{1}{2} \max\{\|\Re_A(T) + \Im_A(T)\|_A^2, \|\Re_A(T) - \Im_A(T)\|_A^2\} \\
&= \frac{1}{2} \left\{ \frac{\|\Re_A(T) + \Im_A(T)\|_A^2 + \|\Re_A(T) - \Im_A(T)\|_A^2}{2} \right. \\
&\quad \left. + \frac{|\|\Re_A(T) + \Im_A(T)\|_A^2 - \|\Re_A(T) - \Im_A(T)\|_A^2|}{2} \right\} \\
&\geq \frac{1}{2} \left\{ \frac{\|(\Re_A(T) + \Im_A(T))^2 + (\Re_A(T) - \Im_A(T))^2\|_A}{2} \right. \\
&\quad \left. + \frac{|\|\Re_A(T) + \Im_A(T)\|_A^2 - \|\Re_A(T) - \Im_A(T)\|_A^2|}{2} \right\} \\
&= \frac{\|T^{\sharp_A}T + TT^{\sharp_A}\|_A}{4} + \frac{|\|\Re_A(T) + \Im_A(T)\|_A^2 - \|\Re_A(T) - \Im_A(T)\|_A^2|}{4}.
\end{aligned}$$

This completes the proof.  $\square$

**Remark 2.12.** (i) Clearly, the inequality in Theorem 2.11 is sharper than the first inequality in (1.2), i.e.,  $w_A^2(T) \geq \frac{1}{4} \|T^{\sharp_A}T + TT^{\sharp_A}\|_A$ .  
(ii) The inequalities obtained in Theorem 2.4 and Theorem 2.11 are not comparable, in general.

### 3. APPLICATIONS

In this section we obtain new inequalities for the  $A$ -numerical radius of the generalized commutators of operators by applying Theorems 2.4 and 2.11. First we prove the following lemma.

**Lemma 3.1.** *If  $T, X, Y \in \mathcal{B}_A(\mathcal{H})$ , then*

$$w_A(TX \pm YT) \leq \max\{\|X\|_A, \|Y\|_A\} \sqrt{2\|T^{\sharp_A}T + TT^{\sharp_A}\|_A}.$$

*Proof.* Let  $x \in \mathcal{H}$  with  $\|x\|_A = 1$  and  $\max\{\|X\|_A, \|Y\|_A\} \leq 1$ . Then by Cauchy Schwarz inequality, we get

$$\begin{aligned} |\langle (TX \pm YT)x, x \rangle_A| &\leq |\langle Xx, T^{\sharp_A}x \rangle_A| + |\langle Tx, Y^{\sharp_A}x \rangle_A| \\ &\leq \|T^{\sharp_A}x\| + \|Tx\| \\ &\leq \sqrt{2} (\|T^{\sharp_A}x\|_A^2 + \|Tx\|_A^2)^{\frac{1}{2}} \\ &\leq \sqrt{2} \|T^{\sharp_A}T + TT^{\sharp_A}\|_A^{\frac{1}{2}}. \end{aligned}$$

Therefore, taking supremum over  $\|x\|_A = 1$ , we get

$$w_A(TX \pm YT) \leq \sqrt{2\|T^{\sharp_A}T + TT^{\sharp_A}\|_A}.$$

If  $X, Y \in \mathcal{B}_A(\mathcal{H})$  are arbitrary with  $\max\{\|X\|_A, \|Y\|_A\} \neq 0$ , then it follows from the above inequality that

$$w_A(TX \pm YT) \leq \max\{\|X\|_A, \|Y\|_A\} \sqrt{2\|T^{\sharp_A}T + TT^{\sharp_A}\|_A}.$$

Also, if  $\max\{\|X\|_A, \|Y\|_A\} = 0$ , then the above inequality holds trivially. This completes the proof.  $\square$

**Theorem 3.2.** *If  $T, X, Y \in \mathcal{B}_A(\mathcal{H})$ , then*

$$(i) \ w_A(TX \pm YT) \leq 2\sqrt{2} \max\{\|X\|_A, \|Y\|_A\} \sqrt{w_A^2(T) - \frac{|\|\Re_A(T)\|_A^2 - \|\Im_A(T)\|_A^2|}{2}}.$$

and

$$(ii) \ w_A(TX \pm YT) \leq 2\sqrt{2} \max\{\|X\|_A, \|Y\|_A\} \sqrt{w_A^2(T) - \frac{|\|\Re_A(T) + \Im_A(T)\|_A^2 - \|\Re_A(T) - \Im_A(T)\|_A^2|}{4}}.$$

*Proof.* By applying the inequalities in Theorem 2.4 and Theorem 2.11 in Lemma 3.1, we have (i) and (ii), respectively.  $\square$

It should be mentioned here that the inequalities (i) and (ii) in Theorem 3.2 are not comparable, in general.

Considering  $X = Y = S$  in Theorem 3.2, we get the following corollary.

**Corollary 3.3.** *If  $T, S \in \mathcal{B}_A(\mathcal{H})$ , then*

$$(i) \quad w_A(TS \pm ST) \leq 2\sqrt{2}\|S\|_A \sqrt{w_A^2(T) - \frac{|\|\Re_A(T)\|_A^2 - \|\Im_A(T)\|_A^2|}{2}}.$$

and

$$(ii) \quad w_A(TS \pm ST) \leq 2\sqrt{2}\|S\|_A \sqrt{w_A^2(T) - \frac{|\|\Re_A(T) + \Im_A(T)\|_A^2 - \|\Re_A(T) - \Im_A(T)\|_A^2|}{4}}.$$

Now, interchanging  $T$  and  $S$  in Corollary 3.3 (i), we get

$$w_A(TS \pm ST) \leq 2\sqrt{2} \min\{\alpha_1, \alpha_2\}, \quad (3.1)$$

where

$$\alpha_1 = \|S\|_A \sqrt{w_A^2(T) - \frac{|\|\Re_A(T)\|_A^2 - \|\Im_A(T)\|_A^2|}{2}},$$

$$\alpha_2 = \|T\|_A \sqrt{w_A^2(S) - \frac{|\|\Re_A(S)\|_A^2 - \|\Im_A(S)\|_A^2|}{2}}.$$

Also, interchanging  $T$  and  $S$  in Corollary 3.3 (ii), we get

$$w_A(TS \pm ST) \leq 2\sqrt{2} \min\{\beta_1, \beta_2\}, \quad (3.2)$$

where

$$\beta_1 = \|S\|_A \sqrt{w_A^2(T) - \frac{|\|\Re_A(T) + \Im_A(T)\|_A^2 - \|\Re_A(T) - \Im_A(T)\|_A^2|}{4}},$$

$$\beta_2 = \|T\|_A \sqrt{w_A^2(S) - \frac{|\|\Re_A(S) + \Im_A(S)\|_A^2 - \|\Re_A(S) - \Im_A(S)\|_A^2|}{4}}.$$

**Remark 3.4.** In [15, Th. 4.2], Zamani proved that if  $T, S \in \mathcal{B}_A(\mathcal{H})$ , then

$$w_A(TS \pm ST) \leq 2\sqrt{2} \min\{\|T\|_A w_A(S), \|S\|_A w_A(T)\}.$$

Clearly, both the inequalities in (3.1) and (3.2) are stronger than the inequality in [15, Th. 4.2].

### Data Availability Statement.

Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

### REFERENCES

1. M.L. Arias, G. Corach and M.C. Gonzalez, Partial isometries in semi-Hilbertian spaces, *Linear Algebra Appl.* 428 (2008) 1460-1475.
2. H. Baklouti, K. Feki and O.A.M. Sid Ahmed, Joint numerical ranges of operators in semi-Hilbertian spaces, *Linear Algebra Appl.* 555 (2018) 266-284.
3. P. Bhunia, K. Feki and K. Paul, A-Numerical radius orthogonality and parallelism of semi-Hilbertian space operators and their applications, *Bull. Iran. Math. Soc.* 47 (2021) 435-457.
4. P. Bhunia, R.K. Nayak and K. Paul, Improvement of A-numerical radius inequalities of semi-Hilbertian space operators, *Results Math.* 76, 120 (2021). <https://doi.org/10.1007/s00025-021-01439-w>
5. P. Bhunia, K. Paul and R.K. Nayak, On inequalities for A-numerical radius of operators, *Electron. J. Linear Algebra* 36 (2020) 143-157.



6. P. Bhunia, R.K. Nayak and K. Paul, Refinements of A-numerical radius inequalities and their applications, Adv. Oper. Theory 5 (2020) 1498-1511.
7. P. Bhunia and K. Paul, Some improvement of numerical radius inequalities of operators and operator matrices, Linear Multilinear Algebra (2020). <https://doi.org/10.1080/03081087.2020.1781037>.
8. P. Bhunia and K. Paul, Development of inequality and characterization of equality conditions for the numerical radius, (2021). <https://arxiv.org/abs/2105.09715>
9. P. Bhunia, S. Jana and K. Paul, Refined inequalities for the numerical radius of Hilbert space operators, (2021). <http://arxiv.org/abs/2106.13949>
10. R.G. Douglas, On majorization, factorization and range inclusion of operators in Hilbert space, Proc. Amer. Math. Soc. 17 (1966) 413-416.
11. K. Feki, Spectral radius of semi-Hilbertian space operators and its applications, Ann. Funct. Anal. 11 (2020) 929-946.
12. K. Feki, A note on the A-numerical radius of operators in semi-Hilbert spaces, Arch. Math. 115 (2020) 535-544.
13. M.S. Moslehian, Q. Xu and A. Zamani, Seminorm and numerical radius inequalities of operators in semi-Hilbertian spaces, Linear Algebra Appl. 591 (2020) 299-321.
14. N.C. Rout, S. Sahoo and D. Mishra, On A-numerical radius inequalities for  $2 \times 2$  operator matrices, Linear Multilinear Algebra (2020). <https://doi.org/10.1080/03081087.2020.1810201>
15. A. Zamani, A-numerical radius inequalities for semi-Hilbertian space operators, Linear Algebra Appl. 578 (2019) 159-183.

(BHUNIA) DEPARTMENT OF MATHEMATICS, JADAVPUR UNIVERSITY, KOLKATA 700032, WEST BENGAL, INDIA

*Email address:* pintubhunia5206@gmail.com; pbhunia.math.rs@jadavpuruniversity.in

(PAUL) DEPARTMENT OF MATHEMATICS, JADAVPUR UNIVERSITY, KOLKATA 700032, WEST BENGAL, INDIA

*Email address:* kalloldada@gmail.com; kallol.paul@jadavpuruniversity.in