

# Vertex and edge metric dimensions of cacti

Jelena Sedlar<sup>1,3</sup>,  
Riste Škrekovski<sup>2,3</sup>

<sup>1</sup> *University of Split, Faculty of civil engineering, architecture and geodesy, Croatia*

<sup>2</sup> *University of Ljubljana, FMF, 1000 Ljubljana, Slovenia*

<sup>3</sup> *Faculty of Information Studies, 8000 Novo Mesto, Slovenia*

January 12, 2022

## Abstract

In a graph  $G$ , a vertex (resp. an edge) metric generator is a set of vertices  $S$  such that any pair of vertices (resp. edges) from  $G$  is distinguished by at least one vertex from  $S$ . The cardinality of a smallest vertex (resp. edge) metric generator is the vertex (resp. edge) metric dimension of  $G$ . In [19] we determined the vertex (resp. edge) metric dimension of unicyclic graphs and that it takes its value from two consecutive integers. Therein, several cycle configurations were introduced and the vertex (resp. edge) metric dimension takes the greater of the two consecutive values only if any of these configurations is present in the graph. In this paper we extend the result to cactus graphs i.e. graphs in which all cycles are pairwise edge disjoint. We do so by defining a unicyclic subgraph of  $G$  for every cycle of  $G$  and applying the already introduced approach for unicyclic graphs which involves the configurations. The obtained results enable us to prove the cycle rank conjecture for cacti. They also yield a simple upper bound on metric dimensions of cactus graphs and we conclude the paper by conjecturing that the same upper bound holds in general.

*Keywords:* vertex metric dimension; edge metric dimension; cactus graphs, zero forcing number, cycle rank conjecture.

*AMS Subject Classification numbers:* 05C12; 05C76

## 1 Introduction

The concept of metric dimension was first studied in the context of navigation system in various graphical networks [7]. There the robot moves from one vertex of the network to another, and some of the vertices are considered to be a landmark which helps a robot to establish its position in a network. Then the problem of establishing the smallest set of landmarks in a network becomes a problem of determining a smallest metric generator in a graph [12].

Another interesting application is in chemistry where the structure of a chemical compound is frequently viewed as a set of functional groups arrayed on a substructure. This can be modeled as a labeled graph where the vertex and edge labels specify the atom and bond types, respectively, and the functional groups and substructure are simply subgraphs of the labeled graph representation. Determining the pharmacological activities related to the feature of compounds relies on the investigation of the same functional groups for two different compounds at the same point [3]. Various other aspects of the notion were studied [1, 5, 13, 15] and a lot of research was dedicated to the behaviour of metric dimension with respect to various graph operations [2, 3, 17, 23].

In this paper, we consider only simple and connected graphs. By  $d(u, v)$  we denote the distance between a pair of vertices  $u$  and  $v$  in a graph  $G$ . A vertex  $s$  from  $G$  *distinguishes* or *resolves* a pair of vertices  $u$  and  $v$  from  $G$  if  $d(s, u) \neq d(s, v)$ . We say that a set of vertices  $S \subseteq V(G)$  is a *vertex metric generator*, if every pair of vertices in  $G$  is distinguished by at least one vertex from  $S$ . The *vertex metric dimension* of  $G$ , denoted by  $\dim(G)$ , is the cardinality of a smallest vertex generator in  $G$ . This variant of metric dimension, as it was introduced first, is sometimes called only metric dimension and the prefix "vertex" is omitted.

In [11] it was noticed that there are graphs in which none of the smallest metric generators distinguishes all pairs of edges, so this was the motivation to introduce the notion of the edge metric generator and dimension, particularly to study the relation between  $\dim(G)$  and  $\text{edim}(G)$ .

The distance  $d(u, vw)$  between a vertex  $u$  and an edge  $vw$  in a graph  $G$  is defined by  $d(u, vw) = \min\{d(u, v), d(u, w)\}$ . Recently, two more variants of metric dimension were introduced, namely the edge metric dimension and the mixed metric dimension of a graph  $G$ . Similarly as above, a vertex  $s \in V(G)$  *distinguishes* two edges  $e, f \in E(G)$  if  $d(s, e) \neq d(s, f)$ . So, a set  $S \subseteq V(G)$  is an *edge metric generator* if every pair of vertices is distinguished by at least one vertex from  $S$ , and the cardinality of a smallest such set is called the *edge metric dimension* and denoted by  $\text{edim}(G)$ . Finally, a set  $S \subseteq V(G)$  is a *mixed metric generator* if it distinguishes all pairs from  $V(G) \cup E(G)$ , and the *mixed metric dimension*, denoted by  $\text{mdim}(G)$ , is defined as the cardinality of a smallest such set in  $G$ .

This new variant also attracted a lot of attention [6, 8, 16, 24, 25, 26], with one particular direction of research being the study of unicyclic graphs and the relation of the two dimensions on them [14, 18, 19]. The mixed metric dimension is then a natural next step, as it unifies these two concepts. It was introduced in [10] and further studied in [21, 20]. A wider and systematic introduction to these three variants of metric dimension can be found in [9].

In this paper we establish the vertex and the edge metric dimension of cactus graph, using the approach from [19] where the two dimensions were established for unicyclic graphs. The extension is not straightforward, as in cactus graphs a problem with indistinguishable pairs of edges and vertices may arise from connecting two cycles, so additional condition will have to be introduced.

## 2 Preliminaries

A *cactus* graph is any graph in which all cycles are pairwise edge disjoint. Let  $G$  be a cactus graph with cycles  $C_1, \dots, C_c$  and let  $g_i$  denote the length of a cycle  $C_i$  in  $G$ . For a vertex  $v$  of a cycle  $C_i$ , denote by  $T_v(C_i)$  the connected component of  $G - E(C_i)$  which contains  $v$ . If  $G$  is a unicyclic graph, then  $T_v(C_i)$  is a tree, otherwise  $T_v(C_i)$  may contain a cycle. When no confusion arises from that, we will use the abbreviated notation  $T_v$ . A *thread* hanging at a vertex  $v \in V(G)$  of degree  $\geq 3$  is any path  $u_1 u_2 \dots u_k$  such that  $u_1$  is a leaf,  $u_2, \dots, u_k$  are of degree 2, and  $u_k$  is connected to  $v$  by an edge. The number of threads hanging at  $v$  is denoted by  $\ell(v)$ .

We say that a vertex  $v \in V(C_i)$  is *branch-active* if  $\deg(v) \geq 4$  or  $T_v$  contains a vertex of degree  $\geq 3$  distinct from  $v$ . We denote the number of branch-active vertices on  $C_i$  by  $b(C_i)$ . If a vertex  $v$  from a cycle  $C_i$  is branch-active, then  $T_v$  contains both a pair of vertices and a pair of edges which are not distinguished by any vertex outside  $T_v$ , see Figure 1.




Figure 1: A cactus graph with two cycles. On the cycle  $C$  vertices  $v$  and  $w$  are branch-active, and a pair of vertices is marked in  $T_v$  and  $T_w$  which is not distinguished by any vertex outside  $T_v$  and  $T_w$ , respectively.

Now, we will introduce a property called "branch-resolving" which a set of vertices  $S \subseteq V(G)$  must possess in order to avoid this problem of non distinguished vertices (resp. edges) due to branching. First, a thread hanging at a vertex  $v$  of degree  $\geq 3$  is  *$S$ -free* if it does not contain a vertex from  $S$ . Now, a set of vertices  $S \subseteq V(G)$  is *branch-resolving* if at most one  $S$ -free thread is hanging at every vertex  $v \in V(G)$  of degree  $\geq 3$ . Therefore, for every branch-resolving set  $S$  it holds that  $|S| \geq L(G)$  where

$$L(G) = \sum_{v \in V(G), \ell(v) > 1} (\ell(v) - 1).$$

It is known in literature [11, 12] that for a tree  $T$  it holds that  $\dim(G) = \text{edim}(G) = L(G)$ .

Also, given a set of vertices  $S \subseteq V(G)$ , we say that a vertex  $v \in V(C_i)$  is  *$S$ -active* if  $T_v$  contains a vertex from  $S$ . The number of  $S$ -active vertices on a cycle  $C_i$  is denoted

by  $a_S(C_i)$ . If  $a_S(C_i) \geq 2$  for every cycle  $C_i$  in  $G$ , then we say the set  $S$  is *biactive*. For a biactive branch-resolving set  $S$  the following holds: if a vertex  $v$  from a cycle  $C_i$  is branch-active, then  $T_v$  contains a vertex with two threads hanging at it or  $T_v$  contains a cycle, either way  $T_v$  contains a vertex from  $S$ , so  $v$  is  $S$ -active. Therefore, for a biactive branch-resolving set  $S$  we have  $a_S(C_i) \geq b(C_i)$  for every  $i$ .

**Lemma 1** *Let  $G$  be a cactus graph and let  $S \subseteq V(G)$  be a set of vertices in  $G$ . If  $S$  is a vertex (resp. an edge) metric generator, then  $S$  is a biactive branch-resolving set.*

**Proof.** Suppose to the contrary that a vertex (resp. an edge) metric generator  $S$  is not a biactive branch-resolving set. If  $S$  is not branch-resolving, then there exists a vertex  $v$  of degree  $\geq 3$  and two threads hanging at  $v$  which do not contain a vertex from  $S$ . Let  $v_1$  and  $v_2$  be two neighbors of  $v$ , each belonging to one of these two threads. Then  $v_1$  and  $v_2$  (resp.  $v_1v$  and  $v_2v$ ) are not distinguished by  $S$ , a contradiction.

Assume now that  $S$  is not biactive. We may assume that  $G$  has at least one cycle, otherwise  $G$  is a tree and there is nothing to prove. Notice that an empty set  $S$  cannot be either a vertex or an edge metric generator in a cactus graph unless  $G = K_2$  but then it is a tree. Therefore, if  $S$  is not biactive, there must exist a cycle  $C_i$  with precisely one  $S$ -active vertex  $v$  and let  $v_1$  and  $v_2$  be the two neighbors of  $v$  on  $C_i$ . Then  $v_1$  and  $v_2$  (resp.  $v_1v$  and  $v_2v$ ) are not distinguished by  $S$ , a contradiction. ■

The above lemma gives us a necessary condition for  $S$  to be a vertex (resp. an edge) metric generator in a cactus graph. In [19], a more elaborate condition for unicyclic graphs was established, which is both necessary and sufficient. In this paper we will extend that condition to cactus graphs, but to do so we first need to introduce the following definitions from [19]. Let  $C_i$  be a cycle in a cactus graph  $G$  and let  $v_i, v_j$  and  $v_k$  be three vertices of  $C_i$ , we say that  $v_i, v_j$  and  $v_k$  are a *geodesic triple* on  $C_i$  if

$$d(v_i, v_j) + d(v_j, v_k) + d(v_i, v_k) = |V(C_i)|.$$

It was shown in [18] that a biactive branch-resolving set with a geodesic triple of  $S$ -active vertices on every cycle is both a vertex and an edge metric generator. This result is useful for bounding the dimensions from above. Also, we need the definition of the five graph configurations from [19].

**Definition 2** *Let  $G$  be a cactus graph,  $C$  a cycle in  $G$  of the length  $g$ , and  $S$  a biactive branch-resolving set in  $G$ . We say that  $C = v_0v_1 \cdots v_{g-1}$  is canonically labeled with respect to  $S$  if  $v_0$  is  $S$ -active and  $k = \max\{i : v_i \text{ is } S\text{-active}\}$  is as small as possible.*

Let us now introduce five configurations which a cactus graph can contain with respect to a biactive branch-resolving set  $S$ . All these configurations are illustrated by Figure 2.

**Definition 3** *Let  $G$  be a cactus graph,  $C$  a canonically labeled cycle in  $G$  of the length  $g$ , and  $S$  a biactive branch-resolving set in  $G$ . We say that the cycle  $C$  with respect to  $S$  contains configurations:*

*A. If  $a_S(C) = 2$ ,  $g$  is even, and  $k = g/2$ ;*




Figure 2: All six graphs shown in this figure are unicyclic graphs with a biactive branch-resolving set  $S$  comprised of vertices  $s_i$ . Vertices on a cycle which are  $S$ -active are marked by a dashed circle and connected to its antipodal vertices by a dashed line. Each graph in this figure contains at least one of the five configurations, namely: a)  $\mathcal{A}$ , b)  $\mathcal{B}$  and also  $\mathcal{D}$ , c)  $\mathcal{C}$ , d)  $\mathcal{D}$ , e)  $\mathcal{E}$  on even cycle and also  $\mathcal{C}$ , f)  $\mathcal{E}$  on odd cycle. A pair  $x$  and  $x'$  of undistinguished vertices and/or edges is highlighted in each of the graphs. Notice that the graph in b) which contains  $\mathcal{B}$  consequently also contains  $\mathcal{D}$ , but graph in d) which contains  $\mathcal{D}$  does not contain  $\mathcal{B}$ . Similarly, the graph in e) which contains  $\mathcal{E}$  also contains  $\mathcal{C}$ , but the graph from c) which contains  $\mathcal{C}$  does not contain  $\mathcal{E}$ .

- B.* If  $k \leq \lfloor g/2 \rfloor - 1$  and there is an  $S$ -free thread hanging at a vertex  $v_i$  for some  $i \in [k, \lfloor g/2 \rfloor - 1] \cup [\lfloor g/2 \rfloor + k + 1, g - 1] \cup \{0\}$ ;
- C.* If  $a_S(C) = 2$ ,  $g$  is even,  $k \leq g/2$  and there is an  $S$ -free thread of the length  $\geq g/2 - k$  hanging at a vertex  $v_i$  for some  $i \in [0, k]$ ;
- D.* If  $k \leq \lceil g/2 \rceil - 1$  and there is an  $S$ -free thread hanging at a vertex  $v_i$  for some  $i \in [k, \lceil g/2 \rceil - 1] \cup [\lceil g/2 \rceil + k + 1, g - 1] \cup \{0\}$ ;
- E.* If  $a_S(C) = 2$  and there is an  $S$ -free thread of the length  $\geq \lfloor g/2 \rfloor - k + 1$  hanging at a vertex  $v_i$  with  $i \in [0, k]$ . Moreover, if  $g$  is even, an  $S$ -free thread must be hanging at the vertex  $v_j$  with  $j = g/2 + k - i$ .




Figure 3: A cactus graph  $G$  from Example 1. A smallest biactive branch-resolving set  $S = \{s_1, s_2, s_3, s_4, s_5\}$  is marked in the figure by squares. Vertices on the cycles which are  $S$ -active are marked by a dashed circle. Since every cycle contains a configuration with respect to  $S$ , for each cycle  $C_i$  there is a pair of vertices and/or edges  $x_i$  and  $x'_i$  which are not distinguished by  $S$ , these pairs are also highlighted in the figure.

Notice that only an even cycle can contain configuration  $\mathcal{A}$  or  $\mathcal{C}$ . Also, configurations  $\mathcal{B}$  and  $\mathcal{D}$  are almost the same, they differ only if  $C$  is odd where the index  $i$  can take

two more values in  $\mathcal{D}$  than in  $\mathcal{B}$ . Finally, for configurations  $\mathcal{A}$ ,  $\mathcal{C}$ , and  $\mathcal{E}$  it holds that  $a_S(C) = 2$ , so there are only two  $S$ -active vertices on the cycle  $C$  and hence no geodesic triple of  $S$ -active vertices. On the other hand, for configurations  $\mathcal{B}$  and  $\mathcal{D}$  there might be more than two  $S$ -active vertices on the cycle  $C$ , but the bounds  $k \leq \lfloor g/2 \rfloor - 1$  and  $k \leq \lceil g/2 \rceil - 1$  again imply there is no geodesic triple of  $S$ -active vertices on  $C$ . Therefore, we can state the following observation which is useful for constructing metric generators.

**Observation 4** *If there is a geodesic triple of  $S$ -active vertices on a cycle  $C$  of a cactus graph  $G$ , then  $C$  does not contain any of the configurations  $\mathcal{A}$ ,  $\mathcal{B}$ ,  $\mathcal{C}$ ,  $\mathcal{D}$ , and  $\mathcal{E}$  with respect to  $S$ .*

The following result regarding configurations  $\mathcal{A}$ ,  $\mathcal{B}$ ,  $\mathcal{C}$ ,  $\mathcal{D}$ , and  $\mathcal{E}$  was established for unicyclic graphs (see Lemmas 6, 7, 13 and 14 from [19]).

**Theorem 5** *Let  $G$  be a unicyclic graph with the cycle  $C$  and let  $S$  be a biactive branch-resolving set in  $G$ . The set  $S$  is a vertex (resp. an edge) metric generator if and only if  $C$  does not contain any of configurations  $\mathcal{A}$ ,  $\mathcal{B}$ , and  $\mathcal{C}$  (resp.  $\mathcal{A}$ ,  $\mathcal{D}$ , and  $\mathcal{E}$ ) with respect to  $S$ .*

In this paper we will extend this result to cactus graphs and then use it to determine the exact value of the vertex and the edge metric dimensions of such graphs. We first give an example how this approach with configurations can be extended to cactus graphs.

**Example 1** *Let  $G$  be the cactus graph from Figure 3. The graph  $G$  contains six cycles and the set  $S = \{s_1, s_2, s_3, s_4, s_5\}$  is a smallest biactive branch-resolving set in  $G$ . In the figure the set of  $S$ -active vertices on each cycle is marked by a dashed circle. The cycle  $C_1$  (resp.  $C_2$ ,  $C_3$ ,  $C_4$ ,  $C_5$ ) with respect to  $S$  contains configuration  $\mathcal{A}$  (resp.  $\mathcal{B}$  and also  $\mathcal{D}$ ,  $\mathcal{C}$ ,  $\mathcal{E}$  on odd cycle,  $\mathcal{E}$  on even cycle), so in each of these cycles there is a pair of vertices and/or edges  $x_i$  and  $x'_i$  which is not distinguished by  $S$ . The cycle  $C_6$  does not contain any of the five configurations as there is a geodesic triple of  $S$ -active vertices on  $C_6$ , so all pairs of vertices and all pair of edges in  $C_6$  are distinguished by  $S$ .*

Besides this configuration approach, in cactus graphs an additional condition will have to be introduced for the situation when a pair of cycles share a vertex.

### 3 Metric generators in cacti

Let  $G$  be a cactus graph with cycles  $C_1, \dots, C_c$ . We say that a vertex  $v \in V(G)$  *gravitates* to a cycle  $C_i$  in  $G$  if there is a path from  $v$  to a vertex from  $C_i$  which does not share any edge nor any internal vertex with any cycle of  $G$ . A *unicyclic region* of the cycle  $C_i$  from  $G$  is the subgraph  $G_i$  of  $G$  induced by all vertices that gravitate to  $C_i$  in  $G$ . The notion of unicyclic region of a cactus graph is illustrated by Figure 4.

Notice that each unicyclic region  $G_i$  is a unicyclic graph with its cycle being  $C_i$ . Also, considering the example from Figure 4, one can easily notice that two distinct unicyclic regions may not be vertex disjoint, as the path connecting vertex  $b_2$  and the cycle  $C_i$  belongs both to  $G_i$  and  $G_j$ . But, it does hold that all unicyclic regions cover the whole  $G$ . We say that a subgraph  $H$  of a graph  $G$  is an *isometric* subgraph if  $d_H(u, v) = d_G(u, v)$  for every pair of vertices  $u, v \in V(H)$ . The following observation is obvious.




Figure 4: A cactus graph  $G$  with three cycles in which the unicyclic region  $G_i$  of the cycle  $C_i$  is distinguished with  $b_1$  and  $b_2$  being the boundary vertices of  $G_i$ . The set  $S = \{s_1, s_2, s_3\}$  is a smallest biactive branch-resolving set in  $G$  for which a set of  $S$ -active vertices is marked on each cycle. For the set  $S$ , the regional set in  $G_i$  is  $S_i = \{s_2, b_1, b_2\}$ .

**Observation 6** *The unicyclic region  $G_i$  of a cycle  $C_i$  is an isometric subgraph of  $G$ .*

Finally, we say that a vertex  $v$  from a unicyclic region  $G_i$  is a *boundary* vertex if  $v \in V(C_j)$  for  $j \neq i$ . In the example from Figure 4, the boundary vertices of the region  $G_i$  are  $b_1$  and  $b_2$ .

Let  $S$  be a biactive branch-resolving set in  $G$  and let  $G_i$  be a unicyclic region in  $G$ . For the set  $S$  we define the *regional set*  $S_i$  as the set obtained from  $S \cap V(G_i)$  by introducing all boundary vertices from  $G_i$  to  $S$ . For example, in Figure 4 the set  $S_i = \{s_2, b_1, b_2\}$  is the regional set in the region of the cycle  $C_i$ .

**Lemma 7** *Let  $G$  be a cactus graph with  $c$  cycles  $C_1, \dots, C_c$  and let  $S \subseteq V(G)$ . If  $S$  is a vertex (resp. edge) metric generator in  $G$ , then the regional set  $S_i$  is a vertex (resp. edge) metric generator in the unicyclic region  $G_i$  for every  $i \in \{1, \dots, c\}$ .*

**Proof.** Suppose first that there is a cycle  $C_i$  in  $G$  such that the regional set  $S_i$  is not a vertex (resp. an edge) metric generator in the unicyclic region  $G_i$ . This implies that there exists a pair of vertices (resp. edges)  $x$  and  $x'$  in  $G_i$  which are not distinguished by  $S_i$ . We will show that  $x$  and  $x'$  are not distinguished by  $S$  in  $G$  either. Suppose the contrary, i.e. there is a vertex  $s \in S$  which distinguishes  $x$  and  $x'$  in  $G$ . If  $s \in V(G_i)$ , then  $s \in S_i$ . Since  $G_i$  is an isometric subgraph of  $G$ , then  $x$  and  $x'$  would be distinguished by  $s \in S_i$  in  $G_i$ , a contradiction. Assume therefore that  $s \notin V(G_i)$ . Notice that the shortest path from every vertex (resp. edge) in  $G_i$  to  $s$  leads through a same boundary vertex  $b$  in  $G_i$ . The definition of  $S_i$  implies  $b \in S_i$ , so  $x$  and  $x'$  are not distinguished by  $b$ . Therefore, we obtain

$$d(x, s) = d(x, b) + d(b, s) = d(x', b) + d(b, s) = d(x', s),$$



so  $x$  and  $x'$  are not distinguished by  $s$  in  $G$ , a contradiction.  $\blacksquare$

Notice that the condition from Lemma 7 is necessary for  $S$  to be a metric generator, but it is not sufficient as is illustrated by the graph shown in Figure 5 in which every regional set  $S_i$  is a vertex (resp. an edge) metric generator in the corresponding region  $G_i$ , but there still exists a pair of vertices (resp. edges) which is not distinguished by  $S$ , so  $S$  is not a vertex (resp. an edge) metric generator in  $G$ .




Figure 5: A cactus graph  $G$  with three cycles and a smallest biactive branch-resolving set  $S = \{s_1, s_2\}$  in  $G$ . Every regional set  $S_i$  is a vertex (resp. an edge) metric generator in the corresponding region  $G_i$ , but still a pair of vertices  $v_1$  and  $v_2$  is not distinguished by  $S$ . The pair of edges  $v_1v$  and  $v_2v$  is also not distinguished by  $S$  and the same holds for the pair of edges  $w_1w$  and  $w_2w$ .

Next, we will introduce notions which are necessary to state a condition which will be both necessary and sufficient for a biactive branch-resolving set  $S$  to be a vertex (resp. an edge) metric generator in a cactus graph  $G$ . An  $S$ -path of the cycle  $C_i$  is any subpath of  $C_i$  which contains all  $S$ -active vertices on  $C_i$  and is of minimum possible length. We denote an  $S$ -path of the cycle  $C_i$  by  $P_i$ . Notice that the end-vertices of an  $S$ -path are always  $S$ -active, otherwise it would not be shortest. For example, on the cycle  $C_2$  in Figure 5 there are two different paths connecting  $S$ -active vertices  $v$  and  $w$ , one is of the length 3 and the other of length 5, so the shorter one is an  $S$ -path. Also, an  $S$ -path  $P_i$  of a cycle  $C_i$  may not be unique, as there may exist several shortest subpaths of  $C_i$  containing all  $S$ -active vertices on  $C_i$ , but if the length of  $P_i$  satisfies  $|P_i| \leq \lceil g_i/2 \rceil - 1$  then  $P_i$  is certainly unique and its end-vertices are  $v_0$  and  $v_k$  in the canonical labelling of  $C_i$ .

**Definition 8** Let  $G$  be a cactus graph with cycles  $C_1, \dots, C_c$  and let  $S$  be a biactive branch-resolving set in  $G$ . We say that a vertex  $v \in V(C_i)$  is vertex-critical (resp. edge-critical) on  $C_i$  with respect to  $S$  if  $v$  is an end-vertex of  $P_i$  and  $|P_i| \leq \lfloor g_i/2 \rfloor - 1$  (resp.  $|P_i| \leq \lceil g_i/2 \rceil - 1$ ).

Notice that the notion of a vertex-critical and an edge-critical vertex differs only on odd cycles. We say that two distinct cycles  $C_i$  and  $C_j$  of a cactus graph  $G$  are vertex-critically incident (resp. edge-critically incident) with respect to  $S$  if  $C_i$  and  $C_j$  share a

vertex  $v$  which is vertex-critical (resp. edge-critical) with respect to  $S$  on both  $C_i$  and  $C_j$ . Notice that on odd cycles the required length of an  $S$ -path  $P_i$  for  $v$  to be vertex-critical differs from the one required for  $v$  to be edge-critical, while on even cycles the required length is the same.

To illustrate this notion, let us consider the cycle  $C_2$  in the graph from Figure 5. Vertices  $v$  and  $w$  are both vertex-critical and edge-critical on  $C_2$  with respect to  $S$  from the figure. Vertex  $v$  belongs also to  $C_1$  and it is also both vertex- and edge-critical on  $C_1$ . Therefore, cycles  $C_1$  and  $C_2$  are both vertex- and edge-critically incident, the consequence of which is that a pair of vertices  $v_1$  and  $v_2$  which are neighbors of  $v$  and a pair of edges  $v_1v$  and  $v_2v$  which are incident to  $v$  are not distinguished by  $S$ . On the other hand, vertex  $w$  belongs also to  $C_3$  on which it is edge-critical, but it is not vertex-critical since  $P_3$  is not short enough. So,  $C_2$  and  $C_3$  are edge-critically incident, but not vertex-critically incident. Consequently, a pair of edges  $w_1w$  and  $w_2w$  is not distinguished by  $S$ , but a pair of vertices  $w_1$  and  $w_2$  is distinguished by  $S$ . We will show in the following lemma that this holds in general.

**Lemma 9** *Let  $G$  be a cactus graph with  $c$  cycles  $C_1, \dots, C_c$  and let  $S$  be a biactive branch-resolving set in  $G$ . If  $S$  is a vertex (resp. an edge) metric generator in  $G$ , then there is no pair of cycles in  $G$  which are vertex-critically (resp. edge critically) incident with respect to  $S$ .*

**Proof.** Let  $S$  be a vertex (resp. an edge) metric generator in  $G$ . Suppose the contrary, i.e. there are two distinct cycles  $C_i$  and  $C_j$  in  $G$  which are vertex-critically (resp. edge-critically) incident with respect to  $S$ . This implies that  $C_i$  and  $C_j$  share a vertex  $v$  which is vertex-critical (resp. edge-critical) on both  $C_i$  and  $C_j$ . Let  $x$  and  $x'$  be a pair of vertices (resp. edges) which are neighbors (resp. incident) to  $v$  on cycles  $C_i$  and  $C_j$  respectively, but which are not contained on paths  $P_i$  and  $P_j$ . The length of paths  $P_i$  and  $P_j$  which is required by the definition of a vertex-critical (resp. edge-critical) vertex implies that a shortest path from both  $x$  and  $x'$  to all vertices from  $P_i$  and  $P_j$  leads through  $v$ . Since  $P_i$  and  $P_j$  contain all  $S$ -active vertices on  $C_i$  and  $C_j$ , this further implies that a shortest path from both  $x$  and  $x'$  to all vertices from  $S$  leads through  $v$ . Since  $d(x, v) = d(x', v)$ , it follows that  $x$  and  $x'$  are not distinguished by  $S$ , a contradiction. ■

Each of Lemmas 7 and 9 gives a necessary condition for a biactive branch-resolving set  $S$  to be a vertex (resp. an edge) metric generator in a cactus graph  $G$ . Let us now show that these two necessary conditions taken together form a sufficient condition for  $S$  to be a vertex (resp. an edge) metric generator.

**Lemma 10** *Let  $G$  be a cactus graph with  $c$  cycles  $C_1, \dots, C_c$  and let  $S$  be a biactive branch-resolving set in  $G$ . If a regional set  $S_i$  is a vertex (resp. an edge) metric generator in the unicyclic region  $G_i$  for every  $i = 1, \dots, c$  and there are no vertex-critically (resp. edge-critically) incident cycles in  $G$ , then  $S$  is a vertex (resp. an edge) metric generator in  $G$ .*

**Proof.** Let  $x$  and  $x'$  be a pair of vertices (resp. edges) from  $G$ . We want to show that  $S$  distinguishes  $x$  and  $x'$ . In order to do so, we distinguish the following two cases.

**Case 1:**  $x$  and  $x'$  belong to a same unicyclic region  $G_i$  of  $G$ . Since the regional set  $S_i$  is a vertex (resp. an edge) metric generator in  $G_i$ , there is a vertex  $s \in S_i$  which distinguishes  $x$  and  $x'$  in  $G_i$ . If  $s \in S$ , then the fact that  $G_i$  is an isometric subgraph of  $G$  implies that the pair  $x$  and  $x'$  is distinguished by the same  $s$  in  $G$  as well. Assume therefore that  $s \notin S$ , so the definition of the regional set  $S_i$  implies that  $s$  is a boundary vertex of  $G_i$ . Let  $s' \in S$  be a vertex in  $G$  such that the shortest path from  $s'$  to both  $x$  and  $x'$  leads through the boundary vertex  $s$ . Recall that such a vertex  $s'$  must exist since  $S$  is biactive. The fact that  $s$  distinguishes  $x$  and  $x'$  in  $G_i$ , implies  $d(x, s) \neq d(x', s)$ , which further implies

$$d(x, s') = d(x, s) + d(s, s') \neq d(x', s) + d(s, s') = d(x', s'),$$

so the pair  $x$  and  $x'$  is distinguished by  $S$  in  $G$ .

**Case 2:**  $x$  and  $x'$  do not belong to a same unicyclic region of  $G$ . Let us assume that  $x$  belongs to  $G_i$  and  $x'$  does not belong to  $G_i$ , and say it belongs to  $G_j$  for  $j \neq i$ . If  $x$  and  $x'$  are distinguished by a vertex  $s \in S \cap V(G_i)$ , then the claim is proven, so let us assume that  $x$  and  $x'$  are not distinguished by any  $s \in S \cap V(G_i)$ . Since  $x$  and  $x'$  do not belong to a same unicyclic region, there exists a boundary vertex  $b$  of the unicyclic region  $G_i$  such that the shortest path from  $x$  to  $x'$  leads through  $b$ . Let  $s_b$  be a vertex from  $S$  such that the shortest path from  $x$  to  $s_b$  also leads through  $b$ , which must exist since  $S$  is biactive. We want to prove that  $x$  and  $x'$  are distinguished by  $s_b$ . Let us suppose the contrary, i.e.  $d(x, s_b) = d(x', s_b)$ . Then we have the following

$$d(x, b) + d(b, s_b) = d(x, s_b) = d(x', s_b) \leq d(x', b) + d(b, s_b),$$

from which we obtain

$$d(x, b) \leq d(x', b). \tag{1}$$

Now, we distinguish the following two subcases.

**Subcase 2.a:**  $b$  does not belong to  $V(C_i)$ . Notice that by the definition of the unicyclic region, any acyclic structure hanging at  $b$  in  $G$  is not included in  $G_i$ , as is illustrated by  $b_2$  from Figure 4, which implies  $b$  is a leaf in  $G_i$ . Let  $b'$  be the only neighbor of  $b$  in  $G_i$ . The inequality (1) further implies  $d(x, b') < d(x', b')$  since  $x$  belongs to  $G_i$  and  $x'$  does not. Let  $v_0$  be the vertex from  $C_i$  closest to  $b$ , which implies  $v_0$  is  $S$ -active on  $C_i$ . Let  $v_k$  be an  $S$ -active vertex on  $C_i$  distinct from  $v_0$ , such a vertex  $v_k$  must exist on  $C_i$  because we assumed  $S$  is biactive. So, we have

$$d(x, v_k) \leq d(x, b') + d(b', v_k) < d(x', b') + d(b', v_k) = d(x', v_k).$$

Let  $s_k$  be a vertex from  $S$  which belongs to the connected component of  $G - E(C_i)$  containing  $v_k$ . Then we have

$$d(x, s_k) \leq d(x, v_k) + d(v_k, s_k) < d(x', v_k) + d(v_k, s_k) = d(x', s_k).$$

Therefore,  $S$  distinguishes  $x$  and  $x'$ , so  $S$  is a vertex (resp. an edge) metric generator in  $G$ .

**Subcase 2.b:**  $b$  belongs to  $V(C_i)$ . Since  $b$  is a boundary vertex of the unicyclic region  $G_i$ , this implies there is a cycle  $C_l$ , for  $l \neq i$ , such that  $b \in V(C_l)$ . Therefore, cycles  $C_i$  and

$C_l$  share the vertex  $b$ . Notice that any acyclic structure hanging at  $b$  belongs to both  $G_i$  and  $G_l$ , as is illustrated by  $b_1$  from Figure 4. If  $x'$  belongs to  $G_l$ , then neither  $x$  nor  $x'$  can belong to an acyclic structure hanging at  $b$ , as we assumed that  $x$  and  $x'$  do not belong to a same component. On the other hand, if  $x'$  does not belong to  $G_l$  and  $x$  belongs to an acyclic structure hanging at  $b$ , then we switch  $G_i$  by  $G_l$  and assume that  $x$  belongs to  $G_l$ . This way we assure that neither  $x$  nor  $x'$  belong to an acyclic structure hanging at  $b$ .

If  $d(x, b) < d(x', b)$ , let  $v_k$  be an  $S$ -active vertex on  $C_i$  distinct from  $b$ , which must exist as  $S$  is biactive. From (1) we obtain

$$d(x, v_k) \leq d(x, b) + d(b, v_k) < d(x', b) + d(b, v_k) = d(x', v_k),$$

so similarly as in previous subcase  $x$  and  $x'$  are distinguished by a vertex  $s_k \in S$  which belongs to the connected component of  $G - E(C_i)$  which contains  $v_k$ .

Therefore, assume that  $d(x, b) = d(x', b)$ . If a shortest path from  $x$  to all  $S$ -active vertices on  $C_i$  and a shortest path from  $x'$  to all  $S$ -active vertices on  $C_l$  leads through  $b$ , then  $x'$  belongs to  $C_l$ , i.e.  $j = l$ , and the pair of cycles  $C_i$  and  $C_l$  are vertex-critically (resp. edge-critically) incident, a contradiction. So, we may assume there is an  $S$ -active vertex  $v_k$ , say on  $C_i$ , such that a shortest path from  $x$  to  $v_k$  does not lead through  $b$ . Therefore,

$$d(x, v_k) < d(x, b) + d(b, v_k) = d(x', b) + d(b, v_k) = d(x', v_k).$$

But now, similarly as in previous cases we have that  $x$  and  $x'$  are distinguished by a vertex  $s_k \in S$  which is contained in the connected component of  $G - E(C_i)$  containing  $v_k$ . ■

Let us now relate these results with configurations  $\mathcal{A}$ ,  $\mathcal{B}$ ,  $\mathcal{C}$ ,  $\mathcal{D}$  and  $\mathcal{E}$ .

**Lemma 11** *Let  $G$  be a cactus graph and let  $S$  be a biactive branch-resolving set in  $G$ . A cycle  $C_i$  of the graph  $G$  contains configuration  $\mathcal{A}$  (or  $\mathcal{B}$  or  $\mathcal{C}$  or  $\mathcal{D}$  or  $\mathcal{E}$ ) with respect to  $S$  in  $G$  if and only if  $C_i$  contains the respective configuration with respect to  $S_i$  in  $G_i$ .*

**Proof.** Let  $G$  be a cactus graph with cycles  $C_1, \dots, C_c$  and let  $S$  be a biactive branch-resolving set in  $G$ . Since  $S$  is a biactive set, for every boundary vertex  $b$  in a unicyclic region  $G_i$ , there is a vertex  $s \in S$  such that the shortest path from  $s$  to  $C_i$  leads through  $b$ , as it is shown in Figure 4. This implies that the set of  $S$ -active vertices on  $C_i$  in  $G$  is the same as the set of  $S_i$ -active vertices on  $C_i$  in  $G_i$ . Since the presence of configurations  $\mathcal{A}$ ,  $\mathcal{B}$ ,  $\mathcal{C}$ ,  $\mathcal{D}$  and  $\mathcal{E}$  on a cycle  $C_i$ , with respect to a set  $S$ , by definition depends on the position of  $S$ -active vertices on  $C_i$ , the claim follows. ■

Notice that Lemmas 7, 9 and 10 give us a condition for  $S$  to be a vertex (resp. an edge) metric generator in a cactus graph, which is both necessary and sufficient. In the light of Lemma 11, we can further apply Theorem 5 to obtain the following result which unifies all our results.

**Theorem 12** *Let  $G$  be a cactus graph with  $c$  cycles  $C_1, \dots, C_c$  and let  $S$  be a biactive branch-resolving set in  $G$ . The set  $S$  is a vertex (resp. an edge) metric generator if and only if each cycle  $C_i$  does not contain any of the configurations  $\mathcal{A}$ ,  $\mathcal{B}$ , and  $\mathcal{C}$  (resp.  $\mathcal{A}$ ,  $\mathcal{D}$ , and  $\mathcal{E}$ ) and there are no vertex-critically (resp. edge-critically) incident cycles in  $G$  with respect to  $S$ .*

**Proof.** Let  $S$  be a vertex (resp. an edge) metric generator in  $G$ . Then Lemma 7 implies that  $S_i$  is a vertex (resp. edge) metric generator in the unicyclic region  $G_i$  and the Theorem 5 further implies that every cycle does not contain any of the configurations  $\mathcal{A}$ ,  $\mathcal{B}$ , and  $\mathcal{C}$  (resp.  $\mathcal{A}$ ,  $\mathcal{D}$ , and  $\mathcal{E}$ ) with respect to  $S$ . Also, Lemma 9 implies that there are no vertex-critically (resp. edge-critically) incident cycles in  $G$  with respect to  $S$ .

The other direction is the consequence of Lemma 10 and Theorem 5.  $\blacksquare$

## 4 Metric dimensions in cacti




Figure 6: A cactus graph with three cycles and two different smallest biactive branch-resolving sets  $S = \{s_1, s_2\}$  and  $S' = \{s'_1, s'_2\}$ . With respect to  $S$  the cycle  $C_2$  contains configuration  $\mathcal{A}$  and  $C_3$  configurations  $\mathcal{B}$ ,  $\mathcal{C}$  and  $\mathcal{D}$ . With respect to  $S'$  cycles  $C_2$  and  $C_3$  contain none of the five configurations. The cycle  $C_1$  has  $b(C_1) = 2$ , so the set of  $S$ -active vertices  $\{u, v\}$  on  $C_1$  is the same for all smallest biactive branch-resolving sets and  $C_1$  contains configurations  $\mathcal{B}$  and  $\mathcal{D}$  with respect to all of them, including  $S$  and  $S'$  shown in the figure. Therefore, in this graph cycle  $C_1$  is both  $\mathcal{ABC}$ - and  $\mathcal{ADE}$ -positive, and cycles  $C_2$  and  $C_3$  are both  $\mathcal{ABC}$ - and  $\mathcal{ADE}$ -negative.

Let  $G$  be a cactus graph and  $S$  a smallest biactive branch-resolving set in  $G$ . Then

$$|S| = L(G) + B(G),$$

where  $B(G) = \sum_{i=1}^c \max\{0, 2 - b(C_i)\}$ . If  $b(C_i) \geq 2$ , then the set of  $S$ -active vertices on  $C_i$  is the same for all smallest biactive branch-resolving sets  $S$ . The set of  $S$ -active vertices may differ only on cycles  $C_i$  with  $b(C_i) < 2$ . Therefore, such a cycle  $C_i$  may contain one of the configurations with respect to one smallest biactive branch-resolving set, but not with respect to another. This is illustrated by Figure 6.

**Definition 13** We say that a cycle  $C_i$  from a cactus graph  $G$  is  $\mathcal{ABC}$ -negative (resp.  $\mathcal{ADE}$ -negative), if there exists a smallest biactive branch-resolving set  $S$  in  $G$  such that  $C_i$  does not contain any of the configurations  $\mathcal{A}$ ,  $\mathcal{B}$ , and  $\mathcal{C}$  (resp.  $\mathcal{A}$ ,  $\mathcal{D}$ , and  $\mathcal{E}$ ) with respect to  $S$ . Otherwise, we say that  $C_i$  is  $\mathcal{ABC}$ -positive (resp.  $\mathcal{ADE}$ -positive). The number of  $\mathcal{ABC}$ -positive (resp.  $\mathcal{ADE}$ -positive) cycles in  $G$  is denoted by  $c_{\mathcal{ABC}}(G)$  (resp.  $c_{\mathcal{ADE}}(G)$ ).

For two distinct smallest biactive branch-resolving sets  $S$ , the set of  $S$ -active vertices may differ only on cycles with  $b(C_i) \leq 1$ . Let  $C_i$  and  $C_j$  be two such cycles in  $G$  and notice that the choice of the vertices included in  $S$  from the region of  $C_i$  is independent of the choice from  $C_j$ . Therefore, there exists at least one smallest biactive branch-resolving set  $S$  such that every  $\mathcal{ABC}$ -negative (resp.  $\mathcal{ADE}$ -negative) avoids the three configurations with respect to  $S$ . Notice that there may exist more than one such set  $S$ , and in that case they all have the same size, so among them we may choose the one with the smallest number of vertex-critical (resp. edge-critical) incidencies. Therefore, we say that a smallest biactive branch-resolving set  $S$  is *nice* if every  $\mathcal{ABC}$ -negative (resp.  $\mathcal{ADE}$ -negative) cycle  $C_i$  does not contain the three configurations with respect to  $S$  and the number of vertex-critically (resp. edge-critically) incident pairs of cycles with respect to  $S$  is the smallest possible. The niceness of a smallest biactive branch-resolving set is illustrated by Figure 7.




Figure 7: A cactus graph with three cycles and two different smallest biactive branch-resolving sets  $S = \{s_1, s_2\}$  and  $S' = \{s'_1, s'_2\}$ . With respect to both  $S$  and  $S'$  all three cycles do not contain any of the five configurations. The difference is that with respect to  $S$  the cycle  $C_2$  is both vertex- and edge-critically incident with both  $C_1$  and  $C_3$ , and with respect to  $S'$  the cycle  $C_2$  is both vertex- and edge-critically incident with  $C_3$ , but only edge-critically incident with  $C_1$ . Therefore, the set  $S$  is not nice. Since the critical incidences of  $S'$  cannot be further avoided the set  $S'$  is nice.

A set  $S \subseteq V(G)$  is a *vertex cover* if it contains a least one end-vertex of every edge in  $G$ . The cardinality of a smallest vertex cover in  $G$  is the *vertex cover number* denoted by  $\tau(G)$ . Now, let  $G$  be a cactus graph and let  $S$  be a nice smallest biactive branch-resolving set in  $G$ . We define the *vertex-incident graph*  $G_{vi}$  (resp. *edge-incident graph*  $G_{ei}$ ) as a graph containing a vertex for every cycle in  $G$ , where two vertices are adjacent if the corresponding cycles in  $G$  are  $\mathcal{ABC}$ -negative and vertex-critically incident (resp.  $\mathcal{ADE}$ -negative and edge-critically incident) with respect to  $S$ . For example, if we consider the cactus graph  $G$  from Figure 8, then  $V(G_{vi}) = V(G_{ei}) = \{C_i : i = 1, \dots, 7\}$ , where  $E(G_{vi}) = \{C_3C_4, C_4C_5\}$  and  $E(G_{ei}) = \{C_2C_3, C_3C_4, C_4C_5\}$ .

We are now in a position to establish the following theorem which gives us the value of the vertex and the edge metric dimensions in a cactus graph.

**Theorem 14** *Let  $G$  be a cactus graph. Then*

$$\dim(G) = L(G) + B(G) + c_{\mathcal{ABC}}(G) + \tau(G_{vi}),$$

and

$$\text{edim}(G) = L(G) + B(G) + c_{\mathcal{ADE}}(G) + \tau(G_{ei}).$$

**Proof.** If there is a cycle in  $G$  with  $b(C) = 0$ , then  $G$  is a unicyclic graph. For unicyclic graphs with  $b(C) = 0$  we have  $B(G) = 2$  and if the three configurations  $\mathcal{A}, \mathcal{B}, \mathcal{C}$  (resp.  $\mathcal{A}, \mathcal{D}, \mathcal{E}$ ) cannot be avoided by choosing two vertices into  $S$ , then  $c_{\mathcal{ABC}}(G) = 1$  (resp.  $c_{\mathcal{ADE}}(G) = 1$ ) and also the third vertex must be introduced to  $S$ , so the claim holds. In all other situations  $B(G)$  equals the number of cycles in  $G$  with  $b(C_i) = 1$ .

Let  $S$  be a smallest vertex (resp. edge) metric generator in  $G$ . Due to Lemma 1 the set  $S$  must be branch-resolving. Let  $S_1 \subseteq S$  be a smallest branch-resolving set contained in  $S$ , so  $|S_1| = L(G)$ . Since according to Lemma 1 the set  $S$  must also be biactive, let  $S_2 \subseteq S \setminus S_1$  be a smallest set such that  $S_1 \cup S_2$  is biactive. Obviously,  $S_1 \cap S_2 = \emptyset$  and  $|S_2| = B(G)$ .

Since  $S_1 \cup S_2$  is a smallest biactive branch-resolving set in  $G$ , it follows that every  $\mathcal{ABC}$ -positive (resp.  $\mathcal{ADE}$ -positive) cycle in  $G$  contains at least one of the three configurations with respect to  $S_1 \cup S_2$ , so according to Theorem 12 the set  $S_1 \cup S_2$  is not a vertex (resp. an edge) metric generator in  $G$ . Therefore, each  $\mathcal{ABC}$ -positive (resp.  $\mathcal{ADE}$ -positive) cycle  $C_i$  must contain a vertex  $s_i \in S \setminus (S_1 \cup S_2)$ , where we may assume that  $s_i$  is chosen so that it forms a geodesic triple on  $C_i$  with vertices from  $S_1 \cup S_2$ , so according to Observation 4 the cycle  $C_i$  will not contain any of the configurations with respect to  $S$ . Denote by  $S_3$  the set of vertices  $s_i$  from every  $\mathcal{ABC}$ -positive (resp.  $\mathcal{ADE}$ -positive) cycle in  $G$ . Obviously,  $S_3 \subseteq S$ ,  $S_1 \cap S_2 \cap S_3 = \emptyset$  and  $|S_3| = c_{\mathcal{ABC}}(G)$  (resp.  $|S_3| = c_{\mathcal{ADE}}(G)$ ).

Notice that  $S_1 \cup S_2 \cup S_3$  is a biactive branch-resolving set in  $G$  such that every cycle  $C_i$  in  $G$  does not contain any of the configurations  $\mathcal{A}, \mathcal{B}, \mathcal{C}$  (resp.  $\mathcal{A}, \mathcal{D}, \mathcal{E}$ ) with respect to it. Notice that  $S_1 \cup S_2 \cup S_3$  still may not be a vertex (resp. an edge) metric generator, as there may exist vertex-critically (resp. edge-critically) incident cycles in  $G$  with respect to  $S_1 \cup S_2 \cup S_3$ . Since  $S$  is a smallest vertex (resp. edge) metric generator, we may assume that  $S_2$  is chosen so that a smallest biactive branch-resolving set  $S_1 \cup S_2$  is nice. Therefore, the graph  $G_{vi}$  (resp.  $G_{ei}$ ) contains an edge for every pair of cycles in  $G$  which are  $\mathcal{ABC}$ -negative and vertex-critically incident (resp.  $\mathcal{ADE}$ -negative and edge-critically incident) with respect to  $S_1 \cup S_2$ . Let us denote  $S_4 = S \setminus (S_1 \cup S_2 \cup S_3)$ . For each edge  $xy$  in  $G_{vi}$  (resp.  $G_{ei}$ ) the set  $S_4$  must contain a vertex from  $C_x$  or  $C_y$ , chosen so that it forms a geodesic triple of  $S$ -active vertices on  $C_x$  or  $C_y$  with other vertices from  $S$ . Therefore,  $S_4$  must contain at least  $\tau(G_{vi})$  (resp.  $\tau(G_{ei})$ ) vertices in order for  $S$  to be a vertex (resp. an edge) metric generator. Since  $S$  is a smallest vertex (resp. edge) metric generator, it must hold  $|S_4| = \tau(G_{vi})$  (resp.  $|S_4| = \tau(G_{ei})$ ).

We have established that  $S = S_1 \cup S_2 \cup S_3 \cup S_4$ , where  $S_1 \cap S_2 \cap S_3 \cap S_4 = \emptyset$ , so  $|S| = |S_1| + |S_2| + |S_3| + |S_4|$ . Since we also established  $|S_1| = L(G)$ ,  $|S_2| = B(G)$ ,  $|S_3| = c_{\mathcal{ABC}}(G)$  (resp.  $|S_3| = c_{\mathcal{ADE}}(G)$ ) and  $|S_4| = \tau(G_{vi})$  (resp.  $|S_4| = \tau(G_{ei})$ ), the proof is finished. ■

The formulas for the calculation of metric dimensions from the above theorem are illustrated by the following examples.

**Example 2** Let us consider the cactus graph  $G$  from Figure 6. The set  $S' = \{s'_1, s'_2\}$  is an optimal smallest biactive branch-resolving set in  $G$ . But, since  $C_1$  is both  $\mathcal{ABC}$ - and  $\mathcal{ADE}$ -positive, the set  $S'$  is neither a vertex nor an edge metric generator. Let  $s'_3$  be any

vertex from  $C_1$  which forms a geodesic triple with two  $S'$ -active vertices on  $C_1$ . Then the set  $S = \{s'_1, s'_2, s'_3\}$  is a smallest vertex (resp. edge) metric generator, so we obtain

$$\dim(G) = L(G) + B(G) + c_{ABC}(G) + \tau(G_{vi}) = 0 + 2 + 1 + 0 = 3.$$

and

$$\text{edim}(G) = L(G) + B(G) + c_{ADE}(G) + \tau(G_{ei}) = 0 + 2 + 1 + 0 = 3.$$




Figure 8: A cactus graph  $G$  from Example 3.

Let us now give an example of determining the vertex and the edge metric dimensions on a cactus graph which is a bit bigger.

**Example 3** Let  $G$  be the cactus graph from Figure 8. The following table gives the choice and the number of vertices for every expression in the formulas for metric dimensions from Theorem 14

	vertices	value
$L(G)$	$s_1, s_2, s_3, s_4$	4
$B(G)$	$s_5$	1
$c_{ABC}(G)$	$s_6$	1
$\tau(G_{vi})$	$s_7$	1
$c_{ADE}(G)$	$s_8, s_9$	2
$\tau(G_{ei})$	$s_{10}, s_{11}$	2



Therefore, the set  $S = \{s_1, s_2, s_3, s_4, s_5, s_6, s_7\}$  is a smallest vertex metric generator, so we obtain

$$\dim(G) = L(G) + B(G) + c_{ABC}(G) + \tau(G_{vi}) = 4 + 1 + 1 + 1 = 7.$$

On the other hand, the set  $S = \{s_1, s_2, s_3, s_4, s_5, s_8, s_9, s_{10}, s_{11}\}$  is a smallest edge metric generator, so we have

$$\text{edim}(G) = L(G) + B(G) + c_{ADE}(G) + \tau(G_{ei}) = 4 + 1 + 2 + 2 = 9.$$

Notice that  $c_{ABC}(G) \leq c$ . Also, if  $\tau(G_{vi}) \geq 1$  then  $c_{ABC}(G) + \tau(G_{vi}) < c$ . The similar holds for  $c_{ADE}(G)$  and  $\tau(G_{ei})$ . From this and Theorem 14 we immediately obtain the following result.

**Corollary 15** *Let  $G$  be a cactus graph with  $c$  cycles. Then  $\dim(G) \leq L(G) + B(G) + c$  and  $\text{edim}(G) \leq L(G) + B(G) + c$ .*

Further, notice that in a cactus graph with at least two cycles every cycle has at least one branch-active vertex. Therefore, in such a cactus graph  $G$ , we have  $B(G) = \sum_{i=1}^c \max\{0, 2 - b(C_i)\} \leq c$  with equality holding only if  $b(C_i) = 1$  for every cycle  $C_i$  in  $G$ . Since  $c_{ABC}(G) + \tau(G_{vi}) = c$  if and only if  $c_{ABC}(G) = c$  and  $\tau(G_{vi}) = 0$ , and similarly holds for the edge version of metric dimension, Theorem 14 immediately implies the following simple upper bound on the vertex and edge metric dimensions of a cactus graph  $G$ .

**Corollary 16** *Let  $G$  be a cactus graph with  $c \geq 2$  cycles. Then*

$$\dim(G) \leq L(G) + 2c$$

*with equality holding if and only if every cycle in  $G$  is  $ABC$ -positive and contains precisely one branch-active vertex.*

**Corollary 17** *Let  $G$  be a cactus graph with  $c \geq 2$  cycles. Then*

$$\text{edim}(G) \leq L(G) + 2c$$

*with equality holding if and only if every cycle in  $G$  is  $ADE$ -positive and contains precisely one branch-active vertex.*

Notice that the upper bound from the above corollary may not hold for  $c = 1$ , i.e. for unicyclic graphs, as for the cycle  $C$  of unicyclic graph it may hold that  $b(C) = 0$ . As for the tightness of these bounds, we have the following proposition.

**Proposition 18** *For every pair of integers  $b \geq 0$  and  $c \geq 2$ , there is a cactus graph  $G$  with  $c$  cycles such that  $L(G) = b$  and  $\dim(G) = \text{edim}(G) = L(G) + 2c$ .*

**Proof.** For a given pair of integers  $b \geq 0$  and  $c \geq 2$ , we construct a cactus graph  $G$  in a following way. Let  $G_0$  be a graph on  $b + 2$  vertices, with one vertex  $u$  of degree  $b + 1$  and all other vertices of degree 1, i.e.  $G_0$  is a star graph. Let  $H$  be a graph obtained from the 6-cycle by introducing a leaf to it and let  $G_1, \dots, G_c$  be  $c$  vertex disjoint copies of  $H$ . Denote by  $v_i$  the only vertex of degree 3 in  $G_i$ . Let  $G$  be a graph obtained from  $G_0, G_1, \dots, G_c$  by connecting them with an edge  $uv_i$  for  $i = 1, \dots, c$ . Obviously,  $G$  is a cactus graph with  $c$  cycles and  $L(G) = b$ . On each of the cycles in  $G$  the vertex  $v_i$  is the only branch-active vertex. If  $S \subseteq V(G)$  is a smallest branch-resolving set in  $G$  such that there is a cycle  $C_i$  in  $G$  with only two  $S$ -active vertices, then because of the leaf pending on  $v_i$  the cycle  $C_i$  contains either configuration  $\mathcal{A}$  if the pair of  $S$ -active vertices on  $C_i$  is an antipodal pair or both configuration  $\mathcal{B}$  and  $\mathcal{D}$ . Either way,  $S$  is not a vertex nor an edge metric generator.

On the other hand, the set  $S$  consisting of  $b$  leaves hanging at  $u$  in  $G_0$  and a pair of vertices from each 6-cycle which form a geodesic triple with  $v_i$  on the cycle is both a vertex and an edge metric generator in  $G$ . Since  $|S| = b + 2c = L(G) + 2c$ , the claims hold. ■

## 5 An application to zero forcing number

The results from previous section enable us to solve for cactus graphs a conjecture posed in literature [4] which involves the vertex metric dimension, the zero forcing number and the cyclomatic number  $c(G) = |E(G)| - |V(G)| + 1$  (which is sometimes called the cycle rank number and denoted by  $r(G)$ ) of a graph  $G$ . Notice that in a cactus graph  $G$  the cyclomatic number  $c(G)$  equals the number of cycles in  $G$ . Let us first define the zero forcing number of a graph.

Assuming that every vertex of a graph  $G$  is assigned one of two colors, say black and white, the set of vertices which are initially black is denoted by  $S$ . If there is a black vertex with only one white neighbor, then the *color-change rule* converts the only white neighbor also to black. This is one iteration of color-change rule, it can be applied iteratively. A *zero forcing set* is any set  $S \subseteq V(G)$  such that all vertices of  $G$  are colored black after applying the color-change rule finitely many times. The cardinality of the smallest zero forcing set in a graph  $G$  is called the *zero forcing number* of  $G$  and it is denoted by  $Z(G)$ . In [4] it was proven that for a unicyclic graph  $G$  it holds that  $\dim(G) \leq Z(G) + 1$ , and it was further conjectured the following.

**Conjecture 19** *For any graph  $G$  it holds that  $\dim(G) \leq Z(G) + c(G)$ .*

Moreover, they proved for cacti with even cycles the bound  $\dim(G) \leq Z(G) + c(G)$ . We will use our results to prove that for cacti the tighter bound from the above conjecture holds.

**Proposition 20** *Let  $G$  be a cactus graph. Then  $\dim(G) \leq Z(G) + c(G)$  and  $\text{edim}(G) \leq Z(G) + c(G)$ .*

**Proof.** Due to Corollary 15 it is sufficient to prove that  $L(G) + B(G) \leq Z(G)$ . Let  $S \subseteq V(G)$  be a zero forcing set in  $G$ . Let us first show that  $S$  must be a branch-resolving set. Assume the contrary, i.e. that  $S$  is not a branch-resolving set and let  $v \in V(G)$  be a vertex of degree  $\geq 3$  with at least two  $S$ -free threads hanging at  $v$ . But then  $v$  has at least two white neighbors, one on each of the  $S$ -free threads hanging at it, which cannot be colored black by  $S$ , so  $S$  is not a zero forcing set, a contradiction.

Let  $S_1 \subseteq S$  be a smallest branch-resolving set contained in  $S$  and let  $S_2 = S \setminus S_1$ . Obviously,  $|S_1| = L(G)$  and  $S_1 \cap S_2 = \emptyset$ . We now wish to prove that  $|S_2| \geq B(G)$ . If  $G$  is a tree, then the claim obviously holds, so let us assume that  $G$  contains at least one cycle. Let  $C_i$  be a cycle in  $G$  such that  $b(C_i) \leq 1$ . If  $b(C_i) = 0$ , then  $G$  is a unicyclic graph and  $C_i$  the only cycle in  $G$ . Since  $b(C_i) = 0$ , we have  $S_1 = \emptyset$ , so  $S_2 = S$ . Since a zero forcing set in unicyclic graph must contain at least two vertices, we obtain  $|S_2| = |S| \geq 2 = B(G)$  and the claim is proven.

Assume now that for every cycle  $C_i$  with  $b(C_i) \leq 1$  it holds that  $b(C_i) = 1$ . Let  $v$  be the branch-active vertex on such a cycle  $C_i$  and notice that  $S_1$  can turn only  $v$  black on  $C_i$ . Therefore, in order for  $S$  to be a zero forcing set it follows that  $S_2$  must contain a vertex from every such cycle, i.e.  $|S_2| \geq B(G)$ . Therefore,  $|S| = |S_1| + |S_2| \geq L(G) + B(G)$ . ■

The above proposition, besides proving for cacti the cycle rank conjecture which was posed for  $\dim(G)$ , also gives a similar result for  $\text{edim}(G)$ . So, this motivates us to pose for  $\text{edim}(G)$  the counterpart conjecture of Conjecture 19.

## 6 Concluding remarks

In [18] it was established that for a unicyclic graph  $G$  both vertex and edge metric dimensions are equal to  $L(G) + \max\{2 - b(G), 0\}$  or  $L(G) + \max\{2 - b(G), 0\} + 1$ . In [19] a characterization under which both of the dimensions take one of the two possible values was further established. In this paper we extend the result to cactus graphs where a similar characterization must hold for every cycle in a graph, and also the additional characterization for the connection of two cycles must be introduced. This result enabled us to prove the cycle rank conjecture for cactus graphs.

Moreover, the results of this paper enabled us to establish a simple upper bound on the value of the vertex and the edge metric dimension of a cactus graph  $G$  with  $c$  cycles

$$\dim(G) \leq L(G) + 2c \quad \text{and} \quad \text{edim}(G) \leq L(G) + 2c.$$

Since the number of cycles can be generalized to all graphs as the cyclomatic number  $c(G) = |E(G)| - |V(G)| + 1$ , we conjecture that the analogous bounds hold in general.

**Conjecture 21** *Let  $G$  be a connected graph. Then,  $\dim(G) \leq L(G) + 2c(G)$ .*

**Conjecture 22** *Let  $G$  be a connected graph. Then,  $\text{edim}(G) \leq L(G) + 2c(G)$ .*

In [21] it was shown that the inequality  $\text{mdim}(G) < 2c(G)$  holds for 3-connected graphs. Since  $\dim(G) \leq \text{mdim}(G)$  and  $\text{edim}(G) \leq \text{mdim}(G)$ , the previous two conjectures obviously hold for 3-connected graphs.

Also, motivated by the bound on edge metric dimension of cacti involving zero forcing number, we state the following conjecture for general graphs, as a counterpart of Conjecture 19.

**Conjecture 23** *Let  $G$  be a connected graph. Then,  $\text{edim}(G) \leq Z(G) + c(G)$ .*

**Acknowledgments.** Both authors acknowledge partial support of the Slovenian research agency ARRS program P1-0383 and ARRS projects J1-1692 and J1-8130. The first author also the support of Project KK.01.1.1.02.0027, a project co-financed by the Croatian Government and the European Union through the European Regional Development Fund - the Competitiveness and Cohesion Operational Programme.

## References

- [1] P. S. Buczkowski, G. Chartrand, C. Poisson, P. Zhang, On  $k$ -dimensional graphs and their bases, *Period. Math. Hungar.* **46**(1) (2003) 9–15.
- [2] J. Caceres, C. Hernando, M. Mora, I. M. Pelayo, M. L. Puertas, C. Seara, D. R. Wood, On the metric dimension of Cartesian products of graphs, *SIAM J. Discrete Math.* **21**(2) (2007) 423–441.
- [3] G. Chartrand, L. Eroh, M. A. Johnson, O. R. Oellermann, Resolvability in graphs and the metric dimension of a graph, *Discrete Appl. Math.* **105** (2000) 99–113.
- [4] L. Eroh, C.X. Kang, E. Yi, A comparison between the metric dimension and zero forcing number of trees and unicyclic graphs, *Acta Math. Sin. (Engl. Ser.)* **33**(6) (2017) 731–747.
- [5] M. Fehr, S. Gosselin, O. R. Oellermann, The metric dimension of Cayley digraphs, *Discrete Math.* **306** (2006) 31–41.
- [6] J. Geneson, Metric dimension and pattern avoidance in graphs, *Discrete Appl. Math.* **284** (2020) 1–7.
- [7] F. Harary, R. A. Melter, On the metric dimension of a graph, *Ars Combin.* **2** (1976) 191–195.
- [8] Y. Huang, B. Hou, W. Liu, L. Wu, S. Rainwater, S. Gao, On approximation algorithm for the edge metric dimension problem, *Theoret. Comput. Sci.* (2020), doi:<https://doi.org/10.1016/j.tcs.2020.05.005>.
- [9] A. Kelenc, Distance-Based Invariants and Measures in Graphs, PhD thesis, University of Maribor, Faculty of Natural Sciences and Mathematics, 2020.
- [10] A. Kelenc, D. Kuziak, A. Taranenko, I. G. Yero, Mixed metric dimension of graphs, *Appl. Math. Comput.* **314**(1) (2017) 42–438.

- [11] A. Kelenc, N. Tratnik, I. G. Yero, Uniquely identifying the edges of a graph: the edge metric dimension, *Discrete Appl. Math.* **251** (2018) 204–220.
- [12] S. Khuller, B. Raghavachari, A. Rosenfeld, Landmarks in graphs, *Discrete Appl. Math.* **70** (1996) 217–229.
- [13] D. J. Klein, E. Yi, A comparison on metric dimension of graphs, line graphs, and line graphs of the subdivision graphs, *Eur. J. Pure Appl. Math.* **5(3)** (2012) 302–316.
- [14] M. Knor, S. Majstorović, A. T. M. Toshi, R. Škrekovski, I. G. Yero, Graphs with the edge metric dimension smaller than the metric dimension, *Appl. Math. Comput.* **401** (2021) 126076.
- [15] R. A. Melter, I. Tomescu, Metric bases in digital geometry, *Comput. Vis. Graph. Image Process.* **25** (1984) 113–121.
- [16] I. Peterin, I. G. Yero, Edge metric dimension of some graph operations, *Bull. Malays. Math. Sci. Soc.* **43** (2020) 2465–2477.
- [17] S. W. Saputro, R. Simanjuntak, S. Uttunggadewa, H. Assiyatun, E. T. Baskoro, A. N. M. Salman, M. Bača, The metric dimension of the lexicographic product of graphs, *Discrete Math.* **313** (2013) 1045–1051.
- [18] J. Sedlar, R. Škrekovski, Bounds on metric dimensions of graphs with edge disjoint cycles, *Appl. Math. Comput.* **396** (2021) 125908.
- [19] J. Sedlar, R. Škrekovski, Vertex and edge metric dimensions of unicyclic graphs, arXiv:2104.00577 [math.CO].
- [20] J. Sedlar, R. Škrekovski, Extremal mixed metric dimension with respect to the cyclomatic number, *Appl. Math. Comput.* **404** (2021) 126238.
- [21] J. Sedlar, R. Škrekovski, Mixed metric dimension of graphs with edge disjoint cycles, *Discr. Appl. Math.* **300** (2021) 1–8.
- [22] P. J. Slater, Leaves of trees, *Congr. Numer.* **14** (1975) 549–559.
- [23] I. G. Yero, D. Kuziak, J. A. Rodriguez-Velazquez, On the metric dimension of corona product graphs, *Comput. Math. Appl.* **61** (2011) 2793–2798.
- [24] Y. Zhang, S. Gao, On the edge metric dimension of convex polytopes and its related graphs, *J. Comb. Optim.* **39(2)** (2020) 334–350.
- [25] E. Zhu, A. Taranenko, Z. Shao, J. Xu, On graphs with the maximum edge metric dimension, *Discrete Appl. Math.* **257** (2019) 317–324.
- [26] N. Zubrilina, On the edge dimension of a graph, *Discrete Math.* **341(7)** (2018) 2083–2088.