

MULTI-PARAMETER FLAG LEIBNIZ RULES OF ARBITRARY COMPLEXITY IN MIXED-NORM SPACES

CRISTINA BENEÀ AND YUJIA ZHAI

ABSTRACT. We prove multi-parameter Leibniz rules corresponding to flag paraproducts of arbitrary complexity in mixed-norm spaces, including endpoint estimates. The proof relies on multi-linear harmonic analysis techniques and a quantitative treatment of the commutators introduced by Bourgain and Li. The argument is robust and applicable to a generic class of multipliers, including (symmetric) Mikhlin multipliers of positive order and asymmetric variants of partial differential operators and Mikhlin multipliers of positive order.

1. INTRODUCTION

1.1. Motivation and Main results. In this work, we study multi-parameter flag Leibniz rules, which include the particular bi-parameter example

$$(1.1) \quad \begin{aligned} & \|D_{(1)}^{\beta_1} D_{(2)}^{\beta_2} (D_{(1)}^{\alpha_1} D_{(2)}^{\alpha_2} (f_1 f_2) f_3 D_{(1)}^{\gamma_1} D_{(2)}^{\gamma_2} (f_4 f_5))\|_{L^{\vec{p}}} \\ & \lesssim \|D_{(1)}^{\alpha_1+\beta_1} D_{(2)}^{\alpha_2+\beta_2} f_1\|_{L^{\vec{p}_1}} \|f_2\|_{L^{\vec{p}_2}} \|f_3\|_{L^{\vec{p}_3}} \|D_{(1)}^{\gamma_1} D_{(2)}^{\gamma_2} f_4\|_{L^{\vec{p}_4}} \|f_5\|_{L^{\vec{p}_5}} + \|f_1\|_{L^{\vec{p}_1}} \|D_{(1)}^{\alpha_1+\beta_1} D_{(2)}^{\alpha_2+\beta_2} f_2\|_{L^{\vec{p}_2}} \|f_3\|_{L^{\vec{p}_3}} \|D_{(1)}^{\gamma_1} D_{(2)}^{\gamma_2} f_4\|_{L^{\vec{p}_4}} \|f_5\|_{L^{\vec{p}_5}} + \\ & \|D_{(1)}^{\alpha_1+\beta_1} f_1\|_{L^{\vec{p}_1}} \|D_{(2)}^{\alpha_2+\beta_2} f_2\|_{L^{\vec{p}_2}} \|D_{(2)}^{\beta_2} f_3\|_{L^{\vec{p}_3}} \|D_{(1)}^{\gamma_1} f_4\|_{L^{\vec{p}_4}} \|D_{(2)}^{\gamma_2} f_5\|_{L^{\vec{p}_5}} + \|D_{(1)}^{\alpha_1} f_1\|_{L^{\vec{p}_1}} \|D_{(2)}^{\alpha_2+\beta_2} f_2\|_{L^{\vec{p}_2}} \|f_3\|_{L^{\vec{p}_3}} \|D_{(1)}^{\gamma_1+\beta_1} f_4\|_{L^{\vec{p}_4}} \|D_{(2)}^{\gamma_2} f_5\|_{L^{\vec{p}_5}} \\ & + \dots \quad 140 \text{ other similar terms.} \end{aligned}$$

Above, $\beta_1, \beta_2, \alpha_1, \alpha_2, \gamma_1, \gamma_2 \geq 0$ and the spaces $L^{\vec{p}_i}(\mathbb{R}^{d_1} \times \mathbb{R}^{d_2})$ represent mixed-norm Lebesgue spaces $L_x^{p_i^1}(L_y^{p_i^2})$ (see Section 2, definition 2.1) with $1 \leq p_i^1, p_i^2 \leq \infty$ for all $1 \leq i \leq 5$. Additionally, we require that

$$\sum_{i=1}^5 \frac{1}{p_i^1} = \frac{1}{r^1}, \quad \sum_{i=1}^5 \frac{1}{p_i^2} = \frac{1}{r^2},$$

and

$$\frac{1}{r^2} < \frac{d_2+\beta_2}{d_2}, \quad \frac{1}{p_1^1} + \frac{1}{p_2^1} < \frac{d_2+\alpha_2}{d_2}, \quad \frac{1}{p_4^1} + \frac{1}{p_5^1} < \frac{d_2+\gamma_2}{d_2},$$

$$\frac{1}{r^1} < \min\left(\frac{d_1+\beta_1}{d_1}, \frac{d_2+\beta_2}{d_2}\right), \quad \frac{1}{p_1^1} + \frac{1}{p_2^1} < \min\left(\frac{d_1+\alpha_1}{d_1}, \frac{d_2+\alpha_2}{d_2}\right), \quad \frac{1}{p_4^1} + \frac{1}{p_5^1} < \min\left(\frac{d_1+\gamma_1}{d_1}, \frac{d_2+\gamma_2}{d_2}\right).$$

The Leibniz rule (1.1) confirms the fact that fractional partial derivatives¹ acting in various ways on products of functions are properly distributed among the functions, provided these are elements of some mixed-norm $L^{\vec{p}}$ spaces, with $\vec{p} = (p^1, \dots, p^N)$, $1 \leq p^1, \dots, p^N \leq \infty$.

The difficulty – and thus the interest – of flag Leibniz rules such as (1.1) resides in the fact that straightaway composition arguments are insufficient when the input functions are too close to L^1 , in spite of them having the form of and distributing the derivatives as compositions of simpler Leibniz rules. This is a feature shared with the *flag paraproducts* introduced by Muscalu in [14], which resemble compositions of Coifman-Meyer multipliers. In fact, Coifman-Meyer multipliers are usually invoked in the study of fractional Leibniz rules, and in particular the boundedness of the one-parameter flag paraproduct from [14] implies the one-parameter flag Leibniz rule.²

When it comes to fractional partial derivatives acting independently on various variables – as in (1.1) – a new layer of difficulty is added since the boundedness of multi-parameter flag paraproducts remains, to our knowledge, a difficult open problem. However, combining multilinear harmonic analysis techniques with the method introduced

¹For $\alpha_1, \alpha_2 \geq 0$, we consider the partial differential operators $D_{(1)}^{\alpha_1}, D_{(2)}^{\alpha_2}$, initially defined on the space $\mathcal{S}(\mathbb{R}^{d_1+d_2})$ of Schwartz functions – via the Fourier transform – by formulas

$$(1.2) \quad D_{(1)}^{\alpha_1} f := \mathcal{F}^{-1}(|\xi|^{\alpha_1} \widehat{f}(\xi, \eta)), \quad D_{(2)}^{\alpha_2} f := \mathcal{F}^{-1}(|\eta|^{\alpha_2} \widehat{f}(\xi, \eta)).$$

In contrast, the homogeneous differential operator D^{β} is defined on the Schwartz space $\mathcal{S}(\mathbb{R}^d)$ by

$$(1.3) \quad D^{\beta} f := \mathcal{F}^{-1}(|\xi|^{\beta} \widehat{f}(\xi)).$$

²Modulo some endpoints.

by Bourgain and Li [4] for proving Leibniz rules for input data in L^∞ (another situation in which Coifman-Meyer multipliers cannot be invoked), we are able to prove multi-parameter flag Leibniz rules of arbitrary complexity.

Differences and similarities between the reduction of Leibniz rules to the boundedness of Coifman-Meyer multipliers and the Bourgain-Li method will be discussed in Section 2.3.

Leibniz rules of various types have been extensively investigated and widely used in nonlinear PDEs. The simplest Leibniz rule, acting on functions defined on \mathbb{R}^d , takes the form

$$(1.4) \quad \|D^\beta(f_1 f_2)\|_{L^r} \lesssim \|D^\beta f_1\|_{L^{p_1}} \|f_2\|_{L^{p_2}} + \|f_1\|_{L^{p_1}} \|D^\beta f_2\|_{L^{p_2}}$$

where

$$(1.5) \quad 1 \leq p_1, p_2 \leq \infty, \quad \frac{d}{d+\alpha} < r \leq \infty, \quad \frac{1}{p_1} + \frac{1}{p_2} = \frac{1}{r}.$$

As mentioned above, the Coifman-Meyer theorem [5] implies the Leibniz rule (1.4), but only in the range

$$1 < p_1, p_2 \leq \infty, \quad \frac{d}{d+\alpha} < r < \infty, \quad \frac{1}{p_1} + \frac{1}{p_2} = \frac{1}{r}.$$

Coifman-Meyer operators are associated in frequency to Mikhlin symbols: they are n -linear operators T_m described³ by

$$T_m(f_1, \dots, f_n) := \int_{\mathbb{R}^{nd}} m(\xi_1, \dots, \xi_n) \hat{f}_1(\xi_1) \cdots \hat{f}_n(\xi_n) e^{2\pi i x(\xi_1 + \dots + \xi_n)} d\xi_1 \dots d\xi_n,$$

where m is a Mikhlin symbol (of order 0) satisfying

$$(1.6) \quad |\partial_{\xi_1}^{\gamma_1} \dots \partial_{\xi_n}^{\gamma_n} m(\xi_1, \dots, \xi_n)| \lesssim (|\xi_1| + \dots + |\xi_n|)^{-|\gamma_1| - \dots - |\gamma_n|}$$

for sufficiently many⁴ multi-indices $\gamma_1, \dots, \gamma_n$. The class of Mikhlin symbols on \mathbb{R}^{nd} will be denoted $\mathcal{M}(\mathbb{R}^{nd})$.

Since multilinear Coifman-Meyer operators (which are particular cases of multilinear Calderon-Zygmund operators) do not satisfy $L^\infty \times L^\infty \mapsto L^\infty$, or (strong type) $L^1 \times L^p \mapsto L^{\frac{p}{p+1}}$ bounds for $1 \leq p \leq \infty$, a different approach is required for dealing with these endpoint estimates. This was introduced in [4], where Bourgain and Li proved that

$$\|D^\beta(f_1 f_2)\|_{L^\infty} \lesssim \|D^\beta f_1\|_{L^\infty} \|f_2\|_{L^\infty} + \|f_1\|_{L^\infty} \|D^\beta f_2\|_{L^\infty},$$

a result conjectured in [7]. In fact, the authors proved in [4] a Kato-Ponce commutator estimate involving Besov norms⁵

$$(1.7) \quad \|D^\beta(f_1 f_2) - (D^\beta f_1) f_2 - f_1 (D^\beta f_2)\|_{L^\infty} \lesssim \|f_1\|_{L^\infty} \|f_2\|_{\dot{B}_{\infty,\infty}^\beta} + \|f_1\|_{\dot{B}_{\infty,\infty}^\beta} \|f_2\|_{L^\infty}.$$

The appearance of Besov norms should already suggest that a scale-by-scale analysis will be performed, and that the estimates obtained in this way will be summed according to their magnitude; this is in sharp contrast with the approach for Coifman-Meyer multipliers, in which the orthogonality between different scales plays a crucial role.

The case when (at least) one of p_1 or p_2 is equal to 1 was proved by Oh and Wu in [18], by applying the methods introduced in [4]. It represented a first instance of a (strong-type) Leibniz rule for input functions in L^1 .

Bi-parameter Leibniz rules, such as

$$(1.8) \quad \|D_{(1)}^{\beta_1} D_{(2)}^{\beta_2}(f_1 f_2)\|_{L^r(\mathbb{R}^{d_1+d_2})} \lesssim \|D_{(1)}^{\beta_1} D_{(2)}^{\beta_2} f_1\|_{L^{p_1}(\mathbb{R}^{d_1+d_2})} \|f_2\|_{L^{p_2}(\mathbb{R}^{d_1+d_2})} + \|f_1\|_{L^{p_1}(\mathbb{R}^{d_1+d_2})} \|D_{(1)}^{\beta_1} D_{(2)}^{\beta_2} f_2\|_{L^{p_2}(\mathbb{R}^{d_1+d_2})} + \|D_{(1)}^{\beta_1} f_1\|_{L^{p_1}(\mathbb{R}^{d_1+d_2})} \|D_{(2)}^{\beta_2} f_2\|_{L^{p_2}(\mathbb{R}^{d_1+d_2})} + \|D_{(2)}^{\beta_2} f_1\|_{L^{p_1}(\mathbb{R}^{d_1+d_2})} \|D_{(1)}^{\beta_1} f_2\|_{L^{p_2}(\mathbb{R}^{d_1+d_2})},$$

at least for

$$(1.9) \quad 1 < p_1, p_2 \leq \infty, \quad \max\left(\frac{d_1}{d_1 + \beta_1}, \frac{d_2}{d_2 + \beta_2}\right) < r < \infty, \quad \frac{1}{p_1} + \frac{1}{p_2} = \frac{1}{r},$$

are a consequence of bi-parameter paraproducts' boundedness within the same range (1.9), as proved in [15]. The endpoints $p_1 = p_2 = r = \infty$ and strong estimates in the case $p_1 = 1$ or $p_2 = 1$ for (1.8) are contained in the work of Oh and Wu [18].

³Given a frequency symbol m , we denote by T_m the associated n -linear operator.

⁴In certain situations, finding minimal regularity conditions for Mikhlin symbols becomes important. This will not be the case for our applications concerning Leibniz rules for homogeneous (or inhomogeneous, as we will see later) partial fractional differential operators, as the symbols involved will either be smooth or will be a suitable superposition of smooth symbols.

⁵The Besov norms $\|\cdot\|_{\dot{B}_{p,\infty}^\beta}$ associated to the real parameter β and to the Lebesgue exponent $1 \leq p \leq \infty$, will be explicitly defined in Section 2.1 – see (2.6).

Mixed-norm estimates for bi-parameter Leibniz rules were obtained more recently, as a consequence of mixed-norm estimates for bi-parameter paraproducts [1], [2]⁶; more exactly, it was proved that the inequality

$$(1.10) \quad \begin{aligned} \|D_{(1)}^{\beta_1} D_{(2)}^{\beta_2} (f_1 f_2)\|_{L_{x^1}^1(L_{y^2}^{r^2})} &\lesssim \|D_{(1)}^{\beta_1} D_{(2)}^{\beta_2} f_1\|_{L_{x^1}^{p_1^1}(L_{y^1}^{p_2^2})} \|f_2\|_{L_{x^2}^{p_1^1}(L_{y^2}^{p_2^2})} + \|f_1\|_{L_{x^1}^{p_1^1}(L_{y^1}^{p_2^2})} \|D_{(1)}^{\beta_1} D_{(2)}^{\beta_2} f_2\|_{L_{x^2}^{p_1^1}(L_{y^2}^{p_2^2})} \\ &+ \|D_{(1)}^{\beta_1} f_1\|_{L_{x^1}^{p_1^1}(L_{y^1}^{p_2^2})} \|D_{(2)}^{\beta_2} f_2\|_{L_{x^2}^{p_1^1}(L_{y^2}^{p_2^2})} + \|D_{(2)}^{\beta_2} f_1\|_{L_{x^1}^{p_1^1}(L_{y^1}^{p_2^2})} \|D_{(1)}^{\beta_1} f_2\|_{L_{x^2}^{p_1^1}(L_{y^2}^{p_2^2})}, \end{aligned}$$

holds for Lebesgue exponents satisfying

$$1 < p_1^i, p_2^i \leq \infty, \quad \frac{1}{p_1^i} + \frac{1}{p_2^i} = \frac{1}{r^i}, \quad \text{for all } 1 \leq i \leq 2, \quad \frac{d_2}{d_2 + \beta_2} < r^2 < \infty, \quad \max\left(\frac{d_1}{d_1 + \beta_1}, \frac{d_2}{d_2 + \beta_2}\right) < r^1 < \infty.$$

The same result, including the endpoints $r^i = \infty$ (which forces $p_1^i = p_2^i = \infty$) for some $1 \leq i \leq 2$, or $p_j^i = 1$ for some $1 \leq i, j \leq 2$, were proved by Oh and Wu [18] using the Bourgain-Li method and thus avoiding mixed-norm estimates for Coifman-Meyer multipliers. Other partial results were obtained in [6].

Multi-parameter multilinear operators are especially interesting; unlike their linear analogues, they take as input several functions, and yield as output only one function, so that linear techniques (freezing a variable, using vector-valued estimates) are not easily applicable. Mixed-norm estimates for multi-parameter multilinear operators present an additional difficulty, and in general they require sharper estimates for the concerned operator (localization, weighted estimates, etc; see [1], [2]).

The results above in (1.10) can be extended to the N -parameter case; except for a few endpoints,⁷ this result is implicit in [1], [2]. The full result, including L^1 and L^∞ endpoints is implicit in [18].

We would like to comment that although (1.4), (1.8) and (1.10) above describe the bi-linear case, the n -linear case for $n > 2$ remains valid and it can be proved by the same methods.

The one-parameter flag Leibniz rule can be perceived as the Leibniz rule for compositions of fractional differential operators. The simplest example is

$$(1.11) \quad \begin{aligned} \|D^\beta (D^\alpha (f_1 f_2) f_3)\|_{L^r} &\lesssim \|D^{\alpha+\beta} f_1\|_{L^{p_1}} \|f_2\|_{L^{p_2}} \|f_3\|_{L^{p_3}} + \|f_1\|_{L^{p_1}} \|D^{\alpha+\beta} f_2\|_{L^{p_2}} \|f_3\|_{L^{p_3}} \\ &+ \|D^\alpha f_1\|_{L^{p_1}} \|f_2\|_{L^{p_2}} \|D^\beta f_3\|_{L^{p_3}} + \|f_1\|_{L^{p_1}} \|D^\alpha f_2\|_{L^{p_2}} \|D^\beta f_3\|_{L^{p_3}}, \end{aligned}$$

where

$$(1.12) \quad 1 \leq p_1, p_2, p_3 \leq \infty, \quad \frac{1}{p_1} + \frac{1}{p_2} + \frac{1}{p_3} = \frac{1}{r}, \quad 0 \leq \frac{1}{r} < \frac{d+\beta}{d}, \quad 0 \leq \frac{1}{p_1} + \frac{1}{p_2} < \frac{d+\alpha}{d}.$$

The Leibniz rule (1.11) (except for endpoints $(p_1, p_2) = (\infty, \infty)$ or strong-type estimates when $p_i = 1$ for some $1 \leq i \leq 3$) is a consequence of the boundedness of the one-parameter flag paraproduct [14]. The endpoint case $(p_1, p_2) = (\infty, \infty)$ can be derived by iteratively applying the endpoint estimate (1.4) due to Bourgain and Li [4].

It is worth pointing out that interpreting the Leibniz rule (1.11) as a composition of two classical Leibniz rules (as described in (1.4)) and iteratively invoking (1.4) only yields a limited range of exponents, namely the case $1 \leq p_1, p_2, p_3 \leq \infty, 0 \leq \frac{1}{p_1} + \frac{1}{p_2} \leq 1$ and $\max(\frac{1}{2}, \frac{d}{d+\beta}) < r \leq \infty$. In order to achieve boundedness in the nontrivial range $1 < \frac{1}{p_1} + \frac{1}{p_2} < \min(2, \frac{d+\alpha}{d})$ and $\max(\frac{1}{3}, \frac{d}{d+\beta}) < r \leq \infty$, one can decompose $D^\beta (D^\alpha (f_1 f_2) f_3)$ into a sum of flag paraproducts whose boundedness was proved by Muscalu [14] and later extended by Miyachi and Tomita [13] to Hardy spaces input data.

The flag paraproduct should be thought of as compositions of Coifman-Meyer multipliers: in frequency, the associated symbol is a product of singular Mihlin symbols

$$(1.13) \quad m(\xi) := \prod_{S \in \{1, \dots, n\}} m_S(\xi_S)$$

where $\xi := (\xi_i)_{i=1}^n \in \mathbb{R}^{dn}$, $\xi_S := (\xi_i)_{i \in S} \in \mathbb{R}^{d \cdot \text{card}(S)}$ and $m_S \in \mathcal{M}(\mathbb{R}^{d \cdot \text{card}(S)})$. Hence the singularity set associated to m consists of unions of subspaces of various dimensions, which can be further organized into a union of ordered subspaces – or flags. Flag paraproducts do not satisfy $L^\infty \times \dots \times L^\infty \rightarrow L^\infty$, or strong type estimates $L^1 \times L^{p_2} \times \dots \times L^{p_n} \rightarrow L^r$; nevertheless, we will see that the flag Leibniz rule remains true even in these particular situations.

Although the boundedness of the generic multi-parameter flag paraproduct is still an open problem, as mentioned previously, a particular case of bi-parameter flag paraproducts was proved independently in [17] and [9].

⁶Although stated for functions defined on $\mathbb{R} \times \mathbb{R}$, the Leibniz rules in [1], [2] remain valid in higher dimensions.

⁷Certain complications appear when one tries to prove mixed-norm estimates for N -parameter paraproducts, when some of the input functions are in mixed-norm Lebesgue spaces involving L^∞ .

That leads to some specific example of bi-parameter Leibniz rules in the full range of boundedness, modulo certain endpoints. However, the generic multi-parameter versions of (1.11) known before were those that could be obtained as a result of compositions of (1.8) or (1.10) – which leaves out a significant range of Lebesgue exponents. We will show that the Bourgain-Li method can be adapted to proving multi-parameter flag Leibniz rules of arbitrary complexity.

Especially in the context of flag Leibniz rules (and of flag paraproducts, as one would expect), it becomes convenient to use *rooted tree* representations. The n -linear Leibniz rule for

$$(1.14) \quad D^\beta(f_1 \dots f_n),$$

corresponds to the simplest tree

$$(1.15) \quad \begin{array}{c} D^\beta \\ \bullet \\ \swarrow \quad \downarrow \quad \searrow \quad \swarrow \\ \bullet \quad \bullet \quad \dots \quad \bullet \\ f_1 \quad f_2 \quad \dots \quad f_n \end{array}$$

where the root of the tree is the vertex associated to the differential operator D^β and the leaves of the tree are the vertices associated to the n functions that the differential operator acts on.

The multilinear expression $D^\beta(D^\alpha(f_1 \cdot f_2)f_3)$ on the left hand side of (1.11) can be represented by the following rooted tree:

$$(1.16) \quad \begin{array}{c} D^\beta \\ \bullet \\ \swarrow \quad \searrow \\ D^\alpha \quad \bullet \\ \swarrow \quad \searrow \quad \downarrow \\ \bullet \quad \bullet \quad f_3 \\ f_1 \quad f_2 \end{array}.$$

In general, a rooted tree \mathcal{G} consists of a collection of vertices which are organized according to their depth; the root – denoted $\mathbf{r}_\mathcal{G}$ – has depth zero, the direct descendants of the root have depth 1, and so on. The vertices which don't have any descendants are called *leaves* – and $\mathcal{L}_\mathcal{G}$ denotes the collection of leaves in the rooted tree \mathcal{G} ; all the other vertices make up \mathcal{V} , the collection of vertices that have at least one descendant.⁸

To each $l \in \mathcal{L}_\mathcal{G}$ we associate a function acting on \mathbb{R}^d , and to each vertex $v \in \mathcal{V}$ we associate a fractional differential operator D^{β^v} for some $\beta^v \geq 0$. With an abuse of notation, we identify the collection $\mathcal{L}_\mathcal{G}$ of leaves with the collection of functions $\{f_1, \dots, f_n\}$, and similarly, the collection of vertices \mathcal{V} is identified with the collection of fractional differential operators $\{D^{\beta^v} : v \in \mathcal{V}\}$.

For each $v \in \mathcal{V}$, we define the set

$$(1.17) \quad \mathcal{L}(v) := \{i \in \{1, \dots, n\} : f_i \text{ is a descendant of } v\}.$$

As a consequence, $\mathcal{L}(\text{root}) = \{1, \dots, n\}$.

It is also easy to verify that for any $v_1, v_2 \in \mathcal{V}$, only one of the following situations can happen:

$$(1.18) \quad (1) \ \mathcal{L}(v_1) \cap \mathcal{L}(v_2) = \emptyset; \quad (2) \ \mathcal{L}(v_1) \subseteq \mathcal{L}(v_2); \quad (3) \ \mathcal{L}(v_2) \subseteq \mathcal{L}(v_1).$$

Last but not least, the *complexity* of the tree \mathcal{G} is defined as the maximal depth among the leaves vertices; or equivalently, as the maximum length of upward paths from a leaf to the root. For example, (1.15) is a tree of complexity 1 while (1.16) is of complexity 2.

We can adapt the tree representation to the multi-parameter setting by substituting the homogenous differential operators (1.3) with partial differential operators (1.2). The 5-linear expression on the left hand side of (1.1) indeed corresponds to a tree of complexity 2 represented by Figure 1(ii) bellow; next to it, Figure 1(i) depicts its one-parameter equivalent.

Building upon the above structures, one can obtain a rooted tree representation for N -parameter flag Leibniz rules as well; in such a situation, the leaves $\mathcal{L}_\mathcal{G}$ correspond to functions $\{f_1, \dots, f_n\}$ defined on $\mathbb{R}^{d_1} \times \dots \times \mathbb{R}^{d_N}$, and the vertices in \mathcal{V} to generic fractional partial differential operators $D_{(1)}^{\beta_1^v} \dots D_{(N)}^{\beta_N^v}$, where each $\beta_j^v \geq 0$ indicates the partial derivatives associated to v in the j -th parameter. As before, we identify a vertex v with its corresponding differential operator, and the set $\mathcal{L}_\mathcal{G}$ with the set of functions $\{f_1, \dots, f_n\}$.

⁸It is more natural to request that every vertex which is not a leaf has at least two descendants.




FIGURE 1. One and bi-parameter flag Leibniz rules

In this paper, we establish multi-parameter flag Leibniz rules of arbitrary complexity for input data in $L^{\vec{p}}$ spaces, with $1 \leq \vec{p} \leq \infty$.⁹ More precisely, we prove an N -parameter, n -linear Leibniz rule associated to a rooted tree \mathcal{G} with n leaves. The possible distribution of derivatives $D_{(1)}^{\beta_1^v} \dots D_{(N)}^{\beta_N^v}$ among the n functions f_1, \dots, f_n is described by the tensor map $\delta_1 \otimes \dots \otimes \delta_N : \mathcal{V}^N \rightarrow \{1, \dots, n\}^N$; for $1 \leq j \leq N$, δ_j represents a map

$$(1.19) \quad \delta_j : \mathcal{V} \rightarrow \{1, \dots, n\}$$

satisfying the following two conditions:

- (i) for any $v \in \mathcal{V}$, $\delta_j(v) \in \mathcal{L}(v)$ (equivalently, $D_{(j)}^{\beta_j^v}$ derivatives are attributed to $f_{\delta_j(v)}$, one of the leaves descending from the vertex v).
- (ii) if $\mathfrak{S}(v)$ denotes the set of non-leaf, direct descendants of the vertex v (for some $v \in \mathcal{V}$), then

$$\delta_j(v) = \delta_j(w), \quad \text{for some } w \in \mathfrak{S}(v).$$

This latter condition ensures that the distribution of derivatives agrees with the composition law. As a consequence of (ii), if $\delta(v) = l$ for some $v \in \mathcal{V}$, then for any non-leaf vertex w along the path from v to the leaf f_l , we have $\delta(w) = l$. Due to observation (1.18), conditions (i) and (ii) can be simultaneously satisfied.

We denote by $\mathcal{D}(\mathcal{V})$ (abbreviated as \mathcal{D}) the collection of maps $\delta_j : \mathcal{V} \rightarrow \{1, \dots, n\}$ satisfying conditions (i) and (ii) above for $1 \leq j \leq N$. We will abbreviate $\underbrace{\mathcal{D} \times \dots \times \mathcal{D}}_{N \text{ copies}}$ as \mathcal{D}^N .

We notice that such maps are well-defined since the leading partial derivatives only hit one function they act on at a time, but δ_j is not necessarily injective nor surjective – in the example (1.1), $D_{(1)}^{\beta_1}$ and $D_{(1)}^{\alpha_1}$ can hit f_1 simultaneously (thus not injective) and it is also possible that no derivative hits f_1 at all (thus not surjective). With abuse of notation, we will denote for any $1 \leq l \leq n$,

$$(1.20) \quad \delta_j^{-1}(l) := \sum_{v \in \mathcal{V} : \delta_j(v) = l} \beta_j^v$$

and if $l \notin \text{range}(\delta_j)$, then $\delta_j^{-1}(l) = 0$; hence $\delta_j^{-1}(l)$ keeps track of the number of partial derivatives attributed to f_l by the map δ_j .

For any $v \in \mathcal{V}$, p_v denotes the Lebesgue exponent defined by


$$(1.21) \quad \frac{1}{p_v} = \sum_{i \in \mathcal{L}(v)} \frac{1}{p_i}.$$

In the case of N -parameters rooted trees, we consider N -tuples $\vec{p}_v = (p_v^1, \dots, p_v^N)$ and each p_v^j , for $1 \leq j \leq N$, is defined by

$$(1.22) \quad \frac{1}{p_v^j} = \sum_{i \in \mathcal{L}(v)} \frac{1}{p_i^j}.$$

Now we are ready to state our main result:

⁹This inequality is to be understood componentwise.

FIGURE 2. A rooted tree of arbitrary complexity associated to an N -parameter flag Leibniz rule.

Theorem 1.1. Let \mathcal{G} be a rooted tree of root $\tau_{\mathcal{G}}$, and to every $v \in \mathcal{V}$ we associate the N -parameter fractional differential operator $D_{(1)}^{\beta_1^v} \dots D_{(N)}^{\beta_N^v}$, with $\beta_1^v, \dots, \beta_N^v \geq 0$. Let \mathcal{D}^N be the collection of maps $\delta_1 \otimes \dots \otimes \delta_N : \mathcal{V}^N \rightarrow \{1, \dots, n\}^N$ satisfying conditions (i) and (ii), which describe the admissible distributions of derivatives. If $T_{\mathcal{G}}$ denotes the n -linear operator indicated by the rooted tree \mathcal{G} , then for any functions $f_1, \dots, f_n \in \mathcal{S}(\mathbb{R}^{d_1} \times \dots \times \mathbb{R}^{d_N})$,

$$(1.23) \quad \|T_{\mathcal{G}}(f_1, \dots, f_n)\|_{\vec{r}} \lesssim \sum_{\delta_1 \otimes \dots \otimes \delta_N \in \mathcal{D}^N} \prod_{l=1}^n \|D_{(1)}^{\delta_1^{-1}(l)} \dots D_{(N)}^{\delta_N^{-1}(l)} f_l\|_{\vec{p}_l},$$

provided that $1 \leq p_l^j \leq \infty$ for all $1 \leq j \leq N$, $1 \leq l \leq n$,

$$(1.24) \quad \frac{1}{r^j} = \sum_{l=1}^n \frac{1}{p_l^j}, \quad \text{for all } 1 \leq j \leq N,$$

$$(1.25) \quad \frac{d_N}{d_N + \beta_N^{\tau_{\mathcal{G}}}} < r^N, \quad \max\left(\frac{d_N}{d_N + \beta_N^{\tau_{\mathcal{G}}}}, \frac{d_{N-1}}{d_{N-1} + \beta_{N-1}^{\tau_{\mathcal{G}}}}\right) < r^{N-1}, \dots, \max\left(\frac{d_N}{d_N + \beta_N^{\tau_{\mathcal{G}}}}, \dots, \frac{d_1}{d_1 + \beta_1^{\tau_{\mathcal{G}}}}\right) < r^1$$

and in general, for any $v \in \mathcal{V}$,

$$(1.26) \quad \frac{d_N}{d_N + \beta_N^v} < p_v^N, \quad \max\left(\frac{d_N}{d_N + \beta_N^v}, \frac{d_{N-1}}{d_{N-1} + \beta_{N-1}^v}\right) < p_v^{N-1}, \dots, \max\left(\frac{d_N}{d_N + \beta_N^v}, \dots, \frac{d_1}{d_1 + \beta_1^v}\right) < p_v^1.$$

Whenever $\beta_j^v \in 2\mathbb{N}$, the corresponding conditions on p_v^j in (1.26) above can be removed.

The only previously known case of the above result corresponds to $N = 1$: the one-parameter flag Leibniz rules associated to trees of arbitrary complexity are a consequence¹⁰ of the boundedness of flag paraproducts of arbitrary complexity from [14].

We remark that the mixed norms

$$\|\cdot\|_{L^{\vec{p}_l}(\mathbb{R}^{d_1} \times \dots \times \mathbb{R}^{d_N})} = \|\dots\|_{L^{p_l^N}(\mathbb{R}^{d_N})} \dots \|_{L^{p_l^1}(\mathbb{R}^{d_1})}$$

in Theorem 1.1 can be further replaced by

$$\|\dots\|_{L^{\vec{p}_l^N}(\mathbb{R}^{d_N})} \dots \|_{L^{\vec{p}_l^1}(\mathbb{R}^{d_1})};$$

that is, each $\|\cdot\|_{L^{p_l^j}(\mathbb{R}^{d_j})}$ norm can be replaced by the mixed norm $\|\cdot\|_{L^{\vec{p}_l^j}(\mathbb{R}^{d_j})}$, as long as $\vec{p}_l^j = (p_{l,1}^j, \dots, p_{l,d_j}^j) \in [1, \infty]^{d_j}$. In this situation we require (component-wise) conditions analogous to (1.24), (1.25) and (1.26):

$$\frac{1}{\vec{r}^j} = \sum_{l=1}^n \frac{1}{\vec{p}_l^j}, \quad \text{for all } 1 \leq j \leq N,$$

$$\frac{d_N}{d_N + \beta_N^{\tau_{\mathcal{G}}}} < \min(r_1^N, \dots, r_{d_N}^N), \dots, \max\left(\frac{d_N}{d_N + \beta_N^{\tau_{\mathcal{G}}}}, \dots, \frac{d_1}{d_1 + \beta_1^{\tau_{\mathcal{G}}}}\right) < \min(r_1^1, \dots, r_{d_1}^1)$$

and for any $v \in \mathcal{V}$,

$$\frac{d_N}{d_N + \beta_N^v} < \min(p_{v,1}^N, \dots, p_{v,d_N}^N), \dots, \max\left(\frac{d_N}{d_N + \beta_N^v}, \dots, \frac{d_1}{d_1 + \beta_1^v}\right) < \min(p_{v,1}^1, \dots, p_{v,d_1}^1).$$

This remains true for all our results, namely Theorem 1.1-1.6.

¹⁰Modulo endpoints.

Moreover, for a rooted tree \mathcal{G} , we can associate to each vertex $v \in \mathcal{V}$ the inhomogeneous differential operator $J_{(1)}^{\beta_1^v} \dots J_{(N)}^{\beta_N^v}$ ¹¹ instead of the homogeneous differential operator $D_{(1)}^{\beta_1^v} \dots D_{(N)}^{\beta_N^v}$. The multilinear operator – denoted by $T_{\mathcal{G}}^J$ – satisfies the same mixed-norm estimates as $T_{\mathcal{G}}$ described in Theorem 1.1 and can be treated in the same fashion; the only notable difference appears at the level of cone/paraproduct decompositions, since the inhomogeneous partial differential operators do not pick out small frequency scales. Details on how to adjust the decompositions can be found in Grafakos-Oh [8] or Oh-Wu [18].

Now we return to our initial examples – the explicit Leibniz rule (1.1) – in order to clarify the notation in our main theorem. In the example (1.1), $n = 5$ and $N = 2$, so that the set of leaves $\mathcal{L}_{\mathcal{G}}$ consists of

$$\mathcal{L}_{\mathcal{G}} = \{f_1, f_2, f_3, f_4, f_5\},$$

and the collection \mathcal{V} of non-leaf vertices of

$$\mathcal{V} = \{\vec{\beta} = (\beta_1, \beta_2), \vec{\alpha} = (\alpha_1, \alpha_2), \vec{\gamma} = (\gamma_1, \gamma_2)\}.$$

Then

$$\mathcal{L}(\vec{\beta}) = \{1, 2, 3, 4, 5\}, \quad \mathcal{L}(\vec{\alpha}) = \{1, 2\}, \quad \mathcal{L}(\vec{\gamma}) = \{4, 5\}$$

and for any $j = 1, 2$, $\delta_j : \{\beta_j, \alpha_j, \gamma_j\} \rightarrow \{1, 2, 3, 4, 5\}$ must satisfy the condition that

$$\delta_j(\alpha_j) \in \{1, 2\}, \quad \delta_j(\gamma_j) \in \{4, 5\},$$

and

$$\delta_j(\beta_j) \in \{1, 2, 3, 4, 5\} \quad \text{with} \quad \delta_j(\beta_j) = \delta_j(\alpha_j) \quad \text{or} \quad \delta_j(\beta_j) = \delta_j(\gamma_j) \quad \text{or} \quad \delta_j(\beta_j) = 3.$$

As a result, there are in total 144 choices of $\delta_1 \otimes \delta_2$! The first term in the right-hand side of (1.1) indeed corresponds to the particular choice of the maps

$$\delta_1(\beta_1) = \delta_1(\alpha_1) = 1, \delta_1(\gamma_1) = 4; \quad \delta_2(\beta_2) = \delta_2(\alpha_2) = 1, \delta_2(\gamma_2) = 4,$$

whereas the fourth term corresponds to

$$\delta_1(\beta_1) = \delta_1(\gamma_1) = 4, \delta_1(\alpha_1) = 1; \quad \delta_2(\beta_2) = \delta_2(\alpha_2) = 2, \delta_2(\gamma_2) = 5.$$

We can equivalently represent the flag Leibniz rules in frequency – since the fractional (partial) differential operators themselves are defined in frequency (see (1.2), (1.3)). Starting from the observation that

$$(1.27) \quad D^\beta(f_1 f_2 \dots f_n) = \mathcal{F}^{-1}(|\xi_1 + \dots + \xi_n|^\beta \hat{f}_1(\xi_1) \dots \hat{f}_n(\xi_n)),$$

we realize that a correspondence can be established between the trees

$$(1.28) \quad \begin{array}{c} D^\beta \\ \bullet \\ \swarrow \quad \downarrow \quad \searrow \\ \bullet \quad \bullet \quad \dots \quad \bullet \\ f_1 \quad f_2 \quad \dots \quad f_n \end{array} \quad \leftrightarrow \quad \begin{array}{c} |\xi_1 + \dots + \xi_n|^\beta \\ \bullet \\ \swarrow \quad \downarrow \quad \searrow \\ \bullet \quad \bullet \quad \dots \quad \bullet \\ \hat{f}_1(\xi_1) \quad \hat{f}_2(\xi_2) \quad \dots \quad \hat{f}_n(\xi_n) \end{array}.$$

One should notice that in the frequency representation of the flag Leibniz rules, the set of leaves $\hat{\mathcal{L}}_{\mathcal{G}}$ consists of $\{\hat{f}_1(\xi_1), \dots, \hat{f}_n(\xi_n)\}$. The vertices $v \in \mathcal{V}$ should be identified, in the one-parameter case, with the symbols

$$(1.29) \quad \left| \sum_{i \in \mathcal{L}(v)} \xi_i \right|^{\beta^v};$$

in the N -parameters case, the non-leaf vertices appearing in the flag Leibniz rules should be identified with

$$(1.30) \quad \prod_{j=1}^N \left| \sum_{i \in \mathcal{L}(v)} \xi_i \right|^{\beta_j^v}.$$

The representation in frequency of the 5-linear flag appearing in (1.1) and of its one-parameter analogue are represented in Figures 3 (ii) and 3(i) below.

¹¹For $1 \leq j \leq N$, we denote by $J_{(j)}^{\beta_j}$, with $\beta_j \geq 0$, the inhomogeneous partial differential operator defined on the space of Schwartz functions $\mathcal{S}(\mathbb{R}^{d_1} \times \dots \times \mathbb{R}^{d_N})$ by

$$J_{(j)}^{\beta_j} f := \mathcal{F}^{-1}((1 + |\xi^j|^2)^{\frac{\beta_j}{2}} \hat{f}(\xi^1, \dots, \xi^N)).$$




FIGURE 3. Frequency representation of one and bi-parameter flag Leibniz rules

Interestingly, the same methods imply the boundedness of multi-parameter flag multipliers that do not correspond directly to multi-parameter flag Leibniz rules, since the structures of the trees are different for different parameters. One such example is the following 5-linear, bi-parameter expression

$$(1.31) \quad T_{\mathcal{G}_1 \otimes \mathcal{G}_2}(f_1, f_2, f_3, f_4, f_5)(x, y) \\ := \int_{\mathbb{R}^{10}} |\xi_1 + \xi_2 + \xi_3 + \xi_4 + \xi_5|^{\beta_1} |\xi_1 + \xi_2|^{\alpha_1} |\xi_3 + \xi_4 + \xi_5|^{\gamma_1} |\xi_3 + \xi_4|^{\zeta_1} |\eta_1 + \eta_2 + \eta_3 + \eta_4 + \eta_5|^{\beta_2} |\eta_1 + \eta_3|^{\alpha_2} |\eta_2 + \eta_4|^{\gamma_2} \\ \prod_{l=1}^5 \hat{f}_l(\xi_l, \eta_l) e^{2\pi i(x, y) \cdot (\xi_1 + \xi_2 + \xi_3 + \xi_4 + \xi_5, \eta_1 + \eta_2 + \eta_3 + \eta_4 + \eta_5)} d\xi d\eta.$$

More generally, we can consider n -linear, N -parameter operators that can be represented as $T_{\mathcal{G}_1 \otimes \dots \otimes \mathcal{G}_N}$, where each \mathcal{G}_j , for $1 \leq j \leq N$, indicates the frequency tree in the j -th parameter.

For any $1 \leq j \leq N$, we denote by \mathcal{V}_j the set of vertices with at least one descendant and by \mathcal{L}_j the set of vertices with no descendants. Whereas \mathcal{L}_j is always going to be identified with the collection of functions $\{f_1, \dots, f_n\}$, the collections \mathcal{V}_j on the other hand can be quite different due to the distinct tree structures associated to each parameter. Let $(\xi_l^j)_{1 \leq l \leq n}$ denote the frequency variables for the j -th parameter. To every vertex $v_j \in \mathcal{V}_j$, we associate a symbol $\left| \sum_{l \in \mathcal{L}(v_j)} \xi_l^j \right|^{\beta_j^{v_j}}$ with $\beta_j^{v_j} \geq 0$. The distribution of derivatives is still described by maps $\delta_1 \otimes \dots \otimes \delta_N$, where for every $1 \leq j \leq N$,

$$\delta_j : \mathcal{V}_j \rightarrow \{1, \dots, n\}$$

satisfies the conditions (i) and (ii), with \mathcal{V} replaced by \mathcal{V}_j . We denote by $\mathcal{D}(\mathcal{V}_j)$ the collection of such δ_j s, so that $\mathcal{D}(\mathcal{V}_1) \times \dots \times \mathcal{D}(\mathcal{V}_N)$ represents the collection of admissible distributions of derivatives among the n functions f_1, \dots, f_n .

We now state the Leibniz-type estimates for multi-parameter flag multipliers with asymmetric symbols generated by partial differential operators:

Theorem 1.2. *Let $T_{\mathcal{G}_1 \otimes \dots \otimes \mathcal{G}_N}$ denote the n -linear operator associated to \mathcal{G}_j for $1 \leq j \leq N$ such that each \mathcal{G}_j denotes the frequency tree for the j -th parameter. Suppose that every vertex $v_j \in \mathcal{V}_j$ is associated to a symbol $\left| \sum_{l \in \mathcal{L}(v_j)} \xi_l^j \right|^{\beta_j^{v_j}}$ with $\beta_j^{v_j} \geq 0$. Then for any functions $f_1, \dots, f_n \in \mathcal{S}(\mathbb{R}^{d_1} \times \dots \times \mathbb{R}^{d_N})$, we have*

$$\|T_{\mathcal{G}_1 \otimes \dots \otimes \mathcal{G}_N}(f_1, \dots, f_n)\|_{\vec{r}} \lesssim \sum_{\delta_1 \otimes \dots \otimes \delta_N \in \mathcal{D}(\mathcal{V}_1) \times \dots \times \mathcal{D}(\mathcal{V}_N)} \prod_{l=1}^n \|D_{(1)}^{\delta_1^{-1}(l)} \dots D_{(N)}^{\delta_N^{-1}(l)} f_l\|_{\vec{p}_l},$$

for any $1 \leq p_l^j \leq \infty$, $1 \leq j \leq N$, $1 \leq l \leq n$,

$$\frac{1}{r^j} = \sum_{l=1}^n \frac{1}{p_l^j}, \quad \text{for all } 1 \leq j \leq N,$$

and

$$(1.32) \quad \max \left\{ \frac{d_j}{d_j + \beta_j^{v_j}} : v_j \in \mathcal{V}_j, 1 \leq j \leq N \right\} < \min \{r^j : 1 \leq j \leq N\}.$$

We remark that the condition (1.32) on the Lebesgue exponents is only sufficient, and that in the case when $\beta_j^{v_j} \in 2\mathbb{N}$ it can be disregarded.

Moreover, the same method allows us to prove mixed-norm Leibniz-type estimates for multilinear Mikhlin multipliers of order $\beta > 0$, by systematically reducing them to estimates for linear Mikhlin multipliers. This extends to flags associated to Mikhlin multipliers of strictly positive order, both in one-parameter and in multi-parameter settings.

Let $\beta \in \mathbb{R}$. We say $m(\xi_1, \dots, \xi_n)$ is a Mikhlin symbol of order β provided that $m : \mathbb{R}^{dn} \rightarrow \mathbb{C}$ is smooth away from the origin $\{(\xi_1, \dots, \xi_n) = 0\}$ and satisfies the condition

$$(1.33) \quad |\partial_{\xi_1}^{\gamma_1} \dots \partial_{\xi_n}^{\gamma_n} m(\xi_1, \dots, \xi_n)| \lesssim (|\xi_1| + \dots + |\xi_n|)^{\beta - |\gamma_1| - \dots - |\gamma_n|}$$

for sufficiently many multi-indices $\gamma_1, \dots, \gamma_n$. We denote¹² by \mathcal{M}_β the class of symbols satisfying the above conditions.

Theorem 1.3. *Let $T_{\mathcal{G}}$ denote the n -linear operator indicated by a rooted tree \mathcal{G} of root $\mathbf{r}_{\mathcal{G}}$, where every vertex $v \in \mathcal{V}$ is associated to a symbol $m_v \in \mathcal{M}_{\beta^v}$ of order $\beta^v > 0$. Let \mathcal{D} be the collection of maps defined in (1.19), satisfying conditions (i) and (ii). Then for any functions $f_1, \dots, f_n \in \mathcal{S}(\mathbb{R}^d)$, we have*

$$(1.34) \quad \|T_{\mathcal{G}}(f_1, \dots, f_n)\|_r \lesssim \sum_{\delta \in \mathcal{D}} \prod_{l=1}^n \|D^{\delta^{-1}(l)} f_l\|_{p_l},$$

provided that $1 < p_l < \infty$, $1 \leq l \leq n$, and $1/n < r < \infty$ satisfy the Hölder condition

$$\frac{1}{r} = \frac{1}{p_1} + \dots + \frac{1}{p_n}.$$

Notice that in this situation we fail to recover precisely the L^∞ and L^1 endpoints.

In the N -parameter case, we consider $\beta_1, \dots, \beta_N \in \mathbb{R}$ and we say that m is an N -parameter Mikhlin symbol of order $(\beta_1, \dots, \beta_N)$, simply written as $m \in \mathcal{M}_{\beta_1, \dots, \beta_N}$, if $m : \mathbb{R}^{(d_1 + \dots + d_N)n} \rightarrow \mathbb{C}$ is smooth away from the region $\bigcup_{j=1}^N \{(\xi_1, \dots, \xi_n) \in \mathbb{R}^{(d_1 + \dots + d_N)n} : (\xi_1^j, \dots, \xi_n^j) = 0 \text{ in } \mathbb{R}^{d_j n}\}$ and satisfies the condition

$$(1.35) \quad |\partial_{\xi_1^1}^{\gamma_1^1} \dots \partial_{\xi_1^N}^{\gamma_1^N} \dots \partial_{\xi_n^1}^{\gamma_n^1} \dots \partial_{\xi_n^N}^{\gamma_n^N} m(\xi_1, \dots, \xi_n)| \lesssim \prod_{j=1}^N (|\xi_1^j| + \dots + |\xi_n^j|)^{\beta_j - |\gamma_1^j| - \dots - |\gamma_n^j|}$$

for sufficiently many multi-indices $\gamma_1^1, \dots, \gamma_1^N, \dots, \gamma_n^1, \dots, \gamma_n^N$.

Then we have the following N -parameter result for symbols which tensorize in each parameter:

Theorem 1.4. *Let $T_{\mathcal{G}}$ denote the n -linear operator indicated by a rooted tree \mathcal{G} of root $\mathbf{r}_{\mathcal{G}}$, where every vertex is associated to a symbol m_v satisfying*

$$(1.36) \quad m_v(\xi_{i_1}^1, \dots, \xi_{i_1}^N, \dots, \xi_{i_k}^1, \dots, \xi_{i_k}^N) = \prod_{j=1}^N m_v^j(\xi_{i_1}^j, \dots, \xi_{i_k}^j),$$

with $m_v^j \in \mathcal{M}_{\beta_j^v}$ for $\beta_j^v > 0$. Let \mathcal{D} be the collection of maps defined in (1.19), satisfying conditions (i) and (ii). Then for any functions $f_1, \dots, f_n \in \mathcal{S}(\mathbb{R}^{d_1} \times \dots \times \mathbb{R}^{d_N})$, we have

$$(1.37) \quad \|T_{\mathcal{G}}(f_1, \dots, f_n)\|_{\vec{r}} \lesssim \sum_{\delta_1 \otimes \dots \otimes \delta_N \in \mathcal{D}^N} \prod_{l=1}^n \|D_{(1)}^{\delta_1^{-1}(l)} \dots D_{(N)}^{\delta_N^{-1}(l)} f_l\|_{\vec{p}_l},$$

provided that $1 < p_1^1, \dots, p_1^N, p_2^1, \dots, p_2^N, \dots, p_n^1, \dots, p_n^N < \infty$, $\frac{1}{n} < r^1, \dots, r^N < \infty$ satisfy component-wise the Hölder condition

$$\frac{1}{\vec{r}} = \frac{1}{\vec{p}_1} + \dots + \frac{1}{\vec{p}_n}.$$

Although we expect the non-tensorized equivalent result to remain true in all its generality, that will be analyzed in an upcoming paper. For now we only examine the depth-1 result:

¹²It should be clear from the context what is the space of variables a symbol $m \in \mathcal{M}_\beta$ acts on. Especially for flags, it will be more convenient to leave this implicit, since the number of variables depends on each vertex of the rooted tree.

Theorem 1.5. *If $\beta_1, \dots, \beta_N > 0$ and $m \in \mathcal{M}_{\beta_1, \dots, \beta_N}$ is an N -parameter Mikhlin symbol satisfying (1.35), then the associated multiplier T_m satisfies*

$$\|T_m(f_1, \dots, f_n)\|_{L^{\vec{r}}} \lesssim \sum_{\substack{\sigma_1^1, \dots, \sigma_n^N \in \{0,1\} \\ \sigma_1^1 + \dots + \sigma_n^N = 1}} \prod_{l=1}^n \|D_{(1)}^{\sigma_l^1 \beta_1} \dots D_{(N)}^{\sigma_l^N \beta_N} f_l\|_{\vec{p}_l},$$

for any $1 < p_1^1, \dots, p_1^N, p_2^1, \dots, p_2^N, \dots, p_n^1, \dots, p_n^N < \infty$, $\frac{1}{n} < r^1, \dots, r^N < \infty$ such that component-wise the following Hölder condition holds

$$\frac{1}{\vec{r}} = \frac{1}{\vec{p}_1} + \dots + \frac{1}{\vec{p}_n}.$$

We notice that the symbol does not have to obey the symmetry in (1.36); in fact, we also obtain Leibniz-type estimates for multipliers associated to asymmetric Mikhlin symbols, as long as they tensorize. One such example is

$$\begin{aligned} & T_{\mathcal{G}_1 \otimes \mathcal{G}_2}(f_1, f_2, f_3, f_4, f_5)(x, y) \\ &:= \int_{\mathbb{R}^6} m_1(\xi_1, \xi_2, \xi_3, \xi_4, \xi_5) m_2(\xi_1, \xi_2) m_3(\xi_3, \xi_4, \xi_5) m_4(\xi_3, \xi_4) \tilde{m}_1(\eta_1, \eta_2, \eta_3, \eta_4, \eta_5) \tilde{m}_2(\eta_1, \eta_3) \tilde{m}_3(\eta_2, \eta_4) \\ & \prod_{l=1}^5 \hat{f}_l(\xi_l, \eta_l) e^{2\pi i(x, y) \cdot (\xi_1 + \xi_2 + \xi_3 + \xi_4 + \xi_5, \eta_1 + \eta_2 + \eta_3 + \eta_4 + \eta_5)} d\xi d\eta, \end{aligned} \quad (1.38)$$

which can be perceived as an extension of (1.31) in the off-diagonal regions.

The following theorem gives a general formulation on the boundedness of multipliers associated to tensorized asymmetric Mikhlin symbols:

Theorem 1.6. *Let $T_{\mathcal{G}_1 \otimes \dots \otimes \mathcal{G}_N}$ denote the n -linear operator associated to \mathcal{G}_j for $1 \leq j \leq N$ such that each \mathcal{G}_j denotes the frequency tree for the j -th parameter. Suppose that every vertex $v_j \in \mathcal{V}_j$ is associated to a symbol $m_{v_j}^j \in \mathcal{M}_{\beta_{v_j}^j}$ with $\beta_{v_j}^j > 0$. Let $\mathcal{D}(\mathcal{V}_j)$ be the collection of maps $\delta_j : \mathcal{V}_j \rightarrow \{1, \dots, n\}$ defined in (1.19), satisfying conditions (i) and (ii). Then for any $f_1, \dots, f_n \in \mathcal{S}(\mathbb{R}^{d_1} \times \dots \times \mathbb{R}^{d_N})$,*

$$\|T_{\mathcal{G}_1 \otimes \dots \otimes \mathcal{G}_N}(f_1, \dots, f_n)\|_{\vec{r}} \lesssim \sum_{\delta_1 \otimes \dots \otimes \delta_N \in \mathcal{D}(\mathcal{V}_1) \times \dots \times \mathcal{D}(\mathcal{V}_N)} \prod_{l=1}^n \|D_{(1)}^{\delta_l^{-1}(l)} \dots D_{(N)}^{\delta_l^{-1}(l)} f_l\|_{\vec{p}_l},$$

for any $1 < p_1^1, \dots, p_1^N, p_2^1, \dots, p_2^N, \dots, p_n^1, \dots, p_n^N < \infty$, $\frac{1}{n} < r^1, \dots, r^N < \infty$ satisfying component-wise the Hölder condition

$$\frac{1}{\vec{r}} = \frac{1}{\vec{p}_1} + \dots + \frac{1}{\vec{p}_n}.$$

Finally, the same type of reasoning allows us also to reprove *smoothing properties* of n -linear Mikhlin multipliers associated to a symbol of negative order, in the mixed-norm multi-parameter setting. Our prototypical example consists of the n -linear N -parameter fractional integral operator, whose frequency symbol is given by

$$(|\xi_1^1|^2 + \dots + |\xi_n^1|^2)^{-\frac{\nu_1}{2}} \dots (|\xi_1^N|^2 + \dots + |\xi_n^N|^2)^{-\frac{\nu_N}{2}},$$

for $0 \leq \nu_j \leq nd_j$, for $1 \leq j \leq N$. This is motivated by the works of Hart-Torres-Wu [10] and Yang-Liu-Wu [19], where smoothing properties for less regular multipliers are studied in the mixed-norm and respectively in the bi-parameter setting. More concrete statements and sketches of proofs will be detailed in Section 6.2.

Although the smoothing properties and the results in Theorem 1.5 are not new, our intention here is to illustrate that a careful, quantitative scale-by-scale analysis (which includes improved estimates thanks to the introduction of certain commutators) offers an alternative route to proving them. We will elaborate on this method in the next section, as well as in Section 2.2.

1.2. Strategy. We provide an overview of our methodology and draw a comparison with the approach based on Coifman-Meyer multipliers. For the sake of simplicity, we will focus on dimension one but the discussion can be easily extended to higher dimensions.

To gain some intuition of the Leibniz rules described in Theorem 1.1, we observe that the derivatives capture a function's oscillation rate, and that can be understood through the Fourier transform. Because of (1.27) and an observation that goes back to Bony [3], it is natural to decompose the frequency space into regions

$$\tilde{R}_l := \{(\xi_1, \dots, \xi_n) : |\xi_1 + \dots + \xi_n| \sim |\xi_l|\},$$

since in that case we expect to have $D^\beta((f_1 \dots f_n)|_{\tilde{R}_j}) \sim f_1 \dots f_{l-1} \cdot (D^\beta f_l) \cdot f_{l+1} \dots f_n$.

Indeed, if we restrict our attention to the region¹³ $R_1 := \{(\xi_1, \dots, \xi_n) : |\xi_1| \gg |\xi_2|, \dots, |\xi_n|\}$, we have that

$$\begin{array}{c} |\xi_1 + \dots + \xi_n|^\beta \\ \swarrow \quad \downarrow \quad \searrow \quad \dots \quad \swarrow \\ \hat{f}_1(\xi_1) \quad \hat{f}_2(\xi_2) \quad \dots \quad \hat{f}_n(\xi_n) \end{array} \sim \begin{array}{c} \bullet \\ \swarrow \quad \downarrow \quad \searrow \quad \dots \quad \swarrow \\ |\xi_1|^\beta \hat{f}_1(\xi_1) \quad \hat{f}_2(\xi_2) \quad \dots \quad \hat{f}_n(\xi_n) \end{array}$$

and if $\tilde{\chi}_{R_1}$ is a smooth function adapted to the region R_1 , then

$$|\xi_1 + \dots + \xi_n|^\beta \cdot \tilde{\chi}_{R_1}(\xi_1, \dots, \xi_n) = m(\xi_1, \dots, \xi_n) |\xi_1|^\beta \cdot \tilde{\chi}_{R_1}(\xi_1, \dots, \xi_n),$$

where

$$m(\xi_1, \dots, \xi_n) := \frac{|\xi_1 + \dots + \xi_n|^\beta}{|\xi_1|^\beta} \cdot \tilde{\chi}_{R_1}(\xi_1, \dots, \xi_n).$$

Now the key point is to notice that $m(\xi_1, \dots, \xi_n)$ is a classical multilinear Mikhlin symbol: it is only singular at $\xi_1 = \xi_2 = \dots = \xi_n = 0$ (in the region R_1 , $|\xi_1| = 0$ is equivalent to $(\xi_1, \dots, \xi_n) = 0$), and it decays fast away from the origin. So once the functions f_1, \dots, f_n are jointly restricted in frequency to the region R_1 ,

$$D^\beta(f_1, f_2, \dots, f_n) = T_m(D^\beta f_1, f_2, \dots, f_n).$$

This is, in short,¹⁴ how the boundedness of Coifman-Meyer multipliers (associated to Mikhlin symbols) imply Leibniz-type estimates such as (1.4), (1.8), (1.10), (1.11), etc. Of course, this excludes certain endpoints.

On the other hand, the methodology that we rely on are *commutators* – originally introduced by Bourgain and Li [4] – which manage to capture a certain cancellation between different scales. To start with, we write R_1 as

$$(1.39) \quad R_1 := \bigcup_{k_1 \gg k_2, \dots, k_n \in \mathbb{Z}} R_{k_1, k_2, \dots, k_n} := \bigcup_{k_1 \gg k_2, \dots, k_n} \{(\xi_1, \dots, \xi_n) : |\xi_l| \sim 2^{k_l} \text{ for } 1 \leq l \leq n\}$$

and notice that on R_1 ,

$$(1.40) \quad |\xi_1 + \dots + \xi_n|^\beta = |\xi_1|^\beta + (|\xi_1 + \dots + \xi_n|^\beta - |\xi_1|^\beta).$$

The first term on the right hand side of the above identity seems to be exactly what we wanted: we can indeed replace $|\xi_1 + \dots + \xi_n|^\beta$ by $|\xi_1|^\beta$; for the second term, we notice that it becomes

$$\begin{aligned} & \left(\int_0^1 \beta |\xi_1 + t(\xi_2 + \dots + \xi_n)|^{\beta-2} (\xi_1 + t(\xi_2 + \dots + \xi_n)) dt \right) (\xi_2 + \dots + \xi_n) \\ (1.41) \quad &= \sum_{l=2}^n \left(\int_0^1 \beta |\xi_1 + t(\xi_2 + \dots + \xi_n)|^{\beta-2} (\xi_1 + t(\xi_2 + \dots + \xi_n)) dt \right) \cdot \xi_l \\ &:= \sum_{l=2}^n m_{C_\beta, 1, l}(\xi_1, \xi_2 + \dots + \xi_n). \end{aligned}$$

We restrict $m_{C_\beta, 1, l}$ to the regions $R_{k_1, k_l, \pm}$ defined by

$$\begin{aligned} R_{k_1, k_l, +} &:= \{(\xi_1, \dots, \xi_n) : \xi_1 \sim 2^{k_1}, |\xi_l| \sim 2^{k_l} \ll 2^{k_1}, |\xi_{\tilde{l}}| \ll 2^{k_1} \text{ for } \tilde{l} \neq 1, l\}, \\ R_{k_1, k_l, -} &:= \{(\xi_1, \dots, \xi_n) : \xi_1 \sim -2^{k_1}, |\xi_l| \sim 2^{k_l} \ll 2^{k_1}, |\xi_{\tilde{l}}| \ll 2^{k_1} \text{ for } \tilde{l} \neq 1, l\}, \end{aligned}$$

which are contained in R_{k_1, \dots, k_n} for $k_1 \gg k_2, \dots, k_n$, and denote by $m_{C_\beta, 1, l, \pm}^{k_1, k_l}$ the localized symbol. We notice that $T_{m_{C_\beta, 1, l}^{k_1, k_l, \pm}}$ – the multilinear operator whose symbols is given precisely by $m_{C_\beta, 1, l}^{k_1, k_l, \pm}$ – satisfies

$$(1.42) \quad \|T_{m_{C_\beta, 1, l}^{k_1, k_l, \pm}}\|_{L^{p_1} \times \dots \times L^{p_n} \rightarrow L^p} \lesssim 2^{k_1(\beta-1)} 2^{k_l}, \quad \text{for } 1 \leq p_1, \dots, p_n \leq \infty.$$

This quantified interaction between different scales will allow us to sum over the scales k_1 and k_l , as we will see in more detail in Section 2.2.

For flag Leibniz rules, such as $D^\beta(D^\alpha(f_1 f_2) f_3 D^\gamma(f_4 f_5))$ whose frequency tree representation appears in Figure 1(i), it is natural to restrict our attention to similar frequency regions: if

$$R'_1 = \{(\xi_1, \dots, \xi_5) : |\xi_1| \gg |\xi_2|, |\xi_3|, |\xi_4| \gg |\xi_5|\},$$

¹³Throughout the paper, we say that two positive expressions E_1 and E_2 are equivalent and we write $E_1 \sim E_2$ if there exists $C > 0$ so that $C^{-1}E_1 \leq E_2 \leq CE_1$. Correspondingly, we say that E_1 is much smaller than E_2 and write $E_1 \ll E_2$ if there exists $C > 0$ (which in general will be implicitly depending on the dimension, number of functions involved) so that $E_1 \leq C^{-1}E_2$.

¹⁴For more details, and a comparison with the Bourgain-Li approach, see Section 2.3.

then

$$\begin{aligned} & |\xi_1 + \xi_2 + \xi_3 + \xi_4 + \xi_5|^\beta |\xi_1 + \xi_2|^\alpha |\xi_4 + \xi_5|^\gamma \tilde{\chi}_{R'_1}(\xi_1, \xi_2, \xi_3, \xi_4, \xi_5) \hat{f}_1(\xi_1) \hat{f}_2(\xi_2) \hat{f}_3(\xi_3) \hat{f}_4(\xi_4) \hat{f}_5(\xi_5) \\ &= \frac{|\xi_1 + \xi_2 + \xi_3 + \xi_4 + \xi_5|^\beta}{|\xi_1|^\beta} \frac{|\xi_1 + \xi_2|^\alpha}{|\xi_1|^\alpha} \frac{|\xi_4 + \xi_5|^\gamma}{|\xi_4|^\gamma} \tilde{\chi}_{R'_1}(\xi_1, \xi_2, \xi_3, \xi_4, \xi_5) \widehat{D^{\alpha+\beta} f_1}(\xi_1) \hat{f}_2(\xi_2) \hat{f}_3(\xi_3) \widehat{D^\gamma f_4}(\xi_4) \hat{f}_5(\xi_5). \end{aligned}$$

This implies that the Leibniz rule is a consequence of the boundedness of the 5-linear operator associated to the symbol

$$\frac{|\xi_1 + \xi_2 + \xi_3 + \xi_4 + \xi_5|^\beta}{|\xi_1|^\beta} \frac{|\xi_1 + \xi_2|^\alpha}{|\xi_1|^\alpha} \frac{|\xi_4 + \xi_5|^\gamma}{|\xi_4|^\gamma} \tilde{\chi}_{R'_1}(\xi_1, \xi_2, \xi_3, \xi_4, \xi_5),$$

which can also be represented as

$$(1.43) \quad m_\beta(\xi_1, \xi_2, \xi_3, \xi_4, \xi_5) \cdot m_\alpha(\xi_1, \xi_2) \cdot m_\gamma(\xi_4, \xi_5),$$

where $m_\beta(\xi_1, \xi_2, \xi_3, \xi_4, \xi_5)$, $m_\alpha(\xi_1, \xi_2)$, $m_\gamma(\xi_4, \xi_5)$ are all Mikhlin multipliers when smoothly restricted to the region R'_1 . Expressions such as (1.43) are prototypes of symbols associated to flag paraproducts – compare to the definition in (1.13).

We will however choose a different path for estimating $D^\beta(D^\alpha(f_1 f_2) f_3 D^\gamma(f_4 f_5))$, which is closer to the Bourgain-Li approach, since it will allow us to also treat the multi-parameter flag Leibniz rules. The main steps of the strategy are:

- (1) splitting of the root symbol and appearance of commutators in the off-diagonal frequency regions;
- (2) Fourier series decomposition for the symbols;
- (3) tensorization into frequency-localized subtrees.

In the treatment of Leibniz rules of complexity 1, Step 1 distributes the derivative to a leaf (1.40) while for a generic flag Leibniz rule, it will pass the derivative from the root to one of its direct descendants. This, together with the application of Step 2, allows us to reduce the estimation of the original tree to subtrees of lower complexities, which leads to Step 3.

The same methodology can be employed to develop Leibniz-type estimates for operators associated to Mikhlin symbols of order β for $\beta > 0$. We observe that in the region R_1 given by (1.39), the condition (1.33) satisfied by the Mikhlin symbol $m_\beta(\xi_1, \dots, \xi_n)$ can be reformulated as

$$|\partial_{\xi_1}^{\gamma_1} \dots \partial_{\xi_n}^{\gamma_n} m_\beta(\xi_1, \dots, \xi_n)| \lesssim (|\xi_1| + \dots + |\xi_n|)^{\beta - |\gamma_1| - \dots - |\gamma_n|} \sim |\xi_1|^{\beta - |\gamma_1| - \dots - |\gamma_n|}.$$

Such a heuristic computation suggests that $m_\beta(\xi_1, \xi_2, \dots, \xi_n)$ behaves like $m_\beta(\xi_1, 0, \dots, 0)$ on R_1 . We can make it rigorous by invoking Step 1 and introducing the commutator

$$m_\beta(\xi_1, 0, \dots, 0) + (m_\beta(\xi_1, \xi_2, \dots, \xi_n) - m_\beta(\xi_1, 0, \dots, 0))$$

and the scale-by-scale analysis is applicable in this setting as well.

We remark that although this strategy works for obtaining Leibniz-type estimates for Mikhlin multipliers with symbols of strictly positive order, it fails in the case of classical Mikhlin symbols, which correspond to order zero.

To extend the methodology to the multi-parameter setting, we notice that for multipliers which are tensor product of symbols in each parameter – such as the ones involved in Theorem 1.1 and Theorem 1.4 – Steps 1 and 2 can be performed independently in each parameter. Step 3 – which allows a decoupling of the rooted tree into subtrees of lower complexities – can be attained thanks to Step 2; further computations concern the distribution of derivatives – this is carried out independently for each parameter so that the mixed Besov and Lebesgue norms (see Section 2.1) naturally appear.

1.3. Structure of the paper. The paper is organized as follows: in Section 2 we introduce the necessary terminology and review the Bourgain-Li approach from [4], putting the accent on certain novel aspects that will be involved in treating the mixed-norm multi-parameter generic flag Leibniz rule. In Section 3 we present a 5-linear flag Leibniz rule, and in Section 4 its bi-parameter version; these particular examples are interesting enough to capture the complexity of the general case, without being too technical. In Section 5 we present in detail the inductive proof of our main Theorem 1.1 and illustrate briefly a modified induction requested by Theorem 1.2. Finally, in Section 6 we discuss Leibniz-type estimates for generic Mikhlin multipliers of positive order (Theorem 1.3, 1.4, 1.5) and recover the smoothing properties described in Theorem 6.3.

1.4. Acknowledgements. C. Benea acknowledges partial support from ANR project RAGE ANR-18-CE40-0012 and research grant PN-III-P1-1.1-TE-2019-2275 from UEFISCDI, Romania. Y. Zhai's research is supported by ERC project FAnFARe no. 637510 and the region Pays de la Loire.

2. NOTATION AND USEFUL RESULTS

In this section we set the notation that will be used throughout the paper, and review the Bourgain-Li method from [4].

2.1. Littlewood-Paley projections and Besov spaces.

Let $N \geq 1$, and let $\vec{p} = (p^1, \dots, p^N)$ be an N -tuple of positive Lebesgue exponents: that is, we assume that $0 < p^1, \dots, p^N \leq \infty$. For functions on $\mathbb{R}^{d_1} \times \dots \times \mathbb{R}^{d_N}$, we define the mixed (quasi)norm

$$(2.1) \quad \|f\|_{L^{\vec{p}}} = \|f\|_{L^{p^1}_{x_1} L^{p^2}_{x_2} \dots L^{p^N}_{x_N}} := \left(\int_{\mathbb{R}^{d_1}} \dots \int_{\mathbb{R}^{d_{N-1}}} \left(\int_{\mathbb{R}^{d_N}} |f(x_1, \dots, x_{N-1}, x_N)|^{p^N} dx_N \right)^{\frac{p^{N-1}}{p^N}} dx_{N-1} \right)^{\frac{p^{N-2}}{p^{N-1}}} \dots dx_1 \right)^{\frac{1}{p^1}}.$$

Whenever one of the p^j is equal to ∞ , the integration in the x_j variable is replaced by taking the essential supremum with respect to x_j .

Then the space $L^{\vec{p}}(\mathbb{R}^{d_1+\dots+d_N})$ consists of all the functions defined on $\mathbb{R}^{d_1+\dots+d_N}$, with finite $\|\cdot\|_{L^{\vec{p}}}$ norm:

$$(2.2) \quad L^{\vec{p}}(\mathbb{R}^{d_1+\dots+d_N}) := \{f : \mathbb{R}^{d_1+\dots+d_N} \rightarrow \mathbb{C} : \|f\|_{L^{\vec{p}}} < \infty\}.$$

Remark 2.1. We record a few useful properties of mixed-norm spaces:

- (1) If $\vec{p} = (p^1, \dots, p^N)$ with $1 \leq p^j \leq \infty$ for all $1 \leq j \leq N$, then $\|\cdot\|_{L^{\vec{p}}}$ is a norm and $L^{\vec{p}}(\mathbb{R}^{d_1+\dots+d_N})$ is a Banach space.
- (2) Generally, $\|\cdot\|_{L^{\vec{p}}}$ is a quasi-norm and $L^{\vec{p}}(\mathbb{R}^{d_1+\dots+d_N})$ is a quasi-Banach space.
- (3) If $\tau > 0$ is so that

$$\tau \leq \min \left(1, \min_{1 \leq j \leq N} p^j \right),$$

then $\|\cdot\|_{L^{\vec{p}}}^\tau$ is subadditive.

Throughout the paper, we make use of the classical Littlewood-Paley decompositions.¹⁵ On \mathbb{R}^d , we start with $\varphi : \mathbb{R}^d \rightarrow [0, \infty)$ a radial function so that $0 \leq \varphi \leq 1$, $\varphi \equiv 1$ on $\{|\xi| \leq 1\}$, $\varphi \equiv 0$ on $\{|\xi| \geq 2\}$; then we define $\psi(\xi) := \varphi(\xi) - \varphi(2\xi)$, which is supported on $\{1/2 \leq |\xi| \leq 2\}$, and we obtain

$$\sum_{k \in \mathbb{Z}} \psi(2^{-k}\xi) = 1, \quad \text{for all } \xi \neq 0.$$

Hence for any Schwartz function $f \in \mathcal{S}(\mathbb{R}^d)$,

$$\hat{f}(\xi) = \sum_{k \in \mathbb{Z}} \psi(2^{-k}\xi) \hat{f}(\xi) \quad \text{for all } \xi \neq 0.$$

If we denote $\psi_k(\xi) := \psi(2^{-k}\xi)$, then the identity above reads in space as

$$f(x) = \sum_{k \in \mathbb{Z}} f * \check{\psi}_k(x).$$

For any $k \in \mathbb{Z}$, Δ_k denotes the Littlewood-Paley projection associated to the frequency region $\{|\xi| \sim 2^k\}$:

$$f \mapsto \Delta_k f := f * \check{\psi}_k = \mathcal{F}^{-1}(\hat{f} \cdot \psi_k).$$

Then $f = \sum_{k \in \mathbb{Z}} \Delta_k f$, and since the functions $\check{\psi}_k$ are L^1 -normalized,¹⁶ we have uniformly in $k \in \mathbb{Z}$,

$$(2.3) \quad \|\Delta_k f\|_p \leq \|\check{\psi}\|_1 \|f\|_p, \quad \text{for any } 1 \leq p \leq \infty.$$

For functions on $\mathbb{R}^{d_1} \times \dots \times \mathbb{R}^{d_N}$, we consider the multi-parameter Littlewood-Paley decomposition

$$f = \sum_{k_1, \dots, k_N \in \mathbb{Z}} \Delta_{k_1}^{(1)} \Delta_{k_2}^{(2)} \dots \Delta_{k_N}^{(N)} f,$$

where

$$(2.4) \quad \Delta_{k_j}^{(j)} f(x_1, \dots, x_{j-1}, x_j, x_{j+1}, \dots, x_N) := f *_{x_j} \check{\psi}_{k_j}(x_1, \dots, x_N) = \int_{\mathbb{R}^{d_j}} f(x_1, \dots, x_{j-1}, x_j - t, x_{j+1}, \dots, x_N) \check{\psi}_{k_j}(t) dt.$$

¹⁵Small perturbations of the base functions φ and ψ will not change the inherent properties of the Littlewood-Paley families $\{\psi(2^k \cdot)\}_{k \in \mathbb{Z}}$. These perturbations will be denoted generically $\tilde{\varphi}$ and $\tilde{\psi}$.

¹⁶That is, $\|\check{\psi}_k\|_1 = \|\check{\psi}\|_1$ uniformly in k .

As before in (2.3), we have $\|\Delta_{k_j}^{(j)} f\|_{L^p(\mathbb{R}^{d_1+\dots+d_N})} \lesssim \|f\|_{L^p(\mathbb{R}^{d_1+\dots+d_N})}$ for any $1 \leq p \leq \infty$, and moreover, the mixed-norm estimate

$$(2.5) \quad \|\Delta_{k_j}^{(j)} f\|_{\vec{p}} \lesssim \|f\|_{\vec{p}}$$

holds for any $\vec{p} = (p^1, \dots, p^N)$ with $1 \leq p^1, \dots, p^N \leq \infty$. This is due to the observation that, for almost every (x_1, \dots, x_j) ,

$$\|\Delta_{k_j}^{(j)} f(x_1, \dots, x_{j-1}, x_j, \cdot)\|_{L_{x_{j+1}}^{p^{j+1}} \dots L_{x_N}^{p^N}} \lesssim \int_{\mathbb{R}^{d_j}} \|f(x_1, \dots, x_{j-1}, x_j - t, \cdot)\|_{L_{x_{j+1}}^{p^{j+1}} \dots L_{x_N}^{p^N}} |\check{\psi}_{k_j}^{(j)}(t)| dt,$$

which is nothing but a direct application of Minkowski's integral inequality (which is appropriate since $1 \leq p^1, \dots, p^N \leq \infty$). Then we use Young's convolution inequality in the x_j variable and integrate in the remaining variables to obtain (2.5).

More generally,¹⁷ we obtain

$$\|\Delta_{k_1}^{(1)} \Delta_{k_2}^{(2)} \dots \Delta_{k_N}^{(N)} f\|_{\vec{p}} \lesssim \|f\|_{\vec{p}}, \quad \text{for any } \vec{p} = (p^1, \dots, p^N) \text{ with } 1 \leq p^1, \dots, p^N \leq \infty.$$

Next, for any $s \in \mathbb{R}$ and any $1 \leq p \leq \infty$, we introduce the homogeneous Besov norms $\|\cdot\|_{\dot{B}_{p,\infty}^s}$ on $\mathcal{S}(\mathbb{R}^d)$ as

$$(2.6) \quad \|f\|_{\dot{B}_{p,\infty}^s} := \sup_{k \in \mathbb{Z}} 2^{ks} \|\Delta_k f\|_p.$$

A straightforward, but important observation is the inequality

$$(2.7) \quad \|f\|_{\dot{B}_{p,\infty}^s} \lesssim \|D^s f\|_p,$$

which holds true whenever the right hand side is finite.

Similarly, the N -parameter Besov norms of functions in $\mathcal{S}(\mathbb{R}^{d_1} \times \dots \times \mathbb{R}^{d_N})$ are defined by

$$(2.8) \quad \|f\|_{\dot{B}_{p^1,\infty}^{s_1} \dots \dot{B}_{p^N,\infty}^{s_N}} := \sup_{k_1, \dots, k_N} 2^{k_1 s_1} \dots 2^{k_N s_N} \|\Delta_{k_1}^{(1)} \Delta_{k_2}^{(2)} \dots \Delta_{k_N}^{(N)} f\|_{\vec{p}}.$$

One can also consider mixed Besov and Lebesgue norms: let $1 \leq m \leq N$ and $i_1, \dots, i_m \in \{1, \dots, N\}$, and $s_{i_1}, \dots, s_{i_m} \in \mathbb{R}$; then for any function $f \in \mathcal{S}(\mathbb{R}^{d_1} \times \dots \times \mathbb{R}^{d_N}) \rightarrow \mathbb{C}$, we define

$$(2.9) \quad \|f\|_{L^{p^1} \dots L^{p^{i_1-1}} \dot{B}_{p^{i_1},\infty}^{s_{i_1}} L^{p^{i_1+1}} \dots L^{p^{i_2-1}} \dot{B}_{p^{i_2},\infty}^{s_{i_2}} L^{p^{i_2+1}} \dots L^{p^N}} := \sup_{k_{i_1}, \dots, k_{i_m}} 2^{k_{i_1} s_{i_1}} \dots 2^{k_{i_m} s_{i_m}} \|\Delta_{k_{i_1}}^{(i_1)} \Delta_{k_{i_2}}^{(i_2)} \dots \Delta_{k_{i_m}}^{(i_m)} f\|_{\vec{p}}.$$

Since

$$2^{k_j s_j} \|\Delta_{k_j}^{(j)} f\|_{\vec{p}} = \|\tilde{\Delta}_{k_j}^{(j)} D_{(j)}^{s_j} f\|_{\vec{p}} \lesssim \|D_{(j)}^{s_j} f\|_{\vec{p}},$$

for a slightly different Littlewood-Paley projection $\tilde{\Delta}_{k_j}^{(j)}$ having similar support properties, we deduce that in general

$$\|f\|_{L^{p^1} \dots L^{p^{i_1-1}} \dot{B}_{p^{i_1},\infty}^{s_{i_1}} L^{p^{i_1+1}} \dots L^{p^{i_2-1}} \dot{B}_{p^{i_2},\infty}^{s_{i_2}} L^{p^{i_2+1}} \dots L^{p^N}} \lesssim \|D_{(i_1)}^{s_{i_1}} \dots D_{(i_m)}^{s_{i_m}} f\|_{\vec{p}},$$

provided all the Lebesgue indices are in the Banach regime: $\vec{p} = (p^1, \dots, p^N)$ with $1 \leq p^1, \dots, p^N \leq \infty$. We will also need a mixed-norm interpolation result: for any $-\infty < s_0 < s_1 < \infty$ and any $0 \leq \theta \leq 1$ so that $s := \theta s_0 + (1 - \theta) s_1$,

$$(2.10) \quad \|f\|_{X_1 \dots X_{j-1} \dot{B}_{p^j,\infty}^{s_j} X_{j+1} \dots X_N} \lesssim \|f\|_{X_1 \dots X_{j-1} \dot{B}_{p^j,\infty}^{s_0} X_{j+1} \dots X_N}^\theta \cdot \|f\|_{X_1 \dots X_{j-1} \dot{B}_{p^j,\infty}^{s_1} X_{j+1} \dots X_N}^{1-\theta},$$

where the norms $\|\cdot\|_{X_i}$, for $i \in \{1, \dots, N\} \setminus \{j\}$, denote either a Lebesgue $\|\cdot\|_{L^{p^i}}$ or a Besov $\|\cdot\|_{\dot{B}_{p^i,\infty}^{s_i}}$ norm. This

is a straightforward consequence of the identity $2^{ks} = (2^{ks_0})^\theta (2^{ks_1})^{1-\theta}$.

Oftentimes, this interpolation inequality will be used in the form

$$(2.11) \quad \|f\|_{X_1 \dots X_{j-1} \dot{B}_{p^j,\infty}^{\epsilon} X_{j+1} \dots X_N} \lesssim \|f\|_{X_1 \dots X_{j-1} \dot{B}_{p^j,\infty}^0 X_{j+1} \dots X_N}^{\frac{\beta-\epsilon}{\beta}} \cdot \|f\|_{X_1 \dots X_{j-1} \dot{B}_{p^j,\infty}^{\beta} X_{j+1} \dots X_N}^{\frac{\epsilon}{\beta}},$$

where¹⁸ $0 < \epsilon < \beta$.

¹⁷This can be further extended by replacing each $\|\cdot\|_{L^{p^j}(\mathbb{R}^{d_j})}$ norm with the mixed-norm $\|\cdot\|_{L^{p_1^j} L^{p_2^j} \dots L^{p_{d_j}^j}}$, as long as all the Lebesgue exponents are between 1 and ∞ .

¹⁸In Section 6.2, the same inequality with $\beta < \epsilon < 0$ will be needed.

Before we proceed, we need to introduce a few extra operators and their properties. Recalling the definition of Δ_k and ψ_k , we define¹⁹

$$(2.12) \quad S_k f(x) := \sum_{\ell \ll k} \Delta_\ell f(x),$$

which is of the form $S_k f(x) = f * \tilde{\varphi}_k(x)$, where $\tilde{\varphi}_k(x) = 2^{dk} \tilde{\varphi}(2^k x)$. As a consequence,

$$\|S_k f\|_p \lesssim \|f\|_p \quad \text{for all } 1 \leq p \leq \infty,$$

uniformly in $k \in \mathbb{Z}$. Similarly, $\Delta_{\leq k} f := \sum_{\ell \leq k} \Delta_\ell f$ satisfies

$$\|\Delta_{\leq k} f\|_p \lesssim \|f\|_p \quad \text{for all } 1 \leq p \leq \infty \text{ and for all } k \in \mathbb{Z}.$$

On the other hand,

$$(2.13) \quad \Delta_{>k} f(x) := \sum_{\ell > k} \Delta_\ell f(x) = \sum_{\ell \geq k-c} \Delta_\ell f(x),$$

and since $\Delta_{>k} f(x) + S_k f(x) = f(x)$, we again have

$$(2.14) \quad \|\Delta_{>k} f\|_p \lesssim \|f\|_p \quad \text{for all } 1 \leq p \leq \infty,$$

uniformly in $k \in \mathbb{Z}$.

For functions on \mathbb{R} , we also define the directional projection operators

$$(2.15) \quad \Delta_{k,+} f(x) := \mathcal{F}^{-1}(\hat{f} \cdot \psi_{k,+})(x), \quad \Delta_{k,-} f(x) := \mathcal{F}^{-1}(\hat{f} \cdot \psi_{k,-})(x)$$

where $\psi_{k,+}(\xi) := \psi_k(\xi) \chi_{\{\xi \geq 0\}}$ and $\psi_{k,-}(\xi) := \psi_k(\xi) \chi_{\{\xi < 0\}}$. In dimension one, the region $\{|\xi| \sim 2^k\}$ naturally splits into two intervals, namely $\{\xi \sim 2^k\}$ and $\{\xi \sim -2^k\}$, which justifies the notion of “directional” projection. We observe that $\Delta_{k,+} f = \tilde{\Delta}_{k,+} \Delta_k f$ for $\tilde{\Delta}_{k,+}$ associated to $\tilde{\psi}_{k,+}$, a function which is $\equiv 1$ on $\{2^{k-1} \leq \xi \leq 2^{k+1}\}$ (similarly for $\Delta_{k,-} f = \tilde{\Delta}_{k,-} \Delta_k f$), so that

$$(2.16) \quad \|\Delta_{k,+} f\|_p, \|\Delta_{k,-} f\|_p \lesssim \|\Delta_k f\|_p \lesssim \|f\|_p \quad \text{for all } 1 \leq p \leq \infty.$$

All these definitions can be reformulated as Fourier projections onto the j th coordinate:

$$(2.17) \quad S_{k_j}^{(j)} f(x) := \sum_{\ell \ll k_j} \Delta_\ell^{(j)} f(x), \quad \text{and} \quad \Delta_{>k_j}^{(j)} f(x) := \sum_{\ell > k_j-c} \Delta_\ell^{(j)} f(x).$$

For any function $f \in \mathcal{S}(\mathbb{R}^N)$, we define the directional projection operator on the j -th parameter by

$$\Delta_{k_j,\pm}^{(j)} f(x_1, \dots, x_{j-1}, x_j, x_{j+1}, \dots, x_N) = \int_{\mathbb{R}^{d_j}} f(x_1, \dots, x_{j-1}, x_j - t, x_{j+1}, \dots, x_N) \tilde{\psi}_{k_j,\pm}(t) dt.$$

As before, for any $\vec{p} = (p^1, \dots, p^N)$ with $1 \leq p^1, \dots, p^N \leq \infty$,

$$\|S_{k_j}^{(j)} f\|_{\vec{p}}, \|\Delta_{>k_j}^{(j)} f\|_{\vec{p}} \lesssim \|f\|_{\vec{p}} \quad \text{and} \quad \|\Delta_{k_j,+}^{(j)} f\|_{\vec{p}}, \|\Delta_{k_j,-}^{(j)} f\|_{\vec{p}} \lesssim \|\Delta_{k_j}^{(j)} f\|_{\vec{p}} \lesssim \|f\|_{\vec{p}}.$$

The Fourier series decomposition, which plays an important role in tensorizing the operator associated to a flag Leibniz rule into subtrees of lower complexity, will introduce certain modulations, which are however inconsequential: if $P_k^{(j)}$ is any of the projections $\Delta_k^{(j)}$, $S_k^{(j)}$ or directional projections $\Delta_{k,+}^{(j)}$, $\Delta_{k,-}^{(j)}$ (so that $P_k^{(j)}$ can also written as a convolution with the function $2^{kd_j} \tilde{\phi}(2^k x_j)$), then for any $a \in \mathbb{R}^{d_j}$, $P_{k,a}^{(j)}$ denotes

$$(2.18) \quad P_{k,a}^{(j)} f(x_1, \dots, x_{j-1}, x_j, x_{j+1}, \dots, x_N) := \int_{\mathbb{R}^{d_j}} \phi(2^{-k} \xi_j) e^{2\pi i a \cdot \frac{\xi_j}{2^k}} (\mathcal{F}^{(j)} f)(x_1, \dots, x_{j-1}, \xi_j, x_{j+1}, \dots, x_N) e^{2\pi i x_j \cdot \xi_j} d\xi_j.$$

But $P_{k,a}^{(j)} f$ is simply the convolution in the j th coordinate between f and $\tilde{\phi}_k$, evaluated at $x_j + \frac{a}{2^k}$:

$$P_{k,a}^{(j)} f(x_1, \dots, x_j, \dots, x_N) = f *_{x_j} \tilde{\phi}_k(x_1, \dots, x_j + \frac{a}{2^k}, \dots, x_N).$$

Due to the trivial identity $\|P_{k,a}^{(j)} f\|_{\vec{p}} = \|P_k^{(j)} f\|_{\vec{p}}$, we deduce

$$(2.19) \quad \|S_{k_j,a}^{(j)} f\|_{\vec{p}} \lesssim \|f\|_{\vec{p}} \quad \text{and} \quad \|\Delta_{k_j,a}^{(j)} f\|_{\vec{p}}, \|\Delta_{k_j,+}^{(j)} f\|_{\vec{p}}, \|\Delta_{k_j,-}^{(j)} f\|_{\vec{p}} \lesssim \|\Delta_{k_j}^{(j)} f\|_{\vec{p}} \lesssim \|f\|_{\vec{p}},$$

for any $\vec{p} = (p^1, \dots, p^N)$ with $1 \leq p^1, \dots, p^N \leq \infty$.

¹⁹In what follows, we want to make sure that the scale 2^ℓ is much smaller than 2^k : $2^\ell \ll 2^k$; this translates into assuming the existence of $c > 0$ large enough (depending implicitly on the dimension, the number of functions involved, etc) so that $\ell \leq k - c$. Moreover, when $2^\ell \ll 2^k$ (so that $\ell > k - c$), we write $2^\ell > 2^k$ or equivalently $\ell > k$.

2.2. A review of the Bourgain-Li approach.

Here we present in dimension one²⁰ some elements of the Bourgain-Li proof of the bilinear Leibniz rule

$$(2.20) \quad \|D^\alpha(fg)\|_{L^p} \lesssim \|D^\alpha f\|_{L^{p_1}} \|g\|_{L^{p_2}} + \|f\|_{L^{p_1}} \|D^\alpha g\|_{L^{p_2}}$$

where $1 \leq p_1, p_2 \leq \infty$, $\frac{1}{1+\alpha} < p \leq \infty$, $\frac{1}{p_1} + \frac{1}{p_2} = \frac{1}{p}$, and $\alpha \geq 0$. Due to the introduction of “commutators”, the use of Coifman-Meyer multipliers can be completely avoided, thus extending the range of Leibniz rules beyond that of Coifman-Meyer multipliers. In distinction to [4], here we quantify the improvement produced by the commutator terms as an interaction between different scales, which in turn requires a suitable double Fourier series decomposition.


We start with Littlewood-Paley decompositions for the functions f and g

$$f = \sum_{k \in \mathbb{Z}} \Delta_k f, \quad g = \sum_{\ell \in \mathbb{Z}} \Delta_\ell g,$$

so that $D^\alpha(fg)$ becomes

$$(2.21) \quad \sum_{k, \ell \in \mathbb{Z}} \int_{\mathbb{R}^2} |\xi_1 + \xi_2|^\alpha \widehat{\Delta_k f}(\xi_1) \widehat{\Delta_\ell g}(\xi_2) e^{2\pi i x(\xi_1 + \xi_2)} d\xi_1 d\xi_2.$$


We have several possibilities, depending whether $|\xi_1 + \xi_2| \sim |\xi_2| \gg |\xi_1|$, $|\xi_1 + \xi_2| \sim |\xi_1| \gg |\xi_2|$, or $|\xi_1| \sim |\xi_2| \geq |\xi_1 + \xi_2|$:



This allows us to decompose $D^\alpha(f \cdot g)$ as

$$\begin{aligned} & \sum_{k \ll \ell} \int_{\mathbb{R}^2} |\xi_1 + \xi_2|^\alpha \widehat{\Delta_k f}(\xi_1) \widehat{\Delta_\ell g}(\xi_2) e^{2\pi i x(\xi_1 + \xi_2)} d\xi_1 d\xi_2 \\ & + \sum_{k \gg \ell} \int_{\mathbb{R}^2} |\xi_1 + \xi_2|^\alpha \widehat{\Delta_k f}(\xi_1) \widehat{\Delta_\ell g}(\xi_2) e^{2\pi i x(\xi_1 + \xi_2)} d\xi_1 d\xi_2 \\ & + \sum_{|k - \ell| \leq 2} \int_{\mathbb{R}^2} |\xi_1 + \xi_2|^\alpha \widehat{\Delta_k f}(\xi_1) \widehat{\Delta_\ell g}(\xi_2) e^{2\pi i x(\xi_1 + \xi_2)} d\xi_1 d\xi_2 := I + II + III, \end{aligned}$$

as suggested by the figure below:



We study each of the cases I and III by taking a closer look at the associated multiplier – this will be sufficient since I and II are symmetric. We highlight the main steps:

(1) In treating I , we approximate

$$|\xi_1 + \xi_2|^\alpha = |\xi_2|^\alpha + \text{“error”},$$

so that it becomes

$$(2.22) \quad \sum_{k \ll \ell} \int_{\mathbb{R}^2} |\xi_2|^\alpha \widehat{\Delta_k f}(\xi_1) \widehat{\Delta_\ell g}(\xi_2) e^{2\pi i x(\xi_1 + \xi_2)} d\xi_1 d\xi_2$$

$$(2.23) \quad + \sum_{k \ll \ell} \int_{\mathbb{R}^2} (|\xi_1 + \xi_2|^\alpha - |\xi_2|^\alpha) \widehat{\Delta_k f}(\xi_1) \widehat{\Delta_\ell g}(\xi_2) e^{2\pi i x(\xi_1 + \xi_2)} d\xi_1 d\xi_2$$

$$(2.24) \quad := \sum_{k \ll \ell} \Delta_k f(x) \cdot (\Delta_\ell D^\alpha g)(x) + \sum_{k \ll \ell} [D^\alpha, \Delta_k f] \Delta_\ell g(x).$$

The term $[D^\alpha, \Delta_k f] \Delta_\ell g(x)$ represents a *commutator*; we will see that it behaves better than the initial $D^\alpha(\Delta_k f \cdot \Delta_\ell g)$, in a way that can be expressed quantitatively.

²⁰This assumption allows for a simplification of the notations, without restricting the method’s generality – see Remark 2.4.

(2) In estimating the first part, it is convenient to switch the order of summation, which produces

$$\begin{aligned} & \sum_{k,\ell} \Delta_k f(x) \cdot (\Delta_\ell D^\alpha g)(x) - \sum_{k>\ell} \Delta_k f(x) \cdot (\Delta_\ell D^\alpha g)(x) \\ &= f(x) \cdot D^\alpha g(x) - \sum_{k>\ell} \Delta_k f(x) \cdot (\Delta_\ell D^\alpha g)(x). \end{aligned}$$

(3) With this, I is converted into

$$\begin{aligned} I &= f(x) \cdot D^\alpha g(x) - \sum_{k>\ell} \Delta_k f(x) \cdot (\Delta_\ell D^\alpha g)(x) + \sum_{k\ll\ell} [D^\alpha, \Delta_k f] \Delta_\ell g(x) \\ &:= f(x) \cdot D^\alpha g(x) - I_A + I_B. \end{aligned}$$

(4) The term III reduces essentially to

$$\sum_{\ell \in \mathbb{Z}} \int_{\mathbb{R}^2} |\xi_1 + \xi_2|^\alpha \widehat{\Delta_\ell f}(\xi_1) \widehat{\Delta_\ell g}(\xi_2) e^{2\pi i x(\xi_1 + \xi_2)} d\xi_1 d\xi_2.$$

Next, we claim that it is sufficient to have precise estimates for the corresponding bilinear operators, with scales k and ℓ fixed.

Lemma 2.2. *Let $1 \leq p_1, p_2 \leq \infty$, $\frac{1}{2} \leq p \leq \infty$ satisfy $\frac{1}{p_1} + \frac{1}{p_2} = \frac{1}{p}$, and $k, \ell \in \mathbb{Z}$. Then we have*

$$(2.25) \quad \|\Delta_k f \cdot (\Delta_\ell D^\alpha g)\|_p \lesssim 2^{\ell\alpha} \|\Delta_k f\|_{p_1} \|\Delta_\ell g\|_{p_2}.$$

If $k \ll \ell$,

$$(2.26) \quad \|[D^\alpha, \Delta_k f] \Delta_\ell g\|_p \lesssim 2^{(\alpha-1)\ell} 2^k \|\Delta_k f\|_{p_1} \|\Delta_\ell g\|_{p_2}$$

and

$$(2.27) \quad \|D^\alpha (S_\ell f \Delta_\ell g)\|_p \lesssim 2^{\ell\alpha} \|S_\ell f\|_{p_1} \|\Delta_\ell g\|_{p_2}.$$

Under the additional assumption that $\frac{1}{\alpha+1} < p \leq \infty$,

$$(2.28) \quad \|D^\alpha (\Delta_\ell f \cdot \Delta_\ell g)\|_p \lesssim 2^{\ell\alpha} \|\Delta_\ell f\|_{p_1} \|\Delta_\ell g\|_{p_2}.$$

Now we show how the fixed-scale estimates listed above allow us to control the terms I and III , and thus prove (2.20). Once that concluded, we will return to the proof of Lemma 2.2, since it allows to illustrate some of the main ideas needed for dealing with the more general Theorem 1.1.

Proof of the Leibniz rule (2.20) assuming Lemma 2.2. We let $\tau \leq \min(1, p)$ so that $\|\cdot\|_{L^p}^\tau$ is subadditive.

- estimating III :

Using (2.28) of Lemma 2.2 and the Besov norms definitions (2.6), we have for $\frac{1}{1+\alpha} < p \leq \infty$ and $\tau \leq \min(1, p)$:

$$\begin{aligned} \|III\|_p^\tau &\lesssim \sum_{\ell \in \mathbb{Z}} 2^{\ell\alpha\tau} \|\Delta_\ell f\|_{p_1}^\tau \|\Delta_\ell g\|_{p_2}^\tau \\ &\lesssim \sum_{\ell \in \mathbb{Z}} \min(2^{\ell\alpha\tau} \|f\|_{\dot{B}_{p_1,\infty}^0}^\tau \|g\|_{\dot{B}_{p_2,\infty}^0}^\tau, 2^{-\ell\alpha\tau} \|f\|_{\dot{B}_{p_1,\infty}^\alpha}^\tau \|g\|_{\dot{B}_{p_2,\infty}^\alpha}^\tau). \end{aligned}$$

Optimizing in ℓ , it is not difficult to see that

$$\begin{aligned} \|III\|_p &\lesssim \|f\|_{\dot{B}_{p_1,\infty}^\alpha}^{\frac{1}{2}} \|g\|_{\dot{B}_{p_2,\infty}^\alpha}^{\frac{1}{2}} \|f\|_{\dot{B}_{p_1,\infty}^0}^{\frac{1}{2}} \|g\|_{\dot{B}_{p_2,\infty}^0}^{\frac{1}{2}} \\ &\lesssim \|D^\alpha f\|_{L^{p_1}} \|g\|_{L^{p_2}} + \|f\|_{L^{p_1}} \|D^\alpha g\|_{L^{p_2}}. \end{aligned}$$

- estimating I_A :

For any $k, \ell \in \mathbb{Z}$ and $0 < \epsilon < \alpha$ we have, as a consequence of (2.25), the following two inequalities:

$$(2.29) \quad \|\Delta_k f \cdot (\Delta_\ell D^\alpha g)\|_p \lesssim 2^{\ell\alpha} \|\Delta_k f\|_{p_1} \|\Delta_\ell g\|_{p_2} \lesssim 2^{\ell\alpha} \|f\|_{\dot{B}_{p_1,\infty}^0} \|g\|_{\dot{B}_{p_2,\infty}^0}$$

$$(2.30) \quad \|\Delta_k f \cdot (\Delta_\ell D^\alpha g)\|_p \lesssim 2^{-k\alpha} 2^{\ell(\alpha-\epsilon)} (2^{k\alpha} \|\Delta_k f\|_{p_1}) (2^{\ell\epsilon} \|\Delta_\ell g\|_{p_2}) \lesssim 2^{-k\alpha} 2^{\ell(\alpha-\epsilon)} \|f\|_{\dot{B}_{p_1,\infty}^\alpha} \|g\|_{\dot{B}_{p_2,\infty}^\epsilon}.$$

If we raise them to the power $\tau \leq \min(1, p)$ and sum over ℓ with $k > \ell$, we obtain

$$\|I_A\|_p^\tau \lesssim \sum_{k \in \mathbb{Z}} \min(2^{k\alpha\tau} \|f\|_{\dot{B}_{p_1,\infty}^0}^\tau \|g\|_{\dot{B}_{p_2,\infty}^0}^\tau, 2^{-k\epsilon\tau} \|f\|_{\dot{B}_{p_1,\infty}^\alpha}^\tau \|g\|_{\dot{B}_{p_2,\infty}^\epsilon}^\tau).$$

Optimizing over $k \in \mathbb{Z}$, we have as before

$$\|I_A\|_p^\tau \lesssim (\|f\|_{\dot{B}_{p_1,\infty}^0}^\tau \|g\|_{\dot{B}_{p_2,\infty}^0}^\tau)^{\frac{\epsilon}{\alpha+\epsilon}} (\|f\|_{\dot{B}_{p_1,\infty}^\alpha}^\tau \|g\|_{\dot{B}_{p_2,\infty}^\epsilon}^\tau)^{\frac{\alpha}{\alpha+\epsilon}}.$$

Using the interpolation of Besov norms mentioned earlier in (2.11), we deduce

$$\|I_A\|_p^\tau \lesssim \|f\|_{\dot{B}_{p_1,\infty}^{\frac{\alpha}{\alpha+\epsilon}}}^{\frac{\epsilon}{\alpha+\epsilon}\tau} \|g\|_{\dot{B}_{p_2,\infty}^{\frac{\epsilon}{\alpha+\epsilon}}}^{\frac{\epsilon}{\alpha+\epsilon}\tau} \|f\|_{\dot{B}_{p_1,\infty}^0}^{\frac{\epsilon}{\alpha+\epsilon}\tau} \|g\|_{\dot{B}_{p_2,\infty}^0}^{\frac{\alpha}{\alpha+\epsilon}\tau} \lesssim (\|D^\alpha f\|_{L^{p_1}} \|g\|_{L^{p_2}} + \|f\|_{L^{p_1}} \|D^\alpha g\|_{L^{p_2}})^\tau.$$

• estimating I_B :

Due to (2.26), I_B is similar to I_A . We start by noticing that, for $k \ll \ell$ fixed,

$$(2.31) \quad \|[D^\alpha, \Delta_k f] \Delta_\ell g\|_p^\tau \lesssim 2^{(\alpha-1)\ell\tau} 2^{k\tau} \|f\|_{\dot{B}_{p_1,\infty}^0}^\tau \|g\|_{\dot{B}_{p_2,\infty}^0}^\tau$$

and on the other hand, if $0 < \epsilon < \min(\alpha, 1)$,

$$(2.32) \quad \begin{aligned} \|[D^\alpha, \Delta_k f] \Delta_\ell g\|_p^\tau &\lesssim 2^{-\ell\tau} 2^{k(1-\epsilon)\tau} 2^{k\epsilon\tau} \|\Delta_k f\|_{p_1}^\tau 2^{\ell\alpha\tau} \|\Delta_\ell g\|_{p_2}^\tau \\ &\lesssim 2^{-\ell\tau} 2^{k(1-\epsilon)\tau} \|f\|_{\dot{B}_{p_1,\infty}^\epsilon}^\tau \|g\|_{\dot{B}_{p_2,\infty}^\alpha}^\tau. \end{aligned}$$

Combining (2.31) and (2.32) for ℓ fixed, and summing over $k \ll \ell$, we obtain

$$(2.33) \quad \sum_{k:k \ll \ell} \|[D^\alpha, \Delta_k f] \Delta_\ell g\|_p^\tau \lesssim \min(2^{\alpha\ell\tau} \|f\|_{\dot{B}_{p_1,\infty}^0}^\tau \|g\|_{\dot{B}_{p_2,\infty}^0}^\tau, 2^{-\epsilon\ell\tau} \|f\|_{\dot{B}_{p_1,\infty}^\epsilon}^\tau \|g\|_{\dot{B}_{p_2,\infty}^\alpha}^\tau).$$

Now the summation in ℓ is straightforward:

$$\begin{aligned} \|I_B\|_p^\tau &\lesssim \sum_{\ell \in \mathbb{Z}} \min(2^{\alpha\ell\tau} \|f\|_{\dot{B}_{p_1,\infty}^0}^\tau \|g\|_{\dot{B}_{p_2,\infty}^0}^\tau, 2^{-\epsilon\ell\tau} \|f\|_{\dot{B}_{p_1,\infty}^\epsilon}^\tau \|g\|_{\dot{B}_{p_2,\infty}^\alpha}^\tau) \\ &\lesssim (\|f\|_{\dot{B}_{p_1,\infty}^0} \|g\|_{\dot{B}_{p_2,\infty}^0})^{\frac{\tau\epsilon}{\alpha+\epsilon}} (\|f\|_{\dot{B}_{p_1,\infty}^\epsilon} \|g\|_{\dot{B}_{p_2,\infty}^\alpha})^{\frac{\tau\alpha}{\alpha+\epsilon}}. \end{aligned}$$

Using again the interpolation of Besov norms from (2.11), we deduce

$$(2.34) \quad \|I_B\|_p^\tau \lesssim \|f\|_{\dot{B}_{p_1,\infty}^{\frac{\epsilon}{\alpha+\epsilon}}}^{\frac{\epsilon}{\alpha+\epsilon}\tau} \|g\|_{\dot{B}_{p_2,\infty}^{\frac{\alpha}{\alpha+\epsilon}}}^{\frac{\alpha}{\alpha+\epsilon}\tau} \|f\|_{\dot{B}_{p_1,\infty}^0}^{\frac{\epsilon}{\alpha+\epsilon}\tau} \|g\|_{\dot{B}_{p_2,\infty}^0}^{\frac{\alpha}{\alpha+\epsilon}\tau} \lesssim (\|D^\alpha f\|_{L^{p_1}} \|g\|_{L^{p_2}} + \|f\|_{L^{p_1}} \|D^\alpha g\|_{L^{p_2}})^\tau.$$

Finally, due to a trivial application of Hölder's inequality ($\|f \cdot D^\alpha g\|_p \lesssim \|f\|_{p_1} \cdot \|D^\alpha g\|_{p_2}$), the Leibniz rule (2.20) follows. \square

The proof above illustrates the main principle of our paper: in order to estimate multilinear operators associated to a symbol of positive order, it is sufficient to obtain quantitative estimates for the associated Littlewood-Paley pieces. We return now to the proof of (2.25)-(2.28).

Proof of Lemma 2.2. The inequality (2.25) is a direct consequence of Hölder's inequality, and (2.27) can be derived from (2.22) and (2.26) in a straightforward manner. So our main focus will be proving (2.28) and (2.26). For this, we appropriately use Fourier series decompositions for the localized symbols – our approach (see 1.2) for proving Theorem 1.1 will also rely on Fourier series decompositions, but in that setting they will allow us to easily tensorize the flag paraproduct into simpler object – see also Remark 2.3.

(1) Fourier series decomposition for the “diagonal” term

Written in frequency, $D^\alpha(\Delta_\ell f \cdot \Delta_\ell g)(x)$ becomes

$$\begin{aligned} &\int_{\mathbb{R}^2} |\xi_1 + \xi_2|^\alpha \widehat{\Delta_\ell f}(\xi_1) \widehat{\Delta_\ell g}(\xi_2) e^{2\pi i x(\xi_1 + \xi_2)} d\xi_1 d\xi_2 \\ &= \int_{\mathbb{R}^2} |\xi_1 + \xi_2|^\alpha \phi(2^{-\ell}(\xi_1 + \xi_2)) \widehat{\Delta_\ell f}(\xi_1) \widehat{\Delta_\ell g}(\xi_2) e^{2\pi i x(\xi_1 + \xi_2)} d\xi_1 d\xi_2, \end{aligned}$$

where ϕ is a smooth, radial function that is equal to 1 on a neighborhood of $\{|\zeta| \leq 4\}$ and is supported on $\{|\zeta| \leq 8\}$. We proceed with the Fourier series decomposition of $|\xi_1 + \xi_2|^\alpha \phi(2^{-\ell}(\xi_1 + \xi_2))$ on $[-2^{\ell+3}, 2^{\ell+3}]$. If we denote $\zeta = \xi_1 + \xi_2$, we have

$$(2.35) \quad |\zeta|^\alpha \phi(2^{-\ell}\zeta) = \sum_{\tilde{L} \in \mathbb{Z}} C_{\tilde{L}}^\ell e^{2\pi i \frac{\tilde{L}}{2^{\ell+4}} \zeta},$$

where

$$\begin{aligned} C_{\tilde{L}}^\ell &= \tilde{C} \frac{1}{2^{\ell+4}} \int_{[-2^{\ell+3}, 2^{\ell+3}]} |\zeta'|^\alpha \phi(2^{-\ell}\zeta') e^{-2\pi i \frac{\tilde{L}}{2^{\ell+4}} \zeta'} d\zeta' = 2^{\ell\alpha} \tilde{C} \int_{[-8, 8]} |\zeta'|^\alpha \phi(\zeta') e^{\frac{-2\pi i \tilde{L} \zeta'}{2^4}} d\zeta' \\ &:= 2^{\ell\alpha} C_{\tilde{L}}. \end{aligned}$$

Because we are integrating $|\zeta|^\alpha$ close to the origin, the coefficients $C_{\tilde{L}}$ only have limited decay (see [8, Lemma 1]):

$$(2.36) \quad |C_{\tilde{L}}| \lesssim \frac{1}{(1 + |\tilde{L}|)^{1+\alpha}}.$$

This means that we can express $D^\alpha(\Delta_\ell f \cdot \Delta_\ell g)$ as

$$(2.37) \quad D^\alpha(\Delta_\ell f \cdot \Delta_\ell g)(x) = \sum_{\tilde{L} \in \mathbb{Z}} 2^{\ell\alpha} C_{\tilde{L}}(\Delta_{\ell, \frac{\tilde{L}}{2^\ell}} f)(x) (\Delta_{\ell, \frac{\tilde{L}}{2^\ell}} g)(x).$$

And this implies, for $\frac{1}{1+\alpha} < \tau \leq \min(1, p)$,

$$\|D^\alpha(\Delta_\ell f \cdot \Delta_\ell g)\|_p^\tau \lesssim \sum_{\tilde{L}} 2^{\ell\alpha\tau} C_{\tilde{L}}^\tau \|\Delta_{\ell, \frac{\tilde{L}}{2^\ell}} f\|_{p_1}^\tau \|\Delta_{\ell, \frac{\tilde{L}}{2^\ell}} g\|_{p_2}^\tau \lesssim 2^{\ell\alpha\tau} \|\Delta_\ell f\|_{p_1}^\tau \|\Delta_\ell g\|_{p_2}^\tau.$$

(2) Fourier series decomposition for the commutator symbol

Now $k \ll \ell$ are fixed and the commutator $[D^\alpha, \Delta_k f] \Delta_\ell g$ is given by

$$(2.38) \quad \begin{aligned} & \int_{\mathbb{R}^2} (|\xi_1 + \xi_2|^\alpha - |\xi_2|^\alpha) \widehat{\Delta_k f}(\xi_1) \widehat{\Delta_\ell g}(\xi_2) e^{2\pi i x(\xi_1 + \xi_2)} d\xi_1 d\xi_2 \\ &= 2^k \int_{\mathbb{R}^2} \frac{|\xi_1 + \xi_2|^\alpha - |\xi_2|^\alpha}{\xi_1} \tilde{\psi}_k(\xi_1) \widehat{\Delta_k f}(\xi_1) \widehat{\Delta_\ell g}(\xi_2) e^{2\pi i x(\xi_1 + \xi_2)} d\xi_1 d\xi_2, \end{aligned}$$

with $\tilde{\psi}_k(\xi_1) = \frac{\xi_1}{2^k} \tilde{\psi}_k(\xi_1)$, and $\tilde{\psi}_k(\xi_1) \equiv 1$ on the support of $\psi_k(\xi_1)$.

We let

$$(2.39) \quad m_{C_\alpha}(\xi_2, \xi_1) := \frac{|\xi_1 + \xi_2|^\alpha - |\xi_2|^\alpha}{\xi_1} = \alpha \int_0^1 |\xi_2 + t\xi_1|^{\alpha-2} (\xi_2 + t\xi_1) dt$$

denote²¹ the symbol measuring the average contribution of the commutator, localized to the cone $R_2 := \{(\xi_1, \xi_2) : |\xi_1| \ll |\xi_2|\}$. Since $k \ll \ell$, we can further (smoothly) localize it to the region $\{|\xi_1| \ll 2^\ell, |\xi_2| \sim 2^\ell\}$:

$$(2.40) \quad m_{C_\alpha}^\ell(\xi_2, \xi_1) = m_{C_\alpha}(\xi_2, \xi_1) \tilde{\varphi}_\ell(\xi_1) \tilde{\psi}_\ell(\xi_2).$$

In view of the fact that we want to use a Fourier series decomposition in both the variables ξ_1 and ξ_2 , $m_{C_\alpha}^\ell$ will be split as

$$(2.41) \quad m_{C_\alpha}^\ell(\xi_2, \xi_1) = m_{C_\alpha}^{\ell,+}(\xi_2, \xi_1) + m_{C_\alpha}^{\ell,-}(\xi_2, \xi_1),$$

where

$$(2.42) \quad m_{C_\alpha}^{\ell,+}(\xi_2, \xi_1) := m_{C_\alpha}(\xi_2, \xi_1) \tilde{\varphi}_\ell(\xi_1) \tilde{\psi}_\ell^+(\xi_2), \quad m_{C_\alpha}^{\ell,-}(\xi_2, \xi_1) := m_{C_\alpha}(\xi_2, \xi_1) \tilde{\varphi}_\ell(\xi_1) \tilde{\psi}_\ell^-(\xi_2).$$

Here $\tilde{\psi}_\ell^+$ is a bump function compactly supported on $[2^{\ell-2}, 2^{\ell+2}]$, $\tilde{\psi}_\ell^-$ a bump function compactly supported on $[-2^{\ell+2}, -2^{\ell-2}]$, and $\tilde{\varphi}_\ell$ on $[-2^{\ell-3}, 2^{\ell-3}]$. Overall we have

$$(2.43) \quad \begin{aligned} [D^\alpha, \Delta_k f] \Delta_\ell g(x) &= 2^k \int_{\mathbb{R}^2} m_{C_\alpha}^{\ell,+}(\xi_2, \xi_1) \tilde{\psi}_k(\xi_1) \widehat{\Delta_k f}(\xi_1) \widehat{\Delta_{\ell,+} g}(\xi_2) e^{2\pi i x(\xi_1 + \xi_2)} d\xi_1 d\xi_2 \\ &\quad + 2^k \int_{\mathbb{R}^2} m_{C_\alpha}^{\ell,-}(\xi_2, \xi_1) \tilde{\psi}_k(\xi_1) \widehat{\Delta_k f}(\xi_1) \widehat{\Delta_{\ell,-} g}(\xi_2) e^{2\pi i x(\xi_1 + \xi_2)} d\xi_1 d\xi_2. \end{aligned}$$

We can now perform a Fourier series decomposition of $m_{C_\alpha}^{\ell,+}(\xi_2, \xi_1)$ on $[2^{\ell-2}, 2^{\ell+2}] \times [-2^{\ell-3}, 2^{\ell-3}]$. Notice that this is a double Fourier series expansion, involving both variables ξ_1 and ξ_2 . Indeed, we obtain

$$m_{C_\alpha}^{\ell,+}(\xi_2, \xi_1) = \sum_{L_1, L_2 \in \mathbb{Z}} C_{L_1, L_2}^{\ell,+} e^{2\pi i \frac{L_1}{2^{\ell-2}} \xi_1} e^{2\pi i \frac{L_2}{2^{\ell+2}} \xi_2},$$

where the Fourier coefficients are described by

$$\begin{aligned} C_{L_1, L_2}^{\ell,+} &= \tilde{C} \frac{1}{2^\ell 2^\ell} \int_{[-2^{\ell-3}, 2^{\ell-3}] \times [2^{\ell-2}, 2^{\ell+2}]} \int_0^1 (\xi_2' + t\xi_1')^{\alpha-1} dt e^{-2\pi i \frac{L_1}{2^{\ell-2}} \xi_1'} e^{-2\pi i \frac{L_2}{2^{\ell+2}} \xi_2'} d\xi_1' d\xi_2' \\ &= 2^{\ell(\alpha-1)} \tilde{C} \int_{[-\frac{1}{8}, \frac{1}{8}] \times [\frac{1}{4}, 4]} \int_0^1 (\xi_2' + t\xi_1')^{\alpha-1} dt e^{-2\pi i 4L_1 \xi_1'} e^{-2\pi i \frac{L_2}{4} \xi_2'} d\xi_1' d\xi_2' \\ &:= 2^{\ell(\alpha-1)} C_{L_1, L_2}^+. \end{aligned}$$

²¹The symbol $m_{C_\alpha}(\xi_2, \xi_1)$, associated to the region $\{|\xi_1| \ll |\xi_2|\}$, is central in the commutator's analysis. On the contrary, if we restrict our attention to the cone $\{|\xi_2| \ll |\xi_1|\}$, we need to study the behavior of $m_{C_\alpha}(\xi_1, \xi_2)$ defined by $\frac{|\xi_1 + \xi_2|^\alpha - |\xi_1|^\alpha}{\xi_2}$.

Similarly, a double Fourier series decomposition can be applied to $m_{C_\alpha}^{\ell,-}(\xi_2, \xi_1)$ on $[-2^{\ell+2}, -2^{\ell-2}] \times [-2^{\ell-3}, 2^{\ell-3}]$ so that

$$m_{C_\alpha}^{\ell,-}(\xi_2, \xi_1) = \sum_{L_1, L_2 \in \mathbb{Z}} C_{L_1, L_2}^{\ell,-} e^{2\pi i \frac{L_1}{2^{\ell-2}} \xi_1} e^{2\pi i \frac{L_2}{2^{\ell+2}} \xi_2},$$

where the Fourier coefficients can be expressed as

$$C_{L_1, L_2}^{\ell,-} = 2^{\ell(\alpha-1)} \tilde{C} \int_{[-\frac{1}{8}, \frac{1}{8}] \times [-4, -\frac{1}{4}]} \int_0^1 (\xi_2' + t\xi_1')^{\alpha-1} dt e^{-2\pi i 4L_1 \xi_1'} e^{-2\pi i \frac{L_2}{4} \xi_2'} d\xi_1' d\xi_2' := 2^{\ell(\alpha-1)} C_{L_1, L_2}^-.$$

A straightforward, but important observation is the fact that the new coefficients C_{L_1, L_2}^+ and C_{L_1, L_2}^- do not depend on the parameter ℓ ; moreover, due to the fact that $\int_0^1 (\xi_2' + t\xi_1')^{\alpha-1} dt$ is smooth for $(\xi_2', \xi_1') \in [\frac{1}{4}, 4] \times [-\frac{1}{8}, \frac{1}{8}] \cup [-4, -\frac{1}{4}] \times [-\frac{1}{8}, \frac{1}{8}]$, we also deduce their fast decay:

$$(2.44) \quad |C_{L_1, L_2}^+|, |C_{L_1, L_2}^-| \lesssim_M \frac{1}{(1 + |L_1| + |L_2|)^M},$$

for any $M > 0$.

Recalling (2.43), $[D^\alpha, \Delta_k f] \Delta_\ell g$ becomes

$$(2.45) \quad \begin{aligned} & \sum_{L_1, L_2} C_{L_1, L_2}^+ 2^{\ell(\alpha-1)} 2^k (\tilde{\Delta}_{k, \frac{L_1}{2^{\ell-3}}} \Delta_k f)(x) (\tilde{\Delta}_{\ell, +, \frac{L_2}{2^{\ell+2}}} \Delta_\ell g)(x) \\ & + \sum_{L_1, L_2} C_{L_1, L_2}^- 2^{\ell(\alpha-1)} 2^k (\tilde{\Delta}_{k, \frac{L_1}{2^{\ell-3}}} \Delta_k f)(x) (\tilde{\Delta}_{\ell, -, \frac{L_2}{2^{\ell+2}}} \Delta_\ell g)(x) = I_+ + I_-. \end{aligned}$$

In other words, the commutator $[D^\alpha, \Delta_k f] \Delta_\ell g$ emerges as a superposition of products of modulated Littlewood-Paley projections, times $2^{\ell(\alpha-1)} 2^k$.

So if $\tau \leq \min(1, p)$,

$$(2.46) \quad \begin{aligned} \|I_\pm\|_p^\tau & \lesssim \sum_{L_1, L_2} |C_{L_1, L_2}^\pm|^\tau 2^{\ell(\alpha-1)\tau} 2^{k\tau} \|\tilde{\Delta}_{k, \frac{L_1}{2^{\ell-3}}} \Delta_k f\|_{p_1}^\tau \|\tilde{\Delta}_{\ell, \pm, \frac{L_2}{2^{\ell+2}}} \Delta_\ell g\|_{p_2}^\tau \\ & \lesssim \sum_{L_1, L_2} |C_{L_1, L_2}^\pm|^\tau 2^{\ell(\alpha-1)\tau} 2^{k\tau} \|\Delta_k f\|_{p_1}^\tau \|\Delta_\ell g\|_{p_2}^\tau. \end{aligned}$$

The fast decay of the C_{L_1, L_2}^\pm coefficients from (2.44) and the estimate (2.16) imply (2.26).

Following the same ideas, one can provide a direct proof for (2.27) without invoking the commutator estimate (2.26). For example, one can perform a double Fourier series decomposition for the smooth function

$$|\xi_1 + \xi_2|^\alpha \tilde{\varphi}_\ell(\xi_1) \tilde{\psi}_{\ell, \pm}(\xi_2)$$

on $[-2^{\ell-3}, 2^{\ell-3}] \times [2^{\ell-2}, 2^{\ell+2}]$ and $[-2^{\ell-3}, 2^{\ell-3}] \times [-2^{\ell+2}, -2^{\ell-2}]$ respectively. □

Remark 2.3. We would like to draw attention to a certain component in the above argument: the Fourier series decomposition (2.45) (and its consequence (2.46)) will be as important as the localized estimates (2.26) in the treatment of generic flags. More concretely, $\|\Delta_k f\|_{p_1}$ can be replaced by $\|D^\alpha(\Delta_k f_1 \cdot S_k f_2)\|_t$ with $t > 0$ (possibly $t < 1$) in the study of a flag $D^\beta(D^\alpha(\Delta_k f_1 \cdot S_k f_2)\Delta_\ell g)$. The Fourier series decomposition reduces the original estimate to superpositions of subtree estimates, thus tensorizing a generic flag into flags of lower complexity. However, quantitative estimates in the spirit of (2.26) for $\|D^\alpha(\Delta_k f_1 \cdot S_k f_2)\|_t$ will be needed in order to conclude by summing up the scales k and ℓ , as before.

Remark 2.4. (1) In higher dimensions, when $\xi_1, \xi_2 \in \mathbb{R}^d$, we cannot use (2.39) anymore; instead, we need to directly handle

$$(2.47) \quad |\xi_1 + \xi_2|^\alpha - |\xi_2|^\alpha = \int_0^1 \xi_1 \cdot (t\xi_1 + \xi_2) |t\xi_1 + \xi_2|^{\alpha-2} dt.$$

in the off-diagonal region $\{(\xi_1, \xi_2) \in \mathbb{R}^{2d} : |\xi_2| \gg |\xi_1|\}$. So when we want to estimate

$$(2.48) \quad \int_{\mathbb{R}^{2d}} (|\xi_1 + \xi_2|^\alpha - |\xi_2|^\alpha) \widehat{\Delta_k f}(\xi_1) \widehat{\Delta_\ell g}(\xi_2) e^{2\pi i x \cdot (\xi_1 + \xi_2)} d\xi_1 d\xi_2,$$

under the assumption that $|\xi_1| \sim 2^k \ll 2^\ell \sim |\xi_2|$, and perform a double Fourier series decomposition for

$$(|\xi_1 + \xi_2|^\alpha - |\xi_2|^\alpha) \tilde{\psi}_k(\xi_1) \tilde{\psi}_\ell(\xi_2),$$

we need to further localize the symbol above onto “Whitney boxes/rectangles” of sizes $\sim 2^k \times 2^\ell$.

This requires a technical (although standard) decomposition of the region $\{\xi \in \mathbb{R}^d : \xi \neq 0\}$ as a finite²² collection \mathfrak{C} of directional cones; moreover, to each $\mathfrak{c} \in \mathfrak{C}$ we associate a collection of Whitney cubes $\{Q_{k,\mathfrak{c}}\}_{k \in \mathbb{Z}}$ covering the conical regions and having the property that

$$\text{sidelength}(Q_{k,\mathfrak{c}}) \sim 2^k \sim \text{dist}(Q_{k,\mathfrak{c}}, 0).$$

In this way, for any $k \in \mathbb{Z}$, we have

$$\psi_k(\xi) = \sum_{\mathfrak{c} \in \mathfrak{C}} \tilde{\psi}_{k,\mathfrak{c}}(\xi) \psi_k(\xi),$$

which should bring back to mind the decomposition performed in (2.40)-(2.42).

So the symbol $|\xi_1 + \xi_2|^\alpha - |\xi_2|^\alpha$, initially localized through Littlewood-Paley projections $\tilde{\psi}_k(\xi_1)$ and $\tilde{\psi}_\ell(\xi_2)$, now becomes

$$(|\xi_1 + \xi_2|^\alpha - |\xi_2|^\alpha) \tilde{\psi}_k(\xi_1) \tilde{\psi}_\ell(\xi_2) = \sum_{\mathfrak{c}_1, \mathfrak{c}_2 \in \mathfrak{C}} (|\xi_1 + \xi_2|^\alpha - |\xi_2|^\alpha) \tilde{\psi}_{k,\mathfrak{c}_1}(\xi_1) \tilde{\psi}_{\ell,\mathfrak{c}_2}(\xi_2) \tilde{\psi}_k(\xi_1) \tilde{\psi}_\ell(\xi_2).$$

We then use a double Fourier series decomposition on each Whitney rectangle $Q_{k,\mathfrak{c}_1} \times Q_{\ell,\mathfrak{c}_2}$ in order to capture the interaction between two different scales. We have

$$(|\xi_1 + \xi_2|^\alpha - |\xi_2|^\alpha) \tilde{\psi}_{k,\mathfrak{c}_1}(\xi_1) \tilde{\psi}_{\ell,\mathfrak{c}_2}(\xi_2) = 2^{\ell(\alpha-1)} 2^k \sum_{L_1, L_2 \in \mathbb{Z}^d} C_{L_1, L_2}^{k, \ell, \mathfrak{c}_1, \mathfrak{c}_2} e^{2\pi i \frac{L_1}{2^k} \xi_1} e^{2\pi i \frac{L_2}{2^\ell} \xi_2},$$

where

$$(2.49) \quad C_{L_1, L_2}^{k, \ell, \mathfrak{c}_1, \mathfrak{c}_2} = \int_{\{|\xi'_1| \sim 1\} \cap c_1} \int_{\{|\xi'_2| \sim 1\} \cap c_2} \tilde{\psi}_{0,\mathfrak{c}_1}(\xi'_1) \tilde{\psi}_{0,\mathfrak{c}_2}(\xi'_2) \left(\int_0^1 \xi'_1 \cdot (2^{k-\ell} t \xi'_1 + \xi'_2) |2^{k-\ell} t \xi'_1 + \xi'_2|^{\alpha-2} dt \right) e^{-2\pi i L_1 \cdot \xi'_1} e^{-2\pi i L_2 \cdot \xi'_2} d\xi'_1 d\xi'_2.$$

In contrast to the one-dimensional case, the renormalized Fourier coefficients do depend on the scales 2^k and 2^ℓ . However, since $2^{k-\ell} \ll 1$ and the multiplier

$$(2.50) \quad \tilde{\psi}_{0,\mathfrak{c}_1}(\xi'_1) \tilde{\psi}_{0,\mathfrak{c}_2}(\xi'_2) \int_0^1 \xi'_1 \cdot (2^{k-\ell} t \xi'_1 + \xi'_2) |2^{k-\ell} t \xi'_1 + \xi'_2|^{\alpha-2} dt$$

is smooth on its support included in $\{(\xi'_1, \xi'_2) \in \mathbb{R}^{2d} : |\xi'_1| \sim 1, |\xi'_2| \sim 1\} \cap c_1 \times c_2$, we do have, *uniformly in k and ℓ* ,

$$(2.51) \quad |C_{L_1, L_2}^{k, \ell, \mathfrak{c}_1, \mathfrak{c}_2}| \lesssim_M \frac{1}{(1 + |L_1| + |L_2|)^M},$$

for any $M > 0$. Indeed, this is a consequence of the uniform boundedness of the derivatives of the multiplier in (2.50). Since the above estimate holds for any tensor product of directional cones $\mathfrak{c}_1 \times \mathfrak{c}_2$ with $\mathfrak{c}_1, \mathfrak{c}_2 \in \mathfrak{C}$ and there are only finitely many directional cones in \mathfrak{C} , we obtain the desired estimate for (2.48).

(2) In the diagonal region $\{(\xi_1, \xi_2) \in \mathbb{R}^{2d} : |\xi_1| \sim |\xi_2|\}$, $|\xi_1 + \xi_2|^\alpha$ is not supported away from the origin, we perform a Fourier series decomposition of $|\xi_1 + \xi_2|^\alpha \phi_\ell(\xi_1 + \xi_2)$ on the cube $[-2^{\ell+3}, 2^{\ell+3}]^d \subseteq \mathbb{R}^d$ and notice that the corresponding Fourier coefficients only have limited decay:

$$(2.52) \quad |C_L| \lesssim \frac{2^{\ell\alpha}}{(1 + |L|)^{\alpha+d}}.$$

The way the Fourier series are being performed is the only point in the proof which is different in higher dimensions. More precisely, the Fourier series decomposition requires a smooth localization of the symbol to Whitney rectangles within the region $\{(\xi_1, \xi_2) : |\xi_1| \sim 2^k, |\xi_2| \sim 2^\ell\}$, and the renormalized Fourier coefficients do depend on the scales k and ℓ , although they are still well-behaved thanks to (2.51). In order to avoid unnecessary complications, in what follows we will focus on the one dimensional case.

2.3. Some contrasting aspects with the Coifman-Meyer multiplier approach.

Alternatively, one might want to use the boundedness of Coifman-Meyer multipliers to deduce the Leibniz rule (2.20). The Littlewood-Paley decompositions $f = \sum_{k \in \mathbb{Z}} \Delta_k f$, $g = \sum_{\ell \in \mathbb{Z}} \Delta_\ell g$ will again be grouped into “paraproducts”, with possibly altered projections:

$$\begin{aligned} f(x)g(x) &= \sum_{k \ll \ell} \Delta_k f(x) \Delta_\ell g(x) + \sum_{\ell \ll k} \Delta_k f(x) \Delta_\ell g(x) + \sum_{|k-\ell| \leq 3} \Delta_k f(x) \Delta_\ell g(x) \\ &= \sum_{\ell \in \mathbb{Z}} S_\ell f(x) \Delta_\ell g(x) + \sum_{k \in \mathbb{Z}} \Delta_k f(x) S_k g(x) + \sum_{k \in \mathbb{Z}} \Delta_k f(x) \tilde{\Delta}_k g(x). \end{aligned}$$

²²The numbers of cones depends on the dimension.

Then $D^\alpha(f \cdot g)$ becomes

$$D^\alpha\left(\sum_{\ell \in \mathbb{Z}} S_\ell f \cdot \Delta_\ell g\right) + D^\alpha\left(\sum_{k \in \mathbb{Z}} \Delta_k f \cdot S_k g\right) + D^\alpha\left(\sum_{k \in \mathbb{Z}} \Delta_k f \cdot \tilde{\Delta}_k g\right);$$

the first two terms being symmetric, it suffices to understand the first and the third one. Now we notice the following:

(1) the first term can be expressed as

$$D^\alpha\left(\sum_{\ell \in \mathbb{Z}} S_\ell f \cdot \Delta_\ell g\right) = \int_{\mathbb{R}^2} \sum_{\ell \in \mathbb{Z}} \varphi(2^\ell \xi) \psi(2^\ell \eta) \frac{|\xi + \eta|^\alpha}{|\eta|^\alpha} \hat{f}(\xi) \widehat{D^\alpha g}(\eta) e^{2\pi i x(\xi + \eta)} d\xi d\eta;$$

where

$$m_1(\xi, \eta) := \sum_{\ell \in \mathbb{Z}} \varphi(2^\ell \xi) \psi(2^\ell \eta) \frac{|\xi + \eta|^\alpha}{|\eta|^\alpha}$$

is a Coifman-Meyer symbol, satisfying (1.6); this is because m_1 is morally constant on Whitney cubes $[-2^{\ell-2}, 2^{\ell-2}] \times [2^\ell, 2^{\ell+1}]$. As a consequence,

$$D^\alpha\left(\sum_{\ell \in \mathbb{Z}} S_\ell f \cdot \Delta_\ell g\right)(x) = T_{m_1}(f, D^\alpha g)(x),$$

where T_{m_1} is the Coifman-Meyer multiplier associated to the Mikhlin symbol $m_1(\xi, \eta)$.

(2) In a similar way, thanks to (2.35), $D^\alpha\left(\sum_{k \in \mathbb{Z}} \Delta_k f \cdot \tilde{\Delta}_k g\right)$ can be represented as a superposition of Coifman-Meyer-type multipliers

$$D^\alpha\left(\sum_{k \in \mathbb{Z}} \Delta_k f \cdot \tilde{\Delta}_k g\right) = \sum_{\tilde{L} \in \mathbb{Z}} C_{\tilde{L}} T_{m_{\tilde{L}}}(f, D^\alpha g),$$

where

$$m_{\tilde{L}}(\xi, \eta) := \sum_{\ell \in \mathbb{Z}} \psi(2^{-\ell} \xi) e^{2\pi i \frac{\tilde{L}}{2^\ell} \xi} \tilde{\psi}(2^{-\ell} \eta) e^{2\pi i \frac{\tilde{L}}{2^\ell} \eta} 2^{\ell\alpha} |\eta|^{-\alpha}$$

is a symbol singular only at the origin, but depending explicitly on \tilde{L} . Since $\|T_{m_{\tilde{L}}}\|_{L^{p_1} \times L^{p_2} \rightarrow L^p}$ can be shown to depend at most logarithmically²³ on $1 + |\tilde{L}|$ whenever $1/p_1 + 1/p_2 = 1/p$, with $1 < p_1, p_2 \leq \infty, 1/2 < p < \infty$, (2.20) follows for $1 < p_1, p_2 \leq \infty, 1/2 < p < \infty$ such that

$$\sum_{\tilde{L}} |C_{\tilde{L}}|^{\min(1, p)}$$

is summable, i.e. for $\frac{1}{1+\alpha} < p < \infty$.

We conclude with a brief comparison between the Coifman-Meyer multiplier approach and the Bourgain-Li approach presented above in Section 2.2.

- (i) the Bourgain-Li approach can deal with endpoints for which Coifman-Meyer multipliers fail to be bounded; this was the original framework in which it was introduced – the $L^\infty \times L^\infty \rightarrow L^\infty$ endpoint;
- (ii) in the Bourgain-Li approach, which involves summation over the scales, the derivatives are allocated jointly to the functions f and g : for small scales, both f and g get no derivatives; in contrast, for large scales f is being attributed α derivatives and g picks up ϵ derivatives for some $\epsilon \in (0, \alpha)$. Interpolation eventually yields, for some $0 < \theta < 1$,

$$(2.53) \quad \|D^\alpha(fg) - D^\alpha f \cdot g - f \cdot D^\alpha g\|_{L^p} \lesssim \left(\|f\|_{\dot{B}_{p_1, \infty}^\alpha} \|g\|_{\dot{B}_{p_2, \infty}^0}\right)^\theta \left(\|f\|_{\dot{B}_{p_1, \infty}^0} \|g\|_{\dot{B}_{p_2, \infty}^\alpha}\right)^{1-\theta},$$

which is a sharper estimate than (2.20);

- (iii) in the Coifman-Meyer multiplier approach, the derivative will always be attached to the function with higher oscillation, as one can see from the identity $D^\alpha\left(\sum_{\ell \in \mathbb{Z}} S_\ell f \cdot \Delta_\ell g\right)(x) = T_{m_1}(f, D^\alpha g)(x)$; as a consequence, one can obtain “off-diagonal” L^p estimates for the Leibniz rule, in the sense that (2.20) can be replaced by

$$(2.54) \quad \|D^\alpha(fg)\|_{L^p} \lesssim \|D^\alpha f\|_{L^{p_1}} \|g\|_{L^{p_2}} + \|f\|_{L^{q_1}} \|D^\alpha g\|_{L^{q_2}},$$

where $\frac{1}{p} = \frac{1}{p_1} + \frac{1}{p_2} = \frac{1}{q_1} + \frac{1}{q_2}$, and $1 < p_1, p_2, q_1, q_2 \leq \infty, \frac{1}{2} < p < \infty$. Interestingly, such off-diagonal estimates proved to be useful in certain applications – for example [11], [12].

²³Indeed, the frequency modulation $e^{2\pi i \tilde{L} \cdot}$ has an \tilde{L} -shifting effect in space (at every scale) and invoking the boundedness of shifted square functions produces the desired result; see [8] or [16].

(iv) the Coifman-Meyer approach yields another type of improvement for the “commutator” in (2.53):

$$\|D^\alpha(fg) - D^\alpha f \cdot g - f \cdot D^\alpha g\|_{L^p} \lesssim \|D^{\alpha_1} f\|_{L^{p_1}} \|D^{\alpha_2} g\|_{L^{p_2}},$$

for any $0 < \alpha_1, \alpha_2 < \alpha$ with $\alpha_1 + \alpha_2 = \alpha$. Because of (ii) and the role played by interpolation, without substantial modifications, the Bourgain-Li approach cannot recover such a result.

Our methodology for proving estimates for generic flag Leibniz rules and other Leibniz-type estimates can be perceived as a generalization of the Bourgain-Li approach presented above, as hinted in Remark 2.3.

3. A FIVE-LINEAR FLAG: ONE-PARAMETER CASE

Before dealing with the generic result, we consider a simpler example: $D^\beta(D^\alpha(f_1 \cdot f_2) \cdot f_3 \cdot D^\gamma(f_4 \cdot f_5))$, which is a one-parameter version of (1.1). This will allow us to emphasize the main ideas without getting too technical. We will assume that $d = 1$, although the argument remains identical in higher dimensions.

We will prove, for $\alpha, \beta, \gamma \geq 0$ that

$$(3.1) \quad \begin{aligned} & \|D^\beta(D^\alpha(f_1 f_2) f_3 D^\gamma(f_4 f_5))\|_{L^r} \\ & \lesssim \|D^{\alpha+\beta} f_1\|_{L^{p_1}} \|f_2\|_{L^{p_2}} \|f_3\|_{L^{p_3}} \|D^\gamma f_4\|_{L^{p_4}} \|f_5\|_{L^{p_5}} + \|f_1\|_{L^{p_1}} \|D^{\alpha+\beta} f_2\|_{L^{p_2}} \|f_3\|_{L^{p_3}} \|D^\gamma f_4\|_{L^{p_4}} \|f_5\|_{L^{p_5}} \\ & + \|D^{\alpha+\beta} f_1\|_{L^{p_1}} \|f_2\|_{L^{p_2}} \|f_3\|_{L^{p_3}} \|f_4\|_{L^{p_4}} \|D^\gamma f_5\|_{L^{p_5}} + \text{other similar terms,} \end{aligned}$$

where $1 \leq p_1, \dots, p_5 \leq \infty$, $1/5 \leq r \leq \infty$ and

$$(3.2) \quad \frac{1}{r} = \frac{1}{p_1} + \dots + \frac{1}{p_5}, \quad \frac{1}{r} < 1 + \beta, \quad \frac{1}{p_{1,2}} := \frac{1}{p_1} + \frac{1}{p_2} < 1 + \alpha, \quad \frac{1}{p_{4,5}} := \frac{1}{p_4} + \frac{1}{p_5} < 1 + \gamma.$$

Whenever α, β or $\gamma \in 2\mathbb{Z}$, the constraint (3.2) can be removed.


Our approach for the above Leibniz rule relies on an iterative argument that in particular makes use of frequency-localized estimates for $D^\alpha(f_1 f_2)$ and $D^\gamma(f_4 f_5)$, in the spirit of Lemma 2.2. The present proof is not as systematic as the one in Section 5, although many elements are contained in the treatment of this particular example.

We use Littlewood-Paley projections to decompose the functions in frequency into dyadic pieces

$$f_l = \sum_{k_l} \Delta_{k_l} f_l, \quad \text{for all } 1 \leq l \leq 5,$$

for which the derivation becomes equivalent to multiplication: $D^\alpha(\Delta_{k_l} f_l) \sim 2^{\alpha k_l} \Delta_{k_l} f_l$. That means that we need to estimate

$$(3.3) \quad \sum_{k_1, \dots, k_5} \int_{\mathbb{R}^5} |\xi_1 + \dots + \xi_5|^\beta \cdot |\xi_1 + \xi_2|^\alpha \cdot |\xi_4 + \xi_5|^\gamma \widehat{\Delta_{k_1} f_1}(\xi_1) \cdot \dots \cdot \widehat{\Delta_{k_5} f_5}(\xi_5) e^{2\pi i x(\xi_1 + \dots + \xi_5)} d\xi_1 \dots d\xi_5.$$



As explained in Sections 1.2 and 2, the frequency space will be split in various conical regions, producing in this way the classical paraproduct decomposition. Restrictions of (3.3) to each of these regions need to be analyzed, and we will see that the leading derivatives will be distributed among two functions: the highest oscillating functions and an auxiliary one.

Due to the structure of the present flag, we will need to consider several conical regions in frequency:

(I) the region where f_1 is the fastest oscillating function:

$$(3.4) \quad R_1 = \{(\xi_1, \dots, \xi_5) : |\xi_1| \gg |\xi_2|, \dots, |\xi_5|\}.$$

This region is symmetric to those where f_2, f_4 or f_5 oscillate much faster than the remaining functions.

(II) the region

$$(3.5) \quad R_3 = \{(\xi_1, \dots, \xi_5) : |\xi_3| \gg |\xi_1|, |\xi_2|, |\xi_4|, |\xi_5|\},$$

where f_3 is the fastest oscillating function.

(III) “diagonal” regions

$$(3.6) \quad \tilde{R}_{l_1, l_2} = \{(\xi_1, \dots, \xi_5) : |\xi_{l_1}| \sim |\xi_{l_2}| \geq |\xi_{l'}| \text{ for } l' \neq l_1, l_2\},$$

for $l_1 \neq l_2$. In this situation, at least two of the functions oscillate at comparable high rates, which might cause $|\xi_1 + \dots + \xi_5|^\beta$ to become more singular than in the previous cases.

3.1. (I): study of the conical region R_1 .

In (3.3), we restrict the summation over $k_1 \ll k_2, \dots, k_5$: we are in the situation when $2^{k_1} \sim |\xi_1| \ll |\xi_l| \sim 2^{k_l}$ for all $2 \leq l \leq 5$. In this case,

$$2^{k_1-1} < |\xi_1 + \dots + \xi_5| < 2^{k_1+1},$$

which we write in short $|\xi_1 + \dots + \xi_5| \sim 2^{k_1}$.

Following the principle introduced by Bourgain and Li, in this scenario we would like to approximate $|\xi_1 + \dots + \xi_5|^\beta$ by $|\xi_1|^\beta$, and use to good advantage the better-behaving commutator $|\xi_1 + \dots + \xi_5|^\beta - |\xi_1|^\beta$. Since a D^α derivative also acts on f_1 , we take an intermediate step in which we approximate $|\xi_1 + \dots + \xi_5|^\beta$ by $|\xi_1 + \xi_2|^\beta$. This is consistent with the overall approach, since $|\xi_1 + \xi_2| \sim 2^{k_1}$ as well. Thus the symbol corresponding to this particular Leibniz rule, which appears in (3.3), breaks down as

$$\begin{aligned} & \frac{|\xi_1 + \dots + \xi_5|^\beta - |\xi_1 + \xi_2|^\beta}{\xi_3 + \xi_4 + \xi_5} \cdot \xi_3 \cdot |\xi_1 + \xi_2|^\alpha \cdot |\xi_4 + \xi_5|^\gamma + \frac{|\xi_1 + \dots + \xi_5|^\beta - |\xi_1 + \xi_2|^\beta}{\xi_3 + \xi_4 + \xi_5} \cdot (\xi_4 + \xi_5) \cdot |\xi_1 + \xi_2|^\alpha \cdot |\xi_4 + \xi_5|^\gamma \\ & + \frac{|\xi_1 + \xi_2|^{\alpha+\beta} - |\xi_1|^{\alpha+\beta}}{\xi_2} \cdot \xi_2 \cdot |\xi_4 + \xi_5|^\gamma + |\xi_1|^{\alpha+\beta} \cdot |\xi_4 + \xi_5|^\gamma := m_{I_A} + m_{I_B} + m_{I_C} + m_{I_D}, \end{aligned}$$

which means that (3.3) restricted to the frequency region R_1 (3.4) reads as

$$\begin{aligned} & \sum_{k_2, \dots, k_5 \ll k_1} T_{m_{I_A}}(\Delta_{k_1} f_1, \dots, \Delta_{k_5} f_5)(x) + \sum_{k_2, \dots, k_5 \ll k_1} T_{m_{I_B}}(\Delta_{k_1} f_1, \dots, \Delta_{k_5} f_5)(x) \\ & + \sum_{k_2, \dots, k_5 \ll k_1} T_{m_{I_C}}(\Delta_{k_1} f_1, \dots, \Delta_{k_5} f_5)(x) + \sum_{k_2, \dots, k_5 \ll k_1} T_{m_{I_D}}(\Delta_{k_1} f_1, \dots, \Delta_{k_5} f_5)(x) := I_A + I_B + I_C + I_D. \end{aligned}$$

This corresponds to the first step of the strategy presented in the introduction: the *splitting of the root symbol*. In Section 5, the splitting will be different: the emphasis will be put on the subtree structures obtained by removing the tree root corresponding to D^β . Here instead we track the root derivatives D^β as they descend towards the leaves represented by the functions f_1, \dots, f_5 .

We will see that, due to the multiplier's shape, in the study of I_A the functions f_1 and f_3 will play a special role and the D^β derivatives will be shared among them; for I_B , it will be f_1 and one of f_4 or f_5 (an extra paraproduct decomposition will be used here, which will also determine the distribution of derivatives); for I_C , f_1 and f_2 ; and finally, for I_D , summing up all the scales will require a change in the order of summation.

I_A) In estimating the I_A term, we will only need the Littlewood-Paley information for the functions f_1 and f_3 ; as a consequence, we can sum over $k_2, k_4, k_5 \ll k_1$ and focus on the five-linear operator

$$\begin{aligned} T_{m_{I_A}}^{k_1, k_3}(f_1, \dots, f_5)(x) &:= \int_{\mathbb{R}^5} \frac{|\xi_1 + \dots + \xi_5|^\beta - |\xi_1 + \xi_2|^\beta}{\xi_3 + \xi_4 + \xi_5} \cdot \xi_3 \cdot |\xi_1 + \xi_2|^\alpha \cdot |\xi_4 + \xi_5|^\gamma \\ &\quad \widehat{\Delta_{k_1} f_1}(\xi_1) \cdot \widehat{S_{k_1} f_2}(\xi_2) \cdot \widehat{\Delta_{k_3} f_3}(\xi_3) \cdot \widehat{S_{k_1} f_4}(\xi_4) \cdot \widehat{S_{k_1} f_5}(\xi_5) e^{2\pi i x(\xi_1 + \dots + \xi_5)} d\xi_1 \dots d\xi_5. \end{aligned}$$

Next, we use *Fourier series decompositions for the symbols* involved – the second step of our strategy. For this we need to place ourselves in a suitable situation, and in particular to smoothly restrict the symbols to intervals where the Fourier series decomposition can be implemented.

First we look at $m_{C_\beta}(\xi_1 + \xi_2, \xi_3 + \xi_4 + \xi_5)$ which, according to definition (2.39), consists of

$$\frac{|\xi_1 + \dots + \xi_5|^\beta - |\xi_1 + \xi_2|^\beta}{\xi_3 + \xi_4 + \xi_5} = \beta \int_0^1 |\xi_1 + \xi_2 + t(\xi_3 + \xi_4 + \xi_5)|^{\beta-2} (\xi_1 + \xi_2 + t(\xi_3 + \xi_4 + \xi_5)) dt,$$

and further restrict it to $[2^{k_1}, 2^{k_1+1}] \times [-2^{k_1-1}, 2^{k_1-1}]$ and $[-2^{k_1+1}, -2^{k_1}] \times [-2^{k_1-1}, 2^{k_1-1}]$, respectively:

$$m_{C_\beta}^{k_1, \pm}(\xi_1 + \xi_2, \xi_3 + \xi_4 + \xi_5) := m_{C_\beta}(\xi_1 + \xi_2, \xi_3 + \xi_4 + \xi_5) \tilde{\psi}_{k_1, \pm}(\xi_1 + \xi_2) \tilde{\varphi}_{k_1}(\xi_3 + \xi_4 + \xi_5).$$

Then we proceed with a double Fourier series decomposition of $m_{C_\beta}^{k_1, \pm}$ on $\pm[2^{k_1}, 2^{k_1+1}] \times [-2^{k_1-1}, 2^{k_1-1}]$:

$$(3.7) \quad m_{C_\beta}^{k_1, \pm}(\xi_1 + \xi_2, \xi_3 + \xi_4 + \xi_5) = \sum_{L_1, L_2 \in \mathbb{Z}} C_{L_1, L_2}^\pm 2^{k_1(\beta-1)} e^{2\pi i L_1 \frac{\xi_1 + \xi_2}{2^{k_1}}} e^{2\pi i L_2 \frac{\xi_3 + \xi_4 + \xi_5}{2^{k_1}}}.$$

Similarly, we perform a Fourier series decomposition of $|\xi_1 + \xi_2|^\alpha$ on the same interval $\pm[2^{k_1}, 2^{k_1+1}]$:

$$(3.8) \quad |\xi_1 + \xi_2|^\alpha \tilde{\psi}_{k_1, \pm}(\xi_1 + \xi_2) = \sum_{\tilde{L} \in \mathbb{Z}} C_{\tilde{L}}^\pm 2^{k_1 \alpha} e^{\frac{2\pi i \tilde{L}(\xi_1 + \xi_2)}{2^{k_1}}},$$

and notice that the coefficients²⁴ C_{L_1, L_2}^\pm and $C_{\tilde{L}}^\pm$ (which depend on β and α , but not on k_1) decay fast enough: see (2.44).

Thanks to these Fourier series decompositions of the symbols, $T_{m_{IA}}(f_1, \dots, f_5)$ becomes a superposition of tensorized operators of the form:

$$\begin{aligned} & \sum_{L_1, L_2 \in \mathbb{Z}} \sum_{\tilde{L} \in \mathbb{Z}} C_{L_1, L_2}^\pm C_{\tilde{L}}^\pm \int_{\mathbb{R}^5} 2^{k_1(\beta-1)} 2^{k_3} 2^{k_1 \alpha} \widehat{\Delta_{k_1, \pm} f_1}(\xi_1) e^{\frac{2\pi i (L_1 + \tilde{L}) \xi_1}{2^{k_1}}} \cdot \widehat{S_{k_1} f_2}(\xi_2) e^{\frac{2\pi i (L_1 + \tilde{L}) \xi_2}{2^{k_1}}} \\ & \cdot \widehat{\Delta_{k_3} f_3}(\xi_3) \tilde{\psi}_{k_3}(\xi_3) \frac{\xi_3}{2^{k_3}} e^{\frac{2\pi i L_2 \xi_3}{2^{k_1}}} \cdot |\xi_4 + \xi_5|^\gamma \cdot \widehat{S_{k_1} f_4}(\xi_4) e^{\frac{2\pi i L_2 \xi_4}{2^{k_1}}} \cdot \widehat{S_{k_1} f_5}(\xi_5) e^{\frac{2\pi i L_2 \xi_5}{2^{k_1}}} e^{2\pi i x(\xi_1 + \dots + \xi_5)} d\xi_1 \dots d\xi_5 \\ & := \sum_{L_1, L_2 \in \mathbb{Z}} \sum_{\tilde{L} \in \mathbb{Z}} C_{L_1, L_2}^\pm C_{\tilde{L}}^\pm 2^{k_1(\beta-1)} 2^{k_3} 2^{k_1 \alpha} (\Delta_{k_1, \pm, \frac{L_1 + \tilde{L}}{2^{k_1}}} f_1)(x) \cdot (S_{k_1, \frac{L_1 + \tilde{L}}{2^{k_1}}} f_2)(x) \cdot (\tilde{\Delta}_{k_3, \frac{L_2}{2^{k_1}}} f_3)(x) \\ & \cdot D^\gamma(S_{k_1, \frac{L_2}{2^{k_1}}} f_4 \cdot S_{k_1, \frac{L_2}{2^{k_1}}} f_5)(x). \end{aligned}$$

This step is precisely the *tensorization into subtrees* part of our strategy.

Now we simply notice that, for $0 < \tau \leq \min(1, r)$, we can use Hölder's inequality with $\frac{1}{p_1} + \frac{1}{p_2} + \frac{1}{p_3} + \frac{1}{p_{4,5}} = \frac{1}{r}$ for each of the tensorized structures:

$$\begin{aligned} \|T_{m_{IA}}^{k_1, k_3}(f_1, \dots, f_5)\|_r^\tau & \lesssim \sum_{L_1, L_2 \in \mathbb{Z}} \sum_{\tilde{L} \in \mathbb{Z}} |C_{L_1, L_2}^\pm|^\tau |C_{\tilde{L}}^\pm|^\tau 2^{k_1(\beta-1)\tau} 2^{k_3\tau} 2^{k_1\alpha\tau} \|\Delta_{k_1, \pm, \frac{L_1 + \tilde{L}}{2^{k_1}}} f_1\|_{p_1}^\tau \|S_{k_1, \frac{L_1 + \tilde{L}}{2^{k_1}}} f_2\|_{p_2}^\tau \\ & \quad \|\tilde{\Delta}_{k_3, \frac{L_2}{2^{k_1}}} \Delta_{k_3} f_3\|_{p_3}^\tau \|D^\gamma(S_{k_1, \frac{L_2}{2^{k_1}}} f_4 \cdot S_{k_1, \frac{L_2}{2^{k_1}}} f_5)\|_{p_{4,5}}^\tau. \end{aligned}$$

For the last term, we invoke the bilinear Leibniz rule (in this particular case, the unified result that first appeared in [18])

$$\|D^\gamma(S_{k_1, \frac{L_2}{2^{k_1}}} f_4 \cdot S_{k_1, \frac{L_2}{2^{k_1}}} f_5)\|_{p_{4,5}} \lesssim \|D^\gamma S_{k_1, \frac{L_2}{2^{k_1}}} f_4\|_{p_4} \cdot \|S_{k_1, \frac{L_2}{2^{k_1}}} f_5\|_{p_5} + \|S_{k_1, \frac{L_2}{2^{k_1}}} f_4\|_{p_4} \cdot \|D^\gamma S_{k_1, \frac{L_2}{2^{k_1}}} f_5\|_{p_5},$$

which holds true whenever $\frac{1}{p_{4,5}} = \frac{1}{p_4} + \frac{1}{p_5}$, $1 \leq p_4, p_5 \leq \infty$, and $\frac{1}{p_{4,5}} < 1 + \gamma$.

Now the estimates (2.16) and (2.19) (and implicitly the fact that $1 \leq p_i \leq \infty$ for all $1 \leq i \leq 5$) imply that

$$\|I_A\|_r^\tau \lesssim \sum_{k_3 < k_1} 2^{k_1(\beta-1)\tau} 2^{k_3\tau} 2^{k_1\alpha\tau} \|\Delta_{k_1} f_1\|_{p_1}^\tau \|f_2\|_{p_2}^\tau \|\Delta_{k_3} f_3\|_{p_3}^\tau (\|D^\gamma f_4\|_{p_4} \cdot \|f_5\|_{p_5} + \|f_4\|_{p_4} \cdot \|D^\gamma f_5\|_{p_5})^\tau.$$

So we are left with proving the estimate

$$(3.9) \quad \sum_{k_3 < k_1} 2^{k_1(\beta-1)\tau} 2^{k_3\tau} 2^{k_1\alpha\tau} \|\Delta_{k_1} f_1\|_{p_1}^\tau \|\Delta_{k_3} f_3\|_{p_3}^\tau \lesssim \|D^{\alpha+\beta} f_1\|_{p_1}^\tau \cdot \|f_3\|_{p_3}^\tau + \|D^\alpha f_1\|_{p_1}^\tau \cdot \|D^\beta f_3\|_{p_3}^\tau.$$

Just like in Section 2.2,

$$\begin{aligned} \sum_{k_3 < k_1} 2^{k_1(\beta-1)\tau} 2^{k_3\tau} 2^{k_1\alpha\tau} \|\Delta_{k_1} f_1\|_{p_1}^\tau \|\Delta_{k_3} f_3\|_{p_3}^\tau & \lesssim \sum_{k_1} \min(2^{k_1\beta\tau} \|D^\alpha f_1\|_{\dot{B}_{p_1, \infty}^0}^\tau \|f_3\|_{\dot{B}_{p_3, \infty}^0}^\tau, 2^{-k_1\epsilon\tau} \|D^\alpha f_1\|_{\dot{B}_{p_1, \infty}^\beta}^\tau \|f_3\|_{\dot{B}_{p_3, \infty}^\epsilon}^\tau) \\ & \lesssim (\|D^\alpha f_1\|_{\dot{B}_{p_1, \infty}^0}^\tau \|f_3\|_{\dot{B}_{p_3, \infty}^0}^\tau)^{\frac{\epsilon}{\beta+\epsilon}} (\|D^\alpha f_1\|_{\dot{B}_{p_1, \infty}^\beta}^\tau \|f_3\|_{\dot{B}_{p_3, \infty}^\epsilon}^\tau)^{\frac{\beta}{\beta+\epsilon}} \\ & \lesssim (\|D^\alpha f_1\|_{\dot{B}_{p_1, \infty}^0}^\tau \|f_3\|_{\dot{B}_{p_3, \infty}^\beta}^\tau)^{\frac{\epsilon}{\beta+\epsilon}} (\|D^\alpha f_1\|_{\dot{B}_{p_1, \infty}^\tau}^\tau \|f_3\|_{\dot{B}_{p_3, \infty}^0}^\tau)^{\frac{\beta}{\beta+\epsilon}}. \end{aligned}$$

Using Young's inequality and standard properties of Besov norms – more specifically (2.7), we deduce that the expression above is bounded by

$$(\|D^\alpha f_1\|_{p_1} \|D^\beta f_3\|_{p_3} + \|D^{\alpha+\beta} f_1\|_{p_1} \|f_3\|_{p_3})^\tau.$$

I_B) In frequency we are still restricted to the region R_1 (3.4), but the shape of the multiplier m_{IB} suggests the important role played by $\xi_4 + \xi_5$ in the current situation, which requires an extra conical decomposition. We assume without loss of generality that we are restricted to the region

$$R_{1,5} = \{(\xi_1, \dots, \xi_5) : |\xi_1| \gg |\xi_2|, \dots, |\xi_5| \text{ and } |\xi_4| \leq |\xi_5|\}.$$

²⁴Notice that the *Fourier coefficients* consist of $C_{L_1, L_2}^\pm 2^{k_1(\beta-1)}$ and $C_{\tilde{L}}^\pm 2^{k_1 \alpha}$, respectively.

In consequence, we can sum in (3.3) over $k_2, k_3 \ll k_1$ and $k_4 \leq k_5$ and focus our attention on

$$T_{m_{IB}}^{k_1, k_5}(f_1, \dots, f_5)(x) := \int_{\mathbb{R}^5} \frac{|\xi_1 + \dots + \xi_5|^\beta - |\xi_1 + \xi_2|^\beta}{\xi_3 + \xi_4 + \xi_5} \cdot (\xi_4 + \xi_5) \cdot |\xi_1 + \xi_2|^\alpha \cdot |\xi_4 + \xi_5|^\gamma \\ \cdot \widehat{\Delta_{k_1} f_1}(\xi_1) \cdot \widehat{S_{k_1} f_2}(\xi_2) \cdot \widehat{S_{k_1} f_3}(\xi_3) \cdot \widehat{\Delta_{\leq k_5} f_4}(\xi_4) \cdot \widehat{\Delta_{k_5} f_5}(\xi_5) e^{2\pi i x(\xi_1 + \dots + \xi_5)} d\xi_1 \dots d\xi_5.$$

As before, we perform a double Fourier series decomposition of $m_{C_\beta}^{k_1, \pm}$ on $\pm[2^{k_1}, 2^{k_1+1}] \times [-2^{k_1-1}, 2^{k_1-1}]$, and similarly to (3.8), we decompose $|\xi_1 + \xi_2|^\alpha$ on $\pm[2^{k_1}, 2^{k_1+1}]$; in both cases, the coefficients have arbitrary decay.

If we denote by d the classical derivative on \mathbb{R} , we have

$$\|T_{m_{IA}}^{k_1, k_5}(f_1, \dots, f_5)\|_r^\tau \lesssim \sum_{L_1, L_2 \in \mathbb{Z}} \sum_{\tilde{L} \in \mathbb{Z}} |C_{L_1, L_2}^\pm|^\tau |C_{\tilde{L}}^\pm|^\tau 2^{k_1(\beta-1)\tau} 2^{k_1\alpha\tau} \|\Delta_{k_1, \pm, \frac{L_1+\tilde{L}}{2^{k_1}}} f_1\|_{p_1}^\tau \|S_{k_1, \frac{L_1+\tilde{L}}{2^{k_1}}} f_2\|_{p_2}^\tau \|S_{k_1, \frac{L_2}{2^{k_1}}} f_3\|_{p_3}^\tau \\ \cdot \|D^\gamma \circ d(\Delta_{\leq k_5, \frac{L_2}{2^{k_1}}} f_4 \cdot \Delta_{k_5, \frac{L_2}{2^{k_1}}} f_5)\|_{p_{4,5}}^\tau.$$

We again use the induction hypothesis for estimating the last term; however, f_4 and f_5 are already well localized in frequency, so we can invoke (2.27) and (2.28):

$$\|D^\gamma \circ d(\Delta_{\leq k_5, \frac{L_2}{2^{k_1}}} f_4 \cdot \Delta_{k_5, \frac{L_2}{2^{k_1}}} f_5)\|_{p_{4,5}} \lesssim 2^{k_5} \|f_4\|_{p_4} \|\Delta_{k_5} D^\gamma f_5\|_{p_5}$$

for $1 \leq p_4, p_5 \leq \infty$, $\frac{1}{1+\gamma} < p_{4,5} \leq \infty$. The last inequality follows from (2.19) and Young's convolution inequality.

With these considerations, and invoking again (2.19) and (2.16), we can simply focus on bounding

$$\|I_B\|_r^\tau \lesssim \sum_{k_5 < k_1} 2^{k_1(\beta-1)\tau} 2^{k_5\tau} 2^{k_1\alpha\tau} \|\Delta_{k_1} f_1\|_{p_1}^\tau \|f_2\|_{p_2}^\tau \|f_3\|_{p_3}^\tau \|f_4\|_{p_4}^\tau \|\Delta_{k_5} f_5\|_{p_5}^\tau \\ \lesssim \|f_2\|_{p_2}^\tau \|f_3\|_{p_3}^\tau \|f_4\|_{p_4}^\tau \left(\sum_{k_5 < k_1} 2^{k_1(\beta-1)\tau} 2^{k_5\tau} \|\Delta_{k_1} D^\alpha f_1\|_{p_1}^\tau \|\Delta_{k_5} D^\gamma f_5\|_{p_5}^\tau \right).$$

As before, we make appear the Besov norms, and we optimize in the 2^{k_1} parameter: the expression in the last display involving k_5 and k_1 is bounded above by

$$\sum_{k_1} \min(2^{k_1\beta\tau} \|D^\alpha f_1\|_{\dot{B}_{p_1, \infty}^0}^\tau \|D^\gamma f_5\|_{\dot{B}_{p_5, \infty}^0}^\tau, 2^{-k_1\epsilon\tau} \|D^\alpha f_1\|_{\dot{B}_{p_1, \infty}^\beta}^\tau \|D^\gamma f_5\|_{\dot{B}_{p_5, \infty}^\epsilon}^\tau) \\ \lesssim \left(\|D^\alpha f_1\|_{\dot{B}_{p_1, \infty}^0}^\tau \|D^\gamma f_5\|_{\dot{B}_{p_5, \infty}^0}^\tau \right)^{\frac{\epsilon}{\beta+\epsilon}} \left(\|D^\alpha f_1\|_{\dot{B}_{p_1, \infty}^\beta}^\tau \|D^\gamma f_5\|_{\dot{B}_{p_5, \infty}^\epsilon}^\tau \right)^{\frac{\beta}{\beta+\epsilon}} \\ \lesssim \left(\|D^\alpha f_1\|_{\dot{B}_{p_1, \infty}^0}^\tau \|D^\gamma f_5\|_{\dot{B}_{p_5, \infty}^\beta}^\tau \right)^{\frac{\epsilon}{\beta+\epsilon}} \left(\|D^\alpha f_1\|_{\dot{B}_{p_1, \infty}^\beta}^\tau \|D^\gamma f_5\|_{\dot{B}_{p_5, \infty}^0}^\tau \right)^{\frac{\beta}{\beta+\epsilon}}.$$

Notice that the D^β derivatives are distributed between the $D^\alpha f_1$ and $D^\gamma f_5$ functions.

I_C) In this case, only the Littlewood-Paley information for f_1 and f_2 will be needed, so we sum over $k_3, k_4, k_5 \ll k_1$ in (3.3), focusing on

$$T_{m_{IC}}^{k_1, k_2}(f_1, \dots, f_5)(x) := \int_{\mathbb{R}^5} \frac{|\xi_1 + \xi_2|^{\alpha+\beta} - |\xi_1|^{\alpha+\beta}}{\xi_2} \cdot \xi_2 \cdot |\xi_4 + \xi_5|^\gamma \cdot \widehat{\Delta_{k_1} f_1}(\xi_1) \cdot \widehat{\Delta_{k_2} f_2}(\xi_2) \\ \cdot \widehat{S_{k_1} f_3}(\xi_3) \cdot \widehat{S_{k_1} f_4}(\xi_4) \cdot \widehat{S_{k_1} f_5}(\xi_5) e^{2\pi i x(\xi_1 + \dots + \xi_5)} d\xi_1 \dots d\xi_5.$$

This factorizes straightway into

$$\left(\int_{\mathbb{R}^2} \frac{|\xi_1 + \xi_2|^{\alpha+\beta} - |\xi_1|^{\alpha+\beta}}{\xi_2} \cdot \xi_2 \cdot \widehat{\Delta_{k_1} f_1}(\xi_1) \cdot \widehat{\Delta_{k_2} f_2}(\xi_2) e^{2\pi i x(\xi_1 + \xi_2)} d\xi_1 d\xi_2 \right) \cdot S_{k_1} f_3(x) \cdot D^\gamma(S_{k_1} f_4 \cdot S_{k_1} f_5)(x).$$

The first term corresponds to $m_{C_{\alpha+\beta}}(\xi_1, \xi_2)$, a commutator symbol as in (2.39), so we simply estimate it as in (2.46). For the remaining terms, we use the bilinear Leibniz rule result and the boundedness of the S_{k_1} operator to deduce

$$\|I_C\|_r^\tau \lesssim \left(\sum_{k_2 < k_1} 2^{k_1(\alpha+\beta-1)\tau} 2^{k_2\tau} \|\Delta_{k_1} f_1\|_{p_1}^\tau \|\Delta_{k_2} f_2\|_{p_2}^\tau \right) \cdot \|f_3\|_{p_3}^\tau \left(\|D^\gamma f_4\|_{p_4} \cdot \|f_5\|_{p_5} + \|f_4\|_{p_4} \cdot \|D^\gamma f_5\|_{p_5} \right)^\tau.$$

The summation over $k_2 < k_1$ is reduced as before to

$$\sum_{k_1} \min(2^{k_1(\alpha+\beta)\tau} \|f_1\|_{\dot{B}_{p_1, \infty}^0}^\tau \|f_2\|_{\dot{B}_{p_2, \infty}^0}^\tau, 2^{-k_1\epsilon\tau} \|f_1\|_{\dot{B}_{p_1, \infty}^{\alpha+\beta}}^\tau \|f_2\|_{\dot{B}_{p_2, \infty}^\epsilon}^\tau) \lesssim \dots \\ \lesssim \left(\|f_1\|_{\dot{B}_{p_1, \infty}^0}^\tau \|f_2\|_{\dot{B}_{p_2, \infty}^{\alpha+\beta}}^\tau \right)^{\frac{\epsilon}{\alpha+\beta+\epsilon}} \left(\|f_1\|_{\dot{B}_{p_1, \infty}^{\alpha+\beta}}^\tau \|f_2\|_{\dot{B}_{p_2, \infty}^0}^\tau \right)^{\frac{\alpha+\beta}{\alpha+\beta+\epsilon}}.$$

I_D) For this last term, we need to deal with $T_{m_{ID}}^{k_1}(f_1, \dots, f_5)$, defined by

$$\int_{\mathbb{R}^5} |\xi_1|^{\alpha+\beta} \cdot |\xi_4 + \xi_5|^\gamma \widehat{\Delta_{k_1} f_1}(\xi_1) \cdot \widehat{S_{k_1} f_2}(\xi_2) \cdot \widehat{S_{k_1} f_3}(\xi_3) \cdot \widehat{S_{k_1} f_4}(\xi_4) \cdot \widehat{S_{k_1} f_5}(\xi_5) e^{2\pi i x(\xi_1 + \dots + \xi_5)} d\xi_1 \dots d\xi_5 \\ = (\Delta_{k_1} D^{\alpha+\beta} f_1)(x) S_{k_1} f_2(x) S_{k_1} f_3(x) D^\gamma(S_{k_1} f_4 \cdot S_{k_1} f_5)(x).$$

The idea is to write each $S_{k_1} f_l$, for $2 \leq l \leq 5$, as

$$S_{k_1} f_l(x) = f_l(x) - \Delta_{>k_1} f_l(x).$$

Then $T_{m_{ID}}^{k_1}(f_1, \dots, f_5)$ becomes, once we sum in k_1 ,

$$\sum_{k_1} (\Delta_{k_1} D^{\alpha+\beta} f_1)(x) (f_2(x) - \Delta_{>k_1} f_2)(x) (f_3(x) - \Delta_{>k_1} f_3)(x) D^\gamma((f_4 - \Delta_{>k_1} f_4) \cdot (f_5 - \Delta_{>k_1} f_5))(x) \\ = (D^{\alpha+\beta} f_1)(x) \cdot f_2(x) \cdot f_3(x) \cdot D^\gamma(f_4 \cdot f_5)(x) - \sum_{k_1} (\Delta_{k_1} D^{\alpha+\beta} f_1)(x) (\Delta_{>k_1} f_2)(x) f_3(x) D^\gamma(f_4 \cdot f_5)(x) \\ - \sum_{k_1} (\Delta_{k_1} D^{\alpha+\beta} f_1)(x) f_2(x) f_3(x) D^\gamma((\Delta_{>k_1} f_4) \cdot f_5)(x) + \text{similar terms}.$$

The first term is bounded thanks to Hölder's inequality and the boundedness of the bilinear Leibniz rule for $D^\gamma(f_4 \cdot f_5)$. For the remaining terms, the D^β derivatives will be shared between $D^\alpha \Delta_{k_1} f_1$ and $\Delta_{>k_1} f_l$, for some $l \in \{2, 3, 4, 5\}$. If there is more than one function with associated projections $\Delta_{>k_1}$, we simply use one of them.

Two situations become apparent: 1) when the function associated to the maximal scale (in this case f_1 which corresponds to k_1) and the function on which $\Delta_{>k_1}$ acts are in the same subtree, and 2) when the functions are in different subtrees.

The term $\sum_{k_1} (\Delta_{k_1} D^{\alpha+\beta} f_1)(\Delta_{>k_1} f_2) f_3 D^\gamma(f_4 \cdot f_5)$ corresponds to the first situation (f_1 and f_2 are contained in the same subtree associated to the Leibniz rule $D^\alpha(f_1 \cdot f_2)$), and $\sum_{k_1} (\Delta_{k_1} D^{\alpha+\beta} f_1) f_2 f_3 D^\gamma((\Delta_{>k_1} f_4) \cdot f_5)$ to the second one (f_1 and f_4 are leaves in different subtrees).

In order to estimate $\sum_{k_1} (\Delta_{k_1} D^{\alpha+\beta} f_1)(x) (\Delta_{>k_1} f_2)(x) f_3(x) D^\gamma(f_4 \cdot f_5)(x)$, we start with the observation that

$$(3.10) \quad \|\Delta_{>k_1} f_2\|_{p_2} \leq \min(\|f_2\|_{p_2}, 2^{-\epsilon k_1} \|f_2\|_{\dot{B}_{p_2, \infty}^\epsilon}).$$

Then we use Hölder's inequality and the bilinear Leibniz rule (2.20) for the $D^\gamma(f_4 \cdot f_5)$ part, to again, reduce ourselves to estimating

$$\sum_{k_1} 2^{k_1 \beta} \|\Delta_{k_1} f_1\|_{p_1}^\tau \|\Delta_{>k_1} f_2\|_{p_2}^\tau \lesssim \sum_{k_1} \min(2^{k_1 \beta \tau} \|D^\alpha f_1\|_{\dot{B}_{p_1, \infty}^0}^\tau \|f_2\|_{p_2}^\tau, 2^{-k_1 \epsilon \tau} \|D^\alpha f_1\|_{\dot{B}_{p_1, \infty}^\beta}^\tau \|f_2\|_{\dot{B}_{p_2, \infty}^\epsilon}^\tau) \\ \lesssim (\|D^\alpha f_1\|_{\dot{B}_{p_1, \infty}^0}^\tau \|f_2\|_{p_2}^\tau)^{\frac{\epsilon}{\beta+\epsilon}} (\|D^\alpha f_1\|_{\dot{B}_{p_1, \infty}^\beta}^\tau \|f_2\|_{\dot{B}_{p_2, \infty}^\epsilon}^\tau)^{\frac{\beta}{\beta+\epsilon}} \lesssim (\|D^\alpha f_1\|_{\dot{B}_{p_1, \infty}^0}^\tau \|f_2\|_{\dot{B}_{p_2, \infty}^\epsilon}^\tau)^{\frac{\epsilon}{\beta+\epsilon}} (\|D^\alpha f_1\|_{\dot{B}_{p_1, \infty}^\beta}^\tau \|f_2\|_{p_2}^\tau)^{\frac{\beta}{\beta+\epsilon}}.$$

Of course, the $D^{\alpha+\beta}$ derivatives could be shared between f_1 and f_2 .

In the second situation, we start by applying Hölder to obtain

$$(3.11) \quad \sum_{k_1} 2^{k_1 \beta \tau} \|D^\alpha \Delta_{k_1} f_1\|_{p_1}^\tau \|f_2\|_{p_2}^\tau \|f_3\|_{p_3}^\tau \|D^\gamma((\Delta_{>k_1} f_4) \cdot f_5)\|_{p_{4,5}}^\tau.$$

Instead of directly applying the bilinear Leibniz rule (2.20) to $D^\gamma((\Delta_{>k_1} f_4) \cdot f_5)$, we perform an extra para-product decomposition. This is because we want the terms appearing in the Leibniz rule to coincide with those obtained by regular composition: we should not have terms such as $\|f_1\|_{p_1} \|D^\beta f_4\|_{p_4} \|D^\gamma f_5\|_{p_5}$ appearing. Hence we write

$$D^\gamma((\Delta_{>k_1} f_4) \cdot f_5) = \sum_{k_4} D^\gamma((\Delta_{k_4} \Delta_{>k_1} f_4) \cdot (\Delta_{\leq k_4} f_5)) + \sum_{k_5} D^\gamma((S_{k_5} \Delta_{>k_1} f_4) \cdot (\Delta_{k_5} f_5)).$$

Now notice that $\Delta_{k_4} \Delta_{>k_1} \neq 0$ only if $k_4 > k_1$, and similarly $S_{k_5} \Delta_{>k_1} \neq 0$ only if $k_5 > k_1$. So in fact

$$\sum_{k_4} D^\gamma((\Delta_{k_4} \Delta_{>k_1} f_4) \cdot (\Delta_{\leq k_4} f_5)) = \sum_{k_4 > k_1} D^\gamma((\Delta_{k_4} \Delta_{>k_1} f_4) \cdot (\Delta_{\leq k_4} f_5))$$

and

$$\sum_{k_5} D^\gamma((S_{k_5} \Delta_{>k_1} f_4) \cdot (\Delta_{k_5} f_5)) = \sum_{k_5 > k_1} D^\gamma((S_{k_5} \Delta_{>k_1} f_4) \cdot (\Delta_{k_5} f_5)).$$

Overall we get, thanks to (2.27) and (2.28),

$$\|D^\gamma((\Delta_{>k_1} f_4) \cdot f_5)\|_{p_{4,5}}^\tau \lesssim \sum_{k_4 > k_1} \|\Delta_{k_4} \Delta_{>k_1} D^\gamma f_4\|_{p_4}^\tau \|\Delta_{\leq k_4} f_5\|_{p_5}^\tau + \sum_{k_5 > k_1} \|S_{k_5} \Delta_{>k_1} f_4\|_{p_4}^\tau \|\Delta_{k_5} D^\gamma f_5\|_{p_5}^\tau \\ \lesssim \sum_{k_4 > k_1} \|\Delta_{k_4} D^\gamma f_4\|_{p_4}^\tau \|f_5\|_{p_5}^\tau + \sum_{k_5 > k_1} \|\Delta_{>k_1} f_4\|_{p_4}^\tau \|\Delta_{k_5} D^\gamma f_5\|_{p_5}^\tau.$$

So that (3.11) is reduced to

$$\|f_2\|_{p_2}^\tau \|f_3\|_{p_3}^\tau \|f_5\|_{p_5}^\tau \left(\sum_{k_4 > k_1} 2^{k_1 \beta \tau} \|D^\alpha \Delta_{k_1} f_1\|_{p_1}^\tau \|\Delta_{k_4} D^\gamma f_4\|_{p_4}^\tau + \|f_2\|_{p_2}^\tau \|f_3\|_{p_3}^\tau \|f_4\|_{p_4}^\tau \left(\sum_{k_5 > k_1} 2^{k_1 \beta \tau} \|D^\alpha \Delta_{k_1} f_1\|_{p_1}^\tau \|\Delta_{k_5} D^\gamma f_5\|_{p_5}^\tau \right) \right).$$

Due to symmetry, we only look at the term $\sum_{k_4 > k_1} 2^{k_1 \beta \tau} \|D^\alpha \Delta_{k_1} f_1\|_{p_1}^\tau \|\Delta_{k_4} D^\gamma f_4\|_{p_4}^\tau$, which can be estimated by

$$\begin{aligned} & \sum_{k_4} \min \left(2^{k_4 \beta \tau} \|D^\alpha f_1\|_{\dot{B}_{p_1, \infty}^0}^\tau \|D^\gamma f_4\|_{\dot{B}_{p_4, \infty}^0}^\tau, 2^{-k_4 \epsilon \tau} \|D^\alpha f_1\|_{\dot{B}_{p_1, \infty}^\epsilon}^\tau \|D^\gamma f_4\|_{\dot{B}_{p_4, \infty}^\beta}^\tau \right) \\ & \lesssim \left(\|D^\alpha f_1\|_{\dot{B}_{p_1, \infty}^0}^\tau \|D^\gamma f_4\|_{\dot{B}_{p_4, \infty}^0}^\tau \right)^{\frac{\epsilon}{\beta + \epsilon}} \left(\|D^\alpha f_1\|_{\dot{B}_{p_1, \infty}^\epsilon}^\tau \|D^\gamma f_4\|_{\dot{B}_{p_4, \infty}^\beta}^\tau \right)^{\frac{\beta}{\beta + \epsilon}} \\ & \lesssim \left(\|D^\alpha f_1\|_{\dot{B}_{p_1, \infty}^\beta}^\tau \|D^\gamma f_4\|_{\dot{B}_{p_4, \infty}^0}^\tau \right)^{\frac{\epsilon}{\beta + \epsilon}} \left(\|D^\alpha f_1\|_{\dot{B}_{p_1, \infty}^0}^\tau \|D^\gamma f_4\|_{\dot{B}_{p_4, \infty}^\beta}^\tau \right)^{\frac{\beta}{\beta + \epsilon}}. \end{aligned}$$

Of course, this is bounded by $(\|D^{\alpha+\beta} f_1\|_{p_1} \|D^\gamma f_4\|_{p_4} + \|D^\alpha f_1\|_{p_1} \|D^{\beta+\gamma} f_4\|_{p_4})^\tau$; we point out that the distribution of derivatives follows the same law as the composition of Leibniz rules, except that now input functions in any L^p spaces, with $1 \leq p \leq \infty$, are admissible.

This exhausts the possible cases corresponding to the restriction to the frequency conical region R_1 (3.4).

3.2. (II): study of the conical region R_3 .

In this case, we restrict our attention to the frequency region R_3 (3.5). Here we split the multiplier associated to the flag in (3.3) into

$$\begin{aligned} & \frac{|\xi_1 + \dots + \xi_5|^\beta - |\xi_3|^\beta}{\xi_1 + \xi_2 + \xi_4 + \xi_5} \cdot (\xi_1 + \xi_2) \cdot |\xi_1 + \xi_2|^\alpha \cdot |\xi_4 + \xi_5|^\gamma + \frac{|\xi_1 + \dots + \xi_5|^\beta - |\xi_3|^\beta}{\xi_1 + \xi_2 + \xi_4 + \xi_5} \cdot |\xi_1 + \xi_2|^\alpha \cdot (\xi_4 + \xi_5) \cdot |\xi_4 + \xi_5|^\gamma \\ & + |\xi_3|^\beta \cdot |\xi_1 + \xi_2|^\alpha \cdot |\xi_4 + \xi_5|^\gamma := m_{II_A} + m_{II_B} + m_{II_C}. \end{aligned}$$

As before, these are combined with Littlewood-Paley projections to obtain that (3.3) restricted to R_3 (3.5) equals

$$\begin{aligned} & \sum_{k_1, \dots, k_5 \ll k_3} T_{m_{II_A}}(\Delta_{k_1} f_1, \dots, \Delta_{k_5} f_5)(x) + \sum_{k_1, \dots, k_5 \ll k_3} T_{m_{II_B}}(\Delta_{k_1} f_1, \dots, \Delta_{k_5} f_5)(x) \\ & + \sum_{k_1, \dots, k_5 \ll k_3} T_{m_{II_C}}(\Delta_{k_1} f_1, \dots, \Delta_{k_5} f_5)(x) := II_A + II_B + II_C. \end{aligned}$$

The multipliers m_{II_A} and m_{II_B} are symmetric, so it will suffice to study II_A and II_C .

II_A) While f_3 is the fastest oscillating function, the shape of m_{II_A} indicates that one of f_1 or f_2 will also be involved in the scale-by-scale analysis. To decide which, an additional paraproduct decomposition concerning the ξ_1 and ξ_2 variables is necessary – for simplicity we assume $|\xi_2| \leq |\xi_1|$.

After summing over $k_4, k_5 \ll k_3$ and $k_2 \leq k_1$ in (3.3), we need to analyze

$$\begin{aligned} T_{m_{II_A}}^{k_3, k_1}(f_1, \dots, f_5)(x) &:= \int_{\mathbb{R}^5} \frac{|\xi_1 + \dots + \xi_5|^\beta - |\xi_3|^\beta}{\xi_1 + \xi_2 + \xi_4 + \xi_5} \cdot (\xi_1 + \xi_2) \cdot |\xi_1 + \xi_2|^\alpha \cdot |\xi_4 + \xi_5|^\gamma \\ & \quad \widehat{\Delta_{k_1} f_1}(\xi_1) \cdot \widehat{\Delta_{\leq k_1} f_2}(\xi_2) \cdot \widehat{\Delta_{k_3} f_3}(\xi_3) \cdot \widehat{S_{k_3} f_4}(\xi_4) \cdot \widehat{S_{k_3} f_5}(\xi_5) e^{2\pi i x(\xi_1 + \dots + \xi_5)} d\xi_1 \dots d\xi_5. \end{aligned}$$

We decompose into Fourier series the symbol

$$m_{C_\beta}^{k_3, \pm}(\xi_3, \xi_1 + \xi_2 + \xi_4 + \xi_5) = \frac{|\xi_1 + \dots + \xi_5|^\beta - |\xi_3|^\beta}{\xi_1 + \xi_2 + \xi_4 + \xi_5} \tilde{\psi}_{k_3, \pm}(\xi_3) \tilde{\varphi}_{k_3}(\xi_1 + \xi_2 + \xi_4 + \xi_5),$$

obtaining arbitrary decay for the Fourier coefficients. This allows to express $T_{m_{II_A}}^{k_3, k_1}(f_1, \dots, f_5)(x)$ as a sum of

$$\sum_{L_1, L_2 \in \mathbb{Z}} C_{L_1, L_2}^\pm 2^{k_3(\beta-1)} D^\alpha \circ d(\Delta_{k_1, \frac{L_2}{2^{k_3}}} f_1 \cdot \Delta_{\leq k_1, \frac{L_2}{2^{k_3}}} f_2)(x) \cdot \Delta_{k_3, \pm, \frac{L_1}{2^{k_3}}} f_3(x) \cdot D^\gamma(S_{k_3, \frac{L_2}{2^{k_3}}} f_4 \cdot S_{k_3, \frac{L_2}{2^{k_3}}} f_5)(x).$$

We will appeal shortly to the boundedness of the bilinear Leibniz rule (2.20), and its frequency-localized versions (2.27) and (2.28),

$$\|D^\alpha \circ d(\Delta_{k_1, \frac{L_2}{2^{k_3}}} f_1 \cdot \Delta_{\leq k_1, \frac{L_2}{2^{k_3}}} f_2)\|_{p_1, 2} \lesssim 2^{k_1} \|\Delta_{k_1} D^\alpha f_1\|_{p_1} \|f_2\|_{p_2},$$

which is true as long as $\frac{1}{p_{1,2}} = \frac{1}{p_1} + \frac{1}{p_2}$, $1 \leq p_1, p_2 \leq \infty$, and $\frac{1}{p_{1,2}} < 1 + \alpha$.

Thus we have, for $0 < \tau \leq \min(1, r)$,

$$\|II_A\|_\tau^\tau \lesssim \sum_{L_1, L_2 \in \mathbb{Z}} |C_{L_1, L_2}^\pm|^\tau \sum_{k_1 < k_3} 2^{k_3(\beta-1)\tau} 2^{k_1\tau} \|\Delta_{k_1} D^\alpha f_1\|_{p_1}^\tau \|f_2\|_{p_2}^\tau \|\Delta_{k_3, \pm} f_3\|_{p_3}^\tau \|D^\gamma(S_{k_3, \frac{L_2}{2^{k_3}}} f_4 \cdot S_{k_3, \frac{L_2}{2^{k_3}}} f_5)\|_{p_{4,5}}^\tau$$

$$\lesssim \sum_{L_1, L_2 \in \mathbb{Z}} |C_{L_1, L_2}^\pm|^\tau \sum_{k_1 < k_3} 2^{k_3(\beta-1)\tau} 2^{k_1\tau} \|\Delta_{k_1} D^\alpha f_1\|_{p_1}^\tau \|f_2\|_{p_2}^\tau \|\Delta_{k_3} f_3\|_{p_3}^\tau \\ \cdot \left(\|S_{k_3, \frac{L_2}{2^{k_3}}} D^\gamma f_4\|_{p_4} \|S_{k_3, \frac{L_2}{2^{k_3}}} f_5\|_{p_5} + \|S_{k_3, \frac{L_2}{2^{k_3}}} f_4\|_{p_4} \|S_{k_3, \frac{L_2}{2^{k_3}}} D^\gamma f_5\|_{p_5} \right)^\tau.$$

Due to the fast decay of the coefficients $|C_{L_1, L_2}^\pm|$, and the boundedness of the $\Delta_{\leq k_1, a}$ and $S_{k_3, a}$ operators (with norms independent of k_1 , k_3 or a), we are left with summing

$$\sum_{k_1 < k_3} 2^{k_3(\beta-1)\tau} 2^{k_1\tau} \|\Delta_{k_1} D^\alpha f_1\|_{p_1}^\tau \|\Delta_{k_3} f_3\|_{p_3}^\tau \lesssim \sum_{k_3} \min \left(2^{k_3\beta\tau} \|D^\alpha f_1\|_{\dot{B}_{p_1, \infty}^\tau}^\tau \|f_3\|_{\dot{B}_{p_3, \infty}^\tau}^\tau, 2^{-k_3\epsilon\tau} \|D^\alpha f_1\|_{\dot{B}_{p_1, \infty}^\tau}^\tau \|f_3\|_{\dot{B}_{p_3, \infty}^\tau}^\tau \right) \\ \lesssim \left(\|D^\alpha f_1\|_{\dot{B}_{p_1, \infty}^\tau}^\tau \|f_3\|_{\dot{B}_{p_3, \infty}^\tau}^\tau \right)^{\frac{\epsilon}{\beta+\epsilon}} \left(\|D^\alpha f_1\|_{\dot{B}_{p_1, \infty}^\tau}^\tau \|f_3\|_{\dot{B}_{p_3, \infty}^\tau}^\tau \right)^{\frac{\beta}{\beta+\epsilon}}.$$

II_C) Now we look at II_C , which for a fixed $k_3 \in \mathbb{Z}$ and after summing in $k_1, k_2, k_4, k_5 \ll k_3$ in (3.3), corresponds to the operator

$$(3.12) \quad D^\alpha(S_{k_3} f_1 \cdot S_{k_3} f_2) \cdot \Delta_{k_3} D^\beta f_3 \cdot D^\gamma(S_{k_3} f_4 \cdot S_{k_3} f_5).$$

The derivatives D^β hit the fastest oscillating function, i.e. f_3 . We still need to sum over k_3 , so although the operator in (3.12) is tensorized, it will not be trivial to estimate. Here we switch the order of summation again as in I_D , and the operator in (3.12) becomes

$$(3.13) \quad D^\alpha((f_1 - \Delta_{>k_3} f_1) \cdot (f_2 - \Delta_{>k_3} f_2)) \cdot \Delta_{k_3} D^\beta f_3 \cdot D^\gamma((f_4 - \Delta_{>k_3} f_4) \cdot (f_5 - \Delta_{>k_3} f_5)).$$

We use the linearity of the derivation operators to sum over k_3 the expressions in (3.13); we have

$$II_C = D^\alpha(f_1 \cdot f_2) \cdot D^\beta f_3 \cdot D^\gamma(f_4 \cdot f_5) - \sum_{k_3} D^\alpha(\Delta_{>k_3} f_1 \cdot f_2) \cdot \Delta_{k_3} D^\beta f_3 \cdot D^\gamma(f_4 \cdot f_5) \\ - \sum_{k_3} D^\alpha(f_1 \cdot \Delta_{>k_3} f_2) \cdot \Delta_{k_3} D^\beta f_3 \cdot D^\gamma(f_4 \cdot f_5) + \text{similar term}$$

The first term can be easily bounded in L^r thanks to Hölder's inequality and the paraproduct Leibniz rule (2.20). For the other terms, it will be sufficient to use one of the functions on which $\Delta_{>k_3}$ acts; and if it acts on several, we pick one of them, which will contribute to the summation in k_3 . For flag Leibniz rules, a special attention is required by the distribution of the derivatives; for that reason, when we are compelled to use a function belonging to a different subtree than f_3 , an intermediate step is necessary to make sure that the D^β derivatives will be distributed according to the composition laws. It will be enough to treat the second term, since the remaining ones are very similar. It will be bounded, in $\|\cdot\|_r^\tau$, by

$$\sum_{k_3} \|D^\alpha(\Delta_{>k_3} f_1 \cdot f_2)\|_{p_{1,2}}^\tau \|\Delta_{k_3} D^\beta f_3\|_{p_3}^\tau \|D^\gamma(\Delta_{>k_3} f_4 \cdot f_5)\|_{p_{4,5}}^\tau \\ \lesssim \sum_{k_3} \|D^\alpha(\Delta_{>k_3} f_1 \cdot f_2)\|_{p_{1,2}}^\tau \|\Delta_{k_3} D^\beta f_3\|_{p_3}^\tau \left(\|D^\gamma f_4\|_{p_4} \cdot \|f_5\|_{p_5} + \|f_4\|_{p_4} \cdot \|D^\gamma f_5\|_{p_5} \right)^\tau.$$

We take a closer look at the factor $D^\alpha(\Delta_{>k_3} f_1 \cdot f_2)$, which is equal to

$$\sum_{k_1} D^\alpha(\Delta_{k_1} \Delta_{>k_3} f_1 \cdot \Delta_{\leq k_1} f_2)(x) + \sum_{k_2} D^\alpha(S_{k_2} \Delta_{>k_3} f_1 \cdot \Delta_{k_2} f_2)(x).$$

We notice that the only way $\Delta_{k_1} \Delta_{>k_3} \neq 0$ is if $k_1 > k_3$, and similarly, $S_{k_2} \Delta_{>k_3} \neq 0$ only if $k_2 > k_3$. Thus the expression above becomes

$$\sum_{k_1 > k_3} D^\alpha(\Delta_{k_1} \Delta_{>k_3} f_1 \cdot \Delta_{\leq k_1} f_2)(x) + \sum_{k_2 > k_3} D^\alpha(S_{k_2} \Delta_{>k_3} f_1 \cdot \Delta_{k_2} f_2)(x).$$

Because of (2.28) and (2.27), which indicate that derivatives tend to move towards higher oscillating functions,

$$\|D^\alpha(\Delta_{>k_3} f_1 \cdot f_2)\|_{p_{1,2}}^\tau \lesssim \sum_{k_1 > k_3} \|D^\alpha(\Delta_{k_1} \Delta_{>k_3} f_1 \cdot \Delta_{\leq k_1} f_2)\|_{p_{1,2}}^\tau + \sum_{k_2 > k_3} \|D^\alpha(S_{k_2} \Delta_{>k_3} f_1 \cdot \Delta_{k_2} f_2)\|_{p_{1,2}}^\tau \\ \lesssim \sum_{k_1 > k_3} \|\Delta_{k_1} D^\alpha f_1\|_{p_1}^\tau \|f_2\|_{p_2}^\tau + \sum_{k_2 > k_3} \|f_1\|_{p_1}^\tau \|\Delta_{k_2} D^\alpha f_2\|_{p_2}^\tau.$$

Hence $\|II_C\|_r^\tau$ is bounded by the sum of several similar terms of the form

$$\sum_{k_1 > k_3} \|\Delta_{k_1} D^\alpha f_1\|_{p_1}^\tau \|f_2\|_{p_2}^\tau 2^{k_3\beta\tau} \|\Delta_{k_3} f_3\|_{p_3}^\tau \|D^\gamma f_4\|_{p_4}^\tau \|f_5\|_{p_5}^\tau.$$

We can put the functions f_2, f_4 and f_5 aside (in the one-parameter case, there will always be two functions involved in this type of summation), so we are left with

$$\begin{aligned} & \sum_{k_1} \min \left(2^{k_1 \beta \tau} \|D^\alpha f_1\|_{\dot{B}_{p_1, \infty}^0}^\tau \|f_3\|_{\dot{B}_{p_3, \infty}^0}^\tau, 2^{-k_1 \epsilon \tau} \|D^\alpha f_1\|_{\dot{B}_{p_1, \infty}^\beta}^\tau \|f_3\|_{\dot{B}_{p_3, \infty}^\epsilon}^\tau \right) \\ & \lesssim \left(\|D^\alpha f_1\|_{\dot{B}_{p_1, \infty}^0}^\tau \|f_3\|_{\dot{B}_{p_3, \infty}^\beta}^\tau \right)^{\frac{\epsilon}{\beta + \epsilon}} \left(\|D^\alpha f_1\|_{\dot{B}_{p_1, \infty}^\beta}^\tau \|f_3\|_{\dot{B}_{p_3, \infty}^0}^\tau \right)^{\frac{\beta}{\beta + \epsilon}}. \end{aligned}$$

3.3. (III): study of “diagonal” conical regions.

Here we restrict the operator in (3.3) to the frequency region (3.6). Due to the structure of the flag and its symmetries, it will be enough to investigate the cases $l_1 = 1, l_2 = 3$ (representing III_A) and $l_1 = 1, l_2 = 5$ (case III_B).

In this situation, we will not make use of commutators; instead, the Littlewood-Paley information of the functions f_{l_1} and f_{l_2} will be sufficient for estimating the summation of the various scales in (3.3).

III_A) In this case, the main contribution will come from the functions f_1 and f_3 . We can sum in (3.3) over $k_2, k_4, k_5 \leq k_1 \sim k_3$ to get

$$\begin{aligned} T_{m_{III_A}}^{k_1}(f_1, \dots, f_5)(x) &:= \int_{\mathbb{R}^5} |\xi_1 + \dots + \xi_5|^\beta \cdot |\xi_1 + \xi_2|^\alpha \cdot |\xi_4 + \xi_5|^\gamma \cdot \widehat{\Delta_{k_1} f_1}(\xi_1) \cdot \widehat{\Delta_{\leq k_1} f_2}(\xi_2) \\ &\quad \cdot \widehat{\Delta_{k_1} f_3}(\xi_3) \cdot \widehat{\Delta_{\leq k_1} f_4}(\xi_4) \cdot \widehat{\Delta_{\leq k_1} f_5}(\xi_5) e^{2\pi i x(\xi_1 + \dots + \xi_5)} d\xi_1 \dots d\xi_5. \end{aligned}$$

Given the assumptions on the scales, we have that $|\xi_1 + \dots + \xi_5| \leq C2^{k_1}$ and $|\xi_1 + \xi_2| \leq 2^{k_1+1}$. Hence²⁵ we use Fourier series on $[-2^{k_1}, 2^{k_1}]$ to tensorize and decompose the $|\xi_1 + \dots + \xi_5|^\beta \tilde{\varphi}_{k_1}(\xi_1 + \dots + \xi_5)$ symbol; the Fourier coefficients will only have limited decay, but that is still okay. $T_{m_{III_A}}^{k_1}(f_1, \dots, f_5)(x)$ becomes

$$\sum_{L \in \mathbb{Z}} C_L 2^{k_1 \beta} D^\alpha(\Delta_{k_1, \frac{L}{2^{k_1}}} f_1 \cdot \Delta_{\leq k_1, \frac{L}{2^{k_1}}} f_2)(x) \cdot \Delta_{k_1, \frac{L}{2^{k_1}}} f_3(x) \cdot D^\gamma(\Delta_{\leq k_1, \frac{L}{2^{k_1}}} f_4 \cdot \Delta_{\leq k_1, \frac{L}{2^{k_1}}} f_5)(x),$$

where $|C_L| \lesssim (1 + |L|)^{-(1+\beta)}$. So for $\tau \leq \min(1, r)$ with $\frac{1}{1+\beta} < \tau$, we have

$$\|III_A\|_r^\tau \lesssim \sum_{k_1} \sum_{L \in \mathbb{Z}} |C_L|^\tau 2^{k_1 \beta \tau} \|D^\alpha(\Delta_{k_1, \frac{L}{2^{k_1}}} f_1 \cdot \Delta_{\leq k_1, \frac{L}{2^{k_1}}} f_2)\|_{p_{1,2}}^\tau \|\Delta_{k_1, \frac{L}{2^{k_1}}} f_3\|_{p_3}^\tau \|D^\gamma(\Delta_{\leq k_1, \frac{L}{2^{k_1}}} f_4 \cdot \Delta_{\leq k_1, \frac{L}{2^{k_1}}} f_5)\|_{p_{4,5}}^\tau.$$

For $p_{1,2}, p_{4,5}$ so that $\frac{1}{1+\alpha} < p_{1,2} = \frac{p_1 p_2}{p_1 + p_2} \leq \infty$, $\frac{1}{1+\gamma} < p_{4,5} = \frac{p_4 p_5}{p_4 + p_5} \leq \infty$, we further deduce

$$\begin{aligned} \|III_A\|_r^\tau &\lesssim \sum_{k_1} \sum_{L \in \mathbb{Z}} C_L^\tau 2^{k_1 \beta \tau} \|D^\alpha f_1\|_{p_1}^\tau \|f_2\|_{p_2}^\tau \|\Delta_{k_1, \frac{L}{2^{k_1}}} f_3\|_{p_3}^\tau \\ &\quad \cdot \left(\|D^\gamma(\Delta_{\leq k_1, \frac{L}{2^{k_1}}} f_4)\|_{p_4} \|\Delta_{\leq k_1, \frac{L}{2^{k_1}}} f_5\|_{p_5} + \|\Delta_{\leq k_1, \frac{L}{2^{k_1}}} f_4\|_{p_4} \|D^\gamma(\Delta_{\leq k_1, \frac{L}{2^{k_1}}} f_5)\|_{p_5} \right)^\tau \\ &\lesssim \sum_{k_1} \|\Delta_{k_1} D^\alpha f_1\|_{p_1}^\tau \|f_2\|_{p_2}^\tau \|\Delta_{k_1} f_3\|_{p_3}^\tau \left(\|D^\gamma f_4\|_{p_4} \|f_5\|_{p_5} + \|f_4\|_{p_4} \|D^\gamma f_5\|_{p_5} \right)^\tau. \end{aligned}$$

Summing in k_1 is by now a formality:

$$\begin{aligned} \sum_{k_1} 2^{k_1 \beta \tau} \|\Delta_{k_1} D^\alpha f_1\|_{p_1}^\tau \|\Delta_{k_1} f_3\|_{p_3}^\tau &\lesssim \sum_{k_1} \min \left(2^{k_1 \beta \tau} \|D^\alpha f_1\|_{\dot{B}_{p_1, \infty}^0}^\tau \|f_3\|_{\dot{B}_{p_3, \infty}^0}^\tau, 2^{-k_1 \beta \tau} \|D^\alpha f_1\|_{\dot{B}_{p_1, \infty}^\beta}^\tau \|f_3\|_{\dot{B}_{p_3, \infty}^\beta}^\tau \right) \\ &\lesssim \left(\|D^\alpha f_1\|_{\dot{B}_{p_1, \infty}^0}^\tau \|f_3\|_{\dot{B}_{p_3, \infty}^\beta}^\tau \right)^{\frac{1}{2}} \left(\|D^\alpha f_1\|_{\dot{B}_{p_1, \infty}^\beta}^\tau \|f_3\|_{\dot{B}_{p_3, \infty}^0}^\tau \right)^{\frac{1}{2}}. \end{aligned}$$

III_B) Now we are in the situation when $|\xi_2|, |\xi_3|, |\xi_4| \leq |\xi_1| \sim |\xi_5|$. Accordingly, we sum over $k_2, k_3, k_4 \leq k_1 \sim k_5$ in (3.3) to obtain

$$\begin{aligned} T_{m_{III_B}}^{k_1}(f_1, \dots, f_5)(x) &:= \int_{\mathbb{R}^5} |\xi_1 + \dots + \xi_5|^\beta \cdot |\xi_1 + \xi_2|^\alpha \cdot |\xi_4 + \xi_5|^\gamma \widehat{\Delta_{k_1} f_1}(\xi_1) \cdot \widehat{\Delta_{\leq k_1} f_2}(\xi_2) \\ &\quad \cdot \widehat{\Delta_{\leq k_1} f_3}(\xi_3) \cdot \widehat{\Delta_{\leq k_1} f_4}(\xi_4) \cdot \widehat{\Delta_{k_1} f_5}(\xi_5) e^{2\pi i x(\xi_1 + \dots + \xi_5)} d\xi_1 \dots d\xi_5. \end{aligned}$$

As before, we smoothly localize and use a Fourier series expansion for $|\xi_1 + \dots + \xi_5|^\beta$ on $[-2^{k_1}, 2^{k_1}]$, with limited decay of the Fourier coefficients. This allows to rewrite

$$T_{m_{III_B}}^{k_1}(f_1, \dots, f_5)(x) = \sum_{L \in \mathbb{Z}} C_L 2^{k_1 \beta} D^\alpha(\Delta_{k_1, \frac{L}{2^{k_1}}} f_1 \cdot \Delta_{\leq k_1, \frac{L}{2^{k_1}}} f_2)(x) \cdot \Delta_{\leq k_1, \frac{L}{2^{k_1}}} f_3(x) \cdot D^\gamma(\Delta_{\leq k_1, \frac{L}{2^{k_1}}} f_4 \cdot \Delta_{k_1, \frac{L}{2^{k_1}}} f_5)(x),$$

²⁵Here we might need to assume a certain amount of separation between the frequency pieces, which is easy to obtain by a sparsification argument that only introduces $O(1)$ new terms.

with $|C_L| \lesssim (1 + |L|)^{-(1+\beta)}$. As before,

$$\|III_B\|_r^\tau \lesssim \|f_2\|_{p_2}^\tau \|f_3\|_{p_3}^\tau \|f_4\|_{p_4}^\tau \left(\|D^\alpha f_1\|_{\dot{B}_{p_1,\infty}^0}^\tau \|D^\gamma f_5\|_{\dot{B}_{p_5,\infty}^\beta}^\tau \right)^{\frac{1}{2}} \left(\|D^\alpha f_1\|_{\dot{B}_{p_1,\infty}^\beta}^\tau \|D^\gamma f_5\|_{\dot{B}_{p_5,\infty}^0}^\tau \right)^{\frac{1}{2}}.$$

This completes the proof of (3.1), which, we recall, follows also from the flag paraproducts' boundedness in [14] – except for certain endpoints.

4. A FIVE-LINEAR FLAG: BI-PARAMETER CASE

Now we move to the bi-parameter version of (3.1), which reads as

$$(4.1) \quad \|D_{(1)}^{\beta_1} D_{(2)}^{\beta_2} (D_{(1)}^{\alpha_1} D_{(2)}^{\alpha_2} (f_1 \cdot f_2) \cdot f_3 \cdot D_{(1)}^{\gamma_1} D_{(2)}^{\gamma_2} (f_4 \cdot f_5))\|_{L^p L^q} \\ \lesssim \|D_{(1)}^{\alpha_1+\beta_1} D_{(2)}^{\alpha_2+\beta_2} f_1\|_{L^{p_1} L^{q_1}} \|f_2\|_{L^{p_2} L^{q_2}} \|f_3\|_{L^{p_3} L^{q_3}} \|D_{(1)}^{\gamma_1} D_{(2)}^{\gamma_2} f_4\|_{L^{p_4} L^{q_4}} \|f_5\|_{L^{p_5} L^{q_5}} + \text{other similar terms}.$$

This is the main motivation of our work, since bi-parameter equivalents of [14] are yet to be proved. For simplicity, we work with functions defined on $\mathbb{R} \times \mathbb{R}$, although all the results remain valid for functions on $\mathbb{R}^{d_1} \times \mathbb{R}^{d_2}$. In order for (4.1) to hold, the conditions on the Lebesgue exponents, already presented in the Introduction, are

$$\frac{1}{q} < 1 + \beta_2, \quad \frac{1}{q_{1,2}} := \frac{1}{q_1} + \frac{1}{q_2} < 1 + \alpha_2, \quad \frac{1}{q_{4,5}} := \frac{1}{q_4} + \frac{1}{q_5} < 1 + \gamma_2, \\ \frac{1}{p} < \min(1 + \beta_1, 1 + \beta_2), \quad \frac{1}{p_{1,2}} := \frac{1}{p_1} + \frac{1}{p_2} < \min(1 + \alpha_1, 1 + \alpha_2), \quad \frac{1}{p_{4,5}} := \frac{1}{p_4} + \frac{1}{p_5} < \min(1 + \gamma_1, 1 + \gamma_2).$$

The asymmetry on the Lebesgue exponents associated to the first and second variables is a consequence of the mixed norm condition. The constraints on p and q are imposed by slower decaying conditions on the associated Fourier coefficients when treating the “diagonal case”: see Section 4.2.

Although conceptually the method employed for proving (4.1) will be similar to that presented in the previous section, technical aspects specific²⁶ to multi-parameter problems will appear; in particular, the number of cases that need to be considered is significantly higher. That is partly due to the asymmetry of the objects: f_2 might not be involved in the summation of the flag acting in the first variable, but we cannot put it aside because it might be involved in the summation of the flag acting in the second variable.

As before in Section 3, in this particular situation we will not systematically apply the inductive procedure presented later in Section 5.2; however, the differences are minor.

In the bi-parameter setting, the multiplier associated to the flag (4.1) is

$$(4.2) \quad (|\xi_1 + \dots + \xi_5|^{\beta_1} |\xi_1 + \xi_2|^{\alpha_1} |\xi_4 + \xi_5|^{\gamma_1}) \cdot (|\eta_1 + \dots + \eta_5|^{\beta_2} |\eta_1 + \eta_2|^{\alpha_2} |\eta_4 + \eta_5|^{\gamma_2}).$$

As before in Section 3, the frequency regions will be decomposed into cones, allowing to determine the fastest oscillating functions in each parameter, and (up to a point) the distribution of derivatives. Depending on the structure of the associated flag/rooted tree and on the conical regions in each parameter, the multiplier will be split into several pieces. However, in each parameter, the decomposition is carried out as in the previous section.

Our aim is to estimate

$$(4.3) \quad \sum_{k_1, \dots, k_5} \sum_{m_1, \dots, m_5} \int_{\mathbb{R}^{10}} (|\xi_1 + \dots + \xi_5|^{\beta_1} |\xi_1 + \xi_2|^{\alpha_1} |\xi_4 + \xi_5|^{\gamma_1}) \cdot (|\eta_1 + \dots + \eta_5|^{\beta_2} |\eta_1 + \eta_2|^{\alpha_2} |\eta_4 + \eta_5|^{\gamma_2}) \mathcal{F}(\Delta_{k_1}^{(1)} \Delta_{m_1}^{(2)} f_1)(\xi_1, \eta_1) \\ \cdot \mathcal{F}(\Delta_{k_2}^{(1)} \Delta_{m_2}^{(2)} f_2)(\xi_2, \eta_2) \cdot \dots \cdot \mathcal{F}(\Delta_{k_5}^{(1)} \Delta_{m_5}^{(2)} f_5)(\xi_5, \eta_5) e^{2\pi i x(\xi_1 + \dots + \xi_5)} e^{2\pi i y(\eta_1 + \dots + \eta_5)} d\xi_1 \dots d\xi_5 d\eta_1 \dots d\eta_5$$

in the mixed (quasi-)norm $\|\cdot\|_{L^p L^q}$. Throughout the section, τ will denote

$$\tau := \min(1, p, q),$$

which renders $\|\cdot\|_{L^p L^q}^\tau$ subadditive.

Next, we will consider various operators that arise from restricting our attention to conical frequency regions $\{|\xi_1| \gg |\xi_2|, \dots, |\xi_5|, |\eta_1| \gg |\eta_1|, \dots, |\eta_5|\}$, etc. This corresponds to a bi-parameter paraproduct decomposition and allows to split (4.3) into pieces that will be independently estimated. We will restrict our attention to certain typical conical regions, as the remaining cases follow from similar arguments.

²⁶These difficulties will be nowhere near as laborious as the technical aspects typical of multi-parameter singular integrals.

4.1. Study of the “off-diagonal” conical region $R_1^{(1)} \times R_1^{(2)}$. When we are in the region where $|\xi_1|$ is the largest frequency variable in the first component and $|\eta_1|$ is the largest one in the second component, we want to approximate $|\xi_1 + \dots + \xi_5|^{\beta_1}$ by $|\xi_1|^{\beta_1}$ (and $|\eta_1 + \dots + \eta_5|^{\beta_2}$ by $|\eta_1|^{\beta_2}$). As before in Section 3, ξ_1 is connected to ξ_2 by the $D_{(1)}^{\alpha_1}$ derivative, so several steps are necessary.

We will have

$$(4.4) \quad \begin{aligned} |\xi_1 + \dots + \xi_5|^{\beta_1} &= |\xi_1 + \dots + \xi_5|^{\beta_1} - |\xi_1 + \xi_2|^{\beta_1} + |\xi_1 + \xi_2|^{\beta_1} - |\xi_1|^{\beta_1} + |\xi_1|^{\beta_1} \\ &= \frac{|\xi_1 + \dots + \xi_5|^{\beta_1} - |\xi_1 + \xi_2|^{\beta_1}}{\xi_3 + \xi_4 + \xi_5} \cdot \xi_3 + \frac{|\xi_1 + \dots + \xi_5|^{\beta_1} - |\xi_1 + \xi_2|^{\beta_1}}{\xi_3 + \xi_4 + \xi_5} \cdot (\xi_4 + \xi_5) + (|\xi_1 + \xi_2|^{\beta_1} - |\xi_1|^{\beta_1}) + |\xi_1|^{\beta_1} \end{aligned}$$

With this decomposition, the part of the multiplier from (4.2) acting on the first coordinate writes as

$$\begin{aligned} &\frac{|\xi_1 + \dots + \xi_5|^{\beta_1} - |\xi_1 + \xi_2|^{\beta_1}}{\xi_3 + \xi_4 + \xi_5} \cdot \xi_3 \cdot |\xi_1 + \xi_2|^{\alpha_1} |\xi_4 + \xi_5|^{\gamma_1} + \frac{|\xi_1 + \dots + \xi_5|^{\beta_1} - |\xi_1 + \xi_2|^{\beta_1}}{\xi_3 + \xi_4 + \xi_5} \cdot (\xi_4 + \xi_5) \cdot |\xi_1 + \xi_2|^{\alpha_1} |\xi_4 + \xi_5|^{\gamma_1} \\ &+ \frac{|\xi_1 + \xi_2|^{\beta_1 + \alpha_1} - |\xi_1|^{\beta_1 + \alpha_1}}{\xi_2} \cdot \xi_2 \cdot |\xi_4 + \xi_5|^{\gamma_1} + |\xi_1|^{\beta_1 + \alpha_1} \cdot |\xi_4 + \xi_5|^{\gamma_1} := m_{I_A^{(1)}} + m_{I_B^{(1)}} + m_{I_C^{(1)}} + m_{I_D^{(1)}}. \end{aligned}$$

The above formulas indicate the presence of three commutators and one multiplier that will require the change in the order of summation. For $m_{I_A^{(1)}}$, the variables involved in the commutator are $\xi_1 + \xi_2$ and ξ_3 , for $m_{I_B^{(1)}}$ - $\xi_1 + \xi_2$ and $\xi_4 + \xi_5$, and for $m_{I_C^{(1)}}$ it will be ξ_1 and ξ_2 that will participate in the summation.

In the second coordinate, the multiplier will split similarly into four terms:


$$\begin{aligned} &\frac{|\eta_1 + \dots + \eta_5|^{\beta_2} - |\eta_1 + \eta_2|^{\beta_2}}{\eta_3 + \eta_4 + \eta_5} \cdot \eta_3 \cdot |\eta_1 + \eta_2|^{\alpha_2} |\eta_4 + \eta_5|^{\gamma_2} + \frac{|\eta_1 + \dots + \eta_5|^{\beta_2} - |\eta_1 + \eta_2|^{\beta_2}}{\eta_3 + \eta_4 + \eta_5} \cdot (\eta_4 + \eta_5) \cdot |\eta_1 + \eta_2|^{\alpha_2} |\eta_4 + \eta_5|^{\gamma_2} \\ &+ \frac{|\eta_1 + \eta_2|^{\beta_2 + \alpha_2} - |\eta_1|^{\beta_2 + \alpha_2}}{\eta_2} \cdot \eta_2 \cdot |\eta_4 + \eta_5|^{\gamma_2} + |\eta_1|^{\beta_2 + \alpha_2} \cdot |\eta_4 + \eta_5|^{\gamma_2} := m_{I_A^{(2)}} + m_{I_B^{(2)}} + m_{I_C^{(2)}} + m_{I_D^{(2)}}. \end{aligned}$$

When the symbol is restricted to the conical region $R_1^{(1)} \times R_1^{(2)}$, (4.3) breaks down as

$$\begin{aligned} &\sum_{\substack{k_2, \dots, k_5 \ll k_1 \\ m_2, \dots, m_5 \ll m_1}} T_{I_A^{(1)} \times I_A^{(2)}}(\Delta_{k_1}^{(1)} \Delta_{m_1}^{(2)} f_1, \dots, \Delta_{k_5}^{(1)} \Delta_{m_5}^{(2)} f_5)(x, y) + \sum_{\substack{k_2, \dots, k_5 \ll k_1 \\ m_2, \dots, m_5 \ll m_1}} T_{I_A^{(1)} \times I_B^{(2)}}(\Delta_{k_1}^{(1)} \Delta_{m_1}^{(2)} f_1, \dots, \Delta_{k_5}^{(1)} \Delta_{m_5}^{(2)} f_5)(x, y) + \\ &\sum_{\substack{k_2, \dots, k_5 \ll k_1 \\ m_2, \dots, m_5 \ll m_1}} T_{I_A^{(1)} \times I_C^{(2)}}(\Delta_{k_1}^{(1)} \Delta_{m_1}^{(2)} f_1, \dots, \Delta_{k_5}^{(1)} \Delta_{m_5}^{(2)} f_5)(x, y) + \dots + \sum_{\substack{k_2, \dots, k_5 \ll k_1 \\ m_2, \dots, m_5 \ll m_1}} T_{I_D^{(1)} \times I_D^{(2)}}(\Delta_{k_1}^{(1)} \Delta_{m_1}^{(2)} f_1, \dots, \Delta_{k_5}^{(1)} \Delta_{m_5}^{(2)} f_5)(x, y) \\ &:= I_A^{(1)} \times I_A^{(2)} + I_A^{(1)} \times I_B^{(2)} + I_A^{(1)} \times I_C^{(2)} + \dots + I_D^{(1)} \times I_D^{(2)}. \end{aligned}$$

In total, we have 16 cases, many of which are similar; so we will only treat some of them, as explained below. Due to the structure of the symbols, $I_A^{(1)} \times I_B^{(2)}$ presents novel attributes, and will be discussed in more detail.

- $I_A^{(1)} \times I_B^{(2)}$



The multiplier suggested by the trees above is $m_{I_A^{(1)}}(\xi_1, \dots, \xi_5) \cdot m_{I_B^{(2)}}(\eta_1, \dots, \eta_5)$, which corresponds to

$$\left(\frac{|\xi_1 + \dots + \xi_5|^{\beta_1} - |\xi_1 + \xi_2|^{\beta_1}}{\xi_3 + \xi_4 + \xi_5} \cdot \xi_3 \cdot |\xi_1 + \xi_2|^{\alpha_1} |\xi_4 + \xi_5|^{\gamma_1} \right) \cdot \left(\frac{|\eta_1 + \dots + \eta_5|^{\beta_2} - |\eta_1 + \eta_2|^{\beta_2}}{\eta_3 + \eta_4 + \eta_5} \cdot (\eta_4 + \eta_5) \cdot |\eta_1 + \eta_2|^{\alpha_2} |\eta_4 + \eta_5|^{\gamma_2} \right).$$

In the first variable the functions involved in the summation over the scales are f_1 and f_3 , so the functions f_2, f_4 and f_5 do not contribute to this process. However, in the second variable, the shape of the commutator indicates that f_1 and one of the functions f_4 or f_5 will play a prominent role; by restricting the symbol further to the region where $|\eta_4| \leq |\eta_5|$, we know the functions f_1 and f_5 will take part in summing up the scales, in the

second parameter. We carefully group the terms contributing to $I_A^{(1)} \times I_B^{(2)}$ by summing over $k_2, k_4, k_5 \ll k_1$ and over $m_2, m_3 \ll m_1, m_4 \leq m_5$:

$$T_{I_A^{(1)}, I_B^{(2)}}^{k_1, k_3; m_1, m_5}(f_1, \dots, f_5)(x, y) = \int_{\mathbb{R}^{10}} m_{I_A^{(1)}}(\xi_1, \dots, \xi_5) \cdot m_{I_B^{(2)}}(\eta_1, \dots, \eta_5) \cdot \mathcal{F}(\Delta_{k_1}^{(1)} \Delta_{m_1}^{(2)} f_1)(\xi_1, \eta_1) \cdot \mathcal{F}(S_{k_1}^{(1)} S_{m_1}^{(2)} f_2)(\xi_2, \eta_2) \\ \cdot \mathcal{F}(\Delta_{k_3}^{(1)} S_{m_1}^{(2)} f_3)(\xi_3, \eta_3) \cdot \mathcal{F}(S_{k_1}^{(1)} \Delta_{\leq m_5}^{(2)} f_4)(\xi_4, \eta_4) \cdot \mathcal{F}(S_{k_1}^{(1)} \Delta_{m_5}^{(2)} f_5)(\xi_5, \eta_5) \cdot e^{2\pi i x(\xi_1 + \dots + \xi_5)} e^{2\pi i y(\eta_1 + \dots + \eta_5)} d\xi_1 \dots d\xi_5 d\eta_1 \dots d\eta_5.$$

As in Section 3, we restrict the symbol

$$\frac{|\xi_1 + \dots + \xi_5|^{\beta_1} - |\xi_1 + \xi_2|^{\beta_1}}{\xi_3 + \xi_4 + \xi_5} = m_{C_{\beta_1}}(\xi_1 + \xi_2, \xi_3 + \xi_4 + \xi_5)$$

to $\pm[2^{k_1}, 2^{k_1+1}] \times [-2^{k_1-1}, 2^{k_1-1}]$ and denote its localized version by $m_{C_{\beta_1}}^{k_1, \pm}(\xi_1 + \xi_2, \xi_3 + \xi_4 + \xi_5)$, on which we perform a double Fourier series decomposition:

$$m_{C_{\beta_1}}^{k_1, \pm}(\xi_1 + \xi_2, \xi_3 + \xi_4 + \xi_5) = \sum_{L_1, L_2} C_{L_1, L_2}^{\pm} 2^{k_1(\beta_1-1)} e^{2\pi i(\xi_1 + \xi_2) \frac{L_1}{2^{k_1}}} e^{2\pi i(\xi_3 + \xi_4 + \xi_5) \frac{L_2}{2^{k_1}}}.$$

Similarly, let $m_{C_{\beta_2}}^{m_1, \pm}(\eta_1 + \eta_2, \eta_3 + \eta_4 + \eta_5)$ denote the symbol $m_{C_{\beta_2}}(\eta_1 + \eta_2, \eta_3 + \eta_4 + \eta_5)$ localized on $\pm[2^{m_1}, 2^{m_1+1}] \times [-2^{m_1-1}, 2^{m_1-1}]$, where

$$m_{C_{\beta_2}}(\eta_1 + \eta_2, \eta_3 + \eta_4 + \eta_5) := \frac{|\eta_1 + \dots + \eta_5|^{\beta_2} - |\eta_1 + \eta_2|^{\beta_2}}{\eta_3 + \eta_4 + \eta_5}.$$

The double Fourier series expansion on $m_{C_{\beta_2}}^{m_1, \pm}$ yields

$$m_{C_{\beta_2}}^{m_1, \pm}(\eta_1 + \eta_2, \eta_3 + \eta_4 + \eta_5) = \sum_{\tilde{L}_1, \tilde{L}_2} C_{\tilde{L}_1, \tilde{L}_2}^{\pm} 2^{m_1(\beta_2-1)} e^{2\pi i(\eta_1 + \eta_2) \frac{\tilde{L}_1}{2^{m_1}}} e^{2\pi i(\eta_3 + \eta_4 + \eta_5) \frac{\tilde{L}_2}{2^{m_1}}}.$$

In both cases, the renormalized Fourier coefficients have arbitrary decay. This implies that

$$\|I_A^{(1)} \times I_B^{(2)}\|_{L^p L^q}^{\tau} \leq \sum_{L_1, L_2} |C_{L_1, L_2}^{\pm}|^{\tau} \sum_{\tilde{L}_1, \tilde{L}_2} |C_{\tilde{L}_1, \tilde{L}_2}^{\pm}|^{\tau} \|D_{(1)}^{\alpha_1} D_{(2)}^{\alpha_2} (\Delta_{k_1, \pm, \frac{L_1}{2^{k_1}}}^{(1)} \Delta_{m_1, \pm, \frac{\tilde{L}_1}{2^{m_1}}}^{(2)} f_1 \cdot S_{k_1, \frac{L_1}{2^{k_1}}}^{(1)} S_{m_1, \frac{\tilde{L}_1}{2^{m_1}}}^{(2)} f_2)\|_{p_{1,2}}^{\tau} \\ \| \Delta_{k_3, \frac{L_2}{2^{k_1}}}^{(1)} S_{m_1, \frac{\tilde{L}_2}{2^{m_1}}}^{(2)} f_3 \|_{L^{p_3} L^{q_3}}^{\tau} \|D_{(1)}^{\gamma_1} D_{(2)}^{\gamma_2} (S_{k_1, \frac{L_2}{2^{k_1}}}^{(1)} \Delta_{\leq m_5, \frac{\tilde{L}_2}{2^{m_1}}}^{(2)} f_4 \cdot S_{k_1, \frac{L_2}{2^{k_1}}}^{(1)} \Delta_{m_5, \frac{\tilde{L}_2}{2^{m_1}}}^{(2)} f_5)\|_{p_{4,5}}^{\tau}.$$

Now we use the boundedness of flag paraproducts of lower complexity: more concretely, the biparameter variant²⁷ of Lemma 2.2 which describes the localized version of Oh and Wu [18] with the additional observation that $D_{(2)}^{\gamma_2}$ derivatives will be attached to f_5 :

$$\|D_{(1)}^{\gamma_1} D_{(2)}^{\gamma_2} (S_{k_1, \frac{L_2}{2^{k_1}}}^{(1)} \Delta_{\leq m_5, \frac{\tilde{L}_2}{2^{m_1}}}^{(2)} f_4 \cdot S_{k_1, \frac{L_2}{2^{k_1}}}^{(1)} \Delta_{m_5, \frac{\tilde{L}_2}{2^{m_1}}}^{(2)} f_5)\|_{L^{p_{4,5}} L^{q_{4,5}}} \lesssim \|D_{(1)}^{\gamma_1} f_4\|_{L^{p_4} L^{q_4}} \| \Delta_{m_5}^{(2)} D_{(2)}^{\gamma_2} f_5 \|_{L^{p_5} L^{q_5}} \\ + \|f_4\|_{L^{p_4} L^{q_4}} \| \Delta_{m_5}^{(2)} D_{(1)}^{\gamma_1} D_{(2)}^{\gamma_2} f_5 \|_{L^{p_5} L^{q_5}}.$$

Here, we need to assume that $p_{4,5} > \frac{1}{1+\gamma_1}$, while $q_{4,5}$ can be any Lebesgue exponent $\geq \frac{1}{2}$. Similarly,²⁸

$$\|D_{(1)}^{\alpha_1} D_{(2)}^{\alpha_2} (\Delta_{k_1, \pm, \frac{L_1}{2^{k_1}}}^{(1)} \Delta_{m_1, \pm, \frac{\tilde{L}_1}{2^{m_1}}}^{(2)} f_1 \cdot S_{k_1, \frac{L_1}{2^{k_1}}}^{(1)} S_{m_1, \frac{\tilde{L}_1}{2^{m_1}}}^{(2)} f_2)\|_{L^{p_{1,2}}} \lesssim \| \Delta_{k_1}^{(1)} \Delta_{m_1}^{(2)} D_{(1)}^{\alpha_1} D_{(2)}^{\alpha_2} f_1 \|_{L^{p_1} L^{q_1}} \|f_2\|_{L^{p_2} L^{q_2}},$$

with $p_{1,2}, q_{1,2} \geq \frac{1}{2}$.

For simplicity, we denote (F_4, F_5) either of the couples $(D_{(1)}^{\gamma_1} f_4, f_5)$ or $(f_4, D_{(1)}^{\gamma_1} f_5)$, and $F_1 := D_{(1)}^{\alpha_1} D_{(2)}^{\alpha_2} f_1$.

Since the renormalized Fourier coefficients C_{L_1, L_2}^{\pm} and $C_{\tilde{L}_1, \tilde{L}_2}^{\pm}$ are summable and all the Fourier projections $P_{k,a}^{(j)}$ are bounded on $L^{p_j} L^{q_j}$ spaces with $1 \leq p_j, q_j \leq \infty$, we are left with summing

$$(4.5) \quad \sum_{k_3 < k_1} \sum_{m_5 < m_1} 2^{k_1(\beta_1-1)\tau} 2^{k_3\tau} 2^{m_1(\beta_2-1)\tau} 2^{m_5\tau} \| \Delta_{k_1}^{(1)} \Delta_{m_1}^{(2)} F_1 \|_{L^{p_1} L^{q_1}}^{\tau} \|f_2\|_{L^{p_2} L^{q_2}}^{\tau} \| \Delta_{k_3}^{(1)} f_3 \|_{L^{p_3} L^{q_3}}^{\tau} \|F_4\|_{L^{p_4} L^{q_4}}^{\tau} \| \Delta_{m_5}^{(2)} D_{(2)}^{\gamma_2} F_5 \|_{L^{p_5} L^{q_5}}^{\tau}.$$

Since f_2 and F_4 have no contribution in the summation, we put them on the side and estimate what is left of (4.5) as

$$(4.6) \quad \sum_{k_1, m_1} \min(2^{k_1\beta_1\tau} 2^{m_1\beta_2\tau} \|F_1\|_{\dot{B}_{p_1, \infty}^0 \dot{B}_{q_1, \infty}^0}^{\tau} \|f_3\|_{\dot{B}_{p_3, \infty}^0 L^{q_3}}^{\tau} \|D_{(2)}^{\gamma_2} F_5\|_{L^{p_5} \dot{B}_{q_5, \infty}^0}^{\tau},$$

²⁷The general statement will be presented in Proposition 5.3.

²⁸Here we use implicitly (2.16) and (2.19)

$$\begin{aligned}
& 2^{k_1 \beta_1 \tau} 2^{-m_1 \epsilon_2 \tau} \|F_1\|_{\dot{B}_{p_1, \infty}^0}^\tau \|\dot{B}_{q_1, \infty}^{\beta_2}\| \|f_3\|_{\dot{B}_{p_3, \infty}^0}^\tau L^{q_3} \|D_{(2)}^{\gamma_2} F_5\|_{L^{p_5} \dot{B}_{q_5, \infty}^{\epsilon_2}}^\tau, \\
& 2^{-k_1 \epsilon_1 \tau} 2^{m_1 \beta_2 \tau} \|F_1\|_{\dot{B}_{p_1, \infty}^{\beta_1}}^\tau \|\dot{B}_{q_1, \infty}^0\| \|f_3\|_{\dot{B}_{p_3, \infty}^{\epsilon_1}}^\tau L^{q_3} \|D_{(2)}^{\gamma_2} F_5\|_{L^{p_5} \dot{B}_{q_5, \infty}^0}^\tau, \\
& 2^{-k_1 \epsilon_1 \tau} 2^{-m_1 \epsilon_2 \tau} \|F_1\|_{\dot{B}_{p_1, \infty}^{\beta_1}}^\tau \|\dot{B}_{q_1, \infty}^{\beta_2}\| \|f_3\|_{\dot{B}_{p_3, \infty}^{\epsilon_1}}^\tau L^{q_3} \|D_{(2)}^{\gamma_2} F_5\|_{L^{p_5} \dot{B}_{q_5, \infty}^{\epsilon_2}}^\tau.
\end{aligned}$$

The aim here is to obtain both positive and negative powers of 2^{k_1} and 2^{m_1} respectively, which will allow to sum both the small and the large scales. Now the key observation is the following:

$$(4.7) \quad \sum_{k_1, m_1} \min(a_{k_1} a_{m_1} A, a_{k_1} b_{m_1} B, b_{k_1} a_{m_1} C, b_{k_1} b_{m_1} D) \leq \sum_{k_1} \min\left(a_{k_1} \sum_{m_1} \min(a_{m_1} A, b_{m_1} B), b_{k_1} \sum_{m_1} \min(a_{m_1} C, b_{m_1} D)\right).$$

We have seen before that

$$\sum_{m_1} \min(2^{a m_1} A, 2^{-b m_1} B) \leq A^{\frac{b}{a+b}} B^{\frac{a}{a+b}},$$

so we have

$$\begin{aligned}
& \sum_{k_1, m_1} \min(2^{a_1 k_1} 2^{a_2 m_1} A, 2^{a_1 k_1} 2^{-b_2 m_1} B, 2^{-b_1 k_1} 2^{a_2 m_1} C, 2^{-b_1 k_1} 2^{-b_2 m_1} D) \\
& \leq \sum_{k_1} \min(2^{a_1 k_1} A^{\frac{b_2}{a_2+b_2}} B^{\frac{a_2}{a_2+b_2}}, 2^{-b_1 k_1} C^{\frac{b_2}{a_2+b_2}} D^{\frac{a_2}{a_2+b_2}}) \\
(4.8) \quad & \leq \left(A^{\frac{b_2}{a_2+b_2}} B^{\frac{a_2}{a_2+b_2}}\right)^{\frac{b_1}{a_1+b_1}} \left(C^{\frac{b_2}{a_2+b_2}} D^{\frac{a_2}{a_2+b_2}}\right)^{\frac{a_1}{a_1+b_1}}.
\end{aligned}$$

Up to this point, the argument is similar to Lemma 4.5 in OhWu [18]. It remains however to justify that we have the correct terms and the correct powers. We have obtained that $\|I_A^{(1)} \times I_B^{(2)}\|_{L^p L^q}$ is bounded by a geometric average of

$$\begin{aligned}
& \|f_2\|_{L^{p_2} L^{q_2}}, \quad \|F_1\|_{\dot{B}_{p_1, \infty}^0} \|\dot{B}_{q_1, \infty}^0\| \|f_3\|_{\dot{B}_{p_3, \infty}^0} L^{q_3} \|D_{(2)}^{\gamma_2} F_5\|_{L^{p_5} \dot{B}_{q_5, \infty}^0}, \quad \|F_1\|_{\dot{B}_{p_1, \infty}^0} \|\dot{B}_{q_1, \infty}^{\beta_2}\| \|f_3\|_{\dot{B}_{p_3, \infty}^0} L^{q_3} \|D_{(2)}^{\gamma_2} F_5\|_{L^{p_5} \dot{B}_{q_5, \infty}^{\epsilon_2}}, \\
& \|F_4\|_{L^{p_4} L^{q_4}}, \quad \|F_1\|_{\dot{B}_{p_1, \infty}^{\beta_1}} \|\dot{B}_{q_1, \infty}^0\| \|f_3\|_{\dot{B}_{p_3, \infty}^{\epsilon_1}} L^{q_3} \|D_{(2)}^{\gamma_2} F_5\|_{L^{p_5} \dot{B}_{q_5, \infty}^0}, \quad \|F_1\|_{\dot{B}_{p_1, \infty}^{\beta_1}} \|\dot{B}_{q_1, \infty}^{\beta_2}\| \|f_3\|_{\dot{B}_{p_3, \infty}^{\epsilon_1}} L^{q_3} \|D_{(2)}^{\gamma_2} F_5\|_{L^{p_5} \dot{B}_{q_5, \infty}^{\epsilon_2}}.
\end{aligned}$$

In our case, $a_1 = \beta_1, a_2 = \beta_2, b_1 = \epsilon_1, b_2 = \epsilon_2$ and

$$\begin{aligned}
A &= \|F_1\|_{\dot{B}_{p_1, \infty}^0} \|\dot{B}_{q_1, \infty}^0\| \|f_3\|_{\dot{B}_{p_3, \infty}^0} L^{q_3} \|D_{(2)}^{\gamma_2} F_5\|_{L^{p_5} \dot{B}_{q_5, \infty}^0}, \quad B = \|F_1\|_{\dot{B}_{p_1, \infty}^0} \|\dot{B}_{q_1, \infty}^{\beta_2}\| \|f_3\|_{\dot{B}_{p_3, \infty}^0} L^{q_3} \|D_{(2)}^{\gamma_2} F_5\|_{L^{p_5} \dot{B}_{q_5, \infty}^{\epsilon_2}}, \\
C &= \|F_1\|_{\dot{B}_{p_1, \infty}^{\beta_1}} \|\dot{B}_{q_1, \infty}^0\| \|f_3\|_{\dot{B}_{p_3, \infty}^{\epsilon_1}} L^{q_3} \|D_{(2)}^{\gamma_2} F_5\|_{L^{p_5} \dot{B}_{q_5, \infty}^0}, \quad D = \|F_1\|_{\dot{B}_{p_1, \infty}^{\beta_1}} \|\dot{B}_{q_1, \infty}^{\beta_2}\| \|f_3\|_{\dot{B}_{p_3, \infty}^{\epsilon_1}} L^{q_3} \|D_{(2)}^{\gamma_2} F_5\|_{L^{p_5} \dot{B}_{q_5, \infty}^{\epsilon_2}}.
\end{aligned}$$

Now we need to understand what happens to the ϵ_1 and ϵ_2 derivatives encoded in the mixed Besov and Lebesgue norms. First, we look at the interaction between A and B ; $\|f_3\|_{\dot{B}_{p_3, \infty}^0} L^{q_3}$ remains unchanged, and similar to the one-parameter Leibniz rule from Section 2.2,

$$\begin{aligned}
& \left(\|F_1\|_{\dot{B}_{p_1, \infty}^0} \|\dot{B}_{q_1, \infty}^0\| \|D_{(2)}^{\gamma_2} F_5\|_{L^{p_5} \dot{B}_{q_5, \infty}^0}\right)^{\frac{\epsilon_2}{\beta_2+\epsilon_2}} \cdot \left(\|F_1\|_{\dot{B}_{p_1, \infty}^0} \|\dot{B}_{q_1, \infty}^{\beta_2}\| \|D_{(2)}^{\gamma_2} F_5\|_{L^{p_5} \dot{B}_{q_5, \infty}^{\epsilon_2}}\right)^{\frac{\beta_2}{\beta_2+\epsilon_2}} \\
& \lesssim \left(\|F_1\|_{\dot{B}_{p_1, \infty}^0} \|\dot{B}_{q_1, \infty}^0\| \|D_{(2)}^{\gamma_2} F_5\|_{L^{p_5} \dot{B}_{q_5, \infty}^{\beta_2}}\right)^{\frac{\epsilon_2}{\beta_2+\epsilon_2}} \cdot \left(\|F_1\|_{\dot{B}_{p_1, \infty}^0} \|\dot{B}_{q_1, \infty}^{\beta_2}\| \|D_{(2)}^{\gamma_2} F_5\|_{L^{p_5} \dot{B}_{q_5, \infty}^0}\right)^{\frac{\beta_2}{\beta_2+\epsilon_2}}.
\end{aligned}$$

For the C and D interaction, initially $\|f_3\|_{\dot{B}_{p_3, \infty}^{\epsilon_1}} L^{q_3}$ remains unchanged and the Besov norm interpolation (2.11) yields

$$\begin{aligned}
& \left(\|F_1\|_{\dot{B}_{p_1, \infty}^{\beta_1}} \|\dot{B}_{q_1, \infty}^0\| \|D_{(2)}^{\gamma_2} F_5\|_{L^{p_5} \dot{B}_{q_5, \infty}^0}\right)^{\frac{\epsilon_2}{\beta_2+\epsilon_2}} \cdot \left(\|F_1\|_{\dot{B}_{p_1, \infty}^{\beta_1}} \|\dot{B}_{q_1, \infty}^{\beta_2}\| \|D_{(2)}^{\gamma_2} F_5\|_{L^{p_5} \dot{B}_{q_5, \infty}^{\epsilon_2}}\right)^{\frac{\beta_2}{\beta_2+\epsilon_2}} \\
& \lesssim \left(\|F_1\|_{\dot{B}_{p_1, \infty}^{\beta_1}} \|\dot{B}_{q_1, \infty}^0\| \|D_{(2)}^{\gamma_2} F_5\|_{L^{p_5} \dot{B}_{q_5, \infty}^{\beta_2}}\right)^{\frac{\epsilon_2}{\beta_2+\epsilon_2}} \cdot \left(\|F_1\|_{\dot{B}_{p_1, \infty}^{\beta_1}} \|\dot{B}_{q_1, \infty}^{\beta_2}\| \|D_{(2)}^{\gamma_2} F_5\|_{L^{p_5} \dot{B}_{q_5, \infty}^0}\right)^{\frac{\beta_2}{\beta_2+\epsilon_2}}.
\end{aligned}$$

So (4.8), for our particular choice of A, \dots, D , a_1, \dots, b_2 , is bounded above by

$$\begin{aligned}
& \left(\left(\|f_3\|_{\dot{B}_{p_3, \infty}^0} L^{q_3} \|F_1\|_{\dot{B}_{p_1, \infty}^0} \|\dot{B}_{q_1, \infty}^0\| \|D_{(2)}^{\gamma_2} F_5\|_{L^{p_5} \dot{B}_{q_5, \infty}^0}\right)^{\frac{\epsilon_2}{\beta_2+\epsilon_2}} \cdot \left(\|f_3\|_{\dot{B}_{p_3, \infty}^0} L^{q_3} \|F_1\|_{\dot{B}_{p_1, \infty}^0} \|\dot{B}_{q_1, \infty}^{\beta_2}\| \|D_{(2)}^{\gamma_2} F_5\|_{L^{p_5} \dot{B}_{q_5, \infty}^0}\right)^{\frac{\beta_2}{\beta_2+\epsilon_2}}\right)^{\frac{\epsilon_1}{\beta_1+\epsilon_1}} \\
& \cdot \left(\left(\|f_3\|_{\dot{B}_{p_3, \infty}^{\epsilon_1}} L^{q_3} \|F_1\|_{\dot{B}_{p_1, \infty}^{\beta_1}} \|\dot{B}_{q_1, \infty}^0\| \|D_{(2)}^{\gamma_2} F_5\|_{L^{p_5} \dot{B}_{q_5, \infty}^{\beta_2}}\right)^{\frac{\epsilon_2}{\beta_2+\epsilon_2}} \cdot \left(\|f_3\|_{\dot{B}_{p_3, \infty}^{\epsilon_1}} L^{q_3} \|F_1\|_{\dot{B}_{p_1, \infty}^{\beta_1}} \|\dot{B}_{q_1, \infty}^{\beta_2}\| \|D_{(2)}^{\gamma_2} F_5\|_{L^{p_5} \dot{B}_{q_5, \infty}^0}\right)^{\frac{\beta_2}{\beta_2+\epsilon_2}}\right)^{\frac{\beta_1}{\beta_1+\epsilon_1}}
\end{aligned}$$

Notice that we removed the ϵ_2 parameter from the Besov norms in the second variable, and that the derivatives have been redistributed thanks to the interpolation result (2.10). Now we regroup the terms, and the expression above becomes

$$\begin{aligned} & \left((\|f_3\|_{\dot{B}_{p_3,\infty}^0 L^{q_3}} \|F_1\|_{\dot{B}_{p_1,\infty}^0 \dot{B}_{q_1,\infty}^0} \|D_{(2)}^{\gamma_2} F_5\|_{L^{p_5} \dot{B}_{q_5,\infty}^{\beta_2}})^{\frac{\epsilon_1}{\beta_1+\epsilon_1}} \cdot (\|f_3\|_{\dot{B}_{p_3,\infty}^{\epsilon_1} L^{q_3}} \|F_1\|_{\dot{B}_{p_1,\infty}^{\beta_1} \dot{B}_{q_1,\infty}^0} \|D_{(2)}^{\gamma_2} F_5\|_{L^{p_5} \dot{B}_{q_5,\infty}^{\beta_2}})^{\frac{\beta_1}{\beta_1+\epsilon_1}} \right)^{\frac{\epsilon_2}{\beta_2+\epsilon_2}} \\ & \cdot \left((\|f_3\|_{\dot{B}_{p_3,\infty}^{\epsilon_1} L^{q_3}} \|F_1\|_{\dot{B}_{p_1,\infty}^{\beta_1} \dot{B}_{q_1,\infty}^0} \|D_{(2)}^{\gamma_2} F_5\|_{L^{p_5} \dot{B}_{q_5,\infty}^{\beta_2}})^{\frac{\epsilon_1}{\beta_1+\epsilon_1}} \cdot (\|f_3\|_{\dot{B}_{p_3,\infty}^{\epsilon_1} L^{q_3}} \|F_1\|_{\dot{B}_{p_1,\infty}^{\beta_1} \dot{B}_{q_1,\infty}^0} \|D_{(2)}^{\gamma_2} F_5\|_{L^{p_5} \dot{B}_{q_5,\infty}^0})^{\frac{\beta_1}{\beta_1+\epsilon_1}} \right)^{\frac{\beta_2}{\beta_2+\epsilon_2}}. \end{aligned}$$

On the first line, $\|D_{(2)}^{\gamma_2} F_5\|_{L^{p_5} \dot{B}_{q_5,\infty}^{\beta_2}}$ remains unchanged and

$$\begin{aligned} & (\|f_3\|_{\dot{B}_{p_3,\infty}^0 L^{q_3}} \|F_1\|_{\dot{B}_{p_1,\infty}^0 \dot{B}_{q_1,\infty}^0})^{\frac{\epsilon_1}{\beta_1+\epsilon_1}} \cdot (\|f_3\|_{\dot{B}_{p_3,\infty}^{\epsilon_1} L^{q_3}} \|F_1\|_{\dot{B}_{p_1,\infty}^{\beta_1} \dot{B}_{q_1,\infty}^0})^{\frac{\beta_1}{\beta_1+\epsilon_1}} \\ & \lesssim (\|f_3\|_{\dot{B}_{p_3,\infty}^{\beta_1} L^{q_3}} \|F_1\|_{\dot{B}_{p_1,\infty}^0 \dot{B}_{q_1,\infty}^0})^{\frac{\epsilon_1}{\beta_1+\epsilon_1}} \cdot (\|f_3\|_{\dot{B}_{p_3,\infty}^0 L^{q_3}} \|F_1\|_{\dot{B}_{p_1,\infty}^{\beta_1} \dot{B}_{q_1,\infty}^0})^{\frac{\beta_1}{\beta_1+\epsilon_1}}. \end{aligned}$$

The second line can be estimated similarly; thus (4.8) is bounded above by

$$\begin{aligned} & \left((\|f_3\|_{\dot{B}_{p_3,\infty}^{\beta_1} L^{q_3}} \|F_1\|_{\dot{B}_{p_1,\infty}^0 \dot{B}_{q_1,\infty}^0} \|D_{(2)}^{\gamma_2} F_5\|_{L^{p_5} \dot{B}_{q_5,\infty}^{\beta_2}})^{\frac{\epsilon_1}{\beta_1+\epsilon_1}} \cdot (\|f_3\|_{\dot{B}_{p_3,\infty}^0 L^{q_3}} \|F_1\|_{\dot{B}_{p_1,\infty}^{\beta_1} \dot{B}_{q_1,\infty}^0} \|D_{(2)}^{\gamma_2} F_5\|_{L^{p_5} \dot{B}_{q_5,\infty}^{\beta_2}})^{\frac{\beta_1}{\beta_1+\epsilon_1}} \right)^{\frac{\epsilon_2}{\beta_2+\epsilon_2}} \\ & \cdot \left((\|f_3\|_{\dot{B}_{p_3,\infty}^0 L^{q_3}} \|F_1\|_{\dot{B}_{p_1,\infty}^{\beta_1} \dot{B}_{q_1,\infty}^0} \|D_{(2)}^{\gamma_2} F_5\|_{L^{p_5} \dot{B}_{q_5,\infty}^{\beta_2}})^{\frac{\epsilon_1}{\beta_1+\epsilon_1}} \cdot (\|f_3\|_{\dot{B}_{p_3,\infty}^0 L^{q_3}} \|F_1\|_{\dot{B}_{p_1,\infty}^{\beta_1} \dot{B}_{q_1,\infty}^0} \|D_{(2)}^{\gamma_2} F_5\|_{L^{p_5} \dot{B}_{q_5,\infty}^0})^{\frac{\beta_1}{\beta_1+\epsilon_1}} \right)^{\frac{\beta_2}{\beta_2+\epsilon_2}} \end{aligned}$$

which can be further estimated by

$$\begin{aligned} & \|f_3\|_{\dot{B}_{p_3,\infty}^{\beta_1} L^{q_3}} \|F_1\|_{\dot{B}_{p_1,\infty}^0 \dot{B}_{q_1,\infty}^0} \|D_{(2)}^{\gamma_2} F_5\|_{L^{p_5} \dot{B}_{q_5,\infty}^{\beta_2}} + \|f_3\|_{\dot{B}_{p_3,\infty}^0 L^{q_3}} \|F_1\|_{\dot{B}_{p_1,\infty}^{\beta_1} \dot{B}_{q_1,\infty}^0} \|D_{(2)}^{\gamma_2} F_5\|_{L^{p_5} \dot{B}_{q_5,\infty}^{\beta_2}} \\ & + \|f_3\|_{\dot{B}_{p_3,\infty}^0 L^{q_3}} \|F_1\|_{\dot{B}_{p_1,\infty}^{\beta_1} \dot{B}_{q_1,\infty}^0} \|D_{(2)}^{\gamma_2} F_5\|_{L^{p_5} \dot{B}_{q_5,\infty}^{\beta_2}} + \|f_3\|_{\dot{B}_{p_3,\infty}^0 L^{q_3}} \|F_1\|_{\dot{B}_{p_1,\infty}^{\beta_1} \dot{B}_{q_1,\infty}^0} \|D_{(2)}^{\gamma_2} F_5\|_{L^{p_5} \dot{B}_{q_5,\infty}^0}. \end{aligned}$$

- $I_A^{(1)} \times I_A^{(2)}$ This case is simpler than the previous one, which is why we will only briefly present the arguments. The multiplier $m_{I_A^{(1)}}(\xi_1, \dots, \xi_5) \cdot m_{I_A^{(2)}}(\eta_1, \dots, \eta_5)$ is given by

$$\left(\frac{|\xi_1 + \dots + \xi_5|^{\beta_1} - |\xi_1 + \xi_2|^{\beta_1}}{\xi_3 + \xi_4 + \xi_5} \cdot \xi_3 \cdot |\xi_1 + \xi_2|^{\alpha_1} |\xi_4 + \xi_5|^{\gamma_1} \right) \cdot \left(\frac{|\eta_1 + \dots + \eta_5|^{\beta_2} - |\eta_1 + \eta_2|^{\beta_2}}{\eta_3 + \eta_4 + \eta_5} \cdot \eta_3 \cdot |\eta_1 + \eta_2|^{\alpha_2} |\eta_4 + \eta_5|^{\gamma_2} \right).$$

After summing over $k_2, k_4, k_5 \ll k_1$ and $m_2, m_4, m_5 \ll m_1$, the associated operator from (4.3) becomes

$$\begin{aligned} & T_{m_{I_A^{(1)}}, m_{I_A^{(2)}}}^{k_1, k_3; m_1, m_3}(f_1, \dots, f_5)(x, y) = \int_{\mathbb{R}^{10}} m_{I_A^{(1)}}(\xi_1, \dots, \xi_5) \cdot m_{I_A^{(2)}}(\eta_1, \dots, \eta_5) \mathcal{F}(\Delta_{k_1}^{(1)} \Delta_{m_1}^{(2)} f_1)(\xi_1, \eta_1) \cdot \mathcal{F}(S_{k_1}^{(1)} S_{m_1}^{(2)} f_2)(\xi_2, \eta_2) \\ & \cdot \mathcal{F}(\Delta_{k_3}^{(1)} \Delta_{m_3}^{(2)} f_3)(\xi_3, \eta_3) \cdot \mathcal{F}(S_{k_1}^{(1)} S_{m_1}^{(2)} f_4)(\xi_4, \eta_4) \cdot \mathcal{F}(S_{k_1}^{(1)} S_{m_1}^{(2)} f_5)(\xi_5, \eta_5) \cdot e^{2\pi i x(\xi_1 + \dots + \xi_5)} e^{2\pi i y(\eta_1 + \dots + \eta_5)} d\xi_1 \dots d\xi_5 d\eta_1 \dots d\eta_5. \end{aligned}$$

Notice that the operator is symmetric in the first and second parameter, in the sense that the conical frequency regions are described by similar inequalities and the multipliers are similar. We continue with the usual Fourier series decomposition, which allows to regard $T_{m_{I_A^{(1)}}, m_{I_A^{(2)}}}(f_1, \dots, f_5)$ as a superposition of (modulated) terms of the form

$$\begin{aligned} & 2^{k_1(\beta_1-1)} 2^{k_3} 2^{m_1(\beta_2-1)} 2^{m_3} (\Delta_{k_1}^{(1)} \Delta_{m_1}^{(2)} D_{(1)}^{\alpha_1} D_{(2)}^{\alpha_2} f_1)(x, y) \cdot S_{k_1}^{(1)} S_{m_1}^{(2)} f_2(x, y) \\ & \cdot \Delta_{k_3}^{(1)} \Delta_{m_3}^{(2)} f_3(x, y) \cdot D_{(1)}^{\gamma_1} D_{(2)}^{\gamma_2} (S_{k_1}^{(1)} S_{m_1}^{(2)} f_4 \cdot S_{k_1}^{(1)} S_{m_1}^{(2)} f_5)(x, y). \end{aligned}$$

When each of these terms is estimated in $\|\cdot\|_{L^p L^q}^\tau$, they are bounded above by

$$\begin{aligned} & 2^{k_1(\beta_1-1)\tau} 2^{k_3\tau} 2^{m_1(\beta_2-1)\tau} 2^{m_3\tau} \|\Delta_{k_1}^{(1)} \Delta_{m_1}^{(2)} D_{(1)}^{\alpha_1} D_{(2)}^{\alpha_2} f_1\|_{L^{p_1} L^{q_1}}^\tau \|S_{k_1}^{(1)} S_{m_1}^{(2)} f_2\|_{L^{p_2} L^{q_2}}^\tau \\ & \|\Delta_{k_3}^{(1)} \Delta_{m_3}^{(2)} f_3\|_{L^{p_3} L^{q_3}}^\tau \|D_{(1)}^{\gamma_1} D_{(2)}^{\gamma_2} (S_{k_1}^{(1)} S_{m_1}^{(2)} f_4 \cdot S_{k_1}^{(1)} S_{m_1}^{(2)} f_5)\|_{L^{p_{4,5}} L^{q_{4,5}}}^\tau \\ & \lesssim 2^{k_1(\beta_1-1)\tau} 2^{k_3\tau} 2^{m_1(\beta_2-1)\tau} 2^{m_3\tau} \|\Delta_{k_1}^{(1)} \Delta_{m_1}^{(2)} D_{(1)}^{\alpha_1} D_{(2)}^{\alpha_2} f_1\|_{L^{p_1} L^{q_1}}^\tau \|\Delta_{k_3}^{(1)} \Delta_{m_3}^{(2)} f_3\|_{L^{p_3} L^{q_3}}^\tau \\ & \|f_2\|_{L^{p_2} L^{q_2}}^\tau (\|D_{(1)}^{\gamma_1} D_{(2)}^{\gamma_2} f_4\|_{L^{p_4} L^{q_4}} \|f_5\|_{L^{p_5} L^{q_5}} + \dots + \|f_4\|_{L^{p_4} L^{q_4}} \|D_{(1)}^{\gamma_1} D_{(2)}^{\gamma_2} f_5\|_{L^{p_5} L^{q_5}})^\tau \end{aligned}$$

The summation over $k_3 < k_1, m_3 < m_1$ will only affect the first line in the term above, and it can be estimated by

$$\sum_{k_1, m_1} \min(2^{k_1\beta_1\tau} 2^{m_1\beta_2\tau} \|D_{(1)}^{\alpha_1} D_{(2)}^{\alpha_2} f_1\|_{\dot{B}_{p_1,\infty}^0 \dot{B}_{q_1,\infty}^0}^\tau \|f_3\|_{\dot{B}_{p_3,\infty}^0 \dot{B}_{q_3,\infty}^0}^\tau, 2^{k_1\beta_1\tau} 2^{-m_1\epsilon_2\tau} \|D_{(1)}^{\alpha_1} D_{(2)}^{\alpha_2} f_1\|_{\dot{B}_{p_1,\infty}^0 \dot{B}_{q_1,\infty}^{\beta_2}}^\tau \|f_3\|_{\dot{B}_{p_3,\infty}^0 \dot{B}_{q_3,\infty}^{\epsilon_2}}^\tau),$$

$$2^{-k_1\epsilon_1\tau}2^{m_1\beta_2\tau}\|D_{(1)}^{\alpha_1}D_{(2)}^{\alpha_2}f_1\|_{\dot{B}_{p_1,\infty}^{\beta_1}\dot{B}_{q_1,\infty}^0}^\tau\|f_3\|_{\dot{B}_{p_3,\infty}^{\epsilon_1}\dot{B}_{q_3,\infty}^0}^\tau, 2^{-k_1\epsilon_1\tau}2^{-m_1\epsilon_2\tau}\|D_{(1)}^{\alpha_1}D_{(2)}^{\alpha_2}f_1\|_{\dot{B}_{p_1,\infty}^{\beta_1}\dot{B}_{q_1,\infty}^0}^\tau\|f_3\|_{\dot{B}_{p_3,\infty}^{\epsilon_1}\dot{B}_{q_3,\infty}^0}^\tau).$$

The procedure described in treating $I_A^{(1)} \times I_B^{(2)}$ will eventually yield that this is majorized by

$$\begin{aligned} & \left(\|D_{(1)}^{\alpha_1}D_{(2)}^{\alpha_2}f_1\|_{\dot{B}_{p_1,\infty}^0\dot{B}_{q_1,\infty}^0}^\tau\|f_3\|_{\dot{B}_{p_3,\infty}^{\beta_1}\dot{B}_{q_3,\infty}^{\beta_2}}^\tau\right)^{\frac{\epsilon_1}{\beta_1+\epsilon_1}\cdot\frac{\epsilon_2}{\beta_2+\epsilon_2}} \cdot \left(\|D_{(1)}^{\alpha_1}D_{(2)}^{\alpha_2}f_1\|_{\dot{B}_{p_1,\infty}^{\beta_1}\dot{B}_{q_1,\infty}^0}^\tau\|f_3\|_{\dot{B}_{p_3,\infty}^0\dot{B}_{q_3,\infty}^{\beta_2}}^\tau\right)^{\frac{\beta_1}{\beta_1+\epsilon_1}\cdot\frac{\epsilon_2}{\beta_2+\epsilon_2}} \\ & \cdot \left(\|D_{(1)}^{\alpha_1}D_{(2)}^{\alpha_2}f_1\|_{\dot{B}_{p_1,\infty}^0\dot{B}_{q_1,\infty}^{\beta_2}}^\tau\|f_3\|_{\dot{B}_{p_3,\infty}^{\beta_1}\dot{B}_{q_3,\infty}^0}^\tau\right)^{\frac{\epsilon_1}{\beta_1+\epsilon_1}\cdot\frac{\beta_2}{\beta_2+\epsilon_2}} \cdot \left(\|D_{(1)}^{\alpha_1}D_{(2)}^{\alpha_2}f_1\|_{\dot{B}_{p_1,\infty}^{\beta_1}\dot{B}_{q_1,\infty}^{\beta_2}}^\tau\|f_3\|_{\dot{B}_{p_3,\infty}^0\dot{B}_{q_3,\infty}^0}^\tau\right)^{\frac{\beta_1}{\beta_1+\epsilon_1}\cdot\frac{\beta_2}{\beta_2+\epsilon_2}}. \end{aligned}$$

- $I_A^{(1)} \times I_C^{(2)}$ We recall that the symbol $m_{I_A^{(1)}}(\xi_1, \dots, \xi_5) \cdot m_{I_C^{(2)}}(\eta_1, \dots, \eta_5)$ is

$$\left(\frac{|\xi_1 + \dots + \xi_5|^{\beta_1} - |\xi_1 + \xi_2|^{\beta_1}}{\xi_3 + \xi_4 + \xi_5} \cdot \xi_3 \cdot |\xi_1 + \xi_2|^{\alpha_1} |\xi_4 + \xi_5|^{\gamma_1}\right) \cdot \left(\frac{|\eta_1 + \eta_2|^{\alpha_2+\beta_2} - |\eta_1|^{\alpha_2+\beta_2}}{\eta_2} \cdot \eta_2 \cdot |\eta_4 + \eta_5|^{\gamma_2}\right)$$

and thus the associated operator, obtained after a suitable regrouping of the terms, is

$$\begin{aligned} T_{m_{I_A^{(1)}}, m_{I_C^{(2)}}}(f_1, \dots, f_5)(x, y) &= \int_{\mathbb{R}^{10}} m_{I_A^{(1)}}(\xi_1, \dots, \xi_5) \cdot m_{I_C^{(2)}}(\eta_1, \dots, \eta_5) \mathcal{F}(\Delta_{k_1}^{(1)} \Delta_{m_1}^{(2)} f_1)(\xi_1, \eta_1) \cdot \mathcal{F}(S_{k_1}^{(1)} \Delta_{m_2}^{(2)} f_2)(\xi_2, \eta_2) \\ &\cdot \mathcal{F}(\Delta_{k_3}^{(1)} S_{m_1}^{(2)} f_3)(\xi_3, \eta_3) \cdot \mathcal{F}(S_{k_1}^{(1)} S_{m_1}^{(2)} f_4)(\xi_4, \eta_4) \cdot \mathcal{F}(S_{k_1}^{(1)} S_{m_1}^{(2)} f_5)(\xi_5, \eta_5) \cdot e^{2\pi i x(\xi_1 + \dots + \xi_5)} e^{2\pi i y(\eta_1 + \dots + \eta_5)} d\xi_1 \dots d\xi_5 d\eta_1 \dots d\eta_5. \end{aligned}$$

Thanks to the usual Fourier series decomposition, the commutator estimates and result on lower complexity flags localized in frequency, we are left with summing

$$\begin{aligned} & \sum_{k_3 < k_1} \sum_{m_2 < m_1} 2^{k_1(\beta_1-1)\tau} 2^{k_3\tau} 2^{k_1\alpha_1\tau} 2^{m_1(\alpha_2+\beta_2-1)\tau} 2^{m_2\tau} \|\Delta_{k_1}^{(1)} \Delta_{m_1}^{(2)} f_1\|_{L^{p_1} L^{q_1}}^\tau \|S_{k_1}^{(1)} \Delta_{m_2}^{(2)} f_2\|_{L^{p_2} L^{q_2}}^\tau \\ & \|\Delta_{k_3}^{(1)} S_{m_1}^{(2)} f_3\|_{L^{p_3} L^{q_3}}^\tau \|D_{(1)}^{\gamma_1} D_{(2)}^{\gamma_2} (S_{k_1}^{(1)} S_{m_1}^{(2)} f_4 \cdot S_{k_1}^{(1)} S_{m_1}^{(2)} f_5)\|_{L^{p_4,5} L^{q_4,5}}^\tau \\ & \lesssim \sum_{k_3 < k_1} \sum_{m_2 < m_1} 2^{k_1(\beta_1-1)\tau} 2^{k_3\tau} 2^{m_1(\alpha_2+\beta_2-1)\tau} 2^{m_2\tau} \|\Delta_{k_1}^{(1)} \Delta_{m_1}^{(2)} D_{(1)}^{\alpha_1} f_1\|_{L^{p_1} L^{q_1}}^\tau \|\Delta_{m_2}^{(2)} f_2\|_{L^{p_2} L^{q_2}}^\tau \|\Delta_{k_3}^{(1)} f_3\|_{L^{p_3} L^{q_3}}^\tau \\ & \cdot \left(\|D_{(1)}^{\gamma_1} D_{(2)}^{\gamma_2} f_4\|_{L^{p_4} L^{q_4}} + \|f_5\|_{L^{p_5} L^{q_5}} + \dots + \|f_4\|_{L^{p_4} L^{q_4}} \|D_{(1)}^{\gamma_1} D_{(2)}^{\gamma_2} f_5\|_{L^{p_5} L^{q_5}}\right)^\tau. \end{aligned}$$

Setting aside the term involving the $D_{(1)}^{\gamma_1} D_{(2)}^{\gamma_2}$ derivatives applied to (projections of) the functions f_4 and f_5 , the term above is bounded by

$$\begin{aligned} & \sum_{k_1, m_1} \min \left(2^{k_1\beta_1\tau} 2^{m_1(\alpha_2+\beta_2)\tau} \|D_{(1)}^{\alpha_1} f_1\|_{\dot{B}_{p_1,\infty}^0 \dot{B}_{q_1,\infty}^0}^\tau \|f_2\|_{L^{p_2} \dot{B}_{q_2,\infty}^0}^\tau \|f_3\|_{\dot{B}_{p_3,\infty}^0 L^{q_3}}^\tau, \right. \\ & 2^{k_1\beta_1\tau} 2^{-m_1\epsilon_2\tau} \|D_{(1)}^{\alpha_1} f_1\|_{\dot{B}_{p_1,\infty}^0 \dot{B}_{q_1,\infty}^{\alpha_2+\beta_2}}^\tau \|f_2\|_{L^{p_2} \dot{B}_{q_2,\infty}^{\epsilon_2}}^\tau \|f_3\|_{\dot{B}_{p_3,\infty}^0 L^{q_3}}^\tau, \\ & 2^{-k_1\epsilon_1\tau} 2^{m_1(\alpha_2+\beta_2)\tau} \|D_{(1)}^{\alpha_1} f_1\|_{\dot{B}_{p_1,\infty}^{\beta_1} \dot{B}_{q_1,\infty}^0}^\tau \|f_2\|_{L^{p_2} \dot{B}_{q_2,\infty}^0}^\tau \|f_3\|_{\dot{B}_{p_3,\infty}^{\epsilon_1} L^{q_3}}^\tau, \\ & \left. 2^{-k_1\epsilon_1\tau} 2^{-m_1\epsilon_2\tau} \|D_{(1)}^{\alpha_1} f_1\|_{\dot{B}_{p_1,\infty}^{\beta_1} \dot{B}_{q_1,\infty}^{\alpha_2+\beta_2}}^\tau \|f_2\|_{L^{p_2} \dot{B}_{q_2,\infty}^{\epsilon_2}}^\tau \|f_3\|_{\dot{B}_{p_3,\infty}^{\epsilon_1} L^{q_3}}^\tau \right). \end{aligned}$$

Eventually, we get

$$\begin{aligned} & \left(\|D_{(1)}^{\alpha_1} f_1\|_{\dot{B}_{p_1,\infty}^0 \dot{B}_{q_1,\infty}^0}^\tau \|f_2\|_{L^{p_2} \dot{B}_{q_2,\infty}^{\alpha_2+\beta_2}}^\tau \|f_3\|_{\dot{B}_{p_3,\infty}^{\beta_1} L^{q_3}}^\tau\right)^{\frac{\epsilon_1}{\beta_1+\epsilon_1}\cdot\frac{\epsilon_2}{\beta_2+\epsilon_2}} \cdot \left(\|D_{(1)}^{\alpha_1} f_1\|_{\dot{B}_{p_1,\infty}^{\beta_1} \dot{B}_{q_1,\infty}^0}^\tau \|f_2\|_{L^{p_2} \dot{B}_{q_2,\infty}^{\alpha_2+\beta_2}}^\tau \|f_3\|_{\dot{B}_{p_3,\infty}^0 L^{q_3}}^\tau\right)^{\frac{\beta_1}{\beta_1+\epsilon_1}\cdot\frac{\epsilon_2}{\beta_2+\epsilon_2}} \\ & \cdot \left(\|D_{(1)}^{\alpha_1} f_1\|_{\dot{B}_{p_1,\infty}^0 \dot{B}_{q_1,\infty}^{\alpha_2+\beta_2}}^\tau \|f_2\|_{L^{p_2} \dot{B}_{q_2,\infty}^0}^\tau \|f_3\|_{\dot{B}_{p_3,\infty}^{\beta_1} L^{q_3}}^\tau\right)^{\frac{\epsilon_1}{\beta_1+\epsilon_1}\cdot\frac{\beta_2}{\beta_2+\epsilon_2}} \cdot \left(\|D_{(1)}^{\alpha_1} f_1\|_{\dot{B}_{p_1,\infty}^{\beta_1} \dot{B}_{q_1,\infty}^{\alpha_2+\beta_2}}^\tau \|f_2\|_{L^{p_2} \dot{B}_{q_2,\infty}^0}^\tau \|f_3\|_{\dot{B}_{p_3,\infty}^0 L^{q_3}}^\tau\right)^{\frac{\beta_1}{\beta_1+\epsilon_1}\cdot\frac{\beta_2}{\beta_2+\epsilon_2}}. \end{aligned}$$

- $I_A^{(1)} \times I_D^{(2)}$ The symbol $m_{I_A^{(1)}}(\xi_1, \dots, \xi_5) \cdot m_{I_D^{(2)}}(\eta_1, \dots, \eta_5)$ is

$$\left(\frac{|\xi_1 + \dots + \xi_5|^{\beta_1} - |\xi_1 + \xi_2|^{\beta_1}}{\xi_3 + \xi_4 + \xi_5} \cdot \xi_3 \cdot |\xi_1 + \xi_2|^{\alpha_1} |\xi_4 + \xi_5|^{\gamma_1}\right) \cdot (|\eta_1|^{\alpha_2+\beta_2} \cdot |\eta_4 + \eta_5|^{\gamma_2})$$

and the operator associated to it, obtained after summing in $k_2, k_4, k_5 \ll k_1$ and $m_2, m_3, m_4, m_5 \ll m_1$, is

$$\begin{aligned} T_{m_{I_A^{(1)}}, m_{I_D^{(2)}}}^{k_1, k_3; m_1}(f_1, \dots, f_5)(x, y) &= \int_{\mathbb{R}^{10}} m_{I_A^{(1)}}(\xi_1, \dots, \xi_5) \cdot m_{I_D^{(2)}}(\eta_1, \dots, \eta_5) \mathcal{F}(\Delta_{k_1}^{(1)} \Delta_{m_1}^{(2)} f_1)(\xi_1, \eta_1) \cdot \mathcal{F}(S_{k_1}^{(1)} S_{m_1}^{(2)} f_2)(\xi_2, \eta_2) \\ &\cdot \mathcal{F}(\Delta_{k_3}^{(1)} S_{m_1}^{(2)} f_3)(\xi_3, \eta_3) \cdot \mathcal{F}(S_{k_1}^{(1)} S_{m_1}^{(2)} f_4)(\xi_4, \eta_4) \cdot \mathcal{F}(S_{k_1}^{(1)} S_{m_1}^{(2)} f_5)(\xi_5, \eta_5) \cdot e^{2\pi i x(\xi_1 + \dots + \xi_5)} e^{2\pi i y(\eta_1 + \dots + \eta_5)} d\xi_1 \dots d\xi_5 d\eta_1 \dots d\eta_5. \end{aligned}$$

In the second parameter, we need to switch the order of summation: each $S_{m_1}^{(2)} f_l$ will be written as

$$S_{m_1}^{(2)} f_l = f_l - \Delta_{>m_1}^{(2)} f_l,^{29} \quad \text{for all } 2 \leq l \leq 5.$$

Then $T_{m_{I_A}^{(1)}, m_{I_D}^{(2)}}^{k_1, k_3; m_1}(f_1, \dots, f_5)$ becomes

$$\begin{aligned} (4.9) \quad & \int_{\mathbb{R}^{10}} m_{I_A}^{(1)}(\xi_1, \dots, \xi_5) \cdot m_{I_D}^{(2)}(\eta_1, \dots, \eta_5) \mathcal{F}(\Delta_{k_1}^{(1)} \Delta_{m_1}^{(2)} f_1)(\xi_1, \eta_1) \cdot \mathcal{F}(S_{k_1}^{(1)} f_2)(\xi_2, \eta_2) \\ & \cdot \mathcal{F}(\Delta_{k_3}^{(1)} f_3)(\xi_3, \eta_3) \cdot \mathcal{F}(S_{k_1}^{(1)} f_4)(\xi_4, \eta_4) \cdot \mathcal{F}(S_{k_1}^{(1)} f_5)(\xi_5, \eta_5) \cdot e^{2\pi i x(\xi_1 + \dots + \xi_5)} e^{2\pi i y(\eta_1 + \dots + \eta_5)} d\xi \\ (4.10) \quad & - \int_{\mathbb{R}^{10}} m_{I_A}^{(1)}(\xi_1, \dots, \xi_5) \cdot m_{I_D}^{(2)}(\eta_1, \dots, \eta_5) \mathcal{F}(\Delta_{k_1}^{(1)} \Delta_{m_1}^{(2)} f_1)(\xi_1, \eta_1) \cdot \mathcal{F}(S_{k_1}^{(1)} \Delta_{>m_1}^{(2)} f_2)(\xi_2, \eta_2) \\ & \cdot \mathcal{F}(\Delta_{k_3}^{(1)} f_3)(\xi_3, \eta_3) \cdot \mathcal{F}(S_{k_1}^{(1)} f_4)(\xi_4, \eta_4) \cdot \mathcal{F}(S_{k_1}^{(1)} f_5)(\xi_5, \eta_5) \cdot e^{2\pi i x(\xi_1 + \dots + \xi_5)} e^{2\pi i y(\eta_1 + \dots + \eta_5)} d\xi \\ (4.11) \quad & - \int_{\mathbb{R}^{10}} m_{I_A}^{(1)}(\xi_1, \dots, \xi_5) \cdot m_{I_D}^{(2)}(\eta_1, \dots, \eta_5) \mathcal{F}(\Delta_{k_1}^{(1)} \Delta_{m_1}^{(2)} f_1)(\xi_1, \eta_1) \cdot \mathcal{F}(S_{k_1}^{(1)} f_2)(\xi_2, \eta_2) \\ & \cdot \mathcal{F}(\Delta_{k_3}^{(1)} f_3)(\xi_3, \eta_3) \cdot \mathcal{F}(S_{k_1}^{(1)} \Delta_{>m_1}^{(2)} f_4)(\xi_4, \eta_4) \cdot \mathcal{F}(S_{k_1}^{(1)} f_5)(\xi_5, \eta_5) \cdot e^{2\pi i x(\xi_1 + \dots + \xi_5)} e^{2\pi i y(\eta_1 + \dots + \eta_5)} d\xi \\ & + \text{similar terms.} \end{aligned}$$

If we sum in m_1 , the first term becomes

$$\begin{aligned} & \int_{\mathbb{R}^{10}} m_{I_A}^{(1)}(\xi_1, \dots, \xi_5) \cdot |\eta_4 + \eta_5|^{\gamma_2} \mathcal{F}(\Delta_{k_1}^{(1)} D_{(2)}^{\alpha_2 + \beta_2} f_1)(\xi_1, \eta_1) \cdot \mathcal{F}(S_{k_1}^{(1)} f_2)(\xi_2, \eta_2) \cdot \mathcal{F}(\Delta_{k_3}^{(1)} f_3)(\xi_3, \eta_3) \\ & \cdot \mathcal{F}(S_{k_1}^{(1)} f_4)(\xi_4, \eta_4) \cdot \mathcal{F}(S_{k_1}^{(1)} f_5)(\xi_5, \eta_5) \cdot e^{2\pi i x(\xi_1 + \dots + \xi_5)} e^{2\pi i y(\eta_1 + \dots + \eta_5)} d\xi. \end{aligned}$$

So besides the $D_{(2)}^{\gamma_2}$ derivatives acting on f_4 and f_5 (thus an object of lower complexity), we have fundamentally a one-parameter Leibniz rule. We use Fourier series for decomposing the symbol $m_{I_A}^{(1)}(\xi_1, \dots, \xi_5)$, and the boundedness of (4.9) in $\|\cdot\|_{L^p L^q}^\tau$ is a consequence of

$$\begin{aligned} & \sum_{k_3 < k_1} 2^{k_1(\beta_1 - 1)\tau} 2^{k_3\tau} 2^{k_1\alpha_1\tau} \|\Delta_{k_1}^{(1)} D_{(2)}^{\alpha_2 + \beta_2} f_1\|_{L^{p_1} L^{q_1}}^\tau \|S_{k_1}^{(1)} f_2\|_{L^{p_2} L^{q_2}}^\tau \|\Delta_{k_3}^{(1)} f_3\|_{L^{p_3} L^{q_3}}^\tau \|D_{(1)}^{\gamma_1} D_{(2)}^{\gamma_2} (S_{k_1}^{(1)} f_4 \cdot S_{k_1}^{(1)} f_5)\|_{L^{p_4,5} L^{q_4,5}}^\tau \\ & \lesssim \sum_{k_3 < k_1} 2^{k_1(\beta_1 - 1)\tau} 2^{k_3\tau} \|\Delta_{k_1}^{(1)} D_{(1)}^{\alpha_1} D_{(2)}^{\alpha_2 + \beta_2} f_1\|_{L^{p_1} L^{q_1}}^\tau \|f_2\|_{L^{p_2} L^{q_2}}^\tau \|\Delta_{k_3}^{(1)} f_3\|_{L^{p_3} L^{q_3}}^\tau \\ & \cdot \left(\|D_{(1)}^{\gamma_1} D_{(2)}^{\gamma_2} f_4\|_{L^{p_4} L^{q_4}}^\tau \|f_5\|_{L^{p_5} L^{q_5}}^\tau + \dots + \|f_4\|_{L^{p_4} L^{q_4}}^\tau \|D_{(1)}^{\gamma_1} D_{(2)}^{\gamma_2} f_5\|_{L^{p_5} L^{q_5}}^\tau \right)^\tau \end{aligned}$$

Only the functions f_1 and f_3 will participate in the summation, and the optimization resembles the one-parameter case discussed in Section 2.2:

$$\begin{aligned} & \sum_{k_1} \min(2^{k_1\beta_1\tau} \|D_{(1)}^{\alpha_1} D_{(2)}^{\alpha_2 + \beta_2} f_1\|_{\dot{B}_{p_1, \infty}^{0, L^{q_1}}}^\tau \|f_3\|_{\dot{B}_{p_3, \infty}^{0, L^{q_3}}}^\tau, 2^{-k_1\epsilon_1\tau} \|D_{(1)}^{\alpha_1} D_{(2)}^{\alpha_2 + \beta_2} f_1\|_{\dot{B}_{p_1, \infty}^{\beta_1, L^{q_1}}}^\tau \|f_3\|_{\dot{B}_{p_3, \infty}^{\epsilon_1, L^{q_3}}}^\tau) \\ & \lesssim (\|D_{(1)}^{\alpha_1} D_{(2)}^{\alpha_2 + \beta_2} f_1\|_{\dot{B}_{p_1, \infty}^{0, L^{q_1}}}^\tau \|f_3\|_{\dot{B}_{p_3, \infty}^{\beta_1, L^{q_3}}}^\tau)^{\frac{\epsilon_1}{\beta_1 + \epsilon_1}} \cdot (\|D_{(1)}^{\alpha_1} D_{(2)}^{\alpha_2 + \beta_2} f_1\|_{\dot{B}_{p_1, \infty}^{\beta_1, L^{q_1}}}^\tau \|f_3\|_{\dot{B}_{p_3, \infty}^{0, L^{q_3}}}^\tau)^{\frac{\beta_1}{\beta_1 + \epsilon_1}}. \end{aligned}$$

We return to (4.10), for which the summation in m_1 is performed outside the $\|\cdot\|_{L^p L^q}^\tau$ quasi-norms:

$$\begin{aligned} & \sum_{k_3 < k_1} \sum_{m_1} 2^{k_1(\beta_1 - 1)\tau} 2^{k_3\tau} 2^{k_1\alpha_1\tau} 2^{m_1(\alpha_2 + \beta_2)\tau} \|\Delta_{k_1}^{(1)} \Delta_{m_1}^{(2)} f_1\|_{L^{p_1} L^{q_1}}^\tau \|S_{k_1}^{(1)} \Delta_{>m_1}^{(2)} f_2\|_{L^{p_2} L^{q_2}}^\tau \\ & \quad \|\Delta_{k_3}^{(1)} f_3\|_{L^{p_3} L^{q_3}}^\tau \|D_{(1)}^{\gamma_1} D_{(2)}^{\gamma_2} (S_{k_1}^{(1)} f_4 \cdot S_{k_1}^{(1)} f_5)\|_{L^{p_4,5} L^{q_4,5}}^\tau \\ & \lesssim \sum_{k_3 < k_1} \sum_{m_1} 2^{k_1(\beta_1 - 1)\tau} 2^{k_3\tau} 2^{m_1(\alpha_2 + \beta_2)\tau} \|\Delta_{k_1}^{(1)} D_{(1)}^{\alpha_1} D_{(2)}^{\alpha_2 + \beta_2} f_1\|_{L^{p_1} L^{q_1}}^\tau \|\Delta_{>m_1}^{(2)} f_2\|_{L^{p_2} L^{q_2}}^\tau \|\Delta_{k_3}^{(1)} f_3\|_{L^{p_3} L^{q_3}}^\tau \\ & \quad \cdot \left(\|D_{(1)}^{\gamma_1} D_{(2)}^{\gamma_2} f_4\|_{L^{p_4} L^{q_4}}^\tau \|f_5\|_{L^{p_5} L^{q_5}}^\tau + \dots + \|f_4\|_{L^{p_4} L^{q_4}}^\tau \|D_{(1)}^{\gamma_1} D_{(2)}^{\gamma_2} f_5\|_{L^{p_5} L^{q_5}}^\tau \right)^\tau. \end{aligned}$$

Using the observation that

$$\|\Delta_{>m_1}^{(2)} f_2\|_{L^{p_2} L^{q_2}} \leq \min(\|f_2\|_{L^{p_2} L^{q_2}}, 2^{-m_1\epsilon} \|f_2\|_{L^{p_2} \dot{B}_{q_2, \infty}^\epsilon}),$$

we notice that the term on the first line is majorized by

$$\sum_{k_1, m_1} \min(2^{k_1\beta_1\tau} 2^{m_1(\alpha_2 + \beta_2)\tau} \|f_1\|_{\dot{B}_{p_1, \infty}^{0, L^{q_1}}}^\tau \|f_2\|_{L^{p_2} L^{q_2}}^\tau \|f_3\|_{\dot{B}_{p_3, \infty}^{0, L^{q_3}}}^\tau, 2^{k_1\beta_1\tau} 2^{-m_1\epsilon_2\tau} \|f_1\|_{\dot{B}_{p_1, \infty}^{\beta_1, L^{q_1}}}^\tau \|f_2\|_{L^{p_2} \dot{B}_{q_2, \infty}^{\epsilon_2}}^\tau \|f_3\|_{\dot{B}_{p_3, \infty}^{0, L^{q_3}}}^\tau),$$

²⁹As in the one-parameter setting, $\Delta_{>m}^{(j)} f_l := \sum_{\tilde{m} > m-3} \Delta_{\tilde{m}}^{(j)} f_l$, and has the same properties as $\Delta_{>m} f_l$ defined in (2.17).

$$2^{-k_1\epsilon_1\tau} 2^{m_1(\alpha_2+\beta_2)\tau} \|f_1\|_{\dot{B}_{p_1,\infty}^{\beta_1}, \dot{B}_{q_1,\infty}^0}^\tau \|f_2\|_{\dot{L}^{p_2} L^{q_2}}^\tau \|f_3\|_{\dot{B}_{p_3,\infty}^{\epsilon_1}, L^{q_3}}^\tau, 2^{-k_1\epsilon_1\tau} 2^{-m_1\epsilon_2\tau} \|f_1\|_{\dot{B}_{p_1,\infty}^{\beta_1}, \dot{B}_{q_1,\infty}^{\alpha_2+\beta_2}}^\tau \|f_2\|_{\dot{L}^{p_2} \dot{B}_{q_2,\infty}^{\epsilon_2}}^\tau \|f_3\|_{\dot{B}_{p_3,\infty}^{\epsilon_1}, L^{q_3}}^\tau).$$

This will eventually produce the expected term – the computations follow the usual pattern.

Finally, we take a quick look at (4.11) as well, since the structure of the subtree associated to $D_{(2)}^{\gamma_2}(f_4 \cdot f_5)$ becomes a part of the analysis. Without loss of generality (since the other case is similar), we further restrict the symbol to the conical region $|\eta_4| \leq |\eta_5|$. Using the usual Fourier series expansion in the first variable in order to tensorize the symbol, we are led ultimately to estimating³⁰

$$\sum_{\substack{k_3 \ll k_1 \\ m_1, m_5}} 2^{(\beta_1-1)k_1\tau} 2^{k_3\tau} 2^{m_1(\alpha_2+\beta_2)\tau} \|\Delta_{k_1}^{(1)} \Delta_{m_1}^{(2)} D_{(1)}^{\alpha_1} f_1\|_{\dot{L}^{p_1} L^{q_1}}^\tau \|f_2\|_{\dot{L}^{p_2} L^{q_2}}^\tau \|\Delta_{k_3}^{(1)} f_3\|_{\dot{L}^{p_3} L^{q_3}}^\tau \|D_{\gamma_2}^{(2)} \left(S_{k_1}^{(1)} \Delta_{m_1}^{(2)} \Delta_{m_5}^{(2)} D_{(1)}^{\gamma_1} f_4 S_{k_1}^{(1)} \Delta_{m_5}^{(2)} f_5 \right)\|_{\dot{L}^{p_4,5} L^{q_4,5}}^\tau.$$

The only way $\Delta_{m_1}^{(2)} S_{k_5}^{(2)}$ is non-zero is if $m_5 > m_1$ so that we can restrict the above expression to the sum over $m_5 > m_1$. We also invoke the localized Leibniz rule:

$$\|D_{\gamma_2}^{(2)} \left(S_{k_1}^{(1)} \Delta_{m_1}^{(2)} \Delta_{m_5}^{(2)} D_{(1)}^{\gamma_1} f_4 \cdot S_{k_1}^{(1)} \Delta_{m_5}^{(2)} f_5 \right)\|_{\dot{L}^{p_4,5} L^{q_4,5}} \lesssim \|S_{k_1}^{(1)} \Delta_{m_1}^{(2)} D_{(1)}^{\gamma_1} f_4\|_{\dot{L}^{p_4} L^{q_4}} \|S_{k_1}^{(1)} \Delta_{m_5}^{(2)} D_{(2)}^{\gamma_2} f_5\|_{\dot{L}^{p_5} L^{q_5}}.$$

This means that the term above can be further estimated by

$$\sum_{\substack{k_3 \ll k_1 \\ m_5 > m_1}} 2^{(\beta_1-1)k_1\tau} 2^{k_3\tau} 2^{m_1(\alpha_2+\beta_2)\tau} \|\Delta_{k_1}^{(1)} \Delta_{m_1}^{(2)} D_{(1)}^{\alpha_1} f_1\|_{\dot{L}^{p_1} L^{q_1}}^\tau \|f_2\|_{\dot{L}^{p_2} L^{q_2}}^\tau \|\Delta_{k_3}^{(1)} f_3\|_{\dot{L}^{p_3} L^{q_3}}^\tau \|D_{(1)}^{\gamma_1} f_4\|_{\dot{L}^{p_4} L^{q_4}}^\tau \|\Delta_{m_5}^{(2)} D_{(2)}^{\gamma_2} f_5\|_{\dot{L}^{p_5} L^{q_5}}^\tau.$$

We put f_2 and f_4 aside, and what is left will be bounded (through the usual process) by

$$\left(\|D_{(1)}^{\alpha_1} f_1\|_{\dot{B}_{p_1,\infty}^0, \dot{B}_{q_1,\infty}^{\alpha_2+\beta_2}}^\tau \|f_3\|_{\dot{B}_{p_3,\infty}^{\beta_1}, L^{q_3}}^\tau \|D_{(2)}^{\gamma_2} f_5\|_{\dot{L}^{p_5} \dot{B}_{q_5,\infty}^0}^\tau \right)^{\frac{\epsilon_1}{\beta_1+\epsilon_1} \cdot \frac{\epsilon_2}{\beta_2+\epsilon_2}} \cdot \left(\|D_{(1)}^{\alpha_1} f_1\|_{\dot{B}_{p_1,\infty}^{\beta_1}, \dot{B}_{q_1,\infty}^{\alpha_2+\beta_2}}^\tau \|f_3\|_{\dot{B}_{p_3,\infty}^0, L^{q_3}}^\tau \|D_{(2)}^{\gamma_2} f_5\|_{\dot{L}^{p_5} \dot{B}_{q_5,\infty}^0}^\tau \right)^{\frac{\beta_1}{\beta_1+\epsilon_1} \cdot \frac{\epsilon_2}{\beta_2+\epsilon_2}} \\ \cdot \left(\|D_{(1)}^{\alpha_1} f_1\|_{\dot{B}_{p_1,\infty}^0, \dot{B}_{q_1,\infty}^0}^\tau \|f_3\|_{\dot{B}_{p_3,\infty}^{\beta_1}, L^{q_3}}^\tau \|D_{(2)}^{\gamma_2} f_5\|_{\dot{L}^{p_5} \dot{B}_{q_5,\infty}^{\alpha_2+\beta_2}}^\tau \right)^{\frac{\epsilon_1}{\beta_1+\epsilon_1} \cdot \frac{\beta_2}{\beta_2+\epsilon_2}} \cdot \left(\|D_{(1)}^{\alpha_1} f_1\|_{\dot{B}_{p_1,\infty}^{\beta_1}, \dot{B}_{q_1,\infty}^0}^\tau \|f_3\|_{\dot{B}_{p_3,\infty}^0, L^{q_3}}^\tau \|D_{(2)}^{\gamma_2} f_5\|_{\dot{L}^{p_5} \dot{B}_{q_5,\infty}^{\alpha_2+\beta_2}}^\tau \right)^{\frac{\beta_1}{\beta_1+\epsilon_1} \cdot \frac{\beta_2}{\beta_2+\epsilon_2}}.$$

The remaining terms can be treated in a similar way.

- $I_B^{(1)} \times I_A^{(2)}$ The case $I_B^{(1)} \times I_A^{(2)}$ is symmetric to $I_A^{(1)} \times I_B^{(2)}$.
- $I_B^{(1)} \times I_B^{(2)}$ The frequency symbol $m_{I_B^{(1)}}(\xi_1, \dots, \xi_5) \cdot m_{I_B^{(2)}}(\eta_1, \dots, \eta_5)$ is described by

$$\left(\frac{|\xi_1 + \dots + \xi_5|^{\beta_1} - |\xi_1 + \xi_2|^{\beta_1}}{\xi_3 + \xi_4 + \xi_5} \cdot (\xi_4 + \xi_5) \cdot |\xi_1 + \xi_2|^{\alpha_1} |\xi_4 + \xi_5|^{\gamma_1} \right) \cdot \left(\frac{|\eta_1 + \dots + \eta_5|^{\beta_2} - |\eta_1 + \eta_2|^{\beta_2}}{\eta_3 + \eta_4 + \eta_5} \cdot (\eta_4 + \eta_5) \cdot |\eta_1 + \eta_2|^{\alpha_2} |\eta_4 + \eta_5|^{\gamma_2} \right).$$

The “low scales” correspond to $\xi_4 + \xi_5$ in the first parameter and to $\eta_4 + \eta_5$ in the second one; for the purpose of deciding which functions will be involved in the optimization part, we need to decide which of $|\xi_4|$ and $|\xi_5|$ is larger (similarly for $|\eta_4|$ and $|\eta_5|$). We assume that $|\xi_5| \leq |\xi_4|$ and $|\eta_4| \leq |\eta_5|$, which is one of the more convoluted situations.

The associated operator of interest is

$$T_{m_{I_B^{(1)}}, m_{I_B^{(2)}}}^{k_1, k_4; m_1, m_5}(f_1, \dots, f_5)(x, y) = \int_{\mathbb{R}^{10}} m_{I_B^{(1)}}(\xi_1, \dots, \xi_5) \cdot m_{I_B^{(2)}}(\eta_1, \dots, \eta_5) \mathcal{F}(\Delta_{k_1}^{(1)} \Delta_{m_1}^{(2)} f_1)(\xi_1, \eta_1) \cdot \mathcal{F}(S_{k_1}^{(1)} S_{m_1}^{(2)} f_2)(\xi_2, \eta_2) \\ \cdot \mathcal{F}(S_{k_1}^{(1)} S_{m_1}^{(2)} f_3)(\xi_3, \eta_3) \cdot \mathcal{F}(\Delta_{k_4}^{(1)} \Delta_{m_5}^{(2)} f_4)(\xi_4, \eta_4) \cdot \mathcal{F}(\Delta_{k_4}^{(1)} \Delta_{m_5}^{(2)} f_5)(\xi_5, \eta_5) \cdot e^{2\pi i x(\xi_1 + \dots + \xi_5)} e^{2\pi i y(\eta_1 + \dots + \eta_5)} d\xi_1 \dots d\xi_5 d\eta_1 \dots d\eta_5.$$

The usual double Fourier series decomposition of the frequency-localized commutator symbols $m_{c_{\beta_1}}(\xi_1 + \xi_2, \xi_3 + \xi_4 + \xi_5)$ and $m_{c_{\beta_2}}(\eta_1 + \eta_2, \eta_3 + \eta_4 + \eta_5)$ allows to reduce the problem concerning the boundedness of $T_{m_{I_B^{(1)}}, m_{I_B^{(2)}}}^{k_1, k_4; m_1, m_5}(f_1, \dots, f_5)$ to subtrees. Using results concerning bi-parameter paraproducts (flags of lower complexity), we deduce

$$\|T_{m_{I_B^{(1)}}, m_{I_B^{(2)}}}^{k_1, k_4; m_1, m_5}(f_1, \dots, f_5)\|_{\dot{L}^p L^q}^\tau \lesssim 2^{k_1(\beta_1-1)\tau} 2^{k_4\tau} 2^{k_1\alpha_1\tau} 2^{k_4\gamma_1\tau} 2^{m_1(\beta_2-1)\tau} 2^{m_5\tau} 2^{m_1\alpha_2\tau} 2^{m_5\gamma_2\tau} \\ \cdot \|\Delta_{k_1}^{(1)} \Delta_{m_1}^{(2)} f_1\|_{\dot{L}^{p_1} L^{q_1}}^\tau \|S_{k_1}^{(1)} S_{m_1}^{(2)} f_2\|_{\dot{L}^{p_2} L^{q_2}}^\tau \|S_{k_1}^{(1)} S_{m_1}^{(2)} f_3\|_{\dot{L}^{p_3} L^{q_3}}^\tau \|\Delta_{k_4}^{(1)} \Delta_{m_5}^{(2)} f_4\|_{\dot{L}^{p_4} L^{q_4}}^\tau \|\Delta_{k_4}^{(1)} \Delta_{m_5}^{(2)} f_5\|_{\dot{L}^{p_5} L^{q_5}}^\tau \\ \lesssim 2^{k_1(\beta_1-1)\tau} 2^{k_4\tau} 2^{m_1(\beta_2-1)\tau} 2^{m_5\tau} \|\Delta_{k_1}^{(1)} \Delta_{m_1}^{(2)} D_{(1)}^{\alpha_1} D_{(2)}^{\alpha_2} f_1\|_{\dot{L}^{p_1} L^{q_1}}^\tau \|f_2\|_{\dot{L}^{p_2} L^{q_2}}^\tau \|f_3\|_{\dot{L}^{p_3} L^{q_3}}^\tau \|\Delta_{k_4}^{(1)} D_{(1)}^{\gamma_1} f_4\|_{\dot{L}^{p_4} L^{q_4}}^\tau \|\Delta_{m_5}^{(2)} D_{(1)}^{\gamma_2} f_5\|_{\dot{L}^{p_5} L^{q_5}}^\tau.$$

Summing now in $k_4 < k_1$, $m_5 < m_1$, we obtain the desired upper bound.

³⁰Several similar terms need to be considered.

- $I_B^{(1)} \times I_C^{(2)}$ The symbol $m_{I_B^{(1)}}(\xi_1, \dots, \xi_5) \cdot m_{I_C^{(2)}}(\eta_1, \dots, \eta_5)$ is

$$\left(\frac{|\xi_1 + \dots + \xi_5|^{\beta_1} - |\xi_1 + \xi_2|^{\beta_1}}{\xi_3 + \xi_4 + \xi_5} \cdot (\xi_4 + \xi_5) \cdot |\xi_1 + \xi_2|^{\alpha_1} |\xi_4 + \xi_5|^{\gamma_1} \right) \cdot \left(\frac{|\eta_1 + \eta_2|^{\alpha_2 + \beta_2} - |\eta_1|^{\alpha_2 + \beta_2}}{\eta_2} \cdot \eta_2 \cdot |\eta_4 + \eta_5|^{\gamma_2} \right).$$

We want to study the multiplier

$$T_{m_{I_B^{(1)}}, m_{I_C^{(2)}}}^{k_1, k_4; m_1, m_2}(f_1, \dots, f_5)(x, y) = \int_{\mathbb{R}^{10}} m_{I_B^{(1)}}(\xi_1, \dots, \xi_5) \cdot m_{I_C^{(2)}}(\eta_1, \dots, \eta_5) \mathcal{F}(\Delta_{k_1}^{(1)} \Delta_{m_1}^{(2)} f_1)(\xi_1, \eta_1) \cdot \mathcal{F}(S_{k_1}^{(1)} \Delta_{m_2}^{(2)} f_2)(\xi_2, \eta_2) \\ \cdot \mathcal{F}(S_{k_1}^{(1)} S_{m_1}^{(2)} f_3)(\xi_3, \eta_3) \cdot \mathcal{F}(\Delta_{k_4}^{(1)} S_{m_1}^{(2)} f_4)(\xi_4, \eta_4) \cdot \mathcal{F}(\Delta_{\leq k_4}^{(1)} S_{m_1}^{(2)} f_5)(\xi_5, \eta_5) \cdot e^{2\pi i x(\xi_1 + \dots + \xi_5)} e^{2\pi i y(\eta_1 + \dots + \eta_5)} d\xi_1 \dots d\xi_5 d\eta_1 \dots d\eta_5.$$

As before, we assume without loss of generality that $|\xi_5| \leq |\xi_4|$. After invoking the Fourier series decomposition,

$T_{m_{I_B^{(1)}}, m_{I_C^{(2)}}}^{k_1, k_4; m_1, m_2}(f_1, \dots, f_5)$ tensorizes as superpositions of the form

$$2^{k_1(\beta_1 - 1)} 2^{k_4} 2^{m_1(\alpha_2 + \beta_2 - 1)} 2^{m_2} \|D_{(1)}^{\alpha_1} (\Delta_{k_1}^{(1)} \Delta_{m_1}^{(2)} f_1 S_{k_1}^{(1)} \Delta_{m_2}^{(2)} f_2)\|_{L^{p_1, 2} L^{q_1, 2}} \|S_{k_1}^{(1)} S_{m_1}^{(2)} f_3\|_{L^{p_3} L^{q_3}} \|D_{(1)}^{\gamma_1} D_{(2)}^{\gamma_2} (\Delta_{k_4}^{(1)} S_{m_1}^{(2)} f_4 \Delta_{\leq k_4}^{(1)} S_{m_1}^{(2)} f_5)\|_{L^{p_4, 5} L^{q_4, 5}}.$$

These imply that $\|I_B^{(1)} \times I_C^{(2)}\|_{L^p L^q}^\tau$ is bounded by

$$\sum_{k_4 < k_1} \sum_{m_2 < m_1} 2^{k_1(\beta_1 - 1)\tau} 2^{k_4\tau} 2^{m_1(\alpha_2 + \beta_2 - 1)\tau} 2^{m_2\tau} \|\Delta_{k_1}^{(1)} \Delta_{m_1}^{(2)} D_{(1)}^{\alpha_1} f_1\|_{L^{p_1} L^{q_1}}^\tau \|\Delta_{m_2}^{(2)} f_2\|_{L^{p_2} L^{q_2}}^\tau \|f_3\|_{L^{p_3} L^{q_3}}^\tau \\ \left(\|\Delta_{k_4}^{(1)} D_{(1)}^{\gamma_1} D_{(2)}^{\gamma_2} f_4\|_{L^{p_4} L^{q_4}} \|f_5\|_{L^{p_5} L^{q_5}} + \|\Delta_{k_4}^{(1)} D_{(1)}^{\gamma_1} f_4\|_{L^{p_4} L^{q_4}} \|D_{(2)}^{\gamma_2} f_5\|_{L^{p_5} L^{q_5}} \right)^\tau.$$

If (F_4, F_5) denotes either of the couples $(D_{(1)}^{\gamma_1} D_{(2)}^{\gamma_2} f_4, f_5)$ or $(D_{(1)}^{\gamma_1} f_4, D_{(2)}^{\gamma_2} f_5)$, we are left with bounding

$$\sum_{k_1, m_1} \min(2^{k_1\beta_1\tau} 2^{m_1(\alpha_2 + \beta_2)\tau} \|f_1\|_{\dot{B}_{p_1, \infty}^0 \dot{B}_{q_1, \infty}^0}^\tau \|f_2\|_{L^{p_2} \dot{B}_{q_2, \infty}^0}^\tau \|F_4\|_{\dot{B}_{p_4, \infty}^0 L^{q_4}}^\tau, 2^{k_1\beta_1\tau} 2^{-m_1\epsilon_2\tau} \|f_1\|_{\dot{B}_{p_1, \infty}^0 \dot{B}_{q_1, \infty}^{\alpha_2 + \beta_2}}^\tau \|f_2\|_{L^{p_2} \dot{B}_{q_2, \infty}^{\epsilon_2}}^\tau \|F_4\|_{\dot{B}_{p_4, \infty}^0 L^{q_4}}^\tau, \\ 2^{-k_1\epsilon_1\tau} 2^{m_1(\alpha_2 + \beta_2)\tau} \|f_1\|_{\dot{B}_{p_1, \infty}^{\beta_1} \dot{B}_{q_1, \infty}^0}^\tau \|f_2\|_{L^{p_2} \dot{B}_{q_2, \infty}^0}^\tau \|F_4\|_{\dot{B}_{p_4, \infty}^{\epsilon_1} L^{q_4}}^\tau, 2^{-k_1\epsilon_1\tau} 2^{-m_1\epsilon_2\tau} \|f_1\|_{\dot{B}_{p_1, \infty}^{\beta_1} \dot{B}_{q_1, \infty}^{\alpha_2 + \beta_2}}^\tau \|f_2\|_{L^{p_2} \dot{B}_{q_2, \infty}^{\epsilon_2}}^\tau \|F_4\|_{\dot{B}_{p_4, \infty}^{\epsilon_1} L^{q_4}}^\tau).$$

Per usual, interpolation and regrouping of the terms produces a desired upper bound.

- $I_D^{(1)} \times I_D^{(2)}$ We want to estimate the operator

$$T_{m_{I_D^{(1)}}, m_{I_D^{(2)}}}^{k_1; m_1}(f_1, \dots, f_5)(x, y) = \int_{\mathbb{R}^{10}} m_{I_D^{(1)}}(\xi_1, \dots, \xi_5) \cdot m_{I_D^{(2)}}(\eta_1, \dots, \eta_5) \mathcal{F}(\Delta_{k_1}^{(1)} \Delta_{m_1}^{(2)} f_1)(\xi_1, \eta_1) \cdot \mathcal{F}(S_{k_1}^{(1)} S_{m_1}^{(2)} f_2)(\xi_2, \eta_2) \\ \cdot \mathcal{F}(S_{k_1}^{(1)} S_{m_1}^{(2)} f_3)(\xi_3, \eta_3) \cdot \mathcal{F}(S_{k_1}^{(1)} S_{m_1}^{(2)} f_4)(\xi_4, \eta_4) \cdot \mathcal{F}(S_{k_1}^{(1)} S_{m_1}^{(2)} f_5)(\xi_5, \eta_5) \cdot e^{2\pi i x(\xi_1 + \dots + \xi_5)} e^{2\pi i y(\eta_1 + \dots + \eta_5)} d\xi_1 \dots d\xi_5 d\eta_1 \dots d\eta_5$$

of symbol $m_{I_D^{(1)}}(\xi_1, \dots, \xi_5) \cdot m_{I_D^{(2)}}(\eta_1, \dots, \eta_5)$, which writes as

$$|\xi_1|^{\alpha_1 + \beta_1} \cdot |\xi_4 + \xi_5|^{\gamma_1} \cdot |\eta_1|^{\alpha_2 + \beta_2} \cdot |\eta_4 + \eta_5|^{\gamma_2}.$$

In fact, $T_{m_{I_D^{(1)}}, m_{I_D^{(2)}}}^{k_1; m_1}(f_1, \dots, f_5)$ is equal to

$$(4.12) \quad \Delta_{k_1}^{(1)} \Delta_{m_1}^{(2)} D_{(1)}^{\alpha_1 + \beta_1} D_{(2)}^{\alpha_2 + \beta_2} f_1(x, y) \cdot S_{k_1}^{(1)} S_{m_1}^{(2)} f_2(x, y) \cdot S_{k_1}^{(1)} S_{m_1}^{(2)} f_3(x, y) \cdot D_{(1)}^{\gamma_1} D_{(2)}^{\gamma_2} (S_{k_1}^{(1)} S_{m_1}^{(2)} f_4 \cdot S_{k_1}^{(1)} S_{m_1}^{(2)} f_5)(x, y).$$

Next, we write every $S_{k_1}^{(1)} S_{m_1}^{(2)} F$ as

$$S_{k_1}^{(1)} S_{m_1}^{(2)} F = F - \Delta_{> k_1}^{(1)} F - \Delta_{> m_1}^{(2)} F + \Delta_{> k_1}^{(1)} \Delta_{> m_1}^{(2)} F,$$

for F being any of the functions f_2, f_3, f_4 or f_5 . When plugging this in (4.12), we obtain four types of terms.

- The first will simply produce

$$(4.13) \quad \Delta_{k_1}^{(1)} \Delta_{m_1}^{(2)} D_{(1)}^{\alpha_1 + \beta_1} D_{(2)}^{\alpha_2 + \beta_2} f_1(x, y) \cdot f_2(x, y) \cdot f_3(x, y) \cdot D_{(1)}^{\gamma_1} D_{(2)}^{\gamma_2} (f_4 \cdot f_5)(x, y)$$

and the summation in k_1 and m_1 – that needs to be performed before taking the $\|\cdot\|_{L^p L^q}$ norms – yields

$$D_{(1)}^{\alpha_1 + \beta_1} D_{(2)}^{\alpha_2 + \beta_2} f_1(x, y) \cdot f_2(x, y) \cdot f_3(x, y) \cdot D_{(1)}^{\gamma_1} D_{(2)}^{\gamma_2} (f_4 \cdot f_5)(x, y),$$

for which we invoke a mixed-norm Hölder's inequality.

- There is at most one function “hit” by the $\Delta_{> k_1}^{(1)}$ projection, and all the functions are unaffected by $\Delta_{> m_1}^{(2)}$: for example

$$\Delta_{k_1}^{(1)} \Delta_{m_1}^{(2)} D_{(1)}^{\alpha_1 + \beta_1} D_{(2)}^{\alpha_2 + \beta_2} f_1(x, y) \Delta_{> k_1}^{(1)} f_2(x, y) \cdot f_3(x, y) \cdot D_{(1)}^{\gamma_1} D_{(2)}^{\gamma_2} (f_4 \cdot f_5)(x, y)$$

or the more involved

$$\Delta_{k_1}^{(1)} \Delta_{m_1}^{(2)} D_{(1)}^{\alpha_1 + \beta_1} D_{(2)}^{\alpha_2 + \beta_2} f_1(x, y) f_2(x, y) \cdot f_3(x, y) \cdot D_{(1)}^{\gamma_1} D_{(2)}^{\gamma_2} (f_4 \cdot \Delta_{> k_1}^{(1)} f_5)(x, y)$$

which requires a further cone decomposition in the subtree corresponding to $D_{(1)}^{\gamma_1} D_{(2)}^{\gamma_2}(f_4 \cdot f_5)$.

In either case, we first sum in m_1 to obtain the “full function” $\Delta_{k_1}^{(1)} D_{(1)}^{\alpha_1+\beta_1} D_{(2)}^{\alpha_2+\beta_2} f_1(x, y)$, and from there on we continue as in the one-parameter situation I_D .

- None of the functions f_2, f_3, f_4, f_5 are affected by $\Delta_{>k_1}^{(1)}$, but at most one of them is hit by $\Delta_{>m_1}^{(2)}$; this situation is symmetric to the previous one.
- At least one of the functions is hit by $\Delta_{>k_1}^{(1)}$, and at least one (possibly a different one) by $\Delta_{>m_1}^{(2)}$; say for example that we have

$$(4.14) \quad 2^{k_1(\alpha_1+\beta_1)} 2^{m_1(\alpha_2+\beta_2)} \Delta_{k_1}^{(1)} \Delta_{m_1}^{(2)} f_1(x, y) f_2(x, y) \cdot \Delta_{>m_1}^{(2)} f_3(x, y) \cdot D_{(1)}^{\gamma_1} D_{(2)}^{\gamma_2} (\Delta_{>k_1}^{(1)} f_4 \cdot f_5)(x, y).$$

Then we can write $\Delta_{>m_1}^{(2)} f_3(x, y)$ as $\sum_{m_3} \Delta_{m_3}^{(2)} \Delta_{>m_1}^{(2)} f_3(x, y)$ and notice that the only non-zero terms correspond to $m_3 > m_1$. Similarly, we re-decompose

$$D_{(1)}^{\gamma_1} D_{(2)}^{\gamma_2} (\Delta_{>k_1}^{(1)} f_4 \cdot f_5)(x, y) = \sum_{k_4, k_5} D_{(1)}^{\gamma_1} D_{(2)}^{\gamma_2} (\Delta_{>k_1}^{(1)} \Delta_{k_4}^{(1)} f_4 \cdot \Delta_{k_5}^{(1)} f_5)(x, y),$$

which we further restrict to the region $|\xi_5| \leq |\xi_4|$; then we need to sum

$$\sum_{k_4 \geq k_5} D_{(1)}^{\gamma_1} D_{(2)}^{\gamma_2} (\Delta_{>k_1}^{(1)} \Delta_{k_4}^{(1)} f_4 \cdot \Delta_{k_5}^{(1)} f_5)(x, y),$$

which can be reduced to

$$\sum_{k_4: k_4 > k_1} D_{(1)}^{\gamma_1} D_{(2)}^{\gamma_2} (\Delta_{>k_1}^{(1)} \Delta_{k_4}^{(1)} f_4 \cdot S_{k_4}^{(1)} f_5)(x, y).$$

All these produce

$$\sum_{\substack{k_4 > k_1 \\ m_3 > m_1}} 2^{k_1 \beta_1 \tau} 2^{m_1 \beta_2 \tau} \|\Delta_{k_1}^{(1)} \Delta_{m_1}^{(2)} f_1\|_{L^{p_1} L^{q_1}}^\tau \|f_2\|_{L^{p_2} L^{q_2}}^\tau \|\Delta_{m_3}^{(2)} f_3\|_{L^{p_3} L^{q_3}}^\tau \|D_{(1)}^{\gamma_1} D_{(2)}^{\gamma_2} (\Delta_{k_4}^{(1)} f_4 \cdot S_{k_4}^{(1)} f_5)\|_{L^{p_4, 5} L^{q_4, 5}}^\tau,$$

which is by now a usual estimate.

- The remaining terms, although not perfectly identical to the ones discussed above, can be treated in a similar way; the details are left to the reader.

4.2. Study of “diagonal” conical regions. Since the strategy used is the same, we will not repeat the computations. We emphasize however that this is the situation where the Fourier coefficients corresponding to $|\xi_1 + \dots + \xi_5|^{\beta_1}$ (or to $|\eta_1 + \dots + \eta_5|^{\beta_2}$), localized to suitable frequency intervals, will only have limited decay. This forces the conditions

$$(4.15) \quad p > \max\left(\frac{1}{1+\beta_1}, \frac{1}{1+\beta_2}\right), \quad q > \frac{1}{1+\beta_2}$$

on the Lebesgue exponents of the target space, as we will shortly see.

To take an example, we assume that $|\xi_1| \sim |\xi_5|, |\xi_2|, |\xi_3|, |\xi_4| \ll |\xi_1|$, and in the second parameter $|\eta_1| \sim |\eta_2|, |\eta_3|, |\eta_4|, |\eta_5| \ll |\eta_1|$. Then (4.3) will be (morally) replaced by

$$(4.16) \quad \sum_{k, m} \int_{\mathbb{R}^{10}} (|\xi_1 + \dots + \xi_5|^{\beta_1} |\xi_1 + \xi_2|^{\alpha_1} |\xi_4 + \xi_5|^{\gamma_1}) \cdot (|\eta_1 + \dots + \eta_5|^{\beta_2} |\eta_1 + \eta_2|^{\alpha_2} |\eta_4 + \eta_5|^{\gamma_2}) \mathcal{F}(\Delta_k^{(1)} \Delta_m^{(2)} f_1)(\xi_1, \eta_1) \\ \cdot \mathcal{F}(S_k^{(1)} \Delta_m^{(2)} f_2)(\xi_2, \eta_2) \cdot \dots \cdot \mathcal{F}(\Delta_k^{(1)} S_m^{(2)} f_5)(\xi_5, \eta_5) e^{2\pi i x(\xi_1 + \dots + \xi_5)} e^{2\pi i y(\eta_1 + \dots + \eta_5)} d\xi_1 \dots d\xi_5 d\eta_1 \dots d\eta_5$$

Since

$$|\xi_1 + \dots + \xi_5| \leq C 2^k, \quad |\eta_1 + \dots + \eta_5| \leq C 2^m,$$

as discussed in Section 2.2 – equation (2.35), we have that

$$|\xi_1 + \dots + \xi_5|^{\beta_1} \phi(2^{-k}(\xi_1 + \dots + \xi_5)) = \sum_{L \in \mathbb{Z}} C_L 2^{k\beta_1} e^{2\pi i \frac{L}{2^k}(\xi_1 + \dots + \xi_5)},$$

where

$$|C_L| \lesssim \frac{1}{(1+|L|)^{1+\beta_1}}.$$

Similarly,

$$|\eta_1 + \dots + \eta_5|^{\beta_2} \phi(2^{-m}(\eta_1 + \dots + \eta_5)) = \sum_{\tilde{L} \in \mathbb{Z}} C_{\tilde{L}} 2^{m\beta_2} e^{2\pi i \frac{\tilde{L}}{2^m}(\eta_1 + \dots + \eta_5)},$$

where

$$|C_{\tilde{L}}| \lesssim \frac{1}{(1 + |\tilde{L}|)^{1+\beta_2}}.$$

So (4.16) becomes a sum over $k, m \in \mathbb{Z}$ of terms of the form

$$(4.17) \quad 2^{k\beta_1} 2^{m\beta_2} \sum_{L, \tilde{L}} C_L C_{\tilde{L}} D_{(1)}^{\alpha_1} D_{(2)}^{\alpha_2} (\Delta_k^{(1)} \Delta_m^{(2)} f_1 \cdot S_k^{(1)} \Delta_m^{(2)} f_2) (x + \frac{L}{2^k}, y + \frac{\tilde{L}}{2^m}) S_k^{(1)} S_m^{(2)} f_3 (x + \frac{L}{2^k}, y + \frac{\tilde{L}}{2^m}) \\ \cdot D_{(1)}^{\gamma_1} D_{(2)}^{\gamma_2} (S_k^{(1)} S_m^{(2)} f_4 \cdot \Delta_k^{(1)} S_m^{(2)} f_5) (x + \frac{L}{2^k}, y + \frac{\tilde{L}}{2^m}).$$

Notice that in this context, we prefer to write

$$D_{(1)}^{\alpha_1} D_{(2)}^{\alpha_2} (\Delta_{k, \frac{L}{2^k}}^{(1)} \Delta_{m, \frac{\tilde{L}}{2^m}}^{(2)} f_1 \cdot S_{k, \frac{L}{2^k}}^{(1)} \Delta_{m, \frac{\tilde{L}}{2^m}}^{(2)} f_2)(x, y)$$

as

$$D_{(1)}^{\alpha_1} D_{(2)}^{\alpha_2} (\Delta_k^{(1)} \Delta_m^{(2)} f_1 \cdot S_k^{(1)} \Delta_m^{(2)} f_2) (x + \frac{L}{2^k}, y + \frac{\tilde{L}}{2^m}).$$

This allows us to first integrate in y without taking into account the effect of the modulation in the L parameter, which only acts in the first variable.

Once this clarified, we return to (4.17) and further write it as

$$2^{k\beta_1} 2^{m\beta_2} \sum_{L, \tilde{L} \in \mathbb{Z}} C_L C_{\tilde{L}} F_{L, \tilde{L}}^{k, m}(x, y) := 2^{k\beta_1} 2^{m\beta_2} \sum_{L, \tilde{L} \in \mathbb{Z}} C_L C_{\tilde{L}} F^{k, m}(x + \frac{L}{2^k}, y + \frac{\tilde{L}}{2^m}).$$

The delicate point about the constraints on p and q appears here. Previously in Section 4.1, the Fourier coefficients had arbitrary decay and we used the estimate

$$\left\| \sum_{L, \tilde{L} \in \mathbb{Z}} C_L C_{\tilde{L}} F_{L, \tilde{L}}^{k, m} \right\|_{L^p L^q}^\tau \leq \sum_{L, \tilde{L} \in \mathbb{Z}} |C_L|^\tau |C_{\tilde{L}}|^\tau \|F_{L, \tilde{L}}^{k, m}\|_{L^p L^q}^\tau;$$

however, this would require that

$$p, q > \max\left(\frac{1}{1 + \beta_1}, \frac{1}{1 + \beta_2}\right),$$

which is stronger than the announced (4.15).

Instead, we use that, for $q_0 \leq \min(1, q)$, $\|\cdot\|_q^{q_0}$ is subadditive, and thus

$$(4.18) \quad \left\| \sum_{L, \tilde{L} \in \mathbb{Z}} C_L C_{\tilde{L}} F_{L, \tilde{L}}^{k, m} \right\|_{L^p L^q}^\tau \lesssim \sum_{L \in \mathbb{Z}} |C_L|^\tau \left(\int_{\mathbb{R}} \left(\sum_{\tilde{L} \in \mathbb{Z}} |C_{\tilde{L}}|^{q_0} \left(\int_{\mathbb{R}} |F^{k, m}(x + \frac{L}{2^k}, y + \frac{\tilde{L}}{2^m})|^q dy \right)^{\frac{q_0}{q}} \right)^{\frac{p}{q_0}} dx \right)^{\frac{\tau}{p}} \\ \lesssim \sum_{L \in \mathbb{Z}} |C_L|^\tau \left(\int_{\mathbb{R}} \left(\sum_{\tilde{L} \in \mathbb{Z}} |C_{\tilde{L}}|^{q_0} \left(\int_{\mathbb{R}} |F^{k, m}(x, y)|^q dy \right)^{\frac{q_0}{q}} \right)^{\frac{p}{q_0}} dx \right)^{\frac{\tau}{p}}.$$

At this point, it is important to notice that

$$\|F^{k, m}(x, \cdot)\|_{L_y^q} \lesssim \|D_{(1)}^{\alpha_1} D_{(2)}^{\alpha_2} (\Delta_k^{(1)} \Delta_m^{(2)} f_1 \cdot S_k^{(1)} \Delta_m^{(2)} f_2)(x, \cdot)\|_{L_y^{q_{1,2}}} \|S_k^{(1)} S_m^{(2)} f_3\|_{L_y^{q_3}} \\ \|D_{(1)}^{\gamma_1} D_{(2)}^{\gamma_2} (S_k^{(1)} S_m^{(2)} f_4 \cdot \Delta_k^{(1)} S_m^{(2)} f_5)(x, \cdot)\|_{L_y^{q_{4,5}}}.$$

So provided that

$$\sum_{L \in \mathbb{Z}} |C_L|^\tau < \infty, \quad \sum_{\tilde{L} \in \mathbb{Z}} |C_{\tilde{L}}|^{q_0} < \infty,$$

which amounts to conditions (4.15) holding, we have that (4.16), estimated in $\|\cdot\|_{L^p L^q}^\tau$ is bounded above by

$$\sum_{k, m} 2^{k\beta_1 \tau} 2^{m\beta_2 \tau} \|D_{(1)}^{\alpha_1} D_{(2)}^{\alpha_2} (\Delta_k^{(1)} \Delta_m^{(2)} f_1 \cdot S_k^{(1)} \Delta_m^{(2)} f_2)\|_{L^{p_{1,2}} L^{q_{1,2}}}^\tau \|S_k^{(1)} S_m^{(2)} f_3\|_{L^{p_3} L^{q_3}}^\tau \\ \|D_{(1)}^{\gamma_1} D_{(2)}^{\gamma_2} (S_k^{(1)} S_m^{(2)} f_4 \cdot \Delta_k^{(1)} S_m^{(2)} f_5)\|_{L^{p_{4,5}} L^{q_{4,5}}}^\tau.$$

From here on the argument follows the usual strategy: using the boundedness of the lower complexity flag paraproducts – in this case the mixed norm estimates for frequency-localized bi-parameter paraproducts, we obtain that the expression above is further bounded by

$$\|f_3\|_{L^{p_3} L^{q_3}}^\tau \|D_{(2)}^{\gamma_2} f_4\|_{L^{p_4} L^{q_4}}^\tau \left(\|D_{(1)}^{\alpha_1} D_{(2)}^{\alpha_2} f_1\|_{\dot{B}_{p_1, \infty}^0 \dot{B}_{q_1, \infty}^0} \|f_2\|_{L^{p_2} \dot{B}_{q_1, \infty}^{\beta_2}} \|D_{(1)}^{\gamma_1} f_5\|_{\dot{B}_{p_5, \infty}^{\beta_1} L^{q_5}} \right)^{\frac{\tau}{4}}$$

$$\left(\|D_{(1)}^{\alpha_1} D_{(2)}^{\alpha_2} f_1\|_{\dot{B}_{p_1, \infty}^{\beta_1} \dot{B}_{q_1, \infty}^0} \|f_2\|_{L^{p_2} \dot{B}_{q_1, \infty}^{\beta_2}} \|D_{(1)}^{\gamma_1} f_5\|_{\dot{B}_{p_5, \infty}^0 L^{q_5}} \right)^{\frac{\tau}{4}} \left(\|D_{(1)}^{\alpha_1} D_{(2)}^{\alpha_2} f_1\|_{\dot{B}_{p_1, \infty}^0 \dot{B}_{q_1, \infty}^{\beta_2}} \|f_2\|_{L^{p_2} \dot{B}_{q_1, \infty}^0} \|D_{(1)}^{\gamma_1} f_5\|_{\dot{B}_{p_5, \infty}^{\beta_1} L^{q_5}} \right)^{\frac{\tau}{4}} \\ + \left(\|D_{(1)}^{\alpha_1} D_{(2)}^{\alpha_2} f_1\|_{\dot{B}_{p_1, \infty}^{\beta_1} \dot{B}_{q_1, \infty}^{\beta_2}} \|f_2\|_{L^{p_2} \dot{B}_{q_1, \infty}^0} \|D_{(1)}^{\gamma_1} f_5\|_{\dot{B}_{p_5, \infty}^0 L^{q_5}} \right)^{\frac{\tau}{4}}$$

+ a similar term, in which the $D_{(2)}^{\gamma_2}$ derivatives acts on the function f_5 .

The remaining cases can be treated in a similar way; the main idea to bear in mind is that a lower decay of the Fourier coefficients requires a regrouping of the information as in (4.18), so that the derivatives acting on the exterior variables will not affect the Lebesgue exponents corresponding to inner variables.

5. GENERIC FLAG: AN INDUCTIVE ARGUMENT

In this section we provide an inductive argument – based on the complexity of the rooted tree – that allows to prove the general result of Theorem 1.1. Our approach integrates many of the ideas already presented: one starts by decomposing the frequency space into cones,³¹ and then further into Whitney rectangles; if the cone is so that the output variable is away from the origin, the symbol smoothly restricted to the Whitney rectangles/cubes will be split as “commutator” + “derivative acting on a lower number of functions”. Next, a Fourier series expansion on each Whitney cube/rectangle is used in order to tensorize the information contained in the root symbol, obtaining in this way similar objects associated to rooted trees of lower complexity. From here, one proceeds as in Section 2.2, although in case of multi-parameter flag Leibniz rules one needs to track more carefully the distribution of derivatives, encoded in various types of mixed Lebesgue and Besov norms.

We will address the various difficulties one at a time. First, in Section 5.1 we present the inductive argument in the one-parameter case, with emphasis on the splitting of the root symbol (depending on the type of cone we are looking at), and the necessary inductive statements that allow to reduce the complexity of the rooted tree. In Section 5.2, the bi-parameter case is presented; the process of splitting the root symbol, already used in the previous section, needs to be performed in each parameter separately, which increases the number of cases to be considered. Similarly, we will have a variety of necessary inductive statements, depending on the tree structures and the configurations of functions that appear in the summation over the scales³² step. Once acquainted with the splitting of the root symbols and the reduction of the tree’s complexity when several parameters are involved, it remains to check that the end result – now expressed as a geometric mean of mixed Lebesgue and Besov norms – indeed corresponds to the desired distribution of derivatives. This last step is carried out in Section 5.3.

In what follows, our analysis will be performed in dimension one; as discussed in Remark 2.4, the employed strategy is easily adaptable to higher dimensions. In the one-parameter case, presented in Section 5.1, we will assume the target space norm $\|\cdot\|_{L^r}$ to be subadditive: when $r < 1$, the subadditivity is achieved by considering $\|\cdot\|_{L^r}^\tau$ with $\tau \leq \min(1, r)$. Similarly, in the mixed-norm multi-parameter case we would need to work with $\|\cdot\|_{L^r}^\tau$ with $\tau \leq \min(1, r^1, \dots, r^N)$ in order to obtain subadditivity; the more involved conditions on the Lebesgue exponents expressed in (1.26) of Theorem 1.1 require a more careful analysis, which was detailed in Section 4.2. In an attempt to remove unnecessary technicalities burdening the notation, we will also assume in Section 5.2 that $\|\cdot\|_{L^r}$ is subadditive.

5.1. One-parameter flag Leibniz rule. We provide a proof for the one-parameter Leibniz rule corresponding to an arbitrary n -linear flag in dimension one using an inductive argument. In what follows, we use the notation introduced in Section 1.2. Let \mathcal{G} be a tree of arbitrary complexity. Due to the paraproduct decomposition described in Section 2.3, the frequency space is split into conical regions, which are generically of two types:

$$(5.1) \quad R_{l_0} := \{(\xi_1, \dots, \xi_n) : |\xi_{l_0}| \gg |\xi_l| \text{ for all } 1 \leq l \neq l_0 \leq n\}$$

and

$$(5.2) \quad \tilde{R}_{l_1, l_2} := \{(\xi_1, \dots, \xi_n) : |\xi_{l_1}| \sim |\xi_{l_2}| \geq |\xi_l| \text{ for all } 1 \leq l \leq n\}.$$

We will introduce the maps \mathfrak{M} and \mathfrak{m} defined on the collection of conical regions:

$$(5.3) \quad \mathfrak{M}(R) := \{1 \leq l \leq n : (\xi_1, \dots, \xi_n) \in R, \quad |\xi_l| \sim \max_{1 \leq l' \leq n} |\xi_{l'}|\}, \\ \mathfrak{m}(R) := \{1 \leq l \leq n : (\xi_1, \dots, \xi_n) \in R, \quad |\xi_l| \ll \max_{1 \leq l' \leq n} |\xi_{l'}|\},$$

where R is a conical region of the form (5.1) or (5.2). The definition of conical regions thus implies that

$$\mathcal{L}(\tau_{\mathcal{G}}) = \{1, \dots, n\} = \mathfrak{M}(R) \cup \mathfrak{m}(R).$$

³¹This is the usual paraproduct decomposition.

³²As in Sections 2.2, 3 and 4, there will always be two functions involved in the summation over the scales step – in each parameter.

Since the frequency space can be decomposed into finitely many such regions, it suffices to derive the same bound for our multilinear expression localized on a fixed conical region in frequency. Let us denote by $T_{\mathcal{G}}^R$ the multilinear operator smoothly restricted to a cone R . Then for $k \in \mathbb{Z}$ and $L \in \mathbb{R}$, we define the projection operators P_k and $P_{k,L}$ ³³:

$$P_k(l) := \begin{cases} \Delta_k & \text{if } l \in \mathfrak{M}(R) \\ S_k & \text{if } l \in \mathfrak{m}(R), \end{cases} \quad \text{and} \quad P_{k,L}(l) := \begin{cases} \Delta_{k,L} & \text{if } l \in \mathfrak{M}(R) \\ S_{k,L} & \text{if } l \in \mathfrak{m}(R). \end{cases}$$

The projection operators themselves depend on the conical region R ; this will be omitted from the notation, but it should be implicit in the analysis.

If we look at the cone R_{l_0} described in (5.1), we notice that $\mathfrak{M}(R_{l_0}) = \{l_0\}$ and $\mathfrak{m}(R_{l_0}) = \{1, \dots, n\} \setminus \{l_0\}$. Moreover, it can be represented as a union of Whitney cubes:

$$(5.4) \quad R_{l_0} = \bigcup_{k_{l_0}} R_{k_{l_0}}^{\pm} := \bigcup_{k_{l_0}} \{(\xi_1, \dots, \xi_n) : \xi_{l_0} \sim \pm 2^{k_{l_0}}, |\xi_l| \ll 2^{k_{l_0}} \text{ for } l \neq l_0\},$$

which corresponds exactly to the projection operators applied to the leaves:

$$\Delta_{k_{l_0}, \pm} f_{l_0} \text{ and } S_{k_{l_0}} f_l \text{ for } l \neq l_0.$$

On the fixed conical region R_{l_0} (as defined in (5.4)), we will refer to k_{l_0} as k_{\max} since it is naturally associated to the variable ξ_{l_0} and the function f_{l_0} . Notice that R_{l_0} becomes the union of all Whitney cubes at scale k_{\max} .

We can therefore denote by

$$T_{\mathcal{G}}^R((P_{k_{\max}} f_l)_{1 \leq l \leq n})$$

the multi-linear expression $T_{\mathcal{G}}(f_1, \dots, f_n)$ localized on the union of Whitney cubes at scale k_{\max} in the conical region R . $T_{\mathcal{G}}(f_1, \dots, f_n)$ restricted to the entire conical region can then be represented as

$$(5.5) \quad \sum_{k_{\max} \in \mathbb{Z}} T_{\mathcal{G}}^R((P_{k_{\max}} f_l)_{1 \leq l \leq n}).$$

In our inductive process, operators associated to subtrees of \mathcal{G} will play an important role; hence, for any vertex v , we denote by \mathcal{G}^v the subtree of \mathcal{G} rooted in v . We will need to consider also paraproduct decompositions on these subtrees. For any non-leaf vertex $v \in \mathcal{V}$, which becomes the root of the subtree \mathcal{G}^v , the new conical region associated to the subtree will be clarified and the corresponding Whitney cubes will be specified by $k_{\max}(v)$. For simplicity of notation, we will use the abbreviation $k_{\max} = k_{\max}(\mathfrak{r}_{\mathcal{G}})$ to denote the maximal scale involved in the definition of $T_{\mathcal{G}}$ restricted to a certain cone.

We observe that when the operator $T_{\mathcal{G}}$ is localized on a conical region R such that $\mathfrak{M}(R) \cap \mathcal{L}(v) \neq \emptyset$ for some $v \in \mathcal{V}$, then $T_{\mathcal{G}^v}$ – the operator associated to the subtree \mathcal{G}^v – is also automatically restricted to the conical region

$$R(v) := \{(\xi_l)_{l \in \mathcal{L}(v)} : |\xi_l| \gg |\xi_{l'}| \text{ for } l \in \mathcal{L}(v) \cap \mathfrak{M}(R), l' \in \mathcal{L}(v) \setminus \mathfrak{M}(R)\}.$$

Such localization on the subtree imposed by the conical decomposition for the original tree will be repetitively used in our inductive process.

The following notation will be useful in the formulation of induction. For any vertex $v \in \mathcal{V}$ which generates the subtree \mathcal{G}^v , define \mathcal{V}^v to be the set of non-leaf vertices associated with \mathcal{G}^v . If v is a leaf, then $\mathcal{G}^v = \{v\}$ and $\mathcal{V}^v = \emptyset$.

We recall that the properties (i) and (ii) from Section 1 were important in describing the derivative distribution function $\delta : \mathcal{V} \rightarrow \mathcal{L}_{\mathcal{G}}$, which has to agree with the composition law. In order to better understand the behavior of δ when restricted to subtrees, we need to define first the collection of non-leaf vertices in the path of f_l (for $1 \leq l \leq n$) to the root v in the subtree \mathcal{G}^v :

$$\mathcal{V}_l^v := \{w : w \in \mathcal{V}^v, l \in \mathcal{L}(w)\},$$

and also its complement with respect to the subtree \mathcal{G}^v :

$$(\mathcal{V}_l^v)^c := \mathcal{V}^v \setminus \mathcal{V}_l^v.$$

With some abuse of notation, if $v = f_l$ is a leaf, then $\mathcal{V}_l^v = \emptyset$.

We include a figure to illustrate the notation: in the first figure, the path highlighted in red represents the path from the vertex v to the leaf l and \mathcal{V}_l^v indeed corresponds to the collection of the non-leaf vertices along the red path, namely $\{\bar{v}_1, \bar{v}_2, \dots, \bar{v}_M\}$, as indicated in Figure 5(i).

We define the common vertices shared by the paths of f_{l_1} and f_{l_2} by

$$\mathcal{V}_{l_1, l_2}^v := \mathcal{V}_{l_1}^v \cap \mathcal{V}_{l_2}^v,$$

³³We recall that $P_{k,L}$ is simply a frequency modulation of P_k – see (2.18).

which can be represented as an ordered set starting from the root v and ending with the vertex denoted by v^{l_1, l_2} ³⁴:

$$\mathcal{V}_{l_1, l_2}^v = \{v =: \tilde{v}_1, \tilde{v}_2, \dots, v^{l_1, l_2} =: \tilde{v}_M\}.$$

in the sense that the latter element is a direct descendant of the former, which further implies that

$$(5.6) \quad \mathcal{L}(v^{l_1, l_2}) = \mathcal{L}(\tilde{v}_M) \subseteq \mathcal{L}(\tilde{v}_{M-1}) \subseteq \dots \subseteq \mathcal{L}(\tilde{v}_2) \subseteq \mathcal{L}(\tilde{v}_1) = \mathcal{L}(v).$$

Moreover, the definitions of the common path and of v^{l_1, l_2} indicate the existence of two vertices w^{l_1}, w^{l_2} such that w^{l_1}, w^{l_2} are direct descendants of v^{l_1, l_2} with $w^{l_1} \neq w^{l_2}$ and the subtree stemming from w^{l_i} , denoted by $\mathcal{G}^{w^{l_i}}$, contains l_i as its leaf (for $i = 1, 2$). Equivalently, $l_1 \in \mathcal{L}(w^{l_1})$ and $l_2 \in \mathcal{L}(w^{l_2})$.

We clarify the notation through Figure 5(ii) – the common path from v to l_1 and to l_2 is marked in red and \mathcal{V}_{l_1, l_2}^v is the collection of all the non-leaf vertices along this path. The subtrees $\mathcal{G}^{w^{l_1}}$ and $\mathcal{G}^{w^{l_2}}$ are highlighted in blue and green respectively.




FIGURE 5. Figures illustrating the notation.

Define the sum of derivatives in the downward path from the vertex v to the leaf f_l by

$$\beta(v, l) := \sum_{w \in \mathcal{V}_l^v} \beta^w.$$

If $v = f_l$ is a leaf, then $\beta(v, l) = 0$. The property (ii) of the derivative distribution function can also be written as follows: if $\delta(v) = l$ for some $v \in \mathcal{V}$, then following the notation in Figure 5(i), $\delta(\tilde{v}_2) = \dots = \delta(\tilde{v}_M) = l$. Equivalently, δ restricted to the non-leaf vertices \mathcal{V}^v of the subtree \mathcal{G}^v satisfies

$$(5.7) \quad (\delta|_{\mathcal{V}^v})^{-1}(l) = \beta(v, l).$$

For any subset of non-leaf vertices $\mathcal{W} \subseteq \mathcal{V}$, we can denote by $\mathcal{D}(\mathcal{W})$ the collection of the maps $\delta|_{\mathcal{W}}$ defined on \mathcal{W} satisfying the conditions (i) and (ii). On the complement $(\mathcal{V}_l^v)^c$, we thus have the condition that $\delta|_{(\mathcal{V}_l^v)^c} \in \mathcal{D}((\mathcal{V}_l^v)^c)$. We will consider $\delta|_{\mathcal{W}} \in \mathcal{D}(\mathcal{W})$ as a default condition for the restricted map $\delta|_{\mathcal{W}}$ with $\mathcal{W} \subseteq \mathcal{V}$, which will be omitted oftentimes for the simplicity of notation.

With the notation set, we are now ready to present the proof of Theorem 1.1, in the one-parameter case. Instead of performing the induction solely on the conclusion statement, we will proceed with multiple inductive

³⁴The vertex v^{l_1, l_2} represents the last common ancestor of l_1 and l_2 in \mathcal{G} .

statements that are useful in deducing (1.23). Let R be a conical region of the form (5.1) or (5.2). Without loss of generality, we will focus on $T_{\mathcal{G}}^R((P_{k_{\max}} f_l)_{1 \leq l \leq n})$ and abbreviate as $T_{\mathcal{G}}((P_{k_{\max}} f_l)_{1 \leq l \leq n})$. The inductive statements, which can be seen as a generalization of (2.27) in Lemma 2.2, are the following:

Proposition 5.1. *Suppose that all the Lebesgue exponents in the inductive statement satisfy the condition described in Theorem 1.1, and that $T_{\mathcal{G}}$ is restricted to a cone denoted by R .*

(1) *Suppose that $l_0 \in \mathfrak{M}(R)$. Then*

$$(5.8) \quad \|T_{\mathcal{G}}((P_{k_{\max}} f_l)_{1 \leq l \leq n})\|_r \lesssim 2^{k_{\max} \beta(\tau_{\mathcal{G}}, l_0)} \|\Delta_{k_{\max}} f_{l_0}\|_{p_{l_0}} \sum_{\delta | (\nu_{l_0}^{\tau_{\mathcal{G}}})^c} \prod_{l \neq l_0} \|D^{\delta^{-1}(l)} f_l\|_{p_l};$$

(2) *Suppose that $l_1, l_2 \in \mathfrak{M}(R)$ with $l_1 \neq l_2$. Then*

$$(5.9) \quad \|T_{\mathcal{G}}((P_{k_{\max}} f_l)_{1 \leq l \leq n})\|_r \lesssim 2^{k_{\max} \beta(\tau_{\mathcal{G}}, v^{l_1, l_2})} \|\Delta_{k_{\max}} D^{\beta(w^{l_1, l_1})} f_{l_1}\|_{p_{l_1}} \|\Delta_{k_{\max}} D^{\beta(w^{l_2, l_2})} f_{l_2}\|_{p_{l_2}} \sum_{\delta | (\nu_{l_1}^{\tau_{\mathcal{G}}})^c \setminus \nu_{l_2}^{w^{l_2}}} \prod_{l \neq l_1, l_2} \|D^{\delta^{-1}(l)} f_l\|_{p_l};$$

(3)

$$(5.10) \quad \left\| \sum_{k_{\max}} T_{\mathcal{G}}((P_{k_{\max}} f_l)_{1 \leq l \leq n}) \right\|_r \lesssim \sum_{\delta} \prod_{l=1}^n \|D^{\delta^{-1}(l)} f_l\|_{p_l}.$$

Remark 5.2. (i) Induction statements (1) and (2) describe the estimates for the multi-linear expression localized to a union of Whitney cubes at scale k_{\max} in the cone R , depending on the configuration of $\mathfrak{M}(R)$. In particular, (5.9) corresponds to the case when the Whitney cubes are located in a cone \tilde{R}_{l_1, l_2} of type (5.2); (5.8) on the other hand holds both for cones of type (5.1) or (5.2). Since (2) describes a special case of (1), it is not surprising to observe that the expression on the right hand side of (5.9) can be majorized by the right hand side of (5.8).

(ii) We would like to emphasize that (5.10) for all possible paraproduct regions implies precisely the conclusion of Theorem 1.1:

$$(5.11) \quad \|T_{\mathcal{G}}((f_l)_{1 \leq l \leq n})\|_r \lesssim \sum_{\delta} \prod_{l=1}^n \|D^{\delta^{-1}(l)} f_l\|_{p_l}.$$

Proof. As mention earlier, we will only focus on the case $r \geq 1$. We proceed by induction on the tree structure: the base case corresponds to trees of complexity 1, thus the base cases for (5.8) and (5.9) are verified by Lemma 2.2.³⁵ The base case for (5.10) is simply the Leibniz rule corresponding to paraproducts and thus is proven to be true – see Section 2.2. We would like to prove (5.8), (5.9) and (5.10) for a tree of complexity \mathcal{C} assuming that (5.8), (5.9) and (5.10) hold for any tree of *any* lower complexity (that is, of complexity $\mathcal{C} - 1, \dots, 1$).

(1) We first verify the inductive statement (5.8). Denote by $(v_i)_{i=1}^{n_1}$ the vertices of depth 1 in \mathcal{G} , and let $m_{\beta \tau_{\mathcal{G}}}$ be the symbol defined by

$$m_{\beta \tau_{\mathcal{G}}} \left(\sum_{l=1}^n \xi_l \right) := \left| \sum_{l=1}^n \xi_l \right|^{\beta \tau_{\mathcal{G}}}.$$

The multilinear expression $T_{\mathcal{G}}$ can be rewritten as

$$(5.12) \quad T_{\mathcal{G}}((P_{k_{\max}} f_l)_{1 \leq l \leq n}) = T_{m_{\beta \tau_{\mathcal{G}}}} \left(T_{\mathcal{G}^{v_1}}((P_{k_{\max}} f_l)_{l \in \mathcal{L}(v_1)}), \dots, T_{\mathcal{G}^{v_{n_1}}}((P_{k_{\max}} f_l)_{l \in \mathcal{L}(v_{n_1})}) \right),$$

where T_m generically denotes a multilinear operator associated to a symbol m .

We recall that the paraproduct decomposition yields the localization to the region $R_{k_{\max}}$ defined by

$$(5.13) \quad R_{k_{\max}} := \{(\xi_1, \dots, \xi_n) : |\xi_l| \sim 2^{k_{\max}} \text{ for } l \in \mathfrak{M}(R) \text{ and } |\xi_l| \ll 2^{k_{\max}} \text{ for } l \in \mathfrak{m}(R)\},$$

on which one has $|\sum_{l=1}^n \xi_l| \lesssim n 2^{k_{\max}}$. We can thus smoothly restrict the symbol $m_{\beta \tau_{\mathcal{G}}}(\sum_{l=1}^n \xi_l)$ to the interval $[-n 2^{k_{\max}}, n 2^{k_{\max}}]$ and denote it by $m_{\beta \tau_{\mathcal{G}}}^{k_{\max}}(\sum_{l=1}^n \xi_l)$. We perform a Fourier series decomposition on the symbol as in Section 2.2 (Fourier series decomposition for the “diagonal” term):

$$m_{\beta \tau_{\mathcal{G}}}^{k_{\max}} \left(\sum_{l=1}^n \xi_l \right) = (n 2^{k_{\max}})^{\beta \tau_{\mathcal{G}}} \sum_{L \in \mathbb{Z}} C_L e^{2\pi i \frac{L}{n 2^{k_{\max}}} \sum_{l=1}^n \xi_l},$$

where the renormalized Fourier coefficients satisfy the decaying condition

$$(5.14) \quad |C_L| \lesssim \frac{1}{(1 + |L|)^{1 + \beta \tau_{\mathcal{G}}}}.$$

³⁵Although the lemma indicates the bi-linear case, it can be easily extended to the n -linear setting

As a result, one can rewrite (5.12) (up to a constant depending implicitly on n and $\beta^{\tau_{\mathcal{G}}}$) as

$$(5.15) \quad \sum_{L \in \mathbb{Z}} C_L 2^{k_{\max} \cdot \beta^{\tau_{\mathcal{G}}}} \prod_{i=1}^{n_1} T_{\mathcal{G}^{v_i}} \left((P_{k_{\max}, \frac{L}{n 2^{k_{\max}}}} f_l)_{l \in \mathcal{L}(v_i)} \right),$$

which can be estimated by

$$(5.16) \quad \| (5.15) \|_r \lesssim \sum_{L \in \mathbb{Z}} |C_L| 2^{k_{\max} \cdot \beta^{\tau_{\mathcal{G}}}} \prod_{i=1}^{n_1} \left\| T_{\mathcal{G}^{v_i}} \left((P_{k_{\max}, \frac{L}{n 2^{k_{\max}}}} f_l)_{l \in \mathcal{L}(v_i)} \right) \right\|_{p_{v_i}}.$$

We observe that for each $1 \leq i \leq n_1$, the following identity holds:

$$(5.17) \quad T_{\mathcal{G}^{v_i}} \left((P_{k_{\max}, \frac{L}{n 2^{k_{\max}}}} f_l)_{l \in \mathcal{L}(v_i)} \right)(x) = T_{\mathcal{G}^{v_i}} \left((P_{k_{\max}} f_l)_{l \in \mathcal{L}(v_i)} \right) \left(x + \frac{L}{n 2^{k_{\max}}} \right)$$

and the translation invariance of the measure yields

$$\left\| T_{\mathcal{G}^{v_i}} \left((P_{k_{\max}, \frac{L}{n 2^{k_{\max}}}} f_l)_{l \in \mathcal{L}(v_i)} \right) \right\|_{p_{v_i}} = \left\| T_{\mathcal{G}^{v_i}} \left((P_{k_{\max}} f_l)_{l \in \mathcal{L}(v_i)} \right) \right\|_{p_{v_i}}.$$

Due to the decay of the Fourier coefficients (5.14),³⁶ (5.16) is majorized by

$$(5.18) \quad 2^{k_{\max} \cdot \beta^{\tau_{\mathcal{G}}}} \prod_{i=1}^{n_1} \left\| T_{\mathcal{G}^{v_i}} \left((P_{k_{\max}} f_l)_{l \in \mathcal{L}(v_i)} \right) \right\|_{p_{v_i}}.$$

A simple observation is that there exists some $1 \leq i_0 \leq n_1$ such that $\mathcal{L}(v_{i_0}) \cap \mathfrak{M}(R) \neq \emptyset$, in which case the subtree $\mathcal{G}^{v_{i_0}}$ is automatically restricted to a conical region and $k_{\max} = k_{\max}(v_{i_0})$ specifies the union of Whitney cubes at scale k_{\max} in such a cone. Assume without loss of generality that $i_0 = 1$. We now apply the inductive hypothesis (5.8):

$$(5.19) \quad \left\| T_{\mathcal{G}^{v_1}} \left((P_{k_{\max}} f_l)_{l \in \mathcal{L}(v_1)} \right) \right\|_{p_{v_1}} \lesssim 2^{k_{\max} \cdot \beta(v_1, l_0)} \|\Delta_{k_{\max}} f_{l_0}\|_{p_{l_0}} \sum_{\delta|_{(\mathcal{V}_{l_0}^{v_1})^c}} \prod_{\substack{l \in \mathcal{L}(v_1) \\ l \neq l_0}} \|D^{\delta^{-1}(l)} f_l\|_{p_l}.$$

Meanwhile, we can invoke the inductive hypothesis (5.10) and thus (5.11) to deduce that for $i \neq 1$,

$$(5.20) \quad \left\| T_{\mathcal{G}^{v_i}} \left((P_{k_{\max}} f_l)_{l \in \mathcal{L}(v_i)} \right) \right\|_{p_{v_i}} \lesssim \sum_{\delta|_{\mathcal{V}^{v_i}}} \prod_{l \in \mathcal{L}(v_i)} \|D^{\delta^{-1}(l)} f_l\|_{p_l}.$$

Combining the estimates (5.19) and (5.20), we derive the following desired estimate for (5.18):

$$2^{k_{\max} \cdot \beta(\tau_{\mathcal{G}}, l_0)} \|\Delta_{k_{\max}} f_{l_0}\|_{p_{l_0}} \sum_{\delta|_{(\mathcal{V}_{l_0}^{\tau_{\mathcal{G}}})^c}} \prod_{l \neq l_0} \|D^{\delta^{-1}(l)} f_l\|_{p_l}.$$

(2) To show the second inductive statement (5.9), we use the Fourier series decomposition for the root symbol applied in the proof of (5.8) and obtain (5.15), whose L^r norm can be estimated by (5.18). There are two possibilities with respect to the positions of l_1, l_2 :

(a) $l_1, l_2 \in \mathcal{L}(v_{i_0})$ for some $1 \leq i_0 \leq n_1$ (or equivalently l_1, l_2 are leaves of the same subtree rooted in v_{i_0} , a direct descendant of the root rg). We now invoke the inductive hypothesis (5.9) on $T_{\mathcal{G}^{v_{i_0}}}$:

$$(5.21) \quad \left\| T_{\mathcal{G}^{v_{i_0}}} \left((P_{k_{\max}} f_l)_{l \in \mathcal{L}(v_{i_0})} \right) \right\|_{p_{v_{i_0}}} \lesssim 2^{k_{\max} \cdot \beta(v_{i_0}, v^{l_1, l_2})} \|\Delta_{k_{\max}} D^{\beta(w^{l_1, l_1})} f_{l_1}\|_{p_{l_1}} \|\Delta_{k_{\max}} D^{\beta(w^{l_2, l_2})} f_{l_2}\|_{p_{l_2}} \sum_{\delta|_{(\mathcal{V}_{l_1}^{v_{i_0}})^c \setminus \mathcal{V}_{l_2}^{w^{l_2}}}} \prod_{\substack{l \in \mathcal{L}(v_{i_0}) \\ l \neq l_1, l_2}} \|D^{\delta^{-1}(l)} f_l\|_{p_l}.$$

We can also apply the inductive hypothesis (5.11) to derive (5.20) for $i \neq i_0$. Combining (5.21) and (5.20), we conclude with (5.9).

³⁶When $r < 1$, we would use subadditivity to deduce

$$\| (5.15) \|_r^r \lesssim \sum_{L \in \mathbb{Z}} |C_L|^r \cdot 2^{k_{\max} \beta^{\tau_{\mathcal{G}}} r} \prod_{i=1}^{n_1} \left\| T_{\mathcal{G}^{v_i}} \left((P_{k_{\max}, \frac{L}{n 2^{k_{\max}}}} f_l)_{l \in \mathcal{L}(v_i)} \right) \right\|_{p_{v_i}}^r = \sum_{L \in \mathbb{Z}} |C_L|^r \cdot 2^{k_{\max} \beta^{\tau_{\mathcal{G}}} r} \prod_{i=1}^{n_1} \left\| T_{\mathcal{G}^{v_i}} \left((P_{k_{\max}} f_l)_{l \in \mathcal{L}(v_i)} \right) \right\|_{p_{v_i}}^r,$$

where the Fourier coefficients satisfy the decay condition (5.14). It is natural to impose the condition

$$r(1 + \beta^{\tau_{\mathcal{G}}}) > 1.$$

For the same reason, for any $v \in \mathcal{L}(\tau_{\mathcal{G}})$, there is an associated differential operator D^{β^v} whose Fourier series decomposition yields Fourier coefficients with limited decay, thus imposing the condition on the Lebesgue exponent

$$p_v(1 + \beta^v) > 1.$$

- (b) $l_1 \in \mathcal{L}(v_{i_1})$ and $l_2 \in \mathcal{L}(v_{i_2})$ for $1 \leq i_1, i_2 \leq n_1$ and $i_1 \neq i_2$ (or equivalently l_1, l_2 are leaves of two different subtrees stemming from direct descendants of the root). In this case, the common path is $\mathcal{V}_{l_1, l_2}^{\tau_{\mathcal{G}}} = \{\mathbf{r}_{\mathcal{G}}\}$, which means that

$$(5.22) \quad v^{l_1, l_2} = \mathbf{r}_{\mathcal{G}}.$$

Moreover, the assumption about the positions of l_1 and l_2 gives the precise vertices w^{l_1} and w^{l_2} :

$$(5.23) \quad w^{l_1} = v_{i_1} \quad \text{and} \quad w^{l_2} = v_{i_2}.$$

The inductive hypothesis (5.8) can be applied to

$$T_{\mathcal{G}^{v_{i_1}}}((P_{k_{\max}} f_l)_{l \in \mathcal{L}(v_{i_1})}) \quad \text{and} \quad T_{\mathcal{G}^{v_{i_2}}}((P_{k_{\max}} f_l)_{l \in \mathcal{L}(v_{i_2})}).$$

In particular,

$$\begin{aligned} \|T_{\mathcal{G}^{v_{i_1}}}((P_{k_{\max}} f_l)_{l \in \mathcal{L}(v_{i_1})})\|_{p_{v_{i_1}}} &\lesssim 2^{k_{\max} \beta(v_{i_1}, l_1)} \|\Delta_{k_{\max}} f_{l_1}\|_{p_{l_1}} \sum_{\delta|_{(\mathcal{V}_{l_1}^{v_{i_1}})^c}} \prod_{\substack{l \in \mathcal{L}(v_{i_1}) \\ l \neq l_1}} \|D^{\delta^{-1}(l)} f_l\|_{p_l} \\ &\lesssim \|\Delta_{k_{\max}} D^{\beta(v_{i_1}, l_1)} f_{l_1}\|_{p_{l_1}} \sum_{\delta|_{(\mathcal{V}_{l_1}^{v_{i_1}})^c}} \prod_{\substack{l \in \mathcal{L}(v_{i_1}) \\ l \neq l_1}} \|D^{\delta^{-1}(l)} f_l\|_{p_l}, \end{aligned}$$

where the last inequality follows from (2.7). A similar reasoning gives

$$\|T_{\mathcal{G}^{v_{i_2}}}((P_{k_{\max}} f_l)_{l \in \mathcal{L}(v_{i_2})})\|_{p_{v_{i_2}}} \lesssim \|\Delta_{k_{\max}} D^{\beta(v_{i_2}, l_2)} f_{l_2}\|_{p_{l_2}} \sum_{\delta|_{(\mathcal{V}_{l_2}^{v_{i_2}})^c}} \prod_{\substack{l \in \mathcal{L}(v_{i_2}) \\ l \neq l_2}} \|D^{\delta^{-1}(l)} f_l\|_{p_l}.$$

For $i \neq i_1, i_2$, we use the estimate (5.20) implied from the inductive hypothesis (5.11). As a consequence, (5.18) can be majorized by

$$2^{k_{\max} \beta^{\tau_{\mathcal{G}}}} \|\Delta_{k_{\max}} D^{\beta(v_{i_1}, l_1)} f_{l_1}\|_{p_{l_1}} \|\Delta_{k_{\max}} D^{\beta(v_{i_2}, l_2)} f_{l_2}\|_{p_{l_2}} \sum_{\delta|_{(\mathcal{V}_{l_1}^{\tau_{\mathcal{G}}})^c \setminus \mathcal{V}_{l_2}^{v_{i_2}}}} \prod_{l \neq l_1, l_2} \|D^{\delta^{-1}(l)} f_l\|_{p_l},$$

which agrees with (5.9) due to the interpretation of notations (5.22) and (5.23).

- (3) We will now prove the third inductive statement corresponding to (5.10); in this case we need to take into account the more precise structure of the conical region $T_{\mathcal{G}}$ is restricted to.

- (a) **Case 1: The conical region R is of the form (5.1).**

Suppose that

$$\mathfrak{M}(R) = \{l_0\}, \quad \mathfrak{m}(R) = \{1 \leq l \leq n : l \neq l_0\},$$

and $l_0 \in \mathcal{L}(v_{i_0})$ for some $1 \leq i_0 \leq n_1$. Then $\sum_{k_{\max}} T_{\mathcal{G}}((P_{k_{\max}} f_l)_{1 \leq l \leq n})$, or more precisely

$$\sum_{k_{\max}} T_{\mathcal{G}}^R((P_{k_{\max}} f_l)_{1 \leq l \leq n})$$

concerns the frequency space localized to the conical region (5.1), on which we apply the *splitting of the root symbol* step introduced in Section 2.2 and used in Section 3.1:

$$(5.24) \quad \left| \sum_{l=1}^n \xi_l \right|^{\beta^{\tau_{\mathcal{G}}}} = \underbrace{\sum_{i \neq i_0} m_{C_{\beta}^{\tau_{\mathcal{G}}}} \left(\sum_{l \in \mathcal{L}(v_{i_0})} \xi_l, \sum_{l \notin \mathcal{L}(v_{i_0})} \xi_l \right)}_{\mathcal{E}_1} \cdot \sum_{l \in \mathcal{L}(v_{i_0})} \xi_l + \underbrace{\sum_{l \in \mathcal{L}(v_{i_0})} \xi_l^{\beta^{\tau_{\mathcal{G}}}}}_{\mathcal{E}_2},$$

where

$$m_{C_{\beta}^{\tau_{\mathcal{G}}}}(\tilde{\xi}_1, \tilde{\xi}_2) := \frac{|\tilde{\xi}_1 + \tilde{\xi}_2|^{\beta^{\tau_{\mathcal{G}}}} - |\tilde{\xi}_1|^{\beta^{\tau_{\mathcal{G}}}}}{\tilde{\xi}_2}.$$

Estimate of \mathcal{E}_1 :

The symbol denoted by \mathcal{E}_1 generates a commutator whose treatment builds on the approach described in Section 2.2. In order to perform the double Fourier series decomposition on one Whitney cube at a time, we need to further decompose $R_{k_{l_0}}$ (5.4) – the union of Whitney cubes at scale k_{l_0} . In particular,

$$R_{k_{l_0}} = R_{k_{l_0}}^+ \cup R_{k_{l_0}}^-,$$

where

$$R_{k_{l_0}}^+ := \{(\xi_1, \dots, \xi_n) : \xi_{l_0} \sim 2^{k_{l_0}}, |\xi_l| \ll 2^{k_{l_0}}\}, \quad R_{k_{l_0}}^- := \{(\xi_1, \dots, \xi_n) : \xi_{l_0} \sim -2^{k_{l_0}}, |\xi_l| \ll 2^{k_{l_0}}\}.$$

We restrict the symbol $m_{C_\beta^{\tau_g}}$ to $R_{k_{l_0}}^+$ and $R_{k_{l_0}}^-$ and denote them by $m_{C_\beta^{\tau_g}}^{k_{l_0},+}$ and $m_{C_\beta^{\tau_g}}^{k_{l_0},-}$ respectively. Then the double Fourier series decomposition yields

$$(5.25) \quad m_{C_\beta^{\tau_g}}^{k_{l_0},\pm} \left(\sum_{l \in \mathcal{L}(v_{i_0})} \xi_l, \sum_{l \notin \mathcal{L}(v_{i_0})} \xi_l \right) = \sum_{L_1, L_2 \in \mathbb{Z}} C_{L_1, L_2}^\pm 2^{k_{l_0}(\beta^{\tau_g} - 1)} e^{2\pi i L_1 \frac{\sum_{l \in \mathcal{L}(v_{i_0})} \xi_l}{2^{k_{l_0}}}} e^{2\pi i L_2 \frac{\sum_{l \notin \mathcal{L}(v_{i_0})} \xi_l}{2^{k_{l_0}}}},$$

which is essentially the same as (3.7) with a few natural adjustments: replacing 1 by l_0 and i_0 , $\xi_1 + \xi_2$ by $\sum_{l \in \mathcal{L}(v_{i_0})} \xi_l$ and $\xi_3 + \xi_4 + \xi_5$ by $\sum_{l \notin \mathcal{L}(v_{i_0})} \xi_l$. We remark that all the renormalized Fourier coefficients involved decay rapidly.

Without loss of generality, assume that

$$\mathfrak{M}(R) = \{1\}, \quad \mathfrak{m}(R) = \{1 \leq l \leq n : l \neq 1\}.$$

We will use k_1 specifically instead of k_{\max} and denote by R_{k_1} the union of Whitney cubes at scale k_1 in the cone R . For any fixed $L \in \mathbb{Z}$,

$$P_{k_1, L}(1) = \Delta_{k_1, L}, \quad P_{k_1, L}(l) = S_{k_1, L} \quad \text{for } l \neq 1.$$

As a consequence of the above two steps, we can rewrite the symbol \mathcal{E}_1 as

$$\mathcal{E}_1 =: \sum_{i \neq i_0} \mathcal{E}_1^i.$$

We focus on \mathcal{E}_1^2 and consider the multiplier associated to it:

$$(5.26) \quad \sum_{k_1} T_{\mathcal{E}_1^2} \left(T_{\mathcal{G}^{v_1}} (\Delta_{k_1} f_1, (S_{k_1} f_l)_{1 \neq l \in \mathcal{L}(v_1)}), T_{\mathcal{G}^{v_2}} ((S_{k_1} f_l)_{l \in \mathcal{L}(v_2)}), \dots, T_{\mathcal{G}^{v_{n_1}}} ((S_{k_1} f_l)_{l \in \mathcal{L}(v_{n_1})}) \right).$$

For the multilinear expression associated to the subtree \mathcal{G}^{v_2} , namely

$$T_{\mathcal{G}^{v_2}} ((S_{k_1} f_l)_{l \in \mathcal{L}(v_2)}),$$

we will further perform the paraproduct decomposition so as to focus on a fixed conical region denoted by $R(v_2)$. Let $\mathfrak{M}(R(v_2))$ denote a subset of $\mathcal{L}(v_2)$ defined similarly to (5.3) and $\mathfrak{m}(R(v_2)) = \mathcal{L}(v_2) \setminus \mathfrak{M}(R(v_2))$. Furthermore, we define

$$P_{k_{\max}(v_2)}(l) = \Delta_{k_{\max}(v_2)}, \quad \text{if } l \in \mathcal{L}(v_2) \cap \mathfrak{M}(R(v_2)) \quad \text{and} \quad P_{k_{\max}(v_2)}(l) = S_{k_{\max}(v_2)} \quad \text{if } l \in \mathcal{L}(v_2) \cap \mathfrak{m}(R(v_2)).$$

The localized multilinear expression can then be written as

$$(5.27) \quad \sum_{k_{\max}(v_2)} T_{\mathcal{G}^{v_2}}^{R(v_2)} ((P_{k_{\max}(v_2)} S_{k_1} f_l)_{l \in \mathcal{L}(v_2)}).$$

Since $\Delta_{k_{\max}(v_2)} S_{k_1} \neq 0$ if and only if $k_{\max}(v_2) \ll k_1$, we can restrict the sum in (5.27):

$$\sum_{k_{\max}(v_2) : k_{\max}(v_2) \ll k_1} T_{\mathcal{G}^{v_2}}^{R(v_2)} ((P_{k_{\max}(v_2)} f_l)_{l \in \mathcal{L}(v_2)}),$$

which abbreviates as

$$\sum_{k_{\max}(v_2) : k_{\max}(v_2) \ll k_1} T_{\mathcal{G}^{v_2}} ((P_{k_{\max}(v_2)} f_l)_{l \in \mathcal{L}(v_2)}).$$

As a consequence, (5.26) takes the form

$$(5.28) \quad \sum_{k_{\max}(v_2) \ll k_1} T_{\mathcal{E}_1^2} \left(T_{\mathcal{G}^{v_1}} (\Delta_{k_1} f_1, (S_{k_1} f_l)_{1 \neq l \in \mathcal{L}(v_1)}), T_{\mathcal{G}^{v_2}} ((P_{k_{\max}(v_2)} f_l)_{l \in \mathcal{L}(v_2)}), \dots, T_{\mathcal{G}^{v_{n_1}}} ((S_{k_1} f_l)_{l \in \mathcal{L}(v_{n_1})}) \right).$$

By applying the Fourier series (5.25) to (5.28), the latter becomes a sum of terms of the form

$$(5.29) \quad \sum_{L_1, L_2 \in \mathbb{Z}} C_{L_1, L_2}^\pm \sum_{k_{\max}(v_2) \ll k_1} 2^{k_{\max}(v_2)} 2^{k_1(\beta^{\tau_g} - 1)} T_{\mathcal{G}^{v_1}} \left(\Delta_{k_1, \pm, \frac{L_1}{2^{k_1}}} f_1, (S_{k_1, \frac{L_1}{2^{k_1}}} f_l)_{l \in \mathcal{L}(v_1)} \right)_{l \neq 1} \\ T_{\mathcal{G}^{v_2}} \left((P_{k_{\max}(v_2), \frac{L_2}{2^{k_1}}} f_l)_{l \in \mathcal{L}(v_2)} \right) \prod_{i=3}^{n_1} T_{\mathcal{G}^{v_i}} \left((S_{k_1, \frac{L_i}{2^{k_1}}} f_l)_{l \in \mathcal{L}(v_i)} \right)$$

Notice that the multipliers generated by \mathcal{E}_1^i , for $i > 2$, behave analogously. By an observation similar to (5.17) and thanks to the decay of the C_{L_1, L_2}^\pm coefficients, the L^r norm of (5.29) can be estimated by

$$(5.30) \quad \sum_{k_{\max}(v_2) \ll k_1} 2^{k_{\max}(v_2)} 2^{k_1 \cdot (\beta^{\tau_{\mathcal{G}} - 1})} \left\| T_{\mathcal{G}^{v_1}} \left(\Delta_{k_1, +} f_1, (S_{k_1} f_l)_{l \in \mathcal{L}(v_1)} \right) \right\|_{p_{v_1}} \\ \left\| T_{\mathcal{G}^{v_2}} \left((P_{k_{\max}(v_2)} f_l)_{l \in \mathcal{L}(v_2)} \right) \right\|_{p_{v_2}} \prod_{i=3}^{n_1} \left\| T_{\mathcal{G}^{v_i}} \left((S_{k_1} f_l)_{l \in \mathcal{L}(v_i)} \right) \right\|_{p_{v_i}}.$$

Since \mathcal{G}^{v_i} for $1 \leq i \leq n_1$ are subtrees of lower complexity, we invoke the inductive hypothesis (5.8) and (2.16), so that

$$(5.31) \quad \left\| T_{\mathcal{G}^{v_1}} \left(\Delta_{k_1, +} f_1, (S_{k_1} f_l)_{l \in \mathcal{L}(v_1)} \right) \right\|_{p_{v_1}} \lesssim \|\Delta_{k_1} D^{\beta(v_1, 1)} f_1\|_{p_1} \sum_{\delta|_{(\mathcal{V}_1^{v_1})^c}} \prod_{\substack{l \in \mathcal{L}(v_1) \\ l \neq 1}} \|D^{\delta^{-1}(l)} f_l\|_{p_l}$$

and

$$(5.32) \quad \left\| T_{\mathcal{G}^{v_2}} \left((P_{k_{\max}(v_2)} f_l)_{l \in \mathcal{L}(v_2)} \right) \right\|_{p_{v_2}} \lesssim \|\Delta_{k_{\max}(v_2)} D^{\beta(v_2, l_0)} f_{l_0}\|_{p_{l_0}} \sum_{\delta|_{(\mathcal{V}_{l_0}^{v_2})^c}} \prod_{l \neq l_0} \|D^{\delta^{-1}(l)} f_l\|_{p_l}.$$

Moreover, the inductive hypotheses (5.10) and thus (5.11) generate

$$(5.33) \quad \left\| T_{\mathcal{G}^{v_i}} \left((S_{k_1} f_l)_{l \in \mathcal{L}(v_i)} \right) \right\|_{p_{v_i}} \lesssim \sum_{\delta|_{\mathcal{V}, v_i}} \prod_{l \in \mathcal{L}(v_i)} \|D^{\delta^{-1}(l)} f_l\|_{p_l} \quad \text{for } i \neq 1, 2.$$

Applying the estimates (5.31), (5.32) and (5.33) to (5.30), we deduce that

$$\sum_{k_{\max}(v_2) \ll k_1} 2^{k_{\max}(v_2)} 2^{k_1 \cdot (\beta^{\tau_{\mathcal{G}} - 1})} \cdot \|\Delta_{k_1} D^{\beta(v_1, 1)} f_1\|_{p_1} \|\Delta_{k_{\max}(v_2)} D^{\beta(v_2, l_0)} f_{l_0}\|_{p_{l_0}} \sum_{\delta|_{\mathcal{V} \setminus \{\tau_{\mathcal{G}}\} \setminus \mathcal{V}_1^{v_1} \setminus \mathcal{V}_{l_0}^{v_2}}} \prod_{l \neq 1, l_0} \|D^{\delta^{-1}(l)} f_l\|_{p_l}.$$

We notice that

$$\sum_{k_{\max}(v_2) \ll k_1} 2^{k_{\max}(v_2)} 2^{k_1 \cdot (\beta^{\tau_{\mathcal{G}} - 1})} \cdot \|\Delta_{k_1} D^{\beta(v_1, 1)} f_1\|_{p_1} \|\Delta_{k_{\max}(v_2)} D^{\beta(v_2, l_0)} f_{l_0}\|_{p_{l_0}} \\ = \sum_{k_{\max}(v_2) \ll k_1} 2^{k_{\max}(v_2) \cdot (1-\epsilon)} 2^{k_1 \cdot (\beta^{\tau_{\mathcal{G}} - 1})} \|\Delta_{k_1} D^{\beta(v_1, 1)} f_1\|_{p_1} 2^{k_{\max}(v_2) \cdot \epsilon} \|\Delta_{k_{\max}(v_2)} D^{\beta(v_2, l_0)} f_{l_0}\|_{p_{l_0}}$$

for any $0 < \epsilon < \min(1, \beta^{\tau_{\mathcal{G}}})$. We can then distribute the derivatives in various ways to bound the above expression by

$$(5.34) \quad \sum_{k_1} \min \left(2^{k_1 \beta^{\tau_{\mathcal{G}}}} \|D^{\beta(v_1, 1)} f_1\|_{\dot{B}_{p_1, \infty}^0} \|D^{\beta(v_2, l_0)} f_{l_0}\|_{\dot{B}_{p_{l_0}, \infty}^0}, 2^{-k_1 \epsilon} \|D^{\beta(v_1, 1)} f_1\|_{\dot{B}_{p_1, \infty}^{\beta^{\tau_{\mathcal{G}}}}} \|D^{\beta(v_2, l_0)} f_{l_0}\|_{\dot{B}_{p_{l_0}, \infty}^{\epsilon}} \right).$$

By further optimizing in k_1 , this becomes

$$(5.35) \quad \left(\|D^{\beta(v_1, 1)} f_1\|_{\dot{B}_{p_1, \infty}^0} \|D^{\beta(v_2, l_0)} f_{l_0}\|_{\dot{B}_{p_{l_0}, \infty}^0} \right)^{\frac{\epsilon}{\beta^{\tau_{\mathcal{G}}} + \epsilon}} \left(\|D^{\beta(v_1, 1)} f_1\|_{\dot{B}_{p_1, \infty}^{\beta^{\tau_{\mathcal{G}}}}} \|D^{\beta(v_2, l_0)} f_{l_0}\|_{\dot{B}_{p_{l_0}, \infty}^{\epsilon}} \right)^{\frac{\beta^{\tau_{\mathcal{G}}}}{\beta^{\tau_{\mathcal{G}}} + \epsilon}}.$$

Now we invoke the interpolation inequality for Besov norms (2.11) – which is essentially a redistribution of the derivatives – to end up with

$$(5.36) \quad \left(\|D^{\beta(v_1, 1)} f_1\|_{\dot{B}_{p_1, \infty}^0} \|D^{\beta(v_2, l_0)} f_{l_0}\|_{\dot{B}_{p_{l_0}, \infty}^{\beta^{\tau_{\mathcal{G}}}}} \right)^{\frac{\epsilon}{\beta^{\tau_{\mathcal{G}}} + \epsilon}} \left(\|D^{\beta(v_1, 1)} f_1\|_{\dot{B}_{p_1, \infty}^{\beta^{\tau_{\mathcal{G}}}}} \|D^{\beta(v_2, l_0)} f_{l_0}\|_{\dot{B}_{p_{l_0}, \infty}^0} \right)^{\frac{\beta^{\tau_{\mathcal{G}}}}{\beta^{\tau_{\mathcal{G}}} + \epsilon}} \\ \lesssim \|D^{\beta^{\tau_{\mathcal{G}}} + \beta(v_1, 1)} f_1\|_{p_1} \|D^{\beta(v_2, l_0)} f_{l_0}\|_{p_{l_0}} + \|D^{\beta(v_1, 1)} f_1\|_{p_1} \|D^{\beta^{\tau_{\mathcal{G}}} + \beta(v_2, l_0)} f_{l_0}\|_{p_{l_0}}.$$

Estimate of \mathcal{E}_2 :

The multiplier generated by the symbol $|\sum_{l \in \mathcal{L}(v_{i_0})} \xi_l|^{\beta^{\tau_{\mathcal{G}}}}$ localized on the conical region (5.1) with $l_0 = 1$ is

$$(5.37) \quad \sum_{k_1} \left(D^{\beta^{\tau_{\mathcal{G}}}} T_{\mathcal{G}^{v_1}} \left(\Delta_{k_1} f_1, (S_{k_1} f_l)_{l \in \mathcal{L}(v_1)} \right) \right) \prod_{i=2}^{n_1} T_{\mathcal{G}^{v_i}} \left((S_{k_1} f_l)_{l \in \mathcal{L}(v_i)} \right).$$

We first simplify (5.37) using the high-low switch technique discussed in Sections 2.2 and 3. Denote by $\tilde{\mathcal{G}}^{v_1}$ the tree having the same structure as \mathcal{G}^{v_1} with the derivative β^{v_1} replaced by $\beta^{v_1} + \beta^{\mathfrak{r}_G}$ so that (5.37) can be rewritten as

$$(5.38) \quad \sum_{k_1} (T_{\tilde{\mathcal{G}}^{v_1}}(\Delta_{k_1} f_1, (S_{k_1} f_l)_{l \in \mathcal{L}(v_1)})) \prod_{i \neq 1} T_{\mathcal{G}^{v_i}}((S_{k_1} f_l)_{l \in \mathcal{L}(v_i)})$$

We then perform a finer³⁷ paraproduct decompositions on the functions in the subtrees \mathcal{G}^{v_i} for $i \neq 1$ so that (5.38) can be written as a finite sum of terms with the following form:

$$(5.39) \quad \begin{aligned} & \sum_{k_1} (T_{\tilde{\mathcal{G}}^{v_1}}(\Delta_{k_1} f_1, (S_{k_1} f_l)_{l \in \mathcal{L}(v_1)})) \prod_{i \neq 1} \sum_{k_{\max}(v_i)} T_{\mathcal{G}^{v_i}}((S_{k_1} P_{k_{\max}(v_i)} f_l)_{l \in \mathcal{L}(v_i)}), \\ & = \sum_{k_1} (T_{\tilde{\mathcal{G}}^{v_1}}(\Delta_{k_1} f_1, (S_{k_1} f_l)_{l \in \mathcal{L}(v_1)})) \prod_{i \neq 1} \sum_{k_{\max}(v_i) \ll k_1} T_{\mathcal{G}^{v_i}}((P_{k_{\max}(v_i)} f_l)_{l \in \mathcal{L}(v_i)}). \end{aligned}$$

We notice that the equation holds because the conical decomposition on the subtree \mathcal{G}^{v_2} gives $\Delta_{k_{\max}(v_2)}$ for $l \in \mathfrak{M}(R(v_2))$ and $S_{k_1} \Delta_{k_{\max}(v_i)} \neq 0$ if and only if $k_{\max}(v_i) \ll k_1$. We can now apply the high-low switch to swap the role of k_1 and $k_{\max}(v_i)$ and rewrite (5.39) as

$$\begin{aligned} & \sum_{k_1} (T_{\tilde{\mathcal{G}}^{v_1}}(\Delta_{k_1} f_1, (S_{k_1} f_l)_{l \in \mathcal{L}(v_1)})) \prod_{l \neq 1} \sum_{k_{\max}(v_i)} T_{\mathcal{G}^{v_i}}((P_{k_{\max}(v_i)} f_l)_{l \in \mathcal{L}(v_i)}) \\ & - \sum_{k_{\max}(v_2) > k_1} (T_{\tilde{\mathcal{G}}^{v_1}}(\Delta_{k_1} f_1, (S_{k_1} f_l)_{l \in \mathcal{L}(v_1)})) T_{\mathcal{G}^{v_2}}((P_{k_{\max}(v_2)} f_l)_{l \in \mathcal{L}(v_2)}) \cdot \prod_{i \neq 1, 2} \sum_{k_{\max}(v_i)} T_{\mathcal{G}^{v_i}}((P_{k_{\max}(v_i)} f_l)_{l \in \mathcal{L}(v_i)}) \end{aligned}$$

\pm similar terms $:= I - II \pm$ similar terms.

The first term I can be estimated by using inductive hypothesis (5.10) on

$$\sum_{k_1} (T_{\tilde{\mathcal{G}}^{v_1}}(\Delta_{k_1} f_1, (S_{k_1} f_l)_{l \in \mathcal{L}(v_1)})),$$

and on

$$\sum_{k_{\max}(v_i)} T_{\mathcal{G}^{v_i}}((P_{k_{\max}(v_i)} f_l)_{l \in \mathcal{L}(v_i)})$$

for $i \neq 1$. The second term II requests a more careful treatment. We recall that

$$T_{\tilde{\mathcal{G}}^{v_1}}(\Delta_{k_1} f_1, (S_{k_1} f_l)_{l \in \mathcal{L}(v_1)}) = D^{\beta^{\mathfrak{r}_G}} T_{\mathcal{G}^{v_1}}(\Delta_{k_1} f_1, (S_{k_1} f_l)_{l \in \mathcal{L}(v_1)}),$$

where $\tilde{\mathcal{G}}^{v_1}$ is a tree with the differential operator associated to the root v_1 being

$$D^{\beta^{v_1} + \beta^{\mathfrak{r}_G}}.$$

We invoke the inductive hypothesis (5.31):

$$\|T_{\tilde{\mathcal{G}}^{v_1}}(\Delta_{k_1} f_1, (S_{k_1} f_l)_{l \in \mathcal{L}(v_1)})\|_{p_{v_1}} \lesssim 2^{k_1 \beta^{\mathfrak{r}_G}} \|\Delta_{k_1} D^{\beta(v_1, 1)} f_1\|_{p_1} \sum_{\delta | (\mathcal{V}_1^{v_1})^c} \prod_{l \in \mathcal{L}(v_1)} \|D^{\delta^{-1}(l)} f_l\|_{p_l}.$$

When combined with (5.32), we deduce that

$$\|II\|_r \lesssim \sum_{k_{\max}(v_2) > k_1} 2^{k_1 \beta^{\mathfrak{r}_G}} \|\Delta_{k_1} D^{\beta(v_1, 1)} f_1\|_{p_1} \|\Delta_{k_{\max}(v_2)} D^{\beta(v_2, l_0)} f_{l_0}\|_{p_{l_0}} \sum_{\delta | (\mathcal{V}_1^{v_1})^c \cup (\mathcal{V}_{l_0}^{v_2})^c} \prod_{l \neq 1, l_0} \|D^{\delta^{-1}(l)} f_l\|_{p_l}.$$

A similar computation specified in Section 3 yields

$$(5.40) \quad \sum_{k_{\max}(v_2) > k_1} 2^{k_1 \beta^{\mathfrak{r}_G}} \|\Delta_{k_1} D^{\beta(v_1, 1)} f_1\|_{p_1} \|\Delta_{k_{\max}(v_2)} D^{\beta(v_2, l_0)} f_{l_0}\|_{p_{l_0}} \lesssim \sum_{k_{\max}(v_2)} 2^{k_{\max}(v_2) \beta^{\mathfrak{r}_G}} \|D^{\beta(v_1, 1)} f_1\|_{\dot{B}_{p_1}^0} \|D^{\beta(v_2, l_0)} f_{l_0}\|_{\dot{B}_{p_{l_0}}^0}.$$

Meanwhile, (5.40) can also be estimated by

$$\begin{aligned} & \sum_{k_{\max}(v_2) > k_1} 2^{k_1(\beta^{\mathfrak{r}_G} - \epsilon)} 2^{-k_{\max}(v_2) \beta^{\mathfrak{r}_G}} (2^{k_1 \epsilon} \|\Delta_{k_1} D^{\beta(v_1, 1)} f_1\|_{p_1}) (2^{k_{\max}(v_2) \beta^{\mathfrak{r}_G}} \|\Delta_{k_{\max}(v_2)} D^{\beta(v_2, l_0)} f_{l_0}\|_{p_{l_0}}) \\ & \lesssim \sum_{k_{\max}(v_2)} 2^{-k_{\max}(v_2) \epsilon} \|D^{\beta(v_1, 1)} f_1\|_{\dot{B}_{p_1}^\epsilon} \|D^{\beta(v_2, l_0)} f_{l_0}\|_{\dot{B}_{p_{l_0}}^{\beta^{\mathfrak{r}_G}}}. \end{aligned}$$

³⁷Notice that as we perform this step, we also restrict ourselves to certain conical regions associated to each subtree \mathcal{G}^{v_i} , for $i \neq 1$.

The optimization and interpolation can be applied to conclude that $\|II\|_r$ is bounded above by

$$\left(\|D^{\beta_{\mathcal{G}} + \beta(v_1, 1)} f_1\|_{p_1} \|D^{\beta(v_2, l_0)} f_{l_0}\|_{p_{l_0}} + \|D^{\beta(v_1, 1)} f_1\|_{p_1} \|D^{\beta_{\mathcal{G}} + \beta(v_2, l_0)} f_{l_0}\|_{p_{l_0}} \right) \sum_{\delta|_{(\mathcal{V}_1^{v_1})^c \cup (\mathcal{V}_{l_0}^{v_2})^c}} \prod_{\substack{l \in \mathcal{L}(v_1) \\ l \neq 1, l_0}} \|D^{\delta^{-1}(l)} f_l\|_{p_l}.$$

We thus have arrived at the expression (5.10) claimed in the inductive statement. With this, we end the proof of Proposition 5.1, if R is of the form (5.1).

(b) **Case 2: The conical region R is of the form (5.2).**

We notice that (5.10) is a direct consequence of (5.9) for the given tree \mathcal{G} . In particular, we apply (5.9) on the union of Whitney cubes at a fixed scale to derive

$$(5.41) \quad \begin{aligned} & \left\| \sum_{k_{\max}} T_{\mathcal{G}}((P_{k_{\max}} f_l)_{1 \leq l \leq n}) \right\|_r \lesssim \sum_{k_{\max}} \|T_{\mathcal{G}}((P_{k_{\max}} f_l)_{1 \leq l \leq n})\|_r \\ & \lesssim \sum_{k_{\max}} 2^{k_{\max} \cdot \beta(\tau_{\mathcal{G}}, v^{l_1, l_2})} \|\Delta_{k_{\max}} D^{\beta(w^{l_1, l_1})} f_{l_1}\|_{p_{l_1}} \|\Delta_{k_{\max}} D^{\beta(w^{l_2, l_2})} f_{l_2}\|_{p_{l_2}} \sum_{\delta|_{(\mathcal{V}_{l_1}^{\tau_{\mathcal{G}}})^c \setminus \mathcal{V}_{l_2}^{w^{l_2}}} } \prod_{l \neq l_1, l_2} \|D^{\delta^{-1}(l)} f_l\|_{p_l}. \end{aligned}$$

We then distribute the derivatives as before:

$$(5.42) \quad \begin{aligned} & \sum_{k_{\max}} 2^{k_{\max} \cdot \beta(\tau_{\mathcal{G}}, v^{l_1, l_2})} \|\Delta_{k_{\max}} D^{\beta(w^{l_1, l_1})} f_{l_1}\|_{p_{l_1}} \|\Delta_{k_{\max}} D^{\beta(w^{l_2, l_2})} f_{l_2}\|_{p_{l_2}} \\ & \leq \sum_{k_{\max}} \min(2^{k_{\max} \cdot \beta(\tau_{\mathcal{G}}, v^{l_1, l_2})} \|D^{\beta(w^{l_1, l_1})} f_{l_1}\|_{\dot{B}_{p_{l_1}}^0} \|D^{\beta(w^{l_2, l_2})} f_{l_2}\|_{\dot{B}_{p_{l_2}}^0}, \\ & \quad 2^{-k_{\max} \cdot \beta(\tau_{\mathcal{G}}, v^{l_1, l_2})} \|D^{\beta(w^{l_1, l_1})} f_{l_1}\|_{\dot{B}_{p_{l_1}}^{\beta(\tau_{\mathcal{G}}, v^{l_1, l_2})}} \|D^{\beta(w^{l_2, l_2})} f_{l_2}\|_{\dot{B}_{p_{l_2}}^{\beta(\tau_{\mathcal{G}}, v^{l_1, l_2})}}) \\ & \lesssim \|D^{\beta(\tau_{\mathcal{G}}, l_1)} f_{l_1}\|_{p_{l_1}} \|D^{\beta(w^{l_2, l_2})} f_{l_2}\|_{p_{l_2}} + \|D^{\beta(w^{l_1, l_1})} f_{l_1}\|_{p_{l_1}} \|D^{\beta(\tau_{\mathcal{G}}, l_2)} f_{l_2}\|_{p_{l_2}}. \end{aligned}$$

We plug (5.42) into (5.41) and obtain the estimate (5.10) claimed in the inductive statement. \square

5.2. Bi-parameter flag Leibniz rule. We will extend our inductive argument to bi-parameter flag Leibniz rules of arbitrary complexity in dimension³⁸ one. We follow the same notation as before, except for the addition of subscription to indicate which parameter is involved. For example, for any $v \in \mathcal{V}$, β_j^v represents the derivative for the j -th parameter (for $j = 1, 2$). As in the one-parameter setting, the frequency space for each parameter can be decomposed into conical regions of the form (5.1) or (5.2). We correspondingly define the maps \mathfrak{M}_j and \mathfrak{m}_j for $j = 1, 2$ on the collection of conical regions for the j -th parameter:

$$\begin{aligned} \mathfrak{M}_1(R) &:= \{1 \leq l \leq n : (\xi_1, \dots, \xi_n) \in R, \quad |\xi_l| \sim \max_{1 \leq l' \leq n} |\xi_{l'}|\}, & \mathfrak{m}_1(R) &:= \{1 \leq l \leq n : (\xi_1, \dots, \xi_n) \in R, \quad |\xi_l| \ll \max_{1 \leq l' \leq n} |\xi_{l'}|\}, \\ \mathfrak{M}_2(R') &:= \{1 \leq l \leq n : (\eta_1, \dots, \eta_n) \in R', \quad |\eta_l| \sim \max_{1 \leq l' \leq n} |\eta_{l'}|\}, & \mathfrak{m}_2(R') &:= \{1 \leq l \leq n : (\eta_1, \dots, \eta_n) \in R', \quad |\eta_l| \ll \max_{1 \leq l' \leq n} |\eta_{l'}|\}, \end{aligned}$$

where R denotes a conical region for the first parameter and R' a conical region for the second parameter.

Fix any integer k ; we define the projections $P_k^{(j)}$ ($j = 1, 2$) depending on l as follows:

$$P_k^{(1)}(l) := \begin{cases} \Delta_k^{(1)} & \text{if } l \in \mathfrak{M}_1(R) \\ S_k^{(1)} & \text{if } l \in \mathfrak{m}_1(R), \end{cases} \quad \text{and} \quad P_k^{(2)}(l) := \begin{cases} \Delta_k^{(2)} & \text{if } l \in \mathfrak{M}_2(R') \\ S_k^{(2)} & \text{if } l \in \mathfrak{m}_2(R'). \end{cases}$$

Similarly, we define $P_{k,L}^{(j)}$ ($j = 1, 2$) for fixed $k \in \mathbb{Z}$ and $L \in \mathbb{R}$ by

$$P_{k,L}^{(1)}(l) := \begin{cases} \Delta_{k,L}^{(1)} & \text{if } l \in \mathfrak{M}_1(R) \\ S_{k,L}^{(1)} & \text{if } l \in \mathfrak{m}_1(R), \end{cases} \quad \text{and} \quad P_{k,L}^{(2)}(l) := \begin{cases} \Delta_{k,L}^{(2)} & \text{if } l \in \mathfrak{M}_2(R') \\ S_{k,L}^{(2)} & \text{if } l \in \mathfrak{m}_2(R'). \end{cases}$$

Let k_{\max} and m_{\max} specify the Whitney cubes at scales k_{\max} and m_{\max} in the cone R for the first parameter and R' for the second parameter. We can thus express the multilinear expression localized to a union of Whitney rectangles³⁹ at the scale $k_{\max} \times m_{\max}$ in the conical region $R \times R'$ as

$$T_{\mathcal{G}}^{R \times R'}((P_{k_{\max}}^{(1)} P_{m_{\max}}^{(2)} f_l)_{1 \leq l \leq n}).$$

Since we will always focus on a certain conical region, the above expression will be abbreviated as

$$T_{\mathcal{G}}((P_{k_{\max}}^{(1)} P_{m_{\max}}^{(2)} f_l)_{1 \leq l \leq n}).$$

³⁸As mentioned in the beginning of Section 5, the methods are adaptable to higher dimensions, in a straightforward manner.

³⁹In this case, a Whitney rectangle is simply the product of two Whitney cubes, one in each parameter.

The multilinear expression restricted to this cone can then be written as

$$\sum_{k_{\max}, m_{\max} \in \mathbb{Z}} T_{\mathcal{G}}((P_{k_{\max}}^{(1)} P_{m_{\max}}^{(2)} f_l)_{1 \leq l \leq n}).$$

Similarly to the one-parameter analysis, the cone decomposition can also be applied to a subtree \mathcal{G}^v with $v \in \mathcal{V}$ for both parameters and let $k_{\max}(v)$ and $m_{\max}(v)$ specify the Whitney cubes in these conical regions for \mathcal{G}^v . We will follow the abbreviation in the one-parameter setting so that k_{\max} and m_{\max} refer to $k_{\max}(\tau_{\mathcal{G}})$ and $m_{\max}(\tau_{\mathcal{G}})$ respectively.

Define the sum of partial derivatives from the vertex v to the leaf f_l by

$$\beta_j(v, l) := \sum_{w \in \mathcal{V}_l^v} \beta_j^w, \quad \text{for } j = 1, 2.$$

With some abuse of notation, if $v = f_l$ is a leaf, then $\beta_j(v, l) = 0$.

Our goal is to prove bi-parameter versions of the inductive statements in Proposition 5.1. The statements (1), (2) and (3) below describe estimates for the multi-linear expression $T_{\mathcal{G}}(P_{k_{\max}}^{(1)} P_{m_{\max}}^{(2)} f_l)_{1 \leq l \leq n}$ on a union of Whitney rectangles (the product of two Whitney cubes) at a fixed scale $k_{\max} \times m_{\max}$ localized to a conical region (the product of two conical regions, one for each parameter); several possibilities need to be investigated:

- the Whitney cubes can be in any conical region for both parameters;
- for at least one parameter its Whitney cube is located in a “diagonal” conical region;
- the Whitney cubes are in diagonal conical regions for both parameters.

One observes that (1) - (3) impose conditions from weak to strong so that (3) implies (2) and (2) leads to (1). The statements (4) and (5) concern Leibniz rules when the frequency space for one parameter is localized on a conical region and on a union of Whitney cubes at a fixed scale for the other parameter. (4) corresponds to the case when the Whitney cubes for one parameter are in any conical region while (5) describes the case when the Whitney cubes lie in a diagonal conical region. (6) is the Leibniz rule when the frequency spaces for both parameters are localized on a conical region.

Proposition 5.3. *Suppose that all the Lebesgue exponents in the inductive statement satisfy the condition described in Theorem 1.1 and that $T_{\mathcal{G}}$ is restricted to the cone denoted by R for the first parameter and R' for the second parameter.*

(1) Suppose that $l_0 \in \mathfrak{M}_1(R)$ and $l'_0 \in \mathfrak{M}_2(R')$.

(a) If $l_0 = l'_0$, then

(5.43)

$$\|T_{\mathcal{G}}(P_{k_{\max}}^{(1)} P_{m_{\max}}^{(2)} f_l)_{1 \leq l \leq n}\|_{\vec{r}} \lesssim 2^{k_{\max} \beta_1(\tau_{\mathcal{G}}, l_0)} 2^{m_{\max} \beta_2(\tau_{\mathcal{G}}, l_0)} \|\Delta_{k_{\max}}^{(1)} \Delta_{m_{\max}}^{(2)} f_{l_0}\|_{\vec{p}_{l_0}} \sum_{\delta_1 \otimes \delta_2|_{(\mathcal{V}_{l_0}^{\tau_{\mathcal{G}}})^c}} \prod_{l \neq l_0} \|D_{(1)}^{\delta_1^{-1}(l)} D_{(2)}^{\delta_2^{-1}(l)} f_l\|_{\vec{p}_l}.$$

(b) If $l_0 \neq l'_0$, then

$$(5.44) \quad \|T_{\mathcal{G}}(P_{k_{\max}}^{(1)} P_{m_{\max}}^{(2)} f_l)_{1 \leq l \leq n}\|_{\vec{r}} \lesssim 2^{k_{\max} \beta_1(\tau_{\mathcal{G}}, l_0)} 2^{m_{\max} \beta_2(\tau_{\mathcal{G}}, l'_0)} \cdot \sum_{\substack{\delta_1|_{(\mathcal{V}_{l_0}^{\tau_{\mathcal{G}}})^c} \\ \delta_2|_{(\mathcal{V}_{l'_0}^{\tau_{\mathcal{G}}})^c}}} \|\Delta_{k_{\max}}^{(1)} D_{(2)}^{\delta_1^{-1}(l_0)} f_{l_0}\|_{\vec{p}_{l_0}} \|\Delta_{m_{\max}}^{(2)} D_{(1)}^{\delta_1^{-1}(l'_0)} f_{l'_0}\|_{\vec{p}_{l'_0}} \prod_{l \neq l_0, l'_0} \|D_{(1)}^{\delta_1^{-1}(l)} D_{(2)}^{\delta_2^{-1}(l)} f_l\|_{\vec{p}_l}.$$

(2) Suppose that $l_1, l_2 \in \mathfrak{M}_1(R)$ with $l_1 \neq l_2$ and $l'_0 \in \mathfrak{M}_2(R')$.

(a) If $l_1 = l'_0$, then

$$(5.45) \quad \|T_{\mathcal{G}}(P_{k_{\max}}^{(1)} P_{m_{\max}}^{(2)} f_l)_{1 \leq l \leq n}\|_{\vec{r}} \lesssim 2^{k_{\max} \beta_1(\tau_{\mathcal{G}}, v^{l_1, l_2})} 2^{m_{\max} \beta_2(\tau_{\mathcal{G}}, l_1)} \|\Delta_{k_{\max}}^{(1)} \Delta_{m_{\max}}^{(2)} D_{(1)}^{\beta_1(w^{l_1, l_1})} f_{l_1}\|_{\vec{p}_{l_1}} \cdot \sum_{\substack{\delta_1|_{(\mathcal{V}_{l_1}^{\tau_{\mathcal{G}}})^c \setminus \mathcal{V}_{l_2}^{w^{l_2}}} \\ \delta_2|_{(\mathcal{V}_{l_1}^{\tau_{\mathcal{G}}})^c}}} \|\Delta_{k_{\max}}^{(1)} D_{(1)}^{\beta_1(w^{l_2, l_2})} D_{(2)}^{\delta_2^{-1}(l_2)} f_{l_2}\|_{\vec{p}_{l_2}} \prod_{l \neq l_1, l_2} \|D_{(1)}^{\delta_1^{-1}(l)} D_{(2)}^{\delta_2^{-1}(l)} f_l\|_{\vec{p}_l}.$$

(b) If $l_1, l_2 \neq l'_0$, then

$$\|T_{\mathcal{G}}(P_{k_{\max}}^{(1)} P_{m_{\max}}^{(2)} f_l)_{1 \leq l \leq n}\|_{\vec{r}} \lesssim 2^{k_{\max} \beta_1(\tau_{\mathcal{G}}, v^{l_1, l_2})} 2^{m_{\max} \beta_2(\tau_{\mathcal{G}}, l'_0)} \prod_{l \neq l_1, l_2} \|D_{(1)}^{\delta_1^{-1}(l)} D_{(2)}^{\delta_2^{-1}(l)} f_l\|_{\vec{p}_l}$$

$$\cdot \sum_{\substack{\delta_1 |_{(\mathcal{V}_{l_1}^{\mathcal{G}})^c \setminus \mathcal{V}_{l_2}^{w^{l_2}}} \\ \delta_2 |_{(\mathcal{V}_{l'_0}^{\mathcal{G}})^c}}} \|\Delta_{k_{\max}}^{(1)} D_{(1)}^{\beta_1(w^{l_1}, l_1)} D_{(2)}^{\delta_2^{-1}(l_1)} f_{l_1}\|_{\bar{p}_{l_1}} \|\Delta_{k_{\max}}^{(1)} D_{(1)}^{\beta_1(w^{l_2}, l_2)} D_{(2)}^{\delta_2^{-1}(l_2)} f_{l_2}\|_{\bar{p}_{l_2}} \|\Delta_{m_{\max}}^{(2)} D_{(1)}^{\delta_1^{-1}(l'_0)} f_{l'_0}\|_{p_{l'_0}}.$$

(3) Suppose that $l_1, l_2 \in \mathfrak{M}_1(R)$ with $l_1 \neq l_2$ and $l'_1, l'_2 \in \mathfrak{M}_2(R')$ with $l'_1 \neq l'_2$.

(a) If $l_1 = l'_1$ and $l_2 = l'_2$, then

$$\|T_{\mathcal{G}}(P_{k_{\max}}^{(1)} P_{m_{\max}}^{(2)} f_l)_{1 \leq l \leq n}\|_{\bar{r}} \lesssim 2^{k_{\max} \beta_1(\mathcal{V}_{\mathcal{G}}, v^{l_1, l_2})} 2^{m_{\max} \beta_2(\mathcal{V}_{\mathcal{G}}, v^{l_1, l_2})} \sum_{\delta_1 \otimes \delta_2 |_{(\mathcal{V}_{l_1}^{\mathcal{G}})^c \setminus \mathcal{V}_{l_2}^{w^{l_2}}}} \prod_{l \neq l_1, l_2} \|D_{(1)}^{\delta_1^{-1}(l)} D_{(2)}^{\delta_2^{-1}(l)} f_l\|_{\bar{p}_l} \\ \cdot \|\Delta_{k_{\max}}^{(1)} \Delta_{m_{\max}}^{(2)} D_{(1)}^{\beta_1(w^{l_1}, l_1)} D_{(2)}^{\beta_2(w^{l_1}, l_1)} f_{l_1}\|_{\bar{p}_{l_1}} \|\Delta_{k_{\max}}^{(1)} \Delta_{m_{\max}}^{(2)} D_{(1)}^{\beta_1(w^{l_2}, l_2)} D_{(2)}^{\beta_2(w^{l_2}, l_2)} f_{l_2}\|_{\bar{p}_{l_2}}.$$

(b) If $l_1 = l'_1$ and $l_2 \neq l'_2$, then

$$\|T_{\mathcal{G}}(P_{k_{\max}}^{(1)} P_{m_{\max}}^{(2)} f_l)_{1 \leq l \leq n}\|_{\bar{r}} \lesssim 2^{k_{\max} \beta_1(\mathcal{V}_{\mathcal{G}}, v^{l_1, l_2})} 2^{m_{\max} \beta_2(\mathcal{V}_{\mathcal{G}}, v^{l_1, l'_2})} \|\Delta_{k_{\max}}^{(1)} \Delta_{m_{\max}}^{(2)} D_{(1)}^{\beta_1(w^{l_1}, l_1)} D_{(2)}^{\beta_2(w^{l_1}, l_1)} f_{l_1}\|_{\bar{p}_{l_1}} \\ \cdot \sum_{\substack{\delta_1 |_{(\mathcal{V}_{l_1}^{\mathcal{G}})^c \setminus \mathcal{V}_{l_2}^{w^{l_2}}} \\ \delta_2 |_{(\mathcal{V}_{l'_1}^{\mathcal{G}})^c \setminus \mathcal{V}_{l'_2}^{w^{l'_2}}}}} \|\Delta_{k_{\max}}^{(1)} D_{(1)}^{\beta_1(w^{l_2}, l_2)} D_{(2)}^{\delta_2^{-1}(l_2)} f_{l_2}\|_{\bar{p}_{l_2}} \|\Delta_{m_{\max}}^{(2)} D_{(1)}^{\delta_1^{-1}(l'_2)} D_{(2)}^{\beta_2(w^{l'_2}, l'_2)} f_{l'_2}\|_{p_{l'_2}} \prod_{l \neq l_1, l_2, l'_2} \|D_{(1)}^{\delta_1^{-1}(l)} D_{(2)}^{\delta_2^{-1}(l)} f_l\|_{\bar{p}_l}.$$

(c) If $l_1 \neq l'_1$ and $l_2 \neq l'_2$, then

$$\|T_{\mathcal{G}}(P_{k_{\max}}^{(1)} P_{m_{\max}}^{(2)} f_l)_{1 \leq l \leq n}\|_{\bar{r}} \lesssim 2^{k_{\max} \beta_1(\mathcal{V}_{\mathcal{G}}, v^{l_1, l_2})} 2^{m_{\max} \beta_2(\mathcal{V}_{\mathcal{G}}, v^{l'_1, l'_2})} \prod_{l \neq l_1, l_2, l'_1, l'_2} \|D_{(1)}^{\delta_1^{-1}(l)} D_{(2)}^{\delta_2^{-1}(l)} f_l\|_{\bar{p}_l} \\ \cdot \sum_{\substack{\delta_1 |_{(\mathcal{V}_{l_1}^{\mathcal{G}})^c \setminus \mathcal{V}_{l_2}^{w^{l_2}}} \\ \delta_2 |_{(\mathcal{V}_{l'_1}^{\mathcal{G}})^c \setminus \mathcal{V}_{l'_2}^{w^{l'_2}}}}} \|\Delta_{k_{\max}}^{(1)} D_{(1)}^{\beta_1(w^{l_1}, l_1)} D_{(2)}^{\delta_2^{-1}(l_1)} f_{l_1}\|_{\bar{p}_{l_1}} \|\Delta_{m_{\max}}^{(2)} D_{(1)}^{\delta_1^{-1}(l'_1)} D_{(2)}^{\beta_2(w^{l'_1}, l'_1)} f_{l'_1}\|_{p_{l'_1}} \\ \cdot \|\Delta_{k_{\max}}^{(1)} D_{(1)}^{\beta_1(w^{l_2}, l_2)} D_{(2)}^{\delta_2^{-1}(l_2)} f_{l_2}\|_{\bar{p}_{l_2}} \|\Delta_{m_{\max}}^{(2)} D_{(1)}^{\delta_1^{-1}(l'_2)} D_{(2)}^{\beta_2(w^{l'_2}, l'_2)} f_{l'_2}\|_{p_{l'_2}}.$$

(4) Suppose that $l_0 \in \mathfrak{M}_1(R)$. Then

$$(5.46) \quad \left\| \sum_{m_{\max} \in \mathbb{Z}} T_{\mathcal{G}}(P_{k_{\max}}^{(1)} P_{m_{\max}}^{(2)} f_l)_{1 \leq l \leq n} \right\|_{\bar{r}} \lesssim 2^{k_{\max} \beta_1(\mathcal{V}_{\mathcal{G}}, l_0)} \sum_{\substack{\delta_2 \\ \delta_2 |_{(\mathcal{V}_{l'_0}^{\mathcal{G}})^c}}} \|\Delta_{k_{\max}}^{(1)} D_{(2)}^{\delta_2^{-1}(l_0)} f_{l_0}\|_{\bar{p}_{l_0}} \prod_{l \neq l_0} \|D_{(1)}^{\delta_1^{-1}(l)} D_{(2)}^{\delta_2^{-1}(l)} f_l\|_{\bar{p}_l}.$$

(5) Suppose that $l_1, l_2 \in \mathfrak{M}_1(R)$ with $l_1 \neq l_2$. Then

$$\left\| \sum_{m_{\max} \in \mathbb{Z}} T_{\mathcal{G}}(P_{k_{\max}}^{(1)} P_{m_{\max}}^{(2)} f_l)_{1 \leq l \leq n} \right\|_{\bar{r}} \lesssim 2^{k_{\max} \beta_1(\mathcal{V}_{\mathcal{G}}, v^{l_1, l_2})} \prod_{l \neq l_1, l_2} \|D_{(1)}^{\delta_1^{-1}(l)} D_{(2)}^{\delta_2^{-1}(l)} f_l\|_{\bar{p}_l} \\ \cdot \sum_{\substack{\delta_2 \\ \delta_2 |_{(\mathcal{V}_{l_1}^{\mathcal{G}})^c \setminus \mathcal{V}_{l_2}^{w^{l_2}}}}} \|\Delta_{k_{\max}}^{(1)} D_{(1)}^{\beta_1(w^{l_1}, l_1)} D_{(2)}^{\delta_2^{-1}(l_1)} f_{l_1}\|_{\bar{p}_{l_1}} \|\Delta_{k_{\max}}^{(1)} D_{(1)}^{\beta_1(w^{l_2}, l_2)} D_{(2)}^{\delta_2^{-1}(l_2)} f_{l_2}\|_{\bar{p}_{l_2}}.$$

(6) If we sum over the whole conical regions, we have

$$(5.47) \quad \left\| \sum_{\substack{k_{\max} \in \mathbb{Z} \\ m_{\max} \in \mathbb{Z}}} T_{\mathcal{G}}(P_{k_{\max}}^{(1)} P_{m_{\max}}^{(2)} f_l)_{1 \leq l \leq n} \right\|_{\bar{r}} \lesssim \sum_{\delta} \prod_l \|D_{(1)}^{\delta_1^{-1}(l)} D_{(2)}^{\delta_2^{-1}(l)} f_l\|_{\bar{p}_l}.$$

Remark 5.4. (i) Observe that the cases summarized in (2) and (3) are not exhaustive but typical: other cases not explicitly stated (such as $l_2 = l'_0$ in (2) and $l_1 = l'_2$, $l_2 \neq l'_1$ in (3)) can be estimated analogously.

(ii) Due to symmetry, the term

$$\sum_{k_{\max} \in \mathbb{Z}} T_{\mathcal{G}}(P_{k_{\max}}^{(1)} P_{m_{\max}}^{(2)} f_l)_{1 \leq l \leq n}$$

is to be treated in the same way as statements (4) or (5) – based on the structure of the conical regions.

(iii) We realize that the semi-localized operator

$$(5.48) \quad T_{\mathcal{G}}((P_{k_{\max}}^{(1)} f_l)_{1 \leq l \leq n})$$

can be written as a finite sum of terms appearing in the left-hand side of (5.46), so that it satisfies the same estimate:

$$(5.49) \quad \| (5.48) \|_{\vec{r}} \lesssim 2^{k_{\max} \cdot \beta_1(\mathbf{r}_G, l_0)} \sum_{\substack{\delta_2 \\ \delta_1 | (\mathbf{v}_{l_0}^{\mathbf{r}_G})^c}} \|\Delta_{k_{\max}}^{(1)} D_{(2)}^{\delta_2^{-1}(l_0)} f_{l_0}\|_{\vec{p}_{l_0}} \prod_{l \neq l_0} \|D_{(1)}^{\delta_1^{-1}(l)} D_{(2)}^{\delta_2^{-1}(l)} f_l\|_{\vec{p}_l}.$$

By symmetry, a similar estimate holds for $T_G((P_{m_{\max}}^{(2)} f_l)_{1 \leq l \leq n})$.

(iv) We notice that as before (5.47) generates the following global Leibniz rule:

$$(5.50) \quad \|T_G((f_l)_{1 \leq l \leq n})\|_{\vec{r}} \lesssim \sum_{\delta} \prod_l \|D_{(1)}^{\delta_1^{-1}(l)} D_{(2)}^{\delta_2^{-1}(l)} f_l\|_{\vec{p}_l}.$$

and the Leibniz rule corresponding to a cone localization in the first parameter:

$$(5.51) \quad \left\| \sum_{k_{\max} \in \mathbb{Z}} T_G((P_{k_{\max}}^{(1)} f_l)_{1 \leq l \leq n}) \right\|_{\vec{r}} \lesssim \sum_{\delta} \prod_l \|D_{(1)}^{\delta_1^{-1}(l)} D_{(2)}^{\delta_2^{-1}(l)} f_l\|_{\vec{p}_l}.$$

Due to the range of different conical regions for each parameter, and hence to the various ways the root symbol can split, more auxiliary inductive statements (necessary for proving Theorem 1.1 in the bi-parameter case) appear. The strategy of the proof is however the same as in the previous Section 5.1, and we will especially focus on two aspects: the *splitting of the root symbol* (when the corresponding cone is of the type (5.1)) and the *Fourier series decompositions*, which allows to systematically reduce the estimation to subtrees of lower complexity. Because of this, the proof of many of these auxiliary statements will be left to the reader.

Proof of Proposition 5.3. (1) We first illustrate the proof for (1), namely (5.43) and (5.44), focusing on the second case (5.44) since (5.43) follows a similar and indeed simpler argument. We observe that the base case for (5.44) can be verified easily by using the Fourier series decomposition and extending the argument in the one-parameter setting.

We assume that all the inductive statements hold for trees of *all* lower complexities. We recall that the multilinear expression $T_G(P_{k_{\max}}^{(1)} P_{m_{\max}}^{(2)} f_l)_{1 \leq l \leq n}$ yields the localization to the frequency region $R_{k_{\max}} \times R'_{m_{\max}}$, where

$$(5.52) \quad R_{k_{\max}} := \{(\xi_1, \dots, \xi_n) : |\xi_l| \sim 2^{k_{\max}} \text{ for } l \in \mathfrak{M}_1(R) \text{ and } |\xi_l| \ll 2^{k_{\max}} \text{ for } l \in \mathfrak{m}_1(R)\},$$

$$(5.53) \quad R'_{m_{\max}} := \{(\eta_1, \dots, \eta_n) : |\eta_l| \sim 2^{m_{\max}} \text{ for } l \in \mathfrak{M}_2(R') \text{ and } |\eta_l| \ll 2^{m_{\max}} \text{ for } l \in \mathfrak{m}_2(R')\},$$

so that $|\sum_{l=1}^n \xi_l| \leq n2^{k_{\max}}$ and $|\sum_{l=1}^n \eta_l| \leq n2^{m_{\max}}$. As in the one-parameter setting, we smoothly restrict the symbol

$$(5.54) \quad m_{\beta_1^{\mathbf{r}_G}}(\sum_{l=1}^n \xi_l) := |\sum_{l=1}^n \xi_l|^{\beta_1^{\mathbf{r}_G}}$$

to the interval $[-n2^{k_{\max}}, n2^{k_{\max}}]$ and denote it by $m_{\beta_1^{\mathbf{r}_G}}^{k_{\max}}(\sum_{l=1}^n \xi_l)$. Similarly, we denote by $m_{\beta_2^{\mathbf{r}_G}}^{m_{\max}}(\sum_{l=1}^n \eta_l)$ the symbol

$$(5.55) \quad m_{\beta_2^{\mathbf{r}_G}}(\sum_{l=1}^n \eta_l) := |\sum_{l=1}^n \eta_l|^{\beta_2^{\mathbf{r}_G}}$$

localized to the interval $[-n2^{m_{\max}}, n2^{m_{\max}}]$. We undertake the Fourier series decomposition of the localized symbols

$$(5.56) \quad m_{\beta_1^{\mathbf{r}_G}}^{k_{\max}}(\sum_{l=1}^n \xi_l) = (2^{k_{\max}})^{\beta_1^{\mathbf{r}_G}} \sum_{L \in \mathbb{Z}} C_L e^{2\pi i \frac{L}{n2^{k_{\max}}} \sum_{l=1}^n \xi_l},$$

$$(5.57) \quad m_{\beta_2^{\mathbf{r}_G}}^{m_{\max}}(\sum_{l=1}^n \eta_l) = (2^{m_{\max}})^{\beta_2^{\mathbf{r}_G}} \sum_{L' \in \mathbb{Z}} C_{L'} e^{2\pi i \frac{L'}{n2^{m_{\max}}} \sum_{l=1}^n \eta_l},$$

where the (renormalized) Fourier coefficients satisfy the decaying conditions

$$|C_L| \lesssim \frac{1}{(1 + |L|)^{1 + \beta_1^{\mathbf{r}_G}}}, \quad |C_{L'}| \lesssim \frac{1}{(1 + |L'|)^{1 + \beta_2^{\mathbf{r}_G}}}.$$

By applying the Fourier series representations on the multiplier, we indeed obtain

$$(5.58)$$

$$T_G(P_{k_{\max}}^{(1)} P_{m_{\max}}^{(2)} f_l)_{1 \leq l \leq n}(x, y) = \sum_{L, L' \in \mathbb{Z}} C_L C_{L'} 2^{k_{\max} \cdot \beta_1^{\mathbf{r}_G}} 2^{m_{\max} \cdot \beta_2^{\mathbf{r}_G}} \cdot \prod_{i=1}^{n_1} T_{G^{v_i}}((P_{k_{\max}}^{(1)} P_{m_{\max}}^{(2)} f_l)_{l \in \mathcal{L}(v_i)})(x + \frac{L}{n2^{k_{\max}}}, y + \frac{L'}{n2^{m_{\max}}}).$$

Therefore (5.58) in its $\|\cdot\|_{\vec{r}}$ norm⁴⁰ can be majorized by

$$(5.59) \quad 2^{k_{\max} \cdot \beta_1^{\tau_{\mathcal{G}}}} 2^{m_{\max} \cdot \beta_2^{\tau_{\mathcal{G}}}} \prod_{i=1}^{n_1} \|T_{\mathcal{G}^{v_i}} \left((P_{k_{\max}}^{(1)} P_{m_{\max}}^{(2)} f_l)_{l \in \mathcal{L}(v_i)} \right)\|_{\vec{p}_{v_i}}.$$

We notice that there are 2 possibilities for the tree structure with respect to l_0 and l'_0 where $l_0 \in \mathfrak{M}_1(R)$, $l'_0 \in \mathfrak{M}_2(R')$ with $l_0 \neq l'_0$:

- (i) l_0, l'_0 belong to the same subtree $\mathcal{V}^{v_{i_0}}$ for some $1 \leq i_0 \leq n_1$. Assume without loss of generality that $l_0, l'_0 \in \mathcal{L}(v_1)$.
- (ii) $l_0 \in \mathcal{L}(v_{i_0})$ and $l'_0 \in \mathcal{L}(v_{i'_0})$ for some $i_0 \neq i'_0$. Assume that $i_0 = 1$ and $i'_0 = 2$.

In Case (i), one observes that $\mathfrak{M}_1(R) \cap \mathcal{L}(v_1) \neq \emptyset$ and $\mathfrak{M}_2(R') \cap \mathcal{L}(v_1) \neq \emptyset$. This implies that the subtree \mathcal{G}^{v_1} is automatically restricted to the conical regions

$$(5.60) \quad R(v_1) := \{(\xi_l)_{l \in \mathcal{L}(v_1)} : |\xi_l| \gg |\xi_{l'}| \text{ for } l \in \mathcal{L}(v_1) \cap \mathfrak{M}_1(R), l' \in \mathcal{L}(v_1) \setminus \mathfrak{M}_1(R)\};$$

$$(5.61) \quad R'(v_1) := \{(\eta_{l'})_{l' \in \mathcal{L}(v_1)} : |\eta_{l'}| \gg |\eta_l| \text{ for } l \in \mathcal{L}(v_1) \cap \mathfrak{M}_2(R'), l' \in \mathcal{L}(v_1) \setminus \mathfrak{M}_2(R')\}.$$

Furthermore, $k_{\max} = k_{\max}(v_1)$ and $m_{\max} = m_{\max}(v_1)$ specify the Whitney cubes in the cones (5.60) and (5.61). One can then invoke the inductive hypothesis (5.44) on $T_{\mathcal{G}^{v_1}}$ localized on Whitney cubes (at fixed scales) for both parameters:

$$(5.62) \quad \|T_{\mathcal{G}^{v_1}} \left((P_{k_{\max}}^{(1)} P_{m_{\max}}^{(2)} f_l)_{l \in \mathcal{L}(v_1)} \right)\|_{\vec{p}_{v_1}} \lesssim 2^{k_{\max} \cdot \beta_1(v_1, l_0)} 2^{m_{\max} \cdot \beta_2(v_1, l'_0)} \sum_{\substack{\delta_1 |_{(\mathcal{V}_{l'_0}^{v_1})^c} \\ \delta_2 |_{(\mathcal{V}_{l_0}^{v_1})^c}}} \|\Delta_{k_{\max}}^{(1)} D_{(2)}^{\delta_2^{-1}(l_0)} f_{l_0}\|_{\vec{p}_{l_0}} \|\Delta_{m_{\max}}^{(2)} D_{(1)}^{\delta_1^{-1}(l'_0)} f_{l'_0}\|_{\vec{p}_{l'_0}} \prod_{l \neq l_0, l'_0} \|D_{(1)}^{\delta_1^{-1}(l)} D_{(2)}^{\delta_2^{-1}(l)} f_l\|_{\vec{p}_l}.$$

Meanwhile the inductive hypothesis (5.47) and thus (5.50) can be invoked to estimate $T_{\mathcal{G}^{v_i}}$ for $i \neq 1$:

$$(5.63) \quad \|T_{\mathcal{G}^{v_i}} \left((P_{k_{\max}}^{(1)} P_{m_{\max}}^{(2)} f_l)_{l \in \mathcal{L}(v_i)} \right)\|_{\vec{p}_{v_i}} \lesssim \sum_{\delta |_{\mathcal{V}^{v_i}}} \prod_{l \in \mathcal{L}(v_i)} \|D_{(1)}^{\delta_1^{-1}(l)} D_{(2)}^{\delta_2^{-1}(l)} f_l\|_{\vec{p}_l}.$$

By applying the estimates (5.62) and (5.63) to (5.58), we obtain the desired estimate claimed in the inductive statement (5.44).

In Case (ii), we define for $l \in \mathcal{L}(v_1)$

$$\tilde{f}_l := P_{m_{\max}}^{(2)} f_l,$$

and for $l \in \mathcal{L}(v_2)$

$$\tilde{f}_l := P_{k_{\max}}^{(1)} f_l.$$

Then we apply the inductive hypothesis ((5.46) and thus (5.49) to estimate

$$(5.64) \quad \|T_{\mathcal{G}^{v_1}} \left((P_{k_{\max}}^{(1)} P_{m_{\max}}^{(2)} f_l)_{l \in \mathcal{L}(v_1)} \right)\|_{\vec{p}_{v_1}} = \|T_{\mathcal{G}^{v_1}} \left((P_{k_{\max}}^{(1)} \tilde{f}_l)_{l \in \mathcal{L}(v_1)} \right)\|_{\vec{p}_{v_1}} \\ \lesssim 2^{k_{\max} \cdot \beta_1(v_1, l_0)} \sum_{\substack{\delta_2 |_{\mathcal{V}^{v_1}} \\ \delta_1 |_{(\mathcal{V}_{l'_0}^{v_1})^c}}} \|\Delta_{k_{\max}}^{(1)} D_{(2)}^{\delta_2^{-1}(l_0)} f_{l_0}\|_{\vec{p}_{l_0}} \prod_{\substack{l \in \mathcal{L}(v_1) \\ l \neq l_0}} \|D_{(1)}^{\delta_1^{-1}(l)} D_{(2)}^{\delta_2^{-1}(l)} f_l\|_{\vec{p}_l},$$

and

$$(5.65) \quad \|T_{\mathcal{G}^{v_2}} \left((P_{k_{\max}}^{(1)} P_{m_{\max}}^{(2)} f_l)_{l \in \mathcal{L}(v_2)} \right)\|_{\vec{p}_{v_2}} = \|T_{\mathcal{G}^{v_2}} \left((P_{m_{\max}}^{(2)} \tilde{f}_l)_{l \in \mathcal{L}(v_2)} \right)\|_{\vec{p}_{v_2}} \\ \lesssim 2^{m_{\max} \cdot \beta_2(v_2, l'_0)} \sum_{\substack{\delta_1 |_{\mathcal{V}^{v_2}} \\ \delta_2 |_{(\mathcal{V}_{l_0}^{v_2})^c}}} \|\Delta_{m_{\max}}^{(2)} D_{(1)}^{\delta_1^{-1}(l'_0)} f_{l'_0}\|_{\vec{p}_{l'_0}} \prod_{\substack{l \in \mathcal{L}(v_2) \\ l \neq l'_0}} \|D_{(1)}^{\delta_1^{-1}(l)} D_{(2)}^{\delta_2^{-1}(l)} f_l\|_{\vec{p}_l}.$$

We also recall the inductive hypothesis (5.47) and hence (5.50) to the other subtrees corresponding to the multilinear forms for $i \neq 1, 2$ and obtain the estimate (5.63). Plugging the estimates (5.64), (5.65) and (5.63) into (5.58), we derive the estimate specified on the right hand side of (5.44) as desired.

⁴⁰We recall that if $\|\cdot\|_{\vec{r}}$ is not subadditive (i.e. if $r^1 < 1$ or $r^2 < 1$), we need to use instead $\|\cdot\|_{\vec{r}}^{\tau}$ with $\tau \leq \min(1, r^1, r^2)$. The analysis is similar to the reasoning in Section 4.2; in particular, we derive analogous estimate to (4.18), which generate the appropriate conditions on the Lebesgue exponents (1.25).

(2) For the inductive statement (2), we will provide a proof for Case (2a). Case (2b) can be verified using a similar (although not perfectly identical) argument – the hypotheses taking part in the inductive argument are different for the two cases. The base case concerns the estimate for

$$D_{(1)}^{\beta_1} D_{(2)}^{\beta_2} ((P_{k_{\max}}^{(1)} P_{m_{\max}}^{(2)} f_l)_{1 \leq l \leq n}).$$

The Fourier series decomposition of the symbol gives

$$\sum_{L, L' \in \mathbb{Z}} C_L C_{L'} 2^{k_{\max} \beta_1} 2^{m_{\max} \beta_2} \prod_{i=1}^n P_{k_{\max}}^{(1)} P_{m_{\max}}^{(2)} f_l \left(x + \frac{L}{n 2^{k_{\max}}}, y + \frac{L'}{n 2^{m_{\max}}} \right)$$

whose $L^{\vec{r}}$ norm for $r^1, r^2 \geq 1$ ⁴¹ can be majorized by

$$2^{k_{\max} \beta_1} 2^{m_{\max} \beta_2} \prod_{i=1}^n \|P_{k_{\max}}^{(1)} P_{m_{\max}}^{(2)} f_l\|_{\vec{p}_i}$$

due to Hölder's inequality. We recall that in the base case v^{l_1, l_2} is the root and $w^{l_1} = l_1$, $w^{l_2} = l_2$ so that

$$\begin{aligned} \beta_1(\mathbf{r}_G, v^{l_1, l_2}) &= \beta_1, \quad \beta_2(\mathbf{r}_G, l_1) = \beta_2, \\ \beta_1(w^{l_1}, l_1) &= 0, \quad \beta_2(w^{l_2}, l_2) = 0. \end{aligned}$$

We have thus verified the base case for (5.45).

To prove the inductive statement, we apply the Fourier series decomposition on the root symbol as before to tensorize the operator into operators associated to subtrees and obtain (5.58); its $L^{\vec{r}}$ norm can now be estimated by (5.59). There are 2 possible positions for $l_1, l_2 \in \mathfrak{M}_1(R)$ and $l'_0 \in \mathfrak{M}_2(R')$ with $l_1 = l'_0$ and $l_1 \neq l_2$.

- (i) $l_1, l_2 \in \mathcal{L}(v_{i_0})$. Assume without loss of generality that $i_0 = 1$.
- (ii) $l_1 \in \mathcal{L}(v_{i_0})$ and $l_2 \in \mathcal{L}(v_{i'_0})$ with $i_0 \neq i'_0$. Assume that $i_0 = 1$ and $i'_0 = 2$.

For Case (i), we deduce that the multilinear expression associated to the subtree \mathcal{G}^{v_1} is automatically restricted to the cones $R(v_1)$ and $R'(v_1)$ of the form (5.60) and (5.61). By assumption, $l_1, l_2 \in \mathfrak{M}_1(R(v_1))$ and $l'_0 = l_1 \in \mathfrak{M}_2(R'(v_1))$. Let $k_{\max} = k_{\max}(v_1)$ and $m_{\max} = m_{\max}(v_1)$ indicate the Whitney cubes in the cones $R(v_1)$ and $R'(v_1)$. By the inductive hypothesis (5.45),

$$\begin{aligned} (5.66) \quad \|T_{\mathcal{G}^{v_1}}(P_{k_{\max}}^{(1)} P_{m_{\max}}^{(2)} f_l)_{l \in \mathcal{L}(v_1)}\|_{\vec{p}_{v_1}} &\lesssim 2^{k_{\max} \beta_1(v_1, v^{l_1, l_2})} 2^{m_{\max} \beta_2(v_1, l_1)} \prod_{\substack{l \in \mathcal{L}(v_1) \\ l \neq l_1, l_2}} \|D_{(1)}^{\delta_1^{-1}(l)} D_{(2)}^{\delta_2^{-1}(l)} f_l\|_{\vec{p}_l} \\ &\quad \cdot \|\Delta_{k_{\max}}^{(1)} \Delta_{m_{\max}}^{(2)} D_{(1)}^{\beta_1(w^{l_1}, l_1)} f_{l_1}\|_{\vec{p}_{l_1}} \sum_{\substack{\delta_1 | (\mathcal{V}_{l_1}^{v_1})^c \setminus \mathcal{V}_{l_2}^{w^{l_2}} \\ \delta_2 | (\mathcal{V}_{l_1}^{v_1})^c}} \|\Delta_{k_{\max}}^{(1)} D_{(1)}^{\beta_1(w^{l_2}, l_2)} D_{(2)}^{\delta_2^{-1}(l_2)} f_{l_2}\|_{\vec{p}_{l_2}}. \end{aligned}$$

We can invoke the inductive hypothesis (5.50) – assumed to hold for trees of lower complexities – to estimate $\|T_{\mathcal{G}^{v_i}}(P_{k_{\max}}^{(1)} P_{m_{\max}}^{(2)} f_l)_{l \in \mathcal{L}(v_i)}\|_{\vec{p}_{v_i}}$ for $i \neq 1$; we obtain the bound

$$(5.67) \quad \sum_{\delta | \mathcal{V}^{v_i}} \prod_{l \in \mathcal{L}(v_i)} \|D_{(1)}^{\delta_1^{-1}(l)} D_{(2)}^{\delta_2^{-1}(l)} f_l\|_{\vec{p}_l}.$$

By plugging the estimates (5.66) and (5.67) into (5.59), we conclude with (5.45).

In Case (ii), the localization of the original T_G to the cones R and R' imposes a similar restriction to conical regions on $T_{\mathcal{G}^{v_1}}$ and $T_{\mathcal{G}^{v_2}}$. More precisely, let $R(v_1)$ and $R'(v_1)$ denote the conical regions for the subtrees \mathcal{G}^{v_1} for the first and second parameters respectively as before. Let $R(v_2)$ represent the conical region for the subtree \mathcal{G}^{v_2} for the first parameter.

Then we have $l_1 \in \mathfrak{M}_1(R(v_1))$, $l_2 \in \mathfrak{M}_1(R(v_2))$ and $l'_0 \in \mathfrak{M}_2(R'(v_1))$ with $l_1 = l'_0$. Also, $k_{\max} = k_{\max}(v_1)$ and $m_{\max} = m_{\max}(v_1)$ indicate the Whitney cubes in the cones $R(v_1)$ and $R'(v_1)$ for the first and second parameters. The inductive hypothesis (5.43) describes exactly the estimate for the subtree \mathcal{G}^{v_1} :

$$\|T_{\mathcal{G}^{v_1}}(P_{k_{\max}}^{(1)} P_{m_{\max}}^{(2)} f_l)_{l \in \mathcal{L}(v_1)}\|_{\vec{p}_{v_1}} \lesssim 2^{k_{\max} \beta_1(v_1, l_1)} 2^{m_{\max} \beta_2(v_1, l_1)} \|\Delta_{k_{\max}}^{(1)} \Delta_{m_{\max}}^{(2)} f_{l_1}\|_{\vec{p}_{l_1}} \sum_{\delta_1 \otimes \delta_2 | (\mathcal{V}_{l_1}^{v_1})^c} \prod_{\substack{l \in \mathcal{L}(v_1) \\ l \neq l_1}} \|D_{(1)}^{\delta_1^{-1}(l)} D_{(2)}^{\delta_2^{-1}(l)} f_l\|_{\vec{p}_l}$$

⁴¹The estimate in $L^{\vec{r}}$ norm when $r^1, r^2 \geq 1$ doesn't hold requests appropriate conditions on the Lebesgue exponents – see (1.25).

$$(5.68) \quad \lesssim 2^{m_{\max} \beta_2(v_1, l_1)} \|\Delta_{k_{\max}}^{(1)} \Delta_{m_{\max}}^{(2)} D_{(1)}^{\beta_1(v_1, l_1)} f_{l_1}\|_{\tilde{p}_{l_1}} \sum_{\delta_1 \otimes \delta_2|_{(\mathcal{V}_{l_1}^{v_1})^c}} \prod_{\substack{l \in \mathcal{L}(v_1) \\ l \neq l_1}} \|D_{(1)}^{\delta_1^{-1}(l)} D_{(2)}^{\delta_2^{-1}(l)} f_l\|_{\tilde{p}_l},$$

where the last inequality follows from (2.7).

Due to the localization to the cone $R(v_2)$ for the first parameter, we also apply the (corollary of the) inductive hypothesis – (5.49) – to derive the following estimate for the subtree \mathcal{G}^{v_2} :

$$(5.69) \quad \|T_{\mathcal{G}^{v_2}}(P_{k_{\max}}^{(1)} P_{m_{\max}}^{(2)} f_l)_{l \in \mathcal{L}(v_2)}\|_{\tilde{p}_{v_2}} \lesssim \sum_{\substack{\delta_2|_{\mathcal{V}^{v_2}} \\ \delta_1|_{(\mathcal{V}_{l_2}^{v_2})^c}}} \|\Delta_{k_{\max}}^{(1)} D_{(1)}^{\beta_1(v_2, l_1)} D_{(2)}^{\delta_2^{-1}(l_2)} f_{l_2}\|_{\tilde{p}_{l_2}} \prod_{\substack{l \in \mathcal{L}(v_2) \\ l \neq l_2}} \|D_{(1)}^{\delta_1^{-1}(l)} D_{(2)}^{\delta_2^{-1}(l)} f_l\|_{\tilde{p}_l}.$$

Last but not least, we use the bound (5.67) for $\|T_{\mathcal{G}^{v_i}}(P_{k_{\max}}^{(1)} P_{m_{\max}}^{(2)} f_l)_{l \in \mathcal{L}(v_i)}\|_{\tilde{p}_{v_i}}$, $i \neq 1, 2$. With the application of (5.68), (5.69) and (5.67) to (5.59), we complete the proof of the inductive statement (5.45).

(6) The base case of the statement (5.47), corresponding to a tree of complexity 1, is contained in [18]; in Section 4, trees of complexity 2 were treated.

We will focus on the case when the multilinear expression is localized on conical regions of type (5.1) for both parameters, as this is the situation which requires the use of commutators – the tools that allow to depart from the usual methods relying on Coifman-Meyer multipliers. The other cases follow similar arguments with application of possibly different inductive hypotheses. The treatment presented here resembles the proof for the bi-parameter 5-linear flag Leibniz rule presented in Section 4.1.

For the symbol in each parameter, we independently carry out the procedure described in the one-parameter setting to derive a similar expression to (5.29). One will first *split the root symbol* and introduce appropriate commutators in both parameters. Let $l_0, l'_0 \in \mathcal{L}(\mathfrak{r}_{\mathcal{G}})$ denote the indices such that

$$(5.70) \quad \mathfrak{M}_1(R) = \{l_0\}, \quad \mathfrak{M}_2(R') = \{l'_0\},$$

and suppose

$$l_0 \in \mathcal{L}(v_{i_0}), \quad l'_0 \in \mathcal{L}(v_{i'_0}).$$

Then the root symbols ((5.54) and (5.55)) localized to the conical regions specified by (5.70) are decomposed as follows:

$$(5.71) \quad \begin{aligned} m_{\beta_1^{\mathfrak{r}_{\mathcal{G}}}} \left(\sum_{l=1}^n \xi_l \right) &= \sum_{\tilde{i} \neq i_0} m_{C_{\beta_1^{\mathfrak{r}_{\mathcal{G}}}}}^{k_{i_0}} \left(\underbrace{\sum_{l \in \mathcal{L}(v_{i_0})} \xi_l, \sum_{l \notin \mathcal{L}(v_{i_0})} \xi_l}_{\mathcal{A}_1} \cdot \sum_{l \in \mathcal{L}(v_{\tilde{i}})} \xi_l + \underbrace{\sum_{l \in \mathcal{L}(v_{i_0})} \xi_l}_{\mathcal{A}_2} \right)^{\beta_1^{\mathfrak{r}_{\mathcal{G}}}}, \\ m_{\beta_2^{\mathfrak{r}_{\mathcal{G}}}} \left(\sum_{l=1}^n \eta_l \right) &= \sum_{\tilde{i}' \neq i'_0} m_{C_{\beta_2^{\mathfrak{r}_{\mathcal{G}}}}}^{m_{i'_0}} \left(\underbrace{\sum_{l \in \mathcal{L}(v_{i'_0})} \eta_l, \sum_{l \notin \mathcal{L}(v_{i'_0})} \eta_l}_{\mathcal{B}_1} \cdot \sum_{l \in \mathcal{L}(v_{\tilde{i}'})} \eta_l + \underbrace{\sum_{l \in \mathcal{L}(v_{i'_0})} \eta_l}_{\mathcal{B}_2} \right)^{\beta_2^{\mathfrak{r}_{\mathcal{G}}}}, \end{aligned}$$

where

$$(5.72) \quad m_{C_{\beta_1^{\mathfrak{r}_{\mathcal{G}}}}}(\tilde{\xi}_1, \tilde{\xi}_2) := \frac{|\tilde{\xi}_1 + \tilde{\xi}_2|^{\beta_1^{\mathfrak{r}_{\mathcal{G}}}} - |\tilde{\xi}_1|^{\beta_1^{\mathfrak{r}_{\mathcal{G}}}}}{\tilde{\xi}_2}, \quad m_{C_{\beta_2^{\mathfrak{r}_{\mathcal{G}}}}}(\tilde{\eta}_1, \tilde{\eta}_2) := \frac{|\tilde{\eta}_1 + \tilde{\eta}_2|^{\beta_2^{\mathfrak{r}_{\mathcal{G}}}} - |\tilde{\eta}_1|^{\beta_2^{\mathfrak{r}_{\mathcal{G}}}}}{\tilde{\eta}_2}.$$

Recall that $R_{k_{\max}}$ and $R'_{m_{\max}}$ are defined in (5.52) and (5.53) and due to the assumption (5.70) on $\mathfrak{M}_1(R)$ and $\mathfrak{M}_2(R')$, they take the form

$$(5.73) \quad R_{k_{\max}} = R_{k_{i_0}} = \{(\xi_1, \dots, \xi_n) : |\xi_{l_0}| \sim 2^{k_{i_0}}, |\xi_l| \ll 2^{k_{i_0}} \text{ for } l \neq l_0\},$$

$$(5.74) \quad R'_{m_{\max}} = R'_{m_{i'_0}} = \{(\eta_1, \dots, \eta_n) : |\eta_{l'_0}| \sim 2^{m_{i'_0}}, |\eta_l| \ll 2^{m_{i'_0}} \text{ for } l \neq l'_0\}.$$

We notice that $R_{k_{i_0}}$ and $R'_{m_{i'_0}}$ can be decomposed as Whitney cubes on which we will perform double Fourier series decompositions. In particular,

$$R_{k_{i_0}} = R_{k_{i_0}}^+ \cup R_{k_{i_0}}^-, \quad R'_{m_{i'_0}} = (R'_{m_{i'_0}})^+ \cup (R'_{m_{i'_0}})^-,$$

where each Whitney cube is defined by

$$(5.75) \quad R_{k_{i_0}}^+ := \{(\xi_1, \dots, \xi_n) : \xi_{l_0} \sim 2^{k_{i_0}}, |\xi_l| \ll 2^{k_{i_0}} \text{ for } l \neq l_0\}, \quad R_{k_{i_0}}^- := \{(\xi_1, \dots, \xi_n) : \xi_{l_0} \sim -2^{k_{i_0}}, |\xi_l| \ll 2^{k_{i_0}} \text{ for } l \neq l_0\},$$

(5.76)

$$R_{m_{l'_0}}^+ := \{(\eta_1, \dots, \eta_n) : \eta_{l'_0} \sim 2^{m_{l'_0}}, |\eta_l| \ll 2^{m_{l'_0}} \text{ for } l \neq l'_0\}, \quad R_{m_{l'_0}}^- := \{(\eta_1, \dots, \eta_n) : \eta_{l'_0} \sim -2^{m_{l'_0}}, |\eta_l| \ll 2^{m_{l'_0}} \text{ for } l \neq l'_0\}.$$

As previously, we smoothly restrict the symbol $m_{C_{\beta_1^{\mathfrak{r}_g}}}$ (5.72) to the Whitney cube $R_{k_{l_0}}^+$ and to $R_{k_{l_0}}^-$ (5.75) and denote the localized symbols by $m_{C_{\beta_1^{\mathfrak{r}_g}}}^{k_{l_0},+}$ and $m_{C_{\beta_1^{\mathfrak{r}_g}}}^{k_{l_0},-}$ respectively. Similarly, the symbol $m_{C_{\beta_2^{\mathfrak{r}_g}}}$ restricted to $(R'_{m_{l'_0}})^\pm$ regions in (5.76) are denoted by $m_{C_{\beta_2^{\mathfrak{r}_g}}}^{m_{l'_0},\pm}$.

We use Fourier series decomposition to rewrite the symbol $m_{C_{\beta_1^{\mathfrak{r}_g}}}^{k_{l_0},\pm} \left(\sum_{l \in \mathcal{L}(v_{i_0})} \xi_l, \sum_{l \notin \mathcal{L}(v_{i_0})} \xi_l \right)$, which is indeed (5.25) with $\beta^{\mathfrak{r}_g}$ replaced by $\beta_1^{\mathfrak{r}_g}$. Similarly,

$$m_{C_{\beta_2^{\mathfrak{r}_g}}}^{m_{l'_0},\pm} \left(\sum_{l \in \mathcal{L}(v_{i'_0})} \eta_l, \sum_{l \notin \mathcal{L}(v_{i'_0})} \eta_l \right) = \sum_{L'_1, L'_2} C_{L'_1, L'_2}^\pm 2^{m_{l'_0}(\beta_2^{\mathfrak{r}_g} - 1)} e^{2\pi i L'_1 \frac{\sum_{l \in \mathcal{L}'(v_{i'_0})} \eta_l}{2^{m_{l'_0}}}} e^{2\pi i L'_2 \frac{\sum_{l \notin \mathcal{L}'(v_{i'_0})} \eta_l}{2^{m_{l'_0}}}}.$$

The estimate for the multiplier in the biparameter setting concerns the different combinations of the symbols involving the commutators and the symbols for differential operators on subtrees of lower complexity:

$$\mathcal{A}_1 \cdot \mathcal{B}_1, \quad \mathcal{A}_1 \cdot \mathcal{B}_2, \quad \mathcal{A}_2 \cdot \mathcal{B}_1, \quad \mathcal{A}_2 \cdot \mathcal{B}_2.$$

where \mathcal{A}_1 and \mathcal{B}_1 are symbols for commutators while \mathcal{A}_2 and \mathcal{B}_2 represent symbols for differential operators on subtrees. There are 3 possibilities with respect to the relation between l_0 and l'_0 :

- (i) $l_0 = l'_0$;
- (ii) $l_0 \neq l'_0$ and $l_0, l'_0 \in \mathcal{L}(v_{i_0})$ for some i_0 . Assume that $i_0 = 1$;
- (iii) $l_0 \in \mathcal{L}(v_{i_0})$ and $l'_0 \in \mathcal{L}(v_{i'_0})$ for some $i_0 \neq i'_0$. Assume that $i_0 = 1$ and $i'_0 = 2$.

Different possibilities generate multipliers that are analogous to the operators discussed in Section 4.1. In the generic induction, estimates for those multipliers are reduced to estimates on subtrees that request various inductive hypotheses. Since the procedure of reduction to subtree estimates is similar in all cases and the computations after the application of the inductive hypotheses are analogous, we will focus on the proof in Case (i).

When $l_0 = l'_0$, assume without loss of generality that $l_0 = 1 \in \mathcal{L}(v_1)$ and thus $i_0 = 1$. The multipliers involved are listed and estimated as follows.

- estimating $\mathcal{A}_1 \cdot \mathcal{B}_1$ (5.71):

Due to the assumption, we will refer to k_{\max} as k_1 and m_{\max} as m_1 . When $\tilde{i} = \tilde{i}' = 2$ in (5.71), the multiplier takes the form:

$$(5.77) \quad \begin{aligned} & \sum_{\substack{L_1, L_2 \in \mathbb{Z} \\ L'_1, L'_2 \in \mathbb{Z}}} C_{L_1, L_2}^\pm C_{L'_1, L'_2}^\pm \sum_{\substack{k_{\max}(v_2) \ll k_1 \\ m_{\max}(v_2) \ll m_1}} 2^{k_{\max}(v_2)} 2^{k_1 \cdot (\beta_1^{\mathfrak{r}_g} - 1)} 2^{m_{\max}(v_2)} 2^{m_1 \cdot (\beta_2^{\mathfrak{r}_g} - 1)} \\ & T_{\mathcal{G}^{v_1}} \left(\Delta_{k_1, \pm}^{(1)} \Delta_{m_1, \pm}^{(2)} f_1, (S_{k_1}^{(1)} S_{m_1}^{(2)} f_l)_{l \in \mathcal{L}(v_1)} \right) \left(x + \frac{L_1}{2^{k_1}}, y + \frac{L'_1}{2^{m_1}} \right) \\ & T_{\mathcal{G}^{v_2}} \left((P_{k_{\max}(v_2)}^{(1)} P_{m_{\max}(v_2)}^{(2)} f_l)_{l \in \mathcal{L}(v_2)} \right) \left(x + \frac{L_2}{2^{k_1}}, y + \frac{L'_2}{2^{m_1}} \right) \\ & \prod_{i=3}^{n_1} T_{\mathcal{G}^{v_i}} \left((S_{k_1}^{(1)} S_{m_1}^{(2)} f_l)_{l \in \mathcal{L}(v_i)} \right) \left(x + \frac{L_2}{2^{k_1}}, y + \frac{L'_2}{2^{m_1}} \right). \end{aligned}$$

The case when $\tilde{i} \neq \tilde{i}'$ follows a similar argument and will not be discussed in details.

Due to the decay of the Fourier coefficients, the $L^{\tilde{r}}$ norm of (5.77) can be bounded by

$$(5.78) \quad \begin{aligned} & \sum_{\substack{k_{\max}(v_2) \ll k_1 \\ m_{\max}(v_2) \ll m_1}} 2^{k_{\max}(v_2)} 2^{k_1 \cdot (\beta_1^{\mathfrak{r}_g} - 1)} 2^{m_{\max}(v_2)} 2^{m_1 \cdot (\beta_2^{\mathfrak{r}_g} - 1)} \|T_{\mathcal{G}^{v_1}} \left(\Delta_{k_1, +}^{(1)} \Delta_{m_1, +}^{(2)} f_1, (S_{k_1}^{(1)} S_{m_1}^{(2)} f_l)_{l \in \mathcal{L}(v_1)} \right)\|_{\vec{p}_{v_1}} \\ & \cdot \|T_{\mathcal{G}^{v_2}} \left((P_{k_{\max}(v_2)}^{(1)} P_{m_{\max}(v_2)}^{(2)} f_l)_{l \in \mathcal{L}(v_2)} \right)\|_{\vec{p}_{v_2}} \prod_{i=3}^{n_1} \|T_{\mathcal{G}^{v_i}} \left((S_{k_1}^{(1)} S_{m_1}^{(2)} f_l)_{l \in \mathcal{L}(v_i)} \right)\|_{\vec{p}_{v_i}}. \end{aligned}$$

We can apply the inductive hypothesis (5.43) on $T_{\mathcal{G}^{v_1}}(\Delta_{k_1, \pm}^{(1)} \Delta_{m_1, \pm}^{(2)} f_1, (S_{k_1}^{(1)} S_{m_1}^{(2)} f_l)_{l \in \mathcal{L}(v_1)})$ and obtain

$$(5.79) \quad \begin{aligned} \|T_{\mathcal{G}^{v_1}}(\Delta_{k_1, \pm}^{(1)} \Delta_{m_1, \pm}^{(2)} f_1, (S_{k_1}^{(1)} S_{m_1}^{(2)} f_l)_{l \in \mathcal{L}(v_1)})\|_{\tilde{p}_{v_1}} &\lesssim \|\Delta_{k_{\max}}^{(1)} \Delta_{m_{\max}}^{(2)} D_{(1)}^{\beta_1(v_1, 1)} D_{(2)}^{\beta_2(v_1, 1)} \Delta_{k_1, \pm}^{(1)} \Delta_{m_1, \pm}^{(2)} f_1\|_{\tilde{p}_1} \sum_{\delta_1 \otimes \delta_2 |_{(\mathcal{V}_1^{v_1})^c}} \prod_{l \neq 1} \|D_{(1)}^{\delta_1^{-1}(l)} D_{(2)}^{\delta_2^{-1}(l)} f_l\|_{\tilde{p}_l} \\ &\lesssim \|\Delta_{k_{\max}}^{(1)} \Delta_{m_{\max}}^{(2)} D_{(1)}^{\beta_1(v_1, 1)} D_{(2)}^{\beta_2(v_1, 1)} f_1\|_{\tilde{p}_1} \sum_{\delta_1 \otimes \delta_2 |_{(\mathcal{V}_1^{v_1})^c}} \prod_{l \neq 1} \|D_{(1)}^{\delta_1^{-1}(l)} D_{(2)}^{\delta_2^{-1}(l)} f_l\|_{\tilde{p}_l}. \end{aligned}$$

Meanwhile, we further decompose the multilinear expression associated to the subtree \mathcal{G}^{v_2} and denote by $R(v_2)$ and $R'(v_2)$ the conical regions for the first and second parameters. We apply (5.43) or (5.44) depending on the type of conical regions on

$$T_{\mathcal{G}^{v_2}}^{R(v_2) \times R'(v_2)}((P_{k_{\max}(v_2)}^{(1)} P_{k_{\max}(v_2)}^{(2)} f_l)_{l \in \mathcal{L}(v_2)}).$$

We also invoke (5.50) on

$$T_{\mathcal{G}^{v_i}}((S_{k_1}^{(1)} S_{m_1}^{(2)} f_l)_{l \in \mathcal{L}(v_i)})$$

for $i \neq 1, 2$. Suppose that $\tilde{l} \in \mathfrak{M}_1(R(v_2))$ and $\tilde{l}' \in \cap \mathfrak{M}_2(R'(v_2))$ with $\tilde{l} \neq \tilde{l}'$. Then (5.44) together with other inductive hypotheses and the estimates (5.79) and (2.16) imply that

$$\begin{aligned} \|(5.78)\|_{\tilde{r}} &\lesssim \sum_{\substack{\delta_1 |_{\mathcal{V} \setminus \{\tau_{\mathcal{G}}\} \setminus \mathcal{V}_1^{v_1} \setminus \mathcal{V}_i^{v_2} \\ \delta_2 |_{\mathcal{V} \setminus \{\tau_{\mathcal{G}}\} \setminus \mathcal{V}_1^{v_1} \setminus \mathcal{V}_{i'}^{v_2}}}} \prod_{l \neq 1, \tilde{l}, \tilde{l}'} \|D_{(1)}^{\delta_1^{-1}(l)} D_{(2)}^{\delta_2^{-1}(l)} f_l\|_{\tilde{p}_l} \cdot \sum_{\substack{k_{\max}(v_2) \ll k_1 \\ m_{\max}(v_2) \ll m_1}} 2^{k_{\max}(v_2)} 2^{k_1 \cdot (\beta_1^{\tau_{\mathcal{G}}} - 1)} 2^{m_{\max}(v_2)} 2^{m_1 \cdot (\beta_2^{\tau_{\mathcal{G}}} - 1)} \\ &\quad \|\Delta_{k_{\max}}^{(1)} \Delta_{m_{\max}}^{(2)} D_{(1)}^{\beta_1(v_1, 1)} D_{(2)}^{\beta_2(v_1, 1)} f_1\|_{\tilde{p}_1} \cdot \|\Delta_{k_{\max}(v_2)}^{(1)} D_{(1)}^{\beta_1(v_2, \tilde{l})} D_{(2)}^{\delta_2^{-1}(\tilde{l})} f_{\tilde{l}}\|_{\tilde{p}_{\tilde{l}}} \|\Delta_{m_{\max}(v_2)}^{(2)} D_{(1)}^{\delta_1^{-1}(\tilde{l}')} D_{(2)}^{\beta_2(v_2, \tilde{l}')} f_{\tilde{l}'}\|_{\tilde{p}_{\tilde{l}'}}. \end{aligned}$$

The similar estimate developed in Section 4.1 can be applied to distribute derivatives as follows: for any fixed $\delta_1 |_{\mathcal{V} \setminus \{\tau_{\mathcal{G}}\} \setminus \mathcal{V}_1^{v_1} \setminus \mathcal{V}_i^{v_2}}$ and $\delta_2 |_{\mathcal{V} \setminus \{\tau_{\mathcal{G}}\} \setminus \mathcal{V}_1^{v_1} \setminus \mathcal{V}_{i'}^{v_2}}$, the inner sum can be bounded by

$$\begin{aligned} \sum_{\substack{k_1 \\ m_1}} \min &\left(2^{k_1 \beta_1^{\tau_{\mathcal{G}}}} 2^{m_1 \beta_2^{\tau_{\mathcal{G}}}} \|D_{(1)}^{\beta_1(v_1, 1)} D_{(2)}^{\beta_2(v_1, 1)} f_1\|_{\dot{B}_{p_1, \infty}^0 \dot{B}_{p_1, \infty}^0} \|D_{(1)}^{\beta_1(v_2, \tilde{l})} D_{(2)}^{\delta_2^{-1}(\tilde{l})} f_{\tilde{l}}\|_{\dot{B}_{p_1, \infty}^0 L_{p_i}^2} \|D_{(1)}^{\delta_1^{-1}(\tilde{l}')} D_{(2)}^{\beta_2(v_2, \tilde{l}')} f_{\tilde{l}'}\|_{L_{p_{i'}}^1 \dot{B}_{p_{i'}, \infty}^0}, \right. \\ &2^{k_1 \beta_1^{\tau_{\mathcal{G}}}} 2^{-m_1 \epsilon_2} \|D_{(1)}^{\beta_1(v_1, 1)} D_{(2)}^{\beta_2(v_1, 1)} f_1\|_{\dot{B}_{p_1, \infty}^0 \dot{B}_{p_2, \infty}^{\tau_{\mathcal{G}}}} \|D_{(1)}^{\beta_1(v_2, \tilde{l})} D_{(2)}^{\delta_2^{-1}(\tilde{l})} f_{\tilde{l}}\|_{\dot{B}_{p_1, \infty}^0 L_{p_i}^2} \|D_{(1)}^{\delta_1^{-1}(\tilde{l}')} D_{(2)}^{\beta_2(v_2, \tilde{l}')} f_{\tilde{l}'}\|_{L_{p_{i'}}^1 \dot{B}_{p_{i'}, \infty}^{\epsilon_2}}, \\ &2^{-k_1 \epsilon_1} 2^{m_1 \beta_2^{\tau_{\mathcal{G}}}} \|D_{(1)}^{\beta_1(v_1, 1)} D_{(2)}^{\beta_2(v_1, 1)} f_1\|_{\dot{B}_{p_1, \infty}^{\beta_1^{\tau_{\mathcal{G}}}} \dot{B}_{p_2, \infty}^0} \|D_{(1)}^{\beta_1(v_2, \tilde{l})} D_{(2)}^{\delta_2^{-1}(\tilde{l})} f_{\tilde{l}}\|_{\dot{B}_{p_1, \infty}^{\epsilon_1} L_{p_i}^2} \|D_{(1)}^{\delta_1^{-1}(\tilde{l}')} D_{(2)}^{\beta_2(v_2, \tilde{l}')} f_{\tilde{l}'}\|_{L_{p_{i'}}^1 \dot{B}_{p_{i'}, \infty}^0}, \\ &2^{-k_1 \epsilon_1} 2^{-m_1 \epsilon_2} \|D_{(1)}^{\beta_1(v_1, 1)} D_{(2)}^{\beta_2(v_1, 1)} f_1\|_{\dot{B}_{p_1, \infty}^{\beta_1^{\tau_{\mathcal{G}}}} \dot{B}_{p_2, \infty}^{\tau_{\mathcal{G}}}} \|D_{(1)}^{\beta_1(v_2, \tilde{l})} D_{(2)}^{\delta_2^{-1}(\tilde{l})} f_{\tilde{l}}\|_{\dot{B}_{p_1, \infty}^{\epsilon_1} L_{p_i}^2} \|D_{(1)}^{\delta_1^{-1}(\tilde{l}')} D_{(2)}^{\beta_2(v_2, \tilde{l}')} f_{\tilde{l}'}\|_{L_{p_{i'}}^1 \dot{B}_{p_{i'}, \infty}^{\epsilon_2}} \Big) \end{aligned}$$

for $0 < \epsilon_j < \min(1, \beta_j^{\tau_{\mathcal{G}}})$, $j = 1, 2$. By the optimization and interpolation procedure specified in Section 5.3, we attain the right hand side of the inductive statement (5.47).

- estimating $\mathcal{A}_1 \cdot \mathcal{B}_2$ (5.71):

The multipliers generated by $\mathcal{A}_1 \cdot \mathcal{B}_2$ are similar to the ones generated by $\mathcal{A}_2 \cdot \mathcal{B}_1$; by symmetry it will be enough to focus on the former. The symbol $\mathcal{A}_1 \cdot \mathcal{B}_2$ with $\tilde{i} = 2$ in (5.71) generates the multiplier

$$\begin{aligned} \sum_{L_1, L_2 \in \mathbb{Z}} C_{L_1, L_2}^{\pm} \sum_{\substack{k_{\max}(v_2) \ll k_1 \\ m_1}} 2^{k_{\max}(v_2)} 2^{k_1 \cdot (\beta_1^{\tau_{\mathcal{G}}} - 1)} D_{(2)}^{\beta_2^{\tau_{\mathcal{G}}}} T_{\mathcal{G}^{v_1}}(\Delta_{k_1, \pm, \frac{L_1}{2k_1}}^{(1)} \Delta_{m_1, \frac{L_2}{2k_1}}^{(2)} f_1, (S_{k_1, \frac{L_1}{2k_1}}^{(1)} S_{m_1}^{(2)} f_l)_{l \in \mathcal{L}(v_1)})) \\ T_{\mathcal{G}^{v_2}}((P_{k_{\max}(v_2), \frac{L_2}{2k_1}}^{(1)} S_{m_1}^{(2)} f_l)_{l \in \mathcal{L}(v_2)}) \prod_{i=3}^{n_1} T_{\mathcal{G}^{v_i}}((S_{k_1, \frac{L_2}{2k_1}}^{(1)} S_{m_1}^{(2)} f_l)_{l \in \mathcal{L}(v_i)}). \end{aligned}$$

Let $\tilde{\mathcal{G}}^{v_1}$ denote the tree having the same structure as \mathcal{G}^{v_1} , except that the differential operator associated to the vertex v_1 is replaced by $D_{(1)}^{\beta_1^{v_1}} D_{(2)}^{\beta_2^{v_1} + \beta_2^{\tau_{\mathcal{G}}}}$. As before, the estimate for the above term is the same as the simpler term when $L_i = 0$ for $i = 1, 2$:

$$\sum_{\substack{k_{\max}(v_2) \ll k_1 \\ m_1}} 2^{k_{\max}(v_2)} 2^{k_1 \cdot (\beta_1^{\tau_{\mathcal{G}}} - 1)} \cdot T_{\tilde{\mathcal{G}}^{v_1}}(\Delta_{k_1, \pm}^{(1)} \Delta_{m_1}^{(2)} f_1, (S_{k_1}^{(1)} S_{m_1}^{(2)} f_l)_{l \in \mathcal{L}(v_1)})) \cdot T_{\mathcal{G}^{v_2}}((P_{k_{\max}(v_2)}^{(1)} S_{m_1}^{(2)} f_l)_{l \in \mathcal{L}(v_2)})$$

$$\cdot \prod_{i=3}^{n_1} T_{\mathcal{G}^{v_i}} \left((S_{k_1}^{(1)} S_{m_1}^{(2)} f_l)_{l \in \mathcal{L}(v_i)} \right).$$

As in the one-parameter setting, we perform finer paraproduct decompositions on the subtrees \mathcal{G}^{v_i} , $i \neq 1$, for the second parameter:

$$(5.80) \quad \sum_{\substack{k_{\max}(v_2) \ll k_1 \\ m_1 \\ m_{\max}(v_2)}} 2^{k_{\max}(v_2)} 2^{k_1(\beta_1^{\tau_{\mathcal{G}}}-1)} T_{\tilde{\mathcal{G}}^{v_1}} \left(\Delta_{k_1, \pm}^{(1)} \Delta_{m_1}^{(2)} f_1, (S_{k_1}^{(1)} S_{m_1}^{(2)} f_l)_{l \in \mathcal{L}(v_1)} \right)_{l \neq 1} \\ T_{\mathcal{G}^{v_2}} \left((P_{k_{\max}(v_2)}^{(1)} P_{m_{\max}(v_2)}^{(2)} S_{m_1}^{(2)} f_l)_{l \in \mathcal{L}(v_2)} \right) \prod_{i=3}^{n_1} \sum_{m_{\max}(v_i)} T_{\mathcal{G}^{v_i}} \left((S_{k_1}^{(1)} P_{m_{\max}(v_i)}^{(2)} S_{m_1}^{(2)} f_l)_{l \in \mathcal{L}(v_i)} \right).$$

Due to the observation that $\Delta_{m_{\max}(v_i)}^{(2)} S_{m_1}^{(2)} \neq 0$ only if $m_{\max}(v_i) \ll m_1$, (5.80) can be simplified as

$$\sum_{\substack{k_{\max}(v_2) \ll k_1 \\ m_{\max}(v_2) \ll m_1}} 2^{k_{\max}(v_2)} 2^{k_1(\beta_1^{\tau_{\mathcal{G}}}-1)} T_{\tilde{\mathcal{G}}^{v_1}} \left(\Delta_{k_1, \pm}^{(1)} \Delta_{m_1}^{(2)} f_1, (S_{k_1}^{(1)} S_{m_1}^{(2)} f_l)_{l \in \mathcal{L}(v_1)} \right)_{l \neq 1} \\ T_{\mathcal{G}^{v_2}} \left((P_{k_{\max}(v_2)}^{(1)} P_{m_{\max}(v_2)}^{(2)} f_l)_{l \in \mathcal{L}(v_2)} \right) \prod_{i=3}^{n_1} \sum_{m_{\max}(v_i): m_{\max}(v_i) \ll m_1} T_{\mathcal{G}^{v_i}} \left((S_{k_1}^{(1)} P_{m_{\max}(v_i)}^{(2)} f_l)_{l \in \mathcal{L}(v_i)} \right).$$

We then apply the high-low switch technique to reduce the expression above to a sum of terms that can be estimated using the inductive hypotheses. In particular,

$$\sum_{\substack{k_{\max}(v_2) \ll k_1 \\ m_1 \\ m_{\max}(v_2)}} 2^{k_{\max}(v_2)} 2^{k_1(\beta_1^{\tau_{\mathcal{G}}}-1)} T_{\tilde{\mathcal{G}}^{v_1}} \left(\Delta_{k_1, \pm}^{(1)} \Delta_{m_1}^{(2)} f_1, (S_{k_1}^{(1)} S_{m_1}^{(2)} f_l)_{l \in \mathcal{L}(v_1)} \right)_{l \neq 1} T_{\mathcal{G}^{v_2}} \left((P_{k_{\max}(v_2)}^{(1)} P_{m_{\max}(v_2)}^{(2)} f_l)_{l \in \mathcal{L}(v_2)} \right) \\ \cdot \prod_{i=3}^{n_1} \sum_{m_{\max}(v_i)} T_{\mathcal{G}^{v_i}} \left((S_{k_1}^{(1)} P_{m_{\max}(v_i)}^{(2)} f_l)_{l \in \mathcal{L}(v_i)} \right) \\ - \sum_{\substack{k_{\max}(v_2) \ll k_1 \\ m_{\max}(v_2) > m_1}} 2^{k_{\max}(v_2)} 2^{k_1(\beta_1^{\tau_{\mathcal{G}}}-1)} T_{\tilde{\mathcal{G}}^{v_1}} \left(\Delta_{k_1, \pm}^{(1)} \Delta_{m_1}^{(2)} f_1, (S_{k_1}^{(1)} S_{m_1}^{(2)} f_l)_{l \in \mathcal{L}(v_1)} \right)_{l \neq 1} T_{\mathcal{G}^{v_2}} \left((P_{k_{\max}(v_2)}^{(1)} P_{m_{\max}(v_2)}^{(2)} f_l)_{l \in \mathcal{L}(v_2)} \right) \\ \cdot \prod_{i=3}^{n_1} \sum_{m_{\max}(v_i)} T_{\mathcal{G}^{v_i}} \left((S_{k_1}^{(1)} P_{m_{\max}(v_i)}^{(2)} f_l)_{l \in \mathcal{L}(v_i)} \right)$$

\pm similar terms $:= I - II \pm$ similar terms.

We will elaborate on the estimates for the first term denoted by I and the second term denoted by II . For I , we first recall the inductive hypothesis (5.46) that allows to control, for k_1 and $k_{\max}(v_2)$ fixed,

$$\sum_{m_1} T_{\tilde{\mathcal{G}}^{v_1}} \left(\Delta_{k_1, \pm}^{(1)} \Delta_{m_1}^{(2)} f_1, (S_{k_1}^{(1)} S_{m_1}^{(2)} f_l)_{l \in \mathcal{L}(v_1)} \right)_{l \neq 1} \quad \text{and} \quad \sum_{m_{\max}(v_2)} T_{\mathcal{G}^{v_2}} \left((P_{k_{\max}(v_2)}^{(1)} P_{m_{\max}(v_2)}^{(2)} f_l)_{l \in \mathcal{L}(v_2)} \right).$$

For $i \geq 3$ and $l \in \mathcal{L}(v_i)$, define

$$\tilde{f}_l := S_{k_1}^{(1)} f_l.$$

and apply the inductive hypothesis (5.51) to

$$\sum_{m_{\max}(v_i)} T_{\mathcal{G}^{v_i}} \left((P_{m_{\max}(v_i)}^{(2)} \tilde{f}_l)_{l \in \mathcal{L}(v_i)} \right).$$

Suppose that $l_0 \in \mathfrak{M}_1(R(v_2))$. Combining the estimates from the inductive hypotheses, we derive that

$$\|I\|_{\tau} \lesssim \sum_{\delta_2} \prod_{l \neq 1, l_0} \|D_{(1)}^{\delta_1^{-1}(l)} D_{(2)}^{\delta_2^{-1}(l)} f_l\|_{\tilde{p}_l} \sum_{k_{\max}(v_2) \ll k_1} 2^{k_{\max}(v_2)} 2^{k_1(\beta_1^{\tau_{\mathcal{G}}}-1)} \\ \cdot \|\Delta_{k_1}^{(1)} D_{(1)}^{\beta_1(v_1, 1)} D_{(2)}^{\delta_2^{-1}(1)} f_1\|_{\tilde{p}_1} \|\Delta_{k_{\max}(v_2)}^{(1)} D_{(1)}^{\beta_1(v_2, \tilde{l})} D_{(2)}^{\delta_2^{-1}(\tilde{l})} f_{\tilde{l}}\|_{\tilde{p}_{\tilde{l}}}.$$

Now we fix δ_2 and $\delta_1|_{\mathcal{V} \setminus \{\tau_{\mathcal{G}}\} \setminus \mathcal{V}_1^{v_1} \setminus \mathcal{V}_{l_0}^{v_2}}$ and distribute the partial derivatives in the first parameter:

$$\sum_{k_{\max}(v_2) \ll k_1} 2^{k_{\max}(v_2)} 2^{k_1(\beta_1^{\tau_{\mathcal{G}}}-1)} \|\Delta_{k_1}^{(1)} D_{(1)}^{\beta_1(v_1, 1)} D_{(2)}^{\delta_2^{-1}(1)} f_1\|_{\tilde{p}_1} \|\Delta_{k_{\max}(v_2)}^{(1)} D_{(1)}^{\beta_1(v_2, \tilde{l})} D_{(2)}^{\delta_2^{-1}(\tilde{l})} f_{\tilde{l}}\|_{\tilde{p}_{\tilde{l}}}$$

$$\lesssim \sum_{k_1} \min \left(2^{k_1 \beta_1^{\tau_{\mathcal{G}}}} \|D_{(1)}^{\beta_1(v_1,1)} D_{(2)}^{\delta_2^{-1}(1)} f_1\|_{\dot{B}_{p_1^1, \infty}^0 L^{p_1^2}} \|D_{(1)}^{\beta_1(v_2, \tilde{l})} D_{(2)}^{\delta_2^{-1}(\tilde{l})} f_{\tilde{l}}\|_{\dot{B}_{p_1^1, \infty}^0 L^{p_1^2}}, \right. \\ \left. 2^{-k_1 \epsilon_1} \|D_{(1)}^{\beta_1(v_1,1)} D_{(2)}^{\delta_2^{-1}(1)} f_1\|_{\dot{B}_{p_1^1, \infty}^{\beta_1^{\tau_{\mathcal{G}}}} L^{p_1^2}} \|D_{(1)}^{\beta_1(v_2, \tilde{l})} D_{(2)}^{\delta_2^{-1}(\tilde{l})} f_{\tilde{l}}\|_{\dot{B}_{p_1^1, \infty}^{\epsilon_1} L^{p_1^2}} \right);$$

using the optimization and interpolation described in Section 5.3, we deduce the inductive statement (5.47). The term II requires the same inductive hypothesis (5.51) on

$$\sum_{m_{\max}(v_i)} T_{\mathcal{G}^{v_i}}((S_{k_1}^{(1)} P_{m_{\max}(v_i)}^{(2)} f_l)_{l \in \mathcal{L}(v_i)})$$

for $i \geq 3$, while (5.43) yields estimates on $T_{\tilde{\mathcal{G}}^{v_1}}(\Delta_{k_1, +}^{(1)} \Delta_{m_1}^{(2)} f_1, (S_{k_1}^{(1)} S_{m_1}^{(2)} f_l)_{l \in \mathcal{L}(v_1)})$.

Recall that $\tilde{l} \in \mathfrak{M}_1(R(v_2))$ and further assume that $\tilde{l}' \in \mathfrak{M}_2(R'(v_2))$ with $\tilde{l} \neq \tilde{l}'$.⁴² Then (5.44) is applicable to

$$T_{\mathcal{G}^{v_2}}((P_{k_{\max}(v_2)}^{(1)} P_{m_{\max}(v_2)}^{(2)} f_l)_{l \in \mathcal{L}(v_2)}).$$

As a consequence,

$$\|II\|_{\tilde{\tau}} \lesssim \sum_{\substack{\delta_1|_{\mathcal{V} \setminus \{\tau_{\mathcal{G}}\} \setminus \mathcal{V}_1^{v_1} \setminus \mathcal{V}_2^{v_2}} \\ \delta_2|_{\mathcal{V} \setminus \{\tau_{\mathcal{G}}\} \setminus \mathcal{V}_{\tilde{l}'}^{v_2}}}} \prod_{l \neq \tilde{l}, \tilde{l}'} \|D_{(1)}^{\delta_1^{-1}(l)} D_{(2)}^{\delta_2^{-1}(l)} f_l\|_{\tilde{p}_l} \sum_{\substack{k_{\max}(v_2) \leq k_1 \\ m_{\max}(v_2) > m_1}} 2^{k_{\max}(v_2)} 2^{k_1(\beta_1^{\tau_{\mathcal{G}}} - 1)} 2^{m_1 \beta_2^{\tau_{\mathcal{G}}}} \\ \| \Delta_{k_1}^{(1)} \Delta_{m_1}^{(2)} D_{(1)}^{\beta_1(v_1,1)} D_{(2)}^{\beta_2(v_1,1)} f_1 \|_{\tilde{p}_1} \| \Delta_{k_{\max}(v_2)}^{(1)} D_{(1)}^{\beta_1(v_2, \tilde{l})} D_{(2)}^{\delta_2^{-1}(\tilde{l})} f_{\tilde{l}} \|_{\tilde{p}_{\tilde{l}}} \| \Delta_{m_{\max}(v_2)}^{(2)} D_{(2)}^{\delta_1^{-1}(\tilde{l}')} D_{(2)}^{\beta_2(v_2, \tilde{l}')} f_{\tilde{l}'} \|_{\tilde{p}_{\tilde{l}'}}.$$

In the inner sum the partial derivatives can be appropriately distributed as before:

$$\sum_{\substack{k_{\max}(v_2) \leq k_1 \\ m_{\max}(v_2) > m_1}} 2^{k_{\max}(v_2)} 2^{k_1(\beta_1^{\tau_{\mathcal{G}}} - 1)} 2^{m_1 \beta_2^{\tau_{\mathcal{G}}}} \| \Delta_{k_1}^{(1)} \Delta_{m_1}^{(2)} D_{(1)}^{\beta_1(v_1,1)} D_{(2)}^{\beta_2(v_1,1)} f_1 \|_{\tilde{p}_1} \\ \| \Delta_{k_{\max}(v_2)}^{(1)} D_{(1)}^{\beta_1(v_2, \tilde{l})} D_{(2)}^{\delta_2^{-1}(\tilde{l})} f_{\tilde{l}} \|_{\tilde{p}_{\tilde{l}}} \| \Delta_{m_{\max}(v_2)}^{(2)} D_{(2)}^{\delta_1^{-1}(\tilde{l}')} D_{(2)}^{\beta_2(v_2, \tilde{l}')} f_{\tilde{l}'} \|_{\tilde{p}_{\tilde{l}'}} \\ \lesssim \sum_{\substack{k_1 \\ m_{\max}(v_2)}} \min(2^{k_1 \beta_1^{\tau_{\mathcal{G}}}} 2^{m_{\max}(v_2) \beta_2^{\tau_{\mathcal{G}}}} \|D_{(1)}^{\beta_1(v_1,1)} D_{(2)}^{\beta_2(v_1,1)} f_1\|_{\dot{B}_{p_1^1, \infty}^0 \dot{B}_{p_1^2, \infty}^0} \|D_{(1)}^{\beta_1(v_2, \tilde{l})} D_{(2)}^{\delta_2^{-1}(\tilde{l})} f_{\tilde{l}}\|_{\dot{B}_{p_1^1, \infty}^0 L^{p_1^2}} \|D_{(2)}^{\delta_1^{-1}(\tilde{l}')} D_{(2)}^{\beta_2(v_2, \tilde{l}')} f_{\tilde{l}'}\|_{L^{p_{\tilde{l}'}} \dot{B}_{p_{\tilde{l}'}^2, \infty}^0}, \\ 2^{k_1 \beta_1^{\tau_{\mathcal{G}}}} 2^{-m_{\max}(v_2) \epsilon_2} \|D_{(1)}^{\beta_1(v_1,1)} D_{(2)}^{\beta_2(v_1,1)} f_1\|_{\dot{B}_{p_1^1, \infty}^0 \dot{B}_{p_1^2, \infty}^{\epsilon_2}} \|D_{(1)}^{\beta_1(v_2, \tilde{l})} D_{(2)}^{\delta_2^{-1}(\tilde{l})} f_{\tilde{l}}\|_{\dot{B}_{p_1^1, \infty}^0 L^{p_1^2}} \|D_{(2)}^{\delta_1^{-1}(\tilde{l}')} D_{(2)}^{\beta_2(v_2, \tilde{l}')} f_{\tilde{l}'}\|_{L^{p_{\tilde{l}'}} \dot{B}_{p_{\tilde{l}'}^2, \infty}^{\beta_2^{\tau_{\mathcal{G}}}}}, \\ 2^{-k_1 \epsilon_1} 2^{m_{\max}(v_2) \beta_2^{\tau_{\mathcal{G}}}} \|D_{(1)}^{\beta_1(v_1,1)} D_{(2)}^{\beta_2(v_1,1)} f_1\|_{\dot{B}_{p_1^1, \infty}^{\beta_1^{\tau_{\mathcal{G}}}} \dot{B}_{p_1^2, \infty}^0} \|D_{(1)}^{\beta_1(v_2, \tilde{l})} D_{(2)}^{\delta_2^{-1}(\tilde{l})} f_{\tilde{l}}\|_{\dot{B}_{p_1^1, \infty}^{\epsilon_1} L^{p_1^2}} \|D_{(2)}^{\delta_1^{-1}(\tilde{l}')} D_{(2)}^{\beta_2(v_2, \tilde{l}')} f_{\tilde{l}'}\|_{L^{p_{\tilde{l}'}} \dot{B}_{p_{\tilde{l}'}^2, \infty}^0}, \\ 2^{-k_1 \epsilon_1} 2^{-m_{\max}(v_2) \epsilon_2} \|D_{(1)}^{\beta_1(v_1,1)} D_{(2)}^{\beta_2(v_1,1)} f_1\|_{\dot{B}_{p_1^1, \infty}^{\beta_1^{\tau_{\mathcal{G}}}} \dot{B}_{p_1^2, \infty}^{\epsilon_2}} \|D_{(1)}^{\beta_1(v_2, \tilde{l})} D_{(2)}^{\delta_2^{-1}(\tilde{l})} f_{\tilde{l}}\|_{\dot{B}_{p_1^1, \infty}^{\epsilon_1} L^{p_1^2}} \|D_{(2)}^{\delta_1^{-1}(\tilde{l}')} D_{(2)}^{\beta_2(v_2, \tilde{l}')} f_{\tilde{l}'}\|_{L^{p_{\tilde{l}'}} \dot{B}_{p_{\tilde{l}'}^2, \infty}^{\beta_2^{\tau_{\mathcal{G}}}}}),$$

to which we can apply optimization and interpolation to obtain the desired estimates described in the right hand side of (5.47).

- estimating $\mathcal{A}_2 \cdot \mathcal{B}_2$ (5.71):

The terms generated by $\mathcal{A}_2 \cdot \mathcal{B}_2$ take the form

$$\sum_{k_1, m_1} D_{(1)}^{\beta_1^{\tau_{\mathcal{G}}}} D_{(2)}^{\beta_2^{\tau_{\mathcal{G}}}} T_{\mathcal{G}^{v_1}}(\Delta_{k_1}^{(1)} \Delta_{m_1}^{(2)} f_1, (S_{k_1}^{(1)} S_{m_1}^{(2)} f_l)_{l \in \mathcal{L}(v_1)}) \prod_{l \neq \tilde{l}}^{n_1} T_{\mathcal{G}^{v_i}}((S_{k_1}^{(1)} S_{m_1}^{(2)} f_l)_{l \in \mathcal{L}(v_i)}).$$

Let $\tilde{\mathcal{G}}^{v_1}$ be the tree having the same configuration as \mathcal{G}^{v_1} , with the original differential operator associated to the vertex v_1

$$D_{(1)}^{\beta_1^{v_1}} D_{(2)}^{\beta_2^{v_1}}$$

replaced by

$$D_{(1)}^{\beta_1^{v_1} + \beta_1^{\tau_{\mathcal{G}}}} D_{(2)}^{\beta_2^{v_1} + \beta_2^{\tau_{\mathcal{G}}}}.$$

⁴²If $l = \tilde{l}'$, then the inductive hypothesis (5.43) will be used instead.

The treatment of the second parameter in last section (estimating $\mathcal{A}_1 \cdot \mathcal{B}_2$) can be used in both parameters here. In particular, the finer paraproduct decompositions on both parameters yield

$$\begin{aligned}
 & \sum_{\substack{k_1 \\ m_1}} T_{\tilde{\mathcal{G}}^{v_1}} (\Delta_{k_1}^{(1)} \Delta_{m_1}^{(2)} f_1, (S_{k_1}^{(1)} S_{m_1}^{(2)} f_l)_{l \in \mathcal{L}(v_1)}) \prod_{i=2}^{n_1} \sum_{\substack{k_{\max}(v_i) \\ m_{\max}(v_i)}} T_{\mathcal{G}^{v_i}} ((P_{k_{\max}(v_i)}^{(1)} P_{m_{\max}(v_i)}^{(2)} S_{k_1}^{(1)} S_{m_1}^{(2)} f_l)_{l \in \mathcal{L}(v_i)}) \\
 (5.81) \quad & = \sum_{\substack{k_1 \\ m_1}} T_{\tilde{\mathcal{G}}^{v_1}} (\Delta_{k_1}^{(1)} \Delta_{m_1}^{(2)} f_1, (S_{k_1}^{(1)} S_{m_1}^{(2)} f_l)_{l \in \mathcal{L}(v_1)}) \prod_{i=2}^{n_1} \sum_{\substack{k_{\max}(v_i): k_{\max}(v_i) \ll k_1 \\ m_{\max}(v_i): m_{\max}(v_i) \ll m_1}} T_{\mathcal{G}^{v_i}} ((P_{k_{\max}(v_i)}^{(1)} P_{m_{\max}(v_i)}^{(2)} f_l)_{l \in \mathcal{L}(v_i)}),
 \end{aligned}$$

where the equality follows from the fact that $\Delta_k^{(j)} S_k^{(j)} \neq 0$ for $j = 1, 2$ if and only if $\tilde{k} \ll k$. We then apply the high-low switch technique to rewrite (5.81) as

$$\begin{aligned}
 & \sum_{\substack{k_1 \\ m_1}} T_{\tilde{\mathcal{G}}^{v_1}} (\Delta_{k_1}^{(1)} \Delta_{m_1}^{(2)} f_1, (S_{k_1}^{(1)} S_{m_1}^{(2)} f_l)_{l \in \mathcal{L}(v_1)}) \prod_{i=2}^{n_1} \sum_{\substack{k_{\max}(v_i) \\ m_{\max}(v_i)}} T_{\mathcal{G}^{v_i}} ((P_{k_{\max}(v_i)}^{(1)} P_{m_{\max}(v_i)}^{(2)} f_l)_{l \in \mathcal{L}(v_i)}) + \\
 & \sum_{\substack{k_{\max}(v_2) > k_1 \\ m_{\max}(v_2) > m_1}} T_{\tilde{\mathcal{G}}^{v_1}} (\Delta_{k_1}^{(1)} \Delta_{m_1}^{(2)} f_1, (S_{k_1}^{(1)} S_{m_1}^{(2)} f_l)_{l \in \mathcal{L}(v_1)}) T_{\mathcal{G}^{v_2}} ((P_{k_{\max}(v_2)}^{(1)} P_{m_{\max}(v_2)}^{(2)} f_l)_{l \in \mathcal{L}(v_2)}) \\
 & \cdot \prod_{i=3}^{n_1} \sum_{\substack{k_{\max}(v_i) \\ m_{\max}(v_i)}} T_{\mathcal{G}^{v_i}} ((P_{k_{\max}(v_i)}^{(1)} P_{m_{\max}(v_i)}^{(2)} f_l)_{l \in \mathcal{L}(v_i)}) \pm \text{similar terms} := I + II \pm \text{similar terms}.
 \end{aligned}$$

By applying the inductive hypothesis (5.47) to both

$$\sum_{k_1, m_1} T_{\tilde{\mathcal{G}}^{v_1}} (\Delta_{k_1}^{(1)} \Delta_{m_1}^{(2)} f_1, (S_{k_1}^{(1)} S_{m_1}^{(2)} f_l)_{l \in \mathcal{L}(v_1)}) \quad \text{and} \quad \sum_{\substack{k_{\max}(v_i) \\ m_{\max}(v_i)}} T_{\mathcal{G}^{v_i}} ((P_{k_{\max}(v_i)}^{(1)} P_{m_{\max}(v_i)}^{(2)} f_l)_{l \in \mathcal{L}(v_i)})$$

for $i \geq 2$, we derive the estimate on the right hand side of (5.47) for I .

On the other hand, different inductive hypotheses are called for in dealing with different subtrees involved in II . More precisely, (5.47) is invoked to estimate

$$\sum_{\substack{k_{\max}(v_i) \\ m_{\max}(v_i)}} T_{\mathcal{G}^{v_i}} ((P_{k_{\max}(v_i)}^{(1)} P_{m_{\max}(v_i)}^{(2)} f_l)_{l \in \mathcal{L}(v_i)})$$

for $i \geq 3$. Meanwhile, (5.43) is used for

$$T_{\tilde{\mathcal{G}}^{v_1}} (\Delta_{k_1}^{(1)} \Delta_{m_1}^{(2)} f_1, (S_{k_1}^{(1)} S_{m_1}^{(2)} f_l)_{l \in \mathcal{L}(v_1)}).$$

Also, we denote by $R(v_2)$ and $R'(v_2)$ the conical regions associated to the subtree \mathcal{G}^{v_2} for the first and second parameters. Suppose that $\tilde{l} \in \mathfrak{M}_1(R(v_2))$, $\tilde{l}' \in \mathfrak{M}_2(R'(v_2))$ with $\tilde{l} \neq \tilde{l}'$, then (5.44) can be applied. Combining all the subtree estimates, we conclude that

$$\begin{aligned}
 \|II\|_{\tilde{\tau}} & \lesssim \sum_{\substack{\delta_1 |_{\mathcal{V} \setminus \{\tau_{\mathcal{G}}\} \setminus \mathcal{V}_1^{v_1} \setminus \mathcal{V}_i^{v_2}} \\ \delta_2 |_{\mathcal{V} \setminus \{\tau_{\mathcal{G}}\} \setminus \mathcal{V}_1^{v_1} \setminus \mathcal{V}_{i'}^{v_2}}}} \prod_{l \neq 1, \tilde{l}, \tilde{l}'} \|D_{(1)}^{\delta_1^{-1}(l)} D_{(2)}^{\delta_2^{-1}(l)} f_l\|_{\tilde{p}_l} \cdot \sum_{\substack{k_{\max}(v_2) > k_1 \\ m_{\max}(v_2) > m_1}} 2^{k_1 \beta_1^{\tau_{\mathcal{G}}}} 2^{m_1 \beta_2^{\tau_{\mathcal{G}}}} \\
 & \cdot \|\Delta_{k_{\max}}^{(1)} \Delta_{m_{\max}}^{(2)} D_{(1)}^{\beta_1(v_1, 1)} D_{(2)}^{\beta_2(v_1, 1)} f_1\|_{\tilde{p}_1} \cdot \|\Delta_{k_{\max}(v_2)}^{(1)} D_{(1)}^{\beta_1(v_2, \tilde{l})} D_{(2)}^{\delta_2^{-1}(\tilde{l})} f_{\tilde{l}}\|_{\tilde{p}_{\tilde{l}}} \|\Delta_{m_{\max}(v_2)}^{(2)} D_{(1)}^{\delta_1^{-1}(\tilde{l}')} D_{(2)}^{\beta_2(v_2, \tilde{l}')} f_{\tilde{l}'}\|_{\tilde{p}_{\tilde{l}'}}.
 \end{aligned}$$

The distribution of partial derivatives on both parameters gives the following estimate of the inner sum with $\delta_1 |_{\mathcal{V} \setminus \{\tau_{\mathcal{G}}\} \setminus \mathcal{V}_1^{v_1} \setminus \mathcal{V}_i^{v_2}}$ and $\delta_2 |_{\mathcal{V} \setminus \{\tau_{\mathcal{G}}\} \setminus \mathcal{V}_1^{v_1} \setminus \mathcal{V}_{i'}^{v_2}}$ fixed:

$$\begin{aligned}
 & \sum_{\substack{k_{\max}(v_2) \\ m_{\max}(v_2)}} \min(2^{k_{\max}(v_2) \beta_1^{\tau_{\mathcal{G}}}} 2^{m_{\max}(v_2) \beta_2^{\tau_{\mathcal{G}}}} \|D_{(1)}^{\beta_1(v_1, 1)} D_{(2)}^{\beta_2(v_1, 1)} f_1\|_{\dot{B}_{p_1^1, \infty}^0 \dot{B}_{p_1^2, \infty}^0} \|D_{(1)}^{\beta_1(v_2, \tilde{l})} D_{(2)}^{\delta_2^{-1}(\tilde{l})} f_{\tilde{l}}\|_{\dot{B}_{p_1^1, \infty}^0 L_{p_1^2}^2} \|D_{(1)}^{\delta_1^{-1}(\tilde{l}')} D_{(2)}^{\beta_2(v_2, \tilde{l}')} f_{\tilde{l}'}\|_{L_{p_1^1}^{p_1^1} \dot{B}_{p_2^2, \infty}^0}, \\
 & 2^{k_{\max}(v_2) \beta_1^{\tau_{\mathcal{G}}}} 2^{-m_{\max}(v_2) \epsilon_2} \|D_{(1)}^{\beta_1(v_1, 1)} D_{(2)}^{\beta_2(v_1, 1)} f_1\|_{\dot{B}_{p_1^1, \infty}^0 \dot{B}_{p_1^2, \infty}^{\epsilon_2}} \|D_{(1)}^{\beta_1(v_2, \tilde{l})} D_{(2)}^{\delta_2^{-1}(\tilde{l})} f_{\tilde{l}}\|_{\dot{B}_{p_1^1, \infty}^0 L_{p_1^2}^2} \|D_{(1)}^{\delta_1^{-1}(\tilde{l}')} D_{(2)}^{\beta_2(v_2, \tilde{l}')} f_{\tilde{l}'}\|_{L_{p_1^1}^{p_1^1} \dot{B}_{p_2^2, \infty}^{\tau_{\mathcal{G}}}}, \\
 & 2^{-k_{\max}(v_2) \epsilon_1} 2^{m_{\max}(v_2) \beta_2^{\tau_{\mathcal{G}}}} \|D_{(1)}^{\beta_1(v_1, 1)} D_{(2)}^{\beta_2(v_1, 1)} f_1\|_{\dot{B}_{p_1^1, \infty}^{\epsilon_1} \dot{B}_{p_1^2, \infty}^0} \|D_{(1)}^{\beta_1(v_2, \tilde{l})} D_{(2)}^{\delta_2^{-1}(\tilde{l})} f_{\tilde{l}}\|_{\dot{B}_{p_1^1, \infty}^{\tau_{\mathcal{G}}} L_{p_1^2}^2} \|D_{(1)}^{\delta_1^{-1}(\tilde{l}')} D_{(2)}^{\beta_2(v_2, \tilde{l}')} f_{\tilde{l}'}\|_{L_{p_1^1}^{p_1^1} \dot{B}_{p_2^2, \infty}^0}, \\
 & 2^{-k_{\max}(v_2) \epsilon_1} 2^{-m_{\max}(v_2) \epsilon_2} \|D_{(1)}^{\beta_1(v_1, 1)} D_{(2)}^{\beta_2(v_1, 1)} f_1\|_{\dot{B}_{p_1^1, \infty}^{\epsilon_1} \dot{B}_{p_1^2, \infty}^{\epsilon_2}} \|D_{(1)}^{\beta_1(v_2, \tilde{l})} D_{(2)}^{\delta_2^{-1}(\tilde{l})} f_{\tilde{l}}\|_{\dot{B}_{p_1^1, \infty}^{\tau_{\mathcal{G}}} L_{p_1^2}^2} \|D_{(1)}^{\delta_1^{-1}(\tilde{l}')} D_{(2)}^{\beta_2(v_2, \tilde{l}')} f_{\tilde{l}'}\|_{L_{p_1^1}^{p_1^1} \dot{B}_{p_2^2, \infty}^{\tau_{\mathcal{G}}}}),
 \end{aligned}$$

so that the optimization and interpolation can be carried out to derive the estimate on the right hand side of (5.47). This completes proof of Case (i) and provides a generic recipe for treating the remaining cases. \square

We make a final remark on the biparameter flag Leibniz-type estimates associated to asymmetric symbols stated in Theorem 1.2. The inductive procedure described in the previous section, while still applicable, requires a certain modification; we only elaborate on this: the main difference arises in the reduction of the frequency trees. We focus on the example (1.31) with the frequency trees specified below:




FIGURE 6. An asymmetric bi-parameter symbol

We further assume that the multilinear expression (1.31) is localized on the conical region

$$\{(\xi_1, \dots, \xi_5) : |\xi_1| \gg |\xi_l| \text{ for } l \neq 1\} \times \{(\eta_1, \dots, \eta_5) : |\eta_4| \gg |\eta_l| \text{ for } l \neq 4\}.$$

Then we can split the root symbol into a commutator and a symbol associated to subtrees of lower complexity for both parameters. However, due to the asymmetry, all the leaves f_1, \dots, f_5 are intertwined and we cannot decouple any subsets of the leaves (previously associated to subtrees) as we did in the symmetric setting. Instead, we obtain a product of subtrees of lower complexity for both parameters as a reduction.

In the conical region above, one term (modulo modulation and after simplifications) that appears in our estimation is the commutator tensorized with a symbol associated to a subtree:

$$(5.82) \quad \sum_{\substack{k_1 \gg k_3 \\ m_3 \gg m_4}} 2^{k_1(\beta_1-1)} 2^{k_3} \int |\xi_1 + \xi_2|^{\alpha_1} |\xi_3 + \xi_4 + \xi_5|^{\gamma_1} |\xi_3 + \xi_4|^{\zeta_1} |\eta_1 + \eta_3|^{\alpha_2} |\eta_2 + \eta_4|^{\gamma_2 + \beta_2} \mathcal{F}(\Delta_{k_1, +}^{(1)} S_{m_3}^{(2)} f_1)(\xi_1, \eta_1) \mathcal{F}(S_{k_1}^{(1)} S_{m_4}^{(2)} f_2)(\xi_2, \eta_2) \mathcal{F}(\Delta_{k_3}^{(1)} \Delta_{m_3}^{(2)} f_3)(\xi_3, \eta_3) \mathcal{F}(S_{k_3}^{(1)} \Delta_{m_4}^{(2)} f_4)(\xi_4, \eta_4) \mathcal{F}(S_{k_3}^{(1)} f_5)(\xi_5, \eta_5) e^{2\pi i(x, y) \cdot (\xi_1 + \xi_2 + \xi_3 + \xi_4 + \xi_5, \eta_1 + \eta_2 + \eta_3 + \eta_4 + \eta_5)} d\xi d\eta.$$


This hints to the necessity of establishing inductive statements associated to disjoint unions of rooted (sub)trees in both parameters, as opposed to inductive statements just for rooted subtrees. Thus the induction is performed based on the maximal complexity of the rooted trees involved.

In our example, we started with a frequency tree \mathcal{G}_1 of complexity 3 in the first parameter (see Figure 6(i)), and a frequency tree \mathcal{G}_2 of complexity 2 in the second parameter (see Figure 6(ii)); so the maximal complexity of $\mathcal{G}_1 \times \mathcal{G}_2$ is 3. By breaking down the root symbols $m_{\tau_{\mathcal{G}_1}}$ and $m_{\tau_{\mathcal{G}_2}}$, we are led, as suggested by (5.82), to considering the frequency forest $\tilde{\mathcal{G}}_1$ of maximal complexity 2 in the first parameter (see Figure 7), and the frequency forest $\tilde{\mathcal{G}}_2$ of maximal complexity 1 in the second parameter (see Figure 8). Overall, the splitting of the roots' symbols reduces the maximal complexity. More concretely, the inductive hypothesis will be applied to the expression

$$(5.83) \quad \int |\xi_1 + \xi_2|^{\alpha_1} |\xi_3 + \xi_4 + \xi_5|^{\gamma_1} |\xi_3 + \xi_4|^{\zeta_1} |\eta_1 + \eta_3|^{\alpha_2} |\eta_2 + \eta_4|^{\gamma_2 + \beta_2} \mathcal{F}(\Delta_{k_1}^{(1)} S_{m_3}^{(2)} f_1)(\xi_1, \eta_1) \mathcal{F}(S_{k_1}^{(1)} S_{m_4}^{(2)} f_2)(\xi_2, \eta_2) \mathcal{F}(\Delta_{k_3}^{(1)} \Delta_{m_3}^{(2)} f_3)(\xi_3, \eta_3) \mathcal{F}(S_{k_3}^{(1)} \Delta_{m_4}^{(2)} f_4)(\xi_4, \eta_4) \mathcal{F}(S_{k_3}^{(1)} f_5)(\xi_5, \eta_5) e^{2\pi i(x, y) \cdot (\xi_1 + \xi_2 + \xi_3 + \xi_4 + \xi_5, \eta_1 + \eta_2 + \eta_3 + \eta_4 + \eta_5)} d\xi d\eta,$$

which is a multiplier appearing in (5.82) with maximal complexity 2 (attained by the subtree with root symbol $|\xi_3 + \xi_4 + \xi_5|^{\gamma_1}$).

The base case of such an inductive procedure involves an analysis of symbols associated to frequency forests of complexity less than or equal to 1 in each parameter. This can be verified directly by implementing the usual

FIGURE 7. Frequency forest $\tilde{\mathcal{G}}_1$ for the first parameter.FIGURE 8. Frequency forest $\tilde{\mathcal{G}}_2$ for the second parameter.

procedure: paraproduct decompositions, splitting of root symbols, Fourier series decompositions, optimization and Besov norm interpolation.

This ends the discussion on the example (1.2), which is generic enough to illustrate the main ingredients in the proof for multi-parameter flag Leibniz-type estimates associated to asymmetric symbols of arbitrary complexity.

5.3. Optimization and interpolation in the N -parameters setting. This section is devoted to the optimization and interpolation procedure which allows to redistribute the derivatives and produce the expected geometric and arithmetic means in the generic N -parameters setting. As a consequence of this procedure, we derive the desired estimates in the statements (1)–(6) of Proposition 5.3.

We first introduce some notation. Let \mathcal{H} and \mathcal{L} denote maps

$$\begin{aligned} \mathcal{H} : i \in \{\text{Index set of the parameters}\} &\mapsto l \in \{\text{Index set for the functions}\} \\ \mathcal{L} : i \in \{\text{Index set of the parameters}\} &\mapsto l \in \{\text{Index set for the functions}\}, \end{aligned}$$

where the map \mathcal{H} indicates which functions would be hit by the full order of derivatives and the map \mathcal{L} illustrates the functions hit by the lower order of derivatives. Equations (5.34)–(5.36) are representative of this action, and in (5.35) in particular we can see that in the region $\{|\xi_1| \gg |\xi_2|, \dots, |\xi_n|\}$ the function f_1 receives $D^{\beta^{\tau_g}}$ derivatives and f_{l_0} receives D^ϵ derivatives.

We denote by Σ_N the set of length- N signatures:

$$(5.84) \quad \Sigma_N := \{\sigma = (\varsigma_1, \dots, \varsigma_N) : \varsigma_1, \dots, \varsigma_N \in \{+, -\}\}$$

Our index set of parameters is $\{1, \dots, N\}$, and the index set of functions $\{1, \dots, n\}$. The above convention implies that, on the specific frequency conical region to which we restrict our operator, in the first parameter the scales $k_{\mathcal{H}(1)} \geq k_{\mathcal{L}(1)}$ (and thus the functions $f_{\mathcal{H}(1)}$ and $f_{\mathcal{L}(1)}$) are involved in the estimation of the flag, in the second parameter the scales $k_{\mathcal{H}(2)} \geq k_{\mathcal{L}(2)}$ (and the functions $f_{\mathcal{H}(2)}$ and $f_{\mathcal{L}(2)}$), and so on.

In order to simplify the notation, let us assume that in estimating an n -linear, N -parameter flag operator $T_{\mathcal{G}}$ associated to a rooted tree with root information $(\beta_1, \dots, \beta_N)$ and restricted⁴³ to frequency conical regions on each subtree, the functions f_1, f_2, \dots, f_m are involved. Given the discussion in the previous section, we are reduced to⁴⁴

$$\begin{aligned} \|T_{\mathcal{G}}(f_1, \dots, f_n)\|_{L^{\vec{p}}} &\lesssim \prod_{l=m+1}^n \|F_l\|_{L^{\vec{p}_l}} \\ \sum_{\ell_1, \dots, \ell_N} \min(2^{\beta_1 \ell_1} \dots 2^{\beta_{N-1} \ell_{N-1}} \cdot 2^{\beta_N \ell_N} \cdot \prod_{l=1}^m \|F_l\|_{\tilde{W}_{l; (+, \dots, +, +)}}, 2^{\beta_1 \ell_1} \dots 2^{\beta_{N-1} \ell_{N-1}} \cdot 2^{-\epsilon \ell_N} \cdot \prod_{l=1}^m \|F_l\|_{\tilde{W}_{l; (+, \dots, +, -)}} \\ &\vdots \end{aligned}$$

⁴³Otherwise $T_{\mathcal{G}}$ will be bounded above by a some of similar terms, each to be estimated through the present analysis.

⁴⁴Again, for simplicity, we omit the exponent $\tau \leq \min_{1 \leq l \leq N} p^l$; at this stage of the proof it plays no role.

$$2^{-\epsilon_1 \ell_1} \dots 2^{-\epsilon_{N-1} \ell_{N-1}} \cdot 2^{\beta_N \ell_N} \cdot \prod_{l=1}^m \|F_l\|_{\tilde{W}_l;(-,\dots,-,+)} \cdot 2^{-\epsilon_1 \ell_1} \dots 2^{-\epsilon_{N-1} \ell_{N-1}} \cdot 2^{-\epsilon \ell_N} \cdot \prod_{l=1}^m \|F_l\|_{\tilde{W}_l;(-,\dots,-,-)},$$

where $\|F_l\|_{\tilde{W}_l;\sigma}$ – for $\sigma \in \Sigma_N$ – is the mixed norm

$$(5.85) \quad \|F_l\|_{\tilde{W}_l;\sigma} := \|F_l\|_{W_l^{1,\varsigma_1} W_l^{2,\varsigma_2} \dots W_l^{N-1,\varsigma_{N-1}} W_l^{N,\varsigma_N}},$$

and for any $1 \leq l \leq n$

$$(5.86) \quad F_l := D_{(1)}^{\alpha_1^1} \dots D_{(N)}^{\alpha_i^N} f_l$$

records the derivatives picked up by the function f_l on the subtrees of \mathcal{G} (which are of lower complexity).

Above, we define for any $1 \leq l \leq n$ and any $1 \leq i \leq N$,

$$(5.87) \quad W_l^{i,\varsigma_i} := \begin{cases} \dot{B}_{p_l^i,\infty}^0, & \text{if } l = \mathcal{H}(i) \text{ or } l = \mathcal{L}(i) \text{ and } \varsigma_i = + \\ \dot{B}_{p_l^i,\infty}^{\beta_i}, & \text{if } l = \mathcal{H}(i) \text{ and } \varsigma_i = - \\ \dot{B}_{p_l^i,\infty}^{\epsilon_i}, & \text{if } l = \mathcal{L}(i) \text{ and } \varsigma_i = - \\ L^{p_l^i} & \text{if } l \notin \{\mathcal{H}(i), \mathcal{L}(i)\}. \end{cases}$$

Then it is not difficult to see⁴⁵ that

$$\begin{aligned} & \sum_{\ell_1, \dots, \ell_N} \min \left(2^{\beta_1 \ell_1} \dots 2^{\beta_{N-1} \ell_{N-1}} \cdot 2^{\beta_N \ell_N} \cdot \prod_{l=1}^m \|F_l\|_{\tilde{W}_l;(+,\dots,+,+)} \cdot 2^{\beta_1 \ell_1} \dots 2^{\beta_{N-1} \ell_{N-1}} \cdot 2^{-\epsilon \ell_N} \cdot \prod_{l=1}^m \|F_l\|_{\tilde{W}_l;(+,\dots,+,-)} \right. \\ & \quad \vdots \\ & \quad \left. 2^{-\epsilon_1 \ell_1} \dots 2^{-\epsilon_{N-1} \ell_{N-1}} \cdot 2^{\beta_N \ell_N} \cdot \prod_{l=1}^m \|F_l\|_{\tilde{W}_l;(-,\dots,-,+)} \cdot 2^{-\epsilon_1 \ell_1} \dots 2^{-\epsilon_{N-1} \ell_{N-1}} \cdot 2^{-\epsilon \ell_N} \cdot \prod_{l=1}^m \|F_l\|_{\tilde{W}_l;(-,\dots,-,-)} \right) \\ & \lesssim \prod_{\sigma=(\varsigma_1, \dots, \varsigma_N) \in \Sigma_N} \left(\prod_{l=1}^m \|F_l\|_{\tilde{W}_l;\sigma} \right)^{\nu_{\varsigma_1} \dots \nu_{\varsigma_N}}, \end{aligned}$$

where for any $1 \leq i \leq N$, ν_{ς_i} is defined by

$$(5.88) \quad \nu_{\varsigma_i} := \begin{cases} \frac{\epsilon_i}{\beta_i + \epsilon_1}, & \text{if } \varsigma_i = + \\ \frac{\beta_i}{\beta_i + \epsilon_1}, & \text{if } \varsigma_i = -. \end{cases}$$

If we denote $\nu_\sigma := \nu_{\varsigma_1} \dots \nu_{\varsigma_N}$, we have

$$\sum_{\sigma \in \Sigma_N} \nu_\sigma = \prod_{i=1}^N \left(\frac{\epsilon_i}{\beta_i + \epsilon_1} + \frac{\beta_i}{\beta_i + \epsilon_1} \right) = 1.$$

This observation and the previous notation allows us to deduce

$$(5.89) \quad \|T_{\mathcal{G}}(f_1, \dots, f_n)\|_{L^{\tilde{p}}} \lesssim \prod_{\sigma=(\varsigma_1, \dots, \varsigma_N) \in \Sigma_N} \left(\prod_{l=1}^n \|F_l\|_{\tilde{W}_l;\sigma} \right)^{\nu_{\varsigma_1} \dots \nu_{\varsigma_N}}.$$

Now our task is to replace $\prod_{l=1}^n \|F_l\|_{\tilde{W}_l;\sigma}$ by more suitable expressions that allow to keep track of the distribution of derivatives. This will be done in N steps, which corresponds to the number of parameters.

Before proceeding, we recall the interpolation result

$$(5.90) \quad \|F_l\|_{\dot{B}_{p_l^1,\infty}^{\epsilon_1}(\tilde{W}_l;\tilde{\sigma})} \leq \|F_l\|_{\dot{B}_{p_l^1,\infty}^0(\tilde{W}_l;\tilde{\sigma})}^{\frac{\beta_1 - \epsilon_1}{\beta_1}} \cdot \|F_l\|_{\dot{B}_{p_l^1,\infty}^{\beta_1}(\tilde{W}_l;\tilde{\sigma})}^{\frac{\epsilon_1}{\beta_1}},$$

where $\tilde{\sigma} \in \Sigma_{N-1}$ is a length- $(N-1)$ signature vector and \tilde{W}_l is the iteration of $N-1$ vector spaces.

We want to replace $\dot{B}_{p_l^1,\infty}^{\epsilon_1}$ appearing in (5.89) either by $\dot{B}_{p_l^1,\infty}^0$ or by $\dot{B}_{p_l^1,\infty}^{\beta_1}$. We notice that $\dot{B}_{p_l^1,\infty}^{\epsilon_1}$ appears in the norm

$$\|F_l\|_{\tilde{W}_l;\sigma}$$

of precisely those l satisfying $l = \mathcal{L}(1)$, and $\varsigma_1 = -$.

⁴⁵Indeed, such a result can very easily be proved via an induction argument on N , the number of parameters.

This observation, together with the fact that the right-hand side of (5.89) equals⁴⁶

$$\begin{aligned}
& \sum_{\tilde{\sigma} \in \Sigma_{N-1}} \left(\|F_{\mathcal{H}(1)}\|_{W_{\mathcal{H}(1)}^{1,+}(\tilde{W}_l; \tilde{\sigma})} \cdot \|F_{\mathcal{L}(1)}\|_{W_{\mathcal{L}(1)}^{1,+}(\tilde{W}_l; \tilde{\sigma})} \prod_{l \neq \mathcal{H}(1), \mathcal{L}(1)}^n \|F_l\|_{L^{p_l^1}(\tilde{W}_l; \tilde{\sigma})} \right)^{\nu_+ \cdot \nu_{\tilde{\sigma}}} \\
& \cdot \left(\|F_{\mathcal{H}(1)}\|_{W_{\mathcal{H}(1)}^{1,-}(\tilde{W}_l; \tilde{\sigma})} \cdot \|F_{\mathcal{L}(1)}\|_{W_{\mathcal{L}(1)}^{1,-}(\tilde{W}_l; \tilde{\sigma})} \prod_{l \neq \mathcal{H}(1), \mathcal{L}(1)}^n \|F_l\|_{L^{p_l^1}(\tilde{W}_l; \tilde{\sigma})} \right)^{\nu_- \cdot \nu_{\tilde{\sigma}}} \\
& = \sum_{\tilde{\sigma} \in \Sigma_{N-1}} \left(\|F_{\mathcal{H}(1)}\|_{\dot{B}_{p_{\mathcal{H}(1)}^1, \infty}^0(\tilde{W}_l; \tilde{\sigma})} \cdot \|F_{\mathcal{L}(1)}\|_{\dot{B}_{p_{\mathcal{L}(1)}^1, \infty}^0(\tilde{W}_l; \tilde{\sigma})} \prod_{l \neq \mathcal{H}(1), \mathcal{L}(1)}^n \|F_l\|_{L^{p_l^1}(\tilde{W}_l; \tilde{\sigma})} \right)^{\nu_+ \cdot \nu_{\tilde{\sigma}}} \\
& \cdot \left(\|F_{\mathcal{H}(1)}\|_{\dot{B}_{p_{\mathcal{H}(1)}^1, \infty}^{\beta_1}(\tilde{W}_l; \tilde{\sigma})} \cdot \|F_{\mathcal{L}(1)}\|_{\dot{B}_{p_{\mathcal{L}(1)}^1, \infty}^{\epsilon_1}(\tilde{W}_l; \tilde{\sigma})} \prod_{l \neq \mathcal{H}(1), \mathcal{L}(1)}^n \|F_l\|_{L^{p_l^1}(\tilde{W}_l; \tilde{\sigma})} \right)^{\nu_- \cdot \nu_{\tilde{\sigma}}},
\end{aligned}$$

naturally suggests the use of interpolation. If $\tilde{\sigma} \in \Sigma_{N-1}$ is fixed, then the expression appearing in the last display equals

$$\begin{aligned}
& \left(\prod_{l \neq \mathcal{H}(1), \mathcal{L}(1)}^n \|F_l\|_{L^{p_l^1}(\tilde{W}_l; \tilde{\sigma})} \right)^{\nu_{\tilde{\sigma}}} \\
& \cdot \left((\|F_{\mathcal{H}(1)}\|_{\dot{B}_{p_{\mathcal{H}(1)}^1, \infty}^0(\tilde{W}_l; \tilde{\sigma})} \cdot \|F_{\mathcal{L}(1)}\|_{\dot{B}_{p_{\mathcal{L}(1)}^1, \infty}^0(\tilde{W}_l; \tilde{\sigma})})^{\frac{\epsilon_1}{\beta_1 + \epsilon_1}} \cdot (\|F_{\mathcal{H}(1)}\|_{\dot{B}_{p_{\mathcal{H}(1)}^1, \infty}^{\beta_1}(\tilde{W}_l; \tilde{\sigma})} \cdot \|F_{\mathcal{L}(1)}\|_{\dot{B}_{p_{\mathcal{L}(1)}^1, \infty}^{\epsilon_1}(\tilde{W}_l; \tilde{\sigma})})^{\frac{\beta_1}{\beta_1 + \epsilon_1}} \right)^{\nu_{\tilde{\sigma}}}.
\end{aligned}$$

The interpolation in (2.11) and direct computations similar to (2.34) allow to bound this expression by

$$\begin{aligned}
& \left(\prod_{l \neq \mathcal{H}(1), \mathcal{L}(1)}^n \|F_l\|_{L^{p_l^1}(\tilde{W}_l; \tilde{\sigma})} \right)^{\nu_{\tilde{\sigma}}} \\
& \cdot \left((\|F_{\mathcal{H}(1)}\|_{\dot{B}_{p_{\mathcal{H}(1)}^1, \infty}^0(\tilde{W}_l; \tilde{\sigma})} \cdot \|F_{\mathcal{L}(1)}\|_{\dot{B}_{p_{\mathcal{L}(1)}^1, \infty}^{\beta_1}(\tilde{W}_l; \tilde{\sigma})})^{\frac{\epsilon_1}{\beta_1 + \epsilon_1}} \cdot (\|F_{\mathcal{H}(1)}\|_{\dot{B}_{p_{\mathcal{H}(1)}^1, \infty}^{\beta_1}(\tilde{W}_l; \tilde{\sigma})} \cdot \|F_{\mathcal{L}(1)}\|_{\dot{B}_{p_{\mathcal{L}(1)}^1, \infty}^0(\tilde{W}_l; \tilde{\sigma})})^{\frac{\beta_1}{\beta_1 + \epsilon_1}} \right)^{\nu_{\tilde{\sigma}}} \\
& = \left(\|F_{\mathcal{H}(1)}\|_{\dot{B}_{p_{\mathcal{H}(1)}^1, \infty}^0(\tilde{W}_l; \tilde{\sigma})} \cdot \|F_{\mathcal{L}(1)}\|_{\dot{B}_{p_{\mathcal{L}(1)}^1, \infty}^{\beta_1}(\tilde{W}_l; \tilde{\sigma})} \prod_{l \neq \mathcal{H}(1), \mathcal{L}(1)}^n \|F_l\|_{L^{p_l^1}(\tilde{W}_l; \tilde{\sigma})} \right)^{\nu_{(+, \tilde{\sigma})}} \\
& \cdot \left(\|F_{\mathcal{H}(1)}\|_{\dot{B}_{p_{\mathcal{H}(1)}^1, \infty}^{\beta_1}(\tilde{W}_l; \tilde{\sigma})} \cdot \|F_{\mathcal{L}(1)}\|_{\dot{B}_{p_{\mathcal{L}(1)}^1, \infty}^0(\tilde{W}_l; \tilde{\sigma})} \prod_{l \neq \mathcal{H}(1), \mathcal{L}(1)}^n \|F_l\|_{L^{p_l^1}(\tilde{W}_l; \tilde{\sigma})} \right)^{\nu_{(-, \tilde{\sigma})}}.
\end{aligned}$$

In conclusion, (5.89) is replaced by

$$(5.91) \quad \|T_{\mathcal{G}}(f_1, \dots, f_n)\|_{L^{\vec{p}}} \lesssim \prod_{\sigma = (\varsigma_1, \dots, \varsigma_N) \in \Sigma_N} \left(\prod_{l=1}^n \|F_l\|_{\tilde{W}_l^1; \sigma} \right)^{\nu_{\varsigma_1} \dots \nu_{\varsigma_N}},$$

where $\tilde{W}_l^1; \sigma := X_l^{1, \varsigma_1} W_l^{2, \varsigma_2} \dots W_l^{N-1, \varsigma_{N-1}} W_l^{N, \varsigma_N}$ and

$$(5.92) \quad X_l^{1, \varsigma_1} := \begin{cases} \dot{B}_{p_l^1, \infty}^0, & \text{if } l = \mathcal{H}(1) \text{ and } \varsigma_1 = + \\ \dot{B}_{p_l^1, \infty}^{\beta_1}, & \text{if } l = \mathcal{L}(1) \text{ and } \varsigma_1 = + \\ \dot{B}_{p_l^1, \infty}^{\beta_1}, & \text{if } l = \mathcal{H}(1) \text{ and } \varsigma_1 = - \\ \dot{B}_{p_l^1, \infty}^0, & \text{if } l = \mathcal{L}(1) \text{ and } \varsigma_1 = - \\ L^{p_l^1} & \text{if } l \notin \{\mathcal{H}(1), \mathcal{L}(1)\}. \end{cases}$$

We iterate the procedure to obtain

$$(5.93) \quad \|T_{\mathcal{G}}(f_1, \dots, f_n)\|_{L^{\vec{p}}} \lesssim \prod_{\sigma = (\varsigma_1, \dots, \varsigma_N) \in \Sigma_N} \left(\prod_{l=1}^n \|F_l\|_{\tilde{W}_l^2; \sigma} \right)^{\nu_{\varsigma_1} \dots \nu_{\varsigma_N}},$$

where $\tilde{W}_l^2; \sigma := X_l^{1, \varsigma_1} X_l^{2, \varsigma_2} W_l^{3, \varsigma_3} \dots W_l^{N-1, \varsigma_{N-1}} W_l^{N, \varsigma_N}$ and X_l^{2, ς_2} is defined in a similar way to (5.93).

⁴⁶We also use the fact that for $l \neq \mathcal{H}(1), \mathcal{L}(1)$, $W_l^1 = L^{p_l^1}$.

After N steps we conclude that

$$(5.94) \quad \|T_{\mathcal{G}}(f_1, \dots, f_n)\|_{L^{\vec{p}}} \lesssim \prod_{\sigma=(\varsigma_1, \dots, \varsigma_N) \in \Sigma_N} \left(\prod_{l=1}^n \|F_l\|_{\vec{W}_l^N; \sigma} \right)^{\nu_{\sigma}},$$

where $\vec{W}_l^N; \sigma := X_l^{1, \varsigma_1} \dots X_l^{N, \varsigma_N}$ and

$$(5.95) \quad X_l^{i, \varsigma_i} := \begin{cases} \dot{B}_{p_i^i, \infty}^0, & \text{if } l = \mathcal{H}(i) \text{ and } \varsigma_i = + \\ \dot{B}_{p_i^i, \infty}^{\beta_i}, & \text{if } l = \mathcal{L}(i) \text{ and } \varsigma_i = + \\ \dot{B}_{p_i^i, \infty}^{\beta_i}, & \text{if } l = \mathcal{H}(i) \text{ and } \varsigma_i = - \\ \dot{B}_{p_i^i, \infty}^0, & \text{if } l = \mathcal{L}(i) \text{ and } \varsigma_i = - \\ L^{p_i^i} & \text{if } l \notin \{\mathcal{H}(i), \mathcal{L}(i)\}. \end{cases}$$

If we carefully read inequality (5.94), we have obtained that the N -parameter n -linear flag $T_{\mathcal{G}}(f_1, \dots, f_n)$ is bounded above in the mixed norm $\|\cdot\|_{L^{\vec{p}}}$ by the geometric mean of 2^N terms (this is because $\sum_{\sigma \in \Sigma_N} \nu_{\sigma} = 1$), and each term is of the form

$$\prod_{l=1}^n \|F_l\|_{X_l^{1, \varsigma_1} \dots X_l^{N, \varsigma_N}}.$$

For every parameter $1 \leq i \leq N$, there are exactly two indices $\mathcal{H}(i)$ and $\mathcal{L}(i)$ in the index set $\{1, \dots, n\}$ for the functions for which

$$(5.96) \quad (X_{\mathcal{H}(i)}^{i, \varsigma_i}, X_{\mathcal{L}(i)}^{i, \varsigma_i}) = (\dot{B}_{p_{\mathcal{H}(i)}^i, \infty}^0, \dot{B}_{p_{\mathcal{L}(i)}^i, \infty}^{\beta_i}) \quad \text{or} \quad (X_{\mathcal{H}(i)}^{i, \varsigma_i}, X_{\mathcal{L}(i)}^{i, \varsigma_i}) = (\dot{B}_{p_{\mathcal{H}(i)}^i, \infty}^{\beta_i}, \dot{B}_{p_{\mathcal{L}(i)}^i, \infty}^0),$$

while for the other indices $l \neq \mathcal{H}(i), \mathcal{L}(i)$ we simply have $X_l^{i, \varsigma_i} = L^{p_i^i}$. The identity (5.96) means precisely that the derivatives are being distributed accordingly: for every $1 \leq i \leq N$, each $\dot{B}_{(i)}^{\beta_i}$ is shared between $F_{\mathcal{H}(i)}$ and $F_{\mathcal{L}(i)}$. Finally, if we undo the definition (5.86), we deduce that $T_{\mathcal{G}}(f_1, \dots, f_n)$, restricted to conical regions according to the subtree structures, is indeed controlled by the geometric mean of terms of the form $\prod_{l=1}^n \|D_{(1)}^{\delta_1^{-1}(l)} \dots D_{(N)}^{\delta_N^{-1}(l)} f_l\|_{\vec{p}_l}$, where $\delta_1 \otimes \dots \otimes \delta_N$ satisfy conditions (i) and (ii) from Section 1.

6. MULTILINEAR OPERATORS OF POSITIVE ORDER

After having worked out the multi-parameter flag Leibniz rules, we discuss in this section Leibniz-type estimates for flag structures associated to Mikhlin symbols of positive order. Our intention here is to provide more examples for which the method introduced by Bourgain and Li [4] offers an alternative to well-established techniques.

We start with the observation that on the region⁴⁷ $R_1 = \{(\xi_1, \dots, \xi_n) \in \mathbb{R}^{dn} : |\xi_1| \gg |\xi_2|, \dots, |\xi_n|\}$, the symbol $|\xi_1 + \dots + \xi_n|^{\beta}$ – which is naturally associated to Leibniz rules for $D^{\beta}(f_1 \dots f_n)$ – is a Mikhlin symbol of order β , satisfying (1.33) for arbitrarily many multi-indices $\gamma_1, \dots, \gamma_n$. Because of this, in what follows we consider Mikhlin symbols of positive order, arbitrarily smooth away from the origin. Under this assumption, we provide a sketch of the proofs of Theorems 1.3, 1.4, 1.5. Afterwards we discuss *smoothing properties* for multilinear operators associated to symbols of negative order, such as multilinear fractional integral operators.

We would like to remark that Theorem 1.3 follows from the boundedness of one-parameter flag paraproducts [14] and Theorem 1.5 is implied by the boundedness of multi-parameter paraproducts [15]. Moreover, the smoothing property described in Theorem 6.3 in the one-parameter mixed-norm setting has been resolved by Hart-Torres-Wu [10], and its bi-parameter, non-mixed-norm variant was established in Yang-Liu-Wu [19]; both cases focus on symbols of limited regularity whereas we handle smooth symbols satisfying pointwise decay conditions.

6.1. Leibniz-type estimates for flag operators associated to Mikhlin symbols of positive order. The natural multi-parameter adaptation of Theorem 1.1 in this context is provided exactly by Theorem 1.4, in which the Mikhlin symbols associated to each vertex are assumed to be products of Mikhlin symbols in each parameter, i.e. they satisfy (1.36). Because the symbols tensorize, the multi-parameter extension will follow closely the procedures described in Section 4, Section 5.2 and Section 5.3. For that reason, we will focus on the one-parameter case – Theorem 1.3.

⁴⁷On the other hand, in the region $\tilde{R}_{1,2} = \{(\xi_1, \dots, \xi_n) \in \mathbb{R}^{dn} : |\xi_1| \sim |\xi_2| \gg |\xi_3|, \dots, |\xi_n|\}$, $|\xi_1 + \dots + \xi_n|^{\beta}$ is less regular than a Mikhlin symbol of order β , unless $\beta \in 2\mathbb{N}$. In general, for a Mikhlin symbol m of order β , $m(\xi_1 + \dots + \xi_n)$ is singular along the subspace $\{(\xi_1, \dots, \xi_n) : \xi_1 + \dots + \xi_n = 0\}$, and it can be seen as a natural extension of $|\xi_1 + \dots + \xi_n|^{\beta}$. Nonetheless, it is not difficult to verify that our method developed in this section also applies to such symbols.

Remark 6.1. Since the multipliers considered are smooth away from the origin (they satisfy (1.33)), they are smooth on every Whitney rectangle and as a consequence the corresponding Fourier coefficients have arbitrary decay. This comes in contrast with the Leibniz rules presented in Theorem 1.1, where conditions (1.25) and (1.26) are necessary. Notice also that in this situation we do not obtain the endpoint estimates corresponding to $p_l = 1$ or $p_l = \infty$, although some of the L^∞ endpoints can be proved through the Coifman-Meyer approach.

Sketch of Proof of Theorem 1.3. Here again the inductive procedure will follow closely the steps described in Section 5.1, and it will be enough to discuss steps (1) and (2) of our strategy presented in Section 1.2: the splitting of the root symbol (and the simultaneous appearance of commutators), and the Fourier series decomposition for the new symbols. Once these steps performed, the operator associated to the rooted tree \mathcal{G} naturally tensorized into operators associated to rooted subtrees of lower complexity; in many situations, the Mikhlin symbol $m_{\beta_{\bar{v}}}(\xi_{i_1}, \dots, \xi_{i_s})$ – associated to the root of a subtree – will be replaced by the product $m_{\beta_{\bar{v}}}(\xi_{i_1}, \dots, \xi_{i_s}) \cdot m_{\beta_{r_{\mathcal{G}}}}(0, \dots, \xi_{i_1}, 0, \dots, \xi_{i_s}, 0, \dots)$, which is again a Mikhlin symbol of order $\beta_{r_{\mathcal{G}}} + \beta_{\bar{v}} > 0$ in the variables $\xi_{i_1}, \dots, \xi_{i_s}$.

For illustrative purposes, we will focus on the one-parameter Leibniz-type estimate of complexity 1 and explain how to achieve the steps (1) and (2) mentioned above. In particular, we consider

$$T_{m_\beta}(f_1, \dots, f_n)(x) := \int_{\mathbb{R}^{nd}} m_\beta(\xi_1, \dots, \xi_n) \hat{f}_1(\xi_1) \cdot \dots \cdot \hat{f}_n(\xi_n) e^{2\pi i x \cdot (\xi_1 + \dots + \xi_n)} d\xi_1 \dots d\xi_n,$$

when restricted to two typical regions:

$$(6.1) \quad \begin{aligned} R_1 &:= \{(\xi_1, \dots, \xi_n) : |\xi_1| \gg |\xi_2|, |\xi_3|, \dots, |\xi_n|\} \\ \tilde{R}_{1,2} &:= \{(\xi_1, \dots, \xi_n) : |\xi_1| \sim |\xi_2| \gg |\xi_3|, \dots, |\xi_n|\}. \end{aligned}$$

The first region corresponds to the situation when one of $|\xi_{j_0}|$ is much larger than the remaining variables, and so $|(\xi_1, \dots, \xi_n)| \sim |\xi_{j_0}|$. For simplicity and without loss of generality, we assume that $j_0 = 1$. The second situation corresponds to the “diagonal case”, when there exist $j_1 \neq j_2$ with $|\xi_{j_1}| \sim |\xi_{j_2}|$ larger than the norm of the remaining variables; then $|(\xi_1, \dots, \xi_n)| \sim |\xi_{j_1}| \sim |\xi_{j_2}|$ and we assume that $j_1 = 1, j_2 = 2$.

(1) The multiplier localized on the off-diagonal conical region $\{(\xi_1, \dots, \xi_n) : |\xi_1| \gg |\xi_2|, \dots, |\xi_n|\}$ can be expressed as

$$(6.2) \quad \sum_{k_1 \gg k_2, \dots, k_n} \int_{\mathbb{R}^{nd}} m_\beta(\xi_1, \dots, \xi_n) \widehat{\Delta_{k_1} f_1}(\xi_1) \cdot \dots \cdot \widehat{\Delta_{k_n} f_n}(\xi_n) e^{2\pi i x \cdot (\xi_1 + \dots + \xi_n)} d\xi_1 \dots d\xi_n.$$

For this, we approximate $m_\beta(\xi_1, \xi_2, \dots, \xi_n)$ by $m_\beta(\xi_1, 0, \dots, 0)$ and in consequence we need to study the “commutator” $m_\beta(\xi_1, \xi_2, \dots, \xi_n) - m_\beta(\xi_1, 0, \dots, 0)$. We notice the following⁴⁸:

$$m_\beta(\xi_1, \xi_2, \dots, \xi_n) - m_\beta(\xi_1, 0, \dots, 0) = \int_0^1 \frac{d}{dt} (m_\beta(\xi_1, t\xi_2, \dots, t\xi_n)) dt = \int_0^1 \sum_{l=2}^n \nabla_{\xi_l} m_\beta(\xi_1, t\xi_2, \dots, t\xi_n) \cdot \xi_l dt.$$

Now we fix $2 \leq l \leq n$ and assume without loss of generality that $l = 2$. The functions f_1 and f_2 will play a prominent role, and for this reason we can sum over $k_3, \dots, k_n \ll k_1$ in (6.2). We would like to implement the Fourier series decomposition of

$$(6.3) \quad \left(\int_0^1 \nabla_2 m_\beta(\xi_1, t\xi_2, \dots, t\xi_n) \cdot \xi_2 dt \right) \tilde{\psi}_{k_1}(\xi_1) \tilde{\psi}_{k_2}(\xi_2) \tilde{\varphi}_{k_1}(\xi_3) \cdot \dots \cdot \tilde{\varphi}_{k_1}(\xi_n),$$

which however requires a further restriction of the symbol. For this, we cover the annuli $\{|\xi_1| \sim 2^{k_1}\}$ and $\{|\xi_2| \sim 2^{k_2}\}$ with Whitney cubes associated to directional cones as described in Remark 2.4:

$$\{|\xi_1| \sim 2^{k_1}\} \subseteq \bigcup_{c_1 \in \mathcal{C}} \{|\xi_1| \sim 2^{k_1}\} \cap c_1, \quad \{|\xi_2| \sim 2^{k_2}\} \subseteq \bigcup_{c_2 \in \mathcal{C}} \{|\xi_2| \sim 2^{k_2}\} \cap c_2.$$

Hence the symbol (6.3) can be rewritten as

$$\sum_{c_1, c_2 \in \mathcal{C}} \int_0^1 \nabla_2 m_\beta(\xi_1, t\xi_2, \dots, t\xi_n) \cdot \xi_2 \tilde{\psi}_{k_1, c_1}(\xi_1) \tilde{\psi}_{k_1}(\xi_1) \tilde{\psi}_{k_2, c_2}(\xi_2) \tilde{\psi}_{k_2}(\xi_2) \cdot \tilde{\varphi}_{k_1}(\xi_3) \dots \tilde{\varphi}_{k_1}(\xi_n) dt.$$

⁴⁸Throughout the section, for a function $m : \mathbb{R}^{dn} \rightarrow \mathbb{C}$ and any $1 \leq j \leq n$, $\partial_{\xi_j} m$ or $\nabla_j m$ denotes the vector

$$\partial_{\xi_j} m(\xi_1, \dots, \xi_n) = (\partial_{\xi_1} m(\xi_1, \dots, \xi_n), \dots, \partial_{\xi_j} m(\xi_1, \dots, \xi_n)).$$

For each summand with $c_1, c_2 \in \mathfrak{C}$ fixed, we apply the Fourier series decomposition to obtain

$$\sum_{L_1, \dots, L_n \in \mathbb{Z}^d} C_{L_1, \dots, L_n}^{k_1, k_2, c_1, c_2} e^{2\pi i L_1 \cdot \frac{\xi_1}{2^{k_1}}} e^{2\pi i L_2 \cdot \frac{\xi_2}{2^{k_2}}} e^{2\pi i L_3 \cdot \frac{\xi_3}{2^{k_1}}} \dots e^{2\pi i L_n \cdot \frac{\xi_n}{2^{k_1}}}$$

on the Whitney rectangle $Q_{k_1, c_1} \times Q_{k_2, c_2} \times \{|\xi_3| \lesssim 2^{k_1}\} \times \dots \times \{|\xi_n| \lesssim 2^{k_1}\}$.

The Fourier coefficients $C_{L_1, \dots, L_n}^{k_1, k_2, c_1, c_2}$ can be written as

$$2^{-k_1 d(n-1)} 2^{-k_2 d} \int_0^1 \int_{\mathbb{R}^{nd}} \nabla_2 m_\beta(\xi_1, t\xi_2, t\xi_3, \dots, t\xi_n) \cdot \xi_2 \tilde{\psi}_{k_1, c_1}(\xi_1) \tilde{\psi}_{k_2, c_2}(\xi_2) \cdot \tilde{\psi}_{k_1}(\xi_1) \cdot \tilde{\psi}_{k_2}(\xi_2) \\ \cdot \tilde{\varphi}_{k_1}(\xi_3) \dots \tilde{\varphi}_{k_1}(\xi_n) \cdot e^{2\pi i L_1 \cdot \frac{\xi_1}{2^{k_1}}} e^{2\pi i L_2 \cdot \frac{\xi_2}{2^{k_2}}} e^{2\pi i L_3 \cdot \frac{\xi_3}{2^{k_1}}} \dots e^{2\pi i L_n \cdot \frac{\xi_n}{2^{k_1}}} d\xi_1 \dots d\xi_n dt.$$

By change of variables $\xi'_l := 2^{-k_1} \xi_l$ for $\tilde{l} \neq 2$ and $\xi'_2 := 2^{-k_2} \xi_2$, it becomes

$$(6.4) \quad 2^{k_2} \int_0^1 \int_{\mathbb{R}^{nd}} \nabla_2 m_\beta(2^{k_1} \xi'_1, t2^{k_2} \xi'_2, t2^{k_1} \xi'_3, \dots, t2^{k_1} \xi'_n) \cdot \xi'_2 \tilde{\psi}_{0, c_1}(\xi'_1) \tilde{\psi}_{0, c_2}(\xi'_2) \tilde{\psi}(\xi'_1) \tilde{\psi}(\xi'_2) \\ \cdot \tilde{\varphi}_0(\xi'_3) \dots \tilde{\varphi}_0(\xi'_n) e^{2\pi i L_1 \cdot \xi'_1} e^{2\pi i L_2 \cdot \xi'_2} e^{2\pi i L_3 \cdot \xi'_3} \dots e^{2\pi i L_n \cdot \xi'_n} d\xi'_1 \dots d\xi'_n dt.$$

Using integration by parts, we can bound (6.4) by

$$(6.5) \quad 2^{k_2} (1 + |L_1| + \dots + |L_n|)^{-|\tilde{M}|} \int_0^1 \int_{\mathbb{R}^{nd}} |\partial_{\xi'_1}^{\tilde{M}} (\nabla_2 m_\beta(2^{k_1} \xi'_1, t2^{k_2} \xi'_2, t2^{k_1} \xi'_3, \dots, t2^{k_1} \xi'_n) \cdot \xi'_2) | \\ |\tilde{\psi}_{0, c_1}(\xi'_1) \tilde{\psi}_{0, c_2}(\xi'_2) \tilde{\psi}(\xi'_1) \tilde{\psi}(\xi'_2) \cdot \varphi_0(\xi'_3) \dots \varphi_0(\xi'_n)| d\xi'_1 \dots d\xi'_n dt,$$

for any multi-indices \tilde{M} , where $\xi' := (\xi'_1, \dots, \xi'_n)$.

Since m_β is of order $\beta > 0$ and its derivatives decay away from 0, we can majorize, for (ξ'_1, \dots, ξ'_n) with $|\xi'_1| \sim 1, |\xi'_2| \sim 1, |\xi'_3| \lesssim 1, \dots, |\xi'_n| \lesssim 1$,

$$(6.6) \quad |\partial_{\xi'_1}^{\tilde{M}} (\nabla_2 m_\beta(2^{k_1} \xi'_1, t2^{k_2} \xi'_2, t2^{k_1} \xi'_3, \dots, t2^{k_1} \xi'_n) \cdot \xi'_2) | \lesssim 2^{k_1(\beta-1)}.$$

By applying (6.6) to (6.5), we conclude that for any $M > 0$,

$$(6.7) \quad |C_{L_1, \dots, L_n}^{k_1, k_2, c_1, c_2}| \lesssim_M \frac{2^{k_1(\beta-1)} 2^{k_2}}{(1 + |L_1| + \dots + |L_n|)^M}.$$

The initial estimate (6.2) becomes the sum of terms of the form:

$$I_A := \sum_{c_1, c_2 \in \mathfrak{C}} \sum_{k_1 \gg k_2} C_{L_1, \dots, L_n}^{k_1, k_2, c_1, c_2} \Delta_{k_1, c_1, \frac{L_1}{2^{k_1}}} f_1(x) \Delta_{k_2, c_2, \frac{L_2}{2^{k_1}}} f_2(x) \cdot S_{k_1, \frac{L_3}{2^{k_1}}} f_3(x) \cdot \dots \cdot S_{k_1, \frac{L_n}{2^{k_1}}} f_n(x)$$

and

$$(6.8) \quad I_B := \sum_{k_1 \gg k_2, \dots, k_n} (T_{m_\beta(\cdot, 0, \dots, 0)} \Delta_{k_1} f_1)(x) \cdot (\Delta_{k_2} f_2)(x) \cdot \dots \cdot (\Delta_{k_n} f_n)(x).$$

Above, the operator Δ_{k_1, c_1} is defined in frequency by

$$(6.9) \quad \widehat{\Delta_{k_1, c_1} f_1}(\xi_1) := \tilde{\psi}_{k_1, c_1}(\xi_1) \tilde{\psi}_{k_1}(\xi_1) \widehat{\Delta_{k_1} f_1}(\xi_1),$$

so that for any $1 \leq p_1 \leq \infty$, we still⁴⁹ have

$$(6.10) \quad \|\Delta_{k_1, c_1, a} f_1\|_{p_1} = \|\Delta_{k_1, c_1} f_1\|_{p_1} \lesssim \|\Delta_{k_1} f_1\|_{p_1}.$$

For I_A , we get the usual estimates analogous to the ones in Section 5.1:

$$\|I_A\|_r^\tau \lesssim \sum_{k_1 \gg k_2} 2^{k_1(\beta-1)\tau} 2^{k_2\tau} \|\Delta_{k_1} f_1\|_{p_1}^\tau \|\Delta_{k_2} f_2\|_{p_2}^\tau \cdot \|f_3\|_{p_3}^\tau \cdot \dots \cdot \|f_n\|_{p_n}^\tau.$$

For I_B , we switch the order of summation to rewrite it as

$$(T_{m_\beta(\cdot, 0, \dots, 0)} f_1)(x) \cdot f_2(x) \cdot \dots \cdot f_n(x) - \sum_{k_1 < k_2} (T_{m_\beta(\cdot, 0, \dots, 0)} \Delta_{k_1} f_1)(x) (\Delta_{k_2} f_2)(x) f_3(x) \cdot \dots \cdot f_n(x) + \text{many similar terms.}$$

The first term is the reason why we cannot obtain endpoint estimates $p_1 = 1$ or $p_1 = \infty$,⁵⁰ but its boundedness reduces to Hölder's inequality and to the fact that

$$m_\beta(\xi_1, 0, \dots, 0) \cdot |\xi_1|^{-\beta}$$

⁴⁹ $\Delta_{k_1, c_1, a}$ represents the a -modulation of Δ_{k_1, c_1} .

⁵⁰In other regions where for example $|\xi_{j_0}| \gg |\xi_1|, \dots, |\xi_n|$, we will miss the endpoints $p_{j_0} = 1$ or $p_{j_0} = \infty$; so overall we simply have the conditions $1 < p_1, \dots, p_n < \infty$.

is a Mihlin symbol of order 0; hence

$$\|T_{m_\beta(\cdot, 0, \dots, 0)} f_1\|_{p_1} \lesssim \|D^\beta f_1\|_{p_1}$$

for any $1 < p_1 < \infty$. The remaining terms reduce to familiar estimates of the form

$$\sum_{k_1 < k_j + n + 10} 2^{k_1 \beta \tau} \|\Delta_{k_1} f_1\|_{p_1}^\tau \|\Delta_{k_j} f_j\|_{p_j}^\tau,$$

which can easily be bounded – see the treatment of I_A in Section 2.2.

- (2) In the diagonal cone $\{(\xi_1, \dots, \xi_n) : |\xi_1| \sim |\xi_l| \geq |\xi_{\tilde{l}}| \text{ for } \tilde{l} \neq 1, l\}$, we assume without loss of generality that $l = 2$ and we would like to estimate

$$(6.11) \quad \sum_{k_1 \sim k_2} \int_{\mathbb{R}^{nd}} m_\beta(\xi_1, \dots, \xi_n) \widehat{\Delta_{k_1} f_1}(\xi_1) \widehat{\Delta_{k_2} f_2}(\xi_2) \widehat{\Delta_{k_3} f_3}(\xi_3) \cdots \widehat{\Delta_{k_n} f_n}(\xi_n) e^{2\pi i x \cdot (\xi_1 + \dots + \xi_n)} d\xi_1 \dots d\xi_n.$$

We apply the Whitney decomposition

$$(6.12) \quad \bigcup_{k_1 \in \mathbb{Z}} \bigcup_{c_1, c_2 \in \mathfrak{C}} (\{|\xi_1| \sim 2^{k_1}\} \cap c_1) \times (\{|\xi_2| \sim 2^{k_1}\} \cap c_2) \times \{|\xi_3| \leq 2^{k_1}\} \times \dots \times \{|\xi_n| \leq 2^{k_1}\}$$

and then perform a Fourier series decomposition of $m_\beta(\xi_1, \dots, \xi_n)$ restricted to each Whitney cube with fixed $k_1 \in \mathbb{Z}$ and $c_1, c_2 \in \mathfrak{C}$. We thus obtain

$$m_\beta(\xi_1, \dots, \xi_n) \tilde{\psi}_{k_1, c_1}(\xi_1) \tilde{\psi}_{k_1, c_2}(\xi_2) \tilde{\varphi}_{k_1}(\xi_3) \cdots \tilde{\varphi}_{k_1}(\xi_n) = \sum_{L_1, \dots, L_n \in \mathbb{Z}^d} C_{L_1, \dots, L_n}^{k_1, c_1, c_2} e^{2\pi i L_1 \cdot \frac{\xi_1}{2^{k_1}}} e^{2\pi i L_2 \cdot \frac{\xi_2}{2^{k_1}}} e^{2\pi i L_3 \cdot \frac{\xi_3}{2^{k_1}}} \dots e^{2\pi i L_n \cdot \frac{\xi_n}{2^{k_1}}}.$$

Since this Whitney cube is away from the origin, we have again for any $M > 0$,

$$|C_{L_1, \dots, L_n}^{k_1}| \lesssim_M \frac{2^{k_1 \beta}}{(1 + |L_1| + \dots + |L_n|)^M}.$$

Hence estimating (6.11) in $\|\cdot\|_r^\tau$ reduces to summing

$$\sum_{k_1} 2^{k_1 \beta \tau} \|\Delta_{k_1} f_1\|_{p_1}^\tau \|\Delta_{k_1} f_2\|_{p_2}^\tau \|f_3\|_{p_3}^\tau \cdots \|f_n\|_{p_n}^\tau,$$

which is a routine computation by now. □

Remark 6.2. For depth-1 trees, we obtain “Kato-Ponce”-type estimates:

$$(6.13) \quad \|T_{m_\beta}(f, g) - (T_{m_\beta(\cdot, 0)} f) \cdot g - f \cdot (T_{m_\beta(0, \cdot)} g)\|_r \lesssim \|D^\beta f\|_{p_1} \|g\|_{p_2} + \|f\|_{p_1} \|D^\beta g\|_{p_2}$$

for any $1 \leq p_1, p_2 \leq \infty$, $1/r = 1/p_1 + 1/p_2$. In this case we can allow for L^1 and L^∞ endpoints. Of course, the right hand side of (6.13) can be replaced by a geometric mean of appropriate Besov norms.

The next interesting situation corresponds to multilinear multi-parameter operators associated to generic Marcinkiewicz symbols of positive orders; that is, we consider trees of complexity 1 for multi-parameter non-tensorized symbols satisfying (1.35). In order to avoid over-burdening the notation, we simply assume that only two parameters and two functions are involved: $N = 2$, $n = 2$.

Sketch of Proof of Theorem 1.5. We start with $m_{\beta_1, \beta_2}(\xi, \eta)$ a symbol in $\mathbb{R}^{2d_1} \times \mathbb{R}^{2d_2}$ that is smooth away from the planes $\{(\xi_1, \xi_2) = 0\}$ and $\{(\eta_1, \eta_2) = 0\}$, and satisfies the Marcinkiewicz condition

$$|\partial_\xi^{\zeta_1} \partial_\eta^{\zeta_2} m_{\beta_1, \beta_2}(\xi, \eta)| \lesssim |\xi|^{\beta_1 - |\zeta_1|} |\eta|^{\beta_2 - |\zeta_2|}$$

for sufficiently many multi-indices ζ_1, ζ_2 , where $\xi = (\xi_1, \xi_2) = ((\xi_1^i)_{i=1}^{d_1}, (\xi_2^i)_{i=1}^{d_1})$ and $\eta = (\eta_1, \eta_2) = ((\eta_1^i)_{i=1}^{d_2}, (\eta_2^i)_{i=1}^{d_2})$.

The associated bilinear operator is given by

$$(6.14) \quad T_{m_{\beta_1, \beta_2}}(f_1, f_2)(x) := \int_{\mathbb{R}^{2d_1} \times \mathbb{R}^{2d_2}} m_{\beta_1, \beta_2}(\xi, \eta) \widehat{f_1}(\xi_1, \eta_1) \widehat{f_2}(\xi_2, \eta_2) e^{2\pi i x \cdot (\xi_1 + \xi_2)} e^{2\pi i y \cdot (\eta_1 + \eta_2)} d\xi d\eta,$$

and our aim is to prove that $T_{m_{\beta_1, \beta_2}}$ satisfies the same estimate described in (1.10) for the Lebesgue exponents $1 < p_1^j, p_2^j < \infty$, $\frac{1}{p_1^j} + \frac{1}{p_2^j} = \frac{1}{r^j}$, $0 < r^j < \infty$, for all $1 \leq j \leq 2$.

As in the one-parameter setting, we first decompose the frequency spaces for both parameters into cones as in (6.1). Depending on the type of cones, the arguments will be different and we will develop a case-by-case study as before.

- (1) In the case when the cones for both parameters are of type (1) as in (6.1), the root symbol split will be two-folded, involving commutators in each parameter.

Assume without loss of generality that the symbol is smoothly restricted to the region

$$R := \{(\xi_1, \xi_2, \eta_1, \eta_2) : |\xi_1| \gg |\xi_2|, |\eta_1| \gg |\eta_2|\}.$$

Let $\tilde{\chi}_R$ denote the smooth restriction to the region R . We split the symbol on this region as

$$\begin{aligned} & m_{\beta_1, \beta_2}(\xi_1, \xi_2, \eta_1, \eta_2) \\ &= (m_{\beta_1, \beta_2}(\xi_1, \xi_2, \eta_1, \eta_2) - m_{\beta_1, \beta_2}(\xi_1, 0, \eta_1, \eta_2)) + (m_{\beta_1, \beta_2}(\xi_1, 0, \eta_1, \eta_2) - m_{\beta_1, \beta_2}(\xi_1, 0, \eta_1, 0)) + m_{\beta_1, \beta_2}(\xi_1, 0, \eta_1, 0) \\ &= \int_0^1 \partial_{\xi_2} m_{\beta_1, \beta_2}(\xi_1, t\xi_2, \eta_1, \eta_2) \cdot \xi_2 dt + \int_0^1 \partial_{\eta_2} m_{\beta_1, \beta_2}(\xi_1, 0, \eta_1, s\eta_2) \cdot \eta_2 ds + m_{\beta_1, \beta_2}(\xi_1, 0, \eta_1, 0) \\ &= \int_0^1 (\partial_{\xi_2} m_{\beta_1, \beta_2}(\xi_1, t\xi_2, \eta_1, \eta_2) - \partial_{\xi_2} m_{\beta_1, \beta_2}(\xi_1, t\xi_2, \eta_1, 0)) \cdot \xi_2 dt + \int_0^1 \partial_{\xi_2} m_{\beta_1, \beta_2}(\xi_1, t\xi_2, \eta_1, 0) \cdot \xi_2 dt + \\ & \quad \int_0^1 \partial_{\eta_2} m_{\beta_1, \beta_2}(\xi_1, 0, \eta_1, s\eta_2) \cdot \eta_2 ds + m_{\beta_1, \beta_2}(\xi_1, 0, \eta_1, 0) \\ &= \int_0^1 \int_0^1 \sum_{i=1}^{d_1} \sum_{i'=1}^{d_2} \xi_2^i \eta_2^{i'} \partial_{\xi_2^i} \partial_{\eta_2^{i'}} m_{\beta_1, \beta_2}(\xi_1, t\xi_2, \eta_1, s\eta_2) dt ds + \int_0^1 \partial_{\xi_2} m_{\beta_1, \beta_2}(\xi_1, t\xi_2, \eta_1, 0) \cdot \xi_2 dt + \\ & \quad \int_0^1 \partial_{\eta_2} m_{\beta_1, \beta_2}(\xi_1, 0, \eta_1, s\eta_2) \cdot \eta_2 ds + m_{\beta_1, \beta_2}(\xi_1, 0, \eta_1, 0) \\ &:= \mathcal{I} + \mathcal{II} + \mathcal{III} + \mathcal{IV}. \end{aligned}$$

In the case when the symbol tensorizes, \mathcal{I} corresponds to the tensor product of two commutator symbols, \mathcal{II} and \mathcal{III} are generalizations of the mix of a commutator symbol and a symbol of lower complexity whereas \mathcal{IV} is a biparameter variant of I_B defined in (6.8).

It suffices to prove the boundedness of the multipliers $T_{\mathcal{I}\tilde{\chi}_R}$, $T_{\mathcal{II}\tilde{\chi}_R}$, $T_{\mathcal{III}\tilde{\chi}_R}$ and $T_{\mathcal{IV}\tilde{\chi}_R}$.

Estimate for $T_{\mathcal{I}\tilde{\chi}_R}$. We recall the Whitney decomposition used in the one parameter setting and decompose frequency spaces for both parameters as

$$\begin{aligned} \{(\xi_1, \xi_2) \in \mathbb{R}^{2d_1} : |\xi_1| \gg |\xi_2|\} &= \bigcup_{c_1, c_2 \in \mathfrak{C}_1} \bigcup_{k_1 \gg k_2} (\{|\xi_1| \sim 2^{k_1}\} \cap c_1) \times (\{|\xi_2| \sim 2^{k_2}\} \cap c_2), \\ \{(\eta_1, \eta_2) \in \mathbb{R}^{2d_2} : |\eta_1| \gg |\eta_2|\} &= \bigcup_{c'_1, c'_2 \in \mathfrak{C}_2} \bigcup_{m_1 \gg m_2} (\{|\eta_1| \sim 2^{m_1}\} \cap c'_1) \times (\{|\eta_2| \sim 2^{m_2}\} \cap c'_2), \end{aligned}$$

where \mathfrak{C}_1 and \mathfrak{C}_2 represent the collections of directional cones in the frequency spaces \mathbb{R}^{d_1} and \mathbb{R}^{d_2} respectively. We then smoothly restrict the symbol

$$m_{C_{\beta_1}, C_{\beta_2}}(\xi_1, \xi_2, \eta_1, \eta_2) := \int_0^1 \int_0^1 \sum_{i=1}^{d_1} \sum_{i'=1}^{d_2} \xi_2^i \eta_2^{i'} \partial_{\xi_2^i} \partial_{\eta_2^{i'}} m_{\beta_1, \beta_2}(\xi_1, t\xi_2, \eta_1, s\eta_2) dt ds$$

to each Whitney rectangle with fixed $k_1 \gg k_2, m_1 \gg m_2, c_1, c_2 \in \mathfrak{C}_1$ and $c'_1, c'_2 \in \mathfrak{C}_2$, and denote it by $m_{C_{\beta_1}, C_{\beta_2}}^{k_1, k_2, m_1, m_2, c_1, c_2, c'_1, c'_2}$. We perform the quadruple Fourier series decomposition to obtain

(6.15)

$$m_{C_{\beta_1}, C_{\beta_2}}^{k_1, k_2, m_1, m_2, c_1, c_2, c'_1, c'_2}(\xi_1, \xi_2, \eta_1, \eta_2) = \sum_{\substack{L_1, L_2 \in \mathbb{Z}^{d_1} \\ L'_1, L'_2 \in \mathbb{Z}^{d_2}}} C_{L_1, L_2, L'_1, L'_2}^{k_1, k_2, m_1, m_2, c_1, c_2, c'_1, c'_2} e^{2\pi i L_1 \cdot \frac{\xi_1}{2^{k_1}}} e^{2\pi i L_2 \cdot \frac{\xi_2}{2^{k_2}}} e^{2\pi i L'_1 \cdot \frac{\eta_1}{2^{m_1}}} e^{2\pi i L'_2 \cdot \frac{\eta_2}{2^{m_2}}},$$

where the Fourier coefficients decay rapidly due to the Marcinkiewicz condition (6.14) on m_{β_1, β_2} :

$$|C_{L_1, L_2, L'_1, L'_2}^{k_1, k_2, m_1, m_2, c_1, c_2, c'_1, c'_2}| \lesssim 2^{k_1(\beta_1-1)} 2^{k_2} 2^{m_1(\beta_2-1)} 2^{m_2} (1 + |L_1| + |L_2|)^{-N} (1 + |L'_1| + |L'_2|)^{-N'}$$

for sufficiently large N and N' .

By applying the Fourier series representation (6.15), we rewrite the multiplier as

$$\begin{aligned} T_{\mathcal{I}\tilde{\chi}_R}(f_1, f_2)(x, y) &= \sum_{\substack{c_1, c_2 \in \mathfrak{C}_1 \\ c'_1, c'_2 \in \mathfrak{C}_2}} \sum_{\substack{L_1, L_2 \in \mathbb{Z}^{d_1} \\ L'_1, L'_2 \in \mathbb{Z}^{d_2}}} \sum_{\substack{k_1 \gg k_2 \\ m_1 \gg m_2}} C_{L_1, L_2, L'_1, L'_2}^{k_1, k_2, m_1, m_2, c_1, c_2, c'_1, c'_2} \\ & \int_{\mathbb{R}^{2d_1} \times \mathbb{R}^{2d_2}} \mathcal{F}(\Delta_{k_1, c_1}^{(1)} \Delta_{m_1, c'_1}^{(2)} f_1)(\xi_1, \eta_1) \mathcal{F}(\Delta_{k_2, c_2}^{(1)} \Delta_{m_2, c'_2}^{(2)} f_2)(\xi_2, \eta_2) e^{2\pi i \xi_1 \cdot (x + \frac{L_1}{2^{k_1}})} e^{2\pi i \xi_2 \cdot (x + \frac{L_2}{2^{k_2}})} e^{2\pi i \eta_1 \cdot (y + \frac{L'_1}{2^{m_1}})} e^{2\pi i \eta_2 \cdot (y + \frac{L'_2}{2^{m_2}})} d\xi d\eta. \end{aligned}$$

We can invoke now the analysis developed in Sections 4 and 5 to conclude the discussion.

Estimate for $T_{\mathcal{I}\mathcal{I}\tilde{\chi}_R}$ and $T_{\mathcal{I}\mathcal{I}\mathcal{I}\tilde{\chi}_R}$. We shall notice that the treatment of the multipliers corresponding to $\mathcal{I}\mathcal{I}$ and $\mathcal{I}\mathcal{I}\mathcal{I}$ are symmetric and for that reason we will focus on $T_{\mathcal{I}\mathcal{I}\tilde{\chi}_R}$. We rewrite the symbol as

$$\mathcal{I}\mathcal{I} = m_{C_{\beta_1}, H}(\xi_1, \xi_2, \eta_1) |\eta_1|^{\beta_2},$$

where

$$m_{C_{\beta_1}, H}(\xi_1, \xi_2, \eta_1) := \int_0^1 \frac{\partial_{\xi_2} m(\xi_1, t\xi_2, \eta_1, 0) \cdot \xi_2}{|\eta_1|^{\beta_2}} dt.$$

As the notation suggests, $m_{C_{\beta_1}, H}(\xi_1, \xi_2, \eta_1)$ is a symbol generating a commutator in the first parameter and a Mihlin symbol of order 0 (the simplest example being the symbol corresponding to the Hilbert transform) in the second parameter localized on the region $\{(\xi_1, \xi_2, \eta_1) : |\xi_1| \gg |\xi_2|\}$. In order to use the Fourier series decomposition on this symbol, we need to decompose

$$\{(\xi_1, \xi_2, \eta_1) : |\xi_1| \gg |\xi_2|\} = \bigcup_{c_1, c_2 \in \mathfrak{C}_1} \bigcup_{c'_1 \in \mathfrak{C}_2} \bigcup_{\substack{k_1 \gg k_2 \\ m_1 \in \mathbb{Z}}} |(\{|\xi_1| \sim 2^{k_1}\} \cap c_1) \times (\{|\xi_2| \sim 2^{k_2}\} \cap c_2) \times (\{|\eta_1| \sim 2^{m_1}\} \cap c'_1)|.$$

We smoothly restrict $m_{C_{\beta_1}, H}(\xi_1, \xi_2, \eta_1)$ to a region with fixed $k_1 \gg k_2, m_1 \in \mathbb{Z}$, $c_1, c_2 \in \mathfrak{C}_1$ and $c'_1 \in \mathfrak{C}_2$, and let $m_{C_{\beta_1}, H}^{k_1, k_2, m_1, c_1, c_2, c'_1}(\xi_1, \xi_2, \eta_1)$ denote the localized symbol. We then perform the triple Fourier series decomposition for

$$(6.16) \quad m_{C_{\beta_1}, H}^{k_1, k_2, m_1, c_1, c_2, c'_1}(\xi_1, \xi_2, \eta_1) = \sum_{\substack{L_1, L_2 \in \mathbb{Z}^{d_1} \\ L' \in \mathbb{Z}^{d_2}}} C_{L_1, L_2, L'}^{k_1, k_2, m_1, c_1, c_2, c'_1} e^{2\pi i L_1 \cdot \frac{\xi_1}{2^{k_1}}} e^{2\pi i L_2 \cdot \frac{\xi_2}{2^{k_2}}} e^{2\pi i L' \cdot \frac{\eta_1}{2^{m_1}}}.$$

Because of the regularity of $m_{C_{\beta_1}, H}(\xi_1, \xi_2, \eta_1)$, the Fourier coefficients satisfy the decay condition

$$(6.17) \quad |C_{L_1, L_2, L'}^{k_1, k_2, m_1, c_1, c_2, c'_1}| \lesssim 2^{k_1(\beta_1-1)} 2^{k_2} (1 + |L_1| + |L_2|)^{-N} (1 + |L'|)^{-N'}$$

for sufficiently large N and N' . Thanks to (6.16), we can rewrite the multiplier as

$$\begin{aligned} T_{\mathcal{I}\mathcal{I}\tilde{\chi}_R} &= \sum_{\substack{c_1, c_2 \in \mathfrak{C}_1 \\ c'_1 \in \mathfrak{C}_2}} \sum_{\substack{L_1, L_2 \in \mathbb{Z}^{d_1} \\ L' \in \mathbb{Z}^{d_2}}} \sum_{\substack{k_1 \gg k_2 \\ m_1 \gg m_2}} C_{L_1, L_2, L'}^{k_1, k_2, m_1, c_1, c_2, c'_1} \int |\eta_1|^{\beta_2} \mathcal{F}(\Delta_{k_1, c_1}^{(1)} \Delta_{m_1, c'_1}^{(2)} f_1)(\xi_1, \eta_1) \mathcal{F}(\Delta_{k_2, c_2}^{(1)} \Delta_{m_2}^{(2)} f_2)(\xi_2, \eta_2) e^{2\pi i \xi_1 \cdot (x + \frac{L_1}{2^{k_1}})} \\ &\quad e^{2\pi i \xi_2 \cdot (x + \frac{L_2}{2^{k_2}})} e^{2\pi i \eta_1 \cdot (y + \frac{L'}{2^{m_1}})} e^{2\pi i \eta_2 \cdot y} d\xi d\eta \\ &= \sum_{\substack{c_1, c_2 \in \mathfrak{C}_1 \\ c'_1 \in \mathfrak{C}_2}} \sum_{\substack{L_1, L_2 \in \mathbb{Z}^{d_1} \\ L' \in \mathbb{Z}^{d_2}}} \sum_{\substack{k_1 \gg k_2 \\ m_1, m_2}} C_{L_1, L_2, L'}^{k_1, k_2, m_1, c_1, c_2, c'_1} \int \mathcal{F}(\Delta_{k_1, c_1}^{(1)} \Delta_{m_1, c'_1}^{(2)} D_{(2)}^{\beta_2} f_1)(\xi_1, \eta_1) \mathcal{F}(\Delta_{k_2, c_2}^{(1)} \Delta_{m_2}^{(2)} f_2)(\xi_2, \eta_2) e^{2\pi i \xi_1 \cdot (x + \frac{L_1}{2^{k_1}})} \\ &\quad e^{2\pi i \xi_2 \cdot (x + \frac{L_2}{2^{k_2}})} e^{2\pi i \eta_1 \cdot (y + \frac{L'}{2^{m_1}})} e^{2\pi i \eta_2 \cdot y} d\xi d\eta \\ &\quad - \sum_{\substack{c_1, c_2 \in \mathfrak{C}_1 \\ c'_1 \in \mathfrak{C}_2}} \sum_{\substack{L_1, L_2 \in \mathbb{Z}^{d_1} \\ L' \in \mathbb{Z}^{d_2}}} \sum_{\substack{k_1 \gg k_2 \\ m_1 < m_2}} C_{L_1, L_2, L'}^{k_1, k_2, m_1, c_1, c_2, c'_1} \int |\eta_1|^{\beta_2} \mathcal{F}(\Delta_{k_1, c_1}^{(1)} \Delta_{m_1, c'_1}^{(2)} f_1)(\xi_1, \eta_1) \mathcal{F}(\Delta_{k_2, c_2}^{(1)} \Delta_{m_2}^{(2)} f_2)(\xi_2, \eta_2) e^{2\pi i \xi_1 \cdot (x + \frac{L_1}{2^{k_1}})} \\ &\quad e^{2\pi i \xi_2 \cdot (x + \frac{L_2}{2^{k_2}})} e^{2\pi i \eta_1 \cdot (y + \frac{L'}{2^{m_1}})} e^{2\pi i \eta_2 \cdot y} d\xi d\eta := \mathcal{A} - \mathcal{B}. \end{aligned}$$

It is not difficult to verify that \mathcal{B} can be treated using the argument presented in Section 4. In contrast, the term \mathcal{A} is trickier to estimate due to the fact that the non-tensorized symbol $m_{C_{\beta_1}, H}(\xi_1, \xi_2, \eta_1)$ exhibits different types of behaviours in the (ξ_1, ξ_2) and η_1 variables.

Nonetheless, we can simplify \mathcal{A} as follows

$$\begin{aligned} &\sum_{\substack{c_1, c_2 \in \mathfrak{C}_1 \\ c'_1 \in \mathfrak{C}_2}} \sum_{\substack{L_1, L_2 \in \mathbb{Z}^{d_1} \\ L' \in \mathbb{Z}^{d_2}}} \sum_{\substack{k_1 \gg k_2 \\ m_1}} \sum_{\substack{L_1, L_2, L'}} C_{L_1, L_2, L'}^{k_1, k_2, m_1, c_1, c_2, c'_1} \Delta_{k_1, c_1}^{(1)} \Delta_{m_1, c'_1}^{(2)} D_{(2)}^{\beta_2} f_1(x + \frac{L_1}{2^{k_1}}, y + \frac{L'}{2^{m_1}}) \sum_{m_2} \Delta_{k_2, c_2}^{(1)} \Delta_{m_2}^{(2)} f_2(x + \frac{L_2}{2^{k_2}}, y) \\ &= \sum_{\substack{c_1, c_2 \in \mathfrak{C}_1 \\ c'_1 \in \mathfrak{C}_2}} \sum_{\substack{L_1, L_2 \in \mathbb{Z}^{d_1} \\ L' \in \mathbb{Z}^{d_2}}} \sum_{\substack{k_1 \gg k_2 \\ m_1}} \sum_{\substack{L_1, L_2, L'}} C_{L_1, L_2, L'}^{k_1, k_2, m_1, c_1, c_2, c'_1} \Delta_{k_1, c_1}^{(1)} \Delta_{m_1, c'_1}^{(2)} D_{(2)}^{\beta_2} f_1(x + \frac{L_1}{2^{k_1}}, y + \frac{L'}{2^{m_1}}) \Delta_{k_2, c_2}^{(1)} f_2(x + \frac{L_2}{2^{k_2}}, y). \end{aligned}$$

Its $\|\cdot\|_{L_x^1(L_y^2)}^\tau$ norm with $\tau \leq \min(1, r^1, r^2)$ can be majorized by

$$(6.18) \quad \sum_{\substack{c_1, c_2 \in \mathfrak{C}_1 \\ c'_1 \in \mathfrak{C}_2}} \sum_{\substack{L_1, L_2 \in \mathbb{Z}^{d_1} \\ L' \in \mathbb{Z}^{d_2}}} \sum_{\substack{k_1 \gg k_2 \\ m_1}} \left\| \sum_{L_1, L_2, L'} C_{L_1, L_2, L'}^{k_1, k_2, m_1, c_1, c_2, c'_1} \Delta_{k_1, c_1}^{(1)} \Delta_{m_1, c'_1}^{(2)} D_{(2)}^{\beta_2} f_1(\cdot, \cdot + \frac{L'}{2^{m_1}}) \right\|_{L_x^1(L_y^2)}^\tau \left\| \Delta_{k_2, c_2}^{(1)} f_2 \right\|_{L_x^{\frac{1}{p_2}}(L_y^{\frac{1}{p_2}})}^\tau.$$

Let $F := \Delta_{k_1, c_1}^{(1)} D_{(2)}^{\beta_2} f_1$. We claim that, for any $1 < p_1^1, p_1^2 < \infty$,

$$(6.19) \quad \left\| \sum_{m_1} C_{L_1, L_2, L'}^{k_1, k_2, m_1, c_1, c_2, c'_1} \Delta_{m_1, c'_1}^{(2)} F(\cdot, \cdot + \frac{L'}{2m_1}) \right\|_{L_x^{p_1^1}(L_y^{p_1^2})} \lesssim (1 + |L_1| + |L_2|)^{-N} (1 + |L'|)^{-N'+1} 2^{k_1(\beta_1-1)} 2^{k_2} \|F\|_{L_x^{p_1^1}(L_y^{p_1^2})}.$$

Applying (6.19) to (6.18), we are left with a familiar expression that can be easily dealt with.

To prove the claim, we first linearize the left hand side of (6.19) by choosing an appropriate function $h \in L_x^{p_1^{1'}}(L_y^{p_1^{2'}})$ with $\|h\|_{L_x^{p_1^{1'}}(L_y^{p_1^{2'}})} = 1$ such that

$$\begin{aligned} & \left\| \sum_{m_1} C_{L_1, L_2, L'}^{k_1, k_2, m_1, c_1, c_2, c'_1} \Delta_{m_1, c'_1}^{(2)} F(\cdot, \cdot + \frac{L'}{2m_1}) \right\|_{L_x^{p_1^1}(L_y^{p_1^2})} = \left\| \sum_{m_1} C_{L_1, L_2, L'}^{k_1, k_2, m_1, c_1, c_2, c'_1} \Delta_{m_1, c'_1}^{(2)} F * \tilde{\psi}_{m_1}(\cdot, \cdot + \frac{L'}{2m_1}) \right\|_{L_x^{p_1^1}(L_y^{p_1^2})} \\ &= \left| \int \sum_{m_1} C_{L_1, L_2, L'}^{k_1, k_2, m_1, c_1, c_2, c'_1} \Delta_{m_1, c'_1}^{(2)} F(x, y + \frac{L'}{2m_1}) \tilde{\Delta}_{m_1}^{(2)} h(x, y) dx dy \right| \\ &\leq \sup_{m_1} |C_{L_1, L_2, L'}^{k_1, k_2, m_1, c_1, c_2, c'_1}| \int \left(\sum_{m_1} |\Delta_{m_1, c'_1}^{(2)} F(x, y + \frac{L'}{2m_1})|^2 \right)^{\frac{1}{2}} \left(\sum_{m_1} |\tilde{\Delta}_{m_1}^{(2)} h(x, y)|^2 \right)^{\frac{1}{2}} dx dy \\ (6.20) \quad &\lesssim \frac{2^{k_1(\beta_1-1)} 2^{k_2}}{(1 + |L_1| + |L_2|)^N (1 + |L'|)^{N'}} \int \left(\int \left(\sum_{m_1} |\Delta_{m_1, c'_1}^{(2)} F(x, y + \frac{L'}{2m_1})|^2 \right)^{\frac{p_1^2}{2}} dy \right)^{\frac{1}{p_1^1}} \left(\int \left(\sum_{m_1} |\tilde{\Delta}_{m_1}^{(2)} h(x, y)|^2 \right)^{\frac{p_1^{2'}}{2}} dy \right)^{\frac{1}{p_1^{1'}}} dx, \end{aligned}$$

where the third inequality follows from Cauchy-Schwartz and the last one holds due to Hölder. We recall the boundedness⁵¹ of the shifted square function

$$(6.21) \quad \left(\int \left(\sum_{m_1} |\Delta_{m_1, c'_1}^{(2)} F(x, y + \frac{L'}{2m_1})|^2 \right)^{\frac{p_1^2}{2}} dy \right)^{\frac{1}{p_1^1}} \lesssim O(1 + |L'|^{100}) \left(\int |F_{c'_1}(x, y)|^{p_1^2} dy \right)^{\frac{1}{p_1^1}},$$

where $F_{c'_1} := \mathcal{F}^{-1}(\hat{F}(\xi_1, \eta_1) \sum_{m_1 \in \mathbb{Z}} \psi_{m_1, c_1}(\eta_1))$ and ψ_{m_1, c_1} is defined in Remark 2.4. We also have that

$$(6.22) \quad \left(\int |F_{c'_1}(x, y)|^{p_1^2} dy \right)^{\frac{1}{p_1^1}} \lesssim \left(\int |F(x, y)|^{p_1^2} dy \right)^{\frac{1}{p_1^1}},$$

for $1 < p_1^2 < \infty$. This ends the proof of the claim.

Estimate for $T_{\mathcal{V}\tilde{\chi}_R}$.

$$m_{\beta_1, \beta_2}(\xi_1, 0, \eta_1, 0) \tilde{\chi}_R = \sum_{\substack{k_1 \gg k_2 \\ m_1 \gg m_2}} m_{\beta_1, \beta_2}(\xi_1, 0, \eta_1, 0) \tilde{\chi}_{R_{k_1, k_2, m_1, m_2}},$$

where $\tilde{\chi}_{R_{k_1, k_2, m_1, m_2}}$ denote the smooth restriction to the region R_{k_1, k_2, m_1, m_2} achieved by the Littlewood-Paley decomposition. Now we apply the high-low switch technique as previously to deduce

$$\begin{aligned} & \sum_{\substack{k_1 \gg k_2 \\ m_1 \gg m_2}} m_{\beta_1, \beta_2}(\xi_1, 0, \eta_1, 0) \tilde{\chi}_{R_{k_1, k_2, m_1, m_2}} = \sum_{\substack{k_1, k_2 \\ m_1, m_2}} m_{\beta_1, \beta_2}(\xi_1, 0, \eta_1, 0) \tilde{\chi}_{R_{k_1, k_2, m_1, m_2}} - \sum_{\substack{k_1 < k_2 \\ m_1, m_2}} m_{\beta_1, \beta_2}(\xi_1, 0, \eta_1, 0) \tilde{\chi}_{R_{k_1, k_2, m_1, m_2}} \\ & - \sum_{\substack{k_1, k_2 \\ m_1 < m_2}} m_{\beta_1, \beta_2}(\xi_1, 0, \eta_1, 0) \tilde{\chi}_{R_{k_1, k_2, m_1, m_2}} + \sum_{\substack{k_1 < k_2 \\ m_1 < m_2}} m_{\beta_1, \beta_2}(\xi_1, 0, \eta_1, 0) \tilde{\chi}_{R_{k_1, k_2, m_1, m_2}} := \mathcal{IV}_a - \mathcal{IV}_b - \mathcal{IV}_c + \mathcal{IV}_d. \end{aligned}$$

It is straightforward to verify that \mathcal{IV}_a generates the symbol for the (linear) bi-parameter Marcinkiewicz multiplier of order (β_1, β_2) , whose boundedness is well-known. \mathcal{IV}_d can be estimated using the routine procedures developed in Section 4; \mathcal{IV}_b and \mathcal{IV}_c are symmetric and follow the similar analysis as \mathcal{II} .

(2) When the cone for the first parameter is of type (1) as in (6.1) and the cone for the second parameter is of type (2),⁵² the argument is a hybrid of the reasoning in cases (1) and (2) developed in the one-parameter setting. In particular, the symbol is smoothly restricted to the region

$$\tilde{R} := \{(\xi_1, \xi_2, \eta_1, \eta_2) : |\xi_1| \gg |\xi_2|, |\eta_1| \sim |\eta_2|\},$$

where we split it as follows:

$$\begin{aligned} m_{\beta_1, \beta_2}(\xi_1, \xi_2, \eta_1, \eta_2) &= m_{\beta_1, \beta_2}(\xi_1, \xi_2, \eta_1, \eta_2) - m_{\beta_1, \beta_2}(\xi_1, 0, \eta_1, \eta_2) + m_{\beta_1, \beta_2}(\xi_1, 0, \eta_1, \eta_2) \\ &= \int_0^1 \partial_{\xi_2} m_{\beta_1, \beta_2}(\xi_1, t\xi_2, \eta_1, \eta_2) \cdot \xi_2 dt + m_{\beta_1, \beta_2}(\xi_1, 0, \eta_1, \eta_2). \end{aligned}$$

⁵¹The bounds provided here are far from being optimal, but they are sufficient for our purpose.

⁵²The other case is similar by symmetry.

The study of the multiplier associated to the symbol

$$m_{C_{\beta_1, \beta_2}}(\xi_1, \xi_2, \eta_1, \eta_2) := \int_0^1 \partial_{\xi_2} m_{\beta_1, \beta_2}(\xi_1, t\xi_2, \eta_1, \eta_2) \cdot \xi_2 dt$$

requests a Fourier series decomposition. With a by now routine decomposition of the frequency space \tilde{R} , we have

$$(6.23) \quad \tilde{R} = \bigcup_{\substack{c_1, c_2 \in \mathfrak{C}_1 \\ c'_1, c'_2 \in \mathfrak{C}_2}} \bigcup_{\substack{k_1 \gg k_2 \\ m_1 \in \mathbb{Z}}} (\{|\xi_1| \sim 2^{k_1}\} \cap c_1) \times (\{|\xi_2| \sim 2^{k_2}\} \cap c_2) \times (\{|\eta_1| \sim 2^{m_1}\} \cap c'_1) \times (\{|\eta_2| \sim 2^{m_1}\} \cap c'_2),$$

where we recall that \mathfrak{C}_1 and \mathfrak{C}_2 are collections of directional cones in the frequency spaces \mathbb{R}^{d_1} and \mathbb{R}^{d_2} , respectively.

We smoothly restrict the symbol to each piece of (6.23) with fixed $k_1 \gg k_2$, $m_1 \in \mathbb{Z}$, $c_1, c_2 \in \mathfrak{C}_1$ and $c'_1, c'_2 \in \mathfrak{C}_2$ and denote it by $m_{C_{\beta_1, \beta_2}}^{k_1, k_2, m_1, c_1, c_2, c'_1, c'_2}$. By applying the Fourier series decomposition of the symbol to the corresponding multiplier, we obtain an expression whose analysis follows the standard procedure of this paper.

The symbol $m_{\beta_1, \beta_2}(\xi_1, 0, \eta_1, \eta_2)$ can be rewritten as

$$(6.24) \quad \frac{m_{\beta_1, \beta_2}(\xi_1, 0, \eta_1, \eta_2)}{|\xi_1|^{\beta_1}} |\xi_1|^{\beta_1} =: m_{H, \beta_2}(\xi_1, \eta_1, \eta_2) |\xi_1|^{\beta_1}.$$

Last but not least, we notice that the symbol $m_{H, \beta_2}(\xi_1, \eta_1, \eta_2)$, as implied by the notation, is of order 0 for the first parameter and order β_2 for the second parameter; because of that, the multiplier corresponding to (6.24) can be treated similarly to \mathcal{II} . \square

6.2. Some remarks on smoothing properties of multipliers. Finally, we present some results in the vein of [10] and [19]. Although our methods do not apply to symbols satisfying only Sobolev conditions,⁵³ we do recover the results from [10] and from [19] for fractional integral operators also in the N -parameter, mixed-norm setting. Once more, the purpose of this section is only illustrative.

Theorem 6.3. *Let $0 \leq \nu_1, \dots, \nu_N < nd$ and let $m \in \mathcal{M}_{-\nu_1, \dots, -\nu_N}(\mathbb{R}^{d_1} \times \dots \times \mathbb{R}^{d_N})$ be a Mihlin symbol of order $(-\nu_1, \dots, -\nu_N)$. Then for any $s_1, \dots, s_N \geq 0$ so that $s_j \neq \nu_j$ for any $1 \leq j \leq N$ and for any functions $f_1, \dots, f_n \in \mathcal{S}(\mathbb{R}^{d_1} \times \dots \times \mathbb{R}^{d_N})$, we have*

$$(6.25) \quad \|D_{(1)}^{s_1} \dots D_{(N)}^{s_N} T_m(f_1, \dots, f_n)\|_{L^r} \lesssim \sum_{\substack{\sigma_1^1, \dots, \sigma_n^N \in \{0, 1\} \\ \sigma_1^j + \dots + \sigma_n^j = 1}} \prod_{l=1}^n \|D_{(1)}^{\sigma_l^1(s_1 - \nu_1)} \dots D_{(N)}^{\sigma_l^N(s_N - \nu_N)} f_l\|_{\tilde{p}_l},$$

provided that $1 < p_1^1, \dots, p_1^N, p_2^1, \dots, p_2^N, \dots, p_n^1, \dots, p_n^N < \infty$, $\frac{1}{n} < r^1, \dots, r^n < \infty$ are satisfying component-wise the Hölder condition

$$\frac{1}{\tilde{r}} = \frac{1}{\tilde{p}_1} + \dots + \frac{1}{\tilde{p}_n}$$

and

$$\frac{d_N}{d_N + s_N} < r^N, \quad \max\left(\frac{d_N}{d_N + s_N}, \frac{d_{N-1}}{d_{N-1} + s_{N-1}}\right) < r^{N-1}, \dots, \max\left(\frac{d_N}{d_N + s_N}, \dots, \frac{d_1}{d_1 + s_1}\right) < r^1.$$

⁵³In the one-parameter bilinear setting, instead of the Sobolev conditions in [10], we require that m is continuously differentiable away from the origin and that it satisfies

$$\begin{aligned} \sup_{c_1, c_2 \in \mathfrak{C}} \sup_k \|m_{k, c_1, c_2}^\nu\|_{W(\tilde{r}, \tilde{r}), 2} < \infty, \quad \sup_{c_1, c_2 \in \mathfrak{C}} \sup_{k_1 \ll k_2} \sup_{t \in [0, 1]} \|m_{k_1, k_2, c_1, c_2}^{\nu, t}\|_{W(\tilde{r}, \tilde{r}), 2} < \infty, \quad \text{and} \\ \sup_{c_1, c_2 \in \mathfrak{C}} \sup_{k_1 \ll k_2} \sup_{t \in [0, 1]} \|m_{k_1, k_2, c_1, c_2}^{\nu, t, \nabla_i}\|_{W(\tilde{r}, \tilde{r}), 2} < \infty \quad \text{for } i = 1, 2, \end{aligned}$$

where \mathfrak{C} is a collection of directional cones in the frequency space \mathbb{R}^d and

$$\begin{aligned} m_{k, c_1, c_2}^\nu &:= 2^{k\nu} m(2^k \xi_1, 2^k \xi_2) \psi_{0, c_1}(\xi_1) \psi_{0, c_2}(\xi_2), \\ m_{k_1, k_2, c_1, c_2}^{\nu, t} &:= 2^{k_1 \nu} m(2^{k_1} \xi_1, t 2^{k_2} \xi_2) \psi_{0, c_1}(\xi_1) \psi_{0, c_2}(\xi_2), \\ m_{k_1, k_2, c_1, c_2}^{\nu, t, \nabla_2} &:= 2^{k_1(\nu+1)} (\nabla_2 m(2^{k_1} \xi_1, t 2^{k_2} \xi_2) \cdot \xi_2) \psi_{0, c_1}(\xi_1) \psi_{0, c_2}(\xi_2), \end{aligned}$$

and $m_{k_1, k_2, c_1, c_2}^{\nu, t, \nabla_1}$ is defined similarly. The discrete Sobolev norm $W(\tilde{r}, \tilde{r}), 2$ for a function f is defined by

$$\|f\|_{W(\tilde{r}, \tilde{r}), 2} := \left(\sum_{n_1, n_2 \in \mathbb{Z}^d} (1 + |n_1|^2)^{\tilde{r}} (1 + |n_2|^2)^{\tilde{r}} |\hat{f}(n_1, n_2)|^2 \right)^{\frac{1}{2}}.$$

Using the argument in the proof of Theorem 6.3, (6.26) can be verified for symbols m satisfying the conditions above for $0 \leq \nu < 2d$, $\max(\frac{d}{\tilde{r}}, \frac{d}{d+s}) < r < \infty$, $1 < p_1, p_2 < \infty$ and $\frac{1}{r} = \frac{1}{p_1} + \frac{1}{p_2}$. A similar Sobolev condition can be formulated in the biparameter setting.

We call attention to the fact that (6.25) remains true whenever $s_1 - \nu_1 \neq 0, \dots, s_N - \nu_N \neq 0$, so even when they are negative numbers.

A typical example of multipliers satisfying the boundedness property in Theorem 6.3 is the N -parameter fractional integral operator $I_\nu(f_1, \dots, f_n)$ given by

$$\int_{\mathbb{R}^{nd_1} \times \dots \times \mathbb{R}^{nd_N}} (|\xi_1^1|^2 + \dots + |\xi_n^1|^2)^{-\frac{\nu_1}{2}} \dots (|\xi_1^N|^2 + \dots + |\xi_n^N|^2)^{-\frac{\nu_N}{2}} \hat{f}_1(\xi_1) \dots \hat{f}_n(\xi_n) e^{2\pi i x \cdot (\xi_1 + \dots + \xi_n)} d\xi^1 \dots d\xi^N,$$

where $\xi^j := (\xi_1^j, \dots, \xi_n^j) \in \mathbb{R}^n$ for $1 \leq j \leq N$. In this particular case we can of course allow for $s_j = \nu_j$ and (6.25) holds for $D_{(1)}^{s_1} \dots D_{(N)}^{s_N} I_\nu(f_1, \dots, f_n)$. The smoothing property for fractional integral operators, naturally implied by Theorem 6.3, seem to be new in the multi-parameter, mixed-norm setting.

Again, for simplicity we only present the case $n = 2$ and $N = 2$ of the proof. First, we start with the one-parameter case, which relies on the results in Theorem 1.3 corresponding to a rooted tree of depth 1; next we discuss the bi-parameter case, which makes use of Theorem 1.5.

Sketch of Proof of Theorem 6.3 when $N = 1$. Here, like in [10], we would like to obtain smoothing properties for operators of order ν , where $0 \leq \nu < 2d$. This simply means that we take $m(\xi, \eta) \in \mathcal{M}_{-\nu}(\mathbb{R}^{2d})$ and would like to prove for any $s \neq \nu$

$$(6.26) \quad \|D^s T_m(f, g)\|_r \lesssim \|D^{s-\nu} f\|_{p_1} \|g\|_{p_2} + \|f\|_{p_1} \|D^{s-\nu} g\|_{p_2}.$$

As usual, $D^s T_m(f, g)$ breaks down as a sum of

$$\begin{aligned} & \sum_{k \ll \ell} \int_{\mathbb{R}^{2d}} |\xi_1 + \xi_2|^s m_\nu(\xi_1, \xi_2) \widehat{\Delta_k f}(\xi_1) \widehat{\Delta_\ell g}(\xi_2) e^{2\pi i x \cdot (\xi_1 + \xi_2)} d\xi_1 d\xi_2 \\ & + \sum_{k \gg \ell} \int_{\mathbb{R}^{2d}} |\xi_1 + \xi_2|^s m_\nu(\xi_1, \xi_2) \widehat{\Delta_k f}(\xi_1) \widehat{\Delta_\ell g}(\xi_2) e^{2\pi i x \cdot (\xi_1 + \xi_2)} d\xi_1 d\xi_2 \\ & + \sum_{|k-\ell| \leq 3} \int_{\mathbb{R}^{2d}} |\xi_1 + \xi_2|^s m_\nu(\xi_1, \xi_2) \widehat{\Delta_k f}(\xi_1) \widehat{\Delta_\ell g}(\xi_2) e^{2\pi i x \cdot (\xi_1 + \xi_2)} d\xi_1 d\xi_2 := I + II + III \end{aligned}$$

and it will be enough to estimate the terms I and III .

(1) For I , we simply notice that on the region

$$R := \bigcup_{k \ll \ell} \{(\xi_1, \xi_2) : |\xi_1| \sim 2^k, |\xi_2| \sim 2^\ell\},$$

the symbol $|\xi_1 + \xi_2|^s m_\nu(\xi_1, \xi_2)$ becomes a Mihlin symbol of order $s - \nu$. So as long as $s > \nu$, the result follows from Theorem 1.3. When $s < \nu$, the proof of Theorem 1.3 still holds: in that case, we would use $\|\cdot\|_{\dot{B}_{p_l, \infty}^{s-\nu}}$ and

$\|\cdot\|_{\dot{B}_{p_l, \infty}^{s-\nu-\epsilon}}$ Besov norms, with $s - \nu < -\epsilon < 0$.

(2) For the diagonal term, we want to estimate

$$III := \sum_k \int_{\mathbb{R}^{2d}} |\xi_1 + \xi_2|^s m_\nu(\xi_1, \xi_2) \widehat{\Delta_k f}(\xi_1) \widehat{\Delta_k g}(\xi_2) e^{2\pi i x \cdot (\xi_1 + \xi_2)} d\xi_1 d\xi_2.$$

For this, we would like to use a Fourier series decomposition for $|\xi_1 + \xi_2|^s m_\nu(\xi_1, \xi_2)$ in the region $\{(\xi_1, \xi_2) : |\xi_1| \sim |\xi_2|\}$, which can be decomposed as (6.12):

$$\{(\xi_1, \xi_2) \in \mathbb{R}^{2d} : |\xi_1| \sim |\xi_2|\} = \bigcup_{c_1, c_2 \in \mathfrak{C}} \bigcup_{k \in \mathbb{Z}} (\{|\xi_1| \sim 2^k\} \cap c_1) \times (\{|\xi_2| \sim 2^k\} \cap c_2),$$

where \mathfrak{C} is the collection of all directional cones in the frequency space \mathbb{R}^d .

We proceed with a Fourier series decomposition of the symbol restricted to the Whitney cube corresponding to a fixed scale $k \in \mathbb{Z}$ and fixed directional cones $c_1, c_2 \in \mathfrak{C}$, namely

$$(6.27) \quad |\xi_1 + \xi_2|^s m_\nu(\xi_1, \xi_2) \psi_{k,c}(\xi_1) \psi_{k,c}(\xi_2).$$

In this region, while $m_\nu(\xi_1, \xi_2)$ is smooth, $|\xi_1 + \xi_2|^s$ is less regular since we cannot exclude that $|\xi_1 + \xi_2| = 0$. So we proceed with Fourier series decompositions on both symbols separately. We first notice that (6.27) can be rewritten as

$$(|\xi_1 + \xi_2|^s \tilde{\varphi}_k(\xi_1 + \xi_2)) (m_\nu(\xi_1, \xi_2) \psi_{k,c}(\xi_1) \psi_{k,c}(\xi_2)).$$

Then the Fourier series of the symbol in the first parenthesis yields

$$(6.28) \quad |\xi_1 + \xi_2|^s \tilde{\varphi}_k(\xi_1 + \xi_2) = \sum_{L \in \mathbb{Z}^d} 2^{ks} C_L e^{2\pi i \frac{L}{2^k} \cdot (\xi_1 + \xi_2)},$$

where the renormalized Fourier coefficients C_L satisfy

$$(6.29) \quad |C_L| \lesssim \frac{1}{(1 + |L|)^{d+s}}.$$

Next, we use a Fourier decomposition on $m_\nu(\xi_1, \xi_2)\psi_{k,c_1}(\xi_1)\psi_{k,c_2}(\xi_2)$:

$$m_\nu(\xi_1, \xi_2)\psi_{k,c_1}(\xi_1)\psi_{k,c_2}(\xi_2) = \sum_{L_1, L_2 \in \mathbb{Z}^d} C_{L_1, L_2}^{k, c_1, c_2} e^{2\pi i \frac{L_1}{2^k} \cdot \xi_1} e^{2\pi i \frac{L_2}{2^k} \cdot \xi_2},$$

with

$$|C_{L_1, L_2}^{k, c_1, c_2}| \lesssim_M \frac{2^{-k\nu}}{(1 + |L_1| + |L_2|)^M}.$$

for $M > 0$ sufficiently large. Overall we get

$$|\xi_1 + \xi_2|^s m_\nu(\xi_1, \xi_2)\psi_{k,c_1}(\xi_1)\psi_{k,c_2}(\xi_2) = \sum_{L \in \mathbb{Z}^d} \sum_{L_1, L_2 \in \mathbb{Z}^d} 2^{ks} C_L C_{L_1, L_2}^{k, c_1, c_2} e^{2\pi i \frac{L+L_1}{2^k} \cdot \xi_1} e^{2\pi i \frac{L+L_2}{2^k} \cdot \xi_2}$$

and in consequence, whenever $0 < \tau \leq \min(1, r)$,

$$\|III\|_r^\tau \lesssim \sum_L |C_L|^\tau \sum_k 2^{k(s-\nu)\tau} \|\Delta_k f\|_{p_1}^\tau \|\Delta_k g\|_{p_2}^\tau.$$

This imposes the restriction $r > \frac{d}{d+s}$, and implies the desired (6.25) for either $s > \nu$ or $s < \nu$. \square

Sketch of Proof of Theorem 6.3 when $N = 2$. Now we prove the bi-parameter version of the smoothing property discussed above; that is, we will show that

$$(6.30) \quad \begin{aligned} \|D_{(1)}^{s_1} D_{(2)}^{s_2} T_m(f_1, f_2)\|_{L_x^{r_1}(L_y^{r_2})} &\lesssim \|D_{(1)}^{s_1-\nu_1} D_{(2)}^{s_2-\nu_2} f_1\|_{L_x^{p_1^1}(L_y^{p_1^2})} \|f_2\|_{L_x^{p_2^1}(L_y^{p_2^2})} + \|f_1\|_{L_x^{p_1^1}(L_y^{p_1^2})} \|D_{(1)}^{s_1-\nu_1} D_{(2)}^{s_2-\nu_2} f_2\|_{L_x^{p_2^1}(L_y^{p_2^2})} \\ &+ \|D_{(1)}^{s_1-\nu_1} f_1\|_{L_x^{p_1^1}(L_y^{p_1^2})} \|D_{(2)}^{s_2-\nu_2} f_2\|_{L_x^{p_2^1}(L_y^{p_2^2})} + \|D_{(2)}^{s_2-\nu_2} f_1\|_{L_x^{p_1^1}(L_y^{p_1^2})} \|D_{(1)}^{s_1-\nu_1} f_2\|_{L_x^{p_2^1}(L_y^{p_2^2})}, \end{aligned}$$

for $1 < p_1^j, p_2^j < \infty$, $\frac{d}{d+s_j} < r^j$ and $\frac{1}{p_1^j} + \frac{1}{p_2^j} = \frac{1}{r^j}$ with $j = 1, 2$.

As indicated by the argument in the one-parameter setting, various considerations are necessary for different regions of the frequency space.

(1) In the “off-diagonal” region

$$\{(\xi_1, \xi_2, \eta_1, \eta_2) : |\xi_1| \gg |\xi_2|, |\eta_1| \gg |\eta_2|\}$$

for both parameters (or other similar regions, obtained by permuting the roles of ξ_1, ξ_2 or η_1, η_2), the symbols $|\xi_1 + \xi_2|^{s_1}$ and $|\eta_1 + \eta_2|^{s_2}$ are Mihlin symbols of order s_1 and s_2 respectively. Hence

$$m_{s_1-\nu_1, s_2-\nu_2}(\xi_1, \xi_2, \eta_1, \eta_2) := |\xi_1 + \xi_2|^{s_1} |\eta_1 + \eta_2|^{s_2} m(\xi_1, \xi_2, \eta_1, \eta_2)$$

is a bi-parameter Marcinkiewicz symbol satisfying

$$|\partial_\xi^{\zeta_1} \partial_\eta^{\zeta_2} m_{\beta_1, \beta_2}(\xi, \eta)| \lesssim |\xi|^{s_1-\nu_1-|\zeta_1|} |\eta|^{s_2-\nu_2-|\zeta_2|}.$$

We can now invoke Theorem 1.5 to conclude the discussion for this case.

(2) In the diagonal region for both parameters

$$\{(\xi_1, \xi_2, \eta_1, \eta_2) : |\xi_1| \sim |\xi_2|, |\eta_1| \sim |\eta_2|\},$$

we no longer need to split the symbol and consider commutator terms. Instead, we perform a quadruple Fourier series decomposition of $m(\xi_1, \xi_2, \eta_1, \eta_2)$ restricted to a Whitney rectangle and a Fourier series decompositions of the symbols $|\xi_1 + \xi_2|^{s_1}$ and $|\eta_1 + \eta_2|^{s_2}$ respectively, restricted accordingly. The subsequent analysis mimics the argument developed in Section 4.

(3) In the region that is “off-diagonal” for the first parameter and “diagonal” for the second parameter (and in the other similar regions, alike)

$$\{(\xi_1, \xi_2, \eta_1, \eta_2) : |\xi_1| \gg |\xi_2|, |\eta_1| \sim |\eta_2|\},$$

we will use a hybrid of the arguments for (1) and (2). In particular,

$$\begin{aligned} &|\xi_1 + \xi_2|^{s_1} m(\xi_1, \xi_2, \eta_1, \eta_2) \\ &= \underbrace{|\xi_1 + \xi_2|^{s_1} m(\xi_1, \xi_2, \eta_1, \eta_2) - |\xi_1|^{s_1} m(\xi_1, 0, \eta_1, \eta_2)}_{=: m_{C_{s_1-\nu_1, -\nu_2}}(\xi_1, \xi_2, \eta_1, \eta_2)} + \underbrace{|\xi_1|^{s_1} m(\xi_1, 0, \eta_1, \eta_2)}_{=: m_{s_1-\nu_1, -\nu_2}(\xi_1, \eta_1, \eta_2)}. \end{aligned}$$

The quadruple Fourier series decomposition can be applied to the symbol $m_{C_{s_1-\nu_1}, -\nu_2}$ smoothly restricted to the region

$$(6.31) \quad (\{|\xi_1| \sim 2^{k_1}\} \cap c_1) \times (\{|\xi_2| \sim 2^{k_2}\} \cap c_2) \times (\{|\eta_1| \sim 2^{m_1}\} \cap c'_1) \times (\{|\eta_1| \sim 2^{m_1}\} \cap c'_2),$$

where $k_1 \gg k_2$ and $c_1, c_2 \in \mathfrak{C}_1$ are directional cones in the frequency space for the first parameter and $c'_1, c'_2 \in \mathfrak{C}_2$ are directional cones in the frequency space for the second parameter.

Meanwhile, we can use the Fourier series decomposition to express the symbol $|\eta_1 + \eta_2|^{s_2}$ localized to the cube $[-2^{m_1+1}, 2^{m_1+1}]^{d_2}$ which yields an expression similar to (6.28). The corresponding Fourier coefficients satisfy a decaying condition similar to (6.29), which imposes the constraint $r^2 > \frac{d_2}{d_2 + s_2}$ on the Lebesgue exponent r^2 . Next, we use the Fourier series representation for the symbol $m_{C_{s_1-\nu_1}, -\nu_2} |\eta_1 + \eta_2|^{s_2}$ in the multiplier; the remaining argument is routine and will be omitted here.

Last but not least, the symbol $m_{s_1-\nu_1, -\nu_2}(\xi_1, \eta_1, \eta_2)$ can be rewritten as

$$m_{s_1-\nu_1, -\nu_2}(\xi_1, \eta_1, \eta_2) = \frac{m_{s_1-\nu_1, -\nu_2}(\xi_1, \eta_1, \eta_2)}{|\xi_1|^{s_1-\nu_1}} |\xi_1|^{s_1-\nu_1} =: m_{H, -\nu_2}(\xi_1, \eta_1, \eta_2) |\xi_1|^{s_1-\nu_1}.$$

We smoothly restrict the symbol $m_{H, -\nu_2}(\xi_1, \eta_1, \eta_2)$ to the region

$$(\{|\xi_1| \sim 2^{k_1}\} \cap c_1) \times (\{|\eta_1| \sim 2^{m_1}\} \cap c'_1) \times (\{|\eta_2| \sim 2^{m_1}\} \cap c'_2),$$

where $k_1, m_1 \in \mathbb{Z}$ are fixed, $c_1 \in \mathfrak{C}_1$ is a directional cone in the frequency space for the first parameter and $c'_1, c'_2 \in \mathfrak{C}_2$ are directional cones in the frequency space for the second parameter. We denote the localized symbol by $m_{H, -\nu_2}^{k_1, m_1}$, on which we perform the triple Fourier series decomposition:

$$m_{H, -\nu_2}^{k_1, m_1, c_1, c'_1, c'_2}(\xi_1, \eta_1, \eta_2) = \sum_{\substack{L \in \mathbb{Z}^{d_1} \\ L'_1, L'_2 \in \mathbb{Z}^{d_2}}} C_{L, L'_1, L'_2}^{k_1, m_1, c_1, c'_1, c'_2} e^{2\pi i L \cdot \frac{\xi_1}{2^{k_1}}} e^{2\pi i L'_1 \cdot \frac{\eta_1}{2^{m_1}}} e^{2\pi i L'_2 \cdot \frac{\eta_2}{2^{m_1}}},$$

where the Fourier coefficients satisfy the decay condition

$$|C_{L, L'_1, L'_2}^{k_1, m_1, c_1, c'_1, c'_2}| \lesssim 2^{-m_1 \nu_2} (1 + |L|)^{-N} (1 + |L'_1| + |L'_2|)^{-N'}$$

for sufficiently large N, N' . When combined with the Fourier series representation for $|\eta_1 + \eta_2|^{s_2}$ on $\{|\eta_1 + \eta_2| \leq 2^{m_1+1}\}$, we obtain for fixed $c_1 \in \mathfrak{C}_1$ and $c'_1, c'_2 \in \mathfrak{C}_2$,

$$\sum_{\substack{L \in \mathbb{Z}^{d_1} \\ L'_1, L'_2 \in \mathbb{Z}^{d_2}}} \sum_{\substack{k_1 \gg k_2 \\ m_1}} C_{L, L'_1, L'_2}^{k_1, m_1, c_1, c'_1, c'_2} C_{L'} 2^{m_1 s_1} \int |\xi_1|^{s_1-\nu_1} \mathcal{F}(\Delta_{k_1, c_1}^{(1)} \Delta_{m_1, c'_1}^{(2)} f_1)(\xi_1, \eta_1) \mathcal{F}(\Delta_{k_2}^{(1)} \Delta_{m_1, c'_2}^{(2)} f_2)(\xi_2, \eta_2) e^{2\pi i \xi_1 \cdot (x + \frac{L}{2^{k_1}})} e^{2\pi i \xi_2 \cdot x} e^{2\pi i \eta_1 \cdot (y + \frac{L'_1 + L'}{2^{m_1}})} e^{2\pi i \eta_2 \cdot (y + \frac{L'_2 + L'}{2^{m_1}})} d\xi d\eta.$$

We then use the high-low switch technique to split the sum in two parts – one with the sum over all k_1, k_2, m_1 and the second sum over $k_2 > k_1$ and all m_1 . The second sum can be estimated using the usual analysis (optimization, Besov norms, etc), while the first can be simplified as

$$\sum_{\substack{L \in \mathbb{Z}^{d_1} \\ L'_1, L'_2, L' \in \mathbb{Z}^{d_2}}} \sum_{k_1, m_1 \in \mathbb{Z}} C_{L, L'_1, L'_2}^{k_1, m_1, c_1, c'_1, c'_2} C_{L'} 2^{m_1 s_1} \Delta_{k_1, c_1}^{(1)} \Delta_{m_1, c'_1}^{(2)} D_{(1)}^{s_1-\nu_1} f_1(x + \frac{L}{2^{k_1}}, y + \frac{L'_1 + L'}{2^{m_1}}) \Delta_{m_1, c'_2}^{(2)} f_2(x, y + \frac{L'_2 + L'}{2^{m_1}}).$$

By subadditivity and Hölder's inequality, its $\|\cdot\|_{L_x^{\tau^1} L_y^{\tau^2}}^{\tau}$ norm with $\tau \leq \min(1, r^1, r^2)$ can be estimated by

$$\sum_{\substack{L \in \mathbb{Z}^{d_1} \\ L'_1, L'_2, L' \in \mathbb{Z}^{d_2}}} \sum_{m_1} |C_{L'}|^{\tau} 2^{m_1 s_1 \tau} \left\| \sum_{k_1} C_{L, L'_1, L'_2}^{k_1, m_1, c_1, c'_1, c'_2} \Delta_{k_1, c_1}^{(1)} \Delta_{m_1, c'_1}^{(2)} D_{(1)}^{s_1-\nu_1} f_1(\cdot + \frac{L}{2^{k_1}}, \cdot) \right\|_{L_x^{\tau^1} L_y^{\tau^2}}^{\tau} \left\| \Delta_{m_1, c'_2}^{(2)} f_2(\cdot, \cdot) \right\|_{L_x^{\tau^1} L_y^{\tau^2}}^{\tau}.$$

The reasoning used for proving the inequality (6.19) also implies that

$$\left\| \sum_{k_1} C_{L, L'_1, L'_2}^{k_1, m_1, c_1, c'_1, c'_2} \Delta_{k_1, c_1}^{(1)} \Delta_{m_1, c'_1}^{(2)} D_{(1)}^{s_1-\nu_1} f_1(\cdot + \frac{L}{2^{k_1}}, \cdot) \right\|_{L_x^{\tau^1} L_y^{\tau^2}}^{\tau} \lesssim (1 + |L|)^{-N+1} (1 + |L'_1| + |L'_2|)^{-N'} 2^{-m_1 \nu_2} \left\| \Delta_{m_1}^{(2)} D_{(1)}^{s_1-\nu_1} f_1 \right\|_{L_x^{\tau^1} L_y^{\tau^2}}.$$

From here on the summation in m_1 is standard.

The reasoning presented above yields the desired estimates in the case when $s_1 > \nu_1, s_2 > \nu_2$. As in the one-parameter case, a similar and even simpler⁵⁴ argument implies (6.30) for any $s_1 \neq \nu_1, s_2 \neq \nu_2$. \square

REFERENCES

- [1] Cristina Benea and Camil Muscalu. Multiple vector-valued inequalities via the helicoidal method. *Anal. PDE*, 9(8):1931–1988, 2016. [3](#)
- [2] Cristina Benea and Camil Muscalu. Quasi-Banach valued inequalities via the helicoidal method. *J. Funct. Anal.*, 273(4):1295–1353, 2017. [3](#)
- [3] Jean-Michel Bony. Calcul symbolique et propagation des singularités pour les équations aux dérivées partielles non linéaires. *Ann. Sci. École Norm. Sup. (4)*, 14(2):209–246, 1981. [10](#)
- [4] Jean Bourgain and Dong Li. On an endpoint Kato-Ponce inequality. *Differential Integral Equations*, 27(11-12):1037–1072, 2014. [2, 3, 11, 12, 13, 16, 67](#)
- [5] R. Coifman and Y. Meyer. *Wavelets, Calderón-Zygmund Operators and Multilinear Operators*. Cambridge University Press, 1997. [2](#)
- [6] Francesco Di Plinio and Yumeng Ou. Banach-valued multilinear singular integrals. *Indiana Univ. Math. J.*, 67(5):1711–1763, 2018. [3](#)
- [7] Loukas Grafakos, Diego Maldonado, and Virginia Naibo. A remark on an endpoint Kato-Ponce inequality. *Differential Integral Equations*, 27(5-6):415–424, 2014. [2](#)
- [8] Loukas Grafakos and Seungly Oh. The Kato-Ponce inequality. *Communications in Partial Differential Equations*, pages 1128–1157, 2014. [7, 19, 22](#)
- [9] Lu Zhang Guozhen Lu, Jill Pipher. Bi-parameter trilinear Fourier multipliers and pseudo-differential operators with flag symbols. <http://arxiv.org/abs/1901.00036>, 2019. Online; accessed June 2021. [3](#)
- [10] Jarod Hart, Rodolfo H. Torres, and Xinfeng Wu. Smoothing properties of bilinear operators and Leibniz-type rules in Lebesgue and mixed Lebesgue spaces. *Trans. Amer. Math. Soc.*, 370(12):8581–8612, 2018. [10, 67, 74, 75](#)
- [11] Tosio Kato and Gustavo Ponce. Commutator estimates and the Euler and Navier-Stokes equations. *Comm. Pure Appl. Math.*, 41(7):891–907, 1988. [22](#)
- [12] Carlos Kenig, Gustavo Ponce, and Luis Vega. Well-posedness and scattering results for the generalized Korteweg-de Vries equation via the contraction principle. *Comm. Pure Appl. Math.*, pages 527–620, 1993. [22](#)
- [13] Akihiko Miyachi and Naohito Tomita. Estimates for trilinear flag paraproducts on L^∞ and Hardy spaces. *Math. Z.*, 282(1-2):577–613, 2016. [3](#)
- [14] Camil Muscalu. Paraproducts with flag singularities. I. A case study. *Rev. Mat. Iberoam.*, 23(2):705–742, 2007. [1, 3, 6, 31, 67](#)
- [15] Camil Muscalu, Jill Pipher, Terence Tao, and Christoph Thiele. Bi-parameter paraproducts. *Acta Mathematica*, pages 269–296, 2004. [2, 67](#)
- [16] Camil Muscalu and Wilhelm Schlag. *Classical and multilinear harmonic analysis. Vol. II*, volume 138 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, 2013. [22](#)
- [17] Camil Muscalu and Yujia Zhai. Five-linear singular integral estimates of Brascamp-Lieb type. <http://arxiv.org/abs/2001.09064>, 2020. Online; accessed June 2021. [3](#)
- [18] Seungly Oh and Xinfeng Wu. On L^1 endpoint Kato – Ponce inequality. *Math. Res. Lett.*, 27(4):1129–1163, 2020. [2, 3, 7, 25, 33, 34, 57](#)
- [19] Jiexing Yang, Zongguang Liu, and Xinfeng Wu. Leibniz-type rules for bilinear and biparameter Fourier multiplier operators with applications. *Potential Analysis*, 370, 2020. [10, 67, 74](#)

(C. Benea) UNIVERSITÉ DE NANTES, LABORATOIRE DE MATHÉMATIQUES JEAN LERAY, NANTES 44322, FRANCE
 Email address: cristina.benea@univ-nantes.fr

(Y. Zhai) CNRS - UNIVERSITÉ DE NANTES, LABORATOIRE DE MATHÉMATIQUES JEAN LERAY, NANTES 44322, FRANCE
 Email address: yujia.zhai@univ-nantes.fr

⁵⁴The high-low switch procedure is no longer necessary if $s_2 < \nu_2$, when one needs to sum

$$\sum_{m_1} 2^{m_1(s_2-\nu_2)\tau} \|\Delta_{m_1}^{(2)} D_{(1)}^{s_1-\nu_1} f_1\|_{L_{x^1}^{p_1^1}(L_{y^1}^{p_1^2})}^\tau \|\Delta_{m_1}^{(2)} f_2\|_{L_{x^2}^{p_2^1}(L_{y^2}^{p_2^2})}^\tau.$$