

TOEPLITZ OPERATORS ON THE FOCK SPACE WITH QUASI-RADIAL SYMBOLS

VISHWA DEWAGE & GESTUR ÓLAFSSON

ABSTRACT. The Fock space $\mathcal{F}(\mathbb{C}^n)$ is the space of holomorphic functions on \mathbb{C}^n that are square-integrable with respect to the Gaussian measure on \mathbb{C}^n . This space plays an important role in several subfields of analysis and representation theory. In particular, it has for a long time been a model to study Toeplitz operators. Esmeral and Maximenko showed in 2016 that radial Toeplitz operators on $\mathcal{F}(\mathbb{C})$ generate a commutative C^* -algebra which is isometrically isomorphic to the C^* -algebra $C_{b,u}(\mathbb{N}_0, \rho_1)$. In this article, we extend the result to k -quasi-radial symbols acting on the Fock space $\mathcal{F}(\mathbb{C}^n)$. We calculate the spectra of the said Toeplitz operators and show that the set of all eigenvalue functions is dense in the C^* -algebra $C_{b,u}(\mathbb{N}_0^k, \rho_k)$ of bounded functions on \mathbb{N}_0^k which are uniformly continuous with respect to the square-root metric. In fact, the C^* -algebra generated by Toeplitz operators with quasi-radial symbols is $C_{b,u}(\mathbb{N}_0^k, \rho_k)$.

CONTENTS

1. Introduction	1
2. Toeplitz operators with bounded k -quasi-radial symbols	3
2.1. A classification of the class of symbols $L^\infty(\mathbb{C}^n)^G$	3
2.2. Diagonalization by Schur's lemma	4
2.3. Computing eigenvalue functions	5
3. The space $C_{b,u}(\mathbb{N}_0^k, \rho_k)$	8
4. Uniform continuity of eigenvalue functions with respect to the square-root metric	10
5. Density of \mathfrak{G}_1 in $C_{b,u}(\mathbb{N}_0^k, \rho_k)$	12
5.1. Approximation by convolutions	12
5.2. The density of \mathfrak{G}_1 in $C_{b,u}(\mathbb{N}_0^k, \rho_k)$	15
6. The C^* -algebra generated by \mathfrak{G}_n	17
7. A comparison of $C_{b,u}(\mathbb{N}_0^k, \rho_k)$ with $C_{b,u}(\mathbb{N}_0, \rho_1) \otimes \cdots \otimes C_{b,u}(\mathbb{N}_0, \rho_1)$	17
7.1. A counter example	17
8. Appendix: The proof of Lemma 3.4	19
8.1. The uniform norm of f	19
8.2. Some useful lemmas	21
8.3. The uniform continuity of f with respect to the square-root metric	23
References	24

1. INTRODUCTION

Reproducing kernel Hilbert spaces \mathcal{H} of square integrable holomorphic functions on complex domains plays an important role in several subfields of analysis and representation theory [B61].

2000 *Mathematics Subject Classification.* 22D25, 30H20, 41A35, 47B35.

Key words and phrases. Toeplitz Operators, Fock Space, Reproducing Kernel Hilbert spaces, Commutative C^* -algebras.

The research of G. Ólafsson was partially supported by Simons grant 586106.

Those spaces lead to a very simple form of quantization, [B75a, B75b, EU10, EU11] turning functions into bounded operators on \mathcal{H} , by using Toeplitz operators $T_\varphi(f) = P(\varphi f)$, where P is the orthogonal projection onto \mathcal{H} . These operators have been studied for a long time, [BC86, BI12, BL11, BS06, C94].

Since the fundamental work [GV02, GQV06], and then later several authors have studied C^* -algebras generated by subclasses of bounded symbols asking the question which of them lead to commutative C^* -algebras as in [DÓQ15, DÓQ18, DÓQ21, DQ18, EM16, GMV13, GKV03, QS11, QV07a, QV07b, QV08, V08]. In the case of commutative C^* -algebras, the next natural question is to determine the spectrum and, if possible, determine the Gelfand transform. This is often achieved by constructing a transform that maps Toeplitz operators to multiplier operators on an L^2 space.

In the special case of the unit ball in \mathbb{C}^n , it was shown that Toeplitz operators with radial symbols acting on the Bergman space generate a commutative C^* -algebra as in [GKV03] and the corresponding C^* -algebra is isometrically isomorphic to the space of bounded sequences that are uniformly continuous with respect to the logarithmic metric [GMV13]. Toeplitz operators with radial symbols on the Fock space $\mathcal{F}(\mathbb{C})$ generate a commutative C^* -algebra [GV02] which is isometrically isomorphic to the C^* -algebra $C_{b,u}(\mathbb{N}_0, \rho_1)$ of bounded sequences that are uniformly continuous with respect to the square-root metric [EM16]. Furthermore, a more general theory for weighted Bergman spaces on any bounded symmetric domain was formulated in [DÓQ15], using representation theory.

In this paper, we consider the C^* -algebra generated by Toeplitz operators with k -quasi-radial symbols, discussed in [BV12, MSR16, V10a, V10b, QS14, QS15], acting on the Fock space $\mathcal{F}(\mathbb{C}^n)$. As an instance of the general theory discussed in [DÓQ15], the generated C^* -algebra is commutative. We compute the eigenvalue functions of Toeplitz operators with quasi-radial symbols and then show that they are dense in the space $C_{b,u}(\mathbb{N}_0^k, \rho_k)$ of bounded functions on \mathbb{N}_0^k that are uniformly continuous with respect to the square-root metric. As a consequence, the C^* -algebra generated by Toeplitz operators with k -quasi-radial symbols is isometrically isomorphic to the C^* -algebra $C_{b,u}(\mathbb{N}_0^k, \rho_k)$.

We now introduce some notations and recall the following well known facts about the Fock space. Denote by $d\lambda_n$ the Gaussian measure $d\lambda_n(z) = \frac{1}{\pi^n} e^{-|z|^2} dz$ for $z \in \mathbb{C}^n$, where dz is the Lebesgue measure on $\mathbb{C}^n \simeq \mathbb{R}^{2n}$. The Fock space $\mathcal{F}(\mathbb{C}^n)$ is the Hilbert space of all holomorphic functions on \mathbb{C}^n that are square integrable with respect to $d\lambda_n$,

$$\mathcal{F}(\mathbb{C}^n) = L^2(\mathbb{C}^n, \lambda_n) \cap \mathcal{O}(\mathbb{C}^n).$$

The point-evaluation maps $f \mapsto f(z)$ are continuous and hence $\mathcal{F}(\mathbb{C}^n)$ is a reproducing kernel Hilbert space with the reproducing kernel K given by $K(w, z) = K_z(w) = e^{w\bar{z}}$, for $(z, w) \in \mathbb{C}^{2n}$ (here $\bar{z}w$ denotes the scalar product $w_1\bar{z}_1 + \dots + w_n\bar{z}_n$). Thus for $z \in \mathbb{C}^n$ and $f \in \mathcal{F}(\mathbb{C}^n)$ we have

$$f(z) = \langle f, K_z \rangle$$

with the inner product of $L^2(\mathbb{C}^n, \lambda_n)$.

The Bergman projection $P : L^2(\mathbb{C}^n) \rightarrow \mathcal{F}(\mathbb{C}^n)$ is given by

$$Pf(z) = \langle f, K_z \rangle.$$

P is a bounded linear operator with $\|P\| = 1$.

Given $\varphi \in L^\infty(\mathbb{C}^n)$, we define the Toeplitz operator $T_\varphi : \mathcal{F}(\mathbb{C}^n) \rightarrow \mathcal{F}(\mathbb{C}^n)$ by

$$T_\varphi f(z) = P(\varphi f)(z) = \int_{\mathbb{C}^n} \varphi(w) f(w) \overline{K_z(w)} d\lambda_n(w).$$

As the multiplier operator $M_\varphi : L^2(\mathbb{C}^n, \lambda) \rightarrow L^2(\mathbb{C}^n, \lambda)$ is bounded of norm $\|\varphi\|_\infty$ and P is bounded, it follows that T_φ is bounded and $\|T_\varphi\| \leq \|\varphi\|_\infty$. The function φ is called the symbol of the Toeplitz operator T_φ .

Grudsky and Vasilevski [GV02] showed that the eigenvalue sequences of radial Toeplitz operators on $\mathcal{F}(\mathbb{C})$ are of the form

$$\gamma_{1,a}(m) = \frac{1}{m!} \int_0^\infty a(\sqrt{r}) r^m e^{-r} dr.$$

We prove that the Toeplitz operators with k -quasi-radial symbols diagonalize with eigenvalue functions given by

$$\gamma_{\mathbf{n},a}(\mathbf{m}) = \frac{1}{(\mathbf{m} + \mathbf{n} - \mathbf{1})!} \int_{\mathbb{R}_+^k} a(\sqrt{r_1}, \dots, \sqrt{r_k}) r^{\mathbf{m} + \mathbf{n} - \mathbf{1}} e^{-(r_1 + \dots + r_k)} dr$$

for any $\mathbf{m} \in \mathbb{N}_0^k$, where $\mathbf{n} = (n_1, \dots, n_k)$.

As a corollary the eigenvalue sequences of radial Toeplitz operators on $\mathcal{F}(\mathbb{C}^n)$ are of the form

$$\gamma_{n,a}(m) = \frac{1}{(m + n - 1)!} \int_0^\infty a(\sqrt{r}) r^{m+n-1} e^{-r} dr$$

which are the $(n - 1)^{\text{th}}$ left shift of $\gamma_{1,a}$.

To prove eigenvalue function of Toeplitz operators with k -quasi-radial symbols is dense in the C^* -algebra $C_{b,u}(\mathbb{N}_0^k, \rho_k)$, we follow some of the ideas in the work by Esmeral and Maximenko [EM16].

Acknowledgements: We sincerely thank the anonymous reviewers for the constructive and positive feedback.

2. TOEPLITZ OPERATORS WITH BOUNDED k -QUASI-RADIAL SYMBOLS

In this section, we introduce quasi-radial symbols and diagonalize the Toeplitz operators with quasi-radial symbols.

We begin with some notations. Let $k \in \mathbb{N}$ and let $\mathbf{n} = \mathbf{n}(k) = (n_1, \dots, n_k) \in \mathbb{N}^k$ and let $n = n_1 + \dots + n_k$. Let $\mathbb{C}^{\mathbf{n}} = \mathbb{C}^{n_1} \times \dots \times \mathbb{C}^{n_k}$. We identify $z \in \mathbb{C}^{\mathbf{n}}$ by $z = (z_{(1)}, \dots, z_{(k)}) \in \mathbb{C}^{\mathbf{n}}$ where $z_{(j)} \in \mathbb{C}^{n_j}$. Consider the group $G = U_{n_1} \times \dots \times U_{n_k}$ where U_{n_i} are $n_i \times n_i$ unitary matrices. They act on f defined on $\mathbb{C}^{\mathbf{n}}$ by $(A_1, \dots, A_k)f(z) = (A_1, \dots, A_k)f(z_{(1)}, \dots, z_{(k)}) := f(A_1^{-1}z_{(1)}, \dots, A_k^{-1}z_{(k)})$ which is the usual action of matrices we get by considering G as a subgroup of U_n . The action leaves the Gaussian measure $d\lambda_n(z) = d\lambda_{n_1}(z_{(1)}) \dots d\lambda_{n_k}(z_{(k)})$ invariant.

Definition 2.1. Let $\varphi : \mathbb{C}^{\mathbf{n}} \rightarrow \mathbb{C}$ be measurable. The function φ is said to be G -invariant if for all $(A_1, \dots, A_k) \in G$ and $z \in \mathbb{C}^{\mathbf{n}}$ we have $(A_1, \dots, A_k)\varphi(z) = \varphi(A_1^{-1}z_{(1)}, \dots, A_k^{-1}z_{(k)}) = \varphi(z)$.

Define the set of all k -quasi-radial symbols, denoted by $L^\infty(\mathbb{C}^{\mathbf{n}})^G$, to be the set of all essentially bounded functions on $\mathbb{C}^{\mathbf{n}}$ that are G -invariant. The case $k = 1$ corresponds to radial symbols and the case $n = k$ corresponds to separately radial symbols. Now we take into consideration the C^* -algebra \mathcal{T}_G generated by Toeplitz operators with symbols in $L^\infty(\mathbb{C}^{\mathbf{n}})^G$. The main objective of this section is to diagonalize the Toeplitz operators with symbols in $L^\infty(\mathbb{C}^{\mathbf{n}})^G$.

2.1. A classification of the class of symbols $L^\infty(\mathbb{C}^{\mathbf{n}})^G$.

Definition 2.2. Let $\varphi : \mathbb{C}^q \rightarrow \mathbb{C}$ be measurable. The function φ is said to be,

- (1) radial if there exists a measurable function $a_\varphi : \mathbb{R}_+ \rightarrow \mathbb{C}$ such that $\varphi(z) = a_\varphi(|z|)$ for $z \neq 0$.
- (2) U_q -invariant if for all $A \in U_q$ and $z \in \mathbb{C}^q$ we have $\varphi(Az) = \varphi(z)$.

Following lemma is a well-known fact.

Lemma 2.3. The measurable function $\varphi : \mathbb{C}^q \rightarrow \mathbb{C}$ is U_q -invariant if and only if there exists $a : \mathbb{R}_+ \rightarrow \mathbb{C}$ such that $\varphi(z) = a(|z|)$.

Proof. As $|Az| = |z|$ for $A \in U_q$, it follows that any function of the form $\varphi(z) = a(|z|)$ is U_q -invariant. As $z \mapsto |z|$ is continuous it follows that φ is measurable if and only if a is measurable.

Assume that φ is U_q -invariant. Define $a_\varphi(r) = \varphi(re_1)$. Then a_φ is measurable. Let $z \in \mathbb{C}^q$. Then, as U_q acts transitively on the sphere S^{2q-1} , there exist $A \in U_q$ such that $Az = |z|e_1$. Thus

$$\varphi(z) = \varphi(Az) = \varphi(|z|e_1) = a_\varphi(|z|).$$

Thus $\varphi(z) = a_\varphi(|z|)$. □

The following lemma is a consequence of Lemma 2.3.

Lemma 2.4. *The measurable function $\varphi : \mathbb{C}^n \rightarrow \mathbb{C}$ is G -invariant if and only if there exists $a : \mathbb{R}_+^k \rightarrow \mathbb{C}$ such that $\varphi(z) = a(|z_{(1)}|, \dots, |z_{(k)}|)$.*

2.2. Diagonalization by Schur's lemma. Let π and σ be unitary representations of a topological group H . A continuous linear map $T : V_\pi \rightarrow V_\sigma$ is an *intertwining operator* if for all $A \in H$ commutating relation $T\pi(A) = \sigma(A)T$ holds. π and σ are equivalent if there exists a unitary isomorphism that intertwines π and σ . Finally if π is irreducible and T is an intertwining operator then Schur's lemma says that $T = \lambda \text{id}$ for some $\lambda \in \mathbb{C}$.

Since the action of G is defined by the action of U_{n_i} on $\mathcal{F}^2(\mathbb{C}^{n_i}), i = 1, \dots, k$, first we consider the action of U_q on $\mathcal{F}^2(\mathbb{C}^q), q \in \mathbb{N}$.

Denote by $P^m[\mathbb{C}^q]$ the space of homogeneous holomorphic polynomials on \mathbb{C}^q of degree m . Note that the space of holomorphic polynomials is dense in $\mathcal{F}(\mathbb{C}^q)$ and any holomorphic polynomial can be written in a unique way as a direct sum of homogeneous polynomials. In fact the following is well known and can be found in [Z12].

Lemma 2.5. *For $\alpha \in \mathbb{N}^q$ let $p_\alpha(z) = z_1^{\alpha_1} \dots z_q^{\alpha_q}$. Then*

$$\|p_\alpha\|_{\mathcal{F}}^2 = \alpha! = \alpha_1! \dots \alpha_q!$$

and the collection

$$\{q_\alpha = \frac{1}{\sqrt{\alpha!}} p_\alpha\}_{\alpha \in \mathbb{N}^q}$$

forms an orthonormal basis for $\mathcal{F}(\mathbb{C}^q)$.

U_q acts on functions on \mathbb{C}^q by $Af(z) = f(A^{-1}z)$. This action leaves $\mathcal{F}(\mathbb{C}^q)$ invariant and, as the measure λ_q is U_q -invariant, it is a unitary representation denoted by π^q . This representation is reducible. In fact this representation leaves the spaces $P^m[\mathbb{C}^q]$ invariant. We denote the corresponding representation by π_m^q . The representation π_m^q is irreducible. The torus $\mathbb{T} \text{id}$ is the center of U_q and acts by the character $z \mapsto z^{-m} \text{id}$. The representations π_m^q and π_k^q are inequivalent if $m \neq k$ as can easily be seen as the action of the center is different. The same conclusion follows from the fact that $d_m = \dim_{\mathbb{C}} P^m[\mathbb{C}^q] \neq d_k$ if $m \neq k$ for $q > 1$ and $d_m = 1$ for $q = 1$. The decomposition of $\mathcal{F}(\mathbb{C}^q)$ in irreducible U_q representations is given by $\mathcal{F}(\mathbb{C}^q) = \bigoplus_{m=0}^{\infty} P^m[\mathbb{C}^q]$.

Definition 2.6. *Let π^{n_i} be the unitary representations of U_{n_i} , acting on $\mathcal{F}(\mathbb{C}^{n_i}), i = 1, \dots, k$, as in the above discussion. The outer tensor product of the representations π_{p_i} , denoted $\pi^{n_1} \otimes \dots \otimes \pi^{n_k}$ is defined by*

$$\pi^{n_1} \otimes \dots \otimes \pi^{n_k}(A_1, \dots, A_k) = \pi^{n_1}(A_1) \otimes \dots \otimes \pi^{n_k}(A_k)$$

for all $(A_1, \dots, A_k) \in U_{n_1} \times \dots \times U_{n_k} = G$. Here $\pi^{n_1}(A_1) \otimes \dots \otimes \pi^{n_k}(A_k)$ is the tensor product of the operators $\pi^{n_i}(A_i)$ acting on the Hilbert spaces $\mathcal{F}(\mathbb{C}^{n_i})$.

The outer tensor product $\pi^{n_1} \otimes \dots \otimes \pi^{n_k}$ is a unitary representation of G acting on $\mathcal{F}(\mathbb{C}^{n_1}) \otimes \dots \otimes \mathcal{F}(\mathbb{C}^{n_k}) = \mathcal{F}(\mathbb{C}^n)$. For more details see [FÓ14].

Lemma 2.7. *Let $\varphi \in L^\infty(\mathbb{C}^n)^G$. Then T_φ intertwines with $\pi^{n_1} \otimes \dots \otimes \pi^{n_k}$.*

Proof. Let $A_i \in U_{n_i}$ and $f_i \in \mathcal{F}(\mathbb{C}^{n_i})$, $i = 1, \dots, m$. Notice that

$$\begin{aligned} T_\varphi(\pi^{n_1} \otimes \dots \otimes \pi^{n_k})(A_1, \dots, A_k) f_1 \otimes \dots \otimes f_k(z) &= T_\varphi f(A_1^{-1} z_{(1)}, \dots, A_k^{-1} z_{(k)}) \\ &= \int_{\mathbb{C}^n} \varphi(w) f(A_1^{-1} z_{(1)}) \dots f(A_k^{-1} z_{(k)}) \overline{k_z(w)} d\lambda_n(w) \\ &= \int_{\mathbb{C}^n} \varphi(A_1 w_{(1)}, \dots, A_k w_{(k)}) f_1 \otimes \dots \otimes f_k(w) \overline{k_z(A_1 w_{(1)}, \dots, A_k w_{(k)})} d\lambda_n(w) \\ &\text{(by the change of variable } (w_{(1)}, \dots, w_{(k)}) \rightarrow (A_1 w_{(1)}, \dots, A_k w_{(k)})\text{)}. \end{aligned}$$

Since

$$\begin{aligned} k_z(A_1 w_{(1)}, \dots, A_k w_{(k)}) &= e^{A_1 w_{(1)} \overline{z_{(1)}} + \dots + A_k w_{(k)} \overline{z_{(k)}}} \\ &= e^{w_{(1)} \overline{A_1^* z_{(1)}} + \dots + w_{(k)} \overline{A_k^* z_{(k)}}} \\ &= e^{w_{(1)} \overline{A_1^{-1} z_{(1)}} + \dots + w_{(k)} \overline{A_k^{-1} z_{(k)}}} \quad (\text{as } A_i \text{ are unitary}) \\ &= k_{(A_1^{-1} z_{(1)}, \dots, A_k^{-1} z_{(k)})}(w) \end{aligned}$$

and φ is G -invariant,

$$\begin{aligned} T_\varphi(\pi^{n_1} \otimes \dots \otimes \pi^{n_k})(A_1, \dots, A_k) f_1 \otimes \dots \otimes f_k(z) &= \int_{\mathbb{C}^n} \varphi(w) f_1 \otimes \dots \otimes f_k(w) \overline{k_{(A_1^{-1} z_{(1)}, \dots, A_k^{-1} z_{(k)})}(w)} d\lambda_n(w) \\ &= (\pi^{n_1} \otimes \dots \otimes \pi^{n_k})(A_1, \dots, A_k) T_\varphi f(z). \end{aligned} \quad \square$$

Since the tensor product of irreducible representations is irreducible (see Proposition 6.75 in [FÓ14] for a proof), $\mathcal{F}(\mathbb{C}^n)$ can be decomposed by irreducible sub-representations of $\pi^{n_1} \otimes \dots \otimes \pi^{n_k}$ as

$$\mathcal{F}(\mathbb{C}^n) = \bigoplus_{m_1, \dots, m_k=0}^{\infty} \mathbb{P}^{m_1}[\mathbb{C}^{n_1}] \otimes \dots \otimes \mathbb{P}^{m_k}[\mathbb{C}^{n_k}].$$

Hence if $\varphi \in L^\infty(\mathbb{C}^n)^G$, $T_\varphi|_{\mathbb{P}^{m_1}[\mathbb{C}^{n_1}] \otimes \dots \otimes \mathbb{P}^{m_k}[\mathbb{C}^{n_k}]} = \gamma_{n, \varphi}(\mathbf{m}) \text{id}$ for some $\gamma_{n, \varphi}(\mathbf{m}) \in \mathbb{C}$ for all $\mathbf{m} = (m_1, \dots, m_k) \in \mathbb{N}_0^k$ by Schur's lemma.

As a consequence of Lemma 2.4, we identify the class of symbols $L^\infty(\mathbb{C}^n)^G$ with essentially bounded functions $a : \mathbb{R}_+^k \rightarrow \mathbb{C}$ and denote the corresponding Toeplitz operator and its spectrum by $T_{n, a}$ and $\gamma_{n, a}$.

Since the above decomposition of $\mathcal{F}(\mathbb{C}^n)$ is multiplicity free and G is compact, the set of all operators that intertwine with $\pi^{n_1} \otimes \dots \otimes \pi^{n_k}$, denoted $\text{End}_G(\mathcal{F}(\mathbb{C}^n))$, is commutative (see in particular Proposition 4.1 and Theorem 6.4 in [DÓQ15] for more details). Then as a consequence of Lemma 2.7 we have the following corollary.

Corollary 2.8. *The C^* -algebra generated by Toeplitz operators with G -invariant symbols is commutative.*

2.3. Computing eigenvalue functions. We denote by $S^q = \{x \in \mathbb{R}^{q+1} \mid |x| = 1\}$ the q -dimensional sphere in \mathbb{R}^{q+1} . We denote by σ the unique U_{q+1} -invariant measure on S^q that satisfy the polar coordinates formula given by

$$\int_{\mathbb{R}^{q+1}} f(x) dx = \int_0^\infty \int_{S^q} f(r\omega) d\sigma(\omega) r^q dr$$

for any $f \in L^1(\mathbb{R}^{q+1}, dx)$ where dx is the Lebesgue measure on \mathbb{R}^{q+1} . See section 2.7 in [F99] for more details.

Lemma 2.9. *Let $f \in L^1(\mathbb{C}^q, dz)$. Then*

$$\int_{\mathbb{C}^q} f(z) dz = \int_0^\infty \int_{S^{2q-1}} f(r\omega) d\sigma(\omega) r^{2q-1} dr.$$

In particular if $f(z) = a(r)$ is $U(q)$ -invariant then

$$\int_{\mathbb{C}^q} f(z) dz = \sigma(S^{2q-1}) \int_0^\infty a(r) r^{2q-1} dr.$$

We recall the following well known fact:

Lemma 2.10. *Let $f, g \in \mathcal{F}^2(\mathbb{C}^q)$ and $\varphi \in L^\infty(\mathbb{C}^q)$. Then*

$$\langle T_\varphi f, g \rangle = \langle \varphi f, g \rangle = \frac{1}{\pi^q} \int_{\mathbb{C}^q} \varphi(z) f(z) \overline{g(z)} e^{-|z|^2} dz.$$

In particular for $f = g = p_\alpha$ we get

$$\langle T_\varphi p_\alpha, p_\alpha \rangle = \frac{1}{\pi^q} \int_{\mathbb{C}^q} \varphi(z) \prod_{j=1}^n |z_j|^{2\alpha_j} e^{-|z_j|^2} dz_1 \cdots dz_n.$$

Proof. Let $P : L^2(\mathbb{C}^q) \rightarrow \mathcal{F}^2(\mathbb{C}^q)$ be the orthogonal projection. Then, as $g \in \mathcal{F}^2(\mathbb{C}^q)$, we get

$$\langle T_\varphi f, g \rangle = \langle P(\varphi f), g \rangle = \langle \varphi f, g \rangle. \quad \square$$

The following is well known and can be found for more general situations in [FK94] and other places. We also point to [F01] for a general discussion on how to integrate polynomials over spheres:

Lemma 2.11. *Let $p_\alpha \in P^m(\mathbb{C}^q)$. Then $\|p_\alpha|_{S^{2q-1}}\|^2 = \frac{2\pi^q}{\Gamma(q+m)} \|p_\alpha\|_{\mathcal{F}}^2$. In particular*

$$\|q_\alpha|_{S^{2q-1}}\|^2 = \frac{2\pi^q}{\Gamma(q+m)}.$$

Proof. We have $p_\alpha(r\omega) = r^m p(\omega)$, $r > 0$. Hence:

$$\begin{aligned} \|p_\alpha\|_{\mathcal{F}}^2 &= \frac{1}{\pi^q} \int_0^\infty \int_{S^{2q-1}} |p_\alpha(r\omega)|^2 d\sigma(\omega) e^{-r^2} r^{2q-1} dr \\ &= \frac{1}{2\pi^q} \left(\int_{S^{2q-1}} |p_\alpha(\omega)|^2 d\sigma(\omega) \right) \left(\int_0^\infty r^{2m+2q-2} e^{-r^2} (2r dr) \right) \\ &= \frac{1}{2\pi^q} \|p_\alpha|_{S^{2q-1}}\|^2 \int_0^\infty u^{m+q-1} e^{-u} du \quad u = r^2, du = 2r dr \\ &= \frac{1}{2\pi^q} \|p_\alpha|_{S^{2q-1}}\|^2 \Gamma(m+q). \quad \square \end{aligned}$$

We recall that the C^* -algebra \mathcal{T}_G generated by G -invariant bounded symbols is commutative and acts by scalars on each of the spaces $P^{m_1}[\mathbb{C}^{n_1}] \otimes \cdots \otimes P^{m_k}[\mathbb{C}^{n_k}]$.

Theorem 2.12. *Assume $\varphi \in L^\infty(\mathbb{C}^n)^G$ let $\mathbf{m} = (m_1, \dots, m_k) \in \mathbb{N}_0^k$. Then*

$$T_\varphi|_{P^{m_1}[\mathbb{C}^{n_1}] \otimes \cdots \otimes P^{m_k}[\mathbb{C}^{n_k}]} = \gamma_{n,\varphi}(\mathbf{m}) \text{id} = \gamma_{n,a}(\mathbf{m}) \text{id}$$

where

$$\gamma_{n,a}(\mathbf{m}) = \frac{1}{(\mathbf{m} + \mathbf{n} - \mathbf{1})!} \int_{\mathbb{R}_+^k} a_\varphi(\sqrt{\mathbf{r}}) \mathbf{r}^{\mathbf{m} + \mathbf{n} - \mathbf{1}} e^{-(r_1 + \cdots + r_k)} dr_1 \cdots dr_k.$$

where we use the notations $\sqrt{\mathbf{r}} = (\sqrt{r_1}, \dots, \sqrt{r_k})$ and $\mathbf{1} = (1, \dots, 1) \in \mathbb{N}^k$.

Proof. Assume that $\varphi(z) = a(|z_{(1)}|, \dots, |z_{(k)}|) = a(r_1, \dots, r_k)$. Let $\alpha_j \in \mathbb{N}_0^{n_j}$ such that $|\alpha_j| = m_j$ and let $q_{\alpha_j} \in \mathbb{P}^{m_j}[\mathbb{C}^{n_j}]$ be the normalized monomials defined in Lemma 2.11. Recall from Lemma 2.11 that

$$\|q_{\alpha_j}|_{S^{2n_j-1}}\|^2 = \frac{2\pi^{n_j}}{\Gamma(n_j + m_j)}$$

and hence

$$\prod_{j=1}^k \|q_{\alpha_j}|_{S^{2n_j-1}}\|^2 = \frac{2^k \pi^n}{(\mathbf{m} + \mathbf{n} - \mathbf{1})!}.$$

Let $\alpha = (\alpha_1, \dots, \alpha_k)$ and define the normalized monomial q_α by

$$q_\alpha(z) = \prod_{j=1}^k q_{\alpha_j}(z_{(j)}).$$

Then $q_\alpha \in \mathbb{P}^{m_1}[\mathbb{C}^{n_1}] \otimes \dots \otimes \mathbb{P}^{m_k}[\mathbb{C}^{n_k}]$ and hence

$$\gamma_{n,\varphi}(\mathbf{m}) = \langle T_\varphi q_\alpha, q_\alpha \rangle.$$

As φ is G -invariant and bounded we can use polar-coordinates on \mathbb{C}^{n_j} , Fubini's theorem, Lemma 2.4, Lemma 2.10 and the fact that q_{α_j} are homogeneous of order m_j , to write

$$\begin{aligned} \langle T_\varphi q_\alpha, q_\alpha \rangle &= \frac{1}{\pi^n} \int_{\mathbb{C}^n} \varphi(z) |q_\alpha(z)|^2 e^{-|z|^2} dz \\ &= \frac{1}{\pi^n} \int_{\mathbb{C}^{n_1}} \dots \int_{\mathbb{C}^{n_k}} a(|z_{(1)}|, \dots, |z_{(k)}|) \prod_{j=1}^k |q_{\alpha_j}(z_{(j)})|^2 e^{-|z_{(j)}|^2} dz_{(1)} \dots dz_{(k)} \\ &= \frac{1}{\pi^n} \int_0^\infty \dots \int_0^\infty a(\mathbf{r}) \prod_{j=1}^k r_j^{2m_j+2n_j-1} e^{-r_j^2} dr_1 \dots dr_k \prod_{j=1}^k \|q_{\alpha_j}|_{S^{2n_j-1}}\|^2 \\ &= \frac{2^k}{(\mathbf{m} + \mathbf{n} - \mathbf{1})!} \int_0^\infty \dots \int_0^\infty a(\mathbf{r}) \prod_{j=1}^k r_j^{2m_j+2n_j-1} e^{-r_j^2} dr_1 \dots dr_k \\ &= \frac{1}{(\mathbf{m} + \mathbf{n} - \mathbf{1})!} \int_0^\infty \dots \int_0^\infty a(\sqrt{\mathbf{r}}) \prod_{j=1}^k r_j^{m_j+n_j-1} e^{-r_j} dr_1 \dots dr_k, \end{aligned}$$

where we in the last line used the substitutions $u = r_j^2$ and hence $du = 2r_j dr_j$. \square

Corollary 2.13. *If $\mathbf{n} = (n_1, \dots, n_k)$ and $\mathbf{m} = (m_1, \dots, m_k)$, then*

$$\gamma_{\mathbf{n},a}(\mathbf{m}) = \gamma_{\mathbf{1},a}(m_1 + n_1 - 1, \dots, m_k + n_k - 1) = \gamma_{\mathbf{1},a}(\mathbf{m} + \mathbf{n} - \mathbf{1}).$$

Note that eigenvalue functions $\gamma_{\mathbf{1},a}$ given above corresponds to Toeplitz operators with separately radial symbols. In particular we obtain the following special cases:

Corollary 2.14 (Grudsky and Vasilevski [GV02]). *Assume that $n = 1$*

$$\gamma_{1,a}(m) = \frac{1}{m!} \int_0^\infty a(\sqrt{r}) r^m e^{-r} dr.$$

If $k = 1$, then $\mathbf{n} = (n)$ and we have the following corollary.

Corollary 2.15. *If $k = 1$ (radial symbols) then*

$$\gamma_{n,a}(m) = \frac{1}{(m + n - 1)!} \int_0^\infty a(\sqrt{r}) r^{m+n-1} e^{-r} dr = \gamma_{1,a}(m + n - 1).$$

If $n = k$, then $\mathbf{n} = (1, \dots, 1) = \mathbf{1}$ and we have the following corollary.

Corollary 2.16. *If $n = k$ (separately radial symbols) then*

$$\gamma_{\mathbf{1},a}(\mathbf{m}) = \frac{1}{\mathbf{m}!} \int_{\mathbb{R}_+^k} a_\varphi(\sqrt{\mathbf{r}}) \mathbf{r}^{\mathbf{m}} e^{-(r_1 + \dots + r_k)} dr_1 \dots dr_k.$$

Corollary 2.17. *If $a(r_1, \dots, r_k) = \prod_{j=1}^k a_j(r_j)$ then*

$$\gamma_{n,a}(m_1, \dots, m_k) = \prod_{j=1}^k \gamma_{1,a_j}(m_j + n_j - 1).$$

3. THE SPACE $C_{b,u}(\mathbb{N}_0^k, \rho_k)$

In this section we introduce C^* -algebra $C_{b,u}(\mathbb{N}_0^k, \rho_k)$, the set of all bounded functions on $\mathbb{N}_0^k \times \mathbb{N}_0^k$ that are uniformly continuous with respect to the square-root metric. These are sometimes called *Square root-slowly oscillating* functions.

For $k \in \mathbb{N}$, let $\rho_k : \mathbb{N}_0^k \times \mathbb{N}_0^k \rightarrow [0, \infty)$ be given by

$$\rho(\mathbf{m}, \mathbf{m}') = |\sqrt{m_1} - \sqrt{m'_1}| + \dots + |\sqrt{m_k} - \sqrt{m'_k}|.$$

for all $\mathbf{m} = (m_1, \dots, m_k), \mathbf{m}' = (m'_1, \dots, m'_k) \in \mathbb{N}_0^k$. Then ρ_k is a metric on \mathbb{N}_0^k . The modulus of continuity with respect to the metric ρ_k of a function $\sigma \in l^\infty(\mathbb{N}_0^k)$ is the function $\omega_{\rho_k, \sigma} : [0, \infty) \rightarrow [0, \infty)$ given by

$$\omega_{\rho_k, \sigma}(\delta) = \sup\{|\sigma(\mathbf{m}) - \sigma(\mathbf{m}')| : \rho(\mathbf{m}, \mathbf{m}') \leq \delta\}.$$

$C_{b,u}(\mathbb{N}_0^k, \rho_k)$ is the set of all bounded functions on \mathbb{N}_0^k that are uniformly continuous with respect to the square-root metric ρ_k :

$$\{\sigma \in l^\infty(\mathbb{N}_0^k) : \lim_{\delta \rightarrow 0} \omega_{\rho_k, \sigma}(\delta) = 0\}.$$

$C_{b,u}(\mathbb{N}_0^k, \rho_k)$ is a closed subalgebra of $l^\infty(\mathbb{N}_0^k)$.

Definition 3.1. *Let $\sigma \in C_{b,u}(\mathbb{N}_0^k, \rho_k)$ and let $\mathbf{s} = (s_1, \dots, s_k) \in \mathbb{N}_0^k$. We define the left and right shift operators on $C_{b,u}(\mathbb{N}_0^k, \rho_k)$ with respect to \mathbf{s} , denoted $\tau_L^{\mathbf{s}}$ and $\tau_R^{\mathbf{s}}$ respectively, by*

$$\tau_L^{\mathbf{s}}\sigma(\mathbf{m}) = \sigma(\mathbf{m} + \mathbf{s}) \quad \text{and} \quad \tau_R^{\mathbf{s}}\sigma(\mathbf{m}) = \begin{cases} \sigma(\mathbf{m} - \mathbf{s}) & ; m_i \geq s_i \ \forall i = 1, \dots, k \\ 0 & ; \text{otherwise} \end{cases}$$

for all $\mathbf{m} = (m_1, \dots, m_k) \in \mathbb{N}_0^k$.

Now we present the following two lemmas that will be used in section 6.

Lemma 3.2. *Let $\sigma \in C_{b,u}(\mathbb{N}_0^k, \rho_k)$ and $\mathbf{s} = (s_1, \dots, s_k) \in \mathbb{N}_0^k$. Then $\tau_L^{\mathbf{s}}\sigma \in C_{b,u}(\mathbb{N}_0^k, \rho_k)$.*

Proof. Boundedness of $\tau_L^{\mathbf{s}}\sigma$ follows easily from the boundedness of σ . Let $\mathbf{m} = (m_1, \dots, m_k), \mathbf{m}' = (m'_1, \dots, m'_k) \in \mathbb{N}_0^k$. By observing the square-root function, we have,

$$|\sqrt{m_i + s_i} - \sqrt{m'_i + s_i}| \leq |\sqrt{m_i} - \sqrt{m'_i}|, \text{ for } i = 1, \dots, k.$$

Hence $\omega_{\rho_k, \tau_L^{\mathbf{s}}\sigma}(\delta) \leq \omega_{\rho_k, \sigma}(\delta)$. □

Lemma 3.3. *Let $\sigma \in C_{b,u}(\mathbb{N}_0^k, \rho_k)$ and $\mathbf{s} = (s_1, \dots, s_k) \in \mathbb{N}_0^k$. Then $\tau_R^{\mathbf{s}}\sigma \in C_{b,u}(\mathbb{N}_0^k, \rho_k)$.*

Proof. Again boundedness of $\tau_R^s \sigma$ follows from the boundedness of σ . Let $s = \max\{s_1, \dots, s_k\}$. Assume $\delta < \sqrt{2s} - \sqrt{2s-1}$ and let $\mathbf{m} = (m_1, \dots, m_k)$, $\mathbf{m}' = (m'_1, \dots, m'_k) \in \mathbb{N}_0^k$ such that $\rho_k(\mathbf{m}, \mathbf{m}') < \delta$. Then $|\sqrt{m_i} - \sqrt{m'_i}| < \sqrt{2s} - \sqrt{2s-1}$ and since

$$\min_{\substack{j, l \in \mathbb{N}_0 \\ i, j \leq 2s}} |\sqrt{j} - \sqrt{l}| = \sqrt{2s} - \sqrt{2s-1},$$

we have $m_i, m'_i > 2s \geq 2s_i$ for all $i = 1, \dots, k$. Notice that

$$\begin{aligned} m_i, m'_i > 2s_i = \frac{s_i}{1 - \frac{1}{2}} &\implies \frac{1}{2}m_i < m_i - s_i, \frac{1}{2}m'_i < m'_i - s_i \\ &\implies \frac{1}{\sqrt{2}}\sqrt{m_i} < \sqrt{m_i - s_i}, \frac{1}{\sqrt{2}}\sqrt{m'_i} < \sqrt{m'_i - s_i} \\ &\implies \frac{1}{\sqrt{2}}(\sqrt{m_i} + \sqrt{m'_i}) < \sqrt{m_i - s_i} + \sqrt{m'_i - s_i} \\ &\implies \frac{|m_i - m'_i|}{\sqrt{m_i - s_i} + \sqrt{m'_i - s_i}} < \sqrt{2} \frac{|m_i - m'_i|}{\sqrt{m_i} + \sqrt{m'_i}} \\ &\implies |\sqrt{m_i - s_i} - \sqrt{m'_i - s_i}| < \sqrt{2} |\sqrt{m_i} - \sqrt{m'_i}|. \end{aligned}$$

Hence $\omega_{\rho_k, \tau_L^s \sigma}(\delta) \leq \omega_{\rho_k, \sigma}(\sqrt{2}\delta)$ for all $\delta < \sqrt{2s} - \sqrt{2s-1}$. \square

The metric ρ_k can be extended to \mathbb{R}_+^k and we denote the set of all functions on \mathbb{R}_+^k that are uniformly continuous with respect to ρ_k by $C_{b,u}(\mathbb{R}_+^k, \rho_k)$. The following lemma shows that any $\sigma \in C_{b,u}(\mathbb{N}_0^k, \rho_k)$ can be extended to some $f \in C_{b,u}(\mathbb{R}_+^k, \rho_k)$. We will use this fact in section 5.

Lemma 3.4. *Let $\sigma \in C_{b,u}(\mathbb{N}_0^k, \rho_k)$. Define f on \mathbb{R}_+^k by*

$$f(x) = f_{\mathbf{m}}(x) := \sigma(\mathbf{m}) + \sum_{l=1}^k \sum_{\substack{i_1, \dots, i_l \in \{1, \dots, k\} \\ i_1 < \dots < i_l}} a_{i_1, \dots, i_l}^l(\mathbf{m}) \prod_{q=1}^l \frac{\sqrt{x_{i_q}} - \sqrt{m_{i_q}}}{\sqrt{m_{i_q} + 1} - \sqrt{m_{i_q}}}$$

where $m_i = \lfloor x_i \rfloor$, $i = 1, \dots, k$ and the coefficients $a_{i_1, \dots, i_l}^l(\mathbf{m})$ are given by

$$a_{i_1, \dots, i_l}^l(\mathbf{m}) = \sum_{q=0}^l (-1)^{l-q} \sum_{\substack{j_1, \dots, j_q \in \{i_1, \dots, i_l\} \\ j_1 < \dots < j_q}} \sigma(\mathbf{m} + e_{j_1} + \dots + e_{j_q})$$

where e_i denotes the standard basis of \mathbb{N}_0^k .

Then $f \in C_{b,u}(\mathbb{N}_0^k, \rho_k)$ and $f|_{\mathbb{N}_0^k} = \sigma$. Moreover $\|f\|_{\infty} = \|\sigma\|_{\infty}$.

The function f defined above is motivated by the interpolation formula for the case $k = 1$, given by

$$f(x) = \sigma(m) + (\sigma(m+1) - \sigma(m)) \frac{\sqrt{x} - \sqrt{m}}{\sqrt{m+1} - \sqrt{m}}$$

where $m = \lfloor x \rfloor$ and $x \in \mathbb{R}_+$. The proof of Lemma 3.4 is a generalization of the proof of Lemma 3.3 in [EM16]. However, the proof depends heavily on combinatorics and induction; we have provided the proof in the appendix.

4. UNIFORM CONTINUITY OF EIGENVALUE FUNCTIONS WITH RESPECT TO THE SQUARE-ROOT METRIC

In this section, we show that the set of all eigenvalue functions is a subset of $C_{b,u}(\mathbb{N}_0^k, \rho_k)$.

Definition 4.1. Let \mathfrak{G}_n be the set of all eigenvalue functions:

$$\mathfrak{G}_n := \{\gamma_{n,a} \mid \varphi \in L^\infty(\mathbb{C}^n)^G\}.$$

Proposition 4.2. The eigenvalue functions $\gamma_{n,a}$ are bounded functions that are uniformly continuous with respect to the square-root metric ρ_k , i.e.,

$$\mathfrak{G}_n \subseteq C_{b,u}(\mathbb{N}_0^k, \rho_k).$$

The boundedness of $\gamma_{n,a}$ follows from the boundedness of the symbol. We proceed to prove the uniform continuity of $\gamma_{n,a}$ with respect to the metric ρ_k . The following lemma is Lemma 4.3 in [EM16] which is used by Esmeral and Maximenko to prove the above theorem for $n = 1$. Then we prove Lemma 4.4 and Lemma 4.5 which are used to prove Proposition 4.6. Theorem 4.2 follows as a corollary of Proposition 4.6.

Lemma 4.3. For any $m \in \mathbb{N}$,

$$\int_0^\infty \left| \frac{r^m}{m!} - \frac{r^{m-1}}{(m-1)!} \right| e^{-r} dr \leq \sqrt{\frac{2}{\pi m}}.$$

As a consequence of the above lemma we prove the following:

Lemma 4.4. For any $m, m' \in \mathbb{N}_0$,

$$\int_0^\infty \left| \frac{r^m}{m!} - \frac{r^{m'}}{m'!} \right| e^{-r} dr \leq 2\sqrt{\frac{2}{\pi}} |\sqrt{m} - \sqrt{m'}|.$$

Proof. Let $\underline{m} = \min\{m, m'\}$ and $\overline{m} = \max\{m, m'\}$. Notice that

$$\begin{aligned} \int_0^\infty \left| \frac{r^m}{m!} - \frac{r^{m'}}{m'!} \right| e^{-r} dr &\leq \sum_{i=\underline{m}+1}^{\overline{m}} \int_0^\infty \left| \frac{r^i}{i!} - \frac{r^{i-1}}{(i-1)!} \right| e^{-r} dr \\ &\leq \sum_{i=\underline{m}+1}^{\overline{m}} \sqrt{\frac{2}{\pi i}} \quad (\text{by Lemma 4.3}) \\ &= \sqrt{\frac{2}{\pi}} \sum_{i=\underline{m}+1}^{\overline{m}} \frac{(\sqrt{i} + \sqrt{i-1})(\sqrt{i} - \sqrt{i-1})}{\sqrt{i}} \\ &\leq \sqrt{\frac{2}{\pi}} \sum_{i=\underline{m}+1}^{\overline{m}} \frac{2\sqrt{i}(\sqrt{i} - \sqrt{i-1})}{\sqrt{i}} = 2\sqrt{\frac{2}{\pi}} |\sqrt{m} - \sqrt{m'}|. \quad \square \end{aligned}$$

Lemma 4.5. Suppose $f_i, g_i : \mathbb{N}_0 \times \mathbb{R}_+ \rightarrow \mathbb{C}$ satisfy $\int_{\mathbb{R}_+} |f_i(m, r)| dr, \int_{\mathbb{R}_+} |g_i(m, r)| dr \leq 1$ for $i = 1, \dots, k$. Then for all $m_i, m'_i \in \mathbb{N}_0$,

$$\int_{\mathbb{R}_+^k} \left| \prod_{i=1}^k f_i(m_i, r_i) - \prod_{i=1}^k g_i(m'_i, r_i) \right| dr_1 \cdots dr_k \leq \sum_{i=1}^k \int_{\mathbb{R}_+} |f_i(m_i, r_i) - g_i(m'_i, r_i)| dr_i.$$

Proof. Notice that the statement is true for $k = 1$. Assume the statement is true for k . Then

$$\begin{aligned}
& \int_{\mathbb{R}_+^{k+1}} \left| \prod_{i=1}^{k+1} f_i(m_i, r_i) - \prod_{i=1}^{k+1} g_i(m'_i, r_i) \right| dr_1 \cdots dr_{k+1} \\
& \leq \int_{\mathbb{R}_+} |f_{k+1}(m_{k+1}, r_{k+1}) - g_{k+1}(m_{k+1}, r_{k+1})| dr_{k+1} \left| \int_{\mathbb{R}_+^k} \prod_{i=1}^k f_i(m_i, r_i) dr_1 \cdots dr_k \right| \\
& \quad + \int_{\mathbb{R}_+} |g_{k+1}(m_{k+1}, r_{k+1})| dr_{k+1} \int_{\mathbb{R}_+^k} \left| \prod_{i=1}^k f_i(m_i, r_i) - \prod_{i=1}^k g_i(m'_i, r_i) \right| dr_1 \cdots dr_k \\
& \leq \sum_{i=1}^{k+1} \int_{\mathbb{R}_+} |f_i(m_i, r_i) - g_i(m'_i, r_i)| dr_i
\end{aligned}$$

by the assumption and because

$$\begin{aligned}
\left| \int_{\mathbb{R}_+^k} \prod_{i=1}^k f_i(m_i, r_i) dr_1 \cdots dr_k \right| & \leq \prod_{i=1}^k \int_{\mathbb{R}_+} |f_i(m_i, r_i)| dr_i \\
& \leq 1.
\end{aligned}$$

Hence the result holds by induction on k . □

Proposition 4.6. *The eigenvalue functions $\gamma_{n,a}$ are Lipchitz with respect to the square-root metric ρ_k : There exists $A_\varphi > 0$ such that*

$$|\gamma_{n,a}(\mathbf{m}) - \gamma_{n,a}(\mathbf{m}')| \leq A_\varphi \rho_k(\mathbf{m}, \mathbf{m}')$$

for all $\mathbf{m} = (m_1, \dots, m_k)$, $\mathbf{m}' = (m'_1, \dots, m'_k) \in \mathbb{N}_0^k$.

Proof. Define $K : \mathbb{N}_0 \times \mathbb{R}_+ \rightarrow [0, \infty)$ by $K(m, r) = \frac{r^m}{m!} e^{-r}$. Then $\int_0^\infty K(m, r) dr = 1$ and

$$\gamma_{\mathbf{1},a}(\mathbf{m}) = \int_{\mathbb{R}_+^k} a_\varphi(\sqrt{r_1}, \dots, \sqrt{r_k}) \prod_{i=1}^k K(m_i, r_i) dr_1 \cdots dr_k$$

where $\mathbf{1} = (1, \dots, 1) \in \mathbb{N}^k$. Notice that

$$\begin{aligned}
& |\gamma_{\mathbf{1},a}(\mathbf{m}) - \gamma_{\mathbf{1},a}(\mathbf{m}')| \\
& \leq \|a_\varphi\|_\infty \int_{\mathbb{R}_+^k} \left| \prod_{i=1}^k K(m_i, r_i) - \prod_{i=1}^k K(m'_i, r_i) \right| dr_1 \cdots dr_k \\
& \leq \|a_\varphi\|_\infty \sum_{i=1}^k \int_0^\infty |K(m_i, r_i) - K(m'_i, r_i)| dr_i \quad (\text{by Lemma 4.5}) \\
& \leq 2\sqrt{\frac{2}{\pi}} \|a_\varphi\|_\infty \rho_k(\mathbf{m}, \mathbf{m}') \quad (\text{by Lemma 4.4}).
\end{aligned}$$

Hence by Corollary 2.13,

$$\begin{aligned}
|\gamma_{n,a}(\mathbf{m}) - \gamma_{n,a}(\mathbf{m}')| &= |\gamma_{1,a}(\mathbf{m} + \mathbf{n} - \mathbf{1}) - \gamma_{1,a}(\mathbf{m}' + \mathbf{n} - \mathbf{1})| \\
&\leq 2\sqrt{\frac{2}{\pi}} \|a_\varphi\|_\infty \rho_k(\mathbf{m} + \mathbf{n} - \mathbf{1}, \mathbf{m}' + \mathbf{n} - \mathbf{1}) \\
&\leq 2\sqrt{\frac{2}{\pi}} \|a_\varphi\|_\infty \rho_k(\mathbf{m}, \mathbf{m}').
\end{aligned}$$

Therefore the function $\gamma_{n,a}$ are uniformly continuous with respect to the metric ρ_k , proving Theorem 4.2. \square

5. DENSITY OF \mathfrak{G}_1 IN $C_{b,u}(\mathbb{N}_0^k, \rho_k)$

In the previous section we showed that $\mathfrak{G}_n \subseteq C_{b,u}(\mathbb{N}_0^k, \rho_k)$. To show that the C^* -algebra generated by \mathfrak{G}_n is $C_{b,u}(\mathbb{N}_0^k, \rho_k)$, it is enough to show that \mathfrak{G}_n is dense in $C_{b,u}(\mathbb{N}_0^k, \rho_k)$ which will be proved in all generality in section 6. In this section we focus on the case $\mathbf{n} = \mathbf{1} = (1, \dots, 1) \in \mathbb{N}^k$. In other words we will show that \mathfrak{G}_1 , the set of eigenvalue functions we get for the case $\mathbf{n} = \mathbf{1}$, is dense in $C_{b,u}(\mathbb{N}_0^k, \rho_k)$. Note that in the above case, the Toeplitz operators under consideration are the Toeplitz operators on $\mathcal{F}(\mathbb{C}^k)$ with separately radial symbols.

This discussion is a generalization of the proof of density for the case $\mathbf{n} = (1)$, presented in [EM16].

5.1. Approximation by convolutions. The main goal of this subsection is to prove Proposition 5.6. For this we present a series of propositions and lemmas.

By a change of variable $(r_1, \dots, r_k) \rightarrow (r_1^2, \dots, r_k^2)$, we have

$$\gamma_{1,a}(\mathbf{m}) = \int_{\mathbb{R}_+^k} a(r_1, \dots, r_k) \prod_{i=1}^k g(m_i, r_i) dr_1 \dots dr_k$$

where $g(m, r) = \frac{2r^{2m+1}e^{-r^2}}{m!}$. Let $h(x) = \sqrt{\frac{2}{\pi}}e^{-2x^2}$.

We have the following lemma from [EM16] (Lemma 6.5).

Lemma 5.1. *Let the functions g and h be as above. Then*

$$\lim_{m \rightarrow \infty} \int_0^\infty |g(m, r) - h(\sqrt{m} - r)| dr = 0.$$

Let $H(x_1, \dots, x_k) = \prod_{i=1}^k h(x_i) = \left(\frac{2}{\pi}\right)^{\frac{k}{2}} e^{-2(x_1^2 + \dots + x_k^2)}$.

Lemma 5.2. *Let $\epsilon > 0$. Then there exists N such that for all $m_i > N$, $i = 1, \dots, k$*

$$\int_{\mathbb{R}_+^k} \left| \prod_{i=1}^k g(m_i, r_i) - H(\sqrt{m_1} - r_1, \dots, \sqrt{m_k} - r_k) \right| dr_1 \dots dr_k < \epsilon.$$

Proof. By Lemma 5.1, there exists N such that for all $m > N$, $\int_0^\infty |g(m, r) - h(\sqrt{m} - r)| dr < \frac{\epsilon}{k}$. Also notice that $\int_0^\infty g(m, r) dr = 1$ and $\int_0^\infty h(\sqrt{m} - r) dr \leq \int_{\mathbb{R}} h(\sqrt{m} - r) dr = \int_{\mathbb{R}} h(r) dr = 1$ for any $m \in \mathbb{N}_0$. Then if $m_i > N$, $i = 1, \dots, k$,

$$\begin{aligned}
& \int_{\mathbb{R}_+^k} \left| \prod_{i=1}^k g(m_i, r_i) - H(\sqrt{m_1} - r_1, \dots, \sqrt{m_k} - r_k) \right| dr_1 \dots dr_k \\
&= \int_{\mathbb{R}_+^k} \left| \prod_{i=1}^k g(m_i, r_i) - \prod_{i=1}^k h(\sqrt{m_i} - r_i) \right| dr_1 \dots dr_k \\
&\leq \sum_{i=1}^k \int_0^\infty |g(m_i, r_i) - h(\sqrt{m_i} - r_i)| dr_i \quad (\text{by Lemma 4.5}) \\
&< k \frac{\epsilon}{k} = \epsilon. \quad \square
\end{aligned}$$

Let $f \in L^1(\mathbb{R}^k, dx)$ and $g \in L^\infty(\mathbb{R}^k, dx)$ where dx denotes the Lebesgue measure on \mathbb{R}^k . Recall that the convolution of f and g , denoted $f * g$, is given by

$$f * g(x) = g * f(x) = \int_{\mathbb{R}^k} f(x - y)g(y)dy, \quad x \in \mathbb{R}^k.$$

The following proposition presents an approximation of the eigenvalue functions by convolutions.

Proposition 5.3. *Let $a \in L^\infty(\mathbb{R}_+^k)$ and let $\epsilon > 0$. Define $H * a$, by considering 'a' as a function on \mathbb{R}^k whose support is \mathbb{R}_+^k . Then there exists N such that for all $m_i > N$, $i = 1, \dots, k$,*

$$|\gamma_{1,a}(m_1, \dots, m_k) - (H * a)(\sqrt{m_1}, \dots, \sqrt{m_k})| < \epsilon.$$

Proof. Since support of a is \mathbb{R}_+^k ,

$$\begin{aligned}
& |\gamma_{1,a}(m_1, \dots, m_k) - (H * a)(\sqrt{m_1}, \dots, \sqrt{m_k})| \\
&= \left| \int_{\mathbb{R}_+^k} \left(\prod_{i=1}^k g(m_i, r_i) - H(\sqrt{m_1} - r_1, \dots, \sqrt{m_k} - r_k) \right) a(y) dr_1 \dots dr_k \right| \\
&\leq \|a\|_\infty \int_{\mathbb{R}_+^k} \left| \prod_{i=1}^k g(m_i, r_i) - H(\sqrt{m_1} - r_1, \dots, \sqrt{m_k} - r_k) \right| dr_1 \dots dr_k.
\end{aligned}$$

Hence the result follows from 5.2 □

Lemma 5.4. *Let $b \in L^\infty(\mathbb{R}^k)$ and $a = \chi_{\mathbb{R}_+^k} b$. Let $\epsilon > 0$. Then there exists M such that for all $x_i > M$, $i = 1, \dots, k$,*

$$|H * a(x_1, \dots, x_k) - H * b(x_1, \dots, x_k)| < \epsilon.$$

Proof. Notice that for $x = (x_1, \dots, x_k) \in \mathbb{R}_+^k$,

$$\begin{aligned}
|H * a(x) - H * b(x)| &\leq \int_{\mathbb{R}^k} |a(y) - b(y)| H(x - y) dy \\
&\leq \|b\|_\infty \int_{\mathbb{R}^k \setminus \mathbb{R}_+^k} H(x - y) dy \\
&\leq \|b\|_\infty \sum_{j=1}^k \int_{\mathbb{R}_-} h(x_j - y_j) dy_j \prod_{\substack{i=1 \\ i \neq j}}^k \int_{\mathbb{R}} h(x_i - y_i) dy_i \\
&= \|b\|_\infty \sum_{j=1}^k \int_{\mathbb{R}_-} h(x_j - y_j) dy_j \\
&= \|b\|_\infty \sum_{j=1}^k \int_{x_j}^\infty h(y_j) dy_j \quad (\text{by a change of variable}).
\end{aligned}$$

The lemma holds as $\int_{x_j}^\infty h(y) dy$ approaches zero as x_j goes to ∞ . □

Proposition 5.5. *Let $g \in C_{b,u}(\mathbb{R}^k)$ and let $\epsilon > 0$. Then there is $b \in L^\infty(\mathbb{R}^k)$ such that*

$$\|H * b - g\|_\infty < \epsilon.$$

Proof. Let \tilde{h} denote the bump function (any compactly supported smooth function would suffice)

$$\tilde{h}(\zeta) = e^{-\frac{1}{1-|\zeta|^2}} \chi_B(\zeta), \quad \zeta \in \mathbb{R}^k$$

where B is the open unit ball in \mathbb{R}^k . Let h denote the normalized Fourier inverse of \tilde{h} , i.e.,

$$h = \frac{\mathcal{F}^{-1}(\tilde{h})}{C}$$

where $C = \|\mathcal{F}^{-1}(\tilde{h})\|_1$. Define the approximate identity h_t on \mathbb{R}^k by

$$h_t(x) = \frac{h(x/t)}{t}, \quad x \in \mathbb{R}^k.$$

Notice that $\lim_{t \rightarrow 0} \|h_t * g - g\|_\infty = 0$. Choose h_{t_0} s.t. $\|h_{t_0} * g - g\|_\infty < \epsilon$. Let $\hat{l} := \frac{\mathcal{F}(h_{t_0})}{\mathcal{F}(H)}$. Then

$$\hat{l}(\zeta) = \frac{\tilde{h}(t_0 \zeta) e^{|\zeta^2|/2}}{C}, \quad \zeta \in \mathbb{R}^k.$$

Since \hat{l} is a Schwartz function and $\mathcal{F}(h_{t_0}) = \mathcal{F}(H) \hat{l}$, $h_{t_0} = H * l$ where $l = \mathcal{F}^{-1}(\hat{l})$. Let $b = l * g$. Then $b \in L^\infty(\mathbb{R}^k)$ and $\|H * b - g\|_\infty = \|h_{t_0} * g - g\|_\infty < \epsilon$. □

Proposition 5.5 can also be proved using Wiener's division lemma, as in the proof of Proposition 5.4 in [EM16].

Proposition 5.6. *Let $\sigma \in C_{b,u}(\mathbb{N}_0^k, \rho_k)$ and let $\epsilon > 0$. Then there exists $a \in L^\infty(\mathbb{R}_+^k)$ and N such that for all $\mathbf{m} = (m_1, \dots, m_k) \in \mathbb{N}_0^k$ with $m_i > N$,*

$$|\sigma(\mathbf{m}) - \gamma_{\mathbf{1},a}(\mathbf{m})| < \epsilon.$$

Proof. This proof is similar to the proof of Proposition 6.8 in [EM16].

By Lemma 3.4, there exists $g \in C_{b,u}(\mathbb{R}_+^k)$ such that $g|_{\mathbb{N}_0^k} = \sigma$ and $\|g\|_\infty = \|\sigma\|_\infty$. Define \tilde{g} on \mathbb{R}^k by $\tilde{g}(x_1, \dots, x_k) = g(x_1^2, \dots, x_k^2)$. Then $\tilde{g} \in C_{b,u}(\mathbb{R}^k)$. Then by Proposition 5.5 there exists $b \in L^\infty(\mathbb{R}^k)$ such that

$$\|H * b - \tilde{g}\|_\infty < \frac{\epsilon}{3}.$$

Define a on \mathbb{R}_+^k by $a = b|_{\mathbb{R}_+^k}$. Then $a \in L^\infty(\mathbb{R}_+^k)$. Then by Proposition 5.3 and Lemma 5.4, there exists N_1, N_2 such that for all $m_i > N_1, i = 1, \dots, k$

$$|\gamma_{1,a}(m_1, \dots, m_k) - (H * a)(\sqrt{m_1}, \dots, \sqrt{m_k})| < \frac{\epsilon}{3}$$

and for all $m_i > N_2$,

$$|H * a(\sqrt{m_1}, \dots, \sqrt{m_k}) - H * b(\sqrt{m_1}, \dots, \sqrt{m_k})| < \frac{\epsilon}{3}.$$

Then for all $\mathbf{m} = (m_1, \dots, m_k)$ with $m_i > N = \max\{N_1, N_2\}$

$$\begin{aligned} |\gamma_{1,a}(\mathbf{m}) - \sigma(\mathbf{m})| &\leq |\gamma_{1,a}(\mathbf{m}) - (H * a)(\sqrt{m_1}, \dots, \sqrt{m_k})| \\ &\quad + |H * a(\sqrt{m_1}, \dots, \sqrt{m_k}) - H * b(\sqrt{m_1}, \dots, \sqrt{m_k})| \\ &\quad + \|H * b - \tilde{g}\|_\infty \\ &< \frac{\epsilon}{3} + \frac{\epsilon}{3} + \frac{\epsilon}{3} = \epsilon. \end{aligned} \quad \square$$

5.2. The density of \mathfrak{G}_1 in $C_{b,u}(\mathbb{N}_0^k, \rho_k)$. The proof of density (Theorem 5.9) requires induction on k . Hence to indicate dependency on k , we identify \mathfrak{G}_1 by \mathfrak{G}_1^k and $\gamma_{1,a}$ by $\gamma_{1,a}^k$ as needed in this subsection.

Lemma 5.7. *Let $k > 1$ and assume \mathfrak{G}_1^{k-1} is dense in $C_{b,u}(\mathbb{N}_0^{k-1}, \rho_{k-1})$. For $i = 1, \dots, k$ and $m_o \in \mathbb{N}_0$, define the set $K_i(m_o) \subset \mathbb{N}_0^k$*

$$K_i(m_o) = \{\mathbf{m} = (m_1, \dots, m_k) \in \mathbb{N}_0^k \mid m_i = m_o\}.$$

Let $\sigma \in C_{b,u}(\mathbb{N}_0^k, \rho_k)$ and let $\epsilon > 0$. Then there exists $a \in L^\infty(\mathbb{R}_+^k)$ s.t.

$$\|\sigma \chi_{K_i(m_o)} - \gamma_{1,a}^k\|_\infty < \epsilon.$$

Proof. Identify $K_i(m_o)$ with \mathbb{N}_0^{k-1} by the map

$$p_{i,m_o} : (m_1, \dots, m_{i-1}, m_o, m_{i+1}, \dots, m_k) \mapsto (m_1, \dots, m_{i-1}, m_{i+1}, \dots, m_k).$$

Define $\sigma_{m_o} : \mathbb{N}_0^{k-1} \rightarrow \mathbb{C}$ by

$$\sigma_{m_o}(m_1, \dots, m_{k-1}) = \sigma \chi_{K_i(m_o)}(p_{i,m_o}^{-1}(m_1, \dots, m_{k-1})).$$

Then $\sigma_{m_o} \in C_{b,u}(\mathbb{N}_0^{k-1}, \rho_{k-1})$. By the assumption there is $b \in L^\infty(\mathbb{R}_+^{k-1})$ s.t.

$$\|\sigma_{m_o} - \gamma_{1,b}^{k-1}\| < \frac{\epsilon}{2}.$$

Since $\chi_{\{m_o\}} \in C_{b,u}(\mathbb{N}_0, \rho_1)$, there is $c \in L^\infty(\mathbb{R}_+)$ s.t.

$$\|\chi_{\{m_o\}} - \gamma_{1,c}\| < \|b\|_\infty \frac{\epsilon}{2}.$$

Now define $a \in L^\infty(\mathbb{R}_+^k)$ by

$$a(\mathbf{x}) = b(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_k)c(x_i) \quad \forall \mathbf{x} \in \mathbb{R}_+^k.$$

Then $\gamma_{\mathbf{1},a}^k(\mathbf{m}) = \gamma_{\mathbf{1},b}^{k-1}(p_i(\mathbf{m}))\gamma_{\mathbf{1},c}(m_i)$ for all $\mathbf{m} \in \mathbb{N}_0^k$. Let $\nu \in C_{b,u}(\mathbb{N}_0^k, \rho_k)$ be defined by $\nu(\mathbf{m}) = \gamma_{\mathbf{1},b}^{k-1}(p_i(\mathbf{m}))\chi_{\{m_0\}}(m_i)$ for all $\mathbf{m} \in \mathbb{N}_0^k$. Then

$$\begin{aligned} \|\sigma\chi_{K_i(m_0)} - \gamma_{\mathbf{1},a}^k\|_\infty &\leq \|\sigma\chi_{K_i(m_0)} - \nu\|_\infty + \|\nu - \gamma_{\mathbf{1},a}^k\|_\infty \\ &\leq \|\sigma_{m_0} - \gamma_{\mathbf{1},b}^{k-1}\|_\infty + \|\gamma_{\mathbf{1},b}^{k-1}\|_\infty \|\gamma_{\mathbf{1},c} - \chi_{\{m_0\}}\|_\infty \\ &< \frac{\epsilon}{2} + \|b\|_\infty \frac{\epsilon}{2\|b\|_\infty} = \epsilon. \end{aligned} \quad \square$$

Lemma 5.8. *Let $k > 1$ and assume \mathfrak{G}_1^{k-1} is dense in $C_{b,u}(\mathbb{N}_0^{k-1}, \rho_{k-1})$. Let $K_N \subset \mathbb{N}_0^k$ be defined by*

$$K_N := \{\mathbf{m} = (m_1, \dots, m_k) \in \mathbb{N}_0^k \mid m_i \leq N \text{ for some } i\}$$

Let $\sigma \in C_{b,u}(\mathbb{N}_0^k, \rho_k)$ and let $\epsilon > 0$. Then there exists $a \in L^\infty(\mathbb{R}_+^k)$ s.t.

$$\|\sigma\chi_{K_N} - \gamma_{\mathbf{1},a}\|_\infty < \epsilon.$$

Proof. For $i = 1, \dots, k$ and $j = 0, 1, \dots, N$, let the sets $K_i(j)$ be as defined in Lemma 5.7. Then there exists $\sigma_{(i,j)} \in C_{b,u}(\mathbb{N}_0^k, \rho_k)$ s.t. $\sigma_{(i,j)}\chi_{K_i(j)} = \sigma_{(i,j)}$ and

$$\sigma\chi_{K_N} = \sum_{i=1}^k \sum_{j=1}^N \sigma_{(i,j)}.$$

The functions $\sigma_{(i,j)}$ can be constructed by considering $\sigma\chi_{K_i(j)}$ and removing up to finitely many points from its support. By Lemma 5.7, there exists $a_{(i,j)} \in L^\infty(\mathbb{R}_+^k)$ s.t.

$$\|\sigma_{(i,j)} - \gamma_{\mathbf{1},a_{(i,j)}}\|_\infty < \frac{\epsilon}{kN}.$$

Define $a \in L^\infty(\mathbb{R}_+^k)$ by $a = \sum_{i=1}^k \sum_{j=1}^N a_{(i,j)}$. Then $\gamma_{\mathbf{1},a} = \sum_{i=1}^k \sum_{j=1}^N \gamma_{\mathbf{1},a_{(i,j)}}$ and

$$\begin{aligned} (5.1) \quad \|\sigma\chi_{K_N} - \gamma_{\mathbf{1},a}\|_\infty &\leq \sum_{i=1}^k \sum_{j=1}^N \|\sigma_{(i,j)} - \gamma_{\mathbf{1},a_{(i,j)}}\|_\infty \\ &< kN \frac{\epsilon}{kN} = \epsilon. \end{aligned} \quad \square$$

Theorem 5.9. \mathfrak{G}_1 is dense in $C_{b,u}(\mathbb{N}_0^k, \rho_k)$.

Proof. The proof is by induction on k . The result for $k = 1$ was proved in [EM16]. Assume the result is true for $k - 1$. Let $\sigma \in C_{b,u}(\mathbb{N}_0^k, \rho_k)$ and let $\epsilon > 0$. By Proposition 5.6 there is N and $b \in L^\infty(\mathbb{R}_+^k)$ s.t. for all $\mathbf{m} \in \mathbb{N}_0^k$ with $m_i > N$,

$$|\sigma(\mathbf{m}) - \gamma_{\mathbf{1},b}(\mathbf{m})| < \frac{\epsilon}{2}.$$

By the induction hypothesis and Lemma 5.8, there exists $c \in L^\infty(\mathbb{R}_+^k)$ s.t.

$$\|(\sigma - \gamma_{\mathbf{1},b})\chi_{K_N} - \gamma_{\mathbf{1},c}\|_\infty < \frac{\epsilon}{2}.$$

Let $a = b + c$. Then $\gamma_{\mathbf{1},a} = \gamma_{\mathbf{1},b} + \gamma_{\mathbf{1},c}$ and

$$\begin{aligned} \|\sigma - \gamma_{\mathbf{1},a}\|_\infty &= \|(\sigma - \gamma_{\mathbf{1},b})\chi_{K_N^c} + (\sigma - \gamma_{\mathbf{1},b})\chi_{K_N} - \gamma_{\mathbf{1},c}\|_\infty \\ &\leq \|(\sigma - \gamma_{\mathbf{1},b})\chi_{K_N^c}\|_\infty + \|(\sigma - \gamma_{\mathbf{1},b})\chi_{K_N} - \gamma_{\mathbf{1},c}\|_\infty \\ &< \sup_{\substack{\mathbf{m} \in \mathbb{N}_0^k \\ m_i > N \forall i}} |\sigma(\mathbf{m}) - \gamma_{\mathbf{1},b}(\mathbf{m})| + \frac{\epsilon}{2} \\ &< \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon. \end{aligned} \quad \square$$

6. THE C^* -ALGEBRA GENERATED BY \mathfrak{G}_n

In this section we prove the density theorem in all generality: the C^* -algebra generated by \mathfrak{G}_n , where $\mathbf{n} = (n_1, \dots, n_k)$, is the C^* -algebra $C_{b,u}(\mathbb{N}_0^k, \rho_k)$. We already noticed that $\gamma_{n,a} \in C_{b,u}(\mathbb{N}_0^k, \rho_k)$ in Proposition 4.2. Recall the shift operators in definition 3.1. Following lemma is a restatement of Corollary 2.13

Lemma 6.1. *The eigenvalue function $\gamma_{n,a}$ is the $(n-1)^{th}$ left shift of $\gamma_{1,a}$, i.e.,*

$$\gamma_{n,a} = \tau_L^{n-1} \gamma_{1,a}.$$

Proposition 6.2. *The set of eigenvalue functions \mathfrak{G}_n is dense in the C^* -algebra $C_{b,u}(\mathbb{N}_0^k, \rho_k)$ for any $\mathbf{n} = (n_1, \dots, n_k) \in \mathbb{N}_0^k$.*

Proof. Let $\sigma \in C_{b,u}(\mathbb{N}_0^k, \rho_k)$ and let $\epsilon > 0$. By Lemma 3.3, $\sigma' := \tau_R^{n-1} \sigma \in C_{b,u}(\mathbb{N}_0^k, \rho_k)$. Also \mathfrak{G}_1 is dense in $C_{b,u}(\mathbb{N}_0^k, \rho_k)$ by Theorem 5.9. Hence there exists $a \in L^\infty(\mathbb{R}_+^k)$ such that $\|\sigma' - \gamma_{1,a}\|_\infty < \epsilon$. Then

$$\begin{aligned} \|\sigma - \gamma_{n,a}\|_\infty &= \|\tau_L^{n-1} \sigma' - \tau_L^{n-1} \gamma_{1,a}\|_\infty \\ &= \|\tau_L^{n-1}\| \|\sigma' - \gamma_{1,a}\|_\infty < \epsilon. \end{aligned}$$

as $\|\tau_L^{n-1}\| < 1$. □

It follows that the C^* -algebra generated by \mathfrak{G}_n is the C^* -algebra $C_{b,u}(\mathbb{N}_0^k, \rho_k)$.

7. A COMPARISON OF $C_{b,u}(\mathbb{N}_0^k, \rho_k)$ WITH $C_{b,u}(\mathbb{N}_0, \rho_1) \otimes \dots \otimes C_{b,u}(\mathbb{N}_0, \rho_1)$

In this section we compare the C^* -algebra $C_{b,u}(\mathbb{N}_0^k, \rho_k)$ with the C^* -tensor product $C_{b,u}(\mathbb{N}_0, \rho_1) \otimes \dots \otimes C_{b,u}(\mathbb{N}_0, \rho_1)$.

By Corollary 2.17 and the density of \mathfrak{G}_1 in $C_{b,u}(\mathbb{N}_0, \rho_1)$, we have the following proposition.

Proposition 7.1. *Let*

$$\tilde{\mathfrak{G}}_n = \{\gamma_{n,a} | a \in L^\infty(\mathbb{R}_+) \otimes \dots \otimes L^\infty(\mathbb{R}_+)\}.$$

Then $\tilde{\mathfrak{G}}_n$ is dense in $C_{b,u}(\mathbb{N}_0, \rho_1) \otimes \dots \otimes C_{b,u}(\mathbb{N}_0, \rho_1)$.

We have the inclusions

$$C_{b,u}(\mathbb{N}_0^{k-1}, \rho_{k-1}) \otimes C_{b,u}(\mathbb{N}_0, \rho_1) \xrightarrow{i_k} C_{b,u}(\mathbb{N}_0^k, \rho_k).$$

In fact

$$C_{b,u}(\mathbb{N}_0, \rho_1) \otimes \dots \otimes C_{b,u}(\mathbb{N}_0, \rho_1) \hookrightarrow C_{b,u}(\mathbb{N}_0^k, \rho_k).$$

However, the above inclusions are not necessarily isomorphisms. We present a counter example for $k = 2$.

7.1. A counter example. Here we construct a counter example to show that $C_{b,u}(\mathbb{N}_0^2, \rho_2)$ is strictly larger than $C_{b,u}(\mathbb{N}_0, \rho_1) \otimes C_{b,u}(\mathbb{N}_0, \rho_1)$. Let

$$I_i : [\sqrt{i}, \sqrt{i} + \pi) \quad ; \quad i \in \mathbb{N}_0.$$

Define $g : \mathbb{N}_0^2 \rightarrow \mathbb{C}$ by

$$g(i, j) = \sin(\sqrt{j} - \sqrt{i}) \chi_{I_i}(\sqrt{j}) \quad ; \quad (i, j) \in \mathbb{N}_0^2.$$

Lemma 7.2. *Let g be the function defined above. Then $g \in C_{b,u}(\mathbb{N}_0^2, \rho_2)$.*

Proof. Clearly g is bounded. First we show that if $|\sqrt{j} - \sqrt{j'}| < \pi$,

$$|g(i, j) - g(i, j')| < |\sqrt{j} - \sqrt{j'}|.$$

If $\sqrt{j}, \sqrt{j'} \notin I_i$,

$$|g(i, j) - g(i, j')| = 0 < |\sqrt{j} - \sqrt{j'}|.$$

W.l.o.g. assume $\sqrt{j} \in I_i$. Then $\sqrt{j'} \in (\sqrt{i} - \pi, \sqrt{i} + 2\pi)$ as $|\sqrt{j} - \sqrt{j'}| < \pi$. Note that if $\sqrt{j'} \in I_i$, $\sin(\sqrt{j'} - \sqrt{i}) = g(j', i)$ and if $\sqrt{j'} \in (\sqrt{i} - \pi, \sqrt{i}) \cup (\sqrt{i} + \pi, \sqrt{i} + 2\pi)$, $\sin(\sqrt{j'} - \sqrt{i}) < 0$. Hence

$$\begin{aligned} |g(i, j) - g(i, j')| &\leq |\sin(\sqrt{j} - \sqrt{i}) - \sin(\sqrt{j'} - \sqrt{i})| \\ &\leq |\sqrt{j} - \sqrt{j'}| \end{aligned}$$

as required. Next we prove that if $|\sqrt{i} - \sqrt{i'}| < \pi$,

$$|g(i, j) - g(i', j)| < |\sqrt{i} - \sqrt{i'}|.$$

If $\sqrt{j} \notin I_i \cup I_{i'}$, $|g(i, j) - g(i', j)| = 0 < |\sqrt{i} - \sqrt{i'}|$. W.l.o.g., assume that $\sqrt{j} \in I_i$. Then $\sqrt{j} \in (\sqrt{i'} - \pi, \sqrt{i'} + 2\pi)$ as $|\sqrt{i} - \sqrt{i'}| < \pi$. Therefore

$$\begin{aligned} |g(i, j) - g(i', j)| &\leq |\sin(\sqrt{j} - \sqrt{i}) - \sin(\sqrt{j} - \sqrt{i'})| \\ &\leq |\sqrt{i} - \sqrt{i'}|, \end{aligned}$$

proving the inequality.

Now notice that for all $(i, j), (i', j') \in \mathbb{N}_0^2$ such that $\rho_2((i, j), (i', j')) < \pi$,

$$\begin{aligned} |g(i, j) - g(i', j')| &\leq |g(i, j) - g(i', j)| + |g(i', j) - g(i', j')| \\ &\leq |\sqrt{i} - \sqrt{i'}| + |\sqrt{j} - \sqrt{j'}| \\ &< \rho_2((i, j), (i', j')), \end{aligned}$$

proving that $g \in C_{b,u}(\mathbb{N}_0^2, \rho_2)$. □

Definition 7.3. If X is a locally compact Hausdorff space and if A is a C^* -algebra equipped with $\|\cdot\|_A$, let $C^b(X, A)$ be the set of all continuous bounded functions $f : X \rightarrow A$ equipped with the norm $\|\cdot\|_\infty$ given by

$$\|f\|_\infty = \sup_{x \in X} \|f(x)\|_A.$$

It is well known that $C^b(X, A)$ is C^* -algebra. Recall that a subset of a topological space is said to be precompact if its closure is compact. The following theorem from Williams [W03] describes a criterion to check whether an element in the C^* -algebra $C^b(X, A)$ belongs to the possibly smaller C^* -algebra $C^b(X) \otimes A$. We will use the Theorem 7.4 to show that $g \notin C_{b,u}(\mathbb{N}_0, \rho_1) \otimes C_{b,u}(\mathbb{N}_0, \rho_1)$.

Theorem 7.4 (Williams [W03]). *If X is a locally compact Hausdorff space and if A is a C^* -algebra, then $f \in C^b(X, A)$ is in $C^b(X) \otimes A$ if and only if the range of f , $R(f) := \{f(x) : x \in X\}$, is precompact.*

In order to use Theorem 7.4 we present several lemmas about g . The proof of Lemma 7.5 is trivial as \mathbb{N}_0 has discrete topology and g is bounded.

Lemma 7.5. *Let g be the function defined above. Define $\tilde{g} : \mathbb{N}_0 \rightarrow C_{b,u}(\mathbb{N}_0, \rho_1)$ by*

$$\tilde{g}(i) = g(i, \cdot).$$

Then $\tilde{g} \in C^b(\mathbb{N}_0, C_{b,u}(\mathbb{N}_0, \rho_1))$.

Lemma 7.6. *Let $\tilde{g} : \mathbb{N}_0 \rightarrow C_{b,u}(\mathbb{N}_0, \rho_1)$ be defined as in Lemma 7.5. Then $\|\tilde{g}(i)\|_\infty \geq \frac{1}{2}$ for all $i \in \mathbb{N}_0$.*

Proof. Notice that for all $i \in \mathbb{N}_0$ and for all $p \in \mathbb{N}$,

$$\sqrt{i+p} - \sqrt{i+p-1} \leq 1 < \frac{2\pi}{3}$$

because $\sqrt{i+p} - \sqrt{i+p-1}$ attains its maximum when $i+p=1$. Fix $i \in \mathbb{N}_0$. Then the sequence $\{\sqrt{i+p}\}_{p=1}^{\infty}$ contains a point in any interval of length $\frac{2\pi}{3}$ and, in particular, it contains a point in $(\sqrt{i} + \frac{\pi}{6}, \sqrt{i} + \frac{5\pi}{6})$. Denote that point by $\sqrt{i+p_0}$. Then $\sqrt{i+p_0} - \sqrt{i} \in (\frac{\pi}{6}, \frac{5\pi}{6})$ and

$$\begin{aligned} \|\tilde{g}(i)\|_{\infty} &\geq |(\tilde{g}(i))(i+p_0)| \\ &= \sin(\sqrt{i+p_0} - \sqrt{i}) \\ &\geq \frac{1}{2}. \end{aligned}$$

□

Lemma 7.7. *Let $\tilde{g} : \mathbb{N}_0 \rightarrow C_{b,u}(\mathbb{N}_0, \rho_1)$ be defined as in Lemma 7.5. Then the range of \tilde{g} is not precompact.*

Proof. Whenever $|\sqrt{i_1} - \sqrt{i_2}| > \pi$, $I_{i_1} \cap I_{i_2} = \emptyset$ and hence

$$\|\tilde{g}(i_1) - \tilde{g}(i_2)\|_{\infty} = \max\{\|\tilde{g}(i_1)\|_{\infty}, \|\tilde{g}(i_2)\|_{\infty}\} \geq \frac{1}{2}.$$

Thus there exists a sequence $\{\tilde{g}(i_s)\}_{s=1}^{\infty}$ s.t. $\|\tilde{g}(i_{s_1}) - \tilde{g}(i_{s_2})\|_{\infty} \geq \frac{1}{2}$ whenever $s_1 \neq s_2$. It follows that the range of \tilde{g} is not totally bounded and hence it is not precompact. □

By Lemma 7.5, Lemma 7.7, and Theorem 7.4, we have that $g \notin C_{b,u}(\mathbb{N}_0, \rho_1) \otimes C_{b,u}(\mathbb{N}_0, \rho_1)$. Also by Lemma 7.2, $g \in C_{b,u}(\mathbb{N}_0^2, \rho_2)$. Hence we have the following proposition.

Proposition 7.8. *$C_{b,u}(\mathbb{N}_0^2, \rho_2)$ is strictly larger than $C_{b,u}(\mathbb{N}_0, \rho_1) \otimes C_{b,u}(\mathbb{N}_0, \rho_1)$.*

8. APPENDIX: THE PROOF OF LEMMA 3.4

In this appendix, we present the proof of Lemma 3.4. It is easy to see that $f|_{\mathbb{N}_0^k} = \sigma$, from the definition of f .

8.1. The uniform norm of f . To show that $\|f\|_{\infty} = \|\sigma\|_{\infty}$, we will use the following lemma.

Lemma 8.1. *Let $s \in \mathbb{N}$ and let $a_i \in (0, 1)$ for all $i = 1, \dots, s$. Then*

$$1 + \sum_{l=1}^s (-1)^l \sum_{\substack{i_1, \dots, i_l \in \{1, \dots, s\} \\ i_1 < \dots < i_l}} \prod_{q=1}^l a_{i_q} \geq 0$$

Proof. Notice that the statement is true for $s = 1$. Assume the result is true for s . Then

$$\begin{aligned}
1 + \sum_{l=1}^{s+1} (-1)^l \sum_{\substack{i_1, \dots, i_l \in \{1, \dots, s+1\} \\ i_1 < \dots < i_l}} \prod_{q=1}^l a_{i_q} &= 1 + \sum_{l=1}^s (-1)^l \sum_{\substack{i_1, \dots, i_l \in \{1, \dots, s\} \\ i_1 < \dots < i_l}} \prod_{q=1}^l a_{i_q} \\
&\quad - a_{s+1} \left(1 + \sum_{l=1}^s (-1)^l \sum_{\substack{i_1, \dots, i_l \in \{1, \dots, s\} \\ i_1 < \dots < i_l}} \prod_{q=1}^l a_{i_q} \right) \\
&= (1 - a_{s+1}) \left(1 + \sum_{l=1}^s (-1)^l \sum_{\substack{i_1, \dots, i_l \in \{1, \dots, s\} \\ i_1 < \dots < i_l}} \prod_{q=1}^l a_{i_q} \right) \\
&\geq 0. \tag*{\square}
\end{aligned}$$

Hence the statement is true by induction on s .

Note that $f_{\mathbf{m}}$ can also be written as

$$f_{\mathbf{m}}(\mathbf{x}) = \sigma(\mathbf{m}) B_0(\mathbf{x}, \mathbf{m}) + \sum_{\substack{j_1, \dots, j_s \in \{1, \dots, k\} \\ j_1 < \dots < j_s}} \sigma(\mathbf{m} + e_{j_1} + \dots + e_{j_s}) B_{j_1, \dots, j_s}(\mathbf{x}, \mathbf{m})$$

where

$$B_{j_1, \dots, j_s}(\mathbf{x}, \mathbf{m}) = \prod_{p=1}^s \frac{\sqrt{x_{j_p}} - \sqrt{m_{j_p}}}{\sqrt{m_{j_p} + 1} - \sqrt{m_{j_p}}} \left(1 + \sum_{l=1}^{k-s} (-1)^l \sum_{\substack{i_1, \dots, i_l \in \{1, \dots, k\} \setminus \{j_1, \dots, j_s\} \\ i_1 < \dots < i_l}} \prod_{q=1}^l \frac{\sqrt{x_{i_q}} - \sqrt{m_{i_q}}}{\sqrt{m_{i_q} + 1} - \sqrt{m_{i_q}}} \right)$$

and

$$B_0(\mathbf{x}, \mathbf{m}) = 1 + \sum_{l=1}^k (-1)^l \sum_{\substack{i_1, \dots, i_l \in \{1, \dots, k\} \\ i_1 < \dots < i_l}} \prod_{q=1}^l \frac{\sqrt{x_{i_q}} - \sqrt{m_{i_q}}}{\sqrt{m_{i_q} + 1} - \sqrt{m_{i_q}}}.$$

The coefficients $B_{j_1, \dots, j_s}(\mathbf{x}, \mathbf{m})$ are computed by summing the products

$$(-1)^{l-s} \prod_{q=1}^l \frac{\sqrt{x_{j_q}} - \sqrt{m_{j_q}}}{\sqrt{m_{j_q} + 1} - \sqrt{m_{j_q}}}$$

over $\{i_1, \dots, i_l\}$ such that $i_1 < \dots < i_l$ and $\{j_1, \dots, j_s\} \subset \{i_1, \dots, i_l\}$.

Consider the sum

$$B_0(\mathbf{x}, \mathbf{m}) + \sum_{\substack{j_1, \dots, j_s \in \{1, \dots, k\} \\ j_1 < \dots < j_s}} B_{j_1, \dots, j_s}(\mathbf{x}, \mathbf{m}).$$

Let $l \in \mathbb{N}$ and let $i_1, \dots, i_l \in \{1, \dots, k\}$ such that $i_1 < \dots < i_l$. Note that $(-1)^{l-s} \prod_{q=1}^l \frac{\sqrt{x_{i_q}} - \sqrt{m_{i_q}}}{\sqrt{m_{i_q} + 1} - \sqrt{m_{i_q}}}$ is a term in $B_{j_1, \dots, j_s}(\mathbf{x}, \mathbf{m})$ whenever $\{j_1, \dots, j_s\} \subset \{i_1, \dots, i_l\}$. Therefore in the above sum, the coefficient of $\prod_{q=1}^l \frac{\sqrt{x_{i_q}} - \sqrt{m_{i_q}}}{\sqrt{m_{i_q} + 1} - \sqrt{m_{i_q}}}$ is given by

$$\sum_{s=0}^l (-1)^{l-s} \binom{l}{s} = (1 - 1)^l = 0.$$

Hence

$$B_0(\mathbf{x}, \mathbf{m}) + \sum_{\substack{j_1, \dots, j_s \in \{1, \dots, k\} \\ j_1 < \dots < j_s}} B_{j_1, \dots, j_s}(\mathbf{x}, \mathbf{m}) = 1.$$

Also, if $\mathbf{x} \in \prod_{i=1}^k [m_i, m_i + 1)$, $B_{j_1, \dots, j_s}(\mathbf{x}, \mathbf{m}) \geq 0$ by Lemma 8.1 and hence

$$\begin{aligned} |f_{\mathbf{m}}(\mathbf{x})| &\leq \|\sigma\|_{\infty} \left(B_0(\mathbf{x}, \mathbf{m}) + \sum_{\substack{j_1, \dots, j_s \in \{1, \dots, k\} \\ j_1 < \dots < j_s}} B_{j_1, \dots, j_s}(\mathbf{x}, \mathbf{m}) \right) \\ &= \|\sigma\|_{\infty}. \end{aligned}$$

Therefore $\|f\|_{\infty} = \|\sigma\|_{\infty}$ as $f|_{\mathbb{N}_0^k} = \sigma$.

8.2. Some useful lemmas. The following lemma is quite useful in the proofs that follow.

Lemma 8.2. *Let $l, s \in \{1, \dots, k\}$. Suppose $i_1, \dots, i_l \in \{1, \dots, k\}$ such that $i_1 < \dots < i_l$. Then*

$$a_{i_1, \dots, s, \dots, i_l}^{l+1}(\mathbf{m}) = a_{i_1, \dots, i_l}^l(\mathbf{m} + e_s) - a_{i_1, \dots, i_l}^l(\mathbf{m})$$

Proof. To keep the notations simple, we will assume $s > i_l$ and we label s by i_{l+1} .

$$\begin{aligned} a_{i_1, \dots, i_{l+1}}^{l+1}(\mathbf{m}) &= \sum_{q=0}^{l+1} (-1)^{l+1-q} \sum_{\substack{j_1, \dots, j_q \in \{i_1, \dots, i_{l+1}\} \\ j_1 < \dots < j_q}} \sigma(\mathbf{m} + e_{j_1} + \dots + e_{j_q}) \\ &= (-1)^{l+1} \sigma(\mathbf{m}) + \sum_{q=1}^l (-1)^{l+1-q} \left(\sum_{\substack{j_1, \dots, j_q \in \{i_1, \dots, i_l\} \\ j_1 < \dots < j_q}} \sigma(\mathbf{m} + e_{j_1} + \dots + e_{j_q}) \right. \\ &\quad \left. + \sum_{\substack{j_1, \dots, j_{q-1} \in \{i_1, \dots, i_l\} \\ j_1 < \dots < j_{q-1}}} \sigma(\mathbf{m} + e_{j_1} + \dots + e_{j_{q-1}} + e_{i_l}) \right) + \sigma(\mathbf{m} + e_{i_1} + \dots + e_{i_l}) \\ &= - \sum_{q=0}^l (-1)^{l-q} \sum_{\substack{j_1, \dots, j_q \in \{i_1, \dots, i_l\} \\ j_1 < \dots < j_q}} \sigma(\mathbf{m} + e_{j_1} + \dots + e_{j_q}) \\ &\quad + \sum_{q=1}^{l+1} (-1)^{l+1-q} \sum_{\substack{j_1, \dots, j_{q-1} \in \{i_1, \dots, i_l\} \\ j_1 < \dots < j_{q-1}}} \sigma(\mathbf{m} + e_{i_l} + e_{j_1} + \dots + e_{j_{q-1}}) \\ &= -a_{i_1, \dots, i_l}^l(\mathbf{m}) + \sum_{q=0}^l (-1)^{l-q} \sum_{\substack{j_1, \dots, j_q \in \{i_1, \dots, i_l\} \\ j_1 < \dots < j_q}} \sigma((\mathbf{m} + e_{i_l}) + e_{j_1} + \dots + e_{j_q}) \\ &= a_{i_1, \dots, i_l}^l(\mathbf{m} + e_{l+1}) - a_{i_1, \dots, i_l}^l(\mathbf{m}). \end{aligned}$$

□

Notice that f is defined on $\prod_{i=1}^k [m_i, m_i + 1)$ by $f_{\mathbf{m}}$. But the following lemma allows f to be defined on $\prod_{i=1}^k [m_i, m_i + 1]$ by $f_{\mathbf{m}}$.

Lemma 8.3. *Let $\mathbf{x} \in \mathbb{R}_+^k$, $\mathbf{m} \in \mathbb{N}_0^k$ and let $s \in \{1, \dots, k\}$. Suppose $[x_i] = m_i$ for all i . Then*

$$f(\mathbf{x} + (1 - x_s)e_s) = f_{\mathbf{m}}(\mathbf{x} + (1 - x_s)e_s).$$

Proof. Note that the s^{th} coordinate of $\mathbf{x} + (1 - x_s)e_s$ is $m_s + 1$. Therefore

$$\begin{aligned}
f_{\mathbf{m}}(\mathbf{x} + (1 - x_s)e_s) &= \sigma(\mathbf{m}) + \sum_{l=1}^{k-1} \sum_{\substack{i_1, \dots, i_l \in \{1, \dots, k\} \setminus \{s\} \\ i_1 < \dots < i_l}} a_{i_1, \dots, i_l}^l(\mathbf{m}) \prod_{q=1}^l \frac{\sqrt{x_{i_q}} - \sqrt{m_{i_q}}}{\sqrt{m_{i_q} + 1} - \sqrt{m_{i_q}}} \\
&\quad + a_s^1 + \sum_{l=2}^k \sum_{\substack{i_1, \dots, i_l \in \{1, \dots, k\} \\ s \in \{i_1, \dots, i_k\} \\ i_1 < \dots < i_l}} a_{i_1, \dots, i_l}^l(\mathbf{m}) \prod_{\substack{q=1 \\ i_q \neq s}}^l \frac{\sqrt{x_{i_q}} - \sqrt{m_{i_q}}}{\sqrt{m_{i_q} + 1} - \sqrt{m_{i_q}}} \\
&= \sigma(\mathbf{m} + e_s) + \left(\sum_{l=1}^{k-1} \sum_{\substack{i_1, \dots, i_l \in \{1, \dots, k\} \setminus \{s\} \\ i_1 < \dots < i_l}} (a_{i_1, \dots, i_l}^l(\mathbf{m}) + a_{i_1, \dots, s, \dots, i_l}^{l+1}(\mathbf{m})) \right. \\
&\quad \left. \times \prod_{q=1}^l \frac{\sqrt{x_{i_q}} - \sqrt{m_{i_q}}}{\sqrt{m_{i_q} + 1} - \sqrt{m_{i_q}}} \right)
\end{aligned}$$

by reindexing the last summand and because $\sigma(\mathbf{m}) + a_s^1 = \sigma(\mathbf{m} + e_s)$. Also,

$$a_{i_1, \dots, i_l}^l(\mathbf{m}) + a_{i_1, \dots, s, \dots, i_l}^{l+1}(\mathbf{m}) = a_{i_1, \dots, i_l}^l(\mathbf{m} + e_s)$$

by Lemma 8.2. Then

$$\begin{aligned}
f_{\mathbf{m}}(\mathbf{x} + (1 - x_s)e_s) &= \sigma(\mathbf{m} + e_s) + \sum_{l=1}^{k-1} \sum_{\substack{i_1, \dots, i_l \in \{1, \dots, k\} \setminus \{s\} \\ i_1 < \dots < i_l}} a_{i_1, \dots, i_l}^l(\mathbf{m} + e_s) \prod_{q=1}^l \frac{\sqrt{x_{i_q}} - \sqrt{m_{i_q}}}{\sqrt{m_{i_q} + 1} - \sqrt{m_{i_q}}} \\
&= f_{\mathbf{m} + e_s}(\mathbf{x} + (1 - x_s)e_s) \\
&= f(\mathbf{x} + (1 - x_s)e_s).
\end{aligned}$$

The second equality in the above computation holds as the s^{th} coordinate of $\mathbf{x} + (1 - x_s)e_s$ is $m_s + 1$ and hence any term in $f_{\mathbf{m} + e_s}(\mathbf{x} + (1 - x_s)e_s)$ indexed by $\{i_1, \dots, i_l\}$ would vanish if $s \in \{i_1, \dots, i_l\}$. \square

As a consequence of above lemma, we have the following corollary.

Corollary 8.4. *Let $\mathbf{m} \in \mathbb{N}_0^k$ and let $\mathbf{x} \in \prod_{i=1}^k [m_i, m_i + 1]$. Then*

$$f(\mathbf{x}) = f_{\mathbf{m}}(\mathbf{x}).$$

Lemma 8.5. *Let $l \in \mathbb{N}$ and let $a_i, b_i \in \mathbb{C}$ such that $|a_i|, |b_i| \leq 1$ for all $i = 1, \dots, l$. Then*

$$\left| \prod_{i=1}^l a_i - \prod_{i=1}^l b_i \right| \leq \sum_{i=1}^l |a_i - b_i|.$$

Proof. Clearly, the result holds for $l = 1$. Assume that the result is true for l . Then by triangle inequality,

$$\begin{aligned} \left| \prod_{i=1}^{l+1} a_i - \prod_{i=1}^{l+1} b_i \right| &\leq |a_{l+1} - b_{l+1}| \left| \prod_{i=1}^l a_i \right| + |b_{l+1}| \left| \prod_{i=1}^l a_i - \prod_{i=1}^l b_i \right| \\ &\leq |a_{l+1} - b_{l+1}| + \left| \prod_{i=1}^l a_i - \prod_{i=1}^l b_i \right| \\ &\leq \sum_{i=1}^{l+1} |a_i - b_i|. \end{aligned}$$

Hence, the result is true by induction on l . □

8.3. The uniform continuity of f with respect to the square-root metric. Let $\mathbf{x}, \mathbf{x}' \in \mathbb{R}_+^k$ s.t. $\rho_k(\mathbf{x}, \mathbf{x}') < \delta$. Let $\mathbf{m} = (m_1, \dots, m_k)$ and $\mathbf{m}' = (m'_1, \dots, m'_k)$ where $m_i = \lfloor x_i \rfloor$ and $m'_i = \lfloor x'_i \rfloor$ for $i = 1, \dots, k$.

Case I. Assume $x'_i \in [m_i, m_{i+1}]$ for all i . Then by Corollary 8.4

$$\begin{aligned} |f(\mathbf{x}) - f(\mathbf{x}')| &\leq \sum_{l=1}^k \sum_{\substack{i_1, \dots, i_l \in \{1, \dots, k\} \\ i_1 < \dots < i_l}} |a_{i_1, \dots, i_l}^l(\mathbf{m})| \\ &\quad \times \left| \prod_{q=1}^l \frac{\sqrt{x_{i_q}} - \sqrt{m_{i_q}}}{\sqrt{m_{i_q} + 1} - \sqrt{m_{i_q}}} - \prod_{q=1}^l \frac{\sqrt{x'_{i_q}} - \sqrt{m_{i_q}}}{\sqrt{m_{i_q} + 1} - \sqrt{m_{i_q}}} \right| \\ &\leq \sum_{l=1}^k \sum_{\substack{i_1, \dots, i_l \in \{1, \dots, k\} \\ i_1 < \dots < i_l}} |a_{i_1, \dots, i_l}^l(\mathbf{m})| \sum_{q=1}^l \frac{|\sqrt{x_{i_q}} - \sqrt{x'_{i_q}}|}{\sqrt{m_{i_q} + 1} - \sqrt{m_{i_q}}} \end{aligned}$$

by Lemma 8.5.

Fix $l \in \{1, \dots, k\}$ and $i_1, \dots, i_l \in \{1, \dots, k\}$ s.t. $i_1 < \dots < i_l$.

If $\sqrt{m_{i_q} + 1} - \sqrt{m_{i_q}} \geq \sqrt{\delta}$ for all $q = 1, \dots, l$,

$$\begin{aligned} |a_{i_1, \dots, i_l}^l(\mathbf{m})| \sum_{q=1}^l \frac{|\sqrt{x_{i_q}} - \sqrt{x'_{i_q}}|}{\sqrt{m_{i_q} + 1} - \sqrt{m_{i_q}}} &\leq |a_{i_1, \dots, i_l}^l(\mathbf{m})| \frac{l\delta}{\sqrt{\delta}} \\ &\leq 2^l l \|\sigma\|_\infty \sqrt{\delta} \end{aligned}$$

as $|a_{i_1, \dots, i_l}^l(\mathbf{m})| \leq 2^l \|\sigma\|_\infty$.

Assume $\sqrt{m_{i_q} + 1} - \sqrt{m_{i_q}} < \sqrt{\delta}$ for some $q \in \{1, \dots, l\}$. W.l.o.g. assume $\sqrt{m_{i_l} + 1} - \sqrt{m_{i_l}} < \sqrt{\delta}$.

If $l \geq 2$,

$$\begin{aligned}
& |a_{i_1, \dots, i_l}^l(\mathbf{m})| \\
&= |a_{i_1, \dots, i_l}^{l-1}(\mathbf{m} + e_{i_l}) - a_{i_1, \dots, i_l}^{l-1}(\mathbf{m})| \\
&\leq \sum_{\substack{j_1, \dots, j_q \in \{i_1, \dots, i_{l-1}\} \\ q \in \{0, \dots, l-1\}}} |\sigma(\mathbf{m} + e_{j_1} + \dots + e_{j_q} + e_{i_l}) - \sigma(\mathbf{m} + e_{j_1} + \dots + e_{j_q})| \\
&\leq 2^{l-1} \omega_{\rho_k, \sigma}(\sqrt{\delta}).
\end{aligned}$$

Hence

$$\begin{aligned}
|f(\mathbf{x}) - f(\mathbf{x}')| &\leq \sum_{l=1}^k \sum_{\substack{i_1, \dots, i_l \in \{1, \dots, k\} \\ i_1 < \dots < i_l}} 2^{l-1} l \max\{2\|\sigma\|_\infty \sqrt{\delta}, \omega_{\rho_k, \sigma}(\sqrt{\delta})\} \\
&= A_k \max\{2\|\sigma\|_\infty \sqrt{\delta}, \omega_{\rho_k, \sigma}(\sqrt{\delta})\}
\end{aligned}$$

where $A_k = k! \sum_{l=1}^k \frac{2^{l-1}}{(l-1)!(k-l)!}$.

Case II. Suppose $x'_i \notin [m_i, m_i + 1]$ for some i . Define $\mathbf{p} = (p_1, \dots, p_k)$, $\mathbf{p}' = (p'_1, \dots, p'_k) \in \mathbb{N}_0^k$ by

$$\begin{aligned}
p_i &= m_i + 1, \quad p'_i = m'_i && \text{if } x_i < x'_i \\
p_i &= m_i, \quad p'_i = m'_i + 1 && \text{if } x_i > x'_i \\
p_i &= m_i = m'_i = p'_i && \text{if } x_i = x'_i
\end{aligned}$$

for $i = 1, \dots, k$. Then $p_i \in [m_i, m_i + 1]$ and $p'_i \in [m'_i, m'_i + 1]$ for all i . Hence by case I and because $\rho_k(\mathbf{p}, \mathbf{p}') \leq \rho_k(\mathbf{x}, \mathbf{x}') < \delta$,

$$\begin{aligned}
|f(\mathbf{x}) - f(\mathbf{x}')| &\leq |f(\mathbf{x}) - f(\mathbf{p})| + |\sigma(\mathbf{p}) - \sigma(\mathbf{p}')| + |f(\mathbf{p}') - f(\mathbf{x}')| \\
&\leq 2A_k \max\{2\|\sigma\|_\infty \sqrt{\delta}, \omega_{\rho_k, \sigma}(\sqrt{\delta})\} + \omega_{\rho_k, \sigma}(\delta).
\end{aligned}$$

In both cases,

$$|f(\mathbf{x}) - f(\mathbf{x}')| \leq 2A_k \max\{2\|\sigma\|_\infty \sqrt{\delta}, \omega_{\rho_k, \sigma}(\sqrt{\delta})\} + \omega_{\rho_k, \sigma}(\delta).$$

Therefore $f \in C_{b,u}(\mathbb{N}_0^k, \rho_k)$. This completes the proof of Lemma 3.4.

Data availability: Data sharing not applicable to this article as no datasets were generated or analysed during the current study.

REFERENCES

- [B75a] F.A. Berezin, *General concept of quantization*. Comm. Math. Phys. **40** (1975), 153–174.
- [B75b] F.A. Berezin, *Quantization in complex symmetric spaces*. Math. USSR-Izv. **9** (1975), 341–379.
- [B61] V. Bargmann, *On a Hilbert space of analytic functions and an associated integral transform I*. Comm. Pure Appl. Math. **14** (1961), 187–214.
- [BC86] C. Berger, L. Coburn, *Toeplitz operators and quantum mechanics*. J. Funct. Anal. **68** (1986), 273–299.
- [BI12] W. Bauer, H. Issa, *Commuting toeplitz operators with quasi-homogeneous symbols on the Segal–Bargmann space*, J. Math. Anal. Appl. **386** (2012), 213–235.
- [BL11] W. Bauer, Y.J. Lee, *Commuting toeplitz operators on the Segal–Bargmann space*. J. Funct. Anal. **260** (2011), 460–489.
- [BS06] A. Böttcher, B. Silbermann, "Analysis of Toeplitz Operators". Second edition. Prepared jointly with Alexei Karlovich. Springer Monographs in Mathematics. Springer-Verlag, Berlin, 2006.
- [BV12] W. Bauer; N. L. Vasilevski, *On the structure of a commutative Banach algebra generated By Toeplitz operators with quasi-radial quasi-homogeneous symbols*. Integral Equations Operator Theory **74** (2012), No. 2, 199–231.
- [C94] L. Coburn, "Berezin–Toeplitz quantization, in Algebraic Methods in Operator Theory", (Birkhauser, Boston, 1994) 101–108.

- [DQ18] M. Dawson, R. Quiroga-Barranco, *Radial Toeplitz operators on the weighted Bergman spaces of Cartan domains*. Representation theory and harmonic analysis on symmetric spaces, 97–114, Contemp. Math., **714**, AMS, 2018.
- [DÓQ15] M. Dawson, G. Ólafsson, R. Quiroga-Barranco, *Commuting Toeplitz operators on bounded symmetric domains and multiplicity-free restrictions of holomorphic discrete series*, Journal of Functional Analysis **268** (2015), No. 7, 1711–1732.
- [DÓQ18] M. Dawson, G. Ólafsson, R. Quiroga-Barranco, *The restriction principle and commuting families of Toeplitz operators on the unit ball*, Sao Paulo J.Math **12** (2018), No. 7, 196–226.
- [DÓQ21] M. Dawson, G. Ólafsson, R. Quiroga-Barranco, *Symbols of some commuting C^* -algebras of Toeplitz operators*. In preparation.
- [EM16] K. Esmeral, E. Maximenko, *Radial Toeplitz operators on the Fock space and square-root-slowly oscillating sequences*, Complex Anal. Oper. Theory **10** (2016), 1655–1677.
- [EU10] M. Englis, H. Upmeyer, *Toeplitz Quantization and Asymptotic Expansions: Peter-Weyl Decomposition*, Integral Equations Operator Theory, **68** No. 3 (2010), 427–449.
- [EU11] M. Englis, H. Upmeyer, *Toeplitz quantization and asymptotic expansions for real bounded symmetric domains*, Math Z, **268** No.3–4 (2011), 931–967.
- [F99] G. B. Folland, "Real analysis: Modern Techniques and Their Applications". New York: Wiley, 1999.
- [F01] G. B. Folland, *How to integrate a polynomial over a sphere*. Amer. Math. Monthly **108** (2001), No 5, 446–448.
- [FK94] J. Faraut, A. Koranyi, "Analysis on Symmetric Cones," Oxford Mathematical Monographs, Oxford University Press, 1994.
- [FÓ14] R. Fabec, G. Ólafsson, "Non-Commutative Harmonic Analysis", Drexville Publishing, St.Gabriel 2014.
- [GKV03] E. A. Grudsky, Karapetyants, A., N. L. Vasilevski, *Toeplitz operators on the unit ball in \mathbb{C}^n with radial symbols*, Journal of Operator Theory **49** (2003), No. 2, 325–346.
- [GMV13] S. M. Grudsky, E. A. Maximenko, N. L. Vasilevski, *Radial Toeplitz operators on the unit ball and slowly oscillating sequences*, Commun. Math. Anal. **14** (2013), No. 2, 77–94.
- [GV02] E. A. Grudsky, N. L. Vasilevski, *Toeplitz operators on the Fock space: Radial component effects*, Integral Equations Operator Theory **44** (2002), No. 1, 10–37.
- [GQV06] S. Grudsky, R. Quiroga-Barranco, N. Vasilevski, *Commutative C^* -algebras of Toeplitz operators and quantization on the unit disk*, J. Funct. Anal. **234** (2006), 1–44.
- [QS11] R. Quiroga-Barranco and A. Sanchez-Nungaray, *Commutative C^* -algebras of Toeplitz operators on complex projective spaces*, Integral Equations Operator Theory **71** (2011), No. 2, 225–243.
- [QS14] R. Quiroga-Barranco, A. Sanchez-Nungaray, *Armando Toeplitz operators with quasi-radial quasi-homogeneous symbols and bundles of Lagrangian frames*. J. Operator Theory **71** (2014), no. 1, 199–222.
- [QS15] R. Quiroga-Barranco, Raul, A. Sanchez-Nungaray, *Toeplitz operators with quasi-homogeneous quasi-radial symbols on some weakly pseudoconvex domains*. Complex Anal. Oper. Theory **9** (2015), no. 5, 1111–1134.
- [QV07a] R. Quiroga-Barranco and N. Vasilevski, *Commutative algebras of Toeplitz operators on the Reinhardt domains*, Integral Equations Operator Theory **59** (2007), No. 1, 67–98.
- [QV07b] R. Quiroga-Barranco and N. Vasilevski, *Commutative C^* -algebras of Toeplitz operators on the unit ball. I. Bargmann-type transforms and spectral representations of Toeplitz operators*, Integral Equations Operator Theory **59** (2007), No. 3, 379–419.
- [QV08] R. Quiroga-Barranco and N. Vasilevski, *Commutative C^* -algebras of Toeplitz operators on the unit ball. II. Geometry of the level sets of symbols*, Integral Equations Operator Theory **60** (2008), No. 1, 89–132.
- [MSR16] M. A. Morales-Ramos, A. Sánchez-Nungaray, J. Ramírez-Ortega, *Toeplitz operators with quasi-separately radial symbols on the complex projective space*. Bol. Soc. Mat. Mex. (3) **22** (2016), no. 1, 213–227.
- [V08] N. L. Vasilevski, *Commutative algebras of Toeplitz operators on the Bergman space*, Operator Theory, Advances and Applications **185**, Birkhäuser, 2008.
- [V10a] N. L. Vasilevski, *Quasi-radial quasi-homogeneous symbols and commutative Banach algebras of Toeplitz operators*. Integral Equations Operator Theory **66** (2010), no. 1, 141–152.
- [V10b] N. L. Vasilevski, *Parabolic quasi-radial quasi-homogeneous symbols and commutative algebras of Toeplitz operators, in Topics in Operator Theory*. Operator Theory: Advances and Applications, Volume 202 . Birkhäuser Basel, 2010.
- [W03] D. P. Williams *Tensor products with bounded continuous functions*, New York Journal of Mathematics **9** (2003), 69–77.
- [Z12] K. Zhu, "Analysis on the Fock Space," Graduate texts in Mathematics, Springer US, 2012.

DEPARTMENT OF MATHEMATICS, LOUISIANA STATE UNIVERSITY, BATON ROUGE, LA 70803, USA, E-MAIL DEWAGE:VDEWAG1@LSU.EDU E-MAIL ÓLAFSSON: OLAFSSON@MATH.LSU.EDU