

# THE MAZUR-ULAM PROPERTY FOR UNIFORM ALGEBRAS

OSAMU HATORI

ABSTRACT. We give a sufficient condition for a Banach space with which the homogeneous extension of a surjective isometry from the unit sphere of it onto another one is real-linear. The condition is satisfied by a uniform algebra and a certain extremely  $C$ -regular space of real-valued continuous functions.

## 1. INTRODUCTION

In 1987 Tingley [25] proposed a problem if a surjective isometry between the unit spheres of Banach spaces is extended to a surjective isometry between whole spaces. Wang [26] seems to be the first to solve Tingley's problem between specific spaces. He dealt with  $C_0(Y)$ , the space of all real (resp. complex) valued continuous functions which vanish at infinity on a locally compact Hausdorff space  $Y$ . Although we do not exhibit each of the literatures, a considerable number of interesting results have shown that Tingley's problem has an affirmative answer. No counterexample is known. Due to [29, p.730] Ding was the first to consider Tingley's problem between different type of spaces [9]. Ding [10, Corollary 2] in fact proved that the real Banach space of all null sequences of real numbers satisfies now we call the Mazur-Ulam property. Later Cheng and Dong [5] introduced the concept of the Mazur-Ulam property. Following Cheng and Dong we say that a real Banach space  $E$  satisfies the Mazur-Ulam property if a surjective isometry between the unit sphere of  $E$  and that of any real Banach space is extended to a surjective real-linear isometry between the whole spaces. Tan [19, 20, 21] showed that the space  $L^p(\mathbb{R})$  for  $\sigma$ -finite positive measure space satisfies the Mazur-Ulam property. In [22] Tan, Huang and Liu introduced the notion of generalized lush spaces and local GL spaces and proved that every local GL space satisfies the Mazur-Ulam property. New achievements by Mori and Ozawa [17] prove that the Mazur-Ulam property is satisfied by unital  $C^*$ -algebras and real von Neumann algebras. Cueto-Avellaneda and Peralta [7] proved that a complex (resp. real) Banach space of all continuous maps with the value in a complex (resp. real) Hilbert space satisfies the

---

2020 *Mathematics Subject Classification.* 46B04, 46B20, 46J10, 46J15 .

*Key words and phrases.* Tingley's problem, the Mazur-Ulam property, surjective isometries, uniform algebras, maximal convex sets, analytic functions.

Mazur-Ulam property (cf. [8]). The result proving that all general JBW\*-triples satisfy the Mazur-Ulam property is established by Becerra-Guerrero, Cueto-Avellaneda, Fernández-Polo and Peralta [2] and Kalenda and Peralta [14]. The study of the Mazur-Ulam property is nowadays a challenging subject of study (cf. [1, 4, 8, 28]).

In this paper we say that a complex Banach space  $B$  satisfies the *complex Mazur-Ulam property*, emphasizing the term ‘complex’, if a surjective isometry between the unit spheres of  $B$  and any complex Banach space is extended to a surjective real-linear isometry between the whole spaces. Jiménez-Vargas, Morales-Compoy, Peralta and Ramírez [13, Theorems 3.8, 3.9] probably provides the first examples of complex Banach spaces satisfying the complex Mazur-Ulam property (cf. [18]). Note that a complex Banach space satisfies the complex Mazur-Ulam property provided that it satisfies the Mazur-Ulam property as a real Banach space (a complex Banach space is a real Banach space simultaneously).

In [12] we proved that a surjective isometry between the unit spheres of uniform algebras is extended to a surjective real-linear isometry between whole of the uniform algebras. In this paper we show the complex Mazur-Ulam property for uniform algebras. Typical examples of a uniform algebra consist of analytic functions of one and several complex-variables such as the disk algebra, the polydisk algebra and the ball algebra. Through the Gelfand transform, the algebra of all bounded analytic functions on a certain domain is considered as a uniform algebra on the maximal ideal space. Hence the main result in this paper provides the first example of a Banach space of analytic functions which satisfies the complex Mazur-Ulam property. For further information about uniform algebras, see [3].

## 2. IS A HOMOGENEOUS EXTENSION LINEAR?

For a real or complex Banach space  $B$ , we denote  $S(B)$  the unit sphere  $\{a \in B : \|a\| = 1\}$  of  $B$ . A maximal convex subset of  $S(B)$  is denoted by  $\mathfrak{S}_B$ . Throughout the paper the map  $T : S(B_1) \rightarrow S(B_2)$  always denotes a surjective isometry with respect to the metric induced by the norm, where  $B_1$  and  $B_2$  are both real Banach spaces or both complex Banach spaces. We define the homogeneous extension  $\tilde{T} : B_1 \rightarrow B_2$  of  $T$  by

$$\tilde{T}(a) = \begin{cases} \|a\|T\left(\frac{a}{\|a\|}\right), & 0 \neq a \in B_1 \\ 0, & a = 0. \end{cases}$$

By the definition  $\tilde{T}$  is a bijection which satisfies that  $\|\tilde{T}(a)\| = \|a\|$  for every  $a \in B_1$ . The Tingley’s problem asks if  $\tilde{T}$  is real-linear or not. In this paper we prove that  $\tilde{T}$  is real-linear for certain Banach spaces  $B_1$  including uniform algebras.

It is well known that for every  $F \in \mathfrak{F}_B$  of a real or complex Banach space  $B$ , there exists an extreme point  $p$  in the closed unit ball  $B(B^*)$  of the dual space  $B^*$  of  $B$  such that  $F = p^{-1}(1) \cap S(B)$  (cf. [24, Lemma 3.3], [12, Lemma 3.1]). Let  $Q$  be the set of all extreme points  $p$  in  $B(B^*)$  such that  $p^{-1}(1) \cap S(B) \in \mathfrak{F}_B$ . We define an equivalence relation  $\sim$  in  $Q$ . We write  $\mathbb{T} = \{z \in \mathbb{C} : |z| = 1\}$  if  $B$  is a complex Banach space, where  $\mathbb{C}$  denotes the space of all complex numbers, and  $\mathbb{T} = \{\pm 1\}$  if  $B$  is a real Banach space.

From Definition 2.1 through Definition 2.6  $B$  is a real or complex Banach space.

**Definition 2.1.** Let  $p_1, p_2 \in Q$ . We denote  $p_1 \sim p_2$  if there exists  $\gamma \in \mathbb{T}$  such that  $p_1^{-1}(1) \cap S(B) = (\gamma p_2)^{-1}(1) \cap S(B)$ .

Note that  $\gamma p \in Q$  provided that  $\gamma \in \mathbb{T}$  and  $p \in Q$ .

**Lemma 2.2.** *The binary relation  $\sim$  is an equivalence relation in  $Q$ .*

A proof is by a routine argument and is omitted.

**Definition 2.3.** A set of all representatives with respect to the equivalence relation  $\sim$  is simply called a set of representatives for  $\mathfrak{F}_B$ .

Note that a set of representatives exists due to the choice axiom. Note also that a set of representatives for  $\mathfrak{F}_B$  is a norming family for  $B$ , hence it is a uniqueness set for  $B$ .

**Example 2.4.** Let  $A$  be a uniform algebra. We assume a uniform algebra as a complex Banach space here and after. We denote the choquet boundary for  $A$  by  $\text{Ch}(A)$ . By the Arens-Kelley theorem (cf. [11, p.29]) we have  $Q = \{\gamma \delta_x : x \in \text{Ch}(A), \gamma \in \mathbb{T}\}$ , where  $\delta_x$  denotes the point evaluation at  $x$ . In this case  $\{\delta_x : x \in \text{Ch}(A)\}$  is a set of representatives for  $\mathfrak{F}_A$ .

**Lemma 2.5.** *Let  $P$  be a set of representatives. For  $F \in \mathfrak{F}_B$  there exists a unique  $(p, \lambda) \in P \times \mathbb{T}$  such that  $F = \{a \in S(B) : p(a) = \lambda\}$ . Conversely, for  $(p, \lambda) \in P \times \mathbb{T}$  we have  $\{a \in S(B) : p(a) = \lambda\}$  is in  $\mathfrak{F}_B$ .*

*Proof.* Let  $F \in \mathfrak{F}_B$ . We first prove the existence of  $(p, \lambda) \in P \times \mathbb{T}$  satisfying the condition. There exists  $p \in Q$  such that  $F = p^{-1}(1) \cap S(B)$ . By the definition, there exists  $q \in P$  such that  $p \sim q$ . Hence there exists  $\gamma \in \mathbb{T}$  such that  $F = (\gamma q)^{-1}(1) \cap S(B)$ . Letting  $\lambda = \bar{\gamma}$  we have  $F = \{a \in S(B) : q(a) = \lambda\}$ . We prove the uniqueness of  $(q, \lambda)$ . Suppose that  $\{a \in S(B) : q(a) = \lambda\} = \{a \in S(B) : q'(a) = \lambda'\}$  for  $(q', \lambda') \in P \times \mathbb{T}$ . Then we have  $q^{-1}(1) \cap S(B) = (\lambda \bar{\lambda}' q')^{-1}(1) \cap S(B)$ , that is  $q \sim q'$ . As  $P$  is the set of all representatives with respect to the equivalence relation  $\sim$  and  $q, q' \in P$  we have  $q = q'$ . It follows that  $\lambda = \lambda'$ .

Conversely, let  $(p, \lambda) \in P \times \mathbb{T}$ . Then  $\bar{\lambda} p \in Q$  and  $\{a \in S(B) : p(a) = \lambda\} = (\bar{\lambda} p)^{-1}(1) \cap S(B) \in \mathfrak{F}_B$ .  $\square$

**Definition 2.6.** For  $(q, \lambda) \in Q \times \mathbb{T}$ , we denote  $F_{q,\lambda} = \{a \in S(B) : q(a) = \lambda\}$ . A map

$$I_B : \mathfrak{F}_B \rightarrow P \times \mathbb{T}$$

is defined by  $I_B(F) = (p, \lambda)$  for  $F = F_{p,\lambda} \in \mathfrak{F}_B$ .

By Lemma 2.5 the map  $I_B$  is well defined and bijective. An important theorem of Cheng, Dong and Tanaka states that a surjective isometry between the unit spheres of Banach spaces preserves maximal convex subsets of the unit spheres. This theorem was first exhibited by Cheng and Dong in [5, Lemma 5.1] and a crystal proof was given by Tanaka [23, Lemma 3.5].

In the following  $P_j$  is a set of representatives for  $\mathfrak{F}_{B_j}$  for  $j = 1, 2$ . Due to the theorem of Cheng, Dong and Tanaka a bijection  $\mathfrak{T} : \mathfrak{F}_{B_1} \rightarrow \mathfrak{F}_{B_2}$  is induced.

**Definition 2.7.** The map  $\mathfrak{T} : \mathfrak{F}_{B_1} \rightarrow \mathfrak{F}_{B_2}$  is defined by  $\mathfrak{T}(F) = T(F)$  for  $F \in \mathfrak{F}_{B_1}$ . The map  $\mathfrak{T}$  is well defined and bijective. Put

$$\Psi = I_{B_2} \circ \mathfrak{T} \circ I_{B_1}^{-1} : P_1 \times \mathbb{T} \rightarrow P_2 \times \mathbb{T}.$$

Define two maps

$$\phi : P_1 \times \mathbb{T} \rightarrow P_2$$

and

$$\tau : P_1 \times \mathbb{T} \rightarrow \mathbb{T}$$

by

$$(1) \quad \Psi(p, \lambda) = (\phi(p, \lambda), \tau(p, \lambda)), \quad (p, \lambda) \in P_1 \times \mathbb{T}.$$

If  $\phi(p, \lambda) = \phi(p, \lambda')$  for every  $p \in P_1$  and  $\lambda, \lambda' \in \mathbb{T}$  we simply write  $\phi(p)$  instead of  $\phi(p, \lambda)$  by discarding the second term  $\lambda$ .

Rewriting the equation (1) we have

$$(2) \quad T(F_{p,\lambda}) = F_{\phi(p,\lambda), \tau(p,\lambda)}, \quad (p, \lambda) \in P_1 \times \mathbb{T}.$$

We point out that

$$(3) \quad \phi(p, -\lambda) = \phi(p, \lambda), \quad \tau(p, -\lambda) = -\tau(p, \lambda)$$

for every  $(p, \lambda) \in P_1 \times \mathbb{T}$ . The reason is as follows. It is well known that  $T(-F) = -T(F)$  for every  $F \in \mathfrak{F}_{B_1}$  (cf. [16, Proposition 2.3]). Hence

$$\begin{aligned} F_{\phi(p,-\lambda), \tau(p,-\lambda)} &= T(F_{p,-\lambda}) = T(-F_{p,\lambda}) = -T(F_{p,\lambda}) \\ &= -F_{\phi(p,\lambda), \tau(p,\lambda)} = F_{\phi(p,\lambda), -\tau(p,\lambda)} \end{aligned}$$

for every  $p \in P_1$ . Since the map  $I_{B_2}$  is a bijection we have (3).

Rewriting (2) we get a basic equation in our argument :

$$(4) \quad \phi(p, \lambda)(T(a)) = \tau(p, \lambda), \quad a \in F_{p,\lambda}.$$

We will prove that under some condition on  $B_1$ , which we will exhibit explicitly later, we have that

$$\phi(p, \lambda) = \phi(p, \lambda'), \quad p \in P_1$$

for every  $\lambda$  and  $\lambda'$  in  $\mathbb{T}$ , and

$$\tau(p, \lambda) = \tau(p, 1) \times \begin{cases} \lambda, & \text{for some } p \in P_1 \\ \bar{\lambda}, & \text{for other } p\text{'s} \end{cases}$$

for  $\lambda \in \mathbb{T}$ . We get, under some condition on  $B_1$ , that

$$(5) \quad \phi(p)(T(a)) = \tau(p, 1) \times \begin{cases} p(a), & \text{for some } p \in P_1 \\ \frac{p(a)}{p(a)}, & \text{for other } p\text{'s} \end{cases}$$

for  $a \in F_{p,p(a)}$ . If the equation (5) holds for any  $a \in S(B_1)$ , without the restriction that  $a \in F_{p,p(a)}$ , then applying the definition of  $\tilde{T}$  we get

$$(6) \quad \phi(p)(\tilde{T}(a)) = \tau(p, 1) \times \begin{cases} p(a), & \text{for some } p \in P_1 \\ \frac{p(a)}{p(a)}, & \text{for other } p\text{'s} \end{cases}$$

for every  $a \in B_1$ , with which we infer that

$$\phi(p)(\tilde{T}(a + rb)) = \phi(p)(\tilde{T}(a)) + \phi(p)(r\tilde{T}(b))$$

for every pair  $a, b \in B_1$  and every real number  $r$ . As  $\phi(P_1) = P_2$  is a norming family, we conclude that  $\tilde{T}$  is real-linear. It means that we arrive at the final positive solution for Tingley's problem under some conditions on  $B_1$ .

### 3. HAUSDORFF DISTANCE BETWEEN THE MAXIMAL CONVEX SUBSETS

Recall that the Hausdorff distance  $d_H(K, L)$  between non-empty closed subsets  $K$  and  $L$  of a metric space with metric  $d(\cdot, \cdot)$  is defined by

$$d_H(K, L) = \max\left\{\sup_{a \in K} d(a, L), \sup_{b \in L} d(b, K)\right\}.$$

**Lemma 3.1.** *Let  $B$  be a complex Banach space and  $P$  a set of representatives for  $\mathfrak{F}_B$ . We consider  $B$  as a metric space induced by the norm. For every  $p \in P$  we have*

$$d_H(F_{p,\lambda}, F_{p,\lambda'}) = |\lambda - \lambda'|$$

for every  $\lambda, \lambda' \in \mathbb{T}$ . Suppose further that  $p' \in P$  is different from  $p$ . If  $\mathfrak{F}_{p,\gamma} \cap \mathfrak{F}_{p',-\gamma} \neq \emptyset$  for some  $\gamma, \gamma' \in \mathbb{T}$ , then we have

$$d_H(F_{p,\gamma}, F_{p',\gamma'}) = 2.$$

*Proof.* Let  $a \in F_{p,\lambda}$ . Then  $\lambda'\bar{\lambda}a \in F_{p,\lambda'}$ . Thus  $d(a, F_{p,\lambda'}) \leq \|a - \lambda'\bar{\lambda}a\| = |\lambda - \lambda'|$ . On the other hand, for every  $b \in F_{p,\lambda'}$  we have  $|\lambda - \lambda'| = |p(a) - p(b)| \leq \|a - b\|$ . Hence  $|\lambda - \lambda'| \leq d(a, F_{p,\lambda'})$ . Therefore  $d(a, F_{p,\lambda'}) = |\lambda - \lambda'|$  holds for every  $a \in F_{p,\lambda}$ . In the similar way we obtain that  $d(b, F_{p,\lambda}) = |\lambda - \lambda'|$  for every  $b \in F_{p,\lambda'}$ . We conclude that  $d_H(F_{p,\lambda}, F_{p,\lambda'}) = |\lambda - \lambda'|$  for every  $\lambda, \lambda' \in \mathbb{T}$ .

Suppose that  $P \ni p' \neq p$  and  $\mathfrak{F}_{p,\gamma} \cap F_{p'-\gamma} \neq \emptyset$  for some  $\gamma, \gamma' \in \mathbb{T}$ . Let  $a \in \mathfrak{F}_{p,\gamma} \cap F_{p'-\gamma}$ . Then for every  $b \in F_{p',\gamma'}$  we have

$$2 = |-\gamma' - \gamma'| = |p'(a) - p'(b)| \leq \|a - b\| \leq 2.$$

Hence  $d(a, F_{p',\gamma'}) = 2$ . It follows that  $d_H(F_{p,\gamma}, F_{p',\gamma'}) = 2$ .  $\square$

Note that the notion of the condition of the Hausdorff distance does not depend on the choice of  $P$ . In fact, we can describe the condition applying the terms of  $Q$ ; the condition of the Hausdorff distance is satisfied by  $B$  if and only if

$$d_H(F_{q,\lambda}, F_{q',\lambda'}) = \begin{cases} |\lambda - \gamma\lambda'|, & q^{-1}(1) \cap S(B) = (\gamma q')^{-1}(1) \cap S(B) \\ 2, & q \not\sim q' \end{cases}$$

for  $q, q' \in Q$ .

**Example 3.2.** Let  $A$  be a uniform algebra and  $P = \{\delta_x : x \in \text{Ch}(A)\}$ , where  $\delta_x$  denotes the point evaluation at  $x$ . Then  $\mathfrak{F}_{\delta_x,\lambda} \cap \mathfrak{F}_{\delta_{x'},\lambda'} \neq \emptyset$  for any pair of different points  $x$  and  $x'$  in  $\text{Ch}(A)$  and any  $\lambda, \lambda' \in \mathbb{T}$  [12, Lemma 4.1]. Thus a uniform algebra satisfies the condition of the Hausdorff distance.

**Lemma 3.3.** *Suppose that  $B_1$  satisfies the condition of the Hausdorff distance. Let  $P_1$  be a set of representatives for  $\mathfrak{F}_{B_1}$ . Then we have  $\phi(p, \lambda) = \phi(p, \lambda')$  for every  $p \in P_1$  and  $\lambda, \lambda' \in \mathbb{T}$ . Put*

$$P_1^+ = \{p \in P_1 : \tau(p, i) = i\tau(p, 1)\}$$

and

$$P_1^- = \{p \in P_1 : \tau(p, i) = \bar{i}\tau(p, 1)\}.$$

Then  $P_1^+$  and  $P_1^-$  are possibly empty disjoint subsets of  $P_1$  such that  $P_1^+ \cup P_1^- = P_1$ . Furthermore we have

$$\tau(p, \lambda) = \lambda\tau(p, 1), \quad p \in P_1^+, \lambda \in \mathbb{T}$$

and

$$\tau(p, \lambda) = \bar{\lambda}\tau(p, 1), \quad p \in P_1^-, \lambda \in \mathbb{T}.$$

*Proof.* First we prove that  $\phi(p, \lambda) = \phi(p, \lambda')$  for every  $p \in P_1$  and  $\lambda, \lambda' \in \mathbb{T}$ . Suppose not: there exist  $p \in P_1$  and  $\lambda, \lambda' \in \mathbb{T}$  such that  $\phi(p, \lambda) \neq \phi(p, \lambda')$ . We may assume that  $|\lambda - \lambda'| < 2$ . (If  $\phi(p, \lambda) \neq \phi(p, \lambda')$  with  $\|\lambda - \lambda'\| =$

2, then  $\phi(p, \lambda) \neq \phi(p, i\lambda)$  or  $\phi(p, \lambda') \neq \phi(p, i\lambda)$ . Replacing  $i\lambda$  by  $\lambda'$  in the first case, and  $i\lambda$  by  $\lambda$  in the later case we have  $\phi(p, \lambda) \neq \phi(p, \lambda')$  with  $|\lambda - \lambda'| = \sqrt{2} < 2$ .) Letting  $\lambda'' = \lambda^2 \bar{\lambda}'$  we have  $|\lambda - \lambda'| = |\lambda - \lambda''|$  and  $\lambda' \neq \lambda''$ . Since  $B_1$  satisfies the condition of the Hausdorff distance, an element  $(q, \alpha) \in P_1 \times \mathbb{T}$  which satisfies

$$(7) \quad d_H(F_{p,\lambda}, F_{q,\alpha}) = |\lambda - \lambda'|$$

is only two elements  $(p, \lambda')$  and  $(p, \lambda'')$ . As  $\mathfrak{T}$  preserves the Hausdorff distance we have

$$d_H(F_{\phi(p,\lambda), \tau(p,\lambda)}, F_{\phi(p,\lambda'), \tau(p,\lambda')}) = d_H(F_{p,\lambda}, F_{p,\lambda'}) = |\lambda - \lambda'|.$$

Applying Lemma 3.1 we have

$$(8) \quad d_H(F_{\phi(p,\lambda), \tau(p,\lambda)}, F_{t,\beta}) = |\lambda - \lambda'|$$

if  $(t, \beta) = (\phi(p, \lambda), \lambda \bar{\lambda}' \tau(p, \lambda)), (\phi(p, \lambda), \lambda \bar{\lambda}'' \tau(p, \lambda)), (\phi(p, \lambda'), \tau(p, \lambda'))$ . As we suppose that  $\phi(p, \lambda) \neq \phi(p, \lambda')$ , the number of points  $(t, \beta) \in P_2 \times \mathbb{T}$  which satisfy (8) is at least three, while the number of points  $(q, \alpha) \in P_1 \times \mathbb{T}$  which satisfies (7) is two since  $B_1$  satisfies the condition of the Hausdorff distance. On the other hand the numbers of  $(q, \alpha)$  and  $(t, \beta)$  which satisfy (7) and (8) respectively must coincide each other because  $\mathfrak{T}$  preserves the Hausdorff distance between the maximal convex subset. We arrive at a contradiction proving that  $\phi(p, \lambda) = \phi(p, \lambda')$  for every  $p \in P_1$  and  $\lambda, \lambda' \in \mathbb{T}$ . In the following we simply write  $\phi(p)$  instead of  $\phi(p, \lambda)$  by discarding the second term.

We prove that  $\tau(p, \lambda) = \lambda \tau(p, 1)$  or  $\bar{\lambda} \tau(p, 1)$  for  $p \in P_1$  and  $\lambda \in \mathbb{T}$ . Since  $\mathfrak{T}$  preserves the Hausdorff distance we have by Lemma 3.1 that

$$\begin{aligned} |\lambda - 1| &= d_H(F_{p,\lambda}, F_{p,1}) = d_H(F_{\phi(p), \tau(p,\lambda)}, F_{\phi(p), \tau(p,1)}) \\ &= |\tau(p, \lambda) - \tau(p, 1)| = |\tau(p, \lambda) \overline{\tau(p, 1)} - 1|, \end{aligned}$$

hence

$$(9) \quad \tau(p, \lambda) \overline{\tau(p, 1)} = \lambda \text{ or } \bar{\lambda}$$

Thus we have  $\tau(p, \lambda) = \lambda \tau(p, 1)$  or  $\bar{\lambda} \tau(p, 1)$ . Letting  $\lambda = i$  we infer that  $P_1^+ \cup P_1^- = P_1$  and  $P_1^+ \cap P_1^- = \emptyset$ . We have

$$2 = d_H(F_{p,i}, F_{p,-i}) = d_H(F_{\phi(p), \tau(p,i)}, F_{\phi(p), \tau(p,-i)}) = |\tau(p, i) - \tau(p, -i)|,$$

hence  $\tau(p, -i) = -\tau(p, i)$  for every  $p \in P_1$ . We show that  $\tau(p, \lambda) = \lambda \tau(p, 1)$  for  $p \in P_1^+$  and  $\lambda \in \mathbb{T}$ . By (9) we may suppose that  $\text{Im } \lambda \neq 0$ . In the case of  $\text{Im } \lambda > 0$  we have

$$\begin{aligned} |i - \lambda| &= d_H(F_{p,i}, F_{p,\lambda}) = d_H(F_{\phi(p), \tau(p,i)}, F_{\phi(p), \tau(p,\lambda)}) \\ &= |\tau(p, i) - \tau(p, \lambda)| = |i\tau(p, 1) - \tau(p, \lambda)| = |i - \tau(p, \lambda) \overline{\tau(p, 1)}|. \end{aligned}$$

Applying (9) we infer that  $\tau(p, \lambda) \overline{\tau(p, 1)} = \lambda$  as  $\text{Im } \lambda > 0$ , hence  $\tau(p, \lambda) = \lambda \tau(p, 1)$ . Next we consider the case of  $\text{Im } \lambda < 0$ . We have

$$\begin{aligned} |-i - \lambda| &= d_H(F_{p, -i}, F_{p, \lambda}) \\ &= d_H(F_{\phi(p), \tau(p, -i)}, F_{\phi(p), \tau(p, \lambda)}) = |\tau(p, -i) - \tau(p, \lambda)|, \end{aligned}$$

as  $\tau(p, -i) = -\tau(p, i)$  we have

$$= |-\tau(p, i) - \tau(p, \lambda)| = |-i\tau(p, 1) - \tau(p, \lambda)| = |-i - \tau(p, \lambda) \overline{\tau(p, 1)}|.$$

As  $\text{Im } \lambda < 0$ , we infer that  $\tau(p, \lambda) = \lambda \tau(p, 1)$  by (9).

Similarly, it can be proved that  $\tau(p, \lambda) = \bar{\lambda} \tau(p, 1)$  if  $p \in P_1^-$  and  $\lambda \in \mathbb{T}$ .  $\square$

#### 4. A SUFFICIENT CONDITION FOR THE COMPLEX MAZUR-ULAM PROPERTY

We define a set  $M_{p, \alpha}$  with which the map  $\Psi$  plays a crucial role to work out the complex Mazur-Ulam property for a uniform algebra. We exhibit the definition of  $M_{p, \alpha}$ . We denote  $\mathbb{D} = \{z \in \mathbb{F} : |z| \leq 1\}$ , where  $\mathbb{F} = \mathbb{R}$  if the corresponding Banach space is a real one and  $\mathbb{F} = \mathbb{C}$  if the corresponding Banach space is a complex one.

**Definition 4.1.** Let  $B$  be a real or complex Banach space and  $P$  a set of representatives for  $\mathfrak{F}_B$ . For  $p \in P$  and  $\alpha \in \mathbb{D}$  we denote

$$M_{p, \alpha} = \{a \in S(B) : d(a, F_{p, \alpha/|\alpha|}) \leq 1 - |\alpha|, d(a, F_{p, -\alpha/|\alpha|}) \leq 1 + |\alpha|\},$$

where we read  $\alpha/|\alpha| = 1$  if  $\alpha = 0$ .

**Lemma 4.2.** *If  $B_j$  is a real Banach space for  $j = 1, 2$ , then we have*

$$T(M_{p, \alpha}) = M_{\phi(p), \alpha \tau(p, 1)}$$

for every  $(p, \alpha) \in P_1 \times \mathbb{T}$ . *If  $B_j$  is a complex Banach space which satisfy the condition of the Hausdorff distance for  $j = 1, 2$ , then we have*

$$T(M_{p, \alpha}) = \begin{cases} M_{\phi(p), \alpha \tau(p, 1)}, & p \in P_1^+ \\ M_{\phi(p), \bar{\alpha} \tau(p, 1)}, & p \in P_1^- \end{cases}$$

for every  $(p, \alpha) \in P_1 \times \mathbb{T}$ .

*Proof.* Due to the definition of the map  $\Psi$  we have

$$T(F_{p, \frac{\alpha}{|\alpha|}}) = F_{\phi(p), \frac{\alpha}{|\alpha|}, \tau(p, \frac{\alpha}{|\alpha|})}$$

and

$$T(F_{p, -\frac{\alpha}{|\alpha|}}) = F_{\phi(p), -\frac{\alpha}{|\alpha|}, \tau(p, -\frac{\alpha}{|\alpha|})}$$



Suppose that  $B_j$  is a real Banach space first. Then by the definition  $\mathbb{T} = \{\pm 1\}$ . By (3) we have  $\phi(p, 1) = \phi(p, -1)$  for every  $p \in P_1$ . Hence  $\phi(p, \lambda)$  does not depend on the second term for a real Banach space. We also have  $\tau(p, -1) = -\tau(p, 1)$  for every  $p \in P_1$  by (3). It follows that

$$T(F_{p, \frac{\alpha}{|\alpha|}}) = F_{\phi(p), \frac{\alpha}{|\alpha|} \tau(p, 1)}$$

and

$$T(F_{p, -\frac{\alpha}{|\alpha|}}) = F_{\phi(p), -\frac{\alpha}{|\alpha|} \tau(p, 1)}$$

As  $T$  is a surjective isometry we have

$$d(a, F_{p, \frac{\alpha}{|\alpha|}}) = d(T(a), F_{\phi(p), \frac{\alpha}{|\alpha|} \tau(p, 1)})$$

and

$$d(a, F_{p, -\frac{\alpha}{|\alpha|}}) = d(T(a), F_{\phi(p), -\frac{\alpha}{|\alpha|} \tau(p, 1)})$$

As  $T$  is a bijection we conclude that

$$T(M_{p, \alpha}) = M_{\phi(p), \alpha \tau(p, 1)}$$

for every  $p \in P_1$  and  $\alpha \in \bar{\mathbb{D}}$ .

Suppose next that  $B_j$  is a complex Banach space which satisfies that the condition of the Hausdorff distance. Let  $p \in P_1^+$ . Then by Lemma 3.3 we have

$$T(F_{p, \frac{\alpha}{|\alpha|}}) = F_{\phi(p), \frac{\alpha}{|\alpha|} \tau(p, 1)}$$

and

$$T(F_{p, -\frac{\alpha}{|\alpha|}}) = F_{\phi(p), -\frac{\alpha}{|\alpha|} \tau(p, 1)}.$$

The left of the proof is similar to the case that  $B_j$  is a real Banach space, hence we see that

$$T(M_{p, \alpha}) = M_{\phi(p), \alpha \tau(p, 1)}$$

for every  $p \in P_1^+$  and  $\alpha \in \bar{\mathbb{D}}$ . The proof for the case of  $p \in P_1^-$  is similar and is omitted.  $\square$

**Lemma 4.3.** *Suppose that  $B$  is a real or complex Banach space and  $P$  is a set of representatives for  $\mathfrak{F}_B$ . For every  $p \in P$  and  $\alpha \in \mathbb{D}$  we have*

$$M_{p, \alpha} \subset \{a \in \mathcal{S}(B) : p(a) = \alpha\}.$$

*Proof.* Let  $a \in M_{p, \alpha}$ . Then for arbitrary  $b \in F_{p, \frac{\alpha}{|\alpha|}}$

$$\left| p(a) - \frac{\alpha}{|\alpha|} \right| = |p(a) - p(b)| \leq \|a - b\|.$$

As  $d(a, F_{p, \frac{\alpha}{|\alpha|}}) \leq 1 - |\alpha|$

$$\left| p(a) - \frac{\alpha}{|\alpha|} \right| \leq d(a, F_{p, \frac{\alpha}{|\alpha|}}) \leq 1 - |\alpha|.$$

In the same way we have

$$\left| p(a) - \left( -\frac{\alpha}{|\alpha|} \right) \right| \leq d(a, F_{p, -\frac{\alpha}{|\alpha|}}) \leq 1 + |\alpha|.$$

Then by the two inequalities  $p(a)$  have to be  $\alpha$ . □

The following is an auxiliary result.

**Proposition 4.4.** *Let  $B_1$  be a complex Banach space and  $P_1$  a set of representatives for  $\mathfrak{S}_{B_1}$ . Assume the following two conditions:*

- i)  $B_1$  satisfies the condition of the Hausdorff distance,
- ii)  $M_{p,\alpha} = \{a \in S(B_1) : p(a) = \alpha\}$  for every  $p \in P_1$  and  $\alpha \in \mathbb{D}$ .

*Then  $B_1$  satisfies the complex Mazur-Ulam property.*

*Proof.* Suppose that  $B_2$  is a complex Banach space and  $T : B_1 \rightarrow B_2$  a surjective isometry. Applying Lemma 3.3 for the equation (4) we get

$$\phi(p)(T(a)) = \tau(p, 1) \times \begin{cases} p(a), & p \in P_1^+ \\ \overline{p(a)}, & p \in P_1^- \end{cases}$$

for  $a \in S(B_1)$  such that  $|p(a)| = 1$ . We prove that

$$\phi(p)(T(a)) = \tau(p, 1) \times \begin{cases} p(a), & p \in P_1^+ \\ \overline{p(a)}, & p \in P_1^- \end{cases}$$

for any  $a \in S(B_1)$ . Let  $a \in S(B_1)$  and  $\alpha = p(a)$ . Then  $|\alpha| \leq 1$ . By condition ii),  $a \in M_{p,\alpha}$  and  $T(a) \in T(M_{p,\alpha})$ . As i) is assumed, we have by Lemma 4.2 that

$$T(M_{p,\alpha}) = \begin{cases} M_{\phi(p), \alpha\tau(p,1)}, & p \in P_1^+ \\ M_{\phi(p), \overline{\alpha}\tau(p,1)}, & p \in P_1^- \end{cases}.$$

Therefore

$$\phi(p)(T(a)) = \alpha\tau(p, 1) = p(a)\tau(p, 1)$$

if  $p \in P_1^+$ . In a similar way we have

$$\phi(p)(T(a)) = \overline{p(a)}\tau(p, 1)$$

if  $p \in P_1^-$ . We conclude that

$$\phi(p)(T(a)) = \tau(p, 1) \times \begin{cases} p(a), & p \in P_1^+ \\ \overline{p(a)}, & p \in P_1^- \end{cases}$$

for every  $a \in S(B_1)$ . Let  $c \in B_1$  and  $c \neq 0$ . Since  $\frac{c}{\|c\|} \in S(B_1)$ , and  $\phi(p)$  and  $p$  are real-linear we have

$$\begin{aligned} \phi(p)(\tilde{T}(c)) &= \phi(p) \left( \|c\| T \left( \frac{c}{\|c\|} \right) \right) = \|c\| \tau(p, 1) \times \begin{cases} p \left( \frac{c}{\|c\|} \right), & p \in P_1^+ \\ p \left( \frac{c}{\|c\|} \right), & p \in P_1^- \end{cases} \\ &= \tau(p, 1) \times \begin{cases} p(c), & p \in P_1^+ \\ p(c), & p \in P_1^- \end{cases} \end{aligned}$$

We infer that

$$\begin{aligned} \phi(p)(\tilde{T}(a+rb)) &= \tau(p, 1) \times \begin{cases} p(a+rb), & p \in P_1^+ \\ p(a+rb), & p \in P_1^- \end{cases} \\ &= \tau(p, 1) \times \begin{cases} p(a), & p \in P_1^+ \\ p(a), & p \in P_1^- \end{cases} + \tau(p, 1) \times \begin{cases} rp(b), & p \in P_1^+ \\ rp(b), & p \in P_1^- \end{cases} \\ &= \phi(p)(\tilde{T}(a) + r\tilde{T}(b)) \end{aligned}$$

for every pair  $a, b \in B_1$  and every real number  $r$ . As  $\phi(P_1) = P_2$  is a norming family we conclude that

$$\tilde{T}(a+rb) = \tilde{T}(a) + r\tilde{T}(b)$$

for every pair  $a, b \in B_1$  and every real number  $r$ . By the definition of  $\tilde{T}$  it is a bijection from  $B_1$  onto  $B_2$  and it satisfies the equality  $\|\tilde{T}(a)\| = \|a\|$  for every  $a \in B_1$ . Thus  $\tilde{T}$  is a surjective real-linear isometry from  $B_1$  onto  $B_2$  which extend  $T$ .  $\square$

The following is the main result in this paper.

**Theorem 4.5.** *A uniform algebra satisfies the complex Mazur-Ulam property.*

*Proof.* Let  $A$  be a uniform algebra. Put  $P = \{\delta_x : x \in \text{Ch}(A)\}$ , where  $\delta_x$  is the point evaluation at  $x \in \text{Ch}(A)$ , the Choquet boundary. Then  $P$  is a set of representatives for  $\mathfrak{F}_A$ . It is known that a uniform algebra satisfies the condition of the Hausdorff distance [12, Lemma 4.1]. By [12, Lemma 6.3] we have  $M_{\delta_x, \alpha} = \{f \in S(A) : f(x) = \alpha\}$  for every  $x \in \text{Ch}(A)$  and  $\alpha \in \mathbb{D}$ . Thus the conditions i) and ii) of Proposition 4.4 holds for  $A$ . Thus  $A$  satisfies the complex Mazur-Ulam property by Proposition 4.4.  $\square$

## 5. THE CASE OF A REAL BANACH SPACE

Throughout the section we denote  $B_j$  a real Banach space,  $P_j$  a set of representatives for  $\mathfrak{F}_{B_j}$  and  $T : S(B_1) \rightarrow S(B_2)$  is a surjective isometry. We

have by (3) that  $\phi(p, 1) = \phi(p, -1)$  for every  $p \in P_1$ . As  $\mathbb{T} = \{\pm 1\}$  for real Banach spaces we have that  $\phi(p, \lambda)$  does not depend the second term for real Banach spaces. We also have  $\tau(p, -1) = -\tau(p, 1)$  for every  $p \in P_1$  by (3). The situation is rather simple than the case of complex Banach spaces, and by (4) we have the following equation (10) without further assumption on  $B_1$ , i.e.,

$$(10) \quad \phi(p)(T(a)) = \tau(p, 1)p(a)$$

for every  $p \in P_1$  and  $a \in S(B_1)$  with  $|p(a)| = 1$ .

**Proposition 5.1.** *Suppose that*

$$(11) \quad M_{p, \alpha} = \{a \in S(B_1) : p(a) = \alpha\}$$

for every  $p \in P$  and  $-1 \leq \alpha \leq 1$ . Then  $T$  is extended to a surjective real-linear isometry form  $B_1$  onto  $B_2$ . Hence  $B_1$  satisfies the Mazur-Ulam property.

*Proof.* We first prove the equation (10) for every  $p \in P_1$  and  $a \in S(B_1)$  without assuming that  $|p(a)| = 1$ . Let  $p \in P_1$  and  $a \in S(B_1)$ . Put  $\alpha = p(a)$ . Then by (11)  $a \in M_{p, \alpha}$ . We have by Lemma 4.2 that

$$\phi(p)(T(a)) = \alpha\tau(p, 1) = \tau(p, 1)p(a).$$

It follows that for the homogeneous extension  $\tilde{T}$  of  $T$  we have

$$\begin{aligned} \phi(p)(\tilde{T}(c)) &= \phi(p)\left(\|c\|T\left(\frac{c}{\|c\|}\right)\right) \\ &= \|c\|\tau(p, 1)p\left(\frac{c}{\|c\|}\right) = \tau(p, 1)p(c) \end{aligned}$$

for every  $0 \neq c \in B_1$ . As the equality  $\phi(p)(\tilde{T}(0)) = \tau(p, 1)p(0)$  holds, we obtain for  $a, b \in B_1$  and a real number  $r$  that

$$\phi(p)(\tilde{T}(a + rb)) = \tau(p, 1)p(a + rb) = \tau(p, 1)p(a) + r\tau(p, 1)p(b)$$

and

$$\begin{aligned} \phi(p)(\tilde{T}(a) + r\tilde{T}(b)) &= \phi(p)(\tilde{T}(a)) + r\phi(p)(\tilde{T}(b)) \\ &= \tau(p, 1)p(a) + r\tau(p, 1)p(b). \end{aligned}$$

It follows that

$$\phi(p)(\tilde{T}(a + rb)) = \phi(p)(\tilde{T}(a) + r\tilde{T}(b))$$

for every  $p \in P_1$ ,  $a, b \in B_1$ , and every real number  $r$ . As  $\phi(P_1) = P_2$  is a norming family we see that  $\tilde{T}$  is real-linear on  $B_1$ . As the homogeneous extension is a norm-preserving bijection as is described in the first part of section 2 we complete the proof.  $\square$

**Definition 5.2.** For a locally compact Hausdorff space, we denote by  $C_0(Y, \mathbb{R})$  a real Banach space of all real-valued continuous functions which vanish at infinity on  $Y$ . Let  $E$  be a closed subspace of  $C_0(Y, \mathbb{R})$  which separates the points of  $Y$ , that is, for any pair  $y_1$  and  $y_2$  of different points in  $Y$  there exists a function  $e \in E$  such that  $e(y_1) \neq e(y_2)$ . In this paper we say that  $E$  satisfies the condition (r) if for any triple  $y \in \text{Ch}(E)$ , a neighborhood  $V$ , and  $\varepsilon > 0$  there exists  $u \in E$  such that  $0 \leq u \leq 1 = u(y)$  on  $Y$  and  $0 \leq u \leq \varepsilon$  on  $Y \setminus V$ .

Note that if  $E$  satisfies the condition (r), then it is extremely  $C$ -regular (cf. [11, Definition 2.3.9]).

**Example 5.3.** The space  $C_0(Y, \mathbb{R})$  satisfies the condition (r) for any locally compact Hausdorff space  $Y$ . Let

$$E = \{f \in C_0((0, 2], \mathbb{R}) : f(t) = at \text{ on } (0, 1] \text{ for some } a \in \mathbb{R}\}.$$

Then  $\text{Ch}(E) = [1, 2]$  and  $E$  satisfies the condition (r).

Liu [15, Corollary 6] established the Mazur-Ulam property for  $C_0(Y, \mathbb{R})$  when  $Y$  is compact. The following generalizes the result of Liu.

**Corollary 5.4.** *Let  $Y$  be a locally compact Hausdorff space. Suppose that  $E$  is a closed subspace of  $C_0(Y, \mathbb{R})$  which separates the points of  $Y$ . Suppose that  $E$  satisfies the condition (r). Then  $E$  satisfies the Mazur-Ulam property. In particular,  $C_0(Y, \mathbb{R})$  satisfies the Mazur-Ulam property.*

*Proof.* Let  $\text{Ch}(E)$  be the Choquet boundary for  $E$ . Then by the Arens-Kelley theorem  $P = \{\delta_x : x \in \text{Ch}(E)\}$  is a set of representatives for  $\mathfrak{S}_E$ . Suppose that  $p \in P$  and  $-1 \leq \alpha \leq 1$ . Proving the inclusion

$$M_{p, \alpha} \supset \{a \in S(E) : p(a) = \alpha\},$$

we get (11) by Lemma 4.3. It will follow by Proposition 5.1 that  $E$  satisfies the Mazur-Ulam property. Let  $a \in S(B_1)$  such that  $p(a) = \alpha$ . First we consider the case of  $\alpha = 1$ . By definition  $a \in F_{p, 1}$ , hence  $d(a, F_{p, 1}) = 0 = 1 - |\alpha|$ . For every  $b \in F_{p, -1}$  we have  $2 = |p(a) - p(b)| \leq \|a - b\| \leq 2$ . Hence  $d(a, F_{p, -1}) = 2 = 1 + |\alpha|$ . Thus  $a \in M_{p, \alpha}$  if  $\alpha = 1$ . A proof for  $\alpha = -1$  is similar and is omitted.

We consider the case that  $-1 < \alpha < 1$ . Let  $a \in S(B_1)$  such that  $p(a) = \alpha$ . Let  $\varepsilon$  be  $0 < \varepsilon < 1 - |\alpha|$ . Put  $K = \{q \in P : q(a) = \alpha\}$  and  $F_0 = \{q \in P : |q(a) - \alpha| \geq \varepsilon/4\}$ . For each positive integer  $n$ , put  $F_n = \{q \in P : \varepsilon/2^{n+2} \leq |q(a) - \alpha| \leq \varepsilon/2^{n+1}\}$ . Then  $P = K \cup (\bigcup_{m=0}^{\infty} F_m)$ . For each positive integer  $n$ , choose  $u_n \in S(B_1)$  such that  $0 \leq u_n \leq 1 = p(u_n)$  on  $Y$  and  $u_n = 0$  on  $F_0 \cup F_n$ . Since  $P = \{\delta_x : x \in \text{Ch}(E)\}$  and  $E$  satisfies the condition (r) such  $u_n \in E$  exists. Put  $u = \sum_{n=1}^{\infty} u_n/2^n$ . As the supremum norm of  $u_n$  is dominated by

1 for every  $n$ ,  $u$  converges uniformly on  $Y$  in  $E$ . Then put

$$g_+ = \left( \frac{\alpha}{|\alpha|} - \alpha \right) u + a,$$

and

$$g_- = \left( -\frac{\alpha}{|\alpha|} - \alpha \right) u + a.$$

We see that  $g_+ \in F_{p, \frac{\alpha}{|\alpha|}}$ ,  $\|g_+ - a\| = 1 - |\alpha|$ , and  $d(a, F_{p, \frac{\alpha}{|\alpha|}}) \leq 1 - |\alpha|$ . A proof is as follows. All what is really needed is to prove that  $|q(g_-)| \leq 1$  for every  $q \in P$  as it is evident that  $g_+ \in E$  and  $p(g_+) = \frac{\alpha}{|\alpha|}$  by the definition of  $g_+$ . If  $q \in F_0$ , then  $q(u) = 0$  asserts that  $|q(g_+)| = |q(a)| \leq 1$ . Suppose that  $q \in F_n$  for some positive integer  $n$ . Then  $q(u_n) = 0$  and  $0 \leq u_k \leq 1$  on  $Y$  we have  $|q(u)| \leq \sum_{k \neq n} 1/2^k = 1 - 1/2^n$ . Thus

$$\begin{aligned} |q(g_+)| &\leq \left| \frac{\alpha}{|\alpha|} - \alpha \right| (1 - 1/2^n) + |q(a) - \alpha| + |\alpha| \\ &\leq (1 - |\alpha|)(1 - 1/2^n) + \varepsilon/2^{n+1} + |\alpha| \\ &\leq (1 - |\alpha|)(1 - 1/2^n) + (1 - |\alpha|)/2^{n+1} + |\alpha| \leq 1 \end{aligned}$$

Suppose that  $q \in K$ . Then

$$|q(g_+)| \leq 1 - |\alpha| + |\alpha| = 1.$$

We conclude that  $g_+ \in F_{p, \frac{\alpha}{|\alpha|}}$ . By the definition of  $g_+$ , we infer that  $\|g_+ - a\| = (1 - |\alpha|)\|u\| = 1 - |\alpha|$ . It follows that

$$(12) \quad d(a, F_{p, \frac{\alpha}{|\alpha|}}) \leq 1 - |\alpha|.$$

We also see that  $g_- \in F_{p, -\frac{\alpha}{|\alpha|}}$ ,  $\|g_- - a\| = 1 + |\alpha|$  and  $d(a, F_{p, -\frac{\alpha}{|\alpha|}}) \leq 1 + |\alpha|$ . A proof is as follows. All what is really needed is to prove that  $|q(g_-)| \leq 1$  for every  $q \in P$  as it is evident that  $g_- \in E$  and  $p(g_-) = -\frac{\alpha}{|\alpha|}$ . If  $q \in F_0$ , then  $q(u) = 0$  asserts that  $|q(g_+)| = |q(a)| \leq 1$ . Suppose that  $q \in F_n$  for some positive integer  $n$ . We have

$$q(g_-) = \left( -\frac{\alpha}{|\alpha|} + \alpha \right) q(u) - 2\alpha q(u) + \alpha + q(a) - \alpha,$$

hence

$$|q(g_-)| \leq (1 - |\alpha|)|q(u)| + |\alpha||1 - 2q(u)| + |q(a) - \alpha|.$$

As  $0 \leq u \leq 1$  on  $Y$  by the definition of  $u$ , we have  $|1 - 2q(u)| \leq 1$  for every  $q \in P = \{\delta_x : x \in \text{Ch}(E)\}$ . As we have shown that  $|q(u)| \leq 1 - 1/2^n$  for

$q \in F_n$

$$\begin{aligned} |q(g_-)| &\leq (1 - |\alpha|)|q(u)| + |\alpha| + |q(a) - \alpha| \\ &\leq (1 - |\alpha|)(1 - 1/2^n) + |\alpha| + \varepsilon/2^{n+1} \\ &\leq (1 - |\alpha|)(1 - 1/2^n) + |\alpha| + (1 - |\alpha|)/2^{n+1} \leq 1. \end{aligned}$$

Suppose that  $q \in K$ . Then we have

$$q(g_-) = \left( -\frac{\alpha}{|\alpha|} + \alpha \right) q(u) - 2\alpha q(u) + \alpha.$$

As  $0 \leq u \leq 1$  on  $Y$  we have

$$|q(g_-)| \leq 1 - |\alpha| + |\alpha||1 - 2q(u)| \leq 1.$$

We conclude that  $g_- \in F_{p, -\frac{\alpha}{|\alpha|}}$ . By the definition of  $g_-$ , we infer that  $\|g_- - a\| = (1 + |\alpha|)\|u\| = 1 + |\alpha|$ . It follows that

$$(13) \quad d(a, F_{p, -\frac{\alpha}{|\alpha|}}) \leq 1 + |\alpha|.$$

By (12) and (13) we conclude that  $a \in M_{p, \alpha}$ .

Finally, as  $C_0(Y, \mathbb{R})$  satisfies the condition (r) by Urysohn's lemma, we see that  $C_0(Y, \mathbb{R})$  satisfies the Mazur-Ulam property.  $\square$

**Example 5.5.** Let  $\ell^\infty(\Gamma, \mathbb{R})$  be the real Banach space of all real-valued bounded functions on a discrete space  $\Gamma$ . Then there is a compact Hausdorff space  $X$  such that  $\ell^\infty(\Gamma, \mathbb{R})$  is isometrically isomorphic to  $C_0(X, \mathbb{R})$  as a real Banach space. Therefore  $\ell^\infty(\Gamma, \mathbb{R})$  satisfies the Mazur-Ulam property (cf. [10, 15]). The Mazur-Ulam property of the space  $c_0(\Gamma, \mathbb{R})$  is established by Ding [10, Corollary 2]. As  $c_0(\Gamma, \mathbb{R})$  is isometrically isomorphic to  $C_0(Y, \mathbb{R})$  for a locally compact Hausdorff space  $Y$ , Corollary 5.4 gives an alternative proof of the result of Ding.

**Example 5.6.** Let  $\mu$  be a positive measure on a  $\sigma$ -algebra  $\Sigma$  of a subsets of a set  $\Omega$ . Let  $L^\infty(\Omega, \Sigma, \mu, \mathbb{R})$  be the usual real Banach space of all real-valued bounded measurable functions on  $(\Omega, \Sigma, \mu)$ . Then there exists a compact Hausdorff space  $X$  such that  $L^\infty(\Omega, \Sigma, \mu, \mathbb{R})$  is isometrically isomorphic to  $C_0(X, \mathbb{R})$  as a real Banach space. Hence the space  $L^\infty(\Omega, \Sigma, \mu, \mathbb{R})$  satisfies the Mazur-Ulam property (cf. [15, Corollary 6]). Note that Tan [19, Theorem 2.5] exhibits the result for the case of the measure being  $\sigma$ -finite by an alternative proof.

## 6. REMARKS

We close the paper with a few remarks. In this paper we merely prove the complex Mazur-Ulam property for a uniform algebra. We conjecture that

a uniform algebra, generally a closed subalgebra of  $C_0(Y, \mathbb{C})$ , satisfies the Mazur-Ulam property.

The second remark concerns Tingley's problem on a Banach space of analytic functions. The main result of this paper concerns with the complex Mazur-Ulam property. We expect that several Banach space of analytic functions including the Hardy spaces satisfy the complex Mazur-Ulam property and the Mazur-Ulam property.

As a final remark we encourage researches on Tingley's problem on Banach algebras of continuous functions. Comparing with the theorem of Wang [27] on the Banach algebra of  $C^{(n)}(X)$ , it is interesting to study a surjective isometry on the unit sphere of a Banach space or algebra of Lipschitz functions, it has already been pointed out by Cueto-Avellaneda [6, Problem 4.0.8].

**Acknowledgments.** The author was supported by JSPS KAKENHI Grant Numbers JP19K03536.

#### REFERENCES

- [1] T. Banach, *Every 2-dimensional Banach space has the Mazur-Ulam property*, preprint
- [2] J. Becerra-Guerrero, M. Cueto-Avellaneda, F. J. Fernández-Polo and A. M. Peralta, *On the extensin of isometries between the unit spheres of a JBW\*-triple and a Banach space*, to appear in J. Inst. Math. Jussieu doi:10.1007/s13324-022-00448-2
- [3] A. Browder, *Introduction to function algebras*, W. A. Benjamin, Inc., New York-Amsterdam 1969 Xii+273 pp
- [4] J. Cabello Sánchez, *A reflection on Tingley's problem and some applications*, J. Math. Anal. Appl. **476** (2019), 319–336 doi:10.1016/j.jmaa.2019.03.041
- [5] L. Cheng and Y. Dong, *On a generalized Mazur-Ulam question: extension of isometries between unit spheres of Banach spaces*, J. Math. Anal. Appl. **377** (2011), 464–470 doi:10.1016/j.jmaa.2020.11.025
- [6] M. Cueto-Avellaneda, *Extension of isometreis and the Mazur-Ulam property*, PhD thesis, Universidad de Almería, 2020
- [7] M. Cueto-Avellaneda and A. M. Peralta, *On the Mazur-Ulam property for the space of Hilbert-space-valued continuous functions*, J. Math. Anal. Appl. **479**(2019), 875–902 doi:10.1016/j.jmaa.2019.06.056
- [8] M. Cueto-Avellaneda and A. M. Peralta, *The Mazur-Ulam property for commutative von Neumann algebras*, Linear Multilinear Algebra **68** (2020), 337–362 doi:10.1080/03081087.2018.1505823
- [9] G. G. Ding, *On extension of isometries between unit spheres of  $E$  and  $C(\Omega)$* , Acta Math. Sin. (Engl. Ser.) **19** (2003), 793–800
- [10] G. G. Ding, *The isometric extension of the into mapping from a  $\mathcal{L}^\infty(\Gamma)$ -type space to some Banach space*, Illinois J. Math. **51** (2007), 445–453
- [11] R. J. Fleming and J. E. Jamison, *Isometries on Banach spaces: function spaces*, Chapman & Hall/CRC Monographs and Surveys in Pure and Applied Mathematics, 129. Chapman & Hall/CRC, Boca Raton, FL, 2003. x+197 pp. ISBN: 1-58488-040-6



- [12] O. Hatori, S. Oi and R. Shindo Togashi, to appear in Jour. Math. Anal. Appl. doi: 10.1016/j.jmaa.2021.125346
- [13] A. Jiménez-Vargas, A. Morales-Campoy, A. M. Peralta and M. I. Ramírez, *The Mazur-Ulam property for the space of complex null sequences*, Linear Multilinear Algebra **67** (2019), 799–816 doi:10.1080/03081087.2018.1433625
- [14] O. F. K. Kalenda and A. M. Peralta, *Extension of isometries from the unit sphere of a rank-2 Cartan factor*, Anal. Math. Phys. **11**, Article number:15 (2021)
- [15] R. Liu, *On extension of isometries between unit spheres of  $\mathcal{L}^\infty(\Gamma)$ -type space and a Banach space  $E$* , J. Math. Anal. Appl. **333** (2007), 959–970 doi:10.1016/j.jmaa.2006.11.044
- [16] M. Mori, *Tingley's problem through the facial structure of operator algebras*, J. Math. Anal. Appl. **466** (2018), 1281–1298 doi:10.1016/j.jmaa.2018.06.050
- [17] M. Mori and N. Ozawa, *Mankiewicz's theorem and the Mazur-Ulam property for  $C^*$ -algebras*, Studia Math. **250** (2020), 265–281 doi:10.4064/sm180727-14-11
- [18] A. M. Peralta, *Extending surjective isometries defined on the unit sphere of  $\ell_\infty(\Gamma)$*  Rev. Mat. Complut. **32** (2019), 99–114 doi:10.1007/s13163-018-0269-2
- [19] D. N. Tan, *Extension of isometries on unit spheres of  $L^\infty$* , Taiwanese J. Math. **15** (2011), 819–827
- [20] D. N. Tan, *On extension of isometries on the unit spheres of  $L^p$ -spaces for  $0 < p \leq 1$* , Nonlinear Anal. **74** (2011), 6981–6987 doi:10.1016/j.na.2011.07.035
- [21] D. N. Tan, *Extension of isometries on the unit sphere of  $L^p$  spaces*, Acta Math. Sin. (Engl. Ser.) **28** (2012), 1197–1208 doi:10.1007/s10114-011-0302-6
- [22] D. Tan, X. Huang and R. Liu, *Generalized-lush spaces and the Mazur-Ulam property*, Studia Math. **219** (2013), 139–153 doi:10.4064/sm219-2-4
- [23] R. Tanaka, *A further property of spherical isometries*, Bull. Aust. Math. Soc. **90** (2014), 304–310 doi:10.1017/S0004972714000185
- [24] R. Tanaka, *The solution of Tingley's problem for the operator norm unit sphere of complex  $n \times n$  matrices*, Linear Algebra Appl. **494** (2016), 274–285 doi:10.1016/j.laa.2016.01.020
- [25] D. Tingley, *Isometries of the unit sphere*, Geom. Dedicata **22** (1987), 371–378
- [26] R. S. Wang, *Isometries between the unit spheres of  $C_0(\Omega)$  type spaces*, Acta Math. Sci. (English Ed.) **14** (1994), 82–89
- [27] Risheng Wang, *Isometries of  $C_0^{(n)}(X)$* , Hokkaido Math. J. **25** (1996), 465–519 doi:10.14492/hokmj/1351516747
- [28] Ruidong Wang and X. Huang, *The Mazur-Ulam property for two dimensional somewhere-flat spaces*, Linear Algebra Appl. **562** (2019), 55–62 doi:10.1016/j.laa.2018.09.024
- [29] X. Yang and X. Zhao, *On the extension problems of isometric and nonexpansive mappings*, In: Mathematics without boundaries. Edited by Themistocles M. Rassias and Panos M. Pardalos, 725–Springer, New York, 2014

INSTITUTE OF SCIENCE AND TECHNOLOGY, NIIGATA UNIVERSITY, NIIGATA 950-2181, JAPAN

Email address: hatori@math.sc.niigata-u.ac.jp