

VOLTERRA TYPE INTEGRATION OPERATORS BETWEEN WEIGHTED BERGMAN SPACES AND HARDY SPACES

YONGJIANG DUAN, SIYU WANG, AND ZIPENG WANG

ABSTRACT. Let \mathcal{D} be the class of radial weights on the unit disk which satisfy both forward and reverse doubling conditions. Let g be an analytic function on the unit disk \mathbb{D} . We characterize bounded and compact Volterra type integration operators

$$J_g(f)(z) = \int_0^z f(\lambda)g'(\lambda)d\lambda$$

between weighted Bergman spaces $L_a^p(\omega)$ induced by \mathcal{D} weights and Hardy spaces H^q for $0 < p, q < \infty$.

1. INTRODUCTION

A *weight* ω in this paper is a positive and integrable function on the unit disk \mathbb{D} . For $0 < p < \infty$, the *weighted Bergman space* $L_a^p(\omega)$ is a linear space of analytic functions on \mathbb{D} satisfying

$$\|f\|_{L^p(\omega)} := \left(\int_{\mathbb{D}} |f(z)|^p \omega(z) dA(z) \right)^{1/p} < \infty,$$

where $dA(z) = dx dy / \pi$ is the normalized area measure on \mathbb{D} . If $\omega(z) = \omega_\alpha(z) := (1 + \alpha)(1 - |z|^2)^\alpha$, $-1 < \alpha < \infty$, then $L_a^p(\omega_\alpha)$ is known as standard weighted Bergman space in literatures. Moreover, $L_a^p(\omega_0)$ is the classical Bergman space, and is denoted by L_a^p [Zhu07]. Recall for $0 < p < \infty$, the *Hardy space* consists of analytic functions f on \mathbb{D} such that

$$\|f\|_{H^p} := \sup_{0 \leq r < 1} \left(\frac{1}{2\pi} \int_0^{2\pi} |f(re^{i\theta})|^p d\theta \right)^{1/p} < \infty.$$

Let g be an analytic function on \mathbb{D} . We consider the *Volterra type integration operator* J_g with symbol g

$$J_g(f)(z) := \int_0^z f(\lambda)g'(\lambda)d\lambda, \quad z \in \mathbb{D},$$

where f is an analytic function on \mathbb{D} .

Pommerenke [Pom77] proved that J_g is bounded on Hardy space H^2 if and only if g is of bounded mean oscillation. The same characterization of the boundedness of J_g on the general Hardy space H^p was obtained by Aleman and Siskakis [AS95] for $p \geq 1$ and by Aleman and Cima [AC01] for $0 < p < 1$. In [AC01], Aleman and Cima further considered any pair of $(p, q) \in (0, \infty) \times (0, \infty)$ and studied the symbol g such that $J_g : H^p \rightarrow H^q$ is bounded. Analogous characterization of the symbol g for J_g to be bounded on Bergman spaces can be found in [Sis06]. Aleman and Siskakis [AS97] established the result on a large class of weighted Bergman spaces, which includes the class of standard weights ω_α as special examples. Pau and Peláez [PP10] described the $L_a^p(\omega)$ - $L_a^q(\omega)$ boundedness of J_g for $0 < p, q < \infty$ when ω is rapidly decreasing, while Peláez and Rättyä [PR14] got the corresponding characterizations when ω is regular or rapidly increasing weight.

Wu [Wu11] established boundedness criterions of J_g from $L_a^p(\omega_\alpha)$ to H^q for all $(p, q) \in (0, \infty) \times (0, \infty)$ but $0 < q < \min\{2, p\} < \infty$. Very recently, this gap was filled by Miihkinen, Pau, Perälä and Wang in [MPPW20] using analysis on tent spaces and some other delicate techniques. Moreover, by modifying the proofs of the boundedness results, one can get the corresponding compactness results ([Sis06, Wu11, CPW21]).

The goal of the present paper is to study Volterra type integration operators between weighted Bergman spaces $L_a^p(\omega)$ and Hardy spaces H^q for $0 < p, q < \infty$. Here the weight ω is a \mathcal{D} weight. Recall that the class of \mathcal{D} weights [PR19] consists of positive integrable radial functions ω on \mathbb{D} such that ω satisfies *both forward and reverse doubling conditions*, i.e., there exist constants $C_1, C_2 \in (1, \infty)$ and $C_3 \in [1, \infty)$,

$$C_1 \widehat{\omega} \left(1 - \frac{1-r}{C_2} \right) \leq \widehat{\omega}(r) \leq C_3 \widehat{\omega} \left(\frac{1+r}{2} \right), \quad r \in [0, 1),$$

where $\widehat{\omega}(z) = \int_{|z|}^1 \omega(r) dr$ for all $z \in \mathbb{D}$. The class of \mathcal{D} weights can be regarded as the largest class of radial weights ω in some sense that most of the results over $L_a^p(\omega_\alpha)$ can still hold on $L_a^p(\omega)$. For instance, the Littlewood-Paley estimate holds for weighted Bergman spaces induced by radial weights if and only if ω is a \mathcal{D} weight [PR19].

Let $H(\mathbb{D})$ be the set of analytic functions on \mathbb{D} . To describe our results, let us introduce two function spaces, which can be regarded as generalized Bloch spaces (little Bloch spaces). For a radial weight ω and $\gamma \in \mathbb{R}$, let

$$\mathcal{B}^{\gamma, \omega} = \left\{ f \in H(\mathbb{D}) : \sup_{z \in \mathbb{D}} \frac{(1-|z|^2)^\gamma}{\omega(z)} |f'(z)| < \infty \right\}$$

and

$$\mathcal{B}_0^{\gamma, \omega} = \left\{ f \in H(\mathbb{D}) : \lim_{|z| \rightarrow 1^-} \frac{(1-|z|^2)^\gamma}{\omega(z)} |f'(z)| = 0 \right\}.$$

Let \mathbb{T} be the unit circle. For an interval $I \subset \mathbb{T}$, we define the *Carleson square*

$$S(I) = \{z \in \mathbb{D} : z/|z| \in I, 1-|I| \leq |z| < 1\},$$

where $|\cdot|$ is the normalized arc measure on \mathbb{T} . For $0 < s < \infty$, a positive measure μ on \mathbb{D} is a *s-Carleson measure (vanishing s-Carleson measure)* [Zhu07, Section 9.2] if

$$\sup_{I \subseteq \mathbb{T}} \frac{\mu(S(I))}{|I|^s} < \infty \quad \left(\lim_{|I| \rightarrow 0} \frac{\mu(S(I))}{|I|^s} = 0 \right).$$

The (vanishing) 1-Carleson measure is called (vanishing) Carleson measure. Let $\xi \in \mathbb{T}$ and $0 < \alpha < \infty$, the *Stolz non-tangential approach region* is given by

$$\Gamma_\alpha(\xi) = \left\{ z \in \mathbb{D} : |z - \xi| < (\alpha + 1)(1 - |z|) \right\}.$$

For convenience, we denote $\Gamma_1(\xi)$ by $\Gamma(\xi)$. Let $0 < p < \infty$, $0 < q \leq \infty$ and ν be a positive Borel measure. The *tent space* $T_{q, \nu}^p$ consists of ν -equivalence classes of ν -measurable functions f on \mathbb{D} such that

$$\|f\|_{T_{q, \nu}^p}^p := \int_{\mathbb{T}} \left(\int_{\Gamma_\alpha(\xi)} |f(z)|^q d\nu(z) \right)^{\frac{p}{q}} d\xi < \infty, \quad q \in (0, \infty)$$

and

$$\|f\|_{T_{q,\nu}^p}^p := \int_{\mathbb{T}} \left(\operatorname{ess\,sup}_{z \in \Gamma_\alpha(\xi)} |f(z)| \right)^p d\xi < \infty, \quad q = \infty,$$

where the essential supremum is taken with respect to ν . Note that the tent spaces $T_{q,\nu}^p$ does not depend on the choice of Γ_α [MPPW20]. We denote $T_{q,\nu}^p$ by $T_{q,\omega,\beta}^p$, where $d\nu(z) = \left(\frac{(1-|z|^2)^2}{\omega(z)} \right)^\beta dA(z)$ and $0 < \beta < \infty$.

Our main results are the followings. The first two results characterize the boundedness and compactness of Volterra type integration operator J_g from weighted Bergman spaces to Hardy spaces.

Theorem 1.1. *Let $0 < p, q < \infty$, $\omega \in \mathcal{D}$ and $g \in H(\mathbb{D})$. The following statements hold:*

- (A): *If $0 < p \leq \min\{2, q\}$ or $2 < p < q < \infty$, then $J_g : L_a^p(\omega) \rightarrow H^q$ is bounded if and only if $g \in \mathcal{B}^{\gamma, \widehat{\omega}^{\frac{1}{p}}}$, where $\gamma = 1 + \frac{1}{q} - \frac{1}{p}$;*
- (B): *If $2 < p = q < \infty$, then $J_g : L_a^p(\omega) \rightarrow H^q$ is bounded if and only if the positive measure μ_g is a Carleson measure, where $d\mu_g(z) = |g'(z)|^{\frac{2p}{p-2}} \frac{(1-|z|^2)^{\frac{p+2}{p-2}}}{\widehat{\omega}(z)^{\frac{2}{p-2}}} dA(z)$;*
- (C): *If $0 < q < \infty$ and $p > \max\{2, q\}$, then $J_g : L_a^p(\omega) \rightarrow H^q$ is bounded if and only if $g' \in T_{\frac{2p}{p-2}, \widehat{\omega}, \frac{2}{p-2}}^{\frac{pq}{p-q}}$;*
- (D): *If $0 < q < p \leq 2$, then $J_g : L_a^p(\omega) \rightarrow H^q$ is bounded if and only if $B_{g,\omega} \in L^{\frac{pq}{p-q}}(\mathbb{T})$, where $B_{g,\omega}(\xi) = \sup_{z \in \Gamma(\xi)} |g'(z)| \frac{(1-|z|^2)^{\frac{1}{p}}}{\widehat{\omega}(z)^{\frac{1}{p}}}$, $\xi \in \mathbb{T}$.*

Theorem 1.2. *Let $0 < p, q < \infty$, $\omega \in \mathcal{D}$ and $g \in H(\mathbb{D})$. The following statements hold:*

- (A): *If $0 < p \leq \min\{2, q\}$ or $2 < p < q < \infty$, then $J_g : L_a^p(\omega) \rightarrow H^q$ is compact if and only if $g \in \mathcal{B}_0^{\gamma, \widehat{\omega}^{\frac{1}{p}}}$, where $\gamma = 1 + \frac{1}{q} - \frac{1}{p}$;*
- (B): *If $2 < p = q < \infty$, then $J_g : L_a^p(\omega) \rightarrow H^q$ is compact if and only if the positive measure μ_g is a vanishing Carleson measure, where $d\mu_g(z) = |g'(z)|^{\frac{2p}{p-2}} \frac{(1-|z|^2)^{\frac{p+2}{p-2}}}{\widehat{\omega}(z)^{\frac{2}{p-2}}} dA(z)$;*
- (C): *If $0 < q < \infty$ and $p > \max\{2, q\}$, then $J_g : L_a^p(\omega) \rightarrow H^q$ is compact if and only if $g' \in T_{\frac{2p}{p-2}, \widehat{\omega}, \frac{2}{p-2}}^{\frac{pq}{p-q}}$;*
- (D): *If $0 < q < p \leq 2$, then $J_g : L_a^p(\omega) \rightarrow H^q$ is compact if and only if*

$$\lim_{r \rightarrow 1^-} \int_{\mathbb{T}} \sup_{z \in \Gamma(\xi) \setminus r\mathbb{D}} |g'(z)|^{\frac{pq}{p-q}} \frac{(1-|z|^2)^{\frac{pq}{p-q}}}{\widehat{\omega}(z)^{\frac{q}{p-q}}} d\xi = 0.$$

The following result gives descriptions for the boundedness and compactness of J_g from Hardy spaces to weighted Bergman spaces. For a positive measure μ , define

$$\tilde{\mu}(\xi) = \int_{\Gamma(\xi)} \frac{d\mu(\lambda)}{1-|\lambda|^2}, \quad \xi \in \mathbb{T}.$$

Theorem 1.3. *Let $0 < p, q < \infty$. For $\omega \in \mathcal{D}$ and $g \in H(\mathbb{D})$, let $d\mu_g(z) = |g'(z)|^q (1-|z|^2)^q \omega(z) dA(z)$. Then*

- (A): *If $0 < p \leq q < \infty$, then $J_g : H^p \rightarrow L_a^q(\omega)$ is bounded (compact) if and only if μ_g is a q/p -Carleson measure (vanishing q/p -Carleson measure);*

(B): If $0 < q < p < \infty$, then $J_g : H^p \rightarrow L_a^q(\omega)$ is compact if and only if $J_g : H^p \rightarrow L_a^q(\omega)$ is bounded if and only if $\tilde{\mu}_g \in L^{\frac{p}{p-q}}(\mathbb{T})$.

Remark 1.1. Theorem 1.1-Theorem 1.3 generalize the recent work of [MPPW20, CPW21] from standard weights $\omega_\alpha(z) = (1 + \alpha)(1 - |z|^2)^\alpha dA(z)$, $-1 < \alpha < \infty$ to \mathcal{D} weights. The main strategies are based on a combination of the basic ideas in [Wu11, MPPW20, CPW21]. Nevertheless, there are two additional obstacles we need to overcome. The first one is the lack of explicit expressions for Bergman kernels of \mathcal{D} weighted Bergman spaces. In addition, the parameter α plays important roles in [MPPW20, CPW21] for standard weighted Bergman spaces. We use some delicate analysis of the weights to capture the suitable information from ω instead of α .

Remark 1.2. Since the proof of boundedness (Theorem 1.1) and compactness (Theorem 1.2) of J_g share several overlaps, we just present detailed proof of Theorem 1.2.

Throughout this paper, for a radial weight ω , we denote $\omega(E) = \int_E \omega(z) dA(z)$, for any measurable set $E \subseteq \mathbb{D}$. And we will use $a \lesssim b$ to represent that there exists a constant $C = C(\cdot) > 0$ satisfying $a \leq Cb$, $a \gtrsim b$ if there exists a constant $C = C(\cdot) > 0$ satisfying $a \geq Cb$, where the constant $C(\cdot)$ depends on the parameters indicated in the parenthesis, varying under different circumstances. Moreover, if $a \lesssim b$ and $b \gtrsim a$, then we denote by $a \asymp b$. Besides, for a number $p > 1$, let p' be the conjugate index of p such that $\frac{1}{p} + \frac{1}{p'} = 1$.

2. FROM BERGMAN SPACE INTO HARDY SPACE

The main goal in this section is to prove Theorem 1.2. The whole proof is divided into four parts.

2.1. Case A. First we are devoted to dealing with the case when $0 < p \leq \min\{2, q\}$ or $2 < p < q < \infty$.

Proposition 2.1. Assume $0 < p \leq \min\{2, q\}$ or $2 < p < q < \infty$. Let ω be a \mathcal{D} weight and $\gamma = 1 + \frac{1}{q} - \frac{1}{p}$. Then the Volterra type integration operator $J_g : L_a^p(\omega) \rightarrow H^q$ is compact if and only if $g \in \mathcal{B}_0^{\gamma, \hat{\omega}^{1/p}}$, where $\hat{\omega}(z) = \int_{|z|}^1 \omega(r) dr$.

To show Proposition 2.1, we first give some useful lemmas. By a disk version of Calderón's theorem, one has the following:

Lemma 2.2. [Pav13, Theorem 1.3] Suppose f is analytic on \mathbb{D} with $f(0) = 0$. If $0 < p < \infty$, then

$$\|f\|_{H^p}^p \asymp \int_{\mathbb{T}} \left(\int_{\Gamma(\xi)} |f'(z)|^2 dA(z) \right)^{\frac{p}{2}} d\xi.$$

The following Dirichlet-type embedding theorem can be found in [MPPW20].

Lemma 2.3. Suppose f is analytic on \mathbb{D} with $f(0) = 0$. If $0 < q \leq 2$, then

$$(2.1) \quad \|f\|_{H^q} \lesssim \|f'\|_{L_a^q(\omega_{q-1})}.$$

If $0 < p < q < \infty$, then

$$(2.2) \quad \|f\|_{H^q} \lesssim \|f'\|_{L_a^p(\omega_{p+\frac{p}{q}-2})}.$$

Now we recall two classes of weights which have close relations with \mathcal{D} weights. A weight ω is called *regular* [PR14] if ω is continuous and radial on \mathbb{D} such that $\psi_\omega(r) \asymp (1-r)$ for $0 \leq r < 1$, where $\psi_\omega(r) = \frac{\widehat{\omega}(r)}{\omega(r)}$ is the distortion function, which was introduced in [Sis00]. We denote the class of regular weights by \mathcal{R} . And a weight ω is a $\widehat{\mathcal{D}}$ weight [PRS18] if there exists a positive constant C such that

$$\widehat{\omega}(r) \leq C\widehat{\omega}\left(\frac{1+r}{2}\right), \quad \text{for all } r \in [0, 1).$$

Note that $\mathcal{R} \subsetneq \mathcal{D} \subsetneq \widehat{\mathcal{D}}$ [PR14, PRS18, PR19]. Here we list some useful properties of $\widehat{\mathcal{D}}$, \mathcal{D} and regular weights which are important in our subsequent work.

Lemma 2.4. [PRS18, Theorem C] *Let $1 < p < \infty$ and ω be a $\widehat{\mathcal{D}}$ weight. Then the reproducing kernel of $L_a^2(\omega)$ at z has the following estimates,*

$$\|K_z^\omega\|_{L_a^2(\omega)}^p \asymp \int_{|z|}^1 \frac{dt}{\widehat{\omega}(t)^{p-1}(1-t)^p}, \quad |z| \rightarrow 1^-.$$

From Lemma 2.4, we get

$$(2.3) \quad \|K_z^\omega\|_{L_a^2(\omega)}^p \asymp \frac{1}{\widehat{\omega}(z)^{p-1}(1-|z|)^{p-1}}, \quad z \in \mathbb{D}.$$

By [PRS18, Proposition 5] and its proof, we have

Lemma 2.5. *Let $0 < p < \infty$ and ω be a \mathcal{D} weight. Then $\kappa(z) = (1-|z|)^{-1}\widehat{\omega}(z)$ is regular and $\|f\|_{L^p(\omega)} \asymp \|f\|_{L^p(\kappa)}$, for each analytic function f on \mathbb{D} .*

Denote by $B(z, r)$ and $D(z, r)$ the Euclidean disk and the pseudo-hyperbolic disk centered at z with radius r respectively. The next result shows a nice property of regular weights, which can be found in page 8 of [PR14].

Lemma 2.6. *Let ω be a regular weight. Then for any $z \in \mathbb{D}$ and $r \in (0, 1)$,*

$$\omega(z) \asymp \omega(u), \quad \text{for all } u \in D(z, r).$$

Combine Lemma 2.5 with Lemma 2.6, we get the following result.

Lemma 2.7. *Let $\omega \in \mathcal{D}$ and $r \in (0, 1)$. Then for any $z \in \mathbb{D}$,*

$$\widehat{\omega}(z) \asymp \widehat{\omega}(u), \quad \text{for all } u \in D(z, r).$$

Now we are ready to prove Proposition 2.1.

Proof of Proposition 2.1. Sufficiency. Let $\{f_j\}_{j=1}^\infty$ be any sequence in $L_a^p(\omega)$ such that $\|f_j\|_{L_a^p(\omega)} \leq 1$ and f_j uniformly converges to 0 on any compact subset of \mathbb{D} . If $g \in \mathcal{B}_0^{\gamma, \widehat{\omega}^{1/p}}$, then we will show

$$\lim_{j \rightarrow \infty} \|J_g(f_j)\|_{H^q} = 0.$$

For any $\varepsilon > 0$, there exists $r \in (0, 1)$ such that for any $z \in \mathbb{D}$ with $r \leq |z| < 1$, it holds that

$$(2.4) \quad \frac{(1-|z|^2)^\gamma}{\widehat{\omega}(z)^{1/p}} |g'(z)| < \varepsilon.$$

Moreover, we can find a constant $k \in \mathbb{N}$ such that for any $j > k$, $|f_j| < \varepsilon$ on $r\mathbb{D} := \{z : |z| < r\}$. By Lemma 2.3, for any $j \in \mathbb{N}$, we obtain

$$\|J_g(f_j)\|_{H^q} \lesssim \|(J_g f_j)'\|_{L_a^p(\omega_{p\gamma-1})}.$$

Together with Lemma 2.5 and (2.4), for any $j > k$, we get

$$\begin{aligned} \|J_g(f_j)\|_{H^q} &\lesssim \left(\int_{r\mathbb{D}} |f_j(z)|^p \frac{\widehat{\omega}(z)}{1-|z|} |g'(z)|^p \frac{(1-|z|^2)^{p\gamma}}{\widehat{\omega}(z)} dA(z) \right)^{\frac{1}{p}} \\ &\quad + \left(\int_{\mathbb{D} \setminus r\mathbb{D}} |f_j(z)|^p \frac{\widehat{\omega}(z)}{1-|z|} |g'(z)|^p \frac{(1-|z|^2)^{p\gamma}}{\widehat{\omega}(z)} dA(z) \right)^{\frac{1}{p}} \\ &\lesssim \varepsilon + \varepsilon \|f_j\|_{L_a^p(\omega)} \lesssim \varepsilon. \end{aligned}$$

Necessity. Assume that $J_g : L_a^p(\omega) \rightarrow H^q$ is compact. Let $\alpha > \frac{1}{p}$ and $\lambda \in \mathbb{D}$. Take testing functions

$$f_\lambda(z) = \frac{(K_\lambda^\omega(z))^\alpha}{\|K_\lambda^\omega\|_{L_a^{\alpha+1-\frac{1}{p}}(\omega)}^{\alpha+1-\frac{1}{p}}}, \quad z \in \mathbb{D}.$$

Let $\omega_{2n+1} = \int_0^1 \omega(r) r^{2n+1} dr$, $n \in \mathbb{N} \cup \{0\}$. It follows from [PR16, formula (20)] that

$$|K_\lambda^\omega(z)| \leq \sum_{n=0}^{\infty} \frac{|z|^n}{2\omega_{2n+1}} \asymp \frac{1}{\omega_1} + \int_0^{|z|} \frac{1}{\widehat{\omega}(t)(1-t)^2} dt \lesssim \frac{1}{\widehat{\omega}(z)(1-|z|)}, \quad |z| \rightarrow 1^-.$$

Together with (2.3), we obtain

$$|f_\lambda(z)| \asymp \widehat{\omega}(\lambda)^{\alpha-\frac{1}{p}} (1-|\lambda|)^{\alpha-\frac{1}{p}} |K_\lambda^\omega(z)|^\alpha \lesssim \frac{\widehat{\omega}(\lambda)^{\alpha-\frac{1}{p}} (1-|\lambda|)^{\alpha-\frac{1}{p}}}{\widehat{\omega}(z)^\alpha (1-|z|)^\alpha}, \quad z \in \mathbb{D}.$$

Then for any compact subset $E \subseteq \mathbb{D}$, we have

$$\lim_{|\lambda| \rightarrow 1^-} \sup_{z \in E} |f_\lambda(z)| = 0.$$

Moreover, from (2.3), we know

$$\|f_\lambda\|_{L_a^p(\omega)} \asymp \widehat{\omega}(\lambda)^{\alpha-\frac{1}{p}} (1-|\lambda|)^{\alpha-\frac{1}{p}} \left(\int_{\mathbb{D}} |K_\lambda^\omega(z)|^{\alpha p} \omega(z) dA(z) \right)^{\frac{1}{p}} \asymp 1.$$

Therefore, we deduce

$$\lim_{|\lambda| \rightarrow 1^-} \|J_g(f_\lambda)\|_{H^q} = 0.$$

For any $z, \lambda \in \mathbb{D}$, by using a classical result of the pointwise estimate for $(J_g f_\lambda)'$ (see [Gar07, Page 85-86]), we get

$$(2.5) \quad |f_\lambda(z)g'(z)| = |(J_g f_\lambda)'(z)| \lesssim \frac{1}{(1-|z|^2)^{1+\frac{1}{q}}} \|J_g f_\lambda\|_{H^q}.$$

By choosing $z = \lambda$ in (2.5), together with (2.3), we deduce

$$\frac{(1-|\lambda|^2)^\gamma}{\widehat{\omega}(\lambda)^{\frac{1}{p}}} |g'(\lambda)| \lesssim \|J_g f_\lambda\|_{H^q},$$

from which it follows that

$$\lim_{|\lambda| \rightarrow 1^-} \frac{(1 - |\lambda|^2)^\gamma}{\widehat{\omega}(\lambda)^{\frac{1}{p}}} |g'(\lambda)| = 0.$$

Thus, we complete the proof. \square

2.2. Case B. Now we turn to deal with the case $2 < p = q < \infty$.

For $k \in \mathbb{N}$, recall the Rademacher functions $r_k(t) := \text{sgn}\{\sin(2^k \pi t)\}$, $t \in [0, 1]$. The following Khinchine's inequality and Kahane's inequality [Lue93] are useful in our subsequent work.

Lemma 2.8. *Let $0 < p < \infty$. Then for any sequence of complex numbers $\{c_k\}_{k=1}^\infty$, it holds that*

$$\left(\sum_{k=1}^\infty |c_k|^2 \right)^{\frac{p}{2}} \asymp \int_0^1 \left| \sum_{k=1}^\infty c_k r_k(t) \right|^p dt.$$

Lemma 2.9. *Let $0 < p, q < \infty$ and X be a quasi-Banach space with quasi-norm $\|\cdot\|_X$. For any sequence $\{x_k\}_{k=1}^\infty \subseteq X$, it holds that*

$$\left(\int_0^1 \left\| \sum_{k=1}^\infty r_k(t) x_k \right\|_X^q dt \right)^{\frac{1}{q}} \asymp \left(\int_0^1 \left\| \sum_{k=1}^\infty r_k(t) x_k \right\|_X^p dt \right)^{\frac{1}{p}}.$$

We first generalize Theorem 1 in [Wu11] to \mathcal{D} weighted Bergman spaces.

Proposition 2.10. *Let $\omega \in \mathcal{D}$, $\kappa(z) = \frac{\widehat{\omega}(z)}{1-|z|}$ and μ be a positive measure on \mathbb{D} . Let $p \in (0, +\infty)$, $s \in [1, +\infty)$ and $\iota \in (0, 1)$. Then the following statements are equivalent:*

- (i) $\left(\frac{\mu(D(z, \delta))}{\kappa(D(z, \delta))} \right)^{\frac{1}{1-\iota}} \kappa(z) dA(z)$ is a vanishing s -Carleson measure, for some $\tilde{\delta} \in (0, 1)$;
- (ii) For any $f \in L_a^p(\omega)$, $|f(z)|^{p\iota} d\mu(z)$ is a vanishing $s(1-\iota)$ -Carleson measure, that is, for any $f \in L_a^p(\omega)$ and any $\varepsilon > 0$, there exists $r_0 \in (0, 1)$ such that for any $I \subseteq \mathbb{T}$ with $0 < |I| \leq r_0$, it holds that

$$\frac{\int_{S(I)} |f(z)|^{p\iota} d\mu(z)}{|I|^{s(1-\iota)}} \lesssim \varepsilon \|f\|_{L_a^p(\omega)}^{p\iota},$$

where the constant of comparison does not depend on the choice of the function f .

Before approaching the proof of Proposition 2.10, we first give some notations. For $0 < \alpha < \infty$ and $\xi \in \mathbb{T}$, we define the arc on \mathbb{T} as follows,

$$I_\alpha(z) = \left\{ \xi \in \mathbb{T} : z \in \Gamma_\alpha(\xi) \right\}, \quad z \in \mathbb{D}.$$

Note that

$$(2.6) \quad |I_\alpha(z)| \asymp 1 - |z|, \quad z \in \mathbb{D},$$

see [Wu11, formula (3.3)]. We denote $I_\alpha(z)$ by $I(z)$ if $\alpha = 1$. For $I \subseteq \mathbb{T}$, denote by the α -tent of I

$$\wedge_\alpha(I) = \left\{ z \in \mathbb{D} : I_\alpha(z) \subseteq I \right\}.$$

First observe the relationship between the Carleson square and the α -tent of an arc on \mathbb{T} . For any fixed $\alpha > 0$, for any $I \subseteq \mathbb{T}$, there exist $I_1, I_2 \subseteq \mathbb{T}$ such that $S(I_1) \subseteq \wedge_\alpha(I) \subseteq S(I_2)$

and $|I_1| \asymp |I| \asymp |I_2|$. This makes us to choose $S(I)$ or $\wedge_\alpha(I)$ freely in the definition of vanishing Carleson measure.

Next we recall some important observations over geometrical regions by Wu in [Wu11].

Lemma 2.11. [Wu11, Lemma 2.2 and Lemma 3.1] *Let $\alpha > 0$, $\xi \in \mathbb{T}$ and $I \subseteq \mathbb{T}$. Let $\tilde{t} = \tanh t$ and $\alpha_t = (\alpha + 2)e^{2t} - 2$ for $t \geq 0$. Then*

$$D(z, \tilde{t}) \subseteq \Gamma_{\alpha_t}(\xi), \quad \text{for any } z \in \Gamma_\alpha(\xi),$$

$$D(z, \tilde{t}) \subseteq \wedge_\alpha(I), \quad \text{for } z \in \wedge_{\alpha_t}(I)$$

and

$$D(z, \tilde{t}) \subseteq \wedge_\alpha(I), \quad \text{for } D(z, \tilde{t}) \cap \wedge_{\alpha_{2t}}(I) \neq \emptyset.$$

Let $\{z_j\}_{j=1}^\infty$ be a sequence in \mathbb{D} , $\delta > 0$ and $r \in (0, 1)$. If $\left| \frac{z_j - z_k}{1 - \bar{z}_j z_k} \right| \geq \delta$ for any $j \neq k$, then we say the sequence $\{z_j\}_{j=1}^\infty$ is δ -separated. And if $\{z_j\}_{j=1}^\infty$ satisfies $\mathbb{D} = \bigcup_{j=1}^\infty D(z_j, r)$ and $\frac{r}{5}$ -separated, we say $\{z_j\}_{j=1}^\infty$ is an r -lattice. Notice that every $z \in \mathbb{D}$ belongs to at most $N = N(r)$ pseudohyperbolic disks $D(z_j, r)$, where $\{z_j\}_{j=1}^\infty$ is a separated sequence (see Lemma 12 of Chapter 2 in [Dur04] or Lemma 3 in [Lue93]).

Let $\{z_k\}$ be a separated sequence. For any $K \in (1, +\infty)$, we re-index the sequence $\{z_k\}$ as follows: For each $j \in \mathbb{N} \cup \{0\}$, we denote the points of the sequence $\{z_k\}$ in the annulus $A_j = \{z \in \mathbb{D} : r_j \leq |z| < r_{j+1}\}$ by $\{z_{j,l}\}_l$, where $r_j = 1 - K^{-j}$. For $p \in (0, \infty)$, note that a discrete sequence $\{\lambda_{j,l}\}_{j,l=1}^\infty \in l^p$ if

$$\left\| \{\lambda_{j,l}\}_{j,l=1}^\infty \right\|_{l^p} := \left(\sum_{j=1}^\infty \sum_{l=1}^\infty |\lambda_{j,l}|^p \right)^{\frac{1}{p}} < \infty.$$

The following (see [PRS21, Theorem 1]) is the atomic decomposition for Bergman spaces induced by \mathcal{D} weights. One can consult [PRS21] for more information.

Lemma 2.12. *Let $0 < p < \infty$, $1 < K < \infty$, $\omega \in \mathcal{D}$ and $\{z_k\}$ be a separated sequence in \mathbb{D} . Denote by $\{z_{j,l}\}_{j,l}$ the sequence re-indexed from $\{z_k\}$ using the method above. Then there exists a positive constant M (M only depends on the choice of ω and p) such that for any sequence $\{\lambda_{j,l}\}_{j,l=1}^\infty \in l^p$, the function*

$$F(z) := \sum_{j,l=1}^\infty \lambda_{j,l} \frac{(1 - |z_{j,l}|)^{M - \frac{1}{p}} \widehat{\omega}(z_{j,l})^{-\frac{1}{p}}}{(1 - \bar{z}_{j,l} z)^M}$$

is analytic in \mathbb{D} and

$$\|F\|_{L_a^p(\omega)} \lesssim \|\{\lambda_{j,l}\}_{j,l=1}^\infty\|_{l^p}.$$

Now we are in a position to prove Proposition 2.10.

Proof of Proposition 2.10. First we prove that (i) \Rightarrow (ii). Choose $\delta \in (0, \infty)$ such that $\tilde{\delta} := \tanh \delta$ satisfies the condition in (i), it follows from Lemma 2.5 that $\kappa(z) = \frac{\widehat{\omega}(z)}{1 - |z|}$ is a regular weight. By Lemma 2.6, we know that

$$(2.7) \quad \kappa(z) \asymp \kappa(\zeta), \quad \zeta \in D(z, \tilde{\delta}).$$

Let $0 < p < \infty$, $1 \leq s < \infty$, $0 < \iota < 1$ and $\alpha > 0$. Let $\alpha_\delta = (\alpha + 2)e^{2\delta} - 2$. For any $f \in L_a^p(\omega)$ and $I \subseteq \mathbb{T}$, using the subharmonicity of $|f|^p$, Lemma 2.11 and (2.7), we obtain

$$\begin{aligned}
|f(z)|^{p\iota} &\lesssim \frac{1}{(1-|z|)^2} \int_{B(z, \tilde{\delta}(1-|z|))} |f(\zeta)|^{p\iota} dA(\zeta) \\
&\lesssim \frac{1}{\kappa(z)(1-|z|)^2} \int_{D(z, \tilde{\delta})} |f(\zeta)|^{p\iota} \kappa(\zeta) dA(\zeta) \\
(2.8) \quad &= \frac{1}{\kappa(z)(1-|z|)^2} \int_{\wedge_\alpha(I) \cap D(z, \tilde{\delta})} |f(\zeta)|^{p\iota} \kappa(\zeta) dA(\zeta), \quad \text{for any } z \in \wedge_{\alpha_\delta}(I).
\end{aligned}$$

Combining Lemma 2.5, Fubini's theorem, Hölder's inequality, (2.7) with (2.8), for any $I \subseteq \mathbb{T}$, we get

$$\begin{aligned}
\int_{\wedge_{\alpha_\delta}(I)} |f(z)|^{p\iota} d\mu(z) &\lesssim \int_{\wedge_{\alpha_\delta}(I)} \frac{1}{\kappa(z)(1-|z|)^2} \int_{\wedge_\alpha(I) \cap D(z, \tilde{\delta})} |f(\zeta)|^{p\iota} \kappa(\zeta) dA(\zeta) d\mu(z) \\
&\asymp \int_{\wedge_\alpha(I)} \frac{1}{\kappa(\zeta)(1-|\zeta|)^2} \int_{\wedge_{\alpha_\delta}(I) \cap D(\zeta, \tilde{\delta})} d\mu(z) |f(\zeta)|^{p\iota} \kappa(\zeta) dA(\zeta) \\
&\lesssim \int_{\wedge_\alpha(I)} \frac{\mu(D(\zeta, \tilde{\delta}))}{\kappa(D(\zeta, \tilde{\delta}))} |f(\zeta)|^{p\iota} \kappa(\zeta) dA(\zeta) \\
(2.9) \quad &\lesssim \|f\|_{L_a^p(\omega)}^{p\iota} \left(\int_{\wedge_\alpha(I)} \left(\frac{\mu(D(\zeta, \tilde{\delta}))}{\kappa(D(\zeta, \tilde{\delta}))} \right)^{\frac{1}{1-\iota}} \kappa(\zeta) dA(\zeta) \right)^{1-\iota}.
\end{aligned}$$

Since $\left(\frac{\mu(D(\zeta, \tilde{\delta}))}{\kappa(D(\zeta, \tilde{\delta}))} \right)^{\frac{1}{1-\iota}} \kappa(\zeta) dA(\zeta)$ is a vanishing s -Carleson measure, then for any $\varepsilon > 0$, there exists $r_1 \in (0, 1)$ such that for any $I \subseteq \mathbb{T}$ with $0 < |I| \leq r_1$, it holds that

$$\frac{\int_{\wedge_\alpha(I)} \left(\frac{\mu(D(\zeta, \tilde{\delta}))}{\kappa(D(\zeta, \tilde{\delta}))} \right)^{\frac{1}{1-\iota}} \kappa(\zeta) dA(\zeta)}{|I|^s} < \varepsilon^{\frac{1}{1-\iota}}.$$

Combining with (2.9), there exists a constant C (C does not depend on the choice of f) such that for any $I \subseteq \mathbb{T}$ with $0 < |I| \leq r_1$, for any $f \in L_a^p(\omega)$, we have

$$\frac{\int_{\wedge_{\alpha_\delta}(I)} |f(z)|^{p\iota} d\mu(z)}{|I|^{s(1-\iota)}} \leq \varepsilon C \|f\|_{L_a^p(\omega)}^{p\iota},$$

as desired.

Next we show (ii) \Rightarrow (i). By the hypothesis, for any $f \in L_a^p(\omega)$ and $\varepsilon > 0$, there exists $r_2 \in (0, 1)$ such that for any $I \subseteq \mathbb{T}$ with $0 < |I| \leq r_2$, it holds that

$$\frac{\int_{\wedge_\alpha(I)} |f(z)|^{p\iota} d\mu(z)}{|I|^{s(1-\iota)}} \lesssim \varepsilon \|f\|_{L_a^p(\omega)}^{p\iota}.$$

Take $\delta > 0$ such that $\tilde{\delta} = \tanh \delta \in (0, \frac{1}{2})$. Let $\{z_k\}_{k=1}^\infty$ be a $\tilde{\delta}$ -lattice. Take the constant M and denote by $\{z_{j,l}\}_{j,l=1}^\infty$ the sequence re-indexed from $\{z_k\}_{k=1}^\infty$ as in Lemma 2.12. Hence, by Lemma 2.12, for any $\{\lambda_{j,l}\}_{j,l=1}^\infty \in l^p$, for any $I \subseteq \mathbb{T}$ with $0 < |I| \leq r_2$, it holds that

$$\frac{\int_{\wedge_\alpha(I)} \left| \sum_{j,l=1}^\infty \lambda_{j,l} f_{j,l}(z) \right|^{p\iota} d\mu(z)}{|I|^{s(1-\iota)}} \lesssim \varepsilon \left\| \sum_{j,l=1}^\infty \lambda_{j,l} f_{j,l}(z) \right\|_{L_a^p(\omega)}^{p\iota} \lesssim \varepsilon \|\{\lambda_{j,l}\}_{j,l=1}^\infty\|_{l^p}^{p\iota},$$

where

$$f_{j,l}(z) = \frac{(1 - |z_{j,l}|)^{M - \frac{1}{p}} \widehat{\omega}(z_{j,l})^{-\frac{1}{p}}}{(1 - \overline{z_{j,l}}z)^M}, \quad z \in \mathbb{D}.$$

For $t \in (0, 1)$, replace $\lambda_{j,l}$ by $\lambda_{j,l} r_{j,l}(t)$, where $r_{j,l}(t)$ is the Rademacher function. Then for any $I \subseteq \mathbb{T}$ with $0 < |I| \leq r_2$, we have

$$\frac{\int_{\wedge_\alpha(I)} \left| \sum_{j,l=1}^{\infty} \lambda_{j,l} r_{j,l}(t) f_{j,l}(z) \right|^{p_\nu} d\mu(z)}{|I|^{s(1-\nu)}} \lesssim \varepsilon \|\{\lambda_{j,l}\}_{j,l=1}^{\infty}\|_{l^p}^{p_\nu}.$$

Integrate both sides of the above inequality over $(0, 1)$ with respect to t , we obtain

$$\frac{\int_{\wedge_\alpha(I)} \int_0^1 \left| \sum_{j,l=1}^{\infty} \lambda_{j,l} r_{j,l}(t) f_{j,l}(z) \right|^{p_\nu} dt d\mu(z)}{|I|^{s(1-\nu)}} \lesssim \varepsilon \|\{\lambda_{j,l}\}_{j,l=1}^{\infty}\|_{l^p}^{p_\nu}.$$

Combining with Lemma 2.8, we get

$$(2.10) \quad \int_{\wedge_\alpha(I)} \left(\sum_{j,l=1}^{\infty} |\lambda_{j,l}|^2 |f_{j,l}(z)|^2 \right)^{\frac{p_\nu}{2}} d\mu(z) \lesssim \varepsilon |I|^{s(1-\nu)} \|\{\lambda_{j,l}\}_{j,l=1}^{\infty}\|_{l^p}^{p_\nu},$$

for $I \subseteq \mathbb{T}$ with $0 < |I| \leq r_2$. It follows from (2.10) that there exists a constant $N = N(\delta)$ such that

$$(2.11) \quad \begin{aligned} \sum_{\{j,l:D(z_{j,l},2\tilde{\delta}) \subseteq \wedge_\alpha(I)\}} |\lambda_{j,l}|^{p_\nu} \frac{\mu(D(z_{j,l},2\tilde{\delta}))}{\widehat{\omega}(z_{j,l})^\nu (1 - |z_{j,l}|)^\nu} &\asymp \sum_{\{j,l:D(z_{j,l},2\tilde{\delta}) \subseteq \wedge_\alpha(I)\}} \int_{D(z_{j,l},2\tilde{\delta})} (|\lambda_{j,l}| \cdot |f_{j,l}(z)|)^{p_\nu} d\mu(z) \\ &\leq \sum_{\{j,l:D(z_{j,l},2\tilde{\delta}) \subseteq \wedge_\alpha(I)\}} \int_{D(z_{j,l},2\tilde{\delta})} \left(\sum_{j,l=1}^{\infty} |\lambda_{j,l}|^2 \cdot |f_{j,l}(z)|^2 \right)^{\frac{p_\nu}{2}} d\mu(z) \\ &\leq N \int_{\wedge_\alpha(I)} \left(\sum_{j,l=1}^{\infty} |\lambda_{j,l}|^2 \cdot |f_{j,l}(z)|^2 \right)^{\frac{p_\nu}{2}} d\mu(z) \\ &\lesssim \varepsilon N |I|^{s(1-\nu)} \|\{\lambda_{j,l}\}_{j,l=1}^{\infty}\|_{l^p}^{p_\nu}, \quad \text{for } I \subseteq \mathbb{T} \text{ with } 0 < |I| \leq r_2. \end{aligned}$$

Since the dual space of $l^{\frac{1}{\nu}}$ can be identified with $l^{\frac{1}{1-\nu}}$, together with (2.11), we deduce

$$(2.12) \quad \left[\sum_{\{j,l:D(z_{j,l},2\tilde{\delta}) \subseteq \wedge_\alpha(I)\}} \left(\frac{\mu(D(z_{j,l},2\tilde{\delta}))}{\widehat{\omega}(z_{j,l})^\nu (1 - |z_{j,l}|)^\nu} \right)^{\frac{1}{1-\nu}} \right]^{1-\nu} \lesssim \varepsilon |I|^{s(1-\nu)}, \quad \text{for } I \subseteq \mathbb{T} \text{ with } 0 < |I| \leq r_2.$$

Take $t = 2 \tanh^{-1}(2\tilde{\delta})$. Let $\alpha_t = (\alpha + 2)e^{2t} - 2$. By Lemma 2.7, Lemma 2.11 and (2.12), for any $I \subseteq \mathbb{T}$ with $0 < |I| \leq r_2$, we get

$$\begin{aligned} \int_{\wedge_{\alpha_t}(I)} \left(\frac{\mu(D(z,\tilde{\delta}))}{\kappa(D(z,\tilde{\delta}))} \right)^{\frac{1}{1-\nu}} \kappa(z) dA(z) &\leq \sum_{\{j,l:D(z_{j,l},\tilde{\delta}) \cap \wedge_{\alpha_t}(I) \neq \emptyset\}} \int_{D(z_{j,l},\tilde{\delta})} \left(\frac{\mu(D(z,\tilde{\delta}))}{\kappa(D(z,\tilde{\delta}))} \right)^{\frac{1}{1-\nu}} \kappa(z) dA(z) \\ &\leq \sum_{\{j,l:D(z_{j,l},2\tilde{\delta}) \subseteq \wedge_\alpha(I)\}} \int_{D(z_{j,l},\tilde{\delta})} \left(\frac{\mu(D(z,\tilde{\delta}))}{\kappa(D(z,\tilde{\delta}))} \right)^{\frac{1}{1-\nu}} \kappa(z) dA(z) \\ &\lesssim \varepsilon^{\frac{1}{1-\nu}} |I|^s, \end{aligned}$$

which yields that $\left(\frac{\mu(D(z,\delta))}{\kappa(D(z,\delta))}\right)^{\frac{1}{1-\alpha}} \kappa(z) dA(z)$ is a vanishing s -Carleson measure, as desired. \square

For a positive measure μ on \mathbb{D} and $\alpha > 0$, define the area integral operator $A_{\mu,\alpha}^2$ acting on $f \in H(\mathbb{D})$ by

$$A_{\mu,\alpha}^2(f)(\xi) = \left(\int_{\Gamma_\alpha(\xi)} |f(z)|^2 \frac{d\mu(z)}{1-|z|} \right)^{\frac{1}{2}}, \quad \xi \in \mathbb{T}.$$

For a \mathcal{D} weight ω , $0 < q, \alpha < \infty$ and $g \in H(\mathbb{D})$, it follows from Lemma 2.2 that the Volterra type integration operator is intimately related to area integral operator as follows,

$$(2.13) \quad \|J_g(f)\|_{H^q} \asymp \|A_{\mu_g,\alpha}^2(f)\|_{L^q(\mathbb{T})}, \quad \text{for any } f \in L_a^p(\omega),$$

where $d\mu_g(z) = |g'(z)|^2(1-|z|)dA(z)$.

Proposition 2.13. *Let $2 < p < \infty$, $\omega \in \mathcal{D}$, $g \in H(\mathbb{D})$ and*

$$d\mu_g(z) = |g'(z)|^{\frac{2p}{p-2}} \frac{(1-|z|^2)^{\frac{p+2}{p-2}}}{\widehat{\omega}(z)^{\frac{2}{p-2}}} dA(z).$$

Then the Volterra type integration operator $J_g : L_a^p(\omega) \rightarrow H^p$ is compact if and only if μ_g is a vanishing Carleson measure.

Proof. Sufficiency. Let $\{f_j\}_{j=1}^\infty$ be any sequence in $L_a^p(\omega)$ with $\|f_j\|_{L_a^p(\omega)} \leq 1$ and f_j converges to 0 uniformly on any compact subset of \mathbb{D} . Since μ_g is a vanishing Carleson measure, for any $\varepsilon > 0$, there exists $r_0 \in (0, 1)$ such that for any $I \subseteq \mathbb{T}$ with $|I| \leq 1 - r_0$, it holds that

$$(2.14) \quad \frac{\mu_g(S(I))}{|I|} < \varepsilon^{\frac{2p}{p-2}}.$$

Moreover, there also exists $k_0 \in \mathbb{N}$ such that $j > k_0$, $|f_j| < \varepsilon$ on $r_0\mathbb{D}$. Let

$$d\mu_{f_j,g}(z) = |f_j(z)|^2 |g'(z)|^2 (1-|z|^2) dA(z).$$

By virtue of (2.14), we obtain

$$(2.15) \quad \frac{\mu_g(S(I) \setminus r_0\mathbb{D})}{|I|} < \varepsilon^{\frac{2p}{p-2}}, \quad \text{for any } I \subseteq \mathbb{T} \text{ with } |I| \leq 1 - r_0.$$

In fact, we have

$$(2.16) \quad \frac{\mu_g(S(I) \setminus r_0\mathbb{D})}{|I|} < 2\varepsilon^{\frac{2p}{p-2}}, \quad \text{for any } I \subseteq \mathbb{T}.$$

For any $|I| > 1 - r_0$, there exists $k \in \mathbb{N}$ such that $(k-1)(1-r_0) < |I| \leq k(1-r_0)$. There also exist intervals I_l on \mathbb{T} , $l = 1, 2, \dots, k$ such that $|I_l| = 1 - r_0$, $l = 1, 2, \dots, k$ and $I \subseteq \bigcup_{l=1}^k I_l$. Notice that $k(1-r_0) < 2|I|$. Combining with (2.15), we deduce

$$\mu_g(S(I) \setminus r_0\mathbb{D}) \leq \sum_{l=1}^k \mu_g(S(I_l) \setminus r_0\mathbb{D}) < \varepsilon^{\frac{2p}{p-2}} k(1-r_0) < \varepsilon^{\frac{2p}{p-2}} 2|I|,$$

which yields that (2.16) holds.

Let

$$\kappa(z) = \frac{\widehat{\omega}(z)}{1 - |z|}, \quad z \in \mathbb{D}.$$

Since $\chi_{\mathbb{D} \setminus r_0 \mathbb{D}} \mu_g$ is a Carleson measure, by [Dur70, Theorem 9.4], Lemma 2.5, Hölder's inequality and (2.16), for any $h \in H^{\frac{p}{p-2}}$, we have

$$\begin{aligned} \int_{\mathbb{D} \setminus r_0 \mathbb{D}} |h(z)| d\mu_{f_j, g}(z) &\leq \left(\int_{\mathbb{D} \setminus r_0 \mathbb{D}} |h(z)|^{\frac{p-2}{p}} d\mu_g(z) \right)^{\frac{p-2}{p}} \|f_j\|_{L_a^p(\kappa)}^2 \\ &\lesssim \|f_j\|_{L_a^p(\omega)}^2 \|h\|_{H^{\frac{p}{p-2}}} \cdot \left(\sup_{I \subseteq \mathbb{T}} \frac{\mu_g(S(I) \setminus r_0 \mathbb{D})}{|I|} \right)^{\frac{p-2}{p}} \\ (2.17) \quad &\lesssim \varepsilon^2 \|h\|_{H^{\frac{p}{p-2}}}. \end{aligned}$$

Furthermore, it follows from Lemma 2.2 that

$$(2.18) \quad \|J_g(f_j)\|_{H^p}^p \asymp \int_{\mathbb{T}} \left(\int_{\Gamma(\xi)} |f_j(z)|^2 |g'(z)|^2 dA(z) \right)^{\frac{p}{2}} d\xi.$$

Recall

$$(2.19) \quad \widetilde{\mu_{f_j, g}}(\xi) = \int_{\Gamma(\xi)} \frac{d\mu_{f_j, g}(z)}{1 - |z|^2}, \quad \xi \in \mathbb{T}.$$

By [Pau16, Theorem E], for any $j \in \mathbb{N}$, we have

$$\|Id\|_{H^{\frac{p}{p-2}} \rightarrow L^1(\mu_{f_j, g} \chi_{\mathbb{D} \setminus r_0 \mathbb{D}})} \asymp \left[\int_{\mathbb{T}} \left(\int_{\Gamma(\xi) \setminus r_0 \mathbb{D}} \frac{d\mu_{f_j, g}(z)}{1 - |z|^2} \right)^{\frac{p}{2}} d\xi \right]^{\frac{2}{p}}.$$

Together with (2.17), (2.18) and (2.19), for $j > k_0$, we deduce

$$\|J_g(f_j)\|_{H^p} \asymp \|\widetilde{\mu_{f_j, g}}\|_{L^{\frac{p}{2}}(\mathbb{T})}^{\frac{1}{2}} \lesssim \varepsilon.$$

Hence, we know that J_g is a compact operator as desired.

Necessity. Assume that $J_g : L_a^p(\omega) \rightarrow H^p$ is compact. Take a positive constant α . Let $d\nu_g(z) = |g'(z)|^2 (1 - |z|) dA(z)$ and $A_{\nu_g, \alpha}^2$ be the area integral operator from $L_a^p(\omega)$ to $L^p(\mathbb{T})$. Since the dual space of $L^{\frac{p}{2}}(\mathbb{T})$ can be identified with $L^{\frac{p}{p-2}}(\mathbb{T})$, together with Fubini's theorem, for any $h \in L^{\frac{p}{p-2}}(\mathbb{T})$ with $h \geq 0$ and $f \in L_a^p(\omega)$, we obtain

$$(2.20) \quad \int_{\mathbb{T}} h(\xi) (A_{\nu_g, \alpha}^2(f)(\xi))^2 d\xi = \int_{\mathbb{D}} \frac{1}{1 - |z|} \int_{I_\alpha(z)} h(\xi) d\xi |f(z)|^2 d\nu_g(z).$$

Notice that for any $I \subseteq \mathbb{T}$, there exists $a \in \mathbb{D}$ such that $I = I_\alpha(a)$. For any $a \in \mathbb{D}$, take testing functions

$$h_a(\xi) = \chi_{I_\alpha(a)}(\xi), \quad \xi \in \mathbb{T}.$$

Notice that

$$(2.21) \quad \|h_a\|_{L^{\frac{p}{p-2}}(\mathbb{T})} \asymp (1 - |a|)^{\frac{p-2}{p}}$$

and

$$(2.22) \quad \int_{I_\alpha(z)} h_a(\xi) d\xi \asymp 1 - |z|, \quad z \in \wedge_\alpha(I_\alpha(a)).$$

Combining (2.20), (2.21), (2.22) with Hölder's inequality, for any $a \in \mathbb{D}$, we deduce

$$\begin{aligned} \int_{\wedge_\alpha(I_\alpha(a))} |f(z)|^2 d\nu_g(z) &\lesssim \int_{\mathbb{D}} \frac{1}{1-|z|} \int_{I_\alpha(z)} h_a(\xi) d\xi |f(z)|^2 d\nu_g(z) \\ &= \int_{\mathbb{T}} h_a(\xi) (A_{\nu_g, \alpha}^2(f)(\xi))^2 d\xi \\ &\leq \|h_a\|_{L^{\frac{p}{p-2}}(\mathbb{T})} \|A_{\nu_g, \alpha}^2(f)\|_{L^p(\mathbb{T})}^2 \\ &\asymp (1-|a|)^{\frac{p-2}{p}} \|A_{\nu_g, \alpha}^2(f)\|_{L^p(\mathbb{T})}^2, \end{aligned}$$

from which it follows that

$$(2.23) \quad \frac{\int_{\wedge_\alpha(I_\alpha(a))} |f(z)|^2 d\nu_g(z)}{(1-|a|)^{\frac{p-2}{p}}} \lesssim \|A_{\nu_g, \alpha}^2(f)\|_{L^p(\mathbb{T})}^2.$$

For any $\zeta \in \mathbb{D}$ and $\beta > 0$, consider test functions

$$f_{\zeta, \beta}(z) := \frac{(1-|\zeta|)^{\frac{\beta+1}{p}}}{(1-\bar{\zeta}z)^{\frac{\beta+1}{p}} \widehat{\omega}(\zeta)^{\frac{1}{p}} (1-|\zeta|)^{\frac{1}{p}}}, \quad z \in \mathbb{D}.$$

Recall Lemma A(vi) in [PRS18] states that there exists $\gamma_0 = \gamma_0(\omega) > 0$ such that for any $\gamma \geq \gamma_0$, it holds that

$$\int_{\mathbb{D}} \frac{\omega(z)}{|1-\bar{\zeta}z|^{\gamma+1}} dA(z) \asymp \frac{\widehat{\omega}(\zeta)}{(1-|\zeta|)^\gamma}, \quad \zeta \in \mathbb{D}.$$

Hence, we can choose γ large enough such that $\|f_{\zeta, \gamma}\|_{L_a^p(\omega)} \asymp 1$ and $f_{\zeta, \gamma}$ converges uniformly on any compact subset of \mathbb{D} as $|\zeta| \rightarrow 1^-$. Combining with (2.23), we obtain that for any $\zeta \in \mathbb{D}$ and $a \in \mathbb{D}$, it holds that

$$(2.24) \quad \frac{\int_{\wedge_\alpha(I_\alpha(a))} |f_{\zeta, \gamma}(z)|^2 d\nu_g(z)}{(1-|a|)^{\frac{p-2}{p}}} \lesssim \|A_{\nu_g, \alpha}^2(f_{\zeta, \gamma})\|_{L^p(\mathbb{T})}^2.$$

In view of the compactness of J_g and (2.13), for any $\varepsilon > 0$, there exists $r_1 \in (0, 1)$ such that for any $\zeta \in \mathbb{D}$ with $r_1 \leq |\zeta| < 1$, it holds that $\|A_{\nu_g, \alpha}^2(f_{\zeta, \gamma})\|_{L^p(\mathbb{T})}^2 < \varepsilon$. Notice that there exists $r_2 \in (r_1, 1)$ such that for any $a \in \mathbb{D}$ with $r_2 \leq |a| < 1$,

$$D(z, r) \subseteq \mathbb{D} \setminus r_1 \mathbb{D}, \quad \text{for any } z \in \wedge_\alpha(I_\alpha(a)).$$

For $a \in \mathbb{D}$ with $r_2 \leq |a| < 1$, take $\zeta \in \mathbb{D}$ with

$$\zeta \in \bigcup_{z \in \wedge_\alpha(I_\alpha(a))} D(z, r).$$

Together with (2.24), for any $a \in \mathbb{D}$ with $r_2 \leq |a| < 1$, we deduce

$$(2.25) \quad \frac{\int_{\wedge_\alpha(I_\alpha(a))} \frac{1}{\widehat{\omega}(z)^{\frac{2}{p}} (1-|z|)^{\frac{2}{p}}} d\nu_g(z)}{(1-|a|)^{\frac{p-2}{p}}} \lesssim \varepsilon.$$

Next we show that for any $f \in L_a^p(\omega)$, $|f(z)|^2 d\nu_g(z)$ is a vanishing $\frac{p-2}{p}$ -Carleson measure. By the subharmonicity of $|f|^p$, Lemma 2.5, Lemma 2.7 and (2.25), for any $a \in \mathbb{D}$ with

$r_2 \leq |a| < 1$, we obtain

$$\begin{aligned}
\frac{\int_{\wedge_\alpha(I_\alpha(a))} |f(z)|^2 d\nu_g(z)}{(1-|a|)^{\frac{p-2}{p}}} &\lesssim \frac{\int_{\wedge_\alpha(I_\alpha(a))} \left(\frac{1}{\widehat{\omega}(z)(1-|z|)} \int_{D(z,r)} |f(u)|^p \frac{\widehat{\omega}(u)}{1-|u|} dA(u) \right)^{\frac{2}{p}} d\nu_g(z)}{(1-|a|)^{\frac{p-2}{p}}} \\
&\leq \|f\|_{L_a^p(\omega)}^2 \frac{\int_{\wedge_\alpha(I_\alpha(a))} \frac{1}{\widehat{\omega}(z)^{\frac{2}{p}}(1-|z|)^{\frac{2}{p}}} d\nu_g(z)}{(1-|a|)^{\frac{p-2}{p}}} \\
(2.26) \qquad \qquad \qquad &\lesssim \varepsilon \|f\|_{L_a^p(\omega)}^2.
\end{aligned}$$

It follows from Proposition 2.10 and (2.26) that there exists $t \in (0, 1)$ such that

$$\left(\frac{\nu_g(D(z, t))}{\widehat{\omega}(z)(1-|z|)} \right)^{\frac{p}{p-2}} \frac{\widehat{\omega}(z)}{1-|z|} dA(z)$$

is a vanishing Carleson measure. By the subharmonicity of $|g'|^2$, we deduce

$$|g'(z)|^2 \frac{(1-|z|)^2}{\widehat{\omega}(z)} \lesssim \frac{1}{\widehat{\omega}(z)(1-|z|)} \int_{D(z,t)} |g'(\zeta)|^2 (1-|\zeta|) dA(\zeta) = \frac{\nu_g(D(z, t))}{\widehat{\omega}(z)(1-|z|)},$$

which yields that

$$\left(|g'(z)|^2 \frac{(1-|z|)^2}{\widehat{\omega}(z)} \right)^{\frac{p}{p-2}} \frac{\widehat{\omega}(z)}{1-|z|} dA(z)$$

is a vanishing Carleson measure. Thus, we conclude that μ_g is a vanishing Carleson measure. \square

2.3. Case C. Now we are devoted to dealing with the case $0 < q < \infty$ and $p > \max\{2, q\}$.

For a separated sequence $\{a_k\}_{k=1}^\infty \subset \mathbb{D}$, let $\nu = \sum_{k=1}^\infty \delta_{a_k}$ where δ_{a_k} is Dirac measure at point a_k , then we denote $T_{q,\nu}^p$ by $T_q^p(\{a_k\})$. For $0 < p < \infty$ and $0 < q \leq \infty$, if

$$\|\{b_k\}\|_{T_q^p(\{a_k\})}^p := \int_{\mathbb{T}} \left(\sum_{a_k \in \Gamma(\xi)} |b_k|^q \right)^{\frac{p}{q}} d\xi < \infty, \quad q \in (0, \infty)$$

then we say the sequence $\{b_k\}_{k=1}^\infty \in T_q^p(\{a_k\})$, and if

$$\|\{b_k\}\|_{T_q^p(\{a_k\})}^p := \int_{\mathbb{T}} \left(\sup_{a_k \in \Gamma(\xi)} |b_k| \right)^p d\xi < \infty, \quad q = \infty.$$

we say the sequence $\{b_k\}_{k=1}^\infty \in T_\infty^p(\{a_k\})$.

The following two results characterize the dual space and the factorization of tent spaces of sequences respectively (see [Ars99] and [MPPW20]).

Lemma 2.14. *Let $1 < p < \infty$ and $\{a_k\}_{k=1}^\infty$ be a separated sequence.*

- *If $1 < q < \infty$, then the dual space of $T_q^p(\{a_k\})$ can be identified with $T_q^{p'}(\{a_k\})$ via the pairing,*

$$\langle \{b_k\}, \{d_k\} \rangle_{T_2^2(\{a_k\})} = \sum_{k=1}^\infty b_k \bar{d}_k (1-|a_k|^2), \quad \{b_k\} \in T_q^p(\{a_k\}), \quad \{d_k\} \in T_q^{p'}(\{a_k\}).$$

- If $0 < q \leq 1$, the the dual space of $T_q^p(\{a_k\})$ can be identified with $T_\infty^{p'}(\{a_k\})$ via the pairing,

$$\langle \{b_k\}, \{d_k\} \rangle_{T_2^2(\{a_k\})} = \sum_{k=1}^{\infty} b_k \bar{d}_k (1 - |a_k|^2), \quad \{b_k\} \in T_q^p(\{a_k\}), \quad \{d_k\} \in T_\infty^{p'}(\{a_k\}).$$

Lemma 2.15. *Let $0 < p, q < \infty$ and $\{a_k\}_{k=1}^\infty$ be an r -lattice. If $p < p_1, p_2 < \infty$, $q < q_1, q_2 < \infty$, $\frac{1}{p_1} + \frac{1}{p_2} = \frac{1}{p}$ and $\frac{1}{q_1} + \frac{1}{q_2} = \frac{1}{q}$. Then*

$$T_q^p(\{a_k\}) = T_{q_1}^{p_1}(\{a_k\}) \cdot T_{q_2}^{p_2}(\{a_k\}),$$

where $T_{q_1}^{p_1}(\{a_k\}) \cdot T_{q_2}^{p_2}(\{a_k\}) = \{\{b_k\} : b_k = c_k d_k, \{c_k\} \in T_{q_1}^{p_1}(\{a_k\}), \{d_k\} \in T_{q_2}^{p_2}(\{a_k\})\}$. Moreover, for any $\{b_k\} \in T_q^p(\{a_k\})$, it holds that

$$\inf \|\{c_k\}\|_{T_{q_1}^{p_1}(\{a_k\})} \cdot \|\{d_k\}\|_{T_{q_2}^{p_2}(\{a_k\})} \lesssim \|\{b_k\}\|_{T_q^p(\{a_k\})},$$

where the infimum above is taken over all possible factorizations of $\{b_k\}$.

We also need the following useful estimates.

Lemma 2.16. [Ars99, Lemma 4] *Let $0 < s < \infty$, $\iota > \max\{1, \frac{1}{s}\}$ and μ be a positive measure on \mathbb{D} . Then*

$$\int_{\mathbb{T}} \left(\int_{\mathbb{D}} \left(\frac{1 - |z|^2}{|1 - z\bar{\xi}|} \right)^\iota d\mu(z) \right)^s d\xi \asymp \int_{\mathbb{T}} \mu(\Gamma(\xi))^s d\xi.$$

Using similar analysis to that in [PRS18, Proposition 14], we get the following result.

Proposition 2.17. *Let $1 < p < \infty$ and $\omega \in \mathcal{D}$. Assume that $\{a_k\}_{k=1}^\infty \subseteq \mathbb{D} \setminus \{0\}$ is a separated sequence. Then*

$$F := \sum_{k=1}^{\infty} c_k \frac{K_{a_k}^\omega}{\|K_{a_k}^\omega\|_{L_a^{2-\frac{1}{p}}(\omega)}} \in H(\mathbb{D}).$$

Moreover, $\|F\|_{L_a^p(\omega)} \lesssim \|\{c_k\}_{k=1}^\infty\|_{l^p}$ for all $\{c_k\}_{k=1}^\infty \in l^p$.

Proof. Let $\kappa(z) = \frac{\widehat{\omega}(z)}{1-|z|}$, $z \in \mathbb{D}$ and $\{c_k\}_{k=1}^\infty \in l^p$. From Lemma 2.5, we know that κ is regular. Assume that $\rho \in (0, 1)$. For any $z \in \mathbb{D}$ with $0 < |z| \leq \rho$, it follows from Lemma 2.6, Hölder's inequality and (2.3) that

$$\begin{aligned} |F(z)| &\leq \left(\sum_{k=1}^{\infty} |c_k|^p \right)^{\frac{1}{p}} \left[\sum_{k=1}^{\infty} \left(\frac{|K_{a_k}^\omega(z)|}{\|K_{a_k}^\omega\|_{L_a^{2-\frac{1}{p}}(\omega)}} \right)^{p'} \right]^{\frac{1}{p'}} \\ &\lesssim \left(\sum_{k=1}^{\infty} |c_k|^p \right)^{\frac{1}{p}} \sup_{k \geq 1} |K_{a_k}^\omega(z)| \left(\sum_{k=1}^{\infty} \widehat{\omega}(a_k) (1 - |a_k|) \right)^{\frac{1}{p'}} \\ &\asymp \left(\sum_{k=1}^{\infty} |c_k|^p \right)^{\frac{1}{p}} \sup_{k \geq 1} |K_{a_k}^\omega(z)| \left(\sum_{k=1}^{\infty} \kappa(D(a_k, r)) \right)^{\frac{1}{p'}} \\ &\leq C \left(\sum_{k=1}^{\infty} |c_k|^p \right)^{\frac{1}{p}}, \end{aligned}$$

where $r \in (0, 1)$ and C is a constant which only depends on p, ρ, r and ω . Thus, we know that F is analytic on \mathbb{D} . Take Lemma 2.5 into account, we deduce $\|F\|_{L_a^p(\omega)} \asymp \|F\|_{L_a^p(\kappa)}$. Since $(L_a^p(\kappa))^* \simeq L_a^{p'}(\kappa)$ (see [PRS18, Theorem D]), together with the property of subharmonicity, similar analysis as above gives the desired result. \square

Now we give the main result of this subsection.

Proposition 2.18. *Let $0 < q < \infty$, $p > \max\{2, q\}$, $\omega \in \mathcal{D}$ and $g \in H(\mathbb{D})$. Then the Volterra type integration operator $J_g : L_a^p(\omega) \rightarrow H^q$ is compact if and only if J_g is bounded if and only if $g' \in T_{\frac{2p}{p-2}, \widehat{\omega}, \frac{2}{p-2}}^{\frac{pq}{p-q}}$. Moreover, $\|J_g\|_{L_a^p(\omega) \rightarrow H^q} \asymp \|g'\|_{T_{\frac{2p}{p-2}, \widehat{\omega}, \frac{2}{p-2}}^{\frac{pq}{p-q}}}$.*

Proof. We divide the proof into two steps.

Step 1. Let g be an analytic function on \mathbb{D} with satisfying

$$g' \in T_{\frac{2p}{p-2}, \widehat{\omega}, \frac{2}{p-2}}^{\frac{pq}{p-q}},$$

we will show that $J_g : L_a^p(\omega) \rightarrow H^q$ is a compact operator. Let $\{f_j\}_{j=1}^\infty$ be any sequence in $L_a^p(\omega)$ with $\|f_j\|_{L_a^p(\omega)} \leq 1$ and f_j converges to 0 uniformly on any compact subset of \mathbb{D} . We are devoted to proving

$$(2.27) \quad \lim_{j \rightarrow \infty} \|J_g(f_j)\|_{H^q} = 0.$$

Since $g' \in T_{\frac{2p}{p-2}, \widehat{\omega}, \frac{2}{p-2}}^{\frac{pq}{p-q}}$, then for any $\varepsilon > 0$, there exists $r_0 \in (0, 1)$ such that

$$(2.28) \quad \left[\int_{\mathbb{T}} \left(\int_{\Gamma(\xi) \setminus r_0 \mathbb{D}} |g'(z)|^{\frac{2p}{p-2}} \left(\frac{(1-|z|^2)^2}{\widehat{\omega}(z)} \right)^{\frac{2}{p-2}} dA(z) \right)^{\frac{q(p-2)}{2(p-q)}} d\xi \right]^{\frac{p-q}{p}} < \varepsilon.$$

Moreover, there exists $k_0 \in \mathbb{N}$ such that for any $j > k_0$, $|f_j|^q < \varepsilon$ on $r_0 \mathbb{D}$. Combining Lemma 2.5, (2.6) with Fubini's theorem, for any $j \in \mathbb{N}$, we deduce

$$(2.29) \quad \int_{\mathbb{T}} \int_{\Gamma(\xi) \setminus r_0 \mathbb{D}} |f_j(z)|^p \frac{\widehat{\omega}(z)}{(1-|z|^2)^2} dA(z) d\xi \lesssim \int_{\mathbb{D}} |f_j(z)|^p \frac{\widehat{\omega}(z)}{(1-|z|^2)^2} \int_{I(z)} d\xi dA(z) \asymp \|f_j\|_{L_a^p(\omega)}^p.$$

By Lemma 2.2, (2.28), (2.29) and Hölder's inequality, we obtain

$$\begin{aligned} \|J_g(f_j)\|_{H^q}^q &\lesssim \int_{\mathbb{T}} \left(\int_{\Gamma(\xi) \cap r_0 \mathbb{D}} |g'(z)|^2 |f_j(z)|^2 dA(z) \right)^{\frac{q}{2}} d\xi + \int_{\mathbb{T}} \left(\int_{\Gamma(\xi) \setminus r_0 \mathbb{D}} |g'(z)|^2 |f_j(z)|^2 dA(z) \right)^{\frac{q}{2}} d\xi \\ &\lesssim \varepsilon + \int_{\mathbb{T}} \left[\left(\int_{\Gamma(\xi) \setminus r_0 \mathbb{D}} |g'(z)|^{\frac{2p}{p-2}} \left(\frac{(1-|z|^2)^2}{\widehat{\omega}(z)} \right)^{\frac{2}{p-2}} dA(z) \right)^{\frac{q(p-2)}{2p}} \right. \\ &\quad \cdot \left. \left(\int_{\Gamma(\xi) \setminus r_0 \mathbb{D}} |f_j(z)|^p \frac{\widehat{\omega}(z)}{(1-|z|^2)^2} dA(z) \right)^{\frac{q}{p}} \right] d\xi \\ &\leq \varepsilon + \left[\int_{\mathbb{T}} \left(\int_{\Gamma(\xi) \setminus r_0 \mathbb{D}} |g'(z)|^{\frac{2p}{p-2}} \left(\frac{(1-|z|^2)^2}{\widehat{\omega}(z)} \right)^{\frac{2}{p-2}} dA(z) \right)^{\frac{q(p-2)}{2(p-q)}} d\xi \right]^{\frac{p-q}{p}} \\ &\quad \cdot \left[\int_{\mathbb{T}} \left(\int_{\Gamma(\xi) \setminus r_0 \mathbb{D}} |f_j(z)|^p \frac{\widehat{\omega}(z)}{(1-|z|^2)^2} dA(z) \right) d\xi \right]^{\frac{q}{p}} \\ &\lesssim \varepsilon + \varepsilon \|f_j\|_{L_a^p(\omega)}^q \lesssim \varepsilon, \quad \text{for any } j > k_0. \end{aligned}$$

Step 2. Assume that J_g is bounded. For $r \in (0, 1)$, let $\tilde{r} = \tanh r$ and $\{a_k\}_{k=1}^\infty$ be an \tilde{r} -lattice. It follows from Lemma 2.7 that

$$(2.30) \quad \widehat{\omega}(z) \asymp \widehat{\omega}(a_k), \quad z \in D(a_k, \tilde{r}).$$

By the subharmonicity of $|g'|^2$ and (2.30), for any $\xi \in \mathbb{T}$, we get

$$\begin{aligned} \int_{\Gamma(\xi)} |g'(z)|^{\frac{2p}{p-2}} \frac{(1-|z|^2)^{\frac{4}{p-2}}}{\widehat{\omega}(z)^{\frac{2}{p-2}}} dA(z) &\lesssim \int_{\Gamma(\xi)} \left(\int_{D(z, \tilde{r})} |g'(\zeta)|^2 dA(\zeta) \right)^{\frac{p}{p-2}} \frac{1}{\widehat{\omega}(z)^{\frac{2}{p-2}} (1-|z|^2)^2} dA(z) \\ &\leq \sum_k \int_{D(a_k, \tilde{r})} \left(\int_{D(z, \tilde{r})} |g'(\zeta)|^2 dA(\zeta) \right)^{\frac{p}{p-2}} \frac{1}{\widehat{\omega}(z)^{\frac{2}{p-2}} (1-|z|^2)^2} dA(z) \\ &\lesssim \sum_k \left(\sup_{\zeta \in D(a_k, 2\tilde{r})} |g'(\zeta)|^{\frac{2p}{p-2}} \right) \frac{(1-|a_k|^2)^{\frac{2p}{p-2}}}{\widehat{\omega}(a_k)^{\frac{2}{p-2}}}, \end{aligned}$$

where the summation runs $\{k : D(a_k, \tilde{r}) \cap \Gamma(\xi) \neq \emptyset\}$. Let $\delta = 3e^{2r} - 2$. By Lemma 2.11, for any $\xi \in \mathbb{T}$, we have

$$D(\zeta, \tilde{r}) \subseteq \Gamma_\delta(\xi), \quad \forall \zeta \in \Gamma(\xi).$$

Hence, for any $\xi \in \mathbb{T}$, we deduce

$$\int_{\Gamma(\xi)} |g'(z)|^{\frac{2p}{p-2}} \frac{(1-|z|^2)^{\frac{4}{p-2}}}{\widehat{\omega}(z)^{\frac{2}{p-2}}} dA(z) \lesssim \sum_{a_k \in \Gamma_\delta(\xi)} \left(\sup_{\zeta \in D(a_k, 2\tilde{r})} |g'(\zeta)|^{\frac{2p}{p-2}} \right) \frac{(1-|a_k|^2)^{\frac{2p}{p-2}}}{\widehat{\omega}(a_k)^{\frac{2}{p-2}}}.$$

Thus, we obtain

$$\begin{aligned} &\int_{\mathbb{T}} \left(\int_{\Gamma(\xi)} |g'(z)|^{\frac{2p}{p-2}} \frac{(1-|z|^2)^{\frac{4}{p-2}}}{\widehat{\omega}(z)^{\frac{2}{p-2}}} dA(z) \right)^{\frac{q(p-2)}{2(p-q)}} d\xi \\ &\lesssim \int_{\mathbb{T}} \left(\sum_{a_k \in \Gamma(\xi)} \left(\sup_{\zeta \in D(a_k, 2\tilde{r})} |g'(\zeta)|^{\frac{2p}{p-2}} \right) \frac{(1-|a_k|^2)^{\frac{2p}{p-2}}}{\widehat{\omega}(a_k)^{\frac{2}{p-2}}} \right)^{\frac{q(p-2)}{2(p-q)}} d\xi. \end{aligned}$$

In what follows, we only need to prove

$$(2.31) \quad \int_{\mathbb{T}} \left(\sum_{a_k \in \Gamma(\xi)} \left(\sup_{\zeta \in D(a_k, 2\tilde{r})} |g'(\zeta)|^{\frac{2p}{p-2}} \right) \frac{(1-|a_k|^2)^{\frac{2p}{p-2}}}{\widehat{\omega}(a_k)^{\frac{2}{p-2}}} \right)^{\frac{q(p-2)}{2(p-q)}} d\xi \lesssim \|J_g\|_{L_a^p(\omega) \rightarrow H^q}^{\frac{pq}{p-q}}.$$

Define a sequence $\{b_k\}_{k=1}^\infty$ by

$$b_k = \left(\sup_{\zeta \in D(a_k, 2\tilde{r})} |g'(\zeta)|^q \right) \frac{(1-|a_k|^2)^q}{\widehat{\omega}(a_k)^{\frac{q}{p}}}.$$

Let $\eta = \frac{2p}{q(p-2)}$. We turn to prove that there exists a positive constant s such that

$$(2.32) \quad \|\{b_k^{\frac{1}{s}}\}\|_{T_{\eta s}^{\frac{ps}{p-s}}(\{a_k\})} \lesssim \|J_g\|_{L_a^p(\omega) \rightarrow H^q}^{\frac{q}{s}}.$$

Choose a constant s with $s > \max\{1, \frac{q}{2}\}$. From Lemma 2.14, the dual space of $T_{(\eta s)'}^{\left(\frac{ps}{p-s}\right)' }(\{a_k\})$ can be identified with $T_{\eta s}^{\frac{ps}{p-s}}(\{a_k\})$. Thus, by (2.6), for any sequence $\{e_k\} \in T_{(\eta s)'}^{\left(\frac{ps}{p-s}\right)' }(\{a_k\})$,

it holds that

$$(2.33) \quad |\langle e_k, b_k^{\frac{1}{s}} \rangle_{T_2^2(\{a_k\})}| \leq \sum_{k=1}^{\infty} |e_k b_k^{\frac{1}{s}}| (1 - |a_k|^2) \asymp \sum_{k=1}^{\infty} |e_k b_k^{\frac{1}{s}}| \int_{I(a_k)} d\xi = \int_{\mathbb{T}} \sum_{a_k \in \Gamma(\xi)} |e_k b_k^{\frac{1}{s}}| d\xi.$$

By Lemma 2.15, we obtain

$$(2.34) \quad T_{(\eta s)'}^{\left(\frac{ps}{p-q}\right)'}(\{a_k\}) = T_{\frac{2s}{2s-q}}^{s'}(\{a_k\}) \cdot T_{\frac{ps}{q}}^{\frac{ps}{q}}(\{a_k\}),$$

from which there exist two sequences $\{x_k\} \in T_{\frac{2s}{2s-q}}^{s'}(\{a_k\})$ and $\{y_k\} \in T_p^p(\{a_k\})$ such that

$$e_k = x_k y_k^{\frac{q}{s}}.$$

Together with (2.33) and Hölder's inequality, we get

$$(2.35) \quad \begin{aligned} |\langle e_k, b_k^{\frac{1}{s}} \rangle_{T_2^2(\{a_k\})}| &\lesssim \int_{\mathbb{T}} \left(\sum_{a_k \in \Gamma(\xi)} |x_k|^{\frac{2s}{2s-q}} \right)^{\frac{2s-q}{2s}} \left(\sum_{a_k \in \Gamma(\xi)} |y_k^2 b_k^{\frac{2}{q}}| \right)^{\frac{q}{2s}} d\xi \\ &\leq \|\{x_k\}\|_{T_{\frac{2s}{2s-q}}^{s'}(\{a_k\})} \left[\int_{\mathbb{T}} \left(\sum_{a_k \in \Gamma(\xi)} |y_k^2 b_k^{\frac{2}{q}}| \right)^{\frac{q}{2}} d\xi \right]^{\frac{1}{s}}. \end{aligned}$$

Let

$$(2.36) \quad f_k(z) = \frac{K_{a_k}^{\omega}(z)}{\|K_{a_k}^{\omega}\|_{L_a^{2-\frac{1}{p}}(\omega)}^{2-\frac{1}{p}}}, \quad z \in \mathbb{D}$$

and $\{r_k : k \in \mathbb{N}\}$ be the set of the Rademacher functions. For $t \in [0, 1]$, set

$$F_t(z) = \sum_{k=1}^{\infty} (1 - |a_k|^2)^{\frac{1}{p}} y_k r_k(t) f_k(z), \quad z \in \mathbb{D}.$$

Then

$$(2.37) \quad \|\{y_k\}\|_{T_p^p(\{a_k\})}^p = \int_{\mathbb{T}} \sum_{a_k \in \Gamma(\xi)} |y_k|^p d\xi = \sum_{k=1}^{\infty} \int_{I(a_k)} |y_k|^p d\xi \asymp \sum_{k=1}^{\infty} |y_k|^p (1 - |a_k|^2).$$

By using Proposition 2.17 and (2.37), we obtain

$$\|J_g(F_t)\|_{H^q}^q \lesssim \|J_g\|_{L_a^p(\omega) \rightarrow H^q}^q \left(\sum_{k=1}^{\infty} |y_k|^p (1 - |a_k|^2) \right)^{\frac{q}{p}} \asymp \|J_g\|_{L_a^p(\omega) \rightarrow H^q}^q \|\{y_k\}\|_{T_p^p(\{a_k\})}^q.$$

Together with Lemma 2.2, we deduce

$$(2.38) \quad \int_{\mathbb{T}} \left(\int_{\Gamma(\xi)} |g'(z)|^2 \left| \sum_{k=1}^{\infty} (1 - |a_k|^2)^{\frac{1}{p}} y_k r_k(t) f_k(z) \right|^2 dA(z) \right)^{\frac{q}{2}} d\xi \lesssim \|J_g\|_{L_a^p(\omega) \rightarrow H^q}^q \cdot \|\{y_k\}\|_{T_p^p(\{a_k\})}^q.$$

Integrate both sides of the inequality (2.38) from 0 to 1 with respect to t , by Fubini's theorem, we have

$$(2.39) \quad \int_{\mathbb{T}} \int_0^1 \left(\int_{\Gamma(\xi)} |g'(z)|^2 \left| \sum_{k=1}^{\infty} (1 - |a_k|^2)^{\frac{1}{p}} y_k r_k(t) f_k(z) \right|^2 dA(z) \right)^{\frac{q}{2}} dt d\xi \lesssim \|J_g\|_{L_a^p(\omega) \rightarrow H^q}^q \cdot \|\{y_k\}\|_{T_p^p(\{a_k\})}^q.$$

By the subharmonicity of $|g^2|$ and Lemma 2.7, for any $k \in \mathbb{N}$, we obtain

$$\begin{aligned}
\sup_{\zeta \in D(a_k, 2\tilde{r})} |g'(\zeta)|^2 &\lesssim \sup_{\zeta \in D(a_k, 2\tilde{r})} \frac{1}{(1 - |\zeta|)^2} \int_{B(\zeta, \tilde{r}(1 - |\zeta|))} |g'(z)|^2 dA(z) \\
&\leq \sup_{\zeta \in D(a_k, 2\tilde{r})} \frac{1}{(1 - |\zeta|)^2} \int_{D(\zeta, \tilde{r})} |g'(z)|^2 dA(z) \\
(2.40) \quad &\leq \frac{1}{\widehat{\omega}(a_k)(1 - |a_k|)} \int_{D(a_k, 3\tilde{r})} |g'(z)|^2 \frac{\widehat{\omega}(z)}{(1 - |z|)} dA(z).
\end{aligned}$$

For $z \in D(a_k, 3\tilde{r})$, it follows from [PRS18, Lemma 8] and (2.3) that for r small enough, we have

$$(2.41) \quad |f_k(z)| \asymp [\widehat{\omega}(a_k)(1 - |a_k|)]^{1 - \frac{1}{p}} |K_{a_k}^\omega(z)| \asymp \frac{1}{\widehat{\omega}(a_k)^{\frac{1}{p}}(1 - |a_k|)^{\frac{1}{p}}}.$$

Take $\iota > \max\{1, \frac{2}{q}\}$. Combining Lemma 2.7, (2.40) with (2.41), for $\xi \in \mathbb{T}$, we deduce

$$\begin{aligned}
\sum_{a_k \in \Gamma(\xi)} |y_k^2 b_k^{\frac{2}{q}}| &\asymp \sum_{a_k \in \Gamma(\xi)} |y_k|^2 \sup_{\zeta \in D(a_k, 2\tilde{r})} |g'(\zeta)|^2 \left(\frac{1 - |a_k|^2}{|1 - \bar{a}_k \xi|} \right)^\iota \frac{(1 - |a_k|^2)^2}{\widehat{\omega}(a_k)^{\frac{2}{p}}} \\
&\lesssim \sum_{k=1}^{\infty} |y_k|^2 \frac{1}{\widehat{\omega}(a_k)(1 - |a_k|)} \int_{D(a_k, 3\tilde{r})} |g'(z)|^2 \frac{\widehat{\omega}(z)}{1 - |z|} \left(\frac{1 - |z|^2}{|1 - \bar{z}\xi|} \right)^\iota \frac{(1 - |z|^2)^2}{\widehat{\omega}(z)^{\frac{2}{p}}} dA(z) \\
&\asymp \sum_{k=1}^{\infty} |y_k|^2 (1 - |a_k|^2)^{\frac{2}{p}} \int_{D(a_k, 3\tilde{r})} |g'(z)|^2 \left(\frac{1 - |z|^2}{|1 - \bar{z}\xi|} \right)^\iota |f_k(z)|^2 dA(z) \\
&\leq \int_{\mathbb{D}} \left(\frac{1 - |z|^2}{|1 - \bar{z}\xi|} \right)^\iota \sum_{k=1}^{\infty} |y_k|^2 (1 - |a_k|^2)^{\frac{2}{p}} |f_k(z)|^2 |g'(z)|^2 dA(z).
\end{aligned}$$

Together with Lemma 2.16, we know

$$\begin{aligned}
\int_{\mathbb{T}} \left(\sum_{a_k \in \Gamma(\xi)} |y_k^2 b_k^{\frac{2}{q}}| \right)^{\frac{q}{2}} d\xi &\lesssim \int_{\mathbb{T}} \left(\int_{\mathbb{D}} \left(\frac{1 - |z|^2}{|1 - \bar{z}\xi|} \right)^\iota \sum_{k=1}^{\infty} |y_k|^2 (1 - |a_k|^2)^{\frac{2}{p}} |f_k(z)|^2 |g'(z)|^2 dA(z) \right)^{\frac{q}{2}} d\xi \\
&\asymp \int_{\mathbb{T}} \left(\int_{\Gamma(\xi)} \sum_{k=1}^{\infty} |y_k|^2 (1 - |a_k|^2)^{\frac{2}{p}} |f_k(z)|^2 |g'(z)|^2 dA(z) \right)^{\frac{q}{2}} d\xi.
\end{aligned}$$

Moreover, using Lemma 2.8, Lemma 2.9 and Fubini's theorem, we deduce

$$\begin{aligned}
&\int_{\mathbb{T}} \left(\int_{\Gamma(\xi)} \sum_{k=1}^{\infty} |y_k|^2 (1 - |a_k|^2)^{\frac{2}{p}} |f_k(z)|^2 |g'(z)|^2 dA(z) \right)^{\frac{q}{2}} d\xi \\
&\asymp \int_{\mathbb{T}} \left(\int_{\Gamma(\xi)} |g'(z)|^2 \int_0^1 \left| \sum_{k=1}^{\infty} y_k r_k(t) (1 - |a_k|^2)^{\frac{1}{p}} f_k(z) \right|^2 dt dA(z) \right)^{\frac{q}{2}} d\xi \\
&= \int_{\mathbb{T}} \left(\int_0^1 \int_{\Gamma(\xi)} \left| g'(z) \sum_{k=1}^{\infty} y_k r_k(t) (1 - |a_k|^2)^{\frac{1}{p}} f_k(z) \right|^2 dA(z) dt \right)^{\frac{q}{2}} d\xi \\
&\asymp \int_{\mathbb{T}} \int_0^1 \left(\int_{\Gamma(\xi)} \left| g'(z) \sum_{k=1}^{\infty} y_k r_k(t) (1 - |a_k|^2)^{\frac{1}{p}} f_k(z) \right|^2 dA(z) \right)^{\frac{q}{2}} dt d\xi.
\end{aligned}$$

Combining with (2.39), we conclude that

$$(2.42) \quad \int_{\mathbb{T}} \left(\sum_{a_k \in \Gamma(\xi)} |y_k^2 b_k^{\frac{2}{q}}| \right)^{\frac{q}{2}} d\xi \lesssim \|J_g\|_{L_a^p(\omega) \rightarrow H^q}^q \cdot \|\{y_k\}\|_{T_p^p(\{a_k\})}^q.$$

According to (2.35) and (2.42), we deduce

$$(2.43) \quad |\langle e_k, b_k^{\frac{1}{s}} \rangle_{T_2^2(\{a_k\})}| \lesssim \|\{x_k\}\|_{T_{\frac{2s}{2s-q}}^{s'}(\{a_k\})} \cdot \|J_g\|_{L_a^p(\omega) \rightarrow H^q}^{\frac{q}{s}} \cdot \|\{y_k\}\|_{T_p^p(\{a_k\})}^{\frac{q}{s}}.$$

Take infimum over all possible factorizations of $\{e_k\}$ as in (2.34), we deduce

$$\inf \|\{x_k\}\|_{T_{\frac{2s}{2s-q}}^{s'}(\{a_k\})} \cdot \|\{y_k\}\|_{T_p^p(\{a_k\})}^{\frac{q}{s}} \lesssim \|\{e_k\}\|_{T_{(\frac{ps}{\eta s})'}(\{a_k\})}.$$

Combining with (2.43), we get

$$|\langle e_k, b_k^{\frac{1}{s}} \rangle_{T_2^2(\{a_k\})}| \lesssim \|J_g\|_{L_a^p(\omega) \rightarrow H^q}^{\frac{q}{s}} \cdot \|\{e_k\}\|_{T_{(\frac{ps}{\eta s})'}(\{a_k\})}.$$

Finally, it follows from Lemma 2.14 that (2.32) holds. Thus, we get the result as desired. \square

2.4. Case D. Last, we will deal with the remaining case $0 < q < p \leq 2$, with the main result as follows.

Proposition 2.19. *Let $0 < q < p \leq 2$, $\omega \in \mathcal{D}$ and $g \in H(\mathbb{D})$. Then the Volterra type integration operator $J_g : L_a^p(\omega) \rightarrow H^q$ is compact if and only if*

$$\lim_{r \rightarrow 1^-} \int_{\mathbb{T}} \sup_{z \in \Gamma(\xi) \setminus r\mathbb{D}} |g'(z)|^{\frac{pq}{p-q}} \frac{(1-|z|^2)^{\frac{pq}{p-q}}}{\widehat{\omega}(z)^{\frac{q}{p-q}}} d\xi = 0.$$

Proof. Sufficiency. Let $\{f_j\}_{j=1}^\infty$ be any sequence in $L_a^p(\omega)$ with $\|f_j\|_{L_a^p(\omega)} \leq 1$ and f_j converges to 0 uniformly on any compact subset of \mathbb{D} . Then we will prove $\|J_g(f_j)\|_{H^q} \rightarrow 0$ as $j \rightarrow \infty$. By the assumption, it follows that for any $\varepsilon > 0$, there exists $R_0 \in (0, 1)$ such that for any $R_1 \in [R_0, 1)$, it holds that

$$(2.44) \quad \int_{\mathbb{T}} \sup_{z \in \Gamma(\xi) \setminus R_1\mathbb{D}} |g'(z)|^{\frac{pq}{p-q}} \frac{(1-|z|^2)^{\frac{pq}{p-q}}}{\widehat{\omega}(z)^{\frac{q}{p-q}}} d\xi < \varepsilon^{\frac{p}{p-q}}.$$

Note that there also exists $k_0 \in \mathbb{N}$ such that

$$(2.45) \quad |f_j| < \varepsilon^{\frac{1}{q}} \text{ on } R_0\mathbb{D}, \quad \text{for any } j > k_0.$$

Choose $\gamma \in (0, \infty)$ with $\tilde{\gamma} := \tanh \gamma \in (0, \frac{1}{2})$ and $\{a_k\}_{k=1}^\infty$ be a $\tilde{\gamma}$ -lattice. Then we know

$$\Gamma(\xi) \subseteq \bigcup_{\{a_k : D(a_k, \tilde{\gamma}) \cap \Gamma(\xi) \neq \emptyset\}} D(a_k, \tilde{\gamma}).$$

Take $\alpha = 3e^{2\gamma} - 2$. By Lemma 2.11, we get

$$D(\zeta, \tilde{\gamma}) \subseteq \Gamma_\alpha(\xi), \quad \zeta \in \Gamma(\xi).$$

Hence, we obtain

$$(2.46) \quad \Gamma(\xi) \subseteq \bigcup_{a_k \in \Gamma_\alpha(\xi)} D(a_k, \tilde{\gamma}).$$

Observe ([MPPW20]) that there exists a constant $\beta > \alpha$ such that

$$(2.47) \quad \bigcup_{a_k \in \Gamma_\alpha(\xi)} D(a_k, \tilde{\gamma}) \subseteq \Gamma_\beta(\xi), \quad \xi \in \mathbb{T}.$$

For any $j \in \mathbb{N}$, by the subharmonicity of $|f_j|^2$, we deduce

$$(2.48) \quad |f_j(z)|^2 \lesssim \left(\frac{1}{(1-|a_k|)^2} \int_{D(a_k, 2\tilde{\gamma})} |f_j(u)|^p dA(u) \right)^{\frac{2}{p}}, \quad z \in D(a_k, \tilde{\gamma}).$$

It follows from Lemma 2.7 that

$$(2.49) \quad \widehat{\omega}(z) \asymp \widehat{\omega}(a_k), \quad z \in D(a_k, 2\tilde{\gamma}).$$

According to [Zhu07, Theorem 3.3], if $p \in (0, 1]$, then for any sequence $\{z_k\}_{k=1}^\infty \in l^p$,

$$(2.50) \quad \left| \sum_{k=1}^\infty z_k \right| \leq \left(\sum_{k=1}^\infty |z_k|^p \right)^{\frac{1}{p}}.$$

Combining (2.46), (2.47), (2.48), (2.49) with (2.50), for any $j \in \mathbb{N}$, we obtain

$$(2.51) \quad \left(\int_{\Gamma(\xi)} |f_j(z)|^2 (1-|z|^2)^{-2} \widehat{\omega}(z)^{\frac{2}{p}} dA(z) \right)^{\frac{q}{2}} \lesssim \int_{\Gamma_\beta(\xi)} |f_j(u)|^p \frac{\widehat{\omega}(u)}{(1-|u|)^2} dA(u).$$

By Lemma 2.5, (2.6) and Fubini's theorem, for any $j \in \mathbb{N}$, we have

$$\int_{\mathbb{T}} \int_{\Gamma_\beta(\xi)} |f_j(u)|^p \frac{\widehat{\omega}(u)}{(1-|u|)^2} dA(u) d\xi = \int_{\mathbb{D}} |f_j(u)|^p \frac{\widehat{\omega}(u)}{(1-|u|)^2} \int_{I_\beta(u)} d\xi dA(u) \asymp \|f_j\|_{L_a^p(\omega)}^p.$$

Together with Lemma 2.2, (2.44), (2.45), (2.51) and Hölder's inequality, for any $j > k_0$, we get

$$\begin{aligned} \|J_g(f_j)\|_{H^q}^q &\lesssim \int_{\mathbb{T}} \left(\int_{\Gamma(\xi) \cap R_0\mathbb{D}} |g'(z)|^2 |f_j(z)|^2 dA(z) \right)^{\frac{q}{2}} d\xi + \int_{\mathbb{T}} \left(\int_{\Gamma(\xi) \setminus R_0\mathbb{D}} |g'(z)|^2 |f_j(z)|^2 dA(z) \right)^{\frac{q}{2}} d\xi \\ &\lesssim \varepsilon + \int_{\mathbb{T}} \left(\sup_{z \in \Gamma(\xi) \setminus R_0\mathbb{D}} |g'(z)|^q \frac{(1-|z|^2)^q}{\widehat{\omega}(z)^{\frac{q}{p}}} \right) \cdot \left(\int_{\Gamma(\xi)} |f_j(z)|^2 \frac{\widehat{\omega}(z)^{\frac{2}{p}}}{(1-|z|^2)^2} dA(z) \right)^{\frac{q}{2}} d\xi \\ &\leq \varepsilon + \left[\int_{\mathbb{T}} \left(\sup_{z \in \Gamma(\xi) \setminus R_0\mathbb{D}} |g'(z)|^q \frac{(1-|z|^2)^q}{\widehat{\omega}(z)^{\frac{q}{p}}} \right)^{\frac{p-q}{p}} d\xi \right]^{\frac{p-q}{p}} \\ &\quad \cdot \left[\int_{\mathbb{T}} \left(\int_{\Gamma(\xi)} |f_j(z)|^2 \frac{\widehat{\omega}(z)^{\frac{2}{p}}}{(1-|z|^2)^2} dA(z) \right)^{\frac{q}{2}} d\xi \right]^{\frac{q}{p}} \\ &\lesssim \varepsilon + \varepsilon \|f_j\|_{L_a^p(\omega)}^q \lesssim \varepsilon, \end{aligned}$$

from which it follows that $J_g : L_a^p(\omega) \rightarrow H^q$ is compact.

Necessity. We divide the proof into two steps.

Step 1. Let δ be a positive constant such that $\tilde{\delta} := \tanh \delta \in (0, \frac{1}{3})$. Take $\{a_k\}_{k=1}^\infty$ be a $\tilde{\delta}$ -lattice. Rearrange the sequence $\{a_k\}_{k=1}^\infty$ such that $|a_1| \leq |a_2| \leq \dots \rightarrow 1^-$. For any $\rho_0 \in (0, 1)$ and $\xi \in \mathbb{T}$, we know that

$$\Gamma(\xi) \setminus \rho_0\mathbb{D} \subseteq \bigcup_{\{a_k : D(a_k, \tilde{\delta}) \cap (\Gamma(\xi) \setminus \rho_0\mathbb{D}) \neq \emptyset\}} D(a_k, \tilde{\delta}).$$

Let $\lambda = 3e^{2\delta} - 2$. By Lemma 2.11, for any $\xi \in \mathbb{T}$, we have

$$D(\zeta, \tilde{\delta}) \subseteq \Gamma_\lambda(\xi), \quad \text{for any } \zeta \in \Gamma(\xi).$$

Hence, for any $\zeta \in D(a_k, \tilde{\delta}) \cap (\Gamma(\xi) \setminus \rho_0 \mathbb{D})$, there exists $\rho_1 \in (0, \rho_0]$ such that

$$(2.52) \quad \Gamma(\xi) \setminus \rho_0 \mathbb{D} \subseteq \bigcup_{a_k \in \Gamma_\lambda(\xi) \setminus \rho_1 \mathbb{D}} D(a_k, \tilde{\delta})$$

and ρ_1 tends to 1 as ρ_0 tends to 1. Recall [Zhu07, Proposition 4.5], for any $z \in D(a_k, \tilde{\delta})$, we know that

$$\frac{1 - \tilde{\delta}}{1 + \tilde{\delta}}(1 - |z|^2) \leq 1 - |a_k|^2 \leq \frac{1 + \tilde{\delta}}{1 - \tilde{\delta}}(1 - |z|^2)$$

and

$$(2.53) \quad A(D(a_k, \tilde{\delta})) = \frac{\tilde{\delta}^2(1 - |a_k|^2)^2}{(1 - |a_k|^2 \tilde{\delta}^2)^2}.$$

Recall (see [PRS18]) there exist positive constants ϱ_1 and ϱ_2 such that

$$(2.54) \quad \left(\frac{1 - t_1}{1 - t_2}\right)^{\varrho_1} \widehat{\omega}(t_2) \lesssim \widehat{\omega}(t_1) \lesssim \left(\frac{1 - t_1}{1 - t_2}\right)^{\varrho_2} \widehat{\omega}(t_2), \quad 0 \leq t_1 \leq t_2 < 1.$$

By the subharmonicity of $|g'|$, for any $z \in D(a_k, \tilde{\delta})$, we have

$$(2.55) \quad \begin{aligned} |g'(z)| &\leq \frac{1}{\tilde{\delta}^2(1 - |z|^2)^2} \int_{B(z, \tilde{\delta}(1 - |z|))} |g'(u)| dA(u) \\ &\leq \frac{1}{\tilde{\delta}^2(1 - |z|^2)^2} \int_{D(z, \tilde{\delta})} |g'(u)| dA(u) \\ &\lesssim \frac{1}{\tilde{\delta}^2(1 - |a_k|^2)^2} \int_{D(a_k, 2\tilde{\delta})} |g'(u)| dA(u). \end{aligned}$$

Combining (2.52), (2.53), (2.54) with (2.55), we obtain

$$\begin{aligned} \sup_{z \in \Gamma(\xi) \setminus \rho_0 \mathbb{D}} \frac{(1 - |z|^2)}{\widehat{\omega}(z)^{\frac{1}{p}}} |g'(z)| &\lesssim \sup_{a_k \in \Gamma_\lambda(\xi) \setminus \rho_1 \mathbb{D}} \frac{1}{\tilde{\delta}^2(1 - |a_k|^2) \widehat{\omega}(a_k)^{\frac{1}{p}}} \int_{D(a_k, 2\tilde{\delta})} |g'(u)| dA(u) \\ &\lesssim \sup_{a_k \in \Gamma_\lambda(\xi) \setminus \rho_1 \mathbb{D}} \frac{(1 - |a_k|^2)}{\widehat{\omega}(a_k)^{\frac{1}{p}}} \left(\sup_{u \in D(a_k, 2\tilde{\delta})} |g'(u)| \right). \end{aligned}$$

Notice that the constants of comparison in the above inequality do not depend on $\tilde{\delta}$, ρ_0 and ρ_1 . Then we deduce there exists a constant $C > 0$ (C does not depend on the choices of $\tilde{\delta}$, ρ_0 and ρ_1) such that

$$\left(\sup_{z \in \Gamma(\xi) \setminus \rho_0 \mathbb{D}} \frac{(1 - |z|^2)}{\widehat{\omega}(z)^{\frac{1}{p}}} |g'(z)| \right)^{\frac{pq}{p-q}} \leq C \left[\sup_{a_k \in \Gamma_\lambda(\xi) \setminus \rho_1 \mathbb{D}} \frac{(1 - |a_k|^2)}{\widehat{\omega}(a_k)^{\frac{1}{p}}} \left(\sup_{u \in D(a_k, 2\tilde{\delta})} |g'(u)| \right) \right]^{\frac{pq}{p-q}}.$$

Thus, by integrating over \mathbb{T} with respect to ξ , we conclude that

$$(2.56) \quad \int_{\mathbb{T}} \left(\sup_{z \in \Gamma(\xi) \setminus \rho_0 \mathbb{D}} \frac{(1 - |z|^2)}{\widehat{\omega}(z)^{\frac{1}{p}}} |g'(z)| \right)^{\frac{pq}{p-q}} d\xi \leq C \int_{\mathbb{T}} \left[\sup_{a_k \in \Gamma_\lambda(\xi) \setminus \rho_1 \mathbb{D}} \frac{(1 - |a_k|^2)}{\widehat{\omega}(a_k)^{\frac{1}{p}}} \left(\sup_{u \in D(a_k, 2\tilde{\delta})} |g'(u)| \right) \right]^{\frac{pq}{p-q}} d\xi.$$

Step 2. We will prove that

$$(2.57) \quad \lim_{r \rightarrow 1^-} \int_{\mathbb{T}} \left[\sup_{a_k \in \Gamma(\xi) \setminus r\mathbb{D}} \frac{(1 - |a_k|^2)^{\frac{1}{p}}}{\widehat{\omega}(a_k)^{\frac{1}{p}}} \left(\sup_{u \in D(a_k, 2\tilde{\delta})} |g'(u)| \right) \right]^{\frac{pq}{p-q}} d\xi = 0.$$

Note that for the any $\varepsilon > 0$, there exists $R_2 \in (0, 1)$ such that for any $R_3 \in (R_2, 1)$, it holds that

$$(2.58) \quad \widehat{\omega}(R_3)(1 - R_3) < \varepsilon.$$

Moreover there exists $k_1 \in \mathbb{N}$ such that for any $k > k_1$, it holds that $|a_k| \in (R_2, 1)$. Combining with (2.58), we have

$$(2.59) \quad \widehat{\omega}(a_k)(1 - |a_k|) < \varepsilon, \quad \text{for any } k > k_1.$$

Since every $z \in \mathbb{D}$ belongs to at most N (N is a constant only depends on δ) $D(a_k, \tilde{\delta})$, then we get

$$\sum_{k=1}^{\infty} \kappa(D(a_k, \tilde{\delta})) \leq N\kappa(\mathbb{D}),$$

where

$$\kappa(z) = \frac{\widehat{\omega}(z)}{1 - |z|}, \quad z \in \mathbb{D}.$$

Hence, there exists $k_2 \in \mathbb{N}$ such that for any $k_3 > k_2$,

$$(2.60) \quad \sum_{k=k_3}^{\infty} \kappa(D(a_k, \tilde{\delta})) < \varepsilon.$$

Let r_k be the Rademacher functions. For $k \in \mathbb{N}$ and $p \in (0, 2]$, consider the functions

$$f_{k,p}(z) = \frac{K_{a_k}^{\omega}(z)^{\frac{1}{q}}}{\|K_{a_k}^{\omega}\|_{L_a^{\frac{p}{q}}(\omega)}^{\frac{1}{q}}} \chi_{\{p:0 < p \leq 1\}} + \frac{K_{a_k}^{\omega}(z)}{\|K_{a_k}^{\omega}\|_{L_a^{2-\frac{1}{p}}(\omega)}^{2-\frac{1}{p}}} \chi_{\{p:1 < p \leq 2\}}, \quad z \in \mathbb{D}.$$

For $t \in (0, 1)$, $k_4 \in \mathbb{N}$ and $\{y_k\}_{k=1}^{\infty} \in T_p^p(\{a_k\})$, choose test functions

$$F_{t,p,k_4,\{y_k\}}(z) = \frac{\sum_{k=k_4}^{\infty} (1 - |a_k|^2)^{\frac{1}{p}} y_k r_k(t) f_{k,p}(z)}{\|\{y_k\}\|_{T_p^p(\{a_k\})}}, \quad z \in \mathbb{D}.$$

Combining Lemma 2.6, (2.3), (2.37), (2.50), (2.59) with (2.60), for any $z \in \overline{D(0, \varrho)}$ with $\varrho \in (0, 1)$, for any $k_4 > \max\{k_1, k_2\}$, we deduce

$$\begin{aligned}
|F_{t,p,k_4,\{y_k\}}(z)| &\leq \frac{1}{\|\{y_k\}\|_{T_p^p(\{a_k\})}} \left(\sum_{k=k_4}^{\infty} (1 - |a_k|^2) |y_k|^p \frac{|K_{a_k}^\omega(z)|^{\frac{p}{q}}}{\|K_{a_k}^\omega\|_{L_a^{\frac{p}{q}}(\omega)}^{\frac{p}{q}}} \right)^{\frac{1}{p}} \\
&\lesssim \frac{\sup_{k \geq k_4} |K_{a_k}^\omega(z)|^{\frac{1}{q}}}{\|\{y_k\}\|_{T_p^p(\{a_k\})}} \left[\sum_{k=k_4}^{\infty} (1 - |a_k|^2) |y_k|^p \left(\widehat{\omega}(a_k)(1 - |a_k|) \right)^{\frac{p-1}{q}} \right]^{\frac{1}{p}} \\
&\leq \sup_{k \geq k_4} |K_{a_k}^\omega(z)|^{\frac{1}{q}} \sup_{k \geq k_4} \left(\widehat{\omega}(a_k)(1 - |a_k|) \right)^{\frac{p-q}{pq}} \frac{\left(\sum_{k=k_4}^{\infty} (1 - |a_k|^2) |y_k|^p \right)^{\frac{1}{p}}}{\|\{y_k\}\|_{T_p^p(\{a_k\})}} \\
&\lesssim \varepsilon^{\frac{p-q}{pq}} \sup_{k \geq k_4} |K_{a_k}^\omega(z)|^{\frac{1}{q}}, \quad p \in (0, 1]
\end{aligned}$$

and

$$\begin{aligned}
|F_{t,p,k_4,\{y_k\}}(z)| &\leq \frac{1}{\|\{y_k\}\|_{T_p^p(\{a_k\})}} \left(\sum_{k=k_4}^{\infty} (1 - |a_k|^2) |y_k|^p \right)^{\frac{1}{p}} \left[\sum_{k=k_4}^{\infty} \left(\frac{|K_{a_k}^\omega(z)|}{\|K_{a_k}^\omega\|_{L_a^{2-\frac{1}{p}}(\omega)}^{2-\frac{1}{p}}} \right)^{\frac{p}{p-1}} \right]^{\frac{p-1}{p}} \\
&\lesssim \sup_{k \geq k_4} |K_{a_k}^\omega(z)| \left(\sum_{k=k_4}^{\infty} \widehat{\omega}(a_k)(1 - |a_k|) \right)^{\frac{p-1}{p}} \\
&\asymp \sup_{k \geq k_4} |K_{a_k}^\omega(z)| \left(\sum_{k=k_4}^{\infty} \kappa(D(a_k, \tilde{\delta})) \right)^{\frac{p-1}{p}} \\
&\leq \varepsilon^{\frac{p-1}{p}} \sup_{k \geq k_4} |K_{a_k}^\omega(z)|, \quad p \in (1, 2],
\end{aligned}$$

which yields that for $p \in (0, 2]$, $F_{t,p,k_4,\{y_k\}}$ uniformly converges to 0 on any compact subset of \mathbb{D} as $k_4 \rightarrow \infty$. Furthermore, by using Proposition 2.17, (2.37) and (2.50), for any $k_4 \in \mathbb{N}$, we obtain

$$\|F_{t,p,k_4,\{y_k\}}\|_{L_a^p(\omega)}^p \leq \frac{1}{\|\{y_k\}\|_{T_p^p(\{a_k\})}^p} \int_{\mathbb{D}} \sum_{k=k_4}^{\infty} (1 - |a_k|^2) |y_k|^p \frac{|K_{a_k}^\omega(z)|^{\frac{p}{q}}}{\|K_{a_k}^\omega\|_{L_a^{\frac{p}{q}}(\omega)}^{\frac{p}{q}}} \omega(z) dA(z) \lesssim 1, \quad p \in (0, 1]$$

and

$$\|F_{t,p,k_4,\{y_k\}}\|_{L_a^p(\omega)}^p \lesssim \frac{\sum_{k=k_4}^{\infty} (1 - |a_k|^2) |y_k|^p}{\|\{y_k\}\|_{T_p^p(\{a_k\})}^p} \lesssim 1, \quad p \in (1, 2].$$

Thus, we know that for $p \in (0, 2]$, $\{F_{t,p,k_4,\{y_k\}}\}$ is a sequence in $L_a^p(\omega)$ with

$$\sup_{k_4 \in \mathbb{N}} \|F_{t,p,k_4,\{y_k\}}\|_{L_a^p(\omega)} \lesssim 1.$$

It follows from the compactness of J_g that

$$\lim_{k_4 \rightarrow \infty} \|J_g(F_{t,p,k_4,\{y_k\}})\|_{H^q}^q = 0.$$

Then there exists $k_5 \in \mathbb{N}$ such that for any $k_4 \in [k_5, \infty)$, it holds that

$$(2.61) \quad \|J_g(F_{t,p,k_4,\{y_k\}})\|_{H^q}^q < \varepsilon.$$

Note that k_5 does not depend on the choice of $\{y_k\}_{k=1}^\infty$ in $T_p^p(\{a_k\})$. Combine with Lemma 2.2, we obtain

$$\int_{\mathbb{T}} \left(\int_{\Gamma(\xi)} |g'(z)|^2 \left| \sum_{k=k_4}^\infty (1-|a_k|^2)^{\frac{1}{p}} y_k r_k(t) f_{k,p}(z) \right|^2 dA(z) \right)^{\frac{q}{2}} d\xi \asymp \|J_g(F_{t,p,k_4,\{y_k\}})\|_{H^q}^q \|\{y_k\}\|_{T_p^p(\{a_k\})}^q.$$

Integrate both sides of the inequality above from 0 to 1 with respect to t , by Fubini's theorem and (2.61), for any $k_4 \in [k_5, \infty)$, we have

$$(2.62) \quad \int_{\mathbb{T}} \int_0^1 \left(\int_{\Gamma(\xi)} |g'(z)|^2 \left| \sum_{k=k_4}^\infty (1-|a_k|^2)^{\frac{1}{p}} y_k r_k(t) f_{k,p}(z) \right|^2 dA(z) \right)^{\frac{q}{2}} dt d\xi \lesssim \varepsilon \|\{y_k\}\|_{T_p^p(\{a_k\})}^q.$$

Notice that there exists $R_4 \in (0, 1)$ such that for all $R_5 \in [R_4, 1)$, we have

$$k \geq k_5, \quad \text{for any } a_k \in \mathbb{D} \setminus R_5 \mathbb{D}.$$

Take a sequence $\{b_k\}_{k=1}^\infty$, where

$$b_k = \frac{(1-|a_k|^2)^q}{\widehat{\omega}(a_k)^{\frac{q}{p}}} \left(\sup_{u \in D(a_k, 2\delta)} |g'(u)| \right)^q \chi_{\{a_k: R_4 \leq |a_k| < 1\}}.$$

Choose $s > 1$. Let $\iota = \frac{2ps}{2ps-pq+2q}$. Then $\iota \leq 1$. Due to Lemma 2.14, the dual space of

$$T_\iota \left(\frac{ps}{p-q} \right)' (\{a_k\})$$

can be identified with

$$T_\infty^{\frac{ps}{p-q}} (\{a_k\}).$$

Take

$$\{e_k\}_{k=1}^\infty \in T_\iota \left(\frac{ps}{p-q} \right)' (\{a_k\}).$$

The factorization in Lemma 2.15 shows that there exist sequences

$$\{x_k\}_{k=1}^\infty \in T_{\frac{2s}{2s-q}}^{s'} (\{a_k\}) \text{ and } \{y_k\}_{k=1}^\infty \in T_p^p (\{a_k\})$$

such that $e_k = x_k y_k^{\frac{q}{s}}$. Using a similar estimation as in (2.35), we get

$$(2.63) \quad |\langle e_k, b_k^{\frac{1}{s}} \rangle_{T_2^2(\{a_k\})}| \lesssim \|\{x_k\}\|_{T_{\frac{2s}{2s-q}}^{s'}(\{a_k\})} \left[\int_{\mathbb{T}} \left(\sum_{a_k \in \Gamma(\xi) \setminus R_4 \mathbb{D}} |y_k^2 b_k^{\frac{2}{q}}| \right)^{\frac{q}{2}} d\xi \right]^{\frac{1}{s}}.$$

Take $\tau > \max\{1, \frac{2}{q}\}$. By the subharmonicity of $|g'^2|$, [PRS18, Lemma 8], Lemma 2.7 and (2.3), we deduce for δ small enough,

$$\begin{aligned}
\sum_{a_k \in \Gamma(\xi) \setminus R_4 \mathbb{D}} |y_k^2 b_k^{\frac{2}{q}}| &\asymp \sum_{a_k \in \Gamma(\xi) \setminus R_4 \mathbb{D}} |y_k|^2 \left(\sup_{u \in D(a_k, 2\delta)} |g'(u)| \right)^2 \left(\frac{1 - |a_k|^2}{|1 - \bar{a}_k \xi|} \right)^\tau \frac{(1 - |a_k|^2)^2}{\widehat{\omega}(a_k)^{\frac{2}{p}}} \\
&\lesssim \sum_{k=k_5}^{\infty} |y_k|^2 \frac{1}{\widehat{\omega}(a_k)(1 - |a_k|)} \int_{D(a_k, 3\delta)} |g'(z)|^2 \frac{\widehat{\omega}(z)}{1 - |z|} \left(\frac{1 - |z|^2}{|1 - \bar{z}\xi|} \right)^\tau \frac{(1 - |z|^2)^2}{\widehat{\omega}(z)^{\frac{2}{p}}} dA(z) \\
&\asymp \sum_{k=k_5}^{\infty} |y_k|^2 (1 - |a_k|^2)^{\frac{2}{p}} \int_{D(a_k, 3\delta)} |g'(z)|^2 \left(\frac{1 - |z|^2}{|1 - \bar{z}\xi|} \right)^\tau |f_{k,p}(z)|^2 dA(z) \\
&\leq \int_{\mathbb{D}} \left(\frac{1 - |z|^2}{|1 - \bar{z}\xi|} \right)^\tau \sum_{k=k_5}^{\infty} |y_k|^2 (1 - |a_k|^2)^{\frac{2}{p}} |g'(z)|^2 |f_{k,p}(z)|^2 dA(z).
\end{aligned}$$

Together with Lemma 2.16, we know

$$\begin{aligned}
\int_{\mathbb{T}} \left(\sum_{a_k \in \Gamma(\xi) \setminus R_4 \mathbb{D}} |y_k^2 b_k^{\frac{2}{q}}| \right)^{\frac{q}{2}} d\xi &\lesssim \int_{\mathbb{T}} \left(\int_{\mathbb{D}} \left(\frac{1 - |z|^2}{|1 - \bar{z}\xi|} \right)^\tau \sum_{k=k_5}^{\infty} |y_k|^2 (1 - |a_k|^2)^{\frac{2}{p}} |g'(z)|^2 |f_{k,p}(z)|^2 dA(z) \right)^{\frac{q}{2}} d\xi \\
(2.64) \quad &\asymp \int_{\mathbb{T}} \left(\int_{\Gamma(\xi)} \sum_{k=k_5}^{\infty} |y_k|^2 (1 - |a_k|^2)^{\frac{2}{p}} |f_{k,p}(z)|^2 |g'(z)|^2 dA(z) \right)^{\frac{q}{2}} d\xi.
\end{aligned}$$

Moreover, according to Lemma 2.8, Lemma 2.9 and Fubini's theorem, we obtain

$$\begin{aligned}
&\int_{\mathbb{T}} \left(\int_{\Gamma(\xi)} \sum_{k=k_5}^{\infty} |y_k|^2 (1 - |a_k|^2)^{\frac{2}{p}} |f_{k,p}(z)|^2 |g'(z)|^2 dA(z) \right)^{\frac{q}{2}} d\xi \\
&\asymp \int_{\mathbb{T}} \left(\int_{\Gamma(\xi)} |g'(z)|^2 \int_0^1 \left| \sum_{k=k_5}^{\infty} y_k r_k(t) (1 - |a_k|^2)^{\frac{1}{p}} f_{k,p}(z) \right|^2 dt dA(z) \right)^{\frac{q}{2}} d\xi \\
&= \int_{\mathbb{T}} \left(\int_0^1 \int_{\Gamma(\xi)} \left| g'(z) \sum_{k=k_5}^{\infty} y_k r_k(t) (1 - |a_k|^2)^{\frac{1}{p}} f_{k,p}(z) \right|^2 dA(z) dt \right)^{\frac{q}{2}} d\xi \\
&\asymp \int_{\mathbb{T}} \int_0^1 \left(\int_{\Gamma(\xi)} \left| g'(z) \sum_{k=k_5}^{\infty} y_k r_k(t) (1 - |a_k|^2)^{\frac{1}{p}} f_{k,p}(z) \right|^2 dA(z) \right)^{\frac{q}{2}} dt d\xi.
\end{aligned}$$

Together with (2.62) and (2.64), we get

$$(2.65) \quad \int_{\mathbb{T}} \left(\sum_{a_k \in \Gamma(\xi) \setminus R_4 \mathbb{D}} |y_k^2 b_k^{\frac{2}{q}}| \right)^{\frac{q}{2}} d\xi \lesssim \varepsilon \|\{y_k\}\|_{T_p^p(\{a_k\})}^q.$$

According to (2.63) and (2.65), it holds that

$$|\langle e_k, b_k^{\frac{1}{s}} \rangle_{T_2^2(\{a_k\})}| \lesssim \|\{x_k\}\|_{T_{\frac{2s}{2s-q}}^{s'}(\{a_k\})} \varepsilon^{\frac{1}{s}} \|\{y_k\}\|_{T_p^p(\{a_k\})}^{\frac{q}{s}}.$$

By taking infimum over all possible factorizations of $\{e_k\}$, we deduce

$$|\langle e_k, b_k^{\frac{1}{s}} \rangle_{T_2^2(\{a_k\})}| \lesssim \varepsilon^{\frac{1}{s}} \|\{e_k\}\|_{T_i(\frac{ps}{p-q})'(\{a_k\})}.$$

Therefore, using Lemma 2.14, we know that for any $\rho_1 \geq R_4$,

$$(2.66) \quad \int_{\mathbb{T}} \left(\sup_{a_k \in \Gamma_\lambda(\xi) \setminus \rho_1 \mathbb{D}} \frac{(1 - |a_k|^2)}{\widehat{\omega}(a_k)^{\frac{1}{p}}} \left(\sup_{u \in D(a_k, 2\delta)} |g'(u)| \right) \right)^{\frac{pq}{p-q}} d\xi \lesssim \varepsilon^{\frac{p}{p-q}}.$$

Hence, (2.57) holds.

Recall in Step 1, we can choose $\rho_1 \in (0, \rho_0]$ such that ρ_1 tends to 1 as ρ_0 tends to 1. Then we can choose $\rho_0 \in (0, 1)$ such that $\rho_1 \geq R_4$. Combining (2.56) with (2.66), we deduce

$$\int_{\mathbb{T}} \left(\sup_{z \in \Gamma(\xi) \setminus \rho_0 \mathbb{D}} \frac{(1 - |z|^2)}{\widehat{\omega}(z)^{\frac{1}{p}}} |g'(z)| \right)^{\frac{pq}{p-q}} d\xi \lesssim \varepsilon^{\frac{p}{p-q}}.$$

Thus, we complete the whole proof of the necessity part. \square

Combining Proposition 2.1, Proposition 2.13, Proposition 2.18 with Proposition 2.19, we get Theorem 1.2. Moreover, one can get Theorem 1.1 by modifying the corresponding proofs of the above four propositions slightly.

3. FROM HARDY SPACE INTO BERGMAN SPACE

In this section, we are devoted to proving Theorem 1.3. To begin with, we state a result (cf. [PR19, Theorem 5]) which is vital to the proof of our main result in this section. It describes that the class of \mathcal{D} weights is the largest class of radial weights such that the Littlewood-Paley estimate holds.

Lemma 3.1. *Let $0 < p < \infty$ and ω be a radial weight. Then*

$$\|f\|_{L_a^p(\omega)}^p \asymp \int_{\mathbb{D}} |f'(z)|^p (1 - |z|^2)^p \omega(z) dA(z) + |f(0)|^p, \quad f \in H(\mathbb{D})$$

holds if and only if $\omega \in \mathcal{D}$.

Proof of Theorem 1.3. Let

$$d\mu_g(z) = |g'(z)|^q (1 - |z|^2)^q \omega(z) dA(z).$$

It follows from Lemma 3.1 that

$$(3.67) \quad \|J_g(f)\|_{L_a^q(\omega)}^q \asymp \int_{\mathbb{D}} |f(z)|^q d\mu_g(z).$$

By (3.67), the boundedness (compactness) of $J_g : H^p \rightarrow L_a^q(\omega)$ is equivalent to that of $Id : H^p \rightarrow L^q(\mu_g)$. For the case when $0 < p \leq q < \infty$, the identity operator $Id : H^p \rightarrow L^q(\mu_g)$ is bounded (compact) if and only if μ_g is a $\frac{q}{p}$ -Carleson measure (vanishing $\frac{q}{p}$ -Carleson measure) (cf. [Zhu07, Section 9.2]).

Now we are in a position to deal with the case when $0 < q < p < \infty$. For $\xi \in \mathbb{T}$, recall

$$\widetilde{\mu}_g(\xi) = \int_{\Gamma(\xi)} \frac{d\mu_g(\lambda)}{1 - |\lambda|^2}.$$

By [Pau16, Theorem E] and (3.67), we know that $J_g : H^p \rightarrow L_a^q(\omega)$ is bounded if and only if $\widetilde{\mu}_g \in L^{\frac{p}{p-q}}(\mathbb{T})$.

It suffices to prove that $J_g : H^p \rightarrow L_a^q(\omega)$ is compact if $\tilde{\mu}_g \in L^{\frac{p}{p-q}}(\mathbb{T})$. Let $\{f_j\}_{j=1}^\infty$ be any sequence in H^p with $\|f_j\|_{H^p} \leq 1$ and f_j converges to 0 uniformly on any compact subset of \mathbb{D} . By Lemma 3.1, we obtain

$$(3.68) \quad \|J_g(f_j)\|_{L_a^q(\omega)}^q \asymp \int_{\mathbb{D}} |f_j(z)|^q d\mu_g(z).$$

Then we only need to show

$$(3.69) \quad \lim_{j \rightarrow \infty} \int_{\mathbb{D}} |f_j(z)|^q d\mu_g(z) = 0.$$

Since $\tilde{\mu}_g \in L^{\frac{p}{p-q}}(\mathbb{T})$, then for any $\varepsilon > 0$, there exists $r_0 \in (0, 1)$ such that

$$(3.70) \quad \left[\int_{\mathbb{T}} \left(\int_{\Gamma(\xi) \setminus r_0\mathbb{D}} \frac{d\mu_g(z)}{1-|z|^2} \right)^{\frac{p}{p-q}} d\xi \right]^{\frac{p-q}{p}} < \varepsilon.$$

Moreover, there exists $k_0 \in \mathbb{N}$ such that for any $j > k_0$, $|f_j|^q < \varepsilon$ on $r_0\mathbb{D}$. It follows from [Gar07, Theorem 3.1 in Chapter II], Hölder's inequality, Fubini's theorem and (2.6), that

$$\begin{aligned} \int_{\mathbb{D}} |f_j(z)|^q d\mu_g(z) &\lesssim \varepsilon + \int_{\mathbb{D} \setminus r_0\mathbb{D}} \frac{|f_j(z)|^q}{1-|z|^2} \int_{I(z)} d\xi d\mu_g(z) \\ &\leq \varepsilon + \left(\int_{\mathbb{T}} \sup_{z \in \Gamma(\xi)} |f_j(z)|^p d\xi \right)^{\frac{q}{p}} \left[\int_{\mathbb{T}} \left(\int_{\Gamma(\xi) \setminus r_0\mathbb{D}} \frac{d\mu_g(z)}{1-|z|^2} \right)^{\frac{p}{p-q}} d\xi \right]^{\frac{p-q}{p}} \\ &\lesssim \varepsilon + \varepsilon \|f_j\|_{H^p}^q \lesssim \varepsilon. \end{aligned}$$

Thus, we get the desired conclusion and complete the proof. \square

ACKNOWLEDGEMENTS

This work is partially supported by National Natural Science Foundation of China.

REFERENCES

- [AC01] Alexandru Aleman, Joseph A. Cima, An integral operator on H^p and Hardy's inequality, *J. Anal. Math.* 85 (2001) 157–176.
- [Ars99] Miloš Arsenović, Embedding derivatives of \mathcal{M} -harmonic functions into L^p -spaces, *Rocky Mountain J. Math.* 29 (1) (1999) 61–76.
- [AS95] Alexandru Aleman, Aristomenis G. Siskakis, An integral operator on H^p , *Complex Variables Theory Appl.* 28 (2) (1995) 149–158.
- [AS97] Alexandru Aleman, Aristomenis G. Siskakis, Integration operators on Bergman spaces, *Indiana Univ. Math. J.* 46 (2) (1997) 337–356.
- [CPW21] Jiale Chen, Jordi Pau, Maofa Wang, Essential norms and Schatten(-Herz) classes of integration operators from Bergman spaces to Hardy spaces, *Results Math.* 76 (2) (2021) Paper No. 88, 33 pp.
- [Dur70] Peter L. Duren, *Theory of H^p Spaces*, Pure and Applied Mathematics, vol. 38, Academic Press, New York-London, 1970.
- [Dur04] Peter L. Duren, Alexander Schuster, *Bergman Spaces*, Math. Surveys Monogr., vol. 100, American Mathematical Society, Providence, RI, 2004.
- [Gar07] John B. Garnett, *Bounded Analytic Functions*, Revised first edition, Graduate Texts in Mathematics, vol. 236, Springer, New York, 2007.
- [Lue93] Daniel H. Luecking, Embedding theorems for spaces of analytic functions via Khinchine's inequality, *Michigan Math. J.* 40 (2) (1993) 333–358.
- [MPPW20] Santeri Miihkinen, Jordi Pau, Antti Perälä, Maofa Wang, Volterra type integration operators from Bergman spaces to Hardy spaces, *J. Funct. Anal.* 279 (4) (2020) 108564, 32 pp.

- [Pau16] Jordi Pau, Integration operators between Hardy spaces on the unit ball of \mathbb{C}^n , *J. Funct. Anal.* 270 (1) (2016) 134–176.
- [Pav13] Miroslav Pavlović, On the Littlewood-Paley g -function and Calderón’s area theorem, *Expo. Math.* 31 (2) (2013) 169–195.
- [Pom77] Ch. Pommerenke, Schlichte Funktionen und analytische Funktionen von beschränkter mittlerer Oszillation, *Comment. Math. Helv.* 52 (4) (1977) 591–602.
- [PP10] Jordi Pau, José Ángel Peláez, Embedding theorems and integration operators on Bergman spaces with rapidly decreasing weights, *J. Funct. Anal.* 259 (10) (2010) 2727–2756.
- [PR14] José Ángel Peláez, Jouni Rättyä, Weighted Bergman spaces induced by rapidly increasing weights, *Mem. Amer. Math. Soc.* 227 (1066) (2014) vi+124.
- [PR16] José Ángel Peláez, Jouni Rättyä, Two weight inequality for Bergman projection, *J. Math. Pures Appl.* (9) 105 (1) (2016) 102–130.
- [PR19] José Ángel Peláez, Jouni Rättyä, Bergman projection induced by radial weight, arXiv:1902.09837v2, Sep 2019.
- [PRS18] José Ángel Peláez, Jouni Rättyä, Kian Sierra, Berezin transform and Toeplitz operators on Bergman spaces induced by regular weights, *J. Geom. Anal.* 28 (1) (2018) 656–687.
- [PRS21] José Ángel Peláez, Jouni Rättyä, Kian Sierra, Atomic decomposition and Carleson measures for weighted mixed norm spaces, *J. Geom. Anal.* 31 (1) (2021) 715–747.
- [Sis00] Aristomenis G. Siskakis, Weighted integrals of analytic functions, *Acta Sci. Math. (Szeged)* 66 (3-4) (2000) 651–664.
- [Sis06] Aristomenis G. Siskakis, Volterra operators on spaces of analytic functions—a survey, In *Proceedings of the First Advanced Course in Operator Theory and Complex Analysis*, Univ. Sevilla Secr. Publ., Seville, (2006) 51–68.
- [Wu11] Zhijian Wu, Volterra operator, area integral and Carleson measures, *Sci. China Math.* 54 (11) (2011) 2487–2500.
- [Zhu07] Kehe Zhu, *Operator Theory in Function Spaces*, second ed., *Math. Surveys Monogr.*, vol. 138, American Mathematical Society, Providence, RI, 2007.

YONGJIANG DUAN: SCHOOL OF MATHEMATICS AND STATISTICS, NORTHEAST NORMAL UNIVERSITY, CHANGCHUN, JILIN 130024, P.R.CHINA
Email address: duanyj086@nenu.edu.cn

SIYU WANG: SCHOOL OF MATHEMATICS AND STATISTICS, NORTHEAST NORMAL UNIVERSITY, CHANGCHUN, JILIN 130024, P.R.CHINA
Email address: wangsy696@nenu.edu.cn

ZIPENG WANG: COLLEGE OF MATHEMATICS AND STATISTICS, CHONGQING UNIVERSITY, CHONGQING, 401331, P.R.CHINA
Email address: zipengwang@cqu.edu.cn