

Algorithms for normalized multiple sequence alignments

Eloi Araujo^{1,*} Diego P. Rubert¹
 Luiz Rozante² Fábio V. Martinez¹

¹ Faculdade de Computação
 Universidade Federal de Mato Grosso do Sul
 Brazil

² Centro de Matemática, Computação e Cognição
 Universidade Federal do ABC
 Brazil

December 6, 2021

Abstract

Sequence alignment supports numerous tasks in bioinformatics, natural language processing, pattern recognition, social sciences, and other fields. While the alignment of two sequences may be performed swiftly in many applications, the simultaneous alignment of multiple sequences proved to be naturally more intricate. Although most multiple sequence alignment (MSA) formulations are NP-hard, several approaches have been developed, as they can outperform pairwise alignment methods or are necessary for some applications. Taking into account not only similarities but also the lengths of the compared sequences (i.e. normalization) can provide better alignment results than both unnormalized or post-normalized approaches. While some normalized methods have been developed for pairwise sequence alignment, none have been proposed for MSA. This work is a first effort towards the development of normalized methods for MSA. We discuss multiple aspects of normalized multiple sequence alignment (NMSA). We define three new criteria for computing normalized scores when aligning multiple sequences, showing the NP-hardness and exact algorithms for solving the NMSA using those criteria. In addition, we provide approximation algorithms for MSA and NMSA for some classes of scoring matrices.

Keywords: Multiple sequence alignment (MSA), Normalized multiple sequence alignment (NMSA), Algorithms and complexity

1 Introduction

Sequence alignment lies at the foundation of bioinformatics. Several procedures rely on alignment methods for a range of distinct purposes, such as detection of sequence homology, secondary structure prediction, phylogenetic analysis, identification of conserved motifs or genome assembly. On the other hand, alignment techniques have also been reshaped and found applications in other fields, such as natural language processing, pattern recognition, or social sciences [AT00, AG97, BL02, MV93].

Given its range of applications in bioinformatics, extensive efforts have been made to improve existing or developing novel methods for sequence alignment. The simpler ones compare a pair of sequences in polynomial time on their lengths, usually trying to find editing operations (insertions, deletions, and substitutions of symbols) that transform one sequence into another while maximizing or minimizing some objective function called edit distance [HAR09]. This

*Corresponding author

concept can naturally be generalized to align multiple sequences [WLXZ15], adding another new layer of algorithmic complexity, though. In this case, most multiple sequence alignment (MSA) formulations lead to NP-hard problems [Eli06]. Nevertheless, a variety of methods suitable for aligning multiple sequences have been developed, as they can outperform pairwise alignment methods on tasks such as phylogenetic inference [OR06], secondary structure prediction [CB99] or identification of conserved regions [SWD+11].

In order to overcome the cost of exact solutions, a number of MSA heuristics have been developed in recent years, most of them using the so-called progressive or iterative methods [HTHI95, SH14, TPP99, WOHN06]. Experimental data suggest that the robustness and accuracy of heuristics can still be improved, however [WLXZ15].

Most approaches for pairwise sequence alignment define edit distances as absolute values, lacking some normalization that would result in edit distances relative to the lengths of the sequences. However, some applications may require sequence lengths to be taken into account. For instance, a difference of one symbol between sequences of length 5 is more significant than between sequences of length 1000. In addition, experiments suggest that normalized edit distances can provide better results than both unnormalized or post-normalized edit distances [MV93]. While normalized edit distances have been developed for pairwise sequence alignment [AE99, MV93], none have been proposed for MSA to the best of our knowledge.

In this work, we propose exact and approximation algorithms for normalized MSA (NMSA). This is a first step towards the development of methods that take into account the lengths of sequences for computing edit distances when multiple sequences are compared. The remainder of this paper is organized as follows. Section 2 introduces concepts related to sequence alignment and presents normalized scores for NMSA, followed by the complexity analysis of NMSA using those scores in Section 3. Next, Sections 4 and 5 describe exact and approximation algorithms, respectively. Section 6 closes the paper with the conclusion and prospects for future work.

2 Preliminaries

An *alphabet* Σ is a finite non-empty set of *symbols*. A finite sequence s with n symbols in Σ is seen as $s(1) \cdots s(n)$. We say that the *length* of s , denoted by $|s|$, is n . The sequence $s(p) \cdots s(q)$, with $1 \leq p \leq q \leq n$, is denoted by $s(p:q)$. If $p > q$, $s(p:q)$ is the *empty* sequence which length is equal to zero and it is denoted by ε . We denote the sequence resulting from the concatenation of sequences s and t by $s \cdot t = st$. A sequence of n symbols \mathbf{a} is denoted by \mathbf{a}^n . A k -tuple S over Σ^* is called a k -*sequence* and we write s_1, \dots, s_k to refer to S , where s_i is the i -th sequence in S . We denote $S = \emptyset$ if every sequence of S is a empty sequence.

Let $\Sigma_- := \Sigma \cup \{-\}$, where $- \notin \Sigma$ and the symbol $-$ is called a *space*. Let $S = s_1, \dots, s_k$ be a k -sequence. An *alignment* of S is a k -tuple $A = [s'_1, \dots, s'_k]$ over Σ_-^* , where

- (a) each sequence s'_h is obtained by inserting spaces in s_h ;
- (b) $|s'_h| = |s'_i|$ for each pair h, i , with $1 \leq h, i \leq k$;
- (c) there is no j in $\{1, \dots, k\}$ such that $s'_1(j) = \dots = s'_k(j) = -$.

Notice that alignments, which are k -tuples over Σ_-^* , are written enclosed by square brackets “[]”. The sequence $A(j) = [s'_1(j), \dots, s'_k(j)] \in \Sigma_-^k$ is the *column* j of the alignment $A = [s'_1, \dots, s'_k]$ and $A[j_1 : j_2]$ is the alignment defined by the columns $j_1, j_1 + 1, \dots, j_2$ of A . We say that the pair $[s'_h(j), s'_i(j)]$ *aligns* in A or, simply, that $s'_h(j)$ and $s'_i(j)$ are *aligned* in A , and $|A| = |s'_i|$ is the *length* of the alignment A . It is easy to check that $\max_i \{|s_i|\} \leq |A| \leq \sum_i |s_i|$. We denote by \mathcal{A}_S the set of all alignments of S .

An alignment can be used to represent *editing operations* of *insertions*, *deletions* and *substitutions* of symbols in sequences, where the symbol $-$ represents insertions or deletions. An alignment can also be represented in the matrix

format by placing one sequence above another. Thus, the alignments

$$[\mathbf{aaa-}, \mathbf{ab--}, -\mathbf{cac}] \quad \text{and} \quad [-\mathbf{aaa-}, \mathbf{ab---}, -\mathbf{ca-c}]$$

of 3-sequence $\mathbf{aaa}, \mathbf{ab}, \mathbf{cac}$ can be represented respectively as

$$\begin{bmatrix} \mathbf{a} & \mathbf{a} & \mathbf{a} & - \\ \mathbf{a} & \mathbf{b} & - & - \\ - & \mathbf{c} & \mathbf{a} & \mathbf{c} \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} - & \mathbf{a} & \mathbf{a} & \mathbf{a} & - \\ \mathbf{a} & \mathbf{b} & - & - & - \\ - & \mathbf{c} & \mathbf{a} & - & \mathbf{c} \end{bmatrix}.$$

Let $I = \{i_1, \dots, i_m\} \subseteq \{1, \dots, k\}$ be a set of indices such that $i_1 < \dots < i_m$ and let $A = [s'_1, \dots, s'_k]$ be an alignment of a k -sequence $S = s_1, \dots, s_k$. We write S_I to denote the m -sequence s_{i_1}, \dots, s_{i_m} . The alignment of S_I induced by A is the alignment A_I obtained from the alignment A , considering only the corresponding sequences in S_I and, from the resulting structure, removing columns where all symbols are $-$. In the previous example, being $A = [\mathbf{aaa-}, \mathbf{ab--}, -\mathbf{cac}]$ an alignment of $\mathbf{aaa}, \mathbf{ab}, \mathbf{cac}$, we have

$$\begin{bmatrix} \mathbf{a} & \mathbf{a} & \mathbf{a} \\ \mathbf{a} & \mathbf{b} & - \end{bmatrix}$$

is an alignment of $\mathbf{aaa}, \mathbf{ab}$ induced by A .

A k -vector $\vec{j} = [j_1, \dots, j_k]$ is a k -tuple, where $j_i \in \mathbb{N} = \{0, 1, 2, \dots\}$. We say that j_i is the i -th element of \vec{j} . The k -vector $\vec{0}$ is such that all its elements are zero. If \vec{j} and \vec{h} are k -vectors, we write $\vec{j} \leq \vec{h}$ if $j_i \leq h_i$ for each i ; and $\vec{j} < \vec{h}$ if $\vec{j} \leq \vec{h}$ and $\vec{j} \neq \vec{h}$. A sequence of k -vectors $\vec{j}_1, \vec{j}_2, \dots$ is in *topological order* if $\vec{j}_i < \vec{j}_h$ implies $i < h$. Given two k -vectors, say $\vec{a} = [a_1, a_2 \dots a_k]$ $\vec{b} = [b_1, b_2 \dots b_k]$, we say that \vec{a} *precedes* \vec{b} when there exists $l \in \mathbb{N}$ such that $a_i = b_i$ for each $i > l$ and $a_l < b_l$. A sequence of k -vectors $\vec{j}_1, \vec{j}_2, \dots$ is in *lexicographical order* if \vec{j}_i precedes \vec{j}_h for each $i < h$. Clearly, a lexicographical order is a special case of topological order.

Consider $S = s_1, \dots, s_k$ a k -sequence with $n_i = |s_i|$ for each i and we call $\vec{n} = [n_1, \dots, n_k]$ the *length of S* . Let $V_{\vec{n}} = \{\vec{j} : \vec{j} \leq \vec{n}\}$. Therefore, $|V_{\vec{n}}| = \prod_i (|s_i| + 1)$ which implies that if $n_i = n$ for all i , then $|V_{\vec{n}}| = (n + 1)^k$. Define $S(\vec{j}) = s_1(j_1), \dots, s_k(j_k)$ a *column \vec{j} in S* and we say that $S(1:\vec{j}) = s_1(1:j_1), \dots, s_k(1:j_k)$ is the *prefix of S ending in \vec{j}* . Thus, $S = S(1:\vec{n})$. Besides that, if A is an alignment and $\vec{j} = [j, j, \dots, j]$, then $A[\vec{j}] = A(j)$.

Denote by \mathcal{B}^k the set of k -vectors $[b_1, \dots, b_k]$, where $b_i \in \{0, 1\}$ for each i . Now, for $\vec{b} \leq \vec{j}$, where $\vec{j} = [j_1, \dots, j_k]$, define

$$\vec{b} \cdot S(\vec{j}) = [x_1, \dots, x_k] \in \Sigma^k,$$

such that

$$x_i = \begin{cases} s_i(j_i), & \text{if } b_i = 1 \\ -, & \text{otherwise.} \end{cases}$$

Therefore, given an alignment A of $S(1:\vec{n})$, there exists $\vec{b} \in \mathcal{B}^k$, with $\vec{b} \leq \vec{n}$, such that $A(|A|) = \vec{b} \cdot S(\vec{n})$. In this case, notice that $b_i = 1$ if and only if $s_i(n_i)$ is in the i -th row of the last column of A and we say that \vec{b} *defines* the column $|A|$ of alignment A . For $\vec{b} \leq \vec{j}$, we also define the operation

$$\vec{j} - \vec{b} = [j_1 - b_1, \dots, j_k - b_k].$$

Notice that $|\mathcal{B}^k| = 2^k$. Figure 1 shows an example of alignment using vectors to define columns and operations.

For a problem \mathbf{P} , we call $\mathbb{I}_{\mathbf{P}}$ the set of instances of \mathbf{P} . If \mathbf{P} is a decision problem, then $\mathbf{P}(I) \in \{\mathbf{Yes}, \mathbf{No}\}$ is the image of an instance I . If \mathbf{P} is an optimization (minimization) problem, there is a set $\text{Sol}(I)$ for each instance I , a function v defining a non-negative rational number for each $X \in \text{Sol}(I)$, and a function $\text{opt}_v(I) = \min_{X \in \text{Sol}(I)} \{v(X)\}$. We use opt instead of opt_v if v is obvious. Let $\mathbf{A}(I) = v(X \in \text{Sol}(I))$ a solution computed by an algorithm \mathbf{A} with input I . We say that \mathbf{A} is an α -*approximation* for \mathbf{P} if $\mathbf{A}(I) \leq \alpha \text{opt}(I)$ for each $I \in \mathbb{I}_{\mathbf{P}}$. We say that α is an *approximation factor* for \mathbf{P} .

$$A = \begin{bmatrix} \mathbf{a} & \mathbf{b} & \mathbf{c} & - & - \\ - & \mathbf{b} & \mathbf{c} & \mathbf{a} & - \\ \mathbf{b} & - & \mathbf{b} & - & \mathbf{a} \\ \mathbf{a} & \mathbf{a} & \mathbf{a} & \mathbf{a} & \mathbf{a} \\ \mathbf{c} & - & - & - & - \end{bmatrix} \Rightarrow \begin{bmatrix} A[1 : 4] \end{bmatrix} \begin{bmatrix} [0, 0, 1, 1, 0] \cdot S(1 : [3, 3, 3, 5, 1]) \end{bmatrix}$$

Figure 1: The length of alignment A of $S = \mathbf{abc}, \mathbf{bca}, \mathbf{bba}, \mathbf{aaaaa}, \mathbf{c}$ is $\vec{n} = [|\mathbf{abc}|, |\mathbf{bca}|, |\mathbf{bba}|, |\mathbf{aaaaa}|, |\mathbf{c}|] = [3, 3, 3, 5, 1]$. Here, $S(\vec{n}) = [\mathbf{c}, \mathbf{a}, \mathbf{a}, \mathbf{a}, \mathbf{c}]$ and since $\vec{b}_5 = [0, 0, 1, 1, 0]$ define the last column of A , we have $S(\vec{n} - \vec{b}_5) = \mathbf{abc}, \mathbf{bca}, \mathbf{bb}, \mathbf{aaaa}, \mathbf{c}$. The alignment defined by the first four columns of A is $A[1 : 4]$ which is an alignment of $S(1 : \vec{n} - \vec{b}_5)$ and the last column of A is $\vec{b}_5 \cdot S(\vec{n})$. It is interesting to note that A can be completely defined by 5-vectors $\vec{b}_1 = [1, 0, 1, 1, 1]$, $\vec{b}_2 = [1, 1, 0, 1, 0]$, $\vec{b}_3 = [1, 1, 1, 1, 0]$, $\vec{b}_4 = [0, 1, 0, 1, 0]$, $\vec{b}_5 = [0, 0, 1, 1, 0]$ where b_i represents column i of alignment A .

The *alignment problem* is a collection of decision and optimization problems whose instances are finite subsets of Σ^* and $\text{Sol}(S) = \mathcal{A}_S$ for each instance S . Function v , used for scoring alignments, is called *criterion* for \mathbf{P} and we call $v[A]$ the *cost of the alignment* A . The *v -optimal alignment* A of S is such that $v[A] = \text{opt}(S)$. Thus, we state the following general optimization problems using the criterion v and integer k :

Problem 1 (Alignment with criterion v). Given a k -sequence S , find the cost of a v -optimal alignment of S .

We also need the decision version of the alignment problem with criterion v where we are given a k -sequence S and a number $d \in \mathbb{Q}_{\geq}$, and we want to decide whether there exists an alignment A of S such that $v[A] \leq d$.

Usually the cost of an alignment v is defined from a scoring matrix. A *scoring matrix* γ is a matrix whose elements are rational numbers. The matrix has rows and columns indexed by symbols in Σ_- . For $\mathbf{a}, \mathbf{b} \in \Sigma_-$ and a scoring matrix γ , we denote by $\gamma_{\mathbf{a} \rightarrow \mathbf{b}}$ the entry of γ in row \mathbf{a} and column \mathbf{b} . The value $\gamma_{\mathbf{a} \rightarrow \mathbf{b}}$ defines the score for a substitution if $\mathbf{a}, \mathbf{b} \in \Sigma$, for an insertion if $\mathbf{a} = -$, and for a deletion if $\mathbf{b} = -$. The entry $\gamma_{- \rightarrow -}$ is not defined.

2.1 vA_γ - and vN_γ -score for scoring alignments of 2-sequences

Consider a scoring matrix γ . Let $S = s, t \in \Sigma^*$ be a 2-sequence whose length is $\vec{n} = [n = |s|, m = |t|]$. A simple criterion for scoring alignments of 2-sequences using the function vA_γ follows. For an alignment $[s', t']$ of s, t we define

$$vA_\gamma[s', t'] = \sum_{j=1}^{[s', t']} \gamma_{s'(j) \rightarrow t'(j)}.$$

We say that $vA_\gamma[s', t']$ is a vA_γ -score of s, t . The optimal function for this criterion is called *unweighted edit distance* and denoted by $\text{opt}A_\gamma$, and an $\text{opt}A_\gamma$ -optimal alignment is also called an *A-optimal alignment*. When $\gamma_{\mathbf{a} \rightarrow \mathbf{a}} = 0$ and $\gamma_{\mathbf{a} \rightarrow \mathbf{b}} = 1$ for each $\mathbf{a} \neq \mathbf{b} \in \Sigma_-$, $\text{opt}A_\gamma$ is also known as *Levenshtein distance* [Lev66].

Now, suppose that $n \geq m$. Needleman and Wunch [NW70] proposed an $O(n^2)$ -time algorithm for computing

$$\text{opt}A_\gamma(s, t) = \text{opt}A_\gamma(S) = \min_{A \in \mathcal{A}_S} vA_\gamma[A] = \begin{cases} 0 & \text{if } S = \emptyset, \\ \min_{\vec{b} \in \mathcal{B}^k, \vec{b} \leq \vec{n}} \left\{ \text{opt}A_\gamma(S[1 : \vec{n} - \vec{b}]) + vA_\gamma[\vec{b} \cdot S] \right\} & \text{otherwise.} \end{cases}$$

If $\text{opt}A_\gamma$ is a Levenshtein distance, Masek and Paterson [MP80] presented an $O(n^2 / \log n)$ -time algorithm using the “*Four Russian’s Method*”. Crochemore, Landau and Ziv-Ukelson [CLZU03] extended $\text{opt}A_\gamma$ -score supporting scoring matrices with real numbers and describing an $O(n^2 / \log n)$ -time algorithm. Indeed, there is no algorithm to determine $\text{opt}A_\gamma(s, t)$ in $O(n^{2-\delta})$ -time for any $\delta > 0$, unless SETH is false [BI18]. Andoni, Krauthgamer and Onak [AKO10] described a nearly linear time algorithm approximating the edit distance within an approximation factor $\text{poly}(\log n)$. Later, Chakraborty et al. [CDG+20] presented an $O(n^{2-2/7})$ -time α -approximation for edit distance, where α is constant.

Marzal and Vidal [MV93] defined another criterion for scoring alignments of two sequences called vN_γ -score, which is a normalization of vA_γ -score, as follows:

$$vN_\gamma[A] = \begin{cases} 0, & \text{if } |A| = 0, \\ vA_\gamma[A]/|A|, & \text{otherwise.} \end{cases}$$

The optimal function for this criterion as known as *normalized edit distance* and it is denoted by $\text{opt}N_\gamma$, and an $\text{opt}N_\gamma$ -optimal alignment is also called a *N-optimal alignment* of $S = s, t$.

A naive dynamic programming algorithm was proposed by Marzal and Vidal [MV93] to obtain an N-optimal alignment of two sequences in $O(n^3)$ -time. Using fractional programming, Vidal, Marzal and Aibar [VMA95] presented an algorithm with running time $O(n^3)$, but requiring only $O(n^2)$ -time in practice which is similarly to the classical (unnormalized) edit distance algorithm. Further, Arslan and Egecioglu [AE99] described an $O(n^2 \log n)$ -time algorithm to this problem.

Let $S = s, t$ a 2-sequence whose length is \vec{n} . A simple heuristic for determining a close value for $\text{opt}N_\gamma(s, t)$ computes first $\text{opt}A_\gamma(s, t)$ and then divides it by the maximum length $L(\vec{n}) = L(S)$ of an optimum alignment of S . In order to find $L(S)$, we use the same dynamic programming strategy for computing $\text{opt}A_\gamma(s, t)$, i.e.,

$$L(S) = L(\vec{n}) = \begin{cases} 0 & \text{if } S = \emptyset \\ \max_{\vec{b} \in \mathcal{B}^k, \vec{b} \leq \vec{n}, \text{opt}A_\gamma(S) = \text{opt}A_\gamma(S - \vec{b}) + vA_\gamma(\vec{b} \cdot S)} \left\{ L(\vec{n} - \vec{b}) \right\} + 1 & \text{otherwise,} \end{cases}$$

in $O(nm)$ -time. The following theorem shows that this heuristic is a simple approximation algorithm to find a N-optimal alignment.

Theorem 2.1. *Let s, t be a 2-sequence of length $[n, m]$ and let $L(s, t)$ be the maximum length of an A-optimal alignment of s, t . Then,*

$$\frac{\text{opt}A_\gamma(s, t)}{L(S)} \leq 2 \text{opt}N_\gamma(s, t),$$

and it can be computed in $O(n^2)$ -time if $n = m$. Moreover, this ratio is tight, i.e., for any positive rational ε , there exists a scoring matrix γ , sequences s, t and an A-optimal alignment of s, t with maximum length A such that

$$\frac{\text{opt}A_\gamma(s, t)}{|A|} = \frac{vA_\gamma[A]}{|A|} = (2 - \varepsilon) \text{opt}N_\gamma(s, t).$$

Proof. Let A be an A-optimal alignment with maximum length computed by the heuristic above in $O(nm)$ -time. Let B be an N-optimal alignment. Thus, $vA_\gamma[A] \leq vA_\gamma[B]$. Moreover, since $|B| \leq n + m$, $n + m \leq 2 \max\{n, m\}$ and $\max\{n, m\} \leq 2|A|$, we have that $|A| \geq |B|/2$. Therefore, $vN_\gamma[A] = \frac{vA_\gamma[A]}{|A|} \leq \frac{vA_\gamma[B]}{|A|} \leq \frac{vA_\gamma[B]}{|B|/2} = 2 \text{opt}N_\gamma(s, t)$.

We present now a 2-sequence s, t and a scoring matrix γ such that the solution given by the heuristic is at least $(2 - \varepsilon) \text{opt}N_\gamma(s, t)$ for any $\varepsilon \in \mathbb{Q}_>$. Let $\Sigma = \{\mathbf{a}, \mathbf{b}\}$, γ be a scoring matrix such that $\gamma_{\mathbf{a} \rightarrow -} = \gamma_{\mathbf{b} \rightarrow -} = 1/\varepsilon$ and $\gamma_{\mathbf{a} \rightarrow \mathbf{b}} = 2/\varepsilon - 1$ and $\mathbf{a}^n, \mathbf{b}^n \in \Sigma^*$, where n in a positive integer. Observe that the vA_γ -score of any alignment of $(\mathbf{a}^n, \mathbf{b}^n)$, where $[\mathbf{a}, \mathbf{b}]$ is aligned in k columns, is equal to $2n/\varepsilon - k$. Thus, $\text{opt}A_\gamma(\mathbf{a}^n, \mathbf{b}^n) = \min_{0 \leq k \leq n} \{2n/\varepsilon - k\} = 2n/\varepsilon - n = (2/\varepsilon - 1)n$ which implies $[\mathbf{a}^n, \mathbf{b}^n]$ is the A-optimal alignment with maximum length. Since $\text{opt}N_\gamma(\mathbf{a}^n, \mathbf{b}^n) \leq vN_\gamma([\mathbf{a}^{n-n}, -^n\mathbf{b}^n]) = 1/\varepsilon$, it follows

$$\frac{\text{opt}A_\gamma(\mathbf{a}^n, \mathbf{b}^n)}{|\mathbf{a}^n, \mathbf{b}^n|} = \frac{vA_\gamma[\mathbf{a}^n, \mathbf{b}^n]}{|\mathbf{a}^n, \mathbf{b}^n|} = \frac{(2/\varepsilon - 1)n}{n} = \frac{2 - \varepsilon}{\varepsilon} = (2 - \varepsilon) vN_\gamma([\mathbf{a}^{n-n}, -^n\mathbf{b}^n]) \geq (2 - \varepsilon) \text{opt}N_\gamma(\mathbf{a}^n, \mathbf{b}^n).$$

□

We define now classes of scoring matrices. The most common class of scoring matrices \mathbb{M}^C has the following properties: for all $\mathbf{a}, \mathbf{b}, \mathbf{c} \in \Sigma_-$, we have (a) $\gamma_{\mathbf{a} \rightarrow \mathbf{b}} > 0$ if $\mathbf{a} \neq \mathbf{b}$, and $\gamma_{\mathbf{a} \rightarrow \mathbf{b}} = 0$ if $\mathbf{a} = \mathbf{b}$; (b) $\gamma_{\mathbf{a} \rightarrow \mathbf{b}} = \gamma_{\mathbf{b} \rightarrow \mathbf{a}}$; and (c) $\gamma_{\mathbf{a} \rightarrow \mathbf{c}} \leq \gamma_{\mathbf{a} \rightarrow \mathbf{b}} + \gamma_{\mathbf{b} \rightarrow \mathbf{c}}$.

The class \mathbb{M}^W of scoring matrices is such that, for all symbols $\mathbf{a}, \mathbf{b}, \mathbf{c} \in \Sigma$, we have (a) $\gamma_{\mathbf{a} \rightarrow -} = \gamma_{- \rightarrow \mathbf{a}} > 0$; (b) $\gamma_{\mathbf{a} \rightarrow \mathbf{b}} > 0$ if $\mathbf{a} \neq \mathbf{b}$, and $\gamma_{\mathbf{a} \rightarrow \mathbf{b}} = 0$ if $\mathbf{a} = \mathbf{b}$; (c) if $\gamma_{\mathbf{a} \rightarrow \mathbf{b}} < \gamma_{\mathbf{a} \rightarrow -} + \gamma_{- \rightarrow \mathbf{b}}$, then $\gamma_{\mathbf{a} \rightarrow \mathbf{b}} = \gamma_{\mathbf{b} \rightarrow \mathbf{a}}$; (d) $\gamma_{\mathbf{a} \rightarrow -} \leq \gamma_{\mathbf{a} \rightarrow \mathbf{b}} + \gamma_{\mathbf{b} \rightarrow -}$; and (e) $\min\{\gamma_{\mathbf{a} \rightarrow \mathbf{c}}, \gamma_{\mathbf{a} \rightarrow -} + \gamma_{- \rightarrow \mathbf{c}}\} \leq \gamma_{\mathbf{a} \rightarrow \mathbf{b}} + \gamma_{\mathbf{b} \rightarrow \mathbf{c}}$. Moreover, the class \mathbb{M}^N is such that (a) $\mathbb{M}^N \subseteq \mathbb{M}^W$ and (b) $\gamma_{\mathbf{a} \rightarrow -} \leq 2\gamma_{\mathbf{b} \rightarrow -}$ for each $\mathbf{a}, \mathbf{b} \in \Sigma$.

For a set S , we say that $f : S \times S \rightarrow \mathbb{R}$ is a *distance function* or *metric on S* if f satisfies for all $s, t, u \in S$:

1. $f(s, s) = 0$ (*reflexivity*);
2. $f(s, t) > 0$ if $s \neq t$ (*strict positiveness*);
3. $f(s, t) = f(t, s)$ (*symmetry*); and
4. $f(s, u) \leq f(s, t) + f(t, u)$ (*triangle inequality*).

If a given criterion v depends on a scoring matrix γ and it is a metric on Σ^* , we say that the scoring matrix γ *induces* a opt_v -distance on Σ^* . Sellers [Sel74] showed that matrices in \mathbb{M}^C induce an opt_{A_γ} -distance on Σ^* . Araujo and Soares [AS06] showed that $\gamma \in \mathbb{M}^W$ and $\gamma \in \mathbb{M}^N$ if and only if γ induces an opt_{A_γ} -distance and an opt_{N_γ} -distance on Σ^* respectively. Figure 2 shows the relationship between these classes.




Figure 2: Relationship between scoring matrices. Araujo and Soares [AS06] showed that $\mathbb{M}^C \subseteq \mathbb{M}^W$, $\mathbb{M}^N \subseteq \mathbb{M}^W$, $\mathbb{M}^C \not\subseteq \mathbb{M}^N$ and $\mathbb{M}^N \not\subseteq \mathbb{M}^C$. Moreover, the scoring matrix γ such that $\gamma_{\mathbf{a} \rightarrow \mathbf{a}} = 0$ for each \mathbf{a} and $\gamma_{\mathbf{a} \rightarrow \mathbf{b}} = 1$ for each $\mathbf{a} \neq \mathbf{b}$ is in $\mathbb{M}^C \cap \mathbb{M}^N$, which implies that $\mathbb{M}^C \cap \mathbb{M}^N \neq \emptyset$.

Given a scoring function v for alignments of 2-sequences s, t that depends on a scoring matrix, we say that two scoring matrices γ and ρ are *equivalent* considering v when $v_\gamma[A] \leq v_\gamma[B]$ if and only if $v_\rho[A] \leq v_\rho[B]$ for any pair of alignments A, B of sequences s, t . If ρ is a matrix obtained from γ by multiplying each entry of γ by a constant $c > 0$, then $v_{A_\rho}[A] = c \cdot v_{A_\gamma}[A]$ and $v_{N_\rho}[A] = c \cdot v_{N_\gamma}[A]$, which implies that γ and ρ are equivalent. As a consequence, when the scoring function is v_{A_γ} or v_{N_γ} and it is convenient, we can suppose that all entries of γ are integers instead of rationals.

2.2 $v\text{SP}_\gamma$ -score for k sequences

Consider a scoring matrix γ . Let $S = s_1, \dots, s_k$ be a k -sequence whose length is \vec{n} and $A = [s'_1, \dots, s'_k]$ be an alignment of S . The criterion $v\text{SP}_\gamma$, also called *SP-score*, for scoring the alignment A is

$$v\text{SP}_\gamma[A] = \sum_{h=1}^{k-1} \sum_{i=h+1}^k v_{A_\gamma}[A_{\{h,i\}}]. \quad (1)$$

We define optSP_γ as the optimal function for the criterion $v\text{SP}_\gamma$. An alignment A of S such that $v\text{SP}_\gamma[A] = \text{optSP}_\gamma(S)$ is called *$v\text{SP}_\gamma$ -optimal alignment*. Regardless its decision or optimization version, we call the associated problem as *multiple sequence alignment problem (MSA)*. Formally,

Problem 2 (Multiple sequence alignment). Let γ be a fixed scoring matrix. Given a k -sequence S , find $\text{optSP}_\gamma(S)$.

In order to compute optSP_γ , we extend the definition of $v\text{SP}_\gamma$ considering a column of an alignment $A = [s'_1, \dots, s'_k]$ as its parameter. Thus, $v\text{SP}_\gamma[A(j)] = v\text{SP}_\gamma[A[\vec{j}]] = \sum_{i < h} \gamma_{s'_i(j) \rightarrow s'_h(j)}$ assuming that $\gamma_{- \rightarrow -} = 0$ and $\vec{j} = [j, \dots, j]$ and

$$\text{optSP}_\gamma(S) = \min_{A \in \mathcal{A}_S} v\text{SP}_\gamma[A] = \begin{cases} 0 & \text{if } S = \emptyset \\ \min_{\vec{b} \in \mathcal{B}^k, \vec{b} \leq \vec{n}} \left\{ \text{optSP}_\gamma(S(1: \vec{n} - \vec{b})) + v\text{SP}_\gamma[\vec{b} \cdot S(\vec{n})] \right\} & \text{otherwise.} \end{cases} \quad (2)$$

Recurrence (2) can be computed using a dynamic programming algorithm, obtaining $D(\vec{j}) = \text{optSP}_\gamma(S(1: \vec{j}))$ for all $\vec{j} \leq \vec{n}$. This task can be performed by generating all indexes of D in lexicographical order, starting with $D(\vec{0}) = 0$, as presented in Algorithm 1.

Algorithm 1

Input: $S = s_1, \dots, s_k \in (\Sigma^*)^k$

Output: $\text{optSP}_\gamma(S)$

- 1: $D(\vec{0}) \leftarrow 0$
 - 2: **for** each $\vec{j} \leq \vec{n}$ in lexicographical order **do**
 - 3: $D(\vec{j}) \leftarrow \min_{\vec{b} \in \mathcal{B}^k, \vec{b} \leq \vec{j}} \{D(\vec{j} - \vec{b}) + v\text{SP}_\gamma[\vec{b} \cdot S(\vec{j})]\}$
 - 4: **return** $D(\vec{n})$
-

Suppose that $|s_i| = n$ for each i . Notice that the space to store the matrix D is $\Theta((n+1)^k)$ and thus Algorithm 1 uses $\Theta((n+1)^k)$ -space. Besides that, Algorithm 1 checks, in the worst case, $\Theta(2^k)$ entries for computing all entries in the matrix D and each computation spends $\Theta(k^2)$ -time. Therefore, its running time is $O(2^k k^2 (n+1)^k)$. Observe that when the distance is small, not all entries in D need to be computed, such as in the Carrillo and Lipman's algorithm [CL88].

We can also describe an obvious but unusual variant of this problem that consider a scoring matrix for each pair of the sequences. Formally, considering an scoring matrix array $\vec{\gamma} = [\gamma^{(12)}, \gamma^{(13)}, \dots, \gamma^{(1k)}, \gamma^{(23)}, \dots, \gamma^{(2k)}, \dots, \gamma^{((k-1)k)}]$ and an alignment A of a k -sequence,

$$v\text{SP}_{\vec{\gamma}}[A] = \sum_{h=1}^{k-1} \sum_{i=h+1}^k vA_{\gamma^{(hi)}}[A_{\{h,i\}}].$$

Thus, given a k -sequence S , we ask for finding $\text{optSP}_{\vec{\gamma}}(S) = \min_{A \in \mathcal{A}_S} \{v\text{SP}_{\vec{\gamma}}[A]\}$. Algorithm 1 can be easily modified in order to solve this extended version with same time and space complexity. This is important here because this version is used as a subroutine of one of the normalized version.

2.3 V_γ^i -score for k sequences

In this section we define a new criteria to normalize the $v\text{SP}_\gamma$ -score of a multiple alignment. The new criteria for aligning sequences takes into account the length of the alignments according to the following:

$$V_\gamma^1[A] = \begin{cases} 0, & \text{if } |A| = 0, \\ v\text{SP}_\gamma[A]/|A|, & \text{otherwise,} \end{cases} \quad (3)$$

$$V_\gamma^2[A] = \sum_{h=1}^{k-1} \sum_{i=h+1}^k vN_\gamma[A_{\{h,i\}}], \quad (4)$$

$$V_\gamma^3[A] = \begin{cases} 0, & \text{if } |A| = 0, \\ v\text{SP}_\gamma[A]/(\sum_{h=1}^{k-1} \sum_{i=h+1}^k |A_{\{h,i\}}|), & \text{otherwise.} \end{cases} \quad (5)$$

We define optNSP_γ^z as the optimal function for the criterion V_γ^z , i.e., for a given k -sequence S

$$\text{optNSP}_\gamma^z(S) = \min_{A \in \mathcal{A}_S} V_\gamma^z[A].$$

$$\begin{array}{c|ccc}
& \mathbf{a} & \mathbf{b} & \mathbf{c} & - \\
\hline
\mathbf{a} & 0 & 9 & 9 & 10 \\
\mathbf{b} & 9 & 0 & 9 & 10 \\
\mathbf{c} & 9 & 9 & 0 & 10 \\
- & 10 & 10 & 10 & 0
\end{array}
\quad
A = \begin{bmatrix} \mathbf{a} \\ \mathbf{b} \\ \mathbf{c} \end{bmatrix}
\quad
B = \begin{bmatrix} \mathbf{a} & - \\ \mathbf{b} & - \\ - & \mathbf{c} \end{bmatrix}
\quad
C = \begin{bmatrix} \mathbf{a} & - & - \\ - & \mathbf{b} & - \\ - & - & \mathbf{c} \end{bmatrix}$$

$$\begin{array}{c|ccc}
& \mathbf{a} & \mathbf{b} & \mathbf{c} & - \\
\hline
\mathbf{a} & 0 & 7 & 7 & 9 \\
\mathbf{b} & 7 & 0 & 7 & 9 \\
\mathbf{c} & 7 & 7 & 0 & 9 \\
- & 9 & 9 & 9 & 0
\end{array}
\quad
D = \begin{bmatrix} \mathbf{a} & \mathbf{b} & \mathbf{c} \\ \mathbf{a} & \mathbf{c} & \mathbf{b} \\ \mathbf{c} & \mathbf{b} & \mathbf{a} \end{bmatrix}
\quad
E = \begin{bmatrix} \mathbf{a} & \mathbf{b} & \mathbf{c} & - \\ \mathbf{a} & - & \mathbf{c} & \mathbf{b} \\ \mathbf{c} & \mathbf{b} & \mathbf{a} & - \end{bmatrix}
\quad
F = \begin{bmatrix} \mathbf{a} & \mathbf{b} & \mathbf{c} & - & - \\ \mathbf{a} & - & \mathbf{c} & \mathbf{b} & - \\ - & - & \mathbf{c} & \mathbf{b} & \mathbf{a} \end{bmatrix}$$

Figure 3: Observe first that A , B or C is an V_γ^z -optimal alignment of 3-sequence $\mathbf{a}, \mathbf{b}, \mathbf{c}$ for each z . Besides, since $V_\gamma^1[A] = 27.0, V_\gamma^2[A] = 27.0, V_\gamma^3[A] = 9.0; V_\gamma^1[B] = 24.5, V_\gamma^2[B] = 29.0, V_\gamma^3[B] = 9.8; V_\gamma^1[C] = 20.0, V_\gamma^2[C] = 30.0, V_\gamma^3[C] = 10.0$, we have that C is an optimal alignment for criterion V_γ^1 but it is not for criteria V_γ^2 or V_γ^3 , which implies that an optimal alignment for criterion V_γ^1 is different when we compare it to criteria V_γ^2 and V_γ^3 . Now, observe that D , E or F is an V_δ^z -optimal alignment of 3-sequence $\mathbf{abc}, \mathbf{acb}, \mathbf{cba}$ for each z . Besides, since $V_\delta^2[D] = 16.33, V_\delta^3[D] = 5.44; V_\delta^2[E] = 17.16, V_\delta^3[E] = 5.81; V_\delta^2[F] = 16.20, V_\delta^3[F] = 5.53$, we have that D is an optimal alignment for criterion V_δ^3 but it is not for criterion V_δ^2 which implies that an optimal alignment for criterion V_δ^2 is different when we compare it to criterion V_δ^3 .

An alignment A of S such that $V_\gamma^z[A] = \text{optNSP}_\gamma^z(S)$ is called V_γ^z -optimal alignment. Moreover, regardless its decision or optimization version, we establish the *criterion V_γ^z for the normalized multiple sequence alignment problem (NMSA- z)*, for $z = 1, 2, 3$. Formally, for a fixed scoring matrix γ and $z \in \{1, 2, 3\}$,

Problem 3 (Normalized multiple sequence alignment with score V_γ^z). Given a k -sequence S , find $\text{optNSP}_\gamma^z(S)$.

An interesting question is whether the definitions above represent the same criterion. i.e., would it be possible that the optimal alignment for a given criterion V_γ^z also represents an optimal alignment for another criterion $V_\gamma^{z'}$, $z \neq z'$, regardless of the sequences and scoring matrices? The answer is no and Figure 3 shows examples that support this claim.

3 Complexity

We study now the complexity of the multiple sequence alignment problem for each new criterion defined in Section 2. We consider the decision version of the computational problems and we prove **NMSA- z** is NP-complete for each z even though the following additional restrictions for the scoring matrix γ hold: $\gamma_{\mathbf{a} \rightarrow \mathbf{b}} = \gamma_{\mathbf{b} \rightarrow \mathbf{a}}$ and $\gamma_{\mathbf{a} \rightarrow \mathbf{b}} = 0$ if and only if $\mathbf{a} = \mathbf{b}$ for each pair $\mathbf{a}, \mathbf{b} \in \Sigma_-$. Elias [Eli06] shows that, even considering such restrictions, **MSA** is NP-complete. We start showing a polynomial time reduction from **MSA** to **NMSA- z** .

For an instance (S, C) of **MSA** (**NMSA- z**), where $S = s_1, \dots, s_k$ is a k -sequence and C is an integer, **MSA**(S, C) (**NMSA- z** (S, C)) is the decision problem version asking whether there exists an alignment A of S such that $v\text{SP}_\gamma[A] \leq C$ ($V_\gamma^i[A] \leq C$).

Consider a fixed alphabet Σ and scoring matrix γ with the restrictions above that are $\gamma_{\mathbf{a} \rightarrow \mathbf{b}} = \gamma_{\mathbf{b} \rightarrow \mathbf{a}}$ and $\gamma_{\mathbf{a} \rightarrow \mathbf{b}} = 0$. We also assume each entry of γ is integer. Let $\sigma \notin \Sigma_-$ be a new symbol and $\Sigma^\sigma = \Sigma \cup \{\sigma\}$ and G the maximum number in γ . We define a scoring matrix γ^σ such that $\gamma_{\mathbf{a} \rightarrow \mathbf{b}}^\sigma = \gamma_{\mathbf{a} \rightarrow \mathbf{b}}, \gamma_{\mathbf{a} \rightarrow \sigma}^\sigma = \gamma_{\sigma \rightarrow \mathbf{a}}^\sigma = G$ and $\gamma_{\sigma \rightarrow \sigma}^\sigma = 0$ for each pair $\mathbf{a}, \mathbf{b} \in \Sigma_-$. Notice that $\gamma_{\mathbf{a} \rightarrow \mathbf{b}}^\sigma = \gamma_{\mathbf{b} \rightarrow \mathbf{a}}^\sigma$ and $\gamma_{\mathbf{a} \rightarrow \mathbf{b}}^\sigma = 0$ if and only if $\mathbf{a} = \mathbf{b}$ for each pair $\mathbf{a}, \mathbf{b} \in \Sigma^\sigma$ and therefore $G \geq 1$. Also, we denote $S^L = s_1\sigma^L, \dots, s_k\sigma^L$, where $L = Nk^2MG$, $M = \max_i\{|s_i|\}$ and $N = \binom{k}{2}M$.

Let A be an alignment of S^L . A σ -column in A is a column where every symbol is equal to σ . The *tail* of A is the alignment $A[j+1 : |A|]$ if each its column is σ -columns but $A(j)$ is not; in this case the *tail length* of A is $|A| - j$ and the column j is the *tail base*. We say that an alignment of S^L is *canonical* if its tail length is L . If $A = [s''_1, \dots, s''_k]$ is the alignment of S , then we denote by A^L the canonical alignment $[s''_1 \sigma^L, \dots, s''_k \sigma^L]$ of S^L , i.e., $A^L[1 : |A|] = A$ and $A^L[|A| + 1 : |A^L|] = [\sigma^L, \sigma^L, \dots, \sigma^L]$. Notice that $|A^L| \geq L$. Then, we establish below a lower bound for $vSP_{\gamma\sigma}[A^L]$ and $vSP_{\gamma}[A]$.

Proposition 3.1. *Let A be an alignment of a k -sequence $S = s_1, s_2, \dots, s_k$. Then, $vSP_{\gamma\sigma}[A^L] = vSP_{\gamma}[A] \leq k^2 MG$.*

Proof. Suppose that $A = [s'_1, \dots, s'_k]$ and then $A^L = [s'_1 \sigma^L, \dots, s'_k \sigma^L]$. Because $M = \max_i |s_i|$, each alignment that is induced by two sequences in A has at most $2M$ columns. Moreover, each entry in γ is at most G . It follows that $vA_{\gamma}[s'_h, s'_i] = vA_{\gamma}[s'_h \sigma^L, s'_i \sigma^L] \leq 2MG$ for each pair h, i and then

$$vSP_{\gamma\sigma}[A^L] = vSP_{\gamma}[A] = \sum_{h=1}^{k-1} \sum_{i=h+1}^k vA_{\gamma}[A_{\{h,i\}}] \leq \sum_{h=1}^{k-1} \sum_{i=h+1}^k 2MG = \binom{k}{2} 2MG \leq k^2 MG.$$

□

The two following result are useful to prove Theorem 3.4, which is the main result of this section. Let $C^1 := C^2 := C/L$, $C^3 := C/\binom{k}{2}L$ and $L := Nk^2 MG$.

Lemma 3.2. *If $C \geq k^2 MG$, then $\mathbf{MSA}(S, C) = \mathbf{NMSA}\text{-}z(S^L, C^z) = \mathbf{Yes}$, for each $z = 1, 2, 3$.*

Proof. Suppose that $C \geq k^2 MG$. Let A be an alignment of S . From Proposition 3.1, $vSP_{\gamma}[A] = vSP_{\gamma\sigma}[A^L] \leq k^2 MG \leq C$ which implies that $\mathbf{MSA}(S, C) = \mathbf{Yes}$. Since $vSP_{\gamma\sigma}[A^L] \leq C$, we have that $V_{\gamma\sigma}^1[A^L] = vSP_{\gamma\sigma}[A^L]/L \leq C/L = C^1$, and then $\mathbf{NMSA}\text{-}1(S^L, C^1) = \mathbf{Yes}$. Since $vSP_{\gamma\sigma}[A^L] \leq C$ and $|A_{\{h,i\}}^L| \geq L$, we have

$$V_{\gamma\sigma}^2[A^L] = \sum_{h=1}^{k-1} \sum_{i=k+1}^k \frac{vA_{\gamma\sigma}[A_{\{h,i\}}^L]}{|A_{\{h,i\}}^L|} \leq \frac{\sum_{h=1}^{k-1} \sum_{i=h+1}^k vA_{\gamma\sigma}[A_{\{h,i\}}^L]}{L} = \frac{vSP_{\gamma\sigma}[A^L]}{L} \leq \frac{C}{L} = C^2,$$

and thus $\mathbf{NMSA}\text{-}2(S^L, C^2) = \mathbf{Yes}$. Again, since $vSP_{\gamma\sigma}[A^L] \leq C$ and $|A_{\{h,i\}}^L| \geq L$, we have

$$V_{\gamma\sigma}^3[A^L] = \frac{vSP_{\gamma\sigma}[A^L]}{\sum_{h=1}^{k-1} \sum_{i=k+1}^k |A_{\{h,i\}}^L|} \leq \frac{vSP_{\gamma\sigma}[A^L]}{\sum_{h=1}^{k-1} \sum_{i=k+1}^k L} = \frac{vSP_{\gamma\sigma}[A^L]}{\binom{k}{2}L} \leq \frac{C}{\binom{k}{2}L} = C^3,$$

and then $\mathbf{NMSA}\text{-}3(S^L, C^3) = \mathbf{Yes}$.

Therefore, if $C \geq k^2 MG$, then $\mathbf{MSA}(S, C) = \mathbf{NMSA}\text{-}z(S^L, C^z) = \mathbf{Yes}$ for each z . □

Lemma 3.3. *There exists a canonical alignment of S^L which is $V_{\gamma\sigma}^z$ -optimal for each z .*

Proof. Suppose by contradiction that any canonical alignment of S^L is not $V_{\gamma\sigma}^z$ -optimal. Let $A = [s'_1, \dots, s'_k]$ be a $V_{\gamma\sigma}^z$ -optimal alignment of S^L with maximum tail length and maximum number of σ in the tail base. Note that, by hypothesis, A is not canonical.

Let q be the index of the tail base of A . Since A is not canonical, the column q contains only symbols $-$ and σ . Let p be the greatest index such that $p < q$ and there exists an integer i where $s'_i(p) = \sigma$ and $s'_i(q) = -$. Let $A' = [s''_1, \dots, s''_k]$

be an alignment of S^L such that A' is almost the same as A , except for columns p and q , that are defined as following. For each h , we have

$$s''_h = \begin{cases} s'_h, & \text{if } s'_h(p) \neq \sigma \text{ or } s'_h(q) \neq -, \\ s'_h(1:p-1) \cdot \dots \cdot s'_h(p+1:q-1) \cdot \sigma \cdot s'_h(q+1:|s'_h|), & \text{otherwise.} \end{cases}$$

Observe that either the tail length of A' is greater than the tail length of A or the tail lengths of A and A' are the same but the number of σ in the tail base of A' is greater than this number in A . Thus, by the choice of A , the alignment A' is not $V_{\gamma\sigma}^z$ -optimal.

Let h, i be integers. We classify the induced alignment $A_{\{h,i\}}$ of A as follows:

- *Type 1:* if $vA_{\gamma\sigma}[A_{\{h,i\}}] = vA_{\gamma\sigma}[A'_{\{h,i\}}]$ and $|A_{\{h,i\}}| = |A'_{\{h,i\}}|$;
- *Type 2:* if $vA_{\gamma\sigma}[A_{\{h,i\}}] \neq vA_{\gamma\sigma}[A'_{\{h,i\}}]$ and $|A_{\{h,i\}}| = |A'_{\{h,i\}}|$. In this case, the only possibility is that in one of the sequences, say s'_h , is such that $s'_h(p) = \sigma$ and $s'_h(q) = -$, and the other, s'_i , is such that $s'_i(p) = x$ and $s'_i(q) = \sigma$, where $x \in \Sigma$. By hypothesis, $G \geq \gamma_{x \rightarrow -} = \gamma_{- \rightarrow x}$. Then,

$$vA_{\gamma\sigma}[A'_{\{h,i\}}] = vA_{\gamma\sigma}[A_{\{h,i\}}] - G + \gamma_{x \rightarrow -} (= \gamma_{- \rightarrow x}) \leq vA_{\gamma\sigma}[A_{\{h,i\}}].$$

Therefore, $vA_{\gamma\sigma}[A'_{\{h,i\}}] \leq vA_{\gamma\sigma}[A_{\{h,i\}}]$;

- *Type 3:* if $|A_{\{h,i\}}| \neq |A'_{\{h,i\}}|$. In this case, the only possibility is that in one of the sequences, say s'_h , is such that $s'_h(p) = \sigma$ and $s'_h(q) = -$, and the other, s'_i , is such that $s'_i(p) = -$ and $s'_i(q) = \sigma$. It follows that

$$vA_{\gamma\sigma}[A'_{\{h,i\}}] = vA_{\gamma\sigma}[A_{\{h,i\}}] - 2G \quad \text{and} \quad |A_{\{h,i\}}| = |A'_{\{h,i\}}| + 1.$$

We consider now the case $z = 1$. Suppose that $|A'| = |A|$. Analyzing the types above, $vA_{\gamma\sigma}[A'_{\{h,i\}}] \leq vA_{\gamma\sigma}[A_{\{h,i\}}]$, which implies that $vSP_{\gamma\sigma}[A'] \leq vSP_{\gamma\sigma}[A]$ and $V_{\gamma\sigma}^1(A') = vSP_{\gamma\sigma}[A']/|A'| \leq vSP_{\gamma\sigma}[A]/|A| = V_{\gamma\sigma}^1[A]$, which contradicts A' is not $V_{\gamma\sigma}^1$ -optimal. Then, we assume $|A'| \neq |A|$ which implies that $|A'| = |A| - 1$ and at least one alignment $A_{\{h,i\}}$ is of type 3, meaning that $vA_{\gamma\sigma}[A'_{\{h,i\}}] = vA_{\gamma\sigma}[A_{\{h,i\}}] - 2G$. It follows that $vSP_{\gamma\sigma}[A'] \leq vSP_{\gamma\sigma}[A] - 2G$. Then,

$$V_{\gamma\sigma}^1[A'] = \frac{vSP_{\gamma\sigma}[A']}{|A'|} \leq \frac{vSP_{\gamma\sigma}[A] - 2G}{|A| - 1}. \quad (6)$$

Let B be a canonical alignment. By Proposition 3.1, we have that $vSP_{\gamma\sigma}[B] \leq k^2MG$. By the choice of A , we have $V_{\gamma\sigma}^1[A] \leq V_{\gamma\sigma}^1[B]$. Since $G \geq 1$ and $|B| \geq L = Nk^2MG$, then

$$V_{\gamma\sigma}^1[A] \leq V_{\gamma\sigma}^1[B] = \frac{vSP_{\gamma\sigma}[B]}{|B|} \leq \frac{k^2MG}{Nk^2MG} = \frac{1}{N} \leq G.$$

Since $vSP_{\gamma\sigma}[A]/|A| \leq G$, we have $(vSP_{\gamma\sigma}[A] - G)/(|A| - 1) \leq vSP_{\gamma\sigma}[A]/|A|$ which implies, by equation (6), $G \geq 1$ and by the definition of $V_{\gamma\sigma}^1[A]$, that

$$V_{\gamma\sigma}^1[A'] \leq \frac{vSP_{\gamma\sigma}[A] - 2G}{|A| - 1} \leq \frac{vSP_{\gamma\sigma}[A] - G}{|A| - 1} \leq \frac{vSP_{\gamma\sigma}[A]}{|A|} = V_{\gamma\sigma}^1[A],$$

which contradicts again that A' is not $V_{\gamma\sigma}^1$ -optimal. Thus, there exists a canonical alignment of S^L which is $V_{\gamma\sigma}^1$ -optimal.

Now, we consider the case $z = 2$. If an induced alignment $A_{\{h,i\}}$ is of type 1 or 2, then $vA_{\gamma\sigma}[A'_{\{h,i\}}] \leq vA_{\gamma\sigma}[A_{\{h,i\}}]$ and $|A'_{\{h,i\}}| = |A_{\{h,i\}}|$, which implies that

$$vN_{\gamma\sigma}[A'_{\{h,i\}}] = \frac{vA_{\gamma\sigma}[A'_{\{h,i\}}]}{|A'_{\{h,i\}}|} \leq \frac{vA_{\gamma\sigma}[A_{\{h,i\}}]}{|A_{\{h,i\}}|} = vN_{\gamma\sigma}[A_{\{h,i\}}]. \quad (7)$$

If $A_{\{h,i\}}$ is of type 3, then $vA_{\gamma^\sigma}[A'_{\{h,i\}}] = vA_{\gamma^\sigma}[A_{\{h,i\}}] - 2G$ and $|A'_{\{h,i\}}| = |A_{\{h,i\}}| - 1$ which implies that

$$vN_{\gamma^\sigma}[A'_{\{h,i\}}] = \frac{vA_{\gamma^\sigma}[A'_{\{h,i\}}]}{|A'_{\{h,i\}}|} = \frac{vA_{\gamma^\sigma}[A_{\{h,i\}}] - 2G}{|A_{\{h,i\}}| - 1} \leq \frac{vA_{\gamma^\sigma}[A_{\{h,i\}}] - G}{|A_{\{h,i\}}| - 1} \leq \frac{vA_{\gamma^\sigma}[A_{\{h,i\}}]}{|A_{\{h,i\}}|} \leq vN_{\gamma^\sigma}[A_{\{h,i\}}], \quad (8)$$

where the first inequality is a consequence of $G \geq 1$ and the second, since G is the maximum value in γ^σ and therefore G is an upper bound to $vN_{\gamma^\sigma}[A_{\{h,i\}}]$, is a consequence of $vA_{\gamma^\sigma}[A_{\{h,i\}}]/|A_{\{h,i\}}| \leq G$. As a consequence of equations (7) and (8) we have that $V_{\gamma^\sigma}^2[A'] \leq V_{\gamma^\sigma}^2[A]$ contradicting the assumption that A' is not $V_{\gamma^\sigma}^2$ -optimal. Thus, there exists a canonical alignment of S^L which is $V_{\gamma^\sigma}^2$ -optimal.

Finally, we show the case when $z = 3$. We denote T_j the set of all pairs (h, i) such that $A_{\{h,i\}}$ is of type j . Recall that each induced alignment $A_{\{h,i\}}$ of types 1 and 2 are such that $vA_{\gamma^\sigma}[A'_{\{h,i\}}] \leq vA_{\gamma^\sigma}[A_{\{h,i\}}]$ and $|A'_{\{h,i\}}| = |A_{\{h,i\}}|$. Thus, the total contribution of the induced alignments of types 1 and 2 to the $V_{\gamma^\sigma}^3$ -score is

$$\sum_{(h,i) \in T_1 \cup T_2} vA_{\gamma^\sigma}[A'_{\{h,i\}}] \leq \sum_{(h,i) \in T_1 \cup T_2} vA_{\gamma^\sigma}[A_{\{h,i\}}] \quad \text{and} \quad \sum_{(h,i) \in T_1 \cup T_2} |A'_{\{h,i\}}| = \sum_{(h,i) \in T_1 \cup T_2} |A_{\{h,i\}}|.$$

And since each alignment $A_{\{h,i\}}$ of type 3 is such that $vA_{\gamma^\sigma}[A'_{\{h,i\}}] = vA_{\gamma^\sigma}[A_{\{h,i\}}] - 2G$ and $|A'_{\{h,i\}}| = |A_{\{h,i\}}| - 1$, we have

$$\sum_{(h,i) \in T_3} vA_{\gamma^\sigma}[A'_{\{h,i\}}] = \sum_{(h,i) \in T_3} (vA_{\gamma^\sigma}[A_{\{h,i\}}] - 2G) \quad \text{and} \quad \sum_{(h,i) \in T_3} |A'_{\{h,i\}}| = \sum_{(h,i) \in T_3} (|A_{\{h,i\}}| - 1).$$

It follows that

$$\begin{aligned} V_{\gamma^\sigma}^3[A'] &= \frac{vSP_{\gamma^\sigma}[A']}{|A'|} = \frac{\sum_{(h,i) \in T_1 \cup T_2} vA_{\gamma^\sigma}[A'_{\{h,i\}}] + \sum_{(h,i) \in T_3} vA_{\gamma^\sigma}[A'_{\{h,i\}}]}{\sum_{(h,i) \in T_1 \cup T_2} |A'_{\{h,i\}}| + \sum_{(h,i) \in T_3} |A'_{\{h,i\}}|} \leq \frac{\sum_{(h,i) \in T_1 \cup T_2} vA_{\gamma^\sigma}[A_{\{h,i\}}] + \sum_{(h,i) \in T_3} (vA_{\gamma^\sigma}[A_{\{h,i\}}] - 2G)}{\sum_{(h,i) \in T_1 \cup T_2} |A_{\{h,i\}}| + \sum_{(h,i) \in T_3} (|A_{\{h,i\}}| - 1)} \\ &= \frac{vSP_{\gamma^\sigma}[A] - 2|T_3|G}{\left(\sum_{h=1}^{k-1} \sum_{i=h+1}^k |A_{\{h,i\}}|\right) - |T_3|} \leq \frac{vSP_{\gamma^\sigma}[A] - |T_3|G}{\left(\sum_{h=1}^{k-1} \sum_{i=h+1}^k |A_{\{h,i\}}|\right) - |T_3|} \leq \frac{vSP_{\gamma^\sigma}[A]}{\sum_{h=1}^{k-1} \sum_{i=h+1}^k |A_{\{h,i\}}|} = V_{\gamma^\sigma}^3[A], \end{aligned}$$

where the second inequality is a consequence of $G \geq 1$, and the last inequality is a consequence of $vSP_{\gamma^\sigma}[A]/|A| \leq G$ since G is the maximum value in γ^σ and then G is an upper bound to $V_{\gamma^\sigma}^3[A]$. Thus, $V_{\gamma^\sigma}^3[A'] \leq V_{\gamma^\sigma}^3[A]$, which contradicts the assumption that A' is not $V_{\gamma^\sigma}^3$ -optimal. Therefore, there exists a canonical alignment of S^L which is $V_{\gamma^\sigma}^3$ -optimal. \square

Theorem 3.4. *NMSA- z is NP-complete for each z .*

Proof. Given a k -sequence S , an alignment A of S and a integer C , it is easy to check in polynomial time on the length of A that $V_{\gamma^\sigma}^z[A] \leq C$ for $z \in \{1, 2, 3\}$. Then, **NMSA- z** is in NP.

Now, we prove that **MSA**(S, C) = Yes if and only if **NMSA- z** (S^L, C^z) = Yes for each $z \in \{1, 2, 3\}$. If $C \geq k^2MG$, from Lemma 3.2 the theorem is proved. Thus, we assume $C < k^2MG$.

Suppose that $\mathbf{MSA}(S, C) = \text{Yes}$ and, hence, there exists an alignment A such that $v\text{SP}_\gamma[A] \leq C$:

$$\begin{aligned} V_{\gamma^\sigma}^1[A^L] &= \frac{v\text{SP}_{\gamma^\sigma}[A^L]}{|A^L|} \leq \frac{v\text{SP}_{\gamma^\sigma}[A^L]}{L} = \frac{v\text{SP}_\gamma[A]}{L} \leq \frac{C}{L} = C^1, \\ V_{\gamma^\sigma}^2[A^L] &= \sum_{h=1}^{k-1} \sum_{i=k+1}^k \frac{vA_{\gamma^\sigma}[A_{\{h,i\}}^L]}{|A_{\{h,i\}}^L|} \leq \sum_{h=1}^{k-1} \sum_{i=h+1}^k \frac{vA_{\gamma^\sigma}[A_{\{h,i\}}^L]}{L} = \frac{v\text{SP}_{\gamma^\sigma}[A^L]}{L} = \frac{v\text{SP}_\gamma[A]}{L} \leq \frac{C}{L} = C^2, \\ V_{\gamma^\sigma}^3[A^L] &= \frac{v\text{SP}_{\gamma^\sigma}[A^L]}{\sum_{h=1}^{k-1} \sum_{i=k+1}^k |A_{\{h,i\}}^L|} \leq \frac{v\text{SP}_{\gamma^\sigma}[A^L]}{\sum_{h=1}^{k-1} \sum_{i=k+1}^k L} = \frac{v\text{SP}_{\gamma^\sigma}[A^L]}{\binom{k}{2}L} = \frac{v\text{SP}_\gamma[A]}{\binom{k}{2}L} \leq \frac{C}{\binom{k}{2}L} = C^3, \end{aligned}$$

where the first inequality in each equation follows from either A^L or each alignment induced by A^L has length at least L and the second inequality follows from $v\text{SP}_\gamma[A] \leq C$. Thus, if $\mathbf{MSA}(S, C) = \text{Yes}$ then $\mathbf{NMSA}\text{-}z(S^L, C^z) = \text{Yes}$.

Conversely, suppose that $\mathbf{NMSA}\text{-}z(S^L, C^z) = \text{Yes}$. It follows from Lemma 3.3 that there exists a canonical alignment A^L of S^L such that $V_{\gamma^\sigma}^z[A^L] \leq C^z$ for each z . Thus, considering $V_{\gamma^\sigma}^1[A^L] \leq C^1$, we have

$$\begin{aligned} v\text{SP}_\gamma[A] &= v\text{SP}_{\gamma^\sigma}[A^L] = (N+L) \frac{v\text{SP}_{\gamma^\sigma}[A^L]}{N+L} \leq (N+L) \frac{v\text{SP}_{\gamma^\sigma}[A^L]}{|A^L|} = (N+L) V_{\gamma^\sigma}^1[A^L] \\ &\leq (N+L) C^1 = (N+L) \frac{C}{L} = \frac{NC}{L} + C < \frac{Nk^2MG}{L} + C = 1 + C, \end{aligned}$$

where the first equality holds since A^L is canonical, the first inequality holds since $|A^L| \leq N+L$ and the second and the third inequalities hold by hypothesis. Considering $V_{\gamma^\sigma}^2[A^L] \leq C^2$, we have

$$\begin{aligned} v\text{SP}_\gamma[A] &= v\text{SP}_{\gamma^\sigma}[A^L] = (N+L) \frac{v\text{SP}_{\gamma^\sigma}[A^L]}{N+L} = (N+L) \sum_{h=1}^{k-1} \sum_{i=h+1}^k \frac{vA_{\gamma^\sigma}[A_{\{h,i\}}^L]}{N+L} \\ &\leq (N+L) \sum_{h=1}^{k-1} \sum_{i=h+1}^k \frac{vA_{\gamma^\sigma}[A_{\{h,i\}}^L]}{|A_{\{h,i\}}^L|} = (N+L) V_{\gamma^\sigma}^2[A^L] \\ &\leq (N+L) C^2 = (N+L) \frac{C}{L} = \frac{NC}{L} + C < \frac{Nk^2MG}{L} + C = 1 + C, \end{aligned}$$

where the first equality holds since A^L is canonical, the first inequality holds since, for each h, i , $|A_{\{h,i\}}^L| \leq N+L$, and the second and the third inequalities hold by hypothesis. And finally, considering $V_{\gamma^\sigma}^3[A^L] \leq C^3$, we have

$$\begin{aligned} v\text{SP}_\gamma[A] &= v\text{SP}_{\gamma^\sigma}[A^L] = (N + \binom{k}{2}L) \frac{v\text{SP}_{\gamma^\sigma}[A^L]}{N + \binom{k}{2}L} \\ &\leq (N + \binom{k}{2}L) \frac{v\text{SP}_{\gamma^\sigma}[A^L]}{\sum_{h=1}^{k-1} \sum_{i=h+1}^k |A_{\{h,i\}}^L|} = (N + \binom{k}{2}L) V_{\gamma^\sigma}^3[A^L] \leq (N + \binom{k}{2}L) C^3 \\ &= (N + \binom{k}{2}L) \frac{C}{\binom{k}{2}L} = \frac{NC}{\binom{k}{2}L} + C < \frac{Nk^2MG}{\binom{k}{2}L} + C = \frac{1}{\binom{k}{2}} + C \leq 1 + C, \end{aligned}$$

where the first equality holds since A^L is canonical, the first inequality holds since the sum of lengths of two sequences induced by a canonical alignment is at most $N + \binom{k}{2}L$ and the second and the third inequalities hold by hypothesis. Therefore, if $\mathbf{NMSA}\text{-}z(S^L, C^z) = \text{Yes}$ then $v\text{SP}_\gamma[A] < 1 + C$ for any $z \in \{1, 2, 3\}$. Since the entries in the scoring matrix are integers, we have that $v\text{SP}_\gamma[A]$ is an integer. And since C is an integer, it follows that $v\text{SP}_\gamma[A] \leq C$.

□

4 Exact algorithms

In the following sections we describe exact dynamic programming algorithms for **NMSA- z** , with $z = 1, 2, 3$.

4.1 NMSA-1

Let $S = s_1, \dots, s_k$ be a k -sequence and $A = [s'_1, \dots, s'_k]$ be an alignment of S . As defined in Equation (3), $V_\gamma^1[A]$ takes into account the length of A , and the optimal function is given by $\text{opt}V_\gamma^1(S) = \min_{A \in \mathcal{A}_S} \{V_\gamma^1[A]\}$. In the optimization version of **NMSA-1**, we are given a k -sequence S and we want to compute $\text{opt}V_\gamma^1(S)$ for a fixed matrix γ . We can solve **NMSA-1** by calculating the minimum SP-score considering every possible length of an alignment. That is, we compute the entries of a table D indexed by $V_{\vec{n}} \times \{0, 1, \dots, N\}$, where $\vec{n} = [s_1, \dots, s_k]$ and $N = \sum_{i=1}^k |s_i|$. The entry $D(\vec{v}, L)$ stores the score of an alignment of $S(\vec{v})$ of length L with minimum SP-score. Notice that $D(\vec{0}, 0) = 0$, $D(\vec{v} \neq \vec{0}, 0) = D(\vec{0}, L \neq 0) = \infty$. Therefore, the table entries can be calculated as:

$$D(\vec{v}, L) = \begin{cases} 0, & \text{if } \vec{v} = \vec{0}, L = 0, \\ \infty, & \text{if } \vec{v} = \vec{0}, L \neq 0 \text{ or } \vec{v} \neq \vec{0}, L = 0, \\ \min_{\vec{b} \in \mathcal{B}_k, \vec{b} \leq \vec{v}} \{D(\vec{v} - \vec{b}, L - 1) + v\text{SP}_\gamma[\vec{b} \cdot S(\vec{v})]\}, & \text{otherwise.} \end{cases}$$

Table D is computed for all possible values of $L = 0, \dots, N$. Consequently, $\text{opt}V_\gamma^1(S) = \min_L \{D(\vec{n}, L)/L\}$ is returned. Algorithm 2 describes this procedure more precisely.

Algorithm 2

Input: k -sequence $S = s_1, \dots, s_k$ such that $n_i = |s_i|$

Output: $\text{opt}V_\gamma^1(S)$

- 1: $D(\vec{0}, 0) \leftarrow 0$
 - 2: **for** each $L \neq 0$ **do** $D(\vec{0}, L) \leftarrow \infty$
 - 3: **for** each $\vec{v} \neq \vec{0}$ **do** $D(\vec{v}, 0) \leftarrow \infty$
 - 4: **for** each $\vec{0} < \vec{v} \leq \vec{n}$ in lexicographical order **do**
 - 5: **for** each $L \leftarrow 1, 2, \dots, N$ **do**
 - 6: $D(\vec{v}, L) \leftarrow \min_{\vec{b} \in \mathcal{B}_k, \vec{b} \leq \vec{v}} \{D(\vec{v} - \vec{b}, L - 1) + v\text{SP}_\gamma[\vec{b} \cdot S(\vec{v})]\}$
 - 7: **return** $\min_L \{D(\vec{n}, L)/L\}$
-

Suppose that $n_i = |s_i| = n$ for each i which implies that $|V_n| = (n+1)^k$ and $N = nk$. Notice that the space to store the matrix D is $\Theta(N \cdot |V_n|) = \Theta(n \cdot (n+1)^k)$. The time consumption of Algorithm 2 corresponds to the time needed to fill the table D up, plus the running time of line 7. Each entry of D can be computed in $O((2^k - 1) \cdot \binom{k}{2}) = O(2^k k^2)$ -time. Therefore, the algorithm spends $O(2^k k^2 \cdot kn(n+1)^k) = O(2^k k^3 (n+1)^{k+1})$ -time to compute the entire table D . Line 7 is computed in $\Theta(N = kn)$ -time. Therefore, the running time of Algorithm 2 is $O(2^k k^3 (n+1)^{k+1}) + \Theta(N) = O(2^k k^3 (n+1)^{k+1})$.

4.2 NMSA-2

Let $S = s_1, \dots, s_k$ be a k -sequence and $A = [s'_1, \dots, s'_k]$ be an alignment of S . As defined in Equation (4), $V_\gamma^2[A]$ takes into account the lengths of the induced alignments in A . In the optimization version of **NMSA-2**, we are given a k -sequence S and we want to compute $\text{opt}V_\gamma^2(S)$ for a fixed scoring matrix γ .

Let $\vec{L} = [L_{12}, \dots, L_{hi}, \dots, L_{(k-1)k}]$ be a $\binom{k}{2}$ -vector indexed by the set of pairs of integers $\{h, i\}$ such that $1 \leq h < i \leq k$ and L_{hi} denotes the element of \vec{L} of index $\{h, i\}$. The lengths of the induced alignments can be represented by a vector \vec{L} .

Thus, if A is an alignment and $|A_{\{h,i\}}| = L_{hi}$ for each pair h, i , we say that \vec{L} is the *induced length* of A . For a k -sequence $S = s_1, \dots, s_k$, where $n_i = |s_i|$ for each i , we define

$$\mathbb{L} = \{\vec{L} = [L_{12}, L_{13}, \dots, L_{1k}, L_{23}, \dots, L_{2k}, \dots, L_{(k-1)k}] : 0 \leq L_{hi} \leq n_h + n_i\}.$$

Therefore, \mathbb{L} contains the induced length of alignment A of S for all $A \in \mathcal{A}_S$. Note that if n is the length of each sequence in S , then $|\mathbb{L}| = (2n+1)^{\binom{k}{2}}$. Let $\vec{b} = [b_1, \dots, b_k]$ be a k -vector of bits. Overloading the minus operator “-”, we define $\vec{L} - \vec{b}$ to be a $\binom{k}{2}$ -vector \vec{L}' such that

$$L'_{hi} = \begin{cases} L_{hi}, & \text{if } b_h = b_i = 0, \\ L_{hi} - 1, & \text{otherwise.} \end{cases}$$

Observe that if \vec{L} is the induced length of an alignment A of $S(\vec{v})$ and \vec{b} is a k -vector of bits such that $\vec{b} \cdot S(\vec{v})$ is the last column of A , then $\vec{L}' = \vec{L} - \vec{b}$ is the induced length of the alignment $A(1:|A| - 1)$.

From a $\binom{k}{2}$ -vector \vec{L} and a scoring matrix γ , we can define an array of $\binom{k}{2}$ scoring matrices $\vec{\gamma} = \gamma \times \vec{L} = [\dots, \gamma^{(hi)}, \dots]$ indexed by $\{1, \dots, k\} \times \{1, \dots, k\}$ such that

$$\gamma^{(hi)}_{\mathbf{a} \rightarrow \mathbf{b}} = \frac{\gamma_{\mathbf{a} \rightarrow \mathbf{b}}}{L_{hi}},$$

for each $\mathbf{a}, \mathbf{b} \in \Sigma^-$. Observe that

$$V_\gamma^2[A] = \sum_{h=1}^{k-1} \sum_{i=h+1}^k \left(vN_\gamma[A_{\{h,i\}}] (= \frac{vA_\gamma[A_{\{h,i\}}]}{|A_{\{h,i\}}|}) \right) = vA_{\gamma^{(hi)}}[A_{\{h,i\}}] = vSP_{\vec{\gamma}}[A]$$

where $\vec{\gamma} = \gamma \times [|A_{\{1,2\}}|, \dots, |A_{\{k-1,k\}}|]$. Besides, if \vec{L} is the induced length of a V_γ^2 -optimal alignment of S , then we can compute $\text{opt}V_\gamma^2$ through the recurrence

$$D_{\vec{L}}(\vec{v}, \vec{L}) = \begin{cases} 0, & \text{if } \vec{v} = \vec{0} \text{ and } \vec{L} = \vec{0}, \\ \infty, & \text{if } \vec{v} = \vec{0} \text{ and } \vec{L} \neq \vec{0}, \\ \infty, & \text{if } \vec{v} \neq \vec{0} \text{ and } \vec{L} = \vec{0}, \\ \min_{\vec{b} \in \mathcal{B}_k, \vec{b} \leq \vec{v}, \vec{b} \leq \vec{L}} \{ D_{\vec{L}}(\vec{v} - \vec{b}, \vec{L} - \vec{b}) + vSP_{\vec{\gamma}}[\vec{b} \cdot S(\vec{v})] \}, & \text{otherwise,} \end{cases}$$

where $\vec{b} \leq \vec{L}$ is also an overloading, meaning that $\vec{L} - \vec{b} \geq \vec{0}$, and $\vec{\gamma} = \gamma \times \vec{L}$. In this case, $\text{opt}V_\gamma^2(S) = D_{\vec{L}}(\vec{v}, \vec{L})$.

Algorithm 3 described below finds $\text{opt}V_\gamma^2(S)$.

Algorithm 3

Input: A k -sequence $S = s_1, \dots, s_k$ such that $n_i = |s_i|$

Output: $\text{opt}V_\gamma^2(S)$

- 1: **for** each $\vec{L} \in \mathbb{L}$ **do**
 - 2: $D_{\vec{L}}(\vec{0}, \vec{0}) \leftarrow 0$
 - 3: **for** each $\vec{L} \neq \vec{0}$ **do** $D_{\vec{L}}(\vec{0}, \vec{L}) \leftarrow \infty$
 - 4: **for** each $\vec{v} \neq \vec{0}$ **do** $D_{\vec{L}}(\vec{v}, \vec{0}) \leftarrow \infty$
 - 5: $\vec{\gamma} \leftarrow \gamma \times \vec{L}$
 - 6: **for** each $\vec{0} < \vec{v} \leq \vec{n}$ in lexicographical order **do**
 - 7: **for** each $\vec{L} \neq \vec{0}$ in lexicographical order **do**
 - 8: $D_{\vec{L}}(\vec{v}, \vec{L}) = \min_{\vec{b} \in \mathcal{B}_k, \vec{b} \leq \vec{v}, \vec{b} \leq \vec{L}} \{ D_{\vec{L}}(\vec{v} - \vec{b}, \vec{L} - \vec{b}) + vSP_{\vec{\gamma}}[\vec{b} \cdot S(\vec{v})] \}$
 - 9: **return** $\min_{\vec{L} \in \mathbb{L}} \{ D_{\vec{L}}(\vec{n}, \vec{L}) \}$
-

For k sequences of length n , Algorithm 3 needs $(2n + 1)^{\binom{k}{2}} \cdot (n + 1)^k$ space to store the table $D_{\vec{\mathcal{L}}}$. For each of the $(2n + 1)^{\binom{k}{2}}$ values $\vec{\mathcal{L}} \in \vec{\mathbb{L}}$, table $D_{\vec{\mathcal{L}}}$ is recalculated. Since the computation of each entry takes $O(2^k k^2)$ -time, the total time is

$$O\left(2^k k^2 \cdot (2n + 1)^{\binom{k}{2}} \cdot (2n + 1)^{\binom{k}{2}} (n + 1)^k = \left(1 + \frac{1}{2n + 1}\right)^k (2n + 1)^{k^2} k^2\right).$$

Therefore, if $k \leq 2n + 1$, the total time is $O((2n + 1)^{k^2} k^2)$, since $(1 + 1/(2n + 1))^k \leq (1 + 1/k)^k \leq e \leq 2.72$ is constant.

Existence of an alignment A for a given $\vec{\mathcal{L}}$

Consider a 3-sequence $S = s_1, s_2, s_3$ with length $\vec{n} = [4, 3, 5]$ and suppose we are interested in an alignment A of S with induced length equal to $\vec{\mathcal{L}} = [4, 7, 5]$, i.e., $|A_{\{1,2\}}| = 4, |A_{\{1,3\}}| = 7, |A_{\{2,3\}}| = 5$. Computing $D_{\vec{\mathcal{L}}}(\vec{v}, \vec{\mathcal{L}})$ as in previous section reveals that $D_{\vec{\mathcal{L}}}(\vec{v}, \vec{\mathcal{L}}) = \infty$ which means that such alignment doesn't exist. In fact, we can check that each symbol in sequence 2 is aligned with one symbol in the first and one symbol in the third sequence which implies that there are at least 3 symbols of the first sequence aligned with symbols in the third sequence. Thus, $|A_{\{1,3\}}| \leq 6$. Because, computing $D_{\vec{\mathcal{L}}}(\vec{v}, \vec{\mathcal{L}})$ spends exponential time, it would be great if we could decide whether there exists an associated alignment with the $\vec{\mathcal{L}}$. before computing $D_{\vec{\mathcal{L}}}(\vec{v}, \vec{\mathcal{L}})$. Thus, an interesting problem arises and we define it below.

Problem 4 (Existence of an alignment that is associated with an induced length). Given the length \vec{n} of a k -sequence and a $\binom{k}{2}$ -vector $\vec{\mathcal{L}}$, decide whether there exists an alignment A of S such that $\vec{\mathcal{L}}$ is the induced length of A .

We denote Problem 4 by **EAIL**. Considering \vec{n} the length of the k sequence $S = s_1, \dots, s_k$, an alternative way to represent \vec{n} is through a matrix of integers M that it is indexed by $\{1, 2, \dots, k\}$ and

$$M(h, i) = \begin{cases} |s_h| & \text{if } h = i, \\ |s_h| + |s_i| - L_{hi} & \text{otherwise} \end{cases}$$

i.e, $M(h, i)$ is the number of symbols in s_h aligned with symbols in s_i . Thus, we decide whether exists a collection of sets c_1, \dots, c_k such that $|c_h \cap c_i| = M[h, i]$, where c_h is a set of indices j of some alignment A of S with $A[h, j] \neq -$. This different way to see the problem is exactly another that is known as *Recognizing Intersection Patterns (RIP)*. Notice that the instance of **RIP** can be obtained in linear time and no extra space from the instance of **EAIL**.

Thus, consider an example where $\vec{n} = [5, 5, 5]$ and $L_{12} = L_{13} = 8$ and $L_{23} = 10$. In this case, **EAIL** returns Yes since the induced alignments by the following alignment with 11 columns

$$\begin{bmatrix} s_1(1) & s_1(2) & s_1(3) & s_1(4) & s_1(5) & - & - & - & - & - & - \\ s_2(1) & s_2(2) & - & - & - & s_2(3) & s_2(4) & s_2(5) & - & - & - \\ - & - & s_3(1) & s_3(2) & - & - & - & - & s_3(3) & s_3(4) & s_3(5) \end{bmatrix}$$

respects required restrictions. On the other hand, in the alternative **RIP** formulation of this instance, we want to find a collection of three sets for the matrix

$$M = \begin{array}{c|ccc} & 1 & 2 & 3 \\ \hline 1 & 5 & 2 & 2 \\ 2 & 2 & 5 & 0 \\ 3 & 2 & 0 & 5 \end{array}$$

that satisfies the aforementioned property. The answer to **RIP** is Yes, as we can see in Figure 4.

In another example, suppose that $\vec{n} = [5, 5, 5]$, $L_{12} = L_{13} = 7$ and $L_{23} = 10$. In this case the answer is No. To check it, suppose by contradiction that there is an alignment A of a 3-tuple s_1, s_2, s_3 for this instance. Since $L_{23} = 10$, we have that s_2 and s_3 has no $(5 + 5 - 10 = 0)$ aligned symbol. Since $L_{12} = 7$, then s_1 must have $5 + 5 - 7 = 3$ symbols aligned




Figure 4: An instance of **RIP** for which the answer is Yes.

with s_2 that cannot be aligned with any symbol of s_3 because s_2 and s_3 has no symbol aligned. On the other hand, since $L_{13} = 7$, we have that s_1 and s_3 must have 3 aligned symbols. Since s_1 has 3 symbols not alignment and 3 symbols aligned with s_3 , it follows that s_1 has at least 6 symbols, which is a contradiction.

Chvátal [Chv80] showed that for a special class of matrices M where $M[i, i] = 3$ for every i , **RIP** is NP-complete. Therefore, we have the following result

Theorem 4.1. ***EAIL** is NP-complete when $|s| = 3$ for each sequence in k -sequence and it is NP-hard if its lengths are arbitrary.*

4.3 NMSA-3

Let $S = s_1, \dots, s_k$ be a k -sequence and $A = [s'_1, \dots, s'_k]$ be an alignment of S . As defined in Equation (5), $V_\gamma^3[A]$ takes into account the lengths of induced alignments of A . In the optimization version of **NMSA-3**, given a k -tuple S , the task is to determine $\text{optV}_\gamma^3(S)$ for a fixed matrix γ .

Let $N = \sum_i |s_i|$. Notice that an alignment A that has exactly only one symbol different of space in each column is such that $\sum_{h=1}^{k-1} \sum_{i=h+1}^k |A_{\{h,i\}}| = (k-1)N$.

Here, each entry $D(\vec{v}, L)$ of D stores the SP-score of an alignment A of the prefix $S(\vec{v})$ with the minimum SP-score such that $\sum_{i < h} |A_{\{i,h\}}| = L$. The Boolean vectors \vec{b} are used to represent the contribution to the sum of the lengths of the induced alignments. Thus, we define $\|\vec{b}\| = \binom{k}{2} - \sum_{h < i, b_h = b_i = 0} 1$. Notice that if $\vec{b} \cdot S(\vec{v})$ is the last column of an alignment A and $L = \sum_{h=1}^{k-1} \sum_{i=h+1}^k |A_{\{h,i\}}|$ is the sum of the lengths of the alignments induced by A , then the sum of the lengths of the alignments induced by $A(1:|A| - 1)$ is $L - \|\vec{b}\|$. Therefore,

$$D(\vec{v}, L) = \begin{cases} 0, & \text{if } \vec{v} = \vec{0}, L = 0, \\ \infty, & \text{if } \vec{v} = \vec{0}, L \neq 0 \text{ or } \vec{v} \neq \vec{0}, L = 0, \\ \min_{\vec{b} \in \mathcal{B}_k, \vec{b} \leq \vec{v}, \|\vec{b}\| \leq L} \left\{ D(\vec{v} - \vec{b}, L - \|\vec{b}\|) + v \text{SP}_\gamma[\vec{b} \cdot S(\vec{v})] \right\}, & \text{otherwise.} \end{cases}$$

Algorithm 4 provides more details about the procedure for computing optV_γ^3 .

Algorithm 4

Input: a k -sequence $S = s_1, \dots, s_k$ such that $n_i = |s_i|$

Output: $\text{opt}V_\gamma^3(S)$

- 1: $D(\vec{0}, 0) \leftarrow 0$
 - 2: **for** each $L \neq 0$ **do** $D(\vec{0}, L) \leftarrow \infty$
 - 3: **for** each $\vec{v} \neq \vec{0}$ **do** $D(\vec{v}, 0) \leftarrow \infty$
 - 4: **for** each $\vec{0} < \vec{v} \leq \vec{n}$ in lexicographical order **do**
 - 5: **for** $L \leftarrow 1, 2, \dots, N(k-1)$ **do**
 - 6: $D(\vec{v}, L) \leftarrow \min_{\vec{b} \in \mathcal{B}^k, \vec{b} \leq \vec{v}, \|\vec{b}\| \leq L} \{D(\vec{v} - \vec{b}, L - \|\vec{b}\|) + v\text{SP}_\gamma[\vec{b} \cdot S(\vec{v})]\}$
 - 7: **return** $\min_L \{D(\vec{n}, L)/L\}$
-

Assume that all sequences in S have length n . In this case, table D has $(nk^2 - nk + 1) \cdot (n+1)^k = O(k^2(n+1)^{k+1})$ entries. Since the time required to determine each entry of D is $O(2^k k^2)$, the running time of Algorithm 4 is $O(2^k k^2 \cdot k^2(n+1)^{k+1}) = O(2^k k^4(n+1)^{k+1})$.

5 Approximation algorithms for MSA and NMSA-2

Gusfield [Gus93] described a 2-approximation algorithm for **MSA** assuming that $\gamma \in \mathbb{M}^{\mathbb{C}}$. In this section, we adapt Gusfield's algorithm, thus proposing a 6-approximation algorithm for **MSA** when $\gamma \in \mathbb{M}^{\mathbb{W}}$ and a 12-approximation algorithm for **NMSA-2** problem when $\gamma \in \mathbb{M}^{\mathbb{N}}$.

We consider here a scoring function $v = vA_\gamma$ and $v = vN_\gamma$ when $\gamma \in \mathbb{M}^{\mathbb{W}}$ and $\gamma \in \mathbb{M}^{\mathbb{N}}$, respectively. Also, for a 2-sequence s, t , define $\text{opt}(s, t) = \min_{A \in \mathcal{A}_{s,t}} \{v[A]\}$ and an v -optimal alignment of s, t is an alignment A such $v[A] = \text{opt}(s, t)$. It follows from [AS06] that opt is a metric on Σ^* . For a k -sequence S , define $V[A] = \sum_{h=1}^{k-1} \sum_{i=h+1}^k v[A_{\{h,i\}}]$ and $\text{OPT}(S) = \min_{A \in \mathcal{A}_S} v(A)$ and a V -optimal alignment of S is an alignment A such that $V[A] = \text{OPT}(S)$.

Let c be an integer, $1 \leq c \leq k$. A *star* X with center c (also called c -star) of the k -sequence $S = s_1, \dots, s_k$ is a collection of $k-1$ alignments: $X_h = [s'_h, s_c^h]$ of s_h, s_c for each $h < c$ and $X_h = [s_c^h, s'_h]$ of s_c, s_h for each $h > c$. The set of all stars with center c is denoted by \mathcal{X}_c . The score of the c -star X is $c\text{Star}(X) = \sum_{h \neq c} v[X_h]$ and an v -optimal star is one whose score is $\text{optStar}(S) = \min_{X \in \mathcal{X}_c, c \in \mathbb{N}} \{c\text{Star}(X)\}$. Notice that in a v -optimal c -star, $v(X_h) = \text{opt}(s_h, s_c)$ if $h < c$ and $v(X_h) = \text{opt}(s_c, s_h)$ if $c < h$ and because the symmetry property of opt , we have

$$\text{optStar}(S) = \min_c \left\{ \sum_{h \neq c} \text{opt}(s_h, s_c) \right\},$$

and if $|s| \leq n$ for each $s \in S$, $\text{optStar}(S)$ can be computed in $O(k^2 n^2)$ -time when $v = vA_\gamma$ and $O(k^2 n^3)$ -time when $v = vN_\gamma$.

We say that alignment A of S and c -star X are *compatible* (A is compatible with X and X is compatible with A) in S when either $A_{\{h,c\}}$ (when $h < c$) or $A_{\{c,h\}}$ (when $c < h$) is equal to X_h for each h . Given the alignment A of S , it is easy to obtain the c -star X compatible with A (and there exists only one) considering fixed c . On the other hand, an important known result in alignment studies from Feng and Doolittle [FD87] is that we can find an alignment A that is compatible with a given c -star X in $O(kn)$, where $n \leq |s|$ for each sequence s in S . In general in this case, there exists many compatible alignments with X .

It is easy to adapt the following result from Gusfield [Gus93] to a k -sequence.

Lemma 5.1. *Given a k -sequence S ,*

$$\text{optStar}(S) \leq \frac{2}{k} \cdot \text{OPT}(S).$$

Proof. Let X be a v -optimal star of S and c its center, and A an v -optimal alignment of S . Then,

$$k \cdot \text{optStar}(S) = k \cdot c\text{Star}(X) = \sum_{h=1}^k c\text{Star}(X) = \sum_{h=1}^k \sum_{h \neq c} v[X_h] = \sum_{h=1}^k \sum_{h \neq c} \text{opt}(s_h, s_c) \quad (9)$$

$$\leq 2 \cdot \sum_{h=1}^{k-1} \sum_{i=h+1}^k \text{opt}(s_h, s_i) \quad (10)$$

$$\leq 2 \cdot \sum_{i=1}^k \sum_{h \neq i} v[A_{\{h,i\}}] = 2V[A] = 2 \cdot \text{OPT}(S), \quad (11)$$

where (9) follows from the definition of a star and from the optimality of star X ; (10) follows from triangle inequality of opt ; and (11) follows from the optimality of alignment A .

Therefore, $\text{optStar} \leq (2/k) \cdot \text{opt}(S)$. \square

Let $A = [s', t']$ be an alignment of a 2-sequence s, t . We say that a column j is *splittable* in A if $s'(j) \neq -$, $t'(j) \neq -$ and $\min\{\gamma_{s'(j) \rightarrow -}, \gamma_{s'(j) \rightarrow -}\} \leq \gamma_{s'(j) \rightarrow t'(j)}$. Let $J := \{j_i \in \mathbb{N} : 1 \leq j_1 < \dots < j_m \leq |A| \text{ and } j_i \text{ is splittable in } A\}$. An A -*splitting* is the alignment

$$\left[\begin{array}{cccccccccc} s'(1:j_1-1) & s'(j_1) & - & s'(j_1+1:j_2-1) & s'(j_2) & - & \dots & s'(j_m) & - & s'(j_m+1:|A|) \\ t'(1:j_1-1) & - & t'(j_1) & t'(j_1+1:j_2-1) & - & t'(j_2) & \dots & - & t'(j_m) & t'(j_m+1:|A|) \end{array} \right].$$

We say that J is *required to split* A . The following proposition is used to check properties of an A -splitting.

Proposition 5.2. *Consider $\gamma \in \mathbb{M}^W$ and $\mathbf{a}, \mathbf{b} \in \Sigma$. If $\gamma_{\mathbf{a} \rightarrow -} > \gamma_{\mathbf{a} \rightarrow \mathbf{b}}$ or $\gamma_{\mathbf{a} \rightarrow -} > \gamma_{\mathbf{b} \rightarrow \mathbf{a}}$, then $\gamma_{\mathbf{a} \rightarrow \mathbf{b}} = \gamma_{\mathbf{b} \rightarrow \mathbf{a}}$.*

Proof. Since $\gamma \in \mathbb{M}^W$, we have that $\gamma_{\mathbf{a} \rightarrow -} = \gamma_{- \rightarrow \mathbf{a}} > 0$ and $\gamma_{\mathbf{b} \rightarrow -} = \gamma_{- \rightarrow \mathbf{b}} > 0$. Suppose that $\gamma_{\mathbf{a} \rightarrow -} > \gamma_{\mathbf{a} \rightarrow \mathbf{b}}$. Then, $\gamma_{\mathbf{a} \rightarrow -} + \gamma_{- \rightarrow \mathbf{b}} > \gamma_{\mathbf{a} \rightarrow -} > \gamma_{\mathbf{a} \rightarrow \mathbf{b}}$, and we have that $\gamma_{\mathbf{a} \rightarrow \mathbf{b}} = \gamma_{\mathbf{b} \rightarrow \mathbf{a}}$ since $\gamma \in \mathbb{M}^W$. Assume now that $\gamma_{\mathbf{a} \rightarrow -} > \gamma_{\mathbf{b} \rightarrow \mathbf{a}}$. It follows that $\gamma_{\mathbf{b} \rightarrow -} + \gamma_{- \rightarrow \mathbf{a}} > \gamma_{- \rightarrow \mathbf{a}} = \gamma_{\mathbf{a} \rightarrow -} > \gamma_{\mathbf{b} \rightarrow \mathbf{a}}$, which implies that $\gamma_{\mathbf{a} \rightarrow \mathbf{b}} = \gamma_{\mathbf{b} \rightarrow \mathbf{a}}$ since $\gamma \in \mathbb{M}^W$. \square

Let $X = \{X_1, \dots, X_{c-1}, X_{c+1}, X_k\}$ be a c -star. A X -*starsplitting* is the c -star $Y = \{Y_1, \dots, Y_{c-1}, Y_{c+1}, Y_k\}$ where Y_j is the X_j -splitting for each j . The next result shows that the v -score of the star Y is bounded by the v -score of star X when $\gamma \in \mathbb{M}^W$ and $v = vA_\gamma$ or $\gamma \in \mathbb{M}^N$ and $v = vN_\gamma$.

Lemma 5.3. *Let S be a k -sequence, X be a star of S , Y be the X -starsplitting and v be a function to score alignments. Consider $\gamma \in \mathbb{M}^W$ and $v = vA_\gamma$ or $v = vN_\gamma$. Then, Y is also a c -star and*

$$c\text{Star}(Y) \leq 3 \cdot c\text{Star}(X).$$

Proof. Consider an alignment $X_h = [s', t'] \in X$ and a set J which is required to split X_h . Then,

$$\begin{aligned} vA_\gamma[Y_h] &= vA_\gamma[X_h] + \sum_{j \in J} (\gamma_{s'(j) \rightarrow -} + \gamma_{- \rightarrow t'(j)} - \gamma_{s'(j) \rightarrow t'(j)}) \\ &\leq vA_\gamma[X_h] + \sum_{j \in J} (\gamma_{s'(j) \rightarrow -} + \gamma_{- \rightarrow t'(j)}) \leq vA_\gamma[X_h] + 2 \cdot \sum_{j \in J} \min\{\gamma_{s'(j) \rightarrow -}, \gamma_{- \rightarrow t'(j)}\} \end{aligned} \quad (12)$$

$$\leq vA_\gamma[X_h] + 2 \cdot \sum_{j \in J} \gamma_{s'(j) \rightarrow t'(j)}, \leq vA_\gamma[X_h] + 2 \cdot vA_\gamma[X_h] = 3 \cdot vA_\gamma[X_h], \quad (13)$$

where (12) hold because, since $\gamma \in \mathbb{M}^W$, $\gamma_{s'(j) \rightarrow t'(j)}$ and (13) hold because j is splittable. Thus, $vA_\gamma[Y_h] \leq 3 \cdot vA_\gamma[X_h]$. Furthermore,

$$vN_\gamma[Y_h] = \frac{vA_\gamma[Y_h]}{|Y_h|} \leq \frac{3 \cdot vA_\gamma[X_h]}{|Y_h|} \leq \frac{3 \cdot vA_\gamma[X_h]}{|X_h|} = 3 \cdot vN_\gamma[X_h].$$

Hence, $v(Y_h) \leq 3 \cdot v(X_h)$ when $v = vA_\gamma$ or $v = vN_\gamma$ and $\gamma \in \mathbb{M}^W$ which implies that

$$cStar(Y) = \sum_{h \neq c} v[Y_h] \leq 3 \cdot \sum_{h \neq c} v[X_h] = 3 \cdot cStar(X).$$

□

Notice that the time consumption for computing an X -splitting from X is $O(kn)$ when $|s| \leq n$ for each $s \in S$.

Considering a star X with center c of $S = s_1, \dots, s_k$, there can exist many compatible alignments with a star Y which is a X -splitting. Let COMPATIBLEALIGN be a subroutine that receives the star Y and returns an alignment A compatible with Y . It is quite simple: if symbols $s_h(j_1)$ and $s_c(j_2)$ are aligned in X_h , they are also aligned in A . Otherwise, $s_h(j)$ aligns only with $-$ in A . This property is enough to guarantee the approximation factor of MSA and NMSA-2. As an example, for $S = \text{aaa, bbbbbb, cc, ddd, eeeeeee}$ and

$$X = \left\{ \left[\begin{array}{cccc} \text{a} & \text{a} & \text{a} & - \\ - & \text{d} & \text{d} & \text{d} \end{array} \right], \left[\begin{array}{cccccc} \text{b} & \text{b} & - & \text{b} & \text{b} & \text{b} \\ - & \text{d} & \text{d} & - & - & \text{d} \end{array} \right], \left[\begin{array}{ccc} \text{c} & \text{c} & - \\ \text{d} & \text{d} & \text{d} \end{array} \right], \left[\begin{array}{cccccc} - & - & - & \text{d} & \text{d} & \text{d} & - \\ \text{e} & \text{e} & \text{e} & \text{e} & - & \text{e} & \text{e} \end{array} \right] \right\}$$

a star with center 4, we obtain the alignment

$$\left[\begin{array}{cccccccccccc} \text{a} & - & - & - & - & \text{a} & \text{a} & - & - & - & - & - \\ - & \text{b} & - & - & - & \text{b} & - & \text{b} & \text{b} & \text{b} & - & - \\ - & - & - & - & - & \text{c} & \text{c} & - & - & - & - & - \\ - & - & - & - & - & \text{d} & \text{d} & - & - & \text{d} & - & - \\ - & - & \text{e} & \text{e} & \text{e} & \text{e} & - & - & - & \text{e} & \text{e} & - \end{array} \right].$$

Let $Q_{\max} := \max_{\mathbf{a} \in \Sigma} \{\gamma_{\mathbf{a} \rightarrow -}, \gamma_{- \rightarrow \mathbf{a}}\}$ and consider the following result.

Proposition 5.4. *Let S be a k -sequence, X be a star of S with center c and Y be the X -starsplitting. Assume that $\gamma \in \mathbb{M}^W$ and that COMPATIBLEALIGN(Y) returns $A = [s'_1, \dots, s'_k]$. If $h \neq c$ and $i \neq c$, we have that*

- (i) $\gamma_{s'_h(j) \rightarrow s'_i(j)} \leq \gamma_{s'_h(j) \rightarrow s'_c(j)} + \gamma_{s'_c(j) \rightarrow s'_i(j)}$ for each $j = 1, \dots, |A|$, and
- (ii) $vN_\gamma[A_{\{h,i\}}] \leq 2 \cdot Q_{\max}$.

Proof. Assume that $s'_h(j) = \mathbf{a}$, $s'_i(j) = \mathbf{b}$ and $s'_c(j) = \mathbf{c}$.

First we show that (i) $\gamma_{\mathbf{a} \rightarrow \mathbf{b}} \leq \gamma_{\mathbf{a} \rightarrow \mathbf{c}} + \gamma_{\mathbf{c} \rightarrow \mathbf{b}}$ for each $j = 1, \dots, |A|$, by analyzing all possible values of \mathbf{a} , \mathbf{b} and \mathbf{c} . The case when $\mathbf{a} = -$ or $\mathbf{b} = -$ can be checked by definition of $\gamma \in \mathbb{M}^W$. Thus, we assume that $\mathbf{a} \neq -$ and $\mathbf{b} \neq -$, which implies by the alignment construction in COMPATIBLEALIGN, that $\mathbf{c} \neq -$. Since $\mathbf{a} \neq -$, $\mathbf{b} \neq -$, $\mathbf{c} \neq -$ and Y is a starsplitting, we have that $\gamma_{\mathbf{a} \rightarrow -} > \gamma_{\mathbf{c} \rightarrow \mathbf{a}}$, $\gamma_{\mathbf{b} \rightarrow -} > \gamma_{\mathbf{c} \rightarrow \mathbf{b}}$ and, since $\gamma \in \mathbb{M}^W$, $\gamma_{- \rightarrow \mathbf{b}} = \gamma_{\mathbf{b} \rightarrow -} > \gamma_{\mathbf{c} \rightarrow \mathbf{b}}$. Since $\gamma_{\mathbf{a} \rightarrow -} > \gamma_{\mathbf{c} \rightarrow \mathbf{a}}$, it follows from Proposition 5.2 that $\gamma_{\mathbf{a} \rightarrow -} > \gamma_{\mathbf{a} \rightarrow \mathbf{c}}$. Hence, $\gamma_{\mathbf{a} \rightarrow -} + \gamma_{- \rightarrow \mathbf{b}} > \gamma_{\mathbf{a} \rightarrow \mathbf{c}} + \gamma_{\mathbf{c} \rightarrow \mathbf{b}}$, which implies from the definition of \mathbb{M}^W that $\gamma_{\mathbf{a} \rightarrow \mathbf{b}} \leq \gamma_{\mathbf{a} \rightarrow \mathbf{c}} + \gamma_{\mathbf{c} \rightarrow \mathbf{b}}$.

Finally, we show (ii). Here, it is enough to prove that $\gamma_{\mathbf{a} \rightarrow \mathbf{b}} \leq 2 \cdot Q_{\max}$ for each column $[\mathbf{a}, \mathbf{b}]$ of $vN_\gamma[A_{\{h,i\}}]$. Again, the case when $\mathbf{a} = -$ or $\mathbf{b} = -$ can easily be checked. Thus, assume that $\mathbf{a} \neq -$ and $\mathbf{b} \neq -$ which implies by construction that $\mathbf{c} \neq -$. Since Y is a splitting, it follows that $\gamma_{\mathbf{c} \rightarrow \mathbf{a}} < \gamma_{\mathbf{a} \rightarrow -}$ and $\gamma_{\mathbf{c} \rightarrow \mathbf{b}} < \gamma_{\mathbf{b} \rightarrow -}$. Since $\gamma_{\mathbf{c} \rightarrow \mathbf{a}} < \gamma_{\mathbf{a} \rightarrow -}$, it follows from Proposition 5.2 that $\gamma_{\mathbf{a} \rightarrow \mathbf{c}} < \gamma_{\mathbf{a} \rightarrow -}$. It follows from (i) that $\gamma_{\mathbf{a} \rightarrow \mathbf{b}} \leq \gamma_{\mathbf{a} \rightarrow \mathbf{c}} + \gamma_{\mathbf{c} \rightarrow \mathbf{b}} < \gamma_{\mathbf{a} \rightarrow -} + \gamma_{\mathbf{b} \rightarrow -} \leq Q_{\max} + Q_{\max} = 2 \cdot Q_{\max}$. Consequently, we have that $vN_\gamma[A_{\{h,i\}}] = vA_\gamma[A_{\{h,i\}}] / |A_{\{h,i\}}| \leq 2 \cdot Q_{\max} |A_{\{h,i\}}| / |A_{\{h,i\}}| = 2 \cdot Q_{\max}$. □

Lemma 5.5. *Let S be a k -sequence, X a star of S with center c , Y a X -starsplitting and $A = \text{COMPATIBLEALIGN}(Y)$. Then,*

$$(i) \ vA_\gamma[A_{\{h,i\}}] \leq vA_\gamma[A_{\{h,c\}}] + vA_\gamma[A_{\{c,i\}}] \text{ when } \gamma \in \mathbb{M}^W \text{ and}$$

$$(ii) \ vN_\gamma[A_{\{h,i\}}] \leq 2 \cdot \left(vN_\gamma[A_{\{h,c\}}] + vN_\gamma[A_{\{c,i\}}] \right) \text{ when } \gamma \in \mathbb{M}^N,$$

for each $h < i$, $h \neq c$, $i \neq c$.

Proof. Consider $A = [s'_1, \dots, s'_k]$, $h, i \in \mathbb{N}$ and $J = \{j : s'_c(j) \neq - \text{ and } s'_h(j) = s'_i(j) = -\}$. Then,

$$\begin{aligned} vA_\gamma[A_{\{h,i\}}] + \sum_{j \in J} (\gamma_{\rightarrow s'_c(j)} + \gamma_{s'_c(j) \rightarrow -}) &= \sum_{j \notin J} \gamma_{s'_h(j) \rightarrow s'_i(j)} + \sum_{j \in J} (\gamma_{\rightarrow s'_c(j)} + \gamma_{s'_c(j) \rightarrow -}) \\ &\leq \sum_{j \notin J} \left(\gamma_{s'_h(j) \rightarrow s'_c(j)} + \gamma_{s'_c(j) \rightarrow s'_i(j)} \right) + \sum_{j \in J} (\gamma_{\rightarrow s'_c(j)} + \gamma_{s'_c(j) \rightarrow -}) \\ &= vA_\gamma[A_{\{h,c\}}] + vA_\gamma[A_{\{c,i\}}], \end{aligned}$$

where the inequality holds due to Proposition 5.4. Therefore,

$$vA_\gamma[A_{\{h,i\}}] \leq vA_\gamma[A_{\{h,c\}}] + vA_\gamma[A_{\{c,i\}}] - \sum_{j \in X} (\gamma_{\rightarrow s'_c(j)} + \gamma_{s'_c(j) \rightarrow -}). \quad (14)$$

Since $\gamma \in \mathbb{M}^W$, we have that $\gamma_{\rightarrow s'_c(j)}, \gamma_{s'_c(j) \rightarrow -} > 0$. It follows from (14) that (i) is proven.

For (ii), observe that, by definition of \mathbb{M}^W , we have that

$$Q_{\max} = \max_{\sigma \in \Sigma} \{ \gamma_{\sigma \rightarrow -}, \gamma_{\rightarrow \sigma} \} = \max_{\sigma \in \Sigma} \{ \gamma_{\sigma \rightarrow -} \} \leq 2 \gamma_{s'_c(j) \rightarrow -} = \gamma_{\rightarrow s'_c(j)} + \gamma_{s'_c(j) \rightarrow -}$$

for every j . Furthermore, following these statements, we have that

$$\begin{aligned} vN_\gamma[A_{\{h,i\}}] &= \frac{vA_\gamma[A_{\{h,i\}}]}{|A_{\{h,i\}}|} \\ &\leq \frac{vA_\gamma[A_{\{h,i\}}] + 2 \cdot Q_{\max} |J|}{|A_{\{h,i\}}| + |J|} \leq 2 \cdot \frac{vA_\gamma[A_{\{h,i\}}] + Q_{\max} |J|}{|A_{\{h,i\}}| + |J|} \end{aligned} \quad (15)$$

$$\leq 2 \cdot \frac{vA_\gamma[A_{\{h,c\}}] + vA_\gamma[A_{\{c,i\}}] - \sum_{j \in J} (\gamma_{\rightarrow s'_c(j)} + \gamma_{s'_c(j) \rightarrow -}) + Q_{\max} |J|}{|A_{\{h,i,c\}}| - |J| + |J|} \quad (16)$$

$$\leq 2 \cdot \frac{vA_\gamma[A_{\{h,c\}}] + vA_\gamma[A_{\{c,i\}}] - Q_{\max} |J| + Q_{\max} |J|}{|A_{\{h,i,c\}}| - |J| + |J|} = 2 \cdot \left(\frac{vA_\gamma[A_{\{h,c\}}]}{|A_{\{h,i,c\}}|} + \frac{vA_\gamma[A_{\{c,i\}}]}{|A_{\{h,i,c\}}|} \right) \quad (17)$$

$$\leq 2 \cdot \left(\frac{vA_\gamma[A_{\{h,c\}}]}{|A_{\{h,c\}}|} + \frac{vA_\gamma[A_{\{c,i\}}]}{|A_{\{c,i\}}|} \right) = 2 \cdot \left(vN_\gamma[A_{\{h,c\}}] + vN_\gamma[A_{\{c,i\}}] \right), \quad (18)$$

where the first inequality of (15) is a consequence of Proposition 5.4 and the second inequality follows since every entry of γ is nonnegative, (16) follows from (14) and from $|A_{\{h,i\}}| = |A_{\{h,i,c\}}| - |J|$, (17) follows as a consequence of $\gamma \in \mathbb{M}^N$, and (18) follows as a consequence of $|A_{\{h,c\}}| \leq |A_{\{h,i,c\}}|$ and $|A_{\{c,i\}}| \leq |A_{\{h,i,c\}}|$. \square

Observe now that the running time of COMPATIBLEALIGN is $O(k^2n)$.

Theorem 5.6. *Let S be a k -sequence and γ be a scoring matrix. Then, Algorithm 5 computes $v[A]$ correctly,*

Algorithm 5

Input: k -sequence S

Output: $v[A]$ such that $A \in \mathcal{A}_S$ and

$v\text{SP}_\gamma[A] \leq 6 \cdot \text{optSP}_\gamma(S)$ if $v = vA_\gamma$ and $\gamma \in \mathbb{M}^W$, and

$V_\gamma^2[A] \leq 12 \cdot \text{optNSP}_\gamma^2(S)$ if $v = vN_\gamma$ and $\gamma \in \mathbb{M}^N$.

- 1: Let X be a v -optimal star of S with center c
 - 2: Compute the X -splitting Y
 - 3: $A \leftarrow \text{COMPATIBLEALIGN}(Y)$
 - 4: **return** $v[A]$
-

(i) in $O(k^2n^2)$ -time such that $v\text{SP}_\gamma[A] \leq 6 \cdot \text{optSP}_\gamma(S)$, if $v = vA_\gamma$ and $\gamma = \mathbb{M}^W$,

(ii) in $O(k^2n^3)$ -time such that $V_\gamma^2[A] \leq 12 \cdot \text{optNSP}_\gamma^2(S)$, if $v = vN_\gamma$ and $\gamma \in \mathbb{M}^N$,

where A is the alignment of S computed by the algorithm.

Proof. Clearly, the value returned by the Algorithm 5 is a score of an alignment of S .

We show then that the approximation factor is as expected. Let c be a center of the stars X and Y found in the first two steps of Algorithm 5. Notice then that

$$\sum_{h=1}^{k-1} \sum_{i=h+1}^k (v[A_{\{h,c\}}] + v[A_{\{c,i\}}]) = (k-1) \cdot c\text{Star}(Y) \quad (19)$$

$$\leq 3 \cdot (k-1) \cdot c\text{Star}(D(X)) \quad (20)$$

$$\leq 3 \cdot (k-1) \cdot \frac{2}{k} \cdot \text{opt}(S) = 6 \cdot \frac{k-1}{h} \cdot \text{opt}(S) \leq 6 \cdot \text{opt}(S), \quad (21)$$

where the equality (19) follows since c is the center of star Y which is compatible with alignment A , (20) follows from Lemma 5.3 and (21) follows from Lemma 5.1.

Suppose then that $v = vA_\gamma$ and $\gamma \in \mathbb{M}^W$. Thus,

$$\begin{aligned} v\text{SP}_\gamma[A] &= \sum_{h=1}^{k-1} \sum_{i=h+1}^k vA_\gamma[A_{\{h,i\}}] \\ &\leq \sum_{h=1}^{k-1} \sum_{i=h+1}^k (vA_\gamma[A_{\{h,c\}}] + vA_\gamma[A_{\{c,i\}}]) \leq 6 \cdot \text{optSP}_\gamma(S), \end{aligned}$$

where the first inequality follows from Lemma 5.5 and the second follows from Equation (21).

Suppose now that $v = vN_\gamma$ and $\gamma \in \mathbb{M}^N$. Thus,

$$\begin{aligned} V_2^\gamma[A] &= \sum_{h=1}^{k-1} \sum_{i=h+1}^k vN_\gamma[A_{\{h,i\}}] \\ &\leq 2 \cdot \sum_{h=1}^{k-1} \sum_{i=h+1}^k (vN_\gamma[A_{\{h,c\}}] + vN_\gamma[A_{\{c,i\}}]) \\ &\leq 2 \cdot 6 \cdot \text{optNSP}_2^\gamma(S) = 12 \cdot \text{optNSP}_2^\gamma(S), \end{aligned}$$

where, similarly, the first inequality follows from Lemma 5.5 and the second follows from Equation (21).

The time required to find an optimal v -star is the time to compute the pairwise alignments of S , which is $\binom{k}{2}O(n^2)$ if $v = vA_\gamma$ and it is $\binom{k}{2}O(n^3)$ if $v = vN_\gamma$. Additionally, we have to consider the time to determine the optimal star, which is $O(k^2)$, implying that the time required to compute line 1 of Algorithm 5 is $\binom{k}{2}O(n^2) + O(k^2) = O(k^2n^2)$ if $v = vA_\gamma$ and $\binom{k}{2}O(n^3) + O(k^2) = O(k^2n^3)$ if $v = vN_\gamma$. The time spent to compute lines 2 and 3 are $O(kn)$ and $O(k^2n)$, respectively, and to compute line 4 is $O(k^3n)$, since we have to compute the score of $\binom{k}{2} = O(k^2)$ pairwise alignments of length $O(kn)$. Therefore, the total time spent by the algorithm is $O(k^2n^2 + k^3n)$ if $v = vA_\gamma$ and $O(k^2n^3 + k^3n)$ if $v = vN_\gamma$. \square

6 Conclusion and future work

We presented and discussed multiple aspects of normalized multiple sequence alignment (NMSA). We defined three new criteria for computing normalized scores when aligning multiple sequences, showing the NP-hardness and exact algorithms for solving the **NMSA- z** given each criterion $z = 1, 2, 3$. In addition, we adapted an existing 2-approximation algorithm for **MSA** when the scoring matrix γ is in the classical class \mathbb{M}^C , leading to a 6-approximation algorithm for **MSA** when γ is in the broader class $\mathbb{M}^W \supseteq \mathbb{M}^C$ and to a 12-approximation for **NMSA-2** when γ is in $\mathbb{M}^N \subseteq \mathbb{M}^W$, a slightly more restricted class such that the cost of deletion for any symbol is at most twice the cost for any other. We summarize these contributions in Table=reftableconclusion.

problems	time of exact algorithms	time of approximation algorithm
MSA	—	$O(k^2n^2 + k^3n)$
NMSA-1	$O(2^k k^3 (n+1)^{k+1})$	—
NMSA-2	$O\left(\left(1 + \frac{1}{2n+1}\right)^k (2n+1)^{k^2} k^2\right)$	$O(k^2n^2 + k^3n)$
NMSA-3	$O(2^k k^4 (n+1)^{k+1})$	—

Table 1: We are considering a k -sequence where each sequence has maximum length of n ; all the problem decision version are NP-complete.

This work is an effort to expand the boundaries of multiple sequence alignment algorithms towards normalization, an unexplored domain that can produce results with higher accuracy in some applications. In future work, we will implement our algorithms in order to verify how large are the sequences our algorithms are able to handle. Also, we plan to perform practical experiments, measuring how well alignments provided by our algorithms and other MSA algorithms agree with multiple alignment benchmarks. In addition, we intend to measure the accuracy of phylogenetic tree reconstruction based on our alignments for simulated and real genomes. Finally, we will work on heuristics and parallel versions of our algorithms in order to faster process large datasets.

References

- [AE99] A. N. Arslan and Ö. Egecioglu. An efficient uniform-cost normalized edit distance algorithm. In *Proc. of SPIRE*, pages 9–15. IEEE, 1999.
- [AG97] A. Apostolico and Z. Galil. *Pattern Matching Algorithms*. Oxford University Press, 1997.
- [AKO10] A. Andoni, R. Krauthgamer, and K. Onak. Polylogarithmic approximation for edit distance and the asymmetric query complexity. In *Proc. of FOCS*, pages 377–386. IEEE, 2010.

- [AS06] E. Araujo and J. Soares. Scoring matrices that induce metrics on sequences. In *Proc. of LATIN*, pages 68–79, 2006.
- [AT00] A. Abbott and A. Tsay. Sequence analysis and optimal matching methods in sociology: Review and prospect. *Sociol Method Res*, 29(1):3–33, 2000.
- [BI18] A. Backurs and P. Indyk. Edit distance cannot be computed in strongly subquadratic time (unless SETH is false). *SIAM J Comput*, 47(3):1087–1097, 2018.
- [BL02] R. Barzilay and L. Lee. Bootstrapping lexical choice via multiple-sequence alignment. In *Proc. of EMNLP*, pages 164–171, USA, 2002. Association for Computational Linguistics.
- [CB99] J. A. Cuff and G. J. Barton. Evaluation and improvement of multiple sequence methods for protein secondary structure prediction. *Proteins*, 34(4):508–519, 1999.
- [CDG⁺20] D. Chakraborty, D. Das, E. Goldenberg, M. Koucký, and M. Saks. Approximating edit distance within constant factor in truly sub-quadratic time. *J ACM*, 67(6):1–22, 2020.
- [Chv80] V. Chvátal. Recognizing intersection patterns. *Ann Discrete Math*, 8:249–251, 1980.
- [CL88] H. Carrillo and D. Lipman. The multiple sequence alignment problem in biology. *SIAM J Appl Math*, 48(5):1073–1082, 1988.
- [CLZU03] Maxime Crochemore, Gad M Landau, and Michal Ziv-Ukelson. A subquadratic sequence alignment algorithm for unrestricted scoring matrices. *SIAM journal on computing*, 32(6):1654–1673, 2003.
- [Eli06] I. Elias. Settling the intractability of multiple alignment. *J Comput Biol*, 13(7):1323–1339, 2006.
- [FD87] D.-F. Feng and R. F. Doolittle. Progressive sequence alignment as a prerequisite to correct phylogenetic trees. *J Mol Evol*, 25(4):351–360, 1987.
- [Gus93] D. Gusfield. Efficient methods for multiple sequence alignment with guaranteed error bounds. *Bull Math Biol*, 55(1):141–154, 1993.
- [HAR09] W. Haque, A. Aravind, and B. Reddy. Pairwise sequence alignment algorithms: A survey. In *Proc. of ISTA*, pages 96–103. ACM Press, 2009.
- [HTHI95] M. Hirosawa, Y. Totoki, M. Hoshida, and M. Ishikawa. Comprehensive study on iterative algorithms of multiple sequence alignment. *Bioinformatics*, 11(1):13–18, 1995.
- [Lev66] V. I. Levenshtein. Binary codes capable of correcting deletions, insertions and reversals. *Dokl. Akad. Nauk.*, 10(8):707–710, 1966.
- [MP80] W. J. Masek and M. S. Paterson. A faster algorithm computing string edit distances. *J Comput Syst Sci*, 20(1):18–31, 1980.
- [MV93] A. Marzal and E. Vidal. Computation of normalized edit distance and applications. *IEEE T Pattern Anal*, 15(9):926–932, 1993.
- [NW70] S. B. Needleman and C. D. Wunsch. A general method applicable to the search for similarities in the amino acid sequence of two proteins. *J Mol Biol*, 48(3):443–453, 1970.
- [OR06] T. H. Ogden and M. S. Rosenberg. Multiple sequence alignment accuracy and phylogenetic inference. *Syst Biol*, 55(2):314–328, 2006.
- [Sel74] P. H. Sellers. On the theory and computation of evolutionary distances. *SIAM J Appl Math*, 26(4):787–793, 1974.

- [SH14] F. Sievers and D. G. Higgins. Clustal Omega. *Curr Protoc Bioinfo*, 48(1):3.13.1–3.13.16, 2014.
- [SWD⁺11] F. Sievers, A. Wilm, D. Dineen, T. J. Gibson, K. Karplus, W. Li, R. Lopez, H. McWilliam, M. Remmert, J. Söding, J. D. Thompson, and D. G. Higgins. Fast, scalable generation of high-quality protein multiple sequence alignments using Clustal Omega. *Mol Syst Biol*, 7(1):539, 2011.
- [TPP99] J. D. Thompson, F. Plewniak, and O. Poch. A comprehensive comparison of multiple sequence alignment programs. *Nucleic Acids Res*, 27(13):2682–2690, 1999.
- [VMA95] E. Vidal, A. Marzal, and P. Aibar. Fast computation of normalized edit distances. *IEEE T Pattern Anal*, 17(9):899–902, 1995.
- [WLXZ15] X.-D. Wang, J.-X. Liu, Y. Xu, and J. Zhang. A survey of multiple sequence alignment techniques. In *Proc. of ICIC*, pages 529–538. Springer, 2015.
- [WOHN06] I. M. Wallace, O. O’Sullivan, D. G. Higgins, and C. Notredame. M-Coffee: combining multiple sequence alignment methods with T-Coffee. *Nucleic Acids Res*, 34(6):1692–1699, 2006.