

Set Values for Mean Field Games*

Melih Iseri[†] Jianfeng Zhang[‡]

October 25, 2021

Abstract

In this paper we study mean field games with possibly multiple mean field equilibria. Instead of focusing on the individual equilibria, we propose to study the set of values over all possible equilibria, which we call the set value of the mean field game. When the mean field equilibrium is unique, typically under certain monotonicity conditions, our set value reduces to the singleton of the standard value function which solves the master equation. The set value is by nature unique, and we shall establish two crucial properties: (i) the dynamic programming principle, also called time consistency; and (ii) the convergence of the set values of the corresponding N -player games. To our best knowledge, this is the first work in the literature which studies the dynamic value of mean field games without requiring the uniqueness of mean field equilibria. We emphasize that the set value is very sensitive to the choice of the admissible controls. In particular, for the convergence one has to restrict to the same type of equilibria for the N -player game and for the mean field game. We shall illustrate this point by investigating three cases, two in finite state space models and the other in a diffusion model.

MSC2020. 91A16, 60H30, 91A25, 91A06, 93E20

Keywords. Mean field games, mean field equilibria, set values, dynamic programming principle, closed loop controls, relaxed controls

*The research is supported in part by NSF grant DMS-1908665, and the authors would like to thank Daniel Lacker for insightful discussions.

[†]Department of Mathematics, University of Southern California, United States, melihise@usc.edu.

[‡]Department of Mathematics, University of Southern California, United States, jianfenz@usc.edu.

1 Introduction

In this paper we study Mean Field Games (MFG, for short) without monotonicity conditions. There are typically multiple Mean Field Equilibria (MFE, for short) with possibly different values. Instead of focusing on the individual equilibria, we propose to study the set of values over all equilibria, which we call the set value of the MFG. Note that the set value always exists (with empty set as a possible value) and is by definition unique. When the MFE is unique, typically under certain monotonicity conditions, our set value is reduced to the singleton of the standard value function of the game, which solves the so called master equation. So the set value can be viewed as the counterpart of the standard value function, but for MFGs without monotonicity conditions, and it indeed enjoys many nice properties of the standard value function. In this paper, we shall focus particularly on two crucial properties of the set value:

- the Dynamic Programming Principle (DPP, for short), or say the time consistency;
- the convergence of the set values of the corresponding N -player games.

For general theory of MFGs, we refer to the classical references Caines-Huang-Malhame [7], Lasry-Lions [33], Lions [35], Cardaliaguet [8], Bensoussan-Frehse-Yam [6], and Camona-Delarue [13, 14].

In standard stochastic control theory, it is well known that the dynamic value function satisfies the DPP. In fact, this is the underlying reason for the PDE approach to work. For MFGs under appropriate monotonicity conditions, the value function (over the unique MFE) also satisfies the DPP, which, together with the Ito formula, leads to the master equation. However, with the presence of multiple equilibria (see, e.g., Bardi-Fischer [2] for some examples), to our best knowledge this is the first work in the literature to study the MFG dynamically and to address the time consistency issue. We show that, when formulated properly, the dynamic set value function satisfies the DPP. This also opens the door to a possible PDE approach for these general games by introducing the so called set valued PDE, which we shall leave for future research. Our set value approach follows from Feinstein-Rudloff-Zhang [24], which studies nonzero sum games with finitely many players. See also the related works Abreu-Pearce-Stacchetti [1] and Sannikov [40] in economics literature, as well as Feinstein [23] which studies the set of equilibria instead of values.

We note that the set value of games relies heavily on the types of admissible controls we use. In this paper we shall consider closed loop controls. The open loop equilibria of games are typically time inconsistent, see e.g. Buckdahn's counterexample in Pham-Zhang [38, Appendix E] for a two person zero sum game, and consequently, the set value of games

with open loop controls would violate the DPP. For the MFG, noting that the required symmetry decomposes the game problem into a standard control problem and a fixed point problem of measures, and that open loop and closed loop controls yield the same value function for a standard control problem, it is possible that the set value with open loop controls still satisfies the DPP. Nevertheless, bearing in mind the DPP of the set value for more general (non-symmetric) games, as well as the practical consideration in terms of the information available to the players, we shall focus on closed loop controls. There is also a very subtle path dependence issue. While the game parameters are state dependent, we may consider both state dependent and path dependent controls. For general non-zero sum games (not mean field type), [24] shows that DPP holds for the set value for path dependent controls, but in general fails for the set value for state dependent controls. For MFGs with closed loop controls, again due to the required symmetric properties, the set values for both state dependent controls and path dependent controls will satisfy the DPP, but they are in general not equal. For MFGs with closed loop relaxed controls, or say closed loop mixed strategies, however, it turns out that the state dependent controls and the path dependent controls induce the same set value which still satisfies the DPP.

For the convergence of the N -player games, in terms of the equilibria instead of the values, there have been many publications on open loop controls. We refer to Camona-Delarue [12], Feleqi [25], Fischer [26], Fischer-Silva [27], Lacker [30], Lasry-Lions [33], Lauriere-Tangpi [34], Nutz-San Martin-Tan [37]. In particular, [30] provides the full characterization for the convergence: any limit of approximate Nash equilibria of N -player games is a weak MFE, and conversely any weak MFE can be obtained as such a limit. The work [26] is also in this direction.

We are interested in the convergence analysis for closed loop controls. In the realm of master equations, again under certain monotonicity conditions, one can show that the empirical measure of the closed loop equilibria of the N -player games converge to the unique MFE, and the corresponding values also converge. See Cardaliaguet-Delarue-Lasry-Lions [10], followed by Bayraktar-Cohen [3], Cardaliaguet [9], Cecchin-Pelino [17], Delarue-Lacker-Ramanan [20, 21], Gangbo-Meszaros [29], and Mou-Zhang [36], to mention a few. The situation becomes very subtle for the general case. Our work is highly inspired by the seminal paper Lacker [31], whose main result reads roughly as follows:

$$\{\text{Strong MFEs}\} \subset \{\text{Limit of } N\text{-player approx. equilibria}\} \subset \{\text{Weak MFEs}\}. \quad (1.1)$$

Here an MFE is strong if it depends only on the state processes, and weak if it allows for additional randomness. However, the equalities in (1.1) were not clear in [31], in particu-

lar, the left inclusion is known to be strict in general. Our goal is to establish the exact convergence, namely equality in (1.1), for closed loop equilibria and more importantly for the corresponding set values.

We emphasize again that the set value is very sensitive to the types of admissible controls. To ensure the convergence, one simple but crucial observation is that the N -player game and the MFG should use the "same" type of controls (more precisely, corresponding types of controls in appropriate sense). We illustrate this point by considering two cases. Note that in the standard literature each player is required to use the same closed loop control along an MFE. For the first case, we will obtain the desired convergence by restricting the N -player game to homogeneous equilibria, namely each player also uses the same closed loop control. In the second case, we remove such restriction and consider heterogenous equilibria for the N -player games. Note that a closed loop control means the control depends only on the state. In this heterogenous case players with the same state may choose different controls, then one can not expect in the limit they will have to use the same control. Indeed, in this case the limit is characterized by the MFG with closed loop relaxed controls, or say closed loop mixed strategies, which exactly means players with the same state may still have a distribution of controls to choose from. For the homogeneous case, we will investigate both a discrete time model with finite state space and a continuous time diffusion model with drift controls. But for the heterogeneous case we will investigate the discrete model only. The diffusion model in such case involves some technical challenges for the convergence and we shall leave it for future research. We shall point out that, however, the DPP would hold in much more general models without significant difficulties.

We note that the relax control we consider for the MFG is still homogeneous, namely each player uses the same relaxed control. Even for the discrete model, the convergence in the heterogenous case is quite involved, because the heterogenous closed loop controls for the N -player game and the homogeneous relaxed control for the MFG appear in different forms. Our main idea is to introduce a new alternative formulation for the MFG, which on one hand mimics the structure of the heterogenous controls of the N -player games and on the other hand shares the same set value as the relaxed control formulation of the MFG.

To ensure the convergence, another main feature is that we define the set value as the limit of the approximate set values over approximate equilibria, rather than the set of values over true equilibria. We call the latter the raw set value, and both the set value and the raw set value satisfy the DPP. However, the raw set value is extremely sensitive to small perturbations of the game parameters, in fact, in general even its measurability is not clear, so one can hardly expect the convergence for the raw set values. In the standard

control theory, the value function is defined as the infimum of controlled values, which is exactly the limit of values over approximate optimal controls, rather than the value over true optimal controls which may not even exist. So our set value, not the raw set value, is the natural extension of the standard value function in control theory. Moreover, since we are considering infinitely many players, an approximate equilibrium means it is approximately optimal for most players, but possibly with a small portion of exceptions, as introduced in Carmona [11].

Moreover, we will impose certain regularity on the controls with respect to the measure variable, which helps greatly for our convergence analysis. This is different from [31] which uses compactness arguments without relying on such regularity. So our set of admissible controls is somewhat smaller.

We shall also mention the very recent paper Lacker-Flem [32], which extends [31] to MFGs with common noise and is circulated right after the first version of our paper. In particular, [32] shows that, by adding additional random signal to the controls of the N -player games, the second inclusion in (1.1) will become an equality. Recall that a weak MFE allows for additional randomness. This confirms further our observation that the convergence would hold true if we use appropriate corresponding types of controls for the N -player game and for the MFG. Furthermore, a more recent paper Djete [22] essentially shows that, in a slightly different setting, the second inclusion in (1.1) is actually an equality. However, we should point out that [31, 32, 22] consider controls with full information, namely the controls may depend on the states of all players. We instead use symmetric controls, where a player's control depends only on his/her own state and the empirical measure of the other players' states. We believe such restriction is reasonable in practice since we are considering very large N .

Finally we would like to mention some other approaches for MFGs with multiple equilibria. One is to add sufficient (possibly infinitely dimensional) noise so that the new game will become non-degenerate and hence have unique MFE, see e.g. Bayraktar-Cecchin-Cohen-Delarue [4, 5], Delarue [18], Delarue-Foguen Tchuendom [19], Foguen Tchuendom [28]. Another approach is to study a special type of MFEs, see e.g. Cecchin-Dai Pra-Fisher-Pelino [15], Cecchin-Delarue [16], and [19]. Another interesting work is Possamai-Tangpi [39] which introduces an additional parameter function Λ such that the MFE corresponding to any fixed Λ is unique and then the desired convergence is obtained.

The rest of the paper is organized as follows. In Section 2 we introduce the set value for an MFG in a discrete time model on finite state space and establish the DPP, and in Section 3 we prove the convergence for the corresponding N -player games with homogeneous

equilibria. Sections 4 and 5 are devoted to MFGs with relaxed controls and the corresponding N -player games with heterogenous equilibria. In Section 6 we study a diffusion model. Finally in Appendix we provide some discussion on the subtle path dependence issue, and complete some technical proofs.

2 Mean field games on finite space with closed loop controls

In this section we consider an MFG on finite space (both time and state are finite) with closed loop controls, and for simplicity we restrict to state dependent setting. Since the game typically has multiple MFEs which may induce different values, see Example 7.1 below for an example, we shall introduce the set value of the game over all MFEs. Our goal is to establish the DPP for the MFG set value, and we shall show in the next section that the set values of the corresponding N -player games converge to the MFG set value.

2.1 The basic setting

Let $\mathbb{T} := \{0, \dots, T\}$ be the set of discrete times; $\mathbb{T}_t := \{t, \dots, T\}$ for $t \in \mathbb{T}$; \mathbb{S} the finite state space¹ with size $|\mathbb{S}| = d$; $\mathcal{P}(\mathbb{S})$ the set of probability measures on \mathbb{S} , equipped with the 1-Wasserstein distance W_1 . Since \mathbb{S} is finite, W_1 is equivalent to the total variation distance² which is more convenient for our purpose: by abusing the notation W_1 ,

$$W_1(\mu, \nu) := \sum_{x \in \mathbb{S}} |\mu(x) - \nu(x)|, \quad \mu, \nu \in \mathcal{P}(\mathbb{S}). \quad (2.1)$$

Let $\mathcal{P}_0(\mathbb{S})$ denote the subset of $\mu \in \mathcal{P}(\mathbb{S})$ which has full support, namely $\mu(x) > 0$ for all $x \in \mathbb{S}$. Moreover, let $\mathbb{A} \subset \mathbb{R}^{d_0}$ be a measurable set from which the controls take values; and $q : \mathbb{T} \times \mathbb{S} \times \mathcal{P}(\mathbb{S}) \times \mathbb{A} \times \mathbb{S} \rightarrow (0, 1)$ be a transition probability function:

$$\sum_{\tilde{x} \in \mathbb{S}} q(t, x, \mu, a; \tilde{x}) = 1, \quad \forall (t, x, \mu, a) \in \mathbb{T} \times \mathbb{S} \times \mathcal{P}(\mathbb{S}) \times \mathbb{A}.$$

We shall use the weak formulation which is more convenient for closed loop controls. That is, we fix the canonical space and consider controlled probability measures on it. To be precise, let $\Omega := \mathbb{X} := \mathbb{S}^{\mathbb{T}+1}$ be the canonical space; $X : \mathbb{T} \times \Omega \rightarrow \mathbb{S}$ the canonical process: $X_t(\omega) = \omega_t$; $\mathbb{F} := \{\mathcal{F}_t\}_{t \in \mathbb{T}} := \mathbb{F}^X$ the filtration generated by X ; and \mathcal{A}_{state} the set of state dependent admissible controls $\alpha : \mathbb{T} \times \mathbb{S} \rightarrow \mathbb{A}$. Introduce the concatenation for controls:

$$(\alpha \oplus_{T_0} \tilde{\alpha})_s := \alpha_s \mathbf{1}_{\{s < T_0\}} + \tilde{\alpha}_s \mathbf{1}_{\{s \geq T_0\}}, \quad \alpha, \tilde{\alpha} \in \mathcal{A}_{state}. \quad (2.2)$$

¹We may allow to have different state space \mathbb{S}_t at different time t and all the results in this paper will remain true.

²More precisely, the total variation distance is $\frac{1}{2}W_1$ for the W_1 in (2.1).

It is clear that $\alpha \oplus_{T_0} \tilde{\alpha} \in \mathcal{A}_{state}$. Given $(t, \mu, \alpha) \in \mathbb{T} \times \mathcal{P}(\mathbb{S}) \times \mathcal{A}_{state}$, let $\mathbb{P}^{t, \mu, \alpha}$ denote the probability measure on \mathcal{F}_T determined recursively by: for $s = t, \dots, T$,

$$\begin{aligned} \mathbb{P}^{t, \mu, \alpha} \circ X_t^{-1} = \mu, \quad \mathbb{P}^{t, \mu, \alpha}(X_{s+1} = \tilde{x} | X_s = x) = q(s, x, \mu_s^\alpha, \alpha(s, x); \tilde{x}); \\ \text{where } \mu_s^\alpha := \mathbb{P}^{t, \mu, \alpha} \circ X_s^{-1}. \end{aligned} \quad (2.3)$$

We note that $\{\mu_s^\alpha\}_{s \in \mathbb{T}_t}$ are uniquely determined and X is a Markov chain on \mathbb{T}_t under $\mathbb{P}^{t, \mu, \alpha}$. However, the distribution of $\{X_s\}_{s=0, \dots, t-1}$ is not specified and is irrelevant, and $\{\alpha_s\}_{0 \leq s < t}$ is also irrelevant. Moreover, given $\{\mu.\} := \{\mu_s\}_{s \in \mathbb{T}_t}$, $x \in \mathbb{S}$, and $\tilde{\alpha} \in \mathcal{A}_{state}$, let $\mathbb{P}^{\{\mu.\}; t, x, \tilde{\alpha}}$ denote the probability measure on \mathcal{F}_T determined recursively by: for $s = t, \dots, T-1$,

$$\mathbb{P}^{\{\mu.\}; t, x, \tilde{\alpha}}(X_t = x) = 1, \quad \mathbb{P}^{\{\mu.\}; t, x, \tilde{\alpha}}(X_{s+1} = \tilde{x} | X_s = \tilde{x}) = q(s, \tilde{x}, \mu_s, \tilde{\alpha}(s, \tilde{x}); \tilde{x}). \quad (2.4)$$

As in the standard MFG literature, here we are assuming that the population uses the common control α while the individual player is allowed to use a different control $\tilde{\alpha}$.

We remark that, since we assume $q > 0$, then for any (t, μ) and $\alpha, \mu_s^\alpha \in \mathcal{P}_0(\mathbb{S})$ for all $s > t$. For the convenience of presentation, in this section we shall restrict our discussion to the case $\mu \in \mathcal{P}_0(\mathbb{S})$. The general case that the initial measure μ is not fully supported can be treated fairly easily, as we will do in Section 6 below. The situation with degenerate q , however, is more subtle and we shall leave for future research.

We finally introduce the cost functional for the MFG: for the μ^α in (2.3),

$$\begin{aligned} J(t, \mu, \alpha; x, \tilde{\alpha}) := J(\mu^\alpha; t, x, \tilde{\alpha}), \quad v(\{\mu.\}; s, x) := \inf_{\tilde{\alpha} \in \mathcal{A}_{state}} J(\{\mu.\}; s, x, \tilde{\alpha}); \\ \text{where } J(\{\mu.\}; s, x, \tilde{\alpha}) := \mathbb{E}^{\mathbb{P}^{\{\mu.\}; s, x, \tilde{\alpha}}} \left[G(X_T, \mu_T) + \sum_{r=s}^{T-1} F(r, X_r, \mu_r, \tilde{\alpha}(r, X_r)) \right]. \end{aligned} \quad (2.5)$$

Here, since \mathbb{T} and \mathbb{S} are finite, F and G are arbitrary measurable functions satisfying

$$\inf_{a \in \mathbb{A}} F(t, x, \mu, a) > -\infty \quad \text{for all } (t, x, \mu).$$

We remark that here $v(\{\mu.\}; \cdot, \cdot)$ is the value function of a standard stochastic control problem with parameter $\{\mu.\}$. In particular, in continuous time models, μ^α and $v(\mu^\alpha; \cdot, \cdot)$ will satisfy the Fokker-Planck equation and the HJB equation, respectively.

Definition 2.1 *Given $(t, \mu) \in \mathbb{T} \times \mathcal{P}_0(\mathbb{S})$, we say $\alpha^* \in \mathcal{A}_{state}$ is a state dependent MFE at (t, μ) , denoted as $\alpha^* \in \mathcal{M}_{state}(t, \mu)$, if*

$$J(t, \mu, \alpha^*; x, \alpha^*) = v(\mu^{\alpha^*}; t, x), \quad \text{for all } x \in \mathbb{S}. \quad (2.6)$$

In this and the next section, we will use the following conditions.

Assumption 2.2 (i) $q \geq c_q$ for some constant $c_q > 0$;

(ii) q is Lipschitz continuous in (μ, a) , with a Lipschitz constant L_q ;

(iii) F, G are bounded by a constant C_0 and uniformly continuous in (μ, a) , with a modulus of continuity function ρ .

2.2 The raw set value \mathbb{V}_0

We introduce the raw set value for the MFG over all state dependent MFEs:

$$\mathbb{V}_0(t, \mu) := \left\{ J(t, \mu, \alpha^*; \cdot, \alpha^*) : \alpha^* \in \mathcal{M}_{state}(t, \mu) \right\} \subset \mathbb{L}^0(\mathbb{S}; \mathbb{R}). \quad (2.7)$$

Here the elements of $\mathbb{V}_0(t, \mu)$ are functions from \mathbb{S} to \mathbb{R} , which coincide with \mathbb{R}^d by identifying $\varphi \in \mathbb{L}^0(\mathbb{S}; \mathbb{R})$ with $(\varphi(x) : x \in \mathbb{S}) \in \mathbb{R}^d$. We call $\mathbb{V}_0(t, \mu)$ the raw set value and we will introduce the set value $\mathbb{V}(t, \mu)$ of the MFG in the next subsection.

Next, for any $T_0 \in \mathbb{T}_t$, $\psi \in \mathbb{L}^0(\mathbb{S} \times \mathcal{P}_0(\mathbb{S}); \mathbb{R})$, we introduce the MFG on $\{t, \dots, T_0\}$:

$$J(T_0, \psi; t, \mu, \alpha; x, \tilde{\alpha}) := \mathbb{E}^{\mathbb{P}^{\mu^\alpha; t, x, \tilde{\alpha}}} \left[\psi(X_{T_0}, \mu_{T_0}^\alpha) + \sum_{s=t}^{T_0-1} F(s, X_s, \mu_s^\alpha, \tilde{\alpha}(s, X_s)) \right]. \quad (2.8)$$

In the obvious sense we define $\alpha^* \in \mathcal{M}_{state}(T_0, \psi; t, \mu)$ by:

$$J(T_0, \psi; t, \mu, \alpha^*; x, \alpha^*) = v(T, \psi; \mu^{\alpha^*}; t, x) := \inf_{\tilde{\alpha} \in \mathcal{A}_{state}} J(T, \psi; t, \mu, \alpha^*; x, \tilde{\alpha}), \quad \forall x \in \mathbb{S}. \quad (2.9)$$

At below we will repeatedly use the following simple fact due to the tower property of conditional expectations:

$$J(t, \mu, \alpha; x, \tilde{\alpha}) = J(T_0, \psi; t, \mu, \alpha; x, \tilde{\alpha}), \quad \text{where} \quad \psi(y, \nu) := J(T_0, \nu, \alpha; y, \tilde{\alpha}). \quad (2.10)$$

The following time consistency of MFE is the essence of the DPP for the raw set value.

Proposition 2.3 Fix $0 \leq t < T_0 \leq T$ and $\mu \in \mathcal{P}_0(\mathbb{S})$. For any $\alpha^*, \tilde{\alpha}^* \in \mathcal{A}_{state}$, denote $\hat{\alpha}^* := \alpha^* \oplus_{T_0} \tilde{\alpha}^*$ and $\psi(y, \nu) := J(T_0, \nu, \tilde{\alpha}^*; y, \tilde{\alpha}^*)$. Then $\hat{\alpha}^* \in \mathcal{M}_{state}(t, \mu)$ if and only if $\alpha^* \in \mathcal{M}_{state}(T_0, \psi; t, \mu)$ and $\tilde{\alpha}^* \in \mathcal{M}_{state}(T_0, \mu_{T_0}^{\alpha^*})$.

Proof (i) We first prove the if part. Let $\alpha^* \in \mathcal{M}_{state}(T_0, \psi; t, \mu)$ and $\tilde{\alpha}^* \in \mathcal{M}_{state}(T_0, \mu_{T_0}^{\alpha^*})$.

For arbitrary $\alpha \in \mathcal{A}_{state}$ and $x \in \mathbb{S}$, by (2.10) we have

$$\begin{aligned}
J(t, \mu, \hat{\alpha}^*; x, \alpha) &= \mathbb{E}^{\mathbb{P}^{\mu^{\alpha^*}; t, x, \alpha}} \left[J(T_0, \mu_{T_0}^{\alpha^*}, \tilde{\alpha}^*; X_{T_0}, \alpha) + \sum_{s=t}^{T_0-1} F(s, X_s, \mu_s^{\alpha^*}, \alpha(s, X_s)) \right] \\
&\geq \mathbb{E}^{\mathbb{P}^{\mu^{\alpha^*}; t, x, \alpha}} \left[J(T_0, \mu_{T_0}^{\alpha^*}, \tilde{\alpha}^*; X_{T_0}, \tilde{\alpha}^*) + \sum_{s=t}^{T_0-1} F(s, X_s, \mu_s^{\alpha^*}, \alpha(s, X_s)) \right] \\
&= \mathbb{E}^{\mathbb{P}^{\mu^{\alpha^*}; t, x, \alpha}} \left[\psi(X_{T_0}, \mu_{T_0}^{\alpha^*}) + \sum_{s=t}^{T_0-1} F(s, X_s, \mu_s^{\alpha^*}, \alpha(s, X_s)) \right] \\
&= J(T_0, \psi; t, \mu, \alpha^*; x, \alpha) \geq J(T_0, \psi; t, \mu, \alpha^*; x, \alpha^*) = J(t, \mu, \hat{\alpha}^*; x, \hat{\alpha}^*),
\end{aligned}$$

where the first inequality is due to $\tilde{\alpha}^* \in \mathcal{M}_{state}(T_0, \mu_{T_0}^{\alpha^*})$ and the second inequality is due to $\alpha^* \in \mathcal{M}_{state}(T_0, \psi; t, \mu)$. Then $\hat{\alpha}^* \in \mathcal{M}_{state}(t, \mu)$.

(ii) We now prove the only if part. Let $\hat{\alpha}^* \in \mathcal{M}_{state}(t, \mu)$. For any $\alpha \in \mathcal{A}_{state}$, we have $\alpha \oplus_{T_0} \tilde{\alpha}^* \in \mathcal{A}_{state}$. Then, since $\hat{\alpha}^* \in \mathcal{M}_{state}(t, \mu)$, for any $x \in \mathbb{S}$, by (2.10) we have

$$J(T_0, \psi; t, \mu, \alpha^*; x, \alpha^*) = J(t, \mu, \hat{\alpha}^*; x, \hat{\alpha}^*) \leq J(t, \mu, \hat{\alpha}^*; x, \alpha \oplus_{T_0} \tilde{\alpha}^*) = J(T_0, \psi; t, \mu, \alpha^*; x, \alpha).$$

This implies that $\alpha^* \in \mathcal{M}_{state}(T_0, \psi; t, \mu)$.

Moreover, note that $\alpha^* \oplus_{T_0} \alpha \in \mathcal{A}_{state}$ and again since $\hat{\alpha}^* \in \mathcal{M}_{state}(t, \mu)$, we have

$$\begin{aligned}
&\mathbb{E}^{\mathbb{P}^{\mu^{\alpha^*}; t, x, \alpha^*}} \left[J(T_0, \mu_{T_0}^{\alpha^*}, \tilde{\alpha}^*; X_{T_0}, \tilde{\alpha}^*) + \sum_{s=t}^{T_0-1} F(s, X_s, \mu_s^{\alpha^*}, \alpha^*(s, X_s)) \right] \\
&= J(t, \mu, \hat{\alpha}^*; x, \hat{\alpha}^*) \leq J(t, \mu, \hat{\alpha}^*; x, \alpha^* \oplus_{T_0} \alpha) \\
&= \mathbb{E}^{\mathbb{P}^{\mu^{\alpha^*}; t, x, \alpha^*}} \left[J(T_0, \mu_{T_0}^{\alpha^*}, \tilde{\alpha}^*; X_{T_0}, \alpha) + \sum_{s=t}^{T_0-1} F(s, X_s, \mu_s^{\alpha^*}, \alpha^*(s, X_s)) \right].
\end{aligned}$$

This implies that, recalling the v in (2.5) and by the standard stochastic control theory,

$$\begin{aligned}
\mathbb{E}^{\mathbb{P}^{\mu^{\alpha^*}; t, x, \alpha^*}} \left[J(T_0, \mu_{T_0}^{\alpha^*}, \tilde{\alpha}^*; X_{T_0}, \tilde{\alpha}^*) \right] &\leq \inf_{\alpha \in \mathcal{A}_{state}} \mathbb{E}^{\mathbb{P}^{\mu^{\alpha^*}; t, x, \alpha^*}} \left[J(T_0, \mu_{T_0}^{\alpha^*}, \tilde{\alpha}^*; X_{T_0}, \alpha) \right] \\
&= \mathbb{E}^{\mathbb{P}^{\mu^{\alpha^*}; t, x, \alpha^*}} \left[v(\mu^{\hat{\alpha}^*}; T_0, X_{T_0}) \right]. \tag{2.11}
\end{aligned}$$

On the other hand, by definition $v(\mu^{\hat{\alpha}^*}; T_0, \tilde{x}) \leq J(T_0, \mu_{T_0}^{\alpha^*}, \tilde{\alpha}^*; \tilde{x}, \tilde{\alpha}^*)$ for all $\tilde{x} \in \mathbb{S}$. Then

$$J(T_0, \mu_{T_0}^{\alpha^*}, \tilde{\alpha}^*; X_{T_0}, \tilde{\alpha}^*) = v(\mu^{\hat{\alpha}^*}; T_0, X_{T_0}), \quad \mathbb{P}^{\mu^{\alpha^*}; t, x, \alpha^*}\text{-a.s.}$$

Since $q > 0$, then clearly $\mathbb{P}^{\mu^{\alpha^*}; t, x, \alpha^*}(X_{T_0} = \tilde{x}) > 0$ for all $\tilde{x} \in \mathbb{S}$. Thus

$$J(T_0, \mu_{T_0}^{\alpha^*}, \tilde{\alpha}^*; \tilde{x}, \tilde{\alpha}^*) = v(\mu^{\hat{\alpha}^*}; T_0, \tilde{x}), \quad \forall \tilde{x} \in \mathbb{S}.$$

This implies that $\tilde{\alpha}^* \in \mathcal{M}_{state}(T_0, \mu_{T_0}^{\alpha^*})$. ■

We then have the following DPP.

Theorem 2.4 For any $0 \leq t < T_0 \leq T$, and $\mu \in \mathcal{P}_0(\mathbb{S})$, we have

$$\mathbb{V}_0(t, \mu) := \left\{ J(T_0, \psi; t, \mu, \alpha^*; \cdot, \alpha^*) : \text{for all } \psi \in \mathbb{L}^0(\mathbb{S} \times \mathcal{P}_0(\mathbb{S}); \mathbb{R}) \text{ and } \alpha^* \in \mathcal{A}_{state} \right. \\ \left. \text{such that } \psi(\cdot, \mu_{T_0}^{\alpha^*}) \in \mathbb{V}_0(T_0, \mu_{T_0}^{\alpha^*}) \text{ and } \alpha^* \in \mathcal{M}_{state}(T_0, \psi; t, \mu) \right\}. \quad (2.12)$$

Proof Let $\tilde{\mathbb{V}}_0(t, \mu)$ denote the right side of (2.12). First, for any $J(T_0, \psi; t, \mu, \alpha^*; \cdot, \alpha^*) \in \tilde{\mathbb{V}}_0(t, \mu)$ with desired ψ, α^* as in (2.12). Since $\psi(\cdot, \mu_{T_0}^{\alpha^*}) \in \mathbb{V}_0(T_0, \mu_{T_0}^{\alpha^*})$, there exists $\tilde{\alpha}^* \in \mathcal{M}_{state}(T_0, \mu_{T_0}^{\alpha^*})$ such that $\psi(\cdot, \mu_{T_0}^{\alpha^*}) = J(T_0, \mu_{T_0}^{\alpha^*}, \tilde{\alpha}^*; \cdot, \tilde{\alpha}^*)$. By Proposition 2.3 we have $\hat{\alpha}^* := \alpha^* \oplus_{T_0} \tilde{\alpha}^* \in \mathcal{M}_{state}(t, \mu)$. Then, by (2.10), $J(T_0, \psi; t, \mu, \alpha^*; \cdot, \alpha^*) = J(t, \mu, \hat{\alpha}^*; \cdot, \hat{\alpha}^*) \in \mathbb{V}_0(t, \mu)$, and thus $\tilde{\mathbb{V}}_0(t, \mu) \subset \mathbb{V}_0(t, \mu)$.

On the other hand, let $J(t, \mu, \alpha^*; \cdot, \alpha^*) \in \mathbb{V}_0(t, \mu)$ with $\alpha^* \in \mathcal{M}_{state}(t, \mu)$. Introduce $\psi(x, \nu) := J(T_0, \nu, \alpha^*; x, \alpha^*)$. By Proposition 2.3 again we see that $\alpha^* \in \mathcal{M}_{state}(T_0, \psi; t, \mu)$ and $\alpha^* \in \mathcal{M}_{state}(T_0, \mu_{T_0}^{\alpha^*})$, and the latter implies further that $\psi(\cdot, \mu_{T_0}^{\alpha^*}) \in \mathbb{V}_0(T_0, \mu_{T_0}^{\alpha^*})$. Then by the definition of $\tilde{\mathbb{V}}_0(t, \mu)$ that $J(t, \mu, \alpha^*; \cdot, \alpha^*) = J(T_0, \psi; t, \mu, \alpha^*; \cdot, \alpha^*) \in \tilde{\mathbb{V}}_0(t, \mu)$. That is, $\mathbb{V}_0(t, \mu) \subset \tilde{\mathbb{V}}_0(t, \mu)$. \blacksquare

2.3 The set value \mathbb{V}_{state}

While Theorem 2.4 is elegant, the raw set value $\mathbb{V}_0(t, \mu)$ is very sensitive to small perturbations of the coefficients F, G and the variable μ . Indeed, even the measurability of the subset $\mathbb{V}_0(t, \mu) \subset \mathbb{R}^d$ and the measurability of the mapping $\mu \mapsto \mathbb{V}_0(t, \mu)$ are not clear to us. Moreover, in general it does not look possible to have the convergence of the raw set value of the corresponding N -player games to $\mathbb{V}_0(t, \mu)$. Therefore, in this subsection we shall modify $\mathbb{V}_0(t, \mu)$ and introduce the set value $\mathbb{V}_{state}(t, \mu)$ of the MFG as follows.

Definition 2.5 (i) For any $(t, \mu) \in \mathbb{T} \times \mathcal{P}_0(\mathbb{S})$ and $\varepsilon > 0$, let $\mathcal{M}_{state}^\varepsilon(t, \mu)$ denote the set of $\alpha^* \in \mathcal{A}_{state}$ such that

$$J(t, \mu, \alpha^*; x, \alpha^*) \leq v(\mu^{\alpha^*}; t, x) + \varepsilon, \quad \text{for all } x \in \mathbb{S}. \quad (2.13)$$

(ii) The set value of the MFG at (t, μ) is defined as:

$$\mathbb{V}_{state}(t, \mu) := \bigcap_{\varepsilon > 0} \mathbb{V}_{state}^\varepsilon(t, \mu), \quad \text{where} \quad (2.14)$$

$$\mathbb{V}_{state}^\varepsilon(t, \mu) := \left\{ \varphi \in \mathbb{L}^0(\mathbb{S}; \mathbb{R}) : \|\varphi - J(t, \mu, \alpha^*; \cdot, \alpha^*)\|_\infty \leq \varepsilon \text{ for some } \alpha^* \in \mathcal{M}_{state}^\varepsilon(t, \mu) \right\}.$$

Recall (2.5), then (2.13) and (2.14) imply that

$$0 \leq J(t, \mu, \alpha^*; x, \alpha^*) - v(\mu^{\alpha^*}; t, x) \leq \varepsilon, \quad \|\varphi - v(\mu^{\alpha^*}; t, \cdot)\|_\infty \leq 2\varepsilon. \quad (2.15)$$

So we may alternatively define $\mathbb{V}_{state}^\varepsilon(t, \mu)$ by using $\|\varphi - v(\mu^{\alpha^*}; t, \cdot)\|_\infty \leq \varepsilon$.

Remark 2.6 (i) In the case that there is only one player, namely q, F, G do not depend on μ , $\mathbb{P}^{\mu^*}; t, x, \alpha = \mathbb{P}^{t, x, \alpha}$ does not depend on μ and α^* . Let

$$V(t, x) := \inf_{\alpha \in \mathcal{A}_{state}} \mathbb{E}^{\mathbb{P}^{t, x, \alpha}} \left[G(X_T) + \sum_{s=t}^{T-1} F(s, X_s, \alpha(s, X_s)) \right]$$

denote the value function of the standard stochastic control problem. One can easily see that, when there exists an optimal control α^* , $\mathbb{V}_0(t, \mu) = \mathbb{V}_{state}(t, \mu) = \{V(t, \cdot)\}$. However, when there is no optimal control, we still have $\mathbb{V}_{state}(t, \mu) = \{V(t, \cdot)\}$ but $\mathbb{V}_0(t, \mu) = \emptyset$. So \mathbb{V}_{state} , not \mathbb{V}_0 , is the natural extension of the value function V .

(ii) We remark that $\bigcap_{\varepsilon > 0} \mathcal{M}_{state}^\varepsilon(t, \mu) = \mathcal{M}_{state}(t, \mu)$, however, in general it is possible that $\mathbb{V}_{state}(t, \mu)$ is strictly larger than $\mathbb{V}_0(t, \mu)$. Indeed, $\mathbb{V}_{state}(t, \mu)$ can be even larger than the closure of $\mathbb{V}_0(t, \mu)$, where the latter is still empty when there is no optimal control.

Similarly, given T_0 and ψ , $\mathcal{M}_{state}^\varepsilon(T_0, \psi; t, \mu)$ denotes the set of $\alpha^* \in \mathcal{A}_{state}$ such that

$$J(T_0, \psi; t, \mu, \alpha^*; x, \alpha^*) \leq \inf_{\alpha \in \mathcal{A}_{state}} J(T_0, \psi; t, \mu, \alpha^*; x, \alpha) + \varepsilon, \quad \forall x \in \mathbb{S}. \quad (2.16)$$

The DPP remains true for \mathbb{V}_{state} after appropriate modifications as follows.

Theorem 2.7 Under Assumption 2.2 (i), for any $0 \leq t < T_0 \leq T$ and $\mu \in \mathcal{P}_0(\mathbb{S})$, we have

$$\begin{aligned} \mathbb{V}_{state}(t, \mu) &:= \bigcap_{\varepsilon > 0} \left\{ \varphi \in \mathbb{L}^0(\mathbb{S}; \mathbb{R}) : \|\varphi - J(T_0, \psi; t, \mu, \alpha^*; \cdot, \alpha^*)\|_\infty \leq \varepsilon \right. \\ &\quad \text{for some } \psi \in \mathbb{L}^0(\mathbb{S} \times \mathcal{P}_0(\mathbb{S}); \mathbb{R}) \text{ and } \alpha^* \in \mathcal{A}_{state} \text{ such that} \\ &\quad \left. \psi(\cdot, \mu_{T_0}^{\alpha^*}) \in \mathbb{V}_{state}^\varepsilon(T_0, \mu_{T_0}^{\alpha^*}), \alpha^* \in \mathcal{M}_{state}^\varepsilon(T_0, \psi; t, \mu) \right\}. \end{aligned} \quad (2.17)$$

This theorem can be proved by modifying the arguments in Theorem 2.4 and Proposition 2.3. However, since the proof is very similar to that of Theorem 4.2 below, except that the latter is in the more complicated path dependent setting, we thus postpone it to Appendix.

3 The N -player game with homogeneous equilibria

In this section we study the N -player game whose set value will converge to \mathbb{V}_{state} .

3.1 The N -player game

Set $\Omega^N := \mathbb{X}^N$ with canonical processes $\vec{X} = (X^1, \dots, X^N)$, where X^i stands for the state process of Player i . The empirical measure of \vec{X} is denoted as: with the Dirac measure δ ,

$$\mu_t^N := \mu_{\vec{X}_t}^N \quad \text{where} \quad \mu_{\vec{x}}^N := \frac{1}{N} \sum_{i=1}^N \delta_{x_i} \in \mathcal{P}(\mathbb{S}), \quad \text{for } \vec{x} = (x_1, \dots, x_N) \in \mathbb{S}^N. \quad (3.1)$$

The player i will have control α^i . In the literature, a closed loop control α^i may depend on the full information \vec{X} . However, since we are talking about large N , in practice it may not be feasible for each player to observe all other players' states individually. Moreover, in the MFG setting the population state is characterized by its distribution, not by each player's individual state. So in this section we consider only symmetric controls, namely α^i depends on his/her own state X^i and on the others through the empirical measure μ^N .

For technical reasons, we introduce another parameter $L \geq 0$. Denote

$$\mathcal{A}_{state}^L := \left\{ \alpha : \mathbb{T} \times \mathbb{S} \times \mathcal{P}(\mathbb{S}) \rightarrow \mathbb{A} : |\alpha(t, x, \mu) - \alpha(t, x, \nu)| \leq LW_1(\mu, \nu), \forall t, x, \mu, \nu \right\}, \quad (3.2)$$

and $\mathcal{A}_{state}^\infty := \bigcup_{L \geq 0} \mathcal{A}_{state}^L$. Given $t \in \mathbb{T}$, $\vec{x} \in \mathbb{S}^N$, and $\vec{\alpha} = (\alpha^1, \dots, \alpha^N) \in (\mathcal{A}_{state}^\infty)^N$, let $\mathbb{P}^{t, \vec{x}, \vec{\alpha}}$ denote the probability measure on $\mathcal{F}_T^{\vec{X}}$ determined recursively by: for $s = t, \dots, T-1$,

$$\mathbb{P}^{t, \vec{x}, \vec{\alpha}}(\vec{X}_t = \vec{x}) = 1, \quad \mathbb{P}^{t, \vec{x}, \vec{\alpha}}(\vec{X}_{s+1} = \vec{x}' | \vec{X}_s = \vec{x}') = \prod_{i=1}^N q(s, x'_i, \mu_s^N, \alpha^i(s, x'_i, \mu_s^N); x''_i), \quad (3.3)$$

and the cost function of Player i is:

$$J_i(t, \vec{x}, \vec{\alpha}) := \mathbb{E}^{\mathbb{P}^{t, \vec{x}, \vec{\alpha}}} \left[G(X_T^i, \mu_T^N) + \sum_{s=t}^{T-1} F(s, X_s^i, \mu_s^N, \alpha^i(s, X_s^i, \mu_s^N)) \right]. \quad (3.4)$$

Remark 3.1 (i) It is obvious that $\mathcal{A}_{state}^0 = \mathcal{A}_{state}$ for the \mathcal{A}_{state} in the previous subsection. For the MFG, there is no need to consider $\mathcal{A}_{state}^\infty$. Indeed, given $(t, \mu) \in \mathbb{T} \times \mathcal{P}_0(\mathbb{S})$, for any $\alpha \in \mathcal{A}_{state}^\infty$, let $\mathbb{P}^{t, \mu, \alpha}$ be defined as in (2.3): again denoting $\mu_s^\alpha := \mathbb{P}^{t, \mu, \alpha} \circ X_s^{-1}$,

$$\mathbb{P}^{t, \mu, \alpha} \circ X_t^{-1} = \mu, \quad \mathbb{P}^{t, \mu, \alpha}(X_{s+1} = \tilde{x} | X_s = x) = q(s, x, \mu_s^\alpha, \alpha(s, x, \mu_s^\alpha); \tilde{x}).$$

Introduce $\tilde{\alpha}(s, x) := \alpha(s, x, \mu_s^\alpha)$. Then $\tilde{\alpha} \in \mathcal{A}_{state}$ and one can easily verify that $\mu^{\tilde{\alpha}} = \mu^\alpha$. In particular, the set value $\mathbb{V}_{state}(t, \mu)$ will remain the same by allowing $\alpha \in \mathcal{A}_{state}^\infty$. For the N -player game, however, since μ^N is random, the dependence on μ^N makes the difference.

(ii) In the literature one typically uses $\mu_t^{N, -i} := \frac{1}{N-1} \sum_{j \neq i} \delta_{X_t^j}$, rather than μ_t^N , in (3.3) and (3.4). The convergence results in this section will remain true if we use $\mu^{N, -i}$ instead. However, we find it more convenient to use μ_t^N .

There is another crucial issue concerning the equilibria. Note that an MFE requires by definition that each player takes the same control α^* . To achieve the desired convergence, for the N -player game it is natural to consider only the homogeneous equilibria: $\alpha_1 = \dots = \alpha_N$, which we will do in the rest of this section. We note that, for a homogeneous control α , the $\mathbb{P}^{t, \vec{x}, \alpha} := \mathbb{P}^{t, \vec{x}, (\alpha, \dots, \alpha)}$ in (3.3) and $J_i(t, \vec{x}, \alpha) := J_i(t, \vec{x}, (\alpha, \dots, \alpha))$ in (3.4) are also symmetric in \vec{x} , or say invariant in terms of its empirical measure:

$$\mathbb{P}^{t, \vec{x}, \alpha} = \mathbb{P}^{t, \mu_{\vec{x}}^N, \alpha}, \quad J_i(t, \vec{x}, \alpha) = J^N(t, x_i, \mu_{\vec{x}}^N, \alpha). \quad (3.5)$$

Definition 3.2 For any $\varepsilon > 0, L \geq 0$, we say $\alpha^* \in \mathcal{A}_{state}^L$ is a homogeneous state dependent (ε, L) -equilibrium of the N -player game at (t, \vec{x}) , denoted as $\alpha^* \in \mathcal{M}_{state}^{N, \varepsilon, L}(t, \vec{x})$, if:

$$J_i(t, \vec{x}, \alpha^*) \leq v_i^{N, L}(t, \vec{x}, \alpha^*) := \inf_{\tilde{\alpha} \in \mathcal{A}_{state}^L} J_i(t, \vec{x}, (\alpha^*, \tilde{\alpha})_i) + \varepsilon, \quad i = 1, \dots, N, \quad (3.6)$$

where $(\alpha, \tilde{\alpha})_i$ denote the vector $\tilde{\alpha}$ such that $\alpha^i = \tilde{\alpha}$ and $\alpha^j = \alpha$ for all $j \neq i$.

In light of (3.5), clearly $\mathcal{M}_{state}^{N, \varepsilon, L}(t, \vec{x})$ is law invariant: $\mathcal{M}_{state}^{N, \varepsilon, L}(t, \vec{x}) = \mathcal{M}_{state}^{N, \varepsilon, L}(t, \vec{x}')$ whenever $\mu_{\vec{x}}^N = \mu_{\vec{x}'}^N$. Thus, by abusing the notation, we may denote $\mathcal{M}_{state}^{N, \varepsilon, L}(t, \vec{x}) = \mathcal{M}_{state}^{N, \varepsilon, L}(t, \mu_{\vec{x}}^N)$ and call such α^* a homogeneous state dependent (ε, L) -equilibrium at $(t, \mu_{\vec{x}}^N)$.

Note again that $q > 0$, then similar to Subsection 2.1, for convenience in this section we restrict to only those \vec{x} such that $\mu_{\vec{x}}^N$ has full support, and we denote

$$\mathbb{S}_0^N := \{\vec{x} \in \mathbb{S}^N : \mu_{\vec{x}}^N \in \mathcal{P}_0(\mathbb{S})\}, \quad \mathcal{P}_N(\mathbb{S}) := \{\mu_{\vec{x}}^N : \vec{x} \in \mathbb{S}_0^N\} \subset \mathcal{P}_0(\mathbb{S}). \quad (3.7)$$

We now define the set value of the homogeneous N -player game: recalling (3.5),

$$\begin{aligned} \mathbb{V}_{state}^N(t, \mu) &:= \bigcap_{\varepsilon > 0} \mathbb{V}_{state}^{N, \varepsilon}(t, \mu) := \bigcap_{\varepsilon > 0} \bigcup_{L \geq 0} \mathbb{V}_{state}^{N, \varepsilon, L}(t, \mu), \quad \forall (t, \mu) \in \mathbb{T} \times \mathcal{P}_N(\mathbb{S}), \quad \text{where} \\ \mathbb{V}_{state}^{N, \varepsilon, L}(t, \mu) &:= \left\{ \varphi \in \mathbb{L}^0(\mathbb{S}; \mathbb{R}) : \exists \alpha^* \in \mathcal{M}_{state}^{N, \varepsilon, L}(t, \mu) \text{ s.t. } \|\varphi - J^N(t, \cdot, \mu, \alpha^*)\|_\infty \leq \varepsilon \right\}. \end{aligned} \quad (3.8)$$

3.2 Convergence of the empirical measures

Theorem 3.3 Let Assumption 2.2 (ii) hold. Then, for any $L \geq 0$, there exists a constant C_L , which depends only on T, d, L_q , and L such that, for any $t \in \mathbb{T}$, $\vec{x} \in \mathbb{S}_0^N$, $\mu \in \mathcal{P}_0(\mathbb{S})$, $\alpha, \tilde{\alpha} \in \mathcal{A}_{state}^L$, and $s \geq t$, $i = 1, \dots, N$,

$$\mathbb{E}^{\mathbb{P}^{t, \vec{x}, (\alpha, \tilde{\alpha})_i}} [\mathcal{W}_1(\mu_s^N, \mu_s^\alpha)] \leq C_L \theta_N, \quad \text{where } \theta_N := W_1(\mu_{\vec{x}}^N, \mu) + \frac{1}{\sqrt{N}}; \quad (3.9)$$

$$\mathcal{W}_1\left(\mathbb{P}^{t, \vec{x}, (\alpha, \tilde{\alpha})_i} \circ (X_s^i)^{-1}, \mathbb{P}^{\mu^\alpha; t, x_i, \tilde{\alpha}} \circ X_s^{-1}\right) \leq C_L \theta_N. \quad (3.10)$$

Proof We first recall Remark 3.1 and extend all the notations in Subsection 2.1 to those $\alpha \in \mathcal{A}_{state}^L$ in the obvious sense. Fix t, i and denote $\mathbb{P}^N := \mathbb{P}^{t, \vec{x}, (\alpha, \tilde{\alpha})_i}$.

Step 1. We first prove (3.9) for $s = t + 1$. Note that $X_{t+1}^1, \dots, X_{t+1}^N$ are independent

under \mathbb{P}^N . By (2.1), we have

$$\begin{aligned}
\mathbb{E}^{\mathbb{P}^N} [W_1(\mu_{t+1}^N, \mu_{t+1}^\alpha)] &= \sum_{\tilde{x} \in \mathbb{S}} \mathbb{E}^{\mathbb{P}^N} [|\mu_{t+1}^N(\tilde{x}) - \mu_{t+1}^\alpha(\tilde{x})|] \\
&\leq \sum_{\tilde{x} \in \mathbb{S}} \left(\mathbb{E}^{\mathbb{P}^N} [|\mu_{t+1}^N(\tilde{x}) - \mu_{t+1}^\alpha(\tilde{x})|^2] \right)^{\frac{1}{2}} \\
&= \sum_{\tilde{x} \in \mathbb{S}} \left[\text{Var}^{\mathbb{P}^N} [\mu_{t+1}^N(\tilde{x})] + \left(\mathbb{E}^{\mathbb{P}^N} [\mu_{t+1}^N(\tilde{x}) - \mu_{t+1}^\alpha(\tilde{x})] \right)^2 \right]^{\frac{1}{2}} \tag{3.11} \\
&= \sum_{\tilde{x} \in \mathbb{S}} \left[\frac{1}{N^2} \sum_{j=1}^N \text{Var}^{\mathbb{P}^N} [\mathbf{1}_{\{X_{t+1}^j = \tilde{x}\}}] + \left(\frac{1}{N} \sum_{j=1}^N \mathbb{P}^N(X_{t+1}^j = \tilde{x}) - \mu_{t+1}^\alpha(\tilde{x}) \right)^2 \right]^{\frac{1}{2}} \\
&\leq \frac{C}{\sqrt{N}} + \sum_{\tilde{x} \in \mathbb{S}} \left| \frac{1}{N} \sum_{j=1}^N \mathbb{P}^N(X_{t+1}^j = \tilde{x}) - \mu_{t+1}^\alpha(\tilde{x}) \right|.
\end{aligned}$$

Note that, by the desired Lipschitz continuity of q in μ and that $|\mathbb{S}| = d$ is finite,

$$\begin{aligned}
&\left| \frac{1}{N} \sum_{j=1}^N \mathbb{P}^N(X_{t+1}^j = \tilde{x}) - \mu_{t+1}^\alpha(\tilde{x}) \right| \\
&= \left| \frac{1}{N} \sum_{x \in \mathbb{S}} \left[\sum_{j \neq i} q(t, x, \mu_x^N, \alpha(t, x, \mu_x^N); \tilde{x}) \mathbf{1}_{\{x_j = x\}} + q(t, x, \mu_x^N, \tilde{\alpha}(t, x, \mu_x^N); \tilde{x}) \mathbf{1}_{\{x_i = x\}} \right] \right. \\
&\quad \left. - \sum_{x \in \mathbb{S}} q(t, x, \mu, \alpha(t, x, \mu); \tilde{x}) \mu(x) \right| \\
&\leq \left| \frac{1}{N} \sum_{x \in \mathbb{S}} \sum_{j=1}^N q(t, x, \mu_x^N, \alpha(t, x, \mu_x^N); \tilde{x}) \mathbf{1}_{\{x_j = x\}} - \sum_{x \in \mathbb{S}} q(t, x, \mu, \alpha(t, x, \mu); \tilde{x}) \mu(x) \right| \\
&\quad + \frac{1}{N} \sum_{x \in \mathbb{S}} |q(t, x, \mu_x^N, \alpha(t, x, \mu_x^N); \tilde{x}) - q(t, x, \mu_x^N, \tilde{\alpha}(t, x, \mu_x^N); \tilde{x})| \mathbf{1}_{\{x_i = x\}} \\
&\leq \left| \sum_{x \in \mathbb{S}} q(t, x, \mu_x^N, \alpha(t, x, \mu_x^N); \tilde{x}) \mu_x^N(x) - \sum_{x \in \mathbb{S}} q(t, x, \mu, \alpha(t, x, \mu); \tilde{x}) \mu(x) \right| + \frac{1}{N} \\
&\leq \sum_{x \in \mathbb{S}} \left[|\mu_x^N(x) - \mu(x)| + C_L W_1(\mu_x^N, \mu) \mu(x) \right] + \frac{1}{N} \leq C_L \theta_N.
\end{aligned}$$

Then, $\mathbb{E}^{\mathbb{P}^N} [W_1(\mu_{t+1}^N, \mu_{t+1}^\alpha)] \leq \frac{C}{\sqrt{N}} + C_L \theta_N \leq C_L \theta_N$.

Step 2. We next prove (3.9) by induction. For any $s = t, \dots, T-1$, by Step 1 we have

$$\mathbb{E}^{\mathbb{P}^N} [W_1(\mu_{s+1}^N, \mu_{s+1}^\alpha) | \mathcal{F}_s^{\tilde{X}}] \leq C_L \left[W_1(\mu_s^N, \mu_s^\alpha) + \frac{1}{\sqrt{N}} \right], \quad \mathbb{P}^N\text{-a.s.}$$

Then

$$\mathbb{E}^{\mathbb{P}^N} [W_1(\mu_{s+1}^N, \mu_{s+1}^\alpha)] = \mathbb{E}^{\mathbb{P}^N} \left[\mathbb{E}^{\mathbb{P}^N} [W_1(\mu_{s+1}^N, \mu_{s+1}^\alpha) | \tilde{X}_s^N] \right] \leq C_L \mathbb{E}^{\mathbb{P}^N} [W_1(\mu_s^N, \mu_s^\alpha)] + \frac{C_L}{\sqrt{N}}.$$

Since T is finite, by induction we obtain (3.9) immediately.

Step 3. We now prove (3.10). Denote

$$\kappa_s := W_1\left(\mathbb{P}^N \circ (X_s^i)^{-1}, \mathbb{P}^i \circ X_s^{-1}\right) \quad \text{where} \quad \mathbb{P}^i := \mathbb{P}^{\mu^\alpha; t, x_i, \tilde{\alpha}}.$$

Then $\kappa_t = 0$, and for $s = t, \dots, T-1$,

$$\begin{aligned} \kappa_{s+1} &= \sum_{\tilde{x} \in \mathbb{S}} \left| \mathbb{P}^N(X_{s+1}^i = \tilde{x}) - \mathbb{P}^i(X_{s+1} = \tilde{x}) \right| \\ &= \sum_{\tilde{x} \in \mathbb{S}} \left| \mathbb{E}^{\mathbb{P}^N} [q(s, X_s^i, \mu_s^N, \tilde{\alpha}(s, X_s^i, \mu_s^N); \tilde{x})] - \mathbb{E}^{\mathbb{P}^i} [q(s, X_s, \mu_s^\alpha, \tilde{\alpha}(s, X_s, \mu_s^\alpha); \tilde{x})] \right| \\ &\leq \sum_{\tilde{x} \in \mathbb{S}} \left| \mathbb{E}^{\mathbb{P}^N} [q(s, X_s^i, \mu_s^N, \tilde{\alpha}(s, X_s^i, \mu_s^N); \tilde{x})] - \mathbb{E}^{\mathbb{P}^N} [q(s, X_s^i, \mu_s^\alpha, \tilde{\alpha}(s, X_s^i, \mu_s^\alpha); \tilde{x})] \right| \\ &\quad + \sum_{\tilde{x} \in \mathbb{S}} \left| \mathbb{E}^{\mathbb{P}^N} [q(s, X_s^i, \mu_s^\alpha, \tilde{\alpha}(s, X_s^i, \mu_s^\alpha); \tilde{x})] - \mathbb{E}^{\mathbb{P}^i} [q(s, X_s, \mu_s^\alpha, \tilde{\alpha}(s, X_s, \mu_s^\alpha); \tilde{x})] \right| \\ &\leq C_L \mathbb{E}^{\mathbb{P}^N} [W_1(\mu_s^N, \mu_s^\alpha)] + \sum_{x, \tilde{x} \in \mathbb{S}} q(s, x, \mu_s^\alpha, \tilde{\alpha}(s, x, \mu_s^\alpha); \tilde{x}) |\mathbb{P}^N(X_s^i = x) - \mathbb{P}^i(X_s = x)| \\ &\leq C_L \theta_N + \kappa_s, \end{aligned}$$

where the last inequality thanks to (3.9). Now by induction one can easily prove (3.10). \blacksquare

3.3 Convergence of the set values

We first study the convergence of the cost functions. Recall the θ_N in (3.9) and the functions v in (2.5) and $v_i^{N,L}$ in (3.6).

Theorem 3.4 *Let Assumption 2.2 (ii) and (iii) hold. For any $L \geq 0$, there exists a modulus of continuity function ρ_L , which depends only on T, d, L_q, C_0, ρ , and L such that, for any $t \in \mathbb{T}$, $\mu_{\vec{x}}^N \in \mathcal{P}_N(\mathbb{S})$, $\mu \in \mathcal{P}_0(\mathbb{S})$, and any $\alpha, \tilde{\alpha} \in \mathcal{A}_{state}^L$, $i = 1, \dots, N$,*

$$\left| J_i(t, \vec{x}, (\alpha, \tilde{\alpha})_i) - J(t, \mu, \alpha; x_i, \tilde{\alpha}) \right| + \left| v_i^{N,L}(t, \vec{x}, \alpha) - v(\mu^\alpha; t, x_i) \right| \leq \rho_L(\theta_N). \quad (3.12)$$

Proof Clearly the uniform estimates for J implies that for v , so we shall only prove the former one. Recall (3.4), (2.5), and the notations $\mathbb{P}^N, \mathbb{P}^i$ in the proof of Theorem 3.3. Then

$$\begin{aligned} \left| J_i(t, \vec{x}, (\alpha, \tilde{\alpha})_i) - J(t, \mu, \alpha; x_i, \tilde{\alpha}) \right| &\leq I_T + \sum_{s=t}^{T-1} I_s, \quad \text{where} \\ I_T &:= \left| \mathbb{E}^{\mathbb{P}^N} [G(X_T^i, \mu_T^N)] - \mathbb{E}^{\mathbb{P}^i} [G(X_T, \mu_T^\alpha)] \right|; \\ I_s &:= \left| \mathbb{E}^{\mathbb{P}^N} [F(s, X_s^i, \mu_s^N, \tilde{\alpha}(s, X_s^i, \mu_s^N))] - \mathbb{E}^{\mathbb{P}^i} [F(s, X_s, \mu_s^\alpha, \tilde{\alpha}(s, X_s, \mu_s^\alpha))] \right|, \quad s < T. \end{aligned}$$

Note that, for $s < T$, by (3.10),

$$\begin{aligned}
I_s &\leq \left| \mathbb{E}^{\mathbb{P}^N} [F(s, X_s^i, \mu_s^N, \tilde{\alpha}(s, X_s^i, \mu_s^N))] - \mathbb{E}^{\mathbb{P}^N} [F(s, X_s^i, \mu_s^\alpha, \tilde{\alpha}(s, X_s^i, \mu_s^\alpha))] \right| \\
&\quad + \left| \mathbb{E}^{\mathbb{P}^N} [F(s, X_s^i, \mu_s^\alpha, \tilde{\alpha}(s, X_s^i, \mu_s^\alpha))] - \mathbb{E}^{\mathbb{P}^i} [F(s, X_s, \mu_s^\alpha, \tilde{\alpha}(s, X_s, \mu_s^\alpha))] \right| \\
&\leq \mathbb{E}^{\mathbb{P}^N} [\rho(C_L W_1(\mu_s^N, \mu_s^\alpha))] + \sum_{x \in \mathbb{S}} |F(s, x, \mu_s^\alpha, \tilde{\alpha}(s, x, \mu_s^\alpha))| |\mathbb{P}^N(X_s^i = x) - \mathbb{P}^i(X_s = x)| \\
&\leq \mathbb{E}^{\mathbb{P}^N} [\rho(C_L W_1(\mu_s^N, \mu_s^\alpha))] + C_L \theta_N.
\end{aligned}$$

Similarly we have the estimate for I_T , and thus

$$\left| J_i(t, \vec{x}, (\alpha, \tilde{\alpha})_i) - J(t, \mu, \alpha; x_i, \tilde{\alpha}) \right| \leq \sum_{s=t}^T \mathbb{E}^{\mathbb{P}^N} [\rho(C_L W_1(\mu_s^N, \mu_s^\alpha))] + C_L \theta_N.$$

This, together with (3.9), implies (3.12) for some appropriately defined modulus of continuity function ρ_L . \blacksquare

Our main result of this section is the following convergence of the set values. Recall, for a sequence of sets $\{E_N\}_{N \geq 1}$, $\overline{\lim}_{N \rightarrow \infty} E_N := \bigcap_{n \geq 1} \bigcup_{N \geq n} E_N$, $\underline{\lim}_{N \rightarrow \infty} E_N := \bigcup_{n \geq 1} \bigcap_{N \geq n} E_N$.

Theorem 3.5 *Let Assumption 2.2 (ii), (iii) hold and $\mu_{\vec{x}}^N \in \mathcal{P}_N(\mathbb{S}) \rightarrow \mu \in \mathcal{P}_0(\mathbb{S})$. Then*

$$\bigcap_{\varepsilon > 0} \bigcup_{L \geq 0} \overline{\lim}_{N \rightarrow \infty} \mathbb{V}_{state}^{N, \varepsilon, L}(t, \mu_{\vec{x}}^N) \subset \mathbb{V}_{state}(t, \mu) \subset \bigcap_{\varepsilon > 0} \underline{\lim}_{N \rightarrow \infty} \mathbb{V}_{state}^{N, \varepsilon, 0}(t, \mu_{\vec{x}}^N) \quad (3.13)$$

In particular, since $\underline{\lim}_{N \rightarrow \infty} \mathbb{V}_{state}^{N, \varepsilon, 0}(t, \mu_{\vec{x}}^N) \subset \bigcup_{L \geq 0} \overline{\lim}_{N \rightarrow \infty} \mathbb{V}_{state}^{N, \varepsilon, L}(t, \mu_{\vec{x}}^N)$, actually equalities hold.

Note that $\vec{x} \in \mathbb{S}_0^N$ obviously depends on N , so more rigorously we should write \vec{x}^N in the above statements. For notational simplicity we omit this N here. We also remark that at above we are not able to switch the order of $\overline{\lim}_{N \rightarrow \infty}$ and $\bigcap_{\varepsilon > 0} \bigcup_{L \geq 0}$ in the left side, or the order of $\underline{\lim}_{N \rightarrow \infty}$ and $\bigcap_{\varepsilon > 0}$ in the right side.

Proof (i) We first prove the right inclusion in (3.13). Fix $\varphi \in \mathbb{V}_{state}(t, \mu)$, $\varepsilon > 0$, and set $\varepsilon_1 := \frac{\varepsilon}{2}$. Note that $\mathcal{A}_{state} = \mathcal{A}_{state}^0$. By (2.14), there exists $\alpha^* \in \mathcal{M}_{state}^{\varepsilon_1}(t, \mu)$ such that $\|\varphi - J(t, \mu, \alpha^*; \cdot, \alpha^*)\|_\infty \leq \varepsilon_1$. Recall (2.13), we have

$$J(t, \mu, \alpha^*; x, \alpha^*) \leq v(\mu^{\alpha^*}; t, x) + \varepsilon_1, \quad \text{for all } x \in \mathbb{S}.$$

For any $\alpha \in \mathcal{A}_{state}^0 = \mathcal{A}_{state}$, by Theorem 3.4 we have

$$\begin{aligned}
J_i(t, \vec{x}, \alpha^*) &\leq J(t, \mu, \alpha^*; x_i, \alpha^*) + \rho_0(\theta_N) \\
&\leq v(\mu^{\alpha^*}; t, x) + \varepsilon_1 + \rho_0(\theta_N) \leq v_i^{N, L}(t, \vec{x}, \alpha^*) + \varepsilon_1 + 2\rho_0(\theta_N).
\end{aligned}$$

Choose N large enough such that $\rho_0(\theta_N) \leq \frac{\varepsilon}{4}$, then $J_i(t, \vec{x}, \alpha^*) \leq v_i^{N,L}(t, \vec{x}, \alpha^*) + \varepsilon$. This implies that $\alpha^* \in \mathcal{M}_{\varepsilon,0}^N(t, \mu_{\vec{x}}^N)$. Moreover,

$$\begin{aligned} \|\varphi - J^N(t, \cdot, \mu_{\vec{x}}^N, \alpha^*)\|_\infty &\leq \varepsilon_1 + \sup_i \left| J_i(t, \vec{x}, \alpha^*) - J(t, \mu, \alpha^*; x_i, \alpha^*) \right| \\ &\leq \varepsilon_1 + \rho_0(\theta_N) \leq \varepsilon_1 + \frac{\varepsilon}{4} \leq \varepsilon. \end{aligned}$$

Then $\varphi \in \mathbb{V}_{state}^{N,\varepsilon,0}(t, \mu_{\vec{x}}^N)$ for all N large enough. That is, $\varphi \in \underline{\lim}_{N \rightarrow \infty} \mathbb{V}_{state}^{N,\varepsilon,0}(t, \mu_{\vec{x}}^N)$. Since $\varphi \in \mathbb{V}_{state}(t, \mu)$ and $\varepsilon > 0$ are arbitrary, we obtain the right inclusion in (3.13).

(ii) We next show the left inclusion in (3.13). Fix $\varphi \in \bigcap_{\varepsilon > 0} \bigcup_{L \geq 0} \overline{\lim}_{N \rightarrow \infty} \mathbb{V}_{state}^{N,\varepsilon,L}(t, \mu_{\vec{x}}^N)$ and $\varepsilon > 0$. Then, for $\varepsilon_1 := \frac{\varepsilon}{2} > 0$, there exist $L_\varepsilon > 0$ and an infinite sequence $\{N_k\}_{k \geq 1}$ such that $\varphi \in \mathbb{V}_{state}^{N_k, \varepsilon_1, L_\varepsilon}(t, \mu_{\vec{x}}^{N_k})$ for all $k \geq 1$. Recall (3.8), for each $k \geq 1$ there exists $\alpha^k \in \mathcal{M}_{state}^{N_k, \varepsilon_1, L_\varepsilon}(t, \mu_{\vec{x}}^{N_k})$ such that $\|\varphi - J^N(t, \cdot, \mu_{\vec{x}}^{N_k}, \alpha^k)\|_\infty \leq \varepsilon_1$. By Definition 3.2, we have $J_i(t, \vec{x}, \alpha^k) \leq v_i^{N_k, L_\varepsilon}(t, \vec{x}, \alpha^k) + \varepsilon_1$. Similar to (i), by Theorem 3.4 we have

$$J(t, \mu, \alpha^k; x_i, \alpha^k) \leq v(\mu^{\alpha^k}; t, x_i) + \varepsilon_1 + 2\rho_{L_\varepsilon}(\theta_{N_k}) \leq v(\mu^{\alpha^k}; t, x_i) + \varepsilon,$$

for k large enough. That is, $\alpha^k \in \mathcal{M}_{state}^\varepsilon(t, \mu)$. Similar to (i) again, for k large enough we have $\|\varphi - J(t, \mu, \alpha^k; \cdot, \alpha^k)\|_\infty \leq \varepsilon$. Then $\varphi \in \mathbb{V}_{state}^\varepsilon(t, \mu)$. Since $\varepsilon > 0$ is arbitrary, we obtain $\varphi \in \mathbb{V}_{state}(t, \mu)$, and hence derive the left inclusion in (3.13). \blacksquare

Remark 3.6 (i) From Theorem 3.5 (i) we see that, for any $\alpha^* \in \mathcal{M}_{state}^{\frac{\varepsilon}{2}}(t, \mu)$, we have $\alpha^* \in \mathcal{M}_{state}^{N,\varepsilon,0}(t, \mu_{\vec{x}}^N)$ when N is large enough. Moreover, by (3.9) we have the desired estimate for the approximate equilibrium measure $\mathbb{E}^{\mathbb{P}^{t, \vec{x}, \alpha^*}}[W_1(\mu_s^N, \mu_s^{\alpha^*})] \leq C_L \theta_N$. This verifies the standard result in the literature that an approximate MFE is an approximate equilibrium of the N -player game.

(ii) From Theorem 3.5 (ii) we see that, for any $\alpha^k \in \mathcal{M}_{state}^{N_k, \frac{\varepsilon}{2}, L_\varepsilon}(t, \mu_{\vec{x}}^{N_k})$, we have $\alpha^k \in \mathcal{M}_{state}^\varepsilon(t, \mu)$ when k is large enough, and we again have the estimate for the approximate equilibrium measure $\mathbb{E}^{\mathbb{P}^{t, \vec{x}, \alpha^k}}[W_1(\mu_s^{N_k}, \mu_s^{\alpha^k})] \leq C_L \theta_{N_k}$. This is in the spirit that any limit point of the N -player equilibrium measures is an MFE measure.

(iii) As we saw above, we have the desired convergence both for set values and for the equilibrium measures. This type of convergence holds true for open loop controls, see [30], and for closed loop controls under monotonicity conditions (where the MFE is unique), see e.g. [10]. However, for general MFG with closed loop controls, to our best knowledge our result is the first one which has the convergence.

Remark 3.7 (i) We should point out that the key to obtain the convergence here is to consider homogeneous equilibria for the N -player games. If we use heterogeneous equilibria

for the N -player games, it turns out that we will have the desired convergence when we consider relaxed controls for the MFG, as we will do in the next two sections.

(ii) Another technical trick we are using is the uniform Lipschitz continuity requirement on the admissible controls. The convergence analysis will become more subtle when we remove such regularity requirement, see e.g. [31].

(iii) We would also like to point out that in general it could be easier to analyze the set values than the equilibria, especially when we do not require the regularities of the controls. One trivial example is that F and G are constants, then the set value is a singleton of constant value, while the equilibria are arbitrary admissible controls.

4 Mean field games on finite space with relaxed controls

In this section we study MFG with relaxed controls, or say mixed strategies. Besides its independent interest, our main motivation is to characterize the limit of N -player games with heterogeneous equilibria. We shall still consider the finite space in Section 2, however, for the purpose of generality in this section we consider path dependent setting.

4.1 The relaxed set value with path dependent controls

We start with some notations for the path dependent setting. For $\mathbf{x} = (\mathbf{x}_t)_{0 \leq t \leq T} \in \mathbb{X}$, denote by $\mathbf{x}_{t\wedge\cdot} = (\mathbf{x}_0, \dots, \mathbf{x}_t, \mathbf{x}_t, \dots, \mathbf{x}_T)$ the path stopping at t and $\mathbb{X}_t := \{\mathbf{x}_{t\wedge\cdot} : \mathbf{x} \in \mathbb{X}\} \subset \mathbb{X}$. For $\mathbf{x}, \tilde{\mathbf{x}} \in \mathbb{X}$, we say $\mathbf{x} =_t \tilde{\mathbf{x}}$ if $\mathbf{x}_{t\wedge\cdot} = \tilde{\mathbf{x}}_{t\wedge\cdot}$. Denote $\mathbb{X}^{t,\mathbf{x}} := \{\tilde{\mathbf{x}} \in \mathbb{X} : \tilde{\mathbf{x}} =_t \mathbf{x}\}$ and $\mathbb{X}_s^{t,\mathbf{x}} := \mathbb{X}^{t,\mathbf{x}} \cap \mathbb{X}_s$, for $s \geq t$. Introduce the concatenation $\mathbf{x} \oplus_t \tilde{\mathbf{x}} \in \mathbb{X}$ by

$$(\mathbf{x} \oplus_t \tilde{\mathbf{x}})_s := \mathbf{x}_s \mathbf{1}_{\{s \leq t\}} + \tilde{\mathbf{x}}_s \mathbf{1}_{\{s > t\}}, \quad \text{and} \quad (\mathbf{x} \oplus_t x)_s := \mathbf{x}_s \mathbf{1}_{\{s \leq t\}} + x \mathbf{1}_{\{s > t\}}, \quad x \in \mathbb{S}.$$

For each $t \in \mathbb{T}$, let $\mathcal{P}(\mathbb{X}_t)$ denote the set of probability measures on $(\Omega, \mathcal{F}_t^X)$, equipped with

$$W_1(\mu, \nu) := \sum_{\mathbf{x} \in \mathbb{X}_t} |\mu(\mathbf{x}) - \nu(\mathbf{x})|, \quad \forall \mu, \nu \in \mathcal{P}(\mathbb{X}_t),$$

and $\mathcal{P}_0(\mathbb{X}_t)$ the subset of $\mu \in \mathcal{P}(\mathbb{X}_t)$ with full support \mathbb{X}_t . Again this is just for convenience of presentation. For a measure $\mu \in \mathcal{P}(\mathbb{X}) = \mathcal{P}(\mathbb{X}_T)$, denote $\mu_{t\wedge\cdot} := \mu \circ X_{t\wedge\cdot}^{-1} \in \mathcal{P}(\mathbb{X}_t)$. We remark that, by abusing the notation μ , here $\mu_{t\wedge\cdot}$ denote the joint law of the stopped process $X_{t\wedge\cdot}$, while in Section 2 $\{\mu_\cdot\}$ denote the family of marginal laws.

For a path dependent function φ on $\mathbb{T} \times \mathbb{X} \times \mathcal{P}(\mathbb{X})$, we say φ is adapted if $\varphi(t, \mathbf{x}, \mu) = \varphi(t, \mathbf{x}_{t\wedge\cdot}, \mu_{t\wedge\cdot})$. Throughout this section, all the path dependent functions are required to be adapted. In particular, the data of the game $q : \mathbb{T} \times \mathbb{X} \times \mathcal{P}(\mathbb{X}) \times \mathbb{A} \times \mathbb{S} \rightarrow (0, 1)$,

$F : \mathbb{T} \times \mathbb{X} \times \mathcal{P}(\mathbb{X}) \times \mathbb{A} \rightarrow \mathbb{R}$, and $G : \mathbb{X} \times \mathcal{P}(\mathbb{X}) \rightarrow \mathbb{R}$ are path dependent with q, F adapted. By adapting to the path dependent setting, we shall still assume Assumption 2.2.

Let \mathcal{A}_{relax} denote the set of path dependent adapted relaxed controls $\gamma : \mathbb{T} \times \mathbb{X} \rightarrow \mathcal{P}(\mathbb{A})$. Given $t \in \mathbb{T}$, $\mu \in \mathcal{P}(\mathbb{X}_t)$, $\gamma \in \mathcal{A}_{relax}$, and $\mathbf{x} \in \mathbb{X}_t$, $\tilde{\mathbf{x}} \in \mathbb{X}^{t, \mathbf{x}}$, $\tilde{\gamma} \in \mathcal{A}_{relax}$, we introduce:

$$\begin{aligned} \mathbb{P}^{t, \mu, \gamma} \circ X_{t \wedge \cdot}^{-1} &= \mu, \quad \mathbb{P}^{t, \mu, \gamma}(X_{s+1} = \tilde{x} | X =_s \mathbf{x}) = \int_{\mathbb{A}} q(s, \mathbf{x}, \mu^\gamma, a; \tilde{x}) \gamma(s, \mathbf{x}; da); \\ &\text{where } \mu_{s \wedge \cdot}^\gamma := \mathbb{P}^{t, \mu, \gamma} \circ X_{s \wedge \cdot}^{-1}, \quad s \geq t; \\ \mathbb{P}^{\mu^\gamma; t, \mathbf{x}, \tilde{\gamma}}(X =_t \mathbf{x}) &= 1, \quad \mathbb{P}^{\mu^\gamma; t, \mathbf{x}, \tilde{\gamma}}(X_{s+1} = \tilde{x} | X =_s \tilde{\mathbf{x}}) = \int_{\mathbb{A}} q(s, \tilde{\mathbf{x}}, \mu^\gamma, a; \tilde{x}) \tilde{\gamma}(s, \tilde{\mathbf{x}}; da); \\ J(\mu^\gamma; s, \tilde{\mathbf{x}}, \tilde{\gamma}) &:= \mathbb{E}^{\mathbb{P}^{\mu^\gamma; t, \mathbf{x}, \tilde{\gamma}}} \left[G(X, \mu^\gamma) + \sum_{r=s}^{T-1} \int_{\mathbb{A}} F(r, X, \mu^\gamma, a) \tilde{\gamma}(r, X, da) \middle| X =_s \tilde{\mathbf{x}} \right]; \\ J(t, \mu, \gamma; \mathbf{x}, \tilde{\gamma}) &:= J(\mu^\gamma; t, \mathbf{x}, \tilde{\gamma}), \quad v(\mu^\gamma; s, \tilde{\mathbf{x}}) := \inf_{\tilde{\gamma} \in \mathcal{A}_{relax}} J(\mu^\gamma; s, \tilde{\mathbf{x}}, \tilde{\gamma}). \end{aligned} \quad (4.1)$$

Definition 4.1 (i) For any $t \in \mathbb{T}$, $\mu \in \mathcal{P}_0(\mathbb{X}_t)$, and $\varepsilon > 0$, let $\mathcal{M}_{relax}^\varepsilon(t, \mu)$ denote the set of relaxed ε -MFE $\gamma^* \in \mathcal{A}_{relax}$ such that

$$J(t, \mu, \gamma^*; \mathbf{x}, \gamma^*) \leq v(\mu^\gamma; t, \mathbf{x}) + \varepsilon, \quad \text{for all } \mathbf{x} \in \mathbb{X}_t. \quad (4.2)$$

(ii) The relaxed set value of the MFG at (t, μ) is defined as:

$$\mathbb{V}_{relax}(t, \mu) := \bigcap_{\varepsilon > 0} \mathbb{V}_{relax}^\varepsilon(t, \mu), \quad \text{where } \|\varphi\|_{\mathbb{X}_t} := \sup_{\mathbf{x} \in \mathbb{X}_t} |\varphi(\mathbf{x})|, \quad \text{and} \quad (4.3)$$

$$\mathbb{V}_{relax}^\varepsilon(t, \mu) := \left\{ \varphi \in \mathbb{L}^0(\mathbb{X}_t; \mathbb{R}) : \exists \gamma^* \in \mathcal{M}_{relax}^\varepsilon(t, \mu) \text{ s.t. } \|\varphi - J(t, \mu, \gamma^*; \cdot, \gamma^*)\|_{\mathbb{X}_t} \leq \varepsilon \right\}.$$

Similarly, given T_0 and $\psi : \mathbb{X}_{T_0} \times \mathcal{P}(\mathbb{X}_{T_0}) \rightarrow \mathbb{R}$, as in (2.8) define

$$J(T_0, \psi; t, \mu, \gamma; \mathbf{x}, \tilde{\gamma}) := \mathbb{E}^{\mathbb{P}^{\mu^\gamma; t, \mathbf{x}, \tilde{\gamma}}} \left[\psi(X_{T_0 \wedge \cdot}, \mu_{T_0 \wedge \cdot}^\gamma) + \sum_{s=t}^{T_0-1} \int_{\mathbb{A}} F(s, X, \mu^\gamma, a) \tilde{\gamma}(s, X, da) \right], \quad (4.4)$$

and let $\mathcal{M}_{relax}^\varepsilon(T_0, \psi; t, \mu)$ denote the set of $\gamma^* \in \mathcal{A}_{relax}$ such that

$$J(T_0, \psi; t, \mu, \gamma^*; \mathbf{x}, \gamma^*) \leq v(T, \psi; \mu^\gamma; s, \mathbf{x}) := \inf_{\gamma \in \mathcal{A}_{relax}} J(T_0, \psi; t, \mu, \gamma; \mathbf{x}, \gamma) + \varepsilon, \quad \forall \mathbf{x} \in \mathbb{X}_t. \quad (4.5)$$

Note that the tower property in (2.10) remains true for relaxed controls:

$$J(t, \mu, \gamma; \mathbf{x}, \tilde{\gamma}) = J(T_0, \psi; t, \mu, \gamma; \mathbf{x}, \tilde{\gamma}), \quad \text{where } \psi(\mathbf{y}, \nu) := J(T_0, \nu, \gamma; \mathbf{y}, \tilde{\gamma}). \quad (4.6)$$

The DPP for \mathbb{V}_{relax} takes the following form.

Theorem 4.2 Under Assumption 2.2 (i), for any $t \in \mathbb{T}$, $T_0 \in \mathbb{T}_t$, and $\mu \in \mathcal{P}_0(\mathbb{X}_t)$, we have

$$\begin{aligned} \mathbb{V}_{relax}(t, \mu) &:= \bigcap_{\varepsilon > 0} \left\{ \varphi \in \mathbb{L}^0(\mathbb{X}_t; \mathbb{R}) : \|\varphi - J(T_0, \psi; t, \mu, \gamma^*; \cdot, \gamma^*)\|_{\mathbb{X}_t} \leq \varepsilon \right. \\ &\quad \text{for some } \psi \in \mathbb{L}^0(\mathbb{X}_{T_0} \times \mathcal{P}_0(\mathbb{X}_{T_0}); \mathbb{R}) \text{ and } \gamma^* \in \mathcal{A}_{relax} \text{ such that} \\ &\quad \left. \psi(\cdot, \mu_{T_0 \wedge \cdot}^{\gamma^*}) \in \mathbb{V}_{relax}^\varepsilon(T_0, \mu_{T_0 \wedge \cdot}^{\gamma^*}), \quad \gamma^* \in \mathcal{M}_{relax}^\varepsilon(T_0, \psi; t, \mu) \right\}. \end{aligned} \quad (4.7)$$

Proof We shall follow the arguments in Theorem 2.4, in particular, we shall extend Proposition 2.3. Let $\tilde{\mathbb{V}}_{relax}(t, \mu) = \bigcap_{\varepsilon > 0} \tilde{\mathbb{V}}_{relax}^\varepsilon(t, \mu)$ denote the right side of (4.7).

(i) We first prove $\tilde{\mathbb{V}}_{relax}(t, \mu) \subset \mathbb{V}_{relax}(t, \mu)$. Fix $\varphi \in \tilde{\mathbb{V}}_{relax}(t, \mu)$, $\varepsilon > 0$, and set $\varepsilon_1 := \frac{\varepsilon}{4}$. Since $\varphi \in \tilde{\mathbb{V}}_{relax}^{\varepsilon_1}(t, \mu)$, then

$$\|\varphi - J(T_0, \psi; t, \mu, \gamma^*; \cdot, \gamma^*)\|_{\mathbb{X}_t} \leq \varepsilon_1 \quad \text{for some desirable } \psi, \gamma^* \text{ as in (4.7).}$$

Since $\psi(\cdot, \mu_{T_0 \wedge \cdot}^{\gamma^*}) \in \mathbb{V}_{relax}^{\varepsilon_1}(T_0, \mu_{T_0 \wedge \cdot}^{\gamma^*})$, there exists $\tilde{\gamma}^* \in \mathcal{M}_{relax}^{\varepsilon_1}(T_0, \mu_{T_0 \wedge \cdot}^{\gamma^*})$ such that

$$\|\psi(\cdot, \mu_{T_0 \wedge \cdot}^{\gamma^*}) - J(T_0, \mu_{T_0 \wedge \cdot}^{\gamma^*}, \tilde{\gamma}^*; \cdot, \tilde{\gamma}^*)\|_{\mathbb{X}_{T_0}} \leq \varepsilon_1.$$

As in (2.2) denote $\hat{\gamma}^* := \gamma^* \oplus_{T_0} \tilde{\gamma}^* := \gamma^* \mathbf{1}_{\{s < T_0\}} + \tilde{\gamma}^* \mathbf{1}_{\{s \geq T_0\}} \in \mathcal{A}_{relax}$. Then, for any $\mathbf{x} \in \mathbb{X}_t$ and $\gamma \in \mathcal{A}_{relax}$, similarly to Proposition 2.3 (i) we have

$$\begin{aligned} & J(t, \mu, \hat{\gamma}^*; \mathbf{x}, \gamma) \\ &= \mathbb{E}^{\mathbb{P}^{\mu^{\gamma^*}}; t, \mathbf{x}, \gamma} \left[J(T_0, \mu_{T_0 \wedge \cdot}^{\gamma^*}, \tilde{\gamma}^*; X_{T_0 \wedge \cdot}, \gamma) + \sum_{s=t}^{T_0-1} \int_{\mathbb{A}} F(s, X, \mu^{\gamma^*}, a) \gamma(s, X, da) \right] \\ &\geq \mathbb{E}^{\mathbb{P}^{\mu^{\gamma^*}}; t, \mathbf{x}, \gamma} \left[J(T_0, \mu_{T_0 \wedge \cdot}^{\gamma^*}, \tilde{\gamma}^*; X_{T_0 \wedge \cdot}, \tilde{\gamma}^*) + \sum_{s=t}^{T_0-1} \int_{\mathbb{A}} F(s, X, \mu^{\gamma^*}, a) \gamma(s, X, da) \right] - \varepsilon_1 \\ &\geq \mathbb{E}^{\mathbb{P}^{\mu^{\gamma^*}}; t, \mathbf{x}, \gamma} \left[\psi(X_{T_0 \wedge \cdot}, \mu_{T_0 \wedge \cdot}^{\gamma^*}) + \sum_{s=t}^{T_0-1} \int_{\mathbb{A}} F(s, X, \mu^{\gamma^*}, a) \gamma(s, X, da) \right] - 2\varepsilon_1 \\ &= J(T_0, \psi; t, \mu, \gamma^*; \mathbf{x}, \gamma) - 2\varepsilon_1 \geq J(T_0, \psi; t, \mu, \gamma^*; \mathbf{x}, \gamma^*) - 3\varepsilon_1 \\ &= \mathbb{E}^{\mathbb{P}^{\mu^{\gamma^*}}; t, \mathbf{x}, \gamma^*} \left[\psi(X_{T_0 \wedge \cdot}, \mu_{T_0 \wedge \cdot}^{\gamma^*}) + \sum_{s=t}^{T_0-1} \int_{\mathbb{A}} F(s, X, \mu^{\gamma^*}, a) \gamma^*(s, X, da) \right] - 3\varepsilon_1 \\ &\geq \mathbb{E}^{\mathbb{P}^{\mu^{\gamma^*}}; t, \mathbf{x}, \gamma^*} \left[J(T_0, \mu_{T_0 \wedge \cdot}^{\gamma^*}, \tilde{\gamma}^*; X_{T_0 \wedge \cdot}, \tilde{\gamma}^*) + \sum_{s=t}^{T_0-1} \int_{\mathbb{A}} F(s, X, \mu^{\gamma^*}, a) \gamma^*(s, X, da) \right] - 4\varepsilon_1 \\ &= J(t, \mu, \hat{\gamma}^*; \mathbf{x}, \hat{\gamma}^*) - 4\varepsilon_1 = J(t, \mu, \hat{\gamma}^*; \mathbf{x}, \hat{\gamma}^*) - \varepsilon. \end{aligned}$$

That is, $\hat{\gamma}^* \in \mathcal{M}_{relax}^\varepsilon(t, \mu)$. Moreover, note that, by (4.6),

$$\begin{aligned} & \|\varphi - J(t, \mu, \hat{\gamma}^*; \cdot, \hat{\gamma}^*)\|_{\mathbb{X}_t} \leq \varepsilon_1 + \|J(T_0, \psi; t, \mu, \gamma^*; \cdot, \gamma^*) - J(t, \mu, \hat{\gamma}^*; \cdot, \hat{\gamma}^*)\|_{\mathbb{X}_t} \\ &= \varepsilon_1 + \sup_{\mathbf{x} \in \mathbb{X}_t} \left| \mathbb{E}^{\mathbb{P}^{\mu^{\gamma^*}}; t, \mathbf{x}, \gamma^*} \left[\psi(X_{T_0 \wedge \cdot}, \mu_{T_0 \wedge \cdot}^{\gamma^*}) - J(T_0, \mu_{T_0 \wedge \cdot}^{\gamma^*}, \tilde{\gamma}^*; X_{T_0 \wedge \cdot}, \tilde{\gamma}^*) \right] \right| \leq 2\varepsilon_1 < \varepsilon. \end{aligned}$$

Then $\varphi \in \mathbb{V}_{relax}^\varepsilon(t, \mu)$. Since $\varepsilon > 0$ is arbitrary, we obtain $\varphi \in \mathbb{V}_{relax}(t, \mu)$.

(ii) We now prove the opposite inclusion. Fix $\varphi \in \mathbb{V}_{relax}(t, \mu)$ and $\varepsilon > 0$. Let $\varepsilon_2 > 0$ be a small number which will be specified later. Since $\varphi \in \mathbb{V}_{relax}^{\varepsilon_2}(t, \mu)$, then

$$\|\varphi - J(t, \mu, \gamma^*; \cdot, \gamma^*)\|_{\mathbb{X}_t} \leq \varepsilon_2 \quad \text{for some } \gamma^* \in \mathcal{M}_{relax}^{\varepsilon_2}(t, \mu).$$

Introduce $\psi(\mathbf{y}, \nu) := J(T_0, \nu, \gamma^*; \mathbf{y}, \gamma^*)$ and recall (4.6). Then

$$\|\varphi - J(T_0, \psi; t, \mu, \gamma^*; \cdot, \gamma^*)\|_{\mathbb{X}_t} = \|\varphi(\mathbf{x}) - J(t, \mu, \gamma^*; \mathbf{x}, \gamma^*)\|_{\mathbb{X}_t} \leq \varepsilon_2.$$

Moreover, since $\gamma^* \in \mathcal{M}_{relax}^{\varepsilon_2}(t, \mu)$, for any $\gamma \in \mathcal{A}_{relax}$ and $\mathbf{x} \in \mathbb{X}_t$, we have

$$\begin{aligned} J(T_0, \psi; t, \mu, \gamma^*; \mathbf{x}, \gamma^*) &= J(t, \mu, \gamma^*; \mathbf{x}, \gamma^*) \\ &\leq J(t, \mu, \gamma^*; \mathbf{x}, \gamma \oplus_{T_0} \gamma^*) + \varepsilon_2 = J(T_0, \psi; t, \mu, \gamma^*; \mathbf{x}, \gamma) + \varepsilon_2. \end{aligned}$$

This implies that $\gamma^* \in \mathcal{M}_{relax}^{\varepsilon_2}(T_0, \psi; t, \mu)$. We claim further that

$$\psi(\cdot, \mu_{T_0 \wedge \cdot}^{\gamma^*}) \in \mathbb{V}_{relax}^{C\varepsilon_2}(T_0, \mu_{T_0 \wedge \cdot}^{\gamma^*}), \quad (4.8)$$

for some constant $C \geq 1$. Then by (4.7) we see that $\varphi \in \tilde{\mathbb{V}}_{relax}^{C\varepsilon_2}(t, \mu) \subset \tilde{\mathbb{V}}_{relax}^{\varepsilon}(t, \mu)$ by setting $\varepsilon_2 \leq \frac{\varepsilon}{C}$. Since $\varepsilon > 0$ is arbitrary, we obtain $\varphi \in \tilde{\mathbb{V}}_{relax}(t, \mu)$.

To see (4.8), recalling (4.1), for any $\gamma \in \mathcal{A}_{relax}$ we have

$$\begin{aligned} &\mathbb{E}^{\mathbb{P}^{\mu^{\gamma^*}; t, \mathbf{x}, \gamma^*}} \left[J(T_0, \mu_{T_0 \wedge \cdot}^{\gamma^*}, \gamma^*; X_{T_0 \wedge \cdot}, \gamma^*) \right] - \mathbb{E}^{\mathbb{P}^{\mu^{\gamma^*}; t, \mathbf{x}, \gamma^*}} \left[J(T_0, \mu_{T_0 \wedge \cdot}^{\gamma^*}, \gamma^*; X_{T_0 \wedge \cdot}, \gamma) \right] \\ &= J(t, \mu, \gamma^*; \mathbf{x}, \gamma^*) - J(t, \mu, \gamma^*; \mathbf{x}, \gamma^* \oplus_{T_0} \gamma) \leq \varepsilon_2. \end{aligned}$$

Then, by taking infimum over $\gamma \in \mathcal{A}_{relax}$, it follows from the standard control theory that

$$\mathbb{E}^{\mathbb{P}^{\mu^{\gamma^*}; t, \mathbf{x}, \gamma^*}} \left[J(T_0, \mu_{T_0 \wedge \cdot}^{\gamma^*}, \gamma^*; X_{T_0 \wedge \cdot}, \gamma^*) \right] \leq \mathbb{E}^{\mathbb{P}^{\mu^{\gamma^*}; t, \mathbf{x}, \gamma^*}} \left[v(\mu^{\gamma^*}; T_0, X_{T_0 \wedge \cdot}) \right] + \varepsilon_2, \quad \forall \mathbf{x} \in \mathbb{X}_t.$$

On the other hand, it is obvious that $v(\mu^{\gamma^*}; T_0, \tilde{\mathbf{x}}) \leq J(T_0, \mu_{T_0 \wedge \cdot}^{\gamma^*}, \gamma^*; \tilde{\mathbf{x}}, \gamma^*)$ for all $\tilde{\mathbf{x}} \in \mathbb{X}_{T_0}$. Moreover, since $q \geq c_q$, clearly $\mathbb{P}^{\mu^{\gamma^*}; t, \mathbf{x}, \gamma^*}(X =_{T_0} \tilde{\mathbf{x}}) \geq c_q^{T_0-t}$, for any $\tilde{\mathbf{x}} \in \mathbb{X}_{T_0}^t$. Thus,

$$\begin{aligned} 0 &\leq J(T_0, \mu_{T_0 \wedge \cdot}^{\gamma^*}, \gamma^*; \tilde{\mathbf{x}}, \gamma^*) - v(\mu^{\gamma^*}; T_0, \tilde{\mathbf{x}}) \\ &\leq C \mathbb{E}^{\mathbb{P}^{\mu^{\gamma^*}; t, \mathbf{x}, \gamma^*}} \left[[J(T_0, \mu_{T_0 \wedge \cdot}^{\gamma^*}, \gamma^*; X_{T_0 \wedge \cdot}, \gamma^*) - v(\mu^{\gamma^*}; T_0, X_{T_0 \wedge \cdot})] \mathbf{1}_{\{X =_{T_0} \tilde{\mathbf{x}}\}} \right] \\ &\leq C \mathbb{E}^{\mathbb{P}^{\mu^{\gamma^*}; t, \mathbf{x}, \gamma^*}} \left[J(T_0, \mu_{T_0 \wedge \cdot}^{\gamma^*}, \gamma^*; X_{T_0 \wedge \cdot}, \gamma^*) - v(\mu^{\gamma^*}; T_0, X_{T_0 \wedge \cdot}) \right] \leq C\varepsilon_2, \end{aligned}$$

where $C := c_q^{t-T_0}$. This implies that $\gamma^* \in \mathcal{M}_{relax}^{C\varepsilon_2}(T_0, \mu_{T_0 \wedge \cdot}^{\gamma^*})$. Then (4.8) follows directly from $\psi(\cdot, \mu_{T_0 \wedge \cdot}^{\gamma^*}) = J(T_0, \mu_{T_0 \wedge \cdot}^{\gamma^*}, \gamma^*; \cdot, \gamma^*)$, and hence $\varphi \in \tilde{\mathbb{V}}_{relax}(t, \mu)$. \blacksquare

Remark 4.3 Consider the setting that q, F, G are state dependent, as in Section 2. There is a very subtle issue between state dependence and path dependence of the controls.

(i) For a standard non-zero sum game problems where the players may have different cost functions F_i, G_i , if one uses state dependent controls, in general the set value does not

satisfy DPP. See a counterexample in [24]. However, with path dependent controls the set value of the game satisfies the DPP.

(ii) In Section 2, since all players have the same cost function, as we saw the set value with state dependent controls satisfies DPP. If we consider path dependent controls $\alpha \in \mathcal{A}_{path}$, the set value will also satisfy DPP. However, the set values in these two settings are in general not equal, see Example 7.1 in Appendix for a counterexample.

(iii) For relaxed controls in this section, again restricting to state dependent q, F, G , it turns out that state dependent and path dependent controls lead to the same set value, see Theorem 7.4 in Appendix. The main reason is that the convex combination of relaxed controls remains a relaxed control, while the controls α in Section 2 does not share this property.

4.2 An alternative formulation of the relaxed mean field game

In this subsection we provide an alternative formulation for the MFG with relaxed controls. This new formulation is motivated from the heterogenous controls for the N -player games, and thus is crucial for the convergence result in the next section.

Let \mathcal{A}_{path} denote the set of adapted path dependent controls $\alpha : \mathbb{T} \times \mathbb{X} \rightarrow \mathbb{A}$, and for each $t \in \mathbb{T}$, $\mathcal{A}_{path}^t = \{(\alpha(t, \cdot), \dots, \alpha(T-1, \cdot)) : \alpha \in \mathcal{A}_{path}\}$. Denote $\Xi_t := \mathcal{P}(\mathbb{X}_t \times \mathcal{A}_{path}^t)$, and for each $\Lambda \in \Xi_t$, define recursively: for $s \geq t$, $\mathbf{x} \in \mathbb{X}_t$, and $\tilde{\mathbf{x}} \in \mathbb{X}^{t, \mathbf{x}}$,

$$\mu_{t \wedge \cdot}^\Lambda(\mathbf{x}) := \Lambda(\mathbf{x}, \mathcal{A}_{path}^t), \quad \mu_{s \wedge \cdot}^\Lambda(\tilde{\mathbf{x}}) := \int_{\mathcal{A}_{path}^t} \prod_{r=t}^{s-1} q(r, \tilde{\mathbf{x}}, \mu^\Lambda, \alpha(r, \tilde{\mathbf{x}}); \tilde{\mathbf{x}}_{r+1}) \Lambda(\mathbf{x}, d\alpha). \quad (4.9)$$

Here, noting that $\alpha \in \mathcal{A}_{path}^t$ can be equivalently expressed as $\{\alpha(s, \tilde{\mathbf{x}}) : t \leq s \leq T-1, \tilde{\mathbf{x}} \in \mathbb{X}_s^{t, \mathbf{x}}\}$, we are using the following interpretation on $d\alpha$: for any $\varphi : \mathcal{A}_{path}^t \rightarrow \mathbb{R}$,

$$\int_{\mathcal{A}_{path}^t} \varphi(\alpha) d\alpha := \int_{\mathbb{A}} \dots \int_{\mathbb{A}} \varphi(\{\alpha(s, \tilde{\mathbf{x}})\}) \prod_{s=t}^{T-1} \prod_{\tilde{\mathbf{x}} \in \mathbb{X}_s^{t, \mathbf{x}}} d\alpha(s, \tilde{\mathbf{x}}). \quad (4.10)$$

Next, for $\mu \in \mathcal{P}_0(\mathbb{X}_t)$, denote $\Xi_t(\mu) := \{\Lambda \in \Xi_t : \mu_{t \wedge \cdot}^\Lambda = \mu\}$. Moreover, recall (4.1),

$$J(t, \Lambda; \mathbf{x}, \alpha) := J(\mu^\Lambda; t, \mathbf{x}, \alpha), \quad v(t, \Lambda; \mathbf{x}) := v(\mu^\Lambda; t, \mathbf{x}), \quad \mathbf{x} \in \mathbb{X}_t, \alpha \in \mathcal{A}_{path}^t. \quad (4.11)$$

To simplify the notations, we introduce:

$$Q_s^t(\{\mu\}; \tilde{\mathbf{x}}, \alpha) := \prod_{r=t}^{s-1} q(r, \tilde{\mathbf{x}}, \mu, \alpha(r, \tilde{\mathbf{x}}); \tilde{\mathbf{x}}_{r+1}). \quad (4.12)$$

In particular, $Q_t^t(\{\mu\}; \mathbf{x}, \alpha) = 1$. Then we have, for any $\tilde{\mathbf{x}} \in \mathbb{X}^{t, \mathbf{x}}$,

$$\mu_s^\Lambda(\tilde{\mathbf{x}}) := \int_{\mathcal{A}_{path}^t} Q_s^t(\mu^\Lambda; \tilde{\mathbf{x}}, \alpha) \Lambda(\mathbf{x}, d\alpha), \quad \mathbb{P}^{\mu^\Lambda; t, \mathbf{x}, \alpha}(X =_s \tilde{\mathbf{x}}) = Q_s^t(\mu^\Lambda; \tilde{\mathbf{x}}, \alpha). \quad (4.13)$$

Definition 4.4 For any $t \in \mathbb{T}$, $\mu \in \mathcal{P}_0(\mathbb{X}_t)$, and $\varepsilon > 0$, we call $\Lambda^* \in \Xi_t(\mu)$ a global ε -MFE at (t, μ) , denoted as $\Lambda^* \in \mathcal{M}_{global}^\varepsilon(t, \mu)$, if

$$\int_{\mathcal{A}_{path}^t} [J(t, \Lambda^*; \mathbf{x}, \alpha) - v(t, \Lambda^*; \mathbf{x})] \Lambda^*(\mathbf{x}, d\alpha) \leq \varepsilon, \quad \forall \mathbf{x} \in \mathbb{X}_t. \quad (4.14)$$

Note that the above α is global in time, so we call Λ^* a global equilibrium. Moreover, since there are infinitely many $\alpha \in \mathcal{A}_{path}^t$, it is hard to require $J(t, \Lambda^*; \mathbf{x}, \alpha) - v(t, \Lambda^*; \mathbf{x}) \leq \varepsilon$ for each $\alpha \in \mathcal{A}_{path}^t$, we thus use the above \mathbb{L}^1 -type of optimality condition. For the \mathbf{x} part, however, since there are only finitely many \mathbf{x} and each of them has positive probability, we may require the optimality for each \mathbf{x} .

The main result of this subsection is the following equivalence result.

Theorem 4.5 For any $t \in \mathbb{T}$ and $\mu \in \mathcal{P}_0(\mathbb{X}_t)$, we have

$$\begin{aligned} \mathbb{V}_{relax}(t, \mu) &= \mathbb{V}_{global}(t, \mu) := \bigcap_{\varepsilon > 0} \mathbb{V}_{global}^\varepsilon(t, \mu), \quad \text{where} \\ \mathbb{V}_{global}^\varepsilon(t, \mu) &:= \left\{ \varphi \in \mathbb{L}^0(\mathbb{X}_t, \mathbb{R}) : \exists \Lambda^* \in \mathcal{M}_{global}^\varepsilon(t, \mu) \text{ s.t. } \|\varphi - v(t, \Lambda^*; \cdot)\|_{\mathbb{X}_t} \leq \varepsilon \right\}. \end{aligned} \quad (4.15)$$

We shall prove the mutual inclusion of the two sides separately. First, given (t, Λ) , we construct a relaxed control as follows: for any $t \in \mathbb{T}$, $\mathbf{x} \in \mathbb{X}_t$, and $s \geq t$, $\tilde{\mathbf{x}} \in \mathbb{X}_s^{t, \mathbf{x}}$,

$$\gamma^\Lambda(s, \tilde{\mathbf{x}}, da) := \frac{1}{\mu_{s \wedge \cdot}^\Lambda(\tilde{\mathbf{x}})} \int_{\mathcal{A}_{path}^t} Q_s^t(\mu^\Lambda; \tilde{\mathbf{x}}; \alpha) \delta_{\alpha(s, \tilde{\mathbf{x}})}(da) \Lambda(\mathbf{x}, d\alpha). \quad (4.16)$$

On the opposite direction, given $t \in \mathbb{T}$, $\mu \in \mathcal{P}_0(\mathbb{X}_t)$, $\gamma \in \mathcal{A}_{relax}$, recalling (4.10) we construct

$$\Lambda^\gamma(\mathbf{x}, d\alpha) := \mu(\mathbf{x}) \prod_{s=t}^{T-1} \prod_{\tilde{\mathbf{x}} \in \mathbb{X}_s^{t, \mathbf{x}}} \gamma(s, \tilde{\mathbf{x}}, d\alpha(s, \tilde{\mathbf{x}})), \quad \forall \mathbf{x} \in \mathbb{X}_t, \alpha \in \mathcal{A}_{path}^t. \quad (4.17)$$

In particular, the following calculation implies $\Lambda^\gamma \in \Xi_t(\mu)$:

$$\Lambda^\gamma(\mathbf{x}, \mathcal{A}_{path}^t) = \mu(\mathbf{x}) \prod_{s=t}^{T-1} \prod_{\tilde{\mathbf{x}} \in \mathbb{X}_s^{t, \mathbf{x}}} \gamma(s, \tilde{\mathbf{x}}, \mathbb{A}) = \mu(\mathbf{x}) \prod_{s=t}^{T-1} \prod_{\tilde{\mathbf{x}} \in \mathbb{X}_s^{t, \mathbf{x}}} 1 = \mu(\mathbf{x}).$$

Lemma 4.6 For any $t \in \mathbb{T}$, $\mu \in \mathcal{P}_0(\mathbb{X}_t)$, and $\Lambda \in \Xi_t(\mu)$, $\gamma \in \mathcal{A}_{relax}$, we have $\mu^{\gamma^\Lambda} = \mu^\Lambda$ and $\mu^{\Lambda^\gamma} = \mu^\gamma$. Moreover,

$$J(t, \mu, \gamma^\Lambda; \mathbf{x}, \gamma^\Lambda) = \frac{1}{\mu(\mathbf{x})} \int_{\mathcal{A}_{path}^t} J(t, \Lambda; \mathbf{x}, \alpha) \Lambda(\mathbf{x}, d\alpha), \quad \forall \mathbf{x} \in \mathbb{X}_t. \quad (4.18)$$

Proof We first prove $\mu_{s\wedge\cdot}^{\gamma^\Lambda} = \mu_{s\wedge\cdot}^\Lambda$ by induction. The case $s = t$ follows from the definitions. Assume it holds for all $r \leq s$. Now for $s + 1$ and $\tilde{\mathbf{x}} \in \mathbb{X}_{s+1}^{t,\mathbf{x}}$, by Fubini Theorem we have

$$\begin{aligned}
\frac{\mu_{(s+1)\wedge\cdot}^{\gamma^\Lambda}(\tilde{\mathbf{x}})}{\mu_{s\wedge\cdot}^{\gamma^\Lambda}(\tilde{\mathbf{x}}_{s\wedge\cdot})} &= \int_{\mathbb{A}} q(s, \tilde{\mathbf{x}}, \mu^{\gamma^\Lambda}, a; \tilde{\mathbf{x}}_{s+1}) \gamma^\Lambda(s, \tilde{\mathbf{x}}, da) \\
&= \int_{\mathbb{A}} q(s, \tilde{\mathbf{x}}, \mu^{\gamma^\Lambda}, a; \tilde{\mathbf{x}}_{s+1}) \frac{1}{\mu_{s\wedge\cdot}^\Lambda(\tilde{\mathbf{x}})} \int_{\mathcal{A}_{path}^t} Q_s^t(\mu^\Lambda; \tilde{\mathbf{x}}; \alpha) \delta_{\alpha(s, \tilde{\mathbf{x}})}(da) \Lambda(\mathbf{x}, d\alpha) \\
&= \frac{1}{\mu_{s\wedge\cdot}^\Lambda(\tilde{\mathbf{x}})} \int_{\mathcal{A}_{path}^t} q(s, \tilde{\mathbf{x}}, \mu^\Lambda, \alpha(s, \tilde{\mathbf{x}}); \tilde{\mathbf{x}}_{s+1}) Q_s^t(\mu^\Lambda; \tilde{\mathbf{x}}; \alpha) \Lambda(\mathbf{x}, d\alpha) \\
&= \frac{1}{\mu_{s\wedge\cdot}^\Lambda(\tilde{\mathbf{x}})} \int_{\mathcal{A}_{path}^t} Q_{s+1}^t(\mu^\Lambda; \tilde{\mathbf{x}}; \alpha) \Lambda(\mathbf{x}, d\alpha) = \frac{\mu_{(s+1)\wedge\cdot}^\Lambda(\tilde{\mathbf{x}})}{\mu_{s\wedge\cdot}^\Lambda(\tilde{\mathbf{x}})}.
\end{aligned}$$

Then $\mu_{(s+1)\wedge\cdot}^{\gamma^\Lambda} = \mu_{(s+1)\wedge\cdot}^\Lambda$, and we complete the induction argument.

We next prove $\mu_{s\wedge\cdot}^{\Lambda^\gamma} = \mu_{s\wedge\cdot}^{\Lambda^\gamma}$ by induction. Again the case $s = t$ is obvious. Assume it holds for all $r < s$. Now for s , recalling (4.10) we have

$$\begin{aligned}
\mu_{s\wedge\cdot}^{\Lambda^\gamma}(\tilde{\mathbf{x}}) &= \int_{\mathcal{A}_{path}^t} \left[\prod_{r=t}^{s-1} q(r, \tilde{\mathbf{x}}, \mu^\gamma, \alpha(r, \tilde{\mathbf{x}}); \tilde{\mathbf{x}}_{r+1}) \right] \left[\mu(\mathbf{x}) \prod_{r=t}^{T-1} \prod_{\bar{\mathbf{x}} \in \mathbb{X}_r^{t,\mathbf{x}}} \gamma(r, \bar{\mathbf{x}}, d\alpha(r, \bar{\mathbf{x}})) \right] \\
&= \mu(\mathbf{x}) \left[\prod_{r=t}^{s-1} \int_{\mathbb{A}} q(r, \tilde{\mathbf{x}}, \mu^\gamma, \alpha(r, \tilde{\mathbf{x}}); \tilde{\mathbf{x}}_{r+1}) \gamma(r, \tilde{\mathbf{x}}, d\alpha(r, \tilde{\mathbf{x}})) \right] \times \\
&\quad \left[\prod_{r=t}^{s-1} \prod_{\bar{\mathbf{x}} \in \mathbb{X}_r^{t,\mathbf{x}} \setminus \{\tilde{\mathbf{x}}\}} \gamma(r, \bar{\mathbf{x}}, \mathbb{A}) \right] \times \left[\prod_{r=s}^{T-1} \prod_{\bar{\mathbf{x}} \in \mathbb{X}_r^{t,\mathbf{x}}} \gamma(r, \bar{\mathbf{x}}, \mathbb{A}) \right] \\
&= \mu(\mathbf{x}) \prod_{r=t}^{s-1} \int_{\mathbb{A}} q(r, \tilde{\mathbf{x}}, \mu^\gamma, a; \tilde{\mathbf{x}}_{r+1}) \gamma(r, \tilde{\mathbf{x}}, da) = \mu_{s\wedge\cdot}^{\Lambda^\gamma}(\tilde{\mathbf{x}}).
\end{aligned}$$

We finally prove (4.18). For each $s \geq t$ and $\tilde{\mathbf{x}} \in \mathbb{X}_s^{t,\mathbf{x}}$, by Fubini Theorem again we have

$$\begin{aligned}
\int_{\mathbb{A}} F(s, \tilde{\mathbf{x}}, \mu^\Lambda, a) \gamma^\Lambda(s, \tilde{\mathbf{x}}, da) &= \int_{\mathbb{A}} \frac{F(s, \tilde{\mathbf{x}}, \mu^\Lambda, a)}{\mu_{s\wedge\cdot}^\Lambda(\tilde{\mathbf{x}})} \int_{\mathcal{A}_{path}^t} Q_s^t(\mu^\Lambda; \tilde{\mathbf{x}}; \alpha) \delta_{\alpha(s, \tilde{\mathbf{x}})}(da) \Lambda(\mathbf{x}, d\alpha) \\
&= \frac{1}{\mu_{s\wedge\cdot}^\Lambda(\tilde{\mathbf{x}})} \int_{\mathcal{A}_{path}^t} F(s, \tilde{\mathbf{x}}, \mu^\Lambda, \alpha(s, \tilde{\mathbf{x}})) Q_s^t(\mu^\Lambda; \tilde{\mathbf{x}}; \alpha) \Lambda(\mathbf{x}, d\alpha)
\end{aligned}$$

By (4.1) we have $\mathbb{P}^{\mu^\Lambda; t, \mathbf{x}, \gamma^\Lambda}(X =_s \tilde{\mathbf{x}}) = \frac{\mu_{s\wedge \cdot}^\Lambda(\tilde{\mathbf{x}})}{\mu(\mathbf{x})}$. Thus

$$\begin{aligned}
& J(t, \mu, \gamma^\Lambda; \mathbf{x}, \gamma^\Lambda) \\
&= \frac{1}{\mu(\mathbf{x})} \left[\sum_{\tilde{\mathbf{x}} \in \mathbb{X}^{t, \mathbf{x}}} G(\tilde{\mathbf{x}}, \mu^\Lambda) \mu_{T\wedge \cdot}^\Lambda(\tilde{\mathbf{x}}) + \sum_{s=t}^{T-1} \sum_{\tilde{\mathbf{x}} \in \mathbb{X}_s^{t, \mathbf{x}}} \mu_{s\wedge \cdot}^\Lambda(\tilde{\mathbf{x}}) \int_{\mathbb{A}} F(s, \tilde{\mathbf{x}}, \mu^\Lambda, a) \gamma^\Lambda(s, \tilde{\mathbf{x}}, da) \right] \\
&= \frac{1}{\mu(\mathbf{x})} \int_{\mathcal{A}_{path}^t} \left[\sum_{\tilde{\mathbf{x}} \in \mathbb{X}^{t, \mathbf{x}}} G(\tilde{\mathbf{x}}, \mu^\Lambda) Q_T^t(\mu^\Lambda; \tilde{\mathbf{x}}; \alpha) \right. \\
&\quad \left. + \sum_{s=t}^{T-1} \sum_{\tilde{\mathbf{x}} \in \mathbb{X}_s^{t, \mathbf{x}}} F(s, \tilde{\mathbf{x}}, \mu^\Lambda, \alpha(s, \tilde{\mathbf{x}})) Q_s^t(\mu^\Lambda; \tilde{\mathbf{x}}; \alpha) \right] \Lambda(\mathbf{x}, d\alpha).
\end{aligned}$$

This implies (4.18) immediately. \blacksquare

Remark 4.7 We can actually show that $\gamma^{(\Lambda^\gamma)} = \gamma$ for all $\gamma \in \mathcal{A}_{relax}$, see Appendix. However, it is not clear that we would have $\Lambda^{(\gamma^\Lambda)} = \Lambda$ for all $\Lambda \in \Xi_t(\mu)$.

Proof of Theorem 4.5. Since $\mu \in \mathcal{P}_0(\mathbb{X}_t)$ has full support, then $c_\mu := \inf_{\mathbf{x} \in \mathbb{X}_t} \mu(\mathbf{x}) > 0$.

(i) We first prove $\mathbb{V}_{global}(t, \mu) \subset \mathbb{V}_{relax}(t, \mu)$. Fix $\varphi \in \mathbb{V}_{global}(t, \mu)$ and $\varepsilon > 0$. Let $\varepsilon_1 > 0$ be a small number which will be specified later. Since $\varphi \in \mathbb{V}_{global}^{\varepsilon_1}(t, \mu)$, there exists $\Lambda^* \in \mathcal{M}_{global}^{\varepsilon_1}(t, \mu)$ such that $\|\varphi - v(t, \Lambda^*; \cdot)\|_{\mathbb{X}_t} \leq \varepsilon_1$. Set $\gamma^* := \gamma^{\Lambda^*}$. For any $\mathbf{x} \in \mathbb{X}_t$, since $\mu^{\gamma^*} = \mu^{\Lambda^*}$, by (4.1), (4.11) we have $v(\mu^{\gamma^*}; t, \mathbf{x}, \gamma^*) = v(t, \Lambda^*; \mathbf{x})$, and, by (4.18), (4.14),

$$J(t, \mu, \gamma^*; \mathbf{x}, \gamma^*) - v(t, \Lambda^*; \mathbf{x}) = \frac{1}{\mu(\mathbf{x})} \int_{\mathcal{A}_{path}^t} [J(t, \Lambda^*; \mathbf{x}, \alpha) - v(t, \Lambda^*; \mathbf{x})] \Lambda^*(\mathbf{x}, d\alpha) \leq \frac{\varepsilon_1}{c_\mu} \leq \varepsilon,$$

provided $\varepsilon_1 > 0$ is small enough. This implies $\gamma^* \in \mathcal{M}_{relax}^\varepsilon(t, \mu)$.

Moreover, it is clear now that, for any $\mathbf{x} \in \mathbb{X}_t$ and for a possibly smaller ε_1 ,

$$|\varphi(\mathbf{x}) - J(t, \mu, \gamma^*; \mathbf{x}, \gamma^*)| \leq \varepsilon_1 + |v(t, \Lambda^*; \mathbf{x}) - J(t, \mu, \gamma^*; \mathbf{x}, \gamma^*)| \leq \varepsilon_1 + \frac{\varepsilon_1}{c_\mu} \leq \varepsilon,$$

Then $\varphi \in \mathbb{V}_{relax}^\varepsilon(t, \mu)$, and since $\varepsilon > 0$ is arbitrary, we obtain $\varphi \in \mathbb{V}_{relax}(t, \mu)$.

(ii) We next prove $\mathbb{V}_{relax}(t, \mu) \subset \mathbb{V}_{global}(t, \mu)$. Fix $\varphi \in \mathbb{V}_{relax}(t, \mu)$, $\varepsilon > 0$, and set $\varepsilon_2 := \frac{\varepsilon}{2}$. Since $\varphi \in \mathbb{V}_{relax}^{\varepsilon_2}(t, \mu)$, there exists $\gamma^* \in \mathcal{M}_{relax}^{\varepsilon_2}(t, \mu)$ such that $\|\varphi - J(t, \mu, \gamma^*; \cdot, \gamma^*)\|_{\mathbb{X}_t} \leq \varepsilon_2$. Set $\Lambda^* := \Lambda^{\gamma^*}$, then $\mu^{\Lambda^*} = \mu^{\gamma^*}$. Since $\gamma^* \in \mathcal{M}_{relax}^{\varepsilon_2}(t, \mu)$, we have

$$|\varphi(\mathbf{x}) - v(t, \Lambda^*; \mathbf{x})| = |\varphi(\mathbf{x}) - v(\mu^{\gamma^*}; t, \mathbf{x})| \leq 2\varepsilon_2 \leq \varepsilon, \quad \forall \mathbf{x} \in \mathbb{X}_t.$$

Moreover, note that, by (4.18) again,

$$\begin{aligned}
& \int_{\mathcal{A}_{path}^t} [J(t, \Lambda^*; \mathbf{x}, \alpha) - v(t, \Lambda^*; \mathbf{x})] \Lambda^*(\mathbf{x}, d\alpha) \\
&= \mu(\mathbf{x}) [J(t, \mu, \gamma^*; \mathbf{x}, \gamma^*) - v(t, \Lambda^*; \mathbf{x})] \leq \mu(\mathbf{x}) \varepsilon_2 \leq \varepsilon_2 \leq \varepsilon.
\end{aligned} \tag{4.19}$$

This implies $\varphi \in \mathbb{V}_{global}^\varepsilon(t, \mu)$, and hence by the arbitrariness of ε , $\varphi \in \mathbb{V}_{global}(t, \mu)$. \blacksquare

5 The N -player game with heterogeneous equilibria

In this section we drop the requirement $\alpha^1 = \dots = \alpha^N$ for the N -player game, and show that the corresponding set value converges to \mathbb{V}_{relax} , which in general is strictly larger than \mathbb{V}_{state} . We note that we shall still use the pure strategies, rather than mixed strategies, for the N -player game. Moreover, since we used path dependent controls in Section 4, we shall also use path dependent controls here.

5.1 The N -player game

Let Ω^N and \vec{X} be as in Section 3, and denote

$$\mu_{t\wedge}^N := \mu_{t, \vec{X}_{t\wedge}}^N, \quad \text{where} \quad \mu_{t, \vec{x}}^N := \frac{1}{N} \sum_{i=1}^N \delta_{\mathbf{x}^i} \in \mathcal{P}(\mathbb{X}_t), \quad \vec{x} = (\mathbf{x}^1, \dots, \mathbf{x}^N) \in \mathbb{X}_t^N. \quad (5.1)$$

Similarly to (3.7), for the convenience of the presentation we introduce

$$\mathbb{X}_{0,t}^N := \left\{ \vec{x} \in \mathbb{X}_t^N : \text{supp}(\mu_{t, \vec{x}}^N) = \mathbb{X}_t \right\}, \quad \mathcal{P}_N(\mathbb{X}_t) := \left\{ \mu_{t, \vec{x}}^N : \vec{x} \in \mathbb{X}_{0,t}^N \right\}. \quad (5.2)$$

We shall consider path dependent symmetric controls: $\mathcal{A}_{path}^{t, \infty} := \bigcup_{L \geq 0} \mathcal{A}_{path}^{t, L}$, where

$$\mathcal{A}_{path}^{t, L} := \left\{ \alpha : \{t, \dots, T-1\} \times \mathbb{X} \times \mathcal{P}(\mathbb{X}) \rightarrow \mathbb{A} \mid \alpha \text{ is adapted and uniformly Lipschitz continuous in } \mu \text{ (under } W_1) \text{ with Lipschitz constant } L \right\}.$$

Given $t \in \mathbb{T}$, $\vec{x} \in \mathbb{X}_{0,t}^N$, and $\vec{\alpha} = (\alpha^1, \dots, \alpha^N) \in (\mathcal{A}_{path}^{t, \infty})^N$, introduce, for $s \geq t$,

$$\begin{aligned} \mathbb{P}^{t, \vec{x}, \vec{\alpha}}(\vec{X} =_t \vec{x}) &= 1, \quad \mathbb{P}^{t, \vec{x}, \vec{\alpha}}(\vec{X}_{s+1} = \vec{x}'' \mid \vec{X} =_s \vec{x}') = \prod_{i=1}^N q(s, \mathbf{x}''^i, \mu^N, \alpha^i(s, \mathbf{x}''^i, \mu^N); x_i''), \\ J_i(t, \vec{x}, \vec{\alpha}) &:= \mathbb{E}^{\mathbb{P}^{t, \vec{x}, \vec{\alpha}}} \left[G(X^i, \mu^N) + \sum_{s=t}^{T-1} F(s, X^i, \mu^N, \alpha^i(s, X^i, \mu^N)) \right]; \\ v_i^{N, L}(t, \vec{x}, \vec{\alpha}) &:= \inf_{\tilde{\alpha} \in \mathcal{A}_{path}^{t, L}} J_i(t, \vec{x}, \vec{\alpha}^{-i}, \tilde{\alpha}), \quad i = 1, \dots, N. \end{aligned} \quad (5.3)$$

Here $(\vec{\alpha}^{-i}, \tilde{\alpha})$ is the vector obtained by replacing α^i in $\vec{\alpha}$ with $\tilde{\alpha}$.

Definition 5.1 For any $\varepsilon > 0, L \geq 0$, we say $\vec{\alpha} \in (\mathcal{A}_{path}^{t, L})^N$ is an (ε, L) -equilibrium of the N -player game at (t, \vec{x}) , denoted as $\vec{\alpha} \in \mathcal{M}_{hetero}^{N, \varepsilon, L}(t, \vec{x})$, if:

$$\frac{1}{N} \sum_{i=1}^N [J_i(t, \vec{x}, \vec{\alpha}) - v_i^{N, L}(t, \vec{x}, \vec{\alpha})] \leq \varepsilon. \quad (5.4)$$

Here, since there are N players and we will send $N \rightarrow \infty$, similar to (4.14) we do not require the optimality for each player. In fact, by (5.4) one can easily show that

$$\frac{1}{N} \left| \{i = 1, \dots, N : J_i(t, \bar{\mathbf{x}}, \bar{\alpha}) - v_i^{N,L}(t, \bar{\mathbf{x}}, \bar{\alpha}) \geq \sqrt{\varepsilon}\} \right| \leq \sqrt{\varepsilon}. \quad (5.5)$$

This is exactly the $(\sqrt{\varepsilon}, \sqrt{\varepsilon})$ -equilibrium in [11].

We then define the set value of the N -player game with heterogeneous equilibria:

$$\begin{aligned} \mathbb{V}_{hetero}^N(t, \bar{\mathbf{x}}) &:= \bigcap_{\varepsilon > 0} \mathbb{V}_{hetero}^{N,\varepsilon}(t, \bar{\mathbf{x}}) := \bigcap_{\varepsilon > 0} \bigcup_{L \geq 0} \mathbb{V}_{hetero}^{N,\varepsilon,L}(t, \bar{\mathbf{x}}), \\ \text{where } \mathbb{V}_{hetero}^{N,\varepsilon,L}(t, \bar{\mathbf{x}}) &:= \left\{ \varphi \in \mathbb{L}^0(\mathbb{X}_t; \mathbb{R}) : \exists \bar{\alpha} \in \mathcal{M}_{hetero}^{N,\varepsilon,L}(t, \bar{\mathbf{x}}) \text{ such that} \right. \\ &\quad \left. \max_{\mathbf{x} \in \mathbb{X}_t} \min_{\{i: \mathbf{x}^i = \mathbf{x}\}} |\varphi(\mathbf{x}) - v_i^{N,L}(t, \bar{\mathbf{x}}, \bar{\alpha})| \leq \varepsilon \right\}. \end{aligned} \quad (5.6)$$

Remark 5.2 (i) An alternative definition of $\mathbb{V}_{hetero}^{N,\varepsilon,L}(t, \bar{\mathbf{x}})$ is to require φ satisfying

$$\max_{i=1, \dots, N} |\varphi(\mathbf{x}^i) - v_i^{N,L}(t, \bar{\mathbf{x}}, \bar{\alpha})| = \max_{\mathbf{x} \in \mathbb{X}_t} \max_{\{i: \mathbf{x}^i = \mathbf{x}\}} |\varphi(\mathbf{x}) - v_i^{N,L}(t, \bar{\mathbf{x}}, \bar{\alpha})| \leq \varepsilon. \quad (5.7)$$

Indeed, the convergence result Theorem 5.3 below remains true if we use (5.7). However, in general it is possible that $\mathbf{x}^i = \mathbf{x}^j$ but $v_i^{N,L}(t, \bar{\mathbf{x}}, \bar{\alpha}) \neq v_j^{N,L}(t, \bar{\mathbf{x}}, \bar{\alpha})$. Then, by fixing N and sending $\varepsilon \rightarrow 0$, under (5.7) we would have $\mathbb{V}_{hetero}^N(t, \bar{\mathbf{x}}) := \bigcap_{\varepsilon > 0} \mathbb{V}_{hetero}^{N,\varepsilon}(t, \bar{\mathbf{x}}) = \emptyset$.

(ii) In the homogeneous case, $v_i^{N,L}(t, \bar{\mathbf{x}}, \bar{\alpha}) = v_j^{N,L}(t, \bar{\mathbf{x}}, \bar{\alpha})$ whenever $\mathbf{x}^i = \mathbf{x}^j$, so we don't have this issue in (3.8).

(iii) Note that $\mu_{t, \bar{\mathbf{x}}}^N = \mu_{t, \bar{\mathbf{x}'}}^N$ if and only if $\bar{\mathbf{x}}$ is a permutation of $\bar{\mathbf{x}'}$, and one can easily verify that $v_i^{N,L}(t, \bar{\mathbf{x}}, \bar{\alpha}) = v_{\pi(i)}^{N,L}(t, (\mathbf{x}_{\pi(1)}, \dots, \mathbf{x}_{\pi(N)}), (\alpha_{\pi(1)}, \dots, \alpha_{\pi(N)}))$ for any permutation π on $\{1, \dots, N\}$. Then, similar to the homogenous case, $\mathbb{V}_{hetero}^{N,\varepsilon,L}(t, \bar{\mathbf{x}})$ is invariant in $\mu_{t, \bar{\mathbf{x}}}^N$ and we will denote it as $\mathbb{V}_{hetero}^{N,\varepsilon,L}(t, \mu_{t, \bar{\mathbf{x}}}^N)$.

The following convergence result of the set value is in the same spirit of Theorem 3.5.

Theorem 5.3 Let Assumption 2.2 hold and $\mu_{t, \bar{\mathbf{x}}}^N \in \mathcal{P}_N(\mathbb{X}_t) \rightarrow \mu \in \mathcal{P}_0(\mathbb{X}_t)$ under W_1 . Then

$$\bigcap_{\varepsilon > 0} \bigcup_{L \geq 0} \overline{\lim}_{N \rightarrow \infty} \mathbb{V}_{hetero}^{N,\varepsilon,L}(t, \mu_{t, \bar{\mathbf{x}}}^N) \subset \mathbb{V}_{relax}(t, \mu) \subset \bigcap_{\varepsilon > 0} \underline{\lim}_{N \rightarrow \infty} \mathbb{V}_{hetero}^{N,\varepsilon,0}(t, \mu_{t, \bar{\mathbf{x}}}^N). \quad (5.8)$$

In particular, since $\underline{\lim}_{N \rightarrow \infty} \mathbb{V}_{hetero}^{N,\varepsilon,0}(t, \mu_{t, \bar{\mathbf{x}}}^N) \subset \bigcup_{L \geq 0} \overline{\lim}_{N \rightarrow \infty} \mathbb{V}_{hetero}^{N,\varepsilon,L}(t, \mu_{t, \bar{\mathbf{x}}}^N)$, actually equalities hold.

Unlike Theorem 3.5, here the N -player game and the MFG take different types of controls $\bar{\alpha}$ and γ , respectively. The key for the convergence is the global formulation in

Subsection 4.2 for MFG. Indeed, given $t \in \mathbb{T}$, $\bar{\mathbf{x}} \in \mathbb{X}_{0,t}^N$, and $\bar{\alpha} \in (\mathcal{A}_{path}^{t,L})^N$, the N -player game is naturally related to the following $\Lambda^N \in \mathcal{P}(\mathbb{X}_t \times \mathcal{A}_{path}^{t,L})$:

$$\Lambda^N(\mathbf{x}, d\alpha) := \frac{1}{N} \sum_{i \in I(\mathbf{x})} \delta_{\alpha_i}(d\alpha), \text{ where } I(\mathbf{x}) := \{i = 1, \dots, N : \mathbf{x}^i = \mathbf{x}\}, \mathbf{x} \in \mathbb{X}_t. \quad (5.9)$$

By the symmetry of the problem, there exists a function J^N , independent of i , such that

$$J_i(t, \bar{\mathbf{x}}, \bar{\alpha}) = J^N(\Lambda^N; t, \mathbf{x}^i, \alpha_i), \quad i = 1, \dots, N. \quad (5.10)$$

We shall use this and Theorem 4.5 to prove Theorem 5.3 in the rest of this section. We also make the following obvious observation:

$$\Lambda^N(\mathbf{x}, \mathcal{A}_{path}^t) = \frac{|I(\mathbf{x})|}{N} = \mu_{t, \bar{\mathbf{x}}}^N(\mathbf{x}), \quad \forall \mathbf{x} \in \mathbb{X}_t. \quad (5.11)$$

Remark 5.4 (i) *In this section we are using symmetric controls and we obtain the convergence in Theorem 5.3. If we use full information controls $\alpha_i(t, \bar{X})$, as observed in [31] in terms of the equilibrium measure, one may expect the limit set value will be strictly larger than \mathbb{V}_{relax} . It will be interesting to find an appropriate notion of MFE so that the corresponding MFG set value will be equal to the above limit, in the sense of Theorem 5.3.*

(ii) *While the convergence in Theorem 5.3 is about set values, the proofs in the rest of this section confirm the convergence of the approximate equilibria as well, exactly in the same manner as in Remark 3.6.*

5.2 From N -player games to mean field games

In this subsection we prove the left inclusion in (5.8). Notice that the Λ^N in (5.9) is defined on $\mathcal{A}_{path}^{t,L}$, rather than $\mathcal{A}_{path}^t = \mathcal{A}_{path}^{t,0}$. For this purpose, recall (4.12) and introduce

$$\begin{aligned} \nu_{i \wedge \cdot}^N(\mathbf{x}) &:= \mu_{t, \bar{\mathbf{x}}}^N(\mathbf{x}), \quad \nu_{s \wedge \cdot}^N(\tilde{\mathbf{x}}) := \frac{1}{N} \sum_{i \in I(\mathbf{x})} Q_s^t(\nu^N; \tilde{\mathbf{x}}, \alpha_i(\cdot, \cdot, \nu^N)), \quad \mathbf{x} \in \mathbb{X}_t, \tilde{\mathbf{x}} \in \mathbb{X}_s^{t, \mathbf{x}}, s \geq t; \\ \bar{\Lambda}^N(\mathbf{x}, d\alpha) &:= \frac{\mu(\mathbf{x})}{|I(\mathbf{x})|} \sum_{i \in I(\mathbf{x})} \delta_{\bar{\alpha}_i}(d\alpha), \quad \text{where } \bar{\alpha}_i(s, \tilde{\mathbf{x}}) := \alpha_i(s, \tilde{\mathbf{x}}, \nu^N). \end{aligned} \quad (5.12)$$

Then it is obvious that $\bar{\alpha}_i \in \mathcal{A}_{path}^t$ and $\bar{\Lambda}^N \in \Xi_t(\mu)$. Moreover, when $\mu = \mu_{t, \bar{\mathbf{x}}}^N$, by (4.13) and (5.11) it is straightforward to verify by induction that $\mu^{\bar{\Lambda}^N} = \nu^N$.

Theorem 5.5 *Let Assumption 2.2 (ii) hold. Then, for any $L \geq 0$, there exists a constant C_L , depending only on T, d, L_q , and L such that, for any $t \in \mathbb{T}$, $\bar{\mathbf{x}} \in \mathbb{X}_{0,t}^N$, $\mu \in \mathcal{P}_0(\mathbb{X}_t)$, $\bar{\alpha} \in (\mathcal{A}_{path}^{t,L})^N$, $\tilde{\alpha} \in \mathcal{A}_{path}^{t,L}$, and for the $\nu^N, \bar{\Lambda}^N$ defined in (5.12), we have*

$$\max_{1 \leq i \leq N} \max_{t \leq s \leq T} \mathbb{E}^{\mathbb{P}^{t, \bar{\mathbf{x}}, (\bar{\alpha}^{-i}, \tilde{\alpha})}} [\mathcal{W}_1(\mu_{s \wedge \cdot}^N, \mu_{s \wedge \cdot}^{\bar{\Lambda}^N})] \leq C_L \theta_N, \text{ where } \theta_N := W_1(\mu_{t, \bar{\mathbf{x}}}^N, \mu) + \frac{1}{\sqrt{N}}. \quad (5.13)$$

Proof Fix i and denote $\tilde{\alpha}_j := \alpha_j$ for $j \neq i$, and $\tilde{\alpha}_i := \tilde{\alpha}_i$. We first show that

$$\kappa_s := \mathbb{E}^{\mathbb{P}^N} [\mathcal{W}_1(\mu_{s\wedge\cdot}^N, \nu_{s\wedge\cdot}^N)] \leq \frac{C_L}{\sqrt{N}}, \quad \text{where } \mathbb{P}^N := \mathbb{P}^{t, \vec{x}, (\tilde{\alpha}^{-i}, \tilde{\alpha})}. \quad (5.14)$$

Indeed, for $s \geq t$, by the conditional independence of $\{X_{s+1}^j\}_{1 \leq j \leq N}$ under \mathbb{P}^N , conditional on \mathcal{F}_s , it follows from the same arguments as in (3.11) that

$$\begin{aligned} \kappa_{s+1} &= \mathbb{E}^{\mathbb{P}^N} \left[\mathbb{E}_{\mathcal{F}_s}^{\mathbb{P}^N} [\mathcal{W}_1(\mu_{(s+1)\wedge\cdot}^N, \nu_{(s+1)\wedge\cdot}^N)] \right] \\ &\leq \frac{C}{\sqrt{N}} + C \sum_{\mathbf{x} \in \mathbb{X}_{s+1}} \mathbb{E}^{\mathbb{P}^N} \left[\left| \frac{1}{N} \sum_{j=1}^N \mathbb{P}^N(X^j =_{s+1} \mathbf{x} | \mathcal{F}_s) - \nu_{(s+1)\wedge\cdot}^N(\mathbf{x}) \right| \right]. \end{aligned}$$

Note that,

$$\begin{aligned} &\left| \frac{1}{N} \sum_{j=1}^N \mathbb{P}^N(X^j =_{s+1} \mathbf{x} | \mathcal{F}_s) - \frac{1}{N} \sum_{j=1}^N \mathbf{1}_{\{X^j =_s \mathbf{x}\}} q(s, \mathbf{x}, \nu^N, \alpha_j(s, \mathbf{x}, \nu^N); \mathbf{x}_{s+1}) \right| \\ &= \left| \frac{1}{N} \sum_{j=1}^N \mathbf{1}_{\{X^j =_s \mathbf{x}\}} [q(s, \mathbf{x}, \mu^N, \tilde{\alpha}_j(s, \mathbf{x}, \mu^N); \mathbf{x}_{s+1}) - q(s, \mathbf{x}, \nu^N, \alpha_j(s, \mathbf{x}, \nu^N); \mathbf{x}_{s+1})] \right| \\ &\leq C_L W_1(\mu_{s\wedge\cdot}^N, \nu_{s\wedge\cdot}^N) + \frac{1}{N} = C_L \kappa_s + \frac{1}{N}, \end{aligned}$$

where in the last inequality, the first term is due to the sum over all $j \neq i$. Then

$$\begin{aligned} \kappa_{s+1} &\leq C_L \kappa_s + \frac{C}{\sqrt{N}} + \mathbb{E}^{\mathbb{P}^N} \left[\sum_{\mathbf{x} \in \mathbb{X}_{s+1}} \left| \frac{1}{N} \sum_{j=1}^N \mathbf{1}_{\{X^j =_s \mathbf{x}\}} q(s, \mathbf{x}, \nu^N, \alpha_j(s, \mathbf{x}, \nu^N); \mathbf{x}_{s+1}) \right. \right. \\ &\quad \left. \left. - \frac{1}{N} \sum_{j \in I(\mathbf{x}_{t\wedge\cdot})} Q_s^t(\nu^N; \mathbf{x}, \tilde{\alpha}_j) q(s, \mathbf{x}, \nu^N, \alpha_j(s, \mathbf{x}, \nu^N); \mathbf{x}_{s+1}) \right| \right] \\ &= C_L \kappa_s + \frac{C}{\sqrt{N}} + \mathbb{E}^{\mathbb{P}^N} \left[\sum_{\mathbf{x} \in \mathbb{X}_s} \left| \frac{1}{N} \sum_{j=1}^N \mathbf{1}_{\{X^j =_s \mathbf{x}\}} - \frac{1}{N} \sum_{j \in I(\mathbf{x}_{t\wedge\cdot})} Q_s^t(\nu^N; \mathbf{x}, \tilde{\alpha}_j) \right| \right] \\ &= C_L \kappa_s + \frac{C}{\sqrt{N}} + \mathbb{E}^{\mathbb{P}^N} \left[\sum_{\mathbf{x} \in \mathbb{X}_s} |\mu_{s\wedge\cdot}^N(\mathbf{x}) - \nu_{s\wedge\cdot}^N(\mathbf{x})| \right] \leq C_L \kappa_s + \frac{C}{\sqrt{N}}. \end{aligned}$$

It is obvious that $\kappa_t = 0$. Then by induction we obtain (5.14).

Next, denote $\bar{\kappa}_s := W_1(\nu_{s\wedge\cdot}^N, \mu_{s\wedge\cdot}^{\bar{\Lambda}^N})$. For $s \geq t$, by (5.12), (4.13), and (4.12), we have

$$\begin{aligned}
\bar{\kappa}_{s+1} &= \sum_{\mathbf{x} \in \mathbb{X}_t} \sum_{\tilde{\mathbf{x}} \in \mathbb{X}_{s+1}^{t, \mathbf{x}}} |\nu_{(s+1)\wedge\cdot}^N(\tilde{\mathbf{x}}) - \mu_{(s+1)\wedge\cdot}^{\bar{\Lambda}^N}(\tilde{\mathbf{x}})| \\
&= \sum_{\mathbf{x} \in \mathbb{X}_t} \sum_{\tilde{\mathbf{x}} \in \mathbb{X}_{s+1}^{t, \mathbf{x}}} \left| \frac{1}{N} \sum_{j \in I(\mathbf{x})} Q_{s+1}^t(\nu^N; \tilde{\mathbf{x}}, \bar{\alpha}_j) - \frac{\mu(\mathbf{x})}{|I(\mathbf{x})|} \sum_{j \in I(\mathbf{x})} Q_{s+1}^t(\mu^{\bar{\Lambda}^N}; \tilde{\mathbf{x}}, \bar{\alpha}_j) \right| \\
&= \sum_{\mathbf{x} \in \mathbb{X}_t} \sum_{\tilde{\mathbf{x}} \in \mathbb{X}_{s+1}^{t, \mathbf{x}}} \left[\frac{1}{N} \sum_{j \in I(\mathbf{x})} |Q_{s+1}^t(\nu^N; \tilde{\mathbf{x}}, \bar{\alpha}_j) - Q_{s+1}^t(\mu^{\bar{\Lambda}^N}; \tilde{\mathbf{x}}, \bar{\alpha}_j)| \right. \\
&\quad \left. + \left| \frac{1}{N} - \frac{\mu(\mathbf{x})}{|I(\mathbf{x})|} \right| \sum_{j \in I(\mathbf{x})} Q_{s+1}^t(\mu^{\bar{\Lambda}^N}; \tilde{\mathbf{x}}, \bar{\alpha}_j) \right] \\
&\leq C \sum_{\mathbf{x} \in \mathbb{X}_t} \sum_{\tilde{\mathbf{x}} \in \mathbb{X}_{s+1}^{t, \mathbf{x}}} \left[\frac{1}{N} \sum_{j \in I(\mathbf{x})} \sum_{r=t}^s W_1(\nu_{r\wedge\cdot}^N, \mu_{r\wedge\cdot}^{\bar{\Lambda}^N}) + \left| \frac{1}{N} - \frac{\mu(\mathbf{x})}{|I(\mathbf{x})|} \right| |I(\mathbf{x})| \right] \\
&\leq C \sum_{r=t}^s \bar{\kappa}_r + C \sum_{\mathbf{x} \in \mathbb{X}_t} |\mu_{t, \tilde{\mathbf{x}}}^N(\mathbf{x}) - \mu(\mathbf{x})| \leq C \sum_{r=t}^s \bar{\kappa}_r.
\end{aligned}$$

Obviously $\bar{k}_t = W_1(\mu_{t, \tilde{\mathbf{x}}}^N, \mu)$. Then by induction we have $\sup_{t \leq s \leq T} \bar{\kappa}_s \leq CW_1(\mu_{t, \tilde{\mathbf{x}}}^N, \mu)$. This, together with (5.14), implies (5.13) immediately. \blacksquare

Theorem 5.6 *For the setting in Theorem 5.5 and assuming further Assumption 2.2 (iii), there exists a modulus of continuity function ρ_L , depending on T, d, L_Q, C_0, ρ, L , such that*

$$\left| J_i(t, \vec{x}, (\vec{\alpha}^{-i}, \vec{\alpha})) - J(t, \bar{\Lambda}^N; \mathbf{x}^i, \tilde{\alpha}(\cdot, \nu^N)) \right| + |v_i^{N,L}(t, \vec{x}, \vec{\alpha}) - v(\mu^{\bar{\Lambda}^N}; t, \mathbf{x}^i)| \leq \rho_L(\theta_N). \quad (5.15)$$

Moreover, assume $\vec{\alpha} \in \mathcal{M}_{hetero}^{N, \varepsilon_1, L}(t, \vec{x})$ for some $\varepsilon_1 > 0$, then

$$\int_{\mathcal{A}_{path}^t} [J(t, \bar{\Lambda}^N; \mathbf{x}, \alpha) - v(t, \bar{\Lambda}^N; \mathbf{x})] \bar{\Lambda}^N(\mathbf{x}, d\alpha) \leq \varepsilon_1 + 2\rho_L(\theta_N), \quad \forall \mathbf{x} \in \mathbb{X}_t. \quad (5.16)$$

In particular, if $\varepsilon_1 + 2\rho_L(\theta_N) \leq \varepsilon$, then $\bar{\Lambda}^N \in \mathcal{M}_{global}^\varepsilon(t, \mu)$.

Proof First, given Theorem 5.5, (5.15) follows from the arguments in Theorem 3.4. Then, for $\vec{\alpha} \in \mathcal{M}_{hetero}^{N, \varepsilon_1, L}(t, \vec{x})$ and $\mathbf{x} \in \mathbb{X}_t$, by (5.4) we have

$$\begin{aligned}
&\int_{\mathcal{A}_{path}^t} [J(t, \bar{\Lambda}^N; \mathbf{x}, \alpha) - v(t, \bar{\Lambda}^N; \mathbf{x})] \bar{\Lambda}^N(\mathbf{x}, d\alpha) = \frac{1}{N} \sum_{i \in I(\mathbf{x})} [J(t, \bar{\Lambda}^N; \mathbf{x}, \bar{\alpha}_i) - v(t, \bar{\Lambda}^N; \mathbf{x})] \\
&\leq \frac{1}{N} \sum_{i \in I(\mathbf{x})} \left[|J(t, \bar{\Lambda}^N; \mathbf{x}^i, \bar{\alpha}_i) - J_i(t, \vec{x}, \vec{\alpha})| + |J_i(t, \vec{x}, \vec{\alpha}) - v_i^{N,L}(t, \vec{x}, \vec{\alpha})| \right. \\
&\quad \left. + |v_i^{N,L}(t, \vec{x}, \vec{\alpha}) - v(\mu^{\bar{\Lambda}^N}; t, \mathbf{x}^i)| \right] \\
&\leq \rho_L(\theta_N) + \varepsilon_1 + \rho_L(\theta_N) = \varepsilon_1 + 2\rho_L(\theta_N),
\end{aligned}$$

completing the proof. \blacksquare

Proof of Theorem 5.3: the left inclusion. Fix $\varphi \in \bigcap_{\varepsilon > 0} \bigcup_{L \geq 0} \overline{\lim}_{N \rightarrow \infty} \mathbb{V}_{hetero}^{N, \varepsilon, L}(t, \mu_{t, \bar{\mathbf{x}}}^N)$, $\varepsilon > 0$, and set $\varepsilon_1 := \frac{\varepsilon}{2}$. Then there exists $L_\varepsilon \geq 0$ and a sequence $N_k \rightarrow \infty$ (possibly depending on ε) such that $\varphi \in \mathbb{V}_{hetero}^{N_k, \varepsilon_1, L_\varepsilon}(t, \mu_{t, \bar{\mathbf{x}}}^{N_k})$, for all $k \geq 1$. Now choose k large enough so that $2\rho_{L_\varepsilon}(\theta_{N_k}) \leq \varepsilon_1$. By (5.6) there exists $\vec{\alpha} \in \mathcal{M}_{hetero}^{N_k, \varepsilon_1, L_\varepsilon}(t, \bar{\mathbf{x}})$ such that $\max_{\mathbf{x} \in \mathbb{X}_t} \min_{i \in I(\mathbf{x})} |\varphi(\mathbf{x}) - v_i^{N, L}(t, \bar{\mathbf{x}}, \vec{\alpha})| \leq \varepsilon_1$. By Theorem 5.6 we see that $\bar{\Lambda}^{N_k} \in \mathcal{M}_{global}^\varepsilon(t, \mu)$ and, by (5.15),

$$\begin{aligned} \|\varphi - v(\mu^{\bar{\Lambda}^N}; t, \cdot)\|_{\mathbb{X}_t} &\leq \max_{\mathbf{x} \in \mathbb{X}_t} \min_{i \in I(\mathbf{x})} \left[|\varphi(\mathbf{x}) - v_i^{N, L}(t, \bar{\mathbf{x}}, \vec{\alpha})| + |v_i^{N, L}(t, \bar{\mathbf{x}}, \vec{\alpha}) - v(\mu^{\bar{\Lambda}^N}; t, \mathbf{x})| \right] \\ &\leq \varepsilon_1 + \rho_{L_\varepsilon}(\theta_N) \leq \varepsilon. \end{aligned}$$

That is, $\varphi \in \mathbb{V}_{global}^\varepsilon(t, \mu)$.

Since $\varepsilon > 0$ is arbitrary, applying Theorem 4.5 we obtain $\varphi \in \mathbb{V}_{relax}(t, \mu)$. \blacksquare

5.3 From mean field games to N -player games

We now turn to the right inclusion in (5.8). Fix $t \in \mathbb{T}$, $\bar{\mathbf{x}} \in \mathbb{X}_{0, t}^N$, $\mu \in \mathcal{P}_0(\mathbb{X}_t)$, and $\gamma \in \mathcal{A}_{relax}$. Our goal is to construct a desired $\vec{\alpha} \in (\mathcal{A}_{path}^{t, 0})^N$. However, since $\vec{\alpha}$, or equivalently the corresponding Λ^N , is discrete, we need to discretize γ first. We note that it is slightly easier to discretize γ than to discretize a general $\Lambda \in \Xi_t(\mu)$.

First, given $\varepsilon > 0$, there exists a partition $\mathbb{A} = \bigcup_{k=0}^{n_\varepsilon} A_k$ with n_ε depending on ε (and γ) such that, for some arbitrarily fixed $a_k \in A_k$, $k = 0, \dots, n_\varepsilon$,

$$\gamma(s, \mathbf{x}, A_0) \leq \varepsilon, \forall s \in \mathbb{T}_t, \mathbf{x} \in \mathbb{X}_s, \quad \text{and} \quad |a - a_k| \leq \varepsilon, \forall a \in A_k, k = 1, \dots, n_\varepsilon. \quad (5.17)$$

Denote by $\mathcal{A}_{path}^{t, \varepsilon}$ the subset of $\alpha \in \mathcal{A}_{path}^{t, 0}$ taking values in $\mathbb{A}_\varepsilon := \{a_k : k = 0, \dots, n_\varepsilon\}$. Define

$$\gamma^\varepsilon(s, \mathbf{x}, da) := \sum_{k=0}^{n_\varepsilon} \gamma(s, \mathbf{x}, A_k) \delta_{a_k}(da). \quad (5.18)$$

Recall (4.17), we see that $\text{supp}(\Lambda^{\gamma^\varepsilon}(\mathbf{x}, d\alpha)) = \mathcal{A}_{path}^{t, \varepsilon} \subset \mathcal{A}_{path}^{t, 0}$ for all $\mathbf{x} \in \mathbb{X}_t$.

Next, recall (5.11) that $N\mu_{t, \bar{\mathbf{x}}}^N(\mathbf{x}) = |I(\mathbf{x})|$ is a positive integer for all $\mathbf{x} \in \mathbb{X}_t$. Let $\Lambda_{t, \bar{\mathbf{x}}}^\varepsilon \in \mathcal{P}(\mathbb{X}_t \times \mathcal{A}_{path}^{t, \varepsilon})$ be a modification of $\Lambda^{\gamma^\varepsilon}$ such that,

$$\begin{aligned} \Lambda_{t, \bar{\mathbf{x}}}^\varepsilon(\mathbf{x}, \mathcal{A}_{path}^{t, \varepsilon}) &= \mu_{t, \bar{\mathbf{x}}}^N(\mathbf{x}) \text{ and } N\Lambda_{t, \bar{\mathbf{x}}}^\varepsilon(\mathbf{x}, \alpha) \text{ is an integer;} \\ |\Lambda_{t, \bar{\mathbf{x}}}^\varepsilon(\mathbf{x}, \alpha) - \Lambda^{\gamma^\varepsilon}(\mathbf{x}, \alpha)| &\leq \frac{1}{N} + |\mu_{t, \bar{\mathbf{x}}}^N(\mathbf{x}) - \mu(\mathbf{x})|; \quad \forall (\mathbf{x}, \alpha) \in \mathbb{X}_t \times \mathcal{A}_{path}^{t, \varepsilon}. \end{aligned} \quad (5.19)$$

Note that, since $\mathcal{A}_{path}^{t, \varepsilon}$ is finite, such a construction is easy.

We now construct $\vec{\alpha} \in (\mathcal{A}_{path}^{t,\varepsilon})^N$, which relies on γ^ε and hence on ε . Note that

$$\sum_{\alpha \in \mathcal{A}_{path}^{t,\varepsilon}} [N\Lambda_{t,\vec{x}}^\varepsilon(\mathbf{x}, \alpha)] = N\Lambda_{t,\vec{x}}^\varepsilon(\mathbf{x}, \mathcal{A}_{path}^{t,\varepsilon}) = N\mu_{t,\vec{x}}^N(\mathbf{x}) = |I(\mathbf{x})|,$$

and each $N\Lambda_{t,\vec{x}}^\varepsilon(\mathbf{x}, \alpha)$ is an integer. Let $I(\mathbf{x}) = \cup_{\alpha \in \mathcal{A}_{path}^{t,\varepsilon}} I(\mathbf{x}, \alpha)$ be a partition of $I(\mathbf{x})$ such that $|I(\mathbf{x}, \alpha)| = N\Lambda_{t,\vec{x}}^\varepsilon(\mathbf{x}, \alpha)$. We then set

$$\alpha_i := \alpha, \quad i \in I(\mathbf{x}, \alpha), \quad (\mathbf{x}, \alpha) \in \mathbb{X}_t \times \mathcal{A}_{path}^{t,\varepsilon}. \quad (5.20)$$

Let Λ^N be the one defined by (5.9) corresponding to this $\vec{\alpha}$. It is clear that $\Lambda^N = \Lambda_{t,\vec{x}}^\varepsilon$.

Theorem 5.7 (i) *Let Assumption 2.2 (ii) hold. Then there exists a constant C , depending only on T, d, L_q , such that, for any $t \in \mathbb{T}$, $\vec{x} \in \mathbb{X}_{0,t}^N$, $\mu \in \mathcal{P}_0(\mathbb{X}_t)$, $\gamma \in \mathcal{A}_{relax}$, $\varepsilon > 0$, and for the $\vec{\alpha} \in (\mathcal{A}_{path}^{t,\varepsilon})^N$ constructed above, we have, for the θ_N in (5.13) and for any $\tilde{\alpha} \in \mathcal{A}_{path}^{t,0}$,*

$$\max_{1 \leq i \leq N} \max_{t \leq s \leq T} \mathbb{E}^{\mathbb{P}^{t,\vec{x},(\vec{\alpha}^{-i}, \tilde{\alpha})}} [W_1(\mu_{s\wedge\cdot}^N, \mu_{s\wedge\cdot}^\gamma)] \leq C\varepsilon + C_\varepsilon\theta_N, \quad (5.21)$$

where C_ε may depend on ε as well.

(ii) *Assume further Assumption 2.2 (iii), then there exists a modulus of continuity function ρ_0 , depending only on T, d, L_q, C_0 , and ρ , such that,*

$$\left| J_i(t, \vec{x}, (\vec{\alpha}^{-i}, \tilde{\alpha})) - J(\mu^\gamma; t, \mathbf{x}^i, \tilde{\alpha}) \right| + \left| v_i^{N,0}(t, \vec{x}, \vec{\alpha}) - v(\mu^\gamma; t, \mathbf{x}^i) \right| \leq \rho_0(C\varepsilon + C_\varepsilon\theta_N). \quad (5.22)$$

Moreover, assume $\gamma \in \mathcal{M}_{relax}^\varepsilon(t, \mu)$, then

$$\frac{1}{N} \sum_{i=1}^N [J_i(t, \vec{x}, \vec{\alpha}) - v_i^{N,0}(t, \vec{x}, \vec{\alpha})] \leq \varepsilon + 2\rho_0(C\varepsilon + C_\varepsilon\theta_N), \quad \forall \mathbf{x} \in \mathbb{X}_t. \quad (5.23)$$

In particular, this means that $\vec{\alpha} \in \mathcal{M}_{hetero}^{N,\varepsilon,0}(t, \vec{x})$ with $\tilde{\varepsilon} := \varepsilon + 2\rho_0(C\varepsilon + C_\varepsilon\theta_N)$.

Proof (i) We first show by induction that

$$\kappa_s := W_1(\mu_{s\wedge\cdot}^\gamma, \mu_{s\wedge\cdot}^{\gamma^\varepsilon}) \leq C\varepsilon, \quad s = t, \dots, T. \quad (5.24)$$

Indeed, it is obvious that $\kappa_t = 0$. For $s \geq t$, by (4.1), (5.17), and (5.18), we have

$$\begin{aligned} \kappa_{s+1} &= \sum_{\mathbf{x} \in \mathbb{X}_{s+1}} \left| \mu_{(s+1)\wedge\cdot}^\gamma(\mathbf{x}) - \mu_{(s+1)\wedge\cdot}^{\gamma^\varepsilon}(\mathbf{x}) \right| \\ &= \sum_{\mathbf{x} \in \mathbb{X}_s, x \in \mathbb{S}} \left| \mu_{s\wedge\cdot}^\gamma(\mathbf{x}) \int_{\mathbb{A}} q(s, \mathbf{x}, \mu^\gamma, a; x) \gamma(s, \mathbf{x}, da) - \mu_{s\wedge\cdot}^{\gamma^\varepsilon}(\mathbf{x}) \int_{\mathbb{A}} q(s, \mathbf{x}, \mu^{\gamma^\varepsilon}, a; x) \gamma^\varepsilon(s, \mathbf{x}, da) \right| \\ &\leq \sum_{\mathbf{x} \in \mathbb{X}_s, x \in \mathbb{S}} \left[\left| \mu_{s\wedge\cdot}^\gamma(\mathbf{x}) - \mu_{s\wedge\cdot}^{\gamma^\varepsilon}(\mathbf{x}) \right| + \sum_{k=1}^{n_\varepsilon} \int_{A_k} \left| q(s, \mathbf{x}, \mu^\gamma, a; x) - q(s, \mathbf{x}, \mu^{\gamma^\varepsilon}, a_k; x) \right| \gamma(s, \mathbf{x}, da) \right. \\ &\quad \left. + \int_{A_0} q(s, \mathbf{x}, \mu^\gamma, a; x) \gamma(s, \mathbf{x}, da) + \int_{A_0} q(s, \mathbf{x}, \mu^{\gamma^\varepsilon}, a; x) \gamma^\varepsilon(s, \mathbf{x}, da) \right] \\ &\leq C\kappa_s + C\varepsilon. \end{aligned}$$

Then by induction we have (5.24).

We next show by induction that, recalling (5.12),

$$\bar{\kappa}_s := W_1(\nu_{s\wedge\cdot}^N, \mu_{s\wedge\cdot}^{\gamma^\varepsilon}) \leq C_\varepsilon \theta_N, \quad s = t, \dots, T. \quad (5.25)$$

Indeed, $\bar{\kappa}_t = W_1(\mu_{t,\tilde{\mathbf{x}}}^N, \mu)$. For $s \geq t$, noting that $\alpha_i \in \mathcal{A}_{path}^{t,\varepsilon} \subset \mathcal{A}_{path}^{t,0}$ and recalling from Lemma 4.6 that $\mu^{\Lambda^{\gamma^\varepsilon}} = \mu^{\gamma^\varepsilon}$, then by (5.12) and (4.13) that

$$\begin{aligned} \bar{\kappa}_{s+1} &= W_1(\nu_{s+1\wedge\cdot}^N, \mu_{(s+1)\wedge\cdot}^{\Lambda^{\gamma^\varepsilon}}) \\ &= \sum_{\mathbf{x} \in \mathbb{X}_t} \sum_{\tilde{\mathbf{x}} \in \mathbb{X}_{s+1}^{t,\mathbf{x}}} \left| \frac{1}{N} \sum_{\alpha \in \mathcal{A}_{path}^{t,\varepsilon}} \sum_{i \in I(\mathbf{x}, \alpha)} Q_{s+1}^t(\nu^N; \tilde{\mathbf{x}}, \alpha) - \int_{\mathcal{A}_{path}^t} Q_{s+1}^t(\mu^{\gamma^\varepsilon}; \tilde{\mathbf{x}}, \alpha) \Lambda^{\gamma^\varepsilon}(\mathbf{x}, d\alpha) \right| \\ &= \sum_{\mathbf{x} \in \mathbb{X}_t} \sum_{\tilde{\mathbf{x}} \in \mathbb{X}_{s+1}^{t,\mathbf{x}}} \left| \sum_{\alpha \in \mathcal{A}_{path}^{t,\varepsilon}} [\Lambda_{t,\tilde{\mathbf{x}}}^\varepsilon(\mathbf{x}, \alpha) Q_{s+1}^t(\nu^N; \tilde{\mathbf{x}}, \alpha) - \Lambda^{\gamma^\varepsilon}(\mathbf{x}, \alpha) Q_{s+1}^t(\mu^{\gamma^\varepsilon}; \tilde{\mathbf{x}}, \alpha)] \right| \\ &\leq \sum_{\mathbf{x} \in \mathbb{X}_t} \sum_{\tilde{\mathbf{x}} \in \mathbb{X}_{s+1}^{t,\mathbf{x}}} \sum_{\alpha \in \mathcal{A}_{path}^{t,\varepsilon}} \left[|\Lambda_{t,\tilde{\mathbf{x}}}^\varepsilon(\mathbf{x}, \alpha) - \Lambda^{\gamma^\varepsilon}(\mathbf{x}, \alpha)| Q_{s+1}^t(\nu^N; \tilde{\mathbf{x}}, \alpha) \right. \\ &\quad \left. + \Lambda^{\gamma^\varepsilon}(\mathbf{x}, \alpha) |Q_{s+1}^t(\nu^N; \tilde{\mathbf{x}}, \alpha) - Q_{s+1}^t(\mu^{\gamma^\varepsilon}; \tilde{\mathbf{x}}, \alpha)| \right]. \end{aligned}$$

Then, by (5.19) and noting that $C_\varepsilon := |\mathcal{A}_{path}^{t,\varepsilon}|$ is independent of N , we have

$$\begin{aligned} \bar{\kappa}_{s+1} &\leq \sum_{\mathbf{x} \in \mathbb{X}_t} \sum_{\tilde{\mathbf{x}} \in \mathbb{X}_{s+1}^{t,\mathbf{x}}} \sum_{\alpha \in \mathcal{A}_{path}^{t,\varepsilon}} \left[\theta_N Q_{s+1}^t(\nu^N; \tilde{\mathbf{x}}, \alpha) + C \Lambda^{\gamma^\varepsilon}(\mathbf{x}, \alpha) \sum_{r=t}^s W_1(\nu_{r\wedge\cdot}^N, \mu_{r\wedge\cdot}^{\gamma^\varepsilon}) \right] \\ &\leq C_\varepsilon \theta_N + C \sum_{r=t}^s \bar{\kappa}_r. \end{aligned}$$

This implies (5.25) immediately.

Finally, combining (5.24), (5.25), and (5.13), we obtain (5.21).

(ii) First, similar to (5.15), by (5.21) we have (5.22) following from the arguments in Theorem 3.4. Next, for $\gamma \in \mathcal{M}_{relax}^\varepsilon(t, \mu)$, by (4.19) we have $\Lambda^\gamma \in \mathcal{M}_{global}^\varepsilon(t, \mu)$. Then (5.23) follows from similar arguments as those for (5.16). \blacksquare

Proof of Theorem 5.3: the right inclusion. Fix $\varphi \in \mathbb{V}_{relax}(t, \mu)$ and $\varepsilon > 0$. Let $\varepsilon_1 > 0$ be a small number which will be specified later. There exists $\gamma \in \mathcal{M}_{relax}^{\varepsilon_1}(t, \mu)$ such that $\|\varphi - J(t, \mu, \gamma; \cdot, \gamma)\|_{\mathbb{X}_t} \leq \varepsilon_1$. Let γ^{ε_1} and $\vec{\alpha}$ be constructed as above. By (5.23) we have

$$\frac{1}{N} \sum_{i=1}^N [J_i(t, \vec{\mathbf{x}}, \vec{\alpha}) - v_i^{N,0}(t, \vec{\mathbf{x}}, \vec{\alpha})] \leq \varepsilon_1 + 2\rho_0(C\varepsilon_1 + C_{\varepsilon_1}\theta_N), \quad \forall \mathbf{x} \in \mathbb{X}_t.$$

Choose ε_1 small enough such that $\varepsilon_1 + 2\rho_0(C\varepsilon_1 + \varepsilon_1) < \varepsilon$. Then, for all N large enough such that $\theta_N \leq \frac{\varepsilon_1}{C_{\varepsilon_1}}$, we have $\frac{1}{N} \sum_{i=1}^N [J_i(t, \vec{\mathbf{x}}, \vec{\alpha}) - v_i^{N,0}(t, \vec{\mathbf{x}}, \vec{\alpha})] \leq \varepsilon$. That is, $\vec{\alpha} \in \mathbb{V}_{hetero}^{N,\varepsilon,0}(t, \mu_{t,\vec{\mathbf{x}}}^N)$

for all N large enough. Then, following the same arguments as those in the proof for the left inclusion, we can easily get $\varphi \in \mathbb{V}_{hetero}^{N,\varepsilon,0}(t, \mu_{t,\bar{x}}^N)$ for all N large enough, and thus $\varphi \in \underline{\lim}_{N \rightarrow \infty} \mathbb{V}_{hetero}^{N,\varepsilon,0}(t, \mu_{t,\bar{x}}^N)$. Since $\varepsilon > 0$ is arbitrary, we get the desired inclusion. \blacksquare

6 A diffusion model with state dependent drift controls

In this section we study a diffusion model with closed loop drift controls, where the laws of the controlled state process are all equivalent. The volatility control case involves mutually singular measures (corresponding to degenerate q in the discrete setting) and is much more challenging. We shall leave that for future research. We will consider state dependent controls only, and to ensure the convergence we consider homogeneous controls for the N -player games, as we did in Section 3.

6.1 The mean field game and the dynamic programming principle

Let $T > 0$ be a fixed terminal time, $(\Omega, \mathcal{F}, \mathbb{F} = \{\mathcal{F}_t\}_{0 \leq t \leq T}, \mathbb{P})$ a filtered probability space where \mathcal{F}_0 is atomless; B a d -dimensional Brownian motion; and the set $\mathbb{A} \subset \mathbb{R}^{d_0}$ a Borel measurable set. The state process X will also take values in \mathbb{R}^d . Its law lies in the space $\mathcal{P}_2 := \mathcal{P}_2(\mathbb{R}^d)$ equipped with the 2-Wasserstein distance W_2 . We remark that in the finite state space case W_1 and W_2 are equivalent, while in diffusion models they are not. In fact, at below we shall require W_1 -regularity, which is stronger than the W_2 -regularity, and obtain W_1 -convergence, which is weaker than the W_2 -convergence. This is not surprising in the mean field literature, see, e.g. [36]. The main advantage of the W_1 -distance is the following well known representation, see e.g. [13]: for any $\mu, \tilde{\mu} \in \mathcal{P}_1(\mathbb{R}^d)$,

$$W_1(\mu, \tilde{\mu}) = \sup \left\{ \int_{\mathbb{R}^d} \varphi(x) [\mu(dx) - \tilde{\mu}(dx)] : \varphi \in C_{Lip}(\mathbb{R}^d) \text{ s.t. } |\varphi(x) - \varphi(\tilde{x})| \leq |x - \tilde{x}| \right\}. \quad (6.1)$$

Here $C_{Lip}(\mathbb{R}^d)$ denote the set of uniformly Lipschitz continuous functions $\varphi : \mathbb{R}^d \rightarrow \mathbb{R}$. Moreover, for each $(t, \mu) \in [0, T] \times \mathcal{P}_2$, let $\mathbb{L}^2(t, \mu)$ denote the set of \mathcal{F}_t -measurable random variables ξ whose law (under \mathbb{P}) $\mathcal{L}_\xi = \mu$.

We consider coefficients $(b, f) : [0, T] \times \mathbb{R}^d \times \mathcal{P}_2 \times \mathbb{A} \rightarrow (\mathbb{R}^d, \mathbb{R})$ and $g : \mathbb{R}^d \times \mathcal{P}_2 \rightarrow \mathbb{R}$. Throughout this section, the following assumptions will always be in force.

Assumption 6.1 (i) b, f, g are Borel measurable in t and bounded by C_0 (for simplicity);
(ii) b, f, g are uniformly Lipschitz continuous in (x, μ, a) with a Lipschitz constant L_0 , where the Lipschitz continuity in μ is under W_1 .

Let \mathcal{A}_{cont} denote the set of admissible controls $\alpha : [0, T] \times \mathbb{R}^d \rightarrow \mathbb{A}$ which is measurable in t and Lipschitz continuous in x , with the Lipschitz constant L_α possibly depending on α . Given $(t, \mu) \in [0, T] \times \mathcal{P}_2$, $\xi \in \mathbb{L}^2(t, \mu)$, and $\alpha \in \mathcal{A}_{cont}$, consider the McKean-Vlasov SDE:

$$X_s^{t, \xi, \alpha} = \xi + \int_t^s b(r, X_r^{t, \xi, \alpha}, \mu_r^\alpha, \alpha(r, X_r^{t, \xi, \alpha})) dr + B_s - B_t, \quad \mu_s^\alpha := \mathcal{L}_{X_s^{t, \xi, \alpha}}. \quad (6.2)$$

By the required Lipschitz continuity, the above SDE is wellposed, and it is obvious that $\mu_t^\alpha = \mu$ and μ_s^α does not depend on the choice of $\xi \in \mathbb{L}^2(t, \mu)$. Then, when only the law is involved, by abusing the notations we may also denote $X^{t, \xi, \alpha}$ as $X^{t, \mu, \alpha}$.

Next, for any $x \in \mathbb{R}^d$, and $\tilde{\alpha} \in \mathcal{A}_{cont}$, we introduce

$$\begin{aligned} J(t, \mu, \alpha; x, \tilde{\alpha}) &:= J(\mu^\alpha; t, x, \tilde{\alpha}), \quad v(\mu^\alpha; s, x) := \inf_{\tilde{\alpha} \in \mathcal{A}_{cont}} J(\mu^\alpha; s, x, \tilde{\alpha}), \quad s \geq t, \quad \text{where} \\ X_r^{\mu^\alpha; s, x, \tilde{\alpha}} &= x + \int_s^r b(l, X_l^{\mu^\alpha; s, x, \tilde{\alpha}}, \mu_l^\alpha, \tilde{\alpha}(l, X_l^{\mu^\alpha; s, x, \tilde{\alpha}})) dl + B_r - B_s, \quad r \geq s; \\ J(\mu^\alpha; s, x, \tilde{\alpha}) &:= \mathbb{E} \left[g(X_T^{\mu^\alpha; s, x, \tilde{\alpha}}, \mu_T^\alpha) + \int_s^T f(r, X_r^{\mu^\alpha; s, x, \tilde{\alpha}}, \mu_r^\alpha, \tilde{\alpha}(r, X_r^{\mu^\alpha; s, x, \tilde{\alpha}})) dr \right]. \end{aligned} \quad (6.3)$$

Here we abuse the notations by using the same notations as in the discrete setting. Clearly $u(s, x) := J(\mu^\alpha; s, x, \tilde{\alpha})$ and $v(s, x) := v(\mu^\alpha; s, x)$ satisfy the following linear PDE and standard HJB equation on $[t, T] \times \mathbb{R}^d$, respectively, with parameter μ^α :

$$\begin{aligned} \partial_s u(s, x) + \frac{1}{2} \text{tr}(\partial_{xx} u(s, x)) + b(s, x, \mu_s^\alpha, \tilde{\alpha}(s, x)) \cdot \partial_x u(s, x) + f(s, x, \mu_s^\alpha, \tilde{\alpha}(s, x)) &= 0; \\ \partial_t v(s, x) + \frac{1}{2} \text{tr}(\partial_{xx} v(s, x)) + \inf_{a \in \mathbb{A}} [b(s, x, \mu_s^\alpha, a) \cdot \partial_x v(s, x) + f(s, x, \mu_s^\alpha, a)] &= 0; \\ u(T, x) = v(T, x) = g(x, \mu_T^\alpha). \end{aligned} \quad (6.4)$$

Definition 6.2 Fix $(t, \mu) \in [0, T] \times \mathcal{P}_2$. For any $\varepsilon > 0$, we say $\alpha^* \in \mathcal{A}_{cont}$ is an ε -MFE at (t, μ) , denoted as $\alpha^* \in \mathcal{M}_{cont}^\varepsilon(t, \mu)$, if

$$\int_{\mathbb{R}^d} [J(t, \mu, \alpha^*; x, \alpha^*) - v(\mu^{\alpha^*}; t, x)] \mu(dx) \leq \varepsilon. \quad (6.5)$$

Similar to (5.4) and (5.5), here we do not require α^* to be optimal for every player x . Indeed, since there are infinitely many players, we shall tolerate that a small portion of players may not be happy for the α^* , as in [11], and their possible deviation from α^* won't change the equilibrium measure μ^{α^*} significantly.

To define the set value, we need the following simple but crucial regularity result, whose proof is postponed to Appendix.

Lemma 6.3 Let Assumption 6.1 hold. There exists a constant $C > 0$, depending only on T, d, C_0, L_0 , such that, for any $t, \mu, \alpha, \tilde{\alpha}$ and $s \geq t$,

$$|J(\mu^\alpha; \tilde{\alpha}, s, x) - J(\mu^\alpha; \tilde{\alpha}, s, \tilde{x})| + |v(\mu^\alpha; s, x) - v(\mu^\alpha; s, \tilde{x})| \leq C|x - \tilde{x}|, \quad \forall x, \tilde{x}. \quad (6.6)$$

We then define the set value of the mean field game:

$$\begin{aligned} \mathbb{V}_{cont}(t, \mu) &:= \bigcap_{\varepsilon > 0} \mathbb{V}_{cont}^\varepsilon(t, \mu), \quad \text{where} \\ \mathbb{V}_{cont}^\varepsilon(t, \mu) &:= \left\{ \varphi \in C_{Lip}(\mathbb{R}^d) : \text{there exists } \alpha^* \in \mathcal{M}_{cont}^\varepsilon(t, \mu) \text{ such that} \right. \\ &\quad \left. \int_{\mathbb{R}^d} |\varphi(x) - J(t, \mu, \alpha^*; x, \alpha^*)| \mu(dx) \leq \varepsilon \right\}. \end{aligned} \quad (6.7)$$

In particular, since $J(t, \mu, \alpha^*; x, \alpha^*) \geq v(\mu^{\alpha^*}; t, x)$, then by (6.6) and (6.5) we see that both $J(t, \mu, \alpha^*; \cdot, \alpha^*)$ and $v(\mu^{\alpha^*}; t, \cdot)$ belong to $\mathbb{V}_{cont}(t, \mu)$. Moreover, again due to (6.5), we may replace the inequality in the last line of (6.7) with $\int_{\mathbb{R}^d} |\varphi(x) - v(\mu^{\alpha^*}; t, x)| \mu(dx) \leq \varepsilon$.

Similarly, given T_0 and $\psi \in C_{Lip}(\mathbb{R}^d)$, we may define the functions $J(T_0, \psi; t, \mu, \alpha; x, \tilde{\alpha})$, $J(T_0, \psi; \mu^\alpha; s, x, \tilde{\alpha})$, $v(T_0, \psi; \mu^\alpha; s, x)$, as well as the sets $\mathcal{M}_{cont}^\varepsilon(T_0, \psi; t, \mu)$, $\mathbb{V}_{cont}^\varepsilon(T_0, \psi; t, \mu)$, $\mathbb{V}_{cont}(T_0, \psi; t, \mu)$ in the obvious sense. In particular, we have the following tower property:

$$\begin{aligned} J(t, \mu, \alpha; x, \tilde{\alpha}) &= J(T_0, \psi; t, \mu, \alpha; x, \tilde{\alpha}), \quad \text{where } \psi(x) := J(T_0, \mu_{T_0}^\alpha, \alpha; x, \tilde{\alpha}); \\ v(\mu^\alpha; t, x) &= v(T_0, \tilde{\psi}; \mu^\alpha; t, x), \quad \text{where } \tilde{\psi}(x) := v(\mu^\alpha; T_0, x). \end{aligned} \quad (6.8)$$

We now establish the DPP for $\mathbb{V}_{cont}(t, \mu)$.

Theorem 6.4 *Let Assumption 6.1 hold. For any $0 \leq t \leq T_0 \leq T$ and $\mu \in \mathcal{P}_2$, it holds that*

$$\begin{aligned} \mathbb{V}_{cont}(t, \mu) &= \tilde{\mathbb{V}}_{cont}(t, \mu) := \bigcap_{\varepsilon > 0} \tilde{\mathbb{V}}_{cont}^\varepsilon(t, \mu), \quad \text{where} \\ \tilde{\mathbb{V}}_{cont}^\varepsilon(t, \mu) &:= \left\{ \varphi \in C_{Lip}(\mathbb{R}^d) : \int_{\mathbb{R}^d} |\varphi(x) - J(T_0, \psi; t, \mu, \alpha^*; x, \alpha^*)| \mu(dx) \leq \varepsilon, \right. \\ &\quad \left. \text{for some } (\psi, \alpha^*) \text{ satisfying: } \psi \in \mathbb{V}_{cont}^\varepsilon(T_0, \mu_{T_0}^{\alpha^*}), \alpha^* \in \mathcal{M}_{cont}^\varepsilon(T_0, \psi; t, \mu) \right\}. \end{aligned} \quad (6.9)$$

Proof (i) We first prove $\mathbb{V}_{cont}(t, \mu) \subset \tilde{\mathbb{V}}_{cont}(t, \mu)$. Fix $\varphi \in \mathbb{V}_{cont}(t, \mu)$, $\varepsilon > 0$, and set $\varepsilon_1 := \frac{\varepsilon}{2}$. Since $\varphi \in \mathbb{V}_{cont}^{\varepsilon_1}(t, \mu)$, there exists $\alpha^* \in \mathcal{M}_{cont}^{\varepsilon_1}(t, \mu)$ satisfying (6.7) for ε_1 . Denote

$$\psi(x) := J(T_0, \mu_{T_0}^{\alpha^*}, \alpha^*; x, \alpha^*), \quad \tilde{\psi}(x) := v(\mu^{\alpha^*}; T_0, x).$$

By (6.8) we have $J(T_0, \psi; t, \mu, \alpha^*; x, \alpha^*) = J(t, \mu, \alpha^*; x, \alpha^*)$ and thus

$$\int_{\mathbb{R}^d} |\varphi(x) - J(T_0, \psi; t, \mu, \alpha^*; x, \alpha^*)| \mu(dx) \leq \varepsilon_1 \leq \varepsilon.$$

We shall show that $\psi \in \mathbb{V}_{cont}^\varepsilon(T_0, \mu_{T_0}^{\alpha^*})$ and $\alpha^* \in \mathcal{M}_{cont}^\varepsilon(T_0, \psi; t, \mu)$. Then $\varphi \in \tilde{\mathbb{V}}_{cont}^\varepsilon(t, \mu)$, and therefore, since $\varepsilon > 0$ is arbitrary, we have $\varphi \in \tilde{\mathbb{V}}(t, \mu)$.

Step 1. In this step we show that

$$\int_{\mathbb{R}^d} [J(T_0, \mu_{T_0}^{\alpha^*}, \alpha^*; x, \alpha^*) - v(\mu^{\alpha^*}; T_0, x)] \mu_{T_0}^{\alpha^*}(dx) = \int_{\mathbb{R}^d} [\psi(x) - \tilde{\psi}(x)] \mu_{T_0}^{\alpha^*}(dx) \leq \varepsilon_1. \quad (6.10)$$

Then $\alpha^* \in \mathcal{M}_{cont}^\varepsilon(T_0, \mu_{T_0}^{\alpha^*})$, which, together with the regularity of ψ from Lemma 6.3, implies immediately that $\psi \in \mathbb{V}_{cont}^\varepsilon(T_0, \mu_{T_0}^{\alpha^*})$.

To see this, we recall (6.2) with $\xi \in \mathbb{L}^2(t, \mu)$. Since $\alpha^* \in \mathcal{M}_{cont}^{\varepsilon_1}(t, \mu)$, by (6.8) we have

$$\begin{aligned} \varepsilon_1 &\geq \mathbb{E} \left[J(t, \mu, \alpha^*; \xi, \alpha^*) - v(\mu^{\alpha^*}; t, \xi) \right] = \mathbb{E} \left[J(T_0, \psi; t, \mu, \alpha^*; \xi, \alpha^*) - v(T_0, \tilde{\psi}; \mu^{\alpha^*}; t, \xi) \right] \\ &\geq \mathbb{E} \left[J(T_0, \psi; t, \mu, \alpha^*; \xi, \alpha^*) - J(T_0, \tilde{\psi}; t, \mu, \alpha^*; \xi, \alpha^*) \right] = \mathbb{E} \left[\psi(X_{T_0}^{t, \xi, \alpha^*}) - \tilde{\psi}(X_{T_0}^{t, \xi, \alpha^*}) \right]. \end{aligned}$$

Note that $\mathcal{L}_{X_{T_0}^{t, \xi, \alpha^*}} = \mu_{T_0}^{\alpha^*}$, then this is exactly (6.10).

Step 2. It remains to show that $\alpha^* \in \mathcal{M}_{cont}^\varepsilon(T_0, \psi; t, \mu)$. By the definition of v and its regularity from Lemma 6.3, there exists $\tilde{\alpha}^* \in \mathcal{A}_{cont}$ such that

$$J(T_0, \psi; t, \mu, \alpha^*; x, \tilde{\alpha}^*) \leq v(T_0, \psi; \mu^{\alpha^*}; t, x) + \varepsilon_1, \quad \forall x \in \mathbb{R}^d.$$

Then, denoting $\hat{\alpha}^* := \tilde{\alpha}^* \oplus_{T_0} \alpha^* \in \mathcal{A}_{cont}$, by (6.8) again we have

$$\begin{aligned} &\mathbb{E} \left[J(T_0, \psi; t, \mu, \alpha^*; \xi, \alpha^*) - v(T_0, \psi; \mu^{\alpha^*}; t, \xi) \right] \\ &\leq \mathbb{E} \left[J(T_0, \psi; t, \mu, \alpha^*; \xi, \alpha^*) - J(T_0, \psi; t, \mu, \alpha^*; \xi, \tilde{\alpha}^*) \right] + \varepsilon_1 \\ &= \mathbb{E} \left[J(t, \mu, \alpha^*; \xi, \alpha^*) - J(t, \mu, \alpha^*; \xi, \hat{\alpha}^*) \right] + \varepsilon_1 \\ &\leq \mathbb{E} \left[J(t, \mu, \alpha^*; \xi, \alpha^*) - v(\mu^{\alpha^*}; t, \xi) \right] + \varepsilon_1 \leq \varepsilon_1 + \varepsilon_1 = \varepsilon, \end{aligned}$$

This means $\alpha^* \in \mathcal{M}_{cont}^\varepsilon(T_0, \psi; t, \mu)$.

(ii) We next prove $\tilde{\mathbb{V}}_{cont}(t, \mu) \subset \mathbb{V}_{cont}(t, \mu)$. Fix $\varphi \in \tilde{\mathbb{V}}_{cont}(t, \mu)$, $\varepsilon > 0$, and set $\varepsilon_1 := \frac{\varepsilon}{4}$. Since $\varphi \in \tilde{\mathbb{V}}_{cont}^{\varepsilon_1}(t, \mu)$, there exist (ψ, α^*) satisfying the desired properties in (6.9) for ε_1 . In particular, since $\psi \in \mathbb{V}_{cont}^{\varepsilon_1}(T_0, \mu_{T_0}^{\alpha^*})$, there exists desired $\tilde{\alpha}^* \in \mathcal{M}_{cont}^{\varepsilon_1}(T_0, \mu_{T_0}^{\alpha^*})$ required in (6.7) for ε_1 . Denote $\hat{\alpha}^* := \alpha^* \oplus_{T_0} \tilde{\alpha}^* \in \mathcal{A}_{cont}$ and

$$\hat{\psi}(x) := J(T_0, \mu_{T_0}^{\alpha^*}, \tilde{\alpha}^*; x, \tilde{\alpha}^*), \quad \tilde{\psi}(x) := v(\mu^{\hat{\alpha}^*}; T_0, x).$$

By (6.9),

$$\begin{aligned} &\mathbb{E} \left[\left| J(T_0, \psi; t, \mu, \alpha^*; \xi, \alpha^*) - J(T_0, \hat{\psi}; t, \mu, \alpha^*; \xi, \alpha^*) \right| \right] \tag{6.11} \\ &= \mathbb{E} \left[\left| \psi(X_{T_0}^{\mu^{\alpha^*}; t, \xi, \alpha^*}) - \hat{\psi}(X_{T_0}^{\mu^{\alpha^*}; t, \xi, \alpha^*}) \right| \right] = \int_{\mathbb{R}^d} |\psi(x) - J(T_0, \mu_{T_0}^{\alpha^*}, \tilde{\alpha}^*; x, \tilde{\alpha}^*)| \mu_{T_0}^{\alpha^*}(dx) \leq \varepsilon_1 \end{aligned}$$

Then, since $\varphi \in \tilde{\mathbb{V}}_{cont}^{\varepsilon_1}(t, \mu)$ with corresponding (ψ, α^*) , by (6.8) and (6.11) we have

$$\mathbb{E} \left[\left| \varphi(\xi) - J(t, \mu, \hat{\alpha}^*; \xi, \hat{\alpha}^*) \right| \right] \leq \mathbb{E} \left[\left| \varphi(\xi) - J(T_0, \psi; t, \mu, \alpha^*; \xi, \alpha^*) \right| \right] + \varepsilon_1 \leq 2\varepsilon_1 \leq \varepsilon,$$

where $\xi \in \mathbb{L}^2(t, \mu)$. We claim further that $\hat{\alpha}^* \in \mathcal{M}_{cont}^\varepsilon(t, \mu)$. Then $\varphi \in \mathbb{V}_{cont}^\varepsilon(t, \mu)$, and thus $\varphi \in \mathbb{V}_{cont}(t, \mu)$, since $\varepsilon > 0$ is arbitrary.

To see the claim, since $\alpha^* \in \mathcal{M}_{cont}^{\varepsilon_1}(T_0, \psi; t, \mu)$, $\tilde{\alpha}^* \in \mathcal{M}_{cont}^{\varepsilon_1}(T_0, \mu_{T_0}^{\alpha^*})$, by (6.8) we have

$$\begin{aligned}
& \mathbb{E} \left[J(t, \mu, \hat{\alpha}^*; \xi, \hat{\alpha}^*) - v(\mu^{\hat{\alpha}^*}; t, \xi) \right] \\
&= \mathbb{E} \left[J(T_0, \hat{\psi}; t, \mu, \alpha^*; \xi, \alpha^*) - v(T_0, \tilde{\psi}; \mu^{\alpha^*}; t, \xi) \right] \\
&\leq \mathbb{E} \left[J(T_0, \psi; t, \mu, \alpha^*; \xi, \alpha^*) - v(T_0, \tilde{\psi}; \mu^{\alpha^*}; t, \xi) \right] + \varepsilon_1 \\
&\leq \mathbb{E} \left[v(T_0, \psi; \mu^{\alpha^*}; t, \xi) - v(T_0, \tilde{\psi}; \mu^{\alpha^*}; t, \xi) \right] + 2\varepsilon_1 \\
&\leq \sup_{\tilde{\alpha} \in \mathcal{A}_{cont}} \mathbb{E} \left[J(T_0, \psi; t, \mu, \alpha^*; \xi, \tilde{\alpha}) - J(T_0, \tilde{\psi}; t, \mu, \alpha^*; \xi, \tilde{\alpha}) \right] + 2\varepsilon_1 \\
&= \mathbb{E} \left[\psi(X_{T_0}^{t, \xi, \alpha^*}) - \tilde{\psi}(X_{T_0}^{t, \xi, \alpha^*}) \right] + 2\varepsilon_1 \leq \mathbb{E} \left[\hat{\psi}(X_{T_0}^{t, \xi, \alpha^*}) - \tilde{\psi}(X_{T_0}^{t, \xi, \alpha^*}) \right] + 3\varepsilon_1 \leq \varepsilon_1 + 3\varepsilon_1 = \varepsilon.
\end{aligned}$$

This means $\hat{\alpha}^* \in \mathcal{M}_{cont}^\varepsilon(t, \mu)$, and hence completes the proof. \blacksquare

6.2 Convergence of the N -player game

By enlarging the filtered probability space $(\Omega, \mathcal{F}, \mathbb{F}, \mathbb{P})$, if necessary, we let B^1, \dots, B^N be independent d -dimensional Brownian motions on it. Set $\mathcal{A}_{cont}^\infty := \cup_{L \geq 0} \mathcal{A}_{cont}^L$, where, for each $L \geq 0$, \mathcal{A}_{cont}^L denotes the set of admissible controls $\alpha : [0, T] \times \mathbb{R}^d \times \mathcal{P}_2 \rightarrow \mathbb{A}$ such that

$$|\alpha(t, x, \mu) - \alpha(t, \tilde{x}, \tilde{\mu})| \leq L_\alpha |x - \tilde{x}| + LW_1(\mu, \tilde{\mu}).$$

Here the Lipschitz constant L_α may depend on α , hence the Lipschitz continuity in x is not uniform in α . We emphasize that the Lipschitz continuity in μ is under W_1 , rather than W_2 , so that we can use the representation (6.1). Note that $\mathcal{A}_{cont} = \mathcal{A}_{cont}^0$, and by Remark 3.1 (i), all the results in the previous subsection remain true if we replace \mathcal{A}_{cont} with $\mathcal{A}_{cont}^\infty$.

Given $t \in [0, T]$, $\vec{x} = (x_1, \dots, x_N) \in \mathbb{R}^{dN}$ and $\vec{\alpha} = (\alpha_1, \dots, \alpha_N) \in (\mathcal{A}_{cont}^L)^N$, consider

$$\begin{aligned}
X_s^{t, \vec{x}, \vec{\alpha}; i} &= x_i + \int_t^s b(r, X_r^{t, \vec{x}, \vec{\alpha}; i}, \mu_r^{t, \vec{x}, \vec{\alpha}}, \alpha_i(r, X_r^{t, \vec{x}, \vec{\alpha}; i}, \mu_r^{t, \vec{x}, \vec{\alpha}})) dr + B_s^i - B_t^i, \quad i = 1, \dots, N; \\
&\text{where } \mu_s^{t, \vec{x}, \vec{\alpha}} := \frac{1}{N} \sum_{i=1}^N \delta_{X_s^{t, \vec{x}, \vec{\alpha}; i}}; \\
J_i(t, \vec{x}, \vec{\alpha}) &:= \mathbb{E} \left[g(X_T^{t, \vec{x}, \vec{\alpha}; i}, \mu_T^{t, \vec{x}, \vec{\alpha}}) + \int_t^T f(s, X_s^{t, \vec{x}, \vec{\alpha}; i}, \mu_s^{t, \vec{x}, \vec{\alpha}}, \alpha_i(s, X_s^{t, \vec{x}, \vec{\alpha}; i}, \mu_s^{t, \vec{x}, \vec{\alpha}})) ds \right], \\
v_i^{N, L}(t, \vec{x}, \vec{\alpha}) &:= \inf_{\tilde{\alpha} \in \mathcal{A}_{cont}^L} J_i(t, \vec{x}, (\vec{\alpha}^{-i}, \tilde{\alpha})).
\end{aligned} \tag{6.12}$$

In light of Lemma 6.3, the following regularity result is interesting in its own right. However, since it will not be used for our main result, we postpone its proof to Appendix.

Proposition 6.5 *Let Assumption 6.1 hold. For any $L \geq 0$, there exists a constant $C_L > 0$, depending only on T, d, C_0, L_0 , and L , such that, for any $(t, \vec{x}) \in [0, T] \times \mathbb{R}^{dN}$, $\bar{x}, \tilde{x} \in \mathbb{R}^d$, and $\vec{\alpha} \in (\mathcal{A}_{cont}^L)^N$, we have*

$$|v_i^{N,L}(t, (\vec{x}^{-i}, \bar{x}), \vec{\alpha}) - v_i^{N,L}(t, (\vec{x}^{-i}, \tilde{x}), \vec{\alpha})| \leq C_L |\bar{x} - \tilde{x}|, \quad i = 1, \dots, N. \quad (6.13)$$

Given $\alpha \in \mathcal{A}_{cont}^L$, by viewing it as the homogeneous control (α, \dots, α) , we may use the simplified notations $X^{t, \vec{x}, \alpha; i}$, $\mu^{t, \vec{x}, \alpha}$, $J_i(t, \vec{x}, \alpha)$, and $v_i^{N,L}(t, \vec{x}, \alpha)$ in the obvious sense.

Definition 6.6 (i) *Fix $(t, \vec{x}) \in [0, T] \times \mathbb{R}^{dN}$, $\varepsilon > 0$, $L \geq 0$. We say $\alpha^* \in \mathcal{A}_{cont}^L$ is a homogeneous (ε, L) -equilibrium of the N -player game at (t, \vec{x}) , denoted as $\alpha^* \in \mathcal{M}_{cont}^{N, \varepsilon, L}(t, \vec{x})$, if*

$$\frac{1}{N} \sum_{i=1}^N [J_i(t, \vec{x}, \alpha^*) - v_i^{N,L}(t, \vec{x}, \alpha^*)] \leq \varepsilon. \quad (6.14)$$

(ii) *The set value for the N -player game is defined as:*

$$\mathbb{V}_{cont}^N(t, \vec{x}) := \bigcap_{\varepsilon > 0} \mathbb{V}_{cont}^{N, \varepsilon}(t, \vec{x}) := \bigcap_{\varepsilon > 0} \bigcup_{L \geq 0} \mathbb{V}_{cont}^{N, \varepsilon, L}(t, \vec{x}), \quad \text{where} \quad (6.15)$$

$$\mathbb{V}_{cont}^{N, \varepsilon, L}(t, \vec{x}) := \left\{ \varphi \in C_{Lip}(\mathbb{R}^d) : \exists \alpha^* \in \mathcal{M}_{cont}^{N, \varepsilon, L}(t, \vec{x}) \text{ s.t. } \frac{1}{N} \sum_{i=1}^N |\varphi(x_i) - J_i(t, \vec{x}, \alpha^*)| \leq \varepsilon \right\}.$$

We remark that, although $\mathbb{V}_{cont}^{N, \varepsilon, L}(t, \vec{x})$ involves only the values $\{\varphi(x_i)\}_{1 \leq i \leq N}$, for the convenience of the convergence analysis we consider its elements as $\varphi \in C_{Lip}(\mathbb{R}^d)$.

Remark 6.7 (i) *Recall (3.1). By the required symmetry, obviously there exist functions $J^N, v^{N,L} : [0, T] \times \mathcal{P}_2 \times \mathcal{A}_{cont}^L \times \mathbb{R}^d \rightarrow \mathbb{R}$ such that*

$$J_i(t, \vec{x}, \alpha) = J^N(t, \mu_{\vec{x}}^N, \alpha; x_i), \quad v_i^{N,L}(t, \vec{x}, \alpha) = v^{N,L}(t, \mu_{\vec{x}}^N, \alpha; x_i), \quad i = 1, \dots, N. \quad (6.16)$$

Moreover, $\mathbb{V}_{cont}^N(t, \vec{x})$ is invariant in $\mu_{\vec{x}}^N$ and thus can be denoted as $\mathbb{V}_{cont}^N(t, \mu_{\vec{x}}^N)$.

(ii) *The required inequalities in Definition 6.6 are equivalent to:*

$$\int_{\mathbb{R}^d} [J^N - v^{N,L}](t, \mu_{\vec{x}}^N, \alpha^*; x) \mu_{\vec{x}}^N(dx) \leq \varepsilon, \quad \int_{\mathbb{R}^d} [\varphi(x) - J^N(t, \mu_{\vec{x}}^N, \alpha^*; x)] \mu_{\vec{x}}^N(dx) \leq \varepsilon.$$

We now turn to the convergence, starting with the convergence of the equilibrium measures. Recall the vector $(\alpha, \tilde{\alpha})_i$ introduced in (3.6).

Theorem 6.8 *Let Assumption 6.1 hold. For any $L \geq 0$, there exists a constant $C_L > 0$, depending only on T, d, C_0, L_0 , and L , such that, for any $t \in [0, T]$, $\vec{x} \in \mathbb{R}^{dN}$, $\mu \in \mathcal{P}_2$, $\alpha, \tilde{\alpha} \in \mathcal{A}_{cont}^L$, and $i = 1, \dots, N$,*

$$\sup_{t \leq s \leq T} \mathbb{E} \left[W_1(\mu_s^{t, \vec{x}, (\alpha, \tilde{\alpha})^i}, \mu_s^\alpha) \right] \leq C_L \theta_N, \quad (6.17)$$

$$\text{where } \theta_N := W_1(\mu_{\vec{x}}^N, \mu) + N^{-\frac{1}{d\vee 3}} \|\vec{x}\|_2 + N^{-1}, \quad \|\vec{x}\|_2^2 := \frac{1}{N} \sum_{i=1}^N |x_i|^2.$$

Proof Recall (6.12) and introduce, for $j = 1, \dots, N$,

$$\begin{aligned} \tilde{X}_s^j &= x_j + \int_t^s b(r, \tilde{X}_r^j, \mu_r^\alpha, \alpha(r, \tilde{X}_r^j, \mu_r^\alpha)) dr + B_s^j - B_t^j, \quad \tilde{\mu}_s^N := \frac{1}{N} \sum_{j=1}^N \delta_{\tilde{X}_s^j}; \\ \tilde{X}_s &= \tilde{\xi} + \int_t^s b(r, \tilde{X}_r, \mu_r^\alpha, \alpha(r, \tilde{X}_r, \mu_r^\alpha)) dr + B_s - B_t, \quad \text{where } \tilde{\xi} \in \mathbb{L}^2(\mathcal{F}_0; \mu_{\vec{x}}^N). \end{aligned} \quad (6.18)$$

Note that $\tilde{X}^1, \dots, \tilde{X}^N$ are independent. We proceed the rest of the proof in two steps.

Step 1. In this step we estimate $\mathbb{E}[W_1(\tilde{\mu}_s^N, \mu_s^\alpha)]$. First, by [36, Lemma 8.4] we have

$$\mathbb{E}[W_1(\tilde{\mu}_s^N, \mathcal{L}_{\tilde{X}_s})] \leq CN^{-\frac{1}{d\vee 3}} \|\vec{x}\|_2.$$

Next, fix an φ in (6.1) and let $u = u_\varphi$ denote the solution to the following PDE on $[t, s]$:

$$\partial_r u + \frac{1}{2} \text{tr}(\partial_{xx} u) + b(r, x, \mu_s^\alpha, \alpha(r, x, \mu_s^\alpha)) \cdot \partial_x u = 0, \quad u(s, x) = \varphi(x). \quad (6.19)$$

Applying Lemma 6.3 with $\tilde{\alpha}(r, x) := \alpha(r, x, \mu_r^\alpha)$ and $f = 0$, we see that u is uniformly Lipschitz continuous in x , with a Lipschitz constant C independent of φ and L . Thus,

$$\mathbb{E}[\varphi(\tilde{X}_s) - \varphi(X_s^\alpha)] = \mathbb{E}[u(t, \tilde{\xi}) - u(t, \xi)] \leq C \mathbb{E}[|\tilde{\xi} - \xi|].$$

Since \mathcal{F}_0 is atomless, we may choose $\xi, \tilde{\xi}$ such that $\mathbb{E}[|\tilde{\xi} - \xi|] = W_1(\mu_{\vec{x}}^N, \mu)$, then (6.1) implies $W_1(\mathcal{L}_{\tilde{X}_s}, \mu_s^\alpha) \leq CW_1(\mu_{\vec{x}}^N, \mu)$. Put together, we have

$$\mathbb{E}[W_1(\tilde{\mu}_s^N, \mu_s^\alpha)] \leq CW_1(\mu_{\vec{x}}^N, \mu) + CN^{-\frac{1}{d\vee 3}} \|\vec{x}\|_2 \leq C\theta_N, \quad t \leq s \leq T. \quad (6.20)$$

Step 2. We next estimate $\mathbb{E}[W_1(\mu_s^{t, \vec{x}, (\alpha, \tilde{\alpha})^i}, \mu_s^\alpha)]$. Denote $\alpha_i := \tilde{\alpha}$, $\alpha_j := \alpha$ for $j \neq i$, and

$$\begin{aligned} \beta_r^j &:= b(s, \tilde{X}_s^j, \tilde{\mu}_s^N, \alpha_j(s, \tilde{X}_s^j, \tilde{\mu}_s^N)) - b(s, \tilde{X}_s^j, \mu_s^\alpha, \alpha(s, \tilde{X}_s^j, \mu_s^\alpha)), \quad 1 \leq j \leq N \\ M_s &:= \prod_{j=1}^N M_s^j, \quad M_s^j := \exp \left(\int_t^s \beta_r^j dB_r^j - \frac{1}{2} \int_t^s |\beta_r^j|^2 dr \right). \end{aligned}$$

Then, by the Girsanov theorem we have

$$\begin{aligned} \mathbb{E}[W_1(\mu_s^{t, \vec{x}, (\alpha, \tilde{\alpha})^i}, \mu_s^\alpha)] &= \mathbb{E}[M_s W_1(\tilde{\mu}_s^N, \mu_s^\alpha)] = \mathbb{E}[(M_s - 1)W_1(\tilde{\mu}_s^N, \mu_s^\alpha)] + \mathbb{E}[W_1(\tilde{\mu}_s^N, \mu_s^\alpha)] \\ &= \sum_{j=1}^N \mathbb{E} \left[\int_t^s M_r \beta_r^j dB_r^j W_1(\tilde{\mu}_s^N, \mu_s^\alpha) \right] + \mathbb{E}[W_1(\tilde{\mu}_s^N, \mu_s^\alpha)]. \end{aligned} \quad (6.21)$$

By the martingale representation theorem, we have

$$W_1(\tilde{\mu}_s^N, \mu_s^\alpha) = \mathbb{E}[W_1(\tilde{\mu}_s^N, \mu_s^\alpha)] + \sum_{j=1}^N \int_t^s Z_r^j dB_r^j. \quad (6.22)$$

Note that \tilde{X}^j are independent. Consider the following linear PDE on $[t, s] \times \mathbb{R}^{dN}$:

$$\begin{aligned} \partial_r u(r, \vec{x}') + \frac{1}{2} \sum_{j=1}^N \text{tr}(\partial_{x_j x_j} u(r, \vec{x}')) + \sum_{j=1}^N b(r, x'_j, \mu_s^\alpha, \alpha(r, x'_j, \mu_r^\alpha)) \cdot \partial_{x_j} u(r, \vec{x}') = 0, \\ u(s, \vec{x}') = W_1(\mu_{\vec{x}'}^N, \mu_s^\alpha). \end{aligned} \quad (6.23)$$

By standard BSDE theory, see e.g. [41, Chapter 5], we have $Z_r^j = \partial_{x_j} u(r, \vec{X}_r^{t, \vec{x}})$, where $X_r^{t, \vec{x}, j} := x_j + B_r^j - B_t^j$. Note that the terminal condition $u(s, \vec{x}')$ is Lipschitz continuous in x'_j with Lipschitz constant $\frac{1}{N}$. Then, similarly to (6.19), by Lemma 6.3 we see that $|Z^j| \leq |\partial_{x_j} u| \leq \frac{C}{N}$ for some constant C independent of α and L . Thus, by (6.21) and (6.22),

$$\mathbb{E}\left[W_1(\mu_s^{t, \vec{x}, (\alpha, \tilde{\alpha})_i}, \mu_s^\alpha) - W_1(\tilde{\mu}_s^N, \mu_s^\alpha)\right] = \sum_{j=1}^N \mathbb{E}\left[\int_t^s M_r \beta_r^j \cdot Z_r^j dr\right] \leq \frac{C}{N} \sum_{j=1}^N \mathbb{E}\left[\int_t^s M_r |\beta_r^j| dr\right].$$

Note that $|\beta^i| \leq C$ and, for $j \neq i$, $|\beta_r^j| \leq C_L W_1(\tilde{\mu}_r^N, \mu_r^\alpha)$. Then, by (6.20),

$$\begin{aligned} \mathbb{E}\left[W_1(\mu_s^{t, \vec{x}, (\alpha, \tilde{\alpha})_i}, \mu_s^\alpha)\right] &\leq \mathbb{E}\left[W_1(\tilde{\mu}_s^N, \mu_s^\alpha)\right] + \frac{C}{N} \mathbb{E}\left[\int_t^s M_r |\beta_r^i| dr + \sum_{j \neq i} \int_t^s M_r |\beta_r^j| dr\right] \\ &\leq \mathbb{E}\left[W_1(\tilde{\mu}_s^N, \mu_s^\alpha)\right] + \frac{C}{N} + \frac{C_L}{N} \sum_{j \neq i} \mathbb{E}\left[\int_t^s M_r W_1(\tilde{\mu}_r^N, \mu_r^\alpha) dr\right] = \frac{C}{N} + C_L \theta_N \leq C_L \theta_N, \end{aligned}$$

proving (6.17). ■

Theorem 6.9 *For the setting in Theorem 6.8, we have*

$$\left|J_i(t, \vec{x}, (\alpha, \tilde{\alpha})_i) - J(t, \mu, \alpha; x_i, \tilde{\alpha})\right| + \left|v_i^{N, L}(t, \vec{x}, \alpha) - v(\mu^\alpha; t, x_i)\right| \leq C_L \theta_N^{\frac{1}{4}}. \quad (6.24)$$

Proof Fix i . First, by taking supremum over $\tilde{\alpha} \in \mathcal{A}_{cont}^L$, the uniform estimate for J implies that for v immediately. So it suffices to prove the former estimate.

For this purpose, recall (6.12) and denote

$$\tilde{J}_i(t, \vec{x}, (\alpha, \tilde{\alpha})_i) := \mathbb{E}^{\mathbb{P}}\left[g(X_T^{t, \vec{x}, (\alpha, \tilde{\alpha})_i; i}, \mu_T^\alpha) + \int_t^T f(s, X_s^{t, \vec{x}, (\alpha, \tilde{\alpha})_i; i}, \mu_s^\alpha, \tilde{\alpha}(s, X_s^{t, \vec{x}, (\alpha, \tilde{\alpha})_i; i}, \mu_s^\alpha)) ds\right].$$

Then one can easily see that, by applying Theorem 6.8,

$$\left|J_i(t, \vec{x}, (\alpha, \tilde{\alpha})_i) - \tilde{J}_i(t, \vec{x}, (\alpha, \tilde{\alpha})_i)\right| \leq C_L \sup_{t \leq s \leq T} \mathbb{E}\left[W_1(\mu_s^{t, \vec{x}, (\alpha, \tilde{\alpha})_i}, \mu_s^\alpha)\right] \leq C_L \theta_N. \quad (6.25)$$

Next, denote

$$\begin{aligned} X_s^i &:= x_i + B_s^i - B_t^i, \quad \tilde{\mu}_s^{N,i} := \frac{1}{N} \left[\sum_{j \neq i} \delta_{X_s^{t,\vec{x},(\alpha,\tilde{\alpha})_i;j}} + \delta_{X_s^i} \right]; \\ \beta_s &:= b(s, X_s^i, \mu_s^\alpha, \tilde{\alpha}(s, X_s^i, \mu_s^\alpha)), \quad M_s := \exp \left(\int_t^s \beta_r dB_r^i - \frac{1}{2} \int_t^s |\beta_r|^2 dr \right); \\ \tilde{\beta}_s &:= b(s, X_s^i, \tilde{\mu}_s^{N,i}, \tilde{\alpha}(s, X_s^i, \tilde{\mu}_s^{N,i})), \quad \tilde{M}_s := \exp \left(\int_t^s \tilde{\beta}_r dB_r^i - \frac{1}{2} \int_t^s |\tilde{\beta}_r|^2 dr \right). \end{aligned}$$

By (6.3) and (6.12), it follows from the Girsanov theorem again that

$$\begin{aligned} & \left| \tilde{J}_i(t, \vec{x}, (\alpha, \tilde{\alpha})_i) - J(t, \mu, \alpha; x_i, \tilde{\alpha}) \right| \tag{6.26} \\ &= \left| \mathbb{E} \left[[\tilde{M}_T - M_T] \left[g(X_T^i, \mu_T^\alpha) + \int_t^T f(s, X_s^i, \mu_s^\alpha, \tilde{\alpha}(s, X_s^i, \mu_s^\alpha)) ds \right] \right] \right| \leq C \mathbb{E} [|\tilde{M}_T - M_T|]. \end{aligned}$$

Denote $\Delta M_s := \tilde{M}_s - M_s$, $\Delta \beta_s := \tilde{\beta}_s - \beta_s$. Then, since b is bounded,

$$\begin{aligned} \mathbb{E} [|\Delta M_s|^2] &= \mathbb{E} \left[\left(\int_t^s [\tilde{M}_r \tilde{\beta}_r - M_r \beta_r] dB_r^i \right)^2 \right] = \mathbb{E} \left[\int_t^s [\tilde{M}_r \tilde{\beta}_r - M_r \beta_r]^2 dr \right] \\ &\leq C \int_t^s \mathbb{E} [|\Delta M_r|^2] dr + C \mathbb{E} \left[\int_t^s |\tilde{M}_r|^2 |\Delta \beta_r|^2 dr \right] \\ &\leq C \int_t^s \mathbb{E} [|\Delta M_r|^2] dr + C \mathbb{E} \left[\int_t^s \tilde{M}_r^{\frac{3}{2}} \tilde{M}_r^{\frac{1}{2}} |\Delta \beta_r|^{\frac{1}{2}} dr \right] \\ &\leq C \int_t^s \mathbb{E} [|\Delta M_r|^2] dr + C \left(\mathbb{E} \left[\int_t^s \tilde{M}_r |\Delta \beta_r| dr \right] \right)^{\frac{1}{2}} \\ &\leq C \int_t^s \mathbb{E} [|\Delta M_r|^2] dr + C_L \left(\mathbb{E} \left[\int_t^s \tilde{M}_r W_1(\tilde{\mu}_r^{N,i}, \mu_r^\alpha) dr \right] \right)^{\frac{1}{2}} \\ &= C \int_t^s \mathbb{E} [|\Delta M_r|^2] dr + C_L \left(\mathbb{E} \left[\int_t^s W_1(\mu_r^{t,\vec{x},(\alpha,\tilde{\alpha})_i}, \mu_r^\alpha) dr \right] \right)^{\frac{1}{2}} \\ &\leq C \int_t^s \mathbb{E} [|\Delta M_r|^2] dr + C_L \theta_N^{\frac{1}{2}}, \end{aligned}$$

where the last inequality thanks to Theorem 6.8. Then, by the Gronwall inequality we obtain $\mathbb{E} [|\Delta M_s|^2] \leq C_L \theta_N^{\frac{1}{2}}$, and thus (6.26) implies

$$\left| \tilde{J}_i(t, \vec{x}, (\alpha, \tilde{\alpha})_i) - J(t, \mu, \alpha; x_i, \tilde{\alpha}) \right| \leq C_L \theta_N^{\frac{1}{4}}.$$

This, together with (6.25), implies the estimate for J in (6.24) immediately. \blacksquare

Theorem 6.10 *Let Assumption 6.1 hold. Assume further that $\lim_{N \rightarrow \infty} W_1(\mu_{\vec{x}}^N, \mu) = 0$, and there exists a constant $C > 0$ such that³ $\|\vec{x}\|_2 \leq C$ for all N . Then*

$$\bigcap_{\varepsilon > 0} \bigcup_{L \geq 0} \overline{\lim}_{N \rightarrow \infty} \mathbb{V}_{cont}^{N,\varepsilon,L}(t, \mu_{\vec{x}}^N) \subset \mathbb{V}_{cont}(t, \mu) \subset \bigcap_{\varepsilon > 0} \underline{\lim}_{N \rightarrow \infty} \mathbb{V}_{cont}^{N,\varepsilon,0}(t, \mu_{\vec{x}}^N) \tag{6.27}$$

In particular, since $\underline{\lim}_{N \rightarrow \infty} \mathbb{V}_{cont}^{N,\varepsilon,0}(t, \mu_{\vec{x}}^N) \subset \bigcup_{L \geq 0} \overline{\lim}_{N \rightarrow \infty} \mathbb{V}_{cont}^{N,\varepsilon,L}(t, \mu_{\vec{x}}^N)$, actually equalities hold.

³Note again that \vec{x} depends on N . Also, the conditions here are slightly weaker than $\lim_{N \rightarrow \infty} W_2(\mu_{\vec{x}}^N, \mu) = 0$.

Proof (i) We first prove the right inclusion in (6.27). Fix $\varphi \in \mathbb{V}_{cont}(t, \mu)$, $\varepsilon > 0$, and set $\varepsilon_1 := \frac{\varepsilon}{2}$. By (6.7) and (6.5), there exists $\alpha^* \in \mathcal{M}_{cont}^{\varepsilon_1}(t, \mu)$ such that

$$\int_{\mathbb{R}^d} [J(t, \mu, \alpha^*; x, \alpha^*) - v(\mu^{\alpha^*}; t, x)] \mu(dx) \leq \varepsilon_1, \quad \int_{\mathbb{R}^d} |\varphi(x) - J(t, \mu, \alpha^*; x, \alpha^*)| \mu(dx) \leq \varepsilon_1.$$

Recall Lemma 6.3 and note that $\varphi \in C_{Lip}(\mathbb{R}^d)$, then by (6.1) we have

$$\begin{aligned} \int_{\mathbb{R}^d} [J(t, \mu, \alpha^*; x, \alpha^*) - v(\mu^{\alpha^*}; t, x)] \mu_{\vec{x}}^N(dx) &\leq \varepsilon_1 + CW_1(\mu_{\vec{x}}^N, \mu), \\ \int_{\mathbb{R}^d} |\varphi(x) - J(t, \mu, \alpha^*; x, \alpha^*)| \mu_{\vec{x}}^N(dx) &\leq \varepsilon_1 + C_\varphi W_1(\mu_{\vec{x}}^N, \mu), \end{aligned}$$

where C_φ may depend on the Lipschitz constant of φ . Moreover, by (6.24) we have

$$\begin{aligned} \frac{1}{N} \sum_{i=1}^N [J_i(t, \vec{x}, \alpha^*) - v_i^{N,L}(t, \vec{x}, \alpha^*)] &\leq \frac{1}{N} \sum_{i=1}^N [J(t, \mu, \alpha^*; x_i, \alpha^*) - v(\mu^{\alpha^*}; t, x_i)] + C_L \theta_N^{\frac{1}{4}} \\ &= \int_{\mathbb{R}^d} [J(t, \mu, \alpha^*; x, \alpha^*) - v(\mu^{\alpha^*}; t, x)] \mu_{\vec{x}}^N(dx) + C_L \theta_N^{\frac{1}{4}} \leq \varepsilon_1 + C_L \theta_N^{\frac{1}{4}}; \\ \frac{1}{N} \sum_{i=1}^N |\varphi(x_i) - J_i(t, \vec{x}, \alpha^*)| &\leq \frac{1}{N} \sum_{i=1}^N |\varphi(x_i) - J(t, \mu, \alpha^*; x_i, \alpha^*)| + C_L \theta_N^{\frac{1}{4}} \\ &= \int_{\mathbb{R}^d} |\varphi(x) - J(t, \mu, \alpha^*; x, \alpha^*)| \mu_{\vec{x}}^N(dx) + C_{L,\varphi} \theta_N^{\frac{1}{4}}. \end{aligned}$$

We emphasize again that $\|\vec{x}\|_2 \leq C$ is independent of N . Then, by choosing N large enough such that $C_L \theta_N^{\frac{1}{4}} \leq \varepsilon_1$, $C_{L,\varphi} \theta_N^{\frac{1}{4}} \leq \varepsilon_1$, we obtain

$$\frac{1}{N} \sum_{i=1}^N [J_i(t, \vec{x}, \alpha^*) - v_i^{N,L}(t, \vec{x}, \alpha^*)] \leq \varepsilon; \quad \frac{1}{N} \sum_{i=1}^N |\varphi(x_i) - J_i(t, \vec{x}, \alpha^*)| \leq \varepsilon.$$

This implies that $\alpha^* \in \mathcal{M}_{cont}^{N,\varepsilon,0}(t, \vec{x})$ and $\varphi \in \mathbb{V}_{cont}^{N,\varepsilon,0}(t, \mu_{\vec{x}}^N)$, for all N large enough. That is, $\varphi \in \varinjlim_{N \rightarrow \infty} \mathbb{V}_{cont}^{N,\varepsilon,0}(t, \vec{x})$ for any $\varepsilon > 0$.

(ii) We next show the left inclusion in (6.27). Fix $\varphi \in \bigcap_{\varepsilon > 0} \bigcup_{L \geq 0} \overline{\lim}_{N \rightarrow \infty} \mathbb{V}_{cont}^{N,\varepsilon,L}(t, \mu_{\vec{x}}^N)$, $\varepsilon > 0$, and set $\varepsilon_1 := \frac{\varepsilon}{2}$. There exist $L_\varepsilon \geq 0$ and an infinite sequence $\{N_k\}_{k \geq 1}$ such that $\varphi \in \mathbb{V}_{cont}^{N_k, \varepsilon_1, L_\varepsilon}(t, \mu_{\vec{x}}^{N_k})$ for all $k \geq 1$. Recall (6.14) and (6.15), there exists $\alpha^k \in \mathcal{A}_{cont}^{L_\varepsilon}$ such that

$$\frac{1}{N_k} \sum_{i=1}^{N_k} [J_i(t, \vec{x}, \alpha^k) - v_i^{N_k, L_\varepsilon}(t, \vec{x}, \alpha^k)] \leq \varepsilon_1; \quad \frac{1}{N_k} \sum_{i=1}^{N_k} |\varphi(x_i) - J_i(t, \vec{x}, \alpha^k)| \leq \varepsilon_1.$$

Note that L_ε is fixed, in particular it is independent of k . In light of Remark 3.1 (i) and denote $\tilde{\alpha}^k(s, x) := \alpha^k(s, x, \mu^{\alpha^k})$, then $\mu^{\tilde{\alpha}^k} = \mu^{\alpha^k}$. Similarly to (i), by (6.24) we have

$$\begin{aligned} \int_{\mathbb{R}^d} [J(t, \mu, \tilde{\alpha}^k; x, \tilde{\alpha}^k) - v(\mu^{\alpha^k}; t, x)] \mu_{\vec{x}}^{N_k}(dx) &\leq \varepsilon_1 + C_{L_\varepsilon} \theta_{N_k}^{\frac{1}{4}}, \\ \int_{\mathbb{R}^d} |\varphi(x) - J(t, \mu, \tilde{\alpha}^k; x, \tilde{\alpha}^k)| \mu_{\vec{x}}^{N_k}(dx) &\leq \varepsilon_1 + C_{L_\varepsilon} \theta_{N_k}^{\frac{1}{4}}. \end{aligned}$$

Then, by Lemma 6.3 and (6.1) we have

$$\begin{aligned} \int_{\mathbb{R}^d} [J(t, \mu, \tilde{\alpha}^k; x, \tilde{\alpha}^k) - v(\mu^{\alpha^k}; t, x)] \mu(dx) &\leq \varepsilon_1 + C_{L_\varepsilon} \theta_{N_k}^{\frac{1}{4}} + CW_1(\mu_{\tilde{x}}^{N_k}, \mu), \\ \int_{\mathbb{R}^d} |\varphi(x) - J(t, \mu, \tilde{\alpha}^k; x, \tilde{\alpha}^k)| \mu(dx) &\leq \varepsilon_1 + C_{L_\varepsilon} \theta_{N_k}^{\frac{1}{4}} + C_\varphi W_1(\mu_{\tilde{x}}^{N_k}, \mu). \end{aligned}$$

Now choose k large enough (possibly depending on ε and φ) such that

$$C_{L_\varepsilon} \theta_{N_k}^{\frac{1}{4}} + CW_1(\mu_{\tilde{x}}^{N_k}, \mu) \leq \varepsilon_1, \quad C_{L_\varepsilon} \theta_{N_k}^{\frac{1}{4}} + C_\varphi W_1(\mu_{\tilde{x}}^{N_k}, \mu) \leq \varepsilon_1.$$

Then we have

$$\int_{\mathbb{R}^d} [J(t, \mu, \tilde{\alpha}^k; x, \tilde{\alpha}^k) - v(\mu^{\alpha^k}; t, x)] \mu(dx) \leq \varepsilon, \quad \int_{\mathbb{R}^d} |\varphi(x) - J(t, \mu, \tilde{\alpha}^k; x, \tilde{\alpha}^k)| \mu(dx) \leq \varepsilon.$$

This implies that $\tilde{\alpha}^k \in \mathcal{M}_{cont}^\varepsilon(t, \mu)$ and $\varphi \in \mathbb{V}_{cont}^\varepsilon(t, \mu)$. Since $\varepsilon > 0$ is arbitrary, we obtain $\varphi \in \mathbb{V}_{cont}(t, \mu)$. ■

Remark 6.11 *In the same manner as in Remark 3.6, we have the convergence of the equilibria as well in the diffusion model of this section.*

7 Appendix

7.1 The subtle path dependence issue in Remark 4.3

In this subsection we elaborate Remark 4.3 (ii) and (iii). Throughout the subsection, q, F, G are state dependent as in Section 2. For simplicity, we compare state dependent controls and path dependent controls only for the raw set values, and we set $t = 0$ and hence $\mu \in \mathcal{P}_0(\mathbb{X}_0) = \mathcal{P}_0(\mathbb{S})$.

We first provide a counterexample to show that the raw set value $\mathbb{V}_0(0, \mu)$ is in general not equal to the corresponding raw set value $\mathbb{V}_{0,path}(0, \mu)$ with controls $\alpha \in \mathcal{A}_{path}$.

Example 7.1 *Set $T = 2$, $\mathbb{S} = \{\underline{x}, \bar{x}\}$, $\mathbb{A} = [a_0, 1 - a_0]$ for some $0 < a_0 < \frac{1}{2}$, and*

$$\begin{aligned} q(0, x, \mu, a; \underline{x}) = q(0, x, \mu, a; \bar{x}) &= \frac{1}{2}, \quad q(1, x, \mu, a; \underline{x}) = a, \quad q(1, x, \mu, a; \bar{x}) = 1 - a; \\ F(0, x, \mu, a) &= 0, \quad F(1, x, \mu, a) = a[1 - a], \quad G(x, \mu) = \mu(\underline{x}). \end{aligned}$$

Then $\mathbb{V}_0(0, \mu) \neq \mathbb{V}_{0,path}(0, \mu)$ for general $\mu \in \mathcal{P}_0(\mathbb{S})$.

Proof Since $|\mathbb{S}| = 2$, for any $\mu \in \mathcal{P}_0(\mathbb{S})$ clearly it suffices to specify $\mu(\underline{x})$. Fix $\mu \in \mathcal{P}_0(\mathbb{S})$.

We first compute $\mathbb{V}_0(0, \mu)$. For any $\alpha, \tilde{\alpha} \in \mathcal{A}_{state}$, it is straightforward to compute:

$$\begin{aligned}
\mu_1^\alpha(\underline{x}) &= \sum_{x_0 \in \mathbb{S}} \mu(x_0) q(0, x_0, \mu, \alpha(0, x_0); \underline{x}) = \sum_{x_0 \in \mathbb{S}} \mu(x_0) \frac{1}{2} = \frac{1}{2}; \\
\mu_2^\alpha(\underline{x}) &= \sum_{x_1 \in \mathbb{S}} \mu_1^\alpha(x_1) q(1, x_1, \mu_1^\alpha, \alpha(1, x_1); \underline{x}) = \frac{1}{2} \sum_{x_1 \in \mathbb{S}} \alpha(1, x_1); \\
\mathbb{P}^{\mu^\alpha; 0, x_0, \tilde{\alpha}}(X_1 = \underline{x}) &= q(0, x_0, \mu, \alpha(0, x_0); \underline{x}) = \frac{1}{2}; \\
J(0, \mu, \alpha; x_0, \tilde{\alpha}) &= \mathbb{E}^{\mathbb{P}^{\mu^\alpha; 0, x_0, \tilde{\alpha}}} \left[G(X_2, \mu_2^\alpha) + \sum_{t=0,1} F(t, X_t, \mu_t^\alpha, \tilde{\alpha}(t, X_t)) \right] \\
&= \mu_2^\alpha(\underline{x}) + \mathbb{E}^{\mathbb{P}^{\mu^\alpha; 0, x_0, \tilde{\alpha}}} \left[\tilde{\alpha}(1, X_1) [1 - \tilde{\alpha}(1, X_1)] \right] \\
&= \frac{1}{2} \sum_{x_1 \in \mathbb{S}} \alpha(1, x_1) + \frac{1}{2} \sum_{x_1 \in \mathbb{S}} \tilde{\alpha}(1, x_1) [1 - \tilde{\alpha}(1, x_1)]
\end{aligned}$$

Given α , we see that $\tilde{\alpha}$ is a minimizer if and only if $\tilde{\alpha}(1, x_1) \in \{a_0, 1 - a_0\}, \forall x_1 \in \mathbb{S}$. Thus, $\alpha^* \in \mathcal{M}_{state}(0, \mu)$ if and only if $\alpha^*(1, x_1) \in \{a_0, 1 - a_0\}, \forall x_1 \in \mathbb{S}$, and

$$J(0, \mu, \alpha^*; x_0, \alpha^*) = \frac{1}{2} \sum_{x_1 \in \mathbb{S}} \alpha^*(1, x_1) [2 - \alpha^*(1, x_1)]$$

is independent of x_0 (and μ). This implies that $\mathbb{V}_0(0, \mu)$ consists of 3 constant values:

$$\mathbb{V}_0(0, \mu) = \left\{ a_0[2 - a_0], \quad \frac{1}{2} + a_0 - a_0^2, \quad 1 - a_0^2 \right\}. \quad (7.1)$$

We next compute $\mathbb{V}_{0,path}(0, \mu)$. For any $\alpha, \tilde{\alpha} \in \mathcal{A}_{path}$, we still have $\mu_1^\alpha(\underline{x}) = \frac{1}{2}$ and $\mathbb{P}^{\mu^\alpha; 0, x_0, \tilde{\alpha}}(X_1 = \underline{x}) = \frac{1}{2}$. Moreover,

$$\begin{aligned}
\mu_2^\alpha(\underline{x}) &= \sum_{x_0, x_1 \in \mathbb{S}} \mu(x_0) q(0, x_0, \mu, \alpha(0, x_0); x_1) q(1, x_1, \mu_1^\alpha, \alpha(1, x_0, x_1); \underline{x}) \\
&= \frac{1}{2} \sum_{x_0, x_1 \in \mathbb{S}} \mu(x_0) \alpha(1, x_0, x_1); \\
J(0, \mu, \alpha; x_0, \tilde{\alpha}) &= \mathbb{E}^{\mathbb{P}^{\mu^\alpha; 0, x_0, \tilde{\alpha}}} \left[G(X_2, \mu_2^\alpha) + F(1, X_1, \mu_1^\alpha, \tilde{\alpha}(1, X_0, X_1)) \right] \\
&= \mu_2^\alpha(\underline{x}) + \mathbb{E}^{\mathbb{P}^{\mu^\alpha; 0, x_0, \tilde{\alpha}}} \left[\tilde{\alpha}(1, X_0, X_1) [1 - \tilde{\alpha}(1, X_0, X_1)] \right] \\
&= \frac{1}{2} \sum_{\tilde{x}_0, x_1 \in \mathbb{S}} \mu(\tilde{x}_0) \alpha(1, \tilde{x}_0, x_1) + \frac{1}{2} \sum_{x_1 \in \mathbb{S}} \tilde{\alpha}(1, x_0, x_1) [1 - \tilde{\alpha}(1, x_0, x_1)].
\end{aligned}$$

Similarly, $\alpha^* \in \mathcal{A}_{path}$ is an MFE if and only if $\alpha^*(1, x_0, x_1) \in \{a_0, 1 - a_0\}, \forall x_0, x_1 \in \mathbb{S}$, and

$$J(0, \mu, \alpha^*; x_0, \alpha^*) = \frac{1}{2} \sum_{\tilde{x}_0, x_1 \in \mathbb{S}} \mu(\tilde{x}_0) \alpha^*(1, \tilde{x}_0, x_1) + \frac{1}{2} \sum_{x_1 \in \mathbb{S}} \alpha^*(1, x_0, x_1) [1 - \alpha^*(1, x_0, x_1)].$$

Choose one particular α^* :

$$\alpha^*(1, \underline{x}, x_1) = a_0, \quad \alpha^*(1, \bar{x}, x_1) = 1 - a_0, \quad \forall x_1 \in \mathbb{S}.$$

Then

$$\begin{aligned} J(0, \mu, \alpha^*; \underline{x}, \alpha^*) &= \frac{1}{2} \left[\sum_{x_1 \in \mathbb{S}} \mu(\underline{x}) \alpha^*(1, \underline{x}, x_1) + \sum_{x_1 \in \mathbb{S}} \mu(\bar{x}) \alpha^*(1, \bar{x}, x_1) \right] \\ &\quad + \frac{1}{2} \sum_{x_1 \in \mathbb{S}} \alpha^*(1, \underline{x}, x_1) [1 - \alpha^*(1, \underline{x}, x_1)] \\ &= \mu(\underline{x}) a_0 + \mu(\bar{x}) [1 - a_0] + a_0 [1 - a_0], \end{aligned}$$

and similarly,

$$J(0, \mu, \alpha^*; \bar{x}, \alpha^*) = \mu(\underline{x}) a_0 + \mu(\bar{x}) [1 - a_0] + [1 - a_0] a_0 = J(0, \mu, \alpha^*; \underline{x}, \alpha^*).$$

So $\mathbb{V}_{0, \text{path}}(0, \mu)$ contains the constant value $\mu(\underline{x}) a_0 + \mu(\bar{x}) [1 - a_0] + [1 - a_0] a_0$, which depends on μ and is in general not in the $\mathbb{V}_0(0, \mu)$ in (7.1). \blacksquare

We next turn to relaxed controls. Let $\mathcal{A}_{\text{relax}}$ be the path dependent ones in Section 4, and $\mathcal{A}_{\text{relax}}^{\text{state}}$ denote the subset taking the form $\gamma(t, x, da)$. We emphasize again that here we are considering state dependent q, F, G . Fix $t = 0$ and $\mu \in \mathcal{P}_0(\mathbb{S})$.

Lemma 7.2 *For any $\gamma \in \mathcal{A}_{\text{relax}}$, define*

$$\tilde{\gamma}(s, x, da) := \frac{1}{\mu_s^{\tilde{\gamma}}(x)} \sum_{\mathbf{x} \in \mathbb{X}_s: \mathbf{x}_s = x} \mu_{s \wedge \cdot}^{\gamma}(\mathbf{x}) \gamma(s, \mathbf{x}, da), \quad \text{where } \mu_s^{\gamma}(x) := \sum_{\mathbf{x} \in \mathbb{X}_s: \mathbf{x}_s = x} \mu_{s \wedge \cdot}^{\gamma}(\mathbf{x}). \quad (7.2)$$

Then $\tilde{\gamma} \in \mathcal{A}_{\text{relax}}^{\text{state}}$ and $\mu_s^{\tilde{\gamma}} = \mu_s^{\gamma}$.

Proof First it is obvious that

$$\tilde{\gamma}(s, x, \mathbb{A}) = \frac{1}{\mu_s^{\tilde{\gamma}}(x)} \sum_{\mathbf{x} \in \mathbb{X}_s: \mathbf{x}_s = x} \mu_{s \wedge \cdot}^{\gamma}(\mathbf{x}) \gamma(s, \mathbf{x}, \mathbb{A}) = \frac{1}{\mu_s^{\gamma}(x)} \sum_{\mathbf{x} \in \mathbb{X}_s: \mathbf{x}_s = x} \mu_{s \wedge \cdot}^{\gamma}(\mathbf{x}) = 1,$$

so $\tilde{\gamma} \in \mathcal{A}_{\text{relax}}^{\text{state}}$. Next, by definition $\mu_0^{\tilde{\gamma}} = \mu = \mu_0^{\gamma}$. Assume $\mu_s^{\tilde{\gamma}} = \mu_s^{\gamma}$, then for $s + 1$,

$$\begin{aligned} \mu_{s+1}^{\tilde{\gamma}}(x) &= \sum_{\tilde{x} \in \mathbb{S}} \mu_s^{\tilde{\gamma}}(\tilde{x}) \int_{\mathbb{A}} q(s, \tilde{x}, \mu_s^{\tilde{\gamma}}, a; x) \tilde{\gamma}(s, \tilde{x}, da) \\ &= \sum_{\tilde{x} \in \mathbb{S}} \mu_s^{\gamma}(\tilde{x}) \int_{\mathbb{A}} q(s, \tilde{x}, \mu_s^{\gamma}, a; x) \frac{1}{\mu_s^{\tilde{\gamma}}(\tilde{x})} \sum_{\mathbf{x} \in \mathbb{X}_s: \mathbf{x}_s = \tilde{x}} \mu_{s \wedge \cdot}^{\gamma}(\mathbf{x}) \gamma(s, \mathbf{x}, da) \\ &= \sum_{\mathbf{x} \in \mathbb{X}_s} \mu_{s \wedge \cdot}^{\gamma}(\mathbf{x}) \int_{\mathbb{A}} q(s, \mathbf{x}_s, \mu_s^{\gamma}, a; x) \gamma(s, \mathbf{x}, da) = \mu_{s+1}^{\gamma}(x). \end{aligned}$$

This completes the induction argument. \blacksquare

Lemma 7.3 *If $\gamma^* \in \mathcal{A}_{relax}$ is a relaxed MFE at $(0, \mu)$, then the corresponding $\tilde{\gamma}^* \in \mathcal{A}_{relax}^{state}$ is a state dependent relaxed MFE at $(0, \mu)$. Moreover, in this case we have*

$$J(0, \mu, \gamma^*; x, \gamma^*) = J(0, \mu, \tilde{\gamma}^*; x, \tilde{\gamma}^*). \quad (7.3)$$

Proof First, by Lemma 7.2 it is straightforward to verify that

$$\int_{\mathbb{S}} J(0, \mu, \gamma; x, \gamma) \mu(dx) = \int_{\mathbb{S}} J(0, \mu, \tilde{\gamma}; x, \tilde{\gamma}) \mu(dx).$$

On the other hand, since $\gamma^* \in \mathcal{A}_{relax}$, by the standard control theory we have

$$\inf_{\gamma \in \mathcal{A}_{relax}} J(0, \mu, \gamma^*; x, \gamma) = v(\mu^{\gamma^*}; 0, x) = v(\mu^{\tilde{\gamma}^*}; 0, x) = \inf_{\gamma' \in \mathcal{A}_{relax}^{state}} J(0, \mu, \tilde{\gamma}^*; x, \gamma'). \quad (7.4)$$

Then

$$\int_{\mathbb{S}} J(0, \mu, \tilde{\gamma}^*; x, \tilde{\gamma}^*) \mu(dx) = \int_{\mathbb{S}} J(0, \mu, \gamma^*; x, \gamma^*) \mu(dx) = \int_{\mathbb{S}} v(\mu^{\tilde{\gamma}^*}; 0, x) \mu(dx).$$

Since $J(0, \mu, \tilde{\gamma}^*; x, \tilde{\gamma}^*) \geq v(\mu^{\tilde{\gamma}^*}; 0, x)$ and $\text{supp}(\mu) = \mathbb{S}$, then $J(0, \mu, \tilde{\gamma}^*; x, \tilde{\gamma}^*) = v(\mu^{\tilde{\gamma}^*}; 0, x)$ for all $x \in \mathbb{S}$. This implies that $\tilde{\gamma}^* \in \mathcal{A}_{relax}^{state}$ is a state dependent relaxed MFE at $(0, \mu)$, and consequently (7.4) leads to (7.3). \blacksquare

Theorem 7.4 *The MFGs with state dependent relaxed controls and path dependent relaxed controls have the same relaxed raw set value.*

Proof By Lemma 7.3, clearly the path dependent raw set value is included in the state dependent raw set value. On the other hand, for any state dependent relaxed control $\hat{\gamma}^* \in \mathcal{A}_{relax}^{state}$, we may still view $\gamma^* := \hat{\gamma}^*$ as a path dependent relaxed control⁴, and it is straightforward to verify that the $\tilde{\gamma}^* \in \mathcal{A}_{relax}^{state}$ corresponding to γ^* is equal to $\hat{\gamma}^*$. Then, following the arguments in Lemma 7.3, in particular (7.4), one can easily show that $J(0, \mu, \gamma^*; x, \gamma^*) = v(\mu^{\gamma^*}; 0, x)$ and thus γ^* is also an MFE among \mathcal{A}_{relax} . Therefore, $J(0, \mu, \gamma^*; \cdot, \gamma^*)$ belong to the path dependent raw set value as well. \blacksquare

7.2 Some postponed proofs

Proof of Theorem 2.7. Let $\tilde{\mathbb{V}}_{state}(t, \mu) = \bigcap_{\varepsilon > 0} \tilde{\mathbb{V}}_{state}^{\varepsilon}(t, \mu)$ denote the right side of (2.17) in the obvious sense. We shall follow the arguments in Theorem 2.4.

⁴While it is trivial that $\mathcal{A}_{relax}^{state} \subset \mathcal{A}_{relax}^{path} := \mathcal{A}_{relax}$, as stated here, in general it is not trivial that $\mathcal{M}_{relax}^{state} \subset \mathcal{M}_{relax}^{path}$, because for the latter one has to compare with other path dependent relax controls, which is a stronger requirement than that for $\mathcal{M}_{relax}^{state}$. The rest of the proof is exactly to prove $\mathcal{M}_{relax}^{state} \subset \mathcal{M}_{relax}^{path}$.

(i) We first prove $\tilde{\mathbb{V}}_{state}(t, \mu) \subset \mathbb{V}_{state}(t, \mu)$. Fix $\varphi \in \tilde{\mathbb{V}}_{state}(t, \mu)$, $\varepsilon > 0$, and set $\varepsilon_1 := \frac{\varepsilon}{4}$. Since $\varphi \in \tilde{\mathbb{V}}_{state}^{\varepsilon_1}(t, \mu)$, there exist desirable ψ and $\alpha^* \in \mathcal{M}_{state}^{\varepsilon_1}(T_0, \psi; t, \mu)$ as in (2.17), and the property $\psi(\cdot, \mu_{T_0}^{\alpha^*}) \in \mathbb{V}_{state}^{\varepsilon_1}(T_0, \mu_{T_0}^{\alpha^*})$ implies further that there exists $\tilde{\alpha}^* \in \mathcal{M}_{state}^{\varepsilon_1}(T_0, \mu_{T_0}^{\alpha^*})$ such that

$$\|\varphi - J(T_0, \psi; t, \mu, \alpha^*; \cdot, \alpha^*)\|_{\infty} \leq \varepsilon_1, \quad \|\psi(\cdot, \mu_{T_0}^{\alpha^*}) - J(T_0, \mu_{T_0}^{\alpha^*}, \tilde{\alpha}^*; \cdot, \tilde{\alpha}^*)\|_{\infty} \leq \varepsilon_1.$$

Denote $\hat{\alpha}^* := \alpha^* \oplus_{T_0} \tilde{\alpha}^* \in \mathcal{A}_{state}$. Then, for any $\alpha \in \mathcal{A}_{state}$ and $x \in \mathbb{S}$, similar to the arguments in Proposition 2.3 (i), we have

$$\begin{aligned} J(t, \mu, \hat{\alpha}^*; x, \alpha) &= \mathbb{E}^{\mathbb{P}^{\mu^{\alpha^*}; t, x, \alpha}} \left[J(T_0, \mu_{T_0}^{\alpha^*}, \tilde{\alpha}^*; X_{T_0}, \alpha) + \sum_{s=t}^{T_0-1} F(s, X_s, \mu_s^{\alpha^*}, \alpha(s, X_s)) \right] \\ &\geq \mathbb{E}^{\mathbb{P}^{\mu^{\alpha^*}; t, x, \alpha}} \left[J(T_0, \mu_{T_0}^{\alpha^*}, \tilde{\alpha}^*; X_{T_0}, \tilde{\alpha}^*) + \sum_{s=t}^{T_0-1} F(s, X_s, \mu_s^{\alpha^*}, \alpha(s, X_s)) \right] - \varepsilon_1 \\ &\geq \mathbb{E}^{\mathbb{P}^{\mu^{\alpha^*}; t, x, \alpha}} \left[\psi(X_{T_0}, \mu_{T_0}^{\alpha^*}) + \sum_{s=t}^{T_0-1} F(s, X_s, \mu_s^{\alpha^*}, \alpha(s, X_s)) \right] - 2\varepsilon_1 \\ &= J(T_0, \psi; t, \mu, \alpha^*; x, \alpha) - 2\varepsilon_1 \geq J(T_0, \psi; t, \mu, \alpha^*; x, \alpha^*) - 3\varepsilon_1 \\ &= \mathbb{E}^{\mathbb{P}^{\mu^{\alpha^*}; t, x, \alpha^*}} \left[\psi(X_{T_0}, \mu_{T_0}^{\alpha^*}) + \sum_{s=t}^{T_0-1} F(s, X_s, \mu_s^{\alpha^*}, \alpha^*(s, X_s)) \right] - 3\varepsilon_1 \\ &\geq \mathbb{E}^{\mathbb{P}^{\mu^{\alpha^*}; t, x, \alpha^*}} \left[J(T_0, \mu_{T_0}^{\alpha^*}, \tilde{\alpha}^*; X_{T_0}, \tilde{\alpha}^*) + \sum_{s=t}^{T_0-1} F(s, X_s, \mu_s^{\alpha^*}, \alpha^*(s, X_s)) \right] - 4\varepsilon_1 \\ &= J(t, \mu, \hat{\alpha}^*; x, \hat{\alpha}^*) - \varepsilon. \end{aligned}$$

That is, $\hat{\alpha}^* \in \mathcal{M}_{state}^{\varepsilon}(t, \mu)$. Moreover, note that

$$\begin{aligned} \|\varphi - J(t, \mu, \hat{\alpha}^*; \cdot, \hat{\alpha}^*)\|_{\infty} &\leq \varepsilon_1 + \|J(T_0, \psi; t, \mu, \alpha^*; \cdot, \alpha^*) - J(t, \mu, \hat{\alpha}^*; \cdot, \hat{\alpha}^*)\|_{\infty} \\ &= \varepsilon_1 + \sup_{x \in \mathbb{S}} \left| \mathbb{E}^{\mathbb{P}^{\mu^{\alpha^*}; t, x, \alpha^*}} \left[\psi(X_{T_0}, \mu_{T_0}^{\alpha^*}) - J(T_0, \mu_{T_0}^{\alpha^*}, \tilde{\alpha}^*; X_{T_0}, \tilde{\alpha}^*) \right] \right| \leq 2\varepsilon_1 \leq \varepsilon. \end{aligned}$$

Then $\varphi \in \mathbb{V}_{state}^{\varepsilon}(t, \mu)$. Since $\varepsilon > 0$ is arbitrary, we obtain $\varphi \in \mathbb{V}_{state}(t, \mu)$.

(ii) We now prove the opposite inclusion. Fix $\varphi \in \mathbb{V}_{state}(t, \mu)$ and $\varepsilon > 0$. Let $\varepsilon_1 > 0$ be a small number which will be specified later. Since $\varphi \in \mathbb{V}_{state}^{\varepsilon_1}(t, \mu)$, there exists $\alpha^* \in \mathcal{M}_{state}^{\varepsilon_1}(t, \mu)$ such that $\|\varphi - J(t, \mu, \alpha^*; \cdot, \alpha^*)\|_{\infty} \leq \varepsilon_1$. Introduce $\psi(x, \nu) := J(T_0, \nu, \alpha^*; x, \alpha^*)$. By (2.10) we have

$$\|\varphi - J(T_0, \psi; t, \mu, \alpha^*; \cdot, \alpha^*)\|_{\infty} = \|\varphi - J(t, \mu, \alpha^*; \cdot, \alpha^*)\|_{\infty} \leq \varepsilon_1.$$

Moreover, since $\alpha^* \in \mathcal{M}_{state}^{\varepsilon_1}(t, \mu)$, for any $\alpha \in \mathcal{A}_{state}$ and $x \in \mathbb{S}$, we have

$$\begin{aligned} J(T_0, \psi; t, \mu, \alpha^*; x, \alpha^*) &= J(t, \mu, \alpha^*; x, \alpha^*) \\ &\leq J(t, \mu, \alpha^*; x, \alpha \oplus_{T_0} \alpha^*) + \varepsilon_1 = J(T, \psi; t, \mu, \alpha^*; x, \alpha) + \varepsilon_1. \end{aligned}$$

This implies that $\alpha^* \in \mathcal{M}_{state}^{\varepsilon_1}(T_0, \psi; t, \mu)$. We claim further that

$$\psi(\cdot, \mu_{T_0}^{\alpha^*}) \in \mathbb{V}_{C\varepsilon_1}(T_0, \mu_{T_0}^{\alpha^*}), \quad (7.5)$$

for some constant $C \geq 1$. Then by (2.17) we see that $\varphi \in \tilde{\mathbb{V}}_{state}^{C\varepsilon_1}(t, \mu) \subset \tilde{\mathbb{V}}_{state}^\varepsilon(t, \mu)$ by setting $\varepsilon_1 \leq \frac{\varepsilon}{C}$. Since $\varepsilon > 0$ is arbitrary, we obtain $\varphi \in \tilde{\mathbb{V}}_{state}(t, \mu)$.

To show (7.5), we follow the arguments in Proposition 2.3 (ii). Recall v in (2.5) and the standard DPP (2.11) for v , for any $x \in \mathbb{S}$ we have

$$\begin{aligned} \mathbb{E}^{\mathbb{P}^{\mu^{\alpha^*}; t, x, \alpha^*}} \left[J(T_0, \mu_{T_0}^{\alpha^*}, \alpha^*; X_{T_0}, \alpha^*) \right] &\leq \inf_{\alpha \in \mathcal{A}_{state}} \mathbb{E}^{\mathbb{P}^{\mu^{\alpha^*}; t, x, \alpha^*}} \left[J(T_0, \mu_{T_0}^{\alpha^*}, \alpha^*; X_{T_0}, \alpha) \right] + \varepsilon_1 \\ &= \mathbb{E}^{\mathbb{P}^{\mu^{\alpha^*}; t, x, \alpha^*}} \left[v(\mu^{\alpha^*}; T_0, X_{T_0}) \right] + \varepsilon_1, \end{aligned}$$

It is obvious that $v(\mu^{\alpha^*}; T_0, \cdot) \leq J(T_0, \mu_{T_0}^{\alpha^*}, \alpha^*; \cdot, \alpha^*)$. Moreover, since $q \geq c_q$, clearly $\mathbb{P}^{\mu^{\alpha^*}; t, x, \alpha^*}(X_{T_0} = \tilde{x}) \geq c_0^{T_0-t}$, for any $\tilde{x} \in \mathbb{S}$. Thus, for $C := c_0^{t-T_0}$,

$$\begin{aligned} 0 &\leq J(T_0, \mu_{T_0}^{\alpha^*}, \alpha^*; \tilde{x}, \alpha^*) - v(\mu^{\alpha^*}; T_0, \tilde{x}) \\ &\leq C \mathbb{E}^{\mathbb{P}^{\mu^{\alpha^*}; t, x, \alpha^*}} \left[\left[J(T_0, \mu_{T_0}^{\alpha^*}, \alpha^*; X_{T_0}, \alpha^*) - v(\mu^{\alpha^*}; T_0, X_{T_0}) \right] \mathbf{1}_{\{X_{T_0} = \tilde{x}\}} \right] \\ &\leq C \mathbb{E}^{\mathbb{P}^{\mu^{\alpha^*}; t, x, \alpha^*}} \left[\left[J(T_0, \mu_{T_0}^{\alpha^*}, \alpha^*; X_{T_0}, \alpha^*) - v(\mu^{\alpha^*}; T_0, X_{T_0}) \right] \right] \leq C\varepsilon_1. \end{aligned}$$

This implies that $\alpha^* \in \mathcal{M}_{state}^{C\varepsilon_1}(T_0, \mu_{T_0}^{\alpha^*})$. Since $\psi(\cdot, \mu_{T_0}^{\alpha^*}) = J(T_0, \mu_{T_0}^{\alpha^*}, \alpha^*; \cdot, \alpha^*)$, we obtain (7.5) immediately, and hence $\varphi \in \tilde{\mathbb{V}}_{state}(t, \mu)$. ■

Proof of the claim in Remark 4.7. By (4.16) and (4.17) we have

$$\begin{aligned}
\gamma^{(\Lambda^\gamma)}(s, \tilde{\mathbf{x}}, da) &:= \frac{1}{\mu_{s\wedge\cdot}^\gamma(\tilde{\mathbf{x}})} \int_{\mathcal{A}_{path}^t} Q_s^t(\mu^\gamma; \tilde{\mathbf{x}}; \alpha) \delta_{\alpha(s, \tilde{\mathbf{x}})}(da) \Lambda^\gamma(\mathbf{x}, da) \\
&= \frac{1}{\mu_{s\wedge\cdot}^\gamma(\tilde{\mathbf{x}})} \int_{\mathbb{A}} \cdots \int_{\mathbb{A}} \left[\prod_{r=t}^{s-1} q(r, \tilde{\mathbf{x}}, \mu^\gamma, \alpha(r, \tilde{\mathbf{x}}); \mathbf{x}_{r+1}) \right] \times \delta_{\alpha(s, \tilde{\mathbf{x}})}(da) \times \\
&\quad \left[\mu(\mathbf{x}) \prod_{r=t}^{T-1} \prod_{\tilde{\mathbf{x}} \in \mathbb{X}_s^{t, \mathbf{x}}} \gamma(r, \tilde{\mathbf{x}}, d\alpha(r, \tilde{\mathbf{x}})) \right] \\
&= \frac{\mu(\mathbf{x})}{\mu_{s\wedge\cdot}^\gamma(\tilde{\mathbf{x}})} \int_{\mathbb{A}} \cdots \int_{\mathbb{A}} \left[\prod_{r=t}^{s-1} q(r, \tilde{\mathbf{x}}, \mu^\gamma, \alpha(r, \tilde{\mathbf{x}}); \mathbf{x}_{r+1}) \gamma(r, \tilde{\mathbf{x}}, d\alpha(r, \tilde{\mathbf{x}})) \right] \times \\
&\quad \left[\delta_{\alpha(s, \tilde{\mathbf{x}})}(da) \gamma(s, \tilde{\mathbf{x}}, d\alpha(s, \tilde{\mathbf{x}})) \prod_{\tilde{\mathbf{x}} \in \mathbb{X}_s^{t, \mathbf{x}} \setminus \{\tilde{\mathbf{x}}\}} \gamma(s, \tilde{\mathbf{x}}, d\alpha(s, \tilde{\mathbf{x}})) \right] \times \\
&\quad \left[\prod_{r=t}^{s-1} \prod_{\tilde{\mathbf{x}} \in \mathbb{X}_s^{t, \mathbf{x}} \setminus \{\tilde{\mathbf{x}}\}} \gamma(r, \tilde{\mathbf{x}}, d\alpha(r, \tilde{\mathbf{x}})) \right] \left[\prod_{r=s}^{T-1} \prod_{\tilde{\mathbf{x}} \in \mathbb{X}_s^{t, \mathbf{x}}} \gamma(r, \tilde{\mathbf{x}}, d\alpha(r, \tilde{\mathbf{x}})) \right] \\
&= \frac{\mu(\mathbf{x})}{\mu_{s\wedge\cdot}^\gamma(\tilde{\mathbf{x}})} \left[\prod_{r=t}^{s-1} \int_{\mathbb{A}} q(r, \tilde{\mathbf{x}}, \mu^\gamma, \bar{a}; \mathbf{x}_{r+1}) \gamma(r, \tilde{\mathbf{x}}, d\bar{a}) \right] \times [\gamma(s, \tilde{\mathbf{x}}, da)] \\
&= \frac{\mu(\mathbf{x})}{\mu_{s\wedge\cdot}^\gamma(\tilde{\mathbf{x}})} Q_s^t(\mu^\gamma; \tilde{\mathbf{x}}, \gamma) \gamma(s, \tilde{\mathbf{x}}, da) = \gamma(s, \tilde{\mathbf{x}}, da).
\end{aligned}$$

That is, $\gamma^{(\Lambda^\gamma)} = \gamma$. ■

Proof of Lemma 6.3. Clearly the uniform estimate for $J(\mu^\alpha; \cdot)$ implies that for $v(\mu^\alpha; \cdot)$, so we shall only prove the former one. Fix $(t, \mu) \in [0, T] \times \mathcal{P}_2$ and $\alpha, \tilde{\alpha} \in \mathcal{A}_{cont}$, and denote $u(s, x) := J(\mu^\alpha; \tilde{\alpha}, s, x)$. By standard PDE theory u is a classical solution to the linear PDE in (6.4) and we have the following formula: denoting $X_r^{s, x} := x + B_r - B_s$,

$$\begin{aligned}
\partial_x u(s, x) &= \mathbb{E}^\mathbb{P} \left[[g(X_T^{s, x}, \mu_T^\alpha) - g(x, \mu_T^\alpha)] \frac{B_T - B_s}{T - s} \right. \\
&\quad \left. + \int_s^T \left[b(r, X_r^{s, x}, \mu_r^\alpha, \tilde{\alpha}(r, X_r^{s, x})) \cdot \partial_x u(r, X_r^{s, x}) + f(r, X_r^{s, x}, \mu_r^\alpha, \tilde{\alpha}(r, X_r^{s, x})) \right] \frac{B_r - B_s}{r - s} dr \right].
\end{aligned}$$

Then, by the Lipschitz continuity of g and the boundedness of b and f ,

$$\begin{aligned}
|\partial_x u(s, x)| &\leq \mathbb{E} \left[L_0 \frac{|B_T - B_s|^2}{T - s} + C_0 \int_s^T [|\partial_x u(r, X_r^{s, x})| + 1] \frac{|B_r - B_s|}{r - s} dr \right] \\
&\leq C + C_0 \mathbb{E} \left[\int_s^T |\partial_x u(r, X_r^{s, x})| \frac{|B_r - B_s|}{r - s} dr \right].
\end{aligned}$$

Denote $K_s := e^{\lambda s} \sup_x |\partial_x u(s, x)|$, $\bar{K} := \sup_{t \leq s \leq T} K_s$, for some constant $\lambda > 0$. Then

$$\begin{aligned} K_s &\leq Ce^{\lambda s} + C_0 \int_s^T \frac{K_r e^{-\lambda(r-s)}}{\sqrt{r-s}} dr \leq Ce^{\lambda s} + C_0 \bar{K} \int_s^T \frac{e^{-\lambda(r-s)}}{\sqrt{r-s}} dr \\ &\leq Ce^{\lambda s} + C_0 \bar{K} \int_s^\infty \frac{e^{-\lambda(r-s)}}{\sqrt{r-s}} dr = Ce^{\lambda s} + C_0 \bar{K} \int_0^\infty \frac{e^{-\lambda r}}{\sqrt{r}} dr = Ce^{\lambda s} + \frac{C_0}{\sqrt{\pi\lambda}} \bar{K}. \end{aligned}$$

Thus $\bar{K} \leq \frac{C_0}{\sqrt{\pi\lambda}} \bar{K} + Ce^{\lambda T}$. Set $\lambda := \frac{4C_0^2}{\pi}$ so that $\frac{C_0}{\sqrt{\pi\lambda}} = \frac{1}{2}$, we obtain $\bar{K} \leq C_1 := 2Ce^{\lambda T}$, which implies the desired estimate immediately. \blacksquare

Proof of Proposition 6.5. Fix $(t, \vec{x}, \vec{\alpha}, \bar{x}, \tilde{x})$ and i . For any $\tilde{\alpha} \in \mathcal{A}_{cont}^L$, introduce $\bar{\alpha}(s, x, \mu) := \tilde{\alpha}(s, x - \bar{x} + \tilde{x}, \mu)$, and denote

$$\begin{aligned} \bar{X}_s^i &:= \bar{x} + B_s^i - B_t^i, \quad X_s^j := x_j + B_s^j - B_t^j, \quad j \neq i, \quad ; \\ \bar{\mu}_s^N &:= \frac{1}{N} [\delta_{\bar{X}_s^i} + \sum_{j \neq i} \delta_{X_s^j}], \quad \bar{M}_s^j := \exp \left(\int_t^s \bar{b}_r^j dB_r^j - \frac{1}{2} \int_t^s |\bar{b}_r^j|^2 dr \right), \quad j \geq 1, \quad \text{where} \\ \bar{b}_s^i &:= b(s, \bar{X}_s^i, \bar{\mu}_s^N, \bar{\alpha}(s, \bar{X}_s^i, \bar{\mu}_s^N)), \quad \bar{b}_s^j := b(s, X_s^j, \bar{\mu}_s^N, \alpha_j(s, X_s^j, \bar{\mu}_s^N)), \quad j \neq i. \end{aligned}$$

By the Girsanov Theorem we have

$$J_i(t, (\bar{x}^{-i}, \bar{x}), (\bar{\alpha}^{-i}, \bar{\alpha})) = \mathbb{E} \left[\left[\prod_{j=1}^N \bar{M}_T^j \right] [g(\bar{X}_T^i, \bar{\mu}_T^N) + \int_t^T f(s, \bar{X}_s^i, \bar{\mu}_s^N, \bar{\alpha}(s, \bar{X}_s^i, \bar{\mu}_s^N))] ds \right].$$

Similarly define $\tilde{X}^i, \tilde{\mu}^N, \tilde{M}^j, \tilde{b}^i, \tilde{b}^j$ corresponding to $(\tilde{x}, \tilde{\alpha})$ in the obvious sense. Then we have a similar expression as above and $\bar{\alpha}(s, \bar{X}_s^i, \mu) = \tilde{\alpha}(s, \tilde{X}_s^i, \mu)$. Therefore,

$$\begin{aligned} &v_i^{N,L}(t, (\bar{x}^{-i}, \bar{x}), \bar{\alpha}) - J_i(t, (\bar{x}^{-i}, \tilde{x}), (\bar{\alpha}^{-i}, \tilde{\alpha})) \\ &\leq J_i(t, (\bar{x}^{-i}, \bar{x}), (\bar{\alpha}^{-i}, \bar{\alpha})) - J_i(t, (\bar{x}^{-i}, \tilde{x}), (\bar{\alpha}^{-i}, \tilde{\alpha})) \leq C \sum_{j=1}^N K_T^j + K_0, \end{aligned} \quad (7.6)$$

where

$$\begin{aligned} K_s^j &:= \mathbb{E} \left[\left[\prod_{k < j} \bar{M}_s^k \right] \left[\prod_{k > j} \tilde{M}_s^k \right] |\bar{M}_s^j - \tilde{M}_s^j| \right], \quad j \geq 1; \\ K_0 &:= \mathbb{E} \left[\prod_{j=1}^N \bar{M}_T^j [|g(\bar{X}_T^i, \bar{\mu}_T^N) - g(\tilde{X}_T^i, \tilde{\mu}_T^N)| \right. \\ &\quad \left. + \int_t^T |f(s, \bar{X}_s^i, \bar{\mu}_s^N, \bar{\alpha}(s, \bar{X}_s^i, \bar{\mu}_s^N)) - f(s, \tilde{X}_s^i, \tilde{\mu}_s^N, \tilde{\alpha}(s, \tilde{X}_s^i, \tilde{\mu}_s^N))| ds \right]. \end{aligned}$$

Denote $\Delta x := \bar{x} - \tilde{x}$. Note that

$$\begin{aligned} \bar{X}_s^i - \tilde{X}_s^i &= \Delta x, \quad W_1(\bar{\mu}_s^N, \tilde{\mu}_s^N) \leq \frac{|\Delta x|}{N}, \\ |\bar{\alpha}(s, \bar{X}_s^i, \bar{\mu}_s^N) - \tilde{\alpha}(s, \tilde{X}_s^i, \tilde{\mu}_s^N)| &= |\bar{\alpha}(s, \tilde{X}_s^i, \bar{\mu}_s^N) - \tilde{\alpha}(s, \tilde{X}_s^i, \tilde{\mu}_s^N)| \leq \frac{L}{N} |\Delta x|. \end{aligned} \quad (7.7)$$

By the required Lipschitz continuity, we have

$$K_0 \leq C \mathbb{E}^{\mathbb{P}} \left[\prod_{j=1}^N \bar{M}_T^j \left[\left[1 + \frac{1}{N} \right] |\Delta x| + \int_t^T \left[1 + \frac{L}{N} \right] |\Delta x| ds \right] \right] \leq C |\Delta x|. \quad (7.8)$$

Next, introduce

$$\Gamma_s^j := \mathbb{E} \left[\left[\prod_{k < j} \bar{M}_s^k \right] \left[\prod_{k > j} \tilde{M}_s^k \right] |\bar{M}_s^j|^2 \right], \quad \Delta \Gamma_s^j := \mathbb{E} \left[\left[\prod_{k < j} \bar{M}_s^k \right] \left[\prod_{k > j} \tilde{M}_s^k \right] |\bar{M}_s^j - \tilde{M}_s^j|^2 \right].$$

Note that B^1, \dots, B^N are independent. By applying the Ito formula, we have

$$\Gamma_s^j = 1 + \int_t^s \mathbb{E} \left[\left[\prod_{k < j} \bar{M}_r^k \right] \left[\prod_{k > j} \tilde{M}_r^k \right] |\bar{M}_r^j \bar{b}_r^j|^2 \right] dr \leq 1 + C \int_t^s \Gamma_r^j dr,$$

Then $\Gamma_s^j \leq C$. Thus, by applying the Ito formula again we have

$$\begin{aligned} \Delta \Gamma_s^j &= \int_t^s \mathbb{E} \left[\left[\prod_{k < j} \bar{M}_r^k \right] \left[\prod_{k > j} \tilde{M}_r^k \right] [\bar{M}_r^j \bar{b}_r^j - \tilde{M}_r^j \tilde{b}_r^j]^2 \right] dr \\ &\leq C \int_t^s \mathbb{E} \left[\left[\prod_{k < j} \bar{M}_r^k \right] \left[\prod_{k > j} \tilde{M}_r^k \right] [|\bar{M}_r^j - \tilde{M}_r^j| + \bar{M}_r^j |\bar{b}_r^j - \tilde{b}_r^j|]^2 \right] dr \\ &\leq C \int_t^s \Delta \Gamma_r^j dr + C \int_t^s \mathbb{E} \left[\left[\prod_{k < j} \bar{M}_r^k \right] \left[\prod_{k > j} \tilde{M}_r^k \right] [\bar{M}_r^j |\bar{b}_r^j - \tilde{b}_r^j|]^2 \right] dr. \end{aligned}$$

Note that, by (7.7),

$$\begin{aligned} |\bar{b}_r^i - \tilde{b}_r^i| &= \left| b(s, \bar{X}_s^i, \bar{\mu}_s^N, \tilde{\alpha}(s, \tilde{X}_s^i, \tilde{\mu}_s^N)) - b(s, \tilde{X}_s^i, \tilde{\mu}_s^N, \tilde{\alpha}(s, \tilde{X}_s^i, \tilde{\mu}_s^N)) \right| \leq C_L |\Delta x| \\ |\bar{b}_r^j - \tilde{b}_r^j| &\leq \frac{C_L}{N} |\Delta x|, \quad j \neq i. \end{aligned}$$

Then, since $\Gamma_s^j \leq C$,

$$\Delta \Gamma_s^i \leq C \int_t^s \Delta \Gamma_r^i dr + C_L |\Delta x|^2, \quad \Delta \Gamma_s^j \leq C \int_t^s \Delta \Gamma_r^j dr + \frac{C_L}{N^2} |\Delta x|^2, \quad j \neq i.$$

and thus

$$\Delta \Gamma_s^i \leq C_L |\Delta x|^2, \quad \Delta \Gamma_s^j \leq \frac{C_L}{N^2} |\Delta x|^2, \quad j \neq i.$$

Note further that

$$K_s^i \leq \frac{|\Delta x|}{2} + \frac{\Delta \Gamma_s^i}{2|\Delta x|} \leq C_L |\Delta x|, \quad K_s^j \leq \frac{|\Delta x|}{2N} + \frac{N \Delta \Gamma_s^j}{2|\Delta x|} \leq \frac{C_L}{N} |\Delta x|, \quad j \neq i. \quad (7.9)$$

Then, by (7.6), (7.8) and (7.9) we have

$$\begin{aligned} v_i^{N,L}(t, (\vec{x}^{-i}, \bar{x}), \vec{\alpha}) - J_i(t, (\vec{x}^{-i}, \tilde{x}), (\vec{\alpha}^{-i}, \tilde{\alpha})) &\leq K_0 + CK_s^i + C \sum_{j \neq i} K_s^j \\ &\leq C|\Delta x| + C_L|\Delta x| + C_L \sum_{j \neq i} \frac{|\Delta x|}{N} \leq C_L|\Delta x|. \end{aligned}$$

Since $\tilde{\alpha} \in \mathcal{A}^L$ is arbitrary, we obtain $v_i^{N,L}(t, (\vec{x}^{-i}, \bar{x}), \vec{\alpha}) - v_i^{N,L}(t, (\vec{x}^{-i}, \tilde{x}), \vec{\alpha}) \leq C_L|\Delta x|$. Similarly we can show that $v_i^{N,L}(t, (\vec{x}^{-i}, \tilde{x}), \vec{\alpha}) - v(t, (\vec{x}^{-i}, \bar{x}), \vec{\alpha}) \leq C_L|\Delta x|$, and thus obtain (6.13). ■

References

- [1] Abreu, D.; Pearce, D.; and Stacchetti, E., *Toward a Theory of Discounted Repeated Games with Imperfect Monitoring*, *Econometrica*, 58 (1990), 1041-1063.
- [2] Bardi, M. and Fischer, M., *On non-uniqueness and uniqueness of solutions in finite-horizon Mean Field Games*, *ESAIM: Control, Optimisation and Calculus of Variations*, 25 (2019), 44.
- [3] Bayraktar, E. and Cohen, A., *Analysis of a finite state many player using its master equation*, *SIAM J. Control Optim.* 56 (2018), 3538-3568.
- [4] Bayraktar, E.; Cecchin, A.; Cohen, A.; and Delarue, F., *Finite state mean field games with Wright-Fisher common noise*, *J. Math. Pures et Appliquées*, 147 (2021), 98-162.
- [5] Bayraktar, E.; Cecchin, A.; Cohen, A.; and Delarue, F., *Finite state mean field games with Wright-Fisher common noise as limits of N-player weighted games*, preprint, arXiv:2012.04845.
- [6] Bensoussan, A.; Frehse, J.; and Yam, S. C. P., *Mean Field Games and Mean Field Type Control Theory*, (2013). New York: Springer Verlag.
- [7] Caines, P. E.; Huang, M.; and Malhamé, R. P., *Large population stochastic dynamic games: closed loop McKean-Vlasov systems and the Nash certainty equivalence principle*, *Commun. Inf. Syst.*, 6 (2006), 221–251.
- [8] Cardaliaguet, P., *Notes on mean field games*, lectures by P.L. Lions, Collège de France, (2010).

- [9] Cardaliaguet, P., *The convergence problem in mean field games with local coupling*, *Applied Mathematics & Optimization*, 76 (2017), 177-215.
- [10] Cardaliaguet, P.; Delarue, F.; Lasry, J.M.; and Lions, P.L., *The master equation and the convergence problem in mean field games*, *Annals of Mathematics Studies*, 201. Princeton University Press, Princeton, NJ, (2019). x+212 pp.
- [11] Carmona, G., *Nash Equilibria of Games with a Continuum of Players*, *Game Theory and Information*, 0412009, University Library of Munich, Germany, 2004.
- [12] Carmona, R. and Delarue, F. *Probabilistic Analysis of Mean-Field Games*, *SIAM Journal on Control and Optimization*, 51 (2013), 2705-2734.
- [13] Carmona, R. and Delarue, F., *Probabilistic theory of mean field games with applications I - Mean field FBSDEs, control, and games*, *Probability Theory and Stochastic Modeling*, 83. Springer, Cham, (2018).
- [14] Carmona, R. and Delarue, F., *Probabilistic theory of mean field games with applications II - Mean field games with common noise and master equations*, *Probability Theory and Stochastic Modeling*, 84. Springer, Cham, (2018).
- [15] Cecchin A.; Dai Pra P.; Fischer M.; and Pelino G., *On the convergence problem in mean field games: a two state model without uniqueness*, *SIAM J. Control Optim.* 57 (2019), 2443-2466.
- [16] Cecchin A. and Delarue F., *Selection by vanishing common noise for potential finite state mean field games*, preprint, arXiv:2005.12153.
- [17] Cecchin, A. and Pelino, G., *Convergence, Fluctuations and Large Deviations for finite state Mean Field Games via the Master Equation*. *Stochastic Process. Appl.* 129 (2019), 4510-4555.
- [18] Delarue, F., *Restoring uniqueness to mean-field games by randomizing the equilibria*, *Stochastic and Partial Differential Equations: Analysis and Computations* 7 (2019), 598-678.
- [19] Delarue, F. and Foguen Tchuendom, R., *Selection of equilibria in a linear quadratic mean-field game*, *Stochastic Process. Appl.* 130 (2020), 1000-1040.

- [20] Delarue, F.; Lacker D.; and Ramanan, K., *From the master equation to mean field game limit theory: Large deviations and concentration of measure*. *Ann. Probab.* 48 (2020), 211-263.
- [21] Delarue, F.; Lacker D.; and Ramanan, K., *From the master equation to mean field game limit theory: A central limit theorem*. *Electron. J. Probab.* 24 (2019), 1-54.
- [22] Djete, M. F., *Large population games with interactions through controls and common noise: convergence results and equivalence between open-loop and closed-loop controls*, preprint, arXiv:2108.02992.
- [23] Feinstein, Z., *Continuity and Sensitivity Analysis of Parameterized Nash Games*. preprint, arXiv:2007.04388.
- [24] Feinstein, Z.; Rudloff, B.; and Zhang, J. *Dynamic set values for nonzero sum games with multiple equilibria*, *Mathematics of Operations Research*, accepted, arXiv:2002.00449.
- [25] Feleqi, E., *The Derivation of Ergodic Mean Field Game Equations for Several Populations of Players*. *Dynamic Games and Applications*, 3 (2013), 523-536.
- [26] Fischer, M., *On the connection between symmetric N-player games and mean field games*. *The Annals of Applied Probability*, 27 (2017), 757-810.
- [27] Fischer, M. and Silva, F.J., *On the Asymptotic Nature of First Order Mean Field Games* *Applied Mathematics & Optimization* (2020), 1432-0606.
- [28] Foguen Tchuendom, R., *Uniqueness for linear-quadratic mean field games with common noise*, *Dynamic Games and Applications* 8 (2018), 199-210.
- [29] Gangbo, W. and Mészáros, A.R., *Global well-posedness of master equations for deterministic displacement convex potential mean field games*, preprint, arXiv:2004.01660.
- [30] Lacker, D., *A general characterization of the mean field limit for stochastic differential games*, *Probab. Th. Rel. Fields* 165 (2016), 581-648.
- [31] Lacker, D., *On the convergence of closed loop Nash equilibria to the mean field game limit*, *Ann. Appl. Probab.* 30 (2020), 1693-1761.
- [32] Lacker, D. and Flem, L.L., *Closed loop convergence for mean field games with common noise*, preprint, arXiv:2107.03273.

- [33] Lasry, J.-M. and Lions, P.-L., Mean field games, *Jpn. J. Math.* 2 (2007), 229–260.
- [34] Lauriere, M. and Tangpi, L. *Convergence of large population games to mean field games with interaction through the controls*, preprint arXiv:2004.08351.
- [35] Lions, P.-L., *Cours au Collège de France*, www.college-de-france.fr.
- [36] Mou, C. and Zhang, J., *Wellposedness of second order master equations for mean field games with nonsmooth data*, preprint, arXiv:1903.09907v2.
- [37] Nutz, M.; San Martin, J.; and Tan, X., *Convergence to the Mean Field Game Limit: A Case Study*, *Annals of Applied Probability*, 30 (2020), 259-286.
- [38] Pham, T. and Zhang, J., *Two person zero-sum game in weak formulation and path dependent Bellman-Isaacs equation*. *SIAM Journal on Control and Optimization*, 52 (2004), 2090-2121.
- [39] Possamai, D. and Tangpi, L., *Non-asymptotic convergence rates for mean-field games: weak formulation and McKean–Vlasov BSDEs*, preprint, arXiv:2105.00484.
- [40] Sannikov, Y., *Games with Imperfectly Observable Actions in Continuous Time*, *Econometrica*, 75 (2007), 1285-1329.
- [41] Zhang, J., *Backward Stochastic Differential Equations – from linear to fully nonlinear theory*, *Probability Theory and Stochastic Modeling* 86, Springer, New York, 2017.