PEARSON EQUATIONS FOR DISCRETE ORTHOGONAL POLYNOMIALS: I. GENERALIZED HYPERGEOMETRIC FUNCTIONS AND TODA EQUATIONS

MANUEL MAÑAS¹, ITSASO FERNÁNDEZ-IRISARRI², AND OMAR F. GONZÁLEZ-HERNÁNDEZ³

ABSTRACT. The Cholesky factorization of the moment matrix is applied to discrete orthogonal polynomials on the homogeneous lattice. In particular, semiclassical discrete orthogonal polynomials, which are built in terms of a discrete Pearson equation, are studied. The Laguerre–Freud structure semi-infinite matrix that models the shifts by ± 1 in the independent variable of the set of orthogonal polynomials is introduced. In the semiclassical case it is proven that this Laguerre–Freud matrix is banded. From the well known fact that moments of the semiclassical weights are logarithmic derivatives of generalized hypergeometric functions, it is shown how the contiguous relations for these hypergeometric functions translate as symmetries for the corresponding moment matrix. It is found that the 3D Nijhoff–Capel discrete Toda lattice describes the corresponding contiguous shifts for the squared norms of the orthogonal polynomials. The continuous Toda for these semiclassical discrete orthogonal polynomials is discussed and the compatibility equations are derived. It also shown that the Kadomtesev– Petvishvilii equation is connected to an adequate deformed semiclassical discrete weight, but in this case the deformation do not satisfy a Pearson equation.

CONTENTS

1. Introduction	2
1.1. Linear functionals and orthogonal polynomials	2
1.2. The shift matrix Λ	3
1.3. The Cholesky factorization of the Gram matrix	3
1.4. The Jacobi matrix	4
1.5. The lower Pascal matrix	5
2. Discrete orthogonal polynomials and discrete Pearson equations	8
2.1. Discrete Pearson equation	8
2.2. Consequences for the Jacobi matrix	g
2.3. Generalized hypergeometric functions and the Pearson equation	10
2.4. The Laguerre–Freud structure matrix and the Cholesky factorization	11
2.5. Contiguous hypergeometric relations	13
2.6. Discrete lattice Toda equation	15
2.7. The Toda flows	18
2.8. Toda–Pearson compatibility	22
3. Conclusions and outlook	23
Acknowledgments	24
References	24

¹Departamento de Física Teórica, Universidad Complutense de Madrid, Plaza Ciencias 1, 28040-Madrid, Spain & Instituto de Ciencias Matematicas (ICMAT), Campus de Cantoblanco UAM, 28049-Madrid, Spain

E-mail addresses: ¹manuel.manas@ucm.es, ²itsasofe@ucm.es, ³omgonzal@math.uc3m.es.

²⁰²⁰ Mathematics Subject Classification. 42C05,33C45,33C47.

Key words and phrases. Discrete orthogonal polynomials, Pearson equations, Cholesky factorization, generalized hypergeometric functions, contigous relations, 3D Nijhoff-Capel discrete Toda equation, Toda hierarchy.

¹Thanks financial support from the Spanish "Agencia Estatal de Investigación" research project [PGC2018-096504-B-C33], Ortogonalidad y Aproximación: Teoría y Aplicaciones en Física Matemática.

1. INTRODUCTION

In this paper, the Gauss–Borel factorization of the moment matrix approach to orthogonality is applied to study discrete orthogonal polynomials subject to a Pearson equation.

Discrete orthogonal polynomials is nowadays a well established subject, the classical case has been treated extensively in [42] and the Riemann–Hilbert problem has been used to study asymptotics and applications, see [16]. Semiclassical reductions, in where a discrete Pearson equation is fulfilled by the weight has been also treated in the literature. See for example [23, 24] and [21, 22] and references therein for an comprehensive account. For some specific type of weights of generalized Charlier and Meixner types, the corresponding Freud–Laguerre type equations for the coefficients of the three term recurrence has been studied, see for example [19, 26, 27, 28, 43]. For a general account see [31, 32, 17] and for discrete orthogonal polynomials and Painlevé equations see [44].

With this introduction we give a fast briefing, of introductory character, of orthogonal polynomials, its discrete version and the Cholesky factorization of the moment matrix. We also discuss for the first time in this context the so called Pascal matrices and its dressed versions.

Then, in §2 we discuss the discrete Pearson equation and give Theorem 1, the first main result of the paper, that describes a new symmetry of the moment matrix, direct consequence of the Pearson equation, in terms of the Pascal matrix. We then deduce the corresponding symmetry for the Jacobi matrix, see Proposition 6. Then, in Theorem 2, the second main result of the paper, we describe a banded semi-infinite matrix, that we called Laguerre–Freud structure matrix, that models the shift in the spectral variable. The name is rooted on the fact that this banded matrix encodes Laguerre–Freud equations for the recurrence coefficients of the orthogonal polynomial sequence, see for example [25]. The number of non-trivial superdiagonals and subdiagonals of this matrix is determined by the Pearson equation. Several properties involving the Jacobi and Laguerre–Freud matrices are derived. Of particular interest are those of compatibility type.

As the Pearson equation leads to generalized hypergeometric moments, we study the contiguous relations of these functions and its description as further symmetries of the moment matrix, see Theorem 4. In Theorem 5 the squared norms of these discrete orthogonal polynomials are shown to be solutions of a well known discrete integrable multidimensional lattice, known as the Nijhoff–Capel lattice, see [30, 41]. We also discuss the Toda hierarchy deformations, with only the first flow preserving the Pearson reduction. The compatibility with this first Toda flow and the Pearson equation is discussed in Proposition 21. We also give the connection of deformations of these weights with solutions of the Kadomtsev–Petviashvilli (KP) equation.

1.1. Linear functionals and orthogonal polynomials. Given a linear functional $\rho_z \in \mathbb{C}^*[z]$, here $\mathbb{C}^*[z]$ denotes the dual of the linear space of complex polynomials $\mathbb{C}[z]$, the corresponding moment matrix is

$$G = (G_{n,m}), \qquad G_{n,m} = \rho_{n+m}, \qquad \rho_n = \langle \rho_z, z^n \rangle, \qquad n, m \in \mathbb{N}_0 := \{0, 1, 2, \dots\},\$$

with ρ_n the *n*-th moment of the linear functional ρ_z . If the moment matrix is such that all its truncations, which are Hankel matrices, $G_{i+1,j} = G_{i,j+1}$,

$$G^{[k]} = \begin{pmatrix} G_{0,0} \cdots \cdots \cdots G_{0,k-1} \\ \vdots & \vdots \\ G_{k-1,0} \cdots \cdots & G_{k-1,k-1} \end{pmatrix} = \begin{pmatrix} \rho_0 & \rho_1 & \rho_2 \cdots \cdots & \rho_{k-1} \\ \rho_1 & \rho_2 & \ddots & \ddots & \rho_k \\ \rho_2 & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & \vdots \\ \rho_{k-1} & \rho_k \cdots & \cdots & \rho_{2k-2} \end{pmatrix}$$

are nonsingular; i.e. the Hankel determinants $\Delta_k := \det G^{[k]}$ do not cancel, $\Delta_k \neq 0$, $k \in \mathbb{N}_0$, then there exists monic polynomials

(1)
$$P_n(z) = z^n + p_n^1 z^{n-1} + \dots + p_n^n, \qquad n \in \mathbb{N}_0,$$

with $p_0^1 = 0$, such that the following orthogonality conditions are fulfilled


$$\langle \rho, P_n(z)z^k \rangle = 0,$$
 $k \in \{0, \dots, n-1\},$ $\langle \rho, P_n(z)z^n \rangle = H_n \neq 0.$

Moreover, the set $\{P_n(z)\}_{n\in\mathbb{N}_0}$ is an orthogonal set of polynomials $\langle \rho, P_n(z)P_m(z)\rangle = \delta_{n,m}H_n, n, m \in \mathbb{N}_0$. In this case, we have a symmetric bilinear form $\langle F, G \rangle_{\rho} := \langle \rho, FG \rangle$, such that the moment matrix is the Gram matrix of this bilinear form and $\langle P_n, P_m \rangle_{\rho} := \delta_{n,m}H_n$. In this paper we will use both denominations, moment matrix or Gram matrix, indistinctly to refer to the matrix G.


1.2. The shift matrix Λ . In terms of the semi-infinite vector of monomials

$$\chi(z) := \begin{pmatrix} 1\\ z\\ z^2\\ \vdots\\ \vdots \end{pmatrix}$$

the Gram matrix is written as $G = \langle \rho, \chi \chi^{\top} \rangle$, and it becomes evident this matrix is symmetric, i.e., $G = G^{\top}$. The vector of monomials χ is an eigenvector of the *shift matrix*



i.e., $\Lambda \chi = x \chi$. From here it immediately follows that $\Lambda G = G \Lambda^{\top}$, i.e., the Gram matrix is a Hankel matrix, as we previously said. The transposed matrix



satisfies $\Lambda\Lambda^{\top} = I$ and $\Lambda^{\top}\Lambda = I - E_{0,0}$, with $(E_{i,j})_{k,l} = \delta_{i,k}\delta_{j,l}$, $i, j \in \mathbb{N}_0$.

1.3. **The Cholesky factorization of the Gram matrix**. Being the Gram matrix symmetric its Borel–Gauss factorization reduces to a Cholesky factorization

$$(2) G = S^{-1}HS^{-\top}$$

where S is a lower unitriangular matrix that can be written as

$$S = \begin{pmatrix} 1 & 0 & \dots & \dots \\ S_{1,0} & 1 & \ddots & \\ S_{2,0} & S_{2,1} & \ddots & \ddots \\ \vdots & \ddots & \ddots & \ddots \\ \vdots & \ddots & \ddots & \ddots \end{pmatrix},$$

and $H = \text{diag}(H_0, H_1, ...)$ is a diagonal matrix, with $H_k \neq 0$, for $k \in \mathbb{N}_0$. The Cholesky factorization hold whenever the principal minors of the moment matrix; i.e., the Hankel determinants Δ_k , do not cancel.

The components $P_n(z)$ of the semi-infinite vector of polynomials

$$P(z) \coloneqq S\chi(z),$$

are the monic orthogonal polynomials of the functional $\rho.$ Indeed, from the Cholesky factorization we know that

$$\left\langle \rho, \chi \chi^{\top} \right\rangle = G = S^{-1} H S^{-\top}$$

so that $S \langle \rho, \chi \chi^{\top} \rangle S^{\top} = H$ and, consequently $\langle \rho, S \chi \chi^{\top} S^{\top} \rangle = H$ and we get $\langle \rho, PP^{\top} \rangle = H$, which recollects the orthogonality of the polynomials $\{P_n(z)\}_{n=0}^{\infty}$.

1.4. **The Jacobi matrix**. Given this construction is natural to introduce the lower Hessenberg semi-infinite matrix

$$(4) J = S\Lambda S^{-1}$$

that has the vector P(z) as eigenvector with eigenvalue z, i.e. JP(z) = zP(z). The Hankel condition of the Gram matrix $\Lambda G = G\Lambda^{\top}$ together with the Cholesky factorization leads to $\Lambda S^{-1}HS^{-\top} = S^{-1}HS^{-\top}\Lambda^{\top}$, or, equivalently, $S\Lambda S^{-1}H = HS^{-\top}\Lambda^{\top}S^{-\top}$, i.e.,

That is, the Hessenberg matrix JH is symmetric, thus being Hessenberg and symmetric we deduce that is tridiagonal. Hence, the Jacobi matrix J given in (4) reads

$$J = \begin{pmatrix} \beta_0 & 1 & 0 & \cdots & \cdots & \ddots \\ \gamma_1 & \beta_1 & 1 & \ddots & \vdots \\ 0 & \gamma_2 & \beta_2 & 1 & \ddots & \vdots \\ \vdots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots \\ \vdots & \ddots & \ddots & \ddots & \ddots & \ddots & \ddots \end{pmatrix}$$

and the eigenvalue equation JP = zP is a three term recursion relation

$$zP_n(z) = P_{n+1}(z) + \beta_n P_n(z) + \gamma_n P_{n-1}(z),$$

that with the initial conditions $P_{-1} = 0$ and $P_0 = 1$ completely determines the sequence of orthogonal polynomials $\{P_n(z)\}_{n \in \mathbb{N}_0}$ in terms of the recursion coefficients β_n, γ_n .

Proposition 1. The recursion coefficients, in terms of the Hankel determinants, are given by

(6)
$$\beta_n = p_n^1 - p_{n+1}^1 = -\frac{\Delta_n}{\Delta_n} + \frac{\Delta_{n+1}}{\Delta_{n+1}}, \qquad \gamma_{n+1} = \frac{H_{n+1}}{H_n} = \frac{\Delta_{n+1}\Delta_{n-1}}{\Delta_n^2}, \qquad n \in \mathbb{N}_0,$$

Proof. For $n \in \mathbb{N} := \{1, 2, ...\}$, from $J^{\top} = H^{-1}JH$, we get that $\gamma_n = \frac{H_n}{H_{n-1}}$. On the other hand, as $J = S\Lambda S^{-1}$ and recalling that for the coefficients $S_{n,n-1}$ of the first subdiagonal of S we have $S_{n,n-1} = p_n^1$, the first nontrivial leading coefficient of the monic polynomial P_n , we get $\beta_n = p_n^1 - p_{n+1}^1$.

It can be easily shown that the second kind functions also satisfy the previous three term recursion relation

$$zQ_n(z) = Q_{n+1}(z) + \beta_n Q_n(z) + \gamma_n Q_{n-1}(z)$$

but now with the initial conditions $Q_{-1} = -H_{-1}$ and $Q_0 = S_\rho$ (the Markov–Stieltjes transform of ρ), with $\beta_n = q_n^1 - q_{n-1}^1$.

For future use we introduce the following diagonal matrices

$$\gamma := \operatorname{diag}(\gamma_1, \gamma_2, \dots), \qquad \beta := \operatorname{diag}(\beta_0, \beta_1, \dots)$$

and

$$J_{-} := \Lambda^{\top} \gamma, \qquad \qquad J_{+} := \beta + \Lambda,$$

so that we have the splitting

$$J = \Lambda^{\top} \gamma + \beta + \Lambda = J_{-} + J_{+}.$$

In general, given any semi-infinite matrix A, we will write $A = A_- + A_+$, where A_- is a strictly lower triangular matrix and A_+ an upper triangular matrix. Moreover, A_0 will denote the diagonal part of A.

1.5. The lower Pascal matrix. The lower Pascal matrix, built up of binomial numbers, is defined by

(7)
$$B = (B_{n,m}), \qquad B_{n,m} := \begin{cases} \binom{n}{m}, & n \ge m\\ 0, & n < m \end{cases}$$

so that

(8)
$$\chi(z+1) = B\chi(z).$$

Moreover,

$$B^{-1} = (\tilde{B}_{n,m}), \qquad \qquad \tilde{B}_{n,m} := \begin{cases} (-1)^{n+m} \binom{n}{m}, & n \ge m, \\ 0, & n < m. \end{cases}$$

and

(9)

$$\chi(z-1) = B^{-1}\chi(z).$$

The lower Pascal matrix and its inverse are explicitly given by

Let us introduce the lower unitriangular semi-infinite matrices, lets us refer to them as dressed Pascal matrices,

$$\Pi := SBS^{-1}, \qquad \qquad \Pi^{-1} := SB^{-1}S^{-1},$$

that are connection matrices; i.e.,

$$P(z+1) = \Pi P(z), \qquad P(z-1) = \Pi^{-1} P(z).$$

The lower Pascal matrix can be expressed in terms of its subdiagonal structure as follows

$$B^{\pm 1} = I \pm \Lambda^{\top} D + (\Lambda^{\top})^2 D^{[2]} \pm (\Lambda^{\top})^3 D^{[3]} + \cdots,$$

where the diagonal matrices $D, D^{[k]}$, with $k \in \mathbb{N}$, $(D = D^{[1]})$ are given by

$$D = \operatorname{diag}(1, 2, 3, \dots), \qquad D^{[k]} = \frac{1}{k} \operatorname{diag}\left(k^{(k)}, (k+1)^{(k)}, (k+2)^{(k)} \cdots\right),$$

in terms of the falling factorials $x^{(k)} = x(x-1)(x-2)\cdots(x-k+1)$. That is,

$$D_n^{[k]} = \frac{(n+k)\cdots(n+1)}{k}, \qquad k \in \mathbb{N}, \qquad n \in \mathbb{N}_0.$$

The lower unitriangular factor can be also written in terms of its subdiagonals

$$S = I + \Lambda^{\top} S^{[1]} + (\Lambda^{\top})^2 S^{[2]} + \cdots$$

with $S^{[k]} = \text{diag}\left(S_0^{[k]}, S_1^{[k]}, \dots\right)$ diagonal matrices. From (3) is clear the following connection between these subdiagonals coefficients and the coefficients of the orthogonal polynomials, given in (1), holds

(10)
$$S_k^{[k]} = p_{n+k}^k.$$

We will use the *shift operators* T_{\pm} acting over the diagonal matrices as follows

$$T_{-} \operatorname{diag}(a_0, a_1, \dots) := \operatorname{diag}(a_1, a_2, \dots), \qquad T_{+} \operatorname{diag}(a_0, a_1, \dots) := \operatorname{diag}(0, a_0, a_1, \dots),$$

where T_{-} is the lowering shift operator and T_{+} the raising shift operator over the diagonal matrices. These shift operators have the following important properties, for any diagonal matrix $A = \text{diag}(A_0, A_1, ...)$

(11)
$$\Lambda A = (T_{-}A)\Lambda, \qquad A\Lambda = \Lambda(T_{+}A), \qquad A\Lambda^{\top} = \Lambda^{\top}(T_{-}A), \qquad \Lambda^{\top}A = (T_{+}A)\Lambda^{\top}.$$

Remark 1. Notice that the standard notation, see [42], for the differences of a sequence $\{f_n\}_{n\in\mathbb{N}_0}$,

$$\Delta f_n := f_{n+1} - f_n, \qquad n \in \mathbb{N}_0,$$

$$\nabla f_n = f_n - f_{n-1}, \qquad n \in \mathbb{N},$$

and $\nabla f_0 = f_0$, connects with the shift operators by means of

$$T_{-} = I + \Delta, \qquad \qquad T_{+} = I - \nabla$$

In terms of these shift operators we find

(12)
$$2D^{[2]} = (T_-D)D,$$
 $3D^{[3]} = (T_-^2D)(T_-D)D = 2(T_-D^{[2]})D = 2D^{[2]}(T_-^2D)$

Proposition 2. The inverse matrix S^{-1} of the matrix S expands in terms of subdiagonals as follows

$$S^{-1} = I + \Lambda^{\top} S^{[-1]} + (\Lambda^{\top})^2 S^{[-2]} + \cdots$$

 $\begin{array}{l} \label{eq:subdiagonals} S^{[-k]} \ are \ explicitly \ given \ in \ terms \ of \ the \ subdiagonals \ of \ S, \ the \ first \ few \ are \\ S^{[-1]} = -S^{[1]}, \\ S^{[-2]} = -S^{[2]} + (T_-S^{[1]})S^{[1]}, \\ S^{[-3]} = -S^{[3]} + (T_-S^{[2]})S^{[1]} + (T_-^2S^{[1]})S^{[2]} - (T_-^2S^{[1]})(T_-S^{[1]})S^{[1]}, \\ S^{[-4]} = -S^{[4]} + (T_-S^{[3]})S^{[1]} + (T_-^2S^{[2]})S^{[2]} - (T_-^2S^{[2]})(T_-S^{[1]})S^{[1]} + (T_-^3S^{[1]})S^{[3]} \\ - (T_-^3S^{[1]})(T_-S^{[2]})S^{[1]} - (T_-^3S^{[1]})(T_-^2S^{[1]})S^{[2]} + (T_-^3S^{[1]})(T_-^2S^{[1]})S^{[1]}. \end{array}$

Proposition 3. The following nonlocal expressions for the polynomial coefficients in terms of the recursion coefficients hold true

(13)
$$p_{n+1}^1 = -\sum_{k=0}^n \beta_k, \qquad p_{n+1}^2 = -\sum_{k=1}^n \gamma_k + \sum_{0 \le l < k < n} \beta_k \beta_l.$$

Moreover,

(14)
$$p_{n+2}^3 - p_{n+3}^3 = \gamma_{n+2}p_{n+1}^1 + (\beta_{n+2} + \beta_{n+1} + \beta_n)p_{n+2}^2 - (\beta_{n+1} + \beta_n)p_{n+2}^1p_{n+1}^1.$$

Proof. We have

$$J = S\Lambda S^{-1} = (I + \Lambda^{\top} S^{[1]} + (\Lambda^{\top})^2 S^{[2]} + \dots)\Lambda (I + \Lambda^{\top} S^{[-1]} + (\Lambda^{\top})^2 S^{[-2]} + \dots)$$

= $\Lambda + T_+ S^{[1]} + S^{[-1]} + \Lambda^{\top} (T_+ S^{[2]} + S^{[-2]} + S^{[1]} S^{[-1]})$
+ $(\Lambda^{\top})^2 (T_+ S^{[3]} + S^{[-3]} + S^{[2]} S^{[-1]} + (T_+ S^{[1]}) S^{[-2]}) + \dots$

Thus, we obtain

(15)
$$\beta = T_+ S^{[1]} - S^{[1]},$$

(16) $\gamma = T_+ S^{[2]} - S^{[2]} + (T_- S^{[1]} - S^{[1]}) S^{[1]}$

and an infinite set of relations among subdiagonals of S, being the first

(17)
$$T_+S^{[3]} + S^{[-3]} + S^{[2]}S^{[-1]} + (T_+S^{[1]})S^{[-2]} = 0$$

The relation (15) component wise is (6). As well, component wise, the relation (16) is

$$\gamma_{n+1} = S_{n-1}^{[2]} - S_n^{[2]} + (S_{n+1}^{[1]} - S_n^{[1]})S_n^{[1]} = p_{n+1}^2 - p_{n+2}^2 - \beta_{n+1}p_{n+1}^1$$

Hence, using telescoping series again, we find (13). Finally, Equation (17) reads

$$T_{+}S^{[3]} - S^{[3]} + (T_{-}S^{[2]})S^{[1]} + (T_{-}^{2}S^{[1]})S^{[2]} - (T_{-}^{2}S^{[1]})(T_{-}S^{[1]})S^{[1]} - S^{[2]}S^{[1]} + (T_{+}S^{[1]})(-S^{[2]} + (T_{-}S^{[1]})S^{[1]}) = 0,$$

that we can write

$$\begin{split} T_{+}S^{[3]} - S^{[3]} + T_{-} \left(S^{[2]} - T_{+}S^{[2]} - (T_{-}S^{[1]} - S^{[1]})S^{[1]} \right) S^{[1]} + (T_{-}^{2}S^{[1]} - T_{+}S^{[1]})S^{[2]} \\ &+ (T_{+}S^{[1]} - T_{-}S^{[1]})(T_{-}S^{[1]})S^{[1]} = 0, \end{split}$$

so that

$$T_{+}S^{[3]} - S^{[3]} = (T_{-}\gamma)S^{[1]} + (T_{-}^{2}\beta + T_{-}\beta + \beta)S^{[2]} - (T_{-}\beta + \beta)(T_{-}S^{[1]})S^{[1]} = 0,$$

and component wise we have (14).

Remark 2. In (14) we can sum up on the LHS, observe that is a telescoping series, to get a nonlocal nonlinear expression, in terms of the recursion coefficients, for p_n^3 . A similar statement holds for every p_n^k with $k = 4, 5, \ldots$

Now, we turn our attention to the dressed Pascal matrix. We also expand the dressed Pascal matrices into subdiagonals

$$\Pi^{\pm 1} = I + \Lambda^{\top} \pi^{[\pm 1]} + (\Lambda^{\top})^2 \pi^{[\pm 2]} + \cdots$$

with $\pi^{[\pm n]} = \text{diag}(\pi_0^{[\pm n]}, \pi_1^{[\pm n]}, \dots)$. Then, for these subdiagonals we find

Proposition 4 (The dressed Pascal matrix coefficients). We have

(18)
$$\pi_n^{[\pm 1]} = \pm (n+1), \quad \pi_n^{[\pm 2]} = \frac{(n+2)(n+1)}{2} \pm p_{n+2}^1 (n+1) \mp (n+2) p_{n+1}^1 = \frac{(n+2)(n+1)}{2} \mp (n+1)\beta_{n+1} \mp p_{n+1}^1,$$

(19)

$$\pi_n^{[\pm 3]} = \pm \frac{(n+3)(n+2)(n+1)}{3} + \frac{(n+2)(n+1)}{2} p_{n+3}^1 - \frac{(n+3)(n+2)}{2} p_{n+1}^1 \\ \pm (n+1) p_{n+3}^2 \mp (n+3) p_{n+2}^2 \pm (n+3) p_{n+2}^1 p_{n+1}^1 \mp (n+2) p_{n+3}^1 p_{n+1}^1.$$

Moreover, the following relations are fulfill

(20)
$$\pi^{[1]} + \pi^{[-1]} = 0, \quad \pi^{[2]} + \pi^{[-2]} = 2D^{[2]}, \quad \pi^{[3]} + \pi^{[-3]} = 2((T_{-}^{2}S^{[1]})D^{[2]} - (T_{-}D^{[2]})S^{[1]}),$$
$$\pi^{[1]} - \pi^{[-1]} = 2D, \quad \pi^{[2]} - \pi^{[-2]} = 2((T_{-}S^{[1]})D - (T_{-}D)S^{[1]}),$$
$$\pi^{[3]} - \pi^{[-3]} = 2(D^{[3]} + (T_{-}S^{[2]})D - (T_{-}^{2}D)S^{[2]} + (T_{-}^{2}D)(T_{-}S^{[1]})S^{[1]} - (T_{-}^{2}S^{[1]})(T_{-}D)S^{[1]}).$$

Proof. From (11) we get

$$\begin{split} \Pi^{\pm 1} &= (I + \Lambda^{\top} S^{[1]} + (\Lambda^{\top})^2 S^{[2]} + \cdots) (I \pm \Lambda^{\top} D + (\Lambda^{\top})^2 D^{[2]} \pm (\Lambda^{\top})^3 D^{[3]} + \cdots) (I + \Lambda^{\top} S^{[-1]} + (\Lambda^{\top})^2 S^{[-2]} + \cdots) \\ &= I + (I + \Lambda^{\top} S^{[1]} + (\Lambda^{\top})^2 S^{[2]} + \cdots) (\pm \Lambda^{\top} D + (\Lambda^{\top})^2 D^{[2]} \pm (\Lambda^{\top})^3 D^{[3]} + \cdots) (I + \Lambda^{\top} S^{[-1]} + (\Lambda^{\top})^2 S^{[-2]} + \cdots) \\ &\text{so that} \end{split}$$

(21)
$$\Pi^{\pm 1} = I \pm \Lambda^{\top} D + (\Lambda^{\top})^{2} (D^{[2]} \pm (T_{-}S^{[1]})D \pm (T_{-}D)S^{[-1]}) + (\Lambda^{\top})^{3} (\pm D^{[3]} + (T_{-}^{2}S^{[1]})D^{[2]} + (T_{-}D^{[2]})S^{[-1]} \\ \pm (T_{-}S^{[2]})D \pm (T_{-}^{2}D)S^{[-2]} \pm (T_{-}^{2}S^{[1]})(T_{-}D)S^{[-1]}) + \cdots$$

From (21) and Proposition 2 we obtain

(22)
$$\pi^{[\pm 1]} = \pm D,$$

(23)
$$\pi^{[\pm 2]} = D^{[2]} \pm (T_- S^{[1]}) D \mp (T_- D) S^{[1]},$$

(24)
$$\pi^{[\pm 3]} = \pm D^{[3]} + (T_{-}^{2}S^{[1]})D^{[2]} - (T_{-}D^{[2]})S^{[1]} \pm (T_{-}S^{[2]})D \mp (T_{-}^{2}D)S^{[2]} \\ \pm (T_{-}^{2}D)(T_{-}S^{[1]})S^{[1]} \mp (T_{-}^{2}S^{[1]})(T_{-}D)S^{[1]}.$$

That component wise gives the desired result once we use the expressions for the β 's in (6).

Proposition 5. For any polynomial R(z) we have

(25)
$$R(\Lambda)B^{\pm 1} = B^{\pm 1}R(\Lambda \pm I), \qquad B^{\pm 1}R(\Lambda) = R(\Lambda \mp I)B^{\pm 1}, \\ \Pi^{\pm 1} = \Pi^{\pm 1}R(J \pm I), \qquad \Pi^{\pm 1}R(J) = R(J \mp I)\Pi^{\pm 1}.$$

Proof. The compatibility condition of

$$\begin{cases} B^{\pm 1}\chi(z) = \chi(z\pm 1), \\ R(\Lambda)\chi(z) = R(z)\chi(z). \end{cases}$$

reads,

$$R(\Lambda)B^{\pm 1}\chi(z) = R(\Lambda)\chi(z\pm 1) = R(z\pm 1)\chi(z\pm 1) = R(z\pm 1)B^{\pm 1}\chi(z) = B^{\pm 1}R(\Lambda\pm I)\chi(z),$$

so that $R(\Lambda)B^{\pm 1} = B^{\pm 1}R(\Lambda \pm I)$ and, consequently, $B^{\pm 1}R(\Lambda) = R(\Lambda \mp I)B^{\pm 1}$. Finally, a similarity transformation $\Lambda = S^{-1}JS$ gives the result.

2. DISCRETE ORTHOGONAL POLYNOMIALS AND DISCRETE PEARSON EQUATIONS

2.1. Discrete Pearson equation. We now assume that the functional is a sum of Dirac delta functions supported \mathbb{N}_0 ,

$$\rho = \sum_{k=0}^{\infty} \delta(z-k)w(k).$$

for some weight function w(z) with finite values w(k) for $k \in \mathbb{N}_0$. Hence, the bilinear form is $\langle F, G \rangle = \sum_{k=0}^{\infty} F(k)G(k)w(k)$. The moments are

(26)
$$\rho_n = \sum_{k=0}^{\infty} k^n w(k),$$

and, in particular, the 0-th moment reads as follows

(27)
$$\rho_0 = \sum_{k=0}^{\infty} w(k).$$

We will study families of weights are those satisfying the following *discrete Pearson equation*

(28)
$$\nabla(\sigma w) = \tau w,$$

that is $\sigma(k)w(k) - \sigma(k-1)w(k-1) = \tau(k)w(k)$, for $k \in \{1, 2, ...\}$, with $\sigma(z), \tau(z) \in \mathbb{R}[z]$ polynomials. If we write $\theta := \sigma - \tau$, the previous Pearson equation reads

(29)
$$\theta(k+1)w(k+1) = \sigma(k)w(k), \qquad k \in \mathbb{N}_0.$$

Theorem 1 (Hypergeometric symmetry of the moment matrix). Let the weight w be subject to the discrete Pearson equation (29), where θ, σ are polynomials with $\theta(0) = 0$. Then, the corresponding moment matrix fulfills

(30)
$$\theta(\Lambda)G = B\sigma(\Lambda)GB^{\top}.$$

Proof. The moment matrix is

 \sim

(31)
$$G = \sum_{k=0}^{\infty} \chi(k) \chi(k)^{\top} w(k)$$

Thus,

$$\begin{split} \theta(\Lambda)G &= \sum_{k=0}^{\infty} \theta(\Lambda)\chi(k)\chi(k)^{\top}w(k) & \text{use } (31) \\ &= \sum_{k=1}^{\infty} \chi(k)\chi(k)^{\top}\theta(k)w(k) & \text{use } \Lambda\chi = z\chi \text{ and } \theta(0) = 0 \\ &= \sum_{k=0}^{\infty} \chi(k+1)\chi(k+1)^{\top}\theta(k+1)w(k+1) & \text{shift the summation variable} \\ &= \sum_{k=0}^{\infty} \chi(k+1)\chi(k+1)^{\top}\sigma(k)w(k) & \text{use Pearson equation } (29) \\ &= \sum_{k=0}^{\infty} B\chi(k)\chi(k)^{\top}B^{\top}\sigma(k)w(k) & \text{use } (8) \\ &= \sum_{k=0}^{\infty} B\sigma(\Lambda)\chi(k)\chi(k)^{\top}w(k)B^{\top} & \text{use } \Lambda\chi = z\chi \text{ again} \\ &= B\sigma(\Lambda)GB^{\top} & \text{use } (31). \end{split}$$

Remark 3. This result extends to the case when θ and σ are entire functions, not necessarily polynomials, and we can ensure some meaning for $\theta(\Lambda)$ and $\sigma(\Lambda)$. Later on, see §2.5, we will show that this symmetry of the Gram matrix is a direct consequence of the generalized hypergeometric ODE satisfied by the the 0-th moment.

2.2. Consequences for the Jacobi matrix. We can use the Cholesky factorization (2) and the Jacobi matrix (4) to get

Proposition 6 (Symmetry of the Jacobi matrix). Let the weight w be subject to the discrete Pearson equation (29), where the functions θ , σ are entire functions, not necessarily polynomials, with $\theta(0) = 0$. Then,

(32)
$$\Pi^{-1}H\theta(J^{\top}) = \sigma(J)H\Pi^{\top}$$

Moreover, $H\theta(J^{\top})$ and $\sigma(J)H$ are symmetric matrices.

Proof. Given the Hankel property, $\Lambda G = G\Lambda^{\top}$, we write (30) as $G\theta(\Lambda^{\top}) = B\sigma(\Lambda)GB^{\top}$, and using the Cholesky factorization (2) we get $S^{-1}HS^{-\top}\theta(\Lambda^{\top}) = B\sigma(\Lambda)S^{-1}HS^{-\top}B^{\top}$, so that $HS^{-\top}\theta(\Lambda^{\top})S^{\top} = SBS^{-1}S\sigma(\Lambda)S^{-1}HS^{-\top}B^{\top}S^{\top}$, and we get Equation (32).

 \square

Let us prove that the matrices $H\theta(J^{\top})$ and $\sigma(J)H$ are symmetric. This fact follows from (5), $JH = HJ^{\top}$. Indeed,

$$(H\theta(J^{\top}))^{\top} = \theta(J)H = \theta(HJ^{\top}H^{-1})H = H\theta(J^{\top})H^{-1}H = H\theta(J^{\top}),$$

$$(\sigma(J)H)^{\top} = H\sigma(J^{\top}) = H\sigma(H^{-1}JH) = HH^{-1}\sigma(J)H = \sigma(J)H.$$

2.3. Generalized hypergeometric functions and the Pearson equation. Let us assume that θ, σ are polynomials, and denote their respective degrees by $N + 1 := \deg \theta(z)$ and $M := \deg \sigma(z)$. The roots of these polynomials are denoted by $\{-b_i + 1\}_{i=1}^N$ and $\{-a_i\}_{i=1}^M$. Following [23] we choose

$$\theta(z) = z(z+b_1-1)\cdots(z+b_N-1), \qquad \sigma(z) = \eta(z+a_1)\cdots(z+a_M).$$

Notice that we have normalized θ to be a monic polynomial, while σ is not monic, being the coefficient of the leading power denoted by η . This parameter η will be instrumental in what follows. Therefore, see [23], we have that the weight satisfying the Pearson equation (29) is proportional to

(33)
$$w(z) = \frac{(a_1)_z \cdots (a_M)_z}{\Gamma(z+1)(b_1)_z \cdots (b_N)_z} \eta^z,$$

where the Pochhammer symbol is $(\alpha)_z = \frac{\Gamma(\alpha + z)}{\Gamma(\alpha)}$, that for $z \in \mathbb{N}$ has the standard expression

$$(\alpha)_n = \alpha(\alpha+1)(\alpha+2)\cdots(\alpha+n-1), \qquad (\alpha)_0 = 1.$$

The moments of this weight are finite whenever, see [23] and references therein,

- i) $M \leq N$ and $\eta \in \mathbb{C}$.
- ii) $M > N, \eta \in \mathbb{C}$ and one or more of the parameters a_i is a nonpositive integer.
- iii) M = N + 1 and $|\eta| < 1$.
- iv) M = N + 1, $|\eta| = 1$ and $\operatorname{Re}(b_1 + \dots + b_{N-1} a_1 \dots a_M) > 0$.

According to (27) the 0-th moment

$$\rho_0 = \sum_{k=0}^{\infty} w(k) = \sum_{k=0}^{\infty} \frac{(a_1)_k \cdots (a_M)_k}{(b_1 + 1)_k \cdots (b_N + 1)_k} \frac{\eta^k}{k!}$$

= $_M F_N(a_1, \dots, a_M; b_1, \dots, b_N; \eta) = {}_M F_N \begin{bmatrix} a_1 & \cdots & a_M \\ b_1 & \cdots & b_N \end{bmatrix},$

is the generalized hypergeometric function, where we are using the two standard notations, see [15]. Then, according to (26), for $n \in \mathbb{N}$, the corresponding higher moments $\rho_n = \sum_{k=0}^{\infty} k^n w(k)$, are

(34)
$$\rho_n = \vartheta_\eta^n \rho_0 = \vartheta_\eta^n \Big({}_M F_N \begin{bmatrix} a_1 & \cdots & a_M \\ b_1 & \cdots & b_N \end{bmatrix} \Big), \qquad \qquad \vartheta_\eta := \eta \frac{\partial}{\partial \eta}$$

Given a function $f(\eta)$, we consider the Wronskian

$$\mathscr{W}_{n}(f) = \det \begin{pmatrix} f & \vartheta_{\eta}f & \vartheta_{\eta}^{2}f \cdots \cdots \vartheta_{\eta}^{k}f \\ \vartheta_{\eta}f & \vartheta_{\eta}^{2}f & \cdots & \vartheta_{\eta}^{k+1}f \\ \vartheta_{\eta}^{2}f & & \cdots & \vdots \\ \vdots & \ddots & \ddots & \vdots \\ \vartheta_{z}^{k}f & \vartheta_{\eta}^{k+1}f & & \vartheta_{\eta}^{2k}f \end{pmatrix}.$$

Then, we have that the Hankel determinants $\Delta_k = \det G^{[k]}$ determined by the truncations of the corresponding Gram matrix are Wronskians of generalized hypergeometric functions

(35)
$$\Delta_k = \tau_k, \qquad \tau_k := \mathscr{W}_k \Big({}_M F_N \begin{bmatrix} a_1 & \cdots & a_M \\ b_1 & \cdots & b_N \end{bmatrix} \Big),$$

(36)
$$\tilde{\Delta}_k = \vartheta_\eta \tau_k$$

We also have

(37)
$$H_k = \frac{\tau_{k+1}}{\tau_k}, \qquad p_k^1 = -\vartheta_\eta \log \tau_k$$

Remark 4. The functions τ_k are known in the theory of integrable systems as τ -functions.

2.4. The Laguerre-Freud structure matrix and the Cholesky factorization.

Theorem 2 (Laguerre–Freud structure matrix). Let us assume that the weight w is subject to the discrete Pearson equation (29) with θ, σ polynomials such that $\theta(0) = 0$, deg $\theta(z) = N + 1$, deg $\sigma(z) = M$. Then, the Laguerre–Freud structure matrix

(38)
$$\Psi := \Pi^{-1} H \theta(J^{\top}) = \sigma(J) H \Pi^{\top} = \Pi^{-1} \theta(J) H = H \sigma(J^{\top}) \Pi^{\top}$$

(39)
$$= \theta(J+I)\Pi^{-1}H = H\Pi^{\top}\sigma(J^{\top}-I),$$

has only N + M + 2 possibly nonzero diagonals (N + 1 superdiagonals, the main diagonal and M subdiagonals) $\Psi = (\Lambda^{\top})^{M} \psi^{(-M)} + \dots + \Lambda^{\top} \psi^{(-1)} + \psi^{(0)} + \psi^{(1)} \Lambda + \dots + \psi^{(N+1)} \Lambda^{N+1},$

for some diagonal matrices $\psi^{(k)}$. In particular, the lowest subdiagonal and highest superdiagonal are given by

(40)
$$\begin{cases} (\Lambda^{\top})^{M}\psi^{(-M)} = \eta(J_{-})^{M}H, \quad \psi^{(-M)} = \eta H \prod_{k=0}^{M-1} T_{-}^{k}\gamma = \eta \operatorname{diag}\left(H_{0} \prod_{k=1}^{M} \gamma_{k}, H_{1} \prod_{k=2}^{M+1} \gamma_{k}, \ldots\right), \\ \psi^{(N+1)}\Lambda^{N+1} = H(J_{-}^{\top})^{N+1}, \quad \psi^{(N+1)} = H \prod_{k=0}^{N} T_{-}^{k}\gamma = \operatorname{diag}\left(H_{0} \prod_{k=1}^{N+1} \gamma_{k}, H_{1} \prod_{k=2}^{N+2} \gamma_{k}, \ldots\right). \end{cases}$$

The vector P(z) of orthogonal polynomials fulfill the following structure equations

(41)
$$\theta(z)P(z-1) = \Psi H^{-1}P(z), \qquad \sigma(z)P(z+1) = \Psi^{\top}H^{-1}P(z).$$

Proof. To get (39) just use (25). Now, notice that the matrices $H\theta(J^{\top})$ and $\sigma(J)H$ are banded matrices with 2N + 3 diagonals (N + 1 superdiagonals and subdiagonals) and 2M + 1 (M superdiagonals and subdiagonals), respectively. Thus, $\Pi^{-1}H\theta(J^{\top})$ has at most N + 1 superdiagonals while $\sigma(J)H\Pi^{\top}$ has at most M subdiagonals. Consequently, Ψ given by (38) is a banded semi-infinite matrix as described. Then, (40) follow from (38) and the mentioned banded structure. To get the lowest subdiagonal we use $\Psi = \sigma(J)H\Pi^{\top}$, so that the lowest subdiagonal will come from the lowest subdiagonal of $\sigma(J)$, namely $\eta(J_{-})^{M}$ right multiplied by the diagonal matrix H and then right multiplied the main diagonal of Π^{\top} , which happens to be the identity. To get the highest superdiagonal we proceed similarly and use $\Psi = \Pi^{-1}H\theta(J^{\top})$, so that the main superdiagonal will come from the product of the three factors I, H and $\theta_N(J_{-}^{\top})^{N+1}$, in this order. The component wise expressions are direct computations.

Finally, we have

$$\begin{split} \Psi H^{-1} P(z) &= \Pi^{-1} \theta(J) H H^{-1} P(z) & \Psi^{\top} H^{-1} P(z) = \Pi \sigma(J) H H^{-1} P(z) \\ &= \Pi^{-1} \theta(J) P(z) &= \Pi \sigma(J) P(z) \\ &= \Pi^{-1} \theta(z) P(z) &= \Pi \sigma(z) P(z) \\ &= \theta(z) P(z-1), &= \sigma(z) P(z+1), \end{split}$$

Remark 5 (Laguerre–Freud equations). As we will see in the following sections (38) will be instrumental in obtaining non linear recurrences for the recursion coefficients

$$\gamma_{n+1} = F_1(n, \gamma_n, \gamma_{n-1}, \dots, \beta_n, \beta_{n-1}, \dots), \qquad \beta_{n+1} = F_2(n, \gamma_{n+1}, \gamma_n, \dots, \beta_n, \beta_{n-1}, \dots),$$

for some functions F_1, F_2 . These recurrences were named by Alphonse Magnus, attending to [33, 29], as Laguerre–Freud see [34, 35, 36, 37]. This is the reason for the given name to Ψ .

Proposition 7. The Laguerre–Freud and Jacobi matrices fulfill

$$\sigma(J)\theta(J+I) = \Psi H^{-1}\Psi^{\top}H^{-1}, \qquad \qquad \theta(J)\sigma(J-I) = \Psi^{\top}H^{-1}\Psi H^{-1}.$$

Proof. We have

$$\begin{split} \sigma(J)\theta(J+I)P(z) &= \sigma(z)\theta(z+1)P(z) = \sigma(z)\Psi H^{-1}P(z+1) = \Psi H^{-1}\sigma(z)P(z+1) \\ &= \Psi H^{-1}\Psi^{\top}H^{-1}P(z), \\ \theta(J)\sigma(J-I)P(z) &= \theta(z)\sigma(z-1)P(z) = \theta(z)\Psi^{\top}H^{-1}P(z-1) = \Psi^{\top}H^{-1}\theta(z)P(z-1) \\ &= \Psi^{\top}H^{-1}\Psi H^{-1}P(z), \end{split}$$

that hold for all $z \in \mathbb{C}$ and, consequently, imply the desired result.

Theorem 3 (Cholesky factorization). Let us assume that $H\theta(J^{\top})$ and $\sigma(J)H$ have the following Cholesky factorizations

(42)
$$H\theta(J^{\top}) = \Theta^{-1}H_{\theta}\Theta^{-\top}, \qquad \sigma(J)H = \Sigma^{-1}H_{\sigma}\Sigma^{-\top}.$$

with Θ and Σ lower unitriangular matrices and H_{θ} and H_{σ} diagonals matrices. Then,

- i) Θ^{-1} and Σ^{-1} have only first N+1 and M subdiagonals possibly nonzero, respectively.
- ii) We have

$$\Pi = \Theta^{-1}\Sigma, \qquad \qquad H_{\theta} = H_{\sigma} =: h.$$

iii) The Laguerre-Freud matrix has the following Gauss-Borel factorization

$$\Psi = \Sigma^{-1} h \Theta^{-\top}.$$

Proof. Recall that according to Proposition 6 the matrices $H\theta(J^{\top})$ and $\sigma(J)H$ are symmetric and, consequently, the corresponding Gauss–Borel factorizations, when they exists, will be Cholesky factorizations.

- i) If follows from the fact that J is tridiagonal and θ and σ polynomials of degrees N + 1 and M, respectively.
- ii) The symmetry (32) yields

$$\Pi^{-1}\Theta^{-1}H_{\theta}\Theta^{-\top} = \Sigma^{-1}H_{\sigma}\Sigma^{-\top}\Pi^{\top}$$

and given the uniqueness of the Gauss–Borel factorization the result follows.

iii) From (38) we have different alternatives to show the result, let us see two of them

$$\Psi = \Pi^{-1} H \theta(J^{\top}) = \Sigma^{-1} \Theta \Theta^{-1} H_{\theta} \Theta^{-\top} = \Sigma^{-1} h \Theta^{-\top}, \quad \Psi = \sigma(J) H \Pi^{\top} = \Sigma^{-1} H_{\sigma} \Sigma^{-\top} \Sigma^{\top} \Theta^{-1} = \Sigma^{-1} h \Theta^{-\top}.$$

The compatibility with the recursion relation, i.e. eigenfunctions of the Jacobi matrix, and the recursion matrix leads to some interesting equations.

Proposition 8. The following compatibility conditions for the Laguerre–Freud and Jacobi matrices hold

(43a)
$$[\Psi H^{-1}, J] = \Psi H^{-1},$$

(43b) $[J, \Psi^{\top} H^{-1}] = \Psi^{\top} H^{-1}.$

Proof. To prove (43a), we evaluate the eigenvalue equation JP(z) = zP(z) in z - 1 to get JP(z - 1) = (z-1)P(z-1). Now multiply it by $\theta(z)$ to obtain $J\theta(z)P(z-1) = (z-1)\theta(z)P(z-1)$. Therefore, recalling (41), $J\Psi H^{-1}P(z) = (z-1)\Psi H^{-1}P(z) = \Psi H^{-1}(J-I)P(z)$ from where the relation follows.

Alternatively, observe that

$$\Psi H^{-1} = SB^{-1}\theta(\Lambda)S^{-1}$$

from where

(44)

$$[\Psi H^{-1}, J] = S[B^{-1}\theta(\Lambda), \Lambda]S^{-1} = S[B^{-1}, \Lambda]\theta(\Lambda)S^{-1} = SB^{-1}\theta(\Lambda)S^{-1} = \Psi H^{-1}.$$

To show (43b) we take the transpose of the proved one to get $[J^{\top}, H^{-1}\Psi^{\top}] = H^{-1}\Psi^{\top}$ so that

$$[HJ^{\top}H^{-1}, HH^{-1}\Psi^{\top}H^{-1}] = HH^{-1}\Psi^{\top}H^{-1}$$

and recalling that $HJ^{\top}H^{-1} = J$, see (5), we get the desired result.

2.5. **Contiguous hypergeometric relations**. As we have seen, the polynomial discrete Pearson equation leads to (34), and the Hankel determinants are Wronskians of a generalized hypergeometric function, see (35) and (36). Hence, we expect that some properties of generalized hypergeometric functions may translate to the Gram matrix. To write this dictionary we require of Proposition 11, discussed latter, that ensures,

$$\vartheta_{\eta}G = \Lambda G = G\Lambda^{\top}.$$

We now study three important relations fulfilled by the generalized hypergeometric function, namely:

(45)
$$(\vartheta_{\eta} + a_{i})_{M} F_{N} \begin{bmatrix} a_{1} \cdots a_{i} \cdots a_{M} \\ b_{1} \cdots b_{N} \end{bmatrix} = a_{i M} F_{N} \begin{bmatrix} a_{1} \cdots a_{i} + 1 \cdots a_{M} \\ b_{1} \cdots b_{N} \end{bmatrix},$$

(46)
$$(\vartheta_{\eta} + b_j - 1)_M F_N \begin{bmatrix} a_1 \cdots a_M \\ b_1 \cdots b_j \cdots b_N; \eta \end{bmatrix} = (b_j - 1)_M F_N \begin{bmatrix} a_1 \cdots a_M \\ b_1 \cdots b_j - 1 \cdots b_N; \eta \end{bmatrix},$$

(47)
$$\frac{\mathrm{d}}{\mathrm{d}\eta} {}_{M}F_{N}\begin{bmatrix}a_{1} & \cdots & a_{M}\\b_{1} & \cdots & b_{N}\end{bmatrix} = \kappa {}_{M}F_{N}\begin{bmatrix}a_{1}+1 & \cdots & a_{M}+1\\b_{1}+1 & \cdots & b_{N}+1\end{bmatrix}, \qquad \kappa := \frac{\prod_{i=1}^{M}a_{i}}{\prod_{j=1}^{N}b_{j}}$$

From these three equations we also derive

(48)
$$\eta \prod_{n=1}^{M} \left(\eta \frac{\mathrm{d}}{\mathrm{d}\eta} + a_n \right) u = \eta \frac{\mathrm{d}}{\mathrm{d}\eta} \prod_{n=1}^{N} \left(\eta \frac{\mathrm{d}}{\mathrm{d}\eta} + b_n - 1 \right) u, \qquad u := {}_M F_N \begin{bmatrix} a_1 & \cdots & a_M \\ b_1 & \cdots & b_N \end{bmatrix},$$

In (45) and (46) we have basic relations between contiguous generalized hypergeometric functions and its derivatives.

For the analysis of these equations let us introduce the shift operators in the parameters $\{a_i\}_{i=1}^M$ and $\{b_j\}_{j=1}^N$. Thus, given a function $f\begin{bmatrix}a_1 & \cdots & a_M\\b_1 & \cdots & b_N\end{bmatrix}$ of these parameters we introduce the shifts $_iT$ and T_j as follows

$${}_{i}Tf\begin{bmatrix}a_{1}\cdots a_{i}\cdots a_{M}\\b_{1}\cdots b_{N}\end{bmatrix} = f\begin{bmatrix}a_{1}\cdots a_{i}+1\cdots a_{M}\\b_{1}\cdots b_{N}\end{bmatrix}, \qquad T_{j}f\begin{bmatrix}a_{1}\cdots a_{M}\\b_{1}\cdots b_{j}\cdots b_{N}\end{bmatrix} = f\begin{bmatrix}a_{1}\cdots a_{M}\\b_{1}\cdots b_{j}-1\cdots b_{N}\end{bmatrix}$$

and a total shift $T = {}_{1}T\cdots {}_{M}TT_{1}^{-1}\cdots T_{N}^{-1}$; i.e,

$$Tf\begin{bmatrix}a_1 & \cdots & a_M\\b_1 & \cdots & b_N\end{bmatrix} := f\begin{bmatrix}a_1+1 & \cdots & a_M+1\\b_1+1 & \cdots & b_N+1\end{bmatrix}$$

Then, we find:

Theorem 4 (Hypergeometric relations). The moment matrix $G = (\rho_{n+m})_{n,n \in \mathbb{N}_0}$ of a weight satisfying (29) fulfills the following hypergeometric relations

(49a)
$$(\Lambda + a_i I)G = a_i \, _i TG,$$

(49b)
$$(\Lambda + (b_j - 1)I)G = (b_j - 1)T_jG,$$

(49c)
$$\Lambda G = \kappa B(TG)B^{\top}$$

with B given in (8).

Proof. We first prove (49a). For that aim we use that $G_{n,m} = \vartheta_{\eta}^{n+m} \left({}_{M}F_{N} \begin{bmatrix} a_{1} & \cdots & a_{M} \\ b_{1} & \cdots & b_{N} \end{bmatrix} \right)$ so that

$$(\vartheta_{\eta} + a_{i})G_{n,m} = (\vartheta_{\eta} + a_{i})\vartheta_{\eta}^{n+m} \Big({}_{M}F_{N} \begin{bmatrix} a_{1} \cdots a_{M} \\ b_{1} \cdots b_{N} \end{bmatrix} \Big)$$
$$= \vartheta_{\eta}^{n+m} (\vartheta_{\eta} + a_{i}) \Big({}_{M}F_{N} \begin{bmatrix} a_{1} \cdots a_{M} \\ b_{1} \cdots b_{N} \end{bmatrix} \Big)$$
$$= a_{i}\vartheta_{\eta}^{n+m} \Big({}_{M}F_{N} \begin{bmatrix} a_{1} \cdots a_{i} + 1 \cdots a_{M} \\ b_{1} \cdots b_{N} \end{bmatrix} \Big)$$
$$= a_{i} {}_{i}TG_{n m}.$$

Thus, $(\vartheta_{\eta} + a_i)G = a_i {}_iTG$, and recalling $\vartheta_{\eta}G = \Lambda G$ we get the result. Relation (49b) is proved similarly. Finally, (49b) follows from (47), the novelty is that in (47) we have $\frac{d}{d\eta} = \eta^{-1}\vartheta_{\eta}$, which do not commute with ϑ_{η} . Observe that the relation

$$\frac{\mathrm{d}}{\mathrm{d}\,\eta}\eta - \eta\frac{\mathrm{d}}{\mathrm{d}\,\eta} = 1$$

can be written $\eta^{-1}\vartheta_\eta\eta = \vartheta_\eta + 1$. Hence, $\eta^{-1}\vartheta_\eta^n\eta = (\eta^{-1}\vartheta_\eta\eta)^n = (\vartheta_\eta + 1)^n$, that, in turn, implies

(50)
$$\frac{\mathrm{d}}{\mathrm{d}\,\eta}\vartheta_{\eta}^{n} = (\vartheta_{\eta} + 1)^{n}\frac{\mathrm{d}}{\mathrm{d}\,\eta},$$

Thus,

$$\begin{aligned} \frac{\mathrm{d}\,G_{n,m}}{\mathrm{d}\,\eta} &= \frac{\mathrm{d}}{\mathrm{d}\,\eta}\vartheta_{\eta}^{n+m} \Big({}_{M}F_{N} \begin{bmatrix} a_{1} & \cdots & a_{M} \\ b_{1} & \cdots & b_{N} \end{bmatrix} \Big) = (\vartheta_{\eta}+1)^{n+m} \frac{\mathrm{d}}{\mathrm{d}\,\eta} \Big({}_{M}F_{N} \begin{bmatrix} a_{1} & \cdots & a_{M} \\ b_{1} & \cdots & b_{N} \end{bmatrix} \Big) \\ &= (\vartheta_{\eta}+1)^{n+m} \kappa_{M}F_{N} \begin{bmatrix} a_{1}+1 & \cdots & a_{M}+1 \\ b_{1}+1 & \cdots & b_{N}+1 \end{bmatrix}; \\ &= \kappa \sum_{k=0}^{n+m} \binom{n+m}{k} \vartheta_{\eta}^{k}{}_{M}F_{N} \begin{bmatrix} a_{1}+1 & \cdots & a_{M}+1 \\ b_{1}+1 & \cdots & b_{N}+1 \end{bmatrix}; \end{aligned}$$

Using the Chu-Vandermonde identity

$$\binom{n+m}{k} = \sum_{l=0}^{k} \binom{n}{l} \binom{m}{k-l},$$

we find

$$\frac{\mathrm{d}\,G_{n,m}}{\mathrm{d}\,\eta} = \kappa \sum_{k=0}^{n+m} \sum_{l=0}^{k} \binom{n}{l} \binom{m}{k-l} \vartheta_{\eta}^{l+k-l}{}_{M}F_{N} \begin{bmatrix} a_{1}+1 & \cdots & a_{M}+1\\ b_{1}+1 & \cdots & b_{N}+1 \end{bmatrix}$$
$$= \kappa \sum_{k=0}^{n+m} \sum_{l=0}^{k} \binom{n}{l} \binom{m}{k-l} TG_{l,k-l}$$
$$= \kappa \sum_{l=0}^{n} \sum_{k=0}^{m} \binom{n}{l} TG_{l,k} \binom{m}{k}$$

from where the result follows.

Finally, using $\vartheta_{\eta} - 1 = \eta \vartheta_{\eta} \eta^{-1}$, we get $(\vartheta_{\eta} - 1)^n \eta = \eta (\vartheta_{\eta})^n$ and we workout (48) as follows $\eta \prod_{n=1}^{M} (\vartheta_{\eta} + a_n) G_{n,m} = \eta \prod_{n=1}^{M} (\vartheta_{\eta} + a_n) \vartheta_{\eta}^{n+m} \Big({}_{M}F_N \begin{bmatrix} a_1 & \cdots & a_M \\ b_1 & \cdots & b_N \end{bmatrix} \Big)$ $= (\vartheta_{\eta} - 1)^{n+m} \eta \prod_{n=1}^{M} (\vartheta_{\eta} + a_n) \Big({}_{M}F_N \begin{bmatrix} a_1 & \cdots & a_M \\ b_1 & \cdots & b_N \end{bmatrix} \Big)$ $= \vartheta_{\eta} \prod_{n=1}^{N} (\vartheta_{\eta} + b_n - 1) (\vartheta_{\eta} - 1)^{n+m} \Big({}_{M}F_N \begin{bmatrix} a_1 & \cdots & a_M \\ b_1 & \cdots & b_N \end{bmatrix} \Big)$ $= \vartheta_{\eta} \prod_{n=1}^{N} (\vartheta_{\eta} + b_n - 1) \sum_{k=0}^{n+m} {n+m \choose k} (-1)^k \vartheta_{\eta}^k \Big({}_{M}F_N \begin{bmatrix} a_1 & \cdots & a_M \\ b_1 & \cdots & b_N \end{bmatrix} \Big)$ $= \vartheta_{\eta} \prod_{n=1}^{N} (\vartheta_{\eta} + b_n - 1) \sum_{l=0}^{n+m} {n+m \choose k} (-1)^l {n \choose l} G_{l,k} (-1)^k {m \choose k},$

and, consequently, we finally deduce

$$\sigma(\Lambda)G = B^{-1}\theta(\Lambda)GB^{-\top},$$

and result follows at once.

Remark 6. From (48) we derive, in an alternative manner, the relation (30).

2.6. **Discrete lattice Toda equation**. The shifts in the hypergeometric parameters induce corresponding transformations on the discrete orthogonal polynomials. In order to describe them we introduce the following semi-infinite matrices

$$_{i}\Omega := S(_{i}TS)^{-1}, \qquad i \in \{1, \dots, M\}, \qquad \Omega_{k} := S(T_{k}S)^{-1}, \qquad k \in \{1, \dots, N\}$$

 $i \in \{1, \ldots, M\},\$

so that the following connection formulas are fulfilled

(51)
$${}_{i}\Omega {}_{i}TP(z) = P(z),$$

(52)
$$\Omega_j T_j P(z) = P(z), \qquad j \in \{1, \dots, N\}.$$

In [39] we proved that

The connection formulas (51) and (52) for contiguous hypergeometric parameters imply a nonlinear compatibility equations that leads to a generalized lattice Toda equation found in [41]. We use the following notation

$$u_n = H_{n-1} \begin{bmatrix} a_1 & \cdots & a_M \\ b_1 & \cdots & b_N \end{bmatrix}$$

We now consider three variables (n, r, s), where r, s are any couple of variables taken from the set of hypergeometric parameters $\{a_1, \ldots, a_M, b_1, \ldots, b_N\}$ and denote the corresponding "shifts" in n, r, s as follows

$$\bar{u}_n(r,s) = u_{n+1}(r,s),$$
 $\hat{u}_n(r,s) = \hat{a}\hat{u}_n(r\pm 1,s),$ $\tilde{u}_n(r,s) = \tilde{a}u(r,s\pm 1),$

where the \pm signs is a + if the corresponding variable belongs to $\{a_i\}_{i=1}^M$ or a - if it belongs to $\{b_j\}_{j=1}^M$, and the constants \hat{a}, \tilde{a} are taken from $\{a_i\}_{i=1}^N$ and $\{b_j - 1\}_{j=1}^N$ according to the choice selected for the variables r and s. We denote by $\Omega^{(r)}$ and $\Omega^{(s)}$, the corresponding matrices Ω 's taken from $\{i\Omega\}_{i=1}^M$ and $\{\Omega_j\}_{j=1}^N$, depending on the variables picked r and s.

Theorem 5 (Nijhoff–Capel discrete lattice Toda equations). The squared norms satisfy the following nonlinear equations linking contiguous hypergeometric parameters

(53)
$$\frac{\hat{\bar{u}} - \tilde{\bar{u}}}{\bar{u}} = \tilde{\bar{u}} \left(\frac{1}{\tilde{u}} - \frac{1}{\hat{u}} \right).$$

Proof. The connection formulas are

$$\Omega^{(r)}\hat{P} = P, \qquad \qquad \Omega^{(s)}\hat{P} = P.$$

Then, we have

$$\tilde{\Omega}^{(r)}\tilde{\hat{P}}=\tilde{P},$$


so that

$$\Omega^{(s)}\tilde{\Omega}^{(r)}\hat{P} = \Omega^{(s)}\tilde{P} = P$$

and the compatibility $\hat{\hat{P}} = \hat{\hat{P}}$ leads to the nonlinear condition

(54)
$$\Omega^{(s)}\tilde{\Omega}^{(r)} = \Omega^{(r)}\hat{\Omega}^{(s)}.$$

Then, as we can write



Equation (54) is

$$\begin{pmatrix} 1 & 0 & \cdots & \cdots \\ \frac{\bar{u}_1}{\hat{u}_1} & 1 & \ddots \\ 0 & \frac{\bar{u}_2}{\hat{u}_2} & \cdots & \ddots \\ \vdots & \ddots & \ddots & \ddots \\ \vdots & \ddots & \ddots & \ddots \end{pmatrix} \begin{pmatrix} 1 & 0 & \cdots & \cdots & \vdots \\ \frac{\bar{u}_1}{\hat{u}_1} & 1 & \ddots & \vdots \\ 0 & \frac{\bar{u}_2}{\hat{u}_2} & \cdots & \ddots \\ \vdots & \ddots & \ddots & \ddots & \ddots \end{pmatrix} = \begin{pmatrix} 1 & 0 & \cdots & \cdots & \vdots \\ \frac{\bar{u}_1}{\hat{u}_1} & 1 & \ddots & \vdots \\ 0 & \frac{\bar{u}_2}{\hat{u}_2} & \cdots & \ddots \\ \vdots & \ddots & \ddots & \ddots & \ddots \end{pmatrix} \begin{pmatrix} 1 & 0 & \cdots & \cdots & \vdots \\ \frac{\bar{u}_1}{\hat{u}_1} & 1 & \ddots & \vdots \\ 0 & \frac{\bar{u}_2}{\hat{u}_2} & \cdots & \ddots \\ \vdots & \ddots & \ddots & \ddots & \ddots \end{pmatrix} \begin{pmatrix} 1 & 0 & \cdots & \cdots & \vdots \\ \frac{\bar{u}_1}{\hat{u}_1} & 1 & \ddots & \vdots \\ 0 & \frac{\bar{u}_2}{\hat{u}_2} & \cdots & \ddots \\ \vdots & \ddots & \ddots & \ddots & \ddots \end{pmatrix} \end{pmatrix},$$
 and, consequently, we deduce

$$\frac{\bar{u}}{\tilde{u}} + \frac{\bar{u}}{\tilde{\hat{u}}} = \frac{\bar{u}}{\hat{u}} + \frac{\bar{u}}{\hat{\tilde{u}}}$$

and the result follows immediately.

Remark 7. Equation (53) is the generalized discrete lattice Toda equation described for the first time by Nijhoff and Capel [41], that taking adequate continuous limits in the r and s variables (hypergeometric parameters) recovers the 2D Toda equation. This is a canonical equation among the difference equations in three variables involving up to second order differences of octahedral type, consistent on the 4D lattice [1], it appears as the type V of the octahedron type integrable discrete equations in §3.9 of the book [30].

Now we study compatibility of the recursion relation and the connection formula for contiguous parameters; i.e., the compatibility of

$$zP = JP,$$

 $\hat{P} = \hat{a} \frac{\omega^{(r)}}{z+r} P.$

These pair of relations lead to

$$\hat{J}\hat{P}(z) = z\hat{P}(z) \qquad \Rightarrow \qquad \hat{J}\frac{\omega^{(r)}}{z+r}P(z) = z\frac{\omega^{(r)}}{z+r}P(z) = \frac{\omega^{(r)}}{z+r}JP(z),$$

and therefore we find the following compatibility equation

$$\hat{J}\omega^{(r)} = \omega^{(r)}J.$$

Using the notation

$$u_n = H_{n-1}, \qquad \qquad v_n = \beta_{n-1}$$

so that $\gamma_n = \frac{\bar{u}_n}{u_n}$ and we have the following expressions

$$J = \begin{pmatrix} v_1 & 1 & 0 & \cdots & \cdots & \vdots \\ \frac{\tilde{u}_1}{u_1} & v_2 & 1 & 0 & \cdots & \cdots \\ 0 & \ddots & \ddots & \ddots & \ddots & \vdots \\ \vdots & \ddots & \ddots & \ddots & \ddots & \ddots \\ \vdots & \ddots & \ddots & \ddots & \ddots & \ddots \\ \vdots & \ddots & \ddots & \ddots & \ddots & \ddots \end{pmatrix}, \qquad \qquad \omega^{(r)} = \begin{pmatrix} \frac{\hat{u}_1}{u_1} & 1 & 0 & \cdots & \cdots \\ 0 & \frac{\hat{u}_2}{u_2} & 1 & \ddots & \vdots \\ \vdots & \ddots & \ddots & \ddots & \ddots & \ddots \\ \vdots & \ddots & \ddots & \ddots & \ddots & \ddots \end{pmatrix},$$

the compatibility reads

and we find $\hat{v}_1\frac{\hat{u}_1}{u_1}=v_1\frac{\hat{u}_1}{u_1}+\frac{\bar{u}_1}{u_1}$ and

$$\begin{aligned} & \frac{\hat{\bar{u}}_n}{\hat{\bar{u}}_n} + \hat{\bar{v}}_n \frac{\hat{\bar{u}}_n}{\bar{\bar{u}}_n} = \frac{\bar{\bar{u}}_n}{\bar{\bar{u}}_n} + \bar{v}_n \frac{\hat{\bar{u}}_n}{\bar{\bar{u}}_n} \\ & \hat{v}_n + \frac{\hat{\bar{u}}_n}{\bar{\bar{u}}_n} = \bar{v}_n + \frac{\hat{u}_n}{u_n} \end{aligned}$$

for $n \in \mathbb{N}$. Thus, we find

Proposition 9. The squared norms $u_n = H_{n-1}$ and the recursion coefficients $v_n = \beta_{n-1} = p_{n-1}^1 - p_n^1$, satisfy the following system of nonlinear difference equations

$$\hat{\bar{v}} - \bar{v} = \frac{\bar{\bar{u}}}{\hat{\bar{u}}} - \frac{\bar{u}}{\hat{u}},$$
$$\hat{v} - \bar{v} = \frac{\hat{u}}{u} - \frac{\hat{\bar{u}}}{\bar{u}}.$$

Finally, recalling $\vartheta_{\eta}P = \Phi P$ and $\Omega \hat{P} = P$, we get

$$\vartheta_{\eta}\Omega = \Phi\Omega - \Omega\tilde{\Phi},$$

so that

Proposition 10. The squared norms $u_n = H_{n-1}$ fulfill

$$\vartheta_\eta \left(\frac{\bar{u}}{\hat{u}}\right) = \frac{\bar{u}}{u} - \frac{\hat{\bar{u}}}{\hat{u}}.$$

2.7. The Toda flows. Given a semi-infinite vector $\boldsymbol{\eta} := \{\eta_l\}_{l=1}^{\infty} \in \mathbb{C}^{\infty}$, for each $z \in \mathbb{C}$ we define $\mathscr{E}(z; \boldsymbol{\eta}) := \prod_{l=1}^{\infty} \eta_l^{z^l}$ and assume we assume a weight of the form $w(z) = v(z)\mathscr{E}(z; \boldsymbol{\eta})$, with $v \eta$ -independent, so that the corresponding moment matrix is

$$G = \sum_{k=0}^{\infty} \chi(k) \chi(k)^{\top} v(k) \mathscr{E}(k; \boldsymbol{\eta}).$$

Observe that only this first flow preserves the Pearson reduction. Notice also that $\mathscr{E} = e^{\sum_{l=1}^{\infty} t_l z^l}$, with $t_l := \log \eta_l$ the standard time flows in an integrable hierarchy. We will write $\eta_1 = \eta$, normally t_1 is denoted by $x, t_1 = x$. We will use

$$\vartheta_l := \eta_l \frac{\partial}{\partial \eta_l} = \frac{\partial}{\partial t_l}$$

In particular $\vartheta_1 = \vartheta_{\eta}$. To ensure the converge of the corresponding series for the moments we require $|\eta_k| \leq 1$ for $k \in \{2, 3, ...\}$.

If we take $\eta_k = 1, k \in \mathbb{N}$, then the corresponding Gram matrix is

$$G_0 = \sum_{k=0}^{\infty} \chi(k) \chi(k)^{\top} v(k).$$

Moreover, we can write for the deformed Gram matrix

$$G = \mathscr{E}(\Lambda; \boldsymbol{\eta}) G_0 = G_0 \mathscr{E}(\Lambda^\top; \boldsymbol{\eta}), \qquad \qquad \mathscr{E}(\Lambda; \boldsymbol{\eta}) := \prod_{l=1}^{\infty} \eta_l^{\Lambda^l} = \exp\big(\sum_{l=1}^{\infty} t_l \Lambda^l\big).$$

Proposition 11. For a functional of the form $\rho = \sum_{k=0}^{\infty} \delta(z-k)v(z)\mathscr{E}(z;\eta)$, given $l \in \mathbb{N}$, the Gram matrix satisfies

(55)
$$\vartheta_l G = (\Lambda)^l G = G(\Lambda^\top)^l.$$

Proof. It follows form $\vartheta_l \mathscr{E}(z; \boldsymbol{\eta}) = z^l \mathscr{E}(z; \boldsymbol{\eta})$, indeed

$$\vartheta_l G = \sum_{k=0}^{\infty} \chi(k) \chi(k)^\top k^l v(k) \mathscr{E}(k; \boldsymbol{\eta}) = \Lambda^l G = G(\Lambda^\top)^l.$$

For $k \in \mathbb{N}$, let us define the strictly lower triangular matrix (strictly because it has zeros on the main diagonal)

$$\Phi_k := (\vartheta_k S) S^{-1}.$$

In particular, we denote $\Phi := \Phi_1 = (\vartheta_\eta S)S^{-1}$.

Proposition 12. The semi-infinite vector P fulfills

(56)
$$\vartheta_k P = \Phi_k P.$$

Proof. As $P = S\chi$ then $\vartheta_k P = (\vartheta_k S)\chi = (\vartheta_k S)S^{-1}S\chi = \Phi P$.

Proposition 13 (Sato–Wilson equations). The following equations holds

(57)
$$-\Phi_k H + \vartheta_k H - H \Phi_k^\top = J^k H, \qquad k \in \mathbb{N}$$

Consequently, for $k \in \mathbb{N}$ and $n \in \mathbb{N}_0$ we have

$$\Phi_k = -(J^k)_{-}, \qquad \qquad \vartheta_k \log H_n = (J^k)_{n,n}$$

Proof. Using the Cholesky factorization (2) for the moment matrix we get

$$\vartheta_k G = \vartheta_k \left(S^{-1} H S^{-\top} \right) = -S^{-1} (\vartheta_k S) S^{-1} H S^{-\top} + S^{-1} (\vartheta_k H) S^{-\top} - S^{-1} H S^{-\top} (\vartheta_k S)^{\top} S^{-\top}$$

Hence, from the symmetry relation (55) we deduce

$$-S^{-1}(\vartheta_{\eta}S)S^{-1}HS^{-\top} + S^{-1}(\vartheta_{\eta}H)S^{-\top} - S^{-1}HS^{-\top}(\vartheta_{\eta}S)^{\top}S^{-\top} = \Lambda^{k}S^{-1}HS^{-\top},$$

from where

$$-(\vartheta_k S)S^{-1}H + \vartheta_k H - H((\vartheta_k S)S^{-1})^\top = J^k H,$$

for $J = S\Lambda S^{-1}$, and the result follows.

For k = 1; i.e. for the first flow $\eta_1 = \eta$, we have formulas (35), (36) and (37) in terms of the τ -function. Moreover,

Proposition 14 (Toda). The following equations hold

(58a)

$$\Phi = (\vartheta_{\eta}S)S^{-1} = -\Lambda^{\top}\gamma$$
(58b)

$$(\vartheta_{\eta}H)H^{-1} = \beta.$$

(58b)
$$(\vartheta_{\eta}H)H^{-1} = \mu$$

In particular, for $n, k-1 \in \mathbb{N}$, we have

(59a)
$$\vartheta_{\eta}p_{n}^{1} = -\gamma_{n}, \quad \vartheta_{\eta}p_{n+k}^{k} = -\gamma_{n+k}p_{n+k-1}^{k-1},$$

(59b) $\vartheta_{\eta}\log H_{n} = \beta_{n}.$

The functions $q_n := \log H_n$, $n \in \mathbb{N}$, satisfy the Toda equations

 $\vartheta_n^2 q_n = \mathrm{e}^{q_{n+1}-q_n} - \mathrm{e}^{q_n-q_{n-1}} \,.$ (60)

For $n \in \mathbb{N}$, we also have

$$\vartheta_{\eta} P_n(z) = -\gamma_n P_{n-1}(z).$$

Proof. Equation (58a) is a direct consequence of (57) for k = 1. To obtain (59a) we we write $\vartheta_{\eta}S = -\Lambda^{\top}\gamma S$ in terms of its subdiagonals, i.e.

$$\vartheta_{\eta} \left(I + \Lambda^{\top} S^{[1]} + (\Lambda^{\top})^2 S^{[2]} + \cdots \right) = -\Lambda^{\top} \gamma \left(I + \Lambda^{\top} S^{[1]} + (\Lambda^{\top})^2 S^{[2]} + \cdots \right)$$

that, for $k \in \mathbb{N}$, gives

$$\vartheta_{\eta} S^{[k]} = -(T^{k-1}_{-}\gamma) S^{[k-1]}$$

with $S^{[0]} = I$. Now, recalling (10) we get (59a). Equation (59b) follows component wise from (58b). As $\beta_n = p_n^1 - p_{n+1}^1$ and $\gamma_n = \frac{H_n}{H_{n-1}}$ we deduce that

$$\vartheta_{\eta}^2 \log H_n = \vartheta_{\eta} p_n^1 - \vartheta_{\eta} p_{n+1}^1 = -\frac{H_n}{H_{n-1}} + \frac{H_{n+1}}{H_n},$$

that is the Toda equation (60) for $H_n = e^{q_n}$. The last equation follows form $\vartheta_\eta P = \Phi P = -\Lambda^\top \gamma P$.

Given the τ -function (35) we find

Proposition 15 (τ -function expressions). In terms of the τ -function

$$\tau_n := \mathscr{W}_n\Big({}_M F_N \begin{bmatrix} a_1 & \cdots & a_M \\ b_1 & \cdots & b_N \end{bmatrix}\Big),$$

we find

(61)
$$H_n = \frac{\tau_{n+1}}{\tau_n}, \qquad p_n^1 = -\vartheta_\eta \log \tau_n, \qquad \gamma_n = \vartheta_\eta^2 \log \tau_n, \qquad \beta_n = \vartheta_\eta \log \frac{\tau_{n+1}}{\tau_n}$$

Proof. Collect together (37) and (59a).

Proposition 16. The following Lax equation holds

$$\vartheta_{\eta}J = [J_+, J].$$

The recursion coefficients satisfy the following Toda system

(62a) $\vartheta_{\eta}\beta_{n} = \gamma_{n+1} - \gamma_{n},$

(62b) $\vartheta_{\eta} \log \gamma_n = \beta_n - \beta_{n-1},$

for $n \in \mathbb{N}_0$ and $\beta_{-1} = 0$. Consequently, we get

(63)
$$\vartheta_{\eta}^{2}\log\gamma_{n} + 2\gamma_{n} = \gamma_{n+1} + \gamma_{n-1}.$$

Moreover,

$$\vartheta_{\eta}^2 \log \tau_n = \frac{\tau_{n+1}\tau_{n-1}}{\tau_n^2}$$

Proof. As $J = S\Lambda S^{-1}$ is clear that,

$$\vartheta_{\eta}J = [\Phi, J] = [-J_{-}, J] = [J_{+} - J, J] = [J_{+}, J]$$

From this Lax equation one could derive, component-wise, the Toda system (62). Alternatively, the Toda system (62) could be derived as follows. From (6) and (59a) we obtain

$$\vartheta_{\eta}\beta_{n} = \vartheta_{\eta}p_{n}^{1} - \vartheta_{\eta}p_{n+1}^{1} = -\gamma_{n} + \gamma_{n+1},$$
$$\vartheta_{\eta}\log\gamma_{n} = \vartheta_{\eta}\log H_{n} - \vartheta_{\eta}\log H_{n-1} = \beta_{n} - \beta_{n-1}.$$

Finally,

$$\vartheta_{\eta}^2 \log \tau_n = \gamma_n = \frac{H_n}{H_{n-1}} = \frac{\tau_{n+1}\tau_{n-1}}{\tau_n}.$$

For higher Toda flows, that generically are not consistent with the hypergeometric reduction, we have **Proposition 17** (Lax equations). *The following Lax equation holds*

$$\vartheta_k J = [(J^k)_+, J].$$

Proof. As $J = S\Lambda S^{-1}$ we have

$$\vartheta_k J = [\Phi_k, J] = [-(J)^k_{-}, J] = [(J^k)_{+} - J^k, J] = [(J^k)_{+}, J]$$

Proposition 18 (Wave functions and Zakharov-Shabat equations). The wave function satisfies the linear system

$$\vartheta_k \Psi = (J^k)_+ \Psi$$

Moreover, the following Zakharov-Shabat equations hold

$$\vartheta_k(J^l)_+ - \vartheta_l(J^k)_+ + [(J^l)_+, (J^k)_+] = 0.$$

Proof. The orthogonal polynomials we have

$$\vartheta_k P = \Phi_k P = ((J^k)_+ - J^k) P = ((J^k)_+ - z^k) P$$

Thus, the wave function fulfills the following linear system

$$\vartheta_k \Psi = \vartheta_k(P)\mathscr{E} + P\vartheta_k(\mathscr{E}) = (J^k)_+ \Psi.$$

Finally, the Zakharov–Shabat equations follow as the compatibility conditions for the previous linear system.

Proposition 19. Let us assume that G_0 , the Gram matrix of the functional $\rho = \sum_{k=0}^{\infty} \delta(z-k)v(z)$, admits a Cholesky factorization, and that two semi-infinite matrices Z_1 and Z_2 are given, such that

i) $Z_1 \mathscr{E}(\Lambda, \eta)^{-1}$ is strictly lower triangular,

- ii) Z_2 is upper triangular and
- iii) $Z_1(\eta)G_0 = Z_2(\eta)$.

Then, we can ensure that $Z_1 = Z_2 = 0$.

Proof. We have

$$Z_1 G_0 = Z_1 \mathscr{E}(\Lambda; \boldsymbol{\eta})^{-1} G = Z_1 \mathscr{E}(\Lambda; \boldsymbol{\eta})^{-1} S^{-1} H S^{-\top}$$

and we get

$$Z_1 \mathscr{E}(\Lambda; \boldsymbol{\eta})^{-1} S^{-1} H S^{-\top} = Z_2.$$

Consequently,

$$Z_1 \mathscr{E}(\Lambda; \boldsymbol{\eta})^{-1} S^{-1} = Z_2 S H^{-1}.$$

But $Z_1 \mathscr{E}(\Lambda; \eta)^{-1} S^{-1}$ is strictly lower triangular while $Z_2 S H^{-1}$ is upper triangular. As both terms are equal the only possibility is that both are the zero matrix

$$Z_1 \mathscr{E}(\Lambda; \eta)^{-1} S^{-1} = Z_2 S H^{-1} = 0,$$

and we get the result.

Using this results one proves that

Proposition 20 (KP equation). The wave function Ψ_n satisfies the linear equations

$$\vartheta_m \Psi_n = \mathscr{P}_m(\vartheta_\eta) \Psi_n$$

with

$$\mathscr{P}_m(\vartheta_\eta) = \vartheta_\eta^m - mp_n^1 \vartheta_\eta^{m-2} + U_{n,m-3} \vartheta_\eta^{m-3} + \dots + U_{n,0}$$

where $\{U_{n,j}\}_{j=0}^{m-3}$ are polynomials in $\vartheta_{\eta}^{l}p_{n}^{1}, \vartheta_{2}p_{n}^{1}, \ldots, \vartheta_{m-1}p_{n}^{1}$. In particular, the following linear equations hold

$$\vartheta_2 \Psi_n = \vartheta_\eta^2 \Psi_n - 2\vartheta_\eta (p_n^1) \Psi_n,$$

$$\vartheta_3 \Psi_n = \vartheta_\eta^3 \Psi_n - 3\vartheta_\eta (p_n^1) \vartheta_\eta \Psi_n - \frac{3}{2} \Big(\vartheta_\eta^2 (p_k^1) + \vartheta_2 (p_k^1) \Big) \Psi_n,$$

and, consequently, its compatibility condition, which is the KP equation

$$\vartheta_{\eta} \Big(4\vartheta_3(p_n^1) + 6 \big(\vartheta_{\eta}(p_n^1) \big)^2 - \vartheta_{\eta}^3(p_n^1) \Big) = \vartheta_2^2(p_n^1),$$

is fulfilled.

Remark 8. The weight w for which this KP equation appears is

$$w(z) = \frac{(a_1)_z \cdots (a_M)_z}{\Gamma(z+1)(b_1)_z \cdots (b_N)_z} \eta^z \eta_2^{z^2} \eta_3^{z^3}, \qquad \qquad |\eta_2|, |\eta_3| < 1.$$

and the corresponding 0-th moment is the series

$$\rho_0 = \sum_{k=0}^{\infty} w(k) = \sum_{k=0}^{\infty} \frac{(a_1)_k \cdots (a_M)_k}{(b_1 + 1)_k \cdots (b_{N-1} + 1)_k} \frac{\eta^k \eta_2^{k^2} \eta_3^{k^3}}{k!}.$$

We see that only for $\eta_2 = \eta_3 = 1$ we recover a hypergeometric function.

2.8. Toda–Pearson compatibility. For the compatibility of (9) and (56), that is for the compatibility of the systems

$$\begin{cases} P(z+1) = \Pi P(z), \\ \vartheta_{\eta}(P(z)) = \Phi P(z). \end{cases}$$

we require

$$\begin{split} \vartheta_{\eta}(P(z+1)) &= \vartheta_{\eta}(\Pi P(z)) = \vartheta_{\eta}(\Pi)P(z) + \Pi\vartheta_{\eta}(P(z)) = (\vartheta_{\eta}(\Pi) + \Pi J\Phi)P(z), \\ (\vartheta_{\eta}P)(z+1) &= \Phi P(z+1) = \Phi \Pi P(z) \end{split}$$

to be equal and, consequently, we obtain

$$\vartheta_{\eta}(\Pi) = [\Phi, \Pi].$$

In the general case the dressed Pascal matrix Π is a lower unitriangular semi-infinite matrix, that possibly has an infinite number of subdiagonals. However, for the case when the weight $w(z) = v(z)\eta^z$ satisfies the Pearson equation (29), with v independent of η , that is

$$\theta(k+1)v(k+1)\eta^{k+1} = \sigma(k)v(k)\eta^k$$

for $k \in \mathbb{N}_0$; i.e.,

$$\theta(k+1)v(k+1)\eta = \sigma(k)v(k),$$

the situation improves as we have the banded semi-infinite matrix Ψ that models the shift in the z variable as in (41). From the previous discrete Pearson equation we see that $\sigma(z) = \eta \kappa(z)$ with $\kappa, \theta \eta$ -independent polynomials in z

(64)
$$\theta(k+1)v(k+1) = \eta \kappa(k)v(k)$$

Proposition 21. Let us assume a weight w satisfying the Pearson equation (29) and, consequently, of the form (33). Then, the Laguerre-Freud structure matrix Ψ given in (38) satisfies

(65a)
$$\vartheta_{\eta}(\eta^{-1}\Psi^{\top}H^{-1}) = [\Phi, \eta^{-1}\Psi^{\top}H^{-1}],$$

(65b)
$$\vartheta_{\eta}(\Psi H^{-1}) = [\Phi, \Psi H^{-1}].$$

Alternatively, the above equations can be written as follows

(66a)
$$\vartheta_{\eta}(\Psi^{\top}H^{-1}) = [J_{+}, \Psi^{\top}H^{-1}],$$

(66b)
$$\vartheta_{\eta}(\eta^{-1}\Psi H^{-1}) = [J_+, \eta^{-1}\Psi H^{-1}]$$

Relations (65a) and (65b) are gauge equivalent.

(67)
$$\begin{cases} \sigma(z)P(z+1) = \Psi^{\top}H^{-1}P(z), \\ \vartheta_{\eta}(P(z)) = \Phi P(z), \end{cases}$$

while for (65b) we consider

(68)
$$\begin{cases} \theta(z)P(z-1) = \Psi H^{-1}P(z), \\ \vartheta_{\eta}(P(z)) = \Phi P(z). \end{cases}$$

From (67), observing that $\vartheta_{\eta}(\sigma) = \sigma$, we deduce

$$\begin{split} \vartheta_{\eta}\big(\sigma(z)P(z+1)\big) &= \vartheta_{\eta}(\Psi^{\top}H^{-1}P(z)) = \vartheta_{\eta}(\Psi^{\top}H^{-1})P(z) + \Psi^{\top}H^{-1}\vartheta_{\eta}(P(z)) \\ &= \big(\vartheta_{\eta}(\Psi^{\top}H^{-1}) + \Psi^{\top}H^{-1}\Phi\big)P(z), \\ \vartheta_{\eta}\big(\sigma(z)P(z+1)\big) &= \sigma(z)P(z+1) + \sigma(z)\Phi P(z+1) = (\Phi+I)\Psi^{\top}H^{-1}P(z). \end{split}$$

so that

$$\vartheta_{\eta}(\Psi^{\top}H^{-1}) - \Psi^{\top}H^{-1} = \left[\Phi, \Psi^{\top}H^{-1}\right],$$

and recalling $\vartheta_{\eta}(\eta^{-1}) = -\eta^{-1}$ we obtain (65a).

From (68) we find $(\vartheta_\eta\theta=0)$

$$\begin{split} \vartheta_{\eta} \big(\theta(z) P(z-1) \big) &= \vartheta_{\eta} (\Psi H^{-1} P(z)) = \vartheta_{\eta} (\Psi H^{-1}) P(z) + \Psi H^{-1} \vartheta_{\eta} (P(z)) \\ &= \big(\vartheta_{\eta} (\Psi H^{-1}) + \Psi H^{-1} \Phi \big) P(z), \\ \vartheta_{\eta} \big(\theta(z) P(z-1) \big) &= \theta(z) \Phi P(z-1) = \Phi \Psi H^{-1} P(z). \end{split}$$

and (65b) follows immediately.

Notice that from (44) this equation (65b) follows immediately:

$$\vartheta_{\eta}(\Psi H^{-1}) = [(\vartheta_{\eta}S)S^{-1}, SB^{-1}\theta(\Lambda)S^{-1}] = [\Phi, \Psi H^{-1}].$$

This comment applies similarly to (65a).

Using $\Phi = -J_{-} = J_{+} - J$ in (65a) and (43a) we easily get (66a), and similarly from (65b) and (43b) we deduce (66b).

Finally, we study the relation between (65a) and (65b). Assume that (65b), $\vartheta_{\eta}(\Psi H^{-1}) = [\Phi, \Psi H^{-1}]$, holds and take its transpose to obtain $\vartheta_{\eta}(H^{-1}\Psi^{\top}) = -[\Phi^{\top}, H^{-1}\Psi^{\top}]$, which is not (65a). Observe that $\Psi^{\top}H^{-1} = H(H^{-1}\Psi^{\top})H^{-1}$

so that, the following gauge type transformation equation arises

$$\begin{split} \vartheta_{\eta}(\Psi^{\top}H^{-1}) &= \vartheta_{\eta}(H)(H^{-1}\Psi^{\top})H^{-1} + H\vartheta_{\eta}(H^{-1}\Psi^{\top})H^{-1} - H(H^{-1}\Psi^{\top})H^{-1}\vartheta_{\eta}(H)H^{-1} \\ &= \left[\vartheta_{\eta}(H)H^{-1}, H(H^{-1}\Psi^{\top})H^{-1}\right] - H[\Phi^{\top}, H^{-1}\Psi^{\top}]H^{-1} \\ &= \left[\vartheta_{\eta}(H)H^{-1} - H\Phi^{\top}H^{-1}, \Psi^{\top}H^{-1}\right]. \end{split}$$

Now, attending to (57), we have $\vartheta_{\eta}(H)H^{-1} - H\Phi^{\top}H^{-1} = J + \Phi$ so that

$$\vartheta_{\eta}(\Psi^{\top}H^{-1}) = \left[J + \Phi, \Psi^{\top}H^{-1}\right] = \left[\Phi, \Psi^{\top}H^{-1}\right] + \Psi^{\top}H^{-1},$$

and (65a) follows immediately.

3. CONCLUSIONS AND OUTLOOK

The use of the Gauss–Borel factorization of the moment matrix has been throughly used by Adler and van Moerbeke in their studies of integrable systems and orthogonal polynomials, see [2, 3, 4]. We have extended these ideas and applied them in different contexts, CMV orthogonal polynomials, matrix orthogonal polynomials, multiple orthogonal polynomials and multivariate orthogonal [6, 5, 7, 8, 9, 10, 11, 12, 13]. For an general overview see [38].

In this paper, we extended those ideas to the discrete world. In particular we applied this approach to the study of the consequences of the Pearson equation on the moment matrix and Jacobi matrices. For that description a new banded matrix is required, the Laguerre– Freud structure matrix that encodes the Laguerre–Freud relations for the recurrence coefficients. We have also found that the contiguous relations fulfilled generalized hypergeometric functions determining the moments of the weight described for the squared norms of the orthogonal polynomials a discrete Toda hierarchy known as Nijhoff–Capel equation, see [41].

Further work in this direction are the study of the role of Christoffel and Geronimus transformations for the description of the mentioned contiguous relations, and the use of the Geronimus–Christoffel transformations to characterize the shifts in the spectral independent variable of the orthogonal polynomials [39]. In [25] these ideas are applied to generalized Charlier, Meixner, and type I Hahn discrete orthogonal polynomials extending the results of [21, 43, 26, 27, 28].

For the future, we will study the type II generalized Hahn polynomials, and extend these techniques to multiple discrete orthogonal polynomials [14] and its relations with the transformations presented in [18] and quadrilateral lattices [20, 40],

ACKNOWLEDGMENTS

This work has its seed in several inspiring conversations with Diego Dominici during a research stay at Johannes Kepler University at Linz.

References

- Vsevolod E. Adler, Alexander I. Bobenko, Yuri B. Suris, *Classification of Integrable Discrete Equations of Octahedron Type*, International Mathematics Research Notices 2012 n°8 (2012) 1822–1889, DOI:10.1093/imrn/rnr083.
- [2] Mark Adler and Pierre van Moerbeke, Vertex operator solutions to the discrete KP hierarchy, Communications in Mathematical Physics 203 (1999) 185-210
- [3] —, Generalized orthogonal polynomials, discrete KP and Riemann–Hilbert problems, Communications in Mathematical Physics 207 (1999) 589-620.
- [4] _____, Darboux transforms on band matrices, weights and associated polynomials, International Mathematics Research Notices 18 (2001) 935-984.
- [5] Carlos Álvarez-Fernández, Ulises Fidalgo Prieto, and Manuel Mañas, Multiple orthogonal polynomials of mixed type: Gauss-Borel factorization and the multi-component 2D Toda hierarchy, Advances in Mathematics 227 (2011) 1451–1525.
- [6] Carlos Álvarez-Fernández, Manuel Mañas, Orthogonal Laurent polynomials on the unit circle, extended CMV ordering and 2D Toda type integrable hierarchies, Advances in Mathematics 240 (2013) 132-193
- [7] Carlos Álvarez-Fernández, Gerardo Ariznabarreta, Juan C. García-Ardila, Manuel Mañas, and Francisco Marcellán, Christoffel transformations for matrix orthogonal polynomials in the real line and the non-Abelian 2D Toda lattice hierarchy, International Mathematics Research Notices 2017 nº5 (2017) 1285-1341, DOI:10.1093/imrn/rnw027.
- [8] Gerardo Ariznabarreta, Juan C. García-Ardila, Manuel Mañas, and Francisco Marcellán, Matrix biorthogonal polynomials on the real line: Geronimus transformations, Bulletin of Mathematical Sciences 9 (2019) 195007 (68 pages) DOI:10.1142/S1664360719500073 (World Scientic) or DOI:10.1007/s13373-018-0128-y3 (Springer-Verlag).
- [9] Gerardo Ariznabarreta, Juan C. García-Ardila, Manuel Mañas, and Francisco Marcellán, *Non-Abelian integrable hierarchies: matrix biorthogonal polynomials and perturbations* Journal of Physics A: Mathematical & Theoretical **51** (2018) 205204.
- [10] Gerardo Ariznabarreta and Manuel Mañas, Matrix orthogonal Laurent polynomials on the unit circle and Toda type integrable systems, Advances in Mathematics 264 (2014) 396-463.
- [11] _____, Multivariate orthogonal polynomials and integrable systems, Advances in Mathematics 302 (2016) 628–739.
- [12] _____, Christoffel transformations for multivariate orthogonal polynomials, Journal of Approximation Theory 225 (2018) 242–283.
- [13] Gerardo Ariznabarreta, Manuel Mañas, and Alfredo Toledano, CMV Biorthogonal Laurent Polynomials: Perturbations and Christoffel Formulas, Studies in Applied Mathematics 140 (2018) 333–400.
- [14] Jorge Arvesú, Jonathan Coussement, and Walter Van Assche, Some discrete multiple orthogonal polynomials, Journal of Computational and Applied Mathematics 153 (2003).
- [15] R. A. Askey and Adri B. Olde Daalhuis, *Generalized hypergeometric function* (2010), in Olver, Frank W. J.; Lozier, Daniel M.; Boisvert, Ronald F.; Clark, Charles W. (eds.), NIST Handbook of Mathematical Functions, Cambridge University Press.
- [16] Jinho Baik, Thomas Kriecherbauer, Kenneth T.-R. McLaughlin, and Peter D. Miller, *Discrete Orthogonal Polynomials*, Annals of Mathematics Studies 164, Princeton University Press, 2007.
- [17] Richard Beals and Roderick Wong, Special functions and orthogonal polynomials, Cambridge Studies in Advanced Mathematics 153, Cambridge University Press, 2016.
- [18] Amílcar Branquinho, Ana Foulquié-Moreno, and Manuel Mañas, Multiple orthogonal polynomials on the step-line, arXiv:2106.12707 [CA] (2021).
- [19] Peter A. Clarkson, Recurrence coefficients for discrete orthonormal polynomials and the Painlevé equations, Journal of Physics A: Mathematical & Theoretical 46 (2013) 185205.
- [20] Adam Doliwa, Paolo Maria Santini, and Manuel Mañas, *Transformations of quadrilateral lattices*, Journal of Mathematical Physics 41 (2000) 944–990.
- [21] Diego Dominici, Laguerre-Freud equations for generalized Hahn polynomials of type I, Journal of Difference Equations and Applications 24 (2018) 916–940.
- [22] _____, Matrix factorizations and orthogonal polynomials, Random Matrices Theory Applications 9 (2020) 2040003, 33 pp.
- [23] Diego Dominici and Francisco Marcellán, Discrete semiclassical orthogonal polynomials of class one, Pacific Journal of Mathematics 268 nº2 (2012) 389-411.

- [24] ——, Discrete semiclassical orthogonal polynomials of class 2 in Orthogonal Polynomials: Current Trends and Applications, edited by E. Huertas and F. Marcellán, SEMA SIMAI Springer Series, 22 (2021) 103-169, Springer.
- [25] Itsaso Fernández-Irrisarri and Manuel Mañas, Pearson Equations for Discrete Orthogonal Polynomials: II. Generalized Hypergeometric Functions and Toda Equations, arXiv: 2107.02177 [CA] (2021).
- [26] Galina Filipuk and Walter Van Assche, Recurrence coefficients of generalized Charlier polynomials and the fifth Painlevé equation, Proceedings of American Mathematical Society 141 (2013) 551–62.
- [27] ——, Recurrence Coefficients of a New Generalization of the Meixner Polynomials, Symmetry, Integrability and Geometry: Methods and Applications (SIGMA) 7 (2011), 068, 11 pages.
- [28] _____, Discrete Orthogonal Polynomials with Hypergeometric Weights and Painlevé VI, Symmetry, Integrability and Geometry: Methods and Applications (SIGMA) 14 (2018), 088, 19 pages.
- [29] Géza Freud. On the coefficients in the recursion formulae of orthogonal polynomials, Proceedings of the Royal Irish Academy Section A 76 nº1 (1976) 1–6.
- [30] Jarmo Hietarinta, Nalini Joshi and Frank W. Nijhoff, Discrete Systems and Integrability, Cambridge Texts in Applied Mathematics, Cambridge University Press, 2016.
- [31] Mourad E. H.Ismail, Classical and Quantum Orthogonal Polynomails in One Variable, Encyclopedia of Mathematics and its Applications 98, Cambridge University Press, 2009.
- [32] Mourad E. H. Ismail and Walter Van Assche, Encyclopedia of Special Functions: The Askey-Bateman Project. Volume I: Univariate Orthogonal Polynomials, Edited by Mourad Ismail, Cambridge University Press, 2020.
- [33] Edmond Laguerre, Sur la réduction en fractions continues d'une fraction qui satisfait à une équation différentialle linéaire du premier ordre dont les coefficients sont rationnels. Journal de Mathématiques Pures et Appliquées 4^e série, tome **1** (1885) 135–165.
- [34] Alphonse P. Magnus, A proof of Freud's conjecture about the orthogonal polynomials related to $|x|\rho \exp(-x^{2m})$, for integer m, in "Orthogonal polynomials and applications (Bar-le-Duc, 1984)", Lecture Notes in Mathematics **1171** 362–372, Springer, 1985.
- [35] ——, On Freud's equations for exponential weights, Journal of Approximation Theory 46(1) (1986) 65–99.
- [36] _____, Painlevé-type differential equations for the recurrence coefficients of semi-classical orthogonal polynomials, Journal of Computational and Applied Mathematics 57 (1995) 215-237.
- [37] ——, Freud's equations for orthogonal polynomials as discrete Painlevé equations, in "Symmetries and integrability of difference equations (Canterbury, 1996)", London Mathematical Society Lecture Note Series 255 228–243, Cambridge University Press, 1999.
- [38] Manuel Mañas, Revisiting Biorthogonal Polynomials. An LU factorization discussion in Orthogonal Polynomials: Current Trends and Applications, edited by E. Huertas and F. Marcellán, SEMA SIMAI Springer Series, 22 (2021) 273-308, Springer.
- [39] —, Pearson Equations for Discrete Orthogonal Polynomials: III. Christoffel and Geronimus transformations, arXiv:2107.02918 [CA] (2021).
- [40] Manuel Mañas, Adam Doliwa, and Paolo Maria Santini, Darboux transformations for multidimensional quadrilateral lattices. I, Physics Letters A 232 (1997) 99–105.
- [41] Frank W. Nijhoff and Hans W. Capel, The direct linearisation approach to hierarchies of integrable PDEs in 2 + 1 dimensions: I. Lattice equations and the differential-difference hierarchies. Inverse Problems 6 (1990) 567-590.
- [42] Arthur F. Nikiforov, Sergei K. Suslov, and Vasilii B. Uvarov, *Classical Orhogonal Polynomials of a Discrete Variable*, Springer Series in Computational Physics, Springer, 1991.
- [43] Christophe Smet and Walter Van Assche, Orthogonal polynomials on a bi-lattice, Constructive Approximation 36 (2012) 215-242.
- [44] Walter Van Assche, Orthogonal Polynomials and Painlevé Equations, Australian Mathematical Society Lecture Series 27, Cambridge University Press, 2018.