Improved Bounds for Codes Correcting Insertions and Deletions

Kenji Yasunaga^{*}

December 14, 2023

Abstract

This paper studies the cardinality of codes correcting insertions and deletions. We give improved upper and lower bounds on code size. Our upper bound is obtained by utilizing the asymmetric property of list decoding for insertions and deletions and can be seen as analogous to the Elias bound in the Hamming metric. Our non-asymptotic bound is better than the existing bounds when the minimum Levenshtein distance is relatively large. The asymptotic bound exceeds the Elias and the MRRW bounds adapted from the Hamming-metric bounds for the binary and the quaternary cases. Our lower bound improves on the bound by Levenshtein, but its effect is limited and vanishes asymptotically.

1 Introduction

We study the existence of codes correcting insertions and deletions. We are interested in deriving upper and lower bounds on the cardinality of codes.

Levenshtein [9] gave asymptotic upper and lower bounds for codes correcting a constant number of insertions and deletions. Later, he presented bounds for correcting any number [11]. Following his work, there have been several studies [3, 7, 8] on the cardinality of codes. However, they mainly focused on codes correcting a constant number of insertions/deletions. Giving better bounds for codes correcting a constant *fraction* of insertions/deletions has been elusive. See [2] for a recent survey.

In this work, we present upper and lower bounds on the cardinality of codes correcting insertions and deletions. First, we give a non-asymptotic upper bound on the cardinality of codes correcting insertions/deletions. Asymptotically, it implies that for any code $C \subseteq \Sigma^n$ of rate R that can correct $\lfloor \delta n \rfloor - 1$ insertions/deletions, it holds that $R \leq (1 - H_q(\delta))/(1 - \delta)$, where $|\Sigma| = q, \ \delta \in [0, 1)$, and $H_q(\cdot)$ is the q-ary entropy function. The bound improves on the asymptotic upper bounds from the literature; The well-known Elias and MRRW

^{*}Tokyo Institute of Technology, Japan. E-mail: yasunaga@c.titech.ac.jp

bounds for the Hamming metric can also be employed as upper bounds in the Levenshtein metric (insertions and deletions). Our bound improves on them for q = 2, 4.

Our bound is obtained by a similar argument to the Elias bound in the Hamming metric. We use the list size upper bound of [6] for insertions and deletions. It is well-known that any s-deletion correcting code can correct any s_1 insertions and s_2 deletions with $s_1 + s_2 = s$. This symmetry in the unique decoding regime does not hold in the list decoding problem. In [6], it is proved that any code with a large Levenshtein distance enables list decoding such that the decoding radius of insertion is superior to that of deletion. We crucially use this property to derive our upper bound.

Next, we give a non-asymptotic lower bound on the cardinality of codes. Our bound improves on the bound by Levenshtein [11] by investigating that every deletion ball contains multiple words that are close to each other in the Levenshtein distance. Asymptotically, our lower bound is the same as in [11].

Finally, we compare our bounds with the existing bounds in the literature. As a nonasymptotic bound, our upper bound is tighter than the bounds in [8,11] when the minimum distance is relatively large. Our asymptotic upper bound is the best compared with the existing bounds from the Hamming metric for the binary and the quaternary codes. Regarding lower bounds, although our bound strictly improves the bound in [11], its effect is limited and vanishes asymptotically.

Related Work

The existence of positive rate codes correcting a constant fraction of insertions/deletions was presented by Schulman and Zuckerman [14]. Their asymptotic lower bounds were later improved by Levenshtein [11].

Cullina and Kiyavash [3] improved Levenshtein's upper bound [11] for correcting a constant number of insertions/deletions using a graph-theoretic approach. Kulkarni and Kiyavash [8] derived non-asymptotic upper bounds by a linear programming argument for graph-matching problems. They also gave upper bounds for correcting a constant fraction of insertions/deletions. Although their asymptotic bound (rate function) improved on the bound in [11], it was not given in the closed-form expressions.

For extreme cases where the deletion fraction is either small or high, Guruswami and Wang [5] gave efficient constructions of codes correcting these cases of deletions. For the case that the coding rate is nearly zero, Kash et al. [7] showed a positive-rate binary code correcting the fraction p of insertions/deletions with $p \ge 0.1737$, which improved on the bound of $p \ge 0.1334$ in [11]. Bukh, Guruswami, and Haståd [1] significantly improved on the previous results by showing the existence of a positive rate q-ary code with $p \ge$ $1 - (2/(q + \sqrt{q}))$, which is $p \ge 0.4142$ for the binary case. Guruswami, He, and Li [4] showed a slight but highly non-trivial improvement to the upper bound on the fraction of correctable insertions/deletions by codes with a non-zero coding rate. They proved that there is a constant $\epsilon > 0$ such that any q-ary code correcting a $(1 - (1 + \epsilon)/q)$ -fraction of insertions/deletions must have a rate approaching 0.

Maringer et al. [12] studied insertion/deletion correcting codes in feedback models and completely determined the maximal asymptotic rate for binary codes.

2 Code Size Upper Bound

Let Σ be a finite alphabet. The Levenshtein distance $d_{\mathrm{L}}(x, y)$ between two words x and y is the minimum number of symbol insertions and deletions needed to transform x into y. For a code $C \subseteq \Sigma^n$, its minimum Levenshtein distance is the minimum distance $d_{\mathrm{L}}(c_1, c_2)$ of every pair of distinct codewords $c_1, c_2 \in C$. Since any two codewords in C are of the same length, the minimum Levenshtein distance of C takes an even number. It is well-known that a code with minimum Levenshtein distance d can correct any s_1 insertions and s_2 deletions as long as $s_1 + s_2 \leq d/2 - 1$. The Levenshtein distance between two words in Σ^n takes integer values from 0 to 2n. Thus, we consider the normalized Levenshtein distance $\delta = d/2n$ in the analysis. The value $\delta \in [0, 1]$ also represents the fraction of insertions/deletions that can be corrected since we require $(s_1 + s_2)/n \leq (d/2 - 1)/n = \delta - 1/n$, which is asymptotically equal to δ .

Let $C \subseteq \Sigma^n$ be a code of minimum Levenshtein distance d with $|\Sigma| = q$. For a word $x \in \Sigma^n$, let $I_t(x) \subseteq \Sigma^{n+t}$ be the set of its supersequences of length n + t. Namely, $I_t(x)$ consists of words obtained from x by inserting t symbols. Similarly, let $D_t(x)$ be the set of words obtained from x by deleting t symbols. It is known [10] that the size of $I_t(x)$ does not depend on x and

$$|I_t(x)| = \sum_{i=0}^t \binom{n+t}{i} (q-1)^i \triangleq I_q(n,t).$$

First, we give a simple sphere packing bound. We use the fact that the number of supersequences, $|I_t(x)|$, is independent of x.

Theorem 1. Let $C \subseteq \Sigma^n$ be a code of minimum Levenshtein distance d and $|\Sigma| = q$. It holds that

$$|C| \le \left\lfloor \frac{q^{n+d/2-1}}{I_q(n,d/2-1)} \right\rfloor.$$
 (1)

Proof. For each codeword $c \in C$, consider the set of supersequences $I_{d/2-1}(c) \subseteq \Sigma^{n+d/2-1}$. Since the code has minimum Levenshtein distance d, each $I_{d/2-1}(c)$ should be disjoint. Thus,

$$\sum_{c \in C} |I_{d/2-1}(c)| \le q^{n+d/2-1}$$

The statement follows by the equality $|I_{d/2-1}(c)| = I_q(n, d/2 - 1)$ for any $c \in C$.

Next, we prove our main theorem, which can be seen as an Elias-type upper bound on the code size in the Levenshtein metric.

Theorem 2. Let $C \subseteq \Sigma^n$ be a code of minimum Levenshtein distance d < 2n and $|\Sigma| = q$. For any integer $t \ge 0$ with

$$t < \frac{nd}{2n-d},\tag{2}$$

it holds that

$$|C| \le \left\lfloor \frac{(n+t)d}{(n+t)d - 2nt} \cdot \frac{q^{n+t}}{I_q(n,t)} \right\rfloor.$$
(3)

Proof. By double counting, the sum of the cardinalities of $I_t(x)$ of all $x \in \Sigma^n$ is equal to the sum of the cardinalities of $D_t(y)$ of all $y \in \Sigma^{n+t}$. Namely,

$$\sum_{y \in \Sigma^{n+t}} |D_t(y)| = \sum_{x \in \Sigma^n} |I_t(x)| = q^n \cdot I_q(n,t).$$

$$\tag{4}$$

By considering the intersection with C,

$$\sum_{y \in \Sigma^{n+t}} |D_t(y) \cap C| = \sum_{x \in C} |I_t(x)| = |C| \cdot I_q(n,t).$$

Thus, by choosing $y \in \Sigma^{n+t}$ uniformly at random,

$$\mathbb{E}_{y}[|D_{t}(y) \cap C|] = \frac{1}{q^{n+t}} \sum_{y \in \Sigma^{n+t}} |D_{t}(y) \cap C| = \frac{|C| \cdot I_{q}(n,t)}{q^{n+t}}.$$

The averaging argument implies that there exists $y \in \Sigma^{n+t}$ such that

$$|D_t(y) \cap C| \ge \frac{|C| \cdot I_q(n,t)}{q^{n+t}}.$$
(5)

We have the following lemma.

Lemma 1. For any non-negative integer t with t < nd/(2n-d) and $y \in \Sigma^{n+t}$, it holds that

$$|D_t(y) \cap C| \le \frac{(n+t)d}{(n+t)d - 2nt}.$$

Proof. In [6], $B_{\mathsf{L}}(v, t_{\mathsf{I}}, t_{\mathsf{D}})$ is defined as the set of words that can be obtained from v by at most t_{I} insertions and at most t_{D} deletions. Then, for $C \subseteq \Sigma^n$ and $y \in \Sigma^{n+t}$, we can see that $B_{\mathsf{L}}(y, 0, t) \cap C = D_t(y) \cap C$. Lemma 1 of [6] with $t_{\mathsf{I}} = t$ and $t_{\mathsf{D}} = 0$ implies that if d/2 > tn/(n+t), $|D_t(y) \cap C| \le (n+t)(d/2)/((n+t)d/2 - tn)$ for every $y \in \Sigma^{n+t}$, which implies the statement.

By combining (5) and Lemma 1, the statement follows.

We analyze asymptotics of Theorems 1 and 2. For a code $C \subseteq \Sigma^n$ of distance d, let $\delta = d/2n$ and $\gamma = t/n$ for $t \ge 0$. Let $A_q(n,d)$ be the maximum size of code $C \subseteq \Sigma^n$ of minimum Levenshtein distance d, where $|\Sigma| = q$. For $\delta \in [0, 1]$, let

$$R_q(\delta) \triangleq \liminf_{n \to \infty} \frac{\log_q A_q(n, \lfloor 2\delta n \rfloor)}{n}$$

be the asymptotic coding rate achievable for normalized Levenshtein distance δ . Note that $R_q(\delta) = 0$ for $\delta \ge 1 - q^{-1}$ (See, for example, [1, Section 1]).

Let $\operatorname{Vol}_q(n, \ell)$ be the volume of the Hamming ball of radius ℓ in \mathbb{F}_q^n . Namely, $\operatorname{Vol}_q(n, \ell) = \sum_{i=0}^{\ell} {n \choose i} (q-1)^i$. It is well-known (cf. [13, Lemma 4.8]) that, for $0 \leq \ell \leq n$,

$$\operatorname{Vol}_q(n,\ell) \ge \frac{1}{n+1} \cdot q^{nH_q(\ell/n)},$$

where $H_q(x) = -x \log_q x - (1-x) \log_q (1-x) + x \log_q (q-1)$. Since $I_q(n,t) = \text{Vol}_q(n+t,t)$,

$$\frac{1}{n} \cdot \log_q I_q(n,t) \ge (1+\gamma)H_q\left(\frac{\gamma}{1+\gamma}\right) - \frac{\log_q((1+\gamma)n+1)}{n}$$

Regarding Theorem 1, it holds that

$$\frac{1}{n} \cdot \log_q I_q(n, d/2 - 1) \ge (1 + \delta) \cdot H_q\left(\frac{\delta}{1 + \delta} - o(1)\right) - o(1).$$

By Theorem 1, the rate R of C is bounded above by

$$R = \frac{\log_q |C|}{n} \le (1+\delta) \left(1 - H_q \left(\frac{\delta}{1+\delta} - o(1) \right) \right) + o(1).$$

Thus, we have the following corollary.

Corollary 1.

$$R_q(\delta) \le (1+\delta)\left(1 - H_q\left(\frac{\delta}{1+\delta}\right)\right).$$

Regarding Theorem 2, condition (2) can be rewritten as

$$\gamma < \frac{\delta}{1-\delta}.\tag{6}$$

Let $\gamma = \delta/(1-\delta) - 1/n$. The bound (3) can be rewritten as

$$C| \leq \frac{(1+\gamma)\delta}{(1+\gamma)\delta-\gamma} \cdot \frac{q^{(1+\gamma)n}}{I_q(n,t)}$$
$$= \frac{\delta/(1-\delta)-\delta/n}{(1-\delta)/n} \cdot \frac{q^{(1+\gamma)n}}{I_q(n,t)}$$
$$= q^{(1+\gamma)n\left(1-H_q\left(\frac{\gamma}{1+\gamma}\right)+o(1)\right)}.$$

Since

$$\frac{\gamma}{1+\gamma} = \frac{\delta/(1-\delta) - 1/n}{1/(1-\delta) - 1/n} = \delta - \frac{(1-\delta)^2}{n - (1-\delta)^2}$$

the rate R of C is bounded above by

$$R = \frac{\log_q |C|}{n} \le (1+\gamma) \left(1 - H_q \left(\frac{\gamma}{1+\gamma} \right) \right) + o(1)$$
$$\le \frac{1}{1-\delta} \left(1 - H_q \left(\delta - o(1) \right) \right) + o(1).$$

We obtain the following corollary.

Corollary 2.

$$R_q(\delta) \le \frac{1}{1-\delta} \left(1 - H_q(\delta)\right)$$

Bounds from Hamming-Metric Bounds

For a code $C \subseteq \Sigma^n$, let d_h be the minimum Hamming distance of C. It is well-known that the minimum Levenshtein distance d of C satisfies $d \leq 2d_h$. Thus, whenever we have an upper bound for a code of minimum Hamming distance d_h , we can use the same bound for a code of minimum Levenshtein distance $d_h/2$. More specifically, if a code $C \subseteq \Sigma^n$ satisfies $|C| \leq f(q, n, d_h)$, then $|C| \leq f(q, n, d/2)$ as long as f is a monotonically decreasing function on the third argument. Let $\delta = d/2n$ and $\delta_h = d_h/n$. Similarly, if we have a bound on the coding rate $R \leq g(q, \delta_h)$, then we have $R_q(\delta) \leq g(q, \delta)$ if g is monotonically decreasing on the second argument.

As far as we know, the best-known upper bounds for coding rate with respect to normalized minimum Levenshtein distance are obtained by the bounds for Hamming metric. The following bounds are well-known in the literature.

Theorem 3. For $0 \le \delta < \theta = 1 - q^{-1}$,

$$R_{q}(\delta) \leq 1 - H_{q}\left(\theta - \sqrt{\theta(\theta - \delta)}\right); \qquad (Elias \ bound)$$

$$R_{q}(\delta) \leq H_{q}\left(\frac{1}{q}\left(q - 1 - (q - 2)\delta - 2\sqrt{\delta(1 - \delta)(q - 1)}\right)\right). \qquad (MRRW \ bound)$$

3 Code Size Lower Bounds

Next, we consider lower bounds on $A_q(n,d)$ for codes $C \subseteq \Sigma^n$ with $|\Sigma| = q$. We assume that d is even. For $x \in \Sigma^n$ and non-negative integers t, s with $t \leq n$, let $L_{t,s}(x)$ be the set of words that can be obtained from x by deleting t symbols and inserting s symbols. By definition, it holds that $L_{t,0}(x) = D_t(x)$ and $L_{0,s}(x) = I_s(x)$.

We would like to derive an upper bound on the average size of $L_{t,t}(x)$ for $x \in \Sigma^n$ and $t \leq n$. This is because it gives a lower bound on $A_q(n,d)$ as discussed in [11, 15]. More specifically, for any $X \subseteq \Sigma^n$, let

$$V_t^{\text{ave}}(X) \triangleq \frac{1}{|X|} \sum_{x \in X} |L_{t,t}(x)|$$

be the average size of $L_{t,t}(x)$ in X. Then, there exists a code $C \subseteq \Sigma^n$ of minimum Levenshtein distance d satisfying

$$|C| \ge \frac{|X|}{V_{d/2-1}^{\text{ave}}(X)} = \frac{|X|^2}{\sum_{x \in X} |L_{d/2-1, d/2-1}(x)|}.$$
(7)

For $x = (x_1, \ldots, x_n) \in \Sigma^n$ and $i \in [1, n - 1]$, we say (x_i, x_{i+1}) is a distinct adjacent pair if $x_i \neq x_{i+1}$. Let $IP(x) = \{(i, i+1) : x_i \neq x_{i+1}\}$ be the index-pair set of distinct adjacent pairs in x. We say $P \subseteq IP(x)$ is disjoint if $P = \{(i_1, i_1 + 1), \ldots, (i_p, i_p + 1)\}$ satisfies $i_j + 1 \neq i_\ell$ for every distinct $j, \ell \in \{1, \ldots, p\}$. We denote by p(x) the maximum size of disjoint index-pair sets of distinct adjacent pairs in x. For example, for x = 01101, we have $IP(x) = \{(1, 2), (3, 4), (4, 5)\}$. The sets $\{(1, 2), (3, 4)\}$ and $\{(1, 2), (4, 5)\}$ are disjoint, but $\{(3, 4), (4, 5)\}$ is not. We have p(01101) = 2. For consistency, we consider the leftmost maximum-sized disjoint index-pair set in x. In the case that x = 01101, the leftmost one is $\{(1, 2), (3, 4)\}$. Similarly, for y = 010100 and y' = 010111, the (leftmost) maximum-sized disjoint index-pair set is $\{(1, 2), (3, 4)\}$, and hence p(y) = p(y') = 2.

We observe that for any positive integer $p' \leq p(x)$ satisfying $p' \leq \min\{t, n-t\}$, there are $2^{p'}$ words in $D_t(x)$ such that they are within a Levenshtein distance of 2p' from each other. The reason is as follows. Since $p' \leq p(x)$, there are n - 2p' indices in $\{1, \ldots, n\}$ such that they are not contained in a disjoint index-pair set of distinct adjacent pairs in x. First, we delete t - p' symbols from x out of the n - 2p' indices. This procedure requires that $p' \leq \min\{t, n-t\}$. Let $y \in \Sigma^{n-t+p'}$ be the resulting word. Second, we delete one of the two symbols in p' distinct adjacent pairs from y. There are $2^{p'}$ possible deletion patterns, resulting in all different words of length n - t. Since each resulting word and y have Levenshtein distance p', the $2^{p'}$ words are within a Levenshtein distance of 2p'. In other words, $D_t(x)$ contains $2^{p'}$ words $\{z_1, \ldots, z_{2p'}\}$ such that all are subsequences of $y \in \Sigma^{n-t+p'}$. We have

$$\begin{vmatrix} 2^{p'} \\ \bigcup_{i=1}^{2^{p'}} I_t(z_i) \end{vmatrix} \le |I_t(z_1)| + \left| \bigcup_{i=2}^{2^{p'}} (I_t(z_i) \setminus I_{t-p'}(y)) \right| \\ \le I_q(n-t,t) + (2^{p'}-1)(I_q(n-t,t) - I_q(n-t+p',t-p')) \\ = 2^{p'} \cdot I_q(n-t,t) - (2^{p'}-1) \cdot I_q(n-t+p',t-p'). \end{aligned}$$

Let $Z = \{z_1, \ldots, z_{2^{p'}}\}$. For any integer $p' \le p(x)$ with $p' \le \min\{t, n-t\}$, it holds that

$$\begin{aligned} |L_{t,t}(x)| &= \left| \bigcup_{z \in D_t(x)} I_t(z) \right| \\ &\leq \sum_{z \in D_t(x) \setminus Z} |I_t(z)| + \left| \bigcup_{z \in Z} I_t(z) \right| \\ &\leq (|D_t(x)| - 2^{p'}) \cdot I_q(n - t, t) + 2^{p'} \cdot I_q(n - t, t) - (2^{p'} - 1) \cdot I_q(n - t + p', t - p') \\ &= |D_t(x)| \cdot I_q(n - t, t) - (2^{p'} - 1) \cdot I_q(n - t + p', t - p') \triangleq L^*_{t,p'}(x). \end{aligned}$$

Note that p' should satisfy $p' \leq \min\{p(x), t, n-t\}$. Since $x \in \Sigma^n$ may have p(x) = 0, it is impossible to choose p' > 0 such that $p' \leq p(x)$ for every $x \in \Sigma^n$. Thus, we consider a subset $X \subseteq \Sigma^n$ and its partition $X = X_0 \cup X_1$ such that $X_1 = \{x \in X : p(x) \ge p'\}$ for some $p' \in [0, \min\{t, n-t\}]$ with p' > 0. Then, as an upper bound on $|L_{t,t}(x)|$, we can use $L_{t,p'}^*(x)$ for $x \in X_1$ and $|D_t(x)| \cdot I_q(n-t,t)$ for $x \in X_0$.

For integers n and $p \in [0, n/2]$, let $N_{n,q}(p)$ be the number of words $x \in \Sigma^n$ such that p(x) = p, where $|\Sigma| = q$. The value of $N_{n,q}(p)$ is given by Theorem 4 below. When X_1 is given by $\{x \in \Sigma^n : p(x) \ge p'\}$, it holds that $|X_1| = \sum_{p=p'}^{\lfloor n/2 \rfloor} N_{n,q}(p)$. For a partition $X = X_0 \cup X_1$ with $X_1 = \{x \in \Sigma^n : p(x) \ge p'\}$, we have that

$$\sum_{x \in X} |L_{t,t}(x)| \leq \sum_{x \in X} L_{t,p'}^*(x)$$

$$\leq \sum_{x \in X} |D_t(x)| \cdot I_q(n-t,t) - \sum_{x \in X_1} (2^{p'}-1) \cdot I_q(n-t+p',t-p')$$

$$= \sum_{x \in X} |D_t(x)| \cdot I_q(n-t,t) - (2^{p'}-1) \cdot I_q(n-t+p',t-p') \sum_{p=p'}^{\lfloor n/2 \rfloor} N_{n,q}(p). \quad (8)$$

We use (8) for our lower bound of Theorem 5.

We determine $N_{n,q}(p)$ for $p \in [0, n/2]$.

Theorem 4. For integers n > 0, $p \in [0, n/2]$, and $q \ge 2$,

$$N_{n,q}(p) = \binom{n-p}{p} \frac{q^p (q-1)^p}{n-p} (p+q(n-2p)).$$

Proof. We count the number of words $x \in \Sigma^n$ such that p(x) = p. When p(x) = 0, x should be a repetition of the same symbol in Σ . Thus, $N_{n,q}(0) = q$. Hereafter, we assume p > 0. Let $(i_1, i_1 + 1), \ldots, (i_p, i_p + 1) \in \{1, \ldots, n\}^2$ be p index-pairs of distinct adjacent pairs such that $i_1 \leq \cdots \leq i_p$. Let $(a_1, a'_1), \ldots, (a_p, a'_p) \in \Sigma^2$ be the corresponding distinct adjacent pairs, where their concrete values are yet determined. Since the order of p pairs (a_i, a'_i)

is fixed, we can construct all words x with p(x) = p by inserting n - 2p symbols into the word $a_1a'_1a_2a'_2\cdots a_pa'_p$. There are p+1 possible places to which symbols can be inserted. Namely, the resulting word should be of the form $w_1a_1a'_1w_2a_2a'_2w_3\cdots w_pa_pa'_pw_{p+1}$, where $w_i \in \Sigma^*$. Here, we consider the leftmost maximum-sized disjoint index-pair set in x. By the property of distinct adjacent pairs, w_1 must be the empty string or repetition of a_1 . Similarly, for $i \in \{2, \ldots, p\}$, w_i must be the empty string or repetition of a_i . Since we consider the *leftmost* disjoint index-pairs, different from the previous cases, w_{p+1} can be the empty string or repetition of any fixed symbol in Σ .

First, suppose that w_{p+1} is the empty string. Then the number of words x with p(x) = pis determined by the number of possible lengths of w_1, \ldots, w_p and possible combinations (a_i, a'_i) . Since the rightmost symbols are fixed to be $a_p a'_p$, the number of possible lengths is equal to the the number of ways of selecting p-1 items from n-2p+(p-1) items, which is $\binom{n-p-1}{p-1}$. There are q(q-1) combinations for each (a_i, a'_i) . Hence, the number $N_{n,q}(p)$ when $|w_{p+1}| = 0$ is equal to $q^p(q-1)^p\binom{n-p-1}{p-1}$. Second, consider the case that $|w_{p+1}| > 0$. In this case, we count the number of possible lengths of $w_1, \ldots, w_p, w_{p+1}$ and possible combinations (a_i, a'_i) . Since $|w_{p+1}| > 0$, we can assume that one symbol (length) has been assigned to w_{p+1} . Then, the number of possible lengths is equal to the number of ways of selecting p items from n-2p-1+p items, which is $\binom{n-p-1}{p}$. As in the previous case, the number of combinations for (a_i, a'_i) 's are $(q(q-1))^p$, and there are q possible symbols corresponding to w_{p+1} . Thus, the number $N_{n,q}(p)$ when $|w_{p+1}| > 0$ is $q^{p+1}(q-1)^p\binom{n-p-1}{p}$.

Therefore, for $p \ge 1$, we have

$$N_{n,q}(p) = q^{p}(q-1)^{p} \binom{n-p-1}{p-1} + q^{p+1}(q-1)^{p} \binom{n-p-1}{p}$$
$$= q^{p}(q-1)^{p} \binom{n-p}{p} \frac{p}{n-p} + q^{p+1}(q-1)^{p} \binom{n-p}{p} \frac{n-2p}{n-p}$$
$$= \binom{n-p}{p} \frac{q^{p}(q-1)^{p}}{n-p} (p+q(n-2p)).$$

When p = 0, the above gives $N_{n,q}(0) = q$. Hence, the statement follows.

By using (7) and (8), we have the following theorem.

Theorem 5. For a set $X \subseteq \Sigma^n$ and its partition $X = X_0 \cup X_1$ such that $X_1 = \{x \in \Sigma^n : p(x) \ge p'\}$ for some $p' \in [0, \min\{t, n-t\}]$, it holds that

$$A_q(n,d) \ge \left\lfloor \frac{|X|^2}{\sum_{x \in X} |D_t(x)| \cdot I_q(n-t,t) - (2^{p'}-1) \cdot I_q(n-t+p',t-p') \sum_{p=p'}^{\lfloor n/2 \rfloor} N_{n,q}(p)} \right\rfloor,$$

where t = d/2 - 1.

We give an explicit expression by choosing $X = \Sigma^n$ and p' = 1,

Corollary 3.

$$A_q(n,d) \ge \left\lfloor \frac{q^n}{q^{-t} \cdot I_q(n-t,t)^2 - (1-q^{-n+1})I_q(n-t+1,t-1)} \right\rfloor,$$

where t = d/2 - 1 > 0.

Proof. When $X = \Sigma^n$, it follows from (4) that $\sum_{x \in X} |D_t(x)| \cdot I_q(n-t,t) = q^{n-t} \cdot I_q(n-t,t)^2$. By choose p' = 1, since $N_{n,q}(0) = q$ and $\sum_{p=0}^{\lfloor n/2 \rfloor} N_{n,q}(p) = q^n$, we have

$$(2^{p'}-1) \cdot I_q(n-t+p',t-p') \sum_{p=p'}^{\lfloor n/2 \rfloor} N_{n,q}(p) = (q^n-q) \cdot I_q(n-t+1,t-1).$$

By combining the above and Theorem 5, the statement follows.

Since Levenshtein [11] gave a lower bound of $A_q(n,d) \ge q^n/(q^{-t} \cdot I_q(n-t,t)^2)$ for t = d/2 - 1, Corollary 3 improves his lower bound. Its effect, however, vanishes when d becomes large; Thus, the asymptotic lower bound on the coding rate given by Corollary 3 is the same as [11].

4 Comparison

First, we compare code-size upper bounds given by [8,11] and Theorems 1 and 2. Table 1 displays numerical values for several parameters q, n, and d. Specifically, we use the bounds in [11, Theorem 2], where we take the minimum over parameter r, and [8, Corollary 4.2]. Though the non-asymptotic bound of [8] achieves the best when d is small such as 4 and 10, Theorem 2 is much better when d is relatively large.

Next, we investigate asymptotic behaviors. Figures 1 and 2 show asymptotic bounds on the rate function $R_q(\delta)$ for q = 2 and q = 4, respectively. Corollary 2 gives the best upper bounds in both cases. For larger q, Corollary 2 is inferior to the MRRW bound, especially for large δ . Regarding lower bounds, [11] gave a bound (displayed as Levenshtein02), and [1] showed that there exist q-ary codes of rate R that achieve $\delta \ge 1 - (2/(q + \sqrt{q}))$ for sufficiently small R > 0 (displayed as BGH17). Also, it is proved in [4] that there is a small constant $\epsilon > 0$ such that R = 0 for $\delta \ge 1 - (1 + \epsilon)/q$ (omitted in the figures). The asymptotic upper bounds of [8, 11] were not given in the closed form. Hence it is not easy to make a clear comparison. By comparing with the plotted bound of [8, Fig. 1], we can see that Corollary 2 gives tighter bounds on $\delta \ge 0.1$ for q = 2 and $\delta \ge 0.2$ for q = 4.

Table 2 shows the numerical values of the lower bounds of [11, Theorem 1] and Corollary 3. As we can see, the improvement by Corollary 3 is small.

Table 1: Upper Bounds on Code Size

q	n	d	UB of [11]	UB of [8]	Theorem 1	Theorem 2
2	20	4	97 453	55 206	95 325	181 643
2	20	10	26 456	2535	$1 \ 295$	2 452
2	20	20	$190 \ 416$	1 059	32	28
2	20	30	$961\ 048$	—	5	4
2	40	4	$47 \ 498 \ 012 \ 376$	$28\ 192\ 605\ 878$	$52 \ 357 \ 696 \ 560$	$102\ 167\ 009\ 660$
2	40	10	$1\ 279\ 636\ 864$	$9\ 880\ 934$	$117 \ 292 \ 187$	$228 \ 473 \ 245$
2	40	20	$1\ 122\ 371\ 648$	$3\ 203\ 459$	215 900	203 859
2	40	30	$13\ 097\ 807\ 352$	298 539	3 735	1 195
2	40	40	$287\ 193\ 094\ 240$	$1 \ 048 \ 713$	231	43
4	20	4	$30\ 003\ 945\ 118$	$19\ 289\ 677\ 788$	$68\ 719\ 476\ 736$	$90\ 174\ 299\ 388$
4	20	10	$360\ 221\ 648$	$25 \ 002 \ 768$	$306 \ 647 \ 351$	$316\ 287\ 316$
4	20	20	$536\ 774\ 720$	$1 \ 645 \ 315$	$1\ 258\ 226$	79 926
4	20	30	$192\ 278\ 071\ 952$		$34 \ 771$	71
4	40	4	≈ 14 843 $\times 10^{18}$	$pprox 10$ 332 $ imes 10^{18}$	$pprox 38$ 997 $ imes 10^{18}$	\approx 51 574 $\times 10^{18}$
4	40	10	$pprox 6 \ 113 \ imes 10^{15}$	$\approx 461 \ 805 \ \times 10^{12}$	$pprox 27$ 238 $ imes 10^{15}$	$pprox$ 36 015 $ imes 10^{15}$
4	40	20	≈ 133 526 $\times 10^{12}$	$\approx 158 \ 374 \ \times 10^9$	$pprox 7$ 269 $ imes 10^{12}$	$\approx 2 \ 172 \ \times 10^{12}$
4	40	40	$\approx 34~641 \times 10^{15}$	$\approx 2 \ 123 \ \times 10^9$	$\approx 173 \ 431 \ \times 10^6$	$\approx 69 \times 10^6$
4	40	60	$\approx 306~026~\times 10^{18}$	—	$164 \ 423 \ 496$	108

Table 2: Lower Bounds on Code Size

q	n	d	LB of [11]	Corollary 3
2	20	6	94	94
2	20	8	4	4
2	40	6	$6\ 524\ 894$	$6\ 526\ 482$
2	40	8	76 814	76 818
2	40	10	1 687	1 687
4	20	6	$5\ 608\ 964$	$5\ 610\ 710$
4	20	8	$66 \ 412$	66 419
4	40	6	$379 \ 316 \ 355 \ 894 \ 427 \ 152$	$379\ 330\ 757\ 315\ 377\ 297$
4	40	8	$1 \ 031 \ 317 \ 510 \ 055 \ 795$	$1 \ 031 \ 323 \ 792 \ 762 \ 824$
4	40	10	$5\ 251\ 871\ 945\ 006$	$5\ 251\ 878\ 194\ 182$




Figure 1: Bounds for q = 2

5 Conclusions

This paper has presented improved upper and lower bounds on code size for correcting insertions and deletions. In particular, our upper bound improves the existing bounds in both non-asymptotic and asymptotic senses. An interesting future work is to develop an upper bound superior to the MRRW bound for large alphabet size q and relative distance δ . Our bound is inferior in some range for $q \geq 5$. A key lemma may be a list-size upper bound of insertions/deletion (as Lemma 1) employing q explicitly in the bound.

Acknowledgment

This work was supported in part by JSPS Grants-in-Aid for Scientific Research Number 18K11159.

References

[1] B. Bukh, V. Guruswami, and J. Håstad. An improved bound on the fraction of correctable deletions. *IEEE Trans. Inf. Theory*, 63(1):93–103, 2017.




Figure 2: Bounds for q = 4

- [2] M. Cheraghchi and J. L. Ribeiro. An overview of capacity results for synchronization channels. *IEEE Trans. Inf. Theory*, 67(6):3207–3232, 2021.
- [3] D. Cullina and N. Kiyavash. An improvement to Levenshtein's upper bound on the cardinality of deletion correcting codes. *IEEE Trans. Inf. Theory*, 60(7):3862–3870, 2014.
- [4] V. Guruswami, X. He, and R. Li. The zero-rate threshold for adversarial bit-deletions is less than 1/2. In 62nd IEEE Annual Symposium on Foundations of Computer Science, FOCS 2021, Denver, CO, USA, February 7-10, 2022, pages 727–738. IEEE, 2021.
- [5] V. Guruswami and C. Wang. Deletion codes in the high-noise and high-rate regimes. *IEEE Trans. Information Theory*, 63(4):1961–1970, 2017.
- [6] T. Hayashi and K. Yasunaga. On the list decodability of insertions and deletions. *IEEE Trans. Inf. Theory*, 66(9):5335–5343, 2020.
- [7] I. A. Kash, M. Mitzenmacher, J. Thaler, and J. R. Ullman. On the zero-error capacity threshold for deletion channels. In *Information Theory and Applications Workshop*, *ITA 2011, San Diego, California, USA, February 6-11, 2011*, pages 285–289. IEEE, 2011.

- [8] A. A. Kulkarni and N. Kiyavash. Nonasymptotic upper bounds for deletion correcting codes. *IEEE Trans. Inf. Theory*, 59(8):5115–5130, 2013.
- [9] V. I. Levenshtein. Binary codes capable of correcting deletions, insertions, and reversals. Soviet Physics Doklady, 10(8):707-710, 1966.
- [10] V. I. Levenshtein. Efficient reconstruction of sequences from their subsequences or supersequences. J. Comb. Theory, Ser. A, 93(2):310–332, 2001.
- [11] V. I. Levenshtein. Bounds for deletion/insertion correcting codes. In Proceedings IEEE International Symposium on Information Theory, page 370, 2002.
- [12] G. Maringer, N. A. Polyanskii, I. V. Vorobyev, and L. Welter. Feedback insertiondeletion codes. *Probl. Inf. Transm.*, 57(3):212–240, 2021.
- [13] R. M. Roth. Introduction to coding theory. Cambridge University Press, 2006.
- [14] L. J. Schulman and D. Zuckerman. Asymptotically good codes correcting insertions, deletions, and transpositions. *IEEE Trans. Information Theory*, 45(7):2552–2557, 1999.
- [15] L. M. G. M. Tolhuizen. The generalized Gilbert-Varshamov bound is implied by Turan's theorem. *IEEE Trans. Inf. Theory*, 43(5):1605–1606, 1997.